

APPLICATION OF MORSE INDEX IN WEAK FORCE N -BODY PROBLEM

GUOWEI YU

ABSTRACT. Due to collision singularities, the Lagrange action functional of the N -body problem in general is not differentiable. Because of this, the usual critical point theory can not be applied to this problem directly. Following ideas from [4], [23] and [2], we introduce a notion called weak critical point for such an action functional, as a generalization of the usual critical point. A corresponding definition of Morse index for such a weak critical point will also be given. Moreover it will be shown that the Morse index gives an upper bound of the number of possible binary collisions in a weak critical point of the N -body problem with weak force potentials including the Newtonian potential.

1. INTRODUCTION

The motion of N point masses, $m_i > 0, i \in \mathbf{N} := \{1, \dots, N\}$, under the universal gravitational force is a classic problem that has been studied by many authors since the time of Newton. Let $q_i \in \mathbb{R}^d$ be the position of mass m_i , then it satisfies the following equation

$$(1) \quad m_i \ddot{q}_i = \frac{\partial U(q)}{\partial q_i} = -\alpha \sum_{j \in \mathbf{N} \setminus \{i\}} m_i m_j \frac{q_i - q_j}{|q_i - q_j|^{\alpha+2}}, \quad \forall i \in \mathbf{N},$$

where $q = (q_i)_{i \in \mathbf{N}} \in \mathbb{R}^{dN}$ and $U(q) = \sum_{\{i < j\} \subset \mathbf{N}} \frac{m_i m_j}{|q_i - q_j|^\alpha}$ is the potential function (the negative potential energy). α here is a positive constant.

Traditionally U is called a *strong force potential*, when $\alpha \geq 2$, and a *weak force potential*, when $0 < \alpha < 2$. The Newtonian potential is a weak force potential corresponding to $\alpha = 1$.

This is a singular Lagrange system with the Lagrangian

$$L(q, \dot{q}) := K(\dot{q}) + U(q), \quad \text{where } K(\dot{q}) = \frac{1}{2} \sum_{i \in \mathbf{N}} m_i |\dot{q}_i|^2.$$

The singularities are caused by collisions between two or more masses

$$(2) \quad \Delta := \{q \in \mathbb{R}^{dN} : q_i = q_j, \text{ for some } \{i \neq j\} \subset \mathbf{N}\}.$$

For any $\mathbf{I} \subset \mathbf{N}$ with $|\mathbf{I}| \geq 2$ ($|\mathbf{I}|$ is the cardinality of \mathbf{I}), we say q has an *\mathbf{I} -cluster collision* at the moment t , when

$$\forall i \in \mathbf{I}, \quad \begin{cases} q_i(t) = q_j(t), & \text{if } j \in \mathbf{I}, \\ q_i(t) \neq q_j(t), & \text{if } j \in \mathbf{N} \setminus \mathbf{I}, \end{cases}$$

An \mathbf{I} -cluster collision is a *binary collision*, when $|\mathbf{I}| = 2$.

The author acknowledges the support of the ERC Advanced Grant 2013 No. 339958 "Complex Patterns for Strongly Interacting Dynamical Systems - COMPAT".

Let $H^1([T_1, T_2], \mathbb{R}^{dN})$ be the space of Sobolev paths defined on $[T_1, T_2]$, and $\hat{\mathbb{R}}^{dN} := \mathbb{R}^{dN} \setminus \Delta$ the set of collision-free configurations, we say a path $q \in H^1([T_1, T_2], \mathbb{R}^{dN})$ is *collision-free*, if $q(t) \in \hat{\mathbb{R}}^{dN}$, for any $t \in [T_1, T_2]$. It is well known the Lagrange action functional

$$(3) \quad \mathcal{A}(q; T_1, T_2) := \int_{T_1}^{T_2} L(q, \dot{q}) dt, \quad \forall q \in H^1([T_1, T_2], \mathbb{R}^{dN}),$$

is C^2 on $H^1([T_1, T_2], \hat{\mathbb{R}}^{dN})$ (see [1]), and a critical point of \mathcal{A} in $H^1([T_1, T_2], \hat{\mathbb{R}}^{dN})$ is a classical solution of (1).

The solution of (1) is invariant under linear translations, in many cases it will be more convenient to fix the center of mass at the origin, so we set

$$\mathcal{X} := \{q \in \mathbb{R}^{dN} : \sum_{i \in \mathbf{N}} m_i q_i = 0\}, \quad \hat{\mathcal{X}} := \mathcal{X} \setminus \Delta.$$

As the action functional is also invariant under linear translation, a critical point of \mathcal{A} in $H^1([T_1, T_2], \hat{\mathcal{X}})$ will be a classical solution of (1) as well.

In general it is much easier to apply variational methods to the N -body, when the potential is a *strong force*, i.e. $\alpha \geq 2$, as in this case any path with a finite action value must be collision-free, see [9]. It is not so, when the potential is a *weak force*, i.e. $\alpha \in (0, 2)$, as the attracting force between the masses are too weak and the action value of a path with collision may still be finite, see [13].

Because of this, for the strong force N -body problem, many results have been obtained by different authors using both minimization and non-minimization variational methods, see [4], [1], [17], [18], [22], [9] and the references within. However the problem is much more difficult for weak force potentials due to the possibility of collision. Set back by this, Bahri and Rabinowitz introduced the so called *generalized solution* in [3] and [4] (see Definition 2.1), where such a solution is allowed to have a non-empty set of collision moments with zero Lebesgue measure. Here we are only interested in the weak force N -body problem, so we assume $\alpha \in (0, 2)$ in the rest of the paper.

For action minimization methods, the breakthrough followed the proofs of the Figure-Eight solution of the three body by Chenciner and Montgomery [10] and the Hip-Hop solution of the four body by Chenciner and Venturelli [11], where both solutions are found as collision-free minimizers of the action functional under proper symmetric constraints. Since then action minimization methods have thrived in the study of the N -body problem with Newtonian potential as well as other weak force potentials. We refer the interesting readers to [8], [12], [7], [27], [28] and the references within.

Now it is more or less well understood, when we can show an action minimizer is collision-free.

Definition 1.1. $q \in H^1([T_1, T_2], \mathcal{X})$ is a **local action minimizer of \mathcal{A} with fixed-ends** in $H^1([T_1, T_2], \mathcal{X})$, if there is a $\delta > 0$ small enough, such that

$$\mathcal{A}(q; T_1, T_2) \leq \mathcal{A}(q + \tilde{q}; T_1, T_2), \quad \forall \tilde{q} \in H_0^1([T_1, T_2], \mathcal{X}) \text{ with } \|\tilde{q}\|_{H^1} \leq \delta,$$

where $H_0^1([T_1, T_2], \mathcal{X})$ is the space of Sobolev paths with compact support in $[T_1, T_2]$.

The following fundamental result is due to Marchal [19] and Chenciner [8], when $\alpha = 1$, and Ferrario and Terracini [12], when $\alpha \in (0, 2)$.

Theorem 1.1. *For any $\alpha \in (0, 2)$, when $d \geq 2$, if q is a local action minimizer of \mathcal{A} with fixed-ends in $H^1([T_1, T_2], \mathcal{X})$, then $q(t)$ is collision-free, for any $t \in (T_1, T_2)$.*

Despite of the above progress, by our knowledge, for the N -body problem no result seems to be available regarding how to rule out collision when the corresponding path is obtained through non-minimization methods, like minimax or mountain-pass. Meanwhile for a special type of singular Lagrange systems with weak force potentials (essentially equivalent to perturbations of the N center problem), using *Morse index theory*, in a series of papers ([23], [24], [25], [26]), Tanaka showed how to rule out collisions when a critical point was obtained by the minimax approach of Bahri and Rabinowitz [3]. The main purpose of our paper is to generalize Tanaka's idea to the N -body problem and show that the Morse index of a critical point can be used to give an upper bound of the number of binary collisions that could occur in it.

To give the precise statement, we recall the following definition according to [2].

Definition 1.2. Let X be a Hilbert space and Ω an open and dense subspace of it, i.e. $\bar{\Omega} = X$. If a functional \mathcal{F} is lower semi-continuous in X and C^2 in Ω , then $c \in \mathbb{R}$ will be called a **critical value** of \mathcal{F} , if there is $q \in \Omega$, such that $\mathcal{F}(q) = c$ and the first derivative of \mathcal{F} vanishes at q , i.e. $d\mathcal{F}(q) = 0$. Moreover such a q will be called a **critical point** of \mathcal{F} .

If $q \in \Omega$ is a critical point of \mathcal{F} , we define its **Morse index** (with respect to \mathcal{F}), $m_{\bar{\Omega}}^-(q, \mathcal{F})$, as the dimension of the largest subspace of Ω , where the second derivative $d^2\mathcal{F}(q)$ is negative definite.

The above definition suits the study of the N -body problem, as $H^1([T_1, T_2], \hat{\mathcal{X}})$ is an open and dense subspace of $H^1([T_1, T_2], \mathcal{X})$ with the action functional \mathcal{A} being C^2 in $H^1([T_1, T_2], \hat{\mathcal{X}})$ and lower semi-continuous in $H^1([T_1, T_2], \mathcal{X})$. Since \mathcal{A} is generally not differentiable at a collision path, such a path can not be a *critical point* and moreover it does not have a well-defined Morse index for a collision path. Although a collision path can still be a *local minimizer*.

To deal with the above problem, following ideas from [3], [23] and [2], let's perturb the weak force N -body problem (1) by a strong force potential

$$(4) \quad \varepsilon \mathcal{U}(q) := \varepsilon \sum_{\{i < j\} \subset \mathbf{N}} \frac{m_i m_j}{|q_i - q_j|^2}, \quad \text{for } \varepsilon > 0 \text{ small enough.}$$

Then the motion of masses satisfies

$$(5) \quad m_i \ddot{q}_i = -\alpha \sum_{j \in \mathbf{N} \setminus \{i\}} \frac{m_i m_j (q_i - q_j)}{|q_i - q_j|^{\alpha+2}} - 2\varepsilon \sum_{j \in \mathbf{N} \setminus \{i\}} \frac{m_i m_j (q_i - q_j)}{|q_i - q_j|^4}, \quad \forall i \in \mathbf{N},$$

which is the Euler-Lagrange equation of the action functional

$$(6) \quad \mathcal{A}^\varepsilon(q; T_1, T_2) := \int_{T_1}^{T_2} L^\varepsilon(q, \dot{q}) dt, \quad \text{where } L^\varepsilon(q, \dot{q}) := L(q, \dot{q}) + \varepsilon \mathcal{U}(q).$$

Furthermore we set $\mathcal{A}^0(q; T_1, T_2) = \mathcal{A}(q; T_1, T_2)$. When $\varepsilon > 0$, any path with a finite action value of \mathcal{A}^ε must be collision-free.

Given an arbitrary path $q \in H^1([T_1, T_2], \mathcal{X})$, set

$$\begin{aligned} H^1(q) &:= \{\tilde{q} \in H^1([T_1, T_2], \mathcal{X}) : \tilde{q}(T_i) = q(T_i), i = 1, 2\}; \\ \hat{H}^1(q) &:= H^1(q) \cap H^1([T_1, T_2], \hat{\mathcal{X}}). \end{aligned}$$

Then $\hat{H}^1(q)$ is an open and dense subset of $H^1(q)$, where \mathcal{A}^ε is lower semi-continuous in $H^1(q)$ and C^2 in $\hat{H}^1(q)$. Hence if q is collision-free and $d\mathcal{A}^\varepsilon(q) = 0$, then it is a critical point of \mathcal{A}^ε , and we will denote its Morse index in $\hat{H}^1(q)$ by $m_{T_1, T_2}^-(q, \mathcal{A}^\varepsilon)$. Now we introduce a notion called *weak critical points* as a generalization of the usual critical points.

Definition 1.3. We say a path $q \in H^1([T_1, T_2], \mathcal{X})$ (which may contain collision) with finite action value, $\mathcal{A}(q; T_1, T_2) < \infty$, is a **weak critical point** of \mathcal{A} , if there exists a sequence of positive numbers $\varepsilon_n \rightarrow 0$ and a sequence of $q^n \in H^1([T_1, T_2], \mathcal{X})$, such that

- (i). $\mathcal{A}^{\varepsilon_n}(q^n; T_1, T_2) < C$, $\forall n$, for some finite constant C ;
- (ii). q^n is a critical point of $\mathcal{A}^{\varepsilon_n}$, for any n ;
- (iii). $q^n \rightarrow q$ weakly in H^1 -norm and strongly in L^∞ -norm.

$c = \mathcal{A}(q; T_1, T_2)$ will be called a **weak critical value** of \mathcal{A} , and the *Morse index* of such a weak critical point q in $H^1(q)$ (with respect to \mathcal{A}) will be defined as

$$(7) \quad m_{T_1, T_2}^-(q, \mathcal{A}) = \inf \liminf_{n \rightarrow \infty} m_{T_1, T_2}^-(q^n, \mathcal{A}^{\varepsilon_n}),$$

where the infimum is taken over all sequences ε_n and q^n satisfying the above conditions.

Remark 1.1. A similar notation was introduced in [2], where it was called **generalized critical point**.

With the above definition, we have the following result, which can be seen as a partial generalization of Theorem 1.1.

Theorem 1.2. *When $d \geq 3$, given a weak critical point $q \in H^1([T_1, T_2], \mathcal{X})$ of \mathcal{A} , let $\mathcal{B}(q)$ represent the number of binary collisions occurring in $q(t)$, $t \in (T_1, T_2)$ (when there are more than one binary collision at a given moment, each of them should be counted separately), then*

$$(8) \quad (d-2)i(\alpha)\mathcal{B}(q) \leq m_{T_1, T_2}^-(q, \mathcal{A}),$$

where

$$(9) \quad i(\alpha) = \max\{k \in \mathbb{Z} : k < \frac{2}{2-\alpha}\}.$$

In particular $q(t)$, $t \in (T_1, T_2)$, is free of binary collision, i.e. $\mathcal{B}(q) = 0$, if

$$m_{T_1, T_2}^-(q, \mathcal{A}) < (d-2)i(\alpha).$$

Remark 1.2. Notice that by (9), $i(\alpha) = 1$, if $\alpha \in (0, 1]$ and $i(\alpha) \geq 2$, if $\alpha \in (1, 2)$. Moreover $i(\alpha)$ goes to infinity, as α goes to 2.

Remark 1.3. It seems the above result is the best we can get based on Tanaka's idea. In particular, we are unable to obtain any nontrivial result, when $d = 2$. For an explanation see Remark 4.1.

The idea of using Morse index to rule out collision should work even when a collision cluster has more than two masses, although the technical difficulty seems very challenging. This is because when two masses approach to a binary collision, they behaves more and more like the two body problem, where the solutions are well understood and their Morse indices are relatively easy to compute. However when the collision cluster has more than two masses, as they approach to collision,

the dynamics is much more complicate (see [20]) and the computation of Morse indices of the relevant solutions is also much more difficult. Despite of this, some progresses have been made recently in [6], [5] and [15].

Theorem 1.2 has the following obvious corollaries.

Corollary 1.1. *When $m_{T_1, T_2}^-(q, \mathcal{A}) = 0$ and $d \geq 3$, $q(t)$, $t \in (T_1, T_2)$, is free of binary collision.*

Corollary 1.2. *When $m_{T_1, T_2}^-(q, \mathcal{A}) = 1$, the following results hold.*

- (a). *If $d \geq 4$ and $\alpha \in (0, 2)$, then $q(t)$, $t \in (T_1, T_2)$ is free of binary collision.*
- (b). *If $d = 3$ and $\alpha \in (1, 2)$, then $q(t)$, $t \in (T_1, T_2)$ is free of binary collision.*
- (c). *If $d = 3$ and $\alpha \in (0, 1]$, then $q(t)$, $t \in (T_1, T_2)$ has at most one binary collision, i.e. $\mathcal{B}(q) \leq 1$.*

Notice that in Corollary 1.2, when $d = 3$ and $\alpha \in (0, 1]$, the weak critical point may very well contains a binary collision and in this case we have the following result.

Theorem 1.3. *When $d = 3$ and $\alpha = 1$, let $q \in H^1([T_1, T_2], \mathcal{X})$ be a weak critical point of \mathcal{A} with $m_{T_1, T_2}^-(q, \mathcal{A}) = 1$, if there is a binary collision between m_{i_1} and m_{i_2} at a moment $t_0 \in (T_1, T_2)$, then both limits $\lim_{t \rightarrow t_0^\pm} \frac{q_{i_2}(t) - q_{i_1}(t)}{|q_{i_2}(t) - q_{i_1}(t)|}$ exist and equal to each other.*

Remark 1.4. The above result is interesting, because it is well-known if a solution of the spatial N -body problem has a single binary collision at a moment (no other partial collision exists at the same moment), then it can be regularized by Kustaanheimo-Stiefel regularization [16]. With the result from the above theorem, under the assumption that there is no other partial collisions, one can show the generalize solution corresponding to the weak critical point is actually a classical solution in the regularized system.

Remark 1.5. Results similar to Theorem 1.3 can be obtained for potentials with $\alpha \neq 1$, see Lemma 3.2. However the problem of regularizing a binary collision is more complicate for non-Newtonian potentials, see [21].

Since the Morse index of a critical point obtained by the mountain pass theorem must be less than or equal to one (see [14]), we believe Corollary 1.1 and 1.2 and Theorem 1.3 could be useful, when mountain pass methods are used in the study the N -body problem. This shall be discussed in a forthcoming paper.

Our paper is organized as follows: in Section 2 we show a weak critical point is a generalized solution, in Section 3 the proofs of the main results will be given, and in Section 4 and 5 we give the proofs of some technical lemmas.

2. GENERALIZED SOLUTIONS

Consider the perturbed N -body problem (5), for any subset of indices $\mathbf{I} \subset \mathbf{N}$, we define the *Lagrangian* and *energy* of the \mathbf{I} -cluster as

$$\begin{aligned} L_{\mathbf{I}}^\varepsilon(q, \dot{q}) &:= K_{\mathbf{I}}(\dot{q}) + U_{\mathbf{I}}(q) + \varepsilon \mathfrak{U}_{\mathbf{I}}(q), \\ E_{\mathbf{I}}^\varepsilon(q, \dot{q}) &:= K_{\mathbf{I}}(\dot{q}) - U_{\mathbf{I}}(q) - \varepsilon \mathfrak{U}_{\mathbf{I}}(q), \end{aligned}$$

where

$$K_{\mathbf{I}}(\dot{q}) := \frac{1}{2} \sum_{i \in \mathbf{I}} m_i |\dot{q}_i|^2, \quad U_{\mathbf{I}}(q) := \sum_{\{i < j\} \subset \mathbf{I}} \frac{m_i m_j}{|q_i - q_j|^\alpha}, \quad \mathfrak{U}_{\mathbf{I}}(q) := \sum_{\{i < j\} \subset \mathbf{I}} \frac{m_i m_j}{|q_i - q_j|^2}.$$

Let $\mathbf{I}' := \mathbf{N} \setminus \mathbf{I}$ denote the complement of \mathbf{I} in \mathbf{N} , then

$$\begin{aligned} L^\varepsilon(q, \dot{q}) &= L_{\mathbf{I}}^\varepsilon(q, \dot{q}) + L_{\mathbf{I}'}^\varepsilon(q, \dot{q}) + U_{\mathbf{I}, \mathbf{I}'}(q) + \varepsilon \mathfrak{U}_{\mathbf{I}, \mathbf{I}'}(q), \\ E^\varepsilon(q, \dot{q}) &= E_{\mathbf{I}}^\varepsilon(q, \dot{q}) + E_{\mathbf{I}'}^\varepsilon(q, \dot{q}) - U_{\mathbf{I}, \mathbf{I}'}(q) - \varepsilon \mathfrak{U}_{\mathbf{I}, \mathbf{I}'}(q), \end{aligned}$$

where

$$U_{\mathbf{I}, \mathbf{I}'}(q) := \sum_{i \in \mathbf{I}, j \in \mathbf{I}'} \frac{m_i m_j}{|q_i - q_j|^\alpha}, \quad \mathfrak{U}_{\mathbf{I}, \mathbf{I}'}(q) := \sum_{i \in \mathbf{I}, j \in \mathbf{I}'} \frac{m_i m_j}{|q_i - q_j|^2}.$$

Definition 2.1. $q \in H^1([T_1, T_2], \mathcal{X})$ is a **generalized solution** of (1), if it satisfies the following conditions:

- (i). $\Delta^{-1}(q) := \{t \in [T_1, T_2] : q(t) \in \Delta\}$ has measure 0 in $[T_1, T_2]$;
- (ii). $q \in C^2$ on $[T_1, T_2] \setminus \Delta^{-1}(q)$ and satisfies (1);
- (iii). the total energy of $q(t)$, $E(t) = E(q(t), \dot{q}(t))$, is a constant, for all $t \in [T_1, T_2] \setminus \Delta^{-1}(q)$;
- (iv). for any subset $\mathbf{I} \subset \mathbf{N}$ and sub-interval $(t_1, t_2) \subset [T_1, T_2]$, if

$$(10) \quad q_i(t) \neq q_j(t), \quad \forall i \in \mathbf{I}, \forall j \in \mathbf{I}', \text{ and } \forall t \in (t_1, t_2),$$

then $E_{\mathbf{I}}(t) = E_{\mathbf{I}}(q(t), \dot{q}(t)) \in H^1((t_1, t_2), \mathbb{R})$. In particular, $E_{\mathbf{I}}(t)$ is continuous in (t_1, t_2) .

Remark 2.1. Condition (iv) in the above definition shows the energy of a \mathbf{I} -cluster is continuous, as long as the masses from the \mathbf{I} -cluster do not collide with masses outside of the cluster, even when there are collisions among the masses inside the cluster. This condition was not required by in the original definition of a generalized solution introduced by Bahri and Rabinowitz, see [4] and [1]. Our definition here is stronger and follows from [12, Definition 4.6].

Proposition 2.1. *A weak critical point $q \in H^1([T_1, T_2], \mathcal{X})$ of \mathcal{A} is a generalized solution of (1).*

Proof. Let $\varepsilon_n > 0$ and $q^n \in H^1(T_1, T_2, \mathcal{X})$ be two sequences satisfying the conditions given in Definition 1.3. Then there is a finite constant $C > 0$, such that

$$(11) \quad \frac{1}{2} \int_{t_1}^{t_2} \sum_{i \in \mathbf{N}} m_i |\dot{q}_i^n(t)|^2 dt \leq \mathcal{A}^{\varepsilon_n}(q^n; T) \leq C, \quad \forall n.$$

The fact that q satisfies the first three conditions given in Definition 2.1 is a standard result, for details see [4] or [1]. In the following, we will show q also satisfies condition (iv). Given an arbitrary $\mathbf{I} \subset \mathbf{N}$, recall that

$$E_{\mathbf{I}}^{\varepsilon_n}(t) = E_{\mathbf{I}}^{\varepsilon_n}(q^n(t), \dot{q}^n(t)) = K_{\mathbf{I}}(\dot{q}^n(t)) - U_{\mathbf{I}}(q^n) - \varepsilon_n \mathfrak{U}_{\mathbf{I}}(q^n).$$

By a direct computation,

$$(12) \quad \frac{dE_{\mathbf{I}}^{\varepsilon_n}}{dt} = \sum_{i \in \mathbf{I}} \left\langle \frac{\partial U_{\mathbf{I}, \mathbf{I}'}(q^n)}{\partial q_i^n} + \varepsilon_n \frac{\partial \mathfrak{U}_{\mathbf{I}, \mathbf{I}'}(q^n)}{\partial q_i^n}, \dot{q}_i^n \right\rangle.$$

Let's assume q satisfies (10) for the above \mathbf{I} and an arbitrary sub-interval $(t_1, t_2) \subset [T_1, T_2]$. Since $q^n(t)$ converges to $q(t)$ uniformly on $[0, T]$,

$$(13) \quad \left| \frac{\partial U_{\mathbf{I}, \mathbf{I}'}(q^n(t))}{\partial q_i^n} \right|, \left| \frac{\partial \mathfrak{U}_{\mathbf{I}, \mathbf{I}'}(q^n(t))}{\partial q_i^n} \right| \leq C_1, \quad \forall t \in [t_1, t_2], \forall i \in \mathbf{I}.$$

Here and in the rest of the proof $C_i, i \in \mathbb{Z}^+$, always represents some positive constant independent of n . With (12) and (13), the Cauchy-Schwarz inequality tells us

$$|E_{\mathbf{I}}^{\dot{\varepsilon}^n}(t)|^2 \leq C_2 \sum_{i \in \mathbf{I}} m_i |\dot{q}_i^n(t)|^2 \leq C_2 \sum_{i \in \mathbf{N}} m_i |q_i^n(t)|^2, \quad \forall t \in [t_1, t_2].$$

Then

$$(14) \quad \int_{t_1}^{t_2} |E_{\mathbf{I}}^{\dot{\varepsilon}^n}(t)|^2 dt \leq C_2 \int_{t_1}^{t_2} \sum_{i \in \mathbf{N}} m_i |q_i^n(t)|^2 dt \leq C_3.$$

Since $U_{\mathbf{I}}, \mathfrak{U}_{\mathbf{I}}$ are always positive, $E_{\mathbf{I}}^{\varepsilon^n}(t) \leq K_{\mathbf{I}}(q^n(t)), \forall t$. Then

$$(15) \quad \int_{t_1}^{t_2} E_{\mathbf{I}}^{\varepsilon^n}(t) dt \leq \frac{1}{2} \int_{t_1}^{t_2} \sum_{i \in \mathbf{N}} m_i |q_i^n(t)|^2 dt \leq C_4.$$

Meanwhile by Poincaré inequality and (14),

$$(16) \quad \int_{t_1}^{t_2} |E_{\mathbf{I}}^{\varepsilon^n}(t) - \int_{t_1}^{t_2} E_{\mathbf{I}}^{\varepsilon^n}(s) ds|^2 dt \leq C_5 \int_{t_1}^{t_2} |E_{\mathbf{I}}^{\dot{\varepsilon}^n}|^2 dt \leq C_6.$$

Then (15) implies

$$\int_{t_1}^{t_2} |E_{\mathbf{I}}^{\varepsilon^n}(t)|^2 dt \leq C_7.$$

The above inequality and (14) implies $E_{\mathbf{I}}^{\varepsilon^n}(t)$ is a bounded sequence in $H^1([t_1, t_2], \mathbb{R})$. After passing to a subsequence, it converges to a $\hat{E}_{\mathbf{I}}(t) \in H^1([t_1, t_2], \mathbb{R})$ weakly in H^1 norm and strongly in L^∞ norm.

Since $\dot{q}(t)$ and $E_{\mathbf{I}}(t) = E_{\mathbf{I}}(q(t), \dot{q}(t))$ are well defined for any $t \notin \Delta^{-1}(q)$, and

$$q^n(t) \rightarrow q(t), \quad \dot{q}^n(t) \rightarrow \dot{q}(t), \quad \text{as } n \rightarrow \infty, \quad \forall t \notin \Delta^{-1}(q),$$

we have $E_{\mathbf{I}}^{\varepsilon^n}(t) \rightarrow E_{\mathbf{I}}(t)$, for any $t \notin \Delta^{-1}(q)$. As a result, $E_{\mathbf{I}}(t) = \hat{E}_{\mathbf{I}}(t)$, for any $t \in [t_1, t_2] \setminus \Delta^{-1}(q)$. Since $\Delta^{-1}(q)$ is a set of measure zero, $E_{\mathbf{I}}(t) = \hat{E}_{\mathbf{I}}(t)$ as a H^1 -Sobolev function, and it is continuous in (t_1, t_2) . \square

Definition 2.2. Given a path $q \in H^1([T_1, T_2], \mathcal{X})$ with an \mathbf{I} -cluster collision at a moment $t_0 \in (T_1, T_2)$, we say it is **isolated**, if there is a constant $a > 0$ small enough, such that for any $i \in \mathbf{I}$,

$$q_i(t) \neq q_j(t), \quad \forall t \in [t_0 - a, t_0 + a] \setminus \{t_0\}, \quad \forall j \in \mathbf{N} \setminus \{i\}.$$

Proposition 2.2. *Given a weak critical point $q \in H^1([T_1, T_2], \mathcal{X})$, if there is a binary collision at the moment $t_0 \in (T_1, T_2)$, then it must be isolated.*

Proof. By Proposition 2.1, q is a generalized solution of (1). In particular it satisfies condition (iv) in Definition 2.1, then the desired result was already proven in [12, Corollary 5.12]. Once the reader notices that every binary collision is a so called *locally minimal collision* defined in [12, Definition 5.2]. \square

3. PROOF OF THEOREM 1.2 AND 1.3

To prove the main theorems, three technical lemmas will be needed. We present them as Lemma 3.1, 3.2 and 3.3 in this section and postpone their proofs until the next two sections.

Let $q \in H^1([T_1, T_2], \mathcal{X})$ be a weak critical point of \mathcal{A} with a binary collision at the moment $t_0 \in (T_1, T_2)$ and $q^n \in H^1([T_1, T_2], \mathcal{X})$ a sequence of critical points of $\mathcal{A}^{\varepsilon_n}$ satisfying the conditions required in Definition 1.3. Without loss of generality, we may assume such a binary collision is between m_1 and m_2 , i.e.

$$q_1(t_0) = q_2(t_0) \neq q_i(t_0), \quad \forall i \in \mathbf{N} \setminus \{1, 2\}.$$

By Proposition 2.2, such an binary collision must be isolated, so we may choose an $a > 0$ small enough, such that $[t_0 - 2a, t_0 + 2a] \subset (T_1, T_2)$ and

$$(17) \quad q_1(t) \neq q_2(t), \quad \forall t \in [t_0 - 2a, t_0 + 2a] \setminus \{t_0\};$$

$$(18) \quad q_i(t) \neq q_j(t), \quad \forall t \in [t_0 - 2a, t_0 + 2a], \quad \forall i \in \{1, 2\}, \quad \forall j \in \mathbf{N} \setminus \{1, 2\}.$$

For each n , we can always find a $t_n \in [t_0 - 2a, t_0 + 2a]$ such that

$$\delta_n := |q_1^n(t_n) - q_2^n(t_n)| = \min\{|q_1^n(t) - q_2^n(t)| : t \in [t_0 - 2a, t_0 + 2a]\}.$$

Obviously δ_n converges to 0, as n goes to infinity. After passing to a subsequence, we may assume the limit of t_n exists. Since $q_1(t_0) = q_2(t_0)$ is an isolated binary collision at the moment t_0 , t_n must converge to t_0 . Then for n large enough, $[t_n - a, t_n + a] \subset [t_0 - 2a, t_0 + 2a]$. As a result,

$$(19) \quad \delta_n = \min\{|q_1^n(t) - q_2^n(t)| : t \in [t_n - a, t_n + a]\}.$$

By Definition 1.3, $q^n(t)$ converges to $q(t)$ uniformly on $[T_1, T_2]$. According to (18), there is constant $C_1 > 0$ independent of n , such that

$$(20) \quad |q_i^n(t) - q_j^n(t)| \geq C_1, \quad \forall t \in [t_n - a, t_n + a], \quad \forall i \in \{1, 2\}, \quad \forall j \in \mathbf{N} \setminus \{1, 2\}.$$

Let

$$\tilde{U}(q^n) := U(q^n) - \frac{m_1 m_2}{|q_1^n - q_2^n|^\alpha}, \quad \tilde{\mathcal{U}}(q^n) := \mathcal{U}(q^n) - \frac{m_1 m_2}{|q_1^n - q_2^n|^2}.$$

There are constants $C_2, C_3 > 0$ independent of n , such that

$$(21) \quad \left| \frac{\partial \tilde{U}(q^n(t))}{\partial q_i^n} \right|, \left| \frac{\partial \tilde{\mathcal{U}}(q^n(t))}{\partial q_i^n} \right| \leq C_2, \quad \forall t \in [t_n - a, t_n + a], \quad \forall 1 \leq i \leq 2,$$

$$(22) \quad \left| \frac{\partial^2 \tilde{U}(q^n(t))}{\partial q_i^n \partial q_i^n} \right|, \left| \frac{\partial^2 \tilde{\mathcal{U}}(q^n(t))}{\partial q_i^n \partial q_i^n} \right| \leq C_3, \quad \forall t \in [t_n - a, t_n + a], \quad \forall 1 \leq i \leq 2.$$

We introduce a new function $\eta^n(t) = (\eta_i^n(t))_{i=1}^N$ by

$$(23) \quad \begin{cases} \eta_1^n(t) &= q_2^n(t) - q_1^n(t), \\ \eta_2^n(t) &= \frac{1}{m_1 + m_2} (m_1 q_1^n(t) + m_2 q_2^n(t)), \\ \eta_i^n(t) &= q_i^n(t), \quad \text{if } i = 3, \dots, N. \end{cases}$$

By a direct computation, $\eta^n(t)$ is a solution of

$$(24) \quad \frac{m_1 m_2}{m_1 + m_2} \ddot{\eta}_1^n = -\alpha m_1 m_2 \frac{\eta_1^n}{|\eta_1^n|^{\alpha+2}} - 2m_1 m_2 \frac{\varepsilon_n \eta_1^n}{|\eta_1^n|^4} + \frac{\partial \tilde{U}(\eta^n)}{\partial \eta_1^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{U}}(\eta^n)}{\partial \eta_1^n};$$

$$(25) \quad (m_1 + m_2) \ddot{\eta}_2^n = \frac{\partial \tilde{U}(\eta^n)}{\partial \eta_2^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{U}}(\eta^n)}{\partial \eta_2^n};$$

$$(26) \quad m_i \ddot{\eta}_i^n = \frac{\partial \tilde{U}(\eta^n)}{\partial \eta_i^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{U}}(\eta^n)}{\partial \eta_i^n}, \quad i = 3, \dots, N.$$

This is the Euler-Lagrange equation of the following Lagrangian

$$(27) \quad L(\eta^n, \dot{\eta}^n) = \frac{m_1 m_2}{2(m_1 + m_2)} |\dot{\eta}_1^n|^2 + \frac{m_1 + m_2}{2} |\dot{\eta}_2^n|^2 + \frac{1}{2} \sum_{i=3}^N m_i |\dot{\eta}_i^n|^2 \\ + \frac{m_1 m_2}{|\eta_1^n|^\alpha} + \frac{\varepsilon_n m_1 m_2}{|\eta_1^n|^2} + \tilde{U}(\eta^n) + \tilde{\mathfrak{U}}(\eta^n).$$

To study the behaviors of the solutions as they approach to the binary collision, Tanaka's *blow-up* technique will be used. The precise argument depends on the limit of $\varepsilon_n / \delta_n^{2-\alpha}$. After passing to subsequence, we may assume such a limit $\lambda = \lim_{n \rightarrow \infty} \varepsilon_n / \delta_n^{2-\alpha}$ always exists. Then two different cases need to be considered: *Case 1*, $\lambda \in [0, \infty)$; *Case 2*, $\lambda = \infty$.

For *Case 1*, we blow up $\eta^n(t)$ according to

$$(28) \quad \xi^n(s) = (\xi_i^n(s))_{i=1}^N := (\delta_n^{-1} \eta_i^n(t(s)))_{i=1}^N, \quad \text{where } t(s) = \delta_n^{1+\frac{\alpha}{2}} s + t_n.$$

By changing the time parameter from t to s , the time interval $[t_n - a, t_n + a]$ is mapped onto $[-a/\delta_n^{1+\alpha/2}, a/\delta_n^{1+\alpha/2}]$. Notice that the latter interval converges to \mathbb{R} , as n goes to infinity. Let $'$ denote derivatives with respect s , then $\xi_1(s)$ satisfies

$$(29) \quad \frac{1}{m_1 + m_2} (\xi_1^n)'' = -\alpha \frac{\xi_1^n}{|\xi_1^n|^{\alpha+2}} - 2 \frac{\varepsilon_n}{\delta_n^{2-\alpha}} \frac{\xi_1^n}{|\xi_1^n|^4} \\ + \frac{\delta_n^{1+\alpha}}{m_1 m_2} \left(\frac{\partial \tilde{U}(\eta^n)}{\partial \eta_1^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{U}}(\eta^n)}{\partial \eta_1^n} \right).$$

Lemma 3.1. *If $\lambda = \lim_{n \rightarrow \infty} \varepsilon_n / \delta_n^{2-\alpha}$ is finite, then the following results hold.*

- (a). *After passing to a subsequence, $\xi_1^n(s)$ converges to a $\xi_1(s)$ in $C^2([-\ell, \ell], \mathbb{R}^d)$, for any $\ell > 0$, where $\xi_1(s)$ is a solution of*

$$(30) \quad \frac{1}{m_1 + m_2} \hat{\xi}_1'' = -\alpha \frac{\hat{\xi}_1}{|\hat{\xi}_1|^{\alpha+2}} - 2 \frac{\lambda \hat{\xi}_1}{|\hat{\xi}_1|^4};$$

$$(31) \quad \frac{1}{2(m_1 + m_2)} |\hat{\xi}_1'|^2 - \frac{1}{|\hat{\xi}_1|^\alpha} - \frac{\lambda}{|\hat{\xi}_1|^2} = 0;$$

$$(32) \quad \langle \hat{\xi}_1(0), \hat{\xi}_1'(0) \rangle = 0.$$

- (b). $\xi_1(t) \in W(\xi_1), \forall t \in \mathbb{R}$, where $W(\xi_1) = \text{span}\{\xi_1(0), \xi_1'(0)\}$ is a 2-dim subspace of \mathbb{R}^d . Moreover the following limits exist

$$\lim_{s \rightarrow \pm\infty} |\xi_1(s)| = +\infty, \quad \lim_{t \rightarrow \pm\infty} \frac{\xi_1(s)}{|\xi_1(s)|} = u^\pm,$$

and

$$\angle(u^-, u^+) = 2\pi \frac{\sqrt{1+\lambda}}{2-\alpha}.$$

For any two unit vectors $u, v \in \mathbb{R}^d$, $\angle(u, v)$ represents the angle between them.

- (c). Let $W^\perp(\xi_1)$ be the orthogonal complement of $W(\xi_1)$ in \mathbb{R}^d and $H(\xi_1)$ the largest subspace of $H_0^1(\mathbb{R}, W^\perp(\xi_1))$, such that

$$d^2\mathcal{I}(\xi_1)(\phi, \phi) < 0, \quad \forall \phi \in H(\xi_1),$$

where \mathcal{I} is the Lagrange action functional corresponding to equation (30):

$$(33) \quad \mathcal{I}(\hat{\xi}_1) = \int \frac{1}{2(m_1 + m_2)} |\hat{\xi}_1'|^2 + \frac{1}{|\hat{\xi}_1|^\alpha} + \frac{\lambda}{|\hat{\xi}_1|^2} dt,$$

then $\dim(H(\xi_1)) \geq (d-2)i(\alpha, \lambda)$, where

$$(34) \quad i(\alpha, \lambda) := \max\{k \in \mathbb{Z} : k < \frac{2\sqrt{1+\lambda}}{2-\alpha}\}.$$

Lemma 3.2. If $\lambda = \lim_{n \rightarrow \infty} \varepsilon_n / \delta_n^{2-\alpha}$ is finite, then

$$\lim_{t \rightarrow t_0^\pm} \frac{q_2(t) - q_1(t)}{|q_2(t) - q_1(t)|} = u^\pm,$$

where u^\pm are the two unit vectors given in property (b), Lemma 3.1.

For Case 2, we define a blow-up of η^n according to

$$(35) \quad \zeta^n(s) = (\zeta_i^n(s))_{i=1}^N = (\delta_n^{-1} \eta_i^n(t(s)))_{i=1}^N, \quad \text{where } t(s) = \varepsilon_n^{-\frac{1}{2}} \delta_n^2 s + t_n.$$

Like the previous case, after changing the time parameter from t to s , the time interval $[t_n - a, t_n + a]$ is mapped onto $[-a\varepsilon_n^{\frac{1}{2}}/\delta_n^2, a\varepsilon_n^{\frac{1}{2}}/\delta_n^2]$, which converges to \mathbb{R} , as n goes to infinity. Again if we let $'$ represents derivatives with respect to s , then $\zeta_1^n(s)$ satisfies

$$(36) \quad \begin{aligned} \frac{1}{m_1 + m_2} (\zeta_1^n)'' &= -2 \frac{\zeta_1^n}{|\zeta_1^n|^4} - \alpha \frac{\delta_n^{2-\alpha}}{\varepsilon_n} \frac{\zeta_1^n}{|\zeta_1^n|^{\alpha+1}} \\ &+ \frac{\varepsilon_n}{\delta_n^3 m_1 m_2} \left(\frac{\partial \tilde{U}(\eta^n)}{\partial \eta_1^n} + \varepsilon_n \frac{\partial \tilde{\Omega}(\eta^n)}{\partial \eta_1^n} \right). \end{aligned}$$

Lemma 3.3. If $\lambda = \lim_{n \rightarrow \infty} \varepsilon_n / \delta_n^{2-\alpha} = \infty$, then the following results hold.

- (a). After passing to a subsequence, $\zeta_1^n(s)$ converges to a $\zeta_1(s)$ in $C^2([-\ell, \ell], \mathbb{R}^d)$, for any $\ell > 0$, where $\zeta_1(s)$ is a solution of

$$(37) \quad \frac{1}{m_1 + m_2} \hat{\zeta}_1'' = -2 \frac{\hat{\zeta}_1}{|\hat{\zeta}_1|^4};$$

$$(38) \quad \frac{1}{2(m_1 + m_2)} |\hat{\zeta}_1'|^2 - \frac{1}{|\hat{\zeta}_1|^2} = 0;$$

$$(39) \quad \langle \hat{\zeta}_1(0), \hat{\zeta}_1'(0) \rangle = 0.$$

- (b). $\zeta_1(t) \in W(\zeta_1), \forall t \in \mathbb{R}$, where $W(\zeta_1) = \text{span}\{\zeta_1(0), \zeta_1'(0)\}$ is a 2-dim subspace of \mathbb{R}^d , and $\lim_{s \rightarrow \pm\infty} |\zeta_1(s)| = +\infty$.

(c). Let $W^\perp(\zeta_1)$ be the orthogonal complement of $W(\zeta_1)$ in \mathbb{R}^d and $H(\zeta_1)$ the largest subspace of $H_0^1(\mathbb{R}, W^\perp(\zeta_1))$, such that

$$d^2\mathcal{J}(\zeta_1)(\phi, \phi) < 0, \quad \forall \phi \in H(\zeta_1),$$

where \mathcal{J} is the Lagrange action functional corresponding to equation (37),

$$(40) \quad \mathcal{J}(\hat{\zeta}) := \int \frac{1}{2(m_1 + m_2)} |\hat{\zeta}'|^2 + \frac{1}{|\hat{\zeta}|^2} ds,$$

then $\dim(H(\hat{\zeta})) = +\infty$.

Proposition 3.1. Under the above notation,

(a). if $\lambda = \lim_{n \rightarrow \infty} \varepsilon_n / \delta_n^{2-\alpha}$ is finite, then

$$(41) \quad \liminf_{n \rightarrow \infty} m_{t_n - a, t_n + a}^-(q^n, \mathcal{A}^{\varepsilon_n}) \geq (d-2)i(\alpha, \lambda);$$

(b). if $\lambda = \lim_{n \rightarrow \infty} \varepsilon_n / \delta_n^{2-\alpha} = +\infty$, then

$$(42) \quad \liminf_{n \rightarrow \infty} m_{t_n - a, t_n + a}^-(q^n, \mathcal{A}^{\varepsilon_n}) = +\infty.$$

Proof. (a). Given an arbitrary $f = (f_i)_{i=1}^N \in H_0^1([t_n - a, t_n + a], \mathbb{R}^{dN})$ satisfying

$$(43) \quad f_i(t) \equiv 0, \quad \forall t, \quad \text{and} \quad \forall i \neq 1,$$

by (23), for any $c > 0$, the center of mass corresponding to the path $\eta^n(t) + cf(t)$ will always be at the origin. In the following, we shall compute the second variation of $\mathcal{A}^{\varepsilon_n}$ at η^n among all $f \in H_0^1([t_n - a, t_n + a], \mathbb{R}^{dN})$ satisfying (43). By a direct computation,

$$(44) \quad \begin{aligned} d^2\mathcal{A}^{\varepsilon_n}(\eta^n)[f, f] &= \int_{t_n - a}^{t_n + a} \frac{m_1 m_2}{m_1 + m_2} |f_1|^2 - \alpha m_1 m_2 \frac{|f_1|^2}{|\eta_1^n|^{\alpha+2}} - 2m_1 m_2 \frac{\varepsilon_n |f_1|^2}{|\eta_1^n|^4} \\ &+ \alpha(\alpha + 2)m_1 m_2 \frac{\langle \eta_1^n, f_1 \rangle}{|\eta_1^n|^{\alpha+4}} + 8m_1 m_2 \varepsilon_n \frac{\langle \eta_1^n, f_1 \rangle}{|\eta_1^n|^6} \\ &+ \left\langle \frac{\partial^2 \tilde{U}(\eta^n)}{\partial \eta_1^n \partial \eta_1^n} f_1, f_1 \right\rangle + \varepsilon_n \left\langle \frac{\partial^2 \tilde{\mathcal{U}}(\eta^n)}{\partial \eta_1^n \partial \eta_1^n} f_1, f_1 \right\rangle dt. \end{aligned}$$

Define the linear operators $T_n : H_0^1(\mathbb{R}, W^\perp(\xi_1)) \rightarrow H_0^1([t_n - a, t_n + a], \mathbb{R}^{dN})$ by $f^n(t) = (f_i^n(t))_{i=1}^N = (T_n \phi)(s(t))$, where $s(t) = \delta_n^{-\frac{2+\alpha}{2}}(t - t_n)$ and

$$f_1^n(t) = \delta_n \phi(s(t)) \quad \text{and} \quad f_i^n(t) \equiv 0, \quad \forall i \neq 1.$$

Then

$$(45) \quad \begin{aligned} \delta_n^{-\frac{2-\alpha}{2}} d^2\mathcal{A}^{\varepsilon_n}(\eta^n)[f^n, f^n] &= \int_{\text{supp}(\phi)} \frac{m_1 m_2}{m_1 + m_2} |\phi'|^2 - \alpha m_1 m_2 \frac{|\phi|^2}{|\xi_1^n|^{\alpha+2}} \\ &- 2m_1 m_2 \frac{\varepsilon_n}{\delta_n^{2-\alpha}} \frac{|\phi|^2}{|\xi_1^n|^4} + \alpha(\alpha + 2)m_1 m_2 \frac{\langle \xi_1^n, \phi \rangle^2}{|\xi_1^n|^{\alpha+4}} + 8m_1 m_2 \frac{\varepsilon_n}{\delta_n^{2-\alpha}} \frac{\langle \xi_1^n, \phi \rangle^2}{|\xi_1^n|^6} \\ &+ \delta_n^{2+\alpha} \left(\left\langle \frac{\partial^2 \tilde{U}(\eta^n)}{\partial \eta_1^n \partial \eta_1^n} \phi, \phi \right\rangle^2 + \varepsilon_n \left\langle \frac{\partial^2 \tilde{\mathcal{U}}(\eta^n)}{\partial \eta_1^n \partial \eta_1^n} \phi, \phi \right\rangle^2 \right) ds \end{aligned}$$

By (22), there is a constant $C > 0$ independent of n , such that

$$(46) \quad \left| \frac{\partial^2 \tilde{U}(\eta^n(t(s)))}{\partial \eta_1^n \partial \eta_1^n} \right|, \left| \frac{\partial^2 \tilde{\mathcal{U}}(\eta^n(t(s)))}{\partial \eta_1^n \partial \eta_1^n} \right| \leq C, \quad \forall s \in \left[-\frac{a}{\delta_n^{1+\alpha/2}}, \frac{a}{\delta_n^{1+\alpha/2}}\right], \quad \forall 1 \leq i \leq 2.$$

As ξ_1^n converges to ξ_1 in $C^2([\ell, \ell], \mathbb{R}^d)$, for any $\ell > 0$, the right hand side of (45) converges to

$$m_1 m_2 \left(\int_{\text{supp}(\phi)} \frac{1}{m_1 + m_2} |\phi'|^2 - \alpha \frac{|\phi|^2}{|\xi_1|^{\alpha+2}} - 2\lambda \frac{|\phi|^2}{|\xi_1|^4} dt \right) = m_1 m_2 (d^2 \mathcal{I}(\xi_1)[\phi, \phi]).$$

Then Lemma 3.1 implies,

$$\delta_n^{-\frac{2-\alpha}{2}} d^2 \mathcal{A}^{\varepsilon_n}(\eta^n)[f^n, f^n] < 0, \quad \text{for } n \text{ large enough, if } \phi \in H(\xi_1).$$

As $\dim(H(\xi_1)) \geq (d-2)i(\alpha, \lambda)$, it immediately implies (41).

(b). If λ is infinity, with Lemma 3.3, the desired property following from a similar argument as above. We will not repeat it again. \square

With the above results, we can now prove Theorem 1.2 and 1.3.

Proof of Theorem 1.2. By Proposition 2.2, if $q(t)$ has a binary collision, then it is isolated. Then $\mathcal{B}(q)$ must be finite.

Assume q has a binary collision at the moment $t_0 \in (T_1, T_2)$, let the corresponding sequences ε_n , $q^n(t)$ and δ_n be defined as before. If $\lim_{n \rightarrow \infty} \varepsilon_n / \delta_n^{2-\alpha} = +\infty$, then property (b) in Proposition 3.1 implies $m_{T_1, T_2}^-(q, \mathcal{A}) = +\infty$, which obviously implies (8).

If the corresponding limit of $\varepsilon_n / \delta_n^{2-\alpha}$ is finite for each binary collision, then (8) follows from the sub-additivity of the Morse index and property (a) in 3.1. Once we notice $i(\alpha, \lambda) \geq i(\alpha)$, for any $\lambda \geq 0$. \square

Proof of Theorem 1.3. Without loss of generality let's assume the binary collision is between m_1 and m_2 with the sequences ε_n , $q^n(t)$ and δ_n defined as before, and $\lambda = \lim_{n \rightarrow +\infty} \varepsilon_n / \delta_n^{2-\alpha}$. Obviously λ must be finite, as otherwise by Proposition 3.1, $m_{T_1, T_2}^-(q, \mathcal{A}) = +\infty$, which is absurd.

Since λ is finite, $\alpha = 1$ and $d = 3$, by Proposition 3.1,

$$1 = m_{T_1, T_2}^-(q, \mathcal{A}) \geq i(1, \lambda) = \max\{k \in \mathbb{Z} : k < 2\sqrt{1 + \lambda}\}.$$

This implies $\lambda = 0$. Then by Lemma 3.2, both limits $\lim_{t \rightarrow t_0^\pm} \frac{q_2(t) - q_1(t)}{|q_2(t) - q_1(t)|}$ exist and the angle between them is 2π , as $\alpha = 1$ and $\lambda = 0$. \square

4. PROOF OF LEMMA 3.1 AND 3.3

Proof of Lemma 3.1. (a). Recall that (19) and (23) imply,

$$|\eta_1^n(t_n)| = \delta_n = \min\{|\eta_1^n(t)| : t \in [t_n - a, t_n + a]\}.$$

Then by the definition of $\xi_1^n(s)$,

$$|\xi_1^n(s)| \geq |\xi_1^n(0)| = 1, \quad \forall s \in [-a/\delta_n^{1+\frac{\alpha}{2}}, a/\delta_n^{1+\frac{\alpha}{2}}].$$

As a result,

$$\langle \xi_1^n(0), (\xi_1^n)'(0) \rangle = 0.$$

Let $\xi_1(s)$ be a solution of (30), with initial condition

$$\xi_1(0) = \lim_{n \rightarrow \infty} \xi_1^n(0), \quad \xi_1'(0) = \lim_{n \rightarrow \infty} (\xi_1^n)'(0).$$

Upon passing to a subsequence, we may assume the above limits always exist. Then $\langle \xi_1(0), \xi_1'(0) \rangle = \lim_{n \rightarrow \infty} \langle \xi_1^n(0), (\xi_1^n)'(0) \rangle = 0$.

Meanwhile by (21) and (23), there is a constant $C_1 > 0$ independent of n with

$$\left| \frac{\partial \tilde{U}(\eta^n(t))}{\partial \eta_1^n} \right|, \left| \frac{\partial \tilde{\mathcal{U}}(\eta^n(t))}{\partial \eta_1^n} \right| \leq C_1, \quad \forall t \in [t_n - a, t_n + a].$$

This means the second line in equation (29) converges to zero, which gives us equation (30), as n goes to infinity. Then by the continuous dependence of solutions on initial conditions and coefficients of differential equations, we have $\xi_1^n(s)$ converges to $\xi_1(s)$ in $C^2([-\ell, \ell], \mathbb{R}^d)$, for any $\ell > 0$.

To show that $\xi_1(s)$ satisfies (31), consider the energy $E_{\mathbf{I}}^{\varepsilon_n}(t) = E_{\mathbf{I}}^{\varepsilon_n}(q^n(t), \dot{q}^n(t))$ and $E_{\mathbf{I}}(t) = E_{\mathbf{I}}(q(t), \dot{q}(t))$ of the \mathbf{I} -cluster with $\mathbf{I} = \{1, 2\}$ corresponding to q^n and q . By the proof of Proposition 2.1, $E_{\mathbf{I}}(t)$ is continuous on $[t_0 - a, t_0 + a]$ and $E_{\mathbf{I}}^{\varepsilon_n}(t)$ converges to it under the L^∞ norm, after passing to a subsequence. Therefore for n large enough, there is a constant $C_2 > 0$ independent of n , such that

$$(47) \quad |E_{\mathbf{I}}^{\varepsilon_n}(t)| \leq C_2, \quad \forall t \in [t_n - a, t_n + a].$$

Meanwhile with (23) and (28), a direct computation shows

$$(48) \quad \frac{1}{2(m_1 + m_2)} |(\xi_1^n)'|^2 - \frac{1}{|\xi_1^n|^\alpha} - \frac{\varepsilon_n}{\delta_n^{2-\alpha}} \frac{1}{|\xi_1^n|^2} = \frac{\delta_n^{2-\alpha}}{m_1 m_2} (E_{\mathbf{I}}^{\varepsilon_n} - \frac{1}{2} |\eta_2^n|^2)$$

To prove that $\xi_1(s)$ satisfies (31), it is enough to show the right hand side of the above equation converges to zero, as n goes to infinity.

To see this, notice that $\eta_2^n(t)$ converges to $\frac{m_1 q_1(t) + m_2 q_2(t)}{m_1 + m_2}$, which is the center of mass of m_1 and m_2 . Although they collide at the moment t_0 , the path of their center of mass, is actually C^2 on $[t_0 - a, t_0 + a]$ (see [12, Remark 4.10]). Hence the convergence of $\eta_2^n(t)$ holds at least under the C^1 norm, and as a result, there is a constant $C_3 > 0$ independent of n , such that

$$(49) \quad |\eta_2^n(t)| \leq C_3, \quad \forall t \in [t_n - a, t_n + a].$$

Now our claim follows directly from (47) and (49).

(b). Notice that (30) describes the motion of a point mass under the attraction of a isotropic central force. As a result, $\langle \xi_1(0), \xi_1'(0) \rangle = 0$ implies $\xi_1(t) \in W(\xi_1) = \text{span}\{\xi_1(0), \xi_1'(0)\}$, for all $t \in \mathbb{R}$. By property (a), $\xi_1(s)$ is a collision-free zero energy solution of (30). Then the rest of the property is well known and a detailed proof can be found in [23, Section 4].

(c). Let $\{\mathbf{e}_i : i = 1, \dots, d\}$ be an orthogonal basis of \mathbb{R}^d , such that $W(\xi_1) = \text{span}\{\mathbf{e}_1, \mathbf{e}_2\}$. Then for any $\varphi \in H_0^1(\mathbb{R}, \mathbb{R})$ and $\mathbf{e}_i, i = 3, \dots, d$, a simple computation shows

$$(50) \quad d^2 \mathcal{I}(\xi_1)[\varphi \mathbf{e}_i, \varphi \mathbf{e}_i] = \int_{-\infty}^{\infty} \frac{|\varphi'|^2}{2(m_1 + m_2)} - \alpha \frac{|\varphi|^2}{|\xi_1|^{\alpha+2}} - 2\lambda \frac{|\varphi|^2}{|\xi_1|^4} dt$$

Let $\Lambda(\xi_1)$ be the largest subspace of $H_0^1(\mathbb{R}, \mathbb{R})$, such that the value in (50) is negative for any $\varphi \in \Lambda(\xi_1)$. Using Sturm Comparison Theorem, in [23, Section 4], [24, Section 4] and [25, Proposition 1.1], Tanaka showed the dimension of $\Lambda(\xi_1)$ is related to the winding number of $\xi_1(s)$, $s \in \mathbb{R}$, in the plane $W(\xi_1)$ with respect to the origin. More precisely

$$(51) \quad \dim(\Lambda(\xi_1)) \geq i(\alpha, \lambda),$$

and this proves property (c). \square

Remark 4.1. Although Tanaka only state (51), a slight modification of his proof should show this is in fact an equality. As a result, if one wants to get a better estimate of the Morse index near a binary collision, one has to compute the Morse index of ξ_1 inside $W(\xi_1)$. However a result in [15, Corollary 5.1] by Hu and the author shows this is actually zero. Because of this we believe with Tanaka's approach, one can not get any nontrivial result for the planar N -body problem.

Proof of Lemma 3.3. With the results given by Tanaka in [24, Section 4], the lemma can be proven following the same argument given in Lemma 3.1. The only difference is the *blow up* should follow (35), instead of (28), and correspondingly (48) needs to be replaced by

$$(52) \quad \frac{1}{2(m_1 + m_2)} |(\zeta_1^n)'|^2 - \frac{1}{|\zeta_1^n|^2} - \frac{\delta_n^{2-\alpha}}{\varepsilon_n} \frac{1}{|\zeta_1^n|^\alpha} = \frac{\delta_n^2}{\varepsilon_n m_1 m_2} (E_{\mathbf{I}}^{\varepsilon_n} - \frac{1}{2} |\dot{\eta}_2^n|^2).$$

Nevertheless the right hand side of the equation still goes to zero, because $\lambda = +\infty$ implies δ_n^2/ε_n converge to zero, as n goes to infinity. \square

5. PROOF OF LEMMA 3.2

Our proof follows the approach given by Tanaka in [25]. First we establish a lemma that corresponds to Lemma 1.3 in [25].

Lemma 5.1. (a). *There is a constant $C_1 > 0$ independent of n , such that*

$$(53) \quad |\dot{\eta}_1^n(t)|^2 \leq \frac{2(m_1 + m_2)}{|\eta_1^n(t)|^\alpha} + \frac{2\varepsilon_n(m_1 + m_2)}{|\eta_1^n(t)|^2} + C_1, \quad \forall t \in [t_n - a, t_n + a].$$

(b). *There is a constant $\ell_0 > 0$, such that for n large enough, if $t \in [t_n - a, t_n + a]$ and $\eta_1^n(t) \in B_{\ell_0}$, then $\frac{d^2}{dt^2} |\eta_1^n(t)|^2 > 0$, where $B_\ell = \{x \in \mathbb{R}^d : |x| \leq \ell\}$ for any $\ell > 0$.*

Proof. (a). Let $\mathbf{I} = \{1, 2\}$. Recall that $E_{\mathbf{I}}^{\varepsilon_n}(t) = E_{\mathbf{I}}^{\varepsilon_n}(q^n(t), \dot{q}^n(t))$ is the energy of the \mathbf{I} -cluster. By (23),

$$(54) \quad E_{\mathbf{I}}^{\varepsilon_n}(t) = \frac{m_1 m_2}{2(m_1 + m_2)} |\dot{\eta}_1^n|^2 + \frac{m_1 + m_2}{2} |\dot{\eta}_2^n|^2 - \frac{m_1 m_2}{|\eta_1^n|^\alpha} - \varepsilon_n \frac{m_1 m_2}{|\eta_1^n|^2}.$$

As a result,

$$|\dot{\eta}_1^n(t)|^2 \leq \frac{2(m_1 + m_2)}{|\eta_1^n(t)|^\alpha} + 2\varepsilon_n \frac{m_1 + m_2}{|\eta_1^n(t)|^2} - \frac{(m_1 + m_2)^2}{m_1 m_2} |\dot{\eta}_2^n(t)|^2 + \frac{2(m_1 + m_2)}{m_1 m_2} E_{\mathbf{I}}^{\varepsilon_n}(t).$$

Then property (a) following from (47) and (49).

(b). By a direct computation,

$$(55) \quad \begin{aligned} \frac{1}{2} \frac{d^2}{dt^2} |\eta_1^n|^2 &= |\dot{\eta}_1^n|^2 + \langle \eta_1^n, \ddot{\eta}_1^n \rangle = |\dot{\eta}_1^n|^2 - \alpha \frac{m_1 + m_2}{|\eta_1^n|^\alpha} - 2\varepsilon_n \frac{m_1 + m_2}{|\eta_1^n|^2} \\ &\quad + \frac{m_1 + m_2}{m_1 m_2} \langle \eta_1^n, \frac{\partial \tilde{U}}{\partial \eta_1^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{A}}}{\partial \eta_1^n} \rangle. \end{aligned}$$

Use (47) and (54), we can find a positive constant C_2 , such that

$$\frac{1}{2} \frac{d^2}{dt^2} |\eta_1^n|^2 \geq (2 - \alpha) \frac{m_1 + m_2}{|\eta_1^n|^\alpha} - \frac{m_1 + m_2}{2} |\dot{\eta}_2^n|^2 + \frac{m_1 + m_2}{m_1 m_2} \langle \eta_1^n, \frac{\partial \tilde{U}}{\partial \eta_1^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{A}}}{\partial \eta_1^n} \rangle - C_2,$$

which clearly implies property (b). \square

Again following [25], we introduce the following functions

$$e_n(t) = \sqrt{|\eta_1^n(t)|^2 |\dot{\eta}_1^n(t)|^2 - \langle \dot{\eta}_1^n(t), \eta_1^n(t) \rangle^2}, \quad \omega(t) = \frac{e_n(t)}{|\eta_1^n(t)| |\dot{\eta}_1^n(t)|}.$$

Notice that $\omega_n(t)$ is well defined, when $\eta_1^n(t) \neq 0$ and $\dot{\eta}_1^n(t) \neq 0$. In particular, $\omega_n(t) = \sin(\angle(\eta_1^n(t)/|\eta_1^n(t)|, \dot{\eta}_1^n(t)/|\dot{\eta}_1^n(t)|))$ and $|\omega_n(t)| \leq 1$. By (24),

$$\frac{de_n}{dt} = \frac{m_1 + m_2}{m_1 m_2} \left\langle \frac{|\eta_1^n|^2 |\dot{\eta}_1^n|^2 - \langle \dot{\eta}_1^n, \eta_1^n \rangle \eta_1^n}{\sqrt{|\eta_1^n|^2 |\dot{\eta}_1^n|^2 - \langle \dot{\eta}_1^n, \eta_1^n \rangle^2}}, \frac{\partial \tilde{U}(\eta^n)}{\partial \eta_1^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{U}}(\eta^n)}{\partial \eta_1^n} \right\rangle.$$

According to (21), there are positive constants C_3, C_4 independent of n , such that (56)

$$|\dot{e}_n(t)| \leq C_3 |\eta_1^n(t)| \cdot \left| \frac{\partial \tilde{U}(\eta_1^n(t))}{\partial \eta_1^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{U}}(\eta_1^n(t))}{\partial \eta_1^n} \right| \leq C_4 |\eta_1^n(t)|, \quad \forall t \in [t_n - a, t_n + a].$$

Again by (24), a direct computation shows

$$\begin{aligned} \frac{d\omega_n}{dt} &= \frac{\dot{e}_n}{|\eta_1^n| |\dot{\eta}_1^n|} - \frac{\omega_n}{|\eta_1^n|^2 |\dot{\eta}_1^n|^2} \left(|\dot{\eta}_1^n|^2 \langle \eta_1^n, \dot{\eta}_1^n \rangle + |\eta_1^n|^2 \langle \dot{\eta}_1^n, \ddot{\eta}_1^n \rangle \right) \\ (57) \quad &= \frac{\dot{e}_n}{|\eta_1^n| |\dot{\eta}_1^n|} - \frac{m_1 + m_2}{m_1 m_2} \frac{\omega_n}{|\eta_1^n|^2} \left\langle \dot{\eta}_1^n, \frac{\partial \tilde{U}}{\partial \eta_1^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{U}}}{\partial \eta_1^n} \right\rangle \\ &\quad - \omega_n \frac{\langle \eta_1^n, \dot{\eta}_1^n \rangle}{|\eta_1^n|^2 |\dot{\eta}_1^n|^2} \left(|\dot{\eta}_1^n|^2 - \alpha \frac{m_1 + m_2}{|\eta_1^n|^\alpha} - 2\varepsilon_n \frac{m_1 + m_2}{|\eta_1^n|^2} \right) \end{aligned}$$

Then (53) and (56) implies,

$$\begin{aligned} (58) \quad \dot{\omega}_n(t) &\leq \frac{C_5}{|\eta_1^n|} \left| \frac{\partial \tilde{U}}{\partial \eta_1^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{U}}}{\partial \eta_1^n} \right| - \omega_n \frac{\langle \eta_1^n, \dot{\eta}_1^n \rangle}{|\eta_1^n|^2 |\dot{\eta}_1^n|^2} \left((2 - \alpha) \frac{m_1 + m_2}{|\eta_1^n|^\alpha} - C_6 \right) \\ &\leq \left(C_5 |\eta_1^n| \left| \frac{\partial \tilde{U}}{\partial \eta_1^n} + \varepsilon_n \frac{\partial \tilde{\mathfrak{U}}}{\partial \eta_1^n} \right| - \omega_n \sqrt{1 - \omega_n^2} \left((2 - \alpha) \frac{m_1 + m_2}{|\eta_1^n|^\alpha} - C_6 \right) \right) \frac{1}{|\eta_1^n| |\dot{\eta}_1^n|}, \end{aligned}$$

where C_5 is positive constant and C_6 is a constant, whose sign depends on the sign of $\langle \eta_1^n, \dot{\eta}_1^n \rangle$, independent of $t \in [t_n - a, t_n + a]$ and n .

With Lemma 5.1 and (58) (this corresponds to (1.8) in [25]), the next result can be proven following the argument given in Proposition 1.4 and 1.5 in [25] line by line, and we will not repeat it here.

Lemma 5.2. *Let ℓ_0 be the constant given in Lemma 5.1, for any $b > 0$ small enough, there exist constants $0 < \ell_2 < \ell_0 < \ell_1$, such that when n is large enough, for any*

$$t_n < t < t^* < t_n + a, \quad \text{or} \quad t_n - a < t < t^* < t_n,$$

if $\eta_1^n(t)$ and $\eta_1^n(t^) \in B_{\ell_2} \setminus B_{\delta_n \ell_1}$, then $\left| \eta_1^n(t)/|\eta_1^n(t)| - \eta_1^n(t^*)/|\eta_1^n(t^*)| \right| < b$.*

Now we give a proof of Lemma 3.2.

Proof of Lemma 3.2. By property (c) in Lemma 3.1, $\lim_{s \rightarrow +\infty} \xi_1(s)/|\xi_1(s)| = u^+$. Fix an arbitrary small $b > 0$, let ℓ_1, ℓ_2 be the constants given in Lemma 5.2. We

can choose an $s^* > 0$ large enough, such that for n large enough,

$$\ell_1 < |\xi_1(s^*)| < \delta_n^{-1} \ell_2, \quad \text{and} \quad \left| \frac{\xi_1(s^*)}{|\xi_1(s^*)|} - u^+ \right| < b.$$

By Lemma 3.1, the same inequalities hold for $\xi_1^n(s^*)$, when n is large enough.

Let $t^* = t_n + \delta_n^{1+\frac{\alpha}{2}} s^*$, by equation (28),

$$\eta_1^n(t^*) \in B_{\ell_2} \setminus B_{\delta_n \ell_1}, \quad \text{and} \quad |\eta_1^n(t^*)/|\eta_1^n(t^*)| - u^+| < b.$$

Since $\ell_2 < \ell_0$, for any $t \in (t^*, t_n + a)$, we claim if $\eta_1^n(t) \in B_{\ell_2}$, then $|\eta_1^n(t)| > \delta_n \ell_1$. Indeed this follows from the fact $d|\eta_1^n(t)|/dt = 0$ and property (b) in Lemma 5.1.

As a result, for any $t \in (t^*, t_n + a)$, $\eta_1^n(t) \in B_{\ell_2}$ implies $\eta_1^n(t) \in B_{\ell_2} \setminus B_{\delta_n \ell_1}$. Then by Lemma 5.2,

$$\left| \frac{\eta_1^n(t)}{|\eta_1^n(t)|} - u^+ \right| \leq \left| \frac{\eta_1^n(t)}{|\eta_1^n(t)|} - \frac{\eta_1^n(t^*)}{|\eta_1^n(t^*)|} \right| + \left| \frac{\eta_1^n(t^*)}{|\eta_1^n(t^*)|} - u^+ \right| \leq 2b.$$

This means

$$(\{\eta_1^n(t) : t \in (t_n + \delta_n^{1+\frac{\alpha}{2}} s^*, t_n + a)\} \cap B_{\ell_2}) \subset (\{x \in B_{\ell_2} : |x/x| - u^+| < 2b\} \cup B_{\delta_n \ell_1}).$$

Recall that when n goes to infinity, t_n converges to t_0 , δ_n converges to zero, and $\eta_1^n(t)$ converges uniformly to $q_2(t) - q_1(t)$. Then

$$(\{q_2(t) - q_1(t) : t \in (t_0, t_0 + a)\} \cap B_{\ell_2}) \subset \{x \in B_{\ell_2} : |x/x| - u^+| < 2b\}.$$

Since the above result hold for any $b > 0$ small enough, we get

$$\lim_{t \rightarrow +\infty} \frac{q_2(t) - q_1(t)}{|q_2(t) - q_1(t)|} = u^+.$$

A similar argument shows

$$\lim_{t \rightarrow -\infty} \frac{q_2(t) - q_1(t)}{|q_2(t) - q_1(t)|} = u^-.$$

□

Acknowledgements. The author thanks Alain Chenciner for a careful reading of an early draft of the paper and Richard Montgomery for pointing out a mistake. Valuable discussions with Vivina Barutello, Jacques Féjóz and Xijun Hu are also appreciated. The main part of the work was done when the author was a postdoc and a visitor at Ceremade in University of Paris-Dauphine, IMCCE in the Paris Observatory and School of Mathematics in Shandong University. He thanks the hospitality of these institutes and the financial support of FSMP and NSFC(No.11425105).

REFERENCES

- [1] A. Ambrosetti and V. Coti Zelati. *Periodic solutions of singular Lagrangian systems*, volume 10 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Boston, Inc., Boston, MA, 1993.
- [2] G. Arioli, V. Barutello, and S. Terracini. A new branch of Mountain Pass solutions for the choreographical 3-body problem. *Comm. Math. Phys.*, 268(2):439–463, 2006.
- [3] A. Bahri and P. H. Rabinowitz. A minimax method for a class of Hamiltonian systems with singular potentials. *J. Funct. Anal.*, 82(2):412–428, 1989.
- [4] A. Bahri and P. H. Rabinowitz. Periodic solutions of Hamiltonian systems of 3-body type. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 8(6):561–649, 1991.
- [5] V. Barutello, X. Hu, A. Portaluri, and S. Terracini. An index theory for asymptotic motions under singular potentials. Preprint, 2017, arxiv 1705.01291.

- [6] V. Barutello and S. Secchi. Morse index properties of colliding solutions to the N -body problem. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 25(3):539–565, 2008.
- [7] K.-C. Chen. Existence and minimizing properties of retrograde orbits to the three-body problem with various choices of masses. *Ann. of Math. (2)*, 167(2):325–348, 2008.
- [8] A. Chenciner. Action minimizing solutions of the Newtonian n -body problem: from homology to symmetry. In *Proceedings of the International Congress of Mathematicians, Vol. III (Beijing, 2002)*, pages 279–294. Higher Ed. Press, Beijing, 2002.
- [9] A. Chenciner, J. Gerver, R. Montgomery, and C. Simó. Simple choreographic motions of N bodies: a preliminary study. In *Geometry, mechanics, and dynamics*, pages 287–308. Springer, New York, 2002.
- [10] A. Chenciner and R. Montgomery. A remarkable periodic solution of the three-body problem in the case of equal masses. *Ann. of Math. (2)*, 152(3):881–901, 2000.
- [11] A. Chenciner and A. Venturelli. Minima de l’intégrale d’action du problème newtonien de 4 corps de masses égales dans \mathbf{R}^3 : orbites “hip-hop”. *Celestial Mech. Dynam. Astronom.*, 77(2):139–152 (2001), 2000.
- [12] D. L. Ferrario and S. Terracini. On the existence of collisionless equivariant minimizers for the classical n -body problem. *Invent. Math.*, 155(2):305–362, 2004.
- [13] W. B. Gordon. A minimizing property of Keplerian orbits. *Amer. J. Math.*, 99(5):961–971, 1977.
- [14] H. Hofer. A geometric description of the neighbourhood of a critical point given by the mountain-pass theorem. *J. London Math. Soc. (2)*, 31(3):566–570, 1985.
- [15] X. Hu and G. Yu. Index theory for zero energy solutions of the planar anisotropic kepler problem. Preprint, 2017, arXiv:1705.05645.
- [16] P. Kustaanheimo and E. Stiefel. Perturbation theory of Kepler motion based on spinor regularization. *J. Reine Angew. Math.*, 218:204–219, 1965.
- [17] P. Majer and S. Terracini. Periodic solutions to some n -body type problems: the fixed energy case. *Duke Math. J.*, 69(3):683–697, 1993.
- [18] P. Majer and S. Terracini. Periodic solutions to some problems of n -body type. *Arch. Rational Mech. Anal.*, 124(4):381–404, 1993.
- [19] C. Marchal. How the method of minimization of action avoids singularities. *Celestial Mech. Dynam. Astronom.*, 83(1-4):325–353, 2002. Modern celestial mechanics: from theory to applications (Rome, 2001).
- [20] R. McGehee. Triple collision in the collinear three-body problem. *Invent. Math.*, 27:191–227, 1974.
- [21] R. McGehee. Double collisions for a classical particle system with nongravitational interactions. *Comment. Math. Helv.*, 56(4):524–557, 1981.
- [22] R. Montgomery. The N -body problem, the braid group, and action-minimizing periodic solutions. *Nonlinearity*, 11(2):363–376, 1998.
- [23] K. Tanaka. Noncollision solutions for a second order singular Hamiltonian system with weak force. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 10(2):215–238, 1993.
- [24] K. Tanaka. A prescribed energy problem for a singular Hamiltonian system with a weak force. *J. Funct. Anal.*, 113(2):351–390, 1993.
- [25] K. Tanaka. A note on generalized solutions of singular Hamiltonian systems. *Proc. Amer. Math. Soc.*, 122(1):275–284, 1994.
- [26] K. Tanaka. A prescribed-energy problem for a conservative singular Hamiltonian system. *Arch. Rational Mech. Anal.*, 128(2):127–164, 1994.
- [27] G. Yu. Shape Space Figure-8 Solution of Three Body Problem with Two Equal Masses. 2015. accepted by *Nonlinearity*, arXiv:1507.02892.
- [28] G. Yu. Simple choreographies of the planar Newtonian N -body problem. *Arch. Ration. Mech. Anal.*, 225(2):901–935, 2017.

E-mail address: guowei.yu@unito.it

DIPARTIMENTO DI MATEMATICA “GIUSEPPE PEANO”, UNIVERSITÀ DEGLI STUDI DI TORINO, ITALY