The Nehari manifold for fractional p-Laplacian system involving concave-convex nonlinearities and sign-changing weight functions

MAODING ZHEN

 School of Mathematics and Statistics, Huazhong University of Science and Technology, Wuhan 430074, China

2) Hubei Key Laboratory of Engineering Modeling and Scientific Computing, Huazhong University of Science and Technology, Wuhan, 430074, China

Abstract

In this paper, we consider a fractional p-Laplacian system (1.1) with both concaveconvex nonlinearities and sign-changing weight functions in bounded domains. With the help of the Nehari manifold, we prove that the system has at least two nontrivial solutions when the pair of the parameters (λ, μ) belongs to a certain subset of \mathbb{R}^n .

Keywords: fractional p-Laplacian system; Nehari manifold; Sign-changing weight functions

1 Introduction

In this paper, we study the following system involving fractional p-Laplacian:

$$\begin{cases} (-\Delta)_p^s u = \lambda f(x) |u|^{q-2} u + \frac{2\alpha}{\alpha+\beta} h(x) |u|^{\alpha-2} u |v|^{\beta} & \text{in } \Omega, \\ (-\Delta)_p^s v = \mu g(x) |v|^{q-2} v + \frac{2\beta}{\alpha+\beta} h(x) |u|^{\alpha} |v|^{\beta-2} v & \text{in } \Omega, \\ u = v = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$
(1.1)

where Ω is a smooth bounded domain in \mathbb{R}^n , $s \in (0, 1)$, n > ps, 1 < q < p and $\alpha > 1, \beta > 1$ satisfy $p < \alpha + \beta < p^* = \frac{np}{n-ps}$, where p^* is the fractional Sobolev exponent and $(-\Delta)_p^s$ is the fractional p-Laplacian operator which is defined as

$$(-\Delta)_{p}^{s}u(x) = 2\lim_{\epsilon \to 0} \int_{\mathbb{R}^{n} \setminus B_{\epsilon}(x)} \frac{|u(y) - u(x)|^{p-2}(u(y) - u(x))}{|x - y|^{n+ps}} dy, \quad x \in \mathbb{R}^{n}.$$

The pair of parameters $(\lambda, \mu) \in \mathbb{R}^n \setminus (0, 0)$ and the weight functions f, g, h satisfy the following conditions;

(A) $f, g \in L^{q^*}(\Omega)$ where $q^* = \frac{\alpha + \beta}{\alpha + \beta - q}$ and $f^+ = \max\{\pm f, 0\} \neq 0$ or $g^+ = \max\{\pm g, 0\} \neq 0$; (B) $h \in C(\overline{\Omega})$ with $\|h\|_{\infty} = 1$ and $h \ge 0$.

Recently, a great deal of attention has been focused on studying of equations or systems involving fractional Laplacian and corresponding nonlocal problems, both for their interesting theoretical structure and their concrete applications (see [2, 10, 15, 5, 6, 4, 25]

^{*} Corresponding author.

AMS Subject Classification: 35J50, 35R11

The authors were supported by the NSFC grant 11571125

E-mails:d201677010@hust.edu.cn

and references therein). This type of operator arises in a quite natural way in many different contexts, such as, the thin obstacle problem, finance, phase transitions, anomalous diffusion, flame propagation and many others(see [11, 16, 22] and references therein).

Compared to the Laplacian problem, the fractional p-Laplacian problem is nonlocal and more challenging. On the one hand, for the fractional elliptic problems when p = 2 has been investigated by many researchs. For example, C. Brändle, E. Colorado, A. de Pablo and U. Sánchez [3] studied the fractional Laplacian equation involving concave-convex nonlinearity for the subcritical case, they prove that there exists a finite parameter $\Lambda > 0$ such that for $0 < \lambda < \Lambda$ there exist at least two solutions, for $\lambda = \Lambda$ there exists at least one solution and for $\lambda > \Lambda$ there is no solution. Furthermore, B. Barrios, E. Colorado, A. de Pablo and U. Sánchez [2] studied the nonhomogeneous equation involving fractional Laplacian and proved the existence and multiplicity of solutions under suitable conditions of s and q. E. Colorado, A. de Pablo and U. Sánchez [10] studied the fractional equation with critical Sobolev exponent, they proved that the existence and the multiplicity of solutions under appropriate conditions on the size of f. For more other advances on this topic, see [18, 19] for the subcritical, [20, 21] for the critical case.

For the fractional p-Laplacian equation, S. Goyal, K. Sreenadh [13] studied the following equation involving concave-convex nonlinearities and sign-changing weight functions.

$$\begin{cases} (-\Delta)_p^s u = \lambda h(x) |u|^{q-1} u + b(x) |u|^{r-1} u & \text{in } \Omega, \\ u = 0 & \text{on } \mathbb{R}^n \setminus \Omega, \end{cases}$$

they showed that the existence and multiplicity of solutions by minimization on the suitable subset of Nehari manifold using the fibering maps and proved that there exists λ_0 such that for $\lambda \in (0, \lambda_0)$, it has at least two non-negative solutions.

B. Cheng, X. Tang [9] studied the existence of solutions for the following fractional p-Laplacian equation with sign-changing potential and nonlinearity

$$(-\Delta)_p^s u + v(x)|u|^{p-2}u = f(x,u) \quad \forall x \in \mathbb{R}^N,$$

where $p \ge 2$, $N \ge 2$, 0 < s < 1, $V \in C(\mathbb{R}^N, \mathbb{R})$ and $f \in C(\mathbb{R}^N \times \mathbb{R}, \mathbb{R})$, under the suitable conditions they prove the equation has at least one nontrivial solution.

It is also natural to study the coupled system of equations. W. Chen, S. Deng [8] considered the special case when f(x) = g(x) = h(x) = 1 for system (1.1). They prove that the system admits at least two nontrivial solutions under proper conditions of λ and μ .

The purpose of this paper is to study system (1.1) in the case of $2 < \alpha + \beta < p^*$, by variational methods and a Nehari manifold decomposition, we prove that the system admits at least two nontrivial solutions when the pair of parameters (λ, μ) belongs to certain subset of \mathbb{R}^2 .

To express the main results, we introduce

$$\Theta = \{ z \in \mathbb{R}^2 \setminus (0,0) \mid 0 < (|\lambda| \| f \|_{L^{q^\star}})^{\frac{p}{p-q}} + (|\mu| \| g \|_{L^{q^\star}})^{\frac{p}{p-q}} < C(\alpha,\beta,p,q,S) \}$$

and $C(\alpha,\beta,p,q,S) = \left[\frac{p-q}{2(\alpha+\beta-q)} S^{\frac{\alpha+\beta}{p}} \right]^{\frac{p}{\alpha+\beta-p}} \left(S^{-\frac{q}{p}} \frac{\alpha+\beta-q}{\alpha+\beta-p} \right)^{-\frac{p}{p-q}}.$
$$\Psi = \{ z \in \mathbb{R}^2 \setminus (0,0) \mid 0 < (|\lambda| \| f \|_{L^{q^\star}})^{\frac{p}{p-q}} + (|\mu| \| g \|_{L^{q^\star}})^{\frac{p}{p-q}} < D(\alpha,\beta,p,q,S) \}$$

and $D(\alpha, \beta, p, q, S) = \left(\frac{q}{p}\right)^{\frac{p}{p-q}} \left[\frac{p-q}{2(\alpha+\beta-q)}S^{\frac{\alpha+\beta}{p}}\right]^{\frac{p}{\alpha+\beta-p}} \left(S^{-\frac{q}{p}}\frac{\alpha+\beta-q}{\alpha+\beta-p}\right)^{-\frac{p}{p-q}}.$

Where S is the best Sobolev constant that will be introduced later. Our main results are:

Theorem 1.1. Suppose that the weight functions f, g, h be satisfied with the conditions (A) and (B), for each $(\lambda, \mu) \in \Theta$, then system (1.1) has at least one nontrivial solution in $W^{s,p}(\Omega) \times W^{s,p}(\Omega)$.

Theorem 1.2. Suppose that the weight functions f, g, h be satisfied with the conditions (A) and (B), for each $(\lambda, \mu) \in \Psi$, then system (1.1) has at least two nontrivial solutions in $W^{s,p}(\Omega) \times W^{s,p}(\Omega)$.

The paper is organized as follows. In section 2, we introduce some preliminaries. In section 3, we define the Nehari manifold and give some Lemmas that will be used later. In section 4, we prove the existence of Palais-Smale sequence. In section 5, we give the results of local minimization problem for system (1.1). Finally, the proofs of Theorem 1.1 and Theorem 1.2 are given in section 6.

2 Preliminaries

In this section, we introduce some preliminaries that will be used to establish the energy functional for system (1.1). Let $s \in (0, 1)$ and $p \in [1, +\infty)$, we define the usual fractional Sobolev space $W^{s,p}(\Omega)$ endowed with the norm

$$\|u\|_{W^{s,p}(\Omega)} = \|u\|_{L^p(\Omega)} + \left(\int_{\Omega \times \Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{n + ps}} dx dy\right)^{\frac{1}{p}}.$$
(2.1)

Let $Q = \mathbb{R}^{2n} \setminus (\mathcal{C}_{\Omega} \times \mathcal{C}_{\Omega})$ with $\mathcal{C}_{\Omega} = \mathbb{R}^n \setminus \Omega$. We define

$$X = \left\{ u \mid u : \mathbb{R}^n \to \mathbb{R} \text{ is measurable and } \int_Q \frac{|u(x) - u(y)|^p}{|x - y|^{n + ps}} dx dy < +\infty \right\}.$$

The space X is endowed with the norm defined by

$$||u||_X = ||u||_{L^p(\Omega)} + \left(\int_Q \frac{|u(x) - u(y)|^p}{|x - y|^{n + ps}} dx dy\right)^{\frac{1}{p}}.$$
(2.2)

The functional space X_0 denotes the closure of $C_0^{\infty}(\Omega)$ in X. By Theorem 6.5 and Theorem 7.1 in [11] the space X_0 is a Hilbert space which can be endowed with the scalar product defined for any $\phi, \psi \in X_0$ as

$$\langle \phi, \psi \rangle_{X_0} = \int_Q \frac{|\phi(x) - \phi(y)|^{p-1}(\psi(x) - \psi(y))}{|x - y|^{n+ps}} dx dy$$
(2.3)

and the norm

$$||u||_{X_0} = \left(\int_Q \frac{|u(x) - u(y)|^p}{|x - y|^{n + ps}} dx dy\right)^{\frac{1}{p}}$$
(2.4)

is equivalent to the usual one defined in (2.1). Since u = 0 a.e. in $\mathbb{R}^n \setminus \Omega$, we have the (2.2), (2.3) and (2.4) can be extended to all \mathbb{R}^n . By results of [13, 11], the embedding $X_0 \hookrightarrow L^r(\Omega)$ is continuous for any $r \in [1, p^*]$ and compact whenever $r \in [1, p^*)$. Let S be the best Sobolev constant for the embedding of $X_0 \hookrightarrow L^{\alpha+\beta}(\Omega)$ defined by

$$S = \inf_{z \in X_0 \setminus 0} \frac{\int_Q \frac{|u(x) - u(y)|^p}{|x - y|^{n + ps}} dx dy}{\left(\int_\Omega |u(x)|^{\alpha + \beta} dx\right)^{\frac{p}{\alpha + \beta}}}.$$

For further details on X and X_0 and also for their properties we refer [11] and the references therein. Let $E = X_0 \times X_0$ be the Cartesian product of two Hilbert spaces, which is a reflexive Banach space endowed with the norm

$$|(u,v)|| = \left(||u||_{X_0}^p + ||v||_{X_0}^p\right)^{\frac{1}{p}} = \left(\int_Q \frac{|u(x) - u(y)|^p}{|x - y|^{n + ps}} dx dy + \int_Q \frac{|v(x) - v(y)|^p}{|x - y|^{n + ps}} dx dy\right)^{\frac{1}{p}}.$$
(2.5)

Definition 2.1. We say that $(u, v) \in E$ is a weak solution of (1.1) if the identity

$$\begin{split} &\int_{Q} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))(\varphi_{1}(x) - \varphi_{1}(y))}{|x - y|^{n+ps}} dx dy \\ &+ \int_{Q} \frac{|v(x) - v(y)|^{p-2} (v(x) - v(y))(\varphi_{2}(x) - \varphi_{2}(y))}{|x - y|^{n+ps}} dx dy \\ &= \int_{\Omega} \left(\lambda f |u|^{q-2} u\varphi_{1} + \mu g |v|^{q-2} v\varphi_{2}\right) dx + \frac{2\alpha}{\alpha + \beta} \int_{\Omega} h |u|^{\alpha - 2} u |v|^{\beta} \varphi_{1} dx + \frac{2\beta}{\alpha + \beta} \int_{\Omega} h |u|^{\alpha} |v|^{\beta - 2} v\varphi_{2} dx \end{split}$$

holds for all $(\varphi_1, \varphi_2) \in E$.

Note that, the energy functional associated with (1.1) is given by

$$I_{\lambda,\mu}(u,v) := \frac{1}{p} \int_{Q} \frac{|u(x) - u(y)|^{p}}{|x - y|^{n + ps}} dx dy + \frac{1}{p} \int_{Q} \frac{|v(x) - v(y)|^{p}}{|x - y|^{n + ps}} dx dy \qquad (2.6)$$
$$- \frac{1}{q} \int_{\Omega} (\lambda f |u|^{q} + \mu g |v|^{q}) dx - \frac{2}{\alpha + \beta} \int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx.$$

In the end of this section, we recall some notations that will be used in the sequel. • $L^p(\Omega)$, $1 \le p \le \infty$ denotes Lebesgue space with norm $\|\cdot\|_p$ and $E = X_0 \times X_0$ is equipped with the norm $\|(u,v)\| = \left(\|u\|_{X_0}^p + \|v\|_{X_0}^p\right)^{\frac{1}{p}} = \left(\int_Q \frac{|u(x)-u(y)|^p}{|x-y|^{n+ps}} dxdy + \int_Q \frac{|v(x)-v(y)|^p}{|x-y|^{n+ps}} dxdy\right)^{\frac{1}{p}}$. • The dual space of a Banach space E will be denoted by E^{-1} . We set t(u,v) = (tu,tv)for all $(u,v) \in E$ and $t \in \mathbb{R}$, z = (u,v) is said to be positive if u(x,y) > 0, v(x,y) > 0 in E and to be non-negative if $u(x,y) \ge 0$, $v(x,y) \ge 0$ in E.

- B(0;r) is the ball at the origin with radius r. $o_n(1)$ denotes $o_n(1) \to 0$ as $n \to +\infty$.
- C, C_i, c will denote various positive constants which may vary from line to line.

3 The Nehari manifold

We consider the Nehari minimization problem: for $(\lambda, \mu) \in \mathbb{R}^2 \setminus (0, 0)$,

$$\theta_{\lambda,\mu} = \inf\{I_{\lambda,\mu}(u,v) \mid (u,v) \in N_{\lambda,\mu}\}$$

where $N_{\lambda,\mu} := \{(u,v) \in E \setminus \{(0,0)\} \mid \langle I'_{\lambda,\mu}(u,v), (u,v) \rangle = 0\}$ and

$$\langle I'_{\lambda,\mu}(u,v),(u,v)\rangle = \|(u,v)\|^p - \int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) \, dx - 2 \int_{\Omega} h |u|^\alpha |v|^\beta \, dx.$$
(3.1)

Note that $N_{\lambda,\mu}$ contains every nonzero solution of problem (1.1).

Define

$$\langle \Phi_{\lambda,\mu}(u,v),(u,v)\rangle = \langle I'_{\lambda,\mu}(u,v),(u,v)\rangle$$

Then

$$\langle \Phi'_{\lambda,\mu}(u,v),(u,v)\rangle = p\|(u,v)\|^p - q \int_{\Omega} \left(\lambda f|u|^q + \mu g|v|^q\right) dx - 2(\alpha+\beta) \int_{\Omega} h|u|^{\alpha}|v|^{\beta} dx.$$

Moreover, if $\int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) dx \neq 0$ and $(u, v) \in N_{\lambda, \mu}$, we have

$$\langle \Phi'_{\lambda,\mu}(u,v), (u,v) \rangle = (p-q) ||(u,v)||^p - 2(\alpha + \beta - q) \int_{\Omega} h|u|^{\alpha} |v|^{\beta} dx.$$
 (3.2)

Similarly to the method used in [24], we split $N_{\lambda,\mu}$ into three parts.

$$\begin{split} N^{+}_{\lambda,\mu} &= \{(u,v) \in N_{\lambda,\mu} \mid \langle \Phi'_{\lambda,\mu}(u,v), (u,v) \rangle > 0\}; \\ N^{0}_{\lambda,\mu} &= \{(u,v) \in N_{\lambda,\mu} \mid \langle \Phi'_{\lambda,\mu}(u,v), (u,v) \rangle = 0\}; \\ N^{-}_{\lambda,\mu} &= \{(u,v) \in N_{\lambda,\mu} \mid \langle \Phi'_{\lambda,\mu}(u,v), (u,v) \rangle < 0\}. \end{split}$$

Then, we have the following results

Lemma 3.1. For each $(\lambda, \mu) \in \Theta$, we have $N^0_{\lambda,\mu} = \emptyset$.

Proof. We consider the following two cases

Case 1: $(u, v) \in N_{\lambda, \mu}$ and $\int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx \leq 0$, we have

$$\int_{\Omega} \left(\lambda f |u|^{q} + \mu g |v|^{q}\right) dx = \|(u, v)\|^{p} - 2 \int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx > 0.$$

Thus $\langle \Phi'_{\lambda,\mu}(u,v),(u,v)\rangle = (p-q)\|(u,v)\|^p - 2(\alpha+\beta-q)\int_{\Omega}h|u|^{\alpha}|v|^{\beta}dx > 0$ and so $(u,v) \notin N^0_{\lambda,\mu}$.

Case 2: $(u,v) \in N_{\lambda,\mu}$ and $\int_{\Omega} h|u|^{\alpha}|v|^{\beta}dx > 0$. Suppose that $N^{0}_{\lambda,\mu} \neq \emptyset$ for all $(\lambda,\mu) \in \mathbb{R}^{2} \setminus (0,0)$. Then for each $(u,v) \in N^{0}_{\lambda,\mu}$, we have

$$\langle \Phi'_{\lambda,\mu}(u,v),(u,v)\rangle = (p-q)\|(u,v)\|^p - 2(\alpha+\beta-q)\int_{\Omega}h|u|^{\alpha}|v|^{\beta}dx = 0.$$
(3.3)

Thus

$$|(u,v)||^p = \frac{2(\alpha+\beta-q)}{p-q} \int_{\Omega} h|u|^{\alpha}|v|^{\beta} dx$$

and

$$\begin{split} \int_{\Omega} \left(\lambda f |u|^{q} + \mu g |v|^{q}\right) dx &= \|(u,v)\|^{p} - 2 \int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx \\ &= \frac{2(\alpha + \beta - p)}{p - q} \int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx > 0. \end{split}$$

By the Hölder and Sobolev inequalities, we have

$$\|(u,v)\| \ge \left[\frac{p-q}{2(\alpha+\beta-q)}S^{\frac{\alpha+\beta}{p}}\right]^{\frac{1}{\alpha+\beta-p}}$$
(3.4)

and

$$\begin{aligned} \frac{\alpha+\beta-p}{\alpha+\beta-q} \|(u,v)\|^p &= \|(u,v)\|^p - 2\int_{\Omega} h|u|^{\alpha}|v|^{\beta}dx = \int_{\Omega} \left(\lambda f|u|^q + \mu g|v|^q\right)dx\\ &\leq |\lambda|\|f\|_{L^{q^{\star}}} \|\|u\|_{L^{\alpha+\beta}}^q + |\mu|\|g\|_{L^{q^{\star}}} \|\|v\|_{L^{\alpha+\beta}}^q\\ &\leq \left[\left(|\lambda|\|f\|_{L^{q^{\star}}}\right)^{\frac{p}{p-q}} + \left(|\mu|\|g\|_{L^{q^{\star}}}\right)^{\frac{p}{p-q}} \frac{p^{-q}}{p}S^{-\frac{q}{p}}\|(u,v)\|^q.\end{aligned}$$

This implies

$$\|(u,v)\| \le \left(S^{-\frac{q}{p}}\frac{\alpha+\beta-q}{\alpha+\beta-p}\right)^{\frac{1}{p-q}} \left[(|\lambda|\|f\|_{L^{q^{\star}}})^{\frac{p}{p-q}} + (|\mu|\|g\|_{L^{q^{\star}}})^{\frac{p}{p-q}} \right]^{\frac{1}{p}}.$$
 (3.5)

By (3.4) and (3.5), we have

$$[(|\lambda|||f||_{L^{q^{\star}}})^{\frac{p}{p-q}} + (|\mu|||g||_{L^{q^{\star}}})^{\frac{p}{p-q}}] \ge \left[\frac{p-q}{2(\alpha+\beta-p)}S^{\frac{\alpha+\beta}{p}}\right]^{\frac{p}{\alpha+\beta-p}} \left(S^{-\frac{q}{p}}\frac{\alpha+\beta-q}{\alpha+\beta-p}\right)^{-\frac{p}{p-q}},$$

contradicting with the assumption.

Lemma 3.1 suggests that for each $(\lambda, \mu) \in \Theta$, we can write $N_{\lambda,\mu} = N^+_{\lambda,\mu} \cup N^-_{\lambda,\mu}$. Next, we define

$$\theta^+_{\lambda,\mu} = \inf_{z \in N^+_{\lambda,\mu}} I_{\lambda,\mu}(z) \quad and \quad \theta^-_{\lambda,\mu} = \inf_{z \in N^-_{\lambda,\mu}} I_{\lambda,\mu}(z).$$

The following Lemma shows that the minimizer on $N_{\lambda,\mu}$ is critical point for $I_{\lambda,\mu}$

Lemma 3.2. For each $(\lambda, \mu) \in \Theta$, let (u_0, v_0) be a local minimizer for $I_{\lambda,\mu}$ on $N_{\lambda,\mu}$, then $I'_{\lambda,\mu}(u_0, v_0) = 0$ in E^{-1} .

Proof. Since (u_0, v_0) is a local minimizer for $I_{\lambda,\mu}$ on $N_{\lambda,\mu}$, that is (u_0, v_0) is a solution of the optimization problem

$$\min\{I_{\lambda,\mu}(u,v) \mid \Phi_{\lambda,\mu}(u,v) = 0\}.$$

Then, by the theory of Lagrange multipliers, there exists a constant $L \in \mathbb{R}$ such that

$$\langle I'_{\lambda,\mu}(u_0, v_0), (u_0, v_0) \rangle = L \langle \Phi'_{\lambda,\mu}(u_0, v_0), (u_0, v_0) \rangle.$$

Since $(u_0, v_0) \notin N^0_{\lambda,\mu}$, we have $\langle \Phi'_{\lambda,\mu}(u_0, v_0), (u_0, v_0) \rangle \neq 0$, thus L = 0, this completes the proof.

Moreover, we have the following properties about the Nehari manifold $N_{\lambda,\mu}$.

Lemma 3.3. we have

(i) If $(u, v) \in N^+_{\lambda,\mu}$, then $\int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) dx > 0$ (ii) If $(u, v) \in N^-_{\lambda,\mu}$, then $\int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx > 0$

Proof. (i) We consider the following two cases. Case 1: If $\int_{\Omega} h|u|^{\alpha}|v|^{\beta}dx \leq 0$, we have

$$\int_{\Omega} \left(\lambda f |u|^{q} + \mu g |v|^{q}\right) dx = \|(u, v)\|^{p} - 2 \int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx > 0.$$

Case 2: If $\int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx > 0$, since

$$||(u,v)||^{p} - \int_{\Omega} (\lambda f|u|^{q} + \mu g|v|^{q}) \, dx - 2 \int_{\Omega} h|u|^{\alpha} |v|^{\beta} \, dx = 0$$

and

$$\langle \Phi_{\lambda,\mu}'(u,v),(u,v)\rangle = p\|(u,v)\|^p - q \int_{\Omega} \left(\lambda f|u|^q + \mu g|v|^q\right) dx - 2(\alpha+\beta) \int_{\Omega} h|u|^{\alpha}|v|^{\beta} dx > 0,$$

it follows that

$$(p-q)\int_{\Omega} \left(\lambda f|u|^{q} + \mu g|v|^{q}\right) dx - 2(\alpha + \beta - q)\int_{\Omega} h|u|^{\alpha}|v|^{\beta} dx > 0,$$

which implies

$$\int_{\Omega} \left(\lambda f|u|^{q} + \mu g|v|^{q}\right) dx > \frac{2(\alpha + \beta - q)}{p - q} \int_{\Omega} h|u|^{\alpha} |v|^{\beta} dx > 0.$$

(ii) We consider the following two cases.

Case 1: If $\int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) dx = 0$, we have

$$2\int_{\Omega} h|u|^{\alpha}|v|^{\beta}dx = \|(u,v)\|^{p} > 0.$$

Case 2: If $\int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) dx \neq 0$, we have

$$(p-q)\|(u,v)\|^p - 2(\alpha+\beta-q)\int_{\Omega}h|u|^{\alpha}|v|^{\beta}dx = \langle \Phi'_{\lambda,\mu}(u,v), (u,v)\rangle < 0.$$

Thus $\int_{\Omega} h|u|^{\alpha}|v|^{\beta}dx > 0.$

Lemma 3.4. The following facts hold

(i) If $(\lambda, \mu) \in \Theta$, then we have $\theta_{\lambda,\mu} \leq \theta^+_{\lambda,\mu} < 0$ (ii If $(\lambda, \mu) \in \Psi$, then we have $\theta^-_{\lambda,\mu} > c_0$ for some positive constant c_0 depending on λ, μ, p, q, S .

(iii) The energy functional $I_{\lambda,\mu}$ is bounded below and coercive on $N_{\lambda,\mu}$.

Proof. (i) Let $(u, v) \in N^+_{\lambda,\mu}$, by (3.2), we have

$$\frac{p-q}{2(\alpha+\beta-q)}\|(u,v)\|^p > \int_{\Omega} h|u|^{\alpha}|v|^{\beta}dx.$$

Hence

$$\begin{split} I_{\lambda,\mu}(u,v) &= \left(\frac{1}{p} - \frac{1}{q}\right) \|(u,v)\|^p + 2\left(\frac{1}{q} - \frac{1}{\alpha+\beta}\right) \int_{\Omega} h|u|^{\alpha}|v|^{\beta} dx\\ &\leq \left[\left(\frac{1}{p} - \frac{1}{q}\right) + \left(\frac{1}{q} - \frac{1}{\alpha+\beta}\right) \frac{p-q}{\alpha+\beta-q}\right] \|(u,v)\|^p\\ &\leq \frac{(q-p)(\alpha+\beta-p)}{pq(\alpha+\beta)} \|(u,v)\|^p < 0. \end{split}$$

Therefore, by the definition of $\theta_{\lambda,\mu}$, $\theta^+_{\lambda,\mu}$, we can deduce that $\theta_{\lambda,\mu} \leq \theta^+_{\lambda,\mu} < 0$. (*ii*) Let $(u, v) \in N^-_{\lambda,\mu}$, by (3.2), we have

$$\frac{p-q}{2(\alpha+\beta-q)}\|(u,v)\|^p < \int_{\Omega} h|u|^{\alpha}|v|^{\beta}dx.$$

By the Hölder inequality and the Sobolev embedding theorem, we have

$$\int_{\Omega} h|u|^{\alpha}|v|^{\beta}dx \le S^{-\frac{\alpha+\beta}{p}} ||(u,v)||^{\alpha+\beta}$$

Hence

$$\|(u,v)\| > \left(\frac{p-q}{2(\alpha+\beta-q)}S^{\frac{\alpha+\beta}{p}}\right)^{\frac{1}{\alpha+\beta-p}} \text{ for all } z \in N^{-}_{\lambda,\mu}.$$
(3.6)

By (3.6), we have

$$\begin{split} I_{\lambda,\mu}(u,v) &= \frac{\alpha + \beta - p}{p(\alpha + \beta)} \|(u,v)\|^p - \frac{\alpha + \beta - q}{q(\alpha + \beta)} \int_{\Omega} \left(\lambda f |u|^q + \mu g |v|^q\right) dx \\ &\geq \|(u,v)\|^q \left[\frac{\alpha + \beta - p}{p(\alpha + \beta)} \|(u,v)\|^{p-q} - \frac{\alpha + \beta - q}{q(\alpha + \beta)} [(|\lambda| \| f \|_{L^{q^*}})^{\frac{p}{p-q}} + (|\mu| \| g \|_{L^{q^*}})^{\frac{p}{p-q}} \right] \\ &> \left\{ -\frac{\alpha + \beta - q}{q(\alpha + \beta)} [(|\lambda| \| f \|_{L^{q^*}})^{\frac{p}{p-q}} + (|\mu| \| g \|_{L^{q^*}})^{\frac{p}{p-q}} \right]^{\frac{p-q}{p}} S^{-\frac{q}{p}} + \frac{\alpha + \beta - p}{p(\alpha + \beta)} \left(\frac{p - q}{2(\alpha + \beta - q)} S^{\frac{\alpha + \beta}{p}} \right)^{\frac{p-q}{\alpha + \beta - p}} \right\} \times \left(\frac{p - q}{2(\alpha + \beta - q)} S^{\frac{\alpha + \beta}{p}} \right)^{\frac{q}{\alpha + \beta - p}} . \end{split}$$

Thus, if $(\lambda, \mu) \in \Psi$, then

$$I_{\lambda,\mu} > c_0$$
, for all $z \in N^-_{\lambda,\mu}$,

for some positive constant $c_0 = c_0(\lambda, \mu, p, q, S)$.

(*iii*) Let $(u, v) \in N_{\lambda,\mu}$, by (3.1), Hölder and Sobolev inequality, we have

$$I_{\lambda,\mu}(u,v) = \frac{\alpha + \beta - p}{p(\alpha + \beta)} \|(u,v)\|^p - \frac{\alpha + \beta - q}{q(\alpha + \beta)} \int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) dx$$

$$\geq \frac{\alpha + \beta - p}{p(\alpha + \beta)} \|(u,v)\|^p - \frac{\alpha + \beta - q}{q(\alpha + \beta)} [(|\lambda| \|f\|_{L^{q^*}})^{\frac{p}{p-q}} + (|\mu| \|g\|_{L^{q^*}})^{\frac{p}{p-q}}]^{\frac{p-q}{p}} S^{-\frac{q}{p}} \|(u,v)\|^q.$$

Since 1 < q < p, then the energy functional $I_{\lambda,\mu}$ is bounded below and coercive on $N_{\lambda,\mu}$.

For each $(\lambda, \mu) \in N^{-}_{\lambda,\mu}$, we write

$$t_{max} = \left(\frac{(p-q)\|(u,v)\|^p}{2(\alpha+\beta-q)\int_{\Omega}h|u|^{\alpha}|v|^{\beta}dx}\right)^{\frac{1}{\alpha+\beta-p}} > 0.$$

Then the following Lemma holds.

Lemma 3.5. For each $(\lambda, \mu) \in \Theta$ and $(u, v) \in N_{\lambda,\mu}^-$, we have (i) If $\int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) dx \leq 0$, then there exists a unique $(t^-u, t^-v) > 0$ such that $(t^-u, t^-v) \in N_{\lambda,\mu}^-$ and $I_{\lambda,\mu}(t^-u, t^-v) = \max_{t>0} I_{\lambda,\mu}(tu, tv)$. (ii) If $\int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) dx > 0$, then there exist unique $0 < t^+ < t_{max} < t^-$, such that $(t^+u, t^+v) \in N_{\lambda,\mu}^+$, $(t^-u, t^-v) \in N_{\lambda,\mu}^-$ and $I_{\lambda,\mu}(t^+u, t^+v) = \min_{0 < t < t_{max}} I_{\lambda,\mu}(tu, tv)$, $I_{\lambda,\mu}(t^-u, t^-v) = \max_{t\geq 0} I_{\lambda,\mu}(tu, tv)$.

Proof. Fix $(u, v) \in N^{-}_{\lambda,\mu}$, by Lemma 3.3, we have $\int_{\Omega} h|u|^{\alpha}|v|^{\beta}dx > 0$. Let

$$m(t) = t^{p-q} ||(u,v)||^p - 2t^{\alpha+\beta-q} \int_{\Omega} h|u|^{\alpha} |v|^{\beta} dx, \text{ for } t \ge 0$$

Clearly, $m(0) = 0, m(t) \to -\infty$ as $t \to \infty$. Since

$$m'(t) = (p-q)t^{p-q-1} ||(u,v)||^p - 2(\alpha + \beta - q)t^{\alpha + \beta - q - 1} \int_{\Omega} h|u|^{\alpha} |v|^{\beta} dx,$$

we have that m(t) is increasing for $t \in [0, t_{max})$, decreasing for $t \in (t_{max}, +\infty)$ and achieves its maximum at t_{max} . Moreover,

$$\begin{split} m(t_{max}) &= \left(\frac{(p-q)\|(u,v)\|^p}{2(\alpha+\beta-q)\int_{\Omega}h|u|^{\alpha}|v|^{\beta}dx}\right)^{\frac{p-q}{\alpha+\beta-p}}\|(u,v)\|^p\\ &- 2\left(\frac{(p-q)\|(u,v)\|^p}{2(\alpha+\beta-q)\int_{\Omega}h|u|^{\alpha}|v|^{\beta}dx}\right)^{\frac{\alpha+\beta-q}{\alpha+\beta-p}}\int_{\Omega}h|u|^{\alpha}|v|^{\beta}dx\\ &= \|(u,v)\|^q\left[\left(\frac{p-q}{2(\alpha+\beta-q)}\right)^{\frac{p-q}{\alpha+\beta-p}} - 2\left(\frac{p-q}{2(\alpha+\beta-q)}\right)^{\frac{\alpha+\beta-q}{\alpha+\beta-p}}\right]\left(\frac{\|(u,v)\|^{\alpha+\beta}}{\int_{\Omega}h|u|^{\alpha}|v|^{\beta}dx}\right)^{\frac{p-q}{\alpha+\beta-p}}\\ &\geq \|(u,v)\|^q\left(\frac{\alpha+\beta-p}{\alpha+\beta-q}\right)\left(S^{\frac{\alpha+\beta}{p}}\frac{p-q}{2(\alpha+\beta-q)}\right)^{\frac{p-q}{\alpha+\beta-p}}.\end{split}$$

That is

$$m(t_{max}) \ge \|(u,v)\|^q \left(\frac{\alpha+\beta-p}{\alpha+\beta-q}\right) \left(S^{\frac{\alpha+\beta}{p}} \frac{p-q}{2(\alpha+\beta-q)}\right)^{\frac{p-q}{\alpha+\beta-p}}.$$
(3.7)

(i) If $\int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) dx \leq 0$, by the property of m(t), there is a unique $t^- > t_{max}$ such that $m(t^-) = \int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) dx$ and $m'(t^-) < 0$. Since

$$\begin{split} \langle \Phi_{\lambda,\mu}'(t^{-}u,t^{-}v),(t^{-}u,t^{-}v)\rangle &= (p-q)(t^{-})^{p} \|(u,v)\|^{p} - 2(\alpha+\beta-q)(t^{-})^{\alpha+\beta} \int_{\Omega} h|u|^{\alpha}|v|^{\beta} dx \\ (3.8) \\ &= (t^{-})^{1+q} \left[(p-q)(t^{-})^{p-q-1} \|(u,v)\|^{p} - 2(\alpha+\beta-q)(t^{-})^{\alpha+\beta-q-1} \int_{\Omega} h|u|^{\alpha}|v|^{\beta} dx \right] \\ &= (t^{-})^{1+q} m'(t^{-}) < 0 \end{split}$$

and

$$\begin{split} \langle I'_{\lambda,\mu}(t^-u,t^-v),(t^-u,t^-v)\rangle &= (t^-)^p \|(u,v)\|^p - (t^-)^q \int_{\Omega} \left(\lambda f |u|^q + \mu g |v|^q\right) dx - 2(t^-)^{\alpha+\beta} \int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx \\ &= (t^-)^q \left[(t^-)^{p-q} \|(u,v)\|^p - \int_{\Omega} \left(\lambda f |u|^q + \mu g |v|^q\right) dx - 2(t^-)^{\alpha+\beta-q} \int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx \right] \\ &= (t^-)^q \left[m(t^-) - \int_{\Omega} \left(\lambda f |u|^q + \mu g |v|^q\right) dx \right] = 0. \end{split}$$

Thus $(t^-u, t^-v) \in N^-_{\lambda,\mu}$.

For $t > t_{max}$, by (3.8), we know

$$(p-q)t^p ||(u,v)||^p - 2(\alpha+\beta-q)t^{\alpha+\beta} \int_{\Omega} h|u|^{\alpha}|v|^{\beta}dx < 0.$$

When $tz \in N_{\lambda,\mu}$, we have

$$||(u,v)||^{p} - t^{q-p} \int_{\Omega} \left(\lambda f|u|^{q} + \mu g|v|^{q}\right) dx - 2t^{\alpha+\beta-p} \int_{\Omega} h|u|^{\alpha}|v|^{\beta} dx = 0$$

and

$$\begin{aligned} \frac{d^2}{dt^2} I_{\lambda,\mu}(tu,tv) &= (p-1)t^{p-2} \|(u,v)\|^p - (q-1)t^{q-2} \int_{\Omega} \left(\lambda f |u|^q + \mu g |v|^q\right) dx - 2(\alpha + \beta - 1)t^{\alpha + \beta - 2} \int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx. \end{aligned}$$
Consequently,
$$\frac{d^2}{dt^2} I_{\lambda,\mu}(tu,tv) &= t^{q-1} m'(t) < 0. \end{aligned}$$

Since

$$\frac{d}{dt}I_{\lambda,\mu}(tu,tv) = t^{p-1} ||(u,v)||^p - t^{q-1} \int_{\Omega} \left(\lambda f |u|^q + \mu g |v|^q\right) dx - 2t^{\alpha+\beta-1} \int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx.$$

We have $\frac{d}{dt}I_{\lambda,\mu}(tu,tv) = 0$ for $t = t^-$. Thus, $I_{\lambda,\mu}(t^-z) = \max_{t>0} I_{\lambda,\mu}(tz)$. (*ii*) If $\int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) dx > 0$.

Since

$$\begin{split} m(0) &= 0 < \int_{\Omega} \left(\lambda f |u|^{q} + \mu g |v|^{q}\right) dx \\ &\leq \left[\left(|\lambda| \|f\|_{L^{q^{\star}}} \right)^{\frac{p}{p-q}} + \left(|\mu| \|g\|_{L^{q^{\star}}} \right)^{\frac{p}{p-q}} \right]^{\frac{p-q}{p}} S^{-\frac{q}{p}} \|(u,v)\|^{q} \\ &\leq \|(u,v)\|^{q} \left(\frac{\alpha+\beta-p}{\alpha+\beta-q}\right) \left(S^{\frac{\alpha+\beta}{p}} \frac{p-q}{2(\alpha+\beta-q)}\right)^{\frac{p-q}{\alpha+\beta-p}} \\ &\leq m(t_{max}) \ for \ (\lambda,\mu) \in \Theta, \end{split}$$

therefore, there are unique t^+ and t^- such that $0 < t^+ < t_{max} < t^-$,

$$m(t^{+}) = \int_{\Omega} (\lambda f |u|^{q} + \mu g |v|^{q}) \, dx = m(t^{-})$$

and

$$m'(t^+) > 0 > m'(t^-).$$

Thus, by the same arguments as (i), we have $(t^+u, t^+v) \in N^+_{\lambda,\mu}$, $(t^-u, t^-v) \in N^-_{\lambda,\mu}$, $I_{\lambda,\mu}(t^-u, t^-v) \geq I_{\lambda,\mu}(tu, tv) \geq I_{\lambda,\mu}(t^+u, t^+v)$ for each $t \in [t^+, t^-]$ and $I_{\lambda,\mu}(t^+u, t^+v) \leq I_{\lambda,\mu}(tu, tv)$ for each $t \in [0, t^+]$.

That is

$$I_{\lambda,\mu}(t^+u, t^+v) = \min_{0 < t < t_{max}} I_{\lambda,\mu}(tu, tv), \quad I_{\lambda,\mu}(t^-u, t^-v) = \max_{t \ge 0} I_{\lambda,\mu}(tu, tv).$$

4 Existence of Palais-Smale sequence

Definition 4.1. We say that $(u_n, v_n) \in E$ is a $(PS)_c$ sequence in E for $I_{\lambda\mu}$, if $I_{\lambda\mu}(u_n, v_n) = c + o_n(1)$ and $I'_{\lambda\mu}(u_n, v_n) = o_n(1)$ strongly in E^{-1} as $n \to \infty$. If any $(PS)_c$ sequence in E for $I_{\lambda\mu}$ admits a convergent subsequence, we say that $I_{\lambda\mu}$ satisfies the $(PS)_c$ condition.

First, we will use the idea of Tarantello [23] to get the following results.

Lemma 4.2. Let z = (u, v) and $(\lambda, \mu) \in \Theta$, then for each $z \in N_{\lambda,\mu}$, there exists r > 0 and a differentiable function $\xi : B(0,r) \subset E \to \mathbb{R}^+$ such that $\xi(0) = 1$ and $\xi(m)(z-m) \in N_{\lambda,\mu}$ for every $m \in B(0,r)$. Let

$$\begin{split} \mathbf{T}_{1} &:= p \int_{Q} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y)) (m_{1}(x) - m_{1}(y))}{|x - y|^{n + ps}} dx dy \\ &+ p \int_{Q} \frac{|v(x) - v(y)|^{p-2} (v(x) - v(y)) (m_{2}(x) - m_{2}(y))}{|x - y|^{n + ps}} dx dy, \\ \mathbf{T}_{2} &:= q \int_{\Omega} \left(\lambda f |u|^{q-2} um_{1} + \mu g |v|^{q-2} vm_{2} \right) dx, \\ \mathbf{T}_{3} &:= 2 \int_{\Omega} \left(\alpha h |u|^{\alpha - 2} um_{1} |v|^{\beta} + \beta h |u|^{\alpha} |v|^{\beta - 2} vm_{2} \right) dx, \end{split}$$

then

$$\langle \xi'(0), m \rangle = \frac{\mathrm{T}_2 + \mathrm{T}_3 - \mathrm{T}_1}{(p-q) \| (u,v) \|^p - 2(\alpha + \beta - q) \int_{\Omega} h |u|^{\alpha} |v|^{\beta} dx}$$
(4.1)

holds for all $m \in E$.

Proof. For $z \in N_{\lambda,\mu}$, define a function $F : \mathbb{R} \times E \to \mathbb{R}$ by

$$\begin{split} F_{z}(\xi,q) &:= \langle I_{\lambda,\mu}'(\xi(z-q)), \xi(z-q) \rangle \\ &= \xi^{p} \int_{Q} \frac{|(u(x)-q_{1}(x))-(u(y)-q_{1}(y))|^{p}}{|x-y|^{n+ps}} dx dy + \xi^{p} \int_{Q} \frac{|(v(x)-q_{2}(x))-(v(y)-q_{2}(y))|^{p}}{|x-y|^{n+ps}} dx dy \\ &- \xi^{q} \int_{\Omega} (\lambda f |u-q_{1}|^{q} + \mu g |v-q_{2}|^{q}) dx - 2\xi^{\alpha+\beta} \int_{\Omega} h |u-q_{1}|^{\alpha} |v-q_{2}|^{\beta} dx. \end{split}$$

Then, $F_z(1,0) = \langle I'_{\lambda,\mu}(u,v), (u,v) \rangle = 0$ and by Lemma 3.1, we have $N^0_{\lambda,\mu} = \emptyset$. That is

$$\frac{dF_z(1,0)}{d\xi} = p \|(u,v)\|^p - q \int_{\Omega} (\lambda f |u|^q + \mu g |v|^q) \, dx - 2(\alpha + \beta) \int_{\Omega} h |u|^\alpha |v|^\beta \, dx$$
$$= (p-q) \|(u,v)\|^p - 2(\alpha + \beta - q) \int_{\Omega} h |u|^\alpha |v|^\beta \, dx \neq 0.$$

According to the implicit function theorem, there exist r > 0 and a differentiable function $\xi : B(0,r) \subset E \to \mathbb{R}^+$ such that $\xi(0) = 1$ and (4.1) holds. Moreover, $F_z(\xi(m),m) = 0$ holds for all $m \in B(0,r)$ is equivalent to $\langle I'_{\lambda,\mu}(\xi(m)(z-m)), \xi(m)(z-m) \rangle = 0$ for all $m \in B(0,r)$. That is $\xi(m)(z-m) \in N_{\lambda,\mu}$.

Lemma 4.3. Let z = (u, v) and $(\lambda, \mu) \in \Theta$, then for each $z \in N_{\lambda,\mu}^-$, there exists r > 0 and a differentiable function $\xi^- : B(0, r) \subset E \to \mathbb{R}^+$ such that $\xi^-(0) = 1$ and $\xi^-(m)(z-m) \in N_{\lambda,\mu}$ for every $m \in B(0, r)$ and formula (4.1) holds.

Proof. Similar to the argument in Lemma 4.2, there exists r > 0 and a differentiable function $\xi^- : B(0,r) \subset E \to \mathbb{R}^+$ such that $\xi^-(0) = 1$ and $\xi^-(m)(z-m) \in N_{\lambda,\mu}$ for every $m \in B(0,r)$ and formula (4.1) holds. Since

$$\langle \Phi'_{\lambda,\mu}(z), z \rangle = (p-q) ||(u,v)||^p - 2(\alpha + \beta - q) \int_{\Omega} h|u|^{\alpha} |v|^{\beta} dx < 0,$$

by the continuity of function $\Phi'_{\lambda,\mu}$ and ξ^- , we have

$$\begin{split} \langle \Phi_{\lambda,\mu}'(\xi^{-}(m)(z-m)), \xi^{-}(m)(z-m) \rangle \\ &= (p-q) \|\xi^{-}(m)(z-m)\|^{2} - 2(\alpha+\beta-q) \int_{\Omega} h |(\xi^{-}(m)(z-m))_{1}|^{\alpha} |(\xi^{-}(m)(z-m))_{2}|^{\beta} dx < 0. \end{split}$$

This implies that $\xi^{-}(m)(z-m) \in N^{-}_{\lambda,\mu}$.

Lemma 4.4. The following facts hold:

(i) If $(\lambda, \mu) \in \Theta$, then there is a $(PS)_{\theta_{\lambda,\mu}}$ -sequence $\{z_n\} = \{(u_n, v_n)\} \subset N_{\lambda,\mu}$ for $I_{\lambda,\mu}$. (ii) If $(\lambda, \mu) \in \Psi$, then there is a $(PS)_{\theta_{\lambda,\mu}^-}$ -sequence $\{z_n\} = \{(u_n, v_n)\} \subset N_{\lambda,\mu}^-$ for $I_{\lambda,\mu}$.

Proof. (1) By Lemma 3.4(iii) and Ekeland Variational Principle [12], there exists a minimizing sequence $\{z_n\} \subset N_{\lambda,\mu}$ such that

$$I_{\lambda,\mu}(z_n) < \theta_{\lambda,\mu} + \frac{1}{n},$$

$$I_{\lambda,\mu}(z_n) < I_{\lambda,\mu}(w) + \frac{1}{n} ||w - z_n||, \ \forall \ w \in N_{\lambda,\mu}.$$

$$(4.2)$$

By taking n large, from Lemma 3.4(i), we have $\theta_{\lambda,\mu} < 0$, thus

$$I_{\lambda,\mu}(z_n) = (\frac{1}{p} - \frac{1}{\alpha + \beta}) \|(u_n, v_n)\|^p - (\frac{1}{q} - \frac{1}{\alpha + \beta}) \int_{\Omega} (\lambda f |u_n|^q + \mu g |v_n|^q) dx \qquad (4.3)$$
$$< \theta_{\lambda,\mu} + \frac{1}{n} < \frac{\theta_{\lambda,\mu}}{2}.$$

This implies

$$-\frac{q(\alpha+\beta)}{2(\alpha+\beta-q)}\theta_{\lambda,\mu} < \int_{\Omega} (\lambda f |u_n|^q + \mu g |v_n|^q) \, dx$$

$$\leq [(|\lambda|||f||_{L^{q^\star}})^{\frac{p}{p-q}} + (|\mu|||g||_{L^{q^\star}})^{\frac{p}{p-q}}]^{\frac{p-q}{p}} S^{-\frac{q}{p}} ||(u_n, v_n)||^q.$$
(4.4)

Consequently, $z_n \neq 0$ and putting together (4.3), (4.4) and the Hölder inequality, we obtain

$$\|(u_n, v_n)\| > \left[-\frac{q(\alpha + \beta)}{2(\alpha + \beta - q)} \theta_{\lambda, \mu} \left[(|\lambda| \|f\|_{L^{q^{\star}}})^{\frac{p}{p-q}} + (|\mu| \|g\|_{L^{q^{\star}}})^{\frac{p}{p-q}} \right]^{\frac{q-p}{p}} S^{\frac{q}{p}} \right]^{\frac{1}{q}}.$$

and

$$\|(u_n, v_n)\| < \left[\frac{p(\alpha + \beta - q)}{q(\alpha + \beta - 2)} \left[(|\lambda| \|f\|_{L^{q^*}})^{\frac{p}{p-q}} + (|\mu| \|g\|_{L^{q^*}})^{\frac{p}{p-q}} \right]^{\frac{p}{p-q}} S^{-\frac{q}{p}} \right]^{\frac{1}{p-q}}.$$
 (4.5)

Now, we will show that

$$||I'_{\lambda,\mu}(z_n)||_{E^{-1}} \to 0 \quad as \quad n \to +\infty.$$

Applying Lemma 4.2 to z_n , we can obtain the function $\xi_n : B(0, r_n) \subset E \to \mathbb{R}^+$ such that $\xi_n(0) = 1$ and $\xi_n(m)(z_n - m) \in N_{\lambda,\mu}$ for every $m \in B(0, r_n)$. Taking $0 < \rho < r_n$, let $w \in E$ with $w \neq 0$ and put $m^* = \frac{\rho w}{\|w\|}$. We set $m_\rho = \xi_n(m^*)(z_n - m^*)$, then $m_\rho \in N_{\lambda,\mu}$. By (4.2), we have

$$I_{\lambda,\mu}(m_{\rho}) - I_{\lambda,\mu}(z_n) \ge -\frac{1}{n} ||m_{\rho} - z_n||.$$

By the Mean Value Theorem, we get

$$\langle I'_{\lambda,\mu}(z_n), m_{\rho} - z_n \rangle + o(||m_{\rho} - z_n||) \ge -\frac{1}{n} ||m_{\rho} - z_n||.$$

Thus, we have

$$\langle I'_{\lambda,\mu}(z_n), -m^* \rangle + (\xi_n(m^*) - 1) \langle I'_{\lambda,\mu}(z_n), z_n - m^* \rangle \ge -\frac{1}{n} \|m_\rho - z_n\| + o(\|m_\rho - z_n\|).$$
(4.6)

From $\xi_n(m^*)(z_n - m^*) \in N_{\lambda,\mu}$ and (4.6), we obtain

$$-\rho \langle I'_{\lambda,\mu}(z_n), \frac{w}{\|w\|} \rangle + (\xi_n(m^*) - 1) \langle I'_{\lambda,\mu}(z_n) - I'_{\lambda,\mu}(m_\rho), z_n - m^* \rangle \ge -\frac{1}{n} \|m_\rho - z_n\| + o(\|m_\rho - z_n\|).$$

So, we get

$$\langle I'_{\lambda,\mu}(z_n), \frac{w}{\|w\|} \rangle \leq \frac{\|m_{\rho} - z_n\|}{n\rho} + \frac{o(\|m_{\rho} - z_n\|)}{\rho} + \frac{(\xi_n(m^{\star}) - 1)}{\rho} \langle I'_{\lambda,\mu}(z_n) - I'_{\lambda,\mu}(m_{\rho}), z_n - m^{\star} \rangle.$$

$$(4.7)$$

Since

$$||m_{\rho} - z_n|| \le \rho |\xi_n(m^*)| + |\xi_n(m^*) - 1|||(u_n, v_n)||$$

and

$$\lim_{\rho \to 0} \frac{|\xi_n(m^*) - 1|}{\rho} \le \|\xi'_n(0)\|.$$

If we let $\rho \to 0$ in (4.7) for fixed $n \in \mathbb{N}$, then by (4.5) we can find a constant C > 0, independent of ρ such that

$$\langle I'_{\lambda,\mu}(z_n), \frac{w}{\|w\|} \rangle \leq \frac{C}{n} \left(1 + \|\xi'_n(0)\| \right).$$

Thus, we are done once we show that $\|\xi'_n(0)\|$ is uniformly bounded. By (4.1), (4.5) and Hölder inequality, we have

$$|\langle \xi'_n(0), m \rangle| \le \frac{C_1 ||m||}{|(p-q)||(u_n, v_n)||^p - 2(\alpha + \beta - q) \int_{\Omega} h|u_n|^{\alpha}|v_n|^{\beta} dx|},$$

for some $C_1 > 0$. We only need to show that

$$\left| (p-q) \| (u_n, v_n) \|^p - 2(\alpha + \beta - q) \int_{\Omega} h |u_n|^{\alpha} |v_n|^{\beta} dx \right| \ge C_2,$$

for some $C_2 > 0$ and n large enough. We argue by contradiction. Assume that there exists a subsequence z_n such that

$$(p-q)\|(u_n, v_n)\|^p - 2(\alpha + \beta - q) \int_{\Omega} h|u_n|^{\alpha}|v_n|^{\beta} dx = o_n(1).$$
(4.8)

By (4.8) and the fact that $z_n \in N_{\lambda,\mu}$, we have

$$||(u_n, v_n)||^p = \frac{2(\alpha + \beta - q)}{p - q} \int_{\Omega} h|u_n|^{\alpha}|v_n|^{\beta} dx + o_n(1)$$

and

$$\|(u_n, v_n)\|^p = \frac{\alpha + \beta - q}{\alpha + \beta - 2} \int_{\Omega} \left(\lambda f |u_n|^q + \mu g |v_n|^q\right) dx + o_n(1).$$

When n is large enough, by the Hölder and Sobolev inequalities, we have

$$\|(u_n, v_n)\| \ge \left[\frac{p-q}{2(\alpha+\beta-q)}S^{\frac{\alpha+\beta}{p}}\right]^{\frac{1}{\alpha+\beta-p}}$$

$$\tag{4.9}$$

and

$$\begin{aligned} \frac{\alpha + \beta - p}{\alpha + \beta - q} \|(u_n, v_n)\|^p &= \int_{\Omega} \left(\lambda f |u_n|^q + \mu g |v_n|^q\right) dx \\ &\leq |\lambda| \|f\|_{L^{q^\star}} \||u_n\|_{L^{\alpha+\beta}}^q + |\mu| \|g\|_{L^{q^\star}} \||v_n\|_{L^{\alpha+\beta}}^q \\ &\leq \left[(|\lambda| \|f\|_{L^{q^\star}})^{\frac{p}{p-q}} + (|\mu| \|g\|_{L^{q^\star}})^{\frac{p}{p-q}} \right]^{\frac{p-q}{p}} S^{-\frac{q}{p}} \|(u_n, v_n)\|^q. \end{aligned}$$

This implies

$$\|(u_n, v_n)\| \le \left(S^{-\frac{q}{p}} \frac{\alpha + \beta - q}{\alpha + \beta - p}\right)^{\frac{1}{p-q}} \left[(|\lambda| \|f\|_{L^{q^{\star}}})^{\frac{p}{p-q}} + (|\mu| \|g\|_{L^{q^{\star}}})^{\frac{p}{p-q}} \right]^{\frac{1}{p}}.$$
 (4.10)

By (4.9) and (4.10), we have

$$[(|\lambda| \|f\|_{L^{q^{\star}}})^{\frac{p}{p-q}} + (|\mu| \|g\|_{L^{q^{\star}}})^{\frac{p}{p-q}}] \ge \left[\frac{p-q}{2(\alpha+\beta-q)}S^{\frac{\alpha+\beta}{p}}\right]^{\frac{p}{\alpha+\beta-p}} \left(S^{-\frac{q}{p}}\frac{\alpha+\beta-q}{\alpha+\beta-p}\right)^{-\frac{p}{p-q}},$$

contradicting the assumption, that is

$$\langle I'_{\lambda,\mu}(z_n), \frac{w}{\|w\|} \rangle \leq \frac{C}{n}.$$

This completes the proof of (1).

Similarly, by Lemma 4.3, we can prove (ii), we omit the details here.

Lemma 4.5. If $\{z_n\} \subset E$ is a $(PS)_c$ -sequence for $I_{\lambda,\mu}$, then $\{z_n\}$ is bounded in E.

Proof. Let $z_n = (u_n, v_n) \subset E$ be a $(PS)_c$ -sequence for $I_{\lambda,\mu}$, suppose by contradiction that $||(u_n, v_n)|| \to +\infty$ as $n \to +\infty$. Let

$$\widetilde{z_n} = (\widetilde{u}_n, \widetilde{v}_n) := \frac{z_n}{\|(u_n, v_n)\|} = (\frac{u_n}{\|(u_n, v_n)\|}, \frac{v_n}{\|(u_n, v_n)\|}).$$

We may assume that $\tilde{z}_n \to \tilde{z} = (\tilde{u}, \tilde{v})$ in E. By the compact embedding theorem, we know $\tilde{u}_n(\cdot, 0) \to \tilde{u}(\cdot, 0)$ and $\tilde{v}_n(\cdot, 0) \to \tilde{v}(\cdot, 0)$ strongly in $L^r(\Omega)$ for all $1 \leq r < 2^*$. Thus, by Hölder inequality and Dominated convergence theorem, we have

$$\int_{\Omega} \left(\lambda f |\widetilde{u}_n|^q + \mu g |\widetilde{v}_n|^q\right) dx = \int_{\Omega} \left(\lambda f |\widetilde{u}|^q + \mu g |\widetilde{v}|^q\right) dx + o_n(1).$$

Since $\{z_n\}$ is a $(PS)_c$ -sequence for $I_{\lambda,\mu}$ and $||(u_n, v_n)|| \to +\infty$, we have

$$\frac{1}{p} \int_{Q} \frac{|\widetilde{u}_{n}(x) - \widetilde{u}_{n}(y)|^{p}}{|x - y|^{n + ps}} dx dy + \frac{1}{p} \int_{Q} \frac{|\widetilde{v}_{n}(x) - \widetilde{v}_{n}(y)|^{p}}{|x - y|^{n + ps}} dx dy$$

$$- \frac{\|(u_{n}, v_{n})\|^{q - p}}{q} \int_{\Omega} (\lambda f(x)|\widetilde{u}_{n}|^{q} + \mu g(x)|\widetilde{v}_{n}|^{q}) dx$$

$$- \frac{2\|(u_{n}, v_{n})\|^{\alpha + \beta - p}}{\alpha + \beta} \int_{\Omega} h(x)|\widetilde{u}_{n}|^{\alpha}|\widetilde{v}_{n}|^{\beta} dx = o_{n}(1)$$
(4.11)

and

$$\int_{Q} \frac{|\widetilde{u}_{n}(x) - \widetilde{u}_{n}(y)|^{p}}{|x - y|^{n + ps}} dx dy + \int_{Q} \frac{|\widetilde{v}_{n}(x) - \widetilde{v}_{n}(y)|^{p}}{|x - y|^{n + ps}} dx dy$$

$$- \|(u_{n}, v_{n})\|^{q - p} \int_{\Omega} (\lambda f(x)|\widetilde{u}_{n}|^{q} + \mu g(x)|\widetilde{v}_{n}|^{q}) dx$$

$$- 2\|(u_{n}, v_{n})\|^{\alpha + \beta - p} \int_{\Omega} h(x)|\widetilde{u}_{n}|^{\alpha}|\widetilde{v}_{n}|^{\beta} dx = o_{n}(1)$$
(4.12)

Combining (4.11) and (4.12), as $n \to \infty$, we obtain

$$\int_{Q} \frac{|\widetilde{u}_{n}(x) - \widetilde{u}_{n}(y)|^{p}}{|x - y|^{n + ps}} dx dy + \int_{Q} \frac{|\widetilde{v}_{n}(x) - \widetilde{v}_{n}(y)|^{p}}{|x - y|^{n + ps}} dx dy$$

$$= \frac{p(\alpha + \beta - q)}{q(\alpha + \beta - p)} \|(u_{n}, v_{n})\|^{q - p} \int_{\Omega} (\lambda f(x)|\widetilde{u}_{n}|^{q} + \mu g(x)|\widetilde{v}_{n}|^{q}) dx + o_{n}(1).$$
(4.13)

Since 1 < q < p and $||(u_n, v_n)|| \to +\infty$ as $n \to +\infty$, (4.13) implies

$$\int_Q \frac{|\widetilde{u}_n(x) - \widetilde{u}_n(y)|^p}{|x - y|^{n + ps}} dx dy + \int_Q \frac{|\widetilde{v}_n(x) - \widetilde{v}_n(y)|^p}{|x - y|^{n + ps}} dx dy \to 0.$$

Which contradicts the fact that $\|\tilde{z}_n\| = 1$ for any $n \ge 1$.

5 Local minimization problem

Now, we establish the existence of a local minimum for $I_{\lambda,\mu}$ on $N^+_{\lambda,\mu}$.

Theorem 5.1. Let $(\lambda, \mu) \in \Theta$, then $I_{\lambda,\mu}$ has a local minimizer z^+ in $N^+_{\lambda,\mu}$ satisfying (i) $I_{\lambda,\mu}(z^+) = \theta_{\lambda,\mu} = \theta^+_{\lambda,\mu}$; (ii) z^+ is a nontrivial solution of (1.1).

Proof. By (i) of Lemma 4.4 there exists a minimizing sequence $\{z_n\} = \{(u_n, v_n)\}$ for $I_{\lambda,\mu}$ in $N_{\lambda,\mu}$ such that

$$I_{\lambda,\mu}(z_n) = \theta_{\lambda,\mu} + o_n(1) \text{ and } I'_{\lambda,\mu}(z_n) = o_n(1) \text{ in } E^{-1}.$$
 (5.1)

By Lemma 3.4, Lemma 4.5 and the compact imbedding theorem, we know there is a subsequence, still denoted by $\{z_n\}$ and $z^+ = (u^+, v^+) \in E$ such that

$$\begin{cases} u_n \rightharpoonup u^+, v_n \rightharpoonup v^+, & weakly \text{ in } X_0^s(\Omega), \\ u_n \rightarrow u^+, v_n \rightarrow v^+, & srongly \text{ in } L^r(\Omega) \text{ for all } 1 \le r < 2^*. \end{cases}$$

As $n \to \infty$, by Hölder inequality and Dominated convergence theorem, we obtain

$$\int_{\Omega} \left(\lambda f |u_n|^q + \mu g |v_n|^q\right) dx = \int_{\Omega} \left(\lambda f |u^+|^q + \mu g |v^+|^q\right) dx + o_n(1)$$
(5.2)

and

$$\int_{\Omega} h|u_n|^{\alpha}|v_n|^{\beta} dx = \int_{\Omega} h|u^+|^{\alpha}|v^+|^{\beta} dx + o_n(1).$$
(5.3)

First, we claim that $\int_{\Omega} (\lambda f |u^+|^q + \mu g |v^+|^q) dx \neq 0$, we argue by contradiction, then we have $\int_{\Omega} (\lambda f |u_n|^q + \mu g |v_n|^q) dx \to 0$ as $n \to \infty$. Thus

$$||(u_n, v_n)||^p = 2 \int_{\Omega} h|u_n|^{\alpha}|v_n|^{\beta} dx + o_n(1)$$

and

$$I_{\lambda,\mu}(z_n) = \frac{1}{p} \|(u_n, v_n)\|^p - \frac{2}{\alpha + \beta} \int_{\Omega} h(x) |u_n|^{\alpha} |v_n|^{\beta} dx + o_n(1)$$
$$= \left(\frac{1}{p} - \frac{1}{\alpha + \beta}\right) \|(u_n, v_n)\|^2 + o_n(1).$$

This contradicts $I_{\lambda,\mu}(z_n) \to \theta_{\lambda,\mu} < 0$ as $n \to \infty$.

Now, we claim z^+ is a nontrivial solution of (1.1). From (5.1), (5.2) and (5.3), we know z^+ is a weak solution of (1.1). From $z_n \in N_{\lambda,\mu}$, we have

$$I_{\lambda,\mu}(z_n) = \frac{\alpha + \beta - p}{p(\alpha + \beta)} \|(u_n, v_n)\|^p - \frac{\alpha + \beta - q}{q(\alpha + \beta)} \int_{\Omega} \left(\lambda f |u_n|^q + \mu g |v_n|^q\right) dx.$$
(5.4)

That is

$$\int_{\Omega} \left(\lambda f |u_n|^q + \mu g |v_n|^q\right) dx = \frac{q(\alpha + \beta - p)}{p(\alpha + \beta - q)} \|(u_n, v_n)\|^p - \frac{q(\alpha + \beta)}{\alpha + \beta - q} I_{\lambda,\mu}(z_n).$$
(5.5)

Let $n \to \infty$ in (5.5), by (5.1), (5.2) and $\theta_{\lambda,\mu} < 0$, we have

$$\int_{\Omega} \left(\lambda f |u^+|^q + \mu g |v^+|^q \right) dx \ge -\frac{q(\alpha + \beta)}{\alpha + \beta - q} \theta_{\lambda,\mu} > 0.$$

Therefore, $z^+ \in N_{\lambda,\mu}$ is a nontrival solution of (1.1). Next, we show that $z_n \to z^+$ strongly in E and $I_{\lambda,\mu}(z^+) = \theta_{\lambda,\mu}$. Since $z^+ \in N_{\lambda,\mu}$, then by (5.4), we obtain

$$\begin{aligned} \theta_{\lambda,\mu} &\leq I_{\lambda,\mu}(z^{+}) = \frac{\alpha + \beta - p}{p(\alpha + \beta)} \|(u^{+}, v^{+})\|^{p} - \frac{\alpha + \beta - q}{q(\alpha + \beta)} \int_{\Omega} \left(\lambda f |u^{+}|^{q} + \mu g |v^{+}|^{q}\right) dx \quad (5.6) \\ &\leq \liminf_{n \to \infty} \left(\frac{\alpha + \beta - p}{p(\alpha + \beta)} \|(u_{n}, v_{n})\|^{p} - \frac{\alpha + \beta - q}{q(\alpha + \beta)} \int_{\Omega} \left(\lambda f |u_{n}|^{q} + \mu g |v_{n}|^{q}\right) dx\right) \\ &\leq \lim_{n \to \infty} \left(\frac{\alpha + \beta - p}{p(\alpha + \beta)} \|(u_{n}, v_{n})\|^{p} - \frac{\alpha + \beta - q}{q(\alpha + \beta)} \int_{\Omega} \left(\lambda f |u_{n}|^{q} + \mu g |v_{n}|^{q}\right) dx\right) \\ &\leq \lim_{n \to \infty} I_{\lambda,\mu}(z_{n}) = \theta_{\lambda,\mu}. \end{aligned}$$

This inplies that $I_{\lambda,\mu}(z^+) = \theta_{\lambda,\mu}$ and $\lim_{n \to \infty} ||(u_n, v_n)||^p = ||(u^+, v^+)||^p$. Hence $z_n \to z^+$ srongly in E.

Finially, we claim that $z^+ \in N^+_{\lambda,\mu}$. Assume by contradiction that $z^+ \in N^-_{\lambda,\mu}$, then by Lemma 3.5, there exist unique t_1^+ and t_1^- , such that $t_1^+(z^+) \in N^+_{\lambda,\mu}$, $t_1^-(z^+) \in N^-_{\lambda,\mu}$. In particular, we have $t_1^+ < t_1^- = 1$. Since

$$\frac{d}{dt}I_{\lambda,\mu}(t_1^+z^+) = 0 \text{ and } \frac{d^2}{dt^2}I_{\lambda,\mu}(t_1^+z^+) > 0,$$

there exists $t_1^+ < t^* < t_1^-$ such that $I_{\lambda,\mu}(t_1^+z^+) < I_{\lambda,\mu}(t^*z^+)$. By Lemma 3.5, we have

$$I_{\lambda,\mu}(t_1^+z^+) < I_{\lambda,\mu}(t^*z^+) \le I_{\lambda,\mu}(t_1^-z^+) = I_{\lambda,\mu}(z^+),$$

a contraction. Since $I_{\lambda,\mu}(z^+) = I_{\lambda,\mu}(|u^+|, |v^+|)$ and $(|u^+|, |v^+|) \in N_{\lambda,\mu}$, by Lemma 3.2, we obtain that z^+ is a nontrivial solution of (1.1).

Next, we establish the existence of a local minimum for $I_{\lambda,\mu}$ on $N_{\lambda,\mu}^-$.

Theorem 5.2. Let $(\lambda, \mu) \in \Psi$, then $I_{\lambda,\mu}$ has a local minimizer z^- in $N^-_{\lambda,\mu}$ satisfying (i) $I_{\lambda,\mu}(z^-) = \theta^-_{\lambda,\mu}$; (ii) z^- is a nontrivial solution of (1.1).

Proof. By (*ii*) of Lemma 4.4 there exists a minimizing sequence $\{z_n\} = \{(u_n, v_n)\}$ for $I_{\lambda,\mu}$ in $N_{\lambda,\mu}^-$ such that

$$I_{\lambda,\mu}(z_n) = \theta_{\lambda,\mu}^- + o_n(1) \text{ and } I'_{\lambda,\mu}(z_n) = o_n(1) \text{ in } E^{-1}.$$

By Lemma 3.4 (*iii*), Lemma 4.5 and the compact imbedding theorem, we know there is a subsequence, still denoted by $\{z_n\}$ and $z^- = (u^-, v^-) \in N^-_{\lambda,\mu}$ such that

$$\begin{cases} u_n \rightharpoonup u^-, v_n \rightharpoonup v^-, & weakly \text{ in } X_0^s(\Omega), \\ u_n \rightarrow u^-, v_n \rightarrow v^-, & srongly \text{ in } L^r(\Omega) \text{ for all } 1 \le r < 2^\star. \end{cases}$$

As $n \to \infty$, by Hölder inequality and Dominated convergence theorem, we obtain

$$\int_{\Omega} \left(\lambda f |u_n|^q + \mu g |v_n|^q\right) dx = \int_{\Omega} \left(\lambda f |u^-|^q + \mu g |v^-|^q\right) dx + o_n(1)$$

and

$$\int_{\Omega} h|u_n|^{\alpha}|v_n|^{\beta}dx = \int_{\Omega} h|u^-|^{\alpha}|v^-|^{\beta}dx + o_n(1).$$

First, we claim that $\int_{\Omega} (\lambda f |u^-|^q + \mu g |v^-|^q) dx \neq 0$, suppose by contradiction, then we have $\int_{\Omega} (\lambda f |u_n|^q + \mu g |v_n|^q) dx \to 0$ as $n \to \infty$. Thus

$$|(u_n, v_n)||^p = 2 \int_{\Omega} h |u_n|^{\alpha} |w_{2,n}|^{\beta} dx + o_n(1)$$

and

$$\begin{split} I_{\lambda,\mu}(z_n) &= \frac{1}{p} \|(u_n, v_n)\|^2 - \frac{2}{\alpha + \beta} \int_{\Omega} h(x) |u_n|^{\alpha} |v_n|^{\beta} dx + o_n(1) \\ &= \left(\frac{1}{p} - \frac{1}{\alpha + \beta}\right) \|(u_n, v_n)\|^2 + o_n(1). \end{split}$$

This contradicts $I_{\lambda,\mu}(z_n) \to \theta_{\lambda,\mu} < 0$ as $n \to \infty$.

Now, we prove that $z_n \to z^-$ strongly in *E*. Othercase, we have

$$\begin{split} \|(u^{-}, v^{-})\|^{p} &- \int_{\Omega} \left(\lambda f |u^{-}|^{q} + \mu g |v^{-}|^{q} \right) dx - 2 \int_{\Omega} h |u^{-}|^{\alpha} |v^{-}|^{\beta} dx \\ &\leq \liminf_{n \to \infty} \left(\|(u_{n}, v_{n})\|^{p} - \int_{\Omega} \left(\lambda f |u_{n}|^{q} + \mu g |v_{n}|^{q} \right) dx - 2 \int_{\Omega} h |u_{n}|^{\alpha} |v_{n}|^{\beta} dx \right) \\ &\leq \lim_{n \to \infty} \left(\|(u_{n}, v_{n})\|^{p} - \int_{\Omega} \left(\lambda f |u_{n}|^{q} + \mu g |v_{n}|^{q} \right) dx - 2 \int_{\Omega} h |u_{n}|^{\alpha} |v_{n}|^{\beta} dx \right) = 0 \end{split}$$

Which contradicts $z^- \in N^-_{\lambda,\mu}$. Hence $z_n \to z^-$ strongly in E. This implies

$$I_{\lambda,\mu}(z_n) \to I_{\lambda,\mu}(z^-) = \theta^-_{\lambda,\mu} \text{ as } n \to +\infty.$$

Since $I_{\lambda,\mu}(z^-) = I_{\lambda,\mu}(|u^-, v^-|)$ and $|u^-, v^-| \in N^-_{\lambda,\mu}$, by Lemma 3.2, we have z^- is a nontrivial solution of (1.1).

6 Proof of Theorem 1.1 and Theorem 1.2

Now, we complete the proof of Theorem 1.1 and Theorem 1.2.

Proof. For $(\lambda, \mu) \in \Theta$, by Theorem 5.1, system (1.1) admits at least one nontrivial solution $z^+ \in N^+_{\lambda,\mu}$. By Theorem 5.1 and Theorem 5.2, we obtain that for $(\lambda, \mu) \in \Psi$, system (1.1) admits at least two nontrivial solutions z^+ and z^- such that $z^+ \in N^+_{\lambda,\mu}$, $z^- \in N^-_{\lambda,\mu}$. Since $N^+_{\lambda,\mu} \cap N^+_{\lambda,\mu} = \emptyset$, then z^+ and z^- are distinct solutions of system (1.1).

References

- C.O. Alves, D.C. de Morais Filho and M.A.S. Souto, On systems of elliptic equations involving subcritical or critical Sobolev exponents, *Nonlinear Anal.*, 42 (2000), no. 5, 771-787.
- [2] B. Barrios, E. Colorado, A. de Pablo and U. Sánchez, On some critical problems for the fractional Laplacian operator, J. Differential Equations, 252 (2012), no. 11, 6133-6162.
- [3] C. Brändle, E. Colorado, A. de Pablo and U. Sánchez, A concave-convex elliptic problem involving the fractional Laplacian, *Proc. Roy. Soc. Edinburgh Sect. A*, 143 (2013), no. 1, 39-71.
- [4] X. Cabré, Y. Sire, Nonlinear equations for fractional Laplacians, I: Regularity, maximum principles, and Hamiltonian estimates, Ann. Inst. H. Poincar Anal. Non Linaire, 31 (2014), no. 1, 23-53.
- [5] L. Caffarelli, L. Silvestre, An extension problem related to the fractional Laplacian, Comm. Partial Differential Equations, 32 (2007), no. 7-9, 1245-1260.
- [6] L. Caffarelli, J. Roquejoffre and Y. Sire, Variational problems with free boundaries for the fractional Laplacian, J. Eur. Math. Soc., 12 (2010), no. 5, 1151-1179.
- [7] A. Capella, J. Dávila, L. Dupaigne and Y. Sire, Regularity of radial extremal solutions for some non-local semilinear equations, *Comm. Partial Differential Equations*, 36 (2011), no. 8, 1353–1384.
- [8] W. Chen, S. Deng, The Nehari manifold for a fractional p-Laplacian system involving concave-convex nonlinearities, *Nonlinear Anal. Real World Appl.*, **27** (2016), 80-92.
- [9] B. Cheng, X. Tang, New existence of solutions for the fractional p-Laplacian equations with sign-changing potential and nonlinearity, *Mediterr. J. Math.*, **13** (2016), no. 5, 3373-3387.
- [10] E. Colorado, A. de Pablo and U. Sánchez, Perturbations of a critical fractional equation, *Pacific J. Math.*, **271** (2014), no. 1, 65-85.

- [11] E. Di Nezza, G. Palatucci and E. Valdinoci, Hitchhikers guide to the fractional Sobolev spaces, Bull. Sci. Math., 136 (2012), no. 5, 521-573.
- [12] I. Ekeland, On the variational principle, J. Math. Anal. Appl., 47 (1974), no. 2, 324-353.
- [13] S. Goyal, K. Sreenadh, Nehari manifold for non-local elliptic operator with concaveconvex nonlinearities and sign-changing weight functions, *Proceedings - Mathematical Sciences*, **125** (2015), no. 4, 545-558.
- [14] X. He, M. Squassina and W. Zou, The Nehari manifold for fractional systems involving critical nonlinearities, *Commun. Pure Appl. Anal.*, 15 (2016), no. 4, 1285-1308.
- [15] J. Marcos, D. Ferraz, Concentration-compactness principle for nonlocal scalar field equations with critical growth, J. Math. Anal. Appl., 449 (2017), no. 2, 1189-1228.
- [16] A. Mellet, S. Mischler and C. Mouhot, Fractional diffusion limit for collisional kinetic equations, Arch. Ration. Mech. Anal., 199 (2011), no. 2, 493-525.
- [17] X. Ros-Oton, Nonlocal elliptic equations in bounded domains: a survey, *Publ. Mat.*, 60 (2016), no. 1, 3-26.
- [18] R. Servadei, E. Valdinoci, Mountain Pass solutions for non-local elliptic operators, J. Math. Anal. Appl., 389 (2012), no. 2, 887-898.
- [19] R. Servadei, E. Valdinoci, Variational methods for non-local operators of elliptic type, Discrete Contin. Dyn. Syst., 33 (2013), no. 5, 2105-2137.
- [20] R. Servadei, E. Valdinoci, The Brezis-Nirenberg result for the fractional Laplacian, *Trans. Amer. Math. Soc.*, 367 (2015), no. 1, 67-102.
- [21] R. Servadei, E. Valdinoci, A Brezis-Nirenberg result for non-local critical equations in low dimension, *Commun. Pure Appl. Anal.*, **12** (2013), no. 6, 2445-2464.
- [22] L. Silvestre, Regularity of the obstacle problem for a fractional power of the Laplace operator, Comm. Pure Appl. Math., 60 (2007), no. 1, 67-112.
- [23] G. Tarantello, On nonhomogeneous elliptic involving critical Sobolev exponent, Ann. Inst. H. Poincaré Anal. Non Linéaire, 9 (1992), no. 3, 281-304.
- [24] T.F. Wu, On semilinear elliptic equations involving concave-convex nonlinearities and sign-changing weight functions, J. Math. Anal. Appl., 318 (2006), no. 1, 253-270.
- [25] X. Zheng, J. Wang, Symmetry results for systems involving fractional Laplacian, Indian J. Pure Appl. Math., 45 (2014), no. 1, 39-51.