

PERSISTENCE BARCODES AND LAPLACE EIGENFUNCTIONS ON SURFACES

IOSIF POLTEROVICH¹, LEONID POLTEROVICH², AND VUKAŠIN STOJISAVLJEVIĆ²

ABSTRACT. We obtain restrictions on the persistence barcodes of Laplace-Beltrami eigenfunctions and their linear combinations on compact surfaces with Riemannian metrics. Some applications to uniform approximation by linear combinations of Laplace eigenfunctions are also discussed.

CONTENTS

1. Introduction and main results	1
1.1. Laplace-Beltrami eigenfunctions	1
1.2. Persistence modules and barcodes	3
1.3. A family of functionals on the space of barcodes	5
1.4. Main results	7
1.5. Sorting the finite bars of functions in \mathcal{F}_λ	9
2. Applications to approximations by eigenfunctions	10
2.1. An obstruction to C^0 -approximations	10
2.2. Modulus of continuity and average length of bars	12
3. Barcodes and the Banach indicatrix	15
3.1. A topological bound on the Banach indicatrix	15
3.2. Proof of Theorem 1.4.1	16
3.3. Proof of Proposition 3.1.1	17
4. Miscellaneous proofs	20
4.1. Proof of Lemma 1.3.1	20
4.2. Proof of Lemma 1.3.5	20
4.3. Proof of Proposition 1.4.10	21
Acknowledgments	22
References	22

1. INTRODUCTION AND MAIN RESULTS

1.1. Laplace-Beltrami eigenfunctions. The past fifteen years have witnessed a number of fascinating applications of the spectral theory of the Laplace-Beltrami operator to data analysis, such as dimensionality reduction and data representation [BN, CL] or shape segmentation in computer graphics [SOCG, Re]. In the present paper we focus on this interaction the other way around and study persistence barcodes, a fundamental notion originated in topological data analysis, of the

¹Partially supported by NSERC, FRQNT and Canada Research Chairs program.

²Partially supported by the European Research Council Advanced grant 338809.

Laplace-Beltrami eigenfunctions and their linear combinations. Our main finding is a constraint on such barcodes in terms of the corresponding eigenvalues. This result turns out to have applications to approximation theory.

Let M be a compact n -dimensional Riemannian manifold, possibly with nonempty boundary. Let Δ be the (positive definite) Laplace-Beltrami operator on M ; if $\partial M \neq \emptyset$ we assume that the Dirichlet condition is imposed on the boundary. The spectrum of the Laplace-Beltrami operator on a compact Riemannian manifold is discrete, and the eigenvalues form a sequence $0 \leq \lambda_1 \leq \lambda_2 \leq \dots \nearrow \infty$, where each eigenvalue is repeated according to its multiplicity. The corresponding eigenfunctions f_k , $\Delta f_k = \lambda_k f_k$, form an orthonormal basis in $L^2(M)$. The properties of Laplace-Beltrami eigenfunctions have fascinated researchers for more than two centuries, starting with the celebrated Chladni's experiments with vibrating plates. We refer to [JNT, Z1, Z2] for a modern overview of the subject. As the examples of trigonometric polynomials and spherical harmonics indicate, the shapes of the eigenfunctions are expected to have an increasingly complex structure as λ goes to infinity. At the same time, various restrictions on the behaviour of eigenfunctions can be formulated in terms of the corresponding eigenvalue. One of the basic facts about eigenfunctions is Courant's nodal domain theorem, stating that the number of nodal domains of an eigenfunction f_k is at most k (see [CH]). There exist also bounds on the $(n-1)$ -dimensional measure of the zero set of eigenfunctions (see [L1, L2, LM] for most recent developments on this topic), on the distribution of nodal extrema ([PS, Po]), on the growth of L^p norms ([So]), and other related results.

In the present paper we focus on topological properties of the sublevel sets of Laplace-Beltrami eigenfunctions, and, more generally, of the linear combinations of eigenfunctions with eigenvalues $\leq \lambda$. There has been a number of important recent advances in the study of topological properties of random linear combinations of Laplace eigenfunctions, with an emphasis on the nodal and critical sets (see, for instance, [NS, Ni, GW1, GW2, SW, CS]). Our approach is deterministic and is based on the study of *persistence barcodes*. In the probabilistic setting, some steps in this direction have been discussed in [CMW, Section 1.4.3]. Roughly speaking, a persistence barcode is a collection of intervals in \mathbb{R} which encodes oscillation of a function (see next subsection for a detailed overview). Our main result (Theorem 1.4.1) implies that the quantity $\Phi_1(f)$, the total length of the barcode of any such linear combination f with unit L^2 -norm, satisfies an upper bound $O(\lambda)$. This inequality is inspired by the ideas introduced in [PS], where a similar bound was proved for the *Banach indicatrix* of f , another measure of oscillation which goes back to the works of Kronrod [Kr] and Yomdin [Yo]. Our central observation (see Proposition 3.1.1 below) is that the length of the barcode admits an upper bound via the Banach indicatrix, which together with [PS] yields the main result.

We believe that discussing eigenfunctions and their linear combinations in the language of barcodes, which originated in topological data analysis, has a number of merits. First, there exists a well developed metric theory of barcodes which highlights their robustness with respect to perturbations of functions in the uniform norm. Some features of this robustness are inherited by the above-mentioned functional Φ_1 . This, in turn, paves the way for applications to the following question of approximation theory (see Section 2): given a function with unit L^2 -norm, how well one can approximate it by a linear combination of Laplace eigenfunctions

with eigenvalues $\leq \lambda$, possibly after composition with a diffeomorphism of the surface? Second, we conjecture that barcodes provide a right framework for a potential extension of our results to higher dimensions (see Conjecture 1.4.8 below).

In a different direction, we present an application to the problem of sorting finite bars of persistence barcodes. This task arises on a number of occasions in topology and data analysis. Our results allow to improve an estimate on the optimal running time of a sorting algorithm for barcodes of linear combinations of Laplace eigenfunctions with eigenvalues $\leq \lambda$, see subsection 1.5.

1.2. Persistence modules and barcodes. In order to describe the topology of the sublevel sets of eigenfunctions and their linear combinations, we use the notions of persistence modules and barcodes, which are briefly reviewed below. We refer the reader to articles [EH, G, C, W, BL] and monographs [E, O, CSGO] for an introduction into this rapidly developing subject and further details.

Let M be a closed, connected, orientable, n -dimensional manifold, possibly with boundary and $f : M \rightarrow \mathbb{R}$ a Morse function (if $\partial M \neq \emptyset$, we assume that f is Morse on $M \setminus \partial M$ and $f|_{\partial M} = \text{const}$, the value on the boundary being regular). For $k = 0, \dots, n$ define a family of finite dimensional real vector spaces depending on a parameter $t \in \mathbb{R}$ by setting

$$V_k^t(f) = H_k(f^{-1}((-\infty, t)); \mathbb{R}).$$

If $s \leq t$ we have that $f^{-1}((-\infty, s)) \subset f^{-1}((-\infty, t))$ and the inclusion of sublevel sets $i_{st} : f^{-1}((-\infty, s)) \rightarrow f^{-1}((-\infty, t))$ induces a map

$$\pi_{st} = (i_{st})_* : V_k^s(f) \rightarrow V_k^t(f) \text{ for } k = 0, \dots, n.$$

These maps are called *comparison maps*. The family of vector spaces $V_k^t(f)$ together with the family of comparison maps π_{st} forms an algebraic structure called a *persistence module*. We call this persistence module a degree k persistence module associated to f . We define the *spectrum* of V_k^t as those points $r \in \mathbb{R}$ for which V_k^t "changes when t passes through r ". More formally, if r has a neighbourhood U_r such that π_{st} are isomorphisms for all $s, t \in U_r$, we say that r is not inside spectrum and we say that r is inside spectrum otherwise. One may readily check that under the above assumptions spectrum is finite and consists of critical values of f (and possibly also the value of f on the boundary of M if $\partial M \neq \emptyset$). Moreover, if $a < b$ are consecutive points of the spectrum, then π_{st} is an isomorphism for all $a < s < t \leq b$.

Persistence modules that we will consider will be associated to a certain function $f : M \rightarrow \mathbb{R}$ as described above. However, one may define a persistence module as an abstract algebraic structure, without the auxiliary function f . In this setting, a persistence module consists of a one-parametric family of finite dimensional³ vector spaces V^t , $t \in \mathbb{R}$ and a family of linear comparison maps $\pi_{st} : V^s \rightarrow V^t$ for $s \leq t$, which satisfy $\pi_{tt} = \mathbf{1}_{V^t}$ and $\pi_{st} \circ \pi_{rs} = \pi_{rt}$ for all real numbers $r \leq s \leq t$. Some of the results that we will refer to, namely the *structure theorem* and the *isometry theorem*, may be formulated and proven in this abstract language.

According to the structure theorem for persistence modules, V^t decomposes into a direct sum of simple persistence modules. Let $I = (a, b]$ or $I = (a, +\infty)$, $a, b \in \mathbb{R}$ and denote by $Q(I) = (Q(I), \pi)$ the persistence module which satisfies $Q^t(I) = \mathbb{R}$

³These are sometimes referred to as *pointwise finite dimensional* persistence modules.




FIGURE 1. Height function on a deformed circle.

for $t \in I$ and $Q^t(I) = 0$ otherwise and $\pi_{st} = id$ for $s, t \in I$ and $\pi_{st} = 0$ otherwise. Now

$$V^t \cong \bigoplus_i (Q^t(I_i))^{m_i},$$

where m_i are finite multiplicities, I_i are intervals of the form $(a_i, b_i]$ or $(a_i, +\infty)$ and $(Q(I), \pi) \oplus (Q(I'), \pi') = (Q(I) \oplus Q(I'), \pi \oplus \pi')$. The above decomposition is unique if we assume that $I_i \neq I_j$ when $i \neq j$. The multiset containing m_i copies of I_i is called the *barcode* of V^t and is denoted by $\mathcal{B}(V^t)$ and intervals I_i are called *bars*. In the case of a module $V^t = V_k^t(f)$ coming from a function f , we have that endpoints a_i, b_i of all of bars belong to the spectrum of V_k^t and the barcode $\mathcal{B}(V_k^t) = \mathcal{B}_k(f)$ is called the *degree k barcode* of f . We also denote by $\mathcal{B}(f) = \cup_k \mathcal{B}_k(f)$ the *full barcode* of f . Under our assumptions $\mathcal{B}(f)$ is finite, i.e. it consists of finitely many distinct intervals with finite multiplicities.

Example 1.2.1. Let $f : \mathbb{S}^1 \rightarrow \mathbb{R}$ be a height function on a deformed circle (see Figure 1). Critical values of f are a, b, c and d , and for a regular value $t \in \mathbb{R}$, the sublevel sets $f^{-1}((-\infty, t))$ are homeomorphic to:

$$f^{-1}((-\infty, t)) = \begin{cases} \emptyset, & \text{for } t < a \\ I, & \text{for } a < t < b \\ I \sqcup I, & \text{for } b < t < c \\ I, & \text{for } c < t < d \\ \mathbb{S}^1, & \text{for } d < t \end{cases}$$

where I stands for an open interval. Degree 1 barcode is now easily seen to contain one infinite bar $\mathcal{B}_1(f) = \{(d, +\infty)\}$, while degree 0 barcode contains one infinite and one finite bar $\mathcal{B}_0(f) = \{(a, +\infty), (b, c]\}$. The finite bar $(b, c]$ corresponds to the

fact that for $b < t < c$, $f^{-1}((-\infty, t))$ has two connected components which merge for $t > c$. The full barcode is given by $\mathcal{B}(f) = \{(a, +\infty), (d, +\infty), (b, c]\}$.

Let us introduce some notation and give a short description of the barcode associated to a function $f : M \rightarrow \mathbb{R}$ under the above assumptions. Denote by $b_i = \dim H_i(M; \mathbb{R})$, $i = 0, \dots, n$ the Betti numbers of M . Bars of infinite lengths in degree k barcode correspond to homology classes in $H_k(M; \mathbb{R})$. This means that the number of infinite bars in degree k barcode is equal to b_k . Since M is connected and orientable, we know that $b_0 = 1$, while $b_n = 1$ if $\partial M = \emptyset$ and $b_n = 0$ otherwise. Thus, if $\partial M = \emptyset$, degree 0 and degree n barcodes each contain exactly one infinite bar $(c^{(0)}, +\infty)$ and $(c^{(n)}, +\infty)$, while degree k barcode contains b_k infinite bars $(c_1^{(k)}, +\infty), \dots, (c_{b_k}^{(k)}, +\infty)$, $c_1^{(k)} \leq \dots \leq c_{b_k}^{(k)}$ for $1 \leq k \leq n-1$. If $\partial M \neq \emptyset$, the barcode is similar, the only difference being that there are no infinite bars in degree n . Numbers $c_i^{(k)}$, $1 \leq i \leq b_k$ are called *spectral invariants* of a function f in degree k and it holds $c^{(0)} = \min f$. If $\partial M = \emptyset$, we also have that $c^{(n)} = \max f$. All finite bars are contained in $(\min f, \max f]$ and there are no finite bars in degree n .

The space of barcodes can be endowed with a natural distance. Let \mathcal{B} and $\tilde{\mathcal{B}}$ be two barcodes. We say that $\mu : U \rightarrow \tilde{U}$ is an ε -matching between \mathcal{B} and $\tilde{\mathcal{B}}$ if it is a bijection between some subsets $U \subset \mathcal{B}$ and $\tilde{U} \subset \tilde{\mathcal{B}}$ which contain all bars of length greater than 2ε and it satisfies

$$\mu((a, b]) = (c, d] \Rightarrow |a - c|, |b - d| \leq \varepsilon.$$

Intuitively, we may delete some bars of length less or equal than 2ε from \mathcal{B} and $\tilde{\mathcal{B}}$, and match the rest up to an error of ε . We may also think of the deleted bars as matched with intervals of length 0. Now, the *bottleneck distance* between \mathcal{B} and $\tilde{\mathcal{B}}$, $d_{bottle}(\mathcal{B}, \tilde{\mathcal{B}})$, is defined as infimum over all $\varepsilon \geq 0$ such that there exists an ε -matching between \mathcal{B} and $\tilde{\mathcal{B}}$. One of the direct consequences of the isometry theorem is the following stability result which we will need later. For two functions f and g it holds

$$(1) \quad d_{bottle}(\mathcal{B}(f), \mathcal{B}(g)) \leq |f - g|_{C^0},$$

where $|f - g|_{C^0} = \max_x |f(x) - g(x)|$.

Remark 1.2.2. Full barcodes $\mathcal{B}(f)$ and $\mathcal{B}(g)$ which we consider do not keep track of the degrees. In particular, this means that we allow matchings between $\mathcal{B}(f)$ and $\mathcal{B}(g)$ to match bars from $\mathcal{B}_i(f)$ with bars from $\mathcal{B}_j(g)$ for $i \neq j$. The isometry theorem can also be applied in each degree independently to obtain

$$\max_k d_{bottle}(\mathcal{B}_k(f), \mathcal{B}_k(g)) \leq |f - g|_{C^0}.$$

From here (1) immediately follows because

$$d_{bottle}(\mathcal{B}(f), \mathcal{B}(g)) \leq \max_k d_{bottle}(\mathcal{B}_k(f), \mathcal{B}_k(g)).$$

1.3. A family of functionals on the space of barcodes. From now on, we assume that M is an orientable surface, possibly with boundary. Let us define, for every positive function $u \in C(\mathbb{R})$, a positive, lower semi-continuous functional Φ_u on the space of Morse functions on M . Let f be a Morse function, denote by $\mathcal{B}'(f) \subset \mathcal{B}(f)$ the multiset of all finite bars in the barcode $\mathcal{B}(f)$ and by $|\mathcal{B}'(f)|$

the total number of finite bars in $\mathcal{B}(f)$. Define a positive functional Φ_u on the set \mathcal{F}_{Morse} of all Morse functions (vanishing on the boundary) by setting

$$(2) \quad \Phi_u(f) = \begin{cases} \int_{\min f}^{\max f} u(t) dt + \sum_{I \in \mathcal{B}'(f)} \int_I u(t) dt & \text{if } \partial M = \emptyset, \\ 0 & \text{if } \partial M \neq \emptyset. \end{cases}$$

In particular, $\Phi_1(f)$ is the sum of the lengths of all the finite bars in the barcode of f and the length of the range of f .

Lemma 1.3.1. *Let*

$$C(u, f) = 2 \cdot (|\mathcal{B}'(f)| + 1) \cdot \left(\max_{[\min f, \max f]} u \right)$$

in the case $\partial M = \emptyset$, or

$$C(u, f) = (2|\mathcal{B}'(f)| + 1) \cdot \left(\max_{[\min f, \max f]} u \right)$$

in the case $\partial M \neq \emptyset$. Then

$$(3) \quad \Phi_u(f) - \Phi_u(h) \leq C(u, f) \cdot d_{bottle}(\mathcal{B}(f), \mathcal{B}(h)).$$

We prove this lemma in subsection 4.1. Combining (3) with (1) yields

$$(4) \quad \Phi_u(f) - \Phi_u(h) \leq C(u, f) \cdot d_{C^0}(f, h),$$

where $d_{C^0}(f, h) = |f - h|_{C^0}$.

Proposition 1.3.2. *The functional Φ_u is lower semi-continuous both as a functional $\Phi_u : (\mathbf{B}, d_{bottle}) \rightarrow \mathbb{R}$ and as a functional $\Phi_u : (\mathcal{F}_{Morse}, d_{C^0}) \rightarrow \mathbb{R}$. Here \mathbf{B} stands for the set of all barcodes corresponding to functions in \mathcal{F}_{Morse} .*

Remark 1.3.3. We slightly abuse the notation here by looking at $\Phi_u(f) = \Phi_u(\mathcal{B}(f))$ as the function of barcode $\mathcal{B}(f)$. However, it is obvious that Φ_u depends only on $\mathcal{B}(f)$ and not on f itself. In the same spirit $\min f$ and $\max f$ should be replaced by the smallest and the largest endpoint of a bar in $\mathcal{B}(f)$.

Proof. Recall that a functional Φ defined on a metric space X is called lower semi-continuous at a point $f \in X$ if $\liminf_{h \rightarrow f} \Phi(h) \geq \Phi(f)$. This relation easily follows from the inequalities (3) and (4) for the functional Φ_u defined on the metric spaces (\mathbf{B}, d_{bottle}) and $(\mathcal{F}_{Morse}, d_{C^0})$, respectively. \square

The inequality (4) could be further strengthened. Let $\text{Diff}(M)$ denote the group of all smooth diffeomorphisms of the surface M (throughout the paper, the term ‘‘smooth’’ stands for C^∞ -smooth).

Corollary 1.3.4. *We have*

$$(5) \quad \Phi_u(f) - \Phi_u(h) \leq C(u, f) \cdot d_{C^0}(f \circ \varphi, h \circ \psi),$$

for any two diffeomorphisms $\varphi, \psi \in \text{Diff}(M)$.

In particular, taking $\varphi = \psi = \mathbf{1}_M$ gives (4).

Proof. Indeed, for any diffeomorphism $\varphi : M \rightarrow M$, the barcodes $\mathcal{B}(f)$ and $\mathcal{B}(f \circ \varphi)$ are the same. Since Φ_u depends only on the barcode and not on the function itself, putting $f \circ \varphi$ and $h \circ \psi$ in (3) yields (5). \square

Let us now extend the functional Φ_u from \mathcal{F}_{Morse} to $C^0(M)$. First, we introduce a “cut-off version” of Φ_u . Define

$$(6) \quad \Phi_{u,k}(f) = \begin{cases} \frac{\max f}{\min f} \int u(t) dt + \sum_{i=1}^k \int_{I_i} u(t) dt & \text{if } \partial M = \emptyset, \\ \int_{\min f}^0 u(t) dt + \sum_{i=1}^k \int_{I_i} u(t) dt & \text{if } \partial M \neq \emptyset. \end{cases}$$

where $I_i \in \mathcal{B}'(f)$ are finite intervals ordered by integral of u , i.e. we have

$$\int_{I_1} u(t) dt \geq \int_{I_2} u(t) dt \geq \dots$$

Lemma 1.3.5. *For every bounded function u the functional $\Phi_{u,k}$ is Lipschitz on \mathcal{F}_{Morse} with respect to d_{bottle} with Lipschitz constant $(2k+2) \cdot \max u$ if $\partial M = \emptyset$ or with Lipschitz constant $(2k+1) \cdot \max u$ if $\partial M \neq \emptyset$.*

The proof of Lemma 1.3.5 is given in subsection 4.2.

Assume now that $f \in C^0(M)$ is an arbitrary continuous function on M . Let $f_n \in \mathcal{F}_{Morse}$ such that $d_{C^0}(f, f_n) \rightarrow 0$ as $n \rightarrow \infty$. Set

$$(7) \quad \Phi_u(f) := \lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \Phi_{u,k}(f_n)$$

Note that $\Phi_u(f_n)$ only depends on $u|_{[\min f_n, \max f_n]}$ and, since for sufficiently large n it holds $[\min f_n, \max f_n] \subset [\min f - 1, \max f + 1]$, we may restrict ourselves to this interval and argue as if u was bounded. Thus due to Lemma 1.3.5 and (1), the double limit on the left hand side of (7) (which could be equal to $+\infty$) does not depend on the choice of the approximating sequence f_n . Therefore, the functional $\Phi_u(f)$ is well defined by (7). Moreover, it is easy to check that the right-hand sides of (7) and (2) coincide for $f \in \mathcal{F}_{Morse}$, and therefore (7) indeed defines an extension of (2) to $C^0(M)$.

1.4. Main results. As before, M is an orientable surface, possibly with boundary, equipped with a Riemannian metric g . Denote by $\|\cdot\|$ the L^2 -norm with respect to Riemannian area σ and by Δ the Laplace-Beltrami operator with respect to g . Slightly abusing the notation, throughout the paper κ_g will denote various constants depending on the Riemannian metric g .

Following⁴ [PS], denote by \mathcal{F}_λ the set of all smooth functions on M (vanishing on the boundary if $\partial M \neq \emptyset$) which satisfy $\|f\| = 1$ and $\|\Delta f\| \leq \lambda$. One may check that \mathcal{F}_λ contains normalized linear combinations of eigenfunctions of Δ with eigenvalues $\lambda_i \leq \lambda$. If $\partial M \neq \emptyset$, \mathcal{F}_λ contains also normalized eigenfunctions of the biharmonic clamped plate boundary value problem on M (see [PS, Example 1.2]).

Theorem 1.4.1. *Let $u \in C(\mathbb{R})$ be a non-negative function and $f \in \mathcal{F}_\lambda$ be a function on an orientable surface (M, g) . Then there exists a constant $\kappa_g > 0$ such that for any $\lambda > 0$,*

$$(8) \quad \Phi_u(f) \leq \kappa_g(\lambda + 1)\|u \circ f\|.$$

⁴Our definition is slightly different from the one in [PS] since we do not assume that $\int_M f \sigma = 0$ if M has no boundary. However, this assumption is not needed for any of the results of [PS] which we use.

In particular, taking $u \equiv 1$ we get the following corollary:

Corollary 1.4.2. *Let (M, g) be an orientable surface without boundary and let $f \in \mathcal{F}_\lambda$ be a Morse function on M . Denote by l_i the lengths of the finite bars of the barcode associated with f . Then*

$$(9) \quad \max f - \min f + \sum_i l_i \leq \kappa_g(\lambda + 1).$$

Example 1.4.3. The order of λ in inequality (9) is sharp. Indeed, consider the flat square torus $\mathbb{T}^2 = \mathbb{R}^2 / (2\pi \cdot \mathbb{Z})^2$. We have a sequence $f_n(x, y) = \frac{1}{\pi} \sin(nx) \cos(ny)$, $n \in \mathbb{N}$ of eigenfunctions of Δ with eigenvalues $2n^2$. By analysing critical points of $f_1 = \sin x \cos x$ and using periodicity, one can compute that the full barcode of f_n contains

- An infinite bar $(-\frac{1}{\pi}, +\infty)$ and $2n^2 - 1$ copies of finite bar $(-\frac{1}{\pi}, 0]$ in degree 0;
- Two copies of infinite bar $(0, +\infty)$ and $2n^2 - 1$ copies of finite bar $(0, \frac{1}{\pi}]$ in degree 1;
- An infinite bar $(\frac{1}{\pi}, +\infty)$ in degree 2.

Putting these values in inequality (9) gives us

$$\frac{4}{\pi} n^2 \leq \kappa_g(2n^2 + 1),$$

which proves that the order of λ in (9) is sharp.

In order to present another application of Theorem 1.4.1 we need the following definition.

Definition 1.4.4. Let $f : M \rightarrow \mathbb{R}$ be a Morse function on a differentiable manifold M and let $\delta > 0$. We say that a critical value $\alpha \in \mathbb{R}$ of the function f is a δ -significant critical value of multiplicity m if the barcode of f contains m bars of length at least δ having α as one of the endpoints.

Given $\delta > 0$ and a Morse function f , let $\mathcal{N}_\delta(f)$ be the number of δ -significant critical values counted with multiplicities. Theorem 1.4.1 then immediately implies:

Corollary 1.4.5. *Let (M, g) be an orientable surface, possibly with boundary, and let $f \in \mathcal{F}_\lambda$ be a Morse function on M . Then*

$$(10) \quad \mathcal{N}_\delta(f) \leq \kappa_{g,\delta}(\lambda + 1)$$

for any $\delta > 0$.

The following example shows that the δ -significance condition for some $\delta > 0$ is essential in Corollary 1.4.5. For simplicity, we present it in one dimension, but it could be easily generalized to any dimension.

Example 1.4.6. Let $M = \mathbb{S}^1$ be a unit circle and let N_i be any sequence of natural numbers tending to infinity. Consider a sequence of functions on M :

$$f_i(x) = \frac{1}{\sqrt{\pi(2 + N_i^{-4})}} \left(1 + \frac{1}{N_i^2} \sin(N_i x) \right).$$

It is easy to check that $\|f_i\|_{L^2(M)} = 1$ and $f_i \in \mathcal{F}_\lambda$, $\lambda \geq \frac{1}{\sqrt{2}}$, for all $i = 1, 2, \dots$. At the same time, the number of critical points, and hence of critical values (counted

with multiplicities) is equal to N_i , which goes to infinity and hence can not be controlled by λ . Note, however, that for any $\delta > 0$, the number of δ -significant critical values is bounded as $i \rightarrow \infty$.

Estimate (10) could be also compared to [Ni, Theorem 1.1], which shows that the expected value of the number of critical points of a random linear combination of Laplace eigenfunctions f_1, \dots, f_m on a Riemannian manifold satisfies an asymptotic expansion with the leading term of order m . Due to Weyl's law, for surfaces this is equivalent to having the number of critical points of order λ_m , which agrees with inequality (10). Inspired in part by this observation, we propose the following generalization of (10) to Riemannian manifolds of arbitrary dimension:

Conjecture 1.4.7. *Let (M, g) be a Riemannian manifold of dimension n , possibly with boundary, and let $f \in \mathcal{F}_\lambda$ be a Morse function on M . Then*

$$\mathcal{N}_\delta(f) \leq \kappa_{g,\delta}(\lambda + 1)^{\frac{n}{2}}$$

for any $\delta > 0$.

Furthermore, for n -dimensional Riemannian manifolds, consider the following generalization of the functional Φ_u : it is defined for Morse functions by an analogue of (2), the sum being taken over all finite bars in $\mathcal{B}(f)$ in all degrees. Similarly to (7) it also could be extended to arbitrary functions in $C^0(M)$.

Conjecture 1.4.8. *Let $u \in C(\mathbb{R})$ be a non-negative function and $f \in \mathcal{F}_\lambda$ be a function on an orientable Riemannian manifold (M, g) . Then there exists a constant $\kappa_g > 0$ such that for any $\lambda > 0$,*

$$\Phi_u(f) \leq \kappa_g(\lambda + 1)^{\frac{n}{2}} \|u \circ f\|.$$

A possible approach to proving this conjecture is discussed in Remark 3.2.2.

Example 1.4.9. Consider the n -dimensional torus $\mathbb{T}^n = \mathbb{R}^n / (2\pi \cdot \mathbb{Z})^n$ equipped with a Euclidean metric $ds^2 = \sum dx_i^2$. Define a sequence of functions

$$f_l(x_1, \dots, x_n) = \frac{\sqrt{2}}{n(2\pi)^{\frac{n}{2}}} (\sin lx_1 + \dots + \sin lx_n), \quad l \in \mathbb{N}.$$

It is easy to check that $\|f_l\| = 1$ and $\Delta f_l = l^2 f_l$. Thus $f_l \in \mathcal{F}_\lambda$ for $\lambda = l^2$. The following proposition shows that the order of λ predicted by Conjecture 1.4.8 is sharp.

Proposition 1.4.10. *There exist constants A_n and B_n such that*

$$\Phi_1(f_l) = A_n \lambda^{\frac{n}{2}} + B_n.$$

The proof of Proposition 1.4.10 uses the Künneth formula for persistence modules [PSS], see subsection 4.3 for details.

1.5. Sorting the finite bars of functions in \mathcal{F}_λ . Given a barcode \mathcal{B} , write the lengths of its finite bars in the descending order, $\beta_1 \geq \beta_2 \geq \dots$. The functions $\beta_i(\mathcal{B})$, which are Lipschitz with respect to the bottleneck distance, are important invariants of barcodes. For instance, β_1 , which was introduced by Usher in [U], is called *the boundary depth* and has various applications in Morse theory and symplectic topology. The functions β_i with $i \geq 2$ are sometimes used in order to

distinguish barcodes, see e.g. [BMMPS]. Fix $\epsilon > 0$ and discard all bars of length $< \epsilon$, i.e., introduce the modified invariant

$$\beta_i^{(\epsilon)}(\mathcal{B}) := \max(\beta_i(\mathcal{B}), \epsilon).$$

Question 1.5.1. *Assume that the barcode \mathcal{B} contains N finite bars. What is the optimal (worst-case scenario) running time T of an algorithm which calculates the ordered sequence $\{\beta_i^{(\epsilon)}(\mathcal{B})\}$, $i \geq 1$?*

Since the sharp lower bound on the running time of any comparison sorting algorithm for an array of N real numbers is $O(N \log N)$ (see [CLRS]), the answer to the above question for a general barcode is $O(N \log N)$. Interestingly enough, in some cases Corollary 1.4.2 enables one to reduce this running time when \mathcal{B} is a barcode of a function from \mathcal{F}_λ . More precisely, there exists a constant $c > 0$ such that for every $\epsilon > 0$, $\lambda > 0$ and any function $f \in \mathcal{F}_\lambda$ whose barcode contains exactly N finite bars, one can find a sorting algorithm for all bars from $\mathcal{B}(f)$ of length $\geq \epsilon$ whose running time satisfies

$$(11) \quad T \leq N + c \cdot \frac{\kappa_g(\lambda + 1)}{\epsilon} \cdot \log \frac{\kappa_g(\lambda + 1)}{\epsilon}.$$

Indeed, consider the following algorithm. First compare the length of each bar with ϵ and pick only those bars whose length is $\geq \epsilon$. This takes time N . Denote by K the number of chosen bars. Next, perform the optimal sorting algorithm for these K bars. This takes time $O(K \log K)$. Finally, notice that by Corollary 1.4.2,

$$K \leq \frac{\kappa_g(\lambda + 1)}{\epsilon},$$

which proves (11). In certain regimes, the running time (11) is shorter than the generic bound $O(N \log N)$. For instance, if λ and ϵ are fixed and $N \rightarrow \infty$, we have $T = O(N)$.

Theorem 1.4.1 also has applications to questions regarding C^0 -approximations by functions from \mathcal{F}_λ , which is the subject of the next section.

2. APPLICATIONS TO APPROXIMATIONS BY EIGENFUNCTIONS

2.1. An obstruction to C^0 -approximations. As before, let M be an orientable surface, possibly with boundary, endowed with the Riemannian metric g , denote by $\text{Diff}(M)$ the set of all diffeomorphisms of M and assume that $f : M \rightarrow \mathbb{R}$ is a Morse function (vanishing on ∂M). We are interested in the question of how well can f be approximated by functions from \mathcal{F}_λ in C^0 -sense. More precisely, we wish to find a lower bound for the quantity

$$d_{C^0}(f, \mathcal{F}_\lambda) = \inf\{d_{C^0}(f, h) \mid h \in \mathcal{F}_\lambda\},$$

where $d_{C^0}(f, h) = \max_x |f(x) - h(x)|$ as before. In fact, we will study a more general question, namely we will give a lower bound for

$$\text{approx}_\lambda(f) = \inf_{\varphi \in \text{Diff}(M)} d_{C^0}(f \circ \varphi, \mathcal{F}_\lambda).$$

Taking $\varphi = \mathbf{1}_M$ one immediately sees that

$$d_{C^0}(f, \mathcal{F}_\lambda) \geq \text{approx}_\lambda(f).$$

We estimate $\text{approx}_\lambda(f)$ from below using the information coming from the barcode $\mathcal{B}(f)$. Recall that the functional $\Phi_1 : \mathcal{F}_{\text{Morse}} \rightarrow \mathbb{R}$ defined by (2) for $u \equiv 1$ gives the sum of the lengths of all finite bars in $\mathcal{B}(f)$.

Proposition 2.1.1. *For every Morse function $f : M \rightarrow \mathbb{R}$, vanishing on the boundary, the following inequality holds*

$$(12) \quad \text{approx}_\lambda(f) \geq \begin{cases} \frac{1}{2 \cdot (|\mathcal{B}'(f)|+1)} \left(\Phi_1(f) - \kappa_g(\lambda + 1) \right) & \text{for } \partial M = \emptyset \\ \frac{1}{2|\mathcal{B}'(f)|+1} \left(\Phi_1(f) - \kappa_g(\lambda + 1) \right) & \text{for } \partial M \neq \emptyset \end{cases}$$

Proof. From (5) and (8), with $\psi = \mathbf{1}_M$ we obtain

$$\kappa_g(\lambda + 1) \cdot \|u \circ h\| \geq \Phi_u(f) - C(u, f) \cdot d_{C^0}(f \circ \varphi, h),$$

for all Morse $h \in \mathcal{F}_\lambda$ and all diffeomorphisms $\varphi \in \text{Diff}(M)$, with constant $C(u, f)$ being equal to $2 \cdot (|\mathcal{B}'(f)|+1) \cdot (\max_{[\min f, \max f]} u)$ or $(2|\mathcal{B}'(f)|+1) \cdot (\max_{[\min f, \max f]} u)$ depending on whether M has a boundary. Putting $u \equiv 1$ we have

$$d_{C^0}(f \circ \varphi, h) \geq \frac{1}{2 \cdot (|\mathcal{B}'(f)|+1)} \left(\Phi_1(f) - \kappa_g(\lambda + 1) \right),$$

if $\partial M = \emptyset$, or

$$d_{C^0}(f \circ \varphi, h) \geq \frac{1}{2|\mathcal{B}'(f)|+1} \left(\Phi_1(f) - \kappa_g(\lambda + 1) \right),$$

if $\partial M \neq \emptyset$. Finally, taking infimum over all h and φ and using the fact that Morse functions in \mathcal{F}_λ are C^0 -dense in \mathcal{F}_λ , finishes the proof. \square

Remark 2.1.2. The following analogue of Proposition 2.1.1 holds for any function $f \in C^0(M)$. For any $k = 1, 2, \dots$, we have

$$(13) \quad \text{approx}_\lambda(f) \geq \begin{cases} \frac{1}{2k+2} \left(\Phi_{1,k}(f) - \kappa_g(\lambda + 1) \right) & \text{for } \partial M = \emptyset \\ \frac{1}{2k+1} \left(\Phi_{1,k}(f) - \kappa_g(\lambda + 1) \right) & \text{for } \partial M \neq \emptyset \end{cases}$$

The proof is the same, with the constant $C(u, f)$ replaced by the Lipschitz constant from Lemma 1.3.5.

Corollary 2.1.3. *Let M be a surface without boundary and $f : M \rightarrow \mathbb{R}$ be a Morse function. Suppose that $\text{approx}_\lambda(f) \leq \epsilon$ for some $\epsilon > 0$, and the barcode $\mathcal{B}(f)$ contains N finite bars of length at least $L + 2\epsilon$ each, for some $L > 0$. Then*

$$(14) \quad \lambda \geq \frac{1}{\kappa_g} (N + 1)L - 1.$$

Proof. Indeed, it follows from the assumptions on the barcode of f that $\Phi_{1,N}(f) \geq (N + 1)(L + 2\epsilon)$, which together with Remark 2.1.2 yields

$$\epsilon \geq \frac{1}{2(N + 1)} \left((N + 1)(L + 2\epsilon) - \kappa_g(\lambda + 1) \right).$$

Rearranging this inequality we obtain (14). \square

Remark 2.1.4. From (14) we see how λ , which is needed to uniformly approximate f by functions from \mathcal{F}_λ , grows with N and L . Informally speaking, one may think of N as a measure of how much f oscillates, while L gives a lower bound on the amplitude of these oscillations. The above inequality should then be understood as a quantitative version of the informal principle that the more the function oscillates and the bigger the oscillations, the larger eigenvalues of the Laplacian are needed to approximate it with a normalized linear combination of the corresponding eigenfunctions.

2.2. Modulus of continuity and average length of bars. Proposition 2.1.1 gives an obstruction to approximating functions by functions from \mathcal{F}_λ . As we mentioned before, \mathcal{F}_λ contains normalized linear combinations of eigenfunctions of Δ with eigenvalues not greater than λ . In the case of flat torus $\mathbb{T}^2 = \mathbb{R}^2/(2\pi \cdot \mathbb{Z})^2$ these eigenfunctions are trigonometric polynomials and Proposition 2.1.1 may be interpreted as an inverse statement about C^0 -approximations by trigonometric polynomials. A direct theorem about C^0 -approximations by trigonometric polynomials on n -dimensional flat torus was proved in [Yu] (theorems of this type are sometimes referred to as Jackson's theorems, see [Pi] for a survey), consequently giving an upper bound for $\text{approx}_\lambda(f)$ in terms of moduli of continuity and smoothness of f . We combine this result with Proposition 2.1.1 to obtain a relation between the average length of a bar in a barcode of a Morse function on \mathbb{T}^2 and its modulus of continuity which is defined below.

Assume $M = \mathbb{T}^2 = \mathbb{R}^2/(2\pi \cdot \mathbb{Z})^2$, let $f : M \rightarrow \mathbb{R}$ be a continuous function, $\delta > 0$ a real number and denote by

$$\omega_1(f, \delta) = \sup_{|t| \leq \delta} \max_x |f(x+t) - f(x)|,$$

the *modulus of continuity* of f and by

$$\omega_2(f, \delta) = \sup_{|t| \leq \delta} \max_x |f(x-t) - 2f(x) + f(x+t)|,$$

the *modulus of smoothness* of f . One readily checks that

$$(15) \quad \omega_2(f, \delta) \leq 2\omega_1(f, \delta).$$

Let

$$\mathcal{T}_\lambda = \left\langle \sin(v_1x + v_2y), \cos(v_1x + v_2y) \mid v_1^2 + v_2^2 \leq \lambda \right\rangle_{\mathbb{R}},$$

be the space of trigonometric polynomials on M whose eigenvalues (as eigenfunctions of Δ) are bounded by λ . The following proposition was proved in [Yu]:

Proposition 2.2.1. *For every continuous function $f : M \rightarrow \mathbb{R}$ it holds*

$$(16) \quad d_{C^0}(f, \mathcal{T}_\lambda) \leq 2\omega_2\left(f, \frac{C_0}{\sqrt{\lambda}}\right),$$

where $C_0 > 0$ is a constant.

By (15) we also have that

$$(17) \quad d_{C^0}(f, \mathcal{T}_\lambda) \leq 4\omega_1\left(f, \frac{C_0}{\sqrt{\lambda}}\right).$$

Remark 2.2.2. Constant C_0 is computed in [Yu] to be $C_0 = \sqrt{\mu_1(D^2(\frac{1}{2}))}$, where $\mu_1(D^2(\frac{1}{2}))$ is the first Dirichlet eigenvalue of Δ inside the 2-dimensional disk $D^2(\frac{1}{2})$ of radius $\frac{1}{2}$.

Our goal is to prove the following result which shows that the average bar length of a Morse function f on a flat torus M could be *uniformly* controlled by the L^2 norm of f and the modulus of continuity of f on the scale $1/\sqrt{|\mathcal{B}'(f)|}$.

Theorem 2.2.3. *There exist constants $C_0, C_1 > 0$ such that for any $f \in \mathcal{F}_{Morse}$ on a flat torus $M = \mathbb{T}^2$,*

$$(18) \quad \frac{\Phi_1(f)}{|\mathcal{B}'(f)| + 1} \leq C_1 \|f\| + 8\omega_1 \left(f, \frac{C_0}{\sqrt{|\mathcal{B}'(f)|}} \right).$$

Proof. Inspecting the proof of Proposition 2.2.1 in [Yu], we observe that it relies on an explicit construction of a function h , depending on f and λ , which satisfies

$$(19) \quad d_{C^0}(f, h) \leq 2\omega_2 \left(f, \frac{C_0}{\sqrt{\lambda}} \right).$$

Our goal is to estimate $d_{C^0}(f, h)$ from below using Proposition 2.1.1. However, a priori we do not have any information about the L^2 -norm of h and Proposition 2.1.1 relates to distance from functions in \mathcal{F}_λ whose L^2 -norm is equal to one. In order to overcome this issue, we present the construction of the approximation-function h and prove that $\|h\| \leq \|f\|$.

For a vector $v \in \mathbb{Z}^2$ let

$$c_v(f) = \frac{1}{(2\pi)^2} \int_{\mathbb{T}^2} f(x) e^{-i\langle v, x \rangle} dx,$$

be the corresponding Fourier coefficient of f . Take U to be the first Dirichlet eigenfunction of Δ inside the disk $D^2(\frac{1}{2})$ of radius $\frac{1}{2}$, normalized by $\|U\|_{L^2(D^2(\frac{1}{2}))} = 1$, and V its extension by zero to the whole plane, i.e.

$$V(x) = \begin{cases} U(x), & \text{if } x \in D^2(\frac{1}{2}) \\ 0, & \text{otherwise.} \end{cases}$$

If we denote by $W = V * V$ the convolution of V with itself, the desired approximation is given by the formula

$$(20) \quad h(x) = \sum_{\substack{v \in \mathbb{Z}^2 \\ |v| \leq \sqrt{\lambda}}} c_v(f) \cdot W \left(\frac{v}{\sqrt{\lambda}} \right) e^{i\langle v, x \rangle},$$

where $|v|$ stands for the standard Euclidean norm on \mathbb{R}^2 . The function h defined by (20) is called the *multidimensional Korovkin's mean*. It satisfies (19), as proved in [Yu], and since $\sum_{v \in \mathbb{Z}^2} c_v(f) e^{i\langle v, x \rangle}$ is the Fourier expansion of f , we have that

$$\|h\| \leq \left(\max_{x \in \mathbb{R}^2} |W(x)| \right) \cdot \|f\|.$$

By using the Cauchy-Schwarz inequality we obtain

$$|W(x)| \leq \int_{\mathbb{R}^2} |V(t)| \cdot |V(x-t)| dt \leq \sqrt{\int_{\mathbb{R}^2} |V(t)|^2 dt} \cdot \sqrt{\int_{\mathbb{R}^2} |V(x-t)|^2 dt} = \|V\|^2 = 1,$$

which yields $\|h\| \leq \|f\|$.

We now proceed with analysing (19). First note that

$$\|h\| \cdot d_{C^0} \left(\frac{f}{\|h\|}, \mathcal{F}_\lambda \right) \leq \|h\| \cdot d_{C^0} \left(\frac{f}{\|h\|}, \frac{h}{\|h\|} \right) \leq 2\omega_2 \left(f, \frac{C_0}{\sqrt{\lambda}} \right),$$

because $\frac{h}{\|h\|} \in \mathcal{F}_\lambda$. The last inequality together with Proposition 2.1.1 gives

$$(21) \quad \frac{\|h\|}{2 \cdot (|\mathcal{B}'(\frac{f}{\|h\|})| + 1)} \left(\Phi_1 \left(\frac{f}{\|h\|} \right) - \kappa_0(\lambda + 1) \right) \leq 2\omega_2 \left(f, \frac{C_0}{\sqrt{\lambda}} \right).$$

Here $\kappa_0 = \kappa_g$ for g being the flat metric on M . Multiplying the function by a positive constant results in multiplying the endpoints of each bar in the barcode by the same constant. Thus, the total number of bars does not change after multiplication, while the lengths of finite bars scale with the same constant. In other words, we have that $|\mathcal{B}'(\frac{f}{\|h\|})| = |\mathcal{B}'(f)|$ and $\|h\| \cdot \Phi_1(\frac{f}{\|h\|}) = \Phi_1(f)$. Substituting these equalities in (21) and using $\|h\| \leq \|f\|$ we obtain

$$\frac{1}{4 \cdot (|\mathcal{B}'(f)| + 1)} \left(\Phi_1(f) - \kappa_0(\lambda + 1)\|f\| \right) \leq \omega_2 \left(f, \frac{C_0}{\sqrt{\lambda}} \right),$$

and by (15) also

$$\frac{1}{8 \cdot (|\mathcal{B}'(f)| + 1)} \left(\Phi_1(f) - \kappa_0(\lambda + 1)\|f\| \right) \leq \omega_1 \left(f, \frac{C_0}{\sqrt{\lambda}} \right).$$

Setting $\lambda = |\mathcal{B}'(f)|$ and $C_1 = \kappa_0$ in the last inequality completes the proof of the theorem. \square

Remark 2.2.4. As follows from Remark 2.1.2 and the proof of Theorem 2.2.3 above, for any $k \geq 1$ and any $f \in C^0(M)$ we have:

$$(22) \quad \frac{\Phi_{1,k}(f)}{k+1} \leq C_1 \|f\| + 8\omega_1 \left(f, \frac{C_0}{\sqrt{k}} \right).$$

The left-hand side of (22) could be interpreted as the average length of a bar among the k longest bars in the barcode of f .

Example 2.2.5. The following example shows that the choice of the scale in the modulus of continuity on the right hand side of (22) is optimal. Take a unit disk B_1 inside the torus M and let χ be a smooth cut-off function supported in B and equal to one in $B_{\frac{1}{2}}$. Let $g_n(x, y) = \chi(x, y) \sin nx \cos ny$, $n \in \mathbb{N}$ be a sequence of functions on the torus. For any $0 < s < 1$, set $g_{n,s}(x, y) = g(\frac{x}{s}, \frac{y}{s})$. Let $\alpha \geq 1$ be some real number. Choose $k = n^2$ and $s = n^{\frac{1-\alpha}{2}}$. It suffices to verify that the inequality

$$(23) \quad \frac{\Phi_{1,n^2}(g_{n,s})}{n^2 + 1} \leq C_1 \|g_{n,s}\| + 8\omega_1 \left(g_{n,s}, \frac{C_0}{n^\alpha} \right).$$

holds for all n only for $\alpha = 1$. Indeed, take any $\alpha > 1$. Note that the left-hand side of (23) is bounded away from zero as $n \rightarrow \infty$, since the number of bars of unit length in the barcode of $g_{n,s}(x, y)$ is of order n^2 . At the same time, $\|g_{n,s}\| = s\|g_n\| \rightarrow 0$

as $n \rightarrow \infty$, since $s = n^{\frac{1-\alpha}{2}} \rightarrow 0$. Moreover, estimating the derivatives of $g_{n,s}$ one can easily check that

$$\omega_1 \left(g_{n,s}, \frac{C_0}{n^\alpha} \right) = O(n^{-\alpha}) O(n \cdot n^{\frac{\alpha-1}{2}}) = O \left(n^{\frac{1-\alpha}{2}} \right) = o(1)$$

for any $\alpha > 1$. Therefore, inequality (23) is violated for $\alpha > 1$ for n large, and hence the choice $\alpha = 1$ is optimal.

Note that, while the functions $g_{n,s}(x, y)$ are compactly supported and hence not Morse, they could be made Morse by adding a small perturbation. A similar argument would then yield optimality of the $1/\sqrt{|\mathcal{B}'(f)|}$ scale in the modulus of continuity on the right-hand side of inequality (18).

Remark 2.2.6. It would be interesting to establish an analogue of Theorem 2.2.3 for Morse functions on an arbitrary Riemannian surface.

3. BARCODES AND THE BANACH INDICATRIX

3.1. A topological bound on the Banach indicatrix. We proceed with some general topological considerations. Let M be a Riemannian manifold and assume that $f|_{\partial M} = 0$ if $\partial M \neq \emptyset$, 0 being a regular value. Let $t \neq 0$ be another regular value of f and denote by $M^t = f^{-1}((-\infty, t])$. $M^t \subset M$ is a submanifold with boundary $\partial M^t = f^{-1}(t)$ for $t < 0$ or $\partial M^t = f^{-1}(t) \sqcup \partial M$ for $t > 0$. Denote by $\beta(t, f)$ the number of connected components of $f^{-1}(t)$. Function $\beta(t, f)$ is called the *Banach indicatrix* of f (see [Kr, Yo]). By description of ∂M^t , one sees that $\beta(t, f)$ essentially counts the number of the boundary components of M^t (up to the boundary components of the whole manifold M). We will exploit this fact to estimate $\beta(t, f)$ from below using information coming from barcode $\mathcal{B}(f)$. If we denote by χ_I the characteristic function of the interval I , the following proposition holds.

Proposition 3.1.1. *Let $f \in \mathcal{F}_{Morse}$ on a Riemannian manifold M of dimension n . Denote by $(x_i^{(k)}, y_i^{(k)}) \in \mathcal{B}_k(f)$ the finite bars in the degree k barcode of f and let $t \neq 0$ be a regular value. If $\partial M = \emptyset$ it holds*

$$(24) \quad \chi_{(\min f, \max f]}(t) + \sum_i \chi_{(x_i^{(0)}, y_i^{(0)})}(t) + \sum_j \chi_{(x_j^{(n-1)}, y_j^{(n-1)})}(t) \leq \beta(t, f),$$

and if $\partial M \neq \emptyset$ it holds

$$(25) \quad \chi_{(\min f, 0]}(t) + \sum_i \chi_{(x_i^{(0)}, y_i^{(0)})}(t) + \sum_j \chi_{(x_j^{(n-1)}, y_j^{(n-1)})}(t) \leq \beta(t, f).$$

We defer proving Proposition 3.1.1 and first deduce Theorem 1.4.1 using it.

Remark 3.1.2. If $\dim M = 2$, integrating inequalities in Proposition 3.1.1 gives an upper bound on the total length of the finite bars in the barcode of a function f in terms of the integral of its Banach indicatrix. The latter quantity admits an interpretation as the total length of the Reeb graph of a function f with respect to a natural metric incorporating the oscillations of f . It is likely that an analogue of the functional Φ_u defined in this setting is robust with respect to the distance on Reeb graphs introduced in [BGW, BMW], and that this way one could get applications to approximation theory similar to the ones obtained in Section 2. We plan to explore this route elsewhere.

3.2. Proof of Theorem 1.4.1. Let us now restrict to the two-dimensional case and assume that M is an orientable surface, possibly with boundary, equipped with Riemannian metric g . First note that it suffices to verify inequality (8) for Morse functions. Indeed, suppose that the inequality is proved for Morse functions in \mathcal{F}_λ for all $\lambda > 0$ and let $f \in \mathcal{F}_\lambda$ be arbitrary. One can easily check that for any $\epsilon > 0$ there exists a sequence of Morse functions $f_n \in \mathcal{F}_{\lambda+\epsilon}$ such that $d_{C^0}(f, f_n) \rightarrow 0$ as $n \rightarrow \infty$. For all $k, n \geq 1$ we have

$$\Phi_{u,k}(f_n) \leq \Phi_u(f_n) \leq \kappa_g(\lambda + 1 + \epsilon),$$

where the first inequality follows from the definition (6) of the functional $\Phi_{u,k}$ and the second inequality holds by the assumption that (8) is true for Morse functions. Taking the limits as k and n go to infinity in definition (7) and using the fact that $\epsilon > 0$ is arbitrary, we obtain that (8) holds for the function f .

It remains to prove inequality (8) when $f \in \mathcal{F}_\lambda$ is Morse. Denote by $\|\cdot\|$ the L^2 -norm with respect to Riemannian area σ and by Δ the Laplace-Beltrami operator with respect to g . The analytical tool that we are going to use is [PS, Theorem 1.5] which gives us that for any continuous function $u \in C(\mathbb{R})$ and any smooth function f on M (which is assumed to be equal to zero on the boundary if $\partial M \neq \emptyset$), the following inequality holds

$$(26) \quad \int_{-\infty}^{+\infty} u(t)\beta(t, f) dt \leq \kappa_g(\|f\| + \|\Delta f\|) \cdot \|u \circ f\|,$$

where κ_g depends on the Riemannian metric g .

If we assume that $f \in \mathcal{F}_\lambda$ in (26), we immediately get

$$(27) \quad \int_{-\infty}^{+\infty} u(t)\beta(t, f) dt \leq \kappa_g(\lambda + 1) \cdot \|u \circ f\|.$$

Since f is Morse we can apply Proposition 3.1.1. Combining this proposition with inequality (27) immediately yields Theorem 1.4.1. \square

Remark 3.2.1. It follows from the proof of Theorem 1.4.1 that inequality (8) holds for any function in the closure of \mathcal{F}_λ in C^0 topology.

Remark 3.2.2. The proof of Theorem 1.4.1 suggests the following approach to proving Conjecture 1.4.8. Recall that, by definition, the Banach indicatrix is given by $\beta(t, f) = b_0(f^{-1}(t))$. In view of Proposition 3.1.1, it is plausible that the following inequality holds in dimension $n \geq 3$:

$$\Phi_u(f) \leq \int_{-\infty}^{+\infty} \left(\sum_{i=0}^{n-2} b_i(f^{-1}(t)) \right) u(t) dt,$$

where b_i is the i -th Betti number. As follows from [Yo], the quantity on the right-hand side could be bounded from above using the uniform norm of the derivatives of f (see also [LL] for related recent developments). In order to establish Conjecture 1.4.8, one would need to prove a higher-dimensional analogue of [PS, Theorem 1.5], allowing to replace the uniform estimates by L^2 bounds.

3.3. Proof of Proposition 3.1.1. Recall the notation previously introduced. For t regular value of f , we denote $M^t = f^{-1}((-\infty, t]) \subset M$. As mentioned before M^t is a submanifold with boundary $\partial M^t = f^{-1}(t)$ for $t < 0$ or $\partial M^t = f^{-1}(t) \sqcup \partial M$ for $t > 0$. Let $b_k(t) = \dim H_k(M^t; \mathbb{R})$ be the Betti numbers of M^t . We assume that $\min f < t < \max f$, since otherwise the inequalities obviously hold (both sides are equal to zero). We will always work with homologies with coefficients in \mathbb{R} and will omit the coefficients from the notation.

Let $j : f^{-1}(t) \rightarrow M^t$ be the inclusion and denote by $j_* : H_{n-1}(f^{-1}(t)) \rightarrow H_{n-1}(M^t)$ the induced map in homology. Since we work over \mathbb{R} , $j_*(H_{n-1}(f^{-1}(t))) \subset H_{n-1}(M^t)$ is a vector subspace. First, we claim that

$$(28) \quad \beta(t, f) = \dim H_n(M^t, f^{-1}(t)) + \dim(j_*(H_{n-1}(f^{-1}(t)))).$$

To prove this, we examine the following part of the long exact sequence of the pair $(M^t, f^{-1}(t))$:

$$H_n(M^t) \rightarrow H_n(M^t, f^{-1}(t)) \xrightarrow{\partial} H_{n-1}(f^{-1}(t)) \xrightarrow{j_*} H_{n-1}(M^t).$$

Since $f^{-1}(t)$ is an $(n-1)$ -dimensional orientable manifold, we have $\beta(t, f) = \dim H_{n-1}(f^{-1}(t))$, and by the Rank-nullity theorem

$$\beta(t, f) = \dim H_{n-1}(f^{-1}(t)) = \dim(\ker j_*) + \dim(j_*(H_{n-1}(f^{-1}(t)))).$$

By the exactness $\dim(\ker j_*) = \dim(\text{im } \partial)$ and, because $H_n(M^t) = 0$, ∂ is an inclusion, which means that $\dim(\text{im } \partial) = \dim H_n(M^t, f^{-1}(t))$ and (28) follows.

Second, we note that

$$(29) \quad \sum_j \chi_{(x_j^{(n-1)}, y_j^{(n-1)})}(t) = \dim(\ker i_*),$$

where $i : M^t \rightarrow M$ is the inclusion and $i_* : H_{n-1}(M^t) \rightarrow H_{n-1}(M)$ induced map on homology. This comes from the fact that finite bars in barcode of f correspond to homology classes which appear throughout filtration process, but do not exist in actual homology of M . Denote by $\widetilde{M}^t = f^{-1}([t, +\infty))$, $\widetilde{M}^t \cap M^t = f^{-1}(t)$, and by $\tilde{j} : f^{-1}(t) \rightarrow \widetilde{M}^t$ and $\tilde{i} : \widetilde{M}^t \rightarrow M$ the inclusions. We examine the following part of the Mayer-Vietoris sequence

$$H_{n-1}(f^{-1}(t)) \xrightarrow{(j_*, \tilde{j}_*)} H_{n-1}(M^t) \oplus H_{n-1}(\widetilde{M}^t) \xrightarrow{i_* - \tilde{i}_*} H_{n-1}(M).$$

From the exactness we have

$$\ker i_* \cong \ker(i_* - \tilde{i}_*) \cap (H_{n-1}(M^t), 0) = \text{im}(j_*, \tilde{j}_*) \cap (H_{n-1}(M^t), 0),$$

while on the other hand

$$\begin{aligned} \text{im}(j_*, \tilde{j}_*) \cap (H_{n-1}(M^t), 0) &= \{(j_*(a), \tilde{j}_*(a)) \mid a \in H_{n-1}(f^{-1}(t)), \tilde{j}_*(a) = 0\} = \\ &= \{(j_*(a), 0) \mid a \in \ker \tilde{j}_* \subset H_{n-1}(f^{-1}(t))\} \cong j_*(\ker \tilde{j}_*), \end{aligned}$$

and thus

$$\ker i_* \cong j_*(\ker \tilde{j}_*).$$

However, since $\tilde{i}_* \circ \tilde{j}_* = i_* \circ j_*$, we see that $j_*(\ker \tilde{j}_*) \subset \ker i_*$ and thus

$$(30) \quad \ker i_* = j_*(\ker \tilde{j}_*).$$

From now on, we distinguish two cases.

Case I - $\partial M^t = f^{-1}(t)$

This case covers the situation when $\partial M = \emptyset$ and when $\partial M \neq \emptyset$, but $t < 0$. The left hand sides of the inequalities (24) and (25) are equal for $t < 0$ and thus we need to prove (24). By the case-assumption, we have that

$$\dim H_n(M^t, f^{-1}(t)) = b_0(t),$$

and from the definition of barcode we know that

$$b_0(t) = \chi_{(\min f, \max f]}(t) + \sum_i \chi_{(x_i^{(0)}, y_i^{(0)})}(t).$$

Combining these equalities with (28) renders the statement into

$$\sum_j \chi_{(x_j^{(n-1)}, y_j^{(n-1)})}(t) \leq \dim(j_*(H_{n-1}(f^{-1}(t)))),$$

which after substituting (29) and (30) becomes

$$\dim(j_*(\ker \tilde{j}_*)) \leq \dim(j_*(H_{n-1}(f^{-1}(t)))).$$

This inequality is obvious because $\ker \tilde{j}_* \subset H_{n-1}(f^{-1}(t))$.

Case II - $\partial M^t = f^{-1}(t) \sqcup \partial M$, $\partial M \neq \emptyset$

This case covers the situation when $\partial M \neq \emptyset$ and $t > 0$. We need to prove (25), which for $t > 0$ becomes

$$\sum_i \chi_{(x_i^{(0)}, y_i^{(0)})}(t) + \sum_j \chi_{(x_j^{(n-1)}, y_j^{(n-1)})}(t) \leq \beta(t, f).$$

Denote by $\partial M = \Sigma_1 \sqcup \dots \sqcup \Sigma_l$ the boundary components of the whole manifold M , where Σ_i are connected, orientable, $(n-1)$ -dimensional manifolds. Now the boundary of M^t is $\partial M^t = f^{-1}(t) \sqcup \Sigma_1 \sqcup \dots \sqcup \Sigma_l$. We may divide connected components of M^t into two groups, one of which consists of all the components whose boundary lies entirely in $f^{-1}(t)$ and the other one consists of all the components whose boundary contains at least one Σ_i (i.e. the boundary of these components is a mix of parts of ∂M and $f^{-1}(t)$). Denote by k the number of connected components of M^t whose boundary contains at least one Σ_i . Now, since

$$\dim H_n(M^t, f^{-1}(t) \sqcup \partial M) = b_0(t),$$

we have that

$$\dim H_n(M^t, f^{-1}(t)) = b_0(t) - k,$$

and thus by (28)

$$\beta(t, f) = b_0(t) - 1 + \dim(j_*(H_{n-1}(f^{-1}(t)))) - (k - 1).$$

Since M is connected

$$b_0(t) - 1 = \sum_i \chi_{(x_i^{(0)}, y_i^{(0)})}(t),$$

and hence we need to prove that

$$\sum_j \chi_{(x_j^{(n-1)}, y_j^{(n-1)})}(t) \leq \dim(j_*(H_{n-1}(f^{-1}(t)))) - (k - 1).$$

Using (29) we transform the statement into

$$\dim(\ker i_*) + k - 1 \leq \dim(j_*(H_{n-1}(f^{-1}(t)))).$$

In order to prove this inequality, we will find $k - 1$ linearly independent vectors in the quotient space $j_*(H_{n-1}(f^{-1}(t)))/\ker i_*$ (note that by (30) we have that $\ker i_* = j_*(\ker \tilde{j}_*) \subset j_*(H_{n-1}(f^{-1}(t)))$). Assume that $k \geq 2$ (because otherwise the statement is trivial) and denote by M_1^t, \dots, M_k^t the connected components of M^t whose boundary contains some Σ_i . We know that for $1 \leq i \leq k$, homology class $0 = [\partial M_i^t] \in H_{n-1}(M^t)$ decomposes as $0 = [\partial M_i^t] = d_i + e_i$, where $d_i = [\Sigma_{i_1}] + \dots + [\Sigma_{i_{m_i}}]$ for some $[\Sigma_{i_1}], \dots, [\Sigma_{i_{m_i}}]$ and $e_i \in j_*(H_{n-1}(f^{-1}(t)))$. Moreover, since M_i^t are disjoint, we have that

$$d_1 + \dots + d_k = [\Sigma_1] + \dots + [\Sigma_l],$$

and d_1, \dots, d_k partition the set $\{[\Sigma_1], \dots, [\Sigma_l]\}$. We have that

$$d_1, \dots, d_k \in j_*(H_{n-1}(f^{-1}(t))),$$

because $d_i = -e_i$, and let $[d_1], \dots, [d_k] \in j_*(H_{n-1}(f^{-1}(t)))/\ker i_*$ be the corresponding classes inside the quotient space. We claim that any $k - 1$ of $[d_1], \dots, [d_k]$ are linearly independent. Once we prove this, choosing any $k - 1$ of these gives us the $k - 1$ classes that we need.

First, we observe that $[\Sigma_1], \dots, [\Sigma_l] \in H_{n-1}(M^t)$ are linearly independent. In order to prove this, consider the following part of the long exact sequence of the pair $(M^t, \partial M)$:

$$H_n(M^t, \partial M) \rightarrow H_{n-1}(\partial M) \rightarrow H_{n-1}(M^t).$$

Note that $H_n(M^t, \partial M) = 0$. Indeed, if $H_n(M^t, \partial M) \neq 0$, then M^t contains a connected component N , such that $\partial N \subset \partial M$. However, this implies that $f^{-1}(t) \cap N = \emptyset$, or equivalently, $N \subset f^{-1}((-\infty, t))$. It is now easy to check that $N \subset M$ is both an open and a closed subset, which contradicts the fact that M is connected. Therefore, $H_n(M^t, \partial M) = 0$, and hence $H_{n-1}(\partial M) \rightarrow H_{n-1}(M^t)$ is an injection, i.e. $[\Sigma_1], \dots, [\Sigma_l]$ are linearly independent. This further implies that d_1, \dots, d_k are linearly independent.

Classes $i_*d_1, \dots, i_*d_k \in H_{n-1}(M)$ satisfy

$$i_*d_1 + \dots + i_*d_k = i_*[\Sigma_1] + \dots + i_*[\Sigma_l] = [\partial M] = 0.$$

By using the exactness of the following part of the long exact sequence of the pair $(M, \partial M)$

$$0 = H_n(M) \rightarrow H_n(M, \partial M) \rightarrow H_{n-1}(\partial M) \rightarrow H_{n-1}(M),$$

we conclude that

$$i_*[\Sigma_1] + \dots + i_*[\Sigma_l] = 0,$$

is the only relation which $i_*[\Sigma_1], \dots, i_*[\Sigma_l]$ satisfy. More formally, restriction of i_* to $\langle [\Sigma_1], \dots, [\Sigma_l] \rangle_{\mathbb{R}} \subset H_{n-1}(M^t)$ has the one-dimensional kernel given by

$$\ker(i_*|_{\langle [\Sigma_1], \dots, [\Sigma_l] \rangle_{\mathbb{R}}}) = \langle [\Sigma_1] + \dots + [\Sigma_l] \rangle_{\mathbb{R}}.$$

This readily implies the only relation which i_*d_1, \dots, i_*d_k satisfy is that their sum is zero, that is

$$\ker(i_*|_{\langle d_1, \dots, d_k \rangle_{\mathbb{R}}}) = \langle d_1 + \dots + d_k \rangle_{\mathbb{R}}.$$

Finally, combining the last equality with the fact that d_1, \dots, d_k are linearly independent immediately gives that any $k - 1$ of $[d_1], \dots, [d_k]$ are linearly independent. \square

4. MISCELLANEOUS PROOFS

4.1. Proof of Lemma 1.3.1. It follows directly from the definitions and non-negativity of u that

$$\sum_{I \in \mathcal{B}'(h)} \int_I u(t) dt \geq \sum_{\tilde{I} \in \mathcal{B}'(f)} \int_{\tilde{I}} u(t) dt - 2|\mathcal{B}'(f)| \left(\max_{[\min f, \max f]} u \right) \cdot d_{bottle}(\mathcal{B}(f), \mathcal{B}(h)),$$

which means that we are left to prove that

$$(31) \quad \int_{\min f}^{\max f} u(t) dt - \int_{\min h}^{\max h} u(t) dt \leq 2 \left(\max_{[\min f, \max f]} u \right) \cdot d_{bottle}(\mathcal{B}(f), \mathcal{B}(h)),$$

if $\partial M = \emptyset$ and

$$(32) \quad \int_{\min f}^0 u(t) dt - \int_{\min h}^0 u(t) dt \leq \left(\max_{[\min f, \max f]} u \right) \cdot d_{bottle}(\mathcal{B}(f), \mathcal{B}(h)),$$

if $\partial M \neq \emptyset$. Let us prove (32). If $\min f \geq \min h$ the left hand side of (32) is non-positive and hence the inequality trivially holds. If $\min f < \min h$ we need to prove that

$$\int_{\min f}^{\min h} u(t) dt \leq \left(\max_{[\min f, \max f]} u \right) \cdot d_{bottle}(\mathcal{B}(f), \mathcal{B}(h)).$$

However, in every $(d_{bottle}(\mathcal{B}(f), \mathcal{B}(h)) + \varepsilon)$ -matching the infinite bar $(\min f, +\infty) \in \mathcal{B}(f)$ has to be matched with some infinite bar $(a, +\infty) \in \mathcal{B}(h)$ and since $\min h$ is the smallest of all endpoints of all infinite bars in $\mathcal{B}(h)$, we have that

$$\min h - \min f \leq a - \min f \leq d_{bottle}(\mathcal{B}(f), \mathcal{B}(h)) + \varepsilon.$$

Since the above holds for all $\varepsilon > 0$ the inequality is proven. To prove (31) one proceeds in the similar fashion, by analysing cases depending on the relative position of $\min f, \min h$ and $\max f, \max h$. This completes the proof of the lemma. \square

4.2. Proof of Lemma 1.3.5. We prove the statement in the case of M without boundary, the other case is treated the same way. Let $\mathcal{B}(f)$ and $\mathcal{B}(\tilde{f})$ be two barcodes associated to two Morse functions and denote finite bars by $I_i \in \mathcal{B}(f)$, $\tilde{I}_j \in \mathcal{B}(\tilde{f})$ where intervals are sorted by integral of u as before. Assume that $\Phi_{u,k}(\mathcal{B}(f)) \geq \Phi_{u,k}(\mathcal{B}(\tilde{f}))$ and $\mu : \mathcal{B}(f) \rightarrow \mathcal{B}(\tilde{f})$ is an ε -matching between these barcodes (we add bars of length 0 if needed and assume that μ is a genuine bijection). For every finite bar $I \in \mathcal{B}(f)$ we have that the distance between endpoints of I and $\mu(I) \in \mathcal{B}(\tilde{f})$ is less or equal than ε and hence

$$\left| \int_I u(t) dt - \int_{\mu(I)} u(t) dt \right| \leq 2\varepsilon \max u.$$

Also $|\min f - \min \tilde{f}| \leq \varepsilon$ and $|\max f - \max \tilde{f}| \leq \varepsilon$ and hence

$$\left| \int_{\min f}^{\max f} u(t) dt - \int_{\min \tilde{f}}^{\max \tilde{f}} u(t) dt \right| \leq 2\varepsilon \max u.$$

Using these estimates and the fact that the integrals of u over \tilde{I}_j decrease with j we get

$$\begin{aligned} 0 \leq \Phi_{u,k}(\mathcal{B}(f)) - \Phi_{u,k}(\mathcal{B}(\tilde{f})) &= \int_{\min f}^{\max f} u(t) dt - \int_{\min \tilde{f}}^{\max \tilde{f}} u(t) dt + \sum_{i=1}^k \int_{I_i} u(t) dt - \\ &- \sum_{j=1}^k \int_{\tilde{I}_j} u(t) dt \leq 2\varepsilon \max u + \sum_{i=1}^k \int_{I_i} u(t) dt - \sum_{i=1}^k \int_{\mu(I_i)} u(t) dt \leq \varepsilon(2k+2) \max u. \end{aligned}$$

Taking infimum over all ε -matchings finishes the proof. \square

4.3. Proof of Proposition 1.4.10. The barcode $\mathcal{B}(f_l)$ of the function

$$f_l(x_1, \dots, x_n) = \frac{\sqrt{2}}{n(2\pi)^{\frac{n}{2}}} (\sin lx_1 + \dots + \sin lx_n), \quad l \in \mathbb{N}.$$

can be computed using the Künneth formula for persistence modules. Below we briefly explain how to apply this formula and refer the reader to [PSS] for a more detailed treatment.

Given two Morse functions $f : M_1 \rightarrow \mathbb{R}$ and $h : M_2 \rightarrow \mathbb{R}$ we define another Morse function $f \oplus h : M_1 \times M_2 \rightarrow \mathbb{R}$ by setting $f \oplus h(x_1, x_2) = f(x_1) + h(x_2)$. Barcode $\mathcal{B}(f \oplus h)$ may be computed from $\mathcal{B}(f)$ and $\mathcal{B}(h)$ via the following procedure:

- An infinite bar $(a, +\infty) \in \mathcal{B}_i(f)$ and an infinite bar $(c, +\infty) \in \mathcal{B}_j(h)$ produce an infinite bar $(a+c, +\infty) \in \mathcal{B}_{i+j}(f \oplus h)$.
- An infinite bar $(a, +\infty) \in \mathcal{B}_i(f)$ and a finite bar $(c, d] \in \mathcal{B}_j(h)$ produce a finite bar $(a+c, a+d] \in \mathcal{B}_{i+j}(f \oplus h)$. The same bar is produced if $(c, d] \in \mathcal{B}_i(f)$ and $(a, +\infty) \in \mathcal{B}_j(h)$.
- A finite bar $(a, b] \in \mathcal{B}_i(f)$ and a finite bar $(c, d] \in \mathcal{B}_j(h)$ produce two finite bars $(a+c, \min\{a+d, b+c\}] \in \mathcal{B}_{i+j}(f \oplus h)$ and $(\max\{a+d, b+c\}, b+d] \in \mathcal{B}_{i+j+1}(f \oplus h)$.

In order to compute $\mathcal{B}(f_l)$ it is enough to compute the barcode of $\sin lx_1 + \dots + \sin lx_n$ and rescale. In the light of the computational procedure described above we wish to look at $\sin lx : \mathbb{S}^1 \rightarrow \mathbb{R}$ and use $\mathbb{T}^n = (\mathbb{S}^1)^n$. One readily checks that

$$\mathcal{B}_0(\sin lx) = \{(-1, +\infty), (-1, 1] \times (l-1)\}, \quad \mathcal{B}_1(\sin lx) = \{(1, +\infty)\}$$

and hence

$$\begin{aligned} \mathcal{B}_0(\sin lx_1 + \sin lx_2) &= \{(-2, +\infty), (-2, 0] \times (l^2 - 1)\}, \\ \mathcal{B}_1(\sin lx_1 + \sin lx_2) &= \{(0, +\infty) \times 2, (0, 2] \times (l^2 - 1)\}, \\ \mathcal{B}_2(\sin lx_1 + \sin lx_2) &= \{(2, +\infty)\}. \end{aligned}$$

We claim that $\mathcal{B}(\sin lx_1 + \dots + \sin lx_n)$ contains 2^n infinite bars and $\frac{1}{2}((2l)^n - 2^n)$ finite bars. To prove the claim we use induction in n . We have already checked that the statement holds for $n = 1, 2$. To complete the induction step note that, in general, if $\mathcal{B}(f)$ contains k_1 infinite and m_1 finite bars, and $\mathcal{B}(h)$ contains k_2 infinite

and m_2 finite bars, then $\mathcal{B}(f \oplus h)$ contains $k_1 k_2$ infinite and $k_1 m_2 + m_1 k_2 + 2m_1 m_2$ finite bars. Taking $k_1 = 2^n$, $m_1 = \frac{1}{2}((2l)^n - 2^n)$ and $k_2 = 2$, $m_2 = l - 1$ yields the proof.

Finally, notice that via the described procedure an infinite bar and a bar of length 2 produce a bar of length 2, as well as that two bars of length 2 produce two new bars of length 2. Since we start with $\mathcal{B}(\sin lx)$ for which all finite bars have length 2, we conclude that all finite bars in $\mathcal{B}(\sin lx_1 + \dots + \sin lx_n)$ have length 2, and thus

$$(33) \quad \Phi_1(\sin lx_1 + \dots + \sin lx_n) = 2n + (2l)^n - 2^n.$$

Rescaling (33) gives us

$$(34) \quad \Phi_1(f_l) = \frac{\sqrt{2}}{n(2\pi)^{\frac{n}{2}}} 2^n l^n + \frac{\sqrt{2}}{n(2\pi)^{\frac{n}{2}}} (2n - 2^n).$$

Since $l = \sqrt{\lambda}$ we have that

$$\Phi_1(f_l) = A_n \lambda^{\frac{n}{2}} + B_n,$$

with the constants A_n, B_n given explicitly by (34). This completes the proof of Proposition 1.4.10. \square

Acknowledgments. The authors would like to thank Yossi Azar, Allan Pinkus and Justin Solomon for useful discussions. Part of this research was accomplished while I.P. was supported by the Weston Visiting Professorship program at the Weizmann Institute of Science.

REFERENCES

- [BGW] U. Bauer, X. Ge and Y. Wang, *Measuring distance between Reeb graphs*, in Proceedings of the Thirtieth Annual Symposium on Computational geometry, 2014, ACM, 464-473.
- [BL] U. Bauer and M. Lesnick, *Induced matchings and the algebraic stability of persistence barcodes*, J. Comput. Geom., 6 (2015), no. 2, 162-191.
- [BMW] U. Bauer, E. Munch and Y. Wang, *Strong equivalence of the interleaving and functional distortion metrics for Reeb graphs*, arXiv:1412.6646.
- [BN] M. Belkin and P. Niyogi, *Laplacian eigenmaps for dimensionality reduction and data representation*, Neural Comput. 15 (2003), 1373-1396.
- [BMMPS] P. Bendich, J.S. Marron, E. Miller, A. Pieloch and S. Skwerer, *Persistent homology analysis of brain artery trees*, Ann. appl. statistics 10, (2016), 198.
- [CMW] V. Cammarota, D. Marinucci and I. Wigman, *On the distribution of the critical values of random spherical harmonics*, J. Geom. Anal. 26 (2016), no. 4, 3252-3324.
- [CS] Y. Canzani and P. Sarnak, *Topology and nesting of the zero set components of monochromatic random waves*, arXiv:1701.00034.
- [C] G. Carlsson, *Topology and data*, Bulletin Amer. Math. Soc. 46 (2009), 255-308.
- [CSGO] F. Chazal, V. de Silva, M. Glisse and S. Oudot, *The structure and stability of persistence modules*, Springer, 2016.
- [CL] R.R. Coifman and S. Lafon, *Diffusion maps*, Appl. Comput. Harmon. Anal. 21, (2006), 5-30.
- [CLRS] T.H. Cormen, C.E. Leiserson, R.L. Rivest, C. Stein, *Introduction to algorithms*, Third edition. MIT Press, Cambridge, MA, 2009
- [CH] R. Courant and D. Hilbert, *Methods of mathematical physics*, Vol. 1, Wiley, 1989.
- [E] H. Edelsbrunner, *A short course in computational geometry and topology*, Springer, 2014.
- [EH] H. Edelsbrunner and J. Harer, *Persistent homology-a survey*, Contemp. Math. 453 (2008), 257-282.
- [GW1] D. Gayet and J.-Y. Welschinger, *Universal components of random nodal sets*, Comm. Math. Phys. 347 (2016), no. 3, 777-797.

- [GW2] D. Gayet and J.-Y. Welschinger, *Betti numbers of random nodal sets of elliptic pseudo-differential operators*, to appear in Asian J. Math, arXiv:1406.0934.
- [G] R. Ghrist, *Barcodes: the persistent topology of data*, Bulletin Amer. Math. Soc. 45 (2008), 61-75.
- [JNT] D. Jakobson, N. Nadirashvili and J. Toth, *Geometric properties of eigenfunctions*, Russ. Math. Surv. 56 (2001), no. 6, 1085-1105.
- [Kr] A. S. Kronrod, *On functions of two variables*, (in Russian), Uspekhi Matem. Nauk (N.S.) 35 (1950), 24-134.
- [LL] F. Lin and D. Liu, *On the Betti numbers of level sets of solutions to elliptic equations*, Dis. Cont. Dyn. Sys. 36 (2016), no. 8, 4517-4529.
- [L1] A. Logunov, *Nodal sets of Laplace eigenfunctions: polynomial upper estimates of the Hausdorff measure*, arXiv:1605.02587.
- [L2] A. Logunov, *Nodal sets of Laplace eigenfunctions: proof of Nadirashvili's conjecture and of the lower bound in Yau's conjecture*, arXiv:1605.02589.
- [LM] A. Logunov and E. Malinnikova, *Nodal sets of Laplace eigenfunctions: estimates of the Hausdorff measure in dimension two and three*, arXiv:1605.02595.
- [NS] F. Nazarov and M. Sodin, *On the number of nodal domains of random spherical harmonics*, Amer. J. Math. 131 (2009), no. 5, 1337-1357.
- [Ni] L. Nicolaescu, *Critical sets of random linear combinations of eigenfunctions*, arXiv:1101.5990.
- [O] S.Y. Oudot, *Persistence theory: from quiver representations to data analysis*, American Mathematical Society, 2015.
- [Pi] A. Pinkus, *Negative theorems in approximation theory*, Amer. Math. Monthly, 110 (2003), no. 10, 900-911.
- [Po] G. Poliquin, *Superlevel sets and nodal extrema of Laplace-Beltrami eigenfunctions*, J. Spec. Theory, vol. 7 (2017), no. 1, 111-136.
- [PSS] L. Polterovich, E. Shelukhin and V. Stojisavljević, *Persistence modules with operators in Morse and Floer theory*, to appear in Moscow Math. J, arXiv:1703.01392.
- [PS] L. Polterovich and M. Sodin, *Nodal inequalities on surfaces*, Math. Proc. Cambridge Philos. Soc. 143 (2007), no. 2, 459-467.
- [Re] M. Reuter, *Hierarchical shape segmentation and registration via topological features of Laplace-Beltrami eigenfunctions*, Int. J. Comp. Vision 89 (2010), no. 2-3, 287-308.
- [SW] P. Sarnak and I. Wigman, *Topologies of nodal sets of random band limited functions*, in: Advances in the theory of automorphic forms and their L-functions, 351-365, Contemp. Math., 664, Amer. Math. Soc., Providence, RI, 2016.
- [SOCG] P. Skraba, M. Ovsjanikov, F. Chazal and L. Guibas, *Persistence-based segmentation of deformable shapes*, in Computer Vision and Pattern Recognition Workshops (CVPRW), 45-52, IEEE, 2010.
- [So] C. Sogge, *Concerning the L^p norm of spectral clusters for second-order elliptic operators on compact manifolds*, J. Func. Anal. 77 (1988), no. 1, 123-138.
- [U] M. Usher, *Boundary depth in Floer theory and its applications to Hamiltonian dynamics and coisotropic submanifolds*, Israel J. Math. 184 (2011), 1-57.
- [W] S. Weinberger, *What is... persistent homology?*, Notices Amer. Math. Soc. 58 (2011), 36-39.
- [Yo] Y. Yomdin, *Global bounds for the Betti numbers of regular fibers of differentiable mappings*, Topology 24 (1985), 145-152.
- [Yu] V. A. Yudin, *A multidimensional Jackson theorem*, Mat. Zametki, 20 (1976), no. 3, 439-444.
- [Z1] S. Zelditch, *Local and global properties of eigenfunctions*, Adv. Lect. Math. 7 (2008), 545-658.
- [Z2] S. Zelditch, *Eigenfunctions and nodal sets*, Surv. Differ. Geom. 18 (2013), 237-308.

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL CP 6128
SUCC CENTRE-VILLE, MONTRÉAL QC H3C 3J7, CANADA.

E-mail address: `iossif@dms.umontreal.ca`

SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, RAMAT AVIV, TEL AVIV 69978
ISRAEL

E-mail address: `polterov@post.tau.ac.il`

SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, RAMAT AVIV, TEL AVIV 69978
ISRAEL

E-mail address: `vukasin@post.tau.ac.il`