

Global Nonlinear Stability of Schwarzschild Spacetime under Polarized Perturbations

Sergiu Klainerman and Jérémie Szeftel

December 24, 2018

Abstract

We prove the nonlinear stability of the Schwarzschild spacetime under axially symmetric polarized perturbations, i.e. solutions of the Einstein vacuum equations for asymptotically flat 1 + 3 dimensional Lorentzian metrics which admit a hypersurface orthogonal spacelike Killing vectorfield with closed orbits. While building on the remarkable advances made in last 15 years on establishing quantitative linear stability, the paper introduces a series of new ideas among which we emphasize the *general covariant modulation* (GCM) procedure which allows us to construct, dynamically, the center of mass frame of the final state. The mass of the final state itself is tracked using the well known Hawking mass relative to a well adapted foliation itself connected to the center of mass frame.

Our work here is the first to prove the nonlinear stability of Schwarzschild in a restricted class of nontrivial perturbations. To a large extent, the restriction to this class of perturbations is only needed to ensure that the final state of evolution is another Schwarzschild space. We are thus confident that our procedure may apply in a more general setting.

Contents

1	INTRODUCTION	17
1.1	Bare-bones version of the main theorem	17
1.2	The Kerr family	18
1.3	Stability of Kerr	20
1.3.1	Stability of Kerr conjecture	20
1.3.2	Formal Mode Analysis	24
1.3.3	Vectorfield Method	24
1.3.4	Linear Stability of the Schwarzschild space-time	25
1.3.5	Nonlinear Stability of Schwarzschild under axially symmetric, polarized, perturbations	26
1.4	Organization	31
1.5	Acknowledgements	31
2	PRELIMINARIES	33
2.1	Axially symmetric polarized spacetimes	33
2.1.1	Axial symmetry	33
2.1.2	Z -frames	35

2.1.3	Axis of symmetry	36
2.1.4	\mathbf{Z} -polarized S - surfaces	38
2.1.5	Invariant S -foliations	57
2.1.6	Schwarzschild spacetime	62
2.2	Main Equations	63
2.2.1	Main equations for general S -foliations	63
2.2.2	Null Bianchi identities	67
2.2.3	Hawking mass	68
2.2.4	Outgoing Geodesic foliations	70
2.2.5	Additional equations	85
2.2.6	Ingoing geodesic foliation	86
2.2.7	Adapted coordinates systems	86
2.3	Perturbations of Schwarzschild and invariant quantities	94
2.3.1	Null frame transformations	94
2.3.2	Schematic notation Γ_g and Γ_b	98
2.3.3	The invariant quantity \mathfrak{q}	99
2.3.4	Several identities for \mathfrak{q}	100
2.4	Invariant wave equations	101
2.4.1	Preliminaries	102
2.4.2	Wave equations for α , $\underline{\alpha}$, and \mathfrak{q}	104
3	MAIN THEOREM	107
3.1	General covariant modulated admissible spacetimes	107

3.1.1	Initial data layer	107
3.1.2	Main Definition	110
3.1.3	Renormalized curvature components and Ricci coefficients	114
3.2	Main norms	115
3.2.1	Main norms in $^{(ext)}\mathcal{M}$	115
3.2.2	Main norms in $^{(int)}\mathcal{M}$	119
3.2.3	Combined norms	120
3.2.4	Initial layer norm	120
3.3	Main Theorem	121
3.3.1	Smallness constants	121
3.3.2	Statement of the main theorem	122
3.4	Bootstrap assumptions and first consequences	125
3.4.1	Main bootstrap assumptions	125
3.4.2	Control of the initial data	126
3.4.3	Control of averages and of the Hawking mass	126
3.4.4	Control of coordinates system	128
3.4.5	Pointwise bounds for high order derivatives	129
3.4.6	Construction of a second frame in $^{(ext)}\mathcal{M}$	130
3.5	Global null frames	132
3.5.1	Extension of frames	132
3.5.2	Construction of the first global frame	132
3.5.3	Construction of the second global frame	134

3.6	Proof of the main theorem	135
3.6.1	Main intermediate results	135
3.6.2	End of the proof of the main theorem	136
3.6.3	Conclusions	138
3.7	The General Covariant Modulation procedure	148
3.7.1	Spacetime assumptions for the GCM procedure	148
3.7.2	Deformations of surfaces	151
3.7.3	Adapted frame transformations	151
3.7.4	GCM results	152
3.7.5	Main ideas	156
3.8	Overview of the proof of Theorem M0-M8	158
3.8.1	Discussion of Theorem M0	158
3.8.2	Discussion of Theorem M1	159
3.8.3	Discussion of Theorem M2	161
3.8.4	Discussion of Theorem M3	162
3.8.5	Discussion of Theorem M4	163
3.8.6	Discussion of Theorem M5	164
3.8.7	Discussion of Theorem M6	165
3.8.8	Discussion of Theorem M7	166
3.8.9	Discussion of Theorem M8	167
3.9	Structure of the rest of the paper	171
4	CONSEQUENCES OF THE BOOTSTRAP ASSUMPTIONS	173

4.1	Proof of Theorem M0	173
4.2	Control of averages and of the Hawking mass	185
4.2.1	Proof of Lemma 3.4.1	186
4.2.2	Proof of Lemma 3.4.2	193
4.3	Control of coordinates systems	196
4.4	Pointwise bounds for high order derivatives	206
4.5	Proof of Proposition 3.4.6	212
4.6	Existence and control of the global frames	221
4.6.1	Proof of Proposition 3.5.2	221
4.6.2	Proof of Lemma 4.6.6	224
4.6.3	Proof of Proposition 3.5.5	233
5	DECAY ESTIMATES FOR \mathfrak{q} (Theorem M1)	239
5.1	Preliminaries	239
5.1.1	The foliation of \mathcal{M} by τ	241
5.1.2	Assumptions for Ricci coefficients and curvature	242
5.1.3	Structure of nonlinear terms	244
5.1.4	Main quantities	245
5.2	Proof of Theorem M1	250
5.2.1	Flux Decay Estimates for \mathfrak{q}	250
5.2.2	Proof of Theorem M1	251
5.2.3	Proof of Proposition 5.2.2	253
5.3	Improved weighted estimates	257

5.3.1	Proof of Theorem 10.2.1	258
5.3.2	Proof of Theorem 5.3.1	260
5.3.3	Proof of Theorem 5.3.2	272
5.4	Decay Estimates	277
5.4.1	First Flux Decay Estimates	278
5.4.2	Flux Decay Estimates for \check{q}	282
5.4.3	Proof of Theorem 5.2.1	284
5.4.4	Proof of Proposition 5.2.4	288
5.4.5	Proof of Proposition 5.2.5	289
6	DECAY ESTIMATES FOR α AND $\underline{\alpha}$ (Theorems M2, M3)	293
6.1	Proof of Theorem M2	293
6.1.1	A renormalized frame on ${}^{(ext)}\mathcal{M}$	293
6.1.2	A transport equation for α	294
6.1.3	Estimates for transport equations in e_3	297
6.1.4	Decay estimates for α	301
6.1.5	End of the proof of Theorem M2	308
6.2	Proof of Theorem M3	309
6.2.1	Estimate for $\underline{\alpha}$ in ${}^{(int)}\mathcal{M}$	309
6.2.2	Estimate for $\underline{\alpha}$ on Σ_*	311
6.2.3	Proof of Proposition 6.2.2	313
6.2.4	Proof of Lemma 6.2.4	317
6.2.5	Proof of Proposition 6.2.6	320

6.2.6	Proof of Lemma 6.2.8	324
7	DECAY ESTIMATES (Theorems M4, M5)	327
7.1	Preliminaries to the proof of Theorem M4	327
7.1.1	Geometric structure of Σ_*	327
7.1.2	Main Assumptions	329
7.1.3	Basic Lemmas	332
7.1.4	Main equations	335
7.1.5	Equations involving \mathfrak{q}	335
7.1.6	Additional equations	339
7.2	Structure of the proof of Theorem M4	343
7.3	Decay estimates on the last slice Σ_*	346
7.3.1	Preliminaries	346
7.3.2	Differential identities involving GCM conditions on Σ_*	349
7.3.3	Control of the flux of some quantities on Σ_*	350
7.3.4	Estimates for the $\ell = 1$ modes on Σ_*	360
7.3.5	Decay of Ricci and curvature components on Σ_*	370
7.4	Control in $^{(ext)}\mathcal{M}$, Part I	376
7.4.1	Preliminaries	376
7.4.2	Proposition 7.4.5	379
7.4.3	Estimates for $\check{\kappa}, \check{\mu}$ in $^{(ext)}\mathcal{M}$	380
7.4.4	Estimates for the $\ell = 1$ modes in $^{(ext)}\mathcal{M}$	381
7.4.5	Completion of the Proof of Proposition 7.4.5	384

7.5	Control in $^{(ext)}\mathcal{M}$, Part II	388
7.5.1	Estimate for η	389
7.5.2	Crucial Lemmas	390
7.5.3	Proof of Proposition 7.5.1, Part I	399
7.5.4	Proof of Proposition 7.5.1, Part II	403
7.6	Conclusion of the Proof of Theorem M4	407
7.7	Proof of Theorem M5	410
8	INITIALIZATION AND EXTENSION (Theorems M6, M7, M8)	417
8.1	Proof of Theorem M6	417
8.2	Proof of Theorem M7	420
8.3	Proof of Theorem M8	432
8.3.1	Main norms	434
8.3.2	Control of the global frame	436
8.3.3	Iterative procedure	438
8.3.4	End of the proof of Theorem M8	440
8.4	Proof of Proposition 8.3.5	443
8.4.1	A wave equation for $\tilde{\rho}$	443
8.4.2	Control of $\square_{\mathbf{g}}(r)$	445
8.4.3	End of the proof of Proposition 8.3.5	451
8.5	Proof of Proposition 8.3.6	453
8.5.1	A wave equations for $\alpha + \Upsilon^2 \underline{\alpha}$	453
8.5.2	End of the proof of Proposition 8.3.6	462

8.6	Proof of Proposition 8.3.7	464
8.6.1	Control of α and $\Upsilon^2 \underline{\alpha}$	464
8.6.2	Control of $\underline{\alpha}$	466
8.6.3	End of the proof of Proposition 8.3.7	469
8.7	Proof of Proposition 8.3.8	470
8.7.1	r -weighted divergence identities for Bianchi pairs	471
8.7.2	End of the proof of Proposition 8.3.8	480
8.8	Proof of Proposition 8.3.9	486
8.8.1	Proof of Proposition 8.8.1	488
8.8.2	Weighted estimates for transport equations along e_4 in $^{(ext)}\mathcal{M}$. . .	498
8.8.3	Several identities	505
8.8.4	Proof of Proposition 8.8.2	509
8.8.5	Proof of Proposition 8.8.3	516
8.9	Proof of Proposition 8.3.10	525
8.9.1	Weighted estimates for transport equations along e_3 in $^{(int)}\mathcal{M}$. . .	526
8.9.2	Proof of Proposition 8.9.1	528
8.10	Proof of Proposition 8.3.11	532
9	GCM PROCEDURE	533
9.1	Preliminaries	533
9.1.1	Main assumptions	535
9.1.2	Elliptic Hodge lemma	536
9.2	Deformations of S surfaces	537

9.2.1	Deformations	537
9.2.2	Pull-back map	538
9.2.3	Comparison of norms between deformations	541
9.2.4	Adapted frame transformations	544
9.3	Frame transformations	548
9.3.1	Main GCM Equations	557
9.3.2	Equation for the average of a	563
9.3.3	Transversality conditions	564
9.3.4	Equation for the $\ell = 1$ mode of $d^{\mathbf{S},*}a$	565
9.4	Existence of GCM spheres	566
9.4.1	A priori estimates	570
9.4.2	Comparison of the Hawking mass	572
9.4.3	Iteration procedure for Theorem 9.4.2	573
9.4.4	Boundedness of the iterates	576
9.4.5	Convergence of the Iterates	577
9.5	Proof of Proposition 9.4.4	581
9.6	Proof of Proposition 9.4.9	594
9.6.1	Pull-back of the main equations	594
9.6.2	Basic Lemmas	596
9.6.3	Proof of the estimates (9.6.5), (9.6.6)	605
9.7	A Corollary to Theorem 9.4.2	609
9.8	Construction of GCM hypersurfaces	614

<i>CONTENTS</i>	13
9.8.1 Definition of Σ_0	617
9.8.2 Extrinsic properties of Σ_0	619
9.8.3 Construction of Σ_0	632
10 REGGE-WHEELER TYPE EQUATIONS	649
10.1 Basic Morawetz Estimates	650
10.1.1 Structure of the proof of Theorem 10.1.1	651
10.1.2 A simplified set of assumptions	652
10.1.3 Functions depending on m and r	652
10.1.4 Deformation tensors of the vectorfields R, T, X	653
10.1.5 Basic Integral Identities	658
10.1.6 Main Morawetz Identity	660
10.1.7 A first estimate	664
10.1.8 Improved lower bound in $^{(ext)}\mathcal{M}$	670
10.1.9 Cut-off Correction in $^{(int)}\mathcal{M}$	678
10.1.10 The red shift vectorfield	685
10.1.11 Combined Estimate	689
10.1.12 Lower bounds for \mathcal{Q}	696
10.1.13 First Morawetz Estimate	698
10.1.14 Analysis of the error term \mathcal{E}_ϵ	705
10.1.15 Proof of Theorem 10.1.1	706
10.2 Dafermos-Rodnianski r^p - weighted estimates	710
10.2.1 Vectorfield $X = f(r)e_4$	713

10.2.2	Energy densities for $X = f(r)e_4$	713
10.2.3	Proof of Theorem 10.2.1	723
10.3	Higher Weighted Estimates	731
10.3.1	Wave equation for $\check{\psi}$	731
10.3.2	The r^p weighted estimates for $\check{\psi}$	732
10.4	Higher Derivative Estimates	738
10.4.1	Basic assumptions	738
10.4.2	Strategy for recovering higher order derivatives	738
10.4.3	Commutation formulas with the wave equation	739
10.4.4	Some weighted estimates for wave equations	753
10.4.5	Proof of Theorem 5.3.4	758
10.4.6	Proof of Theorem 5.3.5	765
10.5	More weighted estimates for wave equations	770
A	APPENDIX TO CHAPTER 2	777
A.1	Proof of Proposition 2.2.9	777
A.2	Proof of Proposition 2.2.16	780
A.3	Proof of Lemma 2.2.17	783
A.4	Proof of Proposition 2.2.18	787
A.5	Proof of Proposition 2.2.19	792
A.6	Proof of Proposition 2.3.4	796
A.7	Proof of Lemma 2.3.6	810
A.8	Proof of Corollary 2.3.7	813

<i>CONTENTS</i>	15
A.9 Proof of Lemma 2.3.5	815
A.10 Proof of Proposition 2.3.13	817
A.11 Proof of Proposition 2.3.14	820
A.12 Proof of the Teukolsky-Starobinski identity	825
A.13 Proof of Proposition 2.4.6	833
A.14 Proof of Theorem 2.4.7	837
A.14.1 The Teukolsky equation for α	841
A.14.2 Commutation lemmas	843
A.14.3 Main commutation	850
A.14.4 Proof of Theorem 2.4.7	859
B APPENDIX TO CHAPTER 8	863
B.1 Proof of Proposition 8.4.1	863
C APPENDIX TO CHAPTER 9	871
C.1 Proof of Lemma 9.2.6	871
D APPENDIX TO CHAPTER 10	885
D.1 Horizontal S -tensors	885
D.1.1 Mixed tensors	886
D.1.2 Invariant Lagrangian	887
D.1.3 Comparison of the Lagrangians	887
D.1.4 Energy-Momentum tensor	889
D.2 Standard Calculation	890

D.3	Vectorfield X_f	891
D.4	Proof of Proposition 10.3.1	894

Chapter 1

INTRODUCTION

1.1 Bare-bones version of the main theorem

The goal of the book is to prove the nonlinear stability of the Schwarzschild spacetime under axially symmetric polarized perturbations, i.e. solutions of the Einstein vacuum equations (1.2.1) for asymptotically flat 1+3 dimensional Lorentzian metrics which admit a hypersurface orthogonal spacelike Killing vectorfield \mathbf{Z} with closed orbits. Recall that the Schwarzschild metric \mathbf{g}_m of mass $m > 0$ is a stationary, spherically symmetric solution of (1.2.1), which takes the form, in standard coordinates,

$$\mathbf{g}_m = - \left(1 - \frac{2m}{r}\right) dt^2 + \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\varphi^2).$$

This class of perturbations allows us to restrict our analysis to the case when the final state of evolution is itself a Schwarzschild spacetime. This is not the case in general, as a typical perturbation of Schwarzschild may approach a member of the Kerr family with small angular momentum.

The simplest version of our main theorem can be stated as follows.

Theorem 1.1.1 (Main Theorem (first version)). *The future globally hyperbolic development of an axially symmetric, polarized¹, asymptotically flat initial data set, sufficiently close (in a specified topology) to a Schwarzschild initial data set of mass $m_0 > 0$, has a complete future null infinity \mathcal{I}^+ and converges in its causal past $\mathcal{J}^{-1}(\mathcal{I}^+)$ to another nearby Schwarzschild solution of mass m_∞ close to m_0 .*

¹See section 2.1.1 for a precise definition of axial symmetry and polarization. This property is preserved by the Einstein equations, i.e. if the data is axially symmetric, polarized, so is its development.

Our theorem is an important step in the long standing effort to prove the full nonlinear stability of Kerr spacetimes $\mathcal{K}(a, m)$, in the sub-extremal regime $|a| < m$. We give a succinct review below of some of the most important results which have been obtained so far in this direction.

1.2 The Kerr family

Consider solutions to the Einstein vacuum equations (EVE),

$$\mathbf{R}_{\alpha\beta} = 0 \tag{1.2.1}$$

with $\mathbf{R}_{\alpha\beta}$ the Ricci curvature of a four dimensional, Lorentzian manifold $(\mathcal{M}, \mathbf{g})$. Solutions of the equations are invariant under general diffeomorphisms of $\Phi : \mathcal{M} \rightarrow \mathcal{M}$, i.e. if \mathbf{g} verifies EVE so does its pull back $\Phi^*\mathbf{g}$. We recall that an initial data set $(\Sigma_{(0)}, g_{(0)}, k_{(0)})$ consists of a 3 dimensional manifold $\Sigma_{(0)}$ together with a complete Riemannian metric $g_{(0)}$ and a symmetric 2-tensor $k_{(0)}$ which verify a well known set of constraint equations (see for instance the introduction in [17]). A Cauchy development of an initial data set is a globally hyperbolic space-time $(\mathcal{M}, \mathbf{g})$, verifying EVE together with an embedding $i : \Sigma_{(0)} \rightarrow \mathcal{M}$ such that $i_*(g_{(0)}), i_*(k_{(0)})$ are the first and second fundamental forms of $i(\Sigma_{(0)})$ in \mathcal{M} . A well known foundational result in GR associates a unique maximal, global hyperbolic, future development to all sufficiently regular initial data sets, see [13], [14]². We further restrict the discussion to asymptotically flat initial data sets, i.e. assume that outside a sufficiently large compact set K , $\Sigma_{(0)} \setminus K$ is diffeomorphic to the complement of the unit ball in \mathbb{R}^3 and admits a system of coordinates in which $g_{(0)}$ is asymptotically euclidean and $k_{(0)}$ vanishes at an appropriate order at infinity.

EVE admits a remarkable two parameter family of explicit solutions, the Kerr spacetimes $\mathcal{K}(a, m)$, $0 \leq a \leq m$, which are stationary, axisymmetric and asymptotically flat. In the usual Boyer-Lindquist coordinates they take the form,

$$\mathbf{g}_{a,m} = -\frac{q^2 \Delta}{\Sigma^2} (dt)^2 + \frac{\Sigma^2 (\sin \theta)^2}{q^2} \left(d\varphi - \frac{2amr}{\Sigma^2} dt \right)^2 + \frac{q^2}{\Delta} (dr)^2 + q^2 (d\theta)^2, \tag{1.2.2}$$

where

$$\begin{cases} \Delta = r^2 + a^2 - 2mr, \\ q^2 = r^2 + a^2 (\cos \theta)^2, \\ \Sigma^2 = (r^2 + a^2) q^2 + 2mra^2 (\sin \theta)^2 = (r^2 + a^2)^2 - a^2 (\sin \theta)^2 \Delta. \end{cases} \tag{1.2.3}$$

²See also [50] for a modern treatment.

Among them one distinguishes the Schwarzschild family of spherically symmetric solutions, of mass $m > 0$,

$$\mathbf{g}_m = - \left(1 - \frac{2m}{r}\right) dt^2 + \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2 d\sigma_{\mathbb{S}^2}. \quad (1.2.4)$$

Though the metric seems singular at $r = 2m$ ($r = r_+$, the largest root of $\Delta(r) = 0$, in the case of Kerr) it turns out that one can glue together two regions $r > 2m$ and two regions $r < 2m$ of the Schwarzschild metric to obtain a metric which is smooth along the null hypersurface $\mathcal{E} = \{r = 2m\}$ called the Schwarzschild event horizon. The portion of $r < 2m$ to the future of the hypersurface $t = 0$ is a black hole whose future boundary $r = 0$ is singular. The region $r > 2m$, free of singularities, is called the domain of outer communication. The more general family of Kerr solutions, which are both stationary and axially symmetric, possesses (in addition to well defined event horizons, black holes and domains of outer communication) Cauchy horizons ($r = r_-$ for the smallest root $r = r_-$ of $\Delta(r) = 0$) inside the black hole region across which predictability seems to fail³. Again, one can easily check, from the precise nature of the Kerr metric, that the region outside the event horizon, i.e. outside the Kerr black hole, is free of singularities⁴. Note that the Kerr spacetimes $\mathcal{K}(a, m)$ possess two Killing vectorfields; the stationary vectorfield $\mathbf{T} = \partial_t$, which is timelike in the asymptotic region, away from the horizon, and the axial symmetric Killing vectorfield $\mathbf{Z} = \partial_\varphi$. In the particular case of Schwarzschild, \mathbf{T} is also orthogonal to the hypersurfaces $t = \text{const}$.

Here are some other important properties of the Kerr family.

- The Kerr solution has a remarkable algebraic feature, encoded in the so called Petrov type D property, according to which it admits, at every point a pair of null vectors (l, \underline{l}) , normalized by the condition $\mathbf{g}(l, \underline{l}) = -2$, called principal null vectors, such that all components of the Riemann curvature tensor vanish identically except for the two independent components

$$\mathbf{R}(l, \underline{l}, l, \underline{l}), \quad \star\mathbf{R}(l, \underline{l}, l, \underline{l})$$

with $\star\mathbf{R}$ the Hodge dual of \mathbf{R} .

- In addition to the symmetries provided by the Killing vectorfields \mathbf{T} and \mathbf{Z} , the Kerr solution possesses a nontrivial Killing tensor i.e. a symmetric 2-covariant tensor \mathbf{C} (the Carter tensor) verifying,

$$\mathbf{D}_{(\alpha} \mathbf{C}_{\beta\gamma)} = 0.$$

³I.e. various smooth extensions are possible.

⁴The generalization of this observation to all but an exceptional set of initial data is the celebrated *weak cosmic censorship conjecture* of Penrose.

- The Kerr family is distinguished among all stationary solutions of EVE by the vanishing of a four tensor called the Mars-Simon tensor, see [42].

1.3 Stability of Kerr

1.3.1 Stability of Kerr conjecture

The nonlinear stability of the Kerr family is one of the most pressing issues in mathematical GR today. Roughly, the problem is to show that all spacetime developments of initial data sets, sufficiently close to the initial data set of a Kerr spacetime, behave in the large like a (typically another) Kerr solution. This is not only a deep mathematical question but one with serious astrophysical implications. Indeed, if the Kerr family would be unstable under perturbations, black holes would be nothing more than mathematical artifacts. Here is a more precise formulation of the conjecture.

Conjecture (Stability of Kerr conjecture). *Vacuum initial data sets, sufficiently close to Kerr initial data, have a maximal development with complete future null infinity⁵ and with domain of outer communication which approaches (globally) a nearby Kerr solution.*

So far, the only space-time for which full nonlinear stability has been established is the Minkowski space, corresponding to the particular case $a = m = 0$. The result was first proved in [17], see also [37], [39], [6] and [28].

Theorem 1.3.1 (Global stability of Minkowski). *Any asymptotically flat initial data set, which is sufficiently close to the trivial one, has a regular, complete, maximal development⁶.*

Here are, very schematically, the main ideas in the proof of stability of Minkowski space.

- (I) Perturbations radiate and decay *sufficiently fast* (just fast enough!) to insure convergence.
- (II) Interpret the Bianchi identities as a Maxwell like system. This is an effective, *invariant*, way to treat the hyperbolic character of the equations.

⁵This means, roughly, that observers which are far away from the black hole may live forever.

⁶The complete result in [17] also provides very precise information about the decay of the curvature tensor along null and timelike directions as well as many other geometric information concerning the causal structure of the corresponding spacetime. Of particular interest are *peeling properties* i.e. the precise decay rates of various components of the curvature tensor along future null geodesics.

(III) Rely on four important PDE advances of late last century:

- (i) Vectorfield approach to get decay based on *approximate* Killing and conformal Killing symmetries of the equations, see [33], [34], [35], [16].
- (ii) Generalized energy estimates using both the Bianchi identities and the approximate Killing and conformal Killing vector fields.
- (iii) The *null condition* identifies the deep mechanism for nonlinear stability, i.e. the specific structure of the nonlinear terms enables stability despite the slow decay rate of the perturbations, see [32], [34], [15].
- (iv) Complex boot-strap argument according to which one makes educated assumptions about the behavior of the space-time and then proceeds to show that they are in fact satisfied. This amounts to a *conceptual linearization*, i.e. a method by which the equations become, essentially, linear without actually linearizing them.

There are three, related, major obstacles in passing from the stability of Minkowski to that of the Kerr family.

1. The first can be understood in the general framework of nonlinear hyperbolic or dispersive equations. Given a nonlinear equation $\mathcal{N}[\phi] = 0$ and a stationary solution ϕ_0 we have two notions of stability, *orbital stability*, according to which small perturbations of ϕ_0 lead to solutions ϕ which remain close, in some norm (typically L^2 based) for all time, and *asymptotical stability*, according to which the perturbed solutions converge, as $t \rightarrow \infty$, to a nearby stationary solution. Note that the second notion is far stronger, and much more precise, than the first and that orbital stability can only be established (without appealing to the the stronger version) only for equations with very weak nonlinearities. For quasilinear equations, such as the Einstein field equations, a proof of stability requires, necessarily, a proof of asymptotic stability. This must then be based on a detailed understanding of the decay properties of the linearized⁷ equations.

One is thus led to study the linearized equations $\mathcal{N}'[\phi_0]\psi = 0$, with $\mathcal{N}'[\phi_0]$ the Fréchet derivative of \mathcal{N} at ϕ_0 , which, in many important cases, are hyperbolic⁸

⁷It is irrelevant whether a specific linearization procedure needs to be implemented; what is important here is to identify the linear mechanism for decay, such as the Maxwell system in the case of the stability of Minkowski space mentioned above.

⁸In the case of EVE the linearized equations are linear hyperbolic only after we mod out the linearized version of general coordinate transformations.

systems with variable coefficients that typically present instabilities. In the exceptional situation, when nonlinear stability can ultimately be established, one can tie all the instability modes of the linearized system to two properties of the nonlinear equation:

- (a) The presence of a continuous^{9,10} family of other stationary solutions of $\mathcal{N}[\phi] = 0$ near ϕ_0 .
- (b) The presence of a continuous family of diffeomorphisms¹¹ of the background manifold which map, by pull back, solutions to solutions.

For a typical stationary solution ϕ_0 , both properties exist and generate nontrivial solutions of the linearized equation $\mathcal{N}'[\phi_0]\psi = 0$. In the case of relatively simple scalar nonlinear equations, where the symmetry group of the equation is small, an effective strategy of dealing with this problem (known under the name of modulation theory) has been developed, see for example [44], [46]. In the case of the Einstein equations this problem is compounded by the large invariance group of the equations, i.e. all diffeomorphisms of the spacetime manifold. To deal with both problems and establish stability one has to

- Track the parameters (a_f, m_f) of the final Kerr spacetime.
 - Track the coordinate system (gauge condition) relative to which we have decay for all linearized quantities. Such a coordinate system cannot be imposed a-priori, it has to emerge dynamically in the construction of the spacetime.
2. As described earlier, the fundamental insight in the stability of the Minkowski space was that we can treat the Bianchi identities as a Maxwell system in a slightly perturbed Minkowski space by using the vectorfield method. This cannot work for perturbations of Kerr due to the fact that some of the null components of the curvature tensor¹² are non-trivial in Kerr.
 3. Even if we can establish a useful version of linearization (i.e. one which addresses the above mentioned problems), there are still major obstacles in understanding their decay properties. Indeed, when one considers the simplest, relevant, linear equation on a fixed Kerr background, i.e. the wave equation $\square_{\mathbf{g}}\psi = 0$ (often referred to as the *poor's man linearization* of EVE), one encounters serious difficulties even to prove the boundedness of solutions for the most reasonable, smooth, compactly supported, data. Below is a very short description of these.

⁹In the case of the stability of Kerr we have a 2 parameter family of solutions $\mathcal{K}(a, m)$.

¹⁰This is responsible of the fact that a small perturbation of the fixed stationary solution ϕ_0 may not converge to ϕ_0 but to another nearby stationary solution.

¹¹In the case of EVE, any diffeomorphism has that property.

¹²With respect to the so called principal null directions.

- *The problem of trapped null geodesics.* This concerns the existence of null geodesics¹³ neither crossing the event horizon nor escaping to null infinity, along which solutions can concentrate for arbitrary long times. This leads to degenerate energy estimates which require a very delicate analysis.
 - *The trapping properties of the horizon.* The horizon itself is ruled by null geodesics, which do not communicate with null infinity and can thus concentrate energy. This problem was solved by understanding the so called red-shift effect associated to the event horizon, which more than counteracts this type of trapping.
 - *The problem of superradiance.* This is essentially the failure of the stationary Killing field $\mathbf{T} = \partial_t$ to be everywhere timelike in the domain of outer communications and, thus, the failure of the associated conserved energy to be positive. Note that this problem is absent in Schwarzschild and, in general, for axially symmetric solutions.
 - *Superposition problem.* This is the problem of combining the estimates in the near region, close to the horizon, (including the ergoregion and trapping) with estimates in the asymptotic region, where the spacetime looks Minkowskian.
4. The full linearized system of EVE around Kerr, usually referred to as the linearized gravity system (LGS), whatever its formulation, presents far more difficulties beyond those mentioned above concerning the poor man's linear scalar wave equation on Kerr, see the discussion below.

Historically, two versions of LGS have been considered.

- (a) At the level of the metric itself, i.e. if \mathbf{G} denotes the Einstein tensor, $\mathbf{G}_{\alpha\beta} = \mathbf{R}_{\alpha\beta} - \frac{1}{2}\mathbf{R}\mathbf{g}_{\alpha\beta}$,

$$\mathbf{G}'(\mathbf{g}_0) \delta\mathbf{g} = 0. \tag{1.3.1}$$

- (b) Via the Newman-Penrose (NP) formalism, based on null frames.

In what follows we review the main known results concerning solutions to the linearized equations on a Kerr background.

¹³In the Schwarzschild case, these geodesics are located on the so-called photon sphere $r = 3m$.

1.3.2 Formal Mode Analysis

The first important results concerning both items (3) and (4) above were obtained by physicists based on the classical method of separation of variables and formal mode analysis. In the particular case where \mathbf{g}_0 is the Schwarzschild metric, the linearized equations (1.3.1) can be formally decomposed into modes, by associating t-derivatives with multiplication by $i\omega$ and angular derivatives with multiplication by l , i.e. the eigenvalues of the spherical laplacian. A similar decomposition, using oblate spheroidal harmonics, can be done in Kerr. The formal study of fixed modes from the point of view of *metric perturbations* as in (1.3.1) was initiated by Regge-Wheeler [49] who discovered the master Regge-Wheeler equation for odd-parity perturbations. This study was completed by Vishveshwara [56] and Zerilli [60]. A gauge-invariant formulation of *metric perturbations* was then given by Moncrief [47]. An alternative approach via the Newman-Penrose (NP) formalism was first undertaken by Bardeen-Press [5]. This latter type of analysis was later extended to the Kerr family by Teukolsky [55] who made the important discovery that the extreme curvature components, relative to a principal null frame, satisfy decoupled, separable, wave equations. These extreme curvature components also turn out to be *gauge invariant* in the sense that small perturbations of the frame lead to quadratic errors in their expression. The full extent of what could be done by mode analysis, in both approaches, can be found in Chandrasekhar's book [11]. Chandrasekhar also introduced (see [12]) a transformation theory relating the two approaches. More precisely, he exhibits a transformation which connects the Teukolsky equations to the Regge-Wheeler one. This transformation was further elucidated and extended by R. Wald [57] and recently by Aksteiner and al [2]. The full mode stability, i.e. lack of exponentially growing modes, for the Teukolsky equation on Kerr is due to Whiting [59] (see also [51] for a stronger quantitative version).

1.3.3 Vectorfield Method

Note that mode stability is far from establishing even boundedness of solutions to the linearized equations. To achieve that and, in addition, to derive realistic decay estimates one needs an entirely different approach based on a far reaching extension of the classical vectorfield method¹⁴ used in the proof of the nonlinear stability of Minkowski [17]. The new vectorfield method compensates for the lack of enough Killing and conformal Killing vectorfields on a Schwarzschild or Kerr background by introducing new vectorfields whose deformation tensors have coercive properties in different regions of spacetime, not nec-

¹⁴Method based on the symmetries of Minkowski space to derive uniform, robust, decay for nonlinear wave equations, see [33], [34], [35], [16].

essarily causal. The new method has emerged in the last 15 years in connection to the study of boundedness and decay for the scalar wave equation in $Kerr(a, m)$,

$$\square_{\mathbf{g}_{a,m}} \psi = 0. \quad (1.3.2)$$

The starting and most demanding part of the new method is the derivation of a global, simultaneous, *Energy–Spacetime Morawetz* estimate which degenerates in the trapping region. This task is somewhat easier in Schwarzschild, or for axially symmetric solutions in Kerr, where the trapping region is restricted to a smooth hypersurface. The first such estimates, in Schwarzschild, were proved by Blue and Soffer in [7], [8] followed by a long sequence of further improvements in [10], [19], [45] etc. See also [31] and [53] for a vectorfield method treatment of the axially symmetric case in Kerr with applications to nonlinear equations. In the absence of axial symmetry the derivation of an Energy–Morawetz estimate in $Kerr(a, m)$, $|a/m| \ll 1$ requires a more refined analysis involving either Fourier decompositions, see [21], [54], or a systematic use of the second order Carter operator, see [3]. The derivation of such an estimate in the full sub-extremal case $|a| < m$ is even more subtle and was recently achieved by Dafermos, Rodnianski and Shlapentokh-Rothman [24] by combining mode decomposition with the vectorfield method.

Once an Energy–Morawetz estimate is established one can commute with the time translation vectorfield and the so called *Red Shift* vectorfield, first introduced in [19], to derive uniform bounds for solutions. The most efficient way to also get decay, and solve the *superposition problem*, is due to Dafermos and Rodnianski, see [20], based on the presence of a family of *r^p-weighted*, quasi-conformal vectorfields defined in the far *r* region of spacetime¹⁵.

1.3.4 Linear Stability of the Schwarzschild space-time

A first quantitative (i.e. which provides precise decay estimates) proof of the linear stability of Schwarzschild spacetime has recently been established¹⁶ by Dafermos, Holzegel and Rodnianski in [22], via the NP formalism (expressed in a double null foliation¹⁷). It is important to note that while the Teukolsky equation (in the NP formalism) is separable, and thus amenable to mode analysis, it is not Lagrangian and thus cannot be treated

¹⁵These replace the scaling and inverted time translation vectorfields used in [33] or their corresponding deformations used in [17]. A recent improvement of the method, relevant to our work here, allowing one to derive higher order decay can be found in [4].

¹⁶A somewhat weaker version of linear stability of Schwarzschild was subsequently proved in [30] by using the original, direct, Regge–Wheeler, Zerilli approach combined with the vectorfield method and adapted gauge choices.

¹⁷This is possible in Schwarzschild where the principal null directions are integrable.

by direct energy type estimates. To overcome this difficulty [22] relies on a new physical space version of the Chandrasekhar transformation [12], which takes solutions of the Teukolsky equations to solutions of Regge-Wheeler, which is manifestly both Lagrangian and coercive. After quantitative decay has been established for this latter equation, based on the new vectorfield method, the physical space form of the transformation allows one to derive quantitative decay for solutions of the original Teukolsky equation. Once decay estimates for the Teukolsky equation have been established, the remaining work in [22] is to bound all other curvature and Ricci coefficients associated to the double null foliation. This last step requires carefully chosen gauge conditions along the event horizon of the fixed Schwarzschild background. This final gauge is itself then quantitatively bounded in terms of the initial data, giving thus a comprehensive statement of linear stability.

1.3.5 Nonlinear Stability of Schwarzschild under axially symmetric, polarized, perturbations

In the passage from linear to nonlinear stability of Schwarzschild one has to overcome major new difficulties. Some are similar to those encountered in the stability of Minkowski [17] such as,

1. Need of an appropriate geometric setting which takes into account the decay and peeling properties of the curvature. In [17] this was achieved with the help of the foliation of the perturbed spacetime given by two optical functions $^{(int)}u$ and $^{(ext)}u$ and a maximal time function t . The exterior optical function $^{(ext)}u$, which was initialized at infinity, was essential to derive the decay and peeling properties along null directions while $^{(int)}u$, initialized on a timelike axis, was responsible for covering the interior, non-radiative, back scattering, decay.
2. The peeling and decay estimates have to be derived by some version of the geometric vectorfield method which relates decay to generalized energy type estimates.
3. The peeling and decay estimates mentioned above should be sufficiently strong to be able to deal with the error terms generated by the vectorfield method. For this to happen, the error terms need to exhibit an appropriate null structure.

The new main difficulties are as follows:

1. One needs a procedure which allows to take into account the change of mass and detect its final value. Note also that we need to restrict the nature of the per-

turbations to insure that the final state of a perturbation of Schwarzschild is still Schwarzschild.

2. While in the stability of Minkowski space all components of the curvature tensor were expected to approach zero, this is no longer true. Indeed, the middle curvature component (relative to an adapted null frame), ought to converge to its respective value in the final Schwarzschild spacetime. This statement is unfortunately hard to quantify since that value depends both on the final mass and on the corresponding Schwarzschild coordinates. Moreover, some of the other curvature components, which are expected to converge to zero, are also ill defined since a small change of the null frame can produce small linear distortion to the basic equation which these curvature components verify. Note that this difficulty was absent in the stability of the Minkowski space where small changes in the frame produce only quadratic errors.
3. The classical vector-field method used in the nonlinear stability of Minkowski space was based on the construction, together with the spacetime, of an adequate family of approximate Killing and conformal Killing vectorfields which mimic the role played by the corresponding vectorfields in Minkowski space in establishing uniform decay estimates. The Schwarzschild space however has a much more limited set of Killing vectorfields and no useful conformal Killing ones. As mentioned above, this problem appears already in the analysis of the standard scalar linear wave equation in Schwarzschild.
4. As in the stability of the Minkowski space, one needs to make gauge conditions to insure that we are measuring decay relative to an appropriate center of mass frame. Yet, as we saw above, it is no longer true that small perturbations of the null frame produce only quadratic errors for the curvature, as was the case in the stability of Minkowski space. In fact, the center of mass frame of the perturbed black hole continuously changes in response to incoming radiation. This, the so called *recoil problem*, does not occur in linear theory.

Here is a very short summary of how we solve these new challenges in our work.


1. We resolve the first difficulty by restricting our analysis to axially symmetric, polarized perturbations and by tracking the mass using a quantity, called the quasi-local Hawking mass, for which we derive simple propagation equations which establish monotonicity of the mass up to errors which are quadratic with respect to the perturbations.
2. We resolve the second difficulty by making use of the fact that the extreme components of the curvature are, up to quadratic terms, invariant under null frame

transformations. As in [22], we also make use of a transformation, similar to that of Chandrasekhar mentioned above, which maps the extreme components of the curvature to a new quantity \mathfrak{q} , defined up to quadratic errors, that verifies a Regge-Wheeler type equation. Once we manage to control \mathfrak{q} , i.e. to derive quantitative decay estimates for it, we can also control, in principle¹⁸, the two extreme curvature invariants α and $\underline{\alpha}$, the first by inverting the Chandrasekhar transformation and the second by using a variant of the Teukolsky- Starobinski identities. One is then left with the arduous task of recovering¹⁹ all other null components of the curvature tensor and all connection coefficients.

3. The third difficulties manifests itself in the most sensitive part of the entire argument, i.e. in the task of deriving quantitative decay estimates for \mathfrak{q} by making use of the Regge-Wheeler type equation it verifies. To do this we rely on the new vectorfield method as outlined in subsection 1.3.3 above. The main new difficulties are:
 - (i) The vectorfield method introduces new error terms, not present in linear theory. To estimate these terms we need precise decay informations, off the final Schwarzschild space, for all connection coefficients and curvature of the perturbation.
 - (ii) The most difficult terms are those due to the quadratic errors made in the derivation of the Regge-Wheeler equation for \mathfrak{q} . As in the proof the stability of the Minkowski space the precise rates of decay for various curvature and connection coefficients, i.e. the peeling properties of the perturbation, and the the precise structure of these error terms is of fundamental importance.
4. We solve the fourth and most important new difficulty by a procedure we call *General Covariant Modulation* (GCM). This procedure, which takes advantage of the full covariance of the Einstein equations, allows us to construct the perturbed spacetime by a continuity argument involving finite GCM admissible spacetimes \mathcal{M} as represented in Figure 1.1. The past boundaries $\underline{\mathcal{C}}_1 \cup \mathcal{C}_1$ are incoming and outgoing null hypersurfaces on which the initial perturbation is prescribed. The future boundaries consists of the union $\mathcal{A} \cup \underline{\mathcal{C}}_* \cup \mathcal{C}_* \cup \Sigma_*$ where \mathcal{A} and Σ_* are spacelike, $\underline{\mathcal{C}}_*$ is incoming null, \mathcal{C}_* outgoing null. The boundary \mathcal{A} is chosen so that, in the limit when \mathcal{M} converges to the final state, is included in the perturbed black hole. The spacelike boundary Σ_* plays a fundamental role in our construction as seen below. The spacetime \mathcal{M} also contains a timelike hypersurface \mathcal{T} which divides \mathcal{M} into an exterior region we call $^{(ext)}\mathcal{M}$ and an interior one $^{(int)}\mathcal{M}$. We say that \mathcal{M} is a GCM admissible spacetime if it verifies the following properties.

¹⁸Provided that one can deal with the nonlinear terms.

¹⁹In the linear setting this was partially achieved in [23].

Figure 1.1: The GCM admissible space-time \mathcal{M}

- (i) The far region $^{(ext)}\mathcal{M}$ is foliated by a geodesic foliation induced by an outgoing optical function u initialized on Σ_*
 - (ii) The near region $^{(int)}\mathcal{M}$ is foliated by a geodesic foliation induced by an incoming optical function \underline{u} initialized at \mathcal{T} such that its level sets on \mathcal{T} coincide with those of u .
 - (iii) The foliation induced on Σ_* is such that specific geometric quantities take Schwarzschildian values. We refer to these as GCM conditions. These conditions are dynamically reset in the continuation process on which our proof is based.
 - (iv) The area radius $r(u)$ of the spheres of constant u along Σ_* is far greater than the corresponding value of u . This condition allows us to simplify somewhat the null structure and Bianchi equations induced on Σ_* and corresponds to the expectation that the spacelike hypersurfaces Σ_* converges to the null infinity of the final state of the perturbation.
5. The GCM conditions together with the control derived on \mathfrak{q} , α and $\underline{\alpha}$ mentioned earlier allows us to control all null connection and curvature coefficients along on Σ_* , i.e. to derive appropriated decay estimates for them. These estimates can then be

transported to ${}^{(ext)}\mathcal{M}$ using the the full scope of the null structure and null Bianchi identities associated to the outgoing geodesic foliation.

6. The decay estimates in ${}^{(ext)}\mathcal{M}$ can then be used as initial condition along the time-like hypersurface \mathcal{T} for the incoming foliation of ${}^{(int)}\mathcal{M}$. These allows us to also derive appropriate decay estimates for all null connection and curvature coefficients of the foliation induced by \underline{u} .
7. The precise decay estimates derived in 5 are sufficiently strong to allow us to control all error terms generated in the process of estimating \mathfrak{q} , as mentioned in 3.

Note that in Figure 1.1, one starts with initial conditions on the union of null hypersurfaces $\mathcal{C}_1 \cup \underline{\mathcal{C}}_1$ rather than an initial spacelike hypersurface $\Sigma_{(0)}$. One can justify this simplification based on the results of [37], [38], see Remark 3.3.1. The full red line \mathcal{H}_+ represents the future event horizon of the perturbed Schwarzschild. The line \mathcal{T} represents the timelike hypersurface separating ${}^{(int)}\mathcal{M}$ from ${}^{(ext)}\mathcal{M}$. In deriving decay estimates the precise choice of \mathcal{T} is irrelevant. A choice, however, needs to be made in order to avoid a derivative loss for our top energy estimates²⁰.

The spacetime is constructed by a continuity argument, i.e. we assume that the spacetime terminating at $\mathcal{C}_* \cup \underline{\mathcal{C}}_*$ saturates a given bootstrap assumption **(BA)** and show, by a long sequence of a-priori estimates which take advantage of the smallness of the initial perturbation, that **(BA)** can be improved and the spacetime extended past $\mathcal{C}_* \cup \underline{\mathcal{C}}_* \cup \Sigma_*$.

Our work here is the first to prove the nonlinear stability of Schwarzschild in a restricted class of nontrivial perturbations, i.e. perturbations for which new ideas, such as our GCM procedure are needed. To a large extent, the restriction to this class of perturbations is only needed to ensure that the final state of evolution is another Schwarzschild space. We are thus confident that our procedure may apply in a more general setting. We would like to single out two other recent important contributions to nonlinear stability of black holes. In the context of asymptotically flat Einstein vacuum equations the result of Dafermos-Holzegel-Rodnianski [23] constructs a class of Kerr black hole solutions starting from future infinity while Hintz-Vasy [26]²¹ prove the nonlinear stability of Kerr-de Sitter, for small angular momentum, in the context of the Einstein vacuum equations with a nontrivial positive cosmological constant. Though the two results are very different they share in common the fact that the perturbations they treat decay exponentially. This makes the analysis significantly easier than in our case when the decay is barely enough to control the nonlinear terms.

²⁰See [17] for a similar situation.

²¹See also [27] for the stability of Kerr-Newman de Sitter.

1.4 Organization

The paper is organized as follow. In Chapter 2 we introduce the main quantities, equations and basic tools needed later. It is our main reference kit providing all main null structure and null Bianchi equations, in general null frames, in the context of axially symmetric polarized spacetimes. Though we work with the reduced equations, i.e the equations reduced by the symmetries, most of the work in the paper does not really depend of the reduction. Besides insuring that the final state is a Schwarzschild space the reduction only plays a significant role in the GCM construction.

Chapter 3, the heart of the paper, contains the precise version of our main theorem, its main conclusions as well as a full strategy of its proof, divided in nine supporting intermediate results, Theorems **M0–M8**. We also give a short description of the proof of each theorem.

In the other chapters of this paper we give complete proofs of Theorems, **M0–M8** and a full description of our GCM procedure.

The reader versed in the formalism of null structure and Bianchi equations, as discussed in [17], is encouraged to glance fast over Chapter 2, to get familiarized with the notation, and then move directly to Chapter 3.

1.5 Acknowledgements

This work would be inconceivable without the remarkable advances made in the last sixty years on black holes. The works of Regge-Weeler, Carter, Teukolsky, Chandrasekhar, Wald etc., made during the so called *golden age* of black hole physics in the sixties and seventies, have greatly influenced our understanding of invariant quantities and the wave equations they satisfy. The advances made in the last fifteen years, quoted earlier, which have led to the development of new mathematical methods to derive the decay of waves on black holes spacetimes, are even more immediately relevant to our work. In particular we would like to single out the direct influence of Dafermos-Holzegel-Rodnianski [22] in the gestation of our own ideas in this paper. Finally, the work on the nonlinear stability of the Minkowski space²² in [17], a milestone in the mathematical GR, has significantly instructed our work here. We would like to thank E. Giorgi for her careful proofreading of various sections of the manuscript. Various discussions we had with S. Aksteiner were very useful. Finally we thank our wives Anca and Emilie for their incredible patience,

²²See also [37].

understanding and support during our many years of work on this project.

The first author has been supported by the NSF grant DMS 1362872. He would like to thank the mathematics departments of Paris 6, Cergy-Pontoise and IHES for their hospitality during his many visits in the last six years. The second author is supported by the ERC grant ERC-2016 CoG 725589 EPGR.

Chapter 2

PRELIMINARIES

2.1 Axially symmetric polarized spacetimes

2.1.1 Axial symmetry

We consider vacuum, four dimensional, simply connected, axially symmetric spacetimes $(\mathcal{M}, \mathbf{g}, \mathbf{Z})$ with \mathbf{g} Lorentzian and \mathbf{Z} an axial Killing vectorfield on \mathcal{M} . We denote by \mathfrak{A} the axis of symmetry, i.e. the points on \mathcal{M} for which $X := \mathbf{g}(\mathbf{Z}, \mathbf{Z}) = 0$. In the case of interest for us we assume $dX \neq 0$ and that \mathfrak{A} is a smooth manifold of codimension 2. The Ernst potential of the spacetime is given by,

$$\sigma_\mu := \mathbf{D}_\mu(-\mathbf{Z}^\alpha \mathbf{Z}_\alpha) - i \epsilon_{\mu\beta\gamma\delta} \mathbf{Z}^\beta \mathbf{D}^\gamma \mathbf{Z}^\delta.$$

The 1-form $\sigma_\mu dx^\mu$ is closed and thus there exists a function $\sigma : \mathcal{M} \rightarrow \mathbb{C}$, called the \mathbf{Z} -Ernst potential, such that $\sigma_\mu = \mathbf{D}_\mu \sigma$. Note also that $\mathbf{D}_\mu \mathbf{g}(\mathbf{Z}, \mathbf{Z}) = 2\mathbf{G}_{\mu\lambda} \mathbf{Z}^\lambda = -\Re(\sigma_\mu)$ where $G_{\mu\nu} = D_\mu Z_\nu$. Hence we can choose the potential σ such that $\Re\sigma = -X$. By a standard calculation one can show that,

$$\square\sigma = -X^{-1} \mathbf{D}_\mu \sigma \mathbf{D}^\mu \sigma.$$

Definition 2.1.1. *An axially symmetric Lorentzian manifold $(\mathcal{M}, \mathbf{g}, \mathbf{Z})$ is said to be polarized if the Ernst potential σ is real, i.e. $\sigma = -X$. In that case the metric \mathbf{g} can be written in the form,*

$$\mathbf{g} = X d\varphi^2 + g_{ab} dx^a dx^b \tag{2.1.1}$$

where X and g are independent of φ . We refer to the orbit space \mathcal{M}/\mathbf{Z} as the reduced space and the metric $g = g_{ab}dx^a dx^b$ as the reduced metric. Note that the reduced space $(\mathcal{M}/\mathbf{Z}, g)$ is smooth away from the axis \mathfrak{A} . Moreover the scalar X verifies the wave equation,

$$\square_{\mathbf{g}}X = X^{-1}\mathbf{D}_\mu X \mathbf{D}^\mu X. \quad (2.1.2)$$

We denote by \mathbf{R} , resp. R the curvature tensor of the spacetime metric \mathbf{g} , respectively g , and by $\square_{\mathbf{g}}$, resp \square_g the d'Alembertian with respect to \mathbf{g} and resp. the reduced metric g . We also denote by $\mathbf{\Gamma}$ the Christoffel symbols of \mathbf{g} and by Γ the ones of g . Note that the only non-vanishing Christoffel symbols are:

$$\mathbf{\Gamma}_{\varphi b}^\varphi = \frac{1}{2}X^{-1}\partial_b X, \quad \mathbf{\Gamma}_{\varphi\varphi}^a = -\frac{1}{2}g^{as}\partial_s X, \quad \mathbf{\Gamma}_{bc}^a = \Gamma_{bc}^a. \quad (2.1.3)$$

One can easily prove the following.

Proposition 2.1.2. *The scalar curvature R of the reduced metric g of an axially symmetric polarized Einstein vacuum spacetime vanishes identically¹. Moreover, setting $\Phi := \frac{1}{2}\log X$ we find,*

$$R_{ab} = D_a D_b \Phi + D_a \Phi D_b \Phi, \quad \square_g \Phi = -D^a \Phi D_a \Phi. \quad (2.1.4)$$

Also,

$$\begin{aligned} \mathbf{R}_{a\varphi b}^\varphi &= -\frac{1}{2}X^{-1}D_a D_b X + \frac{1}{4}X^{-2}D_a X D_b X = -R_{ab}, \\ \mathbf{R}_{acb}^\varphi &= 0, \\ \mathbf{R}_{abc}{}^d &= R_{abc}{}^d, \end{aligned} \quad (2.1.5)$$

and,

$$R_{abcd} = g_{ac}R_{bd} + g_{bd}R_{ac} - g_{ad}R_{bc} - g_{bc}R_{ad}. \quad (2.1.6)$$

Finally, when applied to \mathbf{Z} -invariant functions,

$$\square_{\mathbf{g}} = \square_g + g^{ab}\partial_a \Phi \partial_b. \quad (2.1.7)$$

Remark 2.1.3. *The wave equation in (2.1.4) is equivalent to*

$$\square_{\mathbf{g}}\Phi = 0. \quad (2.1.8)$$

Remark 2.1.4. *Schwarzschild spacetime is axially symmetric polarized with,*

$$X = r^2(\sin \theta)^2, \quad \Phi = \log(r) + \log(\sin \theta).$$

¹This is an easy consequence of the equation (2.1.2).

2.1.2 \mathbf{Z} -frames

We consider orthonormal frames $e_0, e_1, e_\theta = e_2, e_\varphi = X^{-1/2}\mathbf{Z}$, with $X := \mathbf{g}(\mathbf{Z}, \mathbf{Z})$, which are \mathbf{Z} -equivariant, i.e. $[\mathbf{Z}, e_\alpha] = 0$. From now on, the index φ is referring to the frame rather than the coordinates

Lemma 2.1.5. *Setting $(\Lambda_\alpha)_{\beta\gamma} := \mathbf{g}(D_\alpha e_\gamma, e_\beta)$ we have,*

$$(\Lambda_\varphi)_{a\varphi} = -D_a\Phi, \quad (\Lambda_\varphi)_{ab} = (\Lambda_\alpha)_{b\varphi} = 0, \quad \forall a = 0, 1, 2, \quad (2.1.9)$$

and,

$$\begin{aligned} \mathbf{D}_a e_b &= D_a e_b, \\ \mathbf{D}_a e_\varphi &= 0, \\ \mathbf{D}_\varphi e_a &= (\Lambda_\varphi)_{\varphi a} e_\varphi = (D_a \Phi) e_\varphi, \\ \mathbf{D}_\varphi e_\varphi &= (\Lambda_\varphi)_{a\varphi} e_a = -D^a \Phi e_a. \end{aligned} \quad (2.1.10)$$

Proof. Straightforward verification. □

Lemma 2.1.6. *We have,*

$$\begin{aligned} \mathbf{D}_s \mathbf{R}_{abcd} &= D_s R_{abcd}, \\ \mathbf{D}_s \mathbf{R}_{\varphi bcd} &= 0, \\ \mathbf{D}_s \mathbf{R}_{\varphi b\varphi d} &= -D_s R_{bd}, \\ \mathbf{D}_\varphi \mathbf{R}_{abcd} &= 0, \\ \mathbf{D}_\varphi \mathbf{R}_{\varphi bcd} &= D^s \Phi R_{sbcd} + D_c \Phi R_{bd} - D_d \Phi R_{bc}, \\ \mathbf{D}_\varphi \mathbf{R}_{\varphi b\varphi d} &= 0. \end{aligned}$$

Proof. Straightforward verification. □

Definition 2.1.7. *We say that a spacetime tensor \mathbf{U} is \mathbf{Z} -invariant if $\mathcal{L}_{\mathbf{Z}}\mathbf{U} = 0$ and \mathbf{Z} -invariant polarized if its contractions to an odd number of $e_\varphi = X^{-1/2}\mathbf{Z}$ vanish identically.*

Proposition 2.1.8. *All higher covariant derivatives of the Riemann curvature tensor \mathbf{R} of an axially symmetric polarized spacetime $(\mathbf{M}, \mathbf{g}, \mathbf{Z})$ are \mathbf{Z} -invariant, polarized.*

Proof. The statement has been already verified above for both \mathbf{R} and \mathbf{DR} . It suffices to show that, given an arbitrary \mathbf{Z} -invariant, polarized tensor \mathbf{U} , its covariant derivative \mathbf{DU} is also \mathbf{Z} -invariant, polarized. The invariance is immediate. To show polarization

we consider all frame components of \mathbf{DU} with respect to our adapted equivariant frame e_1, e_2, e_3, e_φ . Assume first that the components of \mathbf{DU} contain only one e_φ . These are,

$$\mathbf{D}_\varphi \mathbf{U}_a, \quad \mathbf{D}_a \mathbf{U}_{b\varphi c}$$

with various combinations of horizontal indices a, b, c . Now, in view of the polarization property of \mathbf{U} and the relations $\mathbf{D}_a e_b = D_a e_b$, $\mathbf{D}_a e_\varphi = 0$ we easily deduce,

$$\mathbf{D}_a \mathbf{U}_{b\varphi c} = e_a \mathbf{U}_{b\varphi c} - \mathbf{U}_{\mathbf{D}_a b\varphi c} - \mathbf{U}_{b\mathbf{D}_a \varphi c} - \mathbf{U}_{b\varphi c} = 0.$$

Similarly, since $e_\varphi(U_a) = X^{-1/2} \mathbf{Z}(U_a) = X^{-1/2} \mathcal{L}_{\mathbf{Z}} U_a = 0$ and $\mathbf{D}_\varphi e_a$ is proportional to e_φ ,

$$\mathbf{D}_\varphi \mathbf{U}_a = e_\varphi(\mathbf{U}_a) - \mathbf{U}_{\mathbf{D}_\varphi e_a} = 0.$$

Similarly we can check that the contraction of \mathbf{DU} with any odd number of e_φ must be zero. \square

In what follows we shall refer to \mathbf{Z} -invariant, polarized tensors as simply \mathbf{Z} -polarized.

2.1.3 Axis of symmetry

We denote by \mathfrak{A} the axis of symmetry of \mathbf{Z} , i.e. the set of zeroes of $X = \mathbf{g}(\mathbf{Z}, \mathbf{Z})$. Since we assume $dX \neq 0$, \mathfrak{A} is a smooth timelike submanifold of dimension 2. In view of the definition of axial symmetry every trajectory of \mathbf{Z} is closed and intersects \mathfrak{A} at one point. The following regularity result at \mathfrak{A} holds true.

Lemma 2.1.9. *At the axis of symmetry \mathfrak{A} we have,*

$$\frac{\mathbf{g}^{\mu\nu} \partial_\mu X \partial_\nu X}{4X} = e^{2\Phi} \mathbf{g}^{\mu\nu} \partial_\mu \Phi \partial_\nu \Phi \longrightarrow 1. \quad (2.1.11)$$

Proof. This is a classical result, see for example [43]. We provide a proof for the convenience of the reader. We introduce a coordinates system (x^0, x^1, x^2, x^3) centered at a point $q = (0, 0, 0, 0)$ on the axis such that the Christoffel symbols of the metric vanish at q and $\partial_{x^0}|_q$ and $\partial_{x^1}|_q$ are tangent to the axis at q . In particular, in this coordinates system, the matrix $\partial_\alpha \mathbf{Z}^\mu(q)$ is given by

$$\partial_\alpha \mathbf{Z}^\mu(q) = \begin{pmatrix} 0 & 0 \\ 0 & A \end{pmatrix},$$

where A is an antisymmetric matrix. Note that we used the fact that \mathbf{Z} vanishes on the axis, that q belongs to the axis, and that $\partial_\alpha \mathbf{Z}^\mu(q)$ is antisymmetric since \mathbf{Z} is Killing.

Now, if $x(\varphi)$ denotes an orbit of \mathbf{Z} close to q , and $y = (x^2, x^3)$, we have in particular from Taylor formula

$$\frac{dy}{d\varphi} = Ay + O(y^2).$$

Hence

$$\exp(-\varphi A)y(\varphi) = y(0) + O(\varphi y^2)$$

and since $y(2\pi) = y(0)$ in view of the 2π -periodicity of the orbits of \mathbf{Z} , we infer

$$\exp(-2\pi A)y(0) = y(0) + O(y^2).$$

As $y(0)$ can be taken arbitrarily small, we infer that $\exp(2\pi A)$ is the 2×2 identity matrix. Since A is antisymmetric and non zero, its eigenvalues necessarily are i and $-i$, and hence $A^T A = I$. This yields

$$A^\alpha{}_\mu A_\gamma{}^\nu A_{\alpha\nu} = A^\alpha{}_\mu (A^T A)_{\gamma\alpha} = A_{\gamma\mu}$$

and hence

$$\partial^\alpha \mathbf{Z}_\mu(q) \partial_\gamma \mathbf{Z}^\nu(q) \partial_\alpha \mathbf{Z}_\nu(q) = \partial_\gamma \mathbf{Z}_\mu(q).$$

Finally, since \mathbf{Z} vanishes on the axis, and since the coordinates system we use in this lemma has vanishing Christoffel symbols at q , we have as $|x|$ goes to 0

$$\frac{\mathbf{g}^{\mu\nu} \partial_\mu X \partial_\nu X}{4X} = \frac{\mathbf{Z}^\mu \mathbf{D}^\alpha \mathbf{Z}_\mu \mathbf{Z}^\nu \mathbf{D}_\alpha \mathbf{Z}_\nu}{\mathbf{Z}^\mu \mathbf{Z}_\mu} = \frac{\partial_\beta \mathbf{Z}^\mu(q) x^\beta \partial^\alpha \mathbf{Z}_\mu(q) \partial_\gamma \mathbf{Z}^\nu(q) x^\gamma \partial_\alpha \mathbf{Z}_\nu(q)}{\partial_\beta \mathbf{Z}^\mu(q) x^\beta \partial_\gamma \mathbf{Z}_\mu(q) x^\gamma} + O(x).$$

Together with the previous identity, we infer near any point q on the axis

$$\frac{\mathbf{g}^{\mu\nu} \partial_\mu X \partial_\nu X}{4X} \longrightarrow 1.$$

This concludes the proof of the lemma. □

We note that \mathbf{Z} -polarized, smooth, vectorfields are automatically tangent to \mathfrak{A} . This is the content of the following.

Lemma 2.1.10. *Any, regular (i.e. smooth) \mathbf{Z} -polarized vectorfield \mathbf{U} is tangent to the axis \mathfrak{A} .*

Proof. Let \mathbf{U} a polarized \mathbf{Z} -invariant regular vectorfield. Since it is \mathbf{Z} -invariant, we have

$$0 = [\mathbf{Z}, \mathbf{U}] = \mathbf{Z}^\alpha \mathbf{D}_\alpha \mathbf{U} - \mathbf{U}^\alpha \mathbf{D}_\alpha \mathbf{Z}.$$

Since $\mathbf{Z} = 0$ on the axis and \mathbf{U} is regular (hence bounded on the axis) we infer that $\mathbf{U}^\alpha \mathbf{D}_\alpha \mathbf{Z} = 0$ on \mathfrak{A} . In view of (2.1.10),

$$\mathbf{U}^\alpha \mathbf{D}_\alpha \mathbf{Z} = \mathbf{U}(e^\phi) e_\varphi,$$

and since e_φ is unitary, we infer that

$$\mathbf{U}(X^{1/2}) = \mathbf{U}(e^\phi) = 0 \text{ on } \mathfrak{A}$$

and hence $\mathbf{U}(X) = 0$ when $X = 0$. □

Corollary 2.1.11. *Let u be a smooth regular optical function, i.e. $\mathbf{g}^{\alpha\beta} \mathbf{D}_\alpha u \mathbf{D}_\beta u = 0$, which is \mathbf{Z} -invariant, i.e. $\mathbf{Z}(u) = 0$. Then its associated null geodesic generator $L = -\mathbf{g}^{\alpha\beta} \partial_\alpha u \partial_\beta u$ is \mathbf{Z} -invariant, polarized, tangent to the axis of symmetry \mathfrak{A} .*

Proof. It is easy to check that L is \mathbf{Z} -invariant, polarized. It must therefore be tangent to \mathfrak{A} in view of Lemma 2.1.10. □

2.1.4 \mathbf{Z} -polarized S -surfaces

Throughout our work we shall deal various \mathbf{Z} -polarized, S -foliations i.e. foliations given by compact 2-surfaces S with induced metrics of the form,

$$g = \gamma d\theta^2 + X d\varphi^2, \quad \gamma = \gamma(\theta) > 0, \quad \theta \in [0, \pi]. \quad (2.1.12)$$

Here γ and X are independent of φ and $\Phi = \frac{1}{2} \log X$ vanishes on the poles $\theta = 0$ and $\theta = \pi$. The regularity condition (2.1.11) takes the form,

$$\lim_{\sin\theta \rightarrow 0} \left(e_\theta(e^\Phi) \right)^2 = 1 \quad (2.1.13)$$

where e_θ is the unit vector,

$$e_\theta := \gamma^{-1/2} \partial_\theta.$$

We denote the induced covariant derivative ∇ and define the volume radius of S by the formula

$$|S| = 4\pi r^2$$

where $|S|$ is the volume of the surface using the volume form of the metric g . Note also that the area element on S is given by

$$\sqrt{\gamma}e^\Phi d\theta d\varphi.$$

In this subsection we record some basic general formulas concerning these surfaces. We consider adapted orthonormal frames

$$e_\theta, e_\varphi = X^{-1/2}\mathbf{Z} = X^{-1/2}\partial_\varphi.$$

Note that in view of (2.1.10) we have,

$$\nabla_\varphi e_\varphi = -(e_\theta\Phi)e_\theta, \quad \nabla_\varphi e_\theta = (e_\theta\Phi)e_\varphi, \quad \nabla_\theta e_\theta = \nabla_\theta e_\varphi = 0. \quad (2.1.14)$$

In what follows, we consider \mathbf{Z} -invariant polarized tensors tangent to S or simply polarized k -tensors on S .

In view of Lemma 2.1.10, a regular \mathbf{Z} -polarized tensor on S must vanish on the axis of symmetry i.e. at $\theta = 0$ and $\theta = \pi$. More precisely we have,

Lemma 2.1.12. *The following facts hold true for \mathbf{Z} -polarized tensors on S .*

1. *If U is a 1-form then, on the axis of symmetry² of \mathbf{Z} , (i.e. for $\theta = 0$ and $\theta = \pi$),*

$$U_\theta := U(e_\theta) = 0$$

2. *For a covariant 2-tensor, then, on the axis of symmetry³ of \mathbf{Z} , (i.e. for $\theta = 0$ and $\theta = \pi$),*

$$U_{\theta\theta} = U_{\varphi\varphi} = 0.$$

Similar statements can be deduced for higher order tensors.

Proof. Immediate consequence of Lemma 2.1.10. □

Lemma 2.1.13. *The Gauss curvature K of the metric (2.1.12) can be expressed in terms of the polar function $\Phi := \frac{1}{2} \log X$ by the formula,*

$$\Delta\Phi = -K. \quad (2.1.15)$$

Proof. Direct calculation using the form of the g metric in (2.1.12). □

²Note that the component U_φ must automatically vanish on S .

³Note that the components $U_{\theta\varphi}, U_{\varphi\theta}$ must automatically vanish on S .

Basic operators on S

We recall (see [17] chapter 2) the following operations which preserve the space of fully symmetric traceless tensors:

Definition 2.1.14. We denote by \mathcal{S}_k the set of k -covariant polarized tensors which are fully symmetric and traceless, i.e. which verify,

$$f_{A_1 \dots A_k} = f_{(A_1 \dots A_k)}, \quad \not{g}^{A_1 A_2} f_{A_1 A_2 \dots A_k} = 0.$$

We define the following operators on \mathcal{S}_k -tensors.

1. The operator \mathcal{D}_k which takes \mathcal{S}_k into \mathcal{S}_{k-1} is the divergence operator,

$$(\mathcal{D}_k f)_{A_2, \dots, A_k} := (\text{div} f)_{A_2, \dots, A_k} := \not{g}^{AB} \nabla_B f_{AA_2, \dots, A_k}.$$

2. The operator \mathcal{D}_k^* which takes \mathcal{S}_{k-1} into \mathcal{S}_k is the fully symmetrized, traceless, covariant derivative operator⁴,

$$(\mathcal{D}_k^* f)_{A_1 \dots A_k} := \begin{cases} -\nabla_{A_1} f, & k = 1, \\ -\frac{1}{k} \nabla_{(A_1} f_{A_2 \dots A_k)} + \frac{1}{k(k-1)} \not{g}^{(A_1 A_2} (\text{div} f)_{A_3 \dots A_k)}, & k \geq 2. \end{cases}$$

3. The operator \mathcal{A}_k takes \mathcal{S}_k to \mathcal{S}_k ,

$$(\mathcal{A}_k f)_{A_1 \dots A_k} := \not{g}^{BC} \nabla_B \nabla_C f_{A_1 \dots A_k}.$$

Remark 2.1.15. Note that if $f \in \mathcal{S}_k$ then $\text{curl} f := \epsilon^{BC} \nabla_B f_{CA_1 \dots A_k} = 0$.

Lemma 2.1.16. Given $f \in \mathcal{S}_k$, $k \geq 1$, we have the identity,

$$\nabla_B f_{A_1 \dots A_k} = -(\mathcal{D}_{k+1}^* f)_{BA_1 \dots A_k} + \frac{1}{k} \not{g}^{(BA_1} (\mathcal{D}_k f)_{A_2 \dots A_k)}. \quad (2.1.16)$$

In other words the covariant derivatives of any tensor in \mathcal{S}_k can be expressed as a linear combination of $\mathcal{D}_{k+1}^* f$ and $\not{g} \otimes \mathcal{D}_{k-1} f$.

Proof. The proof follows easily from definitions and the vanishing of the curl. For example, if $k = 2$,

$$\begin{aligned} 3\nabla_B f_{A_1 A_2} &= (\nabla_B f_{A_1 A_2} + \nabla_{A_1} f_{A_2 B} + \nabla_{A_2} f_{BA_1}) \\ &+ (\nabla_B f_{A_1 A_2} - \nabla_{A_1} f_{BA_2}) + (\nabla_B f_{A_1 A_2} - \nabla_{A_2} f_{A_1 B}) \\ &= -3 [(\mathcal{D}_3^* f)_{BA_1 A_2} - \not{g}_{A_1 A_2} (\mathcal{D}_2 f)_B - \not{g}_{A_2 B} (\mathcal{D}_2 f)_{A_1} - \not{g}_{A_1 A_2} (\mathcal{D}_2 f)_B] \\ &= -3 \left[(\mathcal{D}_3^* f)_{BA_1 A_2} - \frac{1}{2} \not{g}^{(BA_1} (\mathcal{D}_3 f)_{A_2)} \right]. \end{aligned}$$

□

⁴For an arbitrary k -tensor, $f_{(A_1 \dots A_k)} = \sum_{\sigma \in \Pi} f_{A_{\sigma(1)} \dots A_{\sigma(k)}}$. In the particular case when $k = 1$ we get $(\mathcal{D}_1^* f)_A = -\nabla_A f$ and when $k = 2$ we get $\mathcal{D}_2^* f_{AB} = -\frac{1}{2} (\nabla_A f_B + \nabla_B f_A - \not{g}_{AB} \text{div} f)$.

We can easily check that \mathcal{D}_k^* is the formal adjoint of \mathcal{D}_k , i.e.,

$$\int_S (\mathcal{D}_k f) g = \int_S f (\mathcal{D}_k^* g).$$

It is also easy to check that the kernels of \mathcal{D}_k are trivial for all $k \geq 1$ (see also Chapter 2 in [17]). The kernel of $\mathcal{D}_1^* : \mathcal{S}_0 \rightarrow \mathcal{S}_1$ consists of constants on S while the kernel of \mathcal{D}_2^* consists of constant multiple of co-vectors f with $f_\theta = C e^\Phi$. Moreover,

$$\begin{aligned} \mathcal{D}_1^* \cdot \mathcal{D}_1 &= -\Delta_1 + K, & \mathcal{D}_1 \cdot \mathcal{D}_1^* &= -\Delta_0, \\ \mathcal{D}_2^* \cdot \mathcal{D}_2 &= -\frac{1}{2}\Delta_2 + K, & \mathcal{D}_2 \cdot \mathcal{D}_2^* &= -\frac{1}{2}(\Delta_1 + K). \end{aligned} \quad (2.1.17)$$

Similar identities also hold for higher k . Using (2.1.17) one can also prove the following (see also Chapter 2 in [17]).

Proposition 2.1.17. *Let (S, g) be a compact manifold with Gauss curvature K . We have,*

i.) *The following identity holds for vectorfields $f \in \mathcal{S}_1$,*

$$\int_S (|\nabla f|^2 + K|f|^2) = \int_S |\mathcal{D}_1 f|^2.$$

ii.) *The following identity holds for symmetric, traceless tensors in \mathcal{S}_2 ,*

$$\int_S (|\nabla f|^2 + 2K|f|^2) = 2 \int_S |\mathcal{D}_2 f|^2.$$

iii.) *The following identity holds for scalars $f \in \mathcal{S}_0$,*

$$\int_S |\nabla f|^2 = \int_S |\mathcal{D}_1^* f|^2.$$

iv.) *The following identity holds for vectors $f \in \mathcal{S}_1$,*

$$\int_S (|\nabla f|^2 - K|f|^2) = 2 \int_S |\mathcal{D}_2^* f|^2.$$

Proof. All statements appear in [17]. □

Proposition 2.1.18. *we have for $f \in \mathcal{S}_0$,*

$$\int_S (|\nabla^2 f|^2 + K|\nabla f|^2) = \int_S |\Delta_0 f|^2.$$

Moreover, under mild assumptions on the curvature such as

$$K = \frac{1}{r^2} + O\left(\frac{\epsilon}{r^2}\right), \quad re_\theta(K) = O\left(\frac{\epsilon}{r^2}\right),$$

for any $f \in \mathcal{S}_k$, $k \geq 1$,

$$\int_S (|\nabla^2 f|^2 + r^{-2}|\nabla f|^2) \lesssim \int_S |\Delta_k f|^2 + O(\epsilon)r^{-4} \int_S |f|^2.$$

Proof. Follows from the standard Bochner identity on S . □

Reduced Picture

Lemma 2.1.19. *The following relations hold true between the spacetime picture and the reduced one.*

1. Let ${}^{(1+3)}f \in \mathcal{S}_k$ such that ${}^{(1+3)}f_{\theta\dots\theta} = f$. Then,

$$(\mathcal{D}_k {}^{(1+3)}f)_{\theta\dots\theta} = e_\theta(f) + ke_\theta(\Phi)f. \quad (2.1.18)$$

2. If $f \in \mathcal{S}_0$ we have,

$$\mathcal{D}_1^* f = -e_\theta(f).$$

3. If ${}^{(1+3)}f \in \mathcal{S}_{k-1}$, $k \geq 2$, such that ${}^{(1+3)}f_{\theta\dots\theta} = f$ we have,

$$2(\mathcal{D}_k^* {}^{(1+3)}f)_{\theta\dots\theta} = -e_\theta(f) + (k-1)e_\theta(\Phi)f. \quad (2.1.19)$$

4. Let ${}^{(1+3)}f \in \mathcal{S}_k$ such that ${}^{(1+3)}f_{\theta\dots\theta} = f$. Then,

$$\Delta_k {}^{(1+3)}f_{\theta_1\dots\theta_k} = e_\theta(e_\theta f) + e_\theta(\Phi)e_\theta f - k^2(e_\theta(\Phi))^2 f.$$

Proof. The proof follows easily from the definitions of \mathcal{D}_k , \mathcal{D}_k^* , Δ_k and the formulae (2.1.14). We check below the formula (2.1.18).

$$\begin{aligned} -(\mathcal{D}_k^* {}^{(1+3)}f)_{\theta\dots\theta} &= e_\theta f - \frac{1}{2}(\mathcal{D}_{k-1} f)_{\theta\dots\theta} = e_\theta(f) - \frac{1}{2}(e_\theta f + (k-1)e_\theta(\Phi)f) \\ &= \frac{1}{2}(e_\theta f - (k-1)e_\theta(\Phi)f) \end{aligned}$$

as desired. □

Definition 2.1.20. We say that a scalar f is a reduced k -scalar on S if there is a \mathbf{Z} -invariant, polarized, k -covector ${}^{(1+3)}f \in \mathcal{S}_k$ such that,

$$f = {}^{(1+3)}f_{\theta\dots\theta}.$$

We denote by \mathfrak{s}_k the set of k reduced scalars.

- Given a k reduced scalar f , reduced from ${}^{(1+3)}f$ we define,

$$|\nabla f|^2 = |\nabla {}^{(1+3)}f|^2, \quad |\nabla^l f|^2 = |\nabla^l {}^{(1+3)}f|^2.$$

- Given a k -reduced scalar f on S we define,

$$\mathcal{A}_k f := e_\theta(f) + k e_\theta(\Phi)f.$$

- Given a $(k-1)$ -reduced scalar $f \in S_{k-1}$ we define,

$$\mathcal{A}_k^* f := -e_\theta(f) + (k-1)e_\theta(\Phi)f.$$

- Given a k -reduced scalar $f \in \mathfrak{s}_k$ we define,

$$\mathcal{A}_k f := e_\theta(e_\theta f) + e_\theta(\Phi)e_\theta f - k^2(e_\theta(\Phi))^2 f.$$

In view of Lemma 2.1.19 we have,

$$\mathcal{A}_k f = (\mathcal{D}_k {}^{(1+3)}f)_{\theta\dots\theta}$$

and,

$$\mathcal{A}_k^* f = \begin{cases} (\mathcal{D}_k^* {}^{(1+3)}f)_{\theta\dots\theta}, & k = 1, \\ 2(\mathcal{D}_k^* {}^{(1+3)}f)_{\theta\dots\theta}, & k \geq 2. \end{cases}$$

Clearly \mathcal{A}_k takes k -reduced scalars into $(k-1)$ -reduced scalars, \mathcal{A}_k^* takes $(k-1)$ -reduced ones into k -reduced and \mathcal{A}_k takes k -reduced scalars into k -reduced scalars.

Remark 2.1.21. Note that, in view of Lemma 2.1.12, any reduced scalar in \mathfrak{s}_k , for $k \geq 1$, must vanish on the axis of symmetry of \mathbf{Z} , i.e. at the two poles.

Remark 2.1.22. The operator \mathcal{A}_k and \mathcal{A}_k^* can only be applied to k -reduced, resp $(k-1)$ -reduced scalars. Thus whenever we write a sequence of operators involving \mathcal{A}_k , \mathcal{A}_k^* we understand from the context to which type of k -reduced scalars they are applied, see for example the proposition below. The same remark applies to \mathcal{A}_k .

Remark 2.1.23. Note that for given reduced scalar $f \in \mathfrak{s}_k$ and $h \in \mathfrak{s}_1$ we can write,

$$he_\theta(f) = \frac{1}{2}h(\not\partial_k f - \not\partial_{k+1}^* f).$$

The term $h\not\partial_k f$ is the reduced form of a tensor product of $(1+3)h$ with $\mathcal{D}_k^{(1+3)}f$ while $h\not\partial_{k+1}^* f$ is the reduced form of a contraction between $(1+3)h$ and $\mathcal{D}_{k+1}^*(1+3)\psi$. This can be formalized precisely using Lemma 2.1.16. The Remark will be useful in what follows, for example in Lemma 2.2.14.

Remark 2.1.24. The duality between the operators $\not\partial_k$ and $\not\partial_k^*$ follows in view of the duality of \mathcal{D}_k and \mathcal{D}_k^* . It can also be interpreted directly in terms of the area element $\sqrt{\gamma}e^\Phi d\theta d\varphi$,

$$\begin{aligned} \int_S (\not\partial_k f g - g \not\partial_k^* g) da_S &= \int_S e_\theta(fg) + e_\theta(\Phi)fg = \int_0^\pi \int_0^{2\pi} (e_\theta(fg) + e_\theta(\Phi)fg) \sqrt{\gamma}e^\Phi d\theta d\varphi \\ &= \int_0^\pi \int_0^{2\pi} \partial_\theta(e^\Phi fg) d\theta d\varphi = 0. \end{aligned}$$

Proposition 2.1.25. The following identities hold true,

$$\begin{aligned} \not\partial_k^* \not\partial_k &= -\not\Delta_k + kK, \\ \not\partial_k \not\partial_k^* &= -\not\Delta_{k-1} - (k-1)K. \end{aligned} \tag{2.1.20}$$

In particular for $k = 1, 2$

$$\not\partial_1^* \not\partial_1 = -\not\Delta_1 + K, \quad \not\partial_1 \not\partial_1^* = -\not\Delta_0, \quad \not\partial_2^* \not\partial_2 = -\not\Delta_2 + 2K, \quad \not\partial_2 \not\partial_2^* = -\not\Delta_1 - K.$$

Moreover, note the following commutation formulas

$$\begin{aligned} \not\partial_k \not\partial_k^* - \not\partial_{k-1}^* \not\partial_{k-1} &= -2(k-1)K, \\ -\not\partial_k \not\Delta_k + \not\Delta_{k-1} \not\partial_k &= K \not\partial_k - ke_\theta(K), \\ -\not\partial_k^* \not\Delta_{k-1} + \not\Delta_k \not\partial_k^* &= (2k-1)K \not\partial_k^* + (k-1)e_\theta(K). \end{aligned}$$

Proof. We have, for a k reduced scalar f ,

$$\begin{aligned} -\not\partial_k^* \not\partial_k f &= (e_\theta - (k-1)e_\theta(\Phi))(e_\theta(f) + ke_\theta(\Phi)f) \\ &= e_\theta(e_\theta(f)) + ke_\theta(\Phi)e_\theta f + k(e_\theta e_\theta \Phi)f - (k-1)e_\theta(\Phi)(e_\theta(f) + ke_\theta(\Phi)f) \\ &= e_\theta(e_\theta(f)) + e_\theta(\Phi)e_\theta f + k(e_\theta e_\theta \Phi)f - k(k-1)(e_\theta(\Phi))^2. \end{aligned}$$

In view of Lemma 2.1.13 we have, since Φ is a scalar

$$-K = \not\Delta\Phi = e_\theta e_\theta(\Phi) + (e_\theta(\Phi))^2.$$

Therefore,

$$\begin{aligned}
-\mathcal{d}_k^* \mathcal{d}_k f &= e_\theta(e_\theta(f)) + e_\theta(\Phi)e_\theta f + k \left(-K - (e_\theta(\Phi))^2 \right) f - k(k-1)(e_\theta(\Phi))^2 f \\
&= e_\theta(e_\theta(f)) + e_\theta(\Phi)e_\theta f - kKf - k^2(e_\theta(\Phi))^2 f \\
&= \mathcal{A}_k f - kKf.
\end{aligned}$$

Similarly, for a $(k-1)$ -reduced f ,

$$\begin{aligned}
-\mathcal{d}_k \mathcal{d}_k^* f &= (e_\theta + ke_\theta(\Phi))(e_\theta(f) - (k-1)e_\theta(\Phi)f) \\
&= e_\theta(e_\theta(f)) + ke_\theta(\Phi)e_\theta f - (k-1)(e_\theta e_\theta \Phi)f - (k-1)e_\theta(\Phi)e_\theta(f) \\
&\quad - k(k-1)(e_\theta(\Phi))^2 f \\
&= e_\theta(e_\theta(f)) + e_\theta(\Phi)e_\theta f - (k-1) \left(-K - (e_\theta(\Phi))^2 \right) f - k(k-1)(e_\theta(\Phi))^2 f \\
&= e_\theta(e_\theta(f)) + e_\theta(\Phi)e_\theta f + (k-1)K - (k-1)^2(e_\theta(\Phi))^2 f \\
&= \mathcal{A}_{k-1} f + (k-1)K.
\end{aligned}$$

Next, we check the commutation formulas. We have

$$\begin{aligned}
\mathcal{d}_k \mathcal{d}_k^* - \mathcal{d}_{k-1}^* \mathcal{d}_{k-1} &= -\mathcal{A}_{k-1} - (k-1)K - \left(-\mathcal{A}_{k-1} + (k-1)K \right) \\
&= -2(k-1)K
\end{aligned}$$

from which we infer

$$\begin{aligned}
\mathcal{d}_k(-\mathcal{A}_k) &= \mathcal{d}_k(\mathcal{d}_k^* \mathcal{d}_k - kK) \\
&= \mathcal{d}_k \mathcal{d}_k^* \mathcal{d}_k - kK \mathcal{d}_k - ke_\theta(K) \\
&= \left(\mathcal{d}_{k-1}^* \mathcal{d}_{k-1} - 2(k-1)K \right) \mathcal{d}_k - kK \mathcal{d}_k - ke_\theta(K) \\
&= \left(-\mathcal{A}_{k-1} + (k-1)K \right) \mathcal{d}_k - (k-2)K \mathcal{d}_k - ke_\theta(K)
\end{aligned}$$

and hence

$$-\mathcal{d}_k \mathcal{A}_k + \mathcal{A}_{k-1} \mathcal{d}_k = K \mathcal{d}_k - ke_\theta(K).$$

Also, we have

$$\begin{aligned}
\mathcal{d}_k^*(-\mathcal{A}_{k-1}) &= \mathcal{d}_k^*(\mathcal{d}_k \mathcal{d}_k^* + (k-1)K) \\
&= \mathcal{d}_k^* \mathcal{d}_k \mathcal{d}_k^* + (k-1)K \mathcal{d}_k^* + (k-1)e_\theta(K) \\
&= \left(-\mathcal{A}_k + kK \right) \mathcal{d}_k^* + (k-1)K \mathcal{d}_k^* + (k-1)e_\theta(K)
\end{aligned}$$

and hence

$$-\mathcal{d}_k^* \mathcal{A}_{k-1} + \mathcal{A}_k \mathcal{d}_k^* = (2k-1)K \mathcal{d}_k^* + (k-1)e_\theta(K)$$

as desired. □

A remarkable identity

First, note the following observation which follows immediately from the form of ϕ_2^* .

Lemma 2.1.26. *The kernel of ϕ_2^* is spanned by e^Φ .*

The above lemma, in connection with a Poincaré inequality for ϕ_2^* , see (2.1.35), will result in the need of a specific treatment for the projection of some of the quantities on the kernel of ϕ_2^* . This motivates the following definition.

Definition 2.1.27 (The $\ell = 1$ mode). *For a 1-reduced scalar f , the $\ell = 1$ mode denotes its projection on the kernel of ϕ_2^* , i.e.*

$$\int_S f e^\Phi.$$

For a 0-reduced scalar f , the $\ell = 1$ mode denotes the projection of $e_\theta(f)$ on the kernel of ϕ_2^* , i.e.

$$\int_S e_\theta(f) e^\Phi.$$

Remark 2.1.28. *The above definition is motivated by the fact that, in Schwarzschild, this corresponds to the projection on the $\ell = 1$ spherical harmonic⁵.*

We are now ready to state the following remarkable identity which will play a crucial role later in the paper.

Lemma 2.1.29 (Vanishing of the $\ell = 1$ mode of the Gauss curvature). *The $\ell = 1$ mode of K vanishes identically, i.e.*

$$\int_S e_\theta(K) e^\Phi = 0. \tag{2.1.21}$$

Proof. To prove (2.1.21) we write,

$$-\int_S e_\theta(K) e^\Phi = \int_S \phi_1^*(K) e^\Phi = \int_S K \phi_1(e^\Phi) = 2 \int_S K e_\theta(\Phi) e^\Phi.$$

⁵In general, there are 3 spherical harmonics corresponding to $\ell = 1$, but only one is axially symmetric. This is why we have only one projection instead of 3 in our case.

Thus, in view of (2.1.15), using in addition $\Delta\Phi = e_\theta(e_\theta(\Phi)) + e_\theta(\Phi)^2$

$$\begin{aligned} - \int_S e_\theta(K)e^\Phi &= 2 \int_S \Delta\Phi e_\theta(\Phi)e^\Phi = 2 \int_S (e_\theta(e_\theta(\Phi)) + e_\theta(\Phi)^2)e_\theta(\Phi)e^\Phi \\ &= \int_S \not\partial_2((e_\theta\Phi)^2)e^\Phi = - \int_S (e_\theta\Phi)^2 \not\partial_2^*(e^\Phi) = 0 \end{aligned}$$

as desired⁶. □

Poincaré inequalities on 2-spheres

Proposition 2.1.17 takes the following reduced form,

Proposition 2.1.30. *The following identities hold true for reduced k -scalars $f \in \mathfrak{s}_k$.*

i.) *If $f \in \mathfrak{s}_1$,*

$$\int_S (|\nabla f|^2 + Kf^2) = \int_S |\not\partial_1 f|^2. \quad (2.1.22)$$

ii.) *If $f \in \mathfrak{s}_2$,*

$$\int_S (|\nabla f|^2 + 4Kf^2) = 2 \int_S |\not\partial_2 f|^2. \quad (2.1.23)$$

iii.) *If $f \in \mathfrak{s}_0$,*

$$\int_S |\nabla f|^2 = \int_S |\not\partial_1^* f|^2. \quad (2.1.24)$$

iv.) *If $f \in \mathfrak{s}_1$,*

$$\int_S (|\nabla f|^2 - Kf^2) = \int_S |\not\partial_2^* f|^2. \quad (2.1.25)$$

v.) *If $f \in \mathfrak{s}_0$,*

$$\int_S |\nabla^2 f|^2 + \int_S K|\nabla f|^2 = \int_S |\Delta_0 f|^2. \quad (2.1.26)$$

Under mild assumptions on the Gauss curvature K , such as

$$K = \frac{1}{r^2} + O\left(\frac{\epsilon}{r^2}\right), \quad re_\theta(K) = O\left(\frac{\epsilon}{r^2}\right).$$

⁶Note that the boundary term which appears from the last integration by parts has the form $(\partial_\theta\Phi)^2 e^{2\Phi}(\pi) - (\partial_\theta\Phi)^2 e^{2\Phi}(0)$ and hence vanishes in view of the regularity condition (2.1.13), see also the computation in Remark 2.1.24.

We also have for $f \in \mathfrak{s}_k$, $k \geq 1$,

$$\|\nabla^2 f\|_{L^2(S)}^2 + r^{-2}\|\nabla f\|_{L^2(S)}^2 \lesssim \|\Delta_k f\|_{L^2(S)}^2 + \epsilon r^{-4}\|f\|_{L^2(S)}^2. \quad (2.1.27)$$

Proof. The proof of the above statements can be either derived from their space-time version or checked directly. \square

Lemma 2.1.31. *The following relations hold between \mathbf{Z} -polarized S -tensors and reduced scalars⁷.*

- If $f \in \mathfrak{s}_0$

$$\begin{aligned} |\nabla f|^2 &= |e_\theta f|^2, \\ |\nabla^2 f|^2 &= |e_\theta(e_\theta f)|^2 + |e_\theta \Phi e_\theta f|^2. \end{aligned}$$

- If $f \in \mathfrak{s}_1$,

$$|\nabla f|^2 = |e_\theta f|^2 + |e_\theta(\Phi)|^2 |f|^2.$$

- If $f \in \mathfrak{s}_2$,

$$|\nabla f|^2 = 2(|e_\theta f|^2 + 4|e_\theta(\Phi)|^2 |f|^2).$$

Proof. If $f \in \mathfrak{s}_0$,

$$|\nabla^2 f|^2 = \nabla_A \nabla_B f \nabla^A \nabla^B f = |\nabla_\theta \nabla_\theta f|^2 + |\nabla_\varphi \nabla_\varphi f|^2 = |e_\theta(e_\theta f)|^2 + |e_\theta \Phi e_\theta f|^2.$$

If $f \in \mathfrak{s}_1$ is reduced from a \mathbf{Z} invariant, polarized vector F ,

$$\begin{aligned} |\nabla f|^2 &= \nabla_A F_B \nabla^A F^B = |\nabla_\theta F_\theta|^2 + |\nabla_\varphi F_\varphi|^2 \\ &= |e_\theta f|^2 + |e_\theta \Phi f|^2. \end{aligned}$$

If $f \in \mathfrak{s}_2$ is reduced from a symmetric, traceless \mathbf{Z} -invariant, polarized tensor $F = {}^{(1+3)}f$ we have,

$$\begin{aligned} |\nabla f|^2 &= |\nabla_\theta F_{\theta\theta}|^2 + 2|\nabla_\theta F_{\varphi\theta}|^2 + |\nabla_\theta F_{\varphi\varphi}|^2 + |\nabla_\varphi F_{\theta\theta}|^2 + 2|\nabla_\varphi F_{\varphi\theta}|^2 + |\nabla_\varphi F_{\varphi\varphi}|^2 \\ &= |\nabla_\theta F_{\theta\theta}|^2 + |\nabla_\theta F_{\varphi\varphi}|^2 + 2|\nabla_\varphi F_{\varphi\theta}|^2 \end{aligned}$$

⁷Note that the expressions on the left of the inequalities below should be interpreted as applying to the spacetime tensor from which f is reduced.

and,

$$\begin{aligned}\nabla_\theta F_{\theta\theta} &= e_\theta f = -\nabla_\theta F_{\varphi\varphi}, \\ \nabla_\varphi F_{\varphi\theta} &= e_\theta \Phi F_{\theta\theta} - e_\theta \Phi F_{\varphi\varphi} = 2e_\theta \Phi f.\end{aligned}$$

Thus,

$$|\nabla f|^2 = 2|e_\theta f|^2 + 8(e_\theta \Phi f)^2$$

as desired. \square

Proposition 2.1.32 (Poincaré). *The following inequalities hold for k -reduced scalars.*

1. If $f \in \mathfrak{s}_0$,

$$\int_S |\nabla^2 f|^2 \geq \int_S K |\phi_1^* f|^2. \quad (2.1.28)$$

2. If $f \in \mathfrak{s}_1$

$$\int_S |\nabla f|^2 \geq \int_S K f^2. \quad (2.1.29)$$

3. If $f \in \mathfrak{s}_2$,

$$\int_S |\nabla f|^2 \geq 4 \int_S K f^2. \quad (2.1.30)$$

Proof. We first prove the result for $f \in \mathfrak{s}_2$. According to Lemma 2.1.31,

$$\begin{aligned}2^{-1} |\nabla f|^2 &= |e_\theta f|^2 + 4|e_\theta(\Phi)|^2 |f|^2 = (e_\theta f - 2e_\theta(\Phi)f)^2 + 4f(e_\theta f)e_\theta(\Phi) \\ &= (e_\theta f - 2e_\theta(\Phi)f)^2 + 2e_\theta(f^2)e_\theta(\Phi).\end{aligned}$$

Hence,

$$2^{-1} \int_S |\nabla F|^2 da_S = \int_S (e_\theta f - 2e_\theta(\Phi)f)^2 da_S + 2 \int_S e_\theta(f^2)e_\theta(\Phi) \sqrt{\gamma} e^\Phi d\theta d\varphi.$$

Now,

$$\begin{aligned}\int_S e_\theta(f^2)e_\theta(\Phi) \sqrt{\gamma} e^\Phi d\theta d\varphi &= \int_0^\pi \int_0^{2\pi} \partial_\theta(f^2)e_\theta(\Phi)e^\Phi d\theta d\varphi = - \int_0^\pi \int_0^{2\pi} f^2 e_\theta(e^\Phi e_\theta \Phi) \sqrt{\gamma} d\theta d\varphi \\ &= - \int_0^\pi \int_0^{2\pi} (e_\theta e_\theta \Phi + (e_\theta \Phi)^2) f^2 e^\Phi \sqrt{\gamma} d\theta d\varphi = \int_S K f^2 da_S.\end{aligned}$$

Hence,

$$\int_S |\nabla f|^2 \geq 4 \int_S K f^2$$

as desired.

The result for $f \in \mathfrak{s}_1$ is proved in the same way.

If $f \in \mathfrak{s}_0$ we write, according to Lemma 2.1.31,

$$\begin{aligned} |\nabla^2 f|^2 &= |e_\theta(e_\theta f)|^2 + |e_\theta \Phi e_\theta f|^2 = |e_\theta h|^2 + |e_\theta \Phi|^2 |e_\theta f|^2 \\ &= (e_\theta e_\theta f - e_\theta(\Phi) e_\theta f)^2 + e_\theta[(e_\theta f)^2] e_\theta(\Phi). \end{aligned}$$

Integrating by parts as before,

$$\int_S e_\theta[(e_\theta f)^2] e_\theta(\Phi) da_S = - \int_0^\pi \int_0^{2\pi} (e_\theta f)^2 e_\theta(e^\Phi e_\theta \Phi) \sqrt{\gamma} d\theta d\varphi = \int_S K (e_\theta f)^2 da_S.$$

Thus,

$$\int_S |\nabla^2 f|^2 \geq \int_S K (e_\theta f)^2.$$

□

As a corollary we deduce the following,

Corollary 2.1.33. *The following hold true for reduced scalars,*

1. *If $f \in \mathfrak{s}_1$,*

$$\int_S |\mathfrak{d}_1 f|^2 \geq \int_S 2K f^2. \quad (2.1.31)$$

2. *If $f \in \mathfrak{s}_2$,*

$$\int_S |\mathfrak{d}_2 f|^2 \geq 4 \int_S K f^2. \quad (2.1.32)$$

Proof. According to (2.1.22),

$$\int_S (|\nabla f|^2 + K f^2) = \int_S |\mathfrak{d}_1 f|^2.$$

We deduce,

$$\int_S |\mathfrak{d}_1 f|^2 \geq 2 \int_S K f^2.$$

According to (2.1.23)

$$\int_S (|\nabla f|^2 + 4K f^2) = 2 \int_S |\mathfrak{d}_2 f|^2.$$

Hence,

$$2 \int_S |\mathfrak{d}_2 f|^2 \geq 8 \int_S K f^2$$

as desired. \square

Corollary 2.1.34. *Under the following mild assumptions on the Gauss curvature*

$$K = \frac{1}{r^2} + O\left(\frac{\epsilon}{r^2}\right), \quad \text{re}_\theta(K) = O\left(\frac{\epsilon}{r^2}\right),$$

the following holds.

1. *If $f \in \mathfrak{s}_0$ is orthogonal to the kernel of \mathfrak{d}_1^* , i.e. $\int_S f = 0$, then, we have*

$$\int_S |\mathfrak{d}_1^* f|^2 \geq 2 \int_S (1 + O(\epsilon)) K f^2. \quad (2.1.33)$$

2. *If $f \in \mathfrak{s}_1$ is orthogonal to the kernel of \mathfrak{d}_2^* , i.e. $\int_S f e^\Phi = 0$, then, we have*

$$\int_S |\mathfrak{d}_1 f|^2 \geq 6 \int_S (1 + O(\epsilon)) K f^2 \quad \text{and} \quad \int_S |\mathfrak{d}_2^* f|^2 \geq 4 \int_S (1 + O(\epsilon)) K f^2. \quad (2.1.34)$$

Proof. We start with the first assertion. If $f \in \mathfrak{s}_0$ satisfies $\int_S f = 0$ then, f is orthogonal to 1 which generates the kernel of \mathfrak{d}_1^* , and hence, f is in the image of \mathfrak{d}_1 , i.e. there exists $h \in \mathfrak{s}_1$ such that

$$f = \mathfrak{d}_1 h.$$

We deduce

$$\int_S (\mathfrak{d}_1^*(f))^2 = \int_S (\mathfrak{d}_1^* \mathfrak{d}_1 h)^2 = \int_S (\mathfrak{d}_1^* \mathfrak{d}_1)^2 h h.$$

Now, the above Poincaré inequality for ϕ_1 and the assumption on K implies a lower bound for the spectrum of the selfadjoint operator $\phi_1^* \phi_1$ by $2K(1 + O(\epsilon))$, and hence

$$\begin{aligned} \int_S (\phi_1^*(f))^2 &\geq 2 \int_S K(1 + O(\epsilon)) (\phi_1^* \phi_1) h h \\ &\geq 2 \int_S (1 + O(\epsilon)) (\phi_1 h)^2 \\ &\geq 2 \int_S (1 + O(\epsilon)) f^2 \end{aligned}$$

which yields the first assertion.

Assume now that $f \in \mathfrak{s}_1$ satisfies $\int_S f e^\Phi = 0$ i.e. f is orthogonal to e^Φ which generates the kernel of ϕ_2^* , and hence, f is in the image of ϕ_2 , i.e. there exists $h \in \mathfrak{s}_2$ such that

$$f = \phi_2 h.$$

We deduce

$$\begin{aligned} \int_S (\phi_1 f)^2 &= \int_S (\phi_1 \phi_2 h)^2 = \int_S \phi_1^* \phi_1 \phi_2 h \phi_2 h = \int_S (\phi_2 \phi_2^* + 2K) \phi_2 h \phi_2 h \\ &= \int_S (\phi_2^* \phi_2)^2 h h + \int_S 2K (\phi_2 h)^2. \end{aligned}$$

Now, the above Poincaré inequality for ϕ_2 and the assumption on K implies a lower bound for the spectrum of the selfadjoint operator $\phi_2^* \phi_2$ by $4K(1 + O(\epsilon))$, and hence

$$\begin{aligned} \int_S (\phi_1 f)^2 &\geq 4 \int_S K(1 + O(\epsilon)) (\phi_2^* \phi_2) h h + \int_S 2K (\phi_2 h)^2 \\ &\geq 6 \int_S (1 + O(\epsilon)) (\phi_2 h)^2 \\ &\geq 6 \int_S (1 + O(\epsilon)) f^2. \end{aligned}$$

Together with the fact that

$$\int_S (\phi_2^* f)^2 = \int_S \phi_2 \phi_2^* f f = \int_S (\phi_1^* \phi_1 - 2K) f f = \int_S (\phi_1 f)^2 - 2 \int_S K f^2,$$

this yields the second assertion and concludes the proof of the corollary. \square

Lemma 2.1.35. *Assume that*

$$K = \frac{1}{r^2} + O\left(\frac{\epsilon}{r^2}\right), \quad re_\theta(K) = O\left(\frac{\epsilon}{r^2}\right), \quad \int_S e^{2\Phi} = r^4 \left(\frac{8\pi}{3} + O(\epsilon)\right).$$

Then, If $f \in \mathfrak{s}_1$, we have the estimate,

$$\int_S |f|^2 \lesssim r^2 \int_S |\not{d}_2^* f|^2 + r^{-4} \left| \int_S e^\Phi f \right|^2. \quad (2.1.35)$$

More precisely,

$$f = \frac{\int_S f e^\Phi}{\int_S e^{2\Phi}} e^\Phi + f^\perp \quad (2.1.36)$$

with

$$\int_S |f^\perp|^2 \lesssim r^2 \int_S |\not{d}_2^* f|^2.$$

Proof. According to Corollary 2.1.34, see 2.1.34, if $f \in \mathfrak{s}_1$ is orthogonal to the kernel of \not{d}_2^* , i.e $\int_S f e^\Phi = 0$, then, we have

$$\int_S |\not{d}_2^* f|^2 \geq 4 \int_S (1 + O(\epsilon)) K f^2.$$

As a consequence $f^\perp = f - \left(\frac{\int_S f e^\Phi}{\int_S e^{2\Phi}} \right) e^\Phi$ verifies,

$$r^{-2} \int_S |f^\perp|^2 \lesssim \int_S |\not{d}_2^*(f^\perp)|^2 = \int_S |\not{d}_2^* f|^2$$

from which we derive,

$$\int_S \left| f - \left(\frac{\int_S f e^\Phi}{\int_S e^{2\Phi}} \right) e^\Phi \right|^2 \lesssim r^2 \int_S |\not{d}_2^* f|^2$$

or,

$$\begin{aligned} \int_S |f|^2 &= r^2 \int_S |\not{d}_2^* f|^2 + \left| \int_S e^\Phi f \right|^2 \frac{1}{\int_S e^{2\Phi}} \\ &\leq r^2 \int_S |\not{d}_2^* f|^2 + r^{-4} \left| \int_S e^\Phi f \right|^2 \end{aligned}$$

as desired. \square

Higher derivative operators and spaces

Definition 2.1.36. Given f a k -reduced scalar and s a positive integer we define,

$$\mathfrak{D}^s f = \begin{cases} r^{2p} \mathbb{A}_k^p, & \text{if } s = 2p, \\ r^{2p+1} \mathbb{A}_k \mathbb{A}_k^p, & \text{if } s = 2p + 1. \end{cases} \quad (2.1.37)$$

We also define the norms,

$$\|f\|_{\mathfrak{h}_s(S)} := \sum_{i=0}^s \|\mathfrak{D}^i f\|_{L^2(S)}. \quad (2.1.38)$$

Lemma 2.1.37. Assume the Gauss curvature K of S verifies the condition,

$$K = \frac{1}{r^2} + O(\epsilon), \quad |r^i \nabla^i K| = O(\epsilon), \quad 1 \leq i \leq [s/2] + 1.$$

Then, the following holds.

1. If f is a k -scalar, reduced from ${}^{(1+3)}f$, we have,

$$\|f\|_{\mathfrak{h}_s(S)} \sim \sum_{j=0}^s r^j \|\nabla^j f\|_{L^2(S)} \quad (2.1.39)$$

where ∇ denotes the usual covariant derivative operator on S .

2. Equivalently, the norm $r^{-s} \|f\|_{\mathfrak{h}_s(S)}$ of a reduced scalar $f \in \mathfrak{s}_s(S)$ can be defined as the sum of L^2 norms of any allowable sequence of Hodge operators $\mathbb{A}_a, \mathbb{A}_a^*$ applied to f .

Proof. For $s = 1, 2$ the proof of the first part follows immediately from Proposition 2.1.30. For higher s the proof follows, step by step, by a simple commutation argument between covariant derivatives and \mathbb{A}_k and applications of Proposition 2.1.30. The proof of the second part follows from our reduced elliptic estimates and definition of the reduced Hodge operators. \square

As a consequence of the lemma we can derive the reduce form of the standard Sobolev and product Sobolev inequalities. Before stating the result we pause to define the product of two reduced scalars.

Definition 2.1.38. Let $f \in \mathfrak{s}_a$ be reduced from an \mathcal{S}_a tensor and $g \in \mathfrak{s}_b$ reduced from an \mathcal{S}_b tensor. We define the product $f \cdot g$ to be the reduction of any product between the corresponding tensors on S , i.e. any contraction of the tensor product between them. Thus $f \cdot g \in \mathfrak{s}_{a+b-2c}$ where c denotes the number of indices affected by the contraction.

Examples. Here are the most relevant examples for us.

- $f \in \mathfrak{s}_0, g \in \mathfrak{s}_k$ in which case $f \cdot g \in \mathfrak{s}_k$ and equals fg .
- $f \in \mathfrak{s}_1, g \in \mathfrak{s}_k$ in which case $f \cdot g \in \mathfrak{s}_{k-1}$ or $f \cdot g \in \mathfrak{s}_{k+1}$ and in both cases $f \cdot g = fg$ as simple product of the reduced scalars.
- $f \in \mathfrak{s}_2, g \in \mathfrak{s}_k$ in which case $f \cdot g \in \mathfrak{s}_{k-2}$ or $f \cdot g \in \mathfrak{s}_k$ or $f \cdot g \in \mathfrak{s}_{k+2}$. In the first case $f \cdot g = 2fg$. In the second case and third cases $f \cdot g = fg$ as simple product of the reduced scalars.

Lemma 2.1.39. Let $f \in \mathfrak{s}_a(S), g \in \mathfrak{s}_b(S), a \geq b, a > 0$, and $f \cdot g \in \mathfrak{s}_{a+b-2c}$ where $0 \leq c \leq \frac{1}{2}(a-b)$ denotes the order of contraction. Then,

$$\begin{aligned} \mathfrak{d}_{a+b-2c}(fg) &= f \mathfrak{d}_b g + g \left(\left(1 - \frac{c}{a}\right) \mathfrak{d}_a f - \frac{c}{a} \mathfrak{d}_{a+1}^* f \right), \\ \mathfrak{d}_{a+b-2c+1}^*(fg) &= f \mathfrak{d}_{b+1}^* g + \left(\left(-1 + \frac{c}{a}\right) \mathfrak{d}_a f + \frac{c}{a} \mathfrak{d}_{a+1}^* f \right). \end{aligned} \tag{2.1.40}$$

Proof. Assume $a \geq b$ and $c \leq \frac{a-b}{2}$. We write,

$$\mathfrak{d}_{a+b-2c}(fg) = f \mathfrak{d}_b g + g(e_\theta f + (a-2c)e_\theta(\Phi)f).$$

We look for reals A, B with $A+B=1$ such that

$$e_\theta f + (a-2c)e_\theta(\Phi)f = A \mathfrak{d}_a f - B \mathfrak{d}_{a+1}^* f = e_\theta f + a(A-B)e_\theta \Phi f.$$

Therefore,

$$a(1-2B) = a-2c$$

i.e. $B = \frac{c}{a}, A = 1 - \frac{c}{a}$ and we derive,

$$\mathfrak{d}_{a+b-2c}(fg) = f \mathfrak{d}_b g + g \left(\left(1 - \frac{c}{a}\right) \mathfrak{d}_a f - \frac{c}{a} \mathfrak{d}_{a+1}^* f \right).$$

Also,

$$\mathfrak{d}_{a+b-2c+1}^*(fg) = f \mathfrak{d}_{b+1}^* g + g(-e_\theta f + (a-2c)e_\theta(\Phi)f).$$

As before we write, with $A + B = -1$

$$-e_\theta(f) + (a - 2c)e_\theta(\Phi)f = A\mathfrak{d}_a f - B\mathfrak{d}_{a+1}^* f = -e_\theta f + a(A - B)e_\theta\Phi f.$$

Hence,

$$-a(-1 - 2B) = (a - 2c)$$

i.e. $B = -\frac{c}{a}$, $A = -1 + \frac{c}{a}$. Hence,

$$\mathfrak{d}_{a+b-2c+1}^*(fg) = f\mathfrak{d}_{b+1}^*g + \left((-1 + \frac{c}{a})\mathfrak{d}_a f + \frac{c}{a}\mathfrak{d}_{a+1}^* f\right)$$

as desired. \square

Proposition 2.1.40. *The following results hold true for k -reduced scalars on S ,*

1. *If $f \in \mathfrak{s}_k$ we have,*

$$\|f\|_{L^\infty(S)} \lesssim r^{-1}\|f\|_{\mathfrak{h}_2(S)}.$$

2. *Given two reduced scalars f, g we have,*

$$\|f \cdot g\|_{\mathfrak{h}_s(S)} \lesssim r^{-1} \left(\|f\|_{\mathfrak{h}_{[s/2]+2}(S)} \|g\|_{\mathfrak{h}_s(S)} + \|g\|_{\mathfrak{h}_{[s/2]+2}(S)} \|f\|_{\mathfrak{h}_s(S)} \right)$$

where $[s/2]$ denotes the largest integer smaller than $s/2$.

Proof. Both statements are classical for $\mathcal{S}_k(S)$ tensors with respect to the norm on the right hand side of (2.1.39). A direct proof can also be derived using Lemma 2.1.39 and the equivalence definition of the $\mathfrak{h}_s(S)$ norms. \square

S -averages

Definition 2.1.41. *Given any $f \in \mathfrak{s}_0$ we denote its average by,*

$$\bar{f} := \frac{1}{|S|} \int_S f, \quad \check{f} := f - \bar{f}.$$

The following follows immediately from the definition.

Lemma 2.1.42. *For any two scalar reduced scalars f and g in \mathfrak{s}_0 we have*

$$\overline{fg} = \overline{f}\overline{g} + \overline{f\check{g}},$$

and,

$$fg - \overline{fg} = \check{f}\overline{g} + \overline{f}\check{g} + (f\check{g} - \overline{f\check{g}}).$$

Remark 2.1.43. *In view of the notations above, we may rewrite the Poincaré inequality for ϕ_1^* as follows. Under mild assumptions on the Gauss curvature ($K = r^{-2} + O(\epsilon r^{-2})$, $re_\theta(K) = O(\epsilon r^{-2})$), we have for any $f \in \mathfrak{s}_0$*

$$\int_S |\phi_1^* f|^2 \geq 2 \int_S (1 + O(\epsilon)) K (f\check{f})^2.$$

2.1.5 Invariant S -foliations

In this section we record the main equations associated to general, \mathbf{Z} -invariant Einstein vacuum spacetimes $(\mathcal{M}, \mathbf{g})$. We start by recalling the spacetime framework of [17] and then we show how the null structure and Bianchi identities simplify in the reduced picture. Throughout this section we consider given an invariant S -foliation⁸ and a fixed adapted null pair e_3, e_4 , i.e. future directed \mathbf{Z} -invariant, polarized, null vectors orthogonal to the leaves S of the foliation such as $\mathbf{g}(e_3, e_4) = -2$.

Definition 2.1.44. *We denote by $\mathcal{S}_k(\mathcal{M})$ the set of k -covariant polarized tensors on \mathcal{M} tangent to the S -foliation and which restrict to $\mathcal{S}_k(S)$ on any S -surface of the foliation and by $\mathfrak{s}_k(\mathcal{M})$ their corresponding reductions.*

Spacetime null decompositions

Following [17] we define the spacetime Ricci coefficients,

$$\begin{aligned} {}^{(1+3)}\chi_{AB} &:= \mathbf{g}(\mathbf{D}_A e_4, e_B), & {}^{(1+3)}\zeta_A &:= \frac{1}{2} \mathbf{g}(\mathbf{D}_4 e_4, e_A), & {}^{(1+3)}\eta_A &:= \frac{1}{2} \mathbf{g}(\mathbf{D}_3 e_4, e_A), \\ {}^{(1+3)}\zeta_A &:= \frac{1}{2} \mathbf{g}(\mathbf{D}_A e_4, e_3), & {}^{(1+3)}\omega &:= \frac{1}{4} \mathbf{g}(\mathbf{D}_4 e_4, e_3), \end{aligned} \tag{2.1.41}$$

and interchanging e_3, e_4 ,

$$\begin{aligned} {}^{(1+3)}\underline{\chi}_{AB} &:= \mathbf{g}(\mathbf{D}_A e_3, e_B), & {}^{(1+3)}\underline{\zeta}_A &:= \frac{1}{2} \mathbf{g}(\mathbf{D}_3 e_3, e_A), & {}^{(1+3)}\underline{\eta}_A &:= \frac{1}{2} \mathbf{g}(\mathbf{D}_4 e_3, e_A), \\ {}^{(1+3)}\zeta_A &:= -\frac{1}{2} \mathbf{g}(\mathbf{D}_A e_3, e_4), & {}^{(1+3)}\underline{\omega} &:= \frac{1}{4} \mathbf{g}(\mathbf{D}_3 e_3, e_4). \end{aligned} \tag{2.1.42}$$

⁸From now on, an invariant S foliation is automatically assumed to be a \mathbf{Z} invariant polarized foliation.

We also define the spacetime null curvature components,

$$\begin{aligned}
{}^{(1+3)}\alpha_{AB} &:= \mathbf{R}_{A4B4}, & {}^{(1+3)}\beta_A &:= \frac{1}{2}\mathbf{R}_{A434}, & {}^{(1+3)}\rho &:= \frac{1}{4}\mathbf{R}_{3434}, \\
{}^{(1+3)}\underline{\alpha}_{AB} &:= \mathbf{R}_{A3B3}, & {}^{(1+3)}\underline{\beta}_A &:= \frac{1}{2}\mathbf{R}_{A334}, & {}^{(1+3)}\star\rho &:= \frac{1}{4}\star\mathbf{R}_{3434}.
\end{aligned} \tag{2.1.43}$$

Reduced null decompositions

We define the spacetime Ricci coefficients as follows

Definition 2.1.45 (*Ricci coefficients*). *Let e_3, e_4, e_θ be a reduced null frame. The following scalars*

$$\begin{aligned}
\chi &= g(D_\theta e_4, e_\theta), & \underline{\chi} &= g(D_\theta e_3, e_\theta), \\
\eta &= \frac{1}{2}g(D_3 e_4, e_\theta), & \underline{\eta} &= \frac{1}{2}g(D_4 e_3, e_\theta), \\
\xi &= \frac{1}{2}g(D_4 e_4, e_\theta), & \underline{\xi} &= \frac{1}{2}g(D_3 e_3, e_\theta), \\
\omega &= \frac{1}{4}g(D_4 e_4, e_3), & \underline{\omega} &= \frac{1}{4}g(D_3 e_3, e_4), \\
\zeta &= \frac{1}{2}g(D_\theta e_4, e_3),
\end{aligned} \tag{2.1.44}$$

are called the Ricci coefficients associated to our canonical null pair.

Lemma 2.1.46. *The following lemma follows easily from the definitions,*

$$\begin{aligned}
D_4 e_4 &= -2\omega e_4 + 2\xi e_\theta, & D_3 e_3 &= -2\underline{\omega} e_3 + 2\underline{\xi} e_\theta, \\
D_4 e_3 &= 2\omega e_3 + 2\underline{\eta} e_\theta, & D_3 e_4 &= 2\underline{\omega} e_4 + 2\underline{\eta} e_\theta, \\
D_4 e_\theta &= \underline{\eta} e_4 + \xi e_3, & D_3 e_\theta &= \underline{\xi} e_4 + \eta e_3, \\
D_\theta e_4 &= -\zeta e_4 + \chi e_\theta, & D_\theta e_3 &= \zeta e_3 + \underline{\chi} e_\theta, \\
D_\theta e_\theta &= \frac{1}{2}\underline{\chi} e_4 + \frac{1}{2}\chi e_3.
\end{aligned} \tag{2.1.45}$$

Definition 2.1.47. *The null components of the Ricci curvature tensor⁹ of the metric g are denoted by*

$$R_{33} = \underline{\alpha}, \quad R_{44} = \alpha, \quad R_{3\theta} = \underline{\beta}, \quad R_{4\theta} = \beta, \quad R_{\theta\theta} = R_{34} = \rho, \quad R_{34} = \rho.$$

⁹Recall that the scalar curvature of the reduced metric g vanishes, $R = 0$, and hence $R_{34} = R_{\theta\theta}$.

Comparison to the space-time frame

Let e_3, e_4, e_θ be a null frame for the reduced metric g and $e_3, e_4, e_\theta, e_\varphi = X^{-1/2}\partial_\varphi$ the augmented adapted 3 + 1 frame for \mathbf{g} . Recall that we have denoted,

$${}^{(1+3)}\chi, {}^{(1+3)}\xi, {}^{(1+3)}\eta, {}^{(1+3)}\underline{\eta}, {}^{(1+3)}\zeta, {}^{(1+3)}\omega, {}^{(1+3)}\underline{\chi}, {}^{(1+3)}\underline{\xi}, {}^{(1+3)}\underline{\omega},$$

the standard (as defined in [17]) space-time Ricci coefficients and by

$${}^{(1+3)}\alpha, {}^{(1+3)}\beta, {}^{(1+3)}\rho, {}^{(1+3)}\star\rho, {}^{(1+3)}\underline{\beta}, {}^{(1+3)}\underline{\alpha},$$

the null decomposition of the curvature tensor \mathbf{R} .

Proposition 2.1.48. *The following relations between the spacetime and reduced Ricci and curvature null components hold true,*

- We have,

$$\begin{aligned} {}^{(1+3)}\chi_{\theta\theta} &= \chi, & {}^{(1+3)}\chi_{\theta\varphi} &= 0, & {}^{(1+3)}\chi_{\varphi\varphi} &= e_4(\Phi), \\ {}^{(1+3)}\underline{\chi}_{\theta\theta} &= \chi, & {}^{(1+3)}\chi_{\theta\varphi} &= 0, & {}^{(1+3)}\chi_{\varphi\varphi} &= e_3(\Phi), \\ {}^{(1+3)}\alpha_{\theta\theta} &= \alpha, & {}^{(1+3)}\alpha_{\theta\varphi} &= 0, & {}^{(1+3)}\alpha_{\varphi\varphi} &= -\alpha, \\ {}^{(1+3)}\underline{\alpha}_{\theta\theta} &= \underline{\alpha}, & {}^{(1+3)}\underline{\alpha}_{\theta\varphi} &= 0, & {}^{(1+3)}\underline{\alpha}_{\varphi\varphi} &= -\alpha. \end{aligned}$$

- All e_φ components of ${}^{(1+3)}\eta, {}^{(1+3)}\underline{\eta}, {}^{(1+3)}\zeta, {}^{(1+3)}\xi, {}^{(1+3)}\underline{\xi}, {}^{(1+3)}\beta, {}^{(1+3)}\underline{\beta}$ vanish and,

$${}^{(1+3)}\eta_\theta = \eta, \quad {}^{(1+3)}\underline{\eta}_\theta = \underline{\eta}, \quad {}^{(1+3)}\zeta_\theta = \zeta, \quad {}^{(1+3)}\xi_\theta = \xi, \quad {}^{(1+3)}\underline{\xi}_\theta = \underline{\xi},$$

and¹⁰

$${}^{(1+3)}\beta_\theta = \beta, \quad {}^{(1+3)}\underline{\beta}_\theta = -\underline{\beta}.$$

Also,

$${}^{(1+3)}\omega = \omega, \quad {}^{(1+3)}\underline{\omega} = \underline{\omega}, \quad {}^{(1+3)}\rho = \rho, \quad {}^{(1+3)}\star\rho = 0.$$

- We have,

$${}^{(1+3)}tr\chi = \chi + e_4(\Phi), \quad {}^{(1+3)}tr\underline{\chi} = \underline{\chi} + e_3(\Phi).$$

Recalling, see definition in [17],

$${}^{(1+3)}\widehat{\chi}_{AB} = {}^{(1+3)}\chi_{AB} - \frac{1}{2}({}^{(1+3)}tr\chi)\not\partial_{AB}, \quad {}^{(1+3)}\widehat{\underline{\chi}}_{AB} = {}^{(1+3)}\underline{\chi}_{AB} - \frac{1}{2}({}^{(1+3)}tr\underline{\chi})\not\partial_{AB},$$

we have

$${}^{(1+3)}\widehat{\chi}_{\theta\theta} = \frac{1}{2}(\chi - e_4(\Phi)), \quad {}^{(1+3)}\widehat{\underline{\chi}}_{\theta\theta} = \frac{1}{2}(\underline{\chi} - e_3(\Phi)).$$

¹⁰Note the change of sign for the $\underline{\beta}$ component.

Proof. We check only the less obvious relations such as those involving the null components of curvature. Using (2.1.5) and (2.1.6) we deduce,

$$\begin{aligned}
{}^{(1+3)}\alpha_{\theta\theta} &= \mathbf{R}_{\theta 4\theta 4} = R_{\theta 4\theta 4} = g_{\theta\theta}R_{44} = \alpha, \\
{}^{(1+3)}\alpha_{\theta\varphi} &= \mathbf{R}_{\theta 4\varphi 4} = 0, \\
{}^{(1+3)}\alpha_{\varphi\varphi} &= \mathbf{R}_{\varphi 4\varphi 4} = -R_{44} = -\alpha, \\
2{}^{(1+3)}\beta_{\theta} &= \mathbf{R}_{\theta 343} = R_{\theta 343} = -g_{34}R_{\theta 3} = 2\beta, \\
2{}^{(1+3)}\beta_{\varphi} &= \mathbf{R}_{\varphi 343} = 0, \\
4{}^{(1+3)}\rho &= \mathbf{R}_{3434} = R_{3434} = -2g_{34}R_{34} = 4\rho, \\
4{}^{(1+3)}\star\rho &= \star\mathbf{R}_{3434} = 0, \\
2{}^{(1+3)}\underline{\beta}_{\theta} &= \mathbf{R}_{\theta 334} = R_{\theta 334} = g_{34}R_{\theta 3} = -2\underline{\beta}.
\end{aligned}$$

□

Definition 2.1.49. We introduce the notation,

$$\begin{aligned}
\vartheta &:= \chi - e_4(\Phi), & \kappa &:= {}^{(1+3)}tr\chi = \chi + e_4(\Phi), \\
\underline{\vartheta} &:= \underline{\chi} - e_3(\Phi), & \underline{\kappa} &:= {}^{(1+3)}tr\underline{\chi} = \underline{\chi} + e_3(\Phi).
\end{aligned}$$

Thus,

$${}^{(1+3)}\widehat{\chi}_{\theta\theta} = {}^{(1+3)}\widehat{\chi}_{\varphi\varphi} = \frac{1}{2}\vartheta, \quad {}^{(1+3)}\widehat{\underline{\chi}}_{\theta\theta} = {}^{(1+3)}\widehat{\underline{\chi}}_{\varphi\varphi} = \frac{1}{2}\underline{\vartheta}.$$

In particular, $\chi = \frac{1}{2}(\vartheta + \kappa)$ and $\underline{\chi} = \frac{1}{2}(\underline{\vartheta} + \underline{\kappa})$.

Remark 2.1.50. In view of Proposition 2.1.48 we have,

1. The quantities $\kappa, \underline{\kappa}, \omega, \underline{\omega}, \rho$ are reduced scalars in \mathfrak{s}_0 .
2. The quantities $\eta, \underline{\eta}, \zeta, \underline{\zeta}, \xi, \underline{\xi}, \beta, \underline{\beta}$ are reduced scalars in \mathfrak{s}_1 .
3. The quantities $\vartheta, \underline{\vartheta}, \alpha, \underline{\alpha}$ are reduced scalars in \mathfrak{s}_2 .

Commutation identities

We record first the commutation relations between the elements of the frame,

$$\begin{aligned}
[e_{\theta}, e_3] &= \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_{\theta} + (\zeta - \eta)e_3 - \underline{\xi}e_4, \\
[e_{\theta}, e_4] &= \frac{1}{2}(\kappa + \vartheta)e_{\theta} - (\zeta + \eta)e_4 - \xi e_3, \\
[e_3, e_4] &= 2\underline{\omega}e_4 - 2\omega e_3 + 2(\eta - \underline{\eta})e_{\theta}.
\end{aligned}$$

Lemma 2.1.51. *The following commutation formulae hold true for reduced scalars.*

1. If $f \in \mathfrak{s}_k$,

$$\begin{aligned}
[\mathfrak{d}_k, e_3]f &= \frac{1}{2}\underline{\kappa}\mathfrak{d}_k f + \underline{Com}_k(f), \\
\underline{Com}_k(f) &= -\frac{1}{2}\underline{\vartheta}\mathfrak{d}_{k+1}^* f + (\zeta - \eta)e_3 f - k\underline{\eta}e_3\Phi f - \underline{\xi}(e_4 f + ke_4(\Phi)f) - k\underline{\beta}f, \\
[\mathfrak{d}_k, e_4] &= \frac{1}{2}\kappa\mathfrak{d}_k f + Com_k(f), \\
Com_k(f) &= -\frac{1}{2}\vartheta\mathfrak{d}_{k+1}^* f - (\zeta + \underline{\eta})e_4 f - k\underline{\eta}e_4\Phi f - \xi(e_3 f + ke_3(\Phi)f) - k\beta f.
\end{aligned} \tag{2.1.46}$$

2. If $f \in \mathfrak{s}_{k-1}$

$$\begin{aligned}
[\mathfrak{d}_k^*, e_3]f &= \frac{1}{2}\underline{\kappa}\mathfrak{d}_k^* f + \underline{Com}_k^*(f), \\
\underline{Com}_k^*(f) &= -\frac{1}{2}\underline{\vartheta}\mathfrak{d}_{k-1} f - (\zeta - \eta)e_3 f - (k-1)\underline{\eta}e_3\Phi f + \underline{\xi}(e_4 f - (k-1)e_4(\Phi)f) \\
&\quad - (k-1)\underline{\beta}f, \\
[\mathfrak{d}_k^*, e_4]f &= \frac{1}{2}\kappa\mathfrak{d}_k^* f + Com_k^*(f), \\
Com_k^*(f) &= -\frac{1}{2}\vartheta\mathfrak{d}_{k-1} f + (\zeta + \underline{\eta})e_4 f - (k-1)\underline{\eta}e_4\Phi f + \xi(e_3 f - (k-1)e_3(\Phi)f) \\
&\quad - (k-1)\beta f.
\end{aligned} \tag{2.1.47}$$

Proof. We write,

$$[e_\theta + ke_\theta(\Phi), e_3]f = [e_\theta, e_3]f - k(e_3 e_\theta \Phi)f.$$

Recall that (see (2.1.4)), $D_a D_b \Phi = R_{ab} - D_a \Phi D_b \Phi$. Hence,

$$e_3 e_\theta \Phi - \underline{\xi}e_4 \Phi - \eta e_3 \Phi = D_3 D_\theta \Phi = R_{3\theta} - D_3 \Phi D_\theta \Phi = \underline{\beta} - e_3 \Phi e_\theta \Phi.$$

Thus,

$$e_3 e_\theta \Phi = \underline{\beta} - e_3 \Phi e_\theta \Phi + \eta e_3(\Phi) + \underline{\xi}e_4 \Phi.$$

We deduce, since $e_3\Phi = \frac{1}{2}(\underline{\kappa} - \underline{\vartheta})$,

$$\begin{aligned}
[e_\theta + ke_\theta(\Phi), e_3]f &= [e_\theta, e_3]f - k(\underline{\beta} - e_3\Phi e_\theta\Phi + \eta e_3(\Phi) + \underline{\xi}e_4\Phi)f \\
&= \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta f + (\zeta - \eta)e_3f - \underline{\xi}e_4f - k(\underline{\beta} - e_3\Phi e_\theta\Phi + \eta e_3(\Phi) + \underline{\xi}e_4\Phi)f \\
&= \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta f + k\frac{1}{2}(\underline{\kappa} - \underline{\vartheta})e_\theta\Phi f \\
&+ (\zeta - \eta)e_3f - k\eta e_3\Phi f - \underline{\xi}(e_4f + ke_4(\Phi)f) - k\underline{\beta}f \\
&= \frac{1}{2}\underline{\kappa}\not{d}_k f + \frac{1}{2}\underline{\vartheta}(e_\theta f - ke_\theta\Phi f) \\
&+ (\zeta - \eta)e_3f - k\eta e_3\Phi f - \underline{\xi}(e_4f + ke_4(\Phi)f) - k\underline{\beta}f
\end{aligned}$$

i.e., recalling the definition of \not{d}_{k+1}^* ,

$$\begin{aligned}
[e_\theta + ke_\theta(\Phi), e_3]f &= \frac{1}{2}\underline{\kappa}\not{d}_k f + \underline{Com}_k(f), \\
\underline{Com}(f)_k &= -\frac{1}{2}\underline{\vartheta}\not{d}_{k+1}^* f + (\zeta - \eta)e_3f - k\eta e_3\Phi f - \underline{\xi}(e_4f + ke_4(\Phi)f) - k\underline{\beta}f.
\end{aligned}$$

The other commutation formulae are proved in the same manner. \square

2.1.6 Schwarzschild spacetime

In standard coordinates the Schwarzschild metric has the form,

$$ds^2 = -\Upsilon dt^2 + \Upsilon^{-1} dr^2 + r^2 d\theta^2 + X^2 d\varphi^2, \quad (2.1.48)$$

where,

$$\Upsilon := 1 - \frac{2m}{r}, \quad X = r^2 \sin^2 \theta.$$

We denote by \mathbf{T} the stationary Killing vectorfield $\mathbf{T} = \partial_t$ and by $\mathbf{Z} = \partial_\varphi$ the axial symmetric one. Recall the regular, \mathbf{Z} -invariant optical functions in the exterior region $r \geq 2m$ of Schwarzschild

$$u = t - r_*, \quad \underline{u} = t + r_*, \quad \frac{dr_*}{dr} = \Upsilon^{-1} \quad (2.1.49)$$

with $r_* = r + 2m \log(\frac{r}{2m} - 1)$. The corresponding null geodesic generators are,

$$\underline{L} := -g^{ab}\partial_a v \partial_b = \Upsilon^{-1}\partial_t - \partial_r, \quad L := -g^{ab}\partial_a u \partial_b = \Upsilon^{-1}\partial_t + \partial_r. \quad (2.1.50)$$

Clearly,

$$g(L, L) = g(\underline{L}, \underline{L}) = 0, \quad g(L, \underline{L}) = -2\Upsilon^{-1}, \quad D_L L = D_{\underline{L}} \underline{L} = 0.$$

Definition 2.1.52. We can use the null geodesic generators L, \underline{L} to define the following canonical null pairs. In all cases all curvature components vanish identically except,

$${}^{(1+3)}\rho = -\frac{2m}{r^3}. \quad (2.1.51)$$

1. The null frame (e_3, e_4) for which e_3 is geodesic (which is regular towards the future for all $r > 0$) is given by

$$e_3 = \underline{L} = \Upsilon^{-1}\partial_t - \partial_r, \quad e_4 = \Upsilon L = \partial_t + \Upsilon\partial_r, \quad \Upsilon = 1 - \frac{2m}{r}. \quad (2.1.52)$$

All Ricci coefficients vanish except,

$$\chi = \frac{\Upsilon}{r}, \quad \underline{\chi} = -\frac{1}{r}, \quad \omega = -\frac{m}{r^2}, \quad \underline{\omega} = 0.$$

2. The null frame (e_3, e_4) for which e_4 is geodesic.

$$e_4 = L = \Upsilon^{-1}\partial_t + \partial_r, \quad e_3 = \Upsilon \underline{L} = \partial_t - \Upsilon\partial_r.$$

All Ricci coefficients vanish except,

$$\chi = \frac{1}{r}, \quad \underline{\chi} = -\frac{\Upsilon}{r}, \quad \omega = 0, \quad \underline{\omega} = \frac{m}{r^2}.$$

Note that the null pair (2.1.52) is regular along the future event horizon as can be easily seen by studying the behavior¹¹. of future directed ingoing null geodesics near $r = 2m$.

2.2 Main Equations

In this section we translate the null structure and null Bianchi identities associated to an S -foliation in the reduced picture. We start with general, \mathbf{Z} -invariant, S foliation . We then consider the special case of geodesic foliations.

2.2.1 Main equations for general S -foliations

We consider a fixed \mathbf{Z} -invariant S -foliation with a fixed \mathbf{Z} -invariant null frame e_3, e_4 .

¹¹i.e. the null geodesics in the direction of \underline{L} reach the horizon in finite proper time. Note that, on the other hand, the past null geodesics in the direction of L still meet the horizon in infinite proper time.

Null structure equations

We simply translate the well known spacetime null structure equations (see¹² proposition 7.4.1 in [17]) in the reduced picture. Thus the spacetime equation¹³,

$$\nabla_3 \widehat{\chi} + \text{tr} \chi \widehat{\chi} = \nabla \widehat{\otimes} \xi - 2\omega \widehat{\chi} + (\eta + \underline{\eta} - 2\zeta) \widehat{\otimes} \xi - \underline{\alpha}$$

becomes¹⁴,

$$e_3(\underline{\vartheta}) + \underline{\kappa} \underline{\vartheta} = 2(e_\theta(\underline{\xi}) - e_\theta(\Phi)\underline{\xi}) - 2\omega \underline{\vartheta} + 2(\eta + \underline{\eta} - 2\zeta) \underline{\xi} - 2\underline{\alpha}. \quad (2.2.1)$$

The spacetime equation,

$$e_3(\text{tr} \chi) + \frac{1}{2} \text{tr} \chi^2 = 2\text{div} \xi - 2\omega \text{tr} \chi + 2\xi \cdot (\eta + \underline{\eta} - 2\zeta) - \widehat{\chi} \cdot \widehat{\chi}$$

becomes,

$$e_3(\underline{\kappa}) + \frac{1}{2} \underline{\kappa}^2 + 2\omega \underline{\kappa} = 2(e_\theta \underline{\xi} + e_\theta(\Phi)\underline{\xi}) + 2(\eta + \underline{\eta} - 2\zeta) \underline{\xi} - \frac{1}{2} \underline{\vartheta} \underline{\vartheta}. \quad (2.2.2)$$

The spacetime equation,

$$\nabla_4 \widehat{\chi} + \frac{1}{2} \text{tr} \chi \widehat{\chi} = \nabla \widehat{\otimes} \underline{\eta} + 2\omega \widehat{\chi} - \frac{1}{2} \text{tr} \chi \widehat{\chi} + \xi \widehat{\otimes} \underline{\xi} + \underline{\eta} \widehat{\otimes} \underline{\eta}$$

becomes,

$$e_4 \underline{\vartheta} + \frac{1}{2} \underline{\kappa} \underline{\vartheta} - 2\omega \underline{\vartheta} = 2(e_\theta \underline{\eta} - e_\theta(\Phi)\underline{\eta}) - \frac{1}{2} \text{tr} \chi \underline{\vartheta} + 2(\xi \underline{\xi} + \underline{\eta}^2).$$

The spacetime equation,

$$\nabla_4 \text{tr} \chi + \frac{1}{2} \text{tr} \chi \text{tr} \chi = 2\text{div} \underline{\eta} + 2\rho + 2\omega \text{tr} \chi - \widehat{\chi} \cdot \widehat{\chi} + 2(\xi \cdot \underline{\xi} + \underline{\eta} \cdot \underline{\eta})$$

becomes,

$$e_4(\underline{\kappa}) + \frac{1}{2} \underline{\kappa} \underline{\kappa} - 2\omega \underline{\kappa} = 2(e_\theta \underline{\eta} + e_\theta(\Phi)\underline{\eta}) + 2\rho - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} + 2(\xi \underline{\xi} + \underline{\eta} \underline{\eta}).$$

The spacetime equation,

$$\nabla_3 \zeta = -\underline{\beta} - 2\nabla \omega - \widehat{\chi} \cdot (\zeta + \eta) - \frac{1}{2} \text{tr} \chi (\zeta + \eta) + 2\omega (\zeta - \eta) + (\widehat{\chi} + \frac{1}{2} \text{tr} \chi) \underline{\xi} + 2\omega \underline{\xi}$$

¹²Note however that the notation in [17] are different, see section 7.3 for the definitions.

¹³For convenience we drop the ⁽¹⁺³⁾ labels in what follows.

¹⁴recall that ⁽¹⁺³⁾ $\widehat{\chi}_{\theta\theta} = \frac{1}{2} \underline{\vartheta}$

becomes (note that ${}^{(1+3)}\underline{\beta} = -\underline{\beta}$!),

$$e_3\underline{\zeta} + \frac{1}{2}\underline{\kappa}(\underline{\zeta} + \underline{\eta}) - 2\underline{\omega}(\underline{\zeta} - \underline{\eta}) = \underline{\beta} - 2e_\theta(\underline{\omega}) + 2\underline{\omega}\underline{\xi} + \frac{1}{2}\underline{\kappa}\underline{\xi} - \frac{1}{2}\underline{\vartheta}(\underline{\zeta} + \underline{\eta}) + \frac{1}{2}\underline{\vartheta}\underline{\xi}.$$

The spacetime equation,

$$\nabla_4\underline{\xi} - \nabla_3\underline{\eta} = -\underline{\beta} + 4\underline{\omega}\underline{\xi} + \widehat{\chi} \cdot (\underline{\eta} - \underline{\eta}) + \frac{1}{2}\text{tr}\underline{\chi}(\underline{\eta} - \underline{\eta})$$

becomes¹⁵,

$$e_4(\underline{\xi}) - e_3(\underline{\eta}) = \underline{\beta} + 4\underline{\omega}\underline{\xi} + \frac{1}{2}\underline{\kappa}(\underline{\eta} - \underline{\eta}) + \frac{1}{2}\underline{\vartheta}(\underline{\eta} - \underline{\eta}).$$

The spacetime equation,

$$\nabla_4\underline{\omega} + \nabla_3\underline{\omega} = \rho + 4\underline{\omega}\underline{\omega} + \underline{\xi} \cdot \underline{\xi} + \underline{\zeta} \cdot (\underline{\eta} - \underline{\eta}) - \underline{\eta} \cdot \underline{\eta}$$

becomes

$$e_4\underline{\omega} + e_3\underline{\omega} = \rho + 4\underline{\omega}\underline{\omega} + \underline{\xi}\underline{\xi} + \underline{\zeta}(\underline{\eta} - \underline{\eta}) - \underline{\eta}\underline{\eta}.$$

The spacetime Codazzi equation,

$${}^{(1+3)}\text{div} {}^{(1+3)}\widehat{\chi} = {}^{(1+3)}\underline{\beta} + \frac{1}{2}({}^{(1+3)}\nabla \text{tr}\underline{\chi} - {}^{(1+3)}\text{tr}\underline{\chi} {}^{(1+3)}\underline{\zeta}) + {}^{(1+3)}\widehat{\chi} \cdot {}^{(1+3)}\underline{\zeta}$$

becomes¹⁶,

$$\frac{1}{2}(e_\theta(\underline{\vartheta}) + 2e_\theta(\Phi)\underline{\vartheta}) = -\underline{\beta} + \frac{1}{2}(e_\theta(\underline{\kappa}) - \underline{\kappa}\underline{\zeta}) + \frac{1}{2}\underline{\vartheta}\underline{\zeta}.$$

The Gauss equation,

$$K = -\frac{1}{4}{}^{(1+3)}\text{tr}\underline{\chi} {}^{(1+3)}\text{tr}\underline{\chi} + \frac{1}{2}{}^{(1+3)}\widehat{\chi} {}^{(1+3)}\widehat{\chi} - {}^{(1+3)}\rho$$

becomes,

$$K = -\frac{1}{4}\underline{\kappa}\underline{\kappa} + \frac{1}{4}\underline{\vartheta}\underline{\vartheta} - \rho.$$

We summarize the results in the following proposition.

¹⁵Note that ${}^{(1+3)}\underline{\beta} = -\underline{\beta}$ and $\not{d}_1 f = -e_\theta(f)$.

¹⁶Note that ${}^{(1+3)}\underline{\beta}_\theta = -\underline{\beta}$, ${}^{(1+3)}\widehat{\chi} = \frac{1}{2}\underline{\vartheta}$.

Proposition 2.2.1.

$$\begin{aligned}
e_3(\underline{\vartheta}) + \underline{\kappa} \underline{\vartheta} + 2\underline{\omega} \underline{\vartheta} &= -2\underline{\alpha} - 2\underline{\phi}_2^* \underline{\xi} + 2(\underline{\eta} + \underline{\eta} - 2\underline{\zeta}) \underline{\xi}, \\
e_3(\underline{\kappa}) + \frac{1}{2} \underline{\kappa}^2 + 2\underline{\omega} \underline{\kappa} &= 2\underline{\phi}_1 \underline{\xi} + 2(\underline{\eta} + \underline{\eta} - 2\underline{\zeta}) \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2, \\
e_4 \underline{\vartheta} + \frac{1}{2} \underline{\kappa} \underline{\vartheta} - 2\underline{\omega} \underline{\vartheta} &= -2\underline{\phi}_2^* \underline{\eta} - \frac{1}{2} \underline{\kappa} \underline{\vartheta} + 2(\underline{\xi} \underline{\xi} + \underline{\eta}^2), \\
e_4(\underline{\kappa}) + \frac{1}{2} \underline{\kappa} \underline{\kappa} - 2\underline{\omega} \underline{\kappa} &= 2\underline{\phi}_1 \underline{\eta} + 2\rho - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} + 2(\underline{\xi} \underline{\xi} + \underline{\eta} \underline{\eta}), \\
e_3 \underline{\zeta} + \frac{1}{2} \underline{\kappa} (\underline{\zeta} + \underline{\eta}) - 2\underline{\omega} (\underline{\zeta} - \underline{\eta}) &= \underline{\beta} + 2\underline{\phi}_1^* \underline{\omega} + 2\underline{\omega} \underline{\xi} + \frac{1}{2} \underline{\kappa} \underline{\xi} - \frac{1}{2} \underline{\vartheta} (\underline{\zeta} + \underline{\eta}) + \frac{1}{2} \underline{\vartheta} \underline{\xi}, \\
e_4(\underline{\xi}) - 4\underline{\omega} \underline{\xi} - e_3(\underline{\eta}) &= \underline{\beta} + \frac{1}{2} \underline{\kappa} (\underline{\eta} - \underline{\eta}) + \frac{1}{2} \underline{\vartheta} (\underline{\eta} - \underline{\eta}), \\
e_4 \underline{\omega} + e_3 \underline{\omega} &= \rho + 4\underline{\omega} \underline{\omega} + \underline{\xi} \underline{\xi} + \underline{\zeta} (\underline{\eta} - \underline{\eta}) - \underline{\eta} \underline{\eta}.
\end{aligned} \tag{2.2.3}$$

In view of the symmetry $e_3 - e_4$, we also derive,

$$\begin{aligned}
e_4(\underline{\vartheta}) + \underline{\kappa} \underline{\vartheta} + 2\underline{\omega} \underline{\vartheta} &= -2\underline{\alpha} - 2\underline{\phi}_2^* \underline{\xi} + 2(\underline{\eta} + \underline{\eta} + 2\underline{\zeta}) \underline{\xi}, \\
e_4(\underline{\kappa}) + \frac{1}{2} \underline{\kappa}^2 + 2\underline{\omega} \underline{\kappa} &= 2\underline{\phi}_1 \underline{\xi} + 2(\underline{\eta} + \underline{\eta} + 2\underline{\zeta}) \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2, \\
e_3 \underline{\vartheta} + \frac{1}{2} \underline{\kappa} \underline{\vartheta} - 2\underline{\omega} \underline{\vartheta} &= -2\underline{\phi}_2^* \underline{\eta} - \frac{1}{2} \underline{\kappa} \underline{\vartheta} + 2(\underline{\xi} \underline{\xi} + \underline{\eta}^2), \\
e_3(\underline{\kappa}) + \frac{1}{2} \underline{\kappa} \underline{\kappa} - 2\underline{\omega} \underline{\kappa} &= 2\underline{\phi}_1 \underline{\eta} + 2\rho - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} + 2(\underline{\xi} \underline{\xi} + \underline{\eta} \underline{\eta}), \\
-e_4 \underline{\zeta} + \frac{1}{2} \underline{\kappa} (-\underline{\zeta} + \underline{\eta}) + 2\underline{\omega} (\underline{\zeta} + \underline{\eta}) &= \underline{\beta} + 2\underline{\phi}_1^* \underline{\omega} + 2\underline{\omega} \underline{\xi} + \frac{1}{2} \underline{\kappa} \underline{\xi} - \frac{1}{2} \underline{\vartheta} (-\underline{\zeta} + \underline{\eta}) - \frac{1}{2} \underline{\vartheta} \underline{\xi}, \\
e_3(\underline{\xi}) - e_4(\underline{\eta}) &= \underline{\beta} + 4\underline{\omega} \underline{\xi} + \frac{1}{2} \underline{\kappa} (\underline{\eta} - \underline{\eta}) + \frac{1}{2} \underline{\vartheta} (\underline{\eta} - \underline{\eta}), \\
e_4 \underline{\omega} + e_3 \underline{\omega} &= \rho + 4\underline{\omega} \underline{\omega} + \underline{\xi} \underline{\xi} + \underline{\zeta} (\underline{\eta} - \underline{\eta}) - \underline{\eta} \underline{\eta}.
\end{aligned} \tag{2.2.4}$$

We also have the Codazzi equations,

$$\begin{aligned}
\underline{\phi}_2 \underline{\vartheta} &= -2\underline{\beta} - \underline{\phi}_1^* \underline{\kappa} - \underline{\zeta} \underline{\kappa} + \underline{\vartheta} \underline{\zeta}, \\
\underline{\phi}_2 \underline{\vartheta} &= -2\underline{\beta} - \underline{\phi}_1^* \underline{\kappa} + \underline{\zeta} \underline{\kappa} - \underline{\vartheta} \underline{\zeta},
\end{aligned}$$

and the Gauss equation,

$$K = -\rho - \frac{1}{4} \underline{\kappa} \underline{\kappa} + \frac{1}{4} \underline{\vartheta} \underline{\vartheta}.$$

2.2.2 Null Bianchi identities

We now translate the spacetime null Bianchi identities of [17] (see proposition 7.3.2.) in the reduced picture. The spacetime equation (note that $\mathcal{D}_2^* \beta := -\frac{1}{2} {}^{(1+3)}\nabla \otimes \beta$),

$$\nabla_3 \alpha + \frac{1}{2} \text{tr} \underline{\chi} \alpha = -2 \mathcal{D}_2^* \beta + 4 \underline{\omega} \alpha - 3(\widehat{\chi} \rho + {}^* \widehat{\chi} {}^* \rho) + (\zeta + 4\eta) \otimes \beta$$

becomes (note that ${}^* \rho = 0$),

$$e_3 \alpha + \frac{1}{2} \underline{\kappa} \alpha = (e_\theta(\beta) - (e_\theta \Phi) \beta) + 4 \underline{\omega} \alpha - \frac{3}{2} \vartheta \rho + (\zeta + 4\eta) \beta. \quad (2.2.5)$$

The spacetime equation,

$$\nabla_4 \beta + 2 \text{tr} \chi \beta = \text{div} \alpha - 2 \omega \beta + (2\zeta + \underline{\eta}) \cdot \alpha + 3(\xi \rho + {}^* \xi {}^* \rho)$$

becomes,

$$e_4 \beta + 2 \kappa \beta = (e_\theta \alpha + 2(e_\theta \Phi) \alpha) - 2 \omega \beta + (2\zeta + \underline{\eta}) \alpha + 3 \xi \rho. \quad (2.2.6)$$

The spacetime equation,

$$\nabla_3 \beta + \text{tr} \underline{\chi} \beta = \mathcal{D}_1^* (-\rho, {}^* \rho) + 2 \widehat{\chi} \cdot \underline{\beta} + 2 \underline{\omega} \beta + \underline{\xi} \cdot \alpha + 3(\eta \rho + {}^* \eta {}^* \rho)$$

becomes (recall ${}^{(1+3)}\underline{\beta}_\theta = -\underline{\beta}$),

$$e_3 \beta + \underline{\kappa} \beta = e_\theta(\rho) + 2 \underline{\omega} \beta + 3 \eta \rho - \vartheta \underline{\beta} + \underline{\xi} \alpha. \quad (2.2.7)$$

The spacetime equation,

$$e_4 \rho + \frac{3}{2} \text{tr} \chi \rho = \text{div} \beta - \frac{1}{2} \widehat{\chi} \cdot \alpha + \zeta \cdot \beta + 2(\underline{\eta} \cdot \beta - \underline{\xi} \cdot \underline{\beta})$$

becomes,

$$e_4 \rho + \frac{3}{2} \kappa \rho = (e_\theta(\beta) + (e_\theta \Phi) \beta) - \frac{1}{2} \vartheta \alpha + \zeta \beta + 2(\underline{\eta} \beta + \underline{\xi} \underline{\beta}). \quad (2.2.8)$$

Indeed note that,

$${}^{(1+3)}\widehat{\chi} \cdot {}^{(1+3)}\alpha = 2 {}^{(1+3)}\widehat{\chi}_{\theta\theta} {}^{(1+3)}\alpha_{\theta\theta} = \vartheta \alpha.$$

All other equations in the proposition below are derived using the $e_3 - e_4$ symmetry. We summarize the results in the following proposition.

Proposition 2.2.2.

$$\begin{aligned}
e_3\alpha + \frac{1}{2}\underline{\kappa}\alpha &= -\underline{\not{d}}_2^*\beta + 4\underline{\omega}\alpha - \frac{3}{2}\underline{\vartheta}\rho + (\zeta + 4\underline{\eta})\beta, \\
e_4\beta + 2\underline{\kappa}\beta &= \underline{\not{d}}_2\alpha - 2\underline{\omega}\beta + (2\zeta + \underline{\eta})\alpha + 3\underline{\xi}\rho, \\
e_3\beta + \underline{\kappa}\beta &= -\underline{\not{d}}_1^*\rho + 2\underline{\omega}\beta + 3\underline{\eta}\rho - \underline{\vartheta}\beta + \underline{\xi}\alpha, \\
e_4\rho + \frac{3}{2}\underline{\kappa}\rho &= \underline{\not{d}}_1\beta - \frac{1}{2}\underline{\vartheta}\alpha + \zeta\beta + 2(\underline{\eta}\beta + \underline{\xi}\beta), \\
e_3\rho + \frac{3}{2}\underline{\kappa}\rho &= \underline{\not{d}}_1\underline{\beta} - \frac{1}{2}\underline{\vartheta}\underline{\alpha} - \zeta\underline{\beta} + 2(\underline{\eta}\underline{\beta} + \underline{\xi}\beta), \\
e_4\underline{\beta} + \underline{\kappa}\underline{\beta} &= -\underline{\not{d}}_1^*\rho + 2\underline{\omega}\underline{\beta} + 3\underline{\eta}\rho - \underline{\vartheta}\beta + \underline{\xi}\alpha, \\
e_3\underline{\beta} + 2\underline{\kappa}\underline{\beta} &= \underline{\not{d}}_2\underline{\alpha} - 2\underline{\omega}\underline{\beta} + (-2\zeta + \underline{\eta})\underline{\alpha} + 3\underline{\xi}\rho, \\
e_4\underline{\alpha} + \frac{1}{2}\underline{\kappa}\underline{\alpha} &= -\underline{\not{d}}_2^*\underline{\beta} + 4\underline{\omega}\underline{\alpha} - \frac{3}{2}\underline{\vartheta}\rho + (-\zeta + 4\underline{\eta})\underline{\beta}.
\end{aligned} \tag{2.2.9}$$

Mass aspect functions

We define the mass aspect functions,

$$\begin{aligned}
\mu &:= -\underline{\not{d}}_1\zeta - \rho + \frac{1}{4}\underline{\vartheta}\underline{\vartheta}, \\
\underline{\mu} &:= \underline{\not{d}}_1\underline{\zeta} - \rho + \frac{1}{4}\underline{\vartheta}\underline{\vartheta}.
\end{aligned} \tag{2.2.10}$$

One can derive useful propagation equations, in the e_4 direction for μ and in the e_3 direction for $\underline{\mu}$ by using the null structure and null Bianchi equations, see [17] and [37]. In the next section we will do this in the context of null-geodesic foliations.

2.2.3 Hawking mass

Definition 2.2.3. *The Hawking mass $m = m(S)$ of S is defined by the formula,*

$$\frac{2m}{r} = 1 + \frac{1}{16\pi} \int_S \kappa\underline{\kappa}. \tag{2.2.11}$$

Proposition 2.2.4. *The following identities hold true.*

1. The average of ρ is given by the formulas,

$$\bar{\rho} = -\frac{2m}{r^3} + \frac{1}{16\pi r^2} \int_S \vartheta \underline{\vartheta}. \quad (2.2.12)$$

2. The average of the mass aspect function is,

$$\bar{\mu} = \underline{\bar{\mu}} = \frac{2m}{r^3}. \quad (2.2.13)$$

3. The average of κ and $\underline{\kappa}$ are related by,

$$\bar{\kappa} \bar{\underline{\kappa}} = -\frac{4\Upsilon}{r^2} - \bar{\underline{\kappa}} \bar{\kappa} \quad (2.2.14)$$

where $\Upsilon = 1 - \frac{2m}{r}$.

Proof. We have from the Gauss equation

$$K = -\frac{1}{4} \kappa \underline{\kappa} + \frac{1}{4} \vartheta \underline{\vartheta} - \rho.$$

Integrating on S and using the Gauss Bonnet formula, we infer

$$4\pi = -\frac{1}{4} \int_S \kappa \underline{\kappa} + \frac{1}{4} \int_S \vartheta \underline{\vartheta} - \int_S \rho.$$

Together with the definition of the Hawking mass, we infer

$$\begin{aligned} \int_S \rho &= -4\pi \left(1 + \frac{1}{16\pi} \int_S \kappa \underline{\kappa} \right) + \frac{1}{4} \int_S \vartheta \underline{\vartheta} \\ &= -\frac{8\pi m}{r} + \frac{1}{4} \int_S \vartheta \underline{\vartheta} \end{aligned}$$

and hence

$$\bar{\rho} = -\frac{2m}{r^3} + \frac{1}{16\pi r^2} \int_S \vartheta \underline{\vartheta}.$$

which proves our first identity. The second identity follows easily from the definition of $\mu, \underline{\mu}$ and the first formula. Thus, for example,

$$\bar{\mu} = \frac{1}{|S|} \int_S \mu = \frac{1}{|S|} \int_S \left(-\not{d}_1 \zeta - \rho + \frac{1}{4} \vartheta \underline{\vartheta} \right) = -\bar{\rho} + \frac{1}{4|S|} \int_S \vartheta \underline{\vartheta} = \frac{2m}{r^3}.$$

To prove the last identity we remark that, in view of the definition of the Hawking mass,

$$-\Upsilon = \frac{2m}{r} - 1 = \frac{1}{16\pi} \int_S \kappa_{\underline{K}} = \frac{1}{16\pi} \left(|S| \overline{\kappa_{\underline{K}}} + \int_S \check{\kappa}_{\check{\underline{K}}} \right)$$

and hence

$$\begin{aligned} \overline{\kappa_{\underline{K}}} &= -\frac{16\pi\Upsilon}{|S|} - \frac{1}{|S|} \int_S \check{\kappa}_{\check{\underline{K}}} \\ &= -\frac{4\Upsilon}{r^2} - \overline{\check{\kappa}_{\check{\underline{K}}}}. \end{aligned}$$

This concludes the proof of the proposition. \square

2.2.4 Outgoing Geodesic foliations

We restrict our attention to geodesic foliations, i.e. geodesic foliations by \mathbf{Z} invariant optical functions.

Basic definitions

Assume given an outgoing optical function u , i.e. \mathbf{Z} -invariant solution of the equation,

$$\mathbf{g}^{\alpha\beta} \partial_\alpha u \partial_\beta u = g^{ab} \partial_a u \partial_b u = 0$$

and $L = -g^{ab} \partial_b u \partial_a$ its null geodesic generator. We choose e_4 such that,

$$e_4 = \varsigma L, \quad L(\varsigma) = 0. \quad (2.2.15)$$

Remark 2.2.5. *In our definition of a GCM admissible spacetime, see section 3.1, we initialize ς on the spacelike hypersurface Σ_* .*

We then choose s such that

$$e_4(s) = 1. \quad (2.2.16)$$

The functions u, s generate what is called an outgoing geodesic foliation. Let $S_{u,s}$ be the 2-surfaces of intersection between the level surfaces of u and s . We choose e_3 the unique \mathbf{Z} -invariant null vectorfield orthogonal to $S_{u,s}$ and such that $g(e_3, e_4) = -2$. We then let e_θ to be unit tangent to $S_{u,s}$, \mathbf{Z} -invariant and orthogonal to \mathbf{Z} . We also introduce

$$\underline{\Omega} := e_3(s). \quad (2.2.17)$$

Lemma 2.2.6. *We have*

$$\omega = \xi = 0, \quad \underline{\eta} = -\zeta, \quad (2.2.18)$$

$$\begin{aligned} \varsigma &= \frac{2}{e_3(u)}, \\ e_4(\varsigma) &= 0, \\ e_\theta(\log \varsigma) &= \eta - \zeta, \\ e_\theta(\underline{\Omega}) &= -\underline{\xi} - (\eta - \zeta)\underline{\Omega}, \\ e_4(\underline{\Omega}) &= -2\underline{\omega}. \end{aligned} \quad (2.2.19)$$

Proof. Recall that L is geodesic, $e_4 = \varsigma L$ and $L(\varsigma) = 0$. This immediately implies that e_4 is geodesic, and hence we have

$$\omega = \xi = 0.$$

Applying the vectorfield

$$[e_4, e_\theta] = (\underline{\eta} + \zeta)e_4 + \xi e_3 - \chi e_\theta$$

to s , and since $e_4(s) = 1$ and $e_\theta(s) = 0$, we derive,

$$\underline{\eta} + \zeta = 0.$$

Next, note that

$$e_3(u) = g(e_3, -L) = -\varsigma^{-1}g(e_3, e_4) = \frac{2}{\varsigma}$$

and hence

$$\varsigma = \frac{2}{e_3(u)}.$$

Applying the vectorfield

$$[e_3, e_\theta] = \underline{\xi}e_4 + (\eta - \zeta)e_3 - \underline{\chi}e_\theta$$

to u and making use of the relation $e_4(u) = e_\theta(u) = 0$ we deduce,

$$(\eta - \zeta)e_3(u) = e_3(e_\theta u) - e_\theta e_3(u) = -e_\theta e_3(u)$$

which together with the identity $\varsigma = 2/e_3(u)$ yields

$$\eta - \zeta = -e_\theta \log(e_3 u) = -e_\theta \log\left(\frac{2}{\varsigma}\right) = e_\theta(\log \varsigma)$$

and hence

$$e_\theta(\log \varsigma) = \eta - \zeta.$$

Applying the vectorfield

$$[e_3, e_\theta] = \underline{\xi}e_4 + (\eta - \zeta)e_3 - \underline{\chi}e_\theta$$

to s we deduce, since $e_4(s) = 1$, $e_\theta(s) = 0$ and $e_3(s) = \underline{\Omega}$,

$$e_\theta(\underline{\Omega}) = -\underline{\xi} - (\eta - \zeta)\underline{\Omega}.$$

Finally applying

$$[e_4, e_3] = -2\underline{\omega}e_4 - 2(\eta - \underline{\eta})e_\theta + 2\underline{\omega}e_3$$

to s , and using $e_4(s) = 1$ and $e_\theta(s) = 0$, we infer $e_4(e_3(s)) = -2\underline{\omega}$, i.e. $e_4(\underline{\Omega}) = -2\underline{\omega}$ as desired. \square

Remark 2.2.7. *In the particular case when ς is constant we have $\eta = \zeta = -\underline{\eta}$. In Schwarzschild, relative to the standard outgoing geodesic frame, we have*

$$\varsigma = 1, \quad \underline{\Omega} = -\Upsilon = -\left(1 - \frac{2m}{r}\right).$$

Basic equations

Proposition 2.2.8. *Relative to an outgoing geodesic foliation we have*

1. The reduced null structure equations take the form,

$$\begin{aligned}
e_4(\vartheta) + \kappa \vartheta &= -2\alpha, \\
e_4(\kappa) + \frac{1}{2}\kappa^2 &= -\frac{1}{2}\vartheta^2, \\
e_4\zeta + \kappa\zeta &= -\beta - \vartheta\zeta, \\
e_4(\eta - \zeta) + \frac{1}{2}\kappa(\eta - \zeta) &= -\frac{1}{2}\vartheta(\eta - \zeta), \\
e_4\underline{\vartheta} + \frac{1}{2}\kappa\underline{\vartheta} &= 2\phi_2^* \zeta - \frac{1}{2}\underline{\kappa}\vartheta + 2\zeta^2, \\
e_4(\underline{\kappa}) + \frac{1}{2}\kappa\underline{\kappa} &= -2\phi_1\zeta + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\zeta^2, \\
e_4\underline{\omega} &= \rho + \zeta(2\eta + \zeta), \\
e_4(\underline{\xi}) &= -e_3(\zeta) + \underline{\beta} - \frac{1}{2}\underline{\kappa}(\zeta + \eta) - \frac{1}{2}\underline{\vartheta}(\zeta + \eta),
\end{aligned}$$

and

$$\begin{aligned}
e_3(\underline{\vartheta}) + \underline{\kappa}\underline{\vartheta} + 2\underline{\omega}\underline{\vartheta} &= -2\underline{\alpha} - 2\phi_2^* \underline{\xi} + 2(\eta - 3\zeta)\underline{\xi}, \\
e_3(\underline{\kappa}) + \frac{1}{2}\underline{\kappa}^2 + 2\underline{\omega}\underline{\kappa} &= 2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\underline{\vartheta}^2, \\
e_3\zeta + \frac{1}{2}\underline{\kappa}(\zeta + \eta) - 2\underline{\omega}(\zeta - \eta) &= \underline{\beta} + 2\phi_1^* \underline{\omega} + \frac{1}{2}\kappa\underline{\xi} - \frac{1}{2}\underline{\vartheta}(\zeta + \eta) + \frac{1}{2}\vartheta\underline{\xi}, \\
e_3\underline{\vartheta} + \frac{1}{2}\underline{\kappa}\underline{\vartheta} - 2\underline{\omega}\underline{\vartheta} &= -2\phi_2^* \eta - \frac{1}{2}\kappa\underline{\vartheta} + 2\eta^2, \\
e_3(\kappa) + \frac{1}{2}\underline{\kappa}\kappa - 2\underline{\omega}\kappa &= 2\phi_1\eta + 2\rho - \frac{1}{2}\underline{\vartheta}\vartheta + 2\eta^2,
\end{aligned}$$

and

$$\begin{aligned}
\phi_2\underline{\vartheta} &= -2\underline{\beta} - \phi_1^* \underline{\kappa} - \zeta\underline{\kappa} + \underline{\vartheta}\zeta, \\
\phi_2\underline{\vartheta} &= -2\underline{\beta} - \phi_1^* \kappa + \zeta\kappa - \vartheta\zeta, \\
K &= -\rho - \frac{1}{4}\kappa\underline{\kappa} + \frac{1}{4}\vartheta\underline{\vartheta}.
\end{aligned}$$

2. The null Bianchi identities are given in this case by

$$\begin{aligned}
e_3\alpha + \frac{1}{2}\underline{\kappa}\alpha &= -\phi_2^* \beta + 4\underline{\omega}\alpha - \frac{3}{2}\vartheta\rho + (\zeta + 4\eta)\beta, \\
e_4\beta + 2\kappa\beta &= \phi_2\alpha + \zeta\alpha, \\
e_3\beta + \underline{\kappa}\beta &= -\phi_1^* \rho + 2\underline{\omega}\beta + 3\eta\rho - \vartheta\underline{\beta} + \underline{\xi}\alpha, \\
e_4\rho + \frac{3}{2}\kappa\rho &= \phi_1\beta - \frac{1}{2}\underline{\vartheta}\alpha - \zeta\beta,
\end{aligned}$$

$$\begin{aligned}
e_3\rho + \frac{3}{2}\underline{\kappa}\rho &= \underline{\phi}_1\underline{\beta} - \frac{1}{2}\vartheta\underline{\alpha} - \zeta\underline{\beta} + 2(\underline{\eta}\underline{\beta} + \underline{\xi}\underline{\beta}), \\
e_4\underline{\beta} + \underline{\kappa}\underline{\beta} &= -\underline{\phi}_1^*\rho - 3\underline{\zeta}\rho - \underline{\vartheta}\underline{\beta}, \\
e_3\underline{\beta} + 2\underline{\kappa}\underline{\beta} &= \underline{\phi}_2\underline{\alpha} - 2\underline{\omega}\underline{\beta} + (-2\underline{\zeta} + \underline{\eta})\underline{\alpha} + 3\underline{\xi}\rho, \\
e_4\underline{\alpha} + \frac{1}{2}\underline{\kappa}\underline{\alpha} &= -\underline{\phi}_2^*\underline{\beta} - \frac{3}{2}\underline{\vartheta}\rho - 5\underline{\zeta}\underline{\beta}.
\end{aligned}$$

3. The mass aspect function $\mu = -\underline{\phi}_1\underline{\zeta} - \rho + \frac{1}{4}\vartheta\underline{\vartheta}$, defined in (2.2.10) verifies the transport equation,

$$\begin{aligned}
e_4(\mu) + \frac{3}{2}\underline{\kappa}\mu &= \text{Err}[e_4\mu], \\
\text{Err}[e_4\mu] &= \frac{1}{2}\underline{\kappa}\zeta^2 + e_\theta(\underline{\kappa})\zeta + \underline{\phi}_1(\vartheta\underline{\zeta}) - \frac{1}{8}\underline{\kappa}\vartheta^2.
\end{aligned}$$

Proof. Concerning the null structure equations we only need to derive the equation for $\eta - \zeta$. According to Proposition (2.2.1) we have,

$$e_3(\underline{\xi}) - e_4(\underline{\eta}) = \underline{\beta} + 4\underline{\omega}\underline{\xi} + \frac{1}{2}\underline{\kappa}(\underline{\eta} - \underline{\eta}) + \frac{1}{2}\vartheta(\underline{\eta} - \underline{\eta})$$

which becomes

$$e_4\underline{\eta} = -\underline{\beta} - \frac{1}{2}\underline{\kappa}(\underline{\eta} - \underline{\eta}) - \frac{1}{2}\vartheta(\underline{\eta} - \underline{\eta})$$

and,

$$-e_4\underline{\zeta} + \frac{1}{2}\underline{\kappa}(-\underline{\zeta} + \underline{\eta}) + 2\underline{\omega}(\underline{\zeta} + \underline{\eta}) = \underline{\beta} + 2\underline{\phi}_1^*\underline{\omega} + 2\underline{\omega}\underline{\xi} + \frac{1}{2}\underline{\kappa}\underline{\xi} - \frac{1}{2}\vartheta(-\underline{\zeta} + \underline{\eta}) - \frac{1}{2}\underline{\vartheta}\underline{\xi}$$

which becomes,

$$e_4\underline{\zeta} = -\underline{\kappa}\underline{\zeta} - \underline{\beta} - \vartheta\underline{\zeta}.$$

Hence,

$$\begin{aligned}
e_4(\underline{\zeta} - \underline{\eta}) &= -\underline{\kappa}\underline{\zeta} - \vartheta\underline{\zeta} + \frac{1}{2}\underline{\kappa}(\underline{\eta} - \underline{\eta}) + \frac{1}{2}\vartheta(\underline{\eta} - \underline{\eta}) \\
&= \underline{\kappa} \left(-\underline{\zeta} + \frac{1}{2}(\underline{\eta} - \underline{\eta}) \right) + \vartheta \left(-\underline{\zeta} + \frac{1}{2}(\underline{\eta} - \underline{\eta}) \right).
\end{aligned}$$

Since $\underline{\zeta} = -\underline{\eta}$ we deduce $-\underline{\zeta} + \frac{1}{2}(\underline{\eta} - \underline{\eta}) = \frac{1}{2}(-\underline{\zeta} + \underline{\eta})$ and thus,

$$e_4(\underline{\zeta} - \underline{\eta}) = -\underline{\kappa}(\underline{\zeta} - \underline{\eta}) - \vartheta(\underline{\zeta} - \underline{\eta})$$

as desired.

The Bianchi equations follow immediately from the general equations derived in the previous section. It only remains to check the equation verified by the mass aspect function μ . We have

$$\begin{aligned}
e_4(\mu) &= -[e_4, \not{d}_1]\zeta - \not{d}_1 e_4(\zeta) - e_4(\rho) + \frac{1}{4}e_4(\vartheta\underline{\vartheta}) \\
&= \frac{1}{2}\kappa \not{d}_1 \zeta - \frac{1}{2}\vartheta \not{d}_2^* \zeta + e_4(\Phi)\zeta^2 - \beta\zeta - \not{d}_1(-\kappa\zeta - \beta - \vartheta\zeta) \\
&\quad - \left(-\frac{3}{2}\kappa\rho + \not{d}_1\beta - \frac{1}{2}\vartheta\underline{\alpha} - \zeta\beta \right) \\
&\quad + \frac{1}{4}\vartheta \left(-\frac{1}{2}\kappa\underline{\vartheta} + 2\not{d}_2^* \zeta - \frac{1}{2}\underline{\kappa}\vartheta + 2\zeta^2 \right) + \frac{1}{4}\underline{\vartheta}(-\kappa\vartheta - 2\alpha) \\
&= \frac{3}{2}\kappa \left(\not{d}_1 \zeta + \rho - \frac{1}{4}\vartheta\underline{\vartheta} \right) - \frac{1}{2}\vartheta \not{d}_2^* \zeta + \frac{1}{2}(\kappa - \vartheta)\zeta^2 + e_\theta(\kappa)\zeta + \not{d}_1(\vartheta\zeta) \\
&\quad + \frac{1}{4}\vartheta \left(2\not{d}_2^* \zeta - \frac{1}{2}\underline{\kappa}\vartheta + 2\zeta^2 \right)
\end{aligned}$$

and hence

$$e_4(\mu) + \frac{3}{2}\kappa\mu = \frac{1}{2}\kappa\zeta^2 + e_\theta(\kappa)\zeta + \not{d}_1(\vartheta\zeta) - \frac{1}{8}\underline{\kappa}\vartheta^2$$

as desired. This concludes the proof of the proposition. \square

Transport equations for S -averages

Proposition 2.2.9. *For any scalar function f , we have,*

$$\begin{aligned}
e_4 \left(\int_S f \right) &= \int_S (e_4(f) + \kappa f), \\
e_3 \left(\int_S f \right) &= \int_S (e_3(f) + \underline{\kappa}f) + \text{Err} \left[e_3 \left(\int_S f \right) \right],
\end{aligned} \tag{2.2.20}$$

where the error term is given by the formula

$$\begin{aligned}
\text{Err} \left[e_3 \left(\int_S f \right) \right] : &= -\varsigma^{-1}\check{\zeta} \int_S (e_3(f) + \underline{\kappa}f) + \varsigma^{-1} \int_S \check{\zeta}(e_3(f) + \underline{\kappa}f) \\
&\quad + (\check{\underline{\Omega}} + \varsigma^{-1}\overline{\underline{\Omega}}\check{\zeta}) \int_S (e_4 f + \kappa f) - \varsigma^{-1}\overline{\underline{\Omega}} \int_S \check{\zeta}(e_4 f + \kappa f) \\
&\quad - \varsigma^{-1} \int_S \check{\underline{\Omega}}\zeta(e_4 f + \kappa f).
\end{aligned}$$

In particular, we have

$$e_4(r) = \frac{r}{2}\bar{\kappa}, \quad e_3(r) = \frac{r}{2}(\bar{\kappa} + \underline{A}) \quad (2.2.21)$$

where

$$\underline{A} := -\varsigma^{-1}\bar{\kappa}\check{\zeta} + \bar{\kappa}(\check{\Omega} + \varsigma^{-1}\bar{\Omega}\check{\zeta}) + \varsigma^{-1}\check{\zeta}\bar{\kappa} - \varsigma^{-1}\bar{\Omega}\check{\zeta}\bar{\kappa} - \varsigma^{-1}\check{\Omega}\check{\zeta}\bar{\kappa}. \quad (2.2.22)$$

Proof. See section A.1. □

Corollary 2.2.10. *For a reduced scalar f , we have*

$$e_4\left(\int_S f e^\Phi\right) = \int_S \left(e_4(f) + \left(\frac{3}{2}\kappa - \frac{1}{2}\vartheta\right) f\right) e^\Phi$$

and

$$e_3\left(\int_S f e^\Phi\right) = \int_S \left(e_3(f) + \left(\frac{3}{2}\kappa - \frac{1}{2}\vartheta\right) f\right) e^\Phi + \text{Err}\left[e_3\left(\int_S f e^\Phi\right)\right].$$

Proof. In view of Proposition 2.2.9, we have

$$\begin{aligned} e_4\left(\int_S f e^\Phi\right) &= \int_S \left(e_4(f e^\Phi) + \kappa f e^\Phi\right) \\ &= \int_S \left(e_4(f) + (\kappa + e_4\Phi) f\right) e^\Phi \\ &= \int_S \left(e_4(f) + \left(\frac{3}{2}\kappa - \frac{1}{2}\vartheta\right) f\right) e^\Phi \end{aligned}$$

as desired.

Also, using again Proposition 2.2.9, we have

$$\begin{aligned} e_3\left(\int_S f e^\Phi\right) &= \int_S \left(e_3(f e^\Phi) + \kappa f e^\Phi\right) + \text{Err}\left[e_3\left(\int_S f e^\Phi\right)\right] \\ &= \int_S \left(e_3(f) + (\kappa + e_3\Phi) f\right) e^\Phi + \text{Err}\left[e_3\left(\int_S f e^\Phi\right)\right] \\ &= \int_S \left(e_3(f) + \left(\frac{3}{2}\kappa - \frac{1}{2}\vartheta\right) f\right) e^\Phi + \text{Err}\left[e_3\left(\int_S f e^\Phi\right)\right] \end{aligned}$$

as desired. □

Corollary 2.2.11. *Given a scalar function f we have,*

$$\begin{aligned} e_4(\bar{f}) &= \overline{e_4(f)} + \bar{\kappa} \bar{f}, \\ e_4(\check{f}) &= e_4(f) - \overline{e_4(f)} - \bar{\kappa} \bar{f}, \end{aligned} \quad (2.2.23)$$

and

$$\begin{aligned} e_3(\bar{f}) &= \overline{e_3(f)} + \text{Err}[e_3 \bar{f}], \\ e_3(\check{f}) &= e_3(f) - \overline{e_3(f)} - \text{Err}[e_3(\bar{f})], \end{aligned} \quad (2.2.24)$$

where,

$$\begin{aligned} \text{Err}[e_3(\bar{f})] &= -\varsigma^{-1} \check{\zeta} (\overline{e_3 f + \kappa f} - \bar{\kappa} \bar{f}) + \varsigma^{-1} (\overline{\check{\zeta}(e_3 f + \kappa f)} - \bar{\check{\zeta}} \bar{\kappa} \bar{f}) \\ &\quad + (\check{\Omega} + \varsigma^{-1} \check{\Omega} \check{\zeta}) (\overline{e_4 f + \kappa f} - \bar{\kappa} \bar{f}) - \varsigma^{-1} \check{\Omega} (\overline{\check{\zeta}(e_4 f + \kappa f)} - \bar{\check{\zeta}} \bar{\kappa} \bar{f}) \\ &\quad - \varsigma^{-1} (\overline{\check{\Omega} \varsigma (e_4 f + \kappa f)} - \bar{\check{\Omega}} \bar{\varsigma} \bar{\kappa} \bar{f}) + \bar{\kappa} \bar{f}. \end{aligned} \quad (2.2.25)$$

Proof. We have, recalling Lemma 2.1.42 and $|S| = 4\pi r^2$,

$$\begin{aligned} e_4(\bar{f}) &= e_4 \left(\frac{\int_S f}{|S|} \right) = \frac{1}{|S|} \int_S (e_4(f) + \kappa f) - \frac{e_4(|S|)}{|S|} \bar{f} = \overline{e_4(f) + \kappa f} - 2 \frac{e_4 r}{r} \bar{f} \\ &= \overline{e_4(f)} + \bar{\kappa} \bar{f} - \bar{\kappa} \bar{f} = \overline{e_4(f)} + \bar{\kappa} \bar{f}. \end{aligned}$$

This also yields

$$e_4(\check{f}) = e_4(f) - e_4(\bar{f}) = e_4(f) - \overline{e_4(f)} - \bar{\kappa} \bar{f}$$

as desired.

Similarly,

$$\begin{aligned} e_3(\bar{f}) &= e_3 \left(\frac{\int_S f}{|S|} \right) = \frac{1}{|S|} e_3 \left(\int_S f \right) - \frac{2e_3(r)}{r} \bar{f} \\ &= \frac{1}{|S|} \int_S (e_3 f + \kappa f) + \frac{1}{|S|} \text{Err} \left[e_3 \left(\int_S f \right) \right] - (\bar{\kappa} + \underline{A}) \bar{f} \\ &= \overline{e_3(f)} + \bar{\kappa} \bar{f} - \bar{\kappa} \bar{f} + \frac{1}{|S|} \text{Err} \left[e_3 \left(\int_S f \right) \right] - \underline{A} \bar{f} \\ &= \overline{e_3(f)} + \bar{\kappa} \bar{f} + \frac{1}{|S|} \text{Err} \left[e_3 \left(\int_S f \right) \right] - \underline{A} \bar{f}. \end{aligned}$$

We deduce,

$$e_3(\bar{f}) = \overline{e_3(f)} + \text{Err}[e_3(\bar{f})]$$

where, recalling the definitions of the error terms $\text{Err}[e_3(\int_S f)]$ and \underline{A} ,

$$\begin{aligned} \text{Err}[e_3(\bar{f})] &= \overline{\check{\kappa}f} + \frac{1}{|S|} \text{Err} \left[e_3 \left(\int_S f \right) \right] - \underline{A} \bar{f} \\ &= \overline{\check{\kappa}f} - \varsigma^{-1} \check{\zeta} \overline{e_3 f + \underline{\kappa}f} + \varsigma^{-1} \check{\zeta} \overline{e_3 f + \underline{\kappa}f} + (\check{\Omega} + \varsigma^{-1} \check{\Omega} \check{\zeta}) \overline{e_4 f + \kappa f} \\ &\quad - \varsigma^{-1} \check{\Omega} \check{\zeta} \overline{e_4 f + \kappa f} - \varsigma^{-1} \check{\Omega} \check{\zeta} \overline{e_4 f + \kappa f} \\ &\quad - \bar{f} \left(-\varsigma^{-1} \check{\kappa} \check{\zeta} + \varsigma^{-1} \check{\zeta} \check{\kappa} + \check{\kappa} (\check{\Omega} + \varsigma^{-1} \check{\Omega} \check{\zeta}) - \varsigma^{-1} \check{\Omega} \check{\zeta} \check{\kappa} - \varsigma^{-1} \check{\Omega} \check{\zeta} \kappa \right), \end{aligned}$$

i.e.,

$$\begin{aligned} \text{Err}[e_3(\bar{f})] &= \overline{\check{\kappa}f} - \varsigma^{-1} \check{\zeta} (\overline{e_3 f + \underline{\kappa}f} - \check{\kappa} \bar{f}) + \varsigma^{-1} (\check{\zeta} \overline{e_3 f + \underline{\kappa}f} - \check{\zeta} \check{\kappa} \bar{f}) \\ &\quad + (\check{\Omega} + \varsigma^{-1} \check{\Omega} \check{\zeta}) (\overline{e_4 f + \kappa f} - \check{\kappa} \bar{f}) - \varsigma^{-1} \check{\Omega} (\check{\zeta} \overline{e_4 f + \kappa f} - \check{\zeta} \check{\kappa} \bar{f}) \\ &\quad - \varsigma^{-1} (\check{\Omega} \check{\zeta} \overline{e_4 f + \kappa f} - \check{\Omega} \check{\zeta} \check{\kappa} \bar{f}) \end{aligned}$$

as stated. Finally

$$e_3(\check{f}) = e_3 f - e_3(\bar{f}) = e_3 f - \overline{e_3(f)} - \text{Err}[e_3 \bar{f}]$$

which ends the proof of the corollary. \square

The following is also an immediate application of Proposition 2.2.9.

Corollary 2.2.12. *If f verifies the scalar equation*

$$e_4(f) + \frac{p}{2} \check{\kappa} f = F,$$

then,

$$e_4(r^p f) = r^p F.$$

Commutation identities revisited

We revisit the general commutation identities of Lemma 2.1.51 in an outgoing geodesic foliation.

Lemma 2.2.13. *The following commutation formulae holds true,*

1. *If $f \in \mathfrak{s}_k$,*

$$\begin{aligned} [r \not{d}_k, e_4]f &= r \left[\text{Com}_k(f) + \frac{1}{2} \check{\kappa} \not{d}_k f \right], \\ [r \not{d}_k, e_3]f &= r \left[\underline{\text{Com}}_k(f) + \frac{1}{2} (-\underline{A} + \check{\kappa}) \not{d}_k f \right]. \end{aligned} \quad (2.2.26)$$

2. *If $f \in \mathfrak{s}_{k-1}$*

$$\begin{aligned} [r \not{d}_k^*, e_4]f &= r \left[\text{Com}_k^*(f) + \frac{1}{2} \check{\kappa} \not{d}_k^* f \right], \\ [r \not{d}_k^*, e_3]f &= r \left[\underline{\text{Com}}_k^*(f) + \frac{1}{2} (-\underline{A} + \check{\kappa}) \not{d}_k^* f \right]. \end{aligned} \quad (2.2.27)$$

Also, we have

$$\begin{aligned} \underline{\text{Com}}_k(f) &= -\frac{1}{2} \vartheta \not{d}_{k+1}^* f + (\zeta - \eta) e_3 f - k \eta e_3 \Phi f - \underline{\xi} (e_4 f + k e_4(\Phi) f) - k \underline{\beta} f, \\ \text{Com}_k(f) &= -\frac{1}{2} \vartheta \not{d}_{k+1}^* f + k \zeta e_4 \Phi f - k \beta f, \\ \underline{\text{Com}}_k^*(f) &= -\frac{1}{2} \vartheta \not{d}_{k-1} f - (\zeta - \eta) e_3 f - (k-1) \eta e_3 \Phi f + \underline{\xi} (e_4 f - (k-1) e_4(\Phi) f) \\ &\quad - (k-1) \underline{\beta} f, \\ \text{Com}_k^*(f) &= -\frac{1}{2} \vartheta \not{d}_{k-1} f + (k-1) \zeta e_4 \Phi f - (k-1) \beta f. \end{aligned}$$

Proof. We make use of the commutation Lemma 2.1.51 and the definition of \underline{A} , see Proposition 2.2.9, to write, for $f \in \mathfrak{s}_k$,

$$\begin{aligned} [r \not{d}_k, e_4]f &= r[\not{d}_k, e_4]f - e_4(r) \not{d}_k f \\ &= \frac{1}{2} r \check{\kappa} \not{d}_k f + r \text{Com}_k(f) - \frac{r}{2} \overline{\kappa} \not{d}_k f \\ &= r \left[\text{Com}_k(f) + \frac{1}{2} \check{\kappa} \not{d}_k f \right] \\ [r \not{d}_k, e_3]f &= r[\not{d}_k, e_3]f - e_3(r) \not{d}_k f \\ &= \frac{1}{2} r \underline{\kappa} \not{d}_k f + r \underline{\text{Com}}_k(f) - \frac{r}{2} (\underline{A} + \overline{\kappa}) \not{d}_k f \\ &= r \left[\underline{\text{Com}}_k(f) + \frac{1}{2} (-\underline{A} + \check{\kappa}) \not{d}_k f \right]. \end{aligned}$$

The remaining formulae are proved in the same manner. Also, the form of $\underline{Com}_k(f)$, $Com_k(f)$, $\underline{Com}_k^*(f)$ and $Com_k^*(f)$ follows from Lemma 2.1.51 and the fact that we have $\xi = \underline{\eta} + \zeta = 0$ in an outgoing geodesic foliation. \square

We also record here for future use the following lemma.

Lemma 2.2.14. *Let $\mathbf{T} = \frac{1}{2}(e_3 + \Upsilon e_4)$, with $\Upsilon = 1 - \frac{2m}{r}$. We have,*

$$\begin{aligned} [\mathbf{T}, e_4] &= \left(\left(\underline{\omega} - \frac{m}{r^2} \right) - \frac{m}{2r} \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{e_4(m)}{r} \right) e_4 + (\eta + \zeta) e_\theta, \\ [\mathbf{T}, e_3] &= \left(-\Upsilon \left(\underline{\omega} - \frac{m}{r^2} \right) - \frac{m}{2r} \left(\bar{\kappa} + \frac{2\Upsilon}{r} \right) - \frac{m}{2r} \underline{A} + \frac{e_3(m)}{r} \right) e_4 - (\eta + \zeta) \Upsilon e_\theta. \end{aligned} \quad (2.2.28)$$

Proof. Recall that $[e_3, e_4] = 2\underline{\omega}e_4 + 2(\eta + \zeta)e_\theta$. Thus,

$$\begin{aligned} [\mathbf{T}, e_4] &= \frac{1}{2}[e_3 + \Upsilon e_4, e_4] = \frac{1}{2} \left(2\underline{\omega}e_4 + 2(\eta + \zeta)e_\theta - e_4 \left(1 - \frac{2m}{r} \right) e_4 \right) \\ &= \left(\underline{\omega} - \frac{m}{r^2} e_4(r) + \frac{e_4(m)}{r} \right) e_4 + (\eta + \zeta) e_\theta \\ &= \left(\left(\underline{\omega} - \frac{m}{r^2} \right) - \frac{m}{r^2} (e_4(r) - 1) + \frac{e_4(m)}{r} \right) e_4 + (\eta + \zeta) e_\theta \\ &= \left(\left(\underline{\omega} - \frac{m}{r^2} \right) - \frac{m}{r^2} \left(\frac{r}{2} \bar{\kappa} - 1 \right) + \frac{e_4(m)}{r} \right) e_4 + (\eta + \zeta) e_\theta \\ &= \left(\left(\underline{\omega} - \frac{m}{r^2} \right) - \frac{m}{2r} \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{e_4(m)}{r} \right) e_4 + (\eta + \zeta) e_\theta \end{aligned}$$

and,

$$\begin{aligned} [\mathbf{T}, e_3] &= \frac{1}{2}[e_3 + \Upsilon e_4, e_3] = \frac{1}{2} \left(\Upsilon (-2\underline{\omega}e_4 - 2(\eta + \zeta)e_\theta) - e_3 \left(1 - \frac{2m}{r} \right) e_4 \right) \\ &= \left(-\Upsilon \underline{\omega} - \frac{m}{r^2} e_3(r) + \frac{e_3(m)}{r} \right) e_4 - \Upsilon (\eta + \zeta) e_\theta \\ &= \left(-\Upsilon \left(\underline{\omega} - \frac{m}{r^2} \right) - \Upsilon \frac{m}{r^2} - \frac{m}{r^2} \frac{r}{2} (\bar{\kappa} + \underline{A}) + \frac{e_3(m)}{r} \right) e_4 - \Upsilon (\eta + \zeta) e_\theta \\ &= \left(-\Upsilon \left(\underline{\omega} - \frac{m}{r^2} \right) - \frac{m}{2r} \left(\bar{\kappa} + \frac{2\Upsilon}{r} \right) - \frac{m}{2r} \underline{A} + \frac{e_3(m)}{r} \right) e_4 - \Upsilon (\eta + \zeta) e_\theta \end{aligned}$$

which concludes the proof of the lemma. \square

Remark 2.2.15. When applying the formulas of Lemma 2.2.14 to a k reduced scalar $f \in \mathfrak{s}_k$, the term $(\eta + \zeta)e_\theta(f)$ should correspond to a reduced scalar. In fact, recalling Remark 2.1.23, we can write,

$$\zeta e_\theta(f) = \frac{1}{2}\zeta(\not\phi_k f - \not\phi_{k+1}^* f)$$

which can indeed be shown to be a k -reduced scalar in \mathfrak{s}_k .

Derivatives of the Hawking mass

Proposition 2.2.16 (Derivatives of the Hawking mass). *We have the following identities for the Hawking mass,*

$$e_4(m) = \frac{r}{32\pi} \int_S Err_1, \quad (2.2.29)$$

and

$$\begin{aligned} e_3(m) &= (1 - \varsigma^{-1}\zeta) \frac{r}{32\pi} \int_S Err_1 + (\check{\Omega} + \varsigma^{-1}\bar{\Omega}\zeta) \frac{r}{32\pi} \int_S Err_1 \\ &\quad + \varsigma^{-1} \frac{r}{32\pi} \int_S \zeta (2\bar{\rho}\check{\kappa} + 2\check{\rho}\bar{\kappa} + 2\kappa \not\phi_1 \eta + 2\kappa \not\phi_1 \xi + Err_2) \\ &\quad - \varsigma^{-1} \frac{r}{32\pi} \int_S (\bar{\Omega}\zeta + \check{\Omega}\varsigma) (2\bar{\rho}\check{\kappa} + 2\check{\rho}\bar{\kappa} - 2\kappa \not\phi_1 \zeta + Err_2) \\ &\quad - \frac{m}{r} \varsigma^{-1} \left[-\bar{\zeta}\check{\kappa} + \bar{\Omega}\check{\zeta}\bar{\kappa} + \bar{\check{\Omega}}\varsigma\kappa \right], \end{aligned} \quad (2.2.30)$$

where

$$\begin{aligned} Err_1 &:= 2\check{\kappa}\check{\rho} + 2e_\theta(\kappa)\zeta - \frac{1}{2}\kappa\vartheta^2 - \frac{1}{2}\check{\kappa}\vartheta\check{\vartheta} + 2\kappa\zeta^2, \\ \underline{Err}_1 &:= 2\check{\rho}\check{\kappa} - 2e_\theta(\kappa)\eta - 2e_\theta(\kappa)\xi - \frac{1}{2}\check{\kappa}\vartheta\check{\vartheta} + 2\kappa\eta^2 + 2\kappa(\eta - 3\zeta)\xi - \frac{1}{2}\kappa\vartheta^2, \\ Err_2 &:= 2\check{\rho}\check{\kappa} - \frac{1}{2}\kappa\vartheta^2 - \frac{1}{2}\kappa\vartheta\check{\vartheta} + 2\kappa\zeta^2, \\ \underline{Err}_2 &:= 2\check{\rho}\check{\kappa} + \kappa \left(2\eta^2 - \frac{1}{2}\vartheta\check{\vartheta} \right) + 2\kappa(\eta - 3\zeta)\xi - \frac{1}{2}\kappa\vartheta^2. \end{aligned}$$

Proof. The proof relies on the definition of the Hawking mass m given by the formula $\frac{2m}{r} = 1 + \frac{1}{16\pi} \int_S \kappa\check{\kappa}$, Proposition 2.2.9, and the the null structure equations for $e_4(\kappa)$, $e_4(\underline{\kappa})$, $e_3(\kappa)$ and $e_3(\underline{\kappa})$ provided by Proposition 2.2.8. We refer to section A.2 for the details. \square

Transport equations for main averaged quantities

Lemma 2.2.17. *The following equations hold true*

$$\begin{aligned}
e_4 \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{1}{2} \bar{\kappa} \left(\bar{\kappa} - \frac{2}{r} \right) &= -\frac{1}{4} \vartheta^2 + \frac{1}{2} \bar{\kappa}^2, \\
e_4 \left(\bar{\omega} - \frac{m}{r^2} \right) &= \bar{\rho} + \frac{2m}{r^3} + \frac{m}{r^2} \left(\bar{\kappa} - \frac{2}{r} \right) - \frac{e_4(m)}{r^2} + 3\zeta(2\eta + \zeta) + \bar{\kappa} \bar{\omega},
\end{aligned} \tag{2.2.31}$$

and

$$\begin{aligned}
&e_3 \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{1}{2} \bar{\kappa} \left(\bar{\kappa} - \frac{2}{r} \right) \\
&= 2\bar{\omega} \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{4}{r} \left(\bar{\omega} - \frac{m}{r^2} \right) + 2 \left(\bar{\rho} + \frac{2m}{r^3} \right) - \varsigma^{-1} \left(-\frac{1}{2} \bar{\kappa} \bar{\kappa} + 2\bar{\omega} \bar{\kappa} + 2\bar{\rho} \right) \zeta \\
&\quad - \frac{1}{2} \bar{\kappa}^2 (\check{\Omega} + \varsigma^{-1} \bar{\Omega} \zeta) - \frac{1}{r} \varsigma^{-1} \bar{\kappa} \check{\zeta} + \frac{1}{r} \bar{\kappa} (\check{\Omega} + \varsigma^{-1} \bar{\Omega} \zeta) + \text{Err} \left[e_3 \left(\bar{\kappa} - \frac{2}{r} \right) \right],
\end{aligned} \tag{2.2.32}$$

where,

$$\begin{aligned}
\text{Err} \left[e_3 \left(\bar{\kappa} - \frac{2}{r} \right) \right] &:= 2\bar{\eta}^2 + 2\bar{\omega} \bar{\kappa} - \frac{1}{2} \bar{\kappa} \bar{\kappa} - \frac{1}{2} \vartheta \vartheta + \frac{1}{r} \varsigma^{-1} \bar{\zeta} \bar{\kappa} - \frac{1}{r} \varsigma^{-1} \bar{\Omega} \bar{\zeta} \bar{\kappa} - \frac{1}{r} \varsigma^{-1} \bar{\Omega} \varsigma \bar{\kappa} \\
&\quad - \varsigma^{-1} \zeta \left(\frac{1}{2} \bar{\kappa} \bar{\kappa} + 2\bar{\omega} \bar{\kappa} - \frac{1}{2} \vartheta \vartheta + 2\eta^2 \right) \\
&\quad + \varsigma^{-1} \left(\zeta \left(\frac{1}{2} \bar{\kappa} \bar{\kappa} + 2\bar{\omega} \bar{\kappa} + 2\bar{\rho} + 2\bar{\rho}_1 \eta - \frac{1}{2} \vartheta \vartheta + 2\eta^2 \right) - \bar{\zeta} \bar{\kappa} \bar{\kappa} \right) \\
&\quad + (\check{\Omega} + \varsigma^{-1} \bar{\Omega} \zeta) \left(\frac{1}{2} \bar{\kappa}^2 - \frac{1}{4} \vartheta^2 \right) - \varsigma^{-1} \bar{\Omega} \left(\zeta \left(\frac{1}{2} \bar{\kappa}^2 - \frac{1}{4} \vartheta^2 \right) - \bar{\zeta} \bar{\kappa} \bar{\kappa} \right) \\
&\quad - \varsigma^{-1} \left(\bar{\Omega} \varsigma \left(\frac{1}{2} \bar{\kappa}^2 - \frac{1}{4} \vartheta^2 \right) - \bar{\Omega} \varsigma \bar{\kappa} \bar{\kappa} \right) + \bar{\kappa} \bar{\kappa}.
\end{aligned} \tag{2.2.33}$$

Proof. The proof relies on Corollary 2.2.11 and the null structure equations for $e_4(\kappa)$ and $e_3(\kappa)$ provided by Proposition 2.2.8. We refer to section A.3 for the details. \square

Transport equations for main checked quantities

Proposition 2.2.18 (Transport equations for checked quantities). *We have the following transport equations in the e_4 direction,*

$$\begin{aligned}
e_4\check{\kappa} + \bar{\kappa}\check{\kappa} &= Err[e_4\check{\kappa}], \\
Err[e_4\check{\kappa}] &:= -\frac{1}{2}\check{\kappa}^2 - \frac{1}{2}\bar{\kappa}^2 - \frac{1}{2}(\vartheta^2 - \bar{\vartheta}^2), \\
e_4\check{\kappa} + \frac{1}{2}\bar{\kappa}\check{\kappa} + \frac{1}{2}\check{\kappa}\bar{\kappa} &= -2\phi_1\zeta + 2\check{\rho} + Err[e_4\check{\kappa}], \\
Err[e_4\check{\kappa}] &:= -\frac{1}{2}\check{\kappa}\check{\kappa} - \frac{1}{2}\bar{\kappa}\bar{\kappa} + \left(-\frac{1}{2}\vartheta\vartheta + 2\zeta^2\right) - \overline{\left(-\frac{1}{2}\vartheta\vartheta + 2\zeta^2\right)}, \\
e_4\check{\omega} &= \check{\rho} + Err[e_4\check{\omega}], \\
Err[e_4\check{\omega}] &:= -\bar{\kappa}\bar{\omega} + (\zeta(2\eta + \zeta) - \overline{\zeta(2\eta + \zeta)}),
\end{aligned} \tag{2.2.34}$$

$$\begin{aligned}
e_4\check{\rho} + \frac{3}{2}\bar{\kappa}\check{\rho} + \frac{3}{2}\check{\rho}\bar{\kappa} &= \phi_1\beta + Err[e_4\check{\rho}], \\
Err[e_4\check{\rho}] &:= -\frac{3}{2}\check{\kappa}\check{\rho} + \frac{1}{2}\bar{\kappa}\check{\rho} - \left(\frac{1}{2}\vartheta\alpha + \zeta\beta\right) + \overline{\left(\frac{1}{2}\vartheta\alpha + \zeta\beta\right)}, \\
e_4\check{\mu} + \frac{3}{2}\bar{\kappa}\check{\mu} + \frac{3}{2}\check{\mu}\bar{\kappa} &= Err[e_4\check{\mu}], \\
Err[e_4\check{\mu}] &:= -\frac{3}{2}\check{\kappa}\check{\mu} + \frac{1}{2}\bar{\kappa}\check{\mu} + Err[e_4\mu] - \overline{Err[e_4\mu]}, \\
e_4(\check{\Omega}) &= -2\check{\omega} + \bar{\kappa}\bar{\Omega}.
\end{aligned} \tag{2.2.35}$$

Also in the e_3 direction,

$$\begin{aligned}
e_3(\check{\kappa}) &= 2\phi_1\eta + 2\check{\rho} - \frac{1}{2}(\underline{\kappa}\check{\kappa} + \kappa\check{\kappa}) + 2(\underline{\omega}\check{\kappa} + \kappa\check{\omega}) \\
&\quad + \varsigma^{-1} \left(-\frac{1}{2}\bar{\kappa}\bar{\kappa} + 2\bar{\omega}\bar{\kappa} + 2\bar{\rho}\right) \zeta + \frac{1}{2}\bar{\kappa}^2 (\check{\Omega} + \varsigma^{-1}\bar{\Omega}\zeta) + Err[e_3\check{\kappa}], \\
e_3(\check{\kappa}) + \bar{\kappa}\check{\kappa} &= 2\phi_1\underline{\xi} - 2(\check{\omega}\bar{\kappa} + \underline{\omega}\check{\kappa}) + \varsigma^{-1}\zeta \left(-\frac{1}{2}\bar{\kappa}^2 - 2\bar{\omega}\bar{\kappa}\right) \\
&\quad - (\check{\Omega} + \varsigma^{-1}\bar{\Omega}\zeta) \left(-\frac{1}{2}\bar{\kappa}\bar{\kappa} + 2\bar{\rho}\right) + Err[e_3(\check{\kappa})], \\
e_3\check{\rho} + \frac{3}{2}\bar{\kappa}\check{\rho} &= -\frac{3}{2}\check{\rho}\bar{\kappa} + \phi_1\underline{\beta} - \frac{3}{2}\bar{\kappa}\bar{\rho}\varsigma^{-1}\zeta + \frac{3}{2}\bar{\kappa}\bar{\rho} (\check{\Omega} + \varsigma^{-1}\bar{\Omega}\zeta) + Err[e_3\check{\rho}],
\end{aligned} \tag{2.2.36}$$

with error terms given by,

$$\begin{aligned}
Err[e_3\check{\kappa}] &:= 2 \left(\eta^2 - \overline{\eta^2} \right) - \frac{1}{2} \overline{\check{\kappa}\check{\kappa}} + 2\overline{\check{\omega}\check{\kappa}} - \frac{1}{2} (\vartheta\vartheta - \overline{\vartheta\vartheta}) \\
&+ \varsigma^{-1} \check{\zeta} \left(\overline{\frac{1}{2} \check{\kappa}\check{\kappa} + 2\check{\omega}\check{\kappa} - \frac{1}{2} \vartheta\vartheta + 2\eta^2} \right) \\
&- \varsigma^{-1} \left(\check{\zeta} \left(\overline{\frac{1}{2} \kappa\kappa + 2\omega\kappa + 2\check{\rho} + 2\phi_1\eta - \frac{1}{2} \vartheta\vartheta + 2\eta^2} \right) - \overline{\check{\zeta}\check{\kappa}\check{\kappa}} \right) \\
&- (\check{\Omega} + \varsigma^{-1} \overline{\check{\Omega}\check{\zeta}}) \left(\overline{\frac{1}{2} \check{\kappa}^2 - \frac{1}{4} \vartheta^2} \right) + \varsigma^{-1} \overline{\check{\Omega}} \left(\check{\zeta} \left(\overline{\frac{1}{2} \kappa^2 - \frac{1}{4} \vartheta^2} \right) - \overline{\check{\zeta}\check{\kappa}\check{\kappa}} \right) \\
&+ \varsigma^{-1} \left(\overline{\check{\Omega}\check{\zeta} \left(\overline{\frac{1}{2} \kappa^2 - \frac{1}{4} \vartheta^2} \right)} - \overline{\check{\Omega}\check{\zeta}\kappa\check{\kappa}} \right) - \overline{\check{\kappa}\check{\kappa}},
\end{aligned} \tag{2.2.37}$$

$$\begin{aligned}
Err[e_3(\check{\kappa})] &:= -\frac{1}{2} \overline{\check{\kappa}^2} - 2\overline{\check{\omega}\check{\kappa}} + 2(\eta - 3\zeta)\underline{\xi} - \overline{2(\eta - 3\zeta)\underline{\xi}} - \frac{1}{2} (\vartheta^2 - \overline{\vartheta^2}) \\
&- \varsigma^{-1} \left(\check{\zeta} \left(\overline{\frac{1}{2} \kappa^2 - 2\omega\kappa + 2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2} \vartheta^2} \right) - \overline{\check{\zeta}\check{\kappa}\check{\kappa}} \right) \\
&+ \varsigma^{-1} \overline{\check{\Omega}} \left(\check{\zeta} \left(\overline{\frac{1}{2} \kappa\kappa - 2\phi_1\zeta + 2\rho - \frac{1}{2} \vartheta\vartheta + 2\zeta^2} \right) - \overline{\check{\zeta}\check{\kappa}\check{\kappa}} \right) \\
&+ \varsigma^{-1} \left(\overline{\check{\Omega}\check{\zeta} \left(\overline{\frac{1}{2} \kappa\kappa - 2\phi_1\zeta + 2\rho - \frac{1}{2} \vartheta\vartheta + 2\zeta^2} \right)} - \overline{\check{\Omega}\check{\zeta}\kappa\check{\kappa}} \right) - \overline{\check{\kappa}^2},
\end{aligned} \tag{2.2.38}$$

and

$$\begin{aligned}
Err[e_3\check{\rho}] &:= - \left(\frac{1}{2} \vartheta\alpha + \zeta\beta - 2\eta\beta - 2\underline{\xi}\beta \right) + \overline{\left(\frac{1}{2} \vartheta\alpha + \zeta\beta - 2\eta\beta - 2\underline{\xi}\beta \right)} - \frac{3}{2} \overline{\check{\kappa}\check{\rho}} \\
&+ \varsigma^{-1} \check{\zeta} \left(\overline{-\frac{1}{2} \check{\kappa}\check{\rho} - \frac{1}{2} \vartheta\alpha - \zeta\beta + 2(\eta\beta + \underline{\xi}\beta)} \right) \\
&- \varsigma^{-1} \left(\check{\zeta} \left(\overline{-\frac{1}{2} \kappa\rho + \phi_1\beta - \frac{1}{2} \vartheta\alpha - \zeta\beta + 2(\eta\beta + \underline{\xi}\beta)} \right) - \overline{\check{\zeta}\check{\kappa}\check{\rho}} \right) \\
&- (\check{\Omega} + \varsigma^{-1} \overline{\check{\Omega}\check{\zeta}}) \left(\overline{-\frac{1}{2} \check{\kappa}\check{\rho} - \frac{1}{2} \vartheta\alpha - \zeta\beta} \right) \\
&+ \varsigma^{-1} \overline{\check{\Omega}} \left(\check{\zeta} \left(\overline{-\frac{1}{2} \kappa\rho + \phi_1\beta - \frac{1}{2} \vartheta\alpha - \zeta\beta} \right) - \overline{\check{\zeta}\check{\kappa}\check{\rho}} \right) \\
&+ \varsigma^{-1} \left(\overline{\check{\Omega}\check{\zeta} \left(\overline{-\frac{1}{2} \kappa\rho + \phi_1\beta - \frac{1}{2} \vartheta\alpha - \zeta\beta} \right)} - \overline{\check{\Omega}\check{\zeta}\kappa\check{\rho}} \right) - \overline{\check{\kappa}\check{\rho}}.
\end{aligned} \tag{2.2.39}$$

Proof. The proof relies on Corollary 2.2.11 and the null structure equations of Proposition 2.2.8. We refer to section A.4 for the details. \square

2.2.5 Additional equations

We derive below additional equations for $\underline{\omega}, \eta, \underline{\xi}$.

Proposition 2.2.19. *The following identities hold true for a general forward geodesic foliation.*

- The scalar $\underline{\omega}$ verifies

$$\begin{aligned} 2\phi_1^*\underline{\omega} &= -\frac{1}{2}\kappa\underline{\xi} + \left(\frac{1}{2}\kappa + 2\underline{\omega} + \frac{1}{2}\vartheta\right)\eta + e_3(\zeta) - \underline{\beta} \\ &\quad + \frac{1}{2}\kappa\zeta - 2\underline{\omega}\zeta + \frac{1}{2}\vartheta\zeta - \frac{1}{2}\vartheta\underline{\xi}. \end{aligned}$$

- The reduced 1-form η verifies

$$\begin{aligned} 2\phi_2\phi_2^*\eta &= \kappa(-e_3(\zeta) + \underline{\beta}) - e_3(e_\theta(\kappa)) - \kappa\left(\frac{1}{2}\kappa\zeta - 2\underline{\omega}\zeta\right) + 6\rho\eta - \underline{\kappa}e_\theta\kappa \\ &\quad - \frac{1}{2}\kappa e_\theta(\underline{\kappa}) + 2\underline{\omega}e_\theta(\kappa) + 2e_\theta(\rho) + \text{Err}[\phi_2\phi_2^*\eta], \\ \text{Err}[\phi_2\phi_2^*\eta] &= \left(2\phi_1\eta - \frac{1}{2}\kappa\vartheta + 2\eta^2\right)\eta + 2e_\theta(\eta^2) - \kappa\left(\frac{1}{2}\vartheta\zeta - \frac{1}{2}\vartheta\underline{\xi}\right) - \frac{1}{2}\vartheta e_\theta(\kappa) \\ &\quad - \left(2\phi_1\eta - \frac{1}{2}\vartheta\vartheta + 2\eta^2\right)\zeta - \frac{1}{2}e_\theta(\vartheta\vartheta) - \frac{1}{2}\vartheta^2\underline{\xi} - \frac{3}{2}\vartheta\vartheta\eta. \end{aligned}$$

- The reduced 1-form $\underline{\xi}$ verifies

$$\begin{aligned} 2\phi_2\phi_2^*\underline{\xi} &= -e_3(e_\theta(\underline{\kappa})) + \underline{\kappa}(e_3(\zeta) - \underline{\beta}) + \underline{\kappa}^2\zeta - \frac{3}{2}\underline{\kappa}e_\theta\underline{\kappa} + 6\rho\underline{\xi} - 2\underline{\omega}e_\theta(\underline{\kappa}) \\ &\quad + \text{Err}[\phi_2\phi_2^*\underline{\xi}], \\ \text{Err}[\phi_2\phi_2^*\underline{\xi}] &= \left(2\phi_1\underline{\xi} + \frac{1}{2}\kappa\vartheta + 2\eta\underline{\xi} - \frac{1}{2}\vartheta^2\right)\eta + 2e_\theta(\eta\underline{\xi}) - \frac{1}{2}e_\theta(\vartheta^2) \\ &\quad + \kappa\left(\frac{1}{2}\vartheta\zeta - \frac{1}{2}\vartheta\underline{\xi}\right) - \frac{1}{2}\vartheta e_\theta\kappa - \frac{1}{2}\vartheta\vartheta\underline{\xi} - \zeta\left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2\right) \\ &\quad + \underline{\xi}\left(-\vartheta\vartheta - 2\phi_1\zeta + 2\zeta^2\right) - 6\eta\zeta\underline{\xi} - 6e_\theta(\zeta\underline{\xi}). \end{aligned}$$

Proof. The proof relies on the null structure equations of Proposition 2.2.8, in particular the ones for $e_3(\zeta)$, $e_3(\kappa)$ and $e_3(\underline{\kappa})$. We refer to section A.5 for the details. \square

2.2.6 Ingoing geodesic foliation

All the equations of section 2.2.4 for outgoing geodesic foliations have a counterpart for ingoing geodesic foliations. The corresponding equations can be easily deduced from the ones in section 2.2.4 by performing the following substitutions

$$\begin{aligned}
u &\rightarrow \underline{u}, & s &\rightarrow s, & \mathcal{C}_u &\rightarrow \mathcal{C}_{\underline{u}}, & S_{u,s} &\rightarrow S_{\underline{u},s}, & r &\rightarrow r, & m &\rightarrow m, \\
e_4 &\rightarrow e_3, & e_3 &\rightarrow e_4, & e_\theta &\rightarrow e_\theta, & e_4(s) = 1 &\rightarrow e_3(s) = -1, \\
\alpha &\rightarrow \underline{\alpha}, & \beta &\rightarrow \underline{\beta}, & \rho &\rightarrow \rho, & \mu &\rightarrow \underline{\mu}, & \underline{\beta} &\rightarrow \beta, & \underline{\alpha} &\rightarrow \alpha, \\
\xi &\rightarrow \underline{\xi}, & \omega &\rightarrow \underline{\omega}, & \kappa &\rightarrow \underline{\kappa}, & \vartheta &\rightarrow \underline{\vartheta}, & \eta &\rightarrow \underline{\eta}, & \underline{\eta} &\rightarrow \eta, & \zeta &\rightarrow -\zeta, & \underline{\kappa} &\rightarrow \kappa, \\
\underline{\vartheta} &\rightarrow \vartheta, & \underline{\omega} &\rightarrow \omega, & \underline{\xi} &\rightarrow \xi, & \underline{\Omega} = e_3(s) &\rightarrow \Omega = e_4(s), & \varsigma = \frac{2}{e_3(u)} &\rightarrow \underline{\varsigma} = \frac{2}{e_4(\underline{u})}, \\
\bar{\kappa} - \frac{2}{r} &\rightarrow \underline{\bar{\kappa}} + \frac{2}{r}, & \bar{\kappa} + \frac{2\Upsilon}{r} &\rightarrow \underline{\bar{\kappa}} - \frac{2\Upsilon}{r}, & \bar{\omega} - \frac{m}{r^2} &\rightarrow \underline{\bar{\omega}} + \frac{m}{r^2}, & \bar{\rho} + \frac{2m}{r^3} &\rightarrow \underline{\bar{\rho}} + \frac{2m}{r^3}, \\
\bar{\mu} - \frac{2m}{r^3} &\rightarrow \underline{\bar{\mu}} - \frac{2m}{r^3}, & \bar{\Omega} + \Upsilon &\rightarrow \underline{\bar{\Omega}} - \Upsilon, & \bar{\varsigma} - 1 &\rightarrow \underline{\bar{\varsigma}} - 1, \\
\underline{A} = \frac{2}{r}e_3(r) - \bar{\kappa} &\rightarrow A = \frac{2}{r}e_4(r) - \bar{\kappa}.
\end{aligned}$$

2.2.7 Adapted coordinates systems

(u, s, θ, φ) coordinates

Proposition 2.2.20. *Consider, in addition to the functions u, s, φ an additional \mathbf{Z} invariant function θ . Then, relative to the coordinates system (u, s, θ, φ) , the following hold true,*

1. *The spacetime metric takes the form,*

$$g = -2\varsigma du ds + \varsigma^2 \underline{\Omega} du^2 + \gamma \left(d\theta - \frac{1}{2}\varsigma(\underline{b} - \underline{\Omega}b)du - bds \right)^2 + e^{2\Phi}(d\varphi)^2 \quad (2.2.40)$$

where,

$$\underline{\Omega} = e_3(s), \quad b = e_4(\theta), \quad \underline{b} = e_3(\theta), \quad \gamma^{-1} = e_\theta(\theta)^2. \quad (2.2.41)$$

2. *In these coordinates the reduced frame takes the form,*

$$\partial_s = e_4 - b\sqrt{\gamma}e_\theta, \quad \partial_u = \varsigma \left(\frac{1}{2}e_3 - \frac{1}{2}\underline{\Omega}e_4 - \frac{1}{2}\sqrt{\gamma}(\underline{b} - b\underline{\Omega})e_\theta \right), \quad \partial_\theta = \sqrt{\gamma}e_\theta. \quad (2.2.42)$$

3. In the particular case when $b = e_4(\theta) = 0$ we have,

$$e_4(\gamma) = 2\chi\gamma, \quad e_4(\underline{b}) = -2(\zeta + \eta)\gamma^{-1/2}. \quad (2.2.43)$$

Proof. First, from the fact that (e_3, e_4, e_θ) forms a null frame, we easily verify that (2.2.42) holds. Then, (2.2.40) immediately follows from (2.2.42) and the fact that (e_3, e_4, e_θ) forms a null frame.

To prove the last statement, when $b = e_4(\theta) = 0$, we start with,

$$[e_4, e_3] = 2\omega e_3 - 2\underline{\omega}e_4 + 2(\underline{\eta} - \eta)e_\theta = -2(\zeta + \eta)e_\theta - 2\underline{\omega}e_4.$$

Applying this to θ we derive,

$$[e_4, e_3](\theta) = (-2(\zeta + \eta)e_\theta - 2\underline{\omega}e_4)(\theta) = -2(\zeta + \eta)e_\theta(\theta) = -2(\zeta + \eta)\gamma^{-1/2}.$$

We deduce,

$$e_4(\underline{b}) = e_4(e_3(\theta)) = -2(\zeta + \eta)\gamma^{-1/2}.$$

To prove the equation for γ we make use of,

$$[e_4, e_\theta] = (\underline{\eta} + \zeta)e_4 + \xi e_3 - \chi e_\theta = -\chi e_\theta$$

so that

$$e_4 e_\theta(\theta) = [e_4, e_\theta](\theta) = -\chi e_\theta(\theta) = -\chi\gamma^{-1/2}.$$

Thus

$$e_4(\gamma^{-1/2}) = -\chi\gamma^{-1/2}$$

from which

$$e_4(\gamma) = 2\chi\gamma.$$

This concludes the proof of the lemma. \square

Remark 2.2.21. In Schwarzschild, relative to the above coordinate system, we have

$$\varsigma = 1, \quad \underline{\Omega} = -\Upsilon, \quad b = \underline{b} = 0, \quad \gamma = r^2, \quad e^\Phi = r \sin \theta,$$

so that we obtain outgoing Eddington-Finkelstein coordinates.

Remark 2.2.22. The (u, s, θ, φ) coordinates system, with the choice $b = 0$ (i.e. θ is transported by $e_4(\theta) = 0$), will be used in section 3.7 and Chapter 9 in connection with our GCM procedure.

(u, r, θ, φ) coordinates

Proposition 2.2.23. *Consider, in addition to the functions u, r, φ an additional \mathbf{Z} invariant function θ . Relative to the coordinates (u, r, θ, φ) the following hold true,*

1. *The spacetime metric takes the form,*

$$g = -\frac{4\varsigma}{r\bar{\kappa}} du dr + \frac{\varsigma^2(\bar{\kappa} + \underline{A})}{\bar{\kappa}} du^2 + \gamma \left(d\theta - \frac{1}{2}\varsigma \underline{b} du - \frac{\underline{b}}{2}\Theta \right)^2 \quad (2.2.44)$$

where,

$$b = e_4(\theta), \quad \underline{b} = e_3(\theta), \quad \gamma = \frac{1}{(e_\theta(\theta))^2} \quad (2.2.45)$$

and,

$$\Theta := \frac{4}{r\bar{\kappa}} dr - \varsigma \left(\frac{\bar{\kappa} + \underline{A}}{\bar{\kappa}} \right) du.$$

2. *The reduced coordinates derivatives take the form,*

$$\begin{aligned} \partial_r &= \frac{2}{r\bar{\kappa}} e_4 - \frac{2\sqrt{\gamma}}{r\bar{\kappa}} b e_\theta, \\ \partial_\theta &= \sqrt{\gamma} e_\theta, \\ \partial_u &= \varsigma \left[\frac{1}{2} e_3 - \frac{1}{2} \frac{\bar{\kappa} + \underline{A}}{\bar{\kappa}} e_4 - \frac{1}{2} \sqrt{\gamma} \left(\underline{b} - \left(\frac{\bar{\kappa} + \underline{A}}{\bar{\kappa}} \right) b \right) e_\theta \right]. \end{aligned} \quad (2.2.46)$$

3. *To control e^Φ , we will rely on the following transport equation*

$$e_4 \left(\frac{e^\Phi}{r \sin \theta} - 1 \right) = \frac{e^\Phi}{2r \sin \theta} (\check{\kappa} - \vartheta). \quad (2.2.47)$$

Proof. First, from the fact that (e_3, e_4, e_θ) forms a null frame, we easily verify that (2.2.46) holds. Then, (2.2.44) immediately follows from (2.2.46) and the fact that (e_3, e_4, e_θ) forms a null frame.

It remains to prove (2.2.47). It follows from

$$\begin{aligned} e_4 \left(\frac{e^\Phi}{r \sin \theta} - 1 \right) &= \frac{e^\Phi}{r \sin \theta} \left(e_4(\Phi) - \frac{e_4(r)}{r} \right) = \frac{e^\Phi}{r \sin \theta} \left(\frac{1}{2}(\kappa - \vartheta) - \frac{\bar{\kappa}}{2} \right) \\ &= \frac{e^\Phi}{2r \sin \theta} (\check{\kappa} - \vartheta) \end{aligned}$$

which concludes the proof of the lemma. \square

Remark 2.2.24. *In Schwarzschild, relative to the above coordinate system, we have*

$$\bar{\kappa} = \frac{2}{r}, \quad \underline{\bar{\kappa}} = -\frac{2\Upsilon}{r}, \quad \varsigma = 1, \quad \underline{A} = 0, \quad b = \underline{b} = 0, \quad \gamma = r^2, \quad e^\Phi = r \sin \theta,$$

so that we obtain outgoing Eddington-Finkelstein coordinates.

Remark 2.2.25. *The (u, r, θ, φ) coordinates system, with the choice (2.2.52) for θ introduced below, will be used in Proposition 3.4.3 to prove the convergence to the outgoing Eddington-Finkelstein coordinates of Schwarzschild.*

$(\underline{u}, r, \theta, \varphi)$ coordinates

We easily deduce an analog statement relative to $(\underline{u}, r, \theta, \varphi)$ coordinates.

Proposition 2.2.26. *Consider, in addition to the functions $\underline{u}, r, \varphi$ an additional \mathbf{Z} invariant function θ . Relative to the coordinates $(\underline{u}, r, \theta, \varphi)$ the following hold true,*

1. *The spacetime metric takes the form,*

$$g = -\frac{4\underline{\varsigma}}{r\underline{\bar{\kappa}}}d\underline{u}dr + \frac{\underline{\varsigma}^2(\bar{\kappa} + A)}{\bar{\kappa}}d\underline{u}^2 + \gamma \left(d\theta - \frac{1}{2}\underline{\varsigma}b d\underline{u} - \frac{b}{2}\underline{\Theta} \right)^2 \quad (2.2.48)$$

where,

$$b = e_4(\theta), \quad \underline{b} = e_3(\theta), \quad \gamma = \frac{1}{(e_\theta(\theta))^2} \quad (2.2.49)$$

and,

$$\underline{\Theta} := \frac{4}{r\underline{\bar{\kappa}}}dr - \underline{\varsigma} \left(\frac{\bar{\kappa} + A}{\bar{\kappa}} \right) d\underline{u}.$$

2. *The reduced coordinates derivatives take the form,*

$$\begin{aligned} \partial_r &= \frac{2}{r\underline{\bar{\kappa}}}e_3 - \frac{2\sqrt{\gamma}}{r\underline{\bar{\kappa}}}\underline{b}e_\theta, \\ \partial_\theta &= \sqrt{\gamma}e_\theta, \\ \partial_{\underline{u}} &= \underline{\varsigma} \left[\frac{1}{2}e_4 - \frac{1}{2}\frac{\bar{\kappa} + A}{\bar{\kappa}}e_3 - \frac{1}{2}\sqrt{\gamma} \left(b - \left(\frac{\bar{\kappa} + A}{\bar{\kappa}} \right) \underline{b} \right) e_\theta \right]. \end{aligned} \quad (2.2.50)$$

3. To control e^Φ , we will rely on the following transport equation

$$e_3 \left(\frac{e^\Phi}{r \sin \theta} - 1 \right) = \frac{e^\Phi}{2r \sin \theta} (\underline{\check{\kappa}} - \underline{\vartheta}). \quad (2.2.51)$$

Remark 2.2.27. In Schwarzschild, relative to the above coordinate system, we have

$$\bar{\kappa} = \frac{2}{r}, \quad \bar{\underline{\kappa}} = -\frac{2\underline{\Upsilon}}{r}, \quad \underline{\varsigma} = 1, \quad A = 0, \quad b = \underline{b} = 0, \quad \gamma = r^2, \quad e^\Phi = r \sin \theta,$$

so that we obtain ingoing Eddington-Finkelstein coordinates.

Remark 2.2.28. The $(\underline{u}, r, \theta, \varphi)$ coordinates system, with the choice (2.2.52) for θ introduced below, will be used in Proposition 3.4.4 to prove the convergence to the ingoing Eddington-Finkelstein coordinates of Schwarzschild.

Initialization of θ

We now introduce the coordinate function θ that will be used for the (u, r, θ, φ) coordinates system and for the $(\underline{u}, r, \theta, \varphi)$ coordinates system, see Remarks 2.2.25 and 2.2.28.

Lemma 2.2.29. Let $\theta \in [0, \pi]$ be the \mathbf{Z} -invariant scalar on \mathcal{M} defined by,

$$\theta := \cot^{-1}(re_\theta(\Phi)). \quad (2.2.52)$$

Then,

$$\frac{e^\Phi}{r \sin \theta} = \sqrt{1 + \mathbf{a}}. \quad (2.2.53)$$

where,

$$\mathbf{a} := \frac{e^{2\Phi}}{r^2} + (e_\theta(e^\Phi))^2 - 1. \quad (2.2.54)$$

Moreover, we have in an outgoing geodesic foliation

$$\begin{aligned} re_\theta(\theta) &= 1 + \frac{r^2(K - \frac{1}{r^2})}{1 + (re_\theta(\Phi))^2}, \\ e_3(\theta) &= -\frac{r\underline{\beta} + \frac{r}{2}(-\underline{\check{\kappa}} + \underline{A} + \underline{\vartheta})e_\theta(\Phi) + r\underline{\xi}e_4(\Phi) + r\underline{\eta}e_3(\Phi)}{1 + (re_\theta(\Phi))^2}, \\ e_4(\theta) &= -\frac{r\underline{\beta} + \frac{r}{2}(-\underline{\check{\kappa}} + \underline{\vartheta})e_\theta(\Phi) - r\underline{\zeta}e_3(\Phi)}{1 + (re_\theta(\Phi))^2}, \end{aligned}$$

and analog identities hold for an ingoing geodesic foliation.

Proof. In view of the definition of θ , we have $\theta \in [0, \pi]$, $\sin \theta \geq 0$ and

$$\begin{aligned} \sin \theta &= \frac{1}{\sqrt{1 + \cot^2 \theta}} = \frac{1}{\sqrt{1 + (re_\theta(\Phi))^2}} = \frac{e^\Phi}{\sqrt{e^{2\Phi} + (re_\theta(e^\Phi))^2}} \\ &= \frac{e^\Phi}{r\sqrt{\frac{e^{2\Phi}}{r^2} + (e_\theta(e^\Phi))^2}} = \frac{e^\Phi}{r\sqrt{1 + \mathfrak{a}}}. \end{aligned}$$

Hence

$$\frac{e^\Phi}{r \sin \theta} = \sqrt{\frac{e^{2\Phi}}{r^2} + (e_\theta(e^\Phi))^2} = \sqrt{1 + \mathfrak{a}}.$$

Also, we compute

$$re_\theta(\theta) = -\frac{r^2 e_\theta e_\theta(\Phi)}{1 + (re_\theta(\Phi))^2}.$$

Next, recall that we have

$$e_\theta e_\theta(\Phi) = -K - (e_\theta(\Phi))^2.$$

We infer

$$re_\theta(\theta) = \frac{r^2(K + (e_\theta(\Phi))^2)}{1 + (re_\theta(\Phi))^2} = 1 + \frac{r^2(K - \frac{1}{r^2})}{1 + (re_\theta(\Phi))^2}.$$

as desired.

Also, we have in an outgoing geodesic foliation

$$\begin{aligned} e_4(\theta) &= -\frac{re_4 e_\theta(\Phi) + e_4(r)e_\theta(\Phi)}{1 + (re_\theta(\Phi))^2} \\ &= -\frac{r(D_4 D_\theta \Phi + D_{D_4 e_\theta} \Phi) + e_4(r)e_\theta(\Phi)}{1 + (re_\theta(\Phi))^2} \\ &= -\frac{r\beta + r\left(\frac{e_4(r)}{r} - e_4(\Phi)\right)e_\theta(\Phi) - r\zeta e_4(\Phi)}{1 + (re_\theta(\Phi))^2} \\ &= -\frac{r\beta + \frac{r}{2}(-\check{\kappa} + \vartheta)e_\theta(\Phi) - r\zeta e_4(\Phi)}{1 + (re_\theta(\Phi))^2}. \end{aligned}$$

Finally, we compute in an outgoing geodesic foliation

$$\begin{aligned}
e_3(\theta) &= -\frac{re_3e_\theta(\Phi) + e_3(r)e_\theta(\Phi)}{1 + (re_\theta(\Phi))^2} \\
&= -\frac{r(D_3D_\theta\Phi + D_{D_3e_\theta}\Phi) + e_3(r)e_\theta(\Phi)}{1 + (re_\theta(\Phi))^2} \\
&= -\frac{r\underline{\beta} + r\left(\frac{e_3(r)}{r} - e_3(\Phi)\right)e_\theta(\Phi) + r\underline{\xi}e_4(\Phi) + r\eta e_3(\Phi)}{1 + (re_\theta(\Phi))^2} \\
&= -\frac{r\underline{\beta} + \frac{r}{2}(-\check{\kappa} + \underline{A} + \underline{\vartheta})e_\theta(\Phi) + r\underline{\xi}e_4(\Phi) + r\eta e_3(\Phi)}{1 + (re_\theta(\Phi))^2}.
\end{aligned}$$

This concludes the proof of the lemma. \square

In view of (2.2.53), we will need to control the quantity \mathbf{a} defined in (2.2.54). To this end, we will need the following lemma.

Lemma 2.2.30. *The quantity \mathbf{a} defined in (2.2.54) vanishes on the axis of symmetry and verifies the following identities in an outgoing geodesic foliation,*

$$\begin{aligned}
e_4(\mathbf{a}) &= \frac{(\check{\kappa} - \vartheta)e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)\left(\beta - e_4(\Phi)\zeta\right)e^\Phi, \\
e_\theta(\mathbf{a}) &= 2e_\theta(\Phi)e^{2\Phi}\left(\left(\rho + \frac{2m}{r^3}\right) + \frac{1}{4}\left(\kappa\underline{\kappa} + \frac{4\underline{\Upsilon}}{r^2}\right) - \frac{1}{4}\vartheta\underline{\vartheta}\right), \\
e_3(\mathbf{a}) &= \frac{(\check{\kappa} - \underline{A} - \underline{\vartheta})e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)\left(\underline{\beta} + e_3(\Phi)\eta + \underline{\xi}e_4(\Phi)\right)e^\Phi,
\end{aligned}$$

and analog identities hold in an outgoing geodesic foliation.

Proof. The vanishing on the axis follow easily from the fact that both $e^{2\Phi}$ and $e_\theta(e^\Phi)^2 - 1$ vanish on the axis (see (2.1.13)). To prove the second part of the lemma we recall that, with respect to the reduced metric (see equation (2.1.4)),

$$R_{ab} = D_aD_b\Phi + D_a\Phi D_b\Phi,$$

and (see Definition 2.1.47)

$$R_{3\theta} = \underline{\beta}, \quad R_{4\theta} = \beta, \quad R_{\theta\theta} = R_{34} = \rho, \quad R_{34} = \rho.$$

Starting with the definition $\mathbf{a} = \frac{e^{2\Phi}}{r^2} + (e_\theta(e^\Phi))^2 - 1$, we compute in an outgoing geodesic foliation

$$\begin{aligned} e_4(\mathbf{a}) &= \frac{2e_4(\Phi)e^{2\Phi}}{r^2} - \frac{2e_4(r)e^{2\Phi}}{r^3} + 2e_\theta(e^\Phi)e_4(e_\theta(e^\Phi)) \\ &= \frac{(\underline{\kappa} - \underline{\vartheta})e^{2\Phi}}{r^2} - \frac{\overline{\kappa}e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)\left(e_4(e_\theta(\Phi)) + e_\theta(\Phi)e_4(\Phi)\right)e^\Phi \\ &= \frac{(\check{\kappa} - \underline{\vartheta})e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)\left(\beta - e_4(\Phi)\zeta\right)e^\Phi. \end{aligned}$$

Also

$$\begin{aligned} e_\theta(\mathbf{a}) &= \frac{2e_\theta(\Phi)e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)e_\theta(e_\theta(e^\Phi)) \\ &= \frac{2e_\theta(\Phi)e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)\left(e_\theta(e_\theta(\Phi)) + e_\theta(\Phi)^2\right)e^\Phi \\ &= \frac{2e_\theta(\Phi)e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)\left(\rho + D_{D_\theta e_\theta}\Phi\right)e^\Phi \\ &= \frac{2e_\theta(\Phi)e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)\left(\rho + \frac{1}{2}\chi e_3\Phi + \frac{1}{2}\underline{\chi}e_4\Phi\right)e^\Phi \\ &= \frac{2e_\theta(\Phi)e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)\left(\rho + \frac{1}{4}\kappa\underline{\kappa} - \frac{1}{4}\underline{\vartheta}\underline{\vartheta}\right)e^\Phi \\ &= 2e_\theta(\Phi)e^{2\Phi}\left(\left(\rho + \frac{2m}{r^3}\right) + \frac{1}{4}\left(\kappa\underline{\kappa} + \frac{4\Upsilon}{r^2}\right) - \frac{1}{4}\underline{\vartheta}\underline{\vartheta}\right). \end{aligned}$$

Finally, we have in an outgoing geodesic foliation

$$\begin{aligned} e_3(\mathbf{a}) &= \frac{2e_3(\Phi)e^{2\Phi}}{r^2} - \frac{2e_3(r)e^{2\Phi}}{r^3} + 2e_\theta(e^\Phi)e_3(e_\theta(e^\Phi)) \\ &= \frac{(\underline{\kappa} - \underline{\vartheta})e^{2\Phi}}{r^2} - \frac{(\overline{\kappa} + \underline{A})e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)\left(e_3(e_\theta(\Phi)) + e_\theta(\Phi)e_3(\Phi)\right)e^\Phi \\ &= \frac{(\check{\kappa} - \underline{A} - \underline{\vartheta})e^{2\Phi}}{r^2} + 2e_\theta(e^\Phi)\left(\underline{\beta} + e_3(\Phi)\eta + \underline{\xi}e_4(\Phi)\right)e^\Phi. \end{aligned}$$

This concludes the proof of the lemma. \square

Remark 2.2.31. *The function θ defined by (3.4.3) defines*

- together with the functions (u, r, φ) , a regular coordinates system with the axis of symmetry corresponding to $\theta = 0, \pi$,
- together with the functions (u, r, φ) , a regular coordinates system with the axis of symmetry corresponding to $\theta = 0, \pi$.

2.3 Perturbations of Schwarzschild and invariant quantities

Recall that in Schwarzschild all Ricci coefficients $\xi, \underline{\xi}, \vartheta, \underline{\vartheta}, \eta, \underline{\eta}, \zeta$ and curvature components $\alpha, \underline{\alpha}, \beta, \underline{\beta}$ vanish identically. In addition the check quantities $\check{\kappa}, \underline{\check{\kappa}}, \check{\omega}, \underline{\check{\omega}}$ and $\check{\rho}$ also vanish. Thus, roughly, we expect that in perturbations of Schwarzschild these quantities stay small, i.e. of order $O(\epsilon)$ for a sufficiently small ϵ . More precisely we say that a smooth, vacuum, \mathbf{Z} -invariant, polarized spacetime is an $O(\epsilon)$ -perturbation of Schwarzschild, or simply $O(\epsilon)$ -Schwarzschild, if the following are true relative to a \mathbf{Z} -invariant null frame e_3, e_4, e_θ ,

$$\xi, \underline{\xi}, \vartheta, \underline{\vartheta}, \eta, \underline{\eta}, \zeta, \check{\kappa}, \underline{\check{\kappa}}, \check{\omega}, \underline{\check{\omega}} \quad \alpha, \underline{\alpha}, \beta, \underline{\beta}, \check{\rho} = O(\epsilon) \quad (2.3.1)$$

Moreover,

$$e_3(r) - \frac{r}{2}\underline{\kappa} = O(\epsilon), \quad e_4(r) - \frac{r}{2}\bar{\kappa} = O(\epsilon), \quad (2.3.2)$$

where r is the area radius of the 2-spheres generated by e_θ, e_φ , see (2.1.12).

In reality, of course, we expect that small perturbations of Schwarzschild, remain not only close to the original Schwarzschild but also converge to a nearby Schwarzschild solution but for the discussion below this will suffice.

2.3.1 Null frame transformations

Our definition of $O(\epsilon)$ -Schwarzschild perturbations does not specify a particular frame. In what follows we investigate how the main Ricci and curvature quantities change relative to frame transformations, i.e linear transformations which take null frames into null frames.

Lemma 2.3.1. *A general null transformation can be written in the form,*

$$\begin{aligned} e'_4 &= \lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right), \\ e'_\theta &= \left(1 + \frac{1}{2} f \underline{f} \right) e_\theta + \frac{1}{2} \underline{f} e_4 + \frac{1}{2} f \left(1 + \frac{1}{4} f \underline{f} \right) e_3, \\ e'_3 &= \lambda^{-1} \left(\left(1 + \frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) e_3 + \underline{f} \left(1 + \frac{1}{4} f \underline{f} \right) e_\theta + \frac{1}{4} \underline{f}^2 e_4 \right). \end{aligned} \quad (2.3.3)$$

Proof. It is straightforward to check that the transformation (2.3.3) takes null frames into null frames. One can also check that it can be written in the form $\text{type}(3) \circ \text{type}(1) \circ$

2.3. PERTURBATIONS OF SCHWARZSCHILD AND INVARIANT QUANTITIES 95

type(2) where the type 1 transformations fix e_3 , i.e. ($\lambda = 1, \underline{f} = 0$), type 2 transformations fix e_4 , i.e. ($\lambda = 1, \underline{f} = 0$) and type 3 transformations keep the directions of e_3, e_4 i.e. ($\underline{f} = \underline{f} = 0$). \square

Remark 2.3.2. Note that f, \underline{f} are reduced from spacetime 1 forms while λ is reduced from a scalar.

Remark 2.3.3. A transformation consistent with $O(\epsilon)$ - Schwarzschild spacetimes must have $f, \underline{f} = O(\epsilon)$ and $a := \log \lambda = O(\epsilon)$.

Proposition 2.3.4 (Transformation formulas). Under a general transformation of type (2.3.3), the Ricci coefficients and curvature components transform as follows:

$$\begin{aligned}\xi' &= \lambda^2 \left(\xi + \frac{1}{2} \lambda^{-1} e'_4(f) + \omega f + \frac{1}{4} f \kappa \right) + \lambda^2 \text{Err}(\xi, \xi'), \\ \text{Err}(\xi, \xi') &= \frac{1}{4} f \vartheta + l.o.t., \\ \underline{\xi}' &= \lambda^{-2} \left(\underline{\xi} + \frac{1}{2} \lambda e'_3(\underline{f}) + \underline{\omega} \underline{f} + \frac{1}{4} \underline{f} \underline{\kappa} \right) + \lambda^{-2} \text{Err}(\underline{\xi}, \underline{\xi}'), \\ \text{Err}(\underline{\xi}, \underline{\xi}') &= -\frac{1}{8} \lambda \underline{f}^2 e'_3(f) + \frac{1}{4} \underline{f} \underline{\vartheta} + l.o.t.,\end{aligned}\tag{2.3.4}$$

$$\begin{aligned}\zeta' &= \zeta - e'_\theta(\log(\lambda)) + \frac{1}{4} (-f \underline{\kappa} + \underline{f} \kappa) + \underline{f} \omega - f \underline{\omega} + \text{Err}(\zeta, \zeta'), \\ \text{Err}(\zeta, \zeta') &= \frac{1}{2} \underline{f} e'_\theta(f) + \frac{1}{4} (-f \underline{\vartheta} + \underline{f} \vartheta) + l.o.t., \\ \eta' &= \eta + \frac{1}{2} \lambda e'_3(f) + \frac{1}{4} \kappa \underline{f} - f \underline{\omega} + \text{Err}(\eta, \eta'), \\ \text{Err}(\eta, \eta') &= \frac{1}{4} \underline{f} \vartheta + l.o.t., \\ \underline{\eta}' &= \underline{\eta} + \frac{1}{2} \lambda^{-1} e'_4(\underline{f}) + \frac{1}{4} \underline{\kappa} f - \underline{f} \omega + \text{Err}(\underline{\eta}, \underline{\eta}'), \\ \text{Err}(\underline{\eta}, \underline{\eta}') &= -\frac{1}{8} \underline{f}^2 \lambda^{-1} e'_4(f) + \frac{1}{4} \underline{f} \underline{\vartheta} + l.o.t.,\end{aligned}\tag{2.3.5}$$

$$\begin{aligned}\kappa' &= \lambda (\kappa + \phi'_1(f)) + \lambda \text{Err}(\kappa, \kappa'), \\ \text{Err}(\kappa, \kappa') &= f(\zeta + \eta) + \underline{f} \xi - \frac{1}{4} f^2 \underline{\kappa} + f \underline{f} \omega - f^2 \underline{\omega} + l.o.t., \\ \underline{\kappa}' &= \lambda^{-1} (\underline{\kappa} + \phi'_1(\underline{f})) + \lambda^{-1} \text{Err}(\underline{\kappa}, \underline{\kappa}'), \\ \text{Err}(\underline{\kappa}, \underline{\kappa}') &= -\frac{1}{4} \underline{f}^2 e'_\theta(f) + \underline{f}(-\zeta + \underline{\eta}) + \underline{f} \underline{\xi} - \frac{1}{4} \underline{f}^2 \underline{\kappa} + \underline{f} \underline{f} \omega - \underline{f}^2 \underline{\omega} + l.o.t.,\end{aligned}\tag{2.3.6}$$

$$\begin{aligned}
\vartheta' &= \lambda(\vartheta - \mathcal{A}_2^*(f)) + \lambda \text{Err}(\vartheta, \vartheta'), \\
\text{Err}(\vartheta, \vartheta') &= f(\zeta + \eta) + \underline{f}\xi + \frac{1}{4}f\underline{f}\kappa + f\underline{f}\omega - f^2\underline{\omega} + l.o.t. \\
\underline{\vartheta}' &= \lambda^{-1}(\underline{\vartheta} - \mathcal{A}_2^*(\underline{f})) + \lambda^{-1} \text{Err}(\underline{\vartheta}, \underline{\vartheta}'), \\
\text{Err}(\underline{\vartheta}, \underline{\vartheta}') &= -\frac{1}{4}\underline{f}^2 e'_\theta(f) + \underline{f}(-\zeta + \eta) + \underline{f}\xi + \frac{1}{4}f\underline{f}\kappa + f\underline{f}\omega - \underline{f}^2\omega + l.o.t.,
\end{aligned} \tag{2.3.7}$$

$$\begin{aligned}
\omega' &= \lambda \left(\omega - \frac{1}{2}\lambda^{-1}e'_4(\log(\lambda)) \right) + \lambda \text{Err}(\omega, \omega'), \\
\text{Err}(\omega, \omega') &= \frac{1}{4}\underline{f}e'_4(f) + \frac{1}{2}\omega\underline{f}\underline{f} - \frac{1}{2}\underline{f}\eta + \frac{1}{2}\underline{f}\xi + \frac{1}{2}\underline{f}\zeta - \frac{1}{8}\underline{\kappa}f^2 + \frac{1}{8}f\underline{f}\kappa - \frac{1}{4}\underline{\omega}f^2 + l.o.t., \\
\underline{\omega}' &= \lambda^{-1} \left(\underline{\omega} + \frac{1}{2}\lambda e'_3(\log(\lambda)) \right) + \lambda^{-1} \text{Err}(\underline{\omega}, \underline{\omega}'), \\
\text{Err}(\underline{\omega}, \underline{\omega}') &= -\frac{1}{4}\underline{f}e'_3(f) + \underline{\omega}\underline{f}\underline{f} - \frac{1}{2}\underline{f}\eta + \frac{1}{2}\underline{f}\xi - \frac{1}{2}\underline{f}\zeta - \frac{1}{8}\underline{\kappa}\underline{f}^2 + \frac{1}{8}f\underline{f}\kappa - \frac{1}{4}\underline{\omega}\underline{f}^2 + l.o.t.
\end{aligned} \tag{2.3.8}$$

The lower order terms we denote by *l.o.t.* are linear with respect $\Gamma = \{\xi, \underline{\xi}, \vartheta, \kappa, \eta, \underline{\eta}, \zeta, \underline{\kappa}, \underline{\vartheta}\}$ and quadratic or higher order in f, \underline{f} , and do not contain derivatives of these latter.

Also,

$$\begin{aligned}
\alpha' &= \lambda^2\alpha + \lambda^2 \text{Err}(\alpha, \alpha'), \\
\text{Err}(\alpha, \alpha') &= 2f\beta + \frac{3}{2}f^2\rho + l.o.t., \\
\beta' &= \lambda \left(\beta + \frac{3}{2}\rho f \right) + \lambda \text{Err}(\beta, \beta'), \\
\text{Err}(\beta, \beta') &= \frac{1}{2}\underline{f}\alpha + l.o.t., \\
\rho' &= \rho + \text{Err}(\rho, \rho'), \\
\text{Err}(\rho, \rho') &= \frac{3}{2}\rho\underline{f}\underline{f} + \underline{f}\beta + f\underline{\beta} + l.o.t., \\
\underline{\beta}' &= \lambda^{-1} \left(\underline{\beta} + \frac{3}{2}\rho\underline{f} \right) + \lambda^{-1} \text{Err}(\underline{\beta}, \underline{\beta}'), \\
\text{Err}(\underline{\beta}, \underline{\beta}') &= \frac{1}{2}f\underline{\alpha} + l.o.t., \\
\underline{\alpha}' &= \lambda^{-2}\underline{\alpha} + \lambda^{-2} \text{Err}(\underline{\alpha}, \underline{\alpha}'), \\
\text{Err}(\underline{\alpha}, \underline{\alpha}') &= 2\underline{f}\underline{\beta} + \frac{3}{2}\underline{f}^2\rho + l.o.t.
\end{aligned} \tag{2.3.9}$$

The lower order terms we denote by *l.o.t.* are linear with respect to the curvature quantities $\alpha, \beta, \rho, \underline{\beta}, \underline{\alpha}$ and quadratic or higher order in f, \underline{f} , and do not contain derivatives of these latter.

2.3. PERTURBATIONS OF SCHWARZSCHILD AND INVARIANT QUANTITIES 97

Proof. See Appendix A.6. □

Lemma 2.3.5. *In the particular case when $\lambda = 1, \underline{f} = 0$, we have*

$$\begin{aligned} e'_4 &= e_4 + f e_\theta + \frac{1}{4} f^2 e_3, \\ e'_\theta &= e_\theta + \frac{1}{2} f e_3, \\ e'_3 &= e_3, \end{aligned}$$

and

$$\begin{aligned} \xi' &= \xi + \frac{1}{2} e'_4 f + \frac{1}{4} \kappa f + f \omega + \frac{1}{4} f \vartheta + \frac{1}{4} f^2 \eta - \frac{1}{4} f^2 \underline{\eta} + \frac{1}{2} f^2 \zeta - \frac{1}{16} f^3 \underline{\kappa} \\ &\quad - \frac{1}{4} f^3 \underline{\omega} - \frac{1}{16} f^3 \underline{\vartheta} - \frac{1}{16} f^4 \underline{\xi}, \\ \omega' &= \omega + \frac{1}{2} f \zeta - \frac{1}{2} \underline{\eta} f - \frac{1}{4} f^2 \underline{\omega} - \frac{1}{8} f^2 \underline{\kappa} - \frac{1}{8} f^2 \underline{\vartheta} - \frac{1}{8} f^3 \underline{\xi}, \\ \zeta' &= \zeta - \left(\frac{1}{4} \underline{\kappa} + \underline{\omega} \right) f - f \left(\frac{1}{4} \underline{\vartheta} + \frac{1}{2} f \underline{\xi} \right), \\ \eta' &= \eta + \frac{1}{2} e'_3(f) - f \underline{\omega} - \frac{1}{4} f^2 \underline{\xi}. \end{aligned}$$

Proof. The proof follows from Proposition 2.3.4 by setting $\lambda = 1, \underline{f} = 0$. Since we need precise formulas for the error terms, we provide a proof in section A.9. □

Lemma 2.3.6 (Transport equations for $(f, \underline{f}, \lambda)$). *Assume that we have in the new null frame (e'_3, e'_4, e'_θ) of type (2.3.3)*

$$\xi' = 0, \quad \omega' = 0, \quad \zeta' + \underline{\eta}' = 0.$$

Then, $(\underline{f}, f, \log(\lambda))$ satisfy the following transport equations

$$\begin{aligned} \lambda^{-1} e'_4(f) + \left(\frac{\kappa}{2} + 2\omega \right) f &= -2\xi + E_1(f, \Gamma), \\ \lambda^{-1} e'_4(\log(\lambda)) &= 2\omega + E_2(f, \Gamma), \\ \lambda^{-1} e'_4(\underline{f}) + \frac{\kappa}{2} \underline{f} &= -2(\zeta + \underline{\eta}) + 2e'_\theta(\log(\lambda)) + 2f \underline{\omega} + E_3(f, \underline{f}, \Gamma), \end{aligned}$$

where E_1, E_2 and E_3 are given by

$$\begin{aligned} E_1(f, \Gamma) &= -\frac{1}{2} \vartheta f + l.o.t., \\ E_2(f, \Gamma) &= f \zeta - \frac{1}{2} f^2 \underline{\omega} - \underline{\eta} f - \frac{1}{4} f^2 \underline{\kappa} + l.o.t., \\ E_3(f, \underline{f}, \Gamma) &= -\underline{f} e'_\theta(f) - \frac{1}{2} \underline{f} \vartheta + l.o.t., \end{aligned}$$

Here, *l.o.t.* denote terms which are cubic or higher order in f, \underline{f} (or in f only in the case of E_1 and E_2) and $\check{\Gamma}$ and do not contain derivatives of these quantities, where Γ and $\check{\Gamma}$ denotes the Ricci coefficients and renormalized Ricci coefficients w.r.t. the original null frame (e_3, e_4, e_θ) .

Proof. See section A.7. □

To avoid a potential log loss for the third equation in Lemma 2.3.6, i.e. the transport equation for \underline{f} , we state the following renormalized version of the lemma.

Corollary 2.3.7. *Assume given a null frame (e_3, e_4, e_θ) associated to an outgoing geodesic foliation as in section 2.2.4, and let r denote the corresponding area radius. Assume that we have in the new null frame (e'_3, e'_4, e'_θ) of type (2.3.3)*

$$\xi' = 0, \quad \omega' = 0, \quad \zeta' + \underline{\eta}' = 0.$$

Then, $(\underline{f}, f, \log(\lambda))$ satisfy the following transport equations

$$\begin{aligned} \lambda^{-1}e'_4(rf) &= E'_1(f, \Gamma), \\ \lambda^{-1}e'_4(\log(\lambda)) &= E'_2(f, \Gamma), \\ \lambda^{-1}e'_4\left(r\underline{f} - 2r^2e'_\theta(\log(\lambda)) + rf\underline{\Omega}\right) &= E'_3(f, \underline{f}, \lambda, \Gamma), \end{aligned}$$

where

$$\begin{aligned} E'_1(f, \Gamma) &= -\frac{r}{2}\check{\kappa}f - \frac{r}{2}\vartheta f + l.o.t., \\ E'_2(f, \Gamma) &= f\zeta - \frac{1}{2}f^2\underline{\omega} - \underline{\eta}f - \frac{1}{4}f^2\underline{\kappa} + l.o.t., \\ E'_3(f, \underline{f}, \lambda, \Gamma) &= -\frac{r}{2}\check{\kappa}\underline{f} + r^2\left(\check{\kappa} - \left(\bar{\kappa} - \frac{2}{r}\right)\right)e'_\theta(\log(\lambda)) + r^2\left(\not{d}'_1(f) + \lambda^{-1}\vartheta'\right)e'_\theta(\log(\lambda)) \\ &\quad - \frac{r}{2}\check{\kappa}\underline{\Omega}f + rE_3(f, \underline{f}, \Gamma) - 2r^2e'_\theta(E_2(f, \Gamma)) + r\underline{\Omega}E_1(f, \Gamma), \end{aligned}$$

and where E_1, E_2 and E_3 are given in Lemma 2.3.6.

Proof. See section A.8. □

2.3.2 Schematic notation Γ_g and Γ_b

Many of the identities which we present below, contain a huge number of $O(\epsilon^2)$ terms. In what follows we introduce schematic notation meant to keep track of the most important

error terms. Note that the decomposition below between the terms Γ_g and Γ_b is consistent with our main bootstrap assumptions **BA-E** on energy and **BA-D** on decay, see section 3.4.1.

Definition 2.3.8. *We divide the small connection coefficient terms (relative to an arbitrary null frame) into*¹⁷

$$\Gamma_g^{(0)} = \left\{ r\xi, \vartheta, \zeta, \underline{\eta}, \frac{2}{r}e_4(r) - \kappa, \frac{1}{r}e_\theta(r) \right\}, \quad \Gamma_b^{(0)} = \left\{ \eta, \underline{\vartheta}, \underline{\xi}, \frac{2}{r}e_3(r) - \underline{\kappa} \right\}.$$

For higher derivatives we introduce,

$$\Gamma_g^{(1)} = \left\{ \mathfrak{d}\Gamma_g^{(0)}, r^2e_\theta(\omega), re_\theta(\kappa), re_\theta(\underline{\kappa}) \right\}, \quad \Gamma_b^{(1)} = \left\{ \mathfrak{d}\Gamma_b^{(0)}, re_\theta(\underline{\omega}) \right\},$$

and for $s \geq 2$,

$$\Gamma_g^{(s)} = \mathfrak{d}^{\leq s}\Gamma_g, \quad \Gamma_b^{(s)} = \mathfrak{d}^{\leq s}\Gamma_b,$$

where we have introduced the notations

$$\mathfrak{d} = \{e_3, re_4, \mathfrak{d}\},$$

with angular derivatives \mathfrak{d} of reduced scalars in \mathfrak{s}_k defined by (2.1.37).

Remark 2.3.9. *According to the main bootstrap assumptions **BA-E**, **BA-D** (see section 3.4.1), the terms Γ_b behave worse in powers of r than the terms in Γ_g . Thus, in the calculations below, we replace the terms of the form $\Gamma_g^{(s)} + \Gamma_b^{(s)}$ by $\Gamma_b^{(s)}$. Given the form of the bootstrap assumptions, we may also replace $r^{-1}\Gamma_b^{(s)}$ by $\Gamma_g^{(s)}$. We will denote l.o.t. the cubic and higher error terms in $\check{\Gamma}, \check{R}$. We also include in l.o.t. terms which decay faster in powers of r than the main quadratic terms.*

2.3.3 The invariant quantity \mathfrak{q}

Note from the transformation formulas of Proposition 2.3.4 that the only quantities which remain invariant up to quadratic or higher order error terms are α , $\underline{\alpha}$ and ρ . Among these only α , $\underline{\alpha}$ vanish in Schwarzschild. We call such quantities $O(\epsilon^2)$ invariant. In what follows we show that, in addition to these two invariants, there exist other important invariants.

¹⁷In the frames we are using, we have in fact $\xi = 0$ for $r \geq 4m_0$ so that it behaves in fact better than the other components of $\Gamma_g^{(0)}$.

Lemma 2.3.10. *The expression,*

$$e_3(e_3(\alpha)) + (2\underline{\kappa} - 6\underline{\omega})e_3(\alpha) + \left(-4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2\right)\alpha$$

is an $O(\epsilon^2)$ invariant. It is also a conformal invariant, i.e. invariant under transformations (2.3.3) with $f = \underline{f} = 0$.

Proof. Clearly the quantity vanishes in Schwarzschild and is an $O(\epsilon^2)$ invariant. For a conformal transformation, the result follows by a straightforward application of the transformation properties of Proposition 2.3.4 in the particular case where $f = \underline{f} = 0$. \square

Remark 2.3.11. *Alternatively one can also define the corresponding quantity obtained by interchanging e_3, e_4 , i.e.*

$$e_4(e_4(\underline{\alpha})) + (2\kappa - 6\omega)e_4(\underline{\alpha}) + \left(-4e_4(\omega) + 8\omega^2 - 8\omega\kappa + \frac{1}{2}\kappa^2\right)\underline{\alpha}.$$

Note that it differs by $O(\epsilon^2)$ from the previous one.

Definition 2.3.12. *Given a general null frame (e_4, e_3, e_θ) , and given a scalar function r satisfying the assumptions for section 2.3.2, i.e.*

$$\frac{2}{r}e_4(r) - \kappa \in \Gamma_g, \quad \frac{1}{r}e_\theta(r) \in \Gamma_g, \quad \frac{2}{r}e_3(r) - \underline{\kappa} \in \Gamma_b,$$

we defined our main quantity \mathfrak{q} as

$$\mathfrak{q} := r^4 \left[e_3(e_3(\alpha)) + (2\underline{\kappa} - 6\underline{\omega})e_3(\alpha) + \left(-4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2\right)\alpha \right]. \quad (2.3.10)$$

2.3.4 Several identities for \mathfrak{q}

In this section, we state three identities involving the quantity \mathfrak{q} defined by (2.3.10). All calculations are made in a general frame.

Proposition 2.3.13. *We have*

$$\mathfrak{q} = r^4 \left(\not\phi_2^* \not\phi_1^* \rho + \frac{3}{4} \underline{\kappa} \rho \not\vartheta + \frac{3}{4} \kappa \rho \not\vartheta \right) + \text{Err}[\mathfrak{q}] \quad (2.3.11)$$

with error term written schematically in the form

$$\text{Err}[\mathfrak{q}] = r^4 e_3 \eta \cdot \beta + r^2 \mathfrak{d}^{\leq 1}(\Gamma_b \cdot \Gamma_g). \quad (2.3.12)$$

Proof. See section A.10 □

The following consequence of Proposition 2.3.13 will prove to be very useful in the sequel.

Proposition 2.3.14. *We have*

$$e_3(r\mathfrak{q}) = r^5 \left\{ \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} - \frac{3}{2} \rho \not\partial_2^* \not\partial_1^* \underline{\kappa} - \frac{3}{2} \underline{\kappa} \rho \not\partial_2^* \zeta - \frac{3}{2} \kappa \rho \underline{\alpha} + \frac{3}{4} (2\rho^2 - \kappa \underline{\kappa} \rho) \underline{\vartheta} \right\} + \text{Err}[e_3(r\mathfrak{q})], \quad (2.3.13)$$

where the error term $\text{Err}[e_3(r\mathfrak{q})]$ is given schematically by

$$\text{Err}[e_3(r\mathfrak{q})] = r\Gamma_b \mathfrak{q} + r^5 \mathfrak{d}^{\leq 1} (e_3 \eta \cdot \beta) + r^3 \mathfrak{d}^{\leq 2} (\Gamma_b \cdot \Gamma_g). \quad (2.3.14)$$

Proof. See section A.11. □

We deduce from Proposition 2.3.14 the following nonlinear version of the Teukolsky-Starobinski identity.

Proposition 2.3.15. *The following identity holds true in $^{(int)}\mathcal{M}$,*

$$e_3(r^2 e_3(r\mathfrak{q})) + 2\underline{\omega} r^2 e_3(r\mathfrak{q}) = r^7 \left\{ \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \underline{\alpha} + \frac{3}{2} \rho (\underline{\kappa} e_4 - \kappa e_3) \underline{\alpha} \right\} + \text{Err}[TS], \quad (2.3.15)$$

where the error term $\text{Err}[TS]$ is given schematically by

$$\begin{aligned} \text{Err}[TS] &= r^4 (\not\partial \Gamma_b + r\Gamma_b \cdot \Gamma_b) \cdot \underline{\alpha} + r^2 (\Gamma_b e_3(r\mathfrak{q}) + (\mathfrak{d}^{\leq 1} \Gamma_b) r\mathfrak{q}) \\ &+ r^7 \mathfrak{d}^{\leq 2} (e_3 \eta \cdot \beta) + r^5 \mathfrak{d}^{\leq 3} (\Gamma_b \cdot \Gamma_g). \end{aligned}$$

Proof. See section A.12. □

2.4 Invariant wave equations

In this section, we write wave equations for the invariant quantities α , $\underline{\alpha}$ and \mathfrak{q} .

2.4.1 Preliminaries

Lemma 2.4.1. *With respect to a general S -foliation we have, for a reduced scalar $\psi \in \mathfrak{s}_0$,*

$$\begin{aligned} \square_{\mathbf{g}}\psi &= -\frac{1}{2}(e_3e_4 + e_4e_3)\psi + \underline{\Delta}\psi + \left(\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi + \left(\omega - \frac{1}{2}\kappa\right)e_3\psi \\ &\quad + (\eta + \underline{\eta})e_\theta\psi. \end{aligned} \quad (2.4.1)$$

Also,

$$\begin{aligned} \square_{\mathbf{g}}\psi &= -e_3e_4\psi + \underline{\Delta}\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi - \frac{1}{2}\kappa e_3\psi + 2\underline{\eta}e_\theta\psi, \\ \square_{\mathbf{g}}\psi &= -e_4e_3\psi + \underline{\Delta}\psi + \left(2\omega - \frac{1}{2}\kappa\right)e_3\psi - \frac{1}{2}\underline{\kappa}e_4\psi + 2\underline{\eta}e_\theta\psi. \end{aligned}$$

Proof. We calculate, in spacetime,

$$\square_{\mathbf{g}}\psi = \mathbf{g}^{34}\mathbf{D}_3\mathbf{D}_4\psi + \mathbf{g}^{43}\mathbf{D}_4\mathbf{D}_3\psi + \delta^{AB}\mathbf{D}_A\mathbf{D}_B\psi = -\frac{1}{2}(\mathbf{D}_3\mathbf{D}_4 + \mathbf{D}_4\mathbf{D}_3)\psi + \mathbf{g}^{AB}\mathbf{D}_A\mathbf{D}_B\psi.$$

Now,

$$\begin{aligned} \delta^{AB}\mathbf{D}_A\mathbf{D}_B\psi &= \underline{\Delta}\psi - \frac{1}{2}{}^{(1+3)}\text{tr}\chi e_3\psi - \frac{1}{2}{}^{(1+3)}\text{tr}\underline{\chi} e_4\psi, \\ \mathbf{D}_3\mathbf{D}_4\psi &= e_3e_4\psi - 2\underline{\omega}e_4\psi - 2\underline{\eta}e_\theta\psi, \\ \mathbf{D}_4\mathbf{D}_3\psi &= e_4e_3\psi - 2\omega e_3\psi - 2\underline{\eta}e_\theta\psi. \end{aligned}$$

Hence,

$$\begin{aligned} \square_{\mathbf{g}}\psi &= -\frac{1}{2}(e_3e_4 + e_4e_3)\psi + \underline{\Delta}\psi - \frac{1}{2}{}^{(1+3)}\text{tr}\chi e_3\psi - \frac{1}{2}{}^{(1+3)}\text{tr}\underline{\chi} e_4\psi \\ &\quad + \underline{\omega}e_4\psi + \underline{\eta}e_\theta\psi + \omega e_3\psi + \underline{\eta}e_\theta\psi \\ &= -\frac{1}{2}(e_3e_4 + e_4e_3)\psi + \underline{\Delta}\psi + \left(\underline{\omega} - \frac{1}{2}{}^{(1+3)}\text{tr}\underline{\chi}\right)e_4\psi + \left(\omega - \frac{1}{2}{}^{(1+3)}\text{tr}\chi\right)e_3\psi \\ &\quad + (\eta + \underline{\eta})e_\theta\psi. \end{aligned}$$

Since,

$$\frac{1}{2}e_4e_3\psi = \frac{1}{2}e_3e_4\psi + \omega e_3\psi - \underline{\omega}e_4\psi + (\underline{\eta} - \eta)e_\theta\psi$$

we also have,

$$\square_{\mathbf{g}}\psi = -e_3e_4\psi + \underline{\Delta}\psi + \left(2\underline{\omega} - \frac{1}{2}{}^{(1+3)}\text{tr}\underline{\chi}\right)e_4\psi - \frac{1}{2}{}^{(1+3)}\text{tr}\chi e_3\psi + 2\underline{\eta}e_\theta\psi.$$

Since $\kappa = {}^{(1+3)}\text{tr}\chi$, $\underline{\kappa} = {}^{(1+3)}\text{tr}\underline{\chi}$, this concludes the proof of the lemma. \square

Definition 2.4.2. Given a reduced k -scalar $\psi \in \mathfrak{s}_k$ we define,

$$\begin{aligned} \square_k \psi &= -\frac{1}{2}(e_3 e_4 + e_4 e_3) \psi + \underline{\Delta}_k \psi + (\underline{\omega} - \frac{1}{2} \text{tr} \underline{\chi}) e_4 \psi + (\omega - \frac{1}{2} \text{tr} \chi) e_3 \psi \\ &\quad - (\eta + \underline{\eta}) e_\theta \psi. \end{aligned} \quad (2.4.2)$$

Equivalently, we have

$$\begin{aligned} \square_k \psi &= -e_3 e_4 \psi + \underline{\Delta}_k \psi + \left(2\underline{\omega} - \frac{1}{2} \underline{\kappa}\right) e_4 \psi - \frac{1}{2} \kappa e_3 \psi + 2\underline{\eta} e_\theta \psi, \\ \square_k \psi &= -e_4 e_3 \psi + \underline{\Delta}_k \psi + \left(2\omega - \frac{1}{2} \kappa\right) e_3 \psi - \frac{1}{2} \underline{\kappa} e_4 \psi + 2\underline{\eta} e_\theta \psi. \end{aligned}$$

Remark 2.4.3. Not that the terms $\eta e_\theta \psi, \underline{\eta} e_\theta \psi$ have to be interpreted as in Remark 2.1.23, i.e.

$$\eta e_\theta \psi = \frac{1}{2} \eta (\not\Delta_k \psi - \not\Delta_{k+1}^* \psi).$$

The term $\eta \not\Delta_k \psi$ is the reduced form of a tensor product of ${}^{(1+3)}\eta$ with $\mathcal{D}_k^{(1+3)} \psi$ while $\eta \not\Delta_{k+1}^* \psi$ is the reduced form of a contraction between the 1 form ${}^{(1+3)}\eta$ and $k+1$ tensor $\mathcal{D}_{k+1}^* {}^{(1+3)} \psi$.

Remark 2.4.4. Recall that (recall Definition 2.1.20),

$$\underline{\Delta}_k f := e_\theta(e_\theta f) + e_\theta(\Phi) e_\theta f - k^2 (e_\theta(\Phi))^2 f.$$

Thus, for a $\psi \in \mathfrak{s}_k$, we have,

$$\underline{\Delta}_k \psi = \Delta \psi - k^2 (e_\theta(\Phi))^2 \psi.$$

Spacetime interpretation of Definition 2.4.2

The linearized equation verified by our main quantity \mathfrak{q} , which will be derived in the next subsection, has the form,

$$\square_2 \psi + V \psi = 0. \quad (2.4.3)$$

with V a scalar potential. In what follows we give simple spacetime interpretation of the equation.

Given a mixed spacetime tensor in $\mathbf{T}^k \mathbf{M} \otimes \mathbf{T}_S^l \mathbf{M}$ of the form $U_{\mu_1 \dots \mu_k, A_1 \dots A_l}$ where e_μ is an orthonormal frame on \mathcal{M} with $(e_A)_{A=1,2}$ tangent to S . We define,

$$\begin{aligned} \dot{\mathbf{D}}_\mu U_{\nu_1 \dots \nu_k, A_1 \dots A_l} &= e_\mu U_{\nu_1 \dots \nu_k, A_1 \dots A_l} - U_{\mathbf{D}_\mu \nu_1 \dots \nu_k, A_1 \dots A_l} - \dots - U_{\nu_1 \dots \nu_k, \mathbf{D}_\mu A_1 \dots A_l} \\ &\quad - U_{\nu_1 \dots \nu_k, \dot{\mathbf{D}}_\mu A_1 \dots A_l} - U_{\nu_1 \dots \nu_k, A_1 \dots \dot{\mathbf{D}}_\mu A_l} \end{aligned}$$

with $\dot{\mathbf{D}}_\mu A$ denoting the projection of $\mathbf{D}_{e_\mu} e_A$ on S . One can easily check the commutator formulae,

$$\begin{aligned} (\dot{\mathbf{D}}_\mu \dot{\mathbf{D}}_\nu - \dot{\mathbf{D}}_\nu \dot{\mathbf{D}}_\mu) \Psi_A &= \mathbf{R}_A{}^B{}_{\mu\nu} \Psi_B, \\ (\dot{\mathbf{D}}_\mu \dot{\mathbf{D}}_\nu - \dot{\mathbf{D}}_\nu \dot{\mathbf{D}}_\mu) \Psi_{\lambda A} &= \mathbf{R}_\lambda{}^\sigma{}_{\mu\nu} \Psi_{\sigma A} + \mathbf{R}_A{}^B{}_{\mu\nu} \Psi_{\lambda B}. \end{aligned}$$

Define,

$$\dot{\square}_{\mathbf{g}} \Psi := \mathbf{g}^{\mu\nu} \dot{\mathbf{D}}_\mu \dot{\mathbf{D}}_\nu \Psi.$$

Consider the following Lagrangian for $\Psi = \Psi_{AB} \in \mathcal{S}_2$.

$$\mathcal{L}[\Psi] = \not{g}^{A_1 B_1} \not{g}^{A_2 B_2} \left(\mathbf{g}^{\mu\nu} \dot{\mathbf{D}}_\mu \Psi_{A_1 A_2} \dot{\mathbf{D}}_\nu \Psi_{B_1 B_2} + V \Psi_{A_1 A_2} \Psi_{B_1 B_2} \right).$$

Proposition 2.4.5. *The Euler- Lagrange equations for the Lagrangian $\mathcal{L}[\Psi]$ above are given by:*

$$\dot{\square} \Psi = V \Psi \tag{2.4.4}$$

and its reduced for $\psi = \Psi_{\theta\theta}$ is precisely (2.4.3).

Proof. Straightforward verification. □

2.4.2 Wave equations for α , $\underline{\alpha}$, and \mathfrak{q}

We start with the wave equations for α and $\underline{\alpha}$, which are derived in a general null frame.

Proposition 2.4.6. *The following identities hold true.*

1. *The invariant quantity $\alpha \in \mathfrak{s}_2$ verifies the Teukolsky wave equation,*

$$\begin{aligned} \square_2 \alpha &= -4\underline{\omega} e_4(\alpha) + (4\omega + 2\kappa) e_3(\alpha) + V\alpha + \text{Err}[\square_{\mathbf{g}} \alpha], \\ V &= -4\rho - 4e_4(\underline{\omega}) - 8\omega \underline{\omega} + 2\omega \underline{\kappa} - 10\kappa \underline{\omega} + \frac{1}{2} \kappa \underline{\kappa}, \end{aligned} \tag{2.4.5}$$

where $\text{Err}[\square_{\mathbf{g}} \alpha]$ is given schematically by

$$\text{Err}(\square_{\mathbf{g}} \alpha) = \Gamma_g e_3(\alpha) + r^{-1} \mathfrak{d}^{\leq 1} \left((\eta, \Gamma_g)(\alpha, \beta) \right) + \xi(e_3(\beta), r^{-1} \mathfrak{d} \check{\rho}) + \text{l.o.t.}$$

where *l.o.t.* denote terms which are quadratic and enjoy better decay properties or are higher order and decay at least as good.

2. The invariant quantity $\underline{\alpha} \in \mathfrak{s}_2$ verifies the Teukolsky wave equation,

$$\begin{aligned} \square_2 \underline{\alpha} &= -4\omega e_3(\underline{\alpha}) + (4\underline{\omega} + 2\underline{\kappa})e_4(\underline{\alpha}) + \underline{V} \underline{\alpha} + \text{Err}[\square_{\mathbf{g}} \underline{\alpha}], \\ \underline{V} &= -4\rho - 4e_3(\omega) - 8\omega \underline{\omega} + 2\underline{\omega} \underline{\kappa} - 10\underline{\kappa} \omega + \frac{1}{2} \underline{\kappa} \underline{\kappa}, \end{aligned} \quad (2.4.6)$$

where

$$\text{Err}[\square_{\mathbf{g}} \underline{\alpha}] = r^{-1} \mathfrak{d}(\Gamma_b \underline{\alpha}) + \mathfrak{d}(\Gamma_b \underline{\beta}) + l.o.t.$$

Proof. See appendix A.13 □

We may now state the wave equation satisfied by \mathbf{q} .

Theorem 2.4.7. *The invariant scalar quantity \mathbf{q} defined in (2.3.10) verifies the equation,*

$$\square_2 \mathbf{q} + \underline{\kappa} \underline{\kappa} \mathbf{q} = \text{Err}[\square_2 \mathbf{q}] \quad (2.4.7)$$

where $\text{Err}[\square_2 \mathbf{q}]$ is $O(\epsilon^2)$.

If \mathbf{q} is defined relative to a null frame satisfying, in addition to the assumptions of section 2.3.2, that $\eta \in \Gamma_g$ and $\xi = 0$ for $r \geq 4m_0$, the error term is then given schematically by

$$\text{Err}[\square_2 \mathbf{q}] = r^2 \mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta)) + e_3 \left(r^3 \mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta)) \right) + \mathfrak{d}^{\leq 1}(\Gamma_g \cdot \mathbf{q}) + l.o.t. \quad (2.4.8)$$

Proof. See appendix A.14. □

Remark 2.4.8. *Note that the main frame used in this paper is an outgoing geodesic null frame in $r \geq 4m_0$ so that $\xi = 0$, but unfortunately, as it turns out, $\eta \in \Gamma_b$. This would not allow us to control the error term appearing in (2.4.7). To overcome this problem, we are forced to define \mathbf{q} relative to a different frame where $\xi = 0$ still holds for $r \geq 4m_0$ and for which we have in addition $\eta \in \Gamma_g$, see Proposition 3.5.5 for the existence of such a frame. See also the discussion at the beginning of section 3.4.6.*

The remark above leads us to the following.

Remark 2.4.9. *The quantity \mathbf{q} we will be working with for the rest of the paper is defined, according to equation (2.3.10), relative to the global frame of Proposition 3.5.5 for which $\eta \in \Gamma_g$. It is only in such a frame that \mathbf{q} verifies the correct decay estimates.*

Chapter 3

MAIN THEOREM

3.1 General covariant modulated admissible spacetimes

Note that all definitions below are consistent with the framework of \mathbf{Z} -invariant polarized spacetimes.

3.1.1 Initial data layer

Recall that $m_0 > 0$ is given as the mass of the Schwarzschild solution to which the initial data is ϵ_0 close. Let $\delta_{\mathcal{H}} > 0$ be a sufficiently small constant which will be specified later.

Definition 3.1.1 (Initial data layer). *We consider a spacetime region (\mathcal{L}_0, g) , sketched below in figure 3.1, where*

- *The metric g is a reduced metric from a Lorentzian spacetime metric \mathbf{g} close to Schwarzschild in a suitable topology¹.*
- $\mathcal{L}_0 = {}^{(ext)}\mathcal{L}_0 \cup {}^{(int)}\mathcal{L}_0$.
- *The intersection ${}^{(ext)}\mathcal{L}_0 \cap {}^{(int)}\mathcal{L}_0$ is non trivial.*

¹This topology will be specified in our initial data layer assumptions, see (3.3.5) as well as section 3.2.4.

Furthermore, our initial data layer (\mathcal{L}_0, g) satisfies

1. **Boundaries.** The future and past boundaries of \mathcal{L}_0 are given by

$$\begin{aligned}\partial^+ \mathcal{L}_0 &= \mathcal{A}_0 \cup \underline{\mathcal{C}}_{(2, \mathcal{L}_0)} \cup \mathcal{C}_{(2, \mathcal{L}_0)}, \\ \partial^- \mathcal{L}_0 &= \mathcal{C}_{(0, \mathcal{L}_0)} \cup \underline{\mathcal{C}}_{(0, \mathcal{L}_0)},\end{aligned}$$

where

- (a) The past outgoing null boundary of the far region ${}^{(ext)}\mathcal{L}_0$ is denoted by $\mathcal{C}_{(0, \mathcal{L}_0)}$.
- (b) The past incoming null boundary of the near region ${}^{(int)}\mathcal{L}_0$ is denoted by $\underline{\mathcal{C}}_{(0, \mathcal{L}_0)}$.
- (c) ${}^{(ext)}\mathcal{L}_0$ is unbounded in the future outgoing null directions.
- (d) The future outgoing null boundary of the far region ${}^{(ext)}\mathcal{L}_0$ is denoted by $\mathcal{C}_{(2, \mathcal{L}_0)}$.
- (e) The future incoming null boundary of the near region ${}^{(int)}\mathcal{L}_0$ is denoted by $\underline{\mathcal{C}}_{(2, \mathcal{L}_0)}$.
- (f) The future spacelike boundary of the near region ${}^{(int)}\mathcal{L}_0$ is denoted by \mathcal{A}_0 .

2. **Foliations of \mathcal{L}_0 and adapted null frames.** The spacetime $\mathcal{L}_0 = {}^{(ext)}\mathcal{L}_0 \cup {}^{(int)}\mathcal{L}_0$ is foliated as follows

- (a) The far region ${}^{(ext)}\mathcal{L}_0$ is foliated by two functions $(u_{\mathcal{L}_0}, {}^{(ext)}s_{\mathcal{L}_0})$ such that
 - $u_{\mathcal{L}_0}$ is an outgoing optical function on ${}^{(ext)}\mathcal{L}_0$ whose leaves are denoted by $\mathcal{C}_{(u_{\mathcal{L}_0}, \mathcal{L}_0)}$.
 - ${}^{(ext)}s_{\mathcal{L}_0}$ is an affine parameter along the level hypersurfaces of $u_{\mathcal{L}_0}$, i.e.

$${}^{(ext)}L_0({}^{(ext)}s_{\mathcal{L}_0}) = 1 \text{ where } {}^{(ext)}L_0 := -g^{ab}\partial_b(u_{\mathcal{L}_0})\partial_a.$$

- We denote by $({}^{(ext)}(e_0)_3, {}^{(ext)}(e_0)_4, {}^{(ext)}(e_0)_\theta)$ the null frame adapted to the outgoing geodesic foliation $(u_{\mathcal{L}_0}, {}^{(ext)}s_{\mathcal{L}_0})$ on ${}^{(ext)}\mathcal{L}_0$.
- Let ${}^{(ext)}r_{\mathcal{L}_0}$ denote the area radius of the 2-spheres $S(u_{\mathcal{L}_0}, {}^{(ext)}s_{\mathcal{L}_0})$ of this foliation.
- The outgoing future null boundary $\mathcal{C}_{(2, \mathcal{L}_0)}$ corresponds precisely to $u_{\mathcal{L}_0} = 1$ and the outgoing past null boundary $\mathcal{C}_{(0, \mathcal{L}_0)}$ corresponds to $u_{\mathcal{L}_0} = -1$.
- The foliation by $u_{\mathcal{L}_0}$ of ${}^{(ext)}\mathcal{L}_0$ terminates at the time like boundary

$$\left\{ {}^{(ext)}r_{\mathcal{L}_0} = 2m_0 \left(1 + \frac{\delta_{\mathcal{H}}}{4} \right) \right\}.$$

- (b) The near region ${}^{(int)}\mathcal{L}_0$ is foliated by two functions $(\underline{u}_{\mathcal{L}_0}, {}^{(int)}s_{\mathcal{L}_0})$ such that

- $\underline{u}_{\mathcal{L}_0}$ is an ingoing optical function on ${}^{(int)}\mathcal{L}_0$ whose leaves are denoted by $\mathcal{C}_{(\underline{u}_{\mathcal{L}_0}, \mathcal{L}_0)}$.
- ${}^{(int)}s_{\mathcal{L}_0}$ is an affine parameter along the level hypersurfaces of $\underline{u}_{\mathcal{L}_0}$, i.e.

$${}^{(int)}\underline{L}_0({}^{(int)}s_{\mathcal{L}_0}) = -1 \text{ where } {}^{(ext)}\underline{L}_0 := -g^{ab}\partial_b(\underline{u}_{\mathcal{L}_0})\partial_a.$$

- We denote by $({}^{(int)}(e_0)_3, {}^{(int)}(e_0)_4, {}^{(int)}(e_0)_\theta)$ the null frame adapted to the outgoing geodesic foliation $(u_{\mathcal{L}_0}, {}^{(int)}s_{\mathcal{L}_0})$ on ${}^{(int)}\mathcal{L}_0$.
- Let ${}^{(int)}r_{\mathcal{L}_0}$ denote the area radius of the 2-spheres $S(\underline{u}_{\mathcal{L}_0}, {}^{(int)}s_{\mathcal{L}_0})$ of this foliation.
- The $(\underline{u}_{\mathcal{L}_0}, {}^{(int)}s)$ foliation is initialized on ${}^{(ext)}r_{\mathcal{L}_0} = 2m_0(1 + \frac{\delta_{\mathcal{H}}}{2})$ as it will be made precise below.
- The foliation by $\underline{u}_{\mathcal{L}_0}$, of ${}^{(int)}\mathcal{L}_0$ terminates at the space like boundary

$$\mathcal{A}_0 = \{ {}^{(int)}r_{\mathcal{L}_0} = 2m_0(1 - 2\delta_{\mathcal{H}}) \}.$$

where m_0 and $\delta_{\mathcal{H}}$ have been defined above.

- The ingoing future null boundary $\underline{\mathcal{C}}_{(2, \mathcal{L}_0)}$ corresponds precisely to $\underline{u}_{\mathcal{L}_0} = 2$ and the ingoing past null boundary $\underline{\mathcal{C}}_{(0, \mathcal{L}_0)}$ corresponds to $\underline{u}_{\mathcal{L}_0} = 0$.
- The foliation by $\underline{u}_{\mathcal{L}_0}$ of ${}^{(int)}\mathcal{L}_0$ terminates at the time like boundary

$$\{ {}^{(int)}r_{\mathcal{L}_0} = 2m_0(1 + 2\delta_{\mathcal{H}}) \}.$$

3. Initializations of the $(\underline{u}_{\mathcal{L}_0}, {}^{(int)}s)$ foliation.

The $(\underline{u}_{\mathcal{L}_0}, {}^{(int)}s_{\mathcal{L}_0})$ foliation is initialized on ${}^{(ext)}r_{\mathcal{L}_0} = 2m_0(1 + \delta_{\mathcal{H}})$ by setting,


$$\underline{u}_{\mathcal{L}_0} = u_{\mathcal{L}_0}, \quad {}^{(int)}s_{\mathcal{L}_0} = {}^{(ext)}s_{\mathcal{L}_0}$$

and, with $\lambda_0 = {}^{(ext)}\lambda_0 = 1 - \frac{2m_0}{{}^{(ext)}r_{\mathcal{L}_0}}$,

$${}^{(int)}(e_0)_4 = \lambda_0 {}^{(ext)}(e_0)_4, \quad {}^{(int)}(e_0)_3 = \lambda_0^{-1} {}^{(ext)}(e_0)_3, \quad {}^{(int)}(e_0)_\theta = {}^{(ext)}(e_0)_\theta.$$

- ### 4. Coordinates system on ${}^{(ext)}\mathcal{L}_0({}^{(ext)}r_{\mathcal{L}_0} \geq 4m_0)$.
- In ${}^{(ext)}\mathcal{L}_0({}^{(ext)}r_{\mathcal{L}_0} \geq 4m_0)$, there exists adapted coordinates $(u_{\mathcal{L}_0}, {}^{(ext)}s_{\mathcal{L}_0}, \theta_{\mathcal{L}_0}, \varphi)$ with $b = 0$, see Proposition 2.2.20, such that the spacetime metric \mathbf{g} takes the form,

$$\mathbf{g} = -2du_{\mathcal{L}_0} d({}^{(ext)}s_{\mathcal{L}_0}) + \underline{\Omega}_{\mathcal{L}_0}(du_{\mathcal{L}_0})^2 + \gamma_{\mathcal{L}_0} \left(d\theta_{\mathcal{L}_0} - \frac{1}{2}\underline{b}_{\mathcal{L}_0} du_{\mathcal{L}_0} \right)^2 + e^{2\Phi} d\varphi^2. \quad (3.1.1)$$

Figure 3.1: The initial data layer \mathcal{L}_0

3.1.2 Main Definition

Recall that $m_0 > 0$ is given as the mass of the Schwarzschild solution to which the initial data is ϵ_0 close, and that $\delta_{\mathcal{H}} > 0$ is a sufficiently small constant which will be specified later.

Definition 3.1.2 (GCM-admissible spacetime). *We consider a spacetime (\mathcal{M}, g) , sketched below in figure 3.2, where*

- *The metric g is a reduced metric from a Lorentzian spacetime metric \mathbf{g} close to Schwarzschild in a suitable topology².*
- $\mathcal{M} = {}^{(ext)}\mathcal{M} \cup {}^{(int)}\mathcal{M}$
- $\mathcal{T} = {}^{(ext)}\mathcal{M} \cap {}^{(int)}\mathcal{M}$ *is a time-like hyper-surface.*

(\mathcal{M}, g) *is called a general covariant modulated admissible (or shortly GCM-admissible) spacetime if it is defined as follows*

²This topology will be specified in our bootstrap assumptions, see (3.3.6) as well as section 3.2.

1. **Boundaries.** *The future and past boundaries of \mathcal{M} are given by*

$$\begin{aligned}\partial^+ \mathcal{M} &= \mathcal{A} \cup \underline{\mathcal{C}}_* \cup \mathcal{C}_* \cup \Sigma_*, \\ \partial^- \mathcal{M} &= \mathcal{C}_1 \cup \underline{\mathcal{C}}_1,\end{aligned}$$

where

- (a) *The past boundary $\mathcal{C}_1 \cup \underline{\mathcal{C}}_1$ is included in the initial data layer \mathcal{L}_0 , defined in section 3.1.1, in which the metric on \mathcal{M} is specified to be a small perturbation of the Schwarzschild data.*
- (b) *The future spacelike boundary of the far region ${}^{(ext)}\mathcal{M}$ is denoted by Σ_* .*
- (c) *The future outgoing null boundary of the far region ${}^{(ext)}\mathcal{M}$ is denoted by \mathcal{C}_* .*
- (d) *The future incoming null boundary of the near region ${}^{(int)}\mathcal{M}$ is denoted by $\underline{\mathcal{C}}_*$.*
- (e) *The future spacelike boundary of the near region ${}^{(int)}\mathcal{M}$ is denoted by \mathcal{A} .*
- (f) *The time-like boundary \mathcal{T} , separating ${}^{(ext)}\mathcal{M}$ from ${}^{(int)}\mathcal{M}$, starts at $\underline{\mathcal{C}}_1 \cap \mathcal{C}_1$ and terminates at $\underline{\mathcal{C}}_* \cap \mathcal{C}_*$.*

2. **Foliations of \mathcal{M} and adapted null frames.** *The spacetime $\mathcal{M} = {}^{(ext)}\mathcal{M} \cup {}^{(int)}\mathcal{M}$ is foliated as follows*

- (a) *The far region ${}^{(ext)}\mathcal{M}$ is foliated by two functions $(u, {}^{(ext)}s)$ such that*
 - *u is an outgoing optical function on ${}^{(ext)}\mathcal{M}$, initialized on Σ_* , whose leaves are denoted by $\mathcal{C}(u)$.*
 - *${}^{(ext)}s$ is an affine parameter along the level hypersurfaces of u , i.e.*

$$L({}^{(ext)}s) = 1 \text{ where } L := -g^{ab} \partial_b u \partial_a.$$

- *The $(u, {}^{(ext)}s)$ foliation is initialized on Σ_* as it will be made precise below.*
- *We denote by $({}^{(ext)}e_3, {}^{(ext)}e_4, {}^{(ext)}e_\theta)$ the null frame adapted to the outgoing geodesic foliation $(u, {}^{(ext)}s)$ on ${}^{(ext)}\mathcal{M}$ where ${}^{(ext)}e_4 = L$.*
- *Let ${}^{(ext)}r$ and ${}^{(ext)}m$ respectively the area radius and the Hawking mass of the 2-spheres $S(u, {}^{(ext)}s)$ of this foliation.*
- *The outgoing future null boundary \mathcal{C}_* corresponds precisely to $u = u_*$ and the outgoing past null boundary \mathcal{C}_1 corresponds to $u = 1$.*
- *The foliation by u of ${}^{(ext)}\mathcal{M}$ terminates at the time like boundary*

$$\mathcal{T} = \{ {}^{(ext)}r = r_{\mathcal{T}} \}$$

where $r_{\mathcal{T}}$ satisfies³

$$2m_0 \left(1 + \frac{\delta_{\mathcal{H}}}{2} \right) \leq r_{\mathcal{T}} \leq 2m_0 \left(1 + \frac{3\delta_{\mathcal{H}}}{2} \right).$$

³A specific choice of $r_{\mathcal{T}}$ will be made in section 3.8.9, see (3.8.7), in the context of a Lebesgue point argument needed to recover the top order derivatives.

(b) The near region ${}^{(int)}\mathcal{M}$ is foliated by two functions $(\underline{u}, {}^{(int)}s)$ such that

- \underline{u} is an ingoing optical function on ${}^{(int)}\mathcal{M}$, initialized on \mathcal{T} , whose leaves are denoted by $\underline{\mathcal{C}}(\underline{u})$.
- ${}^{(int)}s$ is an affine parameter along the level hypersurfaces of \underline{u} , i.e.

$$\underline{L}({}^{(int)}s) = -1 \text{ where } \underline{L} := -g^{ab}\partial_b\underline{u}\partial_a.$$

- The $(\underline{u}, {}^{(int)}s)$ foliation is initialized on \mathcal{T} as it will be made precise below.
- We denote by $({}^{(int)}e_3, {}^{(int)}e_4, {}^{(int)}e_\theta)$ the null frame adapted to the outgoing geodesic foliation $(u, {}^{(int)}s)$ on ${}^{(int)}\mathcal{M}$ where ${}^{(int)}e_3 = \underline{L}$.
- Let ${}^{(int)}r$ and ${}^{(int)}m$ respectively the area radius and the Hawking mass of the 2-spheres $S(\underline{u}, {}^{(int)}s)$ of this foliation.
- The foliation by \underline{u} of ${}^{(int)}\mathcal{M}$ terminates at the space like boundary

$$\mathcal{A} = \{ {}^{(int)}r = 2m_0(1 - \delta_{\mathcal{H}}) \}$$

where m_0 and $\delta_{\mathcal{H}}$ have been defined above.

- The ingoing future null boundary $\underline{\mathcal{C}}_*$ corresponds precisely to $\underline{u} = u_*$ and the ingoing past null boundary $\underline{\mathcal{C}}_1$ corresponds to $\underline{u} = 1$.

3. **GCM foliation of Σ_* .** The $(u, {}^{(ext)}s)$ -foliation of ${}^{(ext)}\mathcal{M}$ restricted to the spacelike hypersurface Σ_* has the following properties

(a) There exists a constant c_{Σ_*} such that

$$\Sigma_* := \{ u + {}^{(ext)}r = c_{\Sigma_*} \}.$$

(b) We have⁴

$$r \gg u_*^4 \text{ on } \Sigma_*. \quad (3.1.2)$$

(c) ${}^{(ext)}s$ satisfies⁵

$${}^{(ext)}s = {}^{(ext)}r \text{ on } \Sigma_*.$$

(d) We say that Σ_* is a general covariant modulated hypersurface⁶ (or shortly GCM hypersurface) if relative to the above defined null frame of ${}^{(ext)}\mathcal{M}$, the following

⁴See (3.3.4) for the precise condition.

⁵Recall that ${}^{(ext)}s$ satisfies on ${}^{(ext)}\mathcal{M}$ the transport equation $L({}^{(ext)}s) = 1$ and thus needs to be initialized on a hypersurface transversal to L , chosen here to be Σ_* .

⁶More generally, a GCM hypersurface is one with the property that we can specify, using the full covariance of the Einstein equations, a number of vanishing conditions (equal to the number of degrees of freedom of the diffeomorphism group) for well-chosen components of $\bar{\Gamma}$.

conditions hold⁷ along Σ_*

$$\begin{aligned} \kappa &= \frac{2}{r}, \quad \not{d}_2^* \not{d}_1^* \kappa = 0, \quad \not{d}_2^* \not{d}_1^* \mu = 0, \\ \int_S \eta e^\Phi &= 0, \quad \int_S \underline{\xi} e^\Phi = 0, \quad a|_{SP} = 1, \end{aligned} \quad (3.1.3)$$

where a is the unique scalar function such that $\nu = e_3 + ae_4$ is tangent to Σ_* , and SP denotes the south poles of the spheres on Σ_* . Moreover we also assume

$$\int_{S_*} \beta e^\Phi = 0, \quad \int_{S_*} e_\theta(\underline{\kappa}) e^\Phi = 0, \quad \text{with } S_* := \Sigma_* \cap \mathcal{C}_*. \quad (3.1.4)$$

Note that the role of the GCM foliation of Σ_* is to initialize the $(u, {}^{(ext)}s)$ -foliation of ${}^{(ext)}\mathcal{M}$.

- (e) In view of the definition of ν and ς , we have $\nu(u) = e_3(u) + ae_4(u) = 2/\varsigma$. ν being tangent to Σ_* , u is thus transported along Σ_* , and hence defined up to a constant. To calibrate u on Σ_* , we fix the value $u = 1$ as follows

$$S_1 = \Sigma_* \cap \{u = 1\} \text{ is such that } S_1 \cap \mathcal{C}_{(1, \mathcal{L}_0)} \cap SP \neq \emptyset, \quad (3.1.5)$$

i.e. S_1 is the unique sphere of Σ_* such that its south pole intersects the south pole of one of the sphere of the outgoing null cone $\mathcal{C}_{(1, \mathcal{L}_0)}$ of the initial data layer.

4. **Initialization the $(\underline{u}, {}^{(int)}s)$ -foliation on \mathcal{T} .** The $(\underline{u}, {}^{(int)}s)$ foliation is initialized on \mathcal{T} such that,

$$\underline{u} = u, \quad {}^{(int)}s = {}^{(ext)}s$$

In particular, the 2-spheres $S(u, {}^{(int)}s)$ coincide on \mathcal{T} with $S(u, {}^{(ext)}s)$ and ${}^{(int)}r = {}^{(ext)}r$. Moreover, the null frame $({}^{(int)}e_3, {}^{(int)}e_4, {}^{(int)}e_\theta)$ is defined on \mathcal{T} by the following renormalization,

$${}^{(int)}e_4 = \lambda {}^{(ext)}e_4, \quad {}^{(int)}e_3 = \lambda^{-1} {}^{(ext)}e_3, \quad {}^{(int)}e_\theta = {}^{(ext)}e_\theta \quad \text{on } \mathcal{T}$$


where

$$\lambda = {}^{(ext)}\lambda = 1 - \frac{2 {}^{(ext)}m}{{}^{(ext)}r}.$$

Remark 3.1.3. In Schwarzschild, $u = t - r_*$, $\underline{u} = t + r_*$, with $\frac{dr_*}{dr} = \Upsilon^{-1}$, and

$$\begin{aligned} {}^{(ext)}e_4 &= \Upsilon^{-1} \partial_t + \partial_r, & {}^{(ext)}e_3 &= \partial_t - \Upsilon \partial_r, \\ {}^{(int)}e_4 &= \partial_t + \Upsilon \partial_r, & {}^{(int)}e_3 &= \Upsilon^{-1} \partial_t - \partial_r. \end{aligned}$$

⁷The existence of such hypersurfaces is an essential part of our construction.

Figure 3.2: The GCM admissible space-time \mathcal{M}

3.1.3 Renormalized curvature components and Ricci coefficients

For convenience, we introduce in this section a notation for renormalized curvature components and Ricci coefficients.

Definition 3.1.4 (Renormalized curvature components and Ricci coefficients in ${}^{(ext)}\mathcal{M}$). We introduce the following notations in ${}^{(ext)}\mathcal{M}$

$${}^{(ext)}\check{R} = \{\alpha, \beta, \check{\rho}, \check{\mu}, \underline{\beta}, \underline{\alpha}\}, \quad {}^{(ext)}\check{\Gamma} = \{\check{\kappa}, \vartheta, \zeta, \eta, \check{\underline{\kappa}}, \underline{\vartheta}, \underline{\omega}, \underline{\xi}\},$$

where, recall,

$$\check{\rho} = \rho - \bar{\rho}, \quad \check{\mu} = \mu - \bar{\mu}, \quad \check{\kappa} = \kappa - \bar{\kappa}, \quad \check{\underline{\kappa}} = \underline{\kappa} - \bar{\underline{\kappa}}, \quad \check{\underline{\omega}} = \underline{\omega} - \bar{\underline{\omega}},$$

and

$$\xi = \omega = 0, \quad \underline{\eta} = -\zeta.$$

Note that all the above quantities are defined with respect to the outgoing geodesic foliation of ${}^{(ext)}\mathcal{M}$ (see section 2.2.4), and that the averages are taken with respect to that corresponding 2-spheres.

Definition 3.1.5 (Renormalized curvature components and Ricci coefficients in $^{(int)}\mathcal{M}$). We introduce the following notations in $^{(int)}\mathcal{M}$

$$\begin{aligned} {}^{(int)}\check{R} &= \left\{ \alpha, \beta, \check{\rho}, \check{\underline{\mu}}, \underline{\beta}, \underline{\alpha} \right\}, \\ {}^{(int)}\check{\Gamma} &= \left\{ \xi, \check{\omega}, \check{\kappa}, \vartheta, \zeta, \underline{\eta}, \check{\underline{\kappa}}, \underline{\vartheta} \right\}, \end{aligned}$$

where we have defined

$$\check{\rho} = \rho - \bar{\rho}, \quad \check{\underline{\mu}} = \underline{\mu} - \bar{\underline{\mu}}, \quad \check{\kappa} = \kappa - \bar{\kappa}, \quad \check{\underline{\kappa}} = \underline{\kappa} - \bar{\underline{\kappa}}, \quad \check{\omega} = \omega - \bar{\omega},$$

and we recall that

$$\underline{\xi} = \underline{\omega} = 0, \quad \eta = \zeta, \quad \bar{\underline{\mu}} - \frac{2m}{r^3} = 0.$$

Note that all the above quantities are defined with respect to the ingoing geodesic foliation of $^{(int)}\mathcal{M}$ (see section 2.2.6), and that the averages are taken with respect to that corresponding 2-spheres.

Remark 3.1.6. In Schwarzschild, we have

$${}^{(ext)}\check{R} = 0, \quad {}^{(int)}\check{R} = 0, \quad {}^{(ext)}\check{\Gamma} = 0, \quad {}^{(int)}\check{\Gamma} = 0.$$

3.2 Main norms

3.2.1 Main norms in ${}^{(ext)}\mathcal{M}$

All quantities appearing in this section are defined relative to the ${}^{(ext)}\mathcal{M}$ frame adapted to the $(u, {}^{(ext)}s)$ foliation. In particular, recall that with respect to this frame, we have

$$\xi = \omega = 0, \quad \underline{\eta} = -\zeta.$$

Recall the definition (2.1.37) of higher order angular derivatives $\check{\partial}^s$ of reduced scalars in \mathfrak{s}_k . We introduce the notations

$$\mathfrak{d} = \{e_3, re_4, \check{\partial}\}.$$

Definition 3.2.1. We introduce the vectorfield \mathbf{T} defined on ${}^{(ext)}\mathcal{M}$ as

$$\mathbf{T} := \frac{1}{2} \left(\left(1 - \frac{2m}{r} \right) e_4 + e_3 \right). \quad (3.2.1)$$

We also introduce the vectorfield \mathbf{N} is defined on ${}^{(ext)}\mathcal{M}$ by

$$\mathbf{N} := \frac{1}{2} \left(\left(1 - \frac{2m}{r} \right) e_4 - e_3 \right). \quad (3.2.2)$$

Remark 3.2.2. *In Schwarzschild, we have*

$$\mathbf{T} = \partial_t, \quad \mathbf{N} = \left(1 - \frac{2m_0}{r}\right) \partial_r$$

in the standard (t, r, θ, φ) coordinates.

We are ready to introduce our norms in ${}^{(ext)}\mathcal{M}$.

L^2 curvature norms in ${}^{(ext)}\mathcal{M}$

Let $\delta_B > 0$ a small constant to be specified later. We introduce the weighted curvature norms,

$$\begin{aligned} \left({}^{(ext)}\mathfrak{R}_0^{\geq 4m_0}[\check{R}]\right)^2 &:= \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq 4m_0)} \left(r^{4+\delta_B} \alpha^2 + r^4 \beta^2\right) \\ &+ \int_{\Sigma_*} \left(r^{4+\delta_B} (\alpha^2 + \beta^2) + r^4 (\check{\rho})^2 + r^2 \underline{\beta}^2 + \underline{\alpha}^2\right) \\ &+ \int_{{}^{(ext)}\mathcal{M}(r \geq 4m_0)} \left(r^{3+\delta_B} (\alpha^2 + \beta^2) + r^{3-\delta_B} (\check{\rho})^2 + r^{1-\delta_B} \underline{\beta}^2 + r^{-1-\delta_B} \underline{\alpha}^2\right), \end{aligned}$$

$$\left({}^{(ext)}\mathfrak{R}_0^{\leq 4m_0}[\check{R}]\right)^2 := \int_{{}^{(ext)}\mathcal{M}(r \leq 4m_0)} \left(1 - \frac{3m}{r}\right)^2 |\check{R}|^2,$$

and

$${}^{(ext)}\mathfrak{R}_0[\check{R}] := {}^{(ext)}\mathfrak{R}_0^{\geq 4m_0}[\check{R}] + {}^{(ext)}\mathfrak{R}_0^{\leq 4m_0}[\check{R}].$$

For any nonzero integer k , we introduce the following higher derivatives norms

$$\left({}^{(ext)}\mathfrak{R}_k[\check{R}]\right)^2 := \left({}^{(ext)}\mathfrak{R}_0[\mathfrak{d}^{\leq k} \check{R}]\right)^2 + \int_{{}^{(ext)}\mathcal{M}(r \leq 4m_0)} \left(|\mathfrak{d}^{\leq k-1} \mathbf{N} \check{R}|^2 + |\mathfrak{d}^{\leq k-1} \check{R}|^2\right).$$

Remark 3.2.3. *Note that the derivative in the \mathbf{N} direction, unlike all other first derivatives of \check{R} , appear in the spacetime integral $\int_{{}^{(ext)}\mathcal{M}(r \leq 4m_0)}$ with top number of derivatives. This reflects the fact the \mathbf{N} -derivatives do not degenerate at $r = 3m$ in the Morawetz estimate.*

L^2 Ricci coefficients norms in ${}^{(ext)}\mathcal{M}$

For any $k \geq 2$, we introduce the following norms

$$\begin{aligned} \left({}^{(ext)}\mathfrak{G}_k^{\geq 4m_0} [\check{\Gamma}] \right)^2 &:= \int_{\Sigma_*} \left[r^2 \left((\mathfrak{d}^{\leq k} \vartheta)^2 + (\mathfrak{d}^{\leq k} \check{\kappa})^2 + (\mathfrak{d}^{\leq k} \zeta)^2 + (\mathfrak{d}^{\leq k} \check{\underline{\kappa}})^2 \right) + (\mathfrak{d}^{\leq k} \underline{\vartheta})^2 \right. \\ &\quad \left. + (\mathfrak{d}^{\leq k} \eta)^2 + (\mathfrak{d}^{\leq k} \check{\omega})^2 + (\mathfrak{d}^{\leq k} \underline{\xi})^2 \right] \\ &\quad + \sup_{\lambda \geq 4m_0} \left(\int_{\{r=\lambda\}} \left[\lambda^2 \left((\mathfrak{d}^{\leq k} \vartheta)^2 + (\mathfrak{d}^{\leq k} \check{\kappa})^2 + (\mathfrak{d}^{\leq k} \zeta)^2 \right) \right. \right. \\ &\quad \left. \left. + \lambda^{2-\delta_B} (\mathfrak{d}^{\leq k} \check{\underline{\kappa}})^2 + (\mathfrak{d}^{\leq k} \underline{\vartheta})^2 + (\mathfrak{d}^{\leq k} \eta)^2 + (\mathfrak{d}^{\leq k} \check{\omega})^2 + \lambda^{-\delta_B} (\mathfrak{d}^{\leq k} \underline{\xi})^2 \right] \right), \end{aligned}$$

$$\left({}^{(ext)}\mathfrak{G}_k^{\leq 4m_0} [\check{\Gamma}] \right)^2 := \int_{{}^{(ext)}\mathcal{M}(\leq 4m_0)} |\mathfrak{d}^{\leq k} (\check{\Gamma})|^2,$$

and

$${}^{(ext)}\mathfrak{G}_k [\check{\Gamma}] := {}^{(ext)}\mathfrak{G}_k^{\leq 4m_0} [\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_k^{\geq 4m_0} [\check{\Gamma}].$$

Decay norms in ${}^{(ext)}\mathcal{M}$

Let $\delta_{dec} > 0$ a small constant to be specified later. We define

$$\begin{aligned} {}^{(ext)}\mathfrak{D}_0[\alpha] &:= \sup_{{}^{(ext)}\mathcal{M}} \left(r^2(2r+u)^{1+\delta_{dec}} + r^3(2r+u)^{\frac{1}{2}+\delta_{dec}} \right) |\alpha|, \\ {}^{(ext)}\mathfrak{D}_0[\beta] &:= \sup_{{}^{(ext)}\mathcal{M}} \left(r^2(2r+u)^{1+\delta_{dec}} + r^3(2r+u)^{\frac{1}{2}+\delta_{dec}} \right) |\beta|, \\ {}^{(ext)}\mathfrak{D}_0[\check{\rho}] &:= \sup_{{}^{(ext)}\mathcal{M}} \left(r^2 u^{1+\delta_{dec}} + r^3 u^{\frac{1}{2}+\delta_{dec}} \right) |\check{\rho}|, \\ {}^{(ext)}\mathfrak{D}_0[\check{\mu}] &:= \sup_{{}^{(ext)}\mathcal{M}} r^3 u^{1+\delta_{dec}} |\check{\mu}|, \\ {}^{(ext)}\mathfrak{D}_0[\underline{\beta}] &:= \sup_{{}^{(ext)}\mathcal{M}} r^2 u^{1+\delta_{dec}} |\underline{\beta}|, \\ {}^{(ext)}\mathfrak{D}_0[\underline{\alpha}] &:= \sup_{{}^{(ext)}\mathcal{M}} r u^{1+\delta_{dec}} |\underline{\alpha}|, \end{aligned}$$

and

$${}^{(ext)}\mathfrak{D}_0[\check{R}] := {}^{(ext)}\mathfrak{D}_0[\alpha] + {}^{(ext)}\mathfrak{D}_0[\beta] + {}^{(ext)}\mathfrak{D}_0[\check{\rho}] + {}^{(ext)}\mathfrak{D}_0[\check{\mu}] + {}^{(ext)}\mathfrak{D}_0[\underline{\beta}] + {}^{(ext)}\mathfrak{D}_0[\underline{\alpha}].$$

Also, we introduce the following higher derivatives norms

$$\begin{aligned} {}^{(ext)}\mathfrak{D}_1[\check{R}] &:= {}^{(ext)}\mathfrak{D}_0[\check{R}] + {}^{(ext)}\mathfrak{D}_0[\mathfrak{D}\check{R}] \\ &+ \sup_{(ext)\mathcal{M}} \left(r^3(2r+u)^{1+\delta_{dec}} + r^4(2r+u)^{\frac{1}{2}+\delta_{dec}} \right) |e_3(\alpha)| \\ &+ \sup_{(ext)\mathcal{M}} \left(r^3u^{1+\delta_{dec}} + r^4u^{\frac{1}{2}+\delta_{dec}} \right) |e_3(\beta)| + \sup_{(ext)\mathcal{M}} r^3u^{1+\delta_{dec}} |e_3(\check{\rho})|, \end{aligned}$$

and for any integer $k \geq 2$

$${}^{(ext)}\mathfrak{D}_k[\check{R}] := {}^{(ext)}\mathfrak{D}_1[\mathfrak{D}^{\leq k-1}\check{R}].$$

Also, we define

$$\begin{aligned} {}^{(ext)}\mathfrak{D}_0[\check{\kappa}] &:= \sup_{(ext)\mathcal{M}} r^2u^{1+\delta_{dec}} |\check{\kappa}|, \\ {}^{(ext)}\mathfrak{D}_0[\vartheta] &:= \sup_{(ext)\mathcal{M}} \left(ru^{1+\delta_{dec}} + r^2u^{\frac{1}{2}+\delta_{dec}} \right) |\vartheta|, \\ {}^{(ext)}\mathfrak{D}_0[\zeta] &:= \sup_{(ext)\mathcal{M}} \left(ru^{1+\delta_{dec}} + r^2u^{\frac{1}{2}+\delta_{dec}} \right) |\zeta|, \\ {}^{(ext)}\mathfrak{D}_0[\check{\kappa}] &:= \sup_{(ext)\mathcal{M}} \left(ru^{1+\delta_{dec}} + r^2u^{\frac{1}{2}+\delta_{dec}} \right) |\check{\kappa}|, \\ {}^{(ext)}\mathfrak{D}_0[\underline{\vartheta}] &:= \sup_{(ext)\mathcal{M}} ru^{1+\delta_{dec}} |\underline{\vartheta}|, \\ {}^{(ext)}\mathfrak{D}_0[\eta] &:= \sup_{(ext)\mathcal{M}} ru^{1+\delta_{dec}} |\eta| + \left(\int_{\Sigma_*} u^{2+2\delta_{dec}} \eta^2 \right)^{\frac{1}{2}}, \\ {}^{(ext)}\mathfrak{D}_0[\check{\omega}] &:= \sup_{(ext)\mathcal{M}} ru^{1+\delta_{dec}} |\check{\omega}|, \\ {}^{(ext)}\mathfrak{D}_0[\underline{\xi}] &:= \sup_{(ext)\mathcal{M}} ru^{1+\delta_{dec}} |\underline{\xi}|, \end{aligned}$$

and

$$\begin{aligned} {}^{(ext)}\mathfrak{D}_0[\check{\Gamma}] &:= {}^{(ext)}\mathfrak{D}_0[\check{\kappa}] + {}^{(ext)}\mathfrak{D}_0[\vartheta] + {}^{(ext)}\mathfrak{D}_0[\zeta] + {}^{(ext)}\mathfrak{D}_0[\check{\kappa}] + {}^{(ext)}\mathfrak{D}_0[\underline{\vartheta}] + {}^{(ext)}\mathfrak{D}_0[\eta] \\ &+ {}^{(ext)}\mathfrak{D}_0[\check{\omega}] + {}^{(ext)}\mathfrak{D}_0[\underline{\xi}]. \end{aligned}$$

Also, we introduce the following higher derivatives norms

$${}^{(ext)}\mathfrak{D}_1[\check{\Gamma}] := {}^{(ext)}\mathfrak{D}_0[\mathfrak{D}\check{\Gamma}] + \sup_{(ext)\mathcal{M}} r^2u^{1+\delta_{dec}} |e_3(\vartheta, \zeta, \check{\kappa})|$$

and for any integer $k \geq 2$

$${}^{(ext)}\mathfrak{D}_k[\check{\Gamma}] := {}^{(ext)}\mathfrak{D}_1[\mathfrak{d}^{\leq k-1}\check{\Gamma}].$$

Remark 3.2.4. *The integral bootstrap assumption on Σ_* for η will only be needed in the proof of Proposition 3.4.6 and recovered in Proposition 7.3.5. In fact, other components satisfy an analog integral estimate on Σ_* : this is the case of $\underline{\vartheta}$, $\underline{\xi}$ and $r\underline{\beta}$, see Proposition 7.3.5. But η is the only component for which we need to make this type of bootstrap assumption.*

3.2.2 Main norms in ${}^{(int)}\mathcal{M}$

All quantities appearing in this section are defined relative to the ${}^{(int)}\mathcal{M}$ frame adapted to the $(\underline{u}, {}^{(int)}s)$ foliation.

L^2 based norms in ${}^{(int)}\mathcal{M}$

We introduce the curvature norms,

$$\left({}^{(int)}\mathfrak{R}_0[\check{R}] \right)^2 := \int_{{}^{(int)}\mathcal{M}} |\check{R}|^2.$$

For any nonzero integer k , we introduce the following higher derivatives norms

$${}^{(int)}\mathfrak{R}_k[\check{R}] := {}^{(int)}\mathfrak{R}_0[\mathfrak{d}^{\leq k}\check{R}].$$

For any $k \geq 0$, we introduce the following norms

$$\left({}^{(int)}\mathfrak{G}_k[\check{\Gamma}] \right)^2 := \int_{{}^{(int)}\mathcal{M}} |\mathfrak{d}^{\leq k}\check{\Gamma}|^2.$$

Decay norms in ${}^{(int)}\mathcal{M}$

We define

$${}^{(int)}\mathfrak{D}_0[\check{R}] := \sup_{{}^{(int)}\mathcal{M}} \underline{u}^{1+\delta_{dec}} |\check{R}|, \quad {}^{(int)}\mathfrak{D}_0[\check{\Gamma}] := \sup_{{}^{(int)}\mathcal{M}} \underline{u}^{1+\delta_{dec}} |\check{\Gamma}|.$$

Also, we introduce the following higher derivatives norms for any integer $k \geq 1$

$${}^{(int)}\mathfrak{D}_k[\check{R}] := {}^{(int)}\mathfrak{D}_0[\mathfrak{d}^{\leq k}\check{R}], \quad {}^{(int)}\mathfrak{D}_k[\check{\Gamma}] := {}^{(int)}\mathfrak{D}_0[\mathfrak{d}^{\leq k}\check{\Gamma}].$$

3.2.3 Combined norms

We define the following norms \mathcal{M} by combining our above norms on ${}^{(ext)}\mathcal{M}$ and ${}^{(int)}\mathcal{M}$

$$\begin{aligned}\mathfrak{N}_k^{(En)} &:= {}^{(ext)}\mathfrak{R}_k[\check{R}] + {}^{(ext)}\mathfrak{G}_k[\check{\Gamma}] + {}^{(int)}\mathfrak{R}_k[\check{R}] + {}^{(int)}\mathfrak{G}_k[\check{\Gamma}], \\ \mathfrak{N}_k^{(Dec)} &:= {}^{(ext)}\mathfrak{D}_k[\check{R}] + {}^{(int)}\mathfrak{D}_k[\check{\Gamma}] + {}^{(ext)}\mathfrak{D}_k[\check{R}] + {}^{(int)}\mathfrak{D}_k[\check{\Gamma}].\end{aligned}$$

3.2.4 Initial layer norm

Recall the notations of section 3.1.1 concerning the initial data layer \mathcal{L}_0 . Recall that the constant $m_0 > 0$ is the mass of the initial Schwarzschild spacetime relative to which our initial perturbation is measured. We define the initial layer norm to be⁸,

$$\mathfrak{J}_k := {}^{(ext)}\mathfrak{J}_k + {}^{(int)}\mathfrak{J}_k + \mathfrak{J}'_k$$

where

$$\begin{aligned}{}^{(ext)}\mathfrak{J}_0 &:= \sup_{{}^{(ext)}\mathcal{L}_0} \left[r^{\frac{7}{2} + \delta_B} (|\alpha| + |\beta|) + r^3 \left| \rho + \frac{2m_0}{r^3} \right| + r^2 |\underline{\beta}| + r |\underline{\alpha}| \right] \\ &+ \sup_{{}^{(ext)}\mathcal{L}_0} r^2 \left(|\vartheta| + \left| \kappa - \frac{2}{r} \right| + |\zeta| + \left| \underline{\kappa} + \frac{2(1 - \frac{2m_0}{r})}{r} \right| \right) \\ &+ \sup_{{}^{(ext)}\mathcal{L}_0} r \left(|\underline{\vartheta}| + \left| \underline{\omega} - \frac{m_0}{r^2} \right| + |\underline{\xi}| \right) \\ &+ \sup_{{}^{(ext)}\mathcal{L}_0 ({}^{(ext)}r_0 \geq 4m_0)} \left(r \left| \frac{\gamma}{r^2} - 1 \right| + r |\underline{b}| + |\underline{\Omega} + \Upsilon| + |\varsigma - 1| + r \left| \frac{e^\Phi}{r \sin \theta} - 1 \right| \right),\end{aligned}$$

$$\begin{aligned}{}^{(int)}\mathfrak{J}_0 &:= \sup_{{}^{(int)}\mathcal{L}_0} \left(|\underline{\alpha}| + |\underline{\beta}| + \left| \rho + \frac{2m_0}{r^3} \right| + |\beta| + |\alpha| \right) \\ &+ \sup_{{}^{(int)}\mathcal{L}_0} \left(|\vartheta| + \left| \kappa - \frac{2(1 - \frac{2m_0}{r})}{r} \right| + |\zeta| + \left| \underline{\kappa} + \frac{2}{r} \right| + |\underline{\vartheta}| + \left| \omega + \frac{m_0}{r^2} \right| + |\xi| \right),\end{aligned}$$

$$\mathfrak{J}'_0 := \sup_{{}^{(int)}\mathcal{L}_0 \cap {}^{(ext)}\mathcal{L}_0} (|f| + |\underline{f}| + |\log(\lambda_0^{-1} \lambda)|), \quad \lambda_0 = {}^{(ext)}\lambda_0 = 1 - \frac{2m_0}{{}^{(ext)}r_{\mathcal{L}_0}},$$

⁸Recall that the initial data layer foliations satisfy $\underline{\eta} + \zeta = 0$, as well as $\xi = \omega = 0$ on ${}^{(ext)}\mathcal{L}_0$ and $\eta = \zeta$ as well as $\underline{\xi} = \underline{\omega} = 0$ on ${}^{(int)}\mathcal{L}_0$.

with \mathfrak{J}_k the corresponding higher derivative norms obtained by replacing each component by $\mathfrak{d}^{\leq k}$ of it. In the definition of \mathfrak{J}'_0 above, $(f, \underline{f}, \lambda)$ denote the transition functions of Lemma 2.3.1 from the frame of the outgoing part ${}^{(ext)}\mathcal{L}_0$ of the initial data layer to the frame of the ingoing part ${}^{(int)}\mathcal{L}_0$ of the initial data layer in the region ${}^{(int)}\mathcal{L}_0 \cap {}^{(ext)}\mathcal{L}_0$.

Remark 3.2.5. *Note that In the definition of ${}^{(ext)}\mathfrak{J}_k$ we allow a higher power of r in front α, β and their derivatives than what it is consistent with the results of [17] and [37]. The additional r^{δ_B} power, for δ_B small, is consistent instead with the result of [38].*

3.3 Main Theorem

3.3.1 Smallness constants

Before stating our main theorem, we first introduce the following constants that will be involved in its statement.

- The constant $m_0 > 0$ is the mass of the initial Schwarzschild spacetime relative to which our initial perturbation is measured.
- The integer k_{large} which corresponds to the maximum number of derivatives of the solution.
- The size of the initial data layer norm is measured by $\epsilon_0 > 0$.
- The size of the bootstrap assumption norms are measured by $\epsilon > 0$.
- $\delta_{\mathcal{H}} > 0$ measures the width of the region $|r - 2m_0| \leq 2m_0\delta_{\mathcal{H}}$ where the redshift estimate holds and which includes in particular the region ${}^{(int)}\mathcal{M}$.
- δ_{dec} is tied to decay estimates in u, \underline{u} for $\check{\Gamma}$ and \check{R} .
- δ_B is involved in the r -power of the r^p weighted estimates for curvature.

In what follows m_0 is a fixed constant, $\delta_{\mathcal{H}}$, δ_B , and δ_{dec} are fixed, sufficiently small, universal constants, and k_{large} is a fixed, sufficiently large, universal constant, chosen such that

$$0 < \delta_{\mathcal{H}}, \delta_{dec}, \delta_B \ll \min\{m_0, 1\}, \quad \delta_B > 2\delta_{dec}, \quad k_{large} \gg \frac{1}{\delta_{dec}}. \quad (3.3.1)$$

Then, ϵ and ϵ_0 are chosen such that

$$\epsilon_0, \epsilon \ll \min \left\{ \delta_{\mathcal{H}}, \delta_{dec}, \delta_B, \frac{1}{k_{large}}, m_0, 1 \right\} \quad (3.3.2)$$

and

$$\epsilon = \epsilon_0^{\frac{2}{3}}. \quad (3.3.3)$$

Using the definition of ϵ_0 , we may now precise the behavior (3.1.2) of r on Σ_*

$$\inf_{\Sigma_*} r \geq \epsilon_0^{-\frac{2}{3}} u_*^4. \quad (3.3.4)$$

From now on, in the rest of the paper, \lesssim means bounded by a constant depending only on geometric universal constants (such as Sobolev embeddings, elliptic estimates,...) as well as the constants

$$m_0, \delta_{\mathcal{H}}, \delta_{dec}, \delta_B, k_{large}$$

but not on ϵ and ϵ_0 .

3.3.2 Statement of the main theorem

We are now ready to give the following precise version of our main theorem.

Main Theorem (Main theorem, version 2). *There exists a sufficiently large integer k_{large} and a sufficiently small constant $\epsilon_0 > 0$ such that given an initial layer defined as in section 3.1.1 and satisfying the bound*

$$\mathfrak{I}_{k_{large}+5} \leq \epsilon_0, \quad (3.3.5)$$

there exists a globally hyperbolic development with a complete future null infinity \mathcal{I}_+ and a complete future horizon \mathcal{H}_+ together with foliations and adapted null frames verifying the admissibility conditions of section 3.1.2 such that following bound is satisfied

$$\mathfrak{N}_{k_{large}}^{(En)} + \mathfrak{N}_{k_{small}}^{(Dec)} \leq C\epsilon_0 \quad (3.3.6)$$

where C is a large enough universal constant and where k_{small} is given by

$$k_{small} = \left\lfloor \frac{1}{2} k_{large} \right\rfloor + 1. \quad (3.3.7)$$

In particular,

- On ${}^{(ext)}\mathcal{M}$, we have

$$\begin{aligned} |\alpha|, |\beta| &\lesssim \min \left\{ \frac{\epsilon_0}{r^3(u+2r)^{\frac{1}{2}+\delta_{dec}}}, \frac{\epsilon_0}{r^2(u+2r)^{1+\delta_{dec}}} \right\}, \\ |\check{\rho}| &\lesssim \min \left\{ \frac{\epsilon_0}{r^3u^{\frac{1}{2}+\delta_{dec}}}, \frac{\epsilon_0}{r^2u^{1+\delta_{dec}}} \right\}, \\ |\underline{\beta}| &\lesssim \frac{\epsilon_0}{r^2u^{1+\delta_{dec}}}, \\ |\underline{\alpha}| &\lesssim \frac{\epsilon_0}{ru^{1+\delta_{dec}}}, \end{aligned}$$

and

$$\begin{aligned} |\check{\kappa}| &\lesssim \frac{\epsilon_0}{r^2u^{1+\delta_{dec}}}, \\ |\vartheta|, |\zeta|, |\check{\kappa}| &\lesssim \min \left\{ \frac{\epsilon_0}{r^2u^{\frac{1}{2}+\delta_{dec}}}, \frac{\epsilon_0}{ru^{1+\delta_{dec}}} \right\}, \\ |\eta|, |\vartheta|, |\check{\omega}|, |\underline{\xi}| &\lesssim \frac{\epsilon_0}{ru^{1+\delta_{dec}}}. \end{aligned}$$

- On ${}^{(int)}\mathcal{M}$ we have, with $\check{\Gamma} = \{\check{\kappa}, \vartheta, \zeta, \eta, \check{\kappa}, \vartheta, \check{\omega}, \xi\}$, $\check{R} = \{\alpha, \beta, \check{\rho}, \underline{\beta}, \underline{\alpha}\}$,

$$|\check{\Gamma}, \check{R}| \lesssim \frac{\epsilon_0}{\underline{u}^{1+\delta_{dec}}}.$$

- The Bondi mass converges as $u \rightarrow +\infty$ along \mathcal{I}_+ to the final Bondi mass which we denote by m_∞ . The final Bondi mass verifies the estimate

$$\left| \frac{m_\infty}{m_0} - 1 \right| \lesssim \epsilon_0.$$

In particular $m_\infty > 0$.

- The Hawking mass m satisfies

$$\frac{|m - m_\infty|}{m_0} \lesssim \begin{cases} \frac{\epsilon_0}{u^{1+\delta_{dec}}} & \text{on } {}^{(ext)}\mathcal{M}, \\ \frac{\epsilon_0}{\underline{u}^{1+\delta_{dec}}} & \text{on } {}^{(int)}\mathcal{M}. \end{cases}$$

- The location of the future Horizon \mathcal{H}_+ satisfies

$$r = 2m_\infty + O\left(\frac{\sqrt{\epsilon_0}}{\underline{u}^{1+\frac{\delta_{dec}}{2}}}\right) \text{ on } \mathcal{H}_+.$$

- On ${}^{(ext)}\mathcal{M}$, we have

$$\begin{aligned} \left| \rho + \frac{2m_\infty}{r^3} \right| &\lesssim \min \left\{ \frac{\epsilon_0}{r^3 u^{\frac{1}{2} + \delta_{dec}}}, \frac{\epsilon_0}{r^2 u^{1 + \delta_{dec}}} \right\}, \\ \left| \kappa - \frac{2}{r} \right| &\lesssim \frac{\epsilon_0}{r^2 u^{1 + \delta_{dec}}}, \\ \left| \underline{\kappa} + \frac{2 \left(1 - \frac{2m_\infty}{r}\right)}{r} \right| &\lesssim \min \left\{ \frac{\epsilon_0}{r^2 u^{\frac{1}{2} + \delta_{dec}}}, \frac{\epsilon_0}{r u^{1 + \delta_{dec}}} \right\}, \\ \left| \underline{\omega} - \frac{m_\infty}{r^2} \right| &\lesssim \frac{\epsilon_0}{r u^{1 + \delta_{dec}}}. \end{aligned}$$

- On ${}^{(int)}\mathcal{M}$, we have.

$$\left| \rho + \frac{2m_\infty}{r^3} \right|, \left| \underline{\kappa} + \frac{2}{r} \right|, \left| \kappa - \frac{2 \left(1 - \frac{2m_\infty}{r}\right)}{r} \right|, \left| \omega + \frac{m_\infty}{r^2} \right| \lesssim \frac{\epsilon_0}{\underline{u}^{1 + \delta_{dec}}}.$$

- On ${}^{(ext)}\mathcal{M}$, the space-time metric \mathbf{g} is given in the (u, r, θ, φ) coordinates system by

$$\mathbf{g} = \mathbf{g}_{m_\infty, (ext)\mathcal{M}} + O\left(\frac{\epsilon_0}{u^{1 + \delta_{dec}}}\right) \left((dr, du, rd\theta)^2, r^2(\sin\theta)^2(d\varphi)^2 \right)$$

where $\mathbf{g}_{m_\infty, (ext)\mathcal{M}}$ denotes the Schwarzschild metric of mass $m_\infty > 0$ in outgoing Eddington-Finkelstein coordinates, i.e.

$$\mathbf{g}_{m_\infty, (ext)\mathcal{M}} := -2dudr - \left(1 - \frac{2m_\infty}{r}\right) (du)^2 + r^2 \left((d\theta)^2 + (\sin\theta)^2 (d\varphi)^2 \right).$$

- On ${}^{(int)}\mathcal{M}$, the space-time metric \mathbf{g} is given in the $(\underline{u}, r, \theta, \varphi)$ coordinates system by

$$\mathbf{g} = \mathbf{g}_{m_\infty, (int)\mathcal{M}} + O\left(\frac{\epsilon_0}{\underline{u}^{1 + \delta_{dec}}}\right) \left((dr, d\underline{u}, rd\theta)^2, r^2(\sin\theta)^2(d\varphi)^2 \right)$$

where $\mathbf{g}_{m_\infty, (int)\mathcal{M}}$ denotes the Schwarzschild metric of mass $m_\infty > 0$ in ingoing Eddington-Finkelstein coordinates, i.e.

$$\mathbf{g}_{m_\infty, (int)\mathcal{M}} := 2d\underline{u}dr - \left(1 - \frac{2m_\infty}{r}\right) (d\underline{u})^2 + r^2 \left((d\theta)^2 + (\sin\theta)^2 (d\varphi)^2 \right).$$

Note that analog statements of the above estimates also hold for \mathfrak{D}^k derivatives with $k \leq k_{small}$.

Remark 3.3.1. *In this paper, we choose to specify the closeness to Schwarzschild of our initial data in the context of the Characteristic Cauchy problem. Note that the conclusions of our main theorem can be immediately extended to the case where the data are specified to be close to Schwarzschild on a spacelike hypersurface Σ . Indeed, one can reduce this latter case to our situation by invoking*

- *The results in [37] [38] which allow us to control the causal region between Σ and the outgoing part of the initial data layer⁹.*
- *A standard local existence result which controls the finite causal region between Σ and the ingoing part of the initial data layer.*

Remark 3.3.2. *In the context of the previous remark, we note that the constant $m_0 > 0$ appearing in the initial data layer norm of the assumption (3.3.5) of our main theorem does not necessarily coincide with the ADM mass of the corresponding initial data set on the spacelike hypersurface Σ . With respect to this ADM mass, we would recover the well known inequality stating that the final Bondi mass is smaller than the ADM mass.*

3.4 Bootstrap assumptions and first consequences

3.4.1 Main bootstrap assumptions

We assume that the combined norms $\mathfrak{N}_k^{(En)}$ and $\mathfrak{N}_k^{(Dec)}$ defined in section 3.2 verifies the following bounds

BA-E (*Bootstrap Assumptions on energies and weighted energies*)

$$\mathfrak{N}_{k_{large}}^{(En)} \leq \epsilon, \quad (3.4.1)$$

BA-D (*Bootstrap Assumptions on decay*)

$$\mathfrak{N}_{k_{small}}^{(Dec)} \leq \epsilon. \quad (3.4.2)$$

In the remaining of section 3.4.1, we state several simple consequences of the bootstrap assumptions which will be proved in Chapter 4.

⁹Note that the results of [38] are consistent with our initial data layer assumptions.

3.4.2 Control of the initial data

While the smallness constant involved in the bootstrap assumptions is $\epsilon > 0$, we need the smallness constant involved in the control of the initial data to be $\epsilon_0 > 0$. This is achieved in the theorem below.

Theorem M0. *Assume that the initial data layer \mathcal{L}_0 , as defined in section 3.1.1, satisfies*

$$\mathfrak{J}_{k_{large}+5} \leq \epsilon_0.$$

Then under the bootstrap assumptions BA-D on decay, the following holds true on the initial data hypersurface $\mathcal{C}_1 \cup \underline{\mathcal{C}}_1$,

$$\begin{aligned} & \max_{0 \leq k \leq k_{large}} \left\{ \sup_{\mathcal{C}_1} \left[r^{\frac{7}{2}+\delta_B} (|\mathfrak{d}^k (ext)\alpha| + |\mathfrak{d}^k (ext)\beta|) + r^{\frac{9}{2}+\delta_B} |\mathfrak{d}^{k-1} e_3((ext)\alpha)| \right] \right. \\ & \quad \left. + \sup_{\mathcal{C}_1} \left[r^3 \left| \mathfrak{d}^k \left((ext)\rho + \frac{2m_0}{r^3} \right) \right| + r^2 |\mathfrak{d}^k (ext)\underline{\beta}| + r |\mathfrak{d}^k (ext)\underline{\alpha}| \right] \right\} \lesssim \epsilon_0, \\ & \max_{0 \leq k \leq k_{large}} \sup_{\underline{\mathcal{C}}_1} \left[|\mathfrak{d}^k (int)\alpha| + |\mathfrak{d}^k (int)\beta| + \left| \mathfrak{d}^k \left((int)\rho + \frac{2m_0}{r^3} \right) \right| \right. \\ & \quad \left. + |\mathfrak{d}^k (int)\underline{\beta}| + |\mathfrak{d}^k (int)\underline{\alpha}| \right] \lesssim \epsilon_0, \end{aligned}$$

and

$$\sup_{\mathcal{C}_1 \cup \underline{\mathcal{C}}_1} \left| \frac{m}{m_0} - 1 \right| \lesssim \epsilon_0.$$

3.4.3 Control of averages and of the Hawking mass

The following two lemma are simple consequence of the bootstrap assumptions and will be proved in section 4.2.

Lemma 3.4.1 (Control of averages). *Assume given a GCM admissible spacetime \mathcal{M} as defined in section 3.1.2 verifying the bootstrap assumption for some sufficiently small $\epsilon > 0$. Then, we have*

$$\begin{aligned} & \sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} \left(r^3 \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\kappa} - \frac{2}{r} \right) \right| + r^3 \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| \right) \lesssim \epsilon_0, \\ & \sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} \left(r^2 \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\kappa} + \frac{2\Upsilon}{r} \right) \right| + r^2 \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\omega} - \frac{m}{r^2} \right) \right| \right) \lesssim \epsilon_0, \end{aligned}$$

$$\begin{aligned}
& \sup_{(ext)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}} \left(r^3 \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\kappa} - \frac{2}{r} \right) \right| + r^3 \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| \right) \lesssim \epsilon_0, \\
& \sup_{(ext)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}} \left(r^2 \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\kappa} + \frac{2\Upsilon}{r} \right) \right| + r^2 \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\omega} - \frac{m}{r^2} \right) \right| \right) \lesssim \epsilon_0, \\
& \sup_{(int)\mathcal{M}} u^{1+\delta_{dec}} \left(\left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\kappa} - \frac{2\Upsilon}{r} \right) \right| + \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| \right) \lesssim \epsilon_0, \\
& \sup_{(int)\mathcal{M}} u^{1+\delta_{dec}} \left(\left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\kappa} + \frac{2}{r} \right) \right| + \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\omega} + \frac{m}{r^2} \right) \right| \right) \lesssim \epsilon_0, \\
& \sup_{(int)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}} \left(\left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\kappa} - \frac{2\Upsilon}{r} \right) \right| + \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| \right) \lesssim \epsilon_0, \\
& \sup_{(int)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}} \left(\left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\kappa} + \frac{2}{r} \right) \right| + \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\omega} + \frac{m}{r^2} \right) \right| \right) \lesssim \epsilon_0.
\end{aligned}$$

Also, we have

$$\begin{aligned}
& \sup_{(ext)\mathcal{M}} \left(u^{1+\delta_{dec}r} \left| \mathfrak{d}^{\leq k_{small}} (\bar{\Omega} + \Upsilon) \right| + u^{\frac{1}{2}+\delta_{dec}r} \left| \mathfrak{d}^{\leq k_{large}} (\bar{\Omega} + \Upsilon) \right| \right) \lesssim \epsilon_0, \\
& \sup_{(int)\mathcal{M}} \left(u^{1+\delta_{dec}} \left| \mathfrak{d}^{\leq k_{small}} (\bar{\Omega} - \Upsilon) \right| + u^{\frac{1}{2}+\delta_{dec}} \left| \mathfrak{d}^{\leq k_{large}} (\bar{\Omega} - \Upsilon) \right| \right) \lesssim \epsilon_0.
\end{aligned}$$

Finally, recall that $\bar{\mu}$ and $\underline{\mu}$ are given by the following formula

$$\bar{\mu} = \frac{2m}{r^3} \quad \text{on } (ext)\mathcal{M}, \quad \underline{\mu} = \frac{2m}{r^3} \quad \text{on } (int)\mathcal{M}.$$

Lemma 3.4.2 (Control of the Hawking mass). *Assume given a GCM admissible spacetime \mathcal{M} as defined in section 3.1.2 verifying the bootstrap assumption for some sufficiently small $\epsilon > 0$. Then, we have*

$$\begin{aligned}
& \max_{0 \leq k \leq k_{large}} \sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} \left(|\mathfrak{d}^k e_3(m)| + r |\mathfrak{d}^k e_4(m)| \right) \lesssim \epsilon_0, \\
& \max_{0 \leq k \leq k_{large}} \sup_{(int)\mathcal{M}} \underline{u}^{1+\delta_{dec}} \left(|\mathfrak{d}^k e_3(m)| + |\mathfrak{d}^k e_4(m)| \right) \lesssim \epsilon_0.
\end{aligned}$$

The e_4 derivatives behave better in powers of r ,

$$\begin{aligned}
& \max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} r^2 u^{1+\delta_{dec}} |\mathfrak{d}^k e_4(m)| \lesssim \epsilon_0, \\
& \max_{0 \leq k \leq k_{large}} \sup_{(ext)\mathcal{M}} r^2 u^{\frac{1}{2}+\delta_{dec}} |\mathfrak{d}^k e_4(m)| \lesssim \epsilon_0.
\end{aligned}$$

Moreover,

$$\sup_{\mathcal{M}} \left| \frac{m}{m_0} - 1 \right| \lesssim \epsilon_0.$$

3.4.4 Control of coordinates system

The following two propositions on the existence of a suitable coordinates system both in ${}^{(ext)}\mathcal{M}$ and in ${}^{(int)}\mathcal{M}$ are also consequences of the bootstrap assumptions and will be proved in section 4.3.

Proposition 3.4.3 (Control of a coordinates system on ${}^{(ext)}\mathcal{M}$). *Let $\theta \in [0, \pi]$ be the \mathbf{Z} -invariant scalar on \mathcal{M} defined by (2.2.52), i.e.*

$$\theta = \cot^{-1}(re_\theta(\Phi)). \quad (3.4.3)$$

Consider the (u, r, θ, φ) coordinates system introduced in Proposition 2.2.23. Then, relative to these (u, r, θ, φ) coordinates,

1. The spacetime metric takes the form,

$$g = -\frac{4\varsigma}{r\bar{\kappa}}dudr + \frac{\varsigma^2(\bar{\kappa} + \underline{A})}{\bar{\kappa}}du^2 + \gamma \left(d\theta - \frac{1}{2}\varsigma\underline{b}du - \frac{\underline{b}}{2}\Theta \right)^2 \quad (3.4.4)$$

where,

$$b = e_4(\theta), \quad \underline{b} = e_3(\theta), \quad \gamma = \frac{1}{(e_\theta(\theta))^2} \quad (3.4.5)$$

and,

$$\Theta = \frac{4}{r\bar{\kappa}}dr - \varsigma \left(\frac{\bar{\kappa} + \underline{A}}{\bar{\kappa}} \right) du.$$

2. The reduced coordinates derivatives take the form,

$$\begin{aligned} \partial_r &= \frac{2}{r\bar{\kappa}}e_4 - \frac{2\sqrt{\gamma}}{r\bar{\kappa}}be_\theta, \\ \partial_\theta &= \sqrt{\gamma}e_\theta, \\ \partial_u &= \varsigma \left[\frac{1}{2}e_3 - \frac{1}{2}\frac{\bar{\kappa} + \underline{A}}{\bar{\kappa}}e_4 - \frac{1}{2}\sqrt{\gamma} \left(\underline{b} - \left(\frac{\bar{\kappa} + \underline{A}}{\bar{\kappa}} \right) b \right) e_\theta \right]. \end{aligned} \quad (3.4.6)$$

3. The following estimates hold true:

$$\begin{aligned} \max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} \left(ru^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}} \right) \left(\left| \mathfrak{d}^k \left(\frac{\gamma}{r^2} - 1 \right) \right| + r \left| \mathfrak{d}^k b \right| \right) &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} u^{1 + \delta_{dec}} \left(\left| \mathfrak{d}^k \underline{\Omega} \right| + \left| \mathfrak{d}^k (\varsigma - 1) \right| + r \left| \mathfrak{d}^k \underline{b} \right| \right) &\lesssim \epsilon. \end{aligned}$$

Also, e^Φ satisfies

$$\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} \left(ru^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}} \right) \left| \mathfrak{d}^k \left(\frac{e^\Phi}{r \sin \theta} - 1 \right) \right| \lesssim \epsilon.$$

Proposition 3.4.4 (Control of a coordinates system on $^{(int)}\mathcal{M}$). *Let $\theta \in [0, \pi]$ be the \mathbf{Z} -invariant scalar on \mathcal{M} defined by (3.4.3). Consider the $(\underline{u}, r, \theta, \varphi)$ coordinates system introduced in Proposition 2.2.26. Then, relative to these $(\underline{u}, r, \theta, \varphi)$ coordinates,*

1. *The spacetime metric takes the form,*

$$g = -\frac{4\underline{\varsigma}}{r\underline{\kappa}}d\underline{u}dr + \frac{\underline{\varsigma}^2(\overline{\kappa} + A)}{\overline{\kappa}}d\underline{u}^2 + \gamma \left(d\theta - \frac{1}{2}\underline{\varsigma}b d\underline{u} - \frac{b}{2}\underline{\Theta} \right)^2 \quad (3.4.7)$$

where,

$$b = e_4(\theta), \quad \underline{b} = e_3(\theta), \quad \gamma = \frac{1}{(e_\theta(\theta))^2} \quad (3.4.8)$$

and,

$$\underline{\Theta} := \frac{4}{r\underline{\kappa}}dr - \underline{\varsigma} \left(\frac{\overline{\kappa} + A}{\overline{\kappa}} \right) d\underline{u}.$$

2. *The reduced coordinates derivatives take the form,*

$$\begin{aligned} \partial_r &= \frac{2}{r\underline{\kappa}}e_3 - \frac{2\sqrt{\gamma}}{r\underline{\kappa}}\underline{b}e_\theta, \\ \partial_\theta &= \sqrt{\gamma}e_\theta, \\ \partial_{\underline{u}} &= \underline{\varsigma} \left[\frac{1}{2}e_4 - \frac{1}{2}\frac{\overline{\kappa} + A}{\overline{\kappa}}e_3 - \frac{1}{2}\sqrt{\gamma} \left(b - \left(\frac{\overline{\kappa} + A}{\overline{\kappa}} \right) \underline{b} \right) e_\theta \right]. \end{aligned} \quad (3.4.9)$$

3. *The following estimates hold true:*

$$\max_{0 \leq k \leq k_{small}} \sup_{(int)\mathcal{M}} \underline{u}^{1+\delta_{dec}} \left(|\mathfrak{d}^k \check{\Omega}| + |\mathfrak{d}^k(\underline{\varsigma} - 1)| + \left| \mathfrak{d}^k \left(\frac{\gamma}{r^2} - 1 \right) \right| + |\mathfrak{d}^k b| + |\mathfrak{d}^k \underline{b}| \right) \lesssim \epsilon.$$

Also, e^Φ satisfies

$$\max_{0 \leq k \leq k_{small}} \sup_{(int)\mathcal{M}} \underline{u}^{1+\delta_{dec}} \left| \mathfrak{d}^k \left(\frac{e^\Phi}{r \sin \theta} - 1 \right) \right| \lesssim \epsilon.$$

3.4.5 Pointwise bounds for high order derivatives

We will need later to interpolate between the estimates provided by the bootstrap assumptions on decay and the bootstrap assumptions on energy. To this end, we will need the following consequence of the bootstrap assumptions on weighted energies.

Proposition 3.4.5. *The Ricci coefficients and curvature components satisfy the following pointwise estimates on \mathcal{M}*

$$\begin{aligned} \max_{k \leq k_{large} - 5} \sup_{\mathcal{M}} & \left\{ r^{\frac{7}{2} + \frac{\delta_B}{2}} (|\mathfrak{d}^k \alpha| + |\mathfrak{d}^k \beta|) + r^3 (|\mathfrak{d}^k \mu| + |\mathfrak{d}^k \check{\rho}|) \right. \\ & + r^2 (|\mathfrak{d}^k \check{\kappa}| + |\mathfrak{d}^k \zeta| + |\mathfrak{d}^k \vartheta| + |\mathfrak{d}^k \check{\kappa}| + |\mathfrak{d}^k \underline{\beta}|) \\ & \left. + r (|\mathfrak{d}^k \underline{\vartheta}| + |\mathfrak{d}^k \underline{\vartheta}| + |\mathfrak{d}^k \underline{\omega}| + |\mathfrak{d}^k \underline{\xi}| + |\mathfrak{d}^k \underline{\alpha}|) \right\} \lesssim \epsilon. \end{aligned}$$

3.4.6 Construction of a second frame in ${}^{(ext)}\mathcal{M}$

Recall that the quantity \mathfrak{q} satisfies the following wave equation, see (2.4.7),

$$\square_2 \mathfrak{q} + \kappa \underline{\kappa} \mathfrak{q} = \text{Err}[\square_2 \mathfrak{q}]$$

where the nonlinear term $\text{Err}[\square_2 \mathfrak{q}]$ has the schematic structure exhibited in (2.4.8). Also, recall that according to our bootstrap assumption on decay and Proposition 3.4.5, η satisfies on ${}^{(ext)}\mathcal{M}$

$$|\mathfrak{d}^{\leq k_{small}} \eta| \leq \frac{\epsilon}{r u^{1 + \delta_{dec}}}, \quad |\mathfrak{d}^{\leq k_{large} - 5} \eta| \lesssim \frac{\epsilon}{r}.$$

As discuss in Remark 2.4.8, this decay in r^{-1} is too weak to derive suitable decay for \mathfrak{q} . We thus need to provide another frame for ${}^{(ext)}\mathcal{M}$. This is the aim of the following proposition.

Proposition 3.4.6. *Let an integer k_{loss} and a small constant $\delta_0 > 0$ satisfying¹⁰*

$$16 \leq k_{loss} \leq \frac{\delta_{dec}}{3} (k_{large} - k_{small}), \quad \delta_0 := \frac{k_{loss}}{k_{large} - k_{small}}. \quad (3.4.10)$$

Let (e_4, e_3, e_θ) the outgoing geodesic null frame of ${}^{(ext)}\mathcal{M}$. There exists another frame (e'_4, e'_3, e'_θ) of ${}^{(ext)}\mathcal{M}$ provided by

$$\begin{aligned} e'_4 &= e_4 + f e_\theta + \frac{1}{4} f^2 e_3, \\ e'_\theta &= e_\theta + \frac{1}{2} f e_3, \\ e'_3 &= e_3, \end{aligned}$$

¹⁰Recall from (3.3.1) and (3.3.7) that we have

$$0 < \delta_{dec} \ll 1, \quad \delta_{dec} k_{large} \gg 1, \quad k_{small} = \left\lfloor \frac{1}{2} k_{large} \right\rfloor + 1.$$

In particular, we have $\delta_{dec} (k_{large} - k_{small}) \gg 1$ and hence the exists an integer k_{loss} satisfying the required constraints.

such that the Ricci coefficients and curvature components with respect to that frame satisfy

$$\xi' = 0,$$

$$\begin{aligned} \max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{(ext)\mathcal{M}} & \left\{ \left(r^2 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r u^{1 + \delta_{dec} - 2\delta_0} \right) |\mathfrak{D}^k \Gamma'_g| + r u^{1 + \delta_{dec} - 2\delta_0} |\mathfrak{D}^k \Gamma'_b| \right. \\ & + r^2 u^{1 + \delta_{dec} - 2\delta_0} \left| \mathfrak{D}^{k-1} e'_3 \left(\kappa' - \frac{2}{r}, \underline{\kappa}' + \frac{2\Upsilon}{r}, \vartheta', \zeta', \underline{\eta}', \eta' \right) \right| \\ & + \left(r^{\frac{7}{2} + \frac{\delta_B}{2}} + r^3 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r^2 u^{1 + \delta_{dec} - 2\delta_0} \right) \left(|\mathfrak{D}^k \alpha'| + |\mathfrak{D}^k \beta'| \right) \\ & + \left(r^{\frac{9}{2} + \frac{\delta_B}{2}} + r^3 u^{1 + \delta_{dec}} + r^4 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} \right) |\mathfrak{D}^{k-1} e'_3(\alpha')| \\ & + \left(r^3 u^{1 + \delta_{dec}} + r^4 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} \right) |\mathfrak{D}^{k-1} e'_3(\beta')| \\ & + \left(r^3 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r^2 r u^{1 + \delta_{dec} - 2\delta_0} \right) |\mathfrak{D}^k \beta'| \\ & \left. + u^{1 + \delta_{dec} - 2\delta_0} \left(r^2 |\mathfrak{D}^k \underline{\beta}'| + r |\mathfrak{D}^k \underline{\alpha}'| \right) \right\} \lesssim \epsilon, \end{aligned}$$

where we have used the notation¹¹

$$\begin{aligned} \Gamma'_g &= \left\{ r\omega', \kappa' - \frac{2}{r}, \vartheta', \zeta', \eta', \underline{\eta}', \underline{\kappa}' + \frac{2\Upsilon}{r}, r^{-1}(e'_4(r) - 1), r^{-1}e'_\theta(r), e'_4(m) \right\}, \\ \Gamma'_b &= \left\{ \underline{\vartheta}', \underline{\omega}' - \frac{m}{r^2}, \underline{\xi}', r^{-1}(e'_3(r) + \Upsilon), r^{-1}e'_3(m) \right\}. \end{aligned}$$

Furthermore, f satisfies

$$\begin{aligned} |\mathfrak{D}^k f| &\lesssim \frac{\epsilon}{r u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + u^{1 + \delta_{dec} - 2\delta_0}}, \quad \text{for } k \leq k_{small} + k_{loss} + 2 \text{ on } (ext)\mathcal{M}, \\ |\mathfrak{D}^{k-1} e'_3 f| &\lesssim \frac{\epsilon}{r u^{1 + \delta_{dec} - 2\delta_0}} \quad \text{for } k \leq k_{small} + k_{loss} + 2 \text{ on } (ext)\mathcal{M}. \end{aligned} \tag{3.4.11}$$

Remark 3.4.7. The crucial point of Proposition 3.4.6 is that in the new frame (e'_4, e'_3, e'_θ) of $(ext)\mathcal{M}$, η' belongs to Γ'_g and thus displays a better decay in r^{-1} than η corresponding to the outgoing geodesic frame (e_4, e_3, e_θ) of $(ext)\mathcal{M}$.

¹¹Here, r and m denote respectively the area radius and the Hawking mass of the outgoing geodesic foliation of $(ext)\mathcal{M}$, i.e. $r = (ext)r$ and $m = (ext)m$. In particular, while $e_\theta(r) = e_\theta(m) = 0$, we have in general $e'_\theta(r) \neq 0$ and $e'_\theta(m) \neq 0$.

3.5 Global null frames

In this section, we construct 2 smooth global frames on \mathcal{M} by matching the frame of ${}^{(int)}\mathcal{M}$ on the one hand with a renormalization of the frame on ${}^{(ext)}\mathcal{M}$, and on the other hand, with a renormalization of the second frame of ${}^{(ext)}\mathcal{M}$ given by Proposition 3.4.6.

3.5.1 Extension of frames

To construct the first global frame, we need to extend the frame $({}^{(int)}e_4, {}^{(int)}e_3, {}^{(int)}e_\theta)$ of ${}^{(int)}\mathcal{M}$ slightly into ${}^{(ext)}\mathcal{M}$, and the frame $({}^{(ext)}e_4, {}^{(ext)}e_3, {}^{(ext)}e_\theta)$ of ${}^{(ext)}\mathcal{M}$ slightly into ${}^{(int)}\mathcal{M}$. We keep the same labels for the extended frame, i.e. $({}^{(int)}e_4, {}^{(int)}e_3, {}^{(int)}e_\theta)$ represents the extended frame of ${}^{(int)}\mathcal{M}$ in ${}^{(ext)}\mathcal{M}$ and vice versa. This convention also applies to the Ricci coefficients, curvature components, area radius and Hawking mass of the extended frames.

Note that these extensions require, in addition to the initialization of the frames on \mathcal{T} , to initialize

1. $({}^{(ext)}e_4, {}^{(ext)}e_3, {}^{(ext)}e_\theta)$ on $\underline{\mathcal{C}}_*$ by

$$({}^{(ext)}e_4, {}^{(ext)}e_3, {}^{(ext)}e_\theta) = (({}^{(int)}\Upsilon)^{-1}({}^{(int)}e_4, {}^{(int)}\Upsilon({}^{(int)}e_3, {}^{(int)}e_\theta)).$$

2. $({}^{(int)}e_4, {}^{(int)}e_3, {}^{(int)}e_\theta)$ on \mathcal{C}_* by

$$({}^{(int)}e_4, {}^{(int)}e_3, {}^{(int)}e_\theta) = ({}^{(ext)}\Upsilon({}^{(ext)}e_4, ({}^{(ext)}\Upsilon)^{-1}({}^{(ext)}e_3, {}^{(ext)}e_\theta)).$$

3.5.2 Construction of the first global frame

We start with the definition of the region where the frame of ${}^{(int)}\mathcal{M}$ and a conformal renormalization of the frame of ${}^{(ext)}\mathcal{M}$ will be matched.

Definition 3.5.1. *We define the matching region as the spacetime region*

$$\text{Match} := \left({}^{(ext)}\mathcal{M} \cap \left\{ {}^{(int)}r \leq 2m_0 \left(1 + \frac{3}{2}\delta_{\mathcal{H}} \right) \right\} \right) \cup \left({}^{(int)}\mathcal{M} \cap \left\{ {}^{(int)}r \geq 2m_0 \left(1 + \frac{1}{2}\delta_{\mathcal{H}} \right) \right\} \right),$$

where, as explained in the previous section, ${}^{(int)}r$ denotes the area radius of the ingoing geodesic foliation of ${}^{(int)}\mathcal{M}$ and its extension to ${}^{(ext)}\mathcal{M}$.

Here is our main proposition concerning our first global frame.

Proposition 3.5.2. *There exists a global null frame defined on $(int)\mathcal{M} \cup (ext)\mathcal{M}$ and denoted by $(glo)e_4, (glo)e_3, (glo)e_\theta$ such that*

(a) *In $(ext)\mathcal{M} \setminus Match$, we have*

$${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta = ({}^{(ext)}\Upsilon {}^{(ext)}e_4, {}^{(ext)}\Upsilon^{-1}({}^{(ext)}e_3, {}^{(ext)}e_\theta).$$

(b) *In $(int)\mathcal{M} \setminus Match$, we have*

$${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta = ({}^{(int)}e_4, {}^{(int)}e_3, {}^{(int)}e_\theta).$$

(c) *In the matching region, we have*

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-2} \sup_{Match \cap (int)\mathcal{M}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k ({}^{(glo)}\check{\Gamma}, {}^{(glo)}\check{R})| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{small}-2} \sup_{Match \cap (ext)\mathcal{M}} u^{1+\delta_{dec}} |\mathfrak{d}^k ({}^{(glo)}\check{\Gamma}, {}^{(glo)}\check{R})| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{large}-1} \left(\int_{Match} |\mathfrak{d}^k ({}^{(glo)}\check{\Gamma}, {}^{(glo)}\check{R})|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon, \end{aligned}$$

where $(glo)\check{R}$ and $(glo)\check{\Gamma}$ are given by

$$\begin{aligned} (glo)\check{R} &= \left\{ \alpha, \beta, \rho + \frac{2m}{r^3}, \underline{\beta}, \underline{\alpha} \right\}, \\ (glo)\check{\Gamma} &= \left\{ \xi, \omega + \frac{m}{r^2}, \kappa - \frac{2\Upsilon}{r}, \vartheta, \zeta, \eta, \underline{\eta}, \underline{\kappa} + \frac{2}{r}, \underline{\vartheta}, \underline{\omega}, \underline{\xi} \right\}. \end{aligned}$$

(d) *Furthermore, we may also choose the global frame such that, in addition, one of the following two possibilities hold,*

i. *We have on all $(ext)\mathcal{M}$*

$${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta = ({}^{(ext)}\Upsilon {}^{(ext)}e_4, {}^{(ext)}\Upsilon^{-1}({}^{(ext)}e_3, {}^{(ext)}e_\theta).$$

ii. *We have on all $(int)\mathcal{M}$*

$${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta = ({}^{(int)}e_4, {}^{(int)}e_3, {}^{(int)}e_\theta).$$

Remark 3.5.3. *The global frame on \mathcal{M} of Proposition 3.5.2 will be used to construct the second global frame in the next section, see Proposition 3.5.5. It will also be used to recover high order derivatives in Theorem M8 (stated in section 3.6.2), see section 8.3.2.*

3.5.3 Construction of the second global frame

We start with the definition of the region where first global frame of \mathcal{M} (i.e. the one of Proposition 3.5.2) and a conformal renormalization of the frame second frame of ${}^{(ext)}\mathcal{M}$ (i.e. the one of Proposition 3.4.6) will be matched.

Definition 3.5.4. *We define the matching region as the spacetime region*

$$Match' := {}^{(ext)}\mathcal{M} \cap \left\{ \frac{7m_0}{2} \leq {}^{(ext)}r \leq 4m_0 \right\},$$

where ${}^{(ext)}r$ denotes the area radius of the outgoing geodesic foliation of ${}^{(ext)}\mathcal{M}$.

Here is our main proposition concerning our second global frame.

Proposition 3.5.5. *Let an integer k_{loss} and a small constant $\delta_0 > 0$ satisfying (3.4.10). There exists a global null frame $({}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta)$ defined on ${}^{(int)}\mathcal{M} \cup {}^{(ext)}\mathcal{M}$ such that*

(a) *In ${}^{(ext)}\mathcal{M} \cap \{{}^{(ext)}r \geq 4m_0\}$, we have*

$$({}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta) = ({}^{(ext)}\Upsilon {}^{(ext)}e'_4, {}^{(ext)}\Upsilon^{-1} {}^{(ext)}e'_3, {}^{(ext)}e'_\theta),$$

where $({}^{(ext)}e'_4, {}^{(ext)}e'_3, {}^{(ext)}e'_\theta)$ denotes the second frame of ${}^{(ext)}\mathcal{M}$, i.e. the frame of Proposition 3.4.6.

(b) *In ${}^{(int)}\mathcal{M} \cup ({}^{(ext)}\mathcal{M} \cap \{{}^{(ext)}r \leq \frac{7m_0}{2}\})$, we have*

$$({}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta) = ({}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta),$$

where $({}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta)$ denotes the first global frame of \mathcal{M} , i.e. the frame of Proposition 3.5.2.

(c) *In the matching region, we have*

$$\max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{Match'} u^{1+\delta_{dec}-2\delta_0} \left| \mathfrak{d}^k ({}^{(glo')}\check{\Gamma}, {}^{(glo')}\check{R}) \right| \lesssim \epsilon,$$

where ${}^{(glo')}\check{R}$ and ${}^{(glo')}\check{\Gamma}$ are given by

$$\begin{aligned} {}^{(glo')}\check{R} &= \left\{ \alpha, \beta, \rho + \frac{2m}{r^3}, \underline{\beta}, \underline{\alpha} \right\}, \\ {}^{(glo')}\check{\Gamma} &= \left\{ \xi, \omega + \frac{m}{r^2}, \kappa - \frac{2\Upsilon}{r}, \vartheta, \zeta, \eta, \underline{\eta}, \underline{\kappa} + \frac{2}{r}, \underline{\vartheta}, \underline{\omega}, \underline{\xi} \right\}. \end{aligned}$$

with the Ricci coefficients and curvature components being the one associated to the frame $({}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta)$.

(d) Furthermore, we may also choose the global frame such that, in addition, one of the following two possibilities hold,

i. We have on ${}^{(ext)}\mathcal{M} \cap \{ {}^{(ext)}r \geq \frac{15m_0}{4} \}$

$$({}^{(gl'o')}e_4, {}^{(gl'o')}e_3, {}^{(gl'o')}e_\theta) = ({}^{(ext)}\Upsilon {}^{(ext)}e'_4, {}^{(ext)}\Upsilon^{-1}({}^{(ext)}e'_3, {}^{(ext)}e'_\theta)).$$

ii. We have on ${}^{(int)}\mathcal{M} \cup ({}^{(ext)}\mathcal{M} \cap \{ {}^{(ext)}r \leq \frac{15m_0}{4} \})$

$$({}^{(gl'o')}e_4, {}^{(gl'o')}e_3, {}^{(gl'o')}e_\theta) = ({}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta).$$

Remark 3.5.6. The global frame on \mathcal{M} of Proposition 3.5.5 will be needed to derive decay estimates for the quantity \mathfrak{q} in Theorem M1 (stated in section 3.6.1).

3.6 Proof of the main theorem

3.6.1 Main intermediate results

We are ready to state our main intermediary results.

Theorem M1. Assume given a GCM admissible spacetime \mathcal{M} as defined in section 3.1.2 verifying the bootstrap assumptions¹² **BA-E** and **BA-D** for some sufficiently small $\epsilon > 0$. Then, if $\epsilon_0 > 0$ is sufficiently small, there exists $\delta_{extra} > \delta_{dec}$ such that we have the following estimates in \mathcal{M} ,

$$\begin{aligned} \max_{0 \leq k \leq k_{small} + 20} \sup_{({}^{(ext)}\mathcal{M})} \left\{ \left(ru^{\frac{1}{2} + \delta_{extra}} + u^{1 + \delta_{extra}} \right) |\mathfrak{d}^k \mathfrak{q}| + ru^{1 + \delta_{extra}} |\mathfrak{d}^k e_3 \mathfrak{q}| \right\} \\ + \max_{0 \leq k \leq k_{small} + 20} \sup_{({}^{(int)}\mathcal{M})} \underline{u}^{1 + \delta_{extra}} |\mathfrak{d}^k \mathfrak{q}| \lesssim \epsilon_0. \end{aligned}$$

Moreover, \mathfrak{q} also satisfies the following estimate

$$\max_{0 \leq k \leq k_{small} + 21} \underline{u}^{2 + 2\delta_{extra}} \int_{({}^{(int)}\mathcal{M})_{(\geq \underline{u})}} |\mathfrak{d}^k \mathfrak{q}|^2 + \max_{0 \leq k \leq k_{small} + 20} u^{2 + 2\delta_{extra}} \int_{\Sigma_{*}(\geq u)} |\mathfrak{d}^k e_3 \mathfrak{q}|^2 \lesssim \epsilon_0^2.$$

Theorem M2. Under the same assumptions as above we have the following decay estimates for ${}^{(ext)}\alpha$

$$\max_{0 \leq k \leq k_{small} + 20} \sup_{({}^{(ext)}\mathcal{M})} \left(\frac{r^2(2r + u)^{1 + \delta_{extra}}}{\log(1 + u)} + r^3(2r + u)^{\frac{1}{2} + \delta_{extra}} \right) \left(|\mathfrak{d}^k ({}^{(ext)}\alpha)| + r |\mathfrak{d}^k e_3 ({}^{(ext)}\alpha)| \right) \lesssim \epsilon_0.$$

¹²Recall in particular that the conclusions of Theorem M0 hold under the bootstrap assumptions **BA-E** and **BA-D**.

Theorem M3. *Under the same assumptions as above we have the following decay estimates for $\underline{\alpha}$*

$${}^{(int)}\mathfrak{D}_{k_{small}+16}[\underline{\alpha}] \lesssim \epsilon_0, \quad \max_{0 \leq k \leq k_{small}+18} \int_{\Sigma_*} u^{2+2\delta_{extra}} |\mathfrak{d}^k \underline{\alpha}|^2 \lesssim \epsilon_0^2.$$

Theorem M4. *Under the same assumptions as above we also have the following decay estimates in ${}^{(ext)}\mathcal{M}$*

$${}^{(ext)}\mathfrak{D}_{k_{small}+8}[\check{R}] + {}^{(ext)}\mathfrak{D}_{k_{small}+8}[\check{\Gamma}] \lesssim \epsilon_0.$$

Theorem M5. *Under the same assumptions as above we also have the following decay estimates for \check{R} and $\check{\Gamma}$ in ${}^{(int)}\mathcal{M}$*

$${}^{(int)}\mathfrak{D}_{k_{small}+5}[\check{R}] + {}^{(int)}\mathfrak{D}_{k_{small}+5}[\check{\Gamma}] \lesssim \epsilon_0.$$

Note that, as an immediate consequence of Theorem M2 to Theorem M5 we have obtained, under the same assumptions as above, the following improvement of our bootstrap assumptions on decay

$$\mathfrak{N}_{k_{small}+5}^{(Dec)} \lesssim \epsilon_0. \quad (3.6.1)$$

3.6.2 End of the proof of the main theorem

Definition 3.6.1 (Definition of $\mathfrak{N}(u_*)$). *Let $\epsilon_0 > 0$ and $\epsilon > 0$ be given small constants satisfying the constraint (3.3.3). Let $\mathfrak{N}(u_*)$ be the set of all GCM admissible spacetimes \mathcal{M} defined in section 3.1.2 such that*

- u_* is the value of u on the last outgoing slice \mathcal{C}_* ,
- u_* satisfies (3.3.4),
- the bootstrap assumptions (3.4.1) (3.4.2) hold true, i.e., relative to the combined norms defined in section 3.2.3, we have

$$\mathfrak{N}_{k_{large}}^{(En)} \leq \epsilon, \quad \mathfrak{N}_{k_{small}}^{(Dec)} \leq \epsilon.$$

Definition 3.6.2. *Let \mathcal{U} be the set of all values of $u_* \geq 0$ such that the spacetime $\mathfrak{N}(u_*)$ exists.*

The following theorem shows that \mathcal{U} is not empty.

Theorem M6. *There exists $\delta_0 > 0$ small enough such that for sufficiently small constants $\epsilon_0 > 0$ and $\epsilon > 0$ satisfying the constraints (3.3.3) (3.3.4), we have $[1, 1 + \delta_0] \subset \mathcal{U}$.*

In view of Theorem M6, we may define U_* as the supremum over all value of u_* that belongs to \mathcal{U} .

$$U_* := \sup_{u_* \in \mathcal{U}} u_*.$$

Assume by contradiction that

$$U_* < +\infty.$$

Then, by the continuity of the flow, $U_* \in \mathcal{U}$. Furthermore, according to the consequence (3.6.1) of Theorem M2 to Theorem M5, the bootstrap assumptions on decay (3.4.2) on any spacetime of $\mathfrak{N}(U_*)$ are improved by

$$\mathfrak{N}_{k_{small}+5}^{(Dec)} \lesssim \epsilon_0.$$

To reach a contradiction, we still need an extension procedure for spacetimes in $\mathfrak{N}(u_*)$ to larger values of u , as well as to improve our bootstrap assumptions on weighted energies (3.4.1). This is done in two steps.

Theorem M7. *Any GCM admissible spacetime in $\mathfrak{N}(u_*)$ for some $0 < u_* < +\infty$ such that*

$$\mathfrak{N}_{k_{small}+5}^{(Dec)} \lesssim \epsilon_0,$$

has a GCM admissible extension (satisfying (3.3.4)), i.e. $u'_ > u_*$, initialized by Theorem M0, which verifies*

$$\mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0.$$

Remark 3.6.3. *Recall that the definition of a GCM admissible spacetime in section 3.1.2 is such that $\mathcal{T} = \{r = r_{\mathcal{T}}\}$ for some $r_{\mathcal{T}}$ satisfying*

$$2m_0 \left(1 + \frac{\delta_{\mathcal{H}}}{2}\right) \leq r_{\mathcal{T}} \leq 2m_0 \left(1 + \frac{3\delta_{\mathcal{H}}}{2}\right). \quad (3.6.2)$$

All results obtained so far, in particular Theorems M0–M7, hold for any choice of $r_{\mathcal{T}}$ satisfying (3.6.2), see Remark 8.3.1 for a more precise statement. It is at this stage, in Theorem M8 below, that we need to make a specific choice of $r_{\mathcal{T}}$ in the context of a Lebesgue point argument required for the control of top order derivatives. This choice will be made in (8.3.2).

Theorem M8. *There exists a choice of $r_{\mathcal{T}}$ satisfying (3.6.2) such that the GCM admissible spacetime exhibited in Theorem M7 satisfies in addition*

$$\mathfrak{N}_{k_{large}}^{(En)} \lesssim \epsilon_0$$

and therefore belongs to $\mathfrak{N}(u'_*)$. In particular u'_* belongs to \mathcal{U} .

In view of Theorem M8, we have reached a contradiction, and hence

$$U_* = +\infty$$

so that the spacetime may be continued forever. This concludes the proof of the main theorem.

3.6.3 Conclusions

The Penrose diagram of \mathcal{M}

Complete future null infinity. We first deduce from our estimate that our spacetime \mathcal{M} has a complete future null infinity \mathcal{I}_+ . The portion of null infinity of \mathcal{M} corresponds to the limit $r \rightarrow +\infty$ along the leaves \mathcal{C}_u of the outgoing geodesic foliation of ${}^{(ext)}\mathcal{M}$. As \mathcal{C}_u exists for all $u \geq 0$ with suitable estimates, it suffices to prove that u is an affine parameter of \mathcal{I}_+ . To this end, recall from our main theorem that the estimates $\mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0$ hold which implies in particular¹³

$$\sup_{(ext)\mathcal{M}} ru^{1+\delta_{dec}} \left(|\underline{\xi}| + \left| \underline{\omega} - \frac{m}{r^2} \right| + r^{-1}|\varsigma - 1| \right) \lesssim \epsilon_0. \quad (3.6.3)$$

As $|m - m_0| \lesssim \epsilon_0 m_0$, see Lemma 3.4.2, m is bounded. We infer that

$$\lim_{\mathcal{C}_u, r \rightarrow +\infty} \underline{\xi}, \underline{\omega} = 0 \text{ for all } 1 \leq u < \infty.$$

In view of the identity

$$D_3 e_3 = -2\underline{\omega} e_3 + 2\underline{\xi} e_\theta,$$

we infer that e_3 is a null geodesic generator of \mathcal{I}_+ . Since we have $e_3(u) = \frac{2}{\xi}$ with $|\varsigma - 1| \lesssim \epsilon_0$ in view of (3.6.3), u is an affine parameter of \mathcal{I}_+ so that \mathcal{I}_+ is indeed complete.

¹³Using also Proposition 3.4.3 for the control of ς .

Existence of a future event horizon. Next, note that the estimates $\mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0$ also imply

$$\sup_{(int)\mathcal{M}} \underline{u}^{1+\delta_{dec}} \left(\left| \underline{\kappa} + \frac{2}{r} \right| + \left| \kappa - \frac{2(1-\frac{2m}{r})}{r} \right| \right) \lesssim \epsilon_0.$$

In particular, considering the spacetime region $r \leq 2m_0(1 - \delta_{\mathcal{H}}/2)$ of $(int)\mathcal{M}$, and in view of the estimate $|m - m_0| \lesssim \epsilon_0 m_0$, we infer, for all $r \leq 2m_0(1 - \delta_{\mathcal{H}}/2)$, that

$$\kappa \leq 2 \frac{r - 2m}{r^2} + O(\epsilon_0) \lesssim \frac{2}{r^2} (r - 2m_0 + 2m_0 - 2m) + O(\epsilon_0) \lesssim \frac{2m_0}{r^2} (-\delta_{\mathcal{H}} + \epsilon_0) + O(\epsilon_0).$$

Thus, since $0 < \epsilon_0 \ll \delta_{\mathcal{H}} \ll 1$, we deduce,

$$\begin{aligned} \sup_{(int)\mathcal{M}(r \leq 2m_0(1-\frac{\delta_{\mathcal{H}}}{2}))} \kappa &\leq -\frac{\delta_{\mathcal{H}}}{2m_0(1-\frac{\delta_{\mathcal{H}}}{2})^2} + O(\epsilon_0) \\ &\leq -\frac{\delta_{\mathcal{H}}}{4m_0}. \end{aligned}$$

Thus, all 2-spheres $S(\underline{u}, s)$ of the ingoing geodesic foliation of $(int)\mathcal{M}$ which are located in the spacetime region $r \leq 2m_0(1 - \delta_{\mathcal{H}}/2)$ of $(int)\mathcal{M}$ are trapped. This implies that the past of \mathcal{I}_+ in \mathcal{M} does not contain this region, and hence \mathcal{M} contains the event horizon \mathcal{H}_+ of a black hole in its interior. Moreover, since the timelike hyper surface \mathcal{T} is foliated by the outgoing null cones \mathcal{C}_u of $(ext)\mathcal{M}$, it is in the past of \mathcal{I}_+ . Hence, since \mathcal{T} is one of the boundaries of $(int)\mathcal{M}$, \mathcal{H}_+ is actually located in the interior of the region $(int)\mathcal{M}$.

Asymptotic stationarity of \mathcal{M} . Recall that we have introduced a vectorfield \mathbf{T} in $(ext)\mathcal{M}$ as well as one in $(int)\mathcal{M}$ by


$$\mathbf{T} = e_3 + \Upsilon e_4 \text{ in } (ext)\mathcal{M}, \quad \mathbf{T} = e_4 + \Upsilon e_3 \text{ in } (int)\mathcal{M}.$$

We can easily express all components of $(\mathbf{T})\pi$ in terms of $\check{\Gamma}, e_3(m), e_4(m)$. Thus, making us of the estimate $\mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0$ of our main theorem, we deduce,

$$|(\mathbf{T})\pi| \lesssim \frac{\epsilon_0}{r\underline{u}^{1+\delta_{dec}}} \text{ in } (ext)\mathcal{M} \text{ and } |(\mathbf{T})\pi| \lesssim \frac{\epsilon_0}{\underline{u}^{1+\delta_{dec}}} \text{ in } (int)\mathcal{M}.$$

In particular, \mathbf{T} is an asymptotically Killing vectorfield and hence our spacetime \mathcal{M} is asymptotically stationary.

The above conclusions regarding \mathcal{I}_+ and \mathcal{H}_+ allow us to draw the Penrose diagram of \mathcal{M} , see Figure 3.3.

Figure 3.3: The Penrose diagram of the space-time \mathcal{M}

Limits at null infinity and Bondi mass

Recall the following formula for the derivative of the Hawking mass in ${}^{(ext)}\mathcal{M}$, see Proposition 2.2.16

$$e_4(m) = \frac{r}{32\pi} \int_S \left(-\frac{1}{2} \kappa \vartheta^2 - \frac{1}{2} \check{\kappa} \vartheta \underline{\vartheta} + 2\check{\kappa} \check{\rho} + 2e_\theta(\kappa) \zeta + 2\kappa \zeta^2 \right).$$

As a simple corollary of the decay estimates of our main theorem, i.e., $\mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0$, we deduce,

$$|e_4(m)| \lesssim \frac{\epsilon_0^2}{r^2 u^{1+2\delta_{dec}}}. \quad (3.6.4)$$

Since r^{-2} is integrable, we infer the existence of a limit to m as $r \rightarrow +\infty$ along \mathcal{C}_u

$$M_B(u) = \lim_{r \rightarrow +\infty} m(u, r) \text{ for all } 1 \leq u < +\infty$$

where $M_B(u)$ is the so-called Bondi mass.

Next, we recall the following formula in ${}^{(ext)}\mathcal{M}$, see Proposition 2.2.8

$$e_4(\underline{\vartheta}) + \frac{1}{2}\kappa\underline{\vartheta} = 2\underline{\vartheta}^*\zeta - \frac{1}{2}\kappa\underline{\vartheta} + 2\underline{\zeta}^2.$$

In view of $\mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0$, we deduce

$$|e_4(r\underline{\vartheta})| \lesssim \frac{\epsilon_0}{r^2 u^{\frac{1}{2} + \delta_{dec}}}.$$

Since r^{-2} is integrable, we infer the existence of a limit to $r\underline{\vartheta}$ as $r \rightarrow +\infty$ along \mathcal{C}_u

$$\underline{\Theta}(u, \cdot) = \lim_{r \rightarrow +\infty} r\underline{\vartheta}(r, u, \cdot) \text{ for all } 1 \leq u < +\infty.$$

On the other hand, in view of $\mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0$ again,

$$r|\underline{\vartheta}| \lesssim \frac{\epsilon_0}{u^{1 + \delta_{dec}}}, \quad \text{on } {}^{(ext)}\mathcal{M}.$$

We infer that

$$|\underline{\Theta}(u, \cdot)| \lesssim \frac{\epsilon_0}{u^{1 + \delta_{dec}}} \text{ for all } 1 \leq u < +\infty.$$

The spheres at null infinity are round

The Gauss curvature is given by the formula,

$$K = -\rho - \frac{1}{4}\kappa\underline{\kappa} + \frac{1}{4}\underline{\vartheta}\underline{\vartheta}.$$

Thus, in view of our estimates in ${}^{(ext)}\mathcal{M}$,

$$\left| K - \frac{1}{r^2} \right| \lesssim \frac{\epsilon_0}{r^3 u^{\frac{1}{2} + \delta_{dec}}}$$

so that

$$\lim_{r \rightarrow +\infty} r^2 K = 1.$$

In particular the spheres at null infinity are round.

A Bondi mass formula

Using the formula for $e_3(m)$ in ${}^{(ext)}\mathcal{M}$, see Proposition 2.2.16, together with the estimates $\mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0$, we deduce

$$\left| e_3(m) + \frac{r}{64\pi} \int_S \kappa \underline{\vartheta}^2 \right| \lesssim \frac{\epsilon_0^2}{ru^{\frac{3}{2}+2\delta_{dec}}}$$

and hence

$$\left| e_3(m) + \frac{1}{8|S|} \int_S (r\underline{\vartheta})^2 \right| \lesssim \frac{\epsilon_0^2}{ru^{\frac{3}{2}+2\delta_{dec}}}.$$

Letting $r \rightarrow +\infty$ along \mathcal{C}_u , and using that the spheres at null infinity are round, we infer in view of the definition of M_B and $\underline{\Theta}$

$$e_3(M_B)(u) = -\frac{1}{8} \int_{\mathbb{S}^2} \underline{\Theta}^2(u, \cdot) \text{ for all } 1 \leq u < +\infty.$$

Since $e_3(u) = \frac{2}{\varsigma}$ and e_3 is orthogonal to the spheres foliating \mathcal{I}_+ , we infer $e_3 = \frac{2}{\varsigma} \partial_u$. Thus, we obtain the following Bondi mass type formula

$$\partial_u M_B(u) = -\frac{\varsigma}{16} \int_{\mathbb{S}^2} \underline{\Theta}^2(u, \cdot) \text{ for all } 1 \leq u < +\infty,$$

with ς satisfying (3.6.3).

Final Bondi mass

In view of the estimate

$$|\underline{\Theta}(u, \cdot)| \lesssim \frac{\epsilon_0}{u^{1+\delta_{dec}}} \text{ for all } 1 \leq u < +\infty,$$

and the control for ς in (3.6.3), we infer that

$$|\partial_u M_B(u)| \lesssim \frac{\epsilon_0^2}{u^{2+2\delta_{dec}}} \text{ for all } 1 \leq u < +\infty.$$

In particular, since $u^{-2-2\delta_{dec}}$ is integrable, the limit along \mathcal{I}_+ exists

$$M_B(+\infty) = \lim_{u \rightarrow +\infty} M_B(u)$$

and is the so-called final Bondi mass. We denote it as m_∞ , i.e. $m_\infty = M_B(+\infty)$.

Control of $m - m_\infty$. We have as a consequence of the above estimate for $\partial_u M_B$ and the definition of m_∞

$$|M_B(u) - m_\infty| \lesssim \frac{\epsilon_0^2}{u^{1+2\delta_{dec}}} \text{ for all } 1 \leq u < +\infty.$$

Also, recall from (3.6.4) that we have obtained in ${}^{(ext)}\mathcal{M}$

$$|e_4(m)| \lesssim \frac{\epsilon_0^2}{r^2 u^{1+2\delta_{dec}}}$$

which yields, together with the definition of $M_B(u)$, by integration in r at fixed u

$$|m(r, u) - M_B(u)| \lesssim \frac{\epsilon_0^2}{r u^{1+2\delta_{dec}}} \text{ in } {}^{(ext)}\mathcal{M}.$$

We infer

$$\sup_{{}^{(ext)}\mathcal{M}} u^{1+2\delta_{dec}} |m - m_\infty| \lesssim \epsilon_0^2. \quad (3.6.5)$$

Also, recall the following formula for the derivative of the Hawking mass in ${}^{(int)}\mathcal{M}$, see Proposition 2.2.16 in the context of an outgoing geodesic foliation,

$$e_3(m) = \frac{r}{32\pi} \int_S \left(-\frac{1}{2} \kappa \vartheta^2 - \frac{1}{2} \check{\kappa} \vartheta \vartheta + 2\check{\kappa} \check{\rho} - 2e_\theta(\check{\kappa}) \zeta + 2\check{\kappa} \zeta^2 \right).$$

Together with the estimates $\mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0$, we deduce

$$|e_3(m)| \lesssim \frac{\epsilon_0^2}{\underline{u}^{2+2\delta_{dec}}} \text{ on } {}^{(int)}\mathcal{M}$$

and hence by integration in r at fixed \underline{u} , for $r \in [2m_0(1 - \delta_{\mathcal{H}}), r_{\mathcal{T}}]$,

$$\left| m(r, \underline{u}) - m(r_{\mathcal{T}}, \underline{u}) \right| \lesssim \frac{\epsilon_0^2}{\underline{u}^{2+2\delta_{dec}}} m_0 \delta_{\mathcal{H}} \quad \text{on } {}^{(int)}\mathcal{M}.$$

According to (3.6.5), since $\{r = r_{\mathcal{T}}\} = \mathcal{T} = {}^{(ext)}\mathcal{M} \cap {}^{(int)}\mathcal{M} \subset {}^{(ext)}\mathcal{M}$, and since $\underline{u} = u$ in \mathcal{T} by the initialization of \underline{u} ,

$$\underline{u}^{1+2\delta_{dec}} \left| m(r_{\mathcal{T}}, \underline{u}) - m_\infty \right| \lesssim \epsilon_0^2.$$

We deduce

$$\sup_{{}^{(int)}\mathcal{M}} \underline{u}^{1+2\delta_{dec}} |m - m_\infty| \lesssim \epsilon_0^2. \quad (3.6.6)$$

Combining (3.6.5) and (3.6.6) with the estimate

$$\sup_{\mathcal{M}} |m - m_0| \lesssim \epsilon_0 m_0,$$

in the statement of our main theorem (see also Lemma 3.4.2), we infer that

$$|m_\infty - m_0| \lesssim \epsilon_0 m_0.$$

In particular we deduce that $m_\infty > 0$ since ϵ_0 can be made arbitrarily small.

Coordinates systems on ${}^{(ext)}\mathcal{M}$ and ${}^{(int)}\mathcal{M}$

In view of Proposition 3.4.3, and together with the control of the averages $\bar{\kappa}$, $\bar{\underline{\kappa}}$ provided by Lemma 3.4.1, the control of $\tilde{\kappa}$ provided by the estimates $\mathfrak{N}_{\tilde{\kappa}_{small}}^{(Dec)} \lesssim \epsilon_0$, and the control of $m - m_\infty$ obtained in (3.6.5), we infer for the space-time metric \mathbf{g} on ${}^{(ext)}\mathcal{M}$ in the (u, r, θ, φ) coordinates system

$$\mathbf{g} = \mathbf{g}_{m_\infty, {}^{(ext)}\mathcal{M}} + O\left(\frac{\epsilon_0}{u^{1+\delta_{dec}}}\right) \left((dr, du, rd\theta)^2, r^2(\sin\theta)^2(d\varphi)^2 \right)$$

where $\mathbf{g}_{m_\infty, {}^{(ext)}\mathcal{M}}$ denotes the Schwarzschild metric of mass $m_\infty > 0$ in outgoing Eddington-Finkelstein coordinates, i.e.

$$\mathbf{g}_{m_\infty, {}^{(ext)}\mathcal{M}} = -2dudr - \left(1 - \frac{2m_\infty}{r}\right) (du)^2 + r^2 \left((d\theta)^2 + (\sin\theta)^2 (d\varphi)^2 \right).$$

Also, in view of Proposition 3.4.4, and together with the control of the averages $\bar{\kappa}$, $\bar{\underline{\kappa}}$ provided by Lemma 3.4.1, the control of $\tilde{\underline{\kappa}}$ provided by the estimates $\mathfrak{N}_{\tilde{\underline{\kappa}}_{small}}^{(Dec)} \lesssim \epsilon_0$, and the control of $m - m_\infty$ obtained in (3.6.6), we infer for the space-time metric \mathbf{g} on ${}^{(int)}\mathcal{M}$ in the $(\underline{u}, r, \theta, \varphi)$ coordinates system

$$\mathbf{g} = \mathbf{g}_{m_\infty, {}^{(int)}\mathcal{M}} + O\left(\frac{\epsilon_0}{\underline{u}^{1+\delta_{dec}}}\right) \left((dr, d\underline{u}, rd\theta)^2, r^2(\sin\theta)^2(d\varphi)^2 \right)$$

where $\mathbf{g}_{m_\infty, {}^{(int)}\mathcal{M}}$ denotes the Schwarzschild metric of mass $m_\infty > 0$ in ingoing Eddington-Finkelstein coordinates, i.e.

$$\mathbf{g}_{m_\infty, {}^{(int)}\mathcal{M}} = 2d\underline{u}dr - \left(1 - \frac{2m_\infty}{r}\right) (d\underline{u})^2 + r^2 \left((d\theta)^2 + (\sin\theta)^2 (d\varphi)^2 \right).$$

Asymptotic of the future event horizon. We show below that \mathcal{H}_+ is located in the following region of ${}^{(int)}\mathcal{M}$

$$2m \left(1 - \frac{\sqrt{\epsilon_0}}{\underline{u}^{1+\delta_{dec}}} \right) \leq r \leq 2m \left(1 + \frac{\sqrt{\epsilon_0}}{\underline{u}^{1+\frac{\delta_{dec}}{2}}} \right) \text{ on } \mathcal{H}_+ \text{ for any } 1 \leq \underline{u} < +\infty. \quad (3.6.7)$$

Note first that the lower bound follows from the fact that

$$\begin{aligned} \sup_{{}^{(int)}\mathcal{M} \left(r \leq 2m \left(1 - \frac{\sqrt{\epsilon_0}}{\underline{u}^{1+\delta_{dec}}} \right) \right)} \kappa &\leq -\frac{\frac{\sqrt{\epsilon_0}}{\underline{u}^{1+\delta_{dec}}}}{m \left(1 - \frac{\sqrt{\epsilon_0}}{\underline{u}^{1+\delta_{dec}}} \right)^2} + O\left(\frac{\epsilon_0}{\underline{u}^{1+\delta_{dec}}}\right) \\ &\leq -\frac{\sqrt{\epsilon_0}}{2m_0 \underline{u}^{1+\delta_{dec}}} < 0. \end{aligned}$$

Concerning the upper bound, we need to show that any 2-sphere

$$S(\underline{u}_1) := S\left(\underline{u}_1, r = 2m\left(1 + \frac{\sqrt{\epsilon_0}}{\underline{u}_1^{1+\frac{\delta_{dec}}{2}}}\right)\right), \quad 1 \leq \underline{u}_1 < +\infty \quad (3.6.8)$$

is in the past of \mathcal{I}_+ . Since ${}^{(ext)}\mathcal{M}$ is in the past of \mathcal{I}_+ , it suffices to show that the forward outgoing null cone emanating from any 2-sphere (3.6.8) reaches ${}^{(ext)}\mathcal{M}$ in finite time.

Assume, by contradiction, that there exists an outgoing null geodesic, denoted by γ , perpendicular to $S(\underline{u}_1)$, that does not reach ${}^{(ext)}\mathcal{M}$ in finite time. Let e'_4 be the geodesic generator of γ . In view of Lemma 2.3.1 on general null frame transformation, and denoting by (e_4, e_3, e_θ) the null frame¹⁴ of ${}^{(int)}\mathcal{M}$, we look for e'_4 under the form

$$e'_4 = \lambda\left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right),$$

and the fact that e'_4 is geodesic implies the following transport equations along γ for f and λ in view of Lemma 2.3.6 (applied¹⁵ with $\underline{f} = 0$)

$$\begin{aligned} \lambda^{-1}e'_4(f) + \left(\frac{\kappa}{2} + 2\omega\right)f &= -2\xi + E_1(f, \Gamma), \\ \lambda^{-1}e'_4(\log(\lambda)) &= 2\omega + E_2(f, \Gamma), \end{aligned}$$

where E_1 and E_2 are given schematically by

$$\begin{aligned} E_1(f, \Gamma) &= -\frac{1}{2}\vartheta f + \text{l.o.t.}, \\ E_2(f, \Gamma) &= f\zeta - \frac{1}{2}f^2\underline{\omega} - \underline{\eta}f - \frac{1}{4}f^2\underline{\kappa} + \text{l.o.t.} \end{aligned}$$

Here, l.o.t. denote terms which are cubic or higher order in f and Γ denotes the Ricci coefficients w.r.t. the original null frame (e_3, e_4, e_θ) of ${}^{(int)}\mathcal{M}$.

We then proceed as follows

1. First, we initialize f and λ as follows on the $\gamma \cap S(\underline{u}_1)$

$$f = 0, \quad \lambda = 1 \text{ on } \gamma \cap S(\underline{u}_1).$$

¹⁴Recall that we assume by contradiction that γ does not reach ${}^{(ext)}\mathcal{M}$ and hence stays in ${}^{(int)}\mathcal{M}$.

¹⁵i.e. we keep the direction of e_3 fixed.

2. Then, we initiate a continuity argument by assuming for some

$$\underline{u}_1 < \underline{u}_2 < \underline{u}_1 + \left(\frac{\underline{u}_1}{\epsilon_0}\right)^{\frac{\delta_{dec}}{2}}$$

that we have

$$|f| \leq \frac{\sqrt{\epsilon_0}}{\underline{u}_1^{\frac{1}{2} + \delta_{dec}}}, \quad \Upsilon \geq \frac{\sqrt{\epsilon_0}}{2\underline{u}_1^{1 + \frac{\delta_{dec}}{2}}}, \quad 0 < \lambda < +\infty \text{ on } \gamma(\underline{u}_1, \underline{u}_2) \cap {}^{(int)}\mathcal{M} \quad (3.6.9)$$

where $\gamma(\underline{u}_1, \underline{u}_2)$ denotes the portion of γ in $\underline{u}_1 \leq \underline{u} \leq \underline{u}_2$.

3. We have

$$\lambda^{-1}e'_4(\underline{u}) = e_4(\underline{u}) + \frac{1}{4}f^2e_3(\underline{u}) = \frac{2}{\zeta}.$$

Relying on our control of the ingoing geodesic foliation of ${}^{(int)}\mathcal{M}$, the above assumption for f and the transport equation for f , we obtain on $\gamma(\underline{u}_1, \underline{u}_2) \cap {}^{(int)}\mathcal{M}$

$$\begin{aligned} \sup_{\gamma(\underline{u}_1, \underline{u}_2) \cap {}^{(int)}\mathcal{M}} |f| &\lesssim \frac{\epsilon_0}{\underline{u}_1^{1 + \delta_{dec}}} (\underline{u}_2 - \underline{u}_1) \\ &\lesssim \frac{\epsilon_0^{1 - \frac{\delta_{dec}}{2}}}{\underline{u}_1^{1 + \frac{\delta_{dec}}{2}}} \end{aligned}$$

which improves our assumption in (3.6.9) on f .

4. We have in view of the control of f

$$\lambda^{-1}e'_4(r) = e_4(r) + \frac{1}{4}f^2e_3(r) = \Upsilon + O\left(\frac{\epsilon_0}{\underline{u}_1^{1 + \delta_{dec}}}\right).$$

This yields

$$\begin{aligned} \lambda^{-1}e'_4(\log(\Upsilon)) &= \frac{\frac{2m}{r^2}e_4(r) - \frac{2}{r}\lambda^{-1}e'_4(m)}{\Upsilon} \\ &= \frac{\frac{2m}{r^2}\Upsilon + O\left(\frac{\epsilon_0}{\underline{u}_1^{1 + \delta_{dec}}}\right)}{\Upsilon}. \end{aligned}$$

Thanks to our assumption on the lower bound of Υ , we infer

$$\lambda^{-1}e'_4(\log(\Upsilon)) = \frac{2m}{r^2}(1 + O(\sqrt{\epsilon_0}))$$

and since we are in ${}^{(int)}\mathcal{M}$

$$\lambda^{-1}e'_4(\log(\Upsilon)) \geq \frac{1}{3m_0}.$$

Integrating from $\underline{u} = \underline{u}_1$, we deduce

$$\Upsilon \geq \frac{\sqrt{\epsilon_0}}{(1 + \sqrt{\epsilon_0})\underline{u}_1^{1+\frac{\delta_{dec}}{2}}} \exp\left(\frac{\underline{u} - \underline{u}_1}{3m_0}\right)$$

which is an improvement of our assumption in (3.6.9) on Υ .

5. In view of the control of f and of the ingoing geodesic foliation of ${}^{(int)}\mathcal{M}$, we rewrite the transport equation for λ as

$$\begin{aligned} \lambda^{-1}e'_4(\log(\lambda)) &= 2\omega + E_2(f, \Gamma) \\ &= -\frac{2m}{r^2} + O\left(\frac{\epsilon_0}{\underline{u}_1^{1+\delta_{dec}}}\right). \end{aligned}$$

Since we have obtained above the other hand

$$\lambda^{-1}e'_4(\log(\Upsilon)) = \frac{2m}{r^2}(1 + O(\sqrt{\epsilon_0}))$$

we immediately infer

$$\lambda^{-1}e'_4(\log(\lambda)\Upsilon^2) > 0, \quad \lambda^{-1}e'_4(\log(\lambda)\sqrt{\Upsilon}) < 0.$$

Integrating from $\underline{u} = \underline{u}_1$, this yields

$$\left(\frac{\sqrt{\epsilon_0}}{(1 + \sqrt{\epsilon_0})\underline{u}_1^{1+\frac{\delta_{dec}}{2}}}\right)^2 \Upsilon^{-2} \leq \lambda \leq \left(\frac{\sqrt{\epsilon_0}}{(1 + \sqrt{\epsilon_0})\underline{u}_1^{1+\frac{\delta_{dec}}{2}}}\right)^{\frac{1}{2}} \Upsilon^{-\frac{1}{2}}.$$

Since Υ has an explicit lower bounded in view of our previous estimate, as well as an explicit upper bound since we are in ${}^{(int)}\mathcal{M}$, this yields an improvement of our assumptions in (3.6.9) for λ .

6. Since we have improved all our bootstrap assumptions (3.6.9), we infer by a continuity argument the following bound

$$\Upsilon \geq \frac{\sqrt{\epsilon_0}}{(1 + \sqrt{\epsilon_0})\underline{u}_1^{1+\frac{\delta_{dec}}{2}}} \exp\left(\frac{\underline{u} - \underline{u}_1}{3m_0}\right) \text{ on } \gamma\left(\underline{u}_1, \underline{u}_1 + \left(\frac{\underline{u}_1}{\epsilon_0}\right)^{\frac{\delta_{dec}}{2}}\right) \cap {}^{(int)}\mathcal{M}.$$

Now, in this \underline{u} interval, we may choose

$$\underline{u}_3 := \underline{u}_1 + 3m_0 \left(1 + \frac{\delta_{dec}}{2}\right) \log\left(\frac{\underline{u}_1}{\epsilon_0}\right)$$

for which we have $\Upsilon \geq 1$. This is a contradiction since $\Upsilon = O(\delta_{\mathcal{H}})$ in ${}^{(int)}\mathcal{M}$. Thus, we deduce that γ reaches ${}^{(ext)}\mathcal{M}$ before $\underline{u} = \underline{u}_3$, a contradiction to our assumption on γ . This concludes the proof of (3.6.7).

3.7 The General Covariant Modulation procedure

The role of this section is to give a short description of the results concerning our General Covariant Modulation (GCM) procedure, which is at the heart of our proof. We will apply it in $^{(ext)}\mathcal{M}$ under our main bootstrap assumptions **BA-E BA-D**. The proof of the results stated in this section will be proved in Chapter 9.

3.7.1 Spacetime assumptions for the GCM procedure

To state our results, which are local in nature it is convenient to consider axially symmetric polarized spacetime regions \mathcal{R} foliated by two functions (u, s) such that

- On \mathcal{R} , (u, s) defines an outgoing geodesic foliation as in section 2.2.4.
- We denote by (e_3, e_4, e_θ) the null frame adapted to the outgoing geodesic foliation (u, s) on \mathcal{R} .
- We denote by $\overset{\circ}{S}$ a fixed sphere of \mathcal{R}

$$\overset{\circ}{S} := S(\overset{\circ}{u}, \overset{\circ}{s}) \quad (3.7.1)$$

and by $\overset{\circ}{r}$ the area radius of $\overset{\circ}{S}$, where $S(u, s)$ denote the 2-spheres of the outgoing geodesic foliation (u, s) on \mathcal{R} .

- In adapted coordinates (u, s, θ, φ) with $b = 0$, see Proposition 2.2.20, the spacetime metric \mathbf{g} in \mathcal{R} takes the form, with $\underline{\Omega} = e_3(s)$, $\underline{b} = e_3(\theta)$,

$$\mathbf{g} = -2\zeta dud s + \zeta^2 \underline{\Omega} du^2 + \gamma \left(d\theta - \frac{1}{2} \zeta \underline{b} du \right)^2 + e^{2\Phi} d\varphi^2, \quad (3.7.2)$$

where θ is chosen such that $b = e_4(\theta) = 0$.

- The spacetime metric induced on $S(u, s)$ is given by,

$$\mathcal{g} = \gamma d\theta^2 + e^{2\Phi} d\varphi^2. \quad (3.7.3)$$

- The relation between the null frame and coordinate system is given by

$$e_4 = \partial_s, \quad e_3 = \frac{2}{\zeta} \partial_u + \underline{\Omega} \partial_s + \underline{b} \partial_\theta, \quad e_\theta = \gamma^{-1/2} \partial_\theta. \quad (3.7.4)$$

- We denote the induced metric on \mathring{S} by

$$\mathring{g} = \mathring{\gamma} d\theta^2 + e^{2\Phi} d\varphi^2.$$

Definition 3.7.1. Let $0 < \delta \leq \mathring{\epsilon}$ two sufficiently small constants. Let $(\mathring{u}, \mathring{s})$ real numbers so that

$$1 \leq \mathring{u} < +\infty, \quad 4m_0 \leq \mathring{s} < +\infty. \quad (3.7.5)$$

We define $\mathcal{R} = \mathcal{R}(\delta, \mathring{\epsilon})$ to be the region

$$\mathcal{R} := \left\{ |u - \mathring{u}| \leq \delta_{\mathcal{R}}, \quad |s - \mathring{s}| \leq \delta_{\mathcal{R}} \right\}, \quad \delta_{\mathcal{R}} := \delta (\mathring{\epsilon})^{-\frac{1}{2}}, \quad (3.7.6)$$

such that assumption **A1-A3** below with constant $\mathring{\epsilon}$ on the background foliation of \mathcal{R} , are verified. The smaller constant δ controls the size of the GCMS quantities as it will be made precise below.

Consider the renormalized Ricci and curvature components associated to the (u, s) geodesic foliation of \mathcal{R}

$$\begin{aligned} \check{\Gamma} &:= \left\{ \check{\kappa}, \vartheta, \zeta, \eta, \bar{\kappa} - \frac{2}{r}, \bar{\kappa} + \frac{2\Upsilon}{r}, \check{\kappa}, \vartheta, \xi, \check{\omega}, \bar{\omega} - \frac{m}{r^2}, \check{\Omega}, (\bar{\Omega} + \Upsilon), (\bar{\varsigma} + 1) \right\}, \\ \check{R} &:= \left\{ \alpha, \beta, \check{\rho}, \bar{\rho} + \frac{2m}{r^3}, \underline{\beta}, \alpha \right\}. \end{aligned}$$

Since our foliation is outgoing geodesic we also have,

$$\xi = \omega = 0, \quad \underline{\eta} + \zeta = 0. \quad (3.7.7)$$

We decompose $\check{\Gamma} = \Gamma_g \cup \Gamma_b$ where,

$$\begin{aligned} \Gamma_g &= \left\{ \check{\kappa}, \vartheta, \zeta, \check{\kappa}, \bar{\kappa} - \frac{2}{r}, \bar{\kappa} + \frac{2\Upsilon}{r} \right\}, \\ \Gamma_b &= \left\{ \eta, \vartheta, \xi, \check{\omega}, \bar{\omega} - \frac{m}{r^2}, r^{-1}\check{\Omega}, r^{-1}\check{\zeta}, r^{-1}(\bar{\Omega} + \Upsilon), r^{-1}(\bar{\varsigma} - 1) \right\}. \end{aligned} \quad (3.7.8)$$

Given an integer s_{max} , we assume the following¹⁶

¹⁶In applications, $s_{max} = k_{small} + 4$ in Theorem M7, and $s_{max} = k_{large} + 5$ in Theorem M0 and Theorem M6.

A1. For $k \leq s_{max}$, we have on \mathcal{R}

$$\begin{aligned} \|\Gamma_g\|_{k,\infty} &\lesssim \mathring{\epsilon} r^{-2}, \\ \|\Gamma_b\|_{k,\infty} &\lesssim \mathring{\epsilon} r^{-1}, \end{aligned} \quad (3.7.9)$$

and,

$$\begin{aligned} \|\alpha, \beta, \check{\rho}, \check{\mu}\|_{k,\infty} &\lesssim \mathring{\epsilon} r^{-3}, \\ \|e_3(\alpha, \beta)\|_{k-1,\infty} &\lesssim \mathring{\epsilon} r^{-4}, \\ \|\underline{\beta}\|_{k,\infty} &\lesssim \mathring{\epsilon} r^{-2}, \\ \|\underline{\alpha}\|_{k,\infty} &\lesssim \mathring{\epsilon} r^{-1}. \end{aligned} \quad (3.7.10)$$

A2. We have, with m_0 denoting the mass of the unperturbed spacetime,

$$\sup_{\mathcal{R}} \left| \frac{m}{m_0} - 1 \right| \lesssim \mathring{\epsilon}. \quad (3.7.11)$$

A3. The metric coefficients are assumed to satisfy the following assumptions in \mathcal{R} , for all $k \leq s_{max}$

$$r \left\| \left(\frac{\gamma}{r^2} - 1, \underline{b}, \frac{e^\Phi}{r \sin \theta} - 1 \right) \right\|_{\infty,k} + \|\underline{\Omega} + \Upsilon\|_{\infty,k} + \|\varsigma - 1\|_{\infty,k} \lesssim \mathring{\epsilon} \quad (3.7.12)$$

We will assume, in addition, that the following small GCM conditions hold true on \mathcal{R} ,

$$\left| \bar{\kappa} - \frac{2}{r} \right| + |\mathfrak{d}^k \check{\kappa}| + r^2 |\mathfrak{d}^{k-2} \mathfrak{d}_2^* \mathfrak{d}_1^* \check{\kappa}| + r^3 |\mathfrak{d}^{k-2} \mathfrak{d}_2^* \mathfrak{d}_1^* \mu| \lesssim \mathring{\delta} r^{-2} \text{ for all } k \leq s_{max}, \quad (3.7.13)$$

$$r^{-2} \left| \int_S \eta e^\Phi \right| \lesssim \mathring{\delta}, \quad r^{-2} \left| \int_S \underline{\xi} e^\Phi \right| \lesssim \mathring{\delta}. \quad (3.7.14)$$

Also,

$$\left| \left(\frac{2}{\varsigma} + \underline{\Omega} \right) \Big|_{SP} - 1 - \frac{2m}{r} \right| \lesssim \mathring{\delta}. \quad (3.7.15)$$

Additionally we may assume on \mathcal{R}

$$r \left| \int_S \beta e^\Phi \right| \lesssim \mathring{\delta}, \quad \left| \int_S e_\theta(\underline{\kappa}) e^\Phi \right| \lesssim \mathring{\delta}. \quad (3.7.16)$$

3.7.2 Deformations of surfaces

Definition 3.7.2. We say that \mathbf{S} is an $O(\epsilon)$ \mathbf{Z} -polarized deformation of \mathring{S} if there exists a map $\Psi : \mathring{S} \rightarrow \mathbf{S}$ of the form,

$$\Psi(\mathring{u}, \mathring{s}, \theta, \varphi) = \left(\mathring{u} + U(\theta), \mathring{s} + S(\theta), \theta, \varphi \right) \quad (3.7.17)$$

where U, S are smooth functions defined on the interval $[0, \pi]$ of amplitude at most ϵ . We denote by ψ the reduce map defined on the interval $[0, \pi]$,

$$\psi(\theta) = (\mathring{u} + U(\theta), \mathring{s} + S(\theta), \theta). \quad (3.7.18)$$

We restrict ourselves to deformations which fix the South Pole, i.e.

$$U(0) = S(0) = 0. \quad (3.7.19)$$

3.7.3 Adapted frame transformations

We consider general null transformations introduced in Lemma 2.3.1,

$$\begin{aligned} e'_4 &= \lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right), \\ e'_\theta &= \left(1 + \frac{1}{2} f \underline{f} \right) e_\theta + \frac{1}{2} \underline{f} e_4 + \frac{1}{2} f \left(1 + \frac{1}{4} f \underline{f} \right) e_3, \\ e'_3 &= \lambda^{-1} \left(\left(1 + \frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) e_3 + \underline{f} \left(1 + \frac{1}{4} f \underline{f} \right) e_\theta + \frac{1}{4} \underline{f}^2 e_4 \right). \end{aligned} \quad (3.7.20)$$

Definition 3.7.3. Given a deformation $\Psi : \mathring{S} \rightarrow \mathbf{S}$ we say that a new frame (e'_3, e'_4, e'_θ) , obtained from the standard frame (e_3, e_4, e_θ) via the transformation (3.7.20), is \mathbf{S} -adapted if we have,

$$e'_\theta = e_\theta^{\mathbf{S}} = \frac{1}{(\gamma^{\mathbf{S}})^{1/2}} \psi_{\#}(\partial_\theta) \quad (3.7.21)$$

where $\psi_{\#}(\partial_\theta)$ is the push-forward defined by the deformation map ψ .

The condition translates into the following relations between the functions U, S defining

the deformation and the transition functions (f, \underline{f}) .

$$\begin{aligned} \varsigma^\# \partial_\theta U &= ((\gamma^\mathbf{S})^\#)^{1/2} f^\# \left(1 + \frac{1}{4} (f \underline{f})^\# \right), \\ \partial_\theta S - \frac{\varsigma^\#}{2} \underline{\Omega}^\# \partial_\theta U &= \frac{1}{2} ((\gamma^\mathbf{S})^\#)^{1/2} \underline{f}^\#, \\ (\gamma^\mathbf{S})^\# &= \gamma^\# + (\varsigma^\#)^2 \left(\underline{\Omega} + \frac{1}{4} \underline{b}^2 \gamma \right)^\# (\partial_\theta U)^2 - 2\varsigma^\# \partial_\theta U \partial_\theta S - (\gamma \underline{\varsigma} \underline{b})^\# \partial_\theta U, \\ U(0) &= S(0) = 0. \end{aligned} \tag{3.7.22}$$

3.7.4 GCM results

Theorem 3.7.4 (GCMS-I). *Consider the region \mathcal{R} as above, verifying the assumptions **A1–A3** and the small GCM conditions¹⁷ (3.7.13). Let \mathring{S} denote the sphere $\mathring{S} = S(\mathring{u}, \mathring{s})$. For any fix $\Lambda, \underline{\Lambda} \in \mathbb{R}$ verifying,*

$$|\Lambda|, |\underline{\Lambda}| \lesssim \delta(\mathring{r})^2, \tag{3.7.23}$$

1. *There exists a unique GCM sphere $\mathbf{S} = \mathbf{S}^{(\Lambda, \underline{\Lambda})}$, which is a deformation¹⁸ of \mathring{S} , and an adapted null frame $e_3^\mathbf{S}, e_\theta^\mathbf{S}, e_4^\mathbf{S}$, such that the following GCMS conditions are verified¹⁹ components.*

$$\not{d}_2^{\mathbf{S}, \star} \not{d}_1^{\mathbf{S}, \star} \underline{\kappa}^\mathbf{S} = \not{d}_2^{\mathbf{S}, \star} \not{d}_1^{\mathbf{S}, \star} \mu^\mathbf{S} = 0, \quad \kappa^\mathbf{S} = \frac{2}{r^\mathbf{S}}. \tag{3.7.24}$$

In addition

$$\int_{\mathbf{S}} f e^\Phi = \Lambda, \quad \int_{\mathbf{S}} \underline{f} e^\Phi = \underline{\Lambda}, \tag{3.7.25}$$

where (f, \underline{f}) belong to the triplet $(f, \underline{f}, \lambda = e^a)$ which denote the change of frame coefficients from the frame of \mathring{S} to the one of \mathbf{S} .

2. *The transition functions $(f, \underline{f}, \log \lambda)$ verify,*

$$\| (f, \underline{f}, \log \lambda) \|_{\mathfrak{h}_k(\mathbf{S})} \lesssim \delta, \quad k \leq s_{max} + 1. \tag{3.7.26}$$

¹⁷Here, the other assumptions (3.7.14) (3.7.15) are not needed.

¹⁸In the sense of Definition 3.7.2.

¹⁹ $\Gamma^\mathbf{S}, R^\mathbf{S}$ denote the Ricci and curvature components with respect to the adapted frame on \mathbf{S} .

3. The area radius $r^{\mathbf{S}}$ and Hawking mass $m^{\mathbf{S}}$ of \mathbf{S} verify,

$$|r^{\mathbf{S}} - \overset{\circ}{r}| \lesssim \delta, \quad |m^{\mathbf{S}} - \overset{\circ}{m}| \lesssim \delta. \quad (3.7.27)$$

The precise version of Theorem 3.7.4 and its proof are given in section 9.4.

Theorem 3.7.5 (GCMS-II). *In addition to the assumptions of Theorem 3.7.4 we also assume that (3.7.16) holds true. Then,*

1. *There exists a unique GCM sphere \mathbf{S} , which is a deformation of $\overset{\circ}{S}$, such that in addition to (3.7.24) the following GCMS conditions also hold true on \mathbf{S} .*

$$\int_{\mathbf{S}} \beta^{\mathbf{S}} e^{\Phi} = 0, \quad \int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}}) e^{\Phi} = 0. \quad (3.7.28)$$

2. *The transition functions $(f, \underline{f}, \log \lambda)$ verify the estimates (3.7.26).*

3. *The area radius $r^{\mathbf{S}}$ and Hawking mass $m^{\mathbf{S}}$ of \mathbf{S} verify (3.7.27).*

The precise version of Theorem 3.7.5 and its proof are given in 9.7.

Theorem 3.7.6 (GCMH). *Consider the region \mathcal{R} as above, verifying the assumptions A1–A3 and the small GCM conditions (3.7.13)–(3.7.15). Let $\mathbf{S}_0 = \mathbf{S}_0[\overset{\circ}{u}, \overset{\circ}{s}, \Lambda_0, \underline{\Lambda}_0]$ the deformation of $\overset{\circ}{S}$ constructed in Theorem GCMS-I above.*

There exists a smooth spacelike hypersurface $\Sigma_0 \subset \mathcal{R}$ passing through \mathbf{S}_0 , a scalar function $u^{\mathbf{S}}$ defined on Σ_0 , whose level surfaces are topological spheres denoted by \mathbf{S} , and a smooth collection of constants $\Lambda^{\mathbf{S}}, \underline{\Lambda}^{\mathbf{S}}$ verifying,

$$\Lambda^{\mathbf{S}_0} = \Lambda_0, \quad \underline{\Lambda}^{\mathbf{S}_0} = \underline{\Lambda}_0,$$

such that the following conditions are verified:

1. *The surfaces \mathbf{S} of constant $u^{\mathbf{S}}$ verifies all the properties stated in Theorem GCMS-I for the prescribed constants $\Lambda^{\mathbf{S}}, \underline{\Lambda}^{\mathbf{S}}$. In particular they come endowed with null frames $(e_4^{\mathbf{S}}, e_{\theta}^{\mathbf{S}}, e_3^{\mathbf{S}})$ such that*

i. For each \mathbf{S} the GCM conditions (3.7.24), (3.7.25) hold with $\Lambda = \Lambda^{\mathbf{S}}, \underline{\Lambda} = \underline{\Lambda}^{\mathbf{S}}$.

iii. The transversality conditions hold true on each \mathbf{S} .

$$\xi^{\mathbf{S}} = 0, \quad \omega^{\mathbf{S}} = 0, \quad \underline{\eta}^{\mathbf{S}} + \zeta^{\mathbf{S}} = 0. \quad (3.7.29)$$

2. We have, for some constant c_{Σ_0} ,

$$u^{\mathbf{S}} + r^{\mathbf{S}} = c_{\Sigma_0}, \quad \text{along } \Sigma_0. \quad (3.7.30)$$

3. Let $\nu^{\mathbf{S}}$ be the unique vectorfield tangent to the hypersurface Σ_0 , normal to \mathbf{S} , and normalized by $g(\nu^{\mathbf{S}}, e_4^{\mathbf{S}}) = -2$. There exists a unique scalar function $a^{\mathbf{S}}$ on Σ_0 such that $\nu^{\mathbf{S}}$ is given by

$$\nu^{\mathbf{S}} = e_3^{\mathbf{S}} + a^{\mathbf{S}} e_4^{\mathbf{S}}.$$

The following normalization condition holds true at the South Pole SP of every sphere \mathbf{S} , i.e. at $\theta = 0$,

$$a^{\mathbf{S}} \Big|_{SP} = -1 - \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}. \quad (3.7.31)$$

4. Under the additional transversality condition²⁰ on Σ_0

$$e_4^{\mathbf{S}}(u^{\mathbf{S}}) = 0, \quad e_4(r^{\mathbf{S}}) = \frac{r^{\mathbf{S}}}{2} \kappa^{\mathbf{S}} = 1. \quad (3.7.32)$$

the Ricci coefficients $\eta^{\mathbf{S}}, \xi^{\mathbf{S}}$ are well defined and verify,

$$\int_{\mathbf{S}} \eta^{\mathbf{S}} e^{\Phi} = \int_{\mathbf{S}} \xi^{\mathbf{S}} e^{\Phi} = 0. \quad (3.7.33)$$

5. The transition functions $(f, \underline{f}, \log \lambda)$ verify the estimates (3.7.26).

6. The area radius $r^{\mathbf{S}}$ and Hawking mass $m^{\mathbf{S}}$ of \mathbf{S} verify (3.7.27).

The precise version of Theorem 3.7.6 and its proof are given in section 9.8.

In addition to the three theorems stated above we also prove three rigidity statements which can be regarded as corollaries.

Corollary 3.7.7 (Rigidity I). *Assume \mathbf{S} is a sphere in \mathcal{R} which verifies the the GCM conditions*

$$\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}, \quad \not{d}_2^{\mathbf{S}} \not{d}_1^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}} = \not{d}_2^{\mathbf{S}} \not{d}_1^{\mathbf{S}} \mu^{\mathbf{S}} = 0. \quad (3.7.34)$$

²⁰Here the average of $\kappa^{\mathbf{S}}$ is taken on \mathbf{S} . In view of the GCM conditions (3.7.24) we deduce $e_4^{\mathbf{S}}(r^{\mathbf{S}}) = 1$.

Assume also that the background foliation verify on \mathcal{R} the conditions **A1-A3**. Then the transition functions $(f, \underline{f}, \log \lambda)$ from the background frame of \mathcal{R} to that of \mathbf{S} verifies the estimates

$$\|(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}_{s_{max}+1}(\mathbf{S})} \lesssim \overset{\circ}{\delta} + \left| \int_{\mathbf{S}} f e^{\Phi} \right| + \left| \int_{\mathbf{S}} \underline{f} e^{\Phi} \right|.$$

Moreover, The area radius $r^{\mathbf{S}}$ and Hawking mass $m^{\mathbf{S}}$ of \mathbf{S} verify (3.7.27).

Corollary 3.7.8 (Rigidity II). Assume given a sphere $\mathbf{S} \subset \mathcal{R}$ endowed with a compatible frame $e_3^{\mathbf{S}}, e_{\theta}^{\mathbf{S}}, e_4^{\mathbf{S}}$ which verifies the GCM conditions

$$\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}, \quad \not\phi_2^{\mathbf{S}} \not\phi_1^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}} = \not\phi_2^{\mathbf{S}} \not\phi_1^{\mathbf{S}} \mu^{\mathbf{S}} = 0. \quad (3.7.35)$$

In addition, we assume

1. We have,

$$r \left| \int_{\mathbf{S}} \beta^{\mathbf{S}} e^{\Phi} \right| + \left| \int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}}) e^{\Phi} \right| \lesssim \overset{\circ}{\delta}. \quad (3.7.36)$$

2. The background foliation verifies on \mathcal{R} the conditions **A1-A3**.

Then the transition functions $(f, \underline{f}, \log \lambda)$ from the background frame of \mathcal{R} to that of \mathbf{S} verifies the estimates

$$\|(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}_{s_{max}+1}(\mathbf{S})} \lesssim \overset{\circ}{\delta}.$$

Moreover, the area radius $r^{\mathbf{S}}$ and Hawking mass $m^{\mathbf{S}}$ of \mathbf{S} verify (3.7.27).

Corollary 3.7.9 (Rigidity III). Assume given a GCM hypersurface $\Sigma_0 \subset \mathcal{R}$ foliated by surfaces \mathbf{S} such that

$$\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}, \quad \not\phi_2^{\mathbf{S}} \not\phi_1^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}} = \not\phi_2^{\mathbf{S}} \not\phi_1^{\mathbf{S}} \mu^{\mathbf{S}} = 0, \quad \int_{\mathbf{S}} \eta^{\mathbf{S}} e^{\Phi} = 0, \quad \int_{\mathbf{S}} \underline{\xi}^{\mathbf{S}} e^{\Phi} = 0.$$

Assume in addition that for a specific sphere \mathbf{S}_0 on Σ_0 we have the following additional information:

1. The transition functions f, \underline{f} from the background foliation to \mathbf{S}_0 verify

$$\int_{\mathbf{S}} f e^{\Phi} = O(\overset{\circ}{\delta}), \quad \int_{\mathbf{S}} \underline{f} e^{\Phi} = O(\overset{\circ}{\delta}). \quad (3.7.37)$$

2. In \mathcal{R} , the Ricci and curvature coefficients of the background foliation verify the assumptions **A1–A3**.

Then, for all derivatives of the transition functions along \mathbf{S} ,

$$\|\mathfrak{d}^{\leq s_{max}+1}(f, \underline{f}, \log \lambda)\|_{L^2(\mathbf{S})} \lesssim \overset{\circ}{\delta}.$$

Moreover, The area radius $r^{\mathbf{S}}$ and Hawking mass $m^{\mathbf{S}}$ of \mathbf{S} verify (3.7.27).

3.7.5 Main ideas

Both theorems GCMS-II and GCMH are based on Theorem GSMS-I. They are heavily based on the transformation formulas for the Ricci and curvature coefficients recorded in Proposition 2.3.4.

Sketch of the proof of Theorems GSMS-I and GSMS-II

A given deformation $\Psi : \overset{\circ}{S} \rightarrow \mathbf{S}$ is fixed by the parameters U, S and transition functions $F = (f, \underline{f}, \lambda)$ connected by the system (3.7.22). Making use of the transformation formulas one can show that the GCMS conditions (3.7.24)-(3.7.25) are if and only if the transition functions F verify a coercive nonlinear elliptic Hodge system of the form $\mathcal{D}_\Psi F = B(\Psi)$, where the operator \mathcal{D}_Ψ depends on the deformation Ψ and the right hand side B , depends on both Ψ and the background foliation (see Proposition 9.4.1 for the precise form of the system). To find a desired GSMS deformation we have to solve a coupled system between the transport type equations in (3.7.22) and the elliptic coercive system $\mathcal{D}_\Psi F = 0$ of Proposition 9.4.1.

The actual proof is thus based on an iteration procedure for a sequence of deformation spheres $\mathbf{S}(n)$ of $\overset{\circ}{S}$ given by the maps $\Psi^{(n)} = (U^{(n)}, S^{(n)}) : \overset{\circ}{S} \rightarrow \mathbf{S}(n)$ and the corresponding transition functions $\log \lambda^{(n)}, f^{(n)}, \underline{f}^{(n)}$. The iteration procedure for the quintets $Q^{(n)} = (U^{(n)}, S^{(n)}, \log \lambda^{(n)}, f^{(n)}, \underline{f}^{(n)})$, starting with the trivial quintet $Q^{(0)}$ corresponding to the zero deformation, is described in section 9.4.3. The main steps in the proof are as follows.

1. Given the triplet $(\log \lambda^{(n)}, f^{(n)}, \underline{f}^{(n)})$ the pair $(U^{(n)}, S^{(n)})$ defines the deformation sphere $\mathbf{S}(n)$ and the corresponding pull back map $\#_n : \overset{\circ}{S} \rightarrow \mathbf{S}(n)$ according to the equation (3.7.22).

2. Given the pair $\Psi^{(n)} = (U^{(n)}, S^{(n)})$ and the deformation sphere $\mathbf{S}(n)$ we define the triplet $(\log \lambda^{(n+1)}, f^{(n+1)}, \underline{f}^{(n+1)})$ by solving the corresponding elliptic system

$$\mathcal{D}_{\Psi^{(n)}} F^{(n+1)} = B(\Psi^{(n)})$$

This step is based on the crucial apriori estimates of section 9.4.1.

3. Given the new pair $(f^{(n+1)}, \underline{f}^{(n+1)})$ we make use of the equations (3.7.22) to find a unique new map $(U^{(n+1)}, S^{(n+1)})$ and thus the new deformation sphere $\mathbf{S}(n+1)$.
4. The convergence of the iterates $Q^{(n)}$, described in subsection 9.4.5 in the boundedness Proposition 9.4.8 and the contraction Proposition 9.4.9. The latter requires us to carefully compare the iterates $Q^{(n)}, Q^{(n+1)}$ by pulling them back to $\overset{\circ}{S}$. One has to be particularly careful with the behavior of the iterates on the axis of symmetry.

Theorem GSMS-II, which is an easy consequence of Theorem GSMS-I is proved in section 9.7 and the transformation formulas which relate $\int_{\mathbf{S}} \beta^{\mathbf{S}} e^{\Phi}$ to $\Lambda = \int_{\mathbf{S}} f e^{\Phi}$ and $\int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}}) e^{\Phi}$ to $\underline{\Lambda} = \int_{\mathbf{S}} \underline{f} e^{\Phi}$. One can show that there exist choices of $\Lambda, \underline{\Lambda}$ such that $\int_{\mathbf{S}} \beta^{\mathbf{S}} e^{\Phi} = \int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}}) e^{\Phi} = 0$.

Sketch of the proof of Theorem GCMH

The proof of Theorem GCMH makes use of Theorem GCMS-I to construct Σ_0 as a union of GCMS spheres.

Step 1. Theorem GCMS-I allows to construct, for every value of the parameters (u, s) in \mathcal{R} (i.e. such that the background spheres $S(u, s) \subset \mathcal{R}$) and every real numbers $(\Lambda, \underline{\Lambda})$, a unique GCM sphere $\mathbf{S}[u, s, \Lambda, \underline{\Lambda}]$, as a \mathbf{Z} -polarized deformation of $S(u, s)$. In particular (3.7.24) and (3.7.25) are verified and $\mathbf{S}_0 = \mathbf{S}_0[\overset{\circ}{u}, \overset{\circ}{s}, \Lambda_0, \underline{\Lambda}_0]$.

Step 2. We look for functions $\Psi(s), \Lambda(s), \underline{\Lambda}(s)$ such that

1. We have,

$$\Psi(\overset{\circ}{s}) = \overset{\circ}{u}, \quad \Lambda(\overset{\circ}{s}) = \Lambda_0, \quad \underline{\Lambda}(\overset{\circ}{s}) = \underline{\Lambda}_0.$$

2. The resulting hypersurface $\Sigma_0 = \cup_s \mathbf{S}[\Psi(s), s, \Lambda(s), \underline{\Lambda}(s)]$ verifies

$$u^{\mathbf{S}} + r^{\mathbf{S}} = c_{\Sigma_0}, \quad \text{along } \Sigma_0.$$

3. The additional GCM conditions (3.7.31) and (3.7.33) of Theorem GCMH are verified.

These conditions lead to a first order differential system for $\Psi(s), \Lambda(s), \underline{\Lambda}(s)$, with prescribed initial conditions at $\overset{\circ}{s}$ which allows us to determine the desired surface. The proof is given in detail in section 9.8.

Sketch of the proof of the Rigidity Corollaries

The Rigidity I corollary, see Corollary 9.7.3, is proved in exactly the same way as Proposition 9.4.4 which is one of the main steps in the proof of Theorem GCMS-I. The rigidity II corollary, see Corollary 9.7.3, is based on Rigidity I and a simple variation of Lemma 9.7.2. The proof of the rigidity III corollary, see Corollary 9.8.2, is essentially part of the proof of Theorem GCMH, once the existence part of the theorem has been established.

3.8 Overview of the proof of Theorem M0-M8

In this section, we provide a brief overview of the proof of Theorem M0-M8. In addition to the null frame adapted to the outgoing foliation of $^{(ext)}\mathcal{M}$ and to the null frame adapted to the ingoing foliation of $^{(int)}\mathcal{M}$, we have also introduced 2 global frames on $\mathcal{M} = ^{(int)}\mathcal{M} \cup ^{(ext)}\mathcal{M}$ as well as associated scalars r and m in section 3.5. Unless otherwise specified, when we discuss a particular spacetime region, i.e. $^{(ext)}\mathcal{M}$, $^{(int)}\mathcal{M}$ or \mathcal{M} , it should be understood that the frame as well as r and m are the ones corresponding to that region.

3.8.1 Discussion of Theorem M0

Step 1. Recall our GCM conditions on $S_* = \Sigma_* \cap \mathcal{C}_*$

$$\int_{S_*} e_{\theta}(\underline{\kappa})e^{\Phi} = 0, \quad \int_{S_*} \beta e^{\Phi} = 0.$$

Recall that $\nu = e_3 + a_*e_4$ is the unique tangent vectorfield to Σ_* which is orthogonal to e_{θ} and normalized by $\mathbf{g}(\nu, e_4) = -2$. Using the null structure equation for $e_3(\underline{\kappa})$ and $e_3(\beta)$, as well as $e_4(\underline{\kappa})$ and $e_4(\beta)$, we obtain transport equations along Σ_* in the ν direction for

$$\int_S e_{\theta}(\underline{\kappa})e^{\Phi} \text{ and } \int_S \beta e^{\Phi} = 0.$$

Integrating these transport equations in ν , we propagate the control on S_* to Σ_* . In particular, we obtain the following estimates on $S_1 = \Sigma_* \cap \mathcal{C}_1$,

$$\left| \int_{S_1} e_\theta(\underline{\kappa}) e^\Phi \right| + \left| \int_{S_1} \beta e^\Phi \right| \lesssim \epsilon^2 + \frac{\epsilon}{r} \lesssim \epsilon_0, \quad (3.8.1)$$

where we used in the last inequality the dominance condition of r on Σ_* , see (3.3.4).

Step 2. We consider the transition functions $(f, \underline{f}, \lambda)$ from the frame of the initial data layer to the frame of ${}^{(ext)}\mathcal{M}$. Since

- S_1 is a sphere of ${}^{(ext)}\mathcal{M}$ in the initial data layer,
- S_1 is a sphere of the GCM hypersurface Σ_* ,
- the estimate (3.8.1) holds on S_1 ,

we can invoke the GCM Corollary Rigidity II and III of section 3.7.4 which yields, together with a Sobolev embedding on S_1

$$\sup_{S_1} \left(r |\mathfrak{d}^{\leq k_{large}+1}(f, \underline{f}, \log \lambda)| + |m - m_0| \right) \lesssim \epsilon_0. \quad (3.8.2)$$

Step 3. Relying on the transport equations in e_4 for $(f, \underline{f}, \lambda)$, see Corollary 2.3.7, and Proposition 2.2.16 for m , we propagate (3.8.2) to \mathcal{C}_1 , and then, proceeding similarly, in the e_3 direction to $\underline{\mathcal{C}}_1$ which yields

$$\sup_{\mathcal{C}_1 \cup \underline{\mathcal{C}}_1} \left(r |\mathfrak{d}^{\leq k_{large}+1}(f, \underline{f}, \log \lambda)| + |m - m_0| \right) \lesssim \epsilon_0.$$

Together with the control of the initial data layer foliation and the transformation formulas of Proposition 2.3.4, we then obtain the desired estimates on $\underline{\mathcal{C}}_1 \cup \underline{\mathcal{C}}_1$ for the curvature components.

3.8.2 Discussion of Theorem M1

Here are the main steps in the proof of Theorem M1.

Step 1. Consider the global frame on \mathcal{M} constructed in Proposition 3.5.5 and the definition of \mathfrak{q} on \mathcal{M} with respect to that frame, see section 2.3.3 for the definition of \mathfrak{q} with respect to any null frame. According to Theorem 2.4.7 we have,

$$\square_2 \mathfrak{q} + V \mathfrak{q} = N, \quad V = \kappa \underline{\kappa} \quad (3.8.3)$$

where the nonlinear term $N = \text{Err}[\square_{\mathbf{g}}\mathbf{q}]$ is a long expression of terms quadratic, or higher order, in $\check{\Gamma}, \check{R}$ involving various powers of r . Making use of the symbolic notation introduced in definition 2.3.8 we have, see (2.4.8),

$$\text{Err}[\square_2\mathbf{q}] = r^2\mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta)) + e_3\left(r^3\mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta))\right) + \mathfrak{d}^{\leq 1}(\Gamma_g \cdot \mathbf{q}) + \text{l.o.t.}$$

where the terms denoted by l.o.t. are higher order in $(\check{\Gamma}, \check{R})$.

Remark 3.8.1. *Recall from Remark 2.4.8 that the above good structure of the error term $\text{Err}[\square_2\mathbf{q}]$ only holds in a frame for which $\xi = 0$ for $r \geq 4m_0$ and $\eta \in \Gamma_g$. This is why, in Theorem M1, \mathbf{q} is defined relative to the global frame of Proposition 3.5.5, see also Remark 2.4.9.*

Step 2. We follow the Dafermos-Rodnianski version of the vector-field method to derive desired decay estimates. We recall that, in the context of a wave equation of the form $\square_{(Sch)}\psi = 0$ on Schwarzschild spacetime, their strategy consists in the following:

- Start by deriving Morawetz-energy type estimates for ψ with nondegenerate flux energies and the usual degeneracy of bulk integrals at $r = 3m$.
- Derive r^p weighted estimates for $0 < p < 2$ and use them, in conjunction to the Morawetz estimates, to derive decay estimates.
- The decay estimates obtained by using the standard r^p weighted approach are too weak to be useful in our nonlinear approach. We improve them by making use of a recent variation of the Dafermos-Rodnianski approach due to Angelopoulos, Aretakis and Gajic [4] which is based on first commuting the wave equation is $\square_{(Sch)}\psi = 0$ with $r^2(e_4 + r^{-1})$ and then repeating the process described for the resulting new equation. This procedure allows to derive the improved decay estimates consistent with our decay norms.
- Derive estimates for higher derivatives by commuting with \mathbf{T} , $r\check{\partial}$, the red-shift vectorfield, and re_4 .

Step 3. The estimates mentioned in step 2 have to be adapted to the case of our equation (3.8.3). There are three main differences to take into account

- The application of the vectorfield method to our context produces various nontrivial commutator terms which have to be absorbed. This is taken care by our bootstrap assumption for $\check{\Gamma}, \check{R}$, as well as, in some cases, by integration by parts.

- The presence of the potential V is mostly advantageous but various modifications have to be nevertheless made, especially near the trapping region²¹.
- The presence of the nonlinear term N is the most important complication. The precise null structure of N is essential and various integrations by parts are needed.
- The quadratic terms involving η in N can only be treated provided the definition of \mathfrak{q} is done with respect to the global frame on \mathcal{M} constructed in Proposition 3.5.5, for with η behaves better in powers of r^{-1} .

3.8.3 Discussion of Theorem M2

Recall from section 2.3.3 that \mathfrak{q} is defined with respect to a general null frame as follows

$$\mathfrak{q} = r^4 \left(e_3(e_3(\alpha)) + (2\underline{\kappa} - 6\underline{\omega})e_3(\alpha) + \left(-4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2 \right) \alpha \right)$$

which yields the following transport equation for α

$$e_3(e_3(\alpha)) + (2\underline{\kappa} - 6\underline{\omega})e_3(\alpha) + \left(-4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2 \right) \alpha = \frac{\mathfrak{q}}{r^4}.$$

Recall also that \mathfrak{q} , controlled in Theorem M1, is defined w.r.t. the global frame of Proposition 3.5.5 whose normalization is such that, in particular, $\underline{\omega}$ is a small quantity. Also, since we have

$$e_3(r) = \frac{r}{2}\underline{\kappa} + \text{l.o.t.}$$

we infer

$$e_3(e_3(r^2\alpha)) = \frac{\mathfrak{q}}{r^2} + \text{l.o.t.}$$

Integrating twice this transport equation from \mathcal{C}_1 where we control the initial data - and in particular α - in view of Theorem M0, and using the decay for \mathfrak{q} provided by Theorem M1, we deduce²²

$$\begin{aligned} \sup_{(ext)\mathcal{M}} \left(\frac{r^2(2r+u)^{1+\delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) |\mathfrak{d}^{\leq k_{small}+20}\alpha| &\lesssim \epsilon_0, \\ \sup_{(ext)\mathcal{M}} \left(\frac{r^3(2r+u)^{1+\delta_{extra}}}{\log(1+u)} + r^4(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) |\mathfrak{d}^{\leq k_{small}+19}e_3(\alpha)| &\lesssim \epsilon_0. \end{aligned}$$

²¹At the linear level, on a Schwarzschild spacetime, this step was also treated (minus the improved decay) in the paper [22].

²²Recall that δ_{extra} has been introduced in Theorem M1 and satisfies $\delta_{extra} > \delta_{dec}$.

Now that we control α in the global frame of Proposition 3.5.5, we need to go back to the frame of $^{(ext)}\mathcal{M}$. By invoking the relationships between our various frame of $^{(ext)}\mathcal{M}$, see Proposition 3.5.5 and Proposition 3.4.6, and the transformation formula for α , we infer

$$^{(ext)}\mathcal{D}_{k_{small}+20} [^{(ext)}\alpha] \lesssim \epsilon_0$$

and hence the conclusion of Theorem M2.

3.8.4 Discussion of Theorem M3

Here are the main steps in the proof of Theorem M3.

Step 1. To derive decay estimates for $\underline{\alpha}$ in \mathcal{M} , we first recall the following Teukolsky-Starobinski identity, see (2.3.15),

$$e_3(r^2 e_3(r\mathbf{q})) + 2\underline{\omega}r^2 e_3(r\mathbf{q}) = r^7 \left\{ \not{d}_2^* \not{d}_1^* \not{d}_1 \not{d}_2 \underline{\alpha} + \frac{3}{2} \underline{\kappa} \rho e_4 \underline{\alpha} - \frac{3}{2} \kappa \rho e_3(\underline{\alpha}) \right\} + \text{l.o.t.}$$

where l.o.t. denotes terms which are quadratic or higher, and where all quantities are defined w.r.t. the global frame of Proposition 3.5.5. Then, introducing the vectorfield

$$\tilde{T} = e_4 - \frac{1}{\underline{\kappa}} \left(\bar{\kappa} + \bar{\kappa} \check{\Omega} - \overline{\check{\kappa} \check{\Omega}} \right) e_3,$$

we rewrite the identity as

$$6m\tilde{T}\underline{\alpha} + r^4 \not{d}_2^* \not{d}_1^* \not{d}_1 \not{d}_2 \underline{\alpha} = \frac{1}{r^3} \left(e_3(r^2 e_3(r\mathbf{q})) + 2\underline{\omega}r^2 e_3(r\mathbf{q}) \right) + \text{l.o.t.} \quad (3.8.4)$$

As it turns out, see Remark 6.2.3, this is a forward parabolic equation on each hyper surface of constant r in $^{(int)}\mathcal{M}$.

Step 2. Thanks to

- the control in $^{(int)}\mathcal{M}$ of the RHS of (3.8.4) which follows from the decay estimates of Theorem M1 for \mathbf{q} , as well as the bootstrap assumptions for the quadratic and higher order terms,
- the control of $\underline{\alpha}$ on $\underline{\mathcal{C}}_1$ - i.e. of the initial data of (3.8.4) - provided by Theorem M0,
- parabolic estimates for the forward parabolic equation (3.8.4),

we obtain the desired decay estimates for $\underline{\alpha}$ in $^{(int)}\mathcal{M}$.

Step 3. It remains to control $\underline{\alpha}$ on Σ_* . Recall that ν denotes the unique tangent vectorfield to Σ_* which can be written as $\nu = e_3 + ae_4$. The Teukolsky-Starobinski identity of Step 1 can then be written as

$$6m\nu\underline{\alpha} + r^4 \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \underline{\alpha} = \frac{1}{r^3} \left(e_3(r^2 e_3(r\mathfrak{q})) + 2\underline{\omega} r^2 e_3(r\mathfrak{q}) \right) + \text{l.o.t.} \quad (3.8.5)$$

where l.o.t. denotes terms which are quadratic of higher, as well as terms which are linear but display additional decay in r . This is a forward parabolic equation along Σ_* . To obtain the desired decay for $\underline{\alpha}$ along Σ_* , one then proceeds as in Step 2, using in addition, for the linear term with extra decay in r , the behavior (3.3.4) of r on Σ_* .

3.8.5 Discussion of Theorem M4

Here are the main steps in the proof of Theorem M4.

Step 1. We derive decay estimates for the spacelike GCM hypersurface Σ_* . More precisely, thanks to

- the GCM conditions on Σ_*

$$\kappa = \frac{2}{r}, \quad \not\partial_2^* \not\partial_1^* \underline{\kappa} = 0, \quad \not\partial_2^* \not\partial_1^* \mu = 0, \quad \int_S \eta e^\Phi = 0, \quad \int_S \underline{\xi} e^\Phi = 0,$$

- the control of \mathfrak{q} in $^{(ext)}\mathcal{M}$, established in Theorem M1, and hence in particular on Σ_* ,
- the control of α of the outgoing geodesic foliation in $^{(ext)}\mathcal{M}$, established in Theorem M2, and hence in particular on Σ_* ,
- the control of $\underline{\alpha}$ on Σ_* , established in Theorem M3,
- the fact that the following condition holds on Σ_*

$$r|_{\Sigma_*} \geq \epsilon_0^{-\frac{2}{3}} u^4,$$

- the identity (2.3.11) relating \mathfrak{q} to derivatives of ρ , i.e.

$$\mathfrak{q} = r^4 \left(\not\partial_2^* \not\partial_1^* \rho + \frac{3}{4} \rho \underline{\kappa} \not\vartheta + \frac{3}{4} \rho \kappa \underline{\vartheta} + \dots \right),$$

- elliptic estimates for Hodge operators on the 2-spheres foliating Σ_* ,

we infer the control with improved decay of all Ricci and curvature components on the spacelike hypersurface Σ_* .

Step 2. We derive decay estimates for the outgoing geodesic foliation of $^{(ext)}\mathcal{M}$. More precisely:

- First, we propagate the estimates involving only $u^{-\frac{1}{2}-\delta_{dec}}$ decay in u from Σ_* to $^{(ext)}\mathcal{M}$.
- We then focus on the harder to recover estimates, i.e. the ones involving $u^{-1-\delta_{dec}}$ decay in u . We proceed as follows.
 - We first propagate the main GCM quantities $\check{\kappa}$, $\check{\mu}$, and a renormalized quantity involving $\check{\xi}$ (see the quantity Ξ in Lemma 7.5.2) from Σ_* to $^{(ext)}\mathcal{M}$.
 - We then recover the estimates involving $u^{-1-\delta_{dec}}$ decay in u on \mathcal{T} . To this end, we use that we control the main GCM quantities, $\underline{\alpha}$ from Theorem M3 (since \mathcal{T} belongs both to $^{(ext)}\mathcal{M}$ and $^{(int)}\mathcal{M}$), \mathfrak{q} and α from Theorem M1–M2, and the estimates are then derived somewhat in the spirit of the ones on Σ_* , in particular by relying on elliptic estimates for Hodge operators on the 2-spheres foliating \mathcal{T} .
 - To recover the remaining estimates in $^{(ext)}\mathcal{M}$ involving $u^{-1-\delta_{dec}}$ decay in u , we integrate the transport equations in e_4 forward from \mathcal{T} , which concludes the proof of Theorem M4.

3.8.6 Discussion of Theorem M5

Here are the main steps in the proof of Theorem M5.

Step 1. We first derive decay estimates for the ingoing geodesic foliation of $^{(int)}\mathcal{M}$ on the timelike hyper surface \mathcal{T} . More precisely, thanks to

- the fact that the null frame of $^{(int)}\mathcal{M}$ is defined on \mathcal{T} as a simple conformal renormalization of the null frame of $^{(ext)}\mathcal{M}$ in view of its initialization, see section 3.1.2,
- the control of the outgoing geodesic foliation of $^{(ext)}\mathcal{M}$ on \mathcal{T} obtained in Theorem M4,

this allows us to transfer the decay estimates for $(^{ext})\check{R}, (^{ext})\check{\Gamma}$ to $(^{int})\check{R}, (^{int})\check{\Gamma}$ on \mathcal{T} .

Step 2. We derive on $(^{int})\mathcal{M}$ decay estimates for the ingoing geodesic foliation of $(^{int})\mathcal{M}$. More precisely, thanks to

- the improve decay estimates for $\underline{\alpha}$ in $(^{int})\mathcal{M}$ derived in Theorem M3,
- the improved decay estimates for $\check{\Gamma}$ and \check{R} on \mathcal{T} derived in the Step 1,
- the null structure equations and Bianchi identities,

we infer $O(\epsilon_0 u^{-1-\delta_{dec}})$ decay estimates for $\check{\Gamma}$ and \check{R} corresponding to the ingoing geodesic foliation of $(^{int})\mathcal{M}$ which concludes the proof of Theorem M5.

3.8.7 Discussion of Theorem M6

Step 1. Using

- (a) The control of the initial data layer,
- (b) Theorem GCMS-II of section 3.7.4,
- (c) Theorem GCMH of section 3.7.4,

we produce a smooth hypersurface Σ_* in the initial data layer starting from a GCM sphere S_* , and satisfying all the required properties for the future spacelike boundary of a GCM admissible spacetime, according to item 3 of definition 3.1.2.

Step 2. We then consider the outgoing geodesic foliation initialized on Σ_* which foliates the region we denote $(^{ext})\mathcal{M}$, to the past of Σ_* , and included in the outgoing part $(^{ext})\mathcal{L}_0$ of the initial data layer. In order to control it, we consider the transition functions $(f, \underline{f}, \lambda)$ from the background frame of the initial data layer to the frame of $(^{ext})\mathcal{M}$. These functions satisfy transport equations in e_4 with right-hand side depending on $(f, \underline{f}, \lambda)$ and the Ricci coefficients of the background foliation. Integrating the transport equations from Σ_* , where $(f, \underline{f}, \lambda)$ are under control as a by product of the use of Theorem GCMH in Step 1, we obtain the control of $(f, \underline{f}, \lambda)$ in $(^{ext})\mathcal{M}$. Using the transformation formulas of Proposition 2.3.4, and using the control of the initial data layer, we then infer the desired control (i.e. with ϵ_0 smallness constant and suitable r -weights) for the Ricci coefficients and curvature components of the foliation of $(^{ext})\mathcal{M}$.

Step 3. $^{(ext)}\mathcal{M}$ terminates on a timelike hypersurface \mathcal{T} of constant area radius²³. We then consider the ingoing geodesic foliation initialized on \mathcal{T} according to item 4 of definition 3.1.2, which foliates the region we denote $^{(int)}\mathcal{M}$, included in the ingoing part $^{(int)}\mathcal{L}_0$ of the initial data layer. Proceeding as in Step 2, relying on transport equations in e_3 instead of e_4 , we then derive the desired control (i.e. with ϵ_0 smallness constant) for the Ricci coefficients and curvature components of the foliation of $^{(int)}\mathcal{M}$, thus concluding the proof of Theorem M6.

3.8.8 Discussion of Theorem M7

From the assumptions of Theorem M7 we are given a GCM admissible spacetime $\mathcal{M} = \mathcal{M}(u_*) \in \mathfrak{N}(u_*)$ verifying the following improved bounds, for a universal constant $C > 0$,

$$\mathfrak{N}_{k_{small}+5}^{(Dec)}(\mathcal{M}) \leq C\epsilon_0$$

provided by Theorems M1-M5. We then proceed as follows.

Step 1. We extend \mathcal{M} by a local existence argument, to a strictly larger spacetime $\mathcal{M}^{(extend)}$, with a naturally extended foliation and the following slightly increased bounds

$$\mathfrak{N}_{k_{small}+5}^{(Dec)}(\mathcal{M}^{(extend)}) \leq 2C\epsilon_0.$$

but which may not verify our admissibility criteria.

Step 2. Using

- (a) The control of the extended spacetime $\mathcal{M}^{(extend)}$,
- (b) Theorem GCMS-II of section 3.7.4,
- (c) Theorem GCMH of section 3.7.4,

we produce a small piece of smooth GCM hypersurface $\tilde{\Sigma}_*$ in $\mathcal{M}^{(extend)} \setminus \mathcal{M}$ starting from a GCM sphere \tilde{S}_* .

Step 3. By a continuity argument based on a priori estimates, we extend $\tilde{\Sigma}_*$ all the way to the initial data layer, while ensuring that it remains in $\mathcal{M}^{(extend)} \setminus \mathcal{M}$ and satisfying all the required properties for the future spacelike boundary of a GCM admissible spacetime, according to item 3 of definition 3.1.2.

²³With respect to the foliation of $^{(ext)}\mathcal{M}$.

Step 4. We then consider the outgoing geodesic foliation initialized on $\widetilde{\Sigma}_*$ which foliates the region we denote ${}^{(ext)}\widetilde{\mathcal{M}}$, included in the outgoing part of $\mathcal{M}^{(extend)}$. In order to control it, we consider the transition functions $(f, \underline{f}, \lambda)$ from the background frame of the initial data layer to the frame of ${}^{(ext)}\widetilde{\mathcal{M}}$. These functions satisfy transport equations in e_4 with right-hand side depending on $(f, \underline{f}, \lambda)$ and the Ricci coefficients of the background foliation. Integrating the transport equations from $\widetilde{\Sigma}_*$, where $(f, \underline{f}, \lambda)$ are under control as a by product of the use of Theorem GCMH in Step 2, we obtain the control of $(f, \underline{f}, \lambda)$ in ${}^{(ext)}\widetilde{\mathcal{M}}$. Using the transformation formulas of Proposition 2.3.4, and using the control of the initial data layer, we then derive the desired control (i.e. with ϵ_0 smallness constant and suitable u and r weights) for the Ricci coefficients and curvature components of the foliation of ${}^{(ext)}\widetilde{\mathcal{M}}$.

Step 5. ${}^{(ext)}\widetilde{\mathcal{M}}$ terminates on a timelike hypersurface $\widetilde{\mathcal{T}}$ of constant area radius²⁴. We then consider the ingoing geodesic foliation initialized on $\widetilde{\mathcal{T}}$ according to item 4 of definition 3.1.2, which foliates the region we denote ${}^{(int)}\widetilde{\mathcal{M}}$, included in the ingoing part of $\mathcal{M}^{(extend)}$. Proceeding as in Step 4, relying on transport equations in e_3 instead of e_4 , we then derive the desired control (i.e. with ϵ_0 smallness constant and suitable u -weights) for the Ricci coefficients and curvature components of the foliation of ${}^{(int)}\widetilde{\mathcal{M}}$, thus concluding the proof of Theorem M7.

3.8.9 Discussion of Theorem M8

So far, we have only improved our bootstrap assumptions on decay estimates. We now improve our bootstrap assumptions on energies and weighted energies for \check{R} and $\check{\Gamma}$ relying on an iterative procedure recovering derivatives one by one²⁵.

Step 0. Let $I_{m_0, \delta_{\mathcal{H}}}$ the interval of \mathbb{R} defined by

$$I_{m_0, \delta_{\mathcal{H}}} := \left[2m_0 \left(1 + \frac{\delta_{\mathcal{H}}}{2} \right), 2m_0 \left(1 + \frac{3\delta_{\mathcal{H}}}{2} \right) \right]. \quad (3.8.6)$$

Recall that $\mathcal{T} = \{r = r_{\mathcal{T}}\}$, where $r_{\mathcal{T}} \in I_{m_0, \delta_{\mathcal{H}}}$, and note, see also Remark 3.6.3, that the results of Theorems M0–M7 hold for any $r_{\mathcal{T}} \in I_{m_0, \delta_{\mathcal{H}}}$.

It is at this stage that we need to make a specific choice of $r_{\mathcal{T}}$ in the context of a Lebesgue

²⁴With respect to the foliation of ${}^{(ext)}\widetilde{\mathcal{M}}$.

²⁵See also [29] for a related strategy to recover higher order derivatives from the control of lower order ones.

point argument. More precisely, we choose $r_{\mathcal{T}}$ such that we have

$$\int_{\{r=r_{\mathcal{T}}\}} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 = \inf_{r_0 \in I_{m_0, \delta \mathcal{H}}} \int_{\{r=r_0\}} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2. \quad (3.8.7)$$

In view of this definition, and since $\mathcal{T} = \{r = r_{\mathcal{T}}\}$, we infer that

$$\int_{\mathcal{T}} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 \lesssim \int_{(ext)\mathcal{M}(r \in I_{m_0, \delta \mathcal{H}})} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2. \quad (3.8.8)$$

Remark 3.8.2. *From now on, we may thus assume that the spacetime \mathcal{M} satisfies the conclusions of Theorem M0 and Theorem M7, as well as (3.8.8), and our goal is to prove Theorem M8, i.e. to prove that $\mathfrak{N}_{k_{large}}^{(En)} \lesssim \epsilon_0$ holds.*

Step 1. The $O(\epsilon_0)$ decay estimates derived in Theorem M7 imply in particular the following (non sharp) consequence

$$\mathfrak{N}_{k_{small}}^{(En)} \lesssim \epsilon_0,$$

where we recall²⁶

$$\mathfrak{N}_k^{(En)} = (ext)\mathfrak{N}_k[\check{R}] + (ext)\mathfrak{G}_k[\check{\Gamma}] + (int)\mathfrak{N}_k[\check{R}] + (int)\mathfrak{G}_k[\check{\Gamma}].$$

This allows us to initialize our iteration scheme in the next step.

Step 2. Next, for J such that $k_{small} \leq J \leq k_{large} - 1$, consider the iteration assumption

$$\mathfrak{N}_J^{(En)} \lesssim \epsilon_{\mathcal{B}}[J], \quad (3.8.9)$$

where

$$\epsilon_{\mathcal{B}}[J] := \sum_{j=k_{small}-2}^J (\epsilon_0)^{\ell(j)} \mathcal{B}^{1-\ell(j)} + \epsilon_0^{\ell(J)} \mathcal{B}, \quad \ell(j) := 2^{k_{small}-2-j}, \quad (3.8.10)$$

$$\mathcal{B} := \left(\int_{(ext)\mathcal{M}(r \in I_{m_0, \delta \mathcal{H}})} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 \right)^{\frac{1}{2}}. \quad (3.8.11)$$

In view of Step 1, (3.8.9) holds for $J = k_{small}$. From now on, we assume that (3.8.9) holds for J such that $k_{small} \leq J \leq k_{large} - 2$, and our goal is to show that this also holds for $J + 1$ derivatives.

²⁶See sections 3.2.1 and 3.2.2 for the definition of our norms measuring energies for curvature components and Ricci coefficients.

Step 3. Using the Teukolsky wave equations for α and $\underline{\alpha}$, as well as a wave equation for $\check{\rho}$, see Proposition 8.4.1, we derive Morawetz type estimates for $J + 1$ derivatives of these quantities in terms of $O(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \mathfrak{N}_{J+1}^{(En)})$.

Step 4. Relying on Bianchi identities, we also derive Morawetz type estimates for $J + 1$ derivatives for β and $\underline{\beta}$. As a consequence, we obtain Morawetz type estimates for $J + 1$ derivatives of all curvature components in terms of $O(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \mathfrak{N}_{J+1}^{(En)})$.

Step 5. As a consequence of Step 4, we immediately obtain, for any $r_0 \geq 4m_0$,

$${}^{(int)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}] \leq {}^{(ext)}\mathfrak{R}_{J+1}^{\geq r_0}[\check{R}] + O(r_0^{10}(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \mathfrak{N}_{J+1}^{(En)})).$$

Step 6. Relying on the Bianchi identities, we derive r^p -weighted estimates for $J + 1$ derivatives of curvature on $r \geq r_0$ with $r_0 \geq 4m_0$. We obtain

$${}^{(ext)}\mathfrak{R}_{J+1}^{\geq r_0}[\check{R}] \lesssim \frac{1}{r_0^{\delta_{\mathcal{B}}}} {}^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}] + r_0^{10}(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \mathfrak{N}_{J+1}^{(En)}).$$

Step 7. Next, we estimate the Ricci coefficients of ${}^{(ext)}\mathcal{M}$. To control them, we rely on the null structure equations in ${}^{(ext)}\mathcal{M}$. Using the null structure equations in ${}^{(ext)}\mathcal{M}$ and the GCM conditions on Σ_* , we derive the following weighted estimates for $J + 1$ derivatives of the Ricci coefficients

$${}^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] \lesssim {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}] + \epsilon_{\mathcal{B}}[J] + \epsilon_0 \mathfrak{N}_{J+1}^{(En)}.$$

Together with the estimates of Step 5 and Step 6, we infer for a large enough choice of r_0

$${}^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}] \lesssim \epsilon_{\mathcal{B}}[J] + \epsilon_0 \mathfrak{N}_{J+1}^{(En)}.$$

Step 8. Next, we estimate the Ricci coefficients of ${}^{(int)}\mathcal{M}$. Using the information on \mathcal{T} induced by Step 7 and the null structure equations in ${}^{(int)}\mathcal{M}$, we derive

$${}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] \lesssim {}^{(int)}\mathfrak{R}_{J+1}[\check{R}] + \epsilon_{\mathcal{B}}[J] + \epsilon_0 \mathfrak{N}_{J+1}^{(En)} + \left(\int_{\mathcal{T}} |\check{\mathfrak{d}}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}.$$

We need to deal with the last term. Relying on a trace theorem in the spacetime region ${}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})$, and the fact that $J + 2 \leq k_{large}$, we obtain

$$\left(\int_{\mathcal{T}} |\check{\mathfrak{d}}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}} \lesssim \left(\int_{{}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\check{\mathfrak{d}}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{4}} ({}^{(ext)}\mathfrak{R}_{J+1}[\check{R}])^{\frac{1}{2}} + {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}].$$

Step 9. The last estimate of Step 7 and the 2 estimates of Step 8 yield, for $\epsilon_0 > 0$ small enough,

$$\mathfrak{N}_{J+1}^{(En)} \lesssim \epsilon_{\mathcal{B}}[J] + \left(\int_{(ext)\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\mathfrak{d}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{4}} \left(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \mathfrak{N}_{J+1}^{(En)} \right)^{\frac{1}{2}}.$$

In view of the definition (3.8.10) of $\epsilon_{\mathcal{B}}[J]$, we infer that

$$\mathfrak{N}_{J+1}^{(En)} \lesssim \epsilon_{\mathcal{B}}[J+1]$$

which is the iteration assumption (3.8.9) for $J+1$ derivatives. We deduce that (3.8.9) holds for all $J \leq k_{large} - 1$, and hence

$$\mathfrak{N}_{k_{large}-1}^{(En)} \lesssim \epsilon_{\mathcal{B}}[k_{large} - 1].$$

Step 10. Relying on the conclusion of Step 9, and arguing as in Step 3 to Step 7, we obtain the conclusion of Step 7 for $J = k_{large} - 1$, i.e.

$${}^{(ext)}\mathfrak{G}_{k_{large}}[\check{\Gamma}] + {}^{(int)}\mathfrak{N}_{k_{large}}[\check{R}] + {}^{(ext)}\mathfrak{N}_{k_{large}}[\check{R}] \lesssim \epsilon_{\mathcal{B}}[k_{large} - 1] + \epsilon_0 \mathfrak{N}_{k_{large}}^{(En)}.$$

We then infer that

$$\epsilon_{\mathcal{B}}[k_{large} - 1] \lesssim \epsilon_0 + \epsilon_0 \mathfrak{N}_{k_{large}}^{(En)}$$

which yields, together with the last estimate of Step 9,

$${}^{(ext)}\mathfrak{G}_{k_{large}}[\check{\Gamma}] + {}^{(int)}\mathfrak{N}_{k_{large}}[\check{R}] + {}^{(ext)}\mathfrak{N}_{k_{large}}[\check{R}] \lesssim \epsilon_0 + \epsilon_0 \mathfrak{N}_{k_{large}}^{(En)}.$$

Step 11. It remains to recover ${}^{(int)}\mathfrak{G}_{k_{large}}[\check{\Gamma}]$. Arguing as for the first estimate of Step 8 with $J = k_{large} - 1$, we have

$${}^{(int)}\mathfrak{G}_{k_{large}}[\check{\Gamma}] \lesssim {}^{(int)}\mathfrak{N}_{k_{large}}[\check{R}] + \epsilon_{\mathcal{B}}[k_{large} - 1] + \epsilon_0 \mathfrak{N}_{k_{large}}^{(En)} + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}.$$

Thanks to the outcome of Step 10, we deduce that

$${}^{(int)}\mathfrak{G}_{k_{large}}[\check{\Gamma}] \lesssim \epsilon_0 + \epsilon_0 \mathfrak{N}_{k_{large}}^{(En)} + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}$$

and hence, for $\epsilon_0 > 0$ small enough, using again the last estimate of Step 10,

$$\mathfrak{N}_{k_{large}}^{(En)} \lesssim \epsilon_0 + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}.$$

It remains to estimate the last term of the RHS of the previous inequality. It is at this stage that we use the choice of $r_{\mathcal{T}}$, or rather its consequence (3.8.8), which implies

$$\left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}((ext)\check{R})|^2 \right)^{\frac{1}{2}} \lesssim \epsilon_0 + \epsilon_0 \mathfrak{N}_{k_{large}}^{(En)}$$

so that we finally obtain, for $\epsilon_0 > 0$ small enough,

$$\mathfrak{N}_{k_{large}}^{(En)} \lesssim \epsilon_0$$

which concludes the proof of Theorem M8.

3.9 Structure of the rest of the paper

The rest of this paper is devoted to the proof of Theorem M0-M8, as well as our GCM procedure. More precisely,

1. Theorem M0, together with other first consequences of the bootstrap assumptions, is proved in Chapter 4.
2. Theorem M1 is proved in Chapter 5.
3. Theorems M2 and M3 are proved in Chapter 6.
4. Theorems M4 and M5 are proved in Chapter 7.
5. Theorems M6, M7 and M8 are proved in Chapter 8.
6. Our GCM procedure is described in details in Chapter 9.
7. Chapter 10 contains estimates for Regge-Wheeler type wave equations used in Theorem M1.
8. Many of the long calculations are to be found in the appendix.

Chapter 4

CONSEQUENCES OF THE BOOTSTRAP ASSUMPTIONS

4.1 Proof of Theorem M0

According to the statement of Theorem M0 we consider given the initial layer $\mathcal{L}_0 = {}^{(ext)}\mathcal{L}_0 \cup {}^{(int)}\mathcal{L}_0$ as defined in Definition 3.1.1. We also assume that the initial layer norm verifies

$$\sup_{k \leq k_{large} + 5} \mathfrak{J}_k \lesssim \epsilon_0 \quad (4.1.1)$$

where $\mathfrak{J}_k = {}^{(ext)}\mathfrak{J}_k + {}^{(int)}\mathfrak{J}_k + \mathfrak{J}'_k$ and,

$$\begin{aligned} {}^{(ext)}\mathfrak{J}_0 &= \sup_{{}^{(ext)}\mathcal{L}_0} \left[r^{\frac{7}{2} + \delta_B} (|\alpha| + |\beta|) + r^3 \left| \rho + \frac{2m_0}{r^3} \right| + r^2 |\underline{\beta}| + r |\underline{\alpha}| \right] \\ &+ \sup_{{}^{(ext)}\mathcal{L}_0} r^2 \left(|\vartheta| + \left| \kappa - \frac{2}{r} \right| + |\zeta| + \left| \underline{\kappa} + \frac{2(1 - \frac{2m_0}{r})}{r} \right| \right) \\ &+ \sup_{{}^{(ext)}\mathcal{L}_0} r \left(|\underline{\vartheta}| + \left| \underline{\omega} - \frac{m_0}{r^2} \right| + |\underline{\xi}| \right) \\ &+ \sup_{{}^{(ext)}\mathcal{L}_0 ({}^{(ext)}r_0 \geq 4m_0)} \left(r \left| \frac{\gamma}{r^2} - 1 \right| + r |\underline{b}| + |\underline{\Omega}| + \Upsilon + |\varsigma - 1| + r \left| \frac{e^\Phi}{r \sin \theta} - 1 \right| \right), \end{aligned}$$

$$\begin{aligned} {}^{(int)}\mathfrak{J}_0 &= \sup_{{}^{(int)}\mathcal{L}_0} \left(|\underline{\alpha}| + |\underline{\beta}| + \left| \rho + \frac{2m_0}{r^3} \right| + |\beta| + |\alpha| \right) \\ &+ \sup_{{}^{(int)}\mathcal{L}_0} \left(|\vartheta| + \left| \kappa - \frac{2(1 - \frac{2m_0}{r})}{r} \right| + |\zeta| + \left| \underline{\kappa} + \frac{2}{r} \right| + |\underline{\vartheta}| + \left| \omega + \frac{m_0}{r^2} \right| + |\xi| \right), \end{aligned}$$

$$\mathfrak{J}'_0 = \sup_{{}^{(int)}\mathcal{L}_0 \cap {}^{(ext)}\mathcal{L}_0} (|f| + |\underline{f}| + |\log(\lambda_0^{-1}\lambda)|), \quad \lambda_0 = {}^{(ext)}\lambda_0 = 1 - \frac{2m_0}{{}^{(ext)}r\mathcal{L}_0},$$

with \mathfrak{J}_k the corresponding higher derivative norms obtained by replacing each component by $\mathfrak{d}^{\leq k}$ of it. In the definition of \mathfrak{J}'_0 above, $(f, \underline{f}, \lambda)$ denote the transition functions of Lemma 2.3.1 from the frame of the outgoing part ${}^{(ext)}\mathcal{L}_0$ of the initial data layer to the frame of the ingoing part ${}^{(int)}\mathcal{L}_0$ of the initial data layer in the region ${}^{(int)}\mathcal{L}_0 \cap {}^{(ext)}\mathcal{L}_0$.

We divide the proof of Theorem M0 in the following steps.

Step 1. We have the following lemma.

Lemma 4.1.1. *We have on ${}^{(ext)}\mathcal{M}$*

$$\begin{aligned} e_4 \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) &= \int_S \left(\frac{1}{2} \kappa e_\theta(\underline{\kappa}) - \frac{1}{2} \underline{\kappa} e_\theta(\kappa) + 4K\zeta + 2e_\theta(\rho) - \frac{1}{2} e_\theta(\vartheta \underline{\vartheta}) \right. \\ &\quad \left. - \vartheta e_\theta(\underline{\kappa}) + 2e_\theta(\zeta^2) \right) e^\Phi, \\ e_4 \left(\int_S \beta e^\Phi \right) &= \int_S \left(-\frac{1}{2} \kappa \beta + \zeta \alpha - \frac{1}{2} \vartheta \beta \right) e^\Phi, \\ e_3 \left(\int_S \beta e^\Phi \right) &= -\frac{1}{4} \int_S e_\theta(\kappa \underline{\kappa}) e^\Phi + 3\bar{\rho} \int_S \eta e^\Phi + \frac{1}{4} \int_S e_\theta(\vartheta \underline{\vartheta}) e^\Phi \\ &\quad + \int_S \left(\frac{1}{2} \underline{\kappa} \beta + 2\underline{\omega} \beta + 3\eta \check{\rho} - \vartheta \underline{\beta} + \underline{\xi} \alpha - \frac{1}{2} \underline{\vartheta} \beta \right) e^\Phi + Err \left[e_3 \left(\int_S \beta e^\Phi \right) \right], \end{aligned}$$

and

$$\begin{aligned} e_3 \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) &= \underline{\kappa} e_3 \left(\int_S \zeta e^\Phi \right) - \underline{\kappa} \int_S \underline{\beta} e^\Phi \\ &\quad + \int_S \left(-\underline{\kappa} \underline{\beta} - \frac{1}{2} \underline{\kappa}^2 \zeta + 6\rho \underline{\xi} - 2\underline{\omega} e_\theta(\underline{\kappa}) - \frac{1}{2} \underline{\vartheta} (e_\theta(\underline{\kappa}) - \underline{\kappa} \zeta) + Err[\underline{\vartheta}_2 \underline{\vartheta}_2^* \underline{\xi}] \right) e^\Phi \\ &\quad + Err \left[e_3 \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) \right] + \int_S \underline{\check{\kappa}} \left(e_3(\zeta) + \left(\frac{3}{2} \underline{\kappa} - \frac{1}{2} \underline{\vartheta} \right) \zeta \right) e^\Phi \\ &\quad - \underline{\check{\kappa}} \int_S \left(e_3(\zeta) + \left(\frac{3}{2} \underline{\kappa} - \frac{1}{2} \underline{\vartheta} \right) \zeta \right) e^\Phi - \underline{\kappa} Err \left[e_3 \left(\int_S \zeta e^\Phi \right) \right]. \end{aligned}$$

Proof. We have in ${}^{(ext)}\mathcal{M}$, see Proposition 2.2.8,

$$e_4(\underline{\kappa}) + \frac{1}{2}\underline{\kappa}\underline{\kappa} = -2\underline{\phi}_1\underline{\zeta} + 2\rho - \frac{1}{2}\underline{\vartheta}\underline{\vartheta} + 2\underline{\zeta}^2.$$

Together with the following commutation relation

$$[e_\theta, e_4] = \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta,$$

we infer

$$e_4(e_\theta(\underline{\kappa})) + \underline{\kappa}e_\theta(\underline{\kappa}) + \frac{1}{2}\underline{\vartheta}e_\theta(\underline{\kappa}) + \frac{1}{2}\underline{\kappa}e_\theta(\underline{\kappa}) = 2\underline{\phi}_1^*\underline{\phi}_1\underline{\zeta} + 2e_\theta(\rho) - \frac{1}{2}e_\theta(\underline{\vartheta}\underline{\vartheta}) + 2e_\theta(\underline{\zeta}^2).$$

Also, we have in view of Proposition 2.2.19 the following identity

$$e_3(e_\theta(\underline{\kappa})) - \underline{\kappa}e_3(\underline{\zeta}) = -2\underline{\phi}_2\underline{\phi}_2^*\underline{\zeta} - \underline{\kappa}\underline{\beta} + \underline{\kappa}^2\underline{\zeta} - \frac{3}{2}\underline{\kappa}e_\theta\underline{\kappa} + 6\rho\underline{\zeta} - 2\underline{\omega}e_\theta(\underline{\kappa}) + \text{Err}[\underline{\phi}_2\underline{\phi}_2^*\underline{\zeta}].$$

Next, in view of Corollary 2.2.10, we have in ${}^{(ext)}\mathcal{M}$

$$\begin{aligned} e_4\left(\int_S e_\theta(\underline{\kappa})e^\Phi\right) &= \int_S \left(e_4(e_\theta(\underline{\kappa})) + \left(\frac{3}{2}\underline{\kappa} - \frac{1}{2}\underline{\vartheta}\right)e_\theta(\underline{\kappa})\right)e^\Phi \\ e_4\left(\int_S \beta e^\Phi\right) &= \int_S \left(e_4(\beta) + \left(\frac{3}{2}\underline{\kappa} - \frac{1}{2}\underline{\vartheta}\right)\beta\right)e^\Phi, \\ e_3\left(\int_S \beta e^\Phi\right) &= \int_S \left(e_3(\beta) + \left(\frac{3}{2}\underline{\kappa} - \frac{1}{2}\underline{\vartheta}\right)\beta\right)e^\Phi + \text{Err}\left[e_3\left(\int_S \beta e^\Phi\right)\right], \end{aligned}$$

and

$$\begin{aligned} &e_3\left(\int_S e_\theta(\underline{\kappa})e^\Phi\right) - \underline{\kappa}e_3\left(\int_S \zeta e^\Phi\right) \\ &= \int_S \left(e_3(e_\theta(\underline{\kappa})) + \left(\frac{3}{2}\underline{\kappa} - \frac{1}{2}\underline{\vartheta}\right)e_\theta(\underline{\kappa})\right)e^\Phi + \text{Err}\left[e_3\left(\int_S e_\theta(\underline{\kappa})e^\Phi\right)\right] \\ &\quad - \underline{\kappa}\int_S \left(e_3(\zeta) + \left(\frac{3}{2}\underline{\kappa} - \frac{1}{2}\underline{\vartheta}\right)\zeta\right)e^\Phi - \underline{\kappa}\text{Err}\left[e_3\left(\int_S \zeta e^\Phi\right)\right] \\ &= \int_S \left(e_3(e_\theta(\underline{\kappa})) - \underline{\kappa}e_3(\zeta) + \left(\frac{3}{2}\underline{\kappa} - \frac{1}{2}\underline{\vartheta}\right)(e_\theta(\underline{\kappa}) - \underline{\kappa}\zeta)\right)e^\Phi + \text{Err}\left[e_3\left(\int_S e_\theta(\underline{\kappa})e^\Phi\right)\right] \\ &\quad + \int_S \underline{\check{\kappa}}\left(e_3(\zeta) + \left(\frac{3}{2}\underline{\kappa} - \frac{1}{2}\underline{\vartheta}\right)\zeta\right)e^\Phi - \underline{\check{\kappa}}\int_S \left(e_3(\zeta) + \left(\frac{3}{2}\underline{\kappa} - \frac{1}{2}\underline{\vartheta}\right)\zeta\right)e^\Phi - \underline{\kappa}\text{Err}\left[e_3\left(\int_S \zeta e^\Phi\right)\right]. \end{aligned}$$

Together with the above identities for $e_4(e_\theta(\underline{\kappa}))$ and $e_3(e_\theta(\underline{\kappa}))$, as well as the Bianchi identities of Proposition 2.2.8 for $e_4(\beta)$ and $e_3(\beta)$, we infer

$$\begin{aligned} e_4 \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) &= \int_S \left(\frac{1}{2} \underline{\kappa} e_\theta(\underline{\kappa}) - \frac{1}{2} \underline{\kappa} e_\theta(\underline{\kappa}) + 2 \underline{\phi}_1^* \underline{\phi}_1 \zeta + 2 e_\theta(\rho) - \frac{1}{2} e_\theta(\vartheta \underline{\vartheta}) \right. \\ &\quad \left. - \vartheta e_\theta(\underline{\kappa}) + 2 e_\theta(\zeta^2) \right) e^\Phi, \\ e_4 \left(\int_S \beta e^\Phi \right) &= \int_S \left(-\frac{1}{2} \underline{\kappa} \beta + \underline{\phi}_2 \alpha + \zeta \alpha - \frac{1}{2} \vartheta \beta \right) e^\Phi, \\ e_3 \left(\int_S \beta e^\Phi \right) &= \int_S \left(\frac{1}{2} \underline{\kappa} \beta + e_\theta(\rho) + 2 \underline{\omega} \beta + 3 \eta \rho - \vartheta \underline{\beta} + \underline{\xi} \alpha - \frac{1}{2} \underline{\vartheta} \beta \right) e^\Phi \\ &\quad + \text{Err} \left[e_3 \left(\int_S \beta e^\Phi \right) \right], \end{aligned}$$

and

$$\begin{aligned} &e_3 \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) - \underline{\kappa} e_3 \left(\int_S \zeta e^\Phi \right) \\ &= \int_S \left(-2 \underline{\phi}_2 \underline{\phi}_2^* \underline{\xi} - \underline{\kappa} \underline{\beta} - \frac{1}{2} \underline{\kappa}^2 \zeta + 6 \rho \underline{\xi} - 2 \underline{\omega} e_\theta(\underline{\kappa}) - \frac{1}{2} \underline{\vartheta} (e_\theta(\underline{\kappa}) - \underline{\kappa} \zeta) + \text{Err}[\underline{\phi}_2 \underline{\phi}_2^* \underline{\xi}] \right) e^\Phi \\ &\quad + \text{Err} \left[e_3 \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) \right] + \int_S \underline{\check{\kappa}} \left(e_3(\zeta) + \left(\frac{3}{2} \underline{\kappa} - \frac{1}{2} \underline{\vartheta} \right) \zeta \right) e^\Phi \\ &\quad - \underline{\check{\kappa}} \int_S \left(e_3(\zeta) + \left(\frac{3}{2} \underline{\kappa} - \frac{1}{2} \underline{\vartheta} \right) \zeta \right) e^\Phi - \underline{\kappa} \text{Err} \left[e_3 \left(\int_S \zeta e^\Phi \right) \right]. \end{aligned}$$

Using in particular the fact that $\underline{\phi}_2^*(e^\Phi) = 0$, that $\underline{\phi}_2^*$ is the adjoint of $\underline{\phi}_2$, and the identity $\underline{\phi}_1^* \underline{\phi}_1 = \underline{\phi}_2 \underline{\phi}_2^* + 2K$, we deduce

$$\begin{aligned} e_4 \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) &= \int_S \left(\frac{1}{2} \underline{\kappa} e_\theta(\underline{\kappa}) - \frac{1}{2} \underline{\kappa} e_\theta(\underline{\kappa}) + 4K \zeta + 2 e_\theta(\rho) - \frac{1}{2} e_\theta(\vartheta \underline{\vartheta}) \right. \\ &\quad \left. - \vartheta e_\theta(\underline{\kappa}) + 2 e_\theta(\zeta^2) \right) e^\Phi, \\ e_4 \left(\int_S \beta e^\Phi \right) &= \int_S \left(-\frac{1}{2} \underline{\kappa} \beta + \zeta \alpha - \frac{1}{2} \vartheta \beta \right) e^\Phi, \\ e_3 \left(\int_S \beta e^\Phi \right) &= \int_S e_\theta(\rho) e^\Phi + 3 \bar{\rho} \int_S \eta e^\Phi + \int_S \left(\frac{1}{2} \underline{\kappa} \beta + 2 \underline{\omega} \beta + 3 \eta \bar{\rho} - \vartheta \underline{\beta} + \underline{\xi} \alpha - \frac{1}{2} \underline{\vartheta} \beta \right) e^\Phi \\ &\quad + \text{Err} \left[e_3 \left(\int_S \beta e^\Phi \right) \right], \end{aligned}$$

and

$$\begin{aligned}
e_3 \left(\int_S e_{\theta(\underline{\kappa})} e^\Phi \right) &= \underline{\kappa} e_3 \left(\int_S \zeta e^\Phi \right) - \bar{\kappa} \int_S \underline{\beta} e^\Phi \\
&+ \int_S \left(-\underline{\check{\kappa}} \underline{\beta} - \frac{1}{2} \underline{\kappa}^2 \zeta + 6\rho \underline{\xi} - 2\underline{\omega} e_{\theta(\underline{\kappa})} - \frac{1}{2} \underline{\vartheta} (e_{\theta(\underline{\kappa})} - \underline{\kappa} \zeta) + \text{Err}[\underline{\vartheta}_2 \underline{\vartheta}_2^* \underline{\xi}] \right) e^\Phi \\
&+ \text{Err} \left[e_3 \left(\int_S e_{\theta(\underline{\kappa})} e^\Phi \right) \right] + \int_S \underline{\check{\kappa}} \left(e_3(\zeta) + \left(\frac{3}{2} \underline{\kappa} - \frac{1}{2} \underline{\vartheta} \right) \zeta \right) e^\Phi \\
&- \underline{\check{\kappa}} \int_S \left(e_3(\zeta) + \left(\frac{3}{2} \underline{\kappa} - \frac{1}{2} \underline{\vartheta} \right) \zeta \right) e^\Phi - \underline{\kappa} \text{Err} \left[e_3 \left(\int_S \zeta e^\Phi \right) \right].
\end{aligned}$$

Finally, from the identity (2.1.21) for $e_\theta(K)$ and the formula for K , we have

$$\int_S e_{\theta(\rho)} e^\Phi = -\frac{1}{4} \int_S e_{\theta(\underline{\kappa} \underline{\kappa})} e^\Phi + \frac{1}{4} \int_S e_{\theta(\underline{\vartheta} \underline{\vartheta})} e^\Phi.$$

We deduce

$$\begin{aligned}
e_3 \left(\int_S \beta e^\Phi \right) &= -\frac{1}{4} \int_S e_{\theta(\underline{\kappa} \underline{\kappa})} e^\Phi + 3\bar{\rho} \int_S \eta e^\Phi + \frac{1}{4} \int_S e_{\theta(\underline{\vartheta} \underline{\vartheta})} e^\Phi \\
&+ \int_S \left(\frac{1}{2} \underline{\kappa} \underline{\beta} + 2\underline{\omega} \underline{\beta} + 3\underline{\eta} \underline{\rho} - \underline{\vartheta} \underline{\beta} + \underline{\xi} \underline{\alpha} - \frac{1}{2} \underline{\vartheta} \underline{\beta} \right) e^\Phi + \text{Err} \left[e_3 \left(\int_S \beta e^\Phi \right) \right]
\end{aligned}$$

which concludes the proof of Lemma 4.1.1. \square

Step 2. Using the transport equations of Lemma 4.1.1 and the bootstrap assumptions on decay for $k = 0, 1$ derivatives in $^{(ext)}\mathcal{M}$, we infer in that region, and in particular on Σ_*

$$\begin{aligned}
\left| e_4 \left(\int_S e_{\theta(\underline{\kappa})} e^\Phi \right) \right| &\lesssim \frac{\epsilon}{r u^{\frac{1}{2} + \delta_{dec}}}, \\
\left| e_4 \left(\int_S \beta e^\Phi \right) \right| &\lesssim \frac{\epsilon}{r^{\frac{3}{2} + \delta_{dec}}}, \\
\left| e_3 \left(\int_S \beta e^\Phi \right) \right| &\lesssim \left| \int_S e_{\theta(\underline{\kappa} \underline{\kappa})} e^\Phi \right| + r^{-3} \left| \int_S \eta e^\Phi \right| + \frac{\epsilon}{r^{\frac{3}{2} + \delta_{dec}}} + \frac{\epsilon^2}{r u^{1 + \delta_{dec}}}, \\
\left| e_3 \left(\int_S e_{\theta(\underline{\kappa})} e^\Phi \right) \right| &\lesssim r^{-1} \left| e_3 \left(\int_S \zeta e^\Phi \right) \right| + r^{-1} \left| \int_S \underline{\beta} e^\Phi \right| + \frac{\epsilon}{r u^{\frac{1}{2} + \delta_{dec}}} + \frac{\epsilon^2}{u^{2 + 2\delta_{dec}}}.
\end{aligned}$$

Next, we focus on the two last estimates. Recall the following GCM conditions

$$\kappa = \frac{2}{r}, \quad \int_S \eta e^\Phi = 0 \quad \text{on } \Sigma_*.$$

We deduce on Σ_*

$$\left| e_3 \left(\int_S \beta e^\Phi \right) \right| \lesssim r^{-1} \left| \int_S e_{\theta(\underline{\kappa})} e^\Phi \right| + \frac{\epsilon}{r^{\frac{3}{2} + \delta_{dec}}} + \frac{\epsilon^2}{ru^{1 + \delta_{dec}}}.$$

Also, projecting both Codazzi on e^Φ , using $\not{d}_2^*(e^\Phi) = 0$ and the fact that \not{d}_2^* is the adjoint of \not{d}_2 , and using also the GCM condition for κ on Σ_* , we have on Σ_*

$$\begin{aligned} \int_S \underline{\beta} e^\Phi &= -\frac{1}{2} \int_S \not{d}_1^* \underline{\kappa} e^\Phi - \frac{1}{2} \int_S \zeta \underline{\kappa} e^\Phi + \frac{1}{2} \int_S \underline{\vartheta} \zeta e^\Phi, \\ \int_S \zeta e^\Phi &= r \int_S \beta e^\Phi + \frac{r}{2} \int_S \vartheta \zeta e^\Phi. \end{aligned}$$

Together with the bootstrap assumptions on decay for $k = 0, 1$ derivatives in $^{(ext)}\mathcal{M}$, we infer on Σ_*

$$\left| e_3 \left(\int_S \zeta e^\Phi \right) \right| + \left| \int_S \underline{\beta} e^\Phi \right| \lesssim \frac{\epsilon}{u^{\frac{1}{2} + \delta_{dec}}}.$$

We have thus on Σ_*

$$\begin{aligned} \left| e_4 \left(\int_S e_{\theta(\underline{\kappa})} e^\Phi \right) \right| &\lesssim \frac{\epsilon}{ru^{\frac{1}{2} + \delta_{dec}}}, \\ \left| e_4 \left(\int_S \beta e^\Phi \right) \right| &\lesssim \frac{\epsilon}{r^{\frac{3}{2} + \delta_{dec}}}, \\ \left| e_3 \left(\int_S \beta e^\Phi \right) \right| &\lesssim r^{-1} \left| \int_S e_{\theta(\underline{\kappa})} e^\Phi \right| + \frac{\epsilon}{r^{\frac{3}{2} + \delta_{dec}}} + \frac{\epsilon^2}{ru^{1 + \delta_{dec}}}, \\ \left| e_3 \left(\int_S e_{\theta(\underline{\kappa})} e^\Phi \right) \right| &\lesssim \frac{\epsilon}{ru^{\frac{1}{2} + \delta_{dec}}} + \frac{\epsilon^2}{u^{2 + 2\delta_{dec}}}. \end{aligned}$$

In view of the behavior (3.3.4) of r on Σ_* , we infer on Σ_*

$$\begin{aligned} \left| e_4 \left(\int_S e_{\theta(\underline{\kappa})} e^\Phi \right) \right| &\lesssim \frac{\epsilon_0}{u^{2 + 2\delta_{dec}}}, \\ \left| e_4 \left(\int_S \beta e^\Phi \right) \right| &\lesssim \frac{\epsilon_0}{ru^{1 + \delta_{dec}}}, \\ \left| e_3 \left(\int_S \beta e^\Phi \right) \right| &\lesssim r^{-1} \left| \int_S e_{\theta(\underline{\kappa})} e^\Phi \right| + \frac{\epsilon_0}{ru^{1 + \delta_{dec}}}, \\ \left| e_3 \left(\int_S e_{\theta(\underline{\kappa})} e^\Phi \right) \right| &\lesssim \frac{\epsilon_0}{u^{2 + 2\delta_{dec}}}. \end{aligned}$$

Step 3. Let ν_* the unique tangent vector to Σ_* which can be written as

$$\nu_* = e_3 + ae_4$$

where a is a scalar function on Σ_* . Recall that there exists a constant c_* such that $\Sigma_* = \{u + r = c_*\}$. We infer $\nu_*(u + r) = 0$ and hence

$$0 = e_3(u + r) + ae_4(u + r) = \frac{2}{\varsigma} + \frac{r}{2}(\bar{\kappa} + \underline{A}) + a\frac{r}{2}\bar{\kappa}$$

which yields

$$a = -\frac{\frac{2}{\varsigma} + \frac{r}{2}(\bar{\kappa} + \underline{A})}{\frac{r}{2}\bar{\kappa}}.$$

In view of the GCM condition on κ and the definition of the Hawking mass m , we have on Σ_*

$$\bar{\kappa} = \frac{2}{r}, \quad \underline{\bar{\kappa}} = -\frac{2\Upsilon}{r}$$

and hence, we have on Σ_*

$$a = -\frac{2}{\varsigma} + \Upsilon - \frac{r}{2}\underline{A}.$$

The bootstrap assumptions on decay for $k = 0$ derivatives in $^{(ext)}\mathcal{M}$, the definition (2.2.22) of \underline{A} , and the estimates for ς and $\underline{\Omega}$ yield the rough estimate¹

$$|a| \lesssim 1.$$

Together with the fact that $\nu_* = e_3 + ae_4$ and the estimates of Step 2, we infer

$$\begin{aligned} \left| \nu_* \left(\int_S e_{\theta}(\underline{\kappa}) e^{\Phi} \right) \right| &\lesssim \frac{\epsilon_0}{u^{2+2\delta_{dec}}}, \\ \left| \nu_* \left(\int_S \beta e^{\Phi} \right) \right| &\lesssim r^{-1} \left| \int_S e_{\theta}(\underline{\kappa}) e^{\Phi} \right| + \frac{\epsilon_0}{ru^{1+\delta_{dec}}}. \end{aligned}$$

Step 4. Now, recall that we have the following GCM on the last sphere $S_* = \Sigma_* \cap \mathcal{C}_*$ of Σ_*

$$\int_{S_*} e_{\theta}(\underline{\kappa}) e^{\Phi} = \int_{S_*} \beta e^{\Phi} = 0.$$

Integrating backward from S_* the estimate for $e_{\theta}(\underline{\kappa})$ of Step 3 yields on Σ_*

$$\left| \int_S e_{\theta}(\underline{\kappa}) e^{\Phi} \right| \lesssim \frac{\epsilon_0}{u^{1+\delta_{dec}}}.$$

¹The estimates for $\underline{\Omega}$ and ς are proved later in Proposition 3.4.3. Since the proof does not rely on Theorem M0, we may use it here.

Plugging in the estimate for β of Step 3, we infer on Σ_*

$$\left| \nu_* \left(\int_S \beta e^\Phi \right) \right| \lesssim \frac{\epsilon_0}{ru^{1+\delta_{dec}}}.$$

Integrating backward from S_* yields on Σ_*

$$\left| \int_S \beta e^\Phi \right| \lesssim \frac{\epsilon_0}{ru^{\delta_{dec}}}.$$

Thus, at the first sphere $S_1 = \Sigma_* \cap \mathcal{C}_1$ of Σ_* , we have obtained

$$\left| \int_{S_1} e_{\theta(\underline{\kappa})} e^\Phi \right| + r \left| \int_{S_1} \beta e^\Phi \right| \lesssim \epsilon_0. \quad (4.1.2)$$

Remark 4.1.2. *Note that the only bootstrap assumptions used in the proof of Theorem M0 are the bootstrap assumption **BA-D** on decay for $k = 0, 1$ derivatives. Indeed, to obtain (4.1.2), we have only used, in Steps 1–4, the bootstrap assumption **BA-D** on decay for $k = 0, 1$ derivatives, while, from now on, we will only rely on (4.1.2) and the assumptions (4.1.1) on the initial data layer. This observation will allow us to use the conclusions of Theorem M0, not only for the bootstrap spacetime \mathcal{M} in Theorem M1–M5, but also for the extended spacetime in the proof of Theorem M8, where the only assumptions are the one on decay (which are established for the extended spacetime in Theorem M7).*

Step 5. On the sphere $S_1 = \Sigma_* \cap \mathcal{C}_1$ of Σ_* , we have in view of the GCM conditions of Σ_* and (4.1.2)

$$\kappa = \frac{2}{r}, \quad \not{d}_2^* \not{d}_1^* \underline{\kappa} = 0, \quad \not{d}_2^* \not{d}_1^* \mu = 0, \quad \left| \int_{S_1} e_{\theta(\underline{\kappa})} e^\Phi \right| + r \left| \int_{S_1} \beta e^\Phi \right| \lesssim \epsilon_0. \quad (4.1.3)$$

We consider the transition functions $(f, \underline{f}, \lambda)$ from the frame of the outgoing part ${}^{(ext)}\mathcal{L}_0$ of the initial data layer to the frame of ${}^{(ext)}\mathcal{M}$. Since

- S_1 is a sphere of ${}^{(ext)}\mathcal{M}$ in the initial data layer,
- S_1 is a sphere of the GCM hypersurface Σ_* ,
- the estimate (4.1.3) holds on S_1 ,

we can invoke the GCM corollary Rigidity II and III of section 3.7.4 with the choice $\overset{\circ}{\epsilon} = \overset{\circ}{\delta} = \epsilon_0$, $s_{max} = k_{large} + 5$, and with the background foliation being the one of the outgoing part ${}^{(ext)}\mathcal{L}_0$ of the initial data layer. We obtain

$$\|\not{d}^{\leq k_{large}+6}(f, \underline{f}, \log \lambda)\|_{L^2(S_1)} + \sup_{S_1} \left| \frac{m}{m_0} - 1 \right| \lesssim \epsilon_0$$

Together with the Sobolev embedding on S_1 , we deduce

$$\sup_{S_1} \left(r |\mathfrak{d}^{\leq k_{large}+4}(f, \underline{f}, \log \lambda)| + \left| \frac{m}{m_0} - 1 \right| + |r - {}^{(ext)}r_{\mathcal{L}_0}| \right) \lesssim \epsilon_0. \quad (4.1.4)$$

Step 6. Recall from Corollary 2.3.7 that $(\underline{f}, f, \log(\lambda))$ satisfy the following transport equations along \mathcal{C}_1

$$\begin{aligned} \lambda^{-1} e'_4(rf) &= E'_1(f, \Gamma), \\ \lambda^{-1} e'_4(\log(\lambda)) &= E'_2(f, \Gamma), \\ \lambda^{-1} e'_4 \left(r \underline{f} - 2r^2 e'_\theta(\log(\lambda)) + r f \underline{\Omega} \right) &= E'_3(f, \underline{f}, \lambda, \Gamma), \end{aligned}$$

where, in view of the form of E'_1, E'_2, E'_3 in Corollary 2.3.7 and the estimates (4.1.1) for the Ricci coefficients of the outgoing part ${}^{(ext)}\mathcal{L}_0$ of the initial data layer, we have

$$|\mathfrak{d}^k E'_1(f, \Gamma)| + |\mathfrak{d}^k E'_2(f, \Gamma)| \lesssim \frac{\epsilon_0^2}{r^2} + |\mathfrak{d}^{\leq k} f|^2 \text{ for } k \leq k_{large} + 5 \text{ on } \mathcal{C}_1$$

and

$$|\mathfrak{d}^k E'_3(f, \underline{f}, \lambda, \Gamma)| \lesssim \frac{\epsilon_0^2}{r^2} + |\mathfrak{d}^{\leq k+1} f|^2 + |\mathfrak{d}^{\leq k+1} \log(\lambda)|^2 \text{ for } k \leq k_{large} + 5 \text{ on } \mathcal{C}_1.$$

Next, recall from Lemma 2.2.14 the following commutator identity

$$[\mathbf{T}, e_4] = \left(\left(\frac{\omega}{r} - \frac{m}{r^2} \right) - \frac{m}{2r} \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{e_4(m)}{r} \right) e_4 + (\eta + \zeta) e_\theta$$

while from Lemma 2.2.13, we have schematically

$$[\mathfrak{D}, e_4] = (\check{\kappa}, \vartheta) \mathfrak{D} + (\zeta, r\beta)$$

Together with the fact that

$$\lambda^{-1} e'_4 = e_4 + f e_\theta + \frac{f^2}{4} e_3,$$

the commutator above identities for $[T, e_4]$ and $[\mathfrak{D}, e_4]$, as well as the estimates (4.1.1) for the Ricci coefficients and curvature components of the outgoing part ${}^{(ext)}\mathcal{L}_0$ of the initial data layer, we infer, for $k \leq k_{large} + 5$,

$$\begin{aligned} & |\mathfrak{d}^k [T, \lambda^{-1} e'_4] h| + |\mathfrak{d}^k [\mathfrak{D}, \lambda^{-1} e'_4] h| \\ & \lesssim \frac{\epsilon_0}{r^2} |\mathfrak{d}^{\leq k+1} h| + \frac{1}{r} |\mathfrak{d}^{\leq k} (f \mathfrak{D} h)| + \frac{1}{r} |\mathfrak{d}^{\leq k} (h \mathfrak{D} f)| + |\mathfrak{d}^{\leq k} (f^2 \mathfrak{D} h)| + |\mathfrak{d}^{\leq k} (h f \mathfrak{D} f)|. \end{aligned}$$

By commuting first the transport equations in the direction $\lambda^{-1}e'_4$ with $(T, \mathfrak{D})^k$, and by using these transport equations to recover the e_4 derivatives, we deduce

$$\begin{aligned}\lambda^{-1}e'_4(r\mathfrak{D}^k f) &= E'_{1,k}(f, \Gamma), \\ \lambda^{-1}e'_4(\mathfrak{D}^k \log(\lambda)) &= E'_{2,k}(f, \Gamma), \\ \lambda^{-1}e'_4\left(r\mathfrak{D}^k(\underline{f} - 2re'_\theta(\log(\lambda)) + f\underline{\Omega})\right) &= E'_{3,k}(f, \underline{f}, \lambda, \Gamma),\end{aligned}$$

where we have

$$|E'_{1,k}(f, \Gamma)| + |E'_{2,k}(f, \Gamma)| \lesssim \frac{\epsilon_0^2}{r^2} + |\mathfrak{D}^{\leq k} f|^2 \text{ for } k \leq k_{large} + 5 \text{ on } \mathcal{C}_1$$

and

$$|E'_{3,k}(f, \underline{f}, \lambda, \Gamma)| \lesssim \frac{\epsilon_0^2}{r^2} + |\mathfrak{D}^{\leq k+1} f|^2 + |\mathfrak{D}^{\leq k+1} \log(\lambda)|^2 + |\mathfrak{D}^{\leq k} \underline{f}|^2 \text{ for } k \leq k_{large} + 5 \text{ on } \mathcal{C}_1.$$

This allows us to propagate the estimates for $(f, \underline{f}, \lambda)$ in (4.1.4) on S_1 to any sphere on \mathcal{C}_1 , and hence

$$\sup_{\mathcal{C}_1} \left(r |\mathfrak{D}^{\leq k_{large}+4}(f, \log \lambda)| \right) + \sup_{\mathcal{C}_1} \left(r |\mathfrak{D}^{\leq k_{large}+3} \underline{f}| \right) \lesssim \epsilon_0. \quad (4.1.5)$$

Step 7. In view of (4.1.5), the change of frame formulas of Proposition 2.3.4, and the estimates (4.1.1) for the Ricci coefficients and curvature components of the outgoing part ${}^{(ext)}\mathcal{L}_0$ of the initial data layer, we obtain

$$\begin{aligned}\max_{0 \leq k \leq k_{large}} \left\{ \sup_{\mathcal{C}_1} \left[r^{\frac{7}{2}+\delta_B} (|\mathfrak{D}^k {}^{(ext)}\alpha| + |\mathfrak{D}^k {}^{(ext)}\beta|) + r^{\frac{9}{2}+\delta_B} |\mathfrak{D}^{k-1} e_3({}^{(ext)}\alpha)| \right] \right. \\ \left. + \sup_{\mathcal{C}_1} \left[r^3 \left| \mathfrak{D}^k \left({}^{(ext)}\rho + \frac{2m_0}{r^3} \right) \right| + r^2 |\mathfrak{D}^k {}^{(ext)}\underline{\beta}| + r |\mathfrak{D}^k {}^{(ext)}\underline{\alpha}| \right] \right\} \lesssim \epsilon_0,\end{aligned} \quad (4.1.6)$$

as well as

$$\max_{0 \leq k \leq k_{large}} \left\{ \sup_{\mathcal{C}_1} \left[r^2 (|\mathfrak{D}^k {}^{(ext)}(\check{\kappa}, \vartheta, \zeta, \check{\underline{\kappa}})| + r |\mathfrak{D}^k {}^{(ext)}\underline{\vartheta}|) \right] \right\} \lesssim \epsilon_0.$$

Now, according to Proposition 2.2.16, we have in ${}^{(ext)}\mathcal{M}$

$$\begin{aligned}{}^{(ext)}e_4({}^{(ext)}m) &= \frac{{}^{(ext)}r}{32\pi} \int_S \left(2 {}^{(ext)}\check{\kappa} {}^{(ext)}\check{\rho} + 2 {}^{(ext)}e_\theta({}^{(ext)}\kappa) {}^{(ext)}\zeta - \frac{1}{2} {}^{(ext)}\check{\underline{\kappa}} ({}^{(ext)}\vartheta)^2 \right. \\ &\quad \left. - \frac{1}{2} {}^{(ext)}\check{\kappa} {}^{(ext)}\vartheta {}^{(ext)}\underline{\vartheta} + 2 {}^{(ext)}\kappa ({}^{(ext)}\zeta)^2 \right),\end{aligned}$$

which together with the above estimates yields

$$\sup_{\mathcal{C}_1} r^2 \left| {}^{(ext)}e_4({}^{(ext)}m) \right| \lesssim \epsilon_0^2.$$

This allows us to propagate the estimates for ${}^{(ext)}m$ in (4.1.4) on S_1 to any sphere on \mathcal{C}_1 , and hence

$$\sup_{\mathcal{C}_1} \left| \frac{{}^{(ext)}m}{m_0} - 1 \right| \lesssim \epsilon_0. \quad (4.1.7)$$

Also, we have

$$\begin{aligned} \lambda^{-1} e'_4 \left(\log \left(\frac{{}^{(ext)}r}{{}^{(ext)}r_{\mathcal{L}_0}} \right) \right) &= \frac{\lambda^{-1} {}^{(ext)}e_4({}^{(ext)}r)}{{}^{(ext)}r} - \frac{{}^{(ext)}(e_0)_4({}^{(ext)}r_{\mathcal{L}_0})}{{}^{(ext)}r_{\mathcal{L}_0}} - \frac{f^2}4 \frac{{}^{(ext)}(e_0)_3({}^{(ext)}r_{\mathcal{L}_0})}{{}^{(ext)}r} \\ &= \frac{1}{2} \left(\lambda^{-1} \kappa' - \kappa \right) + \frac{\lambda^{-1} \check{\kappa}'}2 - \frac{1}{2} \check{\kappa} - \frac{r f^2}{8} (\bar{\kappa} + \underline{A}) \\ &= \frac{1}{2} \left(\phi'_1(f) + \text{Err}(\kappa, \kappa') \right) + \frac{\lambda^{-1} \check{\kappa}'}2 - \frac{1}{2} \check{\kappa} - \frac{r f^2}{8} (\bar{\kappa} + \underline{A}) \end{aligned}$$

where we have denoted with primes the quantities with respect to the frame of ${}^{(ext)}\mathcal{M}$ and without primes the quantities with respect to the frame of ${}^{(ext)}\mathcal{L}_0$. Together with the above estimates, this yields

$$\sup_{\mathcal{C}_1} r^2 \left| \left(\log \left(\frac{{}^{(ext)}r}{{}^{(ext)}r_{\mathcal{L}_0}} \right) \right) \right| \lesssim \epsilon_0.$$

This allows us to propagate the estimates for ${}^{(ext)}r - {}^{(ext)}r_{\mathcal{L}_0}$ in (4.1.4) on S_1 to any sphere on \mathcal{C}_1 , and hence

$$\sup_{\mathcal{C}_1} \left| {}^{(ext)}r - {}^{(ext)}r_{\mathcal{L}_0} \right| \lesssim \epsilon_0.$$

Step 8. Recall that

- $({}^{(ext)}e_4, {}^{(ext)}e_3, {}^{(ext)}e_\theta)$ denotes the null frame of ${}^{(ext)}\mathcal{M}$,
- $({}^{(int)}e_4, {}^{(int)}e_3, {}^{(ext)}e_\theta)$ denotes the null frame of ${}^{(int)}\mathcal{M}$,
- $({}^{(ext)}(e_0)_3, {}^{(ext)}(e_0)_4, {}^{(ext)}(e_0)_\theta)$ denotes the null frame of ${}^{(ext)}\mathcal{L}_0$,
- $({}^{(int)}(e_0)_3, {}^{(int)}(e_0)_4, {}^{(int)}(e_0)_\theta)$ denotes the null frame of ${}^{(int)}\mathcal{L}_0$.

Also, recall that the timelike hyper surface \mathcal{T} is given by

$$\mathcal{T} = \{({}^{ext})r = r_{\mathcal{T}}\} \text{ where } 2m_0 \left(1 + \frac{\delta_{\mathcal{H}}}{2}\right) \leq r_{\mathcal{T}} \leq 2m_0 \left(1 + \frac{3\delta_{\mathcal{H}}}{2}\right)$$

to that $\mathcal{T} \subset ({}^{int})\mathcal{L}_0 \cap ({}^{ext})\mathcal{L}_0$, and recall that the frame of $({}^{int})\mathcal{M}$ is initialed on \mathcal{T} as follows

$$({}^{int})e_4 = \lambda ({}^{ext})e_4, \quad ({}^{int})e_3 = \lambda^{-1} ({}^{ext})e_3, \quad ({}^{int})e_{\theta} = ({}^{ext})e_{\theta} \text{ on } \mathcal{T}$$

where

$$\lambda = ({}^{ext})\lambda = 1 - \frac{2({}^{ext})m}{({}^{ext})r}.$$

Denoting

- by $(f, \underline{f}, \lambda)$ the transition functions from the frame of the outgoing part $({}^{ext})\mathcal{L}_0$ of the initial data layer to the frame of $({}^{ext})\mathcal{M}$ as in Steps 5 to 7,
- by $(f', \underline{f}', \lambda')$ the transition functions from the frame of the ingoing part $({}^{int})\mathcal{L}_0$ of the initial data layer to the frame of $({}^{int})\mathcal{M}$,
- by $(\tilde{f}, \tilde{\underline{f}}, \tilde{\lambda})$ the transition functions on $({}^{int})\mathcal{L}_0 \cap ({}^{ext})\mathcal{L}_0$ from the frame outgoing part $({}^{ext})\mathcal{L}_0$ of the initial data layer to the frame of the ingoing part $({}^{int})\mathcal{L}_0$ of the initial data layer,

we obtain, using also that $\mathcal{C}_1 \cap \underline{\mathcal{C}}_1 \subset \mathcal{T}$,

$$\begin{aligned} \sup_{\mathcal{C}_1 \cap \underline{\mathcal{C}}_1} \left(|\mathfrak{d}^{\leq k_{large}+3}(f', \underline{f}', \log(\lambda'))| \right) &\lesssim \sup_{\mathcal{C}_1 \cap \underline{\mathcal{C}}_1} \left(|\mathfrak{d}^{\leq k_{large}+3}(f, \underline{f}, \log(\lambda))| \right) \\ &+ \sup_{\mathcal{C}_1 \cap \underline{\mathcal{C}}_1} \left(|\mathfrak{d}^{\leq k_{large}+3}(\tilde{f}, \tilde{\underline{f}}, \log(\Upsilon_0^{-1}\tilde{\lambda}))| \right) \\ &+ \sup_{\mathcal{C}_1 \cap \underline{\mathcal{C}}_1} \left(|\mathfrak{d}^{\leq k_{large}+3} \log(\Upsilon_0^{-1}\Upsilon)| \right) \end{aligned}$$

where we have denoted

$$\Upsilon_0 = 1 - \frac{2m_0}{({}^{ext})r_{\mathcal{L}_0}}, \quad \Upsilon = 1 - \frac{2({}^{ext})m}{({}^{ext})r}.$$

Together with the control of $(\tilde{f}, \tilde{\underline{f}}, \log(\Upsilon_0^{-1}\tilde{\lambda}))$ provided on $({}^{int})\mathcal{L}_0 \cap ({}^{ext})\mathcal{L}_0$ by the estimates (4.1.1), the estimates (4.1.5) for $(f, \underline{f}, \lambda)$, and the estimates $({}^{ext})m - m_0$ and $({}^{ext})r - ({}^{ext})r_{\mathcal{L}_0}$ obtained in Step 7, we infer

$$\sup_{\mathcal{C}_1 \cap \underline{\mathcal{C}}_1} \left(|\mathfrak{d}^{\leq k_{large}+3}(f', \underline{f}', \log(\lambda'))| \right) \lesssim \epsilon_0.$$

Step 9. Similarly to Step 6, we propagate the estimate for $(f', \underline{f}', \log(\lambda))$ on $\mathcal{C}_1 \cap \underline{\mathcal{C}}_1$ provided by Step 8 to $\underline{\mathcal{C}}_1$ using the analog of Corollary 2.3.7 in the ingoing direction e_3 . We obtain the following analog of (4.1.5)

$$\sup_{\underline{\mathcal{C}}_1} \left(|\mathfrak{d}^{\leq k_{large}+3}(\underline{f}, \log \lambda)| \right) + \sup_{\mathcal{C}_1} \left(r |\mathfrak{d}^{\leq k_{large}+2} f| \right) \lesssim \epsilon_0.$$

Together with the change of frame formulas of Proposition 2.3.4, and the estimates (4.1.1) for the Ricci coefficients and curvature components of the ingoing part ${}^{(int)}\mathcal{L}_0$ of the initial data layer, we obtain

$$\begin{aligned} \max_{0 \leq k \leq k_{large}} \sup_{\underline{\mathcal{C}}_1} \left[|\mathfrak{d}^k {}^{(int)}\alpha| + |\mathfrak{d}^k {}^{(int)}\beta| + \left| \mathfrak{d}^k \left({}^{(int)}\rho + \frac{2m_0}{r^3} \right) \right| \right. \\ \left. + |\mathfrak{d}^k {}^{(int)}\underline{\beta}| + |\mathfrak{d}^k {}^{(int)}\underline{\alpha}| \right] \lesssim \epsilon_0, \quad (4.1.8) \end{aligned}$$

as well as

$$\max_{0 \leq k \leq k_{large}} \left\{ \sup_{\underline{\mathcal{C}}_1} |\mathfrak{d}^k {}^{(int)}(\check{\kappa}, \underline{\vartheta}, \zeta, \check{\kappa}, \vartheta)| \right\} \lesssim \epsilon_0.$$

Also, since we have as a consequence of the initialization on \mathcal{T} of the ingoing geodesic foliation of ${}^{(int)}\mathcal{M}$

$${}^{(int)}m = {}^{(ext)}m \quad \text{on } \mathcal{C}_1 \cap \underline{\mathcal{C}}_1$$

we infer from the control of ${}^{(ext)}m$ provided by Step 7

$$|{}^{(int)}m - m_0| \lesssim \epsilon_0 \quad \text{on } \mathcal{C}_1 \cap \underline{\mathcal{C}}_1.$$

We then propagate, similarly to Step 7, this bound to $\underline{\mathcal{C}}_1$ and obtain

$$\sup_{\underline{\mathcal{C}}_1} |{}^{(int)}m - m_0| \lesssim \epsilon_0.$$

Together with (4.1.6), (4.1.7) and (4.1.8), this concludes the proof of Theorem M0.

4.2 Control of averages and of the Hawking mass

In this section, we prove Lemma 3.4.1 and Lemma 3.4.2.

4.2.1 Proof of Lemma 3.4.1

Step 1. We start with the control of $\bar{\rho}$ on \mathcal{M} . Recall the identity (2.2.12)

$$\bar{\rho} + \frac{2m}{r^3} = \frac{1}{4} \overline{\vartheta \underline{\vartheta}}.$$

Thus, in view of the bootstrap assumptions **BA-D**, **BA-E**, we have,

$$\begin{aligned} \left| \bar{\rho} + \frac{2m}{r^3} \right| &\lesssim \epsilon^2 \min\{r^{-3} \underline{u}^{-\frac{3}{2}-\delta_{dec}}, r^{-2} \underline{u}^{-2-2\delta_{dec}}\} \quad \text{in } {}^{(ext)}\mathcal{M}, \\ \left| \bar{\rho} + \frac{2m}{r^3} \right| &\lesssim \epsilon^2 \underline{u}^{-2-2\delta_{dec}} \quad \text{in } {}^{(int)}\mathcal{M}. \end{aligned}$$

Differentiating the equation with respect to e_3, e_4 we derive,

$$\begin{aligned} e_4 \left(\bar{\rho} + \frac{2m}{r^3} \right) &= \frac{1}{4} \overline{e_4(\vartheta) \underline{\vartheta} + \vartheta e_4(\underline{\vartheta})} + \text{l.o.t.}, \\ e_3 \left(\bar{\rho} + \frac{2m}{r^3} \right) &= \frac{1}{4} \overline{e_3(\vartheta) \underline{\vartheta} + \vartheta e_3(\underline{\vartheta})} + \text{l.o.t.}, \\ e_\theta \left(\bar{\rho} + \frac{2m}{r^3} \right) &= 0. \end{aligned}$$

Taking higher derivatives in e_3, e_4 and making use of the bootstrap assumptions **BA-D**, **BA-E**, we derive in ${}^{(ext)}\mathcal{M}$,

$$\begin{aligned} \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| &\lesssim \epsilon^2 \min\{r^{-3} \underline{u}^{-\frac{3}{2}-\delta_{dec}}, r^{-2} \underline{u}^{-2-2\delta_{dec}}\}, \\ \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| &\lesssim r^{-3} \underline{u}^{-1/2-\delta_{dec}}, \end{aligned}$$

and in ${}^{(int)}\mathcal{M}$,

$$\begin{aligned} \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| &\lesssim \epsilon^2 \underline{u}^{-2-2\delta_{dec}}, \\ \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| &\lesssim \epsilon^2 \underline{u}^{-1-\delta_{dec}}. \end{aligned}$$

In particular,

$$\begin{aligned} \sup_{{}^{(ext)}\mathcal{M}} \underline{u}^{\frac{3}{2}+\delta_{dec}} r^3 \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| + \sup_{{}^{(ext)}\mathcal{M}} \underline{u}^{\frac{1}{2}+\delta_{dec}} r^3 \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| &\lesssim \epsilon_0, \\ \sup_{{}^{(int)}\mathcal{M}} \underline{u}^{\frac{3}{2}+\delta_{dec}} \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| + \sup_{{}^{(int)}\mathcal{M}} \underline{u}^{\frac{1}{2}+\delta_{dec}} \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\rho} + \frac{2m}{r^3} \right) \right| &\lesssim \epsilon_0 \end{aligned}$$

Step 2. Next, we proceed with the control of $\bar{\kappa}$ in ${}^{(ext)}\mathcal{M}$. Recalling Lemma 2.2.17, we start with

$$e_4 \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{1}{2\bar{\kappa}} \left(\bar{\kappa} - \frac{2}{r} \right) = -\frac{1}{4}\bar{\vartheta}^2 + \frac{1}{2}\bar{\kappa}^2. \quad (4.2.1)$$

In view of Corollary 2.2.12 we deduce, from the first equation,

$$e_4 \left(r \left(\bar{\kappa} - \frac{2}{r} \right) \right) = -r \left(\frac{1}{4}\bar{\vartheta}^2 + \frac{1}{2}\bar{\kappa}^2 \right). \quad (4.2.2)$$

Making use of the GCM condition

$$\kappa = \frac{2}{r} \text{ on } \Sigma_*,$$

which yields

$$\bar{\kappa} = \frac{2}{r} \text{ on } \Sigma_*,$$

we deduce, integrating (4.2.2) with respect to r along C_u from Σ_* ,

$$\sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} r^3 \left| \bar{\kappa} - \frac{2}{r} \right| \lesssim \epsilon^2 \lesssim \epsilon_0.$$

Also, making use of the bootstrap assumptions **BA-D**, **BA-E** we easily deduce,

$$\begin{aligned} \sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} r^3 \left| \mathfrak{D}_{\nearrow}^{\leq k_{small}+1} \left(\bar{\kappa} - \frac{2}{r} \right) \right| &\lesssim \epsilon^2 \lesssim \epsilon_0, \\ \sup_{(ext)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}} r^3 \left| \mathfrak{D}_{\nearrow}^{\leq k_{large}+1} \left(\bar{\kappa} - \frac{2}{r} \right) \right| &\lesssim \epsilon^2 \lesssim \epsilon_0. \end{aligned}$$

We next commute (4.2.2) with e_3 and derive,

$$\begin{aligned} e_4 e_3 \left(r \left(\bar{\kappa} - \frac{2}{r} \right) \right) &= e_3 \left(r \left(\frac{1}{4}\bar{\vartheta}^2 + \frac{1}{2}\bar{\kappa}^2 \right) \right) - [e_3, e_4] \left(r \left(\bar{\kappa} - \frac{2}{r} \right) \right) \\ &= e_3 \left(r \left(\frac{1}{4}\bar{\vartheta}^2 + \frac{1}{2}\bar{\kappa}^2 \right) \right) - 2\underline{\omega} \left(r \left(\bar{\kappa} - \frac{2}{r} \right) \right) - 2\zeta \left(r \left(\bar{\kappa} - \frac{2}{r} \right) \right). \end{aligned}$$

It is thus easy to see that we can prove estimates of the type

$$\begin{aligned} \sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} r^3 \left| \mathfrak{D}_{\nearrow}^{\leq k_{small}+1} \left(\bar{\kappa} - \frac{2}{r} \right) \right| &\lesssim \epsilon^2 \lesssim \epsilon_0, \\ \sup_{(ext)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}} r^3 \left| \mathfrak{D}_{\nearrow}^{\leq k_{large}+1} \left(\bar{\kappa} - \frac{2}{r} \right) \right| &\lesssim \epsilon^2 \lesssim \epsilon_0, \end{aligned}$$

provided that we can check that,

$$\begin{aligned} \sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} r^3 \left| e_3^{\leq k_{small}+1} \left(\bar{\kappa} - \frac{2}{r} \right) \right| &\lesssim \epsilon^2 \lesssim \epsilon_0, \\ \sup_{(ext)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}} r^3 \left| e_3^{\leq k_{large}+1} \left(\bar{\kappa} - \frac{2}{r} \right) \right| &\lesssim \epsilon^2 \lesssim \epsilon_0. \end{aligned}$$

The difficulty in this case is to make sure that we can control terms of the type,

$$e_3^{k+1} \left(r \left(\frac{1}{4} e_3^{k+1}(\vartheta^2) + \frac{1}{2} e_3^{k+1}(\check{\kappa}^2) \right) \right)$$

using only at most k derivatives of $\check{\Gamma}, \check{R}$. To see this we note that,

$$\begin{aligned} e_3(\vartheta^2) &= \overline{e_3 \vartheta^2} - (\check{\underline{\Omega}} \bar{\kappa} - \check{\underline{\Omega}} \check{\kappa}) \overline{\vartheta^2} + \check{\underline{\kappa}} \overline{\vartheta^2}, \\ e_3(\check{\kappa}^2) &= \overline{e_3 \check{\kappa}^2} - (\check{\underline{\Omega}} \bar{\kappa} - \check{\underline{\Omega}} \check{\kappa}) \overline{\check{\kappa}^2} + \check{\underline{\kappa}} \overline{\check{\kappa}^2}, \end{aligned}$$

and,

$$\begin{aligned} e_3(\vartheta) + \frac{1}{2} \underline{\kappa} \vartheta - 2 \underline{\omega} \vartheta &= -2 \not{d}_2^* \zeta - \frac{1}{2} \underline{\kappa} \vartheta + 2 \zeta^2, \\ e_3 \check{\kappa} + \frac{1}{2} \bar{\kappa} \check{\kappa} &= -2 \check{\mu} - \frac{1}{2} \bar{\kappa} \check{\kappa} + 2(\check{\underline{\omega}} \bar{\kappa} + \bar{\omega} \check{\kappa}) + \check{\underline{\Omega}} \bar{\kappa} \bar{\kappa} + \text{Err}[e_3 \check{\kappa}], \\ \text{Err}[e_3 \check{\kappa}] &:= 2(\zeta^2 - \check{\zeta}^2) + 2(\check{\underline{\omega}} \check{\kappa} - \bar{\omega} \bar{\kappa}) - \frac{1}{2} \check{\underline{\kappa}} \check{\kappa} - \frac{1}{2} \bar{\kappa} \check{\kappa} - \check{\underline{\Omega}} \bar{\kappa} \bar{\kappa}. \end{aligned} \tag{4.2.3}$$

We thus derive,

$$\sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} r^3 \left| \mathfrak{d}^{\leq k_{small}+1} \left(\bar{\kappa} - \frac{2}{r} \right) \right| + \sup_{(ext)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}} r^3 \left| \mathfrak{d}^{\leq k_{large}+1} \left(\bar{\kappa} - \frac{2}{r} \right) \right| \lesssim \epsilon_0.$$

Step 3. We next estimate $\bar{\kappa}$ in $(ext)\mathcal{M}$ making use of the identity (2.2.14) derived in connection to the Hawking mass

$$\bar{\kappa} + \frac{2\Upsilon}{r} = \frac{2\Upsilon}{r\bar{\kappa}} \left(\bar{\kappa} - \frac{2}{r} \right) - \frac{1}{\bar{\kappa}} \check{\underline{\kappa}} \check{\underline{\kappa}}.$$

Thus, in view of the estimates for $\bar{\kappa}$ derived in step 2 we easily infer that,

$$\sup_{(ext)\mathcal{M}} u^{\frac{3}{2}+\delta_{dec}} r^2 \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\kappa} + \frac{2\Upsilon}{r} \right) \right| + \sup_{(ext)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}} r^2 \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\kappa} + \frac{2\Upsilon}{r} \right) \right| \lesssim \epsilon_0.$$

as desired.

Step 4. We estimate $\underline{\bar{\omega}}$ in ${}^{(ext)}\mathcal{M}$ based on the following identity in Lemma 2.2.17

$$\begin{aligned} e_3 \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{1}{2\bar{\kappa}} \left(\bar{\kappa} - \frac{2}{r} \right) &= 2\underline{\bar{\omega}} \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{4}{r} \left(\underline{\bar{\omega}} - \frac{m}{r^2} \right) + 2 \left(\bar{\rho} + \frac{2m}{r^3} \right) \\ &\quad - \frac{1}{2\bar{\kappa}} \left(\bar{\kappa} - \frac{2}{r} \right) \check{\underline{\Omega}} + 2\underline{\bar{\omega}}\check{\kappa} - \frac{1}{2}\vartheta\underline{\vartheta} + 2\bar{\zeta}^2 \\ &\quad + \frac{1}{2}\check{\underline{\Omega}} \left(-\vartheta^2 + \check{\kappa}^2 \right) - \overline{\check{\underline{\Omega}}(e_4(\check{\kappa}) + \kappa\check{\kappa})} + \frac{1}{2}\overline{\check{\kappa}\check{\kappa}} - \frac{1}{r}\overline{\check{\underline{\Omega}}\check{\kappa}}, \end{aligned}$$

which we rewrite as

$$\begin{aligned} \underline{\bar{\omega}} - \frac{m}{r^2} &= \frac{r}{4} \left\{ e_3 \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{1}{2\bar{\kappa}} \left(\bar{\kappa} - \frac{2}{r} \right) - 2\underline{\bar{\omega}} \left(\bar{\kappa} - \frac{2}{r} \right) - 2 \left(\bar{\rho} + \frac{2m}{r^3} \right) \right. \\ &\quad \left. + \frac{1}{2\bar{\kappa}} \left(\bar{\kappa} - \frac{2}{r} \right) \check{\underline{\Omega}} - 2\underline{\bar{\omega}}\check{\kappa} + \frac{1}{2}\vartheta\underline{\vartheta} - 2\bar{\zeta}^2 - \frac{1}{2}\check{\underline{\Omega}} \left(-\vartheta^2 + \check{\kappa}^2 \right) \right. \\ &\quad \left. + \overline{\check{\underline{\Omega}}(e_4(\check{\kappa}) + \kappa\check{\kappa})} - \frac{1}{2}\overline{\check{\kappa}\check{\kappa}} + \frac{1}{r}\overline{\check{\underline{\Omega}}\check{\kappa}} \right\}. \end{aligned}$$

Using the estimates of $\bar{\rho}$ in Step 1, the estimates for $\bar{\kappa}$ in Step 2, as well as our bootstrap assumptions on decay and energy, we easily derive

$$\sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}r^2} \left| \mathfrak{d}^{\leq k_{small}} \left(\underline{\bar{\omega}} - \frac{m}{r^2} \right) \right| + \sup_{(ext)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}r^2} \left| \mathfrak{d}^{\leq k_{large}} \left(\underline{\bar{\omega}} - \frac{m}{r^2} \right) \right| \lesssim \epsilon_0.$$

Remark 4.2.1. *It is to estimate k_{large} derivatives of $\underline{\bar{\omega}} - mr^{-2}$ that we had to control $k_{large} + 1$ derivatives of $\bar{\kappa} - 2/r$ in Step 2.*

Step 5. We estimate $\underline{\bar{\Omega}}$ in ${}^{(ext)}\mathcal{M}$. First we need the control of $\underline{\bar{\Omega}}$ on Σ_* . To this end, we recall that s is initialized on Σ_* by $s = r$ so that

$$\nu(s - r) = 0 \text{ on } \Sigma_*, \quad \nu = e_3 + ae_4,$$

where the scalar function a is such that the vectorfield ν is tangent to Σ_* . On the other hand, we have $e_4(s) = 1$ and

$$e_4(r) = \frac{r}{2\bar{\kappa}} = 1 \text{ on } \Sigma_*$$

where we used the GCM condition $\kappa = 2/r$ on Σ_* . We infer $e_3(s) = e_3(r)$ on Σ_* and hence

$$\underline{\bar{\Omega}} = e_3(r) \text{ on } \Sigma_*.$$

This yields

$$\bar{\Omega} = \overline{e_3(r)} = \frac{r\bar{\kappa}}{2} + \bar{A},$$

and hence, in view of the estimate for $\bar{\kappa}$ of step 3, the fact that \bar{A} contains only quadratic terms in view of the formula for \bar{A} , and in view of the bootstrap assumptions on decay and energy, we infer

$$\sup_{\Sigma_*} u^{1+\delta_{dec}r} \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\Omega} - \frac{m}{r^2} \right) \right| + \sup_{\Sigma_*} u^{\frac{1}{2}+\delta_{dec}r} \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\Omega} - \frac{m}{r^2} \right) \right| \lesssim \epsilon_0.$$

Then, we use $e_4(\bar{\Omega}) = -2\bar{\omega}$ and Corollary 2.2.11 to obtain

$$e_4(\bar{\Omega}) = -2\bar{\omega} + \bar{\kappa}\bar{\check{\Omega}}$$

and hence

$$e_4(\bar{\Omega} + \Upsilon) = -2 \left(\bar{\omega} - \frac{m}{r^2} \right) + \frac{m}{r} \left(\bar{\kappa} - \frac{2}{r} \right) + \bar{\kappa}\bar{\check{\Omega}} - \frac{2e_4(m)}{r}.$$

Commuting with \mathfrak{d} , integrating from Σ_* where we have controlled $\bar{\Omega}$ above, and using the estimates of Step 2 for $\bar{\kappa}$, Step 4 for $\bar{\omega}$, the bootstrap assumptions, and the estimates for $e_4(m)$ of Lemma 3.4.2 (which do not depend on the control of $\bar{\Omega}$), we infer

$$\sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}r} \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\Omega} - \frac{m}{r^2} \right) \right| + \sup_{(ext)\mathcal{M}} u^{\frac{1}{2}+\delta_{dec}r} \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\Omega} - \frac{m}{r^2} \right) \right| \lesssim \epsilon_0.$$

Step 6. Next, we control ${}^{(int)}\bar{\kappa}$ on the cylinder \mathcal{T} . From the initialization of the frame of ${}^{(int)}\mathcal{M}$ on \mathcal{T} , we have

$${}^{(int)}r = {}^{(ext)}r, \quad {}^{(int)}\bar{\kappa} = \Upsilon {}^{(ext)}\bar{\kappa}, \quad {}^{(int)}\bar{\check{\kappa}} = \Upsilon^{-1} {}^{(ext)}\bar{\check{\kappa}} \text{ on } \mathcal{T}.$$

Also, making use of the identity (2.2.14) derived in connection to the Hawking mass, we have

$${}^{(ext)}\bar{\check{\kappa}} + \frac{2\Upsilon}{r} = \frac{2\Upsilon}{r {}^{(ext)}\bar{\kappa}} \left({}^{(ext)}\bar{\kappa} - \frac{2}{r} \right) - \frac{1}{{}^{(ext)}\bar{\kappa}} \overline{{}^{(ext)}\check{\kappa} {}^{(ext)}\check{\kappa}}.$$

We deduce

$${}^{(int)}\bar{\check{\kappa}} + \frac{2}{r} = \Upsilon^{-1} \left({}^{(ext)}\bar{\check{\kappa}} + \frac{2\Upsilon}{r} \right) = \frac{2}{r {}^{(ext)}\bar{\kappa}} \left({}^{(ext)}\bar{\kappa} - \frac{2}{r} \right) - \frac{\Upsilon^{-1}}{{}^{(ext)}\bar{\kappa}} \overline{{}^{(ext)}\check{\kappa} {}^{(ext)}\check{\kappa}} \text{ on } \mathcal{T}.$$

To derive higher tangential derivatives along \mathcal{T} we remark that the vectorfield

$$T_{\mathcal{T}} = e_4 - \frac{e_4(r)}{e_3(r)} e_3 = e_4 - \frac{\bar{\kappa} + A}{\bar{\kappa}} e_3,$$

together with e_{θ} , spans the tangent space to \mathcal{T} . The transversal derivatives, on the other hand, can be determined with help of the equation,

$$e_3 \left(\frac{\bar{\kappa}}{\kappa} + \frac{2}{r} \right) + \frac{1}{2\bar{\kappa}} \left(\frac{\bar{\kappa}}{\kappa} + \frac{2}{r} \right) = -\frac{1}{4} \frac{\bar{\vartheta}^2}{\bar{\kappa}} + \frac{1}{2} \frac{\bar{\kappa}^2}{\bar{\kappa}}.$$

adapted to the $(int)\mathcal{M}$ foliation. Making use of the estimates for $(ext)\bar{\kappa}$ in $(ext)\mathcal{M}$ derived in Step 2 and the bootstrap assumptions, we infer that,

$$\begin{aligned} & \sup_{\mathcal{T}} u^{\frac{3}{2}+\delta_{dec}} \left| \mathfrak{D}^{\leq k_{small}+1} \left((int)\frac{\bar{\kappa}}{\kappa} + \frac{2}{r} \right) \right| + \sup_{\mathcal{T}} u^{\frac{1}{2}+\delta_{dec}} \left| \mathfrak{D}^{\leq k_{large}+1} \left((int)\frac{\bar{\kappa}}{\kappa} + \frac{2}{r} \right) \right| \\ & \lesssim \epsilon_0 + \sup_{\mathcal{T}} u^{\frac{1}{2}+\delta_{dec}} \left| \mathfrak{D}^{k_{large}+1} \left(\overline{(ext)\check{\kappa} (ext)\check{\kappa}} \right) \right| \end{aligned}$$

Now, in view of the transport equations for $(ext)e_4((ext)\check{\kappa})$, $(ext)e_3((ext)\check{\kappa})$, $(ext)e_4((ext)\check{\kappa})$ and $(ext)e_3((ext)\check{\kappa})$, as well as the bootstrap assumptions, we have

$$\begin{aligned} & \sup_{\mathcal{T}} u^{\frac{1}{2}+\delta_{dec}} \left| \mathfrak{D}^{k_{large}+1} \left(\overline{(ext)\check{\kappa} (ext)\check{\kappa}} \right) \right| \\ & \lesssim \epsilon_0 + \sup_{\mathcal{T}} u^{\frac{1}{2}+\delta_{dec}} \left| \overline{(ext)\check{\kappa} \mathfrak{D}^{k_{large}} \mathfrak{d}_1((ext)\check{\zeta})} \right| + \sup_{\mathcal{T}} u^{\frac{1}{2}+\delta_{dec}} \left| \overline{(ext)\check{\kappa} \mathfrak{D}^{k_{large}} \mathfrak{d}_1((ext)\check{\zeta})} \right| \\ & \quad + \sup_{\mathcal{T}} u^{\frac{1}{2}+\delta_{dec}} \left| \overline{(ext)\check{\kappa} \mathfrak{D}^{k_{large}} \mathfrak{d}_1((ext)\check{\xi})} \right| \\ & \lesssim \epsilon_0 + \sup_{\mathcal{T}} u^{\frac{1}{2}+\delta_{dec}} \left| \overline{\mathfrak{d}_1^* (ext)\check{\kappa} \mathfrak{D}^{k_{large}} (ext)\check{\zeta}} \right| + \sup_{\mathcal{T}} u^{\frac{1}{2}+\delta_{dec}} \left| \overline{\mathfrak{d}_1^* (ext)\check{\kappa} \mathfrak{D}^{k_{large}} (ext)\check{\zeta}} \right| \\ & \quad + \sup_{\mathcal{T}} u^{\frac{1}{2}+\delta_{dec}} \left| \overline{\mathfrak{d}_1^* (ext)\check{\kappa} \mathfrak{D}^{k_{large}} (ext)\check{\xi}} \right| \\ & \lesssim \epsilon_0 \end{aligned}$$

where we have integrated \mathfrak{d}_1 by parts and used that \mathfrak{d}_1^* is its adjoint. We infer

$$\sup_{\mathcal{T}} u^{\frac{3}{2}+\delta_{dec}} \left| \mathfrak{D}^{\leq k_{small}+1} \left((int)\frac{\bar{\kappa}}{\kappa} + \frac{2}{r} \right) \right| + \sup_{\mathcal{T}} u^{\frac{1}{2}+\delta_{dec}} \left| \mathfrak{D}^{\leq k_{large}+1} \left((int)\frac{\bar{\kappa}}{\kappa} + \frac{2}{r} \right) \right| \lesssim \epsilon_0.$$

Step 7. From now on, we only work with the frame of $(int)\mathcal{M}$. Starting with the equation,

$$e_3 \left(\frac{\bar{\kappa}}{\kappa} + \frac{2}{r} \right) + \frac{1}{2\bar{\kappa}} \left(\frac{\bar{\kappa}}{\kappa} + \frac{2}{r} \right) = -\frac{1}{4} \frac{\bar{\vartheta}^2}{\bar{\kappa}} + \frac{1}{2} \frac{\bar{\kappa}^2}{\bar{\kappa}}.$$

Using the estimates of step 5 we can then proceed precisely as in Step 2 (using the $(int)\mathcal{M}$ counterpart of the equations (4.2.3)) to derive,

$$\sup_{(int)\mathcal{M}} \underline{u}^{1+\delta_{dec}} \left| \mathfrak{d}^{\leq k_{small}+1} \left(\bar{\kappa} + \frac{2}{r} \right) \right| + \sup_{(int)\mathcal{M}} \underline{u}^{\frac{1}{2}+\delta_{dec}} \left| \mathfrak{d}^{\leq k_{large}+1} \left(\bar{\kappa} + \frac{2}{r} \right) \right| \lesssim \epsilon_0.$$

Step 8. Finally, we estimate the remaining averages in $(int)\mathcal{M}$, i.e. $\bar{\kappa}$ and $\bar{\omega}$. To estimate $\bar{\kappa}$ we make use once more of the identity,

$$\bar{\kappa} - \frac{2\Upsilon}{r} = -\frac{2\Upsilon}{r\bar{\kappa}} \left(\bar{\kappa} + \frac{2}{r} \right) - \frac{1}{\bar{\kappa}} \check{\kappa}\check{\kappa}.$$

Making use of the estimates of $\bar{\kappa}$ in Step 5 as well as the bootstrap assumptions for $\check{\kappa}$ and $\check{\kappa}$ we easily derive,

$$\sup_{(int)\mathcal{M}} \underline{u}^{1+\delta_{dec}} \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\kappa} - \frac{2\Upsilon}{r} \right) \right| + \sup_{(int)\mathcal{M}} \underline{u}^{\frac{1}{2}+\delta_{dec}} \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\kappa} - \frac{2\Upsilon}{r} \right) \right| \lesssim \epsilon_0.$$

Step 9. To estimate $\bar{\omega}$ we proceed as in Step 4 by making use of the identity

$$\begin{aligned} \bar{\omega} + \frac{m}{r^2} &= \frac{r}{4} \left\{ e_4 \left(\bar{\kappa} + \frac{2}{r} \right) + \frac{1}{2} \bar{\kappa} \left(\bar{\kappa} + \frac{2}{r} \right) - 2\bar{\omega} \left(\bar{\kappa} + \frac{2}{r} \right) - 2 \left(\bar{\rho} + \frac{2m}{r^3} \right) \right. \\ &\quad + \frac{1}{2} \bar{\kappa} \left(\bar{\kappa} + \frac{2}{r} \right) \check{\Omega} - 2\bar{\omega}\check{\kappa} + \frac{1}{2} \vartheta\check{\vartheta} - 2\check{\zeta}^2 - \frac{1}{2} \check{\Omega} \left(-\check{\vartheta}^2 + \check{\kappa}^2 \right) \\ &\quad \left. + \overline{\check{\Omega}(e_4(\check{\kappa}) + \check{\kappa}\check{\kappa})} - \frac{1}{2} \check{\kappa}\check{\kappa} + \frac{1}{r} \check{\Omega}\check{\kappa} \right\}. \end{aligned}$$

Thus, in view of the estimates of $\bar{\rho}$ in Step 1, the estimates for $\bar{\kappa}$ in Step 5, the estimates of $\bar{\kappa}$ above², as well as the bootstrap assumptions **BA-D** and **BA-E**, we deduce,

$$\sup_{(int)\mathcal{M}} \underline{u}^{1+\delta_{dec}} \left| \mathfrak{d}^{\leq k_{small}} \left(\bar{\omega} + \frac{m}{r^2} \right) \right| + \sup_{(int)\mathcal{M}} \underline{u}^{\frac{1}{2}+\delta_{dec}} \left| \mathfrak{d}^{\leq k_{large}} \left(\bar{\omega} + \frac{m}{r^2} \right) \right| \lesssim \epsilon_0.$$

Step 10. It remains to estimate $\bar{\Omega}$ in $(int)\mathcal{M}$. First we need the control of $\bar{\Omega}$ on \mathcal{T} . To this end, we recall that s is initialized on \mathcal{T} by $s = r$ so that

$$T_{\mathcal{T}}(s-r) = 0 \text{ on } \mathcal{T}, \quad T_{\mathcal{T}} = e_4 - \frac{\bar{\kappa} + A}{\bar{\kappa}} e_3,$$

²It is to estimate k_{large} derivatives of $\bar{\omega} + m/r^2$ that we made sure to control $k_{large} + 1$ derivatives of $\bar{\kappa} + 2/r$.

, where the vectorfield has been introduced above and is tangent to \mathcal{T} . On the other hand, we have $e_3(s) = -1$ and $e_3(r) = r\bar{\kappa}/2$, and hence

$$\Omega = e_4(r) + \frac{\bar{\kappa} + A}{\bar{\kappa}}(-1 - e_3(r)) = \frac{r}{2}(\bar{\kappa} + A) \left(1 + \frac{\bar{\kappa} - \frac{2}{r}}{\bar{\kappa}}\right) \text{ on } \mathcal{T}$$

This yields

$$\bar{\Omega} = \frac{r}{2}(\bar{\kappa} + \bar{A}) \left(1 + \frac{\bar{\kappa} - \frac{2}{r}}{\bar{\kappa}}\right) \text{ on } \mathcal{T},$$

and hence, in view of the estimate for $\bar{\kappa}$ of step 7, the estimate for $\bar{\kappa}$ of step 8, the fact that \bar{A} contains only quadratic terms in view of the formula for A , and in view of the bootstrap assumptions on decay and energy, we infer

$$\sup_{\mathcal{T}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^{\leq k_{small}} (\bar{\Omega} - \Upsilon)| + \sup_{\mathcal{T}} \underline{u}^{\frac{1}{2}+\delta_{dec}} |\mathfrak{d}^{\leq k_{large}} (\bar{\Omega} - \Upsilon)| \lesssim \epsilon_0.$$

Then, we use the analog of the transport equation used to estimate $\bar{\Omega}$ in $^{(ext)}\mathcal{M}$, i.e.

$$e_3(\bar{\Omega} - \Upsilon) = 2 \left(\bar{\omega} + \frac{m}{r^2} \right) - \frac{m}{r} \left(\bar{\kappa} + \frac{2}{r} \right) + \bar{\kappa} \bar{\Omega} + \frac{2e_3(m)}{r}.$$

Commuting with \mathfrak{d} , integrating from \mathcal{T} where we have controlled $\bar{\Omega}$ above, and using the estimates of Step 2 for $\bar{\kappa}$, Step 4 for $\bar{\omega}$, the bootstrap assumptions, and the estimates for $e_4(m)$ of Lemma 3.4.2 (which do not depend on the control of $\bar{\Omega}$), we infer

$$\sup_{^{(int)}\mathcal{M}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^{\leq k_{small}} (\bar{\Omega} - \Upsilon)| + \sup_{^{(int)}\mathcal{M}} \underline{u}^{\frac{1}{2}+\delta_{dec}} |\mathfrak{d}^{\leq k_{large}} (\bar{\Omega} - \Upsilon)| \lesssim \epsilon_0.$$

This concludes the proof of Lemma 3.4.1.

4.2.2 Proof of Lemma 3.4.2

Step 1. We start with the control of $e_3(m)$ and $e_4(m)$ in $^{(ext)}\mathcal{M}$. According to Proposition 2.2.16 we have in $^{(ext)}\mathcal{M}$

$$e_4(m) = \frac{r}{32\pi} \int_S \text{Err}_1, \quad (4.2.4)$$

and

$$\begin{aligned}
e_3(m) &= (1 - \varsigma^{-1}\check{\zeta}) \frac{r}{32\pi} \int_S \underline{\text{Err}}_1 + (\check{\Omega} + \varsigma^{-1}\overline{\Omega}\check{\zeta}) \frac{r}{32\pi} \int_S \text{Err}_1 \\
&\quad + \varsigma^{-1} \frac{r}{32\pi} \int_S \check{\zeta} (2\overline{\rho}\check{\kappa} + 2\check{\rho}\overline{\kappa} + 2\check{\kappa}\phi_1\eta + 2\kappa\phi_1\underline{\xi} + \underline{\text{Err}}_2) \\
&\quad - \varsigma^{-1} \frac{r}{32\pi} \int_S (\overline{\Omega}\check{\zeta} + \check{\Omega}\varsigma) (2\overline{\rho}\check{\kappa} + 2\check{\rho}\overline{\kappa} - 2\kappa\phi_1\zeta + \text{Err}_2) \\
&\quad - \frac{m}{r} \varsigma^{-1} \left[-\check{\zeta}\check{\kappa} + \overline{\Omega}\check{\zeta}\overline{\kappa} + \check{\Omega}\varsigma\kappa \right], \tag{4.2.5}
\end{aligned}$$

where

$$\begin{aligned}
\text{Err}_1 &:= 2\check{\kappa}\check{\rho} + 2e_\theta(\kappa)\zeta - \frac{1}{2}\kappa\underline{\vartheta}^2 - \frac{1}{2}\check{\kappa}\vartheta\underline{\vartheta} + 2\kappa\zeta^2, \\
\underline{\text{Err}}_1 &:= 2\check{\rho}\check{\kappa} - 2e_\theta(\underline{\kappa})\eta - 2e_\theta(\kappa)\underline{\xi} - \frac{1}{2}\check{\kappa}\vartheta\underline{\vartheta} + 2\underline{\kappa}\eta^2 + 2\kappa(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\kappa\underline{\vartheta}^2, \\
\text{Err}_2 &:= 2\check{\rho}\check{\kappa} - \frac{1}{2}\kappa\underline{\vartheta}^2 - \frac{1}{2}\kappa\vartheta\underline{\vartheta} + 2\kappa\zeta^2, \\
\underline{\text{Err}}_2 &:= 2\check{\rho}\check{\kappa} + \underline{\kappa} \left(2\eta^2 - \frac{1}{2}\vartheta\underline{\vartheta} \right) + 2\kappa(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\kappa\underline{\vartheta}^2.
\end{aligned}$$

Thus, according to the bootstrap assumption **BA-D** on decay, we deduce,

$$\begin{aligned}
|e_4(m)| &\lesssim \epsilon^2 r^{-2} u^{-1-\delta_{dec}}, \\
|e_3(m)| &\lesssim \epsilon^2 u^{-2-2\delta_{dec}}.
\end{aligned}$$

Moreover, differentiating the equations with respect to e_3, e_4 and making use of both bootstrap assumptions **BA-D** **BA-E** on decay and energy, and integrating by part once the e_θ derivative for the terms involving $e_\theta(\kappa)$ and $e_\theta(\underline{\kappa})$ when they contain top order derivatives, we infer that,

$$\begin{aligned}
\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} r^2 u^{1+\delta_{dec}} |\mathfrak{d}^k e_4(m)| &\lesssim \epsilon_0, \\
\max_{0 \leq k \leq k_{large}} \sup_{(ext)\mathcal{M}} \left(r^2 u^{\frac{1}{2}+\delta_{dec}} + r u^{1+\delta_{dec}} \right) |\mathfrak{d}^k e_4(m)| &\lesssim \epsilon_0,
\end{aligned}$$

as well as

$$\begin{aligned}
\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} u^{2+2\delta_{dec}} |\mathfrak{d}^k e_3(m)| &\lesssim \epsilon_0, \\
\max_{0 \leq k \leq k_{large}} \sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} |\mathfrak{d}^k e_3(m)| &\lesssim \epsilon_0,
\end{aligned}$$

consistent with the statement of the lemma.

Step 2. We derive the estimates on ${}^{(int)}\mathcal{M}$. According to the analogue of Proposition 2.2.16 in the situation of the incoming geodesic foliations of ${}^{(int)}\mathcal{M}$, and proceeding as in Step 1, we easily derive,

$$\max_{0 \leq k \leq k_{large}} \sup_{(int)\mathcal{M}} \underline{u}^{1+\delta_{dec}} \left(|\mathfrak{d}^k e_3(m)| + |\mathfrak{d}^k e_4(m)| \right) \lesssim \epsilon^2 \lesssim \epsilon_0. \quad (4.2.6)$$

Step 3. We estimate $m - m_0$ in ${}^{(ext)}\mathcal{M}$.

First, recall from Theorem M0 that we have

$$\sup_{\mathcal{C}_1 \cup \underline{\mathcal{C}}_1} |m - m_0| \lesssim \epsilon_0 m_0. \quad (4.2.7)$$

We start with the control in ${}^{(ext)}\mathcal{M}$. Note that ${}^{(ext)}\mathcal{M}$ is covered by integral curves of e_3 starting from \mathcal{C}_1 . Thus, integrating the $e_3 m$ equation and making use of the estimate $\sup_{\mathcal{C}_1} |m - m_0| \lesssim \epsilon_0 m_0$ as well as the fact that $e_3(u) = 2$, we easily deduce that,

$$\sup_{(ext)\mathcal{M}} |m - m_0| \lesssim \epsilon_0 m_0 + \epsilon^2 \lesssim \epsilon_0 m_0.$$

Step 4. We estimate $|m - m_0|$ on \mathcal{T} .

In view of our initialization of the ingoing geodesic foliation of ${}^{(int)}\mathcal{M}$ on \mathcal{T} ,

$${}^{(int)}\underline{\kappa} = {}^{(ext)}\underline{\kappa} \text{ on } \mathcal{T}.$$

Since the spheres of both foliations agree on \mathcal{T} , we infer from the definition of the Hawking mass,

$${}^{(int)}m = {}^{(ext)}m \text{ on } \mathcal{T}.$$

Using the estimate for ${}^{(ext)}m$ we infer that

$$\sup_{\mathcal{T}} |{}^{(int)}m - m_0| \lesssim \epsilon_0 m_0.$$

Step 5. We estimate $|m - m_0|$ on ${}^{(int)}\mathcal{M}$.

Note first that in ${}^{(int)}\mathcal{M}$,

$$e_3(r) + 1 = \frac{r}{2} \bar{\kappa} + 1 = \frac{r}{2} \left(\bar{\kappa} + \frac{2}{r} \right).$$

Thus, in view of the estimate for $\bar{\kappa} + \frac{2}{r}$ derived in Lemma 3.4.1

$$\sup_{(int)\mathcal{M}} |e_3(r) + 1| \lesssim \epsilon^2.$$

Thus integrating the estimate (4.2.6) in $r \in [2m_0(1 - \delta_{\mathcal{H}}), r_{\mathcal{T}}]$, where we recall that $r_{\mathcal{T}} \leq 2m_0(1 + 2\delta_{\mathcal{H}})$, we derive,

$$\sup_{(int)\mathcal{M}} |m - m_0| \lesssim \epsilon_0 m_0.$$

Since $\mathcal{M} = {}^{(ext)}\mathcal{M} \cup {}^{(int)}\mathcal{M}$ we infer that,

$$\sup_{\mathcal{M}} |m - m_0| \lesssim \epsilon_0 m_0.$$

This concludes the proof of Lemma 3.4.2.

4.3 Control of coordinates systems

The goal of this section is to prove Propositions 3.4.3 and 3.4.4. In both cases, the first two claims, on the form of the spacetime metric in the corresponding coordinates system as well as on the expression of the coordinates vectorfield with respect to the null frame (e_4, e_3, e_θ) , is already proved in Propositions 2.2.23 and 2.2.26. So we only focus on the third claim, i.e. on estimating $\check{\underline{\Omega}}, \check{\underline{\Omega}}, \varsigma, \underline{\varsigma}, \gamma, b, \underline{b}$ and e^Φ . The proof of Propositions 3.4.3 and 3.4.4 thus reduces to the proof of the following lemma.

Lemma 4.3.1. *Let $\theta \in [0, \pi]$ be the \mathbf{Z} -invariant scalar on \mathcal{M} defined by (2.2.52), i.e.*

$$\theta = \cot^{-1}(re_\theta(\Phi)). \quad (4.3.1)$$

Let

$$b = e_4(\theta), \quad \underline{b} = e_3(\theta), \quad \gamma = \frac{1}{(e_\theta(\theta))^2}. \quad (4.3.2)$$

Then, we have

$$\begin{aligned} \max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} \left(ru^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}} \right) \left(\left| \mathfrak{d}^k \left(\frac{\gamma}{r^2} - 1 \right) \right| + r \left| \mathfrak{d}^k b \right| \right) &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} u^{1 + \delta_{dec}} \left(\left| \mathfrak{d}^k \check{\underline{\Omega}} \right| + \left| \mathfrak{d}^k (\varsigma - 1) \right| + r \left| \mathfrak{d}^k \underline{b} \right| \right) &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{small}} \sup_{(int)\mathcal{M}} \underline{u}^{1 + \delta_{dec}} \left(\left| \mathfrak{d}^k \check{\underline{\Omega}} \right| + \left| \mathfrak{d}^k (\underline{\varsigma} - 1) \right| + \left| \mathfrak{d}^k \left(\frac{\gamma}{r^2} - 1 \right) \right| + \left| \mathfrak{d}^k b \right| + \left| \mathfrak{d}^k \underline{b} \right| \right) &\lesssim \epsilon. \end{aligned}$$

Also, e^Φ satisfies

$$\begin{aligned} \max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} \left(r u^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}} \right) \left| \mathfrak{d}^k \left(\frac{e^\Phi}{r \sin \theta} - 1 \right) \right| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{small}} \sup_{(int)\mathcal{M}} \frac{u^{1 + \delta_{dec}}}{r} \left| \mathfrak{d}^k \left(\frac{e^\Phi}{r \sin \theta} - 1 \right) \right| &\lesssim \epsilon. \end{aligned}$$

Proof. We prove the estimates in $(ext)\mathcal{M}$. The proof in $(int)\mathcal{M}$ is similar and left to the reader.

Step 1. We start with the estimate for $\underline{\check{\Omega}}$. Recall that

$$\mathfrak{d}_1^* \underline{\check{\Omega}} = \underline{\xi}$$

so that the bootstrap assumptions for $\underline{\xi}$ imply on any 2-sphere of the foliation of $(ext)\mathcal{M}$ and for any $k \leq k_{small}$

$$r^{\frac{1}{2}} \|\mathfrak{d}^k r \mathfrak{d}_1^* \underline{\check{\Omega}}\|_{L^4(S)} + \|\mathfrak{d}^k r \mathfrak{d}_1^* \underline{\check{\Omega}}\|_{L^2(S)} \lesssim r^2 \sup_S |\mathfrak{d}^k \underline{\xi}| \lesssim \epsilon r u^{-1 - \delta_{dec}}.$$

In view of the commutation formulas of Lemma 2.2.13 and of Proposition 2.1.25, together with the bootstrap assumptions, we infer any $k \leq k_{small}$, schematically,

$$[\mathfrak{d}^k, r \mathfrak{d}_1^*] = O(\epsilon) \mathfrak{d}^{\leq k} + O(1) \mathfrak{d}^{\leq k-1},$$

and hence,

$$\begin{aligned} r^{\frac{1}{2}} \|\mathfrak{d}_1^* \mathfrak{d}^k \underline{\check{\Omega}}\|_{L^4(S)} + \|\mathfrak{d}_1^* \mathfrak{d}^k \underline{\check{\Omega}}\|_{L^2(S)} &\lesssim \epsilon r u^{-1 - \delta_{dec}} + \epsilon \|\mathfrak{d}^{\leq k} \underline{\check{\Omega}}\|_{L^2(S)} + \epsilon r^{\frac{1}{2}} \|\mathfrak{d}^{\leq k} \underline{\check{\Omega}}\|_{L^4(S)} \\ &\quad + \|\mathfrak{d}^{\leq k-1} \underline{\check{\Omega}}\|_{L^2(S)} + r^{\frac{1}{2}} \|\mathfrak{d}^{\leq k-1} \underline{\check{\Omega}}\|_{L^4(S)} \\ &\lesssim \epsilon r u^{-1 - \delta_{dec}} + \epsilon \|r \mathfrak{d}_1^* \mathfrak{d}^{\leq k} \underline{\check{\Omega}}\|_{L^2(S)} + \epsilon \|\mathfrak{d}^{\leq k} \underline{\check{\Omega}}\|_{L^2(S)} \\ &\quad + \|\mathfrak{d}^{\leq k-1} \underline{\check{\Omega}}\|_{L^2(S)} + \|\mathfrak{d}^{\leq k} \underline{\check{\Omega}}\|_{L^2(S)}^{\frac{1}{2}} \|\mathfrak{d}^{\leq k-1} \underline{\check{\Omega}}\|_{L^2(S)}^{\frac{1}{2}}, \end{aligned}$$

where we used Gagliardo-Nirenberg on S . Together with the Poincaré inequality of Corollary 2.1.34 for \mathfrak{d}_1^* , we deduce

$$r^{\frac{1}{2}} \|\mathfrak{d}_1^* \mathfrak{d}^k \underline{\check{\Omega}}\|_{L^4(S)} + \|\mathfrak{d}_1^* \mathfrak{d}^k \underline{\check{\Omega}}\|_{L^2(S)} + \|\mathfrak{d}^k \underline{\check{\Omega}}\|_{L^2(S)} \lesssim \epsilon r u^{-1 - \delta_{dec}} + \|\mathfrak{d}^{\leq k-1} \underline{\check{\Omega}}\|_{L^2(S)}.$$

By iteration, and using again Gagliardo-Nirenberg on S , we infer on any 2-sphere of the foliation of $(ext)\mathcal{M}$ and for any $k \leq k_{small}$

$$\|\mathfrak{d}_1^* \mathfrak{d}^k \underline{\check{\Omega}}\|_{L^4(S)} + \|\mathfrak{d}^k \underline{\check{\Omega}}\|_{L^4(S)} \lesssim \epsilon r^{\frac{1}{2}} u^{-1 - \delta_{dec}},$$

and thus, by Sobolev embedding

$$\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} |\mathfrak{d}^k \check{\underline{\Omega}}| \lesssim \epsilon$$

which is the desired estimate for $\check{\underline{\Omega}}$.

Step 2. Next, we estimate ς . First, recall that we have

$$e_\theta(\log \varsigma) = \eta - \zeta.$$

Since the bootstrap assumptions for $\eta - \zeta$ are at least as good as for ξ , we obtain, arguing as in Step 1 the following analog of the above estimate for $\check{\underline{\Omega}}$

$$\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} |\mathfrak{d}^k \check{\zeta}| \lesssim \epsilon.$$

Now that we control $\check{\zeta}$, we turn to the estimate for ς . First, recall from the GCM on Σ_* that we have

$$u + r = c_{\Sigma_*} \text{ and } a|_{SP} = -1 - \frac{2m}{r}, \text{ where } \nu = e_3 + ae_4 \text{ and } \nu \text{ is tangent to } \Sigma_*,$$

with c_{Σ_*} a constant, and SP denoting the south pole of the spheres of Σ_* . We deduce on the south poles of Σ_*

$$0 = \nu(u + r) = e_3(u) + e_3(r) + ae_4(r) = \frac{2}{\varsigma} + e_3(r) - \left(1 + \frac{2m}{r}\right) e_4(r)$$

and hence

$$\frac{2}{\varsigma} - 2 = -\frac{r}{2} \left(\left(\bar{\kappa} + \frac{2\Upsilon}{r} \right) + \underline{A} - \left(1 + \frac{2m}{r} \right) \left(\bar{\kappa} - \frac{2}{r} \right) \right) \text{ on } SP \cap \Sigma_*.$$

Together with the fact that $\bar{\varsigma} = \varsigma - \zeta$, the above control of $\check{\zeta}$, the control of $\bar{\kappa}$ and $\underline{\bar{\kappa}}$ provided by Lemma 3.4.1, the formula for \underline{A} , the control for $\check{\underline{\Omega}}$ in Step 1, the bootstrap assumptions on decay, and the fact that $\bar{\varsigma}$ is constant on the sphere, we infer

$$\max_{0 \leq k \leq k_{small}} \sup_{\Sigma_*} u^{1+\delta_{dec}} |\mathfrak{d}^k (\bar{\varsigma} - 1)| \lesssim \epsilon.$$

Using $\varsigma = \bar{\varsigma} + \zeta$ and the above estimates for $\bar{\varsigma}$ and ζ , we obtain

$$\max_{0 \leq k \leq k_{small}} \sup_{\Sigma_*} u^{1+\delta_{dec}} |\mathfrak{d}^k (\varsigma - 1)| \lesssim \epsilon.$$

Finally, recall

$$e_4(\varsigma) = 0.$$

Commuting with \mathfrak{d} , using the bootstrap assumptions on decay and the above control for $\varsigma - 1$ on Σ_* , we infer

$$\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} u^{1+\delta_{dec}} |\mathfrak{d}^k(\varsigma - 1)| \lesssim \epsilon.$$

Remark 4.3.2. In $^{(int)}\mathcal{M}$, we analogously transport $\underline{\varsigma}$ from the timelike hyper surface \mathcal{T} . To estimate $\underline{\varsigma}$ on \mathcal{T} , one uses the following identity (in the frame of $^{(int)}\mathcal{M}$)

$$\frac{2}{\underline{\varsigma}} - 1 = -\frac{\bar{\kappa} + A}{\Upsilon \bar{\kappa}} \left(\frac{2}{\varsigma} - 1 \right) - \frac{A}{\Upsilon \bar{\kappa}} - \frac{(\bar{\kappa} - \frac{2\Upsilon}{r}) + \Upsilon \left(\bar{\kappa} + \frac{2}{r} \right)}{\Upsilon \bar{\kappa}} \text{ on } \mathcal{T}.$$

This identity follows from the definition of ς and $\underline{\varsigma}$, the identity for $e_3(r)$ and $e_4(r)$ in $^{(int)}\mathcal{M}$, the fact that $\underline{u} = u$ on \mathcal{T} , and that $\mathcal{T} = \{r = r_{\mathcal{T}}\}$ so that the vectorfield

$$T_{\mathcal{T}} = e_4 - \frac{e_4(r)}{e_3(r)} e_3 = e_4 - \frac{\bar{\kappa} + A}{\bar{\kappa}} e_3$$

is tangent to \mathcal{T} .

Step 3. We make the auxiliary bootstrap assumption which will be recovered at the end of Step 5

$$|e^{\Phi}| \leq 2r, \quad |e_{\theta}(e^{\Phi})| \leq 2. \quad (4.3.3)$$

We start with the estimate for e^{Φ} . Recall from (2.2.53) that the following identity holds

$$\frac{e^{\Phi}}{r \sin \theta} = \sqrt{1 + \mathbf{a}}. \quad (4.3.4)$$

where \mathbf{a} has been introduced in (2.2.54) by

$$\mathbf{a} = \frac{e^{2\Phi}}{r^2} + (e_{\theta}(e^{\Phi}))^2 - 1.$$

In order to estimate e^{Φ} , it thus suffices to estimate \mathbf{a} .

Step 4. Now, recall from Lemma 2.2.30 that \mathbf{a} verifies the following identities on $^{(ext)}\mathcal{M}$,

$$\begin{aligned} e_4(\mathbf{a}) &= \frac{(\check{\kappa} - \vartheta)e^{2\Phi}}{r^2} + 2e_{\theta}(e^{\Phi}) \left(\beta - e_4(\Phi)\zeta \right) e^{\Phi}, \\ e_{\theta}(\mathbf{a}) &= 2e_{\theta}(\Phi)e^{2\Phi} \left(\left(\rho + \frac{2m}{r^3} \right) + \frac{1}{4} \left(\kappa \bar{\kappa} + \frac{4\Upsilon}{r^2} \right) - \frac{1}{4} \vartheta \underline{\vartheta} \right), \\ e_3(\mathbf{a}) &= \frac{(\check{\kappa} - \underline{A} - \underline{\vartheta})e^{2\Phi}}{r^2} + 2e_{\theta}(e^{\Phi}) \left(\underline{\beta} + e_3(\Phi)\zeta + \underline{\xi} e_4(\Phi) \right) e^{\Phi}. \end{aligned}$$

Together with our bootstrap assumptions on decay for in ${}^{(ext)}\mathcal{M}$ for $\check{\kappa}, \vartheta, \check{\kappa}, \underline{\vartheta}, \beta, \underline{\beta}, \rho, \zeta, \xi$ and $\check{\Omega}$ and the bootstrap assumption (4.3.3), we infer

$$\max_{1 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} \left(ru^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}} \right) |\mathfrak{d}^k \mathbf{a}| \lesssim \epsilon.$$

In particular, we deduce,

$$\sup_{(ext)\mathcal{M}} \left(ru^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}} \right) |\check{\mathfrak{a}}| \lesssim \epsilon.$$

Step 5. To estimate $\bar{\mathfrak{a}}$ we make use of equation (2.1.13) according to which

$$\left(e_\theta(e^\Phi) \right)^2 = 1 \quad \text{on the axis of symmetry.}$$

Since $e^{2\Phi}$ also vanishes there we infer that $\mathbf{a} = 0$ on the axis. Therefore, on the axis, $\check{\mathfrak{a}} = -\bar{\mathfrak{a}}$, i.e.,

$$\bar{\mathfrak{a}} = -\check{\mathfrak{a}}|_{\text{axis}}$$

and therefore,

$$|\bar{\mathfrak{a}}| \lesssim |\check{\mathfrak{a}}| \lesssim \frac{\epsilon}{ru^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}}}.$$

We conclude that,

$$\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} \left(ru^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}} \right) |\mathfrak{d}^k \mathbf{a}| \lesssim \epsilon. \quad (4.3.5)$$

In view of (4.3.4) and (4.3.5), we immediately infer

$$\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} \left(ru^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}} \right) \left| \mathfrak{d}^k \left(\frac{e^\Phi}{r \sin \theta} - 1 \right) \right| \lesssim \epsilon.$$

Together with (4.3.5) and the definition of \mathbf{a} , this implies

$$|e^\Phi| = (1 + O(\epsilon))r \sin \theta \leq \frac{3r}{2}, \quad |e_\theta(e^\Phi)| = \sqrt{1 - \frac{e^{2\Phi}}{r^2} + \mathbf{a}} \leq |\cos \theta| + O(\epsilon) \leq \frac{3}{2}, \quad (4.3.6)$$

which is an improvement of the bootstrap assumption (4.3.3) which hence holds everywhere on ${}^{(ext)}\mathcal{M}$.

Step 6. We now prove the estimates for b , \underline{b} and γ . Recall from Lemma 2.2.29 that θ defined by (4.3.1) satisfies

$$\begin{aligned} re_\theta(\theta) &= 1 + \frac{r^2(K - \frac{1}{r^2})}{1 + (re_\theta(\Phi))^2}, \\ e_3(\theta) &= -\frac{r\underline{\beta} + \frac{r}{2}(-\check{\kappa} + \underline{A} + \vartheta)e_\theta(\Phi) + r\underline{\xi}e_4(\Phi) + r\underline{\zeta}e_3(\Phi)}{1 + (re_\theta(\Phi))^2}, \\ e_4(\theta) &= -\frac{r\beta + \frac{r}{2}(-\check{\kappa} + \vartheta)e_\theta(\Phi) - r\underline{\zeta}e_3(\Phi)}{1 + (re_\theta(\Phi))^2}. \end{aligned}$$

In view of the definition of b , \underline{b} and γ , we infer

$$\begin{aligned} \frac{r}{\sqrt{\gamma}} &= 1 + \frac{r^2(K - \frac{1}{r^2})}{1 + (re_\theta(\Phi))^2}, \\ \underline{b} &= -\frac{r\underline{\beta} + \frac{r}{2}(-\check{\kappa} + \underline{A} + \vartheta)e_\theta(\Phi) + r\underline{\xi}e_4(\Phi) + r\underline{\zeta}e_3(\Phi)}{1 + (re_\theta(\Phi))^2}, \\ b &= -\frac{r\beta + \frac{r}{2}(-\check{\kappa} + \vartheta)e_\theta(\Phi) - r\underline{\zeta}e_3(\Phi)}{1 + (re_\theta(\Phi))^2}. \end{aligned}$$

Also, we have in view of the definition of \mathbf{a}

$$1 + (re_\theta(\Phi))^2 = 1 + \frac{(e_\theta(e^\Phi))^2}{\frac{e^{2\Phi}}{r^2}} = \frac{r^2}{e^{2\Phi}}(1 + \mathbf{a})$$

and hence

$$\begin{aligned} \frac{r}{\sqrt{\gamma}} &= 1 + \frac{e^{2\Phi}}{r^2} \left(\frac{r^2(K - \frac{1}{r^2})}{1 + \mathbf{a}} \right), \\ \underline{b} &= -\frac{e^{2\Phi}}{r^2} \left(\frac{r\underline{\beta} + \frac{r}{2}(-\check{\kappa} + \underline{A} + \vartheta)e_\theta(\Phi) + r\underline{\xi}e_4(\Phi) + r\underline{\zeta}e_3(\Phi)}{1 + \mathbf{a}} \right), \\ b &= -\frac{e^{2\Phi}}{r^2} \left(\frac{r\beta + \frac{r}{2}(-\check{\kappa} + \vartheta)e_\theta(\Phi) - r\underline{\zeta}e_3(\Phi)}{1 + \mathbf{a}} \right). \end{aligned}$$

The bootstrap assumptions on decay in $^{(ext)}\mathcal{M}$ for $\check{\kappa}$, ϑ , $\check{\kappa}$, $\underline{\vartheta}$, β , $\underline{\beta}$, ζ , $\underline{\xi}$ and $\underline{\Omega}$, the estimate (4.3.5) for \mathbf{a} , the estimate (4.3.6), and the identity

$$\begin{aligned} K - \frac{1}{r^2} &= -\frac{1}{4}\kappa\underline{\kappa} + \frac{1}{4}\vartheta\underline{\vartheta} - \rho - \frac{1}{r^2} \\ &= -\frac{1}{4} \left(\kappa\underline{\kappa} + \frac{4\Upsilon}{r^2} \right) - \left(\rho + \frac{2m}{r^3} \right) + \frac{1}{4}\vartheta\underline{\vartheta} \end{aligned}$$

imply

$$\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} \left(ru^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}} \right) \left(\left| \mathfrak{d}^k \left(\frac{r}{\sqrt{\gamma}} - 1 \right) \right| + r |\mathfrak{d}^k b| \right) \lesssim \epsilon,$$

and

$$\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} ru^{1 + \delta_{dec}} |\mathfrak{d}^k \underline{b}| \lesssim \epsilon.$$

In particular, we also have

$$\max_{0 \leq k \leq k_{small}} \sup_{(ext)\mathcal{M}} \left(ru^{\frac{1}{2} + \delta_{dec}} + u^{1 + \delta_{dec}} \right) \left| \mathfrak{d}^k \left(\frac{\gamma}{r^2} - 1 \right) \right| \lesssim \epsilon.$$

These are the desired estimate for b , \underline{b} and γ in $(ext)\mathcal{M}$. This concludes the proof of the lemma. \square

In this section, we also prove two useful lemmas concerning estimates on 2-spheres of $(ext)\mathcal{M}$ and $(int)\mathcal{M}$.

Lemma 4.3.3. *Let $\theta \in [0, \pi]$ be the \mathbf{Z} -invariant scalar on \mathcal{M} defined by (2.2.52). Then, we have on \mathcal{M}*

$$re_\theta(\Phi) = \frac{\varpi}{\sin \theta}$$

where ϖ is a reduced 1-scalar satisfying

$$\sup_{\mathcal{M}} |\varpi| \leq 2.$$

Also, we have

$$\frac{1}{\sin \theta} \leq 2|re_\theta(\Phi)| + 2 \text{ on } \mathcal{M}.$$

Proof. The proof is similar on $(ext)\mathcal{M}$ and $(int)\mathcal{M}$ so we focus on $(ext)\mathcal{M}$. Recall from (4.3.6) that

$$|e_\theta(e^\Phi)| \leq \frac{3}{2}.$$

Furthermore, in view of Proposition 3.4.3, we have in particular

$$\sup_{(ext)\mathcal{M}} \left| \frac{e^\Phi}{r \sin \theta} - 1 \right| \lesssim \epsilon.$$

Since we have

$$\varpi = r \sin \theta e_\theta(\Phi),$$

we deduce

$$|\varpi| = \frac{r \sin \theta}{e^\Phi} |e_\theta(e^\Phi)| \leq \frac{3}{2}(1 + O(\epsilon)) \leq 2,$$

which is the desired estimate for ϖ .

We now consider the upper bound for $(\sin \theta)^{-1}$. Recall the definition (2.2.54) of \mathbf{a}

$$\mathbf{a} = \frac{e^{2\Phi}}{r^2} + (e_\theta(e^\Phi))^2 - 1.$$

We infer

$$\begin{aligned} r^2 e_\theta(\Phi)^2 &= \frac{r^2 (e_\theta(e^\Phi))^2}{e^{2\Phi}} \\ &= \frac{1 + \mathbf{a}}{\frac{e^{2\Phi}}{r^2}} - 1 \\ &= \frac{1 + \mathbf{a} - (\sin \theta)^2 \left(1 + \left(\frac{e^{2\Phi}}{r^2 (\sin \theta)^2} - 1\right)\right)}{(\sin \theta)^2 \left(1 + \left(\frac{e^{2\Phi}}{r^2 (\sin \theta)^2} - 1\right)\right)} \\ &= \frac{(\cos \theta)^2 + \mathbf{a} - (\sin \theta)^2 \left(\frac{e^{2\Phi}}{r^2 (\sin \theta)^2} - 1\right)}{(\sin \theta)^2 \left(1 + \left(\frac{e^{2\Phi}}{r^2 (\sin \theta)^2} - 1\right)\right)} \end{aligned}$$

and hence

$$\sin \theta |r e_\theta(\Phi)| = \frac{\sqrt{(\cos \theta)^2 + \mathbf{a} - (\sin \theta)^2 \left(\frac{e^{2\Phi}}{r^2 (\sin \theta)^2} - 1\right)}}{\sqrt{1 + \left(\frac{e^{2\Phi}}{r^2 (\sin \theta)^2} - 1\right)}}.$$

Now, in view of (4.3.5), \mathbf{a} satisfies in particular

$$\sup_{(ext)\mathcal{M}} |\mathbf{a}| \lesssim \epsilon.$$

Together with

$$\sup_{(ext)\mathcal{M}} \left| \frac{e^\Phi}{r \sin \theta} - 1 \right| \lesssim \epsilon,$$

we infer

$$\sin \theta |re_\theta(\Phi)| = \frac{\sqrt{(\cos \theta)^2 + O(\epsilon)}}{\sqrt{1 + O(\epsilon)}}.$$

Thus, we deduce

$$\sin \theta |re_\theta(\Phi)| \geq \frac{\sqrt{2}}{2}(1 + O(\epsilon)) \geq \frac{1}{2} \text{ for } 0 \leq \theta \leq \frac{\pi}{4} \text{ and } \frac{3\pi}{4} \leq \theta \leq \pi.$$

On the other hand, we have

$$\sin \theta \geq \frac{\sqrt{2}}{2} \text{ on } \frac{\pi}{4} \leq \theta \leq \frac{3\pi}{4}$$

and hence

$$\frac{1}{\sin \theta} \leq 2|re_\theta(\Phi)| + 2 \text{ on } 0 \leq \theta \leq \pi$$

which is the desired estimate. This concludes the proof of the lemma. \square

Lemma 4.3.4. *Let $\theta \in [0, \pi]$ be the \mathbf{Z} -invariant scalar on \mathcal{M} defined by (2.2.52). Then, for any reduced 1-scalar h , we have on any 2-sphere S on ${}^{(ext)}\mathcal{M}$ and of ${}^{(int)}\mathcal{M}$*

$$\sup_S \frac{|h|}{e^\Phi} \lesssim r^{-1} \sup_S (|h| + |\not\partial h|) \quad \text{and} \quad \left\| \frac{h}{e^\Phi} \right\|_{L^2(S)} \lesssim r^{-1} \|h\|_{b_1(S)}.$$

Proof. The proof is similar on ${}^{(ext)}\mathcal{M}$ and ${}^{(int)}\mathcal{M}$ so we focus on ${}^{(ext)}\mathcal{M}$. Recall that the 2-surface S is parametrized by the coordinate $\theta \in [0, \pi]$, and that the axis corresponds to the 2 poles $\theta = 0$ and $\theta = \pi$. In view of

$$\sup_{{}^{(ext)}\mathcal{M}} \left| \frac{e^\Phi}{r \sin \theta} - 1 \right| \lesssim \epsilon,$$

we have

$$\sup_{S \cap \{\frac{\pi}{4} \leq \theta \leq \frac{3\pi}{4}\}} \frac{|h|}{e^\Phi} \lesssim r^{-1} \sup_S |h| \quad \text{and} \quad \left\| \frac{h}{e^\Phi} \right\|_{L^2(S \cap \{\frac{\pi}{4} \leq \theta \leq \frac{3\pi}{4}\})} \lesssim r^{-1} \|h\|_{L^2(S)}$$

which is the desired estimate for $\pi/4 \leq \theta \leq 3\pi/4$.

It remains to consider the portions $0 \leq \theta \leq \pi/4$ and $3\pi/4 \leq \theta \leq \pi$ of S . These regions can be treated analogously, so we focus on $0 \leq \theta \leq \pi/4$. Recall from Remark 2.1.21 that

any reduced scalar in \mathfrak{s}_k , for $k \geq 1$, must vanish on the axis of symmetry of \mathbf{Z} , i.e. at the two poles. In particular, h must vanish at $\theta = 0$. We deduce

$$\frac{h}{e^\Phi} = \frac{he^\Phi}{e^{2\Phi}} = \frac{\int_0^\theta \partial_\theta(e^\Phi h)}{e^{2\Phi}} = \frac{\int_0^\theta \sqrt{\gamma} \mathfrak{s}_\theta e^\Phi(e^\Phi h)}{e^{2\Phi}} = \frac{\int_0^\theta \sqrt{\gamma} e^\Phi \mathfrak{d}_1 h}{e^{2\Phi}}.$$

Since we have $|\gamma| \lesssim r$, we infer

$$\frac{|h|}{e^\Phi} \lesssim \frac{\int_0^\theta e^\Phi |\mathfrak{d}h|}{e^{2\Phi}}$$

and since

$$\sup_{(ext)\mathcal{M}} \left| \frac{e^\Phi}{r \sin \theta} - 1 \right| \lesssim \epsilon,$$

we deduce

$$\frac{|h|}{e^\Phi} \lesssim r^{-1} \frac{\int_0^\theta \sin(\theta') |\mathfrak{d}h| d\theta'}{(\sin \theta)^2}.$$

This yields

$$\sup_{S \cap \{0 \leq \theta \leq \frac{\pi}{4}\}} \frac{|h|}{e^\Phi} \lesssim r^{-1} \sup_S |\mathfrak{d}h|$$

which is the desired sup norm estimate for $0 \leq \theta \leq \pi/4$.

It remains to control the L^2 -norm on $0 \leq \theta \leq \pi/4$. We have in view of the above

$$\begin{aligned} \left\| \frac{h}{e^\Phi} \right\|_{L^2(S \cap \{0 \leq \theta \leq \frac{\pi}{4}\})}^2 &\lesssim r^{-2} \int_0^{\frac{\pi}{4}} \frac{\left(\int_0^\theta \sin(\theta') |\mathfrak{d}h| d\theta' \right)^2}{(\sin \theta)^4} e^\Phi d\theta \\ &\lesssim r^{-1} \int_0^{\frac{\pi}{4}} \left(\int_0^\theta (\sin(\theta'))^2 |\mathfrak{d}h|^2 d\theta' \right) \frac{d\theta}{(\sin \theta)^2} \\ &\lesssim r^{-1} \int_0^{\frac{\pi}{4}} (\sin \theta)^2 |\mathfrak{d}h|^2 \left(\int_\theta^{\frac{\pi}{4}} \frac{d\theta'}{(\sin(\theta'))^2} \right) d\theta \\ &\lesssim r^{-1} \int_0^{\frac{\pi}{4}} |\mathfrak{d}h|^2 \sin \theta d\theta \\ &\lesssim r^{-2} \int_0^{\frac{\pi}{4}} |\mathfrak{d}h|^2 e^\Phi d\theta \\ &\lesssim r^{-2} \|\mathfrak{d}h\|_{L^2(S)}^2 \end{aligned}$$

and hence

$$\left\| \frac{h}{e^\Phi} \right\|_{L^2(S \cap \{0 \leq \theta \leq \frac{\pi}{4}\})} \lesssim r^{-1} \|\not\partial h\|_{L^2(S)}$$

which is the desired $L^2(S)$ estimate for $0 \leq \theta \leq \pi/4$. This concludes the proof of the lemma. \square

4.4 Pointwise bounds for high order derivatives

The goal of this section is to prove Proposition 3.4.5. We deal first with the region $r \leq 4m_0$ as follows

1. The curvature components and Ricci coefficients satisfy in view of the bootstrap assumptions on energy

$$\max_{k \leq k_{large}} \int_{(int)\mathcal{M}} \left(|\check{R}|^2 + |\check{\Gamma}|^2 \right) + \max_{k \leq k_{large}-1} \int_{(ext)\mathcal{M}(r \leq 4m_0)} \left(|\check{R}|^2 + |\check{\Gamma}|^2 \right) \leq \epsilon^2.$$

2. We first take the trace on the ingoing null cones foliating $(int)\mathcal{M}$ and the outgoing null cones foliating $(ext)\mathcal{M}(r \leq 4m_0)$ which loses one derivative. We thus obtain

$$\max_{k \leq k_{large}-1} \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u} \left(|\check{R}|^2 + |\check{\Gamma}|^2 \right) + \max_{k \leq k_{large}-2} \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \leq 4m_0)} \left(|\check{R}|^2 + |\check{\Gamma}|^2 \right) \lesssim \epsilon^2.$$

3. We then take the trace on the 2-spheres S foliation the null cones in $(int)\mathcal{M}$ and $(ext)\mathcal{M}(r \leq 4m_0)$ to infer

$$\max_{k \leq k_{large}-2} \sup_{(int)\mathcal{M}} \left(\|\check{R}\|_{L^2(S)} + \|\check{\Gamma}\|_{L^2(S)} \right) + \max_{k \leq k_{large}-3} \sup_{(ext)\mathcal{M}(r \leq 4m_0)} \left(|\check{R}|^2 + |\check{\Gamma}|^2 \right) \lesssim \epsilon.$$

4. Finally, using the Sobolev embedding on the 2-sphere S , which loses 2 derivatives, we deduce

$$\max_{k \leq k_{large}-4} \sup_{(int)\mathcal{M}} \left(|\check{R}| + |\check{\Gamma}| \right) + \max_{k \leq k_{large}-5} \sup_{(ext)\mathcal{M}(r \leq 4m_0)} \left(|\check{R}| + |\check{\Gamma}| \right) \lesssim \epsilon,$$

which is the desired estimate in the region $(int)\mathcal{M} \cup (ext)\mathcal{M}(r \leq 4m_0)$.

It remains to consider the region $^{(ext)}\mathcal{M}(r \geq 4m_0)$. We proceed as follows

Step 1. The Ricci coefficients satisfy in view of the bootstrap assumptions on energy

$$\begin{aligned} & \max_{k \leq k_{large}} \int_{\Sigma_*} \left[r^2 \left((\mathfrak{d}^{\leq k} \vartheta)^2 + (\mathfrak{d}^{\leq k} \check{\kappa})^2 + (\mathfrak{d}^{\leq k} \zeta)^2 + (\mathfrak{d}^{\leq k} \check{\underline{\kappa}})^2 \right) + (\mathfrak{d}^{\leq k} \underline{\vartheta})^2 \right. \\ & + \left. (\mathfrak{d}^{\leq k} \eta)^2 + (\mathfrak{d}^{\leq k} \check{\omega})^2 + (\mathfrak{d}^{\leq k} \underline{\xi})^2 \right] \\ & + \sup_{\lambda \geq 4m_0} \left(\int_{\{r=\lambda\}} \left[\lambda^2 \left((\mathfrak{d}^{\leq k} \vartheta)^2 + (\mathfrak{d}^{\leq k} \check{\kappa})^2 + (\mathfrak{d}^{\leq k} \zeta)^2 \right) \right. \right. \\ & \left. \left. + \lambda^{2-\delta_B} (\mathfrak{d}^{\leq k} \check{\underline{\kappa}})^2 + (\mathfrak{d}^{\leq k} \underline{\vartheta})^2 + (\mathfrak{d}^{\leq k} \eta)^2 + (\mathfrak{d}^{\leq k} \check{\omega})^2 + \lambda^{-\delta_B} (\mathfrak{d}^{\leq k} \underline{\xi})^2 \right] \right) \leq \epsilon^2. \end{aligned}$$

We take the trace on the 2-spheres S foliating the timelike cylinders $\{r = r_0\}$, for $r_0 \geq 4m_0$, which loses a derivative, and infer in particular

$$\begin{aligned} & \max_{k \leq k_{large}-1} \sup_{^{(ext)}\mathcal{M}(r \geq 4m_0)} \left\{ r \left(\|\mathfrak{d}^k \check{\kappa}\|_{L^2(S)} + \|\mathfrak{d}^k \zeta\|_{L^2(S)} + \|\mathfrak{d}^k \vartheta\|_{L^2(S)} \right) + r^{1-\frac{\delta_B}{2}} \|\mathfrak{d}^k \check{\underline{\kappa}}\|_{L^2(S)} \right. \\ & \left. + \|\mathfrak{d}^k \eta\|_{L^2(S)} + \|\mathfrak{d}^k \underline{\vartheta}\|_{L^2(S)} + \|\mathfrak{d}^k \check{\omega}\|_{L^2(S)} + r^{-\frac{\delta_B}{2}} \|\mathfrak{d}^k \underline{\xi}\|_{L^2(S)} \right\} \lesssim \epsilon. \end{aligned}$$

Also, we take the trace on the 2-spheres S foliating the spacelike hyper surface Σ_* , which loses a derivative, and infer in particular

$$\max_{k \leq k_{large}-1} \sup_{\Sigma_*} r \|\mathfrak{d}^k \check{\underline{\kappa}}\|_{L^2(S)} \lesssim \epsilon.$$

Step 2. On can easily prove the following trace theorem

$$\max_{k \leq k_{large}-1} \left(\sup_{r \geq 4m_0} r^{5+\delta_B} \int_S (\mathfrak{d}^k \alpha)^2 \right) \lesssim \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u} r^{4+\delta_B} (\mathfrak{d}^{\leq k_{large}} \alpha)^2,$$

which together with the bootstrap assumptions on energy for α in $^{(ext)}\mathcal{M}(r \geq 4m_0)$ implies

$$\max_{k \leq k_{large}-1} \left(\sup_{r \geq 4m_0} r^{5+\delta_B} \int_S (\mathfrak{d}^k \alpha)^2 \right) \lesssim \epsilon^2.$$

Step 3. Using the trace theorem

$$\max_{k \leq k_{large}-1} \left(\sup_{r \geq 4m_0} r^5 \int_S (\mathfrak{d}^k \beta)^2 \right) \lesssim \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u} r^4 (\mathfrak{d}^{\leq k_{large}} \beta)^2,$$

we infer, together with the bootstrap assumptions on energy for β in $^{(ext)}\mathcal{M}(r \geq 4m_0)$,

$$\max_{k \leq k_{iarge} - 1} \left(\sup_{r \geq 4m_0} r^5 \int_S (\mathfrak{d}^k \beta)^2 \right) \lesssim \epsilon^2. \quad (4.4.1)$$

The power of r of the above estimate is not strong enough. To upgrade the estimate, recall that we have the Bianchi identity

$$e_4(\beta) + 2\kappa\beta = \not\partial_2 \alpha + \zeta \alpha.$$

This yields

$$\begin{aligned} e_4 \left(r^{5+\delta_B} \int_S \beta^2 \right) &= \int_S r^{5+\delta_B} \left(2\beta e_4(\beta) + \kappa\beta^2 + b \frac{e_4(r)}{r} \beta^2 \right) \\ &= \int_S r^{5+\delta_B} \left(-\frac{1-\delta_B}{2} \kappa\beta^2 + 2\beta(r^{-1} \not\partial \alpha + \zeta \alpha) - \frac{5+\delta_B}{2} \check{\kappa}\beta^2 \right) \end{aligned}$$

and hence

$$\begin{aligned} e_4 \left(r^{5+\delta_B} \int_S \beta^2 \right) + \frac{1-\delta_B}{2} \int_S r^{5+\delta_B} \kappa\beta^2 &= \int_S r^{4+\delta_B} \left(2\beta(\not\partial \alpha + r\zeta \alpha) - \frac{5+\delta_B}{2} \check{\kappa}\beta^2 \right) \\ &\lesssim \left(\int_S r^{4+\delta_B} (\mathfrak{d}^{\leq 1} \alpha)^2 \right)^{\frac{1}{2}} \left(\int_S r^{4+\delta_B} \beta^2 \right)^{\frac{1}{2}} + \epsilon \int_S r^{4+\delta_B} \beta^2 \end{aligned}$$

where we used the pointwise estimates of Step 1 for $\check{\kappa}$ and ζ . We infer

$$e_4 \left(r^{5+\delta_B} \int_S \beta^2 \right) + \int_S r^{4+\delta_B} \beta^2 \lesssim \int_S r^{4+\delta_B} (\mathfrak{d}^{\leq 1} \alpha)^2.$$

Integrating, from $r \geq 6m_0$, we deduce

$$\begin{aligned} \sup_{r \geq 6m_0} r^{5+\delta_B} \int_S \beta^2 + \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq 6m_0)} r^{4+\delta_B} \beta^2 &\lesssim \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u} r^{4+\delta_B} (\mathfrak{d}^{\leq 1} \alpha)^2 + \int_{S_{r=6m_0}} \beta^2 \\ &\lesssim \epsilon^2, \end{aligned}$$

where we used the bootstrap assumptions on energy for α in $^{(ext)}\mathcal{M}(r \geq 4m_0)$ and the non sharp estimate (4.4.1) for β . Using again (4.4.1), we obtain

$$\sup_{r \geq 4m_0} r^{5+\delta_B} \int_S \beta^2 + \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq 4m_0)} r^{4+\delta_B} \beta^2 \lesssim \epsilon^2.$$

To discuss higher order derivatives, recall from Lemma 2.2.13 the following commutator, written in schematic form,

$$[\not\partial, e_4] = (\check{\kappa}, \vartheta) \not\partial + (\zeta, r\beta).$$

Also, recall from Lemma 2.2.14 the following commutator,

$$[\mathbf{T}, e_4] = \left((\underline{\omega} - \frac{m}{r^2}) - \frac{m}{2r} (\bar{\kappa} - \frac{2}{r}) + \frac{e_4(m)}{r} \right) e_4 + (\eta + \zeta) e_\theta.$$

In view of the estimates of Step 1 for $k_{large} - 1$ derivatives of $\check{\kappa}$, ϑ , ζ , η , $\underline{\omega}$, the pointwise estimates for β in (4.4.1), the control of $\bar{\kappa}$ in Lemma 3.4.1, and the control of $e_4(m)$ in Lemma 3.4.2, we infer, schematically,

$$\left\| \mathfrak{d}^k \left([\mathfrak{d}, e_4]\beta, [\mathbf{T}, e_4]\beta \right) \right\|_{L^2(S)} \lesssim O(\epsilon r^{-2}) \|\mathfrak{d}^{\leq k+1}\beta\|_{L^2(S)} \text{ for } k \leq k_{large} - 2.$$

Thus, commuting the Bianchi identity for $e_4(\beta)$ with \mathbf{T} and \mathfrak{d} together with the above commutator estimate, using the Bianchi identity to recover the e_4 derivatives, we obtain for higher order derivatives

$$\begin{aligned} & \max_{k \leq k_{large} - 1} \left(\sup_{r \geq 4m_0} r^{5+\delta_B} \int_S (\mathfrak{d}^k \beta)^2 + \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq 4m_0)} r^{4+\delta_B} (\mathfrak{d}^k \beta)^2 \right) \\ & \lesssim \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq 4m_0)} r^{4+\delta_B} (\mathfrak{d}^{\leq k_{large}} \alpha)^2 \\ & \lesssim \epsilon^2. \end{aligned}$$

Step 4. Recall from Proposition 2.2.18 that we have

$$\begin{aligned} e_4 \check{\rho} + \frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} &= \mathfrak{d}_1 \beta + \text{Err}[e_4 \check{\rho}], \\ \text{Err}[e_4 \check{\rho}] &= -\frac{3}{2} \check{\kappa} \check{\rho} + \frac{1}{2} \bar{\kappa} \bar{\rho} - \left(\frac{1}{2} \underline{\vartheta} \alpha + \zeta \beta \right) + \overline{\left(\frac{1}{2} \underline{\vartheta} \alpha + \zeta \beta \right)}. \end{aligned}$$

This yields

$$\begin{aligned} e_4 \left(r^4 \int_S (\check{\rho})^2 \right) &= \int_S r^4 \left(2\check{\rho} e_4(\check{\rho}) + \kappa \check{\rho}^2 + 4 \frac{e_4(r)}{r} \check{\rho}^2 \right) \\ &= \int_S r^4 \left(-3\bar{\rho} \check{\kappa} \check{\rho} + 2\check{\rho} (r^{-1} \mathfrak{d} \beta + \text{Err}[e_4 \check{\rho}]) + \check{\kappa}^2 \check{\rho}^2 \right) \end{aligned}$$

and hence

$$e_4 \left[\left(r^4 \int_S (\check{\rho})^2 \right)^{\frac{1}{2}} \right] \lesssim \left[\int_S r^4 \left((\bar{\rho} \check{\kappa})^2 + (r^{-1} \mathfrak{d} \beta)^2 + (\text{Err}[e_4 \check{\rho}])^2 + \check{\kappa}^2 \check{\rho}^2 \right) \right]^{\frac{1}{2}}$$

Using the estimates of Step 1, 2 and 3 for $\check{\kappa}$, ζ , $\underline{\vartheta}$, α and β , and the control of $\bar{\rho}$ in Lemma 3.4.1, we infer

$$e_4 \left[\left(r^4 \int_S (\check{\rho})^2 \right)^{\frac{1}{2}} \right] \lesssim \frac{\epsilon}{r^{\frac{3}{2} + \frac{\delta_B}{2}}} + \frac{\epsilon}{r^2} \left(r^4 \int_S (\check{\rho})^2 \right)^{\frac{1}{2}}.$$

Integrating from $r = 4m_0$, we control $\|\check{\rho}\|_{L^2(S)}$ from the control in $r \leq 4m_0$, we infer

$$\sup_{r \geq 4m_0} r^4 \int_S \check{\rho}^2 \lesssim \epsilon^2.$$

Next, commuting the equation for $e_4(\check{\rho})$ with \mathbf{T} and $\check{\vartheta}$ together with the commutator estimate of Step 3, using the equation for $e_4(\check{\rho})$ to recover the e_4 derivatives, we obtain similarly for higher order derivatives

$$\max_{k \leq k_{large} - 2} \sup_{r \geq 4m_0} r^4 \int_S (\check{\vartheta}^k \check{\rho})^2 \lesssim \epsilon^2.$$

Step 5. Recall from Proposition 2.2.18 that we have the following transport equations in the e_4 direction,

$$\begin{aligned} e_4 \check{\kappa} + \frac{1}{2} \overline{\kappa} \check{\kappa} + \frac{1}{2} \check{\kappa} \overline{\kappa} &= -2 \check{\vartheta}_1 \zeta + 2\check{\rho} + \text{Err}[e_4 \check{\kappa}], \\ \text{Err}[e_4 \check{\kappa}] &= -\frac{1}{2} \check{\kappa} \check{\kappa} - \frac{1}{2} \overline{\kappa} \overline{\kappa} + \left(-\frac{1}{2} \check{\vartheta} \check{\vartheta} + 2\zeta^2 \right) - \overline{\left(-\frac{1}{2} \check{\vartheta} \check{\vartheta} + 2\zeta^2 \right)}. \end{aligned}$$

This yields

$$\begin{aligned} e_4 \left(r \int_S (\check{\kappa})^2 \right) &= \int_S r \left(2\check{\kappa} e_4(\check{\kappa}) + \kappa \check{\kappa}^2 + \frac{e_4(r)}{r} \check{\kappa}^2 \right) \\ &= \int_S r \left(2\check{\kappa} \left(-\frac{1}{2} \check{\kappa} \overline{\kappa} - 2\check{\vartheta}_1 \zeta + 2\check{\rho} + \text{Err}[e_4 \check{\kappa}] \right) + \check{\kappa} \check{\kappa}^2 \right) \end{aligned}$$

and hence, using the the estimates of Step 1 and 4 for $\check{\kappa}$, ζ , $\check{\vartheta}$ and $\check{\rho}$, and the control of $\overline{\kappa}$ and $\overline{\kappa}$ in Lemma 3.4.1, we infer

$$e_4 \left(r \int_S (\check{\kappa})^2 \right) \lesssim \frac{\epsilon}{r^2} \int_S r \check{\kappa}^2 + \epsilon r^{-\frac{3}{2}} \left(\int_S r \check{\kappa}^2 \right)^{\frac{1}{2}}$$

and hence

$$e_4 \left(\left(r \int_S (\check{\kappa})^2 \right)^{\frac{1}{2}} \right) \lesssim \frac{\epsilon}{r^2} \left(r \int_S (\check{\kappa})^2 \right)^{\frac{1}{2}} + \epsilon r^{-\frac{3}{2}}$$

Integrating backward from Σ_* , where $\check{\kappa}$ under control in view of Step 1, we infer

$$\sup_{r \geq 4m_0} r^2 \int_S \check{\kappa}^2 \lesssim \epsilon^2.$$

Next, commuting the equation for $e_4(\check{\underline{\kappa}})$ with \mathbf{T} and $\not\partial$ together with the commutator estimate of Step 3, using the equation for $e_4(\underline{\kappa})$ to recover the e_4 derivatives, we obtain similarly for higher order derivatives

$$\max_{k \leq k_{large} - 2} \sup_{r \geq 4m_0} r^4 \int_S (\not\partial^k \check{\underline{\kappa}})^2 \lesssim \epsilon^2.$$

Step 6. In view of Codazzi for $\underline{\vartheta}$, and the estimates of Step 1 on ζ , and $\underline{\vartheta}$ and of Step 3 on $\check{\underline{\kappa}}$ in $(ext)\mathcal{M}(r \geq 4m_0)$, we infer

$$\max_{k \leq k_{large} - 2} \sup_{(ext)\mathcal{M}(r \geq 4m_0)} r \|\not\partial^k \underline{\beta}\|_{L^2(S)} \lesssim \epsilon.$$

Step 7. In view of the null structure equation for $e_3(\underline{\kappa})$, and the estimates of Step 1 on $\check{\underline{\omega}}$, ζ , η and $\underline{\vartheta}$, and of Step 3 on $\check{\underline{\kappa}}$ in $(ext)\mathcal{M}(r \geq 4m_0)$, we infer

$$\max_{k \leq k_{large} - 3} \sup_{(ext)\mathcal{M}(r \geq 4m_0)} \|\not\partial^k \underline{\xi}\|_{L^2(S)} \lesssim \epsilon.$$

Step 8. In view of the Bianchi identity for $e_3(\underline{\beta})$, and the estimates of Step 1 on $\check{\underline{\omega}}$, ζ , and η , the estimates of Step 2 on $\check{\rho}$, of Step 3 on $\check{\underline{\kappa}}$ and of Step 5 on $\underline{\xi}$ in $(ext)\mathcal{M}(r \geq 4m_0)$, we infer

$$\max_{k \leq k_{large} - 3} \sup_{(ext)\mathcal{M}(r \geq 4m_0)} \|\not\partial^k \underline{\alpha}\|_{L^2(S)} \lesssim \epsilon.$$

Step 9. Gathering the estimates for Step 1 to Step 8, we have obtained

$$\begin{aligned} & \max_{k \leq k_{large} - 1} \sup_{(ext)\mathcal{M}(r \geq 4m_0)} \left\{ r^{\frac{5}{2} + \frac{\delta_B}{2}} (\|\not\partial^k \alpha\|_{L^2(S)} + \|\not\partial^k \beta\|_{L^2(S)}) \right. \\ & + r (\|\not\partial^k \check{\kappa}\|_{L^2(S)} + \|\not\partial^k \zeta\|_{L^2(S)} + \|\not\partial^k \vartheta\|_{L^2(S)}) + \|\not\partial^k \underline{\vartheta}\|_{L^2(S)} + \|\not\partial^k \check{\vartheta}\|_{L^2(S)} + \|\not\partial^k \check{\underline{\omega}}\|_{L^2(S)} \left. \right\} \\ & + \max_{k \leq k_{large} - 2} \sup_{(ext)\mathcal{M}(r \geq 4m_0)} \left\{ r^2 (\|\not\partial^k \mu\|_{L^2(S)} + \|\not\partial^k \check{\rho}\|_{L^2(S)}) + r (\|\not\partial^k \check{\underline{\kappa}}\|_{L^2(S)} + \|\not\partial^k \underline{\beta}\|_{L^2(S)}) \right\} \\ & + \max_{k \leq k_{large} - 3} \sup_{(ext)\mathcal{M}(r \geq 4m_0)} \left\{ \|\not\partial \underline{\xi}\|_{L^2(S)} + \|\not\partial^k \underline{\alpha}\|_{L^2(S)} \right\} \lesssim \epsilon. \end{aligned}$$

Using the Sobolev embedding on the 2-sphere S which loses 2 derivatives, and in view of the previous estimate on $(ext)\mathcal{M}(r \leq 4m_0)$, we infer

$$\begin{aligned} & \max_{k \leq k_{large} - 5} \sup_{\mathcal{M}} \left\{ r^{\frac{7}{2} + \frac{\delta_B}{2}} (|\not\partial^k \alpha| + |\not\partial^k \beta|) + r^3 (|\not\partial^k \check{\mu}| + |\not\partial^k \check{\rho}|) \right. \\ & + r^2 (|\not\partial^k \check{\kappa}| + |\not\partial^k \zeta| + |\not\partial^k \vartheta| + |\not\partial^k \check{\underline{\kappa}}| + |\not\partial^k \underline{\beta}|) \\ & \left. + r (|\not\partial^k \underline{\vartheta}| + |\not\partial^k \check{\vartheta}| + |\not\partial^k \check{\underline{\omega}}| + |\not\partial \underline{\xi}| + |\not\partial^k \underline{\alpha}|) \right\} \lesssim \epsilon \end{aligned}$$

which is the desired estimate on ${}^{(ext)}\mathcal{M}(r \geq 4m_0)$. This concludes the proof of Proposition 3.4.5.

4.5 Proof of Proposition 3.4.6

Let (e_4, e_3, e_θ) the outgoing geodesic null frame of ${}^{(ext)}\mathcal{M}$. We will exhibit another frame (e'_4, e'_3, e'_θ) of ${}^{(ext)}\mathcal{M}$ provided by

$$\begin{aligned} e'_4 &= e_4 + fe_\theta + \frac{1}{4}f^2e_3, \\ e'_\theta &= e_\theta + \frac{1}{2}fe_3, \\ e'_3 &= e_3, \end{aligned} \tag{4.5.1}$$

where f is such that

$$f = 0 \text{ on } \Sigma_* \cap \mathcal{C}_*, \quad \eta' = 0 \text{ on } \Sigma_*, \quad \xi' = 0 \text{ on } {}^{(ext)}\mathcal{M}. \tag{4.5.2}$$

The desired estimates for the Ricci coefficients and curvature components with respect to the new frame (e'_4, e'_3, e'_θ) of ${}^{(ext)}\mathcal{M}$ will be obtained using

- the change of frame formulas of Proposition 2.3.4, applied to the change of frame from (e_4, e_3, e_θ) to (e'_4, e'_3, e'_θ) ,
- the estimates for f on ${}^{(ext)}\mathcal{M}$,
- the estimates for the Ricci coefficients and curvature components with respect to the outgoing geodesic frame (e_4, e_3, e_θ) of ${}^{(ext)}\mathcal{M}$ provided by the bootstrap assumptions on decay and Proposition 3.4.5.

Step 1. We start by deriving an equation for f on ${}^{(ext)}\mathcal{M}$. In view of the condition $\xi' = 0$ on ${}^{(ext)}\mathcal{M}$, see (4.5.2), in view of $\xi = \omega = 0$ and $\underline{\eta} = -\zeta$ satisfied by the outgoing geodesic foliation of ${}^{(ext)}\mathcal{M}$, and in view of Lemma 2.3.5, we have

$$\begin{aligned} e'_4(f) + \frac{1}{2}\kappa f &= -\frac{1}{2}f\vartheta - \frac{1}{2}f^2\eta - \frac{3}{2}f^2\zeta \\ &\quad + \frac{1}{8}f^3\underline{\kappa} + \frac{1}{2}f^3\underline{\omega} + \frac{1}{8}f^3\underline{\vartheta} + \frac{1}{8}f^4\underline{\xi} \text{ on } {}^{(ext)}\mathcal{M}. \end{aligned} \tag{4.5.3}$$

We also derive an equation for f on Σ_* . In view of the condition $\eta' = 0$ on Σ_* , see (4.5.2), and in view of Lemma 2.3.5, we have

$$e'_3(f) = -2\eta + 2f\underline{\omega} + \frac{1}{2}f^2\underline{\xi} \quad \text{on } \Sigma_*. \quad (4.5.4)$$

Now, since $u + r$ is constant on Σ_* , the following vectorfield

$$\nu'_{\Sigma_*} := e'_3 + a'e'_4, \quad a' := -\frac{e'_3(u+r)}{e'_4(u+r)},$$

is tangent to Σ_* . We compute in view of the above

$$\begin{aligned} \nu'_{\Sigma_*}(f) &= e'_3(f) + a'e'_4(f) \\ &= -2\eta + 2f\underline{\omega} + \frac{1}{2}f^2\underline{\xi} + a' \left\{ -\frac{1}{2}\kappa f - \frac{1}{2}f\vartheta - \frac{1}{2}f^2\eta - \frac{3}{2}f^2\zeta \right. \\ &\quad \left. + \frac{1}{8}f^3\underline{\kappa} + \frac{1}{2}f^3\underline{\omega} + \frac{1}{8}f^3\underline{\vartheta} + \frac{1}{8}f^4\underline{\xi} \right\}. \end{aligned}$$

Using (4.5.1), we have

$$\begin{aligned} a' &= -\frac{e'_3(u+r)}{e'_4(u+r)} \\ &= -\frac{e_3(u+r)}{(e_4 + fe_\theta + \frac{1}{4}f^2e_3)(u+r)} \\ &= -\frac{\frac{2}{\varsigma} + \frac{r}{2}(\overline{\kappa} + \underline{A})}{\frac{r}{2}\overline{\kappa} + \frac{1}{4}f^2\left(\frac{2}{\varsigma} + \frac{r}{2}(\overline{\kappa} + \underline{A})\right)} \end{aligned}$$

and hence

$$\begin{aligned} \nu'_{\Sigma_*}(f) &= -2\eta + 2f\underline{\omega} + \frac{1}{2}f^2\underline{\xi} - \frac{\frac{2}{\varsigma} + \frac{r}{2}(\overline{\kappa} + \underline{A})}{\frac{r}{2}\overline{\kappa} + \frac{1}{4}f^2\left(\frac{2}{\varsigma} + \frac{r}{2}(\overline{\kappa} + \underline{A})\right)} \left\{ -\frac{1}{2}\kappa f - \frac{1}{2}f\vartheta \right. \\ &\quad \left. - \frac{1}{2}f^2\eta - \frac{3}{2}f^2\zeta + \frac{1}{8}f^3\underline{\kappa} + \frac{1}{2}f^3\underline{\omega} + \frac{1}{8}f^3\underline{\vartheta} + \frac{1}{8}f^4\underline{\xi} \right\} \quad \text{on } {}^{(ext)}\mathcal{M}. \quad (4.5.5) \end{aligned}$$

Step 2. Next, we estimate f on Σ_* . Introducing an integer k_{loss} and a small constant $\delta_0 > 0$ satisfying

$$16 \leq k_{loss} \leq \frac{\delta_{dec}}{3}(k_{large} - k_{small}), \quad \delta_0 = \frac{k_{loss}}{k_{large} - k_{small}},$$

we assume the following local bootstrap assumption

$$|\mathfrak{d}^{\leq k_{small} + k_{loss} + 2} f| \leq \frac{\sqrt{\epsilon}}{r u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}} \quad \text{on } u_1 \leq u \leq u_* \quad (4.5.6)$$

where

$$1 \leq u_1 < u_*.$$

Since $f = 0$ on $\Sigma_* \cap \mathcal{C}_*$ in view of (4.5.2), (4.5.6) holds for u_1 close enough to u_* , and our goal is to prove that we may in fact choose $u_1 = 1$ and replace $\sqrt{\epsilon}$ with ϵ in (4.5.6).

In view of the estimates for the Ricci coefficients and curvature components with respect to the outgoing geodesic frame (e_4, e_3, e_θ) of $^{(ext)}\mathcal{M}$ provided by Proposition 3.4.5, (4.5.5) yields

$$\nu'_{\Sigma_*}(f) = -2\eta + h, \quad |\mathfrak{d}^k h| \lesssim r^{-1} (|\mathfrak{d}^{\leq k} f| + |\mathfrak{d}^{\leq k} f|^4) \text{ for } k \leq k_{large} - 5.$$

Using commutator identities, using also (4.5.3) and (4.5.4), and in view of (4.5.6), we infer

$$|\nu'_{\Sigma_*}(\mathfrak{d}^k f)| \lesssim |\mathfrak{d}^{\leq k} \eta| + \frac{\sqrt{\epsilon}}{r^2 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2, \quad u_1 \leq u \leq u_*.$$

Since $f = 0$ on $\Sigma_* \cap \mathcal{C}_*$ in view of (4.5.2), and since ν'_{Σ_*} is tangent to Σ_* , we deduce on Σ_* , integrating along the integral curve of ν'_{Σ_*}

$$|\mathfrak{d}^k f| \lesssim \int_u^{u_*} |\mathfrak{d}^{\leq k} \eta| + \frac{\sqrt{\epsilon}}{u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}} \int_u^{u_*} \frac{1}{\nu'_{\Sigma_*}(u') r^2} \text{ for } k \leq k_{small} + k_{loss} + 2, \quad u_1 \leq u \leq u_*.$$

Since

$$\begin{aligned} \nu'_{\Sigma_*}(u) &= e'_3(u) + a' e'_4(u) \\ &= e_3(u) - \frac{\frac{2}{\varsigma} + \frac{r}{2}(\bar{\kappa} + \underline{A})}{\frac{r}{2}\bar{\kappa} + \frac{1}{4}f^2\left(\frac{2}{\varsigma} + \frac{r}{2}(\bar{\kappa} + \underline{A})\right)} \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right) u \\ &= \frac{2}{\varsigma} - \frac{f^2 \frac{2}{\varsigma} + \frac{r}{2}(\bar{\kappa} + \underline{A})}{2\varsigma \frac{r}{2}\bar{\kappa} + \frac{1}{4}f^2\left(\frac{2}{\varsigma} + \frac{r}{2}(\bar{\kappa} + \underline{A})\right)} \end{aligned}$$

we have

$$\nu'_{\Sigma_*}(u) = 2 + O(\epsilon)$$

and hence

$$|\mathfrak{d}^k f| \lesssim \int_u^{u_*} |\mathfrak{d}^{\leq k} \eta| + \frac{\sqrt{\epsilon}}{u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}} \int_u^{u_*} \frac{1}{r^2} \text{ for } k \leq k_{small} + k_{loss} + 2, \quad u_1 \leq u \leq u_*.$$

Together with the behavior (3.3.4) of r on Σ_* , we infer

$$|\mathfrak{d}^k f| \lesssim \int_u^{u_*} |\mathfrak{d}^{\leq k} \eta| + \frac{\epsilon}{r u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2, \quad u_1 \leq u \leq u_*.$$

Next, we estimate η . We have by interpolation, since $k_{loss} \leq k_{large} - k_{small}$,

$$\|\mathfrak{d}^{\leq k_{small} + k_{loss} + 4} \eta\|_{L^2(S)} \lesssim \|\mathfrak{d}^{\leq k_{small}} \eta\|_{L^2(S)}^{1 - \frac{k_{loss} + 4}{k_{large} - k_{small}}} \|\mathfrak{d}^{\leq k_{large}} \eta\|_{L^2(S)}^{\frac{k_{loss} + 4}{k_{large} - k_{small}}},$$

and hence, using $\delta_0 > 0$, we have

$$\begin{aligned} & \int_{\Sigma_*(\geq u)} |\mathfrak{d}^{\leq k_{small} + k_{loss} + 4} \eta| \\ & \lesssim \left(\int_{\Sigma_*(\geq u)} u'^{1 + \delta_0} |\mathfrak{d}^{\leq k_{small} + k_{loss} + 4} \eta|^2 \right)^{\frac{1}{2}} \\ & \lesssim \frac{1}{u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}} \left(\int_{\Sigma_*} u'^{2 + 2\delta_{dec}} |\mathfrak{d}^{\leq k_{small}} \eta|^2 \right)^{\frac{1}{2} - \frac{k_{loss} + 4}{2(k_{large} - k_{small})}} \left(\int_{\Sigma_*} |\mathfrak{d}^{\leq k_{large}} \eta|^2 \right)^{\frac{k_{loss} + 4}{2(k_{large} - k_{small})}}. \end{aligned}$$

where we have used the fact that

$$\frac{k_{loss} + 4}{k_{large} - k_{small}} (1 + \delta_{dec}) + \frac{\delta_0}{2} = \left(\left(1 + \frac{4}{k_{loss}} \right) (1 + \delta_{dec}) + \frac{1}{2} \right) \delta_0 \leq 2\delta_0$$

and

$$\frac{1}{2} + \delta_{dec} - 2\delta_0 = \frac{1}{2} + \delta_{dec} - \frac{4k_{loss}}{k_{large} - k_{small}} \geq \delta_{dec} > 0$$

since $16 \leq k_{loss} \leq \frac{1}{8}(k_{large} - k_{small})$ and $\delta_{dec} > 0$ is small. Now, recall from the bootstrap assumptions on decay and energy for η along Σ_* that we have

$$\int_{\Sigma_*} u^{2 + 2\delta_{dec}} |\mathfrak{d}^{\leq k_{small}} \eta|^2 + \int_{\Sigma_*} |\mathfrak{d}^{\leq k_{large}} \eta|^2 \leq \epsilon^2.$$

We deduce

$$\int_{\Sigma_*(\geq u)} |\mathfrak{d}^{\leq k_{small} + k_{loss} + 4} \eta| \lesssim \frac{\epsilon}{u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}}.$$

Together with the Sobolev embedding on the 2-spheres S foliating Σ_* , as well as the behavior (3.3.4) of r on Σ_* , we infer

$$\int_u^{u_*} |\mathfrak{D}^{\leq k_{small} + k_{loss} + 2} \eta| \lesssim \frac{\epsilon}{u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}}.$$

Plugging in the above estimate for f , we infer

$$|\mathfrak{D}^k f| \lesssim \frac{\epsilon}{r u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2, \quad u_1 \leq u \leq u_*.$$

Together with (4.5.3) and (4.5.4), we recover e_4 and e_3 derivatives to deduce

$$|\mathfrak{D}^k f| \lesssim \frac{\epsilon}{r u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2, \quad u_1 \leq u \leq u_*.$$

This is an improvement of the bootstrap assumption (4.5.6). Thus, we may choose $u_1 = 1$, and f satisfies the following estimate

$$|\mathfrak{D}^k f| \lesssim \frac{\epsilon}{r u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2 \text{ on } \Sigma_*.$$

Together with (4.5.4), as well as the behavior (3.3.4) of r on Σ_* , we infer

$$\begin{aligned} |\mathfrak{D}^{k-1} e'_3 f| &\lesssim |\mathfrak{D}^{k-1} \eta| + \frac{\epsilon}{r^2} \\ &\lesssim \frac{\epsilon}{r u^{1 + \delta_{dec} - 2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2 \text{ on } \Sigma_*. \end{aligned}$$

Collecting the two above estimates, we obtain

$$|\mathfrak{D}^k f| \lesssim \frac{\epsilon}{r u^{\frac{1}{2} + \delta_{dec} - 2\delta_0}}, \quad |\mathfrak{D}^{k-1} e'_3 f| \lesssim \frac{\epsilon}{r u^{1 + \delta_{dec} - 2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2 \text{ on } \Sigma_*. \quad (4.5.7)$$

Step 3. Next, we estimate f on $^{(ext)}\mathcal{M}$. We assume the following local bootstrap assumption

$$|\mathfrak{D}^{\leq k_{small} + k_{loss} + 2} f| \leq \frac{\sqrt{\epsilon}}{r u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + u^{1 + \delta_{dec} - 2\delta_0}} \text{ on } r \geq r_1. \quad (4.5.8)$$

where $r_1 \geq 4m_0$. In view of the control of f on Σ_* provided by (4.5.7), (4.5.8) holds for r_1 sufficiently large, and our goal is to prove that we may in fact choose $r_1 = 4m_0$ and replace $\sqrt{\epsilon}$ with ϵ in (4.5.8).

Recall (4.5.3)

$$\begin{aligned} e'_4(f) + \frac{1}{2}\kappa f &= -\frac{1}{2}f\vartheta - \frac{1}{2}f^2\eta - \frac{3}{2}f^2\zeta \\ &\quad + \frac{1}{8}f^3\underline{\kappa} + \frac{1}{2}f^3\underline{\omega} + \frac{1}{8}f^3\underline{\vartheta} + \frac{1}{8}f^4\underline{\zeta} \text{ on } ^{(ext)}\mathcal{M}. \end{aligned}$$

In view of the estimates for the Ricci coefficients and curvature components with respect to the outgoing geodesic frame (e_4, e_3, e_θ) of $^{(ext)}\mathcal{M}$ provided by Proposition 3.4.5,

$$\begin{aligned} & \left| \mathfrak{d}^k \left(-\frac{1}{2}f\vartheta - \frac{1}{2}f^2\eta - \frac{3}{2}f^2\zeta + \frac{1}{8}f^3\kappa + \frac{1}{2}f^3\omega + \frac{1}{8}f^3\underline{\vartheta} + \frac{1}{8}f^4\underline{\xi} \right) \right| \\ & \lesssim \epsilon r^{-2}u^{-\frac{1}{2}}|\mathfrak{d}^{\leq k}f| + r^{-1}(|\mathfrak{d}^{\leq k}f|^2 + |\mathfrak{d}^{\leq k}f|^4) \text{ for } k \leq k_{large} - 5. \end{aligned}$$

Using commutator identities, using also (4.5.3), and in view of (4.5.8), we infer³

$$e'_4 \left((\mathfrak{d}, T)^k f \right) + \frac{1}{2}\kappa(\mathfrak{d}, T)^k f \leq \frac{\epsilon}{r^3 u^{1+\delta_{dec}-2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2, r \geq r_1.$$

Integrating backwards from Σ_* where we have (4.5.7), we deduce⁴

$$|(\mathfrak{d}, T)^k f| \leq \frac{\epsilon}{r u^{\frac{1}{2}+\delta_{dec}-2\delta_0} + u^{1+\delta_{dec}-2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2, r \geq r_1.$$

Together with (4.5.3), we recover the e_4 derivatives and obtain

$$|\mathfrak{d}^k f| \leq \frac{\epsilon}{r u^{\frac{1}{2}+\delta_{dec}-2\delta_0} + u^{1+\delta_{dec}-2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2, r \geq r_1.$$

This is an improvement of the bootstrap assumption (4.5.8). Thus, we may choose $r_1 = 4m_0$, and we have

$$|\mathfrak{d}^k f| \lesssim \frac{\epsilon}{r u^{\frac{1}{2}+\delta_{dec}-2\delta_0} + u^{1+\delta_{dec}-2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2 \text{ on } ^{(ext)}\mathcal{M}.$$

Also, commuting once (4.5.3) with e'_3 , using the commutator identity $[e'_3, e'_4] = 2\underline{\omega}'e'_4 - 2\omega'e'_3 + (\eta' - \underline{\eta}')e'_\theta$, and proceeding as above to integrate backward from Σ_* where $e'_3 f$ is under control from (4.5.7), we also obtain

$$|\mathfrak{d}^{k-1}e'_3 f| \lesssim \frac{\epsilon}{r u^{1+\delta_{dec}-2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2 \text{ on } ^{(ext)}\mathcal{M}.$$

³Note that

$$\delta_{dec} - 2\delta_0 = \delta_{dec} - \frac{2k_{loss}}{k_{large} - k_{small}} \geq \frac{\delta_{dec}}{3} > 0$$

where we have used the definition of δ_0 and the upper bound on k_{loss} .

⁴Note that (4.5.7) yields

$$|\mathfrak{d}^k f| \lesssim \frac{\epsilon}{u^{1+\delta_{dec}-2\delta_0}} \text{ for } k \leq k_{small} + k_{loss} + 2 \text{ on } \Sigma_*.$$

in view of the behavior (3.3.4) of r on Σ_* .

Collecting the two above estimates, we obtain

$$\begin{aligned} |\mathfrak{d}^k f| &\lesssim \frac{\epsilon}{ru^{\frac{1}{2}+\delta_{dec}-2\delta_0} + u^{1+\delta_{dec}-2\delta_0}}, \quad \text{for } k \leq k_{small} + k_{loss} + 2 \text{ on } {}^{(ext)}\mathcal{M}, \\ |\mathfrak{d}^{k-1} e'_3 f| &\lesssim \frac{\epsilon}{ru^{1+\delta_{dec}-2\delta_0}} \quad \text{for } k \leq k_{small} + k_{loss} + 2 \text{ on } {}^{(ext)}\mathcal{M}, \end{aligned} \quad (4.5.9)$$

which is the desired estimate for f .

Step 4. In view of Proposition 2.3.4 applied to our particular case, i.e. a triplet $(f, \underline{f}, \lambda)$ with $\underline{f} = 0$ and $\lambda = 1$, and the fact that the frame (e_4, e_3, e_θ) is outgoing geodesic, we have

$$\begin{aligned} \underline{\xi}' &= \underline{\xi}, \\ \zeta' &= \zeta - \frac{1}{4}f\underline{\kappa} - f\underline{\omega} - \frac{1}{4}f\underline{\vartheta} + \text{l.o.t.}, \\ \eta' &= \eta + \frac{1}{2}e'_3(f) - f\underline{\omega} + \text{l.o.t.}, \\ \underline{\eta}' &= -\zeta + \frac{1}{4}\underline{\kappa}f + \frac{1}{4}f\underline{\vartheta} + \text{l.o.t.}, \end{aligned}$$

$$\begin{aligned} \kappa' &= \kappa + \mathfrak{d}_1'(f) + f(\zeta + \eta) - \frac{1}{4}f^2\underline{\kappa} - f^2\underline{\omega} + \text{l.o.t.}, \\ \underline{\kappa}' &= \underline{\kappa} + f\underline{\xi} + \text{l.o.t.}, \\ \vartheta' &= \vartheta - \mathfrak{d}_2'(f) + f(\zeta + \eta) - f^2\underline{\omega} + \text{l.o.t.}, \\ \underline{\vartheta}' &= \underline{\vartheta} + f\underline{\xi} + \text{l.o.t.}, \end{aligned}$$

$$\begin{aligned} \omega' &= f\zeta - \frac{1}{8}\underline{\kappa}f^2 - \frac{1}{4}\underline{\omega}f^2 + \text{l.o.t.}, \\ \underline{\omega}' &= \underline{\omega} + \frac{1}{2}f\underline{\xi}, \end{aligned}$$

and

$$\begin{aligned} \alpha' &= \alpha + 2f\beta + \frac{3}{2}f^2\rho + \text{l.o.t.}, \\ \beta' &= \beta + \frac{3}{2}\rho f + \text{l.o.t.}, \\ \rho' &= \rho + f\underline{\beta} + \text{l.o.t.}, \\ \underline{\beta}' &= \underline{\beta} + \frac{1}{2}f\underline{\alpha}, \\ \underline{\alpha}' &= \underline{\alpha}. \end{aligned} \quad (4.5.10)$$

where the lower order terms denoted by l.o.t. are linear with respect to $\xi, \underline{\xi}, \vartheta, \kappa, \eta, \underline{\eta}, \zeta, \underline{\kappa}, \underline{\vartheta}$ and $\alpha, \beta, \rho, \underline{\beta}, \underline{\alpha}$, and quadratic or higher order in f , and do not contain derivatives of the latter. Together with the estimates (4.5.9) for f on $^{(ext)}\mathcal{M}$, and the estimates for the Ricci coefficients and curvature components with respect to the outgoing geodesic frame (e_4, e_3, e_θ) of $^{(ext)}\mathcal{M}$ provided by the bootstrap assumptions on decay and Proposition 3.4.5, we immediately infer

$$\begin{aligned} \max_{0 \leq k \leq k_{small} + k_{loss} + 1} \sup_{(ext)\mathcal{M}} & \left\{ \left(r^2 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r u^{1 + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^k(\Gamma'_g \setminus \{\eta'\})| + r u^{1 + \delta_{dec} - 2\delta_0} |\mathfrak{d}^k \Gamma'_b| \right. \\ & + r^2 u^{1 + \delta_{dec} - 2\delta_0} \left| \mathfrak{d}^{k-1} e'_3 \left(\kappa' - \frac{2}{r}, \underline{\kappa}' + \frac{2\Upsilon}{r}, \vartheta', \zeta', \underline{\eta}' \right) \right| \\ & + \left(r^3 (u + 2r)^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r^2 (u + 2r)^{1 + \delta_{dec} - 2\delta_0} \right) \left(|\mathfrak{d}^k \alpha'| + |\mathfrak{d}^k \beta'| \right) \\ & + \left(r^3 (2r + u)^{1 + \delta_{dec}} + r^4 (2r + u)^{\frac{1}{2} + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^{k-1} e'_3(\alpha')| \\ & + \left(r^3 u^{1 + \delta_{dec}} + r^4 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^{k-1} e'_3(\beta')| \\ & + \left(r^3 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r^2 r u^{1 + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^k \check{\rho}'| \\ & \left. + u^{1 + \delta_{dec} - 2\delta_0} \left(r^2 |\mathfrak{d}^k \underline{\beta}'| + r |\mathfrak{d}^k \underline{\alpha}'| \right) \right\} \lesssim \epsilon \end{aligned} \quad (4.5.11)$$

where we have introduced the notation

$$\Gamma'_g \setminus \{\eta'\} = \left\{ r\omega', \kappa' - \frac{2}{r}, \vartheta', \zeta', \underline{\eta}', \underline{\kappa}' + \frac{2\Upsilon}{r}, r^{-1}(e'_4(r) - 1), r^{-1}e'_\theta(r), e'_4(m) \right\}.$$

Note also, in view of the above transformation formula for ω' , i.e.

$$\omega' = f\zeta - \frac{1}{8}\underline{\kappa}f^2 - \frac{1}{4}\underline{\omega}f^2 + \text{l.o.t.},$$

that we have in fact a gain of r^{-1} for ω' compared to (4.5.11), i.e.

$$\max_{0 \leq k \leq k_{small} + k_{loss} + 1} \sup_{(ext)\mathcal{M}} \left(r^3 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r^2 u^{1 + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^k \omega'| \lesssim \epsilon. \quad (4.5.12)$$

We now focus on estimating η' . Proceeding as for the other Ricci coefficients would yield for η' the same behavior than η and hence a loss of r^{-1} compared to the desired estimate. Instead, we rely on the following null structure equation which follow from Proposition 2.2.1 and the fact that $\xi' = 0$

$$e'_4(\eta' - \zeta') + \frac{1}{2}\kappa'(\eta' - \zeta') = 2\mathfrak{d}_1^* \omega' - \frac{1}{2}\vartheta'(\eta' - \zeta').$$

Next,

- we commute with \mathfrak{D}' and T' , and we rely on the corresponding commutator identities,
- we use the above equation for $e'_4(\eta')$ to recover the e'_4 derivatives,
- we rely on the estimates (4.5.11), as well as the estimate (4.5.12) for ω' ,

which allows us to derive

$$\left| e'_4(\mathfrak{D}^k(\eta' - \zeta')) + \frac{1}{2}\kappa'\mathfrak{D}^k(\eta' - \zeta') \right| \lesssim \frac{\epsilon}{r^4u^{\frac{1}{2}+\delta_{dec}-2\delta_0} + r^3u^{1+\delta_{dec}-2\delta_0}} + \frac{\epsilon}{r^2}|\mathfrak{D}^{\leq k}(\eta' - \zeta')|, \quad k \leq k_{small} + k_{loss}.$$

Integrating backwards from Σ_* where $\eta' = 0$ in view of (4.5.2), and using the control ζ' provided by (4.5.11), we infer

$$\begin{aligned} & \max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{(ext)\mathcal{M}} \left(r^2u^{\frac{1}{2}+\delta_{dec}-2\delta_0} + ru^{1+\delta_{dec}-2\delta_0} \right) |\mathfrak{D}^k \eta'| \\ & \lesssim \epsilon + \max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{(ext)\mathcal{M}} \left(r^2u^{\frac{1}{2}+\delta_{dec}-2\delta_0} + ru^{1+\delta_{dec}-2\delta_0} \right) |\mathfrak{D}^k \zeta'| \\ & \lesssim \epsilon. \end{aligned}$$

Also, commuting first the equation for $e'_4(\eta' - \zeta')$ with e'_3 , using the commutator identity $[e'_3, e'_4] = 2\underline{\omega}'e'_4 - 2\omega'e'_3 + (\eta' - \underline{\eta}')e'_\theta$, and proceeding as above to integrate backward from Σ_* , we also obtain

$$\begin{aligned} & \max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{(ext)\mathcal{M}} r^2u^{1+\delta_{dec}-2\delta_0} |\mathfrak{D}^{k-1} e'_3 \eta'| \\ & \lesssim \epsilon + \max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{(ext)\mathcal{M}} r^2u^{1+\delta_{dec}-2\delta_0} |\mathfrak{D}^{k-1} e'_3 \zeta'| \\ & \lesssim \epsilon. \end{aligned}$$

Thus, together with (4.5.11), we infer

$$\begin{aligned}
& \max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{(ext)\mathcal{M}} \left\{ \left(r^2 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r u^{1 + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^k \Gamma'_g| + r u^{1 + \delta_{dec} - 2\delta_0} |\mathfrak{d}^k \Gamma'_b| \right. \\
& \quad + r^2 u^{1 + \delta_{dec} - 2\delta_0} \left| \mathfrak{d}^{k-1} e'_3 \left(\kappa' - \frac{2}{r}, \underline{\kappa}' + \frac{2\Upsilon}{r}, \vartheta', \zeta', \underline{\eta}', \eta' \right) \right| \\
& \quad + \left(r^3 (u + 2r)^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r^2 (u + 2r)^{1 + \delta_{dec} - 2\delta_0} \right) \left(|\mathfrak{d}^k \alpha'| + |\mathfrak{d}^k \beta'| \right) \\
& \quad + \left(r^3 (2r + u)^{1 + \delta_{dec}} + r^4 (2r + u)^{\frac{1}{2} + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^{k-1} e'_3(\alpha')| \\
& \quad + \left(r^3 u^{1 + \delta_{dec}} + r^4 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^{k-1} e'_3(\beta')| \\
& \quad + \left(r^3 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r^2 r u^{1 + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^k \check{\rho}'| \\
& \quad \left. + u^{1 + \delta_{dec} - 2\delta_0} \left(r^2 |\mathfrak{d}^k \underline{\beta}'| + r |\mathfrak{d}^k \underline{\alpha}'| \right) \right\} \lesssim \epsilon.
\end{aligned}$$

Together with the fact that $\xi' = 0$ in view of (4.5.2), this concludes the proof of Proposition 3.4.6.

4.6 Existence and control of the global frames

4.6.1 Proof of Proposition 3.5.2

To match the frame of $(int)\mathcal{M}$ and a conformal renormalization of the frame of $(ext)\mathcal{M}$, we will need to introduce a cut-off function.

Definition 4.6.1. *Let $\psi : \mathbb{R} \rightarrow \mathbb{R}$ a smooth cut-off function such that $0 \leq \psi \leq 1$, $\psi = 0$ on $(-\infty, 0]$ and $\psi = 1$ on $[1, +\infty)$. We define $\psi_{m_0, \delta_{\mathcal{H}}}$ as follows*

$$\psi_{m_0, \delta_{\mathcal{H}}}(r) = \begin{cases} 1 & \text{if } r \geq 2m_0 \left(1 + \frac{3}{2} \delta_{\mathcal{H}} \right), \\ 0 & \text{if } r \leq 2m_0 \left(1 + \frac{1}{2} \delta_{\mathcal{H}} \right), \end{cases}$$

and

$$\psi_{m_0, \delta_{\mathcal{H}}}(r) = \psi \left(\frac{r - 2m_0 \left(1 + \frac{1}{2} \delta_{\mathcal{H}} \right)}{2m_0 \delta_{\mathcal{H}}} \right) \text{ on } 2m_0 \left(1 + \frac{1}{2} \delta_{\mathcal{H}} \right) \leq r \leq 2m_0 \left(1 + \frac{3}{2} \delta_{\mathcal{H}} \right).$$

We are now ready to define the global frame of the statement of Proposition 3.5.2.

Definition 4.6.2 (Definition of the global frame). *We introduce a global null frame defined on ${}^{(ext)}\mathcal{M} \cup {}^{(int)}\mathcal{M}$ and denoted by $({}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta)$. The global frame is defined as follows*

1. In ${}^{(ext)}\mathcal{M} \setminus Match$, we have

$$({}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta) = ({}^{(ext)}\Upsilon {}^{(ext)}e_4, {}^{(ext)}\Upsilon^{-1} {}^{(ext)}e_3, {}^{(ext)}e_\theta).$$

2. In ${}^{(int)}\mathcal{M} \setminus Match$, we have

$$({}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta) = ({}^{(int)}e_4, {}^{(int)}e_3, {}^{(int)}e_\theta).$$

3. It remains to define the global frame on the matching region. We denote by $(f, \underline{f}, \lambda)$ the reduced scalars such that we have in the matching region

$$\begin{aligned} {}^{(ext)}e_4 &= \lambda \left({}^{(int)}e_4 + f {}^{(int)}e_\theta + \frac{1}{4} f^2 {}^{(int)}e_3 \right), \\ {}^{(ext)}e_\theta &= \left(1 + \frac{1}{2} f \underline{f} \right) {}^{(int)}e_\theta + \frac{f}{2} {}^{(int)}e_4 + \frac{f}{2} \left(1 + \frac{f \underline{f}}{4} \right) {}^{(int)}e_3, \\ {}^{(ext)}e_3 &= \lambda^{-1} \left(\left(1 + \frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) {}^{(int)}e_3 + \underline{f} \left(1 + \frac{f \underline{f}}{4} \right) {}^{(int)}e_\theta + \frac{f^2}{4} {}^{(int)}e_4 \right), \end{aligned}$$

where we recall that the frame of ${}^{(ext)}\mathcal{M}$ has been extended to ${}^{(int)}\mathcal{M}$, see section 3.5.1. Then, in the matching region, the global frame is given by

$$\begin{aligned} {}^{(glo)}e_4 &= \lambda' \left({}^{(int)}e_4 + f' {}^{(int)}e_\theta + \frac{1}{4} f'^2 {}^{(int)}e_3 \right), \\ {}^{(glo)}e_\theta &= \left(1 + \frac{1}{2} f' \underline{f}' \right) {}^{(int)}e_\theta + \frac{f'}{2} {}^{(int)}e_4 + \frac{f'}{2} \left(1 + \frac{f' \underline{f}'}{4} \right) {}^{(int)}e_3, \\ {}^{(glo)}e_3 &= \lambda'^{-1} \left(\left(1 + \frac{1}{2} f' \underline{f}' + \frac{1}{16} f'^2 \underline{f}'^2 \right) {}^{(int)}e_3 + \underline{f}' \left(1 + \frac{f' \underline{f}'}{4} \right) {}^{(int)}e_\theta + \frac{f'^2}{4} {}^{(int)}e_4 \right), \end{aligned}$$

where

$$\begin{aligned} f' &= \psi_{m_0, \delta_{\mathcal{H}}}({}^{(int)}r) f, & \underline{f}' &= \psi_{m_0, \delta_{\mathcal{H}}}({}^{(int)}r) \underline{f}, \\ \lambda' &= 1 - \psi_{m_0, \delta_{\mathcal{H}}}({}^{(int)}r) + \psi_{m_0, \delta_{\mathcal{H}}}({}^{(int)}r) {}^{(ext)}\Upsilon \lambda. \end{aligned} \tag{4.6.1}$$

Remark 4.6.3. *Recall that the smooth cut-off function ψ in Definition 3.5.1, allowing to define $\psi_{m_0, \delta_{\mathcal{H}}}$, is such that we have in particular $\psi = 0$ on $(-\infty, 0]$ and $\psi = 1$ on $[1, +\infty)$. The following two special cases correspond to the properties (d) i. and (d) ii. of Proposition 3.5.2.*

- If the cut-off ψ in Definition 3.5.1 is such that $\psi = 1$ on $[1/2, +\infty)$, then

$${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta = ({}^{(ext)}\Upsilon {}^{(ext)}e_4, {}^{(ext)}\Upsilon^{-1}({}^{(ext)}e_3, {}^{(ext)}e_\theta) \text{ on } ({}^{(ext)})\mathcal{M}.$$

- If the cut-off ψ in Definition 3.5.1 is such that $\psi = 0$ on $(-\infty, 1/2]$, then

$${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta = ({}^{(int)}e_4, {}^{(int)}e_3, {}^{(int)}e_\theta) \text{ on } ({}^{(int)})\mathcal{M}.$$

Definition 4.6.4 (Global area radius and Hawking mass). *We definition an area radius and a Hawking mass on $({}^{(ext)})\mathcal{M} \cup ({}^{(int)})\mathcal{M}$ as follows*

- On $({}^{(ext)})\mathcal{M} \setminus \text{Match}$, we have

$${}^{(glo)}r = ({}^{(ext)}r, \quad {}^{(glo)}m = ({}^{(ext)}m$$

- On $({}^{(int)})\mathcal{M} \setminus \text{Match}$, we have

$${}^{(glo)}r = ({}^{(int)}r, \quad {}^{(glo)}m = ({}^{(int)}m$$

- On the matching region, we have

$$\begin{aligned} {}^{(glo)}r &= (1 - \psi_{m_0, \delta_{\mathcal{H}}}({}^{(int)}r)) {}^{(int)}r + \psi_{m_0, \delta_{\mathcal{H}}}({}^{(int)}r) ({}^{(ext)}r, \\ {}^{(glo)}m &= (1 - \psi_{m_0, \delta_{\mathcal{H}}}({}^{(int)}r)) {}^{(int)}m + \psi_{m_0, \delta_{\mathcal{H}}}({}^{(int)}r) ({}^{(ext)}m. \end{aligned}$$

The following two lemmas provide the main properties of the global frame.

Lemma 4.6.5. *We have in $({}^{(ext)})\mathcal{M} \setminus \text{Match}$ the following relations between the quantities in the respective frames*

$$\begin{aligned} {}^{(glo)}\alpha &= \Upsilon^{2({}^{(ext)})}\alpha, \quad {}^{(glo)}\beta = \Upsilon({}^{(ext)})\beta, \quad {}^{(glo)}\rho + \frac{2m}{r^3} = ({}^{(ext)}\rho + \frac{2m}{r^3}, \quad {}^{(glo)}\underline{\beta} = \Upsilon^{-1}({}^{(ext)})\underline{\beta}, \\ {}^{(glo)}\underline{\alpha} &= \Upsilon^{-2({}^{(ext)})}\underline{\alpha}, \quad {}^{(glo)}\xi = 0, \quad {}^{(glo)}\underline{\xi} = \Upsilon^{-2({}^{(ext)})}\underline{\xi}, \quad {}^{(glo)}\zeta = -{}^{(glo)}\underline{\eta} = ({}^{(ext)})\zeta, \\ {}^{(glo)}\eta &= ({}^{(ext)})\eta, \quad {}^{(glo)}\omega + \frac{m}{r^2} = -\frac{m}{2r} \left(\overline{{}^{(ext)}\kappa} - \frac{2}{r} \right) + \frac{e_4(m)}{r}, \\ {}^{(glo)}\underline{\omega} &= \Upsilon^{-1} \left(({}^{(ext)})\underline{\omega} - \frac{m}{r^2} + \frac{m}{2\Upsilon r} \left(\overline{{}^{(ext)}\kappa} - \frac{2\Upsilon}{r} \right) + \frac{m}{2\Upsilon r} \left(({}^{(ext)})\check{\underline{\Omega}}\overline{{}^{(ext)}\kappa} - \overline{{}^{(ext)}\check{\underline{\Omega}}({}^{(ext)})\check{\kappa}} \right) - \frac{e_3(m)}{\Upsilon r} \right), \\ {}^{(glo)}\kappa - \frac{2\Upsilon}{r} &= \Upsilon \left(({}^{(ext)}\kappa - \frac{2}{r} \right), \quad {}^{(glo)}\underline{\kappa} + \frac{2}{r} = \Upsilon^{-1} \left(({}^{(ext)}\underline{\kappa} + \frac{2\Upsilon}{r} \right), \\ {}^{(glo)}\check{\kappa} &= \Upsilon({}^{(ext)})\check{\kappa}, \quad {}^{(glo)}\check{\underline{\kappa}} = \Upsilon^{-1}({}^{(ext)})\check{\underline{\kappa}}, \quad {}^{(glo)}\vartheta = \Upsilon({}^{(ext)})\vartheta, \quad {}^{(glo)}\underline{\vartheta} = \Upsilon^{-1}({}^{(ext)})\underline{\vartheta}. \end{aligned}$$

Proof. The proof follows immediately from the change of frame formula with the choice ($f = 0, \underline{f} = 0, \lambda = \Upsilon$), the fact that $e_\theta(\Upsilon) = 0$, and the fact that the frame of ${}^{(ext)}\mathcal{M}$ is outgoing geodesic and thus satisfies in particular $\xi = \omega = 0$ and $\underline{\eta} = -\zeta$. \square

Lemma 4.6.6 (Control of the global frame in the matching region). *In the matching region, the following estimates holds for the global frame⁵*

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-2} \left(\sup_{Match \cap {}^{(int)}\mathcal{M}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k({}^{(glo)}\check{\Gamma}, {}^{(glo)}\check{R})| + \sup_{Match \cap {}^{(ext)}\mathcal{M}} u^{1+\delta_{dec}} |\mathfrak{d}^k({}^{(glo)}\check{\Gamma}, {}^{(glo)}\check{R})| \right) \\ + \max_{0 \leq k \leq k_{large}-1} \left(\int_{Match} |\mathfrak{d}^k({}^{(glo)}\check{\Gamma}, {}^{(glo)}\check{R})|^2 \right)^{\frac{1}{2}} \lesssim \epsilon. \end{aligned}$$

and

$$\left(\int_{Match} |\mathfrak{d}^{k_{large}}({}^{(glo)}\check{\Gamma}, {}^{(glo)}\check{R})|^2 \right)^{\frac{1}{2}} \lesssim \epsilon + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}.$$

Remark 4.6.7. *The quantities associated to the global frame can be estimated as follows*

- In ${}^{(int)}\mathcal{M} \setminus Match$, the global frame coincides with the frame of ${}^{(int)}\mathcal{M}$, and hence, the quantities associated to the global frame satisfy the same estimates than the bootstrap assumptions for the frame of ${}^{(int)}\mathcal{M}$.
- In ${}^{(ext)}\mathcal{M} \setminus Match$, estimates for the quantities associated to the global frame follow from the identities of Lemma 4.6.5 together with the bootstrap assumptions for the frame of ${}^{(ext)}\mathcal{M}$.
- In $Match$, the estimates for the quantities associated to the global frame are provided by Lemma 4.6.6.

The proof of Proposition 3.5.2 easily follows from Definition 4.6.2, Remark 4.6.3, and Lemma 4.6.6. Thus, from now on, we focus on the proof of Lemma 4.6.6 which is carried out in the next section.

4.6.2 Proof of Lemma 4.6.6

In this section, we prove Lemma 4.6.6. To ease the exposition, the quantities associated to the the frame of ${}^{(int)}\mathcal{M}$ are unprimed, the quantities associated to the frame of ${}^{(ext)}\mathcal{M}$ are primed, and the quantities associated to the the global frame are double-primed.

⁵We only need the first estimate for the proof of Proposition 3.5.2, but the second estimate will be needed in the proof of Theorem M8.

Step 1. Let (e_3, e_θ, e_4) denote the frame of $^{(int)}\mathcal{M}$ (and its extension) and (e'_3, e'_θ, e'_4) the frame of $^{(ext)}\mathcal{M}$ (and its extension). We denote by $(f, \underline{f}, \lambda)$ the reduced scalars such that

$$\begin{aligned} e'_4 &= \lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right), \\ e'_\theta &= \left(1 + \frac{1}{2} f \underline{f} \right) e_\theta + \frac{f}{2} e_4 + \frac{f}{2} \left(1 + \frac{f \underline{f}}{4} \right) e_3, \\ e'_3 &= \lambda^{-1} \left(\left(1 + \frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) e_3 + \left(\underline{f} + \frac{1}{4} \underline{f}^2 f \right) e_\theta + \frac{f^2}{4} e_4 \right). \end{aligned}$$

Together with the initialization of the frame of $^{(ext)}\mathcal{M}$ and $^{(int)}\mathcal{M}$ on \mathcal{T} in section 3.1.2 (where the spheres coincide), we have in particular

$$f = \underline{f} = 0, \quad \lambda = \Upsilon^{-1} \text{ on } \mathcal{T}. \quad (4.6.2)$$

Also, recall from section 3.5.1 that in order for (e'_3, e'_θ, e'_4) to be defined everywhere on $^{(int)}\mathcal{M} \cap \text{Match}$, we need - in addition to the above initialization of $(f, \underline{f}, \lambda)$ on \mathcal{T} , to initialize it also on $\underline{\mathcal{C}}_* \cap \text{Match}$ by

$$f = \underline{f} = 0, \quad \lambda = \Upsilon^{-1} \text{ on } \underline{\mathcal{C}}_* \cap \text{Match}. \quad (4.6.3)$$

Step 2. Next, we control the change of frame $(f, \underline{f}, \lambda)$ from (e_3, e_θ, e_4) to (e'_3, e'_θ, e'_4) in the region $^{(int)}\mathcal{M} \cap \text{Match}$. To this end, we rely on the transport equation of Lemma 2.3.6 together with the fact that $\omega' = \xi' = \zeta' + \underline{\eta}' = 0$. Then, $(\underline{f}, f, \log(\lambda))$ satisfy the following transport equations

$$\begin{aligned} \lambda^{-1} e'_4(f) + \left(\frac{\kappa}{2} + 2\omega \right) f &= -2\xi + E_1(f, \Gamma), \\ \lambda^{-1} e'_4(\log(\lambda)) &= 2\omega + E_2(f, \Gamma), \\ \lambda^{-1} e'_4(\underline{f}) + \frac{\kappa}{2} \underline{f} &= -2(\zeta + \underline{\eta}) + 2e'_\theta(\log(\lambda)) + 2f\underline{\omega} + E_3(f, \underline{f}, \Gamma), \end{aligned}$$

where E_1, E_2 and E_3 are given by

$$\begin{aligned} E_1(f, \Gamma) &= -\frac{1}{2} \vartheta f + \text{l.o.t.}, \\ E_2(f, \Gamma) &= f\zeta - \frac{1}{2} f^2 \underline{\omega} - \underline{\eta} f - \frac{1}{4} f^2 \underline{\kappa} + \text{l.o.t.}, \\ E_3(f, \underline{f}, \Gamma) &= -\underline{f} e'_\theta(f) - \frac{1}{2} \underline{f} \vartheta + \text{l.o.t.}, \end{aligned}$$

Here, l.o.t. denote terms which are cubic or higher order in f, \underline{f} (or in f only in the case of E_1 and E_2) and $\check{\Gamma}$ and do not contain derivatives of these quantities, where Γ and $\check{\Gamma}$

denotes the Ricci coefficients and renormalized Ricci coefficients w.r.t. the original null frame (e_3, e_4, e_θ) . We rewrite the transport equation for $\log(\lambda)$ as

$$\begin{aligned} & \lambda^{-1} e'_4(\log(\Upsilon\lambda)) \\ = & \lambda^{-1} e'_4(\log(\lambda)) + \lambda^{-1} e'_4(\log(\Upsilon)) \\ = & 2\omega + E_2(f, \Gamma) + \frac{1}{\Upsilon} \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right) \Upsilon \\ = & 2 \left(\omega + \frac{m}{r^2} \right) + E_2(f, \Gamma) + \frac{2}{\Upsilon} \frac{m(e_4(r) - \Upsilon)}{r^2} - \frac{2}{\Upsilon} \frac{e_4(m)}{r} - \frac{1}{\Upsilon} \left(f e_\theta + \frac{1}{4} f^2 e_3 \right) \Upsilon. \end{aligned}$$

In view of the above transport equations for f , \underline{f} and λ , the initialization (4.6.2) (4.6.3) for $(f, \underline{f}, \lambda)$ on $\mathcal{T} \cup (\underline{\mathcal{C}}_* \cap \text{Match})$, and the control of Γ induced by the bootstrap assumptions on ${}^{(int)}\mathcal{M}$, we easily deduce

$$\begin{aligned} & \max_{0 \leq k \leq k_{small}} \sup_{(int)\mathcal{M} \cap \text{Match}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k(f, \log(\Upsilon\lambda))| + \max_{0 \leq k \leq k_{small}-1} \sup_{(int)\mathcal{M} \cap \text{Match}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k \underline{f}| \lesssim \epsilon, \\ & \max_{0 \leq k \leq k_{large}} \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k(f, \log(\Upsilon\lambda))|^2 \right)^{\frac{1}{2}} + \max_{0 \leq k \leq k_{large}-1} \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k \underline{f}|^2 \right)^{\frac{1}{2}} \lesssim \epsilon. \end{aligned}$$

Step 3. We need to improve the number of derivatives in the top order estimate for $(f, \underline{f}, \log(\lambda))$. To this end, note first in view of the transformation formulas of Proposition 2.3.4 and the control of $(f, \underline{f}, \log(\lambda))$ provided by Step 2, we have in particular

$$\max_{0 \leq k \leq k_{large}-1} \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k \check{R}'|^2 \right)^{\frac{1}{2}} \lesssim \epsilon.$$

Relying on this estimate, the control of the Ricci coefficients associated to the outgoing null frame (e'_4, e'_3, e'_θ) on $\mathcal{T} \cup ({}^{(int)}\mathcal{M} \cap \text{Match})$, and the null structure equations, we infer

$$\max_{0 \leq k \leq k_{large}-1} \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k \check{\Gamma}'|^2 \right)^{\frac{1}{2}} \lesssim \epsilon.$$

We refer to section 8.9 for a completely analogous proof where the Ricci coefficients are recovered in ${}^{(int)}\mathcal{M}$ based on the control of the curvature components.

In view of the transformation formulas of Proposition 2.3.4, which can be written schematically as

$$\partial(f, \underline{f}, \log(\lambda)) = F(f, \underline{f}, \lambda, \check{\Gamma}),$$

the control of $(f, \underline{f}, \log(\lambda))$ provided by Step 2, and the above control of Γ' , we infer

$$\max_{0 \leq k \leq k_{large}} \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k(f, \underline{f}, \log(\Upsilon\lambda))|^2 \right)^{\frac{1}{2}} \lesssim \epsilon.$$

Step 4. We still need to control one more derivative of $(f, \underline{f}, \log(\lambda))$. Repeating the process of Step 3, we use again the transformation formulas of Proposition 2.3.4 and then the final estimate of Step 3 for $(f, \underline{f}, \log(\lambda))$ yields the following control for the curvature components

$$\max_{0 \leq k \leq k_{large}} \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k \check{R}'|^2 \right)^{\frac{1}{2}} \lesssim \epsilon.$$

Arguing as in Step 3, we infer⁶

$$\max_{0 \leq k \leq k_{large}} \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k \check{\Gamma}'|^2 \right)^{\frac{1}{2}} \lesssim \epsilon + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}.$$

Using again the transformation formulas of Proposition 2.3.4, this yields the following control for $(f, \underline{f}, \log(\lambda))$

$$\max_{0 \leq k \leq k_{large}+1} \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k(f, \underline{f}, \log(\Upsilon\lambda))|^2 \right)^{\frac{1}{2}} \lesssim \epsilon.$$

We have finally obtained for $(f, \underline{f}, \lambda)$ in $(int)\mathcal{M} \cap \text{Match}$

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-1} \sup_{(int)\mathcal{M} \cap \text{Match}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k(f, \underline{f}, \log(\Upsilon\lambda))| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{large}} \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k(f, \underline{f}, \log(\Upsilon\lambda))|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon, \\ \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^{k_{large}+1}(f, \underline{f}, \log(\Upsilon\lambda))|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Step 5. In addition to the estimate of $(f, \underline{f}, \lambda)$ in $(int)\mathcal{M} \cap \text{Match}$ of Step 4, we need to estimate $(f, \underline{f}, \lambda)$ in $(ext)\mathcal{M} \cap \text{Match}$. To this end, we first control in $(ext)\mathcal{M} \cap \text{Match}$ the

⁶In Step 3, there is no term corresponding to the one integrated on \mathcal{T} . This is due to the fact that for $k \leq k_{large} - 1$, we have thanks to the bootstrap assumptions on energy and a trace estimate

$$\max_{0 \leq k \leq k_{large}-1} \left(\int_{\mathcal{T}} |\mathfrak{d}^k({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}} \lesssim \epsilon.$$

reduced scalar $(f', \underline{f}', \lambda')$ satisfying

$$\begin{aligned} e_3 &= \lambda' \left(e'_3 + \underline{f}' e'_\theta + \frac{1}{4} \underline{f}'^2 e'_4 \right), \\ e_\theta &= \left(1 + \frac{1}{2} \underline{f}' \underline{f}' \right) e'_\theta + \frac{1}{2} \underline{f}' e'_3 + \frac{1}{2} \left(\underline{f}' + \frac{1}{4} \underline{f}' \underline{f}'^2 \right) e'_4, \\ e_4 &= \lambda'^{-1} \left(\left(1 + \frac{1}{2} \underline{f}' \underline{f}' + \frac{1}{16} \underline{f}'^2 \underline{f}'^2 \right) e'_4 + \left(\underline{f}' + \frac{1}{4} \underline{f}'^2 \underline{f}' \right) e'_\theta + \frac{1}{4} \underline{f}'^2 e'_3 \right). \end{aligned}$$

Together with the initialization of the frame of $^{(ext)}\mathcal{M}$ and $^{(int)}\mathcal{M}$ on \mathcal{T} in section 3.1.2 (where the spheres coincide), we have in particular

$$f' = \underline{f}' = 0, \quad \lambda' = \Upsilon^{-1} \text{ on } \mathcal{T}.$$

Also, recall from section 3.5.1 that in order for (e_3, e_θ, e_4) to be defined everywhere on $^{(ext)}\mathcal{M} \cap \text{Match}$, we need - in addition to the above initialization of $(f, \underline{f}, \lambda)$ on \mathcal{T} , to initialize it also on $\mathcal{C}_* \cap \text{Match}$ by

$$f' = \underline{f}' = 0, \quad \lambda' = \Upsilon^{-1} \text{ on } \mathcal{C}_* \cap \text{Match}. \quad (4.6.4)$$

Arguing similarly to Steps 1-4, we estimate $(f', \underline{f}', \lambda')$ and $(\check{\Gamma}, \check{R})$ in $^{(ext)}\mathcal{M} \cap \text{Match}$. We obtain

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-2} \sup_{^{(ext)}\mathcal{M} \cap \text{Match}} u^{1+\delta_{dec}} |\mathfrak{d}^k(\check{\Gamma}, \check{R})| + \max_{0 \leq k \leq k_{large}-1} \left(\int_{^{(ext)}\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k(\check{\Gamma}, \check{R})|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon, \\ \left(\int_{^{(ext)}\mathcal{M} \cap \text{Match}} |\mathfrak{d}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{small}-1} \sup_{^{(ext)}\mathcal{M} \cap \text{Match}} u^{1+\delta_{dec}} |\mathfrak{d}^k(f', \underline{f}', \log(\Upsilon' \lambda'))| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{large}} \left(\int_{^{(ext)}\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k(f', \underline{f}', \log(\Upsilon' \lambda'))|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon, \end{aligned}$$

and

$$\begin{aligned} \left(\int_{^{(ext)}\mathcal{M} \cap \text{Match}} |\mathfrak{d}^{k_{large}} \check{\Gamma}|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}(^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}, \\ \left(\int_{^{(ext)}\mathcal{M} \cap \text{Match}} |\mathfrak{d}^{k_{large}+1}(f', \underline{f}', \log(\Upsilon' \lambda'))|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}(^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Step 6. As mentioned above, in addition to the estimate of $(f, \underline{f}, \lambda)$ in $^{(int)}\mathcal{M} \cap \text{Match}$ of Step 4, we need to estimate $(f, \underline{f}, \lambda)$ in $^{(ext)}\mathcal{M} \cap \text{Match}$. To this end, we derive simple

algebraic relations between $(f, \underline{f}, \lambda)$ and $(f', \underline{f}', \lambda')$ of Step 5. On the one hand, we have from the definition of $(f, \underline{f}, \lambda)$

$$\begin{aligned} g(e'_4, e_3) &= -2\lambda, & g(e'_4, e_\theta) &= \lambda f, & g(e'_\theta, e_4) &= -f \left(1 + \frac{f\underline{f}}{4}\right), & g(e'_\theta, e_3) &= -\underline{f}, \\ g(e'_3, e_4) &= -2\lambda^{-1} \left(1 + \frac{f\underline{f}}{2} + \frac{1}{16} f^2 \underline{f}^2\right), & g(e'_3, e_\theta) &= \lambda^{-1} \underline{f} \left(1 + \frac{f\underline{f}}{4}\right). \end{aligned}$$

On the other hand, we have from the definition of $(f', \underline{f}', \lambda')$

$$\begin{aligned} g(e_3, e'_4) &= -2\lambda', & g(e_3, e'_\theta) &= \lambda' \underline{f}', & g(e_\theta, e'_4) &= -f', & g(e_\theta, e'_3) &= -\underline{f}' \left(1 + \frac{f' \underline{f}'}{4}\right), \\ g(e_4, e'_3) &= -2\lambda'^{-1} \left(1 + \frac{f' \underline{f}'}{2} + \frac{1}{16} f'^2 \underline{f}'^2\right), & g(e_4, e'_\theta) &= \lambda'^{-1} f' \left(1 + \frac{f' \underline{f}'}{4}\right). \end{aligned}$$

We immediately infer

$$\lambda' = \lambda, \quad f' = -\lambda f, \quad \underline{f}' = -\lambda^{-1} \underline{f}.$$

In view of the estimates of Step 5, we infer

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-1} \sup_{(ext)\mathcal{M} \cap \text{Match}} u^{1+\delta_{dec}} |\mathfrak{d}^k(f, \underline{f}, \log(\Upsilon\lambda))| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{large}} \left(\int_{(ext)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k(f, \underline{f}, \log(\Upsilon\lambda))|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon, \end{aligned}$$

and

$$\left(\int_{(ext)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^{k_{large}+1}(f, \underline{f}, \log(\Upsilon\lambda))|^2 \right)^{\frac{1}{2}} \lesssim \epsilon.$$

Together with Step 4, this yields

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-1} \sup_{(int)\mathcal{M} \cap \text{Match}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k(f, \underline{f}, \log(\Upsilon\lambda))| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{small}-1} \sup_{(ext)\mathcal{M} \cap \text{Match}} u^{1+\delta_{dec}} |\mathfrak{d}^k(f, \underline{f}, \log(\Upsilon\lambda))| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{large}} \left(\int_{\text{Match}} |\mathfrak{d}^k(f, \underline{f}, \log(\Upsilon\lambda))|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}((ext)\check{R})|^2 \right)^{\frac{1}{2}}, \end{aligned}$$

and

$$\max_{0 \leq k \leq k_{large}+1} \left(\int_{\text{Match}} |\mathfrak{d}^k(f, \underline{f}, \log(\Upsilon\lambda))|^2 \right)^{\frac{1}{2}} \lesssim \epsilon + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}((ext)\check{R})|^2 \right)^{\frac{1}{2}}.$$

Step 7. Next, we estimate $r' - r$ and $m' - m$. Note first the in view of the initialization of the foliations of $^{(ext)}\mathcal{M}$ and $^{(int)}\mathcal{M}$ on \mathcal{T} , as well as the initializations (4.6.3) on $\underline{\mathcal{C}}_* \cap \text{Match}$ and (4.6.4) on $\mathcal{C}_* \cap \text{Match}$, we have

$$r' = r, \quad m' = m \text{ on } \mathcal{T} \cup \text{Match}. \quad (4.6.5)$$

We start with the region $^{(int)}\mathcal{M} \cap \text{Match}$. We have

$$e'_4(r') = \frac{r'}{2\bar{\kappa}'} = 1 + \frac{r'}{2} \left(\bar{\kappa}' - \frac{2}{r'} \right), \quad e'_3(r') = \frac{r'}{2}(\bar{\kappa}' + \underline{A}') = -\Upsilon' + \frac{r'}{2} \left(\bar{\kappa}' + \frac{2\Upsilon'}{r'} \right) + \frac{r'}{2}\underline{A}',$$

which together with the identities for $e'_4(m')$ and $e'_3(m')$ in the outgoing foliation of $^{(ext)}\mathcal{M}$ and the control of the foliation of $^{(ext)}\mathcal{M}$ in $^{(int)}\mathcal{M} \cap \text{Match}$ established in Step 4 yields, using also $e'_\theta(r') = e'_\theta(m') = 0$,

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-2} \sup_{^{(int)}\mathcal{M} \cap \text{Match}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k(e'_4(r') - 1, e'_3(r') + \Upsilon', e'_\theta(r'), e'_4(m'), e'_3(m'), e'_\theta(m'))| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{large}-1} \left(\int_{^{(int)}\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k(e'_4(r') - 1, e'_3(r') + \Upsilon', e'_\theta(r'), e'_4(m'), e'_3(m'), e'_\theta(m'))|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon. \end{aligned}$$

On the other hand, we have in view of the decomposition of e'_4 , e'_3 and e'_θ of Step 1

$$\begin{aligned} e'_4(r) &= \lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right) r \\ &= \lambda \left(\frac{r}{2}(\bar{\kappa} + A) + \frac{1}{4} f^2 e_3(r) \right) \\ &= 1 + (\lambda\Upsilon - 1) + \lambda \left(\frac{r}{2} \left(\bar{\kappa} - \frac{2\Upsilon}{r} \right) + \frac{r}{2} A + \frac{1}{4} f^2 e_3(r) \right), \end{aligned}$$

$$\begin{aligned} e'_4(m) &= \lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right) m \\ &= \lambda \left(e_4(m) + \frac{1}{4} f^2 e_3(m) \right), \end{aligned}$$

$$\begin{aligned} e'_3(r) &= \lambda^{-1} \left(\left(1 + \frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) e_3 + \left(\underline{f} + \frac{1}{4} f^2 \underline{f} \right) e_\theta + \frac{f^2}{4} e_4 \right) r \\ &= \lambda^{-1} \left(e_3(r) + \left(\frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) e_3(r) + \frac{f^2}{4} e_4(r) \right) \\ &= -\Upsilon + \lambda^{-1}(\lambda\Upsilon - 1) + \lambda^{-1} \left(\frac{r}{2} \left(\bar{\kappa} + \frac{2}{r} \right) + \left(\frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) e_3(r) + \frac{f^2}{4} e_4(r) \right), \end{aligned}$$

$$\begin{aligned}
e'_3(m) &= \lambda^{-1} \left(\left(1 + \frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) e_3 + \left(\underline{f} + \frac{1}{4} \underline{f}^2 f \right) e_\theta + \frac{f^2}{4} e_4 \right) m \\
&= \lambda^{-1} \left(\left(1 + \frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) e_3(m) + \frac{f^2}{4} e_4(m) \right),
\end{aligned}$$

$$\begin{aligned}
e'_\theta(r) &= \left(\left(1 + \frac{1}{2} f \underline{f} \right) e_\theta + \frac{f}{2} e_4 + \frac{f}{2} \left(1 + \frac{f \underline{f}}{4} \right) e_3 \right) r \\
&= \frac{f}{2} e_4(r) + \frac{f}{2} \left(1 + \frac{f \underline{f}}{4} \right) e_3(r),
\end{aligned}$$

and

$$\begin{aligned}
e'_\theta(r) &= \left(\left(1 + \frac{1}{2} f \underline{f} \right) e_\theta + \frac{f}{2} e_4 + \frac{f}{2} \left(1 + \frac{f \underline{f}}{4} \right) e_3 \right) m \\
&= \frac{f}{2} e_4(m) + \frac{f}{2} \left(1 + \frac{f \underline{f}}{4} \right) e_3(m).
\end{aligned}$$

Together with the identities for $e_4(m)$ and $e_3(m)$ in the ingoing foliation of ${}^{(int)}\mathcal{M}$, the final estimates of Step 6 for f and λ , and the bootstrap assumptions for the foliation of ${}^{(int)}\mathcal{M}$, we infer

$$\max_{0 \leq k \leq k_{small}-2} \sup_{(int)\mathcal{M} \cap Match} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k(e'_4(r) - 1, e'_3(r) + \Upsilon, e'_\theta(r), e'_4(m), e'_3(m), e'_\theta(m))| \lesssim \epsilon,$$

$$\max_{0 \leq k \leq k_{large}-1} \left(\int_{(int)\mathcal{M} \cap Match} |\mathfrak{d}^k(e'_4(r) - 1, e'_3(r) + \Upsilon, e'_\theta(r), e'_4(m), e'_3(m), e'_\theta(m))|^2 \right)^{\frac{1}{2}} \lesssim \epsilon.$$

We deduce

$$\max_{0 \leq k \leq k_{small}-2} \sup_{(int)\mathcal{M} \cap Match} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k(e'_4(r' - r), e'_\theta(r - r'), \mathfrak{d}(m' - m))| \lesssim \epsilon,$$

$$\max_{0 \leq k \leq k_{large}-1} \left(\int_{(int)\mathcal{M} \cap Match} |\mathfrak{d}^k(e'_4(r' - r), e'_\theta(r - r'), \mathfrak{d}(m' - m))|^2 \right)^{\frac{1}{2}} \lesssim \epsilon.$$

In particular, we have

$$\sup_{(int)\mathcal{M} \cap Match} \underline{u}^{1+\delta_{dec}} |(e'_4(r' - r), e'_4(m' - m))| \lesssim \epsilon,$$

and together with the initialization (4.6.5), we integrate the transport equation from $\mathcal{T} \cup ({}^{(int)}\mathcal{M} \cap Match)$ and obtain

$$\sup_{(int)\mathcal{M} \cap Match} \underline{u}^{1+\delta_{dec}} |(r' - r, m' - m)| \lesssim \epsilon.$$

Together with the above estimates, and recovering the $e'_3(r' - r)$ using

$$e'_3(r' - r) = \left(e'_3(r') + \Upsilon' \right) - \left(e'_3(r) + \Upsilon \right) + 2 \left(\frac{m'}{r'} - \frac{m}{r} \right),$$

we infer

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-1} \sup_{(int)\mathcal{M} \cap \text{Match}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k(r' - r, m' - m)| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{large}} \left(\int_{(int)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k(r' - r, m' - m)|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon. \end{aligned}$$

Finally, arguing similarly in the region $(ext)\mathcal{M} \cap \text{Match}$, we infer

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-1} \sup_{(ext)\mathcal{M} \cap \text{Match}} u^{1+\delta_{dec}} |\mathfrak{d}^k(r' - r, m' - m)| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{large}} \left(\int_{(ext)\mathcal{M} \cap \text{Match}} |\mathfrak{d}^k(r' - r, m' - m)|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon, \end{aligned}$$

and hence

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-1} \sup_{(int)\mathcal{M} \cap \text{Match}} \underline{u}^{1+\delta_{dec}} |\mathfrak{d}^k(r' - r, m' - m)| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{small}-1} \sup_{(ext)\mathcal{M} \cap \text{Match}} u^{1+\delta_{dec}} |\mathfrak{d}^k(r' - r, m' - m)| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{large}} \left(\int_{\text{Match}} |\mathfrak{d}^k(r' - r, m' - m)|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon. \end{aligned}$$

Step 8. Recall from Definition 4.6.2 that we have defined the global null frame $(e''_4, e''_3, e''_\theta)$ as

- In $(int)\mathcal{M} \setminus \text{Match}$, $(e''_4, e''_3, e''_\theta) = (e_4, e_3, e_\theta)$.
- In $(ext)\mathcal{M} \setminus \text{Match}$, $(e''_4, e''_3, e''_\theta) = (\Upsilon e'_4, \Upsilon^{-1} e'_3, e'_\theta)$.
- In Match , $(e''_4, e''_3, e''_\theta)$ is given by the change of frame formula starting from (e_4, e_3, e_θ) and with change of frame coefficients $(f'', \underline{f}'', \lambda'')$ given by

$$f'' = \psi f, \quad \underline{f}'' = \psi \underline{f}, \quad \lambda'' = 1 - \psi + \psi \Upsilon' \lambda,$$

see (4.6.1).

Also, recall that we have defined r'' and m'' as

$$r'' = (1 - \psi)r + \psi r', \quad m'' = (1 - \psi)m + \psi m'.$$

Step 9. In view of the transformation formulas of Proposition 2.3.4, we have schematically

$$(\check{\Gamma}'', \check{R}'') = (\check{\Gamma}, \check{R}) + \mathfrak{d}(f'', \underline{f}'', \lambda'' - 1) + f'' + \underline{f}'' + (\lambda'' - 1) + (r'' - r) + (m'' - m).$$

In view of the definition of $(f'', \underline{f}'', \lambda'')$ and (r'', m'') in Step 8, we infer

$$(\check{\Gamma}'', \check{R}'') = (\check{\Gamma}, \check{R}) + \mathfrak{d}(f, \underline{f}, \Upsilon\lambda - 1) + f + \underline{f} + (\Upsilon\lambda - 1) + (r' - r) + (m' - m).$$

Together with the bootstrap assumptions in ${}^{(int)}\mathcal{M}$ for (Γ, \check{R}) , the estimates for (Γ, \check{R}) in ${}^{(ext)}\mathcal{M}$ provided by Step 5, the estimates for $(f, \underline{f}, \lambda)$ provided by Step 6 in Match, and the estimates for $r' - r$ and $m' - m$ provided by Step 7, we deduce

$$\begin{aligned} \max_{0 \leq k \leq k_{small} - 2} \sup_{(int)\mathcal{M} \cap \text{Match}} \underline{u}^{1 + \delta_{dec}} |\mathfrak{d}^k(\check{\Gamma}'', \check{R}'')| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{small} - 2} \sup_{(ext)\mathcal{M} \cap \text{Match}} \underline{u}^{1 + \delta_{dec}} |\mathfrak{d}^k(\check{\Gamma}'', \check{R}'')| &\lesssim \epsilon, \\ \max_{0 \leq k \leq k_{large} - 1} \left(\int_{\text{Match}} |\mathfrak{d}^k(\check{\Gamma}'', \check{R}'')|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon, \\ \left(\int_{\text{Match}} |\mathfrak{d}^{k_{large}}(\check{\Gamma}'', \check{R}'')|^2 \right)^{\frac{1}{2}} &\lesssim \epsilon + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Since the double-primed quantities correspond to the quantities associated to the the global frame, this concludes the proof of Lemma 4.6.6.

4.6.3 Proof of Proposition 3.5.5

To match the first global frame of \mathcal{M} of Proposition 3.5.5 with a conformal renormalization of the second frame of ${}^{(ext)}\mathcal{M}$ of Proposition 3.4.6, we will need to introduce a cut-off function.

Definition 4.6.8. Let $\psi : \mathbb{R} \rightarrow \mathbb{R}$ a smooth cut-off function such that $0 \leq \psi \leq 1$, $\psi = 0$ on $(-\infty, 0]$ and $\psi = 1$ on $[1, +\infty)$. We define ψ_{m_0} as follows

$$\psi_{m_0}(r) = \begin{cases} 1 & \text{if } r \geq 4m_0, \\ 0 & \text{if } r \leq \frac{7m_0}{2}, \end{cases}$$

and

$$\psi_{m_0}(r) = \psi \left(\frac{2 \left(r - \frac{7m_0}{2} \right)}{m_0} \right) \text{ on } \frac{7m_0}{2} \leq {}^{(ext)}r \leq 4m_0.$$

We are now ready to define the second global frame, i.e. the global frame of the statement of Proposition 3.5.5.

Definition 4.6.9 (Definition of the second global frame). *We introduce a global null frame defined on ${}^{(ext)}\mathcal{M} \cup {}^{(int)}\mathcal{M}$ and denoted by $({}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta)$. The second global frame is defined as follows*

1. In ${}^{(ext)}\mathcal{M} \cap \{{}^{(ext)}r \geq 4m_0\}$, we have

$$({}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta) = ({}^{(ext)}\Upsilon e'_4, {}^{(ext)}\Upsilon^{-1}e'_3, e'_\theta),$$

where (e'_4, e'_3, e'_θ) denotes the second frame of ${}^{(ext)}\mathcal{M}$, i.e. the one constructed in of Proposition 3.4.6.

2. In ${}^{(int)}\mathcal{M} \cup ({}^{(ext)}\mathcal{M} \cap \{{}^{(ext)}r \leq \frac{7m_0}{2}\})$, we have

$$({}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta) = ({}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta),$$

where $({}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta)$ denotes the first global frame of \mathcal{M} of Proposition 3.5.5.

3. It remains to define the global frame on the matching region $Match'$. We denote by f the reduced scalar introduced in Proposition 3.4.6 such that we have in ${}^{(ext)}\mathcal{M}$

$$\begin{aligned} e'_4 &= {}^{(ext)}e_4 + f {}^{(ext)}e_\theta + \frac{1}{4} f^2 {}^{(ext)}e_3, \\ e'_\theta &= {}^{(ext)}e_\theta + \frac{f}{2} {}^{(ext)}e_3, \\ e'_3 &= {}^{(ext)}e_3. \end{aligned}$$

Then, in the matching region $Match'$, the second global frame of \mathcal{M} is given by

$$\begin{aligned} ({}^{(glo')}e_4) &= \Upsilon' \left(\Upsilon'^{-1} ({}^{(glo)}e_4 + f' ({}^{(glo)}e_\theta) + \frac{1}{4} f'^2 \Upsilon' ({}^{(glo)}e_3) \right), \\ ({}^{(glo')}e_\theta) &= ({}^{(glo)}e_\theta) + \frac{f'}{2} \Upsilon' ({}^{(glo)}e_3), \\ ({}^{(glo')}e_3) &= ({}^{(glo)}e_3), \end{aligned}$$

where

$$f' = \psi_{m_0} ({}^{(ext)}r) f, \quad \Upsilon' = 1 - \psi_{m_0} ({}^{(ext)}r) + \psi_{m_0} ({}^{(ext)}r) ({}^{(ext)}\Upsilon). \quad (4.6.6)$$

Remark 4.6.10. Recall that the smooth cut-off function ψ in Definition 3.5.4, allowing to define ψ_{m_0, δ_H} , is such that we have in particular $\psi = 0$ on $(-\infty, 0]$ and $\psi = 1$ on $[1, +\infty)$. The following two special cases correspond to the properties (d) i. and (d) ii. of Proposition 3.5.5.

- If the cut-off ψ in Definition 3.5.4 is such that $\psi = 1$ on $[1/2, +\infty)$, then

$${}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta = ({}^{(ext)}\Upsilon e'_4, {}^{(ext)}\Upsilon^{-1}e'_3, e'_\theta) \text{ on } {}^{(ext)}\mathcal{M} \left({}^{(ext)}r \geq \frac{15m_0}{4} \right).$$

- If the cut-off ψ in Definition 3.5.4 is such that $\psi = 0$ on $(-\infty, 1/2]$, then

$${}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta = ({}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta) \text{ on } ({}^{(int)}\mathcal{M} \cup {}^{(ext)}\mathcal{M}) \left({}^{(ext)}r \leq \frac{15m_0}{4} \right).$$

Remark 4.6.11. When dealing with the second global frame $({}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta)$, the area radius and Hawking mass that we use are the ones corresponding to the first global frame, i.e. ${}^{(glo)}r$ and ${}^{(glo)}m$.

The following two lemmas provide the main properties of the second global frame of \mathcal{M} .

Lemma 4.6.12. We have in ${}^{(ext)}\mathcal{M}(r \geq 4m_0)$ the following relations between the quantities in the second global frame of \mathcal{M} , i.e. $({}^{(glo')}e_4, {}^{(glo')}e_3, {}^{(glo')}e_\theta)$, and the second frame of ${}^{(ext)}\mathcal{M}$, i.e. (e'_4, e'_3, e'_θ) ,

$$\begin{aligned} {}^{(glo')}\alpha &= \Upsilon^2 \alpha', & {}^{(glo')}\beta &= \Upsilon \beta', & {}^{(glo')}\rho + \frac{2m}{r^3} &= \rho' + \frac{2m}{r^3}, & {}^{(glo')}\underline{\beta} &= \Upsilon^{-1} \underline{\beta}', \\ {}^{(glo')}\underline{\alpha} &= \Upsilon^{-2} \underline{\alpha}', & {}^{(glo')}\xi &= 0, & {}^{(glo')}\underline{\xi} &= \Upsilon^{-2} \underline{\xi}', & {}^{(glo')}\zeta &= -{}^{(glo')}\underline{\eta} = \zeta', \\ {}^{(glo')}\eta &= \eta', & {}^{(glo')}\omega + \frac{m}{r^2} &= \Upsilon \omega' + \frac{m}{r^2} (1 - e'_4(r)) + \frac{e'_4(m)}{r}, \\ {}^{(glo')}\underline{\omega} &= \Upsilon^{-1} \left(\underline{\omega}' - \frac{m}{r^2} + \frac{m}{r^2} \left(1 - \frac{e'_3(r)}{\Upsilon} \right) - \frac{e'_3(m)}{\Upsilon r} \right), & {}^{(glo')}\kappa - \frac{2\Upsilon}{r} &= \Upsilon \left(\kappa' - \frac{2}{r} \right), \\ {}^{(glo')}\underline{\kappa} + \frac{2}{r} &= \Upsilon^{-1} \left(\underline{\kappa}' + \frac{2\Upsilon}{r} \right), & {}^{(glo')}\vartheta &= \Upsilon \vartheta', & {}^{(glo')}\underline{\vartheta} &= \Upsilon^{-1} \underline{\vartheta}'. \end{aligned}$$

Proof. The proof follows immediately from the change of frame formula with the choice $(f = 0, \underline{f} = 0, \lambda = \Upsilon)$, the fact that $e_\theta(\Upsilon) = 0$, and the fact that the frame (e'_4, e'_3, e'_θ) is such that $\xi' = 0$ and $\underline{\eta}' = -\zeta'$. \square

Lemma 4.6.13 (Control of the second global frame in the matching region). *In the matching region, the following estimates holds for the second global frame*

$$\max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{Match'} u^{1+\delta_{dec}-2\delta_0} \left| \mathfrak{D}^k ({}^{(glo')}\check{\Gamma}, {}^{(glo')}\check{R}) \right| \lesssim \epsilon.$$

Remark 4.6.14. The quantities associated to the second global frame can be estimated as follows

- In $(^{int})\mathcal{M} \cup (^{ext})\mathcal{M}({}^{(ext)}r \leq \frac{7m_0}{2})$, the second global frame coincides with the first global frame, and hence, the quantities associated to the second global frame satisfy the same estimates than the corresponding quantities for the first global frame.
- In $(^{ext})\mathcal{M}({}^{(ext)}r \geq 4m_0)$, estimates for the quantities associated to the second global frame follow from the identities of Lemma 4.6.12 together with the estimates of Proposition 3.4.6 for the second frame of $(^{ext})\mathcal{M}$.
- In $Match'$, the estimates for the quantities associated to the global frame are provided by Lemma 4.6.13.

The proof of Proposition 3.5.5 easily follows from Definition 4.6.9, Remark 4.6.10, and Lemma 4.6.13. Thus, from now on, we focus on the proof of Lemma 4.6.13 which is carried out below.

Proof of Lemma 4.6.13. Recall from definition 4.6.9 that we have in the matching region $Match'$

$$\begin{aligned} (^{glo'})e_4 &= \Upsilon' \left(\Upsilon'^{-1} (^{glo})e_4 + f' (^{glo})e_\theta + \frac{1}{4} f'^2 \Upsilon' (^{glo})e_3 \right), \\ (^{glo'})e_\theta &= (^{glo})e_\theta + \frac{f'}{2} \Upsilon' (^{glo})e_3, \\ (^{glo'})e_3 &= (^{glo})e_3, \end{aligned}$$

where

$$f' = \psi_{m_0}({}^{(ext)}r)f, \quad \Upsilon' = 1 - \psi_{m_0}({}^{(ext)}r) + \psi_{m_0}({}^{(ext)}r)({}^{(ext)}\Upsilon).$$

Now, since $(^{ext)}r \geq \frac{7m_0}{2}$ on $Match'$, we also have in that region

$$(^{glo})e_4, (^{glo})e_3, (^{glo})e_\theta = ({}^{(ext)}\Upsilon ({}^{(ext)}e_4, ({}^{(ext)}\Upsilon)^{-1} ({}^{(ext)}e_3, ({}^{(ext)}e_\theta)).$$

We deduce on $Match'$

$$\begin{aligned} (^{glo'})e_4 &= ({}^{(ext)}\Upsilon \left(({}^{(ext)}e_4 + f'' ({}^{(ext)}e_\theta + \frac{1}{4} f''^2 ({}^{(ext)}e_3) \right), \\ (^{glo'})e_\theta &= ({}^{(ext)}e_\theta + \frac{f''}{2} ({}^{(ext)}e_3, \\ (^{glo'})e_3 &= ({}^{(ext)}\Upsilon)^{-1} ({}^{(ext)}e_3, \end{aligned}$$

where

$$\begin{aligned} f'' &= \Upsilon' ({}^{(ext)}\Upsilon)^{-1} f' \\ &= \left(1 - \psi_{m_0}({}^{(ext)}r) + \psi_{m_0}({}^{(ext)}r)({}^{(ext)}\Upsilon) \right) ({}^{(ext)}\Upsilon)^{-1} \psi_{m_0}({}^{(ext)}r) f. \end{aligned}$$

In view of the transformation formulas of Proposition 2.3.4, we deduce, schematically,

$$\left((gl'o')\check{\Gamma}, (gl'o')\check{R} \right) = \left((ext)\check{\Gamma}, (ext)\check{R} \right) + \mathfrak{d}f + f.$$

Together with the bootstrap assumptions on decay and Proposition 3.4.5 for $((ext)\check{\Gamma}, (ext)\check{R})$, and the estimate (3.4.11) for f , we infer

$$\max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{\text{Match}'} u^{1+\delta_{dec}-2\delta_0} \left| \mathfrak{d}^k ((gl'o')\check{\Gamma}, (gl'o')\check{R}) \right| \lesssim \epsilon$$

which concludes the proof of Lemma 4.6.13. \square

Chapter 5

DECAY ESTIMATES FOR \mathfrak{q} (Theorem M1)

The goal of the chapter is to prove Theorem M1, i.e. to derive decay estimates for the quantity \mathfrak{q} for $k \leq k_{small} + 20$ derivatives. To this end, we will make use of the wave equation satisfied by \mathfrak{q} (see (2.4.7))

$$\square_2 \mathfrak{q} + \kappa \underline{\kappa} \mathfrak{q} = N, \quad (5.0.1)$$

where N contains only quadratic or higher order terms. Now, in order to have a suitable right-hand side N , recall from the discussion in Remarks 2.4.8 and 2.4.9 that \mathfrak{q} is defined relative to the global null frame of Proposition 3.5.5 for which $\xi = 0$ for $r \geq 4m_0$ and $\eta \in \Gamma_g$. For such a global frame, N is given schematically by, see (2.4.8),

$$N = r^2 \mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta)) + e_3 \left(r^3 \mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta)) \right) + \mathfrak{d}^{\leq 1}(\Gamma_g \cdot \mathfrak{q}) + \text{l.o.t.} \quad (5.0.2)$$

5.1 Preliminaries

Smallness constants

Recall from the beginning of section 3.3.2 the constant m_0 and the main small constants $\delta_{\mathcal{H}}$, δ_B , δ_{dec} , ϵ and ϵ_0 such that

- The constant $m_0 > 0$ is the mass of the initial Schwarzschild spacetime relative to which our initial perturbation is measured.

- The integer k_{large} which corresponds to the maximum number of derivatives of the solution.
- The size of the initial data layer norm is measured by $\epsilon_0 > 0$.
- The size of the bootstrap assumption norms are measured by $\epsilon > 0$.
- $\delta_{\mathcal{H}} > 0$ measures the width of the region $|r - 2m_0| \leq 2m_0\delta_{\mathcal{H}}$ where the redshift estimate holds and which includes in particular the region $^{(int)}\mathcal{M}$.
- δ_{dec} is tied to decay estimates in u, \underline{u} for $\check{\Gamma}$ and \check{R} .
- δ_B is involved in the r -power of the r^p weighted estimates for curvature.

Recall also that these constants satisfy in view of (3.3.1) (3.3.2) (3.3.3)

$$0 < \delta_{\mathcal{H}}, \delta_{dec}, \delta_B \ll \min\{m_0, 1\}, \quad \delta_B > 2\delta_{dec}, \quad k_{large} \gg \frac{1}{\delta_{dec}},$$

$$\epsilon_0, \epsilon \ll \min\{\delta_{\mathcal{H}}, \delta_{dec}, \delta_B, m_0, 1\},$$

and

$$\epsilon = \epsilon_0^{\frac{2}{3}}.$$

We will need the following additional small constants in this chapter

- $\delta_{extra} > 0$, tied to the decay of \mathbf{q} , and is chosen such that $\delta_{extra} > \delta_{dec}$,
- $\delta > 0$ for various degeneracies,
- $\delta_0 > 0$ which comes from interpolating between $k \leq k_{small}$ derivatives of $(\check{\Gamma}, \check{R})$ and $k \leq k_{large}$ derivatives of $(\check{\Gamma}, \check{R})$, see Lemma 5.1.1,
- $q_0 > 0$ which will allow us to recover the fact that the decay for \mathbf{q} in Theorem M1 has an extra gain $u^{-(\delta_{extra}-\delta_{dec})}$ compared from the expected behavior inferred from the bootstrap assumptions.

We will choose δ_{extra} such that

$$\delta_{dec} < \delta_{extra} < 2\delta_{dec}, \quad \delta_B \geq 2\delta_{extra},$$

δ and δ_0 such that

$$0 < \epsilon, \epsilon_0 \ll \delta, \delta_0 \ll \delta_{dec}, \delta_{extra}, \delta_{\mathcal{H}}, m_0, 1, \quad (5.1.1)$$

and q_0 such that¹

$$2\delta_{dec} < q_0 < 4\delta_{dec} - 4\delta_0 - 4\delta. \quad (5.1.2)$$

5.1.1 The foliation of \mathcal{M} by τ

Recall that the spacetime \mathcal{M} is decomposed as $\mathcal{M} = {}^{(int)}\mathcal{M} \cup {}^{(ext)}\mathcal{M}$ and that u is an outgoing optical function on ${}^{(ext)}\mathcal{M}$ while \underline{u} is an ingoing optical function. In this chapter, we rely on the global frame $(e_3, e_4, e_\theta, e_\varphi)$ defined in section 3.5, and r and m denote the corresponding scalar functions associated to it. Also, we define the trapping region \mathcal{M}_{trap} as,

$$\mathcal{M}_{trap} := \left\{ \frac{5m_0}{2} \leq r \leq \frac{7m_0}{2} \right\}. \quad (5.1.3)$$

Also, let ${}^{(trap)}\mathcal{M} = \mathcal{M} \setminus \mathcal{M}_{trap}$ the complement of \mathcal{M}_{trap} in \mathcal{M} .

We foliate our spacetime domain \mathcal{M} by \mathbf{Z} invariant hypersurfaces $\Sigma(\tau)$ which are:

- Incoming null in ${}^{(int)}\mathcal{M}$, with e_3 as null incoming generator. We denote this portion ${}^{(int)}\Sigma(\tau)$.
- Strictly spacelike in ${}^{(trap)}\mathcal{M}$. We denote this portion by ${}^{(trap)}\Sigma$.
- Outgoing null in $\mathcal{M}_{>4m_0}$. We denote this portion by $\Sigma_{>4m_0}(\tau)$.
- The parameter τ of $\Sigma(\tau)$ can be chosen, smoothly, such that

$$\tau := \begin{cases} u & \text{in } \mathcal{M}_{>4m_0}, \\ u + r & \text{in } \mathcal{M}_{trap}, \\ \underline{u} & \text{in } {}^{(int)}\mathcal{M}. \end{cases} \quad (5.1.4)$$

¹This will allow us to choose in the proof of Theorem M1, see (5.2.10),

$$\delta_{extra} = \frac{q_0 - \delta}{2}$$

which satisfies the desired estimate $\delta_{extra} > \delta_{dec}$ for $\delta > 0$ small enough.

- In particular, the unit normal in the region \mathcal{M}_{trap} , i.e. the normal to $^{(trap)}\Sigma$, satisfies²

$$-2 \leq g(N_\Sigma, e_4) \leq -1, \quad -2 \leq g(N_\Sigma, e_3) \leq -1 \text{ on } \mathcal{M}_{trap}. \quad (5.1.5)$$

5.1.2 Assumptions for Ricci coefficients and curvature

Recall from Remark 2.4.9 that \mathfrak{q} is defined, according to equation (2.3.10) in Lemma 2.3.10, relative to the global frame of Proposition 3.5.5 for which $\eta \in \Gamma_g$ with the notation

$$\begin{aligned} \Gamma_g &= \Gamma_g^{(0)} = \left\{ \xi, \vartheta, \omega + \frac{m}{r^2}, \kappa - \frac{2\Upsilon}{r}, \eta, \underline{\eta}, \zeta, A \right\}, \\ \Gamma_b &= \Gamma_b^{(0)} = \left\{ \underline{\vartheta}, \underline{\kappa} + \frac{2}{r}, \underline{A}, \underline{\omega}, \underline{\xi} \right\}, \end{aligned}$$

where we recall that

$$\Upsilon = 1 - \frac{2m}{r}, \quad A = \frac{2}{r}e_4(r) - \kappa, \quad \underline{A} = \frac{2}{r}e_3(r) - \underline{\kappa}.$$

Note also that ξ vanishes in $^{(ext)}\mathcal{M}$ away from the matching region of Proposition 3.5.5, and in particular for $r \geq 4m_0$.

For higher derivatives we write,

$$\begin{aligned} \Gamma_g^{(1)} &= \left\{ \mathfrak{d}\xi, \mathfrak{d}\vartheta, re_\theta\omega, re_\theta(\kappa), \mathfrak{d}\eta, \mathfrak{d}\underline{\eta}, \mathfrak{d}\zeta, \mathfrak{d}A \right\} \\ \Gamma_b^{(1)} &= \left\{ \mathfrak{d}\underline{\vartheta}, re_\theta(\underline{\kappa}), \mathfrak{d}\underline{\xi}, \mathfrak{d}\underline{A}, re_\theta\underline{\omega}, \mathfrak{d}\underline{\xi} \right\} \end{aligned}$$

and for $s \geq 2$,

$$\Gamma_g^{(s)} = \mathfrak{d}^{s-1}\Gamma_g^{(1)}, \quad \Gamma_b^{(s)} = \mathfrak{d}^{s-1}\Gamma_b^{(1)}$$

Moreover we denote

$$\Gamma_g^{\leq s} = \left\{ \Gamma_g^{(0)}, \Gamma_g^{(1)}, \dots, \Gamma_g^{(s)} \right\}, \quad \Gamma_b^{\leq s} = \left\{ \Gamma_b^{(0)}, \Gamma_b^{(1)}, \dots, \Gamma_b^{(s)} \right\}.$$

² N_Σ is given in view of its definition by

$$\begin{aligned} N_\Sigma &= \frac{1}{\sqrt{2}\sqrt{e_4(r)(e_3(u) + e_3(r))}} \left(e_4(r)e_3 + (e_3(u) + e_3(r))e_4 \right) \\ &= \frac{1}{\sqrt{2}\sqrt{2 - \Upsilon + O(\epsilon)}} \left((1 + O(\epsilon))e_3 + (2 - \Upsilon + O(\epsilon))e_4 \right) \end{aligned}$$

where we used the bootstrap assumptions.

With these notations, we may now state the estimates satisfied by the Ricci coefficients and curvature components.

Lemma 5.1.1. *Consider the global frame of Proposition 3.5.5 and the above definition³ of Γ_g and Γ_b . Let an integer k_{loss} and a small constant $\delta_0 > 0$ satisfying⁴*

$$16 \leq k_{loss} \leq \frac{\delta_{dec}}{3}(k_{large} - k_{small}), \quad \delta_0 = \frac{k_{loss}}{k_{large} - k_{small}}. \quad (5.1.6)$$

Then, the Ricci coefficients and curvature components with respect to the global frame of Proposition 3.5.5 satisfy

$$\begin{aligned} & \xi = 0 \text{ on } r \geq 4m_0, \\ & \max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{\mathcal{M}} \left\{ \left(r^2 \tau^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r \tau^{1 + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^k \Gamma_g| + r \tau^{1 + \delta_{dec} - 2\delta_0} |\mathfrak{d}^k \Gamma_b| \right. \\ & \quad + \left(r^{\frac{7}{2} + \frac{\delta_B}{2}} + r^3 \tau^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r^2 \tau^{1 + \delta_{dec} - 2\delta_0} \right) \left(|\mathfrak{d}^k \alpha| + |\mathfrak{d}^k \beta| \right) \\ & \quad + \left(r^3 \tau^{\frac{1}{2} + \delta_{dec} - 2\delta_0} + r^2 \tau^{1 + \delta_{dec} - 2\delta_0} \right) |\mathfrak{d}^k \check{\rho}| \\ & \quad \left. + \tau^{1 + \delta_{dec} - 2\delta_0} \left(r^2 |\mathfrak{d}^k \underline{\beta}| + r |\mathfrak{d}^k \underline{\alpha}| \right) \right\} \lesssim \epsilon, \\ & \max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{\mathcal{M}} \left\{ r^2 \tau^{1 + \delta_{dec} - 2\delta_0} |\mathfrak{d}^{k-1} e_3(\Gamma_g)| \right. \\ & \quad \left. + r^3 (\tau + 2r)^{1 + \delta_{dec} - 2\delta_0} \left(|\mathfrak{d}^{k-1} e_3(\alpha)| + |\mathfrak{d}^k e_3(\beta)| \right) \right\} \lesssim \epsilon. \end{aligned}$$

Proof. In $r \geq 4m_0$, the global frame of Proposition 3.5.5 coincides with a conformal renormalization of the second frame of $^{(ext)}\mathcal{M}$, see Proposition 3.4.6. The estimates there follow immediately from the ones of Proposition 3.4.6. In the matching region $7/2m_0 \leq r \leq 4m_0$, the estimates are stated in Proposition 3.5.5. Finally, for $^{(ext)}\mathcal{M}(r \leq 7/2m_0)$ and $^{(int)}\mathcal{M}$, the estimates follow directly from interpolation between the bootstrap assumptions on decay for $k \leq k_{small}$ and the pointwise estimates of Proposition 3.4.5 for $k \leq k_{large} - 5$. \square

³Recall in particular that the global frame of Proposition 3.5.5 is such that $\eta \in \Gamma_g$.

⁴Recall that we have

$$0 < \delta_{dec} \ll 1, \quad \delta_{dec} k_{large} \gg 1, \quad k_{small} = \left\lfloor \frac{1}{2} k_{large} \right\rfloor + 1.$$

In particular, we have $\delta_{dec}(k_{large} - k_{small}) \gg 1$ and hence there exists an integer k_{loss} satisfying the required constraints. We will in fact choose $k_{loss} = 33$, see (5.2.3).

5.1.3 Structure of nonlinear terms

The following lemma will be important in what follows.

Lemma 5.1.2. *For the solution \mathbf{q} to the wave equation (5.0.1), the structure of the error term N can be written schematically as follows*

$$N = N_g + e_3(rN_g) + N_m[\mathbf{q}] \quad (5.1.7)$$

where,

$$\begin{aligned} N_g &= r^2 \mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta)), \\ N_m[\mathbf{q}] &= \mathfrak{d}^{\leq 1}(\Gamma_g \cdot \mathbf{q}). \end{aligned} \quad (5.1.8)$$

Moreover, for every $k \leq k_{large} - 3$ we have schematically,

$$\mathfrak{d}^k N = \mathfrak{d}^{\leq k} N_g + e_3(\mathfrak{d}^k(rN_g)) + \mathfrak{d}^k N_m[\mathbf{q}]. \quad (5.1.9)$$

Remark 5.1.3. *In fact, (5.1.7) and (5.1.9) also contain lower order terms which are strictly better in powers of r and contain at most the same number of derivatives. For convenience, we drop them in the rest of the proof of Theorem M1.*

Proof. For $k = 0$, this is an immediate consequence of (5.0.2). For the higher derivatives we write,

$$\mathfrak{d}^k(e_3(rN_g)) = e_3(\mathfrak{d}^k(rN_g)) + [\mathfrak{d}^k, e_3](rN_g).$$

In view of the formula for $[e_3, \mathfrak{d}]$ of Lemma 2.2.13, and the commutator formula for $[e_3, e_4]$, we have, schematically,

$$[e_3, e_3] = 0, \quad [\mathfrak{d}, e_3] = \Gamma_b \mathfrak{d} + \Gamma_b, \quad [re_4, e_3] = \left(\frac{1}{r} + \Gamma_b\right) \mathfrak{d}.$$

In view of our assumptions.

$$|\mathfrak{d}^i(\Gamma_b)| \leq r^{-1}\epsilon, \quad i \leq k_{large} - 4,$$

Γ_b is at least as good as r^{-1} , and hence, we deduce, schematically,

$$[\mathfrak{d}, e_3] = \frac{1}{r} \mathfrak{d} + \frac{1}{r}.$$

On the other hand, we have, schematically,

$$[\mathfrak{d}, r] = r$$

and hence, for $k \leq k_{large} - 3$,

$$\begin{aligned} [\mathfrak{d}^k, e_3](rN_g) &= \sum_{i+j \leq k-1} \mathfrak{d}^i \left(\frac{1}{r} \mathfrak{d} + \frac{1}{r} \right) \mathfrak{d}^j(rN_g) \\ &= \mathfrak{d}^{\leq k} N_g \end{aligned}$$

as desired. \square

5.1.4 Main quantities

We restrict our attention to the region $\mathcal{M}(\tau_1, \tau_2) = \mathcal{M} \cap \{\tau_1 \leq \tau \leq \tau_2\}$. For a given $\psi \in \mathfrak{s}_2(\mathcal{M})$ we introduce the following quantities, for $0 \leq \tau_1 < \tau_2 \leq \tau_*$.

Morawetz bulk quantities

Consider the vectorfields,

$$T := \frac{1}{2}(e_4 + \Upsilon e_3), \quad R := \frac{1}{2}(e_4 - \Upsilon e_3). \quad (5.1.10)$$

Let θ a smooth bump function equal 1 on $|\Upsilon| \leq \delta_{\mathcal{H}}^{\frac{1}{10}}$ vanishing for $|\Upsilon| \geq 2\delta_{\mathcal{H}}^{\frac{1}{10}}$ and define the modified vectorfields,

$$\begin{aligned} \check{R} &:= \theta \frac{1}{2}(e_4 - e_3) + (1 - \theta)\Upsilon^{-1}R = \frac{1}{2} \left[\check{\theta} e_4 - e_3 \right], \\ \check{T} &:= \theta \frac{1}{2}(e_4 + e_3) + (1 - \theta)\Upsilon^{-1}T = \frac{1}{2} \left[\check{\theta} e_4 + e_3 \right], \end{aligned} \quad (5.1.11)$$

where $\check{\theta} = \theta + \Upsilon^{-1}(1 - \theta)$. Note that,

$$\check{\theta} = \begin{cases} 1 & \text{for } |\Upsilon| \leq \delta_{\mathcal{H}}^{\frac{1}{10}}, \\ \Upsilon^{-1} & \text{for } |\Upsilon| \geq 2\delta_{\mathcal{H}}^{\frac{1}{10}}. \end{cases} \quad (5.1.12)$$

Remark 5.1.4. *Note that*

$$\check{R} + \check{T} = e_4, \quad -\check{R} + \check{T} = e_3 \text{ in } {}^{(int)}\mathcal{M} \text{ and } \check{R} + \check{T} = \Upsilon^{-1}e_4, \quad -\check{R} + \check{T} = e_3 \text{ in } \mathcal{M}_{>4m_0}.$$

We define the quantities

$$\text{Mor}[\psi](\tau_1, \tau_2) := \int_{\mathcal{M}(\tau_1, \tau_2)} \frac{1}{r^3} |\check{R}\psi|^2 + \frac{1}{r^4} |\psi|^2 + \left(1 - \frac{3m}{r}\right)^2 \frac{1}{r} \left(|\nabla\psi|^2 + \frac{1}{r^2} |\check{T}\psi|^2 \right), \quad (5.1.13)$$

$$\text{Morr}[\psi](\tau_1, \tau_2) := \text{Mor}[\psi](\tau_1, \tau_2) + \int_{\mathcal{M}_{>4m_0}(\tau_1, \tau_2)} r^{-1-\delta} |e_3(\psi)|^2,$$

with $m = m(\tau, r) = m(u, r)$ the Hawking mass in \mathcal{M} . The constant $\delta > 0$ is a sufficiently small quantity. An equivalent definition for $\text{Morr}[\psi](\tau_1, \tau_2)$ is given below,

$$\begin{aligned} \text{Morr}[\psi](\tau_1, \tau_2) &= \int_{({}^{trap})\mathcal{M}(\tau_1, \tau_2)} |R\psi|^2 + r^{-2} |\psi|^2 + \left(1 - \frac{3m}{r}\right)^2 \left(|\nabla\psi|^2 + \frac{1}{r^2} |T\psi|^2 \right) \\ &\quad + \int_{({}^{trap})^c\mathcal{M}(\tau_1, \tau_2)} r^{-3} (|e_4\psi|^2 + r^{-1} |\psi|^2) + r^{-1} |\nabla\psi|^2 + r^{-1-\delta} |e_3\psi|^2 \end{aligned} \quad (5.1.14)$$

where $({}^{trap})^c\mathcal{M}$ denotes the complement of $({}^{trap})\mathcal{M}$.

Weighted bulk quantities

Define, for $0 < p < 2$,

$$\dot{B}_{p;R}[\psi](\tau_1, \tau_2) := \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{p-1} (p|\check{e}_4(\psi)|^2 + (2-p)|\nabla\psi|^2 + r^{-2} |\psi|^2), \quad (5.1.15)$$

$$B_p[\psi](\tau_1, \tau_2) := \text{Morr}[\psi](\tau_1, \tau_2) + \dot{B}_{p;4m_0}[\psi](\tau_1, \tau_2).$$

The bulk quantity $B_p[\psi](\tau_1, \tau_2)$ is equivalent to⁵

$$B_p[\psi](\tau_1, \tau_2) \simeq \int_{\tau_1}^{\tau_2} M_{p-1}[\psi](\tau) d\tau$$

where,

$$\begin{aligned} M_{p-1}[\psi](\tau) &= \int_{\Sigma_{\leq 4m_0}(\tau)} |\check{R}\psi|^2 + r^{-2} |\psi|^2 + \left(1 - \frac{3m}{r}\right)^2 \left(|\nabla\psi|^2 + \frac{m^2}{r^2} |\check{T}\psi|^2 \right) \\ &\quad + \int_{\Sigma_{\geq 4m_0}(\tau)} r^{p-1} (p|e_4(\psi)|^2 + (2-p)|\nabla\psi|^2 + r^{-2} |\psi|^2) + \int_{\Sigma_{\geq 4m_0}(\tau)} r^{-1-\delta} |e_3\psi|^2. \end{aligned}$$

⁵This equivalence follows from the coarea formula and the fact that the lapse of the τ -foliation is controlled uniformly from above and below.

Remark 5.1.5. Note that, for $\delta \leq p \leq 2 - \delta$,

$$B_p[\psi](\tau_1, \tau_2) := \text{Morr}[\psi](\tau_1, \tau_2) + \dot{B}_{p;4m_0}[\psi](\tau_1, \tau_2)$$

is equivalent to,

$$B_p[\psi](\tau_1, \tau_2) \simeq \text{Morr}[\psi](\tau_1, \tau_2) + \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-1} (|\check{e}_4(\psi)|^2 + |\nabla\psi|^2 + r^{-2}|e_3\psi|^2 + r^{-2}|\psi|^2).$$

Indeed,

$$\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-3}|e_3\psi|^2 \lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{-1-\delta}|e_3\psi|^2.$$

Therefore, since $r^2 (|\check{e}_4(\psi)|^2 + |\nabla\psi|^2) \lesssim |\mathfrak{d}\psi|^2$, we have,

$$B_p[\psi](\tau_1, \tau_2) \simeq \text{Morr}[\psi](\tau_1, \tau_2) + \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-3} (|\mathfrak{d}\psi|^2 + |\psi|^2). \quad (5.1.16)$$

Basic energy-flux quantity

The basic energy-flux quantity on a hypersurface $\Sigma(\tau)$ is defined by

$$E[\psi](\tau) = \int_{\Sigma(\tau)} \left(\frac{1}{2}(N_\Sigma, e_3)^2 |e_4\psi|^2 + \frac{1}{2}(N_\Sigma, e_4)^2 |e_3\psi|^2 + |\nabla\psi|^2 + r^{-2}|\psi|^2 \right). \quad (5.1.17)$$

Here N_Σ denotes a choice for the normal to Σ so that in particular we have

$$N_\Sigma = \begin{cases} N_\Sigma = e_3 & \text{on } (int)\Sigma, \\ N_\Sigma = e_4 & \text{on } (ext)\Sigma, \end{cases} \quad (5.1.18)$$

and, in view of (5.1.5),

$$(N_\Sigma, e_3) \leq -1 \text{ and } (N_\Sigma, e_4) \leq -1 \quad \text{on } (trap)\Sigma. \quad (5.1.19)$$

Weighted Energy-Flux type quantities

We have

$$\dot{E}_{p;R}[\psi](\tau) := \begin{cases} \int_{\Sigma_{\geq R}(\tau)} r^p (|\check{e}_4\psi|^2 + r^{-2}|\psi|^2) & \text{for } p \leq 1 - \delta, \\ \int_{\Sigma_{\geq R}(\tau)} r^p (|\check{e}_4\psi|^2 + r^{-p-1-\delta}|\psi|^2) & \text{for } p > 1 - \delta, \end{cases} \quad (5.1.20)$$

and

$$E_p[\psi](\tau) := E[\psi](\tau) + \dot{E}_{p;4m_0}[\psi](\tau). \quad (5.1.21)$$

Here \check{e}_4 denotes the first order operator

$$\check{e}_4\psi = r^{-1}\Upsilon^{-1}e_4(r\psi). \quad (5.1.22)$$

Remark 5.1.6. *To control the weighted quantities (5.1.21), it will be convenient to introduce in $^{(ext)}\mathcal{M}(r \geq 4m_0)$ the following renormalized frame*

$$e'_4 = \Upsilon^{-1}e_4, \quad e'_3 = \Upsilon e_3, \quad e'_\theta = e_\theta.$$

In particular, this yields

$$\check{e}_4\psi = r^{-1}e'_4(r\psi).$$

Note also that we have the following alternate form

$$\check{e}_4\psi = e'_4\psi + r^{-1}\psi + \frac{e'_4(r) - 1}{r}\psi$$

where $e'_4(r) - 1 = \Upsilon^{-1}e_4(r) - 1 = O(\epsilon r^{-1})$ in view of our assumption on Γ_g .

Flux quantities

The boundary of $\mathcal{M}(\tau_1, \tau_2)$ is given by

$$\partial\mathcal{M}(\tau_1, \tau_2) = \Sigma(\tau_1) \cup \Sigma(\tau_2) \cup \mathcal{A}(\tau_1, \tau_2) \cup \Sigma_*(\tau_1, \tau_2).$$

Our basic flux quantity along the spacelike hypersurfaces \mathcal{A} and Σ_* is given by

$$\begin{aligned} F[\psi](\tau_1, \tau_2) &:= \int_{\mathcal{A}(\tau_1, \tau_2)} \left(\delta_{\mathcal{H}}^{-1}|e_4\Psi|^2 + \delta_{\mathcal{H}}|e_3\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2 \right) \\ &\quad + \int_{\Sigma_*(\tau_1, \tau_2)} \left(|e_4\Psi|^2 + |e_3\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2 \right), \end{aligned} \quad (5.1.23)$$

with $\mathcal{A}(\tau_1, \tau_2) = \mathcal{A} \cap \mathcal{M}(\tau_1, \tau_2)$ and $\Sigma_*(\tau_1, \tau_2) = \Sigma_* \cap \mathcal{M}(\tau_1, \tau_2)$.

Weighted flux quantities

$$\begin{aligned} \dot{F}_p[\psi](\tau_1, \tau_2) &:= \int_{\Sigma_*(\tau_1, \tau_2)} r^p \left(|e_4\psi|^2 + |\nabla\psi|^2 + r^{-2}|\psi|^2 \right), \\ F_p[\psi](\tau_1, \tau_2) &:= F[\psi](\tau_1, \tau_2) + \dot{F}_p[\psi](\tau_1, \tau_2). \end{aligned} \quad (5.1.24)$$

Weighted quantities for the inhomogeneous term N

Recall the decomposition (5.1.7) for the inhomogeneous term N

$$N = N_g + e_3(rN_g) + N_m[\mathbf{q}].$$

We define, for $p \geq \delta$,

$$\begin{aligned} I_p[N_g](\tau_1, \tau_2) &= \left(\int_{\tau_1}^{\tau_2} d\tau \|N_g\|_{L^2(\text{trap}\Sigma(\tau))} \right)^2 + \int_{(\text{tr}q/p)\mathcal{M}(\tau_1, \tau_2)} r^{1+p} |N_g|^2 \\ &\quad + \int_{(\text{tr}q/p)\mathcal{M}(\tau_1, \tau_2)} r^{2+p} |N_g| |e_3(N_g)| + \sup_{\tau \in [\tau_1, \tau_2]} \int_{\Sigma(\tau)} r^{p+2} |N_g|^2 \\ &\quad + \int_{(\text{tr}q/p)\mathcal{M}(\tau_1, \tau_2)} r^{3+\delta} |e_3(N_g)|^2. \end{aligned} \quad (5.1.25)$$

Remark 5.1.7. While $N_m[\mathbf{q}]$ is present in the decomposition of the inhomogeneous term N , (5.1.25) only contains a norm for N_g . In fact, $N_m[\mathbf{q}]$ will always be absorbed by the left hand side wherever it appears.

Higher derivative quantities

We define the higher order derivative quantities $E^s[\psi]$, $\text{Mor}^s[\psi]$, $\text{Morr}^s[\psi]$, $E_p^s[\psi]$, $B_b^s[\psi]$, $M_p^s[\psi]$, $F^s[\psi]$, $F_p^s[\psi]$, $I_p^s[N_g]$ by the obvious procedure,

$$Q^s[\psi] = \sum_{k \leq s} Q[\mathfrak{d}^k \psi].$$

Remark 5.1.8. Note that in view of Remark 5.1.5 we can also write, equivalently, for $p < 2 - \delta$,

$$B_p^s[\psi](\tau_1, \tau_2) = \text{Morr}^s[\psi](\tau_1, \tau_2) + \int_{\mathcal{M}_{>4m_0}(\tau_1, \tau_2)} r^{p-3} |\mathfrak{d}^{\leq 1+s} \psi|^2. \quad (5.1.26)$$

Decay Norms

We introduce,

$$\begin{aligned}
\mathcal{E}_{p,d}^s[\psi] &:= \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d E_p^s[\psi](\tau), \\
\mathcal{B}_{p,d}^s[\psi] &:= \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d B_p^s[\psi](\tau, \tau_*), \\
&\simeq \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d \int_{\tau}^{\tau_*} M_{p-1}^s[\psi](\tau') d\tau', \\
\mathcal{F}_{p,d}^s[\psi] &:= \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d F_p^s[\psi](\tau, \tau_*), \\
\mathcal{I}_{p,d}^s[N_g] &:= \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d I_p^s[N_g](\tau, \tau_*).
\end{aligned} \tag{5.1.27}$$

5.2 Proof of Theorem M1

Recall that we have to prove for $k \leq k_{small} + 20$

$$\begin{aligned}
|\partial^k \mathbf{q}| &\lesssim \epsilon_0 r^{-1} (1 + \tau)^{-\frac{1}{2} - \delta_{extra}}, \\
|\partial^k \mathbf{q}| &\lesssim \epsilon_0 r^{-\frac{1}{2}} (1 + \tau)^{-1 - \delta_{extra}}, \\
|\partial^k e_3(\mathbf{q})| &\lesssim \epsilon_0 r^{-1} (1 + \tau)^{-1 - \delta_{extra}},
\end{aligned}$$

and

$$\int_{(int)\mathcal{M}(\tau, \tau_*)} |\partial^k e_3 \mathbf{q}|^2 + \int_{\Sigma_*(\tau, \tau_*)} |\partial^k e_3 \mathbf{q}|^2 \lesssim \epsilon_0^2 (1 + \tau)^{-2 - 2\delta_{extra}},$$

for some constant δ_{extra} such that $\delta_{dec} < \delta_{extra} < 2\delta_{dec}$.

5.2.1 Flux Decay Estimates for \mathbf{q}

The following result establishes decay of flux estimates for \mathbf{q} .

Theorem 5.2.1. *Let $0 < q_0 < 1$ be a fixed number and $s \leq k_{small} + 25$. Then, for all $\delta > 0$ we have, with a constant C depending only on s , δ and q_0 such that for all $\delta \leq p \leq 2 - \delta$, we have*

$$\begin{aligned}
&\mathcal{E}_{p,2+q_0-p}^s[\mathbf{q}] + \mathcal{B}_{p,2+q_0-p}^s[\mathbf{q}] + \mathcal{F}_{p,2+q_0-p}^s[\mathbf{q}] \\
&\lesssim \mathcal{E}_{q_0}^{s+2}[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+4}[\mathbf{q}](0) + \mathcal{I}_{q_0+2,0}^{s+5}[N_g] + \mathcal{I}_{\delta,2+q_0-\delta}^{s+5}[N_g],
\end{aligned} \tag{5.2.1}$$

where we recall that the decay norms $\mathcal{I}_{p,d}^s[N_g]$ are defined by,

$$\mathcal{I}_{p,d}^s[N_g] = \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d I_p^s[N_g](\tau, \tau_*).$$

Theorem 5.2.1 will be proved in section 5.4.3.

To prove Theorem M1 we have to eliminate the norms $\mathcal{I}_{p,d}^s[N_g]$ on the right hand side of Theorem 5.2.1.

Proposition 5.2.2. *Let $s \leq k_{small} + 30$ and assume*

$$q_0 < 4\delta_{dec} - 4\delta_0 \tag{5.2.2}$$

where

$$\delta_0 = \frac{33}{k_{large} - k_{small}} = \frac{33}{k_{large} - \lfloor \frac{k_{large}}{2} \rfloor - 1} \tag{5.2.3}$$

is the small constant appearing in Lemma 5.1.1. Then, the following estimates hold true,

$$\mathcal{I}_{q_0+2,0}^s[N_g] + \mathcal{I}_{\delta,2+q_0-\delta}^s[N_g] \lesssim \epsilon^4.$$

The proof of Proposition 5.2.2 is postponed to section 5.2.3. Together with Theorem 5.2.1, Proposition 5.2.2 immediately yields the proof of the following corollary.

Corollary 5.2.3. *In addition to the assumptions of Theorem 5.2.1 we assume*

$$2\delta_{dec} < q_0 < 4\delta_{dec} - 4\delta_0 \tag{5.2.4}$$

where $\delta_0 > 0$ is given by (5.2.3). Then for a sufficiently small bootstrap constant $\epsilon > 0$, for all $s \leq k_{small} + 25$ and for all $\delta \leq p \leq 2 - \delta$, we have

$$\mathcal{E}_{p,2+q_0-p}^s[\mathbf{q}] + \mathcal{B}_{p,2+q_0-p}^s[\mathbf{q}] + \mathcal{F}_{p,2+q_0-p}^s[\mathbf{q}] \lesssim \mathcal{E}_{q_0}^{s+2}[\tilde{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+4}[\mathbf{q}](0) + \epsilon^4.$$

5.2.2 Proof of Theorem M1

Since $\epsilon = \epsilon_0^{2/3}$, and in view of the control on \mathbf{q} at $\tau = 0$ provided by Theorem M0, we immediately deduce from Corollary 5.2.3, For all $0 < q \leq q_0$, $\delta \leq p \leq 2 - \delta$, and $s \leq k_{small} + 25$,

$$\mathcal{E}_{p,2+q_0-p}^s[\mathbf{q}] + \mathcal{B}_{p,2+q_0-p}^s[\mathbf{q}] + \mathcal{F}_{p,2+q_0-p}^s[\mathbf{q}] \lesssim \epsilon_0^2. \tag{5.2.5}$$

We will also need the following two propositions concerning L^2 estimates on spheres.

Proposition 5.2.4. *On any $S = S(\tau, r) \subset \Sigma(\tau)$, for $s \leq k_{small} + 25$,*

$$(1 + \tau)^{1+q_0} \int_{S_r} |\mathbf{q}^{(s)}|^2 \lesssim (\mathcal{E}_{1+\delta, 1+q_0-\delta}^s[\mathbf{q}])^{\frac{1}{2}} (\mathcal{E}_{1-\delta, 1+q_0+\delta}^s[\mathbf{q}])^{\frac{1}{2}} \quad (5.2.6)$$

and,

$$r^{-1}(1 + \tau)^{2+q_0-\delta} \int_{S_r} |\mathbf{q}^{(s)}|^2 \lesssim \mathcal{E}_{\delta, 2+q_0-\delta}^s[\mathbf{q}]. \quad (5.2.7)$$

Proposition 5.2.5. *We have for $s \leq k_{small} + 25$*

$$(1 + \tau)^{2+q_0-\delta} \int_{\Sigma_*(\tau, \tau_*)} |e_3 \mathfrak{d}^{\leq s} \mathbf{q}|^2 \lesssim \mathcal{F}_{\delta, 2+q_0-\delta}^s[\mathbf{q}]. \quad (5.2.8)$$

Also, on any $S = S(\tau, r) \subset \Sigma(\tau)$, for $s \leq k_{small} + 23$, we have

$$(1 + \tau)^{2+q_0-\delta} \int_{S_r} |e_3 \mathfrak{d}^{\leq s} \mathbf{q}|^2 \lesssim \epsilon_0^2 + \mathcal{F}_{\delta, 2+q_0-\delta}^{s+1}[\mathbf{q}] + \mathcal{E}_{\delta, 2+q_0-\delta}^{s+2}[\mathbf{q}]. \quad (5.2.9)$$

The proof of Proposition 5.2.4 is postponed to section 5.4.4, and the proof of Proposition 5.2.5 is postponed to section 5.4.5.

We now conclude the proof of Theorem M1. Indeed, in view of (5.2.5), Proposition 5.2.4 and Proposition 5.2.5, we infer for $s \leq k_{small} + 25$

$$\begin{aligned} (1 + \tau)^{2+q_0-\delta} \int_{(int)\mathcal{M}(\tau, \tau_*)} |\mathfrak{d}^{\leq s+1} \mathbf{q}|^2 &\lesssim \epsilon_0^2, \\ (1 + \tau)^{1+q_0} \int_{S_r} |\mathbf{q}^{(s)}|^2 &\lesssim \epsilon_0^2, \\ r^{-1}(1 + \tau)^{2+q_0-\delta} \int_{S_r} |\mathbf{q}^{(s)}|^2 &\lesssim \epsilon_0^2, \\ (1 + \tau)^{2+q_0-\delta} \int_{\Sigma_*(\tau, \tau_*)} |e_3 \mathfrak{d}^{\leq s} \mathbf{q}|^2 &\lesssim \epsilon_0^2, \end{aligned}$$

and for $s \leq k_{small} + 23$

$$(1 + \tau)^{2+q_0-\delta} \int_S |\mathfrak{d}^s e_3 \mathbf{q}|^2 \lesssim \epsilon_0^2.$$

In view of the standard Sobolev inequality on the 2-surfaces S i.e.,

$$\|\psi\|_{L^\infty(S)} \lesssim r^{-1} \|(r \nabla \mathfrak{V})^{\leq 2} \psi\|_{L^2(S)},$$

we immediately infer for $s \leq k_{small} + 23$

$$\begin{aligned} |\mathbf{q}^{(s)}| &\lesssim \epsilon_0 r^{-1} (1 + \tau)^{-\frac{1}{2} - \frac{q_0}{2}}, \\ |\mathbf{q}^{(s)}| &\lesssim \epsilon_0 r^{-\frac{1}{2}} (1 + \tau)^{-1 - \frac{q_0 - \delta}{2}}, \end{aligned}$$

and for $s \leq k_{small} + 21$

$$|\mathfrak{d}^s e_3(\mathbf{q})| \lesssim \epsilon_0 r^{-1} (1 + \tau)^{-1 - \frac{q_0 - \delta}{2}}.$$

Recall that $q_0 > 2\delta_{dec}$ and that $\delta > 0$ can be chosen arbitrarily small so that we have $q_0 - \delta > 2\delta_{dec}$. In particular, we may choose

$$\delta_{extra} := \frac{q_0 - \delta}{2}, \quad \delta_{extra} > \delta_{dec}, \quad (5.2.10)$$

which together with the above estimates for \mathbf{q} implies for $s \leq k_{small} + 25$

$$\begin{aligned} (1 + \tau)^{2+q_0-\delta} \int_{(int)\mathcal{M}(\tau, \tau_*)} |\mathfrak{d}^{\leq s+1} \mathbf{q}|^2 &\lesssim \epsilon_0^2, \\ (1 + \tau)^{2+2\delta_{extra}} \int_{\Sigma_*(\tau, \tau_*)} |e_3 \mathfrak{d}^{\leq s} \mathbf{q}|^2 &\lesssim \epsilon_0^2, \end{aligned}$$

for $s \leq k_{small} + 23$

$$\begin{aligned} |\mathbf{q}^{(s)}| &\lesssim \epsilon_0 r^{-1} (1 + \tau)^{-\frac{1}{2} - \delta_{extra}}, \\ |\mathbf{q}^{(s)}| &\lesssim \epsilon_0 r^{-\frac{1}{2}} (1 + \tau)^{-1 - \delta_{extra}}, \end{aligned}$$

and for $s \leq k_{small} + 21$

$$|\mathfrak{d}^s e_3(\mathbf{q})| \lesssim \epsilon_0 r^{-1} (1 + \tau)^{-1 - \delta_{extra}}$$

as desired. This concludes the proof of Theorem M1.

5.2.3 Proof of Proposition 5.2.2

Recall that,

$$\begin{aligned} I_p[N_g](\tau_1, \tau_2) &= \left(\int_{\tau_1}^{\tau_2} d\tau \|N_g\|_{L^2((trap)\Sigma(\tau))} \right)^2 + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{1+p} |N_g|^2 \\ &\quad + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{2+p} |N_g| |e_3(N_g)| + \sup_{\tau \in [\tau_1, \tau_2]} \int_{\Sigma(\tau)} r^{p+2} |N_g|^2 \\ &\quad + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{3+\delta} |e_3(N_g)|^2 \end{aligned}$$

and,

$$\mathcal{I}_{p,d}^s[N_g] = \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d I_p^s[N_g](\tau, \tau_*).$$

Since we have

$$r^\delta (1 + \tau)^{2+q_0-\delta} \lesssim r^{2+q_0} + (1 + \tau)^{2+q_0},$$

and

$$\int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r^2 |\mathfrak{d}^{\leq s} e_3(N_g)| |\mathfrak{d}^{\leq s} N_g| \lesssim \int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r |\mathfrak{d}^{\leq s} N_g|^2 + \int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r^3 |\mathfrak{d}^{\leq s} e_3(N_g)|^2,$$

we infer

$$\begin{aligned} & \mathcal{I}_{q_0+2,0}^s[N_g] + \mathcal{I}_{\delta,2+q_0-\delta}^s[N_g] \tag{5.2.11} \\ & \lesssim \sup_{0 \leq \tau \leq \tau_*} \left[\int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r^{4+q_0} |\mathfrak{d}^{\leq s+1} N_g|^2 + \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^{4+q_0} |\mathfrak{d}^{\leq s} N_g|^2 \right. \\ & \quad \left. + (1 + \tau)^{2+q_0} \left(\int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r |\mathfrak{d}^{\leq s} N_g|^2 + \int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r^{3+\delta} |\mathfrak{d}^{\leq s} e_3(N_g)|^2 \right) \right. \\ & \quad \left. + \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^2 |\mathfrak{d}^{\leq s} N_g|^2 \right) + (1 + \tau)^{2+q_0} \left(\int_{\tau}^{\tau_*} d\tau' \|\mathfrak{d}^{\leq s} N_g\|_{L^2((trap)\Sigma(\tau'))}^2 \right)^2 \Big]. \end{aligned}$$

In order to prove Proposition 5.2.2, it suffices to estimate the right-hand side of (5.2.11). To this end, we will estimate separately the terms with highest power of r , i.e. the first two terms, and the terms with highest power the τ , i.e. the four last terms.

Terms with highest power of r in (5.2.11)

We estimate the first two terms of (5.2.11). Recall from Lemma 5.1.2 that

$$N_g = r^2 \mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta)).$$

Recall from Lemma 5.1.1 we have

$$\max_{0 \leq k \leq k_{large} - 3} \sup_{(ext) \mathcal{M}(r \geq 4m_0)} r^{\frac{7}{2} + \frac{\delta_B}{2}} \left(|\mathfrak{d}^k \alpha| + |\mathfrak{d}^k \beta| \right) \lesssim \epsilon.$$

We infer for $s \leq k_{large} - 6$

$$|\mathfrak{d}^{\leq s+1} N_g| \lesssim \epsilon r^{-\frac{7}{2} - \frac{\delta_B}{2}} |r^2 \mathfrak{d}^{\leq s+3} \Gamma_g|$$

and hence, for $s \leq k_{large} - 6$, we deduce

$$\begin{aligned} & \int_{(trq/p)_{\mathcal{M}(\tau, \tau_*)}} r^{4+q_0} |\mathfrak{d}^{\leq s+1} N_g|^2 + \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^{4+q_0} |\mathfrak{d}^{\leq s} N_g|^2 \\ & \lesssim \epsilon^2 \int_{(trq/p)_{\mathcal{M}(\tau, \tau_*)}} r^{-3-\delta_B+q_0} (r^2 \mathfrak{d}^{\leq s+3} \Gamma_g)^2 + \epsilon^2 \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^{-3-\delta_B+q_0} (r^2 \mathfrak{d}^{\leq s+2} \Gamma_g)^2. \end{aligned}$$

Since we also have for $s \leq k_{large} - 6$

$$\sup_{r_0 \geq 4m_0} \int_{\{r=r_0\}} (r^2 \mathfrak{d}^{\leq s+3} \Gamma_g)^2 \lesssim \epsilon^2, \quad \int_{\mathcal{M}_{r \leq 4m_0}} (\mathfrak{d}^{\leq s+3} \Gamma_g)^2 \lesssim \epsilon^2, \quad \sup_{\mathcal{M}} |r^2 \mathfrak{d}^{\leq s+2} \Gamma_g| \lesssim \epsilon,$$

we deduce

$$\begin{aligned} & \int_{(trq/p)_{\mathcal{M}(\tau, \tau_*)}} r^{4+q_0} |\mathfrak{d}^{\leq s+1} N_g|^2 + \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^{4+q_0} |\mathfrak{d}^{\leq s} N_g|^2 \\ & \lesssim \epsilon^4 \left(1 + \int_{r \geq 4m_0} \frac{dr}{r^{1+\delta_B-q_0}} \right). \end{aligned}$$

Since $q_0 < 4\delta_{dec}$ and $\delta_B \geq 4\delta_{dec}$, we have $q_0 < \delta_B$ and hence, we obtain for $s \leq k_{large} - 6$

$$\int_{(trq/p)_{\mathcal{M}(\tau, \tau_*)}} r^{4+q_0} |\mathfrak{d}^{\leq s+1} N_g|^2 + \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^{4+q_0} |\mathfrak{d}^{\leq s} N_g|^2 \lesssim \epsilon^4.$$

This is the desired control of the terms with highest power of r in (5.2.11).

Terms with highest power of τ in (5.2.11)

We estimate the four last terms of (5.2.11). In view of Lemma 5.1.1 with $k_{loss} = 33$, so that

$$\delta_0 = \frac{33}{k_{large} - k_{small} - 2} = \frac{33}{k_{large} - \lfloor \frac{k_{large}}{2} \rfloor - 3},$$

we have

$$\begin{aligned} \left| \mathfrak{d}^{\leq k_{small}+33} \Gamma_g \right| & \lesssim \epsilon r^{-2} \tau^{-1/2-\delta_{dec}+2\delta_0}, \\ \left| \mathfrak{d}^{\leq k_{small}+33} \Gamma_g \right| & \lesssim \epsilon r^{-1} \tau^{-1-\delta_{dec}+2\delta_0}, \\ \left| \mathfrak{d}^{\leq S+32} e_3 \Gamma_g \right| & \lesssim \epsilon r^{-2} [\tau^{-1-\delta_{dec}}]^{1-\delta_0} \lesssim \epsilon r^{-2} \tau^{-1-\delta_{dec}+2\delta_0}, \\ \left| \mathfrak{d}^{\leq k_{small}+33} (\alpha, \beta) \right| & \lesssim \epsilon r^{-3} (\tau + r)^{-1/2-\delta_{dec}+2\delta_0}, \\ \left| \mathfrak{d}^{\leq k_{small}+33} (\alpha, \beta) \right| & \lesssim \epsilon r^{-2} (\tau + r)^{-1-\delta_{dec}+2\delta_0}, \\ \left| \mathfrak{d}^{\leq S+32} e_3 (\alpha, \beta) \right| & \lesssim \epsilon r^{-3-\frac{1}{2}\delta_0} [\tau^{-1-\delta_{dec}}]^{1-\delta_0} \lesssim \epsilon r^{-3} \tau^{-1-\delta_{dec}+2\delta_0}. \end{aligned}$$

In particular, together with the bootstrap assumption for $k \leq k_{small}$, the pointwise bound

$$|\mathfrak{d}^{\leq k_{large}-5} \alpha| + |\mathfrak{d}^{\leq k_{large}-5} \beta| \lesssim \epsilon r^{-\frac{7}{2}-\frac{\delta_B}{2}}$$

and since $N_g = r^2 \mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta))$, we infer for $s \leq k_{small} + 30$

$$\begin{aligned} |\mathfrak{d}^s N_g| &\lesssim \epsilon^2 r^{-3} \tau^{-1-2\delta_{dec}+2\delta_0} \\ |\mathfrak{d}^s N_g| &\lesssim \epsilon^2 r^{-1} \tau^{-2-2\delta_{dec}+2\delta_0}, \\ |\mathfrak{d}^s e_3(N_g)| &\lesssim \epsilon^2 r^{-3} \tau^{-\frac{3}{2}-2\delta_{dec}+2\delta_0}, \\ |\mathfrak{d}^s e_3(N_g)| &\lesssim \epsilon^2 r^{-\frac{7}{2}-\frac{\delta_B}{2}} \tau^{-1-\delta_{dec}+2\delta_0}. \end{aligned} \tag{5.2.12}$$

Using these 4 bounds and interpolation, we infer for $\delta > 0$

$$\begin{aligned} &(1+\tau)^{2+q_0} \left(\int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r |\mathfrak{d}^{\leq s} N_g|^2 + \int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r^{3+\delta} |\mathfrak{d}^{\leq s} e_3(N_g)|^2 \right. \\ &+ \left. \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^2 |\mathfrak{d}^{\leq s} N_g|^2 \right) + (1+\tau)^{2+q_0} \left(\int_{\tau}^{\tau_*} d\tau' \|\mathfrak{d}^{\leq s} N_g\|_{L^2((trap)\Sigma(\tau'))} \right)^2 \\ &\lesssim \epsilon^4 (1+\tau)^{2+q_0} \int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r (r^{-3} \tau'^{-1-2\delta_{dec}+2\delta_0})^{1+\delta} (r^{-1} \tau'^{-2-2\delta_{dec}+2\delta_0})^{1-\delta} \\ &+ \epsilon^4 (1+\tau)^{2+q_0} \int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r^{3+\delta} (r^{-3} \tau'^{-\frac{3}{2}-2\delta_{dec}+2\delta_0})^{2-2\delta} (r^{-\frac{7}{2}-\frac{\delta_B}{2}} \tau'^{-1-\delta_{dec}+2\delta_0})^{2\delta} \\ &+ \epsilon^4 (1+\tau)^{2+q_0} \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^2 (r^{-3} \tau'^{-1-2\delta_{dec}+2\delta_0})^2 \\ &+ \epsilon^4 (1+\tau)^{2+q_0} \left(\int_{\tau}^{\tau_*} \tau'^{-2-2\delta_{dec}+2\delta_0} d\tau' \right)^2 \\ &\lesssim \epsilon^4 (1+\tau)^{2+q_0} \int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r^{-3-\delta\delta_B} \tau'^{-3-4\delta_{dec}+\delta+4\delta_0+2\delta\delta_{dec}} \\ &+ \epsilon^4 (1+\tau)^{2+q_0} \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^{-4} \tau'^{-2-4\delta_{dec}+4\delta_0} + \epsilon^4 (1+\tau)^{2+q_0} \left(\int_{\tau}^{\tau_*} \tau'^{-2-2\delta_{dec}+2\delta_0} d\tau' \right)^2 \end{aligned}$$

and since $\delta > 0$, we obtain

$$\begin{aligned} &(1+\tau)^{2+q_0} \left(\int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r |\mathfrak{d}^{\leq s} N_g|^2 + \int_{(trq/p) \mathcal{M}(\tau, \tau_*)} r^{3+\delta} |\mathfrak{d}^{\leq s} e_3(N_g)|^2 \right. \\ &+ \left. \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^2 |\mathfrak{d}^{\leq s} N_g|^2 \right) + (1+\tau)^{2+q_0} \left(\int_{\tau}^{\tau_*} d\tau' \|\mathfrak{d}^{\leq s} N_g\|_{L^2((trap)\Sigma(\tau'))} \right)^2 \\ &\lesssim \epsilon^4 (1+\tau)^{q_0-4\delta_{dec}+\delta+4\delta_0+2\delta\delta_{dec}}. \end{aligned}$$

As we have $q_0 < 4\delta_{dec} - 4\delta_0$, there exists $\delta > 0$ small enough such that

$$q_0 - 4\delta_{dec} + \delta + 4\delta_0 + 2\delta\delta_{dec} \leq 0,$$

and hence

$$\begin{aligned} & (1 + \tau)^{2+q_0} \left(\int_{(trq/p)\mathcal{M}(\tau, \tau_*)} r |\mathfrak{d}^{\leq s} N_g|^2 + \int_{(trq/p)\mathcal{M}(\tau, \tau_*)} r^{3+\delta} |\mathfrak{d}^{\leq s} e_3(N_g)|^2 \right. \\ & \left. + \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^2 |\mathfrak{d}^{\leq s} N_g|^2 \right) + (1 + \tau)^{2+q_0} \left(\int_{\tau}^{\tau_*} d\tau' \|\mathfrak{d}^{\leq s} N_g\|_{L^2((trap)\Sigma(\tau'))} \right)^2 \lesssim \epsilon^4. \end{aligned}$$

This is the desired control of the terms with highest power of τ in (5.2.11). Together with (5.2.11) and the above control of the terms with highest power of r , we infer

$$\begin{aligned} & \mathcal{I}_{q_0+2,0}^s[N_g] + \mathcal{I}_{\delta,2+q_0-\delta}^s[N_g] \\ & \lesssim \sup_{0 \leq \tau \leq \tau_*} \left[\int_{(trq/p)\mathcal{M}(\tau, \tau_*)} r^{4+q_0} |\mathfrak{d}^{\leq s+1} N_g|^2 + \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^{4+q_0} |\mathfrak{d}^{\leq s} N_g|^2 \right. \\ & \quad \left. + (1 + \tau)^{2+q_0} \left(\int_{(trq/p)\mathcal{M}(\tau, \tau_*)} r |\mathfrak{d}^{\leq s} N_g|^2 + \int_{(trq/p)\mathcal{M}(\tau, \tau_*)} r^{3+\delta} |\mathfrak{d}^{\leq s} e_3(N_g)|^2 \right. \right. \\ & \quad \left. \left. + \sup_{\tau' \in [\tau, \tau_*]} \int_{\Sigma(\tau')} r^2 |\mathfrak{d}^{\leq s} N_g|^2 \right) + (1 + \tau)^{2+q_0} \left(\int_{\tau}^{\tau_*} d\tau' \|\mathfrak{d}^{\leq s} N_g\|_{L^2((trap)\Sigma(\tau'))} \right)^2 \right] \\ & \lesssim \epsilon^4 \end{aligned}$$

which is the desired estimate. This concludes the proof of Proposition 5.2.2.

5.3 Improved weighted estimates

The goal of this section is to prove the two following theorems on improved weighted estimates.

Theorem 5.3.1. *Assume \mathfrak{q} verifies following wave equation, see (5.0.1),*

$$\square_2 \mathfrak{q} + \kappa \underline{\kappa} \mathfrak{q} = N$$

with N given, in view of Lemma 5.1.2, by

$$N = N_g + e_3(rN_g) + N_m[\mathfrak{q}].$$

Then, for any $\delta \leq p \leq 2 - \delta$, $0 \leq s \leq k_{small} + 30$,

$$\sup_{\tau \in [\tau_1, \tau_2]} E_p^s[\mathfrak{q}](\tau) + B_p^s[\mathfrak{q}](\tau_1, \tau_2) + F_p^s[\mathfrak{q}](\tau_1, \tau_2) \lesssim E_p^s[\mathfrak{q}](\tau_1) + I_p^{s+1}[N_g](\tau_1, \tau_2). \quad (5.3.1)$$

The next result deals with weighted estimates for the quantity

$$\check{\mathfrak{q}} = f_2 \check{e}_4 \mathfrak{q}, \quad (5.3.2)$$

where f_2 is a fixed smooth function of r defined as follows,

$$f_2(r) = \begin{cases} r^2 & \text{for } r \geq 6m_0, \\ 0 & \text{for } r \leq 4m_0. \end{cases} \quad (5.3.3)$$

Theorem 5.3.2. *Assume \mathfrak{q} verifies equation, see (5.0.1),*

$$\square_2 \mathfrak{q} + \kappa \underline{\kappa} \mathfrak{q} = N$$

with,

$$N = N_g + e_3(rN_g) + N_m[\mathfrak{q}]$$

as in Lemma 5.1.2. Then, for any $-1 + \delta < q \leq 1 - \delta$, $0 \leq s \leq k_{small} + 29$,

$$\sup_{\tau \in [\tau_1, \tau_2]} E_q^s[\check{\mathfrak{q}}](\tau) + B_q^s[\check{\mathfrak{q}}](\tau_1, \tau_2) \lesssim E_q^s[\check{\mathfrak{q}}](\tau_1) + E_{q+1}^{s+1}[\mathfrak{q}](\tau_1) + I_{q+2}^{s+2}[N_g](\tau_1, \tau_2). \quad (5.3.4)$$

Remark 5.3.3. *Note that in (5.3.1) and (5.3.4), the term $N_m[\mathfrak{q}]$ does not appear in the right-hand side since it turns out that it can be absorbed by the left hand side.*

The proof of Theorem 5.3.1 is postponed to section 5.3.2, and the proof of Theorem 5.3.2 is postponed to section 5.3.3. These proofs will rely on weighted energy flux estimates introduced in the next section.

5.3.1 Proof of Theorem 10.2.1

Assume given a spacetime \mathcal{M} verifying the bootstrap assumptions with small constant $\epsilon > 0$. The proof of Theorem 5.3.1 and Theorem 5.3.2 will rely on estimates stated below for solutions $\psi \in \mathfrak{s}_2(\mathcal{M})$ of the equation,

$$\square_2 \psi + V\psi = N, \quad V = \kappa \underline{\kappa}. \quad (5.3.5)$$

Basic weighted estimates

Theorem 5.3.4. *Recall the definitions in (5.1.21), (5.1.15). The following holds for any $0 \leq s \leq k_{small} + 30$. For all $\delta \leq p \leq 2 - \delta$, we have,*

$$\sup_{\tau \in [\tau_1, \tau_2]} E_p^s[\psi](\tau) + B_p^s[\psi](\tau_1, \tau_2) + F_p^s[\psi](\tau_1, \tau_2) \lesssim E_p^s[\psi](\tau_1) + J_p^s[\psi, N](\tau_1, \tau_2), \quad (5.3.6)$$

where, for $p \geq \delta$, we have introduced the notation

$$\begin{aligned}
 J_{p,R}[\psi, N](\tau_1, \tau_2) &:= \left| \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^p \check{e}_4 \psi N \right|, \\
 J_p[\psi, N](\tau_1, \tau_2) &:= \left(\int_{\tau_1}^{\tau_2} d\tau \|N\|_{L^2(\text{trap})\Sigma(\tau)} \right)^2 + \int_{(\text{trap})\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} |N|^2 \\
 &\quad + J_{p,4m_0}[\psi, N](\tau_1, \tau_2),
 \end{aligned} \tag{5.3.7}$$

and

$$J_p^s[\psi, N](\tau_1, \tau_2) := \sum_{k \leq s} J_p[\mathfrak{d}^k \psi, \mathfrak{d}^k N](\tau_1, \tau_2).$$

The proof of Theorem 5.3.4 is postponed to section 10.4.5.

Higher weighted estimates

The next result deals with weighted estimates for the quantity

$$\check{\psi} = f_2 \check{e}_4 \psi, \tag{5.3.8}$$

where f_2 is a fixed smooth function of r defined as follows,

$$f_2(r) = \begin{cases} r^2 & \text{for } r \geq 6m_0, \\ 0 & \text{for } r \leq 4m_0. \end{cases} \tag{5.3.9}$$

Theorem 5.3.5. *The following holds for any $-1 + \delta < q \leq 1 - \delta$, $0 \leq s \leq k_{small} + 29$,*

$$\begin{aligned}
 \sup_{\tau \in [\tau_1, \tau_2]} E_q^s[\check{\psi}](\tau) + B_q^s[\check{\psi}](\tau_1, \tau_2) &\lesssim E_q^s[\check{\psi}](\tau_1) + \check{J}_q^s[\check{\psi}, N](\tau_1, \tau_2) \\
 &\quad + E_{\max(q, \delta)}^{s+1}[\psi](\tau_1) + J_{\max(q, \delta)}^{s+1}[\psi, N],
 \end{aligned} \tag{5.3.10}$$

where we have introduced the notation

$$\begin{aligned}
 \check{J}_q[\check{\psi}, N](\tau_1, \tau_2) &:= J_{q,4m_0} \left[\check{\psi}, r^2 \left(e_4 N + \frac{3}{r} N \right) \right] (\tau_1, \tau_2) \\
 &= \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+2} (\check{e}_4 \check{\psi}) \cdot \left(e_4 N + \frac{3}{r} N \right),
 \end{aligned}$$

and

$$\check{J}_q^s[\check{\psi}, N](\tau_1, \tau_2) := \sum_{k \leq s} \check{J}_q[\mathfrak{d}^k \check{\psi}, \mathfrak{d}^k N](\tau_1, \tau_2).$$

The proof of Theorem 5.3.5 is postponed to section 10.4.6.

We now proceed to the proof of Theorem 5.3.1 and Theorem 5.3.2 in the next 2 sections. The proofs will follow from the structure of the nonlinear term N of \mathfrak{q} provided by Lemma 5.1.2 and the use of Theorem 5.3.4 and Theorem 5.3.5.

5.3.2 Proof of Theorem 5.3.1

Applying Theorem 5.3.4 to the equation for \mathfrak{q} , with N given by Lemma 5.1.2, we derive corresponding estimates with the norm $J_p^s[\mathfrak{q}, N](\tau_1, \tau_2)$ on the right hand side, i.e. for $0 \leq s \leq k_{small} + 30$, and for $\delta \leq p \leq 2 - \delta$,

$$\sup_{\tau \in [\tau_1, \tau_2]} E_p^s[\mathfrak{q}](\tau) + B_p^s[\mathfrak{q}](\tau_1, \tau_2) + F_p^s[\mathfrak{q}](\tau_1, \tau_2) \lesssim E_p^s[\mathfrak{q}](\tau_1) + J_p^s[\mathfrak{q}, N](\tau_1, \tau_2). \quad (5.3.11)$$

To prove Theorem 5.3.1, it suffices, in view of (5.3.11), to estimate $J_p^s[\mathfrak{q}, N](\tau_1, \tau_2)$. Recall that, see (5.3.7) and (5.1.25)

$$\begin{aligned} I_p[N](\tau_1, \tau_2) &= \left(\int_{\tau_1}^{\tau_2} d\tau \|N\|_{L^2((trap)\Sigma(\tau))} \right)^2 + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{1+p} |N|^2 \\ &\quad + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{2+p} |N_g| |e_3(N_g)| + \sup_{\tau \in [\tau_1, \tau_2]} \int_{\Sigma(\tau)} r^{p+2} |N|^2 \\ &\quad + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{3+\delta} |e_3(N_g)|^2 \end{aligned}$$

and,

$$\begin{aligned} J_{p,R}[\mathfrak{q}, N] &= \left| \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^p \check{e}_4(\mathfrak{q}) N \right|, \\ J_p[\mathfrak{q}, N](\tau_1, \tau_2) &= \left(\int_{\tau_1}^{\tau_2} d\tau \|N\|_{L^2((trap)\Sigma(\tau))} \right)^2 + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} |N|^2 \\ &\quad + J_{p,4m_0}^s[\mathfrak{q}, N](\tau_1, \tau_2), \\ J_p^s[\mathfrak{q}, N](\tau_1, \tau_2) &= \sum_{k \leq s} J_p[\mathfrak{d}^k \mathfrak{q}, \mathfrak{d}^k N], \end{aligned}$$

Recall also from (5.1.9)

$$\mathfrak{d}^k N = \mathfrak{d}^{\leq k} N_g + e_3(\mathfrak{d}^k(rN_g)) + \mathfrak{d}^k N_m[\mathfrak{q}] \quad (5.3.12)$$

and consider separately the three terms.

Case of $N_m[\mathbf{q}]$. Recall that $N_m[\mathbf{q}] = \mathfrak{d}^{\leq 1}(\Gamma_g \cdot \mathbf{q})$. We have, schematically,

$$\mathfrak{d}^k N_m[\mathbf{q}] = \mathfrak{d}^{1+k}(\Gamma_g \cdot \mathbf{q}) = \sum_{i+j=k+1} \mathfrak{d}^{\leq i} \Gamma_g \mathfrak{d}^{\leq j} \mathbf{q}.$$

We make use of the following consequence of the bootstrap assumptions for $k \leq k_{large} - 5$

$$|\mathfrak{d}^{\leq k} \Gamma_g| \leq \epsilon r^{-2}$$

to deduce,

$$|\mathfrak{d}^k N_m[\mathbf{q}]| \lesssim \epsilon r^{-2} |\mathfrak{d}^{\leq k+1} \mathbf{q}|. \quad (5.3.13)$$

We deduce,

$$\begin{aligned} J_{p,4m_0}^s[\mathbf{q}, N_m[\mathbf{q}]](\tau_1, \tau_2) &\lesssim \sum_{k \leq s} \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^p |\check{e}_4 \mathbf{q}^{(k)}| |\mathfrak{d}^k N_m[\mathbf{q}]| \\ &\lesssim \epsilon \sum_{k \leq s} \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-3} |\mathfrak{d}^{1+k} \mathbf{q}|^2. \end{aligned}$$

Thus, recalling Remark 5.1.8, we infer

$$J_{p,4m_0}^s[\mathbf{q}, N_m[\mathbf{q}]](\tau_1, \tau_2) \lesssim \epsilon B_p^s[\mathbf{q}](\tau_1, \tau_2). \quad (5.3.14)$$

Next, we estimate in view of (5.3.13)

$$\int_{(tr\mathcal{I}^p)_{\mathcal{M}(\tau_1, \tau_2)}} r^{1+\delta} |\mathfrak{d}^k N_m[\mathbf{q}]|^2 \lesssim \epsilon \int_{(tr\mathcal{I}^p)_{\mathcal{M}(\tau_1, \tau_2)}} r^{\delta-3} |\mathfrak{d}^{\leq k+1} \mathbf{q}|^2$$

which yields, using again Remark 5.1.8,

$$\int_{(tr\mathcal{I}^p)_{\mathcal{M}(\tau_1, \tau_2)}} r^{1+\delta} |\mathfrak{d}^k N_m[\mathbf{q}]|^2 \lesssim \epsilon B_\delta^s[\mathbf{q}](\tau_1, \tau_2). \quad (5.3.15)$$

We next estimate the integral

$$\int_{\tau_1}^{\tau_2} d\tau \|\mathfrak{d}^k N_m[\mathbf{q}]\|_{L^2((tr\mathcal{I}^p)\Sigma(\tau))}.$$

In view of the definition of $N_m[\mathbf{q}] = \mathfrak{d}^{\leq 1}(\Gamma_g \cdot \mathbf{q})$,

$$\begin{aligned} \mathfrak{d}^k N_m[\mathbf{q}] &= \mathfrak{d}^{\leq k+1}(\Gamma_g \cdot \mathbf{q}) = \sum_{i+j=k+1} \mathfrak{d}^{\leq i} \Gamma_g \mathfrak{d}^{\leq j} \mathbf{q} \\ &= \mathfrak{d}^{j \leq (k+1)/2} \Gamma_g \mathfrak{d}^{\leq k+1} \mathbf{q} + \mathfrak{d}^{j \leq (k+1)/2} \mathbf{q} \mathfrak{d}^{\leq k+1} \Gamma_g = J_1 + J_2. \end{aligned}$$

Now, since $\frac{k+1}{2} \leq k_{small}$ we have

$$\left| \mathfrak{d}^{j \leq (k+1)/2} \Gamma_g \right| \lesssim \epsilon (1 + \tau)^{-1 - \delta_{dec}}$$

Hence,

$$\begin{aligned} \|J_1\|_{L^2((trap)\Sigma(\tau))}^2 &= \int_{(trap)\Sigma(\tau)} \left| \mathfrak{d}^{j \leq (k+1)/2} \Gamma_g \right|^2 \left| \mathfrak{d}^{\leq k+1} \mathfrak{q} \right|^2 \\ &\lesssim \epsilon^2 (1 + \tau)^{-2 - 2\delta_{dec}} E^s[\mathfrak{q}](\tau) \end{aligned}$$

i.e.,

$$\|J_1\|_{L^2((trap)\Sigma(\tau))} \lesssim \epsilon (1 + \tau)^{-1 - \delta_{dec}} (E^s[\mathfrak{q}](\tau))^{1/2}.$$

For J_2 we write,

$$\begin{aligned} \|J_2\|_{L^2((trap)\Sigma(\tau))}^2 &= \int_{(trap)\Sigma(\tau)} \left| \mathfrak{d}^{j \leq (k+1)/2} \mathfrak{q} \right|^2 \left| \mathfrak{d}^{\leq k+1} \Gamma_g \right|^2 \\ &\lesssim \left(\sup_{(trap)\Sigma(\tau)} \left| \mathfrak{d}^{j \leq (k+1)/2} \mathfrak{q} \right| \right)^2 \int_{(trap)\Sigma(\tau)} \left| \mathfrak{d}^{\leq k+1} \Gamma_g \right|^2 \\ &\lesssim \int_{(trap)\Sigma(\tau)} \left| \mathfrak{d}^{\leq (k+1)/2 + 2} \mathfrak{q} \right|^2 \int_{(trap)\Sigma(\tau)} \left| \mathfrak{d}^{\leq k+1} \Gamma_g \right|^2 \end{aligned}$$

or, since $(k+1)/2 + 2 \leq s$,

$$\|J_2\|_{L^2((trap)\Sigma(\tau))} \lesssim \left[\int_{(trap)\Sigma(\tau)} \left| \mathfrak{d}^{\leq s} \mathfrak{q} \right|^2 \right]^{1/2} \left[\int_{(trap)\Sigma(\tau)} \left| \mathfrak{d}^{\leq k+1} \Gamma_g \right|^2 \right]^{1/2}.$$

In view of the above estimates for J_1 and J_2 , we deduce, for all $k \leq s \leq k_{large} - 5$

$$\begin{aligned} &\int_{\tau_1}^{\tau_2} d\tau \| \mathfrak{d}^k N_m[\mathfrak{q}] \|_{L^2((trap)\Sigma(\tau))} \\ &\lesssim \epsilon \sup_{\tau_1 \leq \tau \leq \tau_2} (E^s[\mathfrak{q}](\tau))^{1/2} + \int_{\tau_1}^{\tau_2} d\tau \left[\int_{(trap)\Sigma(\tau)} \left| \mathfrak{d}^{\leq s} \mathfrak{q} \right|^2 \right]^{1/2} \left[\int_{(trap)\Sigma(\tau)} \left| \mathfrak{d}^{\leq s} \Gamma_g \right|^2 \right]^{1/2} \\ &\lesssim \epsilon \sup_{\tau_1 \leq \tau \leq \tau_2} (E^s[\mathfrak{q}](\tau))^{1/2} + \left(\int_{(trap)\Sigma(\tau_1, \tau_2)} \left| \mathfrak{d}^{\leq s} \mathfrak{q} \right|^2 \right)^{\frac{1}{2}} \left(\int_{\mathcal{M}_{r \leq 4m_0}} \left| \mathfrak{d}^{\leq s} \Gamma_g \right|^2 \right)^{1/2} \end{aligned}$$

Making use of the following consequence of the bootstrap assumptions

$$\left(\int_{\mathcal{M}_{r \leq 4m_0}} \left| \mathfrak{d}^{\leq s} \Gamma_g \right|^2 \right)^{1/2} \lesssim \epsilon,$$

as well as the fact that

$$\int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} |\mathfrak{d}^{\leq s} \mathbf{q}|^2 \lesssim \text{Morr}^s[\mathbf{q}](\tau_1, \tau_2),$$

we deduce,

$$\left(\int_{\tau_1}^{\tau_2} d\tau \|\mathfrak{d}^k N_m[\mathbf{q}]\|_{L^2((trap)\Sigma(\tau))} \right)^2 \lesssim \epsilon^2 \sup_{\tau_1 \leq \tau \leq \tau_2} E^s[\mathbf{q}](\tau) + \epsilon^2 \text{Morr}^s[\mathbf{q}](\tau_1, \tau_2) \quad (5.3.16)$$

which together with (5.3.15) and (5.3.14) yields for any $p \geq \delta$

$$J_p^s[\mathbf{q}, N_m[\mathbf{q}]](\tau_1, \tau_2) \lesssim \epsilon^2 \sup_{\tau_1 \leq \tau \leq \tau_2} E^s[\mathbf{q}](\tau) + \epsilon B_p^s[\mathbf{q}](\tau_1, \tau_2). \quad (5.3.17)$$

Case of N_g . We write, as before,

$$\begin{aligned} J_{p,4m_0}^s[\mathbf{q}, N_g](\tau_1, \tau_2) &\lesssim \sum_{k \leq s} \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^p |\check{e}_4 \mathbf{q}^{(k)} \mathfrak{d}^k N_g| \\ &\lesssim \sum_{k \leq s} \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-1} |\check{e}_4 \mathbf{q}^{(k)}|^2 \right)^{1/2} \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p+1} |\mathfrak{d}^k N_g|^2 \right)^{1/2}. \end{aligned}$$

Therefore,

$$\begin{aligned} J_{p,4m_0}^s[\mathbf{q}, N_g](\tau_1, \tau_2) &\lesssim (B_p^s[\mathbf{q}](\tau_1, \tau_2))^{1/2} (I_p^s[N_g](\tau_1, \tau_2))^{1/2} \\ &\lesssim \delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2) + \delta_1^{-1} I_p^s[N_g](\tau_1, \tau_2) \end{aligned}$$

where $\delta_1 > 0$ is chosen sufficiently small so that we can later absorb the term $\delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2)$ by the left hand side of our main estimate.

Also, we have in view of the definition of $I_p^s[N](\tau_1, \tau_2)$ and the fact that $p \geq \delta$

$$\left(\int_{\tau_1}^{\tau_2} d\tau \|\mathfrak{d}^{\leq s} N_g\|_{L^2((trap)\Sigma(\tau))} \right)^2 + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} |\mathfrak{d}^{\leq s} N_g|^2 \lesssim I_p^s[N_g](\tau_1, \tau_2).$$

Therefore,

$$\begin{aligned} J_p^s[\mathbf{q}, N_g](\tau_1, \tau_2) &= \left(\int_{\tau_1}^{\tau_2} d\tau \|\mathfrak{d}^{\leq s} N_g\|_{L^2((trap)\Sigma(\tau))} \right)^2 + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} |\mathfrak{d}^{\leq s} N_g|^2 \\ &\quad + J_{p,4m_0}^s[\mathbf{q}, N_g](\tau_1, \tau_2) \\ &\lesssim I_\delta^s[N_g](\tau_1, \tau_2) + \delta_1^{-1} I_p^s[N_g](\tau_1, \tau_2) + \delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2), \end{aligned}$$

i.e.,

$$J_p^s[\mathbf{q}, N_g](\tau_1, \tau_2) \lesssim \delta_1^{-1} I_p^s[N_g](\tau_1, \tau_2) + \delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2). \quad (5.3.18)$$

Case of $e_3(rN_g)$. First, note that we have

$$\begin{aligned} & \left(\int_{\tau_1}^{\tau_2} d\tau \| \mathfrak{d}^{\leq s} e_3(rN_g) \|_{L^2(\text{trap}\Sigma(\tau))} \right)^2 + \int_{(\text{tr}\mathcal{I}^p)_{\mathcal{M}(\tau_1, \tau_2)}} r^{1+\delta} | \mathfrak{d}^{\leq s} e_3(rN_g) |^2 \\ & \lesssim \left(\int_{\tau_1}^{\tau_2} d\tau \| \mathfrak{d}^{\leq s+1} N_g \|_{L^2(\text{trap}\Sigma(\tau))} \right)^2 + \int_{(\text{tr}\mathcal{I}^p)_{\mathcal{M}(\tau_1, \tau_2)}} r^{1+\delta} | \mathfrak{d}^{\leq s} N_g |^2 \\ & \quad + \int_{(\text{tr}\mathcal{I}^p)_{\mathcal{M}(\tau_1, \tau_2)}} r^{3+\delta} | \mathfrak{d}^{\leq s} e_3(N_g) |^2 \end{aligned}$$

where we used the fact that $| \mathfrak{d}^{\leq s} e_3(r) | \lesssim 1$ and $| \mathfrak{d}^{\leq s} r | \lesssim r$. Hence, we infer in view of the definition of $I_p^s[N](\tau_1, \tau_2)$ and the fact that $p \geq \delta$

$$\begin{aligned} & \left(\int_{\tau_1}^{\tau_2} d\tau \| \mathfrak{d}^{\leq s} e_3(rN_g) \|_{L^2(\text{trap}\Sigma(\tau))} \right)^2 + \int_{(\text{tr}\mathcal{I}^p)_{\mathcal{M}(\tau_1, \tau_2)}} r^{1+\delta} | \mathfrak{d}^{\leq s} e_3(rN_g) |^2 \\ & \lesssim I_p^{s+1}[N_g](\tau_1, \tau_2). \end{aligned} \quad (5.3.19)$$

We then estimate

$$J_{p,4m_0}[\mathbf{q}^{(k)}, e_3(\mathfrak{d}^k(rN_g))](\tau_1, \tau_2), \quad k \leq s.$$

To this end, we introduce a smooth cut-off function ϕ_0 vanishing for $r \leq 4m_0$ and equal to 1 for $r \geq 8m_0$. Then, we have

$$\begin{aligned} J_{p,4m_0}[\mathbf{q}^{(k)}, \mathfrak{d}^k(rN_g)](\tau_1, \tau_2) &= \left| \int_{\mathcal{M}(\tau_1, \tau_2)} r^p \check{e}_4 \mathbf{q}^{(k)} e_3 \mathfrak{d}^k(rN_g) \right| \\ &\lesssim J_{p,4m_0}[\mathbf{q}^{(k)}, \phi_0 \mathfrak{d}^k(rN_g)](\tau_1, \tau_2) \\ &\quad + J_{p,4m_0}[\mathbf{q}^{(k)}, (1 - \phi_0)rN_g](\tau_1, \tau_2). \end{aligned} \quad (5.3.20)$$

In view of the fact that $1 - \phi_0$ is supported in $r \leq 8m_0$, we easily obtain

$$J_{p,4m_0}[\mathbf{q}^{(k)}, (1 - \phi_0)rN_g](\tau_1, \tau_2) \lesssim \left(\sup_{\tau_1 \leq \tau \leq \tau_2} E^s[\mathbf{q}](\tau) + B_p^s[\mathbf{q}](\tau_1, \tau_2) \right)^{1/2} \left(I_p^{s+1}[N_g](\tau_1, \tau_2) \right)^{\frac{1}{2}}$$

and hence

$$\begin{aligned} & J_{p,4m_0}[\mathbf{q}^{(k)}, (1 - \phi_0)rN_g](\tau_1, \tau_2) \\ & \lesssim \delta_1 \left(\sup_{\tau_1 \leq \tau \leq \tau_2} E^s[\mathbf{q}](\tau) + B_p^s[\mathbf{q}](\tau_1, \tau_2) \right) + \delta_1^{-1} I_p^{s+1}[N_g](\tau_1, \tau_2) \end{aligned} \quad (5.3.21)$$

where $\delta_1 > 0$ is chosen sufficiently small so that we can later absorb the terms $\delta_1 \sup_{\tau_1 \leq \tau \leq \tau_2} E^s[\mathbf{q}](\tau)$ and $\delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2)$ by the left hand side of our main estimate.

It remains to estimate the terms

$$J_{p,4m_0}[\mathbf{q}^{(k)}, \phi_0 e_3(\mathfrak{d}^k(rN_g))](\tau_1, \tau_2), \quad k \leq s$$

which is supported for $r \geq 4m_0$. Note that $e_3(rN_g)$ behaves like rN_g and therefore the same sequence of estimates as for N_g would lead to a loss of r^{-1} . For this reason we need to integrate by parts by parts in e_3 .

Proposition 5.3.6. *The following estimate holds true, for all $k \leq s \leq k_{large} - 5$,*

$$\sum_{k \leq s} J_{p,4m_0}[\mathbf{q}^{(k)}, \phi_0 e_3(\mathfrak{d}^k(rN_g))](\tau_1, \tau_2) \lesssim \delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2) + \delta_1^{-1} I_p^{s+1}[N_g](\tau_1, \tau_2) \quad (5.3.22)$$

for a sufficiently small $\delta_1 > 0$.

We postponed the proof of Proposition 5.3.6 to the end of the section. We are now in position to conclude the proof of Theorem 5.3.1.

Proof of Theorem 5.3.1. (5.3.21) and (5.3.22) yield

$$\sum_{k \leq s} J_{p,4m_0}[\mathbf{q}^{(k)}, e_3(\mathfrak{d}^k(rN_g))](\tau_1, \tau_2) \lesssim \delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2) + \delta_1^{-1} I_p^{s+1}[N_g](\tau_1, \tau_2).$$

Together with (5.3.17), (5.3.18) and (5.3.19), we infer

$$J_p^s[\mathbf{q}, N](\tau_1, \tau_2) \lesssim (\delta_1 + \epsilon) B_p^s[\mathbf{q}](\tau_1, \tau_2) + \delta_1^{-1} I_p^{s+1}[N_g](\tau_1, \tau_2) + \epsilon^2 \sup_{\tau_1 \leq \tau \leq \tau_2} E^s[\mathbf{q}](\tau).$$

In view of (5.3.11), this concludes the proof of Theorem 5.3.1. □

The proof of Proposition 5.3.6 will rely in particular on the following identity.

Lemma 5.3.7. *The following hold true for any $\psi \in \mathfrak{s}_2$*

- We have, schematically,

$$e_3 e_4(r\psi) = -r\Box_2\psi + r\Delta_2\psi + r^{-1}\mathfrak{d}\psi. \quad (5.3.23)$$

- The following identity holds true, schematically,

$$e_3 e_4(r\mathfrak{d}^k\psi) = -\mathfrak{d}^{\leq k}(r\Box_2\psi) + r\Delta_2(\mathfrak{d}^{\leq k}\psi) + r^{-1}\mathfrak{d}^{\leq k+1}\psi. \quad (5.3.24)$$

Proof. We start with the following identity for $\psi \in \mathfrak{s}_2$, see Definition 2.4.2,

$$\square_2\psi = -e_3e_4\psi + \underline{\Delta}_2\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi - \frac{1}{2}\kappa e_3\psi + 2\eta e_\theta\psi$$

from which we deduce,

$$r\square_2\psi = -re_3e_4\psi + r\left(\underline{\Delta}_2\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi - \frac{1}{2}\kappa e_3\psi + 2\eta e_\theta\psi\right).$$

On the other hand,

$$\begin{aligned} re_3e_4\psi &= e_3(re_4\psi) - (e_3r)e_4\psi = e_3(e_4(r\psi) - e_4(r)\psi) - (e_3r)e_4\psi \\ &= e_3e_4(r\psi) - e_4(r)e_3\psi - (e_3r)e_4\psi - (e_3e_4r)\psi. \end{aligned}$$

Hence,

$$\begin{aligned} r\square_2\psi &= -e_3e_4(r\psi) + e_4(r)e_3\psi + (e_3r)e_4\psi + (e_3e_4r)\psi + r\underline{\Delta}_2\psi \\ &+ r\left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi - \frac{1}{2}r\kappa e_3\psi + 2r\eta e_\theta\psi \\ &= -e_3e_4(r\psi) + r\underline{\Delta}\psi + \left(e_4r - \frac{1}{2}r\kappa\right)e_3\psi + \left(e_3r - \frac{1}{2}r\underline{\kappa} + 2r\underline{\omega}\right)e_4\psi + 2r\eta e_\theta\psi \\ &= -e_3e_4(r\psi) + r\underline{\Delta}\psi + \frac{r}{2}Ae_3\psi + \frac{r}{2}(\underline{A} + 4\underline{\omega})e_4\psi + 2r\eta e_\theta\psi \end{aligned}$$

i.e.,

$$e_3e_4(r\psi) = -r\square_2\psi + r\underline{\Delta}_2\psi + \frac{r}{2}Ae_3\psi + \frac{r}{2}(\underline{A} + 4\underline{\omega})e_4\psi + 2r\eta e_\theta\psi.$$

or, schematically, in view of the definition of $\mathfrak{d}\psi$ and the estimate $|\underline{\omega}| + r|\Gamma_g| + |\Gamma_b| \lesssim r^{-1}$,

$$\begin{aligned} e_3e_4(r\psi) &= -r\square_2\psi + r\underline{\Delta}_2\psi + \left(r\Gamma_g + \Gamma_b + r^{-1}\right)e_3\psi \\ &= -r\square_2\psi + r\underline{\Delta}_2\psi + r^{-1}\mathfrak{d}\psi \end{aligned}$$

which is (5.3.23).

To derive the identity for higher derivatives we write, schematically,

$$\mathfrak{d}^k e_3e_4(r\psi) = -\mathfrak{d}^k(r\square_2\psi) + \mathfrak{d}^k(r\underline{\Delta}_2\psi) + \mathfrak{d}^k(r\Gamma_g\mathfrak{d}\psi).$$

We write,

$$\begin{aligned} \mathfrak{d}^k e_3e_4(r\psi) &= e_3e_4(r\mathfrak{d}^k\psi) + [\mathfrak{d}^k, e_3e_4r]\psi = e_3e_4(r\mathfrak{d}^k\psi) + [\mathfrak{d}^k, e_3]\mathfrak{d}\psi + e_3[\mathfrak{d}^k, e_4r]\psi, \\ \mathfrak{d}^k(r\underline{\Delta}_2\psi) &= r\underline{\Delta}_2\mathfrak{d}^k\psi + [\mathfrak{d}^k, r\underline{\Delta}]\psi = r\underline{\Delta}_2\mathfrak{d}^k\psi + [\mathfrak{d}^k, r^{-1}]\mathfrak{d}^2\psi + r^{-1}[\mathfrak{d}^k, r^2\underline{\Delta}]\psi. \end{aligned}$$

In view of the identities for $[e_3, \mathfrak{D}]$ and $[e_4, \mathfrak{D}]$ of Lemma 2.2.13, the identities of Proposition 2.1.25 for commutation formulas involving \mathfrak{d}_k and \mathfrak{d}_k^* derivatives, and the commutator formula for $[e_3, e_4]$, we have schematically

$$\begin{aligned} [e_3, e_3] &= 0, & [\mathfrak{D}, r^2 \Delta] &= \mathfrak{D} + 1, & [e_3, e_4 r] &= (r^{-1} + \Gamma_g) \mathfrak{D} \\ [e_3, \mathfrak{D}] &= \Gamma_b \mathfrak{D} + \Gamma_b, & [e_4 r, \mathfrak{D}] &= (r^2 \xi + r \Gamma_g) \mathfrak{D} + r \Gamma_g \end{aligned}$$

In view of the estimates for Γ_g , Γ_b , and the fact that $\xi = 0$ for $r \geq 4m_0$, we infer

$$[\mathfrak{D}^k, e_3] = r^{-1} \mathfrak{D}^{\leq k}, \quad [\mathfrak{D}^k, r^2 \Delta] = \mathfrak{D}^{\leq k+1}, \quad [\mathfrak{D}^k, r^{-1}] = r^{-1} \mathfrak{D}^{\leq k-1}$$

and hence

$$\begin{aligned} \mathfrak{D}^k e_3 e_4(r\psi) &= e_3 e_4(r\mathfrak{D}^k \psi) + e_3 [\mathfrak{D}^k, e_4 r] \psi + r^{-1} \mathfrak{D}^{\leq k+1} \psi, \\ \mathfrak{D}^k(r \Delta_2 \psi) &= r \Delta_2 \mathfrak{D}^k \psi + r^{-1} \mathfrak{D}^{\leq k+1} \psi. \end{aligned}$$

Also, we have

$$[re_4, e_4 r] = [re_4, e_4] r + e_4 [re_4, r] = -e_4(r) e_4 r - e_4 r e_4(r) = -2e_4 r + r^{-1} \mathfrak{D}$$

and we infer by induction, schematically,

$$[(re_4)^j, e_4 r] = e_4 r (re_4)^{\leq j-1} + r^{-1} \mathfrak{D}^{\leq j}$$

so that, together with

$$[\mathfrak{D}_{\searrow}^{k-j}, e_4 r] = r^{-1} \mathfrak{D}^{\leq k-j},$$

we infer

$$[\mathfrak{D}^k, e_4 r] = [(re_4)^j \mathfrak{D}_{\searrow}^{k-j}, e_4 r] = e_4 r (re_4)^{\leq j-1} \mathfrak{D}_{\searrow}^{k-j} + r^{-1} \mathfrak{D}^{\leq k}.$$

We deduce

$$e_3 e_4(r(re_4)^j \mathfrak{D}_{\searrow}^{k-j} \psi) = -(re_4)^j \mathfrak{D}_{\searrow}^{k-j}(r \square_2 \psi) + r \Delta_2(\mathfrak{D}^k \psi) + r^{-1} \mathfrak{D}^{\leq k+1} \psi + e_4 r (re_4)^{\leq j-1} \mathfrak{D}_{\searrow}^{k-j} \psi.$$

We infer by induction on j

$$e_3 e_4(r(re_4)^j \mathfrak{D}_{\searrow}^{k-j} \psi) = -(re_4)^{\leq j} \mathfrak{D}_{\searrow}^{k-j}(r \square_2 \psi) + r \Delta_2(\mathfrak{D}^{\leq k} \psi) + r^{-1} \mathfrak{D}^{\leq k+1} \psi$$

and hence

$$e_3 e_4(r \mathfrak{D}^k \psi) = -\mathfrak{D}^{\leq k}(r \square_2 \psi) + r \Delta_2(\mathfrak{D}^{\leq k} \psi) + r^{-1} \mathfrak{D}^{\leq k+1} \psi$$

which is (5.3.24). This concludes the proof of Lemma 5.3.7. \square

We now are in position to prove Proposition 5.3.6.

Proof of Proposition 5.3.6. We integrate by parts,

$$\begin{aligned} J_{p,4m_0}[\mathbf{q}^{(k)}, \phi_0 \mathfrak{d}^k(rN_g)](\tau_1, \tau_2) &\lesssim \left| \int_{\mathcal{M}(\tau_1, \tau_2)} e_3(\phi_0(r)r^p \check{e}_4 \mathbf{q}^{(k)}) \mathfrak{d}^k(rN_g) \right| + |B_p^k(\tau_1)| + |B_p^k(\tau_2)| \\ &\quad + \left| \int_{\mathcal{M}(\tau_1, \tau_2)} \mathbf{Div}(e_3) \phi_0(r)r^p \check{e}_4 \mathbf{q}^{(k)} \mathfrak{d}^k(rN_g) \right| \end{aligned} \quad (5.3.25)$$

where $\mathbf{Div}(e_3)$ denotes the spacetime divergence of e_3 , and where the with boundary terms are bounded by

$$\begin{aligned} |B_p^k(\tau_1)| &\lesssim \int_{\Sigma(\tau_1)} r^p |\check{e}_4 \mathbf{q}^{(k)}| |\mathfrak{d}^k(rN_g)|, \\ |B_p^k(\tau_2)| &\lesssim \int_{\Sigma(\tau_2)} r^p |\check{e}_4 \mathbf{q}^{(k)}| |\mathfrak{d}^k(rN_g)|. \end{aligned}$$

We estimate,

$$\begin{aligned} |B_p^k(\tau)| &\lesssim \int_{\Sigma(\tau)} r^p |\check{e}_4 \mathbf{q}^{(k)}| |\mathfrak{d}^k(rN_g)| \lesssim \left(\int_{\Sigma(\tau)} r^p |\check{e}_4 \mathbf{q}^{(k)}|^2 \right)^{1/2} \left(\int_{\Sigma(\tau)} r^p |\mathfrak{d}^k(rN_g)|^2 \right)^{1/2} \\ &\lesssim \left(E_p^k[\mathbf{q}](\tau) \right)^{1/2} \left(\int_{\Sigma(\tau)} r^{p+2} |\mathfrak{d}^k N_g|^2 \right)^{1/2}. \end{aligned}$$

We deduce, with $\delta_1 > 0$ a sufficiently small constant, for any $\tau \in [\tau_1, \tau_2]$,

$$\begin{aligned} |B_p^k(\tau_1)| &\lesssim \delta_1 \sup_{\tau_1 \leq \tau \leq \tau_2} E_p^k[\mathbf{q}](\tau) + \delta_1^{-1} \sup_{\tau_1 \leq \tau \leq \tau_2} \int_{\Sigma(\tau)} r^{p+2} |N_g^{\leq k}|^2, \\ |B_p^k(\tau_2)| &\lesssim \delta_1 \sup_{\tau_1 \leq \tau \leq \tau_2} E_p^k[\mathbf{q}](\tau) + \delta_1^{-1} \sup_{\tau_1 \leq \tau \leq \tau_2} \int_{\Sigma(\tau)} r^{p+2} |N_g^{\leq k}|^2. \end{aligned} \quad (5.3.26)$$

Next, notice that $\mathbf{Div}(e_3) = \underline{\kappa} - 2\underline{\omega}$ so that

$$|\mathbf{Div}(e_3)| \lesssim r^{-1}.$$

Together with the fact that $e_3(\Phi_0(r))$ is supported in $4m_0 \leq r \leq 8m_0$, the fact that $|e_3(r)| \lesssim 1$ and

$$r \check{e}_4 \mathbf{q}^{(k)} = e_4(r \mathbf{q}^{(k)}) + O(r^{-1}) e_4(\mathbf{q}^{(k)}),$$

we infer

$$\begin{aligned}
& \left| \int_{\mathcal{M}(\tau_1, \tau_2)} e_3 (\phi_0(r) r^p \check{e}_4 \mathbf{q}^{(k)}) \mathfrak{d}^k(r N_g) \right| + \left| \int_{\mathcal{M}(\tau_1, \tau_2)} \mathbf{Div}(e_3) \phi_0(r) r^p \check{e}_4 \mathbf{q}^{(k)} \mathfrak{d}^k(r N_g) \right| \\
& \lesssim \left| \int_{\mathcal{M}(\tau_1, \tau_2)} \phi_0(r) r^{p-1} e_3 e_4(r \mathbf{q}^{(k)}) \mathfrak{d}^k(r N_g) \right| + \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-1} |\check{e}_4(\mathbf{q}^{(k)})| |\mathfrak{d}^k(r N_g)| \\
& \quad + \int_{\mathcal{M}_{4m_0 \leq r \leq 8m_0}(\tau_1, \tau_2)} |\check{e}_4(\mathbf{q}^{(k)})| |\mathfrak{d}^k(r N_g)| \\
& \lesssim \left| \int_{\mathcal{M}(\tau_1, \tau_2)} \phi_0(r) r^{p-1} e_3 e_4(r \mathbf{q}^{(k)}) \mathfrak{d}^k(r N_g) \right| \\
& \quad + \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-1} |\check{e}_4(\mathbf{q}^{(k)})|^2 \right)^{\frac{1}{2}} \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p+1} |\mathfrak{d}^{\leq k} N_g|^2 \right)^{\frac{1}{2}}
\end{aligned}$$

and hence

$$\begin{aligned}
& \left| \int_{\mathcal{M}(\tau_1, \tau_2)} e_3 (\phi_0(r) r^p \check{e}_4 \mathbf{q}^{(k)}) \mathfrak{d}^k(r N_g) \right| + \left| \int_{\mathcal{M}(\tau_1, \tau_2)} \mathbf{Div}(e_3) \phi_0(r) r^p \check{e}_4 \mathbf{q}^{(k)} \mathfrak{d}^k(r N_g) \right| \\
& \lesssim \left| \int_{\mathcal{M}(\tau_1, \tau_2)} \phi_0(r) r^{p-1} e_3 e_4(r \mathbf{q}^{(k)}) \mathfrak{d}^k(r N_g) \right| + (B_p^s[\mathbf{q}](\tau_1, \tau_2))^{1/2} (I_p^s[N_g](\tau_1, \tau_2))^{\frac{1}{2}}
\end{aligned}$$

which yields

$$\begin{aligned}
& \left| \int_{\mathcal{M}(\tau_1, \tau_2)} e_3 (\phi_0(r) r^p \check{e}_4 \mathbf{q}^{(k)}) \mathfrak{d}^k(r N_g) \right| + \left| \int_{\mathcal{M}(\tau_1, \tau_2)} \mathbf{Div}(e_3) \phi_0(r) r^p \check{e}_4 \mathbf{q}^{(k)} \mathfrak{d}^k(r N_g) \right| \\
& \lesssim |L^k| + \delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2) + \delta_1^{-1} I_p^s[N_g](\tau_1, \tau_2) \tag{5.3.27}
\end{aligned}$$

where $\delta_1 > 0$ is chosen sufficiently small so that we can later absorb the term $\delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2)$ by the left hand side of our main estimate, and where we have introduced the notation

$$L^k := \int_{\mathcal{M}(\tau_1, \tau_2)} \phi_0(r) r^{p-1} e_3 e_4(r \mathbf{q}^{(k)}) \mathfrak{d}^k(r N_g). \tag{5.3.28}$$

It remains to estimate the term L^k . Making use of Lemma 5.3.7, we deduce

$$\begin{aligned}
L^k &= \int_{\mathcal{M}(\tau_1, \tau_2)} \phi_0(r) r^{p-1} e_3 e_4(r\mathbf{q}^{(k)}) \mathfrak{d}^k(rN_g) \\
&= - \int_{\mathcal{M}(\tau_1, \tau_2)} \phi_0(r) r^{p-1} \mathfrak{d}^{\leq k}(r\Box_2\mathbf{q}) \mathfrak{d}^k(rN_g) \\
&\quad + \int_{\mathcal{M}(\tau_1, \tau_2)} \phi_0(r) r^p \mathfrak{A}_2(\mathfrak{d}^{\leq k}\mathbf{q}) \mathfrak{d}^k(rN_g) \\
&\quad + \int_{\mathcal{M}(\tau_1, \tau_2)} \phi_0(r) r^{p-2} \mathfrak{d}^{\leq k+1}\mathbf{q} \mathfrak{d}^k(rN_g) \\
&= L_1^k + L_2^k + L_3^k.
\end{aligned}$$

We first estimate L_3^k as follows

$$\begin{aligned}
|L_3^k| &\lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-2} |\mathfrak{d}^{\leq k+1}\mathbf{q}| |\mathfrak{d}^k(rN_g)| \\
&\lesssim \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-3} |\mathfrak{d}^{\leq k+1}\mathbf{q}|^2 \right)^{1/2} \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p+1} |\mathfrak{d}^{\leq k}N_g|^2 \right)^{1/2}
\end{aligned}$$

In view of Remark 5.1.8 we thus deduce,

$$\begin{aligned}
|L_3^k| &\lesssim (B_p^k[\mathbf{q}])^{1/2} \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p+1} |\mathfrak{d}^{\leq k}N_g|^2 \right)^{1/2} \\
&\lesssim (B_p^k[\mathbf{q}](\tau_1, \tau_2))^{1/2} (I_p^k[N_g](\tau_1, \tau_2))^{1/2}
\end{aligned}$$

and hence

$$|L_3^k| \lesssim \delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2) + \delta_1^{-1} I_p^s[N_g](\tau_1, \tau_2) \quad (5.3.29)$$

where $\delta_1 > 0$ is chosen sufficiently small so that we can later absorb the term $\delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2)$ by the left hand side of our main estimate.

We now estimate the term

$$L_2^k = \int_{\mathcal{M}(\tau_1, \tau_2)} \phi_0(r) r^p \mathfrak{A}_2(\mathfrak{d}^k\mathbf{q}) \mathfrak{d}^k(rN_g)$$

by first performing another integration by parts in the angular directions

$$\begin{aligned}
|L_2^k| &\lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-2} |\mathfrak{d}^{k+1}\mathbf{q}| |\mathfrak{d}^{k+1}(rN_g)| \\
&\lesssim \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-3} |\mathfrak{d}^{k+1}\mathbf{q}|^2 \right)^{1/2} \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p+1} |\mathfrak{d}^{\leq k+1}N_g|^2 \right)^{1/2} \\
&\lesssim (B_p^k[\mathbf{q}](\tau_1, \tau_2))^{1/2} (I_p^{k+1}[N_g](\tau_1, \tau_2))^{1/2}.
\end{aligned}$$

Hence,

$$|L_2^k| \lesssim \delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2) + \delta_1^{-1} I_p^{s+1}[N_g](\tau_1, \tau_2) \quad (5.3.30)$$

where $\delta_1 > 0$ is chosen sufficiently small so that we can later absorb the term $\delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2)$ by the left hand side of our main estimate.

It remains to estimate the term,

$$L_1^k = - \int_{\mathcal{M}(\tau_1, \tau_2)} \phi_0(r) r^{p-1} \mathfrak{d}^{\leq k}(r \square_2 \mathbf{q}) \mathfrak{d}^k(r N_g).$$

Making use of the equation verified by \mathbf{q} , i.e., $\square_2 \mathbf{q} = -\kappa \underline{\kappa} \mathbf{q} + N$, we deduce,

$$\mathfrak{d}^k(r \square_2 \mathbf{q}) = -\mathfrak{d}^k(r \kappa \underline{\kappa} \mathbf{q}) + \mathfrak{d}^k(r N).$$

Recall (5.1.9)

$$\mathfrak{d}^k N = \mathfrak{d}^{\leq k} N_g + e_3(\mathfrak{d}^k(r N_g)) + \mathfrak{d}^k N_m[\mathbf{q}].$$

We infer

$$\begin{aligned} \mathfrak{d}^{\leq k}(r N) &= r \mathfrak{d}^{\leq k} N + \mathfrak{d}^{\leq k-1} N \\ &= r \mathfrak{d}^{\leq k} N_g + r e_3(\mathfrak{d}^{\leq k}(r N_g)) + r \mathfrak{d}^{\leq k} N_m[\mathbf{q}] \end{aligned}$$

and hence

$$\begin{aligned} |\mathfrak{d}^k(r \square_2 \mathbf{q})| &\lesssim r^{-1} |\mathfrak{d}^{\leq k} \mathbf{q}| + r |\mathfrak{d}^{\leq k} N_g| + r^2 |\mathfrak{d}^{\leq k} e_3(N_g)| + r |\mathfrak{d}^k N_m[\mathbf{q}]| \\ &\lesssim r^{-1} |\mathfrak{d}^{\leq k+1} \mathbf{q}| + r |\mathfrak{d}^{\leq k} N_g| + r^2 |\mathfrak{d}^{\leq k} e_3(N_g)|. \end{aligned} \quad (5.3.31)$$

Note that we have used in the last inequality the form of $N_m[\mathbf{q}] = \mathfrak{d}^{\leq 1}(\Gamma_g \mathbf{q})$ and the fact that $|\Gamma_g| \leq \epsilon r^{-2}$. We deduce, using (5.3.31),

$$\begin{aligned} |L_1^k| &\lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-1} |\mathfrak{d}^{\leq k+1} \mathbf{q}| |\mathfrak{d}^{\leq k} N_g| + \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p+1} |\mathfrak{d}^{\leq k} N_g|^2 \\ &\quad + \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p+2} |\mathfrak{d}^{\leq k} e_3(N_g)| |\mathfrak{d}^{\leq k} N_g| \\ &\lesssim (B_p^k[\mathbf{q}](\tau_1, \tau_2))^{1/2} (I_p^k[N_g](\tau_1, \tau_2))^{1/2} + I_p^k[N_g](\tau_1, \tau_2). \end{aligned}$$

We deduce

$$|L_1^k| \lesssim \delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2) + \delta_1^{-1} I_p^s[N_g](\tau_1, \tau_2) \quad (5.3.32)$$

where $\delta_1 > 0$ is chosen sufficiently small so that we can later absorb the term $\delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2)$ by the left hand side of our main estimate.

Together with (5.3.29) and (5.3.30) we deduce,

$$|L^k| \lesssim \delta_1 B_p^k[\mathbf{q}](\tau_1, \tau_2) + \delta_1^{-1} I_p^k[N_g](\tau_1, \tau_2). \quad (5.3.33)$$

Together with (5.3.25), (5.3.26) and (5.3.28), we infer,

$$\sum_{k \leq s} J_{p, 4m_0}[\mathbf{q}^{(k)}, \phi_0 \mathfrak{d}^k(rN_g)](\tau_1, \tau_2) \lesssim \delta_1 B_p^s[\mathbf{q}](\tau_1, \tau_2) + \delta_1^{-1} I_p^{s+1}[N_g](\tau_1, \tau_2)$$

which concludes the proof of Proposition 5.3.6. \square

5.3.3 Proof of Theorem 5.3.2

We apply Theorem 5.3.5 to the case when $\psi = \mathbf{q}$. Hence,

$$\begin{aligned} E_q^s[\check{\mathbf{q}}](\tau_2) + B_q^s[\check{\mathbf{q}}](\tau_1, \tau_2) &\lesssim E_q^s[\check{\mathbf{q}}](\tau_1) + \check{J}_q^s[\check{\mathbf{q}}, N](\tau_1, \tau_2) \\ &\quad + E_{\max(q, \delta)}^{s+1}[\mathbf{q}](\tau_1) + J_{\max(q, \delta)}^{s+1}[\mathbf{q}, N](\tau_1, \tau_2). \end{aligned} \quad (5.3.34)$$

Also, recall that

$$\check{\mathbf{q}} = f_2 \check{e}_4 \mathbf{q},$$

where f_2 is a fixed smooth function of r defined as follows,

$$f_2(r) = \begin{cases} r^2 & \text{for } r \geq 6m_0, \\ 0 & \text{for } r \leq 4m_0. \end{cases} \quad (5.3.35)$$

In particular, $\check{\mathbf{q}}$ is supported in $r \geq 4m_0$, and hence, in view of Remark 5.1.5,

$$B_q[\check{\mathbf{q}}](\tau_1, \tau_2) \simeq \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q-3} |\mathfrak{d}\check{\mathbf{q}}|^2, \quad (5.3.36)$$

where we have used the fact that $-1 + \delta \leq q \leq 1 - \delta$.

First, notice that the proof of Theorem 5.3.1 yields

$$J_{\max(q, \delta)}^{s+1}[\mathbf{q}, N](\tau_1, \tau_2) \lesssim \sup_{\tau_1 \leq \tau \leq \tau_2} E^{s+1}[\mathbf{q}](\tau) + B_{\max(q, \delta)}^{s+1}[\mathbf{q}](\tau_1, \tau_2) + I_{\max(q, \delta)}^{s+2}[N_g](\tau_1, \tau_2).$$

Hence, using Theorem 5.3.1, together with the fact that $\max(q, \delta) \leq 1 - \delta$, we infer

$$J_{\max(q, \delta)}^{s+1}[\mathbf{q}, N](\tau_1, \tau_2) \lesssim E_{\max(q, \delta)}^{s+1}[\mathbf{q}](\tau_1) + I_{\max(q, \delta)}^{s+2}[N_g](\tau_1, \tau_2).$$

Since $q \geq -1 + \delta$, we have $\max(q, \delta) \leq \delta \leq q + 1$ and thus

$$J_{\max(q, \delta)}^{s+1}[\mathbf{q}, N](\tau_1, \tau_2) \lesssim E_{q+1}^{s+1}[\mathbf{q}](\tau_1) + I_{q+1}^{s+2}[N_g](\tau_1, \tau_2). \quad (5.3.37)$$

It only remains to estimate the term

$$\check{J}_q^s[\check{\mathbf{q}}, N](\tau_1, \tau_2) = \sum_{k \leq s} \check{J}_q[\mathfrak{d}^k \check{\mathbf{q}}, \mathfrak{d}^k N](\tau_1, \tau_2)$$

with,

$$\begin{aligned} \check{J}_q[\check{\mathbf{q}}, N](\tau_1, \tau_2) &= J_{q, 4m_0} \left[\check{\mathbf{q}}, r^2 \left(e_4 N + \frac{3}{r} N \right) \right] (\tau_1, \tau_2) \\ &= \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+2} (\check{e}_4 \check{\mathbf{q}}) \cdot \left(e_4 N + \frac{3}{r} N \right). \end{aligned}$$

We rewrite in the equivalent form,

$$\check{J}_q[\mathfrak{d}^k \check{\mathbf{q}}, \mathfrak{d}^k N](\tau_1, \tau_2) = \left| \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^q (r \check{e}_4 \mathfrak{d}^k \check{\mathbf{q}}) (\mathfrak{d}^{k+1} N) \right|. \quad (5.3.38)$$

Using the identity (5.1.9), we have

$$\mathfrak{d}^{k+1} N = \mathfrak{d}^{\leq k+1} N_g + e_3 (\mathfrak{d}^{\leq k+1} r N_g) + \mathfrak{d}^{k+1} N_m[\mathbf{q}].$$

The integral due to $\mathfrak{d}^{\leq k+1} N_g$ is treated as follows

$$\begin{aligned} \check{J}_q[\mathfrak{d}^k \check{\mathbf{q}}, \mathfrak{d}^k N_g](\tau_1, \tau_2) &\lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^q |r \check{e}_4 \mathfrak{d}^k \check{\mathbf{q}}| |\mathfrak{d}^{\leq k+1} N_g| \\ &\lesssim \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q-3} |r \check{e}_4 \mathfrak{d}^k \check{\mathbf{q}}|^2 \right)^{1/2} \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+3} |\mathfrak{d}^{\leq k+1} N_g|^2 \right)^{1/2} \end{aligned}$$

Therefore,

$$\begin{aligned} \check{J}_q^s[\check{\mathbf{q}}, N_g](\tau_1, \tau_2) &\lesssim (B_q^s[\check{\mathbf{q}}](\tau_1, \tau_2))^{1/2} (I_{q+2}^{s+1}[N_g](\tau_1, \tau_2))^{1/2} \\ &\lesssim \delta_1 B_q^s[\check{\mathbf{q}}](\tau_1, \tau_2) + \delta_1^{-1} I_{q+2}^{s+1}[N_g](\tau_1, \tau_2) \end{aligned} \quad (5.3.39)$$

where $\delta_1 > 0$ is chosen sufficiently small so that we can later absorb the term $\delta_1 B_q^s[\mathfrak{q}](\tau_1, \tau_2)$ by the left hand side of our main estimate.

The integral due to $\mathfrak{d}^{k+1} N_m[\mathfrak{q}]$ is treated as follows

$$\begin{aligned}
& \check{J}_q[\mathfrak{d}^k \check{\mathfrak{q}}, \mathfrak{d}^k N_m[\mathfrak{q}]](\tau_1, \tau_2) \\
& \lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^q |r \check{e}_4 \mathfrak{d}^k \check{\mathfrak{q}}| |\mathfrak{d}^{k+1} N_m[\mathfrak{q}]| \\
& \lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+1} |\check{e}_4 \mathfrak{d}^k \check{\mathfrak{q}}| |\mathfrak{d}^{\leq k+2} \mathfrak{q}| |\mathfrak{d}^{\leq k+2} \Gamma_g| \\
& \lesssim \epsilon \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q-1} \tau^{-\frac{1}{2} - \delta_{dec} + 2\delta_0} |\check{e}_4 \mathfrak{d}^k \check{\mathfrak{q}}| |\mathfrak{d}^{\leq k+2} \mathfrak{q}| \\
& \lesssim \epsilon \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^q \tau^{-1-2\delta_{dec}+4\delta_0} |\check{e}_4 \mathfrak{d}^k \check{\mathfrak{q}}|^2 \right)^{\frac{1}{2}} \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q-2} |\mathfrak{d}^{\leq k+2} \mathfrak{q}|^2 \right)^{\frac{1}{2}} \\
& \lesssim \epsilon \left(\sup_{\tau_1 \leq \tau \leq \tau_2} E_q^s[\check{\mathfrak{q}}](\tau) \right)^{\frac{1}{2}} \left(B_{q+1}^{s+1}[\mathfrak{q}](\tau_1, \tau_2) \right)^{\frac{1}{2}}
\end{aligned}$$

where we have used $|\Gamma_g| \lesssim \epsilon r^{-2} \tau^{-1/2 - \delta_{dec} + 2\delta_0}$ and $2\delta_0 < \delta_{dec}$. Since $\delta \leq q+1 \leq 2-\delta$ and $s \leq k_{small} + 29$, we have in view of Theorem 5.3.1

$$B_{q+1}^{s+1}[\mathfrak{q}](\tau_1, \tau_2) \lesssim E_{q+1}^{s+1}[\mathfrak{q}](\tau_1) + I_{q+1}^{s+2}[N_g](\tau_1, \tau_2).$$

We infer

$$\sum_{k \leq s} \check{J}_q[\mathfrak{d}^k \check{\mathfrak{q}}, \mathfrak{d}^k N_m[\mathfrak{q}]](\tau_1, \tau_2) \lesssim \epsilon^2 \sup_{\tau_1 \leq \tau \leq \tau_2} E_q^s[\check{\mathfrak{q}}](\tau) + E_{q+1}^{s+1}[\mathfrak{q}](\tau_1) + I_{q+1}^{s+2}[N_g](\tau_1, \tau_2). \quad (5.3.40)$$

It remains to estimate the integral due to $e_3(\mathfrak{d}^{\leq k+1} r N_g)$. We proceed as in Proposition 5.3.6 by integration by parts, and obtain in particular the following analog of (5.3.27)

$$\check{J}_q[\mathfrak{d}^k \check{\mathfrak{q}}, \mathfrak{d}^k e_3(r N_g)](\tau_1, \tau_2) \lesssim |P^k| + \delta_1 B_q^s[\check{\mathfrak{q}}](\tau_1, \tau_2) + \delta_1^{-1} I_{q+2}^{s+1}[N_g](\tau_1, \tau_2) \quad (5.3.41)$$

where $\delta_1 > 0$ is chosen sufficiently small so that we can later absorb the term $\delta_1 B_q^s[\check{\mathfrak{q}}](\tau_1, \tau_2)$ by the left hand side of our main estimate, and where we have introduced the notation P^k for the analog of L^k in (5.3.28), i.e.⁶

$$P^k := \int_{\mathcal{M}(\tau_1, \tau_2)} r^q e_3 e_4 (r \mathfrak{d}^k \check{\mathfrak{q}}) \mathfrak{d}^{\leq k+1} (r N_g). \quad (5.3.42)$$

⁶Recall that $\check{\mathfrak{q}}$ is localized in $r \geq 4m_0$ so that we don't need in (5.3.42) the cutoff function $\phi_0(r)$ introduced in Proposition 5.3.6.

As in Lemma 5.3.7,

$$e_3 e_4(r\mathfrak{d}^k \check{\mathfrak{q}}) = -\mathfrak{d}^{\leq k}(r\Box_2 \check{\mathfrak{q}}) + r\Delta_2(\mathfrak{d}^{\leq k} \check{\mathfrak{q}}) + r^{-1}\mathfrak{d}^{\leq k+1} \check{\mathfrak{q}}. \quad (5.3.43)$$

We infer

$$\begin{aligned} P^k &= \int_{\mathcal{M}(\tau_1, \tau_2)} r^q e_3 e_4(r\mathfrak{d}^k \check{\mathfrak{q}}) \mathfrak{d}^{\leq k+1}(rN_g) \\ &= - \int_{\mathcal{M}(\tau_1, \tau_2)} r^q \mathfrak{d}^{\leq k}(r\Box_2 \check{\mathfrak{q}}) \mathfrak{d}^{\leq k+1}(rN_g) \\ &\quad + \int_{\mathcal{M}(\tau_1, \tau_2)} r^{q+1} \Delta_2(\mathfrak{d}^{\leq k} \check{\mathfrak{q}}) \mathfrak{d}^{\leq k+1}(rN_g) \\ &\quad + \int_{\mathcal{M}(\tau_1, \tau_2)} r^{q-1} \mathfrak{d}^{\leq k+1} \check{\mathfrak{q}} \mathfrak{d}^{\leq k+1}(rN_g) \\ &= P_1^k + P_2^k + P_3^k. \end{aligned}$$

The last two terms on the right can be treated exactly as the the corresponding terms in the treatment of L^k . This yields to the following analog of (5.3.29) and (5.3.30)

$$\begin{aligned} |P_3^k| &\lesssim \delta_1 B_q^s[\check{\mathfrak{q}}](\tau_1, \tau_2) + \delta_1^{-1} I_{q+2}^{s+1}[N_g](\tau_1, \tau_2), \\ |P_2^k| &\lesssim \delta_1 B_q^s[\check{\mathfrak{q}}](\tau_1, \tau_2) + \delta_1^{-1} I_{q+2}^{s+2}[N_g](\tau_1, \tau_2), \end{aligned} \quad (5.3.44)$$

where $\delta_1 > 0$ is chosen sufficiently small so that we can later absorb the term $\delta_1 B_p^s[\check{\mathfrak{q}}](\tau_1, \tau_2)$ by the left hand side of our main estimate.

It thus only remains to consider the analogous of the term L_1^k , i.e.

$$P_1^k = \int_{\mathcal{M}(\tau_1, \tau_2)} r^q \mathfrak{d}^k(r\Box_2 \check{\mathfrak{q}}) \mathfrak{d}^{\leq k+1}(rN_g).$$

Now, in view of Proposition 10.3.1, \mathfrak{q} verifies, schematically,

$$\Box_2 \check{\mathfrak{q}} = r^{-2} \mathfrak{d}^{\leq 1} \check{\mathfrak{q}} + r^{-2} \mathfrak{d}^{\leq 2} \mathfrak{q} + r \mathfrak{d}^{\leq 1} N$$

so that

$$\begin{aligned} \mathfrak{d}^k(r\Box_2 \check{\mathfrak{q}}) &= r^{-1} \mathfrak{d}^{\leq k+1} \check{\mathfrak{q}} + r^{-1} \mathfrak{d}^{\leq k+2} \mathfrak{q} + r^2 \mathfrak{d}^{\leq k+1} N \\ &= r^{-1} \mathfrak{d}^{\leq k+1} \check{\mathfrak{q}} + r^{-1} \mathfrak{d}^{\leq k+2} \mathfrak{q} + r^2 \mathfrak{d}^{\leq k+1} N_g + r^2 \mathfrak{d}^{\leq k+1} N_m[\mathfrak{q}] + r^2 \mathfrak{d}^{\leq k+1} e_3(rN_g). \end{aligned}$$

We infer the following decomposition of P_1^k

$$\begin{aligned}
P_1^k &= \int_{\mathcal{M}(\tau_1, \tau_2)} r^{q-1} \left(\mathfrak{d}^{\leq k+1} \check{\mathfrak{q}} + \mathfrak{d}^{\leq k+2} \mathfrak{q} \right) \mathfrak{d}^{\leq k+1} (rN_g) \\
&\quad + \int_{\mathcal{M}(\tau_1, \tau_2)} r^{q+2} \mathfrak{d}^{\leq k+1} N_m[\mathfrak{q}] \mathfrak{d}^{\leq k+1} (rN_g) \\
&\quad + \int_{\mathcal{M}(\tau_1, \tau_2)} r^{q+2} \left(\mathfrak{d}^{\leq k+1} N_g + \mathfrak{d}^{\leq k+1} e_3(rN_g) \right) \mathfrak{d}^{\leq k+1} (rN_g) \\
&= P_{11}^k + P_{12}^k + P_{13}^k.
\end{aligned}$$

P_{11}^k is estimated as $\check{J}_q^s[\check{\mathfrak{q}}, N_g](\tau_1, \tau_2)$, see (5.3.39), and hence

$$\begin{aligned}
|P_{11}^k| &\lesssim \left(B_q^s[\check{\mathfrak{q}}](\tau_1, \tau_2) \right)^{1/2} \left(I_{q+2}^{s+1}[N_g](\tau_1, \tau_2) \right)^{1/2} \\
&\lesssim \delta_1 B_q^s[\check{\mathfrak{q}}](\tau_1, \tau_2) + B_{\max(q, \delta)}^s[\mathfrak{q}](\tau_1, \tau_2) + \delta_1^{-1} I_{q+2}^{s+1}[N_g](\tau_1, \tau_2)
\end{aligned}$$

which in view of Theorem 5.3.1 yields

$$|P_{11}^k| \lesssim \delta_1 B_q^s[\check{\mathfrak{q}}](\tau_1, \tau_2) + E_{\max(q, \delta)}^{s+1}[\mathfrak{q}](\tau_1) + \delta_1^{-1} I_{q+2}^{s+1}[N_g](\tau_1, \tau_2). \quad (5.3.45)$$

Next, P_{12}^k is estimated as follows

$$\begin{aligned}
|P_{12}^k| &\lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+3} |\mathfrak{d}^{\leq k+1} N_m[\mathfrak{q}]| |\mathfrak{d}^{\leq k+1} N_g| \\
&\lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+3} |\mathfrak{d}^{\leq k+2} \Gamma_g| |\mathfrak{d}^{\leq k+2} \mathfrak{q}| |\mathfrak{d}^{\leq k+1} N_g| \\
&\lesssim \epsilon \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+1} \tau^{-\frac{1}{2} - \delta_{dec} + 2\delta_0} |\mathfrak{d}^{\leq k+2} \mathfrak{q}| |\mathfrak{d}^{\leq k+1} N_g| \\
&\lesssim \epsilon \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q-2} |\mathfrak{d}^{\leq k+2} \mathfrak{q}|^2 \right)^{\frac{1}{2}} \left(\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+4} \tau^{-1-2\delta_{dec}+4\delta_0} |\mathfrak{d}^{\leq k+1} N_g|^2 \right)^{\frac{1}{2}} \\
&\lesssim \epsilon \left(B_{q+1}^{s+1}[\mathfrak{q}](\tau_1, \tau_2) \right)^{\frac{1}{2}} \left(\sup_{\tau \in [\tau_1, \tau_2]} \int_{\Sigma(\tau)} r^{q+4} |\mathfrak{d}^{\leq k+1} N_g|^2 \right)^{\frac{1}{2}}
\end{aligned}$$

where we have used $|\Gamma_g| \lesssim \epsilon r^{-2} \tau^{-1/2 - \delta_{dec} + 2\delta_0}$ and $2\delta_0 < \delta_{dec}$. We infer

$$|P_{12}^k| \lesssim B_{q+1}^{s+1}[\mathfrak{q}](\tau_1, \tau_2) + I_{q+2}^{s+1}[N_g](\tau_1, \tau_2). \quad (5.3.46)$$

Finally, P_{13}^k is estimated as follows

$$\begin{aligned}
|P_{13}^k| &\lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+3} \left(|\mathfrak{d}^{\leq k+1} N_g| + |\mathfrak{d}^{\leq k+1} e_3(rN_g)| \right) |\mathfrak{d}^{\leq k+1} N_g| \\
&\lesssim \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+3} |\mathfrak{d}^{\leq k+1} N_g|^2 + \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{q+4} |\mathfrak{d}^{\leq k+1} e_3(N_g)| |\mathfrak{d}^{\leq k+1} N_g| \\
&\lesssim I_{q+2}^{s+1}[N_g](\tau_1, \tau_2).
\end{aligned}$$

Together with (5.3.45) and (5.3.46), we infer

$$\begin{aligned}
|P_1^k| &\leq |P_{11}^k| + |P_{12}^k| + |P_{13}^k| \\
&\lesssim \delta_1 B_q^s[\check{\mathfrak{q}}](\tau_1, \tau_2) + E_{\max(q, \delta)}^{s+1}[\mathfrak{q}](\tau_1) + \delta_1^{-1} I_{q+2}^{s+1}[N_g](\tau_1, \tau_2) + B_{q+1}^{s+1}[\mathfrak{q}](\tau_1, \tau_2).
\end{aligned}$$

Together with (5.3.44), we deduce

$$\begin{aligned}
|P^k| &\leq |P_1^k| + |P_2^k| + |P_3^k| \\
&\lesssim \delta_1 B_q^s[\check{\mathfrak{q}}](\tau_1, \tau_2) + E_{\max(q, \delta)}^{s+1}[\mathfrak{q}](\tau_1) + \delta_1^{-1} I_{q+2}^{s+2}[N_g](\tau_1, \tau_2) + B_{q+1}^{s+1}[\mathfrak{q}](\tau_1, \tau_2).
\end{aligned}$$

Together with (5.3.34), (5.3.37), (5.3.39), (5.3.40) and (5.3.41), this concludes the proof of Theorem 5.3.2.

5.4 Decay Estimates

In this section we prove the decay estimates. In particular

- In section 5.4.1, we prove first flux decay estimates for \mathfrak{q} .
- In section 5.4.2, we prove flux decay estimates for $\check{\mathfrak{q}}$.
- In section 5.4.3, we prove Theorem 5.2.1.
- In section 5.4.4, we prove Proposition 5.2.4 on pointwise decay estimates for \mathfrak{q} .
- In section 5.4.5, we prove Proposition 5.2.5 on flux estimates on Σ_* and on improved pointwise estimates for $e_3(\mathfrak{q})$.

The decay estimates rely on the norms (5.1.27) which we recall below.

$$\begin{aligned}\mathcal{E}_{p,d}^s[\psi] &= \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d E_p^s[\psi](\tau), \\ \mathcal{B}_{p,d}^s[\psi] &= \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d \int_{\tau}^{\tau_*} M_{p-1}^s[\psi](\tau) d\tau, \\ \mathcal{F}_{p,d}^s[\psi] &= \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d F_p^s[\psi](\tau), \\ \mathcal{I}_{p,d}^s[N_g] &= \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d I_p^s[N_g](\tau, \tau_*).\end{aligned}$$

5.4.1 First Flux Decay Estimates

The goal of this section is to prove the following flux decay estimates for \mathbf{q} .

Theorem 5.4.1. *Assume \mathbf{q} verifies all the estimates of Theorem 5.3.1. Then the following estimates hold true for all $s \leq k_{small} + 30$ and for all $\delta \leq p \leq 2 - \delta$*

$$\mathcal{E}_{p,2-\delta-p}^{s-[2-\delta-p]}[\mathbf{q}] + \mathcal{B}_{p,2-\delta-p}^{s-[2-\delta-p]}[\mathbf{q}] + \mathcal{F}_{p,2-\delta-p}^{s-[2-\delta-p]}[\mathbf{q}] \lesssim \mathcal{E}_{2-\delta}^s[\mathbf{q}](0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{\delta,2-2\delta}^{s+1}[N_g]. \quad (5.4.1)$$

Here $[x]$ denotes the least integer greater or equal to x .

Proof. We make use of Theorem 5.3.1 according to which we have, for $\delta \leq p \leq 2 - \delta$, and $0 \leq k \leq k_{small} + 30$,

$$E_p^s[\mathbf{q}](\tau_2) + B_p^s[\mathbf{q}](\tau_1, \tau_2) + F_p^s[\mathbf{q}](\tau_1, \tau_2) \lesssim E_p^s[\mathbf{q}](\tau_1) + I_p^{s+1}[N_g](\tau_1, \tau_2)$$

which we write in the form,

$$E_p^s(\tau_2) + \int_{\tau_1}^{\tau_2} M_{p-1}^s(\tau) d\tau \lesssim E_p^s(\tau_1) + I_p^{s+1}[N_g](\tau_1, \tau_2), \quad \delta \leq p \leq 2 - \delta. \quad (5.4.2)$$

In particular,

$$E_{2-\delta}^s(\tau) + \int_{\tau/2}^{\tau} M_{1-\delta}^s(\lambda) d\lambda \lesssim E_{2-\delta}^s(\tau/2) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g].$$

By the mean value theorem there exists $\tau_0 \in [\tau/2, \tau]$ such that,

$$M_{1-\delta}^s(\tau_0) \lesssim \frac{1}{\tau} (E_{2-\delta}^s(\tau/2) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g]).$$

Since⁷

$$E_{1-\delta}^{s-1}(\tau) \lesssim M_{1-\delta}^s(\tau),$$

we deduce,

$$E_{1-\delta}^{s-1}(\tau_0) \lesssim \frac{1}{\tau} (E_{2-\delta}^s(\tau/2) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g]).$$

Moreover, applying (5.4.2) again for $p = 1 - \delta$, we deduce,

$$\begin{aligned} E_{1-\delta}^{s-1}(\tau) + \int_{\tau_0}^{\tau} M_{-\delta}^{s-1}(\lambda)d\lambda &\lesssim E_{1-\delta}^{s-1}(\tau_0) + (1 + \tau)^{-1} \mathcal{I}_{1-\delta,1}^s[N_g] \\ &\lesssim (1 + \tau)^{-1} (E_{2-\delta}^s(\tau/2) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{1-\delta,1}^s[N_g]). \end{aligned}$$

In particular,

$$E_{1-\delta}^{s-1}(\tau) \lesssim (1 + \tau)^{-1} (E_{2-\delta}^s(\tau/2) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{1-\delta,1}^s[N_g]). \quad (5.4.3)$$

Interpolating with

$$E_{2-\delta}^s(\tau) \lesssim E_{2-\delta}^s(\tau/2) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g]$$

by using,

$$E_p^s \lesssim (E_{p_1}^s)^{\frac{p_2-p}{p_2-p_1}} (E_{p_2}^s)^{\frac{p-p_1}{p_2-p_1}}, \quad p_1 \leq p \leq p_2,$$

we deduce

$$E_1^{s-1}(\tau) \lesssim (E_{1-\delta}^{s-1}(\tau))^{1-\delta} (E_{2-\delta}^{s-1}(\tau))^{\delta} \lesssim (1 + \tau)^{-1+\delta} (E_{2-\delta}^s(\tau/2) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{1-\delta,1}^s[N_g]).$$

The same inequality holds for τ replaced by $\tau/2$ i.e.,

$$E_1^{s-1}(\tau/2) \lesssim (1 + \tau)^{-1+\delta} (E_{2-\delta}^s(\tau/4) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{1-\delta,1}^s[N_g]). \quad (5.4.4)$$

We now repeat the procedure starting this time with the inequality (5.4.2) for $p = 1$,

$$\begin{aligned} E_1^{s-1}(\tau) + \int_{\tau/2}^{\tau} M_0^{s-1}(\lambda)d\lambda &\lesssim E_1^{s-1}(\tau/2) + I_1^s[N_g](\tau/2, \tau) \\ &\lesssim E_1^{s-1}(\tau/2) + (1 + \tau)^{-1+\delta} \mathcal{I}_{1,1-\delta}^s[N_g]. \end{aligned}$$

⁷Note that the loss of derivative is due to the degeneracy of the bulk integral in the trapping region.

Thus, in view of (5.4.4),

$$\int_{\tau/2}^{\tau} M_0^{s-1}(\lambda) d\lambda \lesssim (1+\tau)^{-1+\delta} (E_{2-\delta}^s(\tau/4) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{1-\delta,1}^s[N_g] + \mathcal{I}_{1,1-\delta}^s[N_g])$$

or, since

$$E_{2-\delta}^s(\tau/4) \lesssim \mathcal{E}_{2-\delta}^s(0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g],$$

we infer that,

$$\int_{\tau/2}^{\tau} M_0^{s-1}(\lambda) d\lambda \lesssim B(1+\tau)^{-1+\delta}$$

where,

$$B := \mathcal{E}_{2-\delta}^s(0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{1-\delta,1}^s[N_g] + \mathcal{I}_{1,1-\delta}^s[N_g]. \quad (5.4.5)$$

Repeating the mean value argument, we can find $\tau_1 \in [\tau/2, \tau]$ such that,

$$M_0^{s-1}(\tau_1) \lesssim \frac{1}{\tau} \int_{\tau/2}^{\tau} M_0^{s-1}(\lambda) d\lambda \lesssim B(1+\tau)^{-2+\delta}.$$

We now make use of the fact that the energy norm E^{s-1} is comparable with M_0^{s-1} everywhere except in the trapping region where we lose a derivative. Thus

$$E^{s-2}(\tau_1) \lesssim M_0^{s-1}(\tau_1)$$

and therefore,

$$E^{s-2}(\tau_1) \lesssim B(1+\tau)^{-2+\delta}. \quad (5.4.6)$$

We would like now to compare $E^{s-2}(\tau)$ with $E^{s-2}(\tau_1)$ using the usual version of the energy inequality and thus derive a similar estimate for the former. Unfortunately⁸, we don't have a closed energy inequality for E and we therefore have instead to rely on E_δ for which we have the inequality,

$$E_\delta^{s-2}(\tau) \lesssim E_\delta^{s-2}(\tau_1) + I_\delta^{s-1}[N_g](\tau_1, \tau). \quad (5.4.7)$$

We also have in view of (5.4.3)

$$E_{1-\delta}^{s-2}(\tau_1) \lesssim (1+\tau)^{-1} (E_{2-\delta}^s(0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{1-\delta,1}^s[N_g]).$$

⁸The loss of δ is due to the fact that we are on a perturbation of Schwarzschild rather than on Schwarzschild.

Interpolating this last inequality with (5.4.6) we deduce, for $\delta > 0$ sufficiently small

$$\begin{aligned} E_\delta^{s-2}(\tau_1) &\lesssim (E^{s-2}(\tau_1))^{\frac{1-2\delta}{1-\delta}} (E_{1-\delta}^{s-2}(\tau_1))^{\frac{\delta}{1-\delta}} \\ &\lesssim (1+\tau)^{-2+2\delta} (B + \mathcal{E}_{1-\delta}^{s-2}(0) + \mathcal{I}_{1-\delta,0}^{s-1}[N_g]) \\ &\lesssim (1+\tau)^{-2+2\delta} B. \end{aligned}$$

Thus, in view of (5.4.7),

$$E_\delta^{s-2}(\tau) \lesssim E_\delta^{s-2}(\tau_1) + I_\delta^{s-1}[N_g](\tau_1, \tau) \lesssim (1+\tau)^{-2+2\delta} (B + \mathcal{I}_{\delta,2-2\delta}^{s-1}[N_g])$$

i.e.,

$$E_\delta^{s-2}(\tau) \lesssim (1+\tau)^{-2+2\delta} (\mathcal{E}_{2-\delta}^{s-2}(0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{1-\delta,1}^s[N_g] + \mathcal{I}_{1,1-\delta}^s[N_g] + \mathcal{I}_{\delta,2-2\delta}^{s-1}[N_g]).$$

We infer

$$\mathcal{E}_{\delta,2-2\delta}^{s-2} \lesssim (\mathcal{E}_{2-\delta}^{s-2}(0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{1-\delta,1}^s[N_g] + \mathcal{I}_{1,1-\delta}^s[N_g] + \mathcal{I}_{\delta,2-2\delta}^{s-1}[N_g])$$

which can be written in the shorter form (by interpolation of the middle terms),

$$\mathcal{E}_{\delta,2-2\delta}^{s-2} \lesssim \mathcal{E}_{2-\delta}^{s-2}(0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{\delta,2-2\delta}^{s+1}[N_g]. \quad (5.4.8)$$

Also, (5.4.3) yields

$$\begin{aligned} \mathcal{E}_{1-\delta,1}^{s-1} &\lesssim \mathcal{E}_{2-\delta}^s(0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{1-\delta,1}^s[N_g] \\ &\lesssim \mathcal{E}_{2-\delta}^s(0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{\delta,2-2\delta}^{s+1}[N_g]. \end{aligned} \quad (5.4.9)$$

while from Theorem 5.3.1, we have

$$\mathcal{E}_{2-\delta,0}^s \lesssim \mathcal{E}_{2-\delta}^s(0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g]. \quad (5.4.10)$$

Interpolating (5.4.8) and (5.4.9), as well as (5.4.9) and (5.4.10), we infer for all $s \leq k_{small} + 30$ and for all $\delta \leq p \leq 2 - \delta$

$$\mathcal{E}_{p,2-\delta-p}^{s-[2-\delta-p]}[\mathbf{q}] \lesssim \mathcal{E}_{2-\delta}^s[\mathbf{q}](0) + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{\delta,2-2\delta}^{s+1}[N_g]. \quad (5.4.11)$$

Finally, making use of Theorem 5.3.1 between τ and τ_* , we have in particular

$$\begin{aligned} B_p^{s-[2-\delta-p]}[\mathbf{q}](\tau, \tau_*) + F_p^{s-[2-\delta-p]}[\mathbf{q}](\tau, \tau_*) &\lesssim E_p^{s-[2-\delta-p]}[\mathbf{q}](\tau) + I_p^{s+1-[2-\delta-p]}[N_g](\tau, \tau_*) \\ &\lesssim (1+\tau)^{-(2-\delta-p)} \left(\mathcal{E}_{p,2-\delta-p}^{s-[2-\delta-p]}[\mathbf{q}] + \mathcal{I}_{p,2-\delta-p}^{s+1}[N_g] \right) \end{aligned}$$

and hence, we infer for all $s \leq k_{small} + 30$ and for all $\delta \leq p \leq 2 - \delta$

$$\mathcal{B}_{p,2-\delta-p}^{s-[2-\delta-p]}[\mathbf{q}] + \mathcal{F}_{p,2-\delta-p}^{s-[2-\delta-p]}[\mathbf{q}] \lesssim \mathcal{E}_{p,2-\delta-p}^{s-[2-\delta-p]}[\mathbf{q}] + \mathcal{I}_{2-\delta,0}^{s+1}[N_g] + \mathcal{I}_{\delta,2-2\delta}^{s+1}[N_g].$$

Together with (5.4.11), this concludes the proof of Theorem 5.4.1. \square

5.4.2 Flux Decay Estimates for $\check{\mathbf{q}}$

The goal of this section is to prove the following flux decay estimates for $\check{\mathbf{q}}$.

Theorem 5.4.2. *The following estimates hold for all $q_0 - 1 \leq q \leq q_0$, where q_0 is a fixed number $\delta < q_0 \leq 1 - \delta$, and $s \leq k_{small} + 28$*

$$\mathcal{E}_{q,q_0-q}^s[\check{\mathbf{q}}] + \mathcal{B}_{q,q_0-q}^s[\check{\mathbf{q}}] \lesssim \mathcal{E}_{q_0}^s[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+2}[\mathbf{q}](0) + \mathcal{I}_{q_0+2,0}^{s+3}[N_g] + \mathcal{I}_{\delta,2+q_0-\delta}^{s+3}[N_g].$$

Proof. Since $\delta < q_0 \leq 1 - \delta$, according to Theorem 5.3.2, $\check{\mathbf{q}} = f_2 \check{e}_4 \mathbf{q}$ verifies, for any $q_0 - 1 \leq q \leq q_0$ and any $s \leq k_{small} + 29$,

$$E_q^s[\check{\mathbf{q}}](\tau_2) + B_q^s[\check{\mathbf{q}}](\tau_1, \tau_2) \lesssim E_q^s[\check{\mathbf{q}}](\tau_1) + E_{q+1}^{s+1}[\mathbf{q}](\tau_1) + I_{q+2}^{s+2}[N_g](\tau_1, \tau_2).$$

According to the definition of our decay norms above we have,

$$I_{q+2}^{s+2}[N_g](\tau_1, \tau_2) \lesssim (1 + \tau_1)^{q-q_0} \mathcal{I}_{q+2,q_0-q}^{s+2}[N_g]. \quad (5.4.12)$$

Also, according to the definition 5.1.27 for the decay norms for \mathbf{q} we also have

$$E_{q+1}^{s+1}[\mathbf{q}](\tau_1) \lesssim (1 + \tau_1)^{q-q_0} \mathcal{E}_{q+1,q_0-q}^{s+2}[\mathbf{q}].$$

We deduce⁹, for all $q_0 - 1 \leq q \leq q_0$,

$$E_q^s[\check{\mathbf{q}}](\tau_2) + \int_{\tau_1}^{\tau_2} M_q^s[\check{\mathbf{q}}](\tau) \lesssim E_q^s[\check{\mathbf{q}}](\tau_1) + (1 + \tau_1)^{q-q_0} \tilde{\mathcal{E}}_{q,q_0-q}^s \quad (5.4.13)$$

where,

$$\tilde{\mathcal{E}}_{q,q_0-q}^s := \mathcal{E}_{q+1,q_0-q}^{s+1}[\mathbf{q}] + \mathcal{I}_{q+2,q_0-q}^{s+2}[N_g]. \quad (5.4.14)$$

In particular,

$$E_{q_0}^s[\check{\mathbf{q}}](\tau_2) + \int_{\tau_1}^{\tau_2} M_{q_0-1}^s[\check{\mathbf{q}}](\tau) d\tau \lesssim E_{q_0}^s[\check{\mathbf{q}}](\tau_1) + \tilde{\mathcal{E}}_{q_0,0}^s. \quad (5.4.15)$$

By the mean value theorem we deduce that there exists $\tau_0 \in [\tau_1, \tau_2]$ such that,

$$M_{q_0-1}^s[\check{\mathbf{q}}](\tau_0) \lesssim \frac{1}{\tau_2 - \tau_1} \left(E_{q_0}^s[\check{\mathbf{q}}](\tau_1) + \tilde{\mathcal{E}}_{q_0,0}^s \right) \lesssim \frac{1}{\tau_2 - \tau_1} \left(\mathcal{E}_{q_0,0}^s[\check{\mathbf{q}}] + \tilde{\mathcal{E}}_{q_0,0}^s \right).$$

⁹Note that it is important in what follows that the r^q weighted estimates hold also for negative values of q .

Thus also,

$$E_{q_0-1}^s[\check{\mathbf{q}}](\tau_0) \lesssim \frac{1}{\tau_2 - \tau_1} \left(\mathcal{E}_{q_0,0}^s[\check{\mathbf{q}}] + \tilde{\mathcal{E}}_{q_0,0}^s \right). \quad (5.4.16)$$

We now make use of (5.4.13) to compare the quantities $E_q[\check{\mathbf{q}}]$ for negative weights ($q = q_0 - 1$) at different values of τ .

$$E_{q_0-1}^s[\check{\mathbf{q}}](\tau_2) \lesssim E_{q_0-1}^s[\check{\mathbf{q}}](\tau_0) + (1 + \tau_0)^{-1} \tilde{\mathcal{E}}_{q_0-1,1}^s.$$

Combining this with (5.4.16) we deduce,

$$E_{q_0-1}^s[\check{\mathbf{q}}](\tau_2) \lesssim \frac{1}{\tau_2 - \tau_1} \left(\mathcal{E}_{q_0,0}^s[\check{\mathbf{q}}] + \tilde{\mathcal{E}}_{q_0,0}^s \right) + (1 + \tau_0)^{-1} \tilde{\mathcal{E}}_{q_0-1,1}^s.$$

Applying this inequality for $\tau_2 = \tau \leq \tau_*$, $\tau_1 = \frac{1}{2}\tau$, $\tau_0 \in [\tau_1, \tau_2]$ we deduce,

$$E_{q_0-1}^s[\check{\mathbf{q}}](\tau) \lesssim (1 + \tau)^{-1} \left(\mathcal{E}_{q_0,0}^s[\check{\mathbf{q}}] + \tilde{\mathcal{E}}_{q_0,0}^s + \tilde{\mathcal{E}}_{q_0-1,1}^s \right). \quad (5.4.17)$$

We now interpolate this last inequality with the following immediate consequence of (5.4.15)

$$E_{q_0}^s[\check{\mathbf{q}}](\tau) \lesssim \mathcal{E}_{q_0,0}^s[\check{\mathbf{q}}] + \tilde{\mathcal{E}}_{q_0,0}^s$$

to deduce, for all $q_0 - 1 \leq q \leq q_0$,

$$E_q^s[\check{\mathbf{q}}](\tau) \lesssim (1 + \tau)^{q-q_0} \left(\mathcal{E}_{q_0,0}^s[\check{\mathbf{q}}] + \tilde{\mathcal{E}}_{q_0,0}^s + \tilde{\mathcal{E}}_{q_0-1,1}^s \right)$$

i.e.,

$$\mathcal{E}_{q,q_0-q}^s[\check{\mathbf{q}}] \lesssim \mathcal{E}_{q_0,0}^s[\check{\mathbf{q}}] + \tilde{\mathcal{E}}_{q_0,0}^s + \tilde{\mathcal{E}}_{q_0-1,1}^s.$$

In view of the definition of $\tilde{\mathcal{E}}_{q,q_0-q}^s$, this yields for all $q_0 - 1 \leq q \leq q_0$,

$$\mathcal{E}_{q,q_0-q}^s[\check{\mathbf{q}}] \lesssim \mathcal{E}_{q_0,0}^s[\check{\mathbf{q}}] + \mathcal{E}_{q_0+1,0}^{s+1}[\mathbf{q}] + \mathcal{E}_{q_0,1}^{s+1}[\mathbf{q}] + \mathcal{I}_{q_0+2,0}^{s+2}[N_g] + \mathcal{I}_{q_0+1,1}^{s+2}[N_g].$$

On the other hand, we have in view of Theorem 5.3.2,

$$\mathcal{E}_{q_0,0}^s[\check{\mathbf{q}}] \lesssim \mathcal{E}_{q_0}^s[\check{\mathbf{q}}](0) + \mathcal{E}_{q_0+1,0}^{s+1}[\mathbf{q}] + \mathcal{I}_{q_0+2,0}^{s+2}[N_g]$$

and hence

$$\mathcal{E}_{q,q_0-q}^s[\check{\mathbf{q}}] \lesssim \mathcal{E}_{q_0}^s[\check{\mathbf{q}}](0) + \mathcal{E}_{q_0+1,0}^{s+1}[\mathbf{q}] + \mathcal{E}_{q_0,1}^{s+1}[\mathbf{q}] + \mathcal{I}_{q_0+2,0}^{s+2}[N_g] + \mathcal{I}_{q_0+1,1}^{s+2}[N_g].$$

Now, since $\delta < q_0 \leq 1 - \delta$, we have $\delta < q_0 < q_0 + 1 \leq 2 - \delta$ and thus, we may apply Theorem 5.4.1 to obtain for all $q_0 - 1 \leq q \leq q_0$

$$\mathcal{E}_{q+1, q_0-q}^{s+1}[\mathbf{q}] \lesssim \mathcal{E}_{2-\delta}^{s+2}[\mathbf{q}](0) + \mathcal{I}_{2-\delta, 0}^{s+3}[N_g] + \mathcal{I}_{\delta, 2-2\delta}^{s+3}[N_g]. \quad (5.4.18)$$

We thus infer

$$\mathcal{E}_{q, q_0-q}^s[\check{\mathbf{q}}] \lesssim \mathcal{E}_{q_0}^s[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+2}[\mathbf{q}](0) + \mathcal{I}_{q_0+2, 0}^{s+3}[N_g] + \mathcal{I}_{q_0+1, 1}^{s+3}[N_g] + \mathcal{I}_{2-\delta, 0}^{s+3}[N_g] + \mathcal{I}_{\delta, 2-2\delta}^{s+3}[N_g]$$

and hence, for all $q_0 - 1 \leq q \leq q_0$,

$$\mathcal{E}_{q, q_0-q}^s[\check{\mathbf{q}}] \lesssim \mathcal{E}_{q_0}^s[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+2}[\mathbf{q}](0) + \mathcal{I}_{q_0+2, 0}^{s+3}[N_g] + \mathcal{I}_{\delta, 2-2\delta}^{s+3}[N_g]. \quad (5.4.19)$$

Finally, making use of Theorem 5.3.2 between τ and τ_* , we have in particular

$$\begin{aligned} B_q^s[\check{\mathbf{q}}](\tau, \tau_*) &\lesssim E_q^s[\check{\mathbf{q}}](\tau) + E_{q+1}^{s+1}[\mathbf{q}](\tau) + I_{q+2}^{s+2}[N_g](\tau, \tau_*) \\ &\lesssim (1 + \tau)^{-(q_0-q)} \left(\mathcal{E}_{q, q_0-q}^s[\check{\mathbf{q}}] + \mathcal{E}_{q+1, q_0-q}^{s+1}[\mathbf{q}] + \mathcal{I}_{q+2, q_0-q}^{s+2}[N_g] \right) \\ &\lesssim (1 + \tau)^{-(q_0-q)} \left(\mathcal{E}_{q, q_0-q}^s[\check{\mathbf{q}}] + \mathcal{E}_{2-\delta}^{s+2}[\mathbf{q}](0) + \mathcal{I}_{q_0+2, 0}^{s+3}[N_g] + \mathcal{I}_{\delta, 2-2\delta}^{s+3}[N_g] \right) \end{aligned}$$

where we used (5.4.18) in the last inequality. Hence, we infer for all $s \leq k_{small} + 28$ and for all $q_0 - 1 \leq q \leq q_0$

$$\mathcal{B}_{q, q_0-q}^s[\check{\mathbf{q}}] \lesssim \mathcal{E}_{q, q_0-q}^s[\check{\mathbf{q}}] + \mathcal{E}_{2-\delta}^{s+2}[\mathbf{q}](0) + \mathcal{I}_{q_0+2, 0}^{s+3}[N_g] + \mathcal{I}_{\delta, 2-2\delta}^{s+3}[N_g].$$

Together with (5.4.19), this concludes the proof of Theorem 5.4.2. \square

5.4.3 Proof of Theorem 5.2.1

In this section, we prove Theorem 5.2.1 by making use of Theorem 5.4.1 and Theorem 5.4.2. We start with the main estimate of Theorem 5.4.2 with $q = -\delta$ which we write in the form,

$$E_{-\delta}^s[\check{\mathbf{q}}] \lesssim (1 + \tau)^{-q_0-\delta} C_{q_0}^s$$

where,

$$C_{q_0}^s := \mathcal{E}_{q_0}^s[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+2}[\mathbf{q}](0) + \mathcal{I}_{q_0, 0}^{s+3}[N_g] + \mathcal{I}_{\delta, q_0+2-\delta}^{s+3}[N_g].$$

In view of the definition (5.1.21) of $E_{-\delta}^s[\check{\mathbf{q}}]$ and since $\check{\mathbf{q}} = f_2 \check{e}_4 \mathbf{q}$,

$$\int_{\Sigma_{\geq 4m_0}(\tau)} r^{-\delta} (|\check{e}_4 \check{\mathbf{q}}|^2 + r^{-2} |\check{\mathbf{q}}|^2) \lesssim (1 + \tau)^{-q_0-\delta} C_{q_0}^s.$$

Hence,

$$\dot{E}_{2-\delta,4m_0}^s[\mathbf{q}] = \int_{\Sigma_{\geq 4m_0}(\tau)} r^{2-\delta} |\check{e}_4 \mathbf{q}|^2 \lesssim (1 + \tau)^{-q_0-\delta} C_{q_0}^s. \quad (5.4.20)$$

In view of the decay estimates (5.4.1) for \mathbf{q} established in Theorem 5.4.1 we have,

$$\begin{aligned} E^s(\tau) &\lesssim (1 + \tau)^{-2+2\delta} B_{2-\delta}^{2+s}, \\ B_{2-\delta}^{2+s} &:= \mathcal{E}_{2-\delta}^{s+2}[\mathbf{q}](0) + \mathcal{I}_{2-\delta,0}^{s+3}[N_g] + \mathcal{I}_{\delta,2-2\delta}^{s+3}[N_g]. \end{aligned}$$

Thus, the quantity

$$E_{2-\delta}^s = E_{2-\delta}^s[\mathbf{q}](\tau) = \dot{E}_{2-\delta,4m_0}^s[\mathbf{q}] + E^s[\mathbf{q}]$$

verifies,

$$E_{2-\delta}^s \lesssim (1 + \tau)^{-q_0-\delta} (C_{q_0}^s + B_{2-\delta}^{2+s}). \quad (5.4.21)$$

On the other hand, $E_{2-\delta}^s$ verifies (5.4.2) for $p = 2 - \delta$, i.e.

$$E_{2-\delta}^s(\tau_2) + \int_{\tau_1}^{\tau_2} M_{1-\delta}^s(\tau) d\tau \lesssim E_{2-\delta}^s(\tau_1) + I_{2-\delta}^{s+1}[N_g](\tau_1, \tau_2).$$

Since

$$I_{2-\delta}^{s+1}[N_g](\tau_1, \tau_2) \lesssim (1 + \tau_1)^{-q_0-\delta} \mathcal{I}_{2-\delta,q_0+\delta}^{s+1}[N_g],$$

we infer

$$\begin{aligned} E_{2-\delta}^s(\tau) + \int_{\tau/2}^{\tau} M_{1-\delta}^s(\tau') d\tau' &\lesssim E_{2-\delta}^s(\tau/2) + I_{2-\delta}^{s+1}[N_g](\tau/2, \tau) \\ &\lesssim (1 + \tau)^{-q_0-\delta} (C_{q_0}^s + B_{2-\delta}^{2+s} + \mathcal{I}_{2-\delta,q_0+\delta}^{s+1}[N_g]). \end{aligned} \quad (5.4.22)$$

Following the same arguments as in the proof of Theorem 5.4.1 we deduce, for a $\tau_0 \in [\tau/2, \tau]$,

$$E_{1-\delta}^{s-1}(\tau_0) \lesssim (1 + \tau)^{-q_0-1-\delta} (C_{q_0}^s + B_{2-\delta}^{2+s} + \mathcal{I}_{2-\delta,q_0+\delta}^{s+1}[N_g])$$

and since,

$$E_{1-\delta}^s(\tau) \lesssim E_{1-\delta}^s(\tau_0) + I_{1-\delta}^{s+1}(\tau_0, \tau)[N_g],$$

we infer that,

$$E_{1-\delta}^s(\tau) \lesssim (1 + \tau)^{-q_0-1-\delta} (C_{q_0}^{s+1} + B_{2-\delta}^{3+s} + \mathcal{I}_{2-\delta,q_0+\delta}^{s+2}[N_g] + \mathcal{I}_{1-\delta,1+q_0+\delta}^{s+1}[N_g]). \quad (5.4.23)$$

Interpolating with (5.4.21), i.e.

$$E_{2-\delta}^s \lesssim (1+\tau)^{-q_0-\delta} (C_{q_0}^s + B_{2-\delta}^{2+s})$$

we deduce,

$$E_1^s \lesssim (E_{1-\delta}^s)^{1-\delta} (E_{2-\delta}^s)^\delta \lesssim (1+\tau)^{-q_0-1} (C_{q_0}^{s+1} + B_{2-\delta}^{3+s} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+2}[N_g] + \mathcal{I}_{1-\delta, 1+q_0+\delta}^{s+1}[N_g]).$$

Hence,

$$E_1^s \lesssim (1+\tau)^{-q_0-1} (C_{q_0}^{s+1} + B_{2-\delta}^{3+s} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+2} + \mathcal{I}_{1-\delta, 1+q_0+\delta}^{s+1}). \quad (5.4.24)$$

As in the proof of Theorem 5.4.1 we repeat the procedure starting with the inequality (5.4.2) for $p = 1$,

$$\begin{aligned} & E_1^s(\tau) + \int_{\tau/2}^{\tau} M_0^s(\lambda) d\lambda \\ & \lesssim E_1^s(\tau/2) + I_1^{s+1}[N_g](\tau/2, \tau) \\ & \lesssim (1+\tau)^{-q_0-1} (C_{q_0}^{s+1} + B_{2-\delta}^{3+s} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+2}[N_g] + \mathcal{I}_{1-\delta, 1+q_0+\delta}^{s+1}[N_g]) + (1+\tau)^{-1-q_0} \mathcal{I}_{1, 1+q_0}^{s+1}[N_g] \\ & \lesssim (1+\tau)^{-q_0-1} (C_{q_0}^{s+1} + B_{2-\delta}^{3+s} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+2}[N_g] + \mathcal{I}_{1-\delta, 1+q_0+\delta}^{s+1}[N_g] + \mathcal{I}_{1, 1+q_0}^{s+1}[N_g]) \end{aligned}$$

from which we infer that, for a $\tau_0 \in [\tau/2, \tau]$,

$$\begin{aligned} & E^s(\tau_0) \\ & \lesssim (1+\tau)^{-q_0-2} (C_{q_0}^{s+2} + B_{2-\delta}^{s+4} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+3}[N_g] + \mathcal{I}_{1-\delta, 1+q_0+\delta}^{s+2}[N_g] + \mathcal{I}_{1, 1+q_0}^{s+2}[N_g]). \end{aligned} \quad (5.4.25)$$

Interpolating (5.4.23) and (5.4.25) we deduce, for $\delta > 0$ sufficiently small

$$\begin{aligned} E_\delta^s(\tau_0) & \lesssim (E^s(\tau_0))^{\frac{1-2\delta}{1-\delta}} (E_{1-\delta}^s(\tau_0))^{\frac{\delta}{1-\delta}} \\ & \lesssim (1+\tau)^{-2-q_0+\delta} (C_{q_0}^{s+2} + B_{2-\delta}^{s+4} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+3}[N_g] + \mathcal{I}_{1-\delta, 1+q_0+\delta}^{s+2}[N_g] + \mathcal{I}_{1, 1+q_0}^{s+2}[N_g]). \end{aligned}$$

Thus, since we have, as in (5.4.7),

$$E_\delta^s(\tau) \lesssim E_\delta^s(\tau_0) + I_\delta^{s+1}[N_g](\tau_0, \tau),$$

we deduce

$$\begin{aligned} E_\delta^s(\tau) & \lesssim (1+\tau)^{-2-q_0+\delta} (C_{q_0}^{s+2} + B_{2-\delta}^{s+4} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+3}[N_g] + \mathcal{I}_{1-\delta, 1+q_0+\delta}^{s+2}[N_g] + \mathcal{I}_{1, 1+q_0}^{s+2}[N_g]) \\ & \quad + (1+\tau)^{-2-q_0+\delta} \mathcal{I}_{\delta, 2+q_0-\delta}^{s+1}[N_g] \end{aligned}$$

i.e.,

$$\begin{aligned} E_\delta^s(\tau) & \lesssim (1+\tau)^{-2-q_0+\delta} \left(C_{q_0}^{s+2} + B_{2-\delta}^{s+4} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+3}[N_g] + \mathcal{I}_{1-\delta, 1+q_0+\delta}^{s+2}[N_g] \right. \\ & \quad \left. + \mathcal{I}_{1, 1+q_0}^{s+2}[N_g] + \mathcal{I}_{\delta, 2+q_0-\delta}^{s+1}[N_g] \right). \end{aligned}$$

By interpolating the middle terms we write,

$$E_\delta^s(\tau) \lesssim (1 + \tau)^{-2-q_0+\delta} (C_{q_0}^{s+2} + B_{2-\delta}^{s+4} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+3}[N_g] + \mathcal{I}_{\delta, 2+q_0-\delta}^{s+3}[N_g]).$$

We now recall,

$$\begin{aligned} C_{q_0}^s &:= \mathcal{E}_{q_0}^s[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+2}[\mathbf{q}](0) + \mathcal{I}_{q_0+2,0}^{s+3}[N_g] + \mathcal{I}_{\delta, q_0+2-\delta}^{s+3}[N_g] \\ B_{2-\delta}^{2+s} &:= \mathcal{E}_{2-\delta}^{s+2}[\mathbf{q}](0) + \mathcal{I}_{2-\delta,0}^{s+3}[N_g] + \mathcal{I}_{\delta, 2-2\delta}^{s+3}[N_g]. \end{aligned}$$

Hence,

$$\begin{aligned} &C_{q_0}^{s+2} + B_{2-\delta}^{s+4} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+3}[N_g] + \mathcal{I}_{\delta, 2+q_0-\delta}^{s+3}[N_g] \\ &= \mathcal{E}_{q_0}^{s+2}[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+4}[\mathbf{q}](0) + \mathcal{I}_{q_0+2,0}^{s+5}[N_g] + \mathcal{I}_{\delta, q_0+2-\delta}^{s+5}[N_g] \\ &+ \mathcal{E}_{2-\delta}^{s+4}[\mathbf{q}](0) + \mathcal{I}_{2-\delta,0}^{s+5}[N_g] + \mathcal{I}_{\delta, 2-2\delta}^{s+5}[N_g] + \mathcal{I}_{2-\delta, q_0+\delta}^{s+3}[N_g] + \mathcal{I}_{\delta, 2+q_0-\delta}^{s+3}[N_g]. \end{aligned}$$

We deduce,

$$\mathcal{E}_{\delta, 2+q_0-\delta}^s[\mathbf{q}] \lesssim \mathcal{E}_{q_0}^{s+2}[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+4}[\mathbf{q}](0) + \mathcal{I}_{q_0+2,0}^{s+5}[N_g] + \mathcal{I}_{\delta, 2+q_0-\delta}^{s+5}[N_g]. \quad (5.4.26)$$

We can also simplify the right hand side of (5.4.24),

$$\begin{aligned} &C_{q_0}^{s+1} + B_{2-\delta}^{3+s} + \mathcal{I}_{2-\delta, q_0+\delta}^{s+2}[N_g] + \mathcal{I}_{1-\delta, 1+q_0+\delta}^{s+1}[N_g] \\ &\lesssim \mathcal{E}_{q_0}^{s+2}[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+4}[\mathbf{q}](0) + \mathcal{I}_{q_0+2,0}^{s+5}[N_g] + \mathcal{I}_{\delta, 2+q_0-\delta}^{s+5}[N_g]. \end{aligned}$$

Thus (5.4.23) becomes,

$$\mathcal{E}_{1-\delta, 1+q_0+\delta}^s \lesssim \mathcal{E}_{q_0}^{s+2}[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+4}[\mathbf{q}](0) + \mathcal{I}_{q_0+2,0}^{s+5}[N_g] + \mathcal{I}_{\delta, 2+q_0-\delta}^{s+5}[N_g]. \quad (5.4.27)$$

Similarly, (5.4.21) yields

$$\mathcal{E}_{2-\delta, q_0-\delta}^s \lesssim \mathcal{E}_{q_0}^{s+2}[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+4}[\mathbf{q}](0) + \mathcal{I}_{q_0+2,0}^{s+5}[N_g] + \mathcal{I}_{\delta, 2+q_0-\delta}^{s+5}[N_g]. \quad (5.4.28)$$

Interpolating (5.4.26) and (5.4.27), as well as (5.4.27) and (5.4.28), we infer for all $s \leq k_{small} + 25$ and for all $\delta \leq p \leq 2 - \delta$

$$\mathcal{E}_{p, 2+q_0-p}^s[\mathbf{q}] \lesssim \mathcal{E}_{q_0}^{s+2}[\check{\mathbf{q}}](0) + \mathcal{E}_{2-\delta}^{s+4}[\mathbf{q}](0) + \mathcal{I}_{q_0+2,0}^{s+5}[N_g] + \mathcal{I}_{\delta, 2+q_0-\delta}^{s+5}[N_g]. \quad (5.4.29)$$

Finally, making use of Theorem 5.3.1 between τ and τ_* , we have in particular

$$\begin{aligned} B_p^s[\mathbf{q}](\tau, \tau_*) + F_p^s[\mathbf{q}](\tau, \tau_*) &\lesssim E_p^s[\mathbf{q}](\tau) + I_p^{s+1}[N_g](\tau, \tau_*) \\ &\lesssim (1 + \tau)^{-(2+q_0-p)} \left(\mathcal{E}_{p, 2+q_0-p}^s[\mathbf{q}] + I_{p, 2+q_0-p}^{s+1}[N_g] \right) \end{aligned}$$

and hence, we infer for all $s \leq k_{small} + 25$ and for all $\delta \leq p \leq 2 - \delta$

$$\mathcal{B}_{p, 2+q_0-p}^s[\mathbf{q}] + \mathcal{F}_{p, 2+q_0-p}^s[\mathbf{q}] \lesssim \mathcal{E}_{p, 2+q_0-p}^s[\mathbf{q}] + \mathcal{I}_{q_0+2,0}^{s+5}[N_g] + \mathcal{I}_{\delta, 2+q_0-\delta}^{s+5}[N_g].$$

Together with (5.4.29), this concludes the proof of Theorem 5.2.1.

5.4.4 Proof of Proposition 5.2.4

Let χ be a smooth cut-off function vanishing for $r \leq 4m_0$ and equal to 1 for $r \geq 6m_0$. To prove estimate (5.2.6) we consider the identity,

$$\begin{aligned} e_4 \left(\int_{S_r} \chi(\mathbf{q}^{(s)})^2 \right) &= \int_{S_r} \left(e_4(\chi(\mathbf{q}^{(s)})^2) + \kappa \chi(\mathbf{q}^{(s)})^2 \right) \\ &= \int_{S_r} \left(\chi(2\mathbf{q}^{(s)} e_4 \mathbf{q}^{(s)} + 2r^{-1}(\mathbf{q}^{(s)})^2) + \chi'(\mathbf{q}^{(s)})^2 + \chi(\kappa - 2r^{-1})|\mathbf{q}^{(s)}|^2 \right) \\ &= \int_{S_r} \left(2\chi \mathbf{q}^{(s)} \check{e}_4 \mathbf{q}^{(s)} + \chi'(\mathbf{q}^{(s)})^2 + O(r^{-2})|\mathbf{q}^{(s)}|^2 \right). \end{aligned}$$

Integrating between $4m_0$ and r for a fixed $r \geq 6m_0$, we deduce, in view of the definitions of $E[\mathbf{q}^{(s)}](\tau)$ and of $E_p[\mathbf{q}^{(s)}](\tau)$,

$$\begin{aligned} \int_{S_r} |\mathbf{q}^{(s)}|^2 &\lesssim \int_{\Sigma(\tau) \geq 4m_0} |\mathbf{q}^{(s)}| |\check{e}_4 \mathbf{q}^{(s)}| + E[\mathbf{q}^{(s)}](\tau) \\ &\lesssim \left(\int_{\Sigma(\tau) \geq 4m_0} r^{1+\delta} |\check{e}_4 \mathbf{q}^{(s)}|^2 \right)^{1/2} \left(\int_{\Sigma(\tau) \geq 4m_0} r^{-1-\delta} |\mathbf{q}^{(s)}|^2 \right)^{1/2} + E[\mathbf{q}^{(s)}](\tau) \\ &\lesssim (E_{1+\delta}[\mathbf{q}^{(s)}](\tau))^{1/2} (E_{1-\delta}[\mathbf{q}^{(s)}](\tau))^{1/2}. \end{aligned}$$

Clearly, this estimate also holds for $r \leq 6m_0$. Together with the definition (5.1.27) of $\mathcal{E}_{p,d}^s[\mathbf{q}^{(s)}]$, we immediately infer

$$(1 + \tau)^{1+q_0} \int_{S_r} |\mathbf{q}^{(s)}|^2 \lesssim (\mathcal{E}_{1+\delta, 1+q_0-\delta}^s[\mathbf{q}])^{\frac{1}{2}} (\mathcal{E}_{1-\delta, 1+q_0+\delta}^s[\mathbf{q}])^{\frac{1}{2}}$$

which is the desired estimate (5.2.6).

To prove (5.2.7) we start instead with the identity,

$$\begin{aligned} e_4 \left(r^{-1} \int_{S_r} \chi(\mathbf{q}^{(s)})^2 \right) &= \int_{S_r} r^{-1} \left(e_4(\chi(\mathbf{q}^{(s)})^2) + \kappa \chi(\mathbf{q}^{(s)})^2 \right) - \frac{e_4(r)}{r^2} \int_{S_r} \chi(\mathbf{q}^{(s)})^2 \\ &= \int_{S_r} r^{-1} \left(\chi(2\mathbf{q}^{(s)} e_4 \mathbf{q}^{(s)} + r^{-1}(\mathbf{q}^{(s)})^2) + \chi'(\mathbf{q}^{(s)})^2 + \chi(\kappa - 2r^{-1})|\mathbf{q}^{(s)}|^2 \right) \\ &\quad - \frac{e_4(r) - 1}{r^2} \int_{S_r} \chi(\mathbf{q}^{(s)})^2 \\ &= \int_{S_r} \left(2r^{-1} \chi e_4(\mathbf{q}^{(s)}) \mathbf{q}^{(s)} + r^{-1} \chi'(\mathbf{q}^{(s)})^2 + O(r^{-2})|\mathbf{q}^{(s)}|^2 \right). \end{aligned}$$

Integrating between $4m_0$ and r for a fixed $r \geq 6m_0$, we deduce, in view of the definitions of $E[\mathbf{q}^{(s)}](\tau)$ and of $E_p[\mathbf{q}^{(s)}](\tau)$,

$$\begin{aligned} r^{-1} \int_{S_r} |\psi|^2 &\lesssim \int_{\Sigma(\tau) \geq 4m_0} r^{-1} |\mathbf{q}^{(s)}| |e_4(\mathbf{q}^{(s)})| + E[\mathbf{q}^{(s)}](\tau) \\ &\lesssim 2 \left(\int_{\Sigma(\tau) \geq 4m_0} |e_4(\mathbf{q}^{(s)})|^2 \right)^{1/2} \left(\int_{\Sigma(\tau) \geq 4m_0} r^{-2} |\mathbf{q}^{(s)}|^2 \right)^{1/2} + E[\mathbf{q}^{(s)}](\tau) \\ &\lesssim E[\mathbf{q}^{(s)}](\tau) \\ &\lesssim E_\delta[\mathbf{q}^{(s)}](\tau). \end{aligned}$$

Clearly, this estimate also holds for $r \leq 6m_0$. Together with the definition (5.1.27) of $\mathcal{E}_{p,d}^s[\mathbf{q}^{(s)}]$, we immediately infer

$$r^{-1} (1 + \tau)^{2+q_0-\delta} \int_{S_r} |\mathbf{q}^{(s)}|^2 \lesssim \mathcal{E}_{\delta,2+q_0-\delta}^s[\mathbf{q}]$$

which is the desired estimate (5.2.7). This concludes the proof of Proposition 5.2.4.

5.4.5 Proof of Proposition 5.2.5

Recall the following definitions

$$\begin{aligned} F[\psi](\tau_1, \tau_2) &= \int_{\mathcal{A}(\tau_1, \tau_2)} \left(\delta_{\mathcal{H}}^{-1} |e_4 \Psi|^2 + \delta_{\mathcal{H}} |e_3 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2 \right) \\ &\quad + \int_{\Sigma_*(\tau_1, \tau_2)} \left(|e_4 \Psi|^2 + |e_3 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2 \right), \\ \dot{F}_p[\psi](\tau_1, \tau_2) &= \int_{\Sigma_*(\tau_1, \tau_2)} r^p \left(|e_4 \psi|^2 + |\nabla \psi|^2 + r^{-2} |\psi|^2 \right), \\ F_p[\psi](\tau_1, \tau_2) &= F[\psi](\tau_1, \tau_2) + \dot{F}_p[\psi](\tau_1, \tau_2), \\ F^s[\psi](\tau_1, \tau_2) &= \sum_{k \leq s} F[\mathfrak{d}^k \psi](\tau_1, \tau_2), \\ F_p^s[\psi](\tau_1, \tau_2) &= \sum_{k \leq s} F_p[\mathfrak{d}^k \psi](\tau_1, \tau_2), \\ \mathcal{F}_{p,d}^s[\psi] &= \sup_{0 \leq \tau \leq \tau_*} (1 + \tau)^d F_p^s[\psi](\tau, \tau_*). \end{aligned}$$

We deduce

$$F^s[\mathbf{q}](\tau, \tau_*) \leq F_\delta^s[\mathbf{q}](\tau, \tau_*) \leq (1 + \tau)^{-2-q_0+\delta} \mathcal{F}_{\delta,2+q_0-\delta}^s[\mathbf{q}]$$

and hence in particular

$$(1 + \tau)^{2+q_0-\delta} \int_{\Sigma_*(\tau, \tau_*)} \left(|e_3 \mathfrak{d}^{\leq s} \mathfrak{q}|^2 + r^{-2} |\mathfrak{d}^{\leq s} \mathfrak{q}|^2 \right) \lesssim \mathcal{F}_{\delta, 2+q_0-\delta}^s[\mathfrak{q}] \quad (5.4.30)$$

which yields the desired estimate (5.2.8).

Next, we focus on the proof of (5.2.9). We start with the following trace estimate

$$\sup_{\Sigma_*(\tau, \tau_*)} \|e_3 \mathfrak{d}^{\leq s} \mathfrak{q}\|_{L^2(S)} \lesssim \|\nu e_3 \mathfrak{d}^{\leq s} \mathfrak{q}\|_{L^2(\Sigma_*(\tau, \tau_*))} + \|e_3 \mathfrak{d}^{\leq s} \mathfrak{q}\|_{L^2(\Sigma_*(\tau, \tau_*))}$$

where we recall that ν is tangent to Σ_* , orthogonal to e_θ and given by

$$\nu = e_3 + a e_4, \quad -2 \leq a \leq -\frac{1}{2}.$$

We infer

$$\begin{aligned} \sup_{\Sigma_*(\tau, \tau_*)} \|e_3 \mathfrak{d}^{\leq s} \mathfrak{q}\|_{L^2(S)} &\lesssim \|e_3 e_3 \mathfrak{d}^{\leq s} \mathfrak{q}\|_{L^2(\Sigma_*(\tau, \tau_*))} + \|e_4 e_3 \mathfrak{d}^{\leq s} \mathfrak{q}\|_{L^2(\Sigma_*(\tau, \tau_*))} \\ &\quad + \|e_3 \mathfrak{d}^{\leq s} \mathfrak{q}\|_{L^2(\Sigma_*(\tau, \tau_*))} \\ &\lesssim \|e_3 \mathfrak{d}^{\leq s+1} \mathfrak{q}\|_{L^2(\Sigma_*(\tau, \tau_*))} + \|r^{-1} \mathfrak{d}^{\leq s+1} \mathfrak{q}\|_{L^2(\Sigma_*(\tau, \tau_*))} \\ &\quad + \|[e_4, e_3] \mathfrak{d}^{\leq s} \mathfrak{q}\|_{L^2(\Sigma_*(\tau, \tau_*))} \\ &\lesssim \|e_3 \mathfrak{d}^{\leq s+1} \mathfrak{q}\|_{L^2(\Sigma_*(\tau, \tau_*))} + \|r^{-1} \mathfrak{d}^{\leq s+1} \mathfrak{q}\|_{L^2(\Sigma_*(\tau, \tau_*))}. \end{aligned}$$

In view of (5.4.30), we deduce

$$\sup_{\Sigma_*} \left\{ (1 + \tau)^{2+q_0-\delta} \|e_3 \mathfrak{d}^{\leq s} \mathfrak{q}\|_{L^2(S)}^2 \right\} \lesssim \mathcal{F}_{\delta, 2+q_0-\delta}^{s+1}[\mathfrak{q}]. \quad (5.4.31)$$

Next, we extend (5.4.31) to $r \geq 4m_0$. In view of (5.3.24), we have schematically

$$\begin{aligned} e_3 e_4(r \mathfrak{d}^k \mathfrak{q}) &= -\mathfrak{d}^{\leq k}(r \square_2 \mathfrak{q}) + r \Delta_2(\mathfrak{d}^{\leq k} \mathfrak{q}) + r^{-1} \mathfrak{d}^{\leq k+1} \mathfrak{q} \\ &= -\mathfrak{d}^{\leq k}(r \square_2 \mathfrak{q}) + r^{-1} \mathfrak{d}^{\leq k+2} \mathfrak{q}. \end{aligned}$$

Also, we have

$$e_4(re_3(\mathfrak{d}^k \mathfrak{q})) = e_3 e_4(r \mathfrak{d}^k \mathfrak{q}) + [e_4, e_3](r \mathfrak{d}^k \mathfrak{q}) - e_4(e_3(r) \mathfrak{d}^k \mathfrak{q})$$

and hence, we infer schematically

$$e_4(re_3(\mathfrak{d}^k \mathfrak{q})) = -\mathfrak{d}^{\leq k}(r \square_2 \mathfrak{q}) + r^{-1} \mathfrak{d}^{\leq k+2} \mathfrak{q}.$$

Now, recall (5.3.31)

$$|\mathfrak{d}^k(r\Box_2\mathfrak{q})| \lesssim r^{-1}|\mathfrak{d}^{\leq k+1}\mathfrak{q}| + r|\mathfrak{d}^{\leq k}N_g| + r^2|\mathfrak{d}^{\leq k}e_3(N_g)|.$$

We deduce

$$|e_4(re_3(\mathfrak{d}^k\mathfrak{q}))| \lesssim r^{-1}|\mathfrak{d}^{\leq k+2}\mathfrak{q}| + r|\mathfrak{d}^{\leq k}N_g| + r^2|\mathfrak{d}^{\leq k}e_3(N_g)|$$

Now, we have

$$\begin{aligned} e_4\left(r^{-2}\int_S(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))^2\right) &= r^{-2}\int_S\left(2e_4(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))re_3(\mathfrak{d}^{\leq s}\mathfrak{q}) + \kappa(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))^2\right) \\ &\quad - \frac{2e_4(r)}{r}r^{-2}\int_S(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))^2 \\ &= r^{-2}\int_S\left(2e_4(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))re_3(\mathfrak{d}^{\leq s}\mathfrak{q}) + (\kappa - 2r^{-1})(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))^2\right) \\ &\quad - 2\frac{e_4(r) - 1}{r}r^{-2}\int_S(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))^2 \end{aligned}$$

and hence

$$\begin{aligned} \left|e_4\left(r^{-2}\int_S(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))^2\right)\right| &\lesssim r^{-2}\int_S\left\{\left(r^{-1}|\mathfrak{d}^{\leq s+2}\mathfrak{q}| + r|\mathfrak{d}^{\leq s}N_g| + r^2|\mathfrak{d}^{\leq s}e_3(N_g)|\right)|re_3(\mathfrak{d}^{\leq s}\mathfrak{q})| \right. \\ &\quad \left. + r^{-2}(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))^2\right\} \\ &\lesssim r^{-2}\int_S\left(r^{-\frac{1}{2}}|\mathfrak{d}^{\leq s+2}\mathfrak{q}|^2 + r^2|\mathfrak{d}^{\leq s}N_g|^2 + r^4|\mathfrak{d}^{\leq s}e_3(N_g)|^2\right) \\ &\quad + r^{-\frac{7}{2}}\int_S(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))^2 \end{aligned}$$

Together with (5.2.7), this yields

$$\begin{aligned} \left|e_4\left(r^{-2}\int_S(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))^2\right)\right| &\lesssim r^{-2}\int_S\left(r^{\frac{7}{2}}|\mathfrak{d}^{\leq s}N_g|^2 + r^{\frac{11}{2}}|\mathfrak{d}^{\leq s}e_3(N_g)|^2\right) \\ &\quad + r^{-4}\int_S(re_3(\mathfrak{d}^{\leq s}\mathfrak{q}))^2 + r^{-\frac{3}{2}}(1 + \tau)^{-2-q_0+\delta}\mathcal{E}_{\delta,2+q_0-\delta}^{s+2}[\mathfrak{q}]. \end{aligned}$$

Now, recall from (5.2.12) that we have for $s \leq k_{small} + 30$

$$\begin{aligned} |\mathfrak{d}^s N_g| &\lesssim \epsilon^2 r^{-3} \tau^{-1-2\delta_{dec}+2\delta_0} \\ |\mathfrak{d}^s N_g| &\lesssim \epsilon^2 r^{-1} \tau^{-2-2\delta_{dec}+2\delta_0}, \\ |\mathfrak{d}^s e_3(N_g)| &\lesssim \epsilon^2 r^{-3} \tau^{-\frac{3}{2}-2\delta_{dec}+2\delta_0}, \\ |\mathfrak{d}^s e_3(N_g)| &\lesssim \epsilon^2 r^{-\frac{7}{2}-\frac{\delta_B}{2}} \tau^{-1-\delta_{dec}+2\delta_0}. \end{aligned}$$

By interpolation, we infer

$$\begin{aligned} r^{-2} \int_S \left(r^{\frac{7}{2}} |\mathfrak{d}^{\leq s} N_g|^2 + r^{\frac{11}{2}} |\mathfrak{d}^{\leq s} e_3(N_g)|^2 \right) &\lesssim \epsilon^4 r^{-\frac{3}{2}} \tau^{-\frac{5}{2}-4\delta_{dec}+4\delta_0} + \epsilon^4 r^{-1-\frac{\delta_B}{2}} \tau^{-\frac{5}{2}-3\delta_{dec}+4\delta_0} \\ &\lesssim \epsilon_0^2 r^{-1-\frac{\delta_B}{2}} \tau^{-\frac{5}{2}-3\delta_{dec}+4\delta_0} \end{aligned}$$

and hence

$$\begin{aligned} \left| e_4 \left(r^{-2} \int_S (r e_3(\mathfrak{d}^{\leq s} \mathbf{q}))^2 \right) \right| &\lesssim r^{-4} \int_S (r e_3(\mathfrak{d}^{\leq s} \mathbf{q}))^2 \\ &\quad + \epsilon_0^2 r^{-1-\frac{\delta_B}{2}} \tau^{-\frac{5}{2}-3\delta_{dec}+4\delta_0} + r^{-\frac{3}{2}} (1+\tau)^{-2+q_0+\delta} \mathcal{E}_{\delta, 2+q_0-\delta}^{s+2}[\mathbf{q}]. \end{aligned}$$

We integrate from Σ_* . By Gronwall, and in view of (5.4.31), we deduce for $r \geq 4m_0$

$$(1+\tau)^{2+q_0-\delta} \int_{S_r} (e_3 \mathfrak{d}^{\leq s} \mathbf{q})^2 \lesssim \epsilon_0^2 + \mathcal{F}_{\delta, 2+q_0-\delta}^{s+1}[\mathbf{q}] + \mathcal{E}_{\delta, 2+q_0-\delta}^{s+2}[\mathbf{q}].$$

On the other hand, we have by a trace estimate for $r \leq 4m_0$

$$(1+\tau)^{2+q_0-\delta} \int_{S_r} (e_3 \mathfrak{d}^{\leq s} \mathbf{q})^2 \lesssim \mathcal{E}_{0, 2+q_0-\delta}^{s+2}[\mathbf{q}].$$

We finally deduce on \mathcal{M}

$$(1+\tau)^{2+q_0-\delta} \int_{S_r} (e_3 \mathfrak{d}^{\leq s} \mathbf{q})^2 \lesssim \epsilon_0^2 + \mathcal{F}_{\delta, 2+q_0-\delta}^{s+1}[\mathbf{q}] + \mathcal{E}_{\delta, 2+q_0-\delta}^{s+2}[\mathbf{q}]$$

which is the desired estimate (5.2.9). This concludes the proof of Proposition 5.2.5.

Chapter 6

DECAY ESTIMATES FOR α AND $\underline{\alpha}$ (Theorems M2, M3)

In this section, we rely on the decay of \mathfrak{q} to prove the decay estimates for α and $\underline{\alpha}$. More precisely, we rely on the results of Theorem M1 to prove Theorem M2 and M3.

6.1 Proof of Theorem M2

6.1.1 A renormalized frame on ${}^{(ext)}\mathcal{M}$

In Theorem M1, decay estimates are derived for \mathfrak{q} defined with respect to the global frame constructed in Proposition 3.5.5. We have the following control for the Ricci coefficients in that frame.

Lemma 6.1.1. *Consider the global null frame (e_3, e_4, e_θ) constructed in Proposition 3.5.5. Then, the Ricci coefficients satisfy the following estimates*

$$\max_{0 \leq k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}} u^{\frac{1}{2}} \left(r^2 \left| \mathfrak{d}^k \left(\omega + \frac{m}{r^2}, \kappa - \frac{2\Upsilon}{r}, \vartheta, \zeta, \eta, \underline{\eta} \right) \right| + r \left| \mathfrak{d}^k \left(\underline{\xi}, \underline{\omega}, \underline{\kappa} + \frac{2}{r}, \underline{\vartheta} \right) \right|, \right. \\ \left. + \left| \mathfrak{d}^k (e_4(r) - \Upsilon, e_3(r) + 1) \right| \right) \lesssim \epsilon.$$

Proof. This follows immediately from the stronger estimates of Lemma 5.1.1 with the

choice $k_{loss} = 20$. □

6.1.2 A transport equation for α

To recover α from \mathfrak{q} , we derive below a transport equation for α where \mathfrak{q} is on the RHS. We are careful to avoid terms of the type $e_3(\underline{\omega})$ as they are anomalous w.r.t. decay in r . Indeed, they only decay linearly in r^{-1} while all comparable term decay like r^{-2} in r .

Lemma 6.1.2. *We have*

$$\begin{aligned} & \underline{\kappa}^2 e_3 \left\{ e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) - \left(8\underline{\omega} - \frac{2}{\underline{\kappa}} \left(2\phi_1 \underline{\xi} - \frac{1}{2} \vartheta^2 \right) \right) \frac{\alpha}{\underline{\kappa}^2} \right\} \\ = & \frac{\mathfrak{q}}{r^4} + \left\{ 10\underline{\omega} + \frac{4}{\underline{\kappa}} \left(-\phi_1 \underline{\xi} - 2(\eta - 3\zeta) \underline{\xi} + \frac{1}{4} \vartheta^2 \right) \right\} e_3 \alpha \\ & + \left\{ -2\phi_1 \underline{\xi} + \left(6\underline{\kappa} - 24\underline{\omega} + \frac{8}{\underline{\kappa}} \left(2\phi_1 \underline{\xi} + 2(\eta - 3\zeta) \underline{\xi} - \frac{1}{2} \vartheta^2 \right) \right) \underline{\omega} + \frac{1}{2} \vartheta^2 - \frac{4}{\underline{\kappa}} e_3((\eta - 3\zeta) \underline{\xi}) \right. \\ & \left. + \left(16 + \frac{48}{\underline{\kappa}} \underline{\omega} - \frac{24}{\underline{\kappa}^2} \left(2\phi_1 \underline{\xi} + 2(\eta - 3\zeta) \underline{\xi} - \frac{1}{2} \vartheta^2 \right) \right) \zeta \underline{\xi} \right\} \alpha. \end{aligned}$$

Proof. We compute

$$\begin{aligned} e_3 e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) &= e_3 \left(\frac{e_3 \alpha}{\underline{\kappa}^2} - \frac{2e_3(\underline{\kappa}) \alpha}{\underline{\kappa}^3} \right) \\ &= \frac{1}{\underline{\kappa}^2} \left(e_3 e_3 \alpha - 4 \frac{e_3(\underline{\kappa})}{\underline{\kappa}} e_3 \alpha - 2\underline{\kappa}^2 e_3 \left(\frac{e_3(\underline{\kappa})}{\underline{\kappa}^3} \right) \alpha \right). \end{aligned}$$

Now, recall the following null structure equation

$$e_3(\underline{\kappa}) + \frac{1}{2} \underline{\kappa}^2 + 2\underline{\omega} \underline{\kappa} = 2\phi_1 \underline{\xi} + 2(\eta - 3\zeta) \underline{\xi} - \frac{1}{2} \vartheta^2.$$

We infer

$$\frac{e_3(\underline{\kappa})}{\underline{\kappa}} = -\frac{1}{2} \underline{\kappa} - 2\underline{\omega} + \frac{1}{\underline{\kappa}} \left(2\phi_1 \underline{\xi} + 2(\eta - 3\zeta) \underline{\xi} - \frac{1}{2} \vartheta^2 \right)$$

and

$$\begin{aligned}
e_3 \left(\frac{e_3(\underline{\kappa})}{\underline{\kappa}^3} \right) &= e_3 \left(-\frac{1}{2\underline{\kappa}} - 2\frac{\underline{\omega}}{\underline{\kappa}^2} + \frac{1}{\underline{\kappa}^3} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) \\
&= \frac{e_3(\underline{\kappa})}{2\underline{\kappa}^2} + e_3 \left(-2\frac{\underline{\omega}}{\underline{\kappa}^2} + \frac{1}{\underline{\kappa}^3} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) \\
&= -\frac{1}{4} + \frac{1}{2\underline{\kappa}} \left(-2\underline{\omega} + \frac{1}{\underline{\kappa}} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) \\
&\quad + e_3 \left(-2\frac{\underline{\omega}}{\underline{\kappa}^2} + \frac{1}{\underline{\kappa}^3} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right)
\end{aligned}$$

and hence

$$\begin{aligned}
\underline{\kappa}^2 e_3 e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) &= e_3 e_3 \alpha - 4\frac{e_3(\underline{\kappa})}{\underline{\kappa}} e_3 \alpha - 2\underline{\kappa}^2 e_3 \left(\frac{e_3(\underline{\kappa})}{\underline{\kappa}^3} \right) \alpha \\
&= e_3 e_3 \alpha + 2\underline{\kappa} e_3 \alpha + \frac{1}{2}\underline{\kappa}^2 \alpha + \left(8\underline{\omega} - \frac{4}{\underline{\kappa}} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) e_3 \alpha \\
&\quad + \left\{ -\underline{\kappa} \left(-2\underline{\omega} + \frac{1}{\underline{\kappa}} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) \right. \\
&\quad \left. - 2\underline{\kappa}^2 e_3 \left(-2\frac{\underline{\omega}}{\underline{\kappa}^2} + \frac{1}{\underline{\kappa}^3} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) \right\} \alpha.
\end{aligned}$$

Next, recall from section 2.3.3 that \mathfrak{q} is defined with respect to a general null frame as follows

$$\mathfrak{q} = r^4 \left(e_3(e_3(\alpha)) + (2\underline{\kappa} - 6\underline{\omega})e_3(\alpha) + \left(-4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2 \right) \alpha \right).$$

We infer

$$\begin{aligned}
\underline{\kappa}^2 e_3 e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) &= \frac{\mathfrak{q}}{r^4} + \left(14\underline{\omega} - \frac{4}{\underline{\kappa}} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) e_3 \alpha \\
&\quad + \left\{ 4e_3(\underline{\omega}) - 8\underline{\omega}^2 + 10\underline{\omega}\underline{\kappa} - \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right. \\
&\quad \left. - 2\underline{\kappa}^2 e_3 \left(-2\frac{\underline{\omega}}{\underline{\kappa}^2} + \frac{1}{\underline{\kappa}^3} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) \right\} \alpha.
\end{aligned}$$

We rewrite the following terms

$$\begin{aligned}
& \left\{ 4e_3(\underline{\omega}) - 2\underline{\kappa}^2 e_3 \left(-2\frac{\underline{\omega}}{\underline{\kappa}^2} + \frac{1}{\underline{\kappa}^3} \left(2\underline{\phi}_1 \underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) \right) \right\} \alpha \\
&= \underline{\kappa}^2 e_3 \left\{ \left(8\frac{\underline{\omega}}{\underline{\kappa}^2} - \frac{2}{\underline{\kappa}^3} \left(2\underline{\phi}_1 \underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) \right) \alpha \right\} - 4\underline{\kappa}^2 \underline{\omega} e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) + \frac{2}{\underline{\kappa}} \left(2\underline{\phi}_1 \underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) e_3(\alpha) \\
&= \underline{\kappa}^2 e_3 \left\{ \left(8\frac{\underline{\omega}}{\underline{\kappa}^2} - \frac{2}{\underline{\kappa}^3} \left(2\underline{\phi}_1 \underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) \right) \alpha \right\} \\
&\quad + \left(-4\underline{\omega} + \frac{2}{\underline{\kappa}} \left(2\underline{\phi}_1 \underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) \right) e_3 \alpha - 4\underline{\kappa}^2 \underline{\omega} e_3 \left(\frac{1}{\underline{\kappa}^2} \right) \alpha
\end{aligned}$$

so that we obtain

$$\begin{aligned}
& \underline{\kappa}^2 e_3 \left\{ e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) - \left(8\frac{\underline{\omega}}{\underline{\kappa}^2} - \frac{2}{\underline{\kappa}^3} \left(2\underline{\phi}_1 \underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) \right) \alpha \right\} \\
&= \frac{\mathfrak{q}}{r^4} + \left\{ 10\underline{\omega} - \frac{4}{\underline{\kappa}} \left(\underline{\phi}_1 \underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{4}\underline{\vartheta}^2 \right) \right\} e_3 \alpha + \left\{ -8\underline{\omega}^2 + 10\underline{\omega} \underline{\kappa} \right. \\
&\quad \left. - \left(2\underline{\phi}_1 \underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) - 4\underline{\kappa}^2 e_3 \left(\frac{1}{\underline{\kappa}^3} (\eta - 3\zeta)\underline{\xi} \right) - 4\underline{\kappa}^2 \underline{\omega} e_3 \left(\frac{1}{\underline{\kappa}^2} \right) \right\} \alpha
\end{aligned}$$

which we rewrite as

$$\begin{aligned}
& \underline{\kappa}^2 e_3 \left\{ e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) - \left(8\underline{\omega} - \frac{2}{\underline{\kappa}} \left(2\underline{\phi}_1 \underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) \right) \frac{\alpha}{\underline{\kappa}^2} \right\} \\
&= \frac{\mathfrak{q}}{r^4} + \left\{ 10\underline{\omega} - \frac{4}{\underline{\kappa}} \left(\underline{\phi}_1 \underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{4}\underline{\vartheta}^2 \right) \right\} e_3 \alpha + \left\{ -8\underline{\omega}^2 + 10\underline{\omega} \underline{\kappa} \right. \\
&\quad \left. - \left(2\underline{\phi}_1 \underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) - \frac{4}{\underline{\kappa}} e_3((\eta - 3\zeta)\underline{\xi}) + 12\frac{e_3(\underline{\kappa})}{\underline{\kappa}^2} (\eta - 3\zeta)\underline{\xi} + 8\frac{e_3(\underline{\kappa})}{\underline{\kappa}} \underline{\omega} \right\} \alpha.
\end{aligned}$$

Now, recall from above that we have

$$\frac{e_3(\underline{\kappa})}{\underline{\kappa}} = -\frac{1}{2}\underline{\kappa} - 2\underline{\omega} + \frac{1}{\underline{\kappa}} \left(2\underline{\phi}_1 \underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right).$$

We finally deduce

$$\begin{aligned}
& \underline{\kappa}^2 e_3 \left\{ e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) - \left(8\underline{\omega} - \frac{2}{\underline{\kappa}} \left(2\underline{\phi}_1 \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \right) \frac{\alpha}{\underline{\kappa}^2} \right\} \\
= & \frac{\mathfrak{q}}{r^4} + \left\{ 10\underline{\omega} + \frac{4}{\underline{\kappa}} \left(-\underline{\phi}_1 \underline{\xi} - 2(\eta - 3\zeta) \underline{\xi} + \frac{1}{4} \underline{\vartheta}^2 \right) \right\} e_3 \alpha \\
& + \left\{ -2\underline{\phi}_1 \underline{\xi} + \left(6\underline{\kappa} - 24\underline{\omega} + \frac{8}{\underline{\kappa}} \left(2\underline{\phi}_1 \underline{\xi} + 2(\eta - 3\zeta) \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \right) \underline{\omega} + \frac{1}{2} \underline{\vartheta}^2 - \frac{4}{\underline{\kappa}} e_3((\eta - 3\zeta) \underline{\xi}) \right. \\
& \left. + \left(16 + \frac{48}{\underline{\kappa}} \underline{\omega} - \frac{24}{\underline{\kappa}^2} \left(2\underline{\phi}_1 \underline{\xi} + 2(\eta - 3\zeta) \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \right) \zeta \underline{\xi} \right\} \alpha.
\end{aligned}$$

This concludes the proof of the lemma. \square

6.1.3 Estimates for transport equations in e_3

The following lemma will be useful to integrate the transport equations in e_3 .

Lemma 6.1.3. *Let $p \in {}^{(ext)}\mathcal{M}$. Let $\gamma[p]$ the unique integral curve of e_3 starting from a point on \mathcal{C}_1 terminating at p . Then, we have for $l \geq 1$*

$$\int_{\gamma[p]} \frac{1}{r'^{2+l} u'^{\frac{1}{2} + \delta_{extra}} + r'^{1+l} u'^{1 + \delta_{extra}}} \lesssim \frac{1}{r^{1+l} (2r + u)^{\frac{1}{2} + \delta_{extra}} + r^l (2r + u)^{1 + \delta_{extra}}}$$

and

$$\int_{\gamma[p]} \frac{1}{r'^2 u'^{\frac{1}{2} + \delta_{extra}} + r' u'^{1 + \delta_{extra}}} \lesssim \frac{1}{r(2r + u)^{\frac{1}{2} + \delta_{extra}} + \frac{1}{\log(1+u)} (2r + u)^{1 + \delta_{extra}}}$$

where (u, r) correspond to p and (r', u') to a point on $\gamma[p]$, and where the integration along $\gamma[p]$ relies on a parametrization of $\gamma[p]$ normalized with respect to e_3 .

Proof. Note first from the construction of ${}^{(ext)}\mathcal{M}$ that $\gamma[p]$ exists for any $p \in {}^{(ext)}\mathcal{M}$ (i.e. any point p can be joined to \mathcal{C}_1 by an integral curve of e_3), and $\gamma[p]$ is included in ${}^{(ext)}\mathcal{M}$.

Next, recall that the integration along $\gamma[p]$ relies on a parametrization of $\gamma[p]$ normalized with respect to e_3 . To parametrize the integration by u or r , we will thus have to derive an upper bound for the corresponding Jacobian of the change of variable, i.e. for

$$\frac{1}{|e_3(u)|}, \quad \frac{1}{|e_3(r)|}.$$

To this end, note that we have on ${}^{(ext)}\mathcal{M}$

$$e_3(u) = \frac{2}{\zeta\Upsilon} \geq \frac{2 + O(\epsilon)}{\Upsilon} \geq \frac{1}{\Upsilon} \geq 1$$

since $\Upsilon \leq 1$ by definition. Also, we have on ${}^{(ext)}\mathcal{M}$ in view of Lemma 6.1.1

$$\begin{aligned} |e_3(r)| &\geq 1 - |e_3(r) + 1| \\ &= 1 + O(\epsilon) \\ &\geq \frac{1}{2}. \end{aligned}$$

Hence, we have obtained on ${}^{(ext)}\mathcal{M}$

$$\frac{1}{|e_3(u)|} \leq \frac{1}{2}, \quad \frac{1}{|e_3(r)|} \leq 1. \quad (6.1.1)$$

Next, since $e_3(u) > 0$ and $e_3(r) < 0$ in ${}^{(ext)}\mathcal{M}$, we have $r' \geq r$ and $1 \leq u' \leq u$. We start with the proof of the first inequality. We consider two cases

- If $r \geq u$, we have

$$\begin{aligned} \int_{\gamma[p]} \frac{1}{r'^{2+l} u'^{\frac{1}{2} + \delta_{extra}} + r'^{1+l} u'^{1 + \delta_{extra}}} &\leq \frac{1}{r^{2+l}} \int_0^u \frac{1}{|e_3(u')|} \frac{du'}{u'^{\frac{1}{2} + \delta_{extra}}} \\ &\lesssim \frac{u^{\frac{1}{2} - \delta_{extra}}}{r^{2+l}} \\ &\lesssim \frac{1}{r^{1+l} (2r + u)^{\frac{1}{2} + \delta_{extra}} + r^l (2r + u)^{1 + \delta_{extra}}}, \end{aligned}$$

where we used (6.1.1).

- If $r \leq u$, we separate the integral in $r' \geq u$, which coincides with $1 \leq u' \leq u$, and

$r \leq r' \leq u$ and compute

$$\begin{aligned}
& \int_{\gamma[p]} \frac{1}{r'^{2+l} u'^{\frac{1}{2}+\delta_{extra}} + r'^{1+l} u'^{1+\delta_{extra}}} \\
&= \int_0^u \frac{1}{r'^{2+l} u'^{\frac{1}{2}+\delta_{extra}} + r'^{1+l} u'^{1+\delta_{extra}}} \frac{du'}{|e_3(u')|} \\
&\quad + \int_r^u \frac{1}{r'^{2+l} u'^{\frac{1}{2}+\delta_{extra}} + r'^{1+l} u'^{1+\delta_{extra}}} \frac{dr'}{|e_3(r')|} \\
&\lesssim \frac{1}{u^{2+l}} \int_0^u \frac{1}{|e_3(u')| u'^{\frac{1}{2}+\delta_{extra}}} du' \\
&\quad + \min \left(\frac{1}{u^{\frac{1}{2}+\delta_{extra}}} \int_r^u \frac{1}{|e_3(r')| r'^{2+l}} dr', \frac{1}{u^{1+\delta_{extra}}} \int_r^u \frac{1}{|e_3(r')| r'^{1+l}} dr' \right) \\
&\lesssim \frac{1}{u^{\frac{5}{2}+\delta_{extra}}} + \min \left(\frac{1}{u^{\frac{1}{2}+\delta_{extra}}} \frac{1}{r^{1+l}}, \frac{1}{u^{1+\delta_{extra}}} \frac{1}{r^l} \right) \\
&\lesssim \frac{1}{r^{1+l} (2r+u)^{\frac{1}{2}+\delta_{extra}} + r^l (2r+u)^{1+\delta_{extra}}},
\end{aligned}$$

where we used (6.1.1).

This proves the first inequality.

The second inequality is obtained similarly as follows

- If $r \geq u$, we have

$$\begin{aligned}
\int_{\gamma[p]} \frac{1}{r'^2 u'^{\frac{1}{2}+\delta_{extra}} + r' u'^{1+\delta_{extra}}} &\leq \frac{1}{r^2} \int_0^u \frac{1}{|e_3(u')| u'^{\frac{1}{2}+\delta_{extra}}} du' \\
&\lesssim \frac{u^{\frac{1}{2}-\delta_{extra}}}{r^2} \\
&\lesssim \frac{1}{r(2r+u)^{\frac{1}{2}+\delta_{extra}} + (2r+u)^{1+\delta_{extra}}},
\end{aligned}$$

where we used (6.1.1).

- If $r \leq u$, we separate the integral in $r' \geq u$, which coincides with $1 \leq u' \leq u$, and

$r \leq r' \leq u$ and compute

$$\begin{aligned}
& \int_{\gamma[p]} \frac{1}{r'^2 u'^{\frac{1}{2} + \delta_{extra}} + r' u'^{1 + \delta_{extra}}} \\
&= \int_0^u \frac{1}{r'^2 u'^{\frac{1}{2} + \delta_{extra}} + r' u'^{1 + \delta_{extra}}} \frac{du'}{|e_3(u')|} + \int_r^u \frac{1}{r'^2 u'^{\frac{1}{2} + \delta_{extra}} + r' u'^{1 + \delta_{extra}}} \frac{dr'}{|e_3(r')|} \\
&\lesssim \frac{1}{u^3} \int_0^u \frac{1}{|e_3(u')|} \frac{du'}{u'^{\frac{1}{2} + \delta_{extra}}} \\
&\quad + \min \left(\frac{1}{u^{\frac{5}{2} + \delta_{extra}}} \int_r^u \frac{1}{|e_3(r')|} \frac{dr'}{r'^2}, \frac{1}{u^{1 + \delta_{extra}}} \int_r^u \frac{1}{|e_3(r')|} \frac{dr'}{r'} \right) \\
&\lesssim \frac{1}{u^{\frac{5}{2} + \delta_{extra}}} + \min \left(\frac{1}{u^{\frac{1}{2} + \delta_{extra}}} \frac{1}{r}, \frac{1}{u^{1 + \delta_{extra}}} \int_{\frac{r}{u}}^1 \frac{dr''}{r''} \right) \\
&\lesssim \frac{1}{r(2r + u)^{\frac{1}{2} + \delta_{extra}} + \frac{(2r + u)^{1 + \delta_{extra}}}{\log(1 + u)}},
\end{aligned}$$

where we used (6.1.1).

This concludes the proof of the lemma. \square

Corollary 6.1.4. *Let ψ a solution of the following transport equation*

$$e_3(\psi) = h \text{ on } {}^{(ext)}\mathcal{M}.$$

Let also $0 < u_1 \leq u_*$. Then

- If h and ψ satisfy for $l \geq 1$

$$|h| \lesssim \frac{\epsilon_0}{r^{2+l} u^{\frac{1}{2} + \delta_{extra}} + r^{1+l} u^{1 + \delta_{extra}}} \text{ on } {}^{(ext)}\mathcal{M}(u \leq u_1) \text{ and } |\psi| \lesssim \frac{\epsilon_0}{r^{l + \frac{3}{2} + \delta_{extra}}} \text{ on } \mathcal{C}_1,$$

we have

$$\sup_{(ext)\mathcal{M}(u \leq u_1)} \left(r^{1+l} (2r + u)^{\frac{1}{2} + \delta_{extra}} + r^l (2r + u)^{1 + \delta_{extra}} \right) |\psi| \lesssim \epsilon_0.$$

- If h and ψ satisfy

$$|h| \lesssim \frac{\epsilon_0}{r^2 u^{\frac{1}{2} + \delta_{extra}} + r u^{1 + \delta_{extra}}} \text{ on } {}^{(ext)}\mathcal{M}(u \leq u_1) \text{ and } |\psi| \lesssim \frac{\epsilon_0}{r^{\frac{3}{2} + \delta_{extra}}} \text{ on } \mathcal{C}_1,$$

we have

$$\sup_{(ext)\mathcal{M}(u \leq u_1)} \left(r(2r + u)^{\frac{1}{2} + \delta_{extra}} + \frac{(2r + u)^{1 + \delta_{extra}}}{\log(1 + u)} \right) |\psi| \lesssim \epsilon_0.$$

Proof. This follows immediately from Lemma 6.1.3. \square

6.1.4 Decay estimates for α

We start with an estimate for α on \mathcal{C}_1 .

Lemma 6.1.5. *We have*

$$\max_{0 \leq k \leq k_{small} + 22} \sup_{\mathcal{C}_1} r^{\frac{7}{2} + \delta_{extra}} |\mathfrak{d}^k \alpha| + \max_{0 \leq k \leq k_{small} + 21} \sup_{\mathcal{C}_1} r^{\frac{9}{2} + \delta_{extra}} |\mathfrak{d}^k e_3 \alpha| \lesssim \epsilon_0.$$

Proof. Recall that on \mathcal{C}_1 , we have obtained in Theorem M0

$$\max_{0 \leq k \leq k_{large}} \left\{ \sup_{\mathcal{C}_1} \left[r^{\frac{7}{2} + \delta_B} (|\mathfrak{d}^k ({}^{(ext)}\alpha)| + |\mathfrak{d}^k ({}^{(ext)}\beta)|) + r^{\frac{9}{2} + \delta_B} |\mathfrak{d}^{k-1} e_3 ({}^{(ext)}\alpha)| \right] \right. \\ \left. + \sup_{\mathcal{C}_1} \left[r^3 \left| \mathfrak{d}^k \left(({}^{(ext)}\rho + \frac{2m_0}{r^3}) \right) \right| + r^2 |\mathfrak{d}^k ({}^{(ext)}\underline{\beta})| + r |\mathfrak{d}^k ({}^{(ext)}\underline{\alpha})| \right] \right\} \lesssim \epsilon_0.$$

Since we have chosen $\delta_B \geq \delta_{extra}$, we deduce

$$\max_{0 \leq k \leq k_{large}} \sup_{\mathcal{C}_1} \left[r^{\frac{7}{2} + \delta_{extra}} |\mathfrak{d}^k ({}^{(ext)}\alpha)| + r^{\frac{9}{2} + \delta_{extra}} |\mathfrak{d}^{k-1} e_3 ({}^{(ext)}\alpha)| \right] \lesssim \epsilon_0.$$

Next, recall that \mathfrak{q} is defined with respect to the global frame constructed in Proposition 3.5.5. In view of Proposition 3.5.5 and Proposition 3.4.6, and the change of frame formula for α in Proposition 2.3.4, we have

$$\alpha = ({}^{(ext)}\Upsilon)^2 \left(({}^{(ext)}\alpha + 2f ({}^{(ext)}\beta + \frac{3}{2} f^2 ({}^{(ext)}\rho + \text{l.o.t.})) \right) \quad (6.1.2)$$

where f satisfies¹, see (3.4.11),

$$|\mathfrak{d}^k f| \lesssim \frac{\epsilon}{ru^{\frac{1}{2}} + u}, \quad \text{for } k \leq k_{small} + 22 \text{ on } ({}^{(ext)}\mathcal{M}), \\ |\mathfrak{d}^{k-1} e_3 f| \lesssim \frac{\epsilon}{ru} \quad \text{for } k \leq k_{small} + 22 \text{ on } ({}^{(ext)}\mathcal{M}). \quad (6.1.3)$$

We easily infer

$$\max_{0 \leq k \leq k_{small} + 22} \sup_{\mathcal{C}_1} r^{\frac{7}{2} + \delta_{extra}} |\mathfrak{d}^k \alpha| + \max_{0 \leq k \leq k_{small} + 21} \sup_{\mathcal{C}_1} r^{\frac{9}{2} + \delta_{extra}} |\mathfrak{d}^k e_3 \alpha| \lesssim \epsilon_0.$$

This concludes the proof of the lemma. \square

¹Here we use (3.4.11) with $k_{loss} = 20$. Note also that the estimates we claim here for f are slightly weaker than those in (3.4.11).

Next, let $0 < u_1 \leq u_*$. We introduce the following bootstrap assumption for α on $(ext)\mathcal{M}(u \leq u_1)$

$$\max_{0 \leq k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r+u)^{1+\delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) \left(|\mathfrak{d}^k \alpha| + r|\mathfrak{d}^k e_3 \alpha| \right) \leq \epsilon. \quad (6.1.4)$$

The goal of this section will be the following proposition, i.e. the improvement of these bootstrap assumptions.

Proposition 6.1.6. *We have*

$$\max_{0 \leq k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r+u)^{1+\delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) \left(|\mathfrak{d}^k \alpha| + r|\mathfrak{d}^k e_3 \alpha| \right) \lesssim \epsilon_0.$$

Proposition 6.1.6 will be proved at the end of this section.

Based on the bootstrap assumptions (6.1.4), we estimate the RHS of the transport equation for α .

Lemma 6.1.7. *We have*

$$e_3 \left\{ e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) - F_1 \right\} = F_2$$

where F_1 and F_2 satisfy

$$\begin{aligned} & \max_{0 \leq k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(r(2r+u)^{1+\delta_{extra}} + r^2(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) |\mathfrak{d}^k F_1| \\ & + \max_{0 \leq k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(r^2 u^{1+\delta_{extra}} + r^3 u^{\frac{1}{2}+\delta_{extra}} \right) |\mathfrak{d}^k F_2| \lesssim \epsilon_0. \end{aligned}$$

Proof. Recall that we have

$$\begin{aligned} & \underline{\kappa}^2 e_3 \left\{ e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) - \left(8\underline{\omega} - \frac{2}{\underline{\kappa}} \left(2\underline{\phi}_1 \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \right) \frac{\alpha}{\underline{\kappa}^2} \right\} \\ = & \frac{\mathfrak{q}}{r^4} + \left\{ 10\underline{\omega} + \frac{4}{\underline{\kappa}} \left(-\underline{\phi}_1 \underline{\xi} - 2(\eta - 3\zeta) \underline{\xi} + \frac{1}{4} \underline{\vartheta}^2 \right) \right\} e_3 \alpha \\ & + \left\{ -2\underline{\phi}_1 \underline{\xi} + \left(6\underline{\kappa} - 24\underline{\omega} + \frac{8}{\underline{\kappa}} \left(2\underline{\phi}_1 \underline{\xi} + 2(\eta - 3\zeta) \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \right) \underline{\omega} + \frac{1}{2} \underline{\vartheta}^2 - \frac{4}{\underline{\kappa}} e_3((\eta - 3\zeta) \underline{\xi}) \right. \\ & \left. + \left(16 + \frac{48}{\underline{\kappa}} \underline{\omega} - \frac{24}{\underline{\kappa}^2} \left(2\underline{\phi}_1 \underline{\xi} + 2(\eta - 3\zeta) \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \right) \zeta \underline{\xi} \right\} \alpha \end{aligned}$$

which we rewrite as

$$e_3 \left\{ e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) - F_1 \right\} = F_2$$

where F_1 and F_2 are defined by

$$F_1 := \left(8\underline{\omega} - \frac{2}{\underline{\kappa}} \left(2\phi_1\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) \frac{\alpha}{\underline{\kappa}^2}$$

and

$$\begin{aligned} F_2 := & \frac{\mathfrak{q}}{r^4\underline{\kappa}^2} + \frac{1}{\underline{\kappa}^2} \left\{ 10\underline{\omega} + \frac{4}{\underline{\kappa}} \left(-\phi_1\underline{\xi} - 2(\eta - 3\zeta)\underline{\xi} + \frac{1}{4}\vartheta^2 \right) \right\} e_3\alpha \\ & + \frac{1}{\underline{\kappa}^2} \left\{ -2\phi_1\underline{\xi} + \left(6\underline{\kappa} - 24\underline{\omega} + \frac{8}{\underline{\kappa}} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) \underline{\omega} + \frac{1}{2}\vartheta^2 - \frac{4}{\underline{\kappa}} e_3((\eta - 3\zeta)\underline{\xi}) \right. \\ & \left. + \left(16 + \frac{48}{\underline{\kappa}}\underline{\omega} - \frac{24}{\underline{\kappa}^2} \left(2\phi_1\underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\vartheta^2 \right) \right) \zeta\underline{\xi} \right\} \alpha. \end{aligned}$$

In view of the bootstrap assumptions (6.1.4) for α , the estimates of Lemma 6.1.1 for the Ricci coefficients, and using Theorem M2 to estimate \mathfrak{q} , we easily infer

$$\begin{aligned} & \max_{0 \leq k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(r(2r + u)^{1+\delta_{extra}} + r^2(2r + u)^{\frac{1}{2}+\delta_{extra}} \right) |\mathfrak{d}^k F_1| \\ \lesssim & \epsilon \max_{0 \leq k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r + u)^{1+\delta_{extra}}}{\log(1 + u)} + r^3(2r + u)^{\frac{1}{2}+\delta_{extra}} \right) \left(|\mathfrak{d}^k \alpha| + r|\mathfrak{d}^k e_3 \alpha| \right) \\ \lesssim & \epsilon^2 \lesssim \epsilon_0. \end{aligned}$$

and

$$\begin{aligned} & \max_{0 \leq k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(r^2 u^{1+\delta_{extra}} + r^3 u^{\frac{1}{2}+\delta_{extra}} \right) |\mathfrak{d}^k F_2| \\ \lesssim & \max_{0 \leq k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(u^{1+\delta_{extra}} + r u^{\frac{1}{2}+\delta_{extra}} \right) |\mathfrak{d}^k \mathfrak{q}| \\ & + \epsilon \max_{0 \leq k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r + u)^{1+\delta_{extra}}}{\log(1 + u)} + r^3(2r + u)^{\frac{1}{2}+\delta_{extra}} \right) \left(|\mathfrak{d}^k \alpha| + r|\mathfrak{d}^k e_3 \alpha| \right) \\ \lesssim & \epsilon_0 + \epsilon^2 \lesssim \epsilon_0. \end{aligned}$$

This concludes the proof of the lemma. \square

Lemma 6.1.8. For $0 \leq k + j \leq k_{small} + 20$, we have

$$e_3 \left\{ e_3 \mathfrak{P}^k e_4^j \left(\frac{\alpha}{\underline{k}^2} \right) - F_{1, \mathfrak{P}^k, e_4^j} \right\} = F_{2, \mathfrak{P}^k, e_4^j}$$

where

$$\begin{aligned} & \max_{0 \leq l \leq k_{small} + 20 - k} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r(2r + u)^{1 + \delta_{extra}} + r^2(2r + u)^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{D}^l F_{1, \mathfrak{P}^k}| \\ & + \max_{0 \leq l \leq k_{small} + 20 - k} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r^2 u^{1 + \delta_{extra}} + r^3 u^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{D}^l F_{2, \mathfrak{P}^k}| \lesssim \epsilon_0, \end{aligned}$$

and for $j \geq 1$

$$\begin{aligned} & \max_{0 \leq l \leq k_{small} + 20 - k - j} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r^{1+j}(2r + u)^{1 + \delta_{extra}} + r^{2+j}(2r + u)^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{D}^l F_{1, \mathfrak{P}^k, e_4^j}| \\ & + \max_{0 \leq l \leq k_{small} + 20 - k - j} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r^{2+j} u^{1 + \delta_{extra}} + r^{3+j} u^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{D}^l F_{2, \mathfrak{P}^k, e_4^j}| \\ & \lesssim \epsilon_0 + \max_{0 \leq j + k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r + u)^{1 + \delta_{extra}}}{\log(1 + u)} + r^3(2r + u)^{\frac{1}{2} + \delta_{extra}} \right) r \\ & \quad \times \left(|\mathfrak{P}^k (re_4)^{j-1} e_3 \alpha| + |\mathfrak{P}^k (re_4)^{j-2} e_3^2 \alpha| \right). \end{aligned}$$

Proof. Recall from Lemma 6.1.7 that we have

$$e_3 \left\{ e_3 \left(\frac{\alpha}{\underline{k}^2} \right) - F_1 \right\} = F_2$$

where F_1 and F_2 satisfy

$$\begin{aligned} & \max_{0 \leq k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}} \left(r(2r + u)^{1 + \delta_{extra}} + r^2(2r + u)^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{D}^k F_1| \\ & + \max_{0 \leq k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}} \left(r^2 u^{1 + \delta_{extra}} + r^3 u^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{D}^k F_2| \lesssim \epsilon_0. \end{aligned}$$

Differentiating with \mathfrak{P}^k , this yields

$$e_3 \left\{ e_3 \mathfrak{P}^k \left(\frac{\alpha}{\underline{k}^2} \right) + [\mathfrak{P}^k, e_3] \left(\frac{\alpha}{\underline{k}^2} \right) - \mathfrak{P}^k F_1 \right\} = \mathfrak{P}^k F_2 - [\mathfrak{P}^k, e_3] \left\{ e_3 \left(\frac{\alpha}{\underline{k}^2} \right) - F_1 \right\}$$

and hence

$$e_3 \left\{ e_3 \mathfrak{P}^k \left(\frac{\alpha}{\underline{k}^2} \right) - F_{1, \mathfrak{P}^k} \right\} = F_{2, \mathfrak{P}^k}$$

where

$$F_{1,\wp^k} := \wp^k F_1 - [\wp^k, e_3] \left(\frac{\alpha}{\underline{\kappa}^2} \right), \quad F_{2,\wp^k} := \wp^k F_2 - [\wp^k, e_3] \left\{ e_3 \left(\frac{\alpha}{\underline{\kappa}^2} \right) - F_1 \right\}.$$

In view of Lemma 2.2.13, we have schematically

$$\begin{aligned} [\wp, e_4] &= \check{\Gamma}_g \mathfrak{d} + \check{\Gamma}_g + r\beta, \\ [\wp, e_3] &= \check{\Gamma}_b \mathfrak{d} + \check{\Gamma}_b + r\underline{\beta}. \end{aligned}$$

Together with the estimates of Lemma 6.1.1 for the Ricci coefficients and curvature components as well as the bootstrap assumptions (6.1.4) for α on $^{(ext)}\mathcal{M}$, we infer

$$\begin{aligned} &\max_{0 \leq j \leq k_{small} + 20 - k} \sup_{(ext)\mathcal{M}} \left(r(2r + u)^{1 + \delta_{extra}} + r^2(2r + u)^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{d}^j F_{1,\wp^k}| \\ &\quad + \max_{0 \leq j \leq k_{small} + 20 - k} \sup_{(ext)\mathcal{M}} \left(r^2 u^{1 + \delta_{extra}} + r^3 u^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{d}^j F_{2,\wp^k}| \lesssim \epsilon_0. \end{aligned}$$

Next, we consider the case $j \geq 1$. We have the commutator

$$[e_4, e_3] = 2\omega e_3 - 2\underline{\omega} e_4 - 4\zeta e_\theta.$$

In view of the estimates of Lemma 6.1.1 for the Ricci coefficients, and in view of the bootstrap assumptions (6.1.4) for α , we infer after commutation by e_4^j for $0 \leq k + j \leq k_{small} + 20$

$$e_3 \left\{ e_3 \wp^k e_4^j \left(\frac{\alpha}{\underline{\kappa}^2} \right) - F_{1,\wp^k, e_4^j} \right\} = F_{2,\wp^k, e_4^j}$$

where

$$\begin{aligned} &\max_{0 \leq l \leq k_{small} + 20 - k - j} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(r^{1+j} (2r + u)^{1 + \delta_{extra}} + r^{2+j} (2r + u)^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{d}^l F_{1,\wp^k, e_4^j}| \\ &\quad + \max_{0 \leq l \leq k_{small} + 20 - k - j} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(r^{2+j} u^{1 + \delta_{extra}} + r^{3+j} u^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{d}^l F_{2,\wp^k, e_4^j}| \\ &\lesssim \epsilon_0 + \max_{0 \leq j+k \leq k_{small} + 20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(\frac{r^2 (2r + u)^{1 + \delta_{extra}}}{\log(1 + u)} + r^3 (2r + u)^{\frac{1}{2} + \delta_{extra}} \right) r \\ &\quad \times \left(|\wp^k (re_4)^{j-1} e_3 \alpha| + |\wp^k (re_4)^{j-2} e_3^2 \alpha| \right) \lesssim \epsilon_0. \end{aligned}$$

This concludes the proof of the lemma. \square

We are now ready to prove Proposition 6.1.6.

Step 1. For $0 \leq k \leq k_{small} + 20$, recall from the above lemma with $j = 0$ that we have

$$e_3 \left\{ e_3 \mathfrak{P}^k \left(\frac{\alpha}{\underline{\kappa}^2} \right) - F_{1, \mathfrak{P}^k} \right\} = F_{2, \mathfrak{P}^k}$$

where

$$\max_{0 \leq j \leq k_{small} + 20 - k} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r^2 u^{1 + \delta_{extra}} + r^3 u^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{D}^j F_{2, \mathfrak{P}^k}| \lesssim \epsilon_0.$$

Also, we have in view of Lemma 6.1.5

$$\max_{0 \leq k \leq k_{large} - 4} \sup_{C_1} r^{\frac{5}{2} + \delta_{extra}} \left| e_3 \mathfrak{P}^k \left(\frac{\alpha}{\underline{\kappa}^2} \right) - F_{1, \mathfrak{P}^k} \right| \lesssim \epsilon_0.$$

In view of Corollary 6.1.4, we immediately infer for any $0 \leq k \leq k_{small} + 20$

$$\max_{0 \leq k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r^2 (2r + u)^{\frac{1}{2} + \delta_{extra}} + r (2r + u)^{1 + \delta_{extra}} \right) \left| e_3 \mathfrak{P}^k \left(\frac{\alpha}{\underline{\kappa}^2} \right) - F_{1, \mathfrak{P}^k} \right| \lesssim \epsilon_0.$$

Since we have from the above lemma that

$$\max_{0 \leq j \leq k_{small} + 20 - k} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r (2r + u)^{1 + \delta_{extra}} + r^2 (2r + u)^{\frac{1}{2} + \delta_{extra}} \right) |\mathfrak{D}^j F_{1, \mathfrak{P}^k}| \lesssim \epsilon_0,$$

we deduce that we have for any $0 \leq k \leq k_{small} + 20$

$$\max_{0 \leq k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r^2 (2r + u)^{\frac{1}{2} + \delta_{extra}} + r (2r + u)^{1 + \delta_{extra}} \right) \left| e_3 \mathfrak{P}^k \left(\frac{\alpha}{\underline{\kappa}^2} \right) \right| \lesssim \epsilon_0. \quad (6.1.5)$$

Step 2. Next, note that we have in view of Lemma 6.1.5

$$\max_{0 \leq k \leq k_{large} - 3} \sup_{C_1} r^{\frac{3}{2} + \delta_{extra}} \left| \mathfrak{P}^k \left(\frac{\alpha}{\underline{\kappa}^2} \right) \right| \lesssim \epsilon_0.$$

Together with the transport equation (6.1.5), and in view of Corollary 6.1.4, we infer

$$\max_{0 \leq k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r (2r + u)^{\frac{1}{2} + \delta_{extra}} + \frac{(2r + u)^{1 + \delta_{extra}}}{\log(1 + u)} \right) \left| \mathfrak{P}^k \left(\frac{\alpha}{\underline{\kappa}^2} \right) \right| \lesssim \epsilon_0.$$

In view of the control of $\underline{\kappa}$ provided by Lemma 6.1.1, we easily deduce

$$\max_{0 \leq k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r^3 (2r + u)^{\frac{1}{2} + \delta_{extra}} + \frac{r^2 (2r + u)^{1 + \delta_{extra}}}{\log(1 + u)} \right) |\mathfrak{P}^k \alpha| \lesssim \epsilon_0.$$

Together with (6.1.5), we infer

$$\max_{0 \leq k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r^3(2r+u)^{\frac{1}{2} + \delta_{extra}} + \frac{r^2(2r+u)^{1 + \delta_{extra}}}{\log(1+u)} \right) \left(|\mathfrak{P}^k \alpha| + r |\mathfrak{P}^k e_3 \alpha| \right) \lesssim \epsilon_0.$$

Step 3. Next, recall from section 2.3.3 that \mathfrak{q} is defined with respect to a general null frame as follows

$$\mathfrak{q} = r^4 \left(e_3(e_3(\alpha)) + (2\underline{\kappa} - 6\underline{\omega})e_3(\alpha) + \left(-4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2 \right) \alpha \right).$$

We infer

$$e_3(e_3(\alpha)) = \frac{\mathfrak{q}}{r^4} - (2\underline{\kappa} - 6\underline{\omega})e_3(\alpha) - \left(-4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2 \right) \alpha.$$

Together with the above estimate for α and $e_3\alpha$, we infer by iteration

$$\begin{aligned} \max_{0 \leq k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} & \left(\frac{r^2(2r+u)^{1 + \delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2} + \delta_{extra}} \right) \\ & \times \left(|(\mathfrak{P}, e_3)^k \alpha| + r |(\mathfrak{P}, e_3)^k e_3 \alpha| \right) \lesssim \epsilon_0. \end{aligned}$$

Step 4. Arguing as for Step 1, but with $j \geq 1$, we infer the following analog of (6.1.5)

$$\begin{aligned} & \max_{0 \leq j+k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r^2(2r+u)^{\frac{1}{2} + \delta_{extra}} + r(2r+u)^{1 + \delta_{extra}} \right) \left| e_3 \mathfrak{P}^k (re_4)^j \left(\frac{\alpha}{\underline{\kappa}^2} \right) \right| \\ \lesssim & \epsilon_0 + \max_{0 \leq j+k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r+u)^{1 + \delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2} + \delta_{extra}} \right) r \\ & \times \left(|\mathfrak{P}^k (re_4)^{j-1} e_3 \alpha| + |\mathfrak{P}^k (re_4)^{j-2} e_3^2 \alpha| \right). \end{aligned}$$

Step 5. Arguing as for Step 2, but with $j \geq 1$, we infer the following analog of the last estimate of Step 2

$$\begin{aligned} & \max_{0 \leq j \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(r^3(2r+u)^{\frac{1}{2} + \delta_{extra}} + \frac{r^2(2r+u)^{1 + \delta_{extra}}}{\log(1+u)} \right) \left(|\mathfrak{P}^k (re_4)^j \alpha| + r |\mathfrak{P}^k (re_4)^j e_3 \alpha| \right) \\ \lesssim & \epsilon_0 + \max_{0 \leq j+k \leq k_{small} + 20} \sup_{(ext) \mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r+u)^{1 + \delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2} + \delta_{extra}} \right) r \\ & \times \left(|\mathfrak{P}^k (re_4)^{j-1} e_3 \alpha| + |\mathfrak{P}^k (re_4)^{j-2} e_3^2 \alpha| \right). \end{aligned}$$

Step 6. Arguing as for Step 3, but with $j \geq 1$, we infer the following analog of the last estimate of Step 3

$$\begin{aligned} & \max_{0 \leq j+k \leq k_{small}+20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r+u)^{1+\delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) \\ & \times \left(|(\not\partial, e_3)^k(re_4)^j \alpha| + r|(\not\partial, e_3)^k(re_4)^j e_3 \alpha| \right) \\ \lesssim & \epsilon_0 + \max_{0 \leq j+k \leq k_{small}+20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r+u)^{1+\delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) r \\ & \times \left(|(\not\partial, e_3)^k(re_4)^{j-1} e_3 \alpha| + r|(\not\partial, e_3)^k(re_4)^{j-2} e_3^2 \alpha| \right). \end{aligned}$$

Step 7. Arguing by iteration on j , noticing that the last estimate of Step 3 corresponds to desired estimate for $j = 0$, and in view of the estimate derived in Step 6, we finally obtain

$$\begin{aligned} & \max_{0 \leq j+k \leq k_{small}+20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r+u)^{1+\delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) \\ & \times \left(|(\not\partial, e_3)^k(re_4)^j \alpha| + r|(\not\partial, e_3)^k(re_4)^j e_3 \alpha| \right) \lesssim \epsilon_0 \end{aligned}$$

and hence

$$\max_{0 \leq k \leq k_{small}+20} \sup_{(ext)\mathcal{M}(u \leq u_1)} \left(\frac{r^2(2r+u)^{1+\delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) \left(|\not\partial^k \alpha| + r|\not\partial^k e_3 \alpha| \right) \lesssim \epsilon_0.$$

This concludes the proof of Proposition 6.1.6.

6.1.5 End of the proof of Theorem M2

First, note in view of the estimates for α on \mathcal{C}_1 provided by Lemma 6.1.5 that the bootstrap assumptions (6.1.4) for α hold by continuity for some sufficiently small $u_1 > 0$. Then, we may in view of Proposition 6.1.6 choose $u_1 = u_*$. We deduce therefore

$$\max_{0 \leq k \leq k_{small}+20} \sup_{(ext)\mathcal{M}} \left(\frac{r^2(2r+u)^{1+\delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) \left(|\not\partial^k \alpha| + r|\not\partial^k e_3 \alpha| \right) \lesssim \epsilon_0.$$

Next, recall from (6.1.2) and (6.1.3) that we have

$$\alpha = ({}^{(ext)}\Upsilon)^2 \left(({}^{(ext)}\alpha + 2f^{(ext)}\beta + \frac{3}{2}f^2({}^{(ext)}\rho + \text{l.o.t.}) \right)$$

where f satisfies

$$\begin{aligned} |\mathfrak{d}^k f| &\lesssim \frac{\epsilon}{ru^{\frac{1}{2}} + u}, \quad \text{for } k \leq k_{small} + 22 \text{ on } {}^{(ext)}\mathcal{M}, \\ |\mathfrak{d}^{k-1} e_3 f| &\lesssim \frac{\epsilon}{ru} \quad \text{for } k \leq k_{small} + 22 \text{ on } {}^{(ext)}\mathcal{M}. \end{aligned}$$

Together with bootstrap assumptions for ${}^{(ext)}\beta$ and ${}^{(ext)}\rho$, we easily infer

$$\max_{0 \leq k \leq k_{small} + 20} \sup_{{}^{(ext)}\mathcal{M}} \left(\frac{r^2(2r+u)^{1+\delta_{extra}}}{\log(1+u)} + r^3(2r+u)^{\frac{1}{2}+\delta_{extra}} \right) \left(|\mathfrak{d}^k {}^{(ext)}\alpha| + r|\mathfrak{d}^k e_3 {}^{(ext)}\alpha| \right) \lesssim \epsilon_0.$$

This concludes the proof of Theorem M2.

6.2 Proof of Theorem M3

Theorem M3 contains decay estimates for $\underline{\alpha}$ in ${}^{(int)}\mathcal{M}$ and on Σ_* . We first proceed with the estimate on ${}^{(int)}\mathcal{M}$ before moving to ${}^{(ext)}\mathcal{M}$.

6.2.1 Estimate for $\underline{\alpha}$ in ${}^{(int)}\mathcal{M}$

Recall that \mathfrak{q} , controlled in Theorem M1, is defined with respect to the global frame of Proposition 3.5.5. Recall also that we may choose the global null frame to coincide with the ingoing geodesic null frame of ${}^{(int)}\mathcal{M}$ in ${}^{(int)}\mathcal{M}$ (see property (b) in Proposition 3.5.5 together with property (d) ii. in Proposition 3.5.2). Thus, in this section, as we only work on ${}^{(int)}\mathcal{M}$, the null frame (e_4, e_3, e_θ) denotes both the frame of ${}^{(int)}\mathcal{M}$ and the global frame with respect to which \mathfrak{q} is defined. We start with the following definition.

Definition 6.2.1. *In ${}^{(int)}\mathcal{M}$, we define with respect to the ingoing geodesic frame of ${}^{(int)}\mathcal{M}$*

$$\tilde{T} := e_4 - \frac{1}{\bar{\kappa}} (\bar{\kappa} + A) e_3. \quad (6.2.1)$$

The estimate for $\underline{\alpha}$ in ${}^{(int)}\mathcal{M}$ relies on the following proposition.

Proposition 6.2.2. *Let $0 \leq k \leq k_{small} + 17$. Then, $\underline{\alpha}$ satisfies in ${}^{(int)}\mathcal{M}$*

$$6m\tilde{T}(\mathfrak{d}^k \underline{\alpha}) + r^4 \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2(\mathfrak{d}^k \underline{\alpha}) = F_k$$

where F_k satisfies

$$\max_{0 \leq k \leq k_{small} + 17} \int_{(int)\mathcal{M}} \underline{u}^{2+2\delta_{dec}} |\mathfrak{d}^{\leq 1} F_k|^2 \lesssim \epsilon_0^2.$$

Remark 6.2.3. In view of the definition of \tilde{T} , we have

$$\tilde{T}(r) = e_4(r) - \frac{1}{\bar{\kappa}} (\bar{\kappa} + A) e_3(r) = 0$$

so that \tilde{T} is tangent to the hypersurfaces of constant r . In particular, (\tilde{T}, e_θ) spans the tangent space of hypersurfaces of constant r . Therefore, in view of Proposition 6.2.2, $\underline{\alpha}$ and its derivatives satisfy on each hyper surface of constant r in $(int)\mathcal{M}$, i.e. on $\{r = r_0\}$ for $2m_0(1 - \delta_{\mathcal{H}}) \leq r \leq r_{\mathcal{T}}$, a forward parabolic equation. Furthermore, since we have $\tilde{T}(\underline{u}) = 2/\underline{\varsigma} = 2 + O(\epsilon)$, \underline{u} plays the role of time in this forward parabolic equation.

We also derive estimates for the control of the parabolic equation appearing in the statement of Proposition 6.2.2.

Lemma 6.2.4. Let f and h reduced 2-scalars such that

$$\left(6m\tilde{T} + r^4 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2\right) f = h.$$

Then, for any real number $n \geq 0$ and any r_0 such that $2m_0(1 - \delta_{\mathcal{H}}) \leq r_0 \leq r_{\mathcal{T}}$, we have

$$\begin{aligned} \sup_{1 \leq \underline{u} \leq u_*} \int_{S(r=r_0, \underline{u})} (1 + \underline{u}^n) f^2 &\lesssim_n \int_{S(r=r_0, 1)} f^2 + \epsilon^2 \int_1^{u_*} \int_{S(r=r_0, \underline{u})} (1 + \underline{u}^{n-2}) (\mathfrak{d}f)^2 \\ &+ \int_{(int)\mathcal{M}} (1 + \underline{u}^n) (\mathfrak{d}^{\leq 1} h)^2. \end{aligned}$$

We are now in position to control $\underline{\alpha}$ in $(int)\mathcal{M}$. Recall from Proposition 6.2.2 that $\underline{\alpha}$ satisfies in $(int)\mathcal{M}$ for $0 \leq k \leq k_{small} + 17$

$$6m\tilde{T}(\mathfrak{d}^k \underline{\alpha}) + r^4 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2(\mathfrak{d}^k \underline{\alpha}) = F_k.$$

Applying Lemma 6.2.4 with $n = 2 + 2\delta_{dec}$, $f = \mathfrak{d}^k \underline{\alpha}$ and $h = F_k$, we infer for any r_0 such that $2m_0(1 - \delta_{\mathcal{H}}) \leq r_0 \leq r_{\mathcal{T}}$

$$\begin{aligned} \sup_{1 \leq \underline{u} \leq u_*} \int_{S(r=r_0, \underline{u})} (1 + \underline{u}^{2+2\delta_{dec}}) (\mathfrak{d}^k \underline{\alpha})^2 &\lesssim \int_{S(r=r_0, 1)} (\mathfrak{d}^k \underline{\alpha})^2 + \epsilon^2 \int_1^{u_*} \int_{S(r=r_0, \underline{u})} (1 + \underline{u}^{2\delta_{dec}}) (\mathfrak{d}^{k+1} \underline{\alpha})^2 \\ &+ \int_{(int)\mathcal{M}} (1 + \underline{u}^{2+2\delta_{dec}}) (\mathfrak{d}^{\leq 1} F_k)^2. \end{aligned}$$

Together with the bounds for $\underline{\alpha}$ on $\underline{\mathcal{C}}_1$ provided by Theorem M0, the bootstrap assumptions on decay and energy for $\underline{\alpha}$ in ${}^{(int)}\mathcal{M}$, and the bound for F_k provided by Proposition 6.2.2, we infer for $0 \leq k \leq k_{small} + 17$ in ${}^{(int)}\mathcal{M}$

$$\sup_{1 \leq \underline{u} \leq u_*} \int_{S(r=r_0, \underline{u})} (1 + \underline{u}^{2+2\delta_{dec}}) (\mathfrak{D}^k \underline{\alpha})^2 \lesssim \epsilon_0^2.$$

In particular, we have obtained

$$\max_{0 \leq k \leq k_{small} + 17} \sup_{{}^{(int)}\mathcal{M}} \underline{u}^{1+\delta_{dec}} \|\mathfrak{D}^k \underline{\alpha}\|_{L^2(S)} \lesssim \epsilon_0.$$

Using the Sobolev embedding on 2-surface and the fact that r is bounded on ${}^{(int)}\mathcal{M}$, we infer

$$\max_{0 \leq k \leq k_{small} + 15} \sup_{{}^{(int)}\mathcal{M}} \underline{u}^{1+\delta_{dec}} |\mathfrak{D}^k \underline{\alpha}| \lesssim \epsilon_0$$

and hence

$${}^{(int)}\mathfrak{D}_{k_{small}+15}[\underline{\alpha}] \lesssim \epsilon_0 \tag{6.2.2}$$

which is the desired estimate for $\underline{\alpha}$ in ${}^{(int)}\mathcal{M}$.

The proof of Proposition 6.2.2 will be given in section 6.2.3, and to the proof of Lemma 6.2.4 which will be given in section 6.2.4. But first, we conclude in the next section the proof of Theorem M3 by controlling $\underline{\alpha}$ on Σ_* .

6.2.2 Estimate for $\underline{\alpha}$ on Σ_*

Recall that \mathfrak{q} , controlled in Theorem M1, is defined with respect to the global frame of Proposition 3.5.5. We will first control $\underline{\alpha}$ in this frame, before coming back to ${}^{(ext)}\mathcal{M}$ at the end of the argument. We start with the following definition.

Definition 6.2.5. *In Σ_* , we define, with respect to the the global frame of Proposition 3.5.5,*

$$\tilde{\nu} := e_3 + ae_4, \tag{6.2.3}$$

where the scalar function a is uniquely defined so that $\tilde{\nu}$ is tangent to Σ_* .

The estimate for $\underline{\alpha}$ on Σ_* relies on the following proposition.

Proposition 6.2.6. *Let $0 \leq k \leq k_{small} + 18$. Then, $\underline{\alpha}$ satisfies on Σ_**

$$6m\tilde{\nu}(\mathfrak{d}^k \underline{\alpha}) + r^4 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2(\mathfrak{d}^k \underline{\alpha}) = F_k$$

where F_k satisfies

$$\max_{0 \leq k \leq k_{small} + 18} \int_{\Sigma_*} u^{2+2\delta_{dec}} |F_k|^2 \lesssim \epsilon_0^2.$$

Remark 6.2.7. *Since $\tilde{\nu}$ is tangent to Σ_* , and since $(\tilde{\nu}, e_\theta)$ spans the tangent space of Σ_* , in view of Proposition 6.2.6, $\underline{\alpha}$ and its derivatives satisfy on Σ_* a forward parabolic equation. Furthermore, since we have $\tilde{\nu}(u) = 2 + O(\epsilon)$, u plays the role of time in this forward parabolic equation.*

We also derive estimates for the control of the parabolic equation appearing in the statement of Proposition 6.2.2.

Lemma 6.2.8. *Let f and h reduced 2-scalars such that*

$$\left(6m\tilde{\nu} + r^4 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2\right) f = h.$$

Then, for any real number $n \geq 0$, we have

$$\int_{\Sigma_*} (1 + u^n) f^2 \lesssim_n \int_{\Sigma_* \cap \mathcal{C}_1} f^2 + \epsilon^2 \int_{\Sigma_*} (1 + u^{n-2}) (\mathfrak{d}f)^2 + \int_{\Sigma_*} (1 + u^n) h^2.$$

We are now in position to control $\underline{\alpha}$ on Σ_* . Recall from Proposition 6.2.6 that $\underline{\alpha}$ satisfies in Σ_* for $0 \leq k \leq k_{small} + 18$

$$6m\tilde{\nu}(\mathfrak{d}^k \underline{\alpha}) + r^4 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2(\mathfrak{d}^k \underline{\alpha}) = F_k.$$

Applying Lemma 6.2.8 with $n = 2 + 2\delta_{dec}$, $f = \mathfrak{d}^k \underline{\alpha}$ and $h = F_k$, we infer

$$\begin{aligned} \int_{\Sigma_*} (1 + u^{2+2\delta_{dec}}) (\mathfrak{d}^k \underline{\alpha})^2 &\lesssim \int_{\Sigma_* \cap \mathcal{C}_1} (\mathfrak{d}^k \underline{\alpha})^2 + \epsilon^2 \int_{\Sigma_*} (1 + u^{2\delta_{dec}}) (\mathfrak{d}^{k+1} \underline{\alpha})^2 \\ &\quad + \int_{\Sigma_*} (1 + u^{2+2\delta_{dec}}) (F_k)^2. \end{aligned}$$

Together with the bounds for $\underline{\alpha}$ on \mathcal{C}_1 provided by Theorem M0, the bootstrap assumptions on decay and energy for $\underline{\alpha}$ in $(ext)\mathcal{M}$, and the bound for F_k provided by Proposition 6.2.6, we infer

$$\int_{\Sigma_*} (1 + u^{2+2\delta_{dec}}) (\mathfrak{d}^k \underline{\alpha})^2 \lesssim \epsilon_0^2.$$

In particular, we have obtained

$$\max_{0 \leq k \leq k_{small} + 18} \int_{\Sigma_*} (1 + u^{2+2\delta_{dec}}) (\mathfrak{D}^k \underline{\alpha})^2 \lesssim \epsilon_0.$$

Now, recall that $\underline{\alpha}$ in the above estimate is defined with respect to the global frame constructed in Proposition 3.5.5. In view of Proposition 3.5.5 and Proposition 3.4.6, and the change of frame formula for $\underline{\alpha}$ in Proposition 2.3.4, we have

$$\underline{\alpha} = ({}^{(ext)}\Upsilon)^{-2} ({}^{(ext)}\underline{\alpha}).$$

Hence, we immediately infer

$$\max_{0 \leq k \leq k_{small} + 18} \int_{\Sigma_*} (1 + u^{2+2\delta_{dec}}) (\mathfrak{D}^{k({}^{(ext)}\underline{\alpha})})^2 \lesssim \epsilon_0.$$

which is the desired estimate in Σ_* . Together with (6.2.2), this concludes the proof of Theorem M3.

The proof of Proposition 6.2.6 will be given in section 6.2.5, and to the proof of Lemma 6.2.8 which will be given in section 6.2.6.

6.2.3 Proof of Proposition 6.2.2

In this section, we infer from the Teukolsky-Starobinski identity, see Proposition 2.3.15, a parabolic equation for $\underline{\alpha}$.

Corollary 6.2.9. $\underline{\alpha}$ satisfies in $({}^{int})\mathcal{M}$ the following equation

$$\begin{aligned} & 6m\tilde{T}\underline{\alpha} + r^4 \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \underline{\alpha} \\ &= \frac{1}{r^3} \left(e_3(r^2 e_3(r\mathfrak{q})) + 2\underline{\omega} r^2 e_3(r\mathfrak{q}) \right) - r^{-3} \text{Err}[TS] - \left\{ \frac{3}{2} r^4 \left(\rho + \frac{2m}{r^3} \right) \underline{\kappa} - 3mr \left(\underline{\kappa} + \frac{2}{r} \right) \right\} e_4 \underline{\alpha} \\ & - \left\{ -\frac{3}{2} r^4 \left(\rho + \frac{2m}{r^3} \right) \kappa + \frac{3mr}{\underline{\kappa}} \left(\underline{\kappa} + \frac{2}{r} \right) \bar{\kappa} + 3mr\check{\kappa} + \frac{6m}{\underline{\kappa}} A \right\} e_3 \underline{\alpha} \end{aligned}$$

where the vectorfield \tilde{T} is defined by (6.2.1).

Proof. Recall from (2.3.15) that we have

$$e_3(r^2 e_3(r\mathfrak{q})) + 2\underline{\omega} r^2 e_3(r\mathfrak{q}) = r^7 \left\{ \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \underline{\alpha} + \frac{3}{2} \rho \left(\underline{\kappa} e_4 - \kappa e_3 \right) \underline{\alpha} \right\} + \text{Err}[TS].$$

This yields

$$\frac{3}{2}r^4\rho\left(\underline{\kappa}e_4 - \kappa e_3\right)\underline{\alpha} + r^4\mathcal{D}_2^*\mathcal{D}_1^*\mathcal{D}_1\mathcal{D}_2\underline{\alpha} = \frac{1}{r^3}\left(e_3(r^2e_3(r\mathbf{q})) + 2\underline{\omega}r^2e_3(r\mathbf{q}))\right) - r^{-3}\text{Err}[TS].$$

Now, we have in view of the definition of \tilde{T}

$$\begin{aligned} & \frac{3}{2}r^4\rho\left(\underline{\kappa}e_4 - \kappa e_3\right) - 6m\tilde{T} \\ = & \left\{ \frac{3}{2}r^4\left(\rho + \frac{2m}{r^3}\right)\underline{\kappa} - 3mr\left(\underline{\kappa} + \frac{2}{r}\right) \right\} e_4 \\ & + \left\{ -\frac{3}{2}r^4\left(\rho + \frac{2m}{r^3}\right)\kappa + \frac{3mr}{\underline{\kappa}}\left(\underline{\kappa} + \frac{2}{r}\right)\bar{\kappa} + 3mr\check{\kappa} + \frac{6m}{\underline{\kappa}}A \right\} e_3. \end{aligned}$$

We infer

$$\begin{aligned} & 6m\tilde{T}\underline{\alpha} + r^4\mathcal{D}_2^*\mathcal{D}_1^*\mathcal{D}_1\mathcal{D}_2\underline{\alpha} \\ = & \frac{1}{r^3}\left(e_3(r^2e_3(r\mathbf{q})) + 2\underline{\omega}r^2e_3(r\mathbf{q}))\right) - r^{-3}\text{Err}[TS] - \left\{ \frac{3}{2}r^4\left(\rho + \frac{2m}{r^3}\right)\underline{\kappa} - 3mr\left(\underline{\kappa} + \frac{2}{r}\right) \right\} e_4\underline{\alpha} \\ & - \left\{ -\frac{3}{2}r^4\left(\rho + \frac{2m}{r^3}\right)\kappa + \frac{3mr}{\underline{\kappa}}\left(\underline{\kappa} + \frac{2}{r}\right)\bar{\kappa} + 3mr\check{\kappa} + \frac{6m}{\underline{\kappa}}A \right\} e_3\underline{\alpha}. \end{aligned}$$

This concludes the proof of the corollary. \square

Corollary 6.2.10. $\underline{\alpha}$ satisfies in ${}^{(int)}\mathcal{M}$

$$6m\tilde{T}\underline{\alpha} + r^4\mathcal{D}_2^*\mathcal{D}_1^*\mathcal{D}_1\mathcal{D}_2\underline{\alpha} = F$$

where F satisfies

$$\max_{0 \leq k \leq k_{small} + 18} \int_{{}^{(int)}\mathcal{M}} \underline{u}^{2+2\delta_{dec}} |\mathfrak{D}^k F|^2 \lesssim \epsilon_0^2.$$

Proof. In view of Corollary 6.2.9, $\underline{\alpha}$ satisfies

$$6m\tilde{T}\underline{\alpha} + r^4\mathcal{D}_2^*\mathcal{D}_1^*\mathcal{D}_1\mathcal{D}_2\underline{\alpha} = F$$

with

$$\begin{aligned} F & := \frac{1}{r^3}\left(e_3(r^2e_3(r\mathbf{q})) + 2\underline{\omega}r^2e_3(r\mathbf{q}))\right) + F_1, \\ F_1 & := -r^{-3}\text{Err}[TS] - \left\{ \frac{3}{2}r^4\left(\rho + \frac{2m}{r^3}\right)\underline{\kappa} - 3mr\left(\underline{\kappa} + \frac{2}{r}\right) \right\} e_4\underline{\alpha} \\ & \quad - \left\{ -\frac{3}{2}r^4\left(\rho + \frac{2m}{r^3}\right)\kappa + \frac{3mr}{\underline{\kappa}}\left(\underline{\kappa} + \frac{2}{r}\right)\bar{\kappa} + 3mr\check{\kappa} + \frac{6m}{\underline{\kappa}}A \right\} e_3\underline{\alpha}. \end{aligned}$$

Using the bootstrap assumptions in $(int)\mathcal{M}$ for decay and energies, and in view of the fact that F_1 contains only quadratic or higher order terms, we easily derive

$$\max_{0 \leq k \leq k_{small}+18} \sup_{(int)\mathcal{M}} \underline{u}^{\frac{3}{2} + \frac{3}{2}\delta_{dec}} |\mathfrak{d}^k F_1| \lesssim \epsilon^2 \lesssim \epsilon_0.$$

In view of the definition of F , this yields

$$\max_{0 \leq k \leq k_{small}+18} \int_{(int)\mathcal{M}} \underline{u}^{2+2\delta_{dec}} |\mathfrak{d}^k F|^2 \lesssim \epsilon_0^2 + \max_{0 \leq k \leq k_{small}+20} \int_{(int)\mathcal{M}} \underline{u}^{2+2\delta_{dec}} |\mathfrak{d}^k \mathfrak{q}|^2.$$

Together with Theorem M1, and the fact that $\delta_{extra} > \delta_{dec}$, we infer

$$\max_{0 \leq k \leq k_{small}+18} \int_{(int)\mathcal{M}} \underline{u}^{2+2\delta_{dec}} |\mathfrak{d}^k F|^2 \lesssim \epsilon_0^2.$$

This concludes the proof of the corollary. \square

We are now ready to prove Proposition 6.2.2. In view of Corollary 6.2.10, $\underline{\alpha}$ satisfies

$$6m\tilde{T}\underline{\alpha} + r^4 \not\phi_2^* \not\phi_1^* \not\phi_1 \not\phi_2 \underline{\alpha} = F.$$

Commuting with \mathfrak{d}^k , we infer

$$6m\tilde{T}(\mathfrak{d}^k \underline{\alpha}) + r^4 \not\phi_2^* \not\phi_1^* \not\phi_1 \not\phi_2 (\mathfrak{d}^k \underline{\alpha}) = F_k$$

where F_k is defined by

$$\begin{aligned} F_k := & -6m[\mathfrak{d}^k, \tilde{T}]\underline{\alpha} - 6 \sum_{j=1}^k \mathfrak{d}^j(m) \mathfrak{d}^{k-j} \tilde{T}\underline{\alpha} - [\mathfrak{d}^k, r \not\phi_2^*] r \not\phi_1^* \not\phi_1 r \not\phi_2 \underline{\alpha} - r \not\phi_2^* [\mathfrak{d}^k, r \not\phi_1^*] r \not\phi_1 r \not\phi_2 \underline{\alpha} \\ & - r \not\phi_2^* r \not\phi_1^* [\mathfrak{d}^k, r \not\phi_1] r \not\phi_2 \underline{\alpha} - r \not\phi_2^* r \not\phi_1^* r \not\phi_1 [\mathfrak{d}^k, r \not\phi_2] \underline{\alpha} + \mathfrak{d}^k F. \end{aligned}$$

Note that we have schematically

$$[\mathfrak{d}, \not\phi] = \check{\Gamma} \not\mathfrak{d}, \quad [\tilde{T}, \not\phi] = (\mathfrak{d}\check{\Gamma} + \check{\Gamma}) \not\mathfrak{d},$$

as well as

$$\begin{aligned}
[\tilde{T}, re_4] &= e_4(r)e_4 - \frac{1}{\underline{\kappa}}(\bar{\kappa} + A)[e_3, re_4] + re_4 \left(\frac{1}{\underline{\kappa}}(\bar{\kappa} + A) \right) e_3 \\
&= \frac{r}{2}(\bar{\kappa} + A)e_4 - \frac{1}{\underline{\kappa}}(\bar{\kappa} + A) \left(\frac{r}{2}\bar{\kappa}e_4 + r(-2\omega e_3 + 4\zeta e_\theta) \right) \\
&\quad + re_4 \left(\frac{\bar{\kappa}}{\underline{\kappa}} \right) e_3 + re_4 \left(\frac{1}{\underline{\kappa}}A \right) e_3 \\
&= \left\{ \frac{2r}{\underline{\kappa}}(\bar{\kappa} + A)\omega + re_4 \left(\frac{\bar{\kappa}}{\underline{\kappa}} \right) + re_4 \left(\frac{1}{\underline{\kappa}}A \right) \right\} e_3 \\
&\quad - \frac{4r}{\underline{\kappa}}(\bar{\kappa} + A)\zeta e_\theta \\
&= \left\{ -\frac{2m}{r} \left(\frac{\bar{\kappa}}{\underline{\kappa}} + \Upsilon \right) + \frac{2r}{\underline{\kappa}}\bar{\kappa} \left(\omega + \frac{m}{r} \right) - \frac{2m(e_4(r) - \Upsilon)}{r} + 2e_4(m) + re_4 \left(\frac{\bar{\kappa}}{\underline{\kappa}} + \Upsilon \right) \right. \\
&\quad \left. + \frac{2r}{\underline{\kappa}}A\omega + re_4 \left(\frac{1}{\underline{\kappa}}A \right) \right\} e_3 - \frac{4r}{\underline{\kappa}}(\bar{\kappa} + A)\zeta e_\theta \\
&= (\partial\check{\Gamma} + \check{\Gamma})\mathfrak{d},
\end{aligned}$$

and

$$\begin{aligned}
[\tilde{T}, e_3] &= [e_4, e_3] + e_3 \left(\frac{1}{\underline{\kappa}}(\bar{\kappa} + A) \right) e_3 \\
&= \left\{ 2\omega + e_3 \left(\frac{\bar{\kappa}}{\underline{\kappa}} \right) + e_3 \left(\frac{1}{\underline{\kappa}}A \right) \right\} e_3 - 4\zeta e_\theta \\
&= \left\{ 2 \left(\omega + \frac{m}{r^2} \right) - \frac{2m(e_3(r) + 1)}{r^2} + \frac{2e_3(m)}{r} + e_3 \left(\frac{\bar{\kappa}}{\underline{\kappa}} + \Upsilon \right) + e_3 \left(\frac{1}{\underline{\kappa}}A \right) \right\} e_3 - 4\zeta e_\theta \\
&= (\partial\check{\Gamma} + \check{\Gamma})\mathfrak{d}.
\end{aligned}$$

Together with the bootstrap assumptions in $^{(int)}\mathcal{M}$ for decay and energies, and in view of the fact that F_1 contains only quadratic or higher order terms, we easily derive

$$\max_{0 \leq k \leq k_{small} + 18} \sup_{(int)\mathcal{M}} \underline{u}^{\frac{3}{2} + \frac{3}{2}\delta_{dec}} \left| \begin{aligned} &-6m[\mathfrak{d}^k, \tilde{T}]\underline{\alpha} - 6 \sum_{j=1}^k \mathfrak{d}^j(m) \mathfrak{d}^{k-j} \tilde{T} \underline{\alpha} - [\mathfrak{d}^k, r \mathfrak{d}_2^*] r \mathfrak{d}_1^* r \mathfrak{d}_1 r \mathfrak{d}_2 \underline{\alpha} \\ &-r \mathfrak{d}_2^* [\mathfrak{d}^k, r \mathfrak{d}_1^*] r \mathfrak{d}_1 r \mathfrak{d}_2 \underline{\alpha} - r \mathfrak{d}_2^* r \mathfrak{d}_1^* [\mathfrak{d}^k, r \mathfrak{d}_1] r \mathfrak{d}_2 \underline{\alpha} \\ &-r \mathfrak{d}_2^* r \mathfrak{d}_1^* r \mathfrak{d}_1 [\mathfrak{d}^k, r \mathfrak{d}_2] \underline{\alpha} \end{aligned} \right| \lesssim \epsilon^2.$$

In view of the definition of F_k , this yields

$$\max_{0 \leq k \leq k_{small} + 18} \int_{(int) \mathcal{M}} \underline{u}^{2+2\delta_{dec}} |F_k|^2 \lesssim \epsilon^4 + \max_{0 \leq k \leq k_{small} + 18} \int_{(int) \mathcal{M}} \underline{u}^{2+2\delta_{dec}} |\mathfrak{D}^k F|^2.$$

Together with the estimate for F of Corollary 6.2.10, we infer

$$\max_{0 \leq k \leq k_{small} + 18} \int_{(int) \mathcal{M}} \underline{u}^{2+2\delta_{dec}} |F_k|^2 \lesssim \epsilon^4 + \epsilon_0^2 \lesssim \epsilon_0^2.$$

This concludes the proof of Proposition 6.2.2.

6.2.4 Proof of Lemma 6.2.4

In this section we prove Lemma 6.2.4, i.e. we derive estimates for the control of the parabolic equation appearing in the statement of Proposition 6.2.2. To this end, we first start with a Poincaré inequality.

Lemma 6.2.11. *We have*

$$\int_S f \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2 f \geq 24 \int_S (1 + O(\epsilon)) K^2 f^2.$$

Proof. We have

$$\begin{aligned} \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2 &= \mathfrak{d}_2^* (-\mathfrak{A}_1 + K) \mathfrak{d}_2 \\ &= -\mathfrak{d}_2^* \mathfrak{A}_1 \mathfrak{d}_2 + K \mathfrak{d}_2^* \mathfrak{d}_2 + \mathfrak{d}_1^*(K) \mathfrak{d}_2 \\ &= -\mathfrak{A}_2 \mathfrak{d}_2^* \mathfrak{d}_2 + \left(\mathfrak{A}_2 \mathfrak{d}_2^* - \mathfrak{d}_2^* \mathfrak{A}_1 \right) \mathfrak{d}_2 + K \mathfrak{d}_2^* \mathfrak{d}_2 + \mathfrak{d}_1^*(K) \mathfrak{d}_2 \\ &= (\mathfrak{d}_2^* \mathfrak{d}_2 - 2K) \mathfrak{d}_2^* \mathfrak{d}_2 + \left(3K \mathfrak{d}_2^* - \mathfrak{d}_1^*(K) \right) \mathfrak{d}_2 + K \mathfrak{d}_2^* \mathfrak{d}_2 + \mathfrak{d}_1^*(K) \mathfrak{d}_2 \\ &= (\mathfrak{d}_2^* \mathfrak{d}_2)^2 + 2K \mathfrak{d}_2^* \mathfrak{d}_2. \end{aligned}$$

Recall also the Poincaré inequality for \mathfrak{d}_2 which holds for any reduced 2-scalar f

$$\int_S |\mathfrak{d}_2 f|^2 \geq 4 \int_S K f^2.$$

Then, we easily infer

$$\begin{aligned} \int_S f \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2 f &= \int_S f (\mathfrak{d}_2^* \mathfrak{d}_2)^2 f + \int_S 2K f \mathfrak{d}_2^* \mathfrak{d}_2 f \\ &\geq 4^2 \int_S (1 + O(\epsilon)) K^2 f^2 + 8 \int_S (1 + O(\epsilon)) K^2 f^2 \\ &\geq 24 \int_S (1 + O(\epsilon)) K^2 f^2 \end{aligned}$$

where we also used the estimates for the Gauss curvature

$$K = \frac{1}{r^2} + O\left(\frac{\epsilon}{r^2}\right), \quad re_\theta(K) = O\left(\frac{\epsilon}{r^2}\right),$$

which follow from the bootstrap assumptions. \square

The following identity will be useful.

Lemma 6.2.12. *We have for any reduced scalar f*

$$\begin{aligned} \tilde{T}\left(\int_S f^2\right) &= 2\int_S f\tilde{T}f + \int_S \left(\frac{2}{\underline{\kappa}}Af e_3(f) + \check{\kappa}f^2\right) - \frac{\bar{\kappa}}{\underline{\kappa}}\left(\int_S \check{\kappa}f^2\right) \\ &\quad - \frac{1}{\underline{\kappa}}A\left(\int_S (2fe_3(f) + \underline{\kappa}f^2)\right) + \text{Err}\left[e_4\left(\int_S f^2\right)\right]. \end{aligned}$$

Proof. Recall from the definition of \tilde{T} that

$$\tilde{T} = e_4 - \frac{1}{\underline{\kappa}}(\bar{\kappa} + A)e_3.$$

We infer, in view of the analog of Proposition 2.2.9 for an ingoing geodesic foliation,

$$\begin{aligned} &\tilde{T}\left(\int_S f^2\right) \\ &= e_4\left(\int_S f^2\right) - \frac{1}{\underline{\kappa}}(\bar{\kappa} + A)e_3\left(\int_S f^2\right) \\ &= \int_S (2fe_4(f) + \kappa f^2) + \text{Err}\left[e_4\left(\int_S f^2\right)\right] - \frac{1}{\underline{\kappa}}(\bar{\kappa} + A)\left(\int_S (2fe_3(f) + \underline{\kappa}f^2)\right) \\ &= \int_S \left(2f\tilde{T}f + \frac{2}{\underline{\kappa}}(\bar{\kappa} + A)fe_3(f) + \kappa f^2\right) + \text{Err}\left[e_4\left(\int_S f^2\right)\right] \\ &\quad - \frac{1}{\underline{\kappa}}(\bar{\kappa} + A)\left(\int_S (2fe_3(f) + \underline{\kappa}f^2)\right) \\ &= 2\int_S f\tilde{T}f + \int_S \left(\frac{2}{\underline{\kappa}}Af e_3(f) + \check{\kappa}f^2\right) - \frac{\bar{\kappa}}{\underline{\kappa}}\left(\int_S \check{\kappa}f^2\right) - \frac{1}{\underline{\kappa}}A\left(\int_S (2fe_3(f) + \underline{\kappa}f^2)\right) \\ &\quad + \text{Err}\left[e_4\left(\int_S f^2\right)\right]. \end{aligned}$$

This concludes the proof of the lemma. \square

We are now ready to prove Lemma 6.2.4. Recall from Lemma 6.2.12 that we have

$$\begin{aligned} \tilde{T} \left(\int_S f^2 \right) &= 2 \int_S f \tilde{T} f + \int_S \left(\frac{2}{\underline{\kappa}} A f e_3(f) + \check{\kappa} f^2 \right) - \frac{\bar{\kappa}}{\underline{\kappa}} \left(\int_S \check{\kappa} f^2 \right) \\ &\quad - \frac{1}{\underline{\kappa}} A \left(\int_S (2f e_3(f) + \underline{\kappa} f^2) \right) + \text{Err} \left[e_4 \left(\int_S f^2 \right) \right]. \end{aligned}$$

In view of the equation satisfied by f , we infer

$$\begin{aligned} \tilde{T} \left(\int_S f^2 \right) &= -\frac{1}{3m} \left(\int_S r^4 f \mathcal{d}_2^* \mathcal{d}_1^* \mathcal{d}_1 \mathcal{d}_2 f \right) + \frac{1}{3m} \left(\int_S h f \right) \\ &\quad + \int_S \left(\frac{2}{\underline{\kappa}} A f e_3(f) + \check{\kappa} f^2 \right) - \frac{\bar{\kappa}}{\underline{\kappa}} \left(\int_S \check{\kappa} f^2 \right) - \frac{1}{\underline{\kappa}} A \left(\int_S (2f e_3(f) + \underline{\kappa} f^2) \right) \\ &\quad + \text{Err} \left[e_4 \left(\int_S f^2 \right) \right]. \end{aligned}$$

Now, from the definition of \tilde{T} , we have $\tilde{T}(\underline{u}) = 2/\underline{\varsigma}$. We deduce

$$\begin{aligned} &\tilde{T} \left(\underline{u} \int_S f^2 \right) + \frac{\underline{u}^n}{3m} \left(\int_S r^4 f \mathcal{d}_2^* \mathcal{d}_1^* \mathcal{d}_1 \mathcal{d}_2 f \right) \\ &= \frac{\underline{u}^n}{3m} \left(\int_S h f \right) + \underline{u} \int_S \left(\frac{2}{\underline{\kappa}} A f e_3(f) + \check{\kappa} f^2 \right) - \underline{u} \frac{\bar{\kappa}}{\underline{\kappa}} \left(\int_S \check{\kappa} f^2 \right) \\ &\quad - \frac{\underline{u}^n}{\underline{\kappa}} A \left(\int_S (2f e_3(f) + \underline{\kappa} f^2) \right) + \underline{u} \text{Err} \left[e_4 \left(\int_S f^2 \right) \right] + \frac{2}{\underline{\varsigma}} n \underline{u}^{n-1} \int_S f^2. \end{aligned}$$

This yields in view of the bootstrap assumptions

$$\begin{aligned} &\tilde{T} \left(\underline{u} \int_S f^2 \right) + \frac{\underline{u}^n}{3m} \left(\int_S r^4 f \mathcal{d}_2^* \mathcal{d}_1^* \mathcal{d}_1 \mathcal{d}_2 f \right) \\ &\lesssim \frac{\underline{u}^n}{3m} \|h\|_{L^2(S)} \|f\|_{L^2(S)} + \epsilon \underline{u}^{n-1} \int_S |f| |\mathfrak{d}^{\leq 1} f| + n \underline{u}^{n-1} \int_S f^2. \end{aligned}$$

Next, we rely on the Poincaré inequality of Lemma 6.2.11 to deduce

$$\tilde{T} \left(\underline{u} \int_S f^2 \right) + \underline{u} \int_S f^2 \lesssim \underline{u} \int_S h^2 + \epsilon^2 \underline{u}^{n-2} \int_S (\mathfrak{d}f)^2 + n \underline{u}^{n-1} \int_S f^2.$$

Integrating in \underline{u} between 1 and u_* , and recalling that $\tilde{T}(\underline{u}) = 2/\underline{\varsigma}$, we infer for any r_0 such that $2m_0(1 - \delta_{\mathcal{H}}) \leq r_0 \leq r_{\mathcal{T}}$

$$\begin{aligned} &\int_{S(r=r_0, \underline{u})} \underline{u}^n f^2 + \int_1^{\underline{u}} \left(\int_{S(r=r_0, \underline{u}')} \underline{u}'^m f^2 \right) d\underline{u}' \\ &\lesssim \int_{S(r=r_0, 1)} f^2 + \int_1^{\underline{u}} \left(\int_{S(r=r_0, \underline{u}')} \underline{u}'^m h^2 \right) d\underline{u}' + \epsilon^2 \int_1^{\underline{u}} \left(\int_{S(r=r_0, \underline{u}')} \underline{u}'^{m-2} (\mathfrak{d}f)^2 \right) d\underline{u}' \\ &\quad + n \int_0^{\underline{u}} \left(\int_{S(r=r_0, \underline{u}')} \underline{u}'^{m-1} f^2 \right) d\underline{u}'. \end{aligned}$$

In particular, we have for $n = 0$

$$\begin{aligned} \sup_{1 \leq \underline{u} \leq u_*} \int_{S(r=r_0, \underline{u})} f^2 + \int_1^{u_*} \left(\int_{S(r=r_0, \underline{u})} f^2 \right) d\underline{u} &\lesssim \int_{S(r=r_0, 1)} f^2 + \int_1^{u_*} \left(\int_{S(r=r_0, \underline{u})} h^2 \right) d\underline{u} \\ &+ \epsilon^2 \int_1^{u_*} \left(\int_{S(r=r_0, \underline{u})} \underline{u}^{-2} (\mathfrak{d}f)^2 \right) d\underline{u}. \end{aligned}$$

Then, starting from the case $n = 0$ and arguing by iteration on the largest integer below n , one immediately deduces for any real $n \geq 0$

$$\begin{aligned} &\sup_{1 \leq \underline{u} \leq u_*} \int_{S(r=r_0, \underline{u})} (1 + \underline{u}^n) f^2 + \int_1^{u_*} \left(\int_{S(r=r_0, \underline{u})} (1 + \underline{u}^n) f^2 \right) d\underline{u} \\ &\lesssim \int_{S(r=r_0, 1)} f^2 + \int_1^{u_*} \left(\int_{S(r=r_0, \underline{u})} (1 + \underline{u}^n) h^2 \right) d\underline{u} + \epsilon^2 \int_1^{u_*} \left(\int_{S(r=r_0, \underline{u})} (1 + \underline{u}^{n-2}) (\mathfrak{d}f)^2 \right) d\underline{u}. \end{aligned}$$

Now, a simple trace estimate yields

$$\int_{S(r=r_0, \underline{u})} (1 + \underline{u}^n) h^2 \lesssim \int_{\underline{\mathcal{C}}_{\underline{u}}} (1 + \underline{u}^n) (|h|^2 + |e_3(h)|^2)$$

so that

$$\begin{aligned} \int_1^{u_*} \left(\int_{S(r=r_0, \underline{u})} (1 + \underline{u}^n) h^2 \right) d\underline{u} &\lesssim \int_1^{u_*} \int_{\underline{\mathcal{C}}_{\underline{u}}} (1 + \underline{u}^n) (|h|^2 + |e_3(h)|^2) d\underline{u} \\ &\lesssim \int_{(int)\mathcal{M}} (1 + \underline{u}^n) (\mathfrak{d}^{\leq 1} h)^2. \end{aligned}$$

We deduce

$$\begin{aligned} &\sup_{1 \leq \underline{u} \leq u_*} \int_{S(r=r_0, \underline{u})} (1 + \underline{u}^n) f^2 + \int_1^{u_*} \left(\int_{S(r=r_0, \underline{u})} (1 + \underline{u}^n) f^2 \right) d\underline{u} \\ &\lesssim \int_{S(r=r_0, 1)} f^2 + \int_{(int)\mathcal{M}} (1 + \underline{u}^n) (\mathfrak{d}^{\leq 1} h)^2 + \epsilon^2 \int_1^{u_*} \left(\int_{S(r=r_0, \underline{u})} (1 + \underline{u}^{n-2}) (\mathfrak{d}f)^2 \right) d\underline{u} \end{aligned}$$

which concludes the proof of Lemma 6.2.4.

6.2.5 Proof of Proposition 6.2.6

In this section, we infer from the Teukolsky-Starobinski identity, see Proposition 2.3.15, a parabolic equation for $\underline{\alpha}$.

Corollary 6.2.13. $\underline{\alpha}$ satisfies in $^{(int)}\mathcal{M}$ the following equation

$$\begin{aligned} & 6m\tilde{\nu}\underline{\alpha} + r^4 \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \mathcal{D}_2 \underline{\alpha} \\ = & \frac{1}{r^3} \left(e_3(r^2 e_3(r\mathbf{q})) + 2\underline{\omega} r^2 e_3(r\mathbf{q}) \right) - r^{-3} \text{Err}[TS] - \left\{ \frac{3}{2} r^4 \rho \underline{\kappa} - 6am \right\} e_4 \underline{\alpha} \\ & - \left\{ -\frac{3}{2} r^4 \left(\rho + \frac{2m}{r^3} \right) \kappa + 3mr \left(\kappa - \frac{2\Upsilon}{r} \right) - \frac{12m}{r} \right\} e_3 \underline{\alpha}. \end{aligned}$$

where the vectorfield $\tilde{\nu}$ is defined by (6.2.3).

Proof. Recall from (2.3.15) that we have

$$e_3(r^2 e_3(r\mathbf{q})) + 2\underline{\omega} r^2 e_3(r\mathbf{q}) = r^7 \left\{ \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \mathcal{D}_2 \underline{\alpha} + \frac{3}{2} \rho (\underline{\kappa} e_4 - \kappa e_3) \underline{\alpha} \right\} + \text{Err}[TS].$$

This yields

$$\frac{3}{2} r^4 \rho (\underline{\kappa} e_4 - \kappa e_3) \underline{\alpha} + r^4 \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \mathcal{D}_2 \underline{\alpha} = \frac{1}{r^3} \left(e_3(r^2 e_3(r\mathbf{q})) + 2\underline{\omega} r^2 e_3(r\mathbf{q}) \right) - r^{-3} \text{Err}[TS].$$

Now, we have in view of the definition of $\tilde{\nu}$

$$\begin{aligned} & \frac{3}{2} r^4 \rho (\underline{\kappa} e_4 - \kappa e_3) - 6m\tilde{\nu} \\ = & \left\{ \frac{3}{2} r^4 \rho \underline{\kappa} - 6am \right\} e_4 + \left\{ -\frac{3}{2} r^4 \left(\rho + \frac{2m}{r^3} \right) \kappa + 3mr \left(\kappa - \frac{2\Upsilon}{r} \right) - \frac{12m}{r} \right\} e_3. \end{aligned}$$

We infer

$$\begin{aligned} & 6m\tilde{\nu}\underline{\alpha} + r^4 \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \mathcal{D}_2 \underline{\alpha} \\ = & \frac{1}{r^3} \left(e_3(r^2 e_3(r\mathbf{q})) + 2\underline{\omega} r^2 e_3(r\mathbf{q}) \right) - r^{-3} \text{Err}[TS] - \left\{ \frac{3}{2} r^4 \rho \underline{\kappa} - 6am \right\} e_4 \underline{\alpha} \\ & - \left\{ -\frac{3}{2} r^4 \left(\rho + \frac{2m}{r^3} \right) \kappa + 3mr \left(\kappa - \frac{2\Upsilon}{r} \right) - \frac{12m}{r} \right\} e_3 \underline{\alpha}. \end{aligned}$$

This concludes the proof of the corollary. □

Corollary 6.2.14. $\underline{\alpha}$ satisfies in $^{(int)}\mathcal{M}$

$$6m\tilde{\nu}\underline{\alpha} + r^4 \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \mathcal{D}_2 \underline{\alpha} = F$$

where F satisfies

$$\max_{0 \leq k \leq k_{small} + 18} \int_{\Sigma_*} \underline{u}^{2+2\delta_{dec}} |\mathfrak{D}^k F|^2 \lesssim \epsilon_0^2.$$

Proof. In view of Corollary 6.2.9, $\underline{\alpha}$ satisfies

$$6m\tilde{\nu}\underline{\alpha} + r^4 \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \underline{\alpha} = F$$

with

$$\begin{aligned} F &:= e_3(e_3(\mathbf{q})) + F_1, \\ F_1 &:= \frac{1}{r^3} \left(e_3(r^2 e_3(r)\mathbf{q}) + e_3(r^3) e_3(\mathbf{q}) + 2\omega r^2 e_3(r\mathbf{q}) \right) - r^{-3} \text{Err}[TS] \\ &\quad - \left\{ \frac{3}{2} r^4 \rho \underline{\kappa} - 6am \right\} e_4 \underline{\alpha} - \left\{ -\frac{3}{2} r^4 \left(\rho + \frac{2m}{r^3} \right) \kappa + 3mr \left(\kappa - \frac{2\Upsilon}{r} \right) - \frac{12m}{r} \right\} e_3 \underline{\alpha}. \end{aligned}$$

Recall also that $\text{Err}[TS]$ is given schematically by, see Proposition 2.3.15,

$$\begin{aligned} \text{Err}[TS] &= r^4 (\not\partial \Gamma_b + r \Gamma_b \cdot \Gamma_b) \cdot \underline{\alpha} + r^2 (\Gamma_b e_3(r\mathbf{q}) + (\not\partial^{\leq 1} \Gamma_b) r \mathbf{q}) \\ &\quad + r^7 \not\partial^{\leq 2} (e_3 \eta \cdot \beta) + r^5 \not\partial^{\leq 3} (\Gamma_b \cdot \Gamma_g). \end{aligned}$$

We infer that F_1 is given schematically by

$$\begin{aligned} F_1 &= r (\not\partial \Gamma_b + r \Gamma_b \cdot \Gamma_b) \cdot \underline{\alpha} + r^{-1} (\Gamma_b e_3(r\mathbf{q}) + (\not\partial^{\leq 1} \Gamma_b) r \mathbf{q}) \\ &\quad + r^4 \not\partial^{\leq 2} (e_3 \eta \cdot \beta) + r^2 \not\partial^{\leq 3} (\Gamma_b \cdot \Gamma_g) + r^{-1} \Gamma_b \\ &= r^{-1} \Gamma_b + r^2 \not\partial^{\leq 3} (\Gamma_b \cdot \Gamma_g) + r^4 \not\partial^{\leq 3} (\Gamma_g \cdot \beta) \end{aligned}$$

where we have used

- The fact we are working here with the global frame of Proposition 3.5.5 which has the property that $\eta \in \Gamma_g$.
- The fact that Γ_b behave better than $r\Gamma_g$.
- The fact that $\underline{\alpha}$ and \mathbf{q} behaves at least as good as Γ_b .
- The fact that $\rho + \frac{2m}{r^3}$ behaves as good as $r^{-1}\Gamma_g$.
- The fact that $e_3(r) + 1$ belongs to $r\Gamma_b$.

Now, recall from Lemma 5.1.1 that the global frame of Proposition satisfies in particular²

$$\max_{0 \leq k \leq k_{\text{small}} + 22} \sup_{\mathcal{M}} \left\{ r^{\frac{7}{2} + \delta_{dec} - 2\delta_0} |\not\partial^k \beta| + r^2 u^{\frac{1}{2} + \delta_{dec} - 2\delta_0} |\not\partial^k \Gamma_g| + r u^{1 + \delta_{dec} - 2\delta_0} |\not\partial^k \Gamma_b| \right\} \lesssim \epsilon. \quad (6.2.4)$$

²Here we use (3.4.11) with $k_{\text{loss}} = 22$.

Together with the schematic for of F_1 and the behavior (3.3.4) of r on Σ_* , and the fact that δ_0 can be chosen to satisfy³ $8\delta_0 \leq \delta_{dec}$, we infer

$$\max_{0 \leq k \leq k_{small}+18} \sup_{\Sigma_*} r u^{\frac{3}{2} + \frac{3}{2}\delta_{dec}} |\mathfrak{D}^k F_1| \lesssim u_*^{\frac{3}{2} + \frac{3}{2}\delta_{dec}} \sup_{\Sigma_*} (r^{-\frac{1}{2}}) \epsilon + \epsilon^2 \lesssim \epsilon_0.$$

In view of the definition of F , this yields

$$\max_{0 \leq k \leq k_{small}+18} \int_{\Sigma_*} u^{2+2\delta_{dec}} |\mathfrak{D}^k F|^2 \lesssim \epsilon_0^2 + \max_{0 \leq k \leq k_{small}+19} \int_{\Sigma_*} u^{2+2\delta_{dec}} |\mathfrak{D}^k e_3(\mathfrak{q})|^2.$$

Together with Theorem M1, and the fact that $\delta_{extra} > \delta_{dec}$, we infer

$$\max_{0 \leq k \leq k_{small}+18} \int_{\Sigma_*} u^{2+2\delta_{dec}} |\mathfrak{D}^k F|^2 \lesssim \epsilon_0^2.$$

This concludes the proof of the corollary. \square

We are now ready to prove Proposition 6.2.6. In view of Corollary 6.2.14, $\underline{\alpha}$ satisfies

$$6m\tilde{\nu}\underline{\alpha} + r^4 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2 \underline{\alpha} = F.$$

Commuting with \mathfrak{D}^k , we infer

$$6m\tilde{\nu}(\mathfrak{D}^k \underline{\alpha}) + r^4 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2 (\mathfrak{D}^k \underline{\alpha}) = F_k$$

where F_k is defined by

$$\begin{aligned} F_k := & -6m[\mathfrak{D}^k, \tilde{\nu}]\underline{\alpha} - 6 \sum_{j=1}^k \mathfrak{D}^j(m) \mathfrak{D}^{k-j} \tilde{\nu} \underline{\alpha} - [\mathfrak{D}^k, r \mathfrak{d}_2^*] r \mathfrak{d}_1^* r \mathfrak{d}_1 r \mathfrak{d}_2 \underline{\alpha} - r \mathfrak{d}_2^* [\mathfrak{D}^k, r \mathfrak{d}_1^*] r \mathfrak{d}_1 r \mathfrak{d}_2 \underline{\alpha} \\ & - r \mathfrak{d}_2^* r \mathfrak{d}_1^* [\mathfrak{D}^k, r \mathfrak{d}_1] r \mathfrak{d}_2 \underline{\alpha} - r \mathfrak{d}_2^* \mathfrak{d}_1^* r \mathfrak{d}_1 [\mathfrak{D}^k, r \mathfrak{d}_2] \underline{\alpha} + \mathfrak{D}^k F. \end{aligned}$$

Note that we have schematically

$$[\mathfrak{D}, \mathfrak{D}] = r\Gamma_b \mathfrak{D}, \quad [\tilde{\nu}, \mathfrak{D}] = \left(O(r^{-1}) + r\Gamma_b \right) \mathfrak{D}, \quad [\tilde{\nu}, r e_4] = O(r^{-1}) \mathfrak{D}, \quad [\tilde{\nu}, e_3] = O(r^{-1}) \mathfrak{D}.$$

³Recall from Lemma 5.1.1 that we have

$$\delta_0 = \frac{k_{loss}}{k_{large} - k_{small}}.$$

Since we have here $k_{loss} = 22$, and since we have $2k_{small} \leq k_{large} + 1$ and $k_{large}\delta_{dec} \gg 1$, we deduce $\delta_0 \ll \delta_{dec}$ and we have indeed $8\delta_0 \leq \delta_{dec}$.

Together with the fact that $\underline{\alpha}$ behaves at least as good as Γ_b , we infer, schematically,

$$F_k = \mathfrak{d}^k F + r^{-1} \mathfrak{d}^{\leq k+4} \Gamma_b + r \mathfrak{d}^{\leq k+4} (\Gamma_b^2).$$

In view of (6.2.4) and the behavior (3.3.4) of r on Σ_* , we have

$$\max_{0 \leq k \leq k_{small} + 18} \sup_{\Sigma_*} r u^{\frac{3}{2} + \frac{3}{2} \delta_{dec}} |r^{-1} \mathfrak{d}^{\leq k+4} \Gamma_b + r \mathfrak{d}^{\leq k+4} (\Gamma_b^2)| \lesssim u_*^{\frac{3}{2} + \frac{3}{2} \delta_{dec}} \sup_{\Sigma_*} (r^{-\frac{1}{2}}) \epsilon + \epsilon^2 \lesssim \epsilon_0.$$

This yields

$$\max_{0 \leq k \leq k_{small} + 18} \int_{\Sigma_*} u^{2+2\delta_{dec}} |F_k|^2 \lesssim \epsilon_0^2 + \max_{0 \leq k \leq k_{small} + 18} \sup_{\Sigma_*} u^{2+2\delta_{dec}} |\mathfrak{d}^k F|^2.$$

Together with the estimate for F of Corollary 6.2.14, we infer

$$\max_{0 \leq k \leq k_{small} + 18} \int_{\Sigma_*} u^{2+2\delta_{dec}} |F_k|^2 \lesssim \epsilon^4 + \epsilon_0^2 \lesssim \epsilon_0^2.$$

This concludes the proof of Proposition 6.2.6.

6.2.6 Proof of Lemma 6.2.8

In this section we prove Lemma 6.2.8, i.e. we derive estimates for the control of the parabolic equation appearing in the statement of Proposition 6.2.6. The following identity will be useful.

Lemma 6.2.15. *We have for any reduced scalar f*

$$\begin{aligned} & \tilde{\nu} \left(\int_S f^2 \right) \\ &= 2 \int_S f \tilde{\nu}(f) + \int_S (-2af e_4(f) + \underline{\kappa} f^2) + a \int_S (2f e_4(f) + \kappa f^2) + \text{Err} \left[e_3 \left(\int_S f^2 \right) \right]. \end{aligned}$$

Proof. Recall from the definition of $\tilde{\nu}$ that

$$\tilde{\nu} = e_3 + a e_4.$$

We infer, in view of Proposition 2.2.9,

$$\begin{aligned}
& \tilde{\nu} \left(\int_S f^2 \right) \\
&= e_3 \left(\int_S f^2 \right) + ae_4 \left(\int_S f^2 \right) \\
&= \int_S (2fe_3(f) + \underline{\kappa}f^2) + \text{Err} \left[e_3 \left(\int_S f^2 \right) \right] + a \int_S (2fe_4(f) + \kappa f^2) \\
&= 2 \int_S f\tilde{\nu}(f) + \int_S (-2afe_4(f) + \underline{\kappa}f^2) + a \int_S (2fe_4(f) + \kappa f^2) + \text{Err} \left[e_3 \left(\int_S f^2 \right) \right].
\end{aligned}$$

This concludes the proof of the lemma. \square

We are now ready to prove Lemma 6.2.4. Recall from Lemma 6.2.15 that we have

$$\begin{aligned}
& \tilde{\nu} \left(\int_S f^2 \right) \\
&= 2 \int_S f\tilde{\nu}(f) + \int_S (-2afe_4(f) + \underline{\kappa}f^2) + a \int_S (2fe_4(f) + \kappa f^2) + \text{Err} \left[e_3 \left(\int_S f^2 \right) \right].
\end{aligned}$$

In view of the equation satisfied by f , we infer

$$\begin{aligned}
\tilde{\nu} \left(\int_S f^2 \right) &= -\frac{1}{3m} \left(\int_S r^4 f \phi_2^* \phi_1^* \phi_1 \phi_2 f \right) + \frac{1}{3m} \left(\int_S hf \right) \\
&\quad + \int_S (-2afe_4(f) + \underline{\kappa}f^2) + a \int_S (2fe_4(f) + \kappa f^2) + \text{Err} \left[e_3 \left(\int_S f^2 \right) \right].
\end{aligned}$$

Now, from the definition of $\tilde{\nu}$, we have $\tilde{\nu}(u) = 2/\varsigma$. We deduce

$$\begin{aligned}
& \tilde{\nu} \left(u^n \int_S f^2 \right) + \frac{u^n}{3m} \left(\int_S r^4 f \phi_2^* \phi_1^* \phi_1 \phi_2 f \right) \\
&= \frac{u^n}{3m} \left(\int_S hf \right) + u^n \int_S (-2afe_4(f) + \underline{\kappa}f^2) + au^n \int_S (2fe_4(f) + \kappa f^2) \\
&\quad + u^n \text{Err} \left[e_3 \left(\int_S f^2 \right) \right] + \frac{2}{\varsigma} nu^{n-1} \int_S f^2.
\end{aligned}$$

This yields in view of the bootstrap assumptions

$$\begin{aligned}
& \tilde{\nu} \left(u^n \int_S f^2 \right) + \frac{u^n}{3m} \left(\int_S r^4 f \phi_2^* \phi_1^* \phi_1 \phi_2 f \right) \\
&\lesssim \frac{u^n}{3m} \|h\|_{L^2(S)} \|f\|_{L^2(S)} + \left(\frac{1}{r} + \epsilon u^{-1} \right) u^n \int_S |f| |\mathfrak{d}^{\leq 1} f| + nu^{n-1} \int_S f^2 \\
&\lesssim \frac{u^n}{3m} \|h\|_{L^2(S)} \|f\|_{L^2(S)} + \epsilon u^{n-1} \int_S |f| |\mathfrak{d}^{\leq 1} f| + nu^{n-1} \int_S f^2
\end{aligned}$$

where we have used in the last inequality the behavior (3.3.4) of r on Σ_* . Next, we rely on the Poincaré inequality of Lemma 6.2.11 to deduce

$$\tilde{\nu} \left(u^n \int_S f^2 \right) + u^n \int_S f^2 \lesssim u^n \int_S h^2 + \epsilon^2 u^{n-2} \int_S (\mathfrak{d}f)^2 + n u^{n-1} \int_S f^2.$$

Integrating in u between 1 and u_* , and recalling that $\tilde{\nu}(u) = 2/\varsigma$, we infer

$$\int_{\Sigma_*} u^n f^2 \lesssim \int_{\Sigma_* \cap \mathcal{C}_1} f^2 + \int_{\Sigma_*} u^n h^2 + \epsilon^2 \int_{\Sigma_*} u^{n-2} (\mathfrak{d}f)^2 + n \int_{\Sigma_*} u^{n-1} f^2.$$

In particular, we have for $n = 0$

$$\int_{\Sigma_*} f^2 \lesssim \int_{\Sigma_* \cap \mathcal{C}_1} f^2 + \int_{\Sigma_*} h^2 + \epsilon^2 \int_{\Sigma_*} u^{-2} (\mathfrak{d}f)^2.$$

Then, starting from the case $n = 0$ and arguing by iteration on the largest integer below n , one immediately deduces for any real $n \geq 0$

$$\int_{\Sigma_*} (1 + u^n) f^2 \lesssim \int_{\Sigma_* \cap \mathcal{C}_1} f^2 + \int_{\Sigma_*} (1 + u^n) h^2 + \epsilon^2 \int_{\Sigma_*} (1 + u^{n-2}) (\mathfrak{d}f)^2$$

which concludes the proof of Lemma 6.2.8.

Chapter 7

DECAY ESTIMATES (Theorems M4, M5)

In this chapter, we rely on the decay of \mathfrak{q} , α and $\underline{\alpha}$ to prove the decay estimates for all the other quantities. More precisely, we rely on the results of Theorem M1, M2 and M3 to prove Theorem M4 and M5.

7.1 Preliminaries to the proof of Theorem M4

In what follows we give a detailed proof of Theorem M4, which, we recall, provides the main decay estimates in ${}^{(ext)}\mathcal{M}$. The proof makes use of the bootstrap assumptions BA-D, BA-E, the results of Theorems M1, M2, M3 and Lemmas 3.4.1, 3.4.2. In this section, we start with some preliminaries.

7.1.1 Geometric structure of Σ_*

The proof of Theorem M4 depends in a fundamental way on the geometric properties of the GCM hypersurface Σ_* , the spacelike future boundary of ${}^{(ext)}\mathcal{M}$ introduced in section 3.1.2. For the convenience of the reader, we recall below its main features.

1. The affine parameter s is initialized on Σ_* such that $s = r$.

2. There exists a constant c_* such that

$$\Sigma_* := \{u + r = c_*\}.$$

3. Let $\nu_* = e_3 + a_*e_4$ be the unique vectorfield tangent to the hypersurface Σ_* , perpendicular to the foliation $S(u)$ induced on Σ_* and normalized by the condition $g(\nu_*, e_4) = -2$. The following normalization condition holds true at the South Pole SP of every sphere S ,

$$a_* \Big|_{SP} = -1 - \frac{2m^S}{r^S}. \quad (7.1.1)$$

4. We have

$$r_* \geq \epsilon_0^{-\frac{2}{3}} u^4 \quad \text{on } \Sigma_*. \quad (7.1.2)$$

5. The following GCM conditions hold on Σ_*

$$\kappa = \frac{2}{r}, \quad \not{d}_2^* \not{d}_1^* \underline{\kappa} = 0, \quad \not{d}_2^* \not{d}_1^* \mu = 0, \quad (7.1.3)$$

$$\int_S \eta e^\Phi = \int_S \underline{\xi} e^\Phi = 0. \quad (7.1.4)$$

Moreover on $S_* = \Sigma_* \cap \mathcal{C}_*$,

$$\int_{S_*} \beta e^\Phi = 0, \quad \int_{S_*} e_\theta(\underline{\kappa}) e^\Phi = 0. \quad (7.1.5)$$

6. According to the definition of the Hawking mass, i.e. $1 - \frac{2m}{r} = -\frac{r^2}{4} \overline{\kappa \underline{\kappa}}$, and the GCM assumption for κ we also have,

$$\underline{\kappa} = -\frac{r}{2} \left(1 - \frac{2m}{r}\right). \quad (7.1.6)$$

Thus on Σ_* ,

$$e_3(r) = \frac{r}{2} (\underline{\kappa} + \underline{A}) = -\Upsilon + \frac{r}{2} \underline{A}, \quad e_4(r) = 1. \quad (7.1.7)$$

7. In view of the definition of ν_* and that of ς we we easily deduce¹ the following relation between a_* and ς on Σ_* .

$$a_* = -\frac{2}{\varsigma} + \Upsilon - \frac{r}{2} \underline{A}. \quad (7.1.8)$$

8. Since on Σ_* we have $r = s$ we deduce,

$$\underline{\Omega} = e_3(r) = -\Upsilon + \frac{r}{2} \underline{A} \quad \text{on } \Sigma_*. \quad (7.1.9)$$

¹Indeed, since ν_* is tangent to Σ_* along which $u = -r + c_*$, using also (7.1.7), $\frac{2}{\varsigma} = e_3(u) = \nu_*(u) = -\nu_*(r) = -e_3(r) - a_*e_4(r) = -a_* + \Upsilon - \frac{r}{2} \underline{A}$.

7.1.2 Main Assumptions

We reformulate below the main bootstrap assumption² in the form needed in the proof of Theorem M4.

Definition 7.1.1. *We make use of the following norms on $S = S(u, r) \subset {}^{(ext)}\mathcal{M}$,*

$$\begin{aligned} \|f\|_\infty(u, r) &:= \|f\|_{L^\infty(S(u, r))}, & \|f\|_2(u, r) &:= \|f\|_{L^2(S(u, r))}, \\ \|f\|_{\infty, k}(u, r) &:= \sum_{i=0}^k \|\mathfrak{d}^i f\|_\infty(u, r), & \|f\|_{2, k}(u, r) &:= \sum_{i=0}^k \|\mathfrak{d}^i f\|_2(u, r). \end{aligned} \quad (7.1.10)$$

To simplify the exposition it also helps to introduce the following schematic notation for the connection coefficients (recall $\omega, \xi = 0$ and $\zeta = -\eta$),

$$\begin{aligned} \Gamma_g &= \left\{ \check{\kappa}, \vartheta, \underline{\eta}, \zeta, \check{\underline{\kappa}} \right\} \cup \left\{ \bar{\kappa} - \frac{2}{r}, \bar{\underline{\kappa}} + \frac{2\Upsilon}{r} \right\}, \\ \Gamma_b &= \left\{ \underline{\vartheta}, \eta, \underline{\check{\omega}}, \underline{\xi}, \underline{A}, r^{-1}\check{\zeta}, r^{-1}\check{\underline{\Omega}} \right\} \cup \left\{ \bar{\omega} - \frac{m}{r^2}, r^{-1}(\bar{\zeta} - 1), r^{-1}(\bar{\underline{\Omega}} + \Upsilon) \right\}. \end{aligned} \quad (7.1.11)$$

Remark 7.1.2. *It is important to note that η belongs to Γ_b rather than Γ_g as it may have been expected. Note also that $\underline{A} \in \Gamma_b$ in view of Proposition 2.2.9 and the fact that $(\check{\zeta}, \check{\underline{\Omega}}) \in r\Gamma_b$. We also note that the averaged quantities $\left\{ \bar{\kappa} - \frac{2}{r}, \bar{\underline{\kappa}} + \frac{2\Upsilon}{r} \right\}$ and $\left\{ \bar{\omega} - \frac{m}{r^2}, r^{-1}(\bar{\zeta} - 1), r^{-1}(\bar{\underline{\Omega}} + \Upsilon) \right\}$ are actually better behaved in view of Lemmas 3.4.1, 3.4.2.*

Ref 1. According to our bootstrap assumptions **BA-D**, and the pointwise estimates of Proposition 3.4.5, which themselves follow from **BA-E**, as well as the control of averages in Lemma 3.4.1 and the control of the Hawking mass in Lemma 3.4.2, we have on ${}^{(ext)}\mathcal{M}$,

1. For $0 \leq k \leq k_{small}$,

$$\begin{aligned} \|\Gamma_g\|_{\infty, k} &\lesssim \epsilon \min \left\{ r^{-2} u^{-\frac{1}{2} - \delta_{dec}}, r^{-1} u^{-1 - \delta_{dec}} \right\}, \\ \|e_3 \Gamma_g\|_{\infty, k-1} &\lesssim \epsilon r^{-2} u^{-1 - \delta_{dec}}, \\ \|\Gamma_b\|_{\infty, k} &\lesssim \epsilon r^{-1} u^{-1 - \delta_{dec}}. \end{aligned} \quad (7.1.12)$$

2. For $k \leq k_{large} - 5$

$$\|\Gamma_g\|_{\infty, k} \lesssim \epsilon r^{-2}, \quad \|\Gamma_b\|_{\infty, k} \lesssim \epsilon r^{-1}. \quad (7.1.13)$$

²Based on bootstrap assumptions **BA-D**, **BA-E**, Theorems **M1**, **M2**, **M3** and Lemmas 3.4.1, 3.4.2.

Ref 2. The quantity³ \mathfrak{q} satisfies on $^{(ext)}\mathcal{M}$, for all $0 \leq k \leq k_{small} + 20$,

$$\begin{aligned} \|\mathfrak{q}\|_{\infty, k} &\lesssim \epsilon_0 \min \left\{ u^{-1-\delta_{extra}}, r^{-1}u^{-\frac{1}{2}-\delta_{extra}} \right\}, \\ \|e_3\mathfrak{q}\|_{\infty, k-1} &\lesssim \epsilon_0 r^{-1}u^{-1-\delta_{extra}}. \end{aligned} \quad (7.1.14)$$

In addition, on the last slice Σ_* , for all $k \leq k_{small} + 20$,

$$\int_{\Sigma_*(\tau, \tau_*)} |e_3\mathfrak{d}^k\mathfrak{q}|^2 + |e_4\mathfrak{d}^k\mathfrak{q}|^2 + r^{-2}|\mathfrak{q}|^2 \lesssim \epsilon_0^2(1+\tau)^{-2-2\delta_{dec}}. \quad (7.1.15)$$

According to Theorem **M2** we have on $^{(ext)}\mathcal{M}$, for all $0 \leq k \leq k_{small} + 20$,

$$\begin{aligned} \|\alpha\|_{\infty, k} &\lesssim \epsilon_0 \min \left\{ r^{-3}(u+2r)^{-\frac{1}{2}-\delta_{extra}}, \log(1+u)r^{-2}(u+2r)^{-1-\delta_{extra}} \right\}, \\ \|e_3\alpha\|_{\infty, k-1} &\lesssim \epsilon_0 \min \left\{ r^{-4}(u+2r)^{-\frac{1}{2}-\delta_{extra}}, \log(1+u)r^{-3}(u+2r)^{-1-\delta_{extra}} \right\}. \end{aligned} \quad (7.1.16)$$

According to Theorem **M3**, the component $\underline{\alpha}$ verifies the following estimate⁴ holds on \mathcal{T} , for $0 \leq k \leq k_{small} + 16$,

$$\sup_{\mathcal{T}} u^{1+\delta_{dec}} |\mathfrak{d}^k \underline{\alpha}| \lesssim \epsilon_0, \quad (7.1.17)$$

and on the last slice Σ_* for all $k \leq k_{small} + 16$

$$\int_{\Sigma_*(\tau, \tau_*)} |\mathfrak{d}^k \underline{\alpha}|^2 \lesssim \epsilon_0^2(1+\tau)^{-2-2\delta_{dec}}. \quad (7.1.18)$$

Ref 3. In view of the bootstrap assumptions **BA-D** and the pointwise estimates of Proposition 3.4.5 for the curvature components, which themselves follow from **BA-E**, we have in $^{(ext)}\mathcal{M}$,

³Recall (see Remark 2.4.9) that the quantity \mathfrak{q} we are working with is defined relative to the global frame of Proposition 3.5.5.

⁴In fact, the corresponding estimate in Theorem M3 holds on $^{(int)}\mathcal{M}$, and hence in particular on \mathcal{T} since $\mathcal{T} \subset ^{(int)}\mathcal{M}$.

i. For all $0 \leq k \leq k_{small}$,

$$\begin{aligned}
\|\beta\|_{\infty,k} &\lesssim \epsilon \min \left\{ r^{-3}(u+2r)^{-\frac{1}{2}-\delta_{dec}}, r^{-2}(u+2r)^{-1-\delta_{dec}} \right\}, \\
\|e_3\beta\|_{\infty,k-1} &\lesssim \epsilon \min \left\{ r^{-4}(u+2r)^{-\frac{1}{2}-\delta_{dec}}, r^{-3}(u+2r)^{-1-\delta_{dec}} \right\}, \\
\left\| \left(\check{\rho}, \bar{\rho} + \frac{2m}{r^3} \right) \right\|_{\infty,k} &\lesssim \epsilon \min \left\{ r^{-3}u^{-\frac{1}{2}-\delta_{dec}}, r^{-2}u^{-1-\delta_{dec}} \right\}, \\
\left\| e_3 \left(\check{\rho}, \bar{\rho} + \frac{2m}{r^3} \right) \right\|_{\infty,k-1} &\lesssim \epsilon r^{-3}u^{-1-\delta_{dec}}, \\
\left\| \check{\mu}, \bar{\mu} - \frac{2m}{r^3} \right\|_{\infty,k} &\lesssim \epsilon r^{-3}u^{-1-\delta_{dec}}, \\
\|\underline{\beta}\|_{\infty,k} &\lesssim \epsilon r^{-2}u^{-1-\delta_{dec}}.
\end{aligned} \tag{7.1.19}$$

Since $K = -\rho - \frac{1}{4}\kappa\underline{\kappa} + \frac{1}{4}\vartheta\underline{\vartheta} = \frac{1}{r^2} - (\rho - \bar{\rho}) - \frac{1}{4}(\kappa\underline{\kappa} - \bar{\kappa}\bar{\kappa}) + \text{l.o.t.}$ we also deduce for all $0 \leq k \leq k_{small}$,

$$\left\| K - \frac{1}{r^2} \right\|_{\infty,k} \lesssim \epsilon \min \left\{ r^{-3}u^{-\frac{1}{2}-\delta_{dec}}, r^{-2}u^{-1-\delta_{dec}} \right\}.$$

ii. For all $k \leq k_{large} - 5$,

$$\begin{aligned}
r^{\frac{7}{2} + \frac{\delta_B}{2}} \left(\|\alpha\|_{\infty,k} + \|\beta\|_{\infty,k} \right) &\lesssim \epsilon, \\
r^3 \|\check{\rho}\|_{\infty,k} + r^2 \|\underline{\beta}\|_{\infty,k} + r \|\underline{\alpha}\|_{\infty,k} &\lesssim \epsilon.
\end{aligned} \tag{7.1.20}$$

Remark 7.1.3. In view of the control of averages Lemma 3.4.1 we have in fact better estimates for the scalars,

$$\bar{\kappa} - \frac{2}{r}, \quad \bar{\kappa} + \frac{2\Upsilon}{r}, \quad \bar{\omega} - \frac{m}{r^2}, \quad \bar{\rho} + \frac{2m}{r^3}.$$

In particular they can be estimated by ϵ replaced by ϵ_0 in **Ref 1**.

Remark 7.1.4. Note that $r(\check{\rho}, \bar{\rho} + \frac{2m}{r^3}), r(K - \frac{1}{r^2})$ behave as Γ_g . For convenience we shall just simply add them to Γ_g . Similarly $(r\underline{\beta}, \underline{\alpha})$ behave as Γ_b . Thus, our extended Γ_g, Γ_b are

$$\begin{aligned}
\Gamma_g &= \left\{ \check{\kappa}, \vartheta, \underline{\eta}, \zeta, \check{\kappa}, r\check{\rho} \right\} \cup \left\{ \bar{\kappa} - \frac{2}{r}, \bar{\kappa} + \frac{2\Upsilon}{r}, r \left(\bar{\rho} + \frac{2m}{r^3} \right) \right\}, \\
\Gamma_b &= \left\{ \underline{\vartheta}, \eta, \underline{\omega}, \underline{\xi}, \underline{A}, r^{-1}\zeta, r^{-1}\check{\Omega}, r\underline{\beta}, \underline{\alpha} \right\} \cup \left\{ \bar{\omega} - \frac{m}{r^2}, r^{-1}(\bar{\varsigma} - 1), r^{-1}(\bar{\Omega} + \Upsilon) \right\}.
\end{aligned}$$

Note also that we can write $e_3(\Gamma_g) = r^{-1}\mathfrak{D}\Gamma_b$.

7.1.3 Basic Lemmas

Commutation identities

Lemma 7.1.5. *We have, schematically,*

$$\begin{aligned} [\check{\rho}, e_4]\psi &= \Gamma_g \mathfrak{D}_{\check{\rho}} \psi + l.o.t., \\ [\check{\rho}, e_3]\psi &= r\Gamma_b e_3 \psi + \Gamma_b^{\leq 1} \mathfrak{D}_{\check{\rho}} \psi + l.o.t. \end{aligned} \quad (7.1.21)$$

Proof. Follows from Lemma 2.2.13 and the symbolic notation introduced in (7.1.11), see also Remark 7.1.4. \square

Product Estimates

We estimate quadratic error terms with the help of the following,

Lemma 7.1.6. *Let $k_{small} < k_{large}$ and $k_{loss} > 0$ three positive integers verifying the conditions,*

$$k_{loss} < k_{small}, \quad k_{loss} \leq \frac{k_{large} - 5 - k_{small}}{4}. \quad (7.1.22)$$

The following product estimates hold true for all $0 \leq k \leq k_{small} + k_{loss}$,

$$\begin{aligned} \|\Gamma_g \cdot \Gamma_g\|_{\infty, k} + r \left\| \left(\check{\rho}, \beta, \alpha \right) \cdot \Gamma_g \right\|_{\infty, k} &\lesssim \epsilon_0 r^{-\frac{7}{2}} u^{-1-\delta_{dec}}, \\ \left\| \Gamma_g \cdot \left(\Gamma_b, \underline{\alpha} \right) \right\|_{\infty, k} + r \|\Gamma_g \cdot \underline{\beta}\|_{\infty, k} + r \|\check{\rho} \cdot \Gamma_b\|_{\infty, k} \\ &\quad + r^{\frac{5}{4}} \left\| \left(\beta, \alpha \right) \cdot \Gamma_b \right\|_{\infty, k} \lesssim \epsilon_0 r^{-3} u^{-1-\delta_{dec}}, \\ \left\| \left(\Gamma_b, \underline{\alpha} \right) \cdot \Gamma_b \right\|_{\infty, k} + r \|\underline{\beta} \cdot \Gamma_b\|_{\infty, k} &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \\ \|e_3(\Gamma \cdot \beta)\|_{\infty, k} &\lesssim r^{-5} u^{-1-\delta_{dec}}. \end{aligned} \quad (7.1.23)$$

Proof. All estimates are easy to prove in the range $0 \leq k \leq k_{small}$. We shall thus assume that $k_{small} \leq k \leq k_{small} + k_{loss}$. Since $k_{loss} < k_{small}$ we have $k/2 < k_{small}$ for all k in that range. We start with the first estimate. Since $r\check{\rho}$ satisfies the same estimates as Γ_g , and as $r\beta$ and $r\alpha$ satisfy even better estimates, it suffices to prove the first estimate for

$\Gamma_g \cdot \Gamma_g$. For simplicity of notation we write $L := k_{large} - 5$, $S := k_{small}$. By standard interpolation inequalities, for all $S \leq k \leq L$,

$$\|\Gamma_g\|_{\infty, k} \lesssim \|\Gamma_g\|_{\infty, L}^{\frac{k-S}{L-S}} \|\Gamma_g\|_{\infty, S}^{\frac{L-k}{L-S}} \lesssim \epsilon r^{-2} \left[u^{-\frac{1}{2} - \delta_{dec}} \right]^{\frac{L-k}{L-S}}.$$

Therefore, for any $S \leq k \leq L$

$$\begin{aligned} \|\Gamma_g \cdot \Gamma_g\|_{\infty, k} &\lesssim \|\Gamma_g\|_{\infty} \|\Gamma_g\|_{\infty, k} \lesssim \|\Gamma_g\|_{\infty} \epsilon r^{-2} \left[u^{-\frac{1}{2} - \delta_{dec}} \right]^{\frac{L-k}{L-S}} \\ &\lesssim \epsilon r^{-2} \|\Gamma_g\|_{\infty}^{\frac{1}{2}} \|\Gamma_g\|_{\infty}^{\frac{1}{2}} u^{-(\frac{1}{2} + \delta_{dec}) \frac{L-k}{L-S}} \\ &\lesssim \epsilon^2 r^{-2} r^{-\frac{1}{2}} u^{-\frac{1}{2}(1 + \delta_{dec})} r^{-1} u^{-\frac{1}{2}(\frac{1}{2} + \delta_{dec})} u^{-(\frac{1}{2} + \delta_{dec}) \frac{L-k}{L-S}} \\ &= \epsilon^2 r^{-\frac{7}{2}} u^{-\frac{3}{4} - \delta_{dec} - (\frac{1}{2} + \delta_{dec}) \frac{L-k}{L-S}}. \end{aligned}$$

Now, we have

$$-\frac{3}{4} - \frac{1}{2} \frac{L-k}{L-S} = -1 - \frac{1}{2} \left(\frac{1}{2} - \frac{k-S}{L-S} \right) = -1 - \frac{1}{2} \left(\frac{1}{2} - \frac{k_{loss}}{k_{large} - 5 - k_{small}} \right) \leq -1$$

where we used the assumption on k_{loss}

$$k_{loss} \leq \frac{k_{large} - 5 - k_{small}}{2}.$$

We infer

$$\|\Gamma_g \cdot \Gamma_g\|_{\infty, k} \lesssim \epsilon^2 r^{-\frac{7}{2}} u^{-1 - \delta_{dec}}.$$

Next, we consider the second estimate. Since $r\check{\rho}$ satisfies the same estimates as Γ_g , and as $\underline{\alpha}$ and $r\check{\beta}$ satisfy the same estimate as Γ_b , it suffices to prove the second estimate for $\Gamma_g \cdot \Gamma_b$ and $(\alpha, \beta) \cdot \Gamma_b$. Starting with,

$$\|\Gamma_b\|_{\infty, k} \lesssim \|\Gamma_b\|_{\infty, L}^{\frac{k-S}{L-S}} \|\Gamma_b\|_{\infty, S}^{\frac{L-k}{L-S}} \lesssim \epsilon r^{-1} \left[u^{-1 - \delta_{dec}} \right]^{\frac{L-k}{L-S}},$$

we deduce,

$$\begin{aligned} \|\Gamma_g \cdot \Gamma_b\|_{\infty, k} &\lesssim \|\Gamma_g\|_{\infty} \|\Gamma_b\|_{\infty, k} + \|\Gamma_b\|_{\infty} \|\Gamma_g\|_{\infty, k} \\ &\lesssim \epsilon^2 r^{-2} u^{-\frac{1}{2} - \delta_{dec}} r^{-1} \left[u^{-1 - \delta_{dec}} \right]^{\frac{L-k}{L-S}} + \epsilon^2 r^{-1} u^{-1 - \delta_{dec}} r^{-2} \left[u^{-\frac{1}{2} - \delta_{dec}} \right]^{\frac{L-k}{L-S}} \\ &\lesssim \epsilon^2 r^{-3} u^{-1 - \delta_{dec}}. \end{aligned}$$

Also,

$$\begin{aligned}
\left\| (\alpha, \beta) \cdot \Gamma_b \right\|_{\infty, k} &\lesssim \|(\alpha, \beta)\|_{\infty} \|\Gamma_b\|_{\infty, k} + \|\Gamma_b\|_{\infty} \|(\alpha, \beta)\|_{\infty, k} \\
&\lesssim \epsilon^2 r^{-\frac{13}{4}} u^{-\frac{1}{4} - \delta_{dec}} r^{-1} \left[u^{-1 - \delta_{dec}} \right]^{\frac{L-k}{L-S}} + \epsilon^2 r^{-1} u^{-1 - \delta_{dec}} r^{-\frac{7}{2}} \\
&\lesssim \epsilon^2 r^{-\frac{17}{4}} u^{-1 - \delta_{dec}}.
\end{aligned}$$

Finally, the estimates⁵ for $\Gamma_b \cdot \Gamma_b$, $\underline{\alpha} \cdot \Gamma_b$, $\underline{\beta} \cdot \Gamma_b$ as well as $e_3(\Gamma \cdot \beta)$ are easier and left to the reader. \square

Elliptic Estimates

We shall often make use of the results of Proposition 2.1.30 and Lemma 2.1.35 which we rewrite as follows with respect to the L^2 based $\mathfrak{h}_k(S)$ spaces introduced in Definition 2.1.36.

Lemma 7.1.7. *Under the assumptions **Ref1** – **Ref3** the following elliptic estimates hold true for the Hodge operators $\mathfrak{d}_1, \mathfrak{d}_2, \mathfrak{d}_1^*, \mathfrak{d}_2^*$, for all $k \leq k_{small} + 20$.*

1. If $f \in \mathfrak{s}_1(S)$,

$$\|\mathfrak{d}f\|_{\mathfrak{h}_k(S)} + \|f\|_{\mathfrak{h}_k(S)} \lesssim r \|\mathfrak{d}_1 f\|_{\mathfrak{h}_k(S)}.$$

2. If $f \in \mathfrak{s}_2(S)$,

$$\|\mathfrak{d}f\|_{\mathfrak{h}_k(S)} + \|f\|_{\mathfrak{h}_k(S)} \lesssim r \|\mathfrak{d}_2 f\|_{\mathfrak{h}_k(S)}.$$

3. If $f \in \mathfrak{s}_0(S)$,

$$\|\mathfrak{d}f\|_{\mathfrak{h}_k(S)} \lesssim r \|\mathfrak{d}_1^* f\|_{\mathfrak{h}_k(S)}.$$

4. If $f \in \mathfrak{s}_1(S)$,

$$\|f\|_{\mathfrak{h}_{k+1}(S)} \lesssim r \|\mathfrak{d}_2^* f\|_{\mathfrak{h}_k(S)} + r^{-2} \left| \int_S e^{\Phi} f \right|.$$

5. If $f \in \mathfrak{s}_1(S)$,

$$\left\| f - \frac{\int_S f e^{\Phi}}{\int_S e^{2\Phi}} e^{\Phi} \right\|_{\mathfrak{h}_{k+1}(S)} \lesssim r \|\mathfrak{d}_2^* f\|_{\mathfrak{h}_k(S)}.$$

⁵Note also that in view of Remark 7.1.4 we can write $\underline{\beta} \in r^{-1}\Gamma_b$, $\underline{\alpha} \in \Gamma_b$.

7.1.4 Main equations

The proof of Theorem **M4** relies heavily on the null structure and null Bianchi identities derived in section 2.2.4, see Propositions 2.2.8. We also rely on Proposition 2.2.18 for equations verified by the check quantities. We rewrite them below in a schematic form.

Proposition 7.1.8 (Transport equations for checked quantities). *We have the following transport equations in the e_4 direction,*

$$\begin{aligned}
e_4\check{\kappa} + \bar{\kappa}\check{\kappa} &= \Gamma_g \cdot \Gamma_g, \\
e_4\check{\underline{\kappa}} + \frac{1}{2}\bar{\kappa}\check{\underline{\kappa}} + \frac{1}{2}\check{\kappa}\bar{\underline{\kappa}} &= -2\check{\phi}_1\zeta + 2\check{\rho} + \Gamma_g \cdot \Gamma_b, \\
e_4\check{\underline{\omega}} &= \check{\rho} + \Gamma_g \cdot \Gamma_b, \\
e_4\check{\rho} + \frac{3}{2}\bar{\kappa}\check{\rho} + \frac{3}{2}\bar{\rho}\check{\kappa} &= \check{\phi}_1\beta + \Gamma_b \cdot \alpha + \Gamma_g \cdot \beta + \check{\kappa} \cdot \check{\rho}, \\
e_4\check{\underline{\mu}} + \frac{3}{2}\bar{\kappa}\check{\underline{\mu}} + \frac{3}{2}\bar{\underline{\mu}}\check{\kappa} &= r^{-1}\Gamma_g \cdot \check{\phi}^{\leq 1}\Gamma_g.
\end{aligned} \tag{7.1.24}$$

Also, we have in the e_3 direction,

$$\begin{aligned}
e_3\check{\underline{\kappa}} &= r^{-1}\check{\phi}^{\leq 1}\Gamma_b + \Gamma_b \cdot \check{\phi}^{\leq 1}\Gamma_b, \\
e_3\check{\rho} &= r^{-2}\check{\phi}^{\leq 1}\Gamma_b + r^{-1}\Gamma_b \cdot \check{\phi}^{\leq 1}\Gamma_b.
\end{aligned} \tag{7.1.25}$$

Proof. The statements follow from the precise formulas of Proposition 2.2.18 and the symbolic notation in (7.1.11). We also use the convention made in Remark 7.1.4 according to which we write $r\check{\rho}, r\check{\underline{\mu}} \in \Gamma_g$, $(r\check{\underline{\beta}}, \check{\underline{\alpha}}) \in \Gamma_b$ and $e_3(\Gamma_g) = r^{-1}(\partial\Gamma_b)$. \square

7.1.5 Equations involving \mathfrak{q}

Recall that our main quantity \mathfrak{q} has been introduced in Definition 2.3.12 with respect to the global frame of Proposition 3.5.5 (see Remark 2.4.9). The passage from the geodesic frame (e_3, e_θ, e_4) of ${}^{(ext)}\mathcal{M}$ to the global frame (e'_3, e'_θ, e'_4) is given by

$$e'_4 = \Upsilon \left(e_4 + fe_\theta + \frac{1}{4}f^2e_3 \right), \quad e'_\theta = e_\theta + \frac{1}{2}fe_3, \quad e'_3 = \Upsilon^{-1}e_3. \tag{7.1.26}$$

with a reduced scalar f which was constructed in Proposition 3.4.6. We recall below the main relevant statements of Proposition 3.4.6 in connection to the construction of the global frame.

Proposition 7.1.9. *Under assumptions Ref 1-2 on $^{(ext)}\mathcal{M}$ there exists a frame transformation of the form, (7.1.26) verifying the following properties⁶:*

1. *Everywhere in $^{(ext)}\mathcal{M}$ we have $\xi' = 0$.*
2. *The transition function f verifies, relative to the background frame (e_3, e_θ, e_4) , the estimates⁷*

$$\begin{aligned} |\mathfrak{d}^k f| &\lesssim \frac{\epsilon}{ru^{\frac{1}{2}+\delta_{dec}-2\delta_0} + u^{1+\delta_{dec}-2\delta_0}}, \quad \text{for } k \leq k_{small} + 20 \text{ on } ^{(ext)}\mathcal{M}, \\ |\mathfrak{d}^{k-1} e'_3 f| &\lesssim \frac{\epsilon}{ru^{1+\delta_{dec}-2\delta_0}} \quad \text{for } k \leq k_{small} + 20 \text{ on } ^{(ext)}\mathcal{M}. \end{aligned} \quad (7.1.27)$$

3. *The primed Ricci coefficients and curvature components verify⁸*

$$\begin{aligned} \max_{0 \leq k \leq k_{small} + k_{loss}} \sup_{^{(ext)}\mathcal{M}} &\left\{ \left(r^2 u^{\frac{1}{2}+\delta_{dec}-2\delta_0} + ru^{1+\delta_{dec}-2\delta_0} \right) |\mathfrak{d}^k \Gamma'_g| + ru^{1+\delta_{dec}-2\delta_0} |\mathfrak{d}^k \Gamma'_b| \right. \\ &+ r^2 u^{1+\delta_{dec}-2\delta_0} \left| \mathfrak{d}^{k-1} e'_3 \left(\kappa' - \frac{2\Upsilon}{r}, \underline{\kappa}' + \frac{2}{r}, \vartheta', \zeta', \underline{\eta}', \eta' \right) \right| \\ &+ \left(r^3 (u+2r)^{\frac{1}{2}+\delta_{dec}-2\delta_0} + r^2 (u+2r)^{1+\delta_{dec}-2\delta_0} \right) \left(|\mathfrak{d}^k \alpha'| + |\mathfrak{d}^k \beta'| \right) \\ &+ \left(r^3 (2r+u)^{1+\delta_{dec}} + r^4 (2r+u)^{\frac{1}{2}+\delta_{dec}-2\delta_0} \right) |\mathfrak{d}^{k-1} e'_3(\alpha')| \\ &+ \left(r^3 u^{1+\delta_{dec}} + r^4 u^{\frac{1}{2}+\delta_{dec}-2\delta_0} \right) |\mathfrak{d}^{k-1} e'_3(\beta')| \\ &+ \left(r^3 u^{\frac{1}{2}+\delta_{dec}-2\delta_0} + r^2 ru^{1+\delta_{dec}-2\delta_0} \right) |\mathfrak{d}^k \check{\rho}'| \\ &\left. + u^{1+\delta_{dec}-2\delta_0} \left(r^2 |\mathfrak{d}^k \underline{\beta}'| + r |\mathfrak{d}^k \underline{\alpha}'| \right) \right\} \lesssim \epsilon. \end{aligned}$$

We have the following analog of Proposition 2.3.13.

Proposition 7.1.10. *We have, relative to the background frame of $^{(ext)}\mathcal{M}$,*

$$r^4 \left(\not\phi_2^* \not\phi_1^* \rho + \frac{3}{4} \underline{\kappa} \rho \vartheta + \frac{3}{4} \kappa \rho \underline{\vartheta} \right) = \mathfrak{q} + Err \quad (7.1.28)$$

with error term expressed schematically in the form

$$Err = r^2 \not\phi^{\leq 2} (\Gamma_b \cdot \Gamma_g). \quad (7.1.29)$$

⁶We denote by primes the Ricci and curvature components w.r.t. to the primed frame.

⁷In fact, the estimates hold for $k_{small} + k_{loss}$, see Proposition 3.4.6, and we choose here $k_{loss} = 20$.

⁸Note that u and r here are the outgoing optical function and area radius of the foliation of $^{(ext)}\mathcal{M}$.

Proof. We make use of Proposition 2.3.13. Recall (see Remark 2.4.9) that the quantity \mathfrak{q} we are working with is defined relative to the global frame of Proposition 3.5.5. We thus write⁹,

$$\begin{aligned}\mathfrak{q} &= r^4 \left((\mathfrak{d}_2^*)' (\mathfrak{d}_1^*)' \rho' + \frac{3}{4} \underline{\kappa}' \rho' \vartheta' + \frac{3}{4} \kappa' \rho' \underline{\vartheta} \right) + \text{Err}', \\ \text{Err}' &= r^4 e_3' \eta' \cdot \beta' + r^2 \mathfrak{d}^{\leq 1}(\Gamma_b \cdot \Gamma_g),\end{aligned}$$

where the primes refer to the global frame in which \mathfrak{q} was defined. Since in that frame $e_3' \eta' \in r^{-1} \mathfrak{d} \Gamma_b$ and $\beta' \in r^{-1} \Gamma_g$ we can simplify and write,

$$\text{Err}' = r^2 \mathfrak{d}^{\leq 1}(\Gamma_b \cdot \Gamma_g).$$

We also have in view of Proposition 2.3.4

$$\begin{aligned}\rho' &= \rho + f \underline{\beta} + O(f^2 \underline{\alpha}), \\ \underline{\beta}' &= \underline{\beta} + \frac{1}{2} f \underline{\alpha}, \\ \underline{\alpha}' &= \underline{\alpha}, \\ \underline{\kappa}' &= \underline{\kappa} + f \underline{\xi}, \\ \kappa' &= \kappa + \mathfrak{d}_1'(f) + f(\zeta + \eta) + O(r^{-1} f^2), \\ \vartheta' &= \vartheta - \mathfrak{d}_2'(f) + f(\zeta + \eta) + O(r^{-1} f^2), \\ \underline{\vartheta}' &= \underline{\vartheta} + f \underline{\xi}.\end{aligned}$$

Note that

$$(\mathfrak{d}_1^*)' \rho = -e_\theta'(\rho) = -e_\theta \rho - \frac{1}{2} f e_3 \rho = \mathfrak{d}_1^* \rho - \frac{1}{2} f e_3 \rho.$$

We deduce,

$$\begin{aligned}(\mathfrak{d}_2^*)' (\mathfrak{d}_1^*)' \rho' &= (\mathfrak{d}_2^*)' (\mathfrak{d}_1^*)' \rho + (\mathfrak{d}_2^*)' (\mathfrak{d}_1^*)' (\Gamma_b \cdot \Gamma_g) + \text{l.o.t.} \\ &= (\mathfrak{d}_2^*)' \left(\mathfrak{d}_1^* - \frac{1}{2} f e_3 \right) \rho + r^{-2} \mathfrak{d}^{\leq 2}(\Gamma_b \cdot \Gamma_g) \\ &= \mathfrak{d}_2^* \left(\mathfrak{d}_1^* - \frac{1}{2} f e_3 \right) \rho + r^{-2} \mathfrak{d}^{\leq 2}(\Gamma_b \cdot \Gamma_g) \\ &= \mathfrak{d}_2^* \mathfrak{d}_1^* \rho - \frac{1}{2} \mathfrak{d}_2^* f e_3 \rho + r^{-2} \mathfrak{d}^{\leq 2}(\Gamma_b \cdot \Gamma_g).\end{aligned}$$

⁹The values of r and r' differ only by lower order terms which do not affect the result.

Similarly,

$$\begin{aligned}\underline{\kappa}'\rho'\vartheta' &= \rho\underline{\kappa}(\vartheta - \phi_2^*f) + r^{-3}\mathfrak{D}^{\leq 1}(\Gamma_g \cdot \Gamma_g), \\ \kappa'\rho'\underline{\vartheta}' &= \kappa\rho\underline{\vartheta} + r^{-3}\mathfrak{D}^{\leq 1}(\Gamma_b \cdot \Gamma_g).\end{aligned}$$

We deduce,

$$\begin{aligned}(\phi_2^*)'(\phi_1^*)'\rho' + \frac{3}{4}\underline{\kappa}'\rho'\vartheta' + \frac{3}{4}\kappa'\rho'\underline{\vartheta}' &= \phi_2^*\phi_1^*\rho + \frac{3}{4}\underline{\kappa}\rho\vartheta + \frac{3}{4}\kappa\rho\underline{\vartheta} - \frac{1}{2}\phi_2^*f\left(e_3\rho + \frac{3}{2}\underline{\kappa}\rho\right) \\ &\quad + r^{-2}\mathfrak{D}^{\leq 2}(\Gamma_b \cdot \Gamma_g).\end{aligned}$$

Note that,

$$\phi_2^*f\left(e_3\rho + \frac{3}{2}\underline{\kappa}\rho\right) = \phi_2^*f\left(\phi_1\underline{\beta} - \frac{1}{2}\vartheta\underline{\alpha} + \text{l.o.t.}\right) = r^{-2}\mathfrak{D}^{\leq 2}(\Gamma_g \cdot \Gamma_b).$$

Hence

$$(\phi_2^*)'(\phi_1^*)'\rho' + \frac{3}{4}\underline{\kappa}'\rho'\vartheta' + \frac{3}{4}\kappa'\rho'\underline{\vartheta}' = \phi_2^*\phi_1^*\rho + \frac{3}{4}\underline{\kappa}\rho\vartheta + \frac{3}{4}\kappa\rho\underline{\vartheta} + r^{-2}\mathfrak{D}^{\leq 2}(\Gamma_b \cdot \Gamma_g).$$

This concludes the proof of Proposition 7.1.10. \square

We shall also need the following analogue of Proposition 2.3.14.

Proposition 7.1.11. *The following identity holds true in $^{(ext)}\mathcal{M}$, with respect to its background frame*

$$\begin{aligned}e_3(r\mathbf{q}) &= r^5\left\{\phi_2^*\phi_1^*\phi_1\underline{\beta} - \frac{3}{2}\rho\phi_2^*\phi_1^*\underline{\kappa} - \frac{3}{2}\underline{\kappa}\rho\phi_2^*\zeta - \frac{3}{2}\kappa\rho\underline{\alpha} + \frac{3}{4}(2\rho^2 - \kappa\underline{\kappa}\rho)\underline{\vartheta}\right\} \\ &\quad + \text{Err}[e_3(r\mathbf{q})],\end{aligned}\tag{7.1.30}$$

where

$$\text{Err}[e_3(r\mathbf{q})] = r^3\mathfrak{D}^{\leq 3}(\Gamma_b \cdot \Gamma_g).\tag{7.1.31}$$

Proof. We start with the result of Proposition 2.3.14 which we write in the form,

$$\begin{aligned}(r')^{-5}e_3'(r'\mathbf{q}) &= (\phi_2^*\phi_1^*\phi_1)'\underline{\beta}' - \frac{3}{2}\kappa'\rho'\underline{\alpha}' - \frac{3}{2}\rho'(\phi_2^*\phi_1^*)'\underline{\kappa}' - \frac{3}{2}\underline{\kappa}'\rho'(\phi_2^*)'\zeta' + \frac{3}{4}(2(\rho')^2 - \kappa'\underline{\kappa}'\rho')\underline{\vartheta}' \\ &\quad + (r')^{-5}\text{Err}[e_3'(r'\mathbf{q})] \\ \text{Err}'[e_3'(r'\mathbf{q})] &= r'\Gamma_b\mathbf{q} + r^5\mathfrak{D}'^{\leq 1}(e_3'\eta' \cdot \beta') + r'^3\mathfrak{D}'^{\leq 2}(\Gamma_b \cdot \Gamma_g).\end{aligned}$$

Since $e'_3\eta' \in r^{-1}\Gamma_b$ and $\mathfrak{q} \in \Gamma_b$, we deduce,

$$\text{Err}'[e'_3(r'\mathfrak{q})] = r^3\mathfrak{d}^{\leq 2}(\Gamma_b \cdot \Gamma_g).$$

Now, in view of Proposition 2.3.4,

$$(\mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1)' \underline{\beta}' = (\mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1)' \left(\beta + \frac{1}{2} f \underline{\alpha} \right) = (\mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1)' \underline{\beta} + r^{-2} \mathfrak{P}^3(\Gamma_b \cdot \Gamma_g).$$

Proceeding in the same manner with all other terms we find,

$$\begin{aligned} & (\mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1)' \underline{\beta}' - \frac{3}{2} \kappa' \rho' \underline{\alpha}' - \frac{3}{2} \rho' (\mathfrak{d}_2^* \mathfrak{d}_1^*)' \underline{\kappa}' - \frac{3}{2} \underline{\kappa}' \rho' (\mathfrak{d}_2^*)' \underline{\zeta}' + \frac{3}{4} (2(\rho')^2 - \kappa' \underline{\kappa}' \rho') \underline{\vartheta}' \\ &= \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \underline{\beta} - \frac{3}{2} \kappa \rho \underline{\alpha} - \frac{3}{2} \rho \mathfrak{d}_2^* \mathfrak{d}_1^* \underline{\kappa} - \frac{3}{2} \underline{\kappa} \rho \mathfrak{d}_2^* \underline{\zeta} + \frac{3}{4} (2\rho^2 - \kappa \underline{\kappa} \rho) \underline{\vartheta} + r^{-2} \mathfrak{P}^{\leq 3}(\Gamma_b \cdot \Gamma_g) \end{aligned}$$

from which the result easily follows. \square

7.1.6 Additional equations

The following proposition is an immediate corollary of Proposition 2.2.19.

Proposition 7.1.12. *We have, schematically,*

$$\begin{aligned} 2 \mathfrak{d}_1^* \underline{\omega} &= \left(\frac{1}{2} \underline{\kappa} + 2 \underline{\omega} \right) \eta + e_3(\zeta) - \underline{\beta} - \frac{1}{2} \kappa \underline{\xi} + r^{-1} \Gamma_g + \Gamma_b^2, \\ 2 \mathfrak{d}_2 \mathfrak{d}_2^* \eta &= \kappa (-e_3(\zeta) + \underline{\beta}) - e_3(e_\theta(\kappa)) + r^{-2} \mathfrak{P}^{\leq 1} \Gamma_g + r^{-1} \mathfrak{P}^{\leq 1}(\Gamma_b \cdot \Gamma_b), \\ 2 \mathfrak{d}_2 \mathfrak{d}_2^* \underline{\xi} &= \underline{\kappa} (e_3(\zeta) - \underline{\beta}) - e_3(e_\theta(\underline{\kappa})) + r^{-2} \mathfrak{P}^{\leq 1} \Gamma_g + r^{-1} \mathfrak{P}^{\leq 1}(\Gamma_b \cdot \Gamma_b). \end{aligned}$$

Remark 7.1.13. *Note that in fact $\Gamma_g = \{\underline{\kappa}, \vartheta, \zeta, \underline{\kappa}, r\dot{\rho}\}$ and $\Gamma_b = \{\underline{\vartheta}, \eta, \underline{\xi}, \underline{\omega}, r\beta, \underline{\alpha}\}$ in the derivation of this proposition. It is important to note also that the terms denoted schematically by $\mathfrak{P}(\Gamma_b \cdot \Gamma_b)$ do not contain derivatives of $\underline{\omega}$.*

The following corollary of Proposition 7.1.12 which will be very useful later on.

Proposition 7.1.14. *The following identities hold true on Σ_* .*

$$\begin{aligned} 2 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2 \mathfrak{d}_2^* \eta &= \kappa \left(e_3(\mathfrak{d}_2^* \mathfrak{d}_1^* \mu) + 2 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \underline{\beta} \right) - \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 e_3(e_\theta(\kappa)) \\ &\quad + r^{-5} \mathfrak{P}^{\leq 4} \Gamma_g + r^{-4} \mathfrak{P}^{\leq 4}(\Gamma_b \cdot \Gamma_b) + l.o.t. \end{aligned} \tag{7.1.32}$$

$$\begin{aligned} 2 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2 \mathfrak{d}_2^* \underline{\xi} &= e_3 \left((\mathfrak{d}_2^* \mathfrak{d}_2 + 2K) \mathfrak{d}_2^* \mathfrak{d}_1^* \underline{\kappa} \right) - \underline{\kappa} \left(e_3(\mathfrak{d}_2^* \mathfrak{d}_1^* \mu) + 2 \mathfrak{d}_2^* \mathfrak{d}_1^* \mathfrak{d}_1 \underline{\beta} \right) \\ &\quad + r^{-5} \mathfrak{P}^{\leq 4} \Gamma_g + r^{-4} \mathfrak{P}^{\leq 4}(\Gamma_b \cdot \Gamma_b) + l.o.t. \end{aligned} \tag{7.1.33}$$

Remark 7.1.15. Here, as in the remark following Proposition 7.1.12, $\Gamma_g = \{\check{\kappa}, \vartheta, \zeta, \check{\kappa}, r\check{\rho}\}$ and $\Gamma_b = \{\underline{\vartheta}, \eta, \underline{\xi}, \underline{\omega}, r\underline{\beta}, \underline{\alpha}\}$. The quadratic terms denoted l.o.t. are lower order both in terms of decay in r, u as well in terms of number of derivatives. They also contain only angular derivatives \mathfrak{D} and not e_3 nor e_4 .

Proof. We make use of Proposition 7.1.12. We shall also make use of the conventions mentioned in Remark 7.1.4, i.e. $\check{\rho}, \check{\mu} \in r^{-1}\Gamma_g$, $\underline{\beta} \in r^{-1}\Gamma_b$, $\underline{\alpha} \in \Gamma_b$.

We start with,

$$2 \mathfrak{d}_2 \mathfrak{d}_2^* \eta = \kappa (-e_3(\zeta) + \underline{\beta}) - e_3(e_\theta(\kappa)) + r^{-2} \mathfrak{D}^{\leq 1} \Gamma_g + r^{-1} \mathfrak{D}(\Gamma_b \cdot \Gamma_b)$$

We apply $\mathfrak{d}_1^* \mathfrak{d}_1$ to derive,

$$\begin{aligned} 2 \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2 \mathfrak{d}_2^* \eta &= \kappa (-\mathfrak{d}_1^* \mathfrak{d}_1 e_3(\zeta) + \mathfrak{d}_1^* \mathfrak{d}_1 \underline{\beta}) - \mathfrak{d}_1^* \mathfrak{d}_1 e_3(e_\theta(\kappa)) + r^{-4} \mathfrak{D}^{\leq 3} \Gamma_g + r^{-3} \mathfrak{D}^3(\Gamma_b \cdot \Gamma_b) \\ &= \kappa (-e_3(\mathfrak{d}_1^* \mathfrak{d}_1(\zeta) + \mathfrak{d}_1^* \mathfrak{d}_1 \underline{\beta}) - \mathfrak{d}_1^* \mathfrak{d}_1 e_3(e_\theta(\kappa))) \\ &\quad - \kappa[\mathfrak{d}_1^* \mathfrak{d}_1, e_3]\zeta + r^{-4} \mathfrak{D}^{\leq 3} \Gamma_g + r^{-3} \mathfrak{D}^3(\Gamma_b \cdot \Gamma_b) \end{aligned}$$

Making use of the commutation formula, see Lemma 7.1.5, and the null structure equations for $e_3\zeta, e_4\zeta$,

$$[\mathfrak{d}_1, e_3]\zeta = -\eta e_3\zeta + r^{-2} \mathfrak{D}\zeta + \Gamma_b e_4\zeta + \text{l.o.t.} = r^{-1}\Gamma_b \cdot \Gamma_b + r^{-2} \mathfrak{D}\Gamma_g + \text{l.o.t.}$$

we deduce, schematically,

$$\begin{aligned} [\mathfrak{d}_1^* \mathfrak{d}_1, e_3]\zeta &= \mathfrak{d}_1^*[\mathfrak{d}_1, e_3]\zeta + [\mathfrak{d}_1^*, e_3]\mathfrak{d}_1\zeta \\ &= r^{-1} \mathfrak{D}(r^{-1}\Gamma_b \cdot \Gamma_b + r^{-2} \mathfrak{D}\zeta + \text{l.o.t.}) + \Gamma_b e_3 \mathfrak{d}_1\zeta + r^{-2} \mathfrak{d}_1\zeta + \text{l.o.t.} \\ &= r^{-2} \mathfrak{D}(\Gamma_b \cdot \Gamma_b) + r^{-3} \mathfrak{D}^2\zeta + \Gamma_b (\mathfrak{d}_1 e_3\zeta + \Gamma_b e_3\zeta + r^{-2} \mathfrak{D}\zeta) + \text{l.o.t.} \\ &= r^{-2} \mathfrak{D}(\Gamma_b \cdot \Gamma_b) + r^{-2} \Gamma_b \mathfrak{D}(\mathfrak{D}\Gamma_b) + r^{-1} \Gamma_b \cdot \Gamma_b \cdot \Gamma_b + r^{-4} \mathfrak{D}^2 \Gamma_g \\ &= r^{-2} \mathfrak{D}(\Gamma_b \mathfrak{D}^{\leq 1} \Gamma_b) + r^{-4} \mathfrak{D}^2 \Gamma_g + \text{l.o.t.} \end{aligned}$$

Hence,

$$\begin{aligned} 2 \mathfrak{d}_1^* \mathfrak{d}_1 \mathfrak{d}_2 \mathfrak{d}_2^* \eta &= \kappa \left(-e_3(\mathfrak{d}_1^* \mathfrak{d}_1 \zeta) + \mathfrak{d}_1^* \mathfrak{d}_1 \underline{\beta} \right) - \mathfrak{d}_1^* \mathfrak{d}_1 e_3(e_\theta(\kappa)) \\ &\quad + r^{-4} \mathfrak{D}^{\leq 3} \Gamma_g + r^{-3} \mathfrak{D}^2(\Gamma_b \cdot \mathfrak{D}\Gamma_b) \end{aligned} \tag{7.1.34}$$

Since $\mu = -\mathfrak{d}_1\zeta - \rho + \frac{1}{4}\vartheta\underline{\vartheta}$, we deduce,

$$\begin{aligned} \mathfrak{d}_1^* \mu &= -\mathfrak{d}_2^* \mathfrak{d}_1\zeta - \mathfrak{d}_1^* \rho + \frac{1}{4} \mathfrak{d}_1^*(\vartheta\underline{\vartheta}), \\ e_3 \mathfrak{d}_1^* \mu &= -e_3(\mathfrak{d}_2^* \mathfrak{d}_1\zeta) - e_3 \mathfrak{d}_1^* \rho + \frac{1}{4} e_3 \mathfrak{d}_1^*(\vartheta\underline{\vartheta}) \\ &= -e_3(\mathfrak{d}_2^* \mathfrak{d}_1\zeta) - \mathfrak{d}_1^* e_3 \rho - [\mathfrak{d}_1^*, e_3]\rho + \frac{1}{4} \mathfrak{d}_1^* e_3(\vartheta\underline{\vartheta}) + \frac{1}{4} [e_3, \mathfrak{d}_1^*](\vartheta \cdot \underline{\vartheta}). \end{aligned}$$

Making use of the equations for $e_3\rho = \underline{\phi}_1\underline{\beta} - \frac{3}{2}\underline{\kappa}\rho + \Gamma_g \cdot \Gamma_b$ and also the equations for¹⁰ $e_4\rho, e_3\vartheta, e_3\underline{\vartheta}, e_4\vartheta, e_4\underline{\vartheta}$ (and writing $\underline{\phi}_1\underline{\beta} = r^{-1}\underline{\phi}\underline{\beta} = r^{-2}\underline{\phi}\Gamma_b$)

$$\begin{aligned} [e_3, \underline{\phi}_1^*]\rho &= \Gamma_b e_3\rho + \Gamma_b e_4\underline{\zeta} + r^{-2}\underline{\phi}\rho = r^{-2}\Gamma_b\underline{\phi}\Gamma_b + r^{-3}\underline{\phi}\Gamma_g + \text{l.o.t.}, \\ [e_3, \underline{\phi}_1^*](\vartheta \cdot \underline{\vartheta}) &= \Gamma_b e_3(\vartheta \cdot \underline{\vartheta}) + \Gamma_b e_4(\vartheta \cdot \underline{\vartheta}) + r^{-2}\underline{\phi}(\vartheta \cdot \underline{\vartheta}) = r^{-2}\underline{\phi}(\Gamma_b \cdot \Gamma_g) + \text{l.o.t.} \end{aligned}$$

We deduce, ignoring the lower order terms,

$$\begin{aligned} e_3 \underline{\phi}_1^* \mu &= -e_3(\underline{\phi}_2^* \underline{\phi}_1 \zeta) - \underline{\phi}_1^* \left(\underline{\phi}_1 \underline{\beta} - \frac{3}{2}\underline{\kappa}\rho + \Gamma_g \cdot \Gamma_b \right) + r^{-2}\Gamma_b \underline{\phi}\Gamma_b + r^{-2}\underline{\phi}(\Gamma_b \Gamma_g) + r^{-3}\underline{\phi}\Gamma_g \\ &= -e_3(\underline{\phi}_2^* \underline{\phi}_1 \zeta) - \underline{\phi}_1^* \underline{\phi}_1 \underline{\beta} + \frac{3}{2}\underline{\kappa} \underline{\phi}_1^* \rho + r^{-3}\underline{\phi}\Gamma_g + r^{-2}\underline{\phi}^{\leq 1}(\Gamma_b \cdot \Gamma_b). \end{aligned}$$

Hence,

$$e_3(\underline{\phi}_1^* \underline{\phi}_1 \zeta) = -e_3(\underline{\phi}_1^* \mu) - \underline{\phi}_1^* \underline{\phi}_1 \underline{\beta} + r^{-3}\underline{\phi}\Gamma_g + r^{-2}\underline{\phi}^{\leq 2}(\Gamma_b \cdot \Gamma_b) + \text{l.o.t.} \quad (7.1.35)$$

and thus, back to (7.1.34),

$$\begin{aligned} 2 \underline{\phi}_1^* \underline{\phi}_1 \underline{\phi}_2 \underline{\phi}_2^* \eta &= \kappa \left(e_3(\underline{\phi}_1^* \mu) + 2 \underline{\phi}_1^* \underline{\phi}_1 \underline{\beta} \right) - \underline{\phi}_1^* \underline{\phi}_1 e_3(e_\theta(\kappa)) \\ &\quad + r^{-4}\underline{\phi}^{\leq 3}\Gamma_g + r^{-3}\underline{\phi}^{\leq 3}(\Gamma_b \cdot \Gamma_b) + \text{l.o.t.} \end{aligned} \quad (7.1.36)$$

Applying $\underline{\phi}_2^*$ and commuting once more with e_3 , i.e.,

$$\begin{aligned} 2 \underline{\phi}_2^* \underline{\phi}_1^* \underline{\phi}_1 \underline{\phi}_2 \underline{\phi}_2^* \eta &= \kappa \left(e_3(\underline{\phi}_2^* \underline{\phi}_1^* \mu) + 2 \underline{\phi}_2^* \underline{\phi}_1^* \underline{\phi}_1 \underline{\beta} \right) - \underline{\phi}_2^* \underline{\phi}_1^* \underline{\phi}_1 e_3(e_\theta(\kappa)) \\ &\quad + \kappa[\underline{\phi}_2^*, e_3] \underline{\phi}_1^* \mu + r^{-1}\underline{\phi}\Gamma_g \cdot \left(e_3(\underline{\phi}_1^* \mu) + 2 \underline{\phi}_1^* \underline{\phi}_1 \underline{\beta} \right) \\ &\quad + r^{-5}\underline{\phi}^{\leq 4}\Gamma_g + r^{-4}\underline{\phi}^{\leq 4}(\Gamma_b \cdot \Gamma_b). \end{aligned} \quad (7.1.37)$$

Note that, in view of (7.1.36) we can write,

$$\begin{aligned} e_3(\underline{\phi}_1^* \mu) &= 2\kappa^{-1} \underline{\phi}_2^* \underline{\phi}_1^* \underline{\phi}_1 e_3(e_\theta(\kappa)) - 2 \underline{\phi}_1^* \underline{\phi}_1 \underline{\beta} + 2\kappa^{-1} \underline{\phi}_1^* \underline{\phi}_1 \underline{\phi}_2 \underline{\phi}_2^* \eta \\ &= r^{-3}\underline{\phi}^{\leq 4}\Gamma_b + \text{l.o.t.} \end{aligned} \quad (7.1.38)$$

Hence,

$$r^{-1}\underline{\phi}\Gamma_g \cdot \left(e_3(\underline{\phi}_1^* \mu) + 2 \underline{\phi}_1^* \underline{\phi}_1 \underline{\beta} \right) = r^{-4}\underline{\phi}\Gamma_g \cdot \underline{\phi}^{\leq 4}\Gamma_b.$$

Similarly,

$$\begin{aligned} [\underline{\phi}_2^*, e_3] \underline{\phi}_1^* \mu &= \Gamma_b \cdot e_3 \underline{\phi}_1^* \mu + \Gamma_b e_4 \underline{\phi}_1^* \mu + r^{-3}\underline{\phi}^2 \mu + \text{l.o.t.} \\ &= r^{-3}\Gamma_b \cdot \underline{\phi}^{\leq 4}\Gamma_b + \Gamma_b (\underline{\phi}_1^* e_4 \mu + [e_4, \underline{\phi}_1^*] \mu) + r^{-4}\underline{\phi}^2 \Gamma_g + \text{l.o.t.} \end{aligned}$$

¹⁰This is to avoid the presence of e_3, e_4 derivatives in the error terms.

Thus, making use of the equation for $e_4\mu$ and combining with the estimate above,

$$\kappa[\not{d}_2^*, e_3] \not{d}_1^* \mu + r^{-1} \not{\partial} \Gamma_g \cdot \left(e_3(\not{d}_1^* \mu) + 2 \not{d}_1^* \not{d}_1 \underline{\beta} \right) = r^{-4} \Gamma_b \cdot \not{\partial}^{\leq 4} \Gamma_b + r^{-5} \not{\partial}^{\leq 2} \Gamma_g.$$

Back to (7.1.37) we deduce,

$$\begin{aligned} 2 \not{d}_2^* \not{d}_1^* \not{d}_1 \not{d}_2 \not{d}_2^* \eta &= \kappa \left(e_3(\not{d}_2^* \not{d}_1^* \mu) + 2 \not{d}_2^* \not{d}_1^* \not{d}_1 \underline{\beta} \right) - \not{d}_2^* \not{d}_1^* \not{d}_1 e_3(e_\theta(\kappa)) \\ &\quad + r^{-5} \not{\partial}^{\leq 4} \Gamma_g + r^{-4} \not{\partial}^{\leq 4}(\Gamma_b \cdot \Gamma_b) \end{aligned}$$

as desired.

To prove the second part we start with the formula for $\not{d}_2 \not{d}_2^* \underline{\xi}$ in Corollary 7.1.12

$$2 \not{d}_2 \not{d}_2^* \underline{\xi} = \underline{\kappa} (e_3(\zeta) - \underline{\beta}) - e_3(e_\theta(\underline{\kappa})) + r^{-2} \not{\partial}^{\leq 1} \Gamma_g + r^{-1} \not{\partial}(\Gamma_b \cdot \Gamma_b).$$

Applying $\not{d}_1^* \not{d}_1$ and proceeding exactly as before in the derivation of (7.1.34) we derive,

$$\begin{aligned} 2 \not{d}_1^* \not{d}_1 \not{d}_2 \not{d}_2^* \underline{\xi} &= -e_3(\not{d}_1^* \not{d}_1 e_\theta(\underline{\kappa})) + \underline{\kappa} (e_3(\not{d}_1^* \not{d}_1 \zeta) - \not{d}_1^* \not{d}_1 \underline{\beta}) \\ &\quad + r^{-4} \not{\partial}^{\leq 3} \Gamma_g + r^{-3} \not{\partial}^2(\Gamma_b \cdot \not{\partial} \Gamma_b). \end{aligned} \tag{7.1.39}$$

Making use of (7.1.35) we deduce, as in (7.1.36),

$$\begin{aligned} 2 \not{d}_1^* \not{d}_1 \not{d}_2 \not{d}_2^* \underline{\xi} &= -e_3(\not{d}_1^* \not{d}_1 e_\theta(\underline{\kappa})) + \underline{\kappa} \left(-e_3(\not{d}_1^* \mu) - 2 \not{d}_1^* \not{d}_1 \underline{\beta} \right) \\ &\quad + r^{-4} \not{\partial}^3 \Gamma_g + r^{-3} \not{\partial}^{\leq 2}(\Gamma_b \cdot \not{\partial} \Gamma_b) + \text{l.o.t.} \end{aligned} \tag{7.1.40}$$

Applying \not{d}_2^* and proceeding as in the derivation of (7.1.37), by making use of (7.1.39) and (7.1.38) we obtain

$$\begin{aligned} 2 \not{d}_2^* \not{d}_1^* \not{d}_1 \not{d}_2 \not{d}_2^* \underline{\xi} &= -e_3(\not{d}_2^* \not{d}_1^* \not{d}_1 e_\theta(\underline{\kappa})) - \underline{\kappa} \left(e_3(\not{d}_2^* \not{d}_1^* \mu) + 2 \not{d}_2^* \not{d}_1^* \not{d}_1 \underline{\beta} \right) \\ &\quad + r^{-5} \not{\partial}^{\leq 4} \Gamma_g + r^{-4} \not{\partial}^{\leq 4}(\Gamma_b \cdot \Gamma_b) + \text{l.o.t.} \end{aligned}$$

The identity $\not{d}_1^* \not{d}_1 = \not{d}_2 \not{d}_2^* + 2K$ yields, together with the bootstrap assumptions,

$$\begin{aligned} 2 \not{d}_2^* \not{d}_1^* \not{d}_1 \not{d}_2 \not{d}_2^* \underline{\xi} &= -e_3((\not{d}_2^* \not{d}_2 + 2K) \not{d}_2^* e_\theta(\underline{\kappa})) - \underline{\kappa} \left(e_3(\not{d}_2^* \not{d}_1^* \mu) + 2 \not{d}_2^* \not{d}_1^* \not{d}_1 \underline{\beta} \right) \\ &\quad + r^{-5} \not{\partial}^{\leq 4} \Gamma_g + r^{-4} \not{\partial}^{\leq 4}(\Gamma_b \cdot \Gamma_b) + \text{l.o.t.} \\ &= e_3((\not{d}_2^* \not{d}_2 + 2K) \not{d}_2^* \not{d}_1^* (\underline{\kappa})) - \underline{\kappa} \left(e_3(\not{d}_2^* \not{d}_1^* \mu) + 2 \not{d}_2^* \not{d}_1^* \not{d}_1 \underline{\beta} \right) \\ &\quad + r^{-5} \not{\partial}^{\leq 4} \Gamma_g + r^{-4} \not{\partial}^{\leq 4}(\Gamma_b \cdot \Gamma_b) + \text{l.o.t.} \end{aligned}$$

as desired. □

7.2 Structure of the proof of Theorem M4

We rephrase the statement of Theorem M4 as follows.

Theorem 7.2.1. *Let $\mathcal{M} = {}^{(int)}\mathcal{M} \cup {}^{(ext)}\mathcal{M}$ be a GCM admissible spacetime¹¹. Under the basic bootstrap assumptions and the results of Theorems M1–M4 (all encoded in **Ref1–Ref4**) the following estimates¹² hold true, for all $k \leq k_{small} + 8$, everywhere on ${}^{(ext)}\mathcal{M}$,*

$$\begin{aligned} \|\Gamma_g\|_{\infty,k} &\lesssim \epsilon_0 \min \left\{ r^{-2} u^{-\frac{1}{2}-\delta_{dec}}, r^{-1} u^{-1-\delta_{dec}} \right\}, \\ \|e_3 \Gamma_g\|_{\infty,k-1} &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \\ \|\Gamma_b\|_{\infty,k} &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \end{aligned} \tag{7.2.1}$$

and,

$$\begin{aligned} \|\beta\|_{\infty,k} &\lesssim \epsilon_0 \min \left\{ r^{-2} (u+2r)^{-1-\delta_{dec}}, r^{-3} (u+2r)^{-\frac{1}{2}-\delta_{dec}} \right\}, \\ \|e_3 \beta\|_{\infty,k-1} &\lesssim \epsilon_0 r^{-3} (u+2r)^{-1-\delta_{dec}}, \\ \|\check{\rho}\|_{\infty,k} &\lesssim \epsilon_0 \min \left\{ r^{-2} u^{-1-\delta_{dec}}, r^{-3} u^{-\frac{1}{2}-\delta_{dec}} \right\}, \\ \|e_3 \check{\rho}\|_{\infty,k} &\lesssim \epsilon_0 r^{-3} u^{-1-\delta_{dec}}, \\ \|\check{\mu}\|_{\infty,k} &\lesssim \epsilon_0 r^{-3} u^{-1-\delta_{dec}}, \\ \|\underline{\beta}\|_{\infty,k} &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}. \end{aligned} \tag{7.2.2}$$

Moreover, everywhere in ${}^{(ext)}\mathcal{M}$,

$$\|\underline{\alpha}\|_{\infty,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}. \tag{7.2.3}$$

Here is a short sketch of the proof of the theorem.

1. *Estimates on Σ_* .* To start with we only have good¹³ estimates for \mathbf{q} , α and $\underline{\alpha}$, according to **Ref2**. To proceed we make use in an essential way of all the GCM conditions (7.1.3)–(7.1.5) on the spacelike boundary Σ_* to estimate all the Ricci and curvature coefficients along Σ_* . The main result is stated in Proposition 7.3.9. The proof is divided in the following intermediary steps.

¹¹In particular the conditions (7.1.1)–(7.1.5) hold on the spacelike boundary Σ_* .

¹²See Remark 7.1.4 for the definition of Γ_g, Γ_b used here.

¹³i.e estimates in terms of ϵ_0 .

- (a) In Proposition 7.3.5 we derive flux type estimates along Σ_* for the quantities $\underline{\beta}, \underline{\theta}, \underline{\eta}, \underline{\xi}$. These estimates take advantage in an essential way of the improved flux estimates for \mathfrak{q} in **Ref 2**, equation (7.1.15). This step also makes use of Proposition 7.1.11 and the identities of Proposition 7.3.4 for $\underline{\eta}, \underline{\xi}$.
- (b) We next estimate the $\ell = 1$ modes of the Ricci and curvature coefficients in Proposition 7.3.8. Besides the information provided by the estimates for $\mathfrak{q}, \alpha, \underline{\alpha}$ and the GCM conditions, an important ingredient in the proof is the vanishing of the $\ell = 1$ mode of $e_\theta(K)$, i.e. $\int e_\theta(K)e^\Phi = 0$. The flux estimates derived in Proposition 7.3.5 play an essential role in deriving the desired estimate for the $\ell = 1$ mode of β .
- (c) We make use of the previous steps to complete the proof all the desired estimates on Σ_* in Proposition 7.3.9. This step also uses, in addition to the GCM conditions, Proposition 7.1.10 relating \mathfrak{q} to $\not{d}_2^* \not{d}_1^* \rho$, the Codazzi equations and elliptic estimates on 2 surfaces.
2. *First Estimates in $^{(ext)}\mathcal{M}$.* We make use of the propagation equations in e_4 and the estimates on Σ_* to derive some of the desired estimates of Theorem 7.2.1, more precisely the better estimates in powers of r for the Γ_g quantities. Note that these estimates decay only like $u^{-1/2-\delta_{dec}}$ in powers of u .
- (a) We first prove the desired estimates for $\check{\kappa}, \check{\mu}$ by simply integrating the corresponding e_4 equations. Note that these estimates are also well behaved in terms of powers of u . This is done in subsection 7.4.3.
- (b) We derive spacetime estimates for all the $\ell = 1$ modes in Lemma 7.4.7. This is done by propagating them from the last slice in the e_4 direction, combined with Codazzi equations and the vanishing of the $\ell = 1$ mode of $e_\theta(K)$.
- (c) We provide all the optimal estimates in terms of powers¹⁴ of r for the quantities $\vartheta, \zeta, \eta, \check{\kappa}, \beta, \check{\rho}$. This is achieved in Proposition 7.4.5 with the help of the estimates on the last slice, the propagation equation for these quantities and the estimates for the $\ell = 1$ modes derived in the previous step.
3. *Optimal u -decay estimates in $^{(ext)}\mathcal{M}$.* We derive all the remaining estimates of Theorem 7.2.1 for all but the quantities $\underline{\xi}, \underline{\omega}, \underline{\check{\Omega}}, \check{\zeta}$. The main remaining difficulty is to get the top decay in powers of u for $\vartheta, \zeta, \eta, \check{\kappa}, \beta, \check{\rho}, \underline{\beta}$. The result is stated in Proposition 7.5.1. We proceed as follows.
- (a) One would like to start with ϑ by using the equation $e_4\vartheta + \kappa\vartheta = -2\alpha$. This unfortunately cannot work by integration¹⁵ starting from the last slice Σ_* .

¹⁴These estimates also provide weak decay in u , i.e. $u^{-\frac{1}{2}-\delta_{dec}}$ decay.

¹⁵It would work however if instead we would integrate from the interior, but we don't possess information about optimal u decay in the interior, for example on the timelike boundary \mathcal{T} of $^{(ext)}\mathcal{M}$.

Similar problems occur for ζ , β , $\check{\rho}$. On the other hand the quantities $\check{\kappa}$ and $\underline{\vartheta}$ could in principle be propagated using their corresponding e_4 equations from Σ_* , but unfortunately they are strongly coupled with the other quantities for which we don't yet have information. For example we have,

$$e_4 \check{\kappa} + \frac{1}{2} \bar{\kappa} \check{\kappa} + \frac{1}{2} \check{\kappa} \bar{\kappa} = -2 \not\partial_1 \zeta + 2 \check{\rho} + \check{\Gamma}_g \cdot \check{\Gamma}_b,$$

and therefore we cannot derive the estimate for $\check{\kappa}$, by integration, before estimating $\not\partial_1 \zeta$ and $\check{\rho}$. To circumvent this difficulty we proceed by an indirect method as follows.

- (b) We can derive optimal decay information on various mixed quantity. For example making use of the equation

$$e_3 \alpha + \left(\frac{1}{2} \underline{\kappa} - 4 \underline{\omega} \right) \alpha = - \not\partial_2^* \beta - \frac{3}{2} \vartheta \rho + 5 \zeta \beta,$$

we infer the desired decay in u for the quantity $\not\partial_2^* \beta - \frac{3}{2} \vartheta \rho$. Other such informations can be derived from the Codazzi equations for ϑ , $\underline{\vartheta}$, the Bianchi identity for $\underline{\beta}$ and the identity (7.1.28) of Lemma 7.1.10.

- (c) We combine the control we have for α , $\check{\kappa}$, $\check{\mu}$ with the control for the mixed quantities mentioned above with a propagation equation for an intermediary quantity,

$$\Xi := r^2 \left(e_\theta(\underline{\kappa}) + 4r \not\partial_1^* \not\partial_1 \zeta - 2r^2 \not\partial_1^* \not\partial_1 \beta \right).$$

We show in the crucial Lemma 7.5.2 that Ξ is also a good mixed quantity, i.e. it has optimal decay in u . It is important to note that this estimate does not depend linearly on $\underline{\alpha}$ for which we only have information on the last slice and \mathcal{T} .

- (d) We can combine the control of Ξ with all other available information mentioned above, to derive good estimates, simultaneously, for $\not\partial_2^* \not\partial_1^* \underline{\kappa}$, $\not\partial_2^* \zeta$ and $\not\partial_2^* \beta$. This is achieved in a sequence of crucial Lemma in subsection 7.5.2. Unfortunately this step is heavily dependent on the estimate of **Ref 2** for $\underline{\alpha}$ and therefore the estimates we derive are only useful on \mathcal{T} .
- (e) We also show that we have good estimates for $\not\partial_2^* \left(\zeta, \not\partial_1^* \check{\kappa}, \beta, \underline{\beta}, \not\partial_1^* \check{\rho} \right)$. To estimate $\check{\kappa}, \zeta, \beta, \underline{\beta}, \check{\rho}$ from $\not\partial_2^* \left(\zeta, \not\partial_1^* \check{\kappa}, \beta, \underline{\beta}, \not\partial_1^* \check{\rho} \right)$ we rely on the elliptic Hodge Lemma 7.1.7 and the control we have for the $\ell = 1$ modes from Lemma 7.4.7 derived earlier. We obtain estimates for $\eta, \vartheta, \underline{\vartheta}$ as well. This establishes all the estimates of Proposition 7.5.1 on \mathcal{T} .

(f) The estimates mentioned above on \mathcal{T} can now be propagated by integrating forward the e_4 null structure and null Bianchi equations. This ends the proof of Proposition 7.5.1 in ${}^{(ext)}\mathcal{M}$.

4. In Proposition 7.6.1 we derive improved decay estimates for $e_3(\beta, \vartheta, \zeta, \underline{\kappa}, \check{\rho})$ and estimates for $\underline{\xi}, \check{\omega}, \check{\Omega}, \check{\zeta}$ in terms of $u^{-1-\delta_{dec}}$ decay. The estimates for $\check{\omega}$ and $\underline{\xi}$ are propagated from the last slice using their e_4 propagation equations. The estimate for $\check{\omega}$ can be easily derived by integrating $e_4(\check{\omega}) = \check{\rho} + \Gamma_g \cdot \Gamma_b$ from the last slice Σ_* . The estimate for $\underline{\xi}$ follows by integrating $e_4(\underline{\xi}) = -e_3(\zeta) + \underline{\beta} - \underline{\kappa}\zeta + \Gamma_b \cdot \Gamma_b$ and making use of the previously derived estimates for $e_3\zeta, \beta, \zeta$. The estimates for $\check{\Omega}, \check{\zeta}$ follow easily from the equations (2.2.19).

7.3 Decay estimates on the last slice Σ_*

7.3.1 Preliminaries

We shall make use of the following norms on Σ_* .

$$\|\psi\|_{\infty, k}^* := \sum_{j \leq k} \|\mathfrak{d}_*^j \psi\|_{L^\infty(S(u, r))}, \quad \mathfrak{d}_*^j = \sum_{j_1 + j_2 \leq j} \mathfrak{d}^{j_1} (\nu_*)^{j_2}. \quad (7.3.1)$$

Recall that $\nu_* = \nu|_{\Sigma_*} = e_3 + a_* e_4$, is the tangent vector to Σ_* and (see (7.1.8) (7.1.9)), along Σ_* ,

$$a_* = -\frac{2}{\varsigma} + \Upsilon - \frac{r}{2} \underline{A} = -\frac{2}{\varsigma} - \underline{\Omega}.$$

Based on our assumptions **Ref 1-2** we deduce

$$\left| a_* + 1 + \frac{2m}{r} \right| \lesssim \epsilon u^{-1-\delta_{dec}}. \quad (7.3.2)$$

As immediate consequence of the commutation Corollary 7.1.5 we derive the following,

Lemma 7.3.1. *We have, schematically,*

$$[\mathfrak{d}, \nu_*] \psi = r \Gamma_b (\nu_* \psi) + \mathfrak{d}^{\leq 1} \Gamma_b \cdot \mathfrak{d} \psi. \quad (7.3.3)$$

Proof. Indeed, see Lemma 7.1.5,

$$\begin{aligned} [\not\partial, e_4]\psi &= \Gamma_g \not\partial_{\nearrow} \psi, \\ [\not\partial, e_3]\psi &= r\Gamma_b e_3 \psi + \Gamma_b \not\partial_{\nearrow} \psi + \text{l.o.t.} \end{aligned} \quad (7.3.4)$$

Hence, since $\not\partial a_* \in r \not\partial \Gamma_b$,

$$\begin{aligned} [\not\partial, \nu_*]\psi &= [\not\partial, e_3 + a_* e_4]\psi = r\Gamma_b e_3 \psi + \Gamma_b \not\partial_{\nearrow} \psi + a_* \Gamma_g \not\partial_{\nearrow} \psi + \not\partial a_* e_4 \psi \\ &= r\Gamma_b (\nu_* \psi - a_* e_4 \psi) + a_* \Gamma_g \not\partial_{\nearrow} \psi + \not\partial a_* e_4 \psi \\ &= r\Gamma_b \nu_* \psi - a_* (\Gamma_b \not\partial_{\nearrow} \psi + \Gamma_g \not\partial_{\nearrow} \psi) + \not\partial \Gamma_b \cdot \not\partial \psi \\ &= r\Gamma_b \nu_* \psi + \not\partial^{\leq 1} \Gamma_b \cdot \not\partial \psi \end{aligned}$$

as desired. \square

To estimate derivatives of the $\ell = 1$ modes on Σ_* we make use of the following.

Lemma 7.3.2. *For every scalar function h we have the formula*

$$\nu_* \left(\int_{\mathbf{S}} h \right) = (\varsigma)^{-1} \int_{\mathbf{S}} \varsigma (\nu_*(h) + (\underline{\kappa} + a_* \kappa) h). \quad (7.3.5)$$

In particular

$$\nu_*(r) = \frac{r}{2} (\varsigma)^{-1} \overline{\varsigma (\underline{\kappa} + a_* \kappa)}. \quad (7.3.6)$$

Proof. We consider the coordinates $u^{\mathbf{S}}, \theta^{\mathbf{S}}$ along Σ_0 with $\nu^{\mathbf{S}}(\theta^{\mathbf{S}}) = 0$. In these coordinates we have,

$$\nu^{\mathbf{S}} = \frac{2}{\varsigma^{\mathbf{S}}} \partial_{u^{\mathbf{S}}}.$$

The lemma follows easily by expressing the volume element of the surfaces $\mathbf{S} \subset \Sigma_0$ with respect to the coordinates $u^{\mathbf{S}}, \theta^{\mathbf{S}}$ (see also the proof of Proposition 2.2.9). \square

Lemma 7.3.3. *Given $\psi \in \mathfrak{s}_1(\mathcal{M})$ we have the formula,*

$$\nu_* \left(\int_{\mathbf{S}} \psi e^{\Phi} \right) = \int_{\mathbf{S}} (\nu_* \psi) e^{\Phi} + \frac{3}{2} (\overline{\kappa} + a_* \overline{\kappa}) \int_{\mathbf{S}} \psi e^{\Phi} + \text{Err}[\psi, \nu_*] \quad (7.3.7)$$

with error term

$$\text{Err}[\psi, \nu_*] = -\varsigma^{-1} \zeta \int_{\mathbf{S}} \left(\nu_* \psi - \frac{4}{r} \psi \right) e^{\Phi} + \varsigma^{-1} \int_{\mathbf{S}} \zeta \left(\nu_* \psi - \frac{4}{r} \psi \right) e^{\Phi} + O(\epsilon_0 u^{-1-\delta_{dec}}) \int_{\mathbf{S}} |\psi|.$$

Proof. We have

$$\begin{aligned}\nu_* \left(\int_S \psi e^\Phi \right) &= \varsigma^{-1} \int_S \varsigma \left(\nu_*(\psi e^\Phi) + (\underline{\kappa} + a_*\kappa) \psi e^\Phi \right) \\ &= \varsigma^{-1} \int_S \varsigma \left(\nu_*\psi e^\Phi + e^{-\Phi} \nu_*(e^\Phi) + \underline{\kappa} + a_*\kappa \right) \psi e^\Phi.\end{aligned}$$

Recalling that $e_4(\Phi) = \frac{1}{2}(\kappa - \vartheta)$, $e_3(\Phi) = \frac{1}{2}(\underline{\kappa} - \underline{\vartheta})$ we deduce

$$e^{-\Phi} \nu_*(e^\Phi) + \underline{\kappa} + a_*\kappa = \frac{3}{2}(\underline{\kappa} + a_*\kappa) - \frac{1}{2}(\underline{\vartheta} - a_*\vartheta).$$

Hence, writing also $\varsigma = \bar{\varsigma} + \check{\varsigma}$ and then $\kappa = \bar{\kappa} + \check{\kappa}$, $\underline{\kappa} = \bar{\underline{\kappa}} + \check{\underline{\kappa}}$

$$\begin{aligned}\nu_* \left(\int_S \psi e^\Phi \right) &= \varsigma^{-1} \int_S \varsigma \left(\nu_*\psi + \frac{3}{2}(\underline{\kappa} + a_*\kappa)\psi \right) e^\Phi - \frac{1}{2}\varsigma^{-1} \int_S \varsigma(\underline{\vartheta} - a_*\vartheta)\psi e^\Phi \\ &= \varsigma^{-1}\bar{\varsigma} \int_S \left(\nu_*\psi + \frac{3}{2}(\underline{\kappa} + a_*\kappa)\psi \right) e^\Phi + \varsigma^{-1} \int_S \check{\varsigma} \left(\nu_*\psi + \frac{3}{2}(\underline{\kappa} + a_*\kappa)\psi \right) e^\Phi \\ &\quad - \frac{1}{2}\varsigma^{-1} \int_S \varsigma(\underline{\vartheta} - a_*\vartheta)\psi e^\Phi \\ &= \int_S \left(\nu_*\psi + \frac{3}{2}(\underline{\kappa} + a_*\kappa)\psi \right) e^\Phi + \varsigma^{-1} \int_S \check{\varsigma} \left(\nu_*\psi + \frac{3}{2}(\underline{\kappa} + a_*\kappa)\psi \right) e^\Phi \\ &\quad + (\varsigma^{-1}\bar{\varsigma} - 1) \int_S \left(\nu_*\psi + \frac{3}{2}(\underline{\kappa} + a_*\kappa)\psi \right) e^\Phi - \frac{1}{2}\varsigma^{-1} \int_S \varsigma(\underline{\vartheta} - a_*\vartheta)\psi e^\Phi \\ &= \int_S (\nu_*\psi) e^\Phi + \frac{3}{2}(\bar{\underline{\kappa}} + a_*\bar{\kappa}) \int_S \psi e^\Phi + \text{Err}[\psi, \nu_*]\end{aligned}$$

where,

$$\begin{aligned}\text{Err}[\psi, \nu_*] &= \varsigma^{-1} \int_S \check{\varsigma} \left(\nu_*\psi + \frac{3}{2}(\underline{\kappa} + a_*\kappa)\psi \right) e^\Phi - \varsigma^{-1}\check{\varsigma} \int_S \left(\nu_*\psi + \frac{3}{2}(\underline{\kappa} + a_*\kappa)\psi \right) e^\Phi \\ &\quad + \frac{3}{2} \int_S (\check{\underline{\kappa}} + a_*\check{\kappa}) \psi e^\Phi - \frac{1}{2}\varsigma^{-1} \int_S \varsigma(\underline{\vartheta} - a_*\vartheta)\psi e^\Phi.\end{aligned}\tag{7.3.8}$$

Using **Ref1-Ref3** and (7.3.2)

$$\underline{\kappa} + a_*\kappa = -\frac{2\Upsilon}{r} + (-1 - \frac{2m}{r})\frac{2}{r} + O(\epsilon r^{-1}u^{-1-\delta_{dec}}) = -\frac{4}{r} + O(\epsilon r^{-1}u^{-1-\delta_{dec}}),$$

we deduce,

$$\text{Err}[\psi, \nu_*] = -\varsigma^{-1}\check{\varsigma} \int_S \left(\nu_*\psi - \frac{4}{r}\psi \right) e^\Phi + \varsigma^{-1} \int_S \check{\varsigma} \left(\nu_*\psi - \frac{4}{r}\psi \right) e^\Phi + O(\epsilon u^{-1-\delta_{dec}}) \int_S |\psi|$$

as stated. \square

7.3.2 Differential identities involving GCM conditions on Σ_*

Recall our our GCM conditions on Σ_*

$$\kappa = \frac{2}{r}, \quad \not\partial_2^* \not\partial_1^* \mu = 0, \quad \not\partial_2^* \not\partial_1^* \underline{\kappa} = 0, \quad \int_S \eta e^\Phi = 0, \quad \int_S \underline{\xi} e^\Phi = 0. \quad (7.3.9)$$

Also, on S_* , the last cut of Σ_* ,

$$\int_{S_*} \beta e^\Phi = 0, \quad \int_{S_*} e_\theta(\underline{\kappa}) e^\Phi = 0. \quad (7.3.10)$$

The goal of the subsection is to derive identities involving the GCM conditions which will be used later, see Lemma 7.3.7.

Proposition 7.3.4. *The following identities hold true on Σ_* .*

$$\begin{aligned} 2 \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \not\partial_2^* \eta &= \kappa \left(\nu_* (\not\partial_2^* \not\partial_1^* \mu) + 2 \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} \right) - \not\partial_2^* \not\partial_1^* \not\partial_1 \nu_* (e_\theta(\kappa)) \\ &+ r^{-5} \not\partial^{\leq 4} \Gamma_g + r^{-4} \not\partial^{\leq 4} (\Gamma_b \cdot \Gamma_b) + l.o.t., \end{aligned} \quad (7.3.11)$$

$$\begin{aligned} 2 \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \not\partial_2^* \underline{\xi} &= \nu_* \left((\not\partial_2^* \not\partial_2 + 2K) \not\partial_2^* \not\partial_1^* \underline{\kappa} \right) - \underline{\kappa} \left(\nu_* (\not\partial_2^* \not\partial_1^* \mu) + 2 \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} \right) \\ &+ r^{-5} \not\partial^{\leq 4} \Gamma_g + r^{-4} \not\partial^{\leq 4} (\Gamma_b \cdot \Gamma_b) + l.o.t. \end{aligned} \quad (7.3.12)$$

The quadratic terms denoted *l.o.t.* are lower order both in terms of decay in r, u as well in terms of number of derivatives.

In particular, if the GCM conditions (7.3.9) are verified, we deduce,

$$\begin{aligned} \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \not\partial_2^* \eta &= \kappa \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} + r^{-5} \not\partial^{\leq 4} \Gamma_g + r^{-4} \not\partial^{\leq 4} (\Gamma_b \cdot \Gamma_b) + l.o.t., \\ \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \not\partial_2^* \underline{\xi} &= -\underline{\kappa} \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} + r^{-5} \not\partial^{\leq 4} \Gamma_g + r^{-4} \not\partial^{\leq 4} (\Gamma_b \cdot \Gamma_b) + l.o.t. \end{aligned} \quad (7.3.13)$$

Proof. The proof is a straightforward application of Proposition 7.1.14. Indeed according to (7.1.32) we have

$$\begin{aligned} 2 \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \not\partial_2^* \eta &= \kappa \left(e_3 (\not\partial_2^* \not\partial_1^* \mu) + 2 \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} \right) - \not\partial_2^* \not\partial_1^* \not\partial_1 e_3 (e_\theta(\kappa)) \\ &+ r^{-5} \not\partial^{\leq 4} \Gamma_g + r^{-4} \not\partial^{\leq 4} (\Gamma_b \cdot \Gamma_b) + l.o.t. \end{aligned}$$

On the other hand since $\nu_* = e_3 + a_* e_4$ with $a_* = \frac{2}{\zeta_*} - \Upsilon + \frac{r}{2} \underline{A}$, see (7.1.8),

$$\begin{aligned} e_3 (\not\partial_2^* \not\partial_1^* \mu) &= \nu_* (\not\partial_2^* \not\partial_1^* \mu) - a_* e_4 (\not\partial_2^* \not\partial_1^* \mu) \\ &= \nu_* (\not\partial_2^* \not\partial_1^* \mu) - a_* (\not\partial_2^* \not\partial_1^* e_4 \mu + [e_4, \not\partial_2^* \not\partial_1^*] \mu). \end{aligned}$$

Also, in the same fashion¹⁶,

$$\begin{aligned}
\mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 e_3(e_\theta(\kappa)) &= \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 [\nu_*(e_\theta(\kappa)) - a_* e_4 e_\theta \kappa] \\
&= \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 [\nu_*(e_\theta(\kappa))] - a_* \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 (e_4 e_\theta \kappa) + r^{-3} \sum_{i+j=2} \mathcal{D}^i a_* \mathcal{D}^j (e_4 e_\theta \kappa) \\
&= \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 [\nu_*(e_\theta(\kappa)) - a_* e_\theta e_4 \kappa - [e_4, e_\theta] \kappa] \\
&= r^{-2} \sum_{i+j=2} \mathcal{D}^i \Gamma_b \mathcal{D}^j (e_\theta(e_4 \kappa) + [e_\theta, e_4] \kappa).
\end{aligned}$$

Thus, after using the transport equations for $e_4 \mu$, $e_4 \kappa$ and the commutator lemma applied to $[e_4, e_\theta]$ we easily deduce,

$$\begin{aligned}
2 \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \mathcal{D}_2 \mathcal{D}_2^* \eta &= \kappa \left(\nu_*(\mathcal{D}_2^* \mathcal{D}_1^* \mu) + 2 \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \underline{\beta} \right) - \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \nu_*(e_\theta(\kappa)) \\
&\quad + r^{-5} \mathcal{D}^{\leq 4} \Gamma_g + r^{-4} \mathcal{D}^{\leq 4} (\Gamma_b \cdot \Gamma_b) + \text{l.o.t.}
\end{aligned}$$

which confirms the first identity of the proposition.

The second part of the proposition can be derived in the same manner starting with the identity (7.1.33)

$$\begin{aligned}
2 \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \mathcal{D}_2 \mathcal{D}_2^* \xi &= e_3 \left((\mathcal{D}_2^* \mathcal{D}_2 + 2K) \mathcal{D}_2^* \mathcal{D}_1^* \kappa \right) - \underline{\kappa} \left(e_3(\mathcal{D}_2^* \mathcal{D}_1^* \mu) + 2 \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \underline{\beta} \right) \\
&\quad + r^{-5} \mathcal{D}^{\leq 4} \Gamma_g + r^{-4} \mathcal{D}^{\leq 4} (\Gamma_b \cdot \Gamma_b) + \text{l.o.t.}
\end{aligned}$$

This concludes the proof of the proposition. \square

7.3.3 Control of the flux of some quantities on Σ_*

The goal of this subsection is to establish the following.

Proposition 7.3.5. *The following estimate holds true for all $k \leq k_{small} + 15$*

$$\int_{\Sigma_*(u, \underline{u})} r^2 |\mathcal{D}_*^k \underline{\beta}|^2 + |\mathcal{D}_*^k(\underline{\vartheta}, \eta, \underline{\xi})|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}. \quad (7.3.14)$$

We also have the weaker estimates for $k \leq k_{small} + 13$.

$$\begin{aligned}
\|\underline{\beta}\|_{\infty, k}^* &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \\
\|(\underline{\vartheta}, \eta, \underline{\xi})\|_{\infty, k}^* &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}.
\end{aligned} \quad (7.3.15)$$

¹⁶Note that in view of (7.1.8) we have $\mathcal{D} a_* \in r \Gamma_b$.

In the process we also prove the following estimates for the $\ell = 1$ modes of ν_* -derivatives of $\underline{\xi}, \eta$ (see (7.3.21))

$$\left| \int_S (\nu_*^k \eta) e^\Phi \right| + \left| \int_S (\nu_*^k \underline{\xi}) e^\Phi \right| \lesssim \epsilon_0 r u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 15.$$

Proof. We concentrate our attention on deriving (7.3.14). Following its proof below one can easily also check the weaker estimates (7.3.15) which require in fact much less work. We divide the proof in the following steps.

Step 1. We first prove the corresponding estimates for $\underline{\beta}$ away from its $\ell = 1$ mode. More precisely we prove.

Lemma 7.3.6. *The following estimates holds true for all $k \leq k_{small} + 15$*

$$\int_{\Sigma_*(u, \underline{u})} r^4 |\not{d}_2^*(\not{d}_*^k \underline{\beta})|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}. \quad (7.3.16)$$

We postpone the proof of the lemma to Step 7 in this subsection.

Step 2. We make use of the result of Lemma 7.3.6 first prove the desired estimate for $\underline{\vartheta}$ i.e.,

$$\int_{\Sigma_*(u, \underline{u})} |\not{d}_*^k \underline{\vartheta}|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}, \quad k \leq k_{small} + 15. \quad (7.3.17)$$

Proof. One starts with the Codazzi equation

$$\not{d}_2 \underline{\vartheta} = -2\underline{\beta} - \not{d}_1^*(\underline{\kappa}) - \zeta \underline{\kappa} + \Gamma_g \cdot \Gamma_b.$$

Differentiating w.r.t. \not{d}_2^* and then taking tangential derivatives \not{d}_*^k we derive,

$$\not{d}_*^k \not{d}_2^* \not{d}_2 \underline{\vartheta} = -2\not{d}_*^k \not{d}_2^* \underline{\beta} - \not{d}_*^k \not{d}_2^* \not{d}_1^*(\underline{\kappa}) - \not{d}_*^k [r^{-2} \not{d} \Gamma_g + r^{-1} \not{d} (\Gamma_g \cdot \Gamma_b)].$$

Making use of the GCM condition $\not{d}_2^* \not{d}_1^* \underline{\kappa} = 0$ along Σ_* and our assumptions **Ref1-Ref2** we deduce¹⁷, for all $k \leq k_{small} + 15$,

$$\begin{aligned} \not{d}_*^k \not{d}_2^* \not{d}_2 \underline{\vartheta} &= -2\not{d}_*^k \not{d}_2^* \underline{\beta} + r^{-2} \not{d}^{k+1} \Gamma_g + r^{-1} \not{d}^{k+1} \Gamma_g \cdot \Gamma_b \\ &= -2\not{d}_*^k \not{d}_2^* \underline{\beta} + \epsilon r^{-4} + \text{l.o.t.} \end{aligned}$$

¹⁷Note that we use (7.1.13) of **Ref1** to estimate the linear term $r^{-2} \not{d} \Gamma_g$.

or, since $r \geq \frac{\epsilon}{\epsilon_0} u^{1+\delta_{dec}}$,

$$\mathfrak{d}_*^k \phi_2^* \phi_2 \underline{\vartheta} = -2\mathfrak{d}_*^k \phi_2^* \underline{\beta} + \epsilon_0 r^{-3} u^{-1-\delta_{dec}}.$$

Moreover,

$$\phi_2^* \phi_2 \mathfrak{d}_*^k \underline{\vartheta} = -2\mathfrak{d}_*^k \phi_2^* \underline{\beta} + \epsilon_0 r^{-3} u^{-3/2-\delta_{dec}} + [\mathfrak{d}_*^k, \phi_2^* \phi_2] \underline{\vartheta}.$$

Using the commutator estimates of Lemma 7.3.1 we derive,

$$\phi_2^* \phi_2 \mathfrak{d}_*^k \underline{\vartheta} = -2\mathfrak{d}_*^k \phi_2^* \underline{\beta} + \epsilon_0 r^{-3} u^{-1-\delta_{dec}}.$$

Integrating and using the previously derived estimate for $\underline{\beta}$ we deduce,

$$\int_{\Sigma_*(u, u_*)} r^4 |\phi_2^* \phi_2 \mathfrak{d}_*^k \underline{\vartheta}|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}, \quad k \leq k_{small} + 15.$$

In view of the coercivity of $\phi_2^* \phi_2$ we infer that,

$$\int_{\Sigma_*(u, u_*)} |\mathfrak{d}_*^k \underline{\vartheta}|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}, \quad k \leq k_{small} + 15$$

as desired. □

Step 3. We next derive a weak¹⁸ estimate for the $\ell = 1$ mode of $\underline{\beta}$ with the help of the Codazzi equation for $\underline{\vartheta}$,

$$\begin{aligned} 2\underline{\beta} &= -\phi_2 \underline{\vartheta} + e_\theta(\underline{\kappa}) - \bar{\kappa} \zeta + \Gamma_g \cdot \Gamma_b \\ &= -\phi_2 \underline{\vartheta} + e_\theta(\check{\kappa}) + \frac{2\Upsilon}{r} \zeta + \Gamma_g \cdot \Gamma_b. \end{aligned}$$

Projecting on the $\ell = 1$ mode and using the bootstrap assumptions **Ref1-Ref2** we deduce,

$$\begin{aligned} 2 \int_S \underline{\beta} e^\Phi &= \frac{2\Upsilon}{r} \int_S \zeta e^\Phi + \int_S e_\theta(\check{\kappa}) e^\Phi + \int_S \Gamma_g \cdot \Gamma_b e^\Phi \\ &= O(\epsilon) + O(\epsilon_0 u^{-3/2-2\delta_{dec}}). \end{aligned}$$

The same estimate holds true for the tangential derivatives. More precisely taking tangential derivatives and projecting on the $\ell = 1$ mode,

$$2 \int_S (\mathfrak{d}_*^k \underline{\beta}) e^\Phi = \int_S \mathfrak{d}_*^k \left[e_\theta(\check{\kappa}) + \frac{2\Upsilon}{r} \zeta + \Gamma_g \cdot \Gamma_b \right] e^\Phi - \int_S [\mathfrak{d}_*^k, \phi_2] \underline{\vartheta} e^\Phi.$$

¹⁸i.e. of order $O(\epsilon)$.

Using the bootstrap assumptions **Ref1-Ref2** and the commutator Lemma 7.3.1 we deduce,

$$\int_S (\mathfrak{d}_*^k \underline{\beta}) e^\Phi = O(\epsilon). \quad (7.3.18)$$

Step 4. We combine the result of Lemma 7.3.6 with 7.3.18 to deduce

$$\int_{\Sigma_*(u, \underline{u})} r^2 |\mathfrak{d}_*^k \underline{\beta}|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}, \quad k \leq k_{small} + 15. \quad (7.3.19)$$

Indeed, according the last elliptic estimate of Lemma 7.1.7,

$$\int_S r^2 |\mathfrak{d}_*^k \underline{\beta}|^2 \lesssim r^4 \int_S |\mathfrak{d}_2^*(\mathfrak{d}_*^k \underline{\beta})|^2 + r^{-2} \left| \int_S (\mathfrak{d}_*^k \underline{\beta}) e^\Phi \right|^2 \lesssim r^4 \int_S |\mathfrak{d}_2^*(\mathfrak{d}_*^k \underline{\beta})|^2 + \epsilon^2 r^{-2}.$$

Since $r \geq (\epsilon \epsilon_0^{-1}) u^{-3/2-2\delta_{dec}}$ on Σ_* we deduce,

$$\int_S r^2 |\mathfrak{d}_*^k \underline{\beta}|^2 \lesssim r^4 \int_S |\mathfrak{d}_2^*(\mathfrak{d}_*^k \underline{\beta})|^2 + \epsilon_0^2 u^{-3-2\delta_{dec}}.$$

Thus, integrating and making use of estimate (7.3.16),

$$\begin{aligned} \int_{\Sigma(u, u_*)} r^2 |\mathfrak{d}_*^k \underline{\beta}|^2 &\lesssim \int_{\Sigma(u, u_*)} r^4 |\mathfrak{d}_2^*(\mathfrak{d}_*^k \underline{\beta})|^2 + \epsilon_0^2 u^{-2-2\delta_{dec}} \\ &\lesssim \epsilon_0^2 u^{-2-2\delta_{dec}} \end{aligned}$$

as stated. This establishes the estimate for $\underline{\beta}$ in Proposition 7.3.5.

Step 5. To finish the proof of Proposition 7.3.5 it remains to establish the estimates for η and $\underline{\xi}$. As for $\underline{\beta}$ we prove separately estimates for $\mathfrak{d}_2^*(\eta, \underline{\xi})$ and estimates for the $\ell = 1$ modes of the tangential derivatives with respect to ν_* of $\eta, \underline{\xi}$ (recall that the $\ell = 1$ modes of $\eta, \underline{\xi}$ were set to zero). We then combine them, as in the case of $\underline{\beta}$, to provide the desired estimates.

Lemma 7.3.7. *We have*

$$\int_{\Sigma_*(u, u_*)} r^2 \left(|\mathfrak{d}_2^*(\mathfrak{d}_*^k \eta)|^2 + |\mathfrak{d}_2^*(\mathfrak{d}_*^k \underline{\xi})|^2 \right) \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}, \quad k \leq k_{small} + 15.$$

We postpone the proof of the lemma to Step 8 of this section.

Step 6. We provide below estimates for the $\ell = 1$ modes of $\underline{\xi}$ and η and use them, in combination with Lemma 7.3.7, to close the estimates of Proposition 7.3.5, i.e. we show that,

$$\int_{\Sigma_*(u, u_*)} \left(|\mathfrak{d}_*^k \eta|^2 + |\mathfrak{d}_*^k \underline{\xi}|^2 \right) \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}, \quad k \leq k_{small} + 15. \quad (7.3.20)$$

In the process we also prove the following estimates for the $\ell = 1$ modes of ν_* - derivatives of $\underline{\xi}, \eta$,

$$\left| \int_S (\nu_*^k \eta) e^\Phi \right| + \left| \int_S (\nu_*^k \underline{\xi}) e^\Phi \right| \lesssim \epsilon_0 r u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 15. \quad (7.3.21)$$

Proof. We prove separately the estimates for $\underline{\xi}$ and η .

Step 6a. We prove first the estimate for $\underline{\xi}$. Since the $\ell = 1$ mode of $\underline{\xi}$ vanishes we have, in view of the elliptic estimates Lemma 7.1.7 and the estimates of Lemma 7.3.7,

$$\int_{\Sigma(u, u_*)} |\mathfrak{d} \underline{\xi}|^2 + |\underline{\xi}|^2 \lesssim \int_{\Sigma(u, u_*)} r^2 |\mathfrak{d}_2^* \underline{\xi}|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}. \quad (7.3.22)$$

According to our GCM conditions we have $\int_S \underline{\xi} e^\Phi = 0$. In view of Lemma 7.3.3 we deduce,

$$\int_S (\nu_* \underline{\xi}) e^\Phi = \text{Err}[\underline{\xi}, \nu_*]$$

where,

$$\text{Err}[\underline{\xi}, \nu_*] = -\varsigma^{-1} \zeta \int_S \left(\nu_* \underline{\xi} - \frac{4}{r} \underline{\xi} \right) e^\Phi + \varsigma^{-1} \int_S \zeta \left(\nu_* \underline{\xi} - \frac{4}{r} \underline{\xi} \right) e^\Phi + O(\epsilon_0 u^{-1-\delta_{dec}}) \int_S |\underline{\xi}|.$$

We deduce,

$$\int_S (\nu_* \underline{\xi}) e^\Phi = \varsigma^{-1} \left[-\zeta \int_S (\nu_* \underline{\xi}) e^\Phi + \int_S \zeta (\nu_* \underline{\xi}) e^\Phi \right] + O(\epsilon_0 u^{-1-\delta_{dec}}) \int_S |\underline{\xi}|.$$

Hence,

$$\left| \int_S (\nu_* \underline{\xi}) e^\Phi - \varsigma^{-1} \int_S \zeta (\nu_* \underline{\xi}) e^\Phi \right| \lesssim O(\epsilon) \left| \int_S (\nu_* \underline{\xi}) e^\Phi \right| + O(\epsilon_0) r u^{-1-\delta_{dec}}$$

i.e., setting $B_1 := \int_S (\nu_* \underline{\xi}) e^\Phi$ and $I_1 = \varsigma^{-1} \int_S \zeta (\nu_* \underline{\xi}) e^\Phi$,

$$|B_1 - I_1| \lesssim \epsilon B_1 + \epsilon_0 r u^{-1-\delta_{dec}}. \quad (7.3.23)$$

To control I_1 we decompose $\nu_*\underline{\xi}$ into its $\ell = 1$ mode $B_1 = \int_S (\nu_*\underline{\xi}) e^\Phi$ and its orthogonal complement

$$\nu_*\underline{\xi} = B_1 \frac{1}{\int_S e^{2\Phi}} e^\Phi + (\nu_*\underline{\xi})^\perp.$$

Thus, in view of our elliptic estimates,

$$\int_S |(\nu_*\underline{\xi})^\perp|^2 \lesssim r^2 \int_S |\not{d}_2^*(\nu_*\underline{\xi})^\perp|^2 = r^2 \int_S |\not{d}_2^*(\nu_*\underline{\xi})|^2.$$

We deduce,

$$\begin{aligned} I_1 &= \int_S \zeta(\nu_*\underline{\xi}) e^\Phi = \frac{B_1}{\int_S e^{2\Phi}} \int \zeta e^{2\Phi} + \int_S \zeta(\nu_*\underline{\xi})^\perp e^\Phi \\ &\lesssim \epsilon |B_1| + \epsilon r^2 \left(\int_S |(\nu_*\underline{\xi})^\perp|^2 \right)^{\frac{1}{2}} \\ &\lesssim \epsilon |B_1| + \epsilon r^3 \left(\int_S |\not{d}_2^*(\nu_*\underline{\xi})|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Hence,

$$|B_1| \lesssim \epsilon r^3 \left(\int_S |\not{d}_2^*(\nu_*\underline{\xi})|^2 \right)^{1/2} + \epsilon_0 r u^{-1-\delta_{dec}}. \quad (7.3.24)$$

Finally,

$$\begin{aligned} \int_S |\nu_*\underline{\xi}|^2 &\lesssim r^2 \int_S |\not{d}_2^*(\nu_*\underline{\xi})|^2 + r^{-4} B_1^2 \\ &\lesssim r^2 \int_S |\not{d}_2^*(\nu_*\underline{\xi})|^2 + \epsilon_0^2 r^{-2} u^{-2-2\delta_{dec}} \end{aligned}$$

or, since $r \geq u^{1/2}$,

$$\int_S |\nu_*\underline{\xi}|^2 \lesssim r^2 \int_S |\not{d}_2^*(\nu_*\underline{\xi})|^2 + \epsilon_0^2 u^{-3-2\delta_{dec}}.$$

Integrating we derive,

$$\int_{\Sigma(u, u_*)} |\nu_*\underline{\xi}|^2 \lesssim \int_{\Sigma(u, u_*)} r^2 |\not{d}_2^*(\nu_*\underline{\xi})|^2 + \epsilon_0^2 u^{-2-2\delta_{dec}}.$$

Thus, making use of (7.3.7),

$$\int_{\Sigma(u, u_*)} |\nu_*\underline{\xi}|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}$$

which, together with (7.3.22), yields

$$\int_{\Sigma(u, u_*)} |\mathfrak{d}_* \underline{\xi}|^2 + |\underline{\xi}|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}.$$

Moreover we as desired. Returning to (7.3.24) we also infer that,

$$|B_1| \lesssim \epsilon r^3 \left(\int_S |\not\phi_2^*(\nu_* \underline{\xi})|^2 \right)^{1/2} + \epsilon_0 r u^{-1-\delta_{dec}} \lesssim \epsilon_0 r u^{-1-\delta_{dec}}.$$

This proves the estimates (7.3.20), (7.3.21) for $\underline{\xi}$ and $k \leq 1$. The proof for the higher tangential derivatives can be derived in the same manner.

Step 6b. To prove the desired estimates for η we make use of the GCM condition $\int_S \eta e^\Phi = 0$. We deduce, by elliptic estimates,

$$\int_S |\eta|^2 + |\not\phi \eta|^2 \lesssim \int_S r^2 |\not\phi_2^* \eta|^2 + r^{-4} \left| \int_S \eta e^\Phi \right|^2 = \int_S r^2 |\not\phi_2^* \eta|^2.$$

Hence, integrating and using the estimates of Lemma 7.3.7,

$$\int_{\Sigma(u, u_*)} |\eta|^2 + |\not\phi \eta|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}} \quad (7.3.25)$$

which is the exact analogue of (7.3.22) in Step 6a.

To estimate $\int_S |\nu_* \eta|^2$ we proceed exactly as in step 6a by making use of the GCM condition $\int_S \underline{\xi} e^\Phi = 0$ and Lemma 7.3.3 to deduce

$$\begin{aligned} \int_S (\nu_* \eta) e^\Phi &= \text{Err}[\eta, \nu_*] \\ &= -\varsigma^{-1} \zeta \int_S \left(\nu_* \eta - \frac{4}{r} \eta \right) e^\Phi + \varsigma^{-1} \int_S \zeta \left(\nu_* \eta - \frac{4}{r} \eta \right) e^\Phi + O(\epsilon_0 u^{-1-\delta_{dec}}) \int_S |\eta|. \end{aligned}$$

The term $\text{Err}[\eta, \nu_*]$ can be dealt with precisely as $\text{Err}[\underline{\xi}, \nu_*]$ in Step 6a. Proceeding exactly as before in Step 6a we deduce,

$$\int_{\Sigma(u, u_*)} |\nu_* \eta|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}.$$

Thus, combined with (7.3.25), we infer that,

$$\int_{\Sigma(u, u_*)} |\eta|^2 + |\mathfrak{d}_* \eta|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}$$

and

$$\left| \int_S (\nu_* \eta) e^\Phi \right| \lesssim \epsilon_0 r u^{-1-\delta_{dec}}$$

which establishes the estimates (7.3.20), (7.3.21) for η and $k \leq 1$. The estimates for the higher tangential derivatives can be proved in the same manner. This completes the proof of both inequalities (7.3.20) and (7.3.21). \square

At this point we have reduced all estimates of Proposition 7.3.5 to the proofs of Lemmas 7.3.6 and 7.3.7.

Step 7. We prove Lemma 7.3.6 i.e. we show that,

$$\int_{\Sigma_*(u, \underline{u})} r^4 |\mathcal{D}_*^k \beta|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}, \quad k \leq k_{small} + 15.$$

We make use of Proposition 7.1.11 according to which

$$\begin{aligned} e_3(r\mathbf{q}) &= r^5 \left\{ \mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \underline{\beta} - \frac{3}{2} \kappa \rho \underline{\alpha} - \frac{3}{2} \rho \mathcal{D}_2^* \mathcal{D}_1^* \underline{\kappa} - \frac{3}{2} \underline{\kappa} \rho \mathcal{D}_2^* \zeta + \frac{3}{4} (2\rho^2 - \kappa \underline{\kappa} \rho) \underline{\vartheta} \right\} \\ &+ \text{Err}[e_3(r\mathbf{q})] \end{aligned}$$

where,

$$\text{Err}[e_3(r\mathbf{q})] = r^3 \mathfrak{D}^{\leq 3} (\Gamma_b \cdot \Gamma_g).$$

It is easy to check, with the help of the estimates **Ref1** and **Ref2** and the product Lemma 7.1.6, that for all $k \leq k_{small} + 16$,

$$\|\text{Err}[e_3(r\mathbf{q})]\|_{\infty, k}(u, r) \lesssim \epsilon_0 (r^{-1/2} u^{-1-\delta_{dec}} + u^{-3/2-\delta_{dec}}). \quad (7.3.26)$$

We can also check, making use of the estimates¹⁹ (7.1.13) of **Ref1** for large r ,

$$\|\rho \mathcal{D}_2^* \mathcal{D}_1^* \underline{\kappa}, \underline{\kappa} \rho \mathcal{D}_2^* \zeta, \kappa \underline{\kappa} \rho \underline{\vartheta}, \rho^2 \underline{\vartheta}\|_{\infty, k} \lesssim \epsilon (r^{-7} + r^{-6} u^{-1-\delta_{dec}}).$$

In view of our assumption for r on Σ_* we have $r \geq \frac{\epsilon}{\epsilon_0} u^{1+\delta_{dec}}$, we thus deduce for all $k \leq k_{small} + 17$

$$\begin{aligned} \left\| e_3(r\mathbf{q}) - r^5 \left(\mathcal{D}_2^* \mathcal{D}_1^* \mathcal{D}_1 \underline{\beta} - \frac{3}{2} \kappa \rho \underline{\alpha} \right) \right\|_{\infty, k} &\lesssim \epsilon r^{-2} + \epsilon_0 (r^{-1/2} u^{-1-\delta_{dec}} + u^{-3/2-\delta_{dec}}) \\ &\lesssim \epsilon_0 u^{-3/2-\delta_{dec}}. \end{aligned}$$

¹⁹which hold for all $k \leq k_{large} - 2$.

We infer that,

$$\|r^4 \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta}\|_{\infty, k} \lesssim \|r^{-1} e_3(r\mathbf{q})\|_{\infty, k} + \|\underline{\alpha}\|_{\infty, k} + \epsilon_0 r^{-1} u^{-3/2 - \delta_{dec}}.$$

Thus integrating on the last slice Σ_* and making use of the assumptions (7.1.15) and (7.1.18), i.e.

$$\int_{\Sigma_*(u, u_*)} |e_3 \not\partial^k \mathbf{q}|^2 + r^{-2} |\mathbf{q}|^2 + |\not\partial^k \underline{\alpha}|^2 \lesssim \epsilon_0^2 (1+u)^{-2-2\delta_{dec}}, \quad k \leq k_{small} + 14,$$

we deduce,

$$\int_{\Sigma_*(u, u_*)} r^8 |\not\partial^k (\not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta})|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}.$$

In particular,

$$\int_{\Sigma_*(u, u_*)} r^8 |\not\partial_*^k (\not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta})|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}$$

where $\not\partial_*^k = \nu_*^{k_1} \not\partial^{k_2}$ denote the tangential derivatives to Σ_* . Taking into account the commutator Lemma 7.3.1 we deduce, for $k \leq k_{small} + 14$,

$$\int_{\Sigma_*(u, u_*)} r^8 |\not\partial_2^* \not\partial_1^* \not\partial_1 (\not\partial_*^k \underline{\beta})|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}. \quad (7.3.27)$$

Since

$$\not\partial_1^* \not\partial_1 = \not\partial_2 \not\partial_2^* + 2K,$$

we infer that

$$\int_{\Sigma_*(u, u_*)} r^8 |(\not\partial_2^* \not\partial_2 + 2K) \not\partial_2^* (\not\partial_*^k \underline{\beta})|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}.$$

In view of the coercivity of $\not\partial_2^* \not\partial_2 + 2K$ we deduce,

$$\int_{\Sigma_*(u, u_*)} r^4 |\not\partial_2^* (\not\partial_*^k \underline{\beta})|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}, \quad k \leq k_{small} + 16.$$

This concludes the proof of Lemma 7.3.6. In view of the results derived in the first three steps this also establishes, unconditionally, the estimate,

$$\int_{\Sigma_*(u, \underline{u})} r^2 |\not\partial_*^k \underline{\beta}|^2 \lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}, \quad k \leq k_{small} + 15 \quad (7.3.28)$$

which will be used in the next step.

Step 8. We prove Lemma 7.3.7 based on the identities of Proposition 7.3.4. To derive the desired flux estimate for η we make use of the first part of Proposition 7.3.4 according to which we have,

$$\begin{aligned} 2 \phi_2^* \phi_1^* \phi_1 \phi_2 \phi_2^* \eta &= \kappa \left(\nu_* (\phi_2^* \phi_1^* \mu) + 2 \phi_2^* \phi_1^* \phi_1 \underline{\beta} \right) - \phi_2^* \phi_1^* \phi_1 \nu_* (e_\theta(\kappa)) \\ &\quad + r^{-5} \mathfrak{F}^{\leq 4} \Gamma_g + r^{-4} \mathfrak{F}^{\leq 4} (\Gamma_b \cdot \Gamma_b) + \text{l.o.t.} \end{aligned}$$

Since, $\phi_1^* \phi_1 = \phi_2 \phi_2^* + 2K$, we deduce,

$$\begin{aligned} \phi_2^* (\phi_2 \phi_2^* + 2K) \phi_2 \phi_2^* \eta &= \frac{1}{2} \left[\kappa \nu_* (\phi_2^* \phi_1^* \mu) - \phi_2^* \phi_1^* \phi_1 \nu_* (e_\theta(\kappa)) \right] + \kappa \phi_2^* (\phi_2 \phi_2^* + 2K) \underline{\beta} \\ &\quad + r^{-5} \mathfrak{F}^{\leq 4} \Gamma_g + r^{-4} \mathfrak{F}^{\leq 4} (\Gamma_b \cdot \Gamma_b) + \text{l.o.t.} \end{aligned}$$

or,

$$\begin{aligned} (\phi_2 \phi_2^* + 2K) \left[\phi_2^* \phi_2 \phi_2^* \eta - \kappa \phi_2^* \underline{\beta} \right] &= \frac{1}{2} \left[\kappa \nu_* (\phi_2^* \phi_1^* \mu) - \phi_2^* \phi_1^* \phi_1 \nu_* (e_\theta(\kappa)) \right] \\ &\quad + r^{-5} \mathfrak{F}^{\leq 4} \Gamma_g + r^{-4} \mathfrak{F}^{\leq 4} (\Gamma_b \cdot \Gamma_b) + \text{l.o.t.} \end{aligned}$$

Taking higher tangential derivatives and using our GCM assumptions on Σ_*

$$\mathfrak{D}_*^k (\phi_2 \phi_2^* + 2K) \left[\phi_2^* \phi_2 \phi_2^* \eta - \kappa \phi_2^* \underline{\beta} \right] = \mathfrak{D}_*^k \left[r^{-5} \mathfrak{F}^{\leq 4} \Gamma_g + r^{-4} \mathfrak{F}^{\leq 3} (\Gamma_b \cdot \mathfrak{D} \Gamma_b) \right] + \text{l.o.t.}$$

Making use of the commutation Lemma 7.3.1 we can rewrite,

$$r^2 (\phi_2 \phi_2^* + 2K) \left[\phi_2^* \phi_2 \phi_2^* (\mathfrak{D}_*^k \eta) - \kappa \phi_2^* (\mathfrak{D}_*^k \underline{\beta}) \right] = \sum_{j \leq k} \mathfrak{F}^{\leq 2} \left[r^{-3} \mathfrak{F}^{\leq 2} \mathfrak{D}_*^j \Gamma_g + r^{-2} \mathfrak{F}^{\leq 1} \mathfrak{D}_*^j (\Gamma_b \cdot \mathfrak{D} \Gamma_b) \right].$$

Using the ellipticity of the operator $(\phi_2 \phi_2^* + 2K)$, assumptions **Ref 2- Ref 4**, interpolation Lemma 7.1.6 and condition $r \geq (\epsilon \epsilon_0^{-1}) u^{3/2 + \delta_{dec}}$ on Σ_* we derive,

$$\begin{aligned} \phi_2^* \phi_2 \phi_2^* (\mathfrak{D}_*^k \eta) &= \kappa \phi_2^* (\mathfrak{D}_*^k \underline{\beta}) + O \left(\epsilon r^{-5} + \epsilon_0 r^{-4} u^{-2 - 2\delta_{dec}} \right) \\ &= \kappa \phi_2^* (\mathfrak{D}_*^k \underline{\beta}) + O \left(\epsilon_0 r^{-4} u^{-3/2 - \delta_{dec}} \right). \end{aligned}$$

Using also the ellipticity of $\phi_2^* \phi_2$ and the assumption $r \geq \epsilon (\epsilon_0^{-1}) u^{3/2 + \delta_{dec}}$ on Σ_* ,

$$\| \phi_2^* (\mathfrak{D}_*^k \eta) \|_{L^2(S)} \lesssim r \| (\mathfrak{D}_*^k \underline{\beta}) \|_{L^2(S)} + \epsilon_0 r^{-1} u^{-3/2 - \delta_{dec}}.$$

Finally, squaring, integrating on Σ_* and taking into account the flux estimate for $\underline{\beta}$ in (7.3.28) we deduce

$$\int_{\Sigma_*(u, u_*)} r^2 | \phi_2^* (\mathfrak{D}_*^k \eta) |^2 \lesssim \int_{\Sigma_*(u, u_*)} r^4 | (\mathfrak{D}_*^k \underline{\beta}) |^2 + \epsilon_0 u^{-2 - 2\delta_{dec}} \lesssim \epsilon_0 u^{-2 - 2\delta_{dec}}.$$

Hence,

$$\int_{\Sigma_*(u, u_*)} r^2 |\phi_2^*(\mathfrak{D}_*^k \eta)|^2 \lesssim \epsilon_0 u^{-2-2\delta_{dec}} \lesssim \epsilon_0 u^{-2-2\delta_{dec}}$$

as stated. This completes the proof of Lemma 7.3.7 for η . The proof for $\underline{\xi}$ is very similar and will thus be omitted. This therefore also completes the proof of Proposition 7.3.5. \square

7.3.4 Estimates for the $\ell = 1$ modes on Σ_*

In the previous subsection we have already derived estimates for the $\ell = 1$ modes of $\underline{\beta}, \eta, \underline{\xi}$. In what follows we derive estimates for the remaining $\ell = 1$ modes. We summarize the results in the following proposition.

Proposition 7.3.8. *The following estimates hold true for all $k \leq k_{small} + 16$,*

$$\begin{aligned} r^{-1} \left| \int_S (\nu_*^k \eta) e^\Phi \right| + r^{-1} \left| \int_S (\nu_*^k \underline{\xi}) e^\Phi \right| &\lesssim \epsilon_0 u^{-1-\delta_{dec}}, \\ \left| \int_S (\nu_*^k \underline{\beta}) e^\Phi \right| + r^{-1} \left| \int_S (\nu_*^k \zeta) e^\Phi \right| + \left| \int_S (\nu_*^k (e_\theta \underline{\kappa})) e^\Phi \right| &\lesssim \epsilon_0 u^{-1-\delta_{dec}}, \\ \left| \int_S \nu_*^k (e_\theta \rho) e^\Phi \right| + \left| \int_S \nu_*^k (e_\theta \mu) e^\Phi \right| + \left| \int_S (\nu_*^k \beta) e^\Phi \right| &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \\ \left| \int_S \nu_*^k (e_\theta \underline{\omega}) e^\Phi \right| &\lesssim \epsilon_0 r u^{-1-\delta_{dec}}. \end{aligned} \tag{7.3.29}$$

Proof. Note that the estimates in the first line of (7.3.29) have already been derived in the previous subsection. The proof of the remaining ones is done in a series of steps starting with the estimate for the $\ell = 1$ mode of $e_\theta(\underline{\kappa})$ and ending with that for the $\ell = 1$ mode of β . To derive the correct estimate for the latter we need to derive in fact stronger intermediary estimates than those stated in the proposition.

Step 1. We prove the desired estimate for $\int_S \zeta e^\Phi$. Indeed, in view of the Codazzi equations and the GCM condition on κ ,

$$\phi_2 \vartheta = -2\beta + (e_\theta(\kappa) + \zeta\kappa) + \Gamma_g \cdot \Gamma_g = -2\beta + \frac{2}{r}\zeta + \Gamma_g \cdot \Gamma_g.$$

Integrating,

$$\int_S \zeta e^\Phi = -r \int_S \beta e^\Phi + r^4 \Gamma_g \cdot \Gamma_g.$$

Thus, in view of **Ref1-3** for β and Γ_g ,

$$\left| \int_S \zeta e^\Phi \right| \lesssim \epsilon r^{1/2-\delta_B/2} + \epsilon_0 u^{-1-2\delta_{dec}}.$$

The higher derivative estimates can be proved in the same manner. Hence,

$$\left| \int_S (\nu_*^k \zeta) e^\Phi \right| \lesssim \epsilon r^{1/2-\delta_B/2} + \epsilon_0 u^{-1-2\delta_{dec}}, \quad k \leq k_{small} + 16. \quad (7.3.30)$$

In particular, for $r \geq \left(\frac{\epsilon}{\epsilon_0}\right)^2 u^{2+2\delta_{dec}}$,

$$\left| \int_S (\nu_*^k \zeta) e^\Phi \right| \lesssim r \epsilon_0 u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 16$$

as stated in the proposition.

Step 2. We prove here the following stronger estimate for the $\ell = 1$ mode of $e_\theta \underline{\kappa}$.

$$\left| \int_S (\nu_*^k e_\theta \underline{\kappa}) e^\Phi \right| \lesssim \epsilon_0 u^{-2-\delta_{dec}}, \quad k \leq k_{small} + 16. \quad (7.3.31)$$

To control the $\ell = 1$ mode of $e_\theta(\underline{\kappa})$ on Σ_* we need the precise identity²⁰ of Proposition 2.2.19

$$\begin{aligned} e_3(e_\theta(\underline{\kappa})) &= -2 \underline{\phi}_2 \underline{\phi}_2^* \underline{\xi} + \underline{\kappa} (e_3(\zeta) - \beta) + \underline{\kappa}^2 \zeta - \frac{3}{2} \underline{\kappa} e_\theta \underline{\kappa} + 6\rho \underline{\xi} - 2\underline{\omega} e_\theta(\underline{\kappa}) + \text{Err}[\underline{\phi}_2 \underline{\phi}_2^* \underline{\xi}], \\ \text{Err}[\underline{\phi}_2 \underline{\phi}_2^* \underline{\xi}] &= \left(2 \underline{\phi}_1 \underline{\xi} + \frac{1}{2} \underline{\kappa} \underline{\vartheta} + 2\underline{\eta} \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \underline{\eta} + 2e_\theta(\underline{\eta} \underline{\xi}) - \frac{1}{2} e_\theta(\underline{\vartheta}^2) \\ &+ \underline{\kappa} \left(\frac{1}{2} \underline{\vartheta} \zeta - \frac{1}{2} \underline{\vartheta} \underline{\xi} \right) - \frac{1}{2} \underline{\vartheta} e_\theta \underline{\kappa} - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} \underline{\xi} - \zeta \left(2 \underline{\phi}_1 \underline{\xi} + 2(\underline{\eta} - 3\underline{\zeta}) \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \\ &+ \underline{\xi} \left(-\underline{\vartheta} \underline{\vartheta} - 2 \underline{\phi}_1 \zeta + 2\underline{\zeta}^2 \right) - 6\underline{\eta} \zeta \underline{\xi} - 6e_\theta(\zeta \underline{\xi}). \end{aligned}$$

The error term can be written schematically as,

$$\begin{aligned} \text{Err}[\underline{\phi}_2 \underline{\phi}_2^* \underline{\xi}] &= \left(2 \underline{\phi}_1 \underline{\xi} + \frac{1}{2} \underline{\kappa} \underline{\vartheta} + 2\underline{\eta} \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \underline{\eta} + 2e_\theta(\underline{\eta} \underline{\xi}) - \frac{1}{2} e_\theta(\underline{\vartheta}^2) \\ &+ r^{-1} \mathfrak{P}^{\leq 1}(\Gamma_g \cdot \Gamma_b) + \Gamma_g \cdot \Gamma_b \cdot \Gamma_b. \end{aligned}$$

Note also that we can write, schematically,

$$\underline{\kappa}^2 \zeta - \frac{3}{2} \underline{\kappa} e_\theta \underline{\kappa} + 6\rho \underline{\xi} - 2\underline{\omega} e_\theta(\underline{\kappa}) = r^{-2} \zeta + r^{-1} e_\theta(\underline{\kappa}) + r^{-3} \underline{\xi} + r^{-1} \mathfrak{P}^{\leq 1}(\Gamma_g \cdot \Gamma_b) + \Gamma_g \cdot \Gamma_b \cdot \Gamma_b.$$

²⁰Note that the schematic form of Corollary 7.1.12 is not suitable here.

Also, using the transport equation for $e_4(\underline{\kappa})$,

$$\begin{aligned}
e_4(e_\theta(\underline{\kappa})) &= e_\theta e_4 \underline{\kappa} + [e_4, e_\theta] \underline{\kappa} \\
&= e_\theta \left[-\frac{1}{2} \kappa \underline{\kappa} - 2 \phi_1 \zeta + 2\rho + \Gamma_g \cdot \Gamma_b \right] + \frac{1}{2} \kappa e_\theta \underline{\kappa} + \text{l.o.t.} \\
&= r^{-1} e_\theta \underline{\kappa} + \phi_1^* \phi_1 \zeta + e_\theta(\rho) + r^{-1} \mathfrak{P}(\Gamma_g \cdot \Gamma_b) \\
&= r^{-1} e_\theta \underline{\kappa} + (\phi_2 \phi_2^* + 2K) \zeta + e_\theta(\rho) + r^{-1} \mathfrak{P}(\Gamma_g \cdot \Gamma_b).
\end{aligned}$$

We can also write, since $\nu_* = e_3 + a_* e_4$

$$e_3(\zeta) = \nu_*(\zeta) - a_* e_4(\zeta) = \nu_*(\zeta) - a_* (-\kappa \zeta - \beta + \Gamma_g \cdot \Gamma_b).$$

Therefore, writing also $\underline{\kappa} = -\frac{2\Upsilon}{r} + \Gamma_g$,

$$\begin{aligned}
\nu_*(e_\theta(\underline{\kappa})) &= -2 \phi_2 \phi_2^* \underline{\zeta} - \frac{2\Upsilon}{r} (\nu_*(\zeta) - \beta) + E_1 + E_2 + E_3, \\
E_1 &= r^{-2} \zeta + r^{-1} e_\theta(\underline{\kappa}) + r^{-3} \underline{\zeta} + \phi_2 \phi_2^* \zeta + r^{-1} \beta + e_\theta(\rho), \\
E_2 &= \left(2 \phi_1 \underline{\zeta} + \frac{1}{2} \underline{\kappa} \underline{\vartheta} + 2\eta \underline{\zeta} - \frac{1}{2} \underline{\vartheta}^2 \right) \eta + 2e_\theta(\eta \underline{\zeta}), -\frac{1}{2} e_\theta(\underline{\vartheta}^2) \\
E_3 &= r^{-1} \mathfrak{P}^{\leq 1}(\Gamma_g \cdot \Gamma_b) + \Gamma_g \cdot \Gamma_b \cdot \Gamma_b.
\end{aligned} \tag{7.3.32}$$

Projecting on the $\ell = 1$ mode we derive,

$$\int_S \nu_* (e_\theta(\underline{\kappa}) - \underline{\kappa} \zeta) e^\Phi = -\frac{2\Upsilon}{r} \int_S \beta e^\Phi + \int_S (E_1 + E_2 + E_3) e^\Phi. \tag{7.3.33}$$

On the other hand, according to Lemma 7.3.3

$$\nu_* \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) = \int_S (\nu_* e_\theta(\underline{\kappa})) e^\Phi + \frac{3}{2} (\bar{\kappa} + a_* \bar{\kappa}) \int_S e_\theta(\underline{\kappa}) e^\Phi + \text{Err}[e_\theta(\underline{\kappa}), \nu_*]$$

with error term

$$\begin{aligned}
\text{Err}[e_\theta(\underline{\kappa}), \nu_*] &= -\zeta^{-1} \int_S \left(\nu_* e_\theta(\underline{\kappa}) - \frac{4}{r} e_\theta(\underline{\kappa}) \right) e^\Phi + \zeta^{-1} \int_S \zeta \left(\nu_* e_\theta(\underline{\kappa}) - \frac{4}{r} e_\theta(\underline{\kappa}) \right) e^\Phi \\
&+ O(\epsilon_0 u^{-1-\delta_{dec}}) \int_S |e_\theta(\underline{\kappa})|.
\end{aligned}$$

We easily deduce

$$\text{Err}[e_\theta(\underline{\kappa}), \nu_*] = r \mathfrak{P}^{\leq 1} \Gamma_b \cdot \Gamma_b + \text{l.o.t.} \tag{7.3.34}$$

i.e.,

$$\begin{aligned} \nu_* \left(\int_S e_{\theta}(\underline{\kappa}) e^{\Phi} \right) &= \int_S (\nu_* e_{\theta}(\underline{\kappa})) e^{\Phi} + O(r^{-1}) \int_S e_{\theta}(\underline{\kappa}) e^{\Phi} + r \mathfrak{O}^{\leq 1} \Gamma_b \cdot \Gamma_b + \text{l.o.t.} \\ &= \int_S (\nu_* e_{\theta}(\underline{\kappa})) e^{\Phi} + r \mathfrak{O} \Gamma_g + r \mathfrak{O}^{\leq 1} \Gamma_b \cdot \Gamma_b + \text{l.o.t.} \end{aligned}$$

Similarly,

$$\nu_* \left(\int_S (\underline{\kappa} \zeta) e^{\Phi} \right) = \int_S (\nu_* \underline{\kappa} \zeta) e^{\Phi} + r \mathfrak{O} \Gamma_g + r \mathfrak{O}^{\leq 1} \Gamma_b \cdot \Gamma_b + \text{l.o.t.}$$

Combining with (7.3.33) we deduce,

$$\begin{aligned} \nu_* \int_S (e_{\theta}(\underline{\kappa}) - \underline{\kappa} \zeta) e^{\Phi} &= -\frac{2\Upsilon}{r} \int_S \underline{\beta} e^{\Phi} + \int_S (E_1 + E_2 + E_3) e^{\Phi} \\ &\quad + r \mathfrak{O} \Gamma_g + r \mathfrak{O}^{\leq 1} \Gamma_b \cdot \Gamma_b + \text{l.o.t.} \end{aligned}$$

Now schematically, in view of **Ref 1-2**,

$$\begin{aligned} \int_S E_1 e^{\Phi} &= \int_S (r^{-2} \zeta + r^{-1} e_{\theta}(\underline{\kappa}) + r^{-1} \beta + e_{\theta}(\rho)) e^{\Phi} = r \mathfrak{O}^{\leq 1} \Gamma_g \\ \int_S E_3 e^{\Phi} &= r^2 \mathfrak{O}^{\leq 1} (\Gamma_g \cdot \Gamma_b) + r^3 \Gamma_g \cdot \Gamma_b \cdot \Gamma_b. \end{aligned}$$

Hence,

$$\begin{aligned} \nu_* \int_S (e_{\theta}(\underline{\kappa}) - \underline{\kappa} \zeta) e^{\Phi} &= -\frac{2\Upsilon}{r} \int_S \underline{\beta} e^{\Phi} + r \mathfrak{O}^{\leq 1} \Gamma_g + \int_S E_2 e^{\Phi} \\ &\quad + r \mathfrak{O}^{\leq 1} \Gamma_b \cdot \Gamma_b + r^3 \Gamma_g \cdot \Gamma_b \cdot \Gamma_b + \text{l.o.t.} \end{aligned} \tag{7.3.35}$$

In particular, in view of the estimate for the $\ell = 1$ mode of $\underline{\beta}$ derived in the previous subsection (see (7.3.18)) and our assumptions on Γ_g, Γ_b ,

$$\nu_* \left(\int_S (e_{\theta}(\underline{\kappa}) - \underline{\kappa} \zeta) e^{\Phi} \right) = \int_S E_2 e^{\Phi} + O(\epsilon r^{-1})$$

or, in view of the condition $r \geq (\epsilon \epsilon_0^{-1}) u^{3+\delta_{dec}}$,

$$\nu_* \left(\int_S (e_{\theta}(\underline{\kappa}) - \underline{\kappa} \zeta) e^{\Phi} \right) = \int_S E_2 e^{\Phi} + O(\epsilon_0 u^{-3-\delta_{dec}}).$$

Integrating along Σ_* from S_* and making use of the vanishing of the $\ell = 1$ mode of $e_\theta(\underline{\kappa})$ on S_* and the estimate for the $\ell = 1$ mode of ζ derived earlier in step 0, see (7.3.30), we deduce,

$$\begin{aligned} \int_{S(u)} \left(e_\theta(\underline{\kappa}) - \underline{\kappa}\zeta \right) e^\Phi &= \int_{S_*} \left(e_\theta(\underline{\kappa}) - \underline{\kappa}\zeta \right) e^\Phi + O\left(\epsilon_0 u^{-2-\delta_{dec}}\right) + \int_{\Sigma(u, u_*)} E_2 e^\Phi \\ &= \frac{2\Upsilon}{r} \int_{S_*} \zeta e^\Phi + O\left(\epsilon_0 u^{-2-\delta_{dec}}\right) + \int_{\Sigma(u, u_*)} E_2 e^\Phi \\ &= O\left(\epsilon r^{-1/2-\frac{1}{2}\delta_B}\right) + O\left(\epsilon_0 u^{-2-\delta_{dec}}\right) + \int_{\Sigma(u, u_*)} E_2 e^\Phi. \end{aligned}$$

Since $\min_{\Sigma_*} r \geq (\epsilon\epsilon_0^{-1})^2 u^4$ and $\delta_B \geq 2\delta_{dec}$ we infer that

$$\left| \int_{S(u)} \left(e_\theta(\underline{\kappa}) - \underline{\kappa}\zeta \right) e^\Phi \right| \lesssim \epsilon_0 u^{-2-\delta_{dec}} + \int_{\Sigma(u, u_*)} r |E_2|.$$

It remains to estimate the term $\int_{\Sigma(u, u_*)} r |E_2|$ where, recall,

$$E_2 = \left(2\underline{\mathcal{A}}_1 \underline{\xi} + \frac{1}{2} \underline{\kappa} \underline{\vartheta} + 2\underline{\eta} \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \eta + 2e_\theta(\underline{\eta} \underline{\xi}) - \frac{1}{2} e_\theta(\underline{\vartheta}^2).$$

Hence, appealing to Proposition 7.3.5

$$\begin{aligned} \int_{\Sigma(u, u_*)} r |E_2| &\lesssim \int_{\Sigma(u, u_*)} \left(|\underline{\vartheta}^{\leq 1}|^2 + |\underline{\eta}^{\leq 1}|^2 + |\underline{\xi}^{\leq 1}|^2 \right) \\ &\lesssim \epsilon_0^2 u^{-2-2\delta_{dec}}. \end{aligned}$$

Thus,

$$\begin{aligned} \left| \int_{S(u)} |e_\theta(\underline{\kappa}) e^\Phi| &\lesssim \left| \int_{S(u)} \underline{\kappa}\zeta e^\Phi \right| + \epsilon_0^2 u^{-2-2\delta_{dec}} \lesssim r^{-1/2} \epsilon + \epsilon_0^2 u^{-2-2\delta_{dec}} \\ &\lesssim \epsilon_0^2 u^{-2-2\delta_{dec}} \end{aligned}$$

which establishes the desired estimate in this case.

To derive a similar estimate for the first tangential derivative we go back to (7.3.33)

$$\int_S \nu_* \left(e_\theta(\underline{\kappa}) - \underline{\kappa}\zeta \right) e^\Phi = -\frac{2\Upsilon}{r} \int_S \underline{\beta} e^\Phi + \int_S (E_1 + E_2 + E_3) e^\Phi$$

from which we easily derive, as before,

$$\left| \int_S \nu_* \left(e_\theta(\underline{\kappa}) - \underline{\kappa}\zeta \right) e^\Phi \right| \lesssim \epsilon r^{-1} + \epsilon_0 u^{-2-2\delta_{dec}} \lesssim \epsilon_0 u^{-2-2\delta_{dec}}$$

and therefore,

$$\left| \int_S \nu_* e_{\theta}(\underline{\kappa}) \right| \lesssim \epsilon r^{-1/2} + \epsilon_0 u^{-2-2\delta_{dec}} \lesssim \epsilon_0 u^{-2-2\delta_{dec}}.$$

To estimate the second tangential derivative we make use once more of Lemma 7.3.3. Setting $\psi := \nu_* (e_{\theta}(\underline{\kappa}) - \underline{\kappa}\zeta)$ we deduce,

$$\begin{aligned} \int_S (\nu_* \psi) e^{\Phi} &= \nu_* \left(\int_S \psi e^{\Phi} \right) - \text{Err}[\psi, \nu_*] \\ &= \nu_* \left(-\frac{2\Upsilon}{r} \int_S \underline{\beta} e^{\Phi} + \int_S E e^{\Phi} \right) - \text{Err}[\psi, \nu_*] \end{aligned}$$

with $E = E_1 + E_2 + E_3$ and error term

$$\text{Err}[\psi, \nu_*] = -\varsigma^{-1} \zeta \int_S (e_3 \psi + \frac{3}{2} \underline{\kappa} \psi) e^{\Phi} + \varsigma^{-1} \int_S \check{\zeta} (e_3 \psi + \frac{3}{2} \underline{\kappa} \psi) e^{\Phi} + O(\epsilon_0 u^{-1-\delta_{dec}}) \int_S |\psi|$$

which can be estimated as in 7.3.34 and shown to be lower order. Note that, using again Lemma 7.3.3,

$$\begin{aligned} \nu_* \left(-\frac{2\Upsilon}{r} \int_S \underline{\beta} e^{\Phi} + \int_S E e^{\Phi} \right) &= -\frac{2\Upsilon}{r} \int_S (\nu_* \underline{\beta}) e^{\Phi} + \int_S (\nu_* E) e^{\Phi} + \text{l.o.t.} \\ &= O(\epsilon r^{-1}) + O(\epsilon_0 u^{-2-2\delta_{dec}}) = O(\epsilon_0 u^{-2-2\delta_{dec}}) \end{aligned}$$

and we then proceed as before. All higher tangential derivatives can be treated in the same manner.

Step 3. We establish the estimate for the $\ell = 1$ mode of β . Recall that we had, see (7.3.18),

$$2 \int_S \underline{\beta} e^{\Phi} = \frac{2\Upsilon}{r} \int_S \zeta e^{\Phi} + \int_S e_{\theta}(\check{\kappa}) e^{\Phi} + \int_S \Gamma_g \cdot \Gamma_b e^{\Phi}.$$

Thus, using the estimates already derived in Steps 0 and 1,

$$\left| \int_S \underline{\beta} e^{\Phi} \right| \lesssim \epsilon_0 u^{-1-\delta_{dec}}.$$

The higher derivative estimates are derived in the same manner.

Step 4. We establish the estimate

$$\left| \int_S \nu_*^k e_{\theta}(\rho) e^{\Phi} \right| \lesssim \epsilon_0 r^{-1} u^{-2-2\delta_{dec}} + \int_S |\nu_*^k \vartheta(\vartheta \underline{\vartheta})|. \tag{7.3.36}$$

We start by differentiating the Gauss equation $K = -\rho - \frac{1}{4}\kappa \underline{\kappa} + \frac{1}{4}\vartheta \underline{\vartheta}$. we derive Using the GCM condition for κ we derive,

$$e_\theta(\rho) = -e_\theta(K) - \frac{1}{2r} e_\theta(\underline{\kappa}) + \frac{1}{4} e_\theta(\vartheta \underline{\vartheta}).$$

We make use of the vanishing of the $\ell = 1$ mode of $e_\theta(K)$ (see Lemma 2.1.29) to derive

$$\int_S e_\theta(\rho) e^\Phi = -\frac{1}{2r} \int_S e_\theta(\underline{\kappa}) e^\Phi + \frac{1}{4} \int_S e_\theta(\vartheta \underline{\vartheta}) e^\Phi. \quad (7.3.37)$$

Using the previously deduced estimate for the $\ell = 1$ mode of $e_\theta(\underline{\kappa})$ we deduce

$$\left| \int_S e_\theta(\rho) e^\Phi \right| \lesssim \epsilon_0 r^{-1} u^{-2-2\delta_{dec}} + \int_S |\underline{\vartheta}| |\vartheta| + |\vartheta| |\underline{\vartheta}|. \quad (7.3.38)$$

Note that in particular we deduce,

$$\left| \int_S e_\theta(\rho) e^\Phi \right| \lesssim \epsilon_0 r^{-1} u^{-3/2-2\delta_{dec}}$$

but we shall need the precise form (7.3.38) in Step 7.

To estimate $\int_S (\nu_* e_\theta(\rho)) e^\Phi$ we apply ν_* to (7.3.38) and make use of Lemma 7.3.3 as in Step 2.

$$\begin{aligned} \int_S (\nu_* e_\theta(\rho)) e^\Phi &= \nu_* \left(\int_S e_\theta(\rho) e^\Phi \right) - \frac{3}{2} (\bar{\kappa} + a_* \bar{\kappa}) \int_S e_\theta(\rho) e^\Phi - \text{Err}[e_\theta(\rho), \nu_*] \\ &= \nu_* \left[-\frac{1}{2r} \int_S e_\theta(\underline{\kappa}) e^\Phi + \frac{1}{4} \int_S e_\theta(\vartheta \underline{\vartheta}) e^\Phi \right] + O(r^{-1}) \int_S e_\theta(\rho) e^\Phi - \text{Err}[e_\theta(\rho), \nu_*] \\ &= -\frac{1}{2r} \nu_* \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) + \frac{1}{4} \int_S \nu_* [e_\theta(\vartheta \underline{\vartheta})] e^\Phi + \text{l.o.t.} \end{aligned}$$

Hence, in view of the results derived in the first step, we deduce (7.3.36) for $k = 1$. The case of higher derivatives can be treated in the same manner.

Step 5. To estimate the $\ell = 1$ mode of μ we differentiate the relation $\mu = -\text{div } \zeta - \rho + \Gamma_g \cdot \Gamma_b$,

$$\begin{aligned} e_\theta(\mu) &= \not\phi_1^* \not\phi_1 \zeta - e_\theta(\rho) + r^{-1} \not\vartheta(\Gamma_g \cdot \Gamma_b) \\ &= (\not\phi_2 \not\phi_2^* + 2K) \zeta - e_\theta(\rho) + r^{-1} \not\vartheta(\Gamma_g \cdot \Gamma_b) \\ &= \not\phi_2 \not\phi_2^* \zeta + \frac{2}{r^2} \zeta - e_\theta(\rho) + r^{-1} \not\vartheta(\Gamma_g \cdot \Gamma_b) + \text{l.o.t.} \end{aligned}$$

Hence,

$$\int_S e_\theta(\mu)e^\Phi = 2r^{-2} \int_S \zeta e^\Phi - \int_S e_\theta(\rho)e^\Phi + r^2 \check{\vartheta}(\Gamma_g \cdot \Gamma_b). \quad (7.3.39)$$

Using the previous estimates for the $\ell = 1$ modes of ζ and $e_\theta\rho$ we deduce,

$$\left| \int_S e_\theta(\mu)e^\Phi \right| \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}.$$

The higher ν_* derivatives can be derived exactly as in the previous steps by differentiating (7.3.39) and applying Lemma 7.3.3.

Step 6. We estimate the $\ell = 1$ mode of $\underline{\omega}$ making use of the first identity in Corollary 7.1.12

$$\begin{aligned} 2\check{\mathcal{I}}_1^* \underline{\omega} &= \left(\frac{1}{2} \underline{\kappa} + 2\underline{\omega} \right) \eta + e_3(\zeta) - \underline{\beta} + \frac{1}{2} \underline{\kappa} \underline{\xi} + r^{-1} \Gamma_g + \Gamma_b^2 \\ &= \left(-\frac{\Upsilon}{r} + 2\frac{m}{r^2} \right) \eta + e_3(\zeta) - \underline{\beta} + \frac{1}{r} \underline{\xi} + r^{-1} \Gamma_g + \Gamma_b^2 + \text{l.o.t.} \\ &= \left(-\frac{\Upsilon}{r} + 2\frac{m}{r^2} \right) \eta + \nu_*(\zeta) - a_* e_4 \zeta - \underline{\beta} + \frac{1}{r} \underline{\xi} + r^{-1} \Gamma_g + \Gamma_b^2 + \text{l.o.t.} \\ &= -\frac{1}{2} \eta + \nu_*(\zeta) - \underline{\beta} + \frac{1}{r} \underline{\xi} + r^{-1} \Gamma_g + \Gamma_b^2 + \text{l.o.t.} \end{aligned}$$

Hence,

$$2 \int_S \check{\mathcal{I}}_1^*(\underline{\omega})e^\Phi = -\frac{1}{2} \int_S \eta e^\Phi + \int_S \nu_* \zeta e^\Phi - \int_S \underline{\beta} e^\Phi + r^{-1} \int_S \underline{\xi} e^\Phi + r^2 \Gamma_g + r^2 \Gamma_b \cdot \Gamma_b.$$

Taking into account the prescribed GCM conditions for the $\ell = 1$ modes of η and $\underline{\xi}$ and the previous results for the $\ell = 1$ modes of $\nu_*(\zeta), \underline{\beta}$ we deduce,

$$\left| \int_S e_\theta(\underline{\omega})e^\Phi \right| \lesssim \epsilon_0 r u^{-3/2-\delta_{dec}}.$$

The higher ν_* derivatives can be treated as in the previous steps.

Step 7. It remains to estimate the $\ell = 1$ mode of β and show

$$\left| \int_S \nu_*^k(\beta)e^\Phi \right| \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}. \quad (7.3.40)$$

We start with the $e_3\beta$ equation which we write in the form

$$e_3(\beta) = e_\theta(\rho) + 3\bar{\rho}\eta + J + O(r^{-1}(\alpha, \beta)), \quad J := 3\eta\check{\rho} - \vartheta\underline{\beta}.$$

Also, taking into account the e_4 equation for β ,

$$\begin{aligned}\nu_*(\beta) &= e_3(\beta) + a_*e_4\beta = e_3(\beta) + a^*(-2\kappa\beta + \mathfrak{d}_1\alpha + \zeta\alpha) \\ &= e_\theta(\rho) + 3\bar{\rho}\eta + J + O(r^{-1}(\beta, \mathfrak{P}^{\leq 1}\alpha)) + \text{l.o.t.}\end{aligned}$$

Projecting on the $\ell = 1$ mode,

$$\int_S \nu_*(\beta)e^\Phi = \int_S e_\theta(\rho)e^\Phi + \int_S Je^\Phi + O(r^2(\beta, \mathfrak{P}^{\leq 1}\alpha)). \quad (7.3.41)$$

On the other hand, making use of Lemma 7.3.3,

$$\begin{aligned}\nu_*\left(\int_S \beta e^\Phi\right) &= \int_S (\nu_*\beta)e^\Phi + \frac{3}{2}(\bar{\kappa} + a_*\bar{\kappa}) \int_S \beta e^\Phi + \text{Err}[\beta, \nu_*] \\ &= \int_S (\nu_*\beta)e^\Phi + O(r^2\beta) + \text{Err}[\beta, \nu_*]\end{aligned}$$

with error term

$$\text{Err}[\beta, \nu_*] = -\varsigma^{-1}\zeta \int_S (e_3\beta + \frac{3}{2}\bar{\kappa}\beta)e^\Phi + \varsigma^{-1} \int_S \zeta(e_3\beta + \frac{3}{2}\bar{\kappa}\beta)e^\Phi + \epsilon_0 u^{-1-\delta_{dec}} \int_S |\beta|.$$

Note that $e_3\beta = e_\theta\rho + \text{l.o.t.} = r^{-2}\mathfrak{P}^{\leq 1}\Gamma_g$. Hence,

$$\text{Err}[\beta, \nu_*] = r^3(r^{-2}\Gamma_b \cdot \mathfrak{P}^{\leq 1}\Gamma_g) + \text{l.o.t.}$$

Hence,

$$\nu_*\left(\int_S \beta e^\Phi\right) = \int_S (\nu_*\beta)e^\Phi + O(r^2\beta) + r\Gamma_b \cdot \mathfrak{P}^{\leq 1}\Gamma_g + \text{l.o.t.} \quad (7.3.42)$$

Thus combining to (7.3.41),

$$\nu_*\left(\int_S \beta e^\Phi\right) = \int_S e_\theta(\rho)e^\Phi + 3\bar{\rho} \int_S \eta e^\Phi + \int_S Je^\Phi + O(r^2(\beta, \mathfrak{P}^{\leq 1}\alpha)) + r\Gamma_b \cdot \mathfrak{P}^{\leq 1}\Gamma_g$$

and, since $\nu_*(r) = O(1)$

$$\nu_*\left(r \int_S \beta e^\Phi\right) = r \int_S e_\theta(\rho)e^\Phi + 3r\bar{\rho} \int_S \eta e^\Phi + r \int_S Je^\Phi + O(r^3(\beta, \mathfrak{P}^{\leq 1}\alpha)) + r^2\Gamma_b \cdot \mathfrak{P}^{\leq 1}\Gamma_g.$$

Note that, based on **Ref1-Ref2** and condition $\min_{\Sigma_*} r \geq (\epsilon\epsilon_0^{-1})^2 r^4$ and $\delta_B \geq 2\delta_{dec}$,

$$\begin{aligned}r^3|(\beta, \mathfrak{P}^{\leq 1}\alpha)| &\lesssim \epsilon r^{-1/2-1/2\delta_B} \lesssim \epsilon_0 u^{-2-\delta_{dec}}, \\ r^2|\Gamma_b \cdot \mathfrak{P}^{\leq 1}\Gamma_g| &\lesssim \epsilon r^{-1} \lesssim \epsilon_0 u^{-2-\delta_{dec}}.\end{aligned}$$

Recall also that the $\ell = 1$ mode of $e_\theta(\rho)$ has been estimated in (7.3.38)

$$\left| r \int_S e_\theta(\rho) e^\Phi \right| \lesssim \epsilon_0 u^{-2-2\delta_{dec}} + r \int_S |\underline{\vartheta}| |\check{\vartheta}| + |\vartheta| |\check{\vartheta}|.$$

Therefore, using also that the $\ell = 1$ mode of η vanishes on Σ_* due to our GCM conditions,

$$\left| \nu_* \left(r \int_S \beta e^\Phi \right) \right| \lesssim \epsilon_0 u^{-2-2\delta_{dec}} + r \int_S |\underline{\vartheta}| |\check{\vartheta}| + |\vartheta| |\check{\vartheta}| + r|J|. \quad (7.3.43)$$

Hence, integrating on Σ_* starting from S_* and using the GCM condition $\int_{S_*} \beta e^\Phi = 0$, we deduce

$$\left| r \int_{S(u)} \beta e^\Phi \right| \lesssim \epsilon_0 u^{-1-\delta_{dec}} + \int_{\Sigma(u, u_*)} r \left(|\underline{\vartheta}| |\check{\vartheta}| + |\vartheta| |\check{\vartheta}| + r|J| \right).$$

Using the flux estimates of Proposition 7.3.5 we infer that

$$\begin{aligned} \int_{\Sigma(u, u_*)} r \left(|\underline{\vartheta}| |\check{\vartheta}| + |\vartheta| |\check{\vartheta}| \right) &\lesssim \left(\int_{\Sigma(u, u_*)} |\underline{\vartheta}|^2 + |\check{\vartheta}|^2 \right)^{1/2} \left(\int_{\Sigma(u, u_*)} r^2 \left(|\vartheta|^2 + |\check{\vartheta}|^2 \right) \right)^{1/2} \\ &\lesssim \epsilon_0 u^{-1-\delta_{dec}} \left(\int_{\Sigma(u, u_*)} r^2 \left(|\vartheta|^2 + |\check{\vartheta}|^2 \right) \right)^{\frac{1}{2}}. \end{aligned}$$

On the other hand, according to **Ref 1**,

$$\int_{\Sigma(u, u_*)} r^2 \left(|\vartheta|^2 + |\check{\vartheta}|^2 \right) \lesssim \epsilon^2 \int_{\Sigma(u, u_*)} r^{-2} u^{-1-2\delta_{dec}} \lesssim \epsilon^2.$$

Hence,

$$\int_{\Sigma(u, u_*)} r \left(|\underline{\vartheta}| |\check{\vartheta}| + |\vartheta| |\check{\vartheta}| \right) \lesssim \epsilon \epsilon_0 u^{-1-\delta_{dec}}.$$

Similarly, recalling the definition of $J = 3\eta\check{\rho} - \vartheta\underline{\beta}$,

$$\begin{aligned} \int_{\Sigma(u, u_*)} r^2 |J| &\lesssim \int_{\Sigma(u, u_*)} r^2 \left(|\eta| |\check{\rho}| + |\vartheta| |\underline{\beta}| \right) \\ &\lesssim \left(\int_{\Sigma(u, u_*)} \left(|\eta|^2 + r^2 |\underline{\beta}|^2 \right) \right)^{1/2} \cdot \left(\int_{\Sigma(u, u_*)} r^2 \left(|\vartheta|^2 + r^2 |\check{\rho}|^2 \right) \right)^{1/2} \\ &\lesssim \epsilon_0 u^{-1-\delta_{dec}} \cdot \left(\int_{\Sigma(u, u_*)} r^2 \left(|\vartheta|^2 + r^2 |\check{\rho}|^2 \right) \right)^{1/2} \lesssim \epsilon_0 \epsilon u^{-1-\delta_{dec}}. \end{aligned}$$

Therefore,

$$\left| \int_S \beta e^\Phi \right| \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}$$

as desired. To derive (7.3.40) for $k = 1$ we go back to (7.3.42) which we write in the form

$$\begin{aligned} \int_S (\nu_* \beta) e^\Phi &= r^{-1} \nu_* \left(r \int_S \beta e^\Phi \right) + O(r^2 \beta) + r \Gamma_b \cdot \not\partial^{\leq 1} \Gamma_g + \text{l.o.t.} \\ &= r^{-1} \nu_* \left(r \int_S \beta e^\Phi \right) + O(\epsilon r^{-3/2}). \end{aligned}$$

Hence, making use of (7.3.43), the assumptions **Ref1-2** and the condition on r , $r \geq (\epsilon \epsilon_0^{-1})^2 u^{2+2\delta_{dec}}$,

$$\begin{aligned} \left| \int_S (\nu_* \beta) e^\Phi \right| &\lesssim \epsilon_0 u^{-2-2\delta_{dec}} + r \int_S (|\underline{\vartheta}| |\not\partial \vartheta| + |\vartheta| |\not\partial \underline{\vartheta}| + r |J|) + \epsilon r^{-3/2} \\ &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}. \end{aligned}$$

The higher derivative estimates are derived in the same manner. This completes the proof of Proposition 7.3.8. \square

7.3.5 Decay of Ricci and curvature components on Σ_*

The goal here is to prove the following proposition.

Proposition 7.3.9. *The following estimates hold true along Σ_* for all $k \leq k_{small} + 19$*

$$\begin{aligned} \|\underline{\vartheta}, \eta, \underline{\xi}, \underline{\omega}, r\underline{\beta}\|_{\infty, k}^* &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \\ \|\underline{\tilde{\Omega}}, \overline{\underline{\Omega}} + \Upsilon, \zeta, \bar{\varsigma} - 1\|_{\infty, k}^* &\lesssim \epsilon_0 u^{-1-\delta_{dec}}, \\ \|\check{\kappa}, \check{\underline{\kappa}}, r\check{\mu}\|_{\infty, k}^* &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \\ \|\vartheta, \zeta, r\check{\rho}\|_{\infty, k}^* &\lesssim \epsilon_0 r^{-2} u^{-1/2-\delta_{dec}}, \\ \|\beta\|_{\infty, k}^* &\lesssim \epsilon_0 r^{-3} (2r + u)^{-1/2-\delta_{dec}}. \end{aligned} \tag{7.3.44}$$

Proof. Recall that we have already derived the desired estimates for $\underline{\beta}, \underline{\vartheta}, \eta, \underline{\xi}$, see (7.3.15), in Proposition 7.3.5. To prove the remaining estimates we proceed in steps as follows.

Step 1. In this step we prove the estimate,

$$\|\check{\underline{\kappa}}\|_{\infty, k}^* \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}. \tag{7.3.45}$$

This makes use of the GCM condition $\not{d}_2^* \not{d}_1^* \check{\kappa} = 0$, the control of the $\ell = 1$ mode of $\not{d}_1^* \check{\kappa}$ provided by Lemma 7.3.8, the bootstrap assumptions and our dominance condition on r along Σ_* . We also make use of the commutation properties of the tangential derivatives on Σ_* with the GCM condition.

According to part 4 of the elliptic Hodge Lemma 7.1.7 we derive,

$$\|\not{d}_1^* \check{\kappa}\|_{\mathfrak{h}_{k+1}(S)} \lesssim r \|\not{d}_2^* \not{d}_1^* \check{\kappa}\|_{\mathfrak{h}_k(S)} + r^{-2} \left| \int_S e^\Phi \not{d}_1^* \check{\kappa} \right| = r^{-2} \left| \int_S e^\Phi \not{d}_1^* \check{\kappa} \right|.$$

According to Proposition 7.3.8 we have, for all $0 \leq k \leq k_{small} + 20$

$$\left| \int_S e^\Phi \not{d}_1^* \check{\kappa} \right| \lesssim \epsilon_0 u^{-1-\delta_{dec}}.$$

Therefore,

$$\|\not{d}_1^* \check{\kappa}\|_{\mathfrak{h}_{k+1}(S)} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}.$$

Thus,

$$\|\check{\kappa}\|_{\mathfrak{h}_{k+2}(S)} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}$$

and therefore, for $k \leq k_{small} + 20$

$$\|\not{d}^{\leq k} \check{\kappa}\|_{L^\infty(S)} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}.$$

We next commute with ν_* . Since ν_* is tangent to Σ_* and $r \not{d}_2^* \not{d}_1^* \check{\kappa} = 0$, making also use of the commutator Lemma 7.3.1,

$$\not{d}_2^*(\nu_* \not{d}_1^* \check{\kappa}) = r^{-1} [\nu_*, r \not{d}_2^*] \not{d}_1^* \check{\kappa} = \Gamma_b \nu_*(\not{d}_1^* \check{\kappa}) + \text{l.o.t.}$$

Proceeding as before,

$$\begin{aligned} \int_S |\nu_*(\not{d}_1^* \check{\kappa})|^2 &\lesssim r^2 \int_S |\not{d}_2^*(\nu_* \not{d}_1^* \check{\kappa})|^2 + r^{-4} \left| \int_S e^\Phi \nu_*(\not{d}_1^* \check{\kappa}) \right|^2 \\ &\lesssim r^2 \int_S \left| \Gamma_b \nu_*(\not{d}_1^* \check{\kappa}) \right|^2 + \epsilon_0^2 r^{-4} u^{-2-2\delta_{dec}} \\ &\lesssim \epsilon^2 \int_S |\nu_*(\not{d}_1^* \check{\kappa})|^2 + \epsilon_0^2 r^{-4} u^{-2-2\delta_{dec}}. \end{aligned}$$

We can then proceed as before making use of the estimate for the $\ell = 1$ mode of $\nu_* e_{\theta \underline{\kappa}}$ in Proposition 7.3.8, i.e. for all $0 \leq k \leq k_{small} + 20$. The higher ν_* estimates are proved in the same manner.

Step 2. In exactly the same manner, making use of the GCM condition $\not\partial_2^* \not\partial_1^* \mu = 0$ and the estimate for the $\ell = 1$ mode of $e_\theta \mu$ in Lemma 7.3.8, we deduce

$$\|\check{\mu}\|_{\infty, k}^* \lesssim \epsilon_0 r^{-3} u^{-1-\delta_{dec}}. \quad (7.3.46)$$

Step 3. At this point we have established the estimates of Proposition 7.3.9 for $\underline{\beta}, \underline{\vartheta}, \underline{\xi}, \underline{\eta}$ and $\check{\kappa}, \check{\mu}$. Next we derive estimates for $\check{\rho}$ by making use of the quantity \mathbf{q} . We make use of Proposition 7.1.10 according to which we have, relative to the background frame of $^{(ext)}\mathcal{M}$,

$$r^4 \left(\not\partial_2^* \not\partial_1^* \rho + \frac{3}{4} \underline{\kappa} \rho \underline{\vartheta} + \frac{3}{4} \kappa \rho \underline{\vartheta} \right) = \mathbf{q} + r^2 \not\partial^{\leq 2}(\Gamma_b \cdot \Gamma_g) + \text{l.o.t.}$$

Using the estimates **Ref1-2** for the Ricci and curvature coefficients as well as the product Lemma 7.1.6 we derive,

$$\begin{aligned} \|\mathbf{q}\|_{\infty, k}^* &\lesssim \epsilon_0 r^{-1} u^{-1/2-\delta_{dec}}, \\ \|\underline{\vartheta}\|_{\infty, k}^* &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \\ \|r^2 \not\partial^{\leq 2}(\Gamma_b \cdot \Gamma_g)\|_{\infty, k}^* &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}. \end{aligned}$$

Also, using our condition on r along Σ_*

$$\|\vartheta\|_{\infty, k}^* \lesssim \epsilon r^{-2} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}.$$

We deduce,

$$\|\not\partial_2^* \not\partial_1^* \check{\rho}\|_{\infty, k}^* \lesssim \epsilon_0 r^{-5} u^{-1/2-\delta_{dec}}.$$

Also, after commuting ν_* with $r \not\partial_2^*$,

$$\|\not\partial_2^*(\nu_*^i \not\partial_1^* \rho)\|_{\infty, k-i}^* \lesssim \epsilon_0 r^{-5} u^{-1/2-\delta_{dec}}.$$

Since, according to Lemma 7.3.8, we also control the $\ell = 1$ modes of $\nu_*^i e_\theta^j(\rho)$ we can proceed as before²¹ to deduce the desired estimate

$$\|\check{\rho}\|_{\infty, k}^* \lesssim \epsilon_0 r^{-3} u^{-1/2-\delta_{dec}}, \quad \forall k \leq k_{small} + 20. \quad (7.3.47)$$

Step 4. From the definition of $\mu = -\not\partial_1 \zeta - \rho + \frac{1}{4} \vartheta \underline{\vartheta}$, we have,

$$\not\partial_1 \zeta = -\check{\mu} - \rho + \Gamma_g \cdot \Gamma_b.$$

²¹i.e. passing to L^2 , using part 4 of Lemma 7.1.7, and then going back to L^∞ norms as before

We deduce, for all $k \leq k_{small} + 19$,

$$\begin{aligned} \|\not\phi_1^* \not\phi_1 \zeta\|_{\infty, k}^* &\lesssim \|\not\phi_1^* \check{\mu}\|_k^* + \|\not\phi_1^* \check{\rho}\|_{\infty, k}^* + \epsilon_0 r^{-4} u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 r^{-4} u^{-1/2-\delta_{dec}}. \end{aligned}$$

Since according to Proposition 7.3.8 we control the $\ell = 1$ modes of ζ and their ν_* derivatives we deduce, passing as before through L^2 norms,

$$\|\zeta\|_{\infty, k}^* \lesssim \epsilon_0 r^{-2} u^{-1/2-\delta_{dec}}, \quad \forall k \leq k_{small} + 19. \quad (7.3.48)$$

Step 5. We next use the Codazzi equations, the bootstrap assumption for β and the above result for ζ to deduce,

$$\begin{aligned} \not\phi_2 \vartheta &= -2\beta + (e_\theta(\kappa) + \zeta \kappa) + \Gamma_g \cdot \Gamma_g, \\ &= -2\beta + \frac{2}{r} \zeta + \Gamma_g \cdot \Gamma_g. \end{aligned}$$

Making use of the bootstrap assumption for β and the previous estimate for ζ ,

$$\begin{aligned} \|\not\phi_2 \vartheta\|_{\infty, k}^* &\lesssim \|\beta\|_{\infty, k} + r^{-1} \|\zeta\|_{\infty, k}^* + \epsilon_0 r^{-3} u^{-1-\delta_{dec}} \\ &\lesssim \epsilon r^{-7/2-\delta_{dec}} + \epsilon_0 r^{-3} u^{-1/2-\delta_{dec}} \end{aligned}$$

from which we derive, using also the condition $\min_{\Sigma_*} r \geq (\epsilon \epsilon_0^{-1})^2 u^{1+2\delta_{dec}}$,

$$\|\vartheta\|_{\infty, k}^* \lesssim \epsilon_0 r^{-2} u^{-1/2-\delta_{dec}}, \quad \forall k \leq k_{small} + 19. \quad (7.3.49)$$

Step 6. We now estimate β from the equation,

$$e_3 \alpha + \left(\frac{1}{2} \underline{\kappa} - 4\underline{\omega} \right) \alpha = -\not\phi_2^* \beta - \frac{3}{2} \vartheta \rho + (\zeta + 4\eta) \beta.$$

Hence,

$$\begin{aligned} \|\not\phi_2^* \beta\|_{\infty, k-1}^* &\lesssim \|e_3 \alpha\|_{\infty, k-1}^* + r^{-1} \|\alpha\|_{\infty, k-1}^* + r^{-3} \|\vartheta\|_{\infty, k-1}^* + \epsilon_0 r^{-4} (2r + u)^{-1/2-\delta_{dec}} \\ &\lesssim \epsilon_0 r^{-4} (2r + u)^{-1/2-\delta_{dec}}. \end{aligned}$$

We deduce, in particular, using our previous estimate for the $\ell = 1$ mode of β

$$\begin{aligned} \|\beta\|_{\mathfrak{h}_{k+1}(S)} &\lesssim r \|\not\phi_2^* \beta\|_{\mathfrak{h}_k(S)} + r^{-2} \left| \int_S \beta e^\Phi \right| \\ &\lesssim \epsilon_0 r^{-2} (2r + u)^{-1/2-\delta_{dec}} + \epsilon_0 r^{-3} u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 r^{-2} (2r + u)^{-1/2-\delta_{dec}}. \end{aligned}$$

We deduce, arguing as before,

$$\|\beta\|_{\infty, k}^* \lesssim \epsilon_0 r^{-3} (2r + u)^{-1/2-\delta_{dec}}, \quad \forall k \leq k_{small} + 19. \quad (7.3.50)$$

Remark 7.3.10. We can in fact prove a stronger estimate by making use of the full strength of the estimate for α in **Ref 2**,

$$\begin{aligned} |\alpha| &\lesssim \log(1+u)r^{-3}(2r+u)^{-1/2-\delta_{extra}}, \\ |e_3\alpha| &\lesssim r^{-4}(2r+u)^{-1/2-\delta_{extra}}. \end{aligned}$$

Hence,

$$\|\phi_2^*\beta\|_{\infty,k}^* \lesssim \epsilon_0 \log(1+u)r^{-4}(2r+u)^{-1/2-\delta_{extra}} + \epsilon_0 r^{-5}u^{-1/2-\delta_{dec}}$$

and thus

$$\|\beta\|_{\infty,k}^* \lesssim \epsilon_0 \log(1+u)r^{-3}(2r+u)^{-1/2-\delta_{extra}}. \quad (7.3.51)$$

Step 7. Next, we estimate $\check{\omega}$ by making use of the first identity²² in Proposition 2.2.19,

$$\phi_1^*\check{\omega} = -\frac{1}{4}\kappa\underline{\xi} + \frac{1}{2}e_3\underline{\zeta} - \frac{1}{2}\underline{\beta} + \frac{1}{2}\kappa\underline{\zeta} + \frac{1}{2}\vartheta\underline{\zeta} - \frac{1}{4}\vartheta\underline{\xi}.$$

We deduce, using (7.3.48) and the dominance condition for r on Σ_* ,

$$\begin{aligned} \|\phi_1^*\check{\omega}\|_{2,k-1} &\lesssim r^{-1}\|\underline{\xi}\|_{2,k-1} + \|e_3\underline{\zeta}\|_{2,k-1} + r^{-1}\|\underline{\zeta}\|_{2,k-1} + \|\underline{\beta}\|_{2,k-1} + \epsilon_0 r^{-2}u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 r^{-1}u^{-1-\delta_{dec}}. \end{aligned}$$

Thus, using the Poincare and Sobolev inequalities,

$$\|\check{\omega}\|_{\infty,k} \lesssim \epsilon_0 r^{-1}u^{-1-\delta_{dec}}. \quad (7.3.52)$$

Step 8. We estimate $\check{\zeta}, \bar{\zeta} - 1$ Recall that, see 7.1.8,

$$a_* = \frac{2}{\varsigma} - \Upsilon + \frac{r}{2}\underline{A}.$$

Also, recall that, see (7.1.9),

$$\underline{\Omega} = e_3(r) = -\Upsilon + \frac{r}{2}\underline{A}, \quad \text{on } \Sigma_*.$$

Step 9. We determine $\check{\zeta}, \check{\omega}$ using the equations, see (2.2.19),

$$\begin{aligned} e_\theta \log \varsigma &= \frac{1}{2}(\eta - \zeta), \\ e_\theta(\check{\Omega}) &= -\underline{\xi} - (\eta - \zeta)\underline{\Omega}. \end{aligned}$$

²²This is a precise version of the identity in Proposition 7.1.12.

Hence, using the estimates for $\eta, \zeta, \underline{\xi}$ and the bootstrap assumptions already derived,

$$\begin{aligned}\|\check{\zeta}\|_{\mathfrak{h}_{k+1}} &\lesssim \epsilon_0 u^{-1-\delta_{dec}}, \\ \|\check{\underline{\Omega}}\|_{\mathfrak{h}_{k+1}} &\lesssim \epsilon_0 u^{-1-\delta_{dec}}.\end{aligned}$$

Step 10. We estimate $\bar{\varsigma} - 1$ and $\bar{\underline{\Omega}} + \Upsilon$ on Σ_* using the equations (7.1.8), (7.1.9),

$$\begin{aligned}a_* &= -\frac{2}{\varsigma} + \Upsilon - \frac{r}{2}\underline{A}, \\ \underline{\Omega} &= -\Upsilon + \frac{r}{2}\underline{A},\end{aligned}$$

and the GCM condition for a_* , see (7.1.1),

$$a_*|_{SP} = -1 - \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}.$$

We deduce,

$$\frac{2}{\varsigma|_{SP}} = 2 - \frac{r}{2}\underline{A}|_{SP} = 2 - (\underline{\Omega} + \Upsilon)|_{SP}$$

i.e.,

$$2\left(1 - \frac{1}{\varsigma|_{SP}}\right) = \underline{\Omega}|_{SP} + \Upsilon.$$

Since

$$\bar{\varsigma} = \varsigma|_{SP} + \check{\zeta}|_{SP}, \quad \bar{\underline{\Omega}} = \underline{\Omega}|_{SP} + \check{\underline{\Omega}}|_{SP}$$

we deduce, in view of the above estimates for $\check{\zeta}, \check{\underline{\Omega}}$,

$$|\bar{\varsigma} - 1| \lesssim \left|\bar{\underline{\Omega}} + \Upsilon\right| + \epsilon_0 u^{-1-\delta_{dec}}.$$

It thus remains to estimate

$$\bar{\underline{\Omega}} + \Upsilon = \frac{r}{2}\bar{\underline{A}}.$$

This can be done with the help of the following identity,

$$\bar{\underline{A}}^{\mathbf{S}} = \frac{1}{\varsigma^{\mathbf{S}}}\left(\bar{\check{\zeta}}^{\mathbf{S}}\bar{\check{\underline{\Omega}}}^{\mathbf{S}} - \bar{\check{\zeta}}^{\mathbf{S}}\bar{\check{\underline{A}}}^{\mathbf{S}}\right) \tag{7.3.53}$$

from which we infer that,

$$\left| \overline{\Omega} + \Upsilon \right| \lesssim \epsilon_0 u^{-1-\delta_{dec}}$$

and therefore also

$$|\overline{\zeta} - 1| \lesssim \epsilon_0 u^{-1-\delta_{dec}}.$$

To prove (7.3.53) we apply Lemma 7.3.2 to deduce,

$$\begin{aligned} e_3(r) + a_* &= \nu_*(r) = \frac{r}{2} \overline{\zeta(\underline{\kappa} + a_* \kappa)} = \frac{r}{2} \zeta^{-1} (\overline{\zeta \underline{\kappa}} + \overline{\zeta \check{\kappa}}) + \zeta^{-1} \overline{\zeta a_*} \\ &= -\Upsilon \zeta^{-1} \overline{\zeta} + \frac{r}{2} \zeta^{-1} \overline{\zeta \check{\kappa}} + \zeta^{-1} \overline{\zeta a_*}. \end{aligned}$$

Since $e_3(r) = -\Upsilon + \frac{r}{2} \underline{A}$ we infer that

$$-\Upsilon + \frac{r}{2} \underline{A} + a_* = -\Upsilon \zeta^{-1} \overline{\zeta} + \frac{r}{2} \zeta^{-1} \overline{\zeta \check{\kappa}} + \zeta^{-1} \overline{\zeta a_*}$$

and therefore

$$\underline{A} = \frac{2}{r} \left(\Upsilon - a_* - \Upsilon \zeta^{-1} \overline{\zeta} + \frac{r}{2} \zeta^{-1} \overline{\zeta \check{\kappa}} + \zeta^{-1} \overline{\zeta a_*} \right).$$

In particular, multiplying by ζ and taking the average, we infer that

$$\overline{\zeta \underline{A}} = \overline{\zeta \check{\kappa}},$$

and hence

$$\overline{A} = \frac{1}{\overline{\zeta}} \left(\overline{\zeta \check{\kappa}} - \overline{\zeta \check{A}} \right)$$

as stated. The estimates for all other derivatives of $\zeta, \check{\Omega}, \overline{\zeta} - 1, \overline{\Omega} + \Upsilon$ can be derived as before. This concludes the proof of Proposition 7.3.9. \square

7.4 Control in $(ext)\mathcal{M}$, Part I

7.4.1 Preliminaries

Commutation Lemmas

Here and below we write schematically

$$\not\partial = r \not\partial, \quad \not\partial_{\nearrow} = \{r e_4, \not\partial\}, \quad \not\partial = (e_3, r e_4, r \not\partial), \quad \mathbf{T} = \frac{1}{2} (e_3 + \Upsilon e_4).$$

Lemma 7.4.1. *We have, schematically,*

$$[\mathbf{T}, e_4] = r^{-1}\Gamma_b \mathfrak{d}_{\nearrow}, \quad [\mathfrak{d}, e_4] = \check{\Gamma}_g \mathfrak{d}_{\nearrow} + \Gamma_g. \quad (7.4.1)$$

Also,

$$\mathbf{T}(r) = \frac{1}{2r} \underline{A} = r^{-1}\Gamma_b.$$

Proof. The identity for $[\mathfrak{d}, e_4]$ has already been discussed in Corollary 7.1.5. According to lemma 2.2.14 we have,

$$[\mathbf{T}, e_4] = \left(\left(\underline{\omega} - \frac{m}{r^2} \right) - \frac{m}{2r} \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{e_4(m)}{r} \right) e_4 + (\eta + \zeta)e_\theta,$$

In view of **Ref 4** and bootstrap assumptions **Ref 2** the factors of e_4 and e_θ , on the right hand side behave at worst like Γ_b . Thus schematically $[\mathbf{T}, e_4] = r^{-1}\Gamma_b \mathfrak{d}_{\nearrow}$. \square

Transport Lemmas

The following lemma will be used repeatedly in what follows.

Lemma 7.4.2. *If f verifies the transport equation*

$$e_4(f) + \frac{p}{2}\bar{\kappa}f = F,$$

we have for fixed u and any $r_0 \leq r \leq r_$,*

$$\begin{aligned} r^p \|f\|_\infty(u, r) &\lesssim r_0^p \|f\|_\infty(u, r_{\mathcal{H}}) + \int_{r_0}^r \lambda^p \|F\|_\infty(u, \lambda) d\lambda, \\ r^p \|f\|_\infty(u, r) &\lesssim r_*^p \|f\|_\infty(u, r_*) + \int_r^{r_*} \lambda^p \|F\|_\infty(u, \lambda) d\lambda, \end{aligned} \quad (7.4.2)$$

where r is the area radius at fixed u .

Proof. According to Corollary 2.2.12 we have $e_4(r^p f) = r^p F$. The desired estimates follow easily by integration with respect to the affine parameter s , recall that $e_4(s) = 1$. \square

Proposition 7.4.3. *The following inequalities hold true for all $k \leq k_{large} - 5$, $r_0 \leq r \leq r_*$*

$$\begin{aligned} r^p \|f\|_{\infty, k}(u, r) &\lesssim r_*^p \|f\|_{\infty, k}(u, r_*) + \int_r^{r_*} \lambda^p \|F\|_{\infty, k}(u, \lambda) d\lambda, \\ r^p \|f\|_{\infty, k}(u, r) &\lesssim r_0^p \|f\|_{\infty, k}(u, r_0) + \int_{r_0}^r \lambda^p \|F\|_{\infty, k}(u, \lambda) d\lambda. \end{aligned} \quad (7.4.3)$$

Proof. Commuting the equation $e_4(r^p f) = r^p F$ with \mathfrak{D} , applying the commutation Lemma 7.4.1 and our bootstrap assumptions on Γ_g we derive,

$$\begin{aligned} e_4(r^p |\mathfrak{D}f|) &\lesssim r^p |\mathfrak{D}F| + r^p |\mathfrak{D}F| + \epsilon r^{-2} r^p (|\mathfrak{D}f| + r e_4(r^p f)) \\ &\lesssim r^p (|\mathfrak{D}F| + |F|) + \epsilon r^{-2} r^p (|\mathfrak{D}f| + |f|). \end{aligned}$$

Similarly, commuting with \mathbf{T} ,

$$e_4(\mathbf{T}(r^p f)) = \mathbf{T}(r^p F) - [\mathbf{T}, e_4](r^p f) = r^p F - p r^{p-1} \mathbf{T}(r) F - r^{-1} \Gamma_b \mathfrak{D}_{\nearrow} r^p f.$$

Hence,

$$e_4(r^p \mathbf{T}f) = r^p \mathbf{T}(F) - p r^{p-1} \mathbf{T}(r) F - r^{-1} \Gamma_b \mathfrak{D}_{\nearrow} r^p f - p e_4(r^{p-1} \mathbf{T}(r) f)$$

i.e.,

$$e_4(r^p |\mathbf{T}f|) \lesssim r^p (|\mathbf{T}F| + |F|) + O(\epsilon r^{-2}) (r^p |F| + |\mathfrak{D}f| + |f|).$$

Similarly, commuting the equation with $r e_4$ we derive,

$$e_4(r^p |r f|) \lesssim r^p (|r e_4 F| + |F|) + O(\epsilon r^{-2}) (r^p |F| + |\mathfrak{D}f| + |f|).$$

Integrating the inequalities,

$$\begin{aligned} e_4(r^p |\mathfrak{D}f|) &\lesssim r^p (|\mathfrak{D}F| + |F|) + \epsilon r^{-2} r^p (|\mathfrak{D}f| + |f|) \\ e_4(r^p |\mathbf{T}f|) &\lesssim r^p (|\mathbf{T}F| + |F|) + \epsilon r^{-2} r^p (|\mathfrak{D}f| + |f|) \\ e_4(r^p |r f|) &\lesssim r^p (|r e_4 F| + |F|) + \epsilon r^{-2} (r^p |F| + |\mathfrak{D}f| + |f|) \end{aligned}$$

and applying Gronwall we derive the desired estimates in (7.4.3) for $k = 1$.

Repeating the procedure for $\tilde{\mathfrak{D}}^k$, any combination of derivatives of the form $\tilde{\mathfrak{D}}^k = \mathbf{T}^{k_1} \mathfrak{D}^{k_2}$ with $k_1 + k_2 = k$, estimating the corresponding commutators using our assumptions **Ref 1**, we deduce for all $0 \leq k \leq k_{large} - 5$,

$$e_4(r^p |\tilde{\mathfrak{D}}^{\leq k} f|) \lesssim r^p |\tilde{\mathfrak{D}}^{\leq k} F| + \epsilon r^{-2} r^p |\tilde{\mathfrak{D}}^{\leq k} f|$$

and the desired estimates follow by integration. \square

Transport equations for $\ell = 1$ modes

To estimate $\ell = 1$ modes we make use of the following.

Lemma 7.4.4. *The following equation holds true for reduced scalars $\psi \in \mathfrak{s}_1^{(ext)}\mathcal{M}$.*

$$e_4 \left(\int_S \psi e^\Phi \right) = \int_S e_4(\psi) e^\Phi + \frac{3}{2} \bar{\kappa} \int_S \psi e^\Phi + \int_S \frac{1}{2} (3\check{\kappa} - \vartheta) \psi e^\Phi. \quad (7.4.4)$$

Proof. This is an immediate consequence of Proposition (2.2.9). Indeed according to it and $e_4\Phi = \frac{1}{2}(\kappa - \vartheta)$,

$$\begin{aligned} e_4 \left(\int_S \psi e^\Phi \right) &= \int_S (e_4(\psi e^\Phi) + \kappa \psi e^\Phi) = \int_S \left(e_4(\psi) + \frac{1}{2}(3\kappa - \vartheta)\psi \right) e^\Phi \\ &= \int_S \left(e_4(\psi) + \frac{3}{2}\bar{\kappa}\psi \right) e^\Phi + \int_S \frac{1}{2} (3\check{\kappa} - \vartheta) \psi e^\Phi \\ &= \int_S e_4(\psi) e^\Phi + \frac{3}{2}\bar{\kappa} \int_S \psi e^\Phi + \int_S \frac{1}{2} (3\check{\kappa} - \vartheta) \psi e^\Phi \end{aligned}$$

as desired. □

7.4.2 Proposition 7.4.5

In what follows we prove the stronger estimates in terms of powers of r for the quantities $\check{\kappa}, \check{\mu}, \vartheta, \zeta, \check{\xi}, \beta, \check{\rho}$. More precisely we establish the following.

Proposition 7.4.5. *The following estimates hold true in $(ext)\mathcal{M}$ for all $k \leq k_{small} + 20$*

$$\begin{aligned} \|\check{\kappa}\|_{\infty, k} &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \\ \|\check{\mu}\|_{\infty, k} &\lesssim \epsilon_0 r^{-3} u^{-1-\delta_{dec}}. \end{aligned} \quad (7.4.5)$$

Also, for all $k \leq k_{small} + 18$

$$\begin{aligned} \|\vartheta, \zeta, \check{\xi}, r\check{\rho}\|_{\infty, k} &\lesssim \epsilon_0 r^{-2} u^{-1/2-\delta_{dec}}, \\ \|\beta\|_{\infty, k} &\lesssim \epsilon_0 r^{-3} (2r + u)^{-1/2-\delta_{dec}}, \\ \|e_3\beta\|_{\infty, k} &\lesssim \epsilon_0 r^{-4} u^{-1/2-\delta_{dec}}, \\ \|e_\theta K\|_{\infty, k} &\lesssim \epsilon_0 r^{-4} u^{-1/2-\delta_{dec}}. \end{aligned} \quad (7.4.6)$$

Remark 7.4.6. *Note that in fact β admits the stronger estimate, for all $k \leq k_{small} + 18$,*

$$\|\beta\|_{\infty, k} \lesssim \epsilon_0 \log(1 + u) r^{-3} (2r + u)^{-1/2-\delta_{extra}}. \quad (7.4.7)$$

7.4.3 Estimates for $\check{\kappa}, \check{\mu}$ in $^{(ext)}\mathcal{M}$

Step 1. We prove the following estimates for $\check{\kappa}$ in $^{(ext)}\mathcal{M}$.

$$\|\check{\kappa}\|_{\infty,k} \lesssim \epsilon_0 r^{-5/2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 20. \quad (7.4.8)$$

We make use of the equation

$$e_4(\check{\kappa}) + \bar{\kappa} \check{\kappa} = F := -\frac{1}{2} \check{\kappa}^2 - \frac{1}{2} \overline{\check{\kappa}^2} - \frac{1}{2} (\vartheta^2 - \overline{\vartheta^2}).$$

In view of our assumptions **Ref1-2** and Lemma 7.1.6

$$\|F\|_{\infty,k}(u, \lambda) \lesssim \epsilon_0 \lambda^{-7/2} u^{-1-\delta_{dec}}.$$

Applying Proposition 7.4.3 we deduce,

$$\begin{aligned} r^2 \|\check{\kappa}\|_{\infty,k}(u, r) &\lesssim r_*^2 \|\check{\kappa}\|_{\infty,k}(u, r_*) + \epsilon_0 u^{-1-\delta_{dec}} \int_r^{r_*} \lambda^2 \lambda^{-7/2} d\lambda \\ &\lesssim r_*^2 \|\check{\kappa}\|_{\infty}(u, r_*) + \epsilon_0 r^{-1/2} u^{-1-\delta_{dec}}. \end{aligned}$$

In view of the control on the last slice we infer that, everywhere in $^{(ext)}\mathcal{M}$,

$$\|\check{\kappa}\|_{\infty,k}(u, r) \lesssim \epsilon_0 r^{-5/2} u^{-1-\delta_{dec}}.$$

Step 2. We prove the estimate,

$$\|\check{\mu}\|_{\infty,k} \lesssim \epsilon_0 r^{-3} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 20. \quad (7.4.9)$$

Recall that we have

$$\begin{aligned} e_4(\check{\mu}) + \frac{3}{2} \bar{\kappa} \check{\mu} &= -\frac{3}{2} \bar{\mu} \check{\kappa} + F, \\ F &:= -\frac{3}{2} \check{\mu} \check{\kappa} + \frac{1}{2} \overline{\check{\mu} \check{\kappa}} + \text{Err}[e_4 \check{\mu}] - \overline{\text{Err}[e_4 \check{\mu}]}, \\ \text{Err}[e_4 \check{\mu}] &= -\frac{1}{8} \bar{\kappa} \vartheta^2 - \vartheta \not\partial_2^* \zeta - \vartheta \zeta^2 + \left(2e_\theta(\kappa) - 2\beta + \frac{3}{2} \kappa \zeta \right) \zeta. \end{aligned}$$

In view of Lemma 7.1.6 we check,

$$\|F\|_{\infty,k}(u, \lambda) \lesssim \epsilon_0 \lambda^{-9/2} u^{-1-\delta_{dec}}.$$

Applying Proposition 7.4.3 and the estimates on the last slice for $\check{\mu}$ we deduce

$$\begin{aligned} r^3 \|\widetilde{\not\partial}^k \check{\mu}\|_{\infty,k}(u, \lambda) &\lesssim r_*^3 \|\widetilde{\not\partial}^k \check{\mu}\|_{\infty,k}(u, r_*) + \epsilon_0 u^{-1-\delta_{dec}} \int_r^{r_*} \lambda^3 \lambda^{-9/2} \\ &\lesssim r_*^3 \|\widetilde{\not\partial}^k \check{\mu}\|_{\infty,k}(u, r_*) + \epsilon_0 u^{-1-\delta_{dec}} r^{-1/2} \\ &\lesssim \epsilon_0 u^{-1-\delta_{dec}} \end{aligned}$$

from which the desired estimate (7.4.9) follows.

7.4.4 Estimates for the $\ell = 1$ modes in $(ext)\mathcal{M}$

We extend the validity of Lemma 7.3.8 to the entire region $(ext)\mathcal{M}$.

Lemma 7.4.7. *The following estimates hold true on $(ext)\mathcal{M}$ or all $k \leq k_{small} + 19$,*

$$\begin{aligned}
\left\| \int_S \beta e^\Phi \right\|_{\infty, k} &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \\
\left\| \int_S \zeta e^\Phi \right\|_{\infty, k} &\lesssim \epsilon_0 r u^{-1-\delta_{dec}}, \\
\left\| \int_S e_\theta(\rho) e^\Phi \right\|_{\infty, k} &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \\
\left\| \int_S e_\theta(\underline{\kappa}) e^\Phi \right\|_{\infty, k} &\lesssim \epsilon_0 u^{-1-\delta_{dec}}, \\
\left\| \int_S \underline{\beta} e^\Phi \right\|_{\infty, k} &\lesssim \epsilon_0 u^{-1-\delta_{dec}}.
\end{aligned} \tag{7.4.10}$$

Proof. We first note that the estimate for the $\ell = 1$ mode of $\check{\mu}$ is an immediate consequence of the estimate (7.4.9). To prove the remaining estimates we proceed in steps as follows.

Step 1. Observe that the estimates of Lemma 7.3.8 remain valid when we replace the norms $\|\cdot\|_{\infty, k}^*$ by $\|\cdot\|_{\infty, k}$. To show this it suffices to prove estimates for re_4 of all $\ell = 1$ modes. This can easily be achieved with the help of Lemma 7.4.4 and our e_4 transport equations for $\zeta, \check{\rho}, \check{\mu}, \check{\kappa}, \underline{\beta}$.

Step 2. We establish the estimate,

$$\left\| \int_S \beta e^\Phi \right\|_{\infty, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 20. \tag{7.4.11}$$

In view of (7.4.4) and the Bianchi identity for $e_4(\beta)$

$$\begin{aligned}
e_4 \left(\int_S \beta e^\Phi \right) &= \int_S e_4 \beta e^\Phi + \frac{3}{2} \bar{\kappa} \int_S \beta e^\Phi + \int_S \frac{1}{2} (3\check{\kappa} - \vartheta) \beta e^\Phi \\
&= \int_S (-2\kappa \beta + \not{d}_2 \alpha + \zeta \alpha) e^\Phi + \frac{3}{2} \bar{\kappa} \int_S \beta e^\Phi + \int_S \frac{1}{2} (3\check{\kappa} - \vartheta) \beta e^\Phi \\
&= -\frac{\bar{\kappa}}{2} \int_S \beta e^\Phi + \int_S \left(\zeta \alpha + \frac{1}{2} (-\check{\kappa} + \vartheta) \beta \right) e^\Phi,
\end{aligned}$$

and hence

$$e_4 \left(\int_S \beta e^\Phi \right) + \frac{\bar{\kappa}}{2} \int_S \beta e^\Phi = \int_S \left(\zeta \alpha + \frac{1}{2}(-\check{\kappa} + \vartheta)\beta \right) e^\Phi. \quad (7.4.12)$$

Recall that

$$|(\alpha, \beta)| \lesssim \epsilon r^{-3} (2r + u)^{-1/2 - \delta_{dec}}.$$

We deduce,

$$\begin{aligned} \left| e_4 \left(r \int_S \beta e^\Phi \right) \right| &\lesssim r \epsilon_0 r^{-2} u^{-1/2 - \delta_{dec}} r^{-3} (2r + u)^{-1/2 - \delta_{dec}} \int_S |e^\Phi| \\ &\lesssim \epsilon_0 r^{-1} u^{-1/2 - \delta_{dec}} (2r + u)^{-1/2 - \delta_{dec}} \\ &\lesssim \epsilon_0 r^{-1 - \delta} u^{-1 - \delta_{dec}} \end{aligned}$$

i.e., in view of the estimate on Σ_* , everywhere on $^{(ext)}\mathcal{M}$,

$$\left| \int_S \beta e^\Phi \right| \lesssim \epsilon r^{-1} u^{-1 - \delta_{dec}}. \quad (7.4.13)$$

Commuting with \mathbf{T} , $\not\partial$ and re_4 we also easily deduce,

$$\left\| \tilde{\delta}^{k_2} (re_4)^{k_1} \int_S \beta e^\Phi \right\|_\infty \lesssim \epsilon r^{-1} u^{-1 - \delta_{dec}}, \quad \forall k_1 + k_2 \leq k_{small} + 20$$

from which (7.4.11) follows.

Step 3. We prove the estimate,

$$\left\| \int_S \zeta e^\Phi \right\|_{\infty, k} \lesssim \epsilon_0 r^{1/2} u^{-1 - \delta_{dec}}, \quad k \leq k_{small} + 19 \quad (7.4.14)$$

which is better than the desired estimate in Lemma 7.4.7. This follows, as for the corresponding estimate on Σ_* , by projecting the Codazzi equation for ϑ on the $\ell = 1$ mode

$$\int_S \zeta e^\Phi = \frac{r}{2\Upsilon} \left(2 \int_S \beta e^\Phi - \int_S e_\theta(\kappa) e^\Phi - \int_S \vartheta \zeta e^\Phi - \int_S \left(\kappa - \frac{2\Upsilon}{r} \right) \zeta e^\Phi \right).$$

Note that in view of the estimates for $\check{\kappa}$ in (7.4.8) already established²³ we have,

$$\left\| \int_S e_\theta(\kappa) e^\Phi \right\|_{\infty, k} \lesssim \epsilon_0 r^{-1/2} u^{-1 - \delta_{dec}}, \quad k \leq k_{small} + 19.$$

²³Note that the estimate for $\check{\kappa}$ is stronger in powers of r than the corresponding bootstrap assumption.

Also, making use of (7.4.11),

$$\left\| \int_S \beta e^\Phi \right\|_{\infty, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 19.$$

Thus, since Υ is bounded away from zero in $(ext)\mathcal{M}$, we easily deduce

$$\left\| \int_S \zeta e^\Phi \right\|_{\infty, k} \lesssim \epsilon_0 r^{1/2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 19.$$

Step 4. We prove the estimate,

$$\left\| \int_S e_\theta(\rho) e^\Phi \right\|_{\infty, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 19. \quad (7.4.15)$$

We proceed as in Step 4 of the proof of Lemma 7.3.8. In view of the definition of μ and the identity $\not{d}_1^* \not{d}_1 = \not{d}_2 \not{d}_2^* + 2K$ we write,

$$\begin{aligned} \int_S e_\theta(\rho) e^\Phi &= - \int_S e_\theta(\mu) e^\Phi + \int_S \not{d}_1^* \not{d}_1 \zeta e^\Phi + \frac{1}{4} \int_S e_\theta(\not{v}\not{v}) e^\Phi \\ &= - \int_S e_\theta(\mu) e^\Phi + \int_S (\not{d}_2 \not{d}_2^* + 2K) \zeta e^\Phi + \frac{1}{4} \int_S e_\theta(\not{v}\not{v}) e^\Phi \\ &= - \int_S e_\theta(\mu) e^\Phi + \frac{2}{r^2} \int_S \zeta e^\Phi + 2 \int_S \left(K - \frac{1}{r^2} \right) \zeta e^\Phi + \frac{1}{4} \int_S e_\theta(\not{v}\not{v}) e^\Phi. \end{aligned}$$

Together with the above estimate for the $\ell = 1$ mode of ζ , the estimate (7.4.9) for $\check{\mu}$ and the bootstrap assumptions, we infer that

$$\left\| \int_S e_\theta(\rho) e^\Phi \right\|_{\infty, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}.$$

Step 5. We prove the estimate,

$$\left\| \int_S e_\theta(\underline{\kappa}) e^\Phi \right\|_{\infty, k} \lesssim \epsilon_0 u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 19. \quad (7.4.16)$$

As in the corresponding estimate on the last slice we make use of the remarkable identity for the $\ell = 1$ mode of $e_\theta(K)$, i.e.

$$\int_S e_\theta(\rho) e^\Phi + \frac{1}{4} \int_S e_\theta(\underline{\kappa}) e^\Phi - \frac{1}{4} \int_S e_\theta(\not{v}\not{v}) e^\Phi = 0.$$

We infer

$$\int_S e_{\theta}(\underline{\kappa})e^{\Phi} = -2r \int_S e_{\theta}(\rho)e^{\Phi} - \frac{r}{2} \int_S \underline{\kappa}e_{\theta}(\kappa)e^{\Phi} + \frac{r}{2} \int_S e_{\theta}(\vartheta\underline{\vartheta})e^{\Phi} - \frac{r}{2} \int_S \left(\kappa - \frac{2}{r}\right) e_{\theta}(\underline{\kappa})e^{\Phi}.$$

The estimate (7.4.16) follows easily from with the above estimate for the $\ell = 1$ mode of $e_{\theta}(\rho)$, the estimate for $\check{\kappa}$ in (7.4.8) and the bootstrap assumptions.

Step 6. We prove the estimate,

$$\left\| \int_S \underline{\beta}e^{\Phi} \right\|_{\infty, k} \lesssim \epsilon_0 u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 19. \quad (7.4.17)$$

Projecting the Codazzi for $\underline{\vartheta}$ on the $\ell = 1$ mode, we have

$$-2 \int_S \underline{\beta}e^{\Phi} + \int_S e_{\theta}(\underline{\kappa})e^{\Phi} - \int_S \underline{\kappa}\zeta e^{\Phi} + \int_S \underline{\vartheta}\zeta e^{\Phi} = 0$$

and hence

$$\int_S \underline{\beta}e^{\Phi} = \frac{1}{2} \int_S e_{\theta}(\underline{\kappa})e^{\Phi} + \frac{\Upsilon}{r} \int_S \zeta e^{\Phi} - \frac{1}{2} \int_S \left(\underline{\kappa} + \frac{2\Upsilon}{r}\right) \zeta e^{\Phi} + \frac{1}{2} \int_S \underline{\vartheta}\zeta e^{\Phi}.$$

The desired estimate follows easily in view of the above estimates for the $\ell = 1$ mode of $e_{\theta}(\underline{\kappa})$, the $\ell = 1$ mode of ζ and the bootstrap assumptions. \square

7.4.5 Completion of the Proof of Proposition 7.4.5

We prove the second part of Proposition 7.4.5, i.e. we prove for all $k \leq k_{small} + 18$

$$\begin{aligned} \|\vartheta, \zeta, \check{\kappa}, r\check{\rho}\|_{\infty, k} &\lesssim \epsilon_0 r^{-2} u^{-1/2-\delta_{dec}}, \\ \|\beta\|_{\infty, k} &\lesssim \epsilon_0 r^{-3} (2r+u)^{-1/2-\delta_{dec}}, \\ \|e_3\beta\|_{k, \infty} &\lesssim \epsilon_0 r^{-4} u^{-1/2-\delta_{dec}}, \\ \|e_{\theta}K\|_{\infty, k} &\lesssim \epsilon_0 r^{-4} u^{-1/2-\delta_{dec}}. \end{aligned} \quad (7.4.18)$$

We also prove the stronger estimate for β (see Remark 7.4.6)

$$\|\beta\|_{\infty, k} \lesssim \epsilon_0 \log(1+u) r^{-3} (2r+u)^{-1/2-\delta_{extra}}. \quad (7.4.19)$$

Proof. We proceed in steps as follows.

Step 1. We derive the estimate,

$$\|\vartheta\|_{\infty,k} \lesssim \epsilon_0 r^{-2} u^{-1/2-\delta_{dec}}, \quad \forall k \leq k_{small} + 19, \quad (7.4.20)$$

with the help of the equation $e_4\vartheta + \bar{\kappa}\vartheta = F := -2\alpha - \bar{\kappa}\vartheta$ and the corresponding estimate on the last slice.

Note that,

$$\|\alpha\|_{\infty,k} \lesssim \epsilon_0 r^{-3-\delta} (2r+u)^{-1/2-\delta_{dec}}$$

where $\delta > 0$ is a small constant, $\delta < \delta_{extra} - \delta_{dec}$. Thus, using also the product estimates of Lemma 7.1.6, we easily check that,

$$\|F\|_{\infty,k} \lesssim \epsilon_0 r^{-3-\delta} u^{-1/2-\delta_{dec}} + \epsilon_0 r^{-7/2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 20.$$

Making use of Proposition 7.4.3 we deduce, for all $k \leq k_{small} + 19$,

$$r^2 \|\mathfrak{d}^k \vartheta\|_{\infty,k}(u, r) \lesssim r_*^2 \|\mathfrak{d}^k \vartheta\|_{\infty,k}(u, r_*) + \epsilon_0 u^{-1/2-\delta_{dec}} \int_r^{r_*} \lambda^{-1-\delta} d\lambda.$$

Thus, in view of the results on the last slice Σ_* , we deduce,

$$\|\mathfrak{d}^k \vartheta\|_{\infty}(u, r) \lesssim r^{-2} u^{-1/2-\delta_{dec}}.$$

Step 2. We derive the estimate,

$$\|\beta\|_{\infty,k} \lesssim \epsilon_0 r^{-3} (2r+u)^{-1/2-\delta_{dec}}, \quad \forall k \leq k_{small} + 19. \quad (7.4.21)$$

We proceed exactly as in the estimates for β on the last slice Σ_* by making use of the Bianchi identity $e_3\alpha + (\frac{1}{2}\underline{\kappa} - 4\underline{\omega})\alpha = -\not{d}_2^*\beta - \frac{3}{2}\vartheta\rho + 5\zeta\beta$, from which we deduce,

$$\|\not{d}_2^*\beta\|_{\infty,k-1} \lesssim \|e_3\alpha\|_{\infty,k-1} + r^{-1}\|\alpha\|_{\infty,k-1} + r^{-3}\|\vartheta\|_{\infty,k-1} + \epsilon_0 r^{-5} u^{-1-\delta_{dec}}.$$

Thus, in view of the above estimate for ϑ and **Ref 2** for α ,

$$\|\not{d}_2^*\beta\|_{\infty,k-1} \lesssim \epsilon_0 r^{-4} (2r+u)^{-1/2-\delta_{dec}} + \epsilon_0 r^{-5} u^{-1/2-\delta_{dec}}.$$

On the other hand we have, according to (7.4.11),

$$\left\| \int_S \beta e^\Phi \right\|_{\infty,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 20.$$

Estimate (7.4.21) follows then easily, according to the part 4 of the elliptic Hodge Lemma 7.1.7.

As mentioned in Remark 7.4.6 we can prove a stronger estimate for β . Indeed we have, in view of **Ref 2**.

$$\begin{aligned} |\alpha| &\lesssim \log(1+u)r^{-3}(2r+u)^{-1/2-\delta_{extra}}, \\ |e_3\alpha| &\lesssim r^{-4}(2r+u)^{-1/2-\delta_{extra}}. \end{aligned}$$

Hence, using the equation $e_3\alpha + (\frac{1}{2}\underline{\kappa} - 4\underline{\omega})\alpha = -\not{d}_2^*\beta - \frac{3}{2}\vartheta\rho + 5\zeta\beta$,

$$\|\not{d}_2^*\beta\|_{\infty,k} \lesssim \epsilon_0 \log(1+u)r^{-4}(2r+u)^{-1/2-\delta_{extra}} + \epsilon_0 r^{-5}u^{-1/2-\delta_{dec}}.$$

According to Lemma 7.1.7

$$\|\beta\|_{\mathfrak{h}_{k+1}(S)} \lesssim r\|\not{d}_2^*\beta\|_{\mathfrak{h}_k(S)} + r^{-2}\left|\int_S e^\Phi\beta\right|$$

and thus, in view of the estimate (7.4.11) for the $\ell = 1$ mode of β ,

$$\begin{aligned} \|\beta\|_{\mathfrak{h}_{k+1}(S)} &\lesssim \epsilon_0 \log(1+u)r^{-2}(2r+u)^{-1/2-\delta_{extra}} + \epsilon_0 r^{-3}u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 \log(1+u)r^{-2}(2r+u)^{-1/2-\delta_{extra}}. \end{aligned}$$

The estimates for the T and e_4 derivatives are derived in the same manner. and hence,

$$\|\beta\|_{\infty,k} \lesssim \epsilon_0 \log(1+u)r^{-3}(2r+u)^{-1/2-\delta_{extra}}, \quad \forall k \leq k_{small} + 19. \quad (7.4.22)$$

This improvement is needed in the next step.

Step 3. We derive the estimate

$$\|\zeta\|_{\infty,k} \lesssim \epsilon_0 r^{-2}u^{-1/2-\delta_{dec}}, \quad \forall k \leq k_{small} + 19. \quad (7.4.23)$$

For this we make use of the transport equation for ζ ,

$$e_4\zeta + \bar{\kappa}\zeta = F := -\beta + \Gamma_g \cdot \Gamma_g$$

and the improved estimate for β in the previous step. Thus, making use of the product Lemma 7.1.6,

$$\begin{aligned} \|F\|_{\infty,k} &\lesssim \|\beta\|_{\infty,k} + \epsilon_0 r^{-7/2}u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 \log(1+u)r^{-3}(2r+u)^{-1/2-\delta_{extra}} + \epsilon_0 r^{-7/2}u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 r^{-3-\delta}u^{-1/2-\delta_{dec}} + \epsilon_0 r^{-7/2}u^{-1-\delta_{dec}}. \end{aligned}$$

Making use of Proposition 7.4.3 we deduce

$$r^2 \|\mathfrak{D}^k \zeta\|_{\infty, k}(u, r) \lesssim r_*^2 \|\mathfrak{D}^k \zeta\|_{\infty, k}(u, r_*) + \epsilon_0 u^{-1/2-\delta_{dec}} \int_r^{r_*} \lambda^{-1-\delta} d\lambda.$$

Thus, in view of the estimates on the last slice,

$$r^2 \|\mathfrak{D}^k \zeta\|_{\infty}(u, r) \lesssim \epsilon_0 u^{-1/2-\delta_{dec}}, \quad k \leq k_{small} + 19$$

as desired.

Step 4. We derive the estimate

$$\|\check{\rho}\|_{\infty, k} \lesssim \epsilon_0 r^{-3} u^{-1/2-\delta_{dec}}, \quad \forall k \leq k_{small} + 18. \quad (7.4.24)$$

We make use of the definition of μ from which we infer that,

$$\check{\mu} = -\not{d}_1 \zeta - \check{\rho} + \Gamma_g \cdot \Gamma_b.$$

Hence, in view of the product Lemma and the estimates already derived, for all $k \leq k_{small} + 18$,

$$\begin{aligned} \|\check{\rho}\|_{\infty, k} &\lesssim r^{-1} \|\zeta\|_{\infty, k+1} + \|\check{\mu}\|_{\infty, k} + \epsilon_0 r^{-3} u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 r^{-3} u^{-1/2-\delta_{dec}} \end{aligned}$$

as desired.

Step 5. We derive the estimate

$$\|\check{\kappa}\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1/2-\delta_{dec}}, \quad \forall k \leq k_{small} + 18. \quad (7.4.25)$$

We make use of the equation

$$e_4 \check{\kappa} + \frac{1}{2} \bar{\kappa} \check{\kappa} = F := -2 \not{d}_1 \zeta - \frac{1}{2} \check{\kappa} \bar{\kappa} + 2\check{\rho} + \Gamma_g \cdot \Gamma_b.$$

In view of the previously derived estimates,

$$\|F\|_{\infty, k} \lesssim \epsilon_0 r^{-3} u^{-1/2-\delta_{dec}}, \quad k \leq k_{small} + 18.$$

Making use of Proposition 7.4.3 we deduce, for all $k \leq k_{small} + 18$,

$$\begin{aligned} r \|\mathfrak{D}^k \check{\kappa}\|_{\infty, k}(u, r) &\lesssim r_* \|\mathfrak{D}^k \check{\kappa}\|_{\infty, k}(u, r_*) + \epsilon_0 u^{-1/2-\delta_{dec}} \int_r^{r_*} \lambda^{-2} d\lambda \\ &\lesssim r_* \|\mathfrak{D}^k \check{\kappa}\|_{\infty, k}(u, r_*) + \epsilon_0 r^{-1} u^{-1/2-\delta_{dec}}. \end{aligned}$$

Thus, in view of the estimates on the last slice,

$$r \|\mathfrak{D}^k \check{\kappa}\|_{\infty}(u, r) \lesssim \epsilon_0 (r_*)^{-1} u^{-1/2-\delta_{dec}} + \epsilon_0 r^{-1} u^{-1/2-\delta_{dec}}$$

from which the desired estimate easily follows. \square

Step 6. We derive the estimate

$$\|e_3\beta\|_{\infty,k} \lesssim \epsilon_0 r^{-4} u^{-1/2-\delta_{dec}}, \quad \forall k \leq k_{small} + 18. \quad (7.4.26)$$

making use of the equation $e_3\beta + (\underline{\kappa} - 2\underline{\omega})\beta = -\not{d}_1^*\rho + 3\zeta\rho + \Gamma_g\underline{\beta} + \Gamma_b\alpha$ and the estimates derived above for β , $\not{d}_1^*\rho$, ζ . Hence,

$$\begin{aligned} \|e_3\beta\|_{\infty,k} &\lesssim r^{-1}\|\beta\|_{\infty,k} + \|\not{d}_1^*\rho\|_{\infty,k} + r^{-3}\|\zeta\|_{\infty,k} + \epsilon_0 r^{-4} u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 r^{-4} u^{-1/2-\delta_{dec}}. \end{aligned}$$

Step 7. As a corollary of the above estimates (see also **Ref 4**) we also derive, in $^{(ext)}\mathcal{M}$,

$$\begin{aligned} \|K - \overline{K}\|_{\infty,k-1} &\lesssim \epsilon_0 r^{-3} u^{-1/2-\delta_{dec}}, \quad k \leq k_{small} + 19, \\ \|K - \frac{1}{r^2}\|_{\infty,k-1} &\lesssim \epsilon_0 r^{-3} u^{-1/2-\delta_{dec}}, \quad k \leq k_{small} + 19. \end{aligned} \quad (7.4.27)$$

In view of the definition of K we have,

$$e_\theta(K) = -e_\theta \left(\check{\rho} - \frac{1}{4}\kappa\check{\underline{\kappa}} - \frac{1}{4}\kappa\check{\underline{\kappa}} + \frac{1}{4}\vartheta\underline{\vartheta} \right).$$

Thus, in view of the above estimates,

$$\|e_\theta K\|_{\infty,k} \lesssim \epsilon_0 r^{-4} u^{-1/2-\delta_{dec}}$$

from which the desired estimate easily follows.

7.5 Control in $^{(ext)}\mathcal{M}$, Part II

We derive the crucial decay estimates which imply, in particular, decay of order $u^{-1-\delta_{dec}}$ for all quantities in Γ and \check{R} (except $\underline{\xi}, \underline{\omega}, \underline{\Omega}$ which will be treated separately) in the interior. More precisely we prove the following,

Proposition 7.5.1. *The following estimates hold true in $^{(ext)}\mathcal{M}$, for all $k \leq k_{small} + 8$.*

$$\|\vartheta, \zeta, \eta, \check{\underline{\kappa}}, \underline{\vartheta}, r\beta, r\check{\rho}, r\underline{\beta}, \underline{\alpha}\|_{\infty,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}. \quad (7.5.1)$$

To prove the proposition we make use of the fact that we already have good decay estimates in terms of powers of u for $\check{\kappa}, \check{\mu}$. We also derive below decay estimates for various renormalized quantities.

7.5.1 Estimate for η

We start with the following simple estimate for η in terms of ζ .

$$\|\eta\|_{\infty,k} \lesssim \|\zeta\|_{\infty,k} + \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 17. \quad (7.5.2)$$

This can be derived by propagation from the last slice with the help of the equation,

$$e_4(\eta - \zeta) + \frac{1}{2}\kappa(\eta - \zeta) = -\frac{1}{2}\vartheta(\eta - \zeta) = \Gamma_g \cdot \Gamma_b.$$

Note that

$$\|\Gamma_g \cdot \Gamma_b\|_{\infty,k} \lesssim \epsilon_0 r^{-3} u^{-1-\delta_{dec}}.$$

Thus making use of Proposition 7.4.3 we deduce

$$\begin{aligned} r\|\eta - \zeta\|_{\infty,k}(u, r) &\lesssim r_*\|\eta - \zeta\|_{\infty,k}(u, r_*) + \int_r^{r_*} \lambda \|\Gamma_g \cdot \Gamma_b\|_{\infty,k}(u, \lambda) \\ &\lesssim r_*\|\eta - \zeta\|_{\infty,k}(u, r_*) + \epsilon_0 u^{-1-\delta_{dec}} \end{aligned}$$

with r_* the value of r on $C(u) \cap \Sigma_*$. On the last slice we have derived the estimates, recorded in Proposition 7.3.5 and Proposition 7.4.5

$$\begin{aligned} \|\eta\|_{\infty,k}^* &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \\ \|\zeta\|_{\infty,k}^* &\lesssim \epsilon_0 r^{-2} u^{-1/2-\delta_{dec}}. \end{aligned}$$

In view of the dominance condition on r on Σ_* we deduce,

$$\|\eta - \zeta\|_{\infty,k}^*(u, r) \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}$$

and therefore also,

$$r_*\|\eta - \zeta\|_{\infty,k}(u, r_*) \lesssim \epsilon_0 u^{-1-\delta_{dec}}.$$

Therefore,

$$r\|\eta\|_{\infty,k}(u, r) \lesssim r\|\zeta\|_{\infty,k}(u, r) + \epsilon_0 u^{-1-\delta_{dec}}$$

as desired.

7.5.2 Crucial Lemmas

We start with the following lemma.

Lemma 7.5.2. *The $\mathfrak{s}_1(\mathcal{M})$ reduced tensor*

$$\Xi : = r^2 (e_\theta(\underline{\kappa}) + 4r \not\partial_1^* \not\partial_1 \zeta - 2r^2 \not\partial_1^* \not\partial_1 \beta) \quad (7.5.3)$$

verifies in $^{(ext)}\mathcal{M}$ the estimate,

$$\|\Xi\|_{\infty, k} \lesssim \epsilon_0 u^{-1-\delta_{dec}}, \quad \forall k \leq k_{small} + 13. \quad (7.5.4)$$

Proof. To calculate $e_4 \Xi$ we make use of the equations,

$$\begin{aligned} e_4(\underline{\kappa}) + \frac{1}{2} \kappa \underline{\kappa} &= -2 \not\partial_1 \zeta + 2\rho - \frac{1}{2} \vartheta \underline{\vartheta} + 2\zeta^2, \\ e_4 \zeta + \kappa \zeta &= -\beta - \vartheta \zeta, \\ e_4 \beta + 2\kappa \beta &= \not\partial_2 \alpha + \zeta \alpha. \end{aligned}$$

Since we already have an estimate for $\check{\mu}$ we re-express $\rho = -\mu - \not\partial_1 \zeta + \frac{1}{4} \vartheta \underline{\vartheta}$ and derive,

$$e_4(\underline{\kappa}) + \frac{1}{2} \kappa \underline{\kappa} = -2\mu - 4 \not\partial_1 \zeta + 2\zeta^2.$$

Commuting with $\not\partial_1^*$ and making use of $[\not\partial_1^*, e_4] = \frac{1}{2}(\kappa + \vartheta) \not\partial_1^*$ we derive,

$$e_4(\not\partial_1^* \underline{\kappa}) + \kappa \not\partial_1^* \underline{\kappa} + \frac{1}{2} \underline{\kappa} \not\partial_1^* \kappa = -\not\partial_1^* \check{\mu} - 4 \not\partial_1^* \not\partial_1 \zeta + 2 \not\partial_1^* (\zeta^2) + \vartheta \not\partial_1^* \underline{\kappa}. \quad (7.5.5)$$

Hence, since $e_4(r) = \frac{r}{2} \bar{\kappa}$,

$$\begin{aligned} e_4(r^2 \not\partial_1^* \underline{\kappa}) &= r^2(\kappa - \bar{\kappa}) \not\partial_1^* \underline{\kappa} - \frac{1}{2} r^2 \underline{\kappa} \not\partial_1^* \check{\kappa} - 4r^2 \not\partial_1^* \not\partial_1 \zeta - r^2 \not\partial_1^* \check{\mu} + r^2 (\not\partial_1^* (\zeta^2) + \vartheta \not\partial_1^* \underline{\kappa}) \\ &= -\frac{1}{2} r^2 \underline{\kappa} \not\partial_1^* \check{\kappa} - 4r^2 \not\partial_1^* \not\partial_1 \zeta + \text{Err}_1 \end{aligned}$$

where,

$$\text{Err}_1 : = -\frac{1}{2} r^2 \underline{\kappa} \not\partial_1^* \check{\kappa} - r^2 \not\partial_1^* \check{\mu} + r^2 (\check{\kappa} \not\partial_1^* \underline{\kappa} + \not\partial_1^* (\zeta^2) + \vartheta \not\partial_1^* \underline{\kappa}).$$

In view of the estimate already established for $\check{\kappa}, \check{\mu}$ and the product Lemma 7.1.6 we check,

$$\|\text{Err}_1\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 19.$$

To simplify notation we introduce the following.

Definition 7.5.3. We say that a quantity $\psi \in \mathfrak{s}_k(\mathcal{M})$ is $r^{-p}Good_a$ provided that it verifies the estimate, everywhere in $(ext)\mathcal{M}$,

$$\|\psi\|_{\infty, k} \lesssim \epsilon_0 r^{-p} u^{-1-\delta_{dec}}, \quad \forall k \leq k_{small} + a. \quad (7.5.6)$$

Using this notation we write,

$$e_4(r^2 \not\partial_1^* \underline{\kappa}) = -4r^2 \not\partial_1^* \not\partial_1 \zeta + r^{-2} Good_{19}. \quad (7.5.7)$$

Using the same notation the transport equation for ζ can be written in the form,

$$e_4 \zeta + \bar{\kappa} \zeta = -\beta - \vartheta \zeta - \check{\kappa} \zeta = -\beta + r^{-7/2} Good_{20}.$$

Commuting with $(r \not\partial_1^*)(r \not\partial_1)$ (making us of Lemma 7.4.1) we derive

$$e_4(r^2 \not\partial_1^* \not\partial_1 \zeta) + \bar{\kappa}(r^2 \not\partial_1^* \not\partial_1 \zeta) = -r^2 \not\partial_1^* \not\partial_1 \beta + r^{-7/2} Good_{18}.$$

Since $e_4(r) = \frac{1}{2} r \bar{\kappa}$ we deduce,

$$e_4(r^3 \not\partial_1^* \not\partial_1 \zeta) = -\frac{1}{2} \bar{\kappa} r^3 \not\partial_1^* \not\partial_1 \zeta - r^3 \not\partial_1^* \not\partial_1 \beta + r^{-5/2} Good_{18}. \quad (7.5.8)$$

Similarly the transport equation for β takes the form

$$e_4 \beta + 2\bar{\kappa} \beta = \not\partial_2 \alpha + \zeta \alpha - 2\check{\kappa} \beta = \not\partial_2 \alpha + r^{-9/2} Good_{20}$$

and,

$$e_4(r^2 \not\partial_1^* \not\partial_1 \beta) + 2\bar{\kappa} r^2 \not\partial_1^* \not\partial_1 \beta = r^2 \not\partial_1^* \not\partial_1 \not\partial_2 \alpha + r^{-9/2} Good_{18}.$$

As before, since $e_4(r) = \frac{1}{2} r \bar{\kappa}$, we deduce,

$$e_4(r^4 \not\partial_1^* \not\partial_1 \beta) = -\bar{\kappa} r^4 \not\partial_1^* \not\partial_1 \beta + r^4 \not\partial_1^* \not\partial_1 \not\partial_2 \alpha + r^{-5/2} Good_{18}. \quad (7.5.9)$$

Combining (7.5.7)–(7.5.9) we deduce,

$$\begin{aligned} e_4 \Xi &= e_4 \left[r^2 \left(-\not\partial_1^* \underline{\kappa} + 4r \not\partial_1^* \not\partial_1 \zeta - 2r^2 \not\partial_1^* \not\partial_1 \beta \right) \right] \\ &= 4r^2 \not\partial_1^* \not\partial_1 \zeta + 4 \left(-\frac{1}{2} \bar{\kappa} r^3 \not\partial_1^* \not\partial_1 \zeta - r^3 \not\partial_1^* \not\partial_1 \beta \right) - 2 \left(-\bar{\kappa} r^4 \not\partial_1^* \not\partial_1 \beta + r^4 \not\partial_1^* \not\partial_1 \not\partial_2 \alpha \right) \\ &+ r^{-2} Good_{18} \\ &= -2 \left(\bar{\kappa} - \frac{2}{r} \right) r^3 \not\partial_1^* \not\partial_1 \zeta + 2r^4 \left(\bar{\kappa} - \frac{2}{r} \right) \not\partial_1^* \not\partial_1 \beta - 2r^4 \not\partial_1^* \not\partial_1 \not\partial_2 \alpha + r^{-2} Good_{18}. \end{aligned}$$

Making use of **Ref 4** estimates for $\bar{\kappa} - \frac{2}{r}$ and the estimates for α in **Ref 2**, i.e.,

$$r^4 |\not{d}_1^* \not{d}_1 \not{d}_2 \alpha| \lesssim \epsilon_0 r^{-1} (2r + u)^{-1 - \delta_{extra}} \lesssim \epsilon_0 r^{-1 - \delta} u^{-1 - \delta_{extra} + \delta}, \quad 0 < \delta < \delta_{extra}$$

i.e.,

$$r^4 \not{d}_1^* \not{d}_1 \not{d}_2 \alpha = r^{-1 - \delta} \text{Good}_{13}$$

we thus deduce,

$$e_4 \Xi = r^{-1 - \delta} \text{Good}_{13}.$$

We deduce,

$$\|\Xi\|_{\infty, k}(u, r) \lesssim \|\Xi\|_{\infty, k}(u, r_*) + \epsilon_0 u^{-1 - \delta_{dec}} \int_r^{r_*} \lambda^{-1 - \delta} d\lambda, \quad \forall k \leq k_{small} + 13.$$

In view of the estimates on the last slice it is easy to check that

$$\|\Xi\|_{\infty, k}(u, r_*) \lesssim \epsilon_0 u^{-1 - \delta_{dec}}, \quad \forall k \leq k_{small} + 13.$$

Indeed, on the last slice,

$$\begin{aligned} \|\not{d}_1^* \underline{\kappa}\|_{\infty, k} &\lesssim \epsilon_0 r^{-3} u^{-1/2 - \delta_{dec}}, \\ \|\not{d}_1^* \not{d}_1 \zeta\|_{\infty, k} &\lesssim \epsilon_0 r^{-4} u^{-1/2 - \delta_{dec}}, \\ \|\not{d}_1^* \not{d}_1 \beta\|_{\infty, k} &\lesssim \epsilon_0 r^{-5} u^{-1/2 - \delta_{dec}}. \end{aligned}$$

Hence, since $r \gg u$ on Σ_* ,

$$\|\Xi\|_{\infty, k}(u, r_*) \lesssim \epsilon_0 r^{-1} u^{-1/2 - \delta_{dec}} \lesssim \epsilon_0 u^{-1 - \delta_{dec}}.$$

Thus everywhere on $^{(ext)}\mathcal{M}$,

$$\|\Xi\|_{\infty, k} \lesssim \epsilon_0 u^{-1 - \delta_{dec}}, \quad \forall k \leq k_{small} + 13 \quad (7.5.10)$$

as desired. \square

In the following lemma, we make use of the control we have already established for $\mathfrak{q}, \alpha, \underline{\alpha}, \check{\kappa}, \check{\mu}$ in $^{(ext)}\mathcal{M}$ to derive two nontrivial relations between angular derivatives of $\zeta, \check{\kappa}$ and β .

Remark 7.5.4. According to Theorem M3 we only have good estimates for $\underline{\alpha}$ along \mathcal{T} and on the last slice Σ_* . To keep track of this fact we denote by $r^{-p} \text{Good}_a(\underline{\alpha})$ those $r^{-p} \text{Good}_a$ terms which depend linearly on $\underline{\alpha}$ and their derivatives.

Lemma 7.5.5. *Let A, B be the operators $A := \not\partial_2^* \not\partial_2 - 3\bar{\rho}$, $B = \not\partial_2^* \not\partial_2 + 2K$. The following identities hold true,*

$$\begin{aligned} AB \not\partial_2^* \zeta - \frac{3}{4} \bar{\kappa} \bar{\rho} \not\partial_2^* e_\theta(\underline{\kappa}) &\in r^{-6} \text{Good}_{14}(\underline{\alpha}) \\ A^2 B \not\partial_2^* \beta + \frac{9}{8} (\bar{\kappa} \bar{\rho})^2 \not\partial_2^* e_\theta(\underline{\kappa}) &\in r^{-9} \text{Good}_9(\underline{\alpha}) \end{aligned} \quad (7.5.11)$$

Proof. In view of the improved control for α in Theorem M2, $\underline{\alpha}$ in Theorem M3, and \mathfrak{q} in Theorem M1, the bootstrap assumptions and product lemma, and the control we have already derived for $\check{\kappa}$ and $\check{\mu}$ in $(ext)\mathcal{M}$, we obtain

$$\begin{aligned} \not\partial_2 \vartheta + 2\beta - \bar{\kappa} \zeta &\in r^{-3} \text{Good}_{20}, && \text{Codazzi and control of } \check{\kappa}, \\ \not\partial_2 \underline{\vartheta} + 2\underline{\beta} - e_\theta(\underline{\kappa}) + \bar{\kappa} \zeta &\in r^{-3} \text{Good}_{20}, && \text{Codazzi,} \\ \not\partial_2^* \beta + \frac{3}{2} \bar{\rho} \vartheta &\in r^{-3} \text{Good}_{15}, && \text{Bianchi and control of } \alpha, \\ \not\partial_2^* \underline{\beta} + \frac{3}{2} \bar{\rho} \underline{\vartheta} &\in r^{-2} \text{Good}_{15}(\underline{\alpha}), && \text{Bianchi and control of } \underline{\alpha}, \\ \not\partial_2^* \not\partial_1^* \rho + \frac{3}{4} \bar{\kappa} \bar{\rho} \vartheta + \frac{3}{4} \bar{\kappa} \bar{\rho} \underline{\vartheta} &\in r^{-4} \text{Good}_{18}, && (7.1.28) \text{ and control of } \mathfrak{q}, \end{aligned} \quad (7.5.12)$$

where we used Codazzi for the two first inequalities, Bianchi for the third and fourth inequalities, the definition of μ for the fifth one, and the identity relating \mathfrak{q} and $\not\partial_2^* \not\partial_1^* \rho$ for the last one.

Combining the first statement with the third and the second with the fourth we infer that,

$$\begin{aligned} (\not\partial_2^* \not\partial_2 - 3\bar{\rho}) \vartheta - \bar{\kappa} \not\partial_2^* \zeta &\in r^{-3} \text{Good}_{14}, \\ (\not\partial_2^* \not\partial_2 - 3\bar{\rho}) \underline{\vartheta} - \not\partial_2^* e_\theta(\underline{\kappa}) + \bar{\kappa} \not\partial_2^* \zeta &\in r^{-2} \text{Good}_{14}(\underline{\alpha}), \end{aligned}$$

or, setting

$$A := \not\partial_2^* \not\partial_2 - 3\bar{\rho},$$

$$\begin{aligned} A \vartheta - \bar{\kappa} \not\partial_2^* \zeta &\in r^{-3} \text{Good}_{14}, \\ A \underline{\vartheta} - \not\partial_2^* e_\theta(\underline{\kappa}) + \bar{\kappa} \not\partial_2^* \zeta &\in r^{-2} \text{Good}_{14}(\underline{\alpha}). \end{aligned} \quad (7.5.13)$$

From the fifth equations we deduce,

$$A \left(\not\partial_2^* \not\partial_1^* \rho + \frac{3}{4} \bar{\kappa} \bar{\rho} \vartheta + \frac{3}{4} \bar{\kappa} \bar{\rho} \underline{\vartheta} \right) \in r^{-6} \text{Good}_{18}$$

i.e.,

$$A \not\partial_2^* \not\partial_1^* \rho + \frac{3}{4} \bar{\kappa} \bar{\rho} A \vartheta + \frac{3}{4} \bar{\kappa} \bar{\rho} A \underline{\vartheta} \in r^{-6} \text{Good}_{18}.$$

Making use of (7.5.13) we deduce,

$$A \not\partial_2^* \not\partial_1^* \rho + \frac{3}{4} \bar{\kappa} \bar{\rho} (\bar{\kappa} \not\partial_2^* \zeta) + \frac{3}{4} \bar{\kappa} \bar{\rho} (\not\partial_2^* e_\theta(\underline{\kappa}) - \bar{\kappa} \not\partial_2^* \zeta) \in r^{-6} \text{Good}_{14}(\underline{\alpha}).$$

Hence, simplifying,

$$A \not\partial_2^* \not\partial_1^* \rho + \frac{3}{4} \bar{\kappa} \bar{\rho} \not\partial_2^* e_\theta(\underline{\kappa}) \in r^{-6} \text{Good}_{14}(\underline{\alpha}). \quad (7.5.14)$$

Next, in view of the identity $\not\partial_1^* \not\partial_1 = \not\partial_2 \not\partial_2^* + 2K$,

$$\begin{aligned} (\not\partial_2^* \not\partial_2 + 2K) \not\partial_2^* \zeta &= \not\partial_2^* \not\partial_2 \not\partial_2^* \zeta + 2K \not\partial_2^* \zeta \\ &= \not\partial_2^* (\not\partial_2 \not\partial_2^* \zeta + 2K \zeta) - 2 \not\partial_2^* K \zeta \\ &= \not\partial_2^* \not\partial_1^* \not\partial_1 \zeta + r^{-9/2} \text{Good}_{19}. \end{aligned}$$

Recalling the definition of $\mu = -\not\partial_1 \zeta - \rho + \frac{1}{4} \vartheta \underline{\vartheta}$ and the product Lemma we write

$$\not\partial_1^* \not\partial_1 \zeta = -\not\partial_1^* \mu - \not\partial_1^* \rho + \frac{1}{4} \not\partial_1^* (\vartheta \underline{\vartheta}) = -\not\partial_1^* \mu - \not\partial_1^* \rho + r^{-4} \text{Good}_{19}.$$

In view of the estimates for $\check{\mu}$ we have already established we deduce,

$$\not\partial_1^* \not\partial_1 \zeta = -\not\partial_1^* \rho + r^{-4} \text{Good}_{19}.$$

Thus,

$$(\not\partial_2^* \not\partial_2 + 2K) \not\partial_2^* \zeta = -\not\partial_2^* \not\partial_1^* \rho + r^{-5} \text{Good}_{18}. \quad (7.5.15)$$

Therefore, making use of (7.5.14)

$$\begin{aligned} A(\not\partial_2^* \not\partial_2 + 2K) \not\partial_2^* \zeta &= -A \not\partial_2^* \not\partial_1^* \rho + r^{-6} \text{Good}_{16}(\underline{\alpha}) \\ &= \frac{3}{4} \bar{\kappa} \bar{\rho} \not\partial_2^* e_\theta(\underline{\kappa}) + r^{-6} \text{Good}_{14}(\underline{\alpha}) \end{aligned}$$

i.e.,

$$A(\not\partial_2^* \not\partial_2 + 2K) \not\partial_2^* \zeta - \frac{3}{4} \bar{\kappa} \bar{\rho} \not\partial_2^* e_\theta(\underline{\kappa}) = r^{-6} \text{Good}_{14}(\underline{\alpha}) \quad (7.5.16)$$

as desired.

To prove the second statement of the lemma we write, using (7.5.13)

$$(\phi_2^* \phi_2 + 2K)A\vartheta = \bar{\kappa} (\phi_2^* \phi_2 + 2K) \phi_2^* \zeta + r^{-5} \text{Good}_{12}(\underline{\alpha}).$$

Hence applying A and making use of (7.5.16),

$$\begin{aligned} A(\phi_2^* \phi_2 + 2K)A\vartheta &= \bar{\kappa} A(\phi_2^* \phi_2 + 2K) \phi_2^* \zeta + r^{-7} \text{Good}_{10}(\underline{\alpha}) \\ &= \frac{3}{4} \bar{\kappa}^2 \bar{\rho} \phi_2^* e_\theta(\underline{\kappa}) + r^{-7} \text{Good}_{10}(\underline{\alpha}). \end{aligned}$$

Finally, making use of the relation $\phi_2^* \beta + \frac{3}{2} \bar{\rho} \vartheta \in r^{-3} \text{Good}_{15}$, we have

$$\begin{aligned} A^2(\phi_2^* \phi_2 + 2K) \phi_2^* \beta &= A(\phi_2^* \phi_2 + 2K)A \phi_2^* \beta + r^{-9} \text{Good}_9(\underline{\alpha}) \\ &= -\frac{3}{2} \bar{\rho} A(\phi_2^* \phi_2 + 2K)A\vartheta + r^{-9} \text{Good}_9(\underline{\alpha}) \\ &= -\frac{3}{2} \bar{\rho} \left(\frac{3}{4} \bar{\kappa}^2 \bar{\rho} \phi_2^* e_\theta(\underline{\kappa}) + r^{-8} \text{Good}_{10}(\underline{\alpha}) \right) + r^{-9} \text{Good}_9 \\ &= -\frac{9}{8} \bar{\kappa}^2 \bar{\rho}^2 \phi_2^* e_\theta(\underline{\kappa}) + r^{-9} \text{Good}_9(\underline{\alpha}) \end{aligned}$$

as desired. This concludes the proof of the lemma. \square

Corollary 7.5.6. *The $\mathfrak{s}_1(\mathcal{M})$ tensor $e_\theta(\underline{\kappa}) = -\phi_1^* \check{\kappa}$ verifies the following fifth order elliptic equation in $(ext)\mathcal{M}$*

$$A^2 \phi_2^* (e_\theta \underline{\kappa}) - \frac{12m}{r^3} A \phi_2^* e_\theta(\underline{\kappa}) + \frac{36m^2}{r^6} \phi_2^* e_\theta(\underline{\kappa}) \in r^{-7} \text{Good}_8(\underline{\alpha}). \quad (7.5.17)$$

Proof. According to Lemma 7.5.5

$$\begin{aligned} AB \phi_2^* \zeta - \frac{3}{4} \bar{\kappa} \bar{\rho} \phi_2^* e_\theta(\underline{\kappa}) &\in r^{-6} \text{Good}_{14}(\underline{\alpha}), \\ A^2 B \phi_2^* \beta + \frac{9}{8} (\bar{\kappa} \bar{\rho})^2 \phi_2^* e_\theta(\underline{\kappa}) &\in r^{-9} \text{Good}_9(\underline{\alpha}), \end{aligned}$$

we have

$$\begin{aligned} A^2 B \phi_2^* \zeta &= \frac{3}{4} \bar{\kappa} \bar{\rho} A \phi_2^* e_\theta(\underline{\kappa}) + r^{-8} \text{Good}_{12}(\underline{\alpha}), \\ A^2 B \phi_2^* \beta &= -\frac{9}{8} (\bar{\kappa} \bar{\rho})^2 \phi_2^* e_\theta(\underline{\kappa}) + r^{-9} \text{Good}_9(\underline{\alpha}). \end{aligned} \quad (7.5.18)$$

In view of Lemma 7.5.2 we have on $(ext)\mathcal{M}$,

$$e_\theta(\underline{\kappa}) + 4r \phi_1^* \phi_1 \zeta - 2r^2 \phi_1^* \phi_1 \beta \in r^{-2} \text{Good}_{13}. \quad (7.5.19)$$

Thus,

$$A^2 \phi_2^* \left(e_\theta(\underline{\kappa}) + 4r \phi_1^* \phi_1 \zeta - 2r^2 \phi_1^* \phi_1 \beta \right) \in r^{-7} \text{Good}_8.$$

Making use of

$$\phi_2^* \phi_1^* \phi_1 = \phi_2^* \left(\phi_2 \phi_2^* + 2K \right) = (\phi_2^* \phi_2 + 2K) \phi_2^* - e_\theta(K),$$

we deduce,

$$\begin{aligned} A^2 \phi_2^* (e_\theta \underline{\kappa}) &= -4r A^2 \phi_2^* \phi_1^* \phi_1 \zeta + 2r^2 A^2 \phi_2^* \phi_1^* \phi_1 \beta + r^{-7} \text{Good}_8 \\ &= -4r A^2 (\phi_2^* \phi_2 + 2K) \phi_2^* \zeta + 2r^2 A^2 (\phi_2^* \phi_2 + 2K) \phi_2^* \beta + r^{-7} \text{Good}_8 \\ &= -4r A^2 B \phi_2^* \zeta + 2r^2 A^2 B \phi_2^* \beta + r^{-7} \text{Good}_8. \end{aligned}$$

Thus, in view of the lemma,

$$\begin{aligned} A^2 \phi_2^* (e_\theta \underline{\kappa}) &= -3r (\bar{\kappa} \bar{\rho} A \phi_2^* e_\theta(\underline{\kappa}) + r^{-8} \text{Good}_{12}) - \frac{9}{4} r^2 \left((\bar{\kappa} \bar{\rho})^2 \phi_2^* e_\theta(\underline{\kappa}) + r^{-9} \text{Good}_9(\underline{\alpha}) \right) \\ &\quad + r^{-7} \text{Good}_8. \end{aligned}$$

We deduce,

$$A^2 \phi_2^* e_\theta \underline{\kappa} + 3r (\bar{\kappa} \bar{\rho}) A \phi_2^* e_\theta(\underline{\kappa}) + \frac{9}{4} r^2 (\bar{\kappa} \bar{\rho})^2 \phi_2^* e_\theta(\underline{\kappa}) \in r^{-7} \text{Good}_8(\underline{\alpha}).$$

Finally,

$$A^2 \phi_2^* e_\theta \underline{\kappa} - \frac{12m}{r^3} A \phi_2^* e_\theta(\underline{\kappa}) + \frac{36m^2}{r^6} \phi_2^* e_\theta(\underline{\kappa}) \in r^{-7} \text{Good}_8(\underline{\alpha})$$

as desired. □

Lemma 7.5.7. *We have the following Poincaré inequality on $^{(ext)}\mathcal{M}$ for $f \in \mathfrak{s}_2(\mathcal{M})$ with $A = (\phi_2^* \phi_2 - 3\bar{\rho})$*

$$\int_S f \left(A^2 - \frac{12m}{r^3} A + \frac{36m^2}{r^6} \right) f \geq \frac{1}{4r^2} \int_S (\phi_2 f)^2 + \frac{9}{r^4} \int_S f^2.$$

Proof. Recall that we have the following Poincaré inequality for ϕ_2

$$\int_S (\phi_2 f)^2 \geq 4 \int_S K f^2.$$

Since $|K - r^{-2}| \lesssim \epsilon r^{-2}$,

$$\begin{aligned} \int_S fAf &= \int_S f(\not{d}_2^* \not{d}_2 - 3\bar{\rho})f \geq \int_S (4K - 3\bar{\rho})f^2 \\ &= \left(\frac{4}{r^2} + \frac{6m}{r^3} + O(r^{-2}\epsilon) \right) \int_S f^2. \end{aligned}$$

Since A is positive self-adjoint,

$$\begin{aligned} \int_S fA^2f &= \int_S (A^{1/2}f)A(A^{1/2}f) = \left(\frac{4}{r^2} + \frac{6m}{r^3} + O(r^{-2}\epsilon) \right) \int_S |A^{1/2}f|^2 \\ &= \left(\frac{4}{r^2} + \frac{6m}{r^3} + O(r^{-2}\epsilon) \right) \int_S fAf \end{aligned}$$

This yields

$$\begin{aligned} \int_S f \left(A^2f - \frac{12m}{r^3}Af \right) &= \left(\frac{4}{r^2} + \frac{6m}{r^3} - \frac{12m}{r^3} + O(r^{-2}\epsilon) \right) \int_S fAf \\ &= \left(\frac{4}{r^2} - \frac{6m}{r^3} + O(r^{-2}\epsilon) \right) \int_S fAf, \end{aligned}$$

and therefore,

$$\begin{aligned} \int_S f \left(A^2 - \frac{12m}{r^3}A + \frac{36m^2}{r^6} \right) &\geq \left(\frac{4}{r^2} - \frac{6m}{r^3} + O(r^{-2}\epsilon) \right) \int_S fAf + \frac{36m^2}{r^6} \int_S f^2 \\ &= \left(\frac{4}{r^2} \left(1 - \frac{3m}{2r} \right) + O(r^{-2}\epsilon) \right) \int_S fAf + \frac{36m^2}{r^6} \int_S f^2. \end{aligned}$$

Note that for $r > 2m$ we have,

$$1 - \frac{3m}{2r} > \frac{1}{4}.$$

We deduce, for sufficiently small ϵ , everywhere in $(ext)\mathcal{M}$,

$$\int_S f \left(A^2 - \frac{12m}{r^3}A + \frac{36m^2}{r^6} \right) > \frac{1}{r^2} \left(\int_S fAf + \frac{36m^2}{r^4} \int_S f^2 \right).$$

Now, since $\left| \bar{\rho} + \frac{2m}{r^3} \right| \lesssim \epsilon_0 r^{-3}$

$$\int_S fAf = \int_S f(\not{d}_2^* \not{d}_2 - 3\bar{\rho})f = \int_S \left(|\not{d}_2f|^2 + \left(\frac{6m}{r^3} + O(r^{-3}\epsilon_0) \right) |f|^2 \right).$$

Hence,

$$\int_S f Af + \frac{36m^2}{r^4} \int_S f^2 > \int_S \left(|\not\phi_2 f|^2 + \left(\frac{6m}{r^3} + \frac{36m^2}{r^6} \right) |f|^2 \right) > \int_S |\not\phi_2 f|^2$$

or, since $\int_S (\not\phi_2 f)^2 \geq 4 \int_S \frac{1}{r^2} \int_S |f|^2 + O(\epsilon_0 r^{-3}) \int_S |f|^2$. We deduce,

$$\int_S f \left(A^2 - \frac{12m}{r^3} A + \frac{36m^2}{r^6} \right) \geq \frac{1}{4r^2} \int_S (\not\phi_2 f)^2 + \frac{9}{r^4} \int_S f^2$$

as desired. This concludes the proof of the lemma. \square

Applying the lemma to $f = \not\phi_2^* e_{\theta \underline{\kappa}}$ in (7.5.17), i.e.

$$A^2 \not\phi_2^* (e_{\theta \underline{\kappa}}) - \frac{12m}{r^3} A \not\phi_2^* e_{\theta}(\underline{\kappa}) + \frac{36m^2}{r^6} \not\phi_2^* e_{\theta}(\underline{\kappa}) \in r^{-7} \text{Good}_8(\underline{\alpha})$$

or, in any region where

$$\|\underline{\alpha}\|_{2,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 16,$$

we have

$$\left\| A^2 \not\phi_2^* (e_{\theta \underline{\kappa}}) - \frac{12m}{r^3} A \not\phi_2^* e_{\theta}(\underline{\kappa}) + \frac{36m^2}{r^6} \not\phi_2^* e_{\theta}(\underline{\kappa}) \right\|_{2,k} \lesssim \epsilon_0 r^{-6} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 8.$$

We deduce, by L^2 -elliptic estimates,

$$\|\not\phi_2^* e_{\theta \check{\kappa}}\|_{2,k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 12. \quad (7.5.20)$$

Since we control the $\ell = 1$ mode of $e_{\theta \check{\kappa}}$ we infer that,

$$\|e_{\theta \check{\kappa}}\|_{2,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 13$$

i.e.,

$$\|\check{\kappa}\|_{2,k} \lesssim \epsilon_0 u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 14.$$

Therefore, using the Sobolev embedding,

$$\|\check{\kappa}\|_{\infty,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}} \quad k \leq k_{small} + 12.$$

This proves the following,

Proposition 7.5.8. *In any region of $^{(ext)}\mathcal{M}$ where,*

$$\|\underline{\alpha}\|_{2,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 16,$$

we also have,

$$\|\check{\kappa}\|_{\infty,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}} \quad k \leq k_{small} + 12. \quad (7.5.21)$$

7.5.3 Proof of Proposition 7.5.1, Part I

We first prove Proposition 7.5.1 in the region where the estimate

$$\|\underline{\alpha}\|_{2,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 16, \quad (7.5.22)$$

holds true.

Step 1. We prove the estimates,

$$\begin{aligned} \|\zeta\|_{\infty,k} &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 15, \\ \|\beta\|_{\infty,k} &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 12. \end{aligned} \quad (7.5.23)$$

According to (7.5.18)

$$\begin{aligned} A^2 B \phi_2^* \zeta &= \frac{3}{4} \bar{\kappa} \bar{\rho} A \phi_2^* e_\theta(\underline{\kappa}) + r^{-8} \text{Good}_{12}(\underline{\alpha}), \\ A^2 B \phi_2^* \beta &= -\frac{9}{8} (\bar{\kappa} \bar{\rho})^2 \phi_2^* e_\theta(\underline{\kappa}) + r^{-9} \text{Good}_9(\underline{\alpha}). \end{aligned}$$

In view of (7.5.20) we deduce, in L^2 norms,

$$\begin{aligned} \|A^2 B \phi_2^* \zeta\|_{2,k} &\lesssim r^{-4} \|A \phi_2^* e_\theta \kappa\|_{2,k} + \epsilon_0 r^{-7} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 12, \\ \|A^2 B \phi_2^* \beta\|_{2,k} &\lesssim r^{-8} \|\phi_2^* e_\theta \kappa\|_{2,k} + \epsilon_0 r^{-8} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 9. \end{aligned}$$

Thus, in view of the estimates for $\check{\kappa}$ derived above,

$$\begin{aligned} \|A^2 B \phi_2^* \zeta\|_{2,k} &\lesssim \epsilon_0 r^{-7} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 12, \\ \|A^2 B \phi_2^* \beta\|_{2,k} &\lesssim \epsilon_0 r^{-8} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 9. \end{aligned}$$

Thus, by elliptic estimates,

$$\begin{aligned} \|\phi_2^* \zeta\|_{2,k} &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 16, \\ \|\phi_2^* \beta\|_{2,k} &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 13. \end{aligned}$$

In view of the estimates for the $\ell = 1$ modes of ζ, β we deduce,

$$\begin{aligned} \|\zeta\|_{2,k} &\lesssim \epsilon_0 u^{-1-\delta_{dec}}, & k &\leq k_{small} + 17, \\ \|\beta\|_{2,k} &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 14. \end{aligned}$$

Passing to L^∞ norms we derive

$$\begin{aligned} \|\zeta\|_{\infty,k} &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 15, \\ \|\beta\|_{\infty,k} &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, & k &\leq k_{small} + 13. \end{aligned} \quad (7.5.24)$$

Step 2. We prove the estimate

$$\|\eta\|_{\infty,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 15.$$

This follows immediately from the estimate from ζ and the previously derived estimate (7.5.2). Indeed,

$$\|\eta\|_{\infty,k} \lesssim \|\zeta\|_{\infty,k} + \epsilon_0 r^{-1} u^{-1-\delta_{dec}} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}.$$

Step 3. We derive the estimate,

$$\|\vartheta\|_{\infty,k} \lesssim r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 11. \quad (7.5.25)$$

This follows easily in view of the equation (see (7.5.12))

$$\not\partial_2 \vartheta + 2\beta - \bar{\kappa} \zeta \in r^{-3} \text{Good}_{20}$$

from which, in view of Step 1,

$$\|\not\partial_2 \vartheta\|_{2,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 12.$$

The desired estimate follows by elliptic estimates and Sobolev.

Step 4. We derive the intermediate estimate for $\underline{\vartheta}$,

$$\|\underline{\vartheta}\|_{\infty,k} \lesssim \epsilon_0 u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 12. \quad (7.5.26)$$

To show this we combine the equations (see (7.5.12))

$$\begin{aligned} \not\partial_2 \underline{\vartheta} + 2\underline{\beta} - e_\theta(\underline{\kappa}) + \bar{\kappa} \zeta &\in r^{-3} \text{Good}_{20}, \\ \not\partial_2^* \underline{\beta} + \frac{3}{2} \bar{\rho} \underline{\vartheta} &\in r^{-2} \text{Good}_{15}, \end{aligned}$$

to deduce,

$$\not\partial_2^* \not\partial_2 \underline{\vartheta} - 3\bar{\rho} \underline{\vartheta} = \not\partial_2^* e_\theta \underline{\kappa} + \bar{\kappa} \not\partial_2^* \zeta + r^{-2} \text{Good}_{15},$$

and hence,

$$\|A\underline{\vartheta}\|_{2,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 12.$$

Thus,

$$\|\underline{\vartheta}\|_{2,k} \lesssim \epsilon_0 r u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 14$$

and hence,

$$\|\underline{\vartheta}\|_{\infty,k} \lesssim \epsilon_0 u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 12$$

as desired.

Step 5. We derive the estimate,

$$\|\check{\rho}\|_{\infty,k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 14. \quad (7.5.27)$$

From,

$$\phi_2^* \phi_1^* \rho + \frac{3}{4} \bar{\kappa} \bar{\rho} \vartheta + \frac{3}{4} \bar{\kappa} \bar{\rho} \underline{\vartheta} \in r^{-4} \text{Good}_{20},$$

we deduce,

$$\begin{aligned} \|\phi_2^* \phi_1^* \rho\|_{2,k} &\lesssim r^{-4} (\|\theta\|_{2,k} + \|\underline{\theta}\|_{2,k}) + \epsilon_0 r^{-3} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 14 \\ &\lesssim \epsilon_0 r^{-3} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 14. \end{aligned}$$

Since we control the $\ell = 1$ mode of $\phi_1^* \rho$ (see Lemma 7.4.7) we infer that,

$$\|\check{\rho}\|_{2,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 16$$

i.e.,

$$\|\check{\rho}\|_{\infty,k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 14$$

as desired.

Step 6. We derive the estimate,

$$\|\underline{\beta}\|_{\infty,k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad \forall k \leq k_{small} + 9 \quad (7.5.28)$$

with the help of the identity

$$e_3(r\mathbf{q}) = r^5 \left\{ \phi_2^* \phi_1^* \phi_1 \underline{\beta} - \frac{3}{2} \kappa \rho \underline{\alpha} - \frac{3}{2} \rho \phi_2^* \phi_1^* \underline{\kappa} - \frac{3}{2} \underline{\kappa} \rho \phi_2^* \zeta + \frac{3}{4} (2\rho^2 - \kappa \underline{\kappa} \rho) \underline{\vartheta} \right\} + \text{Err}[e_3(r\mathbf{q})],$$

$$\text{Err}[e_3(r\mathbf{q})] = r^4 (e_3 \Gamma_b) \cdot \check{\rho}^{\leq 1} \beta + r \Gamma_b \cdot \mathbf{q} + r^2 \check{\rho}^3 (\Gamma_g \cdot \Gamma_b),$$

of Proposition 7.1.11. In view of (7.3.26) we have,

$$\|\text{Err}[e_3(r\mathbf{q})]\|_{\infty,k}(u, r) \lesssim \epsilon_0 u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 16.$$

We can now make use of the estimates for $\underline{\kappa}, \zeta, \vartheta, \underline{\vartheta}$ already derived and the **Ref 2** estimate for $e_3(\mathbf{q})$ and $\underline{\alpha}$ to deduce, for all $k \leq k_{small} + 10$,

$$\begin{aligned} \|\rho \phi_2^* \phi_1^* \underline{\kappa}\|_{\infty, k} &\lesssim \epsilon_0 r^{-6} u^{-1-\delta_{dec}}, \\ \|\underline{\kappa} \rho \phi_2^* \zeta\|_{\infty, k} &\lesssim \epsilon_0 r^{-6} u^{-1-\delta_{dec}}, \\ \|\rho^2 \underline{\vartheta}\|_{\infty, k} &\lesssim \epsilon_0 r^{-6} u^{-1-\delta_{dec}}, \\ \|\kappa \underline{\kappa} \underline{\vartheta}\|_{\infty, k} &\lesssim \epsilon_0 r^{-5} u^{-1-\delta_{dec}}, \\ \|\kappa \rho \underline{\alpha}\|_{\infty, k} &\lesssim \epsilon_0 r^{-5} u^{-1-\delta_{dec}}, \\ \|e_3(r\mathbf{q})\|_{\infty, k} &\lesssim \epsilon_0 u^{-1-\delta_{dec}}. \end{aligned}$$

Therefore,

$$\|\phi_2^* \phi_1^* \phi_1 \underline{\beta}\|_{\infty, k} \lesssim \epsilon_0 r^{-5} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 10,$$

i.e.,

$$\|\phi_2^* \phi_1^* \phi_1 \underline{\beta}\|_{2, k} \lesssim \epsilon_0 r^{-4} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 10.$$

Making use of the identity,

$$\phi_1^* \phi_1 = \phi_2 \phi_2^* + 2K,$$

we deduce

$$\|(\phi_2^* \phi_2 + K) \phi_2^* \underline{\beta}\|_{2, k} \lesssim \epsilon_0 r^{-4} u^{-1-\delta_{dec}}.$$

Since $\phi_2^* \phi_2 + K$ is coercive we deduce,

$$\|\phi_2^* \underline{\beta}\|_{2, k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad \forall k \leq k_{small} + 10.$$

Since we control the $\ell = 1$ mode of $\underline{\beta}$ (see Lemma 7.4.7) according to Lemma 7.3.8,

$$\|\underline{\beta}\|_{2, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad \forall k \leq k_{small} + 11.$$

Hence,

$$\|\underline{\beta}\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad \forall k \leq k_{small} + 9. \quad (7.5.29)$$

Step 7. Using the above estimate for $\underline{\beta}$ we can improve the estimate for $\underline{\vartheta}$ derived in Step 4. We show, in the region where the estimate (7.5.22) for $\underline{\alpha}$ holds,

$$\|\underline{\vartheta}\|_{\infty, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 9. \quad (7.5.30)$$

Indeed in view of the Codazzi equation

$$\not\partial_2 \underline{\vartheta} + 2\underline{\beta} - e_\theta(\underline{\kappa}) + \bar{\kappa}\zeta \in r^{-3}Good_{20},$$

we infer that, for all $k \leq k_{small} + 11$,

$$\begin{aligned} \|\not\partial_2 \underline{\vartheta}\|_{2,k} &\lesssim \|\underline{\beta}\|_{2,k} + r^{-1}\|\check{\kappa}\|_{2,k+1} + r^{-1}\|\zeta\|_{2,k} + \epsilon_0 r^{-2} u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}. \end{aligned}$$

Thus, for all $k \leq k_{small} + 12$

$$\|\underline{\vartheta}\|_{2,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}$$

and hence,

$$\|\underline{\vartheta}\|_{\infty,k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 10. \quad (7.5.31)$$

This ends the proof of Proposition 7.5.1 in the region for which the desired estimate (7.5.22) for $\underline{\alpha}$ holds true.

Since (7.5.22) for $\underline{\alpha}$ holds true on \mathcal{T} in view of²⁴ Theorem M3, this ends the proof of Proposition 7.5.1 on \mathcal{T} .

7.5.4 Proof of Proposition 7.5.1, Part II

We extend the validity of Proposition 7.5.1 to all of $(ext)\mathcal{M}$ propagating the estimates derived in the first part on \mathcal{T} . We also recall that we have good decay estimates for $\check{\kappa}$ and $\check{\mu}$ everywhere on $(ext)\mathcal{M}$.

Step 1. We first derive estimates for ϑ in \mathcal{M}_{ext} making use of the transport equation

$$e_4(\vartheta) + \bar{\kappa}\vartheta = -2\alpha - (\kappa - \bar{\kappa})\vartheta = -2\alpha + \Gamma_g \cdot \Gamma_g.$$

Making use of Proposition 7.4.3 we derive, for all $r \geq r_0 = r_{\mathcal{T}}$,

$$r^2 \|\vartheta\|_{\infty,k}(u, r) \lesssim r_0^2 \|\vartheta\|_{\infty,k}(u, r_0) + \int_{r_0}^r \lambda^2 \|\alpha\|_{\infty,k}(u, \lambda) d\lambda + \epsilon_0 u^{-1-\delta_{dec}}.$$

We now make use of the estimate,

$$\|\alpha\|_{\infty,k}(u, r) \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 20$$

²⁴Recall that r is bounded on \mathcal{T} and that $\mathcal{T} \subset (int)\mathcal{M}$ so that (7.5.22) holds true for $(int)\underline{\alpha}$ on \mathcal{T} in view of Theorem M3. Then, since we have $(ext)\underline{\alpha} = ((ext)\Upsilon)^2 (int)\underline{\alpha}$ on \mathcal{T} , (7.5.22) holds indeed true for $(ext)\underline{\alpha}$ on \mathcal{T} .

and,

$$\|\vartheta\|_{\infty,k}(u, r_0) \lesssim \epsilon_0 u^{-1-\delta_{dec}}$$

derived above in (7.5.25), to derive

$$r^2 \|\vartheta\|_{\infty,k}(u, r) \lesssim \epsilon_0 u^{-1-\delta_{dec}} + \epsilon_0 r u^{-1-\delta_{dec}}.$$

Therefore, everywhere on $^{(ext)}\mathcal{M}$,

$$\|\vartheta\|_{\infty,k}(u, r) \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}. \quad (7.5.32)$$

Step 2. Next, we estimate β from the equation,

$$e_4 \beta + 2\bar{\kappa} \beta = \not{d}_2 \alpha - (\kappa - \bar{\kappa}) \beta + \check{\Gamma}_g \cdot \alpha = \not{d}_2 \alpha + \Gamma_g \cdot (\alpha, \beta)$$

to deduce in the same manner

$$r^4 \|\beta\|_{\infty,k}(u, r) \lesssim r_0^2 \|\beta\|_{\infty,k}(u, r_0) + \int_{r_0}^r \lambda^4 \|\not{d}_2 \alpha\|_{\infty,k}(u, \lambda) d\lambda + \epsilon_0 r u^{-1-\delta_{dec}}.$$

Thus, in view of the estimates for α in (7.5.24) and the estimates for α in **Ref2**, i.e., for $0 \leq k \leq k_{small} + 20$,

$$\|\alpha\|_{\infty,k} \lesssim \epsilon_0 \min\{r^{-2} \log(1+u)(u+2r)^{-1-\delta_{extra}}, r^{-3}(u+2r)^{-\frac{1}{2}-\delta_{extra}}\}.$$

Thus we have with $I(u, r) := \int_{r_0}^r \lambda^4 \|\not{d}_2 \alpha\|_{\infty,k}(u, \lambda) d\lambda$

$$I(u, r) \lesssim \epsilon_0 \min\left\{\log(1+u) \int_{r_0}^r \lambda(u+2\lambda)^{-1-\delta_{extra}} d\lambda, \int_{r_0}^r (u+2\lambda)^{-1/2-\delta_{extra}} d\lambda\right\}.$$

If $r \leq 2u$ we have,

$$\int_{r_0}^r \lambda(u+2\lambda)^{-1-\delta_{extra}} d\lambda \lesssim r^2 u^{-1-\delta_{extra}} \lesssim r^2 (u+2r)^{-1-\delta_{extra}}$$

and

$$r^{-4} I(u, r) \lesssim \epsilon_0 r^{-2} \log(1+u)(u+2r)^{-1-\delta_{extra}}.$$

If $r \geq 2u$ we have,

$$\int_{r_0}^r (u+2\lambda)^{-1/2-\delta_{extra}} d\lambda \lesssim (u+2r)^{1/2+\delta_{extra}}$$

and

$$r^{-4}I(u, r) \lesssim r^{-4}(u+2r)^{1/2+\delta_{extra}} \lesssim r^{-2}(u+2r)^{-1-\delta_{extra}}.$$

We deduce,

$$\|\beta\|_{\infty, k} \lesssim r^{-4}\|\beta\|_{\infty, k}(u, r_0) + \epsilon_0 r^{-2} \log(1+u)(u+2r)^{-1-\delta_{extra}}.$$

Thus in view of (7.5.24),

$$\|\beta\|_{\infty, k} \lesssim \epsilon_0 r^{-2} \log(1+u)(u+2r)^{-1-\delta_{extra}}. \quad (7.5.33)$$

Step 3. We now estimate ζ using the equation

$$e_4(\zeta) + \kappa\zeta = -\beta + \check{\Gamma}_g \cdot \check{\Gamma}_g.$$

This can be done exactly as in Step 1 making use of the estimates already derived for β and the estimate (7.5.24) for ζ along \mathcal{T} . We thus derive,

$$\|\zeta\|_{\infty, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small+15}.$$

Step 4. We estimate $\check{\rho}$ using equation

$$\check{\rho} = -\check{\phi}_1 \zeta - \check{\mu} + \check{\Gamma}_g \cdot \check{\Gamma}_b,$$

the previous estimate for ζ and $\check{\mu}$ in $(ext)\mathcal{M}$. We deduce,

$$\|\check{\rho}\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 14. \quad (7.5.34)$$

Step 5. We estimate $\check{\kappa}$ using the equation,

$$e_4 \check{\kappa} + \frac{1}{2} \bar{\kappa} \check{\kappa} + \frac{1}{2} \check{\kappa} \bar{\kappa} = -2\check{\phi}_1 \zeta + 2\check{\rho} + \check{\Gamma}_g \cdot \check{\Gamma}_b.$$

Making use of the estimates in $(ext)\mathcal{M}$ for $\check{\kappa}$, ζ and $\check{\rho}$ as well as the estimates for $\check{\kappa}$ on \mathcal{T} in Proposition 7.5.8 we derive, everywhere on $(ext)\mathcal{M}$,

$$\|\check{\kappa}\|_{\infty, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 12. \quad (7.5.35)$$

Alternatively we can make use of the estimate for $\Xi = r^2(e_\theta(\check{\kappa}) + 4r\check{\phi}_1^* \check{\phi}_1 \zeta - 2r^2 \check{\phi}_1^* \check{\phi}_1 \beta)$ in Lemma 7.5.2, which holds everywhere on $(ext)\mathcal{M}$, and the above estimates for ζ, β .

Step 6. We estimate $\underline{\beta}$ everywhere on $^{(ext)}\mathcal{M}$ with the help of the equation

$$e_4\underline{\beta} + \bar{\kappa}\underline{\beta} = -\underline{\not{d}}_1\rho - 3\underline{\zeta}\rho - \underline{\vartheta}\beta - (\kappa - \bar{\kappa})\underline{\beta}$$

together with the estimate (7.5.29) for $\underline{\beta}$ on \mathcal{T} and the above derived estimates for $\check{\rho}, \zeta$ in $^{(ext)}\mathcal{M}$ to infer that,

$$\|\underline{\beta}\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad \forall k \leq k_{small} + 9. \quad (7.5.36)$$

Step 7. We extend the for $\underline{\vartheta}$ everywhere on $^{(ext)}\mathcal{M}$ by making use of the Codazzi equation for $\underline{\vartheta}$ in (7.5.12),

$$\underline{\not{d}}_2\underline{\vartheta} + 2\underline{\beta} - e_\theta(\underline{\kappa}) + \bar{\kappa}\underline{\zeta} \in r^{-3}Good_{20}.$$

Using the estimates already derived above, we infer that, for all $k \leq k_{small} + 11$,

$$\begin{aligned} \|\underline{\not{d}}_2\underline{\vartheta}\|_{2, k} &\lesssim \|\underline{\beta}\|_{2, k} + r^{-1}\|\underline{\check{\kappa}}\|_{2, k+1} + r^{-1}\|\underline{\zeta}\|_{2, k} + \epsilon_0 r^{-2} u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}. \end{aligned}$$

Hence, everywhere in $^{(ext)}\mathcal{M}$,

$$\|\underline{\vartheta}\|_{2, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad \text{for all } k \leq k_{small} + 12,$$

and therefore,

$$\|\underline{\vartheta}\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad \text{for all } k \leq k_{small} + 10.$$

Step 7. We estimate $\underline{\alpha}$ everywhere on $^{(ext)}\mathcal{M}$ by making use of the equation

$$e_4\underline{\alpha} + \frac{1}{2}\bar{\kappa}\underline{\alpha} = -\underline{\not{d}}_2^*\underline{\beta} - \frac{3}{2}\underline{\vartheta}\rho - 5\underline{\zeta}\underline{\beta} - \frac{1}{2}(\kappa - \bar{\kappa})\underline{\alpha}$$

as well as the estimate (7.5.22) for $\underline{\alpha}$ on \mathcal{T} and the above estimates in all $^{(ext)}\mathcal{M}$ for $\underline{\beta}$ and $\underline{\vartheta}$. Proceeding as before we derive,

$$\|\underline{\alpha}\|_{\infty, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}} \text{ for all } k \leq k_{small} + 8. \quad (7.5.37)$$

This concludes the proof of Proposition 7.5.1.

7.6 Conclusion of the Proof of Theorem M4

So far we have established the following estimates, for all $k \leq k_{small} + 8$

$$\begin{aligned}
& \|\check{\kappa}, r\check{\mu}\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \\
& \|\vartheta, \zeta, \check{\kappa}, r\check{\rho}\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1/2-\delta_{dec}}, \\
& \|\vartheta, \zeta, \eta, \check{\kappa}, \underline{\vartheta}, r\beta, r\check{\rho}, r\underline{\beta}, \underline{\alpha}\|_{\infty, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \\
& \|\beta, re_3\beta\|_{\infty, k} \lesssim \epsilon_0 r^{-3} (2r + u)^{-1/2-\delta_{dec}}.
\end{aligned} \tag{7.6.1}$$

It only remains to derive improved decay estimates for $e_3(\beta, \vartheta, \zeta, \check{\kappa}, \check{\rho})$ and the estimates for $\underline{\xi}, \underline{\omega}, \check{\zeta}, \underline{\check{\Omega}}$ as well as $\bar{\varsigma} + 1$ and $\bar{\underline{\Omega}} + \Upsilon$ in terms of $u^{-1-\delta_{dec}}$ decay. More precisely it remains to prove the following.

Proposition 7.6.1. *The following estimates hold true on ${}^{(ext)}\mathcal{M}$ for all $k \leq k_{small} + 7$.*

$$\begin{aligned}
& \|e_3(\vartheta, \zeta, \check{\kappa}), re_3\beta, re_3\check{\rho}\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \\
& \|\underline{\xi}, \underline{\omega}\|_{\infty, k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \\
& \|\check{\zeta}, \underline{\check{\Omega}}, \bar{\varsigma} + 1, \bar{\underline{\Omega}} + \Upsilon\|_{\infty, k} \lesssim \epsilon_0 u^{-1-\delta_{dec}}.
\end{aligned}$$

Proof. We proceed in steps as follows.

Step 1. We make use of the equation $e_3\vartheta = -\frac{1}{2}\underline{\kappa}\vartheta + 2\underline{\omega}\vartheta - 2\check{\rho}\eta - \frac{1}{2}\kappa\underline{\vartheta} + 2\eta^2$ and the previously derived estimates to derive,

$$\|e_3\vartheta\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 9. \tag{7.6.2}$$

Step 2. We make use of the equation $e_3\beta + (\underline{\kappa} - 2\underline{\omega})\beta = -\check{\rho}\alpha + 3\eta\rho + \Gamma_g\underline{\beta} + \Gamma_b\alpha$ and the previously derived estimates for $\beta, \check{\rho}, \underline{\beta}$ to derive,

$$\|e_3\beta\|_{\infty, k} \lesssim \epsilon_0 r^{-3} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 9.$$

Step 3. To estimate $e_3\zeta$ in the next step we actually need a stronger estimate for $e_3\beta$ than the one derived above. At the same time we derive an improved estimate for β . We show in fact, for some $0 < \delta$,

$$\begin{aligned}
& \|\beta\|_{\infty, k} \lesssim \epsilon_0 r^{-2-\delta} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 10, \\
& \|e_3\beta\|_{\infty, k-1} \lesssim \epsilon_0 r^{-3-\delta} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 10.
\end{aligned} \tag{7.6.3}$$

This makes use of the equation

$$e_4\beta + 2\kappa\beta = \not{d}_2\alpha + \Gamma_g \cdot \alpha = F := \not{d}_2\alpha + \Gamma_g \cdot \alpha - 2\check{\kappa}\beta$$

and the estimates for α in **Ref 2**. Thus, for some $0 < \delta < \delta_{extra} - \delta_{dec}$,

$$\begin{aligned} \|F\|_{\infty,k} &\lesssim \log(1+u)r^{-3}(2r+u)^{-1-\delta_{extra}} + \epsilon_0 r^{-4} u^{-1-\delta_{dec}} \\ &\lesssim \epsilon_0 u^{-1-\delta_{dec}} r^{-3-\delta}. \end{aligned}$$

Integrating from \mathcal{T} , where $r = r_{\mathcal{T}} = r_0 \lesssim 1$, we deduce with the help of Proposition 7.4.3

$$\begin{aligned} r^4 \|\beta\|_{\infty,k}(u,r) &\lesssim r_0^4 \|\beta\|_{\infty,k}(u,r_0) + \int_{r_0}^r \lambda^4 \|F\|_{\infty,k}(u,\lambda) d\lambda \\ &\lesssim \|\beta\|_{\infty,k}(u,r_0) + \epsilon_0 \int_{r_0}^r \lambda^{1-\delta} d\lambda. \end{aligned}$$

Based on the previously derived estimate for β we have $\|\beta\|_{\infty,k}(u,r_{\mathcal{H}}) \lesssim \epsilon_0 u^{-1-\delta_{dec}}$. Hence,

$$\|\beta\|_{\infty,k}(u,r) \lesssim \epsilon_0 r^{-4} u^{-1-\delta_{dec}} + \epsilon_0 r^{-4} r^{2-\delta} u^{-1-\delta_{dec}} \lesssim \epsilon_0 r^{-2-\delta} u^{-1-\delta_{dec}}$$

as desired.

To prove the second estimate in (7.6.3) we commute the transport equation for β with \mathbf{T} and make use of the corresponding estimate for $\mathbf{T}\alpha$ (which follows from **Ref 2**).

$$\|\mathbf{T}\alpha\|_{\infty,k} \lesssim \epsilon_0 \log(1+u)r^{-4}(2r+u)^{-1-\delta_{extra}} \lesssim \epsilon_0 u^{-1-\delta_{dec}} r^{-4-\delta}$$

as well as the fact that we control $\mathbf{T}\beta$ on \mathcal{T} , i.e. $\|\mathbf{T}\beta\|_{\infty,k-1}(u,r_0) \lesssim \epsilon_0 u^{-1-\delta_{dec}}$.

Step 4. We make use of the equation $e_4\zeta + \bar{\kappa}\zeta = -\beta + \Gamma_g \cdot \Gamma_g$ to derive,

$$\|e_3\zeta\|_{k,\infty} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 9. \quad (7.6.4)$$

Indeed commuting the equation with \mathbf{T} we derive,

$$e_4\mathbf{T}\zeta + \bar{\kappa}\mathbf{T}\zeta = F := -\mathbf{T}\beta + [\mathbf{T}, e_4]\zeta + \zeta\mathbf{T}\bar{\kappa} + \mathbf{T}(\Gamma_g \cdot \Gamma_g).$$

It is easy to check, in view of the commutation Lemma 7.4.1,

$$\|F\|_{\infty,k-1} \lesssim \|\mathbf{T}\beta\|_{\infty,k-1} + \epsilon_0 r^{-4} u^{-1-\delta_{dec}}.$$

Thus, in view of the estimate for $e_3\zeta$ derived in Step 3 and the estimate for $e_4\zeta$ we infer that,

$$\|F\|_{\infty,k-1} \lesssim \epsilon_0 r^{-3-\delta} u^{-1-\delta_{dec}}.$$

Integrating from \mathcal{T} and using the previously derived estimate $\|\zeta\|_{k,\infty} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}$

$$\begin{aligned} r^2 \|\mathbf{T}\zeta\|_{\infty,k-1} &\lesssim r_0^2 \|\mathbf{T}\zeta\|_{\infty,k-1}(u, r_0) + \epsilon_0 u^{-1-\delta_{dec}} \int_{r_0}^r \lambda^{-1-\delta} d\lambda \\ &\lesssim \|\mathbf{T}\zeta\|_{\infty,k-1}(u, r_0) + \epsilon_0 r^{-\delta} u^{-1-\delta_{dec}} \lesssim \epsilon_0 u^{-1-\delta_{dec}}. \end{aligned}$$

Hence

$$\|\mathbf{T}\zeta\|_{\infty,k-1} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}$$

from which the desired estimate easily follows.

Step 5. We make use of the equation $e_4(\underline{\tilde{\omega}}) = \check{\rho} + \Gamma_g \cdot \Gamma_b$ and the previously derived estimates for $\check{\rho}$ as well as the estimates of $\underline{\tilde{\omega}}$ on the last slice (see Proposition 7.3.9) to derive the estimate

$$\|\underline{\tilde{\omega}}\|_{\infty,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 9. \quad (7.6.5)$$

Indeed,

$$\|e_4 \underline{\tilde{\omega}}\|_{\infty,k} \lesssim \|\check{\rho}\|_{\infty,k} + \epsilon_0 r^{-3} u^{-1-\delta_{dec}} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}.$$

Thus, applying Proposition 7.4.3, integrating from Σ_* and using the previously derived estimate for $\underline{\tilde{\omega}}$ on Σ_* ,

$$\begin{aligned} \|\underline{\tilde{\omega}}\|_{\infty,k}(u, r) &\lesssim \|\underline{\tilde{\omega}}\|_{\infty,k}(u, r_*) + \epsilon_0 u^{-1-\delta_{dec}} \int_r^{r_*} \lambda^{-2} d\lambda \\ &\lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}} \end{aligned}$$

as desired.

Step 6. We derive the estimate,

$$\|\underline{\xi}\|_{\infty,k} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 9 \quad (7.6.6)$$

by making use of the transport equation $e_4(\underline{\xi}) = F := -e_3(\zeta) + \underline{\beta} - \frac{1}{2}\underline{\kappa}(\zeta + \eta) + \Gamma_b \cdot \Gamma_b$. In view of the previously derived estimates for $e_3\zeta, \underline{\beta}, \zeta, \eta$ we derive,

$$\|F\|_{\infty,k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}.$$

Integrating from Σ_* and making use of the estimate for $\underline{\xi}$ on Σ_* (see Proposition 7.3.9) we derive,

$$\|\underline{\xi}\|_{\infty,k}(u, r) \lesssim \|\underline{\xi}\|_{\infty,k}(u, r_*) + \epsilon_0 r^{-1} u^{-1-\delta_{dec}} \lesssim \epsilon_0 r^{-1} u^{-1-\delta_{dec}}.$$

Step 7. We derive the estimate

$$\|\check{\underline{\Omega}}\|_{\infty, k} \lesssim \epsilon_0 u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 8. \quad (7.6.7)$$

This follows immediately from the the equation $e_\theta(\underline{\Omega}) = -\underline{\xi} - (\eta - \zeta)\underline{\Omega}$, see (2.2.19), and the previous estimate for $\underline{\xi}$. Note that $\underline{\Omega}$ has been estimated in Lemma 3.4.1.

Step 8. We derive the estimate

$$\|\varsigma - 1\|_{\infty, k} \lesssim \epsilon_0 u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 8. \quad (7.6.8)$$

The estimate follows from the propagation equation $e_4(\varsigma) = 0$ and the estimate for $\varsigma - 1$ on the last slice Σ_* .

Step 9. We derive the estimate,

$$\|e_3\check{\rho}\|_{\infty, k} \lesssim \epsilon_0 r^{-3} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 8 \quad (7.6.9)$$

with the help of the equation (see Proposition 7.1.8)

$$e_3\check{\rho} = r^{-2} \check{\not{D}}^{\leq 1} \Gamma_b + r^{-1} \Gamma_b \cdot \Gamma_b$$

and the previously derived estimates for $\underline{\beta}$, $\underline{\check{\kappa}}$, $\check{\rho}$, $\check{\underline{\Omega}}$, $\check{\zeta}$.

Step 10. We derive the estimate,

$$\|e_3\check{\underline{\kappa}}\|_{\infty, k} \lesssim \epsilon_0 r^{-2} u^{-1-\delta_{dec}}, \quad k \leq k_{small} + 8 \quad (7.6.10)$$

using the equation (see Proposition 7.1.8)

$$e_3\check{\underline{\kappa}} = r^{-1} \check{\not{D}}^{\leq 1} \Gamma_b + \Gamma_b \cdot \Gamma_b$$

and the previously derived estimates for $\underline{\check{\kappa}}$, $\underline{\xi}$, $\underline{\check{\omega}}$, $\check{\underline{\Omega}}$, $\check{\zeta}$. This ends the proof of Proposition 7.6.1 and Theorem M4. \square

7.7 Proof of Theorem M5

Recall from Theorem M3 that we have obtained the following estimate for ${}^{(int)}\underline{\alpha}$ in ${}^{(int)}\mathcal{M}$

$$\max_{0 \leq k \leq k_{small} + 16} \sup_{(int)\mathcal{M}} \underline{u}^{1+\delta_{dec}} |\not{D}^k \underline{\alpha}| \lesssim \epsilon_0. \quad (7.7.1)$$

Step 1. We consider the control of the other curvature components, as well as the Ricci components on \mathcal{T} . Recall that the $(\underline{u}, {}^{(int)}s)$ foliation is initialized on \mathcal{T} as follows

- \underline{u} and ${}^{(int)}s$ are defined on \mathcal{T} by

$$\underline{u} = u \text{ and } {}^{(int)}s = {}^{(ext)}s \text{ on } \mathcal{T}.$$

In particular, the 2-spheres $\mathbf{S}(u, {}^{(int)}s)$ coincide on \mathcal{T} with the 2-sphere $\mathbf{S}(u, {}^{(ext)}s)$.

- In view of the above initialization, and the fact that $\mathcal{T} = \{r = r_{\mathcal{T}}\}$, we infer that

$${}^{(int)}r = {}^{(ext)}r = r_{\mathcal{T}}, \quad {}^{(int)}m = {}^{(ext)}m.$$

- The null frame $({}^{(int)}e_3, {}^{(int)}e_4, {}^{(int)}e_\theta)$ is defined on \mathcal{T} by

$${}^{(int)}e_4 = {}^{(ext)}\lambda {}^{(ext)}e_4, \quad {}^{(int)}e_3 = ({}^{(ext)}\lambda)^{-1} {}^{(ext)}e_3, \quad {}^{(int)}e_\theta = {}^{(ext)}e_\theta \text{ on } \mathcal{T}$$

where

$${}^{(ext)}\lambda = 1 - \frac{2 {}^{(ext)}m}{{}^{(ext)}r}.$$

In particular, we deduce the following identities for the curvature components and Ricci coefficients on \mathcal{T} .

Lemma 7.7.1. *We have on \mathcal{T}*

$$\begin{aligned} {}^{(int)}\underline{\zeta} &= -\frac{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}}{{}^{(ext)}\underline{\kappa}} \lambda^{-1} {}^{(ext)}\underline{\zeta}, \\ {}^{(int)}\underline{\Omega} &= \lambda - \lambda^2 \frac{\overline{{}^{(ext)}\kappa}}{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}} - \lambda \frac{\overline{{}^{(ext)}\kappa}}{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}} {}^{(ext)}\underline{\Omega}. \end{aligned}$$

where

$$\lambda = {}^{(ext)}\lambda = 1 - \frac{2 {}^{(ext)}m}{{}^{(ext)}r}.$$

Moreover, we have on \mathcal{T}

$${}^{(int)}\alpha = \lambda^2 {}^{(ext)}\alpha, \quad {}^{(int)}\beta = \lambda {}^{(ext)}\beta, \quad {}^{(int)}\rho = {}^{(ext)}\rho, \quad {}^{(int)}\underline{\beta} = \lambda^{-1} {}^{(ext)}\underline{\beta}, \quad {}^{(int)}\underline{\alpha} = \lambda^{-2} {}^{(ext)}\underline{\alpha},$$

$$\begin{aligned} {}^{(int)}\underline{\xi} &= 0, \quad {}^{(int)}\underline{\omega} = 0, \quad {}^{(int)}\underline{\zeta} = {}^{(ext)}\underline{\zeta}, \quad {}^{(int)}\underline{\eta} = -{}^{(ext)}\underline{\zeta}, \\ {}^{(int)}\underline{\kappa} &= \lambda {}^{(ext)}\underline{\kappa}, \quad {}^{(int)}\underline{\vartheta} = \lambda {}^{(ext)}\underline{\vartheta}, \quad {}^{(int)}\underline{\kappa} = \lambda^{-1} {}^{(ext)}\underline{\kappa}, \quad {}^{(int)}\underline{\vartheta} = \lambda^{-1} {}^{(ext)}\underline{\vartheta}, \end{aligned}$$

and

$$\begin{aligned} {}^{(int)}\xi &= \frac{\lambda^2 \overline{{}^{(ext)}\kappa}}{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}} ({}^{(ext)}\zeta - {}^{(ext)}\eta), \\ {}^{(int)}\omega &= \frac{\lambda \overline{{}^{(ext)}\kappa}}{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}} {}^{(ext)}\underline{\omega}, \\ {}^{(int)}\eta &= {}^{(ext)}\zeta - \frac{\overline{{}^{(ext)}\kappa}}{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}} {}^{(ext)}\underline{\xi}. \end{aligned}$$

Proof. The following vectorfield is tangent to \mathcal{T}

$$\nu_{\mathcal{T}} := {}^{(ext)}e_3 - \frac{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}}{\overline{{}^{(ext)}\kappa}} {}^{(ext)}e_4,$$

which can also be written as

$$\nu_{\mathcal{T}} = \lambda {}^{(int)}e_3 - \frac{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}}{\overline{{}^{(ext)}\kappa}} \lambda^{-1} {}^{(int)}e_4.$$

Since $\nu_{\mathcal{T}}$ is tangent to \mathcal{T} , and in view of the definition of \underline{u} and ${}^{(int)}s$, we immediately infer

$$\nu_{\mathcal{T}}(\underline{u}) = \nu_{\mathcal{T}}(u) \text{ and } \nu_{\mathcal{T}}({}^{(int)}s) = \nu_{\mathcal{T}}({}^{(ext)}s) \text{ on } \mathcal{T}$$

and hence, using the identities

$${}^{(ext)}e_4(u) = {}^{(int)}e_3(\underline{u}) = 0, \quad {}^{(ext)}e_4({}^{(ext)}s) = 1, \quad {}^{(int)}e_3({}^{(int)}s) = -1,$$

we deduce on \mathcal{T}

$$\begin{aligned} -\frac{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}}{\overline{{}^{(ext)}\kappa}} \lambda^{-1} {}^{(int)}e_4(\underline{u}) &= {}^{(ext)}e_3(u), \\ -\lambda - \frac{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}}{\overline{{}^{(ext)}\kappa}} \lambda^{-1} {}^{(int)}e_4({}^{(int)}s) &= {}^{(ext)}e_3({}^{(ext)}s) - \frac{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}}{\overline{{}^{(ext)}\kappa}}. \end{aligned}$$

In view of the definition of ${}^{(ext)}\zeta$, ${}^{(int)}\underline{\xi}$, ${}^{(ext)}\underline{\Omega}$ and ${}^{(int)}\Omega$, this yields

$$\begin{aligned} {}^{(int)}\underline{\xi} &= -\frac{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}}{\overline{{}^{(ext)}\kappa}} \lambda^{-1} {}^{(ext)}\zeta, \\ {}^{(int)}\Omega &= \lambda - \lambda^2 \frac{\overline{{}^{(ext)}\kappa}}{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}} - \lambda \frac{\overline{{}^{(ext)}\kappa}}{\overline{{}^{(ext)}\kappa} + \overline{{}^{(ext)}\underline{A}}} {}^{(ext)}\underline{\Omega}. \end{aligned}$$

Next, we consider the Ricci coefficients of ${}^{(int)}\mathcal{M}$ on \mathcal{T} . From

$${}^{(int)}e_4 = \lambda {}^{(ext)}e_4, \quad {}^{(int)}e_3 = \lambda^{-1} {}^{(ext)}e_3, \quad {}^{(int)}e_\theta = {}^{(ext)}e_\theta \quad \text{on } \mathcal{T},$$

the fact that λ is constant on \mathcal{T} , and the fact that ${}^{(ext)}e_\theta$ is tangent to \mathcal{T} , we infer on \mathcal{T}

$${}^{(int)}\alpha = \lambda^2 {}^{(ext)}\alpha, \quad {}^{(int)}\beta = \lambda {}^{(ext)}\beta, \quad {}^{(int)}\rho = {}^{(ext)}\rho, \quad {}^{(int)}\underline{\beta} = \lambda^{-1} {}^{(ext)}\underline{\beta}, \quad {}^{(int)}\underline{\alpha} = \lambda^{-2} {}^{(ext)}\underline{\alpha},$$

and

$${}^{(int)}\zeta = {}^{(ext)}\zeta, \quad {}^{(int)}\kappa = \lambda {}^{(ext)}\kappa, \quad {}^{(int)}\vartheta = \lambda {}^{(ext)}\vartheta, \quad {}^{(int)}\underline{\kappa} = \lambda^{-1} {}^{(ext)}\underline{\kappa}, \quad {}^{(int)}\underline{\vartheta} = \lambda^{-1} {}^{(ext)}\underline{\vartheta}.$$

Also, since the foliation of ${}^{(int)}\mathcal{M}$ is ingoing geodesic, we have

$${}^{(int)}\underline{\xi} = 0, \quad {}^{(int)}\underline{\omega} = 0, \quad {}^{(int)}\underline{\eta} = -{}^{(int)}\zeta.$$

It remains to find identities for ${}^{(int)}\xi$, ${}^{(int)}\omega$ and ${}^{(int)}\eta$. Since λ is constant on \mathcal{T} and $\nu_{\mathcal{T}}$ tangent to \mathcal{T} , we have on \mathcal{T}

$$D_{\nu_{\mathcal{T}}} {}^{(int)}e_4 = \lambda D_{\nu_{\mathcal{T}}} {}^{(ext)}e_4, \quad D_{\nu_{\mathcal{T}}} {}^{(int)}e_3 = \lambda^{-1} D_{\nu_{\mathcal{T}}} {}^{(ext)}e_3$$

and hence

$$\begin{aligned} g(D_{\nu_{\mathcal{T}}} {}^{(int)}e_4, {}^{(int)}e_\theta) &= \lambda g(D_{\nu_{\mathcal{T}}} {}^{(ext)}e_4, {}^{(ext)}e_\theta), \\ g(D_{\nu_{\mathcal{T}}} {}^{(int)}e_4, {}^{(int)}e_3) &= g(D_{\nu_{\mathcal{T}}} {}^{(ext)}e_4, {}^{(ext)}e_3), \\ g(D_{\nu_{\mathcal{T}}} {}^{(int)}e_3, {}^{(int)}e_\theta) &= \lambda^{-1} g(D_{\nu_{\mathcal{T}}} {}^{(ext)}e_3, {}^{(ext)}e_\theta). \end{aligned}$$

We deduce

$$\begin{aligned} 2\lambda {}^{(int)}\eta - 2 \frac{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}}{\overline{{}^{(ext)}\kappa}} \lambda^{-1} {}^{(int)}\xi &= \lambda \left(2 {}^{(ext)}\eta - 2 \frac{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}}{\overline{{}^{(ext)}\kappa}} {}^{(ext)}\xi \right), \\ -4\lambda {}^{(int)}\underline{\omega} - 4 \frac{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}}{\overline{{}^{(ext)}\kappa}} \lambda^{-1} {}^{(int)}\omega &= -4 {}^{(ext)}\underline{\omega} - 4 \frac{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}}{\overline{{}^{(ext)}\kappa}} {}^{(ext)}\omega, \\ 2\lambda {}^{(int)}\underline{\xi} - 2 \frac{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}}{\overline{{}^{(ext)}\kappa}} \lambda^{-1} {}^{(int)}\underline{\eta} &= \lambda^{-1} \left(2 {}^{(ext)}\underline{\xi} - 2 \frac{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}}{\overline{{}^{(ext)}\kappa}} {}^{(ext)}\underline{\eta} \right), \end{aligned}$$

and thus

$$\begin{aligned} {}^{(int)}\xi &= \frac{\lambda^2 \overline{{}^{(ext)}\kappa}}{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}} ({}^{(ext)}\zeta - {}^{(ext)}\eta), \\ {}^{(int)}\omega &= \frac{\lambda \overline{{}^{(ext)}\kappa}}{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}} {}^{(ext)}\omega, \\ {}^{(int)}\underline{\eta} &= {}^{(ext)}\zeta - \frac{\overline{{}^{(ext)}\kappa}}{\overline{{}^{(ext)}\kappa} + {}^{(ext)}\underline{A}} {}^{(ext)}\underline{\xi}. \end{aligned}$$

This concludes the proof of the lemma. \square

Remark 7.7.2. Since the 2-spheres $\mathbf{S}(u, {}^{(int)}s)$ coincide on \mathcal{T} with the 2-sphere $\mathbf{S}(u, {}^{(ext)}s)$, the above lemma immediately yields

$$\begin{aligned} {}^{(int)}\check{\rho} &= {}^{(ext)}\check{\rho}, & {}^{(int)}\check{\kappa} &= \lambda {}^{(ext)}\check{\kappa}, & {}^{(int)}\check{\underline{\kappa}} &= \lambda^{-1} {}^{(ext)}\check{\underline{\kappa}} \\ {}^{(int)}\check{\underline{\mu}} &= -{}^{(ext)}\check{\underline{\mu}} - 2 {}^{(ext)}\check{\rho} + \frac{1}{2} {}^{(ext)}\vartheta {}^{(ext)}\check{\underline{\vartheta}} - \frac{1}{2} \overline{{}^{(ext)}\vartheta} \overline{{}^{(ext)}\check{\underline{\vartheta}}}, \\ {}^{(int)}\check{\underline{\omega}} &= \lambda \overline{{}^{(ext)}\kappa} \left(\frac{{}^{(ext)}\underline{\omega}}{\overline{{}^{(ext)}\underline{\kappa}} + \overline{{}^{(ext)}\underline{A}}} - \frac{\overline{{}^{(ext)}\underline{\omega}}}{\overline{{}^{(ext)}\underline{\kappa}} + \overline{{}^{(ext)}\underline{A}}} \right), \end{aligned}$$

and

$$\begin{aligned} {}^{(int)}\check{\underline{\zeta}} &= -\frac{1}{\lambda \overline{{}^{(ext)}\kappa}} \left(\overline{\left(\overline{{}^{(ext)}\underline{\kappa}} + \overline{{}^{(ext)}\underline{A}} \right) {}^{(ext)}\zeta} - \overline{\left(\overline{{}^{(ext)}\underline{\kappa}} + \overline{{}^{(ext)}\underline{A}} \right) {}^{(ext)}\zeta} \right), \\ {}^{(int)}\check{\underline{\Omega}} &= -\lambda^2 \overline{{}^{(ext)}\kappa} \left(\frac{1}{\overline{{}^{(ext)}\underline{\kappa}} + \overline{{}^{(ext)}\underline{A}}} - \frac{1}{\overline{{}^{(ext)}\underline{\kappa}} + \overline{{}^{(ext)}\underline{A}}} \right) \\ &\quad - \lambda \overline{{}^{(ext)}\kappa} \left(\frac{{}^{(ext)}\underline{\Omega}}{\overline{{}^{(ext)}\underline{\kappa}} + \overline{{}^{(ext)}\underline{A}}} - \frac{\overline{{}^{(ext)}\underline{\Omega}}}{\overline{{}^{(ext)}\underline{\kappa}} + \overline{{}^{(ext)}\underline{A}}} \right). \end{aligned}$$

Together with the estimates on \mathcal{T} for the outgoing geodesic foliation of ${}^{(ext)}\mathcal{M}$ derived in Theorem M4, we infer the control of tangential derivatives to \mathcal{T} , i.e. $(e_\theta, T_{\mathcal{T}})$ derivatives. Recovering the traversal derivative thanks to the transport equations in the direction e_3 , we infer for the ingoing geodesic foliation of ${}^{(int)}\mathcal{M}$ on \mathcal{T}

$$\begin{aligned} \max_{0 \leq k \leq k_{small} + 8} \sup_{\mathcal{T}} u^{1+\delta_{dec}} \left\| \mathfrak{D}^k \left({}^{(int)}\alpha, {}^{(int)}\beta, {}^{(int)}\check{\rho}, \overline{{}^{(int)}\beta}, {}^{(int)}\check{\underline{\mu}}, {}^{(int)}\check{\kappa}, {}^{(int)}\vartheta, {}^{(int)}\zeta, \right. \right. \\ \left. \left. {}^{(int)}\check{\underline{\eta}}, {}^{(int)}\check{\underline{\kappa}}, \overline{{}^{(int)}\vartheta}, {}^{(int)}\xi, {}^{(int)}\check{\underline{\omega}}, {}^{(int)}\check{\underline{\zeta}}, {}^{(int)}\check{\underline{\Omega}} \right) \right\|_{L^2(S)} \lesssim \epsilon_0. \end{aligned}$$

Step 2. Relying on the estimates of the ingoing geodesic foliation of ${}^{(int)}\mathcal{M}$ on \mathcal{T} derived in Step 1, we propagate these estimates to ${}^{(int)}\mathcal{M}$ thanks to transport equations in the e_3 direction given by the null structure equations and Bianchi identities. Recalling that $\underline{\alpha}$ has already been estimated in Theorem M3, see (7.7.1), quantities are recovered in the following order

1. We recover $\check{\underline{\kappa}}$, with a control of $k_{small} + 8$ derivatives, from

$$e_3 \check{\underline{\kappa}} + \overline{\kappa} \check{\underline{\kappa}} = \text{Err}[e_3 \check{\underline{\kappa}}].$$

2. We recover $\check{\underline{\vartheta}}$, with a control of $k_{small} + 8$ derivatives, from

$$e_3(\check{\underline{\vartheta}}) + \overline{\kappa} \check{\underline{\vartheta}} = -2\underline{\alpha}.$$

3. We recover $\underline{\beta}$, with a control of $k_{small} + 8$ derivatives, from

$$e_3\underline{\beta} + 2\underline{\kappa}\underline{\beta} = \underline{\mathcal{A}}_2\underline{\alpha} - \underline{\zeta}\underline{\alpha}.$$

4. We recover $\underline{\zeta}$, with a control of $k_{small} + 8$ derivatives, from

$$e_3(\underline{\zeta}) + \underline{\kappa}\underline{\zeta} = \underline{\beta} - \underline{\vartheta}\underline{\zeta}.$$

5. We recover $\underline{\eta}$, with a control of $k_{small} + 8$ derivatives, from

$$e_3(\underline{\eta} + \underline{\zeta}) + \frac{1}{2}\underline{\kappa}(\underline{\eta} + \underline{\zeta}) = -\frac{1}{2}\underline{\vartheta}(\underline{\eta} + \underline{\zeta}).$$

6. We recover $\underline{\check{\mu}}$, with a control of $k_{small} + 8$ derivatives, from

$$e_3\underline{\check{\mu}} + \frac{3}{2}\underline{\bar{\kappa}}\underline{\check{\mu}} + \frac{3}{2}\underline{\check{\mu}}\underline{\bar{\kappa}} = \text{Err}[e_3\underline{\check{\mu}}].$$

7. We recover $\underline{\check{\rho}}$, with a control of $k_{small} + 7$ derivatives, from

$$e_3\underline{\check{\rho}} + \frac{3}{2}\underline{\bar{\kappa}}\underline{\check{\rho}} + \frac{3}{2}\underline{\check{\rho}}\underline{\bar{\kappa}} = \underline{\mathcal{A}}_1\underline{\beta} + \text{Err}[e_3\underline{\check{\rho}}].$$

8. We recover $\underline{\check{\kappa}}$, with a control of $k_{small} + 7$ derivatives, from

$$e_3\underline{\check{\kappa}} + \frac{1}{2}\underline{\bar{\kappa}}\underline{\check{\kappa}} + \frac{1}{2}\underline{\check{\kappa}}\underline{\bar{\kappa}} = 2\underline{\mathcal{A}}_1\underline{\zeta} + 2\underline{\check{\rho}} + \text{Err}[e_3\underline{\check{\kappa}}].$$

9. We recover $\underline{\vartheta}$, with a control of $k_{small} + 7$ derivatives, from

$$e_3\underline{\vartheta} + \frac{1}{2}\underline{\kappa}\underline{\vartheta} = -2\underline{\mathcal{A}}_2^*\underline{\zeta} - \frac{1}{2}\underline{\kappa}\underline{\vartheta} + 2\underline{\zeta}^2.$$

10. We recover $\underline{\beta}$, with a control of $k_{small} + 6$ derivatives, from

$$e_3\underline{\beta} + \underline{\kappa}\underline{\beta} = e_\theta(\underline{\rho}) + 3\underline{\zeta}\underline{\rho} - \underline{\vartheta}\underline{\beta}.$$

11. We recover $\underline{\alpha}$, with a control of $k_{small} + 5$ derivatives, from

$$e_3\underline{\alpha} + \frac{1}{2}\underline{\kappa}\underline{\alpha} = -\underline{\mathcal{A}}_2^*\underline{\beta} - \frac{3}{2}\underline{\vartheta}\underline{\rho} + 5\underline{\zeta}\underline{\beta}.$$

12. We recover $\underline{\check{\omega}}$, with a control of $k_{small} + 7$ derivatives, from

$$e_3\underline{\check{\omega}} = \underline{\check{\rho}} + \text{Err}[e_3\underline{\check{\omega}}].$$

13. We recover $\check{\Omega}$, with a control of $k_{small} + 7$ derivatives, from

$$e_3(\check{\Omega}) = -2\check{\omega} + \overline{\check{\kappa}\check{\Omega}}.$$

14. We recover ξ , with a control of $k_{small} + 6$ derivatives, from

$$e_3(\xi) = e_4(\zeta) + \beta + \frac{1}{2}\kappa(\zeta - \underline{\eta}) + \frac{1}{2}\vartheta(\zeta - \underline{\eta}).$$

15. We recover $\underline{\zeta}$, with a control of $k_{small} + 8$ derivatives, from

$$e_3(\underline{\zeta} - 1) = 0.$$

As the estimates are significantly simpler to derive²⁵ and in the same spirit than the corresponding ones in Theorem M4, we leave the details to the reader. This concludes the proof of Theorem M5.

²⁵Note that r is bounded on ${}^{(int)}\mathcal{M}$ and that all quantities behave the same in ${}^{(int)}\mathcal{M}$.

Chapter 8

INITIALIZATION AND EXTENSION (Theorems M6, M7, M8)

In this chapter, we prove M6 concerning initialization, Theorem M7 concerning extension, and Theorem M8 concerning the improvement of high order weighted energies.

8.1 Proof of Theorem M6

Step 1. Let r_0 such that

$$r_0 := 2\epsilon_0^{-\frac{2}{3}}, \quad (8.1.1)$$

and let $\delta_0 > 0$ sufficiently small. Consider the unique sphere \mathring{S} of the initial data layer on $\mathcal{C}_{(1+\delta_0, \mathcal{L}_0)}$ with area radius r_0 . Then, denoting $S(u_{\mathcal{L}_0}, {}^{(ext)}s_{\mathcal{L}_0})$ the spheres of the outgoing portion of the initial data layer, we have

$$\mathring{S} = S(\mathring{u}, \mathring{s}), \quad \mathring{u} = 1 + \delta_0, \quad |\mathring{s} - r_0| \lesssim \epsilon_0.$$

Relying on the control of the initial data layer given by (3.3.5), i.e.

$$\mathfrak{I}_{k_{large}+5} \leq \epsilon_0,$$

we then invoke Theorem GCMS-II of section 3.7.4 with the choices

$$\mathring{\delta} = \mathring{\epsilon} = \epsilon_0, \quad s_{max} = k_{large} + 5,$$

to produce a unique GCM sphere S_* , which is a deformation of $\overset{\circ}{S}$, satisfying

$$\kappa^{S_*} = \frac{2}{r^{S_*}}, \quad \not\partial_2^{S_*} \not\partial_1^{S_*} \underline{\kappa}^{S_*} = \not\partial_2^{S_*} \not\partial_1^{S_*} \mu^{S_*} = 0, \quad \int_{S_*} \beta^{S_*} e^\Phi = 0, \quad \int_{S_*} e_\theta^{S_*}(\underline{\kappa}^{S_*}) e^\Phi = 0 \text{ on } S_*.$$

Step 2. Starting from S_* constructed in Step 1, and relying on the control of the initial data layer, we then invoke Theorem GCMH of section 3.7.4 to produce a smooth spacelike hypersurface Σ_* included in the initial data layer, passing through the sphere S_* , and a scalar function u defined on Σ_* such that

- The following GCM conditions holds

$$\kappa = \frac{2}{r}, \quad \not\partial_2^* \not\partial_1^* \underline{\kappa} = \not\partial_2^* \not\partial_1^* \mu = 0, \quad \int_S \eta e^\Phi = \int_S \underline{\xi} e^\Phi = 0 \text{ on } \Sigma_*$$

- We have, for some constant c_{Σ_*} ,

$$u + r = c_{\Sigma_*}, \quad \text{along } \Sigma_*.$$

- The following normalization condition holds true at the South Pole SP of every sphere S ,

$$a \Big|_{SP} = -1 - \frac{2m}{r}$$

where a is such that we have

$$\nu = e_3 + a e_4,$$

with ν the unique vectorfield tangent to the hypersurface Σ_* , normal to S , and normalized by $g(\nu, e_4) = -2$.

Furthermore, we have¹

$$\max_{k \leq k_{large} + 4} \sup_{\Sigma_*} r \left(|\not\partial^k f| + |\not\partial^k \underline{f}| + |\not\partial^k \log(\lambda)| \right) \lesssim \epsilon_0, \quad (8.1.2)$$

¹We have in fact

$$\max_{k \leq k_{large} + 6} \sup_{\Sigma_*} \left(\|\not\partial^k f\|_{L^2(S)} + \|\not\partial^k \underline{f}\|_{L^2(S)} + \|\not\partial^k \log(\lambda)\|_{L^2(S)} \right) \lesssim \epsilon_0,$$

and then use the Sobolev embedding on the 2-spheres S foliating Σ_* to deduce (8.1.2).

and

$$\sup_{\Sigma_*} \left(|m - m_0| + |r - r_0| \right) \lesssim \epsilon_0, \quad (8.1.3)$$

where $(f, \underline{f}, \lambda)$ are the transition function from the frame of the initial data layer to the frame of Σ_* .

Step 3. Provided $\delta_0 > 0$ has been chosen sufficiently small, the spacelike hypersurface Σ_* of Step 2 intersects the curve of the south poles of the spheres foliating the outgoing cone $\mathcal{C}_{(1, \mathcal{L}_0)}$ of the initial data layer. We then call S_1 the unique sphere of Σ_* such that its south pole coincides with the south pole of a sphere of $\mathcal{C}_{(1, \mathcal{L}_0)}$, and we calibrate u such that $u = 1$ on S_1 . We then can compare $\overset{\circ}{u} = 1 + \delta_0$ to $u(S_*)$ and obtain

$$|u(S_*) - 1 - \delta_0| \lesssim \epsilon_0 \delta_0,$$

so that

$$1 \leq u \leq u(S_*) \quad \text{on } \Sigma_* \text{ where } 1 < u(S_*) < 1 + 2\delta_0.$$

Together with the estimate (8.1.3), and in view of the choice (8.1.1) for r_0 , we have

$$\inf_{\Sigma_*} r \geq \frac{3}{2} \epsilon_0^{-\frac{2}{3}} \geq \epsilon_0^{-\frac{2}{3}} (u(S_*))^4$$

so that the dominant condition (3.3.4) for r is satisfied since $1 \leq u \leq 1 + 2\delta_0$ on Σ_* .

Step 4. In view of Step 1 to Step 3, Σ_* satisfies all the required properties for the future spacelike boundary of a GCM admissible spacetime, see item 3 of definition 3.1.2. We now control the outgoing geodesic foliation initialized on Σ_* and covering the region we denote by $^{(ext)}\mathcal{M}$, which is included in the initial data layer. Let $(f, \underline{f}, \lambda)$ the transition functions from the frame of the outgoing part of the initial data layer to the frame of $^{(ext)}\mathcal{M}$. Since both frames are outgoing geodesic, we may apply Corollary 2.3.7 which yields for $(\underline{f}, f, \log(\lambda))$ the following transport equations

$$\begin{aligned} \lambda^{-1} e'_4(rf) &= E'_1(f, \Gamma), \\ \lambda^{-1} e'_4(\log(\lambda)) &= E'_2(f, \Gamma), \\ \lambda^{-1} e'_4 \left(r\underline{f} - 2r^2 e'_\theta(\log(\lambda)) + rf\underline{\Omega} \right) &= E'_3(f, \underline{f}, \lambda, \Gamma), \end{aligned}$$

where

$$\begin{aligned} E'_1(f, \Gamma) &= -\frac{r}{2} \tilde{\kappa} f - \frac{r}{2} \vartheta f + \text{l.o.t.}, \\ E'_2(f, \Gamma) &= f\zeta - \frac{1}{2} f^2 \underline{\omega} - \underline{\eta} f - \frac{1}{4} f^2 \underline{\kappa} + \text{l.o.t.}, \\ E'_3(f, \underline{f}, \lambda, \Gamma) &= -\frac{r}{2} \tilde{\kappa} \underline{f} + r^2 \left(\tilde{\kappa} - \left(\bar{\kappa} - \frac{2}{r} \right) \right) e'_\theta(\log(\lambda)) + r^2 \left(\not{d}'_1(f) + \lambda^{-1} \vartheta' \right) e'_\theta(\log(\lambda)) \\ &\quad - \frac{r}{2} \tilde{\kappa} \underline{\Omega} f + r E_3(f, \underline{f}, \Gamma) - 2r^2 e'_\theta(E_2(f, \Gamma)) + r \underline{\Omega} E_1(f, \Gamma), \end{aligned}$$

and where E_1 , E_2 and E_3 are given in Lemma 2.3.6. Integrating these transport equations from Σ_* , using the control (8.1.2) of $(f, \underline{f}, \lambda)$ on Σ_* , and together with the assumption (3.3.5) for the Ricci coefficients of the foliation of the initial data layer, we obtain

$$\sup_{(ext)\mathcal{M}(r \geq 2m_0(1+\delta_{\mathcal{H}}))} r \left(|\mathfrak{d}^{\leq k_{large}+4}(f, \log(\lambda))| + |\mathfrak{d}^{\leq k_{large}+3}\underline{f}| \right) \lesssim \epsilon_0. \quad (8.1.4)$$

Then, let $\mathcal{T} = \{r = 2m_0(1 + \delta_{\mathcal{H}})\}$, i.e. we choose $r_{\mathcal{T}} = 2m_0(1 + \delta_{\mathcal{H}})$. We initialize the ingoing geodesic foliation of $(int)\mathcal{M}$ on \mathcal{T} using the outgoing geodesic foliation of $(ext)\mathcal{M}$ as in item 4 of definition 3.1.2. Using the control of $(f, \underline{f}, \lambda)$ induced on \mathcal{T} by (8.1.4), and using the analog of Corollary 2.3.7 in the e_3 direction for ingoing foliations, we obtain similarly,

$$\sup_{(int)\mathcal{M}} \left(|\mathfrak{d}^{\leq k_{large}+3}(\underline{f}, \log(\lambda))| + |\mathfrak{d}^{\leq k_{large}+2}f| \right) \lesssim \epsilon_0. \quad (8.1.5)$$

Then, in view of (8.1.4) (8.1.5), and the assumption (3.3.5) for the Ricci coefficients and curvature components of the foliation of the initial data layer, and using the transformation formulas of Proposition 2.3.4, we deduce

$$\begin{aligned} \max_{k \leq k_{large}} \left\{ \sup_{(ext)\mathcal{M}} \left(r^{\frac{7}{2}+\delta_B} (|\mathfrak{d}^k \alpha| + |\mathfrak{d}^k \beta|) + r^3 |\mathfrak{d}^k \rho| + r^2 |\mathfrak{d}^k \underline{\beta}| + r |\mathfrak{d}^k \underline{\alpha}| \right) \right. \\ + \sup_{(ext)\mathcal{M}} r^2 (|\mathfrak{d}^k \check{\kappa}| + |\mathfrak{d}^k \vartheta| + |\mathfrak{d}^k \zeta| + |\mathfrak{d}^k \check{\underline{\kappa}}|) \\ \left. + \sup_{(ext)\mathcal{M}} r (|\mathfrak{d}^k \eta| + |\mathfrak{d}^k \vartheta| + |\mathfrak{d}^k \check{\omega}| + |\mathfrak{d}^k \underline{\xi}|) \right\} \lesssim \epsilon_0, \end{aligned}$$

and

$$\max_{k \leq k_{large}} \sup_{(int)\mathcal{M}} \left(|\mathfrak{d}^k \check{R}| + |\mathfrak{d}^k \check{\Gamma}| \right) \lesssim \epsilon_0.$$

In particular, we infer that

$$\mathfrak{N}_{k_{large}}^{(En)} + \mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0$$

which concludes the proof of Theorem M6.

8.2 Proof of Theorem M7

From the assumptions of Theorem M7 we are given a GCM admissible spacetime $\mathcal{M} = \mathcal{M}(u_*) \in \mathfrak{N}(u_*)$ verifying the following improved bounds, for a universal constant $C > 0$,

$$\mathfrak{N}_{k_{small}+5}^{(Dec)}(\mathcal{M}) \leq C\epsilon_0 \quad (8.2.1)$$

provided by Theorems M1-M5. We then proceed as follows.

Step 1. We extend \mathcal{M} by a local existence argument, to a strictly larger spacetime $\mathcal{M}^{(extend)}$, with a naturally extended foliation and the following slightly increased bounds

$$\mathfrak{N}_{k_{small}+5}^{(Dec)}(\mathcal{M}^{(extend)}) \leq 2C\epsilon_0.$$

but which may not verify our admissibility criteria.

Step 2. We then invoke Theorem GCMH of section 3.7.4 to extend Σ_* in $\mathcal{M}^{(extend)} \setminus \mathcal{M}$ as a smooth spacelike hypersurface $\Sigma_*^{(extend)}$, together with a scalar function $u^{(extend)}$, satisfying the same GCM conditions than Σ_* .

Step 3. We consider the outgoing geodesic foliation $(u^{(extend)}, s^{(extend)})$ initialized on $\Sigma_*^{(extend)}$ to the future of $\Sigma_*^{(extend)}$ in $\mathcal{M}^{(extend)}$. Note in particular that we have from the definition of Σ_* and $\Sigma_*^{(extend)}$

$$u^{(extend)} + s^{(extend)} = c_{\Sigma_*}.$$

We define the following spacetime region to the future of $\Sigma_*^{(extend)}$

$$\tilde{\mathcal{R}} := \left\{ u_* \leq u^{(extend)} \leq u_* + \delta_{ext}, \quad c_{\Sigma_*} \leq u^{(extend)} + s^{(extend)} \leq c_{\Sigma_*} + \Delta_{ext} \right\},$$

where

$$\Delta_{ext} := \frac{4r_*}{u_*} \delta_{ext}, \quad r_* := r(S_*), \quad S_* := \Sigma_* \cap \mathcal{C}_*,$$

and $\delta_{ext} > 0$ is chosen sufficiently small so that $\tilde{\mathcal{R}} \subset \mathcal{M}^{(extend)}$. From now on, for convenience, we drop the index $(extend)$ and simply denote $u^{(extend)}$ and $s^{(extend)}$ by u and s .

Step 4. Propagating the GCM quantities in the e_4 direction from $\Sigma_*^{(extend)}$, where they all vanish, we obtain for all $k \leq k_{small} + 4$

$$\sup_{\tilde{\mathcal{R}}} \left(r^2 \left| \mathfrak{d}^k \left(\kappa - \frac{2}{r} \right) \right| + r^2 |\mathfrak{d}^{k-2}(r^2 \mathfrak{d}_2^* \mathfrak{d}_1^* \underline{\kappa})| + r^3 |\mathfrak{d}^{k-2}(r^2 \mathfrak{d}_2^* \mathfrak{d}_1^* \mu)| \right) \lesssim \frac{\epsilon_0}{r} \Delta_{ext}.$$

Similarly, we have for all $k \leq k_{small} + 4$

$$\sup_{\tilde{\mathcal{R}} \cap \{u \geq u_*\}} \left(r \left| \mathfrak{d}^k \left(\int_S \beta e^\Phi \right) \right| + \left| \mathfrak{d}^{k-1} \left(\int_S e_{\theta(\underline{\kappa})} e^\Phi \right) \right| \right) \lesssim \frac{\epsilon_0}{u_*} \delta_{ext} + \frac{\epsilon_0}{r} \Delta_{ext} \lesssim \frac{\epsilon_0}{r} \Delta_{ext},$$

where we used the fact that $u(S_*) = u_*$ and

$$\int_{S_*} \beta e^\Phi = \int_{S_*} e_\theta(\kappa) e^\Phi = 0.$$

Also, recall that $\nu = e_3 + a_* e_4$ denote the unique tangent vectorfield to Σ_* which is orthogonal to e_θ and normalized by $\mathbf{g}(\nu, e_4) = -2$. Then, one has, since $u + r$ is constant on Σ_* and $s = r$ on Σ_*

$$0 = \nu(u + s) = e_3(u) + a e_4(u) + e_3(s) + a e_4(s) = \frac{2}{\zeta} + \underline{\Omega} + a$$

and hence

$$a = -\frac{2}{\zeta} - \underline{\Omega} \text{ on } \Sigma_*.$$

Together with the GCM condition on a , we infer

$$\frac{2}{\zeta} + \underline{\Omega} = 1 + \frac{2m}{r} \text{ on } \Sigma_*.$$

As above, propagating forward in e_4 , we infer

$$\sup_{\tilde{\mathcal{R}}} \left| \frac{2}{\zeta} + \underline{\Omega} - \left(1 + \frac{2m}{r} \right) \right| \lesssim \frac{\epsilon_0}{r} \Delta_{ext}.$$

Step 5. We fix the following sphere of the $(u^{(extend)}, s^{(extend)})$ foliation in $\tilde{\mathcal{R}} \cap \{u \geq u_*\}$

$$\mathring{S} := S(\mathring{u}, \mathring{s}), \quad \mathring{u} := u_* + \frac{\delta_{ext}}{2}, \quad \mathring{s} := r_* + \frac{3r_*}{u_*} \delta_{ext}. \quad (8.2.2)$$

Define

$$\mathring{\delta} := \frac{\epsilon_0}{r} \Delta_{ext} = \frac{\epsilon_0 \delta_{ext}}{u_*}, \quad \mathring{\epsilon} := \frac{\epsilon_0}{u_*^{\frac{1}{2} + \delta_{dec}}}$$

and the small spacetime neighborhood of \mathring{S}

$$\mathcal{R}(\mathring{\epsilon}, \mathring{\delta}) := \left\{ |u - \mathring{u}| \leq \delta_{\mathcal{R}}, \quad |s - \mathring{s}| \leq \delta_{\mathcal{R}} \right\}, \quad \delta_{\mathcal{R}} = \mathring{\delta}(\mathring{\epsilon})^{-\frac{1}{2}}.$$

Note that $\mathcal{R}(\mathring{\epsilon}, \mathring{\delta}) \subset \tilde{\mathcal{R}}$. We are in position to apply Theorem GCMS II of section 3.7.4, with $s_{max} = k_{small} + 4$, which yields the existence of a unique sphere \tilde{S}_* , which is a

deformation of $\overset{\circ}{S}$, is included in $\mathcal{R}(\overset{\circ}{\epsilon}, \overset{\circ}{\delta})$, and is such that the following GCM conditions hold on it

$$\tilde{\phi}_2^* \tilde{\phi}_1^* \tilde{\kappa} = \tilde{\phi}_2^* \tilde{\phi}_1^* \tilde{\mu} = 0, \quad \tilde{\kappa} = \frac{2}{\tilde{r}}, \quad \int_{\tilde{S}_*} \tilde{\beta} e^\Phi = \int_{\tilde{S}_*} \tilde{e}_\theta(\tilde{\kappa}) e^\Phi = 0,$$

where the tilde refer to the quantities and tangential operators on \tilde{S}_* .

Step 6. Starting from \tilde{S}_* constructed in Step 5, we apply Theorem GCMH of section 3.7.4, with $s_{max} = k_{small} + 4$, which yields the existence of a smooth small piece of spacelike $\tilde{\Sigma}_*$ starting from \tilde{S}_* towards the initial data layer, together with a scalar function \tilde{u} defined on $\tilde{\Sigma}_*$, whose level surfaces are topological spheres denoted by \tilde{S} , so that

- the following GCM conditions are verified on $\tilde{\Sigma}_*$

$$\tilde{\phi}_2^* \tilde{\phi}_1^* \tilde{\kappa} = \tilde{\phi}_2^* \tilde{\phi}_1^* \tilde{\mu} = 0, \quad \tilde{\kappa} = \frac{2}{\tilde{r}}, \quad \int_{\tilde{S}} \tilde{\eta} e^\Phi = \int_{\tilde{S}} \tilde{\xi} e^\Phi = 0,$$

where the tilde refer to the quantities and tangential operators on $\tilde{\Sigma}_*$.

- We have, for some constant $c_{\tilde{\Sigma}_*}$,

$$\tilde{u} + \tilde{r} = c_{\tilde{\Sigma}_*}, \quad \text{along } \tilde{\Sigma}_*.$$

- The following normalization condition holds true at the South Pole SP of every sphere \tilde{S} ,

$$\tilde{a} \Big|_{SP} = -1 - \frac{2\tilde{m}}{\tilde{r}}$$

where \tilde{a} is such that we have

$$\tilde{\nu} = \tilde{e}_3 + \tilde{a}\tilde{e}_4,$$

with $\tilde{\nu}$ the unique vectorfield tangent to the hypersurface $\tilde{\Sigma}_*$, normal to \tilde{S} , and normalized by $g(\tilde{\nu}, \tilde{e}_4) = -2$.

- The transition functions $(f, \underline{f}, \lambda)$ from the frame of $\mathcal{M}^{(extend)}$ to the frame of $\tilde{\Sigma}_*$

$$\|(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}_{k_{small}+5}} \lesssim \overset{\circ}{\delta}.$$

Step 7. The spacelike GCM hypersurface $\tilde{\Sigma}_*$ has been constructed in Step 6 in a small neighborhood of \tilde{S}_* . We now focus on proving that it in fact extends all the way to the initial data layer. To this end, we denote by u_1 with

$$1 \leq u_1 < \overset{\circ}{u},$$

the minimal value of u such that:

- We have

$$\tilde{\Sigma}_* \cap \mathcal{C}_u \neq \emptyset \text{ for any } u_1 \leq u \leq \overset{\circ}{u}. \quad (8.2.3)$$

- There exists a large constant $D \geq 1$ such that we have for any sphere \tilde{S} of $\tilde{\Sigma}_*(u \geq u_1)$

$$\|(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}_{k_{small}+5}(\tilde{S})} \leq Du_* \overset{\circ}{\delta}. \quad (8.2.4)$$

- For the same large constant $D \geq 1$ as above, we have along $\tilde{\Sigma}_*(u \geq u_1)$

$$|\psi(s)| \leq Du_* \overset{\circ}{\delta}, \quad (8.2.5)$$

where the function $\psi(s)$ is such that the curve

$$\left(u = -s + c_{\tilde{\Sigma}_*} + \psi(s), s, \theta = 0\right) \text{ with } \psi(\overset{\circ}{s}) = 0, \quad (8.2.6)$$

coincides with the south poles of the sphere \tilde{S} of $\tilde{\Sigma}_*$ and the constant $c_{\tilde{\Sigma}_*}$ is fixed by the condition $\psi(\overset{\circ}{s}) = 0$.

The fact that $\psi(\overset{\circ}{s}) = 0$ together with the bounds of Step 6 implies that (8.2.3) (8.2.4) (8.2.5) hold for $u_1 < \overset{\circ}{u}$ with u_1 close enough to $\overset{\circ}{u}$. By a continuity argument based on reapplying Theorem GCMH, it suffices to show that we may improve the bounds (8.2.4) (8.2.5) independently of the value of u_1 .

Step 8. We now focus on improving the bounds (8.2.4) (8.2.5). We first prove that $\tilde{\Sigma}_*(u \geq u_1)$ is included in $\tilde{\mathcal{R}}$. Indeed, (8.2.4) (8.2.5) imply

$$\begin{aligned} \sup_{\tilde{\Sigma}_*(u \geq u_1)} |u + s - c_{\tilde{\Sigma}_*}| &\lesssim \sup_{\tilde{\Sigma}_*(u \geq u_1)} (|\psi| + r|f| + r|\underline{f}|) \\ &\lesssim Du_* \overset{\circ}{\delta} \\ &\lesssim \frac{Du_*}{r} \epsilon_0 \Delta_{ext} \\ &\lesssim \frac{\epsilon_0^{\frac{2}{3}} D}{u_*^3} \epsilon_0 \Delta_{ext} \\ &\lesssim \epsilon_0 \Delta_{ext}. \end{aligned}$$

On the other hand, by construction, $\psi(\mathring{s}) = 0$ and the south pole of \mathring{S} and \tilde{S}_* coincide, so that we have

$$\begin{aligned} c_{\tilde{\Sigma}_*} &= \mathring{u} + \mathring{s} = u_* + r_* + \frac{\delta_{ext}}{2} + \frac{3r_*}{u_*} \delta_{ext} \\ &= c_{\Sigma_*} + \frac{3}{4} \left(1 + \frac{2u_*}{3r_*} \right) \Delta_{ext} \end{aligned}$$

and hence

$$\begin{aligned} \sup_{\tilde{\Sigma}_*(u \geq u_1)} \left| u + s - c_{\Sigma_*} - \frac{3}{4} \Delta_{ext} \right| &\lesssim \left(\frac{u_*}{2r_*} + \epsilon_0 \right) \Delta_{ext} \\ &\lesssim \epsilon_0^{\frac{2}{3}} \Delta_{ext}. \end{aligned}$$

In view of the definition of $\tilde{\mathcal{R}}$, we infer

$$\tilde{\Sigma}_*(u \geq u_1) \subset \tilde{\mathcal{R}} \quad (8.2.7)$$

as claimed.

Step 9. Since $\tilde{\Sigma}_*(u \geq u_1) \subset \tilde{\mathcal{R}}$, the bound of Step 4 apply, and hence we have

$$\sup_{\tilde{\mathcal{R}}} \left| \frac{2}{\varsigma} + \underline{\Omega} - \left(1 + \frac{2m}{r} \right) \right| \lesssim \frac{\epsilon_0}{r} \Delta_{ext} \lesssim \mathring{\delta},$$

and for all $k \leq k_{small} + 4$

$$\sup_{\tilde{\mathcal{R}}} \left(r^2 \left| \mathfrak{d}^k \left(\kappa - \frac{2}{r} \right) \right| + r^2 |\mathfrak{d}^{k-2}(r^2 \mathfrak{d}_2^* \mathfrak{d}_1^* \kappa)| + r^3 |\mathfrak{d}^{k-2}(r^2 \mathfrak{d}_2^* \mathfrak{d}_1^* \mu)| \right) \lesssim \frac{\epsilon_0}{r} \Delta_{ext} \lesssim \mathring{\delta},$$

as well as

$$\sup_{\tilde{\mathcal{R}} \cap \{u \geq u_*\}} \left(r \left| \mathfrak{d}^k \left(\int_S \beta e^\Phi \right) \right| + \left| \mathfrak{d}^{k-1} \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) \right| \right) \lesssim \frac{\epsilon_0}{r} \Delta_{ext} \lesssim \mathring{\delta}.$$

Together with the a priori estimates of Chapter 9 on the GCM construction, this yields

$$\begin{aligned} |\psi'(s)| &\lesssim \left| 1 + \frac{2\tilde{m}}{\tilde{r}} + \underline{\Omega} + \frac{2}{\varsigma} \right| + |\lambda - 1| \\ &\lesssim \left| \frac{\tilde{m}}{\tilde{r}} - \frac{m}{r} \right| + |\lambda - 1| + \frac{\epsilon_0}{r} \Delta_{ext}. \end{aligned}$$

In view of (8.2.4), we have

$$|\tilde{r} - r| + |\tilde{m} - m| \lesssim \sup_{\tilde{S}} r(|f| + |\underline{f}|) \lesssim Du_* \mathring{\delta} \quad (8.2.8)$$

and we infer

$$\begin{aligned} |\psi'(s)| &\lesssim \frac{Du_* \overset{\circ}{\delta}}{r} + \overset{\circ}{\delta} \\ &\lesssim \left(1 + \frac{\epsilon_0^{\frac{2}{3}}}{(u_*)^3 D}\right) \overset{\circ}{\delta} \\ &\lesssim \overset{\circ}{\delta}. \end{aligned}$$

Integrating from $\overset{\circ}{s}$ where $\psi(\overset{\circ}{s}) = 0$, we infer

$$\begin{aligned} |\psi(s)| &\lesssim |s - \overset{\circ}{s}| \overset{\circ}{\delta} \\ &\lesssim u_* \overset{\circ}{\delta} \end{aligned}$$

which improves (8.2.5) for $D \geq 1$ large enough.

Similarly, we obtain

$$\|(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}_{k_{small}+5}(\tilde{S})} \lesssim r^{-2} \left(\left| \int_S f e^\Phi \right| + \left| \int_S \underline{f} e^\Phi \right| \right) + \overset{\circ}{\delta}$$

and

$$\left| e_3 \left(\int_S f e^\Phi \right) \right| + \left| e_3 \left(\int_S \underline{f} e^\Phi \right) \right| \lesssim r^2 \overset{\circ}{\delta} + \frac{1}{r} \left(\left| \int_S f e^\Phi \right| + \left| \int_S \underline{f} e^\Phi \right| \right).$$

In view of (8.2.4), we infer

$$\left| e_3 \left(\int_S f e^\Phi \right) \right| + \left| e_3 \left(\int_S \underline{f} e^\Phi \right) \right| \lesssim r^2 \overset{\circ}{\delta} + r D u_* \overset{\circ}{\delta}$$

and integrating from \tilde{S}_* , we infer

$$\begin{aligned} r^{-2} \left(\left| \int_S f e^\Phi \right| + \left| \int_S \underline{f} e^\Phi \right| \right) &\lesssim u_* \overset{\circ}{\delta} + \frac{D(u_*)^2 \overset{\circ}{\delta}}{r} \\ &\lesssim \left(1 + \frac{\epsilon_0^{\frac{2}{3}} D}{(u_*)^2}\right) u_* \overset{\circ}{\delta} \\ &\lesssim u_* \overset{\circ}{\delta}. \end{aligned}$$

This yields

$$\|(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}_{k_{small}+5}(\tilde{S})} \lesssim u_* \overset{\circ}{\delta}$$

which improves (8.2.4) for $D \geq 1$ large enough. We thus conclude that $u_1 = 1$, $\tilde{\Sigma}_*$ extends all the way to the initial data layer, $\tilde{\Sigma}_* \subset \tilde{\mathcal{R}}$, and we have the bounds

$$\|(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}^{k_{small}+5}(\tilde{S})} \lesssim u_* \delta^\circ, \quad |\psi(s)| \lesssim u_* \delta^\circ.$$

In view of the definition of δ° , we infer in particular for any sphere \tilde{S} of $\tilde{\Sigma}_*$

$$\|(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}^{k_{small}+5}(\tilde{S})} \lesssim \epsilon_0 \delta_{ext}, \quad |\psi(s)| \lesssim \epsilon_0 \delta_{ext}. \quad (8.2.9)$$

Step 10. As $\tilde{\Sigma}_*$ extends all the way to the initial data layer, this allows us to calibrate \tilde{u} along $\tilde{\Sigma}_*$ by fixing the value $\tilde{u} = 1$ as in (3.1.5):

$$\tilde{S}_1 = \tilde{\Sigma}_* \cap \{\tilde{u} = 1\} \text{ is such that } \tilde{S}_1 \cap \mathcal{C}_{(1, \mathcal{L}_0)} \cap SP \neq \emptyset, \quad (8.2.10)$$

i.e. \tilde{S}_1 is the unique sphere of $\tilde{\Sigma}_*$ such that its south pole intersects the south pole of one of the sphere of the outgoing null cone $\mathcal{C}_{(1, \mathcal{L}_0)}$ of the initial data layer.

Now that \tilde{u} is calibrated, we define

$$\tilde{u}_* := \tilde{u}(\tilde{S}_*). \quad (8.2.11)$$

For the proof of Theorem M7, we need in particular to prove that $\tilde{u}_* > u_*$. First, note that, since $\tilde{u} + \tilde{r}$ is constant along $\tilde{\Sigma}_*$, we have

$$\tilde{\Sigma}_* = \left\{ \tilde{u} + \tilde{r} = 1 + \tilde{r}(\tilde{S}_1) \right\}. \quad (8.2.12)$$

Since $\tilde{S}_* \subset \tilde{\Sigma}_*$, and in view of (8.2.12), (8.2.2), (8.2.6), we infer,

$$\begin{aligned} \left| \tilde{u}(\tilde{S}_*) - \left(u_* + \frac{\delta_{ext}}{2} \right) \right| &= \left| \tilde{u}(\tilde{S}_*) - u(\tilde{S}) \right| \\ &= \left| 1 + \tilde{r}(\tilde{S}_1) - \tilde{r}(\tilde{S}_*) - \left(-s(\tilde{S}) + c_{\tilde{\Sigma}_*} \right) \right|. \end{aligned}$$

Next, note from

$$s = r \text{ on } \Sigma_*, \quad e_4(r - s) = \frac{r}{2} \left(\bar{\kappa} - \frac{2}{r} \right)$$

that we have

$$\sup_{\tilde{\mathcal{R}}} |r - s| \lesssim \frac{\epsilon_0}{r} \Delta_{ext} \lesssim \epsilon_0 \delta_{ext}. \quad (8.2.13)$$

Together with (8.2.8), this yields

$$\left| \tilde{u}(\tilde{S}_*) - \left(u_* + \frac{\delta_{ext}}{2} \right) \right| \lesssim \left| 1 + \tilde{r}(\tilde{S}_1) - c_{\tilde{\Sigma}_*} \right| + \epsilon_0 \delta_{ext}.$$

Since $c_{\tilde{\Sigma}_*}$ in (8.2.6) is a constant, we have in particular

$$c_{\tilde{\Sigma}_*} = u(\tilde{S}_1) + r(\tilde{S}_1) - \psi(s(\tilde{S}_1))$$

and thus

$$\begin{aligned} \left| \tilde{u}(\tilde{S}_*) - \left(u_* + \frac{\delta_{ext}}{2} \right) \right| &\lesssim \left| 1 + \tilde{r}(\tilde{S}_1) - u(\tilde{S}_1) - r(\tilde{S}_1) + \psi(s(\tilde{S}_1)) \right| + \epsilon_0 \delta_{ext} \\ &\lesssim \left| 1 - u(\tilde{S}_1) \right| + \left| \tilde{r}(\tilde{S}_1) - r(\tilde{S}_1) \right| + \left| \psi(s(\tilde{S}_1)) \right| + \epsilon_0 \delta_{ext}. \end{aligned}$$

In view of (8.2.9) and (8.2.8), we infer

$$\left| \tilde{u}(\tilde{S}_*) - \left(u_* + \frac{\delta_{ext}}{2} \right) \right| \lesssim \left| 1 - u(\tilde{S}_1) \right| + \epsilon_0 \delta_{ext}.$$

Also, since (recall in particular (3.1.5))

$$u = 1 \text{ on } S_1 \cap SP, \quad e_4^{\mathcal{L}_0}(u) = O\left(\frac{\epsilon_0}{r^2}\right),$$

and since the south pole of S_1 coincides with the one of the corresponding sphere of $\mathcal{C}_{\mathcal{L}_0,1}$, we infer

$$\sup_{\tilde{\mathcal{R}} \cap \mathcal{C}_{\mathcal{L}_0,1} \cap SP} |u - 1| \lesssim \Delta_{ext} \frac{\epsilon_0}{r^2} \lesssim \epsilon_0 \delta_{ext}.$$

This yields

$$\left| \tilde{u}(\tilde{S}_*) - \left(u_* + \frac{\delta_{ext}}{2} \right) \right| \lesssim \epsilon_0 \delta_{ext}. \quad (8.2.14)$$

In particular, we deduce, for ϵ_0 small enough,

$$\tilde{u}(\tilde{S}_*) > u_* \quad (8.2.15)$$

as desired.

Step 11. We would like to check that the dominant condition (3.3.4) for r holds on $\tilde{\Sigma}_*$, i.e. we need to prove

$$\tilde{r}(\tilde{S}_*) \geq \epsilon_0^{-\frac{2}{3}} (\tilde{u}(\tilde{S}_*))^4.$$

To this end, note that we have in view of (8.2.8), (8.2.13) and (8.2.14)

$$\begin{aligned}\tilde{r}(\tilde{S}_*) - \epsilon_0^{-\frac{2}{3}}(\tilde{u}(\tilde{S}_*))^4 &= s(\mathring{S}) + O(\epsilon_0\delta_{ext}) - \epsilon_0^{-\frac{2}{3}}\left(u_* + \frac{\delta_{ext}}{2} + O(\epsilon_0\delta_{ext})\right)^4 \\ &= s(\mathring{S}) - \epsilon_0^{-\frac{2}{3}}(u_*)^4 - 2\epsilon_0^{-\frac{2}{3}}(u_*)^3\delta_{ext} + \epsilon_0^{-\frac{2}{3}}(u_*)^3\delta_{ext}O\left(\frac{\delta_{ext}}{u_*} + \epsilon_0\right) + O(\epsilon_0\delta_{ext}).\end{aligned}$$

Together with (8.2.2), we infer

$$\begin{aligned}\tilde{r}(\tilde{S}_*) - \epsilon_0^{-\frac{2}{3}}(\tilde{u}(\tilde{S}_*))^4 &= r_* + \frac{3r_*}{u_*}\delta_{ext} - \epsilon_0^{-\frac{2}{3}}(u_*)^4 - 2\epsilon_0^{-\frac{2}{3}}(u_*)^3\delta_{ext} \\ &\quad + \epsilon_0^{-\frac{2}{3}}(u_*)^3\delta_{ext}O\left(\frac{\delta_{ext}}{u_*} + \epsilon_0\right) + O(\epsilon_0\delta_{ext}) \\ &= r_* - \epsilon_0^{-\frac{2}{3}}(u_*)^4 + \left(3r_* - 2\epsilon_0^{-\frac{2}{3}}(u_*)^4\right)\frac{\delta_{ext}}{u_*} \\ &\quad + \epsilon_0^{-\frac{2}{3}}(u_*)^3\delta_{ext}O\left(\frac{\delta_{ext}}{u_*} + \epsilon_0\right) + O(\epsilon_0\delta_{ext}).\end{aligned}$$

Since we have by the control (3.3.4) of r on Σ_*

$$r_* \geq \epsilon_0^{-\frac{2}{3}}(u_*)^4,$$

we deduce

$$\begin{aligned}\tilde{r}(\tilde{S}_*) - \epsilon_0^{-\frac{2}{3}}(\tilde{u}(\tilde{S}_*))^4 &\geq \frac{r_*\delta_{ext}}{u_*} + \epsilon_0^{-\frac{2}{3}}(u_*)^3\delta_{ext}O\left(\frac{\delta_{ext}}{u_*} + \epsilon_0\right) + O(\epsilon_0\delta_{ext}) \\ &= \frac{r_*\delta_{ext}}{u_*}(1 + O(\epsilon_0))\end{aligned}$$

so that, for ϵ_0 small enough, we have

$$\tilde{r}(\tilde{S}_*) \geq \epsilon_0^{-\frac{2}{3}}(\tilde{u}(\tilde{S}_*))^4$$

as desired.

Step 12. We summarize the properties of $\tilde{\Sigma}_*$ obtained so far:

- $\tilde{\Sigma}_*$ is a spacelike hypersurface included in the spacetime region $\tilde{\mathcal{R}}$.
- The scalar function \tilde{u} is defined on $\tilde{\Sigma}_*$ and its level sets are topological 2-spheres denoted by \tilde{S} .

- The following GCM conditions holds on $\tilde{\Sigma}_*$

$$\tilde{\phi}_2^* \tilde{\phi}_1^* \tilde{\kappa} = \tilde{\phi}_2^* \tilde{\phi}_1^* \tilde{\mu} = 0, \quad \tilde{\kappa} = \frac{2}{\tilde{r}}, \quad \int_{\tilde{S}} \tilde{\eta} e^\Phi = \int_{\tilde{S}} \tilde{\xi} e^\Phi = 0.$$

- In addition, the following GCM conditions holds on the sphere \tilde{S}_* of $\tilde{\Sigma}_*$

$$\int_{\tilde{S}_*} \tilde{\beta} e^\Phi = \int_{\tilde{S}_*} \tilde{e}_\theta(\tilde{\kappa}) e^\Phi = 0,$$

- We have, for some constant $c_{\tilde{\Sigma}_*}$,

$$\tilde{u} + \tilde{r} = c_{\tilde{\Sigma}_*}, \quad \text{along } \tilde{\Sigma}_*.$$

- The following normalization condition holds true at the South Pole SP of every sphere \tilde{S} ,

$$\tilde{a} \Big|_{SP} = -1 - \frac{2\tilde{m}}{\tilde{r}}$$

where \tilde{a} is such that we have

$$\tilde{\nu} = \tilde{e}_3 + \tilde{a}\tilde{e}_4,$$

with $\tilde{\nu}$ the unique vectorfield tangent to the hypersurface $\tilde{\Sigma}_*$, normal to \tilde{S} , and normalized by $g(\tilde{\nu}, \tilde{e}_4) = -2$.

- The dominant condition (3.3.4) for r holds on $\tilde{\Sigma}_*$, i.e. we have

$$\tilde{r}(\tilde{S}_*) \geq \epsilon_0^{-\frac{2}{3}} (\tilde{u}(\tilde{S}_*))^4.$$

- \tilde{u} is calibrated along $\tilde{\Sigma}_*$ by fixing the value $\tilde{u} = 1$:

$$\tilde{S}_1 = \tilde{\Sigma}_* \cap \{\tilde{u} = 1\} \text{ is such that } \tilde{S}_1 \cap \mathcal{C}_{(1, \mathcal{L}_0)} \cap SP \neq \emptyset, \quad (8.2.16)$$

i.e. \tilde{S}_1 is the unique sphere of $\tilde{\Sigma}_*$ such that its south pole intersects the south pole of one of the sphere of the outgoing null cone $\mathcal{C}_{(1, \mathcal{L}_0)}$ of the initial data layer.

Thus $\tilde{\Sigma}_*$ satisfies all the required properties for the future spacelike boundary of a GCM admissible spacetime, see item 3 of definition 3.1.2. Furthermore, we have on $\tilde{\Sigma}_*$

$$\tilde{u}(\tilde{S}_*) > u_*, \quad (8.2.17)$$

and $(f, \underline{f}, \lambda)$ satisfy in view of (8.2.9) and Corollary Rigidity III of section 3.7.4

$$\sup_{\tilde{\Sigma}_*} \|\mathfrak{d}^{\leq k_{small}+5}(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}_{k_{small}+5}(\tilde{S})} \lesssim \epsilon_0 \delta_{ext}.$$

Together with the Sobolev embedding on the spheres \tilde{S} , and possibly reducing the size of $\delta_{ext} > 0$, we deduce

$$\sup_{\tilde{\Sigma}_*} \tilde{r} \tilde{u}^{\frac{1}{2} + \delta_{dec}} |\mathfrak{d}^{\leq k_{small}+3}(f, \underline{f}, \log(\lambda))| \lesssim \epsilon_0. \quad (8.2.18)$$

Step 13. We now control the outgoing geodesic foliation initialized on $\tilde{\Sigma}_*$. We denote by ${}^{(ext)}\tilde{\mathcal{M}}$ the region covered by this outgoing geodesic foliation. Let (e_4, e_3, e_θ) of ${}^{(ext)}\mathcal{M}$ extended to the spacetime $\mathcal{M}^{(extend)}$, and satisfying, as discussed in Step 1 to Step 3

$$\mathfrak{R}_{k_{small}+5}^{(Dec)}(\mathcal{M}^{(extend)}) \lesssim \epsilon_0. \quad (8.2.19)$$

Let $(f, \underline{f}, \lambda)$ the transition functions from the frame (e_4, e_3, e_θ) to the frame $(\tilde{e}_4, \tilde{e}_3, \tilde{e}_\theta)$ of ${}^{(ext)}\tilde{\mathcal{M}}$. Since both frames are outgoing geodesic, we may apply Corollary 2.3.7 which yields for $(\underline{f}, f, \log(\lambda))$ the following transport equations

$$\begin{aligned} \lambda^{-1} e'_4(rf) &= E'_1(f, \Gamma), \\ \lambda^{-1} e'_4(\log(\lambda)) &= E'_2(f, \Gamma), \\ \lambda^{-1} e'_4\left(r\underline{f} - 2r^2 e'_\theta(\log(\lambda)) + rf\Omega\right) &= E'_3(f, \underline{f}, \lambda, \Gamma), \end{aligned}$$

where

$$\begin{aligned} E'_1(f, \Gamma) &= -\frac{r}{2} \check{\kappa} f - \frac{r}{2} \vartheta f + \text{l.o.t.}, \\ E'_2(f, \Gamma) &= f\zeta - \frac{1}{2} f^2 \underline{\omega} - \underline{\eta} f - \frac{1}{4} f^2 \underline{\kappa} + \text{l.o.t.}, \\ E'_3(f, \underline{f}, \lambda, \Gamma) &= -\frac{r}{2} \check{\kappa} \underline{f} + r^2 \left(\check{\kappa} - \left(\bar{\kappa} - \frac{2}{r} \right) \right) e'_\theta(\log(\lambda)) + r^2 \left(\mathcal{A}'_1(f) + \lambda^{-1} \vartheta' \right) e'_\theta(\log(\lambda)) \\ &\quad - \frac{r}{2} \check{\kappa} \underline{\Omega} f + r E_3(f, \underline{f}, \Gamma) - 2r^2 e'_\theta(E_2(f, \Gamma)) + r \underline{\Omega} E_1(f, \Gamma), \end{aligned}$$

and where E_1 , E_2 and E_3 are given in Lemma 2.3.6. Integrating these transport equations from $\tilde{\Sigma}_*$, using the control (8.2.18) of $(f, \underline{f}, \lambda)$ on $\tilde{\Sigma}_*$, and together with the control (8.2.19) for the Ricci coefficients of the foliation of $\mathcal{M}^{(extend)}$, we obtain

$$\sup_{{}^{(ext)}\tilde{\mathcal{M}}\left(\tilde{r} \geq 2m_0\left(1 + \frac{\delta_{\mathcal{H}}}{2}\right)\right)} \left(\tilde{r} \tilde{u}^{\frac{1}{2} + \delta_{dec}} + \tilde{u}^{1 + \delta_{dec}} \right) \left(|\mathfrak{d}^{\leq k_{small}+3}(f, \log(\lambda))| + |\mathfrak{d}^{\leq k_{small}+2} \underline{f}| \right) \lesssim \epsilon_0. \quad (8.2.20)$$

Then, for any $r_{\mathcal{T}}$ in the interval

$$2m_0 \left(1 + \frac{\delta_{\mathcal{H}}}{2}\right) \leq r_{\mathcal{T}} \leq 2m_0 \left(1 + \frac{3\delta_{\mathcal{H}}}{2}\right), \quad (8.2.21)$$

we initialize the ingoing geodesic foliation of ${}^{(int)}\widetilde{\mathcal{M}}[r_{\mathcal{T}}]$ on $\tilde{r} = r_{\mathcal{T}}$ using the outgoing geodesic foliation of ${}^{(ext)}\widetilde{\mathcal{M}}$ as in item 4 of definition 3.1.2. Using the control of $(f, \underline{f}, \lambda)$ induced on $\tilde{r} = r_{\mathcal{T}}$ by (8.2.20), and using the analog of Corollary 2.3.7 in the e_3 direction for ingoing foliations, we obtain similarly, for any $r_{\mathcal{T}}$ in the interval (8.2.21),

$$\sup_{{}^{(int)}\widetilde{\mathcal{M}}[r_{\mathcal{T}}]} \tilde{u}^{1+\delta_{dec}} \left(|\mathfrak{d}^{\leq k_{small}+2}(\underline{f}, \log(\lambda))| + |\mathfrak{d}^{\leq k_{small}+1} f| \right) \lesssim \epsilon_0. \quad (8.2.22)$$

Let now, for any $r_{\mathcal{T}}$ in the interval (8.2.21),

$$\mathcal{M}[r_{\mathcal{T}}] := {}^{(ext)}\widetilde{\mathcal{M}}(\tilde{r} \geq r_{\mathcal{T}}) \cup {}^{(int)}\widetilde{\mathcal{M}}[r_{\mathcal{T}}].$$

Then, in view of (8.2.20) (8.2.22), and (8.2.19), and using the transformation formulas of Proposition 2.3.4, we deduce

$$\mathfrak{N}_{k_{small}}^{(Dec)}(\mathcal{M}[r_{\mathcal{T}}]) \lesssim \epsilon_0$$

which concludes the proof of Theorem M7.

8.3 Proof of Theorem M8

So far, we have only improved our bootstrap assumptions on decay estimates. We now improve our bootstrap assumptions on energies and weighted energies for \check{R} and $\check{\Gamma}$ relying on an iterative procedure which recovers derivatives one by one².

Let $I_{m_0, \delta_{\mathcal{H}}}$ the interval of \mathbb{R} defined by

$$I_{m_0, \delta_{\mathcal{H}}} := \left[2m_0 \left(1 + \frac{\delta_{\mathcal{H}}}{2}\right), 2m_0 \left(1 + \frac{3\delta_{\mathcal{H}}}{2}\right) \right]. \quad (8.3.1)$$

Remark 8.3.1. Recall that the results of Theorems M0–M7 hold for any $r_{\mathcal{T}} \in I_{m_0, \delta_{\mathcal{H}}}$, see Remark 3.6.3. More precisely

²See also [29] for a related strategy to recover higher order derivatives from the control of lower order ones.

- they hold on ${}^{(ext)}\mathcal{M}(r \geq 2m_0(1 + \frac{\delta_{\mathcal{H}}}{2}))$, and hence on ${}^{(ext)}\mathcal{M}(r \geq r_{\mathcal{T}})$ for any $r_{\mathcal{T}} \in I_{m_0, \delta_{\mathcal{H}}}$,
- they hold on ${}^{(int)}\mathcal{M}[r_{\mathcal{T}}]$ for any $r_{\mathcal{T}} \in I_{m_0, \delta_{\mathcal{H}}}$, where ${}^{(int)}\mathcal{M}[r_{\mathcal{T}}]$ is initialized on $\mathcal{T} = \{r = r_{\mathcal{T}}\}$ using ${}^{(ext)}\mathcal{M}(r \geq r_{\mathcal{T}})$ as in section 3.1.2.

It is at this stage that we need to make a specific choice of $r_{\mathcal{T}}$ in the context of a Lebesgue point argument. More precisely, we choose $r_{\mathcal{T}}$ such that we have

$$\int_{\{r=r_{\mathcal{T}}\}} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 = \inf_{r_0 \in I_{m_0, \delta_{\mathcal{H}}}} \int_{\{r=r_0\}} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2. \quad (8.3.2)$$

Remark 8.3.2. In case the above infimum is achieved for several values of r , we choose $r_{\mathcal{T}}$ to be the largest of such values, so that $r_{\mathcal{T}}$ is uniquely defined. Note also that the infimum could a priori be infinite, and will only be shown to be finite - and more precisely $O(\epsilon_0)$ -, at the end of the proof of Theorem M8, see section 8.3.4. This could be made rigorous in the context of a continuity argument.

In view of the definition of $r_{\mathcal{T}}$, and since $\mathcal{T} = \{r = r_{\mathcal{T}}\}$, we have

$$\int_{\mathcal{T}} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 \leq \frac{1}{2m_0 \delta_{\mathcal{H}}} \int_{I_{m_0, \delta_{\mathcal{H}}}} \left(\int_{\{r=r_0\}} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 \right) dr_0$$

and hence³

$$\int_{\mathcal{T}} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 \lesssim \int_{{}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2. \quad (8.3.3)$$

From now on, we may thus assume that the spacetime \mathcal{M} satisfies

- the conclusions of Theorem M0, i.e.

$$\max_{0 \leq k \leq k_{large}} \left\{ \sup_{\mathcal{C}_1} \left[r^{\frac{7}{2} + \delta_B} (|\mathfrak{d}^k ({}^{(ext)}\alpha| + |\mathfrak{d}^k ({}^{(ext)}\beta)|) + r^{\frac{9}{2} + \delta_B} |\mathfrak{d}^{k-1} e_3 ({}^{(ext)}\alpha)| \right] \right. \\ \left. + \sup_{\mathcal{C}_1} \left[r^3 \left| \mathfrak{d}^k \left(({}^{(ext)}\rho + \frac{2m_0}{r^3}) \right) \right| + r^2 |\mathfrak{d}^k ({}^{(ext)}\underline{\beta})| + r |\mathfrak{d}^k ({}^{(ext)}\underline{\alpha})| \right] \right\} \lesssim \epsilon_0 \quad (8.3.4)$$

³We use the coarea formula, $d\mathcal{M} = \frac{1}{\sqrt{\mathbf{g}(\mathbf{D}r, \mathbf{D}r)}} d\{r = r_0\} dr_0$ and the fact that, for $r \in I_{m_0, \delta_{\mathcal{H}}}$, $\mathbf{g}(\mathbf{D}r, \mathbf{D}r) = -e_3(r)e_4(r) = \Upsilon + O(\epsilon) \geq \frac{\delta_{\mathcal{H}}}{2} + O(\epsilon + \delta_{\mathcal{H}}^2) \geq \frac{\delta_{\mathcal{H}}}{4}$. Note that \lesssim here depends on $\delta_{\mathcal{H}}^{-1}$, see the convention for \lesssim made at the end of section 3.3.1.

and

$$\begin{aligned} \max_{0 \leq k \leq k_{large}} \sup_{\mathcal{C}_1} \left[& |\mathfrak{d}^k (int) \alpha| + |\mathfrak{d}^k (int) \beta| + \left| \mathfrak{d}^k \left((int) \rho + \frac{2m_0}{r^3} \right) \right| \right. \\ & \left. + |\mathfrak{d}^k (int) \underline{\beta}| + |\mathfrak{d}^k (int) \underline{\alpha}| \right] \lesssim \epsilon_0, \end{aligned} \quad (8.3.5)$$

- the conclusions of Theorem M7, i.e.

$$\mathfrak{N}_{k_{small}}^{(Dec)} \lesssim \epsilon_0, \quad (8.3.6)$$

see section 3.2.3 for the definition of the combined norm on decay $\mathfrak{N}_k^{(Dec)}$,

- the estimate

$$\int_{\mathcal{T}} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 \lesssim \int_{(ext)\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2. \quad (8.3.7)$$

The goal of this section is to prove Theorem M8, i.e. to prove that the following bound holds on \mathcal{M} for the weighted energies

$$\mathfrak{N}_{k_{large}}^{(En)} \lesssim \epsilon_0,$$

see section 3.2.3 for the definition of the combined norm on weighted energies $\mathfrak{N}_k^{(En)}$.

8.3.1 Main norms

We recall below our norms for measuring weighted energies for curvature components and Ricci coefficients, see sections 3.2.1 and 3.2.2. Let $r_0 \geq 4m_0$. Then, we have for $(ext)\mathcal{M}$

$$\begin{aligned} \left((ext)\mathfrak{N}_0^{\geq r_0}[\check{R}] \right)^2 &= \sup_{0 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq 4m_0)} \left(r^{4+\delta_B} \alpha^2 + r^4 \beta^2 \right) \\ &+ \int_{\Sigma_*} \left(r^{4+\delta_B} (\alpha^2 + \beta^2) + r^4 (\check{\rho})^2 + r^2 \underline{\beta}^2 + \underline{\alpha}^2 \right) \\ &+ \int_{(ext)\mathcal{M}(r \geq 4m_0)} \left(r^{3+\delta_B} (\alpha^2 + \beta^2) + r^{3-\delta_B} (\check{\rho})^2 + r^{1-\delta_B} \underline{\beta}^2 + r^{-1-\delta_B} \underline{\alpha}^2 \right), \\ \left((ext)\mathfrak{N}_0^{\leq r_0}[\check{R}] \right)^2 &= \int_{(ext)\mathcal{M}(r \leq 4m_0)} \left(1 - \frac{3m}{r} \right)^2 |\check{R}|^2, \end{aligned}$$

$${}^{(ext)}\mathfrak{R}_0[\check{R}] = {}^{(ext)}\mathfrak{R}_0^{\geq 4m_0}[\check{R}] + {}^{(ext)}\mathfrak{R}_0^{\leq 4m_0}[\check{R}],$$

$$\left({}^{(ext)}\mathfrak{R}_k[\check{R}]\right)^2 = \left({}^{(ext)}\mathfrak{R}_0[\mathfrak{d}^{\leq k}\check{R}]\right)^2 + \int_{{}^{(ext)}\mathcal{M}(r \leq 4m_0)} \left(|\mathfrak{d}^{\leq k-1}\mathbf{N}\check{R}|^2 + |\mathfrak{d}^{\leq k-1}\check{R}|^2\right), \text{ for } k \geq 1,$$

and

$$\begin{aligned} \left({}^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}]\right)^2 &= \int_{\Sigma_*} \left[r^2 \left((\mathfrak{d}^{\leq k}\vartheta)^2 + (\mathfrak{d}^{\leq k}\check{\kappa})^2 + (\mathfrak{d}^{\leq k}\zeta)^2 + (\mathfrak{d}^{\leq k}\check{\underline{\kappa}})^2 \right) + (\mathfrak{d}^{\leq k}\varrho)^2 \right. \\ &\quad \left. + (\mathfrak{d}^{\leq k}\eta)^2 + (\mathfrak{d}^{\leq k}\check{\omega})^2 + (\mathfrak{d}^{\leq k}\underline{\xi})^2 \right] \\ &\quad + \sup_{\lambda \geq 4m_0} \left(\int_{\{r=\lambda\}} \left[\lambda^2 \left((\mathfrak{d}^{\leq k}\vartheta)^2 + (\mathfrak{d}^{\leq k}\check{\kappa})^2 + (\mathfrak{d}^{\leq k}\zeta)^2 \right) \right. \right. \\ &\quad \left. \left. + \lambda^{2-\delta_B} (\mathfrak{d}^{\leq k}\check{\underline{\kappa}})^2 + (\mathfrak{d}^{\leq k}\varrho)^2 + (\mathfrak{d}^{\leq k}\eta)^2 + (\mathfrak{d}^{\leq k}\check{\omega})^2 + \lambda^{-\delta_B} (\mathfrak{d}^{\leq k}\underline{\xi})^2 \right] \right), \\ \left({}^{(ext)}\mathfrak{G}_k^{\leq r_0}[\check{\Gamma}]\right)^2 &= \int_{{}^{(ext)}\mathcal{M}(r \leq 4m_0)} |\mathfrak{d}^{\leq k}(\check{\Gamma})|^2, \\ {}^{(ext)}\mathfrak{G}_k[\check{\Gamma}] &= {}^{(ext)}\mathfrak{G}_k^{\leq 4m_0}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_k^{\geq 4m_0}[\check{\Gamma}]. \end{aligned}$$

Also, we have for ${}^{(int)}\mathcal{M}$

$$\left({}^{(int)}\mathfrak{R}_k[\check{R}]\right)^2 = \int_{{}^{(int)}\mathcal{M}} |\mathfrak{d}^{\leq k}\check{R}|^2,$$

and

$$\left({}^{(int)}\mathfrak{G}_k[\check{\Gamma}]\right)^2 = \int_{{}^{(int)}\mathcal{M}} |\mathfrak{d}^{\leq k}\check{\Gamma}|^2.$$

Finally, we recall the following Morawetz type norms, see section 5.1.4. For $\delta > 0$, we have

$$\begin{aligned} B_\delta[\psi](\tau_1, \tau_2) &= \int_{{}^{(trap)}\mathcal{M}(\tau_1, \tau_2)} |R\psi|^2 + r^{-2}|\psi|^2 + \left(1 - \frac{3m}{r}\right)^2 \left(|\nabla\psi|^2 + \frac{1}{r^2}|T\psi|^2\right) \\ &\quad + \int_{{}^{(trap)}\mathcal{M}(\tau_1, \tau_2)} r^{\delta-3} (|\mathfrak{d}\psi|^2 + |\psi|^2) \end{aligned}$$

where the scalar function τ and the spacetime region ${}^{(trap)}\mathcal{M}$ have been introduced in section 5.1.1, and where ${}^{(trap^c)}\mathcal{M}$ denotes the complement of ${}^{(trap)}\mathcal{M}$. Also, we have

$$\begin{aligned} E_\delta[\psi](\tau) &= \int_{\Sigma(\tau)} \left(\frac{1}{2}(N_\Sigma, e_3)^2 |e_4\psi|^2 + \frac{1}{2}(N_\Sigma, e_4)^2 |e_3\psi|^2 + |\nabla\psi|^2 + r^{-2}|\psi|^2 \right) \\ &\quad + \int_{\Sigma_{\geq 4m_0}(\tau)} r^\delta (|e_4\psi|^2 + r^{-2}|\psi|^2). \end{aligned}$$

Here $\Sigma(\tau)$ denotes the level set of τ , see section 5.1.1, N_Σ denotes a choice for the normal to Σ , and recall that we have

$$N_\Sigma = \begin{cases} N_\Sigma = e_3 & \text{on } {}^{(int)}\Sigma, \\ N_\Sigma = e_4 & \text{on } {}^{(ext)}\Sigma, \end{cases}$$

with ${}^{(int)}\Sigma$ and ${}^{(ext)}\Sigma$ defined in section 5.1.1, and

$$(N_\Sigma, e_3) \leq -1 \text{ and } (N_\Sigma, e_4) \leq -1 \quad \text{on } {}^{(trap)}\Sigma.$$

Moreover, we have

$$\begin{aligned} F_\delta[\psi](\tau_1, \tau_2) &= \int_{\mathcal{A}(\tau_1, \tau_2)} \left(\delta_{\mathcal{H}}^{-1} |e_4\Psi|^2 + \delta_{\mathcal{H}} |e_3\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2 \right) \\ &\quad + \int_{\Sigma_*(\tau_1, \tau_2)} \left(|e_3\Psi|^2 + r^\delta (|e_4\psi|^2 + |\nabla\psi|^2 + r^{-2}|\psi|^2) \right) \end{aligned}$$

with $\mathcal{A}(\tau_1, \tau_2) = \mathcal{A} \cap \mathcal{M}(\tau_1, \tau_2)$ and $\Sigma_*(\tau_1, \tau_2) = \Sigma_* \cap \mathcal{M}(\tau_1, \tau_2)$.

8.3.2 Control of the global frame

Some quantities will be controlled based on the wave equation they satisfy, and will thus need to be defined w.r.t. a global frame, i.e. a smooth frame on \mathcal{M} . To this end, we will rely on the global frame of section 3.5.2. We recall below the main properties of that global frame.

From definition 3.5.1, the region where the frame of ${}^{(int)}\mathcal{M}$ and a conformal renormalization of the frame of ${}^{(ext)}\mathcal{M}$ are matched is given by

$$\text{Match} := \left({}^{(ext)}\mathcal{M} \cap \left\{ {}^{(int)}r \leq 2m_0 \left(1 + \frac{3}{2}\delta_{\mathcal{H}} \right) \right\} \right) \cup \left({}^{(int)}\mathcal{M} \cap \left\{ {}^{(int)}r \geq 2m_0 \left(1 + \frac{1}{2}\delta_{\mathcal{H}} \right) \right\} \right),$$

where ${}^{(int)}r$ denotes the area radius of the ingoing geodesic foliation of ${}^{(int)}\mathcal{M}$ and its extension to ${}^{(ext)}\mathcal{M}$.

The following proposition concerning the global frame is an immediate consequence of Proposition 3.5.2 and the decay estimates (8.3.6).

Proposition 8.3.3. *Assume (8.3.6). Then, there exists a global null frame defined on ${}^{(int)}\mathcal{M} \cup {}^{(ext)}\mathcal{M}$ and denoted by ${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta$ such that*

(a) *In ${}^{(ext)}\mathcal{M} \setminus Match$, we have*

$${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta = ({}^{(ext)}\Upsilon {}^{(ext)}e_4, {}^{(ext)}\Upsilon^{-1}{}^{(ext)}e_3, {}^{(ext)}e_\theta).$$

(b) *In ${}^{(int)}\mathcal{M} \setminus Match$, we have*

$${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta = ({}^{(int)}e_4, {}^{(int)}e_3, {}^{(int)}e_\theta).$$

(c) *In the matching region, we have*

$$\begin{aligned} \max_{0 \leq k \leq k_{small}-2} \sup_{Match \cap {}^{(int)}\mathcal{M}} \underline{u}^{1+\delta_{dec}} |\partial^k ({}^{(glo)}\check{\Gamma}, {}^{(glo)}\check{R})| &\lesssim \epsilon_0, \\ \max_{0 \leq k \leq k_{small}-2} \sup_{Match \cap {}^{(ext)}\mathcal{M}} u^{1+\delta_{dec}} |\partial^k ({}^{(glo)}\check{\Gamma}, {}^{(glo)}\check{R})| &\lesssim \epsilon_0, \end{aligned}$$

where ${}^{(glo)}\check{R}$ and ${}^{(glo)}\check{\Gamma}$ are given by

$$\begin{aligned} {}^{(glo)}\check{R} &= \left\{ \alpha, \beta, \rho + \frac{2m}{r^3}, \underline{\beta}, \underline{\alpha} \right\}, \\ {}^{(glo)}\check{\Gamma} &= \left\{ \xi, \omega + \frac{m}{r^2}, \kappa - \frac{2\Upsilon}{r}, \vartheta, \zeta, \eta, \underline{\eta}, \underline{\kappa} + \frac{2}{r}, \underline{\vartheta}, \underline{\omega}, \underline{\xi} \right\}. \end{aligned}$$

(d) *Furthermore, we may also choose the global frame such that, in addition, one of the following two possibilities hold,*

i. *We have on all ${}^{(ext)}\mathcal{M}$*

$${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta = ({}^{(ext)}\Upsilon {}^{(ext)}e_4, {}^{(ext)}\Upsilon^{-1}{}^{(ext)}e_3, {}^{(ext)}e_\theta).$$

ii. *We have on all ${}^{(int)}\mathcal{M}$*

$${}^{(glo)}e_4, {}^{(glo)}e_3, {}^{(glo)}e_\theta = ({}^{(int)}e_4, {}^{(int)}e_3, {}^{(int)}e_\theta).$$

8.3.3 Iterative procedure

Recall our norms for measuring energies for curvature components and Ricci coefficients which are given respectively by ${}^{(int)}\mathfrak{R}_k[\check{R}]$, ${}^{(ext)}\mathfrak{R}_k[\check{R}]$ and ${}^{(int)}\mathfrak{G}_k[\check{\Gamma}]$, ${}^{(ext)}\mathfrak{G}_k[\check{\Gamma}]$, see sections 3.2.1 and 3.2.2. Recall also our combined weighted energy norm

$$\mathfrak{N}_k^{(En)} = {}^{(ext)}\mathfrak{R}_k[\check{R}] + {}^{(ext)}\mathfrak{G}_k[\check{\Gamma}] + {}^{(int)}\mathfrak{R}_k[\check{R}] + {}^{(int)}\mathfrak{G}_k[\check{\Gamma}].$$

We also introduce the following norm controlling on the matching region the Ricci coefficients and curvature components of the global frame of Proposition 8.3.3

$$\mathcal{N}_k^{(match)} := \left(\int_{Match} |\mathfrak{d}^{\leq k}({}^{(glo)}\check{\Gamma}, {}^{(glo)}\check{R})|^2 \right)^{\frac{1}{2}}. \quad (8.3.8)$$

The estimate (8.3.6) and Proposition 8.3.3 imply in particular

$$\mathfrak{N}_{k_{small}}^{(En)} + \mathcal{N}_{k_{small}-2}^{(match)} \lesssim \epsilon_0. \quad (8.3.9)$$

Next, for J such that $k_{small} - 2 \leq J \leq k_{large} - 1$, consider the iteration assumption

$$\mathfrak{N}_J^{(En)} + \mathcal{N}_J^{(match)} \lesssim \epsilon_{\mathcal{B}}[J], \quad (8.3.10)$$

where

$$\epsilon_{\mathcal{B}}[J] := \sum_{j=k_{small}-2}^J (\epsilon_0)^{\ell(j)} \mathcal{B}^{1-\ell(j)} + \epsilon_0^{\ell(J)} \mathcal{B}, \quad \ell(j) := 2^{k_{small}-2-j}, \quad (8.3.11)$$

$$\mathcal{B} := \left(\int_{{}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 \right)^{\frac{1}{2}}. \quad (8.3.12)$$

Lemma 8.3.4. *The following estimate holds true for $\epsilon_{\mathcal{B}}[J]$ as defined above*

$$\epsilon_{\mathcal{B}}[J] + \mathcal{B}^{\frac{1}{2}} (\epsilon_{\mathcal{B}}[J])^{\frac{1}{2}} + \epsilon_0 \mathcal{B} \lesssim \epsilon_{\mathcal{B}}[J+1]. \quad (8.3.13)$$

Proof. We clearly have

$$\epsilon_{\mathcal{B}}[J] + \epsilon_0 \mathcal{B} \lesssim \epsilon_{\mathcal{B}}[J+1]. \quad (8.3.14)$$

Also, we have, using $\ell(j) = 2\ell(j+1)$,

$$\begin{aligned}
\mathcal{B}\epsilon_{\mathcal{B}}[J] &\lesssim \sum_{j=k_{small}-2}^J (\epsilon_0)^{\ell(j)} \mathcal{B}^{2-\ell(j)} + \epsilon_0^{\ell(J)} \mathcal{B}^2 \\
&\lesssim \sum_{j=k_{small}-1}^{J+1} (\epsilon_0)^{2\ell(j)} \mathcal{B}^{2-2\ell(j)} + \epsilon_0^{2\ell(J+1)} \mathcal{B}^2 \\
&\lesssim \left(\sum_{j=k_{small}-2}^{J+1} (\epsilon_0)^{\ell(j)} \mathcal{B}^{1-\ell(j)} + \epsilon_0^{\ell(J+1)} \mathcal{B} \right)^2 \\
&= (\epsilon_{\mathcal{B}}[J+1])^2
\end{aligned}$$

which concludes the proof of the lemma. \square

In view of (8.3.9), (8.3.10) holds for $J = k_{small} - 2$. The propositions below will allow us to prove Theorem M8 in the next section.

Proposition 8.3.5. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Consider the global frame constructed in Proposition 8.3.3. In that frame, let*

$$\tilde{\rho} := r^2 \rho + 2mr^{-1}. \quad (8.3.15)$$

Then, under the iteration assumption (8.3.10), we have

$$\sup_{\tau \in [1, \tau_*]} E_{\delta}^J[\tilde{\rho}](\tau) + B_{\delta}^J[\tilde{\rho}](1, \tau_*) + F_{\delta}^J[\tilde{\rho}](1, \tau_*) \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Proposition 8.3.6. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Consider the global frame constructed in Proposition 8.3.3. In that frame, under the iteration assumption (8.3.10), we have*

$$\begin{aligned}
&\sup_{\tau \in [1, \tau_*]} E_{\delta}^J[\alpha + \Upsilon^2 \underline{\alpha}](\tau) + B_{\delta}^J[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) + F_{\delta}^J[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) \\
&\lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.
\end{aligned}$$

Proposition 8.3.7. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Consider the global frame constructed in Proposition 8.3.3. In that frame, under the iteration assumption (8.3.10), we have*

$$B_{-2}^J[\check{\rho}, \alpha, \underline{\alpha}, \beta, \underline{\beta}](1, \tau_*) \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Proposition 8.3.8. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Under the iteration assumption (8.3.10), we have for $r_0 \geq 4m_0$*

$${}^{(int)}\mathfrak{N}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{N}_{J+1}[\check{R}] \leq {}^{(ext)}\mathfrak{N}_{J+1}^{\geq r_0}[\check{R}] + O\left(r_0^{10} \left(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)}\right)\right)\right)$$

and

$${}^{(ext)}\mathfrak{N}_{J+1}^{\geq r_0}[\check{R}] \lesssim r_0^{-\delta_B} {}^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}] + r_0^{10} \left(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)}\right)\right).$$

Proposition 8.3.9. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Under the iteration assumption (8.3.10), we have*

$${}^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{N}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{N}_{J+1}[\check{R}] \lesssim \epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)}\right).$$

Proposition 8.3.10. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Under the iteration assumption (8.3.10), we have*

$${}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] \lesssim \epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)}\right) + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2\right)^{\frac{1}{2}}.$$

Proposition 8.3.11. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Under the iteration assumption (8.3.10), we have*

$$\mathcal{N}_{J+1}^{(match)} \lesssim \mathfrak{N}_{J+1}^{(En)} + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2\right)^{\frac{1}{2}}.$$

The proof of Propositions 8.3.5, 8.3.6, 8.3.7, 8.3.8, 8.3.9, 8.3.10 and 8.3.11 are postponed respectively to sections 8.4, 8.5, 8.6, 8.7, 8.8, 8.9 and 8.10.

8.3.4 End of the proof of Theorem M8

To prove Theorem M8, we rely on Propositions 8.3.9, 8.3.10 and 8.3.11. Note that among them only the second two involve the dangerous boundary term $\left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2\right)^{\frac{1}{2}}$. We proceed as follows.

Step 1. As mentioned earlier, the estimate (8.3.9) trivially implies the iteration assumption (8.3.10) with $J = k_{small} - 2$. We assume that the iteration assumption (8.3.10) holds for any fixed J such that $k_{small} - 2 \leq J \leq k_{large} - 2$. In view of Proposition 8.3.10, we have

$${}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] \lesssim \epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)}\right) + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2\right)^{\frac{1}{2}}. \quad (8.3.16)$$

We need to deal with the last term in the RHS of (8.3.16). Relying on a trace theorem in the spacetime region ${}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})$, and the fact that $J + 2 \leq k_{large}$, we obtain

$$\left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}} \lesssim \left(\int_{{}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\mathfrak{d}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{4}} ({}^{(ext)}\mathfrak{R}_{J+1}[\check{R}])^{\frac{1}{2}} + {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}]. \quad (8.3.17)$$

Proposition 8.3.9, (8.3.16) and (8.3.17) yield, for $\epsilon_0 > 0$ small enough so that we can absorb some of the terms to the left,

$$\mathfrak{N}_{J+1}^{(En)} \lesssim \epsilon_{\mathcal{B}}[J] + \left(\int_{{}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\mathfrak{d}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{4}} \left(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right)^{\frac{1}{2}} + \epsilon_0 \mathcal{N}_{J+1}^{(match)},$$

and using also Proposition 8.3.11,

$$\begin{aligned} \mathcal{N}_{J+1}^{(match)} &\lesssim \mathfrak{N}_{J+1}^{(En)} + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}} \\ &\lesssim \epsilon_{\mathcal{B}}[J] + \left(\int_{{}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\mathfrak{d}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{4}} \left(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right)^{\frac{1}{2}} + \epsilon_0 \mathcal{N}_{J+1}^{(match)}. \end{aligned}$$

For $\epsilon_0 > 0$ small enough, we infer, by absorbing the appropriate terms to the left,

$$\begin{aligned} &\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \\ &\lesssim \epsilon_{\mathcal{B}}[J] + \left(\int_{{}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\mathfrak{d}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{4}} \left(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right)^{\frac{1}{2}} \\ &\lesssim \epsilon_{\mathcal{B}}[J] + \left(\int_{{}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\mathfrak{d}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{4}} (\epsilon_{\mathcal{B}}[J])^{\frac{1}{2}} \\ &\quad + \left(\int_{{}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}})} |\mathfrak{d}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{4}} \left(\epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right)^{\frac{1}{2}} \end{aligned}$$

and hence

$$\begin{aligned} \mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} &\lesssim \epsilon_{\mathcal{B}}[J] + \left(\int_{(ext)\mathcal{M}}(r \in I_{m_0, \delta_{\mathcal{H}}}) |\mathfrak{d}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{4}} \left(\epsilon_{\mathcal{B}}[J] \right)^{\frac{1}{2}} \\ &\quad + \epsilon_0 \left(\int_{(ext)\mathcal{M}}(r \in I_{m_0, \delta_{\mathcal{H}}}) |\mathfrak{d}^{k_{large}} \check{R}|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

In view of Lemma 8.3.4, we deduce

$$\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \lesssim \epsilon_{\mathcal{B}}[J+1]$$

which is (8.3.10) for $J+1$ derivatives. We deduce that (8.3.10) holds for all $J \leq k_{large} - 1$, and hence

$$\mathfrak{N}_{k_{large}-1}^{(En)} + \mathcal{N}_{k_{large}-1}^{(match)} \lesssim \epsilon_{\mathcal{B}}[k_{large} - 1]. \quad (8.3.18)$$

Step 2. Next, Proposition 8.3.9 implies in view of (8.3.18)

$$\begin{aligned} (ext)\mathfrak{G}_{k_{large}}[\check{\Gamma}] + (int)\mathfrak{A}_{k_{large}}[\check{R}] + (ext)\mathfrak{A}_{k_{large}}[\check{R}] &\lesssim \epsilon_{\mathcal{B}}[k_{large} - 1] \\ &\quad + \epsilon_0 \left(\mathfrak{N}_{k_{large}}^{(En)} + \mathcal{N}_{k_{large}}^{(match)} \right). \end{aligned} \quad (8.3.19)$$

In particular, we have

$$\left(\int_{(ext)\mathcal{M}}(r \in I_{m_0, \delta_{\mathcal{H}}}) |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 \right)^{\frac{1}{2}} \leq (ext)\mathfrak{A}_{k_{large}}[\check{R}] \lesssim \epsilon_{\mathcal{B}}[k_{large} - 1] + \epsilon_0 \left(\mathfrak{N}_{k_{large}}^{(En)} + \mathcal{N}_{k_{large}}^{(match)} \right).$$

In view of the definition of $\epsilon_{\mathcal{B}}[k_{large} - 1]$, we infer for $\epsilon_0 > 0$ small enough

$$\left(\int_{(ext)\mathcal{M}}(r \in I_{m_0, \delta_{\mathcal{H}}}) |\mathfrak{d}^{\leq k_{large}} \check{R}|^2 \right)^{\frac{1}{2}} \lesssim \epsilon_0 + \epsilon_0 \left(\mathfrak{N}_{k_{large}}^{(En)} + \mathcal{N}_{k_{large}}^{(match)} \right)$$

and hence

$$\epsilon_{\mathcal{B}}[k_{large} - 1] \lesssim \epsilon_0 + \epsilon_0 \left(\mathfrak{N}_{k_{large}}^{(En)} + \mathcal{N}_{k_{large}}^{(match)} \right)$$

which yields, together with (8.3.19),

$$(ext)\mathfrak{G}_{k_{large}}[\check{\Gamma}] + (int)\mathfrak{A}_{k_{large}}[\check{R}] + (ext)\mathfrak{A}_{k_{large}}[\check{R}] \lesssim \epsilon_0 + \epsilon_0 \left(\mathfrak{N}_{k_{large}}^{(En)} + \mathcal{N}_{k_{large}}^{(match)} \right). \quad (8.3.20)$$

Step 3. Next, Proposition 8.3.10 implies in view of (8.3.20),

$${}^{(int)}\mathfrak{G}_{k_{large}}[\check{\Gamma}] \lesssim \epsilon_0 + \epsilon_0 \left(\mathfrak{N}_{k_{large}}^{(En)} + \mathcal{N}_{k_{large}}^{(match)} \right) + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}$$

and hence, for $\epsilon_0 > 0$ small enough, using again (8.3.20),

$$\mathfrak{N}_{k_{large}}^{(En)} \lesssim \epsilon_0 + \epsilon_0 \mathcal{N}_{k_{large}}^{(match)} + \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}.$$

Together with Proposition 8.3.11, we infer for $\epsilon_0 > 0$ small enough

$$\mathfrak{N}_{k_{large}}^{(En)} + \mathcal{N}_{k_{large}}^{(match)} \lesssim \epsilon_0 + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}.$$

Step 4. It remains to estimate the last term of the RHS of the previous inequality. Now, in view of (8.3.7) and (8.3.20), we have

$$\begin{aligned} \left(\int_{\mathcal{T}} |\mathfrak{d}^{k_{large}}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}} &\lesssim \left(\int_{({}^{(ext)}\mathcal{M}(r \in I_{m_0, \delta_{\mathcal{H}}}))} |\mathfrak{d}^{\leq k_{large}}\check{R}|^2 \right)^{\frac{1}{2}} \\ &\lesssim {}^{(ext)}\mathfrak{N}_{k_{large}}[\check{R}] \\ &\lesssim \epsilon_0 + \epsilon_0 \mathfrak{N}_{k_{large}}^{(En)} \end{aligned}$$

so that we finally obtain, for $\epsilon_0 > 0$ small enough,

$$\mathfrak{N}_{k_{large}}^{(En)} \lesssim \epsilon_0.$$

This concludes the proof of Theorem M8.

8.4 Proof of Proposition 8.3.5

8.4.1 A wave equation for $\tilde{\rho}$

Proposition 8.4.1. *The following wave equations hold true.*

1. *The curvature component ρ verifies the identity*

$$\square_{\mathbf{g}}\rho = \underline{\kappa}e_4\rho + \kappa e_3\rho + \frac{3}{2}(\underline{\kappa}\kappa + 2\rho)\rho + Err[\square_{\mathbf{g}}\rho],$$

where

$$\begin{aligned}
Err[\square_{\mathbf{g}}\rho] &:= \frac{3}{2}\rho \left(-\frac{1}{2}\vartheta\vartheta + 2(\underline{\xi}\xi + \eta\eta) \right) + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) \left(\frac{1}{2}\vartheta\alpha - \zeta\beta - 2(\underline{\eta}\beta + \xi\underline{\beta}) \right) \\
&\quad - \frac{1}{2}\vartheta\phi_2^*\beta + (\zeta - \eta)e_3\beta - \eta e_3(\Phi)\beta - \underline{\xi}(e_4\beta + e_4(\Phi)\beta) - \underline{\beta}\beta \\
&\quad - e_3 \left(-\frac{1}{2}\vartheta\alpha + \zeta\beta + 2(\underline{\eta}\beta + \xi\underline{\beta}) \right) \\
&\quad - \phi_1^*(\underline{\kappa})\beta + 2\phi_1^*(\underline{\omega})\beta + 3\eta\phi_1^*(\rho) - \phi_1 \left(-\vartheta\underline{\beta} + \underline{\xi}\alpha \right) - 2\eta e_\theta\rho.
\end{aligned}$$

2. The small curvature quantity,

$$\tilde{\rho} := r^2 \left(\rho + \frac{2m}{r^3} \right)$$

verifies the wave equation,

$$\begin{aligned}
\square_{\mathbf{g}}(\tilde{\rho}) + \frac{8m}{r^3}\tilde{\rho} &= -6m \frac{\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right)}{r^2} - \frac{3m}{r} \left(\kappa\underline{\kappa} + \frac{4\underline{\Upsilon}}{r^2} \right) \\
&\quad - \frac{3m}{r} (A\underline{\kappa} + \underline{A}\kappa) + Err[\square_g\tilde{\rho}],
\end{aligned}$$

where

$$\begin{aligned}
Err[\square_g\tilde{\rho}] &:= -\frac{6m}{r}A\underline{A} + \frac{3}{r^2}\tilde{\rho}^2 + \frac{3}{2} \left(\frac{4}{3}A \frac{e_3(r)}{r} + \frac{4}{3}\underline{A} \frac{e_4(r)}{r} \right) \tilde{\rho} \\
&\quad + \left(\frac{3}{2} \left(\kappa\underline{\kappa} - \frac{8m}{r^3} + \frac{2}{3r^2}\square_{\mathbf{g}}(r^2) \right) + \frac{8m}{r^3} \right) \tilde{\rho} \\
&\quad - Ae_3(\tilde{\rho}) - \underline{A}e_4(\tilde{\rho}) + \frac{2}{r}Ae_3(m) + \frac{2}{r}\underline{A}e_4(m) \\
&\quad + 4D^a(m)D_a \left(\frac{1}{r} \right) + \frac{2}{r}\square_{\mathbf{g}}(m) + 4r\phi_1^*(r)\phi_1^*(\rho) + r^2 Err[\square_{\mathbf{g}}\rho],
\end{aligned}$$

and where we recall that,

$$A = \frac{2}{r}e_4(r) - \kappa, \quad \underline{A} = \frac{2}{r}e_3(r) - \underline{\kappa}.$$

Proof. See appendix B.1. □

8.4.2 Control of $\square_{\mathbf{g}}(r)$

Lemma 8.4.2. *Let r the function on \mathcal{M} associated to the global frame constructed in Proposition 8.3.3, see definition 4.6.4. Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Under the iteration assumption (8.3.10), we have*

$$\int_{(int)\mathcal{M} \cup (ext)\mathcal{M}(r \leq 4m_0)} \left(\mathfrak{d}^J \left(\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right) \right) \right)^2 + \sup_{r_0 \geq 4m_0} \int_{\{r=r_0\}} \left(\mathfrak{d}^J \left(\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right) \right) \right)^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2$$

and

$$\int_{(trap)\mathcal{M}} \left(\mathfrak{d}^J e_4 \left(\square_{\mathbf{g}}((ext)r) - \left(\frac{2}{(ext)r} - \frac{2^{(ext)}m}{((ext)r)^2} \right) \right) \right)^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Proof. Recall that, according to definition 4.6.4, r is defined on $(ext)\mathcal{M} \cup (int)\mathcal{M}$ as follows

- on $(ext)\mathcal{M} \setminus \text{Match}$, we have

$$(glo)_r = (ext)_r,$$

- on $(int)\mathcal{M} \setminus \text{Match}$, we have

$$(glo)_r = (int)_r,$$

- on the matching region, we have

$$(glo)_r = (1 - \psi_{m_0, \delta_{\mathcal{H}}}(^{(int)}r))^{(int)}r + \psi_{m_0, \delta_{\mathcal{H}}}(^{(int)}r)^{(ext)}r,$$

where the matching region of Proposition 8.3.3 is given by

$$\text{Match} := \left((ext)\mathcal{M} \cap \left\{ (int)r \leq 2m_0 \left(1 + \frac{3}{2}\delta_{\mathcal{H}} \right) \right\} \right) \cup \left((int)\mathcal{M} \cap \left\{ (int)r \geq 2m_0 \left(1 + \frac{1}{2}\delta_{\mathcal{H}} \right) \right\} \right),$$

and where $\psi_{m_0, \delta_{\mathcal{H}}}$ is given by

$$\psi_{m_0, \delta_{\mathcal{H}}}(r) = \psi \left(\frac{r - 2m_0 \left(1 + \frac{1}{2}\delta_{\mathcal{H}} \right)}{2m_0 \delta_{\mathcal{H}}} \right) \text{ on } 2m_0 \left(1 + \frac{1}{2}\delta_{\mathcal{H}} \right) \leq r \leq 2m_0 \left(1 + \frac{3}{2}\delta_{\mathcal{H}} \right).$$

with $\psi : \mathbb{R} \rightarrow \mathbb{R}$ a smooth cut-off function such that $0 \leq \psi \leq 1$, $\psi = 0$ on $(-\infty, 0]$ and $\psi = 1$ on $[1, +\infty)$.

We have on ${}^{(ext)}\mathcal{M}$

$$\square_{\mathbf{g}}({}^{(ext)}r) = -e_3 e_4({}^{(ext)}r) + \underline{\Delta}({}^{(ext)}r) + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) e_4({}^{(ext)}r) - \frac{1}{2}\kappa e_3({}^{(ext)}r) + 2\eta e_\theta({}^{(ext)}r).$$

Here, (e_4, e_3, e_θ) denotes the frame of ${}^{(ext)}\mathcal{M}$ and the Ricci coefficients are computed w.r.t. frame, so we have

$$e_4({}^{(ext)}r) = \frac{{}^{(ext)}r}{2}\bar{\kappa}, \quad e_3({}^{(ext)}r) = \frac{{}^{(ext)}r}{2}(\bar{\kappa} + \underline{A}), \quad e_\theta({}^{(ext)}r) = 0$$

and hence

$$\begin{aligned} \square_{\mathbf{g}}({}^{(ext)}r) &= -e_3 \left(\frac{{}^{(ext)}r}{2}\bar{\kappa} \right) + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa} \right) \frac{{}^{(ext)}r}{2}\bar{\kappa} - \frac{1}{2}\kappa \frac{{}^{(ext)}r}{2}(\bar{\kappa} + \underline{A}) \\ &= -\frac{{}^{(ext)}r}{2}e_3(\bar{\kappa}) - \frac{e_3({}^{(ext)}r)}{2}\bar{\kappa} + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa} \right) \frac{{}^{(ext)}r}{2}\bar{\kappa} - \frac{{}^{(ext)}r}{4}\kappa\bar{\kappa} - \frac{{}^{(ext)}r}{4}\kappa\underline{A} \\ &= -\frac{{}^{(ext)}r}{2}e_3(\bar{\kappa}) - \frac{1}{2}\bar{\kappa} \frac{{}^{(ext)}r}{2}(\bar{\kappa} + \underline{A}) + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa} \right) \frac{{}^{(ext)}r}{2}\bar{\kappa} - \frac{{}^{(ext)}r}{4}\kappa\bar{\kappa} - \frac{{}^{(ext)}r}{4}\kappa\underline{A}. \end{aligned}$$

Now, we have

$$\begin{aligned} e_3(\bar{\kappa}) &= \overline{e_3(\kappa)} + \text{Err}[e_3\bar{\kappa}] \\ &= \overline{-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa + 2\rho + 2\underline{\theta}_1\eta - \frac{1}{2}\vartheta\underline{\vartheta} + 2\eta^2} + \text{Err}[e_3\bar{\kappa}] \\ &= \overline{-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\eta^2} + \text{Err}[e_3\bar{\kappa}] \end{aligned}$$

and hence

$$\begin{aligned} \square_{\mathbf{g}}({}^{(ext)}r) &= -\frac{{}^{(ext)}r}{2} \left(\overline{-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\eta^2} + \text{Err}[e_3\bar{\kappa}] \right) \\ &\quad - \frac{1}{2}\bar{\kappa} \frac{{}^{(ext)}r}{2}(\bar{\kappa} + \underline{A}) + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa} \right) \frac{{}^{(ext)}r}{2}\bar{\kappa} - \frac{{}^{(ext)}r}{4}\kappa\bar{\kappa} - \frac{{}^{(ext)}r}{4}\kappa\underline{A}. \end{aligned}$$

Together with (8.3.4) and the iteration assumption (8.3.10), we easily infer⁴

$$\begin{aligned} & \int_{(ext)\mathcal{M}(r \leq 4m_0)} \left(\mathfrak{d}^J \left(\square_{\mathbf{g}}((ext)_r) - \left(\frac{2}{(ext)_r} - \frac{2^{(ext)m}}{(ext)_r^2} \right) \right) \right)^2 \\ & + \sup_{r_0 \geq 4m_0} \int_{\{r=r_0\}} \left(\mathfrak{d}^J \left(\square_{\mathbf{g}}((ext)_r) - \left(\frac{2}{(ext)_r} - \frac{2^{(ext)m}}{(ext)_r^2} \right) \right) \right)^2 \\ & \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2. \end{aligned} \quad (8.4.1)$$

Also, using again (8.3.4) and the iteration assumption (8.3.10), we have

$$\begin{aligned} & \int_{(trap)\mathcal{M}} \left(\mathfrak{d}^J e_4 \left(\square_{\mathbf{g}}((ext)_r) - \left(\frac{2}{(ext)_r} - \frac{2^{(ext)m}}{(ext)_r^2} \right) \right) \right)^2 \\ & \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2, \end{aligned} \quad (8.4.2)$$

where we have used the null structure equations for $e_4(\underline{\kappa})$, $e_4(\underline{\kappa})$, $e_4(\underline{\omega})$, $e_4(\underline{\vartheta})$, $e_4(\underline{\vartheta})$, $e_4(\underline{\eta})$, the equations for $e_4(\underline{\Omega})$, $e_4(\underline{\varsigma})$, $e_4(r)$, and the Bianchi identity for $e_4(\rho)$.

Remark 8.4.3. *Note that we have used in the last estimate the following observations to avoid a potential loss of one derivative*

$$\begin{aligned} e_4(\underline{\kappa}) &= -2\mathfrak{d}_1\zeta + \dots = 2 \left(\rho + \mu - \frac{1}{4}\vartheta\underline{\vartheta} \right) + \dots, \\ \overline{e_4(\rho)} &= \overline{\mathfrak{d}_1\beta} + \dots = \dots, \\ e_4(\text{Err}[e_3\overline{\kappa}]) &= 2e_4(\zeta^{-1}\overline{\zeta}\mathfrak{d}_1\eta) + \dots = 2\zeta^{-1}\overline{\zeta}\mathfrak{d}_1 e_4\eta + \dots = -2\zeta^{-1}e_\theta(\zeta)e_4\eta + \dots \end{aligned}$$

Note also that there is no term involving $\mathfrak{d}^J\rho$ (without average) as such a term appears only in the null structure equations for $e_4(\underline{\kappa})$, as well as $e_4(\underline{\omega})$ and vanishes due to the cancellation

$$\begin{aligned} e_4 \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa} \right) &= 2e_4(\underline{\omega}) - \frac{1}{2}e_4(\underline{\kappa}) \\ &= 2\rho + \dots - \frac{1}{2}(-2\mathfrak{d}_1\zeta + 2\rho) + \dots \\ &= 2\mu + \dots \end{aligned}$$

This is important as such a term would otherwise violate (8.4.2) at $r = 3m$.

Remark 8.4.4. *Recall that the global frame constructed in Proposition 8.3.3*

⁴Recall in particular that $\overline{\rho}$ is under control in view of Lemma 3.4.1.

- coincides with the frame of ${}^{(int)}\mathcal{M}$ in ${}^{(int)}\mathcal{M} \setminus Match$,
- coincides with a conformal renormalization of the frame of ${}^{(ext)}\mathcal{M}$ in ${}^{(ext)}\mathcal{M} \setminus Match$.

Thus, $J + 1$ derivatives of its Ricci coefficients and curvature components are controlled

- by $\mathcal{N}_{J+1}^{(match)}$ in $Match$,
- by $\mathfrak{N}_{J+1}^{(En)}$ in $\mathcal{M} \setminus Match$,

and hence by $\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)}$ on \mathcal{M} . This explains the occurrence of $\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)}$ on the right-hand side of numerous estimates, see for example (8.4.1) (8.4.2).

Arguing similarly for ${}^{(int)}r$, we obtain the following analog of (8.4.1)

$$\int_{{}^{(int)}\mathcal{M}} \left(\mathfrak{d}^J \left(\square_{\mathbf{g}}({}^{(int)}r) - \left(\frac{2}{{}^{(int)}r} - \frac{2({}^{(int)}m)}{({}^{(int)}r)^2} \right) \right) \right)^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2. \quad (8.4.3)$$

Then, since

- on ${}^{(ext)}\mathcal{M} \setminus Match$, we have

$$\square_{\mathbf{g}}(r) = \square_{\mathbf{g}}({}^{(ext)}r), \quad m = {}^{(ext)}m,$$

- on ${}^{(int)}\mathcal{M} \setminus Match$, we have

$$\square_{\mathbf{g}}(r) = \square_{\mathbf{g}}({}^{(int)}r), \quad m = {}^{(int)}m,$$

we immediately infer from (8.4.1), (8.4.2) and (8.4.3)

$$\begin{aligned} & \int_{\left({}^{(int)}\mathcal{M} \cup {}^{(ext)}\mathcal{M}(r \leq 4m_0) \right) \setminus Match} \left(\mathfrak{d}^J \left(\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right) \right) \right)^2 \\ & + \sup_{r_0 \geq 4m_0} \int_{\{r=r_0\}} \left(\mathfrak{d}^J \left(\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right) \right) \right)^2 \\ & \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \end{aligned}$$

and

$$\int_{(trap)\mathcal{M}} \left(\mathfrak{d}^J e_4 \left(\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right) \right) \right)^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2$$

which are the desired estimates outside of the matching region. Note that we have used the fact that $(trap)\mathcal{M} \cap Match = \emptyset$.

It remains to derive the desired estimates in the matching region. To this end, we need to estimate $(ext)_r - (int)_r$ and $(int)m - (ext)m$ in the matching region. Step 7 or the proof of Lemma 4.6.6 in section 4.6.2 yields⁵

$$\int_{(int)\mathcal{M}} \left(\mathfrak{d}^{J+1} \left((ext)_r - (int)_r, (ext)m - (int)m \right) \right)^2 \lesssim (\mathfrak{N}_J^{(En)})^2 + (\mathcal{N}_J^{(match)})^2.$$

We infer, in view of the iteration assumption (8.3.10),

$$\int_{(int)\mathcal{M}} \left(\mathfrak{d}^{J+1} \left((ext)_r - (int)_r, (ext)m - (int)m \right) \right)^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2. \quad (8.4.4)$$

Then, since we have on the matching region,

$$\begin{aligned} r &= (1 - \psi_{m_0, \delta_{\mathcal{H}}}((int)_r)) (int)_r + \psi_{m_0, \delta_{\mathcal{H}}}((int)_r) (ext)_r, \\ m &= (1 - \psi_{m_0, \delta_{\mathcal{H}}}((int)_r)) (int)m + \psi_{m_0, \delta_{\mathcal{H}}}((int)_r) (ext)m, \\ \square_{\mathbf{g}}(r) &= (1 - \psi_{m_0, \delta_{\mathcal{H}}}((int)_r)) \square_{\mathbf{g}}((int)_r) + \psi_{m_0, \delta_{\mathcal{H}}}((int)_r) \square_{\mathbf{g}}((ext)_r) \\ &\quad + 2\psi'_{m_0, \delta_{\mathcal{H}}}((int)_r) \mathbf{D}^{\alpha}((int)_r) \mathbf{D}_{\alpha}((ext)_r - (int)_r) \\ &\quad + ((ext)_r - (int)_r) \square_{\mathbf{g}}(\psi_{m_0, \delta_{\mathcal{H}}}), \end{aligned}$$

we deduce there

$$\begin{aligned} &\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right) \\ &= (1 - \psi_{m_0, \delta_{\mathcal{H}}}((int)_r)) \left(\square_{\mathbf{g}}((int)_r) - \left(\frac{2}{(int)_r} - \frac{2(int)m}{((int)_r)^2} \right) \right) \\ &\quad + \psi_{m_0, \delta_{\mathcal{H}}}((int)_r) \left(\square_{\mathbf{g}}((ext)_r) - \left(\frac{2}{(ext)_r} - \frac{2(ext)m}{((ext)_r)^2} \right) \right) \\ &\quad + (1 - \psi_{m_0, \delta_{\mathcal{H}}}((int)_r)) \left(\frac{2}{(int)_r} - \frac{2}{r} - \frac{2(int)m}{((int)_r)^2} + \frac{2m}{r^2} \right) \\ &\quad + \psi_{m_0, \delta_{\mathcal{H}}}((int)_r) \left(\frac{2}{(ext)_r} - \frac{2}{r} - \frac{2(ext)m}{((ext)_r)^2} + \frac{2m}{r^2} \right) \\ &\quad + 2\psi'_{m_0, \delta_{\mathcal{H}}}((int)_r) \mathbf{D}^{\alpha}((int)_r) \mathbf{D}_{\alpha}((ext)_r - (int)_r) + ((ext)_r - (int)_r) \square_{\mathbf{g}}(\psi_{m_0, \delta_{\mathcal{H}}}) \end{aligned}$$

⁵The proof of Lemma 4.6.6 in section 4.6.2 is done in the particular case $J = k_{large} - 1$ but extends immediately to the case $k_{small} - 2 \leq J \leq k_{large} - 1$.

and thus, in view of (8.4.1), (8.4.3) and (8.4.4), we have on the matching region

$$\int_{\text{Match}} \left(\mathfrak{d}^J \left(\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right) \right) \right)^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2$$

as desired. This concludes the proof of the lemma. \square

Corollary 8.4.5. *Let N_0 the RHS of the wave equation for $\tilde{\rho}$ provided by Proposition 8.4.1, i.e.*

$$N_0 = -6m \frac{\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right)}{r^2} - \frac{3m}{r} \left(\kappa_{\underline{\kappa}} + \frac{4\Upsilon}{r^2} \right) - \frac{3m}{r} (A_{\underline{\kappa}} + \underline{A}_{\kappa}) + \text{Err}[\square_g \tilde{\rho}].$$

Then, $N_0 - \text{Err}[\square_g \tilde{\rho}]$ satisfies

$$\begin{aligned} & \int_{(int)\mathcal{M} \cup (ext)\mathcal{M}(r \leq 4m_0)} (\mathfrak{d}^J (N_0 - \text{Err}[\square_g \tilde{\rho}]))^2 \\ & + \sup_{r_0 \geq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^J (N_0 - \text{Err}[\square_g \tilde{\rho}]))^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \end{aligned}$$

and

$$\mathfrak{d}^J e_4 \left(N_0 - \text{Err}[\square_g \tilde{\rho}] \right) = -\frac{12m\kappa}{r} \mathfrak{d}^J \rho + a_J \text{ on } (trap)\mathcal{M}$$

where a^J satisfies

$$\int_{(trap)\mathcal{M}} (\mathfrak{d}^J e_4 a^J)^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Proof. The first estimate is an immediate consequence of Lemma 8.4.2, (8.3.4) and the iteration assumption (8.3.10).

Concerning the second estimate, note that the term $\mathfrak{d}^J \rho$ is due to the null structure equations for $e_4(\underline{\kappa})$, i.e.

$$\begin{aligned} e_4(\underline{\kappa}) &= -2\mathfrak{d}_1 \zeta + 2\rho + \dots \\ &= 4\rho + \dots \end{aligned}$$

Then, the estimate for a_J follows from Lemma 8.4.2, (8.3.4) and the iteration assumption (8.3.10). \square

8.4.3 End of the proof of Proposition 8.3.5

In view of Proposition 8.4.1, $\tilde{\rho}$ satisfies

$$(\square_0 + V_0)\tilde{\rho} = N_0, \quad V_0 = \frac{8m}{r^3},$$

where

$$N_0 := -6m \frac{\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2}\right)}{r^2} - \frac{3m}{r} \left(\kappa \underline{\kappa} + \frac{4\Upsilon}{r^2} \right) - \frac{3m}{r} (A_{\underline{\kappa}} + \underline{A}\kappa) + \text{Err}[\square_g \tilde{\rho}].$$

We may thus apply the estimate (10.5.2) of Theorem 10.5.2 with $\phi = \tilde{\rho}$ and $s = J$ to obtain for any $k_{small} \leq J \leq k_{large} - 1$

$$\begin{aligned} & \sup_{\tau \in [1, \tau_*]} E_\delta^J[\tilde{\rho}](\tau) + B_\delta^J[\tilde{\rho}](1, \tau_*) + F_\delta^J[\tilde{\rho}](1, \tau_*) \\ & \lesssim E_\delta^J[\tilde{\rho}](1) + \sup_{\tau \in [1, \tau_*]} E_\delta^{J-1}[\tilde{\rho}](\tau) + B_\delta^{J-1}[\tilde{\rho}](1, \tau_*) + F_\delta^{J-1}[\tilde{\rho}](1, \tau_*) \\ & \quad + D_J[\Gamma] \left(\sup_{\mathcal{M}} r u_{trap}^{\frac{1}{2} + \delta_{dec}} |\mathfrak{d}^{\leq k_{small}} \tilde{\rho}| \right)^2 + \int_{\Sigma(\tau_*)} \frac{(\mathfrak{d}^{\leq J} \tilde{\rho})^2}{r^3} \\ & \quad + \int_{\mathcal{M}} r^{1+\delta} |\mathfrak{d}^{\leq J} N_0|^2 + \left| \int_{(trap)\mathcal{M}} T(\mathfrak{d}^J \tilde{\rho}) \mathfrak{d}^J N_0 \right|, \end{aligned}$$

where $D_J[\Gamma]$ is defined by

$$\begin{aligned} D_J[\Gamma] & := \int_{(int)\mathcal{M} \cup (ext)\mathcal{M} (r \leq 4m_0)} (\mathfrak{d}^{\leq J} \check{\Gamma})^2 \\ & \quad + \sup_{r_0 \geq 4m_0} \left(r_0 \int_{\{r=r_0\}} |\mathfrak{d}^{\leq J} \Gamma_g|^2 + r_0^{-1} \int_{\{r=r_0\}} |\mathfrak{d}^{\leq J} \Gamma_b|^2 \right). \end{aligned}$$

Next we use the iteration assumption (8.3.10) which yields in particular

$$D_J[\Gamma] \lesssim (\epsilon_{\mathcal{B}}[J])^2.$$

Also, we have

$$\tilde{\rho} = r^2 \left(\bar{\rho} - \frac{2m}{r^3} \right) + r^2 \check{\rho}$$

and hence, using again the iteration assumption (8.3.10), as well as the control on averages provided by Lemma 3.4.1, we infer

$$\sup_{\tau \in [1, \tau_*]} E_\delta^{J-1}[\tilde{\rho}](\tau) + B_\delta^{J-1}[\tilde{\rho}](1, \tau_*) + F_\delta^{J-1}[\tilde{\rho}](1, \tau_*) + \int_{\Sigma(\tau_*)} \frac{(\mathfrak{d}^{\leq J} \tilde{\rho})^2}{r^3} \lesssim (\epsilon_{\mathcal{B}}[J])^2.$$

Together with the control of $\mathfrak{d}^{\leq k_{small}} \tilde{\rho}$ provided by the decay estimate (8.3.6), we infer from the above estimates

$$\begin{aligned} & \sup_{\tau \in [1, \tau_*]} E_\delta^J[\tilde{\rho}](\tau) + B_\delta^J[\tilde{\rho}](1, \tau_*) + F_\delta^J[\tilde{\rho}](1, \tau_*) \\ & \lesssim E_\delta^J[\tilde{\rho}](1) + (\epsilon_{\mathcal{B}}[J])^2 + \int_{\mathcal{M}} r^{1+\delta} |\mathfrak{d}^{\leq J} N_0|^2 + \left| \int_{(trap)\mathcal{M}} T(\mathfrak{d}^J \tilde{\rho}) \mathfrak{d}^J N_0 \right|. \end{aligned}$$

Next, using the form of N_0 , as well as Corollary 8.4.5, we derive

$$\int_{\mathcal{M}} r^{1+\delta} |\mathfrak{d}^{\leq J} N_0|^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Also, decomposing T as a combination of R and e_4 , integrating e_4 by parts, using again the form of N_0 , as well as Corollary 8.4.5, we have

$$\begin{aligned} & \left| \int_{(trap)\mathcal{M}} T(\mathfrak{d}^J \tilde{\rho}) \mathfrak{d}^J N_0 \right| \\ & \lesssim \left| \int_{(trap)\mathcal{M}} R(\mathfrak{d}^J \tilde{\rho}) \mathfrak{d}^J (N_0 - \text{Err}[\square_g \tilde{\rho}]) \right| + \left| \int_{(trap)\mathcal{M}} e_4(N_0 - \text{Err}[\square_g \tilde{\rho}]) \mathfrak{d}^J N_0 \right| \\ & \quad + \int_{(trap)\mathcal{M}} |T(\mathfrak{d}^J \tilde{\rho})| |\mathfrak{d}^J \text{Err}[\square_g \tilde{\rho}]| \\ & \lesssim \left| \int_{(trap)\mathcal{M}} \mathfrak{d}^J \tilde{\rho} e_4(\mathfrak{d}^J (N_0 - \text{Err}[\square_g \tilde{\rho}])) \right| \\ & \quad + \left(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right) \left(\sup_{\tau \in [1, \tau_*]} E_\delta^J[\tilde{\rho}](\tau) + B_\delta^J[\tilde{\rho}](1, \tau_*) \right)^{\frac{1}{2}} \\ & \lesssim \int_{(trap)\mathcal{M}} (\mathfrak{d}^J \tilde{\rho})^2 + \left(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right) \left(\sup_{\tau \in [1, \tau_*]} E_\delta^J[\tilde{\rho}](\tau) + B_\delta^J[\tilde{\rho}](1, \tau_*) \right)^{\frac{1}{2}}. \end{aligned}$$

In view of the above, we infer

$$\begin{aligned} & \sup_{\tau \in [1, \tau_*]} E_\delta^J[\tilde{\rho}](\tau) + B_\delta^J[\tilde{\rho}](1, \tau_*) + F_\delta^J[\tilde{\rho}](1, \tau_*) \\ & \lesssim \int_{(trap)\mathcal{M}} (\mathfrak{d}^J \tilde{\rho})^2 + (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2. \end{aligned}$$

Next, note that we have on

$$R(r - 3m) = \frac{1}{2}(e_4(r) - \Upsilon e_3(r)) - \frac{3}{2}(e_4(m) - \Upsilon e_3(m)) = \Upsilon + O(\epsilon_0) \geq \frac{1}{6} \text{ on } (trap)\mathcal{M},$$

and hence, using also integration by parts,

$$\begin{aligned}
\int_{(trap)\mathcal{M}} (\mathfrak{d}^J \tilde{\rho})^2 &\lesssim \int_{(trap)\mathcal{M}} R(r-3m) (\mathfrak{d}^J \tilde{\rho})^2 \\
&\lesssim \int_{(trap)\mathcal{M}} \left| 1 - \frac{3m}{r} \right| |\mathfrak{d}^J \tilde{\rho}| |R \mathfrak{d}^J \rho| \\
&\lesssim \epsilon_{\mathcal{B}}[J] \left(\sup_{\tau \in [1, \tau_*]} E_{\delta}^J[\tilde{\rho}](\tau) + B_{\delta}^J[\tilde{\rho}](1, \tau_*) \right)^{\frac{1}{2}}.
\end{aligned}$$

We deduce

$$\sup_{\tau \in [1, \tau_*]} E_{\delta}^J[\tilde{\rho}](\tau) + B_{\delta}^J[\tilde{\rho}](1, \tau_*) + F_{\delta}^J[\tilde{\rho}](1, \tau_*) \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2$$

as desired. This concludes the proof of Proposition 8.3.5.

8.5 Proof of Proposition 8.3.6

8.5.1 A wave equations for $\alpha + \Upsilon^2 \underline{\alpha}$

Lemma 8.5.1. *We have*

$$\begin{aligned}
\Box_2(\alpha + \Upsilon^2 \underline{\alpha}) &= \frac{4}{r} \left(1 - \frac{3m}{r} \right) \left(e_3(\alpha) - \Upsilon e_4(\underline{\alpha}) \right) + \left(-\frac{2}{r^2} + \frac{16m}{r^3} \right) \alpha \\
&\quad - \frac{2\Upsilon}{r^2} \left(1 - \frac{2m}{r} - \frac{8m^2}{r^2} \right) \underline{\alpha} + Err[\Box_2(\alpha + \Upsilon^2 \underline{\alpha})]
\end{aligned}$$

where

$$\begin{aligned}
& \text{Err}[\square_2(\alpha + \Upsilon^2 \underline{\alpha})] \\
= & \left(\Upsilon^2 \underline{V} + \frac{4m}{r^2} \Upsilon \square_g(r) - \frac{8m}{r^3} \Upsilon \mathbf{D}^\alpha(r) \mathbf{D}_\alpha(r) + \frac{8m^2}{r^4} \mathbf{D}^\alpha(r) \mathbf{D}_\alpha(r) \right) \underline{\alpha} \\
& + \left(4 \left(\omega + \frac{m}{r^2} \right) + 2 \left(\kappa - \frac{2\Upsilon}{r} \right) \right) e_3(\alpha) - 4\underline{\omega} e_4(\alpha) \\
& - 4\Upsilon \left(\Upsilon \left(\omega + \frac{m}{r^2} \right) + \frac{m}{r^2} (e_4(r) - 1) - \frac{e_4(m)}{r} \right) e_3(\underline{\alpha}) \\
& + \left(4\Upsilon^2 \underline{\omega} + 2\Upsilon^2 \left(\underline{\kappa} + \frac{2}{r} \right) - 4m\Upsilon \frac{(e_3(r) + 1)}{r^2} + 4\Upsilon \frac{e_3(m)}{r} \right) e_4(\underline{\alpha}) \\
& + \left(\left(-4\rho - \frac{8m}{r^3} \right) + 2 \left(\omega \underline{\kappa} - \frac{2m}{r^3} \right) + \frac{1}{2} \left(\kappa \underline{\kappa} + \frac{4\Upsilon}{r^2} \right) - 4e_4(\underline{\omega}) - 8\omega \underline{\omega} - 10\kappa \underline{\omega} \right) \alpha \\
& + \left(\frac{8\Upsilon}{r^2} \mathbf{D}^\alpha(m) \mathbf{D}_\alpha(r) - \frac{4\Upsilon}{r} \square_g(m) - \frac{8m}{r^3} \mathbf{D}^\alpha(r) \mathbf{D}_\alpha(m) + \frac{8m}{r} \mathbf{D}^\alpha(m) \mathbf{D}_\alpha \left(\frac{m}{r} \right) \right) \underline{\alpha} \\
& + \Upsilon^2 \left(\left(-4\rho - \frac{8m}{r^3} \right) - 4 \left(e_3(\omega) - \frac{2m}{r^3} \right) - 10 \left(\underline{\kappa} \omega - \frac{2m}{r^3} \right) + \frac{1}{2} \left(\kappa \underline{\kappa} + \frac{4\Upsilon}{r^2} \right) - 8\omega \underline{\omega} + 2\kappa \underline{\omega} \right) \underline{\alpha} \\
& + \frac{4m}{r^2} \Upsilon \left(\square_g(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right) \right) \underline{\alpha} - \frac{8m}{r^3} \left(1 - \frac{3m}{r} \right) \left(-e_4(r) e_3(r) - \Upsilon + (e_\theta(r))^2 \right) \underline{\alpha} \\
& + 4\Upsilon e_\theta(\Upsilon) e_\theta(\underline{\alpha}) + \text{Err}[\square_g \alpha] + \Upsilon^2 \text{Err}[\square_g \underline{\alpha}].
\end{aligned}$$

Proof. Recall from Proposition 2.4.6 that the curvature components α and $\underline{\alpha}$ verify the following Teukolsky equations

$$\begin{aligned}
\square_2 \alpha &= -4\underline{\omega} e_4(\alpha) + (4\omega + 2\kappa) e_3(\alpha) + V \alpha + \text{Err}[\square_g \alpha], \\
V &= -4\rho - 4e_4(\underline{\omega}) - 8\omega \underline{\omega} + 2\omega \underline{\kappa} - 10\kappa \underline{\omega} + \frac{1}{2} \kappa \underline{\kappa},
\end{aligned}$$

where

$$\begin{aligned}
\text{Err}(\square_g \alpha) &= \frac{1}{2} \vartheta e_3(\alpha) + \frac{3}{4} \vartheta^2 \rho + e_\theta(\Phi) \vartheta \beta - \frac{1}{2} \kappa (\zeta + 4\eta) \beta - (\zeta + \underline{\eta}) e_4(\beta) - \xi e_3(\beta) \\
&+ e_\theta(\Phi) (2\zeta + \underline{\eta}) \alpha + \beta^2 + e_4(\Phi) \underline{\eta} \beta + e_3(\Phi) \xi \beta - (\zeta + 4\eta) e_4(\beta) \\
&- (e_4(\zeta) + 4e_4(\eta)) \beta - 2(\kappa + \omega) (\zeta + 4\eta) \beta + 2e_\theta(\kappa + \omega) \beta - e_\theta((2\zeta + \underline{\eta}) \alpha) \\
&- 3\xi e_\theta(\rho) + 2\underline{\eta} e_\theta(\alpha) + \frac{3}{2} \vartheta \not\!{d}_1 \beta + 3\rho (\underline{\eta} + \eta + 2\zeta) \xi + \not\!{d}_1 \underline{\eta} \alpha + \frac{1}{4} \underline{\kappa} \vartheta \alpha - 2\underline{\omega} \vartheta \alpha \\
&- \frac{1}{2} \vartheta \underline{\vartheta} \alpha + \xi \underline{\xi} \alpha + \underline{\eta}^2 \alpha + \frac{3}{2} \vartheta \zeta \beta + 3\vartheta (\underline{\eta} \beta + \xi \beta) - \frac{1}{2} \vartheta (\zeta + 4\eta) \beta,
\end{aligned}$$

and

$$\begin{aligned}\square_2 \underline{\alpha} &= -4\underline{\omega} e_3(\underline{\alpha}) + (4\underline{\omega} + 2\underline{\kappa}) e_4(\underline{\alpha}) + \underline{V} \underline{\alpha} + \text{Err}[\square_{\mathbf{g}} \underline{\alpha}], \\ \underline{V} &= -4\rho - 4e_3(\underline{\omega}) - 8\underline{\omega} \underline{\omega} + 2\underline{\omega} \underline{\kappa} - 10\underline{\kappa} \underline{\omega} + \frac{1}{2} \underline{\kappa} \underline{\kappa},\end{aligned}$$

where

$$\begin{aligned}\text{Err}(\square_{\mathbf{g}} \underline{\alpha}) &= \frac{1}{2} \underline{\vartheta} e_4(\underline{\alpha}) + \frac{3}{4} \underline{\vartheta}^2 \rho + e_\theta(\Phi) \underline{\vartheta} \underline{\beta} - \frac{1}{2} \underline{\kappa} (-\zeta + 4\underline{\eta}) \underline{\beta} - (-\zeta + \eta) e_3(\underline{\beta}) - \underline{\xi} e_4(\underline{\beta}) \\ &+ e_\theta(\Phi) (-2\zeta + \eta) \underline{\alpha} + \underline{\beta}^2 + e_3(\Phi) \eta \underline{\beta} + e_4(\Phi) \underline{\xi} \underline{\beta} - (-\zeta + 4\underline{\eta}) e_3(\underline{\beta}) \\ &- (-e_3(\zeta) + 4e_3(\underline{\eta})) \underline{\beta} - 2(\underline{\kappa} + \underline{\omega}) (-\zeta + 4\underline{\eta}) \underline{\beta} + 2e_\theta(\underline{\kappa} + \underline{\omega}) \underline{\beta} - e_\theta((-2\zeta + \eta) \underline{\alpha}) \\ &- 3\underline{\xi} e_\theta(\rho) + 2\underline{\eta} e_\theta(\underline{\alpha}) + \frac{3}{2} \underline{\vartheta} \underline{\phi}_1 \underline{\beta} + 3\rho(\eta + \underline{\eta} - 2\underline{\zeta}) \underline{\xi} + \underline{\phi}_1 \eta \underline{\alpha} + \frac{1}{4} \underline{\kappa} \underline{\vartheta} \underline{\alpha} - 2\underline{\omega} \underline{\vartheta} \underline{\alpha} \\ &- \frac{1}{2} \underline{\vartheta} \underline{\vartheta} \underline{\alpha} + \underline{\xi} \underline{\xi} \underline{\alpha} + \eta^2 \underline{\alpha} - \frac{3}{2} \underline{\vartheta} \underline{\zeta} \underline{\beta} + 3\underline{\vartheta}(\eta \underline{\beta} + \underline{\xi} \underline{\beta}) - \frac{1}{2} \underline{\vartheta} (-\zeta + 4\underline{\eta}) \underline{\beta}.\end{aligned}$$

We infer from the above wave equations

$$\begin{aligned}\square_2(\alpha + \Upsilon^2 \underline{\alpha}) &= \square_2(\alpha) + \Upsilon^2 \square_2(\underline{\alpha}) + 2\mathbf{D}^\mu(\Upsilon^2) \mathbf{D}_\mu(\underline{\alpha}) + \square_0(\Upsilon^2) \underline{\alpha} \\ &= -4\underline{\omega} e_4(\alpha) + (4\underline{\omega} + 2\underline{\kappa}) e_3(\alpha) \\ &+ \Upsilon^2 \left(-4\underline{\omega} e_3(\underline{\alpha}) + (4\underline{\omega} + 2\underline{\kappa}) e_4(\underline{\alpha}) \right) - 2\underline{\Upsilon} e_3(\underline{\Upsilon}) e_4(\underline{\alpha}) - 2\underline{\Upsilon} e_4(\underline{\Upsilon}) e_3(\underline{\alpha}) \\ &+ V \alpha + \left(\Upsilon^2 \underline{V} + \square_0(\Upsilon^2) \right) \underline{\alpha} + 4\underline{\Upsilon} e_\theta(\underline{\Upsilon}) e_\theta(\underline{\alpha}) + \text{Err}[\square_{\mathbf{g}} \alpha] + \Upsilon^2 \text{Err}[\square_{\mathbf{g}} \underline{\alpha}]\end{aligned}$$

and hence

$$\begin{aligned}\square_2(\alpha + \Upsilon^2 \underline{\alpha}) &= \frac{4}{r} \left(1 - \frac{3m}{r} \right) \left(e_3(\alpha) - \underline{\Upsilon} e_4(\underline{\alpha}) \right) \\ &+ V \alpha + \left(\Upsilon^2 \underline{V} + \frac{4m}{r^2} \underline{\Upsilon} \square_g(r) - \frac{8m}{r^3} \underline{\Upsilon} \mathbf{D}^\alpha(r) \mathbf{D}_\alpha(r) + \frac{8m^2}{r^4} \mathbf{D}^\alpha(r) \mathbf{D}_\alpha(r) \right) \underline{\alpha} \\ &+ \left(4 \left(\omega + \frac{m}{r^2} \right) + 2 \left(\kappa - \frac{2\underline{\Upsilon}}{r} \right) \right) e_3(\alpha) - 4\underline{\omega} e_4(\alpha) \\ &- 4\underline{\Upsilon} \left(\underline{\Upsilon} \left(\omega + \frac{m}{r^2} \right) + \frac{m}{r^2} (e_4(r) - 1) - \frac{e_4(m)}{r} \right) e_3(\underline{\alpha}) \\ &+ \left(4\underline{\Upsilon}^2 \underline{\omega} + 2\underline{\Upsilon}^2 \left(\underline{\kappa} + \frac{2}{r} \right) - 4m \underline{\Upsilon} \frac{(e_3(r) + 1)}{r^2} + 4\underline{\Upsilon} \frac{e_3(m)}{r} \right) e_4(\underline{\alpha}) \\ &+ \left(\frac{8\underline{\Upsilon}}{r^2} \mathbf{D}^\alpha(m) \mathbf{D}_\alpha(r) - \frac{4\underline{\Upsilon}}{r} \square_g(m) - \frac{8m}{r^3} \mathbf{D}^\alpha(r) \mathbf{D}_\alpha(m) + \frac{8m}{r} \mathbf{D}^\alpha(m) \mathbf{D}_\alpha \left(\frac{m}{r} \right) \right) \underline{\alpha} \\ &+ 4\underline{\Upsilon} e_\theta(\underline{\Upsilon}) e_\theta(\underline{\alpha}) + \text{Err}[\square_{\mathbf{g}} \alpha] + \Upsilon^2 \text{Err}[\square_{\mathbf{g}} \underline{\alpha}].\end{aligned}$$

Next, we have in view of the formula for V

$$\begin{aligned} V &= -4\rho - 4e_4(\underline{\omega}) - 8\underline{\omega\underline{\omega}} + 2\underline{\omega\underline{\kappa}} - 10\underline{\kappa\underline{\omega}} + \frac{1}{2}\underline{\kappa\underline{\kappa}} \\ &= -\frac{2}{r^2} + \frac{16m}{r^3} + \left(-4\rho - \frac{8m}{r^3}\right) + 2\left(\underline{\omega\underline{\kappa}} - \frac{2m}{r^3}\right) + \frac{1}{2}\left(\underline{\kappa\underline{\kappa}} + \frac{4\underline{\Upsilon}}{r^2}\right) \\ &\quad - 4e_4(\underline{\omega}) - 8\underline{\omega\underline{\omega}} - 10\underline{\kappa\underline{\omega}}. \end{aligned}$$

Also, we have in view of the formula for \underline{V}

$$\begin{aligned} \underline{V} &= -4\rho - 4e_3(\omega) - 8\underline{\omega\underline{\omega}} + 2\underline{\omega\underline{\kappa}} - 10\underline{\kappa\underline{\omega}} + \frac{1}{2}\underline{\kappa\underline{\kappa}} \\ &= -\frac{2}{r^2} + \left(-4\rho - \frac{8m}{r^3}\right) - 4\left(e_3(\omega) + \frac{2m}{r^3}\right) - 10\left(\underline{\kappa\underline{\omega}} - \frac{2m}{r^3}\right) \\ &\quad + \frac{1}{2}\left(\underline{\kappa\underline{\kappa}} + \frac{4\underline{\Upsilon}}{r^2}\right) - 8\underline{\omega\underline{\omega}} + 2\underline{\kappa\underline{\omega}}. \end{aligned}$$

Moreover, we have

$$\begin{aligned} &\frac{4m}{r^2}\Upsilon\underline{\square}_g(r) - \frac{8m}{r^3}\Upsilon\underline{\mathbf{D}}^\alpha(r)\underline{\mathbf{D}}_\alpha(r) + \frac{8m^2}{r^4}\underline{\mathbf{D}}^\alpha(r)\underline{\mathbf{D}}_\alpha(r) \\ &= \frac{4m\underline{\Upsilon}}{r^2}\left(\frac{2}{r} - \frac{2m}{r^2}\right) + \frac{4m}{r^2}\Upsilon\left(\underline{\square}_g(r) - \left(\frac{2}{r} - \frac{2m}{r^2}\right)\right) \\ &\quad - \frac{8m}{r^3}\left(1 - \frac{3m}{r}\right)\left(-e_4(r)e_3(r) + (e_\theta(r))^2\right) \\ &= \frac{16m^2\underline{\Upsilon}}{r^4} + \frac{4m}{r^2}\Upsilon\left(\underline{\square}_g(r) - \left(\frac{2}{r} - \frac{2m}{r^2}\right)\right) - \frac{8m}{r^3}\left(1 - \frac{3m}{r}\right)\left(-e_4(r)e_3(r) - \Upsilon + (e_\theta(r))^2\right) \end{aligned}$$

and hence

$$\begin{aligned} &\Upsilon^2\underline{V} + \frac{4m}{r^2}\Upsilon\underline{\square}_g(r) - \frac{8m}{r^3}\Upsilon\underline{\mathbf{D}}^\alpha(r)\underline{\mathbf{D}}_\alpha(r) + \frac{8m^2}{r^4}\underline{\mathbf{D}}^\alpha(r)\underline{\mathbf{D}}_\alpha(r) \\ &= -\frac{2\underline{\Upsilon}}{r^2}\left(1 - \frac{2m}{r} - \frac{8m^2}{r^2}\right) \\ &\quad + \Upsilon^2\left(\left(-4\rho - \frac{8m}{r^3}\right) - 4\left(e_3(\omega) + \frac{2m}{r^3}\right) - 10\left(\underline{\kappa\underline{\omega}} - \frac{2m}{r^3}\right) + \frac{1}{2}\left(\underline{\kappa\underline{\kappa}} + \frac{4\underline{\Upsilon}}{r^2}\right) - 8\underline{\omega\underline{\omega}} + 2\underline{\kappa\underline{\omega}}\right) \\ &\quad + \frac{4m}{r^2}\Upsilon\left(\underline{\square}_g(r) - \left(\frac{2}{r} - \frac{2m}{r^2}\right)\right) - \frac{8m}{r^3}\left(1 - \frac{3m}{r}\right)\left(-e_4(r)e_3(r) - \Upsilon + (e_\theta(r))^2\right). \end{aligned}$$

We deduce

$$\begin{aligned} \underline{\square}_2(\alpha + \Upsilon^2\underline{\alpha}) &= \frac{4}{r}\left(1 - \frac{3m}{r}\right)\left(e_3(\alpha) - \Upsilon e_4(\underline{\alpha})\right) + \left(-\frac{2}{r^2} + \frac{16m}{r^3}\right)\alpha \\ &\quad - \frac{2\underline{\Upsilon}}{r^2}\left(1 - \frac{2m}{r} - \frac{8m^2}{r^2}\right)\underline{\alpha} + \text{Err}\left[\underline{\square}_2(\alpha + \Upsilon^2\underline{\alpha})\right] \end{aligned}$$

where

$$\begin{aligned}
& \text{Err}[\square_2(\alpha + \Upsilon^2 \underline{\alpha})] \\
= & \left(\Upsilon^2 \underline{V} + \frac{4m}{r^2} \Upsilon \square_g(r) - \frac{8m}{r^3} \Upsilon \mathbf{D}^\alpha(r) \mathbf{D}_\alpha(r) + \frac{8m^2}{r^4} \mathbf{D}^\alpha(r) \mathbf{D}_\alpha(r) \right) \underline{\alpha} \\
& + \left(4 \left(\omega + \frac{m}{r^2} \right) + 2 \left(\kappa - \frac{2\Upsilon}{r} \right) \right) e_3(\alpha) - 4\underline{\omega} e_4(\alpha) \\
& - 4\Upsilon \left(\Upsilon \left(\omega + \frac{m}{r^2} \right) + \frac{m}{r^2} (e_4(r) - 1) - \frac{e_4(m)}{r} \right) e_3(\underline{\alpha}) \\
& + \left(4\Upsilon^2 \underline{\omega} + 2\Upsilon^2 \left(\underline{\kappa} + \frac{2}{r} \right) - 4m\Upsilon \frac{(e_3(r) + 1)}{r^2} + 4\Upsilon \frac{e_3(m)}{r} \right) e_4(\underline{\alpha}) \\
& + \left(\left(-4\rho - \frac{8m}{r^3} \right) + 2 \left(\omega \underline{\kappa} - \frac{2m}{r^3} \right) + \frac{1}{2} \left(\kappa \underline{\kappa} + \frac{4\Upsilon}{r^2} \right) - 4e_4(\underline{\omega}) - 8\omega \underline{\omega} - 10\kappa \underline{\omega} \right) \alpha \\
& + \left(\frac{8\Upsilon}{r^2} \mathbf{D}^\alpha(m) \mathbf{D}_\alpha(r) - \frac{4\Upsilon}{r} \square_g(m) - \frac{8m}{r^3} \mathbf{D}^\alpha(r) \mathbf{D}_\alpha(m) + \frac{8m}{r} \mathbf{D}^\alpha(m) \mathbf{D}_\alpha \left(\frac{m}{r} \right) \right) \underline{\alpha} \\
& + \Upsilon^2 \left(\left(-4\rho - \frac{8m}{r^3} \right) - 4 \left(e_3(\omega) + \frac{2m}{r^3} \right) - 10 \left(\underline{\kappa} \omega - \frac{2m}{r^3} \right) + \frac{1}{2} \left(\kappa \underline{\kappa} + \frac{4\Upsilon}{r^2} \right) - 8\omega \underline{\omega} + 2\kappa \underline{\omega} \right) \underline{\alpha} \\
& + \frac{4m}{r^2} \Upsilon \left(\square_g(r) - \left(\frac{2}{r} - \frac{2m}{r^2} \right) \right) \underline{\alpha} - \frac{8m}{r^3} \left(1 - \frac{3m}{r} \right) \left(-e_4(r) e_3(r) - \Upsilon + (e_\theta(r))^2 \right) \underline{\alpha} \\
& + 4\Upsilon e_\theta(\Upsilon) e_\theta(\underline{\alpha}) + \text{Err}[\square_g \alpha] + \Upsilon^2 \text{Err}[\square_g \underline{\alpha}]
\end{aligned}$$

as desired. This concludes the proof of the lemma. \square

Lemma 8.5.2. *We have*

$$\begin{aligned}
e_3(\alpha) = & -\frac{1}{2} \underline{\kappa} \alpha - \not{d}_2^* \not{d}_1^{-1} \left\{ e_4 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa - \frac{2\Upsilon}{r} \right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} - \frac{2e_4(m)}{r^3} \right. \\
& \left. + \frac{1}{2} \underline{\vartheta} \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta) \right\} + 4\underline{\omega} \alpha - \frac{3}{2} \underline{\vartheta} \rho + (\zeta + 4\underline{\eta}) \beta.
\end{aligned}$$

and

$$\begin{aligned}
e_4(\underline{\alpha}) = & -\frac{1}{2} \kappa \underline{\alpha} - \not{d}_2^* \not{d}_1^{-1} \left\{ e_3 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \underline{\kappa} \tilde{\rho} - \frac{3m}{r^3} \left(\underline{\kappa} + \frac{2}{r} \right) + \frac{6m(e_3(r) + 1)}{r^4} - \frac{2e_3(m)}{r^3} \right. \\
& \left. + \frac{1}{2} \underline{\vartheta} \alpha + \zeta \underline{\beta} - 2(\underline{\eta} \underline{\beta} + \underline{\xi} \underline{\beta}) \right\} + 4\omega \underline{\alpha} - \frac{3}{2} \underline{\vartheta} \rho + (-\zeta + 4\underline{\eta}) \underline{\beta}.
\end{aligned}$$

Proof. Recall that we have

$$\begin{aligned}\square_2(\alpha + \Upsilon^2 \underline{\alpha}) &= \frac{4}{r} \left(1 - \frac{3m}{r}\right) \left(e_3(\alpha) - \Upsilon e_4(\underline{\alpha})\right) + \left(-\frac{2}{r^2} + \frac{16m}{r^3}\right) \alpha \\ &\quad - \frac{2\Upsilon}{r^2} \left(1 - \frac{2m}{r} - \frac{8m^2}{r^2}\right) \underline{\alpha} + \text{Err} \left[\square_2(\alpha + \Upsilon^2 \underline{\alpha})\right].\end{aligned}$$

We first express $e_3(\alpha) - \Upsilon e_4(\underline{\alpha})$ in terms of $\tilde{\rho}$, where we recall that $\tilde{\rho} = r^2 \rho + \frac{2m}{r}$. Using Bianchi, we have

$$\begin{aligned}e_3(\alpha) &= -\frac{1}{2} \underline{\kappa} \alpha - \underline{\mathfrak{d}}_2^* \beta + 4 \underline{\omega} \alpha - \frac{3}{2} \vartheta \rho + (\zeta + 4\underline{\eta}) \beta, \\ \underline{\mathfrak{d}}_1 \beta &= e_4(\rho) + \frac{3}{2} \underline{\kappa} \rho + \frac{1}{2} \vartheta \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta) \\ &= e_4 \left(\frac{\tilde{\rho}}{r^2} - \frac{2m}{r^3} \right) + \frac{3}{2} \underline{\kappa} \rho + \frac{1}{2} \vartheta \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta) \\ &= e_4 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \underline{\kappa} \tilde{\rho} - \frac{3m}{r^3} \left(\underline{\kappa} - \frac{2\Upsilon}{r} \right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} - \frac{2e_4(m)}{r^3} \\ &\quad + \frac{1}{2} \vartheta \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta)\end{aligned}$$

and hence

$$\begin{aligned}e_3(\alpha) &= -\frac{1}{2} \underline{\kappa} \alpha - \underline{\mathfrak{d}}_2^* \underline{\mathfrak{d}}_1^{-1} \left\{ e_4 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \underline{\kappa} \tilde{\rho} - \frac{3m}{r^3} \left(\underline{\kappa} - \frac{2\Upsilon}{r} \right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} - \frac{2e_4(m)}{r^3} \right. \\ &\quad \left. + \frac{1}{2} \vartheta \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta) \right\} + 4 \underline{\omega} \alpha - \frac{3}{2} \vartheta \rho + (\zeta + 4\underline{\eta}) \beta.\end{aligned}$$

Similarly, we have

$$\begin{aligned}e_4(\underline{\alpha}) &= -\frac{1}{2} \underline{\kappa} \underline{\alpha} - \underline{\mathfrak{d}}_2^* \underline{\beta} + 4 \underline{\omega} \underline{\alpha} - \frac{3}{2} \vartheta \rho + (-\zeta + 4\underline{\eta}) \underline{\beta}, \\ \underline{\mathfrak{d}}_1 \underline{\beta} &= e_3(\rho) + \frac{3}{2} \underline{\kappa} \rho + \frac{1}{2} \vartheta \underline{\alpha} + \zeta \underline{\beta} - 2(\underline{\eta} \underline{\beta} + \underline{\xi} \beta) \\ &= e_3 \left(\frac{\tilde{\rho}}{r^2} - \frac{2m}{r^3} \right) + \frac{3}{2} \underline{\kappa} \rho + \frac{1}{2} \vartheta \underline{\alpha} + \zeta \underline{\beta} - 2(\underline{\eta} \underline{\beta} + \underline{\xi} \beta) \\ &= e_3 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \underline{\kappa} \tilde{\rho} - \frac{3m}{r^3} \left(\underline{\kappa} + \frac{2}{r} \right) + \frac{6m(e_3(r) + 1)}{r^4} - \frac{2e_3(m)}{r^3} \\ &\quad + \frac{1}{2} \vartheta \underline{\alpha} + \zeta \underline{\beta} - 2(\underline{\eta} \underline{\beta} + \underline{\xi} \beta)\end{aligned}$$

and hence

$$e_4(\underline{\alpha}) = -\frac{1}{2}\kappa\underline{\alpha} - \phi_2^* \phi_1^{-1} \left\{ e_3 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa + \frac{2}{r} \right) + \frac{6m(e_3(r) + 1)}{r^4} - \frac{2e_3(m)}{r^3} \right. \\ \left. + \frac{1}{2} \vartheta \underline{\alpha} + \zeta \underline{\beta} - 2(\eta \underline{\beta} + \xi \underline{\beta}) \right\} + 4\omega \underline{\alpha} - \frac{3}{2} \vartheta \rho + (-\zeta + 4\eta) \underline{\beta}.$$

This concludes the proof of the lemma. \square

Corollary 8.5.3. *We have*

$$\square_2(\alpha + \Upsilon^2 \underline{\alpha}) - \frac{2}{r^2} \left(1 + \frac{2m}{r} \right) (\alpha + \Upsilon^2 \underline{\alpha}) \\ = -\frac{8}{r} \left(1 - \frac{3m}{r} \right) \phi_2^* \phi_1^{-1} R \left(\frac{\tilde{\rho}}{r^2} \right) - \frac{6}{r} \left(1 - \frac{3m}{r} \right) (\vartheta - \Upsilon \vartheta) \rho \\ - \frac{4}{r} \left(1 - \frac{3m}{r} \right) \phi_2^* \phi_1^{-1} \left\{ \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa - \frac{2\Upsilon}{r} \right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} \right\} \\ + \frac{4\Upsilon}{r} \left(1 - \frac{3m}{r} \right) \phi_2^* \phi_1^{-1} \left\{ \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa + \frac{2}{r} \right) + \frac{6m(e_3(r) + 1)}{r^4} \right\} + Err_1,$$

where

$$Err_1 := \frac{4}{r} \left(1 - \frac{3m}{r} \right) \left[4\omega \alpha + (\zeta + 4\eta) \beta - \Upsilon(-\zeta + 4\eta) \underline{\beta} + [\Upsilon, \phi_2^* \phi_1^{-1}] e_3 \left(\frac{\tilde{\rho}}{r^2} \right) \right. \\ \left. - \phi_2^* \phi_1^{-1} \left\{ -\frac{2e_4(m)}{r^3} + \frac{1}{2} \vartheta \alpha - \zeta \beta - 2(\eta \underline{\beta} + \xi \underline{\beta}) \right\} \right. \\ \left. + \Upsilon \phi_2^* \phi_1^{-1} \left\{ -\frac{2e_3(m)}{r^3} + \frac{1}{2} \vartheta \alpha + \zeta \underline{\beta} - 2(\eta \underline{\beta} + \xi \underline{\beta}) \right\} \right] \\ - \frac{2}{r} \left(1 - \frac{3m}{r} \right) \left(\kappa + \frac{2}{r} \right) \alpha + \frac{4\Upsilon}{r} \left(1 - \frac{3m}{r} \right) \left(\frac{1}{2} \left(\kappa - \frac{2\Upsilon}{r} \right) - 4 \left(\omega + \frac{m}{r^2} \right) \right) \underline{\alpha} \\ + Err[\square_2(\alpha + \Upsilon^2 \underline{\alpha})].$$

Proof. Recall from Lemma 8.5.1 that we have

$$\square_2(\alpha + \Upsilon^2 \underline{\alpha}) = \frac{4}{r} \left(1 - \frac{3m}{r} \right) (e_3(\alpha) - \Upsilon e_4(\underline{\alpha})) + \left(-\frac{2}{r^2} + \frac{16m}{r^3} \right) \alpha \\ - \frac{2\Upsilon}{r^2} \left(1 - \frac{2m}{r} - \frac{8m^2}{r^2} \right) \underline{\alpha} + Err[\square_2(\alpha + \Upsilon^2 \underline{\alpha})].$$

In view of Lemma 8.5.2, we have

$$\begin{aligned}
e_3(\alpha) - \Upsilon e_4(\underline{\alpha}) &= -2 \phi_2^* \phi_1^{-1} R \left(\frac{\tilde{\rho}}{r^2} \right) - \frac{1}{2} \underline{\kappa} \alpha + \Upsilon \left(\frac{1}{2} \kappa - 4\omega \right) \underline{\alpha} - \frac{3}{2} (\vartheta - \Upsilon \underline{\vartheta}) \rho \\
&\quad - \phi_2^* \phi_1^{-1} \left\{ \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa - \frac{2\Upsilon}{r} \right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} \right\} \\
&\quad + \Upsilon \phi_2^* \phi_1^{-1} \left\{ \frac{3}{2r^2} \underline{\kappa} \tilde{\rho} - \frac{3m}{r^3} \left(\underline{\kappa} + \frac{2}{r} \right) + \frac{6m(e_3(r) + 1)}{r^4} \right\} \\
&\quad + 4\underline{\omega} \alpha + (\zeta + 4\underline{\eta}) \beta - \Upsilon(-\zeta + 4\underline{\eta}) \underline{\beta} + [\Upsilon, \phi_2^* \phi_1^{-1}] e_3 \left(\frac{\tilde{\rho}}{r^2} \right) \\
&\quad - \phi_2^* \phi_1^{-1} \left\{ -\frac{2e_4(m)}{r^3} + \frac{1}{2} \underline{\vartheta} \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta) \right\} \\
&\quad + \Upsilon \phi_2^* \phi_1^{-1} \left\{ -\frac{2e_3(m)}{r^3} + \frac{1}{2} \underline{\vartheta} \alpha + \zeta \underline{\beta} - 2(\underline{\eta} \underline{\beta} + \underline{\xi} \underline{\beta}) \right\}.
\end{aligned}$$

We infer

$$\begin{aligned}
\Box_2(\alpha + \Upsilon^2 \underline{\alpha}) &= -\frac{8}{r} \left(1 - \frac{3m}{r} \right) \phi_2^* \phi_1^{-1} R \left(\frac{\tilde{\rho}}{r^2} \right) - \frac{2}{r} \left(1 - \frac{3m}{r} \right) \underline{\kappa} \alpha + \left(-\frac{2}{r^2} + \frac{16m}{r^3} \right) \alpha \\
&\quad + \frac{4\Upsilon}{r} \left(1 - \frac{3m}{r} \right) \left(\frac{1}{2} \kappa - 4\omega \right) \underline{\alpha} - \frac{2\Upsilon}{r^2} \left(1 - \frac{2m}{r} - \frac{8m^2}{r^2} \right) \underline{\alpha} \\
&\quad - \frac{6}{r} \left(1 - \frac{3m}{r} \right) (\vartheta - \Upsilon \underline{\vartheta}) \rho \\
&\quad - \frac{4}{r} \left(1 - \frac{3m}{r} \right) \phi_2^* \phi_1^{-1} \left\{ \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa - \frac{2\Upsilon}{r} \right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} \right\} \\
&\quad + \frac{4\Upsilon}{r} \left(1 - \frac{3m}{r} \right) \phi_2^* \phi_1^{-1} \left\{ \frac{3}{2r^2} \underline{\kappa} \tilde{\rho} - \frac{3m}{r^3} \left(\underline{\kappa} + \frac{2}{r} \right) + \frac{6m(e_3(r) + 1)}{r^4} \right\} \\
&\quad + \frac{4}{r} \left(1 - \frac{3m}{r} \right) \left[4\underline{\omega} \alpha + (\zeta + 4\underline{\eta}) \beta - \Upsilon(-\zeta + 4\underline{\eta}) \underline{\beta} + [\Upsilon, \phi_2^* \phi_1^{-1}] e_3 \left(\frac{\tilde{\rho}}{r^2} \right) \right. \\
&\quad \left. - \phi_2^* \phi_1^{-1} \left\{ -\frac{2e_4(m)}{r^3} + \frac{1}{2} \underline{\vartheta} \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta) \right\} \right. \\
&\quad \left. + \Upsilon \phi_2^* \phi_1^{-1} \left\{ -\frac{2e_3(m)}{r^3} + \frac{1}{2} \underline{\vartheta} \alpha + \zeta \underline{\beta} - 2(\underline{\eta} \underline{\beta} + \underline{\xi} \underline{\beta}) \right\} \right] \\
&\quad + \text{Err} \left[\Box_2(\alpha + \Upsilon^2 \underline{\alpha}) \right].
\end{aligned}$$

Since we have

$$\begin{aligned}
& -\frac{2}{r} \left(1 - \frac{3m}{r}\right) \underline{\kappa}\alpha + \frac{4\Upsilon}{r} \left(1 - \frac{3m}{r}\right) \left(\frac{1}{2}\kappa - 4\omega\right) \underline{\alpha} \\
= & \frac{4}{r^2} \left(1 - \frac{3m}{r}\right) \alpha + \frac{4\Upsilon}{r^2} \left(1 - \frac{3m}{r}\right) \left(1 + \frac{2m}{r}\right) \underline{\alpha} - \frac{2}{r} \left(1 - \frac{3m}{r}\right) \left(\frac{\underline{\kappa}}{r} + \frac{2}{r}\right) \alpha \\
& + \frac{4\Upsilon}{r} \left(1 - \frac{3m}{r}\right) \left(\frac{1}{2} \left(\kappa - \frac{2\Upsilon}{r}\right) - 4 \left(\omega + \frac{m}{r^2}\right)\right) \underline{\alpha},
\end{aligned}$$

this yields

$$\begin{aligned}
\Box_2(\alpha + \Upsilon^2 \underline{\alpha}) = & -\frac{8}{r} \left(1 - \frac{3m}{r}\right) \not\phi_2^* \not\phi_1^{-1} R \left(\frac{\tilde{\rho}}{r^2}\right) + \frac{2}{r^2} \left(1 + \frac{2m}{r}\right) \alpha + \frac{2\Upsilon}{r^2} \left(1 - \frac{4m^2}{r^2}\right) \underline{\alpha} \\
& - \frac{6}{r} \left(1 - \frac{3m}{r}\right) (\vartheta - \Upsilon \underline{\vartheta}) \rho \\
& - \frac{4}{r} \left(1 - \frac{3m}{r}\right) \not\phi_2^* \not\phi_1^{-1} \left\{ \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa - \frac{2\Upsilon}{r}\right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} \right\} \\
& + \frac{4\Upsilon}{r} \left(1 - \frac{3m}{r}\right) \not\phi_2^* \not\phi_1^{-1} \left\{ \frac{3}{2r^2} \underline{\kappa} \tilde{\rho} - \frac{3m}{r^3} \left(\frac{\underline{\kappa}}{r} + \frac{2}{r}\right) + \frac{6m(e_3(r) + 1)}{r^4} \right\} + \text{Err}_1,
\end{aligned}$$

where

$$\begin{aligned}
\text{Err}_1 = & \frac{4}{r} \left(1 - \frac{3m}{r}\right) \left[4\underline{\omega}\alpha + (\zeta + 4\underline{\eta})\beta - \Upsilon(-\zeta + 4\underline{\eta})\underline{\beta} + [\Upsilon, \not\phi_2^* \not\phi_1^{-1}] e_3 \left(\frac{\tilde{\rho}}{r^2}\right) \right. \\
& - \not\phi_2^* \not\phi_1^{-1} \left\{ -\frac{2e_4(m)}{r^3} + \frac{1}{2} \underline{\vartheta}\alpha - \zeta\beta - 2(\underline{\eta}\beta + \underline{\xi}\beta) \right\} \\
& \left. + \Upsilon \not\phi_2^* \not\phi_1^{-1} \left\{ -\frac{2e_3(m)}{r^3} + \frac{1}{2} \vartheta\underline{\alpha} + \zeta\underline{\beta} - 2(\eta\underline{\beta} + \underline{\xi}\beta) \right\} \right] \\
& - \frac{2}{r} \left(1 - \frac{3m}{r}\right) \left(\frac{\underline{\kappa}}{r} + \frac{2}{r}\right) \alpha + \frac{4\Upsilon}{r} \left(1 - \frac{3m}{r}\right) \left(\frac{1}{2} \left(\kappa - \frac{2\Upsilon}{r}\right) - 4 \left(\omega + \frac{m}{r^2}\right)\right) \underline{\alpha} \\
& + \text{Err} \left[\Box_2(\alpha + \Upsilon^2 \underline{\alpha}) \right].
\end{aligned}$$

Now, since we have

$$\frac{2}{r^2} \left(1 + \frac{2m}{r}\right) \alpha + \frac{2\Upsilon}{r^2} \left(1 - \frac{4m^2}{r^2}\right) \underline{\alpha} = \frac{2}{r^2} \left(1 + \frac{2m}{r}\right) (\alpha + \Upsilon^2 \underline{\alpha}),$$

we infer

$$\begin{aligned}
& \square_2(\alpha + \Upsilon^2 \underline{\alpha}) - \frac{2}{r^2} \left(1 + \frac{2m}{r}\right) (\alpha + \Upsilon^2 \underline{\alpha}) \\
= & -\frac{8}{r} \left(1 - \frac{3m}{r}\right) \not\phi_2^* \not\phi_1^{-1} R \left(\frac{\tilde{\rho}}{r^2}\right) - \frac{6}{r} \left(1 - \frac{3m}{r}\right) (\vartheta - \Upsilon \underline{\vartheta}) \rho \\
& - \frac{4}{r} \left(1 - \frac{3m}{r}\right) \not\phi_2^* \not\phi_1^{-1} \left\{ \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa - \frac{2\Upsilon}{r}\right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} \right\} \\
& + \frac{4\Upsilon}{r} \left(1 - \frac{3m}{r}\right) \not\phi_2^* \not\phi_1^{-1} \left\{ \frac{3}{2r^2} \underline{\kappa} \tilde{\rho} - \frac{3m}{r^3} \left(\underline{\kappa} + \frac{2}{r}\right) + \frac{6m(e_3(r) + 1)}{r^4} \right\} + \text{Err}_1,
\end{aligned}$$

as desired. This concludes the proof of the corollary. \square

8.5.2 End of the proof of Proposition 8.3.6

In view of Corollary 8.5.3, $\alpha + \Upsilon^2 \underline{\alpha}$ satisfies

$$(\square_2 + V_2)(\alpha + \Upsilon^2 \underline{\alpha}) = N_2, \quad V_2 = -\frac{2}{r^2} \left(1 + \frac{2m}{r}\right),$$

where

$$\begin{aligned}
N_2 := & -\frac{8}{r} \left(1 - \frac{3m}{r}\right) \not\phi_2^* \not\phi_1^{-1} R \left(\frac{\tilde{\rho}}{r^2}\right) - \frac{6}{r} \left(1 - \frac{3m}{r}\right) (\vartheta - \Upsilon \underline{\vartheta}) \rho \\
& - \frac{4}{r} \left(1 - \frac{3m}{r}\right) \not\phi_2^* \not\phi_1^{-1} \left\{ \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa - \frac{2\Upsilon}{r}\right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} \right\} \\
& + \frac{4\Upsilon}{r} \left(1 - \frac{3m}{r}\right) \not\phi_2^* \not\phi_1^{-1} \left\{ \frac{3}{2r^2} \underline{\kappa} \tilde{\rho} - \frac{3m}{r^3} \left(\underline{\kappa} + \frac{2}{r}\right) + \frac{6m(e_3(r) + 1)}{r^4} \right\} + \text{Err}_1.
\end{aligned}$$

We may thus apply the estimate (10.5.1) of Theorem 10.5.2 with $\psi = \alpha + \Upsilon^2 \underline{\alpha}$ and $s = J$ to obtain for any $k_{small} \leq J \leq k_{large} - 1$

$$\begin{aligned}
& \sup_{\tau \in [1, \tau_*]} E_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](\tau) + B_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) + F_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) \\
\lesssim & E_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](1) + \sup_{\tau \in [1, \tau_*]} E_\delta^{J-1}[\alpha + \Upsilon^2 \underline{\alpha}](\tau) + B_\delta^{J-1}[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) \\
& + F_\delta^{J-1}[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) + D_J[\Gamma] \left(\sup_{\mathcal{M}} r u_{trap}^{\frac{1}{2} + \delta_{dec}} |\mathfrak{d}^{\leq k_{small}}(\alpha + \Upsilon^2 \underline{\alpha})| \right)^2 \\
& + \int_{\mathcal{M}} r^{1+\delta} |\mathfrak{d}^{\leq J} N_2|^2 + \left| \int_{(trap)_{\mathcal{M}}} T(\mathfrak{d}^J(\alpha + \Upsilon^2 \underline{\alpha})) \mathfrak{d}^J N_2 \right|,
\end{aligned}$$

where $D_J[\Gamma]$ is defined by

$$D_J[\Gamma] := \int_{(int)\mathcal{M} \cup (ext)\mathcal{M}(r \leq 4m_0)} (\mathfrak{d}^{\leq J} \check{\Gamma})^2 + \sup_{r_0 \geq 4m_0} \left(r_0 \int_{\{r=r_0\}} |\mathfrak{d}^{\leq J} \Gamma_g|^2 + r_0^{-1} \int_{\{r=r_0\}} |\mathfrak{d}^{\leq J} \Gamma_b|^2 \right).$$

Next we use the iteration assumption (8.3.10) which yields in particular

$$D_J[\Gamma] \lesssim (\epsilon_{\mathcal{B}}[J])^2$$

and

$$\sup_{\tau \in [1, \tau_*]} E_\delta^{J-1}[\alpha + \Upsilon^2 \underline{\alpha}](\tau) + B_\delta^{J-1}[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) + F_\delta^{J-1}[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) \lesssim (\epsilon_{\mathcal{B}}[J])^2.$$

Together with the control of $\mathfrak{d}^{\leq k_{small}}(\alpha + \Upsilon^2 \underline{\alpha})$ provided by the decay estimate (8.3.6), we infer from the above estimates

$$\begin{aligned} & \sup_{\tau \in [1, \tau_*]} E_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](\tau) + B_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) + F_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) \\ & \lesssim E_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](1) + (\epsilon_{\mathcal{B}}[J])^2 + \int_{\mathcal{M}} r^{1+\delta} |\mathfrak{d}^{\leq J} N_2|^2 + \left| \int_{(trap)\mathcal{M}} T(\mathfrak{d}^J(\alpha + \Upsilon^2 \underline{\alpha})) \mathfrak{d}^J N_2 \right|. \end{aligned}$$

Next, using the form of N_2 , as well as the control of $\tilde{\rho}$ provided by Proposition 8.3.5, we derive

$$\int_{\mathcal{M}} r^{1+\delta} |\mathfrak{d}^{\leq J} N_2|^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2$$

and

$$\begin{aligned} & \left| \int_{(trap)\mathcal{M}} T(\mathfrak{d}^J(\alpha + \Upsilon^2 \underline{\alpha})) \mathfrak{d}^J N_2 \right| \\ & \lesssim \int_{(trap)\mathcal{M}} \left| 1 - \frac{3m}{r} \right| |T(\mathfrak{d}^J(\alpha + \Upsilon^2 \underline{\alpha}))| \left(|R(\mathfrak{d}^J \tilde{\rho})| + |\mathfrak{d}^J \tilde{\rho}| + |\mathfrak{d}^J \check{\Gamma}| \right) \\ & \quad + \int_{(trap)\mathcal{M}} |T(\mathfrak{d}^J(\alpha + \Upsilon^2 \underline{\alpha}))| |\text{Err}_1| \\ & \lesssim \left(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right) \left(\sup_{\tau \in [1, \tau_*]} E_\delta^J[\tilde{\rho}](\tau) + B_\delta^J[\tilde{\rho}](1, \tau_*) \right)^{\frac{1}{2}}. \end{aligned}$$

In view of the above, we infer

$$\begin{aligned} & \sup_{\tau \in [1, \tau_*]} E_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](\tau) + B_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) + F_\delta^J[\alpha + \Upsilon^2 \underline{\alpha}](1, \tau_*) \\ & \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \end{aligned}$$

as desired. This concludes the proof of Proposition 8.3.6.

8.6 Proof of Proposition 8.3.7

8.6.1 Control of α and $\Upsilon^2 \underline{\alpha}$

We initiate the proof of Proposition 8.3.7 by deriving a suitable control for α and $\Upsilon^2 \underline{\alpha}$. Recall from Lemma 8.5.2 that we have

$$\begin{aligned} e_4(\underline{\alpha}) &= -\frac{1}{2} \kappa \underline{\alpha} - \mathfrak{d}_2^* \mathfrak{d}_1^{-1} \left\{ e_3 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa + \frac{2}{r} \right) + \frac{6m(e_3(r) + 1)}{r^4} - \frac{2e_3(m)}{r^3} \right. \\ & \quad \left. + \frac{1}{2} \vartheta \underline{\alpha} + \zeta \underline{\beta} - 2(\underline{\eta} \underline{\beta} + \underline{\xi} \underline{\beta}) \right\} + 4\omega \underline{\alpha} - \frac{3}{2} \vartheta \rho + (-\zeta + 4\underline{\eta}) \underline{\beta}. \end{aligned}$$

We infer

$$\begin{aligned} & e_4(\alpha - \Upsilon^2 \underline{\alpha}) \\ &= e_4(\alpha + \Upsilon^2 \underline{\alpha}) - 2e_4(\Upsilon^2 \underline{\alpha}) \\ &= e_4(\alpha + \Upsilon^2 \underline{\alpha}) - 2\Upsilon^2 e_4(\underline{\alpha}) - 2e_4(\Upsilon^2 \underline{\alpha}) \\ &= e_4(\alpha + \Upsilon^2 \underline{\alpha}) + 2\Upsilon^2 \mathfrak{d}_2^* \mathfrak{d}_1^{-1} \left\{ e_3 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa + \frac{2}{r} \right) + \frac{6m(e_3(r) + 1)}{r^4} - \frac{2e_3(m)}{r^3} \right. \\ & \quad \left. + \frac{1}{2} \vartheta \underline{\alpha} + \zeta \underline{\beta} - 2(\underline{\eta} \underline{\beta} + \underline{\xi} \underline{\beta}) \right\} + \Upsilon^2 \kappa \underline{\alpha} - 8\Upsilon^2 \omega \underline{\alpha} + 3\Upsilon^2 \vartheta \rho - 2\Upsilon^2 (-\zeta + 4\underline{\eta}) \underline{\beta} \\ & \quad - \frac{8m\Upsilon e_4(r)}{r^2} \underline{\alpha} + \frac{8\Upsilon e_4(m)}{r} \underline{\alpha}. \end{aligned}$$

Also, recall from Lemma 8.5.2 that we have

$$\begin{aligned} e_3(\alpha) &= -\frac{1}{2} \kappa \alpha - \mathfrak{d}_2^* \mathfrak{d}_1^{-1} \left\{ e_4 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa - \frac{2\Upsilon}{r} \right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} - \frac{2e_4(m)}{r^3} \right. \\ & \quad \left. + \frac{1}{2} \vartheta \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta) \right\} + 4\omega \alpha - \frac{3}{2} \vartheta \rho + (\zeta + 4\underline{\eta}) \beta. \end{aligned}$$

We infer

$$\begin{aligned}
& e_3(\alpha - \Upsilon^2 \underline{\alpha}) \\
&= -e_3(\alpha + \Upsilon^2 \underline{\alpha}) + 2e_3(\alpha) \\
&= -e_3(\alpha + \Upsilon^2 \underline{\alpha}) - 2 \mathfrak{d}_2^* \mathfrak{d}_1^{-1} \left\{ e_4 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa - \frac{2\Upsilon}{r} \right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} - \frac{2e_4(m)}{r^3} \right. \\
&\quad \left. + \frac{1}{2} \underline{\vartheta} \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta) \right\} - \underline{\kappa} \alpha + 8\underline{\omega} \alpha - 3\underline{\vartheta} \rho + 2(\zeta + 4\underline{\eta}) \beta.
\end{aligned}$$

In view of the above identities for $e_4(\alpha - \Upsilon^2 \underline{\alpha})$ and $e_3(\alpha - \Upsilon^2 \underline{\alpha})$, and using the control for $\tilde{\rho}$ provided by Proposition 8.3.5 as well as the control for $\alpha + \Upsilon^2 \underline{\alpha}$ provided by Proposition 8.3.6, and the iteration assumption (8.3.10), we obtain

$$B_\delta^{J-1}[e_3(\alpha - \Upsilon^2 \underline{\alpha})](1, \tau_*) + B_\delta^{J-1}[r e_4(\alpha - \Upsilon^2 \underline{\alpha})](1, \tau_*) \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Also, using the Bianchi identity for $\mathfrak{d}_2 \alpha$ and $\mathfrak{d}_1 \beta$, we have

$$\begin{aligned}
\mathfrak{d}_1 \mathfrak{d}_2 \alpha &= \mathfrak{d}_1 \left(e_4 \beta + 2(\kappa + \omega) \beta - (2\zeta + \eta) \alpha - 3\xi \rho \right) \\
&= e_4(\mathfrak{d}_1 \beta) + [\mathfrak{d}_1, e_4] \beta + \mathfrak{d}_1 \left(2(\kappa + \omega) \beta - (2\zeta + \eta) \alpha - 3\xi \rho \right) \\
&= e_4 \left(e_4 \rho + \frac{3}{2} \rho + \frac{1}{2} \underline{\vartheta} \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta) \right) + [\mathfrak{d}_1, e_4] \beta \\
&\quad + \mathfrak{d}_1 \left(2(\kappa + \omega) \beta - (2\zeta + \eta) \alpha - 3\xi \rho \right) \\
&= e_4 \left[e_4 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\kappa - \frac{2\Upsilon}{r} \right) + \frac{6m(e_4(r) - \Upsilon)}{r^4} - \frac{2e_4(m)}{r^3} \right. \\
&\quad \left. + \frac{1}{2} \underline{\vartheta} \alpha - \zeta \beta - 2(\underline{\eta} \beta + \underline{\xi} \beta) \right] + [\mathfrak{d}_1, e_4] \beta + \mathfrak{d}_1 \left(2(\kappa + \omega) \beta - (2\zeta + \eta) \alpha - 3\xi \rho \right).
\end{aligned}$$

Using the control for $\tilde{\rho}$ provided by Proposition 8.3.5 as well as the iteration assumption (8.3.10), we obtain

$$B_\delta^{J-2}[r^2 \mathfrak{d}_1 \mathfrak{d}_2 \alpha](1, \tau_*) \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Using the control for $\alpha + \Upsilon^2 \underline{\alpha}$ provided by Proposition 8.3.6, we infer

$$\begin{aligned}
B_\delta^{J-2}[r^2 \mathfrak{d}_1 \mathfrak{d}_2(\alpha - \Upsilon^2 \underline{\alpha})](1, \tau_*) &\lesssim B_\delta^{J-2}[r^2 \mathfrak{d}_1 \mathfrak{d}_2 \alpha](1, \tau_*) + B_\delta^{J-2}[r^2 \mathfrak{d}_1 \mathfrak{d}_2(\alpha + \Upsilon^2 \underline{\alpha})](1, \tau_*) \\
&\lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.
\end{aligned}$$

Using a Poincaré inequality for \not{d}_1 and for \not{d}_2 , we deduce

$$B_\delta^{J-2}[\not{d}^2(\alpha - \Upsilon^2\underline{\alpha})](1, \tau_*) \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Together with the above estimate for $e_3(\alpha - \Upsilon^2\underline{\alpha})$ and $re_4(\alpha - \Upsilon^2\underline{\alpha})$, we deduce

$$B_\delta^J[\alpha - \Upsilon^2\underline{\alpha}](1, \tau_*) \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Together with the control for $\alpha + \Upsilon^2\underline{\alpha}$ provided by Proposition 8.3.6, we finally obtain

$$B_\delta^J[\alpha](1, \tau_*) + B_\delta^J[\Upsilon^2\underline{\alpha}](1, \tau_*) \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2. \quad (8.6.1)$$

8.6.2 Control of $\underline{\alpha}$

(8.6.1) provides in particular the control of $\Upsilon^2\underline{\alpha}$. In this section, we infer a suitable control for $\underline{\alpha}$ using the wave equation satisfied by $\underline{\alpha}$ and the redshift vectorfield.

Let $Y_{(0)}$ the vectorfield given by

$$Y_{(0)} := \left(1 + \frac{5}{4m}(r - 2m) + \Upsilon \right) e_3 + \left(1 + \frac{5}{4m}(r - 2m) \right) e_4,$$

where $Y_{(0)}$ has been introduced in Proposition 10.1.29 in connection with the redshift vectorfield.

Lemma 8.6.1. *We have*

$$\square_{2\underline{\alpha}} = \frac{4m}{r^2 \left(1 + \frac{5}{4m}(r - 2m) + \Upsilon \right)} Y_{(0)}\underline{\alpha} + \tilde{N}_2$$

where \tilde{N}_2 is given by

$$\begin{aligned} \tilde{N}_2 := & -\frac{4}{r} \left(1 + \frac{m \left(1 + \frac{5}{4m}(r - 2m) \right)}{r \left(1 + \frac{5}{4m}(r - 2m) + \Upsilon \right)} \right) \left[-\frac{1}{2} \kappa \underline{\alpha} \right. \\ & - \not{d}_2^* \not{d}_1^{-1} \left\{ e_3 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\frac{\kappa}{r} + \frac{2}{r} \right) + \frac{6m(e_3(r) + 1)}{r^4} - \frac{2e_3(m)}{r^3} \right. \\ & \left. \left. + \frac{1}{2} \not{d} \underline{\alpha} + \zeta \underline{\beta} - 2(\eta \underline{\beta} + \xi \underline{\beta}) \right\} + 4\omega \underline{\alpha} - \frac{3}{2} \not{d} \rho + (-\zeta + 4\underline{\eta}) \underline{\beta} \right] \\ & + \underline{V} \underline{\alpha} - 4 \left(\omega + \frac{m}{r^2} \right) e_3(\underline{\alpha}) + \left(4\underline{\omega} + 2 \left(\frac{\kappa}{r} + \frac{2}{r} \right) \right) e_4(\underline{\alpha}) + Err[\square_{\mathbf{g}} \underline{\alpha}]. \end{aligned}$$

Proof. Recall from Proposition 2.4.6 that $\underline{\alpha}$ verifies the following Teukolsky equation

$$\begin{aligned}\square_2 \underline{\alpha} &= -4\omega e_3(\underline{\alpha}) + (4\underline{\omega} + 2\underline{\kappa})e_4(\underline{\alpha}) + \underline{V}\underline{\alpha} + \text{Err}[\square_{\mathbf{g}}\underline{\alpha}], \\ \underline{V} &= -4\rho - 4e_3(\omega) - 8\underline{\omega}\underline{\omega} + 2\underline{\omega}\underline{\kappa} - 10\underline{\kappa}\underline{\omega} + \frac{1}{2}\underline{\kappa}\underline{\kappa},\end{aligned}$$

where

$$\begin{aligned}\text{Err}(\square_{\mathbf{g}}\underline{\alpha}) &= \frac{1}{2}\vartheta e_4(\underline{\alpha}) + \frac{3}{4}\vartheta^2 \rho + e_\theta(\Phi)\vartheta \underline{\beta} - \frac{1}{2}\underline{\kappa}(-\zeta + 4\underline{\eta})\underline{\beta} - (-\zeta + \eta)e_3(\underline{\beta}) - \underline{\xi}e_4(\underline{\beta}) \\ &\quad + e_\theta(\Phi)(-2\zeta + \eta)\underline{\alpha} + \underline{\beta}^2 + e_3(\Phi)\eta\underline{\beta} + e_4(\Phi)\underline{\xi}\underline{\beta} - (-\zeta + 4\underline{\eta})e_3(\underline{\beta}) \\ &\quad - (-e_3(\zeta) + 4e_3(\underline{\eta}))\underline{\beta} - 2(\underline{\kappa} + \underline{\omega})(-\zeta + 4\underline{\eta})\underline{\beta} + 2e_\theta(\underline{\kappa} + \underline{\omega})\underline{\beta} - e_\theta((-2\zeta + \eta)\underline{\alpha}) \\ &\quad - 3\underline{\xi}e_\theta(\rho) + 2\underline{\eta}e_\theta(\underline{\alpha}) + \frac{3}{2}\vartheta \underline{\phi}_1 \underline{\beta} + 3\rho(\eta + \underline{\eta} - 2\underline{\zeta})\underline{\xi} + \underline{\phi}_1 \eta \underline{\alpha} + \frac{1}{4}\underline{\kappa}\vartheta \underline{\alpha} - 2\underline{\omega}\vartheta \underline{\alpha} \\ &\quad - \frac{1}{2}\vartheta\vartheta \underline{\alpha} + \underline{\xi}\underline{\xi}\underline{\alpha} + \eta^2 \underline{\alpha} - \frac{3}{2}\vartheta \underline{\zeta}\underline{\beta} + 3\vartheta(\eta\underline{\beta} + \underline{\xi}\underline{\beta}) - \frac{1}{2}\vartheta(-\zeta + 4\underline{\eta})\underline{\beta}.\end{aligned}$$

We deduce

$$\begin{aligned}\square_2 \underline{\alpha} &= \frac{4m}{r^2}e_3(\underline{\alpha}) - \frac{4}{r}e_4(\underline{\alpha}) + \underline{V}\underline{\alpha} - 4\left(\omega + \frac{m}{r^2}\right)e_3(\underline{\alpha}) + \left(4\underline{\omega} + 2\left(\underline{\kappa} + \frac{2}{r}\right)\right)e_4(\underline{\alpha}) \\ &\quad + \text{Err}[\square_{\mathbf{g}}\underline{\alpha}].\end{aligned}$$

In view of the definition of $Y_{(0)}$, we infer

$$\begin{aligned}\square_2 \underline{\alpha} &= \frac{4m}{r^2\left(1 + \frac{5}{4m}(r - 2m) + \Upsilon\right)}Y_{(0)}\underline{\alpha} - \frac{4}{r}\left(1 + \frac{m\left(1 + \frac{5}{4m}(r - 2m)\right)}{r\left(1 + \frac{5}{4m}(r - 2m) + \Upsilon\right)}\right)e_4(\underline{\alpha}) \\ &\quad + \underline{V}\underline{\alpha} - 4\left(\omega + \frac{m}{r^2}\right)e_3(\underline{\alpha}) + \left(4\underline{\omega} + 2\left(\underline{\kappa} + \frac{2}{r}\right)\right)e_4(\underline{\alpha}) + \text{Err}[\square_{\mathbf{g}}\underline{\alpha}].\end{aligned}$$

Next, recall from Lemma 8.5.2 that we have

$$\begin{aligned}e_4(\underline{\alpha}) &= -\frac{1}{2}\underline{\kappa}\underline{\alpha} - \underline{\phi}_2^* \underline{\phi}_1^{-1} \left\{ e_3\left(\frac{\tilde{\rho}}{r^2}\right) + \frac{3}{2r^2}\underline{\kappa}\tilde{\rho} - \frac{3m}{r^3}\left(\underline{\kappa} + \frac{2}{r}\right) + \frac{6m(e_3(r) + 1)}{r^4} - \frac{2e_3(m)}{r^3} \right. \\ &\quad \left. + \frac{1}{2}\vartheta \underline{\alpha} + \underline{\zeta}\underline{\beta} - 2(\eta\underline{\beta} + \underline{\xi}\underline{\beta}) \right\} + 4\underline{\omega}\underline{\alpha} - \frac{3}{2}\vartheta \rho + (-\zeta + 4\underline{\eta})\underline{\beta}.\end{aligned}$$

We infer

$$\square_2 \underline{\alpha} = \frac{4m}{r^2\left(1 + \frac{5}{4m}(r - 2m) + \Upsilon\right)}Y_{(0)}\underline{\alpha} + \tilde{N}_2$$

where \tilde{N}_2 is given by

$$\begin{aligned} \tilde{N}_2 = & -\frac{4}{r} \left(1 + \frac{m \left(1 + \frac{5}{4m}(r-2m) \right)}{r \left(1 + \frac{5}{4m}(r-2m) + \Upsilon \right)} \right) \left[-\frac{1}{2} \kappa \underline{\alpha} \right. \\ & - \not\partial_2^* \not\partial_1^{-1} \left\{ e_3 \left(\frac{\tilde{\rho}}{r^2} \right) + \frac{3}{2r^2} \kappa \tilde{\rho} - \frac{3m}{r^3} \left(\frac{\kappa}{r} + \frac{2}{r} \right) + \frac{6m(e_3(r)+1)}{r^4} - \frac{2e_3(m)}{r^3} \right. \\ & \left. \left. + \frac{1}{2} \vartheta \underline{\alpha} + \zeta \underline{\beta} - 2(\eta \underline{\beta} + \xi \underline{\beta}) \right\} + 4\omega \underline{\alpha} - \frac{3}{2} \vartheta \rho + (-\zeta + 4\eta) \underline{\beta} \right] \\ & + \underline{V} \underline{\alpha} - 4 \left(\omega + \frac{m}{r^2} \right) e_3(\underline{\alpha}) + \left(4\underline{\omega} + 2 \left(\frac{\kappa}{r} + \frac{2}{r} \right) \right) e_4(\underline{\alpha}) + \text{Err}[\square_{\mathbf{g}} \underline{\alpha}]. \end{aligned}$$

This concludes the proof of the lemma. \square

Lemma 8.6.2. \tilde{N}_2 , in the RHS of the wave equation for $\underline{\alpha}$ introduced in Lemma 8.6.1, satisfies

$$\int_{(int)\mathcal{M}} |\not\partial^J \tilde{N}_2|^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Proof. The proof of the lemma follows immediately from the form of \tilde{N}_2 , see Lemma 8.6.1, as well as the control for $\tilde{\rho}$ provided by Proposition 8.3.5, (8.3.6), and the iteration assumption (8.3.10). \square

In view of Lemma 8.6.1, we may apply Proposition 10.5.4 with

$$\psi = \underline{\alpha}, \quad f_2(r, m) = \frac{4m}{r^2 \left(1 + \frac{5}{4m}(r-2m) + \Upsilon \right)}.$$

We infer

$$\begin{aligned} \int_{(int)\mathcal{M}(1, \tau_*)} (\not\partial^{J+1} \underline{\alpha})^2 & \lesssim E_\delta^J[\underline{\alpha}](\tau=1) + \int_{(ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}(1, \tau_*)} (\not\partial^{J+1} \underline{\alpha})^2 \\ & + D_J[\Gamma] \left(\sup_{(int)\mathcal{M}(1, \tau_*) \cup (ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}} r |\not\partial^{\leq k_{small}} \underline{\alpha}| \right)^2 \\ & + \int_{(int)\mathcal{M}(1, \tau_*) \cup (ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}} \left((\not\partial^{\leq s} \underline{\alpha})^2 + (\not\partial^{\leq J+1} \tilde{N}_2)^2 \right). \end{aligned}$$

Next we use the iteration assumption (8.3.10) which yields in particular

$$D_J[\Gamma] \lesssim (\epsilon_{\mathcal{B}}[J])^2$$

together with the control of $\mathfrak{d}^{\leq k_{small}} \underline{\alpha}$ provided by the decay estimate (8.3.6), as well as the iteration assumption and the control for \tilde{N}_2 provided by Lemma 8.6.2 to deduce

$$\int_{(int)\mathcal{M}(1,\tau_*)} (\mathfrak{d}^{J+1} \underline{\alpha})^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 + \int_{(ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}(1,\tau_*)} (\mathfrak{d}^{J+1} \underline{\alpha})^2.$$

Note that $\Upsilon^2 \gtrsim \delta_{\mathcal{H}}^2 > 0$ on $(ext)\mathcal{M}$ and hence

$$\int_{(ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}(1,\tau_*)} (\mathfrak{d}^{J+1} \underline{\alpha})^2 \lesssim \int_{(ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}(1,\tau_*)} (\mathfrak{d}^{J+1}(\Upsilon^2 \underline{\alpha}))^2$$

which together with the control of $\Upsilon^2 \underline{\alpha}$ provided by (8.6.1) yields

$$\int_{(ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}(1,\tau_*)} (\mathfrak{d}^{J+1} \underline{\alpha})^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2$$

and hence

$$\int_{(int)\mathcal{M}(1,\tau_*)} (\mathfrak{d}^{J+1} \underline{\alpha})^2 \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Since

$$B_{\delta}^J[\underline{\alpha}](1, \tau_*) \lesssim \int_{(int)\mathcal{M}(1,\tau_*)} (\mathfrak{d}^{\leq J+1} \underline{\alpha})^2 + B_{\delta}^J[\Upsilon^2 \underline{\alpha}](1, \tau_*),$$

using again (8.6.1), we finally obtain

$$B_{\delta}^J[\underline{\alpha}](1, \tau_*) \lesssim (\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2. \quad (8.6.2)$$

8.6.3 End of the proof of Proposition 8.3.7

We have

$$\check{\rho} = \frac{\tilde{\rho}}{r^2} - \left(\bar{\rho} - \frac{2m}{r^3} \right).$$

Together with the control for $\tilde{\rho}$ provided by Proposition 8.3.5, as well as the control on averages provided by Lemma 3.4.1, we infer

$$B_\delta^J[\check{\rho}](1, \tau_*) \lesssim (\epsilon_B[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Together with the control for α provided by (8.6.1) and the control for $\underline{\alpha}$ provided by (8.6.2), we infer

$$B_\delta^J[\alpha, \check{\rho}, \underline{\alpha}](1, \tau_*) \lesssim (\epsilon_B[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2.$$

Together with the Bianchi identities for $e_4(\beta)$, $e_3(\beta)$, $\phi_1\beta$, $e_4(\underline{\beta})$, $e_3(\underline{\beta})$, $\phi_1\underline{\beta}$, as well as the iteration assumption (8.3.10), we infer

$$B_{\delta-2}^J[\beta, \underline{\beta}](1, \tau_*) \lesssim (\epsilon_B[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2$$

and hence

$$B_{-2}^J[\alpha, \beta, \check{\rho}, \underline{\beta}, \underline{\alpha}](1, \tau_*) \lesssim (\epsilon_B[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2$$

as desired. This concludes the proof of Proposition 8.3.7.

8.7 Proof of Proposition 8.3.8

First, note that, by definition of the norms B_{-2}^J , ${}^{(int)}\mathfrak{R}_{J+1}[\check{R}]$ and ${}^{(ext)}\mathfrak{R}_{J+1}[\check{R}]$, we have for any $r_0 \geq 4m_0$

$${}^{(int)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{R}_{J+1}^{\leq r_0}[\check{R}] \lesssim r^{10} B_{-2}^J[\alpha, \beta, \check{\rho}, \underline{\beta}, \underline{\alpha}](1, \tau_*).$$

Together with Proposition 8.3.7, this implies

$${}^{(int)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{R}_{J+1}^{\leq r_0}[\check{R}] \lesssim r_0^{10} \left((\epsilon_B[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \right).$$

Since we have

$${}^{(int)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}] = {}^{(int)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{R}_{J+1}^{\leq r_0}[\check{R}] + {}^{(ext)}\mathfrak{R}_{J+1}^{\geq r_0}[\check{R}],$$

we deduce for any $r_0 \geq 4m_0$

$${}^{(int)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}] \leq {}^{(ext)}\mathfrak{R}_{J+1}^{\geq r_0}[\check{R}] + O \left(r_0^{10} \left(\epsilon_B[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right) \right).$$

Thus, to prove Proposition 8.3.8, it suffices to establish the following inequality

$${}^{(ext)}\mathfrak{R}_{J+1}^{\geq r_0}[\check{R}] \lesssim r_0^{-\delta_B} {}^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}] + r_0^{10} \left(\epsilon_B[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right).$$

This will follow from r^p weighted estimates for the curvature components.

8.7.1 r -weighted divergence identities for Bianchi pairs

Lemma 8.7.1. *Let $k \geq 1$, let $a_{(1)}$ and $a_{(2)}$ real numbers. We consider the following equations.*

- If $\psi_{(1)}, h_{(1)} \in \mathfrak{s}_k$, $\psi_{(2)}, h_{(2)} \in \mathfrak{s}_{k-1}$, let $(\psi_{(1)}, \psi_{(2)})$ such that

$$\begin{cases} e_3(\psi_{(1)}) + a_{(1)}\underline{\kappa}\psi_{(1)} &= -\underline{\mathfrak{d}}_k^*\psi_{(2)} + h_{(1)}, \\ e_4(\psi_{(2)}) + a_{(2)}\kappa\psi_{(2)} &= \underline{\mathfrak{d}}_k\psi_{(1)} + h_{(2)}, \end{cases} \quad (8.7.1)$$

- If $\psi_{(1)}, h_{(1)} \in \mathfrak{s}_{k-1}$, $\psi_{(2)}, h_{(2)} \in \mathfrak{s}_k$, let $(\psi_{(1)}, \psi_{(2)})$ such that

$$\begin{cases} e_3(\psi_{(1)}) + a_{(1)}\underline{\kappa}\psi_{(1)} &= \underline{\mathfrak{d}}_k\psi_{(2)} + h_{(1)}, \\ e_4(\psi_{(2)}) + a_{(2)}\kappa\psi_{(2)} &= -\underline{\mathfrak{d}}_k^*\psi_{(1)} + h_{(2)}. \end{cases} \quad (8.7.2)$$

Then, the pair $(\psi_{(1)}, \psi_{(2)})$ satisfies for any real number b

$$\begin{aligned} & \mathbf{Div}\left(r^b\psi_{(1)}^2 e_3\right) + \mathbf{Div}\left(r^b\psi_{(2)}^2 e_4\right) - \frac{1}{2}r^b\underline{\kappa}\left(-4a_{(1)} + b + 2\right)\psi_{(1)}^2 + \frac{1}{2}r^b\kappa\left(4a_{(2)} - b - 2\right)\psi_{(2)}^2 \\ &= 2r^b\underline{\mathfrak{d}}_1(\psi_{(1)}\psi_{(2)}) - 2r^b\underline{\omega}\psi_{(1)}^2 - 2r^b\omega\psi_{(2)}^2 + 2r^b\psi_{(1)}h_{(1)} + 2r^b\psi_{(2)}h_{(2)} \\ & \quad + br^{b-1}\left(e_3(r) - \frac{r}{2}\underline{\kappa}\right)\psi_{(1)}^2 + br^{b-1}\left(e_4(r) - \frac{r}{2}\kappa\right)\psi_{(2)}^2. \end{aligned} \quad (8.7.3)$$

Remark 8.7.2. *Note that the Bianchi identities can be written as systems of equations of the type (8.7.1) (8.7.2). In particular*

- the Bianchi pair (α, β) satisfies (8.7.1) with $k = 2$, $a_{(1)} = \frac{1}{2}$, $a_{(2)} = 2$,
- the Bianchi pair (β, ρ) satisfies (8.7.1) with $k = 1$, $a_{(1)} = 1$, $a_{(2)} = \frac{3}{2}$,
- the Bianchi pair $(\rho, \underline{\beta})$ satisfies (8.7.2) with $k = 1$, $a_{(1)} = \frac{3}{2}$, $a_{(2)} = 1$,
- the Bianchi pair $(\underline{\beta}, \underline{\alpha})$ satisfies (8.7.2) with $k = 2$, $a_{(1)} = 2$, $a_{(2)} = \frac{1}{2}$.

Proof of Lemma 8.7.1. The proof being identical for (8.7.1) and (8.7.2), it suffices to prove it in the case where $(\psi_{(1)}, \psi_{(2)})$ satisfies (8.7.1).

We compute

$$\begin{aligned} \mathbf{D}_\gamma e_4^\gamma &= -\frac{1}{2}\mathbf{g}(\mathbf{D}_4 e_4, e_3) - \frac{1}{2}\mathbf{g}(\mathbf{D}_3 e_4, e_4) + \mathbf{g}(D_\theta e_4, e_\theta) + \mathbf{g}(D_\varphi e_4, e_\varphi) \\ &= \kappa - 2\omega \end{aligned}$$

and

$$\begin{aligned}\mathbf{D}_\gamma e_3^\gamma &= -\frac{1}{2}\mathbf{g}(\mathbf{D}_4 e_3, e_3) - \frac{1}{2}\mathbf{g}(\mathbf{D}_3 e_3, e_4) + \mathbf{g}(D_\theta e_3, e_\theta) + \mathbf{g}(D_\varphi e_3, e_\varphi) \\ &= \underline{\kappa} - 2\underline{\omega}.\end{aligned}$$

We infer in view of (8.7.1)

$$\begin{aligned}&\mathbf{D}_\gamma \left(r^b \psi_{(1)}^2 e_3^\gamma \right) \\ &= 2r^b \psi_{(1)} e_3(\psi_{(1)}) + br^{b-1} e_3(r) \psi_{(1)}^2 + r^b \psi_{(1)}^2 \mathbf{D}_\gamma e_3^\gamma \\ &= 2r^b \psi_{(1)} \left(-a_{(1)} \underline{\kappa} \psi_{(1)} - \underline{\mathfrak{d}}_k^* \psi_{(2)} + h_{(1)} \right) + br^{b-1} e_3(r) \psi_{(1)}^2 + r^b \psi_{(1)}^2 (\underline{\kappa} - 2\underline{\omega}) \\ &= -2r^b \psi_{(1)} \underline{\mathfrak{d}}_k^* \psi_{(2)} + r^b \left(-2a_{(1)} + \frac{b}{2} + 1 \right) \underline{\kappa} \psi_{(1)}^2 + br^{b-1} \left(e_3(r) - \frac{r}{2} \underline{\kappa} \right) \psi_{(1)}^2 - 2\underline{\omega} r^b \psi_{(1)}^2 + 2r^b \psi_{(1)} h_{(1)}\end{aligned}$$

and

$$\begin{aligned}&\mathbf{D}_\gamma \left(r^b \psi_{(2)}^2 e_4^\gamma \right) \\ &= 2r^b \psi_{(2)} e_4(\psi_{(2)}) + br^{b-1} e_4(r) \psi_{(2)}^2 + r^b \psi_{(2)}^2 \mathbf{D}_\gamma e_4^\gamma \\ &= 2r^b \psi_{(2)} \left(-a_{(2)} \kappa \psi_{(2)} + \underline{\mathfrak{d}}_k \psi_{(1)} + h_{(2)} \right) + br^{b-1} e_4(r) \psi_{(2)}^2 + r^b \psi_{(2)}^2 (\kappa - 2\omega) \\ &= 2r^b \psi_{(2)} \underline{\mathfrak{d}}_k \psi_{(1)} + r^b \left(-2a_{(2)} + \frac{b}{2} + 1 \right) \kappa \psi_{(2)}^2 + br^{b-1} \left(e_4(r) - \frac{r}{2} \kappa \right) \psi_{(2)}^2 - 2r^b \omega \psi_{(2)}^2 + 2r^b \psi_{(2)} h_{(2)}.\end{aligned}$$

We sum the two identities

$$\begin{aligned}&\mathbf{D}_\gamma \left(r^b \psi_{(1)}^2 e_3^\gamma \right) + \mathbf{D}_\gamma \left(r^b \psi_{(2)}^2 e_4^\gamma \right) \\ &= -2r^b \psi_{(1)} \underline{\mathfrak{d}}_k^* \psi_{(2)} + 2r^b \psi_{(2)} \underline{\mathfrak{d}}_k \psi_{(1)} + r^b \left(-2a_{(1)} + \frac{b}{2} + 1 \right) \underline{\kappa} \psi_{(1)}^2 + r^b \left(-2a_{(2)} + \frac{b}{2} + 1 \right) \kappa \psi_{(2)}^2 \\ &\quad + br^{b-1} \left(e_3(r) - \frac{r}{2} \underline{\kappa} \right) \psi_{(1)}^2 + br^{b-1} \left(e_4(r) - \frac{r}{2} \kappa \right) \psi_{(2)}^2 - 2r^b \underline{\omega} \psi_{(1)}^2 - 2r^b \omega \psi_{(2)}^2 \\ &\quad + 2r^b \psi_{(2)} h_{(2)} + 2r^b \psi_{(1)} h_{(1)}\end{aligned}$$

and hence

$$\begin{aligned}&\mathbf{D}_\gamma \left(r^b \psi_{(1)}^2 e_3^\gamma \right) + \mathbf{D}_\gamma \left(r^b \psi_{(2)}^2 e_4^\gamma \right) - r^b \underline{\kappa} \left(-2a_{(1)} + \frac{b}{2} + 1 \right) \psi_{(1)}^2 + r^b \kappa \left(2a_{(2)} - \frac{b}{2} - 1 \right) \psi_{(2)}^2 \\ &= 2r^b \underline{\mathfrak{d}}_1(\psi_{(1)} \psi_{(2)}) + br^{b-1} \left(e_3(r) - \frac{r}{2} \underline{\kappa} \right) \psi_{(1)}^2 + br^{b-1} \left(e_4(r) - \frac{r}{2} \kappa \right) \psi_{(2)}^2 - 2r^b \underline{\omega} \psi_{(1)}^2 - 2r^b \omega \psi_{(2)}^2 \\ &\quad + 2r^b \psi_{(1)} h_{(1)} + 2r^b \psi_{(2)} h_{(2)}.\end{aligned}$$

This concludes the proof of Lemma 8.7.1. □

To obtain r^p weighted estimates for higher order derivatives of the curvature components, we will need several lemmas.

Lemma 8.7.3. *Let $k \geq 1$ and $s \geq 1$ two integers. Let $\psi_{(1)} \in \mathfrak{s}_k$ and $\psi_{(2)} \in \mathfrak{s}_{k-1}$. Then, we have*

$$-\mathfrak{D}^s \psi_{(1)} \mathfrak{D}^s \mathfrak{d}_k^* \psi_{(2)} + \mathfrak{D}^s \psi_{(2)} \mathfrak{D}^s \mathfrak{d}_k \psi_{(1)} = \mathfrak{d}_1 \left(\mathfrak{D}^s \psi_{(1)} \mathfrak{D}^s \psi_{(2)} \right) + E[\mathfrak{D}, s, k, \psi_{(1)}, \psi_{(2)}]$$

where

$$\begin{aligned} |E[s, k, \psi_{(1)}, \psi_{(2)}]| &\lesssim r |\mathfrak{D}^s \psi_{(1)}| \sum_{j=0}^{s-1} |\mathfrak{D}^{s-1-j}(\psi_{(2)})| |\mathfrak{D}^j(K)| \\ &\quad + r |\mathfrak{D}^s \psi_{(2)}| \sum_{j=0}^{s-1} |\mathfrak{D}^{s-1-j}(\psi_{(1)})| |\mathfrak{D}^j(K)|. \end{aligned}$$

Proof. Recall our definition \mathfrak{D}^s for higher angular derivatives. Given f a k -reduced scalar and s a positive integer we define,

$$\mathfrak{D}^s f = \begin{cases} r^{2p} \mathfrak{A}_k^p, & \text{if } s = 2p, \\ r^{2p+1} \mathfrak{d}_k \mathfrak{A}_k^p, & \text{if } s = 2p + 1. \end{cases}$$

We start with the case $s = 2p$, i.e. s is even. Since $\psi_{(1)} \in \mathfrak{s}_k$ and $\psi_{(2)} \in \mathfrak{s}_{k-1}$, we have

$$\begin{aligned} &-\mathfrak{D}^s \psi_{(1)} \mathfrak{D}^s \mathfrak{d}_k^* \psi_{(2)} + \mathfrak{D}^s \psi_{(2)} \mathfrak{D}^s \mathfrak{d}_k \psi_{(1)} \\ &= r^{4p} \left(-\mathfrak{A}_k^p \psi_{(1)} \mathfrak{A}_k^p \mathfrak{d}_k^* \psi_{(2)} + \mathfrak{A}_{k-1}^p \psi_{(2)} \mathfrak{A}_{k-1}^p \mathfrak{d}_k \psi_{(1)} \right). \end{aligned}$$

Next, recall the commutation formulas

$$\begin{aligned} -\mathfrak{d}_k \mathfrak{A}_k + \mathfrak{A}_{k-1} \mathfrak{d}_k &= K \mathfrak{d}_k - k e_\theta(K), \\ -\mathfrak{d}_k^* \mathfrak{A}_{k-1} + \mathfrak{A}_k \mathfrak{d}_k^* &= (2k - 1) K \mathfrak{d}_k^* + (k - 1) e_\theta(K). \end{aligned}$$

We infer

$$\begin{aligned} \mathfrak{A}_{k-1}^p \mathfrak{d}_k &= \mathfrak{d}_k \mathfrak{A}_k^p + \sum_{j=1}^p \mathfrak{A}_{k-1}^{p-j} \left(\mathfrak{A}_{k-1} \mathfrak{d}_k - \mathfrak{d}_k \mathfrak{A}_k \right) \mathfrak{A}_k^{j-1} \\ &= \mathfrak{d}_k \mathfrak{A}_k^p + \sum_{j=1}^p \mathfrak{A}_{k-1}^{p-j} \left(K \mathfrak{d}_k - k e_\theta(K) \right) \mathfrak{A}_k^{j-1} \end{aligned}$$

and

$$\begin{aligned}\mathbb{A}_k^p \mathbb{d}_k^* &= \mathbb{d}_k^* \mathbb{A}_{k-1}^p + \sum_{j=1}^p \mathbb{A}_k^{p-j} \left(\mathbb{A}_k \mathbb{d}_k^* - \mathbb{d}_k^* \mathbb{A}_{k-1} \right) \mathbb{A}_{k-1}^{j-1} \\ &= \mathbb{d}_k^* \mathbb{A}_{k-1}^p + \sum_{j=1}^p \mathbb{A}_k^{p-j} \left((2k-1)K \mathbb{d}_k^* + (k-1)e_\theta(K) \right) \mathbb{A}_{k-1}^{j-1}\end{aligned}$$

This yields

$$\begin{aligned}& -\mathfrak{P}^s \psi_{(1)} \mathfrak{P}^s \mathbb{d}_k^* \psi_{(2)} + \mathfrak{P}^s \psi_{(2)} \mathfrak{P}^s \mathbb{d}_k \psi_{(1)} \\ &= r^{4p} \left\{ -\mathbb{A}_k^p \psi_{(1)} \mathbb{d}_k^* \mathbb{A}_{k-1}^p \psi_{(2)} - \sum_{j=1}^p \mathbb{A}_k^p \psi_{(1)} \mathbb{A}_k^{p-j} \left((2k-1)K \mathbb{d}_k^* + (k-1)e_\theta(K) \right) \mathbb{A}_{k-1}^{j-1} \psi_{(2)} \right. \\ &\quad \left. + \mathbb{A}_{k-1}^p \psi_{(2)} \mathbb{d}_k \mathbb{A}_k^p \psi_{(1)} + \sum_{j=1}^p \mathbb{A}_{k-1}^p \psi_{(2)} \mathbb{A}_{k-1}^{p-j} \left(K \mathbb{d}_k - k e_\theta(K) \right) \mathbb{A}_k^{j-1} \psi_{(1)} \right\} \\ &= \mathbb{d}_1 \left(\mathfrak{P}^s \psi_{(1)} \mathfrak{P}^s \psi_{(2)} \right) \\ &\quad - \sum_{j=1}^p \mathfrak{P}^s \psi_{(1)} \mathfrak{P}^{s-2j} \left((2k-1)r^2 K \mathbb{d}_k^* + (k-1)r^2 e_\theta(K) \right) \mathfrak{P}^{2j-2} \psi_{(2)} \\ &\quad + \sum_{j=1}^p \mathfrak{P}^s \psi_{(2)} \mathfrak{P}^{s-2j} \left(r^2 K \mathbb{d}_k - k r^2 e_\theta(K) \right) \mathfrak{P}^{2j-2} \psi_{(1)}.\end{aligned}$$

Hence, we infer

$$-\mathfrak{P}^s \psi_{(1)} \mathfrak{P}^s \mathbb{d}_k^* \psi_{(2)} + \mathfrak{P}^s \psi_{(2)} \mathfrak{P}^s \mathbb{d}_k \psi_{(1)} = \mathbb{d}_1 \left(\mathfrak{P}^s \psi_{(1)} \mathfrak{P}^s \psi_{(2)} \right) + E[s, k, \psi_{(1)}, \psi_{(2)}]$$

where

$$\begin{aligned}|E[s, k, \psi_{(1)}, \psi_{(2)}]| &\lesssim r^2 |\mathfrak{P}^s \psi_{(1)}| \sum_{j=0}^{s-1} |\mathfrak{P}^{s-1-j}(\psi_{(2)})| |\mathfrak{P}^j(K)| \\ &\quad + r^2 |\mathfrak{P}^s \psi_{(2)}| \sum_{j=0}^{s-1} |\mathfrak{P}^{s-1-j}(\psi_{(1)})| |\mathfrak{P}^j(K)|.\end{aligned}$$

Next, we deal with the case $s = 2p + 1$, i.e. s odd. Since $\psi_{(1)} \in \mathfrak{s}_k$ and $\psi_{(2)} \in \mathfrak{s}_{k-1}$, we have

$$\begin{aligned}& -\mathfrak{P}^s \psi_{(1)} \mathfrak{P}^s \mathbb{d}_k^* \psi_{(2)} + \mathfrak{P}^s \psi_{(2)} \mathfrak{P}^s \mathbb{d}_k \psi_{(1)} \\ &= r^{4p} \left(-\mathbb{d}_k \mathbb{A}_k^p \psi_{(1)} \mathbb{d}_k \mathbb{A}_k^p \mathbb{d}_k^* \psi_{(2)} + \mathbb{d}_{k-1} \mathbb{A}_{k-1}^p \psi_{(2)} \mathbb{d}_{k-1} \mathbb{A}_{k-1}^p \mathbb{d}_k \psi_{(1)} \right)\end{aligned}$$

In view of the case $s = 2p$ above, we infer

$$\begin{aligned}
& -\mathfrak{P}^s \psi_{(1)} \mathfrak{P}^s \mathfrak{d}_k^* \psi_{(2)} + \mathfrak{P}^s \psi_{(2)} \mathfrak{P}^s \mathfrak{d}_k \psi_{(1)} \\
= & r^{4p+2} \left\{ -\mathfrak{d}_k \mathbb{A}_k^p \psi_{(1)} \mathfrak{d}_k \mathfrak{d}_k^* \mathbb{A}_{k-1}^p \psi_{(2)} - \sum_{j=1}^p \mathfrak{d}_k \mathbb{A}_k^p \psi_{(1)} \mathfrak{d}_k \mathbb{A}_k^{p-j} \left((2k-1)K \mathfrak{d}_k^* + (k-1)e_\theta(K) \right) \mathbb{A}_{k-1}^{j-1} \psi_{(2)} \right. \\
& \left. + \mathfrak{d}_{k-1} \mathbb{A}_{k-1}^p \psi_{(2)} \mathfrak{d}_{k-1} \mathfrak{d}_k \mathbb{A}_k^p \psi_{(1)} + \sum_{j=1}^p \mathfrak{d}_{k-1} \mathbb{A}_{k-1}^p \psi_{(2)} \mathfrak{d}_{k-1} \mathbb{A}_{k-1}^{p-j} \left(K \mathfrak{d}_k - k e_\theta(K) \right) \mathbb{A}_k^{j-1} \psi_{(1)} \right\}.
\end{aligned}$$

Next, recall the commutation formula

$$\mathfrak{d}_k \mathfrak{d}_k^* - \mathfrak{d}_{k-1}^* \mathfrak{d}_{k-1} = -2(k-1)K.$$

We infer

$$\begin{aligned}
& -\mathfrak{P}^s \psi_{(1)} \mathfrak{P}^s \mathfrak{d}_k^* \psi_{(2)} + \mathfrak{P}^s \psi_{(2)} \mathfrak{P}^s \mathfrak{d}_k \psi_{(1)} \\
= & r^{4p+2} \left\{ -\mathfrak{d}_k \mathbb{A}_k^p \psi_{(1)} \left(\mathfrak{d}_{k-1}^* \mathfrak{d}_{k-1} - 2(k-1)K \right) \mathbb{A}_{k-1}^p \psi_{(2)} \right. \\
& - \sum_{j=1}^p \mathfrak{d}_k \mathbb{A}_k^p \psi_{(1)} \mathfrak{d}_k \mathbb{A}_k^{p-j} \left((2k-1)K \mathfrak{d}_k^* + (k-1)e_\theta(K) \right) \mathbb{A}_{k-1}^{j-1} \psi_{(2)} \\
& \left. + \mathfrak{d}_{k-1} \mathbb{A}_{k-1}^p \psi_{(2)} \mathfrak{d}_{k-1} \mathfrak{d}_k \mathbb{A}_k^p \psi_{(1)} + \sum_{j=1}^p \mathfrak{d}_{k-1} \mathbb{A}_{k-1}^p \psi_{(2)} \mathfrak{d}_{k-1} \mathbb{A}_{k-1}^{p-j} \left(K \mathfrak{d}_k - k e_\theta(K) \right) \mathbb{A}_k^{j-1} \psi_{(1)} \right\} \\
= & \mathfrak{d}_1 \left(\mathfrak{P}^s \psi_{(1)} \mathfrak{P}^s \psi_{(2)} \right) \\
& + 2(k-1)r^2 K \mathfrak{P}^s \psi_{(1)} \mathfrak{P}^{s-1} \psi_{(2)} - \sum_{j=1}^p \mathfrak{P}^s \psi_{(1)} \mathfrak{P}^{s-2j} \left((2k-1)r^2 K \mathfrak{d}_k^* + (k-1)r^2 e_\theta(K) \right) \mathfrak{P}^{2j-2} \psi_{(2)} \\
& + \sum_{j=1}^p \mathfrak{P}^s \psi_{(2)} \mathfrak{P}^{s-2j} \left(r^2 K \mathfrak{d}_k - k r^2 e_\theta(K) \right) \mathfrak{P}^{2j-2} \psi_{(1)}.
\end{aligned}$$

Hence, we obtain

$$-\mathfrak{P}^s \psi_{(1)} \mathfrak{P}^s \mathfrak{d}_k^* \psi_{(2)} + \mathfrak{P}^s \psi_{(2)} \mathfrak{P}^s \mathfrak{d}_k \psi_{(1)} = \mathfrak{d}_1 \left(\mathfrak{P}^s \psi_{(1)} \mathfrak{P}^s \psi_{(2)} \right) + E[\mathfrak{P}, s, k, \psi_{(1)}, \psi_{(2)}]$$

where

$$\begin{aligned}
|E[s, k, \psi_{(1)}, \psi_{(2)}]| & \lesssim r^2 |\mathfrak{P}^s \psi_{(1)}| \sum_{j=0}^{s-1} |\mathfrak{P}^{s-1-j}(\psi_{(2)})| |\mathfrak{P}^j(K)| \\
& \quad + r^2 |\mathfrak{P}^s \psi_{(2)}| \sum_{j=0}^{s-1} |\mathfrak{P}^{s-1-j}(\psi_{(1)})| |\mathfrak{P}^j(K)|.
\end{aligned}$$

This concludes the proof of the lemma. \square

Corollary 8.7.4. *Let $k \geq 1$, let $a_{(1)}$ and $a_{(2)}$ real numbers and let $0 \leq s \leq k_{large}$. Consider the outgoing geodesic foliation of $^{(ext)}\mathcal{M}$. We consider the following equations.*

- If $\psi_{(1)} \in \mathfrak{s}_k$, $\psi_{(2)} \in \mathfrak{s}_{k-1}$, let $(\psi_{(1)}, \psi_{(2,s)})$ such that

$$\begin{cases} e_3(\not\partial^s \psi_{(1)}) + a_{(1)\underline{\kappa}} \not\partial^s \psi_{(1)} &= -\not\partial^s \not\partial_k^* \psi_{(2)} + h_{(1,s)}, \\ e_4(\not\partial^s \psi_{(2)}) + a_{(2)\kappa} \not\partial^s \psi_{(2)} &= \not\partial^s \not\partial_k \psi_{(1)} + h_{(2,s)}, \end{cases}$$

- If $\psi_{(1)} \in \mathfrak{s}_{k-1}$, $\psi_{(2)}, h_{(2)} \in \mathfrak{s}_k$, let $(\psi_{(1)}, \psi_{(2)})$ such that

$$\begin{cases} e_3(\not\partial^s \psi_{(1)}) + a_{(1)\underline{\kappa}} \not\partial^s \psi_{(1)} &= \not\partial^s \not\partial_k \psi_{(2)} + h_{(1,s)}, \\ e_4(\not\partial^s \psi_{(2)}) + a_{(2)\kappa} \not\partial^s \psi_{(2)} &= -\not\partial^s \not\partial_k^* \psi_{(1)} + h_{(2,s)}. \end{cases}$$

Then, the pair $(\psi_{(1)}, \psi_{(2)})$ satisfies for any real number b

$$\begin{aligned} & \mathbf{Div}\left(r^b(\not\partial^s \psi_{(1)})^2 e_3\right) + \mathbf{Div}\left(r^b(\not\partial^s \psi_{(2)})^2 e_4\right) \\ & - \frac{1}{2}r^b \underline{\kappa} \left(-4a_{(1)} + b + 2\right) (\not\partial^s \psi_{(1)})^2 + \frac{1}{2}r^b \kappa \left(4a_{(2)} - b - 2\right) (\not\partial^s \psi_{(2)})^2 \\ & = 2r^b \not\partial_1 \left(\not\partial^s \psi_{(1)} \not\partial^s \psi_{(2)}\right) + 2r^b E[\not\partial, s, k, \psi_{(1)}, \psi_{(2)}] - 2r^b \underline{\omega} (\not\partial^s \psi_{(1)})^2 \\ & + 2r^b \not\partial^s \psi_{(1)} h_{(1,s)} + 2r^b \not\partial^s \psi_{(2)} h_{(2,s)} + br^{b-1} \left(e_3(r) - \frac{r}{2}\underline{\kappa}\right) (\not\partial^s \psi_{(1)})^2 \\ & + br^{b-1} \left(e_4(r) - \frac{r}{2}\kappa\right) (\not\partial^s \psi_{(2)})^2. \end{aligned}$$

where $E[\not\partial, s, k, \psi_{(1)}, \psi_{(2)}]$ has been introduced in Lemma 8.7.3.

Proof. The proof follows immediately from combining Lemma 8.7.1 and Lemma 8.7.3. \square

Lemma 8.7.5. *Let j, k, l three integers. Consider a Bianchi $(\psi_{(1)}, \psi_{(2)})$ satisfying (8.7.1) or (8.7.2). Then, the pair $(\psi_{(1)}, \psi_{(2)})$ satisfies for any real number b*

$$\begin{aligned} & \mathbf{Div}\left(r^b(\not\partial^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 e_3\right) + \mathbf{Div}\left(r^b(\not\partial^j(re_4)^k \mathbf{T}^l \psi_{(2)})^2 e_4\right) \\ & - \frac{1}{2}r^b \underline{\kappa} \left(-4a_{(1)} + 2k + b + 2\right) (\not\partial^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 \\ & + \frac{1}{2}r^b \kappa \left(4a_{(2)} - 2k - b - 2\right) (\not\partial^j(re_4)^k \mathbf{T}^l \psi_{(2)})^2 \\ & = 2r^b \not\partial_1 \left(\not\partial^j(re_4)^k \mathbf{T}^l \psi_{(1)} \not\partial^j(re_4)^k \mathbf{T}^l \psi_{(2)}\right) + 2r^b E[\not\partial, j, k, (re_4)^k \mathbf{T}^l \psi_{(1)}, (re_4)^k \mathbf{T}^l \psi_{(2)}] \\ & - 2r^b \underline{\omega} (\not\partial^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 + 2r^b \not\partial^j(re_4)^k \mathbf{T}^l \psi_{(1)} h_{(1),j,k,l} + 2r^b \not\partial^j(re_4)^k \mathbf{T}^l \psi_{(2)} h_{(2),j,k,l} \\ & + br^{b-1} \left(e_3(r) - \frac{r}{2}\underline{\kappa}\right) (\not\partial^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 + br^{b-1} \left(e_4(r) - \frac{r}{2}\kappa\right) (\not\partial^j(re_4)^k \mathbf{T}^l \psi_{(2)})^2. \end{aligned}$$

where $E[\mathfrak{J}, s, k, (re_4)^k \mathbf{T}^l \psi_{(1)}, (re_4)^k \mathbf{T}^l \psi_{(2)}]$ has been introduced in Lemma 8.7.3, and where $h_{(1),j,k,l}$ and $h_{(2),j,k,l}$ are given, schematically, by

$$\begin{aligned} h_{(1),j,k,l} &= \mathfrak{J}^{\leq j+k+l}(h_{(1)}) + kr^{-1} \mathfrak{J}^{j+1}(re_4)^{k-1} \mathbf{T}^l \psi_{(2)} \\ &\quad + r \mathfrak{D}^{j+k+l} \left(\Gamma_g(\psi_{(1)}, \psi_{(2)}) \right) + O(r^{-1}) \mathfrak{D}^{\leq j+k+l-1}(\psi_{(1)}, \psi_{(2)}) \end{aligned}$$

and

$$\begin{aligned} h_{(2),j,k,l} &= \mathfrak{J}^{\leq j+k+l}(h_{(2)}) + kr^{-1} \mathfrak{J}^{j+1}(re_4)^{k-1} \mathbf{T}^l \psi_{(1)} + r \mathfrak{D}^{j+k+l} \left(\Gamma_g(\psi_{(1)}, \psi_{(2)}) \right) \\ &\quad + O(r^{-1}) \mathfrak{D}^{\leq j+k+l-1}(\psi_{(1)}, \psi_{(2)}). \end{aligned}$$

Proof. We have the following simple schematic consequences of the commutator identities

$$\begin{aligned} [T, e_4], [T, e_3] &= r^{-1} \Gamma_b \mathfrak{D}, \quad [T, \mathfrak{A}_k] = -\eta e_3 + \Gamma_g \mathfrak{D}, \\ [\mathfrak{J}, e_4] &= \Gamma_g \mathfrak{D} + \Gamma_g, \quad [\mathfrak{J}, e_3] = -r\eta e_3 + r\Gamma_g \mathfrak{D}, \\ [re_4, e_4] &= -\frac{r}{2} \kappa e_4 + \Gamma_g \mathfrak{D}, \quad [re_4, e_3] = -\frac{r}{2} \underline{\kappa} e_4 + \Gamma_b \mathfrak{D}, \quad [re_4, \mathfrak{J}_k] = r^{-1} \mathfrak{J} + \Gamma_g \mathfrak{D} + \Gamma_g. \end{aligned}$$

Then, differentiating with $\mathfrak{J}^j (re_4)^k \mathbf{T}^l$ the equations

$$\begin{cases} e_3(\psi_{(1)}) + a_{(1)} \underline{\kappa} \psi_{(1)} &= -\mathfrak{A}_k^* \psi_{(2)} + h_{(1)}, \\ e_4(\psi_{(2)}) + a_{(2)} \kappa \psi_{(2)} &= \mathfrak{A}_k \psi_{(1)} + h_{(2)}, \end{cases}$$

and using the above commutator identities we infer

$$\begin{cases} e_3(\mathfrak{J}^j (re_4)^k \mathbf{T}^l \psi_{(1)}) + (a_{(1)} - \frac{\kappa}{2}) \underline{\kappa} \mathfrak{J}^j (re_4)^k \mathbf{T}^l \psi_{(1)} &= -\mathfrak{J}^j \mathfrak{A}_k^* ((re_4)^k \mathbf{T}^l \psi_{(2)}) + h_{(1),j,k,l}, \\ e_4(\mathfrak{J}^j (re_4)^k \mathbf{T}^l \psi_{(2)}) + (a_{(2)} - \frac{\kappa}{2}) \kappa \mathfrak{J}^j (re_4)^k \mathbf{T}^l \psi_{(2)} &= \mathfrak{J}^j \mathfrak{A}_k ((re_4)^k \mathbf{T}^l \psi_{(1)}) + h_{(2),j,k,l}, \end{cases}$$

were

$$\begin{aligned} h_{(1),j,k,l} &= \mathfrak{J}^j (re_4)^k \mathbf{T}^l (h_{(1)}) + kr^{-1} \mathfrak{J}^{j+1} (re_4)^{k-1} \mathbf{T}^l \psi_{(2)} + jr\eta \mathfrak{D}^{j+k+l-1} e_3 \psi_{(1)} \\ &\quad + r \mathfrak{D}^{j+k+l} \left(\Gamma_g(\psi_{(1)}, \psi_{(2)}) \right) + O(r^{-1}) \mathfrak{D}^{\leq j+k+l-1}(\psi_{(1)}, \psi_{(2)}) \end{aligned}$$

and

$$\begin{aligned} h_{(2),j,k,l} &= \mathfrak{J}^j (re_4)^k \mathbf{T}^l (h_{(2)}) + kr^{-1} \mathfrak{J}^{j+1} (re_4)^{k-1} \mathbf{T}^l \psi_{(1)} + r \mathfrak{D}^{j+k+l} \left(\Gamma_g(\psi_{(1)}, \psi_{(2)}) \right) \\ &\quad + O(r^{-1}) \mathfrak{D}^{\leq j+k+l-1}(\psi_{(1)}, \psi_{(2)}). \end{aligned}$$

Also, using the equation

$$e_3(\psi_{(1)}) = -a_{(1)} \underline{\kappa} \psi_{(1)} - \mathfrak{A}_k^* \psi_{(2)} + h_{(1)},$$

we obtain

$$jr\eta\mathfrak{d}^{j+k+l-1}e_3\psi_{(1)} = r\mathfrak{d}^{j+k+l}\left(\Gamma_g(\psi_{(1)},\psi_{(2)})\right) + O(r^{-1})\mathfrak{d}^{\leq j+k+l-1}(\psi_{(1)},\psi_{(2)}) + r\eta\mathfrak{d}^{k+j+l-1}(h_{(1)})$$

and hence,

$$\begin{aligned} h_{(1),j,k,l} &= \mathfrak{d}^{\leq j+k+l}(h_{(1)}) + kr^{-1}\mathfrak{d}^{j+1}(re_4)^{k-1}\mathbf{T}^l\psi_{(2)} \\ &\quad + r\mathfrak{d}^{j+k+l}\left(\Gamma_g(\psi_{(1)},\psi_{(2)})\right) + O(r^{-1})\mathfrak{d}^{\leq j+k+l-1}(\psi_{(1)},\psi_{(2)}). \end{aligned}$$

We have thus obtained the desired form for $h_{(1),j,k,l}$ and $h_{(2),j,k,l}$.

The divergence identity now follows from the equations

$$\begin{cases} e_3(\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(1)}) + (a_{(1)} - \frac{k}{2})\underline{\kappa}\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(1)} = -\mathfrak{d}^j\mathfrak{d}_k^*(re_4)^k\mathbf{T}^l\psi_{(2)} + h_{(1),j,k,l}, \\ e_4(\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(2)}) + (a_{(2)} - \frac{k}{2})\kappa\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(2)} = \mathfrak{d}^j\mathfrak{d}_k((re_4)^k\mathbf{T}^l\psi_{(1)}) + h_{(2),j,k,l}, \end{cases}$$

together with Corollary 8.7.4. This concludes the proof of the lemma. \square

Corollary 8.7.6. *Let $r_0 \geq 4m_0$ and $1 \leq u_0 \leq u_*$. We introduce the spacetime region*

$$\mathcal{R}_{u_0} = {}^{(ext)}\mathcal{M} \cap \{r \geq 4m_0\} \cap \{1 \leq u \leq u_0\},$$

Let j, k, l three integers. Assume that the frame of ${}^{(ext)}\mathcal{M}$ satisfies

$$\sup_{{}^{(ext)}\mathcal{M}} \left(\left| e_3(r) - \frac{r}{2}\underline{\kappa} \right| + r \left(|\underline{\omega}| + \left| e_4(r) - \frac{r}{2}\kappa \right| \right) \right) \lesssim \epsilon_0.$$

Consider a pair $(\psi_{(1)}, \psi_{(2)})$ satisfying (8.7.1) or (8.7.2). Then, $(\psi_{(1)}, \psi_{(2)})$ satisfies for any real number b

(a) If

$$-4a_{(1)} + 2k + b + 2 > 0 \text{ and } 4a_{(2)} - 2k - b - 2 > 0,$$

then, we have

$$\begin{aligned} & \int_{\mathcal{C}_{u_0}(r \geq r_0)} r^b (\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(1)})^2 + \int_{\Sigma_*(\leq u_0)} r^b \left((\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(1)})^2 + (\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(2)})^2 \right) \\ & + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^{b-1} \left((\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(1)})^2 + (\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(2)})^2 \right) \\ & \lesssim \int_{{}^{(ext)}\mathcal{M}(\frac{r_0}{2} \leq r \leq r_0)} r^{b-1} \left((\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(1)})^2 + (\mathfrak{d}^j(re_4)^k\mathbf{T}^l\psi_{(2)})^2 \right) \\ & + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^{b+1} \left((h_{(1),j,k,l})^2 + (h_{(2),j,k,l})^2 \right) \\ & + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^b E[\mathfrak{d}, j, k, (re_4)^k\mathbf{T}^l\psi_{(1)}, (re_4)^k\mathbf{T}^l\psi_{(2)}]. \end{aligned}$$

(b) If

$$-4a_{(1)} + 2k + b + 2 > 0 \text{ and } 4a_{(2)} - 2k - b - 2 = 0,$$

then, we have

$$\begin{aligned} & \int_{\mathcal{C}_{u_0}(r \geq r_0)} r^b (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 + \int_{\Sigma_*(\leq u_0)} r^b \left((\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 + (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(2)})^2 \right) \\ \lesssim & \int_{(ext)\mathcal{M}(\frac{r_0}{2} \leq r \leq r_0)} r^{b-1} \left((\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 + (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 \right) \\ & + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^{b+1-\delta_B} (h_{(1),j,k,l})^2 + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^{b+1+\delta_B} (h_{(2),j,k,l})^2 \\ & + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^{b-1+\delta_B} (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^{b-1-\delta_B} (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(2)})^2 \\ & + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^b E[\mathfrak{P}, j, k, (re_4)^k \mathbf{T}^l \psi_{(1)}, (re_4)^k \mathbf{T}^l \psi_{(2)}]. \end{aligned}$$

(c) If

$$-4a_{(1)} + 2k + b + 2 > 0 \text{ and } 4a_{(2)} - 2k - b - 2 < 0,$$

then, we have

$$\begin{aligned} & \int_{\mathcal{C}_{u_0}(r \geq r_0)} r^b (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 + \int_{\Sigma_*(\leq u_0)} r^b \left((\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 + (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(2)})^2 \right) \\ & + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^{b-1} (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 \\ \lesssim & \int_{(ext)\mathcal{M}(\frac{r_0}{2} \leq r \leq r_0)} r^{b-1} \left((\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 + (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(1)})^2 \right) \\ & + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^{b+1} (h_{(1),j,k,l})^2 + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^{b+1} (h_{(2),j,k,l})^2 \\ & + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^{b-1} \left((\mathfrak{P}^j(re_4)^k \mathbf{T}^l \psi_{(2)})^2 \right) \\ & + \int_{\mathcal{R}_{u_0}(r \geq r_0)} r^b E[\mathfrak{P}, j, k, (re_4)^k \mathbf{T}^l \psi_{(1)}, (re_4)^k \mathbf{T}^l \psi_{(2)}]. \end{aligned}$$

Proof. We multiply the pair $(\psi_{(1)}, \psi_{(2)})$ by a smooth cut-off function in r supported in $r \geq \frac{r_0}{2}$ and identically one for $r \geq r_0$. We obtain again a solution to (8.7.1) or (8.7.2) up to error terms that are supported in the region $\frac{r_0}{2} \leq r \leq r_0$. We then integrate the divergence identities of Lemma 8.7.5 on the region \mathcal{R}_{u_0} and the corollary follows. \square

8.7.2 End of the proof of Proposition 8.3.8

Let $r_0 \geq 4m_0$. Recall that, to prove Proposition 8.3.8, it suffices to establish the following inequality

$${}^{(ext)}\mathfrak{R}_{J+1}^{\geq r_0}[\check{R}] \lesssim r_0^{-\delta_B} {}^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}] + r_0^{10} \left(\epsilon_B[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right).$$

To this end, we will rely on the r^p -weighted estimates derived in Corollary 8.7.6 applied to the Bianchi pairs, where we recall Remark 8.7.2.

Remark 8.7.7. For the Bianchi pair (β, ρ) , we replace the Bianchi identities for $e_4(\rho)$ by its analog for $e_4(\check{\rho})$, i.e.

$$e_4\check{\rho} + \frac{3}{2}\bar{\kappa}\check{\rho} = \not{d}_1\beta - \frac{3}{2}\bar{\rho}\check{\kappa} + Err[e_4\check{\rho}],$$

while for the Bianchi pair $(\rho, \underline{\beta})$, we replace the Bianchi identities for $e_3(\rho)$ by its analog for $e_3(\check{\rho})$, i.e.

$$e_3\check{\rho} + \frac{3}{2}\bar{\kappa}\check{\rho} = \not{d}_1\underline{\beta} - \frac{3}{2}\bar{\rho}\check{\kappa} - \frac{3}{2}\bar{\kappa}\bar{\rho}\varsigma^{-1}\zeta + \frac{3}{2}\bar{\kappa}\bar{\rho}(\check{\Omega} + \varsigma^{-1}\bar{\Omega}\zeta) + Err[e_3\check{\rho}],$$

see Proposition 2.2.18 for the derivation of these equations.

Let j, k, l three integers such that

$$j + k + l = J + 1.$$

To derive r^p weighted curvature estimates for $\not{\partial}^j(re_4)^k\mathbf{T}^l$ derivatives in the region $r \geq r_0$, we proceed as follows.

Step 1. We start with the case $k = 0$, i.e. we derive r^p weighted curvature estimates for $\not{\partial}^j\mathbf{T}^l$ derivatives with $j + l = J + 1$. First, we apply Corollary 8.7.6

- to the Bianchi pair (α, β) with the choice $b = 4 + \delta_B$,
- to the Bianchi pair (β, ρ) with the choice $b = 4 - \delta_B$,
- to the Bianchi pair $(\rho, \underline{\beta})$ with the choice $b = 2 - \delta_B$,
- to the Bianchi pair $(\underline{\beta}, \underline{\alpha})$ with the choice $b = -\delta_B$.

All the above choices are such that we have in case (a) of Corollary 8.7.6. In particular, we obtain

$$\begin{aligned}
& \sum_{j+l=J+1} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} \left(r^{4+\delta_B} (\mathfrak{P}^j \mathbf{T}^l \alpha)^2 + r^{4-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \beta)^2 + r^{2-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 \right. \right. \\
& + r^{-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 \left. \right) + \int_{\Sigma_*} \left(r^{4+\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \alpha)^2 + (\mathfrak{P}^j \mathbf{T}^l \beta)^2 \right) + r^{4-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 \right. \\
& + r^{2-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + r^{-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \left. \right) + \int_{(ext)\mathcal{M}(r \geq r_0)} \left(r^{3+\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \alpha)^2 + (\mathfrak{P}^j \mathbf{T}^l \beta)^2 \right) \right. \\
& \left. \left. + r^{3-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 + r^{1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + r^{-1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \right) \right\} \\
& \lesssim r_0^{8+\delta_B} \int_{(ext)\mathcal{M}(\frac{r_0}{2} \leq r \leq r_0)} \frac{(\mathfrak{D}^{J+1} \check{R})^2}{r^5} + \int_{(ext)\mathcal{M}(r \geq r_0)} \left\{ r^{-1+\delta_B} (\mathfrak{P}^j \mathbf{T}^l \vartheta)^2 \right. \\
& + r^{-1-\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \eta)^2 + (\mathfrak{P}^j \mathbf{T}^l \check{\kappa})^2 \right) + r^{-3-\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \check{\kappa})^2 + (\mathfrak{P}^j \mathbf{T}^l \zeta)^2 \right) \\
& \left. + r^{-5-\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \xi)^2 + (\mathfrak{P}^j \mathbf{T}^l \vartheta)^2 + (\mathfrak{P}^j \mathbf{T}^l \zeta)^2 + (\mathfrak{P}^j \mathbf{T}^l \check{\Omega})^2 \right) \right\} + (\epsilon_B[J])^2 + \epsilon_0^2 (\mathfrak{N}_{J+1}^{(En)})^2.
\end{aligned}$$

Using Proposition 8.3.7 to bound the first term on the right-hand side, and using also the definition of the norm ${}^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}]$, we infer that

$$\begin{aligned}
& \sum_{j+l=J+1} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} \left(r^{4+\delta_B} (\mathfrak{P}^j \mathbf{T}^l \alpha)^2 + r^{4-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \beta)^2 + r^{2-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 \right. \right. \\
& + r^{-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 \left. \right) + \int_{\Sigma_*} \left(r^{4+\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \alpha)^2 + (\mathfrak{P}^j \mathbf{T}^l \beta)^2 \right) + r^{4-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 \right. \\
& + r^{2-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + r^{-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \left. \right) + \int_{(ext)\mathcal{M}(r \geq r_0)} \left(r^{3+\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \alpha)^2 + (\mathfrak{P}^j \mathbf{T}^l \beta)^2 \right) \right. \\
& \left. \left. + r^{3-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 + r^{1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + r^{-1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \right) \right\} \\
& \lesssim \left(\int_{r_0}^{+\infty} \frac{dr}{r^{1+\delta_B}} \right) ({}^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}]) + r_0^{10} \left((\epsilon_B[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \right)
\end{aligned}$$

and hence

$$\begin{aligned}
 & \sum_{j+l=J+1} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} \left(r^{4+\delta_B} (\not\partial^j \mathbf{T}^l \alpha)^2 + r^{4-\delta_B} (\not\partial^j \mathbf{T}^l \beta)^2 + r^{2-\delta_B} (\not\partial^j \mathbf{T}^l \check{\rho})^2 \right. \right. \\
 & \left. \left. + r^{-\delta_B} (\not\partial^j \mathbf{T}^l \underline{\beta})^2 \right) + \int_{\Sigma_*} \left(r^{4+\delta_B} \left((\not\partial^j \mathbf{T}^l \alpha)^2 + (\not\partial^j \mathbf{T}^l \beta)^2 \right) + r^{4-\delta_B} (\not\partial^j \mathbf{T}^l \check{\rho})^2 \right. \right. \\
 & \left. \left. + r^{2-\delta_B} (\not\partial^j \mathbf{T}^l \underline{\beta})^2 + r^{-\delta_B} (\not\partial^j \mathbf{T}^l \underline{\alpha})^2 \right) + \int_{(ext)\mathcal{M}(r \geq r_0)} \left(r^{3+\delta_B} \left((\not\partial^j \mathbf{T}^l \alpha)^2 + (\not\partial^j \mathbf{T}^l \beta)^2 \right) \right. \right. \\
 & \left. \left. + r^{3-\delta_B} (\not\partial^j \mathbf{T}^l \check{\rho})^2 + r^{1-\delta_B} (\not\partial^j \mathbf{T}^l \underline{\beta})^2 + r^{-1-\delta_B} (\not\partial^j \mathbf{T}^l \underline{\alpha})^2 \right) \right\} \\
 & \lesssim r_0^{-\delta_B} {}^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}] + r_0^{10} \left((\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \right). \tag{8.7.4}
 \end{aligned}$$

Step 2. We derive additional r^p weighted curvature estimates for $\not\partial^j \mathbf{T}^l$ derivatives with $j+l=J+1$. To this end, we apply Corollary 8.7.6

- to the Bianchi pair (β, ρ) with the choice $b = 4$,
- to the Bianchi pair $(\rho, \underline{\beta})$ with the choice $b = 2$,
- to the Bianchi pair $(\underline{\beta}, \underline{\alpha})$ with the choice $b = 0$.

All the above choices are such that we have in case (b) of Corollary 8.7.6. In particular,

we obtain

$$\begin{aligned}
& \sum_{j+l=J+1} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} \left(r^4 (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + r^2 (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 + (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 \right) \right. \\
& \left. + \int_{\Sigma_*} \left(r^4 \left((\mathfrak{P}^j \mathbf{T}^l \beta)^2 + (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 \right) + r^2 (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \right) \right\} \\
\lesssim & r_0^8 \int_{(ext)\mathcal{M}(\frac{r_0}{2} \leq r \leq r_0)} \frac{(\mathfrak{d}^{J+1} \check{R})^2}{r^5} \\
& + \sum_{j+l=J+1} \left\{ \int_{(ext)\mathcal{M}(r \geq r_0)} \left(r^{3+\delta_B} (\mathfrak{P}^j \mathbf{T}^l \beta)^2 + r^{3-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 + r^{1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 \right. \right. \\
& \left. \left. + r^{-1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \right) \right\} + \int_{(ext)\mathcal{M}(r \geq r_0)} \left\{ r^{-1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \eta)^2 + r^{-1+\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\kappa})^2 \right. \\
& \left. + r^{-3+\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \check{\kappa})^2 + (\mathfrak{P}^j \mathbf{T}^l \zeta)^2 \right) \right. \\
& \left. + r^{-5+\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \underline{\xi})^2 + (\mathfrak{P}^j \mathbf{T}^l \underline{\vartheta})^2 + (\mathfrak{P}^j \mathbf{T}^l \zeta)^2 + (\mathfrak{P}^j \mathbf{T}^l \check{\Omega})^2 \right) \right\} + (\epsilon_B[J])^2 + \epsilon_0^2 (\mathfrak{N}_{J+1}^{(En)})^2.
\end{aligned}$$

Using Proposition 8.3.7 to bound the first term on the right-hand side, and using also the definition of the norm ${}^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}]$, we infer that

$$\begin{aligned}
& \sum_{j+l=J+1} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} \left(r^4 (\mathfrak{P}^j \mathbf{T}^l \beta)^2 + r^2 (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 + (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 \right) \right. \\
& \left. + \int_{\Sigma_*} \left(r^4 \left((\mathfrak{P}^j \mathbf{T}^l \beta)^2 + (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 \right) + r^2 (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \right) \right\} \\
\lesssim & \left(\int_{r_0}^{+\infty} \frac{dr}{r^{1+\delta_B}} \right) {}^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}] + r_0^{10} \left((\epsilon_B[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \right) \\
& + \sum_{j+l=J+1} \left\{ \int_{(ext)\mathcal{M}(r \geq r_0)} \left(r^{3+\delta_B} (\mathfrak{P}^j \mathbf{T}^l \beta)^2 + r^{3-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 + r^{1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 \right. \right. \\
& \left. \left. + r^{-1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \right) \right\}
\end{aligned}$$

and hence

$$\begin{aligned}
& \sum_{j+l=J+1} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} \left(r^4 (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + r^2 (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 + (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 \right) \right. \\
& \left. + \int_{\Sigma_*} \left(r^4 \left((\mathfrak{P}^j \mathbf{T}^l \beta)^2 + (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 \right) + r^2 (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \right) \right\} \\
\lesssim & r_0^{-\delta_B} (ext) \mathfrak{G}_k^{\geq r_0} [\check{\Gamma}] + r_0^{10} \left((\epsilon_B[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \right) \\
& + \sum_{j+l=J+1} \left\{ \int_{(ext) \mathcal{M}(r \geq r_0)} \left(r^{3+\delta_B} (\mathfrak{P}^j \mathbf{T}^l \beta)^2 + r^{3-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 + r^{1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 \right. \right. \\
& \left. \left. + r^{-1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \right) \right\}.
\end{aligned}$$

Together with (8.7.4), we deduce

$$\begin{aligned}
& \sum_{j+l=J+1} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} \left(r^{4+\delta_B} (\mathfrak{P}^j \mathbf{T}^l \alpha)^2 + r^4 (\mathfrak{P}^j \mathbf{T}^l \beta)^2 + r^2 (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 \right. \right. \\
& \left. \left. + (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 \right) + \int_{\Sigma_*} \left(r^{4+\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \alpha)^2 + (\mathfrak{P}^j \mathbf{T}^l \beta)^2 \right) + r^4 (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 \right. \right. \\
& \left. \left. + r^2 (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \right) + \int_{(ext) \mathcal{M}(r \geq r_0)} \left(r^{3+\delta_B} \left((\mathfrak{P}^j \mathbf{T}^l \alpha)^2 + (\mathfrak{P}^j \mathbf{T}^l \beta)^2 \right) \right. \right. \\
& \left. \left. + r^{3-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \check{\rho})^2 + r^{1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\beta})^2 + r^{-1-\delta_B} (\mathfrak{P}^j \mathbf{T}^l \underline{\alpha})^2 \right) \right\} \\
\lesssim & r_0^{-\delta_B} (ext) \mathfrak{G}_k^{\geq r_0} [\check{\Gamma}] + r_0^{10} \left((\epsilon_B[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \right). \tag{8.7.5}
\end{aligned}$$

Step 3. We now argue by iteration on k . For $0 \leq k \leq J$, we consider the following

iteration assumption

$$\begin{aligned}
& \sum_{j+l=J+1-k} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} \left(r^{4+\delta_B} (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \alpha)^2 + r^4 (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \beta)^2 + r^2 (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \check{\rho})^2 \right. \right. \\
& + (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \underline{\beta})^2 \left. \right) + \int_{\Sigma_*} \left(r^{4+\delta_B} \left((\mathfrak{P}^j(re_4)^k \mathbf{T}^l \alpha)^2 + (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \beta)^2 \right) + r^4 (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \check{\rho})^2 \right. \\
& + r^2 (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \underline{\beta})^2 + (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \underline{\alpha})^2 \left. \right) + \int_{(ext)\mathcal{M}(r \geq r_0)} \left(r^{3+\delta_B} \left((\mathfrak{P}^j(re_4)^k \mathbf{T}^l \alpha)^2 + (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \beta)^2 \right) \right. \\
& \left. \left. + r^{3-\delta_B} (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \check{\rho})^2 + r^{1-\delta_B} (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \underline{\beta})^2 + r^{-1-\delta_B} (\mathfrak{P}^j(re_4)^k \mathbf{T}^l \underline{\alpha})^2 \right) \right\} \\
& \lesssim r_0^{-\delta_B} (ext) \mathfrak{G}_k^{\geq r_0} [\check{\Gamma}] + r_0^{10} \left((\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \right). \tag{8.7.6}
\end{aligned}$$

(8.7.6) holds true for $k = 0$ in view of (8.7.5). We now assume that (8.7.6) holds true for k such that $0 \leq k \leq J$, and our goal is to prove that it also holds for $k + 1$.

First, note that the Bianchi identities for $e_4(\beta)$, $e_4(\check{\rho})$, $e_4(\underline{\beta})$ and $e_4(\underline{\alpha})$, together with (8.7.6), yields

$$\begin{aligned}
& \sum_{j+l=J+1-(k+1)} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} \left(r^4 (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \beta)^2 + r^2 (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \check{\rho})^2 \right. \right. \\
& + (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \underline{\beta})^2 \left. \right) + \int_{\Sigma_*} \left(r^{4+\delta_B} (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \beta)^2 + r^4 (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \check{\rho})^2 \right. \\
& + r^2 (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \underline{\beta})^2 + (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \underline{\alpha})^2 \left. \right) + \int_{(ext)\mathcal{M}(r \geq r_0)} \left(r^{3+\delta_B} (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \beta)^2 \right. \\
& \left. \left. + r^{3-\delta_B} (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \check{\rho})^2 + r^{1-\delta_B} (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \underline{\beta})^2 + r^{-1-\delta_B} (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \underline{\alpha})^2 \right) \right\} \\
& \lesssim r_0^{-\delta_B} (ext) \mathfrak{G}_k^{\geq r_0} [\check{\Gamma}] + r_0^{10} \left((\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \right). \tag{8.7.7}
\end{aligned}$$

We still need to estimate $\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \alpha$. To this end, we apply Corollary 8.7.6 to the Bianchi pair (α, β) with the choice $b = 4 + \delta_B$. Since $k + 1 \geq 1$, we are in case (c) of

Corollary 8.7.6. In particular, we obtain, arguing similarly as above,

$$\begin{aligned}
& \sum_{j+l=J+1-(k+1)} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} r^{4+\delta_B} (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \alpha)^2 + \int_{\Sigma_*} r^{4+\delta_B} (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \alpha)^2 \right. \\
& \left. + \int_{(ext)\mathcal{M}(r \geq r_0)} r^{3+\delta_B} (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \alpha)^2 \right\} \\
\lesssim & \sum_{j+l=J+1-(k+1)} \left\{ \int_{(ext)\mathcal{M}(r \geq r_0)} r^{3+\delta_B} (\mathfrak{P}^j(re_4)^{k+1} \mathbf{T}^l \beta)^2 \right\} + r_0^{-\delta_B} (ext) \mathfrak{G}_k^{\geq r_0}[\check{\Gamma}] \\
& + r_0^{10} \left((\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \right).
\end{aligned}$$

Together with (8.7.7), this implies (8.7.6) for $k+1$. Hence, by iteration, (8.7.6) holds for any $0 \leq k \leq J+1$. This implies

$$\begin{aligned}
& \sum_{k \leq J+1} \left\{ \sup_{1 \leq u \leq u_*} \int_{\mathcal{C}_u(r \geq r_0)} \left(r^{4+\delta_B} (\mathfrak{D}^k \alpha)^2 + r^4 (\mathfrak{D}^k \beta)^2 + r^2 (\mathfrak{D}^k \rho)^2 + (\mathfrak{D}^k \underline{\beta})^2 \right) \right. \\
& + \int_{\Sigma_*} \left(r^{4+\delta_B} \left((\mathfrak{D}^k \alpha)^2 + (\mathfrak{D}^k \beta)^2 \right) + r^4 (\mathfrak{D}^k \rho)^2 + r^2 (\mathfrak{D}^k \underline{\beta})^2 + (\mathfrak{D}^k \underline{\alpha})^2 \right) \\
& \left. + \int_{(ext)\mathcal{M}(r \geq r_0)} \left(r^{3+\delta_B} \left((\mathfrak{D}^k \alpha)^2 + (\mathfrak{D}^k \beta)^2 \right) + r^{3-\delta_B} (\mathfrak{D}^k \rho)^2 + r^{1-\delta_B} (\mathfrak{D}^k \underline{\beta})^2 + r^{-1-\delta_B} (\mathfrak{D}^k \underline{\alpha})^2 \right) \right\} \\
\lesssim & r_0^{-\delta_B} (ext) \mathfrak{G}_k^{\geq r_0}[\check{\Gamma}] + r_0^{10} \left((\epsilon_{\mathcal{B}}[J])^2 + \epsilon_0^2 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right)^2 \right).
\end{aligned}$$

Hence, we have obtained

$$(ext) \mathfrak{R}_{J+1}^{\geq r_0}[\check{R}] \lesssim r_0^{-\delta_B} (ext) \mathfrak{G}_k^{\geq r_0}[\check{\Gamma}] + r_0^{10} \left(\epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right)$$

which concludes the proof of Proposition 8.3.8.

8.8 Proof of Proposition 8.3.9

To prove Proposition 8.3.9, we rely on the following three propositions.

Proposition 8.8.1. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Then, we have*

$$(\Sigma_*) \mathfrak{G}_{J+1}[\check{\Gamma}] + (\Sigma_*) \mathfrak{G}'_{J+1}[\check{\Gamma}] \lesssim (\Sigma_*) \mathfrak{R}_{J+1}[\check{R}] + (\Sigma_*) \mathfrak{G}_J[\check{\Gamma}],$$

where we have introduced the notations

$$\begin{aligned} {}^{(\Sigma^*)}\mathfrak{G}_k[\check{\Gamma}] &:= \int_{\Sigma^*} \left[r^2 \left((\mathfrak{d}^{\leq k} \vartheta)^2 + (\mathfrak{d}^{\leq k} \check{\kappa})^2 + (\mathfrak{d}^{\leq k} \zeta)^2 + (\mathfrak{d}^{\leq k} \check{\underline{\kappa}})^2 \right) + (\mathfrak{d}^{\leq k} \underline{\vartheta})^2 \right. \\ &\quad \left. + (\mathfrak{d}^{\leq k} \eta)^2 + (\mathfrak{d}^{\leq k} \check{\underline{\omega}})^2 + (\mathfrak{d}^{\leq k} \underline{\xi})^2 \right], \end{aligned}$$

$${}^{(\Sigma^*)}\mathfrak{G}'_k[\check{\Gamma}] := \int_{\Sigma^*} \left[r^2 \left((\mathfrak{d}^{k+1} \check{\vartheta})^2 + (\mathfrak{d}^{k+1} \check{\kappa})^2 + (\mathfrak{d}^{\leq k+1} \check{\mu})^2 + (\mathfrak{d}^{k+1} \check{\underline{\kappa}})^2 + (\mathfrak{d}^{k+1} \zeta)^2 \right) \right],$$

and

$${}^{(\Sigma^*)}\mathfrak{R}_k[\check{R}] := \int_{\Sigma^*} \left(r^{4+\delta_B} \left((\mathfrak{d}^{\leq k} \alpha)^2 + (\mathfrak{d}^{\leq k} \beta)^2 \right) + r^4 (\mathfrak{d}^{\leq k} \check{\rho})^2 + r^2 (\mathfrak{d}^{\leq k} \underline{\beta})^2 + (\mathfrak{d}^{\leq k} \underline{\alpha})^2 \right).$$

Proposition 8.8.2. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Then, we have*

$${}^{(ext)}\mathfrak{G}_{J+1}^{\geq 4m_0}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_{J+1}^{\geq 4m_0'}[\check{\Gamma}] \lesssim {}^{(\Sigma^*)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(\Sigma^*)}\mathfrak{G}'_{J+1}[\check{\Gamma}] + {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{G}_J[\check{\Gamma}],$$

where we have introduced the notation

$$\begin{aligned} {}^{(ext)}\mathfrak{G}_k^{\geq 4m_0'}[\check{\Gamma}] &:= \sup_{\lambda \geq 4m_0} \left(\int_{\{r=\lambda\}} \left[\lambda^6 \left(\mathfrak{d}^k \left(\mathfrak{d}_1 \mathfrak{d}_1^* \kappa - \vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} \right) \right)^2 \right. \right. \\ &\quad \left. \left. + \lambda^2 (\mathfrak{d}^{k+1} \check{\kappa})^2 + \lambda^6 \left(\mathfrak{d}^k \left(e_\theta(\mu) + \vartheta \mathfrak{d}_2 \mathfrak{d}_2^* (\mathfrak{d}_1^* \mathfrak{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) \right)^2 \right. \right. \\ &\quad \left. \left. + \lambda^4 (\mathfrak{d}^{\leq k} \check{\mu})^2 + \lambda^2 \left(\mathfrak{d}^k \left(e_\theta(\underline{\kappa}) - 4\underline{\beta} \right) \right)^2 + \lambda^2 \left(\mathfrak{d}^k \left(e_3(\zeta) + \underline{\beta} \right) \right)^2 \right] \right). \end{aligned}$$

Proposition 8.8.3. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Then, we have*

$${}^{(ext)}\mathfrak{G}_{J+1}^{\leq 4m_0}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_{J+1}^{\leq 4m_0'}[\check{\Gamma}] \lesssim {}^{(ext)}\mathfrak{G}_{J+1}^{\geq 4m_0}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_{J+1}^{\geq 4m_0'}[\check{\Gamma}] + {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{G}_J[\check{\Gamma}],$$

where we have introduced the notation

$$\begin{aligned} {}^{(ext)}\mathfrak{G}_k^{\leq 4m_0'}[\check{\Gamma}] &:= \sup_{r\tau \leq \lambda \leq 4m_0} \left(\int_{\{r=\lambda\}} \left[\lambda^6 \left(\mathfrak{d}^k \left(\mathfrak{d}_1 \mathfrak{d}_1^* \kappa - \vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} \right) \right)^2 \right. \right. \\ &\quad \left. \left. + \lambda^2 (\mathfrak{d}^{k+1} \check{\kappa})^2 + \lambda^6 \left(\mathfrak{d}^k \left(e_\theta(\mu) + \vartheta \mathfrak{d}_2 \mathfrak{d}_2^* (\mathfrak{d}_1^* \mathfrak{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) \right)^2 \right. \right. \\ &\quad \left. \left. + \lambda^4 (\mathfrak{d}^{\leq k} \check{\mu})^2 + \lambda^2 \left(\mathfrak{d}^k \left(e_\theta(\underline{\kappa}) - 4\underline{\beta} \right) \right)^2 + \lambda^2 \left(\mathfrak{d}^{k-1} \mathbf{N} \left(e_3(\zeta) + \underline{\beta} \right) \right)^2 \right] \right). \end{aligned}$$

The proof of Proposition 8.8.1 is postponed to section 8.8.1, the proof of Proposition 8.8.2 is postponed to section 8.8.4, and the proof of Proposition 8.8.3 is postponed to section 8.8.5. The proof of the two latter propositions will rely in particular on basic weighted estimates for transport equations along e_4 in $^{(ext)}\mathcal{M}$ derived in section 8.8.2, as well as several renormalized identities derived in section 8.8.3

We now conclude the proof of Proposition 8.3.9. In view of Propositions 8.8.1, 8.8.2 and 8.8.3, we have, for J such that $k_{small} - 2 \leq J \leq k_{large} - 1$,

$$^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] \lesssim ^{(ext)}\mathfrak{R}_{J+1}[\check{R}] + ^{(ext)}\mathfrak{G}_J[\check{\Gamma}],$$

where we have used the fact that

$$^{(\Sigma_*)}\mathfrak{R}_{J+1}[\check{R}] \leq ^{(ext)}\mathfrak{R}_{J+1}[\check{R}], \quad ^{(\Sigma_*)}\mathfrak{G}_J[\check{\Gamma}] \leq ^{(ext)}\mathfrak{G}_J[\check{\Gamma}].$$

In view of the iteration assumption (8.3.10), we infer

$$^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] \lesssim ^{(ext)}\mathfrak{R}_{J+1}[\check{R}] + \epsilon_B[J].$$

Since the estimates in Proposition 8.8.2 are integrated from Σ_* , we obtain similarly, for any $r_0 \geq 4m_0$,

$$^{(ext)}\mathfrak{G}_{J+1}^{\geq r_0}[\check{\Gamma}] \lesssim ^{(ext)}\mathfrak{R}_{J+1}^{\geq r_0}[\check{R}] + \epsilon_B[J].$$

On the other hand, we have in view of Proposition 8.3.8, for any $r_0 \geq 4m_0$,

$$^{(ext)}\mathfrak{R}_{J+1}^{\geq r_0}[\check{R}] \lesssim r_0^{-\delta_B} ^{(ext)}\mathfrak{G}_k^{\geq r_0}[\check{\Gamma}] + r_0^{10} \left(\epsilon_B[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right).$$

and

$$^{(int)}\mathfrak{R}_{J+1}[\check{R}] + ^{(ext)}\mathfrak{R}_{J+1}[\check{R}] \leq ^{(ext)}\mathfrak{R}_{J+1}^{\geq r_0}[\check{R}] + O \left(r_0^{10} \left(\epsilon_B[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) \right) \right).$$

Choosing $r_0 \geq 4m_0$ large enough, we infer from the above estimates

$$^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] + ^{(int)}\mathfrak{R}_{J+1}[\check{R}] + ^{(ext)}\mathfrak{R}_{J+1}[\check{R}] \lesssim \epsilon_B[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right).$$

This concludes the proof of Proposition 8.3.9.

8.8.1 Proof of Proposition 8.8.1

Step 1. We control κ on Σ_* . Recall the GCM conditions $\kappa = 2/r$ on Σ_* . Since ν_{Σ_*} and e_θ are tangent, we infer

$$(\emptyset, \nu_{\Sigma_*})^k \left(\kappa - \frac{2}{r} \right) = 0.$$

Together with Raychadhuri, we infer

$$\begin{aligned} & \max_{k \leq J+2} \int_{\Sigma_*} \left(r^2 \left(\mathfrak{d}^k \left(\kappa - \frac{2}{r} \right) \right)^2 + r^4 \left(\mathfrak{d}^k e_\theta(\kappa) \right)^2 \right) \\ & \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2, \end{aligned}$$

where we have used the fact that e_3 is in the span of e_4 and ν_{Σ_*} . Note that we have used Codazzi for ϑ to control the term $\mathfrak{d}^{J+1} e_4(e_\theta(\kappa))$.

Step 2. We control the $\ell = 1$ modes on Σ_* . In view of the GCM conditions for κ , and projecting the Codazzi for ϑ on the $\ell = 1$ mode, we infer on Σ_*

$$\int_S \zeta e^\Phi = r \int_S \beta e^\Phi + \frac{r}{2} \int_S \vartheta \zeta e^\Phi.$$

Since the vectorfield ν is tangent to Σ_* , we infer

$$\begin{aligned} \nu^{J+2} \left(\int_S \zeta e^\Phi \right) &= r \int_S \nu^{J+2} \beta e^\Phi + \frac{r}{2} \int_S \nu^{J+2} (\vartheta \zeta) e^\Phi + \text{l.o.t.} \\ &= r \int_S \nu^{J+2} \beta e^\Phi + \frac{r}{2} \int_S \zeta \nu^{J+2} (\vartheta) e^\Phi + \frac{r}{2} \int_S \vartheta \nu^{J+2} (\zeta) e^\Phi + \text{l.o.t.} \end{aligned}$$

where l.o.t. denote, here and below, terms that

- either are linear and contain at most $J + 1$ derivatives of curvature components and J derivatives of Ricci coefficients,
- or are quadratic and contain at most $J + 1$ derivatives of Ricci coefficients and curvature components.

Using Bianchi identities and the null structure equations, we deduce

$$\begin{aligned} & \nu^{J+2} \left(\int_S \zeta e^\Phi \right) \\ &= r \int_S \nu^{J+1} (\mathfrak{d}_2 \alpha, \mathfrak{d}_1^* \rho - 3\rho\eta) e^\Phi + \frac{r}{2} \int_S \zeta \nu^{J+1} \mathfrak{d}_2^* \eta e^\Phi + \frac{r}{2} \int_S \vartheta \nu^{J+1} \mathfrak{d}_1^* \underline{\omega} e^\Phi + \text{l.o.t.} \\ &= r \int_S (\mathfrak{d}_2 \nu^{J+1} \alpha, \nu^J \mathfrak{d}_1^* \mathfrak{d}_1(\beta, \underline{\beta})) e^\Phi - 3\bar{\rho} \int_S \nu^{J+1} \eta e^\Phi + \frac{r}{2} \int_S \zeta \mathfrak{d}_2^* \nu^{J+1} \eta e^\Phi \\ & \quad + \frac{r}{2} \int_S \vartheta \mathfrak{d}_1^* \nu^{J+1} \underline{\omega} e^\Phi + \text{l.o.t.} \\ &= r \int_S (\mathfrak{d}_2 \nu^{J+1} \alpha, \mathfrak{d}_1^* \mathfrak{d}_1 \nu^J(\beta, \underline{\beta})) e^\Phi + \frac{r}{2} \int_S \zeta \mathfrak{d}_2^* \nu^{J+1} \eta e^\Phi + \frac{r}{2} \int_S \vartheta \mathfrak{d}_1^* \nu^{J+1} \underline{\omega} e^\Phi + \text{l.o.t.}, \end{aligned}$$

where we have used, in the last equality, a cancellation due to the fact that ν is tangent to Σ_* and $\int_S \eta e^\Phi = 0$ on Σ_* . Using the identity $\mathfrak{d}_1^* \mathfrak{d}_1 = \mathfrak{d}_2 \mathfrak{d}_2^* + 2K$, integration by parts for all terms, and the fact that $\mathfrak{d}_2^*(e^\Phi) = 0$ so that the top order linear term vanish, we infer

$$\nu^{J+2} \left(\int_S \zeta e^\Phi \right) = \text{l.o.t.}$$

with the above convention for the lower order terms. Also, relying on the null equation for $e_4(\zeta)$, i.e.

$$e_4(\zeta) = -\kappa\zeta - \beta - \vartheta\zeta$$

we obtain, with more ease since this estimate is at one lower level of derivatives

$$(re_4, \nu)^{J+2} \left(\int_S \zeta e^\Phi \right) = \text{l.o.t.}$$

We infer

$$\max_{k \leq J+2} \int_1^{u_*} r^{-2} \left(\mathfrak{d}^k \left(\int_S \zeta e^\Phi \right) \right)^2 \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2.$$

Next, we have in view of the definition of μ and the identity $\mathfrak{d}_1^* \mathfrak{d}_1 = \mathfrak{d}_2 \mathfrak{d}_2^* + 2K$

$$\begin{aligned} \int_S e_\theta(\mu) e^\Phi &= \int_S \mathfrak{d}_1^* \mathfrak{d}_1 \zeta e^\Phi - \int_S e_\theta(\rho) e^\Phi + \frac{1}{4} \int_S e_\theta(\vartheta \underline{\vartheta}) e^\Phi \\ &= 2 \int_S K \zeta e^\Phi - \int_S e_\theta(\rho) e^\Phi + \frac{1}{4} \int_S e_\theta(\vartheta \underline{\vartheta}) e^\Phi \\ &= \frac{2}{r^2} \int_S \zeta e^\Phi - \int_S e_\theta(\rho) e^\Phi + \int_S \left(K - \frac{2}{r^2} \right) \zeta e^\Phi + \frac{1}{4} \int_S e_\theta(\vartheta \underline{\vartheta}) e^\Phi. \end{aligned}$$

To estimate the RHS, we use in particular

- for the second term, in view of Bianchi

$$\begin{aligned}
& (e_3, re_4)^{J+2} e_\theta(\rho) \\
= & (e_3, re_4)^{J+1} \mathcal{D}_1^* \mathcal{D}_1(r\beta, \underline{\beta}) - \frac{3}{2} (e_3, re_4)^{J+1} \mathcal{D}_1^*(r\kappa\rho, \underline{\kappa}\rho) \\
& + (e_3, re_4)^{J+1} \mathcal{D}_1^*(r\underline{\vartheta}\alpha, (r\underline{\zeta}, \underline{\xi})\beta, (\zeta, \eta)\underline{\beta}, \vartheta\underline{\alpha}) + \text{l.o.t.} \\
= & (e_3, re_4)^{J+1} \mathcal{D}_2 \mathcal{D}_2^*(r\beta, \underline{\beta}) + \frac{3}{2} \rho (e_3, re_4)^{J+1} (re_\theta(\kappa), e_\theta(\underline{\kappa})) \\
& + (r\underline{\vartheta}(e_3, re_4)^{J+1} \mathcal{D}_1^* \alpha, (r\underline{\zeta}, \underline{\xi})(e_3, re_4)^{J+1} \mathcal{D}_1^* \beta, (\zeta, \eta)(e_3, re_4)^{J+1} \mathcal{D}_1^* \underline{\beta}, \vartheta(e_3, re_4)^{J+1} \mathcal{D}_1^* \underline{\alpha}) \\
& + (r\alpha(e_3, re_4)^{J+1} \mathcal{D}_1^* \underline{\vartheta}, \beta(e_3, re_4)^{J+1} \mathcal{D}_1^*(r\underline{\zeta}, \underline{\xi}), \underline{\beta}(e_3, re_4)^{J+1} \mathcal{D}_1^*(\zeta, \eta), \underline{\alpha}(e_3, re_4)^{J+1} \mathcal{D}_1^* \underline{\vartheta}) + \text{l.o.t.} \\
= & \mathcal{D}_2(e_3, re_4)^{J+1} \mathcal{D}_2^*(r\beta, \underline{\beta}) + [(e_3, re_4)^{J+1}, \mathcal{D}_2] \mathcal{D}_2^*(r\beta, \underline{\beta}) + \frac{3}{2} \bar{\rho} (e_3, re_4)^{J+1} (re_\theta(\kappa), e_\theta(\underline{\kappa})) \\
& - \frac{3}{2} \check{\rho} \mathcal{D}_1^*(e_3, re_4)^{J+1} (r\check{\kappa}, \check{\underline{\kappa}}) \\
& + (r\underline{\vartheta} \mathcal{D}_1^*(e_3, re_4)^{J+1} \alpha, (r\underline{\zeta}, \underline{\xi}) \mathcal{D}_1^*(e_3, re_4)^{J+1} \beta, (\zeta, \eta) \mathcal{D}_1^*(e_3, re_4)^{J+1} \underline{\beta}, \vartheta \mathcal{D}_1^*(e_3, re_4)^{J+1} \underline{\alpha}) \\
& + (r\alpha \mathcal{D}_1^*(e_3, re_4)^{J+1} \underline{\vartheta}, \beta \mathcal{D}_1^*(e_3, re_4)^{J+1} (r\underline{\zeta}, \underline{\xi}), \underline{\beta} \mathcal{D}_1^*(e_3, re_4)^{J+1} (\zeta, \eta), \underline{\alpha} \mathcal{D}_1^*(e_3, re_4)^{J+1} \underline{\vartheta}) + \text{l.o.t.}
\end{aligned}$$

- for the third term

$$\begin{aligned}
& (e_3, re_4)^{J+2} \left(\left(K - \frac{2}{r^2} \right) \zeta \right) \\
= & \zeta (e_3, re_4)^{J+2} \left(K - \frac{2}{r^2} \right) + \left(K - \frac{2}{r^2} \right) (e_3, re_4)^{J+2} \zeta + \text{l.o.t.} \\
= & \zeta (e_3, re_4)^{J+1} \left(\mathcal{D}_1(r\beta, \underline{\beta}, \eta, r^{-1}\underline{\xi}) \right) + \left(K - \frac{2}{r^2} \right) (e_3, re_4)^{J+1} e_\theta(\underline{\omega}) + \text{l.o.t.} \\
= & \zeta (e_3, re_4)^{J+1} \mathcal{D}_1 \left(r\beta, \underline{\beta}, \eta, r^{-1}\underline{\xi} \right) + \left(K - \frac{2}{r^2} \right) (e_3, re_4)^{J+1} e_\theta(\underline{\omega}) + \text{l.o.t.} \\
= & \zeta \mathcal{D}_1(e_3, re_4)^{J+1} \left(r\beta, \underline{\beta}, \eta, r^{-1}\underline{\xi} \right) + \left(K - \frac{2}{r^2} \right) \mathcal{D}_1^*(e_3, re_4)^{J+1} \underline{\omega} + \text{l.o.t.},
\end{aligned}$$

- for the fourth term

$$\begin{aligned}
(e_3, re_4)^{J+2} e_\theta(\vartheta\underline{\vartheta}) &= (e_3, re_4)^{J+1} e_\theta(\vartheta \mathcal{D}_2^*(\underline{\xi}, r\underline{\zeta})) + (e_3, re_4)^{J+1} e_\theta(\vartheta \mathcal{D}_2^* \eta) + \text{l.o.t.} \\
&= \vartheta \mathcal{D}_1^* \mathcal{D}_2^*(e_3, re_4)^{J+1} (\underline{\xi}, r\underline{\zeta}) + \vartheta \mathcal{D}_1^* \mathcal{D}_2^*(e_3, re_4)^{J+1} \eta + \text{l.o.t.}
\end{aligned}$$

We infer

$$\begin{aligned}
& (e_3, re_4)^{J+2} \left(\int_S e_\theta(\mu) e^\Phi \right) \\
= & \frac{2}{r^2} (e_3, re_4)^{J+2} \left(\int_S \zeta e^\Phi \right) + \int_S \mathfrak{d}_2(e_3, re_4)^{J+1} \mathfrak{d}_2^*(r\beta, \underline{\beta}) e^\Phi \\
& + \int_S [(e_3, re_4)^{J+1}, \mathfrak{d}_2] \mathfrak{d}_2^*(r\beta, \underline{\beta}) e^\Phi + \frac{3}{2} \bar{\rho} \int_S (e_3, re_4)^{J+1} (re_\theta(\kappa), e_\theta(\underline{\kappa})) e^\Phi \\
& - \frac{3}{2} \int_S \check{\rho} \mathfrak{d}_1^*(e_3, re_4)^{J+1} (r\check{\kappa}, \check{\underline{\kappa}}) e^\Phi \\
& + \int_S (r\vartheta \mathfrak{d}_1^*(e_3, re_4)^{J+1} \alpha, (r\zeta, \underline{\xi}) \mathfrak{d}_1^*(e_3, re_4)^{J+1} \beta, (\zeta, \eta) \mathfrak{d}_1^*(e_3, re_4)^{J+1} \underline{\beta}, \vartheta \mathfrak{d}_1^*(e_3, re_4)^{J+1} \underline{\alpha}) e^\Phi \\
& + \int_S (r\alpha \mathfrak{d}_1^*(e_3, re_4)^{J+1} \underline{\vartheta}, \beta \mathfrak{d}_1^*(e_3, re_4)^{J+1} (r\zeta, \underline{\xi}), \underline{\beta} \mathfrak{d}_1^*(e_3, re_4)^{J+1} (\zeta, \eta), \underline{\alpha} \mathfrak{d}_1^*(e_3, re_4)^{J+1} \vartheta) e^\Phi \\
& + \int_S \zeta \mathfrak{d}_1^*(e_3, re_4)^{J+1} (r\beta, \underline{\beta}, \zeta, r^{-1}\underline{\xi}) e^\Phi + \int_S \left(K - \frac{2}{r^2} \right) \mathfrak{d}_1^*(e_3, re_4)^{J+1} \check{\omega} e^\Phi \\
& + \int_S \vartheta \mathfrak{d}_1^* \mathfrak{d}_2^*(e_3, re_4)^{J+1} (\underline{\xi}, r\zeta) e^\Phi + \int_S \underline{\vartheta} \mathfrak{d}_1^* \mathfrak{d}_2^*(e_3, re_4)^{J+1} \zeta e^\Phi + \text{l.o.t.}
\end{aligned}$$

and after integrations by parts and the fact that

$$\mathfrak{d}_k(Fe^\Phi) = \mathfrak{d}_{k+1}(F)e^\Phi, \quad \mathfrak{d}_k^*(Fe^\Phi) = \mathfrak{d}_{k-1}^*(F)e^\Phi,$$

we obtain

$$\begin{aligned}
& (e_3, re_4)^{J+2} \left(\int_S e_\theta(\mu) e^\Phi \right) \\
= & \frac{2}{r^2} (e_3, re_4)^{J+2} \left(\int_S \zeta e^\Phi \right) + \int_S \mathfrak{d}_1^* \left([(e_3, re_4)^{J+1}, \mathfrak{d}_2] \right) (r\beta, \underline{\beta}) e^\Phi \\
& + \frac{3}{2} \bar{\rho} \int_S (e_3, re_4)^{J+1} (re_\theta(\kappa), e_\theta(\underline{\kappa})) e^\Phi + \frac{3}{2} \int_S \mathfrak{d}_2 \rho(e_3, re_4)^{J+1} (r\check{\kappa}, \check{\underline{\kappa}}) e^\Phi \\
& + \int_S (r \mathfrak{d}_2 \underline{\vartheta}(e_3, re_4)^{J+1} \alpha, \mathfrak{d}_2(r\zeta, \underline{\xi})(e_3, re_4)^{J+1} \beta, \mathfrak{d}_2(\zeta, \eta)(e_3, re_4)^{J+1} \underline{\beta}, \mathfrak{d}_2 \vartheta(e_3, re_4)^{J+1} \underline{\alpha}) e^\Phi \\
& + \int_S (r \mathfrak{d}_2 \alpha(e_3, re_4)^{J+1} \underline{\vartheta}, \mathfrak{d}_2 \beta(e_3, re_4)^{J+1} (r\zeta, \underline{\xi}), \mathfrak{d}_2 \underline{\beta}(e_3, re_4)^{J+1} (\zeta, \eta), \mathfrak{d}_2 \underline{\alpha}(e_3, re_4)^{J+1} \vartheta) e^\Phi \\
& + \int_S \mathfrak{d}_1 \zeta(e_3, re_4)^{J+1} (r\beta, \underline{\beta}, \zeta, r^{-1}\underline{\xi}) e^\Phi + \int_S \mathfrak{d}_2 \left(K - \frac{2}{r^2} \right) (e_3, re_4)^{J+1} \check{\omega} e^\Phi \\
& + \int_S \mathfrak{d}_3 \mathfrak{d}_2 \vartheta(e_3, re_4)^{J+1} (\underline{\xi}, r\zeta) e^\Phi + \int_S \mathfrak{d}_3 \mathfrak{d}_2 \underline{\vartheta}(e_3, re_4)^{J+1} \zeta e^\Phi + \text{l.o.t.}
\end{aligned}$$

Together with the above estimate for the $\ell = 1$ mode of ζ and the estimate of Step 1 for κ , we infer

$$\begin{aligned} \max_{k \leq J+2} \int_1^{u_*} r^2 \left(\mathfrak{d}^k \left(\int_S e_\theta(\mu) e^\Phi \right) \right)^2 &\lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2 \\ &+ \max_{k \leq J+1} \int_1^{u_*} r^{-4} \left(\mathfrak{d}^k \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) \right)^2. \end{aligned}$$

In view of the dominant condition (3.3.4) for r on Σ_* , we infer

$$\begin{aligned} \max_{k \leq J+2} \int_1^{u_*} r^2 \left(\mathfrak{d}^k \left(\int_S e_\theta(\mu) e^\Phi \right) \right)^2 &\lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2 \\ &+ \epsilon_0 \max_{k \leq J+1} \int_1^{u_*} \left(\mathfrak{d}^k \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) \right)^2. \end{aligned}$$

Next, in view of the remarkable identity for the $\ell = 1$ mode of $e_\theta(K)$, we have

$$-\int_S e_\theta(\rho) e^\Phi - \frac{1}{4} \int_S e_\theta(\kappa \underline{\kappa}) e^\Phi + \frac{1}{4} \int_S e_\theta(\vartheta \underline{\vartheta}) e^\Phi = 0$$

and hence

$$\int_S e_\theta(\underline{\kappa}) e^\Phi = -2r \int_S e_\theta(\rho) e^\Phi - \frac{r}{2} \int_S \left(\kappa - \frac{2}{r} \right) e_\theta(\underline{\kappa}) e^\Phi - \frac{r}{2} \int_S \underline{\kappa} e_\theta(\kappa) e^\Phi + \frac{r}{2} \int_S e_\theta(\vartheta \underline{\vartheta}) e^\Phi.$$

Arguing as for the estimate of the $\ell = 1$ mode of $e_\theta(\mu)$, and using the smallness of ϵ_0 , we infer

$$\begin{aligned} &\max_{k \leq J+2} \int_1^{u_*} r^2 \left(\mathfrak{d}^k \left(\int_S e_\theta(\mu) e^\Phi \right) \right)^2 + \max_{k \leq J+2} \int_1^{u_*} \left(\mathfrak{d}^k \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) \right)^2 \\ &\lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2. \end{aligned}$$

We have thus obtained

$$\begin{aligned} &\max_{k \leq J+2} \int_1^{u_*} \left(r^{-2} \left(\mathfrak{d}^k \left(\int_S \zeta e^\Phi \right) \right)^2 + r^2 \left(\mathfrak{d}^k \left(\int_S e_\theta(\mu) e^\Phi \right) \right)^2 + \left(\mathfrak{d}^k \left(\int_S e_\theta(\underline{\kappa}) e^\Phi \right) \right)^2 \right) \\ &\lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2. \end{aligned}$$

Step 3. Recall the GCM conditions $\phi_2^* \phi_1^* \underline{\kappa} = \phi_2^* \phi_1^* \mu = 0$ on Σ_* . This yields on Σ_*

$$e_\theta(\mu) = \frac{\int_S e_\theta(\mu) e^\Phi}{\int_S e^{2\Phi}} e^\Phi, \quad e_\theta(\underline{\kappa}) = \frac{\int_S e_\theta(\underline{\kappa}) e^\Phi}{\int_S e^{2\Phi}} e^\Phi.$$

Together with Step 2, we infer

$$\begin{aligned} & \max_{k \leq J+2} \int_{\Sigma_*} \left(r^4 ((\vartheta, \nu_{\Sigma_*})^k \check{\mu})^2 + r^2 ((\vartheta, \nu_{\Sigma_*})^k \check{\kappa})^2 \right) \\ & \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2. \end{aligned}$$

Then, in view of the null structure equations for $e_4(\check{\mu})$ and $e_4(\check{\kappa})$,

$$\begin{aligned} e_4(\check{\mu}) &= -\frac{3}{2} \bar{\kappa} \check{\mu} - \frac{3}{2} \bar{\mu} \check{\kappa} + \text{Err}[e_4 \check{\mu}] \\ e_4(\check{\kappa}) &= -\frac{1}{2} \bar{\kappa} \check{\kappa} - \frac{1}{2} \bar{\kappa} \check{\kappa} + 2\check{\mu} + 4\check{\rho} + \text{Err}[e_4 \check{\kappa}], \end{aligned}$$

we infer, together with the control of $\check{\kappa}$ provided by Step 1,

$$\begin{aligned} & \max_{k \leq J+2} \int_{\Sigma_*} \left(r^4 (\mathfrak{d}^k \check{\mu})^2 + r^2 (\mathfrak{d}^k \check{\kappa})^2 \right) \\ & \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2. \end{aligned}$$

Step 4. Recall that we have

$$\mathfrak{d}_1 \zeta = -\check{\mu} - \check{\rho} + \frac{1}{4} \vartheta \underline{\vartheta}.$$

Differentiating, and using the Bianchi identities for $e_4(\check{\rho})$ and $e_3(\check{\rho})$, and the null structure equations for $e_4(\vartheta)$, $e_3(\vartheta)$, $e_4(\underline{\vartheta})$ and $e_3(\underline{\vartheta})$, we infer

$$\begin{aligned} \mathfrak{d}_1 \mathfrak{d}^k \zeta &= -\mathfrak{d}^k \check{\mu} - \mathfrak{d}^{k-1} \vartheta \left(\check{\rho}, \beta, r^{-1} \underline{\beta} \right) + \frac{1}{4} \mathfrak{d}^{k-1} \left(\vartheta(\vartheta \underline{\vartheta}), r^{-1} \underline{\vartheta} \vartheta \eta, \vartheta \vartheta \zeta, r^{-1} \vartheta \vartheta \xi \right) + \text{l.o.t.} \\ &= -\mathfrak{d}^k \check{\mu} - \vartheta \mathfrak{d}^{k-1} \left(\check{\rho}, \beta, r^{-1} \underline{\beta} \right) + \frac{1}{4} \vartheta \left(\underline{\vartheta} \mathfrak{d}^{k-1}(\vartheta, r^{-1} \eta) \right) + \frac{1}{4} \vartheta \left(\vartheta \mathfrak{d}^{k-1}(\underline{\vartheta}, \zeta, r^{-1} \underline{\xi}) \right) + \text{l.o.t.} \end{aligned}$$

We infer, since \mathfrak{d}_1 is invertible in view of the corresponding Poincaré inequality,

$$\mathfrak{d}^k \zeta = -r \vartheta^{-1} \mathfrak{d}^k \check{\mu} - \mathfrak{d}^{k-1} \left(\check{\rho}, \beta, r^{-1} \underline{\beta} \right) + \frac{1}{4} \left(\underline{\vartheta} \mathfrak{d}^{k-1}(\vartheta, r^{-1} \eta) \right) + \frac{1}{4} \left(\vartheta \mathfrak{d}^{k-1}(\underline{\vartheta}, \zeta, r^{-1} \underline{\xi}) \right) + \text{l.o.t.}$$

Together with the estimate for $\check{\mu}$ of Step 3, this yields

$$\max_{k \leq J+2} \int_{\Sigma_*} r^2 (\mathfrak{d}^k \zeta)^2 \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2.$$

Step 5. Recall from the GCM condition that we have on Σ_*

$$\int_S \eta e^\Phi = 0.$$

Together with the transport equation

$$e_4(\eta - \zeta) = -\frac{1}{2}\kappa(\eta - \zeta) - \frac{1}{2}\vartheta(\eta - \zeta),$$

we infer in view of the the estimates for ζ of Step 4,

$$\max_{k \leq J+1} \int_1^{u_*} r^{-4} \left(\mathfrak{d}^k \left(\int_S \eta e^\Phi \right) \right)^2 \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2.$$

Next, recall from Proposition 2.2.19 that η verifies

$$\begin{aligned} 2 \mathfrak{d}_2 \mathfrak{d}_2^* \eta &= \kappa (-e_3(\zeta) + \underline{\beta}) - e_3(e_\theta(\kappa)) - \kappa \left(\frac{1}{2} \underline{\kappa} \zeta - 2 \underline{\omega} \zeta \right) + 6 \rho \eta - \underline{\kappa} e_\theta \kappa \\ &\quad - \frac{1}{2} \kappa e_\theta(\underline{\kappa}) + 2 \underline{\omega} e_\theta(\kappa) + 2 e_\theta(\rho) + \text{Err}[\mathfrak{d}_2 \mathfrak{d}_2^* \eta], \\ \text{Err}[\mathfrak{d}_2 \mathfrak{d}_2^* \eta] &= \left(2 \mathfrak{d}_1 \eta - \frac{1}{2} \kappa \underline{\vartheta} + 2 \eta^2 \right) \eta + 2 e_\theta(\eta^2) - \kappa \left(\frac{1}{2} \underline{\vartheta} \zeta - \frac{1}{2} \underline{\vartheta} \underline{\xi} \right) - \frac{1}{2} \underline{\vartheta} e_\theta(\kappa) \\ &\quad - \left(2 \mathfrak{d}_1 \eta - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} + 2 \eta^2 \right) \zeta - \frac{1}{2} e_\theta(\underline{\vartheta} \vartheta) - \frac{1}{2} \underline{\vartheta}^2 \underline{\xi} - \frac{3}{2} \underline{\vartheta} \underline{\vartheta} \eta. \end{aligned}$$

Together with the estimates for κ of Step 1, the estimates for $\underline{\kappa}$ of Step 3, and the estimates for ζ of Step 4,

$$\begin{aligned} &\max_{k \leq J+1} \int_{\Sigma_*} \left(\mathfrak{d}^k \left(r^2 \mathfrak{d}_2 \mathfrak{d}_2^* \eta - r^2 e_\theta(\rho) - \frac{r^2}{2} \mathfrak{d}_2(\eta^2) - r^2 e_\theta(\eta^2) + r^2 \mathfrak{d}_1(\zeta \eta) + \frac{1}{4} r^2 e_\theta(\underline{\vartheta} \vartheta) \right) \right)^2 \\ &\lesssim \max_{k \leq J+1} \int_{\Sigma_*} r^{-2} |\mathfrak{d}^k \eta|^2 + \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2. \end{aligned}$$

In view of the dominant condition (3.3.4) for r on Σ_* , we infer

$$\begin{aligned} &\max_{k \leq J+1} \int_{\Sigma_*} \left(\mathfrak{d}^k \left(r^2 \mathfrak{d}_2 \mathfrak{d}_2^* \eta - r^2 e_\theta(\rho) - \frac{r^2}{2} \mathfrak{d}_2(\eta^2) - r^2 e_\theta(\eta^2) + r^2 \mathfrak{d}_1(\zeta \eta) + \frac{1}{4} r^2 e_\theta(\underline{\vartheta} \vartheta) \right) \right)^2 \\ &\lesssim \epsilon_0^{\frac{2}{3}} \max_{k \leq J+1} \int_{\Sigma_*} |\mathfrak{d}^k \eta|^2 + \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2. \end{aligned}$$

This yields

$$\begin{aligned} &\max_{k \leq J+1} \int_{\Sigma_*} \left(r^2 \mathfrak{d}_2 \mathfrak{d}_2^* \mathfrak{d}^k \eta + r \mathfrak{d}_2^* [\mathfrak{d}^k, r \mathfrak{d}_2] \eta + r \mathfrak{d}_2 [\mathfrak{d}^k, r \mathfrak{d}_2^*] \eta \right. \\ &\quad \left. - r^2 e_\theta(\mathfrak{d}^k \rho) - \frac{r^2}{2} \mathfrak{d}_2 \mathfrak{d}^k(\eta^2) - r^2 e_\theta \mathfrak{d}^k(\eta^2) + r^2 \mathfrak{d}_1 \mathfrak{d}^k(\zeta \eta) + \frac{1}{4} r^2 e_\theta \mathfrak{d}^k(\underline{\vartheta} \vartheta) \right)^2 \\ &\lesssim \epsilon_0^{\frac{2}{3}} \max_{k \leq J+1} \int_{\Sigma_*} |\mathfrak{d}^k \eta|^2 + \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2. \end{aligned}$$

We deduce, using a Poincaré inequality for $\not\phi_2$,

$$\max_{k \leq J+1} \int_{\Sigma_*} (r \not\phi_2^* \not\partial^k \eta)^2 \lesssim \epsilon_0^{\frac{2}{3}} \max_{k \leq J+1} \int_{\Sigma_*} |\not\partial^k \eta|^2 + \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2.$$

Together with a Poincaré inequality for $r \not\phi_2^*$ and the above control of the $\ell = 1$ mode of η , we infer

$$\max_{k \leq J+1} \int_{\Sigma_*} (\not\partial^k \eta)^2 \lesssim \epsilon_0^{\frac{2}{3}} \max_{k \leq J+1} \int_{\Sigma_*} |\not\partial^k \eta|^2 + \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2,$$

and hence, for ϵ_0 small enough,

$$\max_{k \leq J+1} \int_{\Sigma_*} (\not\partial^k \eta)^2 \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2.$$

Step 6. Recall from the GCM condition that we have on Σ_*

$$\int_S \underline{\xi} e^\Phi = 0.$$

Together with the transport equation

$$e_4(\underline{\xi}) = -e_3(\underline{\zeta}) + \underline{\beta} - \underline{\kappa} \underline{\zeta} - \underline{\zeta} \underline{\vartheta},$$

we infer in view of the the estimates for $\underline{\zeta}$ of Step 4, the estimates for $\underline{\beta}$, and the bootstrap assumptions

$$\max_{k \leq J+1} \int_1^{u_*} r^{-4} \left(\not\partial^k \left(\int_S \underline{\xi} e^\Phi \right) \right)^2 \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2.$$

Next, from Proposition 2.2.19 that we have

$$\begin{aligned} 2 \not\phi_2 \not\phi_2^* \underline{\xi} &= -e_3(e_\theta(\underline{\kappa})) + \underline{\kappa} (e_3(\underline{\zeta}) - \underline{\beta}) + \underline{\kappa}^2 \underline{\zeta} - \frac{3}{2} \underline{\kappa} e_\theta \underline{\kappa} + 6\rho \underline{\xi} - 2\underline{\omega} e_\theta(\underline{\kappa}) + \text{Err}[\not\phi_2 \not\phi_2^* \underline{\xi}], \\ \text{Err}[\not\phi_2 \not\phi_2^* \underline{\xi}] &= \left(2 \not\phi_1 \underline{\xi} + \frac{1}{2} \underline{\kappa} \underline{\vartheta} + 2\eta \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \eta + 2e_\theta(\eta \underline{\xi}) - \frac{1}{2} e_\theta(\underline{\vartheta}^2) \\ &+ \underline{\kappa} \left(\frac{1}{2} \underline{\vartheta} \underline{\zeta} - \frac{1}{2} \underline{\vartheta} \underline{\xi} \right) - \frac{1}{2} \underline{\vartheta} e_\theta \underline{\kappa} - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} \underline{\xi} - \underline{\zeta} \left(2 \not\phi_1 \underline{\xi} + 2(\eta - 3\underline{\zeta}) \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \\ &+ \underline{\xi} \left(-\underline{\vartheta} \underline{\vartheta} - 2 \not\phi_1 \underline{\zeta} + 2\underline{\zeta}^2 \right) - 6\eta \underline{\zeta} \underline{\xi} - 6e_\theta(\underline{\zeta} \underline{\xi}). \end{aligned}$$

Together with the estimates for κ of Step 1, the estimates for $\underline{\kappa}$ of Step 3, and the estimates for ζ of Step 4,

$$\begin{aligned} & \max_{k \leq J+1} \int_{\Sigma_*} \left(\mathfrak{d}^k \left(r^2 \mathfrak{d}_2 \mathfrak{d}_2^* \underline{\xi} + \frac{1}{2} e_\theta(e_3(\underline{\kappa})) - \eta \mathfrak{d}_1 \underline{\xi} - e_\theta(\eta \underline{\xi}) + \frac{1}{4} e_\theta(\underline{\vartheta}^2) + \mathfrak{d}_2(\zeta \underline{\xi}) + 3e_\theta(\zeta \underline{\xi}) \right) \right)^2 \\ & \lesssim \max_{k \leq J+1} \int_{\Sigma_*} r^{-2} |\mathfrak{d}^k \underline{\xi}|^2 + \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2. \end{aligned}$$

In view of the dominant condition (3.3.4) for r on Σ_* , we infer

$$\begin{aligned} & \max_{k \leq J+1} \int_{\Sigma_*} \left(\mathfrak{d}^k \left(r^2 \mathfrak{d}_2 \mathfrak{d}_2^* \underline{\xi} + \frac{1}{2} e_\theta(e_3(\underline{\kappa})) - \eta \mathfrak{d}_1 \underline{\xi} - e_\theta(\eta \underline{\xi}) + \frac{1}{4} e_\theta(\underline{\vartheta}^2) + \mathfrak{d}_2(\zeta \underline{\xi}) + 3e_\theta(\zeta \underline{\xi}) \right) \right)^2 \\ & \lesssim \epsilon_0^{\frac{2}{3}} \max_{k \leq J+1} \int_{\Sigma_*} |\mathfrak{d}^k \underline{\xi}|^2 + \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2. \end{aligned}$$

This yields

$$\begin{aligned} & \max_{k \leq J+1} \int_{\Sigma_*} \left(r^2 \mathfrak{d}_2 \mathfrak{d}_2^* \mathfrak{d}^k \underline{\xi} + r \mathfrak{d}_2^* [\mathfrak{d}^k, r \mathfrak{d}_2] \underline{\xi} + r \mathfrak{d}_2 [\mathfrak{d}^k, r \mathfrak{d}_2^*] \underline{\xi} \right. \\ & \quad \left. + \frac{1}{2} e_\theta(\mathfrak{d}^k e_3(\underline{\kappa})) - \mathfrak{d}_1(\eta \mathfrak{d}^k \underline{\xi}) - e_\theta \mathfrak{d}^k(\eta \underline{\xi}) + \frac{1}{4} e_\theta \mathfrak{d}^k(\underline{\vartheta}^2) + \mathfrak{d}_2 \mathfrak{d}^k(\zeta \underline{\xi}) + 3e_\theta \mathfrak{d}^k(\zeta \underline{\xi}) \right)^2 \\ & \lesssim \epsilon_0^{\frac{2}{3}} \max_{k \leq J+1} \int_{\Sigma_*} |\mathfrak{d}^k \underline{\xi}|^2 + \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2. \end{aligned}$$

We deduce, using a Poincaré inequality for \mathfrak{d}_2 and the estimates for $\underline{\kappa}$ of Step 3,

$$\max_{k \leq J+1} \int_{\Sigma_*} (r \mathfrak{d}_2^* \mathfrak{d}^k \underline{\xi})^2 \lesssim \epsilon_0^{\frac{2}{3}} \max_{k \leq J+1} \int_{\Sigma_*} |\mathfrak{d}^k \underline{\xi}|^2 + \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2.$$

Together with a Poincaré inequality for $r \mathfrak{d}_2^*$ and the above control of the $\ell = 1$ mode of $\underline{\xi}$, we infer

$$\max_{k \leq J+1} \int_{\Sigma_*} (\mathfrak{d}^k \underline{\xi})^2 \lesssim \epsilon_0^{\frac{2}{3}} \max_{k \leq J+1} \int_{\Sigma_*} |\mathfrak{d}^k \underline{\xi}|^2 + \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2,$$

and hence, for ϵ_0 small enough,

$$\max_{k \leq J+1} \int_{\Sigma_*} (\mathfrak{d}^k \underline{\xi})^2 \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0 {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2.$$

Step 7. Using the Codazzi for ϑ and $\underline{\vartheta}$, the transport equation for ϑ and $\underline{\vartheta}$ in the e_4 and e_3 direction, the control of $\check{\kappa}$ of Step 1, the control of $\underline{\kappa}$ of Step 3, the control of ζ of Step

4, the control of η of Step 5, the control of $\underline{\xi}$ of Step 6, and a Poincaré inequality for $\not\partial_2$, we infer

$$\max_{k \leq J+1} \int_{\Sigma_*} \left(r^2 (\not\partial^k \vartheta)^2 + (\not\partial^k \underline{\vartheta})^2 \right) \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0^{\frac{2}{3}} {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2.$$

Step 8. Recall from Proposition 2.2.19 that $\underline{\omega}$ verifies

$$\begin{aligned} 2 \not\partial_1^* \underline{\omega} &= -\frac{1}{2} \kappa \underline{\xi} + \left(\frac{1}{2} \kappa + 2\underline{\omega} + \frac{1}{2} \underline{\vartheta} \right) \eta + e_3(\zeta) - \underline{\beta} \\ &\quad + \frac{1}{2} \kappa \zeta - 2\underline{\omega} \zeta + \frac{1}{2} \underline{\vartheta} \zeta - \frac{1}{2} \underline{\vartheta} \underline{\xi}. \end{aligned}$$

Together with a Poincaré inequality for $\not\partial_1^*$, the control of $\underline{\xi}$ from Step 6, the control of η from Step 5, and the control of ζ from Step 4, we infer

$$\max_{k \leq J+1} \int_{\Sigma_*} |\not\partial^k \underline{\omega}|^2 \lesssim \left({}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0^{\frac{2}{3}} {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] \right)^2.$$

Finally, gathering the estimates of Step 1 to Step 8, we infer

$${}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(\Sigma_*)} \mathfrak{G}'_{J+1}[\check{\Gamma}] \lesssim {}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}] + \epsilon_0^{\frac{2}{3}} {}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}].$$

and hence, for ϵ_0 small enough,

$${}^{(\Sigma_*)} \mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(\Sigma_*)} \mathfrak{G}'_{J+1}[\check{\Gamma}] \lesssim {}^{(\Sigma_*)} \mathfrak{R}_{J+1}[\check{R}] + {}^{(\Sigma_*)} \mathfrak{G}_J[\check{\Gamma}]$$

as desired. This concludes the proof of Proposition 8.8.1.

8.8.2 Weighted estimates for transport equations along e_4 in ${}^{(ext)}\mathcal{M}$

Lemma 8.8.4. *Let the following transport equation in ${}^{(ext)}\mathcal{M}$*

$$e_4(f) + \frac{a}{2} \kappa f = h$$

where $a \in \mathbb{R}$ is a given constant, and f and h are scalar functions. Also, let and $\delta_B > 0$. Then, f satisfies

$$\sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} f^2 \right) \lesssim \int_{\Sigma_*} r^{2a-2} f^2 + \int_{{}^{(ext)}\mathcal{M}(\geq 4m_0)} r^{2a-1+\delta_B} h^2.$$

Proof. Multiply by f to obtain

$$\frac{1}{2}e_4(f^2) + \frac{a}{2}\kappa f^2 = hf.$$

Next, integrate over $S_{u,r}$ to obtain

$$\begin{aligned} \frac{1}{2}e_4 \left(\int_{S_{u,r}} f^2 \right) &= \int_{S_{u,r}} \frac{1}{2}(e_4(f^2) + \kappa f^2) \\ &= - \int_{S_{u,r}} \frac{a-1}{2}\kappa f^2 + \int_{S_{u,r}} hf \\ &= -\frac{a-1}{2}\bar{\kappa} \int_{S_{u,r}} f^2 - \frac{a-1}{2} \int_{S_{u,r}} \check{\kappa} f^2 + \int_{S_{u,r}} hf \end{aligned}$$

and hence

$$\frac{1}{2}e_4 \left(\int_{S_{u,r}} f^2 \right) + \frac{a-1}{2}\bar{\kappa} \int_{S_{u,r}} f^2 = -\frac{a-1}{2} \int_{S_{u,r}} \check{\kappa} f^2 + \int_{S_{u,r}} hf.$$

Also, we multiply by r^{2a-2} which yields

$$\frac{1}{2}e_4 \left(r^{2a-2} \int_{S_{u,r}} f^2 \right) = -\frac{a-1}{2}r^{2a-2} \int_{S_{u,r}} \check{\kappa} f^2 + r^{2a-2} \int_{S_{u,r}} hf$$

where we used the fact that $2e_4(r) = r\bar{\kappa}$. This yields

$$-e_4 \left(r^{2a-2} \int_{S_{u,r}} f^2 \right) \leq r^{2a-3-\delta_B} \int_{S_{u,r}} f^2 + \frac{1}{4}r^{2a-1+\delta_B} \int_{S_{u,r}} h^2$$

and hence

$$-e_4 \left(e^{-\delta_B^{-1}r^{-\delta_B}} r^{2a-2} \int_{S_{u,r}} f^2 \right) \lesssim e^{-\delta_B^{-1}r^{-\delta_B}} r^{2a-1+\delta_B} \int_{S_{u,r}} h^2$$

where we used the fact that $2e_4(r) = r\bar{\kappa} = 2 + O(\epsilon_0)$. Integrating between $r = r_0$ and $r = r_*(u)$, where $r_*(u)$ is such that $S_{u,r_*(u)} \subset \Sigma_*$, we infer

$$r_0^{2a-2} \int_{S_{u,r_0}} f^2 \lesssim r_*(u)^{2a-2} \int_{S_{u,r_*(u)}} f^2 + \int_{r_0}^{r_*(u)} r^{2a-1+\delta_B} \int_{S_{u,r}} h^2. \quad (8.8.1)$$

Remark 8.8.5. Note that we have the following consequences of the coarea formula

$$d\Sigma_* = \varsigma \sqrt{\frac{2}{\varsigma} - \Upsilon + \frac{r}{2}\underline{A}} d\mu_{u,\Sigma_*} du, \quad d\{r = r_0\} = \frac{\varsigma \sqrt{-\bar{\kappa} - \underline{A}}}{\sqrt{\bar{\kappa}}} d\mu_{u,r_0} du,$$

where we used in particular that $\Sigma_* = \{u + r = c_{\Sigma_*}\}$. Also, we have in ${}^{(ext)}\mathcal{M}$

$$d\mathcal{M} = \frac{4\zeta^2}{r^2\kappa^2} d\mu_{u,r} du dr.$$

We infer, in ${}^{(ext)}\mathcal{M}$, using in particular the dominant condition of r on Σ_* ,

$$d\Sigma_* = \left(1 + O\left(\epsilon_0^{\frac{2}{3}}\right)\right) d\mu_{u,\Sigma_*} du, \quad d\{r = r_0\} = \sqrt{1 - \frac{2m_0}{r_0}} (1 + O(\epsilon_0)) d\mu_{u,r_0} du,$$

and

$$d\mathcal{M} = (1 + O(\epsilon_0)) d\mu_{u,r} du dr.$$

Integrating (8.8.1) in $u \in [1, u_*]$, and relying on Remark 8.8.5 we deduce for $r_0 \geq 4m_0$

$$r_0^{2a-2} \int_{\{r=r_0\}} f^2 \lesssim \int_{\Sigma_*} r^{2a-2} f^2 + \int_{{}^{(ext)}\mathcal{M}(r \geq 4m_0)} r^{2a-\frac{1}{2}} h^2$$

as desired. This concludes the proof of the lemma. \square

Corollary 8.8.6. *Let the following transport equation in ${}^{(ext)}\mathcal{M}$*

$$e_4(f) + \frac{a}{2}\kappa f = h$$

where $a \in \mathbb{R}$ is a given constant, and f and h are scalar functions. Also, let and $\delta_B > 0$. Then, f satisfies for $5 \leq k \leq k_{large} + 1$

$$\begin{aligned} & \sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} (\mathfrak{d}^k f)^2 \right) \\ & \lesssim \int_{\Sigma_*} r^{2a-2} (\mathfrak{d}^{\leq k} f)^2 + \sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} (\mathfrak{d}^{\leq k-1} f)^2 \right) + \int_{{}^{(ext)}\mathcal{M}(\geq 4m_0)} r^{2a-1+\delta_B} (\mathfrak{d}^{\leq k} h)^2 \\ & + \left(\sup_{{}^{(ext)}\mathcal{M}(\geq 4m_0)} (r^a |\mathfrak{d}^{\leq k-5} f|) \right)^2 \left({}^{(ext)}\mathfrak{G}_{k-1}^{\geq 4m_0}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_k^{\geq 4m_0}[\check{\kappa}] \right)^2. \end{aligned}$$

Proof. We commute first differentiate the equation for f with $(\mathfrak{d}, \mathbf{T})^l$ and obtain

$$\begin{aligned} e_4((\mathfrak{d}, \mathbf{T})^l f) + \frac{a}{2}\kappa(\mathfrak{d}, \mathbf{T})^l f &= h_l, \\ h_l &:= (\mathfrak{d}, \mathbf{T})^l h - [(\mathfrak{d}, \mathbf{T})^l, e_4]f - \frac{a}{2}[(\mathfrak{d}, \mathbf{T})^l, \kappa]f. \end{aligned}$$

In view of Lemma 8.8.4, we deduce

$$\sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} ((\emptyset, \mathbf{T})^l f)^2 \right) \lesssim \int_{\Sigma_*} r^{2a-2} (\mathfrak{d}^l f)^2 + \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{2a-1+\delta_B} h_l^2.$$

Now, we have the following schematic commutation formulas

$$[\emptyset, e_4] = \Gamma_g \mathfrak{d} + \Gamma_g, \quad [\mathbf{T}, e_4] = r^{-1} \Gamma_b \mathfrak{d},$$

Together with the definition of h_l and for $5 \leq l \leq k_{large} + 1$, we deduce

$$\begin{aligned} \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{2a-1+\delta_B} h_l^2 &\lesssim \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{2a-1+\delta_B} (\mathfrak{d}^l h)^2 + \epsilon_0^2 \int_{(ext)\mathcal{M}} r^{2a-5+\delta_B} (\mathfrak{d}^{\leq l} f)^2 \\ &\quad + \left(\sup_{(ext)\mathcal{M}} (r^a |\mathfrak{d}^{\leq l-5} f|) \right)^2 \left(({}^{(ext)}\mathfrak{G}_{l-1}[\check{\Gamma}] + ({}^{(ext)}\mathfrak{G}_l[\check{\kappa}]) \right)^2 \end{aligned}$$

and hence

$$\begin{aligned} &\sup_{r_0 \geq r_T} \left(r_0^{2a-2} \int_{\{r=r_0\}} ((\emptyset, \mathbf{T})^l f)^2 \right) \\ &\lesssim \int_{\Sigma_*} r^{2a-2} (\mathfrak{d}^l f)^2 + \int_{(ext)\mathcal{M}} r^{2a-1+\delta_B} (\mathfrak{d}^l h)^2 + \epsilon_0^2 \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{2a-5+\delta_B} (\mathfrak{d}^{\leq l} f)^2 \\ &\quad + \left(\sup_{(ext)\mathcal{M}(\geq 4m_0)} (r^a |\mathfrak{d}^{\leq l-5} f|) \right)^2 \left(({}^{(ext)}\mathfrak{G}_{l-1}^{\geq 4m_0}[\check{\Gamma}] + ({}^{(ext)}\mathfrak{G}_l^{\geq 4m_0}[\check{\kappa}]) \right)^2 \end{aligned}$$

or,

$$\begin{aligned} &\sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} ((\emptyset, \mathbf{T})^l f)^2 \right) \\ &\lesssim \int_{\Sigma_*} r^{2a-2} (\mathfrak{d}^l f)^2 + \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{2a-1+\delta_B} (\mathfrak{d}^l h)^2 + \epsilon_0^2 \sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} (\mathfrak{d}^{\leq l} f)^2 \right) \\ &\quad + \left(\sup_{(ext)\mathcal{M}(\geq 4m_0)} (r^a |\mathfrak{d}^{\leq l-5} f|) \right)^2 \left(({}^{(ext)}\mathfrak{G}_{l-1}^{\geq 4m_0}[\check{\Gamma}] + ({}^{(ext)}\mathfrak{G}_l^{\geq 4m_0}[\check{\kappa}]) \right)^2. \end{aligned}$$

Together with the first equation which yields

$$re_4((\emptyset, \mathbf{T})^l f) + \frac{a}{2} r \kappa (\emptyset, \mathbf{T})^l f = r h_l,$$

and hence

$$(re_4)^j ((\emptyset, \mathbf{T})^l f) + \frac{a}{2} (re_4)^{j-1} (r \kappa (\emptyset, \mathbf{T})^l f) = (re_4)^{j-1} (r h_l),$$

we infer, for $\epsilon_0 > 0$ small enough, and for $5 \leq k \leq k_{large} + 1$,

$$\begin{aligned} & \sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} (\mathfrak{d}^k f)^2 \right) \\ \lesssim & \int_{\Sigma_*} r^{2a-2} (\mathfrak{d}^{\leq k} f)^2 + \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{2a-1+\delta_B} (\mathfrak{d}^{\leq k} h)^2 + \sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} (\mathfrak{d}^{\leq k-1} h)^2 \right) \\ & + \sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} (\mathfrak{d}^{\leq k-1} f)^2 \right) \\ & + \left(\sup_{(ext)\mathcal{M}(\geq 4m_0)} (r^a |\mathfrak{d}^{\leq k-5} f|) \right)^2 \left({}^{(ext)}\mathfrak{G}_{k-1}^{\geq 4m_0}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_k^{\geq 4m_0}[\check{\kappa}] \right)^2. \end{aligned}$$

Using a trace estimate, we infer

$$\begin{aligned} & \sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} (\mathfrak{d}^k f)^2 \right) \\ \lesssim & \int_{\Sigma_*} r^{2a-2} (\mathfrak{d}^{\leq k} f)^2 + \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{2a-1+\delta_B} (\mathfrak{d}^{\leq k} h)^2 + \sup_{r_0 \geq 4m_0} \left(r_0^{2a-2} \int_{\{r=r_0\}} (\mathfrak{d}^{\leq k-1} f)^2 \right) \\ & + \left(\sup_{(ext)\mathcal{M}(\geq 4m_0)} (r^a |\mathfrak{d}^{\leq k-5} f|) \right)^2 \left({}^{(ext)}\mathfrak{G}_{k-1}^{\geq 4m_0}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_k^{\geq 4m_0}[\check{\kappa}] \right)^2 \end{aligned}$$

as desired. This concludes the proof of the corollary. \square

Lemma 8.8.7. *Let the following transport equation in $(ext)\mathcal{M}$*

$$e_4(f) + \frac{a}{2}\kappa f = h$$

where $a \in \mathbb{R}$ is a given constant, and f and h are scalar functions. Let $b > 2a - 2$. Then, f satisfies

$$\sup_{r_0 \geq 4m_0} \left(r_0^b \int_{\{r=r_0\}} f^2 \right) + \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{b-1} f^2 \lesssim \int_{\Sigma_*} r^b f^2 + \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{b+1} h^2.$$

Proof. Recall from Lemma 8.8.4 the following identity

$$\frac{1}{2}e_4 \left(\int_{S_{u,r}} f^2 \right) + \frac{a-1}{2}\bar{\kappa} \int_{S_{u,r}} f^2 = -\frac{a-1}{2} \int_{S_{u,r}} \check{\kappa} f^2 + \int_{S_{u,r}} h f.$$

We multiply by r^b which yields

$$\frac{1}{2}e_4 \left(r^b \int_{S_{u,r}} f^2 \right) + \frac{1}{2} \left(a-1-\frac{b}{2} \right) \bar{\kappa} \int_{S_{u,r}} r^b f^2 = -\frac{a-1}{2} r^b \int_{S_{u,r}} \check{\kappa} f^2 + r^b \int_{S_{u,r}} h f$$

where we used the fact that $2e_4(r) = r\bar{\kappa}$. We choose $b > 2a - 2$ and integrate between $r = r_0$ and $r = r_*(u)$, where $r_*(u)$ is such that $S_{u,r_*(u)} \subset \Sigma_*$, which yields

$$\int_{S_{u,r_0}} r^b f^2 + \int_{r_0}^{r_*} \int_{S_{u,r}} r^{b-1} f^2 \lesssim \int_{S_{u,r_*}} r^b f^2 + \int_{r_0}^{r_*} \int_{S_{u,r}} r^{b+1} h^2.$$

Then, integrating in u in $u \in [1, u_*]$, and relying on Remark 8.8.5 we deduce for $r_0 \geq 4m_0$,

$$r_0^b \int_{\{r=r_0\}} f^2 + \int_{(ext)\mathcal{M} \cap \{r \geq r_0\}} r^{b-1} f^2 \lesssim \int_{\Sigma_*} r^b f^2 + \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{b+1} h^2.$$

This concludes the proof of the lemma. \square

Corollary 8.8.8. *Let the following transport equation in $(ext)\mathcal{M}$*

$$e_4(f) + \frac{a}{2}\kappa f = h$$

where $a \in \mathbb{R}$ is a given constant, and f and h are scalar functions. Let $b > 2a - 2$. Then, f satisfies for $5 \leq l \leq k_{large} + 1$

$$\begin{aligned} & \sup_{r_0 \geq 4m_0} \left(r_0^b \int_{\{r=r_0\}} (\mathfrak{d}^k f)^2 \right) + \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{b-1} (\mathfrak{d}^k f)^2 \\ & \lesssim \int_{\Sigma_*} r^b (\mathfrak{d}^{\leq k} f)^2 + \sup_{r_0 \geq 4m_0} \left(r_0^b \int_{\{r=r_0\}} (\mathfrak{d}^{\leq k-1} f)^2 \right) + \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{b-1} (\mathfrak{d}^{\leq k} h)^2 \\ & \quad + \left(\sup_{(ext)\mathcal{M}(\geq 4m_0)} (r^b |\mathfrak{d}^{\leq k-5} f|) \right)^2 \left((ext)\mathfrak{G}_{k-1}^{\geq 4m_0}[\check{\Gamma}] + (ext)\mathfrak{G}_k^{\geq 4m_0}[\check{\kappa}] \right)^2. \end{aligned}$$

Proof. The proof is based on Lemma 8.8.7. It is similar to the one of Corollary 8.8.6 and left to the reader. \square

Lemma 8.8.9. *Let the following transport equation in $(ext)\mathcal{M}$*

$$e_4(f) + \frac{a}{2}\kappa f = h$$

where $a \in \mathbb{R}$ is a given constant, and f and h are scalar functions. Then, f satisfies

$$\sup_{r_T \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} f^2 \lesssim \int_{\{r=4m_0\}} f^2 + \int_{(ext)\mathcal{M}(\leq 4m_0)} h^2.$$

Proof. Let $b > 2a - 2$. Recall from Lemma 8.8.7 the following identity

$$\frac{1}{2}e_4 \left(r^b \int_{S_{u,r}} f^2 \right) + \frac{1}{2} \left(a - 1 - \frac{b}{2} \right) \bar{\kappa} \int_{S_{u,r}} r^b f^2 = -\frac{a-1}{2} r^b \int_{S_{u,r}} \check{\kappa} f^2 + r^b \int_{S_{u,r}} h f.$$

Choosing $b = 2a$, we obtain

$$\frac{1}{2}e_4 \left(r^b \int_{S_{u,r}} f^2 \right) - \frac{1}{2\bar{\kappa}} \int_{S_{u,r}} r^b f^2 = -\frac{a-1}{2} r^b \int_{S_{u,r}} \check{\kappa} f^2 + r^b \int_{S_{u,r}} h f.$$

Next, let $1 \leq u \leq u_*$ and $r_{\mathcal{T}} \leq r_0 \leq 4m_0$. We now integrate in $r_0 \leq r \leq 4m_0$ and along \mathcal{C}_u in ${}^{(ext)}\mathcal{M}$. Since r is bounded on ${}^{(ext)}\mathcal{M}(r \leq 4m_0)$ from above and below, we obtain, for $\epsilon_0 > 0$ small enough,

$$\int_{S_{u,r_0}} f^2 \lesssim \int_{S_{u,4m_0}} f^2 + \int_{r_{\mathcal{T}}}^{4m_0} \int_{S_{u,r}} h^2.$$

We may now integrate in u to deduce

$$\int_1^{u_*} \int_{S_{u,r_0}} f^2 \lesssim \int_1^{u_*} \int_{S_{u,4m_0}} f^2 + \int_1^{u_*} \int_{r_{\mathcal{T}}}^{4m_0} \int_{S_{u,r}} h^2.$$

Relying on Remark 8.8.5 we deduce

$$\sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} f^2 \lesssim \int_{\{r=4m_0\}} f^2 + \int_{{}^{(ext)}\mathcal{M}(\leq 4m_0)} h^2.$$

as desired. This concludes the proof of the lemma. \square

Corollary 8.8.10. *Let the following transport equation in ${}^{(ext)}\mathcal{M}$*

$$e_4(f) + \frac{a}{2}\kappa f = h$$

where $a \in \mathbb{R}$ is a given constant, and f and h are scalar functions. Then, f satisfies for $5 \leq l \leq k_{large} + 1$

$$\begin{aligned} & \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\partial^k f)^2 \\ & \lesssim \int_{\{r=4m_0\}} (\partial^{\leq k} f)^2 + \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\partial^{\leq k-1} f)^2 + \int_{{}^{(ext)}\mathcal{M}(\leq 4m_0)} (\partial^{\leq k} h)^2 \\ & \quad + \left(\sup_{{}^{(ext)}\mathcal{M}(r \leq 4m_0)} (|\partial^{\leq k-5} f|) \right)^2 \left({}^{(ext)}\mathfrak{G}_{k-1}^{\leq 4m_0}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_k^{\leq 4m_0}[\check{\kappa}] \right)^2. \end{aligned}$$

Proof. The proof is based on Lemma 8.8.9. It is similar to the one of Corollary 8.8.6 and left to the reader. \square

8.8.3 Several identities

The goal of this section is to prove the identities below that will be used to avoid losing derivatives when controlling the weighted energies of the Ricci coefficients.

Lemma 8.8.11. *We have*

$$\begin{aligned}
& e_4 \left(\not{d}_1 \not{d}_1^* \kappa - \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right) + 2\kappa \left(\not{d}_1 \not{d}_1^* \kappa - \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right) \\
= & - \left(-\frac{1}{2} \vartheta \not{d}_2^* + \zeta e_4(\Phi) - \beta \right) \not{d}_1^* \kappa - \frac{1}{2} \vartheta \not{d}_1 \not{d}_1^* \kappa + \frac{1}{2} \not{d}_1^* (\kappa + \vartheta) \not{d}_1^* \kappa + (\not{d}_1^* \kappa)^2 \\
& - 2\kappa \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} + (\kappa \vartheta + 2\alpha) \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \\
& + \vartheta \left[\left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1}, e_4 \right] \check{\rho} + \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \left(\frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\
& - \frac{1}{2} \vartheta \not{d}_4 \not{d}_3^* \not{d}_2^{-1} (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) - \frac{1}{2} \vartheta \not{d}_2^* (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \\
& + \frac{1}{2} \not{d}_3^* \not{d}_2^{-1} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_3^* \vartheta + \frac{1}{2} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_2 \vartheta,
\end{aligned}$$

$$\begin{aligned}
& e_4 \left(e_\theta(\mu) + \vartheta \not{d}_2 \not{d}_2^* (\not{d}_1^* \not{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) + 2\kappa \left(e_\theta(\mu) + \vartheta \not{d}_2 \not{d}_2^* (\not{d}_1^* \not{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) \\
= & -\frac{3}{2} \mu e_\theta(\kappa) - \frac{1}{2} \vartheta e_\theta(\mu) - (\kappa \vartheta + 2\alpha) \not{d}_2 \not{d}_2^* (\not{d}_1^* \not{d}_1)^{-1} \underline{\beta} \\
& - \vartheta \left[\not{d}_2 \not{d}_2^* (\not{d}_1^* \not{d}_1)^{-1}, e_4 \right] \underline{\beta} - \vartheta \not{d}_2 \not{d}_2^* (\not{d}_1^* \not{d}_1)^{-1} (\kappa \underline{\beta} + 3\rho \zeta + \underline{\vartheta} \beta) \\
& + \not{d}_3^* \vartheta \not{d}_2^* \not{d}_1^{-1} \check{\rho} - e_\theta \left(\vartheta \not{d}_2^* \not{d}_1^{-1} \left(-\check{\mu} + \frac{1}{4} \vartheta \underline{\vartheta} \right) \right) - e_\theta \left(\vartheta \left(\frac{1}{8} \underline{\kappa} \vartheta + \zeta^2 \right) \right) \\
& - 2\zeta \not{d}_1 \not{d}_1^* \kappa - 2e_\theta(\kappa) \not{d}_2^* \zeta - 2(\kappa \zeta + \beta + \vartheta \zeta) \check{\rho} - 2\zeta \left(\frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\
& + 2\beta \not{d}_2^* \zeta + \frac{3}{2} e_\theta(\kappa \zeta^2) + 2\kappa \left(\vartheta \not{d}_2 \not{d}_2^* (\not{d}_1^* \not{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right),
\end{aligned}$$

and

$$\begin{aligned}
& e_4(e_\theta(\underline{\kappa}) - 4\underline{\beta}) + \kappa(e_\theta(\underline{\kappa}) - 4\underline{\beta}) \\
= & 2e_\theta(\mu) + 12\rho\zeta - \frac{1}{2} \underline{\kappa} e_\theta(\kappa) + 4\underline{\vartheta} \beta - \frac{1}{2} \vartheta e_\theta(\underline{\kappa}) - e_\theta(\vartheta \underline{\vartheta}) + 2e_\theta(\zeta^2) \\
= & 2 \left(e_\theta(\mu) + \vartheta \not{d}_2 \not{d}_2^* (\not{d}_1^* \not{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) + 12\rho\zeta - \frac{1}{2} \underline{\kappa} e_\theta(\kappa) \\
& - 2\vartheta \not{d}_2 \not{d}_2^* (\not{d}_1^* \not{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} + 4\underline{\vartheta} \beta - \frac{1}{2} \vartheta (e_\theta(\underline{\kappa}) - 4\underline{\beta}) - 2\vartheta \underline{\beta} - e_\theta(\vartheta \underline{\vartheta}) + 2e_\theta(\zeta^2).
\end{aligned}$$

Proof. Recall Raychadhuri

$$e_4(\kappa) + \frac{1}{2}\kappa^2 = -\frac{1}{2}\vartheta^2.$$

We commute with $\not{d}_1 \not{d}_1^*$ which yields

$$\begin{aligned} & e_4(\not{d}_1 \not{d}_1^* \kappa) + 2\kappa \not{d}_1 \not{d}_1^* \kappa \\ &= -\left(-\frac{1}{2}\vartheta \not{d}_2^* + \zeta e_4(\Phi) - \beta\right) \not{d}_1^* \kappa - \frac{1}{2}\vartheta \not{d}_1 \not{d}_1^* \kappa + \frac{1}{2} \not{d}_1^* (\kappa + \vartheta) \not{d}_1^* \kappa + (\not{d}_1^* \kappa)^2 - \frac{1}{2} \not{d}_1 \not{d}_1^* (\vartheta^2). \end{aligned}$$

We have in view of Codazzi for ϑ

$$\begin{aligned} \frac{1}{2} \not{d}_1 \not{d}_1^* (\vartheta^2) &= \not{d}_1 (\vartheta \not{d}_1^* \vartheta) \\ &= \frac{1}{2} \not{d}_1 (\vartheta \not{d}_3^* \vartheta - \vartheta \not{d}_2 \vartheta) \\ &= \frac{1}{2} \not{d}_1 \left(\vartheta \not{d}_3^* \not{d}_2^{-1} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) - \vartheta (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \right) \\ &= \frac{1}{2} \vartheta \not{d}_4 \not{d}_3^* \not{d}_2^{-1} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) + \frac{1}{2} \vartheta \not{d}_2^* (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \\ &\quad - \frac{1}{2} \not{d}_3^* \not{d}_2^{-1} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_3^* \vartheta - \frac{1}{2} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_2 \vartheta \\ &= -\vartheta \not{d}_4 \not{d}_3^* \not{d}_2^{-1} \beta - \vartheta \not{d}_2^* \beta + \frac{1}{2} \vartheta \not{d}_4 \not{d}_3^* \not{d}_2^{-1} (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) + \frac{1}{2} \vartheta \not{d}_2^* (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \\ &\quad - \frac{1}{2} \not{d}_3^* \not{d}_2^{-1} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_3^* \vartheta - \frac{1}{2} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_2 \vartheta. \end{aligned}$$

Together with Bianchi for $e_4(\check{\rho})$, we infer

$$\begin{aligned} \frac{1}{2} \not{d}_1 \not{d}_1^* (\vartheta^2) &= -\vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \left(e_4(\check{\rho}) + \frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\ &\quad + \frac{1}{2} \vartheta \not{d}_4 \not{d}_3^* \not{d}_2^{-1} (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) + \frac{1}{2} \vartheta \not{d}_2^* (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \\ &\quad - \frac{1}{2} \not{d}_3^* \not{d}_2^{-1} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_3^* \vartheta - \frac{1}{2} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_2 \vartheta \\ &= -\vartheta e_4 \left(\left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right) - \vartheta \left[\left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1}, e_4 \right] \check{\rho} \\ &\quad - \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \left(\frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\ &\quad + \frac{1}{2} \vartheta \not{d}_4 \not{d}_3^* \not{d}_2^{-1} (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) + \frac{1}{2} \vartheta \not{d}_2^* (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \\ &\quad - \frac{1}{2} \not{d}_3^* \not{d}_2^{-1} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_3^* \vartheta - \frac{1}{2} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_2 \vartheta. \end{aligned}$$

In view of the null structure equation for $e_4(\vartheta)$, we infer

$$\begin{aligned} \frac{1}{2} \not{d}_1 \not{d}_1^*(\vartheta^2) &= -e_4 \left(\vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right) - (\kappa\vartheta + 2\alpha) \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \\ &\quad - \vartheta \left[\left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1}, e_4 \right] \check{\rho} - \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \left(\frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\ &\quad + \frac{1}{2} \vartheta \not{d}_4 \not{d}_3^* \not{d}_2^{-1} (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) + \frac{1}{2} \vartheta \not{d}_2^* (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \\ &\quad - \frac{1}{2} \not{d}_3^* \not{d}_2^{-1} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_3^* \vartheta - \frac{1}{2} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_2 \vartheta. \end{aligned}$$

This yields

$$\begin{aligned} &e_4 \left(\not{d}_1 \not{d}_1^* \kappa - \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right) + 2\kappa \left(\not{d}_1 \not{d}_1^* \kappa - \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right) \\ &= - \left(-\frac{1}{2} \vartheta \not{d}_2^* + \zeta e_4(\Phi) - \beta \right) \not{d}_1^* \kappa - \frac{1}{2} \vartheta \not{d}_1 \not{d}_1^* \kappa + \frac{1}{2} \not{d}_1^* (\kappa + \vartheta) \not{d}_1^* \kappa + (\not{d}_1^* \kappa)^2 \\ &\quad - 2\kappa \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} + (\kappa\vartheta + 2\alpha) \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \\ &\quad + \vartheta \left[\left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1}, e_4 \right] \check{\rho} + \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \left(\frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\ &\quad - \frac{1}{2} \vartheta \not{d}_4 \not{d}_3^* \not{d}_2^{-1} (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) - \frac{1}{2} \vartheta \not{d}_2^* (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \\ &\quad + \frac{1}{2} \not{d}_3^* \not{d}_2^{-1} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_3^* \vartheta + \frac{1}{2} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_2 \vartheta. \end{aligned}$$

Next, recall that we have

$$e_4 \mu + \frac{3}{2} \kappa \mu = -\vartheta \not{d}_2^* \zeta - \vartheta \left(\frac{1}{8} \underline{\kappa} \vartheta + \zeta^2 \right) + \left(2e_\theta(\kappa) - 2\beta + \frac{3}{2} \kappa \zeta \right) \zeta$$

We commute with e_θ which yields

$$\begin{aligned} &e_4(e_\theta(\mu)) + 2\kappa e_\theta(\mu) \\ &= -\frac{3}{2} \mu e_\theta(\kappa) - \frac{1}{2} \vartheta e_\theta(\mu) - e_\theta \left(\vartheta \not{d}_2^* \not{d}_1^{-1} \left(-\check{\mu} - \check{\rho} + \frac{1}{4} \vartheta \underline{\vartheta} \right) \right) - e_\theta \left(\vartheta \left(\frac{1}{8} \underline{\kappa} \vartheta + \zeta^2 \right) \right) \\ &\quad + e_\theta \left(\left(2e_\theta(\kappa) - 2\beta + \frac{3}{2} \kappa \zeta \right) \zeta \right) \\ &= -\frac{3}{2} \mu e_\theta(\kappa) - \frac{1}{2} \vartheta e_\theta(\mu) + \vartheta \not{d}_2 \not{d}_2^* (\not{d}_1^* \not{d}_1)^{-1} \not{d}_1^* \rho + \not{d}_3^* \vartheta \not{d}_2^* \not{d}_1^{-1} \check{\rho} \\ &\quad - e_\theta \left(\vartheta \not{d}_2^* \not{d}_1^{-1} \left(-\check{\mu} + \frac{1}{4} \vartheta \underline{\vartheta} \right) \right) - e_\theta \left(\vartheta \left(\frac{1}{8} \underline{\kappa} \vartheta + \zeta^2 \right) \right) \\ &\quad - 2\zeta \not{d}_1 \not{d}_1^* \kappa - 2e_\theta(\kappa) \not{d}_2^* \zeta - 2\zeta \not{d}_1 \beta + 2\beta \not{d}_2^* \zeta + \frac{3}{2} e_\theta(\kappa \zeta^2). \end{aligned}$$

Now, using the Bianchi identities for $e_4(\underline{\beta})$ and $e_4(\underline{\rho})$, we have

$$\begin{aligned}
\vartheta \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} \not\phi_1^* \rho &= -\vartheta \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} (e_4 \underline{\beta} + \kappa \underline{\beta} + 3\rho \zeta + \vartheta \underline{\beta}) \\
&= -e_4 \left(\vartheta \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} \underline{\beta} \right) + e_4(\vartheta) \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} \underline{\beta} - \vartheta \left[\not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1}, e_4 \right] \underline{\beta} \\
&\quad - \vartheta \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} (\kappa \underline{\beta} + 3\rho \zeta + \vartheta \underline{\beta}) \\
&= -e_4 \left(\vartheta \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} \underline{\beta} \right) - (\kappa \vartheta + 2\alpha) \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} \underline{\beta} \\
&\quad - \vartheta \left[\not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1}, e_4 \right] \underline{\beta} - \vartheta \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} (\kappa \underline{\beta} + 3\rho \zeta + \vartheta \underline{\beta})
\end{aligned}$$

and

$$\begin{aligned}
\zeta \not\phi_1 \beta &= \zeta \left(e_4(\underline{\rho}) + \frac{3}{2} \bar{\kappa} \underline{\rho} + \frac{3}{2} \bar{\rho} \bar{\kappa} - \text{Err}[e_4 \underline{\rho}] \right) \\
&= e_4(\zeta \underline{\rho}) - e_4(\zeta) \underline{\rho} + \zeta \left(\frac{3}{2} \bar{\kappa} \underline{\rho} + \frac{3}{2} \bar{\rho} \bar{\kappa} - \text{Err}[e_4 \underline{\rho}] \right) \\
&= e_4(\zeta \underline{\rho}) + (\kappa \zeta + \beta + \vartheta \zeta) \underline{\rho} + \zeta \left(\frac{3}{2} \bar{\kappa} \underline{\rho} + \frac{3}{2} \bar{\rho} \bar{\kappa} - \text{Err}[e_4 \underline{\rho}] \right).
\end{aligned}$$

We infer

$$\begin{aligned}
&e_4 \left(e_\theta(\mu) \right) + \vartheta \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} \underline{\beta} + 2\zeta \underline{\rho} \Big) + 2\kappa e_\theta(\mu) \\
&= -\frac{3}{2} \mu e_\theta(\kappa) - \frac{1}{2} \vartheta e_\theta(\mu) - (\kappa \vartheta + 2\alpha) \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} \underline{\beta} \\
&\quad - \vartheta \left[\not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1}, e_4 \right] \underline{\beta} - \vartheta \not\phi_2 \not\phi_2^* (\not\phi_1^* \not\phi_1)^{-1} (\kappa \underline{\beta} + 3\rho \zeta + \vartheta \underline{\beta}) \\
&\quad + \not\phi_3^* \vartheta \not\phi_2^* \not\phi_1^{-1} \underline{\rho} - e_\theta \left(\vartheta \not\phi_2^* \not\phi_1^{-1} \left(-\check{\mu} + \frac{1}{4} \vartheta \vartheta \right) \right) - e_\theta \left(\vartheta \left(\frac{1}{8} \kappa \vartheta + \zeta^2 \right) \right) \\
&\quad - 2\zeta \not\phi_1 \not\phi_1^* \kappa - 2e_\theta(\kappa) \not\phi_2^* \zeta - 2(\kappa \zeta + \beta + \vartheta \zeta) \underline{\rho} - 2\zeta \left(\frac{3}{2} \bar{\kappa} \underline{\rho} + \frac{3}{2} \bar{\rho} \bar{\kappa} - \text{Err}[e_4 \underline{\rho}] \right) \\
&\quad + 2\beta \not\phi_2^* \zeta + \frac{3}{2} e_\theta(\kappa \zeta^2)
\end{aligned}$$

and hence

$$\begin{aligned}
& e_4\left(e_\theta(\mu) + \vartheta \phi_2 \phi_2^* (\phi_1^* \phi_1)^{-1} \underline{\beta} + 2\zeta \check{\rho}\right) + 2\kappa\left(e_\theta(\mu) + \vartheta \phi_2 \phi_2^* (\phi_1^* \phi_1)^{-1} \underline{\beta} + 2\zeta \check{\rho}\right) \\
= & -\frac{3}{2}\mu e_\theta(\kappa) - \frac{1}{2}\vartheta e_\theta(\mu) - (\kappa\vartheta + 2\alpha) \phi_2 \phi_2^* (\phi_1^* \phi_1)^{-1} \underline{\beta} \\
& -\vartheta \left[\phi_2 \phi_2^* (\phi_1^* \phi_1)^{-1}, e_4 \right] \underline{\beta} - \vartheta \phi_2 \phi_2^* (\phi_1^* \phi_1)^{-1} (\kappa \underline{\beta} + 3\rho \zeta + \vartheta \underline{\beta}) \\
& + \phi_3^* \vartheta \phi_2^* \phi_1^{-1} \check{\rho} - e_\theta \left(\vartheta \phi_2^* \phi_1^{-1} \left(-\check{\mu} + \frac{1}{4} \vartheta \vartheta \right) \right) - e_\theta \left(\vartheta \left(\frac{1}{8} \kappa \vartheta + \zeta^2 \right) \right) \\
& - 2\zeta \phi_1 \phi_1^* \kappa - 2e_\theta(\kappa) \phi_2^* \zeta - 2(\kappa \zeta + \beta + \vartheta \zeta) \check{\rho} - 2\zeta \left(\frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\
& + 2\beta \phi_2^* \zeta + \frac{3}{2} e_\theta(\kappa \zeta^2) + 2\kappa \left(\vartheta \phi_2 \phi_2^* (\phi_1^* \phi_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right).
\end{aligned}$$

Finally, recall that we have

$$\begin{aligned}
e_4(\underline{\kappa}) + \frac{1}{2} \kappa \underline{\kappa} &= -2 \phi_1 \zeta + 2\rho - \frac{1}{2} \vartheta \vartheta + 2\zeta^2 \\
&= 2\mu + 4\rho - \vartheta \vartheta + 2\zeta^2.
\end{aligned}$$

We commute with e_θ which yields

$$e_4(e_\theta(\underline{\kappa})) + \kappa e_\theta(\underline{\kappa}) = 2e_\theta(\mu) + 4e_\theta(\rho) - \frac{1}{2} \kappa e_\theta(\kappa) - \frac{1}{2} \vartheta e_\theta(\underline{\kappa}) - e_\theta(\vartheta \vartheta) + 2e_\theta(\zeta^2).$$

Together with Bianchi for $e_4(\underline{\beta})$, we infer

$$\begin{aligned}
& e_4(e_\theta(\underline{\kappa}) - 4\underline{\beta}) + \kappa(e_\theta(\underline{\kappa}) - 4\underline{\beta}) \\
= & 2e_\theta(\mu) + 12\rho\zeta - \frac{1}{2} \kappa e_\theta(\kappa) + 4\underline{\beta} - \frac{1}{2} \vartheta e_\theta(\underline{\kappa}) - e_\theta(\vartheta \vartheta) + 2e_\theta(\zeta^2) \\
= & 2\left(e_\theta(\mu) + \vartheta \phi_2 \phi_2^* (\phi_1^* \phi_1)^{-1} \underline{\beta} + 2\zeta \check{\rho}\right) + 12\rho\zeta - \frac{1}{2} \kappa e_\theta(\kappa) \\
& - 2\vartheta \phi_2 \phi_2^* (\phi_1^* \phi_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} + 4\underline{\beta} - \frac{1}{2} \vartheta (e_\theta(\underline{\kappa}) - 4\underline{\beta}) - 2\vartheta \underline{\beta} - e_\theta(\vartheta \vartheta) + 2e_\theta(\zeta^2).
\end{aligned}$$

This concludes the proof of the lemma. \square

8.8.4 Proof of Proposition 8.8.2

We introduce the following notation which will constantly appear on the RHS of the equalities below

$$\begin{aligned}
N^{\geq 4m_0}[J, \check{\Gamma}, \check{R}] &:= \binom{\Sigma_*}{\Sigma_*} \mathfrak{G}_{J+1}[\check{\Gamma}] + \binom{\Sigma_*}{\Sigma_*} \mathfrak{G}'_{J+1}[\check{\Gamma}] + \binom{ext}{ext} \mathfrak{R}_{J+1}[\check{R}] + \binom{ext}{ext} \mathfrak{G}_J[\check{\Gamma}] \\
&+ \epsilon_0 \left(\binom{ext}{ext} \mathfrak{G}_{J+1}^{\geq 4m_0}[\check{\Gamma}] + \binom{ext}{ext} \mathfrak{G}_{J+1}^{\geq 4m_0'}[\check{\Gamma}] \right). \tag{8.8.2}
\end{aligned}$$

Step 1. Recall that

$$e_4(\vartheta) + \kappa\vartheta = -2\alpha.$$

In view of Corollary 8.8.6 with $a = 2$, we have for any $r_0 \geq 4m_0$

$$\max_{k \leq J+1} \sup_{r_0 \geq 4m_0} r_0^2 \int_{\{r=r_0\}} (\mathfrak{d}^k \vartheta)^2 \lesssim \left(N^{\geq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2.$$

Step 2. Next, recall that

$$e_4(\check{\kappa}) + \bar{\kappa} \check{\kappa} = -\frac{1}{4}\vartheta^2 + \frac{1}{4}\bar{\vartheta}^2 - \bar{\kappa}^2.$$

In view of Corollary 8.8.6 with $a = 2$, we have for any $r_0 \geq 4m_0$

$$\max_{k \leq J+2} \sup_{r_0 \geq 4m_0} r_0^2 \int_{\{r=r_0\}} (\mathfrak{d}^k \check{\kappa})^2 \lesssim \left(N^{\geq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the null structure equations for $e_4(\vartheta)$, $e_3(\vartheta)$ and $\not\partial_2 \vartheta$ to avoid a loss of one derivative for the RHS.

Step 3. Next, recall that

$$e_4(\zeta) + \kappa\zeta = -\beta - \vartheta\zeta.$$

In view of Corollary 8.8.6 with $a = 2$, we have for any $r_0 \geq 4m_0$

$$\max_{k \leq J+1} \sup_{r_0 \geq 4m_0} r_0^2 \int_{\{r=r_0\}} (\mathfrak{d}^k \zeta)^2 \lesssim \left(N^{\geq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2.$$

Step 4. Next, recall that

$$e_4(\check{\mu}) + \frac{3}{2}\bar{\kappa}\check{\mu} = -\frac{3}{2}\bar{\mu}\check{\kappa} + \text{Err}[e_4\check{\mu}].$$

In view of Corollary 8.8.6 with $a = 3$, commuting with $\not\partial$ and \mathbf{T} , we have for any $r_0 \geq 4m_0$

$$\max_{k \leq J+1} \sup_{r_0 \geq 4m_0} r_0^4 \int_{\{r=r_0\}} (\mathfrak{d}^k \check{\mu})^2 \lesssim \left(N^{\geq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we used the estimates for $\check{\kappa}$ on $^{(ext)}\mathcal{M}$ derived in Step 2.

Step 5. Next, recall that

$$e_4(\check{\underline{\kappa}}) + \frac{1}{2}\bar{\kappa}\check{\underline{\kappa}} = -\frac{1}{2}\bar{\kappa}\check{\underline{\kappa}} + 2\check{\rho} - 2\not\partial_1 \zeta + \text{Err}[e_4\check{\underline{\kappa}}].$$

In view of Corollary 8.8.8 with $a = 1$ and $b = 2 - \delta_B$ which satisfy the constraint $b > 2a - 2$, we have for any $r_0 \geq 4m_0$

$$\max_{k \leq J+1} \sup_{r_0 \geq 4m_0} \left(r_0^{2-\delta_B} \int_{\{r=r_0\}} (\mathfrak{d}^k \check{\kappa})^2 \right) + \int_{(ext)\mathcal{M}(r \geq 4m_0)} r^{1-\delta_B} (\mathfrak{d}^k \check{\kappa})^2 \lesssim \left(N^{\geq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we used the estimates for $\check{\kappa}$ and $\check{\mu}$ on $(ext)\mathcal{M}$ derived respectively in Step 2 and Step 4.

Step 6. Next, recall that

$$\begin{aligned} e_4(\vartheta) + \frac{1}{2} \kappa \vartheta &= 2 \phi_2^* \zeta - \frac{1}{2} \kappa \vartheta + 2\zeta^2 \\ &= 2 \phi_2^* \phi_1^{-1} \left(-\mu - \rho + \frac{1}{4} \vartheta \vartheta \right) - \frac{1}{2} \kappa \vartheta + 2\zeta^2. \end{aligned}$$

In view of Corollary 8.8.6 with $a = 1$, we have for any $r_0 \geq 4m_0$

$$\max_{k \leq J+1} \sup_{r_0 \geq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \vartheta)^2 \lesssim \left(N^{\geq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we used the estimates for ϑ and $\check{\mu}$ on $(ext)\mathcal{M}$ derived respectively in Step 1 and Step 4.

Step 7. Next, recall that

$$e_4(\check{\omega}) = \check{\rho} + 3\zeta^2 - 3\bar{\zeta}^2 - \bar{\kappa} \check{\omega}.$$

In view of Corollary 8.8.8 with $a = 0$ and $b = 0$ which satisfy the constraint $b > 2a - 2$, we have for any $r_0 \geq 4m_0$

$$\max_{k \leq J+1} \sup_{r_0 \geq 4m_0} \left(\int_{\{r=r_0\}} (\mathfrak{d}^k \check{\omega})^2 \right) + \int_{(ext)\mathcal{M}(\geq 4m_0)} r^{-1} (\mathfrak{d}^k \check{\omega})^2 \lesssim \left(N^{\geq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2.$$

Step 8. In order to estimate $\underline{\xi}$ in Step 9, we derive an estimate for $e_3(\zeta) + \underline{\beta}$. Recall that we have

$$e_4(\zeta) + \kappa \zeta = -\beta - \vartheta \zeta.$$

Commuting with e_3 , we infer

$$e_4(e_3(\zeta)) + [e_3, e_4]\zeta + \kappa e_3(\zeta) + e_3(\kappa)\zeta = -e_3(\beta) - \vartheta \zeta.$$

In view of the null structure equation for $e_3(\kappa)$, the Bianchi identity for $e_3(\beta)$ and the commutator identity for $[e_3, e_4]$, we infer

$$\begin{aligned} & e_4(e_3(\zeta)) + \left(2\underline{\omega}e_4 + 4\zeta e_\theta\right)\zeta + \kappa e_3(\zeta) + \left(-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa + 2\underline{\phi}_1\zeta + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\zeta^2\right)\zeta \\ &= (\underline{\kappa} - 2\underline{\omega})\beta + \underline{\phi}_1^*\rho - 3\zeta\rho + \vartheta\underline{\beta} - \underline{\xi}\alpha - \vartheta\zeta. \end{aligned}$$

Together with the null structure equation for $e_4(\zeta)$, the Bianchi identity for $e_4(\underline{\beta})$ to get rid of the term $\underline{\phi}_1^*\rho$, and the definition of μ , we infer

$$\begin{aligned} & e_4(e_3(\zeta)) + 2\underline{\omega}(-\kappa\zeta - \beta - \vartheta\zeta) + 4\zeta\underline{\phi}_1^*\underline{\phi}_1^{-1}\left(\check{\mu} + \check{\rho} - \frac{1}{4}\vartheta\underline{\vartheta} + \frac{1}{4}\overline{\vartheta\underline{\vartheta}}\right) \\ & + \kappa e_3(\zeta) + \left(-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa - 2\mu + 2\zeta^2\right)\zeta \\ &= (\underline{\kappa} - 2\underline{\omega})\beta - e_4(\underline{\beta}) - \kappa\underline{\beta} - 3\zeta\rho - \vartheta\underline{\beta} - 3\zeta\rho + \vartheta\underline{\beta} - \underline{\xi}\alpha - \vartheta\zeta. \end{aligned}$$

and hence

$$\begin{aligned} & e_4(e_3(\zeta) + \underline{\beta}) + \kappa(e_3(\zeta) + \underline{\beta}) \\ &= \underline{\kappa}\beta + \left(\frac{1}{2}\kappa\underline{\kappa} + 2\mu - 6\rho\right)\zeta \\ & \quad - \vartheta\underline{\beta} + \vartheta\underline{\beta} - \underline{\xi}\alpha - \vartheta\zeta + 2\underline{\omega}\vartheta\zeta - 4\zeta\underline{\phi}_1^*\underline{\phi}_1^{-1}\left(\check{\mu} + \check{\rho} - \frac{1}{4}\vartheta\underline{\vartheta} + \frac{1}{4}\overline{\vartheta\underline{\vartheta}}\right) - 2\zeta^3 \end{aligned}$$

In view of Corollary 8.8.6 with $a = 2$, we have for any $r_0 \geq 4m_0$

$$\max_{k \leq J+1} \sup_{r_0 \geq 4m_0} r_0^2 \int_{\{r=r_0\}} (\mathfrak{d}^k(e_3(\zeta) + \underline{\beta}))^2 \lesssim \left(N^{\geq 4m_0}[J, \check{\Gamma}, \check{R}]\right)^2$$

where we used the estimates for ζ derived in Step 3.

Step 9. Next, recall that we have

$$\begin{aligned} e_4(\underline{\xi}) &= -e_3(\zeta) + \underline{\beta} - \underline{\kappa}\zeta - \zeta\underline{\vartheta} \\ &= -(e_3(\zeta) + \underline{\beta}) + 2\underline{\beta} - \underline{\kappa}\zeta - \zeta\underline{\vartheta}. \end{aligned}$$

In view of Corollary 8.8.8 with $a = 0$ and $b = -\delta_B$ which satisfy the constraint $b > 2a - 2$, we have for any $r_0 \geq 4m_0$

$$\max_{k \leq J+1} \sup_{r_0 \geq 4m_0} r_0^{-\delta_B} \left(\int_{\{r=r_0\}} (\mathfrak{d}^k \underline{\xi})^2 \right) + \int_{(ext)\mathcal{M}(r \geq 4m_0)} r^{-1-\delta_B} (\mathfrak{d}^k \underline{\xi})^2 \lesssim \left(N^{\geq 4m_0}[J, \check{\Gamma}, \check{R}]\right)^2$$

where we used the estimates for ζ and $e_3(\zeta) + \underline{\beta}$ on $(ext)\mathcal{M}$ derived respectively in Step 3 and Step 8.

Step 10. Recall that we have

$$\begin{aligned}
& e_4 \left(\not{d}_1 \not{d}_1^* \kappa - \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right) + 2\kappa \left(\not{d}_1 \not{d}_1^* \kappa - \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right) \\
= & - \left(-\frac{1}{2} \vartheta \not{d}_2^* + \zeta e_4(\Phi) - \beta \right) \not{d}_1^* \kappa - \frac{1}{2} \vartheta \not{d}_1 \not{d}_1^* \kappa + \frac{1}{2} \not{d}_1^* (\kappa + \vartheta) \not{d}_1^* \kappa + (\not{d}_1^* \kappa)^2 \\
& - 2\kappa \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} + (\kappa \vartheta + 2\alpha) \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \\
& + \vartheta \left[\left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1}, e_4 \right] \check{\rho} + \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \left(\frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\
& - \frac{1}{2} \vartheta \not{d}_4 \not{d}_3^* \not{d}_2^{-1} (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) - \frac{1}{2} \vartheta \not{d}_2^* (-\not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \\
& + \frac{1}{2} \not{d}_3^* \not{d}_2^{-1} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_3^* \vartheta + \frac{1}{2} (-2\beta - \not{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \not{d}_2 \vartheta.
\end{aligned}$$

In view of Corollary 8.8.6 with $a = 4$, we have

$$\max_{k \leq J+1} \sup_{r_0 \geq 4m_0} \int_{\{r=r_0\}} r_0^6 \left(\not{d}^k \left(\not{d}_1 \not{d}_1^* \kappa - \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right) \right)^2 \lesssim \left(N^{\geq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used

- the fact that $\not{\vartheta} \vartheta = \not{\vartheta} \not{d}_2^{-1} \not{d}_2 \vartheta$ and Codazzi for ϑ to estimate the terms of the RHS with one angular derivative of ϑ ,
- the estimates of Step 2 to estimate the terms of the RHS with one derivative of $\check{\kappa}$,
- the fact that $\not{\vartheta} \zeta = \not{\vartheta} \not{d}_1^{-1} \not{d}_1 \zeta$ and the definition of μ to estimate terms of the RHS with one angular derivative of ζ ,
- the identity

$$\begin{aligned}
\not{\vartheta} \not{d}_1^* \kappa &= \not{\vartheta} \not{d}_1^{-1} \left(\vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right) \\
&\quad + \not{\vartheta} \not{d}_1^{-1} \left(\not{d}_1 \not{d}_1^* \kappa - \vartheta \left(\not{d}_4 \not{d}_3^* \not{d}_2^{-1} + \not{d}_2^* \right) \not{d}_1^{-1} \check{\rho} \right)
\end{aligned}$$

to estimate the terms of the RHS with two angular derivatives of $\check{\kappa}$.

Step 11. Recall that we have

$$\begin{aligned}
& e_4 \left(e_\theta(\mu) + \vartheta \mathfrak{d}_2 \mathfrak{d}_2^* (\mathfrak{d}_1^* \mathfrak{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) + 2\kappa \left(e_\theta(\mu) + \vartheta \mathfrak{d}_2 \mathfrak{d}_2^* (\mathfrak{d}_1^* \mathfrak{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) \\
= & -\frac{3}{2} \mu e_\theta(\kappa) - \frac{1}{2} \vartheta e_\theta(\mu) - (\kappa \vartheta + 2\alpha) \mathfrak{d}_2 \mathfrak{d}_2^* (\mathfrak{d}_1^* \mathfrak{d}_1)^{-1} \underline{\beta} \\
& - \vartheta \left[\mathfrak{d}_2 \mathfrak{d}_2^* (\mathfrak{d}_1^* \mathfrak{d}_1)^{-1}, e_4 \right] \underline{\beta} - \vartheta \mathfrak{d}_2 \mathfrak{d}_2^* (\mathfrak{d}_1^* \mathfrak{d}_1)^{-1} (\kappa \underline{\beta} + 3\rho \zeta + \underline{\vartheta} \beta) \\
& + \mathfrak{d}_3^* \vartheta \mathfrak{d}_2^* \mathfrak{d}_1^{-1} \check{\rho} - e_\theta \left(\vartheta \mathfrak{d}_2^* \mathfrak{d}_1^{-1} \left(-\check{\mu} + \frac{1}{4} \vartheta \underline{\vartheta} \right) \right) - e_\theta \left(\vartheta \left(\frac{1}{8} \underline{\kappa} \vartheta + \zeta^2 \right) \right) \\
& - 2\zeta \mathfrak{d}_1 \mathfrak{d}_1^* \kappa - 2e_\theta(\kappa) \mathfrak{d}_2^* \zeta - 2(\kappa \zeta + \beta + \vartheta \zeta) \check{\rho} - 2\zeta \left(\frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\
& + 2\beta \mathfrak{d}_2^* \zeta + \frac{3}{2} e_\theta(\kappa \zeta^2) + 2\kappa \left(\vartheta \mathfrak{d}_2 \mathfrak{d}_2^* (\mathfrak{d}_1^* \mathfrak{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right).
\end{aligned}$$

In view of Corollary 8.8.6 with $a = 4$, we have

$$\begin{aligned}
& \max_{k \leq J+1} \sup_{r_0 \geq 4m_0} r_0^6 \int_{\{r=r_0\}} \left(\mathfrak{d}^k \left(e_\theta(\mu) + \vartheta \mathfrak{d}_2 \mathfrak{d}_2^* (\mathfrak{d}_1^* \mathfrak{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) \right)^2 \\
& \lesssim \left(N^{\geq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2 + \epsilon^2 \int_{(ext) \mathcal{M}(\geq 4m_0)} \left(\mathfrak{d}^{\leq J+1} \left(e_\theta(\underline{\kappa}) - 4\underline{\beta} \right) \right)^2,
\end{aligned}$$

where we have used

- the fact that $\mathfrak{d}\vartheta = \mathfrak{d}\mathfrak{d}_2^{-1} \mathfrak{d}_2 \vartheta$ and Codazzi for ϑ to estimate the terms of the RHS with one angular derivative of ϑ ,
- the estimates of Step 2 to estimate the terms of the RHS with one derivative of $\check{\kappa}$,
- the fact that $\mathfrak{d}\zeta = \mathfrak{d}\mathfrak{d}_1^{-1} \mathfrak{d}_1 \zeta$ and the definition of μ to estimate terms of the RHS with one angular derivative of ζ ,
- the fact that $e_\theta(\underline{\kappa}) = (e_\theta(\underline{\kappa}) - 4\underline{\beta}) + 4\underline{\beta}$ to estimate the term with one angular derivative of $\underline{\kappa}$,
- the identity

$$\begin{aligned}
\mathfrak{d} \mathfrak{d}_1^* \kappa &= \mathfrak{d} \mathfrak{d}_1^{-1} \left(\vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} \right) \\
&+ \mathfrak{d} \mathfrak{d}_1^{-1} \left(\mathfrak{d}_1 \mathfrak{d}_1^* \kappa - \vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} \right)
\end{aligned}$$

and the estimates of Step 10 to estimate the terms of the RHS with two angular derivatives of $\check{\kappa}$.

Step 12. Recall that we have

$$\begin{aligned}
& e_4(e_\theta(\underline{\kappa}) - 4\underline{\beta}) + \kappa(e_\theta(\underline{\kappa}) - 4\underline{\beta}) \\
&= 2e_\theta(\mu) + 12\rho\zeta - \frac{1}{2}\underline{\kappa}e_\theta(\kappa) + 4\underline{\vartheta}\underline{\beta} - \frac{1}{2}\vartheta e_\theta(\underline{\kappa}) - e_\theta(\vartheta\underline{\vartheta}) + 2e_\theta(\zeta^2) \\
&= 2\left(e_\theta(\mu) + \vartheta \not\!{d}_2 \not\!{d}_2^*(\not\!{d}_1^* \not\!{d}_1)^{-1}\underline{\beta} + 2\zeta\check{\rho}\right) + 12\rho\zeta - \frac{1}{2}\underline{\kappa}e_\theta(\kappa) \\
&\quad - 2\vartheta \not\!{d}_2 \not\!{d}_2^*(\not\!{d}_1^* \not\!{d}_1)^{-1}\underline{\beta} + 2\zeta\check{\rho} + 4\underline{\vartheta}\underline{\beta} - \frac{1}{2}\vartheta(e_\theta(\underline{\kappa}) - 4\underline{\beta}) - 2\vartheta\underline{\beta} - e_\theta(\vartheta\underline{\vartheta}) + 2e_\theta(\zeta^2).
\end{aligned}$$

In view of Corollary 8.8.6 with $a = 2$, we have

$$\begin{aligned}
& \max_{k \leq J+1} \sup_{r_0 \geq 4m_0} r_0^2 \int_{\{r=r_0\}} (\not\!{d}^k (e_\theta(\underline{\kappa}) - 4\underline{\beta}))^2 \\
& \lesssim \left(N^{\geq 4m_0}[J, \check{\Gamma}, \check{R}]\right)^2 + \int_{(e_{xt})\mathcal{M}(\geq 4m_0)} r^4 (\not\!{d}^{\leq J+1} (e_\theta(\mu) + \vartheta \not\!{d}_2 \not\!{d}_2^*(\not\!{d}_1^* \not\!{d}_1)^{-1}\underline{\beta} + 2\zeta\check{\rho}))^2,
\end{aligned}$$

where we have used

- the fact that $\not\!{d}\vartheta = \not\!{d}\not\!{d}_2^{-1}\not\!{d}_2\vartheta$ and Codazzi for ϑ to estimate the terms of the RHS with one angular derivative of ϑ ,
- the fact that $\not\!{d}\underline{\vartheta} = \not\!{d}\not\!{d}_2^{-1}\not\!{d}_2\underline{\vartheta}$ and Codazzi for $\underline{\vartheta}$ to estimate the terms of the RHS with one angular derivative of $\underline{\vartheta}$,
- the estimates of Step 2 to estimate the terms of the RHS with one derivative of $\check{\kappa}$,
- the fact that $\not\!{d}\zeta = \not\!{d}\not\!{d}_1^{-1}\not\!{d}_1\zeta$ and the definition of μ to estimate terms of the RHS with one angular derivative of ζ ,
- the estimate for ζ of Step 3.

Together with the estimate of Step 11, we infer

$$\begin{aligned}
& \max_{k \leq J+1} \sup_{r_0 \geq 4m_0} r_0^6 \int_{\{r=r_0\}} (\not\!{d}^k (e_\theta(\mu) + \vartheta \not\!{d}_2 \not\!{d}_2^*(\not\!{d}_1^* \not\!{d}_1)^{-1}\underline{\beta} + 2\zeta\check{\rho}))^2 \\
& \quad + \max_{k \leq J+1} \sup_{r_0 \geq 4m_0} r_0^2 \int_{\{r=r_0\}} (\not\!{d}^k (e_\theta(\underline{\kappa}) - 4\underline{\beta}))^2 \lesssim \left(N^{\geq 4m_0}[J, \check{\Gamma}, \check{R}]\right)^2.
\end{aligned}$$

Finally, we have obtained

$$\begin{aligned}
& \max_{k \leq J+1} \sup_{r_0 \geq 4m_0} \left(\int_{\{r=r_0\}} \left(r_0^4 (\not\!{d}^k \check{\mu})^2 + r_0^2 (\not\!{d}^k \vartheta)^2 + r_0^2 (\not\!{d}^k \zeta)^2 + r_0^2 (\not\!{d}^k (e_3(\zeta) + \underline{\beta}))^2 \right. \right. \\
& \quad \left. \left. + r_0^{2-\delta_B} (\not\!{d}^k \check{\kappa})^2 + (\not\!{d}^k \underline{\vartheta})^2 + (\not\!{d}^k \check{\omega})^2 + r_0^{-\delta_B} (\not\!{d}^k \underline{\xi})^2 \right) \right) \lesssim \left(N^{\geq 4m_0}[J, \check{\Gamma}, \check{R}]\right)^2,
\end{aligned}$$

$$\max_{k \leq J+2} \sup_{r_0 \geq 4m_0} \left(r_0^2 \int_{\{r=r_0\}} \left(\mathfrak{d}^k \left(\kappa - \frac{2}{r} \right) \right)^2 \right) \lesssim \left(N^{\geq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2,$$

and

$$\begin{aligned} \max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \left(\int_{\{r=r_0\}} \left\{ r_0^6 \left(\mathfrak{d}^k \left(\mathfrak{d}_1 \mathfrak{d}_1^* \kappa - \vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \mathfrak{d}_1^{-1} \check{\rho} \right) \right)^2 \right. \right. \\ \left. \left. + r_0^6 \left(\mathfrak{d}^k \left(e_\theta(\mu) + \vartheta \mathfrak{d}_2 \mathfrak{d}_2^* (\mathfrak{d}_1^* \mathfrak{d}_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) \right)^2 \right. \right. \\ \left. \left. + r_0^2 \left(\mathfrak{d}^k \left(e_\theta(\underline{\kappa}) - 4\underline{\beta} \right) \right)^2 \right\} \right) \lesssim \left(N^{\geq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2. \end{aligned}$$

In view of the definition (8.8.2) of $N^{\geq 4m_0}[J, \check{\Gamma}, \check{R}]$, and of the various norms, we infer

$$\begin{aligned} (ext) \mathfrak{G}_{J+1}^{\geq 4m_0}[\check{\Gamma}] + (ext) \mathfrak{G}_{J+1}^{\geq 4m_0'}[\check{\Gamma}] &\lesssim (\Sigma_*) \mathfrak{G}_{J+1}[\check{\Gamma}] + (\Sigma_*) \mathfrak{G}'_{J+1}[\check{\Gamma}] + (ext) \mathfrak{R}_{J+1}[\check{R}] + (ext) \mathfrak{G}_J[\check{\Gamma}] \\ &+ \epsilon_0 \left((ext) \mathfrak{G}_{J+1}^{\geq 4m_0}[\check{\Gamma}] + (ext) \mathfrak{G}_{J+1}^{\geq 4m_0'}[\check{\Gamma}] \right) \end{aligned}$$

and hence, for ϵ_0 small enough,

$$(ext) \mathfrak{G}_{J+1}^{\geq 4m_0}[\check{\Gamma}] + (ext) \mathfrak{G}_{J+1}^{\geq 4m_0'}[\check{\Gamma}] \lesssim (\Sigma_*) \mathfrak{G}_{J+1}[\check{\Gamma}] + (\Sigma_*) \mathfrak{G}'_{J+1}[\check{\Gamma}] + (ext) \mathfrak{R}_{J+1}[\check{R}] + (ext) \mathfrak{G}_J[\check{\Gamma}].$$

This concludes the proof of Proposition 8.8.2.

8.8.5 Proof of Proposition 8.8.3

In the proof below, we will repeatedly use the following estimate

$$\begin{aligned} &\max_{k \leq J+1} \int_{(ext) \mathcal{M}(r \leq 4m_0)} (\mathfrak{d}^k f)^2 \\ &\lesssim \max_{k \leq J} \int_{(ext) \mathcal{M}(r \leq 4m_0)} \left((\mathfrak{d}^k f)^2 + (\mathfrak{d}^k \mathbf{N}f)^2 + (\mathfrak{d}^k e_4 f)^2 + (\mathfrak{d}^k \check{\rho} f)^2 \right) \end{aligned} \quad (8.8.3)$$

which follows from the fact that $\mathfrak{d} = (\check{\rho}, re_4, e_3)$ and $e_3 = \Upsilon e_4 - 2\mathbf{N}$, where we recall that

$$\mathbf{N} = \frac{1}{2} (\Upsilon e_4 - e_3).$$

Also, we introduce the following notation which will constantly appear on the RHS of the equalities below

$$\begin{aligned} N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] &:= (ext) \mathfrak{G}_{J+1}^{\geq 4m_0}[\check{\Gamma}] + (ext) \mathfrak{G}_{J+1}^{\geq 4m_0'}[\check{\Gamma}] + (ext) \mathfrak{R}_{J+1}[\check{R}] + (ext) \mathfrak{G}_J[\check{\Gamma}] \\ &+ \epsilon_0 \left((ext) \mathfrak{G}_{J+1}^{\leq 4m_0}[\check{\Gamma}] + (ext) \mathfrak{G}_{J+1}^{\leq 4m_0'}[\check{\Gamma}] \right). \end{aligned} \quad (8.8.4)$$

Step 1. Recall that

$$e_4(\check{\kappa}) + \kappa\check{\kappa} = \text{Err}[e_4\check{\kappa}].$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J+2} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \check{\kappa})^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the null structure equations for $e_4(\vartheta)$, $e_3(\vartheta)$ and $\not{d}_2\vartheta$ to avoid losing one derivative.

Step 2. Next, recall that

$$e_4(\check{\mu}) + \frac{3}{2}\bar{\kappa}\check{\mu} = -\frac{3}{2}\bar{\mu}\check{\kappa} + \text{Err}[e_4\check{\mu}].$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \check{\mu})^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimates for $\check{\kappa}$ of Step 1.

Step 3. Next, recall that

$$e_4(\zeta) + \kappa\zeta = -\beta - \vartheta\zeta.$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} \left((\mathfrak{d}^k e_4\zeta)^2 + (\mathfrak{d}^k \zeta)^2 \right) \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2.$$

Also, commuting first with \mathbf{N} , and proceeding analogously, we infer

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \mathbf{N}\zeta)^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2$$

Furthermore, in view of the definition of μ and a Poincaré inequality for \not{d}_1 , we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \not{d}\zeta)^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used a trace estimate and the estimate for $\check{\mu}$ of Step 2. The above estimates, together with (8.8.3), imply

$$\max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \zeta)^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2.$$

Step 4. Recall that

$$e_4(\vartheta) + \kappa\vartheta = -2\alpha.$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} \left((\mathfrak{d}^k e_4 \vartheta)^2 + (\mathfrak{d}^k \vartheta)^2 \right) \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2.$$

Also, commuting first one time with \mathbf{N} , and proceeding analogously, we infer

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \mathbf{N} \vartheta)^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2.$$

Furthermore, in view of Codazzi for ϑ , and a Poincaré inequality for \not{d}_2 , we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \not{d} \vartheta)^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used a trace estimate, and the estimate for $\check{\kappa}$ and ζ respectively in Step 1 and Step 3. The above estimates, together with (8.8.3), imply

$$\max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \vartheta)^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2.$$

Step 5. Recall that we have

$$\begin{aligned} e_4(\check{\kappa}) + \frac{1}{2} \bar{\kappa} \check{\kappa} &= -\frac{1}{2} \bar{\kappa} \check{\kappa} - 2 \not{d}_1 \zeta + 2\check{\rho} + \text{Err}[e_4 \check{\kappa}] \\ &= -\frac{1}{2} \bar{\kappa} \check{\kappa} + 2\check{\mu} + 4\check{\rho} - \frac{1}{2} \vartheta \not{d} \vartheta + \text{Err}[e_4 \check{\kappa}]. \end{aligned}$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} \left((\mathfrak{d}^k e_4 \check{\kappa})^2 + (\mathfrak{d}^k \check{\kappa})^2 \right) \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimates for $\check{\kappa}$ and $\check{\mu}$ derived respectively in Step 1 and Step 2. Also, commuting first one time with \mathbf{N} , and proceeding analogously, we infer

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \mathbf{N} \check{\kappa})^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimates for $\check{\kappa}$ and $\check{\mu}$ derived respectively in Step 1 and Step 2. Furthermore, commuting the equation for $e_4(\kappa)$ once with e_θ , we have

$$e_4(e_\theta(\underline{\kappa})) + \kappa e_\theta(\underline{\kappa}) = -\frac{1}{2} \underline{\kappa} e_\theta(\kappa) + 2e_\theta(\mu) + 4e_\theta(\rho) - e_\theta(\vartheta \not{d} \vartheta) + 2e_\theta(\zeta^2) - \frac{1}{2} \vartheta e_\theta(\underline{\kappa}).$$

Together with the Bianchi identity for $e_4(\underline{\beta})$, we infer

$$\begin{aligned} e_4(e_\theta(\underline{\kappa}) - 4\underline{\beta}) + \kappa(e_\theta(\underline{\kappa}) - 4\underline{\beta}) &= -\frac{1}{2}\underline{\kappa}e_\theta(\kappa) + 2e_\theta(\mu) + 12\rho\zeta \\ &\quad + 4\underline{\vartheta}\beta - e_\theta(\vartheta\underline{\vartheta}) + 2e_\theta(\zeta^2) - \frac{1}{2}\vartheta e_\theta(\underline{\kappa}). \end{aligned}$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} \left((\mathfrak{d}^k(e_4(e_\theta(\underline{\kappa}) - 4\underline{\beta}))^2 + (\mathfrak{d}^k(e_\theta(\underline{\kappa}) - 4\underline{\beta}))^2 \right) \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimates for $\check{\kappa}$, $\check{\mu}$ and ζ derived respectively in Step 1, Step 2 and Step 3.

The above estimates, together with (8.8.3), imply

$$\begin{aligned} \max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \check{\underline{\kappa}})^2 &\lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2 + \max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \underline{\beta})^2 \\ &\lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2 \end{aligned}$$

where we have used a trace estimate on $\{r = r_0\}$ for $r_{\mathcal{T}} \leq r_0 \leq 4m_0$.

Step 6. Recall that we have

$$e_4(\check{\underline{\omega}}) = \check{\rho} + \text{Err}[e_4\check{\underline{\omega}}].$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} \left((\mathfrak{d}^k e_4\check{\underline{\omega}})^2 + (\mathfrak{d}^k \check{\underline{\omega}})^2 \right) \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2.$$

Also, commuting first one time with \mathbf{N} , and proceeding analogously, we infer

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \mathbf{N}\check{\underline{\omega}})^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2.$$

Step 7. Recall that we have

$$\begin{aligned} &e_4(e_3(\zeta) + \underline{\beta}) + \kappa(e_3(\zeta) + \underline{\beta}) \\ &= \underline{\kappa}\beta + \left(\frac{1}{2}\kappa\underline{\kappa} + 2\mu - 6\rho \right) \zeta \\ &\quad - \underline{\vartheta}\beta + \vartheta\underline{\beta} - \underline{\xi}\alpha - \vartheta\zeta + 2\underline{\omega}\vartheta\zeta - 4\zeta \mathfrak{d}_1^* \mathfrak{d}_1^{-1} \left(\check{\mu} + \check{\rho} - \frac{1}{4}\vartheta\underline{\vartheta} + \frac{1}{4}\vartheta\overline{\vartheta} \right) - 2\zeta^3 \end{aligned}$$

Commuting first one time with \mathbf{N} , and in view of Corollary 8.8.10, we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \mathbf{N}(e_3(\zeta) + \underline{\beta}))^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimate for ζ in Step 3.

Step 8. Recall that we have

$$e_4(\underline{\xi}) = -e_3(\zeta) + \underline{\beta} - \underline{\kappa}\zeta - \zeta\underline{\vartheta}.$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} \left((\mathfrak{d}^k e_4 \underline{\xi})^2 + (\mathfrak{d}^k \underline{\xi})^2 \right) \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimates for ζ derived in Step 3. Also, commuting first one time with \mathbf{N} , and proceeding analogously, we infer

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \mathbf{N} \underline{\xi})^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimates for $e_3(\zeta) + \underline{\beta}$ derived in Step 7.

Step 9. Recall

$$2 \mathfrak{d}_1^* \underline{\omega} = e_3 \zeta + \underline{\kappa} \zeta - \underline{\beta} - \frac{1}{2} \underline{\kappa} \underline{\xi} + \underline{\vartheta} \zeta - \frac{1}{2} \underline{\vartheta} \underline{\xi}.$$

Using a Poincaré inequality for \mathfrak{d}_1^* , we infer

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \mathfrak{d} \underline{\omega})^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used a trace estimate and the estimate for ζ and $\underline{\xi}$ respectively in Step 3 and Step 8. The above estimates, together with the estimates for $\underline{\omega}$ of Step 6 and (8.8.3), imply

$$\max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \underline{\omega})^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}] \right)^2.$$

Step 10. Recall that we have

$$e_4(\check{\underline{\Omega}}) = -2\check{\underline{\omega}} + \overline{\check{\kappa}\check{\underline{\Omega}}}.$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \check{\underline{\Omega}})^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimates for $\check{\underline{\omega}}$ derived in Step 9.

Step 11. Recall

$$2 \mathfrak{d}_1 \underline{\xi} = e_3(\check{\underline{\kappa}}) + \bar{\kappa} \check{\underline{\kappa}} + 2\bar{\omega} \check{\underline{\kappa}} + 2\bar{\kappa} \check{\underline{\omega}} - \left(\frac{1}{2} \bar{\kappa} \bar{\kappa} - 2\bar{\rho} \right) \check{\underline{\Omega}} - \text{Err}[e_3 \check{\underline{\kappa}}].$$

Using a Poincaré inequality for \mathfrak{d}_1 , we infer

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \mathfrak{d} \underline{\xi})^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimates for $\check{\underline{\kappa}}$, $\check{\underline{\omega}}$ and $\check{\underline{\Omega}}$ respectively in Step 5, Step 9 and Step 10. The above estimates, together with the estimates for $\underline{\xi}$ of Step 8 and (8.8.3), imply

$$\max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \underline{\xi})^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2.$$

Step 12. Recall that

$$\begin{aligned} e_4(\underline{\vartheta}) + \frac{1}{2} \kappa \underline{\vartheta} &= 2 \mathfrak{d}_2^* \zeta - \frac{1}{2} \kappa \underline{\vartheta} + 2\zeta^2 \\ &= 2 \mathfrak{d}_2^* \mathfrak{d}_1^{-1} \left(-\check{\mu} - \check{\rho} + \frac{1}{4} \underline{\vartheta} \underline{\vartheta} - \frac{1}{4} \overline{\underline{\vartheta}} \overline{\underline{\vartheta}} \right) - \frac{1}{2} \kappa \underline{\vartheta} + 2\zeta^2. \end{aligned}$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} \left((\mathfrak{d}^k e_4 \underline{\vartheta})^2 + (\mathfrak{d}^k \underline{\vartheta})^2 \right) \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimate for $\check{\mu}$ and $\underline{\vartheta}$ respectively in Step 2 and Step 4. Also, commuting first one time with \mathbf{N} , and proceeding analogously, we infer

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \mathbf{N} \underline{\vartheta})^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used the estimate for $\check{\mu}$ and $\underline{\vartheta}$ respectively in Step 2 and Step 4. Furthermore, in view of Codazzi for $\underline{\vartheta}$, and a Poincaré inequality for \mathfrak{d}_2 , we have

$$\max_{k \leq J} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k \mathfrak{d} \underline{\vartheta})^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used a trace estimate and the estimate for $\check{\kappa}$ and ζ respectively in Step 5 and Step 3. The above estimates, together with (8.8.3), imply

$$\max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{D}^k \underline{\vartheta})^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2.$$

Step 13. Recall that we have

$$\begin{aligned} & e_4 \left(\mathfrak{d}_1 \mathfrak{d}_1^* \kappa - \vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} \right) + 2\kappa \left(\mathfrak{d}_1 \mathfrak{d}_1^* \kappa - \vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} \right) \\ = & - \left(-\frac{1}{2} \vartheta \mathfrak{d}_2^* + \zeta e_4(\Phi) - \beta \right) \mathfrak{d}_1^* \kappa - \frac{1}{2} \vartheta \mathfrak{d}_1 \mathfrak{d}_1^* \kappa + \frac{1}{2} \mathfrak{d}_1^* (\kappa + \vartheta) \mathfrak{d}_1^* \kappa + (\mathfrak{d}_1^* \kappa)^2 \\ & - 2\kappa \vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} + (\kappa \vartheta + 2\alpha) \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} \\ & + \vartheta \left[\left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1}, e_4 \right] \check{\rho} + \vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \left(\frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\ & - \frac{1}{2} \vartheta \mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} (-\mathfrak{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) - \frac{1}{2} \vartheta \mathfrak{d}_2^* (-\mathfrak{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \\ & + \frac{1}{2} \mathfrak{d}_3^* \mathfrak{d}_2^{-1} (-2\beta - \mathfrak{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \mathfrak{d}_3^* \vartheta + \frac{1}{2} (-2\beta - \mathfrak{d}_1^* \kappa + \kappa \zeta - \vartheta \zeta) \mathfrak{d}_2 \vartheta. \end{aligned}$$

In view of Corollary 8.8.10, we have

$$\max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} \left(\mathfrak{D}^k \left(\mathfrak{d}_1 \mathfrak{d}_1^* \kappa - \vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} \right) \right)^2 \lesssim \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2$$

where we have used

- the fact that $\mathfrak{D}\vartheta = \mathfrak{D}\mathfrak{d}_2^{-1} \mathfrak{d}_2 \vartheta$ and Codazzi for ϑ to estimate the terms of the RHS with one angular derivative of ϑ ,
- the estimates of Step 1 to estimate the terms of the RHS with one derivative of $\check{\kappa}$,
- the fact that $\mathfrak{D}\zeta = \mathfrak{D}\mathfrak{d}_1^{-1} \mathfrak{d}_1 \zeta$ and the definition of μ to estimate terms of the RHS with one angular derivative of ζ ,
- the identity

$$\begin{aligned} \mathfrak{D} \mathfrak{d}_1^* \kappa &= \mathfrak{D} \mathfrak{d}_1^{-1} \left(\vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} \right) \\ &\quad + \mathfrak{D} \mathfrak{d}_1^{-1} \left(\mathfrak{d}_1 \mathfrak{d}_1^* \kappa - \vartheta \left(\mathfrak{d}_4 \mathfrak{d}_3^* \mathfrak{d}_2^{-1} + \mathfrak{d}_2^* \right) \mathfrak{d}_1^{-1} \check{\rho} \right) \end{aligned}$$

to estimate the terms of the RHS with two angular derivatives of $\check{\kappa}$.

Step 14. Recall that we have

$$\begin{aligned}
& e_4 \left(e_\theta(\mu) + \vartheta \not\!d_2 \not\!d_2^* (\not\!d_1^* \not\!d_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) + 2\kappa \left(e_\theta(\mu) + \vartheta \not\!d_2 \not\!d_2^* (\not\!d_1^* \not\!d_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) \\
= & -\frac{3}{2} \mu e_\theta(\kappa) - \frac{1}{2} \vartheta e_\theta(\mu) - (\kappa \vartheta + 2\alpha) \not\!d_2 \not\!d_2^* (\not\!d_1^* \not\!d_1)^{-1} \underline{\beta} \\
& - \vartheta \left[\not\!d_2 \not\!d_2^* (\not\!d_1^* \not\!d_1)^{-1}, e_4 \right] \underline{\beta} - \vartheta \not\!d_2 \not\!d_2^* (\not\!d_1^* \not\!d_1)^{-1} (\kappa \underline{\beta} + 3\rho \zeta + \underline{\vartheta} \beta) \\
& + \not\!d_3^* \vartheta \not\!d_2^* \not\!d_1^{-1} \check{\rho} - e_\theta \left(\vartheta \not\!d_2^* \not\!d_1^{-1} \left(-\check{\mu} + \frac{1}{4} \vartheta \underline{\vartheta} \right) \right) - e_\theta \left(\vartheta \left(\frac{1}{8} \underline{\kappa} \vartheta + \zeta^2 \right) \right) \\
& - 2\zeta \not\!d_1 \not\!d_1^* \kappa - 2e_\theta(\kappa) \not\!d_2^* \zeta - 2(\kappa \zeta + \beta + \vartheta \zeta) \check{\rho} - 2\zeta \left(\frac{3}{2} \bar{\kappa} \check{\rho} + \frac{3}{2} \bar{\rho} \check{\kappa} - \text{Err}[e_4 \check{\rho}] \right) \\
& + 2\beta \not\!d_2^* \zeta + \frac{3}{2} e_\theta(\kappa \zeta^2) + 2\kappa \left(\vartheta \not\!d_2 \not\!d_2^* (\not\!d_1^* \not\!d_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right).
\end{aligned}$$

In view of Corollary 8.8.10, we have

$$\begin{aligned}
& \max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} \left(\not\!d^k \left(e_\theta(\mu) + \vartheta \not\!d_2 \not\!d_2^* (\not\!d_1^* \not\!d_1)^{-1} \underline{\beta} + 2\zeta \check{\rho} \right) \right)^2 \\
\lesssim & \left(N^{\leq 4m_0} [J, \check{\Gamma}, \check{R}] \right)^2 + \epsilon^2 \int_{(ext) \mathcal{M}(\leq 4m_0)} \left(\not\!d^{\leq J+1} \left(e_\theta(\underline{\kappa}) - 4\underline{\beta} \right) \right)^2,
\end{aligned}$$

where we have used

- the fact that $\not\!d \vartheta = \not\!d \not\!d_2^{-1} \not\!d_2 \vartheta$ and Codazzi for ϑ to estimate the terms of the RHS with one angular derivative of ϑ ,
- the estimates of Step 1 to estimate the terms of the RHS with one derivative of $\check{\kappa}$,
- the fact that $\not\!d \zeta = \not\!d \not\!d_1^{-1} \not\!d_1 \zeta$ and the definition of μ to estimate terms of the RHS with one angular derivative of ζ ,
- the fact that $e_\theta(\underline{\kappa}) = (e_\theta(\underline{\kappa}) - 4\underline{\beta}) + 4\underline{\beta}$ to estimate the term with one angular derivative of $\underline{\kappa}$,
- the identity

$$\begin{aligned}
\not\!d \not\!d_1^* \kappa &= \not\!d \not\!d_1^{-1} \left(\vartheta \left(\not\!d_4 \not\!d_3^* \not\!d_2^{-1} + \not\!d_2^* \right) \not\!d_1^{-1} \check{\rho} \right) \\
&+ \not\!d \not\!d_1^{-1} \left(\not\!d_1 \not\!d_1^* \kappa - \vartheta \left(\not\!d_4 \not\!d_3^* \not\!d_2^{-1} + \not\!d_2^* \right) \not\!d_1^{-1} \check{\rho} \right)
\end{aligned}$$

and the estimates of Step13 to estimate the terms of the RHS with two angular derivatives of $\check{\kappa}$.

Step 15. Recall that we have

$$\begin{aligned}
& e_4(e_\theta(\underline{\kappa}) - 4\underline{\beta}) + \kappa(e_\theta(\underline{\kappa}) - 4\underline{\beta}) \\
&= 2e_\theta(\mu) + 12\rho\zeta - \frac{1}{2}\underline{\kappa}e_\theta(\kappa) + 4\underline{\vartheta}\beta - \frac{1}{2}\vartheta e_\theta(\underline{\kappa}) - e_\theta(\vartheta\underline{\vartheta}) + 2e_\theta(\zeta^2) \\
&= 2\left(e_\theta(\mu) + \vartheta \phi_2 \phi_2^*(\phi_1^* \phi_1)^{-1}\underline{\beta} + 2\zeta\check{\rho}\right) + 12\rho\zeta - \frac{1}{2}\underline{\kappa}e_\theta(\kappa) \\
&\quad - 2\vartheta \phi_2 \phi_2^*(\phi_1^* \phi_1)^{-1}\underline{\beta} + 2\zeta\check{\rho} + 4\underline{\vartheta}\beta - \frac{1}{2}\vartheta(e_\theta(\underline{\kappa}) - 4\underline{\beta}) - 2\vartheta\underline{\beta} - e_\theta(\vartheta\underline{\vartheta}) + 2e_\theta(\zeta^2).
\end{aligned}$$

In view of Corollary 8.8.10, we have

$$\begin{aligned}
& \max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k (e_\theta(\underline{\kappa}) - 4\underline{\beta}))^2 \\
& \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}]\right)^2 + \int_{(ext)\mathcal{M}(\leq 4m_0)} (\mathfrak{d}^{\leq J+1} (e_\theta(\mu) + \vartheta \phi_2 \phi_2^*(\phi_1^* \phi_1)^{-1}\underline{\beta} + 2\zeta\check{\rho}))^2.
\end{aligned}$$

where we have used

- the fact that $\check{\vartheta}\vartheta = \check{\vartheta}\phi_2^{-1}\phi_2\vartheta$ and Codazzi for ϑ to estimate the terms of the RHS with one angular derivative of ϑ ,
- the fact that $\check{\vartheta}\underline{\vartheta} = \check{\vartheta}\phi_2^{-1}\phi_2\underline{\vartheta}$ and Codazzi for $\underline{\vartheta}$ to estimate the terms of the RHS with one angular derivative of $\underline{\vartheta}$,
- the estimates of Step 1 to estimate the terms of the RHS with one derivative of $\check{\kappa}$,
- the fact that $\check{\vartheta}\zeta = \check{\vartheta}\phi_1^{-1}\phi_1\zeta$ and the definition of μ to estimate terms of the RHS with one angular derivative of ζ ,
- the estimate for ζ of Step 3.

Together with the estimate of Step 14, we infer

$$\begin{aligned}
& \max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k (e_\theta(\mu) + \vartheta \phi_2 \phi_2^*(\phi_1^* \phi_1)^{-1}\underline{\beta} + 2\zeta\check{\rho}))^2 \\
& \quad + \max_{k \leq J+1} \sup_{r_{\mathcal{T}} \leq r_0 \leq 4m_0} \int_{\{r=r_0\}} (\mathfrak{d}^k (e_\theta(\underline{\kappa}) - 4\underline{\beta}))^2 \lesssim \left(N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}]\right)^2.
\end{aligned}$$

In view of Step 1 to Step 15, of the definition (8.8.4) of $N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}]$, and of the various norms, we infer

$$\begin{aligned}
({}^{ext})\mathfrak{G}_{J+1}^{\leq 4m_0}[\check{\Gamma}] + ({}^{ext})\mathfrak{G}_{J+1}^{\leq 4m_0'}[\check{\Gamma}] & \lesssim ({}^{ext})\mathfrak{G}_{J+1}^{\geq 4m_0}[\check{\Gamma}] + ({}^{ext})\mathfrak{G}_{J+1}^{\geq 4m_0'}[\check{\Gamma}] + ({}^{ext})\mathfrak{R}_{J+1}[\check{R}] + ({}^{ext})\mathfrak{G}_J[\check{\Gamma}] \\
& \quad + \epsilon_0 \left(({}^{ext})\mathfrak{G}_{J+1}^{\leq 4m_0}[\check{\Gamma}] + ({}^{ext})\mathfrak{G}_{J+1}^{\leq 4m_0'}[\check{\Gamma}] \right).
\end{aligned}$$

and hence, for ϵ_0 small enough,

$${}^{(ext)}\mathfrak{G}_{J+1}^{\leq 4m_0}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_{J+1}^{\leq 4m_0'}[\check{\Gamma}] \lesssim {}^{(ext)}\mathfrak{G}_{J+1}^{\geq 4m_0}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}_{J+1}^{\geq 4m_0'}[\check{\Gamma}] + {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{G}_J[\check{\Gamma}].$$

This concludes the proof of Proposition 8.8.3.

8.9 Proof of Proposition 8.3.10

To prove Proposition 8.3.10, we rely on the following proposition.

Proposition 8.9.1. *Let J such that $k_{small} - 2 \leq J \leq k_{large} - 1$. Then, we have*

$$\begin{aligned} {}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{G}'_{J+1}[\check{\Gamma}] &\lesssim {}^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}'_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{R}_{J+1}[\check{R}] \\ &\quad + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}, \end{aligned}$$

where the notation ${}^{(ext)}\mathfrak{G}'_{J+1}[\check{\Gamma}]$ has been introduced in Proposition 8.8.2, and where we have introduced the notation

$${}^{(int)}\mathfrak{G}'_k[\check{\Gamma}] := \int_{(int)\mathcal{M}} \left[(\mathfrak{d}^k e_\theta(\underline{\kappa}))^2 + (\mathfrak{d}^{\leq k} \underline{\mu})^2 + (\mathfrak{d}^k (e_4(\zeta) - \beta))^2 \right].$$

The proof of Proposition 8.9.1 is postponed to section 8.9.2. It will rely in particular on basic weighted estimates for transport equations along e_3 in ${}^{(int)}\mathcal{M}$ derived in section 8.9.1.

We now conclude the proof of Proposition 8.3.10. In view of Proposition 8.9.1, we have

$$\begin{aligned} {}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] &\lesssim {}^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}'_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{R}_{J+1}[\check{R}] \\ &\quad + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Also, we have in view of Proposition 8.8.1, Proposition 8.8.2 and the iteration assumption (8.3.10)

$${}^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}'_{J+1}[\check{\Gamma}] \lesssim {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}] + \epsilon_{\mathcal{B}}[J].$$

We infer

$${}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] \lesssim {}^{(int)}\mathfrak{R}_{J+1}[\check{R}] + {}^{(ext)}\mathfrak{R}_{J+1}[\check{R}] + \epsilon_{\mathcal{B}}[J] + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}.$$

Together with Proposition 8.3.9, we deduce

$${}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] \lesssim \epsilon_{\mathcal{B}}[J] + \epsilon_0 \left(\mathfrak{N}_{J+1}^{(En)} + \mathcal{N}_{J+1}^{(match)} \right) + \left(\int_{\mathcal{T}} |\mathfrak{D}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}$$

which concludes the proof of Proposition 8.3.10.

The rest of this section is dedicated to the proof of Proposition 8.9.1.

8.9.1 Weighted estimates for transport equations along e_3 in ${}^{(int)}\mathcal{M}$

Lemma 8.9.2. *Let the following transport equation in ${}^{(int)}\mathcal{M}$*

$$e_3(f) + \frac{a}{2}\kappa f = h$$

where $a \in \mathbb{R}$ is a given constant, and f and h are scalar functions. Then, f satisfies

$$\int_{{}^{(int)}\mathcal{M}} f^2 \lesssim \int_{\mathcal{T}} f^2 + \int_{{}^{(int)}\mathcal{M}} h^2.$$

Proof. Multiply by f to obtain

$$\frac{1}{2}e_3(f^2) + \frac{a}{r}f^2 = hf.$$

Next, integrate over $S_{\underline{u},r}$ to obtain

$$\begin{aligned} \frac{1}{2}e_3 \left(\int_{S_{\underline{u},r}} f^2 \right) &= \int_{S_{\underline{u},r}} \frac{1}{2}(e_3(f^2) + \underline{\kappa}f^2) \\ &= - \int_{S_{\underline{u},r}} \frac{a-1}{2}\underline{\kappa}f^2 + \int_{S_{\underline{u},r}} hf \\ &= -\frac{a-1}{2}\underline{\bar{\kappa}} \int_{S_{\underline{u},r}} f^2 - \frac{a-1}{2} \int_{S_{\underline{u},r}} \check{\underline{\kappa}}f^2 + \int_{S_{\underline{u},r}} hf \end{aligned}$$

and hence

$$\frac{1}{2}e_3 \left(r^b \int_{S_{\underline{u},r}} f^2 \right) + \frac{1}{2} \left(a+1 + \frac{b}{2} \right) \underline{\bar{\kappa}} r^b \int_{S_{\underline{u},r}} f^2 = -\frac{a-1}{2} r^b \int_{S_{\underline{u},r}} \check{\underline{\kappa}}f^2 + r^b \int_{S_{\underline{u},r}} hf$$

where we used the fact that $2e_3(r) = r\bar{\kappa}$. Also, choosing $b = -2a$, we obtain

$$\frac{1}{2}e_3 \left(r^{-2a} \int_{S_{\underline{u},r}} f^2 \right) + \frac{1}{2\bar{\kappa}} r^{-2a} \int_{S_{\underline{u},r}} f^2 = -\frac{a-1}{2} r^{-2a} \int_{S_{\underline{u},r}} \check{\kappa} f^2 + r^{-2a} \int_{S_{\underline{u},r}} h f.$$

Next, let $1 \leq \underline{u} \leq u_*$. We now integrate in r and along $\mathcal{C}_{\underline{u}}$ in ${}^{(int)}\mathcal{M}$. Since r is bounded on ${}^{(int)}\mathcal{M}$ from above and below, we obtain, for $\epsilon_0 > 0$ small enough,

$$\int_{2m_0-2m_0\delta_0}^{r_{\mathcal{T}}} \int_{S_{\underline{u},r}} f^2 \lesssim \int_{S_{\underline{u},r_{\mathcal{T}}}} f^2 + \int_{2m_0-2m_0\delta_0}^{r_{\mathcal{T}}} \int_{S_{\underline{u},r}} h^2.$$

We may now integrate in \underline{u} to deduce

$$\int_1^{u_*} \int_{2m_0-2m_0\delta_0}^{r_{\mathcal{T}}} \int_{S_{\underline{u},r}} f^2 \lesssim \int_1^{u_*} \int_{S_{\underline{u},r_{\mathcal{T}}}} f^2 + \int_1^{u_*} \int_{2m_0-2m_0\delta_0}^{r_{\mathcal{T}}} \int_{S_{\underline{u},r}} h^2. \quad (8.9.1)$$

Remark 8.9.3. *Note that we have the following consequence of the coarea formula*

$$d\mathcal{T} = \frac{\varsigma\sqrt{\bar{\kappa}} + A}{\sqrt{-\bar{\kappa}}} d\mu_{\underline{u},r_{\mathcal{T}}} d\underline{u},$$

where we used that $\mathcal{T} = \{r = r_{\mathcal{T}}\}$. Also, we have in ${}^{(int)}\mathcal{M}$

$$d\mathcal{M} = \frac{4\varsigma^2}{r^2\bar{\kappa}^2} d\mu_{\underline{u},r} d\underline{u} dr.$$

We infer, in ${}^{(int)}\mathcal{M}$,

$$d\mathcal{T} = \sqrt{1 - \frac{2m_0}{r_{\mathcal{T}}}} (1 + O(\epsilon_0)) d\mu_{\underline{u},r_0} d\underline{u},$$

and

$$d\mathcal{M} = (1 + O(\epsilon_0)) d\mu_{\underline{u},r} d\underline{u} dr.$$

Relying on Remark 8.9.3 we deduce from (8.9.1)

$$\int_{{}^{(int)}\mathcal{M}} f^2 \lesssim \int_{\mathcal{T}} f^2 + \int_{{}^{(int)}\mathcal{M}} h^2$$

as desired. This concludes the proof of the lemma. \square

Corollary 8.9.4. *Let the following transport equation in ${}^{(int)}\mathcal{M}$*

$$e_3(f) + \frac{a}{2}\underline{\kappa}f = h$$

where $a \in \mathbb{R}$ is a given constant, and f and h are scalar functions. Then, f satisfies for $5 \leq l \leq k_{large} + 1$

$$\begin{aligned} \int_{{}^{(int)}\mathcal{M}} (\mathfrak{d}^k f)^2 &\lesssim \int_{\mathcal{T}} (\mathfrak{d}^{\leq k} f)^2 + \int_{{}^{(int)}\mathcal{M}} (\mathfrak{d}^{\leq k-1} f)^2 \\ &\quad + \int_{{}^{(int)}\mathcal{M}} (\mathfrak{d}^{\leq k} h)^2 + \left(\sup_{{}^{(int)}\mathcal{M}} |\mathfrak{d}^{\leq k-5} f| \right)^2 \left({}^{(int)}\mathfrak{G}_{k-1}[\check{\Gamma}] + {}^{(int)}\mathfrak{G}_k[\check{\underline{\kappa}}] \right)^2. \end{aligned}$$

Proof. The proof is based on Lemma 8.9.2. It is similar to the one of Corollary 8.8.6 and left to the reader. \square

8.9.2 Proof of Proposition 8.9.1

We introduce the following notation which will constantly appear on the RHS of the equalities below

$$\begin{aligned} N^{(int)}[J, \check{\Gamma}, \check{R}] &:= {}^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}'_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{R}_{J+1}[\check{R}] \\ &\quad + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}} + \epsilon_0 \left({}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{G}'_{J+1}[\check{\Gamma}] \right) \end{aligned} \quad (8.9.2)$$

Step 1. In view of Lemma 7.7.1 relating the Ricci coefficients and curvature components of ${}^{(int)}\mathcal{M}$ to the ones of ${}^{(ext)}\mathcal{M}$ on the timelike hypersurface \mathcal{T} , we have

$$\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(int)}\check{\Gamma})|^2 \lesssim \int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{\Gamma})|^2.$$

Also, using again Lemma 7.7.1, we have

$$\begin{aligned} &\int_{\mathcal{T}} \left| \mathfrak{d}^{J+1} \left({}^{(int)}e_{\theta}({}^{(int)}\underline{\kappa}), {}^{(int)}\check{\underline{\mu}}, {}^{(int)}e_4({}^{(int)}\zeta - {}^{(int)}\beta) \right) \right|^2 \\ &\lesssim \int_{\mathcal{T}} \left| \mathfrak{d}^{J+1} \left({}^{(ext)}e_{\theta}({}^{(int)}\underline{\kappa}) - 4{}^{(ext)}\underline{\beta}, {}^{(int)}\check{\underline{\mu}}, {}^{(int)}e_3({}^{(int)}\zeta) + {}^{(int)}\underline{\beta} \right) \right|^2 + \int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \end{aligned}$$

We deduce, using that $\mathcal{T} = \{r = r_{\mathcal{T}}\}$ and the definitions of the various norms on ${}^{(ext)}\mathcal{M}$,

$$\begin{aligned} &\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(int)}\check{\Gamma})|^2 + \int_{\mathcal{T}} \left| \mathfrak{d}^{J+1} \left({}^{(int)}e_{\theta}({}^{(int)}\underline{\kappa}), {}^{(int)}\check{\underline{\mu}}, {}^{(int)}e_4({}^{(int)}\zeta - {}^{(int)}\beta) \right) \right|^2 \\ &\lesssim {}^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}'_{J+1}[\check{\Gamma}] + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

and hence, in view of (8.9.2),

$$\begin{aligned} & \int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(int)}\check{\Gamma})|^2 + \int_{\mathcal{T}} \left| \mathfrak{d}^{J+1} \left({}^{(int)}e_{\theta}({}^{(int)}\check{\underline{\kappa}}), {}^{(int)}\check{\underline{\mu}}, {}^{(int)}e_4({}^{(int)}\zeta - {}^{(int)}\beta) \right) \right|^2 \\ & \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2. \end{aligned}$$

From now on, we only consider the frame of ${}^{(int)}\mathcal{M}$. The previous estimate can be written as

$$\begin{aligned} \max_{k \leq J+1} \left(\int_{\mathcal{T}} \left((\mathfrak{d}^k \check{\underline{\mu}})^2 + (\mathfrak{d}^k \zeta)^2 + (\mathfrak{d}^k \check{\kappa})^2 + (\mathfrak{d}^k \vartheta)^2 + (\mathfrak{d}^k \check{\underline{\kappa}})^2 + (\mathfrak{d}^k \underline{\vartheta})^2 \right. \right. \\ \left. \left. + (\mathfrak{d}^k (e_4(\zeta) - \beta))^2 + (\mathfrak{d}^k \xi)^2 + (\mathfrak{d}^k \check{\omega})^2 + (\mathfrak{d}^k \check{\Omega})^2 \right) \right) \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2 \end{aligned}$$

and

$$\max_{k \leq J+1} \int_{\mathcal{T}} (\mathfrak{d}^k e_{\theta}(\underline{\kappa}))^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

Step 2. We have obtained all the desired estimates on \mathcal{T} for the foliation of ${}^{(int)}\mathcal{M}$ in Step 1. We now derive the desired estimates on ${}^{(int)}\mathcal{M}$. To this end, we rely on the transport equations in the e_3 directions which we estimate thanks to Corollary 8.9.4. The initial data on \mathcal{T} is estimated thanks to Step 1. In particular, we proceed in the following order

- From

$$e_3(\check{\underline{\kappa}}) + \bar{\underline{\kappa}} \check{\underline{\kappa}} = \text{Err}[e_3 \check{\underline{\kappa}}]$$

and the bootstrap assumptions, we infer

$$\max_{k \leq J+1} \int_{{}^{(int)}\mathcal{M}} (\mathfrak{d}^k \check{\underline{\kappa}})^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

- From

$$e_3(e_{\theta}(\underline{\kappa})) + \frac{3}{2} \underline{\kappa} e_{\theta}(\underline{\kappa}) = -\frac{1}{2} \underline{\vartheta} e_{\theta}(\underline{\kappa}) - \frac{1}{2} e_{\theta}(\underline{\vartheta}^2)$$

and the bootstrap assumptions, we infer

$$\max_{k \leq J+1} \int_{{}^{(int)}\mathcal{M}} (\mathfrak{d}^k e_{\theta}(\underline{\kappa}))^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

- From

$$e_3(\underline{\check{\mu}}) + \frac{3}{2}\overline{\kappa}\underline{\check{\mu}} = -\frac{3}{2}\underline{\mu}\underline{\check{\kappa}} + \text{Err}[e_3\underline{\check{\mu}}],$$

the above control of $\underline{\check{\kappa}}$ and $e_\theta(\underline{\kappa})$ (the control of $e_\theta(\underline{\kappa})$ is needed to estimate $\text{Err}[e_3\underline{\check{\mu}}]$), and the bootstrap assumptions, we infer

$$\max_{k \leq J+1} \int_{(int)\mathcal{M}} (\mathfrak{d}^k \underline{\check{\mu}})^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

- From

$$e_3(\underline{\vartheta}) + \underline{\kappa}\underline{\vartheta} = -2\underline{\alpha}$$

and the control of $\underline{\alpha}$, we infer

$$\max_{k \leq J+1} \int_{(int)\mathcal{M}} (\mathfrak{d}^k \underline{\vartheta})^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

- From

$$e_3(\underline{\zeta}) + \underline{\kappa}\underline{\zeta} = \underline{\beta} - \underline{\vartheta}\underline{\zeta}$$

the control of $\underline{\beta}$, and the bootstrap assumptions, we infer

$$\max_{k \leq J+1} \int_{(int)\mathcal{M}} (\mathfrak{d}^k \underline{\zeta})^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

- From

$$\begin{aligned} e_3(\underline{\check{\kappa}}) + \frac{1}{2}\overline{\kappa}\underline{\check{\kappa}} &= -\frac{1}{2}\overline{\kappa}\underline{\check{\kappa}} + 2\mathfrak{d}_1\underline{\zeta} + 2\check{\rho} + \text{Err}[e_3\underline{\check{\kappa}}] \\ &= -\frac{1}{2}\overline{\kappa}\underline{\check{\kappa}} + 2\underline{\check{\mu}} + 4\check{\rho} - \frac{1}{2}\underline{\vartheta}\underline{\vartheta} + \frac{1}{2}\overline{\vartheta}\underline{\vartheta} + \text{Err}[e_3\underline{\check{\kappa}}], \end{aligned}$$

the control of $\check{\rho}$, the above control of $\underline{\check{\kappa}}$ and $\underline{\check{\mu}}$, and the bootstrap assumptions, we infer

$$\max_{k \leq J+1} \int_{(int)\mathcal{M}} (\mathfrak{d}^k \underline{\check{\kappa}})^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

- From

$$\begin{aligned} e_3(\underline{\vartheta}) + \frac{1}{2}\underline{\kappa}\underline{\vartheta} &= 2\mathfrak{d}_2^*\underline{\zeta} - \frac{1}{2}\underline{\kappa}\underline{\vartheta} + 2\underline{\zeta}^2 \\ &= 2\mathfrak{d}_2^*\mathfrak{d}_1^{-1} \left(\underline{\check{\mu}} + \check{\rho} - \frac{1}{4}\underline{\vartheta}\underline{\vartheta} + \frac{1}{4}\overline{\vartheta}\underline{\vartheta} \right) - \frac{1}{2}\underline{\kappa}\underline{\vartheta} + 2\underline{\zeta}^2, \end{aligned}$$

the control of $\check{\rho}$, the above control of $\underline{\vartheta}$ and $\check{\mu}$, and the bootstrap assumptions, we infer

$$\max_{k \leq J+1} \int_{(int)\mathcal{M}} (\mathfrak{d}^k \vartheta)^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

- From

$$e_3(\check{\omega}) = \check{\rho} + \text{Err}[e_3 \check{\omega}],$$

the control of $\check{\rho}$, and the bootstrap assumptions, we infer

$$\max_{k \leq J+1} \int_{(int)\mathcal{M}} (\mathfrak{d}^k \check{\omega})^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

- From

$$\begin{aligned} & e_3(e_4(\zeta) - \beta) + \underline{\kappa}(e_4(\zeta) - \beta) \\ = & -\underline{\kappa}\underline{\beta} + \left(\frac{1}{2}\underline{\kappa}\underline{\kappa} + 2\underline{\mu} - 6\rho \right) \zeta \\ & + \underline{\vartheta}\underline{\beta} - \underline{\vartheta}\beta + \underline{\xi}\underline{\alpha} - \underline{\vartheta}\zeta + 2\omega\underline{\vartheta}\zeta - 4\zeta \underline{\vartheta}_1^* \underline{\vartheta}_1^{-1} \left(\check{\mu} + \check{\rho} - \frac{1}{4}\underline{\vartheta}\underline{\vartheta} + \frac{1}{4}\overline{\underline{\vartheta}\underline{\vartheta}} \right) - 2\zeta^3, \end{aligned}$$

the control of $\underline{\beta}$, the above control of ζ , and the bootstrap assumptions, we infer

$$\max_{k \leq J+1} \int_{(int)\mathcal{M}} (\mathfrak{d}^k (e_4(\zeta) - \beta))^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

- From

$$e_3(\xi) = (e_4(\zeta) - \beta) + 2\beta + \underline{\kappa}\zeta + \underline{\vartheta}\zeta,$$

the control of $\underline{\beta}$, the above control of $e_4(\zeta) - \beta$ and ζ , and the bootstrap assumptions, we infer

$$\max_{k \leq J+1} \int_{(int)\mathcal{M}} (\mathfrak{d}^k \xi)^2 \lesssim \left(N^{(int)}[J, \check{\Gamma}, \check{R}] \right)^2.$$

In view of the above estimates, of the definition (8.9.2) of $N^{\leq 4m_0}[J, \check{\Gamma}, \check{R}]$, and of the various norms, we infer

$$\begin{aligned} {}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{G}'_{J+1}[\check{\Gamma}] & \lesssim {}^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}'_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{R}_{J+1}[\check{R}] \\ & + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}} + \epsilon_0 \left({}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{G}'_{J+1}[\check{\Gamma}] \right) \end{aligned}$$

and hence, for ϵ_0 small enough,

$$\begin{aligned} {}^{(int)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{G}'_{J+1}[\check{\Gamma}] &\lesssim {}^{(ext)}\mathfrak{G}_{J+1}[\check{\Gamma}] + {}^{(ext)}\mathfrak{G}'_{J+1}[\check{\Gamma}] + {}^{(int)}\mathfrak{R}_{J+1}[\check{R}] \\ &\quad + \left(\int_{\mathcal{T}} |\mathfrak{d}^{J+1}({}^{(ext)}\check{R})|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

This concludes the proof of Proposition 8.9.1.

8.10 Proof of Proposition 8.3.11

Lemma 4.6.6 corresponds to the particular case $J = k_{large} - 1$ of Proposition 8.3.11. Its proof in section 4.6.2 extends immediately to the case $k_{small} - 2 \leq J \leq k_{large} - 1$ which thus yields the proof of Proposition 8.3.11.

Chapter 9

GCM PROCEDURE

9.1 Preliminaries

We consider an axially symmetric polarized spacetime regions \mathcal{R} foliated by two functions (u, s) such that

- On \mathcal{R} , (u, s) defines an outgoing geodesic foliation as in section 2.2.4.
- We denote by (e_3, e_4, e_θ) the null frame adapted to the outgoing geodesic foliation (u, s) on \mathcal{R} .
- Let

$$\overset{\circ}{S} := S(\overset{\circ}{u}, \overset{\circ}{s}) \quad (9.1.1)$$

and $\overset{\circ}{r}$ the area radius of $\overset{\circ}{S}$, where $S(u, s)$ denote the 2-spheres of the outgoing geodesic foliation (u, s) on \mathcal{R} .

- In adapted coordinates (u, s, θ, φ) with $b = 0$, see Proposition 2.2.20, the spacetime metric \mathbf{g} in \mathcal{R} takes the form, with $\underline{\Omega} = e_3(s)$, $\underline{b} = e_3(\theta)$,

$$\mathbf{g} = -2\varsigma duds + \varsigma^2 \underline{\Omega} du^2 + \gamma \left(d\theta - \frac{1}{2} \varsigma \underline{b} du \right)^2 + e^{2\Phi} d\varphi^2, \quad (9.1.2)$$

where θ is chosen such that $b = e_4(\theta) = 0$.

- The spacetime metric induced on $S(u, s)$ is given by,

$$\mathring{g} = \gamma d\theta^2 + e^{2\Phi} d\varphi^2. \quad (9.1.3)$$

- The relation between the null frame and coordinate system is given by

$$e_4 = \partial_s, \quad e_3 = \frac{2}{\zeta} \partial_u + \underline{\Omega} \partial_s + \underline{b} \partial_\theta, \quad e_\theta = \gamma^{-1/2} \partial_\theta. \quad (9.1.4)$$

- We denote the induced metric on \mathring{S} by

$$\mathring{g} = \mathring{\gamma} d\theta^2 + e^{2\mathring{\Phi}} d\varphi^2.$$

Definition 9.1.1. Let $0 < \delta \leq \mathring{\epsilon}$ two sufficiently small constants. Let $(\mathring{u}, \mathring{s})$ real numbers so that

$$1 \leq \mathring{u} < +\infty, \quad 4m_0 \leq \mathring{s} < +\infty. \quad (9.1.5)$$

We define $\mathcal{R} = \mathcal{R}(\mathring{\delta}, \mathring{\epsilon})$ to be the region

$$\mathcal{R} := \left\{ |u - \mathring{u}| \leq \delta_{\mathcal{R}}, \quad |s - \mathring{s}| \leq \delta_{\mathcal{R}} \right\}, \quad \delta_{\mathcal{R}} := \mathring{\delta}(\mathring{\epsilon})^{-\frac{1}{2}}, \quad (9.1.6)$$

such that assumption **A1-A3** below with constant $\mathring{\epsilon}$ on the background foliation of \mathcal{R} , are verified. The smaller constant $\mathring{\delta}$ controls the size of the GCMS quantities as it will be made precise below.

In this section we define the renormalized Ricci and curvature components,

$$\begin{aligned} \check{\Gamma} &:= \left\{ \check{\kappa}, \vartheta, \zeta, \eta, \bar{\kappa} - \frac{2}{r}, \bar{\kappa} + \frac{2\Upsilon}{r}, \check{\underline{\kappa}}, \underline{\vartheta}, \underline{\xi}, \underline{\omega}, \underline{\bar{\omega}} - \frac{m}{r^2}, \check{\underline{\Omega}}, (\underline{\bar{\Omega}} + \Upsilon), (\bar{\zeta} + 1) \right\}, \\ \check{R} &:= \left\{ \alpha, \beta, \check{\rho}, \bar{\rho} + \frac{2m}{r^3}, \underline{\beta}, \underline{\alpha} \right\}. \end{aligned}$$

Since our foliation is outgoing geodesic we also have,

$$\xi = \omega = 0, \quad \underline{\eta} + \zeta = 0. \quad (9.1.7)$$

We decompose $\check{\Gamma} = \Gamma_g \cup \Gamma_b$ where,

$$\begin{aligned} \Gamma_g &= \left\{ \check{\kappa}, \vartheta, \zeta, \check{\underline{\kappa}}, \bar{\kappa} - \frac{2}{r}, \bar{\kappa} + \frac{2\Upsilon}{r}, \right\}, \\ \Gamma_b &= \left\{ \eta, \underline{\vartheta}, \underline{\xi}, \underline{\omega}, \underline{\bar{\omega}} - \frac{m}{r^2}, r^{-1} \check{\underline{\Omega}}, r^{-1} \check{\zeta}, r^{-1} (\underline{\bar{\Omega}} + \Upsilon), r^{-1} (\bar{\zeta} + 1) \right\}. \end{aligned} \quad (9.1.8)$$

Given a p -reduced scalar $f \in \mathfrak{s}_p(\mathcal{M})$, with respect to the given geodesic foliation on \mathcal{R} , we consider the following norms on spheres $S = S(u, r) \subset \mathcal{R}$,

$$\begin{aligned} \|f\|_\infty(u, r) &:= \|f\|_{L^\infty(S(u,r))}, & \|f\|_2(u, r) &:= \|f\|_{L^2(S(u,r))}, \\ \|f\|_{\infty,k}(u, r) &= \sum_{i=0}^k \|\mathfrak{d}^i f\|_\infty(u, r), & \|f\|_{2,k}(u, r) &= \sum_{i=0}^k \|\mathfrak{d}^i f\|_2(u, r). \end{aligned} \tag{9.1.9}$$

where, we recall, that \mathfrak{d}^i stands for any combination of length i of operators of the form e_3, re_4, \mathfrak{d} . Recall that,

$$\mathfrak{d}^s f = \begin{cases} r^{2p} \mathbb{A}_k^p, & \text{if } s = 2p, \\ r^{2p+1} \mathbb{d}_k \mathbb{A}_k^p, & \text{if } s = 2p + 1. \end{cases} \tag{9.1.10}$$

On a given polarized surface $\mathbf{S} \subset \mathcal{R}$, not necessarily a leaf S of the given foliation, we define

$$\|f\|_{\mathfrak{h}_s^q(\mathbf{S})} := \sum_{i=0}^s \|(\mathfrak{d}^{\mathbf{S}})^i f\|_{L^q(\mathbf{S})}. \tag{9.1.11}$$

where $\mathfrak{d}^{\mathbf{S}}$ is defined as above with respect to the intrinsic metric on \mathbf{S} . In the particular case when $q = 2$ we omit the upper index i.e., $\mathfrak{h}_s(\mathbf{S}) = \mathfrak{h}_s^2(\mathbf{S})$.

9.1.1 Main assumptions

Given an integer s_{max} , we assume the following¹

A1. For all $k \leq s_{max}$, we have on \mathcal{R}

$$\begin{aligned} \|\Gamma_g\|_{k,\infty} &\lesssim \mathring{\epsilon} r^{-2}, \\ \|\Gamma_b\|_{k,\infty} &\lesssim \mathring{\epsilon} r^{-1}, \end{aligned} \tag{9.1.12}$$

and,

$$\begin{aligned} \|\alpha, \beta, \check{\rho}, \check{\mu}\|_{k,\infty} &\lesssim \mathring{\epsilon} r^{-3}, \\ \|e_3(\alpha, \beta)\|_{k-1,\infty} &\lesssim \mathring{\epsilon} r^{-4}, \\ \|\underline{\beta}\|_{k,\infty} &\lesssim \mathring{\epsilon} r^{-2}, \\ \|\underline{\alpha}\|_{k,\infty} &\lesssim \mathring{\epsilon} r^{-1}. \end{aligned} \tag{9.1.13}$$

¹In applications, $s_{max} = k_{small} + 4$ in Theorem M7, and $s_{max} = k_{large} + 5$ in Theorem M0 and Theorem M6.

A2. We have, with m_0 denoting the mass of the unperturbed spacetime,

$$\sup_{\mathcal{R}} \left| \frac{m}{m_0} - 1 \right| \lesssim \overset{\circ}{\epsilon}. \quad (9.1.14)$$

A3. The metric coefficients are assumed to satisfy the following assumptions in \mathcal{R} , for all $k \leq s_{max}$

$$r \left\| \left(\frac{\gamma}{r^2} - 1, \underline{b}, \frac{e^\Phi}{r \sin \theta} - 1 \right) \right\|_{\infty, k} + \|\underline{\Omega} + \Upsilon\|_{\infty, k} + \|\varsigma - 1\|_{\infty, k} \lesssim \overset{\circ}{\epsilon}. \quad (9.1.15)$$

Remark 9.1.2. *The above assumptions imply in particular the following*

$$|e_4(r)|, |e_3(r)| \lesssim 1, \quad e_4(s) = 1 + O(\overset{\circ}{\epsilon}), \quad e_3(u) = 2 + O(\overset{\circ}{\epsilon}), \quad e_4(u) = 0.$$

Hence, since $r = \overset{\circ}{r}$ at $(\overset{\circ}{u}, \overset{\circ}{s})$, we infer

$$|r - \overset{\circ}{r}| \lesssim |s - \overset{\circ}{s}| + |u - \overset{\circ}{u}|,$$

and thus, in view of the definition (9.1.6) of \mathcal{R} ,

$$\sup_{\mathcal{R}} |r - \overset{\circ}{r}| \lesssim \overset{\circ}{\delta}(\overset{\circ}{\epsilon})^{-\frac{1}{2}}. \quad (9.1.16)$$

We will make use of the following lemma, see Lemmas 4.3.3 and 4.3.4.

Lemma 9.1.3. *Under the assumption **A3** for the metric coefficients we have,*

$$r |e_\theta(\Phi)| \leq \frac{2}{\sin \theta}, \quad \frac{1}{\sin \theta} \leq 2(r |e_\theta \Phi| + 1). \quad (9.1.17)$$

Moreover, for any reduced 1-scalar h , we have

$$\sup_S \frac{|h|}{e^\Phi} \lesssim r^{-1} \sup_S (|h| + |\not\partial h|), \quad \left\| \frac{h}{e^\Phi} \right\|_{L^2(S)} \lesssim r^{-1} \|h\|_{\mathfrak{h}_1(S)}. \quad (9.1.18)$$

9.1.2 Elliptic Hodge lemma

We shall often make use of the results of Proposition 2.1.30 and Lemma 2.1.35 which we rewrite as follows.

Lemma 9.1.4. *Under the assumptions **A1**, **A3** the following elliptic estimates hold true for the Hodge operators $\not\partial_1, \not\partial_2, \not\partial_1^*, \not\partial_2^*$, for all $k \leq s_{max}$*

1. If $f \in \mathfrak{s}_1(S)$

$$\|\not\partial f\|_{\mathfrak{h}_k(S)} + \|f\|_{\mathfrak{h}_k(S)} \lesssim r \|\not\partial_1 f\|_{\mathfrak{h}_k(S)}.$$

2. If $f \in \mathfrak{s}_2(S)$

$$\|\not\partial f\|_{\mathfrak{h}_k(S)} + \|f\|_{\mathfrak{h}_k(S)} \lesssim r \|\not\partial_2 f\|_{\mathfrak{h}_k(S)}.$$

3. If $f \in \mathfrak{s}_0(S)$

$$\|\not\partial f\|_{\mathfrak{h}_k(S)} \lesssim r \|\not\partial_1^* f\|_{\mathfrak{h}_k(S)}.$$

4. If $f \in \mathfrak{s}_1(S)$

$$\|f\|_{\mathfrak{h}_{k+1}(S)} \lesssim r \|\not\partial_2^* f\|_{\mathfrak{h}_k(S)} + r^{-2} \left| \int_S e^\Phi f \right|.$$

5. If $f \in \mathfrak{s}_1(S)$

$$\left\| f - \frac{\int_S f e^\Phi}{\int_S e^{2\Phi}} e^\Phi \right\|_{\mathfrak{h}_{k+1}(S)} \lesssim r \|\not\partial_2^* f\|_{\mathfrak{h}_k(S)}.$$

We shall often make use of the following non-sharp product estimate on \mathbf{S} , see Proposition 2.1.40.

Lemma 9.1.5. *The following estimates hold true on a given polarized surface $\mathbf{S} \subset \mathcal{R}$, for any contraction between two reduced scalars ψ_1, ψ_2 , $k \geq 2$,*

$$\|\psi_1 \cdot \psi_2\|_{\mathfrak{h}_k(\mathbf{S})} \lesssim r^{-1} \|\psi_1\|_{\mathfrak{h}_k(\mathbf{S})} \|\psi_2\|_{\mathfrak{h}_k(\mathbf{S})}.$$

9.2 Deformations of S surfaces

9.2.1 Deformations

Recall that $\mathring{S} = S(\mathring{u}, \mathring{s})$ is a fixed sphere of the (u, s) outgoing geodesic foliation of a fixed spacetime region $\mathcal{R} = \mathcal{R}(\mathring{\epsilon}, \mathring{\delta})$.

Definition 9.2.1. We say that \mathbf{S} is an $O(\dot{\epsilon})$ \mathbf{Z} -polarized deformation of \dot{S} if there exists a map $\Psi : \dot{S} \rightarrow \mathbf{S}$ of the form,

$$\Psi(\dot{u}, \dot{s}, \theta, \varphi) = \left(\dot{u} + U(\theta), \dot{s} + S(\theta), \theta, \varphi \right) \quad (9.2.1)$$

where U, S are smooth functions defined on the interval $[0, \pi]$ of amplitude at most $\dot{\epsilon}$. We denote by ψ the reduce map defined on the interval $[0, \pi]$,

$$\psi(\theta) = (\dot{u} + U(\theta), \dot{s} + S(\theta), \theta). \quad (9.2.2)$$

We restrict ourselves to deformations which fix the South Pole, i.e.

$$U(0) = S(0) = 0. \quad (9.2.3)$$

9.2.2 Pull-back map

We recall that given a scalar function f on \mathbf{S} one defines its pull-back on \dot{S} to be the function,

$$f^\# := \Psi^\# f = f \circ \Psi.$$

On the other hand, given a vectorfield X on \dot{S} one defines its push-forward $\Psi_\# X$ to be the vectorfield on \mathbf{S} defined by,

$$\Psi_\# X(f) = X(\Psi^\# f) = X(f \circ \Psi).$$

Given a covariant tensor U on \mathbf{S} , one defines its pull back to \dot{S} to be the tensor

$$\Psi^\# U(X_1, \dots, X_k) = U(\Psi_\# X_1, \dots, \Psi_\# X_k).$$

Lemma 9.2.2. Given a \mathbf{Z} -invariant deformation $\Psi : \dot{S} \rightarrow \mathbf{S}$, we have,

1. Let $g^\mathbf{S}$ the induced metric on \mathbf{S} and $g^{\mathbf{S},\#} = \gamma^{\mathbf{S},\#} d\theta^2 + e^{2\Phi^\#} d\varphi^2$ its pull-back to \dot{S} . The metric coefficients $\gamma^\mathbf{S}$ and $\gamma^{\mathbf{S},\#}$ are related by,

$$\gamma^{\mathbf{S},\#}(\theta) = \gamma^\mathbf{S}(\psi(\theta)) = \gamma^\mathbf{S}(\dot{u} + U(\theta), \dot{s} + S(\theta), \theta) \quad (9.2.4)$$

where $\gamma^\mathbf{S}$ is defined implicitly by,

$$(\gamma^\mathbf{S})^\# = \gamma^\# + (\varsigma^\#)^2 \left(\underline{\Omega} + \frac{1}{4} b^2 \gamma \right)^\# (U')^2 - 2\varsigma^\# U' S' - (\gamma_{\varsigma b})^\# U', \quad (9.2.5)$$

that is,

$$\begin{aligned} \gamma^{\mathbf{S}}(\psi(\theta)) &= \gamma(\psi(\theta)) + \varsigma^2(\psi(\theta)) \left(\underline{\Omega}(\psi(\theta)) + \frac{1}{4}(\underline{b}(\psi(\theta)))^2 \gamma(\psi(\theta)) \right) (U'(\theta))^2 \\ &\quad - 2\varsigma(\psi(\theta))U'(\theta)S'(\theta) - \gamma(\psi(\theta))\varsigma(\psi(\theta))\underline{b}(\psi(\theta))U'(\theta). \end{aligned}$$

2. The \mathbf{Z} -invariant vectorfield $\partial_\theta^{\mathbf{S}} := \Psi_\#(\partial_\theta)$ is tangent to \mathbf{S} and

$$\partial_\theta^{\mathbf{S}}|_{\Psi(p)} = \left[\left(\partial_\theta S - \frac{\varsigma}{2}\underline{\Omega}\partial_\theta U \right) e_4 + \frac{\varsigma}{2}\partial_\theta U e_3 + \sqrt{\gamma} \left(1 - \frac{\varsigma}{2}\underline{b}\partial_\theta U \right) e_\theta \right] \Big|_{\Psi(p)}. \quad (9.2.6)$$

3. If $f \in \mathfrak{s}_k(\mathbf{S})$ and $P^{\mathbf{S}}$ is a geometric operator acting on f then,

$$(P^{\mathbf{S}}[f])^\# = P^{\mathbf{S},\#}[f^\#] \quad (9.2.7)$$

where, $P^{\mathbf{S},\#}$ is the corresponding geometric operator on \mathring{S} with respect to the metric $\mathring{g}^{\mathbf{S},\#}$ and $f^\# = \psi^\# f$.

4. The L^2 norm of $f^\# = \psi^\# f$ with respect to the metric $\mathring{g}^{\mathbf{S},\#}$ is the same as as the L^2 norm of f with respect to the metric $g^{\mathbf{S}}$, i.e.,

$$\int_{\mathring{S}} |f^\#|^2 da_{\mathring{g}^{\mathbf{S},\#}} = \int_{\mathbf{S}} |f|^2 da_{g^{\mathbf{S}}}.$$

5. If $f \in \mathfrak{h}_k(\mathbf{S})$ and $f^\#$ is its pull-back by ψ then,

$$\|f^\#\|_{\mathfrak{h}_k(\mathring{S}, \mathring{g}^{\mathbf{S},\#})} = \|f\|_{\mathfrak{h}_k(\mathbf{S})}.$$

Proof. If ∂_θ denotes the coordinate derivative $\partial_\theta = \frac{\partial}{\partial\theta}$ then, at every point $p \in \mathring{S}$,

$$\Psi_\#(\partial_\theta)|_{\Psi(p)} = \partial_\theta U \partial_u|_{\Psi(p)} + \partial_\theta S \partial_s|_{\Psi(p)} + \partial_\theta|_{\Psi(p)}, \quad \Psi_\#(\partial_\varphi) = \partial_\varphi.$$

In view of (9.1.4) we have

$$\partial_s = e_4, \quad \partial_u = \frac{\varsigma}{2} (e_3 - \underline{\Omega}e_4 - \underline{b}\gamma^{1/2}e_\theta), \quad \partial_\theta = \sqrt{\gamma}e_\theta.$$

Hence, at a point $\Psi(p)$ on \mathbf{S} we have,

$$\Psi_\#(\partial_\theta) = \left(\partial_\theta S - \frac{\varsigma}{2}\underline{\Omega}\partial_\theta U \right) e_4 + \frac{\varsigma}{2}\partial_\theta U e_3 + \sqrt{\gamma} \left(1 - \frac{\varsigma}{2}\underline{b}\partial_\theta U \right) e_\theta.$$

We denote by $\mathring{g}^\# = \Psi^\#(\mathring{g}^\mathbf{S})$ the pull back to \mathring{S} of the metric $\mathring{g}^\mathbf{S}$ on \mathbf{S} , i.e. at any point $p \in \mathring{S}$,

$$\begin{aligned}\mathring{g}^\#(\partial_\theta, \partial_\theta) &= \mathring{g}^\mathbf{S}(\Psi_\# \partial_\theta, \Psi_\# \partial_\theta) = \mathbf{g}(\partial_\theta U \partial_u + \partial_\theta S \partial_s + \partial_\theta, \partial_\theta U \partial_u + \partial_\theta S \partial_s + \partial_\theta) \\ &= (\partial_\theta U)^2 \mathbf{g}_{uu} + 2\partial_\theta U \partial_\theta S \mathbf{g}_{us} + 2\partial_\theta U \mathbf{g}_{u\theta} + \mathbf{g}_{\theta\theta}, \\ \mathring{g}^\#(\partial_\theta, \partial_\varphi) &= 0, \\ \mathring{g}^\#(\partial_\varphi, \partial_\varphi) &= e^{2\Phi^\#},\end{aligned}$$

where,

$$\mathbf{g}_{uu} = \varsigma^2 \left(\underline{\Omega} + \frac{1}{4} \gamma \underline{b}^2 \right), \quad \mathbf{g}_{us} = -\varsigma, \quad \mathbf{g}_{u\theta} = -\frac{\varsigma}{2} \gamma \underline{b}, \quad \mathbf{g}_{ss} = \mathbf{g}_{s\theta} = 0, \quad \mathbf{g}_{\theta\theta} = \gamma.$$

Hence the pull-back metric $\Psi^\#(\mathring{g}^\mathbf{S})$ on \mathring{S} is given by,

$$\gamma^{\mathbf{S},\#} d\theta^2 + e^{2\Phi^\#} d\varphi^2$$

where

$$\gamma^{\mathbf{S},\#} = (\gamma^\mathbf{S})^\#, \quad (9.2.8)$$

with $\gamma^\mathbf{S}$ is defined by,

$$(\gamma^\mathbf{S})^\# = \gamma^\# + (\varsigma^\#)^2 \left(\underline{\Omega} + \frac{1}{4} \underline{b}^2 \gamma \right)^\# (U')^2 - 2\varsigma^\# U' S' - (\gamma \varsigma \underline{b})^\# U'. \quad (9.2.9)$$

Note that the vectorfield,

$$e_\theta^\mathbf{S} := \frac{1}{(\gamma^\mathbf{S})^{1/2}} \psi_\#(\partial_\theta)$$

is tangent, \mathbf{Z} invariant and forms together with e_φ an orthonormal frame on \mathbf{S} . Note that we can also write,

$$e_\theta^\mathbf{S} := \frac{(\mathring{\gamma})^{1/2}}{(\gamma^\mathbf{S})^{1/2}} \Psi_\#(e_\theta)$$

where $\mathring{\gamma}$ is the coefficient in front of $d\theta^2$ of the metric induced by \mathbf{g} on \mathring{S} ,

$$\mathring{g} = \mathring{\gamma} d\theta^2 + e^{2\Phi} d\varphi^2.$$

In general, any geometric calculation on \mathbf{S} can be reduced to a geometric calculation on \mathring{S} with respect to the metric $\mathring{g}^{\mathbf{S},\#}$. Moreover the L^2 norm on \mathbf{S} with respect to the metric $\mathring{g}^\mathbf{S}$ is the same as the L^2 norm of $f^\# = \psi^\# f$ with respect to the norm $\mathring{g}^{\mathbf{S},\#}$. This concludes the proof of the lemma. \square

9.2.3 Comparison of norms between deformations

Lemma 9.2.3. *Let $\Psi : \mathring{S} \rightarrow \mathbf{S}$ a \mathbf{Z} -invariant deformation in $\mathcal{R}(\mathring{\epsilon}, \mathring{\delta})$ with U, V verifying the bounds*

$$\sup_{0 \leq \theta \leq \pi} \left(|U'(\theta)| + |S'(\theta)| \right) \lesssim \mathring{\delta}, \quad (9.2.10)$$

as well as the bound (9.1.15) for the coordinates system (u, s, θ, φ) of \mathcal{R} . The following hold true

1. We have,

$$|\gamma^{\mathbf{S}, \#} - \mathring{\gamma}| \lesssim \mathring{\delta} \mathring{r}. \quad (9.2.11)$$

2. For every $f \in \mathfrak{s}_k(\mathbf{S})$ we have,

$$\|f^\#\|_{L^2(\mathring{S}, \mathring{g}^{\mathbf{S}, \#})} = \|f^\#\|_{L^2(\mathring{S}, \mathring{g})} \left(1 + O(r^{-1} \mathring{\delta}) \right). \quad (9.2.12)$$

3. As a corollary of (9.2.12) (choosing $f = 1$) we deduce²,

$$\frac{r^{\mathbf{S}}}{\mathring{r}} = 1 + O(\mathring{r}^{-1} \mathring{\delta}) \quad (9.2.13)$$

where $r^{\mathbf{S}}$ is the area radius of \mathbf{S} and \mathring{r} that of \mathring{S} .

Proof. Recall,

$$\begin{aligned} \gamma^{\mathbf{S}, \#}(\mathring{u}, \mathring{s}, \theta) &= \gamma(\psi(\theta)) + \varsigma^2(\psi(\theta)) \left(\underline{\Omega}(\psi(\theta)) + \frac{1}{4}(\underline{b}(\psi(\theta)))^2 \gamma(\psi(\theta)) \right) (U'(\theta))^2 \\ &\quad - 2\varsigma(\psi(\theta))U'(\theta)S'(\theta) - \gamma(\psi(\theta))\varsigma(\psi(\theta))\underline{b}(\psi(\theta))U'(\theta). \end{aligned}$$

In view of our assumptions on U' and S' as well as our estimates (9.1.15) for γ , $\underline{\Omega}$ and \underline{b} and ς , we infer

$$|\gamma^{\mathbf{S}, \#} - \gamma| \lesssim |\gamma^\# - \gamma| + \mathring{r} \mathring{\epsilon}^{1/2} \mathring{\delta}.$$

²Recall also from (9.1.16) that $r - \mathring{r} = O(\mathring{\delta} \mathring{\epsilon}^{-1/2})$.

Also, we have

$$\begin{aligned}
\gamma^\#(\overset{\circ}{u}, \overset{\circ}{s}, \theta) - \gamma(\overset{\circ}{u}, \overset{\circ}{s}, \theta) &= \gamma(\overset{\circ}{u} + U(\theta), \overset{\circ}{s} + S(\theta), \theta) - \gamma(\overset{\circ}{u}, \overset{\circ}{s}, \theta) \\
&= \int_0^1 \frac{d}{d\lambda} \left[\gamma(\overset{\circ}{u} + \lambda U(\theta), \overset{\circ}{s} + \lambda S(\theta), \theta) \right] d\lambda \\
&= U(\theta) \int_0^1 \partial_u \gamma(\overset{\circ}{u} + \lambda U(\theta), \overset{\circ}{s} + \lambda S(\theta), \theta) d\lambda \\
&\quad + S(\theta) \int_0^1 \partial_s \gamma(\overset{\circ}{u} + \lambda U(\theta), \overset{\circ}{s} + \lambda S(\theta), \theta) d\lambda.
\end{aligned}$$

In view of our estimates (9.1.15) for γ , the assumption (9.2.10) on (U', V') and the fact that

$$\partial_s = e_4, \quad \partial_u = \frac{\zeta}{2} (e_3 - \underline{\Omega} e_4 - \underline{b} \gamma^{1/2} e_\theta),$$

we infer³

$$|\gamma^\# - \gamma| \lesssim \overset{\circ}{r} \overset{\circ}{\delta}.$$

We have finally, obtained

$$|\gamma^{\mathbf{S}, \#} - \gamma| \lesssim |\gamma^\# - \gamma| + \overset{\circ}{r} \overset{\circ}{\delta} \lesssim \overset{\circ}{r} \overset{\circ}{\delta}.$$

To prove the second part of the lemma we write,

$$\int_{\overset{\circ}{S}} |f^\#|^2 da_{\mathcal{J}^{\mathbf{S}, \#}} = \int_{\overset{\circ}{S}} |f^\#|^2 \frac{\sqrt{\gamma^{\mathbf{S}, \#}}}{\sqrt{\overset{\circ}{\gamma}}} da_{\overset{\circ}{\mathcal{J}}} = \int_{\overset{\circ}{S}} |f^\#|^2 da_{\overset{\circ}{\mathcal{J}}} + \int_{\overset{\circ}{S}} |f^\#|^2 \left(\frac{\sqrt{\gamma^{\mathbf{S}, \#}}}{\sqrt{\overset{\circ}{\gamma}}} - 1 \right) da_{\overset{\circ}{\mathcal{J}}}$$

which yields, in view of the first part,

$$\int_{\overset{\circ}{S}} |f^\#|^2 da_{\mathcal{J}^{\mathbf{S}, \#}} = \int_{\overset{\circ}{S}} |f^\#|^2 da_{\overset{\circ}{\mathcal{J}}} \left(1 + O(\overset{\circ}{r}^{-1} \overset{\circ}{\delta}) \right).$$

This concludes the proof of the lemma. \square

Remark 9.2.4. In view of (9.2.13) and (9.1.16), $\overset{\circ}{r}$, $r^{\mathbf{S}}$ and the value of r along \mathbf{S} are all comparable.

Corollary 9.2.5. Under the assumptions of Lemma 9.2.3 the following estimate⁴ holds true for an arbitrary scalar $f \in \mathfrak{s}_0(\mathcal{R})$,

$$\left| \int_{\mathbf{S}} f - \int_{\overset{\circ}{S}} f \right| \lesssim \overset{\circ}{\delta} \overset{\circ}{r} \left(\sup_{\mathcal{R}} |\mathfrak{d}_{\overrightarrow{\mathbf{S}}}^{\leq 1} f| + \sup_{\mathcal{R}} r |e_3 f| \right).$$

³Note that we also use the assumption $U(0) = S(0) = 0$ to estimate (U, S) from (U', S') .

⁴Recall that $\mathcal{R} := \{|u - \overset{\circ}{u}| \leq \delta_{\mathcal{R}}, |s - \overset{\circ}{s}| \leq \delta_{\mathcal{R}}\}$, see (9.1.6).

Proof. We have,

$$\int_{\mathbf{S}} f - \int_{\mathring{S}} f = \int_{\mathring{S}} f^\# \frac{\sqrt{\gamma^{\mathbf{S},\#}}}{\sqrt{\gamma}} - \int_{\mathring{S}} f = \int_{\mathring{S}} f^\# \left(\frac{\sqrt{\gamma^{\mathbf{S},\#}}}{\sqrt{\gamma}} - 1 \right) + \int_{\mathring{S}} (f^\# - f).$$

Hence,

$$\left| \int_{\mathbf{S}} f - \int_{\mathring{S}} f \right| \lesssim \mathring{\delta} r \sup_{\mathbf{S}} |f| + \int_{\mathring{S}} |f^\# - f|.$$

Now, proceeding as in the proof of (9.2.11),

$$\begin{aligned} f(\mathring{u} + U(\theta), \mathring{s} + S(\theta)) - f(\mathring{u}, \mathring{s}) &\lesssim \int_0^1 \frac{d}{d\lambda} \left[f(\mathring{u} + \lambda U(\theta), \mathring{s} + \lambda S(\theta), \theta) \right] d\lambda \\ &= U(\theta) \int_0^1 \partial_u f(\mathring{u} + \lambda U(\theta), \mathring{s} + \lambda S(\theta), \theta) d\lambda \\ &\quad + S(\theta) \int_0^1 \partial_s f(\mathring{u} + \lambda U(\theta), \mathring{s} + \lambda S(\theta), \theta) d\lambda. \end{aligned}$$

Therefore ,

$$\begin{aligned} \left| \int_{\mathbf{S}} f - \int_{\mathring{S}} f \right| &\lesssim \mathring{r} \mathring{\delta} \sup_{\mathbf{S}} |f| + \mathring{\delta} r \left(\sup_{\mathcal{R}} \mathring{r} |\mathring{\partial}_\lambda f| + \sup_{\mathcal{R}} r |e_3 f| \right) \\ &\lesssim \mathring{\delta} \mathring{r} \left(\sup_{\mathcal{R}} |\mathring{\partial}_\lambda^{\leq 1} f| + \sup_{\mathcal{R}} r |e_3 f| \right) \end{aligned}$$

as stated. □

To compare higher order Sobolev spaces, we will need the following lemma.

Lemma 9.2.6. *Let $\mathring{S} \subset \mathcal{R} = \mathcal{R}(\mathring{\epsilon}, \mathring{\delta})$ as in Definition 9.1.1 verifying the assumptions **A1-A3**. Let $\Psi : \mathring{S} \rightarrow \mathbf{S}$ be \mathbf{Z} -invariant deformation. Assume the bound*

$$\|(U', S')\|_{L^\infty(\mathring{S})} + \max_{0 \leq s \leq s_{max}} (\mathring{r})^{-1} \|(U', S')\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})} \lesssim \mathring{\delta}. \quad (9.2.14)$$

Then, we have for any reduced scalar h defined on \mathcal{R}

$$\|h\|_{\mathfrak{h}_s(\mathbf{S})} \lesssim r \sup_{\mathcal{R}} |\mathring{\partial}^{\leq s} h|, \quad \text{for } 0 \leq s \leq s_{max}.$$

Also, if $f \in \mathfrak{h}_s(\mathbf{S})$ and $f^\#$ is its pull-back by ψ , we have

$$\|f\|_{\mathfrak{h}_s(\mathbf{S})} = \|f^\#\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g}^{\mathbf{S},\#})} = \|f^\#\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})} (1 + O(r^{-1} \mathring{\delta})) \quad \text{for } 0 \leq s \leq s_{max} - 2.$$

Proof. See appendix C.1. □

Corollary 9.2.7. *Under the same assumptions as Lemma 9.2.6, we have, for all $0 \leq k \leq s_{max} - 2$,*

$$\begin{aligned} \|\mathfrak{d}^{\leq 2} \Gamma_g\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-1}, \\ \|\mathfrak{d}^{\leq 2} \Gamma_b\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon}, \end{aligned} \tag{9.2.15}$$

$$\begin{aligned} \|\mathfrak{d}^{\leq 2} (\alpha, \beta, \check{\rho}, \check{\mu})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-2}, \\ \|\mathfrak{d}^{\leq 2} \underline{\beta}\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-1}, \\ \|\mathfrak{d}^{\leq 2} \underline{\alpha}\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon}, \end{aligned} \tag{9.2.16}$$

$$\begin{aligned} \left\| \mathfrak{d}^{\leq 2} \left(\frac{\gamma}{r^2} - 1, \underline{b}, \frac{e^\Phi}{r \sin \theta} - 1 \right) \right\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon}, \\ \|\mathfrak{d}^{\leq 2} (\underline{\Omega} + \Upsilon)\|_{\mathfrak{h}_k(\mathbf{S})} + \|\mathfrak{d}^{\leq 2} (\varsigma + 1)\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r. \end{aligned} \tag{9.2.17}$$

Proof. In view of Lemma 9.2.6 and assumptions **A1-A3** we have, for $0 \leq s \leq s_{max} - 2$,

$$\|\mathfrak{d}^{\leq 2} \check{\Gamma}_g\|_{\mathfrak{h}_s(\mathbf{S})} \lesssim r \sup_{\mathcal{R}} |\mathfrak{d}^{\leq s+2} \check{\Gamma}_g| \lesssim r^{-1} \overset{\circ}{\epsilon}.$$

The other estimates are proved in the same manner. □

9.2.4 Adapted frame transformations

We consider general null transformations introduced in Lemma 2.3.1,

$$\begin{aligned} e'_4 &= \lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right), \\ e'_\theta &= \left(1 + \frac{1}{2} f \underline{f} \right) e_\theta + \frac{1}{2} \underline{f} e_4 + \frac{1}{2} f \left(1 + \frac{1}{4} f \underline{f} \right) e_3, \\ e'_3 &= \lambda^{-1} \left(\left(1 + \frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) e_3 + \underline{f} \left(1 + \frac{1}{4} f \underline{f} \right) e_\theta + \frac{1}{4} \underline{f}^2 e_4 \right). \end{aligned} \tag{9.2.18}$$

Definition 9.2.8. *Given a deformation $\Psi : \overset{\circ}{S} \rightarrow \mathbf{S}$ we say that a new frame (e'_3, e'_4, e'_θ) , obtained from the standard frame (e_3, e_4, e_θ) via the transformation (9.2.18), is \mathbf{S} -adapted if we have,*

$$e'_\theta = e_\theta^{\mathbf{S}} = \frac{1}{(\gamma^{\mathbf{S}})^{1/2}} \psi_{\#}(\partial_\theta). \tag{9.2.19}$$

Proposition 9.2.9. *Consider a deformation $\Psi : \overset{\circ}{S} \rightarrow \mathbf{S}$ in $\mathcal{R} = \mathcal{R}(\overset{\circ}{\epsilon}, \overset{\circ}{\delta})$ verifying the assumption **A3**. The following statements hold true.*

1. *A new frame e'_3, e'_θ, e'_4 generated by $(f, \underline{f}, \lambda = e^a)$ according to (9.2.18) is adapted to $\mathbf{S} = \mathbf{S}(\overset{\circ}{u} + U, \overset{\circ}{s} + S)$ provided that, at all points $\theta \in [0, \pi]$,*

$$\begin{aligned} \sqrt{\gamma^\#} \left(1 - \frac{\varsigma}{2} \underline{b}^\# U'\right) &= ((\gamma^{\mathbf{S}})^\#)^{1/2} \left(1 + \frac{1}{2} (f \underline{f})^\#\right), \\ \varsigma U' &= ((\gamma^{\mathbf{S}})^\#)^{1/2} f^\# \left(1 + \frac{1}{4} (f \underline{f})^\#\right), \\ 2 \left(S' - \frac{\varsigma}{2} \underline{\Omega}^\# U'\right) &= ((\gamma^{\mathbf{S}})^\#)^{1/2} \underline{f}^\#, \end{aligned} \quad (9.2.20)$$

where,

$$(\gamma^{\mathbf{S}})^\# = \gamma^\# + (\varsigma^\#)^2 \left(\underline{\Omega} + \frac{1}{4} \underline{b}^2 \gamma\right)^\# (\partial_\theta U)^2 - 2\varsigma^\# \partial_\theta U \partial_\theta S - (\gamma \varsigma \underline{b})^\# \partial_\theta U$$

and $\#$ denotes the pull back by ψ of the corresponding reduced scalars, i.e. for example, $f^\#(\theta) = f(\overset{\circ}{u} + U(\theta), \overset{\circ}{s} + S(\theta), \theta)$.

2. *There exists a small enough constant⁵ δ_1 such that for given f, \underline{f} on \mathcal{R} satisfying*

$$\sup_{\mathcal{R}} (|f| + |\underline{f}|) \leq r^{-1} \delta_1,$$

we can uniquely solve the system (9.2.20) for U, S subject to the initial conditions,

$$U(0) = 0, \quad S(0) = 0.$$

Thus, if $(\overset{\circ}{u}, \overset{\circ}{s}, 0)$ corresponds to the south pole of $\overset{\circ}{S}$ and f, \underline{f} are given there exists a unique deformation $\mathbf{S} \subset \mathcal{R}$, given by $U, S : [0, \pi] \rightarrow \mathbb{R}$, adapted to frames generated by⁶ (f, \underline{f}) which passes through the same south pole. Moreover,

$$\sup_{[0, \pi]} |(U', S')| \lesssim \overset{\circ}{r} \sup_{\mathbf{S}} (|f| + |\underline{f}|) \quad (9.2.21)$$

⁵In later applications, we will have

$$\sup_{\mathcal{R}} (|f| + |\underline{f}|) \lesssim r^{-1} \overset{\circ}{\delta}.$$

⁶Note that a is not restricted in this result.

and, for $2 \leq s \leq s_{max}$,

$$\|(U', S')\|_{L^\infty(\overset{\circ}{S})} + (\overset{\circ}{r})^{-1} \|(U', S')\|_{\mathfrak{h}_s(\overset{\circ}{S}, \overset{\circ}{g})} \lesssim \|f, \underline{f}\|_{\mathfrak{h}_s(\mathbf{S})} \quad (9.2.22)$$

with $\|f, \underline{f}\|_{\mathfrak{h}_s(\mathbf{S})} = \|f\|_{\mathfrak{h}_s(\mathbf{S})} + \|\underline{f}\|_{\mathfrak{h}_s(\mathbf{S})}$.

3. As a consequence of (9.2.22) the deformation thus obtained verifies the conclusions of Lemmas 9.2.3-9.2.6 and Corollary 9.2.7. In particular,

(a) We have,

$$\left| \gamma^{\mathbf{S}, \#} - \overset{\circ}{\gamma} \right| \lesssim \delta_1 \overset{\circ}{r}.$$

(b) We have

$$\left| \frac{r^{\mathbf{S}}}{\overset{\circ}{r}} - 1 \right| \lesssim \overset{\circ}{r}^{-1} \delta_1.$$

Proof. In view of Lemma 9.2.2, The \mathbf{Z} -invariant vectorfield $e_\theta^{\mathbf{S}} := \frac{1}{(\gamma^{\mathbf{S}})^{1/2}} \Psi_{\#}(\partial_\theta)$ can be expressed by the formula,

$$e_\theta^{\mathbf{S}} = \frac{1}{(\gamma^{\mathbf{S}})^{1/2}} \left[\left(\partial_\theta S - \frac{\varsigma}{2} \underline{\Omega} \partial_\theta U \right) e_4 + \frac{\varsigma}{2} \partial_\theta U e_3 + \sqrt{\gamma} \left(1 - \frac{\varsigma}{2} \underline{b} \partial_\theta U \right) e_\theta \right].$$

where $\psi(p) = (\overset{\circ}{u} + U(\theta), \overset{\circ}{s} + S(\theta), \theta)$ and $U' = \partial_\theta U(\theta)$, $S' = \partial_\theta S(\theta)$. On the other hand, according to (9.2.18), at $\Psi(p) \in \mathbf{S}$,

$$e'_\theta = \left(1 + \frac{1}{2} f \underline{f} \right) e_\theta + \frac{1}{2} f \left(1 + \frac{1}{4} f \underline{f} \right) e_3 + \frac{1}{2} \underline{f} e_4.$$

We deduce, at every $\theta \in [0, \pi]$,

$$\begin{aligned} \sqrt{\gamma^{\#}} \left(1 - \frac{\varsigma^{\#}}{2} \underline{b}^{\#} U' \right) &= ((\gamma^{\mathbf{S}})^{\#})^{1/2} \left(1 + \frac{1}{2} (f \underline{f})^{\#} \right), \\ \varsigma^{\#} U' &= ((\gamma^{\mathbf{S}})^{\#})^{1/2} f^{\#} \left(1 + \frac{1}{4} (f \underline{f})^{\#} \right), \\ 2 \left(S' - \frac{\varsigma^{\#}}{2} \underline{\Omega}^{\#} U' \right) &= ((\gamma^{\mathbf{S}})^{\#})^{1/2} \underline{f}^{\#}, \end{aligned}$$

as desired.

To prove the second part of the lemma we first check for the compatibility of the three equations in (9.2.20). Note that, if we denote,

$$A = 1 + \frac{1}{2}(f\underline{f})^\#, \quad B = f^\# \left(1 + \frac{1}{4}(f\underline{f})^\# \right), \quad C = \underline{f}^\#,$$

we have $A^2 - BC = 1$. Hence, squaring the first equation and subtracting the product of the other two we derive,

$$\begin{aligned} (\gamma^{\mathbf{S}})^\# &= \left(\sqrt{\gamma^\#} \left(1 - \frac{\varsigma^\#}{2} \underline{b}^\# U' \right) \right)^2 - 2U' \varsigma^\# \left(S' - \frac{\varsigma^\#}{2} \underline{\Omega}^\# U' \right) \\ &= \gamma^\# \left(1 - (\varsigma \underline{b})^\# U' + \frac{1}{4} ((\varsigma \underline{b})^\# U')^2 \right) - 2\varsigma^\# U' S' + (\varsigma^\#)^2 \underline{\Omega}^\# (U')^2 \\ &= \gamma^\# + (\varsigma^\#)^2 \left(\underline{\Omega}^\# + \frac{1}{4} (\underline{b}^\#)^2 \gamma^\# \right) (U')^2 - 2\varsigma^\# U' S' - \gamma^\# \varsigma^\# \underline{b}^\# U' \end{aligned} \quad (9.2.23)$$

which coincides with the formula (9.2.5). It thus suffices to only consider the last two equations in (9.2.20) which we write in the form,

$$\begin{aligned} U' &= (\varsigma^\#)^{-1} ((\gamma^{\mathbf{S}})^\#)^{1/2} f^\# \left(1 + \frac{1}{4}(f\underline{f})^\# \right), \\ S' &= \frac{1}{2} ((\gamma^{\mathbf{S}})^\#)^{1/2} \underline{f}^\# + \frac{1}{2} \underline{\Omega}^\# ((\gamma^{\mathbf{S}})^\#)^{1/2} f^\# \left(\underline{f}^\# + \frac{1}{4}(f\underline{f})^\# \right), \end{aligned} \quad (9.2.24)$$

i.e.,

$$\begin{aligned} U'(\theta) &= \left[(\varsigma^\#)^{-1} (\gamma^{\mathbf{S}})^{1/2} f \left(1 + \frac{1}{4}(f\underline{f}) \right) \right] (\overset{\circ}{u} + U(\theta), \overset{\circ}{s} + S(\theta), \theta), \\ S'(\theta) &= \left[\frac{1}{2} (\gamma^{\mathbf{S}})^{1/2} \underline{f} + \frac{1}{2} \underline{\Omega} (\gamma^{\mathbf{S}})^{1/2} f \left(1 + \frac{1}{4} f\underline{f} \right) \right] (\overset{\circ}{u} + U(\theta), \overset{\circ}{s} + S(\theta), \theta). \end{aligned}$$

Thus under the assumption $\sup_{\mathcal{R}}(|f| + |\underline{f}|) \leq \overset{\circ}{r}^{-1} \delta_1$, with δ_1 sufficiently small, making also use of the expression (9.2.23) of γ^S , and the estimates (9.1.15) for $(\gamma, \underline{b}, \underline{\Omega})$, for $\overset{\circ}{\epsilon}$ sufficiently small, we can uniquely solve for U, S subject to the initial conditions,

$$U(0) = 0, \quad S(0) = 0.$$

Moreover the solution verifies,

$$\sup_{[0, \pi]} |(U', S')| \lesssim \overset{\circ}{r} \sup_{\mathbf{S}} (|f| + |\underline{f}|)$$

according to the Definition 9.2.1. Estimate (9.2.22) can be easily derived by taking higher derivatives and using **A1-A3**. This concludes the proof of the lemma. \square

9.3 Frame transformations

For the convenience of the reader we start by recalling the transformation formulas recorded in Proposition 2.3.4.

Proposition 9.3.1 (Transformation formulas-GCM). *Under a general transformation of type (9.2.18) with $\lambda = e^a$ the Ricci coefficients and curvature components transform as follows:*

$$\begin{aligned}
 \xi' &= \lambda^2 \left(\xi + \frac{1}{2} \lambda^{-1} e'_4(f) + \omega f + \frac{1}{4} f \kappa \right) + \lambda^2 \text{Err}(\xi, \xi'), \\
 \text{Err}(\xi, \xi') &= \frac{1}{4} f \vartheta + l.o.t., \\
 \underline{\xi}' &= \lambda^{-2} \left(\underline{\xi} + \frac{1}{2} \lambda e'_3(\underline{f}) + \underline{\omega} \underline{f} + \frac{1}{4} \underline{f} \underline{\kappa} \right) + \lambda^{-2} \text{Err}(\underline{\xi}, \underline{\xi}'), \\
 \text{Err}(\underline{\xi}, \underline{\xi}') &= -\frac{1}{8} \lambda \underline{f}^2 e'_3(f) + \frac{1}{4} \underline{f} \underline{\vartheta} + l.o.t.,
 \end{aligned} \tag{9.3.1}$$

$$\begin{aligned}
 \zeta' &= \zeta - e'_\theta(\log(\lambda)) + \frac{1}{4} (-f \underline{\kappa} + \underline{f} \kappa) + \underline{f} \omega - f \underline{\omega} + \text{Err}(\zeta, \zeta'), \\
 \text{Err}(\zeta, \zeta') &= \frac{1}{2} \underline{f} e'_\theta(f) + \frac{1}{4} (-f \underline{\vartheta} + \underline{f} \vartheta) + l.o.t., \\
 \eta' &= \eta + \frac{1}{2} \lambda e'_3(f) + \frac{1}{4} \kappa \underline{f} - f \underline{\omega} + \text{Err}(\eta, \eta'), \\
 \text{Err}(\eta, \eta') &= \frac{1}{4} \underline{f} \vartheta + l.o.t., \\
 \underline{\eta}' &= \underline{\eta} + \frac{1}{2} \lambda^{-1} e'_4(\underline{f}) + \frac{1}{4} \underline{\kappa} \underline{f} - \underline{f} \omega + \text{Err}(\underline{\eta}, \underline{\eta}'), \\
 \text{Err}(\underline{\eta}, \underline{\eta}') &= -\frac{1}{8} \underline{f}^2 \lambda^{-1} e'_4(f) + \frac{1}{4} \underline{f} \underline{\vartheta} + l.o.t.,
 \end{aligned} \tag{9.3.2}$$

$$\begin{aligned}
 \kappa' &= \lambda (\kappa + \phi'_1(f)) + \lambda \text{Err}(\kappa, \kappa'), \\
 \text{Err}(\kappa, \kappa') &= f(\zeta + \eta) + \underline{f} \xi - \frac{1}{4} f^2 \underline{\kappa} + f \underline{f} \omega - f^2 \underline{\omega} + l.o.t., \\
 \underline{\kappa}' &= \lambda^{-1} (\underline{\kappa} + \phi'_1(\underline{f})) + \lambda^{-1} \text{Err}(\underline{\kappa}, \underline{\kappa}'), \\
 \text{Err}(\underline{\kappa}, \underline{\kappa}') &= -\frac{1}{4} \underline{f}^2 e'_\theta(f) + \underline{f}(-\zeta + \eta) + \underline{f} \underline{\xi} - \frac{1}{4} \underline{f}^2 \underline{\kappa} + \underline{f} \underline{f} \omega - \underline{f}^2 \underline{\omega} + l.o.t.,
 \end{aligned} \tag{9.3.3}$$

$$\begin{aligned}
\vartheta' &= \lambda(\vartheta - \mathcal{A}_2^*(f)) + \lambda \text{Err}(\vartheta, \vartheta'), \\
\text{Err}(\vartheta, \vartheta') &= f(\zeta + \eta) + \underline{f}\xi + \frac{1}{4}f\underline{f}\kappa + f\underline{f}\omega - f^2\underline{\omega} + l.o.t. \\
\underline{\vartheta}' &= \lambda^{-1}(\underline{\vartheta} - \mathcal{A}_2^*(\underline{f})) + \lambda^{-1} \text{Err}(\underline{\vartheta}, \underline{\vartheta}'), \\
\text{Err}(\underline{\vartheta}, \underline{\vartheta}') &= -\frac{1}{4}\underline{f}^2 e'_\theta(f) + \underline{f}(-\zeta + \eta) + \underline{f}\xi + \frac{1}{4}f\underline{f}\kappa + f\underline{f}\omega - \underline{f}^2\omega + l.o.t.,
\end{aligned} \tag{9.3.4}$$

$$\begin{aligned}
\omega' &= \lambda \left(\omega - \frac{1}{2}\lambda^{-1}e'_4(\log(\lambda)) \right) + \lambda \text{Err}(\omega, \omega'), \\
\text{Err}(\omega, \omega') &= \frac{1}{4}\underline{f}e'_4(f) + \frac{1}{2}\omega\underline{f}\underline{f} - \frac{1}{2}\underline{f}\eta + \frac{1}{2}\underline{f}\xi + \frac{1}{2}\underline{f}\zeta - \frac{1}{8}\kappa\underline{f}^2 + \frac{1}{8}f\underline{f}\kappa - \frac{1}{4}\omega\underline{f}^2 + l.o.t., \\
\underline{\omega}' &= \lambda^{-1} \left(\underline{\omega} + \frac{1}{2}\lambda e'_3(\log(\lambda)) \right) + \lambda^{-1} \text{Err}(\underline{\omega}, \underline{\omega}'), \\
\text{Err}(\underline{\omega}, \underline{\omega}') &= -\frac{1}{4}\underline{f}e'_3(f) + \underline{\omega}\underline{f}\underline{f} - \frac{1}{2}\underline{f}\eta + \frac{1}{2}\underline{f}\xi - \frac{1}{2}\underline{f}\zeta - \frac{1}{8}\kappa\underline{f}^2 + \frac{1}{8}f\underline{f}\kappa - \frac{1}{4}\omega\underline{f}^2 + l.o.t.
\end{aligned} \tag{9.3.5}$$

The lower order terms we denote by *l.o.t.* are linear with respect to $\{\xi, \underline{\xi}, \vartheta, \kappa, \eta, \underline{\eta}, \zeta, \underline{\kappa}, \underline{\vartheta}\}$ and quadratic or higher order in f, \underline{f} , and do not contain derivatives of these latter.

Also,

$$\begin{aligned}
\alpha' &= \lambda^2\alpha + \lambda^2 \text{Err}(\alpha, \alpha'), \\
\text{Err}(\alpha, \alpha') &= 2f\beta + \frac{3}{2}f^2\rho + l.o.t., \\
\beta' &= \lambda \left(\beta + \frac{3}{2}\rho f \right) + \lambda \text{Err}(\beta, \beta'), \\
\text{Err}(\beta, \beta') &= \frac{1}{2}\underline{f}\alpha + l.o.t., \\
\rho' &= \rho + \text{Err}(\rho, \rho'), \\
\text{Err}(\rho, \rho') &= \frac{3}{2}\rho\underline{f}\underline{f} + \underline{f}\beta + f\underline{\beta} + l.o.t., \\
\underline{\beta}' &= \lambda^{-1} \left(\underline{\beta} + \frac{3}{2}\rho\underline{f} \right) + \lambda^{-1} \text{Err}(\underline{\beta}, \underline{\beta}'), \\
\text{Err}(\underline{\beta}, \underline{\beta}') &= \frac{1}{2}\underline{f}\alpha + l.o.t., \\
\underline{\alpha}' &= \lambda^{-2}\underline{\alpha} + \lambda^{-2} \text{Err}(\underline{\alpha}, \underline{\alpha}'), \\
\text{Err}(\underline{\alpha}, \underline{\alpha}') &= 2\underline{f}\underline{\beta} + \frac{3}{2}\underline{f}^2\rho + l.o.t.
\end{aligned} \tag{9.3.6}$$

The lower order terms we denote by *l.o.t.* are linear with respect to the curvature quantities $\alpha, \beta, \rho, \underline{\beta}, \underline{\alpha}$ and quadratic or higher order in f, \underline{f} , and do not contain derivatives of these latter.

In the following lemma we rewrite a subset of these transformations in a more useful form,

Lemma 9.3.2. *Under a general transformation of type (9.2.18) with $\lambda = e^a$ we have, in particular,*

$$\begin{aligned} \zeta' &= \zeta - e'_\theta(a) - f\underline{\omega} + \underline{f}\omega - \frac{1}{2}f\underline{\chi} + \frac{1}{2}\underline{f}\chi + Err(\zeta, \zeta'), \\ Err(\zeta, \zeta') &= \frac{1}{2}\underline{f} \left(1 + \frac{1}{4}f\underline{f} \right) e'_\theta(f) - \frac{1}{16}\underline{f}^2 e'_\theta(f^2) + \frac{1}{4}(-f\underline{\vartheta} + \underline{f}\vartheta) + l.o.t. \end{aligned} \quad (9.3.7)$$

$$\begin{aligned} \kappa' &= e^a (\kappa + \not{d}'_1 f) + e^a Err(\kappa, \kappa'), \\ Err(\kappa, \kappa') &= \frac{1}{2}f\underline{f} e'_\theta(f) - \frac{1}{4}\underline{f} e'_\theta(f^2) + f(\zeta + \eta) + \underline{f}\xi - \frac{1}{4}f^2\underline{\kappa} + f\underline{f}\omega - f^2\underline{\omega} + l.o.t. \end{aligned} \quad (9.3.8)$$

$$\begin{aligned} \underline{\kappa}' &= e^{-a} (\underline{\kappa} + \not{d}'_1 \underline{f}) + e^{-a} Err(\underline{\kappa}, \underline{\kappa}'), \\ Err(\underline{\kappa}, \underline{\kappa}') &= -\frac{1}{2}\underline{f} e'_\theta \left(\underline{f}\underline{f} + \frac{1}{8}f^2\underline{f}^2 \right) + \left(\frac{3}{4}\underline{f}\underline{f} + \frac{1}{8}(f\underline{f})^2 \right) e'_\theta(\underline{f}) \\ &\quad + \frac{1}{4} \left(1 + \frac{1}{2}\underline{f}\underline{f} \right) \underline{f} e'_\theta(f\underline{f}) - \frac{1}{4}\underline{f} \left(1 + \frac{1}{4}\underline{f}\underline{f} \right) e'_\theta(\underline{f}^2) \\ &\quad + \underline{f}(-\zeta + \eta) + \underline{f}\xi - \frac{1}{4}\underline{f}^2\underline{\kappa} + f\underline{f}\omega - \underline{f}^2\omega + l.o.t. \end{aligned} \quad (9.3.9)$$

Also,

$$\begin{aligned} \vartheta' &= \lambda (\vartheta - \not{d}'_2(f)) + \lambda Err(\vartheta, \vartheta'), \\ Err(\vartheta, \vartheta') &= \frac{1}{2}f\underline{f} e'_\theta(f) - \frac{1}{4}\underline{f} e'_\theta(f^2) + f(\zeta + \eta) + \underline{f}\xi + \frac{1}{4}f\underline{f}\kappa + f\underline{f}\omega - f^2\underline{\omega} + l.o.t. \\ \underline{\vartheta}' &= \lambda^{-1} (\underline{\vartheta} - \not{d}'_2(\underline{f})) + \lambda^{-1} Err(\underline{\vartheta}, \underline{\vartheta}'), \\ Err(\underline{\vartheta}, \underline{\vartheta}') &= -\frac{1}{2}\underline{f} e'_\theta \left(\underline{f}\underline{f} + \frac{1}{8}f^2\underline{f}^2 \right) + \left(\frac{3}{4}\underline{f}\underline{f} + \frac{1}{8}(f\underline{f})^2 \right) e'_\theta(\underline{f}) \\ &\quad + \frac{1}{4} \left(1 + \frac{1}{2}\underline{f}\underline{f} \right) \underline{f} e'_\theta(f\underline{f}) - \frac{1}{4}\underline{f} \left(1 + \frac{1}{4}\underline{f}\underline{f} \right) e'_\theta(\underline{f}^2) + \underline{f}(-\zeta + \eta) \\ &\quad + \underline{f}\xi + \frac{1}{4}f\underline{f}\kappa + f\underline{f}\omega - \underline{f}^2\omega + l.o.t., \end{aligned} \quad (9.3.10)$$

The lower order terms we denote by *l.o.t.* are cubic or higher order in the small quantities $\xi, \underline{\xi}, \vartheta, \eta, \underline{\eta}, \zeta, \underline{\zeta}$ as well as f, \underline{f} , and do not contain derivatives of these quantities.

We also have,

$$\begin{aligned}
\beta' &= \lambda \left(\beta + \frac{3}{2} \rho f \right) + \lambda \text{Err}(\beta, \beta'), \\
\text{Err}(\beta, \beta') &= \frac{1}{2} \underline{f} \alpha + \text{l.o.t.}, \\
\rho' &= \rho + \text{Err}(\rho, \rho'), \\
\text{Err}(\rho, \rho') &= \frac{3}{2} \rho f \underline{f} + \underline{f} \beta + f \underline{\beta} + \text{l.o.t.}
\end{aligned} \tag{9.3.11}$$

The lower order terms above denoted by *l.o.t.* are cubic or higher order in the small quantities $\xi, \underline{\xi}, \vartheta, \eta, \underline{\eta}, \zeta, \underline{\vartheta}$ as well as a, f, \underline{f} .

Lemma 9.3.3. *The following transformation formula holds true*

$$\begin{aligned}
\mu' &= \mu + (\mathcal{A}_1)' \left(-(\mathcal{A}_1^*)' a + f \underline{\omega} - \underline{f} \omega + \frac{1}{4} f \underline{\kappa} - \frac{1}{4} \underline{f} \kappa \right) + \text{Err}(\mu, \mu'), \\
\text{Err}(\mu, \mu') &= -\mathcal{A}'_1 \text{Err}(\zeta, \zeta') - \text{Err}(\rho, \rho') + \frac{1}{4} (\vartheta' \underline{\vartheta}' - \vartheta \underline{\vartheta}).
\end{aligned}$$

The error term $\text{Err}(\mu, \mu')$ is quadratic or higher order with respect to $(f, \underline{f}, a, \check{\Gamma}, \check{R})$ and depends only on at most two angular derivatives e'_θ of f and one angular derivative e'_θ of a, \underline{f} .

Proof. Recall that

$$\mu = -\mathcal{A}_1 \zeta - \rho + \frac{1}{4} \vartheta \underline{\vartheta}.$$

Therefore,

$$\begin{aligned}
\mu' &= -\mathcal{A}'_1 \zeta' - \rho' + \frac{1}{4} \vartheta' \underline{\vartheta}' \\
&= -\mathcal{A}'_1 \left(\zeta - e'_\theta(a) - f \underline{\omega} + \underline{f} \omega - \frac{1}{4} f \underline{\kappa} + \frac{1}{4} \underline{f} \kappa + \text{Err}(\zeta, \zeta') \right) - \rho - \text{Err}(\rho, \rho') + \frac{1}{4} \vartheta' \underline{\vartheta}' \\
&= -\mathcal{A}'_1 \zeta - \rho + \frac{1}{4} \vartheta \underline{\vartheta} - \mathcal{A}'_1 \left((\mathcal{A}_1^*)' a - f \underline{\omega} + \underline{f} \omega - \frac{1}{4} f \underline{\kappa} + \frac{1}{4} \underline{f} \kappa \right) \\
&\quad - \mathcal{A}'_1 \text{Err}(\zeta, \zeta') - \text{Err}(\rho, \rho') + \frac{1}{4} (\vartheta' \underline{\vartheta}' - \vartheta \underline{\vartheta}).
\end{aligned}$$

Note that,

$$\begin{aligned}
-\mathcal{A}'_1 \zeta - \rho + \frac{1}{4} \vartheta \underline{\vartheta} &= -\mathcal{A}_1 \zeta - \rho + \frac{1}{4} \vartheta \underline{\vartheta} + f e_3 \zeta + \underline{f} e_z e + \text{l.o.t.} \\
&= \mu + f e_3 \zeta + \underline{f} e_4 \zeta + \text{l.o.t.}
\end{aligned}$$

Hence,

$$\mu' = \mu + \phi_1' \left(-(\phi_1^*)'a + f\underline{\omega} - \underline{f}\omega + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) + \text{Err}(\mu, \mu')$$

where,

$$\text{Err}(\mu, \mu') = -\phi_1' \text{Err}(\zeta, \zeta') - \text{Err}(\rho, \rho') + \frac{1}{4}(\vartheta'\underline{\vartheta}' - \vartheta\underline{\vartheta}) + fe_3\zeta + \underline{f}e_4\zeta + \text{l.o.t.}$$

In view of the transformation formulas for $\theta, \underline{\theta}$ and the structure of the error terms $\text{Err}(\zeta, \zeta')$, $\text{Err}(\rho, \rho')$, $\text{Err}(\vartheta, \vartheta')$, $\text{Err}(\underline{\vartheta}, \underline{\vartheta}')$ in Lemma 9.3.2 we easily deduce that the error term $\text{Err}(\mu, \mu')$ depends only on at most two angular derivatives e'_θ of f and one angular derivative e'_θ of a, \underline{f} . \square

We shall also make use of the following,

Lemma 9.3.4. *We have the transformation equations,*

$$\begin{aligned} e'_\theta(\kappa') &= e_\theta\kappa + e'_\theta\phi_1'f + \kappa e'_\theta a - \frac{1}{4}\kappa(f\underline{\kappa} + \underline{f}\kappa) + \kappa(f\underline{\omega} - \omega\underline{f}) + f\rho \\ &\quad + \text{Err}(e'_\theta\kappa', e_\theta\kappa), \\ e'_\theta(\underline{\kappa}') &= e_\theta\underline{\kappa} + e'_\theta\phi_1'\underline{f} - \underline{\kappa}e'_\theta a - \frac{1}{4}\underline{\kappa}(f\underline{\kappa} + \underline{f}\kappa) + \underline{\kappa}(f\underline{\omega} - \omega\underline{f}) + \underline{f}\rho \\ &\quad + \text{Err}(e'_\theta\underline{\kappa}', e_\theta\underline{\kappa}), \\ e'_\theta(\mu') &= e_\theta\mu + e'_\theta(\phi_1)' \left(-(\phi_1^*)'a + f\underline{\omega} - \underline{f}\omega + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) + \frac{3}{4}\rho(f\underline{\kappa} + \underline{f}\kappa) \\ &\quad + \text{Err}(e'_\theta\mu', e_\theta\mu), \end{aligned} \tag{9.3.12}$$

where,

$$\begin{aligned} \text{Err}(e'_\theta\kappa', e_\theta\kappa) &= (e^a - 1) \left(e_\theta\kappa + e'_\theta\phi_1'f + \frac{1}{2}\underline{f}e_4\kappa + \frac{1}{2}fe_3\kappa \right) \\ &\quad + e^a \left[e'_\theta \text{Err}(\kappa, \kappa') + e'_\theta(a) \left(\phi_1'f + \text{Err}(\kappa, \kappa') \right) + \frac{1}{2}f\underline{f}e_\theta\kappa + \frac{1}{8}f^2\underline{f}e_3\kappa \right] \\ &\quad + \frac{1}{2}f \left(2\phi_1\eta - \frac{1}{2}\underline{\vartheta}\vartheta + 2(\xi\underline{\xi} + \eta^2) \right) \\ &\quad + \frac{1}{2}f\underline{f}e_\theta\kappa + \frac{1}{8}f^2\underline{f}e_3\kappa + \frac{1}{2}\underline{f} \left(2\phi_1\xi - \frac{1}{2}\vartheta^2 + 2(\eta + \underline{\eta} + 2\underline{\zeta})\xi \right), \end{aligned}$$

$$\begin{aligned}
Err(e'_\theta \underline{\kappa}', e_\theta \underline{\kappa}) &= (e^{-a} - 1) \left(e_\theta \underline{\kappa} + e'_\theta \underline{\mathcal{A}}'_1 f + \frac{1}{2} f e_3 \underline{\kappa} + \frac{1}{2} f e_4 \underline{\kappa} \right) \\
&+ e^{-a} \left[e'_\theta Err(\underline{\kappa}, \underline{\kappa}') + e'_\theta(a) \left(\underline{\mathcal{A}}'_1 f + Err(\underline{\kappa}, \underline{\kappa}') \right) + \frac{1}{2} f f e_\theta \underline{\kappa} + \frac{1}{8} f^2 f e_3 \underline{\kappa} \right] \\
&+ \frac{1}{2} f \left(2 \underline{\mathcal{A}}_1 \underline{\eta} - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} + 2(\underline{\xi} \underline{\xi} + \underline{\eta}^2) \right) \\
&+ \frac{1}{2} f f e_\theta \underline{\kappa} + \frac{1}{8} f^2 f e_3 \underline{\kappa} + \frac{1}{2} f \left(2 \underline{\mathcal{A}}_1 \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 + 2(\underline{\eta} + \underline{\eta} - 2\underline{\zeta}) \underline{\xi} \right),
\end{aligned}$$

and,

$$\begin{aligned}
Err(e'_\theta \underline{\mu}', e_\theta \underline{\mu}) &= e'_\theta Err(\underline{\mu}, \underline{\mu}') + \frac{1}{2} f f e_\theta \underline{\mu} + \frac{1}{8} f^2 f e_3 \underline{\mu} \\
&- \frac{1}{2} f \left(\underline{\mathcal{A}}_1 \underline{\beta} - \frac{1}{2} \underline{\vartheta} \underline{\alpha} - \underline{\zeta} \underline{\beta} + 2(\underline{\eta} \underline{\beta} + \underline{\xi} \underline{\beta}) \right) \\
&- \frac{1}{2} f \left(\underline{\mathcal{A}}_1 \underline{\beta} - \frac{1}{2} \underline{\vartheta} \underline{\alpha} + \underline{\zeta} \underline{\beta} + 2(\underline{\eta} \underline{\beta} + \underline{\xi} \underline{\beta}) \right) \\
&+ \frac{1}{2} f e_4 \left(-\underline{\mathcal{A}}_1 \underline{\zeta} + \frac{1}{4} \underline{\vartheta} \underline{\vartheta} \right) + \frac{1}{2} f e_3 \left(-\underline{\mathcal{A}}_1 \underline{\zeta} + \frac{1}{4} \underline{\vartheta} \underline{\vartheta} \right).
\end{aligned}$$

Proof. Applying the vectorfield e'_θ to

$$\underline{\kappa}' = e^a (\underline{\kappa} + \underline{\mathcal{A}}'_1 f + Err(\underline{\kappa}, \underline{\kappa}'))$$

we deduce,

$$e'_\theta(\underline{\kappa}') = e^a \left(e'_\theta \underline{\kappa} + e'_\theta \underline{\mathcal{A}}'_1 f + e'_\theta(Err(\underline{\kappa}, \underline{\kappa}')) \right) + e^a e'_\theta(a) \left(\underline{\kappa} + \underline{\mathcal{A}}'_1 f + Err(\underline{\kappa}, \underline{\kappa}') \right).$$

Hence,

$$e^{-a} e'_\theta(\underline{\kappa}') = e'_\theta \underline{\kappa} + e'_\theta \underline{\mathcal{A}}'_1 f + e'_\theta(Err(\underline{\kappa}, \underline{\kappa}')) + e'_\theta(a) \left(\underline{\kappa} + \underline{\mathcal{A}}'_1 f + Err(\underline{\kappa}, \underline{\kappa}') \right)$$

and thus

$$e'_\theta(\underline{\kappa}') = e_\theta \underline{\kappa} + e'_\theta(a) \underline{\kappa} + e'_\theta \underline{\mathcal{A}}'_1 f + \frac{1}{2} f e_4 \underline{\kappa} + \frac{1}{2} f e_3 \underline{\kappa} + Err_1[e_\theta(\underline{\kappa}), e'_\theta(\underline{\kappa}')]]$$

with error term,

$$\begin{aligned}
Err_1[e_\theta(\underline{\kappa}), e'_\theta(\underline{\kappa}')] &= (e^a - 1) \left(e_\theta \underline{\kappa} + e'_\theta \underline{\mathcal{A}}'_1 f + \frac{1}{2} f e_4 \underline{\kappa} + \frac{1}{2} f e_3 \underline{\kappa} \right) \\
&+ e^a \left[e'_\theta(Err(\underline{\kappa}, \underline{\kappa}')) + e'_\theta(a) \left(\underline{\mathcal{A}}'_1 f + Err(\underline{\kappa}, \underline{\kappa}') \right) + \frac{1}{2} f f e_\theta \underline{\kappa} + \frac{1}{8} f^2 f e_3 \underline{\kappa} \right].
\end{aligned}$$

Now, making use of

$$\begin{aligned} e'_\theta \kappa &= \left(1 + \frac{1}{2} \underline{f} \underline{f}\right) e_\theta \kappa + \frac{1}{2} \underline{f} e_4 \kappa + \frac{1}{2} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f}\right) e_3 \kappa \\ &= e_\theta \kappa + \frac{1}{2} \underline{f} e_4 \kappa + \frac{1}{2} \underline{f} e_3 \kappa + \frac{1}{2} \underline{f} \underline{f} e_\theta \kappa + \frac{1}{8} \underline{f}^2 \underline{f} e_3 \kappa \end{aligned}$$

and the null structure equations,

$$\begin{aligned} e_3(\kappa) + \frac{1}{2} \underline{\kappa} \kappa - 2 \underline{\omega} \kappa &= 2 \underline{\phi}_1 \eta + 2 \rho - \frac{1}{2} \underline{\vartheta} \vartheta + 2(\underline{\xi} \underline{\xi} + \eta^2), \\ e_4 \kappa + \frac{1}{2} \kappa^2 + 2 \omega \kappa &= 2 \underline{\phi}_1 \xi - \frac{1}{2} \vartheta^2 + 2(\eta + \underline{\eta} + 2 \zeta) \xi, \end{aligned}$$

we deduce,

$$\begin{aligned} e'_\theta \kappa &= e_\theta \kappa + \frac{1}{2} \underline{f} \left(-\frac{1}{2} \kappa^2 - 2 \omega \kappa\right) + \frac{1}{2} \underline{f} \left(-\frac{1}{2} \underline{\kappa} \kappa + 2 \underline{\omega} \kappa + 2 \rho\right) \\ &+ \frac{1}{2} \underline{f} \underline{f} e_\theta \kappa + \frac{1}{8} \underline{f}^2 \underline{f} e_3 \kappa + \frac{1}{2} \underline{f} \left(2 \underline{\phi}_1 \xi - \frac{1}{2} \vartheta^2 + 2(\eta + \underline{\eta} + 2 \zeta) \xi\right) \\ &+ \frac{1}{2} \underline{f} \left(2 \underline{\phi}_1 \eta - \frac{1}{2} \underline{\vartheta} \vartheta + 2(\underline{\xi} \underline{\xi} + \eta^2)\right). \end{aligned}$$

Hence,

$$e'_\theta(\kappa') = e_\theta \kappa + e'_\theta(a) \kappa + e'_\theta \underline{\phi}_1 f + \kappa e'_\theta a - \frac{1}{4} \kappa (f \underline{\kappa} + \underline{f} \kappa) + \kappa (f \underline{\omega} - \underline{\omega} f) + f \rho + \text{Err}(e'_\theta \kappa', e_\theta \kappa)$$

where,

$$\begin{aligned} \text{Err}(e'_\theta \kappa', e_\theta \kappa) &= \text{Err}_1(e'_\theta \kappa', e_\theta \kappa) + \frac{1}{2} \underline{f} \left(2 \underline{\phi}_1 \eta - \frac{1}{2} \underline{\vartheta} \vartheta + 2(\underline{\xi} \underline{\xi} + \eta^2)\right) \\ &+ \frac{1}{2} \underline{f} \underline{f} e_\theta \kappa + \frac{1}{8} \underline{f}^2 \underline{f} e_3 \kappa + \frac{1}{2} \underline{f} \left(2 \underline{\phi}_1 \xi - \frac{1}{2} \vartheta^2 + 2(\eta + \underline{\eta} + 2 \zeta) \xi\right) \end{aligned}$$

as desired. The formula for $e'_\theta(\underline{\kappa}')$ is easily derived by symmetry from the one on $e'_\theta(\kappa')$. Note however that a becomes $-a$ in the transformation.

Applying the operator $e'_\theta = (1 + \frac{1}{2} \underline{f} \underline{f}) e_\theta + \frac{1}{2} \underline{f} e_4 + \frac{1}{2} \underline{f} (1 + \frac{1}{4} \underline{f} \underline{f}) e_3$ to the transformation formula for μ ,

$$\mu' = \mu + (\underline{\phi}_1)' \left(-(\underline{\phi}_1^*)' a + f \underline{\omega} - \underline{\omega} f + \frac{1}{4} f \underline{\kappa} - \frac{1}{4} \underline{f} \kappa\right) + \text{Err}(\mu, \mu')$$

we derive,

$$\begin{aligned}
e'_\theta(\mu') &= e'_\theta(\mu) + e'_\theta(\phi_1)' \left(-(\phi_1^*)'a + f\underline{\omega} - \underline{f}\omega + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) + e'_\theta \text{Err}(\mu, \mu') \\
&= e_\theta(\mu) + \frac{1}{2}\underline{f}e_4\mu + \frac{1}{2}fe_3\mu + e'_\theta(\phi_1)' \left(-(\phi_1^*)'a + f\underline{\omega} - \underline{f}\omega + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) \\
&\quad + e'_\theta \text{Err}(\mu, \mu') + \frac{1}{2}ff\underline{e}_\theta\mu + \frac{1}{8}f^2\underline{f}e_3\mu.
\end{aligned}$$

Recalling that $\mu = -\phi_1\zeta - \rho + \frac{1}{4}\vartheta\underline{\vartheta}$ we find,

$$\frac{1}{2}\underline{f}e_4\mu + \frac{1}{2}fe_3\mu = -\frac{1}{2}(fe_3 + \underline{f}e_4)\rho + \frac{1}{2}\underline{f}e_4 \left(-\phi_1\zeta + \frac{1}{4}\vartheta\underline{\vartheta} \right) + \frac{1}{2}fe_3 \left(-\phi_1\zeta + \frac{1}{4}\vartheta\underline{\vartheta} \right).$$

Recalling the Bianchi equations for $e_3\rho, e_4\rho$

$$\begin{aligned}
e_4\rho + \frac{3}{2}\kappa\rho &= \phi_1\underline{\beta} - \frac{1}{2}\underline{\vartheta}\alpha + \zeta\underline{\beta} + 2(\eta\underline{\beta} + \xi\underline{\beta}), \\
e_3\rho + \frac{3}{2}\underline{\kappa}\rho &= \phi_1\underline{\beta} - \frac{1}{2}\vartheta\underline{\alpha} - \zeta\underline{\beta} + 2(\eta\underline{\beta} + \xi\underline{\beta}),
\end{aligned}$$

we further deduce,

$$\begin{aligned}
\frac{1}{2}\underline{f}e_4\mu + \frac{1}{2}fe_3\mu &= \frac{3}{4}\rho(f\underline{\kappa} + \underline{f}\kappa) \\
&\quad - \frac{1}{2}f \left(\phi_1\underline{\beta} - \frac{1}{2}\underline{\vartheta}\alpha - \zeta\underline{\beta} + 2(\eta\underline{\beta} + \xi\underline{\beta}) \right) \\
&\quad - \frac{1}{2}\underline{f} \left(\phi_1\underline{\beta} - \frac{1}{2}\vartheta\underline{\alpha} + \zeta\underline{\beta} + 2(\eta\underline{\beta} + \xi\underline{\beta}) \right) \\
&\quad + \frac{1}{2}\underline{f}e_4 \left(-\phi_1\zeta + \frac{1}{4}\vartheta\underline{\vartheta} \right) + \frac{1}{2}fe_3 \left(-\phi_1\zeta + \frac{1}{4}\vartheta\underline{\vartheta} \right).
\end{aligned}$$

Therefore,

$$\begin{aligned}
e'_\theta(\mu') &= e_\theta(\mu) + \frac{3}{4}\rho(f\underline{\kappa} + \underline{f}\kappa) + e'_\theta(\phi_1)' \left(-(\phi_1^*)'a + f\underline{\omega} - \underline{f}\omega + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) \\
&\quad + \text{Err}(e_\theta\mu, e_\theta\mu)
\end{aligned}$$

with,

$$\begin{aligned}
\text{Err}(e'_\theta\mu', e_\theta\mu) &= e'_\theta \text{Err}(\mu, \mu') + \frac{1}{2}ff\underline{e}_\theta\mu + \frac{1}{8}f^2\underline{f}e_3\mu \\
&\quad - \frac{1}{2}f \left(\phi_1\underline{\beta} - \frac{1}{2}\underline{\vartheta}\alpha - \zeta\underline{\beta} + 2(\eta\underline{\beta} + \xi\underline{\beta}) \right) \\
&\quad - \frac{1}{2}\underline{f} \left(\phi_1\underline{\beta} - \frac{1}{2}\vartheta\underline{\alpha} + \zeta\underline{\beta} + 2(\eta\underline{\beta} + \xi\underline{\beta}) \right) \\
&\quad + \frac{1}{2}\underline{f}e_4 \left(-\phi_1\zeta + \frac{1}{4}\vartheta\underline{\vartheta} \right) + \frac{1}{2}fe_3 \left(-\phi_1\zeta + \frac{1}{4}\vartheta\underline{\vartheta} \right)
\end{aligned}$$

as desired. \square

Finally recalling the definition of the Hodge operators $\not{d}_1, \not{d}_1^*, (\not{d}_1)', (\not{d}_1)'$ and noticing that

$$\begin{aligned} (\not{d}_1^*)'(\kappa') &= (\not{d}_1^*)'(\check{\kappa}'), & (\not{d}_1^*)(\kappa) &= (\not{d}_1^*)(\check{\kappa}), \\ (\not{d}_1^*)'(\underline{\kappa}') &= (\not{d}_1^*)'(\check{\underline{\kappa}}'), & (\not{d}_1^*)(\underline{\kappa}) &= (\not{d}_1^*)(\check{\underline{\kappa}}), \\ (\not{d}_1^*)'(\mu') &= (\not{d}_1^*)'(\check{\mu}'), & (\not{d}_1^*)(\mu) &= (\not{d}_1^*)(\check{\mu}), \\ (\not{d}_1^*)'(\underline{\mu}') &= (\not{d}_1^*)'(\check{\underline{\mu}}'), & (\not{d}_1^*)(\underline{\mu}) &= (\not{d}_1^*)(\check{\underline{\mu}}), \end{aligned}$$

we recast the results of Lemma 9.3.4 in the following form.

Lemma 9.3.5. *We have the transformation equations,*

$$\begin{aligned} (\not{d}_1^*)'(\check{\kappa}') &= \not{d}_1^*(\check{\kappa}) + (\not{d}_1^*)'(\not{d}_1)'f + \kappa(\not{d}_1^*)'a - \rho f + \frac{1}{4}\kappa(f\underline{\kappa} + \underline{f}\kappa) \\ &\quad - \kappa(f\underline{\omega} - \underline{f}\omega) - Err_1, \\ (\not{d}_1^*)'(\check{\underline{\kappa}}') &= \not{d}_1^*(\check{\underline{\kappa}}) + (\not{d}_1^*)'(\not{d}_1)'\underline{f} - \underline{\kappa}(\not{d}_1^*)'a - \rho\underline{f} + \frac{1}{4}\underline{\kappa}(f\underline{\kappa} + \underline{f}\kappa) \\ &\quad - \underline{\kappa}(f\underline{\omega} - \underline{f}\omega) - Err_2, \\ (\not{d}_1^*)'(\check{\mu}') &= \not{d}_1^*(\check{\mu}) + (\not{d}_1^*)'(\not{d}_1)' \left(-(\not{d}_1^*)'a + f\underline{\omega} - \underline{f}\omega + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) \\ &\quad - \frac{3}{4}\rho(f\underline{\kappa} + \underline{f}\kappa) - Err_3, \end{aligned} \tag{9.3.13}$$

where,

$$\begin{aligned} Err_1 &= Err(e'_\theta \kappa', e_\theta \kappa) = e'_\theta Err(\kappa, \kappa') + ae'_\theta \not{d}'_1 f + e'_\theta(a) \not{d}'_1 f \\ &\quad + a \left(e_\theta \kappa + \frac{1}{2}(f e_{4\kappa} + f e_{3\kappa}) \right) + f \not{d}_1 \eta + \underline{f} \not{d}_1 \xi + l.o.t., \\ Err_2 &= Err(e'_\theta \underline{\kappa}', e_\theta \underline{\kappa}) = e'_\theta Err(\underline{\kappa}, \underline{\kappa}') - ae'_\theta \not{d}'_1 \underline{f} - e'_\theta(a) \not{d}'_1 \underline{f} \\ &\quad - a \left(e_\theta \underline{\kappa} + \frac{1}{2}(\underline{f} e_{4\underline{\kappa}} + \underline{f} e_{3\underline{\kappa}}) \right) + \underline{f} \not{d}_1 \eta + f \not{d}_1 \underline{\xi} + l.o.t., \\ Err_3 &= Err(e'_\theta \mu', e_\theta \mu) = e'_\theta Err(\mu, \mu') - \frac{1}{2}(\underline{f} \not{d}_1 \beta + f \not{d}_1 \underline{\beta}) - \frac{1}{2}(f e_3 + \underline{f} e_4) \not{d}_1 \zeta + l.o.t., \end{aligned} \tag{9.3.14}$$

where the terms denoted by *l.o.t.* are cubic or higher order in $a, f, \underline{f}, \check{\Gamma}, \check{R}$ and contain no derivatives of (a, f, \underline{f}) .

9.3.1 Main GCM Equations

Given a deformation $\Psi : \mathring{S} \rightarrow \mathbf{S}$ and adapted frame (e'_3, e'_4, e'_θ) with $e'_\theta = e_\theta^{\mathbf{S}}$ we derive an elliptic system for the transition parameters (a, f, \underline{f}) . The system will later be used in the construction of GCM surfaces.

In what follows we denote by $\not{d}_1^{\mathbf{S}}, \not{d}_2^{\mathbf{S}}, \not{d}_1^{\mathbf{S},*}, \not{d}_2^{\mathbf{S},*}$ the basic Hodge operators on \mathbf{S} . Noting that the transformation formulae in (9.3.13)–(9.3.14) contain only the operators $(\not{d}_1)^{\mathbf{S}}, (\not{d}_1^*)^{\mathbf{S}} = \not{d}_1^{\mathbf{S},*}$ applied to a, f, \underline{f} we introduce the simplified notation,

$$\not{d}^{\mathbf{S}} := (\not{d}_1)^{\mathbf{S}}, \quad \not{d}^{\mathbf{S},*} := (\not{d}_1^*)^{\mathbf{S}}, \quad A^{\mathbf{S}} := \not{d}^{\mathbf{S},*} \not{d}^{\mathbf{S}}, \quad \not{d}^* := \not{d}_1^*. \quad (9.3.15)$$

With these notation (9.3.13) takes the following form,

$$\begin{aligned} \not{d}^{\mathbf{S},*} \check{\kappa}^{\mathbf{S}} &= \not{d} \check{\kappa} + A^{\mathbf{S}} f + \kappa \not{d}^{\mathbf{S},*} a - \rho f + \frac{1}{4} \kappa (f \underline{\kappa} + \underline{f} \kappa) - \kappa (f \underline{\omega} - \underline{f} \omega) - \text{Err}_1, \\ \not{d}^{\mathbf{S},*} \check{\underline{\kappa}}^{\mathbf{S}} &= \not{d} \check{\underline{\kappa}} + A^{\mathbf{S}} \underline{f} - \underline{\kappa} \not{d}^{\mathbf{S},*} a - \rho \underline{f} + \frac{1}{4} \underline{\kappa} (f \underline{\kappa} + \underline{f} \kappa) - \underline{\kappa} (\underline{f} \omega - f \underline{\omega}) - \text{Err}_2, \\ \not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}} &= \not{d}^* \check{\mu} + A^{\mathbf{S}} \left(-\not{d}^{\mathbf{S},*} a + f \underline{\omega} - \underline{f} \omega + \frac{1}{4} f \underline{\kappa} - \frac{1}{4} \underline{f} \kappa \right) - \frac{3}{4} \rho (f \underline{\kappa} + \underline{f} \kappa) - \text{Err}_3, \end{aligned}$$

or,

$$\begin{aligned} A^{\mathbf{S}} \left(-\not{d}^{\mathbf{S},*} a + f \underline{\omega} - \underline{f} \omega + \frac{1}{4} f \underline{\kappa} - \frac{1}{4} \underline{f} \kappa \right) - \frac{3}{4} \rho (\kappa \underline{f} + \underline{\kappa} f) &= \not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}} - \not{d}^* \check{\mu} + \text{Err}_3, \\ A^{\mathbf{S}} f + \kappa \not{d}^{\mathbf{S},*} a - \rho f + \frac{1}{4} \kappa (f \underline{\kappa} + \underline{f} \kappa) - \kappa (f \underline{\omega} - \underline{f} \omega) &= \not{d}^{\mathbf{S},*} \check{\kappa}^{\mathbf{S}} - \not{d}^* \check{\kappa} + \text{Err}_1, \\ A^{\mathbf{S}} \underline{f} - \underline{\kappa} \not{d}^{\mathbf{S},*} a - \rho \underline{f} + \frac{1}{4} \underline{\kappa} (f \underline{\kappa} + \underline{f} \kappa) - \underline{\kappa} (\underline{f} \omega - f \underline{\omega}) &= \not{d}^{\mathbf{S},*} \check{\underline{\kappa}}^{\mathbf{S}} - \not{d}^* \check{\underline{\kappa}} + \text{Err}_2. \end{aligned} \quad (9.3.16)$$

Since $A^{\mathbf{S}}$ is invertible⁷ we can write, setting $z := \kappa \underline{f} + \underline{\kappa} f$,

$$\not{d}^{\mathbf{S},*} a = f \underline{\omega} - \underline{f} \omega + \frac{1}{4} f \underline{\kappa} - \frac{1}{4} \underline{f} \kappa - \frac{3}{4} (A^{\mathbf{S}})^{-1} (\rho z) + (A^{\mathbf{S}})^{-1} (-\not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}} + \not{d}^* \check{\mu} - \text{Err}_3).$$

⁷We have $\int_{\mathbf{S}} f A^{\mathbf{S}} f = \int_{\mathbf{S}} (\not{d}^{\mathbf{S}} f)^2$ which in view of the identity (2.1.22) for $\not{d}_1^{\mathbf{S}}$ and the definition of $\not{d}^{\mathbf{S}}$ implies that $A^{\mathbf{S}}$ is invertible.

We can thus eliminate $\mathcal{A}^{\mathbf{S},*}a$ from the last two equations,

$$\begin{aligned} A^{\mathbf{S}}f + \left(\frac{1}{2}\kappa\underline{\kappa} - \rho\right)f - \frac{3}{4}\kappa(A^{\mathbf{S}})^{-1}(\rho z) &= \mathcal{A}^{\mathbf{S},*\check{\kappa}}^{\mathbf{S}} - \mathcal{A}^{\check{\kappa}} - \kappa(A^{\mathbf{S}})^{-1}(-\mathcal{A}^{\mathbf{S},*\check{\mu}}^{\mathbf{S}} + \mathcal{A}^{\check{\mu}}) \\ &+ \text{Err}_4, \end{aligned}$$

$$\begin{aligned} A^{\mathbf{S}}\underline{f} + \left(\frac{1}{2}\kappa\underline{\kappa} - \rho\right)\underline{f} + \frac{3}{4}\underline{\kappa}(A^{\mathbf{S}})^{-1}(\rho z) &= \mathcal{A}^{\mathbf{S},*\check{\underline{\kappa}}}^{\mathbf{S}} - \mathcal{A}^{\check{\underline{\kappa}}} + \underline{\kappa}(A^{\mathbf{S}})^{-1}(-\mathcal{A}^{\mathbf{S},*\check{\underline{\mu}}}^{\mathbf{S}} + \mathcal{A}^{\check{\underline{\mu}}}) \\ &+ \text{Err}_5, \end{aligned}$$

where,

$$\text{Err}_4 = \text{Err}_1 + \kappa(A^{\mathbf{S}})^{-1}\text{Err}_3, \quad \text{Err}_5 = \text{Err}_2 - \underline{\kappa}(A^{\mathbf{S}})^{-1}\text{Err}_3.$$

Therefore the system (9.3.16) is equivalent to the system,

$$\begin{aligned} A^{\mathbf{S}}f + \left(\frac{1}{2}\kappa\underline{\kappa} - \rho\right)f - \frac{3}{4}\kappa(A^{\mathbf{S}})^{-1}(\rho z) &= \mathcal{A}^{\mathbf{S},*\check{\kappa}}^{\mathbf{S}} - \mathcal{A}^{\check{\kappa}} - \kappa(A^{\mathbf{S}})^{-1}(-\mathcal{A}^{\mathbf{S},*\check{\mu}}^{\mathbf{S}} + \mathcal{A}^{\check{\mu}}) \\ &+ \text{Err}_4, \end{aligned}$$

$$\begin{aligned} A^{\mathbf{S}}\underline{f} + \left(\frac{1}{2}\kappa\underline{\kappa} - \rho\right)\underline{f} + \frac{3}{4}\underline{\kappa}(A^{\mathbf{S}})^{-1}(\rho z) &= \mathcal{A}^{\mathbf{S},*\check{\underline{\kappa}}}^{\mathbf{S}} - \mathcal{A}^{\check{\underline{\kappa}}} + \underline{\kappa}(A^{\mathbf{S}})^{-1}(-\mathcal{A}^{\mathbf{S},*\check{\underline{\mu}}}^{\mathbf{S}} + \mathcal{A}^{\check{\underline{\mu}}}) \\ &+ \text{Err}_5, \end{aligned}$$

$$\mathcal{A}^{\mathbf{S},*}a + \frac{3}{4}(A^{\mathbf{S}})^{-1}(\rho z) - \underline{f}\omega + \underline{f}\omega - \frac{1}{4}\underline{f}\underline{\kappa} + \frac{1}{4}\underline{f}\kappa = (A^{\mathbf{S}})^{-1}(-\mathcal{A}^{\mathbf{S},*\check{\mu}}^{\mathbf{S}} + \mathcal{A}^{\check{\mu}}) - (A^{\mathbf{S}})^{-1}\text{Err}_3.$$

Furthermore, we have

$$A^{\mathbf{S}}z = A^{\mathbf{S}}(\underline{\kappa}f + \kappa f) = \underline{\kappa}A^{\mathbf{S}}f + \kappa A^{\mathbf{S}}\underline{f} + [A^{\mathbf{S}}, \underline{\kappa}]f + [A^{\mathbf{S}}, \kappa]\underline{f}.$$

In view of the above equations for $A^{\mathbf{S}}f$ and $A^{\mathbf{S}}\underline{f}$, we infer

$$\begin{aligned} A^{\mathbf{S}}z &= \underline{\kappa} \left\{ -\left(\frac{1}{2}\kappa\underline{\kappa} - \rho\right)f + \frac{3}{4}\kappa(A^{\mathbf{S}})^{-1}(\rho z) + \mathcal{A}^{\mathbf{S},*\check{\kappa}}^{\mathbf{S}} - \mathcal{A}^{\check{\kappa}} - \kappa(A^{\mathbf{S}})^{-1}(-\mathcal{A}^{\mathbf{S},*\check{\mu}}^{\mathbf{S}} + \mathcal{A}^{\check{\mu}}) \right\} \\ &+ \kappa \left\{ -\left(\frac{1}{2}\kappa\underline{\kappa} - \rho\right)\underline{f} - \frac{3}{4}\underline{\kappa}(A^{\mathbf{S}})^{-1}(\rho z) + \mathcal{A}^{\mathbf{S},*\check{\underline{\kappa}}}^{\mathbf{S}} - \mathcal{A}^{\check{\underline{\kappa}}} + \underline{\kappa}(A^{\mathbf{S}})^{-1}(-\mathcal{A}^{\mathbf{S},*\check{\underline{\mu}}}^{\mathbf{S}} + \mathcal{A}^{\check{\underline{\mu}}}) \right\} \\ &+ [A^{\mathbf{S}}, \underline{\kappa}]f + [A^{\mathbf{S}}, \kappa]\underline{f} + \underline{\kappa}\text{Err}_4 + \kappa\text{Err}_5. \end{aligned}$$

Furthermore

$$\begin{aligned} A^{\mathbf{S}}z &= -\left(\frac{1}{2}\kappa\underline{\kappa} - \rho\right)z + \underline{\kappa} \left\{ \mathcal{A}^{\mathbf{S},*\check{\kappa}}^{\mathbf{S}} - \mathcal{A}^{\check{\kappa}} - \kappa(A^{\mathbf{S}})^{-1}(-\mathcal{A}^{\mathbf{S},*\check{\mu}}^{\mathbf{S}} + \mathcal{A}^{\check{\mu}}) \right\} \\ &+ \kappa \left\{ \mathcal{A}^{\mathbf{S},*\check{\underline{\kappa}}}^{\mathbf{S}} - \mathcal{A}^{\check{\underline{\kappa}}} + \underline{\kappa}(A^{\mathbf{S}})^{-1}(-\mathcal{A}^{\mathbf{S},*\check{\underline{\mu}}}^{\mathbf{S}} + \mathcal{A}^{\check{\underline{\mu}}}) \right\} \\ &+ [A^{\mathbf{S}}, \underline{\kappa}]f + [A^{\mathbf{S}}, \kappa]\underline{f} + \underline{\kappa}\text{Err}_4 + \kappa\text{Err}_5 \\ &= -\left(\frac{1}{2}\kappa\underline{\kappa} - \rho\right)z + \underline{\kappa}\mathcal{A}^{\mathbf{S},*\check{\kappa}}^{\mathbf{S}} + \kappa\mathcal{A}^{\mathbf{S},*\check{\underline{\kappa}}}^{\mathbf{S}} - \underline{\kappa}\mathcal{A}^{\check{\kappa}} - \kappa\mathcal{A}^{\check{\underline{\kappa}}} \\ &+ \underline{\kappa}\text{Err}_4 + \kappa\text{Err}_5 + [A^{\mathbf{S}}, \underline{\kappa}]f + [A^{\mathbf{S}}, \kappa]\underline{f}. \end{aligned}$$

We summarize the results of the above calculation in the following lemma.

Lemma 9.3.6. *The original system (9.3.13) in (a, f, \underline{f}) associated to a deformation sphere \mathbf{S} is equivalent to the following*

$$\begin{aligned}
(A^{\mathbf{S}} + V)z &= \underline{\kappa} \not{d}^{\mathbf{S},*} \check{\kappa}^{\mathbf{S}} + \kappa \not{d}^{\mathbf{S},*} \check{\underline{\kappa}}^{\mathbf{S}} - \underline{\kappa} \not{d}^* \check{\kappa} - \kappa \not{d}^* \check{\underline{\kappa}} \\
&\quad + \underline{\kappa} \text{Err}_4 + \kappa \text{Err}_5 + [A^{\mathbf{S}}, \underline{\kappa}]f + [A^{\mathbf{S}}, \kappa] \underline{f}, \\
(A^{\mathbf{S}} + V)f &= \frac{3}{4} \kappa (A^{\mathbf{S}})^{-1}(\rho z) + \not{d}^{\mathbf{S},*} \check{\kappa}^{\mathbf{S}} - \not{d}^* \check{\kappa} - \kappa (A^{\mathbf{S}})^{-1}(-\not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}} + \not{d}^* \check{\mu}) \\
&\quad + \text{Err}_4, \\
(A^{\mathbf{S}} + V)\underline{f} &= -\frac{3}{4} \underline{\kappa} (A^{\mathbf{S}})^{-1}(\rho z) + \not{d}^{\mathbf{S},*} \check{\underline{\kappa}}^{\mathbf{S}} - \not{d}^* \check{\underline{\kappa}} + \underline{\kappa} (A^{\mathbf{S}})^{-1}(-\not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}} + \not{d}^* \check{\mu}) \\
&\quad + \text{Err}_5, \\
\not{d}^{\mathbf{S},*} a &= -\frac{3}{4} (A^{\mathbf{S}})^{-1}(\rho z) + f \underline{\omega} - \underline{f} \omega + \frac{1}{4} f \underline{\kappa} - \frac{1}{4} \underline{f} \kappa + (A^{\mathbf{S}})^{-1}(-\not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}} + \not{d}^* \check{\mu}) \\
&\quad - (A^{\mathbf{S}})^{-1} \text{Err}_3,
\end{aligned} \tag{9.3.17}$$

where,

$$z := \underline{\kappa} f + \kappa \underline{f}, \quad V := -\frac{1}{2} \kappa \underline{\kappa} + \rho. \tag{9.3.18}$$

The error terms are given by $\text{Err}_1, \text{Err}_2, \text{Err}_3$, defined in Lemma 9.3.5, and

$$\text{Err}_4 = \text{Err}_1 + \kappa (A^{\mathbf{S}})^{-1} \text{Err}_3, \quad \text{Err}_5 = \text{Err}_2 - \underline{\kappa} (A^{\mathbf{S}})^{-1} \text{Err}_3. \tag{9.3.19}$$

Remark 9.3.7. *We note the following remarks concerning the system (9.3.17).*

1. *The right hand side of the equations is linear in the quantities,*

$$\not{d}^{\mathbf{S},*} \check{\kappa}^{\mathbf{S}}, \quad \not{d}^{\mathbf{S},*} \check{\underline{\kappa}}^{\mathbf{S}}, \quad \not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}}, \quad \text{as well as} \quad \not{d}^* \check{\kappa}, \quad \not{d}^* \check{\underline{\kappa}}, \quad \not{d}^* \check{\mu}.$$

The first group is to be constrained by our GCM conditions in the next section while the second group depends on assumptions regarding the background foliation of \mathcal{R} .

2. *The error terms contain only \mathbf{S} -angular derivatives of (a, f, \underline{f}) of order at most equal to the order of the corresponding operators on the left hand sides, see Lemma 9.3.8 below. Thus the system is in a standard quasilinear elliptic system form.*
3. *In order to uniquely solve the equations for z , and then f and \underline{f} , we need to the coercivity of the operator $A^{\mathbf{S}} + V$. One can easily show that the potential V is positive for small values of r , i.e. r near $r_{\mathcal{H}} = 2m_0(1 + \delta_{\mathcal{H}})$ but negative for large r . In fact*

$A^{\mathbf{S}} + V$ has a nontrivial kernel for large r as one can easily see from the following calculation. Since,

$$A^{\mathbf{S}} = \phi_1^* \phi_1 = \phi_2^{\mathbf{S}} \phi_2^{\mathbf{S}} + 2K, \quad K = -\rho - \frac{1}{4}\kappa\kappa + \frac{1}{4}\vartheta\vartheta$$

we deduce,

$$A^{\mathbf{S}} + V = A^{\mathbf{S}} + \frac{1}{2}\kappa\kappa - \rho = \phi_2^{\mathbf{S}} \phi_2^{\mathbf{S}} - 3\rho + \frac{1}{2}\vartheta\vartheta.$$

Thus for large enough r the operator $A^{\mathbf{S}} + V$ behaves like $\phi_2^{\mathbf{S}} \phi_2^{\mathbf{S},*}$ which has a nontrivial kernel.

4. To be able to correct for the lack of coercivity of the system we need to control the $\ell = 1$ modes of $f, \underline{f}, \phi^{\mathbf{S},*}a$. In subsection 9.3.4 we derive an equation for the last. The $\ell = 1$ modes of (f, \underline{f}) , on the other hand, have to be prescribed.
5. The equations do not provide information on the average of a . For this we will need yet another equation derived in subsection 9.3.2.

Lemma 9.3.8. *The error terms Err_1, \dots, Err_5 can be written schematically as follows,*

$$\begin{aligned} r^2 Err_1 &= (\vartheta^{\mathbf{S}})^2 ((f, \underline{f}, a)^2) + \vartheta^{\mathbf{S}} ((f, \underline{f}, a)(r\check{\Gamma}_g)), \\ r Err_2 &= r^{-1} (\vartheta^{\mathbf{S}})^2 ((f, \underline{f}, a)^2) + \vartheta^{\mathbf{S}} ((f, \underline{f}, a)(\check{\Gamma})), \\ r^3 Err_3 &= (\vartheta^{\mathbf{S}})^3 ((f, \underline{f}, a)^2) + (\vartheta^{\mathbf{S}})^2 ((f, \underline{f}, a)(r\check{\Gamma}_g)), \\ Err_4, Err_5 &= Err_1 + (A^{\mathbf{S}})^{-1} Err_3, \end{aligned} \tag{9.3.20}$$

where the lower order terms denoted l.o.t. are cubic with respect to $a, f, \underline{f}, \check{\Gamma}, \check{R}$ and may involve fewer angular (along \mathbf{S}) derivatives of a, f, \underline{f} .

Remark 9.3.9. *Note that Err_2 behaves worse in powers of r than Err_1 . The reason is the presence of the terms $f e_{\theta} \underline{\xi}, e_{\theta}(f \underline{\xi})$ in the formula for $e_{\theta}^{\mathbf{S}}(Err(\underline{\kappa}', \underline{\kappa}))$.*

Proof. Note that in the spacetime region \mathcal{R} of interest r and $r^{\mathbf{S}}$ are comparable. Recall, see (9.3.14),

$$\begin{aligned} Err_1 &= Err(e'_{\theta} \kappa', e_{\theta} \kappa) = e'_{\theta} Err(\kappa, \kappa') + a e'_{\theta} \phi'_1 f + e'_{\theta}(a) \phi'_1 f \\ &\quad + a \left(e_{\theta} \kappa + \frac{1}{2} (\underline{f} e_4 \kappa + f e_3 \kappa) \right) + f \phi_1 \eta + \underline{f} \phi_1 \xi + \text{l.o.t.}, \\ Err_2 &= Err(e'_{\theta} \underline{\kappa}', e_{\theta} \underline{\kappa}) = e'_{\theta} Err(\underline{\kappa}, \underline{\kappa}') - a e'_{\theta} \phi'_1 \underline{f} - e'_{\theta}(a) \phi'_1 \underline{f} \\ &\quad - a \left(e_{\theta} \underline{\kappa} + \frac{1}{2} (\underline{f} e_4 \underline{\kappa} + f e_3 \underline{\kappa}) \right) + \underline{f} \phi_1 \eta + f \phi_1 \xi + \text{l.o.t.}, \\ Err_3 &= Err(e'_{\theta} \mu', e_{\theta} \mu) = e'_{\theta} Err(\mu, \mu') - \frac{1}{2} (\underline{f} \phi_1 \beta + f \phi_1 \underline{\beta}) - \frac{1}{2} (f e_3 + \underline{f} e_4) \phi_1 \zeta + \text{l.o.t.}, \end{aligned}$$

and⁸,

$$\begin{aligned} \text{Err}(\kappa, \kappa') &= \frac{1}{2} \underline{f} \underline{f} e'_\theta(f) - \frac{1}{4} \underline{f} e'_\theta(f^2) + f(\zeta + \eta) + \underline{f} \xi - \frac{1}{4} f^2 \underline{\kappa} + \underline{f} \underline{f} \omega - f^2 \underline{\omega} + \text{l.o.t.}, \\ \text{Err}(\underline{\kappa}, \underline{\kappa}') &= -\frac{1}{2} \underline{f} e'_\theta \left(\underline{f} \underline{f} + \frac{1}{8} f^2 \underline{f} \right) + \left(\frac{3}{4} \underline{f} \underline{f} + \frac{1}{8} (f \underline{f})^2 \right) e'_\theta(\underline{f}) \\ &\quad + \frac{1}{4} \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) \underline{f} e'_\theta(f \underline{f}) - \frac{1}{4} f \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_\theta(f^2) \\ &\quad + \underline{f}(-\zeta + \eta) + \underline{f} \xi - \frac{1}{4} f^2 \underline{\kappa} + \underline{f} \underline{f} \omega - f^2 \underline{\omega} + \text{l.o.t.} \end{aligned}$$

Also,

$$\begin{aligned} \text{Err}(\mu, \mu') &= -e'_\theta \text{Err}(\zeta, \zeta') - \text{Err}(\rho, \rho') + \frac{1}{4} (\vartheta' \underline{\vartheta}' - \vartheta \underline{\vartheta}), \\ \text{Err}(\zeta, \zeta') &= \frac{1}{2} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_\theta(f) - \frac{1}{16} f^2 e'_\theta(f^2) + \frac{1}{4} (-f \underline{\vartheta} + \underline{f} \vartheta) + \text{l.o.t.}, \\ \text{Err}(\rho, \rho') &= \frac{3}{2} \rho \underline{f} \underline{f} + \underline{f} \beta + f \underline{\beta} + \text{l.o.t.} \end{aligned}$$

We write schematically⁹,

$$\begin{aligned} \text{Err}_1 &= (f, \underline{f}, a) (r^{-2} \mathfrak{P}^{\mathbf{S}})^2 (f, \underline{f}, a) + (r^{-1} \mathfrak{P}^{\mathbf{S}}(f, \underline{f}, a))^2 + r^{-1} \mathfrak{P}^{\mathbf{S}}((f, \underline{f}, a) \check{\Gamma}_g) \\ &\quad + r^{-2} \mathfrak{P}^{\mathbf{S}}(f^2) + \frac{1}{2} a (\underline{f} e_4 \kappa + f e_3 \kappa) + \text{l.o.t.} \end{aligned}$$

Making use of

$$\begin{aligned} e_3(\kappa) + \frac{1}{2} \underline{\kappa} \kappa - 2 \underline{\omega} \kappa &= 2 \mathfrak{A}_1 \eta + 2 \rho - \frac{1}{2} \underline{\vartheta} \vartheta + 2(\xi \underline{\xi} + \eta^2), \\ e_4 \kappa + \frac{1}{2} \kappa^2 + 2 \omega \kappa &= 2 \mathfrak{A}_1 \xi - \frac{1}{2} \vartheta^2 + 2(\eta + \underline{\eta} + 2\zeta) \xi, \end{aligned}$$

and treating the curvature terms that appear as $\check{\Gamma}_g$ we easily derive,

$$r^2 \text{Err}_1 = (\mathfrak{P}^{\mathbf{S}})^2 ((f, \underline{f}, a)^2) + \mathfrak{P}^{\mathbf{S}}((f, \underline{f}, a) (r \check{\Gamma}_g)).$$

We obtain a worse estimate for Err_2 because of the presence $e'_\theta(f \underline{\xi})$, since $\underline{\xi} \in \check{\Gamma}_b$. In fact,

$$r \text{Err}_2 = r^{-1} (\mathfrak{P}^{\mathbf{S}})^2 ((f, \underline{f}, a)^2) + \mathfrak{P}^{\mathbf{S}}((f, \underline{f}, a) \check{\Gamma}).$$

⁸Recall also the outgoing geodesic conditions i.e. $\xi = 0$, $\zeta + \underline{\eta} = 0$, $\zeta - \eta = 0$, $\omega = 0$.

⁹The last term $r^{-2} \mathfrak{P}^{\mathbf{S}}(f^2)$ on the right of the identity below is due to the term $e'_\theta(f^2 \underline{\omega})$ in the expression of $e'_\theta \text{Err}(\kappa, \kappa')$.

For Err_3 we write similarly, treating the curvature terms that appear as $\check{\Gamma}_g$,

$$e_\theta(\mu, \mu') = r^{-3}(\vartheta^{\mathbf{S}})^3 ((f, \underline{f}, a)^2) + r^{-3}(\vartheta^{\mathbf{S}})^2 ((f, \underline{f}, a)\check{\Gamma}_g) + \text{l.o.t.}$$

Using the null structure equations for ζ we infer that,

$$\begin{aligned} \text{Err}_3 &= e_\theta(\mu, \mu') - \frac{1}{2}(\underline{f}\phi_1\beta + f\phi_1\underline{\beta}) - \frac{1}{2}(fe_3 + \underline{f}e_4)\phi_1\zeta + \text{l.o.t.} \\ &= r^{-3}(\vartheta^{\mathbf{S}})^3 ((f, \underline{f}, a)^2) + r^{-3}(\vartheta^{\mathbf{S}})^2 ((f, \underline{f}, a)\check{\Gamma}_g) + \text{l.o.t.} \end{aligned}$$

as stated. \square

Making use of the above lemma and the assumptions **A1-A3** we can derive the following.

Lemma 9.3.10. *Assume given a deformation $\Psi : \mathring{S} \rightarrow \mathbf{S}$ in \mathcal{R} and adapted frame (e'_3, e'_4, e'_θ) with $e'_\theta = e_\theta^{\mathbf{S}}$ with transition parameters a, f, \underline{f} defined on \mathbf{S} . Assume that¹⁰ the following holds true*

$$\|(U', S')\|_{L^\infty(\mathring{S})} + \max_{0 \leq s \leq s_{max}} (\mathring{r})^{-1} \|(U', S')\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})} \lesssim \mathring{\epsilon}.$$

Then, for $s = s_{max} + 1$, with $s_{max} \geq 4$,

$$\begin{aligned} \|\text{Err}_1, \text{Err}_2\|_{\mathfrak{h}_{s-2}(\mathbf{S})} &\lesssim r^{-2} \left\| (f, \underline{f}, a) \right\|_{\mathfrak{h}_s(\mathbf{S})} \left(\mathring{\epsilon} + \left\| f, \underline{f}, a \right\|_{\mathfrak{h}_s(\mathbf{S})} \right), \\ \|\text{Err}_3\|_{\mathfrak{h}_{s-3}(\mathbf{S})} &\lesssim r^{-3} \left\| (f, \underline{f}, a) \right\|_{\mathfrak{h}_s(\mathbf{S})} \left(\mathring{\epsilon} + \left\| f, \underline{f}, a \right\|_{\mathfrak{h}_s(\mathbf{S})} \right), \\ \|\text{Err}_4, \text{Err}_5\|_{\mathfrak{h}_{s-2}(\mathbf{S})} &\lesssim r^{-2} \left\| (f, \underline{f}, a) \right\|_{\mathfrak{h}_s(\mathbf{S})} \left(\mathring{\epsilon} + \left\| f, \underline{f}, a \right\|_{\mathfrak{h}_s(\mathbf{S})} \right). \end{aligned} \tag{9.3.21}$$

Proof. The proof follows easily from Lemma 9.3.8, Corollary 9.2.7, coercivity of $A^{\mathbf{S}}$ and obvious product estimates on \mathbf{S} . Consider for example the term

$$\text{Err}_2 = r^{-2}(\vartheta^{\mathbf{S}})^2 ((f, \underline{f}, a)^2) + r^{-1}\vartheta^{\mathbf{S}} ((f, \underline{f}, a)(\check{\Gamma})).$$

We write,

$$(\vartheta^{\mathbf{S}})^k \text{Err}_2 = r^{-2}(\vartheta^{\mathbf{S}})^{2+k} ((f, \underline{f}, a)^2) + r^{-1}(\vartheta^{\mathbf{S}})^{1+k} ((f, \underline{f}, a)(\check{\Gamma})) + \text{l.o.t.}$$

and

$$(\vartheta^{\mathbf{S}})^{2+k} ((f, \underline{f}, a)^2) = \sum_{i+j=k+2} \vartheta^i(f, \underline{f}, a) \cdot \vartheta^j(f, \underline{f}, a).$$

¹⁰This is assumption (9.2.14) of Lemma 9.2.6 with $\mathring{\delta}$ replaced to $\mathring{\epsilon}$.

Thus, dividing the sum into terms with $i \geq [\frac{k+2}{2}]$ and $i < [\frac{k+2}{2}]$ and using Sobolev estimates for the terms involving fewer derivatives we derive, for $[\frac{k+2}{2}] + 2 \leq k + 2$,

$$\|(\varphi^{\mathbf{S}})^{2+k} ((f, \underline{f}, a)^2)\|_{L^2(\mathbf{S})} \lesssim r^{-1} \|(a, f, \underline{f})\|_{\mathfrak{h}^{k+2}(\mathbf{S})}^2.$$

Similarly, making use of our assumptions for $\check{\Gamma}$,

$$\begin{aligned} (\varphi^{\mathbf{S}})^{1+k} ((f, \underline{f}, a)(\check{\Gamma})) &\lesssim r^{-1} \|(a, f, \underline{f})\|_{\mathfrak{h}^{k+1}(\mathbf{S})} \|\check{\Gamma}\|_{\mathfrak{h}^{k+1}(\mathbf{S})} \\ &\lesssim \overset{\circ}{\epsilon} r^{-1} \|(a, f, \underline{f})\|_{\mathfrak{h}^{k+1}(\mathbf{S})}. \end{aligned}$$

Thus, for all $2 \leq k \leq s_{max} - 1$

$$\|(\varphi^{\mathbf{S}})^k \text{Err}_2\|_{L^2(\mathbf{S})} \lesssim r^{-2} \|(f, \underline{f}, a)\|_{\mathfrak{h}^{k+2}(\mathbf{S})} \left(\epsilon_0 + \|(f, \underline{f}, a)\|_{\mathfrak{h}^{k+2}(\mathbf{S})} \right)$$

i.e., for $4 \leq s \leq s_{max} + 1$,

$$\|\text{Err}_2\|_{\mathfrak{h}_{s-2}(\mathbf{S})} \lesssim r^{-2} \|(f, \underline{f}, a)\|_{\mathfrak{h}_s(\mathbf{S})} \left(\overset{\circ}{\epsilon} + \|(f, \underline{f}, a)\|_{\mathfrak{h}_s(\mathbf{S})} \right).$$

All other terms can be treated similarly. □

9.3.2 Equation for the average of a

In the proof of existence and uniqueness of GCMS, see Theorem 9.4.2 we will need, in addition of the equations derived so far, an equation for the average of a . To achieve this we make use of the transformation formula for κ of Lemma 9.3.2

$$\begin{aligned} \kappa' &= e^a (\kappa + \not{d}'_1 f) + e^a \text{Err}(\kappa, \kappa'), \\ \text{Err}(\kappa, \kappa') &= f(\zeta + \eta) + \underline{f}\xi + \frac{1}{2}\underline{f}e_4 f + \frac{1}{2}f e_3 f + \frac{1}{4}f \underline{f} \kappa + f \underline{f} \omega - \underline{\omega} f^2 + \text{l.o.t.} \end{aligned}$$

which we rewrite in the form,

$$\kappa^{\mathbf{S}} = e^a \left(\frac{2}{r} + \check{\kappa} + \left(\bar{\kappa} - \frac{2}{r} \right) + \not{d}^{\mathbf{S}} f + \text{Err}(\kappa, \kappa') \right).$$

We deduce,

$$\begin{aligned} (e^a - 1) \frac{2}{r} &= \kappa^{\mathbf{S}} - \frac{2}{r} - e^a \left(\check{\kappa} + \bar{\kappa} - \frac{2}{r} + \not{d}^{\mathbf{S}} f \right) - e^a \text{Err}(\kappa, \kappa') \\ &= \kappa^{\mathbf{S}} - \frac{2}{r^{\mathbf{S}}} + \left(\frac{2}{r^{\mathbf{S}}} - \frac{2}{r} \right) - \left(\check{\kappa} + \bar{\kappa} - \frac{2}{r} + \not{d}^{\mathbf{S}} f \right) \\ &\quad - e^a \text{Err}(\kappa, \kappa') - (e^a - 1) \left(\check{\kappa} + \bar{\kappa} - \frac{2}{r} + \not{d}^{\mathbf{S}} f \right) \end{aligned}$$

or,

$$\begin{aligned} a \frac{2}{r} &= \kappa^{\mathbf{S}} - \frac{2}{r^{\mathbf{S}}} + \left(\frac{2}{r^{\mathbf{S}}} - \frac{2}{r} \right) - \left(\check{\kappa} + \bar{\kappa} - \frac{2}{r} + \not{d}^{\mathbf{S}} f \right) \\ &\quad - e^a \text{Err}(\kappa, \kappa') - (e^a - 1) \left(\check{\kappa} + \bar{\kappa} - \frac{2}{r} + \not{d}^{\mathbf{S}} f \right) - (e^a - 1 - a) \frac{2}{r}. \end{aligned}$$

We deduce,

$$\begin{aligned} a &= \frac{r^{\mathbf{S}}}{2} \left(\kappa^{\mathbf{S}} - \frac{2}{r^{\mathbf{S}}} \right) + \left(1 - \frac{r^{\mathbf{S}}}{r} \right) - \frac{r^{\mathbf{S}}}{2} \left(\check{\kappa} + \bar{\kappa} - \frac{2}{r} + \not{d}^{\mathbf{S}} f \right) + \text{Err}_6 \\ \text{Err}_6 &= -\frac{r^{\mathbf{S}}}{2} \left[e^a \text{Err}(\kappa, \kappa') - (e^a - 1) \left(\check{\kappa} + \bar{\kappa} - \frac{2}{r} + \not{d}^{\mathbf{S}} f \right) - (e^a - 1 - a) \frac{2}{r} \right] \\ &\quad - a \left(\frac{r^{\mathbf{S}}}{r} - 1 \right). \end{aligned}$$

Taking the average on \mathbf{S} we infer that,

$$\bar{a}^{\mathbf{S}} = \frac{r^{\mathbf{S}}}{2} \left(\overline{\kappa^{\mathbf{S}}} - \frac{2}{r^{\mathbf{S}}} \right) + \overline{\left(1 - \frac{r^{\mathbf{S}}}{r} \right)^{\mathbf{S}}} - \frac{r^{\mathbf{S}}}{2} \overline{\left(\check{\kappa} + \bar{\kappa} - \frac{2}{r} \right)^{\mathbf{S}}} + \overline{\text{Err}_6}^{\mathbf{S}} \quad (9.3.22)$$

where $\bar{h}^{\mathbf{S}}$ denotes the average of h on \mathbf{S} .

9.3.3 Transversality conditions

Lemma 9.3.11. *Assume given a deformed sphere $\mathbf{S} \subset \mathcal{R}$ with adapted null frame $e_3^{\mathbf{S}}, e_4^{\mathbf{S}}, e_\theta^{\mathbf{S}}$ and transition functions (a, f, \underline{f}) . We can extend a, f, \underline{f} , and thus the frame $e_3^{\mathbf{S}}, e_4^{\mathbf{S}}, e_\theta^{\mathbf{S}}$, in a small neighborhood of \mathbf{S} such that the following hold true*

$$\xi^{\mathbf{S}} = 0, \quad \omega^{\mathbf{S}} = 0, \quad \underline{\eta}^{\mathbf{S}} + \zeta^{\mathbf{S}} = 0. \quad (9.3.23)$$

Proof. According to Proposition 9.3.1 we have,

$$\begin{aligned} \xi^{\mathbf{S}} &= e^{2a} \left(\xi + \frac{1}{2}e^{-a}e_4^{\mathbf{S}}(f) + \frac{1}{4}f\kappa + f\omega \right) + e^{2a}\text{Err}(\xi, \xi^{\mathbf{S}}), \\ \underline{\xi}^{\mathbf{S}} &= e^{-2a} \left(\underline{\xi} + \frac{1}{2}e^ae_3^{\mathbf{S}}(\underline{f}) + \frac{1}{4}\underline{f}\underline{\kappa} + \underline{f}\underline{\omega} \right) + e^{-2a}\text{Err}(\underline{\xi}, \underline{\xi}^{\mathbf{S}}), \\ \zeta^{\mathbf{S}} &= \zeta - e_{\theta}^{\mathbf{S}}(a) - f\underline{\omega} + \underline{f}\omega - \frac{1}{4}f\underline{\kappa} + \frac{1}{4}\underline{f}\kappa + \text{Err}(\zeta, \zeta^{\mathbf{S}}), \\ \eta^{\mathbf{S}} &= \eta + \frac{1}{2}e^ae_3^{\mathbf{S}}f - f\underline{\omega} + \frac{1}{4}f\underline{\kappa} + \text{Err}(\eta, \eta^{\mathbf{S}}), \\ \underline{\eta}^{\mathbf{S}} &= \underline{\eta} + \frac{1}{2}e^{-a}e_4^{\mathbf{S}}\underline{f} - \underline{f}\omega + \frac{1}{4}\underline{f}\kappa + \text{Err}(\underline{\eta}, \underline{\eta}^{\mathbf{S}}), \\ \omega^{\mathbf{S}} &= e^a \left(\omega - \frac{1}{2}e^{-a}e_4a \right) + e^a\text{Err}(\omega, \omega^{\mathbf{S}}), \\ \underline{\omega}^{\mathbf{S}} &= e^{-a} \left(\underline{\omega} + \frac{1}{2}e^ae_3a \right) + e^{-a}\text{Err}(\underline{\omega}, \underline{\omega}^{\mathbf{S}}). \end{aligned}$$

Clearly the conditions $\xi^{\mathbf{S}} = 0, \omega^{\mathbf{S}} = 0$ allows us to determine $e_4^{\mathbf{S}}f$ and $e_4^{\mathbf{S}}a$ on \mathbf{S} while the condition $\underline{\eta}^{\mathbf{S}} + \zeta^{\mathbf{S}} = 0$ allows us to determine $e_4^{\mathbf{S}}\underline{f}$ on \mathbf{S} . \square

Remark 9.3.12. Note that the equations above also allow us to impose, in addition, vanishing conditions on $\underline{\xi}$ and ω along \mathbf{S} . Indeed these are determined by $e_3^{\mathbf{S}}f$ and $e_3^{\mathbf{S}}a$.

9.3.4 Equation for the $\ell = 1$ mode of $\not{d}^{\mathbf{S},*}a$

Recall the following change of frame formulas

$$e'_{\theta}(\kappa') = e_{\theta}\kappa + e'_{\theta}\not{d}_1'f + \kappa e'_{\theta}a - \frac{1}{4}\kappa(f\underline{\kappa} + \underline{f}\kappa) + \kappa(f\underline{\omega} - \omega\underline{f}) + f\rho + \text{Err}(e'_{\theta}\kappa', e_{\theta}\kappa).$$

where $e'_{\theta} = -\not{d}^{\mathbf{S},*}$, $\not{d}_1' = \not{d}^{\mathbf{S}}$, $\kappa' = \kappa^{\mathbf{S}}$, $\beta' = \beta^{\mathbf{S}}$. Making use of

$$\not{d}_1^* \not{d}_1' = \not{d}_2' \not{d}_2^* + 2K^{\mathbf{S}}, \quad K = -\rho - \frac{1}{4}\kappa\underline{\kappa} + \frac{1}{4}\vartheta\underline{\vartheta},$$

we deduce,

$$\begin{aligned} e'_{\theta}(\kappa') &= e_{\theta}\kappa - (\not{d}_2)'(\not{d}_2^*)'f + \kappa e'_{\theta}(a) - \kappa \left(\frac{1}{4}\kappa + \omega \right) \underline{f} + \left(-\frac{1}{4}\kappa\underline{\kappa} + \kappa\underline{\omega} + \rho - 2K \right) f \\ &+ \text{Err}(e'_{\theta}\kappa', e_{\theta}\kappa) - 2(K^{\mathbf{S}} - K)f \\ &= e_{\theta}\kappa - (\not{d}_2)'(\not{d}_2^*)'f + \kappa e'_{\theta}(a) - \kappa \left(\frac{1}{4}\kappa + \omega \right) \underline{f} + \left(\frac{1}{4}\kappa\underline{\kappa} + \kappa\underline{\omega} + 3\rho \right) f \\ &+ \text{Err}(e'_{\theta}\kappa', e_{\theta}\kappa) - 2(K^{\mathbf{S}} - K)f - \frac{1}{2}\vartheta\underline{\vartheta}f, \end{aligned}$$

or, in view of the condition $\kappa' = \kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}$ we have $e'_\theta(\kappa') = -\not{d}^{\mathbf{S},*}(\kappa^{\mathbf{S}}) = 0$,

$$\begin{aligned} e'_\theta(\kappa') &= e_\theta\kappa - (\not{d}_2)'(\not{d}_2^*)'f + \frac{2}{r'}e'_\theta(a) - \kappa\left(\frac{1}{4}\kappa + \omega\right)\underline{f} + \left(\frac{1}{4}\kappa\underline{\kappa} + \kappa\underline{\omega} + 3\rho\right)f \\ &+ \text{Err}(e'_\theta\kappa', e_\theta\kappa) + \left(\kappa - \frac{2}{r'}\right)e'_\theta(a) - 2(K^{\mathbf{S}} - K)f - \frac{1}{2}\vartheta\underline{\vartheta}f. \end{aligned}$$

Multiplying by e^Φ and integrating on \mathbf{S} we derive,

$$\begin{aligned} \int_{\mathbf{S}} e'_\theta(\kappa')e^\Phi &= \int_{\mathbf{S}} e_\theta(\kappa)e^\Phi + \frac{2}{r^{\mathbf{S}}} \int_{\mathbf{S}} e_\theta^{\mathbf{S}}(a)e^\Phi - \int_{\mathbf{S}} \kappa\left(\frac{1}{4}\kappa + \omega\right)\underline{f}e^\Phi + \int_{\mathbf{S}} \left(\frac{1}{4}\kappa\underline{\kappa} + \kappa\underline{\omega} + 3\rho\right)fe^\Phi \\ &+ \int_{\mathbf{S}} \left(\text{Err}(e_\theta^{\mathbf{S}}\kappa^{\mathbf{S}}, e_\theta\kappa) - 2(K^{\mathbf{S}} - K)f - \left(\kappa - \frac{2}{r^{\mathbf{S}}}\right)\not{d}^{\mathbf{S},*}a - \frac{1}{2}\vartheta\underline{\vartheta}f\right)e^\Phi. \end{aligned}$$

Therefore (recall that $\omega = 0$),

$$\begin{aligned} \frac{2}{r^{\mathbf{S}}} \int_{\mathbf{S}} e_\theta^{\mathbf{S}}(a)e^\Phi &= \int_{\mathbf{S}} e_\theta^{\mathbf{S}}(\kappa^{\mathbf{S}})e^\Phi - \int_{\mathbf{S}} e_\theta(\kappa)e^\Phi + \frac{1}{4} \int_{\mathbf{S}} \kappa^2\underline{f}e^\Phi - \int_{\mathbf{S}} \left(\frac{1}{4}\kappa\underline{\kappa} + \kappa\underline{\omega} + 3\rho\right)fe^\Phi \\ &+ \text{Err}_7, \\ \text{Err}_7 &= \int_{\mathbf{S}} \left(-\text{Err}(e_\theta^{\mathbf{S}}\kappa^{\mathbf{S}}, e_\theta\kappa) + 2(K^{\mathbf{S}} - K)f + (\kappa - \kappa^{\mathbf{S}})\not{d}^{\mathbf{S},*}a + \frac{1}{2}\vartheta\underline{\vartheta}f\right)e^\Phi. \end{aligned}$$

We summarize the result in the following lemma.

Lemma 9.3.13. *We have,*

$$\begin{aligned} \frac{2}{r^{\mathbf{S}}} \int_{\mathbf{S}} e_\theta^{\mathbf{S}}(a)e^\Phi &= \int_{\mathbf{S}} e_\theta^{\mathbf{S}}(\kappa^{\mathbf{S}})e^\Phi - \int_{\mathbf{S}} e_\theta(\kappa)e^\Phi + \frac{1}{4} \int_{\mathbf{S}} \kappa^2\underline{f}e^\Phi \\ &- \int_{\mathbf{S}} \left(\frac{1}{4}\kappa\underline{\kappa} + \kappa\underline{\omega} + 3\rho\right)fe^\Phi + \text{Err}_7, \end{aligned} \tag{9.3.24}$$

where,

$$\text{Err}_7 = \int_{\mathbf{S}} \left(-\text{Err}(e_\theta^{\mathbf{S}}\kappa^{\mathbf{S}}, e_\theta\kappa) + 2(K^{\mathbf{S}} - K)f + \left(\kappa - \frac{2}{r^{\mathbf{S}}}\right)\not{d}^{\mathbf{S},*}a + \frac{1}{2}\vartheta\underline{\vartheta}f\right)e^\Phi.$$

9.4 Existence of GCM spheres

We now impose the GCM conditions on the deformed sphere \mathbf{S}

$$\begin{aligned} \not{d}_2^{\mathbf{S},*}\not{d}_1^{\mathbf{S},*}\underline{\kappa}^{\mathbf{S}} &= \not{d}_2^{\mathbf{S},*}\not{d}_1^{\mathbf{S},*}\underline{\mu}^{\mathbf{S}} = 0, & \kappa^{\mathbf{S}} &= \frac{2}{r^{\mathbf{S}}}, \\ \int_{\mathbf{S}} fe^\Phi &= \underline{\Lambda}, & \int_{\mathbf{S}} \underline{f}e^\Phi &= \underline{\Lambda}, \end{aligned} \tag{9.4.1}$$

where (f, \underline{f}) belong to the triplet $(f, \underline{f}, \lambda = e^a)$ which denote the change of frame coefficients from the frame of $\overset{\circ}{S}$ to the one of \mathbf{S} . We show that under these conditions the deformation parameters a, f, \underline{f} verify a coercive elliptic system.

We start with the following simple adaptation of Lemmas 9.3.17, 9.3.13 and the result of subsection 9.3.2.

Proposition 9.4.1. *Assume that the deformed sphere \mathbf{S} verifies the GCMS conditions (9.4.1). Then the deformation parameters a, f, \underline{f} verify the system (recall that $z = \underline{\kappa}f + \kappa\underline{f}$ and $V = \frac{1}{2}\kappa\underline{\kappa} - \rho$)*

$$\begin{aligned}
(A^{\mathbf{S}} + V)z &= \kappa \not{d}^{\mathbf{S},*} \underline{\check{\kappa}}^{\mathbf{S}} - \underline{\kappa} \not{d}^* \check{\kappa} - \kappa \not{d}^* \check{\kappa} \\
&\quad + \underline{\kappa} \text{Err}_4 + \kappa \text{Err}_5 + [A^{\mathbf{S}}, \underline{\kappa}]f + [A^{\mathbf{S}}, \kappa]\underline{f}, \\
(A^{\mathbf{S}} + V)f &= \frac{3}{4}\kappa(A^{\mathbf{S}})^{-1}(\rho z) - \not{d}^* \check{\kappa} - \kappa(A^{\mathbf{S}})^{-1}(-\not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}} + \not{d}^* \check{\mu}) \\
&\quad + \text{Err}_4, \\
(A^{\mathbf{S}} + V)\underline{f} &= -\frac{3}{4}\underline{\kappa}(A^{\mathbf{S}})^{-1}(\rho z) + \not{d}^{\mathbf{S},*} \check{\kappa}^{\mathbf{S}} + \not{d}^* \check{\kappa} - \underline{\kappa}(A^{\mathbf{S}})^{-1}(-\not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}} + \not{d}^* \check{\mu}) \\
&\quad + \text{Err}_5, \\
\not{d}^{\mathbf{S},*}a &= -\frac{3}{4}(A^{\mathbf{S}})^{-1}(\rho z) + f\underline{\omega} + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa + (A^{\mathbf{S}})^{-1}(-\not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}} + \not{d}^* \check{\mu}) \\
&\quad - (A^{\mathbf{S}})^{-1} \text{Err}_3, \\
\bar{a}^{\mathbf{S}} &= \left(1 - \frac{r^{\mathbf{S}}}{r}\right)^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2} \overline{\left(\check{\kappa} + \bar{\kappa} - \frac{2}{r}\right)^{\mathbf{S}}} + \overline{\text{Err}_6}^{\mathbf{S}},
\end{aligned} \tag{9.4.2}$$

and,

$$\begin{aligned}
\frac{2}{r^{\mathbf{S}}} \int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(a) e^{\Phi} &= \int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(\kappa^{\mathbf{S}}) e^{\Phi} - \int_{\mathbf{S}} e_{\theta}(\kappa) e^{\Phi} + \frac{1}{4} \int_{\mathbf{S}} \kappa^2 \underline{f} e^{\Phi} \\
&\quad - \int_{\mathbf{S}} \left(\frac{1}{4} \kappa \underline{\kappa} + \kappa \underline{\omega} + 3\rho \right) f e^{\Phi} + \text{Err}_7,
\end{aligned} \tag{9.4.3}$$

$$\int_{\mathbf{S}} e^{\Phi} f = \Lambda, \quad \int_{\mathbf{S}} e^{\Phi} \underline{f} = \underline{\Lambda}. \tag{9.4.4}$$

The error terms are given by $\text{Err}_1, \text{Err}_2, \text{Err}_3$, defined in Lemma 9.3.5, and

$$\text{Err}_4 = \text{Err}_1 + \kappa(A^{\mathbf{S}})^{-1} \text{Err}_3, \quad \text{Err}_5 = \text{Err}_2 - \underline{\kappa}(A^{\mathbf{S}})^{-1} \text{Err}_3,$$

$$\begin{aligned}
Err_6 &= -\frac{r^{\mathbf{S}}}{2} \left[e^a Err(\kappa, \kappa') - (e^a - 1) \left(\check{\kappa} + \bar{\kappa} - \frac{2}{r} + \not{d}^{\mathbf{S}} f \right) - (e^a - 1 - a) \frac{2}{r} \right] \\
&\quad - a \left(\frac{r^{\mathbf{S}}}{r} - 1 \right), \\
Err_7 &= \int_{\mathbf{S}} \left(-Err(e_{\theta}^{\mathbf{S}} \kappa^{\mathbf{S}}, e_{\theta} \kappa) + 2(K^{\mathbf{S}} - K)f + \left(\frac{2}{r} - \frac{2}{r^{\mathbf{S}}} \right) \not{d}^{\mathbf{S}, \star} a + \frac{1}{2} \vartheta \not{d} f \right) e^{\Phi}.
\end{aligned}$$

We are ready to state the first main result of this chapter.

Theorem 9.4.2 (Existence of GCM spheres). *Let $\mathring{S} = S(\mathring{u}, \mathring{s})$ be a fixed sphere of the (u, s) outgoing geodesic foliation of a fixed spacetime region \mathcal{R} . Assume in addition to **A1–A3** that the following estimates hold true on \mathcal{R} , for all $k \leq s_{max}$,*

$$\begin{aligned}
|\not{d}^{k-2} \not{d}_2^{\star} \not{d}_1^{\star} \check{\kappa}| &\lesssim \mathring{\delta} r^{-4}, \\
|\not{d}^{k-2} \not{d}_2^{\star} \not{d}_1^{\star} \mu| &\lesssim \mathring{\delta} r^{-5}, \\
|\not{d}^k \check{\kappa}| &\lesssim \mathring{\delta} r^{-2}, \\
\left| \bar{\kappa} - \frac{2}{r} \right| + \left| \bar{\kappa} + \frac{2(1 - \frac{2m}{r})}{r} \right| &\lesssim \mathring{\delta} r^{-2}.
\end{aligned} \tag{9.4.5}$$

For any fix $\Lambda, \underline{\Lambda} \in \mathbb{R}$ verifying,

$$|\Lambda|, |\underline{\Lambda}| \lesssim \mathring{\delta} r^2 \tag{9.4.6}$$

there exists a unique GCM sphere $\mathbf{S} = \mathbf{S}^{(\Lambda, \underline{\Lambda})}$, which is a deformation of \mathring{S} , such that the GCMS conditions (9.4.1) are verified. Moreover the following estimates hold true.

1. We have

$$\left| \frac{r^{\mathbf{S}}}{\mathring{r}} - 1 \right| \lesssim r^{-1} \mathring{\delta}. \tag{9.4.7}$$

In particular r, \mathring{r} and $r^{\mathbf{S}}$ are all comparable in \mathcal{R} .

2. The unique functions $(\lambda, \underline{f}, \underline{f})$ on \mathbf{S} , which relate the original frame e_3, e_4, e_{θ} to the new frame on $e_3^{\mathbf{S}}, e_4^{\mathbf{S}}, e_{\theta}^{\mathbf{S}}$ according to (9.2.18), verify the estimates

$$\left\| f, \underline{f}, \log \lambda \right\|_{\mathfrak{h}_k(\mathbf{S})} \lesssim \mathring{\delta}, \quad k \leq s_{max} + 1. \tag{9.4.8}$$

3. The parameters U, S of the deformation, see Definition 9.2.1, verify the estimate

$$\|(U', S')\|_{L^\infty(\mathring{S})} + \max_{0 \leq s \leq s_{max}} r^{-1} \|(U', S')\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})} \lesssim \mathring{\delta}. \quad (9.4.9)$$

4. The Hawking mass $m^{\mathbf{S}}$ verifies the estimate,

$$|m^{\mathbf{S}} - \mathring{m}| \lesssim \mathring{\delta}. \quad (9.4.10)$$

5. The curvature components $(\alpha^{\mathbf{S}}, \beta^{\mathbf{S}}, \rho^{\mathbf{S}}, \underline{\beta}^{\mathbf{S}}, \underline{\alpha}^{\mathbf{S}})$, as well as $\mu^{\mathbf{S}}$ and the Ricci coefficients¹¹ $(\kappa^{\mathbf{S}}, \vartheta^{\mathbf{S}}, \zeta^{\mathbf{S}}, \underline{\kappa}^{\mathbf{S}}, \underline{\vartheta}^{\mathbf{S}})$ on \mathbf{S} , verify, for all $k \leq s_{max}$,

$$\begin{aligned} \|\check{\kappa}^{\mathbf{S}}, \vartheta^{\mathbf{S}}, \zeta^{\mathbf{S}}, \check{\underline{\kappa}}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon} r^{-1}, \\ \|\underline{\vartheta}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon}, \\ \|\alpha^{\mathbf{S}}, \beta^{\mathbf{S}}, \check{\rho}^{\mathbf{S}}, \mu^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon} r^{-2}, \\ \|\underline{\beta}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon} r^{-1}, \\ \|\underline{\alpha}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon}. \end{aligned} \quad (9.4.11)$$

6. The functions, $(\lambda, \underline{f}, \underline{f})$ uniquely defined above, can be smoothly extended to a small neighborhood of \mathbf{S} in such a way that the corresponding Ricci coefficients verify the following transversality conditions

$$\xi^{\mathbf{S}} = 0, \quad \omega^{\mathbf{S}} = 0, \quad \underline{\eta}^{\mathbf{S}} + \zeta^{\mathbf{S}} = 0. \quad (9.4.12)$$

In that case, the following estimates hold¹² for all $k \leq s_{max} - 1$

$$\|e_4(f, \underline{f}, \log \lambda)\|_{\mathfrak{h}_{k+1}(\mathbf{S})} \lesssim r^{-1} \mathring{\delta} + r^{-3} (|\Lambda| + |\underline{\Lambda}|), \quad (9.4.13)$$

and,

$$\begin{aligned} \|e_4^{\mathbf{S}}(\check{\kappa}^{\mathbf{S}}, \vartheta^{\mathbf{S}}, \zeta^{\mathbf{S}}, \check{\underline{\kappa}}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon} r^{-2}, \\ \|e_4^{\mathbf{S}}(\underline{\vartheta}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon} r^{-1}, \\ \|e_4^{\mathbf{S}}(\alpha^{\mathbf{S}}, \beta^{\mathbf{S}}, \check{\rho}^{\mathbf{S}}, \mu^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon} r^{-3}, \\ \|e_4^{\mathbf{S}}(\underline{\beta}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon} r^{-2}, \\ \|e_4^{\mathbf{S}}(\underline{\alpha}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon} r^{-1}. \end{aligned} \quad (9.4.14)$$

¹¹All other Ricci coefficients involve the transversal derivatives $e_3^{\mathbf{S}}, e_4^{\mathbf{S}}$ of the frame.

¹²To be more precise one should replace r by \mathring{r} in the estimates below. Of course r and \mathring{r} are comparable in \mathcal{R} , in particular on \mathbf{S} .

Remark 9.4.3. *In view of Propositions 9.2.9 and 9.4.1, to find a GCM sphere amounts to solve the following coupled system*

$$\begin{aligned} \zeta^\# U' &= ((\gamma^\mathbf{S})^\#)^{1/2} f^\# \left(1 + \frac{1}{4} (f \underline{f})^\# \right), \\ S' - \frac{\zeta^\#}{2} \underline{\Omega}^\# U' &= \frac{1}{2} ((\gamma^\mathbf{S})^\#)^{1/2} \underline{f}^\#, \\ (\gamma^\mathbf{S})^\# &= \gamma^\# + (\zeta^\#)^2 \left(\underline{\Omega} + \frac{1}{4} \underline{b}^2 \gamma \right)^\# (\partial_\theta U)^2 - 2\zeta^\# \partial_\theta U \partial_\theta S - (\gamma \zeta \underline{b})^\# \partial_\theta U, \\ U(0) &= S(0) = 0, \end{aligned} \tag{9.4.15}$$

where the inputs (a, f, \underline{f}) verifies (9.4.2) and (9.4.3). Recall that for a reduced scalar h defined on \mathbf{S} we write

$$h^\#(\overset{\circ}{u}, \overset{\circ}{s}, \theta) = h(\overset{\circ}{u} + U(\theta), \overset{\circ}{s} + S(\theta), \theta).$$

We will solve the coupled system of equations (9.4.15), (9.4.2), (9.4.3) and (9.4.4) by an iteration argument which will be introduced below. Before doing this however it pays to observe that the system (9.4.2) and (9.4.3) can be interpreted as an elliptic system on a fixed surface \mathbf{S} for (a, f, \underline{f}) . In the next subsection we state a result which establishes the coercivity of the system. The full proof of the theorem is detailed in the remaining part of this section, see subsection 9.4.1 – 9.6.3.

9.4.1 A priori estimates

The following result plays the main role in the proof of Theorem 9.4.2.

Proposition 9.4.4 (A priori Estimates for GCMS). *Let a fixed spacetime region \mathcal{R} verifying assumptions **A1** – **A3** and (9.4.5). Assume \mathbf{S} is a given surface in \mathcal{R} such that the area radius of \mathbf{S} verifies¹³*

$$\left| \frac{r^\mathbf{S}}{\overset{\circ}{r}} - 1 \right| \lesssim r^{-1} \overset{\circ}{\delta}. \tag{9.4.16}$$

Then, for every $\Lambda, \underline{\Lambda}$, $|\Lambda| + |\underline{\Lambda}| \lesssim \overset{\circ}{\delta}$, there exists a unique solution $(\lambda, f, \underline{f})$ of the system (9.4.2)–(9.4.3) verifying the estimate,

$$\|(\log \lambda, f, \underline{f})\|_{\mathfrak{h}_{s_{max}+1}(\mathbf{S})} \lesssim \overset{\circ}{\delta}. \tag{9.4.17}$$

¹³Recall also from (9.1.16) that $r - \overset{\circ}{r} = O(\overset{\circ}{\delta} \epsilon^{-1/2})$. Thus $r, \overset{\circ}{r}$ and $r^\mathbf{S}$ are all comparable.

More precisely,

$$\|f, \underline{f}, \log \lambda\|_{\mathfrak{h}_{s_{max}+1}(\mathbf{S})} \lesssim \overset{\circ}{\delta} + |\Lambda| + |\underline{\Lambda}|. \tag{9.4.18}$$

As an immediate corollary we derive the following rigidity result for GCM spheres.

Corollary 9.4.5 (Rigidity I). *Let a fixed spacetime region \mathcal{R} verifying assumptions **A1** – **A3** and (9.4.5). Assume \mathbf{S} is a sphere in \mathcal{R} which verifies the the GCM conditions*

$$\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}, \quad \phi_2^{\star \mathbf{S}} \phi_1^{\star \mathbf{S}} \underline{\kappa}^{\mathbf{S}} = \phi_2^{\star \mathbf{S}} \phi_1^{\star \mathbf{S}} \mu^{\mathbf{S}} = 0. \tag{9.4.19}$$

Then the transition functions $(f, \underline{f}, \log \lambda)$ from the background frame of \mathcal{R} to that of \mathbf{S} verifies the estimates

$$\|(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}_{s_{max}+1}(\mathbf{S})} \lesssim \overset{\circ}{\delta} + \left| \int_{\mathbf{S}} f e^{\Phi} \right| + \left| \int_{\mathbf{S}} \underline{f} e^{\Phi} \right|.$$

The proof of Proposition 9.4.4 will be given in the next section. The proof of Corollary 9.4.5 is an immediate consequence of the proposition.

We assume in addition that \mathbf{S} is a deformation of $\overset{\circ}{S}$ and prove the following corollary of Proposition 9.4.4.

Corollary 9.4.6. *Let a fixed spacetime region \mathcal{R} verifying assumptions **A1** – **A3** and (9.4.5), and let $\Psi : \overset{\circ}{S} \rightarrow \mathbf{S}$ be a fixed deformation in \mathcal{R} given by the scalars U, S . We assume that the deformation verifies, see (9.2.14),*

$$\|(U', S')\|_{L^\infty(\overset{\circ}{S})} \lesssim \overset{\circ}{\delta}. \tag{9.4.20}$$

Then (a, f, \underline{f}) verify the estimate

$$\|(a, f, \underline{f})\|_{\mathfrak{h}_{s_{max}+1}(\mathbf{S})} \lesssim \overset{\circ}{\delta}. \tag{9.4.21}$$

Proof. In view of Proposition 9.4.4, it suffices to prove that (9.4.16) holds. (9.4.16) actually follows from the comparison Lemma 9.2.3. □

9.4.2 Comparison of the Hawking mass

We establish the estimate (9.4.10) concerning the Hawking mass $m^{\mathbf{S}}$. Recall that,

$$\begin{aligned} m^{\mathbf{S}} &= \frac{r^{\mathbf{S}}}{2} \left(1 + \frac{1}{16\pi} \int_{\mathbf{S}} \kappa^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}} \right), \\ \mathring{m} &= \frac{\mathring{r}}{2} \left(1 + \frac{1}{16\pi} \int_{\mathring{\mathbf{S}}} \mathring{\kappa} \mathring{\underline{\kappa}} \right). \end{aligned}$$

We write

$$\begin{aligned} 2 \left(\frac{m^{\mathbf{S}}}{r^{\mathbf{S}}} - \frac{\mathring{m}}{\mathring{r}} \right) &= \frac{1}{16\pi} \left[\int_{\mathbf{S}} (\kappa^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}} - \kappa \underline{\kappa}) + \left(\int_{\mathbf{S}} \kappa \underline{\kappa} - \int_{\mathring{\mathbf{S}}} \kappa \underline{\kappa} \right) - \int_{\mathring{\mathbf{S}}} (\kappa \underline{\kappa} - \mathring{\kappa} \mathring{\underline{\kappa}}) \right] \\ &= I_1 + I_2 + I_3. \end{aligned}$$

In view of Proposition 9.2.9 we have $|r^{\mathbf{S}} - \mathring{r}| \lesssim \mathring{\delta}$ and $|\gamma^{\mathbf{S},\#} - \mathring{\gamma}| \lesssim \mathring{\delta} \mathring{r}$. Making use of Corollary 9.2.5 and the assumptions **A1-A3** for $\kappa, \underline{\kappa}$ we deduce,

$$|I_2| = \left| \int_{\mathbf{S}} \kappa \underline{\kappa} - \int_{\mathring{\mathbf{S}}} \kappa \underline{\kappa} \right| \lesssim \mathring{\delta} r^{-1}.$$

Similarly, taking into account the definition of $\mathcal{R} := \left\{ |u - \mathring{u}| \leq \mathring{\delta}, \quad |s - \mathring{s}| \leq \mathring{\delta} \right\}$,

$$|I_3| \lesssim \mathring{\delta} r^{-1}.$$

Finally, making also use of the transformation formula from the original frame (e_4, e_3, e_θ) to the frame $(e_4^{\mathbf{S}}, e_3^{\mathbf{S}}, e_\theta^{\mathbf{S}})$ of \mathbf{S}

$$\kappa^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}} = (\kappa + \mathcal{L}^{\mathbf{S}} f + \text{Err}(\kappa, \kappa^{\mathbf{S}})) (\underline{\kappa} + \mathcal{L}^{\mathbf{S}} \underline{f} + \text{Err}(\underline{\kappa}, \underline{\kappa}^{\mathbf{S}}))$$

and the estimates for $(f, \underline{f}, a = \log \lambda)$ we deduce,

$$|\kappa^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}} - \kappa \underline{\kappa}| \lesssim r^{-3} \mathring{\delta}.$$

Hence,

$$|I_1| \lesssim \mathring{\delta} r^{-1}.$$

We infer that,

$$\left| \frac{m^{\mathbf{S}}}{r^{\mathbf{S}}} - \frac{\mathring{m}}{\mathring{r}} \right| \lesssim \mathring{\delta} r^{-1}$$

from which the desired estimate (9.4.10) easily follows.

9.4.3 Iteration procedure for Theorem 9.4.2

We solve the coupled system of equations (9.4.15), (9.4.2) and (9.4.3) by an iteration argument as follows.

Starting with the trivial quintet

$$Q^{(0)} := (U^{(0)}, S^{(0)}, a^{(0)}, f^{(0)}, \underline{f}^{(0)}) = (0, 0, 0, 0, 0),$$

corresponding to the undeformed sphere \mathring{S} , we define iteratively the quintet

$$Q^{(n+1)} = (U^{(n+1)}, S^{(n+1)}, a^{(n+1)}, f^{(n+1)}, \underline{f}^{(n+1)})$$

from

$$Q^{(n)} = (U^{(n)}, S^{(n)}, a^{(n)}, f^{(n)}, \underline{f}^{(n)})$$

as follows.

1. The pair $(U^{(n)}, S^{(n)})$ defines the deformation sphere $\mathbf{S}(n)$ and the corresponding pull back map $\#_n$ given by the map $\Psi^{(n)} : \mathring{S} \rightarrow \mathbf{S}(n)$,

$$(\mathring{u}, \mathring{s}, \theta, \varphi) \longrightarrow (\mathring{u} + U^{(n)}(\theta), \mathring{s} + S^{(n)}(\theta), \theta, \varphi). \quad (9.4.22)$$

By induction we assume that the following estimates hold true:

$$\left\| (a^{(n)}, f^{(n)}, \underline{f}^{(n)}) \right\|_{\mathfrak{h}_{s_{max}-2}(\mathbf{S}(n))} \lesssim \mathring{\delta}, \quad (9.4.23)$$

and

$$\|\partial_\theta (U^{(n)}, S^{(n)})\|_{L^\infty(\mathring{S})} + \max_{0 \leq s \leq s_{max}-2} r^{-1} \|\partial_\theta (U^{(n)}, S^{(n)})\|_{\mathfrak{h}_s(\mathring{S}, \mathring{\vartheta})} \lesssim \mathring{\delta}. \quad (9.4.24)$$

2. We then define the triplet $(a^{(n+1)}, f^{(n+1)}, \underline{f}^{(n+1)})$ by solving the system on $\mathbf{S}(n)$

consisting of the equations (9.4.25), (9.4.28) and (9.4.29) below.

$$\begin{aligned}
(A^{\mathbf{S}^{(n)}} + V) z^{(n+1)} &= E^{(n+1)}, \\
E^{(n+1)} &:= \kappa \not{d}^{\mathbf{S}^{(n),\star} \check{\kappa}^{\mathbf{S}^{(n)}}} - \underline{\kappa} \not{d}^{\star} \check{\kappa} - \kappa \not{d}^{\star} \check{\kappa} \\
&\quad + \underline{\kappa} \text{Err}_4^{(n+1)} + \kappa \text{Err}_5^{(n+1)} + [A^{\mathbf{S}^{(n)}}, \underline{\kappa}] f^{(n+1)} + [A^{\mathbf{S}^{(n)}}, \kappa] \underline{f}^{(n+1)}, \\
(A^{\mathbf{S}^{(n)}} + V) f^{(n+1)} &= \frac{3}{4} \kappa (A^{\mathbf{S}^{(n)}})^{-1} (\rho z^{(n+1)}) + E^{(n+1)}, \\
E^{(n+1)} &:= -\not{d}^{\star} \check{\kappa} - \kappa (A^{\mathbf{S}^{(n)}})^{-1} (-\not{d}^{\mathbf{S}^{(n),\star} \check{\mu}^{\mathbf{S}^{(n)}}} + \not{d}^{\star} \check{\mu}) + \text{Err}_4^{(n+1)},
\end{aligned} \tag{9.4.25}$$

$$\begin{aligned}
(A^{\mathbf{S}^{(n)}} + V) \underline{f}^{(n+1)} &= -\frac{3}{4} \underline{\kappa} (A^{\mathbf{S}^{(n)}})^{-1} (\rho z^{(n+1)}) + \underline{E}^{(n+1)}, \\
\underline{E}^{(n+1)} &:= \not{d}^{\mathbf{S}^{(n),\star} \check{\kappa}^{\mathbf{S}^{(n)}}} - \not{d}^{\star} \check{\kappa} + \underline{\kappa} (A^{\mathbf{S}^{(n)}})^{-1} (-\not{d}^{\mathbf{S}^{(n),\star} \check{\mu}^{\mathbf{S}^{(n)}}} + \not{d}^{\star} \check{\mu}) \\
&\quad + \text{Err}_5^{(n+1)}, \\
\not{d}^{\mathbf{S}^{(n)}} a^{(n+1)} &= -\frac{3}{4} (A^{\mathbf{S}^{(n)}})^{-1} (\rho z^{(n+1)}) + f^{(n+1)} \underline{\omega} + \frac{1}{4} f^{(n+1)} \underline{\kappa} - \frac{1}{4} \underline{f}^{(n+1)} \kappa \\
&\quad \tilde{E}^{(n+1)}, \\
\tilde{E}^{(n+1)} &:= (A^{\mathbf{S}^{(n)}})^{-1} (-\not{d}^{\mathbf{S}^{(n),\star} \check{\mu}^{\mathbf{S}^{(n)}}} + \not{d}^{\star} \check{\mu}) - (A^{\mathbf{S}^{(n)}})^{-1} (\text{Err}_3^{(n+1)}),
\end{aligned}$$

where,

$$z^{(n+1)} = \underline{\kappa} f^{(n+1)} + \kappa \underline{f}^{(n+1)}, \tag{9.4.26}$$

and the error terms,

$$\text{Err}_1^{(n+1)}, \text{Err}_2^{(n+1)}, \text{Err}_3^{(n+1)}, \text{Err}_4^{(n+1)}, \text{Err}_5^{(n+1)}, \tag{9.4.27}$$

are obtained from the error terms $\text{Err}_1, \text{Err}_2, \text{Err}_3, \text{Err}_4, \text{Err}_5$ by setting $(a, f, \underline{f}) = (a^{(n+1)}, f^{(n+1)}, \underline{f}^{(n+1)})$ and their derivatives by the corresponding ones on the sphere $\mathbf{S}^{(n)}$.

We also set,

$$\begin{aligned}
\frac{2}{r^{\mathbf{S}^{(n)}}} \int_{\mathbf{S}^{(n)}} e_{\theta}^{\mathbf{S}^{(n)}}(a^{(n+1)}) e^{\Phi} &= \frac{1}{4} \int_{\mathbf{S}^{(n)}} \kappa^2 \underline{f}^{(n+1)} e^{\Phi} - \int_{\mathbf{S}^{(n)}} \left(\frac{1}{4} \kappa \underline{\kappa} + \kappa \underline{\omega} + 3\rho \right) f^{(n+1)} e^{\Phi} \\
&\quad - \int_{\mathbf{S}^{(n)}} e_{\theta}(\kappa) e^{\Phi} + \text{Err}_7^{(n+1)}, \\
\int_{\mathbf{S}^{(n)}} e^{\Phi} f^{(n+1)} &= \underline{\Lambda}, \quad \int_{\mathbf{S}^{(n)}} e^{\Phi} \underline{f}^{(n+1)} = \underline{\Lambda},
\end{aligned} \tag{9.4.28}$$

and,

$$\overline{a^{(n+1)}}^{\mathbf{S}(n)} = \overline{\left(1 - \frac{r^{\mathbf{S}(n)}}{r}\right)^{\mathbf{S}(n)}} - \frac{r^{\mathbf{S}(n)}}{2} \overline{\left(\check{\kappa} + \bar{\kappa} - \frac{2}{r}\right)^{\mathbf{S}(n)}} + \overline{\text{Err}_6^{(n+1)}}^{\mathbf{S}(n)}, \quad (9.4.29)$$

where $\text{Err}_6^{(n+1)}$, $\text{Err}_7^{(n+1)}$ are obtained from the error terms Err_6 , Err_7 as above in (9.4.27), by setting $(a, f, \underline{f}) = (a^{(n+1)}, f^{(n+1)}, \underline{f}^{(n+1)})$ and their derivatives by the corresponding ones on the sphere $\mathbf{S}(n)$.

3. In view of the induction hypothesis (9.4.24) we are thus in a position to apply Proposition 9.4.4, more precisely its Corollary 9.4.6, to construct a unique solution verifying the estimate, uniformly for $n \in \mathbb{N}$,

$$\left\| (a^{(n+1)}, f^{(n+1)}, \underline{f}^{(n+1)}) \right\|_{\mathfrak{h}_{s_{max}-2}(\mathbf{S}(n))} \lesssim \mathring{\delta}. \quad (9.4.30)$$

4. We use the new pair $(f^{(n+1)}, \underline{f}^{(n+1)})$ to solve the equations on \mathring{S} ,

$$\begin{aligned} \varsigma^{\#_n} \partial_\theta U^{(1+n)} &= (\gamma^{(n)})^{1/2} (f^{(1+n)})^{\#_n} \left(1 + \frac{1}{4} (f^{(1+n)} \underline{f}^{(1+n)})^{\#_n} \right), \\ \partial_\theta S^{(1+n)} - \frac{1}{2} \varsigma^{\#_n} \underline{\Omega}^{\#_n} \partial_\theta U^{(1+n)} &= \frac{1}{2} (\gamma^{(n)})^{1/2} (\underline{f}^{(1+n)})^{\#_n}, \\ \gamma^{(n)} &= \gamma^{\#_n} + (\varsigma^{\#_n})^2 \left(\underline{\Omega} + \frac{1}{4} \underline{b}^2 \gamma \right)^{\#_n} (\partial_\theta U^{(n)})^2 \\ &\quad - 2 \varsigma^{\#_n} \partial_\theta U^{(n)} \partial_\theta S^{(n)} - (\gamma \varsigma \underline{b})^{\#_n} \partial_\theta U^{(n)}, \\ U^{(1+n)}(0) &= S^{(1+n)}(0) = 0, \end{aligned} \quad (9.4.31)$$

where, we repeat, the pull back $\#_n$ is defined with respect to the map

$$\Psi^{(n)}(\mathring{u}, \mathring{s}, \theta) = (\mathring{u} + U^{(n)}(\theta), \mathring{s} + S^{(n)}(\theta), \theta),$$

and

$$\gamma^{(n)} := \gamma^{\mathbf{S}(n), \#_n}.$$

The equation (9.4.31) admits a unique solution $U^{(1+n)}, S^{(1+n)}$, according to the proposition below. The new pair $(U^{(n+1)}, S^{(n+1)})$ defines the new polarized sphere $\mathbf{S}(n+1)$ and we can proceed with the next step of the iteration.

9.4.4 Boundedness of the iterates

Proposition 9.4.7. *The equation (9.4.31) admits a unique solution $U^{(1+n)}, S^{(1+n)}$ verifying the estimate,*

$$\|\partial_\theta (U^{(n+1)}, S^{(n+1)})\|_{L^\infty(\mathring{S})} + r^{-1} \|\partial_\theta (U^{(n+1)}, S^{(n+1)})\|_{\mathfrak{h}_{s_{max}-2}(\mathring{S}, \mathring{g})} \lesssim \mathring{\delta}$$

uniformly for all $n \in \mathbb{N}$.

Proof. The existence and uniqueness part of the proposition is an immediate consequence of the standard results for ODE's.

To prove the desired estimate, we use the equations for $(U^{(1+n)}, S^{(1+n)})$ and infer, for $s = s_{max} - 2$,

$$\|\partial_\theta U^{(n+1)}\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})} \lesssim \left\| (\gamma^{(n)})^{1/2} (\varsigma^{\#n})^{-1} (f^{(1+n)})^{\#n} \left(1 + \frac{1}{4} (f^{(1+n)} \underline{f}^{(1+n)})^{\#n} \right) \right\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})}.$$

Together with the non sharp product estimate on $(\mathring{S}, \mathring{g})$, see Lemma 9.1.5, we infer that, for $s = s_{max} - 2$,

$$\begin{aligned} \|\partial_\theta U^{(n+1)}\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})} &\lesssim r^{-1} \left\| (f^{(1+n)})^{\#n}, (\underline{f}^{(1+n)})^{\#n} \right\|_{\mathfrak{h}_{s_{max}-2}(\mathring{S}, \mathring{g})} \left\| (\varsigma^{\#n})^{-1} (\gamma^{(n)})^{1/2} \right\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})} \\ &\quad \times \left(1 + \left\| (f^{(1+n)})^{\#n}, (\underline{f}^{(1+n)})^{\#n} \right\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})}^2 \right). \end{aligned}$$

In view of Lemma 9.2.6, Corollary 9.2.7, and (9.4.30), we deduce

$$\|\partial_\theta U^{(n+1)}\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})} \lesssim \mathring{\delta} r^{-1} \left\| (\varsigma^{\#n})^{-1} (\gamma^{(n)})^{1/2} \right\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g})}.$$

We recall that,

$$\gamma^{(n)} = \gamma^{\#n} + (\varsigma^{\#n})^2 \left(\underline{\Omega} + \frac{1}{4} \underline{b}^2 \gamma \right)^{\#n} (\partial_\theta U^{(n)})^2 - 2\varsigma^{\#n} \partial_\theta U^{(n)} - \left(\gamma \varsigma \underline{b} \right)^{\#n} \partial_\theta U^{(n)}.$$

In view of our assumptions on the Ricci coefficients and the non-sharp product estimates of Lemma 9.1.5

$$\left\| \left(\varsigma \left(\underline{\Omega} + \frac{1}{4} \underline{b}^2 \gamma \right) \right)^{\#n} \right\|_{\mathfrak{h}_{s_{max}-2}(\mathring{S}, \mathring{g})} + \left\| \left(\gamma \underline{b} \right)^{\#n} \right\|_{\mathfrak{h}_{s_{max}-2}(\mathring{S}, \mathring{g})} \lesssim \mathring{\epsilon} r$$

we deduce,

$$\left\| (\varsigma^{\#n})^{-1} \gamma^{(n)} \right\|_{\mathfrak{h}_{smax-2}(\mathring{S}, \mathring{\mathcal{G}})} \lesssim \left\| \gamma^{\#n} \right\|_{\mathfrak{h}_{smax-2}(\mathring{S}, \mathring{\mathcal{G}})} + \mathring{\delta} r^2.$$

Together with Lemma 9.2.6 and Corollary 9.2.7, we deduce

$$\left\| (\varsigma^{\#n})^{-1} \gamma^{(n)} \right\|_{\mathfrak{h}_{smax-2}(\mathring{S}, \mathring{\mathcal{G}})} \lesssim r^2$$

and therefore,

$$\left\| \partial_\theta U^{(n+1)} \right\|_{\mathfrak{h}_{smax-2}(\mathring{S}, \mathring{\mathcal{G}})} \lesssim \mathring{\delta} r.$$

Proceeding in the same manner with equation

$$\partial_\theta S^{(1+n)} - \frac{1}{2} \varsigma^{\#n} \underline{\Omega}^{\#n} \partial_\theta U^{(1+n)} = \frac{1}{2} (\gamma^{(n)})^{1/2} (\underline{f}^{(1+n)})^{\#n}$$

we infer that,

$$r^{-1} \left\| \partial_\theta U^{(n+1)}, \partial_\theta S^{(n+1)} \right\|_{\mathfrak{h}_{smax-2}(\mathring{S}, \mathring{\mathcal{G}})} \lesssim \mathring{\delta}.$$

This, together with the Sobolev inequality, concludes the proof of Proposition 9.4.7. \square

9.4.5 Convergence of the Iterates

To finish the proof of Theorem 9.4.2 it remains to prove convergence of the iterates.

Step 1. In order to prove the convergence of the iterative scheme, we introduce the following quintet $P^{(n)}$

$$P^{(0)} = (0, 0, 0, 0, 0), \quad P^{(n)} = \left(U^{(n)}, S^{(n)}, (a^{(n)})^{\#_{n-1}}, (f^{(n)})^{\#_{n-1}}, (\underline{f}^{(n)})^{\#_{n-1}} \right), \quad n \geq 1.$$

Since $(a^{(n)}, f^{(n)}, \underline{f}^{(n)})$ are defined on $\mathbf{S}(n-1)$, their respective pullback by $\Psi^{(n-1)}$ is defined on \mathring{S} so that $P^{(n)}$ is a quintet of functions on \mathring{S} for any n and we may introduce the following norms to compare the elements of the sequence

$$\begin{aligned} \|P^{(n)}\|_k &:= \left\| \partial_\theta (U^{(n)}, S^{(n)}) \right\|_{L^\infty(\mathring{S})} + r^{-1} \left\| \partial_\theta (U^{(n)}, S^{(n)}) \right\|_{\mathfrak{h}_{k-2}(\mathring{S})} \\ &+ \left\| \left((a^{(n)})^{\#_{n-1}}, (f^{(n)})^{\#_{n-1}}, (\underline{f}^{(n)})^{\#_{n-1}} \right) \right\|_{\mathfrak{h}_{k-2}(\mathring{S})}. \end{aligned} \tag{9.4.32}$$

Here are the substeps needed to implement a convergence argument.

1. The quintets $P^{(n)}$ are bounded with respect to the norm (9.4.32) for the choice $k = s_{max}$.
2. The quintets $P^{(n)}$ are contractive with respect to the norm (9.4.32) for the choice $k = 3$.

The precise statements are given in the following propositions.

Proposition 9.4.8. *We have, uniformly for all $n \in \mathbb{N}$,*

$$\|P^{(n)}\|_{s_{max}} \lesssim \overset{\circ}{\delta}.$$

Proof. The proof is an immediate consequence of Propositions 9.4.4, 9.4.7 and the estimate,

$$\left\| (\Psi^{(n-1)})^\# \left(f^{(n)}, \underline{f}^{(n)}, a^{(n)} \right) \right\|_{\mathfrak{h}_{s_{max}-2}(\overset{\circ}{S})} \lesssim \left\| \left(f^{(n)}, \underline{f}^{(n)}, a^{(n)} \right) \right\|_{\mathfrak{h}_{s_{max}-2}(\mathbf{S}^{(n-1)})} \quad (9.4.33)$$

which is a consequence of Lemma 9.2.6. □

Proposition 9.4.9. *We have, uniformly for all $n \in \mathbb{N}$, the contraction estimate,*

$$\|P^{(n+1)} - P^{(n)}\|_3 \lesssim \overset{\circ}{\delta} \|P^{(n)} - P^{(n-1)}\|_3.$$

The proof of Proposition 9.4.9 is postponed to section 9.6.

Step 2. In view of Proposition 9.4.9, we have

$$\|P^{(n+1)} - P^{(n)}\|_3 \lesssim (\overset{\circ}{\delta})^n \|P^{(1)} - P^{(0)}\|_3$$

which in view of Proposition 9.4.8 yields

$$\|P^{(n+1)} - P^{(n)}\|_3 \lesssim (\overset{\circ}{\delta})^{n+1}.$$

Together with a simple interpolation argument on $\overset{\circ}{S}$ and Proposition 9.4.8, we infer

$$\|P^{(n+1)} - P^{(n)}\|_k \lesssim (\overset{\circ}{\delta})^{1 + \left(\frac{s_{max}-k}{s_{max}-3}\right)n}, \quad 3 \leq k \leq s_{max}.$$

We infer the existence of a quintet of functions $P^{(\infty)}$ on $\overset{\circ}{S}$ such that

$$\|P^{(\infty)}\|_{s_{max}} \lesssim \overset{\circ}{\delta} \quad (9.4.34)$$

and

$$\lim_{n \rightarrow +\infty} \|P^{(n)} - P^{(\infty)}\|_{s_{max}-1} = 0. \tag{9.4.35}$$

Also, we have

$$P^{(\infty)} = \left(U^{(\infty)}, S^{(\infty)}, a_0^{(\infty)}, f_0^{(\infty)}, \underline{f}_0^{(\infty)} \right),$$

where all function are defined on $\overset{\circ}{S}$. The functions $(U^{(\infty)}, S^{(\infty)})$ defines a sphere $\mathbf{S}(\infty)$ and we introduce the map

$$\Psi^{(\infty)}(\overset{\circ}{u}, \overset{\circ}{s}, \theta, \varphi) = \left(\overset{\circ}{u} + U^{(\infty)}(\theta), \overset{\circ}{s} + S^{(\infty)}(\theta), \theta, \varphi \right)$$

so that $\Psi^{(\infty)}$ is a map from $\overset{\circ}{S}$ to $\mathbf{S}(\infty)$. Then, let

$$a^{(\infty)} = a_0^{(\infty)} \circ (\Psi^{(\infty)})^{-1}, \quad f^{(\infty)} = f_0^{(\infty)} \circ (\Psi^{(\infty)})^{-1}, \quad \underline{f}^{(\infty)} = \underline{f}_0^{(\infty)} \circ (\Psi^{(\infty)})^{-1}$$

so that $a^{(\infty)}, f^{(\infty)}, \underline{f}^{(\infty)}$ are defined on $\mathbf{S}(\infty)$ and

$$a_0^{(\infty)} = (a^{(\infty)})\#_{\infty}, \quad f_0^{(\infty)} = (f^{(\infty)})\#_{\infty}, \quad \underline{f}_0^{(\infty)} = (\underline{f}^{(\infty)})\#_{\infty}.$$

From these definitions, the above control of $P^{(\infty)}$ and Lemma 9.2.6, we infer

$$r^{-1} \|(U^{(\infty)}, S^{(\infty)})\|_{\mathfrak{h}_{s_{max}-1}(\overset{\circ}{S})} + \|(a^{(\infty)}, f^{(\infty)}, \underline{f}^{(\infty)})\|_{\mathfrak{h}_{s_{max}-2}(\mathbf{S}(\infty))} \lesssim \overset{\circ}{\delta}.$$

In particular, using the Sobolev embedding on $\overset{\circ}{S}$, we have $\|(U^{(\infty)'}, S^{(\infty)'})\|_{L^{\infty}(\overset{\circ}{S})} \lesssim \overset{\circ}{\delta}$, and hence, in view of Corollary 9.4.6, we deduce

$$\|(a^{(\infty)}, f^{(\infty)}, \underline{f}^{(\infty)})\|_{\mathfrak{h}_{s_{max}+1}(\mathbf{S}(\infty))} \lesssim \overset{\circ}{\delta}.$$

This estimate allows to argue as in Proposition 9.4.7 with $(U^{(n+1)}, S^{(n+1)})$ replaced by $(U^{(\infty)}, S^{(\infty)})$ and $s_{max} - 2$ replaced by s_{max} . We finally obtain

$$r^{-1} \|(U^{(\infty)'}, S^{(\infty)'})\|_{\mathfrak{h}_{s_{max}}(\overset{\circ}{S})} + \|(a^{(\infty)}, f^{(\infty)}, \underline{f}^{(\infty)})\|_{\mathfrak{h}_{s_{max}+1}(\mathbf{S}(\infty))} \lesssim \overset{\circ}{\delta}. \tag{9.4.36}$$

Step 3. We proceed to control the area radius $r^{\mathbf{S}(\infty)}$ and the Hawking mass $m^{\mathbf{S}(\infty)}$ of the sphere $\mathbf{S}(\infty)$. First, note from (9.4.36) and the Sobolev embedding on $\overset{\circ}{S}$ that we have

$$\|(U^{(\infty)}, S^{(\infty)})\|_{L^{\infty}(\overset{\circ}{S})} \lesssim \overset{\circ}{\delta}. \tag{9.4.37}$$

Together with Lemma 9.2.3, we infer that¹⁴

$$\left| \frac{r^{\mathbf{S}(\infty)}}{r} - 1 \right| \lesssim \overset{\circ}{\delta}. \quad (9.4.38)$$

Next, we denote by $\Gamma^{\mathbf{S}(\infty)}$ the connection coefficients of $\mathbf{S}(\infty)$. We have in view of the transformation formula from the original frame (e_4, e_3, e_θ) to the frame $(e_4^{\mathbf{S}(\infty)}, e_3^{\mathbf{S}(\infty)}, e_\theta^{\mathbf{S}(\infty)})$ of $\mathbf{S}(\infty)$

$$\kappa^{\mathbf{S}(\infty)} \underline{\kappa}^{\mathbf{S}(\infty)} = \left(\kappa + \not{d}^{\mathbf{S}(\infty)} f^{(\infty)} + \text{Err}(\kappa, \kappa^{\mathbf{S}(\infty)}) \right) \left(\underline{\kappa} + \not{d}^{\mathbf{S}(\infty)} \underline{f}^{(\infty)} + \text{Err}(\underline{\kappa}, \underline{\kappa}^{\mathbf{S}(\infty)}) \right).$$

Together with the estimate (9.4.36) for $f^{(\infty)}$ and $\underline{f}^{(\infty)}$ and the assumptions **A1-A3** for $\check{\Gamma}$ corresponding to the original frame (e_4, e_3, e_θ) , we infer

$$\left| \kappa^{\mathbf{S}(\infty)} \underline{\kappa}^{\mathbf{S}(\infty)} - \kappa \underline{\kappa} \right| \lesssim \overset{\circ}{\delta} r^{-3}.$$

Recall that (see (9.4.5))

$$\left| \kappa - \frac{2}{r} \right| \lesssim \overset{\circ}{\delta} r^{-2}, \quad \left| \bar{\kappa} + \frac{2(1 - \frac{2m}{r})}{r} \right| \lesssim \overset{\circ}{\delta}.$$

Thus, since $\underline{\kappa} = \bar{\kappa} + \check{\kappa}$,

$$\kappa \underline{\kappa} = -\frac{4(1 - \frac{2m}{r})}{r^2} + \frac{2}{r} \check{\kappa} + O(\overset{\circ}{\delta}) r^{-2}.$$

We deduce,

$$\left| \kappa^{\mathbf{S}(\infty)} \underline{\kappa}^{\mathbf{S}(\infty)} + \frac{4(1 - \frac{2m}{r})}{r^2} - \frac{2}{r} \check{\kappa} \right| \lesssim \overset{\circ}{\delta} r^{-3}.$$

Thus, in view of (9.4.38),

$$\int_{\mathbf{S}(\infty)} \kappa^{\mathbf{S}(\infty)} \underline{\kappa}^{\mathbf{S}(\infty)} = - \int_{\mathbf{S}(\infty)} \frac{4(1 - \frac{2m}{r})}{r^2} + O(\overset{\circ}{\delta}) r^{-1}.$$

Making use of the definition of the Hawking mass $m^{\mathbf{S}(\infty)} = \frac{r^{\mathbf{S}(\infty)}}{2} \left(1 + \frac{1}{16\pi} \int_{\mathbf{S}(\infty)} \kappa^{\mathbf{S}(\infty)} \underline{\kappa}^{\mathbf{S}(\infty)} \right)$ we easily deduce¹⁵

$$\left| m^{\mathbf{S}(\infty)} - m \right| \lesssim \overset{\circ}{\delta}. \quad (9.4.39)$$

¹⁴Here, we also use the fact that, on $S^{(\infty)}$, we have

$$|r - \overset{\circ}{r}| \lesssim \|(U^{(\infty)}, S^{(\infty)})\|_{L^\infty(\overset{\circ}{S})} \lesssim \overset{\circ}{\delta}.$$

¹⁵See also subsection 9.4.2.

Step 4. We make use of Lemma 9.3.11 to extend $(a^\infty, f^\infty, \underline{f}^\infty)$ as well as the frame $(e_3^{\mathbf{S}(\infty)}, e_4^{\mathbf{S}(\infty)}, e_\theta^{\mathbf{S}(\infty)})$ in a small neighborhood of $\mathbf{S}(\infty)$ such that we have,

$$\xi^{\mathbf{S}(\infty)} = 0, \quad \omega^{\mathbf{S}(\infty)} = 0, \quad \underline{\eta}^{\mathbf{S}(\infty)} + \zeta^{\mathbf{S}(\infty)} = 0, \tag{9.4.40}$$

and then provide estimates for the corresponding Ricci coefficients and curvature components $\check{\Gamma}^{\mathbf{S}(\infty)}, \check{R}^{\mathbf{S}(\infty)}$. More precisely we make use of the assumption **A1**, the estimates in (9.4.36) for $(a^\infty, f^\infty, \underline{f}^\infty)$, and the transformation formulae to derive the desired estimates (9.4.11) for s_{max} derivative of the Ricci coefficients and curvature components of $\mathbf{S}(\infty)$.

Step 5. Thanks to (9.4.35), we can pass to the limit in (9.4.25) (9.4.29). We deduce that all equations recorded in Proposition 9.4.1 hold true and thus that all the desired GCMS hold true.

9.5 Proof of Proposition 9.4.4

Since all estimates below take place on \mathbf{S} , we simplify our notation and denote the norms $\|\cdot\|_{\mathfrak{h}_k(\mathbf{S})}$ simply by $\|\cdot\|_k$ in what follows¹⁶. In the particular case $k = 0$ we also write $\|\cdot\|_0 = \|\cdot\|_{\mathbf{S}}$. The sup norms $\|\cdot\|_{\mathfrak{h}_k^\infty(\mathbf{S})}$, though rarely needed, will be denoted by $|\cdot|_k$. Since r and $r^{\mathbf{S}}$ are comparable we will freely choose one or the other throughout the proof.

We introduce the notation (recall that $a = \log \lambda$)

$$\begin{aligned} \|F\|_k &= \|a\|_k + \|f\|_k + \|\underline{f}\|_k, \\ I(f, \underline{f}) &= \left| \int_{\mathbf{S}} f e^\Phi \right| + \left| \int_{\mathbf{S}} \underline{f} e^\Phi \right|, \\ I(a, f, \underline{f}) &= I(f, \underline{f}) + r^{\mathbf{S}} \left| \int_{\mathbf{S}} (e_\theta a) e^\Phi \right|. \end{aligned} \tag{9.5.1}$$

We assume that a solution exists verifying the auxiliary bootstrap assumption¹⁷

$$\|(a, f, \underline{f})\|_{s_{max}+1} \lesssim \sqrt{\overset{\circ}{\epsilon}} \tag{9.5.2}$$

based on which we will prove the stronger estimate,

$$\|(a, f, \underline{f})\|_{s_{max}+1} \lesssim \overset{\circ}{\delta}. \tag{9.5.3}$$

¹⁶To remind the reader about our convention we shall in fact alternately use both notations.

¹⁷This is not really needed, we only use it to simplify the exposition.

Step 1. We write the first equation in (9.4.2) in the form

$$\begin{aligned} (A^S + V)z &= \frac{2}{r} (\not{d}^{\mathbf{S}, \star \underline{\kappa}} \mathbf{S} - \not{d}^{\star \underline{\kappa}}) - \frac{2\Upsilon}{r} \not{d}^{\star \check{\kappa}} + \text{Err}(z), \\ \text{Err}(z) &:= \underline{\kappa} \text{Err}_4 + \kappa \text{Err}_5 + [A^S, \underline{\kappa}]f + [A^S, \kappa]f \\ &\quad + \left(\kappa - \frac{2}{r} \right) (\not{d}^{\mathbf{S}, \star \check{\kappa}} \mathbf{S} - \not{d}^{\star \check{\kappa}}) + \left(\underline{\kappa} + \frac{2\Upsilon}{r} \right) \not{d}^{\star \check{\kappa}}. \end{aligned}$$

Thus,

$$\|(A^S + V)z\|_k \lesssim r^{-1} \|\not{d}^{\mathbf{S}, \star \check{\kappa}} \mathbf{S} - \not{d}^{\star \check{\kappa}}\|_k + r^{-1} \|\not{d}^{\star \check{\kappa}}\|_k + \|\text{Err}(z)\|_k.$$

In view of Lemma 9.3.10 and the auxiliary bootstrap assumption (9.5.2) we have¹⁸, for $k \leq s_{max} - 1$,

$$\begin{aligned} \|\text{Err}_4, \text{Err}_5\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim r^{-2} \|f, \underline{f}, a\|_{\mathfrak{h}_{k+2}(\mathbf{S})} \left(\overset{\circ}{\epsilon} + \|f, \underline{f}, a\|_{\mathfrak{h}_{k+2}(\mathbf{S})} \right) \\ &\lesssim r^{-2} \sqrt{\overset{\circ}{\epsilon}} \|F\|_{k+2}. \end{aligned}$$

Also,

$$\|[A^S, \underline{\kappa}]f + [A^S, \kappa]f\|_{\mathfrak{h}_k(\mathbf{S})} \lesssim r^{-3} \sqrt{\overset{\circ}{\epsilon}} \|(f, \underline{f})\|_{\mathfrak{h}_{k+1}(\mathbf{S})}.$$

In view of (9.4.5) we have $\|\check{\kappa}, \bar{\kappa} - \frac{2}{r}\|_{\mathfrak{h}_k(\mathbf{S})} \lesssim \overset{\circ}{\delta} r^{-1}$ and $\|\not{d}^{\star \check{\kappa}}\|_{\mathfrak{h}_{k-1}(\mathbf{S})} \lesssim \overset{\circ}{\delta} r^{-2}$ for all $k \leq s_{max}$, and thus

$$\begin{aligned} \left\| \left(\kappa - \frac{2}{r} \right) (\not{d}^{\mathbf{S}, \star \check{\kappa}} \mathbf{S} - \not{d}^{\star \check{\kappa}}) \right\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\delta} r^{-2} \|\not{d}^{\mathbf{S}, \star \check{\kappa}} \mathbf{S} - \not{d}^{\star \check{\kappa}}\|_{\mathfrak{h}_k(\mathbf{S})}, \\ \left\| \left(\underline{\kappa} + \frac{2\Upsilon}{r} \right) \not{d}^{\star \check{\kappa}} \right\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\delta} \epsilon r^{-4}, \end{aligned}$$

i.e.,

$$\|\text{Err}(z)\|_k \lesssim r^{-3} \sqrt{\overset{\circ}{\epsilon}} \|F\|_{k+2} + \overset{\circ}{\delta} r^{-2} \|\not{d}^{\mathbf{S}, \star \check{\kappa}} \mathbf{S} - \not{d}^{\star \check{\kappa}}\|_{\mathfrak{h}_k(\mathbf{S})} + \overset{\circ}{\delta} \epsilon r^{-4}.$$

Hence,

$$\|(A^S + V)z\|_k \lesssim r^{-1} \|\not{d}^{\mathbf{S}, \star \check{\kappa}} \mathbf{S} - \not{d}^{\star \check{\kappa}}\|_k + r^{-3} \overset{\circ}{\delta} + r^{-3} \sqrt{\overset{\circ}{\epsilon}} \|F\|_{k+2}. \quad (9.5.4)$$

¹⁸We also assume tacitly, throughout the estimates below, that $k \geq s_{max}/2 + 2$ so that we have product estimates in $\|\cdot\|_k$.

To estimate $\|\phi^{\mathbf{S},*}\check{\underline{\kappa}}^{\mathbf{S}} - \phi^*\check{\underline{\kappa}}\|_k$ we proceed as follows. Note, using the GCMS conditions, that

$$\phi_2^{\mathbf{S},*} \left(\phi_1^{\mathbf{S},*}\check{\underline{\kappa}}^{\mathbf{S}} - \phi_1^*\check{\underline{\kappa}} \right) = -\phi_2^{\mathbf{S},*}\phi_1^*\check{\underline{\kappa}} = -\phi_2^*\phi_1^*\check{\underline{\kappa}} - (\phi_2^{\mathbf{S},*} - \phi_2^*)\phi_1^*\check{\underline{\kappa}}.$$

According to the first inequality in (9.4.5) $\|\phi_2^*\phi_1^*\check{\underline{\kappa}}\|_{\mathfrak{h}_k(\mathbf{s})} \lesssim \delta r^{-3}$. Hence,

$$\begin{aligned} \left\| \phi_2^{\mathbf{S},*} \left(\phi_1^{\mathbf{S},*}\check{\underline{\kappa}}^{\mathbf{S}} - \phi_1^*\check{\underline{\kappa}} \right) \right\|_k &\lesssim \|\phi_2^*\phi_1^*\check{\underline{\kappa}}\|_k + \|(\phi_2^{\mathbf{S},*} - \phi_2^*)\phi_1^*\check{\underline{\kappa}}\|_k \\ &\lesssim \delta r^{-3} + \|(\phi_2^{\mathbf{S},*} - \phi_2^*)\phi_1^*\check{\underline{\kappa}}\|_k. \end{aligned}$$

Recalling the transformation formulas (9.2.18) we have schematically,

$$\phi_2^{\mathbf{S},*} - \left(1 + \frac{1}{2}f\underline{f}\right)\phi_2^* = \frac{1}{2}\underline{f}e_4 + \frac{1}{2}f\left(1 + \frac{1}{4}f\underline{f}\right)e_3 + \text{l.o.t.}$$

Therefore,

$$\|(\phi_2^{\mathbf{S},*} - \phi_2^*)\phi_1^*\check{\underline{\kappa}}\|_k \lesssim r^{-3}\sqrt{\overset{\circ}{\epsilon}}\|(f, \underline{f})\|_k$$

and thus setting

$$B := \phi_1^{\mathbf{S},*}\check{\underline{\kappa}}^{\mathbf{S}} - \phi_1^*\check{\underline{\kappa}}, \tag{9.5.5}$$

we have

$$\|\phi_2^{\mathbf{S},*}B\|_k \lesssim r^{-3}\left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}}\|(f, \underline{f})\|_k\right). \tag{9.5.6}$$

According to the elliptic estimate given by Lemma 9.1.4 we have,

$$\|B\|_k \lesssim r\|\phi_2^*B\|_k + r^{-2}\left|\int_{\mathbf{S}} Be^\Phi\right|$$

i.e.,

$$\|B\|_k \lesssim r^{-2}\left|\int_{\mathbf{S}} Be^\Phi\right| + r^{-2}\left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}}\|(f, \underline{f})\|_k\right). \tag{9.5.7}$$

To estimate the $\ell = 1$ mode of B , i.e. its projection on the kernel of ϕ_2^* , we recall the transformation formula, see Lemma 9.3.4,

$$e'_\theta(\underline{\kappa}') = e_{\theta\underline{\kappa}} + e'_\theta\underline{\phi}_1\underline{f} - \underline{\kappa}e'_\theta a - \frac{1}{4}\underline{\kappa}(f\underline{\kappa} + \underline{f}\kappa) - \underline{\kappa}\omega\underline{f} + \underline{f}\rho + \text{Err}(e'_\theta\underline{\kappa}', e_{\theta\underline{\kappa}})$$

i.e.,

$$\begin{aligned} B &= \underline{d}_1^{\mathbf{S},*} \underline{d}_1^{\mathbf{S}} f + \underline{\kappa} \underline{d}_1^{\mathbf{S}} f + \left(\frac{1}{4} \underline{\kappa}^2 - \underline{\kappa} \underline{\omega} \right) f + \left(\frac{1}{4} \underline{\kappa} \underline{\kappa} - \rho \right) \underline{f} - \text{Err}(e'_\theta \underline{\kappa}', e_\theta \underline{\kappa}) \\ &= A^{\mathbf{S}} f + \underline{\kappa} \underline{d}_1^{\mathbf{S}} f + c_1^{\mathbf{S}} f + c_2^{\mathbf{S}} \underline{f} + \text{Err}(B) \end{aligned}$$

where,

$$c_1^{\mathbf{S}} = \left(\frac{\Upsilon^{\mathbf{S}}}{r^{\mathbf{S}}} \right)^2 - \frac{2m^{\mathbf{S}}}{(r^{\mathbf{S}})^3}, \quad c_2^{\mathbf{S}} = -\frac{\Upsilon^{\mathbf{S}}}{(r^{\mathbf{S}})^2} + \frac{2m^{\mathbf{S}}}{(r^{\mathbf{S}})^3}$$

are constants on S verifying

$$\left| (c_1^{\mathbf{S}}, c_2^{\mathbf{S}}) \right| \lesssim r^{-2}.$$

We also denote,

$$c_1 = \overline{\left(\frac{1}{4} \underline{\kappa}^2 - \underline{\kappa} \underline{\omega} \right)} = \left(\frac{\Upsilon}{r} \right)^2 - \frac{2m}{r^3} \quad c_2 = \overline{\left(\frac{1}{4} \underline{\kappa} \underline{\kappa} - \rho \right)} = -\frac{\Upsilon}{r^2} + \frac{2m}{r^3}.$$

The error term $\text{Err}(B)$ is given by

$$\text{Err}(B) = -\text{Err}(e'_\theta \underline{\kappa}', e_\theta \underline{\kappa}) + f \left[\left(\frac{1}{4} \underline{\kappa}^2 - \underline{\kappa} \underline{\omega} \right) - c_1 \right] + \underline{f} \left[\left(\frac{1}{4} \underline{\kappa} \underline{\kappa} - \rho \right) - c_2 \right].$$

We deduce,

$$\left| \int_{\mathbf{S}} B e^\Phi \right| \lesssim \left| \int_{\mathbf{S}} A^{\mathbf{S}} f e^\Phi \right| + r^{-2} \left(\left| \int_{\mathbf{S}} f e^\Phi \right| + \left| \int_{\mathbf{S}} \underline{f} e^\Phi \right| \right) + \left| \int_{\mathbf{S}} \text{Err}(B) e^\Phi \right|. \quad (9.5.8)$$

We can easily check, using the assumptions of the theorem,

$$\begin{aligned} \left| \frac{1}{4} \underline{\kappa}^2 - \underline{\kappa} \underline{\omega} - \overline{\left(\frac{1}{4} \underline{\kappa}^2 - \underline{\kappa} \underline{\omega} \right)} \right| &\lesssim \overset{\circ}{\epsilon} r^{-3}, \\ \left| \frac{1}{4} \underline{\kappa} \underline{\kappa} - \rho - \overline{\left(\frac{1}{4} \underline{\kappa} \underline{\kappa} - \rho \right)} \right| &\lesssim \overset{\circ}{\epsilon} r^{-3}, \\ \left| c_1^{\mathbf{S}} - c_1 \right| + \left| c_2^{\mathbf{S}} - c_2 \right| &\lesssim \overset{\circ}{\epsilon} r^{-2}. \end{aligned}$$

For the last inequality we note that

$$\left| \frac{1}{(r^{\mathbf{S}})^2} - \frac{1}{r^2} \right| \lesssim (r^{\mathbf{S}})^{-2} \left| \frac{r^{\mathbf{S}}}{r} - 1 \right| \left| \frac{r^{\mathbf{S}}}{r} + 1 \right| \lesssim r^{-2} \overset{\circ}{\epsilon}.$$

Thus, by the estimate¹⁹ for Err_2 in Lemma 9.3.10,

$$\left| \int_{\mathbf{S}} \text{Err}(B)e^\Phi \right| \lesssim r^2 \|\text{Err}(B)\|_{\mathbf{S}} \lesssim r^2 \|\text{Err}_2\|_{\mathbf{S}} + \|(f, \underline{f})\|_{\mathbf{S}} \lesssim \sqrt{\overset{\circ}{\epsilon}} \|F\|_2.$$

Consequently,

$$\left| \int_{\mathbf{S}} B e^\Phi \right| \lesssim \left| \int_{\mathbf{S}} A^{\mathbf{S}} f e^\Phi \right| + r^{-2} \left(\left| \int_{\mathbf{S}} f e^\Phi \right| + \left| \int_{\mathbf{S}} \underline{f} e^\Phi \right| \right) + \sqrt{\overset{\circ}{\epsilon}} \|F\|_2.$$

We can evaluate the integral $\int_{\mathbf{S}} A^{\mathbf{S}} f e^\Phi$ by using the identity $A^{\mathbf{S}} = \not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} + K$. Thus,

$$\begin{aligned} \int_{\mathbf{S}} A^{\mathbf{S}} f e^\Phi &= \int_{\mathbf{S}} \not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} f e^\Phi + \int_{\mathbf{S}} K f e^\Phi \\ &= \int_{\mathbf{S}} \not{d}_2^{\mathbf{S},*} f \not{d}_2^{\mathbf{S},*}(e^\Phi) + \int_{\mathbf{S}} K f e^\Phi = \int_{\mathbf{S}} K f e^\Phi \\ &= (r^{\mathbf{S}})^{-2} \int_{\mathbf{S}} f e^\Phi + \int_{\mathbf{S}} \left(\left(K - \frac{1}{r^2} \right) + \frac{1}{r^2} - \frac{1}{(r^{\mathbf{S}})^2} \right) f e^\Phi. \end{aligned}$$

Hence,

$$\left| \int_{\mathbf{S}} A^{\mathbf{S}} f e^\Phi \right| \lesssim r^{-2} \left| \int_{\mathbf{S}} f e^\Phi \right| + \sqrt{\overset{\circ}{\epsilon}} \|f\|_0.$$

Therefore,

$$\left| \int_{\mathbf{S}} B e^\Phi \right| \lesssim r^{-2} \left(\left| \int_{\mathbf{S}} f e^\Phi \right| + \left| \int_{\mathbf{S}} \underline{f} e^\Phi \right| \right) + \sqrt{\overset{\circ}{\epsilon}} \|F\|_2. \tag{9.5.9}$$

Recalling (9.5.7),

$$\begin{aligned} \|B\|_k &\lesssim r^{-2} \left| \int_{\mathbf{S}} B e^\Phi \right| + r^{-2} \left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_k \right) \\ &\lesssim r^{-4} \left(\left| \int_{\mathbf{S}} f e^\Phi \right| + \left| \int_{\mathbf{S}} \underline{f} e^\Phi \right| \right) + r^{-2} \left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_k \right). \end{aligned}$$

Returning to (9.5.4) and recalling the definition of $B = \not{d}^{\mathbf{S},*} \check{\kappa}^{\mathbf{S}} - \not{d}^* \check{\kappa}$,

$$\begin{aligned} \|(A^{\mathbf{S}} + V)z\|_k &\lesssim r^{-1} \|B\|_k + r^{-3} \overset{\circ}{\delta} + r^{-3} \sqrt{\overset{\circ}{\epsilon}} \|F\|_{k+2} \\ &\lesssim r^{-5} \left(\left| \int_{\mathbf{S}} f e^\Phi \right| + \left| \int_{\mathbf{S}} \underline{f} e^\Phi \right| \right) + r^{-3} \left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_{k+2} \right) \end{aligned}$$

¹⁹Note that $\text{Err}(e'_{\theta \underline{\kappa}'}, e_{\theta \underline{\kappa}}) = \text{Err}_2 + \text{l.o.t.}$ from the definition of Err_2 .

or, recalling the definition of $I(f, \underline{f})$,

$$\|(A^{\mathbf{S}} + V)z\|_k \lesssim r^{-5}I(f, \underline{f}) + r^{-3} \left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_{k+2} \right). \quad (9.5.10)$$

To estimate z we recall that,

$$A^{\mathbf{S}} + V = \not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} - 3\rho + \frac{1}{2}\not{v}\not{v}.$$

Thus,

$$\left((A^{\mathbf{S}} + V)z, z \right)_{\mathbf{S}} = \|\not{d}_s^{\mathbf{S},*}z\|_{\mathbf{S}}^2 + \left(\frac{6m}{r^3} + O(r^{-3}\sqrt{\overset{\circ}{\epsilon}}) \right) \|z\|_{\mathbf{S}}.$$

We deduce,

$$\|\not{d}_s^{\mathbf{S},*}z\|_{\mathbf{S}}^2 \lesssim \left((A^{\mathbf{S}} + V)z, z \right)_{\mathbf{S}} \lesssim \|(A^{\mathbf{S}} + V)z\|_{\mathbf{S}} \|z\|_{\mathbf{S}}.$$

We make use of the elliptic estimate, see Lemma 9.1.4,

$$\begin{aligned} \|z\|_{\mathbf{S}}^2 &\lesssim r^2 \|\not{d}_s^{\mathbf{S},*}z\|_{\mathbf{S}}^2 + r^{-4} \left| \int_{\mathbf{S}} z e^{\Phi} \right|^2 \\ &\lesssim r^2 \|(A^{\mathbf{S}} + V)z\|_{\mathbf{S}} \|z\|_{\mathbf{S}} + r^{-4} \left| \int_{\mathbf{S}} z e^{\Phi} \right|^2 \end{aligned}$$

from which we infer that

$$\|z\|_{\mathbf{S}} \lesssim r^2 \|(A^{\mathbf{S}} + V)z\|_{\mathbf{S}} + r^{-2} \left| \int_{\mathbf{S}} z e^{\Phi} \right|. \quad (9.5.11)$$

Recalling the definition of z ,

$$\begin{aligned} z &= \kappa \underline{f} + \underline{\kappa} f = \frac{2}{r} \underline{f} - \frac{2\Upsilon}{r} f + \left(\kappa - \frac{2}{r} \right) \underline{f} + \left(\underline{\kappa} + \frac{2\Upsilon}{r} \right) f \\ &= \frac{2}{r^{\mathbf{S}}} \underline{f} - \frac{2\Upsilon^{\mathbf{S}}}{r^{\mathbf{S}}} f - \left(\frac{2}{r^{\mathbf{S}}} - \frac{2}{r} \right) \underline{f} + \left(\frac{2\Upsilon^{\mathbf{S}}}{r^{\mathbf{S}}} - \frac{2\Upsilon}{r} \right) f + \left(\kappa - \frac{2}{r} \right) \underline{f} + \left(\underline{\kappa} + \frac{2\Upsilon}{r} \right) f, \end{aligned}$$

and making use of our assumption (in particular $\left| \frac{r^{\mathbf{S}}}{r} - 1 \right| \lesssim r^{-1}\overset{\circ}{\epsilon}$) we deduce,

$$\left| \int_{\mathbf{S}} z e^{\Phi} - \frac{2}{r^{\mathbf{S}}} \int_{\mathbf{S}} \underline{f} e^{\Phi} + \frac{2\Upsilon^{\mathbf{S}}}{r^{\mathbf{S}}} \int_{\mathbf{S}} f e^{\Phi} \right| \lesssim \sqrt{\overset{\circ}{\epsilon}} r (\|f\|_0 + \|\underline{f}\|_0).$$

Hence,

$$\left| \int_{\mathbf{S}} z e^{\Phi} \right| \lesssim r^{-1} I(f, \underline{f}) + \sqrt{\overset{\circ}{\epsilon}} r \|F\|_0$$

and therefore in view of (9.5.11) and (9.5.10) (for $k = 0$)

$$\begin{aligned} \|z\|_0 &\lesssim r^2 \|(A^{\mathbf{S}} + V)z\|_0 + r^{-2} \left| \int_{\mathbf{S}} z e^{\Phi} \right| \\ &\lesssim r^2 \left(r^{-5} I(f, \underline{f}) + r^{-3} \left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_2 \right) \right) + r^{-2} \left(r^{-1} I(f, \underline{f}) + \sqrt{\overset{\circ}{\epsilon}} r \|F\|_0 \right) \\ &\lesssim r^{-3} I(f, \underline{f}) + r^{-1} \left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_2 \right). \end{aligned}$$

By standard elliptic theory,

$$\|z\|_{k+2} \lesssim \|z\|_0 + r^2 \|(A^{\mathbf{S}} + V)z\|_k.$$

We infer that, for $k \leq s_{max} - 1$,

$$\|z\|_{k+2} \lesssim r^{-3} I(f, \underline{f}) + r^{-1} \left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_{2+k} \right). \quad (9.5.12)$$

Step 2. We now proceed in the same manner to derive an estimate²⁰ for f . Using the second equation in (9.4.2)

$$(A^{\mathbf{S}} + V) f = \frac{3}{4} \kappa (A^{\mathbf{S}})^{-1} (\rho z) - \not{d}^{\star} \check{\kappa} - \kappa (A^{\mathbf{S}})^{-1} (-\not{d}^{\mathbf{S}, \star} \check{\mu}^{\mathbf{S}} + \not{d}^{\star} \check{\mu}) + \text{Err}_4,$$

we derive (note that $\|(A^{\mathbf{S}})^{-1} h\|_k \lesssim r^2 \|h\|_{k-2}$),

$$\begin{aligned} \|(A^{\mathbf{S}} + V) f\|_k &\lesssim r \|\rho z\|_{k-2} + \|\not{d}^{\star} \check{\kappa}\|_k + r \|\not{d}^{\mathbf{S}, \star} \check{\mu}^{\mathbf{S}} - \not{d}^{\star} \check{\mu}\|_{k-2} + \|\text{Err}_4\|_k + \sqrt{\overset{\circ}{\epsilon}} r^{-2} \|F\|_k \\ &\lesssim r^{-2} \|z\|_{k-2} + \overset{\circ}{\delta} r^{-2} + r \|C\|_{k-2} + \sqrt{\overset{\circ}{\epsilon}} r^{-2} \|F\|_{k+2} \end{aligned} \quad (9.5.13)$$

where,

$$C := \not{d}^{\mathbf{S}, \star} \check{\mu}^{\mathbf{S}} - \not{d}^{\star} \check{\mu}. \quad (9.5.14)$$

To estimate C we write, proceeding as in the estimate for B is Step 1. Making use of the GCM condition $\not{d}_2^{\mathbf{S}, \star} \not{d}_1^{\mathbf{S}, \star} \mu^{\mathbf{S}} = 0$ we derive, as in (9.5.6),

$$\|\not{d}_2^{\mathbf{S}, \star} C\|_k \lesssim r^{-4} \left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_k \right) \quad (9.5.15)$$

²⁰Since most estimates are similar as in Step 1 above we skip some of the intermediary steps.

which implies, as in (9.5.7),

$$\|C\|_k \lesssim r^{-2} \left| \int_{\mathbf{S}} C e^\Phi \right| + r^{-3} \left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_k \right). \quad (9.5.16)$$

To estimate the $\ell = 1$ mode of C we recall the transformation formula, see Lemma 9.3.4,

$$\begin{aligned} e'_\theta(\mu') &= e_{\theta\mu} + e'_\theta(\not{d}_1)' \left(-(\not{d}_1^*)' a + f\underline{\omega} - \underline{f}\omega + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) + \frac{3}{4}\rho(f\underline{\kappa} + \underline{f}\kappa) \\ &\quad + \text{Err}(e'_\theta\mu', e_{\theta\mu}), \end{aligned}$$

i.e.,

$$C = A^{\mathbf{S}} \left(-\not{d}_1^{\mathbf{S},*} a + f\underline{\omega} + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) - \frac{3}{4}\rho z + \text{Err}(e'_\theta\mu', e_{\theta\mu}).$$

Recall that,

$$A^{\mathbf{S}} = \not{d}_1^{\mathbf{S}} \not{d}_1^{\mathbf{S},*} = \not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} + K.$$

Therefore, for any $h \in \mathfrak{s}_1$,

$$\int_{\mathbf{S}} A^{\mathbf{S}} h e^\Phi = \int_{\mathbf{S}} h K e^\Phi.$$

We deduce,

$$\begin{aligned} \int_{\mathbf{S}} C e^\Phi &= \int_{\mathbf{S}} K \left(-\not{d}_1^{\mathbf{S},*} a + f\underline{\omega} + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) e^\Phi - \frac{3}{4} \int_{\mathbf{S}} (\rho z) e^\Phi + \int_{\mathbf{S}} \text{Err}(e'_\theta\mu', e_{\theta\mu}) e^\Phi \\ &= (r^{\mathbf{S}})^{-2} \int_{\mathbf{S}} \left(-\not{d}_1^{\mathbf{S},*} a + f\underline{\omega} + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) e^\Phi - \frac{3}{4} \int_{\mathbf{S}} (\rho z) e^\Phi + \int_{\mathbf{S}} \text{Err}(e'_\theta\mu', e_{\theta\mu}) e^\Phi \\ &\quad + \int_{\mathbf{S}} ((K - r^{-2}) + (r^{-2} - (r^{\mathbf{S}})^{-2})) \left(-\not{d}_1^{\mathbf{S},*} a + f\underline{\omega} + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) e^\Phi. \end{aligned}$$

Also,

$$\begin{aligned} \int_{\mathbf{S}} \left(f\underline{\omega} + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa \right) e^\Phi &= \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^2} \int_{\mathbf{S}} f e^\Phi - \frac{\Upsilon^{\mathbf{S}}}{2r^{\mathbf{S}}} \int_{\mathbf{S}} f e^\Phi - \frac{1}{2r^{\mathbf{S}}} \int_{\mathbf{S}} \underline{f} e^\Phi + \text{l.o.t.} \\ \int_{\mathbf{S}} \rho z e^\Phi &= -\frac{2m^{\mathbf{S}}}{(r^{\mathbf{S}})^3} \int_{\mathbf{S}} z e^\Phi + \text{l.o.t.} \end{aligned}$$

Note that, since $\text{Err}(e'_\theta\mu', e_{\theta\mu}) = \text{Err}_3$, we have in view of Lemma 9.3.10,

$$\left| \int_{\mathbf{S}} \text{Err}(e'_\theta\mu', e_{\theta\mu}) e^\Phi \right| \lesssim r^2 \|\text{Err}_3\|_0 \lesssim r^{-1} \|F\|_3.$$

We deduce,

$$\left| \int_{\mathbf{S}} C e^{\Phi} \right| \lesssim r^{-3} I(a, f, \underline{f}) + r^{-1} \|F\|_3. \quad (9.5.17)$$

Back to (9.5.16) we infer that,

$$\|C\|_k \lesssim r^{-5} I(a, f, \underline{f}) + r^{-3} \left(\overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_k \right). \quad (9.5.18)$$

Back to (9.5.13),

$$\begin{aligned} \|(A^{\mathbf{S}} + V) f\|_k &\lesssim r^{-2} \|z\|_{k-2} + \overset{\circ}{\delta} r^{-2} + r \|C\|_{k-2} + \sqrt{\overset{\circ}{\epsilon}} r^{-2} \|F\|_{k+2} \\ &\lesssim r^{-2} \|z\|_{k-2} + \overset{\circ}{\delta} r^{-2} + \sqrt{\overset{\circ}{\epsilon}} r^{-2} \|F\|_{k+2} + r^{-4} I(a, f, \underline{f}) \end{aligned}$$

i.e., recalling the estimate (9.5.12) for $\|z\|_k$ in Step 1,

$$\|(A^{\mathbf{S}} + V) f\|_k \lesssim \overset{\circ}{\delta} r^{-2} + \sqrt{\overset{\circ}{\epsilon}} r^{-2} \|F\|_{k+2} + r^{-4} I(a, f, \underline{f}). \quad (9.5.19)$$

To estimate $\|f\|_0 = \|f\|_{\mathbf{S}}$ we proceed as in the proof of the estimate for $\|z\|_0$ in Step 1 and deduce, as in (9.5.11),

$$\|f\|_{\mathbf{S}} \lesssim r^2 \|(A^{\mathbf{S}} + V) f\|_{\mathbf{S}} + r^{-2} \left| \int_{\mathbf{S}} f e^{\Phi} \right|. \quad (9.5.20)$$

Hence,

$$\|f\|_0 \lesssim \overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_2 + r^{-2} I(a, f, \underline{f}) + r^{-2} \left| \int_{\mathbf{S}} f e^{\Phi} \right|$$

i.e.,

$$\|f\|_0 \lesssim \overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_2 + r^{-2} I(a, f, \underline{f}).$$

Finally,

$$\begin{aligned} \|f\|_{k+2} &\lesssim \|f\|_0 + r^2 \|(A^{\mathbf{S}} + V) f\|_k \\ &\lesssim \overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_{2+k} + r^{-2} I(a, f, \underline{f}). \end{aligned}$$

Hence,

$$\|f\|_{k+2} \lesssim \overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_{k+2} + r^{-2} I(a, f, \underline{f}). \quad (9.5.21)$$

Since by the definition of z we have,

$$\underline{f} = \kappa^{-1}(z - \underline{\kappa}f)$$

we also derive,

$$\|\underline{f}\|_{k+2} \lesssim r^{-1}\|z\|_{k+2} + \|f\|_{k+2}.$$

Thus, making use of the estimate (9.5.12) for z , we also deduce the estimate for \underline{f} ,

$$\|\underline{f}\|_{k+2} \lesssim \delta^\circ + \sqrt{\epsilon^\circ} \|F\|_{k+2} + r^{-2}I(a, f, \underline{f})$$

i.e.,

$$\|(f, \underline{f})\|_{k+2} \lesssim \delta^\circ + \sqrt{\epsilon^\circ} \|F\|_{k+2} + r^{-2}I(a, f, \underline{f}). \quad (9.5.22)$$

Step 3. It remains to estimate a by making use of the equation (see Proposition 9.4.2 and definition of $C = \not{d}^{\mathbf{S},*} \check{\mu}^{\mathbf{S}} - \not{d}^* \check{\mu}$ in (9.5.14))

$$\not{d}^{\mathbf{S},*} a = -\frac{3}{4}(A^{\mathbf{S}})^{-1}(\rho z) + f\underline{\omega} + \frac{1}{4}f\underline{\kappa} - \frac{1}{4}\underline{f}\kappa - (A^{\mathbf{S}})^{-1} C - (A^{\mathbf{S}})^{-1}\text{Err}_3.$$

Proceeding as before we deduce,

$$\|\not{d}^{\mathbf{S},*} a\|_k \lesssim r^{-1}\|z\|_{k-2} + r^{-1}\|(f, \underline{f})\|_k + r^2\|C\|_{k-2} + r^2\|\text{Err}_3\|_{k-2}.$$

Making use of (9.5.12) and,(9.5.18) and the estimate of Lemma 9.3.10 for Err_3 ,

$$\begin{aligned} \|z\|_{k-2} &\lesssim r^{-3}I(f, \underline{f}) + r^{-1} \left(\delta^\circ + \sqrt{\epsilon^\circ} \|F\|_{k-2} \right), \\ \|C\|_{k-2} &\lesssim r^{-5}I(a, f, \underline{f}) + r^{-3} \left(\delta^\circ + \sqrt{\epsilon^\circ} \|F\|_{k-2} \right), \\ \|\text{Err}_3\|_{k-2} &\lesssim r^{-3} \sqrt{\epsilon^\circ} \|F\|_{k+1}, \end{aligned}$$

we further deduce,

$$\|\not{d}^{\mathbf{S},*} a\|_k \lesssim r^{-1}(\delta^\circ + \sqrt{\epsilon^\circ} \|F\|_{k+1}) + r^{-3}I(a, f, \underline{f}). \quad (9.5.23)$$

Combining this with (9.5.22),

$$r^{\mathbf{S}} \|\not{d}^{\mathbf{S},*} a\|_{k+1} + \|(f, \underline{f})\|_{k+2} \lesssim \delta^\circ + \sqrt{\epsilon^\circ} \|F\|_{k+2} + r^{-2}I(a, f, \underline{f}). \quad (9.5.24)$$

Step 4. We make use of the equations (9.4.3) to estimate $I(a, f, \underline{f})$. Recall that,

$$I(a, f, \underline{f}) = I(f, \underline{f}) + r^{\mathbf{S}} \left| \int_{\mathbf{S}} (e_{\theta}^{\mathbf{S}} a) e^{\Phi} \right|.$$

Step 4a. We estimate the integral $\int_{\mathbf{S}} (e_{\theta}^{\mathbf{S}} a) e^{\Phi}$ making use of the the first equation²¹ in (9.4.3).

$$\begin{aligned} \frac{2}{r^{\mathbf{S}}} \int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(a) e^{\Phi} &= - \int_{\mathbf{S}} e_{\theta}(\kappa) e^{\Phi} + \frac{1}{4} \int_{\mathbf{S}} \kappa^2 \underline{f} e^{\Phi} - \int_{\mathbf{S}} \left(\frac{1}{4} \kappa \underline{\kappa} + \kappa \underline{\omega} + 3\rho \right) f e^{\Phi} + \text{Err}_7 \\ &= -J_1 + J_2 - J_3 + \text{Err}_7 \end{aligned}$$

where,

$$\text{Err}_7 = \int_{\mathbf{S}} \left(-\text{Err}(e_{\theta}^{\mathbf{S}} \kappa^{\mathbf{S}}, e_{\theta} \kappa) + 2(K^{\mathbf{S}} - K)f + \left(\kappa - \frac{2}{r^{\mathbf{S}}} \right) \not{d}^{\mathbf{S}, \star} a + \frac{1}{2} \vartheta \underline{\vartheta} f \right) e^{\Phi}.$$

We deduce,

$$\left| \int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(a) e^{\Phi} \right| \lesssim r (|J_1| + |J_2| + |J_3|) + r |\text{Err}_6|.$$

We make use of

$$e_{\theta}^{\mathbf{S}} = \left(1 + \frac{1}{2} f \underline{f} \right) e_{\theta} + \frac{1}{2} \underline{f} e_4 + \frac{1}{2} f \left(1 + \frac{1}{4} f \underline{f} \right) e_3$$

and the assumptions of Theorem 9.4.4, in particular, $\|e_{\theta}^{\mathbf{S}} \kappa\|_{\mathbf{S}} \lesssim \mathring{\delta} r^{-2}$, to deduce,

$$|J_1| \lesssim r \int_{\mathbf{S}} (|e_{\theta}^{\mathbf{S}}(\kappa)| + |e_{\theta}^{\mathbf{S}}(\kappa) - e_{\theta}(\kappa)|) \lesssim \mathring{\delta} + \sqrt{\mathring{\epsilon}} \|(f, \underline{f})\|_{\mathbf{S}}.$$

Also,

$$\begin{aligned} |J_2| &\lesssim \frac{1}{(r^{\mathbf{S}})^2} \left| \int_{\mathbf{S}} \underline{f} e^{\Phi} \right| + r \int_{\mathbf{S}} |\underline{f}| \left[\left| \kappa^2 - \frac{4}{r^2} \right| + \left| \frac{4}{(r^{\mathbf{S}})^2} - \frac{4}{r^2} \right| \right] \\ &\lesssim r^{-2} \left| \int_{\mathbf{S}} \underline{f} e^{\Phi} \right| + \sqrt{\mathring{\epsilon}} \|f\|_{\mathbf{S}}. \end{aligned}$$

Similarly,

$$|J_3| \lesssim r^{-2} \left| \int_{\mathbf{S}} f e^{\Phi} \right| + \sqrt{\mathring{\epsilon}} \|f\|_{\mathbf{S}}.$$

²¹recall that $\omega = 0$.

Also,

$$|\text{Err}_7| \lesssim r^2 \|\text{Err}(e'_\theta \kappa', e_\theta \kappa)\|_{\mathbf{S}} + r \left(\|K^{\mathbf{S}} - K\|_{\mathbf{S}} \|f\|_{\mathbf{S}} + \|\kappa - \frac{2}{r^{\mathbf{S}}}\|_{\mathbf{S}} \|\not{d}^{\mathbf{S},*} a\|_{\mathbf{S}} + \|\vartheta \underline{\vartheta}\|_{\mathbf{S}} \|f\|_{\mathbf{S}} \right).$$

Thus, proceeding as before,

$$|\text{Err}_7| \lesssim \sqrt{\overset{\circ}{\epsilon}} \|F\|_2.$$

We deduce,

$$\begin{aligned} \left| \int_{\mathbf{S}} e_\theta^{\mathbf{S}}(a) e^\Phi \right| &\lesssim r (|J_1| + |J_2| + |J_3|) + r |\text{Err}_6| \\ &\lesssim r \overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} r \|(f, \underline{f})\|_{\mathbf{S}} + r^{-1} \left[\left| \int_{\mathbf{S}} \underline{f} e^\Phi \right| + \left| \int_{\mathbf{S}} f e^\Phi \right| \right] + r \sqrt{\overset{\circ}{\epsilon}} \|F\|_2 \end{aligned}$$

i.e.,

$$\left| \int_{\mathbf{S}} e_\theta^{\mathbf{S}}(a) e^\Phi \right| \lesssim r \overset{\circ}{\delta} + r^{-1} I(f, \underline{f}) + r \sqrt{\overset{\circ}{\epsilon}} \|F\|_2. \quad (9.5.25)$$

As a consequence we deduce,

$$I(a, f, \underline{f}) \lesssim I(f, \underline{f}) + r^2 \overset{\circ}{\delta}. \quad (9.5.26)$$

Step 5. We combine estimate (9.5.24) with (9.5.26) with

$$r^{\mathbf{S}} \|\not{d}^{\mathbf{S},*} a\|_{k+1} + \|(f, \underline{f})\|_{k+2} \lesssim \overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|F\|_{k+2} + r^{-2} I(f, \underline{f}).$$

Also, taking into account that

$$\|F\|_{k+2}^2 = \|a\|_0 + r^{\mathbf{S}} \|\not{d}^{\mathbf{S},*} a\|_{k+1} + \|(f, \underline{f})\|_{k+2}$$

and the smallness of $\overset{\circ}{\epsilon}$, we deduce,

$$r^{\mathbf{S}} \|\not{d}^{\mathbf{S},*} a\|_{k+1} + \|(f, \underline{f})\|_{k+2} \lesssim \overset{\circ}{\delta} + \sqrt{\overset{\circ}{\epsilon}} \|a\|_0 + r^{-2} I(f, \underline{f}). \quad (9.5.27)$$

Step 6. It remains to get an estimate for $\|a\|_0 = \|a\|_{L^2(\mathbf{S})}$. For this we need to make use of the last equation in for the average of a in (9.4.2) as well as the GCM condition $\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}$,

$$\bar{a}^{\mathbf{S}} = \overline{\left(1 - \frac{r^{\mathbf{S}}}{r}\right)^{\mathbf{S}}} - \frac{r^{\mathbf{S}}}{2} \overline{\left(\check{\kappa} + \bar{\kappa} - \frac{2}{r}\right)^{\mathbf{S}}} + \overline{\text{Err}_6}^{\mathbf{S}}$$

where,

$$\begin{aligned} \text{Err}_6 &= -\frac{r^{\mathbf{S}}}{2} \left[e^a \text{Err}(\kappa, \kappa') - (e^a - 1) \left(\check{\kappa} + \bar{\kappa} - \frac{2}{r} + \not{d}^{\mathbf{S}} f \right) - (e^a - 1 - a) \frac{2}{r} \right] \\ &\quad - a \left(\frac{r^{\mathbf{S}}}{r} - 1 \right). \end{aligned}$$

Hence,

$$|\bar{a}^{\mathbf{S}}| \lesssim \left| \left(1 - \frac{r^{\mathbf{S}}}{r} \right)^{\mathbf{S}} \right| + r \left(|\check{\kappa}| + \left| \bar{\kappa} - \frac{2}{r} \right| \right) + r |\text{Err}_6|.$$

Making use of the assumption²² 9.4.16, (9.4.5) and proceeding as before, it is easy to derive the estimate,

$$|\bar{a}^{\mathbf{S}}| \lesssim r^{-1} \mathring{\delta}. \tag{9.5.28}$$

Step 7. By the standard Poincare inequality on \mathbf{S} we have,

$$\|a - \bar{a}^{\mathbf{S}}\|_{\mathbf{S}} \lesssim r^{\mathbf{S}} \|\not{d}^{\mathbf{S},*} a\|_{\mathbf{S}}.$$

Combining this with (9.5.27) and (9.5.28) we deduce,

$$\|(a, f, \underline{f})\|_{k+2} \lesssim \mathring{\delta} + r^{-2} I(f, \underline{f}). \tag{9.5.29}$$

Finally, making use of the condition

$$\int_{\mathbf{S}} f e^{\Phi} = \Lambda, \quad \int_{\mathbf{S}} \underline{f} e^{\Phi} = \underline{\Lambda}, \quad \Lambda, \underline{\Lambda} = O(\mathring{\delta} r^2)$$

we derive the desired estimate,

$$\|(a, f, \underline{f})\|_{k+2} \lesssim \mathring{\delta}. \tag{9.5.30}$$

The uniqueness part is obvious and left to the reader. This ends the proof of Theorem 9.4.4.

²²Note that full strength of the assumption is only used here, otherwise the weaker assumption, with $\mathring{\delta}$ replaced by $\mathring{\epsilon}$, suffices in the rest of the argument.

9.6 Proof of Proposition 9.4.9

9.6.1 Pull-back of the main equations

According to Proposition 9.4.8 we may assume valid the uniform bounds for the quintets $P^{(n)}$. To establish a contraction estimate we need to compare the quantities,

$$\begin{aligned} h^{(n)} &:= (\Psi^{(n-1)})\# f^{(n)}, & \underline{h}^{(n)} &:= (\Psi^{(n-1)})\# \underline{f}^{(n)}, & w^{(n)} &:= (\Psi^{(n-1)})\# z^{(n)}, \\ e^{(n)} &:= (\Psi^{(n-1)})\# a^{(n)}, \end{aligned}$$

and,

$$\begin{aligned} h^{(n+1)} &:= (\Psi^{(n)})\# f^{(n+1)}, & \underline{h}^{(n+1)} &:= (\Psi^{(n)})\# \underline{f}^{(n+1)}, & w^{(n+1)} &:= (\Psi^{(n)})\# z^{(n+1)}, \\ e^{(n+1)} &:= (\Psi^{(n)})\# a^{(n+1)}. \end{aligned}$$

According to Lemma 9.2.2 we have,

$$\begin{aligned} (\Psi^{(n)})\# (\not{d}^{\mathbf{S}^{(n)}} f^{(n+1)}) &= \not{d}^{(n)} h^{(n+1)}, \\ (\Psi^{(n)})\# (A^{\mathbf{S}^{(n)}} f^{(n+1)}) &= A^{(n)} h^{(n+1)}, \end{aligned}$$

where $\not{d}^{(n)}$, $\not{d}^{*(n)}$, $A^{(n)}$ are the corresponding Hodge operators on \mathring{S} defined with respect to the metric $\not{g}^{(n)} := (\Psi^{(n)})\# (\not{g}^{\mathbf{S}^{(n)}})$ given by,

$$\not{g}^{(n)} = \gamma^{(n)} d\theta^2 + e^{2\Phi^{(n)}} d\varphi^2.$$

Consequently the system (9.4.25) takes the form,

$$\begin{aligned} A^{(n)} w^{(n+1)} + V^{(n)} w^{(n+1)} &= (\Psi^{(n)})\# E^{(n+1)}, \\ A^{(n)} h^{(n+1)} + V^{(n)} h^{(n+1)} &= \frac{3}{4} \kappa^{(n)} (A^{(n)})^{-1} (\rho^{(n)} w^{(n+1)}) + (\Psi^{(n)})\# E^{(n+1)}, \\ A^{(n)} \underline{h}^{(n+1)} + V^{(n)} \underline{h}^{(n+1)} &= -\frac{3}{4} \kappa^{(n)} (A^{(n)})^{-1} (\rho^{(n)} w^{(n+1)}) + (\Psi^{(n)})\# \underline{E}^{(n+1)}, \\ \not{d}^{*(n)} e^{(n+1)} &= \frac{3}{4} (A^{(n)})^{-1} (\rho^{(n)} w^{(n+1)}) - h^{(n+1)} \underline{\omega}^{(n)} - \frac{1}{4} h^{(n+1)} \underline{\kappa}^{(n)} + \frac{1}{4} \underline{h}^{(n+1)} \kappa^{(n)} \\ &\quad + (\Psi^{(n)})\# \tilde{E}^{(n+1)}, \end{aligned} \tag{9.6.1}$$

where, $\kappa^{(n)}$, $\underline{\kappa}^{(n)}$, $\rho^{(n)}$, $\omega^{(n)}$, $\underline{\omega}^{(n)}$ are the pull backs by $\Psi^{(n)}$ of κ , $\underline{\kappa}$, ρ , ω , $\underline{\omega}$. Also $V^{(n)}$ is the pull back by $\Psi^{(n)}$ of the potential $V = \frac{1}{2} \kappa \underline{\kappa} - \rho$.

Equation (9.4.28) takes the form

$$\begin{aligned}
 \frac{2}{r\mathbf{S}^{(n)}} \int_{\mathring{S}, \mathring{g}^{(n)}} \mathring{d}^{(n)} e^{(n+1)} e^{\Phi^{\#n}} &= -\frac{1}{4} \int_{\mathring{S}, \mathring{g}^{(n)}} (\kappa^{(n)})^2 \underline{h}^{(n+1)} e^{\Phi^{\#n}} \\
 &+ \int_{\mathring{S}, \mathring{g}^{(n)}} \left(\frac{1}{4} \kappa \underline{\kappa} + \kappa \underline{\omega} + 3\rho \right)^{(n)} h^{(n+1)} e^{\Phi^{\#n}} \\
 &- \int_{\mathring{S}, \mathring{g}^{(n)}} \mathring{d}^{(n)} (\kappa^{(n)}) e^{\Phi^{\#n}} da_{\mathring{g}^{(n)}} + (\Psi^{(n)})^{\#} \text{Err}_6^{(n+1)}, \\
 \int_{\mathring{S}, \mathring{g}^{(n)}} h^{(n+1)} e^{\Phi^{\#n}} &= \underline{\Lambda}, \quad \int_{\mathring{S}, \mathring{g}^{(n)}} \underline{h}^{(n+1)} e^{\Phi^{\#n}} = \underline{\Lambda},
 \end{aligned} \tag{9.6.2}$$

Equation (9.4.29) takes the form,

$$\frac{e^{(n+1)}}{\mathring{S}, \mathring{g}^{(n)}} = \frac{e^{(n+1)}}{\left(1 - \frac{r\mathbf{S}^{(n)}}{r}\right)^{(n)} \mathring{S}, \mathring{g}^{(n)}} - \frac{r\mathbf{S}^{(n)}}{2} \frac{e^{(n+1)}}{\left(\check{\kappa} + \bar{\kappa} - \frac{2}{r}\right)^{(n)} \mathring{S}, \mathring{g}^{(n)}} + \frac{e^{(n+1)}}{\text{Err}_8^{(n+1)} \mathring{S}, \mathring{g}^{(n)}}. \tag{9.6.3}$$

Finally the system (9.4.31) takes the form,

$$\begin{aligned}
 \varsigma^{\#n} \partial_{\theta} U^{(1+n)} &= (\gamma^{(n)})^{1/2} h^{(1+n)} \left(1 + \frac{1}{4} h^{(1+n)} \underline{h}^{(1+n)} \right), \\
 \partial_{\theta} S^{(1+n)} - \frac{1}{2} \varsigma^{\#n} \underline{\Omega}^{\#n} \partial_{\theta} U^{(1+n)} &= \frac{1}{2} (\gamma^{(n)})^{1/2} \underline{h}^{(1+n)}, \\
 \gamma^{(n)} &= \gamma^{\#n} + (\varsigma^{\#n})^2 \left(\underline{\Omega}^{\#n} + \frac{1}{4} (\underline{b}^{\#n})^2 \gamma^{\#n} \right) (\partial_{\theta} U^{(n)})^2 \\
 &\quad - 2\varsigma^{\#n} \partial_{\theta} U^{(n)} \partial_{\theta} S^{(n)} - \gamma^{\#n} \varsigma^{\#n} \underline{b}^{\#n} \partial_{\theta} U^{(n)}, \\
 U^{(1+n)}(0) &= S^{(1+n)}(0) = 0.
 \end{aligned} \tag{9.6.4}$$

We recall, see (9.4.32), the definition of the norm for the quintets $P^{(n)}$ in the particular case $k = 3$

$$\begin{aligned}
 \|P^{(n)}\|_3 &:= \|\partial_{\theta} (U^{(n)}, S^{(n)})\|_{L^{\infty}(\mathring{S})} + r^{-1} \|\partial_{\theta} (U^{(n)}, S^{(n)})\|_{\mathfrak{h}_1(\mathring{S})} \\
 &\quad + \left\| \left((a^{(n)})^{\#n-1}, (f^{(n)})^{\#n-1}, (\underline{f}^{(n)})^{\#n-1} \right) \right\|_{\mathfrak{h}_1(\mathring{S})}.
 \end{aligned}$$

To prove the estimate

$$\|P^{(n+1)} - P^{(n)}\|_3 \lesssim \overset{\circ}{\delta} \|P^{(n)} - P^{(n-1)}\|_3$$

we set,

$$\begin{aligned}
 \delta w^{(n+1)} &= w^{(n+1)} - w^{(n)}, & \delta h^{(n+1)} &= h^{(n+1)} - h^{(n)}, & \delta \underline{h}^{(n+1)} &= \underline{h}^{(n+1)} - \underline{h}^{(n)}, \\
 \delta e^{(n+1)} &= e^{(n+1)} - e^{(n)}, & \delta U^{(n+1)} &= U^{(n+1)} - U^{(n)}, & \delta S^{(n+1)} &= S^{(n+1)} - S^{(n)},
 \end{aligned}$$

and,

$$\begin{aligned}\delta w^{(n)} &= w^{(n)} - w^{(n-1)}, & \delta h^{(n)} &= h^{(n)} - h^{(n-1)}, & \delta \underline{h}^{(n)} &= \underline{h}^{(n)} - \underline{h}^{(n-1)}, \\ \delta e^{(n)} &= e^{(n)} - e^{(n-1)}, & \delta U^{(n)} &= U^{(n)} - U^{(n-1)}, & \delta S^{(n)} &= S^{(n)} - S^{(n-1)}.\end{aligned}$$

We will derive below the following estimates

$$\|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}, \delta e^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \lesssim \mathring{\delta} r^{-1} \|\partial_\theta(\delta U^{(n)}, \delta S^{(n)})\|_{\mathfrak{h}_1(\mathring{S})} \quad (9.6.5)$$

and

$$r^{-1} \|\partial_\theta(\delta U^{(n+1)}, \delta S^{(n+1)})\|_{\mathfrak{h}_1(\mathring{S})} \lesssim \|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \mathring{\epsilon} r^{-1} \|\partial_\theta(\delta U^{(n)}, \delta S^{(n)})\|_{\mathfrak{h}_1(\mathring{S})}. \quad (9.6.6)$$

Proposition 9.4.9 is then an immediate consequence of (9.6.5) (9.6.6). Thus, from now on, we focus on the proof of (9.6.5) (9.6.6). To this end, we will rely on the following lemmas.

9.6.2 Basic Lemmas

Lemma 9.6.1. *Let F be a reduced scalar function defined in a neighborhood of \mathring{S} in \mathcal{R} and define its pull back $F^{(n)} = (\Psi^{(n)})^\# F$ to \mathring{S} , i.e.,*

$$\begin{aligned}F^{(n)}(\theta) &= F(\mathring{u} + U^{(n)}(\theta), \mathring{s} + S^{(n)}(\theta), \theta), \\ F^{(n-1)}(\theta) &= F(\mathring{u} + U^{(n-1)}(\theta), \mathring{s} + S^{(n-1)}(\theta), \theta).\end{aligned}$$

Then²³, for all $1 \leq p \leq \infty$, with $\delta_n U = U^{(n+1)} - U^{(n)}$, $\delta_n S = S^{(n+1)} - S^{(n)}$

$$\|\delta_n F\|_{L^p(\mathring{S})} \lesssim \left(\|\delta_n U\|_{L^p(\mathring{S})} + \|\delta_n S\|_{L^p(\mathring{S})} \right) \sup_{\mathcal{R}} (|\partial_s F| + |\partial_u F|). \quad (9.6.7)$$

Also,

$$\|\delta_n F\|_{\mathfrak{h}_1(\mathring{S})} \lesssim \left(\|\delta_n U\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta_n S\|_{\mathfrak{h}_1(\mathring{S})} \right) \sup_{\mathcal{R}} |\mathfrak{d}^{\leq 1} \mathfrak{d} F| \quad (9.6.8)$$

where $\delta_n U = U^{(n+1)} - U^{(n)}$, $\delta_n S = S^{(n+1)} - S^{(n)}$.

²³Recall $\partial_s = e_4$, $\partial_u = \frac{\zeta}{2} (e_3 - \underline{\Omega} e_4 - \underline{b} \gamma^{1/2} e_\theta)$.

Proof. We write,

$$\begin{aligned}\delta_n F &:= F(u_0 + U^{(n)}(\theta), s_0 + S^{(n)}(\theta), \theta) - F(u_0 + U^{(n-1)}(\theta), s_0 + S^{(n-1)}(\theta), \theta) \\ &= \int_0^1 \frac{d}{dt} F(u_0 + tU^{(n)}(\theta) + (1-t)U^{(n-1)}(\theta), s_0 + tS^{(n)}(\theta) + (1-t)S^{(n-1)}(\theta), \theta),\end{aligned}$$

i.e., denoting $\delta_n U = U^{(n)} - U^{(n-1)}$, $\delta_n S = S^{(n)} - S^{(n-1)}$,

$$\begin{aligned}|\delta_n F| &\lesssim |\delta_n U| \int_0^1 |\partial_u F(u_0 + tU^{(n)}(\theta) + (1-t)U^{(n-1)}(\theta), s_0 + tS^{(n)}(\theta) + (1-t)S^{(n-1)}(\theta), \theta)| \\ &\quad + |\delta_n S| \int_0^1 |\partial_s F(u_0 + tU^{(n)}(\theta) + (1-t)U^{(n-1)}(\theta), s_0 + tS^{(n)}(\theta) + (1-t)S^{(n-1)}(\theta), \theta)|\end{aligned}$$

i.e.,

$$|\delta_n F| \lesssim |U^{(n)}(\theta) - U^{(n-1)}(\theta)| \sup_{\overset{\circ}{S} + \delta \overset{\circ}{S}} |\partial_u F| + |S^{(n)}(\theta) - S^{(n-1)}(\theta)| \sup_{\overset{\circ}{S} + \epsilon \overset{\circ}{S}} |\partial_s F|$$

from which (9.6.7) easily follows.

Also, in view of $\partial_s = e_4$, $\partial_u = \frac{1}{2}(e_3 - \underline{\Omega}e_4 - \underline{b}\gamma^{1/2}e_\theta)$ and our assumptions for $\underline{\Omega}$ and \underline{b} ,

$$|\delta_n F| \lesssim (|U^{(n)} - U^{(n-1)}| + |S^{(n)} - S^{(n-1)}|) \sup_{\mathcal{R}} \mathfrak{d}F|$$

and by integration,

$$\|\delta_n F\|_{L^p(\overset{\circ}{S})} \lesssim \left(\|\delta_n U\|_{L^p(\overset{\circ}{S})} + \|\delta_n S\|_{L^p(\overset{\circ}{S})} \right) \sup_{\mathcal{R}} |\mathfrak{d}F|.$$

Similarly,

$$\|\mathfrak{d}\delta_n F\|_{L^2(\overset{\circ}{S})} \lesssim \left(\|\delta_n U\|_{\mathfrak{h}^1(\overset{\circ}{S})} + \|\delta_n S\|_{\mathfrak{h}^1(\overset{\circ}{S})} \right) \sup_{\mathcal{R}} |\mathfrak{d}^{\leq 1} \mathfrak{d}F|.$$

Hence,

$$\|\delta_n F\|_{\mathfrak{h}^1(\overset{\circ}{S})} \lesssim \left(\|\delta_n U\|_{\mathfrak{h}^1(\overset{\circ}{S})} + \|\delta_n S\|_{\mathfrak{h}^1(\overset{\circ}{S})} \right) \sup_{\mathcal{R}} |\mathfrak{d}^{\leq 1} \mathfrak{d}F|$$

as desired. \square

Lemma 9.6.2. *Let $\psi, h \in \mathfrak{s}_1(\overset{\circ}{S})$, and $\delta A^{(n)} = A^{(n)} - A^{(n-1)}$. The following formula holds true.*

$$\left| \int_{(\overset{\circ}{S}, \mathfrak{g}^{(n)})} \psi \delta A^{(n)} h \right| \lesssim r^{-2} \|\partial_\theta (\Psi^{(n)} - \Psi^{(n-1)})\|_{\mathfrak{h}^1(\overset{\circ}{S})} \left(\|\psi\|_{\mathfrak{h}^1(\overset{\circ}{S})} \|h\|_{\mathfrak{h}^2(\overset{\circ}{S})} + \|\psi\|_{\mathfrak{h}^2(\overset{\circ}{S})} \|h\|_{\mathfrak{h}^1(\overset{\circ}{S})} \right).$$

Proof. Recall that the metric $g^{(n)}$ is given by

$$g^{(n)} = \gamma^{(n)} d\theta^2 + e^{2\Phi^{#n}} d\varphi^2$$

so that the operator $A^{(n)} = -\mathfrak{d}^{*(n)} \mathfrak{d}^{(n)}$, applied to \mathfrak{s}_1 tensors h on \mathring{S} is given by

$$A^{(n)}h = -\frac{1}{\sqrt{\gamma^{(n)}}} \partial_\theta \left(\frac{1}{\sqrt{\gamma^{(n)}}} (\partial_\theta h + \partial_\theta(\Phi^{#n})h) \right).$$

This yields

$$\begin{aligned} \delta A^{(n)}h &= -\left(\frac{1}{\sqrt{\gamma^{(n)}}} - \frac{1}{\sqrt{\gamma^{(n-1)}}} \right) \partial_\theta \left(\frac{1}{\sqrt{\gamma^{(n)}}} (\partial_\theta h + \partial_\theta(\Phi^{#n})h) \right) \\ &\quad - \frac{1}{\sqrt{\gamma^{(n-1)}}} \partial_\theta \left(\left(\frac{1}{\sqrt{\gamma^{(n)}}} - \frac{1}{\sqrt{\gamma^{(n-1)}}} \right) (\partial_\theta h + \partial_\theta(\Phi^{#n})h) \right) \\ &\quad - \frac{1}{\sqrt{\gamma^{(n-1)}}} \partial_\theta \left(\frac{1}{\sqrt{\gamma^{(n-1)}}} \partial_\theta(\Phi^{#n} - \Phi^{#n-1})h \right). \end{aligned}$$

Using the previous formula to integrate $\psi \delta A^{(n)}h$ on \mathring{S} with the volume of $g^{(n)}$, and after integration by parts, we infer

$$\begin{aligned} &\int_{(\mathring{S}, g^{(n)})} \psi \delta A^{(n)}h \\ &= \int_{(\mathring{S}, g^{(n)})} \frac{1}{\sqrt{\gamma^{(n)}}} \partial_\theta \left(\left(1 - \frac{\sqrt{\gamma^{(n)}}}{\sqrt{\gamma^{(n-1)}}} \right) \psi \right) \frac{1}{\sqrt{\gamma^{(n)}}} (\partial_\theta h_2 + \partial_\theta(\Phi^{#n})h) \\ &\quad + \int_{(\mathring{S}, g^{(n)})} \frac{1}{\sqrt{\gamma^{(n)}}} \partial_\theta \left(\frac{\sqrt{\gamma^{(n)}}}{\sqrt{\gamma^{(n-1)}}} \psi \right) \left(\frac{1}{\sqrt{\gamma^{(n)}}} - \frac{1}{\sqrt{\gamma^{(n-1)}}} \right) (\partial_\theta h_2 + \partial_\theta(\Phi^{#n})h) \\ &\quad + \int_{(\mathring{S}, g^{(n)})} \frac{1}{\sqrt{\gamma^{(n)}}} \partial_\theta \left(\frac{\sqrt{\gamma^{(n)}}}{\sqrt{\gamma^{(n-1)}}} \psi \right) \frac{1}{\sqrt{\gamma^{(n-1)}}} \partial_\theta(\Phi^{#n} - \Phi^{#n-1})h. \end{aligned}$$

We now make us of the bounds (9.1.15) for $(\underline{\Omega}, \underline{b}, \gamma)$ involved in the definition of $\gamma^{(n-1)}$ and $\gamma^{(n)}$, the uniform bound of $P^{(n)}$ provided by Proposition 9.4.8 and the Sobolev inequality to deduce,

$$\begin{aligned} \left| \int_{(\mathring{S}, g^{(n)})} \psi \delta A^{(n)}h \right| &\lesssim r^{-4} \|\gamma^{(n)} - \gamma^{(n-1)}\|_{\mathfrak{h}_1(\mathring{S})} \left(\|\psi\|_{\mathfrak{h}_1(\mathring{S})} \|h\|_{\mathfrak{h}_2(\mathring{S})} + (\|\psi\|_{\mathfrak{h}_2(\mathring{S})} \|h\|_{\mathfrak{h}_1(\mathring{S})}) \right) \\ &\quad + r^{-3} \|\partial_\theta(\Phi^{#n} - \Phi^{#n-1})h\|_{L^2(\mathring{S})} \|\psi\|_{\mathfrak{h}_1(\mathring{S})}. \end{aligned} \quad (9.6.9)$$

To estimate the term $\gamma^{(n)} - \gamma^{(n-1)}$ we recall that,

$$\begin{aligned} \gamma^{(n)} &= \gamma^{\#n} + (\zeta^{\#n})^2 \left(\underline{\Omega}^{\#n} + \frac{1}{4}(\underline{b}^{\#n})^2 \gamma^{\#n} \right) (\partial_\theta U^{(n)})^2 \\ &\quad - 2\zeta^{\#n} \partial_\theta U^{(n)} \partial_\theta S^{(n)} - \gamma^{\#n} \zeta^{\#n} \underline{b}^{\#n} \partial_\theta U^{(n)} \\ \gamma^{(n-1)} &= \gamma^{\#n-1} + (\zeta^{\#n-1})^2 \left(\underline{\Omega}^{\#n-1} + \frac{1}{4}(\underline{b}^{\#n-1})^2 \gamma^{\#n-1} \right) (\partial_\theta U^{(n-1)})^2 \\ &\quad - 2\zeta^{\#n-1} \partial_\theta U^{(n-1)} \partial_\theta S^{(n-1)} - \gamma^{\#n-1} \zeta^{\#n-1} \underline{b}^{\#n-1} \partial_\theta U^{(n-1)}. \end{aligned}$$

The principal term $\gamma^{\#n} - \gamma^{\#n-1}$ can be estimated with the help of Lemma 9.6.1, the uniform bound of $P^{(n)}$ provided by Proposition 9.4.8, and the bounds provided by **A3**. All other terms can be estimated in a similar fashion. We derive,

$$\|\gamma^{(n)} - \gamma^{(n-1)}\|_{\mathfrak{h}_1(\mathring{S})} \lesssim r^2 \|\partial_\theta (\Psi^{(n)} - \Psi^{(n-1)})\|_{\mathfrak{h}_1(\mathring{S})} \quad (9.6.10)$$

where,

$$\|\partial_\theta (\Psi^{(n)} - \Psi^{(n-1)})\|_{\mathfrak{h}_1(\mathring{S})} := \|\partial_\theta (U^{(n)} - U^{(n-1)})\|_{\mathfrak{h}_1(\mathring{S})} + \|\partial_\theta (S^{(n)} - S^{(n-1)})\|_{\mathfrak{h}_1(\mathring{S})}.$$

We deduce,

$$\begin{aligned} \left| \int_{(\mathring{S}, \mathring{g}^{(n)})} \psi \delta A^{(n)} h \right| &\lesssim r^{-2} \|\partial_\theta (\Psi^{(n)} - \Psi^{(n-1)})\|_{\mathfrak{h}_1(\mathring{S})} \left(\|\psi\|_{\mathfrak{h}_1(\mathring{S})} \|h\|_{\mathfrak{h}_2(\mathring{S})} + (\|\psi\|_{\mathfrak{h}_2(\mathring{S})} \|h\|_{\mathfrak{h}_1(\mathring{S})}) \right) \\ &\quad + r^{-3} \|\partial_\theta (\Phi^{\#n} - \Phi^{\#n-1}) h\|_{L^2(\mathring{S})} \|\psi\|_{\mathfrak{h}_1(\mathring{S})}. \end{aligned} \quad (9.6.11)$$

The proof of 9.6.2. is now an immediate consequence of the following.

Lemma 9.6.3. *The following estimate holds true for a reduced scalar $h \in \mathfrak{s}_1(\mathring{S})$*

$$\|\partial_\theta (\Phi^{\#n} - \Phi^{\#n-1}) h\|_{L^2(\mathring{S})} \lesssim r \|\partial_\theta (\Psi^{(n)} - \Psi^{(n-1)})\|_{\mathfrak{h}_1(\mathring{S})} \|h\|_{\mathfrak{h}_2(\mathring{S})}. \quad (9.6.12)$$

Proof. We write,

$$\begin{aligned} &\partial_\theta (\Phi^{\#n} - \Phi^{\#n-1}) \\ &= \left\{ \left(\partial_\theta S^{(n)} - \frac{1}{2} \underline{\Omega} \partial_\theta U^{(n)} \right) e_4 \Phi + \frac{1}{2} \partial_\theta U^{(n)} e_3 \Phi + \sqrt{\gamma} \left(1 - \frac{1}{2} \underline{b} \partial_\theta U^{(n)} \right) e_\theta \Phi \right\}^{\#n} \\ &\quad - \left\{ \left(\partial_\theta S^{(n-1)} - \frac{1}{2} \underline{\Omega} \partial_\theta U^{(n-1)} \right) e_4 \Phi + \frac{1}{2} \partial_\theta U^{(n-1)} e_3 \Phi + \sqrt{\gamma} \left(1 - \frac{1}{2} \underline{b} \partial_\theta U^{(n-1)} \right) e_\theta \Phi \right\}^{\#n-1} \\ &= \left(\partial_\theta S^{(n)} - \frac{1}{2} \underline{\Omega}^{\#n} \partial_\theta U^{(n)} \right) (e_4 \Phi)^{\#n} + \frac{1}{2} \partial_\theta U^{(n)} (e_3 \Phi)^{\#n} + \sqrt{\gamma}^{\#n} \left(1 - \frac{1}{2} \underline{b}^{\#n} \partial_\theta U^{(n)} \right) (e_\theta \Phi)^{\#n} \\ &\quad - \left(\partial_\theta S^{(n-1)} - \frac{1}{2} \underline{\Omega}^{\#n-1} \partial_\theta U^{(n-1)} \right) (e_4 \Phi)^{\#n-1} - \frac{1}{2} \partial_\theta U^{(n-1)} (e_3 \Phi)^{\#n-1} \\ &\quad - \sqrt{\gamma}^{\#n-1} \left(1 - \frac{1}{2} \underline{b}^{\#n-1} \partial_\theta U^{(n-1)} \right) (e_\theta \Phi)^{\#n-1} \end{aligned}$$

i.e., grouping the terms appropriately,

$$\begin{aligned} \partial_\theta (\Phi^{\#n} - \Phi^{\#n-1}) &= J_1 + J_2 + J_3, \\ J_1 &= \left(\partial_\theta S^{(n)} - \frac{1}{2} \underline{\Omega}^{\#n} \partial_\theta U^{(n)} \right) (e_4 \Phi)^{\#n} - \left(\partial_\theta S^{(n-1)} - \frac{1}{2} \underline{\Omega}^{\#n-1} \partial_\theta U^{(n-1)} \right) (e_4 \Phi)^{\#n-1}, \\ J_2 &= \frac{1}{2} \partial_\theta U^{(n)} (e_3 \Phi)^{\#n} - \frac{1}{2} \partial_\theta U^{(n-1)} (e_3 \Phi)^{\#n-1}, \\ J_3 &= \sqrt{\gamma}^{\#n} \left(1 - \frac{1}{2} \underline{b}^{\#n} \partial_\theta U^{(n)} \right) (e_\theta \Phi)^{\#n} - \sqrt{\gamma}^{\#n-1} \left(1 - \frac{1}{2} \underline{b}^{\#n-1} \partial_\theta U^{(n-1)} \right) (e_\theta \Phi)^{\#n-1}, \end{aligned}$$

and,

$$\begin{aligned} J_3 &= J_{31} + J_{32}, \\ J_{31} &= (e_\theta \Phi)^{\#n-1} \left(\sqrt{\gamma}^{\#n} \left(1 - \frac{1}{2} \underline{b}^{\#n} \partial_\theta U^{(n)} \right) - \sqrt{\gamma}^{\#n-1} \left(1 - \frac{1}{2} \underline{b}^{\#n-1} \partial_\theta U^{(n-1)} \right) \right), \\ J_{32} &= \sqrt{\gamma}^{\#n} \left(1 - \frac{1}{2} \underline{b}^{\#n} \partial_\theta U^{(n)} \right) ((e_\theta \Phi)^{\#n} - (e_\theta \Phi)^{\#n-1}). \end{aligned}$$

The contribution to the estimate of of Lemma 9.6.3 given by J_1, J_2, J_{31} can be easily estimated by making use of the uniform bound of $P^{(n)}$ provided by Proposition 9.4.8, the bound (9.1.15) for $(\underline{\Omega}, \underline{b}, \gamma)$, Lemma 9.2.6 as well as Lemma 9.6.1. We thus derive,

$$\|(J_1, J_2, J_{31})h\|_{L^2(\mathring{S})} \lesssim r \|\partial_\theta \Psi^{(n)} - \partial_\theta \Psi^{(n-1)}\|_{\mathfrak{h}_1(\mathring{S})} \|h\|_{\mathfrak{h}_2(\mathring{S})}.$$

It remains to estimate the term $\|J_{32}h\|_{L^2(\mathring{S})}$ which presents a difficulty at the axis of symmetry where $\sin \theta = 0$. We can write,

$$\begin{aligned} J_3 &= J_{31} + J_{32}, \\ J_{31} &= (e_\theta \Phi)^{\#n-1} \left(\sqrt{\gamma}^{\#n} \left(1 - \frac{1}{2} \underline{b}^{\#n} \partial_\theta U^{(n)} \right) - \sqrt{\gamma}^{\#n-1} \left(1 - \frac{1}{2} \underline{b}^{\#n-1} \partial_\theta U^{(n-1)} \right) \right), \\ J_{32} &= \sqrt{\gamma}^{\#n} \left(1 - \frac{1}{2} \underline{b}^{\#n} \partial_\theta U^{(n)} \right) ((e_\theta \Phi)^{\#n} - (e_\theta \Phi)^{\#n-1}). \end{aligned}$$

Clearly, $\|J_{32}h\|_{L^2(\mathring{S})} \lesssim \|((e_\theta \Phi)^{\#n} - (e_\theta \Phi)^{\#n-1})h\|_{L^2(\mathring{S})}$. We are thus left to estimate the term $\|((e_\theta \Phi)^{\#n} - (e_\theta \Phi)^{\#n-1})h\|_{L^2(\mathring{S})}$. Proceeding as in the proof of Lemma 9.6.1 we write, for $F = e_\theta \Phi$,

$$\begin{aligned} |\delta_n F| &\lesssim |\delta_n U| \int_0^1 |\partial_u F(u_0 + tU^{(n)}(\theta) + (1-t)U^{(n-1)}(\theta), s_0 + tS^{(n)}(\theta) + (1-t)S^{(n-1)}(\theta), \theta)| \\ &\quad + |\delta_n S| \int_0^1 |\partial_s F(u_0 + tU^{(n)}(\theta) + (1-t)U^{(n-1)}(\theta), s_0 + tS^{(n)}(\theta) + (1-t)S^{(n-1)}(\theta), \theta)|. \end{aligned}$$

We need to pay special attention on the axis²⁴, where $\sin \theta = 0$, to the integral term involving

$$\partial_u(e_\theta \Phi) = \frac{1}{2} (e_3 - \underline{\Omega}e_4 - \underline{b}\gamma^{1/2}e_\theta) e_\theta \Phi.$$

This leads us to consider the integral,

$$\int_0^1 [\underline{b}e_\theta(e_\theta(\Phi))] (\overset{\circ}{u} + tU^{(n)}(\theta) + (1-t)U^{(n-1)}(\theta), \overset{\circ}{s}, \theta) dt$$

and the L^2 norm of its product with h on $\overset{\circ}{S}$. We recall (see Lemma 2.1.13) that $\Delta \Phi = -K$. and Therefore, $|e_\theta(e_\theta \Phi)| \lesssim r^{-2} + |e_\theta \Phi|^2$ The contribution due to K does not present any difficulties on the axis therefore we are led to consider the integral

$$I(\theta) := \int_0^1 [b(e_\theta(\Phi))^2] (\overset{\circ}{u} + tU^{(n)}(\theta) + (1-t)U^{(n-1)}(\theta), \overset{\circ}{s}, \theta) dt$$

and the L^2 norm of its product with h on $\overset{\circ}{S}$. Making use of (9.1.17) and then the first estimate of (9.1.18) of Lemma 9.1.3 together with our assumption **A3** we derive the bound,

$$\begin{aligned} |I(\theta)h(\theta)| &\lesssim \frac{1}{\sin^2 \theta} \left(\int_0^1 \left| \underline{b}(\overset{\circ}{u} + tU^{(n)}(\theta) + (1-t)U^{(n-1)}(\theta), \overset{\circ}{s}, \theta) \right| dt \right) |h(\theta)| \\ &\lesssim \left| \frac{h(\theta)}{\sin \theta} \right| \sup_{\mathcal{R}} \left| \frac{\underline{b}}{\sin \theta} \right| \lesssim r^2 \left| \frac{h(\theta)}{e^\Phi} \right| \sup_{\mathcal{R}} \left| \frac{\underline{b}}{e^\Phi} \right| \lesssim \overset{\circ}{\epsilon} r \left| \frac{h(\theta)}{e^\Phi} \right|. \end{aligned}$$

Making use of the second estimate in (9.1.17) we then derive,

$$\|I \cdot h\|_{L^2(\overset{\circ}{S})} \lesssim \overset{\circ}{\epsilon} r \left\| \frac{h}{e^\Phi} \right\|_{L^2(S)} \lesssim \overset{\circ}{\epsilon} \|h\|_{\mathfrak{b}_1(S)}.$$

This shows that the behavior along the axis in (9.6.12) is not an issue. This ends the proof of both Lemma 9.6.3 and Lemma 9.6.2. □

□

Lemma 9.6.4. Consider equations on $\overset{\circ}{S}$, of the form,

$$\begin{aligned} A^{(n)}h^{(n+1)} + V^{(n)}h^{(n+1)} &= H^{(n)}, \\ A^{(n-1)}h^{(n)} + V^{(n-1)}h^{(n)} &= H^{(n-1)}. \end{aligned} \tag{9.6.13}$$

²⁴Indeed the term $e_\theta(e_\theta \Phi)$ is quite singular on the axis.

Also,

$$\begin{aligned} \int_{\mathring{S}, \mathring{g}^{(n)}} h^{(n+1)} e^{\Phi^{\#n}} &= B^{(n)}, \\ \int_{\mathring{S}, \mathring{g}^{(n-1)}} h^{(n)} e^{\Phi^{\#n-1}} &= B^{(n-1)}. \end{aligned}$$

Then we have,

$$\begin{aligned} \|h^{(n+1)} - h^{(n)}\|_{\mathfrak{h}_1(\mathring{S})} &\lesssim \overset{\circ}{\delta} r^{-1} \|\partial_\theta (\Psi^{(n)} - \Psi^{(n-1)})\|_{\mathfrak{h}_1(\mathring{S})} \\ &\quad + r \|H^{(n+1)} - H^{(n)}\|_{L^2(\mathring{S})} + r^{-2} |B^{(n)} - B^{(n-1)}|. \end{aligned} \quad (9.6.14)$$

Proof. Setting

$$\begin{aligned} \delta h^{(n+1)} &:= h^{(n+1)} - h^{(n)}, \quad \delta A^{(n)} := A^{(n)} - A^{(n-1)}, \quad \delta V^{(n)} := V^{(n)} - V^{(n-1)}, \\ \delta H^{(n+1)} &:= H^{(n+1)} - H^{(n)}, \quad \delta B^{(n)} := B^{(n)} - B^{(n-1)}, \end{aligned}$$

and subtracting the equations (9.6.13) we derive,

$$(A^{(n)} + V^{(n)})\delta h^{(n+1)} = -(\delta A^{(n)})h^{(n)} - \delta V^{(n)}h^{(n)} + \delta H^{(n+1)}. \quad (9.6.15)$$

Also writing $\int_{\mathring{S}, \mathring{g}^{(n)}} f = \int_{\mathring{S}} f da_{\mathring{g}^{(n)}}$

$$\int_{\mathring{S}, \mathring{g}^{(n)}} \delta h^{(n+1)} e^{\Phi^{\#n}} = \delta B_n - \int_{\mathring{S}} h^{(n)} \left(e^{\Phi^{\#n}} da_{\mathring{g}^{(n)}} - e^{\Phi^{\#n-1}} da_{\mathring{g}^{(n-1)}} \right). \quad (9.6.16)$$

Throughout the proof below we make systematic use of the boundedness result of Proposition 9.4.8 and comparison Lemma 9.2.3, according to which the spaces $\mathfrak{h}_k(\mathring{S}, \mathring{g}^{(n)})$ are all uniformly equivalent to $\mathfrak{h}_k(\mathring{S})$.

Step 1. Estimates for $\|\mathring{\phi}_2^{(n)}(\delta h^{(n+1)})\|_{\mathfrak{h}_1(\mathring{S})}$.

As in Remark 9.3.7, see also the proof of Proposition 9.4.4 in the previous section, we write,

$$A^{(n)} = \mathring{\phi}_1^{(n)} \mathring{\phi}_1^{(n)} = \mathring{\phi}_2^{(n)} \mathring{\phi}_2^{(n)} - (\Psi^{(n)})^\# \left(-3\rho + \frac{1}{2}\vartheta \underline{\vartheta} \right).$$

Hence,

$$\begin{aligned} \int_{(\mathring{S}, \mathring{g}^{(n)})} \delta h^{(n+1)} (A^{(n)} + V^{(n)}) \delta h^{(n+1)} &= \int_{(\mathring{S}, \mathring{g}^{(n)})} |\mathring{\phi}_2^{(n)}(\delta h^{(n+1)})|^2 \\ &\quad + \left(\frac{6m}{r^3} + O(r^{-3}\overset{\circ}{\delta}) \right) \int_{(\mathring{S}, \mathring{g}^{(n)})} |\delta h^{(n+1)}|^2. \end{aligned}$$

Since $\overset{\circ}{\delta}$ is small,

$$\begin{aligned} \int_{(\overset{\circ}{S}, \overset{\circ}{g}^{(n)})} |\phi_2^{(n)}(\delta h^{(n+1)})|^2 &\lesssim \int_{(\overset{\circ}{S}, \overset{\circ}{g}^{(n)})} \delta h^{(n+1)}(A^{(n)} + V^{(n)})\delta h^{(n+1)} \\ &\lesssim \left| \int_{(\overset{\circ}{S}, \overset{\circ}{g}^{(n)})} \delta h^{(n+1)}(\delta A^{(n)}h^{(n)}) \right| + \int_{(\overset{\circ}{S}, \overset{\circ}{g}^{(n)})} \left| \delta h^{(n+1)}\delta V^{(n)}h^{(n)} \right| + \int_{(\overset{\circ}{S}, \overset{\circ}{g}^{(n)})} \left| \delta h^{(n+1)}\delta H^{(n+1)} \right|. \end{aligned}$$

We deduce, making use of the comparison Lemma and the boundedness result of Proposition 9.4.8,

$$\begin{aligned} \|\phi_2^{(n)}\delta h^{(n+1)}\|_{\mathfrak{h}_1(\overset{\circ}{S})}^2 &\lesssim \left| \int_{(\overset{\circ}{S}, \overset{\circ}{g}^{(n)})} \delta h^{(n+1)}(\delta A^{(n)}h^{(n)}) \right| \\ &\quad + \left(\overset{\circ}{\delta}r^{-1}\|\delta V^{(n)}\|_{L^2(\overset{\circ}{S})} + \|\delta H^{(n+1)}\|_{L^2(\overset{\circ}{S})} \right) \|\delta h^{(n+1)}\|_{L^2(\overset{\circ}{S})}. \end{aligned} \quad (9.6.17)$$

Step 1a. We estimate the term $\left| \int_{(\overset{\circ}{S}, \overset{\circ}{g}^{(n)})} \delta h^{(n+1)}(\delta A^{(n)}h^{(n)}) \right|$ using Lemma 9.6.2 with $\psi = \delta h^{(n+1)}$ and $h = h^{(n)}$. We deduce, making use of the boundedness

$$\begin{aligned} \left| \int_{(\overset{\circ}{S}, \overset{\circ}{g}^{(n)})} \delta h^{(n+1)}(\delta A^{(n)}h^{(n)}) \right| &\lesssim r^{-2}\|\partial_\theta(\Psi^{(n)} - \Psi^{(n-1)})\|_{\mathfrak{h}_1(\overset{\circ}{S})}\|\delta h^{(n+1)}\|_{\mathfrak{h}_2(\overset{\circ}{S})}\|h^{(n)}\|_{\mathfrak{h}_2(\overset{\circ}{S})} \\ &\lesssim r^{-2}\overset{\circ}{\delta}\|\delta h^{(n+1)}\|_{\mathfrak{h}_2(\overset{\circ}{S})}\|\partial_\theta(\Psi^{(n)} - \Psi^{(n-1)})\|_{\mathfrak{h}_1(\overset{\circ}{S})}. \end{aligned}$$

Back to (9.6.17) we deduce,

$$\begin{aligned} \|\phi_2^{(n)}\delta h^{(n+1)}\|_{\mathfrak{h}_1(\overset{\circ}{S})}^2 &\lesssim \left(\overset{\circ}{\delta}r^{-1}\|\delta V^{(n)}\|_{L^2(\overset{\circ}{S})} + r^{-2}\overset{\circ}{\delta}\|\partial_\theta(\delta\Psi^{(n)})\|_{\mathfrak{h}_1(\overset{\circ}{S})} + \|\delta H^{(n+1)}\|_{L^2(\overset{\circ}{S})} \right) \|\delta h^{(n+1)}\|_{L^2(\overset{\circ}{S})}. \end{aligned} \quad (9.6.18)$$

Step 1b. We estimate $\|\delta V^{(n)}\|_{L^2(\overset{\circ}{S})}$. Recall that

$$\delta V^{(n)} = \left(\frac{1}{2}\kappa_{\underline{K}} - \rho \right)^{\#_n} - \left(\frac{1}{2}\kappa_{\underline{K}} - \rho \right)^{\#_{n-1}}.$$

In view of Lemma 9.6.1 and our assumptions **A1-A3** we derive,

$$\|\delta V^{(n)}\|_{L^2(\overset{\circ}{S})} \lesssim r^{-2}\|(\delta U^{(n)}, \delta S^{(n)})\|_{L^2(\overset{\circ}{S})}.$$

Therefore, back to (9.6.18), we deduce

$$\| \mathcal{A}_2^{(n)} \delta h^{(n+1)} \|_{\mathring{b}_1(\mathring{S})}^2 \lesssim \left(r^{-2} \mathring{\delta} \| \partial_\theta (\delta \Psi^{(n)}) \|_{\mathring{b}_1(\mathring{S})} + \| \delta H^{(n+1)} \|_{L^2(\mathring{S})} \right) \| \delta h^{(n+1)} \|_{L^2(\mathring{S})}. \quad (9.6.19)$$

Step 2. We make use of the equation (9.6.16)

$$\int_{\mathring{S}, \mathring{g}^{(n)}} \delta h^{(n+1)} e^{\Phi^{\#n}} = \delta B_n - \int_{\mathring{S}} h^{(n)} \left(e^{\Phi^{\#n}} da_{\mathring{g}^{(n)}} - e^{\Phi^{\#n-1}} da_{\mathring{g}^{(n-1)}} \right)$$

from which we deduce,

$$\begin{aligned} \left| \int_{\mathring{S}, \mathring{g}^{(n)}} \delta h^{(n+1)} e^{\Phi^{\#n}} \right| &\lesssim \left| \int_{\mathring{S}} h^{(n)} \left(e^{\Phi^{\#n}} da_{\mathring{g}^{(n)}} - e^{\Phi^{\#n-1}} da_{\mathring{g}^{(n-1)}} \right) \right| + |\delta B_n| \\ &\lesssim \| h^{(n)} \|_{L^2(\mathring{S})} \left\| \left(e^{\Phi^{\#n}} \frac{\sqrt{\gamma^{(n)}}}{\sqrt{\gamma^{(0)}}} - e^{\Phi^{\#n-1}} \frac{\sqrt{\gamma^{(n-1)}}}{\sqrt{\gamma^{(0)}}} \right) \right\|_{L^2(\mathring{S})} + |\delta B_n|. \end{aligned}$$

On the other hand,

$$\left(e^{\Phi^{\#n}} \frac{\sqrt{\gamma^{(n)}}}{\sqrt{\gamma^{(0)}}} - e^{\Phi^{\#n-1}} \frac{\sqrt{\gamma^{(n-1)}}}{\sqrt{\gamma^{(0)}}} \right) = \left(e^{\Phi^{\#n}} - e^{\Phi^{\#n-1}} \right) \frac{\sqrt{\gamma^{(n)}}}{\sqrt{\gamma^{(0)}}} + e^{\Phi^{\#n-1}} \left(\frac{\sqrt{\gamma^{(n)}}}{\sqrt{\gamma^{(0)}}} - \frac{\sqrt{\gamma^{(n-1)}}}{\sqrt{\gamma^{(0)}}} \right).$$

Hence, proceeding as before,

$$\begin{aligned} \left\| \left(e^{\Phi^{\#n}} \frac{\sqrt{\gamma^{(n)}}}{\sqrt{\gamma^{(0)}}} - e^{\Phi^{\#n-1}} \frac{\sqrt{\gamma^{(n-1)}}}{\sqrt{\gamma^{(0)}}} \right) \right\|_{L^2(\mathring{S})} &\lesssim \| e^{\Phi^{\#n}} - e^{\Phi^{\#n-1}} \|_{L^2(\mathring{S})} + r^{-1} \| \gamma^{\#n} - \gamma^{\#n-1} \|_{L^2(\mathring{S})} \\ &\lesssim r \| \partial_\theta (\Psi^{(n)} - \Psi^{(n-1)}) \|_{\mathring{b}_1(\mathring{S})}. \end{aligned}$$

Therefore,

$$\left| \int_{\mathring{S}, \mathring{g}^{(n)}} \delta h^{(n+1)} e^{\Phi^{\#n}} \right| \lesssim \mathring{\delta} r \| \partial_\theta (\Psi^{(n)} - \Psi^{(n-1)}) \|_{\mathring{b}_1(\mathring{S})} + |\delta B_n|. \quad (9.6.20)$$

Step 3. Making use of Lemma 9.1.4 together with (9.6.19), (9.6.20) and the comparison Lemma and we deduce,

$$\begin{aligned} \| \delta h^{(n+1)} \|_{L^2(\mathring{S}, \mathring{g}^{(n)})} &\lesssim r \| \mathcal{A}_2^{(n)} \delta h^{(n+1)} \|_{\mathring{b}_1(\mathring{S}, \mathring{g}^{(n)})} + r^{-2} \left| \int_{\mathring{S}, \mathring{g}^{(n)}} \delta h^{(n+1)} e^{\Phi^{\#n}} \right| \\ &\lesssim r \| \mathcal{A}_2^{(n)} \delta h^{(n+1)} \|_{\mathring{b}_1(\mathring{S})} + \mathring{\delta} r^{-1} \| \partial_\theta (\Psi^{(n)} - \Psi^{(n-1)}) \|_{\mathring{b}_1(\mathring{S})} + r^{-2} |\delta B_n| \\ &\lesssim r \left(r^{-2} \mathring{\delta} \| \partial_\theta (\delta \Psi^{(n)}) \|_{\mathring{b}_1(\mathring{S})} + \| \delta H^{(n+1)} \|_{L^2(\mathring{S})} \right)^{1/2} \| \delta h^{(n+1)} \|_{L^2(\mathring{S})}^{1/2} \\ &\quad + \mathring{\delta} r^{-1} \| \partial_\theta (\Psi^{(n)} - \Psi^{(n-1)}) \|_{\mathring{b}_1(\mathring{S})} + r^{-2} |\delta B_n|. \end{aligned}$$

We infer that,

$$\|\delta h^{(n+1)}\|_{L^2(\mathring{S}, \mathring{\mathcal{G}}^{(n)})} \lesssim \overset{\circ}{\delta} r^{-1} \|\partial_\theta (\Psi^{(n)} - \Psi^{(n-1)})\|_{\mathfrak{h}_1(\mathring{S})} + r \|\delta H^{(n+1)}\|_{L^2(\mathring{S})} + r^{-2} |\delta B_n|$$

which, together with (9.6.19), ends the proof of Lemma 9.6.4. \square

9.6.3 Proof of the estimates (9.6.5), (9.6.6)

We are now in position to prove (9.6.5) (9.6.6).

Step 1. We start by estimating $\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}$. To this end, we need to apply Lemma 9.6.4 to the equations for $\delta w^{(n+1)}, \delta h^{(n+1)}, \delta \underline{h}^{(n+1)}$, derived from the first three equations in (9.6.1), and estimate the corresponding $\delta H^{(n+1)}$ on the right-hand side. This is tedious but straightforward and one derives, in the case of estimates of $\delta w^{(n+1)}$

$$\begin{aligned} \|\delta H^{(n+1)}\|_{L^2(\mathring{S})} &\lesssim \overset{\circ}{\epsilon} \left\{ \|U^{(n)} - U^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} + \|S^{(n)} - S^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} \right. \\ &\quad \left. + \|\delta h^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta e^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \right\} \end{aligned}$$

and in the case of $\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}$

$$\begin{aligned} \|\delta H^{(n+1)}\|_{L^2(\mathring{S})} &\lesssim \|\delta w^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \overset{\circ}{\epsilon} \left\{ \|U^{(n)} - U^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} + \|S^{(n)} - S^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} \right. \\ &\quad \left. + \|\delta h^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta e^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \right\}. \end{aligned}$$

Remark 9.6.5. Note that the presence of the inverse operators $(A^{(n)})^{-1}$ in the right-hand side of the equations for $\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}$ do not create any difficulties when taking differences. Indeed we can write,

$$(A^{(n)})^{-1} - (A^{(n-1)})^{-1} = (A^{(n)})^{-1} (A^{(n-1)} - A^{(n)}) (A^{(n-1)})^{-1}$$

and estimate the difference $\delta A^{(n)} = A^{(n)} - A^{(n-1)}$ as in the proof of Lemma 9.6.4.

We infer from Lemma 9.6.4 and the above estimates

$$\begin{aligned} \|\delta w^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} &\lesssim \overset{\circ}{\epsilon} \left\{ \|U^{(n)} - U^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} + \|S^{(n)} - S^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} \right. \\ &\quad \left. + \|\delta h^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta e^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \right\} \end{aligned}$$

and

$$\begin{aligned} \|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} &\lesssim \|\delta w^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \overset{\circ}{\epsilon} \left\{ \|U^{(n)} - U^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} + \|S^{(n)} - S^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} \right. \\ &\quad \left. + \|\delta h^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta e^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \right\}. \end{aligned}$$

This thus yields

$$\begin{aligned} \|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} &\lesssim \overset{\circ}{\epsilon} \|U^{(n)} - U^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} + \overset{\circ}{\epsilon} \|S^{(n)} - S^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} \\ &\quad + \overset{\circ}{\epsilon} \|\delta e^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})}. \end{aligned} \quad (9.6.21)$$

Step 2. Next, we estimate $\not{d}^* \delta e^{(n+1)}$. We make use of the last equation in (9.6.1) which we write into the form

$$\not{d}^{*(n)} e^{(n+1)} = H^{(n+1)}$$

with,

$$\begin{aligned} H^{(n+1)} &= \frac{3}{4} (A^{(n)})^{-1} (\rho^{(n)} w^{(n+1)}) - h^{(n+1)} \underline{\omega}^{(n)} + \underline{h}^{(n+1)} \omega^{(n)} - \frac{1}{4} h^{(n+1)} \underline{\kappa}^{(n)} + \frac{1}{4} \underline{f}^{(n+1)} \kappa^{(n)} \\ &\quad + (A^{(n)})^{-1} (-\not{d}^{*(n)} \check{\mu}^{(n)} + (\Psi^{(n)})^\# \text{Err}_3^{(n+1)}). \end{aligned}$$

Hence,

$$\not{d}^{*(n)} (\delta e^{(n+1)}) + (\not{d}^{*(n)} - \not{d}^{*(n-1)}) e^{(n)} = \delta H^{(n+1)}$$

which can be written in the form,

$$\begin{aligned} \frac{\sqrt{\gamma^{(0)}}}{\sqrt{\gamma^{(n)}}} \not{d}^* (\delta e^{(n+1)}) &= -(\not{d}^{*(n)} - \not{d}^{*(n-1)}) e^{(n)} + \delta H^{(n+1)} \\ &= -\left(\frac{\sqrt{\gamma^{(0)}}}{\sqrt{\gamma^{(n)}}} - \frac{\sqrt{\gamma^{(0)}}}{\sqrt{\gamma^{(n-1)}}} \right) \not{d}^* e^{(n)} + \delta H^{(n+1)} \end{aligned}$$

or,

$$\not{d}^* (\delta e^{(n+1)}) = \left(1 - \frac{\sqrt{\gamma^{(n)}}}{\sqrt{\gamma^{(n-1)}}} \right) \not{d}^* e^{(n)} + \frac{\sqrt{\gamma^{(n)}}}{\sqrt{\gamma^{(0)}}} \delta H^{(n+1)}.$$

This yields

$$\|\not{d}^* (\delta e^{(n+1)})\|_{L^2(\mathring{S})} \lesssim \overset{\circ}{\epsilon} \|\gamma^{(n)} - \gamma^{(n-1)}\|_{L^2(\mathring{S})} + \|\delta H^{(n+1)}\|_{L^2(\mathring{S})}$$

where we used the uniform bound of $P^{(n)}$ provided by Proposition 9.4.8 and the bound (9.1.15) for $(\underline{\Omega}, \underline{b}, \gamma)$. $\gamma^{(n)} - \gamma^{(n-1)}$ can be estimated as in the proof of Lemma 9.6.4, and hence

$$\|\phi^*(\delta e^{(n+1)})\|_{L^2(\mathring{S})} \lesssim \mathring{\epsilon} \left(\|U^{(n)} - U^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} + \|S^{(n)} - S^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} \right) + \|\delta H^{(n+1)}\|_{L^2(\mathring{S})}.$$

The differences $H^{(n+1)} - H^{(n)}$ can also be easily estimated and deduce,

$$\begin{aligned} \|H^{(n)} - H^{(n-1)}\|_{L^2(\mathring{S})} &\lesssim \|\delta h^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \\ &+ \mathring{\epsilon} \left(\|U^{(n)} - U^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} + \|S^{(n)} - S^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} \right). \end{aligned}$$

We obtain

$$\begin{aligned} \|\phi^*(\delta e^{(n+1)})\|_{L^2(\mathring{S})} &\lesssim \|\delta h^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \\ &+ \mathring{\epsilon} \left(\|U^{(n)} - U^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} + \|S^{(n)} - S^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} \right). \end{aligned} \quad (9.6.22)$$

Step 3. Next, we estimate the average of $\delta e^{(n+1)}$. Recall from (9.6.3)

$$\overline{e^{(n+1)}}^{\mathring{S}, \mathring{g}^{(n)}} = \overline{\left(1 - \frac{r^{\mathbf{S}(n)}}{r}\right)^{(n)}_{\mathring{S}, \mathring{g}^{(n)}}} - \frac{r^{\mathbf{S}(n)}}{2} \overline{\left(\check{\kappa} + \bar{\kappa} - \frac{2}{r}\right)^{(n)}_{\mathring{S}, \mathring{g}^{(n)}}} + \overline{\text{Err}_8^{(n+1)}}^{\mathring{S}, \mathring{g}^{(n)}}.$$

and

$$\overline{e^{(n)}}^{\mathring{S}, \mathring{g}^{(n)}} = \overline{\left(1 - \frac{r^{\mathbf{S}(n-1)}}{r}\right)^{(n-1)}_{\mathring{S}, \mathring{g}^{(n-1)}}} - \frac{r^{\mathbf{S}(n-1)}}{2} \overline{\left(\check{\kappa} + \bar{\kappa} - \frac{2}{r}\right)^{(n)}_{\mathring{S}, \mathring{g}^{(n-1)}}} + \overline{\text{Err}_8^{(n)}}^{\mathring{S}, \mathring{g}^{(n-1)}}.$$

Taking the difference, recalling that we have in the (θ, φ) coordinates system

$$d\text{vol} \mathring{g}^{(n)} = \sqrt{\gamma^{(n)}} e^{\Phi^{\#n}} d\theta d\varphi, \quad r^{\mathbf{S}(n)} = \int_{\mathring{S}} \sqrt{\gamma^{(n)}} e^{\Phi^{\#n}} d\theta d\varphi$$

and

$$d\text{vol} \mathring{g}^{(n-1)} = \sqrt{\gamma^{(n-1)}} e^{\Phi^{\#n-1}} d\theta d\varphi, \quad r^{\mathbf{S}(n-1)} = \int_{\mathring{S}} \sqrt{\gamma^{(n-1)}} e^{\Phi^{\#n-1}} d\theta d\varphi$$

and using the uniform bound of $P^{(n)}$ provided by Proposition 9.4.8 and the bounds **A1** for $\check{\Gamma}$, we infer

$$\left| \overline{\delta e^{(n+1)}}^{\mathring{S}} \right| \lesssim \mathring{\epsilon} \left\{ \|\gamma^{(n)} - \gamma^{(n-1)}\|_{L^2(\mathring{S})} + \|\Phi^{\#n} - \Phi^{\#n-1}\|_{L^2(\mathring{S})} + \|\delta \text{Err}_6^{(n+1)}\|_{L^2(\mathring{S})} \right\}.$$

Arguing as above, we deduce

$$\left| \overline{\delta e^{(n+1)}}^{\circ \mathring{S}} \right| \lesssim \left\{ \begin{aligned} & \overset{\circ}{\epsilon} \left\{ \|\delta h^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \|\delta e^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \right. \\ & \left. + \|U^{(n)} - U^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} + \|S^{(n)} - S^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} \right\}. \end{aligned} \right.$$

Together with (9.6.21) (9.6.22), and an elliptic estimate for \not{d}^* we infer

$$\|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}, \delta e^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \lesssim \overset{\circ}{\epsilon} \|U^{(n)} - U^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} + \overset{\circ}{\epsilon} \|S^{(n)} - S^{(n-1)}\|_{\mathfrak{h}_2(\mathring{S})} \quad (9.6.23)$$

which concludes the proof of (9.6.5).

Step 4. Finally, we focus on (9.6.6). Recall (9.6.4)

$$\begin{aligned} \varsigma^{\#n} \partial_{\theta} U^{(1+n)} &= (\gamma^{(n)})^{1/2} h^{(1+n)} \left(1 + \frac{1}{4} h^{(1+n)} \underline{h}^{(1+n)} \right), \\ \partial_{\theta} S^{(1+n)} - \frac{1}{2} \varsigma^{\#n} \underline{\Omega}^{\#n} \partial_{\theta} U^{(1+n)} &= \frac{1}{2} (\gamma^{(n)})^{1/2} \underline{h}^{(1+n)}, \\ \gamma^{(n)} &= \gamma^{\#n} + (\varsigma^{\#n})^2 \left(\underline{\Omega}^{\#n} + \frac{1}{4} (\underline{b}^{\#n})^2 \gamma^{\#n} \right) (\partial_{\theta} U^{(n)})^2 \\ &\quad - 2 \varsigma^{\#n} \partial_{\theta} U^{(n)} \partial_{\theta} S^{(n)} - \gamma^{\#n} \varsigma^{\#n} \underline{b}^{\#n} \partial_{\theta} U^{(n)}, \\ U^{(1+n)}(0) &= S^{(1+n)}(0) = 0. \end{aligned} \quad (9.6.24)$$

Taking the difference and arguing as above, we derive

$$\begin{aligned} \|\partial_{\theta} \delta U^{(1+n)}\|_{\mathfrak{h}_1(\mathring{S})} &\lesssim \|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \overset{\circ}{\epsilon} \|\gamma^{(n)} - \gamma^{(n-1)}\|_{\mathfrak{h}_1(\mathring{S})}, \\ \|\partial_{\theta} \delta S^{(1+n)}\|_{\mathfrak{h}_1(\mathring{S})} &\lesssim \|\delta U^{(1+n)}\|_{\mathfrak{h}_2(\mathring{S})} + \overset{\circ}{\epsilon} \|\delta S^{(1+n)}\|_{\mathfrak{h}_2(\mathring{S})} + \|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \\ &\quad + \overset{\circ}{\epsilon} \|\gamma^{(n)} - \gamma^{(n-1)}\|_{\mathfrak{h}_1(\mathring{S})}. \end{aligned}$$

Since $\delta U^{(1+n)} = \delta S^{(1+n)} = 0$, we deduce

$$\begin{aligned} \|\delta U^{(1+n)}\|_{\mathfrak{h}_2(\mathring{S})} &\lesssim \|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \overset{\circ}{\epsilon} \|\gamma^{(n)} - \gamma^{(n-1)}\|_{\mathfrak{h}_1(\mathring{S})}, \\ \|\delta S^{(1+n)}\|_{\mathfrak{h}_2(\mathring{S})} &\lesssim \|\delta U^{(1+n)}\|_{\mathfrak{h}_2(\mathring{S})} + \overset{\circ}{\epsilon} \|\delta S^{(1+n)}\|_{\mathfrak{h}_2(\mathring{S})} + \|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} \\ &\quad + \overset{\circ}{\epsilon} \|\gamma^{(n)} - \gamma^{(n-1)}\|_{\mathfrak{h}_1(\mathring{S})} \end{aligned}$$

and hence

$$\|\delta U^{(n+1)}, \delta S^{(n+1)}\|_{\mathfrak{h}_2(\mathring{S})} \lesssim \|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \overset{\circ}{\epsilon} \|\delta U^{(n)}, \delta S^{(n)}\|_{\mathfrak{h}_2(\mathring{S})} + \overset{\circ}{\epsilon} \|\gamma^{(n)} - \gamma^{(n-1)}\|_{\mathfrak{h}_1(\mathring{S})}.$$

Estimating $\gamma^{(n)} - \gamma^{(n-1)}$ as above, we infer

$$\|\delta U^{(n+1)}, \delta S^{(n+1)}\|_{\mathfrak{h}_2(\mathring{S})} \lesssim \|\delta h^{(n+1)}, \delta \underline{h}^{(n+1)}\|_{\mathfrak{h}_1(\mathring{S})} + \mathring{\epsilon} \|\delta U^{(n)}, \delta S^{(n)}\|_{\mathfrak{h}_2(\mathring{S})} + \mathring{\epsilon} \|\delta U^{(n)}, \delta S^{(n)}\|_{\mathfrak{h}_2(\mathring{S})}.$$

This is the desired estimate (9.6.6) and hence concludes the proof of Proposition 9.4.9.

9.7 A Corollary to Theorem 9.4.2

The following result is a simple Corollary of Theorem 9.4.2.

Theorem 9.7.1 (Existence of GCM spheres). *In addition to the assumptions of Theorem 9.4.2 we assume that, for any background sphere S in \mathcal{R} ,*

$$r \left| \int_S \beta e^\Phi \right| + \left| \int_S e_\theta(\underline{\kappa}) e^\Phi \right| \lesssim \mathring{\delta}. \tag{9.7.1}$$

Then there exists a unique GCM sphere \mathbf{S} , which is a deformation of \mathring{S} , such that the GCMS conditions hold true

$$\begin{aligned} \mathring{\phi}_2^{\mathbf{S},*} \mathring{\phi}_1^{\mathbf{S},*} \underline{\kappa}^{\mathbf{S}} &= \mathring{\phi}_2^{\mathbf{S},*} \mathring{\phi}_1^{\mathbf{S},*} \mu^{\mathbf{S}} = 0, & \kappa^{\mathbf{S}} &= \frac{2}{r\mathbf{S}}, \\ \int_{\mathbf{S}} \beta^{\mathbf{S}} e^\Phi &= 0, & \int_{\mathbf{S}} e_\theta^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}}) e^\Phi &= 0. \end{aligned} \tag{9.7.2}$$

Moreover all other estimates of Theorem 9.4.2 hold true.

Proof. The proof of the theorem follows easily in view of Theorem 9.4.2 and the following lemma.

Lemma 9.7.2. *Let \mathbf{S} be a deformation of \mathring{S} as in Theorem 9.4.2 with $\Lambda = \int_{\mathbf{S}} f e^\Phi, \underline{\Lambda} = \int_{\mathbf{S}} \underline{f} e^\Phi$. The following identities hold true.*

$$\begin{aligned} \Lambda &= \frac{r^3}{3m} \left(\int_{\mathring{S}} \beta e^\Phi - \int_{\mathbf{S}} \beta^{\mathbf{S}} e^\Phi \right) + F_1(\Lambda, \underline{\Lambda}), \\ \underline{\Lambda} &= \frac{r^2}{2(1 + \frac{m}{r})} \left(\int_{\mathring{S}} (e_\theta \underline{\kappa}) e^\Phi - \int_{\mathbf{S}} (e_\theta^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}}) e^\Phi + \Upsilon \int_{\mathring{S}} e_\theta(\kappa) e^\Phi \right) \\ &\quad + \frac{\Upsilon}{2(1 + \frac{m}{r})} \Lambda + F_2(\Lambda, \underline{\Lambda}), \end{aligned} \tag{9.7.3}$$

where F_1, F_2 are continuous²⁵ in $\Lambda, \underline{\Lambda}$, with $F_1(0, 0) = F_2(0, 0) = 0$, verifying the estimates,

$$|F_1| + |F_2| \lesssim \mathring{\epsilon} \delta r^2.$$

²⁵In fact smooth.

Proof. To prove (9.7.3) we start with the change of frame formula,

$$\begin{aligned}\beta^{\mathbf{S}} &= e^a \left(\beta + \frac{3}{2} \rho f \right) + e^a \text{Err}(\beta, \beta^{\mathbf{S}}), \\ \text{Err}(\beta, \beta^{\mathbf{S}}) &= \frac{1}{2} \underline{f} \alpha + \text{l.o.t.}\end{aligned}$$

We write²⁶

$$\begin{aligned}\beta^{\mathbf{S}} &= \beta + \frac{3}{2} \rho f + (e^a - 1) \left(\beta + \frac{3}{2} \rho f \right) + e^a \text{Err}(\beta, \beta^{\mathbf{S}}) \\ &= \beta + \frac{3}{2} \left(-\frac{2m}{r^3} \right) f + \frac{3}{2} \left(\rho + \frac{2m}{r^3} \right) f + (e^a - 1) \left(\beta + \frac{3}{2} \rho f \right) + e^a \text{Err}(\beta, \beta^{\mathbf{S}})\end{aligned}$$

and deduce,

$$\beta^{\mathbf{S}} + \frac{3m^{\mathbf{S}}}{(r^{\mathbf{S}})^3} f = \beta + \text{Err}'(\beta, \beta^{\mathbf{S}})$$

with error term $\text{Err}'(\beta, \beta^{\mathbf{S}})$,

$$\text{Err}'(\beta, \beta^{\mathbf{S}}) = \left(\frac{3m^{\mathbf{S}}}{(r^{\mathbf{S}})^3} - \frac{3m}{r^3} f \right) + \frac{3}{2} \left(\rho + \frac{2m}{r^3} \right) f + (e^a - 1) \left(\beta + \frac{3}{2} \rho f \right) + e^a \text{Err}(\beta, \beta^{\mathbf{S}}).$$

Making use of the assumptions **A1-A3**, the estimates of Theorem 9.4.2 for (f, \underline{f}, a) as well as the bounds for $\overset{\circ}{r} - r^{\mathbf{S}}$, $\overset{\circ}{m} - m^{\mathbf{S}}$ we deduce

$$\left| \text{Err}'(\beta, \beta^{\mathbf{S}}) \right| \lesssim r^{-1} \overset{\circ}{\delta} \epsilon.$$

Thus,

$$\begin{aligned}\frac{3m^{\mathbf{S}}}{(r^{\mathbf{S}})^3} \int_{\mathbf{S}} f e^{\Phi} &= \int_{\mathbf{S}} \beta e^{\Phi} - \int_{\mathbf{S}} \beta^{\mathbf{S}} e^{\Phi} + \int_{\mathbf{S}} \text{Err}'(\beta, \beta^{\mathbf{S}}) e^{\Phi} \\ &= \int_{\overset{\circ}{S}} \beta e^{\Phi} - \int_{\mathbf{S}} \beta^{\mathbf{S}} e^{\Phi} + \int_{\mathbf{S}} \text{Err}'(\beta, \beta^{\mathbf{S}}) e^{\Phi} + \left(\int_{\mathbf{S}} \beta e^{\Phi} - \int_{\overset{\circ}{S}} \beta e^{\Phi} \right)\end{aligned}$$

or,

$$\begin{aligned}\frac{3m}{r^3} \int_{\mathbf{S}} f e^{\Phi} &= \int_{\mathbf{S}} \beta e^{\Phi} - \int_{\mathbf{S}} \beta^{\mathbf{S}} e^{\Phi} + \int_{\mathbf{S}} \text{Err}'(\beta, \beta^{\mathbf{S}}) e^{\Phi} + \left(\int_{\mathbf{S}} \beta e^{\Phi} - \int_{\mathbf{S}} \beta e^{\Phi} \right) \\ &\quad + \left(\frac{3m}{r^3} - \frac{3m^{\mathbf{S}}}{(r^{\mathbf{S}})^3} \right) \int_{\mathbf{S}} f e^{\Phi}.\end{aligned}$$

²⁶Here (r, m) represents the area radius and Hawking of $\overset{\circ}{S}$ while $(r^{\mathbf{S}}, m^{\mathbf{S}})$ represent the area radius and Hawking of \mathbf{S} . Since $|\frac{r^{\mathbf{S}}}{r} - 1| \lesssim \overset{\circ}{\delta}$, $|m^{\mathbf{S}} - m| \lesssim \overset{\circ}{\delta}$ we can interchange freely $r^{\mathbf{S}}$ with r and $m^{\mathbf{S}}$ with m .

Clearly,

$$\left| \int_{\mathbf{S}} \text{Err}'(\beta, \beta^{\mathbf{S}}) e^{\Phi} \right| \lesssim r^2 \delta \overset{\circ}{\epsilon}.$$

Also, proceeding exactly as in Corollary 9.2.5 we deduce,

$$\left| \int_{\mathbf{S}} \beta e^{\Phi} - \int_{\overset{\circ}{\mathbf{S}}} \beta e^{\Phi} \right| \lesssim \delta \left(\sup_{\mathcal{R}} \overset{\circ}{r} |\mathfrak{D}_{\mathcal{R}}^{\leq 1}(\beta e^{\Phi})| + \sup_{\mathcal{R}} \overset{\circ}{r}^2 |e_3(\beta e^{\Phi})| \right).$$

Thus, in view of the assumptions **A1-A3**,

$$\left| \int_{\mathbf{S}} \beta e^{\Phi} - \int_{\overset{\circ}{\mathbf{S}}} \beta e^{\Phi} \right| \lesssim \delta \overset{\circ}{\epsilon} r^2. \quad (9.7.4)$$

We deduce,

$$\Lambda = \frac{r^3}{3m} \left(\int_{\overset{\circ}{\mathbf{S}}} \beta e^{\Phi} - \int_{\mathbf{S}} \beta^{\mathbf{S}} e^{\Phi} \right) + F_1(\Lambda, \underline{\Lambda})$$

where the error term $F_1(\Lambda, \underline{\Lambda})$ is a continuous function of $\Lambda, \underline{\Lambda}$ verifying the estimate,

$$\left| F_1(\Lambda, \underline{\Lambda}) \right| \lesssim \overset{\circ}{\epsilon} \delta r^2.$$

We also recall, see Lemma 9.3.4

$$\begin{aligned} e_{\theta}^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}}) &= e_{\theta \underline{\kappa}} - \mathfrak{d}_1^{\mathbf{S},*} \mathfrak{d}_1^{\mathbf{S}} \underline{f} - \underline{\kappa} e_{\theta}^{\mathbf{S}} a - \frac{1}{4} \underline{\kappa} (f \underline{\kappa} + \underline{f} \underline{\kappa}) + \underline{\kappa} (f \underline{\omega} - \underline{\omega} f) + \underline{f} \rho \\ &\quad + \text{Err}(e_{\theta}^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}}, e_{\theta \underline{\kappa}}) \end{aligned}$$

where,

$$\begin{aligned} \text{Err}(e_{\theta}^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}}, e_{\theta \underline{\kappa}}) &= (e^{-a} - 1) \left(e_{\theta \underline{\kappa}} - \mathfrak{d}_1^{\mathbf{S},*} \mathfrak{d}_1^{\mathbf{S}} \underline{f} + \frac{1}{2} f e_{3 \underline{\kappa}} + \frac{1}{2} \underline{f} e_{4 \underline{\kappa}} \right) \\ &\quad + e^{-a} \left[e_{\theta}^{\mathbf{S}} \text{Err}(\underline{\kappa}, \underline{\kappa}^{\mathbf{S}}) + e_{\theta}^{\mathbf{S}}(a) \left(\mathfrak{d}_1^{\mathbf{S}} \underline{f} + \text{Err}(\underline{\kappa}, \underline{\kappa}^{\mathbf{S}}) \right) + \frac{1}{2} f \underline{f} e_{\theta \underline{\kappa}} + \frac{1}{8} f^2 \underline{f} e_{3 \underline{\kappa}} \right] \\ &\quad + \frac{1}{2} \underline{f} \left(2 \mathfrak{d}_1 \underline{\eta} - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} + 2(\underline{\xi} \underline{\xi} + \underline{\eta}^2) \right) \\ &\quad + \frac{1}{2} \underline{f} \underline{f} e_{\theta \underline{\kappa}} + \frac{1}{8} f^2 \underline{f} e_{3 \underline{\kappa}} + \frac{1}{2} \underline{f} \left(2 \mathfrak{d}_1 \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 + 2(\underline{\eta} + \underline{\eta} - 2\underline{\zeta}) \underline{\xi} \right). \end{aligned}$$

Making use of the identity $\mathfrak{d}_1^{\mathbf{S},*} \mathfrak{d}_1^{\mathbf{S}} = \mathfrak{d}_2^{\mathbf{S}} \mathfrak{d}_2^{\mathbf{S},*} + 2K$ we deduce,

$$e_{\theta}^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}}) + \left(\frac{1}{4} \underline{\kappa} \underline{\kappa} - \rho + 2K \right) \underline{f} = e_{\theta \underline{\kappa}} - \mathfrak{d}_2^{\mathbf{S}} \mathfrak{d}_2^{\mathbf{S},*} \underline{f} + \underline{\kappa} e_{\theta}^{\mathbf{S}} a - \frac{1}{4} \underline{\kappa}^2 f - \underline{\kappa} \underline{\omega} f + \text{Err}(e_{\theta}^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}}, e_{\theta \underline{\kappa}}).$$

Writing, $k = \frac{2}{r} + (\kappa - \frac{2}{r})$, $\underline{\kappa} = \frac{-2\Upsilon}{r} + (\underline{\kappa} + \frac{2\Upsilon}{r})$, $\rho = -\frac{2m}{r^3} + (\rho + \frac{2m}{r^3})$, $K = \frac{1}{r^2} + (K - \frac{1}{r^2})$

$$\begin{aligned} \frac{1}{4}\underline{\kappa}\underline{\kappa} - \rho + 2K &= \frac{1}{r^2} + \frac{4m}{r^3} + \frac{1}{2r} \left(\underline{\kappa} + \frac{2\Upsilon}{r} \right) - \frac{\Upsilon}{2r} \left(\kappa - \frac{2}{r} \right) + \left(\rho + \frac{2m}{r^3} \right) + 2 \left(K - \frac{1}{r^2} \right) \\ &= \frac{1}{r^2} + \frac{4m}{r^3} + O(\epsilon r^{-2}). \end{aligned}$$

Also,

$$\begin{aligned} \underline{\kappa} &= -\frac{2\Upsilon}{r} + \left(\underline{\kappa} + \frac{2\Upsilon}{r} \right) = -\frac{2\Upsilon}{r} + O(r^{-2}\epsilon), \\ \underline{\kappa}\underline{\omega} &= -\frac{2m\Upsilon}{r^3} + O(r^{-2}\epsilon), \end{aligned}$$

and, since $a, f, \underline{f} = O(r^{-1}\delta)$,

$$\text{Err}(e_{\theta\underline{\kappa}}^{\mathbf{S}\mathbf{S}}, e_{\theta\underline{\kappa}}) = O(r^{-3}\delta\epsilon).$$

We deduce,

$$e_{\theta\underline{\kappa}}^{\mathbf{S}\mathbf{S}} + \left(\frac{1}{r^2} + \frac{4m}{r^3} \right) \underline{f} = e_{\theta\underline{\kappa}} - \mathcal{D}_2 \mathcal{D}_2^* \underline{f} - \frac{2\Upsilon}{r} e_{\theta}^{\mathbf{S}} a + \frac{2m\Upsilon}{r^3} f + \text{Err}_1$$

with error term

$$|\text{Err}_1| \lesssim \epsilon \delta r^{-3}.$$

Projecting on e^{Φ} and proceeding as before,

$$\begin{aligned} \left(\frac{1}{r^2} + \frac{4m}{r^3} \right) \int_{\mathbf{S}} \underline{f} e^{\Phi} &= \int_{\mathring{\mathbf{S}}} (e_{\theta\underline{\kappa}}) e^{\Phi} - \int_{\mathbf{S}} (e_{\theta\underline{\kappa}}^{\mathbf{S}\mathbf{S}}) e^{\Phi} - \frac{2\Upsilon}{r} \int_{\mathbf{S}} (e_{\theta}^{\mathbf{S}} a) e^{\Phi} + \frac{2m\Upsilon}{r^3} \int_{\mathbf{S}} f e^{\Phi} \\ &\quad + I_1(\Lambda, \underline{\Lambda}) \end{aligned} \quad (9.7.5)$$

where the error term I_1 is continuous in $\Lambda, \underline{\Lambda}$ and verifies the estimate

$$|I_1| \lesssim \epsilon \delta.$$

We now calculate $\int_{\mathbf{S}} (e_{\theta}^{\mathbf{S}} a) e^{\Phi}$. In view of formula (9.4.3) we have,

$$\begin{aligned} \frac{2}{r\mathbf{S}} \int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(a) e^{\Phi} &= - \int_{\mathbf{S}} e_{\theta}(\kappa) e^{\Phi} + \frac{1}{4} \int_{\mathbf{S}} \kappa^2 \underline{f} e^{\Phi} - \int_{\mathbf{S}} \left(\frac{1}{4} \underline{\kappa}\underline{\kappa} + \underline{\kappa}\underline{\omega} + 3\rho \right) f e^{\Phi} + \text{Err}_7 \\ &= - \int_{\mathring{\mathbf{S}}} e_{\theta}(\kappa) e^{\Phi} + \frac{1}{r^2} \underline{\Lambda} - \frac{1}{r^2} \left(1 + \frac{m}{r} \right) \Lambda + I_2(\Lambda, \underline{\Lambda}) \end{aligned}$$

where,

$$\left| I_2(\Lambda, \underline{\Lambda}) \right| \lesssim \overset{\circ}{\epsilon} \delta.$$

Indeed, using once more Corollary 9.2.5, we note that

$$\left| \int_{\mathbf{S}} e_{\theta}(\kappa) e^{\Phi} - \int_{\overset{\circ}{\mathbf{S}}} e_{\theta}(\kappa) e^{\Phi} \right| \lesssim \overset{\circ}{\epsilon} \delta.$$

All other error terms are easily estimated.

Back to (9.7.5) we deduce,

$$\begin{aligned} \left(\frac{1}{r^2} + \frac{4m}{r^3} \right) \underline{\Lambda} &= \int_{\overset{\circ}{\mathbf{S}}} (e_{\theta} \underline{\kappa}) e^{\Phi} - \int_{\mathbf{S}} (e_{\theta}^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}}) e^{\Phi} - \frac{2\Upsilon}{r} \int_{\mathbf{S}} (e_{\theta}^{\mathbf{S}} a) e^{\Phi} + \frac{2m\Upsilon}{r^3} \Lambda + I_1(\Lambda, \underline{\Lambda}) \\ &= \int_{\overset{\circ}{\mathbf{S}}} (e_{\theta} \underline{\kappa}) e^{\Phi} - \int_{\mathbf{S}} (e_{\theta}^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}}) e^{\Phi} + \frac{2m\Upsilon}{r^3} \Lambda + I_1(\Lambda, \underline{\Lambda}) \\ &\quad - \Upsilon \left(- \int_{\overset{\circ}{\mathbf{S}}} e_{\theta}(\kappa) e^{\Phi} + \frac{1}{r^2} \underline{\Lambda} - \frac{1}{r^2} \left(1 + \frac{m}{r} \right) \Lambda + I_2(\Lambda, \underline{\Lambda}) \right) \\ &= \int_{\overset{\circ}{\mathbf{S}}} (e_{\theta} \underline{\kappa}) e^{\Phi} - \int_{\mathbf{S}} (e_{\theta}^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}}) e^{\Phi} + \Upsilon \int_{\overset{\circ}{\mathbf{S}}} e_{\theta}(\kappa) e^{\Phi} - \frac{\Upsilon}{r^2} \underline{\Lambda} + \frac{\Upsilon}{r^2} \left(1 + \frac{3m}{r} \right) \Lambda \\ &\quad + I_1(\Lambda, \underline{\Lambda}) - \Upsilon I_2(\Lambda, \underline{\Lambda}). \end{aligned}$$

Hence,

$$\begin{aligned} \frac{2}{r^2} \left(1 + \frac{m}{r} \right) \underline{\Lambda} &= \int_{\overset{\circ}{\mathbf{S}}} (e_{\theta} \underline{\kappa}) e^{\Phi} - \int_{\mathbf{S}} (e_{\theta}^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}}) e^{\Phi} + \Upsilon \int_{\overset{\circ}{\mathbf{S}}} e_{\theta}(\kappa) e^{\Phi} + \frac{\Upsilon}{r^2} \left(1 + \frac{3m}{r} \right) \Lambda \\ &\quad + I_1(\Lambda, \underline{\Lambda}) - \Upsilon I_2(\Lambda, \underline{\Lambda}). \end{aligned}$$

Thus,

$$\underline{\Lambda} = \frac{r^2}{2(1 + \frac{m}{r})} \left(\int_{\overset{\circ}{\mathbf{S}}} (e_{\theta} \underline{\kappa}) e^{\Phi} - \int_{\mathbf{S}} (e_{\theta}^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}}) e^{\Phi} + \Upsilon \int_{\overset{\circ}{\mathbf{S}}} e_{\theta}(\kappa) e^{\Phi} \right) + \frac{\Upsilon}{2(1 + \frac{m}{r})} \Lambda + F_2(\Lambda, \underline{\Lambda})$$

with error term $F_2(\Lambda, \underline{\Lambda})$ continuous in $\Lambda, \underline{\Lambda}$ and verifying the estimate,

$$\left| F_2(\Lambda, \underline{\Lambda}) \right| \lesssim \overset{\circ}{\epsilon} \delta r^2.$$

This ends the proof of the lemma. \square

Under the assumptions of the theorem the system

$$\begin{aligned} \Lambda &= \frac{r^3}{3m} \int_{\overset{\circ}{\mathbf{S}}} \beta e^{\Phi} + F_1(\Lambda, \underline{\Lambda}), \\ \underline{\Lambda} &= \frac{r^2}{2(1 + \frac{m}{r})} \left(\int_{\overset{\circ}{\mathbf{S}}} (e_{\theta}^{\mathbf{S}} \underline{\kappa}) e^{\Phi} + \Upsilon \int_{\overset{\circ}{\mathbf{S}}} e_{\theta}(\kappa) e^{\Phi} \right) + \frac{\Upsilon}{2(1 + \frac{m}{r})} \Lambda + F_2(\Lambda, \underline{\Lambda}), \end{aligned}$$

has a unique solution $\Lambda_0, \underline{\Lambda}_0$ verifying the estimate

$$|\Lambda_0| + |\underline{\Lambda}_0| \lesssim \overset{\circ}{\delta} r^2.$$

Taking $\Lambda = \Lambda_0, \underline{\Lambda} = \underline{\Lambda}_0$ in (9.7.3) we deduce,

$$\int_{\mathbf{S}} \beta^{\mathbf{S}} e^{\Phi} = 0, \quad \int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}}) e^{\Phi} = 0,$$

as stated. □

Corollary 9.7.3 (Rigidity II). *Assume that the spacetime region \mathcal{R} satisfies assumptions **A1** – **A3** and (9.4.5), as well as (9.7.1). Assume given a sphere $\mathbf{S} \subset \mathcal{R}$ endowed with a compatible frame $e_3^{\mathbf{S}}, e_{\theta}^{\mathbf{S}}, e_4^{\mathbf{S}}$ which verifies the GCM conditions*

$$\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}, \quad \not\phi_2^{\mathbf{S}} \not\phi_1^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}} = \not\phi_2^{\mathbf{S}} \not\phi_1^{\mathbf{S}} \mu^{\mathbf{S}} = 0. \quad (9.7.6)$$

In addition, we have

$$r \left| \int_{\mathbf{S}} \beta^{\mathbf{S}} e^{\Phi} \right| + \left| \int_{\mathbf{S}} e_{\theta}^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}}) e^{\Phi} \right| \lesssim \overset{\circ}{\delta}. \quad (9.7.7)$$

Then the transition functions $(f, \underline{f}, \log \lambda)$ from the background frame of \mathcal{R} to that of \mathbf{S} verifies the estimates

$$\|(f, \underline{f}, \log(\lambda))\|_{\mathfrak{h}_{\text{max}-2}(\mathbf{S})} \lesssim \overset{\circ}{\delta}.$$

The proof is an immediate consequence of Lemma 9.7.2 and the rigidity result of Corollary 9.4.5. Note that in the corollary, the sphere \mathbf{S} is reinterpreted as a deformation sphere from the unique background sphere sharing the same south pole.

9.8 Construction of GCM hypersurfaces

We are ready to state our main result concerning the construction of GCM hypersurfaces.

Theorem 9.8.1. *Let a fixed spacetime region \mathcal{R} verifying assumptions **A1** – **A3** and (9.4.5). In addition we assume that,*

$$\left| \int_{S(u,s)} \eta e^{\phi} \right| \lesssim r^2 \overset{\circ}{\delta}, \quad \left| \int_{S(u,s)} \underline{\xi} e^{\phi} \right| \lesssim r^2 \overset{\circ}{\delta}, \quad (9.8.1)$$

and, everywhere on Σ_0 ,

$$\left| \left(\frac{2}{s} + \underline{\Omega} \right) \Big|_{SP} - 1 - \frac{2m}{r} \right| \lesssim \delta \quad (9.8.2)$$

where SP denotes the South pole, i.e. $\theta = 0$ relative to the adapted geodesic coordinates u, s, θ .

Let $\mathbf{S}_0 = \mathbf{S}_0[\overset{\circ}{u}, \overset{\circ}{s}, \Lambda_0, \underline{\Lambda}_0]$ be a fixed GCMS provided by Theorem 9.4.2. Then, there exists then a unique, local²⁷, smooth, \mathbf{Z} -invariant spacelike hypersurface Σ_0 passing through \mathbf{S}_0 , a scalar function $u^{\mathbf{S}}$ defined on Σ_0 , whose level surfaces are topological spheres denoted by \mathbf{S} , and a smooth collection of constants $\Lambda^{\mathbf{S}}, \underline{\Lambda}^{\mathbf{S}}$ verifying,

$$\Lambda^{\mathbf{S}_0} = \Lambda_0, \quad \underline{\Lambda}^{\mathbf{S}_0} = \underline{\Lambda}_0,$$

such that the following conditions are verified:

1. The surfaces \mathbf{S} of constant $u^{\mathbf{S}}$ verifies all the properties stated in Theorem 9.4.2 for the prescribed constants $\Lambda^{\mathbf{S}}, \underline{\Lambda}^{\mathbf{S}}$. In particular they come endowed with null frames $(e_4^{\mathbf{S}}, e_\theta^{\mathbf{S}}, e_3^{\mathbf{S}})$ such that
 - i. For each \mathbf{S} the GCM conditions (9.4.1) with $\Lambda = \Lambda^{\mathbf{S}}, \underline{\Lambda} = \underline{\Lambda}^{\mathbf{S}}$, are verified.
 - ii. The transition functions $(f, \underline{f}, a = \log \lambda)$ verify the estimates (9.4.8).
 - iii. The transversality conditions (9.4.12) are verified.
 - iv. The corresponding Ricci and curvature coefficients verify the estimates (9.4.11) and (9.4.14).
2. Denoting $r^{\mathbf{S}}$ to be the area radius of the spheres \mathbf{S} we have, for some constant c_* ,

$$u^{\mathbf{S}} + r^{\mathbf{S}} = c_*, \quad \text{along } \Sigma_0. \quad (9.8.3)$$

3. Let $\nu^{\mathbf{S}}$ be the unique vectorfield tangent to the hypersurface Σ_0 , normal to \mathbf{S} , and normalized by $g(\nu^{\mathbf{S}}, e_4^{\mathbf{S}}) = -2$. There exists a unique scalar function $a^{\mathbf{S}}$ on Σ_0 such that $\nu^{\mathbf{S}}$ is given by

$$\nu^{\mathbf{S}} = e_3^{\mathbf{S}} + a^{\mathbf{S}} e_4^{\mathbf{S}}.$$

The following normalization condition holds true at the South Pole SP of every sphere \mathbf{S} , i.e. at $\theta = 0$,

$$a^{\mathbf{S}} \Big|_{SP} = -1 - \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}. \quad (9.8.4)$$

²⁷i.e. in a neighborhood of \mathbf{S}_0 .

4. We extend $u^{\mathbf{S}}$ and $r^{\mathbf{S}}$ in a small neighborhood of Σ_0 such that the following transversality conditions are verified²⁸ on Σ_0 ,

$$e_4^{\mathbf{S}}(u^{\mathbf{S}}) = 0, \quad e_4(r^{\mathbf{S}}) = \frac{r^{\mathbf{S}}}{2\kappa^{\mathbf{S}}} = 1. \quad (9.8.5)$$

5. In view of (9.8.5) the Ricci coefficients $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}$ are well defined for every $\mathbf{S} \subset \Sigma_0$ and verify

$$\int_{\mathbf{S}} \eta^{\mathbf{S}} e^{\Phi} = \int_{\mathbf{S}} \underline{\xi}^{\mathbf{S}} e^{\Phi} = 0. \quad (9.8.6)$$

6. The following estimates hold true for all $k \leq s_{max}$,

$$\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} \lesssim \overset{\circ}{\epsilon}, \quad (9.8.7)$$

$$\|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} \lesssim \overset{\circ}{\epsilon}, \quad (9.8.8)$$

$$\left\| a^{\mathbf{S}} + 1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} \right\|_{\mathfrak{h}_k(\mathbf{S})} \lesssim \overset{\circ}{\epsilon}. \quad (9.8.9)$$

The $e_3^{\mathbf{S}}$ derivatives of $\check{\kappa}^{\mathbf{S}}, \vartheta^{\mathbf{S}}, \zeta^{\mathbf{S}}, \check{\kappa}^{\mathbf{S}}, \underline{\vartheta}^{\mathbf{S}}, \alpha^{\mathbf{S}}, \beta^{\mathbf{S}}, \check{\rho}^{\mathbf{S}}, \mu^{\mathbf{S}}, \underline{\beta}^{\mathbf{S}}$ are well defined on Σ_0 and we have, for all $k \leq s_{max} - 1$

$$\begin{aligned} \|e_3^{\mathbf{S}}(\check{\kappa}^{\mathbf{S}}, \vartheta^{\mathbf{S}}, \zeta^{\mathbf{S}}, \check{\kappa}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-1}, \\ \|e_3^{\mathbf{S}}(\underline{\vartheta}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon}, \\ \|e_3^{\mathbf{S}}(\alpha^{\mathbf{S}}, \beta^{\mathbf{S}}, \check{\rho}^{\mathbf{S}}, \mu^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-2}, \\ \|e_3^{\mathbf{S}}(\underline{\beta}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-1}, \\ \|e_3^{\mathbf{S}}(\underline{\alpha}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon}. \end{aligned} \quad (9.8.10)$$

7. The transition functions from the background foliation to that of Σ_0 verify

$$\|\mathfrak{d}^{\leq s_{max}+1}(f, \underline{f}, \log \lambda)\|_{L^2(\mathbf{S})} \lesssim \overset{\circ}{\delta}. \quad (9.8.11)$$

Corollary 9.8.2 (Rigidity III). *Let a fixed spacetime region \mathcal{R} verifying assumptions **A1** – **A3** and the small GCM conditions (9.4.5). Assume given a GCM hypersurface $\Sigma_0 \subset \mathcal{R}$ foliated by surfaces \mathbf{S} such that*

$$\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}, \quad \phi_2^{\mathbf{S}} \phi_1^{\mathbf{S}} \underline{\kappa}^{\mathbf{S}} = \phi_2^{\mathbf{S}} \phi_1^{\mathbf{S}} \mu^{\mathbf{S}} = 0, \quad \int_{\mathbf{S}} \eta^{\mathbf{S}} e^{\Phi} = 0, \quad \int_{\mathbf{S}} \underline{\xi}^{\mathbf{S}} e^{\Phi} = 0.$$

²⁸Here the average of $\kappa^{\mathbf{S}}$ is taken on \mathbf{S} . In view of the GCM conditions (9.8.13) we deduce $e_4^{\mathbf{S}}(r^{\mathbf{S}}) = 1$.

Assume in addition that for a specific sphere \mathbf{S}_0 on Σ_0 , the transition functions f, \underline{f} from the background foliation to \mathbf{S}_0 verify

$$\int_{\mathbf{S}_0} f e^\Phi = O(\overset{\circ}{\delta}), \quad \int_{\mathbf{S}_0} \underline{f} e^\Phi = O(\overset{\circ}{\delta}). \quad (9.8.12)$$

Then, for all derivatives of the transition functions along \mathbf{S} ,

$$\|\mathfrak{d}^{\leq s_{max}+1}(f, \underline{f}, \log \lambda)\|_{L^2(\mathbf{S}_0)} \lesssim \overset{\circ}{\delta}.$$

We give below the proof of Theorem 9.8.1 and a short discussion of the proof of the Corollary.

9.8.1 Definition of Σ_0

As stated in the theorem we assume given a spacetime region $\mathcal{R} = \{|u - \overset{\circ}{u}| \leq \delta_{\mathcal{R}}, |s - \overset{\circ}{s}| \leq \delta_{\mathcal{R}}\}$ (see definition(9.1.6)) endowed with a background foliation such that the condition **A1-A3** hold true. We also assume given a deformation sphere

$$\mathbf{S}_0 := \mathbf{S}[\overset{\circ}{u}, \overset{\circ}{s}, \Lambda_0, \underline{\Lambda}_0]$$

of a given sphere $\overset{\circ}{S} = S(\overset{\circ}{u}, \overset{\circ}{s})$ of the background foliation which verify the conclusions of Theorem 9.4.2. We then proceed to construct, in a small neighborhood of \mathbf{S}_0 , a spacelike hypersurface Σ_0 initiating at \mathbf{S}_0 verifying all the desired properties mentioned above. In what follows we outline the main steps in the construction.

Step 1. According to Theorem 9.4.2, for every value of the parameters (u, s) in \mathcal{R} (i.e. such that the background spheres $S(u, s) \subset \mathcal{R}$) and every real numbers $(\Lambda, \underline{\Lambda})$, there exists a unique GCM sphere $\mathbf{S}[u, s, \Lambda, \underline{\Lambda}]$, as a \mathbf{Z} -polarized deformation of $S(u, s)$. In particular the following are verified:

- \mathbf{S} coincides with $S(u, s)$ at their south poles (i.e. for $\theta = 0$ in the adapted coordinates).
- On \mathbf{S} , the following GCMS conditions hold

$$\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}, \quad \not\phi_2^{\mathbf{S},*} \not\phi_1^{\mathbf{S},*} \underline{\kappa}^{\mathbf{S}} = 0, \quad \not\phi_2^{\mathbf{S},*} \not\phi_1^{\mathbf{S},*} \mu^{\mathbf{S}} = 0, \quad (9.8.13)$$

$$\int_{\mathbf{S}} f e^\Phi = \Lambda^{\mathbf{S}}, \quad \int_{\mathbf{S}} \underline{f} e^\Phi = \underline{\Lambda}^{\mathbf{S}}, \quad (9.8.14)$$

where $(f, \underline{f}, \lambda)$ are the transition parameters of the frame transformation from the background frame (e_3, e_θ, e_4) to the adapted frame $(e_3^{\mathbf{S}}, e_\theta^{\mathbf{S}}, e_4^{\mathbf{S}})$. The constants $\Lambda^{\mathbf{S}}, \underline{\Lambda}^{\mathbf{S}}$ depend smoothly on the surfaces \mathbf{S} and

$$\Lambda^{\mathbf{S}_0} = \Lambda_0, \quad \underline{\Lambda}^{\mathbf{S}_0} = \underline{\Lambda}_0.$$

- There is a map $\Xi : S(u, s) \rightarrow \mathbf{S}$ given by

$$\Xi : (u, s, \theta) = \left(u + U(\theta, u, s, \Lambda, \underline{\Lambda}), s + S(\theta, u, s, \Lambda, \underline{\Lambda}), \theta \right) \quad (9.8.15)$$

with U, S vanishing at $\theta = 0$.

- The transversality conditions (9.4.12) hold, i.e. $\underline{\xi}^{\mathbf{S}} = \omega^{\mathbf{S}} = \zeta^{\mathbf{S}} + \underline{\eta}^{\mathbf{S}} = 0$. Note that these specify the $e_4^{\mathbf{S}}$ derivatives of $(f, \underline{f}, \lambda)$ on \mathbf{S} .
- The Ricci coefficients²⁹ $\kappa^{\mathbf{S}}, \underline{\kappa}^{\mathbf{S}}, \vartheta^{\mathbf{S}}, \underline{\vartheta}^{\mathbf{S}}, \zeta^{\mathbf{S}}$ are well defined on each sphere \mathbf{S} of Σ_0 , and hence on Σ_0 . The same holds true for all curvature coefficients $\alpha^{\mathbf{S}}, \beta^{\mathbf{S}}, \rho^{\mathbf{S}}, \underline{\beta}^{\mathbf{S}}, \underline{\alpha}^{\mathbf{S}}$. Taking into account our transversality condition we remark that the only ill defined Ricci coefficients are $\eta^{\mathbf{S}}, \xi^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}}$.
- Let $\nu^{\mathbf{S}}$ be the unique vectorfield tangent to the hypersurface Σ_0 , normal to \mathbf{S} , and normalized by $g(\nu^{\mathbf{S}}, e_4^{\mathbf{S}}) = -2$. There exists a unique scalar function $a^{\mathbf{S}}$ on Σ_0 such that $\nu^{\mathbf{S}}$ is given by

$$\nu^{\mathbf{S}} = e_3^{\mathbf{S}} + a^{\mathbf{S}} e_4^{\mathbf{S}}.$$

We deduce that the quantities

$$\begin{aligned} g(D_{\nu^{\mathbf{S}}} e_4^{\mathbf{S}}, e_\theta^{\mathbf{S}}) &= 2\eta^{\mathbf{S}} + 2a^{\mathbf{S}} \xi^{\mathbf{S}} = 2\eta^{\mathbf{S}}, \\ g(D_{\nu^{\mathbf{S}}} e_3^{\mathbf{S}}, e_\theta^{\mathbf{S}}) &= 2\underline{\xi}^{\mathbf{S}} + 2a^{\mathbf{S}} \underline{\eta}^{\mathbf{S}} = 2(\underline{\xi}^{\mathbf{S}} - a^{\mathbf{S}} \zeta^{\mathbf{S}}), \\ g(D_{\nu^{\mathbf{S}}} e_3^{\mathbf{S}}, e_4^{\mathbf{S}}) &= 4\underline{\omega}^{\mathbf{S}} - 4a^{\mathbf{S}} \omega^{\mathbf{S}} = 4\underline{\omega}^{\mathbf{S}}, \end{aligned}$$

are well defined on Σ_0 . Thus the scalar $a^{\mathbf{S}}$ allows us to specify the remaining Ricci coefficients, $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}}$ along Σ_0 , which we do below.

²⁹Consequently the Hawking mass $m^{\mathbf{S}}$ is also well defined.

9.8.2 Extrinsic properties of Σ_0

We analyze the extrinsic properties of the hypersurfaces Σ_0 defined in Step 1.

Step 2. We define the scalar function $u^{\mathbf{S}}$ on Σ_0 as

$$u^{\mathbf{S}} := c_0 - r^{\mathbf{S}}, \quad (9.8.16)$$

where $r^{\mathbf{S}}$ is the area radius of \mathbf{S} and the constant c_0 is such that $u^{\mathbf{S}}|_{\mathbf{S}_0} = \overset{\circ}{u}$, i.e. $c_0 = \overset{\circ}{u} + r^{\mathbf{S}}|_{\mathbf{S}_0}$.

Step 3. We extend $u^{\mathbf{S}}$ and $r^{\mathbf{S}}$ in a small neighborhood of Σ_0 such that the following transversality conditions are verified.

$$e_4^{\mathbf{S}}(u^{\mathbf{S}}) = 0, \quad e_4^{\mathbf{S}}(r^{\mathbf{S}}) = \frac{r^{\mathbf{S}}}{2} \overline{\kappa^{\mathbf{S}}}, \quad (9.8.17)$$

where the average of $\kappa^{\mathbf{S}}$ is taken on \mathbf{S} . In view of the GCM conditions (9.8.13) we deduce $e_4^{\mathbf{S}}(r^{\mathbf{S}}) = 1$.

Step 4. Note that $e_3^{\mathbf{S}}(u^{\mathbf{S}}, r^{\mathbf{S}})$ remain undetermined. On the other hand, since $e_{\theta}^{\mathbf{S}}(u^{\mathbf{S}}) = e_{\theta}^{\mathbf{S}}(r^{\mathbf{S}}) = 0$, we deduce in view of (9.8.17)

$$\begin{aligned} e_{\theta}^{\mathbf{S}}(e_3^{\mathbf{S}}(u^{\mathbf{S}})) &= [e_{\theta}^{\mathbf{S}}, e_3^{\mathbf{S}}]u^{\mathbf{S}} = \left[\frac{1}{2}(\underline{\kappa}^{\mathbf{S}} + \underline{\vartheta}^{\mathbf{S}})e_{\theta}^{\mathbf{S}} + (\zeta^{\mathbf{S}} - \eta^{\mathbf{S}})e_3^{\mathbf{S}} + \underline{\xi}^{\mathbf{S}}e_4^{\mathbf{S}} \right] u^{\mathbf{S}} \\ &= (\zeta^{\mathbf{S}} - \eta^{\mathbf{S}})e_3^{\mathbf{S}}(u^{\mathbf{S}}), \\ e_{\theta}^{\mathbf{S}}(e_3^{\mathbf{S}}(r^{\mathbf{S}})) &= [e_{\theta}^{\mathbf{S}}, e_3^{\mathbf{S}}]r^{\mathbf{S}} = \left[\frac{1}{2}(\underline{\kappa}^{\mathbf{S}} + \underline{\vartheta}^{\mathbf{S}})e_{\theta}^{\mathbf{S}} + (\zeta^{\mathbf{S}} - \eta^{\mathbf{S}})e_3^{\mathbf{S}} + \underline{\xi}^{\mathbf{S}}e_4^{\mathbf{S}} \right] r^{\mathbf{S}} \\ &= (\zeta^{\mathbf{S}} - \eta^{\mathbf{S}})e_3^{\mathbf{S}}(r^{\mathbf{S}}) + \underline{\xi}^{\mathbf{S}}. \end{aligned}$$

Thus introducing the scalars

$$\zeta^{\mathbf{S}} := \frac{2}{e_3^{\mathbf{S}}(u^{\mathbf{S}})}, \quad (9.8.18)$$

and,

$$\underline{A}^{\mathbf{S}} := \frac{2}{r^{\mathbf{S}}}(e_3^{\mathbf{S}}(r^{\mathbf{S}}) + \Upsilon^{\mathbf{S}}), \quad (9.8.19)$$

we deduce,

$$e_{\theta}^{\mathbf{S}}(\log \zeta^{\mathbf{S}}) = (\eta^{\mathbf{S}} - \zeta^{\mathbf{S}}), \quad (9.8.20)$$

$$e_{\theta}^{\mathbf{S}}(\underline{A}^{\mathbf{S}}) = -\frac{2\Upsilon^{\mathbf{S}}}{r^{\mathbf{S}}}(\zeta^{\mathbf{S}} - \eta^{\mathbf{S}}) - \frac{2}{r^{\mathbf{S}}}\underline{\xi}^{\mathbf{S}} + (\zeta^{\mathbf{S}} - \eta^{\mathbf{S}})\underline{A}^{\mathbf{S}}. \quad (9.8.21)$$

We infer that $e_\theta^{\mathbf{S}}(\log \varsigma^{\mathbf{S}})$ and $e_\theta^{\mathbf{S}}(\underline{A}^{\mathbf{S}})$ are determined in terms of $\eta, \underline{\xi}$.

Step 5. In view of the definition of $\nu^{\mathbf{S}}$ and $\varsigma^{\mathbf{S}}$ we make use of (9.8.17) to deduce

$$\nu^{\mathbf{S}}(u^{\mathbf{S}}) = e_3^{\mathbf{S}}(u^{\mathbf{S}}) + a^{\mathbf{S}}e_4^{\mathbf{S}}(u^{\mathbf{S}}) = \frac{2}{\varsigma^{\mathbf{S}}}.$$

On the other hand, since $u^{\mathbf{S}} := c_0 - r^{\mathbf{S}}$ along Σ_0 ,

$$\nu^{\mathbf{S}}(u^{\mathbf{S}}) = -\nu^{\mathbf{S}}(r^{\mathbf{S}}) = -e_3^{\mathbf{S}}(r^{\mathbf{S}}) - a^{\mathbf{S}}e_4^{\mathbf{S}}(r^{\mathbf{S}}) = \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2}\underline{A}^{\mathbf{S}} - a^{\mathbf{S}}$$

and therefore,

$$a^{\mathbf{S}} = -\frac{2}{\varsigma^{\mathbf{S}}} + \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2}\underline{A}^{\mathbf{S}} = -\frac{2}{\varsigma^{\mathbf{S}}} - \underline{\Omega}^{\mathbf{S}} \quad (9.8.22)$$

where,

$$\underline{\Omega}^{\mathbf{S}} := e_3^{\mathbf{S}}(r^{\mathbf{S}}) = -\Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2}\underline{A}^{\mathbf{S}}. \quad (9.8.23)$$

Step 6. The following lemma will be used, in particular³⁰, to determine the $\overline{A}^{\mathbf{S}}$.

Lemma 9.8.3. *For every scalar function h we have the formula*

$$\nu^{\mathbf{S}}\left(\int_{\mathbf{S}} h\right) = (\varsigma^{\mathbf{S}})^{-1} \int_{\mathbf{S}} \varsigma^{\mathbf{S}} (\nu^{\mathbf{S}}(h) + (\underline{\kappa}^{\mathbf{S}} + a^{\mathbf{S}}\kappa^{\mathbf{S}})h). \quad (9.8.24)$$

In particular

$$\nu^{\mathbf{S}}(r^{\mathbf{S}}) = \frac{r^{\mathbf{S}}}{2}(\varsigma^{\mathbf{S}})^{-1} \overline{\varsigma^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}} + a^{\mathbf{S}}\kappa^{\mathbf{S}})}$$

where the average is with respect to \mathbf{S} .

Proof. We consider the coordinates $u^{\mathbf{S}}, \theta^{\mathbf{S}}$ along Σ_0 with $\nu^{\mathbf{S}}(\theta^{\mathbf{S}}) = 0$. In these coordinates we have,

$$\nu^{\mathbf{S}} = \frac{2}{\varsigma^{\mathbf{S}}}\partial_{u^{\mathbf{S}}}.$$

The lemma follows easily by expressing the volume element of the surfaces $\mathbf{S} \subset \Sigma_0$ with respect to the coordinates $u^{\mathbf{S}}, \theta^{\mathbf{S}}$ (see also the proof of Proposition 2.2.9). \square

³⁰It will also be used below to derive equations for $\Lambda, \underline{\Lambda}$.

Step 7. Note that the GCM condition $\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}$ together with the definition of the Hawking mass implies that,

$$\overline{\underline{\kappa}^{\mathbf{S}}} = -\frac{2\Upsilon^{\mathbf{S}}}{r^{\mathbf{S}}}, \quad \Upsilon^{\mathbf{S}} = 1 - \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}},$$

where the average is taken with respect to \mathbf{S} . Thus in view of Lemma 9.8.3 we deduce

$$\begin{aligned} e_3^{\mathbf{S}}(r^{\mathbf{S}}) + a^{\mathbf{S}} &= \nu^{\mathbf{S}}(r^{\mathbf{S}}) = \frac{r^{\mathbf{S}}}{2}(\zeta^{\mathbf{S}})^{-1}\overline{\zeta^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}} + a^{\mathbf{S}}\kappa^{\mathbf{S}})} = \frac{r^{\mathbf{S}}}{2}(\zeta^{\mathbf{S}})^{-1}(\overline{\zeta^{\mathbf{S}}\underline{\kappa}^{\mathbf{S}}} + \overline{\zeta^{\mathbf{S}}\check{\underline{\kappa}}^{\mathbf{S}}}) + (\zeta^{\mathbf{S}})^{-1}\overline{\zeta^{\mathbf{S}}a^{\mathbf{S}}} \\ &= -\Upsilon^{\mathbf{S}}(\zeta^{\mathbf{S}})^{-1}\overline{\zeta^{\mathbf{S}}} + \frac{r^{\mathbf{S}}}{2}(\zeta^{\mathbf{S}})^{-1}\overline{\zeta^{\mathbf{S}}\check{\underline{\kappa}}^{\mathbf{S}}} + (\zeta^{\mathbf{S}})^{-1}\overline{\zeta^{\mathbf{S}}a^{\mathbf{S}}}. \end{aligned}$$

Since according to (9.8.19) $e_3^{\mathbf{S}}(r^{\mathbf{S}}) = -\Upsilon^{\mathbf{S}} + \frac{r^{\mathbf{S}}}{2}\underline{A}^{\mathbf{S}}$, we deduce

$$\underline{A}^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}} \left(\Upsilon^{\mathbf{S}} - a^{\mathbf{S}} - \Upsilon^{\mathbf{S}}(\zeta^{\mathbf{S}})^{-1}\overline{\zeta^{\mathbf{S}}} + \frac{r^{\mathbf{S}}}{2}(\zeta^{\mathbf{S}})^{-1}\overline{\zeta^{\mathbf{S}}\check{\underline{\kappa}}^{\mathbf{S}}} + (\zeta^{\mathbf{S}})^{-1}\overline{\zeta^{\mathbf{S}}a^{\mathbf{S}}} \right).$$

In particular, multiplying by $\zeta^{\mathbf{S}}$ and taking the average, we infer

$$\overline{\zeta^{\mathbf{S}}\underline{A}^{\mathbf{S}}} = \overline{\zeta^{\mathbf{S}}\check{\underline{\kappa}}^{\mathbf{S}}},$$

and hence

$$\underline{A}^{\mathbf{S}} = \frac{1}{\zeta^{\mathbf{S}}} \left(\overline{\zeta^{\mathbf{S}}\check{\underline{\kappa}}^{\mathbf{S}}} - \overline{\zeta^{\mathbf{S}}\underline{A}^{\mathbf{S}}} \right). \tag{9.8.25}$$

Step 8. We summarize the results in Steps 1-7 in the following.

Proposition 9.8.4. *Let Σ_0 be a smooth spacelike hypersurface foliated by framed³¹ spheres $(\mathbf{S}, e_4^{\mathbf{S}}, e_\theta^{\mathbf{S}}, e_3^{\mathbf{S}})$ whose Ricci coefficients verify the GCM condition $\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}$ and transversality condition (9.4.12). Define $u^{\mathbf{S}}$ as in (9.8.16) such that $u^{\mathbf{S}} + r^{\mathbf{S}}$ is constant on Σ_0 with $r^{\mathbf{S}}$ the area radius of the spheres \mathbf{S} . Extend $u^{\mathbf{S}}$ and $r^{\mathbf{S}}$ in a neighborhood of Σ_0 such that the transversality conditions (9.8.17) are verified. Then, defining the scalars $\zeta^{\mathbf{S}}, \underline{A}^{\mathbf{S}}$ as in (9.8.18), (9.8.19) we establish the following relations between $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}$ and $\zeta^{\mathbf{S}}, \underline{A}^{\mathbf{S}}$ and $a^{\mathbf{S}}$, where the latter scalar is defined in Step 1,*

$$\begin{aligned} e_\theta^{\mathbf{S}}(\log \zeta^{\mathbf{S}}) &= (\eta^{\mathbf{S}} - \zeta^{\mathbf{S}}), \\ e_\theta^{\mathbf{S}}(\underline{A}^{\mathbf{S}}) &= -\frac{2\Upsilon^{\mathbf{S}}}{r^{\mathbf{S}}}(\zeta^{\mathbf{S}} - \eta^{\mathbf{S}}) - \frac{2}{r^{\mathbf{S}}}\underline{\xi}^{\mathbf{S}} + (\zeta^{\mathbf{S}} - \eta^{\mathbf{S}})\underline{A}^{\mathbf{S}}, \\ \underline{A}^{\mathbf{S}} &= \frac{1}{\zeta^{\mathbf{S}}} \left(\overline{\zeta^{\mathbf{S}}\check{\underline{\kappa}}^{\mathbf{S}}} - \overline{\zeta^{\mathbf{S}}\underline{A}^{\mathbf{S}}} \right), \\ a^{\mathbf{S}} &= -\frac{2}{\zeta^{\mathbf{S}}} + \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2}\underline{A}^{\mathbf{S}}. \end{aligned} \tag{9.8.26}$$

³¹i.e. differentiable spheres \mathbf{S} endowed with adapted null frames $(e_4^{\mathbf{S}}, e_\theta^{\mathbf{S}}, e_3^{\mathbf{S}})$.

Remark 9.8.5. Note that we lack equations for $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}$ and the average of $a^{\mathbf{S}}$. The latter can be fixed by fixing the value of $a^{\mathbf{S}}|_{SP}$ and observing that

$$\bar{a}^{\mathbf{S}} = a^{\mathbf{S}}|_{SP} - \check{a}^{\mathbf{S}}|_{SP}. \quad (9.8.27)$$

In what follows we state a result which ties $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}$ to the other GCM conditions in (9.8.13)–(9.8.14).

Step 9. To state the proposition below we split the Ricci coefficients into the following groups.

$$\begin{aligned} \Gamma_g^{\mathbf{S}} &= \left\{ \check{\kappa}^{\mathbf{S}}, \vartheta^{\mathbf{S}}, \zeta^{\mathbf{S}}, \check{\kappa}^{\mathbf{S}}, r\check{\rho}^{\mathbf{S}}, \overline{\kappa}^{\mathbf{S}} - \frac{2}{r^{\mathbf{S}}}, \overline{\rho}^{\mathbf{S}} + \frac{2m^{\mathbf{S}}}{(r^{\mathbf{S}})^3} \right\}, \\ \Gamma_b^{\mathbf{S}} &= \left\{ \eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}, \check{\omega}^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}} - \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^2}, \underline{\beta}^{\mathbf{S}}, \underline{\alpha}^{\mathbf{S}} \right\}. \end{aligned}$$

Proposition 9.8.6. The following statements hold true³²,

1. Under the same assumptions as in Proposition 9.8.4, the Ricci coefficients $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}}$ verify the following identities.

$$\begin{aligned} 2\check{\phi}_2^{\mathbf{S},*} \check{\phi}_1^{\mathbf{S},*} \check{\phi}_1^{\mathbf{S}} \check{\phi}_2^{\mathbf{S}} \check{\phi}_2^{\mathbf{S},*} \eta^{\mathbf{S}} &= \kappa^{\mathbf{S}} \left(C_1 + 2\check{\phi}_2^{\mathbf{S},*} \check{\phi}_1^{\mathbf{S},*} \check{\phi}_1^{\mathbf{S}} \underline{\beta}^{\mathbf{S}} \right) - r^{\mathbf{S}-3} \check{\rho}^3 C_2 \\ &\quad + r^{\mathbf{S}-5} (\check{\rho}^{\mathbf{S}})^{\leq 4} \Gamma_g^{\mathbf{S}} + r^{\mathbf{S}-4} (\check{\rho}^{\mathbf{S}})^{\leq 4} (\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}) + l.o.t., \\ 2\check{\phi}_2^{\mathbf{S},*} \check{\phi}_1^{\mathbf{S},*} \check{\phi}_1^{\mathbf{S}} \check{\phi}_2^{\mathbf{S}} \check{\phi}_2^{\mathbf{S},*} \underline{\xi} &= C_3 - \check{\kappa}^{\mathbf{S}} \left(C_1 + 2\check{\phi}_2^{\mathbf{S},*} \check{\phi}_1^{\mathbf{S},*} \check{\phi}_1^{\mathbf{S}} \underline{\beta}^{\mathbf{S}} \right) \\ &\quad + r^{\mathbf{S}-5} (\check{\rho}^{\mathbf{S}})^{\leq 4} \Gamma_g^{\mathbf{S}} + r^{\mathbf{S}-4} (\check{\rho}^{\mathbf{S}})^{\leq 4} (\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}) + l.o.t., \quad (9.8.28) \\ \check{\phi}_1^{\mathbf{S},*} \underline{\omega}^{\mathbf{S}} &= \left(\frac{1}{4} \check{\kappa}^{\mathbf{S}} + \underline{\omega}^{\mathbf{S}} \right) \eta^{\mathbf{S}} - (\kappa^{\mathbf{S}})^{-1} \check{\phi}_2^{\mathbf{S}} \check{\phi}_2^{\mathbf{S},*} \eta^{\mathbf{S}} \\ &\quad + \frac{1}{4} \kappa^{\mathbf{S}} \underline{\xi}^{\mathbf{S}} - \frac{1}{2} (\kappa^{\mathbf{S}})^{-1} C_2 + r^{\mathbf{S}-1} (\check{\rho}^{\mathbf{S}})^{\leq 1} \Gamma_g^{\mathbf{S}} + \check{\rho}^{\mathbf{S}} (\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}), \end{aligned}$$

where,

$$\begin{aligned} C_1 &= e_3^{\mathbf{S}}(\check{\phi}_2^{\mathbf{S},*} \check{\phi}_1^{\mathbf{S},*} \mu^{\mathbf{S}}), \\ C_2 &= e_3^{\mathbf{S}}(e_{\theta}^{\mathbf{S}} \kappa^{\mathbf{S}}), \\ C_3 &= e_3^{\mathbf{S}}\left((\check{\phi}_2^{\mathbf{S},*} \check{\phi}_2^{\mathbf{S}} + 2K^{\mathbf{S}}) \check{\phi}_2^{\mathbf{S},*} \check{\phi}_1^{\mathbf{S},*} \check{\kappa}^{\mathbf{S}} \right). \end{aligned} \quad (9.8.29)$$

The quadratic terms denoted *l.o.t.* are lower order both in terms of decay in $r^{\mathbf{S}}$ as well as in terms of number of derivatives. They also contain only angular derivatives $\check{\rho}^{\mathbf{S}}$

³² $r_{\mathbf{S}}$ here denotes $r^{\mathbf{S}}$ the area radius of \mathbf{S} .

and not $e_3^{\mathbf{S}}$ or $e_4^{\mathbf{S}}$. We also note that the error terms $r_{\mathbf{S}}^{-5}(\vartheta^{\mathbf{S}})^{\leq 4}\Gamma_g^{\mathbf{S}} + r_{\mathbf{S}}^{-4}(\vartheta^{\mathbf{S}})^{\leq 4}(\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}})$ does not in fact contain more than 3 derivatives of $\underline{\omega}^{\mathbf{S}}$.

2. If in addition (9.4.11) of Theorem 9.4.2 hold true then, for $k \leq s_{max} - 7$,

$$\begin{aligned} \|\phi_2^{\mathbf{S},*}\eta^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} &\lesssim r_{\mathbf{S}}^3\|C_1\|_{\mathfrak{h}_k(\mathbf{S})} + r\|C_2\|_{\mathfrak{h}_{3+k}(\mathbf{S})} + \overset{\circ}{e}r_{\mathbf{S}}^{-1} \\ &\quad + r_{\mathbf{S}}^{-1}\|\Gamma_g^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + l.o.t., \\ \|\phi_2^{\mathbf{S},*}\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} &\lesssim r_{\mathbf{S}}^4\|C_3\|_{\mathfrak{h}_k(\mathbf{S})} + r_{\mathbf{S}}^3\|C_1\|_{\mathfrak{h}_{3+k}(\mathbf{S})} + \overset{\circ}{e}r_{\mathbf{S}}^{-1} \\ &\quad + r_{\mathbf{S}}^{-1}\|\Gamma_g^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + l.o.t., \\ \|\phi_1^{\mathbf{S},*}\underline{\omega}\|_{\mathfrak{h}_{2+k}(\mathbf{S})} &\lesssim r_{\mathbf{S}}^{-1}\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + r_{\mathbf{S}}^{-1}\|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{2+k}(\mathbf{S})} + r\|C_2\|_{\mathfrak{h}_{2+k}(\mathbf{S})} \\ &\quad + r_{\mathbf{S}}^{-1}\|\Gamma_g^{\mathbf{S}}\|_{\mathfrak{h}_{3+k}(\mathbf{S})} + \|\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}\|_{\mathfrak{h}_{3+k}(\mathbf{S})} + l.o.t. \end{aligned} \tag{9.8.30}$$

3. If in addition the GCM conditions (9.8.13) hold true along Σ_0 and the estimates (9.4.14) are also verified then, for $k \leq s_{max} - 7$,

$$\begin{aligned} \|C_1\|_{\mathfrak{h}_{k-2}(\mathbf{S})} &\lesssim \overset{\circ}{e}r^{-5} \left(|\overline{a^{\mathbf{S}}} + 1 + \frac{2m^{\mathbf{S}}}{r_{\mathbf{S}}}| + r^{-1}\|\check{a}^{\mathbf{S}}\|_{\mathfrak{h}_{k-2}(\mathbf{S})} \right), \\ \|C_2\|_{\mathfrak{h}_{k-1}(\mathbf{S})} &\lesssim \overset{\circ}{e}r^{-3} \left(|\overline{a^{\mathbf{S}}} + 1 + \frac{2m^{\mathbf{S}}}{r_{\mathbf{S}}}| + r^{-1}\|\check{a}^{\mathbf{S}}\|_{\mathfrak{h}_{k-1}(\mathbf{S})} \right), \\ \|C_3\|_{\mathfrak{h}_{k-4}(\mathbf{S})} &\lesssim \overset{\circ}{e}r^{-5} \left(|\overline{a^{\mathbf{S}}} + 1 + \frac{2m^{\mathbf{S}}}{r_{\mathbf{S}}}| + r^{-1}\|\check{a}^{\mathbf{S}}\|_{\mathfrak{h}_{k-4}(\mathbf{S})} \right), \end{aligned} \tag{9.8.31}$$

where $a^{\mathbf{S}}$ was defined in Step 1 and can be expressed in terms of $\zeta^{\mathbf{S}}$ and $\underline{A}^{\mathbf{S}}$ by formula (9.8.22).

Proof. The proof³³ of the first two identities in (9.8.28) were derived in Proposition 7.3.4 in connection to the proof³⁴ of Theorem M4, starting with the following³⁵

$$\begin{aligned} 2\phi_1^{\mathbf{S},*}\underline{\omega}^{\mathbf{S}} &= \left(\frac{1}{2}\underline{\kappa}^{\mathbf{S}} + 2\underline{\omega}^{\mathbf{S}} \right) \eta^{\mathbf{S}} + e_3^{\mathbf{S}}(\zeta^{\mathbf{S}}) - \underline{\beta}^{\mathbf{S}} + \frac{1}{2}\kappa\underline{\xi}^{\mathbf{S}} + r^{-1}\Gamma_g^{\mathbf{S}} + \Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}, \\ 2\phi_2^{\mathbf{S}}\phi_2^{\mathbf{S},*}\eta^{\mathbf{S}} &= \kappa^{\mathbf{S}}(-e_3(\zeta^{\mathbf{S}}) + \underline{\beta}^{\mathbf{S}}) - e_3^{\mathbf{S}}(e_{\theta}^{\mathbf{S}}(\kappa^{\mathbf{S}})) + r_{\mathbf{S}}^{-2}(\vartheta^{\mathbf{S}})^{\leq 1}\Gamma_g^{\mathbf{S}} + r_{\mathbf{S}}^{-1}\vartheta(\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}), \\ 2\phi_2^{\mathbf{S}}\phi_2^{\mathbf{S},*}\underline{\xi}^{\mathbf{S}} &= \underline{\kappa}^{\mathbf{S}}(e_3(\zeta^{\mathbf{S}}) - \underline{\beta}^{\mathbf{S}}) - e_3^{\mathbf{S}}(e_{\theta}^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}})) + r_{\mathbf{S}}^{-2}(\vartheta^{\mathbf{S}})^{\leq 1}\Gamma_g^{\mathbf{S}} + r_{\mathbf{S}}^{-1}\vartheta^{\mathbf{S}}(\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}). \end{aligned} \tag{9.8.32}$$

³³The equations used in the derivation of these identities only require the transversality conditions (9.4.12).

³⁴Strictly speaking Proposition 7.3.4 requires the e_3 Ricci and Bianchi identities of a geodesic foliation. It is easy to justify the application of these equations in our context by using the transversality conditions to generate a geodesic foliation in a neighborhood of Σ_0 .

³⁵These identities were recorded in Proposition 7.1.12 which was itself a corollary Proposition 2.2.19. Note also that $\vartheta(\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}})$ does not contain derivatives of $\underline{\omega}$.

The last identity in (9.8.28) follows by combining the first two identities in (9.8.32).

To prove the estimates for $\eta^{\mathbf{S}}$ in the second part of the proposition we make use of the identity $\not{d}_1^{\mathbf{S},*} \not{d}_1^{\mathbf{S}} = \not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} + 2K^{\mathbf{S}}$ to deduce,

$$\begin{aligned} \not{d}_2^{\mathbf{S},*} (\not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} + 2K^{\mathbf{S}}) \not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} \eta^{\mathbf{S}} &= \frac{1}{2} \kappa^{\mathbf{S}} C_1 + \kappa^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} (\not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} + 2K^{\mathbf{S}}) \underline{\beta}^{\mathbf{S}} - r_{\mathbf{S}}^{-3} (\not{\vartheta}^{\mathbf{S}})^3 C_2 \\ &+ r_{\mathbf{S}}^{-5} (\not{\vartheta}^{\mathbf{S}})^{\leq 4} \Gamma_g^{\mathbf{S}} + r_{\mathbf{S}}^{-4} (\not{\vartheta}^{\mathbf{S}})^{\leq 4} (\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}) + \text{l.o.t.} \end{aligned}$$

i.e.,

$$\begin{aligned} \not{d}_2^{\mathbf{S},*} (\not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} + 2K^{\mathbf{S}}) \left(\not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} \eta^{\mathbf{S}} - \kappa^{\mathbf{S}} \underline{\beta}^{\mathbf{S}} \right) &= \frac{1}{2} \kappa^{\mathbf{S}} C_1 - \frac{1}{2} r_{\mathbf{S}}^{-3} (\not{\vartheta}^{\mathbf{S}})^3 C_2 \\ &+ r_{\mathbf{S}}^{-5} (\not{\vartheta}^{\mathbf{S}})^{\leq 4} \Gamma_g^{\mathbf{S}} + r_{\mathbf{S}}^{-4} (\not{\vartheta}^{\mathbf{S}})^{\leq 4} (\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}) + \text{l.o.t.} \end{aligned}$$

Similarly for $\underline{\xi}^{\mathbf{S}}$

$$\begin{aligned} \not{d}_2^{\mathbf{S},*} (\not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} + 2K^{\mathbf{S}}) \left(\not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} \underline{\xi}^{\mathbf{S}} + \underline{\kappa}^{\mathbf{S}} \underline{\beta}^{\mathbf{S}} \right) &= \frac{1}{2} C_3 - \frac{1}{2} \underline{\kappa}^{\mathbf{S}} C_1 \\ &+ r_{\mathbf{S}}^{-5} (\not{\vartheta}^{\mathbf{S}})^{\leq 4} \Gamma_g^{\mathbf{S}} + r_{\mathbf{S}}^{-4} (\not{\vartheta}^{\mathbf{S}})^{\leq 4} (\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}) + \text{l.o.t.} \end{aligned}$$

The desired estimates for $\eta^{\mathbf{S}}$ and $\underline{\xi}^{\mathbf{S}}$ follow then by making use of the coercivity of the operator $\not{d}_2^{\mathbf{S},*} (\not{d}_2^{\mathbf{S}} \not{d}_2^{\mathbf{S},*} + 2K^{\mathbf{S}})$. and the estimate for $\underline{\beta} = \underline{\beta}^{\mathbf{S}}$ in (9.4.14). The estimate for $\not{d}_1^{\mathbf{S},*} \underline{\omega}^{\mathbf{S}}$ is straightforward from the last identity in (9.8.32).

To prove the last part of the proposition we make use of the GCM conditions (9.8.13) on Σ_0 to deduce that

$$\nu^{\mathbf{S}} (\not{d}_2^{\mathbf{S},*} \not{d}_1^{\mathbf{S},*} \mu^{\mathbf{S}}) = 0, \quad \nu^{\mathbf{S}} (e_{\theta}^{\mathbf{S}} \kappa^{\mathbf{S}}) = 0, \quad \nu^{\mathbf{S}} \left((\not{d}_2^{\mathbf{S},*} \not{d}_2^{\mathbf{S}} + 2K^{\mathbf{S}}) \not{d}_2^{\mathbf{S},*} \not{d}_1^{\mathbf{S},*} \underline{\kappa}^{\mathbf{S}} \right) = 0.$$

Hence, the quantities C_1, C_2, C_3 in (9.8.29) can be expressed in the form

$$\begin{aligned} C_1 &= -a^{\mathbf{S}} e_4^{\mathbf{S}} (\not{d}_2^{\mathbf{S},*} \not{d}_1^{\mathbf{S},*} \mu^{\mathbf{S}}), \\ C_2 &= -a^{\mathbf{S}} e_4^{\mathbf{S}} (e_{\theta}^{\mathbf{S}} \kappa^{\mathbf{S}}), \\ C_3 &= a^{\mathbf{S}} e_4^{\mathbf{S}} \left((\not{d}_2^{\mathbf{S},*} \not{d}_2^{\mathbf{S}} + 2K^{\mathbf{S}}) \not{d}_2^{\mathbf{S},*} \not{d}_1^{\mathbf{S},*} \underline{\kappa}^{\mathbf{S}} \right). \end{aligned}$$

Making use of our commutation formulas of Lemma 2.2.13 and the estimates (9.4.14) and (9.4.11) we easily deduce,

$$\begin{aligned} \|e_4^{\mathbf{S}} (\not{d}_2^{\mathbf{S},*} \not{d}_1^{\mathbf{S},*} \mu^{\mathbf{S}})\|_{\mathfrak{h}_{k-2}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-5}, \\ \|e_4^{\mathbf{S}} (e_{\theta}^{\mathbf{S}} \kappa^{\mathbf{S}})\|_{\mathfrak{h}_{k-1}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-3}. \end{aligned}$$

Similarly,

$$\left\| e_4^{\mathbf{S}} \left((\phi_2^{\mathbf{S},*} \phi_2^{\mathbf{S}} + 2K^{\mathbf{S}}) \phi_2^{\mathbf{S},*} \phi_1^{\mathbf{S},*} \underline{\kappa}^{\mathbf{S}} \right) \right\|_{\mathfrak{h}_{k-4}(\mathbf{S})} \lesssim \overset{\circ}{\epsilon} r^{-5}.$$

Writing $a^{\mathbf{S}} = \overline{a^{\mathbf{S}}} + \check{a}^{\mathbf{S}}$ and making use of product estimates we deduce

$$\begin{aligned} \|C_1\|_{\mathfrak{h}_{k-2}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-5} \left(|\overline{a^{\mathbf{S}}} + 1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}| + r^{-1} \|\check{a}^{\mathbf{S}}\|_{\mathfrak{h}_{k-2}(\mathbf{S})} \right), \\ \|C_2\|_{\mathfrak{h}_{k-1}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-3} \left(|\overline{a^{\mathbf{S}}} + 1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}| + r^{-1} \|\check{a}^{\mathbf{S}}\|_{\mathfrak{h}_{k-1}(\mathbf{S})} \right), \\ \|C_3\|_{\mathfrak{h}_{k-4}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r^{-5} \left(|\overline{a^{\mathbf{S}}} + 1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}| + r^{-1} \|\check{a}^{\mathbf{S}}\|_{\mathfrak{h}_{k-4}(\mathbf{S})} \right), \end{aligned}$$

as stated. □

Step 10. Propositions 9.8.4 and 9.8.6 provide us with potential³⁶ estimates for $\phi_2^{\mathbf{S},*} \eta^{\mathbf{S}}$, $\phi_2^{\mathbf{S},*} \underline{\xi}^{\mathbf{S}}$, $\phi_1^{\mathbf{S},*} \underline{\omega}^{\mathbf{S}}$, $\phi_1^{\mathbf{S},*} \underline{\varsigma}^{\mathbf{S}}$. To close we also need to control the $\ell = 1$ modes of $\eta^{\mathbf{S}}$, $\underline{\xi}^{\mathbf{S}}$ the average of $\underline{\omega}^{\mathbf{S}}$ and the average³⁷ of $a^{\mathbf{S}}$. Note that the average of $\underline{\omega}^{\mathbf{S}}$ can in fact be derived from the equation,

$$e_3^{\mathbf{S}}(\kappa^{\mathbf{S}}) + \frac{1}{2} \underline{\kappa}^{\mathbf{S}} \kappa^{\mathbf{S}} - 2 \underline{\omega}^{\mathbf{S}} \kappa^{\mathbf{S}} = 2 \phi_1^{\mathbf{S}} \eta^{\mathbf{S}} + 2 \rho^{\mathbf{S}} - \frac{1}{2} \underline{\vartheta}^{\mathbf{S}} \vartheta^{\mathbf{S}} + 2(\eta^{\mathbf{S}})^2$$

in terms of $\overline{A^{\mathbf{S}}}$ and $\eta^{\mathbf{S}}$. Indeed, making use of the GCM condition $\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}$,

$$\begin{aligned} \underline{\omega}^{\mathbf{S}} &= \frac{1}{2\kappa^{\mathbf{S}}} \left[e_3^{\mathbf{S}}(\kappa^{\mathbf{S}}) + \frac{1}{2} \underline{\kappa}^{\mathbf{S}} \kappa^{\mathbf{S}} - 2 \phi_1^{\mathbf{S}} \eta^{\mathbf{S}} - 2 \rho^{\mathbf{S}} + \frac{1}{2} \underline{\vartheta}^{\mathbf{S}} \vartheta^{\mathbf{S}} - 2(\eta^{\mathbf{S}})^2 \right] \\ &= -\frac{1}{2} e_3(r^{\mathbf{S}}) - \frac{\gamma^{\mathbf{S}}}{2r^{\mathbf{S}}} + \frac{r^{\mathbf{S}}}{4} \left[-2 \phi_1^{\mathbf{S}} \eta^{\mathbf{S}} - 2 \rho^{\mathbf{S}} + \frac{1}{2} \underline{\vartheta}^{\mathbf{S}} \vartheta^{\mathbf{S}} - 2(\eta^{\mathbf{S}})^2 \right] \\ &= -\frac{1}{4} \overline{A^{\mathbf{S}}} + \frac{r^{\mathbf{S}}}{4} \left[-2 \phi_1^{\mathbf{S}} \eta^{\mathbf{S}} - 2 \rho^{\mathbf{S}} + \frac{1}{2} \underline{\vartheta}^{\mathbf{S}} \vartheta^{\mathbf{S}} - 2(\eta^{\mathbf{S}})^2 \right]. \end{aligned}$$

Thus, recalling the definition of $\mu^{\mathbf{S}}$,

$$\overline{\omega}^{\mathbf{S}} = -\frac{1}{4} \overline{A^{\mathbf{S}}} + \frac{r^{\mathbf{S}}}{2} \overline{\mu^{\mathbf{S}} - (\eta^{\mathbf{S}})^2}$$

³⁶We cannot close the estimates without being also able to estimate the $\ell = 1$ modes of $\eta^{\mathbf{S}}$, $\underline{\xi}^{\mathbf{S}}$, $\overline{\omega}^{\mathbf{S}}$ and the average $\overline{a^{\mathbf{S}}}$.

³⁷The quantity $\check{a}^{\mathbf{S}}$ can be determined using Proposition 9.8.4.

or,

$$\underline{\omega}^{\mathbf{S}} - \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^2} = -\frac{1}{4}\underline{A}^{\mathbf{S}} + \frac{r^{\mathbf{S}}}{2} \left(\overline{\mu^{\mathbf{S}} - \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^3}} - \overline{\eta^{\mathbf{S}} \cdot \eta^{\mathbf{S}}} \right). \quad (9.8.33)$$

Step 11. In view of the above we can determine $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}}, \varsigma^{\mathbf{S}}, \underline{A}^{\mathbf{S}}$ provided that we control the $\ell = 1$ modes of $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}$ and the average of $\varsigma^{\mathbf{S}}$. For this reason we introduce³⁸, along Σ_0 ,

$$B^{\mathbf{S}} = \int_{\mathbf{S}} \eta^{\mathbf{S}} e^{\Phi}, \quad \underline{B}^{\mathbf{S}} = \int_{\mathbf{S}} \underline{\xi}^{\mathbf{S}} e^{\Phi}, \quad D^{\mathbf{S}} = a^{\mathbf{S}} \Big|_{SP} + 1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}. \quad (9.8.34)$$

We are now ready to prove the following

Proposition 9.8.7. *Let Σ_0 be a smooth spacelike hypersurface foliated by framed spheres $(\mathbf{S}, e_4^{\mathbf{S}}, e_{\theta}^{\mathbf{S}}, e_3^{\mathbf{S}})$ which verify the GCM conditions (9.8.13), transversality condition (9.4.12) and the estimates (9.4.11)–(9.4.14) of Theorem 9.4.2. Let $u^{\mathbf{S}}$ as in (9.8.16) such that $u^{\mathbf{S}} + r^{\mathbf{S}}$ is constant on Σ_0 . Extend $u^{\mathbf{S}}$ and $r^{\mathbf{S}}$ in a neighborhood of Σ_0 such that the transversality conditions (9.8.17) are verified. As shown above these allow us to define $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}}, \varsigma^{\mathbf{S}}, \underline{A}^{\mathbf{S}}, a^{\mathbf{S}}$ and the constants $B^{\mathbf{S}}, \underline{B}^{\mathbf{S}}, D^{\mathbf{S}}$ as in (9.8.34). Finally we assume that,*

$$r^{-2}(|B^{\mathbf{S}}| + |\underline{B}^{\mathbf{S}}|) + |D^{\mathbf{S}}| \leq \overset{\circ}{\epsilon}^{1/2}. \quad (9.8.35)$$

Under these assumptions the following estimates hold true for all $k \leq s_{max} - 7$,

1. The Ricci coefficients $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}}$ verify

$$\begin{aligned} \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_{5+k}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-2} |B^{\mathbf{S}}|, \\ \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{5+k}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-2} |\underline{B}^{\mathbf{S}}|, \\ \|\underline{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{3+k}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-2} (|\underline{B}^{\mathbf{S}}| + |B^{\mathbf{S}}|), \\ \left| \underline{\omega}^{\mathbf{S}} - \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^2} \right| &\lesssim \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-2} (|\underline{B}^{\mathbf{S}}| + |B^{\mathbf{S}}|). \end{aligned} \quad (9.8.36)$$

2. The scalar $a^{\mathbf{S}}$ verifies,

$$r_{\mathbf{S}}^{-1} \|\check{a}^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} + \left| \bar{a}^{\mathbf{S}} + 1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} \right| \lesssim \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-2} |B^{\mathbf{S}}| + |D^{\mathbf{S}}|. \quad (9.8.37)$$

³⁸Note that to prove our main theorem we have to construct our hypersurface Σ_0 such that in fact $B = \underline{B} = D = 0$.

3. We also have

$$\begin{aligned} \|\underline{A}^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-2} (|\underline{B}^{\mathbf{S}}| + |B^{\mathbf{S}}|) + |D^{\mathbf{S}}|, \\ r_{\mathbf{S}}^{-1} \|\zeta^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} + |\overline{\zeta}^{\mathbf{S}} - 1| &\lesssim \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-2} (|\underline{B}^{\mathbf{S}}| + |B^{\mathbf{S}}|) + |D^{\mathbf{S}}|. \end{aligned} \quad (9.8.38)$$

4. We also have, for all $k \leq s_{max} - 4$

$$\begin{aligned} \|e_3^{\mathbf{S}}(\check{\kappa}^{\mathbf{S}}, \vartheta^{\mathbf{S}}, \zeta^{\mathbf{S}}, \check{\kappa}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r_{\mathbf{S}}^{-1}, \\ \|e_3^{\mathbf{S}}(\underline{\vartheta}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon}, \\ \|e_3^{\mathbf{S}}(\alpha^{\mathbf{S}}, \beta^{\mathbf{S}}, \check{\rho}^{\mathbf{S}}, \mu^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r_{\mathbf{S}}^{-2}, \\ \|e_3^{\mathbf{S}}(\underline{\beta}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} r_{\mathbf{S}}^{-1}, \\ \|e_3^{\mathbf{S}}(\underline{\alpha}^{\mathbf{S}})\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon}. \end{aligned}$$

Proof. To simplify the exposition below we make the auxiliary bootstrap assumptions,

$$\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_{5+k}(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{5+k}(\mathbf{S})} \lesssim \overset{\circ}{\epsilon}^{1/2}. \quad (9.8.39)$$

We start with the following lemma.

Lemma 9.8.8. *The following estimates hold true*

$$r_{\mathbf{S}}^{-1} \|\check{a}^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} + \left| \overline{a}^{\mathbf{S}} + 1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} \right| \lesssim |D^{\mathbf{S}}| + \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \overset{\circ}{\epsilon}. \quad (9.8.40)$$

Proof. Since $a^{\mathbf{S}} = \overline{a}^{\mathbf{S}} + \check{a}^{\mathbf{S}}$ we deduce $a^{\mathbf{S}}|_{SP} = \overline{a}^{\mathbf{S}} + \check{a}^{\mathbf{S}}|_{SP}$. Hence,

$$\overline{a}^{\mathbf{S}} = D^{\mathbf{S}} - 1 - \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} - \check{a}^{\mathbf{S}}|_{SP}. \quad (9.8.41)$$

We also have (see Proposition 9.8.4)

$$a^{\mathbf{S}} = -\frac{2}{\zeta^{\mathbf{S}}} + \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2} \underline{A}^{\mathbf{S}}.$$

Hence,

$$a^{\mathbf{S}} = -\frac{2}{\overline{\zeta}^{\mathbf{S}} + \zeta^{\mathbf{S}}} + \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2} \underline{A}^{\mathbf{S}} = -\frac{2}{\overline{\zeta}^{\mathbf{S}}} \left(1 - \frac{\zeta^{\mathbf{S}}}{\overline{\zeta}^{\mathbf{S}}} + O\left(\left(\frac{\zeta^{\mathbf{S}}}{\overline{\zeta}^{\mathbf{S}}}\right)^2\right) \right) + \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2} \underline{A}^{\mathbf{S}}.$$

Taking the average on \mathbf{S} we deduce,

$$\overline{a}^{\mathbf{S}} = -\frac{2}{\overline{\zeta}^{\mathbf{S}}} + \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2} \overline{A}^{\mathbf{S}} + O\left(\left(\frac{\zeta^{\mathbf{S}}}{\overline{\zeta}^{\mathbf{S}}}\right)^2\right). \quad (9.8.42)$$

Also, using (9.8.41),

$$\check{a}^{\mathbf{S}} = 2\zeta^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2}\check{\underline{A}}^{\mathbf{S}} + \text{l.o.t.} \quad (9.8.43)$$

where l.o.t. denotes higher order terms in $\zeta^{\mathbf{S}}$ and $\overline{\zeta^{\mathbf{S}}} - 1$. Indeed

$$\begin{aligned} \check{a}^{\mathbf{S}} = a^{\mathbf{S}} - \overline{a^{\mathbf{S}}} &= -\frac{2}{\zeta^{\mathbf{S}}} + \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2}\underline{A}^{\mathbf{S}} - \left(-\frac{2}{\overline{\zeta^{\mathbf{S}}}} + \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2}\overline{\underline{A}}^{\mathbf{S}} \right) \\ &= -\frac{2}{\zeta^{\mathbf{S}}} + \frac{2}{\overline{\zeta^{\mathbf{S}}}} - \frac{r^{\mathbf{S}}}{2}\check{\underline{A}}^{\mathbf{S}} = \frac{2\zeta^{\mathbf{S}}}{\zeta^{\mathbf{S}}\overline{\zeta^{\mathbf{S}}}} - \frac{r^{\mathbf{S}}}{2}\check{\underline{A}}^{\mathbf{S}} = 2\zeta^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2}\check{\underline{A}}^{\mathbf{S}} + \text{l.o.t.} \end{aligned}$$

Thus to estimate $\check{a}^{\mathbf{S}}$ and $\overline{a^{\mathbf{S}}}$ we first need to estimate $\underline{A}^{\mathbf{S}}$, $\zeta^{\mathbf{S}}$ and $\overline{\zeta^{\mathbf{S}}}$. Using the equations (see Proposition 9.8.4)

$$\begin{aligned} e_{\theta}^{\mathbf{S}}(\underline{A}^{\mathbf{S}}) &= -\frac{2\Upsilon^{\mathbf{S}}}{r^{\mathbf{S}}}(\zeta^{\mathbf{S}} - \eta^{\mathbf{S}}) - \frac{2}{r^{\mathbf{S}}}\xi^{\mathbf{S}} + (\zeta^{\mathbf{S}} - \eta^{\mathbf{S}})\underline{A}^{\mathbf{S}}, \\ \overline{\underline{A}}^{\mathbf{S}} &= \frac{1}{\overline{\zeta^{\mathbf{S}}}} \left(\overline{\zeta^{\mathbf{S}}\check{\underline{A}}^{\mathbf{S}}} - \overline{\zeta^{\mathbf{S}}\underline{A}^{\mathbf{S}}} \right), \end{aligned}$$

and the auxiliary assumption we derive,

$$\|\underline{A}^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} \lesssim \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \|\xi^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \mathring{\epsilon}r^{-1} \left(1 + \|\zeta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + |\overline{\zeta^{\mathbf{S}}} - 1| \right). \quad (9.8.44)$$

From the equation

$$e_{\theta}^{\mathbf{S}}(\log \zeta^{\mathbf{S}}) = (\eta^{\mathbf{S}} - \zeta^{\mathbf{S}}).$$

we also derive,

$$r_{\mathbf{S}}^{-1}\|\zeta^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} \lesssim \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \mathring{\epsilon} + \mathring{\epsilon}|\overline{\zeta^{\mathbf{S}}} - 1|. \quad (9.8.45)$$

To estimate $\overline{\zeta^{\mathbf{S}}} - 1$ we derive from (9.8.42) and (9.8.43),

$$\begin{aligned} \frac{2}{\overline{\zeta^{\mathbf{S}}}} &= -\overline{a^{\mathbf{S}}} + \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2}\overline{\underline{A}}^{\mathbf{S}} = -\left(D^{\mathbf{S}} - 1 - \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} - \check{a}^{\mathbf{S}}|_{SP} \right) + \Upsilon^{\mathbf{S}} - \frac{r^{\mathbf{S}}}{2}\overline{\underline{A}}^{\mathbf{S}} + \text{l.o.t.} \\ &= -D^{\mathbf{S}} + 2 + \check{a}^{\mathbf{S}}|_{SP} - \frac{r^{\mathbf{S}}}{2}\overline{\underline{A}}^{\mathbf{S}} + \text{l.o.t.} \\ &= -D^{\mathbf{S}} + 2 + 2\zeta^{\mathbf{S}}|_{SP} - \frac{r^{\mathbf{S}}}{2} \left(\overline{\underline{A}}^{\mathbf{S}} + \check{\underline{A}}^{\mathbf{S}}|_{SP} \right) + \text{l.o.t.} \end{aligned}$$

and therefore,

$$\frac{2(1 - \overline{\zeta^{\mathbf{S}}})}{\overline{\zeta^{\mathbf{S}}}} = -D^{\mathbf{S}} + 2\zeta^{\mathbf{S}}|_{SP} - \frac{r^{\mathbf{S}}}{2} \left(\overline{\underline{A}}^{\mathbf{S}} + \check{\underline{A}}^{\mathbf{S}}|_{SP} \right) + \text{l.o.t.}$$

i.e.,

$$\overline{\zeta^{\mathbf{S}}} - 1 = \frac{1}{2}D^{\mathbf{S}} - \zeta^{\mathbf{S}}|_{SP} + \frac{r^{\mathbf{S}}}{4} \left(\overline{\underline{A}}^{\mathbf{S}} + \check{\underline{A}}^{\mathbf{S}}|_{SP} \right) + \text{l.o.t.}$$

where l.o.t. denote higher order terms in $\zeta^{\mathbf{S}}$ and $\overline{\zeta^{\mathbf{S}}} - 1$. Thus,

$$|\overline{\zeta^{\mathbf{S}}} - 1| \lesssim |D^{\mathbf{S}}| + \|\zeta^{\mathbf{S}}\|_{L^\infty(\mathbf{S})} + r^{\mathbf{S}}\|\underline{A}^{\mathbf{S}}\|_{L^\infty(\mathbf{S})}.$$

Hence, back to (9.8.45) we derive,

$$r_{\mathbf{S}}^{-1}\|\zeta^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} + |\overline{\zeta^{\mathbf{S}}} - 1| \lesssim |D^{\mathbf{S}}| + r^{\mathbf{S}}\|\underline{A}^{\mathbf{S}}\|_{L^\infty(\mathbf{S})} + r\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \mathring{\epsilon}.$$

Combining with (9.8.44) we deduce,

$$\begin{aligned} \|\underline{A}^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} &\lesssim \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \mathring{\epsilon}, \\ r_{\mathbf{S}}^{-1}\|\zeta^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} + |\overline{\zeta^{\mathbf{S}}} - 1| &\lesssim r\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \mathring{\epsilon}. \end{aligned} \quad (9.8.46)$$

In view of (9.8.43) we also deduce,

$$\begin{aligned} r_{\mathbf{S}}^{-1}\|\check{\underline{a}}^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} &\lesssim r_{\mathbf{S}}^{-1}\|\zeta^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} + \|\underline{A}^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} \\ &\lesssim \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \mathring{\epsilon}. \end{aligned}$$

From (9.8.41) we further deduce

$$\left| \overline{a}^{\mathbf{S}} + 1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} \right| \lesssim |D^{\mathbf{S}}| + \|\check{\underline{a}}^{\mathbf{S}}\|_{L^\infty(\mathbf{S})} \lesssim |D^{\mathbf{S}}| + \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \mathring{\epsilon}.$$

Hence,

$$r_{\mathbf{S}}^{-1}\|\check{\underline{a}}^{\mathbf{S}}\|_{\mathfrak{h}_{k+1}(\mathbf{S})} + \left| \overline{a}^{\mathbf{S}} + 1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} \right| \lesssim |D^{\mathbf{S}}| + \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \mathring{\epsilon} \quad (9.8.47)$$

as stated. \square

In view of the lemma above and the assumption $|D^{\mathbf{S}}| \lesssim \mathring{\epsilon}^{1/2}$ the estimates (9.8.31) become,

$$\begin{aligned} \|C_1\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon}r_{\mathbf{S}}^{-4} \left(\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \mathring{\epsilon}^{1/2} \right), \\ \|C_2\|_{\mathfrak{h}_{k+3}(\mathbf{S})} &\lesssim \mathring{\epsilon}r_{\mathbf{S}}^{-2} \left(\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_{k+3}(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{k+3}(\mathbf{S})} + \mathring{\epsilon}^{1/2} \right), \\ \|C_3\|_{\mathfrak{h}_k(\mathbf{S})} &\lesssim \mathring{\epsilon}r_{\mathbf{S}}^{-4} \left(\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_k(\mathbf{S})} + \mathring{\epsilon}^{1/2} \right). \end{aligned} \quad (9.8.48)$$

To prove the desired estimate for $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}}$ we make use of (9.8.30) and the following lemma.

Lemma 9.8.9. *The error term*

$$E_k = r_{\mathbf{S}}^{-1} \|\Gamma_g^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})}, \quad k \leq s_{max} - 7,$$

appearing in (9.8.30) verifies the estimate

$$E_k \lesssim r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon}^{1/2} \left(\|(\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}})\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\check{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{k+3}(\mathbf{S})} \right).$$

Proof. Since $\Gamma_g^{\mathbf{S}}$ contains only terms estimated by (9.4.11),

$$\|\Gamma_g^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} \lesssim r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon}$$

$\Gamma_b^{\mathbf{S}}$ contains $\vartheta^{\mathbf{S}}$, which is estimated by (9.4.11), as well as $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}, \check{\omega}^{\mathbf{S}}, \overline{\omega}^{\mathbf{S}} - \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^2}$. Thus, in view of the auxiliary estimates $\|\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{5+k}(\mathbf{S})} \lesssim \overset{\circ}{\epsilon}^{1/2}$ and the fact that the quadratic error terms contain one less derivative of $\check{\omega}^{\mathbf{S}}$, we deduce,

$$\|\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} \lesssim r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon}^{1/2} \left(\|(\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}})\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\check{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{k+3}(\mathbf{S})} + r^{\mathbf{S}} \left| \overline{\omega}^{\mathbf{S}} - \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^2} \right| \right).$$

In view of equation (9.8.33), $\overline{\omega}^{\mathbf{S}} - \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^2} = -\frac{1}{4} \overline{A}^{\mathbf{S}} + \frac{r^{\mathbf{S}}}{2} \left(\overline{\mu}^{\mathbf{S}} - \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^3} - \overline{\eta^{\mathbf{S}} \cdot \eta^{\mathbf{S}}} \right)$,

$$\begin{aligned} \left| \overline{\omega}^{\mathbf{S}} - \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^2} \right| &\lesssim |\overline{A}^{\mathbf{S}}| + r^{\mathbf{S}} \left| \overline{\mu} - \frac{m^{\mathbf{S}}}{(r^{\mathbf{S}})^3} \right| + |\eta^{\mathbf{S}}|^2 \\ &\lesssim r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon} |D^{\mathbf{S}}| + r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon}^{1/2} \left(\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_2(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_2(\mathbf{S})} \right) + \overset{\circ}{\epsilon} r^{-2} \\ &\lesssim r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon}^{1/2} \left(\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_2(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_2(\mathbf{S})} + \overset{\circ}{\epsilon} \right). \end{aligned}$$

Hence,

$$\|\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} \lesssim r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon}^{1/2} \left(\|(\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}})\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\check{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{k+3}(\mathbf{S})} + \overset{\circ}{\epsilon} \right)$$

and,

$$\begin{aligned} E_k &= r_{\mathbf{S}}^{-1} \|\Gamma_g^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\Gamma_b^{\mathbf{S}} \cdot \Gamma_b^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} \\ &\lesssim r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon}^{1/2} \left(\|(\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}})\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\check{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{k+3}(\mathbf{S})} \right) \end{aligned}$$

as stated. □

In view of the lemma and estimates (9.8.48) for C_1, C_2, C_3 the estimates (9.8.30) of Proposition 9.8.6 become,

$$\begin{aligned} \|\mathcal{d}_2^{\mathbf{S},*} \eta^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} &\lesssim r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon}^{1/2} \left(\|\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\underline{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{k+3}(\mathbf{S})} \right), \\ \|\mathcal{d}_2^{\mathbf{S},*} \underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} &\lesssim r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon}^{1/2} \left(\|\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\underline{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{k+3}(\mathbf{S})} \right), \\ \|\mathcal{d}_1^{\mathbf{S},*} \underline{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{2+k}(\mathbf{S})} &\lesssim r_{\mathbf{S}}^{-1} \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + r_{\mathbf{S}}^{-1} \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{2+k}(\mathbf{S})} \\ &\quad + r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-1} \overset{\circ}{\epsilon}^{1/2} \left(\|\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{3+k}(\mathbf{S})} + \|\underline{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{2+k}(\mathbf{S})} \right). \end{aligned} \tag{9.8.49}$$

From the last equation we derive,

$$\|\underline{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{3+k}(\mathbf{S})} \lesssim \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{2+k}(\mathbf{S})} + \overset{\circ}{\epsilon}.$$

Thus the first two equations in (9.8.49) become

$$\begin{aligned} r^{\mathbf{S}} \|\mathcal{d}_2^{\mathbf{S},*} \eta^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} + \overset{\circ}{\epsilon}^{1/2} \left(\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} \right), \\ r^{\mathbf{S}} \|\mathcal{d}_2^{\mathbf{S},*} \underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} + \overset{\circ}{\epsilon}^{1/2} \left(\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} \right), \end{aligned} \tag{9.8.50}$$

from which we deduce,

$$\begin{aligned} \|\eta^{\mathbf{S}}\|_{\mathfrak{h}_{5+k}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-2} |B^{\mathbf{S}}|, \\ \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{5+k}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-2} |\underline{B}^{\mathbf{S}}|, \\ \|\underline{\omega}^{\mathbf{S}}\|_{\mathfrak{h}_{3+k}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon} + r_{\mathbf{S}}^{-2} (|B^{\mathbf{S}}| + |\underline{B}^{\mathbf{S}}|), \end{aligned}$$

as stated. We can then go back to the preliminary estimates obtained above for $\zeta^{\mathbf{S}}, \underline{A}^{\mathbf{S}}$ and $a^{\mathbf{S}}$ to derive the remaining statements (1-4) of Proposition 9.8.7. To prove the last part of the Proposition we make use of the corresponding Ricci and Bianchi equations in the $e_3^{\mathbf{S}}$ direction. \square

Corollary 9.8.10. *Under the same assumptions as in the proposition above we have the more precise estimates, with $d(\mathbf{S}) = \int_{\mathbf{S}} e^{2\Phi}$,*

$$\begin{aligned} \left\| \eta^{\mathbf{S}} - \frac{1}{d(\mathbf{S})} B^{\mathbf{S}} e^{\Phi} \right\|_{\mathfrak{h}_{5+k}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon}, \\ \left\| \underline{\xi}^{\mathbf{S}} - \frac{1}{d(\mathbf{S})} \underline{B}^{\mathbf{S}} e^{\Phi} \right\|_{\mathfrak{h}_{5+k}(\mathbf{S})} &\lesssim \overset{\circ}{\epsilon}. \end{aligned}$$

Note also that,

$$d(\mathbf{S}) = (r^{\mathbf{S}})^4 \left(\frac{8\pi}{3} + O(\overset{\circ}{\epsilon}) \right).$$

Proof. In view of (9.8.50), (9.8.36) and auxiliary assumption (9.8.35) we deduce,

$$\begin{aligned} \left\| \eta^{\mathbf{S}} - \left(\frac{\int_{\mathbf{S}} \eta^{\mathbf{S}} e^{\Phi}}{\int_{\mathbf{S}} e^{2\Phi}} \right) e^{\Phi} \right\|_{\mathfrak{h}_{5+k}(\mathbf{S})} &\lesssim r \|\not{d}_2^{\mathbf{S},*} \eta^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} \\ &\lesssim \overset{\circ}{\epsilon} + \overset{\circ}{\epsilon}^{1/2} (\|\eta^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})} + \|\underline{\xi}^{\mathbf{S}}\|_{\mathfrak{h}_{4+k}(\mathbf{S})}) \\ &\lesssim \overset{\circ}{\epsilon} + \overset{\circ}{\epsilon}^{1/2} \left(\overset{\circ}{\epsilon} + r^{-2} (|\underline{B}^{\mathbf{S}}| + |B^{\mathbf{S}}|) \right) \\ &\lesssim \overset{\circ}{\epsilon}. \end{aligned}$$

We deduce,

$$\left\| \eta^{\mathbf{S}} - B^{\mathbf{S}} \frac{1}{\int_{\mathbf{S}} e^{2\Phi}} e^{\Phi} \right\|_{\mathfrak{h}_{5+k}(\mathbf{S})} \lesssim \overset{\circ}{\epsilon}.$$

Similarly,

$$\left\| \underline{\xi}^{\mathbf{S}} - \underline{B}^{\mathbf{S}} \frac{1}{\int_{\mathbf{S}} e^{2\Phi}} e^{\Phi} \right\|_{\mathfrak{h}_{5+k}(\mathbf{S})} \lesssim \overset{\circ}{\epsilon}$$

as desired. □

9.8.3 Construction of Σ_0

To construct the spacelike hypersurface of Theorem 9.8.1 we proceed as follows.

Step 12. Let $\Psi(s), \Lambda(s), \underline{\Lambda}(s)$ real valued functions that will be carefully chosen later. We look for the hypersurface Σ_0 in the form,

$$\Sigma_0 = \bigcup_{s \geq \overset{\circ}{s}} \mathbf{S}[P(s)] = \bigcup_{s \geq \overset{\circ}{s}} \mathbf{S}[\Psi(s), s, \Lambda(s), \underline{\Lambda}(s)] \quad (9.8.51)$$

where $P(s)$ is a curve in the parameter space P given by,

$$P(s) = (\Psi(s), s, \Lambda(s), \underline{\Lambda}(s)). \quad (9.8.52)$$

In order for Σ_0 to start at $\mathbf{S}_0 = \mathbf{S}[\overset{\circ}{u}, \overset{\circ}{s}, \Lambda_0, \underline{\Lambda}_0]$ we impose the conditions

$$\Psi(\overset{\circ}{s}) = \overset{\circ}{u}, \quad \Lambda(\overset{\circ}{s}) = \Lambda_0, \quad \underline{\Lambda}(\overset{\circ}{s}) = \underline{\Lambda}_0. \quad (9.8.53)$$

Step 13. We expect Σ_0 to be a perturbation of the timelike surface $u + s = c_0$ for some constant c_0 . We thus introduce the notation

$$\psi(s) := \Psi(s) + s - c_0, \text{ so that } \Psi(s) = -s + c_0 + \psi(s)$$

and expect $\psi(s) = O(\overset{\circ}{\delta})$.

Step 14. In view of (9.8.15) we can express the collection of spheres Σ_0 in the form

$$\Sigma_0 = \left\{ \Xi(s, \theta), \quad s \geq \overset{\circ}{s}, \quad \theta \in [0, \pi] \right\} \quad (9.8.54)$$

where the map $\Xi(s, \theta) = \Xi(\Psi(s), s, \theta)$ is defined as

$$\Xi(s, \theta) := \left(\Psi(s) + U(\theta, P(s)), \quad s + S(\theta, P(s)), \quad \theta \right). \quad (9.8.55)$$

At the South Pole, i.e. $\theta = 0$, where $U(0, P) = S(0, P) = 0$

$$\Xi(s, 0) = \left(\Psi(s), \quad s, \quad 0 \right). \quad (9.8.56)$$

Clearly,

$$\begin{aligned} \partial_s \Xi(s, \theta) &= \left(\Psi'(s) + \partial_P U(\theta, P(s)) P'(s), \quad 1 + \partial_P S(\theta, P(s)) P'(s), \quad 0 \right), \\ \partial_\theta \Xi(s, \theta) &= \left(\partial_\theta U(\theta, P(s)), \quad \partial_\theta S(\theta, P(s)), \quad 1 \right), \end{aligned}$$

where,

$$\begin{aligned} \partial_P U(\cdot) P'(s) &= \Psi'(s) \partial_u U(\cdot) + \partial_s U(\cdot) + \Lambda'(s) \partial_\Lambda U(\cdot) + \underline{\Lambda}'(s) \partial_{\underline{\Lambda}} U(\cdot), \\ \partial_P S(\cdot) P'(s) &= \Psi'(s) \partial_u S(\cdot) + \partial_s S(\cdot) + \Lambda'(s) \partial_\Lambda S(\cdot) + \underline{\Lambda}'(s) \partial_{\underline{\Lambda}} S(\cdot). \end{aligned}$$

Given f a function on Σ_* we have,

$$\begin{aligned} \frac{d}{ds} f(\Xi(s, \theta)) &= \left(\Psi'(s) + \partial_P U(\theta, P(s)) P'(s) \right) \partial_u f + \left(1 + \partial_P S(\theta, P(s)) P'(s) \right) \partial_s f \\ &= X_* f, \\ \frac{d}{ds} f(\Xi(s, \theta)) &= \partial_\theta U(\theta, P(s)) \partial_s f + \partial_\theta S(\theta, P(s)) \partial_s + \partial_\theta f \\ &= Y_* f, \end{aligned}$$

where X_*, Y_* are the following tangent vectorfields along Σ_* ,

$$\begin{aligned} X_*(s, \theta) &:= \left(\Psi'(s) + \partial_P U(\theta, P(s)) P'(s) \right) \partial_u + \left(1 + \partial_P S(\theta, P(s)) P'(s) \right) \partial_s, \\ Y_*(s, \theta) &:= \partial_\theta U(\theta, P(s)) \partial_s + \partial_\theta S(\theta, P(s)) \partial_s + \partial_\theta, \end{aligned} \quad (9.8.57)$$

or,

$$\begin{aligned} X_*(s, \theta) &:= \left(\Psi'(s) + \check{A}(s, \theta) \right) \partial_u + \left(1 + \check{B}(s, \theta) P'(s) \right) \partial_s, \\ Y_*(s, \theta) &:= \check{C}(s, \theta) \partial_u + \check{D}(s, \theta) \partial_s + \partial_\theta, \end{aligned} \quad (9.8.58)$$

where,

$$\begin{aligned} \check{A}(s, \theta) &:= \partial_P U(\theta, P(s)) P'(s) \\ &= \partial_u U(\theta, P(s)) \Psi'(s) + \partial_s U(\theta, P(s)) + \partial_\Lambda U(\theta, P(s)) \Lambda'(s) + \partial_{\underline{\Lambda}} U(\theta, P(s)) \underline{\Lambda}'(s), \\ \check{B}(s, \theta) &:= \partial_P S(\theta, P(s)) P'(s) \\ &= \partial_u S(\theta, P(s)) \Psi'(s) + \partial_s S(\theta, P(s)) + \partial_\Lambda U(\theta, P(s)) \Lambda'(s) + \partial_{\underline{\Lambda}} S(\theta, P(s)) \underline{\Lambda}'(s), \\ \check{C}(s, \theta) &:= \partial_\theta U(\theta, P(s)), \\ \check{D}(s, \theta) &:= \partial_\theta S(\theta, P(s)). \end{aligned}$$

Step 15. Define the vectorfield, along the South Pole of each $\mathbf{S} \subset \Sigma_0$,

$$X_* \Big|_{SP} h = \frac{d}{ds} h(\Xi(s, 0)). \quad (9.8.59)$$

Lemma 9.8.11. *At the South Pole we have the relations (recall $\nu^{\mathbf{S}} = e_3^{\mathbf{S}} + a^{\mathbf{S}} e_4^{\mathbf{S}}$)*

$$X_* \Big|_{SP} = \frac{1}{2\lambda} \varsigma \Psi' \nu^{\mathbf{S}} \Big|_{SP}, \quad (9.8.60)$$

$$a^{\mathbf{S}} \Big|_{SP} = \frac{2\lambda^2}{\Psi'(s)\varsigma} \left(1 - \frac{1}{2} \Psi'(s)\varsigma \underline{\Omega} \right) \Big|_{SP}, \quad (9.8.61)$$

or, more precisely,

$$a^{\mathbf{S}}(\Psi(s), s, 0) = \frac{1}{\Psi'(s)} \frac{2\lambda^2}{\varsigma} \left(1 - \frac{1}{2} \Psi'(s)\varsigma \underline{\Omega} \right) (\Psi(s), s, 0).$$

Here $f, \underline{f}, \lambda$ are the transition functions and $\varsigma, \underline{\Omega}$ correspond to the background foliation.

Proof. Note that

$$\check{A}(s, 0) = \check{B}(s, 0) = \check{C}(s, 0) = \check{D}(s, 0) = 0.$$

Thus, at the South Pole SP,

$$X_*(s, 0) = \Psi'(s)\partial_u + \partial_s.$$

Recall that

$$\partial_s = e_4, \quad \partial_u = \frac{1}{2}\varsigma(e_3 - \underline{\Omega}e_4 - \underline{b}\gamma^{1/2}e_\theta), \quad \partial_\theta = \sqrt{\gamma}e_\theta,$$

or, since \underline{b} vanishes at the South Pole,

$$X_*(s, 0) = \Psi' \frac{1}{2}\varsigma(e_3 - \underline{\Omega}e_4) + e_4 = (1 - \frac{1}{2}\Psi'(s)\varsigma\underline{\Omega})e_4 + \frac{1}{2}\Psi'(s)\varsigma e_3.$$

On the other hand, since the transition functions f, \underline{f} vanish at the South Pole,

$$e_4^{\mathbf{S}} = \lambda e_4, \quad e_3^{\mathbf{S}} = \lambda^{-1}e_3.$$

Hence,

$$\begin{aligned} X_*(s, 0) &= \lambda(1 - \frac{1}{2}\Psi'(s)\varsigma\underline{\Omega})e_4^{\mathbf{S}} + \frac{1}{2}\lambda^{-1}\Psi'(s)\varsigma e_3^{\mathbf{S}} \\ &= \frac{1}{2}\lambda^{-1}\Psi'(s)\varsigma \left(e_3^{\mathbf{S}} + \frac{2\lambda^2}{\Psi'(s)\varsigma} (1 - \frac{1}{2}\Psi'(s)\varsigma\underline{\Omega})e_4^{\mathbf{S}} \right). \end{aligned}$$

In view of the definition of $\nu^{\mathbf{S}}$ we deduce,

$$X_*(s, 0) = \frac{1}{2}\lambda^{-1}\Psi'(s)\varsigma \nu^{\mathbf{S}}|_{SP}$$

and³⁹,

$$a^{\mathbf{S}}(s, 0) = \frac{2\lambda^2}{\Psi'(s)\varsigma} (1 - \frac{1}{2}\Psi'(s)\varsigma\underline{\Omega})$$

as stated. □

Step 16. The transition functions $(f, \underline{f}, \lambda)$ are uniquely determined on \mathbf{S} by the results of Theorem 9.4.2 in terms of $\Lambda, \underline{\Lambda}$. The same holds true for all curvature components and the Ricci coefficients $\kappa^{\mathbf{S}}, \vartheta^{\mathbf{S}}, \zeta^{\mathbf{S}}, \underline{\kappa}^{\mathbf{S}}, \underline{\vartheta}^{\mathbf{S}}$. One can easily see from the transformation

³⁹Note that $a^{\mathbf{S}}(s, 0) = a^{\mathbf{S}}(\Xi(s, 0))$.

formulas that the values of the $e_3^{\mathbf{S}}$ derivatives of $(f, \underline{f}, \lambda)$ are determined by the transversal Ricci coefficients $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}, \underline{\eta}^{\mathbf{S}}$. Indeed, schematically, from the transformation formulas for $\eta, \underline{\xi}, \underline{\omega}$ in Proposition 9.3.1,

$$\begin{aligned} e_3^{\mathbf{S}} f &= 2(\eta^{\mathbf{S}} - \eta) - \frac{1}{2} \kappa \underline{f} + f \underline{\omega} + F \cdot \Gamma_b + \text{l.o.t.}, \\ e_3^{\mathbf{S}} \underline{f} &= 2(\underline{\xi}^{\mathbf{S}} - \underline{\xi}) - \frac{1}{2} \underline{f} (\underline{\kappa} + 4\underline{\omega}) + F \cdot \Gamma_b + \text{l.o.t.}, \\ e_3^{\mathbf{S}}(\log \lambda) &= 2(\underline{\omega}^{\mathbf{S}} - \underline{\omega}) + \Gamma_b \cdot F + \text{l.o.t.}, \end{aligned} \tag{9.8.62}$$

where $F = (f, \underline{f}, \log \lambda)$ and l.o.t. denotes terms which are linear in Γ_g, Γ_b and linear and higher order in F . Recall also that the $e_4^{\mathbf{S}}$ derivatives of F are fixed by our transversality condition (9.4.12) More precisely we have,

$$\begin{aligned} e_4^{\mathbf{S}}(f) &= -\frac{1}{2} \kappa f + \text{l.o.t.}, \\ e_4^{\mathbf{S}}(\underline{f}) &= 2e_\theta^{\mathbf{S}}(\log \lambda) - \underline{f} \kappa + 2(\underline{\omega} + \frac{1}{4} \underline{\kappa}) f + \text{l.o.t.}, \\ e_4^{\mathbf{S}}(\log \lambda) &= \text{l.o.t.} \end{aligned}$$

It follows that $\eta^{\mathbf{S}}, \underline{\xi}^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}}$ can be determined by $\nu^{\mathbf{S}}(f, \underline{f}, \lambda)$ and the scalar $a^{\mathbf{S}}$. More precisely,

$$\begin{aligned} \nu^{\mathbf{S}}(f) &= 2(\eta^{\mathbf{S}} - \eta) - \frac{1}{2} (\kappa \underline{f} + a^{\mathbf{S}} \underline{\kappa} f) + f \underline{\omega} + F \cdot \Gamma_b + \text{l.o.t.}, \\ \nu^{\mathbf{S}}(\underline{f}) &= 2(\underline{\xi}^{\mathbf{S}} - \underline{\xi}) - \frac{1}{2} (\underline{\kappa} + 4\underline{\omega})(\underline{f} - a^{\mathbf{S}} f) + a^{\mathbf{S}} (2e_\theta^{\mathbf{S}}(\log \lambda) - \underline{f} \kappa) + F \cdot \Gamma_b + \text{l.o.t.} \end{aligned} \tag{9.8.63}$$

Step 17. We derive equations for $\Lambda(s) = \Lambda(\Psi(s), s, 0)$, $\underline{\Lambda}(s) = \underline{\Lambda}(\Psi(s), s, 0)$ as follows.

Lemma 9.8.12. *We have the following identities*

$$\begin{aligned} c(s) \frac{1}{\Psi'(s)} \Lambda'(s) &= \int_{\mathbf{S}} \nu^{\mathbf{S}}(f) e^\Phi - \frac{6}{r^{\mathbf{S}}} \Lambda(s) + E(s), \\ c(s) \frac{1}{\Psi'(s)} \underline{\Lambda}'(s) &= \int_{\mathbf{S}(s)} \nu^{\mathbf{S}}(\underline{f}) e^\Phi - \frac{6}{r^{\mathbf{S}}} \underline{\Lambda}(s) + \underline{E}(s), \end{aligned} \tag{9.8.64}$$

where,

$$c(s) = \left(\frac{2\lambda}{\varsigma} \right) \Big|_{SP}(s) = \left(\frac{2\lambda}{\varsigma} \right) (\Psi(s), s, 0)$$

and error terms,

$$\begin{aligned}
E(s) &= \frac{1}{2} \int_{\mathbf{S}(s)} \left(3\underline{\kappa}^{\mathbf{S}} - \underline{\vartheta}^{\mathbf{S}} - a^{\mathbf{S}}\vartheta^{\mathbf{S}} + \frac{3}{r^{\mathbf{S}}}(a^{\mathbf{S}} + (1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}})) \right) f e^{\Phi} \\
&+ (\zeta^{\mathbf{S}})^{-1} \Big|_{SP} \int_{\mathbf{S}(s)} \left(\zeta^{\mathbf{S}} - \zeta^{\mathbf{S}} \Big|_{SP} \right) \left(\nu^{\mathbf{S}}(f) - \frac{6}{r^{\mathbf{S}}}\Lambda(s) \right) e^{\Phi} + l.o.t., \\
\underline{E}(s) &= \frac{1}{2} \int_{\mathbf{S}(s)} \left(3\underline{\kappa}^{\mathbf{S}} - \underline{\vartheta}^{\mathbf{S}} - a^{\mathbf{S}}\vartheta^{\mathbf{S}} + \frac{3}{r^{\mathbf{S}}}(a^{\mathbf{S}} + (1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}})) \right) \underline{f} e^{\Phi} \\
&+ (\zeta^{\mathbf{S}})^{-1} \Big|_{SP} \int_{\mathbf{S}(s)} \left(\zeta^{\mathbf{S}} - \zeta^{\mathbf{S}} \Big|_{SP} \right) \left(\nu^{\mathbf{S}}(\underline{f}) - \frac{6}{r^{\mathbf{S}}}\underline{\Lambda}(s) \right) e^{\Phi} + l.o.t.
\end{aligned}$$

Proof. According to Lemma 9.8.3 we have

$$\nu^{\mathbf{S}} \left(\int_{\mathbf{S}} h \right) = (\zeta^{\mathbf{S}})^{-1} \int_{\mathbf{S}} \zeta^{\mathbf{S}} (\nu^{\mathbf{S}}(h) + (\underline{\kappa}^{\mathbf{S}} + a^{\mathbf{S}}\kappa^{\mathbf{S}})h).$$

Thus, applying the vectorfield $\nu^{\mathbf{S}}|_{SP} = \frac{2\lambda}{\zeta^{\mathbf{S}}} X_*|_{SP}$ to the formulas (9.8.14),

$$\begin{aligned}
\frac{1}{\Psi'(s)} \left(\frac{2\lambda}{\zeta} \right) \Big|_{SP} \frac{d}{ds} \Lambda(s) &= \nu^{\mathbf{S}} \Big|_{SP} (\Lambda) = \nu^{\mathbf{S}}(\Lambda) \Big|_{SP} = \nu^{\mathbf{S}} \left(\int_{\mathbf{S}} f e^{\Phi} \right) \Big|_{SP} \\
&= (\zeta^{\mathbf{S}})^{-1} \Big|_{SP} \int_{\mathbf{S}(s)} \zeta^{\mathbf{S}} \left(\nu^{\mathbf{S}}(f e^{\Phi}) + (\underline{\kappa}^{\mathbf{S}} + a^{\mathbf{S}}\kappa^{\mathbf{S}}) f e^{\Phi} \right).
\end{aligned}$$

Introducing

$$J(f) = e^{-\Phi} \nu^{\mathbf{S}}(f e^{\Phi}) + (\underline{\kappa}^{\mathbf{S}} + a^{\mathbf{S}}\kappa^{\mathbf{S}}) f \quad (9.8.65)$$

we deduce,

$$\begin{aligned}
c(s) \frac{1}{\Psi'(s)} &= (\zeta^{\mathbf{S}})^{-1} \Big|_{SP} \int_{\mathbf{S}(s)} \zeta^{\mathbf{S}} J(f) e^{\Phi} \\
&= \int_{\mathbf{S}(s)} J(f) e^{\Phi} + (\zeta^{\mathbf{S}})^{-1} \Big|_{SP} \int_{\mathbf{S}(s)} \left(\zeta^{\mathbf{S}} - \zeta^{\mathbf{S}} \Big|_{SP} \right) J(f).
\end{aligned}$$

On the other hand, since $e_3\Phi = \frac{1}{2}(\underline{\kappa} - \underline{\vartheta})$, $e_4\Phi = \frac{1}{2}(\kappa - \vartheta)$

$$\begin{aligned}
J(f) &= \nu^{\mathbf{S}}(f) + (e_3^{\mathbf{S}}\Phi + a^{\mathbf{S}}e_4^{\mathbf{S}}\Phi + \underline{\kappa}^{\mathbf{S}} + a^{\mathbf{S}}\kappa^{\mathbf{S}}) f \\
&= \nu^{\mathbf{S}}(f) + \frac{1}{2} (3\underline{\kappa}^{\mathbf{S}} - \underline{\vartheta}^{\mathbf{S}} + a^{\mathbf{S}}(3\kappa^{\mathbf{S}} - \vartheta^{\mathbf{S}})) f \\
&= \nu^{\mathbf{S}}(f) + \frac{3}{2} (\underline{\kappa}^{\mathbf{S}} + a^{\mathbf{S}}\kappa^{\mathbf{S}}) - \frac{1}{2} (\underline{\vartheta}^{\mathbf{S}} + a^{\mathbf{S}}\vartheta^{\mathbf{S}}).
\end{aligned}$$

Since $\kappa^{\mathbf{S}} = \frac{2}{r^{\mathbf{S}}}$ and $\underline{\kappa}^{\mathbf{S}} = \overline{\underline{\kappa}^{\mathbf{S}}} + \check{\kappa}^{\mathbf{S}} = -\frac{2\Upsilon^{\mathbf{S}}}{r^{\mathbf{S}}} + \check{\kappa}^{\mathbf{S}}$ we deduce,

$$\begin{aligned} J(f) &= \nu^{\mathbf{S}}(f) + \frac{3}{r^{\mathbf{S}}} (-\Upsilon^{\mathbf{S}} + a^{\mathbf{S}}) f + \frac{1}{2} (3\check{\kappa}^{\mathbf{S}} - \underline{\vartheta}^{\mathbf{S}} - a^{\mathbf{S}}\vartheta^{\mathbf{S}}) f \\ &= \nu^{\mathbf{S}}(f) + \frac{3}{r^{\mathbf{S}}} \left(-\Upsilon^{\mathbf{S}} - \left(1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}\right) \right) f + \frac{1}{2} \left(3\check{\kappa}^{\mathbf{S}} - \underline{\vartheta}^{\mathbf{S}} - a^{\mathbf{S}}\vartheta^{\mathbf{S}} + \frac{3}{r^{\mathbf{S}}} \left(a^{\mathbf{S}} + \left(1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}\right) \right) \right) f \\ &= \nu^{\mathbf{S}}(f) - \frac{6}{r^{\mathbf{S}}} \Lambda(s) + \frac{1}{2} \left(3\check{\kappa}^{\mathbf{S}} - \underline{\vartheta}^{\mathbf{S}} - a^{\mathbf{S}}\vartheta^{\mathbf{S}} + \frac{3}{r^{\mathbf{S}}} \left(a^{\mathbf{S}} + \left(1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}\right) \right) \right) f. \end{aligned}$$

We deduce,

$$c(s) \frac{1}{\Psi'(s)} = \int_{\mathbf{S}} \nu^{\mathbf{S}}(f) e^{\Phi} - \frac{6}{r^{\mathbf{S}}} \Lambda(s) + E(s)$$

where,

$$\begin{aligned} E(s) &= \frac{1}{2} \int_{\mathbf{S}(s)} \left(3\check{\kappa}^{\mathbf{S}} - \underline{\vartheta}^{\mathbf{S}} - a^{\mathbf{S}}\vartheta^{\mathbf{S}} + \frac{3}{r^{\mathbf{S}}} \left(a^{\mathbf{S}} + \left(1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}\right) \right) \right) f e^{\Phi} \\ &\quad + (\varsigma^{\mathbf{S}})^{-1} \Big|_{SP} \int_{\mathbf{S}(s)} \left(\varsigma^{\mathbf{S}} - \varsigma^{\mathbf{S}} \Big|_{SP} \right) J(f) \\ &= \frac{1}{2} \int_{\mathbf{S}(s)} \left(3\check{\kappa}^{\mathbf{S}} - \underline{\vartheta}^{\mathbf{S}} - a^{\mathbf{S}}\vartheta^{\mathbf{S}} + \frac{3}{r^{\mathbf{S}}} \left(a^{\mathbf{S}} + \left(1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}\right) \right) \right) f e^{\Phi} \\ &\quad + (\varsigma^{\mathbf{S}})^{-1} \Big|_{SP} \int_{\mathbf{S}(s)} \left(\varsigma^{\mathbf{S}} - \varsigma^{\mathbf{S}} \Big|_{SP} \right) \left(\nu^{\mathbf{S}}(f) - \frac{6}{r^{\mathbf{S}}} \Lambda(s) \right) e^{\Phi} + \text{l.o.t.} \end{aligned}$$

The proof for $\underline{\Lambda}$ is exactly the same. □

Step 18. We make use of the estimates for $F = (f, \underline{f}, \log \lambda)$ and $e_4^{\mathbf{S}}(F)$ derived in Theorem 9.4.2 as well as the estimates for $a^{\mathbf{S}}, \varsigma^{\mathbf{S}}, \eta^{\mathbf{S}}, \xi^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}}$ derived in Proposition 9.8.7 to evaluate the right hand sides of (9.8.64). Recall that in Proposition 9.8.7 we have made the auxiliary assumption (9.8.35) i.e.

$$r_{\mathbf{S}}^{-2} (|B^{\mathbf{S}}| + |\underline{B}^{\mathbf{S}}|) + |D^{\mathbf{S}}| \leq \epsilon^{1/2}.$$

Proposition 9.8.13. *The following equations hold true for the functions⁴⁰*

$$\begin{aligned} \Lambda(s) &= \Lambda(\Psi(s), s, 0), & \underline{\Lambda}(s) &= \underline{\Lambda}(\Psi(s), s, 0), \\ B(s) &= \Lambda(\Psi(s), s, 0), & \underline{B}(s) &= \Lambda(\Psi(s), s, 0), & r(s) &= r(\Psi(s), s, 0), \end{aligned}$$

⁴⁰Note also that $r^{\mathbf{S}(s)} = r^{\mathbf{S}(s)} = r|_{SP(\mathbf{S}(s))} = r(s)$.

$$\begin{aligned} \frac{1}{-1 + \psi'(s)} \Lambda'(s) &= B(s) - \frac{1}{2} r(s)^{-1} \underline{\Lambda}(s) - \frac{7}{2} r(s)^{-1} \Lambda(s) + O(r^{-1}) \Lambda(s) \\ &\quad + N(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi)(s), \\ \frac{1}{-1 + \psi'(s)} \underline{\Lambda}'(s) &= \underline{B}(s) - \frac{7}{2} r(s)^{-1} \underline{\Lambda}(s) + \frac{1}{2} r(s)^{-1} \Lambda(s) + O(r^{-1}) (\Lambda(s) + \underline{\Lambda}(s)) \\ &\quad + \underline{N}(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi)(s). \end{aligned} \tag{9.8.66}$$

The expressions N, \underline{N} verify the following properties.

- They depend on $B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi, F = (f, \underline{f}, \lambda - 1)$, the background Ricci coefficients Γ_b, Γ_g and curvature $\check{R} = \{\alpha, \beta, \check{\rho}, \underline{\beta}, \underline{\alpha}\}$.
- N, \underline{N} vanish at $(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi) = (0, 0, 0, 0, 0, 0)$. In fact,

$$|N, \underline{N}| \lesssim r^2 \overset{\circ}{\delta}.$$

- The linear part in B, \underline{B}, D has $O(\overset{\circ}{\epsilon})$ coefficients, i.e. coefficients which depend on the quantities $\Gamma_b, \Gamma_g, \check{R}, F$ and $\Lambda, \underline{\Lambda}, \psi$.
- The linear part in $\Lambda, \underline{\Lambda}, \psi$ has $O(\overset{\circ}{\epsilon})$ coefficients.

Proof. To prove the desired result we make use of (9.8.63) to check the following,

$$\begin{aligned} \int_{\mathbf{S}(s)} \nu^{\mathbf{S}}(f) e^{\Phi} &= 2B(s) - r^{-1} \underline{\Lambda}(s) - r^{-1} \Lambda(s) + O(r^{-1}) \Lambda(s) + O(r^2 \overset{\circ}{\delta}), \\ \int_{\mathbf{S}(s)} \nu^{\mathbf{S}}(\underline{f}) e^{\Phi} &= 2\underline{B}(s) - r^{-1} \underline{\Lambda}(s) + r^{-1} \Lambda(s) + O(r^{-1}) (\Lambda(s) + \underline{\Lambda}(s)) + O(r^2 \overset{\circ}{\delta}). \end{aligned} \tag{9.8.67}$$

Combining this with (9.8.64),

$$\begin{aligned} c(s) \frac{1}{\Psi'(s)} \Lambda'(s) &= \int_{\mathbf{S}} \nu^{\mathbf{S}}(f) e^{\Phi} - \frac{6}{r\mathbf{S}} \Lambda(s) + E(s), \\ c(s) \frac{1}{\Psi'(s)} \underline{\Lambda}'(s) &= \int_{\mathbf{S}(s)} \nu^{\mathbf{S}}(\underline{f}) e^{\Phi} - \frac{6}{r\mathbf{S}} \underline{\Lambda}(s) + \underline{E}(s), \end{aligned}$$

and the following estimates for the error terms E, \underline{E} ,

$$|E(s)| + |\underline{E}(s)| \lesssim r^2 \overset{\circ}{\delta}, \tag{9.8.68}$$

we deduce,

$$\begin{aligned}\frac{1}{\Psi'(s)}\Lambda'(s) &= \frac{1}{c(s)} \left(2B(s) - r^{-1}\underline{\Lambda}(s) - 7r^{-1}\Lambda(s) + O(r^{-1})\Lambda(s) + O(r^2\overset{\circ}{\delta}) \right), \\ \frac{1}{\Psi'(s)}\underline{\Lambda}'(s) &= \frac{1}{c(s)} \left(2\underline{B}(s) - 7r^{-1}\underline{\Lambda}(s) + r^{-1}\Lambda(s) + O(r^{-1})(\Lambda(s) + \underline{\Lambda}(s)) + O(r^2\overset{\circ}{\delta}) \right).\end{aligned}$$

According to our assumptions $\varsigma = 1 + O(\overset{\circ}{\epsilon})$. Also according to Theorem 9.4.2 $\lambda = 1 + O(r^{-1}\overset{\circ}{\epsilon})$. Thus,

$$c(s) = \left(\frac{2\lambda}{\varsigma} \right) \Big|_{SP}(s) = \frac{2(1 + Or^{-1}(\overset{\circ}{\epsilon}))}{1 + O(\overset{\circ}{\epsilon})} = 2 + O(\overset{\circ}{\epsilon}).$$

Hence,

$$\begin{aligned}\frac{1}{\Psi'(s)}\Lambda'(s) &= \frac{1}{2} \left(2B(s) - r^{-1}\underline{\Lambda}(s) - 7r^{-1}\Lambda(s) + O(r^{-1})\Lambda(s) + O(r^2\overset{\circ}{\delta}) \right) \\ &= B(s) - \frac{1}{2}r^{-1}\underline{\Lambda}(s) - \frac{7}{2}r^{-1}\Lambda(s) + O(r^{-1})\Lambda(s) + O(r^2\overset{\circ}{\delta}).\end{aligned}$$

Setting $\Psi(s) = -s + \psi(s) + c_0$ and recalling the structure of the error terms we have denoted by $O(r^2\overset{\circ}{\delta})$

$$\frac{1}{-1 + \psi'(s)}\Lambda'(s) = B(s) - \frac{1}{2}r^{-1}\underline{\Lambda}(s) - \frac{7}{2}r^{-1}\Lambda(s) + O(r^{-1})\Lambda(s) + N(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi)(s)$$

where N verifies the properties mentioned in the proposition. In the same manner we derive

$$\begin{aligned}\frac{1}{-1 + \psi'(s)}\underline{\Lambda}'(s) &= \underline{B}(s) - \frac{7}{2}r^{-1}\underline{\Lambda}(s) + \frac{1}{2}r^{-1}\Lambda(s) + O(r^{-1})(\Lambda(s) + \underline{\Lambda}(s)) \\ &\quad + \underline{N}(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi)(s)\end{aligned}$$

as stated in the proposition.

It remains to check (9.8.67) and (9.8.68) According to (9.8.63) and our assumptions on the Ricci coefficients $\kappa, \underline{\kappa}, \underline{\omega}$, we have along the sphere \mathbf{S}

$$\begin{aligned}\nu^{\mathbf{S}}(f) &= 2(\eta^{\mathbf{S}} - \eta) - \frac{1}{2} \left(\frac{2}{r}\underline{f} - a^{\mathbf{S}}\frac{2\Upsilon}{r}f \right) + f\frac{m}{r^2} + F \cdot \Gamma_b + \text{l.o.t.} \\ &= 2(\eta^{\mathbf{S}} - \eta) - r^{-1}\underline{f} + r^{-1} \left(\frac{m}{r} + a^{\mathbf{S}}(1 - \frac{2m}{r}) \right) f + F \cdot \Gamma.\end{aligned}$$

According to (9.8.40) and auxiliary assumption (9.8.35)

$$\left| a^{\mathbf{S}} + \left(1 + \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} \right) \right| \lesssim \overset{\circ}{\epsilon} + r^{-2} (|\underline{B}^{\mathbf{S}}| + |B^{\mathbf{S}}|) + |D^{\mathbf{S}}| \lesssim \overset{\circ}{\epsilon}^{1/2}.$$

Thus,

$$\nu^{\mathbf{S}}(f) = 2(\eta^{\mathbf{S}} - \eta) - r^{-1}\underline{f} - r^{-1} \left(1 - \frac{m}{r} - \frac{m^2}{r^2} \right) f + r^{-2} O \left(\overset{\circ}{\delta} \overset{\circ}{\epsilon}^{1/2} \right).$$

Since r and $r^{\mathbf{S}}$ are comparable along \mathbf{S} , i.e. $|r - r^{\mathbf{S}}| \leq \overset{\circ}{\delta}$, we deduce, recalling the definition of B ,

$$\int_{\mathbf{S}(s)} \nu^{\mathbf{S}}(f) e^{\Phi} = 2B(s) - 2 \int_{\mathbf{S}(s)} \eta e^{\Phi} - r^{-1} \underline{\Lambda}(s) - r^{-1} \left(1 - \frac{m}{r} - \frac{m^2}{r^2} \right) \Lambda(s) + r O \left(\overset{\circ}{\delta} \overset{\circ}{\epsilon}^{1/2} \right).$$

Making use of the assumption (9.8.1) for η as well as Corollary 9.2.5 we easily deduce,

$$\left| \int_{\mathbf{S}(s)} \eta e^{\Phi} \right| \lesssim r^2 \overset{\circ}{\delta}. \quad (9.8.69)$$

Hence,

$$\begin{aligned} \int_{\mathbf{S}(s)} \nu^{\mathbf{S}}(f) e^{\Phi} &= 2B(s) - r^{-1} \underline{\Lambda}(s) - r^{-1} (1 + O(r^{-1})) \Lambda(s) + O(r^2 \overset{\circ}{\delta}) \\ &= 2B(s) - r^{-1} \underline{\Lambda}(s) - r^{-1} \Lambda(s) + O(r^{-1}) \Lambda(s) + O(r^2 \overset{\circ}{\delta}). \end{aligned}$$

Similarly, starting with,

$$\nu^{\mathbf{S}}(\underline{f}) = 2(\underline{\xi}^{\mathbf{S}} - \underline{\xi}) - \frac{1}{2}(\underline{\kappa} + 4\underline{\omega})(\underline{f} - a^{\mathbf{S}}f) + a^{\mathbf{S}}(2e_{\theta}^{\mathbf{S}}(\log \lambda) - \underline{f}\kappa) + F \cdot \Gamma_b + \text{l.o.t.}$$

we deduce,

$$\begin{aligned} \int_{\mathbf{S}(s)} \nu^{\mathbf{S}}(\underline{f}) e^{\Phi} &= 2\underline{B}(s) - 2 \int_{\mathbf{S}(s)} \underline{\xi} e^{\Phi} + r^{-1} \left(1 + \frac{8m}{r} \right) \underline{\Lambda}(s) + r^{-1} \left(1 - \frac{2m}{r} - \frac{8m^2}{r^2} \right) \Lambda(s) \\ &\quad - 2 \left(1 + \frac{2m}{r} \right) \int_{\mathbf{S}(s)} e_{\theta}^{\mathbf{S}}(\log \lambda) e^{\Phi} + r O \left(\overset{\circ}{\delta} \overset{\circ}{\epsilon}^{1/2} \right). \end{aligned}$$

Making use of the assumption (9.8.1) for $\underline{\xi}$, as well as Corollary 9.2.5,

$$\left| \int_{\mathbf{S}(s)} \underline{\xi} e^{\Phi} \right| \lesssim r^2 \overset{\circ}{\delta}. \quad (9.8.70)$$

Also, in view of the estimates of Theorem 9.4.2,

$$\left| \int_{\mathbf{S}(s)} e_{\theta}^{\mathbf{S}}(\log \lambda) e^{\Phi} \right| \lesssim r^2 \delta^{\circ}.$$

We deduce,

$$\int_{\mathbf{S}(s)} \nu^{\mathbf{S}}(\underline{f}) e^{\Phi} = 2\underline{B}(s) - r^{-1}(1 + O(r^{-1}))\underline{\Lambda}(s) + r^{-1}(1 + O(r^{-1}))\Lambda(s) + O(r^2 \delta^{\circ})$$

as stated. The estimates for E, \underline{E} in (9.8.68) can also be easily checked. This ends the proof of Proposition 9.8.13. \square

Step 19. We derive an equation for ψ . The main result is stated in the proposition below.

Proposition 9.8.14. *The function $\psi(s) = \Psi(s) + s - c_0$ defined in Step 13 verifies the following equation*

$$\psi'(s) = -\frac{1}{2}D(s) + O(D(s)^2) + M(s) \quad (9.8.71)$$

where $M(s)$ is a function which depends only on Γ, R of the background foliation, ψ and $(f, \underline{f}, \lambda - 1)$ such that,

$$|M(s)| \lesssim \delta^{\circ} r(s)^{-1}.$$

Proof. In view of (9.8.61) and the definition of $c(s) = \left(\frac{2\lambda}{\varsigma}\right)\Big|_{SP}(s)$ we have,

$$\Psi'(s) = \frac{1}{a^{\mathbf{S}}\Big|_{SP}(s)} \cdot \frac{2\lambda^2}{\varsigma} \left(1 - \frac{1}{2}\Psi'_{\varsigma\underline{\Omega}}\right)\Big|_{SP}(s)$$

or

$$\Psi'(s) = \left[\frac{2\lambda^2}{\varsigma} \frac{1}{a^{\mathbf{S}} + \lambda^2 \underline{\Omega}} \right]\Big|_{SP}. \quad (9.8.72)$$

Now, we have

$$\begin{aligned} \frac{2\lambda^2}{\varsigma} \frac{1}{a^{\mathbf{S}} + \lambda^2 \underline{\Omega}} &= \frac{2}{\varsigma} \frac{1}{a^{\mathbf{S}} + \underline{\Omega}} + O(\lambda - 1) \\ &= -1 + \frac{a^{\mathbf{S}} + \frac{2}{\varsigma} + \underline{\Omega}}{a^{\mathbf{S}} + \underline{\Omega}} + O(\lambda - 1). \end{aligned}$$

Hence,

$$\psi'(s) = \Psi'(s) + 1 = \left[\frac{a^{\mathbf{S}} + \frac{2}{\underline{\zeta}} + \underline{\Omega}}{a^{\mathbf{S}} + \underline{\Omega}} \right] \Big|_{SP} + O(\lambda - 1) = \left[\frac{a^{\mathbf{S}} + \frac{2}{\underline{\zeta}} + \underline{\Omega}}{a^{\mathbf{S}} + \underline{\Omega}} \right] \Big|_{SP} + O(r^{-1}\overset{\circ}{\delta}).$$

We have, see (9.8.34),

$$a^{\mathbf{S}} \Big|_{SP}(s) = D(s) - 1 - \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}}.$$

Hence,

$$\begin{aligned} (a^{\mathbf{S}} + \underline{\Omega}) \Big|_{SP}(s) &= D(s) - 1 - \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} + \underline{\Omega} \Big|_{SP}(s) = D(s) - 1 - \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} - \left(1 - \frac{2m}{r}\right) + O(\overset{\circ}{\epsilon}) \\ &= D(s) - 2 + O(\overset{\circ}{\epsilon}). \end{aligned}$$

In view of the assumption (9.8.2)

$$\left| \left(\frac{2}{\underline{\zeta}} + \underline{\Omega} \right) \Big|_{SP} - 1 - \frac{2m}{r} \right| \lesssim r^{-1}\overset{\circ}{\delta}$$

we deduce

$$\begin{aligned} \left(a^{\mathbf{S}} + \frac{2}{\underline{\zeta}} + \underline{\Omega} \right) \Big|_{SP}(s) &= a^{\mathbf{S}} \Big|_{SP} + 1 + \frac{2m}{r} + O(r^{-1}\overset{\circ}{\delta}) \\ &= D(s) + \frac{2m}{r} - \frac{2m^{\mathbf{S}}}{r^{\mathbf{S}}} + O(r^{-1}\overset{\circ}{\delta}) \\ &= D(s) + O(r^{-1}\overset{\circ}{\delta}). \end{aligned}$$

Hence,

$$\psi'(s) = \left[\frac{a^{\mathbf{S}} + \frac{2}{\underline{\zeta}} + \underline{\Omega}}{a^{\mathbf{S}} + \underline{\Omega}} \right] \Big|_{SP} + O(r^{-1}\overset{\circ}{\delta}) = -\frac{1}{2}D(s) + O(D(s)^2) + O(r^{-1}\overset{\circ}{\delta})$$

as stated. □

Step 20. We combine Propositions 9.8.13 and 9.8.14 to derive the closed system of equations in $\Lambda, \underline{\Lambda}, \psi$,

$$\begin{aligned} \frac{1}{-1 + \psi'(s)} \Lambda'(s) &= B(s) - \frac{1}{2}r(s)^{-1}\underline{\Lambda}(s) - \frac{7}{2}r(s)^{-1}\Lambda(s) + O(r^{-1})\Lambda(s) \\ &\quad + N(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi)(s), \\ \frac{1}{-1 + \psi'(s)} \underline{\Lambda}'(s) &= \underline{B}(s) - \frac{7}{2}r(s)^{-1}\underline{\Lambda}(s) + \frac{1}{2}r(s)^{-1}\Lambda(s) + O(r^{-1})(\Lambda(s) + \underline{\Lambda}(s)) \\ &\quad + \underline{N}(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi)(s), \\ \psi'(s) &= -\frac{1}{2}D(s) + O(D(s)^2) + M(s), \end{aligned} \tag{9.8.73}$$

with initial conditions

$$\psi(\overset{\circ}{s}) = 0, \quad \Lambda(\overset{\circ}{s}) = \Lambda_0, \quad \underline{\Lambda}(\overset{\circ}{s}) = \underline{\Lambda}_0. \quad (9.8.74)$$

Recall also that $r(s)$ is a smooth function of $\psi(s)$.

The system (9.8.73) is verified for all choices of $(\Lambda, \underline{\Lambda}, \Psi)$. We now make a suitable particular choice for $(\Lambda, \underline{\Lambda}, \Psi)$ as follows.

Consider in particular the system obtained from (9.8.73) by setting B, \underline{B}, D to zero

$$\begin{aligned} \psi'(s) &= M(s), \\ \frac{1}{-1 + \psi'(s)} \Lambda'(s) &= -\frac{1}{2} r(s)^{-1} \underline{\Lambda}(s) - \frac{7}{2} r(s)^{-1} \Lambda(s) + O(r^{-1}) \Lambda(s) + \tilde{N}(\Lambda, \underline{\Lambda}, \psi)(s), \\ \frac{1}{-1 + \psi'(s)} \underline{\Lambda}'(s) &= -\frac{7}{2} r(s)^{-1} \underline{\Lambda}(s) + \frac{1}{2} r(s)^{-1} \Lambda(s) + O(r^{-1}) (\Lambda(s) + \underline{\Lambda}(s)) + \underline{\tilde{N}}(\Lambda, \underline{\Lambda}, \psi)(s), \end{aligned} \quad (9.8.75)$$

where,

$$\begin{aligned} \tilde{N}(\Lambda, \underline{\Lambda}, \psi) &= N(0, 0, 0, \Lambda, \underline{\Lambda}, \psi), \\ \underline{\tilde{N}}(\Lambda, \underline{\Lambda}, \psi) &= \underline{N}(0, 0, 0, \Lambda, \underline{\Lambda}, \psi). \end{aligned}$$

We initialize the system at $s = \overset{\circ}{s}$ as in (9.8.74), i.e.,

$$\Lambda(\overset{\circ}{s}) = \psi(\overset{\circ}{s}) = 0, \quad \Lambda_0, \quad \underline{\Lambda}(\overset{\circ}{s}) = \underline{\Lambda}_0.$$

The system admits a unique solution $\psi(s)$ defined in a small neighborhood $\overset{\circ}{I}$ of $\overset{\circ}{s}$. The function $\Psi(s) = -s + \psi(s) + c_0$ defines the desired hypersurface Σ_0 .

Step 21. It remains to show that the function B, \underline{B}, D vanish on the hypersurface Σ_0 defined above. Since the system (9.8.73) is verified for all functions $\Lambda, \underline{\Lambda}, \psi$ we deduce, along Σ_0 ,

$$\begin{aligned} D &= 0, \\ B &= N(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi)(s) - N(0, 0, 0, \Lambda, \underline{\Lambda}, \psi)(s), \\ \underline{B} &= \underline{N}(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi)(s) - \underline{N}(0, 0, 0, \Lambda, \underline{\Lambda}, \psi)(s). \end{aligned}$$

In view of the properties of N, \underline{N} we deduce,

$$\begin{aligned} \left| N(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi)(s) - N(0, 0, 0, \Lambda, \underline{\Lambda}, \psi)(s) \right| &\lesssim \overset{\circ}{\epsilon} \sup_{\overset{\circ}{I}} (|B(s)| + |\underline{B}(s)|), \\ \left| \underline{N}(B, \underline{B}, D, \Lambda, \underline{\Lambda}, \psi)(s) - \underline{N}(0, 0, 0, \Lambda, \underline{\Lambda}, \psi)(s) \right| &\lesssim \overset{\circ}{\epsilon} \sup_{\overset{\circ}{I}} (|B(s)| + |\underline{B}(s)|). \end{aligned}$$

Hence,

$$\sup_{\overset{\circ}{I}} |B(s)| + \sup_{\overset{\circ}{I}} |\underline{B}(s)| \lesssim \overset{\circ}{\delta} \left(\sup_{\overset{\circ}{I}} |B(s)| + \sup_{\overset{\circ}{I}} |\underline{B}(s)| \right).$$

Hence B, \underline{B}, D vanish identically on Σ_0 .

Step 22. We have,

$$\left| \frac{dr}{ds} - 1 \right| \lesssim \overset{\circ}{\epsilon}. \tag{9.8.76}$$

Indeed, according to Step 15 and Lemma 9.8.3 we have

$$\begin{aligned} \frac{d}{ds} r(s) &= X_* \Big|_{SP} r^{\mathbf{S}} = \frac{1}{2\lambda} \varsigma \Psi' \nu^{\mathbf{S}}(r^{\mathbf{S}}) \Big|_{SP} = (-1 + \psi'(s)) \frac{1}{2\lambda} \varsigma \nu^{\mathbf{S}}(r^{\mathbf{S}}) \Big|_{SP} \\ &= (-1 + \psi'(s)) \left(\frac{1}{2\lambda} \varsigma \frac{r^{\mathbf{S}}}{2} (\varsigma^{\mathbf{S}})^{-1} \overline{\varsigma^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}} + a^{\mathbf{S}} \kappa^{\mathbf{S}})} \right) \Big|_{SP}. \end{aligned}$$

In view of Proposition 9.8.14, with $D = 0$, $|\psi'| \lesssim r^{-1} \overset{\circ}{\delta}$. We deduce,

$$\frac{d}{ds} r(s) = - \left(\frac{1}{2\lambda} \varsigma \frac{r^{\mathbf{S}}}{2} (\varsigma^{\mathbf{S}})^{-1} \overline{\varsigma^{\mathbf{S}}(\underline{\kappa}^{\mathbf{S}} + a^{\mathbf{S}} \kappa^{\mathbf{S}})} \right) \Big|_{SP} + O(\overset{\circ}{\delta}).$$

Step 23. Therefore the functions B, \underline{B}, D vanish identically on the hypersurface Σ_0 defined by the function $\Psi(s) = -s + \psi(s) + c_0$ which accomplishes the main task of Theorem 9.8.1. More precisely we have produced a local hypersurface Σ_0 , as defined in Step 12, foliated by the function $u^{\mathbf{S}}$, defined in Step 2 and extended in Step 3, such that the items 2-5 of the theorem are verified. The estimates in items 6-7 are an immediate consequence of Proposition 9.8.7. It only remains to prove the smoothness of the function $\Xi(s, \theta)$ in (9.8.54), Step 14 and the estimates for $F = (f, \underline{f}, \log \lambda)$ in the last part of the theorem. To check the differentiability properties recall that,

$$\begin{aligned} \partial_s \Xi(s, \theta) &= \left(\Psi'(s) + \partial_P U(\theta, P(s)) P'(s), 1 + \partial_P S(\theta, P(s)) P'(s), 0 \right), \\ \partial_\theta \Xi(s, \theta) &= \left(\partial_\theta U(\theta, P(s)), \partial_\theta S(\theta, P(s)), 1 \right), \end{aligned}$$

where,

$$\begin{aligned} \partial_P U(\cdot) P'(s) &= \Psi'(s) \partial_u U(\cdot) + \partial_s U(\cdot) + \Lambda'(s) \partial_\Lambda U(\cdot) + \underline{\Lambda}'(s) \partial_{\underline{\Lambda}} U(\cdot), \\ \partial_P S(\cdot) P'(s) &= \Psi'(s) \partial_u S(\cdot) + \partial_s S(\cdot) + \Lambda'(s) \partial_\Lambda S(\cdot) + \underline{\Lambda}'(s) \partial_{\underline{\Lambda}} S(\cdot). \end{aligned}$$

Thus to prove the smoothness of Ξ we need to appeal to the smoothness of U, S with respect to the parameters $\Lambda, \underline{\Lambda}$ and u, s . Though tedious, this can be easily done, by

appealing back to the coupled system of equations, (9.4.2), (9.4.3) and (9.4.4) (9.4.15), as in the proof of Theorem 9.4.2 and studying its dependence on these parameters.

Step 24. It only remains to derive the estimates (9.8.11) for the transition functions $F = (f, \underline{f}, \log \lambda)$. To start with we have, in view of the construction of Σ_0 and the estimates for $F = (f, \underline{f}, \log \lambda)$ of Theorem 9.4.2, for every $\mathbf{S} \subset \Sigma_0$

$$\|F\|_{\mathfrak{h}_{s_{\max}+1}(\mathbf{S})} \lesssim \overset{\circ}{\delta}. \quad (9.8.77)$$

To derive the remaining tangential derivatives of F along Σ_0 we commute the GCM system (9.4.2) of Proposition 9.4.1 with respect to $\nu = \nu^{\mathbf{S}} = e_3^{\mathbf{S}} + a^{\mathbf{S}} e_4^{\mathbf{S}}$ and then proceed, as in the proof of the a priori estimates of Theorem 9.4.4 to derive recursively the estimates, for $K = k_{\max} + 1$,

$$\begin{aligned} \|\nu^l(F)\|_{\mathfrak{h}_{K-l}(\mathbf{S})} &\lesssim \overset{\circ}{\delta} + \left| \int_{\mathbf{S}} \nu^l(f) e^{\Phi} \right| + \left| \int_{\mathbf{S}} \nu^l(\underline{f}) e^{\Phi} \right| \\ &+ \overset{\circ}{\delta} \|\nu^{\leq l-1} a^{\mathbf{S}}\|_{\mathfrak{h}_{K-l+1}(\mathbf{S})} + \|\nu^{\leq l-1} F\|_{\mathfrak{h}_{K-l+1}(\mathbf{S})}. \end{aligned} \quad (9.8.78)$$

We already have estimates for the $\ell = 1$ modes of $F = (f, \underline{f})$. To estimate the $\ell = 1$ modes of $\nu^l(f, \underline{f})$, $l \geq 1$, we make use of the equations (9.8.63) and the vanishing of the $\ell = 1$ modes of $\eta^{\mathbf{S}}$, $\xi^{\mathbf{S}}$ along Σ_0 to derive, recursively, for all $1 \leq l \leq K$,

$$\begin{aligned} \left| \int_{\mathbf{S}} \nu^l(f) e^{\Phi} \right| + \left| \int_{\mathbf{S}} \nu^l(\underline{f}) e^{\Phi} \right| &\lesssim \overset{\circ}{\delta} + \overset{\circ}{\delta} \left\| \nu^{\leq l-1} \left(a, \underline{\Omega}^{\mathbf{S}}, \varsigma^{\mathbf{S}}, \xi^{\mathbf{S}}, \eta^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}} \right) \right\|_{\mathfrak{h}_{K-l+1}(\mathbf{S})} \\ &+ \|\nu^{\leq l-1}(F)\|_{\mathfrak{h}_{K-l+1}(\mathbf{S})}. \end{aligned} \quad (9.8.79)$$

We can then proceed as in the proof of Proposition 9.8.7 derive, recursively, the estimates

$$\left\| \nu^{\leq l-1} \left(a^{\mathbf{S}}, \underline{\Omega}^{\mathbf{S}}, \varsigma^{\mathbf{S}}, \xi^{\mathbf{S}}, \eta^{\mathbf{S}}, \underline{\omega}^{\mathbf{S}} \right) \right\|_{\mathfrak{h}_{K-l+1}(\mathbf{S})} \lesssim 1 + \sum_{j=0}^l \|\nu^j(F)\|_{\mathfrak{h}_{K-j}(\mathbf{S})}. \quad (9.8.80)$$

Combining (9.8.78) (9.8.79) (9.8.80), we obtain

$$\|\nu^l(F)\|_{\mathfrak{h}_{K-l}(\mathbf{S})} \lesssim \overset{\circ}{\delta} + \sum_{j=0}^{l-1} \|\nu^j(F)\|_{\mathfrak{h}_{K-j}(\mathbf{S})},$$

which, together with (9.8.77), yields the desired estimate for all tangential derivatives

$$\sum_{j=0}^K \|\nu^j(F)\|_{\mathfrak{h}_{K-j}(\mathbf{S})} \lesssim \overset{\circ}{\delta}.$$

To complete the desired estimate for all derivatives we make use of the equations for $e_4^{\mathbf{S}}(F)$, due to the transversality conditions (9.4.12). The $e_3^{\mathbf{S}}$ derivatives can then be derived from $\nu^{\mathbf{S}} = e_3^{\mathbf{S}} + a^{\mathbf{S}}e_4^{\mathbf{S}}$ and the estimates for $a^{\mathbf{S}}$. This ends the proof of Theorem 9.8.1.

Step 25. To prove Corollary 9.8.2 we start with the rigidity statement of Corollary 9.4.5. Making use of the assumption $\int_{\mathbf{S}_0} f e^{\Phi} = O(\delta)$, $\int_{\mathbf{S}_0} \bar{f} e^{\Phi} = O(\delta)$ we infer that the estimate (9.8.77) holds true for \mathbf{S}_0 . We then proceed exactly as in Step 24 to derive the estimates (9.8.78) (9.8.79) (9.8.80) for our distinguished sphere \mathbf{S}_0 . Note that \mathbf{S}_0 can be viewed as a deformation of the unique background sphere sharing the same south pole.

Chapter 10

REGGE-WHEELER TYPE EQUATIONS

The goal of this chapter is to prove Theorem 5.3.4 and Theorem 5.3.5 concerning the weighted estimates for the solution ψ to

$$\square_2 \psi + V\psi = N, \quad V = \kappa \underline{\kappa}.$$

Recall that these theorems were used in Chapter 5 to prove Theorem M1.

The structure of the chapter is as follows.

- In section 10.1, we prove basic Morawetz estimates for ψ .
- In section 10.2, we prove r^p -weighted estimates in the spirit of Dafermos-Rodnianski [20] for ψ . In particular, we obtain as an immediate corollary the proof of Theorem 5.3.4 in the case $s = 0$ (i.e. without commuting the equation of ψ with derivatives).
- In section 10.3, we use a variation of the method of [4] to derive slightly stronger weighted estimates and prove Theorem 5.3.5 in the case $s = 0$ (i.e. without commuting the equation of $\check{\psi}$ with derivatives).
- In section 10.4, commuting the equation of ψ with derivatives, we complete the proof of Theorem 5.3.4 by controlling higher order derivatives of ψ , i.e. for $s \leq k_{small} + 30$. Also, commuting the equation of $\check{\psi}$ with derivatives, we complete the proof of Theorem 5.3.5 by controlling higher order derivatives of $\check{\psi}$, i.e. for $s \leq k_{small} + 29$.

10.1 Basic Morawetz Estimates

Recall

- the definitions in section 5.1.1 of ${}^{(trap)}\mathcal{M}$, ${}^{(trq\dot{p})}\mathcal{M}$, τ , $\Sigma(\tau)$ and ${}^{(trap)}\Sigma$,
- the main quantities involved in the energy and Morawetz estimates, e.g. $E[\psi](\tau)$, $\text{Mor}[\psi](\tau_1, \tau_2)$, $\text{Morr}[\psi](\tau_1, \tau_2)$, $F[\psi](\tau_1, \tau_2)$, $J_\delta[\psi, N](\tau_1, \tau_2)$ and $\dot{B}_{p;R}^s[\psi](\tau_1, \tau_2)$, introduced in section 5.1.4.

The following theorem claims basic Morawetz estimates for the solution ψ of the wave equation (5.3.5).

Theorem 10.1.1 (Morawetz). *Let ψ a reduced 2-scalar solution to*

$$\square_2\psi + V\psi = N, \quad V = \kappa\underline{\kappa}.$$

Let $\delta > 0$ be a fixed small constant verifying $0 < \epsilon \ll \delta$. The following estimates hold true in $\mathcal{M}(\tau_1, \tau_2)$, $0 \leq \tau_1 < \tau_2 \leq \tau_$,*

$$E[\psi](\tau_2) + \text{Morr}[\psi](\tau_1, \tau_2) + F[\psi](\tau_1, \tau_2) \lesssim E[\psi](\tau_1) + J_\delta[\psi, N](\tau_1, \tau_2) + O(\epsilon)\dot{B}_{\delta;4m_0}^s[\psi](\tau_1, \tau_2). \quad (10.1.1)$$

Remark 10.1.2. *Note that the bulk term $\dot{B}_{\delta;4m_0}^s[\psi](\tau_1, \tau_2)$ cannot yet be absorbed on the left hand side of the inequality. To do that we will rely on the r^p weighted estimates of Theorem 10.2.1.*

Remark 10.1.3. *In addition to ϵ and δ , the proof of Theorem 10.1.1 will involve several smallness constants: C^{-1} , $\widehat{\delta}$, δ_1 , $\delta_{\mathcal{H}}$, $\epsilon_{\mathcal{H}}$, $\Lambda_{\mathcal{H}}^{-1}$ and Λ^{-1} . These smallness constants will be chosen such that*

$$0 < \epsilon \ll \widehat{\delta}, \delta_{\mathcal{H}}, \epsilon_{\mathcal{H}}, \Lambda_{\mathcal{H}}^{-1}, \Lambda^{-1} \ll \delta_1 \ll C^{-1}. \quad (10.1.2)$$

In addition, $\widehat{\delta}$, $\epsilon_{\mathcal{H}}$, $\Lambda_{\mathcal{H}}^{-1}$ and Λ^{-1} will in fact be chosen towards the end of the proof as explicit powers of $\delta_{\mathcal{H}}$, see (10.1.62), (10.1.64) and Proposition 10.1.30.

The goal of this section is to prove Theorem 10.1.1. This will be achieved in section 10.1.15.

10.1.1 Structure of the proof of Theorem 10.1.1

To prove Theorem 10.1.1, we proceed as follows

- In section 10.1.2, we introduce a simplified set of assumptions of the Ricci coefficients which is sufficient in order to prove Theorem 10.1.1.
- In section 10.1.3, we discuss notations concerning functions depending on m and r .
- In section 10.1.4, we compute the deformation tensor of the vectorfields R , T , and $X = f(r, m)R$.
- In section 10.1.5, we introduce the basic integral identities for wave equations that will be used repeatedly in the proof of Theorem 10.1.1.
- In section 10.1.6, we derive the main Morawetz identity.
- In section 10.1.7, we derive a first estimate. This estimate is insufficient due to
 - a lack of positivity of the bulk in the region $3m \leq r \leq 4m$,
 - a log divergence of a suitable choice of vectorfield at $r = 2m$,
 - a degeneracy at $r = 2m$.
- In section 10.1.8, we add a correction and rely on a Poincaré inequality to obtain a positive estimate also on the region $3m \leq r \leq 4m$.
- In section 10.1.9, we perform a cut-off to remove above mentioned log divergence at $r = 2m$.
- In section 10.1.10, we introduce the red shift vectorfield to remove the above mentioned degeneracy at $r = 2m$.
- In section 10.1.11, we combine the previous estimates with the redshift vectorfield to obtain a bulk term suitable on the whole spacetime \mathcal{M} .
- In section 10.1.12, we prove the positivity of the boundary terms arising from adding a large multiple of the energy estimate to the Morawetz estimate.
- In section 10.1.13, combining the good properties of the bulk and of the boundary terms established so far, we obtain a first Morawetz estimate providing in particular the control of the quantity $\text{Mor}[\psi]$.
- In section 10.1.14, we analyse an error term appearing in the right-hand side of the above mentioned Morawetz estimate.

- Finally, in section 10.1.15, we add a correction to upgrade the control of $\text{Mor}[\psi]$ to the control of the quantity $\text{Morr}[\psi]$, hence concluding the proof of Theorem 10.1.1.

10.1.2 A simplified set of assumptions

To prove Theorem 10.1.1, it suffices to make a simplified set of assumptions. Define

$$u_{\text{trap}} = \begin{cases} 1 + \tau & \text{for } r \in \left[\frac{5m_0}{2}, \frac{7m_0}{2}\right], \\ 1 & \text{for } r \notin \left[\frac{5m_0}{2}, \frac{7m_0}{2}\right]. \end{cases} \quad (10.1.3)$$

For $k = 0, 1$, we assume the following.

Mor1. The renormalized Ricci coefficients $\check{\Gamma}^{\leq k}$ verify on $\mathcal{M} = {}^{(int)}\mathcal{M} \cup {}^{(ext)}\mathcal{M}$,

$$\begin{aligned} |\check{\Gamma}^{\leq k}| &\lesssim \epsilon r^{-1} u_{\text{trap}}^{-1-\delta_{dec}}, \\ \left| \mathfrak{d}^{\leq k} \left(\omega + \frac{m}{r^2}, \xi \right) \right| &\lesssim \epsilon r^{-2} u_{\text{trap}}^{-1-\delta_{dec}}. \end{aligned} \quad (10.1.4)$$

Mor2. The Gauss curvature K of S and ρ verify,

$$\begin{aligned} \left| \mathfrak{d}^{\leq k} \left(\rho + \frac{2m}{r^3} \right) \right| &\lesssim \epsilon r^{-2} u_{\text{trap}}^{-1-\delta_{dec}}, \\ \left| \mathfrak{d}^{\leq k} \left(K - \frac{1}{r^2} \right) \right| &\lesssim \epsilon r^{-2} u_{\text{trap}}^{-1-\delta_{dec}}. \end{aligned} \quad (10.1.5)$$

Mor3. We also assume

$$\begin{aligned} |m - m_0| &\lesssim \epsilon m_0, \\ |\mathfrak{d}^{\leq k}(e_3 m, r^2 e_4 m)| &\lesssim \epsilon u_{\text{trap}}^{-1-\delta_{dec}}. \end{aligned} \quad (10.1.6)$$

Remark 10.1.4. Note that in the case when the bootstrap constant $\epsilon = 0$, i.e. in Schwarzschild, the assumptions made above are consistent with the behavior relative to the regular frame (near horizon)

$$e_3 = \Upsilon^{-1} \partial_t - \partial_r, e_4 = \partial_t + \Upsilon \partial_r.$$

10.1.3 Functions depending on m and r

In order to prove Theorem 10.1.1, we will adapt the derivation of the Morawetz estimate for the wave equation in Schwarzschild. In particular, we will need to consider various

scalar functions, used to define suitable analogs of the vectorfields in Schwarzschild, which depend on m and r . Now, m is now a scalar function unlike the Schwarzschild case where it is constant. To take this into account, we will rely on the following lemma.

Lemma 10.1.5. *Let $f = f(r, m)$ a C^1 function of r and m . Then, we have*

$$\begin{aligned} e_4(f(r, m)) &= \partial_r f(r, m)e_4(r) + O(\epsilon r^{-2} u_{trap}^{-1-\delta_{dec}} |\partial_m f|), \\ e_3(f(r, m)) &= \partial_r f(r, m)e_3(r) + O(\epsilon u_{trap}^{-1-\delta_{dec}} |\partial_m f|), \\ e_4(e_3(f(r, m))) &= \partial_r^2 f(r, m)e_4(r)e_3(r) + \partial_r f(r, m)e_4(e_3(r)) \\ &\quad + O(\epsilon r^{-2} u_{trap}^{-1-\delta_{dec}} (r |\partial_r \partial_m f| + |\partial_m^2 f|)), \\ e_3(e_4(f(r, m))) &= \partial_r^2 f(r, m)e_4(r)e_3(r) + \partial_r f(r, m)e_3(e_4(r)) \\ &\quad + O(\epsilon r^{-2} u_{trap}^{-1-\delta_{dec}} (r |\partial_r \partial_m f| + |\partial_m^2 f|)), \\ e_\theta(f(r, m)) &= 0. \end{aligned}$$

Proof. Straightforward verification using (10.1.6). □

Remark 10.1.6. *Note that in the sequel, $\partial_r f$ will not denote a spacetime coordinate vectorfield applied to f , but instead the partial derivative with respect to the variable r of the function $f(r, m)$.*

10.1.4 Deformation tensors of the vectorfields R, T, X

Recall the definition (5.1.10) of the regular vectorfields¹,

$$T = \frac{1}{2}(e_4 + \Upsilon e_3), \quad R = \frac{1}{2}(e_4 - \Upsilon e_3).$$

Note that,

$$-g(T, T) = g(R, R) = \Upsilon, \quad g(T, R) = 0.$$

Note also that,

$$R(r) = 1 - \frac{2m}{r} + O(\epsilon u_{trap}^{-1-\delta_{dec}}), \quad T(r) = O(\epsilon u_{trap}^{-1-\delta_{dec}}).$$

Lemma 10.1.7. *The following hold true.*

¹In Schwarzschild, in standard coordinates, we have $T = \partial_t$, $R = \Upsilon \partial_r$ which are regular near the horizon.

1. The components of the deformation tensor of $R = \frac{1}{2}(e_4 - \Upsilon e_3)$ are given by,

$$\begin{aligned} \left| {}^{(R)}\pi_{34} + \frac{4m}{r^2} \right| &\lesssim \epsilon r^{-1} u_{\text{trap}}^{-1-\delta_{\text{dec}}}, \\ \left| {}^{(R)}\pi(e_A, e_B) - \frac{2}{r}\Upsilon\delta_{AB} \right| &\lesssim \epsilon r^{-1} u_{\text{trap}}^{-1-\delta_{\text{dec}}}, \\ \left| {}^{(R)}\pi_{33} \right| &\lesssim \epsilon r^{-1} u_{\text{trap}}^{-1-\delta_{\text{dec}}}, \\ \left| {}^{(R)}\pi_{3\theta} \right| &\lesssim \epsilon r^{-1} u_{\text{trap}}^{-1-\delta_{\text{dec}}}, \\ \left| {}^{(R)}\pi_{4\theta} \right| &\lesssim \epsilon r^{-1} u_{\text{trap}}^{-1-\delta_{\text{dec}}}. \end{aligned}$$

Moreover,

$$\left| {}^{(R)}\pi_{44} \right| \lesssim \epsilon r^{-2} u_{\text{trap}}^{-1-\delta_{\text{dec}}}.$$

2. If $V = -\kappa \underline{\kappa}$ we have,

$$\begin{aligned} e_3(V) &= \frac{8}{r^3} \left(1 - \frac{3m}{r} \right) + O(\epsilon) r^{-3} u_{\text{trap}}^{-1-\delta_{\text{dec}}}, \\ e_4(V) &= -\frac{8\Upsilon}{r^3} \left(1 - \frac{3m}{r} \right) + O(\epsilon) r^{-3} u_{\text{trap}}^{-1-\delta_{\text{dec}}}, \end{aligned} \tag{10.1.7}$$

and,

$$\begin{aligned} R(V) &= -\frac{8\Upsilon}{r^3} \left(1 - \frac{3m}{r} \right) + O(\epsilon) r^{-3} u_{\text{trap}}^{-1-\delta_{\text{dec}}}, \\ T(V) &= O(\epsilon) r^{-3} u_{\text{trap}}^{-1-\delta_{\text{dec}}}. \end{aligned}$$

3. All components of the deformation tensor of $T = \frac{1}{2}(e_4 + \Upsilon e_3)$ can be bounded by $O(\epsilon r^{-1} u_{\text{trap}}^{-1-\delta_{\text{dec}}})$. Moreover,

$$\left| {}^{(T)}\pi_{44} \right| \lesssim \epsilon r^{-2} u_{\text{trap}}^{-1-\delta_{\text{dec}}}.$$

Proof. We have

$$\begin{aligned}
{}^{(R)}\pi_{44} &= \mathbf{g}(\mathbf{D}_4(e_4 - \Upsilon e_3), e_4) = 2e_4(\Upsilon) + 4\Upsilon\omega, \\
{}^{(R)}\pi_{34} &= \frac{1}{2}\mathbf{g}(\mathbf{D}_3(e_4 - \Upsilon e_3), e_4) + \frac{1}{2}\mathbf{g}(\mathbf{D}_4(e_4 - \Upsilon e_3), e_3) \\
&= e_3(\Upsilon) - 2\Upsilon\omega + 2\omega, \\
{}^{(R)}\pi_{33} &= \mathbf{g}(\mathbf{D}_3(e_4 - \Upsilon e_3), e_3) = -4\omega, \\
{}^{(R)}\pi_{AB} &= \frac{1}{2}\mathbf{g}(\mathbf{D}_A(e_4 - \Upsilon e_3), e_B) + \frac{1}{2}\mathbf{g}(\mathbf{D}_B(e_4 - \Upsilon e_3), e_A), \\
&= {}^{(1+3)}\chi_{AB} - \Upsilon {}^{(1+3)}\underline{\chi}_{AB} \\
&= \frac{1}{2}(\kappa - \Upsilon\underline{\kappa})\delta_{AB} + {}^{(1+3)}\widehat{\chi}_{AB} - \Upsilon {}^{(1+3)}\widehat{\underline{\chi}}_{AB}.
\end{aligned}$$

Note that,

$$\begin{aligned}
e_3(\Upsilon) &= e_3\left(1 - \frac{2m}{r}\right) = \frac{2m}{r^2}e_3(r) - \frac{2e_3m}{r} = \frac{m}{r}(\underline{\kappa} + \underline{A}) + O(\epsilon r^{-1}u_{trap}^{-1-\delta_{dec}}) \\
&= \frac{m}{r}\underline{\kappa} + O(\epsilon r^{-1}u_{trap}^{-1-\delta_{dec}}) = -\frac{2m}{r^2} + O(\epsilon r^{-1}u_{trap}^{-1-\delta_{dec}}), \\
e_4(\Upsilon) &= e_4\left(1 - \frac{2m}{r}\right) = \frac{2m}{r^2}e_4(r) - \frac{2e_4m}{r} = \frac{m}{r}(\kappa + A) + O(\epsilon r^{-2}u_{trap}^{-1-\delta_{dec}}) \\
&= \frac{m}{r}\kappa + O(\epsilon r^{-2}u_{trap}^{-1-\delta_{dec}}) = \frac{2m}{r^2}\Upsilon + O(\epsilon r^{-2}u_{trap}^{-1-\delta_{dec}}).
\end{aligned}$$

Thus

$$\begin{aligned}
{}^{(R)}\pi_{44} &= O(\epsilon r^{-2}u_{trap}^{-1-\delta_{dec}}), \\
{}^{(R)}\pi_{33} &= O(\epsilon r^{-1}u_{trap}^{-1-\delta_{dec}}), \\
{}^{(R)}\pi_{34} &= -4\frac{m}{r^2} + O(\epsilon r^{-1}u_{trap}^{-1-\delta_{dec}}), \\
{}^{(R)}\pi_{AB} &= \frac{2\Upsilon}{r}\delta_{AB} + O(\epsilon r^{-1}u_{trap}^{-1-\delta_{dec}}).
\end{aligned}$$

Also, in view of,

$$\left| \xi, \underline{\xi}, \eta, \underline{\eta}, \zeta \right| \lesssim \epsilon r^{-1}u_{trap}^{-1-\delta_{dec}},$$

we deduce,

$$\left| {}^{(R)}\pi_{3\theta}, {}^{(R)}\pi_{4\theta} \right| \lesssim \epsilon r^{-1}u_{trap}^{-1-\delta_{dec}}$$

as desired.

To prove the second part of the lemma we write,

$$\begin{aligned}
e_3(V) &= -e_3(\kappa)\underline{\kappa} - \kappa e_3(\underline{\kappa}) \\
&= -\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa + 2\rho + O(\epsilon r^{-2}u_{trap}^{-1-\delta_{dec}})\right)\underline{\kappa} - \kappa\left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega}\underline{\kappa} + O(\epsilon r^{-2}u_{trap}^{-1-\delta_{dec}})\right) \\
&= (\kappa\underline{\kappa} - 2\rho)\underline{\kappa} + O(\epsilon r^{-3}u_{trap}^{-1-\delta_{dec}}).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\kappa\underline{\kappa} - 2\rho &= -\left(\frac{2\Upsilon}{r} + O(\epsilon)r^{-1}u_{trap}^{-1-\delta_{dec}}\right)\left(\frac{2}{r} + O(\epsilon)r^{-1}u_{trap}^{-1-\delta_{dec}}\right) + \frac{4m}{r^3} + O(\epsilon r^{-3}u_{trap}^{-1-\delta_{dec}}) \\
&= -\frac{4}{r^2}\left(1 - \frac{3m}{r}\right) + O(\epsilon r^{-2}u_{trap}^{-1-\delta_{dec}}).
\end{aligned}$$

Hence,

$$\begin{aligned}
e_3(V) &= (\kappa\underline{\kappa} - 2\rho)\underline{\kappa} + O(\epsilon r^{-3}u_{trap}^{-1-\delta_{dec}}) \\
&= \frac{8}{r^3}\left(1 - \frac{3m}{r}\right) + O(\epsilon r^{-3}u_{trap}^{-1-\delta_{dec}})
\end{aligned}$$

and similarly for $e_4(V)$. Thus,

$$\begin{aligned}
R(V) &= \frac{1}{2}(e_4 - \Upsilon e_3)V = -\frac{8\Upsilon}{r^3}\left(1 - \frac{3m}{r}\right) + O(\epsilon r^{-3}u_{trap}^{-1-\delta_{dec}}), \\
T(V) &= \frac{1}{2}(e_4 + \Upsilon e_3)V = O(\epsilon)r^{-3}u_{trap}^{-1-\delta_{dec}},
\end{aligned}$$

as desired.

To prove the last part of the lemma we write,

$$\begin{aligned}
{}^{(T)}\pi_{44} &= \mathbf{g}(\mathbf{D}_4(e_4 + \Upsilon e_3), e_4) = -2e_4(\Upsilon) - 4\Upsilon\omega, \\
{}^{(T)}\pi_{34} &= \frac{1}{2}\mathbf{g}(\mathbf{D}_3(e_4 + \Upsilon e_3), e_4) + \frac{1}{2}\mathbf{g}(\mathbf{D}_4(e_4 + \Upsilon e_3), e_3) \\
&= -e_3(\Upsilon) + 2\Upsilon\underline{\omega} + 2\omega, \\
{}^{(T)}\pi_{33} &= \mathbf{g}(\mathbf{D}_3(e_4 + \Upsilon e_3), e_3) = -4\underline{\omega}, \\
{}^{(T)}\pi_{AB} &= \frac{1}{2}\mathbf{g}(\mathbf{D}_A(e_4 + \Upsilon e_3), e_B) + \frac{1}{2}\mathbf{g}(\mathbf{D}_B(e_4 + \Upsilon e_3), e_A), \\
&= {}^{(1+3)}\chi_{AB} + \Upsilon {}^{(1+3)}\underline{\chi}_{AB} \\
&= \frac{1}{2}(\kappa + \Upsilon\underline{\kappa})\delta_{AB} + {}^{(1+3)}\widehat{\chi}_{AB} + \Upsilon {}^{(1+3)}\widehat{\underline{\chi}}_{AB},
\end{aligned}$$

and the proof continues as above in view of our assumptions. \square

Consider now $X = f(r, m)R$ and ${}^{(X)}\pi$ its deformation tensor. We have the following lemma.

Lemma 10.1.8. *Let $X = f(r, m)R$ and ${}^{(X)}\pi$ its deformation tensor. We have,*

$${}^{(X)}\pi = {}^{(X)}\dot{\pi} + \epsilon {}^{(X)}\ddot{\pi}$$

where²

- The only nonvanishing components of ${}^{(X)}\dot{\pi}$ are

$$\begin{aligned} {}^{(X)}\dot{\pi}_{33} &= 2\partial_r f, \\ {}^{(X)}\dot{\pi}_{44} &= 2\partial_r f \Upsilon^2, \\ {}^{(X)}\dot{\pi}_{34} &= -\frac{4m}{r^2} f - 2\partial_r f \Upsilon, \\ {}^{(X)}\dot{\pi}_{AB} &= f \frac{2\Upsilon}{r} \delta_{AB}. \end{aligned}$$

- All components of ${}^{(X)}\ddot{\pi}$ verify,

$$\left| {}^{(X)}\ddot{\pi} \right| \lesssim r^{-1} u_{trap}^{-1-\delta_{dec}} (|f| + r|\partial_m f| + r^2|\partial_r f|).$$

Moreover,

$$\left| {}^{(X)}\ddot{\pi}_{44} \right| \lesssim r^{-2} u_{trap}^{-1-\delta_{dec}} (|f| + r|\partial_m f| + r^2|\partial_r f|).$$

Proof. Clearly,

$${}^{(X)}\pi_{\mu\nu} = f {}^{(R)}\pi_{\mu\nu} + e_\mu f R_\nu + e_\nu f R_\mu.$$

Therefore, since $\mathbf{g}(R, e_3) = -1$, $\mathbf{g}(R, e_4) = \Upsilon$ and,

$$\left| e_4(r) - \Upsilon, e_3(r) + 1 \right| \lesssim \epsilon u_{trap}^{-1-\delta_{dec}},$$

²Recall from Remark 10.1.6 that $\partial_r f$ does not denote a spacetime coordinate vectorfield applied to f , but instead the partial derivative with respect to the variable r of the function $f(r, m)$.

and using Lemma 10.1.5, we deduce,

$$\begin{aligned}
{}^{(X)}\pi_{33} &= f^{(R)}\pi_{33} - 2e_3(f) = f^{(R)}\pi_{33} - 2\partial_r f e_3(r) - 2\partial_m f e_3(m) \\
&= 2\partial_r f + O\left(\epsilon r^{-1} u_{trap}^{-1-\delta_{dec}} (|f| + r|\partial_m f| + r^2|\partial_r f|)\right) \\
{}^{(X)}\pi_{44} &= f^{(R)}\pi_{44} + 2\Upsilon e_4(f) = f^{(R)}\pi_{44} + 2\Upsilon\partial_r f e_4(r) + 2\Upsilon\partial_m f e_4(m) \\
&= 2\partial_r f \Upsilon^2 + O\left(\epsilon r^{-2} u_{trap}^{-1-\delta_{dec}} (|f| + r|\partial_m f| + r^2|\partial_r f|)\right) \\
{}^{(X)}\pi_{34} &= f^{(R)}\pi_{34} + e_3(f)\Upsilon - e_4(f) \\
&= f^{(R)}\pi_{34} + (\partial_r f e_3(r) + \partial_m f e_3(m))\Upsilon - (\partial_r f e_4(r) + \partial_m f e_4(m)) \\
&= -\frac{4m}{r^2} f - 2\partial_r f \Upsilon + O\left(\epsilon r^{-1} u_{trap}^{-1-\delta_{dec}} (|f| + r|\partial_m f| + r^2|\partial_r f|)\right).
\end{aligned}$$

This concludes the proof of the lemma. \square

10.1.5 Basic Integral Identities

We recall, see section 2.4.1, that wave equations for $\psi \in \mathfrak{s}_2(\mathcal{M})$ of the form

$$\square_2 \psi = V\psi + N[\psi], \quad V = -\kappa \underline{\kappa}, \quad (10.1.8)$$

can be lifted to the spacetime version,

$$\dot{\square} \Psi + V\Psi = N[\Psi] \quad (10.1.9)$$

where $\Psi \in \mathcal{S}_2(\mathcal{M})$ and $N[\Psi] \in \mathcal{S}_2(\mathcal{M})$ are defined according to Proposition 2.4.5. In fact,

$$\begin{aligned}
\Psi_{\theta\theta} &= -\Psi_{\varphi\varphi} = \psi, & \Psi_{\theta\varphi} &= 0. \\
N_{\theta\theta}[\Psi] &= N_{\varphi\varphi}[\Psi] = N(\psi), & N[\Psi]_{\theta\varphi} &= 0.
\end{aligned}$$

All estimates for (10.1.9) derived in this section can be easily transferred to estimates for (10.1.8) and vice versa.

Consider wave equations of the form,

$$\dot{\square}_{\mathbf{g}} \Psi = V\Psi + N \quad (10.1.10)$$

with $\Psi \in \mathcal{S}_2(\mathcal{M})$ and N a given symmetric traceless tensor, i.e. $N \in \mathcal{S}_2(\mathcal{M})$.

Proposition 10.1.9. *Assume $\Psi \in \mathcal{S}_2(\mathcal{M})$ verifies (10.1.9). Then,*

1. The energy momentum tensor $\mathcal{Q} = \mathcal{Q}[\Psi]$ given by,

$$\begin{aligned}\mathcal{Q}_{\mu\nu} &:= \dot{\mathbf{D}}_\mu \Psi \cdot \dot{\mathbf{D}}_\nu \Psi - \frac{1}{2} \mathbf{g}_{\mu\nu} \left(\dot{\mathbf{D}}_\lambda \Psi \cdot \dot{\mathbf{D}}^\lambda \Psi + V \Psi \cdot \Psi \right) \\ &= \dot{\mathbf{D}}_\mu \Psi \cdot \dot{\mathbf{D}}_\nu \Psi - \frac{1}{2} \mathbf{g}_{\mu\nu} \mathcal{L}(\Psi)\end{aligned}$$

verifies,

$$\mathbf{D}^\nu \mathcal{Q}_{\mu\nu} = \dot{\mathbf{D}}_\mu \Psi \cdot \mathcal{N}[\Psi] + \dot{\mathbf{D}}^\nu \Psi^A \mathbf{R}_{AB\nu\mu} \Psi^B - \frac{1}{2} \mathbf{D}_\mu V \Psi \cdot \Psi.$$

2. The null components of \mathcal{Q} are given by,

$$\begin{aligned}\mathcal{Q}_{33} &= |e_3 \Psi|^2, \\ \mathcal{Q}_{44} &= |e_4 \Psi|^2, \\ \mathcal{Q}_{34} &= |\nabla \Psi|^2 + V |\Psi|^2,\end{aligned}$$

and,

$$\mathbf{g}^{\mu\nu} \mathcal{Q}_{\mu\nu} = -\mathcal{L}(\Psi) - V |\Psi|^2.$$

Also,

$$|\mathcal{L}(\Psi)| \lesssim |e_3 \Psi| |e_4 \Psi| + |\nabla \Psi|^2 + V |\Psi|^2$$

and

$$\begin{aligned}|\mathcal{Q}_{AB}| &\leq |e_3 \Psi| |e_4 \Psi| + |\nabla \Psi|^2 + |V| |\Psi|^2, \\ |\mathcal{Q}_{A3}| &\leq |e_3 \Psi| |\nabla \Psi|, \\ |\mathcal{Q}_{A4}| &\leq |e_4 \Psi| |\nabla \Psi|.\end{aligned}$$

3. Introducing

$$\widehat{\mathcal{Q}}_{34} := \mathcal{Q}_{34} - V |\Psi|^2 = |\nabla \Psi|^2$$

we have,

$$-\widehat{\mathcal{Q}}_{34} + \mathcal{Q}_{\theta\theta} + \mathcal{Q}_{\varphi\varphi} = -\mathcal{L}(\Psi).$$

4. Let $X = ae_3 + be_4$. Then, since $\mathbf{R}_{AB34} = 0$ in an axially symmetric polarized spacetime,

$$\mathbf{D}^\mu (\mathcal{Q}_{\mu\nu} X^\nu) = \frac{1}{2} \mathcal{Q} \cdot {}^{(X)}\pi + X(\Psi) \cdot \mathcal{N}[\Psi] - \frac{1}{2} X(V) \Psi \cdot \Psi.$$

5. Let $X = ae_3 + be_4$ as above, w a scalar function and M a one form. Define,

$$\mathcal{P}_\mu = \mathcal{P}_\mu[X, w, M] = \mathcal{Q}_{\mu\nu}X^\nu + \frac{1}{2}w\Psi\dot{\mathbf{D}}_\mu\Psi - \frac{1}{4}|\Psi|^2\partial_\mu w + \frac{1}{4}|\Psi|^2M_\mu.$$

Then,

$$\begin{aligned} \mathbf{D}^\mu\mathcal{P}_\mu[X, w, M] &= \frac{1}{2}\mathcal{Q} \cdot {}^{(X)}\pi - \frac{1}{2}X(V)\Psi \cdot \Psi + \frac{1}{2}w\mathcal{L}[\Psi] - \frac{1}{4}|\Psi|^2\Box_{\mathbf{g}}w \\ &\quad + \frac{1}{4}\dot{\mathbf{D}}^\mu(|\Psi|^2M_\mu) + \left(X(\Psi) + \frac{1}{2}w\Psi\right) \cdot \mathcal{N}[\Psi]. \end{aligned} \quad (10.1.11)$$

Proof. See sections D.1.4 and D.2 in the appendix. \square

Notation. For convenience we introduce the notation,

$$\mathcal{E}[X, w, M](\Psi) := \mathbf{D}^\mu\mathcal{P}_\mu[X, w, M] - \left(X(\Psi) + \frac{1}{2}w\Psi\right) \cdot \mathcal{N}[\Psi]. \quad (10.1.12)$$

Thus equation (10.1.11) becomes,

$$\begin{aligned} \mathcal{E}[X, w, M](\Psi) &= \frac{1}{2}\mathcal{Q} \cdot {}^{(X)}\pi - \frac{1}{2}X(V)\Psi \cdot \Psi + \frac{1}{2}w\mathcal{L}[\Psi] \\ &\quad - \frac{1}{4}|\Psi|^2\Box_{\mathbf{g}}w + \frac{1}{4}\dot{\mathbf{D}}^\mu(|\Psi|^2M_\mu). \end{aligned} \quad (10.1.13)$$

When $M = 0$ we simply write $\mathcal{E}[X, w](\Psi)$.

10.1.6 Main Morawetz Identity

Lemma 10.1.10. *Let $f(r, m)$ a function of r and m , and let X a vectorfield defined by $X = f(r, m)R$. Then, we have³,*

$$\mathcal{Q} \cdot {}^{(X)}\dot{\pi} = f \left(-\frac{2m}{r^2} + \frac{2\Upsilon}{r} \right) |\nabla\Psi|^2 + 2\partial_r f |R\Psi|^2 - \left(\frac{2\Upsilon}{r}f + \Upsilon\partial_r f \right) \mathcal{L} - \frac{2m}{r^2} fV|\Psi|^2.$$

where ${}^{(X)}\dot{\pi}$ has been defined in Lemma 10.1.8.

³Recall from Remark 10.1.6 that $\partial_r f$ does not denote a spacetime coordinate vectorfield applied to f , but instead the partial derivative with respect to the variable r of the function $f(r, m)$.

Proof. In view of Lemma 10.1.8, we have

$$\begin{aligned}\mathcal{Q} \cdot (X)\dot{\pi} &= \frac{1}{2}\mathcal{Q}_{34}\dot{\pi}_{34} + \frac{1}{4}\mathcal{Q}_{44}\dot{\pi}_{33} + \frac{1}{4}\mathcal{Q}_{33}\dot{\pi}_{44} + \mathcal{Q}_{AB}\dot{\pi}_{AB} \\ &= -\frac{2m}{r^2}f\mathcal{Q}_{34} - \partial_r f \Upsilon \mathcal{Q}_{34} + \frac{1}{2}\mathcal{Q}_{44}\partial_r f + \frac{1}{2}\mathcal{Q}_{33}\Upsilon^2\partial_r f + \frac{2\Upsilon}{r}f\delta^{AB}\mathcal{Q}_{AB} \\ &= -\frac{2m}{r^2}f\mathcal{Q}_{34} + \frac{2\Upsilon}{r}f\delta^{AB}\mathcal{Q}_{AB} + \frac{1}{2}\partial_r f (\mathcal{Q}_{44} - 2\Upsilon\mathcal{Q}_{34} + \Upsilon^2\mathcal{Q}_{33})\end{aligned}$$

Note that,

$$(\mathcal{Q}_{44} - 2\Upsilon\mathcal{Q}_{34} + \Upsilon^2\mathcal{Q}_{33}) = 4\mathcal{Q}_{RR}$$

and, since $\mathbf{g}^{\mu\nu}\mathcal{Q}_{\mu\nu} = -\mathcal{L}(\Psi) - V|\Psi|^2$,

$$\delta^{AB}\mathcal{Q}_{AB} = \mathcal{Q}_{34} - \mathcal{L} - V|\Psi|^2 = \widehat{\mathcal{Q}}_{34} - \mathcal{L}.$$

Hence,

$$\begin{aligned}\mathcal{Q} \cdot (X)\dot{\pi} &= -\frac{2m}{r^2}f\mathcal{Q}_{34} + \frac{2\Upsilon}{r}f(\widehat{\mathcal{Q}}_{34} - \mathcal{L}) + 2\partial_r f \mathcal{Q}_{RR} \\ &= -\frac{2m}{r^2}f(\widehat{\mathcal{Q}}_{34} + V|\Psi|^2) + \frac{2\Upsilon}{r}f(\widehat{\mathcal{Q}}_{34} - \mathcal{L}) + 2\partial_r f \mathcal{Q}_{RR} \\ &= f\left(-\frac{2m}{r^2} + \frac{2\Upsilon}{r}\right)\widehat{\mathcal{Q}}_{34} + 2\partial_r f \mathcal{Q}_{RR} - \frac{2\Upsilon}{r}f\mathcal{L} - \frac{2m}{r^2}fV|\Psi|^2.\end{aligned}$$

Finally,

$$\mathcal{Q}_{RR} = |R\Psi|^2 - \frac{1}{2}\mathbf{g}(R, R)\mathcal{L} = |R\Psi|^2 - \frac{1}{2}\Upsilon\mathcal{L}.$$

Hence,

$$\begin{aligned}\mathcal{Q} \cdot (X)\dot{\pi} &= 2f\left(-\frac{m}{r^2} + \frac{\Upsilon}{r}\right)\widehat{\mathcal{Q}}_{34} + 2\partial_r f \left(|R\Psi|^2 - \frac{1}{2}\Upsilon\mathcal{L}\right) - \frac{2\Upsilon}{r}f\mathcal{L} - \frac{2m}{r^2}fV|\Psi|^2 \\ &= 2f\left(-\frac{m}{r^2} + \frac{\Upsilon}{r}\right)|\nabla\Psi|^2 + 2\partial_r f |R\Psi|^2 - \left(\frac{2\Upsilon}{r}f + \partial_r f \Upsilon\right)\mathcal{L} - \frac{2m}{r^2}fV|\Psi|^2.\end{aligned}$$

This concludes the proof of the lemma. \square

We shall also make use of the following lemma.

Lemma 10.1.11. *If $f = f(r, m)$, then*

$$\begin{aligned}\square_{\mathbf{g}}(f(r, m)) &= r^{-2}\partial_r(r^2\Upsilon\partial_r f) + O(\epsilon r^{-2}u_{trap}^{-1-\delta_{dec}})\left[r^2|\partial_r^2 f(r, m)| + r|\partial_r f(r, m)|\right. \\ &\quad \left.+ r|\partial_r\partial_m f(r, m)| + |\partial_m^2 f(r, m)|\right].\end{aligned}$$

Proof. Recall from Lemma 2.4.1 that, for a general scalar f ,

$$\square_{\mathbf{g}}f = -\frac{1}{2}(e_3e_4 + e_4e_3)f + \underline{\Delta}f + \left({}^{(1+3)}\underline{\omega} - \frac{1}{2} {}^{(1+3)}\text{tr}\underline{\chi} \right) e_4f + \left({}^{(1+3)}\omega - \frac{1}{2} {}^{(1+3)}\text{tr}\chi \right) e_3f.$$

Recall that,

$${}^{(1+3)}\text{tr}\chi = 2\chi - \vartheta, \quad {}^{(1+3)}\text{tr}\underline{\chi} = 2\underline{\chi} - \underline{\vartheta}, \quad {}^{(1+3)}\omega = \omega, \quad {}^{(1+3)}\underline{\omega} = \underline{\omega}$$

and

$$\underline{\Delta}f = e_\theta e_\theta f + (e_\theta \Phi)^2 e_\theta f.$$

Using Lemma 10.1.5, we deduce, for a function $f = f(r, m)$,

$$\begin{aligned} \square_{\mathbf{g}}f &= -\frac{1}{2}(e_3e_4 + e_4e_3)f + \left(\underline{\omega} - \frac{1}{2}\underline{\kappa} \right) e_4f + \left(\omega - \frac{1}{2}\kappa \right) e_3f \\ &= -\partial_r^2 f(r, m) e_3(r) e_4(r) - \frac{1}{2} \partial_r f(r, m) (e_3 e_4(r) + e_4 e_3(r)) \\ &\quad - \frac{1}{2} \underline{\kappa} \partial_r f(r, m) e_4 r + \left(\omega - \frac{1}{2} \kappa \right) \partial_r f(r, m) e_3(r) + O(\epsilon r^{-2} u_{\text{trap}}^{-1-\delta_{\text{dec}}}) \left[r^2 |\partial_r^2 f(r, m)| \right. \\ &\quad \left. + r |\partial_r f(r, m)| + r |\partial_r \partial_m f(r, m)| + |\partial_m^2 f(r, m)| \right] \\ &= -\partial_r^2 f(r, m) \left(-\Upsilon + O(\epsilon u_{\text{trap}}^{-1-\delta_{\text{dec}}}) \right) + \partial_r f(r, m) \frac{m}{r^2} + \partial_r f(r, m) \frac{\Upsilon}{r} + \frac{r-m}{r^2} \partial_r f(r, m) \\ &\quad + O(\epsilon r^{-2} u_{\text{trap}}^{-1-\delta_{\text{dec}}}) \left[r^2 |\partial_r^2 f(r, m)| + r |\partial_r f(r, m)| + r |\partial_r \partial_m f(r, m)| + |\partial_m^2 f(r, m)| \right] \\ &= \Upsilon \partial_r^2 f(r, m) + \partial_r f(r, m) \left(\frac{2}{r} - \frac{2m}{r^2} \right) \\ &\quad + O(\epsilon r^{-2} u_{\text{trap}}^{-1-\delta_{\text{dec}}}) \left[r^2 |\partial_r^2 f(r, m)| + r |\partial_r f(r, m)| + r |\partial_r \partial_m f(r, m)| + |\partial_m^2 f(r, m)| \right] \\ &= r^{-2} \partial_r (r^2 \Upsilon \partial_r f) \\ &\quad + O(\epsilon r^{-2} u_{\text{trap}}^{-1-\delta_{\text{dec}}}) \left[r^2 |\partial_r^2 f(r, m)| + r |\partial_r f(r, m)| + r |\partial_r \partial_m f(r, m)| + |\partial_m^2 f(r, m)| \right] \end{aligned}$$

as desired. \square

According to equation (10.1.13) we have,

$$\mathcal{E}[X, w](\Psi) = \frac{1}{2} \mathcal{Q} \cdot {}^{(X)}\pi - \frac{1}{2} X(V) |\Psi|^2 + \frac{1}{2} w \mathcal{L}(\Psi) - \frac{1}{4} |\Psi|^2 \square_{\mathbf{g}} w.$$

In the next proposition we choose X to be of the form $X = f(r, m)R$ and make a choice of w as a function of f .

Proposition 10.1.12. *Assume*

$$X = f(r, m)R \text{ and } w(r, m) = r^{-2}\Upsilon\partial_r(r^2f).$$

Then,

$$\mathcal{E}[X, w](\Psi) = \dot{\mathcal{E}}[X, w] + \mathcal{E}_\epsilon[X, w]$$

where, with $\widehat{\mathcal{Q}}_{34} := \mathcal{Q}_{34} - V|\Psi|^2 = |\nabla\Psi|^2$,

$$\begin{aligned} \dot{\mathcal{E}}[fR, w](\Psi) &= \frac{1}{r} \left(1 - \frac{3m}{r}\right) f \widehat{\mathcal{Q}}_{34} + \partial_r f |R(\Psi)|^2 - \frac{1}{4} r^{-2} \partial_r (r^2 \Upsilon \partial_r w) |\Psi|^2 \\ &\quad + 4\Upsilon \frac{r-4m}{r^4} f |\Psi|^2, \\ \mathcal{E}_\epsilon[fR, w](\Psi) &= \epsilon \frac{1}{2} \mathcal{Q} \cdot {}^{(X)}\ddot{\pi} + O\left(\epsilon r^{-3} u_{trap}^{-1-\delta_{dec}} (|f| + r^2 |\partial_r w| + r^3 |\partial_r^2 w| \right. \\ &\quad \left. + r^2 |\partial_r \partial_m w| + r |\partial_m^2 w|)\right) |\Psi|^2. \end{aligned} \tag{10.1.14}$$

Proof. According to Lemma 10.1.8 and equation (10.1.13) we have,

$$\mathcal{E}[X, w](\Psi) = \frac{1}{2} \mathcal{Q} \cdot ({}^{(X)}\dot{\pi} + \epsilon {}^{(X)}\ddot{\pi}) - \frac{1}{2} X(V) |\Psi|^2 + \frac{1}{2} w \mathcal{L}(\Psi) - \frac{1}{4} |\Psi|^2 \square_{\mathbf{g}} w.$$

Hence, in view of lemmas 10.1.8 and 10.1.10,

$$\begin{aligned} \mathcal{E}[X, w](\Psi) - \epsilon \frac{1}{2} \mathcal{Q} \cdot {}^{(X)}\ddot{\pi} &= \frac{1}{2} \mathcal{Q} \cdot {}^{(X)}\dot{\pi} - \frac{1}{2} X(V) |\Psi|^2 + \frac{1}{2} w \mathcal{L}(\Psi) - \frac{1}{4} |\Psi|^2 \square_{\mathbf{g}} w \\ &= f \left(-\frac{m}{r^2} + \frac{\Upsilon}{r} \right) |\nabla\Psi|^2 + \partial_r f |R\Psi|^2 - \left(\frac{\Upsilon}{r} f + \frac{1}{2} \Upsilon \partial_r f \right) \mathcal{L}(\Psi) \\ &\quad - \frac{m}{r^2} f V |\Psi|^2 - \frac{1}{2} X(V) |\Psi|^2 + \frac{1}{2} w \mathcal{L}(\Psi) - \frac{1}{4} |\Psi|^2 \square_{\mathbf{g}} w. \end{aligned}$$

Thus, assuming $w = r^{-2}\Upsilon\partial_r(r^2f) = \frac{2\Upsilon}{r}f + \partial_r f \Upsilon$,

$$\begin{aligned} \mathcal{E}[X, w](\Psi) - \epsilon \frac{1}{2} \mathcal{Q} \cdot {}^{(X)}\ddot{\pi} &= r^{-1} f \left(1 - \frac{3m}{r}\right) |\nabla\Psi|^2 + \partial_r f |R\Psi|^2 \\ &\quad - \left(\frac{m}{r^2} f V + \frac{1}{2} X(V) + \frac{1}{4} \square_{\mathbf{g}} w \right) |\Psi|^2. \end{aligned}$$

Note that, in view of Lemma 10.1.7,

$$X(V) = fR(V) = -8\Upsilon f \frac{r-3m}{r^4} + O(\epsilon r^{-3} u_{trap}^{-1-\delta_{dec}} |f|)$$

and,

$$\begin{aligned} \frac{m}{r^2}fV + \frac{1}{2}X(V) &= f \left(\frac{4m}{r^4}\Upsilon - 4\Upsilon \frac{r-3m}{r^4} \right) + O(\epsilon r^{-3} u_{trap}^{-1-\delta_{dec}} |f|) \\ &= -4f\Upsilon \frac{r-4m}{r^4} + O(\epsilon r^{-3} u_{trap}^{-1-\delta_{dec}} |f|). \end{aligned}$$

Note also that, in view of Lemma 10.1.11

$$\square_{\mathbf{g}}(w) = r^{-2}\partial_r(r^2\Upsilon\partial_r w) + O(\epsilon r^{-2} u_{trap}^{-1-\delta_{dec}}) \left[r^2|\partial_r^2 w| + r|\partial_r w| + r|\partial_r\partial_m w| + |\partial_m^2 w| \right].$$

Thus,

$$\begin{aligned} \frac{m}{r^2}fV + \frac{1}{2}X(V) + \frac{1}{4}\square_{\mathbf{g}}w &= -4\Upsilon \frac{r-4m}{r^4}f + \frac{1}{4}r^{-2}\partial_r(r^2\Upsilon\partial_r w) \\ &\quad + O(\epsilon r^{-3} u_{trap}^{-1-\delta_{dec}}) \left[|f| + r^3|\partial_r^2 w| + r^2|\partial_r w| \right. \\ &\quad \left. + r^2|\partial_r\partial_m w| + r|\partial_m^2 w| \right] \end{aligned}$$

and hence

$$\begin{aligned} \mathcal{E}[X, w](\Psi) - \epsilon \frac{1}{2} \mathcal{Q} \cdot (X) \ddot{\pi} &= r^{-1}f \left(1 - \frac{3m}{r} \right) |\nabla\Psi|^2 + \partial_r f |R\Psi|^2 - \frac{1}{4}|\Psi|^2 r^{-2}\partial_r(r^2\Upsilon\partial_r w) \\ &\quad + 4\Upsilon \frac{r-4m}{r^4} f |\Psi|^2 \\ &\quad + O\left(\epsilon r^{-3} u_{trap}^{-1-\delta_{dec}} (|f| + r^2|\partial_r w| + r^3|\partial_r^2 w| \right. \\ &\quad \left. + r^2|\partial_r\partial_m w| + r|\partial_m^2 w|) \right) |\Psi|^2 \end{aligned}$$

as desired. \square

10.1.7 A first estimate

We concentrate our attention on the principal term

$$\dot{\mathcal{E}}[fR, w](\Psi) = \frac{1}{r} \left(1 - \frac{3m}{r} \right) f \widehat{\mathcal{Q}}_{34} + \partial_r f |R(\Psi)|^2 - \frac{1}{4} r^{-2} \partial_r (r^2 \Upsilon \partial_r w) |\Psi|^2 + 4\Upsilon \frac{r-4m}{r^4} f |\Psi|^2$$

and choose $f = f(r, m)$ such that the right hand side is positive definite.

Consider the quadratic forms,

$$\begin{aligned} \dot{\mathcal{E}}_0(\Psi) &:= A \widehat{\mathcal{Q}}_{34} + B |R\Psi|^2 + r^{-2} W |\Psi|^2, \\ \dot{\mathcal{E}}(\Psi) &:= \dot{\mathcal{E}}_0(\Psi) + 4\Upsilon \frac{r-4m}{r^4} f |\Psi|^2, \end{aligned} \tag{10.1.15}$$

with the coefficients

$$A := r^{-1}f \left(1 - \frac{3m}{r}\right), \quad B := \partial_r f, \quad W := -\frac{1}{4}\partial_r(r^2\Upsilon\partial_r w). \quad (10.1.16)$$

The goal is to show that there exist choices of f, w verifying the condition of Proposition 10.1.12, i.e. $w = r^{-2}\partial_r(r^2 f)$, which makes $\mathcal{E}(\Psi)$ positive definite, for all smooth S -valued tensorfields Ψ defined in the region $r \geq 2m_0(1 - \delta_{\mathcal{H}})$, which decay reasonable fast at infinity. We look first for choices of f, w such that the coefficient A, B, W are non-negative. Note in particular that f must be increasing as a function of r and $f = 0$ on $r = 3m$. Following J. Stogin [53] we choose w first to ensure that W is non-negative and then choose f , compatible with the equation,

$$\partial_r(r^2 f) = \frac{r^2}{\Upsilon}w, \quad f = 0 \text{ on } r = 3m. \quad (10.1.17)$$

To ensure that $A = r^{-2}f(r - 3m)$ is positive we need a non-negative w which verifies (modulo error terms⁴) $W = -\frac{1}{4}\partial_r(r^2\Upsilon\partial_r w) \geq 0$. It is more difficult to choose w such that $B = \partial_r f$ is also non-negative.

Stogin defines w based on the following lemma.

Lemma 10.1.13. *The scalar function w defined by*

$$w(r, m) = \begin{cases} \frac{1}{4m}, & \text{if } r \leq 4m, \\ \frac{2\Upsilon}{r}, & \text{if } r \geq 4m, \end{cases}$$

is C^1 , non-negative and such that $W = -\frac{1}{4}\partial_r(r^2\Upsilon\partial_r w)$ verifies,

$$W(r, m) = \begin{cases} 0, & \text{if } r < 4m, \\ \frac{m}{r^2} \left(3 - \frac{8m}{r}\right), & \text{if } r > 4m. \end{cases} \quad (10.1.18)$$

Proof. For $r \geq 4m$, we have

$$w(r, m) = \frac{2\Upsilon}{r}, \quad \partial_r w(r, m) = -\frac{2}{r^2} + \frac{8m}{r^3}, \quad \partial_r^2 w(r, m) = \frac{4}{r^3} - \frac{24m}{r^4}.$$

In particular, we have

$$w = \frac{1}{4m}, \quad \partial_r w = 0 \quad \text{at } r = 4m$$

⁴i.e. terms which vanish in Schwarzschild.

so that w is indeed C^1 . Furthermore, we also have

$$\begin{aligned} r^{-2}\partial_r(r^2\Upsilon\partial_r w) &= \Upsilon\partial_r^2 w(r) + \partial_r w(r) \left(\frac{2}{r} - \frac{2m}{r^2}\right) \\ &= \Upsilon \left(\frac{4}{r^3} - \frac{24m}{r^4}\right) + \left(-\frac{2}{r^2} + \frac{8m}{r^3}\right) \left(\frac{2}{r} - \frac{2m}{r^2}\right) \\ &= -\frac{4m}{r^4} \left(3 - \frac{8m}{r}\right) \end{aligned}$$

so that, for $r \geq 4m$,

$$W = -\frac{1}{4}\partial_r(r^2\Upsilon\partial_r w) = \frac{m}{r^2} \left(3 - \frac{8m}{r}\right)$$

as desired. □

Once w is defined we can evaluate f as follows.

Lemma 10.1.14. *Let $w(r, m)$ defined as in Lemma 10.1.13. Then, the function $f(r, m)$ given by,*

$$r^2 f(r, m) := \begin{cases} 2m^2 \log\left(\frac{r-2m}{m}\right) + (r-3m)\frac{r^2+6mr+30m^2}{12m}, & \text{for } r \leq 4m, \\ C_* m^2 + r^2 - (4m)^2, & \text{for } r \geq 4m, \end{cases} \quad (10.1.19)$$

with the constant C_* given by⁵

$$C_* := 2 \log(2) + \frac{35}{6}, \quad C_* \sim 7.22,$$

is C^2 and satisfies (10.1.17), i.e. we have

$$\partial_r(r^2 f) = r^2 \Upsilon^{-1} w, \quad f = 0 \text{ on } r = 3m.$$

Proof. By direct check⁶, we have for $r \leq 4m$

$$\begin{aligned} \partial_r(r^2 f)(r, m) &= \frac{2m^2}{(r-2m)} + \frac{r^2 + 6mr + 30m^2}{12m} + (r-3m)\frac{2r+6m}{12m} \\ &= \frac{r^3}{4m(r-2m)} \\ &= \frac{r^2}{\Upsilon} \frac{1}{4m} \end{aligned}$$

⁵ C_* is chosen so that f is continuous across $r = 4m$.

⁶Recall from Remark 10.1.6 that $\partial_r f$ does not denote a spacetime coordinate vectorfield applied to f , but instead the partial derivative with respect to the variable r of the function $f(r, m)$.

and for $r \geq 4m$

$$\partial_r(r^2 f)(r, m) = 2r,$$

as well as $f = 0$ on $r = 3m$ so that, in view of the definition of $w(r)$ in Lemma 10.1.13, we infer

$$\partial_r(r^2 f) = r^2 \Upsilon^{-1} w, \quad f = 0 \text{ on } r = 3m$$

as desired. Note also that w being C^1 , f is thus indeed C^2 . \square

Next, we derive a lower bound on $\partial_r f$ for $r \leq 4m$.

Lemma 10.1.15. *We have for all r and m*

$$r^3 \partial_r f \geq 16m^2.$$

Also, there exists a constant $C > 0$ such that for all r and m

$$\left(1 - \frac{3m}{r}\right) f \geq C^{-1} \left(1 - \frac{3m}{r}\right)^2.$$

Proof. We have

$$\partial_r(r^3 \partial_r f) = \partial_r(r \partial_r(r^2 f)) - 2\partial_r(r^2 f).$$

Using the identity $\partial_r(r^2 f) = r^2 \Upsilon^{-1} w$, we infer

$$\partial_r(r^3 \partial_r f) = \partial_r(r^3 \Upsilon^{-1} w) - 2r^2 \Upsilon^{-1} w.$$

For $r \leq 4m$, we have $w = (4m)^{-1}$ and hence

$$\begin{aligned} \partial_r(r^3 \partial_r f) &= \frac{1}{4m} (\partial_r(r^3 \Upsilon^{-1}) - 2r^2 \Upsilon^{-1}) \\ &= \frac{1}{4m} r^3 \partial_r(\Upsilon^{-1}) \\ &= -\frac{r}{2\Upsilon^2} \end{aligned}$$

In particular, $r^3 \partial_r f$ is decreasing in r on $r \leq 4m$ and hence

$$r^3 \partial_r f \geq (4m)^3 \partial_r f(r = 4m, m) \text{ on } r \leq 4m.$$

On the other hand, we have, in view of the definition 10.1.19 of f

$$\begin{aligned}\partial_r(r^2 f)(r = 4m, m) &= (4m)^2 \partial_r f(r = 4m, m) + 8m f(r = 4m, m) \\ &= (4m)^2 \partial_r f(r = 4m, m) + \frac{m}{2} C_*$$

and hence

$$(4m)^2 \partial_r f(r = 4m, m) = \left(8 - \frac{C_*}{2}\right) m$$

so that

$$r^3 \partial_r f \geq 2(16 - C_*)m^2 \text{ on } r \leq 4m.$$

Since $C_* \sim 7.22 < 8$, we deduce

$$r^3 \partial_r f \geq 16m^2 \text{ on } r \leq 4m.$$

Also, for $r \geq 4m$, we have

$$f = 1 - \frac{(16 - C_*)m^2}{r^2}$$

so that

$$\partial_r f = \frac{2(16 - C_*)m^2}{r^3}.$$

Since $C_* \sim 7.22 < 8$, we deduce

$$r^3 \partial_r f \geq 16m^2 \text{ on } r \geq 4m$$

which together with the case $r \leq 4m$ above yields for all r and m the desired estimate for $\partial_r f$

$$r^3 \partial_r f \geq 16m^2.$$

In particular, $\partial_r f > 0$ and hence is strictly increasing. On the other hand, $f = 0$ on $r = 3$ and converges to 1 as $r \rightarrow +\infty$. We deduce the existence of a constant $C > 0$ such that

$$\left(1 - \frac{3m}{r}\right) f \geq C^{-1} \left(1 - \frac{3m}{r}\right)^2$$

as desired. □

We summarize the results in the following.

Proposition 10.1.16. *There exist functions $f \in C^2, w \in C^1$ verifying the relation $w = r^{-2}\Upsilon\partial_r(r^2f)$ and such that,*

$$r^2f = \begin{cases} 2m^2 \log\left(\frac{r-2m}{m}\right) + (r-3m)\frac{r^2+6mr+30m^2}{12m}, & \text{for } r \leq 4m, \\ C_*m^2 + r^2 - (4m)^2, & \text{for } r \geq 4m, \end{cases} \quad (10.1.20)$$

where C_* is a constant satisfying $7 < C_* < 8$. In particular,

$$f = \begin{cases} \frac{2m^2}{r^2} \log\left(\frac{r-2m}{m}\right) + O\left(\frac{r-3m}{m}\right), & \text{for } r \leq 4m, \\ 1 + O\left(\frac{m^2}{r^2}\right), & \text{for } r \geq 4m, \end{cases} \quad (10.1.21)$$

and, for some $C > 0$ and all $r \geq 2m$

$$\left(1 - \frac{3m}{r}\right) f \geq C^{-1} \left(1 - \frac{3m}{r}\right)^2, \quad \partial_r f \geq \frac{16m^2}{r^3}. \quad (10.1.22)$$

Also, w is given by

$$w = \begin{cases} \frac{1}{4m}, & \text{for } r \leq 4m, \\ \frac{2}{r} \left(1 - \frac{2m}{r}\right), & \text{for } r \geq 4m. \end{cases} \quad (10.1.23)$$

Moreover $W = -\frac{1}{4}\partial_r(r^2\Upsilon\partial_r w)$ verifies,

$$W = \begin{cases} 0, & \text{if } r < 4m, \\ \frac{m}{r^2} \left(3 - \frac{8m}{r}\right), & \text{if } r > 4m, \end{cases} \quad (10.1.24)$$

and,

$$\begin{aligned} \dot{\mathcal{E}}_0[fR, w](\Psi) &= \partial_r f |R(\Psi)|^2 + r^{-2}W|\Psi|^2 + r^{-1} \left(1 - \frac{3m}{r}\right) f \widehat{\mathcal{Q}}_{34}, \\ \dot{\mathcal{E}}[fR, w](\Psi) &= \mathcal{E}_0[fR, w](\Psi) + 4\Upsilon \frac{r-4m}{r^4} f |\Psi|^2. \end{aligned} \quad (10.1.25)$$

Recall also that,

$$\widehat{\mathcal{Q}}_{34} = |\nabla\Psi|^2.$$

Remark 10.1.17. *The estimates obtained so far have two major deficiencies*

1. *The quadratic form $\dot{\mathcal{E}}_0[fR, w](\Psi) + 4\Upsilon \frac{r-4m}{r^4} f |\Psi|^2$ fails to be positive definite in the region $3m \leq r \leq 4m$ because of the potential term $\Upsilon \frac{r-4m}{r^4} f |\Psi|^2$.*
2. *The function f blows up logarithmically at $r = 2m$ in $^{(int)}\mathcal{M}$.*

In the next section we deal with the first issue. We handle the second problem in the following two sections.

10.1.8 Improved lower bound in $^{(ext)}\mathcal{M}$

Note that the term $4f\Upsilon\frac{r-4m}{r^4}$ is negative for $3m \leq r \leq 4m$ and positive everywhere else. An improvement can be obtained by using the following Poincaré inequality.

Lemma 10.1.18. *We have for $\Psi \in \mathcal{S}_2(\mathcal{M})$,*

$$\int_S |\nabla\Psi|^2 \geq 2r^{-2} \left(1 - O(\epsilon)\right) \int_S \Psi^2 da_S. \quad (10.1.26)$$

Proof. See Proposition 2.1.32. □

According to Proposition 10.1.16 we deduce,

$$\begin{aligned} \int_S \dot{\mathcal{E}}[fR, w](\Psi) &\geq \int_S \dot{\mathcal{E}}_1 - O(\epsilon r^{-3}) \int_S \Psi^2 da_S, \\ \dot{\mathcal{E}}_1 &:= \partial_r f |R(\Psi)|^2 + r^{-2} W |\Psi|^2 + 2r^{-3} \left(1 - \frac{3m}{r}\right) f |\Psi|^2 + 4\Upsilon \frac{r-4m}{r^4} f |\Psi|^2, \end{aligned} \quad (10.1.27)$$

with W defined in (10.1.24). It is easy to see however that $\dot{\mathcal{E}}_1$ still fails to be positive for $3m < r < 4m$. To achieve positivity we also need to modify the original energy density $\mathcal{E}[fR, w](\Psi)$ by considering instead the modified energy density $\mathcal{E}[fR, w, M](\Psi)$ (see (10.1.11) and notation (10.1.12)) with $M = 2hR$ for a function $h = h(r, m)$ supported for $r \geq 3m$ and constant for $r \geq 4m$.

$$\begin{aligned} \mathcal{E}[fR, w, M](\Psi) &= \mathcal{E}[fR, w](\Psi) + \frac{1}{4} \mathbf{D}^\mu (|\Psi|^2 M_\mu) = \mathcal{E}[fR, w](\Psi) + \frac{1}{4} (\mathbf{D}^\mu M_\mu) |\Psi|^2 + \frac{1}{2} \Psi M(\Psi) \\ &= \mathcal{E}[fR, w](\Psi) + \frac{1}{2} \mathbf{D}^\mu (hR_\mu) |\Psi|^2 + h\Psi R(\Psi). \end{aligned}$$

To take into account the additional terms in the modified $\mathcal{E}[fR, w, M](\Psi)$ we first derive the following.

Lemma 10.1.19. *Let $h(r, m)$ a C^1 function of r and m . We have,*

$$\mathbf{D}^\mu (hR_\mu) = r^{-2} \partial_r (\Upsilon r^2 h) + O\left(\epsilon r^{-1} u_{trap}^{-1-\delta_{dec}} (r|\partial_r h| + |h| + r|\partial_m h|)\right). \quad (10.1.28)$$

Proof. In view of Lemma 10.1.7, which computes the components of $^{(R)}\pi$, as well as

Lemma 10.1.5 to compute $R(h)$, we calculate,

$$\begin{aligned} \mathbf{D}^\mu(hR_\mu) &= R(h) + h(\mathbf{D}^\mu R_\mu) = \frac{1}{2}(e_4(h) - \Upsilon e_3(h)) + h\frac{1}{2}\text{tr}({}^{(R)}\pi) \\ &= \frac{1}{2}(e_4(r) - \Upsilon e_3(r))\partial_r h + O(\epsilon u_{\text{trap}}^{-1-\delta_{dec}}|\partial_m h|) + \frac{1}{2}h(-{}^{(R)}\pi_{34} + {}^{(R)}\pi_{\theta\theta} + {}^{(R)}\pi_{\varphi\varphi}) \\ &= \Upsilon\partial_r h + \frac{1}{2}\left(\frac{4m}{r^2} + 4\frac{\Upsilon}{r}\right)h + O\left(\epsilon u_{\text{trap}}^{-1-\delta_{dec}}(|\partial_r h| + r^{-1}|h| + |\partial_m h|)\right) \\ &= r^{-2}\partial_r(\Upsilon r^2 h) + O\left(\epsilon u_{\text{trap}}^{-1-\delta_{dec}}(|\partial_r h| + r^{-1}|h| + |\partial_m h|)\right) \end{aligned}$$

as desired. □

In view of the lemma we write,

$$\begin{aligned} \mathcal{E}[fR, w, 2hR](\Psi) &= \dot{\mathcal{E}}[fR, w, 2hR](\Psi) + \mathcal{E}_\epsilon[fR, w, 2hR](\Psi), \\ \dot{\mathcal{E}}[fR, w, 2hR](\Psi) &:= \dot{\mathcal{E}}[fR, w](\Psi) + \frac{1}{2}r^{-2}\partial_r(\Upsilon r^2 h)|\Psi|^2 + h\Psi R(\Psi), \\ \mathcal{E}_\epsilon[fR, w, 2hR](\Psi) &:= \mathcal{E}_\epsilon[fR, w](\Psi) + O\left(r^{-1}\epsilon u_{\text{trap}}^{-1-\delta_{dec}}(r|\partial_r h| + |h| + r|\partial_m h|)\right)|\Psi|^2. \end{aligned} \tag{10.1.29}$$

The main result of this section is stated below.

Proposition 10.1.20. *There exists a function $h = h(r, m)$ with bounded derivative h' , supported in $r \geq 3m$ such that $h = O(r^{-2}), h' = O(r^{-3})$ for $r \geq 4m$ such that,*

$$\begin{aligned} \mathcal{E}[fR, w, 2hR](\Psi) &= \dot{\mathcal{E}}[fR, w, 2hR](\Psi) + \mathcal{E}_\epsilon[fR, w, 2hR](\Psi), \\ \mathcal{E}_\epsilon[fR, w, 2hR](\Psi) &= \epsilon\frac{1}{2}\mathcal{Q} \cdot {}^{(X)}\ddot{\pi} + O\left(\epsilon r^{-3}u_{\text{trap}}^{-1-\delta_{dec}}(|f| + 1)\right)|\Psi|^2, \end{aligned} \tag{10.1.30}$$

and, for sufficiently large universal constant $C > 0$, in the region $r \geq \frac{5m}{2}$,

$$\int_S \dot{\mathcal{E}}[fR, w, 2hR](\Psi) \geq C^{-1} \int_S \left(\frac{m^2}{r^3}|R(\Psi)|^2 + r^{-1}\left(1 - \frac{3m}{r}\right)^2 |\nabla\Psi|^2 + \frac{m}{r^4}|\Psi|^2 \right). \tag{10.1.31}$$

Proof. We first derive the weaker inequality,

$$\int_S \dot{\mathcal{E}}[fR, w, 2hR](\Psi) \geq C^{-1} \int_S \left(\frac{m^2}{r^3}|R(\Psi)|^2 + \frac{m}{r^4}|\Psi|^2 \right) \quad \text{on } r \geq \frac{5m}{2}$$

by making full use of the Poincaré inequality above, i.e.,

$$\int_S r^{-1} \left(1 - \frac{3m}{r}\right) f(r, m) |\nabla \Psi|^2 \geq \int_S (2 - O(\epsilon)) r^{-3} \left(1 - \frac{3m}{r}\right) f(r, m) |\Psi|^2.$$

The result will the easily follow by writing instead, with a sufficiently small $\mu > 0$,

$$\begin{aligned} & \int_S r^{-1} \left(1 - \frac{3m}{r}\right) f(r, m) |\nabla \Psi|^2 \\ &= \mu \int_S r^{-1} \left(1 - \frac{3m}{r}\right) f(r, m) |\nabla \Psi|^2 + (1 - \mu) \int_S r^{-1} \left(1 - \frac{3m}{r}\right) f(r, m) |\nabla \Psi|^2 \\ &\geq \mu \int_S r^{-1} \left(1 - \frac{3m}{r}\right) f(r, m) |\nabla \Psi|^2 + (1 - \mu) \int_S 2r^{-3} \left(1 - \frac{3m}{r}\right) f(r, m) |\Psi|^2 \end{aligned}$$

and then proceeding exactly as below.

We start with,

$$\dot{\mathcal{E}}[fR, w, 2hR](\Psi) = \dot{\mathcal{E}}[fR, w](\Psi) + \frac{1}{2} r^{-2} \partial_r (\Upsilon r^2 h) |\Psi|^2 + h \Psi R(\Psi).$$

Recalling the definition of $\dot{\mathcal{E}}_1$ in (10.1.27),

$$\dot{\mathcal{E}}_1 := \partial_r f |R(\Psi)|^2 + r^{-2} W |\Psi|^2 + 2r^{-3} \left(1 - \frac{3m}{r}\right) f |\Psi|^2 + 4\Upsilon \frac{r - 4m}{r^4} f |\Psi|^2$$

and setting,

$$\begin{aligned} \dot{\mathcal{E}}_2 &:= \dot{\mathcal{E}}_1 + \frac{1}{2} r^{-2} (\Upsilon r^2 h)' |\Psi|^2 + h \Psi R(\Psi) & (10.1.32) \\ &= \partial_r f |R(\Psi)|^2 + 2r^{-3} \left(1 - \frac{3m}{r}\right) f |\Psi|^2 + 4\Upsilon \frac{r - 4m}{r^4} f |\Psi|^2 + r^{-2} W |\Psi|^2 \\ &\quad + \frac{1}{2} r^{-2} (\Upsilon r^2 h)' |\Psi|^2 + h \Psi R(\Psi) \end{aligned}$$

we deduce, from (10.1.27)

$$\int_S \dot{\mathcal{E}}[fR, w, 2hR](\Psi) \geq \int_S \dot{\mathcal{E}}_2 - O(\epsilon r^{-3}) \int_S |\Psi|^2.$$

We now substitute,

$$h = 4\Upsilon r^{-4} \tilde{h}.$$

Hence,

$$\begin{aligned} \frac{1}{2}r^{-2}\partial_r(\Upsilon r^2h)|\Psi|^2 + h\Psi R(\Psi) &= \frac{1}{2}r^{-2}\partial_r(4\Upsilon^2r^{-2}\tilde{h})|\Psi|^2 + 4\Upsilon r^{-4}\tilde{h}\Psi R(\Psi) \\ &= \frac{1}{2}r^{-2}\partial_r(4\Upsilon^2r^{-2})\tilde{h}|\Psi|^2 + 2r^{-4}\Upsilon^2\partial_r\tilde{h}|\Psi|^2 + 4\Upsilon r^{-4}\tilde{h}\Psi R(\Psi) \end{aligned}$$

or, since $\frac{1}{2}r^{-2}\partial_r(4\Upsilon^2r^{-2}) = -4r^{-2}\Upsilon\frac{r-4m}{r^4}$,

$$\frac{1}{2}r^{-2}\partial_r(\Upsilon r^2h)|\Psi|^2 + h\Psi R(\Psi) = -4r^{-2}\Upsilon\frac{r-4m}{r^4}\tilde{h}|\Psi|^2 + 2r^{-4}\Upsilon^2\partial_r\tilde{h}|\Psi|^2 + 4\Upsilon r^{-4}\tilde{h}\Psi R(\Psi).$$

Thus we have,

$$\begin{aligned} \dot{\mathcal{E}}_2 &= \partial_r f |R(\Psi)|^2 + 2r^{-3} \left(1 - \frac{3m}{r}\right) f |\Psi|^2 + 4\Upsilon \frac{r-4m}{r^4} (f - r^{-2}\tilde{h}) |\Psi|^2 \\ &\quad + 2r^{-4}\Upsilon^2\partial_r\tilde{h}|\Psi|^2 + 4\Upsilon r^{-4}\tilde{h}\Psi R(\Psi) + r^{-2}W|\Psi|^2. \end{aligned}$$

We also express,

$$4\Upsilon r^{-4}\tilde{h}\Psi R(\Psi) = \frac{2\tilde{h}}{r^3}(R(\Psi) + \Upsilon r^{-1}\Psi)^2 - \frac{2\tilde{h}}{r^3}|R(\Psi)|^2 - \frac{2\tilde{h}}{r^5}\Upsilon^2|\Psi|^2$$

and therefore,

$$\begin{aligned} \dot{\mathcal{E}}_2 &= (\partial_r f - 2r^{-3}\tilde{h})|R(\Psi)|^2 + \frac{2\tilde{h}}{r^3}(R(\Psi) + \Upsilon r^{-1}\Psi)^2 + r^{-2}W|\Psi|^2 \\ &\quad + \left[2r^{-3} \left(1 - \frac{3m}{r}\right) f + 4\Upsilon \frac{r-4m}{r^4} (f - r^{-2}\tilde{h}) + 2r^{-4}\Upsilon\partial_r\tilde{h} - 2r^{-5}\Upsilon^2\tilde{h} \right] |\Psi|^2. \end{aligned}$$

We choose $\tilde{h}(r, m)$ as the following continuous and piecewise C^1 function,

$$\tilde{h} = \begin{cases} 0, & r \leq \frac{5m}{2}, \\ \delta_{\tilde{h}} \left(\frac{5m}{2} - r\right), & \frac{5m}{2} \leq r \leq \frac{11m}{4}, \\ \delta_{\tilde{h}}(r - 3m), & \frac{11m}{4} \leq r \leq 3m, \\ r^2 f, & 3m \leq r \leq 4m, \\ (4m)^2 f(4m, m), & r \geq 4m. \end{cases}$$

where the constant $\delta_{\tilde{h}} > 0$ will be chosen small enough. We consider the following cases:

Case 1 ($\frac{5m}{2} \leq r \leq 3m$). In view of the definition of \tilde{h} and since $W = 0$, we deduce,

$$\begin{aligned} \dot{\mathcal{E}}_2 &= \partial_r f |R(\Psi)|^2 + \left[2r^{-3} \left(1 - \frac{3m}{r} \right) f + 4\Upsilon \frac{r-4m}{r^4} (f - r^{-2}\tilde{h}) \right. \\ &\quad \left. + 2\delta_{\tilde{h}} r^{-4} \Upsilon^2 \left(1_{\frac{11m}{4} \leq r \leq 3m} - 1_{\frac{5m}{2} \leq r \leq \frac{11m}{4}} \right) \right] |\Psi|^2 + \delta_{\tilde{h}} O(1) \Psi R(\Psi) 1_{\frac{5m}{2} \leq r \leq 3m}. \end{aligned}$$

In view of (10.1.22), we may assume, choosing for $\delta_{\tilde{h}} > 0$ small enough, that

$$f - \tilde{h} \leq -\frac{1}{2}|f| \text{ on } r \leq 3m. \quad (10.1.33)$$

We infer, using also that $f < 0$ on $r \leq 3m$,

$$\begin{aligned} \dot{\mathcal{E}}_2 &\geq \partial_r f |R(\Psi)|^2 + \left[2r^{-3} \left(1 - \frac{3m}{r} \right) f + 2\Upsilon \frac{r-4m}{r^4} f \right. \\ &\quad \left. + 2\delta_{\tilde{h}} r^{-4} \Upsilon^2 \left(1_{\frac{11m}{4} \leq r \leq 3m} - 1_{\frac{5m}{2} \leq r \leq \frac{11m}{4}} \right) \right] |\Psi|^2 + \delta_{\tilde{h}} O(1) \Psi R(\Psi) 1_{\frac{5m}{2} \leq r \leq 3m}. \end{aligned}$$

Since we have

$$\partial_r f \gtrsim 1, \quad 2r^{-3} \left(1 - \frac{3m}{r} \right) + 4\Upsilon \frac{r-4m}{r^4} \lesssim -1, \quad f \lesssim - \left| 1 - \frac{3m}{r} \right| \text{ on } r \leq 3m,$$

where we have used in particular Lemma 10.1.15 and Proposition 10.1.16, we infer

$$\begin{aligned} \dot{\mathcal{E}}_2 &\gtrsim |R(\Psi)|^2 + \left(\left| 1 - \frac{3m}{r} \right| + \delta_{\tilde{h}} 1_{\frac{11m}{4} \leq r \leq 3m} - O(1) \delta_{\tilde{h}} 1_{\frac{5m}{2} \leq r \leq \frac{11m}{4}} \right) |\Psi|^2 \\ &\quad - \delta_{\tilde{h}} O(1) \Psi R(\Psi) 1_{\frac{5m}{2} \leq r \leq 3m} \\ &\geq \frac{1}{2} |R(\Psi)|^2 + \left(\left| 1 - \frac{3m}{r} \right| + \delta_{\tilde{h}} \left(1 - O(1) \delta_{\tilde{h}} \right) 1_{\frac{11m}{4} \leq r \leq 3m} - O(1) \delta_{\tilde{h}} 1_{\frac{5m}{2} \leq r \leq \frac{11m}{4}} \right) |\Psi|^2. \end{aligned}$$

Thus, for $\delta_{\tilde{h}} > 0$ small enough, there exists some large $C > 0$ such that

$$\mathcal{E}_2 \geq C^{-1} [|R(\Psi)|^2 + |\Psi|^2] \text{ on } \frac{5m}{2} \leq r \leq 3m. \quad (10.1.34)$$

Case 2 ($3m \leq r \leq 4m$). Since $\tilde{h} = r^2 f$ and $W = 0$, using in particular $\tilde{h} \geq 0$ on $3m \leq r \leq 4m$, we deduce,

$$\begin{aligned} \dot{\mathcal{E}}_2 &\geq (\partial_r f - 2r^{-3}(r^2 f)) |R(\Psi)|^2 + \left[2r^{-3} \left(1 - \frac{3m}{r} \right) f + 2r^{-4} \Upsilon \partial_r (r^2 f) - 2r^{-5} \Upsilon^2 (r^2 f) \right] |\Psi|^2 \\ &= (\partial_r f - 2r^{-1} f) |R(\Psi)|^2 + \left[2r^{-3} \left(1 - \frac{3m}{r} \right) f + 2r^{-4} \Upsilon^2 (2r f + r^2 \partial_r f) - 2r^{-3} \Upsilon^2 f \right] |\Psi|^2 \\ &= (\partial_r f - 2r^{-1} f) |R(\Psi)|^2 + \left[2r^{-3} \left(1 - \frac{3m}{r} \right) f + 2r^{-2} \Upsilon^2 \partial_r f + 2r^{-3} \Upsilon^2 f \right] |\Psi|^2. \end{aligned}$$

Note that the second term is strictly positive. It remains to analyze the first term.

Lemma 10.1.21. *In the interval $[3m, 4m]$ we have,*

$$\partial_r f - 2r^{-1}f > 0.$$

Proof. Recall from Proposition 10.1.16 that $w = r^{-2}\Upsilon\partial_r(r^2f) = \frac{1}{4m}$ in the interval $[3m, 4m]$. Using also $f = 0$ on $r = 3m$, we deduce

$$\partial_r(r^2f) = \frac{r^2}{\Upsilon} \frac{1}{4m}.$$

We compute

$$\begin{aligned} \partial_r \left(r^2f - \frac{(r-3m)r^2}{4m\Upsilon} \right) &= -\frac{(r-3m)}{4m} \partial_r \left(\frac{r^2}{\Upsilon} \right) \\ &= -\frac{(r-3m)(r-4m)}{2m\Upsilon^2} \\ &\leq 0 \quad \text{on } 3m \leq r \leq 4m, \end{aligned}$$

so that the differentiated quantity decays in r on $[3m, 4m]$. Since it vanishes on $r = 3m$, we infer

$$f \leq \frac{(r-3m)}{4m\Upsilon} \quad \text{on } 3m \leq r \leq 4m.$$

Thus, we deduce, using again $\partial_r(r^2f) = \frac{r^2}{\Upsilon} \frac{1}{4m}$,

$$\begin{aligned} \partial_r f - \frac{2}{r}f &= r^{-2} \left(\partial_r(r^2f) - 4rf \right) \\ &= \frac{1}{4m\Upsilon} - \frac{4}{r}f \\ &\geq \frac{1}{4m\Upsilon} - \frac{(r-3m)}{rm\Upsilon} \\ &\geq \frac{1}{4m\Upsilon} \left(1 - 4 \left(1 - \frac{3m}{r} \right) \right) \\ &> 0 \quad \text{on } 3m \leq r < 4m. \end{aligned}$$

On the other hand, we have by direct check at $r = 4m$, using (10.1.20),

$$\left(\partial_r f - \frac{2}{r}f \right)_{r=4m} = \frac{1}{2m} - \frac{1}{m}f_{r=4m} = \frac{1}{2m} \left(1 - \frac{C_*}{8} \right) > 0$$

since $C_* < 8$. Hence, we infer

$$\partial_r f - 2r^{-1}f > 0 \quad \text{on } 3m \leq r \leq 4m$$

as desired. □

We thus conclude, for some $C > 0$, in the interval $[3m, 4m]$

$$\dot{\mathcal{E}}_2 \geq C^{-1} [|R(\Psi)|^2 + |\Psi|^2]. \quad (10.1.35)$$

Case 3 ($r \geq 4m$). Since \tilde{h} is constant and positive on $r \geq 4m$, we deduce,

$$\begin{aligned} \dot{\mathcal{E}}_2 &\geq (\partial_r f - 2r^{-3}\tilde{h})|R(\Psi)|^2 \\ &+ \left[2r^{-3} \left(1 - \frac{3m}{r} \right) f + 4\Upsilon \frac{r-4m}{r^4} (f - r^{-2}\tilde{h}) - 2r^{-5}\Upsilon^2\tilde{h} + r^{-2}W \right] |\Psi|^2. \end{aligned}$$

We examine the first term. In view of the formula for f for $r \geq 4m$, see (10.1.20),

$$\partial_r f = \frac{2}{r^3}(16 - C_*)m^2, \quad \tilde{h} = (4m)^2 f(4m, m) = C_* m^2$$

and hence

$$\partial_r f - 2r^{-3}\tilde{h} = \frac{2(16 - 2C_*)m^2}{r^3}$$

and hence, since $C_* < 8$, we have

$$\partial_r f - 2r^{-3}\tilde{h} \gtrsim \frac{m^2}{r^3} \quad \text{for } r \geq 4m.$$

It remains to analyze the sign of

$$\begin{aligned} &2r^{-3} \left(1 - \frac{3m}{r} \right) f + 4\Upsilon \frac{r-4m}{r^4} (f - r^{-2}\tilde{h}) - 2r^{-5}\Upsilon^2\tilde{h} \\ &= \left[2r^{-3} \left(1 - \frac{3m}{r} \right) + 4\Upsilon \frac{r-4m}{r^4} \right] (f - r^{-2}\tilde{h}) + \left[2r^{-3} \left(1 - \frac{3m}{r} \right) - 2r^{-3}\Upsilon^2 \right] r^{-2}\tilde{h}. \end{aligned}$$

The first term, which can be written in the form,

$$\left[2r^{-3} \left(1 - \frac{3m}{r} \right) + 4\Upsilon \frac{r-4m}{r^4} \right] r^{-2} (r^2 f(r, m) - (4m)^2 f(4m, m))$$

is manifestly positive for $r \geq 4m$. To evaluate the sign of the second term we calculate,

$$2r^{-3} \left(1 - \frac{3m}{r} \right) - 2r^{-3}\Upsilon^2 = 2mr^{-5}(r - 4m).$$

Thus, for $r \geq 4m$,

$$2r^{-3} \left(1 - \frac{3m}{r} \right) f + 4\Upsilon \frac{r-4m}{r^4} (f - r^{-2}\tilde{h}) - 2r^{-5}\Upsilon^2\tilde{h} \geq 0.$$

Also, since $W = \frac{m}{r^2} \left(3 - \frac{8m}{r}\right)$, we have

$$r^{-2}W \gtrsim \frac{1}{r^4}.$$

Thus, in view of the above, we have for some $C > 0$ and for $r \geq 4m$,

$$\dot{\mathcal{E}}_2 \geq C^{-1} \left[\frac{1}{r^3} |R(\Psi)|^2 + \frac{1}{r^4} |\Psi|^2 \right]. \tag{10.1.36}$$

Gathering (10.1.34), (10.1.35) and (10.1.36), we infer for some $C > 0$,

$$\dot{\mathcal{E}}_2 \geq C^{-1} \left[\frac{1}{r^3} |R(\Psi)|^2 + \frac{1}{r^4} |\Psi|^2 \right] \quad \text{on } r \geq \frac{5m}{2}.$$

Recalling

$$\int_S \dot{\mathcal{E}}[fR, w, 2hR](\Psi) \geq \int_S \dot{\mathcal{E}}_2 - O(\epsilon r^{-3}) \int_S |\Psi|^2,$$

we infer

$$\int_S \dot{\mathcal{E}}[fR, w, 2hR](\Psi) \geq C^{-1} \int_S \left[\frac{1}{r^3} |R(\Psi)|^2 + \frac{1}{r^4} |\Psi|^2 \right] - O(\epsilon r^{-3}) \int_S |\Psi|^2$$

and hence, for $\epsilon > 0$ small enough,

$$\int_S \dot{\mathcal{E}}[fR, w, 2hR](\Psi) \geq \frac{1}{2} C^{-1} \int_S \left[\frac{1}{r^3} |R(\Psi)|^2 + \frac{1}{r^4} |\Psi|^2 \right] \quad \text{on } r \geq \frac{5m}{2}$$

as desired.

It remains to analyze the error term,

$$\begin{aligned} \mathcal{E}_\epsilon[fR, w, 2hR](\Psi) &= \mathcal{E}_\epsilon[fR, w](\Psi) + O\left(r^{-1} \epsilon u_{trap}^{-1-\delta_{dec}} (r|\partial_r h| + |h| + r|\partial_m h|)\right) |\Psi|^2 \\ &= \epsilon \frac{1}{2} \mathcal{Q}^{(X)} \ddot{\pi} + O(\epsilon r^{-3} u_{trap}^{-1-\delta_{dec}} (|f| + r^2 |\partial_r w| + r^3 |\partial_r^2 w| \\ &\quad + r^2 |\partial_r \partial_m w| + r |\partial_m^2 w|)) |\Psi|^2 \\ &\quad + O\left(r^{-3} \epsilon u_{trap}^{-1-\delta_{dec}} (r^3 |\partial_r h| + r^2 |h| + r^3 |\partial_m h|)\right) |\Psi|^2. \end{aligned}$$

Recall that,

$$w = \begin{cases} \frac{1}{4m}, & \text{for } r \leq 4m, \\ \frac{2}{r} \left(1 - \frac{2m}{r}\right), & \text{for } r \geq 4m, \end{cases}$$

and $h = 4\Upsilon r^{-4}\tilde{h}$, with

$$\tilde{h} = \begin{cases} 0, & r \leq \frac{5m}{2}, \\ \delta_{\tilde{h}}\left(\frac{5m}{2} - r\right), & \frac{5m}{2} \leq r \leq \frac{11m}{4}, \\ \delta_{\tilde{h}}(r - 3m), & \frac{11m}{4} \leq r \leq 3m, \\ r^2 f, & 3m \leq r \leq 4m, \\ (4m)^2 f(4m, m), & r \geq 4m. \end{cases}$$

We deduce,

$$\mathcal{E}_\epsilon[fR, w, 2hR](\Psi) = \epsilon \mathcal{Q}^{(X)}\ddot{\pi} + O\left(\epsilon r^{-3} u_{trap}^{-1-\delta_{dec}}(|f| + 1)\right)$$

which concludes the proof of Proposition 10.1.20. \square

10.1.9 Cut-off Correction in $^{(int)}\mathcal{M}$

So far we have found a triplet $(X = fR, w = r^{-2}\Upsilon\partial_r(r^2 f), M = 2hR)$ with f defined in Proposition 10.1.16 and h in Proposition 10.1.20 allowing for the lower bound (10.1.31) on $\int_S \dot{\mathcal{E}}[fR, w, M](\Psi)$. The main problem which remains to be addressed is that

1. f blows up logarithmically near $r = 2m$.
2. The lower bound for $\int_S \dot{\mathcal{E}}[fR, w, 2hR](\Psi)$ does not control $e_3(\Psi)$ near $r = 2m$.

In this section, we deal with the first problem, while the second problem will be treated in section 10.1.10. To correct for the first problem, i.e. the fact that f blows up logarithmically near $r = 2m$, we have to modify our choice of f and w there. Introducing

$$u := r^2 f,$$

we have,

$$f = r^{-2}u, \quad w = r^{-2}\Upsilon\partial_r u. \quad (10.1.37)$$

Warning. The auxiliary function u introduced here, and used only in this section, has of course nothing to do with our previously defined optical function on $^{(ext)}\mathcal{M}$.

Definition 10.1.22. For a given $\widehat{\delta} > 0$ we define the following functions of (r, m)

$$\begin{aligned} u_{\widehat{\delta}} &:= -\frac{m^2}{\widehat{\delta}} F\left(-\frac{\widehat{\delta}}{m^2}u\right), & f_{\widehat{\delta}} &:= r^{-2}u_{\widehat{\delta}}, \\ w_{\widehat{\delta}} &:= r^{-2}\Upsilon\partial_r u_{\widehat{\delta}}, & W_{\widehat{\delta}} &:= -\frac{1}{4}\partial_r(r^2\Upsilon\partial_r w_{\widehat{\delta}}), \end{aligned}$$

where $F : \mathbb{R} \rightarrow \mathbb{R}$ is a fixed, increasing, smooth function such that

$$F(x) = \begin{cases} x & \text{for } x \leq 1, \\ 2 & \text{for } x \geq 3. \end{cases}$$

We now derive useful properties satisfied by $f_{\widehat{\delta}}$, $w_{\widehat{\delta}}$ and $W_{\widehat{\delta}}$.

Lemma 10.1.23. Let $f_{\widehat{\delta}}$, $w_{\widehat{\delta}}$ and $W_{\widehat{\delta}}$ introduced in definition 10.1.22. Then, $f_{\widehat{\delta}} \in C^2(r > 0)$, $w_{\widehat{\delta}} \in C^1(r > 0)$, and we have for $\widehat{\delta} > 0$ sufficiently small

$$f_{\widehat{\delta}} = f \quad w_{\widehat{\delta}} = w, \quad W_{\widehat{\delta}} = W \quad \text{for } r \geq \frac{5m}{2}.$$

Also, we have for all $r > 0$

$$r^{-1}f_{\widehat{\delta}}\left(1 - \frac{3m}{r}\right) \geq C^{-1}r^{-1}\left(1 - \frac{3m}{r}\right)^2 \tag{10.1.38}$$

and

$$\partial_r(f_{\widehat{\delta}}) \geq \frac{16m^2}{r^3}. \tag{10.1.39}$$

Proof. Note first that

$$w_{\widehat{\delta}} = r^{-2}\Upsilon\partial_r u_{\widehat{\delta}} = r^{-2}\Upsilon F'\left(-\frac{\widehat{\delta}}{m^2}u\right)\partial_r u = wF'\left(-\frac{\widehat{\delta}}{m^2}u\right).$$

In view of the definition of $u_{\widehat{\delta}}$, $f_{\widehat{\delta}}$, $w_{\widehat{\delta}}$ and $W_{\widehat{\delta}}$, we have

$$\begin{aligned} u_{\widehat{\delta}} = u, & & f_{\widehat{\delta}} = f, & & w_{\widehat{\delta}} = w, & & W_{\widehat{\delta}} = W & & \text{for } u \geq -\frac{m^2}{\widehat{\delta}}, \\ u_{\widehat{\delta}} = -\frac{2m^2}{\widehat{\delta}}, & & f_{\widehat{\delta}} = -\frac{2m^2}{\widehat{\delta}r^2}, & & w_{\widehat{\delta}} = 0, & & W_{\widehat{\delta}} = 0 & & \text{for } u \leq -\frac{3m^2}{\widehat{\delta}}. \end{aligned}$$

Also, according to (10.1.21)

$$u = \begin{cases} 2m^2 \log \frac{r-2m}{m} + O(m(r-3m)), & \text{for } r \leq 4m, \\ r^2 + O(m^2), & \text{for } r \geq 4m, \end{cases}$$

and hence, for $\widehat{\delta} > 0$ sufficiently small

$$\left\{ r \geq 2m + e^{-\frac{1}{3\widehat{\delta}}} \right\} \cup \left\{ u \geq -\frac{m^2}{\widehat{\delta}} \right\}, \quad \left\{ r \leq 2m + e^{-\frac{2}{\widehat{\delta}}} \right\} \subset \left\{ u \leq -\frac{3m^2}{\widehat{\delta}} \right\}.$$

This yields

$$f_{\widehat{\delta}} = f \quad w_{\widehat{\delta}} = w, \quad W_{\widehat{\delta}} = W \quad \text{for } r \geq \frac{5m}{2}.$$

Also, we have

$$f_{\widehat{\delta}} = \begin{cases} -\frac{2m^2}{\widehat{\delta}r^2}, & \text{for } r \leq 2m + e^{-\frac{2}{\widehat{\delta}}}, \\ f, & \text{for } r \geq 2m + e^{-\frac{1}{3\widehat{\delta}}}, \end{cases}$$

and

$$f_{\widehat{\delta}} \gtrsim \frac{1}{\widehat{\delta}} \quad \text{on} \quad 2m + e^{-\frac{2}{\widehat{\delta}}} \leq r \leq 2m + e^{-\frac{1}{3\widehat{\delta}}},$$

and thus, there exists $C > 0$ such that, for all $r > 0$,

$$r^{-1} f_{\widehat{\delta}} \left(1 - \frac{3m}{r} \right) \geq C^{-1} r^{-1} \left(1 - \frac{3m}{r} \right)^2$$

which is (10.1.38).

For $u \leq -\frac{3m^2}{\widehat{\delta}}$,

$$\partial_r(f_{\widehat{\delta}}) = \partial_r(r^{-2}u_{\widehat{\delta}}) = -2r^{-3}u_{\widehat{\delta}} + r^{-2}\partial_r(u_{\widehat{\delta}}) = \frac{4m^2}{\widehat{\delta}}r^{-3}.$$

For $-\frac{3m^2}{\widehat{\delta}} \leq u \leq -\frac{m^2}{\widehat{\delta}}$

$$\begin{aligned} \partial_r(f_{\widehat{\delta}}) &= \partial_r(r^{-2}u_{\widehat{\delta}}) = -2r^{-3}u_{\widehat{\delta}} + r^{-2}\partial_r(u_{\widehat{\delta}}) \\ &= -2r^{-3}u_{\widehat{\delta}} + r^{-2}F' \left(-\frac{\widehat{\delta}}{m^2}u \right) \partial_r u \\ &= -2r^{-3}u_{\widehat{\delta}} + r^{-2}F' \left(-\frac{\widehat{\delta}}{m^2}u \right) r^2 \Upsilon^{-1} w, \end{aligned}$$

and since $w \geq 0$ and $F' \geq 0$, we deduce

$$\partial_r(f_{\widehat{\delta}}) \geq -2r^{-3}u_{\widehat{\delta}} \geq 2\widehat{\delta}^{-1}m^2r^{-3}.$$

For $u \geq -\frac{m^2}{\widehat{\delta}}$, using Lemma 10.1.15, we have

$$\partial_r(f_{\widehat{\delta}}) = \partial_r f \geq \frac{16m^2}{r^3}.$$

Hence, for all $r \geq 2m$, $\widehat{\delta} > 0$ sufficiently small,

$$\partial_r(f_{\widehat{\delta}}) \geq \frac{16m^2}{r^3}$$

which is (10.1.39). □

It remains to evaluate $W_{\widehat{\delta}}$. This is done in the following lemma.

Lemma 10.1.24. *Let*

$$\overline{W}_{\widehat{\delta}}(r, m) := 1_{r \leq \frac{5m}{2}} |W_{\widehat{\delta}}|. \tag{10.1.40}$$

Then, $\overline{W}_{\widehat{\delta}}$ is supported, for $\delta > 0$ small enough, in the region

$$2m + e^{-\frac{2}{\delta}} \leq r \leq \frac{9m}{4}.$$

Moreover its primitive,

$$\widetilde{W}_{\widehat{\delta}}(r, m) := \int_{2m}^r \overline{W}_{\widehat{\delta}}(r', m) dr' \tag{10.1.41}$$

verifies the pointwise estimate

$$\widetilde{W}_{\widehat{\delta}}(r, m) \lesssim \widehat{\delta}. \tag{10.1.42}$$

Proof. Recall that we have chosen $w = \frac{1}{4m}$ to be constant in the region $r \leq 4m$. Hence, in that region,

$$w_{\widehat{\delta}} = \frac{1}{4m} F' \left(-\frac{\widehat{\delta}}{m^2} u \right), \quad \partial_r w_{\widehat{\delta}} = \frac{1}{4m} \partial_r \left(F' \left(-\frac{\widehat{\delta}}{m^2} u \right) \right).$$

Hence,

$$W_{\widehat{\delta}} = -\frac{1}{4} r^{-2} \partial_r \left(\frac{1}{4m} r^2 \Upsilon \partial_r \left(F' \left(-\frac{\widehat{\delta}}{m^2} u \right) \right) \right) = -\frac{1}{16m} r^{-2} \partial_r \left(r^2 \Upsilon \partial_r \left(F' \left(-\frac{\widehat{\delta}}{m^2} u \right) \right) \right).$$

Now, setting $\delta_0 = \frac{\widehat{\delta}}{m^2}$ for convenience below,

$$r^{-2}\partial_r (r^2\Upsilon\partial_r (F'(-\delta_0 u))) = -\delta_0 F''(-\delta_0 u)r^{-2}\partial_r (r^2\Upsilon\partial_r u) + \delta_0^2 F'''(-\delta_0 u)\Upsilon(\partial_r u)^2.$$

Note that, since $r^{-2}\Upsilon\partial_r u = w$ and $w = (4m)^{-1}$ is constant in r in the region of interest

$$r^{-2}\partial_r (r^2\Upsilon\partial_r u) = r^{-2}\partial_r (r^4 r^{-2}\Upsilon\partial_r u) = r^{-2}\partial_r \left(\frac{r^4}{4m} \right) = \frac{r}{m}.$$

Hence,

$$\begin{aligned} r^{-2}\partial_r (r^2\Upsilon\partial_r (F'(-\delta_0 u))) &= -\delta_0 F''(-\delta_0 u)r^{-2}\partial_r (r^2\Upsilon\partial_r u) + \delta_0^2 F'''(-\delta_0 u)\Upsilon(\partial_r u)^2 \\ &= -\delta_0 F''(-\delta_0 u)\frac{r}{m} + \delta_0^2 F'''(-\delta_0 u)\Upsilon(\partial_r u)^2. \end{aligned}$$

Hence, for $r \leq 4m$, with $\delta_0 = \frac{\widehat{\delta}}{m^2}$,

$$|W_{\widehat{\delta}}| \lesssim \delta_0^2 |\Upsilon| |F'''(-\delta_0 u)| (\partial_r u)^2 + \delta_0 |F''(-\delta_0 u)|$$

or, since $|\partial_r u| \lesssim \frac{1}{r-2m}$, in the region of interest,

$$|W_{\widehat{\delta}}| \lesssim \frac{\delta_0^2}{|r-2m|} |F'''(-\delta_0 u)| + \delta_0 |F''(-\delta_0 u)|.$$

Since $F''(-\delta_0 u), F'''(-\delta_0 u)$ are supported in the region $1 \leq -\delta_0 u \leq 3$, i.e. $-\frac{3}{\delta_0} \leq u \leq -\frac{1}{\delta_0}$, for $\widehat{\delta} > 0$ sufficiently small

$$e^{-\frac{2}{\widehat{\delta}}} \leq r - 2m \leq e^{-\frac{1}{3\widehat{\delta}}} \leq \frac{m}{4}.$$

Hence,

$$\overline{W}_{\widehat{\delta}} = 1_{r \leq \frac{5}{2}m} |W_{\widehat{\delta}}| \lesssim \widehat{\delta} \left(\frac{\delta}{r-2m} + 1 \right) \kappa_{\widehat{\delta}}(r-2m)$$

with $\kappa_{\widehat{\delta}}(x)$ the characteristic function of the interval $[e^{-\frac{2}{\widehat{\delta}}}, e^{-\frac{1}{3\widehat{\delta}}}]$. Note that the primitive of $\overline{W}_{\widehat{\delta}}$, i.e.

$$\widetilde{W}_{\widehat{\delta}}(r, m) = \int_{2m}^r \overline{W}_{\widehat{\delta}}(r', m) dr',$$

is a positive, increasing function. Moreover,

$$\widetilde{W}_{\widehat{\delta}}(r) \lesssim \int_{2m}^{4m} \overline{W}_{\widehat{\delta}}(r) dr \lesssim \delta + \delta^2 \int_{e^{-\frac{2}{\widehat{\delta}}}}^{e^{-\frac{1}{3\widehat{\delta}}}} \frac{1}{x} dx \lesssim \delta$$

as desired. \square

We now recall that, see (10.1.15),

$$\begin{aligned} \dot{\mathcal{E}}[fR, w](\Psi) &= \dot{\mathcal{E}}_0[fR, w](\Psi) + 4\Upsilon \frac{r-4m}{r^4} f|\Psi|^2, \\ \dot{\mathcal{E}}_0[fR, w](\Psi) &= \partial_r f |R(\Psi)|^2 + r^{-2} W |\Psi|^2 + r^{-1} \left(1 - \frac{3m}{r}\right) f \widehat{\mathcal{Q}}_{34}. \end{aligned}$$

Using the functions $f_{\widehat{\delta}}$, $w_{\widehat{\delta}}$ and $W_{\widehat{\delta}}$ introduced in definition 10.1.22, we have

$$\dot{\mathcal{E}}_0[f_{\widehat{\delta}}R, w_{\widehat{\delta}}](\Psi) = \frac{1}{r} f_{\widehat{\delta}} \left(1 - \frac{3m}{r}\right) \widehat{\mathcal{Q}}_{34} + \partial_r(f_{\widehat{\delta}}) |R\Psi|^2 + W_{\widehat{\delta}} |\Psi|^2.$$

Note that in view of the estimates (10.1.38) (10.1.39), and Lemma 10.1.24, we immediately deduce the existence of a constant $C > 0$ independent of $\widehat{\delta}$ such that

$$\dot{\mathcal{E}}[f_{\widehat{\delta}}R, w_{\widehat{\delta}}](\Psi) \geq C^{-1} [|R\Psi|^2 + |\nabla\Psi|^2 + \Upsilon|\Psi|^2] - \overline{W}_{\widehat{\delta}} |\Psi|^2 \quad \text{on } r \leq \frac{5m}{2} \quad (10.1.43)$$

where $\overline{W}_{\widehat{\delta}}$ is a non-negative potential supported in the region $2m + e^{-\frac{2}{\widehat{\delta}}} \leq r \leq \frac{9m}{4}$, whose primitive $\widetilde{W}_{\widehat{\delta}}(r) = \int_{2m}^r \overline{W}_{\widehat{\delta}}(r'm) dr'$ verifies $\widetilde{W}_{\widehat{\delta}} \lesssim \widehat{\delta}$. Combining this with estimates of the previous section we derive the following.

Proposition 10.1.25. *There exists a constant $C > 0$, and for any small enough $\widehat{\delta} > 0$, there exists functions $f_{\widehat{\delta}} \in C^2(r > 0)$, $w_{\widehat{\delta}} \in C^1(r > 0)$ and $h \in C^2(r > 0)$ verifying, for all $r > 0$,*

$$|f_{\widehat{\delta}}(r)| \lesssim \widehat{\delta}^{-1}, \quad w_{\widehat{\delta}} \lesssim r^{-1}, \quad h \lesssim r^{-4},$$

such that

$$\mathcal{E}[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR](\Psi) = \dot{\mathcal{E}}[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR] + \mathcal{E}_\epsilon[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR](\Psi)$$

satisfies

$$\begin{aligned} \int_S \dot{\mathcal{E}}[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR] &\geq C^{-1} \int_S \left(\frac{m^2}{r^3} |R(\Psi)|^2 + r^{-1} \left(1 - \frac{3m}{r}\right)^2 \left(|\nabla\Psi|^2 + \frac{m^2}{r^2} |T\Psi|^2 \right) + \frac{m}{r^4} |\Psi|^2 \right) \\ &\quad - \int_S \overline{W}_{\widehat{\delta}} |\Psi|^2, \\ \mathcal{E}_\epsilon[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR] &= \epsilon \frac{1}{2} \mathcal{Q} \cdot (f_{\widehat{\delta}}R) \ddot{\pi} + O(r^{-3} u_{trap}^{-1-\delta_{dec}} (1 + |f_{\widehat{\delta}}|)) |\Psi|^2, \end{aligned}$$

where $\overline{W}_{\widehat{\delta}}$ is non-negative, supported in the region $2m + e^{-\frac{2}{\widehat{\delta}}} \leq r \leq \frac{9m}{4}$, and such that its primitive $\widetilde{W}_{\widehat{\delta}}(r) = \int_{2m}^r \overline{W}_{\widehat{\delta}}$ verifies $\widetilde{W}_{\widehat{\delta}} \lesssim \widehat{\delta}$.

Proof. We choose h to be the function of (r, m) introduced in Proposition 10.1.20, $f_{\widehat{\delta}}$ to be the function of (r, m) introduced in definition 10.1.22, and $\overline{W}_{\widehat{\delta}}$, introduced in Lemma 10.1.24. Also, by an abuse of notation, we denote by $w_{\widehat{\delta},0}$ the function denoted by $w_{\widehat{\delta}}$ in definition 10.1.22. Then, combining Proposition 10.1.20 in the region $r \geq \frac{5m}{2}$ with the estimate (10.1.43) in the region $r \leq \frac{5m}{2}$, we immediately obtain

$$\begin{aligned} \int_S \dot{\mathcal{E}}[f_{\widehat{\delta}}R, w_{\widehat{\delta},0}, 2hR] &\geq C^{-1} \int_S \left(\frac{m^2}{r^3} |R(\Psi)|^2 + r^{-1} \left(1 - \frac{3m}{r}\right)^2 |\nabla\Psi|^2 + \frac{m}{r^4} |\Psi|^2 \right) \\ &\quad - \int_S \overline{W}_{\widehat{\delta}} |\Psi|^2. \end{aligned} \quad (10.1.44)$$

(10.1.44) corresponds to the desired estimate without the presence of the term $|T\Psi|^2$ on the right hand side. To get the improved estimate of Proposition 10.1.25, we set

$$w_{\widehat{\delta}} := w_{\widehat{\delta},0} - \delta_1 w_1, \quad (10.1.45)$$

for a small parameter $\delta_1 > 0$ to be chosen later, where $w_{\widehat{\delta},0}$ is our previous choice introduced in definition 10.1.22, and where

$$w_1(r, m) := r^{-1} \frac{m^2}{r^2} \Upsilon \left(1 - \frac{3m}{r}\right)^2. \quad (10.1.46)$$

We evaluate (modulo the same type of error terms as before which we include in \mathcal{E}_ϵ),

$$\dot{\mathcal{E}}[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR](\Psi) = \dot{\mathcal{E}}[X_{\widehat{\delta}}, w_{\widehat{\delta},0}, 2hR] - \frac{1}{2} \delta_1 w_1 \mathcal{L}(\Psi) + \frac{\delta_1}{4} |\Psi|^2 r^{-2} \partial_r (r^2 \Upsilon \partial_r w_1).$$

Now, since

$$\begin{aligned} \mathcal{L}(\Psi) &= -e_3 \Psi \cdot e_4 \Psi + |\nabla\Psi|^2 + V|\Psi|^2 \\ &= \Upsilon^{-1} (-|T\Psi|^2 + |R\Psi|^2) + |\nabla\Psi|^2 + V|\Psi|^2, \end{aligned}$$

we have,

$$\begin{aligned} &-\frac{1}{2} \delta_1 w_1 \mathcal{L}(\Psi) + \frac{\delta_1}{4} |\Psi|^2 r^{-2} \partial_r (r^2 \Upsilon \partial_r w_1) \\ &= -\frac{1}{2} \delta_1 r^{-1} \frac{m^2}{r^2} \Upsilon \left(1 - \frac{3m}{r}\right)^2 \mathcal{L}(\Psi) + \frac{\delta_1}{4} |\Psi|^2 r^{-2} \partial_r \left(r^2 \Upsilon \partial_r \left(r^{-1} \frac{m^2}{r^2} \Upsilon \left(1 - \frac{3m}{r}\right)^2 \right) \right) \\ &= \frac{1}{2} \delta_1 r^{-1} \left(1 - \frac{3m}{r}\right)^2 \frac{m^2}{r^2} |T\Psi|^2 + O(\delta_1) \left(\frac{m^2}{r^3} |R(\Psi)|^2 + r^{-1} \left(1 - \frac{3m}{r}\right)^2 |\nabla\Psi|^2 + \frac{m}{r^4} |\Psi|^2 \right) \end{aligned}$$

and hence

$$\begin{aligned} \dot{\mathcal{E}}[f_{\delta}R, w_{\delta}, 2hR](\Psi) &= \dot{\mathcal{E}}[X_{\delta}, w_{\delta,0}, 2hR] + \frac{1}{2}\delta_1 r^{-1} \left(1 - \frac{3m}{r}\right)^2 \frac{m^2}{r^2} |T\Psi|^2 \\ &+ O(\delta_1) \left(\frac{m^2}{r^3} |R(\Psi)|^2 + r^{-1} \left(1 - \frac{3m}{r}\right)^2 |\nabla\Psi|^2 + \frac{m}{r^4} |\Psi|^2 \right). \end{aligned} \tag{10.1.47}$$

The desired estimate now follows from (10.1.44) and (10.1.47) provided $\delta_1 > 0$ is chosen small enough compared to the constant $C > 0$ of (10.1.44) so that the last term $O(\delta_1)$ in the above identity can be absorbed. \square

10.1.10 The red shift vectorfield

Note that the vectorfields T and R become both proportional to e_4 for $\Upsilon = 0$ which means that the estimate of Proposition 10.1.25 degenerates along $\Upsilon = 0$, i.e. it does not control $e_3(\Psi)$ there. In this section we make use of the Dafermos-Rodnianski red shift vectorfield to compensate for this degeneracy. The crucial ingredient here is the favorable sign of ω in a small neighborhood of $r = 2m$.

Lemma 10.1.26. *Let $\pi^{(3)}, \pi^{(4)}$ denote the deformation tensors of e_3, e_4 . In the region $r \leq 3m$ all components are $O(\epsilon)$ with the exception of,*

$$\begin{aligned} \pi_{44}^{(3)} &= -8\omega = \frac{8m}{r^2} + O(\epsilon), & \pi_{\theta\theta}^{(3)} &= \underline{\kappa} + \underline{\vartheta} = -\frac{2}{r} + O(\epsilon), \\ \pi_{\varphi\varphi}^{(3)} &= \underline{\kappa} - \underline{\vartheta} = -\frac{2}{r} + O(\epsilon), \\ \pi_{34}^{(4)} &= 4\omega = -\frac{4m}{r^2} + O(\epsilon), & \pi_{\theta\theta}^{(4)} &= \kappa + \vartheta = \frac{2\Upsilon}{r} + O(\epsilon), \\ \pi_{\varphi\varphi}^{(4)} &= \kappa - \vartheta = \frac{2\Upsilon}{r} + O(\epsilon). \end{aligned}$$

Proof. Immediate verification in view of our assumptions. \square

Lemma 10.1.27. *Given the vectorfield,*

$$Y = a(r, m)e_3 + b(r, m)e_4, \tag{10.1.48}$$

and assuming

$$\sup_{r \leq 3m} \left(|a| + |\partial_r a| + |\partial_m a| + |b| + |\partial_r b| + |\partial_m b| \right) \lesssim 1,$$

we have, for $r \leq 3m$,

$$\begin{aligned} \mathcal{Q}^{\alpha\beta(Y)}\pi_{\alpha\beta} &= \left(\frac{2m}{r^2}a - \Upsilon\partial_r a\right)\mathcal{Q}_{33} + \partial_r b\mathcal{Q}_{44} + \left(\partial_r a - \frac{2m}{r^2}b - \Upsilon\partial_r b\right)\mathcal{Q}_{34} \\ &+ \frac{2}{r}(b\Upsilon - a)e_3\Psi \cdot e_4\Psi + 8\frac{\Upsilon}{r^3}(a - \Upsilon b)|\Psi|^2 \\ &+ O(\epsilon)(|\mathcal{Q}(\Psi)| + r^{-2}|\Psi|^2). \end{aligned}$$

Moreover, with the notation (10.1.13),

$$\mathcal{E}[Y, 0](\Psi) = \frac{1}{2}\mathcal{Q}^{\alpha\beta(Y)}\pi_{\alpha\beta} + 4\frac{r-3m}{r^4}(-a + b\Upsilon)|\Psi|^2 + O(\epsilon)r^{-2}|\Psi|^2. \quad (10.1.49)$$

Proof. In view of

$$|e_4(r) - \Upsilon, e_3(r) + 1| \lesssim \epsilon$$

Lemma 10.1.5, and the assumptions on the derivatives of a and b w.r.t. (r, m) , we have

$$\begin{aligned} e_4(a) &= \Upsilon\partial_r a + O(\epsilon), & e_3(a) &= -\partial_r a + O(\epsilon), \\ e_4(b) &= \Upsilon\partial_r b + O(\epsilon), & e_3(b) &= -\partial_r b + O(\epsilon), & e_\theta(a) &= e_\theta(b) = 0. \end{aligned}$$

We infer,

$$\begin{aligned} \mathcal{Q}^{\alpha\beta(Y)}\pi_{\alpha\beta} &= a\mathcal{Q}^{\alpha\beta}\pi_{\alpha\beta}^{(3)} - (\mathcal{Q}_{33}e_4a + \mathcal{Q}_{43}e_3a) + b\mathcal{Q}^{\alpha\beta}\pi_{\alpha\beta}^{(4)} - (\mathcal{Q}_{34}e_4b + \mathcal{Q}_{44}e_3b) + O(\epsilon)|\mathcal{Q}(\Psi)| \\ &= a\mathcal{Q}^{\alpha\beta}\pi_{\alpha\beta}^{(3)} + b\mathcal{Q}^{\alpha\beta}\pi_{\alpha\beta}^{(4)} - \mathcal{Q}_{33}\Upsilon\partial_r a - \mathcal{Q}_{34}(-\partial_r a + \Upsilon\partial_r b) + \mathcal{Q}_{44}\partial_r b + O(\epsilon)|\mathcal{Q}(\Psi)|. \end{aligned}$$

Note that,

$$\mathcal{Q}_{\theta\theta} + \mathcal{Q}_{\varphi\varphi} = e_3\Psi \cdot e_4\Psi - V|\Psi|^2 = e_3\Psi \cdot e_4\Psi - 4\frac{\Upsilon}{r^2}|\Psi|^2 + O(\epsilon)r^{-2}|\Psi|^2. \quad (10.1.50)$$

Hence,

$$\begin{aligned} \mathcal{Q}^{\alpha\beta}\pi_{\alpha\beta}^{(3)} &= \mathcal{Q}^{44}\pi_{44}^{(3)} + \mathcal{Q}^{\theta\theta}\pi_{\theta\theta}^{(3)} + \mathcal{Q}^{\varphi\varphi}\pi_{\varphi\varphi}^{(3)} + O(\epsilon)|\mathcal{Q}(\Psi)| \\ &= \frac{1}{4}\mathcal{Q}_{33}\frac{8m}{r^2} - \frac{2}{r}(\mathcal{Q}_{\theta\theta} + \mathcal{Q}_{\varphi\varphi}) + O(\epsilon)|\mathcal{Q}(\Psi)| \\ &= \frac{2m}{r^2}\mathcal{Q}_{33} - \frac{2}{r}e_3\Psi \cdot e_4\Psi + 8\frac{\Upsilon}{r^3}|\Psi|^2 + O(\epsilon)(|\mathcal{Q}(\Psi)| + r^{-2}|\Psi|^2), \\ \mathcal{Q}^{\alpha\beta}\pi_{\alpha\beta}^{(4)} &= 2\mathcal{Q}^{34}\pi_{34}^{(4)} + \mathcal{Q}^{\theta\theta}\pi_{\theta\theta}^{(4)} + \mathcal{Q}^{\varphi\varphi}\pi_{\varphi\varphi}^{(4)} + O(\epsilon)|\mathcal{Q}(\Psi)| \\ &= \frac{1}{2}\mathcal{Q}_{34}\left(-4\frac{m}{r^2}\right) + \frac{2\Upsilon}{r}(\mathcal{Q}_{\theta\theta} + \mathcal{Q}_{\varphi\varphi}) + O(\epsilon)|\mathcal{Q}(\Psi)| \\ &= -\frac{2m}{r^2}\mathcal{Q}_{34} + \frac{2\Upsilon}{r}e_3\Psi \cdot e_4\Psi - 8\frac{\Upsilon^2}{r^3}|\Psi|^2 + O(\epsilon)(|\mathcal{Q}(\Psi)| + r^{-2}|\Psi|^2). \end{aligned}$$

Therefore,

$$\begin{aligned}
\mathcal{Q}^{\alpha\beta(Y)}\pi_{\alpha\beta} &= a \left[\frac{2m}{r^2} \mathcal{Q}_{33} - \frac{2}{r} e_3 \Psi e_4 \Psi + 8 \frac{\Upsilon}{r^3} |\Psi|^2 \right] + b \left[-\frac{2m}{r^2} \mathcal{Q}_{34} + \frac{2\Upsilon}{r} e_3 \Psi \cdot e_4 \Psi - 8 \frac{\Upsilon^2}{r^3} |\Psi|^2 \right] \\
&- \mathcal{Q}_{33} \Upsilon \partial_r a - \mathcal{Q}_{34} (-\partial_r a + \Upsilon \partial_r b) + \mathcal{Q}_{44} \partial_r b + O(\epsilon) (|\mathcal{Q}(\Psi)| + r^{-2} |\Psi|^2) \\
&= \left(\frac{2m}{r^2} a - \Upsilon \partial_r a \right) \mathcal{Q}_{33} + \mathcal{Q}_{44} \partial_r b + \frac{2}{r} (b\Upsilon - a) e_3 \Psi e_4 \Psi + \left(\partial_r a - \frac{2m}{r^2} b - \Upsilon \partial_r b \right) \mathcal{Q}_{34} \\
&+ 8 \frac{\Upsilon}{r^3} (a - \Upsilon b) |\Psi|^2 + O(\epsilon) (|\mathcal{Q}(\Psi)| + r^{-2} |\Psi|^2).
\end{aligned}$$

To prove the second part of the lemma we recall (see (10.1.13)),

$$\mathcal{E}[Y, 0](\Psi) = \frac{1}{2} \mathcal{Q}^{\alpha\beta(Y)} \pi_{\alpha\beta} - \frac{1}{2} Y(V) |\Psi|^2$$

and, relying on Lemma 10.1.5, we have on $r \leq 3m$

$$Y(V) = (-a + b\Upsilon) \partial_r V + O(\epsilon) = (-a + b\Upsilon) \left(-8 \frac{r - 3m}{r^4} \right) + O(\epsilon)$$

which concludes the proof of the lemma. \square

Corollary 10.1.28. *If we choose,*

$$a(2m, m) = 1, \quad b(2m, m) = 0, \quad \partial_r a(2m, m) \geq \frac{1}{4m}, \quad \partial_r b(2m, m) \geq \frac{5}{4m},$$

then, at $r = 2m$, we have

$$\mathcal{Q}^{\alpha\beta(Y)} \pi_{\alpha\beta} \geq \frac{1}{4m} (|e_3 \Psi|^2 + |e_4 \Psi|^2 + \mathcal{Q}_{34}) + O(\epsilon) (|\mathcal{Q}(\Psi)| + r^{-2} |\Psi|^2) \quad (10.1.51)$$

and,

$$\mathcal{E}[Y, 0](\Psi) \geq \frac{1}{8m} \left(|e_3 \Psi|^2 + |e_4 \Psi|^2 + \mathcal{Q}_{34} + \frac{1}{m^2} |\Psi|^2 \right) + O(\epsilon) (|\mathcal{Q}(\Psi)| + r^{-2} |\Psi|^2). \quad (10.1.52)$$

Moreover the estimates remain true if we add to Y a multiple of $T = \frac{1}{2} (e_4 + \Upsilon e_3)$.

Proof. Recall from Lemma 10.1.27 that we have, for $r \leq 3m$,

$$\begin{aligned}
\mathcal{Q}^{\alpha\beta(Y)} \pi_{\alpha\beta} &= \left(\frac{2m}{r^2} a - \Upsilon \partial_r a \right) \mathcal{Q}_{33} + \partial_r b \mathcal{Q}_{44} + \left(\partial_r a - \frac{2m}{r^2} b - \Upsilon \partial_r b \right) \mathcal{Q}_{34} \\
&+ \frac{2}{r} (b\Upsilon - a) e_3 \Psi \cdot e_4 \Psi + 8 \frac{\Upsilon}{r^3} (a - \Upsilon b) |\Psi|^2 + O(\epsilon) (|\mathcal{Q}(\Psi)| + r^{-2} |\Psi|^2).
\end{aligned}$$

Hence, at $r = 2m$, using $\Upsilon = 0$, $a = 1$, $b = 0$, $\partial_r a \geq (4m)^{-1}$ and $\partial_r b \geq 5(4m)^{-1}$, we deduce

$$\begin{aligned} \mathcal{Q}^{\alpha\beta(Y)}\pi_{\alpha\beta} &= \frac{1}{2m}\mathcal{Q}_{33} + \partial_r b\mathcal{Q}_{44} + \partial_r a\mathcal{Q}_{34} - \frac{1}{m}e_3\Psi \cdot e_4\Psi + O(\epsilon)(|\mathcal{Q}(\Psi)| + r^{-2}|\Psi|^2) \\ &\geq \frac{1}{2m}|e_3(\Psi)|^2 + \frac{5}{4m}|e_4(\Psi)|^2 + \frac{1}{4m}\mathcal{Q}_{34} - \frac{1}{m}e_3\Psi \cdot e_4\Psi + O(\epsilon)(|\mathcal{Q}(\Psi)| + r^{-2}|\Psi|^2) \end{aligned}$$

from which the desired lower bound in (10.1.51) follows.

Also, at $r = 2m$, using (10.1.49), $\Upsilon = 0$, $a = 1$, and $b = 0$, we have

$$\begin{aligned} \mathcal{E}[Y, 0](\Psi) &= \frac{1}{2}\mathcal{Q}^{\alpha\beta(Y)}\pi_{\alpha\beta} + 4\frac{r-3m}{r^4}(-a+b\Upsilon)|\Psi|^2 + O(\epsilon)r^{-2}|\Psi|^2 \\ &= \frac{1}{2}\mathcal{Q}^{\alpha\beta(Y)}\pi_{\alpha\beta} + \frac{1}{4m^3}|\Psi|^2 \\ &\geq \frac{1}{8m}\left(|e_3\Psi|^2 + |e_4\Psi|^2 + \mathcal{Q}_{34} + \frac{1}{m^2}|\Psi|^2\right) + O(\epsilon)(|\mathcal{Q}(\Psi)| + r^{-2}|\Psi|^2) \end{aligned}$$

which yields (10.1.52). □

We are now ready to prove the following result.

Proposition 10.1.29. *Given a small parameter $\delta_{\mathcal{H}} > 0$ there exists a smooth vectorfield $Y_{\mathcal{H}}$ supported in the region $|\Upsilon| \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}$ such that the following estimate holds,*

$$\begin{aligned} \mathcal{E}[Y_{\mathcal{H}}, 0](\Psi) &\geq \frac{1}{16m}1_{|\Upsilon| \leq \delta_{\mathcal{H}}^{\frac{1}{10}}}\left(|e_3\Psi|^2 + |e_4\Psi|^2 + \widehat{\mathcal{Q}}_{34} + m^{-2}|\Psi|^2\right) \\ &\quad - \frac{1}{m}\delta_{\mathcal{H}}^{-\frac{1}{10}}1_{\delta_{\mathcal{H}}^{\frac{1}{10}} \leq \Upsilon \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}}\left(|e_3\Psi|^2 + |e_4\Psi|^2 + \widehat{\mathcal{Q}}_{34} + m^{-2}|\Psi|^2\right) \\ &\quad + O(\epsilon)1_{|\Upsilon| \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}}\left(|\mathcal{Q}(\Psi)| + m^{-2}|\Psi|^2\right). \end{aligned}$$

Moreover, for $|\Upsilon| \leq \delta_{\mathcal{H}}^{\frac{1}{10}}$, we have

$$Y_{\mathcal{H}} = e_3 + e_4 + O(\delta_{\mathcal{H}}^{\frac{1}{10}})(e_3 + e_4).$$

Proof. We introduce the vectorfield

$$Y_{(0)} := ae_3 + be_4 + 2T, \quad a(r, m) := 1 + \frac{5}{4m}(r - 2m), \quad b(r, m) := \frac{5}{4m}(r - 2m),$$

with $T = \frac{1}{2}(e_4 + \Upsilon e_3)$. Also, we pick positive bump function $\kappa = \kappa(r)$, supported in the region in $[-2, 2]$ and equal to 1 for $[-1, 1]$ and define, for sufficiently small $\delta_{\mathcal{H}} > 0$.

$$Y_{\mathcal{H}} := \kappa_{\mathcal{H}} Y_{(0)}, \quad \kappa_{\mathcal{H}} := \kappa \left(\frac{\Upsilon}{\delta_{\mathcal{H}}^{\frac{1}{10}}} \right). \tag{10.1.53}$$

We have

$$\begin{aligned} \mathcal{E}[Y_{\mathcal{H}}, 0](\Psi) &= \mathcal{Q} \cdot {}^{(Y_{\mathcal{H}})}\pi - Y_{\mathcal{H}}(V)|\Psi|^2 \\ &= \kappa_{\mathcal{H}} \mathcal{Q} \cdot {}^{(Y_0)}\pi + \mathcal{Q}(Y_{(0)}, d\kappa_{\mathcal{H}}) + \kappa_{\mathcal{H}} Y_{(0)}(V)|\Psi|^2 \\ &= \kappa_{\mathcal{H}} \mathcal{E}[Y_{(0)}, 0](\Psi) + O(\delta_{\mathcal{H}}^{-\frac{1}{10}}) 1_{\delta_{\mathcal{H}}^{\frac{1}{10}} \leq \Upsilon \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}} \left(|e_3 \Psi|^2 + |e_4 \Psi|^2 + \widehat{\mathcal{Q}}_{34} + m^{-2} |\Psi|^2 \right). \end{aligned}$$

Note from the definition of $Y_{(0)}$ and the choice of a and b that Corollary 10.1.28 applies to $Y_{(0)}$. In particular, we deduce from (10.1.52) for $\delta_{\mathcal{H}} > 0$ small enough,

$$\begin{aligned} \mathcal{E}[Y_{\mathcal{H}}, 0](\Psi) &\geq \frac{1}{16m} 1_{|\Upsilon| \leq \delta_{\mathcal{H}}^{\frac{1}{10}}} \left(|e_3 \Psi|^2 + |e_4 \Psi|^2 + \widehat{\mathcal{Q}}_{34} + m^{-2} |\Psi|^2 \right) \\ &\quad - \frac{1}{m} \delta_{\mathcal{H}}^{-\frac{1}{10}} 1_{\delta_{\mathcal{H}}^{\frac{1}{10}} \leq \Upsilon \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}} \left(|e_3 \Psi|^2 + |e_4 \Psi|^2 + \widehat{\mathcal{Q}}_{34} + m^{-2} |\Psi|^2 \right) \\ &\quad + O(\epsilon) 1_{|\Upsilon| \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}} \left(|\mathcal{Q}(\Psi)| + m^{-2} |\Psi|^2 \right) \end{aligned}$$

as desired. □

10.1.11 Combined Estimate

We consider the combined Morawetz triplet

$$(X, w, M) := (X_{\widehat{\delta}}, w_{\widehat{\delta}}, 2hR) + \epsilon_{\mathcal{H}}(Y_{\mathcal{H}}, 0, 0), \tag{10.1.54}$$

with $\epsilon_{\mathcal{H}} > 0$ sufficiently small to be determined later. Here $(X_{\widehat{\delta}} = f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR)$ is the triplet given by Proposition 10.1.25 and $Y_{\mathcal{H}}$ the vectorfield of Proposition 10.1.29.

Recall, see Proposition 10.1.25, that $\dot{\mathcal{E}}_{\widehat{\delta}} := \dot{\mathcal{E}}[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR](\Psi)$ verifies,

$$\begin{aligned} \int_S \dot{\mathcal{E}}_{\widehat{\delta}} &\geq C^{-1} \int_S \left(\frac{m^2}{r^3} |R(\Psi)|^2 + \left(1 - \frac{3m}{r} \right)^2 r^{-1} \left(\widehat{\mathcal{Q}}_{34} + \frac{m^2}{r^2} |T\Psi|^2 \right) + \Upsilon \frac{m}{r^4} |\Psi|^2 \right) \\ &\quad - \int_S \overline{W}_{\widehat{\delta}} |\Psi|^2. \end{aligned}$$

According to Proposition 10.1.29, we write for $\mathcal{E}_{\mathcal{H}} = \mathcal{E}(Y_{\mathcal{H}}, 0, 0)(\Psi)$,

$$\begin{aligned}\mathcal{E}_{\mathcal{H}} &= \dot{\mathcal{E}}_{\mathcal{H}} + \mathcal{E}_{\mathcal{H},\epsilon}, \\ \dot{\mathcal{E}}_{\mathcal{H}} &\geq \frac{1}{8m} 1_{|\Upsilon| \leq \delta_{\mathcal{H}}^{\frac{1}{10}}} \left(|e_3 \Psi|^2 + |e_4 \Psi|^2 + \widehat{\mathcal{Q}}_{34} + m^{-2} |\Psi|^2 \right) \\ &\quad - \frac{1}{m} \delta_{\mathcal{H}}^{-\frac{1}{10}} 1_{\delta_{\mathcal{H}}^{\frac{1}{10}} \leq \Upsilon \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}} \left(|e_3 \Psi|^2 + |e_4 \Psi|^2 + \widehat{\mathcal{Q}}_{34} + m^{-2} |\Psi|^2 \right), \\ \mathcal{E}_{\mathcal{H},\epsilon} &= O(\epsilon) (|\mathcal{Q}(\Psi)| + m^{-2} |\Psi|^2) 1_{|\Upsilon| \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}}.\end{aligned}$$

Note that, for $|\Upsilon| \geq \delta_{\mathcal{H}}^{\frac{1}{10}}$ we have,

$$|R\Psi|^2 + |T\Psi|^2 = \frac{1}{2} (|e_4 \Psi|^2 + \Upsilon^2 |e_3 \Psi|^2) \geq \frac{1}{2} \delta_{\mathcal{H}}^{\frac{1}{5}} (|e_4 \Psi|^2 + |e_3 \Psi|^2).$$

We now proceed to find a lower bound for the expression $\dot{\mathcal{E}}_{\widehat{\delta}} + \epsilon_{\mathcal{H}} \dot{\mathcal{E}}_{\mathcal{H}}$. For brevity the S integration is omitted below.

Region $\delta_{\mathcal{H}}^{\frac{1}{10}} \leq |\Upsilon| \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}$.

$$\begin{aligned}\dot{\mathcal{E}}_{\widehat{\delta}} + \epsilon_{\mathcal{H}} \dot{\mathcal{E}}_{\mathcal{H}} &\geq m^{-1} C^{-1} \left[\delta_{\mathcal{H}}^{\frac{1}{5}} (|e_4 \Psi|^2 + |e_3 \Psi|^2) + m^{-2} |\Psi|^2 + |\nabla \Psi|^2 \right] - \overline{W}_{\widehat{\delta}} |\Psi|^2 \\ &\quad - \epsilon_{\mathcal{H}} \frac{1}{m} \delta_{\mathcal{H}}^{-\frac{1}{10}} (|e_3 \Psi|^2 + |e_4 \Psi|^2 + |\nabla \Psi|^2 + m^{-2} |\Psi|^2).\end{aligned}$$

Therefore, choosing $\epsilon_{\mathcal{H}} \leq (2C)^{-1} \delta_{\mathcal{H}}^{\frac{3}{10}}$, we deduce,

$$\dot{\mathcal{E}}_{\widehat{\delta}} + \epsilon_{\mathcal{H}} \dot{\mathcal{E}}_{\mathcal{H}} \geq m^{-1} \delta_{\mathcal{H}}^{\frac{1}{5}} (2C)^{-1} (|e_4 \Psi|^2 + |e_3 \Psi|^2 + |\nabla \Psi|^2 + m^{-2} |\Psi|^2) - \overline{W}_{\widehat{\delta}} |\Psi|^2.$$

Region $|\Upsilon| \leq \delta_{\mathcal{H}}^{\frac{1}{10}}$.

$$\epsilon_{\mathcal{H}} \dot{\mathcal{E}}_{\mathcal{H}} + \dot{\mathcal{E}}_{\widehat{\delta}} \geq \epsilon_{\mathcal{H}} \frac{1}{16m} \left(|e_3 \Psi|^2 + |e_4 \Psi|^2 + \widehat{\mathcal{Q}}_{34} + m^{-2} |\Psi|^2 \right) - \overline{W}_{\widehat{\delta}} |\Psi|^2.$$

Region $\Upsilon \geq 2\delta_{\mathcal{H}}^{\frac{1}{10}}$. In this region $\dot{\mathcal{E}}_{\widehat{\delta}} + \epsilon_{\mathcal{H}} \dot{\mathcal{E}}_{\mathcal{H}} = \dot{\mathcal{E}}_{\widehat{\delta}}$. Hence (ignoring the S -integration),

$$\begin{aligned}\dot{\mathcal{E}}_{\widehat{\delta}} + \epsilon_{\mathcal{H}} \dot{\mathcal{E}}_{\mathcal{H}} &\geq C^{-1} \left(\frac{m^2}{r^3} |R(\Psi)|^2 + \left(1 - \frac{3m}{r} \right)^2 r^{-1} \left(\widehat{\mathcal{Q}}_{34} + \frac{m^2}{r^2} |T\Psi|^2 \right) + \frac{m}{r^4} |\Psi|^2 \right) \\ &\quad - \overline{W}_{\widehat{\delta}} |\Psi|^2.\end{aligned}$$

To combine these three cases together we modify the vectorfields R, T near $r = 2m$ according to (5.1.11), i.e.

$$\begin{aligned}\check{R} &:= \theta \frac{1}{2}(e_4 - e_3) + (1 - \theta)\Upsilon^{-1}R = \frac{1}{2} \left[\check{\theta}e_4 - e_3 \right], \\ \check{T} &:= \theta \frac{1}{2}(e_4 + e_3) + (1 - \theta)\Upsilon^{-1}T = \frac{1}{2} \left[\check{\theta}e_4 + e_3 \right],\end{aligned}$$

where θ a smooth bump function equal 1 on $|\Upsilon| \leq \delta_{\mathcal{H}}^{\frac{1}{10}}$ vanishing for $|\Upsilon| \geq 2\delta_{\mathcal{H}}^{\frac{1}{10}}$, and where

$$\check{\theta} = \theta + \Upsilon^{-1}(1 - \theta) = \begin{cases} 1, & \text{for } |\Upsilon| \leq \delta_{\mathcal{H}}^{\frac{1}{10}}, \\ \Upsilon^{-1}, & \text{for } |\Upsilon| \geq 2\delta_{\mathcal{H}}^{\frac{1}{10}}. \end{cases}$$

Note that

$$2(|\check{R}\Psi|^2 + |\check{T}\Psi|^2) = |e_3\Psi|^2 + \check{\theta}^2|e_4\Psi|^2.$$

Thus in the region $|\Upsilon| \leq \delta_{\mathcal{H}}^{\frac{1}{10}}$ we have $|e_3\Psi|^2 + |e_4\Psi|^2 = 2(|\check{R}\Psi|^2 + |\check{T}\Psi|^2)$ and therefore,

$$\begin{aligned}\dot{\mathcal{E}}_{\hat{\delta}} + \epsilon_{\mathcal{H}}\dot{\mathcal{E}}_{\mathcal{H}} &\geq \epsilon_{\mathcal{H}}\frac{1}{16m} \left(|e_3\Psi|^2 + |e_4\Psi|^2 + \widehat{\mathcal{Q}}_{34} + m^{-2}|\Psi|^2 \right) - \overline{W}_{\hat{\delta}}|\Psi|^2 \\ &= \epsilon_{\mathcal{H}}\frac{1}{16m} \left(|\check{R}\Psi|^2 + |\check{T}\Psi|^2 + \widehat{\mathcal{Q}}_{34} + m^{-2}|\Psi|^2 \right) - \overline{W}_{\hat{\delta}}|\Psi|^2.\end{aligned}$$

In the region $\delta_{\mathcal{H}}^{\frac{1}{10}} \leq |\Upsilon| \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}$, we have $|\check{R}\Psi|^2 + |\check{T}\Psi|^2 \lesssim |e_3\Psi|^2 + \delta_{\mathcal{H}}^{-\frac{1}{5}}|e_4\Psi|^2$. Hence, for $\epsilon_{\mathcal{H}} \leq (2C)^{-1}\delta_{\mathcal{H}}^{\frac{3}{10}}$, we deduce,

$$\begin{aligned}\dot{\mathcal{E}}_{\hat{\delta}} + \epsilon_{\mathcal{H}}\dot{\mathcal{E}}_{\mathcal{H}} &\geq m^{-1}\delta_{\mathcal{H}}^{\frac{1}{5}}(2C)^{-1} \left(|e_4\Psi|^2 + |e_3\Psi|^2 + |\nabla\Psi|^2 + m^{-2}|\Psi|^2 \right) - \overline{W}_{\hat{\delta}}|\Psi|^2 \\ &\geq m^{-1}\delta_{\mathcal{H}}^{\frac{1}{5}}(2C)^{-1} \left(\delta_{\mathcal{H}}^{\frac{1}{5}} \left(|\check{R}\Psi|^2 + |\check{T}\Psi|^2 \right) + |\nabla\Psi|^2 + m^{-2}|\Psi|^2 \right) - \overline{W}_{\hat{\delta}}|\Psi|^2.\end{aligned}$$

Finally, for $|\Upsilon| \geq 2\delta_{\mathcal{H}}^{\frac{1}{10}}$ we have $\check{R} = \Upsilon^{-1}R$, $\check{T} = \Upsilon^{-1}T$. Hence,

$$\begin{aligned}\dot{\mathcal{E}}_{\hat{\delta}} + \epsilon_{\mathcal{H}}\dot{\mathcal{E}}_{\mathcal{H}} &\geq C^{-1} \left(\frac{m^2}{r^3}|R(\Psi)|^2 + \left(1 - \frac{3m}{r}\right)^2 r^{-1} \left(\widehat{\mathcal{Q}}_{34} + \frac{m^2}{r^2}|T\Psi|^2 \right) + \Upsilon\frac{m}{r^4}|\Psi|^2 \right) \\ &\quad - \overline{W}_{\hat{\delta}}|\Psi|^2 \\ &\geq C^{-1} \left(\delta_{\mathcal{H}}^{\frac{1}{5}}\frac{m^2}{r^3}|\check{R}(\Psi)|^2 + \left(1 - \frac{3m}{r}\right)^2 r^{-1} \left(\widehat{\mathcal{Q}}_{34} + \delta_{\mathcal{H}}^{\frac{2}{10}}\frac{m^2}{r^2}|\check{T}\Psi|^2 \right) + \Upsilon\frac{m}{r^4}|\Psi|^2 \right) \\ &\quad - \overline{W}_{\hat{\delta}}|\Psi|^2.\end{aligned}$$

We deduce the following.

Proposition 10.1.30. *Let $C > 0$ the constant of Proposition 10.1.25. Consider the combined Morawetz triplet*

$$(X, w, M) := (X_{\widehat{\delta}}, w_{\widehat{\delta}}, 2hR) + \epsilon_{\mathcal{H}}(Y_{\mathcal{H}}, 0, 0), \quad (10.1.55)$$

with $C^{-1}\delta_{\mathcal{H}}^{\frac{2}{5}} \leq \epsilon_{\mathcal{H}} \leq (2C)^{-1}\delta_{\mathcal{H}}^{\frac{3}{10}}$ where, for given fixed $\widehat{\delta} > 0$, $(X_{\widehat{\delta}}, w_{\widehat{\delta}}, 2hR)$ is the triplet of Proposition 10.1.25 and $Y_{\mathcal{H}}$ the vectorfield of Proposition 10.1.29, supported in $|\Upsilon| \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}$ with $\delta_{\mathcal{H}} > 0$ sufficiently small, independent of $\widehat{\delta}$. Let $\dot{\mathcal{E}}_{\widehat{\delta}}, \dot{\mathcal{E}}_{\mathcal{H}}$ be the principal parts of $\mathcal{E}[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR](\Psi)$ and respectively $\mathcal{E}_{\mathcal{H}}[Y_{\mathcal{H}}, 0, 0](\Psi)$ and $\mathcal{E}_{\widehat{\delta}, \epsilon}, \mathcal{E}_{\mathcal{H}, \epsilon}$ the corresponding error terms, i.e.,

$$\mathcal{E}[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR](\Psi) = \dot{\mathcal{E}}_{\widehat{\delta}} + \mathcal{E}_{\widehat{\delta}, \epsilon}, \quad \mathcal{E}_{\mathcal{H}}[Y_{\mathcal{H}}, 0, 0](\Psi) = \dot{\mathcal{E}}_{\mathcal{H}} + \mathcal{E}_{\mathcal{H}, \epsilon}.$$

Then, provided $\delta_{\mathcal{H}} > 0$ is sufficiently small, we have

1. In the region $-2\delta_{\mathcal{H}}^{\frac{1}{10}} \leq \Upsilon$, $r \leq \frac{5m}{2}$, we have with a constant $\Lambda_{\mathcal{H}}^{-1} := C^{-1}\delta_{\mathcal{H}}^{\frac{2}{5}} > 0$

$$\int_S (\dot{\mathcal{E}}_{\widehat{\delta}} + \epsilon_{\mathcal{H}}\dot{\mathcal{E}}_{\mathcal{H}}) \geq m^{-1}\Lambda_{\mathcal{H}}^{-1} \int_S \left(|\check{R}(\Psi)|^2 + |\check{T}\Psi|^2 + |\check{\nabla}\Psi|^2 + m^{-2}|\Psi|^2 \right) - \int_S \overline{W}_{\widehat{\delta}}|\Psi|^2.$$

2. In the region $r \geq \frac{5m}{2}$, where $\dot{\mathcal{E}}_{\widehat{\delta}} + \epsilon_{\mathcal{H}}\dot{\mathcal{E}}_{\mathcal{H}} = \dot{\mathcal{E}}_{\widehat{\delta}}$ and $\overline{W}_{\widehat{\delta}} = 0$, we have the same estimate as in Proposition 10.1.25, i.e.

$$\int_S (\dot{\mathcal{E}}_{\widehat{\delta}} + \epsilon_{\mathcal{H}}\dot{\mathcal{E}}_{\mathcal{H}}) \geq C^{-1} \int_S \left(\frac{m^2}{r^3} |R(\Psi)|^2 + r^{-1} \left(1 - \frac{3m}{r} \right)^2 \left(|\check{\nabla}\Psi|^2 + \frac{m^2}{r^2} |T\Psi|^2 \right) + \frac{m}{r^4} |\Psi|^2 \right).$$

3. The ϵ -error terms verify the upper bound estimate,

$$\begin{aligned} \mathcal{E}_{\widehat{\delta}, \epsilon} + \epsilon_{\mathcal{H}}\mathcal{E}_{\mathcal{H}, \epsilon} &\lesssim C\widehat{\delta}^{-1}u_{trap}^{-1-\delta_{dec}} \left[r^{-2}|e_3\Psi|^2 + r^{-1}(|e_4\Psi|^2 + |\check{\nabla}\Psi|^2) \right] \\ &+ C\widehat{\delta}^{-1}u_{trap}^{-1-\delta_{dec}} r^{-1}|e_3\Psi| (|e_4\Psi| + |\check{\nabla}\Psi|) + C\widehat{\delta}^{-1}u_{trap}^{-1-\delta_{dec}} r^{-3}|\Psi|^2. \end{aligned}$$

Proof. It only remains to check the last part. In view of Proposition 10.1.20 we have,

$$\mathcal{E}_{\widehat{\delta}, \epsilon} = \mathcal{E}_{\epsilon}[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR](\Psi) = \epsilon \frac{1}{2} \mathcal{Q} \cdot {}^{(X_{\widehat{\delta}})}\ddot{\pi} + O(\epsilon r^{-3}u_{trap}^{-1-\delta_{dec}}(|f_{\widehat{\delta}}| + 1))|\Psi|^2$$

and $|f_{\widehat{\delta}}| \lesssim \widehat{\delta}^{-1}$. Hence,

$$\mathcal{E}_{\widehat{\delta}, \epsilon} = \mathcal{E}_{\epsilon}[f_{\widehat{\delta}}R, w_{\widehat{\delta}}, 2hR](\Psi) = \epsilon \left(\frac{1}{2} \mathcal{Q} \cdot {}^{X_{\widehat{\delta}}}\ddot{\pi} + O(\widehat{\delta}^{-1}r^{-3}u_{trap}^{-1-\delta_{dec}})|\Psi|^2 \right).$$

We write with $\ddot{\pi} = {}^{(X_\delta)} \ddot{\pi}$ for simplicity,

$$\mathcal{Q} \cdot \ddot{\pi} = \frac{1}{4} (\mathcal{Q}_{33} \ddot{\pi}_{44} + 2\mathcal{Q}_{34} \ddot{\pi}_{34} + \mathcal{Q}_{44} \ddot{\pi}_{33}) - \frac{1}{2} (\mathcal{Q}_{A3} \ddot{\pi}_{A4} + \mathcal{Q}_{A4} \ddot{\pi}_{A3}) + \mathcal{Q}_{AB} \ddot{\pi}_{AB}.$$

Thus, recalling part 1 and 2 of Proposition 10.1.9, and Lemma 10.1.8,

$$\begin{aligned} \mathcal{Q} \cdot \ddot{\pi} &\lesssim r^{-2} u_{trap}^{-1-\delta_{dec}} |e_3 \Psi|^2 + r^{-1} u_{trap}^{-1-\delta_{dec}} (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2) \\ &\quad + r^{-1} u_{trap}^{-1-\delta_{dec}} |e_3 \Psi| (|e_4 \Psi| + |\nabla \Psi|). \end{aligned}$$

Finally, since $r \sim 2m$ and $u_{trap} = 1$ on $|\Upsilon| \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}$, the error terms generated by the red shift vectorfield $Y_{\mathcal{H}}$,

$$\mathcal{E}_{\mathcal{H},\epsilon} = O(\epsilon) 1_{|\Upsilon| \leq 2\delta_{\mathcal{H}}^{\frac{1}{10}}} (|\mathcal{Q}(\Psi)| + m^{-2} |\Psi|^2)$$

can easily be absorbed on the right hand side to derive the desired estimate. □

Elimination of \overline{W}_δ

We now proceed to eliminate the potential \overline{W}_δ by a procedure analogous to that used in section 10.1.8. More precisely we set, in view of (10.1.13),

$$\mathcal{E}_\delta = \mathcal{E}[f_\delta R, w_\delta, 2hR](\Psi), \quad \mathcal{E}'_\delta = \mathcal{E}[f_\delta R, w_\delta, 2(hR + h_2 \check{R})](\Psi),$$

and,

$$\mathcal{E}'_\delta = \mathcal{E}_\delta + h_2 \Psi \check{R} \Psi + \frac{1}{2} \mathbf{D}^\mu (h_2 \check{R}_\mu) |\Psi|^2,$$

where h_2 is a smooth, compactly supported function supported⁷ in the region $r \leq \frac{9m}{4}$.

Thus, we have in view of Proposition 10.1.30, ignoring the integration on S ,

$$\begin{aligned} \dot{\mathcal{E}}'_\delta + \epsilon_{\mathcal{H}} \dot{\mathcal{E}}_{\mathcal{H}} &= \dot{\mathcal{E}}_\delta + \epsilon_{\mathcal{H}} \dot{\mathcal{E}}_{\mathcal{H}} + h_2 \Psi \check{R} \Psi + \frac{1}{2} \mathbf{D}^\mu (h_2 \check{R}_\mu) |\Psi|^2 \\ &\geq \mathcal{I}(\Psi) + m^{-1} \Lambda_{\mathcal{H}}^{-1} \left(\frac{1}{2} |\check{R}(\Psi)|^2 + |\check{T}\Psi|^2 + |\nabla \Psi|^2 + m^{-2} |\Psi|^2 \right) \end{aligned} \tag{10.1.56}$$

where,

$$\mathcal{I}(\Psi) := \frac{1}{2} \Lambda_{\mathcal{H}}^{-1} m^{-1} |\check{R}(\Psi)|^2 + \Psi h_2 \check{R} \Psi + \frac{1}{2} \mathbf{D}^\mu (h_2 \check{R}_\mu) |\Psi|^2 - \overline{W}_\delta |\Psi|^2$$

⁷Recall that \overline{W}_δ is supported in the region $2m < r \leq \frac{5m}{2}$.

so that we have

$$\mathcal{I}(\Psi) \geq \frac{1}{2} \left[\mathbf{D}^\mu(h_2 \check{R}_\mu) - 2\overline{W}_\delta - m\Lambda_{\mathcal{H}} h_2^2 \right] |\Psi|^2. \quad (10.1.57)$$

We focus on the coefficient in front of $|\Psi|^2$ on the RHS of (10.1.57). Ignoring the error terms in ϵ (which can easily be incorporated in the upper bound for $\mathcal{E}_{\delta, \epsilon} + \epsilon_{\mathcal{H}} \mathcal{E}_{\mathcal{H}, \epsilon}$ of the previous proposition), we have,

$$\mathbf{Div} \check{R} = \frac{1}{2} \left(\mathbf{D}^\mu(\check{\theta}(e_4)_\mu) - \mathbf{D}^\mu((e_3)_\mu) \right) = \frac{1}{4} \left(\check{\theta} \text{tr} \pi^{(4)} - \text{tr} \pi^{(3)} \right) + \frac{1}{2} e_4(\check{\theta}) = O(\delta_{\mathcal{H}}^{-\frac{1}{10}})$$

and, using in particular Lemma 10.1.5,

$$\begin{aligned} \mathbf{D}^\mu(h_2 \check{R}_\mu) &= \check{R} h_2 + h_2 \mathbf{Div} \check{R} = \frac{1}{2} \partial_r h_2 (\check{\theta} e_4 r - e_3 r) + h_2 \mathbf{Div} \check{R} \\ &= \frac{1}{2} \partial_r h_2 (\check{\theta} \Upsilon + 1) + h_2 O(\delta_{\mathcal{H}}^{-\frac{1}{10}}) \\ &\geq \frac{1}{2} \partial_r h_2 + h_2 O(\delta_{\mathcal{H}}^{-\frac{1}{10}}). \end{aligned}$$

Together with (10.1.57), we infer

$$\mathcal{I}(\Psi) \geq \frac{1}{4} \left[\partial_r h_2 - 4\overline{W}_\delta - 4m\Lambda_{\mathcal{H}} h_2^2 + h_2 O(\delta_{\mathcal{H}}^{-\frac{1}{10}}) \right] |\Psi|^2. \quad (10.1.58)$$

We now consider the choice of the function $h_2 = h_2(r, m)$. Recall (see Lemma 10.1.24) that \overline{W}_δ is supported in the region $2m + e^{-\frac{2}{\delta}} \leq r \leq \frac{9m}{4}$ and that its primitive $\widetilde{W}_\delta(r) := \int_{2m}^r \overline{W}_\delta$ verifies $\widetilde{W}_\delta \lesssim m^{-2} \widehat{\delta}$. We choose

$$h_2 =: \begin{cases} 4\widetilde{W}_\delta, & \text{for } r \leq \frac{9m}{4} \\ 0, & \text{for } r \geq \frac{5m}{2} \end{cases} \quad (10.1.59)$$

and since $\widetilde{W}_\delta \lesssim m^{-2} \widehat{\delta}$, we may extend h_2 in $\frac{9m}{4} \leq r \leq \frac{5m}{2}$ such that h_2 is C^1 and we have for all $r > 0$

$$|h_2| \lesssim m^{-2} \widehat{\delta}, \quad |\partial_r h_2| \lesssim m^{-3} \widehat{\delta}. \quad (10.1.60)$$

In view of (10.1.58), this choice of h_2 yields

$$\mathcal{I}(\Psi) \geq -\frac{1}{4} O\left(m^{-1} \widehat{\delta} + \Lambda_{\mathcal{H}} m^{-1} \widehat{\delta}^2 + \widehat{\delta} (\delta_{\mathcal{H}})^{-\frac{1}{10}}\right) |\Psi|^2.$$

Hence, for $\widehat{\delta} \ll \delta_{\mathcal{H}}^{\frac{1}{10}} \Lambda_{\mathcal{H}}^{-1}$, i.e. $\widehat{\delta} \ll \delta_{\mathcal{H}}^{\frac{1}{2}}$ (recall that $\Lambda_{\mathcal{H}}^{-1} = C^{-1} \delta_{\mathcal{H}}^{\frac{2}{5}}$) and h_2 defined as above, we infer

$$\mathcal{I}(\Psi) \geq -\frac{1}{2} m^{-1} \Lambda_{\mathcal{H}}^{-1} m^{-2} |\Psi|^2$$

which together with (10.1.56) finally yields

$$\begin{aligned} \int_S (\dot{\mathcal{E}}'_\delta + \epsilon_{\mathcal{H}} \dot{\mathcal{E}}_{\mathcal{H}}) &\geq \int_S \mathcal{I}(\Psi) + m^{-1} \Lambda_{\mathcal{H}}^{-1} \int_S \left(\frac{1}{2} |\check{R}(\Psi)|^2 + |\check{T}\Psi|^2 + |\check{\nabla}\Psi|^2 + m^{-2} |\Psi|^2 \right) \\ &\geq m^{-1} \Lambda_{\mathcal{H}}^{-1} \int_S \left(\frac{1}{2} |\check{R}(\Psi)|^2 + |\check{T}\Psi|^2 + |\check{\nabla}\Psi|^2 + \frac{1}{2} m^{-2} |\Psi|^2 \right). \end{aligned}$$

Summary of results so far

We summarize the result in the following,

Proposition 10.1.31. *Consider the combined Morawetz triplet*

$$(X, w, M) := (f_{\widehat{\delta}} R, w_{\widehat{\delta}}, 2hR) + \epsilon_{\mathcal{H}}(Y_{\mathcal{H}}, 0, 0) + (0, 0, 2h_2 \check{R}) \tag{10.1.61}$$

with $(f_{\widehat{\delta}} R, w_{\widehat{\delta}}, 2hR)$ the triplet of Proposition 10.1.25, $Y_{\mathcal{H}}$ the red shift vectorfield of Proposition 10.1.29 (corresponding to the small parameter $\delta_{\mathcal{H}}$) and h_2 the C^1 function above satisfying (10.1.59) (10.1.60). Let $\dot{\mathcal{E}}[X, w, M]$ the principal part of $\mathcal{E}[X, w, M]$ (independent of ϵ) and $\mathcal{E}_\epsilon[X, w, M]$ the error term in ϵ such that $\mathcal{E} = \dot{\mathcal{E}} + \mathcal{E}_\epsilon$.

We choose the small strictly positive parameters $\epsilon_{\mathcal{H}}, \delta_{\mathcal{H}}, \delta$ such that⁸,

$$\epsilon_{\mathcal{H}} = \delta_{\mathcal{H}}^{\frac{7}{20}}, \quad \widehat{\delta} = \delta_{\mathcal{H}}^{\frac{3}{5}}. \tag{10.1.62}$$

Then, there holds⁹

$$\int_S \mathcal{E}[X, w, M](\Psi) \geq \delta_{\mathcal{H}}^{\frac{1}{2}} \int_S \left[\frac{m^2}{r^3} |\check{R}\Psi|^2 + r^{-1} \left(1 - \frac{3m}{r} \right)^2 \left(\widehat{\mathcal{Q}}_{34} + \frac{m^2}{r^2} |\check{T}\Psi|^2 \right) + \frac{m}{r^4} |\Psi|^2 \right], \tag{10.1.63}$$

$$\begin{aligned} \mathcal{E}_\epsilon[X, w, M](\Psi) &\leq \delta_{\mathcal{H}}^{-1} \epsilon u_{trap}^{-1-\delta_{dec}} \left[r^{-2} |e_3 \Psi|^2 + r^{-1} (|e_4 \Psi|^2 + |\check{\nabla} \Psi|^2) \right] \\ &\quad + \delta_{\mathcal{H}}^{-1} \epsilon u_{trap}^{-1-\delta_{dec}} r^{-1} |e_3 \Psi| (|e_4 \Psi| + |\check{\nabla} \Psi|) + \delta_{\mathcal{H}}^{-1} \epsilon u_{trap}^{-1-\delta_{dec}} r^{-3} |\Psi|^2. \end{aligned}$$

⁸Note that (10.1.62) verifies all the restrictions we have encountered so far, i.e. $\delta_{\mathcal{H}}^{\frac{2}{5}} \ll \epsilon_{\mathcal{H}} \ll \delta_{\mathcal{H}}^{\frac{3}{10}}$ and $0 < \widehat{\delta} \ll \delta_{\mathcal{H}}^{\frac{1}{2}}$.

⁹Note that $\delta_{\mathcal{H}}^{\frac{1}{2}} \ll \Lambda_{\mathcal{H}}^{-1}$ (recall that $\Lambda_{\mathcal{H}}^{-1} = C^{-1} \delta_{\mathcal{H}}^{\frac{2}{5}}$) and $\delta_{\mathcal{H}}^{-1} \gg \widehat{\delta}^{-1}$ in view of (10.1.62).

10.1.12 Lower bounds for \mathcal{Q}

In this section we prove lower bounds for $\mathcal{Q}(X + 2\Lambda T, e_3)$ and $\mathcal{Q}(X + 2\Lambda T, e_4)$ in the region $r_{\mathcal{H}} \leq r$, for $r_{\mathcal{H}}$ to be determined and Λ sufficiently large.

Proposition 10.1.32. *Under the assumptions of Proposition 10.1.31, and with the choice*

$$\Lambda := \frac{1}{4} \delta_{\mathcal{H}}^{-\frac{13}{20}}, \quad (10.1.64)$$

the following inequalities hold true for $r \geq 2m_0(1 - \delta_{\mathcal{H}})$.

1. For the region such that $r \geq 2m_0(1 - \delta_{\mathcal{H}})$ and $\Upsilon \leq \delta_{\mathcal{H}}^{\frac{1}{10}}$, we have

$$\begin{aligned} \mathcal{Q}(X + \Lambda T, e_3) &\geq \frac{1}{4} \epsilon_{\mathcal{H}} \mathcal{Q}_{33} + \frac{1}{2} \Lambda \mathcal{Q}_{34}, \\ \mathcal{Q}(X + \Lambda T, e_4) &\geq \frac{1}{4} \epsilon_{\mathcal{H}} \mathcal{Q}_{34} + \frac{1}{2} \Lambda \mathcal{Q}_{44}. \end{aligned}$$

2. For the region $\delta_{\mathcal{H}}^{\frac{1}{10}} \leq \Upsilon \leq \frac{1}{3}$, we have

$$\begin{aligned} \mathcal{Q}(X + \Lambda T, e_3) &\geq \delta_{\mathcal{H}}^{-\frac{1}{2}} (\mathcal{Q}_{33} + \mathcal{Q}_{34}), \\ \mathcal{Q}(X + \Lambda T, e_4) &\geq \delta_{\mathcal{H}}^{-\frac{1}{2}} (\mathcal{Q}_{44} + \mathcal{Q}_{34}). \end{aligned}$$

3. For the region $r \geq 3m$, we have

$$\begin{aligned} \mathcal{Q}(X + \Lambda T, e_3) &\geq \frac{1}{4} \Lambda (\mathcal{Q}_{33} + \mathcal{Q}_{34}), \\ \mathcal{Q}(X + \Lambda T, e_4) &\geq \frac{1}{4} \Lambda (\mathcal{Q}_{44} + \mathcal{Q}_{34}). \end{aligned}$$

4. The null components of \mathcal{Q} are given by (recall Proposition 10.1.9),

$$\mathcal{Q}_{33} = |e_3 \Psi|^2, \quad \mathcal{Q}_{44} = |e_4 \Psi|^2, \quad \mathcal{Q}_{34} = |\nabla \Psi|^2 + \frac{4\Upsilon}{r^2} (1 + O(\epsilon)) |\Psi|^2.$$

Proof. Since $X = f_{\hat{\delta}} R + \epsilon_{\mathcal{H}} Y_{\mathcal{H}}$ and $T = \frac{1}{2}(e_4 + \Upsilon e_3)$, $R = \frac{1}{2}(e_4 - \Upsilon e_3)$, we write,

$$\begin{aligned} \mathcal{Q}(X + 2\Lambda T, e_3) &= \mathcal{Q}(X, e_3) + \Lambda \mathcal{Q}(e_4 + \Upsilon e_3, e_3) = \mathcal{Q}(X, e_3) + \Lambda (\mathcal{Q}_{34} + \Upsilon \mathcal{Q}_{33}) \\ &= \epsilon_{\mathcal{H}} \mathcal{Q}(Y_{\mathcal{H}}, e_3) + \Lambda (\mathcal{Q}_{34} + \Upsilon \mathcal{Q}_{33}) + \frac{1}{2} f_{\hat{\delta}} (\mathcal{Q}_{34} - \Upsilon \mathcal{Q}_{33}). \end{aligned}$$

In the region $2m_0(1 - \delta_{\mathcal{H}}) \leq r \leq 2m$ we have $Y_{\mathcal{H}} = e_3 + e_4 + O(\delta_{\mathcal{H}})(e_3 + e_4)$, $\Upsilon \geq 0$ and $f_{\hat{\delta}} < 0$. Hence, in that region,

$$\begin{aligned} \mathcal{Q}(X + 2\Lambda T, e_3) &\geq \frac{1}{2}\epsilon_{\mathcal{H}}(\mathcal{Q}_{33} + \mathcal{Q}_{34}) + \left(\Lambda - \frac{1}{2}|f_{\hat{\delta}}|\right)\mathcal{Q}_{34} - |\Upsilon| \left(\Lambda + \frac{1}{2}|f_{\hat{\delta}}|\right)\mathcal{Q}_{33} \\ &\geq \left(\frac{1}{2}\epsilon_{\mathcal{H}} - |\Upsilon| \left(\Lambda + \frac{1}{2}|f_{\hat{\delta}}|\right)\right)\mathcal{Q}_{33} + \left(\frac{1}{2}\epsilon_{\mathcal{H}} + \Lambda - \frac{1}{2}|f_{\hat{\delta}}|\right)\mathcal{Q}_{34}. \end{aligned}$$

Thus, we need to choose Λ such that

$$\frac{1}{2}|f_{\hat{\delta}}| \leq \Lambda \leq \frac{1}{4}\frac{\epsilon_{\mathcal{H}}}{\delta_{\mathcal{H}}} - \frac{1}{2}|f_{\hat{\delta}}|$$

Now, recall (10.1.62) as well as the fact that $|f_{\hat{\delta}}|$ is of size $O((\hat{\delta})^{-1})$. Thus it suffices to choose Λ such that

$$O(\delta_{\mathcal{H}}^{-\frac{3}{5}}) \leq \Lambda \leq \frac{1}{2}\delta_{\mathcal{H}}^{-\frac{13}{20}} - O(\delta_{\mathcal{H}}^{-\frac{3}{5}}),$$

i.e. it suffices to choose, for $\delta_{\mathcal{H}} > 0$ small enough,

$$\Lambda = \frac{1}{4}\delta_{\mathcal{H}}^{-\frac{13}{20}},$$

to deduce the inequality,

$$\mathcal{Q}(X + 2\Lambda T, e_3) \geq \frac{1}{4}\epsilon_{\mathcal{H}}\mathcal{Q}_{33} + \frac{1}{2}\Lambda\mathcal{Q}_{34}.$$

Next, in the region $0 \leq \Upsilon \leq \delta_{\mathcal{H}}^{\frac{1}{10}}$, the sign of Υ is more favorable and we have

$$\begin{aligned} \mathcal{Q}(X + 2\Lambda T, e_3) &\geq \frac{1}{2}\epsilon_{\mathcal{H}}(\mathcal{Q}_{33} + \mathcal{Q}_{34}) + \left(\Lambda - \frac{1}{2}|f_{\hat{\delta}}|\right)\mathcal{Q}_{34} + |\Upsilon| \left(\Lambda + \frac{1}{2}|f_{\hat{\delta}}|\right)\mathcal{Q}_{33} \\ &\geq \left(\frac{1}{2}\epsilon_{\mathcal{H}} + |\Upsilon| \left(\Lambda + \frac{1}{2}|f_{\hat{\delta}}|\right)\right)\mathcal{Q}_{33} + \left(\frac{1}{2}\epsilon_{\mathcal{H}} + \Lambda - \frac{1}{2}|f_{\hat{\delta}}|\right)\mathcal{Q}_{34}. \end{aligned}$$

In particular, we simply need $\Lambda \gg \hat{\delta}^{-1}$, which is in particular satisfied by (10.1.64), to deduce the same inequality,

$$\mathcal{Q}(X + 2\Lambda T, e_3) \geq \frac{1}{4}\epsilon_{\mathcal{H}}\mathcal{Q}_{33} + \frac{1}{2}\Lambda\mathcal{Q}_{34}.$$

In the region $\delta_{\mathcal{H}}^{\frac{1}{10}} \leq \Upsilon \leq \frac{1}{3}$, where $f_{\hat{\delta}} \leq 0$, and using the fact that $|f_{\hat{\delta}}|$ is of size $O((\hat{\delta})^{-1})$

$$\begin{aligned} \mathcal{Q}(X + 2\Lambda T, e_3) &= \epsilon_{\mathcal{H}}\mathcal{Q}(Y_{\mathcal{H}}, e_3) + \Lambda(\mathcal{Q}_{34} + \Upsilon\mathcal{Q}_{33}) + \frac{1}{2}f_{\hat{\delta}}(\mathcal{Q}_{34} - \Upsilon\mathcal{Q}_{33}) \\ &\geq \Lambda\left(\mathcal{Q}_{34} + \delta_{\mathcal{H}}^{\frac{1}{10}}\mathcal{Q}_{33}\right) - O(\hat{\delta}^{-1})\mathcal{Q}_{34}. \end{aligned}$$

Hence, for the choice (10.1.64), and in view of (10.1.62), we infer

$$\mathcal{Q}(X + 2\Lambda T, e_3) \geq \delta_{\mathcal{H}}^{-\frac{1}{2}} (\mathcal{Q}_{33} + \mathcal{Q}_{34}).$$

Finally, for $r \geq 3m$ where we have $0 \leq f_{\hat{\delta}} \lesssim 1$, $\frac{1}{3} \leq \Upsilon \leq 1$ and $Y_{\mathcal{H}} = 0$,

$$\begin{aligned} \mathcal{Q}(X + 2\Lambda T, e_3) &= \Lambda (\mathcal{Q}_{34} + \Upsilon \mathcal{Q}_{33}) + \frac{1}{2} f_{\hat{\delta}} (\mathcal{Q}_{34} - \Upsilon \mathcal{Q}_{33}) \\ &\geq \Lambda \left(\mathcal{Q}_{34} + \frac{1}{3} \mathcal{Q}_{33} \right) - O(1) \mathcal{Q}_{33} \end{aligned}$$

and hence, (10.1.64) implies

$$\mathcal{Q}(X + 2\Lambda T, e_3) \geq \frac{1}{4} \Lambda (\mathcal{Q}_{34} + \mathcal{Q}_{33})$$

as desired. The proof for $\mathcal{Q}(X + \Lambda T, e_4)$ is similar. \square

10.1.13 First Morawetz Estimate

We are now ready to state our first Morawetz estimate which is simply obtained by integrating the pointwise inequality in Proposition 10.1.30 on our domain $\mathcal{M} = {}^{(int)}\mathcal{M} \cup {}^{(ext)}\mathcal{M}$ described at the beginning of the section, with X replaced by $X + \Lambda T$ for $\Lambda > 0$ sufficiently large. In view of the choice of τ , note that we have

$$N_{\Sigma} = ae_3 + be_4, \quad 0 \leq a, b \leq 1, \quad a + b \geq 1, \quad (10.1.65)$$

with

$$b = 0, a = 1 \text{ on } {}^{(int)}\mathcal{M}, \quad a = 0, b = 1 \text{ on } \mathcal{M}_{r \geq 4m_0}, \quad a, b \geq \frac{1}{4} \text{ on } {}^{(trap)}\mathcal{M}.$$

We recall the following quantities for Ψ in regions $\mathcal{M}(\tau_1, \tau_2) \subset \mathcal{M}$ in the past of $\Sigma(\tau_2)$ and future of $\Sigma(\tau_1)$.

1. Morawetz bulk quantity

$$\text{Mor}[\Psi](\tau_1, \tau_2) = \int_{\mathcal{M}(\tau_1, \tau_2)} \frac{m^2}{r^3} |\check{R}\psi|^2 + \frac{m}{r^4} |\Psi|^2 + \left(1 - \frac{3m}{r}\right)^2 \frac{1}{r} \left(|\nabla\psi|^2 + \frac{m^2}{r^2} |\check{T}\psi|^2 \right).$$

2. Basic energy quantity

$$E[\Psi](\tau) = \int_{\Sigma(\tau)} \left(\frac{1}{2} (N_{\Sigma}, e_3)^2 |e_4\Psi|^2 + \frac{1}{2} (N_{\Sigma}, e_4)^2 |e_3\Psi|^2 + |\nabla\Psi|^2 + r^{-2} |\Psi|^2 \right).$$

3. Flux through \mathcal{A} and Σ_*

$$F[\Psi](\tau_1, \tau_2) = \int_{\mathcal{A}(\tau_1, \tau_2)} (\delta_{\mathcal{H}}^{-1}|e_4\Psi|^2 + \delta_{\mathcal{H}}|e_3\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2) + \int_{\Sigma_*(\tau_1, \tau_2)} (|e_4\Psi|^2 + |e_3\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2),$$

with $\mathcal{A}(\tau_1, \tau_2) = \mathcal{A} \cap \mathcal{M}(\tau_1, \tau_2)$ and $\Sigma_*(\tau_1, \tau_2) = \Sigma_* \cap \mathcal{M}(\tau_1, \tau_2)$.

The following theorem is our first Morawetz estimate.

Theorem 10.1.33. *Consider the equation (10.1.9), i.e. $\dot{\square}\Psi = V\Psi + \mathcal{N}$, with $V = -\kappa\underline{\kappa}$ and a domain $\mathcal{M}(\tau_1, \tau_2) \subset \mathcal{M}$. Then, we have*

$$E[\Psi](\tau_2) + Mor[\Psi](\tau_1, \tau_2) + F[\Psi](\tau_1, \tau_2) \lesssim (E[\Psi](\tau_1) + J[N, \Psi](\tau_1, \tau_2) + Err_\epsilon(\tau_1, \tau_2)[\Psi]),$$

$$J[N, \Psi](\tau_1, \tau_2) := \int_{\mathcal{M}(\tau_1, \tau_2)} (|\check{R}\Psi| + |\check{T}\Psi| + r^{-1}|\Psi|)|N|, \tag{10.1.66}$$

$$Err_\epsilon[\Psi](\tau_1, \tau_2) = \int_{\mathcal{M}(\tau_1, \tau_2)} \mathcal{E}_\epsilon[\Psi],$$

where,

$$\mathcal{E}_\epsilon[\Psi] \lesssim \epsilon u_{trap}^{-1-\delta_{dec}} [r^{-2}|e_3\Psi|^2 + r^{-1}(|e_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2 + |e_3\Psi|(|e_4\Psi| + |\nabla\Psi|))].$$

Proof. Recall that, see (10.1.12)

$$\mathcal{E}[X, w, M](\Psi) := \mathbf{D}^\mu \mathcal{P}_\mu[X, w, M] - \left(X(\Psi) + \frac{1}{2}w\Psi \right) \cdot \mathcal{N}[\Psi]$$

where,

$$\mathcal{P}_\mu = \mathcal{P}_\mu[X, w, M] = \mathcal{Q}_{\mu\nu}X^\nu + \frac{1}{2}w\Psi\dot{\mathbf{D}}_\mu\Psi - \frac{1}{4}|\Psi|^2\partial_\mu w + \frac{1}{4}|\Psi|^2M_\mu$$

with triplet,

$$(X, w, M) := (f_{\check{\delta}}R, w_{\check{\delta}}, 2hR) + \epsilon_{\mathcal{H}}(Y_{\mathcal{H}}, 0, 0) + (0, 0, 2h_2\check{R})$$

given in Proposition 10.1.30. Replacing X by $\check{X} = X + \Lambda T$ in the calculation above we deduce,

$$\check{\mathcal{P}}_\mu = \mathcal{P}_\mu[\check{X}, w, M] = \mathcal{Q}_{\mu\nu}\check{X}^\nu + \frac{1}{2}w\Psi\dot{\mathbf{D}}_\mu\Psi - \frac{1}{4}|\Psi|^2\partial_\mu w + \frac{1}{4}|\Psi|^2M_\mu.$$

By the divergence theorem we have,

$$\begin{aligned} & \int_{\mathcal{A}} \check{\mathcal{P}} \cdot N_{\mathcal{A}} + \int_{\Sigma_2} \check{\mathcal{P}} \cdot N_{\Sigma} + \int_{\mathcal{M}(\tau_1, \tau_2)} \mathcal{E} + \int_{\Sigma_*} \check{\mathcal{P}} \cdot N_{\Sigma_*} = \int_{\Sigma_1} \check{\mathcal{P}} \cdot N_{\Sigma} \\ & - \int_{\mathcal{M}(\tau_1, \tau_2)} \left(\check{X}(\Psi) + \frac{1}{2} w \Psi \right) N[\Psi] \end{aligned} \quad (10.1.67)$$

where $\mathcal{E} = \mathcal{E}[\check{X}, w, M](\Psi)$. Now,

$$\mathcal{E}[\check{X}, w, M](\Psi) = \mathcal{E}[X, w, M](\Psi) + \frac{1}{2} \Lambda \mathcal{Q} \cdot {}^{(T)}\pi - \frac{1}{2} T(V) |\Psi|^2.$$

According to Lemma 10.1.7 $T(V) = O(\epsilon) r^{-3} u_{trap}^{-1-\delta_{dec}}$, and all components of ${}^{(T)}\pi$ are $O(\epsilon r^{-1} u_{trap}^{-1-\delta_{dec}})$ except for ${}^{(T)}\pi_{44}$ which is $O(\epsilon r^{-2} u_{trap}^{-1-\delta_{dec}})$. We easily deduce,

$$\Lambda |\mathcal{Q} \cdot {}^{(T)}\pi| + |T(V)| |\Psi|^2 \lesssim \Lambda \mathcal{E}_{\epsilon}.$$

Thus in view of to Proposition 10.1.31, we have¹⁰,

$$\begin{aligned} \int_{\mathcal{M}(\tau_1, \tau_2)} \mathcal{E} & \geq \delta_{\mathcal{H}}^{\frac{1}{2}} \int_{\mathcal{M}(\tau_1, \tau_2)} \left[\frac{m^2}{r^3} |\check{R}\Psi|^2 + r^{-1} \left(1 - \frac{3m}{r} \right)^2 \left(|\nabla\Psi|^2 + \frac{m^2}{r^2} |\check{T}\Psi|^2 \right) + \frac{m}{r^4} |\Psi|^2 \right] \\ & - O(\delta_{\mathcal{H}}^{-1}) \int_{\mathcal{M}(\tau_1, \tau_2)} \mathcal{E}_{\epsilon} \end{aligned}$$

i.e.,

$$\int_{\mathcal{M}(\tau_1, \tau_2)} \mathcal{E} \geq \delta_{\mathcal{H}}^{\frac{1}{2}} \text{Mor}[\Psi](\tau_1, \tau_2) - O(\delta_{\mathcal{H}}^{-1}) \text{Err}_{\epsilon}(\tau_1, \tau_2). \quad (10.1.68)$$

We now analyze the boundary terms in (10.1.67).

Boundary term along \mathcal{A}

Along the spacelike hypersurface \mathcal{A} , i.e. $r = 2m_0(1 - \delta_{\mathcal{H}})$, the unit normal $N_{\mathcal{A}}$ is given by

$$\begin{aligned} N_{\mathcal{A}} & = \frac{1}{2\sqrt{\frac{e_4(r)}{e_3(r)}}} \left(e_4 + \frac{e_4(r)}{e_3(r)} e_3 \right) \\ & = \frac{1}{2\sqrt{\delta_{\mathcal{H}} + O(\epsilon)}} \left(e_4 + (\delta_{\mathcal{H}} + O(\epsilon)) e_3 \right), \end{aligned}$$

¹⁰Recall from (10.1.64) that we have $\Lambda = \frac{1}{4} \delta_{\mathcal{H}}^{-\frac{13}{20}} \ll \delta_{\mathcal{H}}^{-1}$.

and we have $h, h_2 = 0$ as well as $w = -\delta_1 w_1$ where $\delta_1 > 0$ is a small constant and w_1 is given by (10.1.46)

$$w_1(r, m) = r^{-1} \frac{m^2}{r^2} \Upsilon \left(1 - \frac{3m}{r} \right)^2.$$

In particular, we have on \mathcal{A} in view of the formula for w_1 and for $N_{\mathcal{A}}$

$$|w_1| \lesssim \delta_{\mathcal{H}}, \quad |N_{\mathcal{A}}(w_1)| \lesssim \sqrt{\delta_{\mathcal{H}}}.$$

Hence,

$$\begin{aligned} \mathcal{P} \cdot N_{\mathcal{A}} &= \mathcal{Q}(X + \Lambda T, N_{\mathcal{A}}) - \frac{\delta_1}{2} w_1 \Psi N_{\mathcal{A}}(\Psi) + \frac{\delta_1}{4} |\Psi|^2 N_{\mathcal{A}}(w_1) \\ &= \frac{2}{\sqrt{\delta_{\mathcal{H}} + O(\epsilon)}} \mathcal{Q}(X + \Lambda T, e_4) + 2\sqrt{\delta_{\mathcal{H}} + O(\epsilon)} \mathcal{Q}(X + \Lambda T, e_3) \\ &\quad - O(\sqrt{\delta_{\mathcal{H}}}) \Psi e_4(\Psi) - O(\delta_{\mathcal{H}}^{\frac{3}{2}}) \Psi e_3(\Psi) - O(\sqrt{\delta_{\mathcal{H}}}) |\Psi|^2. \end{aligned}$$

Thus, in view of Proposition 10.1.32, we infer

$$\begin{aligned} \mathcal{P} \cdot N_{\mathcal{A}} &\geq \frac{2}{\sqrt{\delta_{\mathcal{H}} + O(\epsilon)}} \left(\frac{1}{4} \epsilon_{\mathcal{H}} \mathcal{Q}_{34} + \frac{1}{2} \Lambda \mathcal{Q}_{44} \right) + 2\sqrt{\delta_{\mathcal{H}} + O(\epsilon)} \left(\frac{1}{4} \epsilon_{\mathcal{H}} \mathcal{Q}_{33} + \frac{1}{2} \Lambda \mathcal{Q}_{34} \right) \\ &\quad - O(\sqrt{\delta_{\mathcal{H}}}) \Psi e_4(\Psi) - O(\delta_{\mathcal{H}}^{\frac{3}{2}}) \Psi e_3(\Psi) - O(\sqrt{\delta_{\mathcal{H}}}) |\Psi|^2 \end{aligned}$$

Using in particular (10.1.62) and (10.1.64), we deduce

$$\begin{aligned} \mathcal{P} \cdot N_{\mathcal{A}} &\geq \frac{2}{\sqrt{\delta_{\mathcal{H}} + O(\epsilon)}} \left(\frac{1}{4} \delta_{\mathcal{H}}^{\frac{7}{20}} \left(|\nabla \Psi|^2 + O(\delta_{\mathcal{H}}) |\Psi|^2 \right) + \frac{1}{8} \delta_{\mathcal{H}}^{-\frac{13}{20}} |e_4 \Psi|^2 \right) \\ &\quad + 2\sqrt{\delta_{\mathcal{H}} + O(\epsilon)} \left(\frac{1}{4} \delta_{\mathcal{H}}^{\frac{7}{20}} |e_3 \Psi|^2 + \frac{1}{8} \delta_{\mathcal{H}}^{-\frac{13}{20}} \left(|\nabla \Psi|^2 + O(\delta_{\mathcal{H}}) |\Psi|^2 \right) \right) \\ &\quad - O(\sqrt{\delta_{\mathcal{H}}}) \Psi e_4(\Psi) - O(\delta_{\mathcal{H}}^{\frac{3}{2}}) \Psi e_3(\Psi) - O(\sqrt{\delta_{\mathcal{H}}}) |\Psi|^2 \\ &\geq \frac{1}{2} \delta_{\mathcal{H}}^{-\frac{3}{20}} |\nabla \Psi|^2 + \frac{1}{8} \delta_{\mathcal{H}}^{-\frac{23}{20}} |e_4 \Psi|^2 + \frac{1}{4} \delta_{\mathcal{H}}^{\frac{17}{20}} |e_3 \Psi|^2 \\ &\quad - O(\sqrt{\delta_{\mathcal{H}}}) \Psi e_4(\Psi) - O(\delta_{\mathcal{H}}^{\frac{3}{2}}) \Psi e_3(\Psi) - O(\sqrt{\delta_{\mathcal{H}}}) |\Psi|^2. \end{aligned}$$

Recalling the Poincaré inequality (10.1.26),

$$\int_S |\nabla \Psi|^2 \geq 2r^{-2} (1 - O(\epsilon)) \int_S \Psi^2 da_S,$$

we deduce, in this region,

$$\int_{\mathcal{A}(\tau_1, \tau_2)} \mathcal{P} \cdot N_{\mathcal{A}} \geq \frac{1}{8} \int_{\mathcal{A}(\tau_1, \tau_2)} \left(\delta_{\mathcal{H}}^{-1} |e_4 \Psi|^2 + \delta_{\mathcal{H}} |e_3 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2 \right)$$

as desired in view of the definition of the flux along \mathcal{A} .

Boundary terms along $\Sigma(\tau_1), \Sigma(\tau_2)$

Along a hypersurface $\Sigma(\tau)$ with timelike unit future normal $N_{\Sigma(\tau)} = ae_3 + be_4$, we have,

$$\mathcal{P} \cdot N_{\Sigma} = \mathcal{Q}(X + \Lambda T, N_{\Sigma}) + \frac{1}{2}w\Psi N_{\Sigma}(\Psi) - \frac{1}{4}N_{\Sigma}(w)|\Psi|^2 + \frac{1}{2}N_{\Sigma} \cdot (hR + h_2\check{R})|\Psi|^2$$

and

$$E[\Psi](\tau) = \int_{\Sigma(\tau)} \left(2b^2 |e_4\Psi|^2 + 2a^2 |e_3\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2 \right).$$

1. In the region $r \geq 2m_0(1 - \delta_{\mathcal{H}})$, $\Upsilon \leq \delta_{\mathcal{H}}^{\frac{1}{10}}$ we have $h = 0$, $h_2 = O(\widehat{\delta})$ and $N_{\Sigma} = e_3$ (i.e. $a = 1$, $b = 0$). Also, we have $w = -\delta_1 w_1$, where $\delta_1 > 0$ is a small constant and w_1 is given by (10.1.46)

$$w_1(r, m) = r^{-1} \frac{m^2}{r^2} \Upsilon \left(1 - \frac{3m}{r} \right)^2.$$

In particular, we have in the region of interest, in view of the formula for w_1 and for N_{Σ}

$$|w_1| \lesssim \delta_{\mathcal{H}}^{\frac{1}{10}}, \quad |N_{\Sigma}(w_1)| = |e_3(w_1)| \lesssim 1.$$

We infer

$$\begin{aligned} \mathcal{P} \cdot N_{\Sigma} &= \mathcal{Q}(X + \Lambda T, e_3) - \frac{\delta_1}{2}w_1\Psi e_3(\Psi) + \frac{\delta_1}{4}e_3(w_1)|\Psi|^2 + \frac{1}{2}h_2e_3 \cdot \check{R}|\Psi|^2 \\ &= \mathcal{Q}(X + \Lambda T, e_3) - O(\delta_{\mathcal{H}}^{\frac{1}{10}})w_1\Psi e_3(\Psi) - O(1)|\Psi|^2. \end{aligned}$$

where we used the fact that $\check{R} = \frac{1}{2}(e_4 - e_3)$ in the region of interest. Thus, according to Proposition 10.1.32,

$$\mathcal{P} \cdot N_{\Sigma} \geq \frac{1}{4}\epsilon_{\mathcal{H}}\mathcal{Q}_{33} + \frac{1}{2}\Lambda\mathcal{Q}_{34} - O(\delta_{\mathcal{H}}^{\frac{1}{10}})|\Psi||e_3(\Psi)| - O(1)|\Psi|^2.$$

Using in particular (10.1.62) and (10.1.64), we deduce

$$\mathcal{P} \cdot N_{\Sigma} \geq \frac{1}{4}\delta_{\mathcal{H}}^{\frac{7}{20}}|e_3\Psi|^2 + \frac{1}{8}\delta_{\mathcal{H}}^{-\frac{13}{20}}(|\nabla\Psi|^2 + O(\epsilon)|\Psi|^2) - O(\delta_{\mathcal{H}}^{\frac{1}{10}})|\Psi||e_3\Psi| - O(1)|\Psi|^2$$

Together with the Poincaré inequality (10.1.26), we deduce

$$\begin{aligned} \int_{\Sigma} \mathcal{P} \cdot N_{\Sigma} &\geq \frac{1}{8}\delta_{\mathcal{H}}^{\frac{7}{20}} \int_{\Sigma} \left(|e_3\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2 \right) \\ &\geq \frac{1}{8}\delta_{\mathcal{H}}^{\frac{7}{20}} E_{r \geq 2m_0(1-\delta_{\mathcal{H}}), \Upsilon \leq \delta_{\mathcal{H}}^{\frac{1}{10}}}[\Psi](\tau). \end{aligned}$$

2. In the region $\Upsilon \geq \delta_{\mathcal{H}}^{\frac{1}{10}}$, we have $w = O(r^{-1})$, $N_{\Sigma}(w) = O(r^{-2})$, $h = O(r^{-4})$ and $h_2 = O(r^{-4})$. We infer

$$\mathcal{P} \cdot N_{\Sigma} = a\mathcal{Q}(X + \Lambda T, e_3) + b\mathcal{Q}(X + \Lambda T, e_4) - O(r^{-1})|\Psi|(a|e_3\Psi| + b|e_4\Psi|) - O(r^{-2})|\Psi|^2.$$

Thus, according to Proposition 10.1.32,

$$\begin{aligned} \mathcal{P} \cdot N_{\Sigma} &\geq \delta_{\mathcal{H}}^{-\frac{1}{2}}(a\mathcal{Q}_{33} + b\mathcal{Q}_{44} + (a+b)\mathcal{Q}_{34}) - O(1)(a^2|e_3\Psi|^2 + b^2|e_4\Psi|^2) - O(r^{-2})|\Psi|^2 \\ &= \delta_{\mathcal{H}}^{-\frac{1}{2}}\left(a|e_3\Psi|^2 + b|e_4\Psi|^3 + (a+b)\left(|\nabla\Psi|^2 + \frac{4\Upsilon}{r^2}|\Psi|^2\right)\right) \\ &\quad - O(1)\left(a^2|e_3\Psi|^2 + b^2|e_4\Psi|^2 + r^{-2}|\Psi|^2\right) \\ &\geq \delta_{\mathcal{H}}^{-\frac{1}{2}}\left(a|e_3\Psi|^2 + b|e_4\Psi|^3 + (a+b)\left(|\nabla\Psi|^2 + \frac{4\delta_{\mathcal{H}}^{\frac{1}{10}}}{r^2}|\Psi|^2\right)\right) \\ &\quad - O(1)\left(a^2|e_3\Psi|^2 + b^2|e_4\Psi|^2 + r^{-2}|\Psi|^2\right). \end{aligned}$$

Hence, for $\delta_{\mathcal{H}} > 0$ sufficiently small, and since $a^2 \leq a$, $b^2 \leq b$ and $a + b \geq 1$, we infer in this region

$$\begin{aligned} \mathcal{P} \cdot N_{\Sigma} &\geq \delta_{\mathcal{H}}^{-\frac{1}{5}} \int_{\Sigma_{r \geq \delta_{\mathcal{H}}^{\frac{1}{10}}(\tau)}} \left(2b^2|e_4\Psi|^2 + 2a^2|e_3\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2\right) \\ &= \delta_{\mathcal{H}}^{-\frac{1}{5}} E[\Psi]_{\Upsilon \geq \delta_{\mathcal{H}}^{\frac{1}{10}}}(\tau) \end{aligned}$$

In view of the above estimates in $r \geq 2m_0(1 - \delta_{\mathcal{H}})$, $\Upsilon \leq \delta_{\mathcal{H}}^{\frac{1}{10}}$ and in $\Upsilon \geq \delta_{\mathcal{H}}^{\frac{1}{10}}$, we deduce, everywhere,

$$\int_{\Sigma(\tau)} \mathcal{P} \cdot N_{\Sigma} \geq \frac{1}{8} \delta_{\mathcal{H}}^{\frac{7}{20}} E[\Psi](\tau). \quad (10.1.69)$$

Boundary terms along Σ_*

On Σ_* , we have

$$N_{\Sigma_*} = T + O\left(\epsilon + \frac{m}{r}\right)(e_3 + e_4),$$

$w = O(r^{-1})$, $N_{\Sigma_*}(w) = O(\epsilon r^{-2})$, $h = O(r^{-4})$ and $h_2 = 0$.

Proceeding as before, we have along Σ_* ,

$$\begin{aligned} \mathcal{P} \cdot N_\Sigma &= \left(\frac{1}{2} + O\left(\epsilon + \frac{m}{r}\right) \right) \mathcal{Q}(X + \Lambda T, e_3) + \left(\frac{1}{2} + O\left(\epsilon + \frac{m}{r}\right) \right) \mathcal{Q}(X + \Lambda T, e_4) \\ &\quad - O(r^{-1})|\Psi|(|e_3\Psi| + |e_4\Psi|) - O(r^{-2})|\Psi|^2 \\ &\geq \frac{1}{4}\mathcal{Q}(X + \Lambda T, e_3) + \frac{1}{4}\mathcal{Q}(X + \Lambda T, e_4) - O\left(|e_3\Psi|^2 + |e_4\Psi|^2 + r^{-2}|\Psi|^2\right). \end{aligned}$$

Thus, according to Proposition 10.1.32, we have

$$\begin{aligned} \mathcal{P} \cdot N_\Sigma &\geq \frac{1}{16}\Lambda(\mathcal{Q}_{33} + \mathcal{Q}_{44} + 2\mathcal{Q}_{34}) - O\left(|e_3\Psi|^2 + |e_4\Psi|^2 + r^{-2}|\Psi|^2\right) \\ &= \frac{1}{16}\Lambda\left(|e_3\Psi|^2 + |e_4\Psi|^2 + 2\left(|\nabla\Psi|^2 + \frac{4\Upsilon}{r^2}|\Psi|^2\right)\right) - O\left(|e_3\Psi|^2 + |e_4\Psi|^2 + r^{-2}|\Psi|^2\right) \\ &\geq \frac{1}{64}\delta_{\mathcal{H}}^{-\frac{13}{20}}\left(|e_3\Psi|^2 + |e_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2\right) - O\left(|e_3\Psi|^2 + |e_4\Psi|^2 + r^{-2}|\Psi|^2\right) \\ &\geq \delta_{\mathcal{H}}^{-\frac{1}{2}}\left(|e_3\Psi|^2 + |e_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2\right) \end{aligned}$$

and hence

$$\int_{\Sigma_*(\tau_1, \tau_2)} \mathcal{P} \cdot N_{\Sigma_*} \geq \delta_{\mathcal{H}}^{-\frac{1}{2}} \int_{\Sigma_*(\tau_1, \tau_2)} \left(|e_3\Psi|^2 + |e_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2 \right). \quad (10.1.70)$$

The inhomogeneous term $\int_{\mathcal{M}(\tau_1, \tau_2)} (\check{X}(\Psi) + \frac{1}{2}w\Psi)N[\Psi]$

Recall that, $\check{X} = X + \Lambda T = f_{\check{\delta}}R + Y_{\mathcal{H}} + \Lambda T$. We easily check, recalling the properties of $f_{\check{\delta}}$, w and Λ and the definition of $J[N, \Psi]$,

$$\begin{aligned} \left| \int_{\mathcal{M}(\tau_1, \tau_2)} \left(\check{X}(\Psi) + \frac{1}{2}w\Psi \right) N[\Psi] \right| &\leq \delta_{\mathcal{H}}^{-\frac{3}{4}} \int_{\mathcal{M}(\tau_1, \tau_2)} \left(|\check{R}\Psi| + |\check{T}\Psi| + r^{-2}|\Psi|^2 \right) |N(\Psi)| \\ &= \delta_{\mathcal{H}}^{-\frac{3}{4}} J[N, \Psi](\tau_1, \tau_2). \end{aligned} \quad (10.1.71)$$

Going back to (10.1.67) we deduce,

$$E[\Psi](\tau_2) + \int_{\mathcal{M}(\tau_1, \tau_2)} \mathcal{E} + F[\Psi](\tau_1, \tau_2) \leq \delta_{\mathcal{H}}^{-\frac{7}{20}} (E[\Psi](\tau_1) + J[N, \Psi](\tau_1, \tau_2)).$$

In view of (10.1.68) we obtain,

$$\begin{aligned} E[\Psi](\tau_2) + \text{Mor}[\Psi](\tau_1, \tau_2) + F[\Psi](\tau_1, \tau_2) &\leq \delta_{\mathcal{H}}^{-1} (E[\Psi](\tau_1) + J[N, \Psi](\tau_1, \tau_2)) \\ &\quad + O\left(\delta_{\mathcal{H}}^{-\frac{3}{2}}\right) \text{Err}_\epsilon(\tau_1, \tau_2). \end{aligned}$$

This concludes the proof of Theorem 10.1.33. \square

10.1.14 Analysis of the error term \mathcal{E}_ϵ

Recall that $\text{Err}_\epsilon(\tau_1, \tau_2) = \int_{\mathcal{M}(\tau_1, \tau_2)} \mathcal{E}_\epsilon$ where,

$$\mathcal{E}_\epsilon \lesssim \epsilon u_{trap}^{-1-\delta_{dec}} \left[r^{-2} |e_3 \Psi|^2 + r^{-1} (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2 + |e_3 \Psi| (|e_4 \Psi| + |\nabla \Psi|)) \right].$$

- In the trapping region \mathcal{M}_{trap} , i.e. $\frac{5m}{2} \leq r \leq \frac{7m}{2}$, where $u_{trap} = 1 + \tau$ and $\Sigma(\tau)$ is strictly spacelike, we have

$$\begin{aligned} \int_{\Sigma_{trap}(\tau)} \mathcal{E}_\epsilon &\lesssim \epsilon \tau_{trap}^{-1-\delta_{dec}} \int_{\Sigma_{trap}(\tau)} (|e_3 \Psi|^2 + |e_4 \Psi|^2 + |\nabla \Psi|^2 + |\Psi|^2) \\ &\lesssim \epsilon \tau_{trap}^{-1-\delta_{dec}} E[\Psi](\tau). \end{aligned}$$

Thus,

$$\begin{aligned} \int_{\mathcal{M}_{trap}(\tau_1, \tau_2)} \mathcal{E}_\epsilon &\lesssim \epsilon \int_{\tau_1}^{\tau_2} \tau_{trap}^{-1-\delta_{dec}} E[\Psi](\tau) \\ &\lesssim \epsilon \left(\int_{\tau_1}^{\tau_2} (1 + \tau)^{-1-\delta} \right) \sup_{\tau \in [\tau_1, \tau_2]} \mathcal{E}[\Psi](\tau) \\ &\lesssim \epsilon \sup_{\tau \in [\tau_1, \tau_2]} \mathcal{E}[\Psi](\tau) \end{aligned}$$

and therefore, for small $\epsilon > 0$, the integral $\int_{\mathcal{M}_{trap}(\tau_1, \tau_2)} \mathcal{E}_\epsilon$ can be absorbed on the left hand side of (10.1.66).

- In the non trapping region \mathcal{M}_{trqp} we write, with a fixed $\delta > 0$,

$$\mathcal{E}_\epsilon \lesssim \epsilon r^{-1-\delta} |e_3 \Psi|^2 + r^{-1+\delta} (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2).$$

Hence,

$$\begin{aligned} \int_{\mathcal{M}_{trqp}(\tau_1, \tau_2)} \mathcal{E}_\epsilon &\lesssim \epsilon \int_{\mathcal{M}_{r \geq 4m_0}(\tau_1, \tau_2)} r^{-1-\delta} |e_3 \Psi|^2 \\ &\quad + \epsilon \int_{\mathcal{M}_{r \geq 4m_0}(\tau_1, \tau_2)} r^{-1+\delta} (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2) \\ &\quad + \epsilon \int_{(trqp)\mathcal{M}_{r \leq 4m_0}(\tau_1, \tau_2)} (|e_3 \Psi|^2 + |e_4 \Psi|^2 + |\nabla \Psi|^2 + |\Psi|^2). \end{aligned}$$

Note that for $\epsilon > 0$ sufficiently small, the last integral, on $(trqp)\mathcal{M}_{r \leq 4m_0}$, can be absorbed by the left hand side of (10.1.66).

As a consequence we deduce the following.

Corollary 10.1.34. *The statement of Theorem 10.1.33 remains true if we replace Err_ϵ in the statement of the theorem with*

$$\begin{aligned} Err_\epsilon &= \int_{\mathcal{M}_{r \geq 4m_0}(\tau_1, \tau_2)} \mathcal{E}_\epsilon, \\ \mathcal{E}_\epsilon &\lesssim \epsilon r^{-1-\delta} |e_3 \Psi|^2 + \epsilon r^{-1+\delta} (|e_4 \Psi|^2 + |\nabla \Psi|^2 + |r^{-2} \Psi|^2), \end{aligned}$$

for a fixed $\delta > 0$.

Remark 10.1.35. *Note that the error terms Err_ϵ cannot yet be absorbed to the left hand side of (10.1.66). In fact we need additional estimates. The Morawetz bulk quantity (5.1.13),*

$$Mor[\Psi](\tau_1, \tau_2) := \int_{\mathcal{M}(\tau_1, \tau_2)} \frac{m^2}{r^3} |\check{R}\Psi|^2 + \frac{m}{r^4} |\Psi|^2 + \left(1 - \frac{3m}{r}\right)^2 \frac{1}{r} \left(|\nabla \Psi|^2 + \frac{m^2}{r^2} |\check{T}\Psi|^2 \right)$$

is quite weak for r large with regard to the terms $|\check{R}\Psi|^2$ and $|\check{T}\Psi|^2$, while, using the Poincaré inequality, $Mor[\Psi]$ controls the term $\int_{\mathcal{M}_{r \geq 4m_0}(\tau_1, \tau_2)} r^{-1} (|\nabla \Psi|^2 + r^{-2} |\Psi|^2)$. In the next section we show how we can estimate $\int_{\mathcal{M}_{\geq R_0}(\tau_1, \tau_2)} r^{-1-\delta} |e_3 \Psi|^2$ by $\int_{\mathcal{M}_{\geq R_0}(\tau_1, \tau_2)} r^{-1-\delta} |e_4 \Psi|^2$ and then, we provide estimates for the remaining terms. Note also that the weight $r^{-1-\delta}$ is optimal in estimating $e_3 \Psi$ in the wave zone region.

10.1.15 Proof of Theorem 10.1.1

We are now ready to prove Theorem 10.1.1. Note that it suffices to improve the previous Morawetz estimate of Theorem 10.1.33 by replacing the quantity $Mor[\Psi](\tau_1, \tau_2)$ with

$$Morr[\Psi](\tau_1, \tau_2) := Mor[\Psi](\tau_1, \tau_2) + \int_{\mathcal{M}_{far}(\tau_1, \tau_2)} r^{-1-\delta} |e_3(\Psi)|^2.$$

In view of the Morawetz estimate (10.1.66) and corollary 10.1.34 we have

$$\begin{aligned} E[\Psi](\tau_2) + Mor[\Psi](\tau_1, \tau_2) + F[\Psi](\tau_1, \tau_2) &\lesssim E[\Psi](\tau_2) + J[N, \Psi](\tau_1, \tau_2) + Err_\epsilon(\tau_1, \tau_2), \\ J[N, \Psi](\tau_1, \tau_2) &:= \int_{\mathcal{M}(\tau_1, \tau_2)} (|\check{R}\Psi| + |\check{T}\Psi| + r^{-1} |\Psi|) |N|, \end{aligned} \quad (10.1.72)$$

with error term,

$$\text{Err}_\epsilon \lesssim \epsilon \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{-1-\delta} |e_3 \Psi|^2 + r^{-1+\delta} (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2).$$

We divide $J[N] = J[N, \Psi]$ as follows:

$$J[N] = J[N]_{\text{trap}} + J[N]_{\text{tr}\not{d}p}$$

where,

$$\begin{aligned} J[N]_{\text{trap}} &:= \int_{\mathcal{M}_{\text{trap}}} (|\check{R}\Psi| + |\check{T}\Psi| + r^{-1}|\Psi|)|N|, \\ J[N]_{\text{tr}\not{d}p} &:= \int_{(\text{tr}\not{d}p)_{\mathcal{M}}} (|\check{R}\Psi| + |\check{T}\Psi| + r^{-1}|\Psi|)|N|. \end{aligned}$$

For the trapping region, where the hypersurfaces $\Sigma(\tau)$ are strictly spacelike, we write,

$$\begin{aligned} J[N]_{\text{trap}}(\tau_1, \tau_2) &= \int_{\tau_1}^{\tau_2} d\tau \int_{\Sigma_{\text{trap}}(\tau)} (|\check{R}\Psi| + |\check{T}\Psi| + r^{-1}|\Psi|)|N| \\ &\leq \int_{\tau_1}^{\tau_2} E[\Psi](\tau)^{1/2} \left(\int_{\Sigma_{\text{trap}}(\tau)} |N|^2 \right)^{1/2} \\ &\leq \sup_{\tau \in [\tau_1, \tau_2]} E[\Psi](\tau)^{1/2} \int_{\tau_1}^{\tau_2} \left(\int_{\Sigma_{\text{trap}}(\tau)} |N|^2 \right)^{1/2} \\ &\lesssim \lambda \sup_{\tau \in [\tau_1, \tau_2]} E[\Psi](\tau) + \lambda^{-1} \left(\int_{\tau_1}^{\tau_2} \|N\|_{L^2(\Sigma_{\text{trap}}(\tau))} \right)^2. \end{aligned}$$

Hence, for $\lambda > 0$ sufficiently small, we deduce,

$$\begin{aligned} E[\Psi](\tau_2) + \text{Mor}[\Psi](\tau_1, \tau_2) + F[\Psi](\tau_1, \tau_2) &\lesssim E[\Psi](\tau_2) + \text{Err}_\epsilon(\tau_1, \tau_2) \\ &\quad + J_{\text{tr}\not{d}p}[N, \Psi](\tau_1, \tau_2) + \left(\int_{\tau_1}^{\tau_2} \|N\|_{L^2(\Sigma_{\text{trap}}(\tau))} \right)^2. \end{aligned}$$

On the other hand we have,

$$\begin{aligned} J[N]_{\text{tr}\not{d}p}(\tau_1, \tau_2) &= \int_{\mathcal{M}_{\text{tr}\not{d}p}(\tau_1, \tau_2)} (|\check{R}\Psi| + |\check{T}\Psi| + r^{-1}|\Psi|)|N| \\ &\leq \lambda \int_{\mathcal{M}_{\text{tr}\not{d}p}} r^{-1-\delta} (|\check{R}\Psi|^2 + |\check{T}\Psi|^2 + r^{-2}|\Psi|^2) + \lambda^{-1} \int_{\mathcal{M}_{\text{tr}\not{d}p}} r^{1+\delta} |N|^2. \end{aligned}$$

The first integral on the right can be divided further into integrals for $r \leq 4m_0$ and $r \geq 4m_0$. The first integral can be easily absorbed by the term $Mor[\Psi](\tau_1, \tau_2)$, if $\lambda > 0$ is sufficiently small. We are thus led to the estimate,

$$\begin{aligned} E[\Psi](\tau_2) + Mor[\Psi](\tau_1, \tau_2) + F[\Psi](\tau_1, \tau_2) &\lesssim E[\Psi](\tau_2) + Err_\epsilon(\tau_1, \tau_2) + \mathcal{I}_\delta[N](\tau_1, \tau_2) \\ &\quad + \int_{\mathcal{M}_{r \geq 4m_0}} r^{-1-\delta} (|e_3\Psi|^2 + |e_4\Psi|^2 + r^{-2}|\Psi|^2) \end{aligned}$$

where,

$$\mathcal{I}_\delta[N](\tau_1, \tau_2) := \int_{\mathcal{M}_{trap}(\tau_1, \tau_2)} r^{1+\delta} |N|^2 + \left(\int_{\tau_1}^{\tau_2} d\tau \|N\|_{L^2(\Sigma_{trap}(\tau))} \right)^2.$$

Recalling the definition of Err_ϵ in Corollary 10.1.34, we deduce,

$$E[\Psi](\tau_2) + Mor[\Psi](\tau_1, \tau_2) + F[\Psi](\tau_1, \tau_2) \lesssim E[\Psi](\tau_2) + Err_\epsilon(\tau_1, \tau_2) + \mathcal{I}_\delta[N](\tau_1, \tau_2). \quad (10.1.73)$$

To eliminate the term in $e_3\Psi$ from the error term we appeal to the following proposition.

Proposition 10.1.36. *Assume $\square\Psi = V\Psi + N$ and consider the vectorfield $X = f_{-\delta}T$ with $f_{-\delta} := r^{-\delta}$ for $r \geq 4m_0$ and compactly supported in $r \geq \frac{7m_0}{2}$. With the notation of Proposition 10.1.9, let*

$$\begin{aligned} \mathcal{P}_\mu[f_{-\delta}T, 0, 0] &= f_{-\delta} \mathcal{Q}_{\alpha\mu} T^\mu, \\ \mathcal{E}[f_{-\delta}T, 0, 0] &= \mathbf{D}^\mu \mathcal{P}_\mu[f_{-\delta}T, 0, 0] - f_{-\delta} T(\Psi)N. \end{aligned}$$

Then,

1. We have, for $r \geq 4m_0$

$$\mathcal{E}[f_{-\delta}T, 0, 0] = \frac{\Upsilon^2}{4} \delta r^{-1-\delta} |e_3\Psi|^2 - \frac{1}{4} \delta r^{-1-\delta} |e_4\Psi|^2 + O(\epsilon r^{-1-\delta} (|\mathbf{D}\Psi|^2 + r^{-2}|\Psi|^2)).$$

2. We have,

$$\mathcal{P}[f_{-\delta}T, 0, 0] \cdot e_4 = f_{-\delta} \mathcal{Q}(T, e_4) \geq 0, \quad \mathcal{P}[f_{-\delta}T, 0, 0] \cdot e_3 = f_{-\delta} \mathcal{Q}(T, e_3) \geq 0.$$

We postponed the proof of Proof of Proposition 10.1.36 and continue the proof of Theorem 10.1.1. By integration, the proposition provides a bound for¹¹

$$\int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{-1-\delta} |e_3\Psi|^2$$

¹¹Note that $\Upsilon^2 \geq \frac{1}{4}$ in $r \geq 4m_0$.

in terms of $E[\Psi](\tau_1)$, the integrals $\int_{\mathcal{M}_{\geq \frac{7m_0}{2}}(\tau_1, \tau_2)} r^{-1-\delta} |e_4 \Psi|^2$ and $\int_{\mathcal{M}_{\geq \frac{7m_0}{2}}(\tau_1, \tau_2)} r^{-\delta} T(\Psi)N$, as well as the error terms. The second bulk integral involving the inhomogeneous term N can be estimated exactly like before. Thus combining the new estimate with that in the corollary 10.1.34 we derive the desired estimate hence concluding the proof of Theorem 10.1.1.

Proof of Proposition 10.1.36

We consider vectorfields of the form $X = f(r)T$ with $T = \frac{1}{2}(\Upsilon e_3 + e_4)$. Recall, see Lemma 10.1.7, that all components of the deformation tensor ${}^{(X)}\pi$ of $T = \frac{1}{2}(e_4 + \Upsilon e_3)$ can be bounded by $O(\epsilon r^{-1})$. Since $f = O(r^{-\delta})$, we deduce,

$${}^{(X)}\pi_{\alpha\beta} = f {}^{(T)}\pi_{\alpha\beta} + \mathbf{D}_\alpha f T_\beta + \mathbf{D}_\beta f T_\alpha = \mathbf{D}_\alpha f T_\beta + \mathbf{D}_\beta f T_\alpha + O(\epsilon r^{-1-\delta}).$$

Also,

$$e_3(f) = f' e_3(r) = -f' + O(\epsilon r^{-1-\delta}), \quad e_4(f) = f' e_4(r) = \Upsilon f' + O(\epsilon r^{-1-\delta}).$$

Thus, modulo error terms of the form $O(\epsilon) r^{-1-\delta} (|e_3 \Psi|^2 + |e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2)$, we have

$$\begin{aligned} \mathcal{Q} \cdot {}^{(X)}\pi &= 2\mathcal{Q}^{\alpha\beta} T_\alpha \mathbf{D}_\beta f = 2(\mathcal{Q}^{3\beta} T_\beta e_3 f + \mathcal{Q}^{4\beta} T_\beta e_4 f) = -\mathcal{Q}(e_4, T) e_3 f - \mathcal{Q}(e_3, T) e_4 f \\ &= \frac{1}{2} \mathcal{Q}(e_4, e_4 + \Upsilon e_3) f' - \frac{1}{2} f' \Upsilon \mathcal{Q}(e_3, e_4 + \Upsilon e_3) \\ &= \frac{1}{2} f' (|e_4 \Psi|^2 - \Upsilon^2 |e_3 \Psi|^2). \end{aligned}$$

We now apply Proposition 10.1.9, as well as (10.1.12) (10.1.13), with $X = f_{-\delta}(r)T$, $w = 0$, $M = 0$ so that

$$\mathcal{P}_\mu[f_{-\delta}T, 0, 0] = f_{-\delta} \mathcal{Q}_{\alpha\mu} T^\mu, \quad \mathcal{E}[f_{-\delta}T, 0, 0] := \mathbf{D}^\mu \mathcal{P}_\mu[f_{-\delta}T, 0, 0] - f_{-\delta} T(\Psi)N$$

and

$$\begin{aligned} \mathcal{E}[f_{-\delta}T, 0, 0] &= \frac{1}{2} \mathcal{Q} \cdot {}^{(X)}\pi - \frac{1}{2} f_{-\delta} T(V) |\Psi|^2 \\ &= \frac{1}{4} f'_{-\delta}(r) (|e_4 \Psi|^2 - \Upsilon^2 |e_3 \Psi|^2) + O(\epsilon r^{-1-\delta} (|\mathbf{D}\Psi|^2 + r^{-2} |\Psi|^2)) \end{aligned}$$

with $|\mathbf{D}\Psi|^2 = |e_3 \Psi|^2 + |e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2$. Since $f_{-\delta}(r) = r^{-\delta}$ for $r \geq 4m_0$, we deduce, for $r \geq 4m_0$,

$$\mathcal{E}[f_{-\delta}T, 0, 0] = \frac{\Upsilon^2}{4} \delta r^{-1-\delta} |e_3 \Psi|^2 - \frac{1}{4} \delta r^{-1-\delta} |e_4 \Psi|^2 + O(\epsilon r^{-1-\delta} (|\mathbf{D}\Psi|^2 + r^{-2} |\Psi|^2)).$$

On the other hand,

$$\begin{aligned}\mathcal{P}[f_{-\delta}T, 0, 0] \cdot e_4 &= f_{-\delta}\mathcal{Q}(T, e_4) \geq 0, \\ \mathcal{P}[f_{-\delta}T, 0, 0] \cdot e_3 &= f_{-\delta}\mathcal{Q}(T, e_3) \geq 0,\end{aligned}$$

as desired. This concludes the proof of Proposition 10.1.36.

10.2 Dafermos-Rodnianski r^p - weighted estimates

For convenience, we work in this section with the renormalized frame (e'_3, e'_4) defined in (10.2.7) instead of the original frame (e_3, e_4) . To simplify the exposition, we still denote it as (e_3, e_4) . Recall that the two are frames are equivalent up to lower terms in m/r .

In this section we rely on the Morawetz estimates proved in the previous section to establish r^p -weighted estimates in the spirit of Dafermos-Rodnianski [20]. The following theorem claims r^p -weighted estimates for the solution ψ of the wave equation (5.3.5).

Theorem 10.2.1 (r^p -weighted estimates). *Consider a fixed $\delta > 0$ and let $R \gg \frac{m_0}{\delta}$, $\epsilon \ll \delta$. The following estimates hold true and for all $\delta \leq p \leq 2 - \delta$,*

$$\dot{E}_{p;R}[\psi](\tau_2) + \dot{B}_{p;R}[\psi](\tau_1, \tau_2) + \dot{F}_p[\psi](\tau_1, \tau_2) \lesssim E_p[\psi](\tau_1) + J_p[\psi, N](\tau_1, \tau_2). \quad (10.2.1)$$

Remark 10.2.2. *Note that Theorem 10.1.1 on Morawetz estimates and Theorem 10.2.1 on r^p -weighted estimates immediately yield for all $\delta \leq p \leq 2 - \delta$,*

$$\sup_{\tau \in [\tau_1, \tau_2]} E_p[\psi](\tau) + B_p[\psi](\tau_1, \tau_2) + F_p[\psi](\tau_1, \tau_2) \lesssim E_p[\psi](\tau_1) + J_p[\psi, N](\tau_1, \tau_2), \quad (10.2.2)$$

which corresponds to Theorem 5.3.4 in the case $s = 0$.

Theorem 10.2.1 will be proved in section 5.3.1. We will need in this section stronger assumptions in the region $r \geq 4m_0$, away from trapping, than those in (10.1.4)–(10.1.6) of the previous section. For convenience we express our conditions with respect to the weights¹²,

$$w_{p,q}(u, r) = r^{-p}(1 + \tau)^{-q - \delta_{dec} + 2\delta_0}.$$

¹²The assumptions are consistent with the global frame used in Theorem M1, see Lemma 5.1.1. In particular, $\delta_0 > 0$ is such that $\delta_{dec} - 2\delta_0 > 0$ which is the only needed property of $\delta_{dec} - 2\delta_0$ to derive the r^p weighted estimates.

RP0. The assumptions **Mor1–Mor4** made in the previous section hold true.

RP1. The Ricci coefficients verify, for $r \geq 4m_0$

$$\begin{aligned} |\underline{\xi}, \vartheta, \underline{\vartheta}, \eta, \underline{\eta}, \zeta, \underline{\omega}| &\lesssim \epsilon w_{1,1}, \\ \left| \underline{\kappa} + \frac{2}{r} \right|, \left| \underline{\chi} + \frac{1}{r} \right|, \left| e_3 \Phi - \underline{\chi} \right| &\lesssim \epsilon w_{1,1}, \\ \left| \kappa - \frac{2\Upsilon}{r} \right|, \left| \chi - \frac{\Upsilon}{r} \right|, \left| e_4 \Phi - \chi \right| &\lesssim \epsilon \min\{w_{1,1}, w_{2,1/2}\}, \\ \left| \omega + \frac{m}{r^2} \right|, |\xi| &\lesssim \epsilon \min\{w_{2,1}, w_{3,1/2}\}. \end{aligned} \tag{10.2.3}$$

RP2. The derivatives of r verify, for $r \geq 4m_0$,

$$\begin{aligned} |e_3(r) + 1| &\lesssim \epsilon w_{0,1}, \\ |e_4(r) - \Upsilon| &\lesssim \epsilon \min\{w_{0,1}, w_{1,1/2}\}, \\ \left| e_3 e_4(r) + \frac{2m}{r^2}, e_4 e_3(r) \right| &\lesssim \epsilon w_{1,1}. \end{aligned} \tag{10.2.4}$$

RP3. For $r \geq 4m_0$,

$$\begin{aligned} \left| \rho + \frac{2m}{r^3} \right| &\lesssim \epsilon w_{3,1}, \\ \left| K - \frac{1}{r^2} \right| &\lesssim \epsilon r^{-2}, \\ |e_\theta(\Phi)| &\lesssim r^{-1}. \end{aligned} \tag{10.2.5}$$

RP4. We also assume, for $r \geq 4m_0$,

$$\begin{aligned} |m - m_0| &\lesssim \epsilon, \\ |e_3 m, r e_4 m| &\lesssim \epsilon w_{0,1}, \\ |e_3 e_4(m), e_4 e_3(m)| &\lesssim \epsilon w_{1,1}. \end{aligned} \tag{10.2.6}$$

Since the estimates we are establishing are restricted to the far region $r > R$ it is convenient, in this section, to work with the with renormalized frame

$$e'_3 = \Upsilon e_3, \quad e'_4 = \Upsilon^{-1} e_4, \quad e'_\theta = e_\theta. \tag{10.2.7}$$

Relative to the new frame (e'_3, e'_4, e'_θ) we have,

$$\xi' = \Upsilon^{-2} \xi, \quad \underline{\xi}' = \Upsilon^2 \underline{\xi}, \quad \zeta' = \zeta, \quad \eta' = \eta, \quad \chi' = \Upsilon^{-1} \chi, \quad \underline{\chi}' = \Upsilon \underline{\chi}$$

and,

$$\begin{aligned}
\omega' &= \Upsilon^{-1} \left(\omega + \frac{1}{2} e_4(\log \Upsilon) \right) = \Upsilon^{-1} \left(\omega + \frac{1}{2} \Upsilon^{-1} \Upsilon' e_4(r) \right) \\
&= \Upsilon^{-1} \left(-\frac{m}{r^2} + O(\epsilon r^{-2} (1 + |u|)^{-1/2 - \delta_{dec}}) + \Upsilon^{-1} \frac{m}{r^2} (\Upsilon + O(\epsilon r^{-1} (1 + |u|)^{-1/2 - \delta_{dec}})) \right) \\
&= O(\epsilon r^{-2}), \\
\underline{\omega}' &= \Upsilon \left(\underline{\omega} - \frac{1}{2} e_3(\log \Upsilon) \right) = \Upsilon \left(\underline{\omega} - \frac{1}{2} \Upsilon^{-1} \Upsilon' e_3(r) \right) \\
&= \Upsilon \left(\underline{\omega} - \frac{1}{2} \Upsilon^{-1} \Upsilon' (-1 + O(\epsilon (1 + |u|)^{-1 - \delta_{dec}})) \right) \\
&= \frac{m}{r^2} + O(\epsilon r^{-1} (1 + |u|)^{-1 - \delta_{dec}}).
\end{aligned}$$

Thus in the new frame we have, for $r \geq 4m_0$,

RP1'. The Ricci coefficients with respect to the null frame (e'_3, e'_4, e'_θ) verify, for $r \geq 4m_0$:

$$\begin{aligned}
|\underline{\xi}', \vartheta', \underline{\vartheta}, \eta', \underline{\eta}', \zeta'|, |\underline{\omega}' - \frac{m}{r^2}| &\lesssim \epsilon w_{1,1}, \\
\left| \underline{\kappa}' + \frac{2\Upsilon}{r} \right|, \left| \underline{\chi}' + \frac{\Upsilon}{r} \right|, |e'_3 \Phi - \underline{\chi}'| &\lesssim \epsilon w_{1,1}, \\
\left| \kappa' - \frac{2}{r} \right|, \left| \chi' - \frac{1}{r} \right|, |e'_4 \Phi - \chi'| &\lesssim \epsilon \min\{w_{1,1}, w_{2,1/2}\}, \\
|\omega'|, |\xi'| &\lesssim \epsilon \min\{w_{2,1}, w_{3,1/2}\}.
\end{aligned} \tag{10.2.8}$$

RP2'. The derivatives of r verify,

$$\begin{aligned}
|e'_3(r) + \Upsilon| &\lesssim \epsilon w_{0,1}, \\
|e'_4(r) - 1| &\lesssim \epsilon w_{1,1}, \\
\left| e'_3 e'_4(r), e'_4 e'_3(r) + \frac{2m}{r^2} \right| &\lesssim \epsilon w_{1,1}.
\end{aligned} \tag{10.2.9}$$

RP3'. The Gauss curvature K of S and ρ verify,

$$\begin{aligned}
\left| \rho + \frac{2m}{r^3} \right| &\lesssim \epsilon r^{-3}, \\
\left| K - \frac{1}{r^2} \right| &\lesssim \epsilon r^{-2}.
\end{aligned} \tag{10.2.10}$$

RP4'. We also assume

$$\begin{aligned}
|m - m_0| &\lesssim \epsilon, \\
|e'_3 m, r e'_4 m| &\lesssim \epsilon w_{0,1}, \\
|e'_3 e'_4(m), e'_4 e'_3(m)| &\lesssim \epsilon w_{1,1}.
\end{aligned} \tag{10.2.11}$$

Remark 10.2.3. *In the far region $r \geq 4m_0$ all norms we are using in our estimates are equivalent when expressed relative to the null frame (e_3, e_4, e_θ) or (e'_3, e'_4, e'_θ) .*

Convention. For the remaining of this section we shall do all calculations with respect to the renormalized frame (e'_3, e'_4, e'_θ) . For convenience we shall drop the primes, throughout this section, since there is no danger of confusion. Note however that the main results, which include the interior region $r \leq R$, are always expressed with respect the original frame.

10.2.1 Vectorfield $X = f(r)e_4$

Lemma 10.2.4. *Consider the vectorfield $X = f(r)e_4$.*

1. *We have the decomposition,*

$${}^{(X)}\pi = {}^{(X)}\Lambda \mathbf{g} + {}^{(X)}\tilde{\pi}, \quad {}^{(X)}\Lambda = \frac{2}{r}f$$

with symmetric tensor ${}^{(X)}\tilde{\pi}$ which verifies

$$\begin{aligned} {}^{(X)}\tilde{\pi}_{43} &= -2f' + \frac{4f}{r} + O(\epsilon)w_{1,1}(|f| + r|f'|) \\ {}^{(X)}\tilde{\pi}_{33} &= 4f'\Upsilon - 4\Upsilon' + O(\epsilon)w_{1,1}(|f| + r|f'|) \\ {}^{(X)}\tilde{\pi}_{4\theta} &= O(\epsilon)w_{2,1/2}|f| \\ {}^{(X)}\tilde{\pi}_{AB} &= O(\epsilon)w_{2,1/2}|f| \\ {}^{(X)}\tilde{\pi}_{3\theta} &= O(\epsilon)w_{1,1}|f| \end{aligned} \tag{10.2.12}$$

2. *We have,*

$$\square {}^{(X)}\Lambda = \frac{2}{r}f'' + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right)(|f| + r|f'| + r^2|f''|) \tag{10.2.13}$$

Proof. See Lemma D.3.1 in appendix. □

10.2.2 Energy densities for $X = f(r)e_4$

We start with the following proposition.

Proposition 10.2.5. *Assume Ψ verifies the equation $\square_{\mathbf{g}}\Psi = V\Psi + N$ and let $X = fe_4$ and $w = {}^{(X)}\Lambda = \frac{2f}{r}$ and let $\mathcal{E} := \mathcal{E}[X, w] = \mathcal{E}[X = fe_4, w = \frac{2f}{r}]$.*

1. *We have,*

$$\mathcal{E} = \frac{1}{2}f'|e_4\Psi|^2 + \frac{1}{2}\left(-f' + \frac{2f}{r}\right)(|\nabla\Psi|^2 + V|\Psi|^2) - \frac{1}{2r}f''|\Psi|^2 + \text{Err}\left(\epsilon, \frac{m}{r}, f\right)(\Psi)$$

where,

$$\begin{aligned} \text{Err}\left(\epsilon, \frac{m}{r}, f\right)(\Psi) &= O\left(\frac{m}{r^2}\right)(|f| + r|f'|)|e_4\Psi|^2 + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right)(|f| + r|f'| + r^2|f''|)|\Psi|^2 \\ &+ O(\epsilon)w_{1,1}(|f| + r|f'|)(|e_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2) \\ &+ O(\epsilon)w_{2,1/2}|f|\left(|e_3\Psi|(|e_4\Psi| + r^{-1}|\nabla\Psi|) + |\nabla\Psi|^2 + r^{-2}|\Psi|^2\right). \end{aligned}$$

2. *The current,*

$$\mathcal{P}_\mu = \mathcal{P}_\mu[X, w] = \mathcal{Q}_{\mu\nu}X^\nu + \frac{1}{2}w\Psi \cdot \mathbf{D}_\mu\Psi - \frac{1}{4}|\Psi|^2\partial_\mu w$$

verifies,

$$\begin{aligned} \mathcal{P} \cdot e_4 &= f\left|e_4\Psi + \frac{1}{r}\Psi\right|^2 - \frac{1}{2}r^{-2}e_4(rf|\Psi|^2) + O(\epsilon r^{-3})f|\Psi|^2, \\ \mathcal{P} \cdot e_3 &= f\mathcal{Q}_{34} + \frac{1}{2}r^{-2}e_3(rf\psi^2) + r^{-1}f'\psi^2 + O(mr^{-3} + \epsilon r^{-2})|rf'|\Psi|^2. \end{aligned}$$

3. *Let $\theta = \theta(r)$ supported for $r \geq R/2$ with $\theta = 1$ for $r \geq R$ such that $f_p = \theta(r)r^p$. Let ${}^{(p)}\mathcal{P} := \mathcal{P}[f_p e_4, w_p]$. Then, for all $r \geq R$,*

$${}^{(p)}\mathcal{P} \cdot e_4 + \frac{p}{2}r^{-2}e_4(\theta r^{p+1}|\Psi|^2) \geq \frac{1}{8}r^{p-2}(p-1)^2|\Psi|^2.$$

Before proceeding with the proof of Proposition 10.2.5, we first establish the following lemma.

Lemma 10.2.6. *We have,*

$$\begin{aligned} \mathcal{Q} \cdot {}^{(X)}\tilde{\pi} &= \left(f' + O\left(\frac{m}{r^2}\right)(|f| + r|f'|)\right)|e_4\Psi|^2 + \left(-f' + \frac{2f}{r}\right)(|\nabla\Psi|^2 + V|\Psi|^2) \\ &+ O(\epsilon)w_{1,1}(|f| + r|f'|)(|e_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2) \\ &+ O(\epsilon)w_{2,1/2}|f|\left(|e_3\Psi|(|e_4\Psi| + r^{-1}|\nabla\Psi|) + |\nabla\Psi|^2 + r^{-2}|\Psi|^2\right). \end{aligned}$$

Proof. Recall from Proposition 10.1.9 that we have

$$\mathcal{Q}_{33} = |e_3\Psi|^2, \quad \mathcal{Q}_{44} = |e_4\Psi|^2, \quad \mathcal{Q}_{34} = |\nabla\Psi|^2 + V|\Psi|^2,$$

and,

$$|\mathcal{Q}_{AB}| \leq |e_3\Psi||e_4\Psi| + |\nabla\Psi|^2 + |V||\Psi|^2, \quad |\mathcal{Q}_{A3}| \leq |e_3\Psi||\nabla\Psi|, \quad |\mathcal{Q}_{A4}| \leq |e_4\Psi||\nabla\Psi|.$$

Hence, in view of Lemma D.3.1 for ${}^{(X)}\tilde{\pi}$, we have

$$\begin{aligned} \mathcal{Q} \cdot {}^{(X)}\tilde{\pi} &= \frac{1}{4}\mathcal{Q}_{44} {}^{(X)}\tilde{\pi}_{33} + \frac{1}{2}\mathcal{Q}_{34} {}^{(X)}\tilde{\pi}_{34} - \frac{1}{2}\mathcal{Q}_{4A} {}^{(X)}\tilde{\pi}_{3A} - \frac{1}{2}\mathcal{Q}_{3A} {}^{(X)}\tilde{\pi}_{4A} + \mathcal{Q}_{AB} {}^{(X)}\tilde{\pi}_{AB} \\ &= \left(f'\Upsilon - \Upsilon'f + O(\epsilon)w_{1,1}(|f| + r|f'|)\right)\mathcal{Q}_{44} \\ &+ \left(-f' + \frac{2f}{r} + O(\epsilon)w_{1,1}(|f| + r|f'|)\right)\mathcal{Q}_{34} \\ &+ O(\epsilon)w_{1,1}|f|\mathcal{Q}_{4A} + O(\epsilon)w_{2,1/2}|f|(\mathcal{Q}_{3A} + \mathcal{Q}_{AB}) \\ &= \left(f' + O\left(\frac{m}{r^2}\right)(|f| + r|f'|)\right)\mathcal{Q}_{44} + \left(-f' + \frac{2f}{r}\right)\mathcal{Q}_{34} \\ &+ O(\epsilon)(|f| + r|f'|)w_{1,1}(\mathcal{Q}_{44} + \mathcal{Q}_{4A}) + O(\epsilon)w_{2,1/2}|f|(\mathcal{Q}_{AB} + \mathcal{Q}_{34}) \\ &+ O(\epsilon)w_{3,1/2}|f|\mathcal{Q}_{3A} \end{aligned}$$

from which we deduce,

$$\begin{aligned} \mathcal{Q} \cdot {}^{(X)}\tilde{\pi} &= \left(f' + O\left(\frac{m}{r^2}\right)(|f| + r|f'|)\right)|e_4\Psi|^2 + \left(-f' + \frac{2f}{r}\right)(|\nabla\Psi|^2 + V|\Psi|^2) \\ &+ O(\epsilon)w_{1,1}(|f| + r|f'|)\left(|e_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2\right) \\ &+ O(\epsilon)w_{2,1/2}|f|\left(|e_3\Psi|(|e_4\Psi| + r^{-1}|\nabla\Psi|) + |\nabla\Psi|^2 + r^{-2}|\Psi|^2\right) \end{aligned}$$

as desired. □

We are now ready to prove Proposition 10.2.5.

Proof of Proposition 10.2.5. If $\mathcal{Q} = \mathcal{Q}[\Psi]$ is the energy momentum tensor of Ψ (recall $\dot{\square}\Psi = V\Psi + N$) and

$${}^{(X)}\pi = {}^{(X)}\Lambda \mathbf{g} + {}^{(X)}\tilde{\pi}$$

we deduce,

$$\mathcal{Q} \cdot {}^{(X)}\pi = {}^{(X)}\Lambda \text{tr}\mathcal{Q} + \mathcal{Q} \cdot {}^{(X)}\tilde{\pi} = {}^{(X)}\Lambda(-\mathcal{L}(\Psi) - V|\Psi|^2) + \mathcal{Q} \cdot {}^{(X)}\tilde{\pi}.$$

Hence, for $X = fe_4$ and $w = {}^{(X)}\Lambda = \frac{2f}{r}$,

$$\frac{1}{2}\mathcal{Q} \cdot {}^{(X)}\pi + \frac{1}{2}w\mathcal{L}[\Psi] = -\frac{1}{2}{}^{(X)}\Lambda V|\Psi|^2 + \frac{1}{2}\mathcal{Q} \cdot {}^{(X)}\tilde{\pi}.$$

In view of (10.1.13), we infer

$$\begin{aligned} \mathcal{E} &:= \mathcal{E}[X, w = {}^{(X)}\Lambda, M = 0] \\ &= \frac{1}{2}\mathcal{Q} \cdot {}^{(X)}\tilde{\pi} - \frac{1}{4}|\Psi|^2\Box_{\mathbf{g}}{}^{(X)}\Lambda - \frac{1}{2}(X(V) + {}^{(X)}\Lambda V)|\Psi|^2. \end{aligned}$$

Recall that $V = -\kappa\underline{\kappa}$. Hence,

$$\begin{aligned} X(V) + {}^{(X)}\Lambda V &= fe_4(V) + \frac{2f}{r}V = -f \left(e_4(\kappa\underline{\kappa}) + \frac{2}{r}\kappa\underline{\kappa} \right) = f \left(\kappa^2\underline{\kappa} - 2\kappa\rho - \frac{2}{r}\kappa\underline{\kappa} + O(\epsilon)w_{3,1} \right) \\ &= O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) f. \end{aligned}$$

Hence, in view of the computation (10.2.13) of $\Box{}^{(X)}\Lambda$

$$\begin{aligned} &-\frac{1}{4}|\Psi|^2\Box_{\mathbf{g}}{}^{(X)}\Lambda - \frac{1}{2}(X(V) + {}^{(X)}\Lambda V)|\Psi|^2 \\ &= -\frac{1}{2r}f''|\Psi|^2 + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) (|f| + r|f'| + r^2|f''|) |\Psi|^2. \end{aligned}$$

We deduce,

$$\mathcal{E} = \frac{1}{2}\mathcal{Q} \cdot {}^{(X)}\tilde{\pi} - \frac{1}{2r}f''|\Psi|^2 + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) (|f| + r|f'| + r^2|f''|) |\Psi|^2$$

Using Lemma 10.2.6, we deduce,

$$\mathcal{E} = \frac{1}{2}f'|e_4\Psi|^2 + \frac{1}{2}\left(-f' + \frac{2f}{r}\right) (|\nabla\Psi|^2 + V|\Psi|^2) - \frac{1}{2r}f''|\Psi|^2 + \text{Err}\left(\epsilon, \frac{m}{r}, f\right)(\Psi)$$

where,

$$\begin{aligned} \text{Err}\left(\epsilon, \frac{m}{r}, f\right)(\Psi) &= O\left(\frac{m}{r^2}\right) (|f| + r|f'|)|e_4\Psi|^2 + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) (|f| + r|f'| + r^2|f''|) |\Psi|^2 \\ &\quad + O(\epsilon)w_{1,1}(|f| + r|f'|)\left(|e_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2\right) \\ &\quad + O(\epsilon)w_{2,1/2}|f|\left(|e_3\Psi|(|e_4\Psi| + r^{-1}|\nabla\Psi|) + |\nabla\Psi|^2 + r^{-2}|\Psi|^2\right) \end{aligned}$$

which is the first part of Proposition 10.2.5.

To prove the second part of Proposition 10.2.5, we compute

$$\begin{aligned}
\mathcal{P} \cdot e_4 &= f\mathcal{Q}_{44} + \frac{1}{r}f\Psi \cdot e_4\Psi - \frac{1}{2}e_4(r^{-1}f)|\Psi|^2 \\
&= f\left(|e_4\Psi|^2 + \frac{1}{r}\Psi \cdot e_4\Psi\right) - \frac{1}{2}e_4(r^{-1}f)|\Psi|^2 \\
&= f\left|e_4\Psi + \frac{1}{r}\Psi\right|^2 - \frac{1}{r}f\Psi \cdot e_4\Psi - r^{-2}f|\Psi|^2 - \frac{1}{2}e_4(r^{-1}f)|\Psi|^2 \\
&= f\left|e_4\Psi + \frac{1}{r}\Psi\right|^2 - \frac{1}{2}r^{-2}e_4(rf|\Psi|^2) + \frac{1}{2}r^{-2}e_4(rf)|\Psi|^2 - r^{-2}f|\Psi|^2 - \frac{1}{2}e_4(r^{-1}f)|\Psi|^2 \\
&= f\left|e_4\Psi + \frac{1}{r}\Psi\right|^2 - \frac{1}{2}r^{-2}e_4(rf|\Psi|^2) + r^{-2}(e_4(r) - 1)f|\Psi|^2.
\end{aligned}$$

Since

$$e_4(r) = \frac{r}{2}(\kappa + A),$$

we have¹³

$$e_4(r) - 1 = O(\epsilon r^{-1}).$$

Thus, as desired,

$$\mathcal{P} \cdot e_4 = f\left|e_4\Psi + \frac{1}{r}\Psi\right|^2 - \frac{1}{2}r^{-2}e_4(rf|\Psi|^2) + O(\epsilon r^{-3})f|\Psi|^2.$$

Also,

$$\begin{aligned}
\mathcal{P} \cdot e_3 &= f\mathcal{Q}_{34} + r^{-1}f\Psi \cdot e_3\Psi - \frac{1}{2}e_3(r^{-1}f)|\Psi|^2 \\
&= f\mathcal{Q}_{34} + \frac{1}{2}r^{-1}fe_3(|\Psi|^2) - \frac{1}{2}e_3(r^{-1}f)|\Psi|^2 \\
&= f\mathcal{Q}_{34} + \frac{1}{2}\left[r^{-2}e_3(rf|\Psi|^2) - r^{-2}e_3(rf)|\Psi|^2\right] - \frac{1}{2}e_3(r^{-1}f)|\Psi|^2 \\
&= f\mathcal{Q}_{34} + \frac{1}{2}r^{-2}e_3(rf|\Psi|^2) + r^{-1}f'\Upsilon|\Psi|^2 - r^{-1}f'(e_3(r) + \Upsilon)|\Psi|^2 \\
&= f\mathcal{Q}_{34} + \frac{1}{2}r^{-2}e_3(rf|\Psi|^2) + r^{-1}f'|\Psi|^2 + O\left(mr^{-3} + \epsilon r^{-2}\right)(r|f'|)|\Psi|^2
\end{aligned}$$

as desired.

¹³Note that so far we have only used the weaker version $e_4(r) - 1 = O(\epsilon)$. This is the first time we need the stronger version of the estimate in this chapter.

It remains to prove the last part of Proposition 10.2.5. We have, for $r \geq R$

$${}^{(p)}\mathcal{P} \cdot e_4 = r^p |e_4 \Psi|^2 + r^{p-1} \Psi e_4 \Psi - \frac{1}{2} e_4(r^{p-1}) |\Psi|^2$$

and,

$$\begin{aligned} {}^{(p)}\mathcal{P} \cdot e_4 + \frac{p}{2} r^{-2} e_4(\theta r^{p+1} |\Psi|^2) &= r^p |e_4 \Psi|^2 + r^{p-1} \Psi \cdot e_4 \Psi - \frac{1}{2} e_4(r^{p-1}) |\Psi|^2 \\ &+ p r^{p-1} \Psi \cdot e_4 \Psi + \frac{p(p+1)}{2} r^{p-2} e_4(r) |\Psi|^2 \\ &= r^p |e_4 \Psi|^2 + (p+1) r^{p-1} \Psi \cdot e_4 \Psi + \frac{p^2+1}{2} e_4(r) r^{p-2} |\Psi|^2 \\ &= r^p \left[\left| e_4 \Psi + \frac{p+1}{2r} \Psi \right|^2 + \frac{(p-1)^2}{4r^2} |\Psi|^2 \right] \\ &+ \frac{p^2+1}{2} (e_4(r) - 1) r^{p-2} |\Psi|^2 \\ &\geq r^{p-2} \frac{(p-1)^2}{4} |\Psi|^2 - O(\epsilon) \frac{p^2+1}{2} r^{p-3} |\Psi|^2. \end{aligned}$$

This concludes the proof of Proposition 10.2.5. \square

In applications we would like to apply Proposition 10.2.5 to $f = r^p$, $0 < p < 2$. We note however that the presence of the term $-\frac{1}{2} r^{-1} f'' |\Psi|^2$ on the right hand side of the \mathcal{E} identity requires an additional correction if $p > 1$. This additional correction is taken into account by the following proposition.

Proposition 10.2.7. *Assume Ψ verifies the equation $\square_{\mathbf{g}} \Psi = V \Psi + N$ and let $X = f(r) e_4$, $w = {}^{(X)}\Lambda = \frac{2f}{r}$ and $M = 2r^{-1} f' e_4$. Then,*

1. We have, with $\check{e}_4 = r^{-1} e_4(r \cdot)$,

$$\mathcal{E}[X, w, M] = \frac{1}{2} f' |\check{e}_4(\Psi)|^2 + \frac{1}{2} \left(\frac{2f}{r} - f' \right) \mathcal{Q}_{34} + \text{Err} \left(\epsilon, \frac{m}{r}; f \right) [\Psi] \quad (10.2.14)$$

with error term,

$$\begin{aligned} \text{Err} \left(\epsilon, \frac{m}{r}, f \right) (\Psi) &= O \left(\frac{m}{r^2} \right) (|f| + r|f'|) |e_4 \Psi|^2 + O \left(\frac{m}{r^4} + \epsilon w_{3,1} \right) (|f| + r|f'| + r^2 |f''|) |\Psi|^2 \\ &+ O(\epsilon) w_{1,1} (|f| + r|f'|) (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2) \\ &+ O(\epsilon) w_{2,1/2} |f| \left(|e_3 \Psi| (|e_4 \Psi| + r^{-1} |\nabla \Psi|) + |\nabla \Psi|^2 + r^{-2} |\Psi|^2 \right). \end{aligned}$$

2. The current,

$$\mathcal{P}_\mu = \mathcal{P}_\mu[X, w, M] = \mathcal{Q}_{\mu\nu}X^\nu + \frac{1}{2}w\Psi\mathbf{D}_\mu\Psi - \frac{1}{4}|\Psi|^2\partial_\mu w + \frac{1}{4}M_\mu|\Psi|^2$$

verifies,

$$\begin{aligned} \mathcal{P} \cdot e_4 &= f(\check{e}_4\Psi)^2 - \frac{1}{2}r^{-2}e_4(rf|\Psi|^2) + O(\epsilon r^{-1})f(|e_4\Psi|^2 + r^{-2}|\Psi|^2), \\ \mathcal{P} \cdot e_3 &= f\mathcal{Q}_{34} + \frac{1}{2}r^{-2}e_3(rf|\Psi|^2) + O(mr^{-3} + \epsilon r^{-2})(|f| + r|f'|)|\Psi|^2. \end{aligned}$$

3. Let $\theta = \theta(r)$ supported for $r \geq R/2$ with $\theta = 1$ for $r \geq R$ such that $f_p = \theta(r)r^p$. Let ${}^{(p)}\mathcal{P} := \mathcal{P}[f_p e_4, w_p, M_p]$. Then, for all $r \geq R$,

$${}^{(p)}\mathcal{P} \cdot e_4 + \frac{p}{2}r^{-2}e_4(\theta r^{p+1}|\Psi|^2) \geq \frac{1}{8}r^{p-2}(p-1)^2|\Psi|^2.$$

Proof. We start with the first part of Proposition 10.2.7. To this end, we use

$${}^{(X)}\pi_{43} = -2e_4f + 4f\omega, \quad {}^{(X)}\pi_{AB} = 2f^{(1+3)}\chi_{AB}$$

so that

$$\begin{aligned} \text{tr}{}^{(X)}\pi &= -{}^{(X)}\pi_{43} + \mathbf{g}^{AB}{}^{(X)}\pi_{AB} \\ &= 2e_4f - 4f\omega + 2f\kappa, \end{aligned}$$

and we compute

$$\begin{aligned} \mathbf{D}^\mu M_\mu &= \mathbf{D}^\mu(2r^{-1}f'e_4)_\mu = \mathbf{D}^\mu\left(\frac{2f'}{rf}X_\mu\right) = \frac{2f'}{rf}\mathbf{Div}X + X\left(\frac{2f'}{rf}\right) \\ &= \frac{f'}{rf}\text{tr}{}^{(X)}\pi + X\left(\frac{2f'}{rf}\right) \\ &= \frac{f'}{rf}(2e_4f - 4f\omega + 2f\kappa) + 2fe_4\left(\frac{f'}{rf}\right) \\ &= \frac{f'}{rf}\left(2e_4(r)f' + \frac{4f}{r} + O\left(\frac{m}{r^3} + \epsilon w_{2,1}\right)\right) + 2f\left(\frac{f''rf - f'(f + rf')}{r^2f^2}\right)e_4(r) \\ &= \frac{4f'}{r^2} + \frac{2(f')^2}{rf} + 2\frac{f''}{r} - \frac{2f'}{r^2} - \frac{2(f')^2}{rf} + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right)(|f| + r|f'| + r^2|f''|) \\ &= \frac{2f'}{r^2} + \frac{2f''}{r} + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right)(|f| + r|f'| + r^2|f''|). \end{aligned}$$

We also have

$$\frac{1}{2}\Psi \cdot \mathbf{D}_\mu \Psi M^\mu = r^{-1} f' \Psi \cdot \mathbf{D}_4 \Psi.$$

Since we have

$$\mathcal{E}[X, w, M] = \mathcal{E}[X, w] + \frac{1}{4}(\mathbf{D}^\mu M_\mu)|\Psi|^2 + \frac{1}{2}\Psi \cdot \mathbf{D}_\mu \Psi M^\mu,$$

we infer

$$\mathcal{E}[X, w, M] = \mathcal{E}[X, w] + \left(\frac{f'}{2r^2} + \frac{f''}{2r} + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) (|f| + r|f'| + r^2|f''|) \right) |\Psi|^2 + r^{-1} f' \Psi \cdot \mathbf{D}_4 \Psi.$$

Together with Proposition 10.2.5, this yields

$$\begin{aligned} \mathcal{E}[X, w, M] &= \frac{1}{2}f' \left(|e_4 \Psi|^2 + 2r^{-1} \Psi \cdot \mathbf{D}_4 \Psi + r^{-2} |\Psi|^2 \right) + \frac{1}{2} \left(-f' + \frac{2f}{r} \right) (|\nabla \Psi|^2 + V|\Psi|^2) \\ &\quad + \text{Err} \left(\epsilon, \frac{m}{r}, f \right) (\Psi) + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) (|f| + r|f'| + r^2|f''|) |\Psi|^2 \\ &= \frac{1}{2}f' |e_4 \Psi + r^{-1} \Psi|^2 + \frac{1}{2} \left(-f' + \frac{2f}{r} \right) (|\nabla \Psi|^2 + V|\Psi|^2) \\ &\quad + \text{Err} \left(\epsilon, \frac{m}{r}, f \right) (\Psi) + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) (|f| + r|f'| + r^2|f''|) |\Psi|^2 \\ &= \frac{1}{2}f' |\check{e}_4 \Psi + r^{-1}(1 - e_4(r))\Psi|^2 + \frac{1}{2} \left(-f' + \frac{2f}{r} \right) (|\nabla \Psi|^2 + V|\Psi|^2) \\ &\quad + \text{Err} \left(\epsilon, \frac{m}{r}, f \right) (\Psi) + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) (|f| + r|f'| + r^2|f''|) |\Psi|^2 \end{aligned}$$

and hence

$$\mathcal{E}[X, w, M] = \frac{1}{2}f' |\check{e}_4 \Psi|^2 + \frac{1}{2} \left(-f' + \frac{2f}{r} \right) (|\nabla \Psi|^2 + V|\Psi|^2) + \text{Err} \left(\epsilon, \frac{m}{r}, f \right) (\Psi)$$

where

$$\begin{aligned} \text{Err} \left(\epsilon, \frac{m}{r}, f \right) (\Psi) &= O\left(\frac{m}{r^2}\right) (|f| + r|f'|) |e_4 \Psi|^2 + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) (|f| + r|f'| + r^2|f''|) |\Psi|^2 \\ &\quad + O(\epsilon) w_{1,1} (|f| + r|f'|) (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2) \\ &\quad + O(\epsilon) w_{2,1/2} |f| \left(|e_3 \Psi| (|e_4 \Psi| + r^{-1} |\nabla \Psi|) + |\nabla \Psi|^2 + r^{-2} |\Psi|^2 \right). \end{aligned}$$

This is the desired estimate (10.2.14).

Next, we consider the second part of Proposition 10.2.7.

$$\mathcal{P}_\mu[X, w, M] = \mathcal{P}_\mu[X, w] + \frac{1}{4} |\Psi|^2 M_\mu = \mathcal{P}_\mu[X, w] + \frac{1}{2} r^{-1} f' |\Psi|^2 e_4.$$

Hence, in view of the results in part 2 of Proposition 10.2.5,

$$\begin{aligned}
\mathcal{P}_4[X, w, M] &= \mathcal{P}_4[X, w] = f \left| \check{e}_4 \Psi + (1 - e_4(r)) \Psi \right|^2 - \frac{1}{2} r^{-2} e_4(r f |\Psi|^2) + O(\epsilon r^{-3}) f |\Psi|^2 \\
&= f |\check{e}_4 \Psi|^2 - \frac{1}{2} r^{-2} e_4(r f |\Psi|^2) + O(\epsilon r^{-1}) f (|e_4 \Psi|^2 + r^{-2} |\Psi|^2), \\
\mathcal{P}_3[X, w, M] &= \mathcal{P}_3[X, w] - r^{-1} f' |\Psi|^2 = f \mathcal{Q}_{34} + \frac{1}{2} r^{-2} e_3(r f |\Psi|^2) + r^{-1} f' |\Psi|^2 - r^{-1} f' |\Psi|^2 \\
&+ O(\epsilon r^{-2}) (|f| + r |f'|) |\Psi|^2 \\
&= f \mathcal{Q}_{34} + \frac{1}{2} r^{-2} e_3(r f |\Psi|^2) + O(\epsilon r^{-2}) (|f| + r |f'|) |\Psi|^2
\end{aligned}$$

as desired. The last part follows from the third part of Proposition 10.2.5. \square

Lemma 10.2.8. *On Σ_* , we have*

$$\begin{aligned}
\mathcal{P} \cdot N_{\Sigma_*} &= \frac{1}{2} f \mathcal{Q}_{34} + \frac{1}{2} f (\check{e}_4 \Psi)^2 + \frac{1}{2} \operatorname{div}_{\Sigma_*} \left(r^{-1} f |\Psi|^2 \nu_{\Sigma_*} \right) \\
&+ O(mr^{-1} + \epsilon) (|f| + r |f'|) (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2)
\end{aligned}$$

Proof. Recall that there exists a constant c_* such that $u + r = c_*$ on Σ_* . In particular, the unit normal N_{Σ_*} is collinear to

$$\begin{aligned}
-2\mathbf{g}^{\alpha\beta} \partial_\alpha (u + r) \partial_\beta &= e_4(u + r) e_3 + e_3(u + r) e_4 \\
&= e_4(r) e_3 + (e_3(u) + e_3(r)) e_4
\end{aligned}$$

and since

$$g \left(e_4(r) e_3 + (e_3(u) + e_3(r)) e_4, e_4(r) e_3 + (e_3(u) + e_3(r)) e_4 \right) = -4e_4(r) (e_3(u) + e_3(r)),$$

we infer

$$N_{\Sigma_*} = \frac{\sqrt{e_4(r)}}{2\sqrt{e_3(u) + e_3(r)}} e_3 + \frac{\sqrt{e_3(u) + e_3(r)}}{2\sqrt{e_4(r)}} e_4.$$

In particular, we have

$$\mathcal{P} \cdot N_{\Sigma_*} = \frac{\sqrt{e_4(r)}}{2\sqrt{e_3(u) + e_3(r)}} \mathcal{P} \cdot e_3 + \frac{\sqrt{e_3(u) + e_3(r)}}{2\sqrt{e_4(r)}} \mathcal{P} \cdot e_4.$$

Now, recall from Proposition 10.2.7 that we have

$$\begin{aligned}
\mathcal{P} \cdot e_4 &= f (\check{e}_4 \Psi)^2 - \frac{1}{2} r^{-2} e_4(r f |\Psi|^2) + O(\epsilon r^{-1}) f (|e_4 \Psi|^2 + r^{-2} |\Psi|^2), \\
\mathcal{P} \cdot e_3 &= f \mathcal{Q}_{34} + \frac{1}{2} r^{-2} e_3(r f |\Psi|^2) + O(mr^{-3} + \epsilon r^{-2}) (|f| + r |f'|) |\Psi|^2.
\end{aligned}$$

We deduce

$$\begin{aligned}
\mathcal{P} \cdot N_{\Sigma_*} &= \frac{\sqrt{e_4(r)}}{2\sqrt{e_3(u) + e_3(r)}} \left(f\mathcal{Q}_{34} + \frac{1}{2}r^{-2}e_3(rf|\Psi|^2) + O(mr^{-3} + \epsilon r^{-2})(|f| + r|f'|)|\Psi|^2 \right) \\
&\quad + \frac{\sqrt{e_3(u) + e_3(r)}}{2\sqrt{e_4(r)}} \left(f(\check{e}_4\Psi)^2 - \frac{1}{2}r^{-2}e_4(rf|\Psi|^2) + O(\epsilon r^{-1})f(|e_4\Psi|^2 + r^{-2}|\Psi|^2) \right) \\
&= \frac{\sqrt{e_4(r)}}{2\sqrt{e_3(u) + e_3(r)}} f\mathcal{Q}_{34} + \frac{\sqrt{e_3(u) + e_3(r)}}{2\sqrt{e_4(r)}} f(\check{e}_4\Psi)^2 \\
&\quad + \frac{\sqrt{e_4(r)}}{2\sqrt{e_3(u) + e_3(r)}} \frac{1}{2}r^{-2}e_3(rf|\Psi|^2) - \frac{\sqrt{e_3(u) + e_3(r)}}{2\sqrt{e_4(r)}} \frac{1}{2}r^{-2}e_4(rf|\Psi|^2) \\
&\quad + O(mr^{-3} + \epsilon r^{-2})(|f| + r|f'|)|\Psi|^2 + O(\epsilon r^{-1})f(|e_4\Psi|^2 + r^{-2}|\Psi|^2) \\
&= \frac{1}{2\sqrt{2-\Upsilon}}(1 + O(\epsilon))f\mathcal{Q}_{34} + \frac{\sqrt{2-\Upsilon}}{2}(1 + O(\epsilon))f(\check{e}_4\Psi)^2 + \frac{1}{2}r^{-2}\nu_{\Sigma_*}(rf|\Psi|^2) \\
&\quad + O(mr^{-3} + \epsilon r^{-2})(|f| + r|f'|)|\Psi|^2 + O(\epsilon r^{-1})f(|e_4\Psi|^2 + r^{-2}|\Psi|^2) \\
&= \frac{1}{2}f\mathcal{Q}_{34} + \frac{1}{2}f(\check{e}_4\Psi)^2 + \frac{1}{2}r^{-2}\nu_{\Sigma_*}(rf|\Psi|^2) \\
&\quad + O(mr^{-1} + \epsilon)(|f| + r|f'|)(|e_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2)
\end{aligned}$$

where we used

$$e_4(r) = 1 + O(\epsilon), \quad e_3(r) = -\Upsilon + O(\epsilon), \quad e_3(u) = 2 + O(\epsilon),$$

and where ν_{Σ_*} denotes the vectorfield

$$\nu_{\Sigma_*} = \frac{\sqrt{e_4(r)}}{2\sqrt{e_3(u) + e_3(r)}}e_3 - \frac{\sqrt{e_3(u) + e_3(r)}}{2\sqrt{e_4(r)}}e_4.$$

Next, note from the formula that ν_{Σ_*} is unitary and orthogonal to N_{Σ_*} so that ν_{Σ_*} is a unit vectorfield, tangent to Σ_* and normal to e_θ . Furthermore, since $(\nu_{\Sigma_*}, e_\theta, e_\varphi)$ is an orthonormal frame of Σ_* , we have

$$\operatorname{div}_{\Sigma_*}(\nu_{\Sigma_*}) = \mathbf{g}(\mathbf{D}_{\nu_{\Sigma_*}}\nu_{\Sigma_*}, \nu_{\Sigma_*}) + \mathbf{g}(\mathbf{D}_{e_\theta}\nu_{\Sigma_*}, e_\theta) + \mathbf{g}(\mathbf{D}_{e_\varphi}\nu_{\Sigma_*}, e_\varphi).$$

Since ν_{Σ_*} is a unit vector, the first term vanishes, and hence

$$\begin{aligned}
\operatorname{div}_{\Sigma_*}(\nu_{\Sigma_*}) &= \mathbf{g}(\mathbf{D}_{e_\theta}\nu_{\Sigma_*}, e_\theta) + \mathbf{g}(\mathbf{D}_{e_\varphi}\nu_{\Sigma_*}, e_\varphi) \\
&= \frac{\sqrt{e_4(r)}}{2\sqrt{e_3(u) + e_3(r)}}g(D_\theta e_3, e_\theta) - \frac{\sqrt{e_3(u) + e_3(r)}}{2\sqrt{e_4(r)}}g(D_\theta e_4, e_\theta) + \nu_{\Sigma_*}(\Phi) \\
&= \frac{\sqrt{e_4(r)}}{2\sqrt{e_3(u) + e_3(r)}}\kappa - \frac{\sqrt{e_3(u) + e_3(r)}}{2\sqrt{e_4(r)}}\kappa.
\end{aligned}$$

In particular, we have

$$\begin{aligned}
\operatorname{div}_{\Sigma_*} \left(r^{-1} f |\Psi|^2 \nu_{\Sigma_*} \right) &= r^{-2} \nu_{\Sigma_*} (r f |\Psi|^2) + \nu_{\Sigma_*} (r^{-2}) r f |\Psi|^2 + \operatorname{div}_{\Sigma_*} (\nu_{\Sigma_*}) r^{-1} f |\Psi|^2 \\
&= r^{-2} \nu_{\Sigma_*} (r f |\Psi|^2) + \left(\operatorname{div}_{\Sigma_*} (\nu_{\Sigma_*}) - \frac{2\nu_{\Sigma_*}(r)}{r} \right) r^{-1} f |\Psi|^2 \\
&= r^{-2} \nu_{\Sigma_*} (r f |\Psi|^2) + \left(\frac{\sqrt{e_4(r)}}{2\sqrt{e_3(u) + e_3(r)}} \left(\underline{\kappa} - \frac{2e_3(r)}{r} \right) \right. \\
&\quad \left. - \frac{\sqrt{e_3(u) + e_3(r)}}{2\sqrt{e_4(r)}} \left(\kappa - \frac{2e_4(r)}{r} \right) \right) r^{-1} f |\Psi|^2
\end{aligned}$$

and hence

$$r^{-2} \nu_{\Sigma_*} (r f |\Psi|^2) = \operatorname{div}_{\Sigma_*} \left(r^{-1} f |\Psi|^2 \nu_{\Sigma_*} \right) + O(\epsilon r^{-2}) f |\Psi|^2$$

We finally obtain

$$\begin{aligned}
\mathcal{P} \cdot N_{\Sigma_*} &= \frac{1}{2} f \mathcal{Q}_{34} + \frac{1}{2} f (\check{e}_4 \Psi)^2 + \frac{1}{2} r^{-2} \nu_{\Sigma_*} (r f |\Psi|^2) \\
&\quad + O(mr^{-1} + \epsilon) (|f| + r|f'|) (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2) \\
&= \frac{1}{2} f \mathcal{Q}_{34} + \frac{1}{2} f (\check{e}_4 \Psi)^2 + \frac{1}{2} \operatorname{div}_{\Sigma_*} \left(r^{-1} f |\Psi|^2 \nu_{\Sigma_*} \right) \\
&\quad + O(mr^{-1} + \epsilon) (|f| + r|f'|) (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2)
\end{aligned}$$

which concludes the proof of the lemma. \square

10.2.3 Proof of Theorem 10.2.1

Consider the function $f_p = f_{p,R}$ defined by,

$$f_p = \begin{cases} r^p, & \text{if } r \geq R, \\ 0, & \text{if } r \leq \frac{R}{2}, \end{cases} \quad (10.2.15)$$

where R is a fixed, sufficiently large constant which will be chosen in the proof. We also consider

$$X_p = f_p e_4, \quad w_p = \frac{2f_p}{r}, \quad M_p = \frac{2f'_p}{r} e_4.$$

The proof relies on Proposition 10.2.7.

Step 0. (Reduction to the region $r \geq R$) In view of the definition of $\mathcal{E}[X_p, w_p, M_p]$, see (10.1.12), and in view of the choice of X_p and w_p , we have

$$\mathbf{D}^\mu \mathcal{P}_\mu[X_p, w_p, M_p] = \mathcal{E}[X_p, w_p, M_p] + f_p(r) \check{e}_4 \Psi \cdot N.$$

We integrate this identity on the domain $\mathcal{M}(\tau_1, \tau_2)$ to derive,

$$\int_{\Sigma(\tau_2)} \mathcal{P} \cdot N_\Sigma + \int_{\Sigma_*(\tau_1, \tau_2)} \mathcal{P} \cdot N_{\Sigma_*} + \int_{\mathcal{M}(\tau_1, \tau_2)} \mathcal{E} = \int_{\Sigma(\tau_1)} \mathcal{P} \cdot N_\Sigma - \int_{\mathcal{M}(\tau_1, \tau_2)} f_p \check{e}_4 \Psi \cdot N.$$

Denoting the boundary terms,

$$\begin{aligned} K_{\geq R}(\tau_1, \tau_2) &:= \int_{\Sigma_{\geq R}(\tau_2)} \mathcal{P} \cdot e_4 + \int_{\Sigma_*(\tau_1, \tau_2)} \mathcal{P} \cdot N_{\Sigma_*} - \int_{\Sigma_{\geq R}(\tau_1)} \mathcal{P} \cdot e_4, \\ K_{\leq R}(\tau_1, \tau_2) &:= \int_{\Sigma_{\leq R}(\tau_1)} \mathcal{P} \cdot N_\Sigma - \int_{\Sigma_{\leq R}(\tau_2)} \mathcal{P} \cdot N_\Sigma, \end{aligned}$$

we write,

$$K_{\geq R}(\tau_1, \tau_2) + \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} \mathcal{E} = K_{\leq R}(\tau_1, \tau_2) - \int_{\mathcal{M}_{\leq R}(\tau_1, \tau_2)} \mathcal{E} - \int_{\mathcal{M}(\tau_1, \tau_2)} f_p \check{e}_4 \Psi \cdot N. \quad (10.2.16)$$

We have the following lemma.

Lemma 10.2.9. *For $p \geq \delta$, we have*

$$K_{\geq R}(\tau_1, \tau_2) + \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} \mathcal{E} \lesssim R^{p+2} \left(E[\Psi](\tau_1) + J_p[N, \psi](\tau_1, \tau_2) + O(\epsilon) \dot{B}_{\delta; 4m_0}^s[\psi](\tau_1, \tau_2) \right).$$

Proof of Lemma 10.2.9. The terms $\int_{\Sigma_{\leq R}(\tau)} \mathcal{P} \cdot N_\Sigma$ and $\int_{\mathcal{M}_{\leq R}(\tau_1, \tau_2)} \mathcal{E}$ on the right can be estimated as follows

$$\begin{aligned} \left| \int_{\Sigma_{\leq R}(\tau_1)} \mathcal{P} \cdot N_\Sigma \right| &\lesssim R^p E[\Psi](\tau_1), \\ \left| \int_{\Sigma_{\leq R}(\tau_2)} \mathcal{P} \cdot N_\Sigma \right| &\lesssim R^p E[\Psi](\tau_2), \\ \left| \int_{\mathcal{M}_{\leq R}(\tau_1, \tau_2)} \mathcal{E} \right| &\lesssim R^{p+2} \text{Mor}[\Psi](\tau_1, \tau_2). \end{aligned}$$

Hence,

$$K_{\leq R}(\tau_1, \tau_2) - \int_{\mathcal{M}_{\leq R}(\tau_1, \tau_2)} \mathcal{E} \lesssim R^{p+2} (E[\Psi](\tau_1) + E[\Psi](\tau_2) + \text{Mor}[\Psi](\tau_1, \tau_2)).$$

In view of the improved Morawetz Theorem 10.1.1 we have, for fixed $\delta > 0$,

$$E[\Psi](\tau_2) + \text{Morr}[\Psi](\tau_1, \tau_2) + F[\Psi](\tau_1, \tau_2) \lesssim E[\Psi](\tau_1) + J_\delta[N, \psi](\tau_1, \tau_2) + O(\epsilon)\dot{B}_{\delta; 4m_0}^s[\psi](\tau_1, \tau_2)$$

which implies

$$K_{\geq R}(\tau_1, \tau_2) + \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} \mathcal{E} \lesssim R^{p+2} \left(E[\Psi](\tau_1) + J_\delta[N, \psi](\tau_1, \tau_2) + O(\epsilon)\dot{B}_{\delta; 4m_0}^s[\psi](\tau_1, \tau_2) \right) + \left| \int_{\mathcal{M}(\tau_1, \tau_2)} f_p \check{e}_4 \Psi \cdot N \right|.$$

Together with the definition (5.3.7) of J_p and the fact that $p \geq \delta$, we infer

$$K_{\geq R}(\tau_1, \tau_2) + \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} \mathcal{E} \lesssim R^{p+2} \left(E[\Psi](\tau_1) + J_p[N, \psi](\tau_1, \tau_2) + O(\epsilon)\dot{B}_{\delta; 4m_0}^s[\psi](\tau_1, \tau_2) \right)$$

which concludes the proof of Lemma 10.2.9. \square

The proof of Theorem 10.2.1 now proceeds according to the following steps.

Step 1. (Bulk terms for $r \geq R$) We prove the following lower bound for $\int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} \mathcal{E}$.

Lemma 10.2.10. *Given a fixed $\delta > 0$ we have for all $\delta \leq p \leq 2 - \delta$ and $R \gg \frac{m}{\delta}$, $\epsilon \ll \delta$,*

$$\int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} \mathcal{E} \geq \frac{1}{4} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{p-1} \left(p |\check{e}_4(\Psi)|^2 + (2-p)(|\nabla \Psi|^2 + r^{-2} |\Psi|^2) \right) - O(\epsilon) \text{Morr}[\Psi](\tau_1, \tau_2). \quad (10.2.17)$$

Proof of Lemma 10.2.10. We make use of Proposition 10.2.7 according to which,

$$\begin{aligned} \mathcal{E}[X, w, M] &= \frac{1}{2} f'_p |\check{e}_4(\Psi)|^2 + \frac{1}{2} \left(\frac{2f_p}{r} - f'_p \right) \mathcal{Q}_{34} + \text{Err} \left(\epsilon, \frac{m}{r}; f_p \right) [\Psi] \\ &= r^{p-1} \left[\frac{p}{2} |\check{e}_4(\Psi)|^2 + \frac{1}{2} (2-p)(|\nabla \Psi|^2 + V |\Psi|^2) \right] + \text{Err} \left(\epsilon, \frac{m}{r}; f_p \right) [\Psi] \\ &\geq r^{p-1} \left[\frac{p}{2} |\check{e}_4(\Psi)|^2 + \frac{1}{2} (2-p)(|\nabla \Psi|^2 + r^{-2} |\Psi|^2) \right] + \text{Err} \left(\epsilon, \frac{m}{r}; f_p \right) [\Psi] \end{aligned}$$

where,

$$\begin{aligned}
\text{Err}\left(\epsilon, \frac{m}{r}, f_p\right)(\Psi) &= r^p O\left(\frac{m}{r^2}\right) [|e_4\Psi|^2 + r^{-2}|\Psi|^2] + r^p O(\epsilon) w_{1,1} (|e_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2) \\
&+ r^p O(\epsilon) w_{2,1/2} \left(|e_3\Psi| (|e_4\Psi| + r^{-1}|\nabla\Psi|) + |\nabla\Psi|^2 + r^{-2}|\Psi|^2 \right) \\
&\lesssim \text{Err}\left(\frac{m}{r}\right) + \text{Err}(\epsilon), \\
\text{Err}\left(\frac{m}{r}\right) &= O\left(\frac{m}{r}\right) r^{p-1} [|\check{e}_4\Psi|^2 + r^{-2}|\Psi|^2], \\
\text{Err}(\epsilon) &= O(\epsilon) r^{p-1} (|\check{e}_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2 + r^{-2}|e_3\Psi|^2).
\end{aligned}$$

Thus,

$$\begin{aligned}
\int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} \mathcal{E} &\geq \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{p-1} \left(\frac{p}{2} |\check{e}_4(\Psi)|^2 + \frac{1}{2} (2-p) (|\nabla\Psi|^2 + r^{-2}|\Psi|^2) \right) \\
&- O\left(\frac{m}{R}\right) \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{p-1} [|\check{e}_4\Psi|^2 + r^{-2}|\Psi|^2] \\
&- O(\epsilon) \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{p-1} (|\check{e}_4\Psi|^2 + |\nabla\Psi|^2 + r^{-2}|\Psi|^2) \\
&- O(\epsilon) \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{p-3} |e_3\Psi|^2.
\end{aligned}$$

For $\delta \leq p \leq 2 - \delta$, $R \gg \frac{m}{\delta}$ and $\epsilon \ll \delta$ we can absorb all error terms except the last, i.e.

$$\begin{aligned}
\int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} \mathcal{E} &\geq \frac{1}{4} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{p-1} \left(p |\check{e}_4(\Psi)|^2 + (2-p) (|\nabla\Psi|^2 + r^{-2}|\Psi|^2) \right) \\
&- O(\epsilon) \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{p-3} |e_3\Psi|^2.
\end{aligned}$$

Note also that for all $\delta \leq p \leq 2 - \delta$ we have,

$$\int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{p-3} |e_3\Psi|^2 \lesssim \text{Morr}(\tau_1, \tau_2).$$

Hence, for all $\delta \leq p \leq 2 - \delta$ and $R \gg \frac{m}{\delta}$, $\epsilon \ll \delta$,

$$\begin{aligned}
\int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} \mathcal{E} &\geq \frac{1}{4} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{p-1} \left(p |\check{e}_4(\Psi)|^2 + (2-p) (|\nabla\Psi|^2 + r^{-2}|\Psi|^2) \right) \\
&- O(\epsilon) \text{Morr}[\Psi](\tau_1, \tau_2)
\end{aligned}$$

as desired. □

Combining (10.2.17) with Lemma 10.2.9, we deduce,

$$K_{\geq R}(\tau_1, \tau_2) + \dot{B}_{p,R}[\Psi](\tau_1, \tau_2) \lesssim R^{p+2} \left(E[\Psi](\tau_1) + J_p[N, \psi](\tau_1, \tau_2) \right) \quad (10.2.18) \\ + O(\epsilon) \dot{B}_{\delta; 4m_0}^s[\psi](\tau_1, \tau_2).$$

Step 2. (Boundary terms for $r \geq R$.) Recall that according to Proposition 10.2.7,

$$\mathcal{P} \cdot e_4 = f_p |\check{e}_4 \Psi|^2 - \frac{1}{2} r^{-2} e_4 (r f_p |\Psi|^2) + O(\epsilon r^{-1}) f_p (|e_4 \Psi|^2 + r^{-2} |\Psi|^2),$$

and according to Lemma 10.2.8

$$\mathcal{P} \cdot N_{\Sigma_*} = \frac{1}{2} f \mathcal{Q}_{34} + \frac{1}{2} f (\check{e}_4 \Psi)^2 + \frac{1}{2} \operatorname{div}_{\Sigma_*} \left(r^{-1} f |\Psi|^2 \nu_{\Sigma_*} \right) \\ + O(mr^{-1} + \epsilon) (|f| + r|f'|) (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2)$$

Recalling the definition of

$$K_{\geq R} = \int_{\Sigma_{\geq R}(\tau_2)} \mathcal{P} \cdot e_4 + \int_{\Sigma_*(\tau_1, \tau_2)} \mathcal{P} \cdot N_{\Sigma_*} - \int_{\Sigma_{\geq R}(\tau_1)} \mathcal{P} \cdot e_4$$

we write,

$$K_{\geq R} = \int_{\Sigma_{\geq R}(\tau_2)} f_p |\check{e}_4 \Psi|^2 + \frac{1}{2} \int_{\Sigma_*(\tau_1, \tau_2)} r^p \left(\mathcal{Q}_{34} + (\check{e}_4 \Psi)^2 \right) - \int_{\Sigma_{\geq R}(\tau_1)} f_p (\check{e}_4 \Psi)^2 \\ - \frac{1}{2} \int_{\Sigma_{\geq R}(\tau_2)} r^{-2} e_4 (r f_p |\Psi|^2) + \frac{1}{2} \int_{\Sigma_*(\tau_1, \tau_2)} \operatorname{div}_{\Sigma_*} \left(r^{-1} f |\Psi|^2 \nu_{\Sigma_*} \right) \\ + \frac{1}{2} \int_{\Sigma_{\geq R}(\tau_1)} r^{-2} e_4 (r f_p |\Psi|^2) + \int_{\Sigma_*(\tau_1, \tau_2)} O(mr^{-1} + \epsilon) r^{p-2} (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2) \\ + O(\epsilon) \left(\int_{\Sigma_{\geq R}(\tau_2)} r^{p-1} (|e_4 \Psi|^2 + r^{-2} |\Psi|^2) - \int_{\Sigma_{\geq R}(\tau_1)} r^{p-1} (|e_4 \Psi|^2 + r^{-2} |\Psi|^2) \right).$$

Now, the following integrations by parts hold true

$$\begin{aligned}
& \int_{\Sigma_{\geq R}(\tau)} r^{-2} e_4(r f_p |\Psi|^2) \\
&= \int_{r \geq R} \left(\int_{S_r} r^{-2} e_4(r f_p |\Psi|^2) \right) \frac{1}{e_4(r)} \\
&= \int_{r \geq R} \frac{1}{e_4(r)} e_4 \left(\int_{S_r} r^{-1} f_p |\Psi|^2 \right) - \int_{\Sigma_{\geq R}(\tau)} (e_4(r^{-2}) + \kappa r^{-2}) r f_p |\Psi|^2 \\
&= \int_{S_*(\tau)} r^{p-1} |\Psi|^2 - \int_{S_R(\tau)} r^{p-1} |\Psi|^2 - \int_{\Sigma_{\geq R}(\tau)} \left(\kappa - \frac{2e_4(r)}{r} \right) r^{-1} f_p |\Psi|^2 \\
&= \int_{S_*(\tau)} r^{p-1} |\Psi|^2 - \int_{S_R(\tau)} r^{p-1} |\Psi|^2 + O(\epsilon) \int_{\Sigma_{\geq R}(\tau)} r^{p-3} |\Psi|^2
\end{aligned}$$

and

$$\int_{\Sigma_*(\tau_1, \tau_2)} \operatorname{div}_{\Sigma_*} \left(r^{-1} f |\Psi|^2 \nu_{\Sigma_*} \right) = \int_{S_*(\tau_2)} r^{p-1} |\Psi|^2 - \int_{S_*(\tau_1)} r^{p-1} |\Psi|^2$$

where $S_*(\tau)$ denotes the 2-sphere $\Sigma_* \cap \Sigma(\tau)$. Note that the boundary terms cancel, except the one on $r = R$, and hence

$$\begin{aligned}
K_{\geq R} &= \int_{\Sigma_{\geq R}(\tau_2)} f_p |\check{e}_4 \Psi|^2 + \frac{1}{2} \int_{\Sigma_*(\tau_1, \tau_2)} r^p \left(\mathcal{Q}_{34} + (\check{e}_4 \Psi)^2 \right) - \int_{\Sigma_{\geq R}(\tau_1)} f_p |\check{e}_4 \Psi|^2 \\
&+ \int_{\Sigma_*(\tau_1, \tau_2)} O(mr^{-1} + \epsilon) r^{p-2} (|e_4 \Psi|^2 + |\nabla \Psi|^2 + r^{-2} |\Psi|^2) \\
&+ O(\epsilon) \left(\int_{\Sigma_{\geq R}(\tau_2)} r^{p-1} (|e_4 \Psi|^2 + r^{-2} |\Psi|^2) - \int_{\Sigma_{\geq R}(\tau_1)} r^{p-1} (|e_4 \Psi|^2 + r^{-2} |\Psi|^2) \right) \\
&+ \frac{1}{2} \int_{S_R(\tau_2)} r^{p-1} |\Psi|^2 - \frac{1}{2} \int_{S_R(\tau_1)} r^{p-1} |\Psi|^2.
\end{aligned}$$

Using

$$\begin{aligned}
\mathcal{Q}_{34} + |\check{e}_4 \Psi|^2 &= |\nabla \Psi|^2 + \frac{4\Upsilon}{r^2} |\Psi|^2 + \left| e_4 \Psi + \frac{1}{r} \Psi \right|^2 \\
&= |\nabla \Psi|^2 + \frac{4\Upsilon}{r^2} |\Psi|^2 + (e_4 \Psi)^2 + \frac{1}{r^2} |\Psi|^2 + \frac{2}{r} \Psi \cdot e_4(\Psi) \\
&\geq |\nabla \Psi|^2 + \frac{4\Upsilon - 3}{r^2} |\Psi|^2 + \frac{2}{3} |e_4 \Psi|^2
\end{aligned}$$

and the fact that $4\Upsilon \geq 3 + 2/3$ for $r \geq R$ and R large enough, we infer

$$\begin{aligned} K_{\geq R} &\geq \frac{1}{2} \left(\int_{\Sigma_{\geq R}(\tau_2)} r^p |\check{e}_4 \Psi|^2 - \int_{\Sigma_{\geq R}(\tau_1)} r^p |\check{e}_4 \Psi|^2 + \dot{F}_p[\Psi](\tau_1, \tau_2) \right) \\ &+ O(\epsilon) \left(\int_{\Sigma_{\geq R}(\tau_2)} r^{p-3} |\Psi|^2 - \int_{\Sigma_{\geq R}(\tau_1)} r^{p-3} |\Psi|^2 \right) \\ &+ \frac{1}{2} \int_{S_R(\tau_2)} r^{p-1} |\Psi|^2 - \frac{1}{2} \int_{S_R(\tau_1)} r^{p-1} |\Psi|^2 \end{aligned} \quad (10.2.19)$$

Next, recall that according to Proposition 10.2.7, we have

$$\mathcal{P} \cdot e_4 \geq \frac{1}{8} r^{p-2} (p-1)^2 |\Psi|^2 - \frac{p}{2} r^{-2} e_4 (r f_p |\Psi|^2).$$

We infer

$$\int_{\Sigma_{\geq R}(\tau_2)} \mathcal{P} \cdot e_4 \geq \frac{1}{8} \int_{\Sigma_{\geq R}(\tau_2)} r^{p-2} (p-1)^2 |\Psi|^2 - \frac{p}{2} \int_{\Sigma_{\geq R}(\tau_2)} r^{-2} e_4 (r f_p |\Psi|^2).$$

Integrating by parts similarly as before, we infer

$$\begin{aligned} \int_{\Sigma_{\geq R}(\tau_2)} \mathcal{P} \cdot e_4 &\geq \frac{1}{8} \int_{\Sigma_{\geq R}(\tau_2)} r^{p-2} (p-1)^2 |\Psi|^2 - \frac{p}{2} \int_{S_*(\tau)} r^{p-1} |\Psi|^2 \\ &+ \frac{p}{2} \int_{S_R(\tau)} r^{p-1} |\Psi|^2 + O(\epsilon) \int_{\Sigma_{\geq R}(\tau)} r^{p-3} |\Psi|^2. \end{aligned}$$

Arguing as for the proof of (10.2.19) except for the boundary term on $\Sigma_{\geq R}(\tau_2)$ for which we use the above estimate, we deduce

$$\begin{aligned} K_{\geq R} &\geq \frac{1}{8} \int_{\Sigma_{\geq R}(\tau_2)} r^{p-2} (p-1)^2 |\Psi|^2 + \frac{1}{2} \left(- \int_{\Sigma_{\geq R}(\tau_1)} r^p |\check{e}_4 \Psi|^2 + \dot{F}_p[\Psi](\tau_1, \tau_2) \right) \\ &+ O(\epsilon) \left(\int_{\Sigma_{\geq R}(\tau_2)} r^{p-1} (|e_4 \Psi|^2 + r^{-2} |\Psi|^2) - \int_{\Sigma_{\geq R}(\tau_1)} r^{p-1} (|e_4 \Psi|^2 + r^{-2} |\Psi|^2) \right) \\ &+ \frac{1-p}{2} \int_{S_*(\tau_2)} r^{p-1} |\Psi|^2 + \frac{p}{2} \int_{S_R(\tau_2)} r^{p-1} |\Psi|^2 - \frac{1}{2} \int_{S_R(\tau_1)} r^{p-1} |\Psi|^2. \end{aligned}$$

We first focus on the case $\delta \leq p \leq 1 - \delta$, in which case the previous estimate yields

$$\begin{aligned} K_{\geq R} &\geq \frac{\delta^2}{8} \int_{\Sigma_{\geq R}(\tau_2)} r^{p-2} |\Psi|^2 + \frac{1}{2} \left(- \int_{\Sigma_{\geq R}(\tau_1)} r^p |\check{e}_4 \Psi|^2 + \dot{F}_p[\Psi](\tau_1, \tau_2) \right) \\ &+ O(\epsilon) \left(\int_{\Sigma_{\geq R}(\tau_2)} r^{p-1} (|e_4 \Psi|^2 + r^{-2} |\Psi|^2) - \int_{\Sigma_{\geq R}(\tau_1)} r^{p-3} |\Psi|^2 \right) \\ &+ \frac{1}{2} \int_{S_R(\tau_2)} r^{p-1} |\Psi|^2 - \frac{1}{2} \int_{S_R(\tau_1)} r^{p-1} |\Psi|^2. \end{aligned}$$

Together with (10.2.19) and the fact that $\epsilon \ll \delta^2$ by assumption, we infer in view of the definition of $\dot{E}_{p,R}[\Psi]$ for $\delta \leq p \leq 1 - \delta$,

$$\dot{E}_{p,R}[\Psi](\tau_2) + \dot{F}_p[\Psi](\tau_1, \tau_2) \lesssim K_{\geq R} + \dot{E}_{p,R}[\Psi](\tau_1) + \int_{S_R(\tau_2)} r^{p-1} |\Psi|^2.$$

Together with (10.2.18), we deduce for $\delta \leq p \leq 1 - \delta$

$$\begin{aligned} &\dot{E}_{p,R}[\Psi](\tau_2) + \dot{F}_p[\Psi](\tau_1, \tau_2) + \dot{B}_{p,R}[\Psi](\tau_1, \tau_2) \\ &\lesssim \dot{E}_{p,R}[\Psi](\tau_1) + \int_{S_R(\tau_2)} r^{p-1} |\Psi|^2 + R^{p+2} \left(E[\Psi](\tau_1) + J_p[N, \psi](\tau_1, \tau_2) \right. \\ &\quad \left. + O(\epsilon) \dot{B}_{\delta; 4m_0}^s[\psi](\tau_1, \tau_2) \right). \end{aligned}$$

In view of the improved Morawetz Theorem 10.1.1, and thanks also to the term $\dot{B}_{p,R}[\Psi](\tau_1, \tau_2)$ on the left hand side, we may absorb the term $O(\epsilon) \dot{B}_{\delta; 4m_0}^s[\psi](\tau_1, \tau_2)$ and obtain

$$\begin{aligned} &\dot{E}_{p,R}[\Psi](\tau_2) + \dot{F}_p[\Psi](\tau_1, \tau_2) + \dot{B}_{p,R}[\Psi](\tau_1, \tau_2) \\ &\lesssim R^{p+2} \left(E_p[\Psi](\tau_1) + J_p[N, \psi](\tau_1, \tau_2) \right) \end{aligned} \quad (10.2.20)$$

which is the desired estimate in the case $\delta \leq p \leq 1 - \delta$.

Finally, we focus on the remaining case, i.e. $1 - \delta \leq p \leq 2 - \delta$. Combining (10.2.19) and (10.2.18), arguing as in the proof of (10.2.20), and in view of the definition of $\dot{E}_{p,R}[\Psi]$ for $1 - \delta \leq p \leq 2 - \delta$, we obtain

$$\begin{aligned} &\dot{E}_{p,R}[\Psi](\tau_2) + \dot{F}_p[\Psi](\tau_1, \tau_2) + \dot{B}_{p,R}[\Psi](\tau_1, \tau_2) \\ &\lesssim R^{p+2} (E_p[\Psi](\tau_1) + J_p[N, \psi](\tau_1, \tau_2)) \\ &\quad + O(\epsilon) \int_{\Sigma_{\geq R}(\tau_2)} r^{p-3} |\Psi|^2 + \int_{\Sigma_{\geq R}(\tau_2)} r^{-1-\delta} |\Psi|^2 \\ &\lesssim R^{p+2} (E_p[\Psi](\tau_1) + J_p[N, \psi](\tau_1, \tau_2)) + E_{1-\delta}[\Psi](\tau_2) \end{aligned}$$

where we also used the fact that $p \leq 2 - \delta$ so that $p - 3 \leq -1 - \delta$. Together with the fact that

$$\dot{E}_{p,R}[\Psi](\tau) \geq \dot{E}_{1-\delta,R}[\Psi](\tau) \text{ for } p \geq 1 - \delta$$

and (10.2.20), we infer

$$\dot{E}_{p,R}[\Psi](\tau_2) + \dot{F}_p[\Psi](\tau_1, \tau_2) + \dot{B}_{p,R}[\Psi](\tau_1, \tau_2) \lesssim R^{p+2} (E_p[\Psi](\tau_1) + J_p[N, \psi](\tau_1, \tau_2))$$

for all $\delta \leq p \leq 2 - \delta$ as desired. This concludes the proof of Theorem 10.2.1.

10.3 Higher Weighted Estimates

We use a variation of the method of [4] to derive slightly stronger weighted estimates. This allows us to prove Theorem 5.3.5 for $s = 0$ in section 10.4.6. The proof for higher order derivatives $s \leq k_{small} + 29$ will be provided in section 10.4.6.

As in the previous section we rely on the assumptions (10.2.8)–(10.2.11) to which we add,

RP5. The assumptions **RP0–RP4** hold true for one extra derivative with respect to \mathfrak{d} .

RP6. $e_4(m)$ satisfies the following improvement of **RP4**

$$|\mathfrak{d}^{\leq 2} e_4(m)| \lesssim \epsilon w_{2,1}. \tag{10.3.1}$$

10.3.1 Wave equation for $\check{\psi}$

Proposition 10.3.1. *Assume ψ verifies $\square_2 \psi = -\kappa \underline{\kappa} \psi + N$. Then $\check{\psi} = f_2 \check{e}_4 \psi$ verifies:*

1. *In the region $r \geq 6m_0$,*

$$\begin{aligned} (\square_2 + \kappa \underline{\kappa}) \check{\psi} &= r^2 \left(e_4(N) + \frac{3}{r} N \right) + \frac{2}{r} \left(1 - \frac{3m}{r} \right) e_4 \check{\psi} + O(r^{-2}) \mathfrak{d}^{\leq 1} \psi \\ &\quad + r \Gamma_b e_4 \mathfrak{d} \psi + \mathfrak{d}^{\leq 1}(\Gamma_b) \mathfrak{d}^{\leq 1} \psi + r \mathfrak{d}^{\leq 1}(\Gamma_g) e_3 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^2 \psi. \end{aligned}$$

2. *In the region $4m_0 \leq r \leq 6m_0$,*

$$(\square_2 + \kappa \underline{\kappa}) \check{\psi} = f_2 \left(e_4(N) + \frac{3}{r} N \right) + O(1) \mathfrak{d}^{\leq 2} \psi.$$

The proof of Proposition 10.3.1 is postponed to Appendix D.4.

10.3.2 The r^p weighted estimates for $\check{\psi}$

The goal of this section is to prove Theorem 5.3.5 in the case $s = 0$. The proof for higher order derivatives $s \leq k_{small} + 29$ will be provided in section 10.4.6.

Proof of Theorem 5.3.5 in the case $s = 0$. We write, in accordance to Proposition 10.3.1

$$\square \check{\psi} - V \check{\psi} = \check{N} + f_2 \left(e_4 + \frac{3}{r} \right) N$$

where,

$$\check{N} = \begin{cases} \frac{2}{r} \left(1 - \frac{3m}{r} \right) e_4 \check{\psi} + O(r^{-2}) \mathfrak{d}^{\leq 1} \psi \\ + r \Gamma_b e_4 \mathfrak{d} \psi + \mathfrak{d}^{\leq 1}(\Gamma_b) \mathfrak{d}^{\leq 1} \psi + r \mathfrak{d}^{\leq 1}(\Gamma_g) e_3 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^2 \psi, & r \geq 6m_0, \\ O(1) \mathfrak{d}^{\leq 2} \psi, & 4m_0 \leq r \leq 6m_0. \end{cases} \quad (10.3.2)$$

We apply the first part of Proposition 10.2.7 to ψ replaced by $\check{\psi}$. This yields, using also (10.1.12),

$$\mathbf{Div} \mathcal{P}_q(\check{\psi}) = \mathcal{E}_q(\check{\psi}) + f_q \check{e}_4 \check{\psi} \cdot \check{N} + f_q \check{e}_4 \check{\psi} \cdot f_2 \left(e_4 + \frac{3}{r} \right) N,$$

where, with $f = f_q$, $X_q = f_q e_4$, $w_q = \frac{2f_q}{r}$, $M_q = 2r^{-1} f'_q e_4$,

$$\mathcal{E}_q(\check{\psi}) = \mathcal{E}[X_q, w_q, M_q] = \frac{1}{2} f'_q |\check{e}_4(\check{\psi})|^2 + \frac{1}{2} \left(\frac{2f_q}{r} - f'_q \right) \mathcal{Q}_{34}(\check{\psi}) + \text{Err}_q(\check{\psi}),$$

$$\begin{aligned} \text{Err}_q(\check{\psi}) &:= \text{Err} \left(\epsilon, \frac{m}{r}; f_q \right) [\check{\psi}] \\ &= O \left(\frac{m}{r^2} \right) r^q |e_4 \check{\psi}|^2 + O \left(\frac{m}{r^4} + \epsilon w_{3,1} \right) r^q |\check{\psi}|^2 + O(\epsilon) w_{1,1} r^q (|e_4 \check{\psi}|^2 + |\nabla \check{\psi}|^2 + r^{-2} |\check{\psi}|^2) \\ &\quad + O(\epsilon) w_{2,1/2} r^q \left((|e_4 \check{\psi}| + r^{-1} |\nabla \check{\psi}|) |e_3 \check{\psi}| + |\nabla \check{\psi}|^2 + r^{-2} |\check{\psi}|^2 \right), \end{aligned}$$

$$\mathcal{P}_k(\check{\psi}) = \mathcal{P}[X_q, w_q, M_q](\check{\psi}).$$

We then integrate on the domain $\mathcal{M}(\tau_1, \tau_2)$ to derive, exactly as in the proof of Theorem 10.2.1 (see section 10.2.3),

$$\begin{aligned} &\int_{\Sigma(\tau_2)} \mathcal{P}_q \cdot e_4 + \int_{\Sigma_*(\tau_1, \tau_2)} \mathcal{P}_q \cdot N_{\Sigma_*} + \int_{\mathcal{M}(\tau_1, \tau_2)} (\mathcal{E}_q + f_q \check{e}_4 \check{\psi} \check{N}) \\ &= \int_{\Sigma(\tau_1)} \mathcal{P}_q \cdot e_4 - \int_{\mathcal{M}(\tau_1, \tau_2)} f_q \check{e}_4 \check{\psi} \cdot f_2 \left(e_4 + \frac{3}{r} \right) N. \end{aligned} \quad (10.3.3)$$

All terms can be treated exactly as in the proof of Theorem 10.2.1, except for the bulk term, i.e. we obtain the following analog of (10.2.1)

$$\begin{aligned} & \dot{E}_{q,R}[\check{\psi}](\tau_2) + \int_{\mathcal{M}(\tau_1, \tau_2)} (\mathcal{E}_q + r^q \check{e}_4(\check{\psi})\check{N}) + \dot{F}_q[\check{\psi}](\tau_1, \tau_2) \\ & \lesssim E_q[\check{\psi}](\tau_1) + J_q \left[\check{\psi}, f_2 \left(e_4 + \frac{3}{r} \right) N \right] (\tau_1, \tau_2). \end{aligned}$$

Since all terms for $r \leq R$ can be controlled by one derivative of ψ , we infer

$$\begin{aligned} & \dot{E}_q[\check{\psi}](\tau_2) + \text{Morr}[\check{\psi}](\tau_1, \tau_2) + \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} (\mathcal{E}_q + r^q \check{e}_4(\check{\psi})\check{N}) + \dot{F}_q[\check{\psi}](\tau_1, \tau_2) \\ & \lesssim E_q[\check{\psi}](\tau_1) + \check{J}_q[\check{\psi}, N](\tau_1, \tau_2) + R^{q+3} (E^1[\psi](\tau_2) + \text{Morr}^1[\psi](\tau_1, \tau_2)). \end{aligned} \quad (10.3.4)$$

Also, since $\delta \leq \max(q, \delta) \leq 1 - \delta$, we have in view of Theorem 5.3.4 in the case $s = 1^4$

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_{\max(q, \delta)}^1[\psi](\tau) + B_{\max(q, \delta)}^1[\psi](\tau_1, \tau_2) \\ & \lesssim E_{\max(q, \delta)}^1[\psi](\tau_1) + J_{\max(q, \delta)}^1[\psi, N](\tau_1, \tau_2), \end{aligned} \quad (10.3.5)$$

In view of (10.3.4) and (10.3.5), it thus only remains to estimate the integral

$$\int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} (\mathcal{E}_q + r^q \check{e}_4(\check{\psi})\check{N}),$$

i.e. we need to derive the analog of (10.2.17) used in the proof of Theorem 10.2.1. This is achieved in Proposition 10.3.2 below, which together with (10.3.4) and (10.3.5) immediately yields the proof of Theorem 5.3.5 in the case $s = 0$. \square

Proposition 10.3.2. *The following estimate holds true,*

$$\begin{aligned} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} (\mathcal{E}_q + r^q \check{e}_4(\check{\psi})\check{N}) & \geq \frac{1}{8} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-1} ((2+q)|\check{e}_4 \check{\psi}|^2 + (2-q)|\check{\nabla} \check{\psi}|^2 + 2r^{-2}|\check{\psi}|^2) \\ & \quad - O(\epsilon) \sup_{\tau_1 \leq \tau \leq \tau_2} \dot{E}_{q,R}[\check{\psi}](\tau) \\ & \quad - O(1) (E_{\max(q, \delta)}^1[\psi](\tau_1) + J_{\max(q, \delta)}^1[\psi, N]). \end{aligned} \quad (10.3.6)$$

We now focus on the proof of Proposition 10.3.2. In view of the definition of \check{N} , we have

¹⁴The proof of Theorem 5.3.4 for higher derivatives $s \geq 1$, even though proved later in section 10.4.5, is in fact independent of the proof of Theorem 5.3.5 and can thus be invoked here.

for $r \geq R$,

$$\begin{aligned}\check{N} &= A_0 + A_1 + A_2, \\ A_0 &= \frac{2}{r}e_4\check{\psi} = \frac{2}{r}(\check{e}_4\check{\psi} - r^{-1}\check{\psi}), \\ A_1 &= -\frac{6m}{r^2}e_4\check{\psi} + O(r^{-2})\mathfrak{d}^{\leq 1}\psi, \\ A_2 &= \text{Err}[\square_{\mathbf{g}}\check{\psi}], \\ \text{Err}[\square_{\mathbf{g}}\check{\psi}] &= r^2\Gamma_g e_4\mathfrak{d}\psi + r\mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 1}\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi.\end{aligned}$$

Also, recall that we have for $r \geq R$

$$\mathcal{E}_q(\check{\psi}) = \mathcal{E}[X_q, w_q, M_q] = \frac{q}{2}r^{q-1}|\check{e}_4(\check{\psi})|^2 + \frac{2-q}{2}r^{q-1}\mathcal{Q}_{34}(\check{\psi}) + \text{Err}_q(\check{\psi}).$$

Consequently, we write,

$$\begin{aligned}(\mathcal{E}_q + r^q\check{e}_4(\check{\psi})\check{N}) &= I_0 + I_1 + I_2, \\ I_0 &:= \frac{1}{2}r^{q-1}(q|\check{e}_4\check{\psi}|^2 + (2-q)|\check{\nabla}\check{\psi}|^2 + 4(2-q)r^{-2}\check{\psi}^2) + 2r^{q-1}\check{e}_4\check{\psi}(\check{e}_4\check{\psi} - r^{-1}\check{\psi}) \\ &= \frac{1}{2}r^{q-1}((q+4)|\check{e}_4\check{\psi}|^2 + (2-q)|\check{\nabla}\check{\psi}|^2 + 4(2-q)r^{-2}\check{\psi}^2 - 4r^{-1}\check{e}_4\check{\psi}\check{\psi}), \quad (10.3.7) \\ I_1 &:= r^{q-2}\check{e}_4(\check{\psi})[-6me_4\check{\psi} + O(1)\mathfrak{d}^{\leq 1}\psi] + O\left(\frac{m}{r}\right)r^{q-3}(\check{\psi})^2, \\ I_2 &:= \text{Err}_q(\check{\psi}) + r^q\check{e}_4(\check{\psi})A_2.\end{aligned}$$

We will rely on the following two lemmas.

Lemma 10.3.3. *The following lower bound estimate holds true for $q \leq 1 - \delta$ and $r \geq R$, where R is sufficiently large,*

$$I_0 + I_1 \geq \frac{1}{4}r^{q-1}((2+q)|\check{e}_4\check{\psi}|^2 + (2-q)|\check{\nabla}\check{\psi}|^2 + 2r^{-2}|\check{\psi}|^2) - O(1)r^{q-3}(\mathfrak{d}^{\leq 1}\psi)^2 \quad (10.3.8)$$

Lemma 10.3.4. *The following estimate holds true for the error term I_2*

$$\begin{aligned}\int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} |I_2| &\lesssim \epsilon \sup_{\tau_1 \leq \tau \leq \tau_2} \dot{E}_{q,R}[\check{\psi}](\tau) + \left(\frac{m_0}{R} + \epsilon\right) \dot{B}_{q,R}[\check{\psi}](\tau_1, \tau_2) \\ &\quad + \epsilon \left(\sup_{\tau_1 \leq \tau \leq \tau_2} E_q^1[\psi](\tau) + B_q^1[\psi](\tau_1, \tau_2) + J_q[\psi, N](\tau_1, \tau_2) \right).\end{aligned}$$

We postpone the proof of Lemma 10.3.3 and Lemma 10.3.4 to finish the proof of Proposition 10.3.2.

Proof of Proposition 10.3.2. In view of Lemma 10.3.3 and Lemma 10.3.4, we have

$$\begin{aligned}
 \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} (\mathcal{E}_q + r^q \check{e}_4(\check{\psi})\check{N}) &= \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} (I_0 + I_1) + \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} I_2 \\
 &\geq \frac{1}{4} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-1} ((2+q)|\check{e}_4 \check{\psi}|^2 + (2-q)|\nabla \check{\psi}|^2 + 2r^{-2}|\check{\psi}|^2) \\
 &\quad - O(1) \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-3}(\psi)^2 \\
 &\quad - O(\epsilon) \sup_{\tau_1 \leq \tau \leq \tau_2} \dot{E}_{q,R}[\check{\psi}](\tau) + O\left(\frac{m_0}{R} + \epsilon\right) \dot{B}_{q,R}[\check{\psi}](\tau_1, \tau_2) \\
 &\quad - O(\epsilon) \left(\sup_{\tau_1 \leq \tau \leq \tau_2} E_q^1[\psi](\tau) + B_q^1[\psi](\tau_1, \tau_2) + J_q[\psi, N](\tau_1, \tau_2) \right)
 \end{aligned}$$

so that, since $1 - \delta < q \leq 1 - \delta$, and for R sufficiently large and small ϵ ,

$$\begin{aligned}
 \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} (\mathcal{E}_q + r^q \check{e}_4(\check{\psi})\check{N}) &\geq \frac{1}{8} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-1} ((2+q)|\check{e}_4 \check{\psi}|^2 + (2-q)|\nabla \check{\psi}|^2 + 2r^{-2}|\check{\psi}|^2) \\
 &\quad - O(\epsilon) \sup_{\tau_1 \leq \tau \leq \tau_2} \dot{E}_{q,R}[\check{\psi}](\tau) \\
 &\quad - O(\epsilon) \left(\sup_{\tau_1 \leq \tau \leq \tau_2} E_q^1[\psi](\tau) + B_q^1[\psi](\tau_1, \tau_2) + J_q[\psi, N](\tau_1, \tau_2) \right).
 \end{aligned}$$

In view of (10.3.5), we infer

$$\begin{aligned}
 \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} (\mathcal{E}_q + r^q \check{e}_4(\check{\psi})\check{N}) &\geq \frac{1}{8} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-1} ((2+q)|\check{e}_4 \check{\psi}|^2 + (2-q)|\nabla \check{\psi}|^2 + 2r^{-2}|\check{\psi}|^2) \\
 &\quad - O(\epsilon) \sup_{\tau_1 \leq \tau \leq \tau_2} \dot{E}_{q,R}[\check{\psi}](\tau) \\
 &\quad - O(1) (E_{\max(q,\delta)}^1[\psi](\tau_1) + J_{\max(q,\delta)}^1[\psi, N])
 \end{aligned}$$

which concludes the proof. □

It finally remains to prove Lemma 10.3.3 and Lemma 10.3.4.

Proof of Lemma 10.3.3. Note that,

$$\begin{aligned}
 &(q+4)|\check{e}_4 \check{\psi}|^2 - 4r^{-1}(\check{e}_4 \check{\psi}) \check{\psi} + 4(2-q)r^{-2}|\check{\psi}|^2 \\
 = &(q+2)|\check{e}_4 \check{\psi}|^2 + (6-4q)r^{-2}|\check{\psi}|^2 + 2(\check{e}_4 \check{\psi} - r^{-1}\check{\psi})^2 \\
 \geq &(q+2)|\check{e}_4 \check{\psi}|^2 + (6-4q)r^{-2}|\check{\psi}|^2 \\
 \geq &(q+2)|\check{e}_4 \check{\psi}|^2 + 2r^{-2}|\check{\psi}|^2,
 \end{aligned}$$

where we used the fact that $q \leq 1 - \delta$. Hence,

$$I_0 \geq \frac{1}{2} r^{q-1} \left((2+q) |\check{e}_4 \check{\psi}|^2 + (2-q) |\check{\nabla} \check{\psi}|^2 + 2r^{-2} |\check{\psi}|^2 \right).$$

We also have,

$$I_1 \leq O\left(\frac{m}{r}\right) r^{q-1} (|\check{e}_4 \check{\psi}|^2 + r^{-2} |\check{\psi}|^2) + O(1) (r^{q-1} (\check{e}_4 \check{\psi})^2)^{\frac{1}{2}} (r^{q-3} (\mathfrak{d}^{\leq 1} \psi)^2)^{\frac{1}{2}}.$$

Thus if m_0/R is sufficiently small, and since $q \leq 1 - \delta$, we deduce, for $r \geq R$,

$$I_0 + I_1 \geq \frac{1}{4} r^{q-1} \left((2+q) |\check{e}_4 \check{\psi}|^2 + (2-q) |\check{\nabla} \check{\psi}|^2 + 2r^{-2} |\check{\psi}|^2 \right) - O(1) r^{q-3} (\mathfrak{d}^{\leq 1} \psi)^2$$

as desired. \square

Proof of Lemma 10.3.4. Recall that

$$\begin{aligned} I_2 &= \text{Err}_q(\check{\psi}) + r^q \check{e}_4(\check{\psi}) A_2, \\ A_2 &= \text{Err}[\square_{\mathbf{g}} \check{\psi}], \\ \text{Err}[\square_{\mathbf{g}} \check{\psi}] &= r \Gamma_b e_4 \mathfrak{d} \psi + \mathfrak{d}^{\leq 1}(\Gamma_b) \mathfrak{d}^{\leq 1} \psi + r \mathfrak{d}^{\leq 1}(\Gamma_g) e_3 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^2 \psi, \\ \text{Err}_q(\check{\psi}) &= O\left(\frac{m}{r^2}\right) r^q |e_4 \check{\psi}|^2 + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) r^q |\check{\psi}|^2 + O(\epsilon) w_{1,1} r^q (|e_4 \check{\psi}|^2 + |\check{\nabla} \check{\psi}|^2 + r^{-2} |\check{\psi}|^2) \\ &\quad + O(\epsilon) w_{2,1/2} r^q \left((|e_4 \check{\psi}| + r^{-1} |\check{\nabla} \check{\psi}|) |e_3 \check{\psi}| + |\check{\nabla} \check{\psi}|^2 + r^{-2} |\check{\psi}|^2 \right). \end{aligned}$$

Hence,

$$\begin{aligned} |I_2| &\lesssim \epsilon r^q |\check{e}_4(\check{\psi})| \left(\tau^{-1-\delta_{dec}} (|\check{e}_4 \mathfrak{d}^{\leq 1} \psi| + |\mathfrak{d}^{\leq 1} \psi|) + r^{-1} \tau^{-\frac{1}{2}-\delta_{dec}} (|e_3 \psi| + r^{-1} |\mathfrak{d}^2 \psi|) \right) \\ &\quad + \left(\frac{m}{r} + \epsilon \right) r^{q-1} (|\check{e}_4 \check{\psi}|^2 + |\check{\nabla} \check{\psi}|^2 + r^{-2} |\check{\psi}|^2) + \epsilon r^{q-2} \tau^{-\frac{1}{2}-\delta_{dec}} |e_4 \check{\psi}| |e_3 \check{\psi}| \\ &\quad + \epsilon r^{q-3} |\check{\nabla} \check{\psi}| |e_3 \check{\psi}|. \end{aligned}$$

This yields, using $q \leq 1 - \delta$,

$$\begin{aligned} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} |I_2| &\lesssim \epsilon \left(\sup_{\tau_1 \leq \tau \leq \tau_2} \dot{E}_{q,R}[\check{\psi}](\tau) \right)^{\frac{1}{2}} \left(\sup_{\tau_1 \leq \tau \leq \tau_2} E_q^1[\psi](\tau) + B_q^1[\psi](\tau_1, \tau_2) \right)^{\frac{1}{2}} \\ &\quad + \left(\frac{m_0}{R} + \epsilon \right) \dot{B}_{q,R}[\check{\psi}](\tau_1, \tau_2) \\ &\quad + \epsilon \left(\sup_{\tau_1 \leq \tau \leq \tau_2} \dot{E}_{q,R}[\check{\psi}](\tau) + \dot{B}_q[\check{\psi}](\tau_1, \tau_2) \right)^{\frac{1}{2}} \left(\int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-4} |e_3 \check{\psi}|^2 \right)^{\frac{1}{2}} \\ &\lesssim \epsilon \sup_{\tau_1 \leq \tau \leq \tau_2} \dot{E}_{q,R}[\check{\psi}](\tau) + \left(\frac{m_0}{R} + \epsilon \right) \dot{B}_{q,R}[\check{\psi}](\tau_1, \tau_2) \\ &\quad + \epsilon \left(\sup_{\tau_1 \leq \tau \leq \tau_2} E_q^1[\psi](\tau) + B_q^1[\psi](\tau_1, \tau_2) \right) + \epsilon \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-4} |e_3 \check{\psi}|^2. \end{aligned}$$

Next, we estimate the term involving $e_3 \check{\psi}$. For this we need to appeal to the formula in Lemma 5.3.7 which we recall below,

$$\square_2 \psi = -e_3 e_4 \psi + \mathbb{A}_2 \psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) e_4 \psi - \frac{1}{2}\kappa e_3 \psi + 2\eta e_\theta \psi$$

We have for $r \geq 6m_0$

$$\begin{aligned} e_3 \check{\psi} &= e_3(re_4(r\psi)) = re_3(re_4\psi) + e_3(r)e_4(r\psi) = r^2 e_3 e_4 \psi + 2re_3(r)e_4 \psi + e_3(r)e_4(r)\psi \\ &= r^2 \left(-\square_2 \psi + \mathbb{A}_2 \psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) e_4 \psi - \frac{1}{2}\kappa e_3 \psi + 2\eta e_\theta \psi \right) + 2re_3(r)e_4 \psi + e_3(r)e_4(r)\psi \end{aligned}$$

so that

$$|e_3 \check{\psi}| \lesssim r^2 |N| + r |e_3 \psi| + |\mathfrak{d}^{\leq 2} \psi|$$

and hence

$$\int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-4} |e_3 \check{\psi}|^2 \lesssim \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-4} \left(r^4 |N|^2 + r^2 |e_3 \psi|^2 + |\mathfrak{d}^{\leq 2} \psi|^2 \right).$$

Since $q \leq 1 - \delta$, we infer

$$\begin{aligned} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-4} |e_3 \check{\psi}|^2 &\lesssim \int_{(trq\bar{p})\mathcal{M}(\tau_1, \tau_2)} r^{1-\delta} |N|^2 + B_q^1[\psi](\tau_1, \tau_2) \\ &\lesssim J_q[\psi, N](\tau_1, \tau_2) + B_q^1[\psi](\tau_1, \tau_2) \end{aligned}$$

and thus

$$\begin{aligned} \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} |I_2| &\lesssim \epsilon \sup_{\tau_1 \leq \tau \leq \tau_2} \dot{E}_{q,R}[\check{\psi}](\tau) + \left(\frac{m_0}{R} + \epsilon\right) \dot{B}_{q,R}[\check{\psi}](\tau_1, \tau_2) \\ &\quad + \epsilon \left(\sup_{\tau_1 \leq \tau \leq \tau_2} E_q^1[\psi](\tau) + B_q^1[\psi](\tau_1, \tau_2) \right) + \epsilon \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^{q-4} |e_3 \check{\psi}|^2 \\ &\lesssim \epsilon \sup_{\tau_1 \leq \tau \leq \tau_2} \dot{E}_{q,R}[\check{\psi}](\tau) + \left(\frac{m_0}{R} + \epsilon\right) \dot{B}_{q,R}[\check{\psi}](\tau_1, \tau_2) \\ &\quad + \epsilon \left(\sup_{\tau_1 \leq \tau \leq \tau_2} E_q^1[\psi](\tau) + B_q^1[\psi](\tau_1, \tau_2) + J_q[\psi, N](\tau_1, \tau_2) \right). \end{aligned}$$

which concludes the proof of Lemma 10.3.4. \square

10.4 Higher Derivative Estimates

We have proved, respectively in section 10.2 and section 10.3.2, Theorem 5.3.4 on basic weighted estimates (see Remark 10.2.2) and Theorem 5.3.5 on higher weighted estimates only in the case $s = 0$. In this section, we conclude the proof of these theorems by recovering higher order derivatives¹⁵ one by one.

10.4.1 Basic assumptions

Recall that any Ricci coefficient either belongs to Γ_g or Γ_b , where Γ_g and Γ_b are defined in section 5.1.2. We make use of the following non sharp consequence of the estimates of Lemma 5.1.1. We assume, concerning the Ricci coefficients

$$\begin{aligned} |\mathfrak{d}^k(\Gamma_g)| &\lesssim \frac{\epsilon}{r^2 u_{trap}^{1+\delta_{dec}-2\delta_0}} && \text{for } k \leq k_{small} + 30, \\ |\mathfrak{d}^k(\Gamma_b)| &\lesssim \frac{\epsilon}{r u_{trap}^{1+\delta_{dec}-2\delta_0}} && \text{for } k \leq k_{small} + 30, \\ |\mathfrak{d}^k(\alpha, \beta, \check{\rho})| &\lesssim \frac{\epsilon}{r^3 u_{trap}^{1+\delta_{dec}-2\delta_0}} && \text{for } k \leq k_{small} + 30, \\ |\mathfrak{d}^k \underline{\alpha}| + r |\mathfrak{d}^k \underline{\beta}| &\lesssim \frac{\epsilon}{r u_{trap}^{1+\delta_{dec}-2\delta_0}} && \text{for } k \leq k_{small} + 30, \end{aligned}$$

where we recall that δ_{dec} and δ_0 are such that we have in particular $0 < 2\delta_0 < \delta_{dec}$.

10.4.2 Strategy for recovering higher order derivatives

So far, we have proved Theorem 5.3.4 in the case $s = 0$ ¹⁶ in section 10.2, and Theorem 5.3.5 on higher weighted estimates in the case $s = 0$ in section 10.3. We now conclude the proof of these theorems by recovering higher order derivatives one by one. Since going from $s = 0$ to $s = 1$ is analogous to going from s to $s + 1$, we will in fact consider only the former. More precisely, we assume the following bounds proved respectively in section 10.2 and section 10.3,

$$\sup_{\tau \in [\tau_1, \tau_2]} E_p[\psi](\tau) + B_p[\psi](\tau_1, \tau_2) + F_p[\psi](\tau_1, \tau_2) \lesssim E_p[\psi](\tau_1) + J_p[\psi, N](\tau_1, \tau_2), \quad (10.4.1)$$

¹⁵Respectively $s \leq k_{small} + 30$ in the case of Theorem 5.3.4, and $s \leq k_{small} + 29$ in the case of Theorem 5.3.5.

¹⁶Recall that Theorem 5.3.4 in the case $s = 0$ is obtained as a consequence of Theorem 10.1.1 on Morawetz and energy estimates, and Theorem 10.2.1 on r^p -weighted estimates, see Remark 10.2.2.

and

$$\begin{aligned} \sup_{\tau \in [\tau_1, \tau_2]} E_q[\check{\psi}](\tau) + B_q[\check{\psi}](\tau_1, \tau_2) &\lesssim E_q[\check{\psi}](\tau_1) + \check{J}_q[\check{\psi}, N](\tau_1, \tau_2) \\ &+ E^1_{\max(q, \delta)}[\psi](\tau_1) + J^1_{\max(q, \delta)}[\psi, N], \end{aligned} \tag{10.4.2}$$

and our goal is to prove the corresponding estimates for $s = 1$. We will proceed as follows

1. we first commute the wave equation for ψ and $\check{\psi}$ with T and derive (10.4.1) for $T\psi$ instead of ψ , and (10.4.2) for $T\check{\psi}$ instead of $\check{\psi}$,
2. we then commute the wave equation for ψ and $\check{\psi}$ with $r\check{\not{d}}_2$ and derive (10.4.1) for $r\check{\not{d}}_2\psi$ instead of ψ , and (10.4.2) for $r\check{\not{d}}_2\check{\psi}$ instead of $\check{\psi}$,
3. we then use the wave equation satisfied by ψ to derive an estimate for $R^2\psi$ in $r \leq 6m_0$ ¹⁷ with a degeneracy at $r = 3m$,
4. we then commute the wave equation for ψ with R and remove the degeneracy at $r = 3m$ for $R^2\psi$,
5. we then commute the wave equation for ψ with the redshift vectorfield $Y_{\mathcal{H}}$ and derive (10.4.1) for $Y_{\mathcal{H}}\psi$ instead of ψ ,
6. we then commute the wave equation for ψ and $\check{\psi}$ with f_1e_4 and derive (10.4.1) for $re_4\psi$ instead of ψ , and (10.4.2) for $f_1e_4\check{\psi}$ instead of $\check{\psi}$, where $f_1 = r$ for $r \geq 6m_0$ and $f_1 = 0$ for $r \leq 4m_0$,
7. we finally gather all estimates and conclude.

We will follow the above strategy in section 10.4.5 to prove Theorem 5.3.4, and in section 10.4.6 to prove Theorem 5.3.5. To this end, we first derive several commutator identities and estimates.

10.4.3 Commutation formulas with the wave equation

Commutation with T

Lemma 10.4.1. *We have, schematically, the following commutator formulae*

$$[T, e_4] = \Gamma_g \not{d}, \quad [T, e_3] = \Gamma_b \not{d}, \quad [T, \not{d}_k] = \Gamma_b \not{d} + \Gamma_b, \quad [T, \not{d}_k^*] = \Gamma_b \not{d} + \Gamma_b.$$

¹⁷Note that any finite region in r strictly containing the trapping region would suffice.

Proof. Recall that we have

$$[e_3, e_4] = 2\underline{\omega}e_4 - 2\underline{\omega}e_3 + 2(\underline{\eta} - \underline{\eta})e_\theta.$$

We infer

$$\begin{aligned} 2[T, e_4] &= [e_4 + \Upsilon e_3, e_4] \\ &= \Upsilon[e_3, e_4] - e_4(\Upsilon)e_3 \\ &= \Upsilon(2\underline{\omega}e_4 - 2\underline{\omega}e_3 + 2(\underline{\eta} - \underline{\eta})e_\theta) - e_4(\Upsilon)e_3 \\ &= \Upsilon\left(2\underline{\omega}e_4 - 2\left(\omega + \frac{1}{2}\Upsilon^{-1}e_4(\Upsilon)\right)e_3 + 2(\underline{\eta} - \underline{\eta})e_\theta\right) \\ &= (r^{-1}\Gamma_b + \Gamma_g)\mathfrak{d} \\ &= \Gamma_g\mathfrak{d}, \end{aligned}$$

and

$$\begin{aligned} 2[T, e_3] &= [e_4 + \Upsilon e_3, e_3] \\ &= [e_4, e_3] - e_3(\Upsilon)e_3 \\ &= -2\underline{\omega}e_4 + 2\underline{\omega}e_3 - 2(\underline{\eta} - \underline{\eta})e_\theta - e_3(\Upsilon)e_3 \\ &= -2\underline{\omega}e_4 + 2\left(\omega - \frac{1}{2}e_3(\Upsilon)\right)e_3 - 2(\underline{\eta} - \underline{\eta})e_\theta \\ &= -2\underline{\omega}e_4 + 2\left(\omega + \frac{1}{2}\Upsilon^{-1}e_4(\Upsilon) - \frac{1}{2\Upsilon}T(\Upsilon)\right)e_3 - 2(\underline{\eta} - \underline{\eta})e_\theta \\ &= (\Gamma_g + \Gamma_b)\mathfrak{d} \\ &= \Gamma_b\mathfrak{d}. \end{aligned}$$

Next, recall in view of Lemma 2.1.51, the following commutation formulae for reduced scalars.

1. If $f \in \mathfrak{s}_k$,

$$\begin{aligned} [\mathfrak{d}_k, e_3]f &= \frac{1}{2}\underline{\kappa}\mathfrak{d}_k f + \underline{Com}_k(f), \\ \underline{Com}_k(f) &= -\frac{1}{2}\underline{\vartheta}\mathfrak{d}_{k+1}^* f + (\zeta - \underline{\eta})e_3 f - k\underline{\eta}e_3\Phi f - \underline{\xi}(e_4 f + ke_4(\Phi)f) - k\underline{\beta}f, \\ [\mathfrak{d}_k, e_4] &= \frac{1}{2}\kappa\mathfrak{d}_k f + \text{Com}_k(f), \\ \text{Com}_k(f) &= -\frac{1}{2}\vartheta\mathfrak{d}_{k+1}^* f - (\zeta + \underline{\eta})e_4 f - k\underline{\eta}e_4\Phi f - \underline{\xi}(e_3 f + ke_3(\Phi)f) - k\beta f. \end{aligned}$$

2. If $f \in \mathfrak{s}_{k-1}$

$$\begin{aligned} [\mathfrak{d}_k^*, e_3]f &= \frac{1}{2}\underline{\kappa}\mathfrak{d}_k^*f + \underline{Com}_k^*(f), \\ \underline{Com}_k^*(f) &= -\frac{1}{2}\underline{\vartheta}\mathfrak{d}_{k-1}f - (\zeta - \eta)e_3f - (k-1)\eta e_3\Phi f + \underline{\xi}(e_4f - (k-1)e_4(\Phi)f) \\ &\quad - (k-1)\underline{\beta}f, \\ [\mathfrak{d}_k^*, e_4]f &= \frac{1}{2}\kappa\mathfrak{d}_k^*f + \text{Com}_k^*(f), \\ \text{Com}_k^*(f) &= -\frac{1}{2}\vartheta\mathfrak{d}_{k-1}f + (\zeta + \eta)e_4f - (k-1)\eta e_4\Phi f + \xi(e_3f - (k-1)e_3(\Phi)f) \\ &\quad - (k-1)\beta f. \end{aligned}$$

We infer, schematically,

$$\begin{aligned} 2[T, \mathfrak{d}_k] &= [e_4 + \Upsilon e_3, \mathfrak{d}_k] \\ &= [e_4, \mathfrak{d}_k] + \Upsilon[e_3, \mathfrak{d}_k] - e_\theta(\Upsilon)e_3 \\ &= -\frac{1}{2}(\kappa + \Upsilon\underline{\kappa})\mathfrak{d}_k + \Gamma_b\mathfrak{d} + r^{-1}\Gamma_b + 2e_\theta\left(\frac{m}{r}\right)e_3 \\ &= \Gamma_b\mathfrak{d} + \Gamma_b. \end{aligned}$$

The estimate for $[T, \mathfrak{d}_k^*]$ is similar and left to the reader. This concludes the proof of the lemma. \square

Lemma 10.4.2. *We have*

$$T(\kappa) = \mathfrak{d}^{\leq 1}\Gamma_g, \quad T\left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) = \mathfrak{d}^{\leq 1}\Gamma_b, \quad T(K) = \mathfrak{d}^{\leq 1}\Gamma_g.$$

Proof. We have

$$\begin{aligned} 2T(\kappa) &= (e_4 + \Upsilon e_3)\kappa \\ &= -\frac{1}{2}\kappa^2 - 2\omega\kappa + 2\mathfrak{d}_1\xi + 2(\underline{\eta} + \eta + 2\zeta)\xi - \frac{1}{2}\vartheta^2 \\ &\quad + \Upsilon\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa + 2\mathfrak{d}_1\eta + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2(\xi\underline{\xi} + \eta^2)\right) \\ &= -\frac{1}{2}\kappa^2 - 2\omega\kappa - \frac{1}{2}\underline{\kappa}\kappa\Upsilon + 2\underline{\omega}\kappa\Upsilon + 2\Upsilon\rho \\ &\quad + 2\mathfrak{d}_1\xi + 2\Upsilon\mathfrak{d}_1\eta + 2(\underline{\eta} + \eta + 2\zeta)\xi - \frac{1}{2}\vartheta^2 + \Upsilon\left(-\frac{1}{2}\vartheta\underline{\vartheta} + 2(\xi\underline{\xi} + \eta^2)\right) \\ &= r^{-1}\mathfrak{d}\Gamma_b + r^{-1}\Gamma_b \\ &= \mathfrak{d}^{\leq 1}\Gamma_g. \end{aligned}$$

Also, we have

$$T(r) = e_4(r) + \Upsilon e_3(r) = e_4(r) - \Upsilon + \Upsilon(e_3(r) - 1) \in r\Gamma_b.$$

We infer

$$\begin{aligned} T\left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) &= T\left(\frac{1}{r}\right) + \Gamma_b - \frac{1}{2}T\left(\frac{\underline{\kappa}}{r} + \frac{2}{r}\right) \\ &= -\frac{T(r)}{r^2} + \mathfrak{d}\Gamma_b \\ &= \mathfrak{d}^{\leq 1}\Gamma_b \end{aligned}$$

and

$$\begin{aligned} T(K) &= T\left(\frac{1}{r^2}\right) + T\left(K - \frac{1}{r^2}\right) \\ &= -\frac{2T(r)}{r^3} + r^{-1}\Gamma_b \\ &= r^{-1}\mathfrak{d}(\Gamma_b) + r^{-1}\Gamma_b \\ &= \mathfrak{d}^{\leq 1}\Gamma_g. \end{aligned}$$

This concludes the proof of the lemma. □

Corollary 10.4.3. *We have*

$$[T, \square_2]\psi = \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi.$$

Proof. Recall that we have

$$\square_2\psi = -e_3e_4\psi + \mathbb{A}_2\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi - \frac{1}{2}\kappa e_3\psi + 2\eta e_\theta\psi.$$

We infer

$$\begin{aligned} [T, \square_2]\psi &= -[T, e_3]e_4\psi - e_3[T, e_4]\psi + [T, \mathbb{A}_2]\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)[T, e_4]\psi \\ &\quad + T\left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi - \frac{1}{2}\kappa[T, e_3]\psi - \frac{1}{2}T(\kappa)e_3\psi + 2\eta[T, e_\theta]\psi + 2T(\eta)e_\theta\psi. \end{aligned}$$

and hence, using also $\mathbb{A}_2 = -\mathfrak{d}_2^* \mathfrak{d}_2 + 2K$,

$$\begin{aligned} [T, \square_2]\psi &= -[T, e_3](r^{-1}\mathfrak{d}\psi) - \mathfrak{d}[T, e_4]\psi - r^{-1}\mathfrak{d}[T, \mathfrak{d}_2]\psi - [T, \mathfrak{d}_2^*]r^{-1}\mathfrak{d}\psi + 2T(K)\psi \\ &\quad + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)[T, e_4]\psi + T\left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)r^{-1}\mathfrak{d}\psi - \frac{1}{2}\kappa[T, e_3]\psi - \frac{1}{2}T(\kappa)\mathfrak{d}\psi \\ &\quad + 2\eta[T, e_\theta]\psi + 2T(\eta)r^{-1}\mathfrak{d}\psi. \end{aligned}$$

In view of

$$[T, e_4] = \Gamma_g \mathfrak{d}, \quad [T, e_3] = \Gamma_b \mathfrak{d}, \quad [T, \phi_k] = \Gamma_b \mathfrak{d} + \Gamma_b, \quad [T, \phi_k^*] = \Gamma_b \mathfrak{d} + \Gamma_b$$

and

$$T(\kappa) = \mathfrak{d}^{\leq 1} \Gamma_g, \quad T\left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) = \mathfrak{d}^{\leq 1} \Gamma_b, \quad T(K) = \mathfrak{d}^{\leq 1} \Gamma_g,$$

we deduce, schematically,

$$\begin{aligned} [T, \square_2] \psi &= \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \psi + r^{-1} \mathfrak{d}^{\leq 1}(\Gamma_b) \mathfrak{d}^{\leq 2} \psi \\ &= \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \psi. \end{aligned}$$

This concludes the proof of the corollary. □

Commutation with angular derivatives

Lemma 10.4.4. *We have, schematically,*

$$\begin{aligned} [r \phi_k, e_4] f, [r \phi_k^*, e_4] f &= \Gamma_g \mathfrak{d}^{\leq 1} f, \quad [r^2 \underline{\Delta}_k, e_4] f = \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} f \\ [r \phi_k, e_3] f &= -r \eta e_3(f) + \Gamma_b \mathfrak{d}^{\leq 1} f, \quad [r \phi_k^*, e_3] f = r \eta e_3(f) + \Gamma_b \mathfrak{d}^{\leq 1} f. \end{aligned}$$

Proof. Recall from Lemma 2.2.13 that the following commutation formulae holds true,

1. If $f \in \mathfrak{s}_k$,

$$\begin{aligned} [r \phi_k, e_4] &= r \left[\text{Com}_k(f) - \frac{1}{2} A \phi_k f \right], \\ [r \phi_k, e_3] f &= r \left[\underline{\text{Com}}_k(f) - \frac{1}{2} \underline{A} \phi_k f \right]. \end{aligned}$$

2. If $f \in \mathfrak{s}_{k-1}$

$$\begin{aligned} [r \phi_k^*, e_4] f &= r \left[\text{Com}_k^*(f) - \frac{1}{2} A \phi_k^* f \right], \\ [r \phi_k^*, e_3] f &= r \left[\underline{\text{Com}}_k^*(f) - \frac{1}{2} \underline{A} \phi_k^* f \right], \end{aligned}$$

where $A = 2/re_4(r) - \kappa$ and $\underline{A} = 2/re_3(r) - \underline{\kappa}$. Now, we have

$$\begin{aligned} \text{Com}_k(f) &= r^{-1}\Gamma_g \mathfrak{d}^{\leq 1} f, & \text{Com}_k^*(f) &= r^{-1}\Gamma_g \mathfrak{d}^{\leq 1} f, \\ \underline{\text{Com}}_k(f) &= -\eta e_3(f) + r^{-1}\Gamma_b \mathfrak{d}^{\leq 1} f, & \underline{\text{Com}}_k^*(f) &= \eta e_3(f) + r^{-1}\Gamma_b \mathfrak{d}^{\leq 1} f, \end{aligned}$$

which together with the fact that $A \in \Gamma_g$ and $\underline{A} \in \Gamma_b$ implies, schematically,

$$\begin{aligned} [r \not{d}_k, e_4]f, [r \not{d}_k^*, e_4]f &= \Gamma_g \mathfrak{d}^{\leq 1} f, \\ [r \not{d}_k, e_3]f &= -r\eta e_3(f) + \Gamma_b \mathfrak{d}^{\leq 1} f, \quad [r \not{d}_k^*, e_3]f = r\eta e_3(f) + \Gamma_b \mathfrak{d}^{\leq 1} f. \end{aligned}$$

Since $\not{d}_k = -\not{d}_k^* \not{d}_k + kK$, We infer

$$\begin{aligned} [r^2 \not{d}_k, e_4] &= [-r^2 \not{d}_k^* \not{d}_k + kr^2 K, e_4] \\ &= -[r \not{d}_k^*, e_4]r \not{d}_2 - r \not{d}_k^* [r \not{d}_k, e_4] + \end{aligned}$$

This concludes the proof of the lemma. □

Corollary 10.4.5. *We have*

$$r \not{d}_2(\square_2 \psi) - (\square_1 - K)(r \not{d}_2 \psi) = -r\eta \square_2 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \psi$$

and

$$r \not{d}_2^*(\square_1 \phi) - (\square_2 - 3K)(r \not{d}_2^* \phi) = r\eta \square_1 \phi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \phi.$$

Proof. Recall that we have

$$\square_2 \psi = -e_3 e_4 \psi + \not{d}_2 \psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) e_4 \psi - \frac{1}{2}\kappa e_3 \psi + 2\eta e_\theta \psi$$

and

$$\square_1 \phi = -e_3 e_4 \phi + \not{d}_2 \phi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) e_4 \phi - \frac{1}{2}\kappa e_3 \phi + 2\eta e_\theta \phi$$

We infer

$$\begin{aligned} r \not{d}_2(\square_2 \psi) - \square_1(r \not{d}_2 \psi) &= -[r \not{d}_2, e_3]e_4 \psi - e_3[r \not{d}_2, e_4]\psi + r(\not{d}_2 \not{d}_2 - \not{d}_1 \not{d}_2)\psi + [r, \not{d}_1] \not{d}_2 \psi \\ &\quad + r e_\theta \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) e_4 \psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) [r \not{d}_2, e_4]\psi - \frac{1}{2}r e_\theta(\kappa) e_3 \psi \\ &\quad - \frac{1}{2}\kappa [r \not{d}_2, e_3]\psi + 2r \not{d}_2(\eta e_\theta \psi) - 2\eta e_\theta(r \not{d}_2 \psi), \end{aligned}$$

and

$$\begin{aligned} r \not{d}_2^*(\square_1 \phi) - \square_2(r \not{d}_2^* \phi) &= -[r \not{d}_2^*, e_3]e_4 \phi - e_3[r \not{d}_2^*, e_4]\phi + r(\not{d}_2^* \not{A}_1 - \not{A}_2 \not{d}_2^*)\phi + [r, \not{A}_2] \not{d}_2^* \phi \\ &\quad - re_\theta \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) e_4 \phi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) [r \not{d}_2^*, e_4]\phi + \frac{1}{2}re_\theta(\kappa)e_3 \phi \\ &\quad - \frac{1}{2}\kappa[r \not{d}_2^*, e_3]\phi + 2r \not{d}_2(\eta e_\theta \phi) - 2\eta e_\theta(r \not{d}_2 \phi), \end{aligned}$$

and hence, using also in particular the following identities from Proposition 2.1.25

$$\begin{aligned} \not{d}_2 \not{A}_2 - \not{A}_1 \not{d}_2 &= -K \not{d}_2 + 2e_\theta(K), \\ \not{d}_2^* \not{A}_1 - \not{A}_2 \not{d}_2^* &= -3K \not{d}_2^* - e_\theta(K), \end{aligned}$$

we infer,

$$\begin{aligned} r \not{d}_2(\square_2 \psi) - (\square_1 - K)(r \not{d}_2 \psi) &= -[r \not{d}_2, e_3]e_4 \psi - \not{d}[r \not{d}_2, e_4]\psi + 2re_\theta(K)\psi + [r, \not{A}_1](r^{-1} \not{d}\psi) \\ &\quad + re_\theta \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) r^{-1} \not{d}\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) [r \not{d}_2, e_4]\psi - \frac{1}{2}re_\theta(\kappa) \not{d}\psi \\ &\quad - \frac{1}{2}\kappa[r \not{d}_2, e_3]\psi + 2\not{d}(r^{-1} \eta \not{d}\psi) - 2r^{-1} \eta \not{d}(\not{d}\psi), \end{aligned}$$

and

$$\begin{aligned} r \not{d}_2^*(\square_1 \phi) - (\square_2 - 3K)(r \not{d}_2^* \phi) &= -[r \not{d}_2^*, e_3]e_4 \phi - \not{d}[r \not{d}_2^*, e_4]\phi - re_\theta(K)\phi + [r, \not{A}_2](r^{-1} \not{d}\phi) \\ &\quad - re_\theta \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) r^{-1} \not{d}\phi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) [r \not{d}_2^*, e_4]\phi + \frac{1}{2}re_\theta(\kappa) \not{d}\phi \\ &\quad - \frac{1}{2}\kappa[r \not{d}_2^*, e_3]\phi + 2\not{d}(r^{-1} \eta \not{d}\phi) - 2r^{-1} \eta \not{d}(\not{d}\phi). \end{aligned}$$

This yields, schematically,

$$\begin{aligned} &r \not{d}_2(\square_2 \psi) - (\square_1 - K)(r \not{d}_2 \psi) \\ &= -[r \not{d}_2, e_3]e_4 \psi - \not{d}[r \not{d}_2, e_4]\psi + r^{-1}[r \not{d}_2, e_4]\psi - \frac{1}{2}\kappa[r \not{d}_2, e_3]\psi + \not{d}^{\leq 1}(\Gamma_g) \not{d}^{\leq 2} \psi \end{aligned}$$

and

$$\begin{aligned} &r \not{d}_2^*(\square_1 \phi) - (\square_2 - 3K)(r \not{d}_2^* \phi) \\ &= -[r \not{d}_2^*, e_3]e_4 \phi - \not{d}[r \not{d}_2^*, e_4]\phi + r^{-1}[r \not{d}_2^*, e_4]\phi - \frac{1}{2}\kappa[r \not{d}_2^*, e_3]\phi + \not{d}^{\leq 1}(\Gamma_g) \not{d}^{\leq 2} \phi \end{aligned}$$

where we used the fact that $r^{-1} \not{d}^{\leq 1} \Gamma_b$ is at least as good as $\not{d}^{\leq 1} \Gamma_g$ and the fact that $r^{-1} e_\theta(r)$ is Γ_g .

Next, we rely on

$$\begin{aligned} [r \not{d}_k, e_4]f, [r \not{d}_k^*, e_4]f &= \Gamma_g \mathfrak{d}^{\leq 1} f, \\ [r \not{d}_k, e_3]f &= -r\eta e_3(f) + \Gamma_b \mathfrak{d}^{\leq 1} f, [r \not{d}_k^*, e_3]f = r\eta e_3(f) + \Gamma_b \mathfrak{d}^{\leq 1} f \end{aligned}$$

to infer

$$r \not{d}_2(\square_2 \psi) - (\square_1 - K)(r \not{d}_2 \psi) = r\eta e_3 e_4 \psi + \frac{1}{2} r \kappa \eta e_3 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \psi$$

and

$$r \not{d}_2^*(\square_1 \phi) - (\square_2 - 3K)(r \not{d}_2^* \phi) = -r\eta e_3 e_4 \phi - \frac{1}{2} r \kappa \eta e_3 \phi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \phi.$$

This yields

$$\begin{aligned} r \not{d}_2(\square_2 \psi) - (\square_1 - K)(r \not{d}_2 \psi) &= r\eta \left(-\square_2 \psi + \not{d}_2 \psi + \left(2\underline{\omega} - \frac{1}{2} \underline{\kappa} \right) e_4 \psi + 2\eta e_\theta \psi \right) \\ &\quad + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \psi \\ &= -r\eta \square_2 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \psi \end{aligned}$$

and

$$\begin{aligned} r \not{d}_2^*(\square_1 \phi) - (\square_2 - 3K)(r \not{d}_2^* \phi) &= -r\eta \left(-\square_1 \phi + \not{d}_2 \phi + \left(2\underline{\omega} - \frac{1}{2} \underline{\kappa} \right) e_4 \phi + 2\eta e_\theta \phi \right) \\ &\quad + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \phi \\ &= r\eta \square_1 \phi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \phi. \end{aligned}$$

where we used the fact that $r - 1\mathfrak{d}^{\leq 1}\Gamma_b$ is at least as good as $\mathfrak{d}^{\leq 1}\Gamma_g$. This concludes the proof of the corollary. \square

Commutation with R in the region $r \leq r_0$

We derive in the following lemma commutator identities that are non sharp as far as decay in r is concerned. This is sufficient for our needs since we will commute the wave equation with R only in the region $r \leq r_0$ for a fixed $r_0 \geq 4m_0$ large enough. We will use in particular the following estimate Also, recall that

$$\max_{k \leq k_{small} + 30} |\mathfrak{d}(\Gamma_g)| \lesssim \frac{\epsilon}{r^2 u_{trap}^{1 + \delta_{dec} - 2\delta_0}}. \quad (10.4.3)$$

Lemma 10.4.6. *We have*

$$\begin{aligned} [R, e_4] &= O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d}, & [R, e_3] &= -\frac{2m}{r^2}e_3 + O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d}, \\ [r \not\Delta_k, R]f, [r \not\Delta_k^*, R]f &= O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d}^{\leq 1}f, & [r^2 \not\Delta_k, R]f &= O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d}^{\leq 2}f. \end{aligned}$$

Proof. Recall that R is defined by

$$R = \frac{1}{2}(e_4 - \Upsilon e_3)$$

and that we have

$$[e_3, e_4] = 2\underline{\omega}e_4 - 2\underline{\omega}e_3 + 2(\eta - \underline{\eta})e_\theta.$$

We infer

$$[R, e_4] = \frac{1}{2}[-\Upsilon e_3, e_4] = -\frac{\Upsilon}{2}[e_3, e_4] + \frac{1}{2}e_4(\Upsilon)e_3 = \left(\Upsilon\underline{\omega}e_3 + \frac{m}{r^2}e_4(r)\right)e_3 + O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d},$$

and

$$[R, e_3] = \frac{1}{2}[e_4 - \Upsilon e_3, e_3] = \frac{1}{2}[e_4, e_3] + \frac{1}{2}e_3(\Upsilon)e_3 = \left(\underline{\omega}e_3 + \frac{m}{r^2}e_3(r)\right)e_3 + O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d},$$

and hence,

$$[R, e_4] = O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d}, \quad [R, e_3] = -\frac{2m}{r^2}e_3 + O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d}.$$

Also, recall that we have

$$\begin{aligned} [r \not\Delta_k, e_4]f, [r \not\Delta_k^*, e_4]f &= \Gamma_g \mathfrak{d}^{\leq 1}f, & [r^2 \not\Delta_k, e_4]f &= \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2}f \\ [r \not\Delta_k, e_3]f &= -r\underline{\eta}e_3(f) + \Gamma_b \mathfrak{d}^{\leq 1}f, & [r \not\Delta_k^*, e_3]f &= r\underline{\eta}e_3(f) + \Gamma_b \mathfrak{d}^{\leq 1}f. \end{aligned}$$

We infer

$$\begin{aligned} [r \not\Delta_k, e_4]f, [r \not\Delta_k^*, e_4]f &= O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d}^{\leq 1}f, & [r^2 \not\Delta_k, e_4]f &= O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d}^{\leq 2}f \\ [r \not\Delta_k, e_3]f &= O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d}^{\leq 1}f, & [r \not\Delta_k^*, e_3]f &= O\left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}}\right) \mathfrak{d}^{\leq 1}f. \end{aligned}$$

Together with the definition for R , we deduce

$$[r \not{d}_k, R]f, [r \not{d}_k^*, R]f = O\left(\frac{\epsilon}{u_{trap}^{1+\delta_{dec}-2\delta_0}}\right) \mathfrak{d}^{\leq 1} f, [r^2 \not{d}_k, R]f = O\left(\frac{\epsilon}{u_{trap}^{1+\delta_{dec}-2\delta_0}}\right) \mathfrak{d}^{\leq 2} f.$$

This concludes the proof of the lemma. \square

Corollary 10.4.7. *We have in the region $r \leq r_0$*

$$\square_2(R\psi) = \left(1 - \frac{3m}{r}\right) \mathfrak{d}^2 \psi + O\left(\frac{\epsilon}{u_{trap}^{1+\delta_{dec}-2\delta_0}}\right) \mathfrak{d}^2 \psi + O(1) \mathfrak{d}^{\leq 1} \psi + O(1) \mathfrak{d}^{\leq 1} N.$$

Proof. Recall that we have

$$\square_2 \psi = -e_4(e_3(\psi)) + \not{d}_2 \psi - \frac{1}{2} \underline{\kappa} e_4 \psi + \left(-\frac{1}{2} \kappa + 2\omega\right) e_3 \psi + 2\underline{\eta} e_\theta \psi.$$

Multiplying by r^2 , we infer

$$r^2 \square_2 \psi = -r^2 e_4(e_3(\psi)) + r^2 \not{d}_2 \psi - \frac{1}{2} r^2 \underline{\kappa} e_4 \psi + r^2 \left(-\frac{1}{2} \kappa + 2\omega\right) e_3 \psi + 2r \underline{\eta} e_\theta \psi$$

and hence

$$\begin{aligned} R(r^2 \square_2 \psi) &= r^2 \square_2(R\psi) - [R, r^2 e_4 e_3] \psi + [R, r^2 \not{d}_2] \psi - \frac{1}{2} R(r^2 \underline{\kappa}) e_4 \psi - \frac{1}{2} r^2 \underline{\kappa} [R, e_4] \psi \\ &\quad + R\left(r^2 \left(-\frac{1}{2} \kappa + 2\omega\right)\right) e_3 \psi + r^2 \left(-\frac{1}{2} \kappa + 2\omega\right) [R, e_3] \psi + 2R(r \underline{\eta}) r e_\theta \psi \\ &\quad + 2r \underline{\eta} [R, r e_\theta] \psi. \end{aligned}$$

Using the commutation identities of the previous lemma, we infer in the region $r \leq r_0$

$$R(r^2 \square_2 \psi) = r^2 \square_2(R\psi) - [R, r^2 e_4 e_3] \psi + O\left(\frac{\epsilon}{u_{trap}^{1+\delta_{dec}-2\delta_0}}\right) \mathfrak{d}^2 \psi + O(1) \mathfrak{d} \psi.$$

Also, since ψ satisfies $\square_2 \psi = V\psi + N$, we infer in the region $r \leq r_0$

$$r^2 \square_2(R\psi) = [R, r^2 e_4 e_3] \psi + O\left(\frac{\epsilon}{u_{trap}^{1+\delta_{dec}-2\delta_0}}\right) \mathfrak{d}^2 \psi + O(1) \mathfrak{d}^{\leq 1} \psi + O(1) \mathfrak{d}^{\leq 1} N.$$

Next, recall that we have

$$[R, e_4] = O\left(\frac{\epsilon}{u_{trap}^{1+\delta_{dec}-2\delta_0}}\right) \mathfrak{d}, \quad [R, e_3] = -\frac{2m}{r^2} e_3 + O(\epsilon) \mathfrak{d}.$$

We infer

$$\begin{aligned}
 [R, r^2 e_4 e_3] \psi &= R(r^2) e_4 e_3 + r^2 [R, e_4] e_3 + r^2 e_4 [R, e_3] \\
 &= \frac{1}{2} \left(e_4(r^2) - \Upsilon e_3(r^2) \right) e_4 e_3 \psi + r^2 e_4 \left(-\frac{2m}{r^2} e_3 \psi \right) + O \left(\frac{\epsilon}{u_{trap}^{1+\delta_{dec}-2\delta_0}} \right) \mathfrak{d}^2 \psi \\
 &= 2(r-3m) e_4 e_3 \psi + O \left(\frac{\epsilon}{u_{trap}^{1+\delta_{dec}-2\delta_0}} \right) \mathfrak{d}^2 \psi + O(1) \mathfrak{d} \psi
 \end{aligned}$$

and thus, in the region $r \leq r_0$,

$$\square_2(R\psi) = \left(1 - \frac{3m}{r} \right) \mathfrak{d}^2 \psi + O \left(\frac{\epsilon}{u_{trap}^{1+\delta_{dec}-2\delta_0}} \right) \mathfrak{d}^2 \psi + O(1) \mathfrak{d}^{\leq 1} \psi + O(1) \mathfrak{d}^{\leq 1} N$$

as desired. □

Commutation with the redshift vectorfield

Let a positive bump function $\kappa = \kappa(r)$, supported in the region in $[-2, 2]$ and equal to 1 for $[-1, 1]$. Recall that the redshift vectorfield is given by

$$Y_{\mathcal{H}} = \kappa_{\mathcal{H}} Y_{(0)}, \quad \kappa_{\mathcal{H}} := \kappa \left(\frac{\Upsilon}{\delta_{\mathcal{H}}} \right)$$

where $Y_{(0)}$ is defined by

$$Y_{(0)} = a e_3 + b e_4 + 2T, \quad a = 1 + \frac{5}{4m}(r-2m), \quad b = \frac{5}{4m}(r-2m).$$

Lemma 10.4.8. *We have*

$$[\square_2, e_3] \psi = -2\omega e_3(e_3 \psi) + \underline{\kappa} e_4(e_3 \psi) + \underline{\kappa} \square_2 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^2 \psi + r^{-2} \mathfrak{d}^{\leq 1} \psi.$$

Proof. Recall that we have

$$\square_2 \psi = -e_4(e_3(\psi)) + \underline{\Delta}_2 \psi - \frac{1}{2} \underline{\kappa} e_4 \psi + \left(-\frac{1}{2} \kappa + 2\omega \right) e_3 \psi + 2\underline{\eta} e_{\theta} \psi.$$

Since we have

$$\begin{aligned}
 [e_4, e_3] &= 2\omega e_3 + r^{-1} \Gamma_b \not\partial, & [\not\partial_k, e_3] &= \frac{1}{2} \underline{\kappa} \not\partial_k + \Gamma_b \not\partial + r^{-1} \Gamma_b, \\
 [\not\partial_k^*, e_3] &= \frac{1}{2} \underline{\kappa} \not\partial_k^* + \Gamma_b \not\partial + r^{-1} \Gamma_b
 \end{aligned}$$

We infer

$$\begin{aligned}
[\square_2, e_3]\psi &= -[e_4, e_3](e_3(\psi)) + [\underline{\Delta}_2, e_3] - \frac{1}{2}\underline{\kappa}[e_4, e_3](\psi) \\
&\quad + \frac{1}{2}e_3(\underline{\kappa})e_4(\psi) - e_3\left(-\frac{1}{2}\kappa + 2\omega\right)e_3(\psi) + 2\underline{\eta}[e_\theta, e_3]\psi - 2e_3(\underline{\eta})e_\theta(\psi) \\
&= -2\omega e_3(e_3\psi) + \underline{\kappa}\underline{\Delta}_2\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}^{\leq 1}\psi.
\end{aligned}$$

Using again

$$\square_2\psi = -e_4(e_3(\psi)) + \underline{\Delta}_2\psi - \frac{1}{2}\underline{\kappa}e_4\psi + \left(-\frac{1}{2}\kappa + 2\omega\right)e_3\psi + 2\underline{\eta}e_\theta\psi,$$

we deduce

$$\begin{aligned}
[\square_2, e_3]\psi &= -2\omega e_3(e_3\psi) + \underline{\kappa}\left(\square_2\psi + e_4(e_3\psi)\right) + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}^{\leq 1}\psi \\
&= -2\omega e_3(e_3\psi) + \underline{\kappa}e_4(e_3\psi) + \underline{\kappa}\square_2\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}^{\leq 1}\psi.
\end{aligned}$$

This concludes the proof of the lemma. \square

Lemma 10.4.9. *The exists a scalar function d_0 satisfying the bound*

$$d_0 = \frac{1}{2m_0} + O(\delta_{\mathcal{H}}) \text{ on the support of } \kappa_{\mathcal{H}},$$

such that we have, schematically

$$\begin{aligned}
[\square_2, Y_{\mathcal{H}}]\psi &= d_0 Y_{(0)}(Y_{\mathcal{H}}\psi) + 1_{\Upsilon \leq 2\delta_{\mathcal{H}}} \left(\square_2\psi + \mathfrak{d}T\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi + \frac{1}{\delta_{\mathcal{H}}^2}\mathfrak{d}^{\leq 1}\psi \right) \\
&\quad + \frac{1}{\delta_{\mathcal{H}}} 1_{\delta_{\mathcal{H}} \leq \Upsilon \leq 2\delta_{\mathcal{H}}} \mathfrak{d}^{\leq 2}\psi.
\end{aligned}$$

Proof. We have

$$\begin{aligned}
Y_{(0)} = ae_3 + be_4 + 2T &= ae_3 + b(2T - \Upsilon e_3) + 2T \\
&= (a - \Upsilon b)e_3 + 2(1 + b)T.
\end{aligned}$$

Thus, in view of the commutator identities

$$\begin{aligned}
[T, \square_2]\psi &= \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi, \\
[\square_2, e_3]\psi &= -2\omega e_3(e_3\psi) + \underline{\kappa}e_4(e_3\psi) + \underline{\kappa}\square_2\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}^{\leq 1}\psi,
\end{aligned}$$

we deduce, schematically,

$$\begin{aligned}
[\square_2, Y_{(0)}]\psi &= [\square_2, (a - \Upsilon b)e_3]\psi + [\square_2, 2(1 + b)T]\psi \\
&= (a - \Upsilon b)[\square_2, e_3]\psi + \mathbf{g}^{\alpha\beta}\mathbf{D}_\alpha(a)\mathbf{D}_\beta e_3\psi + 2(1 + b)[\square_2, T]\psi \\
&\quad + 2\mathbf{g}^{\alpha\beta}\mathbf{D}_\alpha(b)\mathbf{D}_\beta T\psi + \mathfrak{d}^{\leq 1}\psi \\
&= (a - \Upsilon b)\left(-2\omega e_3(e_3\psi) + \underline{\kappa}e_4(e_3\psi) + \underline{\kappa}\square_2\psi\right) - \frac{1}{2}e_3(a)e_4(e_3\psi) - \frac{1}{2}e_4(a)e_3(e_3\psi) \\
&\quad + \mathfrak{d}T\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi + \mathfrak{d}^{\leq 1}\psi.
\end{aligned}$$

Since $e_4 = -\Upsilon e_3 + 2T$, we infer schematically

$$\begin{aligned}
[\square_2, Y_{(0)}]\psi &= \left((a - \Upsilon b)(-2\omega - \Upsilon\underline{\kappa}) + \frac{\Upsilon}{2}e_3(a) - \frac{1}{2}e_4(a)\right) e_3(e_3\psi) \\
&\quad + \square_2\psi + \mathfrak{d}T\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi + \mathfrak{d}^{\leq 1}\psi.
\end{aligned}$$

We deduce,

$$\begin{aligned}
[\square_2, Y_{\mathcal{H}}]\psi &= [\square_2, \kappa_{\mathcal{H}}Y_{(0)}]\psi \\
&= \kappa_{\mathcal{H}}[\square_2, Y_{(0)}]\psi + \kappa'_{\mathcal{H}}\mathfrak{d}^{\leq 2}\psi + \kappa''_{\mathcal{H}}\mathfrak{d}^{\leq 1}\psi \\
&= \kappa_{\mathcal{H}}\left((a - \Upsilon b)(-2\omega - \Upsilon\underline{\kappa}) + \frac{\Upsilon}{2}e_3(a) - \frac{1}{2}e_4(a)\right) e_3(e_3\psi) \\
&\quad + 1_{\Upsilon \leq 2\delta_{\mathcal{H}}}\left(\square_2\psi + \mathfrak{d}T\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi + \frac{1}{\delta_{\mathcal{H}}^2}\mathfrak{d}^{\leq 1}\psi\right) + \frac{1}{\delta_{\mathcal{H}}}1_{\delta_{\mathcal{H}} \leq \Upsilon \leq 2\delta_{\mathcal{H}}}\mathfrak{d}^{\leq 2}\psi.
\end{aligned}$$

Now, we have

$$\begin{aligned}
\kappa_{\mathcal{H}}e_3(e_3\psi) &= \frac{1}{a - \Upsilon b}\kappa_{\mathcal{H}}Y_{(0)}(e_3\psi) + T\mathfrak{d}\psi \\
&= \frac{1}{(a - \Upsilon b)^2}\kappa_{\mathcal{H}}Y_{(0)}(Y_{(0)}\psi) + \mathfrak{d}T\psi + \mathfrak{d}^{\leq 1}\psi \\
&= \frac{1}{(a - \Upsilon b)^2}Y_{(0)}(Y_{\mathcal{H}}\psi) + \mathfrak{d}T\psi + \frac{1}{\delta_{\mathcal{H}}}\mathfrak{d}^{\leq 1}\psi
\end{aligned}$$

and hence

$$\begin{aligned}
[\square_2, Y_{\mathcal{H}}]\psi &= \frac{(a - \Upsilon b)(-2\omega - \Upsilon\underline{\kappa}) + \frac{\Upsilon}{2}e_3(a) - \frac{1}{2}e_4(a)}{(a - \Upsilon b)^2}Y_{(0)}(Y_{\mathcal{H}}\psi) \\
&\quad + 1_{\Upsilon \leq 2\delta_{\mathcal{H}}}\left(\square_2\psi + \mathfrak{d}T\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi + \frac{1}{\delta_{\mathcal{H}}^2}\mathfrak{d}^{\leq 1}\psi\right) + \frac{1}{\delta_{\mathcal{H}}}1_{\delta_{\mathcal{H}} \leq \Upsilon \leq 2\delta_{\mathcal{H}}}\mathfrak{d}^{\leq 2}\psi.
\end{aligned}$$

Now, we have in view of the definition of a and b ,

$$\begin{aligned} \frac{(a - \Upsilon b)(-2\omega - \Upsilon \underline{\kappa}) + \frac{\Upsilon}{2}e_3(a) - \frac{1}{2}e_4(a)}{(a - \Upsilon b)^2} &= \frac{(1 + O(\Upsilon))(-2\omega + O(\Upsilon)) + O(\Upsilon)}{(1 + O(\Upsilon))^2} \\ &= \frac{1}{2m} + O(\epsilon) + O(\Upsilon) \\ &= \frac{1}{2m_0} + O(\epsilon) + O(\Upsilon) \end{aligned}$$

where we used also our assumptions on ω and m . Thus, we have on the support of $\kappa_{\mathcal{H}}$

$$\begin{aligned} \frac{(a - \Upsilon b)(-2\omega - \Upsilon \underline{\kappa}) + \frac{\Upsilon}{2}e_3(a) - \frac{1}{2}e_4(a)}{(a - \Upsilon b)^2} &= \frac{1}{2m_0} + O(\epsilon + \delta_{\mathcal{H}}) \\ &= \frac{1}{2m_0} + O(\delta_{\mathcal{H}}) \end{aligned}$$

where we used the fact that $\epsilon \ll \delta_{\mathcal{H}}$ by assumption. Setting

$$d_0 := \frac{(a - \Upsilon b)(-2\omega - \Upsilon \underline{\kappa}) + \frac{\Upsilon}{2}e_3(a) - \frac{1}{2}e_4(a)}{(a - \Upsilon b)^2},$$

this concludes the proof of the lemma. \square

Commutation with re_4

Lemma 10.4.10. *We have, schematically,*

$$[\square_2, re_4]\psi = \frac{\Upsilon}{r} \left(1 + \frac{2m}{r\Upsilon^2} \right) \check{e}_4(re_4\psi) + \square_2\psi + \Gamma_g \mathfrak{d}^2\psi + \frac{1}{\Upsilon} r^{-2} \mathfrak{d}T\psi + r^{-2} \mathfrak{d}^2\psi + r^{-2} \mathfrak{d}\psi.$$

Proof. Recall that we have

$$\square_2\psi = -e_3(e_4(\psi)) + \mathbb{A}_2\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa} \right) e_4\psi - \frac{1}{2}\kappa e_3\psi + 2\eta e_\theta\psi.$$

Since we have

$$\begin{aligned} [re_4, e_3] &= 2r\omega e_3 - \frac{r}{2}\underline{\kappa}e_4 + \Gamma_b\mathfrak{d}, & [re_4, e_4] &= -\frac{r}{2}\kappa e_4 + \Gamma_g\mathfrak{d}, \\ [\mathfrak{d}_k, re_4] &= \frac{1}{2}r\kappa \mathfrak{d}_k + \Gamma_g\mathfrak{d} + \Gamma_g, & [\mathfrak{d}_k^*, re_4] &= \frac{1}{2}r\kappa \mathfrak{d}_k^* + \Gamma_g\mathfrak{d} + \Gamma_g, \end{aligned}$$

we infer, schematically,

$$\begin{aligned}
[\square_2, re_4]\psi &= -[e_3, re_4]e_4\psi - e_3[e_4, re_4]\psi + [\mathbb{A}_2, re_4]\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)[e_4, re_4]\psi \\
&\quad - re_4\left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi - \frac{1}{2}\kappa[e_3, re_4]\psi + \frac{1}{2}re_4(\kappa)e_3\psi + 2\eta[e_\theta, re_4]\psi - 2re_4(\eta)e_\theta\psi \\
&= \left(2r\omega e_3 - \frac{r}{2}\underline{\kappa}e_4\right)e_4\psi - \frac{1}{2}e_3(r\kappa e_4\psi) + r\kappa\mathbb{A}_2\psi + \frac{1}{2}re_4(\kappa)e_3\psi + \Gamma_g\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}\psi \\
&= \left(2r\omega e_3 - \frac{r}{2}\underline{\kappa}e_4\right)e_4\psi - \frac{1}{2}r\kappa e_3e_4\psi - \frac{1}{4}r\kappa^2e_3\psi + \Gamma_g\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}\psi.
\end{aligned}$$

Using again

$$\square_2\psi = -e_3(e_4(\psi)) + \mathbb{A}_2\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi - \frac{1}{2}\kappa e_3\psi + 2\eta e_\theta\psi.$$

we have

$$-\frac{1}{2}r\kappa e_3e_4\psi - \frac{1}{4}r\kappa^2e_3\psi = \frac{1}{2}r\kappa\square_2\psi + r^{-2}\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}\psi$$

and hence

$$\begin{aligned}
[\square_2, re_4]\psi &= \left(2r\omega e_3 - \frac{r}{2}\underline{\kappa}e_4\right)e_4\psi + \square_2\psi + \Gamma_g\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}\psi \\
&= \left(2r\omega\frac{1}{\Upsilon}(2T - e_4) - \frac{r}{2}\underline{\kappa}e_4\right)e_4\psi + \square_2\psi + \Gamma_g\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}\psi \\
&= \left(-2r\omega\frac{1}{\Upsilon} - \frac{r}{2}\underline{\kappa}\right)e_4(e_4\psi) + \square_2\psi + \Gamma_g\mathfrak{d}^2\psi + \frac{1}{\Upsilon}r^{-2}\mathfrak{d}T\psi + r^{-2}\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}\psi \\
&= \left(\Upsilon + \frac{2m}{r\Upsilon}\right)e_4(e_4\psi) + \square_2\psi + \Gamma_g\mathfrak{d}^2\psi + \frac{1}{\Upsilon}r^{-2}\mathfrak{d}T\psi + r^{-2}\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}\psi \\
&= \frac{\Upsilon}{r}\left(1 + \frac{2m}{r\Upsilon^2}\right)\check{e}_4(re_4\psi) + \square_2\psi + \Gamma_g\mathfrak{d}^2\psi + \frac{1}{\Upsilon}r^{-2}\mathfrak{d}T\psi + r^{-2}\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}\psi.
\end{aligned}$$

This concludes the proof of the lemma. \square

10.4.4 Some weighted estimates for wave equations

Recall from Corollary 10.4.5 that we have the following commutator identity

$$r\mathfrak{d}_2(\square_2\psi) - (\square_1 - K)(r\mathfrak{d}_2\psi) = -r\eta\square_2\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi.$$

In particular, to derive weighted estimates for $r \not{d}_2$, we need to derive weighted estimates for solutions ϕ to wave equations of the type

$$(\square_1 - V_1)\phi = N,$$

where ϕ is a reduced 1-scalar and the potential V_1 is given by $V_1 = V + K = -\kappa \underline{\kappa} + K$. This is done in the following theorem.

Theorem 10.4.11. *Let ϕ a reduced 1-scalar solution to*

$$(\square_1 - V_1)\phi = N, \quad V_1 = -\kappa \underline{\kappa} + K.$$

Then, ϕ satisfies for all $\delta \leq p \leq 2 - \delta$,

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_p[\phi](\tau) + B_p[\phi](\tau_1, \tau_2) + F_p[\phi](\tau_1, \tau_2) \\ & \lesssim E_p[\phi](\tau_1) + J_p[\phi, N](\tau_1, \tau_2) + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} \left| 1 - \frac{3m}{r} \right| |\phi|(|\phi| + |R\phi|) \\ & \quad + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |\phi|(|\phi| + |\not{d}\phi|), \end{aligned} \quad (10.4.4)$$

and $\check{\phi} = f_2 \check{e}_4 \phi$ satisfies for all $1 - \delta < q \leq 1 - \delta$,

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_q[\check{\phi}](\tau) + B_q[\check{\phi}](\tau_1, \tau_2) \\ & \lesssim E_q[\check{\phi}](\tau_1) + \check{J}_q[\check{\phi}, N](\tau_1, \tau_2) + E_{\max(q, \delta)}^1[\phi](\tau_1) + J_{\max(q, \delta)}^1[\phi, N] \\ & \quad + \int_{\mathcal{M}(\tau_1, \tau_2)} r^{q-3} |\check{\phi}|(|\check{\phi}| + |\not{d}\check{\phi}|). \end{aligned} \quad (10.4.5)$$

Remark 10.4.12. *Although we will not need it, we expect that the last 2 terms in the right-hand side of (10.4.4) and the last term in the right-hand side of (10.4.5) could be removed.*

Proof. We start with the following observations.

- (10.4.5) is the analog of (10.4.1), i.e. of Theorem 5.3.5 in the case $s = 0$, with V replaced by V_1 , and with the reduced 2-scalar ψ replaced by the reduced 1-scalar ϕ . The proof is in fact significantly easier in view of the presence of the term

$$\int_{\mathcal{M}(\tau_1, \tau_2)} r^{q-3} |\check{\phi}|(|\check{\phi}| + |\not{d}\check{\phi}|)$$

on the right-hand side of (10.4.5).

- (10.4.4) is the analog of (10.4.2), i.e. of Theorem 5.3.4 in the case $s = 0$, with V replaced by V_1 , and with the reduced 2-scalar ψ replaced by the reduced 1-scalar ϕ . The proof is in fact significantly easier in view of the presence of the terms

$$\int_{(trap)\mathcal{M}(\tau_1, \tau_2)} \left| 1 - \frac{3m}{r} \right| |\phi| (|\phi| + |R\phi|) + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |\phi| (|\phi| + |\mathfrak{d}\phi|)$$

on the right-hand side of (10.4.4).

- The boundary terms can be treated as in the proof of (10.4.1) and (10.4.2) in view of the fact that V_1 is a positive potential¹⁸.
- The only place where there might a potential difficulty concerns the proof of (10.4.5) in $(trap)\mathcal{M}$ where the second to last term on the right-hand side is required to have a more precise structure.

In view of the above observations, and in particular of the last one, we focus on recovering the bulk term leading to (10.4.5) in $(trap)\mathcal{M}$. To this end, we choose f and w as in Proposition 10.1.16. This yields¹⁹

$$\dot{\mathcal{E}}[fR, w](\Psi) \geq f'|R(\Psi)|^2 + r^{-1} \left(1 - \frac{3m}{r}\right) f|\nabla\Psi|^2 + O\left(\frac{1-\frac{3m}{r}}{r^3}\right) |\Psi|^2.$$

We infer

$$\begin{aligned} \dot{\mathcal{E}}[fR, w, M = 2hR](\Psi) &\geq f'|R(\Psi)|^2 + r^{-1} \left(1 - \frac{3m}{r}\right) f|\nabla\Psi|^2 + O\left(\frac{1-\frac{3m}{r}}{r^3}\right) |\Psi|^2 \\ &\quad + \frac{1}{2}r^{-2}(\Upsilon r^2 h)'|\Psi|^2 + h\Psi R(\Psi). \end{aligned}$$

We now choose a smooth h , compactly supported in $[5/2m_0, 7/2m_0]$, such that $h(3m) = 0$ and $h'(3m) = 1$ ²⁰. We infer $r^{-2}(\Upsilon r^2 h)'(3m) = 1/3 > 0$ and hence

$$\begin{aligned} \dot{\mathcal{E}}[fR, w, M](\Psi) &\geq f'|R(\Psi)|^2 + r^{-1} \left(1 - \frac{3m}{r}\right) f|\nabla\Psi|^2 + \frac{\Upsilon}{r^3} |\Psi|^2 \\ &\quad + O\left(\frac{1-\frac{3m}{r}}{r^3}\right) |\Psi| (|\Psi| + |R(\Psi)|). \end{aligned}$$

¹⁸We have

$$V_1 = -\kappa\underline{\kappa} + K = \frac{4\Upsilon + 1 + O(\epsilon)}{r^2}$$

in view of the assumptions so that V_1 is indeed a positive potential.

¹⁹Note that Proposition 10.1.16 does not use the particular form of the potential and the type of the reduced scalar ϕ and hence holds in our more general case.

²⁰This differs from the choice of h in the proof of (10.4.1) in order to avoid using a Poincaré inequality (which depends of the type of the reduced scalar) and the particular form of the potential V_1 .

In view of the choice of f in Proposition 10.1.16, we have

$$f' \gtrsim \frac{1}{r^3}, \quad \left(1 - \frac{3m}{r}\right) f \geq \left(1 - \frac{3m}{r}\right)^2,$$

and hence, there exists two constants $c_0 > 0$ and $C_0 > 0$ such that

$$\begin{aligned} \dot{\mathcal{E}}[fR, w_0, M](\Psi) &\geq c_0 \left(\frac{1}{r^3} |R(\Psi)|^2 + r^{-1} \left(1 - \frac{3m}{r}\right)^2 |\nabla \Psi|^2 + \frac{\Upsilon}{r^3} |\Psi|^2 \right) \\ &\quad - C_0 \frac{|1 - \frac{3m}{r}|}{r^3} |\Psi| (|\Psi| + |R(\Psi)|). \end{aligned}$$

The last term above is responsible for the second to last term on the right-hand side of (10.4.5). \square

Next, we have the following consequence of (10.4.1) and Theorem 10.4.11.

Corollary 10.4.13. *Let ϕ be a reduced k -scalar for $k = 1, 2$ such that ϕ satisfies²¹*

$$(\square_k - W)\phi = O\left(\frac{\epsilon}{r^2 u_{trap}^{1+\delta_{dec}-2\delta_0}}\right) \mathfrak{d}\phi_1 + \phi_2$$

where ϕ_1 and ϕ_2 are given reduced scalars, and where $W = V$ in the case $k = 2$ and $W = V_1$ in the case $k = 1$. Then, ϕ satisfies for all $\delta \leq p \leq 2 - \delta$,

$$\begin{aligned} &\sup_{\tau \in [\tau_1, \tau_2]} E_p[\phi](\tau) + B_p[\phi](\tau_1, \tau_2) + F_p[\phi](\tau_1, \tau_2) \\ &\lesssim E_p[\phi](\tau_1) + \epsilon^2 \left(\sup_{[\tau_1, \tau_2]} E[\phi_1](\tau) + B_p[\phi_1](\tau_1, \tau_2) \right) + J_p[\phi, \phi_2](\tau_1, \tau_2) \\ &\quad + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} \left| 1 - \frac{3m}{r} \right| |\phi| (|\phi| + |R\phi|) + \int_{(trap^p)\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |\phi| (|\phi| + |\mathfrak{d}\phi|). \end{aligned}$$

Proof. The wave equation for ϕ satisfies the assumptions of (10.4.1) and Theorem 10.4.11 with

$$N = O\left(\frac{\epsilon}{r^2 u_{trap}^{1+\delta_{dec}-2\delta_0}}\right) \mathfrak{d}\phi_1 + \phi_2.$$

²¹Recall that we have $\delta_0 \ll \delta_{dec}$ in view of (5.1.1), and hence $\delta_{dec} - 2\delta_0 > 0$.

We deduce

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_p[\phi](\tau) + B_p[\phi](\tau_1, \tau_2) + F_p[\phi](\tau_1, \tau_2) \\ & \lesssim E_p[\phi](\tau_1) + J_p \left[\phi, O \left(\frac{\epsilon}{r^2 u_{tr\text{ap}}^{1+\delta_{dec}-2\delta_0}} \right) \mathfrak{d}\phi_1 + \phi_2 \right] (\tau_1, \tau_2) \\ & \quad + \int_{(tr\text{ap})\mathcal{M}(\tau_1, \tau_2)} \left| 1 - \frac{3m}{r} \right| |\phi| (|\phi| + |R\phi|) + \int_{(tr\text{q}/p)\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |\phi| (|\phi| + |\mathfrak{d}\phi|). \end{aligned}$$

Now, in view of the definition

$$\begin{aligned} J_{p,R}[\psi, N](\tau_1, \tau_2) &= \left| \int_{\mathcal{M}_{\geq R}(\tau_1, \tau_2)} r^p \check{e}_4 \psi N \right|, \\ J_p[\psi, N](\tau_1, \tau_2) &= \left(\int_{\tau_1}^{\tau_2} d\tau \|N\|_{L^2((tr\text{ap})\Sigma(\tau))} \right)^2 + \int_{(tr\text{q}/p)\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} |N|^2 \\ & \quad + J_{p,4m_0}[\psi, N](\tau_1, \tau_2), \end{aligned}$$

we have

$$\begin{aligned} & J_p \left[\phi, O \left(\frac{\epsilon}{r^2 u_{tr\text{ap}}^{1+\delta_{dec}-2\delta_0}} \right) \mathfrak{d}\phi_1 + \phi_2 \right] (\tau_1, \tau_2) \\ & \lesssim J_p \left[\phi, O \left(\frac{\epsilon}{r^2 u_{tr\text{ap}}^{1+\delta_{dec}-2\delta_0}} \right) \mathfrak{d}\phi_1 \right] (\tau_1, \tau_2) + J_p[\phi, \phi_2](\tau_1, \tau_2) \end{aligned}$$

and, for $\delta \leq p \leq 2 - \delta$, using also $\delta_{dec} - 2\delta_0 > 0$, we have

$$\begin{aligned} & J_p \left[\phi, O \left(\frac{\epsilon}{r^2 u_{tr\text{ap}}^{1+\delta_{dec}-2\delta_0}} \right) \mathfrak{d}\phi_1 \right] (\tau_1, \tau_2) \\ & \lesssim \epsilon^2 \left(\int_{\tau_1}^{\tau_2} \|\mathfrak{d}\phi_1\|_{L^2((tr\text{ap})\Sigma(\tau))} \frac{d\tau}{\tau^{1+\delta_{dec}-2\delta_0}} \right)^2 + \epsilon^2 \int_{(tr\text{q}/p)\mathcal{M}(\tau_1, \tau_2)} r^{\delta-3} |\mathfrak{d}\phi_1|^2 \\ & \quad + \epsilon \left| \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^{p-2} \check{e}_4(\psi) \mathfrak{d}\phi_1 \right| \\ & \lesssim \epsilon^2 \sup_{[\tau_1, \tau_2]} \|\mathfrak{d}\phi_1\|_{L^2((tr\text{ap})\Sigma(\tau))}^2 + \epsilon^2 \int_{(tr\text{q}/p)\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |\mathfrak{d}\phi_1|^2 \\ & \quad + \epsilon \left(\int_{(tr\text{q}/p)\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |\mathfrak{d}\phi|^2 \right)^{\frac{1}{2}} \left(\int_{(tr\text{q}/p)\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |\mathfrak{d}\phi_1|^2 \right)^{\frac{1}{2}} \\ & \lesssim \epsilon^2 \left(\sup_{[\tau_1, \tau_2]} E^1[\phi_1](\tau) + B_p^1[\phi_1](\tau_1, \tau_2) \right) + \epsilon (B_p^1[\phi_1](\tau_1, \tau_2))^{\frac{1}{2}} (B_p^1[\phi](\tau_1, \tau_2))^{\frac{1}{2}}. \end{aligned}$$

We immediately deduce

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_p[\phi](\tau) + B_p[\phi](\tau_1, \tau_2) + F_p[\phi](\tau_1, \tau_2) \\ & \lesssim E_p[\phi](\tau_1) + \epsilon^2 \left(\sup_{[\tau_1, \tau_2]} E[\phi_1](\tau) + B_p[\phi_1](\tau_1, \tau_2) \right) + J_p[\phi, \phi_2](\tau_1, \tau_2) \\ & \quad + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} \left| 1 - \frac{3m}{r} \right| |\phi| (|\phi| + |R\phi|) + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |\phi| (|\phi| + |\mathfrak{d}\phi|). \end{aligned}$$

This concludes the proof of the corollary. \square

Finally, we end this section with the following lemma.

Lemma 10.4.14. *Let ϕ be a reduced k -scalar for $k = 1, 2$, and let X a vectorfield. We have for all $\delta \leq p \leq 2 - \delta$,*

$$\begin{aligned} & \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} \left| 1 - \frac{3m}{r} \right| |\mathfrak{d}\phi| (|X\phi| + |R(X\phi)|) + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |\mathfrak{d}\phi| (|X\phi| + |\mathfrak{d}(X\phi)|) \\ & \lesssim (B_p[X\phi](\tau_1, \tau_2))^{\frac{1}{2}} (B_p[\phi](\tau_1, \tau_2))^{\frac{1}{2}}. \end{aligned}$$

Proof. The proof follows immediately from the definition of $B_p[\phi](\tau_1, \tau_2)$. \square

10.4.5 Proof of Theorem 5.3.4

We now conclude the proof of Theorem 5.3.4 for all $0 \leq s \leq k_{small} + 30$ by recovering higher derivatives $s \geq 1$ one by one starting from the estimate $s = 0$ provided by (10.4.1). As explained in section 10.4.2, it suffices to recover the estimates for $s = 1$ from the one for $s = 0$ as the procedure to recover the estimate for $s + 1$ from the one for s is completely analogous. We now follow the strategy outlined in section 10.4.2.

Recovering estimates for $T\psi$

Recall that ψ satisfies

$$\square_2 \psi + V\psi = N, \quad V = \kappa \underline{\kappa},$$

and recall also from Corollary 10.4.3 that we have

$$[T, \square_2] \psi = \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \psi.$$

We infer

$$\square_2(T\psi) + VT(\psi) = T(N) + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi.$$

In view of Corollary 10.4.13 with $\phi = T(\psi)$, $\phi_1 = \mathfrak{d}^{\leq 1}\psi$ and $\phi_2 = T(N)$, and in view of (10.4.3), we deduce

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_p[T\psi](\tau) + B_p[T\psi](\tau_1, \tau_2) + F_p[T\psi](\tau_1, \tau_2) \\ \lesssim & E_p[T\psi](\tau_1) + \epsilon^2 \left(\sup_{[\tau_1, \tau_2]} E[\mathfrak{d}^{\leq 1}\psi](\tau) + B_p[\mathfrak{d}^{\leq 1}\psi](\tau_1, \tau_2) \right) + J_p[T(\psi), T(N)](\tau_1, \tau_2) \\ & + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} \left| 1 - \frac{3m}{r} \right| |T\phi| (|T\phi| + |R(T\phi)|) \\ & + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |T\phi| (|T\phi| + |\mathfrak{d}(T\phi)|), \end{aligned}$$

and hence, using Lemma 10.4.14 with $X = T$, we infer for any $\delta \leq p \leq 2 - \delta$,

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_p[T\psi](\tau) + B_p[T\psi](\tau_1, \tau_2) + F_p[T\psi](\tau_1, \tau_2) \tag{10.4.6} \\ \lesssim & E_p[T\psi](\tau_1) + J_p^1[\psi, N](\tau_1, \tau_2) + \epsilon^2 \left(\sup_{[\tau_1, \tau_2]} E^1[\psi](\tau) + B_p^1[\psi](\tau_1, \tau_2) \right) + B_p[\psi](\tau_1, \tau_2). \end{aligned}$$

Recovering estimates for $r \not\! d_2 \psi$

Recall that ψ satisfies

$$\square_2\psi + V\psi = N, \quad V = \kappa \underline{\kappa},$$

and recall also from Corollary 10.4.5 that we have

$$r \not\! d_2(\square_2\psi) - (\square_1 - K)(r \not\! d_2\psi) = -r\eta \square_2\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi$$

We infer

$$\begin{aligned} \square_1(r \not\! d_2\psi) + (V - K)r \not\! d_2\psi &= r\eta \square_2\psi + r \not\! d_2(N) + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi \\ &= -r\eta N + r \not\! d_2(N) + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi. \end{aligned}$$

and hence

$$\square_1(r \not\! d_2\psi) + (V - K)r \not\! d_2\psi = -r\eta N + r \not\! d_2(N) + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi.$$

In view of Corollary 10.4.13 with $\phi = r \not\!d_2 \psi$, $\phi_1 = \mathfrak{d}^{\leq 1} \psi$ and $\phi_2 = -r\eta N + r \not\!d_2(N)$, and in view of (10.4.3), we deduce

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_p[r \not\!d_2 \psi](\tau) + B_p[r \not\!d_2 \psi](\tau_1, \tau_2) + F_p[r \not\!d_2 \psi](\tau_1, \tau_2) \\ \lesssim & E_p[r \not\!d_2 \psi](\tau_1) + \epsilon^2 \left(\sup_{[\tau_1, \tau_2]} E[\mathfrak{d}^{\leq 1} \psi](\tau) + B_p[\mathfrak{d}^{\leq 1} \psi](\tau_1, \tau_2) \right) \\ & + J_p \left[r \not\!d_2 \psi, -r\eta N + r \not\!d_2(N) \right](\tau_1, \tau_2) + \int_{(\text{trap})\mathcal{M}(\tau_1, \tau_2)} \left| 1 - \frac{3m}{r} \right| |r \not\!d_2 \phi| (|r \not\!d_2 \phi| + |R(r \not\!d_2 \phi)|) \\ & + \int_{(\text{trap})\mathcal{M}(\tau_1, \tau_2)} r^{p-3} |r \not\!d_2 \phi| (|r \not\!d_2 \phi| + |\mathfrak{d}(r \not\!d_2 \phi)|), \end{aligned}$$

and hence, using Lemma 10.4.14 with $X = r \not\!d_2$, we infer for any $\delta \leq p \leq 2 - \delta$,

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_p[r \not\!d_2 \psi](\tau) + B_p[r \not\!d_2 \psi](\tau_1, \tau_2) + F_p[r \not\!d_2 \psi](\tau_1, \tau_2) \tag{10.4.7} \\ \lesssim & E_p[r \not\!d_2 \psi](\tau_1) + J_p^1[\psi, N](\tau_1, \tau_2) + \epsilon^2 \left(\sup_{[\tau_1, \tau_2]} E^1[\psi](\tau) + B_p^1[\psi](\tau_1, \tau_2) \right) + B_p[\psi](\tau_1, \tau_2). \end{aligned}$$

Recovering estimates for $R\psi$ in $r \leq r_0$

We start with the following lemma.

Lemma 10.4.15. *Let ψ satisfy*

$$\square_2 \psi = V\psi + N, \quad V = \kappa \underline{\kappa}.$$

Then, $R^2\psi$ satisfies

$$R^2\psi = -\Upsilon N + T^2\psi + O(r^{-2})\not\!d^2\psi + O(r^{-1})\mathfrak{d}\psi + O(r^{-2})\psi.$$

Proof. Recall that we have

$$\square_2 \psi = -e_3(e_4(\psi)) + \not\!d_2 \psi - \frac{1}{2} \kappa e_3 \psi + \left(-\frac{1}{2} \underline{\kappa} + 2\underline{\omega} \right) e_4 \psi + 2\eta e_\theta \psi$$

and

$$e_4 = T + R, \quad \Upsilon e_3 = (T - R).$$

We infer

$$\begin{aligned} \Upsilon \square_2 \psi &= -(T - R)(T + R)\psi + \Upsilon \Delta_2 \psi - \frac{\Upsilon}{2} \kappa e_3 \psi + \Upsilon \left(-\frac{1}{2} \underline{\kappa} + 2\underline{\omega} \right) e_4 \psi + 2\Upsilon \eta e_\theta \psi \\ &= -T^2 \psi + R^2 \psi - [T, R]\psi + O(r^{-2}) \mathfrak{I}^2 \psi + O(r^{-1}) \mathfrak{D} \psi \end{aligned}$$

and hence

$$\begin{aligned} R^2 \psi &= -\Upsilon \square_2 \psi + T^2 \psi - [T, R]\psi + O(r^{-2}) \mathfrak{I}^2 \psi + O(r^{-1}) \mathfrak{D} \psi \\ &= -\Upsilon N + T^2 \psi - [T, R]\psi + O(r^{-2}) \mathfrak{I}^2 \psi + O(r^{-1}) \mathfrak{D} \psi + O(r^{-2}) \psi \end{aligned}$$

where we used the fact that $\square_2 \psi = V\psi + N$ and $V = \kappa \underline{\kappa} = O(r^{-2})$. Also, we have

$$\begin{aligned} [T, R]\psi &= \frac{1}{4} [e_4 + \Upsilon e_3, e_4 - \Upsilon e_3] \psi \\ &= \frac{1}{2} \left(-e_4(\Upsilon) e_3 + \Upsilon [e_3, e_4] \right) \psi \\ &= O(r^{-2}) \mathfrak{D} \psi \end{aligned}$$

and thus

$$R^2 \psi = -\Upsilon N + T^2 \psi + O(r^{-2}) \mathfrak{I}^2 \psi + O(r^{-1}) \mathfrak{D} \psi + O(r^{-2}) \psi.$$

This concludes the proof of the lemma. □

We now estimate $R\psi$ in $r \leq r_0$ for a fixed $r_0 \geq 4m_0$ that will be chosen large enough. First, in view of the identity of the previous lemma, i.e.

$$R^2 \psi = -\Upsilon N + T^2 \psi + O(r^{-2}) \mathfrak{I}^2 \psi + O(r^{-1}) \mathfrak{D} \psi + O(r^{-2}) \psi,$$

we infer

$$\begin{aligned} &\sup_{[\tau_1, \tau_2]} \int_{\Sigma_{r \leq r_0}(\tau)} |R^2 \psi|^2 + \int_{\mathcal{M}_{r \leq r_0}(\tau_1, \tau_2)} \left(1 - \frac{3m}{r} \right)^2 (R^2 \psi)^2 \tag{10.4.8} \\ &\lesssim \sup_{[\tau_1, \tau_2]} \left(E[T\psi] + E[r \mathfrak{I}_2 \psi] + E[\psi] + \int_{\Sigma_{r \leq r_0}(\tau)} N^2 \right) \\ &\quad + \int_{\mathcal{M}_{r \leq r_0}(\tau_1, \tau_2)} N^2 + \text{Morr}[T\psi](\tau_1, \tau_2) + \text{Morr}[r \mathfrak{I}_2 \psi](\tau_1, \tau_2) + \text{Morr}[\psi](\tau_1, \tau_2). \end{aligned}$$

Next, we remove the degeneracy of the above estimate at $r = 3m$. Recall from Corollary 10.4.7 that we have in the region $r \leq 4m_0$

$$\square_2(R\psi) = \left(1 - \frac{3m}{r} \right) \mathfrak{D}^2 \psi + O \left(\frac{\epsilon}{u_{\text{trap}}^{1+\delta_{\text{dec}}-2\delta_0}} \right) \mathfrak{D}^2 \psi + O(1) \mathfrak{D}^{\leq 1} \psi + O(1) \mathfrak{D}^{\leq 1} N.$$

Then,

1. multiplying $R\psi$ with a cut-off function equal to one on $[5/2m_0, 7/2m_0]$ and vanishing on $[9/4m_0, 4m_0]$ and inferring the corresponding wave equation from the above one for $R\psi$,
2. relying on the Morawetz estimate of Proposition 10.1.12 with the particular choice $f(r) = r - 3m$,
3. adding a large multiple of the energy estimate,
4. using Proposition 10.1.32 for the boundary terms,

we easily infer the following estimate

$$\int_{(trap)\mathcal{M}(\tau_1, \tau_2)} (R^2\psi)^2 \lesssim \int_{\mathcal{M}_{r \leq 4m_0}(\tau_1, \tau_2)} \left(\left(1 - \frac{3m}{r}\right)^2 (\mathfrak{d}^2\psi)^2 + (\mathfrak{d}\psi)^2 + (\mathfrak{d}^{\leq 1}N)^2 \right) + E[R\psi](\tau_1) + \epsilon \sup_{[\tau_1, \tau_2]} E^1[\psi](\tau).$$

Together with (10.4.8), we infer

$$\begin{aligned} & \sup_{[\tau_1, \tau_2]} \int_{\Sigma_{r \leq r_0}(\tau)} |R^2\psi|^2 + \int_{\mathcal{M}_{r \leq r_0}(\tau_1, \tau_2)} |R^2\psi|^2 & (10.4.9) \\ & \lesssim E[R\psi](\tau_1) + \sup_{[\tau_1, \tau_2]} \left(\epsilon E^1[\psi](\tau) + E[T\psi](\tau) + E[r \not{d}_2\psi](\tau) + E[\psi](\tau) \right) \\ & \quad + J_p^1[\psi, N](\tau_1, \tau_2) + \text{Morr}[T\psi](\tau_1, \tau_2) + \text{Morr}[r \not{d}_2\psi](\tau_1, \tau_2) + \text{Morr}[\psi](\tau_1, \tau_2). \end{aligned}$$

Recovering estimates for $Y_{\mathcal{H}}\psi$

Recall that ψ satisfies

$$\square_2\psi + V\psi = N, \quad V = \kappa \underline{\kappa},$$

and recall also from Lemma 10.4.9

$$\begin{aligned} [\square_2, Y_{\mathcal{H}}]\psi &= d_0 Y_{(0)}(Y_{\mathcal{H}}\psi) + 1_{\Upsilon \leq 2\delta_{\mathcal{H}}} \left(\square_2\psi + \mathfrak{d}T\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi + \frac{1}{\delta_{\mathcal{H}}^2}\mathfrak{d}^{\leq 1}\psi \right) \\ & \quad + \frac{1}{\delta_{\mathcal{H}}} 1_{\delta_{\mathcal{H}} \leq \Upsilon \leq 2\delta_{\mathcal{H}}} \mathfrak{d}^{\leq 2}\psi \end{aligned}$$

where the scalar function d_0 satisfying the bound

$$d_0 = \frac{1}{2m_0} + O(\delta_{\mathcal{H}}) \text{ on the support of } \kappa_{\mathcal{H}}.$$

We infer

$$\begin{aligned} \square_2(Y_{\mathcal{H}}\psi) + VY_{\mathcal{H}}(\psi) &= d_0Y_{(0)}(Y_{\mathcal{H}}\psi) + 1_{\Upsilon \leq 2\delta_{\mathcal{H}}} \left(N + \mathfrak{d}T\psi + \epsilon\mathfrak{d}^2\psi + \frac{1}{\delta_{\mathcal{H}}^2}\mathfrak{d}^{\leq 1}\psi \right) \\ &\quad + \frac{1}{\delta_{\mathcal{H}}}1_{\delta_{\mathcal{H}} \leq \Upsilon \leq 2\delta_{\mathcal{H}}}\mathfrak{d}^{\leq 2}\psi + Y_{\mathcal{H}}(N). \end{aligned}$$

Then,

1. we use the redshift vectorfield $Y_{\mathcal{H}}$ as a multiplier,
2. we rely on Proposition 10.1.29,
3. we use the fact that $d_0 \geq 0$,
4. we add a large multiple of the energy,
5. we use Proposition 10.1.32 for the boundary terms.

We easily infer

$$\begin{aligned} \sup_{[\tau_1, \tau_2]} E[Y_{\mathcal{H}}\psi] + \text{Morr}[Y_{\mathcal{H}}\psi](\tau_1, \tau_2) &\lesssim E[Y_{\mathcal{H}}\psi](\tau_1) + J_p^1[\psi, N](\tau_1, \tau_2) + \epsilon\text{Morr}[\mathfrak{d}\psi](\tau_1, \tau_2) \\ &\quad + \text{Morr}[R\psi](\tau_1, \tau_2) + \text{Morr}[T\psi](\tau_1, \tau_2) \\ &\quad + \text{Morr}[r\mathfrak{d}_2\psi](\tau_1, \tau_2) + \text{Morr}[\psi](\tau_1, \tau_2). \end{aligned} \quad (10.4.10)$$

Recovering estimates for $re_4\psi$ in $r \geq r_0$

Recall that ψ satisfies

$$\square_2\psi + V\psi = N, \quad V = \kappa\underline{\kappa},$$

and recall also from Lemma 10.4.10

$$[\square_2, re_4]\psi = \frac{\Upsilon}{r} \left(1 + \frac{2m}{r\Upsilon^2} \right) \check{e}_4(re_4\psi) + \square_2\psi + \Gamma_g\mathfrak{d}^2\psi + \frac{1}{\Upsilon}r^{-2}\mathfrak{d}T\psi + r^{-2}\mathfrak{d}^2\psi + r^{-2}\mathfrak{d}\psi.$$

We infer

$$\begin{aligned} \square_2(re_4\psi) + Vre_4(\psi) &= \frac{\Upsilon}{r} \left(1 + \frac{2m}{r\Upsilon^2} \right) \check{e}_4(re_4\psi) + O\left(\frac{\epsilon}{r^2}\right)\mathfrak{d}^2\psi + \frac{1}{\Upsilon}r^{-2}\mathfrak{d}T\psi + r^{-2}\mathfrak{d}^2\psi \\ &\quad + r^{-2}\mathfrak{d}^{\leq 1}\psi + N + re_4(N). \end{aligned}$$

Then,

1. as in section 10.2.3, we use the vectorfield $f_p e_4$ as a multiplier, where $f_p = \theta_{r_0}(r) r^p e_4$ and the cut-off $\theta_{r_0}(r)$ is equal to one in the region $r \geq r_0$ and vanishes in the region $r \leq r_0/2$,
2. we rely on Proposition 10.2.7 to control the bulk and the boundary terms,
3. we use the fact that the prefactor of the term $\check{\epsilon}_4(re_4\psi)$ on the right-hand side is positive for $r \geq 4m_0$, i.e.

$$\frac{\Upsilon}{r} \left(1 + \frac{2m}{r\Upsilon^2} \right) \geq 0 \text{ for } r \geq 4m_0.$$

We easily infer

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_{p,r \geq r_0}[re_4\psi](\tau) + B_{p,r \geq r_0}[re_4\psi](\tau_1, \tau_2) + F_{p,r \geq r_0}[re_4\psi](\tau_1, \tau_2) \\ & \lesssim E_p^1[re_4\psi](\tau_1) + J_p^1[\psi, N](\tau_1, \tau_2) + B_{p,r_0/2 \leq r < r_0}^1[\psi](\tau_1, \tau_2) + \epsilon B_p^1[\psi](\tau_1, \tau_2) \\ & \quad + B_p[T\psi](\tau_1, \tau_2) + B_p[r \not\! d_2 \psi](\tau_1, \tau_2) + B_p[\psi](\tau_1, \tau_2). \end{aligned} \quad (10.4.11)$$

Conclusion of the proof of Theorem 5.3.4

Gathering the estimates (10.4.6), (10.4.7), (10.4.9), (10.4.10) and (10.4.11), we infer for any $\delta \leq p \leq 2 - \delta$,

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_p^1[\psi](\tau) + B_p^1[\psi](\tau_1, \tau_2) + F_p^1[\psi](\tau_1, \tau_2) \\ & \lesssim E_p^1[\psi](\tau_1) + J_p^1[\psi, N](\tau_1, \tau_2) + \epsilon^2 \left(\sup_{[\tau_1, \tau_2]} E_p^1[\psi](\tau) + B_p^1[\psi](\tau_1, \tau_2) \right) \\ & \quad + \sup_{[\tau_1, \tau_2]} E_p[\psi](\tau) + B_p[\psi](\tau_1, \tau_2), \end{aligned}$$

and hence

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_p^1[\psi](\tau) + B_p^1[\psi](\tau_1, \tau_2) + F_p^1[\psi](\tau_1, \tau_2) \\ & \lesssim E_p^1[\psi](\tau_1) + J_p^1[\psi, N](\tau_1, \tau_2) + \sup_{[\tau_1, \tau_2]} E_p[\psi](\tau) + B_p[\psi](\tau_1, \tau_2). \end{aligned}$$

In view of (10.4.1), we deduce

$$\sup_{\tau \in [\tau_1, \tau_2]} E_p^1[\psi](\tau) + B_p^1[\psi](\tau_1, \tau_2) + F_p^1[\psi](\tau_1, \tau_2) \lesssim E_p^1[\psi](\tau_1) + J_p^1[\psi, N](\tau_1, \tau_2)$$

which is Theorem 5.3.4 in the case $s = 1$. We have thus deduced Theorem 5.3.4 in the case $s = 1$ from the case $s = 0$, i.e. (10.4.1). Since going from $s = 0$ to $s = 1$ is analogous to going from s to $s + 1$, higher order derivatives $k \leq k_{small} + 30$ are recovered in the same fashion. This concludes the proof of Theorem 5.3.4.

10.4.6 Proof of Theorem 5.3.5

We now conclude the proof of Theorem 5.3.5 for all $0 \leq s \leq k_{small} + 29$ by recovering higher derivatives $s \geq 1$ one by one starting from the estimate $s = 0$ provided by (10.4.2). As explained in section 10.4.2, it suffices to recover the estimates for $s = 1$ from the one for $s = 0$ as the procedure to recover the estimate for $s + 1$ from the one for s is completely analogous. We now follow the strategy outlined in section 10.4.2.

Recovering estimates for $T\check{\psi}$

Recall from Proposition 10.3.1 that $\check{\psi} = f_2\check{e}_4\psi$ satisfies

$$\square\check{\psi} - V\check{\psi} = \frac{2}{r} \left(1 - \frac{3m}{r}\right) e_4\check{\psi} + \check{N} + f_2 \left(e_4 + \frac{3}{r}\right) N$$

where,

$$\check{N} = \begin{cases} O(r^{-2})\mathfrak{d}^{\leq 1}\psi + r\Gamma_b e_4\mathfrak{d}\psi + \mathfrak{d}^{\leq 1}(\Gamma_b)\mathfrak{d}^{\leq 1}\psi + r\mathfrak{d}^{\leq 1}(\Gamma_g)e_3\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi, & r \geq 6m_0, \\ O(1)\mathfrak{d}^{\leq 2}\psi, & 4m_0 \leq r \leq 6m_0, \end{cases}$$

and recall also from Corollary 10.4.3 that we have

$$[T, \square_2]\check{\psi} = \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\check{\psi}.$$

We infer

$$\square(T\check{\psi}) - VT(\check{\psi}) = \frac{2}{r} \left(1 - \frac{3m}{r}\right) e_4(T\check{\psi}) + N^T + T \left(f_2 \left(e_4 + \frac{3}{r} \right) N \right),$$

where we have, in view of the estimates²² of Lemma 5.1.1 for $\mathfrak{d}^k \Gamma_g$ and $\mathfrak{d}^k \Gamma_b$ with $k \leq k_{small} + 30$,

$$N^T = \begin{cases} O\left(\frac{1}{\tau^{1+\delta_{dec}-2\delta_0}}\right) \left(e_4 \mathfrak{d}^2 \psi + r^{-1} \mathfrak{d}^{\leq 2} \psi\right) \\ + O\left(\frac{1}{r\tau^{\frac{1}{2}+\delta_{dec}-2\delta_0}}\right) e_3 \mathfrak{d}^{\leq 1} \psi + O\left(\frac{1}{r^2}\right) \left(\mathfrak{d}^{\leq 3} \psi + \epsilon \mathfrak{d}^{\leq 2} \check{\psi}\right), & r \geq 6m_0, \\ O(1) \mathfrak{d}^{\leq 3} \psi, & 4m_0 \leq r \leq 6m_0. \end{cases}$$

In view of (10.4.2) with $T\check{\psi}$ instead of $\check{\psi}$ and with

$$N^T + T \left(f_2 \left(e_4 + \frac{3}{r} \right) N \right)$$

instead of $\check{N} + f_2 \left(e_4 + \frac{3}{r} \right) N$, we deduce

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_q[T\check{\psi}](\tau) + B_q[T\check{\psi}](\tau_1, \tau_2) \\ & \lesssim E_q[T\check{\psi}](\tau_1) + J_q \left[T\check{\psi}, N^T + T \left(f_2 \left(e_4 + \frac{3}{r} \right) N \right) \right] (\tau_1, \tau_2) \\ & \quad + E_{\max(q, \delta)}^1 [T\psi](\tau_1) + J_{\max(q, \delta)}^1 [T\psi, TN] \\ & \lesssim E_q[T\check{\psi}](\tau_1) + \check{J}_q^1 [\check{\psi}, N] (\tau_1, \tau_2) + J_q [T\check{\psi}, N^T] (\tau_1, \tau_2) \\ & \quad + E_{\max(q, \delta)}^2 [\psi](\tau_1) + J_{\max(q, \delta)}^2 [\psi, N], \end{aligned}$$

so that it remains to estimate

$$J_q [T\check{\psi}, N^T] (\tau_1, \tau_2) = J_{q, 4m_0} [T\check{\psi}, N^T] (\tau_1, \tau_2) = \left| \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^q (\check{e}_4(T\check{\psi})) N^T \right|.$$

²²Here, unlike the proof of Theorem 5.3.4 above, the non sharp estimates of section 10.4.1 are not enough, and we need instead to rely on the stronger estimates provided by Lemma 5.1.1.

We have, in view of the definition of N^T ,

$$\begin{aligned}
 & J_q[T \check{\psi}, N^T](\tau_1, \tau_2) \\
 \lesssim & \left| \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^q (\check{e}_4(T \check{\psi})) \frac{1}{\tau^{1+\delta_{dec}-2\delta_0}} \left(e_4 \mathfrak{d}^2 \psi + r^{-1} \mathfrak{d}^{\leq 2} \psi \right) \right| \\
 & + \left| \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^q (\check{e}_4(T \check{\psi})) \frac{1}{r\tau^{\frac{1}{2}+\delta_{dec}-2\delta_0}} e_3 \mathfrak{d}^{\leq 1} \psi \right| \\
 & + \left| \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^q (\check{e}_4(T \check{\psi})) \frac{1}{r^2} \mathfrak{d}^{\leq 3} \psi \right| + \left| \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^q (\check{e}_4(T \check{\psi})) \frac{\epsilon}{r^2} \mathfrak{d}^{\leq 2} \check{\psi} \right| \\
 \lesssim & \left(\sup_{[\tau_1, \tau_2]} \int_{\Sigma(\tau)} r^q (\check{e}_4(T \check{\psi}))^2 \right)^{\frac{1}{2}} \left\{ \left(\sup_{[\tau_1, \tau_2]} \int_{\Sigma(\tau)} r^q \left((e_4 \mathfrak{d}^2 \psi)^2 + r^{-2} (\mathfrak{d}^{\leq 2} \psi)^2 \right) \right)^{\frac{1}{2}} \right. \\
 & + \left. \left(\int_{\mathcal{M}_{r \geq 4m_0}(\tau_1, \tau_2)} r^{q-2} (e_3 \mathfrak{d}^{\leq 1} \psi)^2 \right)^{\frac{1}{2}} \right\} + \epsilon \int_{\mathcal{M}(\tau_1, \tau_2)} r^{q-3} (\mathfrak{d}^{\leq 2} \check{\psi})^2 \\
 & + \left(\int_{\mathcal{M}(\tau_1, \tau_2)} r^{q-1} (\check{e}_4(T \check{\psi}))^2 \right)^{\frac{1}{2}} \left(\int_{\mathcal{M}(\tau_1, \tau_2)} r^{q-3} (\mathfrak{d}^{\leq 3} \psi)^2 \right)^{\frac{1}{2}}
 \end{aligned}$$

which yields, using in particular the fact that $q \leq 1 - \delta$,

$$\begin{aligned}
 & J_q[T \check{\psi}, N^T](\tau_1, \tau_2) \\
 \lesssim & \left(\sup_{[\tau_1, \tau_2]} E_q[T \check{\psi}](\tau) \right)^{\frac{1}{2}} \left\{ \sup_{[\tau_1, \tau_2]} E_{\max(q, \delta)}^2[\psi](\tau) + B_{\max(q, \delta)}^1[\psi](\tau_1, \tau_2) \right\}^{\frac{1}{2}} + \epsilon B_q^1[\check{\psi}](\tau_1, \tau_2) \\
 & + \left(B_q[T \check{\psi}](\tau_1, \tau_2) \right)^{\frac{1}{2}} \left(B_{\max(q, \delta)}^2[\psi](\tau_1, \tau_2) \right)^{\frac{1}{2}}.
 \end{aligned}$$

We deduce

$$\begin{aligned}
 & \sup_{\tau \in [\tau_1, \tau_2]} E_q[T \check{\psi}](\tau) + B_q[T \check{\psi}](\tau_1, \tau_2) \\
 \lesssim & E_q[T \check{\psi}](\tau_1) + \check{J}_q^1[\check{\psi}, N](\tau_1, \tau_2) + \epsilon B_q^1[\check{\psi}](\tau_1, \tau_2) \\
 & + \sup_{[\tau_1, \tau_2]} E_{\max(q, \delta)}^2[\psi](\tau) + B_{\max(q, \delta)}^2[\psi](\tau_1, \tau_2) + J_{\max(q, \delta)}^2[\psi, N].
 \end{aligned}$$

Together with Theorem 5.3.4, this yields

$$\begin{aligned}
 \sup_{\tau \in [\tau_1, \tau_2]} E_q[T \check{\psi}](\tau) + B_q[T \check{\psi}](\tau_1, \tau_2) \lesssim & E_q[T \check{\psi}](\tau_1) + \check{J}_q^1[\check{\psi}, N](\tau_1, \tau_2) + \epsilon B_q^1[\check{\psi}](\tau_1, \tau_2) \\
 & + E_{\max(q, \delta)}^2[\psi](\tau_1) + J_{\max(q, \delta)}^2[\psi, N]. \quad (10.4.12)
 \end{aligned}$$

Recovering estimates for $r \not\!d_2 \check{\psi}$

Recall from Proposition 10.3.1 that $\check{\psi} = f_2 \check{e}_4 \psi$ satisfies

$$\square \check{\psi} - V \check{\psi} = \frac{2}{r} \left(1 - \frac{3m}{r}\right) e_4 \check{\psi} + \check{N} + f_2 \left(e_4 + \frac{3}{r}\right) N.$$

Recall also from Corollary 10.4.5 that we have

$$r \not\!d_2(\square_2 \check{\psi}) - (\square_1 - K)(r \not\!d_2 \check{\psi}) = -r\eta \square_2 \check{\psi} + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \check{\psi}$$

We infer

$$\square_1(r \not\!d_2 \check{\psi}) + (V - K)r \not\!d_2 \check{\psi} = \frac{2}{r} \left(1 - \frac{3m}{r}\right) e_4(r \not\!d_2 \check{\psi}) + N^{r \not\!d_2} + (r \not\!d_2 - r\eta) \left(f_2 \left(e_4 + \frac{3}{r}\right) N\right)$$

where

$$N^{r \not\!d_2} = -r\eta \check{N} + r \not\!d_2(\check{N}) + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^{\leq 2} \check{\psi}.$$

In view of (10.4.5) with $r \not\!d_2 \check{\psi}$ instead of $\check{\psi}$ and with

$$N^{r \not\!d_2} + (r \not\!d_2 - r\eta) \left(f_2 \left(e_4 + \frac{3}{r}\right) N\right)$$

instead of $\check{N} + f_2 \left(e_4 + \frac{3}{r}\right) N$, we deduce

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_q[r \not\!d_2 \check{\psi}](\tau) + B_q[r \not\!d_2 \check{\psi}](\tau_1, \tau_2) \\ & \lesssim E_q[r \not\!d_2 \check{\psi}](\tau_1) + \check{J}_q \left[r \not\!d_2 \check{\psi}, N^{r \not\!d_2} + (r \not\!d_2 - r\eta) \left(f_2 \left(e_4 + \frac{3}{r}\right) N\right) \right] (\tau_1, \tau_2) \\ & \quad + E_{\max(q, \delta)}^1[r \not\!d_2 \psi](\tau_1) + J_{\max(q, \delta)}^1[r \not\!d_2 \psi, r \not\!d_2 N] \\ & \quad + \int_{\mathcal{M}(\tau_1, \tau_2)} r^{q-3} |r \not\!d_2 \check{\psi}| (|r \not\!d_2 \check{\psi}| + |\mathfrak{d}(r \not\!d_2 \check{\psi})|) \\ & \lesssim E_q[r \not\!d_2 \check{\psi}](\tau_1) + \check{J}_q [r \not\!d_2 \check{\psi}, N^{r \not\!d_2}] (\tau_1, \tau_2) \\ & \quad + E_{\max(q, \delta)}^2[\psi](\tau_1) + J_{\max(q, \delta)}^2[\psi, N] + \left(B_q[r \not\!d_2 \check{\psi}](\tau_1, \tau_2)\right)^{\frac{1}{2}} \left(B_q[\check{\psi}](\tau_1, \tau_2)\right)^{\frac{1}{2}} \end{aligned}$$

so that it remains to estimate

$$\check{J}_q [r \not\!d_2 \check{\psi}, N^{r \not\!d_2}] (\tau_1, \tau_2) = J_{q, 4m_0} [r \not\!d_2 \check{\psi}, N^{r \not\!d_2}] (\tau_1, \tau_2) = \left| \int_{\mathcal{M}_{\geq 4m_0}(\tau_1, \tau_2)} r^q (\check{e}_4(r \not\!d_2 \check{\psi})) N^{r \not\!d_2} \right|.$$

The estimate follows along the same lines as the above one for $J_q[T\check{\psi}, N^T](\tau_1, \tau_2)$ so we leave the details to the reader. In the end, we arrive at the following analog of (10.4.12)

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_q[r\check{\phi}_2\check{\psi}](\tau) + B_q[r\check{\phi}_2\check{\psi}](\tau_1, \tau_2) \\ \lesssim & E_q[r\check{\phi}_2\check{\psi}](\tau_1) + B_q[\check{\psi}](\tau_1, \tau_2) + \check{J}_q^1[\check{\psi}, N](\tau_1, \tau_2) + \epsilon B_q^1[\check{\psi}](\tau_1, \tau_2) \\ & + E_{\max(q, \delta)}^2[\psi](\tau_1) + J_{\max(q, \delta)}^2[\psi, N]. \end{aligned} \tag{10.4.13}$$

Recovering estimates for $re_4\check{\psi}$

Recall from Proposition 10.3.1 that $\check{\psi} = f_2\check{e}_4\psi$ satisfies

$$\square\check{\psi} - V\check{\psi} = \frac{2}{r} \left(1 - \frac{3m}{r}\right) e_4\check{\psi} + \check{N} + f_2 \left(e_4 + \frac{3}{r}\right) N.$$

Recall also from Lemma 10.4.10 that we have

$$[\square_2, re_4]\check{\psi} = \frac{\Upsilon}{r} \left(1 + \frac{2m}{r\Upsilon^2}\right) \check{e}_4(re_4\check{\psi}) + \square_2\check{\psi} + \Gamma_g\check{\mathfrak{d}}^2\check{\psi} + \frac{1}{\Upsilon}r^{-2}\check{\mathfrak{d}}T\check{\psi} + r^{-2}\check{\mathfrak{d}}^2\check{\psi} + r^{-2}\check{\mathfrak{d}}\check{\psi}.$$

We infer²³

$$\begin{aligned} \square(re_4\check{\psi}) - Vre_4(\check{\psi}) &= \left(\frac{2}{r} \left(1 - \frac{3m}{r}\right) + \frac{\Upsilon}{r} \left(1 + \frac{2m}{r\Upsilon^2}\right)\right) e_4(re_4\check{\psi}) + N^{re_4} \\ &\quad + re_4 \left(f_2 \left(e_4 + \frac{3}{r}\right) N\right), \end{aligned}$$

where

$$N^{re_4} = re_4(\check{N}) + \check{N} + \Gamma_g\check{\mathfrak{d}}^2\check{\psi} + \frac{1}{\Upsilon}r^{-2}\check{\mathfrak{d}}T\check{\psi} + r^{-2}\check{\mathfrak{d}}^2\check{\psi} + r^{-2}\check{\mathfrak{d}}\check{\psi}.$$

The rest follows along the same lines as the estimate for $T\check{\psi}$ and we arrive at the following analog of (10.4.12)

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_q[re_4\check{\psi}](\tau) + B_q[re_4\check{\psi}](\tau_1, \tau_2) \tag{10.4.14} \\ \lesssim & E_q[re_4\check{\psi}](\tau_1) + B_q[\check{\psi}](\tau_1, \tau_2) + B_q[T\check{\psi}](\tau_1, \tau_2) + B_q[r\check{\phi}_2\check{\psi}](\tau_1, \tau_2) \\ & + \check{J}_q^1[\check{\psi}, N](\tau_1, \tau_2) + \epsilon B_q^1[\check{\psi}](\tau_1, \tau_2) + E_{\max(q, \delta)}^2[\psi](\tau_1) + J_{\max(q, \delta)}^2[\psi, N]. \end{aligned}$$

²³Notice that the coefficient in front of the term $e_4(re_4\check{\psi})$ in the RHS of the wave equation for $re_4\check{\psi}$ differs from the one in front of the term $e_4\check{\psi}$ in the RHS of the wave equation for $\check{\psi}$. Nevertheless, we may apply (10.4.2) with $re_4\check{\psi}$ instead of $\check{\psi}$ since the only property of this coefficient which is used in that it is positive on $r \geq 4m_0$, i.e.

$$\frac{2}{r} \left(1 - \frac{3m}{r}\right) + \frac{\Upsilon}{r} \left(1 + \frac{2m}{r\Upsilon^2}\right) \geq 0 \text{ on } r \geq 4m_0.$$

Conclusion of the proof of Theorem 5.3.5

Gathering the estimates (10.4.12), (10.4.13) and (10.4.14), we infer for any $-1 + \delta < q \leq 1 - \delta$,

$$\begin{aligned} \sup_{\tau \in [\tau_1, \tau_2]} E_q^1[\check{\psi}](\tau) + B_q^1[\check{\psi}](\tau_1, \tau_2) &\lesssim E_q^1[\check{\psi}](\tau_1) + \check{J}_q^1[\check{\psi}, N](\tau_1, \tau_2) + B_q[\check{\psi}](\tau_1, \tau_2) \\ &\quad + \epsilon B_q^1[\check{\psi}](\tau_1, \tau_2) + E_{\max(q, \delta)}^2[\psi](\tau_1) + J_{\max(q, \delta)}^2[\psi, N] \end{aligned}$$

and hence

$$\begin{aligned} \sup_{\tau \in [\tau_1, \tau_2]} E_q^1[\check{\psi}](\tau) + B_q^1[\check{\psi}](\tau_1, \tau_2) &\lesssim E_q^1[\check{\psi}](\tau_1) + \check{J}_q^1[\check{\psi}, N](\tau_1, \tau_2) + B_q[\check{\psi}](\tau_1, \tau_2) \\ &\quad + E_{\max(q, \delta)}^2[\psi](\tau_1) + J_{\max(q, \delta)}^2[\psi, N]. \end{aligned}$$

In view of (10.4.2), we deduce

$$\begin{aligned} \sup_{\tau \in [\tau_1, \tau_2]} E_q^1[\check{\psi}](\tau) + B_q^1[\check{\psi}](\tau_1, \tau_2) &\lesssim E_q^1[\check{\psi}](\tau_1) + \check{J}_q^1[\check{\psi}, N](\tau_1, \tau_2) \\ &\quad + E_{\max(q, \delta)}^2[\psi](\tau_1) + J_{\max(q, \delta)}^2[\psi, N] \end{aligned}$$

which is Theorem 5.3.5 in the case $s = 1$. We have thus deduced Theorem 5.3.5 in the case $s = 1$ from the case $s = 0$, i.e. (10.4.2). Since going from $s = 0$ to $s = 1$ is analogous to going from s to $s + 1$, higher order derivatives $k \leq k_{small} + 29$ are recovered in the same fashion. This concludes the proof of Theorem 5.3.5.

10.5 More weighted estimates for wave equations

The goal of this section is to derive Theorem 10.5.2 and Proposition 10.5.4, see below, which is needed for the proof of Theorem M8 in Chapter 8. Recall that we have used so far in Chapter 10 the global frame of Proposition 3.5.5. For this last section of Chapter 10, we rely instead on the global frame used in Theorem M8, i.e. the one of Proposition 3.5.2, as it is more regular and allows us to derive estimates for up to k_{large} derivatives.

Remark 10.5.1. *Recall that in the frame of Proposition 3.5.2, we only have²⁴ $\eta \in \Gamma_b$. Note that the assumptions on the frame used in Chapter 10 are all consistent with $\eta \in \Gamma_b$, so that all results in this chapter apply for the frame of Proposition 3.5.2.*

²⁴Unlike the frame of Proposition 3.5.5 for which $\eta \in \Gamma_g$.

Theorem 10.5.2. *Let ψ a reduced 2-scalar, and ϕ a reduced 0-scalar satisfying respectively*

$$(\square_2 + V_2)\psi = N_2, \quad (\square_0 + V_0)\phi = N_0, \quad V_2 = -\frac{2}{r^2} \left(1 + \frac{2m}{r}\right), \quad V_0 = \frac{8m}{r^3}.$$

Also, assume that the Ricci coefficients and curvature components associated to the global null frame we are using satisfy the estimates of section 10.4.1 for $k \leq k_{small}$ derivatives. Then, for any $1 \leq s \leq k_{large} - 1$, we have

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_\delta^s[\psi](\tau) + B_\delta^s[\psi](\tau_1, \tau_2) + F_\delta^s[\psi](\tau_1, \tau_2) \\ \lesssim & E_\delta^s[\psi](\tau_1) + \sup_{\tau \in [\tau_1, \tau_2]} E_\delta^{s-1}[\psi](\tau) + B_\delta^{s-1}[\psi](\tau_1, \tau_2) + F_\delta^{s-1}[\psi](\tau_1, \tau_2) \\ & + D_s[\Gamma] \left(\sup_{\mathcal{M}(\tau_1, \tau_2)} r u_{trap}^{\frac{1}{2} + \delta_{dec}} |\mathfrak{d}^{\leq k_{small}} \psi| \right)^2 \\ & + \int_{\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} |\mathfrak{d}^{\leq s} N_2|^2 + \left| \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} T(\mathfrak{d}^s \phi) \mathfrak{d}^s N_2 \right| \end{aligned} \tag{10.5.1}$$

and

$$\begin{aligned} & \sup_{\tau \in [\tau_1, \tau_2]} E_\delta^s[\phi](\tau) + B_\delta^s[\phi](\tau_1, \tau_2) + F_\delta^s[\phi](\tau_1, \tau_2) \\ \lesssim & E_\delta^s[\phi](\tau_1) + \sup_{\tau \in [\tau_1, \tau_2]} E_\delta^{s-1}[\phi](\tau) + B_\delta^{s-1}[\phi](\tau_1, \tau_2) + F_\delta^{s-1}[\phi](\tau_1, \tau_2) \\ & + D_s[\Gamma] \left(\sup_{\mathcal{M}(\tau_1, \tau_2)} r u_{trap}^{\frac{1}{2} + \delta_{dec}} |\mathfrak{d}^{\leq k_{small}} \phi| \right)^2 + \int_{\Sigma(\tau_2)} \frac{(\mathfrak{d}^{\leq s} \phi)^2}{r^3} \\ & + \int_{\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} |\mathfrak{d}^{\leq s} N_0|^2 + \left| \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} T(\mathfrak{d}^s \phi) \mathfrak{d}^s N_0 \right|, \end{aligned} \tag{10.5.2}$$

where $D_s[\Gamma]$ is defined by

$$\begin{aligned} D_s[\Gamma] & := \int_{(int)\mathcal{M} \cup (ext)\mathcal{M}(r \leq 4m_0)} (\mathfrak{d}^{\leq s} \check{\Gamma})^2 \\ & + \sup_{r_0 \geq 4m_0} \left(r_0 \int_{\{r=r_0\}} |\mathfrak{d}^{\leq s} \Gamma_g|^2 + r_0^{-1} \int_{\{r=r_0\}} |\mathfrak{d}^{\leq s} \Gamma_b|^2 \right). \end{aligned}$$

The proof of Theorem 10.5.2 relies on the following theorem.

Theorem 10.5.3. *Let ψ a reduced scalar, and ϕ a reduced 0-scalar satisfying respectively*

$$(\square_2 + V_2)\psi = N_2, \quad (\square_0 + V_0)\phi = N_0, \quad V_2 = -\frac{2}{r^2} \left(1 + \frac{2m}{r}\right), \quad V_0 = \frac{8m}{r^3}.$$

Then, we have

$$\begin{aligned}
& \sup_{\tau \in [\tau_1, \tau_2]} E_\delta[\psi](\tau) + B_\delta[\psi](\tau_1, \tau_2) + F_\delta[\psi](\tau_1, \tau_2) \\
\lesssim & E_\delta[\psi](\tau_1) + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} \left| 1 - \frac{3m}{r} \right| |\psi|(|\psi| + |R\psi|) + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{\delta-3} |\psi|(|\psi| + |\mathfrak{d}\psi|) \\
& + \int_{\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} |N_2|^2 + \left| \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} T(\psi)N_2 \right|, \tag{10.5.3}
\end{aligned}$$

and

$$\begin{aligned}
& \sup_{\tau \in [\tau_1, \tau_2]} E_\delta[\phi](\tau) + B_\delta[\phi](\tau_1, \tau_2) + F_\delta[\phi](\tau_1, \tau_2) \\
\lesssim & E_\delta[\phi](\tau_1) + \int_{\mathcal{A}(\tau_1, \tau_2) \cup \Sigma(\tau_2) \cup \Sigma_*(\tau_1, \tau_2)} \frac{\phi^2}{r^3} \tag{10.5.4} \\
& + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} \left| 1 - \frac{3m}{r} \right| |\phi|(|\phi| + |R\phi|) + \int_{(trq/p)\mathcal{M}(\tau_1, \tau_2)} r^{\delta-3} |\phi|(|\phi| + |\mathfrak{d}\phi|) \\
& + \int_{\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} |N_0|^2 + \left| \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} T(\phi)N_0 \right|.
\end{aligned}$$

Proof. The proof of Theorem 10.5.3 is analogous to the one of Theorem 10.4.11. The only differences are

- The treatment of the right-hand sides N_0 and N_2 in the spacetime region $(trap)\mathcal{M}$.
- The boundary term on $\mathcal{A}(\tau_1, \tau_2) \cup \Sigma(\tau_2) \cup \Sigma_*(\tau_1, \tau_2)$ appearing in the right-hand side of²⁵ (10.5.4).

The treatment of N_0 and N_2 is similar, so we focus on the one of N_2 . The only estimate in which N_2 appear in the trapping region is the Morawetz estimate. More precisely, it appear under the form, see (10.1.71),

$$\left| \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} \left(f_\delta R(\Psi) + \Lambda T(\Psi) + \frac{1}{2} w \Psi \right) N_2 \right|,$$

²⁵This boundary term, as discussed below, is due to the fact that V_0 is positive, which explains why no such term is present in (10.5.3) due to the negativity of the potential V_2 for the wave equation satisfied by ψ .

where we recall that Λ is a constant, and $f_{\widehat{\delta}}, w$ are functions which are in particular bounded on $^{(trap)}\mathcal{M}$. We infer

$$\begin{aligned} & \left| \int_{^{(trap)}\mathcal{M}(\tau_1, \tau_2)} \left(f_{\widehat{\delta}} R(\Psi) + \Lambda T(\Psi) + \frac{1}{2} w \Psi \right) N_2 \right| \\ & \lesssim \int_{^{(trap)}\mathcal{M}(\tau_1, \tau_2)} (|R(\Psi)| + |\Psi|) |N_2| + \left| \int_{^{(trap)}\mathcal{M}(\tau_1, \tau_2)} T(\Psi) N_2 \right| \\ & \lesssim \lambda B_{\delta}[\psi](\tau_1, \tau_2) + \lambda^{-1} \int_{^{(trap)}\mathcal{M}(\tau_1, \tau_2)} |N_2|^2 + \left| \int_{^{(trap)}\mathcal{M}(\tau_1, \tau_2)} T(\Psi) N_2 \right| \end{aligned}$$

which yields the desired control provided $\lambda > 0$ is chosen small enough so that the term $\lambda B_{\delta}[\psi](\tau_1, \tau_2)$ can be absorbed by the LHS in (10.5.3).

Concerning the boundary terms on $\mathcal{A}(\tau_1, \tau_2) \cup \Sigma(\tau_2) \cup \Sigma_*(\tau_1, \tau_2)$ appearing in the right-hand side of (10.5.4), the potential V_0 does not appear in the boundary term of the r^p weighted estimates, but it does appear in the boundary term of the energy estimates²⁶. More precisely, it appears in

$$\int_{\mathcal{A}(\tau_1, \tau_2) \cup \Sigma(\tau_2) \cup \Sigma_*(\tau_1, \tau_2)} Q_{34} = \int_{\mathcal{A}(\tau_1, \tau_2) \cup \Sigma(\tau_2) \cup \Sigma_*(\tau_1, \tau_2)} (|\nabla \phi|^2 + V_0 \phi^2).$$

Now, we have in view of the definition of V_0

$$\int_{\mathcal{A}(\tau_1, \tau_2) \cup \Sigma(\tau_2) \cup \Sigma_*(\tau_1, \tau_2)} Q_{34} \geq \int_{\mathcal{A}(\tau_1, \tau_2) \cup \Sigma(\tau_2) \cup \Sigma_*(\tau_1, \tau_2)} |\nabla \phi|^2 - O(1) \int_{\mathcal{A}(\tau_1, \tau_2) \cup \Sigma(\tau_2) \cup \Sigma_*(\tau_1, \tau_2)} \frac{\phi^2}{r^3}$$

and the control of the boundary terms follows. This concludes the proof of 10.5.3. \square

We are now in position to prove Theorem 10.5.2. Note first that we have

$$\int_{\mathcal{A}(\tau_1, \tau_2) \cup \Sigma_*(\tau_1, \tau_2)} \frac{(\mathfrak{d}^{\leq s} \phi)^2}{r^3} \lesssim F_{\delta}^{s-1}[\phi](\tau_1, \tau_2)$$

which explains why the term $\int_{\mathcal{A}(\tau_1, \tau_2) \cup \Sigma_*(\tau_1, \tau_2)} \frac{(\mathfrak{d}^{\leq s} \phi)^2}{r^3}$, that one would a priori would expect in view of (10.5.4), is not present on the right-hand side of (10.5.2). Also, the estimates for ψ and ϕ are similar, so we focus on the estimate for ψ .

Proof of Theorem 10.5.2. The proof of Theorem 10.5.2 follows along the same lines as the one of Theorem 5.3.4. More precisely, following the strategy in section 10.4.2, we recover

²⁶The boundary term of the r^p weighted estimates involves only $Q_{44} = (e_4 \phi)^2$, while the one of the energy estimate involves also $Q_{34} = |\nabla \phi|^2 + V_0 \phi^2$.

derivatives one by one starting from Theorem 10.5.3 and use it iteratively in conjunction with the commutator estimates of section 10.4.3. The only difference is the treatment of the derivatives for $s \geq k_{small} + 1$ as we assume that the estimates of section 10.4.1 for the Ricci coefficients and curvature components only hold for $k \leq k_{small}$ derivatives. Thus, to conclude, we need to consider the terms for which at least $k_{small} + 1$ derivatives fall on the Ricci coefficients and curvature components. Since on the other hand we have $s \leq k_{large} - 1$, in view of the definition (3.3.7) of k_{small} in terms of k_{large} , and in view of the commutator estimates of section 10.4.3, one easily checks that these terms are bounded in absolute value from above by

$$\left(|\mathfrak{d}^{\leq s}(\Gamma_g)| + r^{-1} |\mathfrak{d}^{\leq s}(\Gamma_b)| \right) |\mathfrak{d}^{\leq k_{small}} \psi|.$$

We thus need, in view of Theorem 10.5.3, to estimate

$$\begin{aligned} & \int_{\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} \left(|\mathfrak{d}^{\leq s}(\Gamma_g)| + r^{-1} |\mathfrak{d}^{\leq s}(\Gamma_b)| \right)^2 |\mathfrak{d}^{\leq k_{small}} \psi|^2 \\ & + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} |T\mathfrak{d}^s \psi| |\mathfrak{d}^{\leq s}(\check{\Gamma})| |\mathfrak{d}^{\leq k_{small}} \psi| \\ \lesssim & \sup_{\mathcal{M}(\tau_1, \tau_2)} \left(r^2 |\mathfrak{d}^{\leq k_{small}} \psi|^2 \right) \int_{\mathcal{M}(\tau_1, \tau_2)} r^{-1+\delta} \left(|\mathfrak{d}^{\leq s}(\Gamma_g)| + r^{-1} |\mathfrak{d}^{\leq s}(\Gamma_b)| \right)^2 \\ & + \left(\sup_{(trap)\mathcal{M}(\tau_1, \tau_2)} u^{\frac{1}{2}+\delta_{dec}} |\mathfrak{d}^{\leq k_{small}} \psi| \right) \left(\sup_{\tau \in [\tau_1, \tau_2]} E_\delta^s[\psi](\tau) \right)^{\frac{1}{2}} \left(\int_{(trap)\mathcal{M}(\tau_1, \tau_2)} |\mathfrak{d}^{\leq s} \check{\Gamma}|^2 \right)^{\frac{1}{2}} \\ \lesssim & \left(\sup_{\mathcal{M}(\tau_1, \tau_2)} r u_{trap}^{\frac{1}{2}+\delta_{dec}} |\mathfrak{d}^{\leq k_{small}} \psi| \right)^2 D_s[\Gamma] \\ & + \left(\sup_{\mathcal{M}(\tau_1, \tau_2)} r u_{trap}^{\frac{1}{2}+\delta_{dec}} |\mathfrak{d}^{\leq k_{small}} \psi| \right) \left(\sup_{\tau \in [\tau_1, \tau_2]} E_\delta^s[\psi](\tau) \right)^{\frac{1}{2}} \sqrt{D_s[\Gamma]} \end{aligned}$$

where we have used the definition of $D_s[\Gamma]$. We infer

$$\begin{aligned} & \int_{\mathcal{M}(\tau_1, \tau_2)} r^{1+\delta} \left(|\mathfrak{d}^{\leq s}(\Gamma_g)| + r^{-1} |\mathfrak{d}^{\leq s}(\Gamma_b)| \right)^2 |\mathfrak{d}^{\leq k_{small}} \psi|^2 \\ & + \int_{(trap)\mathcal{M}(\tau_1, \tau_2)} |T\mathfrak{d}^s \psi| |\mathfrak{d}^{\leq s}(\check{\Gamma})| |\mathfrak{d}^{\leq k_{small}} \psi| \\ \lesssim & \lambda^{-1} \left(\sup_{\mathcal{M}(\tau_1, \tau_2)} r u_{trap}^{\frac{1}{2}+\delta_{dec}} |\mathfrak{d}^{\leq k_{small}} \psi| \right)^2 D_s[\Gamma] + \lambda \sup_{\tau \in [\tau_1, \tau_2]} E_\delta^s[\psi](\tau) \end{aligned}$$

for any $\lambda > 0$ and the last term is then absorbed from the left-hand side of the desired estimate by choosing $\lambda > 0$ small enough which concludes the proof of Theorem 10.5.2. \square

Proposition 10.5.4. *Let ψ a reduced 2-scalar satisfying*

$$\square_2 \psi = f_2(r, m)Y_{(0)}\psi + \tilde{N}_2,$$

where the function f_2 is smooth and positive, and where the vectorfield $Y_{(0)}$ has been introduced in Proposition 10.1.29 in connection with the redshift vectorfield and is given by

$$Y_{(0)} := \left(1 + \frac{5}{4m}(r - 2m) + \Upsilon\right) e_3 + \left(1 + \frac{5}{4m}(r - 2m)\right) e_4.$$

Also, assume that the Ricci coefficients and curvature components associated to the global null frame we are using satisfy the estimates of section 10.4.1 for $k \leq k_{small}$ derivatives. Then, for any $1 \leq s \leq k_{large} - 1$, we have

$$\begin{aligned} \int_{(int)\mathcal{M}(\tau_1, \tau_2)} (\mathfrak{d}^{s+1}\psi)^2 &\lesssim E_\delta^s[\psi](\tau_1) + \int_{(ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}(\tau_1, \tau_2)} (\mathfrak{d}^{s+1}\psi)^2 \\ &\quad + D_s[\Gamma] \left(\sup_{(int)\mathcal{M}(\tau_1, \tau_2) \cup (ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}} r |\mathfrak{d}^{\leq k_{small}} \psi| \right)^2 \\ &\quad + \int_{(int)\mathcal{M}(\tau_1, \tau_2) \cup (ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}} \left((\mathfrak{d}^{\leq s}\psi)^2 + (\mathfrak{d}^{\leq s+1}\tilde{N}_2)^2 \right). \end{aligned}$$

Proof. Recall from Proposition 10.1.29 that the redshift vectorfield is given by

$$Y_{\mathcal{H}} := \kappa_{\mathcal{H}} Y_{(0)}, \quad \kappa_{\mathcal{H}} := \kappa \left(\frac{\Upsilon}{\delta_{\frac{1}{10}\mathcal{H}}} \right),$$

where κ is a positive bump function $\kappa = \kappa(r)$, supported in the region in $[-2, 2]$ and equal to 1 for $[-1, 1]$.

To estimate ψ in $(int)\mathcal{M}$, we consider

$$\tilde{\psi} := \tilde{\kappa} \left(\frac{r - 2m_0(1 + 2\delta_{\mathcal{H}})}{2m_0\delta_{\mathcal{H}}} \right)$$

where $\tilde{\kappa}$ is a positive bump function $\kappa = \kappa(r)$, supported in the region in $(-\infty, 1]$ and equal to 1 for $(-\infty, 0]$. Since $(int)\mathcal{M}$ is included in $r \leq 2m_0(1 + 2\delta_{\mathcal{H}})$, we infer in particular

$$\tilde{\psi} = \psi \text{ on } (int)\mathcal{M}, \quad \text{supp}(\tilde{\psi}) \subset (int)\mathcal{M}(\tau_1, \tau_2) \cup (ext)\mathcal{M}_{r \leq 2m_0(1+3\delta_{\mathcal{H}})}.$$

Also, we have, in view of the wave equation for ψ ,

$$\square_2 \tilde{\psi} = f_2(r, m) Y_{(0)} \tilde{\psi} + \tilde{N}'_2$$

where \tilde{N}'_2 satisfies

$$\begin{aligned} \int_{(int)\mathcal{M}(\tau_1, \tau_2) \cup (ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}} (\mathfrak{d}^{\leq s+1} \tilde{N}'_2)^2 &\lesssim \int_{(ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}(\tau_1, \tau_2)} (\mathfrak{d}^{\leq s+1} \psi)^2 \\ &+ \int_{(int)\mathcal{M}(\tau_1, \tau_2) \cup (ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}} (\mathfrak{d}^{\leq s+1} \tilde{N}'_2)^2. \end{aligned}$$

Since $\tilde{\psi} = \psi$ on $(int)\mathcal{M}$, it thus suffices to prove for $\tilde{\psi}$ the following estimate

$$\begin{aligned} \int_{\mathcal{M}(\tau_1, \tau_2)} (\mathfrak{d}^{s+1} \tilde{\psi})^2 &\lesssim E_\delta^s[\tilde{\psi}](\tau_1) + D_s[\Gamma] \left(\sup_{(int)\mathcal{M}(\tau_1, \tau_2) \cup (ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}} r |\mathfrak{d}^{\leq k_{small}} \tilde{\psi}| \right)^2 \\ &+ \int_{(int)\mathcal{M}(\tau_1, \tau_2) \cup (ext)\mathcal{M}_{r \leq \frac{5}{2}m_0}} \left((\mathfrak{d}^{\leq s} \tilde{\psi})^2 + (\mathfrak{d}^{\leq s+1} \tilde{N}'_2)^2 \right). \end{aligned}$$

This estimate follows from first deriving the corresponding estimate for $s = 0$ by using the redshift as a multiplier, and then by recover derivatives one by one using commutation with T , $\not\partial$ and the redshift vectorfield. Note that

- $\tilde{\psi}$ is supported on $r \leq 2m_0(1 + 2\delta_{\mathcal{H}})$ and hence is estimated on

$$2m_0(1 - 2\delta_{\mathcal{H}}) \leq r \leq 2m_0(1 + 2\delta_{\mathcal{H}})$$

so that the redshift vectorfield $Y_{\mathcal{H}}$ has good properties, both as a multiplier and as a commutator, on the support of $\tilde{\psi}$.

- The term $f_2(r, m)Y_{(0)}$ yields a good sign when using $Y_{\mathcal{H}}$ as a multiplier since the function $f_2(r, m)$ is positive, and since $Y_{\mathcal{H}} = \kappa_{\mathcal{H}}Y_{(0)}$.

This concludes the proof of the proposition. □

Appendix A

APPENDIX TO CHAPTER 2

A.1 Proof of Proposition 2.2.9

In a neighborhood of a given sphere S , we consider a (u, s, θ, φ) coordinates system, where θ is such that $e_4(\theta) = 0$. Then, in this coordinates system, we have

$$\partial_s = e_4.$$

Since we have

$$\partial_s \left(\int_S f \right) = \int_S \left(\partial_s f + g(\mathbf{D}_{e_\theta} \partial_s, e_\theta) f + g(\mathbf{D}_{e_\varphi} \partial_s, e_\varphi) f \right),$$

we infer

$$e_4 \left(\int_S f \right) = \int_S (e_4(f) + \kappa f).$$

In particular, choosing $f = 1$, we deduce

$$\frac{1}{|S|} e_4(|S|) = \bar{\kappa}$$

and since $|S| = 4\pi r^2$,

$$e_4(r) = \frac{r\bar{\kappa}}{2}.$$

Next, let ∂_u the coordinates vectorfield in the (u, s, θ, φ) coordinates system. We have

$$\begin{aligned} \partial_u \left(\int_S f \right) &= \int_S \left(\partial_u f + g(\mathbf{D}_{e_\theta} \partial_u, e_\theta) f + g(\mathbf{D}_{e_\varphi} \partial_u, e_\varphi) f \right) \\ &= \int_S \left(\partial_u f + g(\mathbf{D}_{e_\theta} \partial_u, e_\theta) f - g(\partial_u, \mathbf{D}_{e_\varphi} e_\varphi) f \right) \\ &= \int_S \left(\partial_u f + g(\mathbf{D}_{e_\theta} \partial_u, e_\theta) f + g(\partial_u, D^a(\Phi) e_a) f \right) \end{aligned}$$

On the other hand, we have we have, see (2.2.42),

$$\partial_u = \varsigma \left(\frac{1}{2} e_3 - \frac{1}{2} \underline{\Omega} e_4 - \frac{1}{2} \sqrt{\gamma} \underline{b} e_\theta \right).$$

We infer

$$g(D_{e_\theta} \partial_u, e_\theta) + g(\partial_u, D^a(\Phi) e_a) = \frac{1}{2} \varsigma \underline{\kappa} - \frac{1}{2} \varsigma \underline{\Omega} \kappa - \frac{1}{2} \not\!d_1(\varsigma \sqrt{\gamma} \underline{b})$$

and thus

$$\begin{aligned} \varsigma \left(\frac{1}{2} e_3 - \frac{1}{2} \underline{\Omega} e_4 \right) \left(\int_S f \right) &= \int_S \left(\varsigma \left(\frac{1}{2} e_3 - \frac{1}{2} \underline{\Omega} e_4 - \frac{1}{2} \sqrt{\gamma} \underline{b} e_\theta \right) f + \frac{1}{2} \varsigma \underline{\kappa} f - \frac{1}{2} \varsigma \underline{\Omega} \kappa f \right. \\ &\quad \left. - \frac{1}{2} \not\!d_1(\sqrt{\gamma} \underline{b}) f \right). \end{aligned}$$

We deduce

$$e_3 \left(\int_S f \right) = \underline{\Omega} e_4 \left(\int_S f \right) + \varsigma^{-1} \int_S \left(\varsigma e_3 f - \varsigma \underline{\Omega} e_4 f + \varsigma \underline{\kappa} f - \varsigma \underline{\Omega} \kappa f - \not\!d_1(\varsigma \sqrt{\gamma} \underline{b}) f \right).$$

Next, we use

$$e_4 \left(\int_S f \right) = \int_S (e_4(f) + \kappa f)$$

and

$$\int_S \not\!d_1(\varsigma \sqrt{\gamma} \underline{b}) f = 0.$$

We infer

$$\begin{aligned} e_3 \left(\int_S f \right) &= \underline{\Omega} \int_S (e_4(f) + \kappa f) + \varsigma^{-1} \int_S \left(\varsigma e_3 f - \varsigma \underline{\Omega} e_4 f + \varsigma \underline{\kappa} f - \varsigma \underline{\Omega} \kappa f \right) \\ &= \varsigma^{-1} \int_S \varsigma (e_3 f + \underline{\kappa} f) + (\check{\underline{\Omega}} + \varsigma^{-1} \overline{\underline{\Omega}} \check{\varsigma}) \int_S (e_4 f + \kappa f) \\ &\quad - \varsigma^{-1} \overline{\underline{\Omega}} \int_S \check{\varsigma} (e_4 f + \kappa f) - \varsigma^{-1} \int_S \check{\underline{\Omega}} \varsigma (e_4 f + \kappa f). \end{aligned}$$

We further write,

$$\begin{aligned} \varsigma^{-1} \int_S \varsigma (e_3 f + \underline{\kappa} f) &= \varsigma^{-1} \overline{\varsigma} \int_S (e_3 f + \underline{\kappa} f) + \varsigma^{-1} \int_S \check{\varsigma} (e_3 f + \underline{\kappa} f) \\ &= \int_S (e_3 f + \underline{\kappa} f) + (\varsigma^{-1} \overline{\varsigma} - 1) \int_S (e_3 f + \underline{\kappa} f) + \varsigma^{-1} \int_S \check{\varsigma} (e_3 f + \underline{\kappa} f) \\ &= \int_S (e_3 f + \underline{\kappa} f) - \varsigma^{-1} \check{\varsigma} \int_S (e_3 f + \underline{\kappa} f) + \varsigma^{-1} \int_S \check{\varsigma} (e_3 f + \underline{\kappa} f). \end{aligned}$$

Hence,

$$\begin{aligned} e_3 \left(\int_S f \right) &= \int_S (e_3 f + \underline{\kappa} f) + \text{Err} \left[e_3 \left(\int_S f \right) \right], \\ \text{Err} \left[e_3 \left(\int_S f \right) \right] &= -\varsigma^{-1} \check{\varsigma} \int_S (e_3 f + \underline{\kappa} f) + \varsigma^{-1} \int_S \check{\varsigma} (e_3 f + \underline{\kappa} f) \\ &\quad + (\check{\underline{\Omega}} + \varsigma^{-1} \overline{\underline{\Omega}} \check{\varsigma}) \int_S (e_4 f + \kappa f) - \varsigma^{-1} \overline{\underline{\Omega}} \int_S \check{\varsigma} (e_4 f + \kappa f) \\ &\quad - \varsigma^{-1} \int_S \check{\underline{\Omega}} \varsigma (e_4 f + \kappa f) \end{aligned}$$

as desired.

In particular, choosing $f = 1$, we infer

$$\begin{aligned} \frac{1}{|S|} e_3(|S|) &= \overline{\underline{\kappa}} - \varsigma^{-1} \check{\varsigma} \overline{\underline{\kappa}} + \varsigma^{-1} \overline{\underline{\kappa}} + (\check{\underline{\Omega}} + \varsigma^{-1} \overline{\underline{\Omega}} \check{\varsigma}) \overline{\underline{\kappa}} - \varsigma^{-1} \overline{\underline{\Omega}} \overline{\underline{\kappa}} - \varsigma^{-1} \overline{\underline{\Omega}} \overline{\underline{\kappa}} \\ &= \overline{\underline{\kappa}} - \varsigma^{-1} \check{\varsigma} \overline{\underline{\kappa}} + \varsigma^{-1} \overline{\underline{\kappa}} + (\check{\underline{\Omega}} + \varsigma^{-1} \overline{\underline{\Omega}} \check{\varsigma}) \overline{\underline{\kappa}} - \varsigma^{-1} \overline{\underline{\Omega}} \overline{\underline{\kappa}} - \varsigma^{-1} \overline{\underline{\Omega}} \overline{\underline{\kappa}}. \end{aligned}$$

Hence, since $|S| = 4\pi r^2$, recalling the definition of \underline{A} ,

$$\begin{aligned} \frac{2e_3(r)}{r} &= \overline{\underline{\kappa}} - \varsigma^{-1} \check{\varsigma} \overline{\underline{\kappa}} + \varsigma^{-1} \overline{\underline{\kappa}} + (\check{\underline{\Omega}} + \varsigma^{-1} \overline{\underline{\Omega}} \check{\varsigma}) \overline{\underline{\kappa}} - \varsigma^{-1} \overline{\underline{\Omega}} \overline{\underline{\kappa}} - \varsigma^{-1} \overline{\underline{\Omega}} \overline{\underline{\kappa}} \\ &= \overline{\underline{\kappa}} + \underline{A}. \end{aligned}$$

This concludes the proof of Proposition 2.2.9.

A.2 Proof of Proposition 2.2.16

We start with the proof for $e_4(m)$. Recall that the Hawking mass m is given by the formula $\frac{2m}{r} = 1 + \frac{1}{16\pi} \int_S \kappa \underline{\kappa}$. Differentiating in the e_4 direction, we deduce,

$$\frac{2e_4(m)}{r} - \frac{2me_4(r)}{r^2} = \frac{1}{16\pi} e_4 \left(\int_S \kappa \underline{\kappa} \right) = \frac{1}{16\pi} \int_S \left(e_4(\kappa \underline{\kappa}) + \underline{\kappa} \kappa^2 \right).$$

Now, making use of the e_4 transport equations of Proposition 2.2.8,

$$\begin{aligned} e_4(\kappa \underline{\kappa}) &= \underline{\kappa} \left(-\frac{1}{2} \kappa^2 - \frac{1}{2} \vartheta^2 \right) + \kappa \left(-\frac{1}{2} \kappa \underline{\kappa} + 2\rho - 2\phi_1 \zeta - \frac{1}{2} \vartheta \underline{\vartheta} + 2\zeta^2 \right) \\ &= -\underline{\kappa} \kappa^2 + 2\kappa \rho - 2\kappa \phi_1 \zeta - \frac{1}{2} \underline{\kappa} \vartheta^2 - \frac{1}{2} \kappa \vartheta \underline{\vartheta} + 2\kappa \zeta^2. \end{aligned}$$

we infer

$$\begin{aligned} \frac{2e_4(m)}{r} - \frac{m}{r} \bar{\kappa} &= \frac{1}{16\pi} \int_S \left(2\kappa \rho - 2\kappa \phi_1 \zeta - \frac{1}{2} \underline{\kappa} \vartheta^2 - \frac{1}{2} \kappa \vartheta \underline{\vartheta} + 2\kappa \zeta^2 \right) \\ &= \frac{1}{8\pi} |S| \bar{\kappa} \bar{\rho} + \frac{1}{16\pi} \int_S \left(2\check{\kappa} \check{\rho} + 2e_\theta(\kappa) \zeta - \frac{1}{2} \underline{\kappa} \vartheta^2 - \frac{1}{2} \kappa \vartheta \underline{\vartheta} + 2\kappa \zeta^2 \right) \\ &= \frac{r^2}{2} \bar{\kappa} \bar{\rho} + \frac{1}{16\pi} \int_S \left(2\check{\kappa} \check{\rho} + 2e_\theta(\kappa) \zeta - \frac{1}{2} \underline{\kappa} \vartheta^2 - \frac{1}{2} \kappa \vartheta \underline{\vartheta} + 2\kappa \zeta^2 \right) \end{aligned}$$

and hence

$$e_4(m) = \frac{r^3}{4} \bar{\kappa} \left(\bar{\rho} + \frac{2m}{r^3} \right) + \frac{r}{32\pi} \int_S \left(-\frac{1}{2} \underline{\kappa} \vartheta^2 - \frac{1}{2} \kappa \vartheta \underline{\vartheta} + 2\check{\kappa} \check{\rho} + 2e_\theta(\kappa) \zeta + 2\kappa \zeta^2 \right).$$

Using the identity $\bar{\rho} = -\frac{2m}{r^3} + \frac{1}{16\pi r^2} \int_S \vartheta \underline{\vartheta}$ (see (2.2.12) of Proposition 2.2.4), we deduce

$$\begin{aligned} e_4(m) &= \frac{r}{32\pi} \int_S \left(-\frac{1}{2} \underline{\kappa} \vartheta^2 - \frac{1}{2} (\kappa - \bar{\kappa}) \vartheta \underline{\vartheta} + 2\check{\kappa} \check{\rho} + 2e_\theta(\kappa) \zeta + 2\kappa \zeta^2 \right) \\ &= \frac{r}{32\pi} \int_S \left(-\frac{1}{2} \underline{\kappa} \vartheta^2 - \frac{1}{2} \check{\kappa} \vartheta \underline{\vartheta} + 2\check{\kappa} \check{\rho} + 2e_\theta(\kappa) \zeta + 2\kappa \zeta^2 \right) \\ &= \frac{r}{32\pi} \int_S \text{Err}_1 \end{aligned}$$

as desired.

In the same vein,

$$\frac{2e_3(m)}{r} - \frac{2me_3(r)}{r^2} = \frac{1}{16\pi} e_3 \left(\int_S \kappa \underline{\kappa} \right) = \frac{1}{16\pi} \int_S \left(e_3(\kappa \underline{\kappa}) + \underline{\kappa} \kappa^2 \right) + E_1,$$

with E_1 the error term defined in Proposition 2.2.9

$$E_1 = \frac{1}{16\pi} \text{Err} \left[e_3 \left(\int_S \underline{\kappa} \underline{\kappa} \right) \right].$$

We make use of the e_3 transport equations of Proposition 2.2.8,

$$\begin{aligned} e_3(\underline{\kappa} \underline{\kappa}) &= \underline{\kappa} \left(-\frac{1}{2} \underline{\kappa} \underline{\kappa} + 2 \underline{\omega} \underline{\kappa} + 2 \underline{\phi}_1 \underline{\eta} + 2 \underline{\rho} - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} + 2 \underline{\eta}^2 \right) \\ &+ \underline{\kappa} \left(-\frac{1}{2} \underline{\kappa}^2 - 2 \underline{\omega} \underline{\kappa} + 2 \underline{\phi}_1 \underline{\xi} + 2(\underline{\eta} - 3\underline{\zeta}) \underline{\xi} - \frac{1}{2} \underline{\vartheta}^2 \right) \\ &= -\underline{\kappa} \underline{\kappa}^2 + 2 \underline{\kappa} \underline{\phi}_1 \underline{\eta} + 2 \underline{\kappa} \underline{\phi}_1 \underline{\xi} + 2 \underline{\rho} \underline{\kappa} + \underline{\kappa} \left(2 \underline{\eta}^2 - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} \right) + 2 \underline{\kappa} (\underline{\eta} - 3 \underline{\zeta}) \underline{\xi} - \frac{1}{2} \underline{\kappa} \underline{\vartheta}^2. \end{aligned}$$

Therefore, setting $E_2 = \frac{1}{16\pi} \int_S \left(\underline{\kappa} (2 \underline{\eta}^2 - \frac{1}{2} \underline{\vartheta} \underline{\vartheta}) + 2 \underline{\kappa} (\underline{\eta} - 3 \underline{\zeta}) \underline{\xi} - \frac{1}{2} \underline{\kappa} \underline{\vartheta}^2 \right)$,

$$\begin{aligned} \frac{2e_3(m)}{r} - \frac{m}{r} (\underline{\bar{\kappa}} + \underline{A}) &= \frac{1}{16\pi} \int_S \left(2 \underline{\kappa} \underline{\phi}_1 \underline{\eta} + 2 \underline{\kappa} \underline{\phi}_1 \underline{\xi} + 2 \underline{\rho} \underline{\kappa} \right) + E_1 + E_2 \\ &= \frac{1}{16\pi} \int_S \left(-2e_\theta(\underline{\kappa}) \underline{\eta} - 2e_\theta(\underline{\kappa}) \underline{\xi} + 2(\underline{\bar{\rho}} + \underline{\check{\rho}}) (\underline{\bar{\kappa}} + \underline{\check{\kappa}}) \right) + E_1 + E_2 \\ &= \frac{1}{2} r^2 \underline{\bar{\rho}} \underline{\bar{\kappa}} + \frac{1}{16\pi} \int_S \left(-2e_\theta(\underline{\kappa}) \underline{\eta} - 2e_\theta(\underline{\kappa}) \underline{\xi} + 2 \underline{\check{\rho}} \underline{\check{\kappa}} \right) + E_1 + E_2 \\ &= \frac{1}{2} r^2 \underline{\bar{\kappa}} \left(-\frac{2m}{r^3} + \frac{1}{16\pi r^2} \int_S \underline{\vartheta} \underline{\vartheta} \right) \\ &+ \frac{1}{16\pi} \int_S \left(-2e_\theta(\underline{\kappa}) \underline{\eta} - 2e_\theta(\underline{\kappa}) \underline{\xi} + 2 \underline{\check{\rho}} \underline{\check{\kappa}} \right) + E_1 + E_2. \end{aligned}$$

We deduce

$$\begin{aligned} \frac{2e_3(m)}{r} &= \frac{1}{16\pi} \int_S \left(-2e_\theta(\underline{\kappa}) \underline{\eta} - 2e_\theta(\underline{\kappa}) \underline{\xi} + 2 \underline{\check{\rho}} \underline{\check{\kappa}} + \frac{1}{2} \underline{\bar{\kappa}} \underline{\vartheta} \underline{\vartheta} \right) + E_1 \\ &+ \frac{1}{16\pi} \int_S \left(\underline{\kappa} \left(2 \underline{\eta}^2 - \frac{1}{2} \underline{\vartheta} \underline{\vartheta} \right) + 2 \underline{\kappa} (\underline{\eta} - 3 \underline{\zeta}) \underline{\xi} - \frac{1}{2} \underline{\kappa} \underline{\vartheta}^2 \right) + \frac{m}{r} \underline{A} \\ &= \frac{1}{16\pi} \int_S \left(-2e_\theta(\underline{\kappa}) \underline{\eta} + 2 \underline{\kappa} \underline{\eta}^2 - 2e_\theta(\underline{\kappa}) \underline{\xi} + 2 \underline{\kappa} \underline{\eta} \underline{\xi} - \frac{1}{2} \underline{\kappa} \underline{\vartheta}^2 \right) \\ &+ \frac{1}{16\pi} \int_S \left(2 \underline{\check{\rho}} \underline{\check{\kappa}} - 6 \underline{\kappa} \underline{\zeta} \underline{\xi} - \frac{1}{2} \underline{\check{\kappa}} \underline{\vartheta} \underline{\vartheta} \right) + E_1 + \frac{m}{r} \underline{A}, \end{aligned}$$

i.e.,

$$\begin{aligned} e_3(m) &= \frac{r}{32\pi} \int_S \left(-2e_\theta(\underline{\kappa}) \underline{\eta} + 2 \underline{\kappa} \underline{\eta}^2 - 2e_\theta(\underline{\kappa}) \underline{\xi} + 2 \underline{\kappa} \underline{\eta} \underline{\xi} - \frac{1}{2} \underline{\kappa} \underline{\vartheta}^2 \right) \\ &+ \frac{r}{32\pi} \int_S \left(2 \underline{\check{\rho}} \underline{\check{\kappa}} - 6 \underline{\kappa} \underline{\zeta} \underline{\xi} - \frac{1}{2} \underline{\check{\kappa}} \underline{\vartheta} \underline{\vartheta} \right) + \frac{r}{2} \left(E_1 + \frac{m}{r} \underline{A} \right). \end{aligned}$$

It remains to calculate $E_1 + \frac{m}{r}\underline{A}$. Using the definitions of E_1 and \underline{A} and grouping similar terms appropriately we find

$$\begin{aligned} E_1 + \frac{m}{r}\underline{A} &= -\varsigma^{-1}\check{\zeta} \left[\frac{1}{16\pi} \int_S (e_3(\kappa\underline{\kappa}) + \kappa\underline{\kappa}^2) + \frac{m}{r}\overline{\underline{\kappa}} \right] + \varsigma^{-1} \left[\frac{1}{16\pi} \int_S \check{\zeta} (e_3(\kappa\underline{\kappa}) + \kappa\underline{\kappa}^2) + \frac{m}{r}\overline{\check{\zeta}\underline{\kappa}} \right] \\ &+ (\check{\underline{\Omega}} + \varsigma^{-1}\overline{\underline{\Omega}}\check{\zeta}) \left[\frac{1}{16\pi} \int_S (e_4(\kappa\underline{\kappa}) + \kappa^2\underline{\kappa}) + \frac{m}{r}\overline{\underline{\kappa}} \right] - \varsigma^{-1}\overline{\underline{\Omega}} \left[\frac{1}{16\pi} \int_S \check{\zeta} (e_4(\kappa\underline{\kappa}) + \kappa^2\underline{\kappa}) + \frac{m}{r}\overline{\check{\zeta}\underline{\kappa}} \right] \\ &- \varsigma^{-1} \left[\frac{1}{16\pi} \int_S \check{\underline{\Omega}}\varsigma (e_4(\kappa\underline{\kappa}) + \kappa^2\underline{\kappa}) + \frac{m}{r}\overline{\check{\underline{\Omega}}\varsigma\underline{\kappa}} \right]. \end{aligned}$$

Now, we have from above calculations

$$\begin{aligned} e_4(\kappa\underline{\kappa}) + \underline{\kappa}\kappa^2 &= 2\kappa\rho - 2\kappa\phi_1\zeta + \text{Err}[e_4(\kappa\underline{\kappa})], \\ \text{Err}[e_4(\kappa\underline{\kappa})] &= -\frac{1}{2}\underline{\kappa}\vartheta^2 - \frac{1}{2}\kappa\vartheta\underline{\vartheta} + 2\kappa\zeta^2, \\ e_3(\kappa\underline{\kappa}) + \underline{\kappa}\kappa^2 &= 2\rho\underline{\kappa} + 2\underline{\kappa}\phi_1\eta + 2\kappa\phi_1\underline{\xi} + \text{Err}[e_3(\kappa\underline{\kappa})], \\ \text{Err}[e_3(\kappa\underline{\kappa})] &= \underline{\kappa} \left(2\eta^2 - \frac{1}{2}\vartheta\underline{\vartheta} \right) + 2\kappa(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\kappa\underline{\vartheta}^2. \end{aligned}$$

We infer

$$\begin{aligned} E_1 + \frac{m}{r}\underline{A} &= -\varsigma^{-1}\check{\zeta} \left[\frac{1}{16\pi} \int_S (2\rho\underline{\kappa} + 2\underline{\kappa}\phi_1\eta + 2\kappa\phi_1\underline{\xi} + \text{Err}[e_3(\kappa\underline{\kappa})]) + \frac{m}{r}\overline{\underline{\kappa}} \right] \\ &+ \varsigma^{-1} \left[\frac{1}{16\pi} \int_S \check{\zeta} (2\rho\underline{\kappa} + 2\underline{\kappa}\phi_1\eta + 2\kappa\phi_1\underline{\xi} + \text{Err}[e_3(\kappa\underline{\kappa})]) + \frac{m}{r}\overline{\check{\zeta}\underline{\kappa}} \right] \\ &+ (\check{\underline{\Omega}} + \varsigma^{-1}\overline{\underline{\Omega}}\check{\zeta}) \left[\frac{1}{16\pi} \int_S (2\kappa\rho - 2\kappa\phi_1\zeta + \text{Err}[e_4(\kappa\underline{\kappa})]) + \frac{m}{r}\overline{\underline{\kappa}} \right] \\ &- \varsigma^{-1}\overline{\underline{\Omega}} \left[\frac{1}{16\pi} \int_S \check{\zeta} (2\kappa\rho - 2\kappa\phi_1\zeta + \text{Err}[e_4(\kappa\underline{\kappa})]) + \frac{m}{r}\overline{\check{\zeta}\underline{\kappa}} \right] \\ &- \varsigma^{-1} \left[\frac{1}{16\pi} \int_S \check{\underline{\Omega}}\varsigma (2\kappa\rho - 2\kappa\phi_1\zeta + \text{Err}[e_4(\kappa\underline{\kappa})]) + \frac{m}{r}\overline{\check{\underline{\Omega}}\varsigma\underline{\kappa}} \right] \end{aligned}$$

and hence

$$\begin{aligned}
 E_1 + \frac{m}{r} \underline{A} &= -\varsigma^{-1} \check{\zeta} \left[\frac{1}{16\pi} \int_S \left(2\check{\rho}\check{\kappa} - 2e_\theta(\underline{\kappa})\eta - 2e_\theta(\kappa)\underline{\xi} + \frac{1}{2}\overline{\kappa}\vartheta\underline{\vartheta} + \text{Err}[e_3(\underline{\kappa}\kappa)] \right) \right] \\
 &+ \varsigma^{-1} \left[\frac{1}{16\pi} \int_S \check{\zeta} (2\overline{\rho}\check{\kappa} + 2\check{\rho}\overline{\kappa} + 2\check{\rho}\check{\kappa} + 2\underline{\kappa}\phi_1\eta + 2\kappa\phi_1\underline{\xi} + \text{Err}[e_3(\underline{\kappa}\kappa)]) + \frac{m}{r} \overline{\check{\zeta}\check{\kappa}} \right] \\
 &+ (\check{\underline{\Omega}} + \varsigma^{-1}\overline{\underline{\Omega}}\check{\zeta}) \left[\frac{1}{16\pi} \int_S \left(2\check{\kappa}\check{\rho} + 2e_\theta(\kappa)\zeta + \frac{1}{2}\overline{\kappa}\vartheta\underline{\vartheta} + \text{Err}[e_4(\underline{\kappa}\kappa)] \right) \right] \\
 &- \varsigma^{-1}\overline{\underline{\Omega}} \left[\frac{1}{16\pi} \int_S \check{\zeta} (2\overline{\rho}\check{\kappa} + 2\check{\rho}\overline{\kappa} + 2\check{\rho}\check{\kappa} - 2\kappa\phi_1\zeta + \text{Err}[e_4(\underline{\kappa}\kappa)]) + \frac{m}{r} \overline{\check{\zeta}\check{\kappa}} \right] \\
 &- \varsigma^{-1} \left[\frac{1}{16\pi} \int_S \check{\underline{\Omega}}\zeta (2\overline{\rho}\check{\kappa} + 2\check{\rho}\overline{\kappa} + 2\check{\rho}\check{\kappa} - 2\kappa\phi_1\zeta + \text{Err}[e_4(\underline{\kappa}\kappa)]) + \frac{m}{r} \overline{\check{\underline{\Omega}}\zeta\kappa} \right].
 \end{aligned}$$

We deduce

$$\begin{aligned}
 e_3(m) &= (1 - \varsigma^{-1}\check{\zeta}) \frac{r}{32\pi} \int_S \underline{\text{Err}}_1 + (\check{\underline{\Omega}} + \varsigma^{-1}\overline{\underline{\Omega}}\check{\zeta}) \frac{r}{32\pi} \int_S \text{Err}_1 \\
 &+ \varsigma^{-1} \frac{r}{32\pi} \int_S \check{\zeta} (2\overline{\rho}\check{\kappa} + 2\check{\rho}\overline{\kappa} + 2\underline{\kappa}\phi_1\eta + 2\kappa\phi_1\underline{\xi} + \underline{\text{Err}}_2) \\
 &- \varsigma^{-1} \frac{r}{32\pi} \int_S (\overline{\underline{\Omega}}\check{\zeta} + \check{\underline{\Omega}}\zeta) (2\overline{\rho}\check{\kappa} + 2\check{\rho}\overline{\kappa} - 2\kappa\phi_1\zeta + \text{Err}_2) \\
 &- \frac{m}{r} \varsigma^{-1} \left[-\overline{\check{\zeta}\check{\kappa}} + \overline{\underline{\Omega}}\overline{\check{\zeta}\check{\kappa}} + \overline{\check{\underline{\Omega}}\zeta\kappa} \right],
 \end{aligned}$$

where we have introduced

$$\begin{aligned}
 \text{Err}_1 &= 2\check{\kappa}\check{\rho} + 2e_\theta(\kappa)\zeta + \frac{1}{2}\overline{\kappa}\vartheta\underline{\vartheta} + \text{Err}[e_4(\underline{\kappa}\kappa)], \\
 \underline{\text{Err}}_1 &= 2\check{\rho}\check{\kappa} - 2e_\theta(\underline{\kappa})\eta - 2e_\theta(\kappa)\underline{\xi} + \frac{1}{2}\overline{\kappa}\vartheta\underline{\vartheta} + \text{Err}[e_3(\underline{\kappa}\kappa)], \\
 \text{Err}_2 &= 2\check{\rho}\check{\kappa} + \text{Err}[e_4(\underline{\kappa}\kappa)], \\
 \underline{\text{Err}}_2 &= 2\check{\rho}\check{\kappa} + \text{Err}[e_3(\underline{\kappa}\kappa)].
 \end{aligned}$$

In view of the definition of $\text{Err}[e_4(\underline{\kappa}\kappa)]$ and $\text{Err}[e_3(\underline{\kappa}\kappa)]$, this concludes the proof of Proposition 2.2.16.

A.3 Proof of Lemma 2.2.17

Recall that we have

$$e_4(\kappa) = -\frac{1}{2}\kappa^2 - \frac{1}{4}\vartheta^2.$$

We infer

$$\begin{aligned} e_4(\bar{\kappa}) &= \overline{e_4(\kappa)} + \bar{\kappa}^2 = \overline{-\frac{1}{2}\kappa^2 - \frac{1}{4}\vartheta^2} + \bar{\kappa}^2 \\ &= -\frac{1}{2}\bar{\kappa}^2 - \frac{1}{4}\bar{\vartheta}^2 + \frac{1}{2}\bar{\kappa}^2 \end{aligned}$$

and hence

$$\begin{aligned} e_4\left(\bar{\kappa} - \frac{2}{r}\right) &= -\frac{1}{2}\bar{\kappa}^2 - \frac{1}{4}\bar{\vartheta}^2 + \frac{1}{2}\bar{\kappa}^2 + \frac{2}{r} \frac{e_4(r)}{r} = -\frac{1}{2}\bar{\kappa}^2 + \frac{1}{r}\bar{\kappa} - \frac{1}{4}\bar{\vartheta}^2 + \frac{1}{2}\bar{\kappa}^2 \\ &= -\frac{1}{2}\bar{\kappa} \left(\bar{\kappa} - \frac{2}{r}\right) - \frac{1}{4}\bar{\vartheta}^2 + \frac{1}{2}\bar{\kappa}^2. \end{aligned}$$

Next, using

$$e_4(\underline{\omega}) = \rho + \zeta(2\eta + \zeta)$$

we infer that

$$e_4(\underline{\omega}) = \overline{e_4(\underline{\omega})} + \bar{\kappa}\bar{\omega} = \bar{\rho} + \overline{\zeta(2\eta + \zeta)} + \bar{\kappa}\bar{\omega},$$

and hence

$$\begin{aligned} e_4\left(\underline{\omega} - \frac{m}{r^2}\right) &= e_4(\underline{\omega}) + \frac{2me_4(r)}{r^3} - \frac{e_4(m)}{r^2} \\ &= \bar{\rho} + \frac{2m}{r^3} + \frac{m}{r^2} \left(\bar{\kappa} - \frac{2}{r}\right) - \frac{e_4(m)}{r^2} + \overline{3\zeta(2\eta + \zeta)} + \bar{\kappa}\bar{\omega} \end{aligned}$$

as stated.

Next, using

$$e_3(\kappa) + \frac{1}{2}\kappa\bar{\kappa} - 2\omega\kappa = 2\phi_1\eta + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\eta^2$$

we deduce

$$\begin{aligned} \overline{e_3(\kappa)} &= -\frac{1}{2}\bar{\kappa}\bar{\kappa} + 2\bar{\omega}\bar{\kappa} + 2\bar{\rho} - \frac{1}{2}\bar{\vartheta}\bar{\vartheta} + 2\bar{\eta}^2 \\ &= -\frac{1}{2}\bar{\kappa}\bar{\kappa} + 2\bar{\omega}\bar{\kappa} + 2\bar{\rho} + 2\bar{\omega}\bar{\kappa} - \frac{1}{2}\bar{\kappa}\bar{\kappa} - \frac{1}{2}\bar{\vartheta}\bar{\vartheta} + 2\bar{\eta}^2. \end{aligned}$$

Making use of Corollary 2.2.11

$$\begin{aligned} e_3(\bar{\kappa}) &= \overline{e_3(\kappa)} + \text{Err}[e_3\bar{\kappa}] \\ &= -\frac{1}{2}\bar{\kappa}\bar{\kappa} + 2\bar{\omega}\bar{\kappa} + 2\bar{\rho} + 2\bar{\omega}\bar{\kappa} - \frac{1}{2}\bar{\kappa}\bar{\kappa} - \frac{1}{2}\bar{\vartheta}\bar{\vartheta} + 2\bar{\eta}^2 + \text{Err}[e_3\bar{\kappa}] \end{aligned}$$

and,

$$\begin{aligned} e_3\left(\bar{\kappa} - \frac{2}{r}\right) &= e_3(\bar{\kappa}) + \frac{2}{r^2} r (\bar{\kappa} + \underline{A}) \\ &= -\frac{1}{2} \bar{\kappa} \bar{\kappa} + 2\bar{\omega} \bar{\kappa} + 2\bar{\rho} + 2\bar{\omega} \check{\kappa} - \frac{1}{2} \check{\kappa} \check{\kappa} - \frac{1}{2} \vartheta \vartheta + 2\bar{\eta}^2 + \frac{1}{r} \bar{\kappa} + \frac{1}{r} \underline{A} + \text{Err}[e_3 \bar{\kappa}] \\ &= -\frac{1}{2} \bar{\kappa} \left(\bar{\kappa} - \frac{2}{r}\right) + 2\bar{\omega} \bar{\kappa} + 2\bar{\rho} + 2\bar{\omega} \check{\kappa} - \frac{1}{2} \check{\kappa} \check{\kappa} - \frac{1}{2} \vartheta \vartheta + 2\bar{\eta}^2 + \frac{1}{r} \underline{A} + \text{Err}[e_3 \bar{\kappa}]. \end{aligned}$$

Now,

$$\begin{aligned} 2\bar{\omega} \bar{\kappa} + 2\bar{\rho} &= 2\bar{\omega} \left(\bar{\kappa} - \frac{2}{r}\right) + \frac{4}{r} \bar{\omega} + 2\bar{\rho} \\ &= 2\bar{\omega} \left(\bar{\kappa} - \frac{2}{r}\right) + \frac{4}{r} \left(\bar{\omega} - \frac{m}{r^2}\right) + 2 \left(\bar{\rho} + \frac{2m}{r^3}\right). \end{aligned}$$

Hence,

$$\begin{aligned} e_3\left(\bar{\kappa} - \frac{2}{r}\right) + \frac{1}{2} \bar{\kappa} \left(\bar{\kappa} - \frac{2}{r}\right) &= 2\bar{\omega} \left(\bar{\kappa} - \frac{2}{r}\right) + \frac{4}{r} \left(\bar{\omega} - \frac{m}{r^2}\right) + 2 \left(\bar{\rho} + \frac{2m}{r^3}\right) \\ &\quad + 2\bar{\eta}^2 + 2\bar{\omega} \check{\kappa} - \frac{1}{2} \check{\kappa} \check{\kappa} - \frac{1}{2} \vartheta \vartheta + \frac{1}{r} \underline{A} + \text{Err}[e_3 \bar{\kappa}]. \end{aligned}$$

In view of Corollary 2.2.11 the error term $\text{Err}[e_3(\bar{\kappa})]$ is given by

$$\begin{aligned} \text{Err}[e_3(\bar{\kappa})] &= -\varsigma^{-1} \check{\zeta} \left(\overline{e_3 \kappa + \kappa \kappa} - \bar{\kappa} \bar{\kappa}\right) + \varsigma^{-1} \left(\overline{\check{\zeta}(e_3 \kappa + \kappa \kappa)} - \check{\zeta} \bar{\kappa} \bar{\kappa}\right) \\ &\quad + \left(\check{\Omega} + \varsigma^{-1} \bar{\Omega} \check{\zeta}\right) \left(\overline{e_4 \kappa + \kappa^2} - \bar{\kappa}^2\right) - \varsigma^{-1} \bar{\Omega} \left(\overline{\check{\zeta}(e_4 \kappa + \kappa^2)} - \check{\zeta} \bar{\kappa} \bar{\kappa}\right) \\ &\quad - \varsigma^{-1} \left(\overline{\check{\Omega} \varsigma(e_4 \kappa + \kappa^2)} - \check{\Omega} \varsigma \bar{\kappa} \bar{\kappa}\right) + \bar{\kappa} \bar{\kappa}. \end{aligned}$$

Together with the null structure equations for $e_3(\kappa)$ and $e_4(\kappa)$, we infer

$$\begin{aligned} \text{Err}[e_3(\bar{\kappa})] &= -\varsigma^{-1} \check{\zeta} \left(\overline{\frac{1}{2} \kappa \kappa + 2\bar{\omega} \kappa + 2\rho + 2\phi_1 \eta - \frac{1}{2} \vartheta \vartheta + 2\eta^2} - \bar{\kappa} \bar{\kappa}\right) \\ &\quad + \varsigma^{-1} \left(\check{\zeta} \left(\overline{\frac{1}{2} \kappa \kappa + 2\bar{\omega} \kappa + 2\rho + 2\phi_1 \eta - \frac{1}{2} \vartheta \vartheta + 2\eta^2}\right) - \check{\zeta} \bar{\kappa} \bar{\kappa}\right) \\ &\quad + \left(\check{\Omega} + \varsigma^{-1} \bar{\Omega} \check{\zeta}\right) \left(\overline{\frac{1}{2} \kappa^2 - \frac{1}{4} \vartheta^2} - \bar{\kappa}^2\right) - \varsigma^{-1} \bar{\Omega} \left(\check{\zeta} \left(\overline{\frac{1}{2} \kappa^2 - \frac{1}{4} \vartheta^2}\right) - \check{\zeta} \bar{\kappa} \bar{\kappa}\right) \\ &\quad - \varsigma^{-1} \left(\overline{\check{\Omega} \varsigma \left(\frac{1}{2} \kappa^2 - \frac{1}{4} \vartheta^2\right)} - \check{\Omega} \varsigma \bar{\kappa} \bar{\kappa}\right) + \bar{\kappa} \bar{\kappa}. \end{aligned}$$

and hence

$$\begin{aligned}
\text{Err}[e_3(\bar{\kappa})] &= -\varsigma^{-1} \left(-\frac{1}{2}\bar{\kappa}\bar{\kappa} + 2\bar{\omega}\bar{\kappa} + 2\bar{\rho} \right) \zeta - \frac{1}{2}\bar{\kappa}^2 (\check{\Omega} + \varsigma^{-1}\bar{\Omega}\zeta) \\
&\quad -\varsigma^{-1}\zeta \left(\frac{1}{2}\check{\kappa}\check{\kappa} + 2\check{\omega}\check{\kappa} - \frac{1}{2}\vartheta\vartheta + 2\eta^2 \right) \\
&\quad +\varsigma^{-1} \left(\zeta \left(\frac{1}{2}\kappa\kappa + 2\omega\kappa + 2\check{\rho} + 2\phi_1\eta - \frac{1}{2}\vartheta\vartheta + 2\eta^2 \right) - \check{\zeta}\check{\kappa}\bar{\kappa} \right) \\
&\quad + (\check{\Omega} + \varsigma^{-1}\bar{\Omega}\zeta) \left(\frac{1}{2}\check{\kappa}^2 - \frac{1}{4}\vartheta^2 \right) - \varsigma^{-1}\bar{\Omega} \left(\zeta \left(\frac{1}{2}\kappa^2 - \frac{1}{4}\vartheta^2 \right) - \check{\zeta}\check{\kappa}\bar{\kappa} \right) \\
&\quad -\varsigma^{-1} \left(\check{\Omega}\varsigma \left(\frac{1}{2}\kappa^2 - \frac{1}{4}\vartheta^2 \right) - \check{\Omega}\varsigma\kappa\bar{\kappa} \right) + \bar{\kappa}\bar{\kappa} \tag{A.3.1}
\end{aligned}$$

so that, in view of the definition of \underline{A} , we obtain

$$\begin{aligned}
&e_3 \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{1}{2}\bar{\kappa} \left(\bar{\kappa} - \frac{2}{r} \right) \\
&= 2\bar{\omega} \left(\bar{\kappa} - \frac{2}{r} \right) + \frac{4}{r} \left(\bar{\omega} - \frac{m}{r^2} \right) + 2 \left(\bar{\rho} + \frac{2m}{r^3} \right) - \varsigma^{-1} \left(-\frac{1}{2}\bar{\kappa}\bar{\kappa} + 2\bar{\omega}\bar{\kappa} + 2\bar{\rho} \right) \zeta \\
&\quad - \frac{1}{2}\bar{\kappa}^2 (\check{\Omega} + \varsigma^{-1}\bar{\Omega}\zeta) - \frac{1}{r}\varsigma^{-1}\bar{\kappa}\zeta + \frac{1}{r}\bar{\kappa} (\check{\Omega} + \varsigma^{-1}\bar{\Omega}\zeta) + \text{Err} \left[e_3 \left(\bar{\kappa} - \frac{2}{r} \right) \right],
\end{aligned}$$

with

$$\begin{aligned}
\text{Err} \left[e_3 \left(\bar{\kappa} - \frac{2}{r} \right) \right] &= 2\bar{\eta}^2 + 2\check{\omega}\check{\kappa} - \frac{1}{2}\check{\kappa}\check{\kappa} - \frac{1}{2}\vartheta\vartheta + \frac{1}{r}\varsigma^{-1}\check{\zeta}\check{\kappa} - \frac{1}{r}\varsigma^{-1}\bar{\Omega}\check{\zeta}\check{\kappa} - \frac{1}{r}\varsigma^{-1}\check{\Omega}\varsigma\kappa \\
&\quad -\varsigma^{-1}\zeta \left(\frac{1}{2}\check{\kappa}\check{\kappa} + 2\check{\omega}\check{\kappa} - \frac{1}{2}\vartheta\vartheta + 2\eta^2 \right) \\
&\quad +\varsigma^{-1} \left(\zeta \left(\frac{1}{2}\kappa\kappa + 2\omega\kappa + 2\check{\rho} + 2\phi_1\eta - \frac{1}{2}\vartheta\vartheta + 2\eta^2 \right) - \check{\zeta}\check{\kappa}\bar{\kappa} \right) \\
&\quad + (\check{\Omega} + \varsigma^{-1}\bar{\Omega}\zeta) \left(\frac{1}{2}\check{\kappa}^2 - \frac{1}{4}\vartheta^2 \right) - \varsigma^{-1}\bar{\Omega} \left(\zeta \left(\frac{1}{2}\kappa^2 - \frac{1}{4}\vartheta^2 \right) - \check{\zeta}\check{\kappa}\bar{\kappa} \right) \\
&\quad -\varsigma^{-1} \left(\check{\Omega}\varsigma \left(\frac{1}{2}\kappa^2 - \frac{1}{4}\vartheta^2 \right) - \check{\Omega}\varsigma\kappa\bar{\kappa} \right) + \bar{\kappa}\bar{\kappa}.
\end{aligned}$$

This concludes the proof of Lemma 2.2.17.

A.4 Proof of Proposition 2.2.18

In view of Corollary 2.2.11 applied to

$$e_4(\kappa) + \frac{1}{2}\kappa^2 = -\frac{1}{2}\vartheta^2,$$

we deduce,

$$e_4\check{\kappa} + \bar{\kappa}\check{\kappa} = -\frac{1}{2}\check{\kappa}^2 - \frac{1}{2}\overline{\check{\kappa}^2} - \frac{1}{2}(\vartheta^2 - \overline{\vartheta^2}).$$

In view of Corollary 2.2.11 applied to

$$e_4(\underline{\kappa}) + \frac{1}{2}\kappa\underline{\kappa} = -2\phi_1\underline{\zeta} + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\underline{\zeta}^2$$

we deduce,

$$e_4\check{\underline{\kappa}} + \frac{1}{2}\bar{\kappa}\check{\underline{\kappa}} + \frac{1}{2}\check{\underline{\kappa}}\bar{\underline{\kappa}} = -\frac{1}{2}\check{\underline{\kappa}}\check{\underline{\kappa}} - \frac{1}{2}\overline{\check{\underline{\kappa}}\check{\underline{\kappa}}} + F - \bar{F}$$

where,

$$\begin{aligned} F - \bar{F} &= \left(-2\phi_1\underline{\zeta} + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\underline{\zeta}^2\right) - \overline{\left(-2\phi_1\underline{\zeta} + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\underline{\zeta}^2\right)} \\ &= -2\phi_1\underline{\zeta} + 2\check{\rho} + \left(-\frac{1}{2}\vartheta\underline{\vartheta} + 2\underline{\zeta}^2\right) - \overline{\left(-\frac{1}{2}\vartheta\underline{\vartheta} + 2\underline{\zeta}^2\right)}. \end{aligned}$$

Hence,

$$\begin{aligned} e_4\check{\underline{\kappa}} + \frac{1}{2}\bar{\kappa}\check{\underline{\kappa}} + \frac{1}{2}\check{\underline{\kappa}}\bar{\underline{\kappa}} &= -2\phi_1\underline{\zeta} + 2\check{\rho} + \text{Err}[e_4\check{\underline{\kappa}}] \\ \text{Err}[e_4\check{\underline{\kappa}}] &:= -\frac{1}{2}\check{\underline{\kappa}}\check{\underline{\kappa}} - \frac{1}{2}\overline{\check{\underline{\kappa}}\check{\underline{\kappa}}} + \left(-\frac{1}{2}\vartheta\underline{\vartheta} + 2\underline{\zeta}^2\right) - \overline{\left(-\frac{1}{2}\vartheta\underline{\vartheta} + 2\underline{\zeta}^2\right)}. \end{aligned}$$

In view of Corollary 2.2.11 applied to $e_4(\underline{\omega}) = \rho + 3\underline{\zeta}^2$ we deduce,

$$e_4\check{\underline{\omega}} = -\bar{\kappa}\check{\underline{\omega}} + (\rho + 3\underline{\zeta}^2) - \overline{(\rho + 3\underline{\zeta}^2)} = \check{\rho} - \bar{\kappa}\check{\underline{\omega}} + 3(\underline{\zeta}^2 - \overline{\underline{\zeta}^2}).$$

In view of Corollary 2.2.11 applied to

$$e_4(\rho) + \frac{3}{2}\kappa\rho = \phi_1\beta - \frac{1}{2}\vartheta\alpha - \zeta\beta$$

we deduce,

$$e_4\check{\rho} + \frac{3}{2}\overline{\kappa}\check{\rho} + \frac{3}{2}\overline{\rho}\check{\kappa} = -\frac{3}{2}\check{\kappa}\check{\rho} + \frac{1}{2}\overline{\kappa}\check{\rho} + \phi_1\beta - \left(\frac{1}{2}\vartheta\alpha + \zeta\beta\right) + \overline{\left(\frac{1}{2}\vartheta\alpha + \zeta\beta\right)}.$$

$$e_4\mu + \frac{3}{2}\kappa\mu = \text{Err}[e_4\mu],$$

we deduce

$$e_4\check{\mu} + \frac{3}{2}\overline{\kappa}\check{\mu} + \frac{3}{2}\overline{\mu}\check{\kappa} = -\frac{3}{2}\check{\kappa}\check{\mu} + \frac{1}{2}\overline{\kappa}\check{\mu} + \text{Err}[e_4\mu] - \overline{\text{Err}[e_4\mu]}.$$

In view of Corollary 2.2.11 applied to $e_4(\Omega) = -2\underline{\omega}$ we deduce,

$$-e_4(\check{\Omega}) = 2\underline{\omega} - \overline{\kappa\check{\Omega}}$$

as stated.

In view of Corollary 2.2.11 applied to the equation

$$e_3(\kappa) + \frac{1}{2}\underline{\kappa}\kappa = 2\phi_1\eta + 2\rho + 2\eta^2 + 2\underline{\omega}\kappa - \frac{1}{2}\vartheta\vartheta$$

to deduce,

$$\begin{aligned} e_3(\check{\kappa}) &= e_3(\kappa) - \overline{e_3(\kappa)} - \text{Err}[e_3(\overline{\kappa})] \\ &= -\frac{1}{2}\underline{\kappa}\kappa + 2\phi_1\eta + 2\rho + 2\eta^2 + 2\underline{\omega}\kappa - \frac{1}{2}\vartheta\vartheta \\ &\quad + \frac{1}{2}\overline{\kappa}\kappa - 2\overline{\rho} - 2\overline{\eta^2} - 2\overline{\omega}\kappa + \frac{1}{2}\overline{\vartheta\vartheta} - \text{Err}[e_3\overline{\kappa}] \\ &= 2\phi_1\eta + 2\check{\rho} - \frac{1}{2}(\underline{\kappa}\check{\kappa} + \kappa\check{\kappa}) + 2(\underline{\omega}\check{\kappa} + \kappa\check{\omega}) \\ &\quad + 2\left(\eta^2 - \overline{\eta^2}\right) - \frac{1}{2}\overline{\kappa}\check{\kappa} + 2\overline{\omega}\check{\kappa} - \frac{1}{2}(\vartheta\vartheta - \overline{\vartheta\vartheta}) - \text{Err}[e_3\overline{\kappa}]. \end{aligned}$$

Now, recall that we have in view of (A.3.1),

$$\begin{aligned} \text{Err}[e_3(\bar{\kappa})] &= -\varsigma^{-1} \left(-\frac{1}{2} \bar{\kappa} \bar{\kappa} + 2\bar{\omega} \bar{\kappa} + 2\bar{\rho} \right) \zeta - \frac{1}{2} \bar{\kappa}^2 (\check{\Omega} + \varsigma^{-1} \bar{\Omega} \zeta) \\ &\quad - \varsigma^{-1} \zeta \left(\overline{\frac{1}{2} \check{\kappa} \check{\kappa} + 2\check{\omega} \check{\kappa} - \frac{1}{2} \vartheta \vartheta + 2\eta^2} \right) \\ &\quad + \varsigma^{-1} \left(\overline{\zeta \left(\frac{1}{2} \kappa \kappa + 2\omega \kappa + 2\check{\rho} + 2\phi_1 \eta - \frac{1}{2} \vartheta \vartheta + 2\eta^2 \right) - \check{\zeta} \bar{\kappa} \bar{\kappa}} \right) \\ &\quad + (\check{\Omega} + \varsigma^{-1} \bar{\Omega} \zeta) \left(\overline{\frac{1}{2} \check{\kappa}^2 - \frac{1}{4} \vartheta^2} \right) - \varsigma^{-1} \bar{\Omega} \left(\overline{\zeta \left(\frac{1}{2} \kappa^2 - \frac{1}{4} \vartheta^2 \right) - \check{\zeta} \bar{\kappa} \bar{\kappa}} \right) \\ &\quad - \varsigma^{-1} \left(\overline{\check{\Omega} \varsigma \left(\frac{1}{2} \kappa^2 - \frac{1}{4} \vartheta^2 \right) - \check{\Omega} \varsigma \kappa \bar{\kappa}} \right) + \bar{\kappa} \check{\kappa}. \end{aligned}$$

We deduce

$$\begin{aligned} e_3(\check{\kappa}) &= 2\phi_1 \eta + 2\check{\rho} - \frac{1}{2} (\kappa \check{\kappa} + \kappa \check{\kappa}) + 2(\omega \check{\kappa} + \kappa \check{\omega}) \\ &\quad + \varsigma^{-1} \left(-\frac{1}{2} \bar{\kappa} \bar{\kappa} + 2\bar{\omega} \bar{\kappa} + 2\bar{\rho} \right) \zeta + \frac{1}{2} \bar{\kappa}^2 (\check{\Omega} + \varsigma^{-1} \bar{\Omega} \zeta) \\ &\quad + 2 \left(\eta^2 - \bar{\eta}^2 \right) - \frac{1}{2} \bar{\kappa} \bar{\kappa} + 2\bar{\omega} \bar{\kappa} - \frac{1}{2} (\vartheta \vartheta - \bar{\vartheta} \bar{\vartheta}) + \varsigma^{-1} \zeta \left(\overline{\frac{1}{2} \check{\kappa} \check{\kappa} + 2\check{\omega} \check{\kappa} - \frac{1}{2} \vartheta \vartheta + 2\eta^2} \right) \\ &\quad - \varsigma^{-1} \left(\overline{\zeta \left(\frac{1}{2} \kappa \kappa + 2\omega \kappa + 2\check{\rho} + 2\phi_1 \eta - \frac{1}{2} \vartheta \vartheta + 2\eta^2 \right) - \check{\zeta} \bar{\kappa} \bar{\kappa}} \right) \\ &\quad - (\check{\Omega} + \varsigma^{-1} \bar{\Omega} \zeta) \left(\overline{\frac{1}{2} \check{\kappa}^2 - \frac{1}{4} \vartheta^2} \right) + \varsigma^{-1} \bar{\Omega} \left(\overline{\zeta \left(\frac{1}{2} \kappa^2 - \frac{1}{4} \vartheta^2 \right) - \check{\zeta} \bar{\kappa} \bar{\kappa}} \right) \\ &\quad + \varsigma^{-1} \left(\overline{\check{\Omega} \varsigma \left(\frac{1}{2} \kappa^2 - \frac{1}{4} \vartheta^2 \right) - \check{\Omega} \varsigma \kappa \bar{\kappa}} \right) - \bar{\kappa} \check{\kappa} \end{aligned}$$

as desired.

In view of Corollary 2.2.11 applied to the equation

$$e_3(\underline{\kappa}) + \frac{1}{2} \underline{\kappa}^2 = 2\phi_1 \underline{\xi} - 2\omega \underline{\kappa} + 2(\eta - 3\zeta) \underline{\xi} - \frac{1}{2} \vartheta^2$$

we deduce,

$$\begin{aligned} e_3(\check{\kappa}) + \kappa \check{\kappa} &= 2\phi_1 \underline{\xi} - 2(\check{\omega} \underline{\kappa} + \omega \check{\kappa}) \\ &\quad - \frac{1}{2} \bar{\kappa}^2 - 2\bar{\omega} \bar{\kappa} + 2(\eta - 3\zeta) \underline{\xi} - \overline{2(\eta - 3\zeta) \underline{\xi}} - \frac{1}{2} (\vartheta^2 - \bar{\vartheta}^2) - \text{Err}[e_3 \bar{\kappa}] \end{aligned}$$

where,

$$\begin{aligned} -\text{Err}[e_3\bar{\kappa}] &= \varsigma^{-1}\check{\zeta} \left(\overline{e_3\kappa + \kappa^2} - \overline{\kappa\kappa} \right) - \varsigma^{-1} \left(\overline{\check{\zeta}(e_3\kappa + \kappa^2)} - \overline{\check{\zeta}\kappa\kappa} \right) \\ &\quad - \left(\check{\Omega} + \varsigma^{-1}\overline{\check{\Omega}\check{\zeta}} \right) \left(\overline{e_4\kappa + \kappa\kappa} - \overline{\kappa\kappa} \right) + \varsigma^{-1}\overline{\check{\Omega}} \left(\overline{\check{\zeta}(e_4\kappa + \kappa\kappa)} - \overline{\check{\zeta}\kappa\kappa} \right) \\ &\quad + \varsigma^{-1} \left(\overline{\check{\Omega}\check{\zeta}(e_4\kappa + \kappa\kappa)} - \overline{\check{\Omega}\check{\zeta}\kappa\kappa} \right) - \overline{\check{\kappa}^2}. \end{aligned}$$

In view of the null structure equations for $e_3(\underline{\kappa})$ and $e_4(\underline{\kappa})$, we infer

$$\begin{aligned} -\text{Err}[e_3\bar{\kappa}] &= \varsigma^{-1}\check{\zeta} \left(-\frac{1}{2}\overline{\kappa^2} - 2\overline{\omega\kappa} \right) - \left(\check{\Omega} + \varsigma^{-1}\overline{\check{\Omega}\check{\zeta}} \right) \left(-\frac{1}{2}\overline{\kappa\kappa} + 2\overline{\rho} \right) \\ &\quad + \varsigma^{-1}\check{\zeta} \left(\overline{\frac{1}{2}\check{\kappa}^2 - 2\check{\omega}\check{\kappa} + 2(\eta - 3\check{\zeta})\check{\xi} - \frac{1}{2}\check{\vartheta}^2} \right) - \left(\check{\Omega} + \varsigma^{-1}\overline{\check{\Omega}\check{\zeta}} \right) \left(\overline{\frac{1}{2}\check{\kappa}\check{\kappa} - \frac{1}{2}\check{\vartheta}\check{\vartheta} + 2\check{\zeta}^2} \right) \\ &\quad - \varsigma^{-1} \left(\overline{\check{\zeta} \left(\frac{1}{2}\overline{\kappa^2} - 2\overline{\omega\kappa} + 2\overline{\phi_1\xi} + 2(\eta - 3\check{\zeta})\check{\xi} - \frac{1}{2}\check{\vartheta}^2 \right)} - \overline{\check{\zeta}\kappa\kappa} \right) \\ &\quad + \varsigma^{-1}\overline{\check{\Omega}} \left(\overline{\check{\zeta} \left(\frac{1}{2}\overline{\kappa\kappa} - 2\overline{\phi_1\zeta} + 2\overline{\rho} - \frac{1}{2}\check{\vartheta}\check{\vartheta} + 2\check{\zeta}^2 \right)} - \overline{\check{\zeta}\kappa\kappa} \right) \\ &\quad + \varsigma^{-1} \left(\overline{\check{\Omega}\check{\zeta} \left(\frac{1}{2}\overline{\kappa\kappa} - 2\overline{\phi_1\zeta} + 2\overline{\rho} - \frac{1}{2}\check{\vartheta}\check{\vartheta} + 2\check{\zeta}^2 \right)} - \overline{\check{\Omega}\check{\zeta}\kappa\kappa} \right) - \overline{\check{\kappa}^2} \end{aligned}$$

and hence

$$\begin{aligned} e_3(\check{\kappa}) + \underline{\kappa}\check{\kappa} &= 2\overline{\phi_1\xi} - 2(\check{\omega}\underline{\kappa} + \underline{\omega}\check{\kappa}) + \varsigma^{-1}\check{\zeta} \left(-\frac{1}{2}\overline{\kappa^2} - 2\overline{\omega\kappa} \right) \\ &\quad - \left(\check{\Omega} + \varsigma^{-1}\overline{\check{\Omega}\check{\zeta}} \right) \left(-\frac{1}{2}\overline{\kappa\kappa} + 2\overline{\rho} \right) + \text{Err}[e_3(\check{\kappa})], \\ \text{Err}[e_3(\check{\kappa})] &= -\frac{1}{2}\overline{\check{\kappa}^2} - 2\overline{\check{\omega}\check{\kappa}} + 2(\eta - 3\check{\zeta})\check{\xi} - \overline{2(\eta - 3\check{\zeta})\check{\xi}} - \frac{1}{2}(\check{\vartheta}^2 - \overline{\check{\vartheta}^2}) \\ &\quad - \varsigma^{-1} \left(\overline{\check{\zeta} \left(\frac{1}{2}\overline{\kappa^2} - 2\overline{\omega\kappa} + 2\overline{\phi_1\xi} + 2(\eta - 3\check{\zeta})\check{\xi} - \frac{1}{2}\check{\vartheta}^2 \right)} - \overline{\check{\zeta}\kappa\kappa} \right) \\ &\quad + \varsigma^{-1}\overline{\check{\Omega}} \left(\overline{\check{\zeta} \left(\frac{1}{2}\overline{\kappa\kappa} - 2\overline{\phi_1\zeta} + 2\overline{\rho} - \frac{1}{2}\check{\vartheta}\check{\vartheta} + 2\check{\zeta}^2 \right)} - \overline{\check{\zeta}\kappa\kappa} \right) \\ &\quad + \varsigma^{-1} \left(\overline{\check{\Omega}\check{\zeta} \left(\frac{1}{2}\overline{\kappa\kappa} - 2\overline{\phi_1\zeta} + 2\overline{\rho} - \frac{1}{2}\check{\vartheta}\check{\vartheta} + 2\check{\zeta}^2 \right)} - \overline{\check{\Omega}\check{\zeta}\kappa\kappa} \right) - \overline{\check{\kappa}^2} \end{aligned}$$

as desired.

In view of Corollary 2.2.11 applied to equation

$$e_3(\rho) + \frac{3}{2}\underline{\kappa}\rho = \underline{\phi}_1\underline{\beta} - \frac{1}{2}\underline{\vartheta}\underline{\alpha} - \underline{\zeta}\underline{\beta} + 2\underline{\eta}\underline{\beta} + 2\underline{\xi}\underline{\beta}$$

we deduce,

$$\begin{aligned} e_3\check{\rho} + \frac{3}{2}\underline{\kappa}\check{\rho} + \frac{3}{2}\underline{\check{\kappa}}\check{\rho} &= \underline{\phi}_1\underline{\beta} - \left(\frac{1}{2}\underline{\vartheta}\underline{\alpha} + \underline{\zeta}\underline{\beta} - 2\underline{\eta}\underline{\beta} - 2\underline{\xi}\underline{\beta} \right) + \overline{\left(\frac{1}{2}\underline{\vartheta}\underline{\alpha} + \underline{\zeta}\underline{\beta} - 2\underline{\eta}\underline{\beta} - 2\underline{\xi}\underline{\beta} \right)} \\ &\quad - \frac{3}{2}\underline{\check{\kappa}}\check{\rho} - \text{Err}[e_3\check{\rho}] \end{aligned}$$

where

$$\begin{aligned} -\text{Err}[e_3(\check{\rho})] &= \varsigma^{-1}\check{\zeta} \left(\overline{e_3\rho + \underline{\kappa}\rho} - \underline{\kappa}\check{\rho} \right) - \varsigma^{-1} \left(\overline{\check{\zeta}(e_3\rho + \underline{\kappa}\rho)} - \check{\zeta}\underline{\check{\kappa}}\check{\rho} \right) \\ &\quad - \left(\check{\underline{\Omega}} + \varsigma^{-1}\check{\underline{\Omega}}\check{\zeta} \right) \left(\overline{e_4\rho + \underline{\kappa}\rho} - \underline{\kappa}\check{\rho} \right) + \varsigma^{-1}\check{\underline{\Omega}} \left(\overline{\check{\zeta}(e_4\rho + \underline{\kappa}\rho)} - \check{\zeta}\underline{\check{\kappa}}\check{\rho} \right) \\ &\quad + \varsigma^{-1} \left(\overline{\check{\underline{\Omega}}\varsigma(e_4\rho + \underline{\kappa}\rho)} - \check{\underline{\Omega}}\varsigma\underline{\kappa} \right) - \underline{\check{\kappa}}\check{\rho}. \end{aligned}$$

In view of the null structure equations for $e_3(\rho)$ and $e_4(\rho)$, we infer

$$\begin{aligned} -\text{Err}[e_3(\check{\rho})] &= -\frac{3}{2}\underline{\kappa}\check{\rho}\varsigma^{-1}\check{\zeta} + \frac{3}{2}\underline{\kappa}\check{\rho} \left(\check{\underline{\Omega}} + \varsigma^{-1}\check{\underline{\Omega}}\check{\zeta} \right) \\ &\quad + \varsigma^{-1}\check{\zeta} \left(\overline{-\frac{1}{2}\underline{\check{\kappa}}\check{\rho} - \frac{1}{2}\underline{\vartheta}\underline{\alpha} - \underline{\zeta}\underline{\beta} + 2(\underline{\eta}\underline{\beta} + \underline{\xi}\underline{\beta})} \right) \\ &\quad - \varsigma^{-1} \left(\overline{\check{\zeta} \left(-\frac{1}{2}\underline{\kappa}\rho + \underline{\phi}_1\underline{\beta} - \frac{1}{2}\underline{\vartheta}\underline{\alpha} - \underline{\zeta}\underline{\beta} + 2(\underline{\eta}\underline{\beta} + \underline{\xi}\underline{\beta}) \right)} - \check{\zeta}\underline{\check{\kappa}}\check{\rho} \right) \\ &\quad - \left(\check{\underline{\Omega}} + \varsigma^{-1}\check{\underline{\Omega}}\check{\zeta} \right) \left(\overline{-\frac{1}{2}\underline{\check{\kappa}}\check{\rho} - \frac{1}{2}\underline{\vartheta}\underline{\alpha} - \underline{\zeta}\underline{\beta}} \right) \\ &\quad + \varsigma^{-1}\check{\underline{\Omega}} \left(\overline{\check{\zeta} \left(-\frac{1}{2}\underline{\kappa}\rho + \underline{\phi}_1\underline{\beta} - \frac{1}{2}\underline{\vartheta}\underline{\alpha} - \underline{\zeta}\underline{\beta} \right)} - \check{\zeta}\underline{\check{\kappa}}\check{\rho} \right) \\ &\quad + \varsigma^{-1} \left(\overline{\check{\underline{\Omega}}\varsigma \left(-\frac{1}{2}\underline{\kappa}\rho + \underline{\phi}_1\underline{\beta} - \frac{1}{2}\underline{\vartheta}\underline{\alpha} - \underline{\zeta}\underline{\beta} \right)} - \check{\underline{\Omega}}\varsigma\underline{\kappa} \right) - \underline{\check{\kappa}}\check{\rho}. \end{aligned}$$

and hence

$$\begin{aligned}
e_3\check{\rho} + \frac{3}{2}\bar{\kappa}\check{\rho} &= -\frac{3}{2}\bar{\rho}\check{\kappa} + \check{\mu}_1\check{\beta} - \frac{3}{2}\bar{\kappa}\bar{\rho}\varsigma^{-1}\check{\zeta} + \frac{3}{2}\bar{\kappa}\bar{\rho}(\check{\Omega} + \varsigma^{-1}\bar{\Omega}\check{\zeta}) + \text{Err}[e_3\check{\rho}], \\
\text{Err}[e_3\check{\rho}] &= -\left(\frac{1}{2}\vartheta\check{\alpha} + \zeta\check{\beta} - 2\eta\check{\beta} - 2\xi\check{\beta}\right) + \overline{\left(\frac{1}{2}\vartheta\check{\alpha} + \zeta\check{\beta} - 2\eta\check{\beta} - 2\xi\check{\beta}\right)} - \frac{3}{2}\check{\kappa}\check{\rho} \\
&\quad + \varsigma^{-1}\check{\zeta} \left(\overline{-\frac{1}{2}\check{\kappa}\check{\rho} - \frac{1}{2}\vartheta\check{\alpha} - \zeta\check{\beta} + 2(\eta\check{\beta} + \xi\check{\beta})} \right) \\
&\quad - \varsigma^{-1} \left(\check{\zeta} \left(\overline{-\frac{1}{2}\kappa\rho + \check{\mu}_1\check{\beta} - \frac{1}{2}\vartheta\check{\alpha} - \zeta\check{\beta} + 2(\eta\check{\beta} + \xi\check{\beta})} \right) - \check{\zeta}\bar{\kappa}\bar{\rho} \right) \\
&\quad - (\check{\Omega} + \varsigma^{-1}\bar{\Omega}\check{\zeta}) \left(\overline{-\frac{1}{2}\check{\kappa}\check{\rho} - \frac{1}{2}\vartheta\check{\alpha} - \zeta\check{\beta}} \right) \\
&\quad + \varsigma^{-1}\bar{\Omega} \left(\check{\zeta} \left(\overline{-\frac{1}{2}\kappa\rho + \check{\mu}_1\check{\beta} - \frac{1}{2}\vartheta\check{\alpha} - \zeta\check{\beta}} \right) - \check{\zeta}\bar{\kappa}\bar{\rho} \right) \\
&\quad + \varsigma^{-1} \left(\overline{\check{\Omega}\varsigma \left(\overline{-\frac{1}{2}\kappa\rho + \check{\mu}_1\check{\beta} - \frac{1}{2}\vartheta\check{\alpha} - \zeta\check{\beta}} \right)} - \bar{\Omega}\varsigma\kappa \right) - \bar{\kappa}\bar{\rho},
\end{aligned}$$

which ends the proof of Proposition 2.2.18.

A.5 Proof of Proposition 2.2.19

In view of the null structure equation for $e_3(\zeta)$, we have

$$\frac{1}{2}\kappa\underline{\xi} + 2\check{\mu}_1^*\underline{\omega} = e_3(\zeta) + \frac{1}{2}\kappa(\zeta + \eta) - 2\underline{\omega}(\zeta - \eta) - \underline{\beta} + \frac{1}{2}\vartheta(\zeta + \eta) - \frac{1}{2}\vartheta\underline{\xi}$$

and hence

$$\begin{aligned}
\frac{1}{2}\kappa\underline{\xi} + 2\check{\mu}_1^*\underline{\omega} &= \left(\frac{1}{2}\kappa + 2\underline{\omega} + \frac{1}{2}\vartheta \right) \eta + e_3(\zeta) - \underline{\beta} \\
&\quad + \frac{1}{2}\kappa\zeta - 2\underline{\omega}\zeta + \frac{1}{2}\vartheta\zeta - \frac{1}{2}\vartheta\underline{\xi}
\end{aligned}$$

which is the first desired identity.

To prove the second identity we start with

$$e_3(\kappa) + \frac{1}{2}\kappa\kappa - 2\underline{\omega}\kappa = 2\check{\mu}_1\eta + 2\rho - \frac{1}{2}\vartheta\vartheta + 2\eta^2.$$

Applying e_θ ,

$$\begin{aligned} & e_3(e_\theta(\kappa)) + [e_\theta, e_3]\kappa + \frac{1}{2}\underline{\kappa}e_\theta(\kappa) + \frac{1}{2}\kappa e_\theta(\underline{\kappa}) - 2\underline{\omega}e_\theta(\kappa) - 2\kappa e_\theta(\underline{\omega}) \\ &= 2e_\theta(\underline{\phi}_1\eta) + 2e_\theta(\rho) - \frac{1}{2}e_\theta(\underline{\vartheta}\vartheta) + 2e_\theta(\eta^2). \end{aligned}$$

Since $[e_\theta, e_3]\kappa = \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta\kappa + (\zeta - \eta)e_3\kappa - \underline{\xi}e_4\kappa$ we deduce,

$$\begin{aligned} 2e_\theta(\underline{\phi}_1\eta) + \eta e_3(\kappa) + 2e_\theta(\eta^2) &= -\underline{\xi}e_4(\kappa) - 2\kappa e_\theta(\underline{\omega}) + e_3(e_\theta(\kappa)) \\ &+ \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta(\kappa) + \zeta e_3(\kappa) + \frac{1}{2}\underline{\kappa}e_\theta(\kappa) + \frac{1}{2}\kappa e_\theta(\underline{\kappa}) \\ &- 2\underline{\omega}e_\theta(\kappa) - 2e_\theta(\rho) + \frac{1}{2}e_\theta(\underline{\vartheta}\vartheta), \end{aligned}$$

or, making use of the equations for $e_3\kappa$ and $e_4\kappa$ in Proposition 2.2.8,

$$\begin{aligned} & 2e_\theta(\underline{\phi}_1\eta) + \left(-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa + 2\underline{\phi}_1\eta + 2\rho - \frac{1}{2}\underline{\vartheta}\underline{\vartheta} + 2\eta^2\right)\eta + 2e_\theta(\eta^2) \\ &= -\left(-\frac{1}{2}\kappa^2 - \frac{1}{2}\underline{\vartheta}^2\right)\underline{\xi} - 2\kappa e_\theta(\underline{\omega}) + e_3(e_\theta(\kappa)) \\ &+ \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta(\kappa) + \left(-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa + 2\underline{\phi}_1\eta + 2\rho - \frac{1}{2}\underline{\vartheta}\underline{\vartheta} + 2\eta^2\right)\zeta \\ &+ \frac{1}{2}\underline{\kappa}e_\theta(\kappa) + \frac{1}{2}\kappa e_\theta(\underline{\kappa}) - 2\underline{\omega}e_\theta(\kappa) - 2e_\theta(\rho) + \frac{1}{2}e_\theta(\underline{\vartheta}\vartheta). \end{aligned}$$

Since $e_\theta = -\underline{\phi}_1^*$, $\underline{\phi}_1^*\underline{\phi}_1 = \underline{\phi}_2\underline{\phi}_2^* + 2K$ and $K = -\rho - \frac{1}{4}\kappa\underline{\kappa} + \frac{1}{4}\underline{\vartheta}\underline{\vartheta}$, we infer that

$$\begin{aligned} & \left(-2\underline{\phi}_2\underline{\phi}_2^* + \frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa + 2\underline{\phi}_1\eta + 6\rho - \frac{3}{2}\underline{\vartheta}\underline{\vartheta} + 2\eta^2\right)\eta + 2e_\theta(\eta^2) \\ &= \kappa\left(\frac{1}{2}\kappa\underline{\xi} + 2\underline{\phi}_1^*\underline{\omega}\right) + e_3(e_\theta(\kappa)) \\ &+ \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta(\kappa) + \left(-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\kappa + 2\underline{\phi}_1\eta + 2\rho - \frac{1}{2}\underline{\vartheta}\underline{\vartheta} + 2\eta^2\right)\zeta \\ &+ \frac{1}{2}\underline{\kappa}e_\theta(\kappa) + \frac{1}{2}\kappa e_\theta(\underline{\kappa}) - 2\underline{\omega}e_\theta(\kappa) - 2e_\theta(\rho) + \frac{1}{2}e_\theta(\underline{\vartheta}\vartheta) + \frac{1}{2}\underline{\vartheta}^2\underline{\xi}. \end{aligned}$$

Making use of the previously derived identity,

$$\begin{aligned} 2\underline{\phi}_1^*\underline{\omega} + \frac{1}{2}\kappa\underline{\xi} &= \left(\frac{1}{2}\underline{\kappa} + 2\underline{\omega} + \frac{1}{2}\underline{\vartheta}\right)\eta + e_3(\zeta) - \underline{\beta} \\ &+ \frac{1}{2}\underline{\kappa}\zeta - 2\underline{\omega}\zeta + \frac{1}{2}\underline{\vartheta}\zeta - \frac{1}{2}\underline{\vartheta}\underline{\xi}, \end{aligned}$$

we infer that,

$$\begin{aligned}
& \left(-2\phi_2\phi_2^* + \frac{1}{2}\kappa\kappa + 2\underline{\omega}\kappa + 2\phi_1\eta + 6\rho - \frac{3}{2}\vartheta\underline{\vartheta} + 2\eta^2 \right) \eta + 2e_\theta(\eta^2) \\
= & \kappa \left(\left(\frac{1}{2}\kappa + 2\underline{\omega} + \frac{1}{2}\vartheta \right) \eta + e_3(\zeta) - \underline{\beta} \right) \\
& + \kappa \left(\frac{1}{2}\kappa\zeta - 2\underline{\omega}\zeta + \frac{1}{2}\vartheta\zeta - \frac{1}{2}\vartheta\underline{\xi} \right) + e_3(e_\theta(\kappa)) \\
& + \frac{1}{2}(\kappa + \vartheta)e_\theta(\kappa) + \left(-\frac{1}{2}\kappa\kappa + 2\underline{\omega}\kappa + 2\phi_1\eta + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\eta^2 \right) \zeta \\
& + \frac{1}{2}\kappa e_\theta(\kappa) + \frac{1}{2}\kappa e_\theta(\kappa) - 2\underline{\omega}e_\theta(\kappa) - 2e_\theta(\rho) + \frac{1}{2}e_\theta(\vartheta\vartheta) + \frac{1}{2}\vartheta^2\underline{\xi},
\end{aligned}$$

or,

$$\begin{aligned}
& \left(-2\phi_2\phi_2^* + 6\rho + 2\phi_1\eta - \frac{1}{2}\kappa\vartheta - \frac{3}{2}\vartheta\underline{\vartheta} + 2\eta^2 \right) \eta + 2e_\theta(\eta^2) \\
= & \kappa \left(e_3(\zeta) - \underline{\beta} \right) + e_3(e_\theta(\kappa)) \\
& + \kappa \left(\frac{1}{2}\kappa\zeta - 2\underline{\omega}\zeta + \frac{1}{2}\vartheta\zeta - \frac{1}{2}\vartheta\underline{\xi} \right) \\
& + \frac{1}{2}(\kappa + \vartheta)e_\theta(\kappa) + \left(-\frac{1}{2}\kappa\kappa + 2\underline{\omega}\kappa + 2\phi_1\eta + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\eta^2 \right) \zeta \\
& + \frac{1}{2}\kappa e_\theta(\kappa) + \frac{1}{2}\kappa e_\theta(\kappa) - 2\underline{\omega}e_\theta(\kappa) - 2e_\theta(\rho) + \frac{1}{2}e_\theta(\vartheta\vartheta) + \frac{1}{2}\vartheta^2\underline{\xi}
\end{aligned}$$

and hence

$$\begin{aligned}
& \left(2\phi_2\phi_2^* - 2\phi_1\eta + \frac{1}{2}\kappa\vartheta - 2\eta^2 \right) \eta - 2e_\theta(\eta^2) \\
= & \kappa \left(-e_3(\zeta) + \underline{\beta} \right) - e_3(e_\theta(\kappa)) \\
& - \kappa \left(\frac{1}{2}\kappa\zeta - 2\underline{\omega}\zeta + \frac{1}{2}\vartheta\zeta - \frac{1}{2}\vartheta\underline{\xi} \right) + 6\rho\eta - \frac{3}{2}\vartheta\underline{\vartheta}\eta \\
& - \frac{1}{2}(\kappa + \vartheta)e_\theta(\kappa) - \left(-\frac{1}{2}\kappa\kappa + 2\underline{\omega}\kappa + 2\phi_1\eta + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2\eta^2 \right) \zeta \\
& - \frac{1}{2}\kappa e_\theta(\kappa) - \frac{1}{2}\kappa e_\theta(\kappa) + 2\underline{\omega}e_\theta(\kappa) + 2e_\theta(\rho) - \frac{1}{2}e_\theta(\vartheta\vartheta) - \frac{1}{2}\vartheta^2\underline{\xi}
\end{aligned}$$

which is the second desired identity.

To prove the third identity we start with,

$$e_3(\kappa) + \frac{1}{2}\kappa^2 + 2\underline{\omega}\kappa = 2\phi_1\underline{\xi} + 2(\eta - 3\underline{\zeta})\underline{\xi} - \frac{1}{2}\vartheta^2.$$

Taking $e_\theta = -\phi_1^*$ and using $\phi_1^* \phi_1 = \phi_2 \phi_2^* + 2K$ as before,

$$\begin{aligned} & e_3(e_\theta(\underline{\kappa})) + [e_\theta, e_3]\underline{\kappa} + \underline{\kappa}e_\theta(\underline{\kappa}) + 2\underline{\omega}e_\theta(\underline{\kappa}) + 2\underline{\kappa}e_\theta(\underline{\omega}) \\ &= -2\phi_1^* \phi_1 \underline{\xi} + 2e_\theta\left((\eta - 3\zeta)\underline{\xi}\right) - \frac{1}{2}e_\theta(\underline{\vartheta}^2) \\ &= -2(\phi_2 \phi_2^* + 2K)\underline{\xi} + 2e_\theta\left((\eta - 3\zeta)\underline{\xi}\right) - \frac{1}{2}e_\theta(\underline{\vartheta}^2). \end{aligned}$$

Thus, since $[e_\theta, e_3]\underline{\kappa} = \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta \underline{\kappa} + \zeta - \eta)e_3 \underline{\kappa} - \underline{\xi}e_4 \underline{\kappa}$,

$$\begin{aligned} -2(\phi_2 \phi_2^* + 2K)\underline{\xi} &= e_3(e_\theta(\underline{\kappa})) + 2\underline{\kappa}e_\theta(\underline{\omega}) + \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta \underline{\kappa} + (\zeta - \eta)e_3 \underline{\kappa} - \underline{\xi}e_4 \underline{\kappa} \\ &\quad + \underline{\kappa}e_\theta(\underline{\kappa}) + 2\underline{\omega}e_\theta(\underline{\kappa}) - 2e_\theta\left((\eta - 3\zeta)\underline{\xi}\right) + \frac{1}{2}e_\theta(\underline{\vartheta}^2). \end{aligned}$$

Making use of the equations for $e_3 \underline{\kappa}, e_4 \underline{\kappa}$ in Proposition 2.2.8

$$\begin{aligned} 2(\phi_2 \phi_2^* + 2K)\underline{\xi} &= 2\underline{\kappa} \phi_1^* \underline{\omega} - e_3(e_\theta(\underline{\kappa})) - \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta \underline{\kappa} \\ &\quad - (\zeta - \eta) \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega} \underline{\kappa} + 2\phi_1 \underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) \\ &\quad + \underline{\xi} \left(-\frac{1}{2}\underline{\kappa} \underline{\kappa} - 2\phi_1 \zeta + 2\rho - \frac{1}{2}\underline{\vartheta} \underline{\vartheta} + 2\zeta^2 \right) \\ &\quad - \underline{\kappa}e_\theta(\underline{\kappa}) - 2\underline{\omega}e_\theta(\underline{\kappa}) + 2e_\theta\left((\eta - 3\zeta)\underline{\xi}\right) - \frac{1}{2}e_\theta(\underline{\vartheta}^2). \end{aligned}$$

We deduce,

$$\begin{aligned} & 2(\phi_2 \phi_2^* + K)\underline{\xi} \\ &= 2\underline{\kappa} \phi_1^* \underline{\omega} + \eta \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega} \underline{\kappa} + 2\phi_1 \underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) + 2e_\theta\left((\eta - 3\zeta)\underline{\xi}\right) \\ &\quad - e_3(e_\theta(\underline{\kappa})) - \frac{1}{2}e_\theta(\underline{\vartheta}^2) - \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta(\underline{\kappa}) - \zeta \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega} \underline{\kappa} + 2\phi_1 \underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) \\ &\quad + \underline{\xi} \left(-\frac{1}{2}\underline{\kappa} \underline{\kappa} - 2K - 2\phi_1 \zeta + 2\rho - \frac{1}{2}\underline{\vartheta} \underline{\vartheta} + 2\zeta^2 \right) - \underline{\kappa}e_\theta(\underline{\kappa}) - 2\underline{\omega}e_\theta(\underline{\kappa}) - 6\eta \zeta \underline{\xi} - 6e_\theta(\zeta \underline{\xi}). \end{aligned}$$

Making use of $K = -\rho - \frac{1}{4}\underline{\kappa} \underline{\kappa} + \frac{1}{4}\underline{\vartheta} \underline{\vartheta}$ and reorganizing we deduce,

$$\begin{aligned} & 2(\phi_2 \phi_2^* + K)\underline{\xi} \\ &= 2\underline{\kappa} \phi_1^* \underline{\omega} + \eta \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega} \underline{\kappa} + 2\phi_1 \underline{\xi} + 2\eta \underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) + 2e_\theta(\eta \underline{\xi}) - e_3(e_\theta(\underline{\kappa})) - \frac{1}{2}e_\theta(\underline{\vartheta}^2) \\ &\quad - \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})e_\theta(\underline{\kappa}) - \zeta \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega} \underline{\kappa} + 2\phi_1 \underline{\xi} + 2(\eta - 3\zeta)\underline{\xi} - \frac{1}{2}\underline{\vartheta}^2 \right) \\ &\quad + \underline{\xi} \left(4\rho - \underline{\vartheta} \underline{\vartheta} - 2\phi_1 \zeta + 2\zeta^2 \right) - \underline{\kappa}e_\theta(\underline{\kappa}) - 2\underline{\omega}e_\theta(\underline{\kappa}) - 6\eta \zeta \underline{\xi} - 6e_\theta(\zeta \underline{\xi}). \end{aligned}$$

We make use again of the identity,

$$\begin{aligned} 2\phi_1^*\omega + \frac{1}{2}\kappa\xi &= \left(\frac{1}{2}\kappa + 2\omega + \frac{1}{2}\vartheta\right)\eta + e_3(\zeta) - \underline{\beta} \\ &\quad + \frac{1}{2}\kappa\zeta - 2\omega\zeta + \frac{1}{2}\vartheta\zeta - \frac{1}{2}\vartheta\xi, \end{aligned}$$

to derive,

$$\begin{aligned} &2(\phi_2\phi_2^* + K)\xi \\ &= \kappa\left(-\frac{1}{2}\kappa\xi + \left(\frac{1}{2}\kappa + 2\omega + \frac{1}{2}\vartheta\right)\eta + e_3(\zeta) - \underline{\beta}\right) + \kappa\left(\frac{1}{2}\kappa\zeta - 2\omega\zeta + \frac{1}{2}\vartheta\zeta - \frac{1}{2}\vartheta\xi\right) \\ &+ \eta\left(-\frac{1}{2}\kappa^2 - 2\omega\kappa + 2\phi_1\xi + 2\eta\xi - \frac{1}{2}\vartheta^2\right) + 2e_\theta(\eta\xi) - e_3(e_\theta(\kappa)) - \frac{1}{2}e_\theta(\vartheta^2) \\ &\quad - \frac{1}{2}(\kappa + \vartheta)e_\theta(\kappa) - \zeta\left(-\frac{1}{2}\kappa^2 - 2\omega\kappa + 2\phi_1\xi + 2(\eta - 3\zeta)\xi - \frac{1}{2}\vartheta^2\right) \\ &\quad + \xi\left(4\rho - \vartheta\vartheta - 2\phi_1\zeta + 2\zeta^2\right) - \kappa e_\theta(\kappa) - 2\omega e_\theta(\kappa) - 6\eta\zeta\xi - 6e_\theta(\zeta\xi). \end{aligned}$$

Grouping terms and using once more the identity $K = -\rho - \frac{1}{4}\kappa\kappa + \frac{1}{4}\vartheta\vartheta$ we deduce,

$$\begin{aligned} &2\phi_2\phi_2^*\xi \\ &= -e_3(e_\theta(\kappa)) + \kappa(e_3(\zeta) - \underline{\beta}) + \left(2\phi_1\xi + \frac{1}{2}\kappa\vartheta + 2\eta\xi - \frac{1}{2}\vartheta^2\right)\eta + 2e_\theta(\eta\xi) - \frac{1}{2}e_\theta(\vartheta^2) \\ &\quad + \kappa\left(\frac{1}{2}\kappa\zeta - 2\omega\zeta + \frac{1}{2}\vartheta\zeta - \frac{1}{2}\vartheta\xi\right) - \frac{1}{2}(\kappa + \vartheta)e_\theta(\kappa) - \frac{1}{2}\vartheta\vartheta\xi \\ &\quad - \zeta\left(-\frac{1}{2}\kappa^2 - 2\omega\kappa + 2\phi_1\xi + 2(\eta - 3\zeta)\xi - \frac{1}{2}\vartheta^2\right) \\ &\quad + \xi\left(6\rho - \vartheta\vartheta - 2\phi_1\zeta + 2\zeta^2\right) - \kappa e_\theta(\kappa) - 2\omega e_\theta(\kappa) - 6\eta\zeta\xi - 6e_\theta(\zeta\xi) \end{aligned}$$

which is the third desired identity. This concludes the proof of Proposition 2.2.19.

A.6 Proof of Proposition 2.3.4

The proof follows by straightforward calculations using the definition of Ricci coefficients and curvature components with respect to the two frames. Recall the transformation

(2.3.3)

$$\begin{aligned}
e'_4 &= \lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right), \\
e'_\theta &= \left(1 + \frac{1}{2} f \underline{f} \right) e_\theta + \frac{1}{2} \underline{f} e_4 + \frac{1}{2} f \left(1 + \frac{1}{4} f \underline{f} \right) e_3, \\
e'_3 &= \lambda^{-1} \left(\left(1 + \frac{1}{2} f \underline{f} + \frac{1}{16} f^2 \underline{f}^2 \right) e_3 + \underline{f} \left(1 + \frac{1}{4} f \underline{f} \right) e_\theta + \frac{1}{4} \underline{f}^2 e_4 \right).
\end{aligned}$$

We first derive the transformation formulae for κ . We have, under a transformation of type (2.3.3),

$$\begin{aligned}
\chi' &= g(D_{e'_\theta} e'_4, e_{\theta'}) \\
&= g \left(D_{e'_\theta} \left(\lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right) \right), e'_\theta \right) \\
&= \lambda g \left(D_{e'_\theta} \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right), e'_\theta \right) \\
&= \lambda g(D_{e'_\theta} e_4, e'_\theta) + \lambda e'_\theta(f) g(e_\theta, e'_\theta) + \frac{\lambda}{4} e'_\theta(f^2) g(e_3, e'_\theta) + \lambda f g(D_{e'_\theta} e_\theta, e'_\theta) + \frac{\lambda}{4} f^2 g(D_{e'_\theta} e_3, e'_\theta) \\
&= \lambda g(D_{e'_\theta} e_4, e'_\theta) + \lambda \left(1 + \frac{1}{2} f \underline{f} \right) e'_\theta(f) - \frac{\lambda}{4} \underline{f} e'_\theta(f^2) + \lambda f g(D_{e'_\theta} e_\theta, e'_\theta) + \frac{\lambda}{4} f^2 \underline{\chi} + \text{l.o.t.}
\end{aligned}$$

We recall that the lower order terms we denote by l.o.t., here and throughout the proof, are linear with respect $\Gamma = \{\xi, \underline{\xi}, \vartheta, \kappa, \eta, \underline{\eta}, \zeta, \underline{\zeta}, \underline{\vartheta}\}$ and quadratic or higher order in f, \underline{f} , and do not contain derivatives of these latter. We also recall that $\chi = \frac{1}{2}(\kappa + \vartheta)$, $\underline{\chi} = \frac{1}{2}(\underline{\kappa} + \underline{\vartheta})$.

Next, we compute

$$\begin{aligned}
g(D_{e'_\theta} e_4, e'_\theta) &= g \left(D_{e'_\theta} e_4, \left(1 + \frac{1}{2} f \underline{f} \right) e_\theta + \frac{1}{2} f e_3 \right) + \text{l.o.t.} \\
&= \left(1 + \frac{1}{2} f \underline{f} \right) g \left(D_{(1+\frac{1}{2}f\underline{f})e_\theta + \frac{1}{2}\underline{f}e_4 + \frac{1}{2}fe_3} e_4, e_\theta \right) \\
&\quad + \frac{1}{2} f g \left(D_{e_\theta + \frac{1}{2}\underline{f}e_4 + \frac{1}{2}fe_3} e_4, e_3 \right) + \text{l.o.t.} \\
&= (1 + f \underline{f}) \chi + \underline{f} \xi + f \eta + f \zeta + f \underline{f} \omega - f^2 \underline{\omega} + \text{l.o.t.},
\end{aligned}$$

and

$$\begin{aligned}
f g(D_{e'_\theta} e_\theta, e'_\theta) &= \frac{1}{2} f \underline{f} g(D_{e'_\theta} e_\theta, e_4) + \frac{1}{2} f^2 g(D_{e'_\theta} e_\theta, e_3) \\
&= -\frac{1}{2} f \underline{f} \chi - \frac{1}{2} f^2 \underline{\chi} + \text{l.o.t.}
\end{aligned}$$

This yields

$$\begin{aligned}
\chi' &= \lambda g(D_{e'_\theta} e_4, e'_\theta) + \lambda \left(1 + \frac{1}{2} f \underline{f}\right) e'_\theta(f) - \frac{\lambda}{4} \underline{f} e'_\theta(f^2) + \lambda f g(D_{e'_\theta} e_\theta, e'_\theta) + \frac{\lambda}{4} f^2 \underline{\chi} + \text{l.o.t.} \\
&= \lambda \left(\chi + \left(1 + \frac{1}{2} f \underline{f}\right) e'_\theta(f) - \frac{1}{4} \underline{f} e'_\theta(f^2) + f(\zeta + \eta) + \underline{f} \xi - \frac{1}{4} f^2 \underline{\chi} + \frac{1}{2} f \underline{f} \chi + f \underline{f} \omega - f^2 \underline{\omega} \right. \\
&\quad \left. + \text{l.o.t.} \right).
\end{aligned}$$

Hence,

$$\begin{aligned}
\kappa' &= \chi' + e'_4 \Phi = \chi' + \lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right) \Phi \\
&= \lambda \left(\kappa + e'_\theta(f) + e_\theta(\Phi) f + \frac{1}{8} (\underline{\kappa} - \underline{\vartheta}) f^2 + \frac{1}{2} f \underline{f} e'_\theta(f) - \frac{1}{4} \underline{f} e'_\theta(f^2) + f(\zeta + \eta) + \underline{f} \xi \right. \\
&\quad \left. - \frac{1}{4} f^2 \underline{\chi} + \frac{1}{2} f \underline{f} \chi + f \underline{f} \omega - f^2 \underline{\omega} + \text{l.o.t.} \right) \\
&= \lambda \left(\kappa + \phi_1'(f) + \frac{1}{2} f \underline{f} e'_\theta(f) - \frac{1}{4} \underline{f} e'_\theta(f^2) - \frac{1}{4} f^2 \underline{\kappa} + f(\zeta + \eta) + \underline{f} \xi + f \underline{f} \omega - f^2 \underline{\omega} + \text{l.o.t.} \right)
\end{aligned}$$

and

$$\begin{aligned}
\vartheta' &= \chi' - e'_4 \Phi = \chi' - \lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right) \Phi \\
&= \lambda \left(\vartheta + e'_\theta(f) - e_\theta(\Phi) f - \frac{1}{8} (\underline{\kappa} - \underline{\vartheta}) f^2 + \frac{1}{2} f \underline{f} e'_\theta(f) - \frac{1}{4} \underline{f} e'_\theta(f^2) + f(\zeta + \eta) + \underline{f} \xi \right. \\
&\quad \left. - \frac{1}{4} f^2 \underline{\chi} + \frac{1}{2} f \underline{f} \chi + f \underline{f} \omega - f^2 \underline{\omega} + \text{l.o.t.} \right) \\
&= \lambda \left(\vartheta - \phi_2'(f) + \frac{1}{2} f \underline{f} e'_\theta(f) - \frac{1}{4} \underline{f} e'_\theta(f^2) + \frac{1}{4} f \underline{f} \kappa + f(\zeta + \eta) + \underline{f} \xi + f \underline{f} \omega - f^2 \underline{\omega} + \text{l.o.t.} \right).
\end{aligned}$$

This yields

$$\begin{aligned}
\kappa' &= \lambda(\kappa + \phi_1'(f)) + \lambda \text{Err}(\kappa, \kappa'), \\
\text{Err}(\kappa, \kappa') &= \frac{1}{2} f \underline{f} e'_\theta(f) - \frac{1}{4} \underline{f} e'_\theta(f^2) + f(\zeta + \eta) + \underline{f} \xi - \frac{1}{4} f^2 \underline{\kappa} + f \underline{f} \omega - f^2 \underline{\omega} + \text{l.o.t.} \\
&= f(\zeta + \eta) + \underline{f} \xi - \frac{1}{4} f^2 \underline{\kappa} + f \underline{f} \omega - f^2 \underline{\omega} + \text{l.o.t.}
\end{aligned}$$

and

$$\begin{aligned}
\vartheta' &= \lambda(\vartheta - \mathcal{A}_2'(f)) + \lambda \text{Err}(\vartheta, \vartheta'), \\
\text{Err}(\vartheta, \vartheta') &= \frac{1}{2} \underline{f} \underline{f} e'_\theta(f) - \frac{1}{4} \underline{f} e'_\theta(f^2) + f(\zeta + \eta) + \underline{f} \underline{\xi} + \frac{1}{4} \underline{f} \underline{f} \underline{\kappa} + \underline{f} \underline{f} \underline{\omega} - f^2 \underline{\omega} + \text{l.o.t.} \\
&= f(\zeta + \eta) + \underline{f} \underline{\xi} + \frac{1}{4} \underline{f} \underline{f} \underline{\kappa} + \underline{f} \underline{f} \underline{\omega} - f^2 \underline{\omega} + \text{l.o.t.}
\end{aligned}$$

Next, we derive the transformation formula for $\underline{\kappa}$ and $\underline{\vartheta}$. We have, under a transformation of type (2.3.3),

$$\begin{aligned}
\underline{\chi}' &= g(D_{e'_\theta} e'_3, e_{\theta'}) \\
&= g\left(D_{e'_\theta} \left(\lambda^{-1} \left(\left(1 + \frac{1}{2} \underline{f} \underline{f} + \frac{1}{16} \underline{f}^2 \underline{f}^2\right) e_3 + \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f}\right) e_\theta + \frac{1}{4} \underline{f}^2 e_4 \right)\right), e'_\theta\right) \\
&= \lambda^{-1} g\left(D_{e'_\theta} \left(\left(1 + \frac{1}{2} \underline{f} \underline{f} + \frac{1}{16} \underline{f}^2 \underline{f}^2\right) e_3 + \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f}\right) e_\theta + \frac{1}{4} \underline{f}^2 e_4 \right), e'_\theta\right) \\
&= \frac{\lambda^{-1}}{2} e'_\theta \left(\underline{f} \underline{f} + \frac{1}{8} \underline{f}^2 \underline{f}^2 \right) g(e_3, e'_\theta) + \lambda^{-1} e'_\theta \left(\underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f}\right) \right) g(e_\theta, e'_\theta) + \frac{\lambda^{-1}}{4} e'_\theta (\underline{f}^2) g(e_4, e'_\theta) \\
&\quad + \lambda^{-1} \left(1 + \frac{1}{2} \underline{f} \underline{f} + \frac{1}{16} \underline{f}^2 \underline{f}^2\right) g(D_{e'_\theta} e_3, e'_\theta) + \lambda^{-1} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f}\right) g(D_{e'_\theta} e_\theta, e'_\theta) \\
&\quad + \frac{1}{4} \lambda^{-1} \underline{f}^2 g(D_{e'_\theta} e_4, e'_\theta) \\
&= -\frac{\lambda^{-1}}{2} \underline{f} e'_\theta \left(\underline{f} \underline{f} + \frac{1}{8} \underline{f}^2 \underline{f}^2 \right) + \lambda^{-1} \left(1 + \frac{1}{2} \underline{f} \underline{f}\right) e'_\theta \left(\underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f}\right) \right) \\
&\quad - \frac{\lambda^{-1}}{4} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f}\right) e'_\theta (\underline{f}^2) + \lambda^{-1} \left(1 + \frac{1}{2} \underline{f} \underline{f}\right) g(D_{e'_\theta} e_3, e'_\theta) + \lambda^{-1} \underline{f} g(D_{e'_\theta} e_\theta, e'_\theta) \\
&\quad + \frac{1}{4} \lambda^{-1} \underline{f}^2 \chi + \text{l.o.t.}
\end{aligned}$$

Then, we easily derive by symmetry from the formula for $\underline{\kappa}$ and $\underline{\vartheta}$

$$\begin{aligned}
\underline{\kappa}' &= \lambda^{-1} (\underline{\kappa} + \mathcal{A}_1'(f)) + \lambda^{-1} \text{Err}(\underline{\kappa}, \underline{\kappa}'), \\
\text{Err}(\underline{\kappa}, \underline{\kappa}') &= -\frac{1}{2} \underline{f} e'_\theta \left(\underline{f} \underline{f} + \frac{1}{8} \underline{f}^2 \underline{f}^2 \right) + \left(\frac{3}{4} \underline{f} \underline{f} + \frac{1}{8} (\underline{f} \underline{f})^2 \right) e'_\theta(\underline{f}) + \frac{1}{4} \left(1 + \frac{1}{2} \underline{f} \underline{f}\right) \underline{f} e'_\theta(\underline{f} \underline{f}) \\
&\quad - \frac{1}{4} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f}\right) e'_\theta(\underline{f}^2) + \underline{f}(-\zeta + \eta) + \underline{f} \underline{\xi} - \frac{1}{4} \underline{f}^2 \underline{\kappa} + \underline{f} \underline{f} \underline{\omega} - \underline{f}^2 \underline{\omega} + \text{l.o.t.} \\
&= -\frac{1}{4} \underline{f}^2 e'_\theta(\underline{f}) + \underline{f}(-\zeta + \eta) + \underline{f} \underline{\xi} - \frac{1}{4} \underline{f}^2 \underline{\kappa} + \underline{f} \underline{f} \underline{\omega} - \underline{f}^2 \underline{\omega} + \text{l.o.t.}
\end{aligned}$$

and

$$\begin{aligned}
\vartheta' &= \lambda (\vartheta - \phi_2^*(f)) + \lambda^{-1} \text{Err}(\vartheta, \vartheta'), \\
\text{Err}(\vartheta, \vartheta') &= -\frac{1}{2} \underline{f} e'_\theta \left(\underline{f} \underline{f} + \frac{1}{8} \underline{f}^2 \underline{f}^2 \right) + \left(\frac{3}{4} \underline{f} \underline{f} + \frac{1}{8} (\underline{f} \underline{f})^2 \right) e'_\theta(\underline{f}) + \frac{1}{4} \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) \underline{f} e'_\theta(\underline{f} \underline{f}) \\
&\quad - \frac{1}{4} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_\theta(\underline{f}^2) + \underline{f}(-\zeta + \eta) + \underline{f} \underline{\xi} + \frac{1}{4} \underline{f} \underline{f} \underline{\kappa} + \underline{f} \underline{f} \underline{\omega} - \underline{f}^2 \omega + \text{l.o.t.} \\
&= -\frac{1}{4} \underline{f}^2 e'_\theta(\underline{f}) + \underline{f}(-\zeta + \eta) + \underline{f} \underline{\xi} + \frac{1}{4} \underline{f} \underline{f} \underline{\kappa} + \underline{f} \underline{f} \underline{\omega} - \underline{f}^2 \omega + \text{l.o.t.}
\end{aligned}$$

Next, we derive the transformation formula for ζ . We have, under a transformation of type (2.3.3),

$$\begin{aligned}
2\zeta' &= g(D_{e'_\theta} e'_4, e'_3) \\
&= g\left(D_{e'_\theta} \left(\lambda \left(e_4 + \underline{f} e_\theta + \frac{1}{4} \underline{f}^2 e_3 \right) \right), e'_3\right) \\
&= -2e'_\theta(\log(\lambda)) + \lambda g\left(D_{e'_\theta} \left(e_4 + \underline{f} e_\theta + \frac{1}{4} \underline{f}^2 e_3 \right), e'_3\right) \\
&= -2e'_\theta(\log(\lambda)) + \lambda e'_\theta(\underline{f}) g(e_\theta, e'_3) + \frac{1}{4} \lambda e'_\theta(\underline{f}^2) g(e_3, e'_3) + \lambda g(D_{e'_\theta} e_4, e'_3) \\
&\quad + \lambda \underline{f} g(D_{e'_\theta} e_\theta, e'_3) + \frac{1}{4} \lambda \underline{f}^2 g(D_{e'_\theta} e_3, e'_3) \\
&= -2e'_\theta(\log(\lambda)) + \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_\theta(\underline{f}) - \frac{1}{8} \underline{f}^2 e'_\theta(\underline{f}^2) + \lambda g(D_{e'_\theta} e_4, e'_3) + \lambda \underline{f} g(D_{e'_\theta} e_\theta, e'_3) + \text{l.o.t.}
\end{aligned}$$

We compute

$$\begin{aligned}
\lambda g(D_{e'_\theta} e_4, e'_3) &= g(D_{e'_\theta} e_4, e_3 + \underline{f} e_\theta) + \text{l.o.t.} \\
&= g\left(D_{e_\theta + \frac{1}{2} \underline{f} e_4 + \frac{1}{2} \underline{f} e_3} e_4, e_3\right) + \underline{f} g\left(D_{e_\theta + \frac{1}{2} \underline{f} e_4 + \frac{1}{2} \underline{f} e_3} e_4, e_\theta\right) + \text{l.o.t.} \\
&= 2\zeta + 2\omega \underline{f} - 2\underline{\omega} \underline{f} + \underline{f} \underline{\chi} + \text{l.o.t.}
\end{aligned}$$

and

$$\begin{aligned}
\lambda \underline{f} g(D_{e'_\theta} e_\theta, e'_3) &= \underline{f} g(D_{e'_\theta} e_\theta, e_3) + \text{l.o.t.} \\
&= \underline{f} g\left(D_{e_\theta + \frac{1}{2} \underline{f} e_4 + \frac{1}{2} \underline{f} e_3} e_\theta, e_3\right) + \text{l.o.t.} \\
&= -\underline{f} \underline{\chi} + \text{l.o.t.}
\end{aligned}$$

This yields

$$\begin{aligned} 2\zeta' &= -2e'_\theta(\log(\lambda)) + \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_\theta(f) - \frac{1}{8} \underline{f}^2 e'_\theta(f^2) + \lambda g(D_{e'_\theta} e_4, e'_3) + \lambda f g(D_{e'_\theta} e_\theta, e'_3) + \text{l.o.t.} \\ &= 2\zeta - 2e'_\theta(\log(\lambda)) + \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_\theta(f) - \frac{1}{8} \underline{f}^2 e'_\theta(f^2) + 2\omega \underline{f} - 2\underline{\omega} \underline{f} + \underline{f} \chi - \underline{f} \underline{\chi} + \text{l.o.t.} \end{aligned}$$

and hence

$$\begin{aligned} \zeta' &= \zeta - e'_\theta(\log(\lambda)) + \frac{1}{4}(-\underline{f} \underline{\kappa} + \underline{f} \kappa) + \underline{f} \omega - \underline{f} \underline{\omega} + \text{Err}(\zeta, \zeta'), \\ \text{Err}(\zeta, \zeta') &= \frac{1}{2} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_\theta(f) - \frac{1}{16} \underline{f}^2 e'_\theta(f^2) + \frac{1}{4}(-\underline{f} \underline{\vartheta} + \underline{f} \vartheta) + \text{l.o.t.} \\ &= \frac{1}{2} \underline{f} e'_\theta(f) + \frac{1}{4}(-\underline{f} \underline{\vartheta} + \underline{f} \vartheta) + \text{l.o.t.} \end{aligned}$$

Next, we derive the transformation formulae for η . We have, under a transformation of type (2.3.3),

$$\begin{aligned} 2\eta' &= g(D_{e'_3} e'_4, e'_\theta) \\ &= g\left(D_{e'_3} \left(\lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right) \right), e'_\theta\right) \\ &= \lambda g\left(D_{e'_3} \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right), e'_\theta\right) \\ &= \lambda g(D_{e'_3} e_4, e'_\theta) + \lambda e'_3(f) g(e_\theta, e'_\theta) + \lambda f g(D_{e'_3} e_\theta, e'_\theta) + \frac{1}{4} \lambda e'_3(f^2) g(e_3, e'_\theta) + \frac{1}{4} \lambda f^2 g(D_{e'_3} e_3, e'_\theta) \\ &= \lambda \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) e'_3(f) - \frac{1}{4} \lambda \underline{f} e'_3(f^2) + \lambda g(D_{e'_3} e_4, e'_\theta) + \lambda f g(D_{e'_3} e_\theta, e'_\theta) + \text{l.o.t.} \end{aligned}$$

We compute

$$\begin{aligned} \lambda g(D_{e'_3} e_4, e'_\theta) &= \lambda g\left(D_{e'_3} e_4, e_\theta + \frac{1}{2} f e_3\right) + \text{l.o.t.} \\ &= g\left(D_{e_3 + \underline{f} e_\theta} e_4, e_\theta\right) + \frac{1}{2} f g(D_{e_3} e_4, e_3) + \text{l.o.t.} \\ &= 2\eta + \underline{f} \chi - 2\underline{\omega} \underline{f} + \text{l.o.t.} \end{aligned}$$

and

$$\lambda f g(D_{e'_3} e_\theta, e'_\theta) = \text{l.o.t.}$$

This yields

$$\begin{aligned} 2\eta' &= \lambda \left(1 + \frac{1}{2}f\underline{f} \right) e'_3(f) - \frac{1}{4}\lambda\underline{f}e'_3(f^2) + \lambda g(D_{e'_3}e_4, e'_\theta) + \lambda f g(D_{e'_3}e_\theta, e'_\theta) + \text{l.o.t.} \\ &= \lambda \left(1 + \frac{1}{2}f\underline{f} \right) e'_3(f) - \frac{1}{4}\lambda\underline{f}e'_3(f^2) + 2\underline{\eta} + \underline{f}\underline{\chi} - 2\underline{\omega}f + \text{l.o.t.} \end{aligned}$$

and hence

$$\begin{aligned} \eta' &= \eta + \frac{1}{2}\lambda e'_3(f) + \frac{1}{4}\kappa\underline{f} - \underline{f}\underline{\omega} + \text{Err}(\eta, \eta'), \\ \text{Err}(\eta, \eta') &= \frac{1}{4}\lambda\underline{f}e'_3(f) - \frac{1}{8}\lambda\underline{f}e'_3(f^2) + \frac{1}{4}\underline{f}\vartheta + \text{l.o.t.} \\ &= \frac{1}{4}\underline{f}\vartheta + \text{l.o.t.} \end{aligned}$$

Next, we derive the transformation formulae for $\underline{\eta}$. We have, under a transformation of type (2.3.3),

$$\begin{aligned} 2\underline{\eta}' &= g(D_{e'_4}e'_3, e'_\theta) \\ &= g\left(D_{e'_4}\left(\lambda^{-1}\left(\left(1 + \frac{1}{2}f\underline{f} + \frac{1}{16}f^2\underline{f}^2\right)e_3 + \underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)e_\theta + \frac{1}{4}f^2e_4\right)\right), e'_\theta\right) \\ &= \lambda^{-1}g\left(D_{e'_4}\left(\left(1 + \frac{1}{2}f\underline{f} + \frac{1}{16}f^2\underline{f}^2\right)e_3 + \underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)e_\theta + \frac{1}{4}f^2e_4\right), e'_\theta\right) \\ &= \frac{1}{2}\lambda^{-1}e'_4\left(\underline{f}\underline{f} + \frac{1}{8}f^2\underline{f}^2\right)g(e_3, e'_\theta) + \lambda^{-1}\left(1 + \frac{1}{2}f\underline{f} + \frac{1}{16}f^2\underline{f}^2\right)g(D_{e'_4}e_3, e'_\theta) \\ &\quad + \lambda^{-1}e'_4\left(\underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)\right)g(e_\theta, e'_\theta) + \lambda^{-1}\underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)g(D_{e'_4}e_\theta, e'_\theta) \\ &\quad + \frac{1}{4}\lambda^{-1}e'_4(\underline{f}^2)g(e_4, e'_\theta) + \frac{1}{4}\lambda^{-1}\underline{f}^2g(D_{e'_4}e_4, e'_\theta) \\ &= -\frac{1}{2}\lambda^{-1}\underline{f}e'_4\left(\underline{f}\underline{f} + \frac{1}{8}f^2\underline{f}^2\right) + \lambda^{-1}\left(1 + \frac{1}{2}f\underline{f}\right)e'_4\left(\underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)\right) \\ &\quad - \frac{1}{4}\lambda^{-1}\underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)e'_4(\underline{f}^2) + \lambda^{-1}g(D_{e'_4}e_3, e'_\theta) + \lambda^{-1}\underline{f}g(D_{e'_4}e_\theta, e'_\theta) + \text{l.o.t.} \end{aligned}$$

We compute

$$\begin{aligned} \lambda^{-1}g(D_{e'_4}e_3, e'_\theta) &= \lambda^{-1}g\left(D_{e'_4}e_3, e_\theta + \frac{1}{2}\underline{f}e_4\right) + \text{l.o.t.} \\ &= 2\underline{\eta} + \underline{f}\underline{\chi} - 2\underline{f}\underline{\omega} + \text{l.o.t.} \end{aligned}$$

and

$$\lambda^{-1} \underline{f} g (D_{e'_4} e_\theta, e'_\theta) = \text{l.o.t.}$$

This yields

$$\begin{aligned} 2\underline{\eta}' &= -\frac{1}{2} \lambda^{-1} \underline{f} e'_4 \left(\underline{f} \underline{f} + \frac{1}{8} f^2 \underline{f}^2 \right) + \lambda^{-1} \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) e'_4 \left(\underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) \right) \\ &\quad - \frac{1}{4} \lambda^{-1} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_4(\underline{f}^2) + \lambda^{-1} g (D_{e'_4} e_3, e'_\theta) + \lambda^{-1} \underline{f} g (D_{e'_4} e_\theta, e'_\theta) + \text{l.o.t.} \\ &= -\frac{1}{2} \lambda^{-1} \underline{f} e'_4 \left(\underline{f} \underline{f} + \frac{1}{8} f^2 \underline{f}^2 \right) + \lambda^{-1} \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) e'_4 \left(\underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) \right) \\ &\quad - \frac{1}{4} \lambda^{-1} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_4(\underline{f}^2) + 2\underline{\eta} + \underline{f} \underline{\chi} - 2\underline{f} \underline{\omega} + \text{l.o.t.} \end{aligned}$$

and hence

$$\begin{aligned} \underline{\eta}' &= \underline{\eta} + \frac{1}{2} \lambda^{-1} e'_4(\underline{f}) + \frac{1}{4} \underline{\kappa} \underline{f} - \underline{f} \underline{\omega} + \text{Err}(\underline{\eta}, \underline{\eta}'), \\ \text{Err}(\underline{\eta}, \underline{\eta}') &= \frac{1}{4} \lambda^{-1} \underline{f} \underline{f} e'_4(\underline{f}) - \frac{1}{4} \lambda^{-1} \underline{f} e'_4 \left(\underline{f} \underline{f} + \frac{1}{8} f^2 \underline{f}^2 \right) + \lambda^{-1} \frac{1}{8} \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) e'_4(\underline{f} \underline{f}^2) \\ &\quad - \frac{1}{8} \lambda^{-1} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_4(\underline{f}^2) + \frac{1}{4} \underline{f} \underline{\vartheta} + \text{l.o.t.} \\ &= -\frac{1}{8} \underline{f}^2 \lambda^{-1} e'_4(\underline{f}) + \frac{1}{4} \underline{f} \underline{\vartheta} + \text{l.o.t.} \end{aligned}$$

Next, we derive the transformation formulae for ξ . We have, under a transformation of type (2.3.3),

$$\begin{aligned} 2\underline{\xi}' &= g (D_{e'_4} e'_4, e'_\theta) \\ &= g \left(D_{e'_4} \left(\lambda \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right) \right), e'_\theta \right) \\ &= \lambda g \left(D_{e'_4} \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right), e'_\theta \right) \\ &= \lambda g (D_{e'_4} e_4, e'_\theta) + \lambda e'_4(\underline{f}) g (e_\theta, e'_\theta) + \lambda \underline{f} g (D_{e'_4} e_\theta, e'_\theta) + \frac{1}{4} \lambda e'_4(\underline{f}^2) g (e_3, e'_\theta) + \frac{1}{4} \lambda \underline{f}^2 g (D_{e'_4} e_3, e'_\theta) \\ &= \lambda \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) e'_4(\underline{f}) - \frac{1}{4} \lambda \underline{f} e'_4(\underline{f}^2) + \lambda g (D_{e'_4} e_4, e'_\theta) + \lambda \underline{f} g (D_{e'_4} e_\theta, e'_\theta) + \text{l.o.t.} \end{aligned}$$

We compute

$$\begin{aligned}
\lambda g(D_{e'_4} e_4, e'_\theta) &= \lambda g\left(D_{e'_4} e_4, e_\theta + \frac{1}{2} f e_3\right) + \text{l.o.t.} \\
&= \lambda^2 g(D_{e_4 + f e_\theta} e_4, e_\theta) + \frac{1}{2} \lambda^2 f g(D_{e_4} e_4, e_3) + \text{l.o.t.} \\
&= 2\lambda^2 \xi + \lambda^2 f \chi + 2\lambda^2 f \omega + \text{l.o.t.}
\end{aligned}$$

and

$$\lambda f g(D_{e'_4} e_\theta, e'_\theta) = \text{l.o.t.}$$

This yields

$$\begin{aligned}
2\xi' &= \lambda \left(1 + \frac{1}{2} f \underline{f}\right) e'_4(f) - \frac{1}{4} \lambda \underline{f} e'_4(f^2) + \lambda g(D_{e'_4} e_4, e'_\theta) + \lambda f g(D_{e'_4} e_\theta, e'_\theta) + \text{l.o.t.} \\
&= \lambda \left(1 + \frac{1}{2} f \underline{f}\right) e'_4(f) - \frac{1}{4} \lambda \underline{f} e'_4(f^2) + 2\lambda^2 \xi + \lambda^2 f \chi + 2\lambda^2 f \omega + \text{l.o.t.}
\end{aligned}$$

and hence

$$\begin{aligned}
\xi' &= \lambda^2 \left(\xi + \frac{1}{2} \lambda^{-1} e'_4(f) + \omega f + \frac{1}{4} f \kappa\right) + \lambda^2 \text{Err}(\xi, \xi'), \\
\text{Err}(\xi, \xi') &= \frac{1}{4} \lambda^{-1} f \underline{f} e'_4(f) - \frac{1}{8} \lambda^{-1} \underline{f} e'_4(f^2) + \frac{1}{4} f \vartheta + \text{l.o.t.} \\
&= \frac{1}{4} f \vartheta + \text{l.o.t.}
\end{aligned}$$

In the particular case when $\lambda = 1$, $\underline{f} = 0$, see Remark 2.3.5, the error term takes the form,

$$\text{Err}(\xi, \xi') = \frac{1}{4} f \vartheta + \frac{1}{4} f^2 (\eta + 2\zeta - \underline{\eta}) - \frac{1}{4} f^3 \left(\underline{\omega} + \frac{1}{2} \underline{\chi}\right) - \frac{3}{32} f^4 \underline{\xi}.$$

Next, we derive the transformation formulae for $\underline{\xi}$. We have, under a transformation of

type (2.3.3),

$$\begin{aligned}
2\underline{\xi}' &= g(D_{e'_3}e'_3, e'_\theta) \\
&= g\left(D_{e'_3}\left(\lambda^{-1}\left(\left(1 + \frac{1}{2}f\underline{f} + \frac{1}{16}f^2\underline{f}^2\right)e_3 + \underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)e_\theta + \frac{1}{4}\underline{f}^2e_4\right)\right), e'_\theta\right) \\
&= \lambda^{-1}g\left(D_{e'_3}\left(\left(1 + \frac{1}{2}f\underline{f} + \frac{1}{16}f^2\underline{f}^2\right)e_3 + \underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)e_\theta + \frac{1}{4}\underline{f}^2e_4\right), e'_\theta\right) \\
&= \frac{1}{2}\lambda^{-1}e'_3\left(\underline{f}\underline{f} + \frac{1}{8}f^2\underline{f}^2\right)g(e_3, e'_\theta) + \lambda^{-1}g(D_{e'_3}e_3, e'_\theta) \\
&\quad + \lambda^{-1}e'_3\left(\underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)\right)g(e_\theta, e'_\theta) + \lambda^{-1}\underline{f}g(D_{e'_3}e_\theta, e'_\theta) + \frac{1}{4}\lambda^{-1}e'_3(\underline{f}^2)g(e_4, e'_\theta) + \text{l.o.t.} \\
&= -\frac{1}{2}\lambda^{-1}\underline{f}e'_3\left(\underline{f}\underline{f} + \frac{1}{8}f^2\underline{f}^2\right) + \lambda^{-1}\left(1 + \frac{1}{2}f\underline{f}\right)e'_3\left(\underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)\right) \\
&\quad - \frac{1}{4}\lambda^{-1}f\left(1 + \frac{1}{4}f\underline{f}\right)e'_3(\underline{f}^2) + \lambda^{-1}g(D_{e'_3}e_3, e'_\theta) + \lambda^{-1}\underline{f}g(D_{e'_3}e_\theta, e'_\theta) + \text{l.o.t.}
\end{aligned}$$

We compute

$$\begin{aligned}
\lambda^{-1}g(D_{e'_3}e_3, e'_\theta) &= \lambda^{-1}g\left(D_{e'_3}e_3, e_\theta + \frac{1}{2}\underline{f}e_4\right) + \text{l.o.t.} \\
&= \lambda^{-2}g\left(D_{e_3+\underline{f}e_\theta}e_3, e_\theta\right) + \frac{1}{2}\lambda^{-2}\underline{f}g(D_{e_3}e_3, e_4) + \text{l.o.t.} \\
&= 2\lambda^{-2}\underline{\xi} + \lambda^{-2}\underline{f}\underline{\chi} + 2\lambda^{-2}\underline{f}\underline{\omega} + \text{l.o.t.}
\end{aligned}$$

and

$$\lambda^{-1}\underline{f}g(D_{e'_3}e_\theta, e'_\theta) = \text{l.o.t.}$$

This yields

$$\begin{aligned}
2\underline{\xi}' &= -\frac{1}{2}\lambda^{-1}\underline{f}e'_3\left(\underline{f}\underline{f} + \frac{1}{8}f^2\underline{f}^2\right) + \lambda^{-1}\left(1 + \frac{1}{2}f\underline{f}\right)e'_3\left(\underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)\right) \\
&\quad - \frac{1}{4}\lambda^{-1}f\left(1 + \frac{1}{4}f\underline{f}\right)e'_3(\underline{f}^2) + \lambda^{-1}g(D_{e'_3}e_3, e'_\theta) + \lambda^{-1}\underline{f}g(D_{e'_3}e_\theta, e'_\theta) + \text{l.o.t.} \\
&= -\frac{1}{2}\lambda^{-1}\underline{f}e'_3\left(\underline{f}\underline{f} + \frac{1}{8}f^2\underline{f}^2\right) + \lambda^{-1}\left(1 + \frac{1}{2}f\underline{f}\right)e'_3\left(\underline{f}\left(1 + \frac{1}{4}f\underline{f}\right)\right) \\
&\quad - \frac{1}{4}\lambda^{-1}f\left(1 + \frac{1}{4}f\underline{f}\right)e'_3(\underline{f}^2) + 2\lambda^{-2}\underline{\xi} + \lambda^{-2}\underline{f}\underline{\chi} + 2\lambda^{-2}\underline{f}\underline{\omega} + \text{l.o.t.}
\end{aligned}$$

and hence

$$\begin{aligned}
\underline{\xi}' &= \lambda^{-2} \left(\underline{\xi} + \frac{1}{2} \lambda e'_3(\underline{f}) + \underline{\omega} \underline{f} + \frac{1}{4} \underline{f} \underline{\kappa} \right) + \lambda^{-2} \text{Err}(\underline{\xi}, \underline{\xi}'), \\
\text{Err}(\underline{\xi}, \underline{\xi}') &= -\frac{1}{4} \lambda \underline{f} e'_3 \left(\underline{f} \underline{f} + \frac{1}{8} \underline{f}^2 \underline{f}^2 \right) + \frac{1}{4} \lambda \underline{f} e'_3(\underline{f}) + \frac{1}{8} \lambda \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) e'_3(\underline{f} \underline{f}^2) \\
&\quad - \frac{1}{8} \lambda \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_3(\underline{f}^2) + \frac{1}{4} \underline{f} \underline{\vartheta} + \text{l.o.t.} \\
&= -\frac{1}{8} \lambda \underline{f}^2 e'_3(\underline{f}) + \frac{1}{4} \underline{f} \underline{\vartheta} + \text{l.o.t.}
\end{aligned}$$

Next, we derive the transformation formulae for ω . We have, under a transformation of type (2.3.3),

$$\begin{aligned}
4\omega' &= g(D_{e'_4} e'_4, e'_3) \\
&= g \left(D_{e'_4} \left(\lambda \left(e_4 + \underline{f} e_\theta + \frac{1}{4} \underline{f}^2 e_3 \right) \right), e'_3 \right) \\
&= -2e'_4(\log \lambda) + \lambda g \left(D_{e'_4} \left(e_4 + \underline{f} e_\theta + \frac{1}{4} \underline{f}^2 e_3 \right), e'_3 \right) \\
&= -2e'_4(\log \lambda) + \lambda g(D_{e'_4} e_4, e'_3) + \lambda e'_4(\underline{f}) g(e_\theta, e'_3) + \lambda \underline{f} g(D_{e'_4} e_\theta, e'_3) \\
&\quad + \frac{1}{4} \lambda e'_4(\underline{f}^2) g(e_3, e'_3) + \frac{1}{4} \lambda \underline{f}^2 g(D_{e'_4} e_3, e'_3) \\
&= -2e'_4(\log \lambda) + \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_4(\underline{f}) - \frac{1}{8} \underline{f}^2 e'_4(\underline{f}^2) + \lambda g(D_{e'_4} e_4, e'_3) + \lambda \underline{f} g(D_{e'_4} e_\theta, e'_3) + \text{l.o.t.}
\end{aligned}$$

We compute

$$\begin{aligned}
\lambda g(D_{e'_4} e_4, e'_3) &= g \left(D_{e'_4} e_4, \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) e_3 + \underline{f} e_\theta \right) + \text{l.o.t.} \\
&= \lambda \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) g \left(D_{e_4 + \underline{f} e_\theta + \frac{1}{4} \underline{f}^2 e_3} e_4, e_3 \right) + \lambda \underline{f} g(D_{e_4 + \underline{f} e_\theta} e_4, e_\theta) + \text{l.o.t.} \\
&= 4\lambda \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) \omega + 2\lambda \underline{f} \zeta - \lambda \underline{f}^2 \underline{\omega} + 2\lambda \underline{f} \underline{\xi} + \lambda \underline{f} \underline{f} \underline{\chi} + \text{l.o.t.}
\end{aligned}$$

and

$$\begin{aligned}
\lambda \underline{f} g(D_{e'_4} e_\theta, e'_3) &= \underline{f} g(D_{e'_4} e_\theta, e_3) + \text{l.o.t.} \\
&= \lambda \underline{f} g(D_{e_4 + \underline{f} e_\theta} e_\theta, e_3) + \text{l.o.t.} \\
&= -2\lambda \underline{f} \underline{\eta} - \lambda \underline{f}^2 \underline{\chi} + \text{l.o.t.}
\end{aligned}$$

This yields

$$\begin{aligned}
4\omega' &= -2e'_4(\log \lambda) + \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_4(f) - \frac{1}{8} \underline{f}^2 e'_4(f^2) + \lambda g(D_{e'_4} e_4, e'_3) + \lambda f g(D_{e'_4} e_\theta, e'_3) + \text{l.o.t.} \\
&= -2e'_4(\log \lambda) + \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_4(f) - \frac{1}{8} \underline{f}^2 e'_4(f^2) + 4\lambda \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) \omega + 2\lambda f \zeta - \lambda f^2 \underline{\omega} \\
&\quad + 2\lambda \underline{f} \xi + \lambda \underline{f} \chi - 2\lambda \underline{f} \eta - \lambda f^2 \underline{\chi} + \text{l.o.t.}
\end{aligned}$$

and hence

$$\begin{aligned}
\omega' &= \lambda \left(\omega - \frac{1}{2} \lambda^{-1} e'_4(\log(\lambda)) \right) + \lambda \text{Err}(\omega, \omega'), \\
\text{Err}(\omega, \omega') &= \frac{1}{4} \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e'_4(f) - \frac{1}{32} \underline{f}^2 e'_4(f^2) + \frac{1}{2} \omega \underline{f} \underline{f} - \frac{1}{2} \underline{f} \eta + \frac{1}{2} \underline{f} \xi + \frac{1}{2} \underline{f} \zeta \\
&\quad - \frac{1}{8} \underline{\kappa} \underline{f}^2 + \frac{1}{8} \underline{f} \underline{f} \underline{\kappa} - \frac{1}{4} \underline{\omega} \underline{f}^2 + \text{l.o.t.} \\
&= \frac{1}{4} \underline{f} e'_4(f) + \frac{1}{2} \omega \underline{f} \underline{f} - \frac{1}{2} \underline{f} \eta + \frac{1}{2} \underline{f} \xi + \frac{1}{2} \underline{f} \zeta - \frac{1}{8} \underline{\kappa} \underline{f}^2 + \frac{1}{8} \underline{f} \underline{f} \underline{\kappa} - \frac{1}{4} \underline{\omega} \underline{f}^2 + \text{l.o.t.}
\end{aligned}$$

In the particular case, see Remark 2.3.5, when $\lambda = 1$, $\underline{f} = 0$ we have the more precise formula,

$$\omega' = \omega + \frac{1}{2} f(\zeta - \eta) - \frac{1}{8} f^2 (2\omega + \underline{\kappa} + \underline{\vartheta} + f \underline{\xi})$$

Next, we derive the transformation formulae for $\underline{\omega}$. We have, under a transformation of type (2.3.3),

$$\begin{aligned}
4\underline{\omega}' &= g(D_{e'_3} e'_3, e'_4) \\
&= g \left(D_{e'_3} \left(\lambda^{-1} \left(\left(1 + \frac{1}{2} \underline{f} \underline{f} + \frac{1}{16} \underline{f}^2 \underline{f}^2 \right) e_3 + \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e_\theta + \frac{1}{4} \underline{f}^2 e_4 \right) \right), e'_4 \right) \\
&= 2e'_3(\log(\lambda)) \\
&\quad + \lambda^{-1} g \left(D_{e'_3} \left(\left(1 + \frac{1}{2} \underline{f} \underline{f} + \frac{1}{16} \underline{f}^2 \underline{f}^2 \right) e_3 + \underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) e_\theta + \frac{1}{4} \underline{f}^2 e_4 \right), e'_4 \right) \\
&= 2e'_3(\log(\lambda)) + \frac{1}{2} \lambda^{-1} e'_3 \left(\underline{f} \underline{f} + \frac{1}{8} \underline{f}^2 \underline{f}^2 \right) g(e_3, e'_4) + \lambda^{-1} \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) g(D_{e'_3} e_3, e'_4) \\
&\quad + \lambda^{-1} e'_3 \left(\underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) \right) g(e_\theta, e'_4) + \lambda^{-1} \underline{f} g(D_{e'_3} e_\theta, e'_4) + \frac{1}{4} \lambda^{-1} e'_3(\underline{f}^2) g(e_4, e'_4) + \text{l.o.t.} \\
&= 2e'_3(\log(\lambda)) - e'_3 \left(\underline{f} \underline{f} + \frac{1}{8} \underline{f}^2 \underline{f}^2 \right) + \underline{f} e'_3 \left(\underline{f} \left(1 + \frac{1}{4} \underline{f} \underline{f} \right) \right) - \frac{1}{8} \underline{f}^2 e'_3(\underline{f}^2) \\
&\quad + \lambda^{-1} \left(1 + \frac{1}{2} \underline{f} \underline{f} \right) g(D_{e'_3} e_3, e'_4) + \lambda^{-1} \underline{f} g(D_{e'_3} e_\theta, e'_4) + \text{l.o.t.}
\end{aligned}$$

We compute

$$\begin{aligned}
\lambda^{-1} \left(1 + \frac{1}{2} f \underline{f} \right) g(D_{e'_3} e_3, e'_4) &= \left(1 + \frac{1}{2} f \underline{f} \right) g(D_{e'_3} e_3, e_4 + f e_\theta) \\
&= \lambda^{-1} \left(1 + \frac{1}{2} f \underline{f} \right) g \left(D_{(1+\frac{1}{2} f \underline{f}) e_3 + f e_\theta + \frac{1}{4} f^2 e_4} e_3, e_4 + f e_\theta \right) + \text{l.o.t.} \\
&= 4\lambda^{-1} \left(1 + \frac{1}{2} f \underline{f} \right) \underline{\omega} - 2\lambda^{-1} f \underline{\zeta} - \lambda^{-1} f^2 \underline{\omega} + 2\lambda^{-1} f \underline{f} \underline{\omega} \\
&\quad + 2\lambda^{-1} f \underline{\xi} + \lambda^{-1} f \underline{f} \underline{\chi} + \text{l.o.t.}
\end{aligned}$$

and

$$\begin{aligned}
\lambda^{-1} \underline{f} g(D_{e'_3} e_\theta, e'_4) &= \underline{f} g(D_{e'_3} e_\theta, e_4) + \text{l.o.t.} \\
&= \lambda^{-1} \underline{f} g(D_{e_3 + f e_\theta} e_\theta, e_4) + \text{l.o.t.} \\
&= -2\lambda^{-1} f \underline{\eta} - \lambda^{-1} f^2 \underline{\chi} + \text{l.o.t.}
\end{aligned}$$

This yields

$$\begin{aligned}
4\underline{\omega}' &= 2e'_3(\log(\lambda)) - e'_3 \left(f \underline{f} + \frac{1}{8} f^2 \underline{f}^2 \right) + f e'_3 \left(\underline{f} \left(1 + \frac{1}{4} f \underline{f} \right) \right) - \frac{1}{8} f^2 e'_3(f^2) \\
&\quad + 4\lambda^{-1} \left(1 + \frac{1}{2} f \underline{f} \right) \underline{\omega} - 2\lambda^{-1} f \underline{\zeta} - \lambda^{-1} f^2 \underline{\omega} + 2\lambda^{-1} f \underline{f} \underline{\omega} \\
&\quad + 2\lambda^{-1} f \underline{\xi} + \lambda^{-1} f \underline{f} \underline{\chi} - 2\lambda^{-1} f \underline{\eta} - \lambda^{-1} f^2 \underline{\chi} + \text{l.o.t.}
\end{aligned}$$

and hence

$$\begin{aligned}
\underline{\omega}' &= \lambda^{-1} \left(\underline{\omega} + \frac{1}{2} \lambda e'_3(\log(\lambda)) \right) + \lambda^{-1} \text{Err}(\underline{\omega}, \underline{\omega}'), \\
\text{Err}(\underline{\omega}, \underline{\omega}') &= -\frac{1}{4} e'_3 \left(f \underline{f} + \frac{1}{8} f^2 \underline{f}^2 \right) + \frac{1}{4} f e'_3 \left(\underline{f} \left(1 + \frac{1}{4} f \underline{f} \right) \right) - \frac{1}{32} f^2 e'_3(f^2) \\
&\quad + \underline{\omega} f \underline{f} - \frac{1}{2} f \underline{\eta} + \frac{1}{2} f \underline{\xi} - \frac{1}{2} f \underline{\zeta} - \frac{1}{8} \kappa f^2 + \frac{1}{8} f \underline{f} \underline{\kappa} - \frac{1}{4} \omega f^2 + \text{l.o.t.} \\
&= -\frac{1}{4} f e'_3(f) + \underline{\omega} f \underline{f} - \frac{1}{2} f \underline{\eta} + \frac{1}{2} f \underline{\xi} - \frac{1}{2} f \underline{\zeta} - \frac{1}{8} \kappa f^2 + \frac{1}{8} f \underline{f} \underline{\kappa} - \frac{1}{4} \omega f^2 + \text{l.o.t.}
\end{aligned}$$

Next we derive the formula for α . We have

$$\begin{aligned}
\alpha' &= R(e'_4, e'_4) = \lambda^2 R \left(e_4 + f e_\theta + \frac{1}{4} f^2 e_3, e_4 + f e_\theta + \frac{1}{4} f^2 e_3 \right) \\
&= \lambda^2 \left(R_{44} + 2f R_{4\theta} + f^2 R_{\theta\theta} + \frac{1}{2} f^2 R_{34} \right) + \text{l.o.t.} \\
&= \lambda^2 \left(\alpha + 2f\beta + \frac{3}{2} f^2 \rho \right) + \text{l.o.t.}
\end{aligned}$$

and hence

$$\begin{aligned}\alpha' &= \lambda^2\alpha + \lambda^2\text{Err}(\alpha, \alpha'), \\ \text{Err}(\alpha, \alpha') &= 2f\beta + \frac{3}{2}f^2\rho + \text{l.o.t.}\end{aligned}$$

The formula for $\underline{\alpha}$ is easily derived by symmetry from the one on α .

Next we derive the formula for β . We have

$$\begin{aligned}\beta' &= R(e'_4, e'_\theta) = \lambda R\left(e_4 + fe_\theta + \frac{1}{4}f^2e_3, \left(1 + \frac{1}{2}f\underline{f}\right)e_\theta + \frac{1}{2}(\underline{f}e_4 + fe_3)\right) + \text{l.o.t.} \\ &= \lambda\left(R_{4\theta} + fR_{\theta\theta} + \frac{1}{2}\underline{f}R_{44} + \frac{1}{2}fR_{43}\right) + \text{l.o.t.} \\ &= \lambda\left(\beta + \frac{3}{2}f\rho + \frac{1}{2}\underline{f}\alpha\right) + \text{l.o.t.}\end{aligned}$$

and hence

$$\begin{aligned}\beta' &= \lambda\left(\beta + \frac{3}{2}f\rho\right) + \lambda\text{Err}(\beta, \beta'), \\ \text{Err}(\beta, \beta') &= \frac{1}{2}\underline{f}\alpha + \text{l.o.t.}\end{aligned}$$

The formula for $\underline{\beta}$ is easily derived by symmetry from the one on β .

Finally, we derive the formula for ρ . We have

$$\begin{aligned}\rho' &= R(e'_4, e'_3) = R\left(e_4 + fe_\theta + \frac{1}{4}f^2e_3, \left(1 + \frac{1}{2}f\underline{f}\right)e_3 + \underline{f}e_\theta + \frac{1}{4}\underline{f}^2e_4\right) + \text{l.o.t.} \\ &= R_{43} + \frac{1}{2}f\underline{f}R_{43} + \underline{f}R_{4\theta} + fR_{\theta 3} + f\underline{f}R_{\theta\theta} + \text{l.o.t.} \\ &= \rho + \frac{3}{2}\rho f\underline{f} + \underline{f}\beta + f\underline{\beta} + \text{l.o.t.}\end{aligned}$$

and hence

$$\begin{aligned}\rho' &= \rho + \text{Err}(\rho, \rho'), \\ \text{Err}(\rho, \rho') &= \frac{3}{2}\rho f\underline{f} + \underline{f}\beta + f\underline{\beta} + \text{l.o.t.}\end{aligned}$$

This concludes the proof of Proposition 2.3.4.

A.7 Proof of Lemma 2.3.6

For ξ' and ω' , we need more precise transformation formula than the ones of Proposition 2.3.4. We have

$$\begin{aligned}
2\xi' &= g(D_{e'_4}e'_4, e'_\theta) \\
&= \lambda^2 g(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e'_\theta) \\
&= \lambda^2 g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_\theta + \frac{1}{2}fe_3\right) + \lambda^2 \frac{f}{2} g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_4 + fe_\theta + \frac{1}{4}f^2e_3\right) \\
&= \lambda^2 g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_\theta + \frac{1}{2}fe_3\right) + \lambda^2 \frac{f}{2} g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), \lambda^{-1}e'_4\right) \\
&= \lambda^2 g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_\theta + \frac{1}{2}fe_3\right).
\end{aligned}$$

Also, we have

$$\begin{aligned}
4\omega' &= g(D_{e'_4}e'_4, e'_3) \\
&= -2e'_4(\log(\lambda)) + \lambda g(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), \lambda e'_3) \\
&= -2e'_4(\log(\lambda)) + \lambda g(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_3) + \lambda \underline{f} g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_\theta + \frac{1}{2}fe_3\right) \\
&\quad + \frac{1}{4}\lambda \underline{f}^2 g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_4 + fe_\theta + \frac{1}{4}f^2e_3\right) \\
&= -2e'_4(\log(\lambda)) + \lambda g(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_3) + \lambda \underline{f} g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_\theta + \frac{1}{2}fe_3\right) \\
&\quad + \frac{1}{4}\lambda \underline{f}^2 g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), \lambda^{-1}e'_4\right) \\
&= -2e'_4(\log(\lambda)) + \lambda g(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_3) + \lambda \underline{f} g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_\theta + \frac{1}{2}fe_3\right).
\end{aligned}$$

In view of the change of frame formula for ξ' , we infer

$$4\omega' = -2e'_4(\log(\lambda)) + \lambda g(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_3) + \lambda^{-1}\underline{f}\xi'.$$

Next, we compute

$$\begin{aligned}
g(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_\theta) &= g\left(D_{\lambda^{-1}e'_4}\left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right), e_\theta\right) \\
&= g(D_{\lambda^{-1}e'_4}e_4, e_\theta) + \lambda^{-1}e'_4(f) + \frac{1}{4}f^2g(D_{\lambda^{-1}e'_4}e_3, e_\theta) \\
&= g\left(D_{e_4+fe_\theta+\frac{1}{4}f^2e_3}e_4, e_\theta\right) + \left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right) f \\
&\quad + \frac{1}{4}f^2g\left(D_{e_4+fe_\theta+\frac{1}{4}f^2e_3}e_3, e_\theta\right) \\
&= 2\xi + f\chi + \frac{1}{2}f^2\eta + \left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right) f + \frac{1}{2}f^2\underline{\eta} + \frac{1}{4}f^3\underline{\chi} \\
&\quad + \frac{1}{8}f^4\underline{\xi}.
\end{aligned}$$

Also, we have

$$\begin{aligned}
g(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_3) &= g\left(D_{\lambda^{-1}e'_4}\left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right), e_3\right) \\
&= g(D_{\lambda^{-1}e'_4}e_4, e_3) + fg(D_{\lambda^{-1}e'_4}e_\theta, e_3) \\
&= g\left(D_{e_4+fe_\theta+\frac{1}{4}f^2e_3}e_4, e_3\right) + fg\left(D_{e_4+fe_\theta+\frac{1}{4}f^2e_3}e_\theta, e_3\right) \\
&= 4\omega + 2f\zeta - f^2\underline{\omega} - 2\underline{\eta}f - f^2\underline{\chi} - \frac{1}{2}f^3\underline{\xi}.
\end{aligned}$$

We deduce

$$\begin{aligned}
2\xi' &= \lambda^2g\left(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_\theta + \frac{1}{2}fe_3\right) \\
&= \lambda^2\left\{2\xi + f\chi + \frac{1}{2}f^2\eta + \left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right) f + \frac{1}{2}f^2\underline{\eta} + \frac{1}{4}f^3\underline{\chi} \right. \\
&\quad \left. + \frac{1}{8}f^4\underline{\xi} + \frac{1}{2}f\left(4\omega + 2f\zeta - f^2\underline{\omega} - 2\underline{\eta}f - f^2\underline{\chi} - \frac{1}{2}f^3\underline{\xi}\right)\right\} \\
&= \lambda^2\left\{2\xi + \left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right) f + f\chi + 2f\omega + \frac{1}{2}f^2\eta - \frac{1}{2}f^2\underline{\eta} + f^2\zeta - \frac{1}{4}f^3\underline{\chi} \right. \\
&\quad \left. - \frac{1}{2}f^3\underline{\omega} - \frac{1}{8}f^4\underline{\xi}\right\}
\end{aligned}$$

and

$$\begin{aligned} 4\omega' &= -2e'_4(\log(\lambda)) + \lambda g(D_{\lambda^{-1}e'_4}(\lambda^{-1}e'_4), e_3) + \lambda^{-1}f\underline{\xi}' \\ &= \lambda \left\{ 4\omega - 2 \left(e_4 + fe_\theta + \frac{1}{4}f^2e_3 \right) \log(\lambda) + 2f\underline{\zeta} - f^2\underline{\omega} - 2\underline{\eta}f - f^2\underline{\chi} - \frac{1}{2}f^3\underline{\xi} \right\} \\ &\quad + \lambda^{-1}f\underline{\xi}'. \end{aligned}$$

If $\xi' = 0$, we infer

$$\begin{aligned} 2\underline{\xi} + \left(e_4 + fe_\theta + \frac{1}{4}f^2e_3 \right) f + f\underline{\chi} + 2f\underline{\omega} + \frac{1}{2}f^2\underline{\eta} - \frac{1}{2}f^2\underline{\eta} + f^2\underline{\zeta} - \frac{1}{4}f^3\underline{\chi} \\ - \frac{1}{2}f^3\underline{\omega} - \frac{1}{8}f^4\underline{\xi} = 0 \end{aligned}$$

and hence

$$\begin{aligned} \lambda^{-1}e'_4(f) + \left(\frac{\kappa}{2} + 2\omega \right) f &= -2\underline{\xi} - \frac{1}{2}\vartheta f - \frac{1}{2}f^2\underline{\eta} + \frac{1}{2}f^2\underline{\eta} - f^2\underline{\zeta} + \frac{1}{8}f^3\underline{\kappa} + \frac{1}{2}f^3\underline{\omega} \\ &\quad + \frac{1}{8}f^3\underline{\vartheta} + \frac{1}{8}f^4\underline{\xi} \end{aligned}$$

which yields the desired transport equation for f

$$\begin{aligned} \lambda^{-1}e'_4(f) + \left(\frac{\kappa}{2} + 2\omega \right) f &= -2\underline{\xi} + E_1(f, \Gamma), \\ E_1(f, \Gamma) &= -\frac{1}{2}\vartheta f - \frac{1}{2}f^2\underline{\eta} + \frac{1}{2}f^2\underline{\eta} - f^2\underline{\zeta} + \frac{1}{8}f^3\underline{\kappa} + \frac{1}{2}f^3\underline{\omega} + \frac{1}{8}f^3\underline{\vartheta} + \frac{1}{8}f^4\underline{\xi}. \end{aligned}$$

Also, if $\xi' = 0$ and $\omega' = 0$, we infer

$$0 = 4\omega - 2 \left(e_4 + fe_\theta + \frac{1}{4}f^2e_3 \right) \log(\lambda) + 2f\underline{\zeta} - f^2\underline{\omega} - 2\underline{\eta}f - f^2\underline{\chi} - \frac{1}{2}f^3\underline{\xi}$$

and hence

$$\lambda^{-1}e'_4(\log(\lambda)) = 2\omega + f\underline{\zeta} - \frac{1}{2}f^2\underline{\omega} - \underline{\eta}f - \frac{1}{4}f^2\underline{\kappa} - \frac{1}{4}f^2\underline{\vartheta} - \frac{1}{4}f^3\underline{\xi}$$

which yields the desired transport equation for $\log(\lambda)$

$$\begin{aligned} \lambda^{-1}e'_4(\log(\lambda)) &= 2\omega + E_2(f, \Gamma), \\ E_2(f, \Gamma) &= f\underline{\zeta} - \frac{1}{2}f^2\underline{\omega} - \underline{\eta}f - \frac{1}{4}f^2\underline{\kappa} - \frac{1}{4}f^2\underline{\vartheta} - \frac{1}{4}f^3\underline{\xi}. \end{aligned}$$

Finally, we derive the transport equation for \underline{f} . In view of the transformation formulas of Proposition 2.3.4 for $\underline{\zeta}'$ and $\underline{\eta}'$, and the fact that we assume $\underline{\zeta}' + \underline{\eta}' = 0$, we have

$$\begin{aligned} \frac{1}{2}\lambda^{-1}e'_4(\underline{f}) &= -(\underline{\zeta} + \underline{\eta}) + e'_\theta(\log(\lambda)) - \frac{1}{4}\underline{f}\kappa + \underline{f}\underline{\omega} - \frac{1}{2}\underline{f}e'_\theta(\underline{f}) + \frac{1}{8}\underline{f}^2\lambda^{-1}e'_4(\underline{f}) \\ &\quad - \frac{1}{4}\underline{f}\vartheta + \text{l.o.t.} \end{aligned}$$

Together with the above identity for $\lambda^{-1}e'_4(\underline{f})$, we infer

$$\begin{aligned} \lambda^{-1}e'_4(\underline{f}) + \frac{\kappa}{2}\underline{f} &= -2(\underline{\zeta} + \underline{\eta}) + 2e'_\theta(\log(\lambda)) + 2\underline{f}\underline{\omega} + E_3(\underline{f}, \underline{f}, \Gamma), \\ E_3(\underline{f}, \underline{f}, \Gamma) &= -\underline{f}e'_\theta(\underline{f}) - \frac{1}{2}\underline{f}\vartheta + \text{l.o.t.}, \end{aligned}$$

which yields the third identity of the statement. This concludes the proof of Lemma 2.3.6.

A.8 Proof of Corollary 2.3.7

In view of Lemma 2.3.6 and the fact that (e_3, e_4, e_θ) emanates from an outgoing geodesic foliation and hence

$$\xi = 0, \quad \omega = 0, \quad \zeta + \eta = 0,$$

we have

$$\begin{aligned} \lambda^{-1}e'_4(\underline{f}) + \frac{\kappa}{2}\underline{f} &= E_1(\underline{f}, \Gamma), \\ \lambda^{-1}e'_4(\log(\lambda)) &= E_2(\underline{f}, \Gamma), \\ \lambda^{-1}e'_4(\underline{f}) + \frac{\kappa}{2}\underline{f} &= 2e'_\theta(\log(\lambda)) + 2\underline{f}\underline{\omega} + E_3(\underline{f}, \underline{f}, \Gamma). \end{aligned}$$

The second equation is the desired identity for $\log(\lambda)$.

We still need to derive the first and the third identities. We start with the first one. We have

$$\begin{aligned} \lambda^{-1}e'_4(rf) &= r\left(-\frac{\kappa}{2}f + E_1(\underline{f}, \Gamma)\right) + \lambda^{-1}e'_4(r)f \\ &= -\frac{r}{2}\left(\kappa - \frac{2\lambda^{-1}e'_4(r)}{r}\right)f + rE_1(\underline{f}, \Gamma). \end{aligned}$$

$$\lambda^{-1}e'_4 = e_4 + fe_\theta + \frac{f^2}{4}e_3,$$

we infer

$$\lambda^{-1}e'_4(r) = \frac{r}{2}\bar{\kappa} + \frac{f^2}{4}e_3(r)$$

and hence

$$\lambda^{-1}e'_4(rf) = -\frac{r}{2}\left(\check{\kappa} - \frac{e_3(r)}{2r}f^2\right)f + rE_1(f, \Gamma)$$

as desired.

Next, we have

$$\begin{aligned} & \lambda^{-1}e'_4\left(r\underline{f} - 2r^2e'_\theta(\log(\lambda)) + rf\underline{\Omega}\right) \\ = & r\left(-\frac{\kappa}{2}\underline{f} + 2e'_\theta(\log(\lambda)) + 2f\underline{\omega} + E_3(f, \underline{f}, \Gamma)\right) - 2r^2e'_\theta(E_2(f, \Gamma)) + r\underline{\Omega}\left(-\frac{\kappa}{2}f + E_1(f, \Gamma)\right) \\ & + \lambda^{-1}e'_4(r)\underline{f} - 2r^2[\lambda^{-1}e'_4, e'_\theta]\log(\lambda) - 4r\lambda^{-1}e'_4(r)e'_\theta(\log(\lambda)) + r\lambda^{-1}e'_4(\underline{\Omega})f + \lambda^{-1}e'_4(r)f\underline{\Omega} \\ = & -\frac{r}{2}\left(\kappa - \frac{2\lambda^{-1}e'_4(r)}{r}\right)\underline{f} + 2r\left(1 - 2\lambda^{-1}e'_4(r)\right)e'_\theta(\log(\lambda)) - 2r^2\lambda^{-1}[e'_4, e'_\theta]\log(\lambda) \\ & + r\left(\lambda^{-1}e'_4(\underline{\Omega}) + 2\underline{\omega}\right)f - \frac{r}{2}\left(\kappa - \frac{2\lambda^{-1}e'_4(r)}{r}\right)\underline{\Omega}f - 2r^2e'_\theta(\log(\lambda))\lambda^{-1}e'_4(\log(\lambda)) \\ & + rE_3(f, \underline{f}, \Gamma) - 2r^2e'_\theta(E_2(f, \Gamma)) + r\underline{\Omega}E_1(f, \Gamma). \end{aligned}$$

Since we have

$$\lambda^{-1}e'_4 = e_4 + fe_\theta + \frac{f^2}{4}e_3,$$

we infer

$$\begin{aligned} \lambda^{-1}e'_4(r) &= \frac{r}{2}\bar{\kappa} + \frac{f^2}{4}e_3(r), \\ \lambda^{-1}e'_4(\underline{\Omega}) &= e_4(\underline{\Omega}) + fe_\theta(\underline{\Omega}) + \frac{f^2}{4}e_3(\underline{\Omega}) \\ &= -2\underline{\omega} + fe_\theta(\underline{\Omega}) + \frac{f^2}{4}e_3(\underline{\Omega}). \end{aligned}$$

Together with the transport equation for $\log(\lambda)$ and the commutator identity for $[e'_4, e'_\theta]$, we infer

$$\begin{aligned} & \lambda^{-1}e'_4\left(r\underline{f} - 2r^2e'_\theta(\log(\lambda)) + rf\underline{\Omega}\right) \\ = & -\frac{r}{2}\left(\check{\kappa} - \frac{e_3(r)}{2r}f^2\right)\underline{f} + 2r\left(1 - r\bar{\kappa} - \frac{e_3(r)}{2}f^2\right)e'_\theta(\log(\lambda)) + r^2\lambda^{-1}(\kappa' + \vartheta')e'_\theta(\log(\lambda)) \\ & + r\left(e_\theta(\underline{\Omega}) + \frac{f}{4}e_3(\underline{\Omega})\right)f^2 - \frac{r}{2}\left(\check{\kappa} - \frac{e_3(r)}{2r}f^2\right)\underline{\Omega}f \\ & - 2r^2e'_\theta(\log(\lambda))E_2(f, \Gamma) + rE_3(f, \underline{f}, \Gamma) - 2r^2e'_\theta(E_2(f, \Gamma)) + r\underline{\Omega}E_1(f, \Gamma). \end{aligned}$$

Now, recall the following transformation formulas

$$\begin{aligned}\lambda^{-1}\kappa' &= \kappa + \phi_1'(f) + \text{Err}(\kappa, \kappa'), \\ \text{Err}(\kappa, \kappa') &= f(\zeta + \eta) - \frac{1}{4}f^2\underline{\kappa} - f^2\underline{\omega} + \text{l.o.t.}\end{aligned}$$

We infer

$$\begin{aligned}& \lambda^{-1}e_4'(rf - 2r^2e_\theta'(\log(\lambda)) + rf\underline{\Omega}) \\ &= -\frac{r}{2}\left(\check{\kappa} - \frac{e_3(r)}{2r}f^2\right)\underline{f} + r^2\left(\check{\kappa} - \left(\bar{\kappa} - \frac{2}{r}\right) - \frac{e_3(r)}{r}f^2\right)e_\theta'(\log(\lambda)) \\ & \quad + r^2\left(\phi_1'(f) + \text{Err}(\kappa, \kappa') + \lambda^{-1}\vartheta'\right)e_\theta'(\log(\lambda)) \\ & \quad + r\left(e_\theta(\underline{\Omega}) + \frac{f}{4}e_3(\underline{\Omega})\right)f^2 - \frac{r}{2}\left(\check{\kappa} - \frac{e_3(r)}{2r}f^2\right)\underline{\Omega}f \\ & \quad - 2r^2e_\theta'(\log(\lambda))E_2(f, \Gamma) + rE_3(f, \underline{f}, \Gamma) - 2r^2e_\theta'(E_2(f, \Gamma)) + r\underline{\Omega}E_1(f, \Gamma).\end{aligned}$$

This concludes the proof of Corollary 2.3.7.

A.9 Proof of Lemma 2.3.5

Recall that we have obtained in section A.7

$$\begin{aligned}2\xi' &= \lambda^2\left\{2\xi + \left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right)f + \left(\frac{1}{2}\kappa + 2\omega\right)f + \frac{1}{2}\vartheta f + \frac{1}{2}f^2\eta - \frac{1}{2}f^2\underline{\eta} + f^2\zeta - \frac{1}{4}f^3\underline{\chi}\right. \\ & \quad \left. - \frac{1}{2}f^3\underline{\omega} - \frac{1}{8}f^4\underline{\xi}\right\}, \\ 4\omega' &= \lambda\left\{4\omega - 2\left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right)\log(\lambda) + 2f\zeta - f^2\underline{\omega} - 2\underline{\eta}f - f^2\underline{\chi} - \frac{1}{2}f^3\underline{\xi}\right\} \\ & \quad + \lambda^{-1}f\xi'.\end{aligned}$$

In the case where $\lambda = 1$ and $\underline{f} = 0$, we immediately infer

$$\begin{aligned}2\xi' &= 2\xi + \left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right)f + \left(\frac{1}{2}\kappa + 2\omega\right)f + \frac{1}{2}\vartheta f + \frac{1}{2}f^2\eta - \frac{1}{2}f^2\underline{\eta} + f^2\zeta - \frac{1}{4}f^3\underline{\chi} \\ & \quad - \frac{1}{2}f^3\underline{\omega} - \frac{1}{8}f^4\underline{\xi}, \\ 4\omega' &= 4\omega + 2f\zeta - f^2\underline{\omega} - 2\underline{\eta}f - f^2\underline{\chi} - \frac{1}{2}f^3\underline{\xi},\end{aligned}$$

and hence

$$\begin{aligned}\xi' &= \xi + \frac{1}{2}e'_4(f) + \left(\frac{1}{4}\kappa + \omega\right) f + \frac{1}{4}\vartheta f + \frac{1}{4}f^2\eta - \frac{1}{4}f^2\underline{\eta} + \frac{1}{2}f^2\zeta - \frac{1}{8}f^3\underline{\chi} \\ &\quad - \frac{1}{4}f^3\underline{\omega} - \frac{1}{16}f^4\underline{\xi}, \\ \omega' &= \omega + \frac{1}{2}f\zeta - \frac{1}{4}f^2\underline{\omega} - \frac{1}{2}\underline{\eta}f - \frac{1}{8}f^2\underline{\kappa} - \frac{1}{8}f^2\underline{\vartheta} - \frac{1}{8}f^3\underline{\xi}.\end{aligned}$$

Finally, we compute the change of frame formula for ζ' and η' when $\lambda = 1, \underline{f} = 0$. We have in this case

$$\begin{aligned}e'_4 &= e_4 + fe_\theta + \frac{1}{4}f^2e_3, \\ e'_\theta &= e_\theta + \frac{1}{2}fe_3, \\ e'_3 &= e_3,\end{aligned}$$

and hence

$$\begin{aligned}2\zeta' &= g(D_{e'_\theta}e'_4, e'_3) \\ &= g\left(D_{e'_\theta}\left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right), e_3\right) \\ &= g(D_{e'_\theta}e_4, e_3) + fg(D_{e'_\theta}e_\theta, e_3) \\ &= g\left(D_{e_\theta + \frac{1}{2}fe_3}e_4, e_3\right) + fg\left(D_{e_\theta + \frac{1}{2}fe_3}e_\theta, e_3\right) \\ &= 2\zeta - 2\underline{\omega}f - \underline{\chi}f - \underline{\xi}f^2 \\ &= 2\zeta - \left(\frac{1}{2}\underline{\kappa} + 2\underline{\omega}\right)f - f\left(\frac{1}{2}\underline{\vartheta} + f\underline{\xi}\right)\end{aligned}$$

and

$$\begin{aligned}2\eta' &= g(D_{e'_3}e'_4, e'_\theta) \\ &= g\left(D_{e'_3}\left(e_4 + fe_\theta + \frac{1}{4}f^2e_3\right), e'_\theta\right) \\ &= g(D_{e'_3}e_4, e'_\theta) + e'_3(f)g(e_\theta, e'_\theta) + fg(D_{e'_3}e_\theta, e'_\theta) + \frac{1}{4}e'_3(f^2)g(e_3, e'_\theta) + \frac{1}{4}f^2g(D_{e'_3}e_3, e'_\theta) \\ &= e'_3(f) + g\left(D_{e_3}e_4, e_\theta + \frac{1}{2}fe_3\right) + fg\left(D_{e_3}e_\theta, \frac{1}{2}fe_3\right) + \frac{1}{4}f^2g(D_{e_3}e_3, e_\theta) \\ &= 2\eta + e'_3(f) - 2f\underline{\omega} - \frac{1}{2}f^2\underline{\xi}\end{aligned}$$

which yields the desired change of frame formula for ζ' and η' . This concludes the proof of Lemma 2.3.5.

A.10 Proof of Proposition 2.3.13

Recall that we have

$$\mathfrak{q} = r^4 \left(e_3(e_3(\alpha)) + (2\underline{\kappa} - 6\underline{\omega})e_3(\alpha) + \left(-4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2 \right) \alpha \right),$$

which we write in the form $\mathfrak{q} = r^4 J$ where,

$$\begin{aligned} J &= e_3(e_3(\alpha)) + (2\underline{\kappa} - 6\underline{\omega})e_3(\alpha) + V\alpha, \\ V &= -4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2. \end{aligned}$$

We make use of the general¹ Bianchi equations, see Proposition 2.2.2

$$\begin{aligned} e_3\alpha + \frac{1}{2}\underline{\kappa}\alpha &= -\underline{\phi}_2^*\beta + 4\underline{\omega}\alpha - \frac{3}{2}\vartheta\rho + \text{Err}[e_3(\alpha)], \\ e_3\beta + \underline{\kappa}\beta &= -\underline{\phi}_1^*\rho + 2\underline{\omega}\beta + 3\eta\rho + \text{Err}[e_3(\beta)], \\ e_3\rho + \frac{3}{2}\underline{\kappa}\rho &= \underline{\phi}_1\underline{\beta} + \text{Err}[e_3(\rho)] \end{aligned}$$

as well as the null structure equations (see Proposition 2.2.1)

$$\begin{aligned} e_3\vartheta + \frac{1}{2}\underline{\kappa}\vartheta - 2\underline{\omega}\vartheta &= -2\underline{\phi}_2^*\eta - \frac{1}{2}\underline{\kappa}\underline{\vartheta} + \text{Err}[e_3(\vartheta)] \\ e_3(\underline{\kappa}) + \frac{1}{2}\underline{\kappa}^2 + 2\underline{\omega}\underline{\kappa} &= 2\underline{\phi}_1\underline{\xi} + \text{Err}[e_3(\underline{\kappa})] \end{aligned}$$

where $\text{Err}[e_3(\alpha)], \text{Err}[e_3(\beta)], +\text{Err}[e_3(\rho)], \text{Err}[e_3(\vartheta)], \text{Err}[e_3(\underline{\kappa})]$ denote the corresponding quadratic terms in each equation. We also make use of the commutation formula (see Lemma 2.1.51)

$$\begin{aligned} [e_3, \underline{\phi}_2^*\beta] &= -\frac{1}{2}\underline{\kappa}\underline{\phi}_2^*\beta - \underline{Com}_2^*(\beta) \\ \underline{Com}_2^*(\beta) &= -\frac{1}{2}\underline{\vartheta}\underline{\phi}_1\beta - (\zeta - \eta)e_3\beta - \eta e_3\Phi\beta + \underline{\xi}(e_4\beta - e_4(\Phi)\beta) - \underline{\beta} \cdot \beta \end{aligned}$$

¹In an arbitrary \mathbf{Z} -invariant frame.

Thus,

$$\begin{aligned}
J &= e_3 \left(-\frac{1}{2}\underline{\kappa}\alpha - \underline{\phi}_2^*\beta + 4\underline{\omega}\alpha - \frac{3}{2}\vartheta\rho + \text{Err}[e_3(\alpha)] \right) + (2\underline{\kappa} - 6\underline{\omega})e_3(\alpha) + V\alpha \\
&= \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) e_3\alpha + \left(-\frac{1}{2}e_3\underline{\kappa} + 4e_3(\underline{\omega}) + V \right) \alpha - \underline{\phi}_2^*e_3\underline{\beta} - [e_3, \underline{\phi}_2^*]\beta - \frac{3}{2}\vartheta e_3\rho - \frac{3}{2}\rho e_3\vartheta \\
&\quad + e_3\text{Err}[e_3(\alpha)] \\
&= \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) \left(-\frac{1}{2}\underline{\kappa}\alpha - \underline{\phi}_2^*\beta + 4\underline{\omega}\alpha - \frac{3}{2}\vartheta\rho + \text{Err}[e_3(\alpha)] \right) + \left(-\frac{1}{2}e_3\underline{\kappa} + 4e_3(\underline{\omega}) + V \right) \alpha \\
&\quad - \underline{\phi}_2^*e_3\beta + \frac{1}{2}\underline{\kappa}\underline{\phi}_2^*\beta - \frac{3}{2}\vartheta e_3\rho - \frac{3}{2}\rho e_3\vartheta + e_3\text{Err}[e_3(\alpha)] + \underline{Com}_2^*(\beta) \\
&= -\underline{\phi}_2^*e_3\beta + (-\underline{\kappa} + 2\underline{\omega})\underline{\phi}_2^*\beta + \left(-\frac{1}{2}e_3\underline{\kappa} + 4e_3(\underline{\omega}) + V + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) \left(-\frac{1}{2}\underline{\kappa} + 4\underline{\omega} \right) \right) \alpha \\
&\quad - \frac{3}{2} \left(\vartheta e_3\rho + \rho e_3\vartheta + \vartheta\rho \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) \right) + e_3\text{Err}[e_3(\alpha)] + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) \text{Err}[e_3(\alpha)] + \underline{Com}_2^*(\beta)
\end{aligned}$$

Hence,

$$\begin{aligned}
J &= -\underline{\phi}_2^*e_3\beta + (-\underline{\kappa} + 2\underline{\omega})\underline{\phi}_2^*\beta - \frac{3}{2} \left(\vartheta e_3\rho + \rho e_3\vartheta + \vartheta\rho \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) \right) + W\alpha + E \\
W &:= -\frac{1}{2}e_3\underline{\kappa} + 4e_3(\underline{\omega}) + V + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) \left(-\frac{1}{2}\underline{\kappa} + 4\underline{\omega} \right) \tag{A.10.1} \\
E &:= e_3\text{Err}[e_3(\alpha)] + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) \text{Err}[e_3(\alpha)] + \underline{Com}_2^*(\beta)
\end{aligned}$$

Now, ignoring cubic and higher order terms,

$$\begin{aligned}
-\underline{\phi}_2^*e_3\beta + (-\underline{\kappa} + 2\underline{\omega})\underline{\phi}_2^*\beta &= -\underline{\phi}_2^* \left(-\underline{\kappa}\beta - \underline{\phi}_1^*\rho + 2\underline{\omega}\beta + 3\eta\rho + \text{Err}[e_3(\beta)] \right) + (-\underline{\kappa} + 2\underline{\omega})\underline{\phi}_2^*\beta \\
&= \underline{\phi}_2^*\underline{\phi}_1^*\rho - 3\rho\underline{\phi}_2^*\eta + \beta\underline{\phi}_1^*(\underline{\kappa} - 2\underline{\omega}) - 3\eta\underline{\phi}_1^*\rho - \underline{\phi}_2^*\text{Err}[e_3(\beta)]
\end{aligned}$$

Also,

$$\begin{aligned}
\vartheta e_3\rho + \rho e_3\vartheta + \vartheta\rho \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) &= \vartheta \left(-\frac{3}{2}\underline{\kappa}\rho + \underline{\phi}_1^*\beta \right) + \vartheta\rho \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) \\
&\quad + \rho \left(-\frac{1}{2}\underline{\kappa}\vartheta + 2\underline{\omega}\vartheta - 2\underline{\phi}_2^*\eta - \frac{1}{2}\underline{\kappa}\vartheta + \text{Err}[e_3(\vartheta)] \right) \\
&= -\frac{1}{2}\underline{\kappa}\rho\vartheta - \frac{1}{2}\underline{\kappa}\rho\vartheta - 2\rho\underline{\phi}_2^*\eta + \vartheta\underline{\phi}_1^*\beta + \rho\text{Err}[e_3(\rho)]
\end{aligned}$$

and,

$$\begin{aligned}
W &= -\frac{1}{2}e_3\underline{\kappa} + 4e_3(\underline{\omega}) + \left(-4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2\right) + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega}\right)\left(-\frac{1}{2}\underline{\kappa} + 4\underline{\omega}\right) \\
&= -\frac{1}{2}e_3\underline{\kappa} + \left(8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2\right) + \left(-\frac{3}{4}\underline{\kappa}^2 - 8\underline{\omega}^2 + 7\underline{\omega}\underline{\kappa}\right) \\
&= -\frac{1}{2}e_3\underline{\kappa} - \frac{1}{4}\underline{\kappa}^2 - \underline{\omega}\underline{\kappa} \\
&= -\frac{1}{2}\left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega}\underline{\kappa} + 2\phi_1\underline{\xi} + \text{Err}[e_3(\underline{\kappa})]\right) - \frac{1}{4}\underline{\kappa}^2 - \underline{\omega}\underline{\kappa} \\
&= -\phi_1\underline{\xi} - \frac{1}{2}\text{Err}[e_3(\underline{\kappa})]
\end{aligned}$$

Thus, back to (A.10.1),

$$\begin{aligned}
J &= \phi_2^* \phi_1^* \rho - 3\rho \phi_2^* \eta + \beta \phi_1^* (\underline{\kappa} - 2\underline{\omega}) - 3\eta \phi_1^* \rho - \phi_2^* \text{Err}[e_3(\beta)] \\
&\quad - \frac{3}{2}\left(-\frac{1}{2}\underline{\kappa}\rho\vartheta - \frac{1}{2}\kappa\rho\underline{\vartheta} - 2\rho \phi_2^* \eta + \vartheta \phi_1 \underline{\beta} + \rho \text{Err}[e_3(\rho)]\right) - \phi_1 \underline{\xi} \alpha + E \\
&= \phi_2^* \phi_1^* \rho + \frac{3}{4}\rho(\underline{\kappa}\vartheta + \kappa\underline{\vartheta}) + \beta \phi_1^* (\underline{\kappa} - 2\underline{\omega}) - 3\eta \phi_1^* \rho - \frac{3}{2}\vartheta \phi_1 \underline{\beta} - \phi_1 \underline{\xi} \alpha - \frac{3}{2}\rho \text{Err}[e_3(\rho)] + E
\end{aligned}$$

In other words,

$$\begin{aligned}
J &= \phi_2^* \phi_1^* \rho + \frac{3}{4}\rho(\underline{\kappa}\vartheta + \kappa\underline{\vartheta}) + \text{Err} \\
\text{Err} &:= \beta \phi_1^* (\underline{\kappa} - 2\underline{\omega}) - 3\eta \phi_1^* \rho - \frac{3}{2}\vartheta \phi_1 \underline{\beta} - \phi_1 \underline{\xi} \alpha \\
&\quad - \frac{3}{2}\rho \text{Err}[e_3(\rho)] + e_3 \text{Err}[e_3(\alpha)] + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega}\right) \text{Err}[e_3(\alpha)] + \underline{\text{Com}}_2^*(\beta) + \text{l.o.t.}
\end{aligned}$$

It remains to analyze the lower order terms according to our convention in Definition 2.3.8. Note that we can write the first line in the expression of Err

$$\begin{aligned}
\text{Err}_1 &= r^{-1}\Gamma_b \cdot \beta + r^{-2}\Gamma_g \cdot \vartheta \Gamma_g + r^{-2}\Gamma_g \vartheta \Gamma_b + r^{-1}\vartheta \Gamma_b \cdot \alpha \\
&= r^{-1}\vartheta^{\leq 1}\Gamma_b \cdot \beta + r^{-2}\Gamma_g \vartheta \Gamma_b + \text{l.o.t.}
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\text{Err}[e_3(\rho)] &= -\frac{1}{2}\vartheta \underline{\alpha} - \zeta \underline{\beta} + 2(\eta \underline{\beta} + \underline{\xi} \beta) = \Gamma_g \cdot \Gamma_b + \Gamma_b \cdot \beta \\
\text{Err}[e_3(\alpha)] &= (\zeta + 4\eta)\beta = \Gamma_g \cdot \beta \\
e_3 \text{Err}[e_3(\alpha)] &= e_3(\zeta + 4\eta)\beta + (\zeta + 4\eta)e_3(\beta) \\
&= e_3(\zeta + 4\eta) \cdot \beta + (\zeta + 4\eta)(-\underline{\kappa}\beta - \phi_1^* \rho + 2\underline{\omega}\beta + 3\eta\rho) \\
&= e_3(\zeta + 4\eta) \cdot \beta + r^{-1}\Gamma_g \beta + r^{-2}\Gamma_g \vartheta \Gamma_g + r^{-3}\Gamma_g \cdot \Gamma_g
\end{aligned}$$

$$\begin{aligned}
\underline{Com}_2^*(\beta) &= -\frac{1}{2}\vartheta\phi_1\beta - (\zeta - \eta)e_3\beta - \eta e_3\Phi\beta + \underline{\xi}(e_4\beta - e_4(\Phi)\beta) - \underline{\beta} \cdot \beta \\
&= -\frac{1}{2}\vartheta\phi_1\beta - (\zeta - \eta)(-\underline{\kappa}\beta - \phi_1^*\rho + 2\underline{\omega}\beta + 3\eta\rho) - \eta e_3(\Phi)\beta \\
&\quad + \underline{\xi}(-2\kappa\beta + \phi_2^*\alpha - 2\omega\beta) - \underline{\xi}e_4\Phi\beta - \underline{\beta} \cdot \beta + \text{l.o.t.} \\
&= r^{-1}\Gamma_b \cdot \vartheta^{\leq 1}\beta + r^{-2}\Gamma_g \cdot \vartheta\Gamma_b + \text{l.o.t.}
\end{aligned}$$

Therefore, schematically,

$$\text{Err} = e_3(\zeta + 4\eta) \cdot \beta + r^{-1}\Gamma_b \vartheta^{\leq 1}\beta + r^{-2}\Gamma_g \vartheta\Gamma_b + \text{l.o.t.}$$

and therefore,

$$\text{Err}[\mathbf{q}] = r^4 \text{Err} = r^4 (e_3(\zeta + 4\eta) \cdot \beta + r^{-1}\Gamma_b \cdot \vartheta^{\leq 1}\beta + r^{-2}\Gamma_g \cdot \vartheta\Gamma_b) + \text{l.o.t.}$$

Since $e_3\zeta \in r^{-1}\mathfrak{d}\Gamma_b$ and $\beta \in r^{-1}\Gamma_g$ we rewrite in the form,

$$\text{Err}[\mathbf{q}] = r^4 e_3\eta \cdot \beta + r^2 \mathfrak{d}^{\leq 1}(\Gamma_b \cdot \Gamma_g).$$

This concludes the proof of Proposition 2.3.13.

A.11 Proof of Proposition 2.3.14

We start with the formula (2.3.11)

$$r\mathbf{q} = r^5 \left(\phi_2^* \phi_1^* \rho + \frac{3}{4} \underline{\kappa} \rho \vartheta + \frac{3}{4} \kappa \rho \underline{\vartheta} \right) + r \text{Err}[\mathbf{q}].$$

with $\text{Err}[\mathbf{q}]$ given by (2.3.12). Taking the e_3 derivative we deduce,

$$\begin{aligned}
e_3(r\mathbf{q}) &= r^5 L + 5e_3(r) \mathbf{q} + e_3(r \text{Err}[\mathbf{q}]) - 5e_3(r) \text{Err}[\mathbf{q}], \\
L &:= e_3 \left\{ \phi_2^* \phi_1^* \rho + \frac{3}{4} e_3(\underline{\kappa} \rho \vartheta) + \frac{3}{4} e_3(\kappa \rho \underline{\vartheta}) \right\}
\end{aligned} \tag{A.11.1}$$

We calculate L as follows,

$$\begin{aligned}
L &= e_3 \phi_2^* \phi_1^* \rho + \frac{3}{4} e_3(\underline{\kappa} \rho \vartheta) + \frac{3}{4} e_3(\kappa \rho \underline{\vartheta}) \\
&= \phi_2^* \phi_1^* e_3(\rho) + [e_3, \phi_2^* \phi_1^*] \rho + \frac{3}{4} e_3(\underline{\kappa} \rho \vartheta) + \frac{3}{4} e_3(\kappa \rho \underline{\vartheta}).
\end{aligned}$$

Ignoring cubic and higher order terms

$$\begin{aligned}
e_3(\underline{\kappa}\rho\vartheta) &= \underline{\kappa}\rho e_3(\vartheta) + e_3(\underline{\kappa})\rho\vartheta + \underline{\kappa}e_3(\rho)\vartheta \\
&= \underline{\kappa}\rho \left(-\frac{1}{2}\underline{\kappa}\vartheta + 2\underline{\omega}\vartheta - 2\mathfrak{d}_2^*\eta - \frac{1}{2}\underline{\kappa}\vartheta + \text{Err}[e_3\vartheta] \right) \\
&\quad + \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega}\underline{\kappa} + 2\mathfrak{d}_1\underline{\xi} + \text{Err}[e_3\underline{\kappa}] \right) \rho\vartheta \\
&\quad + \underline{\kappa} \left(-\frac{3}{2}\underline{\kappa}\rho + \mathfrak{d}_1\underline{\beta} - \frac{1}{2}\vartheta\underline{\alpha} + \text{Err}[e_3(\rho)] \right) \vartheta \\
&= \underline{\kappa}\rho \left(-\frac{5}{2}\underline{\kappa}\vartheta - 2\mathfrak{d}_2^*\eta - \frac{1}{2}\underline{\kappa}\vartheta \right) + \underline{\kappa}\rho\text{Err}[e_3\vartheta] + 2\mathfrak{d}_1\underline{\xi}\rho\vartheta + \underline{\kappa}(\mathfrak{d}_1\underline{\beta})\vartheta
\end{aligned}$$

We deduce,

$$\begin{aligned}
e_3(\underline{\kappa}\rho\vartheta) &= \underline{\kappa}\rho \left(-\frac{5}{2}\underline{\kappa}\vartheta - 2\mathfrak{d}_2^*\eta - \frac{1}{2}\underline{\kappa}\vartheta \right) + E_1 \\
E_1 &= 2\mathfrak{d}_1\underline{\xi}\rho\vartheta + \underline{\kappa}(\mathfrak{d}_1\underline{\beta})\vartheta + \underline{\kappa}\rho\text{Err}[e_3(\vartheta)]
\end{aligned} \tag{A.11.2}$$

where $\text{Err}[e_3(\vartheta)]$, $\text{Err}[e_3(\underline{\kappa})]$, $\text{Err}[e_3(\rho)]$ denote the quadratic error terms in the corresponding equations. Also,

$$\begin{aligned}
e_3(\kappa\rho\underline{\vartheta}) &= \kappa\rho e_3(\underline{\vartheta}) + e_3(\kappa)\rho\underline{\vartheta} + \kappa e_3(\rho)\underline{\vartheta} \\
&= \kappa\rho \left(-\underline{\kappa}\underline{\vartheta} - 2\underline{\omega}\underline{\vartheta} - 2\underline{\alpha} - 2\mathfrak{d}_2^*\underline{\xi} + \text{Err}[e_3(\underline{\vartheta})] \right) \\
&\quad + \left(-\frac{1}{2}\underline{\kappa}\kappa + 2\underline{\omega}\kappa + 2\mathfrak{d}_1\underline{\eta} + 2\rho + \text{Err}[e_3(\kappa)] \right) \rho\underline{\vartheta} \\
&\quad + \left(-\frac{3}{2}\underline{\kappa}\rho + \mathfrak{d}_1\underline{\beta} + \text{Err}[e_3(\rho)] \right) \kappa\underline{\vartheta}
\end{aligned}$$

Hence, ignoring the higher order terms,

$$\begin{aligned}
e_3(\kappa\rho\underline{\vartheta}) &= \kappa\rho \left(-3\underline{\kappa}\underline{\vartheta} - 2\underline{\alpha} - 2\mathfrak{d}_2^*\underline{\xi} \right) + 2\rho^2\underline{\vartheta} + E_2 \\
E_2 &:= 2\rho\mathfrak{d}_1\underline{\eta}\underline{\vartheta} + \kappa\mathfrak{d}_1\underline{\beta}\underline{\vartheta} + \kappa\rho\text{Err}[e_3(\underline{\vartheta})]
\end{aligned} \tag{A.11.3}$$

Also, we have

$$\begin{aligned}
\mathfrak{d}_2^*\mathfrak{d}_1^*e_3(\rho) &= \mathfrak{d}_2^*\mathfrak{d}_1^* \left(-\frac{3}{2}\underline{\kappa}\rho + \mathfrak{d}_1\underline{\beta} + \text{Err}[e_3(\rho)] \right) \\
&= \mathfrak{d}_2^*\mathfrak{d}_1^*\mathfrak{d}_1\underline{\beta} - \frac{3}{2}\rho\mathfrak{d}_2^*\mathfrak{d}_1^*\underline{\kappa} - \frac{3}{2}\underline{\kappa}\mathfrak{d}_2^*\mathfrak{d}_1^*\rho + E_3 \\
E_3 &= \mathfrak{d}_2^*\mathfrak{d}_1^*\text{Err}[e_3(\rho)] - 3\mathfrak{d}_1^*\underline{\kappa} \cdot \mathfrak{d}_1^*\rho
\end{aligned}$$

i.e.,

$$\begin{aligned}
e_3(\underline{\kappa}\rho\vartheta) &= \underline{\kappa}\rho \left(-\frac{5}{2}\underline{\kappa}\vartheta - 2\phi_2^*\eta - \frac{1}{2}\underline{\kappa}\vartheta \right) + E_1, \\
e_3(\underline{\kappa}\rho\underline{\vartheta}) &= \underline{\kappa}\rho \left(-3\underline{\kappa}\underline{\vartheta} - 2\underline{\alpha} - 2\phi_2^*\underline{\xi} \right) + 2\rho^2\underline{\vartheta} + E_2 \\
\phi_2^*\phi_1^*e_3(\rho) &= \phi_2^*\phi_1^*\phi_1\underline{\beta} - \frac{3}{2}\rho\phi_2^*\phi_1^*\underline{\kappa} - \frac{3}{2}\underline{\kappa}\phi_2^*\phi_1^*\rho + E_3
\end{aligned} \tag{A.11.4}$$

Now, in view of Lemma 2.1.51 we have (for $f = \phi_1^*\rho \in \mathfrak{s}_{2-1}$),

$$[e_3, \phi_2^*]\phi_1^*\rho = -\frac{1}{2}\underline{\kappa}\phi_2^*\phi_1^*\rho - \underline{Com}_2^*(\phi_1^*\rho)$$

and

$$\begin{aligned}
[e_3, \phi_1^*]\rho &= -[e_3, e_\theta]\rho = -\frac{1}{2}\underline{\kappa}\phi_1^*\rho - \frac{1}{2}\vartheta e_\theta\rho + (\zeta - \eta)e_3\rho - \underline{\xi}e_4\rho \\
&= -\frac{1}{2}\underline{\kappa}\phi_1^*\rho - \frac{1}{2}\vartheta e_\theta\rho + (\zeta - \eta)\left(-\frac{3}{2}\underline{\kappa}\rho + \phi_1\underline{\beta} + \text{Err}[e_3(\rho)]\right) \\
&\quad - \underline{\xi}\left(-\frac{3}{2}\underline{\kappa}\rho + \phi_1\underline{\beta} + \text{Err}[e_4(\rho)]\right) \\
&= -\frac{1}{2}\underline{\kappa}\phi_1^*\rho - \frac{3}{2}\rho\left[(\zeta - \eta)\underline{\kappa} - \underline{\xi}\underline{\kappa}\right] + E_{41} \\
E_{41} &= (\zeta - \eta)\phi_1\underline{\beta} - \underline{\xi}\phi_1\underline{\beta} + (\zeta - \eta)e_3[\rho] - \frac{1}{2}\vartheta e_\theta\rho - \underline{\xi}\text{Err}[e_4(\rho)]
\end{aligned}$$

We deduce

$$\begin{aligned}
\phi_2^*[e_3, \phi_1^*]\rho &= \phi_2^*\left(-\frac{1}{2}\underline{\kappa}\phi_1^*\rho - \frac{3}{2}\rho\left[\underline{\kappa}(\zeta - \eta) - \underline{\kappa}\underline{\xi}\right] + E_{41}\right) \\
&= -\frac{1}{2}\underline{\kappa}\phi_2^*\phi_1^*\rho - \frac{3}{2}\rho\left(\underline{\kappa}\phi_2^*(\zeta - \eta) - \underline{\kappa}\phi_2^*\underline{\xi}\right) + E_4 \\
E_4 &= \phi_2^*E_{41} - \frac{1}{2}\phi_1^*\underline{\kappa} \cdot \phi_1^*\rho - \frac{3}{2}(\zeta - \eta)\phi_1^*(\underline{\kappa}\rho) + \frac{3}{2}\underline{\xi}\phi_1^*(\underline{\kappa}\rho)
\end{aligned}$$

Hence, since $[e_3, \phi_2^*\phi_1^*]\rho = [e_3, \phi_2^*]\phi_1^*\rho + \phi_2^*[e_3, \phi_1^*]\rho$,

$$[e_3, \phi_2^*\phi_1^*]\rho = -\underline{\kappa}\phi_2^*\phi_1^*\rho - \frac{3}{2}\left(\phi_2^*(\zeta - \eta)\underline{\kappa}\rho - \phi_2^*\underline{\xi}\underline{\kappa}\rho\right) - \underline{Com}_2^*(\phi_1^*\rho) + E_4 \tag{A.11.5}$$

We deduce, recalling (A.11.4),

$$\begin{aligned}
L &= \phi_2^* \phi_1^* e_3(\rho) + [e_3, \phi_2^* \phi_1^*] \rho + \frac{3}{4} e_3(\underline{\kappa} \rho \vartheta) + \frac{3}{4} e_3(\kappa \rho \underline{\vartheta}) \\
&= \phi_2^* \phi_1^* \phi_1 \underline{\beta} - \frac{3}{2} \rho \phi_2^* \phi_1^* \underline{\kappa} - \frac{3}{2} \underline{\kappa} \phi_2^* \phi_1^* \rho + E_3 \\
&\quad - \underline{\kappa} \phi_2^* \phi_1^* \rho - \frac{3}{2} \left(\phi_2^* (\zeta - \eta) \underline{\kappa} \rho - \phi_2^* \underline{\xi} \kappa \rho \right) - \underline{Com}_2^*(\phi_1^* \rho) + E_4 \\
&\quad + \frac{3}{4} \left(\underline{\kappa} \rho \left(-\frac{5}{2} \underline{\kappa} \vartheta - 2 \phi_2^* \eta - \frac{1}{2} \kappa \underline{\vartheta} \right) \right) + E_1 \\
&\quad + \frac{3}{4} \left(\kappa \rho \left(-3 \underline{\kappa} \underline{\vartheta} - 2 \underline{\alpha} - 2 \phi_2^* \underline{\xi} \right) + 2 \rho^2 \underline{\vartheta} + \text{Err}_1 \right) \\
&= \phi_2^* \phi_1^* \phi_1 \underline{\beta} - \frac{5}{2} \underline{\kappa} \phi_2^* \phi_1^* \rho - \frac{3}{2} \rho \phi_2^* \phi_1^* \underline{\kappa} - \frac{3}{2} \rho \underline{\kappa} \phi_2^* \zeta \\
&\quad + \frac{3}{4} \underline{\kappa} \rho \left(-\frac{5}{2} \underline{\kappa} \vartheta - \frac{1}{2} \kappa \underline{\vartheta} \right) + \frac{3}{4} \kappa \rho \left(-3 \underline{\kappa} \underline{\vartheta} - 2 \underline{\alpha} \right) + \frac{3}{2} \rho^2 \underline{\vartheta} \\
&\quad + E_1 + E_2 + E_3 + E_4 - \underline{Com}_2^*(\phi_1^* \rho)
\end{aligned}$$

i.e.,

$$\begin{aligned}
L &= \phi_2^* \phi_1^* \phi_1 \underline{\beta} - \frac{5}{2} \underline{\kappa} \phi_2^* \phi_1^* \rho - \frac{3}{2} \rho \phi_2^* \phi_1^* \underline{\kappa} - \frac{3}{2} \rho \underline{\kappa} \phi_2^* \zeta \\
&\quad - \frac{3}{4} \underline{\kappa} \rho \left(\frac{5}{2} \underline{\kappa} \vartheta + \frac{1}{2} \kappa \underline{\vartheta} \right) - \frac{3}{4} \kappa \rho (3 \underline{\kappa} \underline{\vartheta} + 2 \underline{\alpha}) + \frac{3}{2} \rho^2 \underline{\vartheta} + E \\
E &= E_1 + E_2 + E_3 + E_4 - \underline{Com}_2^*(\phi_1^* \rho)
\end{aligned} \tag{A.11.6}$$

On the other hand, in view of (2.3.11), writing $e_3 r = \frac{r}{2} (\underline{\kappa} + \underline{A})$,

$$\begin{aligned}
5e_3(r) \mathfrak{q} &= r \frac{5}{2} \underline{\kappa} \mathfrak{q} + 5r \underline{A} \mathfrak{q} \\
&= \frac{5}{2} r^5 \underline{\kappa} \left\{ \phi_2^* \phi_1^* \rho + \frac{3}{4} \underline{\kappa} \rho \vartheta + \frac{3}{4} \kappa \rho \underline{\vartheta} + \text{Err} \right\} + 5r \underline{A} \mathfrak{q}
\end{aligned}$$

Hence, in view of (A.11.1) and (A.11.6),

$$\begin{aligned}
e_3(r\mathbf{q}) &= r^5 L + 5e_3(r)\mathbf{q} + e_3(r\text{Err}) - 5e_3(r)\text{Err} \\
&= r^5 \left\{ \left(\underline{\phi}_2^* \underline{\phi}_1^* \underline{\phi}_1 \underline{\beta} - \frac{5}{2} \underline{\kappa} \underline{\phi}_2^* \underline{\phi}_1^* \rho - \frac{3}{2} \rho \underline{\phi}_2^* \underline{\phi}_1^* \underline{\kappa} - \frac{3}{2} \rho \underline{\kappa} \underline{\phi}_2^* \zeta \right) \right. \\
&\quad \left. - \frac{3}{4} \underline{\kappa} \rho \left(\frac{5}{2} \underline{\kappa} \underline{\vartheta} + \frac{1}{2} \underline{\kappa} \underline{\vartheta} \right) - \frac{3}{4} \underline{\kappa} \rho (3\underline{\kappa} \underline{\vartheta} + 2\underline{\alpha}) + \frac{3}{2} \rho^2 \underline{\vartheta} \right\} \\
&\quad + \frac{5}{2} r^5 \underline{\kappa} \left\{ \underline{\phi}_2^* \underline{\phi}_1^* \rho + \frac{3}{4} \underline{\kappa} \rho \underline{\vartheta} + \frac{3}{4} \underline{\kappa} \rho \underline{\vartheta} \right\} + r^5 E + e_3(r^5 \text{Err}) - 5 \frac{e_3 r}{r} r^5 \text{Err} + 5r \underline{A} \mathbf{q} \\
&= r^5 \left(\underline{\phi}_2^* \underline{\phi}_1^* \underline{\phi}_1 \underline{\beta} - \frac{3}{2} \rho \underline{\phi}_2^* \underline{\phi}_1^* \underline{\kappa} - \frac{3}{2} \underline{\kappa} \rho \underline{\phi}_2^* \zeta - \frac{3}{2} \underline{\kappa} \rho \underline{\alpha} + \frac{3}{4} (2\rho^2 - \underline{\kappa} \underline{\kappa} \rho) \underline{\vartheta} \right) \\
&\quad + \text{Err}[e_3(r\mathbf{q})].
\end{aligned}$$

where,

$$\text{Err}[e_3(r\mathbf{q})] = e_3(r\text{Err}[\mathbf{q}]) - 5e_3 r \text{Err}[\mathbf{q}] + 5r \underline{A} \mathbf{q} + r^5 E \quad (\text{A.11.7})$$

$$= r e_3(\text{Err}[\mathbf{q}]) + \text{Err}[\mathbf{q}] + 5r \underline{A} \mathbf{q} + r^5 E \quad (\text{A.11.8})$$

and

$$E = E_1 + E_2 + E_3 + E_4 - \underline{Com}_2^*(\underline{\phi}_1^* \rho)$$

with,

$$E_1 = 2 \underline{\phi}_1 \underline{\xi} \rho \underline{\vartheta} + \underline{\kappa} (\underline{\phi}_1 \underline{\beta}) \underline{\vartheta} + \underline{\kappa} \rho \text{Err}[e_3(\underline{\kappa})]$$

$$E_2 = 2\rho \underline{\phi}_1 \underline{\eta} \underline{\vartheta} + \underline{\kappa} \underline{\phi}_1 \underline{\beta} \underline{\vartheta} + \underline{\kappa} \rho \text{Err}[e_3(\underline{\vartheta})]$$

$$E_3 = \underline{\phi}_2^* \underline{\phi}_1^* \text{Err}[e_3(\rho)] - 3 \underline{\phi}_1^* \underline{\kappa} \cdot \underline{\phi}_1^* \rho$$

$$E_4 = \underline{\phi}_2^* E_{41} - \frac{1}{2} \underline{\phi}_1^* \underline{\kappa} \cdot \underline{\phi}_1^* \rho - \frac{3}{2} (\zeta - \eta) \underline{\phi}_1^* (\underline{\kappa} \rho) + \frac{3}{2} \underline{\xi} \underline{\phi}_1^* (\underline{\kappa} \rho)$$

$$E_{41} = (\zeta - \eta) \underline{\phi}_1 \underline{\beta} - \underline{\xi} \underline{\phi}_1 \underline{\beta} - \frac{1}{2} \underline{\vartheta} e_{\theta} \rho$$

$$\underline{Com}_2^*(\underline{\phi}_1^* \rho) = -\frac{1}{2} \underline{\vartheta} \underline{\phi}_1 \underline{\phi}_1^* \rho - (\zeta - \eta) e_3 \underline{\phi}_1^* \rho - \eta e_3 \Phi \underline{\phi}_1^* \rho + \underline{\xi} (e_4 \underline{\phi}_1^* \rho - e_4(\Phi) \underline{\phi}_1^* \rho) - \underline{\beta} \underline{\phi}_1^* \rho.$$

Note also that,

$$(\zeta - \eta) e_3 \underline{\phi}_1^* \rho = (\zeta - \eta) \cdot \underline{\phi}_1^* e_3 \rho + (\zeta - \eta) \left(-\frac{1}{2} \underline{\kappa} \underline{\phi}_1^* \rho - \frac{1}{2} \underline{\vartheta} e_{\theta} \rho + (\zeta - \eta) e_3 \rho - \underline{\xi} e_4 \rho \right)$$

Using our schematic notation

$$\text{Err}[e_3(\underline{\kappa})] = \Gamma_b \cdot \Gamma_b + \text{l.o.t.}$$

$$\text{Err}[e_3(\underline{\vartheta})] = \Gamma_b \cdot \Gamma_b + \text{l.o.t.}$$

$$\text{Err}[e_3(\rho)] = \Gamma_g \cdot \underline{\alpha} + \text{l.o.t.} = \Gamma_g \cdot \Gamma_b + \text{l.o.t.}$$

and

$$\begin{aligned}
E_1 &= r^{-4}\Gamma_b \cdot \not{\partial}^{\leq 1}\Gamma_b + \text{l.o.t.} \\
E_2 &= r^{-4}\Gamma_b \cdot \not{\partial}^{\leq 1}\Gamma_b + r^{-2}\Gamma_b \cdot \beta + \text{l.o.t.} \\
E_3 &= r^{-2}\not{\partial}^2(\Gamma_g \cdot \Gamma_b) + r^{-3}(\not{\partial}\Gamma_g) \cdot (\not{\partial}\Gamma_g) \\
E_{41} &= r^{-2}\Gamma_g \cdot (\not{\partial}\Gamma_b) + r^{-1}\Gamma_b \cdot \not{\partial}\beta + r^{-2}\Gamma_b \not{\partial} \cdot \Gamma_g \\
&= r^{-2}\not{\partial}(\Gamma_g \cdot \Gamma_b) + \text{l.o.t.} \\
E_4 &= r^{-3}\not{\partial}^2(\Gamma_g \cdot \Gamma_b) + \text{l.o.t.} \\
\text{Com}_2^*(\not{\partial}_1^*\rho) &= r^{-3}\Gamma_b \cdot \not{\partial}^{\leq 2}\Gamma_g + r^{-2}\not{\partial}\Gamma_b \cdot \Gamma_g + \text{l.o.t.}
\end{aligned}$$

and, since $r^{-1}\Gamma_b$ can be replaced by Γ_g and $\not{\partial}\beta$ can be replaced by $r^{-1}\Gamma_g$,

$$E = r^{-3}\not{\partial}^2(\Gamma_g \cdot \Gamma_b) + \text{l.o.t.}$$

Taking into account the expression of $\text{Err}[\mathbf{q}]$ in Proposition 2.3.13 we write

$$\begin{aligned}
re_3(\text{Err}[\mathbf{q}]) + \text{Err}[\mathbf{q}] &= re_3\left[r^4e_3\eta \cdot \beta + r^2\mathfrak{d}^{\leq 1}(\Gamma_b \cdot \Gamma_g)\right] + r^4e_3\eta \cdot \beta + r^2\mathfrak{d}^{\leq 1}(\Gamma_b \cdot \Gamma_g) \\
&= r^5\mathfrak{d}^{\leq 1}(e_3\eta \cdot \beta) + r^3\mathfrak{d}^{\leq 2}(\Gamma_b \cdot \Gamma_g)
\end{aligned}$$

and therefore, back to (A.11.7),

$$\begin{aligned}
\text{Err}[e_3(r\mathbf{q})] &= e_3(r\text{Err}[\mathbf{q}]) + \text{Err}[\mathbf{q}] + r\underline{A}\mathbf{q} + r^5E \\
&= r^5\mathfrak{d}^{\leq 1}(e_3\eta \cdot \beta) + r^3\mathfrak{d}^{\leq 2}(\Gamma_b \cdot \Gamma_g) + r\Gamma_b\mathbf{q} + r^2\not{\partial}^2(\Gamma_g \cdot \Gamma_b) + \text{l.o.t.} \\
&= r\Gamma_b\mathbf{q} + r^5\mathfrak{d}^{\leq 1}(e_3\eta \cdot \beta) + r^3\mathfrak{d}^{\leq 2}(\Gamma_b \cdot \Gamma_g).
\end{aligned}$$

This concludes the proof of Proposition 2.3.14.

A.12 Proof of the Teukolsky-Starobinski identity

According to Proposition 2.3.14 we have

$$e_3(r\mathbf{q}) = r^5 \left\{ \not{\partial}_2^* \not{\partial}_1^* \not{\partial}_1 \underline{\beta} - \frac{3}{2}\rho \not{\partial}_2^* \not{\partial}_1^* \underline{\kappa} - \frac{3}{2}\underline{\kappa}\rho \not{\partial}_2^* \zeta - \frac{3}{2}\kappa\rho \underline{\alpha} + \frac{3}{4}(2\rho^2 - \kappa\underline{\kappa}\rho)\underline{\varrho} \right\} + \text{Err}[e_3\mathbf{q}].$$

We infer that

$$\begin{aligned}
e_3(r^2e_3(r\mathbf{q})) &= r^7 \left\{ e_3 \not{\partial}_2^* \not{\partial}_1^* \not{\partial}_1 \underline{\beta} - \frac{3}{2}e_3(\rho \not{\partial}_2^* \not{\partial}_1^* \underline{\kappa}) - \frac{3}{2}e_3(\underline{\kappa}\rho \not{\partial}_2^* \zeta) - \frac{3}{2}e_3(\kappa\rho \underline{\alpha}) \right. \\
&\quad \left. + \frac{3}{4}e_3\left((2\rho^2 - \kappa\underline{\kappa}\rho)\underline{\varrho}\right) \right\} + 7re_3(r)e_3(r\mathbf{q}) \\
&\quad + r^2e_3(\text{Err}[e_3\mathbf{q}]) + r\text{Err}[e_3\mathbf{q}] + \text{l.o.t.}
\end{aligned} \tag{A.12.1}$$

We first compute

$$\begin{aligned}
e_3 \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} &= \not\partial_2^* \not\partial_1^* \not\partial_1 (e_3 \underline{\beta}) + [e_3, \not\partial_2^*] \not\partial_1^* \not\partial_1 \underline{\beta} + \not\partial_2^* [e_3, \not\partial_1^*] \not\partial_1 \underline{\beta} + \not\partial_2^* \not\partial_1^* [e_3, \not\partial_1] \underline{\beta} \\
&= \not\partial_2^* \not\partial_1^* \not\partial_1 (e_3 \underline{\beta}) + \left(-\frac{1}{2} \underline{\kappa} \not\partial_2^* + \frac{1}{2} \underline{\vartheta} \not\partial_1 + (\zeta - \eta) e_3 + \eta e_3 \Phi - \underline{\xi} (e_4 - e_4(\Phi)) + \underline{\beta} \right) \not\partial_1^* \not\partial_1 \underline{\beta} \\
&\quad + \not\partial_2^* \left(\left(-\frac{1}{2} \underline{\kappa} \not\partial_1^* - \frac{1}{2} \underline{\vartheta} \not\partial_1^* + (\zeta - \eta) e_3 - \underline{\xi} e_4 \right) \not\partial_1 \underline{\beta} \right) \\
&\quad + \not\partial_2^* \not\partial_1^* \left(\left(-\frac{1}{2} \underline{\kappa} \not\partial_1 + \frac{1}{2} \underline{\vartheta} \not\partial_2^* - (\zeta - \eta) e_3 + \eta e_3 \Phi + \underline{\xi} (e_4 + e_4(\Phi)) + \underline{\beta} \right) \underline{\beta} \right)
\end{aligned}$$

In view of our general commutation formulas in Lemma 2.1.51 and our notation convention for error terms² we have³,

$$\begin{aligned}
[e_3, \not\partial_2^*] \not\partial_1^* \not\partial_1 \underline{\beta} &= \left(-\frac{1}{2} \underline{\kappa} \not\partial_2^* + \frac{1}{2} \underline{\vartheta} \not\partial_1 + (\zeta - \eta) e_3 + \eta e_3 \Phi - \underline{\xi} (e_4 - e_4(\Phi)) + \underline{\beta} \right) \not\partial_1^* \not\partial_1 \underline{\beta} \\
&= -\frac{1}{2} \underline{\kappa} \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} + r^{-4} \Gamma_b \cdot \not\partial^{\leq 3} \Gamma_b + \text{l.o.t.}
\end{aligned}$$

$$\begin{aligned}
\not\partial_2^* [e_3, \not\partial_1^*] \not\partial_1 \underline{\beta} &= \not\partial_2^* \left(\left(-\frac{1}{2} \underline{\kappa} \not\partial_1^* - \frac{1}{2} \underline{\vartheta} \not\partial_1^* + (\zeta - \eta) e_3 - \underline{\xi} e_4 \right) \not\partial_1 \underline{\beta} \right) \\
&= -\frac{1}{2} \underline{\kappa} \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} + r^{-4} (\Gamma_b \cdot \not\partial^{\leq 3} \Gamma_b + \Gamma_b^{\leq 1} \cdot \not\partial^{\leq 2} \Gamma_b) + \text{l.o.t.}
\end{aligned}$$

$$\begin{aligned}
\not\partial_2^* \not\partial_1^* [e_3, \not\partial_1] \underline{\beta} &= \not\partial_2^* \not\partial_1^* \left(\left(-\frac{1}{2} \underline{\kappa} \not\partial_1 + \frac{1}{2} \underline{\vartheta} \not\partial_2^* - (\zeta - \eta) e_3 + \eta e_3 \Phi + \underline{\xi} (e_4 + e_4(\Phi)) + \underline{\beta} \right) \underline{\beta} \right) \\
&= -\frac{1}{2} \underline{\kappa} \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} + r^{-4} (\Gamma_b \cdot \not\partial^{\leq 3} \Gamma_b + \Gamma_b^{\leq 1} \cdot \not\partial^{\leq 2} \Gamma_b) + \text{l.o.t.}
\end{aligned}$$

Hence, schematically,

$$e_3 \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} = \not\partial_2^* \not\partial_1^* \not\partial_1 (e_3 \underline{\beta}) - \frac{3}{2} \underline{\kappa} \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} + r^{-4} \not\partial^{\leq 3} (\Gamma_b \cdot \Gamma_b)$$

Using the Bianchi identity $e_3 \underline{\beta} = \not\partial_2 \underline{\alpha} - 2(\underline{\kappa} + \underline{\omega}) \underline{\beta} + (-2\zeta + \eta) \underline{\alpha} + 3\underline{\xi} \rho$ we further deduce,

$$\not\partial_2^* \not\partial_1^* \not\partial_1 (e_3 \underline{\beta}) = \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \underline{\alpha} - 2(\underline{\kappa} + \underline{\omega}) \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} + 3\rho \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\xi} + r^{-4} \not\partial^{\leq 3} (\Gamma_b \cdot \Gamma_b)$$

i.e.,

$$\begin{aligned}
e_3 \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} &= \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \underline{\alpha} - 2(\underline{\kappa} + \underline{\omega}) \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} + 3\rho \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\xi} - \frac{3}{2} \underline{\kappa} \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} \\
&\quad + r^{-4} \not\partial^{\leq 3} (\Gamma_b \cdot \Gamma_b)
\end{aligned} \tag{A.12.2}$$

²In particular we write $\underline{\beta} \in r^{-1} \Gamma_b$.

³We also commute once more e_3 and e_4 with $\not\partial_1^*$, $\not\partial_1$, $\not\partial_2^*$, $\not\partial_2$ and use Bianchi.

We next calculate the second term $e_3(\rho \phi_2^* \phi_1^* \kappa)$ on the right hand side of (A.12.1)

$$e_3(\rho \phi_2^* \phi_1^* \kappa) = \rho \phi_2^* \phi_1^* e_3 \kappa + \rho [e_3, \phi_2^*] \phi_1^* \kappa + \rho \phi_2^* [e_3, \phi_1^*] \kappa + e_3(\rho) \phi_2^* \phi_1^* \kappa.$$

Using the equation for $e_3 \kappa$ in Proposition 2.2.1 we derive,

$$\begin{aligned} \rho \phi_2^* \phi_1^* e_3 \kappa &= \rho \phi_2^* \phi_1^* \left(-\frac{1}{2} \kappa^2 - 2\omega \kappa + 2\phi_1 \xi + \Gamma_b \cdot \Gamma_b \right), \\ &= -\rho(\kappa + 2\omega) \phi_2^* \phi_1^* \kappa - 2\rho \kappa \phi_2^* \phi_1^* \omega + 2\rho \phi_2^* \phi_1^* \phi_1 \xi + r^{-5} \mathfrak{P}^{\leq 2} \Gamma_b \cdot \Gamma_b \end{aligned}$$

Also,

$$\begin{aligned} [e_3, \phi_2^*] \phi_1^* \kappa &= \left(-\frac{1}{2} \kappa \phi_2^* + \frac{1}{2} \vartheta \phi_1 + (\zeta - \eta) e_3 + \eta e_3 \Phi - \xi(e_4 - e_4(\Phi)) + \underline{\beta} \right) \phi_1^* \kappa \\ &= -\frac{1}{2} \kappa \phi_2^* \phi_1^* \kappa + r^{-2} \Gamma_b \cdot \mathfrak{P}^{\leq 2} \Gamma_g, \\ \phi_2^* [e_3, \phi_1^*] \kappa &= \phi_2^* \left(\left(-\frac{1}{2} \kappa \phi_1^* - \frac{1}{2} \vartheta \phi_1 + (\zeta - \eta) e_3 - \xi e_4 \right) \kappa \right), \\ &= -\frac{1}{2} \kappa \phi_2^* \phi_1^* \kappa + \phi_2^* (\zeta - \eta) e_3 \kappa - \phi_2^* \xi e_4 \kappa + r^{-2} \mathfrak{P}^{\leq 2} (\Gamma_b \cdot \Gamma_g) \end{aligned}$$

Using also. the Bianchi equation

$$e_3 \rho = -\frac{3}{2} \kappa \rho + \phi_1 \underline{\beta} - \frac{1}{2} \vartheta \alpha - \zeta \underline{\beta} + 2(\eta \underline{\beta} + \xi \beta)$$

We deduce,

$$\begin{aligned} e_3(\rho \phi_2^* \phi_1^* \kappa) &= -\rho(\kappa + 2\omega) \phi_2^* \phi_1^* \kappa - 2\rho \kappa \phi_2^* \phi_1^* \omega + 2\rho \phi_2^* \phi_1^* \phi_1 \xi \\ &\quad - \rho \kappa \phi_2^* \phi_1^* \kappa + \rho \left(\phi_2^* (\zeta - \eta) e_3 \kappa - \phi_2^* (\xi) e_4 \kappa \right) - \frac{3}{2} \rho \kappa \phi_2^* \phi_1^* \kappa + r^{-5} \mathfrak{P}^{\leq 2} (\Gamma_b \cdot \Gamma_g) \end{aligned}$$

i.e.,

$$\begin{aligned} e_3(\rho \phi_2^* \phi_1^* \kappa) &= -\frac{7}{2} \rho \kappa \phi_2^* \phi_1^* \kappa - 2\rho \omega \phi_2^* \phi_1^* \kappa - 2\rho \kappa \phi_2^* \phi_1^* \omega + 2\rho \phi_2^* \phi_1^* \phi_1 \xi \\ &\quad + \rho \left(\phi_2^* (\zeta - \eta) e_3 \kappa - \phi_2^* (\xi) e_4 \kappa \right) + r^{-5} \mathfrak{P}^{\leq 2} (\Gamma_b \cdot \Gamma_g). \end{aligned} \tag{A.12.3}$$

Now,

$$\begin{aligned}
& \phi_2^*(\zeta - \eta)e_{3\underline{\kappa}} - \phi_2^*(\xi)e_{4\underline{\kappa}} \\
&= \phi_2^*(\zeta - \eta) \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega}\underline{\kappa} + 2\phi_1\underline{\xi} + \Gamma_b \cdot \Gamma_b \right) \\
&\quad - \phi_2^*\underline{\xi} \left(-\frac{1}{2}\underline{\kappa}\underline{\kappa} + 2\underline{\omega}\underline{\kappa} + 2\phi_1\underline{\eta} + 2\rho + \Gamma_g \cdot \Gamma_b \right) \\
&= \phi_2^*(\zeta - \eta) \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega}\underline{\kappa} \right) - \phi_2^*\underline{\xi} \left(-\frac{1}{2}\underline{\kappa}\underline{\kappa} + 2\underline{\omega}\underline{\kappa} + 2\rho \right) + r^{-2} \mathfrak{O}^{\leq 1}\Gamma_b \cdot \mathfrak{O}^{\leq 1}\Gamma_b.
\end{aligned}$$

Therefore, back to (A.12.3),

$$\begin{aligned}
e_3(\rho \phi_2^* \phi_1^* \underline{\kappa}) &= -\frac{7}{2}\rho \underline{\kappa} \phi_2^* \phi_1^* (\underline{\kappa}) - 2\rho \underline{\omega} \phi_2^* \phi_1^* \underline{\kappa} - 2\rho \underline{\kappa} \phi_2^* \phi_1^* \underline{\omega} + 2\rho \phi_2^* \phi_1^* \phi_1 \underline{\xi} \\
&\quad + \rho \phi_2^*(\zeta - \eta) \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega}\underline{\kappa} \right) - \rho \phi_2^*\underline{\xi} \left(-\frac{1}{2}\underline{\kappa}\underline{\kappa} + 2\underline{\omega}\underline{\kappa} + 2\rho \right) \quad (\text{A.12.4}) \\
&\quad + r^{-5} \mathfrak{O}^{\leq 2}(\Gamma_b \cdot \Gamma_g)..
\end{aligned}$$

We next estimate the third term $e_3(\underline{\kappa}\rho \phi_2^*\zeta)$ on the right hand side of (A.12.1),

$$e_3(\underline{\kappa}\rho \phi_2^*\zeta) = \underline{\kappa}\rho \phi_2^*(e_3\zeta) + \underline{\kappa}\rho[e_3, \phi_2^*]\zeta + e_3(\underline{\kappa})\rho \phi_2^*\zeta + \underline{\kappa}e_3(\rho) \phi_2^*\zeta$$

Using again the equations

$$\begin{aligned}
e_3(\underline{\kappa}) &= -\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega}\underline{\kappa} + 2\phi_1\underline{\xi} + \Gamma_b \cdot \Gamma_b \\
e_3\rho &= -\frac{3}{2}\underline{\kappa}\rho + \phi_1\underline{\beta} + \Gamma_g \cdot \Gamma_b
\end{aligned}$$

i.e.,

$$\begin{aligned}
e_3(\underline{\kappa})\rho \phi_2^*\zeta &= \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega}\underline{\kappa} \right) \rho \phi_2^*\zeta + r^{-5} \mathfrak{O}\Gamma_b \cdot \mathfrak{O}\Gamma_g \\
\underline{\kappa}e_3\rho \phi_2^*\zeta &= -\frac{3}{2}\underline{\kappa}^2\rho + r^{-4} \mathfrak{O}\Gamma_b \cdot \mathfrak{O}\Gamma_g
\end{aligned}$$

the equation,

$$e_3\zeta = -\frac{1}{2}\underline{\kappa}(\zeta + \eta) + 2\underline{\omega}(\zeta - \eta) + \underline{\beta} + 2\phi_1^*\underline{\omega} + 2\underline{\omega}\underline{\xi} + \frac{1}{2}\underline{\kappa}\underline{\xi} + \Gamma_b \cdot \Gamma_b$$

and the commutator formula,

$$\begin{aligned} [e_3, \not{d}_2^*] \zeta &= \left(-\frac{1}{2} \underline{\kappa} \not{d}_2^* + \frac{1}{2} \not{v} \not{d}_1 + (\zeta - \eta) e_3 + \eta e_3 \Phi - \underline{\xi} (e_4 - e_4(\Phi)) + \underline{\beta} \right) \zeta \\ &= -\frac{1}{2} \underline{\kappa} \not{d}_2^* \zeta + r^{-1} \Gamma_b \cdot \not{v}^{\leq 1} \Gamma_g \end{aligned}$$

we deduce

$$\begin{aligned} e_3(\underline{\kappa} \rho \not{d}_2^* \zeta) &= \underline{\kappa} \rho \not{d}_2^* \left(-\frac{1}{2} \underline{\kappa} (\zeta + \eta) + 2\underline{\omega} (\zeta - \eta) + \underline{\beta} + 2 \not{d}_1^* \underline{\omega} + 2\underline{\omega} \underline{\xi} + \frac{1}{2} \underline{\kappa} \underline{\xi} + \Gamma_b \cdot \Gamma_b \right) \\ &\quad - \frac{1}{2} \underline{\kappa}^2 \rho \not{d}_2^* \zeta + r^{-5} \Gamma_b \cdot \not{v}^{\leq 1} \Gamma_g \\ &\quad - \frac{1}{2} \underline{\kappa}^2 \rho \not{d}_2^* \zeta - 2\underline{\omega} \underline{\kappa} \rho \not{d}_2^* \zeta + r^{-5} \not{v}^{\leq 1} \Gamma_b \cdot \not{v}^{\leq 1} \Gamma_g \\ &\quad - \frac{3}{2} \underline{\kappa}^2 \rho \not{d}_2^* \zeta + r^{-4} \not{v} \Gamma_b \cdot \not{v} \Gamma_g \\ &= \underline{\kappa} \rho \left(-3 \underline{\kappa} \not{d}_2^* \zeta - \frac{1}{2} \underline{\kappa} \not{d}_2^* \eta - 2\underline{\omega} \not{d}_2^* \eta + \not{d}_2^* \underline{\beta} + 2 \not{d}_2^* \not{d}_1^* \underline{\omega} + 2\underline{\omega} \not{d}_2^* \underline{\xi} + \frac{1}{2} \underline{\kappa} \not{d}_2^* \underline{\xi} \right) \\ &\quad + r^{-4} \not{v}^{\leq 1} \Gamma_b \cdot \not{v}^{\leq 1} \Gamma_g + r^{-5} \not{v}^{\leq 1} \Gamma_b \cdot \not{v}^{\leq 1} \Gamma_b \end{aligned}$$

i.e.,

$$\begin{aligned} e_3(\underline{\kappa} \rho \not{d}_2^* \zeta) &= \underline{\kappa} \rho \left(-3 \underline{\kappa} \not{d}_2^* \zeta - \frac{1}{2} \underline{\kappa} \not{d}_2^* \eta - 2\underline{\omega} \not{d}_2^* \eta + \not{d}_2^* \underline{\beta} + 2 \not{d}_2^* \not{d}_1^* \underline{\omega} + 2\underline{\omega} \not{d}_2^* \underline{\xi} + \frac{1}{2} \underline{\kappa} \not{d}_2^* \underline{\xi} \right) \\ &\quad + r^{-4} \not{v}^{\leq 1} \Gamma_b \cdot \not{v}^{\leq 1} \Gamma_g + r^{-5} \not{v}^{\leq 1} \Gamma_b \cdot \not{v}^{\leq 1} \Gamma_b \end{aligned} \tag{A.12.5}$$

For the fourth term on the right hand side of (A.12.1) we have

$$\begin{aligned} e_3(\underline{\kappa} \rho \underline{\alpha}) &= \underline{\kappa} \rho e_3(\underline{\alpha}) + e_3(\underline{\kappa}) \rho \underline{\alpha} + \underline{\kappa} e_3(\rho) \underline{\alpha} \\ &= \underline{\kappa} \rho e_3(\underline{\alpha}) + \left(-\frac{1}{2} \underline{\kappa} \underline{\kappa} + 2\underline{\omega} \underline{\kappa} + 2 \not{d}_1 \eta + 2\rho - \frac{1}{2} \not{v} \not{v} + 2(\underline{\xi} \underline{\xi} + \eta \eta) \right) \rho \underline{\alpha} \\ &\quad + \underline{\kappa} \left(-\frac{3}{2} \underline{\kappa} \rho + \not{d}_1 \underline{\beta} - \frac{1}{2} \not{v} \underline{\alpha} - \zeta \underline{\beta} + 2(\eta \underline{\beta} + \underline{\xi} \beta) \right) \underline{\alpha} \\ &= \underline{\kappa} \rho e_3(\underline{\alpha}) + (-2 \underline{\kappa} \underline{\kappa} + 2\underline{\omega} \underline{\kappa} + 2\rho) \rho \underline{\alpha} + (r^{-3} \not{v} \Gamma_b + r^{-2} \Gamma_b \cdot \Gamma_b) \cdot \underline{\alpha} \end{aligned}$$

i.e.,

$$e_3(\underline{\kappa} \rho \underline{\alpha}) = \underline{\kappa} \rho e_3(\underline{\alpha}) + (-2 \underline{\kappa} \underline{\kappa} + 2\underline{\omega} \underline{\kappa} + 2\rho) \rho \underline{\alpha} + (r^{-3} \not{v} \Gamma_b + r^{-2} \Gamma_b \cdot \Gamma_b) \cdot \underline{\alpha}. \tag{A.12.6}$$

Finally, for the fifth term on the right hand side of (A.12.1), using the e_3 equations for

$\underline{\vartheta}, \rho, \underline{\kappa}, \kappa,$

$$\begin{aligned}
e_3\left((2\rho^2 - \kappa\underline{\kappa}\rho)\underline{\vartheta}\right) &= (2\rho^2 - \kappa\underline{\kappa}\rho)e_3\underline{\vartheta} + 4\rho e_3(\rho)\underline{\vartheta} - e_3(\kappa)\underline{\kappa}\rho\underline{\vartheta} - \kappa e_3(\underline{\kappa})\rho\underline{\vartheta} - \kappa\underline{\kappa}e_3(\rho)\underline{\vartheta} \\
&= (2\rho^2 - \kappa\underline{\kappa}\rho)\left(-\underline{\kappa}\underline{\vartheta} - 2\underline{\omega}\underline{\vartheta} - 2\underline{\alpha} - 2\underline{\phi}_2^*\underline{\xi} + \Gamma_b \cdot \Gamma_b\right) \\
&\quad + 4\rho\left(-\frac{3}{2}\underline{\kappa}\rho + \underline{\phi}_1\underline{\beta} + \Gamma_g \cdot \Gamma_b\right)\underline{\vartheta} \\
&\quad - \left(-\frac{1}{2}\underline{\kappa}\kappa + 2\underline{\omega}\kappa + 2\rho + 2\underline{\phi}_1\underline{\eta} + \Gamma_b \cdot \Gamma_b\right)\underline{\kappa}\rho\underline{\vartheta} \\
&\quad - \kappa\left(-\frac{1}{2}\kappa^2 - 2\underline{\omega}\kappa + 2\underline{\phi}_1\underline{\xi} + \Gamma_b \cdot \Gamma_b\right)\rho\underline{\vartheta} \\
&\quad - \kappa\underline{\kappa}\left(-\frac{3}{2}\underline{\kappa}\rho + \underline{\phi}_1\underline{\beta} + \Gamma_g \cdot \Gamma_b\right)\underline{\vartheta}
\end{aligned}$$

i.e.,

$$\begin{aligned}
e_3\left((2\rho^2 - \kappa\underline{\kappa}\rho)\underline{\vartheta}\right) &= (2\rho^2 - \kappa\underline{\kappa}\rho)\left(-\underline{\kappa}\underline{\vartheta} - 2\underline{\omega}\underline{\vartheta} - 2\underline{\alpha} - 2\underline{\phi}_2^*\underline{\xi}\right) - 6\underline{\kappa}\rho^2\underline{\vartheta} \\
&\quad - \left(-\frac{1}{2}\underline{\kappa}\kappa + 2\underline{\omega}\kappa + 2\rho\right)\underline{\kappa}\rho\underline{\vartheta} + \kappa\left(\frac{1}{2}\kappa^2 + 2\underline{\omega}\kappa\right)\rho\underline{\vartheta} + \frac{3}{2}\kappa^2\rho \\
&\quad + r^{-5}\underline{\vartheta}^{\leq 1}\Gamma_b \cdot \Gamma_b
\end{aligned}$$

from which,

$$\begin{aligned}
e_3\left((2\rho^2 - \kappa\underline{\kappa}\rho)\underline{\vartheta}\right) &= (2\rho^2 - \kappa\underline{\kappa}\rho)\left(-2\underline{\alpha} - 2\underline{\phi}_2^*\underline{\xi}\right) + \left(\frac{7}{2}\kappa\underline{\kappa}^2\rho - 10\underline{\kappa}\rho^2 + 2\kappa\underline{\kappa}\rho\underline{\omega} - 4\underline{\omega}\rho^2\right)\underline{\vartheta} \\
&\quad + r^{-5}\underline{\vartheta}^{\leq 1}\Gamma_b \cdot \Gamma_b. \tag{A.12.7}
\end{aligned}$$

Recalling (A.12.1)

$$\begin{aligned}
r^{-7}e_3(r^2e_3(r\mathbf{q})) &= e_3\underline{\phi}_2^*\underline{\phi}_1^*\underline{\phi}_1\underline{\beta} - \frac{3}{2}e_3(\rho\underline{\phi}_2^*\underline{\phi}_1^*\underline{\kappa}) - \frac{3}{2}e_3(\underline{\kappa}\rho\underline{\phi}_2^*\underline{\zeta}) - \frac{3}{2}e_3(\kappa\rho\underline{\alpha}) \\
&\quad + \frac{3}{4}e_3\left((2\rho^2 - \kappa\underline{\kappa}\rho)\underline{\vartheta}\right) + 7r^{-6}e_3(r)e_3(r\mathbf{q}) \\
&\quad + r^2e_3(\text{Err}[e_3\mathbf{q}]) + r\text{Err}[e_3\mathbf{q}] + \text{l.o.t.}
\end{aligned}$$

and making use of (A.12.4)–(A.12.7) we deduce,

$$\begin{aligned}
r^{-7}e_3(r^2e_3(r\mathbf{q})) &= \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \underline{\alpha} - 2(\underline{\kappa} + \underline{\omega}) \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} + 3\rho \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\xi} - \frac{3}{2}\underline{\kappa} \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} \\
&- \frac{3}{2} \left\{ -\frac{7}{2}\rho \underline{\kappa} \not\partial_2^* \not\partial_1^* (\underline{\kappa}) - 2\rho \underline{\omega} \not\partial_2^* \not\partial_1^* \underline{\kappa} - 2\rho \underline{\kappa} \not\partial_2^* \not\partial_1^* \underline{\omega} + 2\rho \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\xi} \right. \\
&\quad \left. + \rho \not\partial_2^* (\zeta - \eta) \left(-\frac{1}{2}\underline{\kappa}^2 - 2\underline{\omega} \underline{\kappa} \right) - \rho \not\partial_2^* \underline{\xi} \left(-\frac{1}{2}\underline{\kappa} \underline{\kappa} + 2\underline{\omega} \underline{\kappa} + 2\rho \right) \right\} \\
&- \frac{3}{2} \left\{ \underline{\kappa} \rho \left(-3\underline{\kappa} \not\partial_2^* \zeta - \frac{1}{2}\underline{\kappa} \not\partial_2^* \eta - 2\underline{\omega} \not\partial_2^* \eta + \not\partial_2^* \underline{\beta} + 2 \not\partial_2^* \not\partial_1^* \underline{\omega} + 2\underline{\omega} \not\partial_2^* \underline{\xi} + \frac{1}{2}\underline{\kappa} \not\partial_2^* \underline{\xi} \right) \right\} \\
&- \frac{3}{2} \left\{ \underline{\kappa} \rho e_3(\underline{\alpha}) + (-2\underline{\kappa} \underline{\kappa} + 2\underline{\omega} \underline{\kappa} + 2\rho) \rho \underline{\alpha} + (r^{-3} \not\partial \Gamma_b + r^{-2} \Gamma_b \cdot \Gamma_b) \cdot \underline{\alpha} \right\} \\
&+ \frac{3}{4} \left\{ (2\rho^2 - \underline{\kappa} \underline{\kappa} \rho) \left(-2\underline{\alpha} - 2 \not\partial_2^* \underline{\xi} \right) + \left(\frac{7}{2}\underline{\kappa} \underline{\kappa}^2 \rho - 10\underline{\kappa} \rho^2 + 2\underline{\kappa} \underline{\kappa} \rho \underline{\omega} - 4\underline{\omega} \rho^2 \right) \underline{\vartheta} \right\} \\
&+ 7r^{-6}e_3(r)e_3(r\mathbf{q}) + \text{Err} + r^{-7} \left(r^2e_3(\text{Err}[e_3\mathbf{q}]) + r\text{Err}[e_3\mathbf{q}] \right) + \text{l.o.t.}
\end{aligned}$$

where, the error term Err is given by

$$\text{Err} = (r^{-3} \not\partial \Gamma_b + r^{-2} \Gamma_b \cdot \Gamma_b) \cdot \underline{\alpha} + r^{-4} \not\partial^{\leq 1} \Gamma_b \cdot \not\partial^{\leq 1} \Gamma_g + r^{-5} \not\partial^{\leq 1} \Gamma_b \cdot \not\partial^{\leq 1} \Gamma_b \quad (\text{A.12.8})$$

Denoting the expression of left hand side of the identity (2.3.15) by I , i.e.

$$I := e_3(r^2e_3(r\mathbf{q})) + 2\underline{\omega}r^2e_3(r\mathbf{q})$$

we deduce,

$$\begin{aligned}
r^{-7}I &= \not\partial_2^* \not\partial_1^* \not\partial_1 \not\partial_2 \underline{\alpha} - \left(\frac{7}{2}\underline{\kappa} + 2\underline{\omega} \right) \not\partial_2^* \not\partial_1^* \not\partial_1 \underline{\beta} + \frac{3}{2} \left(\frac{7}{2}\underline{\kappa} + 2\underline{\omega} \right) \rho \not\partial_2^* \not\partial_1^* \underline{\kappa} + \frac{3}{2} \left(\frac{7}{2}\underline{\kappa} + 2\underline{\omega} \right) \underline{\kappa} \rho \not\partial_2^* \zeta \\
&- \frac{3}{2}\underline{\kappa} \rho \not\partial_2^* \underline{\beta} - \frac{3}{2}\underline{\kappa} \rho e_3(\underline{\alpha}) - \frac{3}{2} \left(-3\underline{\kappa} \underline{\kappa} + 2\underline{\omega} \underline{\kappa} + 4\rho \right) \rho \underline{\alpha} \\
&+ \frac{3}{4} \left(\frac{7}{2}\underline{\kappa} \underline{\kappa}^2 \rho - 10\underline{\kappa} \rho^2 + 2\underline{\kappa} \underline{\kappa} \rho \underline{\omega} - 4\underline{\omega} \rho^2 \right) \underline{\vartheta} + r^{-7} [7re_3(r)e_3(r\mathbf{q}) + 2\underline{\omega}r^2e_3(r\mathbf{q})] + \widetilde{\text{Err}}.
\end{aligned}$$

where the new error term $\widetilde{\text{Err}}$ is given by

$$\widetilde{\text{Err}} = \text{Err} + r^{-7} \left(r^2e_3(\text{Err}[e_3\mathbf{q}]) + r\text{Err}[e_3\mathbf{q}] \right) + 2\underline{\omega}r^{-5}\text{Err}[e_3\mathbf{q}]$$

To calculate the term $J := 7re_3(r)e_3(r\mathbf{q}) + 2\omega r^2 e_3(r\mathbf{q})$ in the last row we make use once more of the identity of Lemma 2.3.14 to derive

$$\begin{aligned} J &= r^2 \left(\frac{7}{2}\underline{\kappa} + 2\underline{\omega} \right) e_3(r\mathbf{q}) + 7r \left(e_3(r) - \frac{r}{2}\underline{\kappa} \right) e_3(r\mathbf{q}) \\ &= r^2 \left(\frac{7}{2}\underline{\kappa} + 2\underline{\omega} \right) e_3(r\mathbf{q}) + r^2 \Gamma_b e_3(r\mathbf{q}) \\ &= r^7 \left(\frac{7}{2}\underline{\kappa} + 2\underline{\omega} \right) \left\{ \phi_2^* \phi_1^* \phi_1 \underline{\beta} - \frac{3}{2} \rho \phi_2^* \phi_1^* \underline{\kappa} - \frac{3}{2} \underline{\kappa} \rho \phi_2^* \zeta - \frac{3}{2} \kappa \rho \underline{\alpha} + \frac{3}{4} (2\rho^2 - \kappa \underline{\kappa} \rho) \underline{\vartheta} \right\} \\ &+ r^2 \left(\frac{7}{2}\underline{\kappa} + 2\underline{\omega} \right) \text{Err}[e_3(r\mathbf{q})] + r^2 \Gamma_b e_3(r\mathbf{q}) \end{aligned}$$

i.e.,

$$\begin{aligned} r^{-7} J &= \left(\frac{7}{2}\underline{\kappa} + 2\underline{\omega} \right) \left\{ \phi_2^* \phi_1^* \phi_1 \underline{\beta} - \frac{3}{2} \rho \phi_2^* \phi_1^* \underline{\kappa} - \frac{3}{2} \underline{\kappa} \rho \phi_2^* \zeta - \frac{3}{2} \kappa \rho \underline{\alpha} + \frac{3}{4} (2\rho^2 - \kappa \underline{\kappa} \rho) \underline{\vartheta} \right\} \\ &+ r^{-5} \left(\frac{7}{2}\underline{\kappa} + 2\underline{\omega} \right) \text{Err}[e_3(r\mathbf{q})] + r^{-5} \Gamma_b e_3(r\mathbf{q}) \end{aligned}$$

Combining and simplifying,

$$r^{-7} I = \phi_2^* \phi_1^* \phi_1 \phi_2 \underline{\alpha} - \frac{3}{2} \underline{\kappa} \rho \phi_2^* \underline{\beta} - \frac{3}{2} \kappa \rho e_3(\underline{\alpha}) - \frac{3}{2} \left(\frac{1}{2} \kappa \underline{\kappa} + 4\underline{\omega} \kappa + 4\rho \right) \rho \underline{\alpha} - \frac{9}{4} \underline{\kappa} \rho^2 \underline{\vartheta} + \widetilde{\text{Err}}.$$

where,

$$\widetilde{\text{Err}} = \widetilde{\text{Err}} + r^{-5} \left(\frac{7}{2}\underline{\kappa} + 2\underline{\omega} \right) \text{Err}[e_3(r\mathbf{q})] + r^{-5} \Gamma_b e_3(r\mathbf{q})$$

Using Bianchi to replace $\phi_2^* \underline{\beta}$, we deduce

$$\begin{aligned} r^{-7} I &= \phi_2^* \phi_1^* \phi_1 \phi_2 \underline{\alpha} - \frac{3}{2} \underline{\kappa} \rho \left(-e_4 \underline{\alpha} - \frac{1}{2} \kappa \underline{\alpha} + 4\underline{\omega} \underline{\alpha} - \frac{3}{2} \rho \underline{\vartheta} + r^{-1} \Gamma_g \cdot \Gamma_b \right) \\ &- \frac{3}{2} \kappa \rho e_3(\underline{\alpha}) - \frac{3}{2} \left(\frac{1}{2} \kappa \underline{\kappa} + 4\underline{\omega} \kappa + 4\rho \right) \rho \underline{\alpha} - \frac{9}{4} \underline{\kappa} \rho^2 \underline{\vartheta} + r^{-5} \Gamma_g \cdot \Gamma_b + \text{l.o.t.} \\ &= \phi_2^* \phi_1^* \phi_1 \phi_2 \underline{\alpha} + \frac{3}{2} \underline{\kappa} \rho e_4 \underline{\alpha} - \frac{3}{2} \kappa \rho e_3(\underline{\alpha}) - 6 \left(\underline{\kappa} \omega + \underline{\omega} \kappa + \rho \right) \rho \underline{\alpha} + r^{-5} \Gamma_g \cdot \Gamma_b. \end{aligned}$$

$$\begin{aligned} I &= e_3(r^2 e_3(r\mathbf{q})) + 2\omega r^2 e_3(r\mathbf{q}) \\ &= r^7 \left\{ \phi_2^* \phi_1^* \phi_1 \phi_2 \underline{\alpha} + \frac{3}{2} \underline{\kappa} \rho e_4 \underline{\alpha} - \frac{3}{2} \kappa \rho e_3(\underline{\alpha}) \right\} + \text{Err}[ST] \end{aligned}$$

where,

$$\begin{aligned} \text{Err}[ST] &= r^7 \widetilde{\text{Err}} + r^2 \left(\frac{7}{2} \underline{\kappa} + 2\underline{\omega} \right) \text{Err}[e_3(r\mathbf{q})] + r^2 \Gamma_b e_3(r\mathbf{q}) + r^2 \Gamma_g \cdot \Gamma_b \\ &\quad + r^7 \text{Err} + \left(r^2 e_3(\text{Err}[e_3\mathbf{q}]) + r \text{Err}[e_3\mathbf{q}] \right) + 2\underline{\omega} r^2 \text{Err}[e_3\mathbf{q}] + r^2 \Gamma_b e_3(r\mathbf{q}) + r^2 \Gamma_g \cdot \Gamma_b \end{aligned}$$

Recall that, see (A.12.8),

$$\text{Err} = \left(r^{-3} \mathfrak{I} \Gamma_b + r^{-2} \Gamma_b \cdot \Gamma_b \right) \cdot \underline{\alpha} + r^{-4} \mathfrak{I}^{\leq 1} \Gamma_b \cdot \mathfrak{I}^{\leq 1} \Gamma_g + r^{-5} \mathfrak{I}^{\leq 1} \Gamma_b \cdot \mathfrak{I}^{\leq 1} \Gamma_b$$

Hence,

$$\begin{aligned} \text{Err}[ST] &= r^4 \left(\mathfrak{I} \Gamma_b + r \Gamma_b \cdot \Gamma_b \right) \cdot \underline{\alpha} + r^3 \mathfrak{I}^{\leq 1} \Gamma_b \cdot \mathfrak{I}^{\leq 1} \Gamma_g + r^2 \mathfrak{I}^{\leq 1} \Gamma_b \cdot \mathfrak{I}^{\leq 1} \Gamma_b + r^2 \Gamma_b e_3(r\mathbf{q}) \\ &\quad + r^2 e_3(\text{Err}[e_3\mathbf{q}]) + r \text{Err}[e_3\mathbf{q}] + 2\underline{\omega} r^2 \text{Err}[e_3\mathbf{q}] \end{aligned}$$

Recall that, see Proposition 2.3.14,

$$\text{Err}[e_3(r\mathbf{q})] = r \Gamma_b \mathbf{q} + r^5 \mathfrak{d}^{\leq 1}(e_3 \eta \cdot \beta) + r^3 \mathfrak{d}^{\leq 2}(\Gamma_b \cdot \Gamma_g).$$

Therefore,

$$\begin{aligned} E &= r^2 e_3(\text{Err}[e_3\mathbf{q}]) + r \text{Err}[e_3\mathbf{q}] + 2\underline{\omega} r^2 \text{Err}[e_3\mathbf{q}] \\ &= r^2 (\Gamma_b e_3(r\mathbf{q}) + e_3(\Gamma_b) r \mathbf{q}) + r^7 \mathfrak{d}^{\leq 2}(e_3 \eta \cdot \beta) + r^5 \mathfrak{d}^{\leq 3}(\Gamma_b \cdot \Gamma_g) \\ &\quad + r^2 \Gamma_b \mathbf{q} + r^6 \mathfrak{d}^{\leq 1}(e_3 \eta \cdot \beta) + r^4 \mathfrak{d}^{\leq 2}(\Gamma_b \cdot \Gamma_g) \\ &= r^2 (\Gamma_b e_3(r\mathbf{q}) + (\mathfrak{d}^{\leq 1} \Gamma_b) r \mathbf{q}) + r^7 \mathfrak{d}^{\leq 2}(e_3 \eta \cdot \beta) + r^5 \mathfrak{d}^{\leq 3}(\Gamma_b \cdot \Gamma_g) \end{aligned}$$

Thus,

$$\begin{aligned} \text{Err}[TS] &= r^4 \left(\mathfrak{I} \Gamma_b + r \Gamma_b \cdot \Gamma_b \right) \cdot \underline{\alpha} + r^2 (\Gamma_b e_3(r\mathbf{q}) + (\mathfrak{d}^{\leq 1} \Gamma_b) r \mathbf{q}) \\ &\quad + r^7 \mathfrak{d}^{\leq 2}(e_3 \eta \cdot \beta) + r^5 \mathfrak{d}^{\leq 3}(\Gamma_b \cdot \Gamma_g) \end{aligned}$$

which end the proof of Proposition 2.3.15.

A.13 Proof of Proposition 2.4.6

In this section we give a proof of Proposition 2.4.6, i.e. we derive the wave equation for the extreme curvature component α ,

$$\begin{aligned} \square_{\mathbf{g}} \alpha &= -4\underline{\omega} e_4(\alpha) + (2\underline{\kappa} + 4\underline{\omega}) e_3(\alpha) + V\alpha + \text{Err}(\square_{\mathbf{g}} \alpha), \\ V &:= \left(-4e_4(\underline{\omega}) + \frac{1}{2} \underline{\kappa} \underline{\kappa} - 10\underline{\kappa} \underline{\omega} + 2\underline{\kappa} \underline{\omega} - 8\underline{\omega} \underline{\omega} - 4\rho + 4e_\theta(\Phi)^2 \right) \alpha, \end{aligned} \quad (\text{A.13.1})$$

where

$$\begin{aligned}
\text{Err}(\square_{\mathbf{g}}\alpha) &= \frac{1}{2}\vartheta e_3(\alpha) + \frac{3}{4}\vartheta^2\rho + e_\theta(\Phi)\vartheta\beta - \frac{1}{2}\kappa(\zeta + 4\underline{\eta})\beta - (\zeta + \underline{\eta})e_4(\beta) - \xi e_3(\beta) \\
&+ e_\theta(\Phi)(2\zeta + \underline{\eta})\alpha + \beta^2 + e_4(\Phi)\underline{\eta}\beta + e_3(\Phi)\xi\beta - (\zeta + 4\underline{\eta})e_4(\beta) - (e_4(\zeta) + 4e_4(\underline{\eta}))\beta \\
&- 2(\kappa + \omega)(\zeta + 4\underline{\eta})\beta + 2e_\theta(\kappa + \omega)\beta - e_\theta((2\zeta + \underline{\eta})\alpha) - 3\xi e_\theta(\rho) + 2\underline{\eta}e_\theta(\alpha) \\
&+ \frac{3}{2}\vartheta(e_\theta(\beta) + e_\theta(\Phi)\beta) + 3\rho(\underline{\eta} + \eta + 2\zeta)\xi + (e_\theta(\underline{\eta}) + e_\theta(\Phi)\underline{\eta})\alpha + \frac{1}{4}\kappa\vartheta\alpha - 2\underline{\omega}\vartheta\alpha \\
&- \frac{1}{2}\vartheta\underline{\vartheta}\alpha + \xi\underline{\xi}\alpha + \underline{\eta}^2\alpha + \frac{3}{2}\vartheta\zeta\beta + 3\vartheta(\underline{\eta}\beta + \xi\beta) - \frac{1}{2}\vartheta(\zeta + 4\underline{\eta})\beta.
\end{aligned}$$

The equation for $\underline{\alpha}$ can then be easily inferred by symmetry.

Proof. We make use of the Bianchi identities

$$\begin{aligned}
e_\theta(\beta) - e_\theta(\Phi)\beta &= e_3(\alpha) + \left(\frac{\kappa}{2} - 4\underline{\omega}\right)\alpha + \frac{3}{2}\vartheta\rho - (\zeta + 4\underline{\eta})\beta, \\
e_4(\beta) + 2(\kappa + \omega)\beta &= e_\theta(\alpha) + 2e_\theta(\Phi)\alpha + (2\zeta + \underline{\eta})\alpha + 3\xi\rho.
\end{aligned}$$

to infer that

$$\begin{aligned}
e_4(e_3(\alpha)) &= e_4(e_\theta(\beta)) - e_\theta(\Phi)e_4(\beta) - e_4(e_\theta(\Phi))\beta - \left(\frac{\kappa}{2} - 4\underline{\omega}\right)e_4(\alpha) - \left(\frac{e_4(\kappa)}{2} - 4e_4(\underline{\omega})\right)\alpha \\
&- \frac{3}{2}\vartheta e_4(\rho) - \frac{3}{2}e_4(\vartheta)\rho + (\zeta + 4\underline{\eta})e_4(\beta) + (e_4(\zeta) + 4e_4(\underline{\eta}))\beta \\
&= e_4(e_\theta(\beta)) - e_\theta(\Phi)\left(e_\theta(\alpha) + 2e_\theta(\Phi)\alpha - 2(\kappa + \omega)\beta + (2\zeta + \underline{\eta})\alpha + 3\xi\rho\right) \\
&- (D_4D_\theta\Phi + D_{D_4e_\theta}\Phi)\beta - \left(\frac{\kappa}{2} - 4\underline{\omega}\right)e_4(\alpha) - \left(\frac{e_4(\kappa)}{2} - 4e_4(\underline{\omega})\right)\alpha \\
&- \frac{3}{2}\vartheta e_4(\rho) - \frac{3}{2}e_4(\vartheta)\rho + (\zeta + 4\underline{\eta})e_4(\beta) + (e_4(\zeta) + 4e_4(\underline{\eta}))\beta.
\end{aligned}$$

Hence,

$$\begin{aligned}
e_4(e_3(\alpha)) &= e_4(e_\theta(\beta)) - e_\theta(\Phi)(e_\theta(\alpha) + 2e_\theta(\Phi)\alpha) + 2e_\theta(\Phi)(\kappa + \omega)\beta - 3e_\theta(\Phi)\xi\rho \\
&+ e_4(\Phi)e_\theta(\Phi)\beta - \left(\frac{\kappa}{2} - 4\underline{\omega}\right)e_4(\alpha) - \left(\frac{e_4(\kappa)}{2} - 4e_4(\underline{\omega})\right)\alpha \\
&- \frac{3}{2}\vartheta e_4(\rho) - \frac{3}{2}e_4(\vartheta)\rho - e_\theta(\Phi)(2\zeta + \underline{\eta})\alpha - \beta^2 - e_4(\Phi)\underline{\eta}\beta - e_3(\Phi)\xi\beta \\
&+ (\zeta + 4\underline{\eta})e_4(\beta) + (e_4(\zeta) + 4e_4(\underline{\eta}))\beta
\end{aligned}$$

and

$$\begin{aligned}
e_\theta(e_\theta(\alpha)) &= e_\theta(e_4(\beta)) + 2(\kappa + \omega)e_\theta(\beta) + 2e_\theta(\kappa + \omega)\beta - 2e_\theta(\Phi)e_\theta(\alpha) - 2e_\theta(e_\theta(\Phi))\alpha \\
&\quad - e_\theta((2\underline{\zeta} + \underline{\eta})\alpha) - 3e_\theta(\xi\rho) \\
&= e_\theta(e_4(\beta)) + 2(\kappa + \omega)\left(e_\theta(\Phi)\beta + e_3(\alpha) + \left(\frac{\kappa}{2} - 4\underline{\omega}\right)\alpha + \frac{3}{2}\vartheta\rho - (\zeta + 4\underline{\eta})\beta\right) \\
&\quad + 2e_\theta(\kappa + \omega)\beta - 2e_\theta(\Phi)e_\theta(\alpha) - 2(D_\theta D_\theta \Phi + D_{D_\theta e_\theta} \Phi)\alpha - e_\theta((2\underline{\zeta} + \underline{\eta})\alpha) - 3e_\theta(\xi\rho) \\
&= e_\theta(e_4(\beta)) + 2(\kappa + \omega)e_\theta(\Phi)\beta + 2(\kappa + \omega)e_3(\alpha) + 2(\kappa + \omega)\left(\frac{\kappa}{2} - 4\underline{\omega}\right)\alpha + 3(\kappa + \omega)\vartheta\rho \\
&\quad - 2e_\theta(\Phi)e_\theta(\alpha) - 2\left(\rho - e_\theta(\Phi)^2 + \frac{1}{2}\underline{\chi}e_4(\Phi) + \frac{1}{2}\chi e_3(\Phi)\right)\alpha - 3e_\theta(\xi)\rho \\
&\quad - 2(\kappa + \omega)(\zeta + 4\underline{\eta})\beta + 2e_\theta(\kappa + \omega)\beta - e_\theta((2\underline{\zeta} + \underline{\eta})\alpha) - 3\underline{\xi}e_\theta(\rho).
\end{aligned}$$

In view of Lemma 2.4.1, we have

$$\square_{\mathbf{g}}f = -e_4(e_3(f)) + e_\theta(e_\theta(f)) - \frac{1}{2}\underline{\kappa}e_4(f) + \left(-\frac{1}{2}\kappa + 2\omega\right)e_3(f) + e_\theta(\Phi)e_\theta(f) + 2\underline{\eta}e_\theta(f).$$

We infer

$$\begin{aligned}
\square_{\mathbf{g}}\alpha &= -e_4(e_3(\alpha)) + e_\theta(e_\theta(\alpha)) - \frac{1}{2}\underline{\kappa}e_4(\alpha) + \left(-\frac{1}{2}\kappa + 2\omega\right)e_3(\alpha) + e_\theta(\Phi)e_\theta(\alpha) + 2\underline{\eta}e_\theta(\alpha) \\
&= [e_\theta, e_4](\beta) - e_4(\Phi)e_\theta(\Phi)\beta + \frac{3}{2}\vartheta e_4(\rho) + \frac{3}{2}e_4(\vartheta)\rho + 3(\kappa + \omega)\vartheta\rho - 3(e_\theta(\xi) - e_\theta(\Phi)\xi)\rho \\
&\quad - 4\underline{\omega}e_4(\alpha) + \left(\frac{3}{2}\kappa + 4\omega\right)e_3(\alpha) \\
&\quad + \left(\frac{e_4(\underline{\kappa})}{2} - 4e_4(\underline{\omega}) + \kappa\underline{\kappa} - 8\underline{\kappa}\omega + \underline{\kappa}\omega - 8\omega\underline{\omega} - 2\rho + 4e_\theta(\Phi)^2 - \underline{\chi}e_4(\Phi) - \chi e_3(\Phi)\right)\alpha \\
&\quad + e_\theta(\Phi)(2\underline{\zeta} + \underline{\eta})\alpha + \beta^2 + e_4(\Phi)\underline{\eta}\beta + e_3(\Phi)\xi\beta - (\zeta + 4\underline{\eta})e_4(\beta) - (e_4(\zeta) + 4e_4(\underline{\eta}))\beta \\
&\quad - 2(\kappa + \omega)(\zeta + 4\underline{\eta})\beta + 2e_\theta(\kappa + \omega)\beta - e_\theta((2\underline{\zeta} + \underline{\eta})\alpha) - 3\underline{\xi}e_\theta(\rho) + 2\underline{\eta}e_\theta(\alpha).
\end{aligned}$$

Next, we have

$$\begin{aligned}
[e_\theta, e_4](\beta) &= \chi e_\theta(\beta) - (\zeta + \underline{\eta})e_4(\beta) - \xi e_3(\beta) \\
&= \chi\left(e_\theta(\Phi)\beta + e_3(\alpha) + \left(\frac{\kappa}{2} - 4\underline{\omega}\right)\alpha + \frac{3}{2}\vartheta\rho - (\zeta + 4\underline{\eta})\beta\right) \\
&\quad - (\zeta + \underline{\eta})e_4(\beta) - \xi e_3(\beta)
\end{aligned}$$

and hence

$$\begin{aligned}\square_{\mathbf{g}}\alpha &= -4\underline{\omega}e_4(\alpha) + \left(\frac{3}{2}\kappa + \chi + 4\underline{\omega}\right) e_3(\alpha) + V_1\alpha \\ &+ \frac{3}{2}\vartheta e_4(\rho) + \frac{3}{2}e_4(\vartheta)\rho + 3(\kappa + \omega)\vartheta\rho - 3(e_\theta(\xi) - e_\theta(\Phi)\xi)\rho + \frac{3}{2}\chi\vartheta\rho \\ &+ \text{Err}_1\end{aligned}$$

where,

$$\begin{aligned}V_1 &:= \frac{e_4(\underline{\kappa})}{2} - 4e_4(\underline{\omega}) + \kappa\underline{\kappa} - 8\kappa\underline{\omega} + \underline{\kappa}\omega - 8\underline{\omega}\omega - 2\rho + 4e_\theta(\Phi)^2 - \underline{\chi}e_4(\Phi) - \chi e_3(\Phi) \\ &+ \chi\frac{\underline{\kappa}}{2} - 4\underline{\chi}\omega,\end{aligned}$$

$$\begin{aligned}\text{Err}_1 &:= e_\theta(\Phi)\vartheta\beta - \chi(\zeta + 4\underline{\eta})\beta - (\zeta + \underline{\eta})e_4(\beta) - \xi e_3(\beta) \\ &+ e_\theta(\Phi)(2\zeta + \underline{\eta})\alpha + \beta^2 + e_4(\Phi)\underline{\eta}\beta + e_3(\Phi)\xi\beta - (\zeta + 4\underline{\eta})e_4(\beta) - (e_4(\zeta) + 4e_4(\underline{\eta}))\beta \\ &- 2(\kappa + \omega)(\zeta + 4\underline{\eta})\beta + 2e_\theta(\kappa + \omega)\beta - e_\theta((2\zeta + \underline{\eta})\alpha) - 3\xi e_\theta(\rho) + 2\underline{\eta}e_\theta(\alpha).\end{aligned}$$

Next, we make use of

$$\begin{aligned}e_4(\vartheta) + \kappa\vartheta + 2\underline{\omega}\vartheta &= -2\alpha + 2(e_\theta(\xi) - e_\theta(\Phi)\xi) + 2(\underline{\eta} + \eta + 2\zeta)\xi, \\ e_4(\rho) + \frac{3}{2}\kappa\rho &= e_\theta(\beta) + e_\theta(\Phi)\beta - \frac{1}{2}\vartheta\alpha + \zeta\beta + 2(\underline{\eta}\beta + \xi\beta),\end{aligned}$$

to calculate the term

$$\begin{aligned}I &:= \frac{3}{2}\vartheta e_4(\rho) + \frac{3}{2}e_4(\vartheta)\rho + 3(\kappa + \omega)\vartheta\rho - 3(e_\theta(\xi) - e_\theta(\Phi)\xi)\rho + \frac{3}{2}\chi\vartheta\rho \\ &= \frac{3}{2}\vartheta\left(-\frac{3}{2}\kappa\rho + \underline{\phi}_1\beta\right) + \frac{3}{2}\rho(-\kappa\vartheta - 2\underline{\omega}\vartheta - 2\alpha + 2(e_\theta(\xi) - e_\theta(\Phi)\xi)) + 3(\kappa + \omega)\vartheta\rho \\ &- 3(e_\theta(\xi) - e_\theta(\Phi)\xi)\rho + \frac{3}{2}\frac{\kappa + \vartheta}{2}\vartheta\rho + \text{l.o.t.} \\ &= -3\rho\alpha + \frac{3}{2}\vartheta\underline{\phi}_1\beta + \frac{3}{4}\vartheta^2\rho.\end{aligned}$$

Hence,

$$\begin{aligned}\square_{\mathbf{g}}\alpha &= -4\underline{\omega}e_4(\alpha) + \left(\frac{3}{2}\kappa + \chi + 4\underline{\omega}\right) e_3(\alpha) + (V_1 - 3\rho)\alpha \\ &+ \text{Err}_1 + \frac{3}{2}\vartheta\underline{\phi}_1\beta + \frac{3}{4}\vartheta^2\rho.\end{aligned}$$

Using also,

$$e_4(\underline{\kappa}) + \frac{1}{2}\kappa\underline{\kappa} - 2\underline{\omega}\underline{\kappa} = 2(e_\theta(\underline{\eta}) + e_\theta(\Phi)\underline{\eta}) + 2\rho - \frac{1}{2}\vartheta\underline{\vartheta} + 2(\underline{\xi}\underline{\xi} + \underline{\eta}^2)$$

and the identities, $2\chi = \kappa + \vartheta$, as well as $2\underline{\chi} = \underline{\kappa} + \underline{\vartheta}$, we finally obtain

$$\square_{\mathbf{g}}\alpha = -4\underline{\omega}e_4(\alpha) + (2\kappa + 4\omega)e_3(\alpha) + V\alpha + \text{Err}[\square\alpha]$$

as desired.

We write schematically the error term,

$$\begin{aligned} \text{Err}[\square\alpha] &= \frac{1}{2}\vartheta e_3(\alpha) + \frac{3}{4}\vartheta^2\rho + e_\theta(\Phi)\vartheta\beta - \frac{1}{2}\kappa(\zeta + 4\underline{\eta})\beta - (\zeta + \underline{\eta})e_4(\beta) - \xi e_3(\beta) \\ &+ e_\theta(\Phi)(2\zeta + \underline{\eta})\alpha + \beta^2 + e_4(\Phi)\underline{\eta}\beta + e_3(\Phi)\xi\beta - (\zeta + 4\underline{\eta})e_4(\beta) - (e_4(\zeta) + 4e_4(\underline{\eta}))\beta \\ &- 2(\kappa + \omega)(\zeta + 4\underline{\eta})\beta + 2e_\theta(\kappa + \omega)\beta - e_\theta((2\zeta + \underline{\eta})\alpha) - 3\xi e_\theta(\rho) + 2\underline{\eta}e_\theta(\alpha) \\ &+ \frac{3}{2}\vartheta(e_\theta(\beta) + e_\theta(\Phi)\beta) + 3\rho(\underline{\eta} + \eta + 2\zeta)\xi + (e_\theta(\underline{\eta}) + e_\theta(\Phi)\underline{\eta})\alpha + \frac{1}{4}\underline{\kappa}\vartheta\alpha - 2\underline{\omega}\vartheta\alpha \\ &- \frac{1}{2}\vartheta\underline{\vartheta}\alpha + \xi\underline{\xi}\alpha + \underline{\eta}^2\alpha + \frac{3}{2}\vartheta\zeta\beta + 3\vartheta(\underline{\eta}\beta + \xi\beta) - \frac{1}{2}\vartheta(\zeta + 4\underline{\eta})\beta \end{aligned}$$

as follows,

$$\begin{aligned} \text{Err}[\square_{\mathbf{g}}\alpha] &= \left(\frac{1}{r}\Gamma_g + \frac{1}{r}\vartheta\Gamma_g\right)\alpha + \Gamma_g e_3(\alpha) + \frac{1}{r}\Gamma_g\vartheta\alpha \\ &+ \left(\beta + \frac{1}{r}\Gamma_g + \frac{1}{r}\vartheta\Gamma_g\right)\beta + \frac{1}{r}\Gamma_g\vartheta(\beta) + \Gamma_g e_3(\beta) \\ &+ (\Gamma_g)^2\rho + \frac{1}{r}\Gamma_g\vartheta(\rho) \\ &= \Gamma_g e_3(\alpha, \beta) + r^{-1}\Gamma_g^{\leq 1} \cdot \vartheta^{\leq 1}(\alpha, \beta, \rho) + \beta^2 + \Gamma_g^2\rho. \end{aligned}$$

This concludes the proof of Proposition 2.4.6. \square

A.14 Proof of Theorem 2.4.7

Recall the symbolic notation used in the statement of the theorem.

$$\begin{aligned} \Gamma_g &= \{\vartheta, \eta, \underline{\eta}, \zeta, A\}, & \Gamma_b &= \{\underline{\vartheta}, \underline{\xi}, \underline{A}\}, \\ \vartheta\Gamma_g &= \{\vartheta\vartheta, re_\theta(\kappa), \vartheta\eta, \vartheta\underline{\eta}, \vartheta\zeta, \vartheta A\}, & \vartheta\Gamma_b &= \{\vartheta\underline{\vartheta}, e_\theta(\underline{\kappa}), \vartheta\underline{\xi}, \vartheta\underline{A}\}, \end{aligned}$$

where $A = \frac{2}{r}e_4(r) - \kappa$, $\underline{A} = \frac{2}{r}e_3(r) - \underline{\kappa}$. We also denote, for $s \geq 2$,

$$\vartheta^s\Gamma_g = \vartheta^{s-1}\vartheta\Gamma_g, \quad \vartheta^s\Gamma_b = \vartheta^{s-1}\vartheta\Gamma_b,$$

for higher derivatives with respect to $\mathfrak{d} = (e_3, re_4, \check{\vartheta})$ (see definition 2.1.37 for the notation $\check{\vartheta}$ and $\check{\vartheta}^s$).

We also recall Remark 2.3.9.

Remark A.14.1. *According to the main bootstrap assumptions BA-E, BA-D (see section 3.4.1.) the terms Γ_b behave worse in powers of r than the terms in Γ_g . Thus, in the symbolic expressions below, we replace the terms of the form $\Gamma_g + \Gamma_b$ by Γ_b . We also replace $r^{-1}\Gamma_b$ by Γ_g . We will denote l.o.t. all cubic and higher error terms in $\check{\Gamma}, \check{R}$. We also include in l.o.t. terms which decay faster in powers of r than those taking into account by the main quadratic terms.*

Recall that

$$\mathfrak{q} = r^4 Q(\alpha), \quad (\text{A.14.1})$$

where Q is the operator

$$Q := e_3 e_3 + (2\underline{\kappa} - 6\underline{\omega})e_3 + W, \quad W := -4e_3(\underline{\omega}) + 8\underline{\omega}^2 - 8\underline{\omega}\underline{\kappa} + \frac{1}{2}\underline{\kappa}^2. \quad (\text{A.14.2})$$

Lemma A.14.2. *The quantity \mathfrak{q} is fully invariant with respect to the conformal frame transformations*

$$e'_3 = \lambda^{-1}e_3, \quad e_4 = \lambda e_4, \quad e'_\theta = e_\theta.$$

Proof. The proof is an immediate consequence of Definition A.14.3 and Lemmas A.14.5, A.14.4 below. \square

We recall that under the above mentioned frame transformation we have

$$\begin{aligned} \alpha' &= \lambda^2 \alpha, \quad \beta' = \lambda \beta, \quad \rho' = \rho, \quad \underline{\kappa}' = \lambda^{-1} \underline{\kappa}, \quad \kappa' = \lambda \kappa, \quad \eta' = \eta, \quad \underline{\eta}' = \underline{\eta}, \\ \underline{\omega}' &= \lambda^{-1} \left(\underline{\omega} + \frac{1}{2} e_3(\log \lambda) \right), \quad \omega' = \lambda \left(\omega - \frac{1}{2} e_4(\log \lambda) \right), \quad \zeta' = \zeta - e_\theta(\log \lambda). \end{aligned}$$

Definition A.14.3. *We say that a reduced tensor is conformal invariant of type⁴ a , i.e. a -conformal invariant, if under the conformal change of frames $e'_3 = \lambda^{-1}e_3, e'_4 = \lambda e_4$ it transforms by*

$$f' = \lambda^a f.$$

⁴Note that for a given Ricci or curvature coefficient a coincides with the signature of the component.

Lemma A.14.4. *Let f be an a -conformal invariant tensor.*

1. *The tensor*

$$\nabla_3 f := e_3 f - 2a\underline{\omega} f \quad (\text{A.14.3})$$

is $a - 1$ conformal invariant.

2. *The tensor*

$$\nabla_4 f := e_4 f + 2a\underline{\omega} f \quad (\text{A.14.4})$$

is $a + 1$ conformal invariant.

3. *The tensor,*

$$\nabla_A^{(c)} f = \nabla_A f + \alpha \zeta_A f \quad (\text{A.14.5})$$

is a -conformal invariant.

Proof. Immediate verification. □

Lemma A.14.5. *We have*

$$Q(\alpha) = \nabla_3(\nabla_3\alpha) + 2\underline{\kappa}\nabla_3\alpha + \frac{1}{2}\underline{\kappa}^2\alpha.$$

Proof. We have,

$$\begin{aligned} \nabla_3(\nabla_3\alpha) &= \nabla_3(e_3\alpha - 4\underline{\omega}\alpha) = e_3(e_3\alpha - 4\underline{\omega}\alpha) - 2\underline{\omega}(e_3\alpha - 4\underline{\omega}\alpha) \\ &= e_3e_3\alpha - 4e_3\underline{\omega}\alpha - 4\underline{\omega}e_3\alpha - 2\underline{\omega}e_3\alpha + 8\underline{\omega}^2\alpha. \end{aligned}$$

Hence,

$$\begin{aligned} Q(\alpha) &= \nabla_3(\nabla_3\alpha) + 2\underline{\kappa}\nabla_3\alpha + \frac{1}{2}\underline{\kappa}^2\alpha \\ &= e_3e_3\alpha - 4e_3\underline{\omega}\alpha - 4\underline{\omega}e_3\alpha - 2\underline{\omega}e_3\alpha + 8\underline{\omega}^2\alpha + 2\underline{\kappa}(e_3\alpha - 4\underline{\omega}\alpha) + \frac{1}{2}\underline{\kappa}^2\alpha \\ &= e_3e_3\alpha + (2\underline{\kappa} - 6\underline{\omega})e_3\alpha + \left(-4e_3\underline{\omega} + 8\underline{\omega}^2 - 8\underline{\kappa}\underline{\omega} + \frac{1}{2}\underline{\kappa}^2\right)\alpha \end{aligned}$$

as stated. □

Remark A.14.6. Using the definitions of ∇_3, ∇_4 the null structure equations for $\kappa, \underline{\kappa}$ take the form,

$$\begin{aligned}\nabla_3 \underline{\kappa} + \frac{1}{2} \underline{\kappa}^2 &= 2 \not\Delta_1 \underline{\xi} + \Gamma_b \cdot \Gamma_b = r^{-1} \not\partial + l.o.t., \\ \nabla_4 \underline{\kappa} + \frac{1}{2} \kappa \underline{\kappa} &= 2 \not\Delta_1 \underline{\eta} + 2\rho + \Gamma_g \cdot \Gamma_b = 2\rho + r^{-1} \not\partial \Gamma_g, \\ \nabla_3 \kappa + \frac{1}{2} \underline{\kappa} \kappa &= 2 \not\Delta_1 \eta + 2\rho + \Gamma_g \cdot \Gamma_b = 2\rho + r^{-1} \not\partial \Gamma_g, \\ \nabla_4 \kappa + \frac{1}{2} \kappa^2 &= 2 \not\Delta_1 \xi + \Gamma_g \cdot \Gamma_g = r^{-1} \not\partial \Gamma_g.\end{aligned}\tag{A.14.6}$$

Also, since ρ is 0-conformal

$$\nabla_3 \rho + \frac{3}{2} \underline{\kappa} \rho = \not\Delta_1 \underline{\beta} + \Gamma_g \cdot \Gamma_b = r^{-1} \not\partial \Gamma_g.\tag{A.14.7}$$

Definition A.14.7. Given f an a -conformal S -tangent tensor we define its a -conformal Laplacian to be

$${}^{(c)}\not\Delta f = {}^{(c)}\nabla_A {}^{(c)}\nabla^A f.$$

Lemma A.14.8. The following formula holds true for a 2-conformal tensor f

$${}^{(c)}\not\Delta f = \not\Delta_2 f + 4\zeta \nabla f + 2(\operatorname{div} \zeta + 2|\zeta|^2) f.$$

In particular we have,

$${}^{(c)}\not\Delta f = \not\Delta_2 f + r^{-1} \not\partial^{\leq 1}(\Gamma_g \cdot f).$$

Proof. Immediate verification. □

The goal of this section is to prove Theorem 2.4.7 which we recall below for the convenience of the reader.

Theorem A.14.9. The invariant scalar quantity \mathfrak{q} defined in (2.3.10) verifies the equation,

$$\square_2 \mathfrak{q} + \kappa \underline{\kappa} \mathfrak{q} = \operatorname{Err}[\square_2 \mathfrak{q}]\tag{A.14.8}$$

where, schematically,

$$\operatorname{Err}[\square_2 \mathfrak{q}] := r^2 \mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta)) + e_3 \left(r^3 \mathfrak{d}^{\leq 2}(\Gamma_g \cdot (\alpha, \beta)) \right) + \mathfrak{d}^{\leq 1}(\Gamma_g \cdot \mathfrak{q}) + l.o.t.\tag{A.14.9}$$

Definition A.14.10. *Given a quadratic or higher order E we say the following*

1. $E \in \text{Good}$ if $r^4 E$ can be expressed in the form (2.4.8).
2. $E \in \text{Good}_1$ if after applying $r^4 e_3$ or r^3 it can be expressed in the form (2.4.8).
3. $E \in \text{Good}_2$ if after applying $r^4 e_3 e_3, r^4 e_3$ or r^3 it can be expressed in the form (2.4.8).

In view of the definition we note that,

$$(e_3 + r^{-1})\text{Good}_1 = \text{Good}, \quad Q\text{Good}_2 = \text{Good}.$$

To prove the theorem we have to check that $\text{Err}[\square_2 \mathbf{q}] = r^4 \text{Good}$.

A.14.1 The Teukolsky equation for α

We recall below Proposition 2.4.6.

Lemma A.14.11. *We have*

$$\begin{aligned} \square_2 \alpha &= -4\underline{\omega} e_4(\alpha) + (4\omega + 2\kappa) e_3(\alpha) + V\alpha + \text{Err}[\square_{\mathbf{g}} \alpha], \\ V &= -4\rho - 4e_4(\underline{\omega}) - 8\omega \underline{\omega} + 2\omega \underline{\kappa} - 10\kappa \underline{\omega} + \frac{1}{2}\kappa \underline{\kappa}, \end{aligned}$$

where $\text{Err}[\square_{\mathbf{g}} \alpha]$ is given schematically by

$$\text{Err}(\square_{\mathbf{g}} \alpha) := \Gamma_g e_3(\alpha) + r^{-1} \mathfrak{d}^{\leq 1} \left((\eta, \Gamma_g)(\alpha, \beta) \right) + \xi(e_3(\beta), r^{-1} \mathfrak{d} \rho).$$

Remark A.14.12. *Since ξ vanishes for $r \geq 4m_0$, $\eta \in \Gamma_g$ and $e_3 \alpha = r^{-1} \mathfrak{d} \alpha$ we deduce,*

$$\text{Err}(\square_{\mathbf{g}} \alpha) \in \text{Good}_2.$$

Lemma A.14.13. *The Teukolsky equation for α can be written in the form,*

$$\mathcal{L}(\alpha) = \text{Good}_2 \tag{A.14.10}$$

where \mathcal{L} is the operator

$$\mathcal{L} \alpha = -\nabla_4 \nabla_3 \alpha + {}^{(c)}\Delta_2 \alpha - \frac{5}{2} \kappa \nabla_3 \alpha - \frac{1}{2} \underline{\kappa} \nabla_4 \alpha - \left(-4\rho + \frac{1}{2} \kappa \underline{\kappa} \right) \alpha. \tag{A.14.11}$$

We also note that, for a 2-conformal tensor f ,

$$\square_2 f = -\nabla_4 \nabla_3 f + {}^{(c)}\Delta_2 f - \frac{1}{2} \kappa \nabla_3 f - \frac{1}{2} \underline{\kappa} \nabla_4 f + r^{-1} \Gamma_g \cdot \mathfrak{d} f. \tag{A.14.12}$$

Proof. Recall that we have (see Definition 2.4.2)

$$\square_2\alpha = -e_4(e_3(\alpha)) + \underline{\Delta}_2\alpha - \frac{1}{2}\underline{\kappa}e_4(\alpha) + \left(-\frac{1}{2}\kappa + 2\omega\right)e_3(\alpha) + 2\underline{\eta}e_\theta(\alpha).$$

Therefore,

$$\begin{aligned} \mathcal{L}(\alpha) &= -e_4(e_3(\alpha)) + \underline{\Delta}_2\alpha - \frac{1}{2}\underline{\kappa}e_4(\alpha) + \left(-\frac{1}{2}\kappa + 2\omega\right)e_3(\alpha) + 2\underline{\eta}e_\theta(\alpha) \\ &\quad + 4\underline{\omega}e_4(\alpha) - (4\omega + 2\kappa)e_3(\alpha) - V\alpha \\ &= -e_4(e_3(\alpha)) + \underline{\Delta}_2\alpha - \left(\frac{1}{2}\underline{\kappa} - 4\underline{\omega}\right)e_4\alpha - \left(\frac{5}{2}\kappa + 2\omega\right)e_3\alpha + 2\underline{\eta}e_\theta(\alpha) - V\alpha \\ &= -e_4(e_3(\alpha)) + {}^{(c)}\underline{\Delta}_2\alpha - \left(\frac{1}{2}\underline{\kappa} - 4\underline{\omega}\right)e_4\alpha - \left(\frac{5}{2}\kappa + 2\omega\right)e_3\alpha - V\alpha + \text{Good}_2. \end{aligned}$$

On the other hand,

$$\begin{aligned} \nabla_4(\nabla_3(\alpha)) &= \nabla_4(e_3\alpha - 4\underline{\omega}\alpha) = e_4(e_3\alpha - 4\underline{\omega}\alpha) + 2\omega(e_3\alpha - 4\underline{\omega}\alpha) \\ &= e_4e_3\alpha - 4\underline{\omega}e_4\alpha - 4e_4\underline{\omega}\alpha + 2\omega e_3\alpha - 8\omega\underline{\omega}\alpha. \end{aligned}$$

Hence,

$$\begin{aligned} -\nabla_4\nabla_3\alpha - \frac{5}{2}\kappa\nabla_3\alpha - \frac{1}{2}\underline{\kappa}\nabla_4\alpha &= -e_4e_3\alpha + 4\underline{\omega}e_4\alpha + 4e_4\underline{\omega}\alpha - 2\omega e_3\alpha + 8\omega\underline{\omega}\alpha \\ &\quad - \frac{5}{2}\kappa(e_3\alpha - 4\underline{\omega}\alpha) - \frac{1}{2}\underline{\kappa}(e_4\alpha + 4\omega\alpha) \\ &= -e_4e_3\alpha - \frac{1}{2}(\underline{\kappa} - 4\underline{\omega})e_4\alpha - \left(\frac{5}{2}\kappa + 2\omega\right)e_3\alpha \\ &\quad + (4e_4\underline{\omega} + 8\omega\underline{\omega} + 10\kappa\underline{\omega} - 2\omega\underline{\kappa})\alpha. \end{aligned}$$

We deduce, with $V' = -4\rho + \frac{1}{2}\kappa\underline{\kappa}$,

$$\begin{aligned} -\nabla_4\nabla_3\alpha - \frac{5}{2}\kappa\nabla_3\alpha - \frac{1}{2}\underline{\kappa}\nabla_4\alpha - V'\alpha &= -e_4e_3\alpha - \frac{1}{2}(\underline{\kappa} - 4\underline{\omega})e_4\alpha - \left(\frac{5}{2}\kappa + 2\omega\right)e_3\alpha \\ &\quad + \left(4e_4\underline{\omega} + 8\omega\underline{\omega} + 10\kappa\underline{\omega} - 2\omega\underline{\kappa} + 4\rho - \frac{1}{2}\kappa\underline{\kappa}\right)\alpha \\ &= -e_4e_3\alpha - \frac{1}{2}(\underline{\kappa} - 4\underline{\omega})e_4\alpha - \left(\frac{5}{2}\kappa + 2\omega\right)e_3\alpha - V\alpha. \end{aligned}$$

Hence,

$$\mathcal{L}\alpha = -\nabla_4\nabla_3\alpha + {}^{(c)}\underline{\Delta}_2\alpha - \frac{5}{2}\kappa\nabla_3\alpha - \frac{1}{2}\underline{\kappa}\nabla_4\alpha - \left(-4\rho + \frac{1}{2}\kappa\underline{\kappa}\right)\alpha \in \text{Good}_2$$

as desired. The proof of the second part of the lemma follows in the same manner. \square

A.14.2 Commutation lemmas

The goal of the following lemmas is to calculate the commutator of Q with \mathcal{L} .

Lemma A.14.14. *Give f an a -conformal tensor we have,*

$$[\nabla_3, \nabla_4]f = 2a\rho f + r^{-1}\Gamma_g \mathfrak{P}^{\leq 1}f. \quad (\text{A.14.13})$$

Proof. We have

$$\begin{aligned} [\nabla_3, \nabla_4]f &= \nabla_3\nabla_4f - \nabla_4\nabla_3f \\ &= (e_3 - 2(a+1)\underline{\omega})(e_4f + 2a\omega f) - (e_4 + 2(a-1)\omega)(e_3f - 2a\underline{\omega}f) \\ &= e_3e_4f - 2(a+1)\underline{\omega}e_4f + 2ae_3(\omega f) - 4a(a+1)\underline{\omega}\omega f \\ &\quad - e_4e_3f - 2(a-1)\omega e_3f + 2ae_4(\underline{\omega}f) + 4a(a-1)\omega\underline{\omega} \\ &= [e_3, e_4]f - 2\underline{\omega}e_4f + 2\omega e_3(f) + 2a(e_3\omega + e_4\underline{\omega} - 4\underline{\omega}\omega)f. \end{aligned}$$

Recall that,

$$\begin{aligned} [e_3, e_4] &= -2\omega e_3 + 2\underline{\omega}e_4 + 2(\underline{\eta} - \eta)e_\theta, \\ e_3\omega + e_4\underline{\omega} - 4\underline{\omega}\omega &= \rho + \Gamma_g \cdot \Gamma_b. \end{aligned}$$

We deduce⁵,

$$[\nabla_3, \nabla_4]f = 2a\rho + r^{-1}\Gamma_g \mathfrak{P}^{\leq 1}f$$

as stated. □

Lemma A.14.15. *Assume f a -conformal and g is b -conformal. Then fg is $a+b$ -conformal and*

$$\begin{aligned} \nabla_3(fg) &= f\nabla_3g + g\nabla_3f, \\ \nabla_4(fg) &= f\nabla_4g + g\nabla_4f. \end{aligned}$$

Proof. Indeed

$$\nabla_3(fg) = e_3(fg) - 2(a+b)\underline{\omega}fg = fe_3g + ge_3f - 2(a+b)\underline{\omega}fg = f\nabla_3g + g\nabla_3f$$

as stated. □

⁵Recall that $\eta \in \Gamma_g$ in the frame we are using.

Lemma A.14.16. *We have,*

$$\begin{aligned} [Q, \nabla_3]\alpha &= \underline{\kappa}^2 \nabla_3 \alpha + \frac{1}{2} \underline{\kappa}^3 \alpha + \text{Good}_1, \\ [Q, \nabla_4]\alpha &= (2\rho + \underline{\kappa} \underline{\kappa}) \nabla_3 \alpha + \frac{1}{2} \underline{\kappa} \underline{\kappa}^2 \alpha + \text{Good}_1. \end{aligned} \tag{A.14.14}$$

Also,

$$\begin{aligned} [Q, \nabla_4 \nabla_3]\alpha &= (-2\rho + \underline{\kappa} \underline{\kappa}) \nabla_3 \nabla_3 \alpha + \underline{\kappa}^2 \nabla_4 \nabla_3 \alpha + \frac{1}{2} \underline{\kappa}^3 \nabla_4 \alpha \\ &\quad + \left(3\rho \underline{\kappa} - \frac{1}{2} \underline{\kappa} \underline{\kappa}^2\right) \nabla_3 \alpha + \frac{3}{2} \underline{\kappa}^2 \left(-\frac{1}{2} \underline{\kappa} \underline{\kappa} + 2\rho\right) \alpha + \text{Good}. \end{aligned} \tag{A.14.15}$$

Proof. We have⁶,

$$\begin{aligned} [Q, \nabla_3]\alpha &= \left(\nabla_3 \nabla_3 + 2\underline{\kappa} \nabla_3 + \frac{1}{2} \underline{\kappa}^2\right) \nabla_3 \alpha - \nabla_3 \left(\left(\nabla_3 \nabla_3 + 2\underline{\kappa} \nabla_3 + \frac{1}{2} \underline{\kappa}^2\right) \alpha\right) \\ &= -2\nabla_3(\underline{\kappa}) \nabla_3 \alpha - \underline{\kappa}(\nabla_3 \underline{\kappa}) \alpha \\ &= -2\left(-\frac{1}{2} \underline{\kappa}^2 + r^{-1} \Gamma_b\right) \nabla_3 \alpha - \underline{\kappa} \left(-\frac{1}{2} \underline{\kappa}^2 + r^{-1} \Gamma_b\right) \alpha \\ &= \underline{\kappa}^2 \nabla_3 \alpha + \frac{1}{2} \underline{\kappa}^3 \alpha + r^{-1} \Gamma_g \mathfrak{d}^{\leq 1} \alpha \end{aligned}$$

and,

$$\begin{aligned} [Q, \nabla_4]\alpha &= \left(\nabla_3 \nabla_3 + 2\underline{\kappa} \nabla_3 + \frac{1}{2} \underline{\kappa}^2\right) \nabla_4 \alpha - \nabla_4 \left(\left(\nabla_3 \nabla_3 + 2\underline{\kappa} \nabla_3 + \frac{1}{2} \underline{\kappa}^2\right) \alpha\right) \\ &= (\nabla_3 \nabla_3 \nabla_4 - \nabla_4 \nabla_3 \nabla_3) \alpha + 2\underline{\kappa} (\nabla_3 \nabla_4 - \nabla_4 \nabla_3) \alpha - 2\nabla_4 \underline{\kappa} \nabla_3 \alpha - \underline{\kappa} (\nabla_4 \underline{\kappa}) \alpha \\ &= \nabla_3([\nabla_3, \nabla_4] \alpha) + [\nabla_3, \nabla_4] \nabla_3 \alpha + 2\underline{\kappa} [\nabla_3, \nabla_4] \alpha - 2\nabla_4 \underline{\kappa} \nabla_3 \alpha - \underline{\kappa} (\nabla_4 \underline{\kappa}) \alpha. \end{aligned}$$

In view of Lemma A.14.14 we have,

$$\begin{aligned} [\nabla_3, \nabla_4] \alpha &= 4\rho \alpha + r^{-1} \Gamma_g \cdot \mathfrak{d}^{\leq 1} \alpha, \\ [\nabla_3, \nabla_4] \nabla_3 \alpha &= 2\rho \nabla_3 \alpha + r^{-1} \Gamma_g \cdot \mathfrak{d}^{\leq 1} \nabla_3 \alpha. \end{aligned}$$

Hence,

$$\begin{aligned} [Q, \nabla_4]\alpha &= \nabla_3(4\rho \alpha + r^{-1} \Gamma_g \mathfrak{d}^{\leq 1} \alpha) + (2\rho + r^{-1} \Gamma_g \mathfrak{d}^{\leq 1}) \nabla_3 \alpha + 2\underline{\kappa} (4\rho \alpha + r^{-1} \Gamma_g \mathfrak{d}^{\leq 1} \alpha) \\ &\quad - 2\nabla_4 \underline{\kappa} \nabla_3 \alpha - \underline{\kappa} (\nabla_4 \underline{\kappa}) \alpha \\ &= (6\rho - 2\nabla_4 \underline{\kappa}) \nabla_3 \alpha + (4\nabla_3 \rho + 8\underline{\kappa} \rho - \underline{\kappa} \nabla_4 \underline{\kappa}) \alpha + \text{Good}_1. \end{aligned}$$

⁶Recall that $r^{-1} \Gamma_b = \Gamma_g$.

We now note, using the equations for $\nabla_4\rho$ and $\nabla_4\underline{\kappa}$,

$$\begin{aligned} 4\nabla_3\rho + 8\underline{\kappa}\rho - \underline{\kappa}\nabla_4\underline{\kappa} &= 4\left(-\frac{3}{2}\underline{\kappa}\rho + r^{-1}\wp\Gamma_g\right) + 8\underline{\kappa}\rho - \underline{\kappa}\left(-\frac{1}{2}\underline{\kappa}\underline{\kappa} + 2\rho + r^{-1}\wp\Gamma_g\right) \\ &= \frac{1}{2}\underline{\kappa}\underline{\kappa}^2 + r^{-1}\wp\Gamma_g \\ 6\rho - 2\nabla_4\underline{\kappa} &= 6\rho - 2\left(-\frac{1}{2}\underline{\kappa}\underline{\kappa} + 2\rho + r^{-1}\wp\Gamma_g\right) \\ &= 2\rho + \underline{\kappa}\underline{\kappa} + r^{-1}\wp\Gamma_g. \end{aligned}$$

Hence,

$$[Q, \nabla_4]\alpha = (2\rho + \underline{\kappa}\underline{\kappa})\nabla_3\alpha + \frac{1}{2}\underline{\kappa}\underline{\kappa}^2\alpha + \text{Good}_1$$

as stated.

Also,

$$[Q, \nabla_4\nabla_3]\alpha = [Q, \nabla_4]\nabla_3\alpha + \nabla_4([Q, \nabla_3]\alpha). \quad (\text{A.14.16})$$

We first calculate, as above, for $f = \nabla_3\alpha$

$$\begin{aligned} [Q, \nabla_4]f &= \left(\nabla_3\nabla_3 + 2\underline{\kappa}\nabla_3 + \frac{1}{2}\underline{\kappa}^2\right)\nabla_4f - \nabla_4\left(\left(\nabla_3\nabla_3 + 2\underline{\kappa}\nabla_3 + \frac{1}{2}\underline{\kappa}^2\right)f\right) \\ &= (\nabla_3\nabla_3\nabla_4 - \nabla_4\nabla_3\nabla_3)f + 2\underline{\kappa}(\nabla_3\nabla_4 - \nabla_4\nabla_3)f - 2\nabla_4\underline{\kappa}\nabla_3\alpha - \underline{\kappa}(\nabla_4\underline{\kappa})f \\ &= \nabla_3([\nabla_3, \nabla_4]f) + [\nabla_3, \nabla_4]\nabla_3f + 2\underline{\kappa}[\nabla_3, \nabla_4]f - 2\nabla_4\underline{\kappa}\nabla_3f - \underline{\kappa}(\nabla_4\underline{\kappa})f. \end{aligned}$$

In view of Lemma A.14.14, since $f = \nabla_3\alpha$ is 1-conformal and ∇_3f is 0-conformal, we have

$$\begin{aligned} [\nabla_3, \nabla_4]f &= 2\rho f + r^{-1}\Gamma_g\wp^{\leq 1}f, \\ [\nabla_3, \nabla_4]\nabla_3f &= r^{-1}\Gamma_g\wp^{\leq 1}\nabla_3f. \end{aligned}$$

Hence

$$\begin{aligned} [Q, \nabla_4]f &= \nabla_3(2\rho f + r^{-1}\Gamma_g\wp^{\leq 1}f) + (r^{-1}\Gamma_g\wp^{\leq 1})\nabla_3f + 2\underline{\kappa}(2\rho f + r^{-1}\Gamma_g\wp^{\leq 1}f) \\ &\quad - 2\nabla_4\underline{\kappa}\nabla_3f - \underline{\kappa}(\nabla_4\underline{\kappa})f \\ &= (2\rho - 2\nabla_4\underline{\kappa})\nabla_3f + (2\nabla_3\rho + 4\underline{\kappa}\rho - \underline{\kappa}\nabla_4\underline{\kappa})f \\ &\quad + r^{-1}\Gamma_g\wp^{\leq 1}\nabla_3f + r^{-2}\Gamma_g\wp^{\leq 1}f. \end{aligned}$$

Therefore,

$$[Q, \nabla_4] \nabla_3 \alpha = (2\rho - 2\nabla_4 \underline{\kappa}) \nabla_3 \nabla_3 \alpha + (2\nabla_3 \rho + 4\underline{\kappa} \rho - \underline{\kappa} \nabla_4 \underline{\kappa}) \nabla_3 \alpha + r^{-2} \Gamma_g \mathfrak{d}^{\leq 2} \alpha.$$

As above,

$$\begin{aligned} 2\rho - 2\nabla_4 \underline{\kappa} &= 2\rho - 2 \left(-\frac{1}{2} \underline{\kappa} \underline{\kappa} + 2\rho + r^{-1} \Gamma_g \right) = -2\rho + \underline{\kappa} \underline{\kappa} + r^{-1} \Gamma_g, \\ 2\nabla_3 \rho + 4\underline{\kappa} \rho - \underline{\kappa} \nabla_4 \underline{\kappa} &= 2 \left(-\frac{3}{2} \underline{\kappa} \rho + r^{-1} \Gamma_g \right) + 4\underline{\kappa} \rho - \underline{\kappa} \left(-\frac{1}{2} \underline{\kappa} \underline{\kappa} + 2\rho + r^{-1} \Gamma_g \right) \\ &= \frac{1}{2} \underline{\kappa} \underline{\kappa}^2 - \rho \underline{\kappa} + r^{-1} \Gamma_g. \end{aligned}$$

Hence, since $r^{-1} \Gamma_g (\nabla_3 \nabla_3 \alpha, \nabla_3 \alpha) = r^{-2} \Gamma_g \cdot \mathfrak{d}^{\leq 2} \alpha = \text{Good}$,

$$[Q, \nabla_4] \nabla_3 \alpha = (-2\rho + \underline{\kappa} \underline{\kappa}) \nabla_3 \nabla_3 \alpha + \left(\frac{1}{2} \underline{\kappa} \underline{\kappa}^2 - \rho \underline{\kappa} \right) \nabla_3 \alpha + \text{Good}. \quad (\text{A.14.17})$$

We deduce,

$$\begin{aligned} [Q, \nabla_4 \nabla_3] \alpha &= [Q, \nabla_4] \nabla_3 \alpha + \nabla_4 ([Q, \nabla_3] \alpha) \\ &= (-2\rho + \underline{\kappa} \underline{\kappa}) \nabla_3 \nabla_3 \alpha + \left(\frac{1}{2} \underline{\kappa} \underline{\kappa}^2 - \rho \underline{\kappa} \right) \nabla_3 \alpha \\ &\quad + \nabla_4 \left(\underline{\kappa}^2 \nabla_3 \alpha + \frac{1}{2} \underline{\kappa}^3 \alpha + \text{Good}_1 \right) + \text{Good} \\ &= (-2\rho + \underline{\kappa} \underline{\kappa}) \nabla_3 \nabla_3 \alpha + \underline{\kappa}^2 \nabla_4 \nabla_3 \alpha + \frac{1}{2} \underline{\kappa}^3 \nabla_4 \alpha \\ &\quad + \left(\nabla_4 (\underline{\kappa}^2) + \frac{1}{2} \underline{\kappa} \underline{\kappa}^2 - \rho \underline{\kappa} \right) \nabla_3 \alpha + \frac{3}{2} \underline{\kappa}^2 \nabla_4 \underline{\kappa} \alpha + \text{Good}. \end{aligned}$$

Note that

$$\begin{aligned} \nabla_4 (\underline{\kappa}^2) + \frac{1}{2} \underline{\kappa} \underline{\kappa}^2 - \rho \underline{\kappa} &= 2\underline{\kappa} \left(-\frac{1}{2} \underline{\kappa} \underline{\kappa} + 2\rho + r^{-1} \mathfrak{d} \Gamma_g \right) + \frac{1}{2} \underline{\kappa} \underline{\kappa}^2 - \rho \underline{\kappa} \\ &= 3\rho \underline{\kappa} - \frac{1}{2} \underline{\kappa} \underline{\kappa}^2 + r^{-2} \mathfrak{d} \Gamma_g, \\ \frac{3}{2} \underline{\kappa}^2 \nabla_4 \underline{\kappa} &= \frac{3}{2} \underline{\kappa}^2 \left(-\frac{1}{2} \underline{\kappa} \underline{\kappa} + 2\rho + r^{-1} \mathfrak{d} \Gamma_g \right). \end{aligned}$$

Hence,

$$\begin{aligned} [Q, \nabla_4 \nabla_3] \alpha &= (-2\rho + \underline{\kappa} \underline{\kappa}) \nabla_3 \nabla_3 \alpha + \underline{\kappa}^2 \nabla_4 \nabla_3 \alpha + \frac{1}{2} \underline{\kappa}^3 \nabla_4 \alpha \\ &\quad + \left(3\rho \underline{\kappa} - \frac{1}{2} \underline{\kappa} \underline{\kappa}^2 \right) \nabla_3 \alpha + \frac{3}{2} \underline{\kappa}^2 \left(-\frac{1}{2} \underline{\kappa} \underline{\kappa} + 2\rho \right) \alpha + \text{Good} \end{aligned}$$

as stated. \square

Lemma A.14.17. *Given f a 2-conformal tensor in \mathfrak{s}_2 we have*

$$[\nabla_3, {}^{(c)}\Delta]f = -\underline{\kappa} {}^{(c)}\Delta f + r^{-1}\mathfrak{d}^{\leq 2}(\Gamma_b \cdot f).$$

Proof. Recall that for a 2-conformal spacetime tensor f we have

$${}^{(c)}\Delta f = \Delta f + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_g \cdot f).$$

Hence,

$$[\nabla_3, {}^{(c)}\Delta]f = [\nabla_3, \Delta]f + \nabla_3(r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_g \cdot f)) + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_g \cdot \nabla_3 f).$$

On the other hand, since $\nabla \underline{\omega} = r^{-1}\mathfrak{d}\Gamma_b$, $\nabla^2 \underline{\omega} = r^{-2}\mathfrak{d}^2\Gamma_b$,

$$[\nabla_3, \Delta]f = [e_3 - 4\underline{\omega}, \Delta]f = [e_3, \Delta]f + r^{-2}\mathfrak{d}^{\leq 2}(\Gamma_b \cdot f).$$

We deduce,

$$[\nabla_3, {}^{(c)}\Delta]f = [e_3, \Delta]f + r^{-2}\mathfrak{d}^{\leq 2}(\Gamma_b \cdot f) + e_3(r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_g \cdot f)).$$

In the reduced form, for an \mathfrak{s}_2 tensor f ,

$$[\nabla_3, {}^{(c)}\Delta]f = [e_3, \Delta_2]f + r^{-2}\mathfrak{d}^{\leq 2}(\Gamma_b \cdot f) + e_3(r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_g \cdot f)).$$

We now recall that $\Delta_2 = -\mathfrak{d}_2^* \mathfrak{d}_2 + 2K$. Hence, applying the commutation Lemma⁷ 2.1.51,

$$\begin{aligned} [\Delta_2, e_3]f &= [-\mathfrak{d}_2^* \mathfrak{d}_2 + 2K, e_3]f = -\mathfrak{d}_2^*[\mathfrak{d}_2, e_3]f - [\mathfrak{d}_2^*, e_3]\mathfrak{d}_2 f - 2e_3(K)f \\ &= -\mathfrak{d}_2^* \left(\frac{1}{2}\underline{\kappa} \mathfrak{d}_2 + \underline{Com}_2(f) \right) - \left(\frac{1}{2}\underline{\kappa} \mathfrak{d}_2^* + \underline{Com}_2^*(\mathfrak{d}_2 f) \right) - 2e_3(K) \\ &= -\underline{\kappa} \mathfrak{d}_2^* \mathfrak{d}_2 f - 2e_3(K)f + e_\theta(\underline{\kappa}) \mathfrak{d}_2 f - \mathfrak{d}_2^*(\underline{Com}_2(f)) - \underline{Com}_2^*(\mathfrak{d}_2 f) \\ &= -\underline{\kappa} \mathfrak{d}_2^* \mathfrak{d}_2 f - 2e_3(K)f + r^{-2}\mathfrak{d}^{\leq 2}(\Gamma_b \cdot f) + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_g \cdot e_3 f) \\ &= \underline{\kappa} \Delta_2 f - 2(e_3 K + \underline{\kappa} K)f + r^{-2}\mathfrak{d}^{\leq 2}(\Gamma_b \cdot f) + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_g \cdot e_3 f). \end{aligned}$$

⁷Recall that we have

$$\begin{aligned} \underline{Com}_2(f) &= -\frac{1}{2}\mathfrak{d} \mathfrak{d}_3^* f + (\zeta - \eta)e_3 f - 2\eta e_3 \Phi f - \underline{\xi}(e_4 f + \kappa e_4(\Phi)f) - 2\underline{\beta} f, \\ \underline{Com}_2^*(f) &= -\frac{1}{2}\mathfrak{d} \mathfrak{d}_1 f - (\zeta - \eta)e_3 f - \eta e_3 \Phi f + \underline{\xi}(e_4 f - e_4(\Phi)f) - \underline{\beta} f. \end{aligned}$$

Note that, ignoring the quadratic terms,

$$\begin{aligned}
e_3 K + \underline{\kappa} K &= -e_3 \left(\rho + \frac{1}{4} \underline{\kappa} \underline{\kappa} \right) - \underline{\kappa} \left(\rho + \frac{1}{4} \underline{\kappa} \underline{\kappa} \right) \\
&= -e_3 \rho - \underline{\kappa} \rho - \frac{1}{4} \left(e_3(\underline{\kappa} \underline{\kappa}) + \underline{\kappa} \underline{\kappa}^2 \right) \\
&= \frac{1}{2} \underline{\kappa} \rho - \underline{\phi}_1 \underline{\beta} - \frac{1}{4} \left\{ \underline{\kappa} \left(-\frac{1}{2} \underline{\kappa}^2 - 2 \underline{\omega} \underline{\kappa} \right) + \underline{\kappa} \left(-\frac{1}{2} \underline{\kappa} \underline{\kappa} + 2 \underline{\omega} \underline{\kappa} + 2 \underline{\phi}_1 \eta + 2 \rho \right) + \underline{\kappa} \underline{\kappa}^2 \right\} \\
&= -\underline{\phi}_1 \underline{\beta} - \frac{1}{2} \underline{\kappa} \underline{\phi}_1 \eta.
\end{aligned}$$

We deduce,

$$[e_3, \underline{\Delta}_2] = -[\underline{\Delta}_2, e_3] f = -\underline{\kappa} \underline{\Delta}_2 f + r^{-1} \mathfrak{d}^{\leq 2}(\Gamma_b \cdot f).$$

Consequently,

$$[\nabla_3, {}^{(c)}\underline{\Delta}] f = -\underline{\kappa} {}^{(c)}\underline{\Delta} f + r^{-1} \mathfrak{d}^{\leq 2}(\Gamma_b \cdot f)$$

as stated. \square

Lemma A.14.18. *We have,*

$$[Q, {}^{(c)}\underline{\Delta}] \alpha = -2 \underline{\kappa} \nabla_3 {}^{(c)}\underline{\Delta} \alpha - \frac{5}{2} \underline{\kappa}^2 {}^{(c)}\underline{\Delta} \alpha + \text{Good}. \quad (\text{A.14.18})$$

Proof. We have

$$\begin{aligned}
[Q, {}^{(c)}\underline{\Delta}_2] \alpha &= \left[\nabla_3 \nabla_3 + 2 \underline{\kappa} \nabla_3 + \frac{1}{2} \underline{\kappa}^2 \right] {}^{(c)}\underline{\Delta} \alpha - {}^{(c)}\underline{\Delta} \left[\nabla_3 \nabla_3 \alpha + 2 \underline{\kappa} \nabla_3 \alpha + \frac{1}{2} \underline{\kappa}^2 \alpha \right] \\
&= \nabla_3 [\nabla_3, {}^{(c)}\underline{\Delta}] \alpha + [\nabla_3, {}^{(c)}\underline{\Delta}] e_3 \alpha + [2 \underline{\kappa} \nabla_3, {}^{(c)}\underline{\Delta}] \alpha + \left[\frac{1}{2} \underline{\kappa}^2, {}^{(c)}\underline{\Delta}_2 \right] \alpha.
\end{aligned}$$

Note that

$$\begin{aligned}
[2 \underline{\kappa} \nabla_3, {}^{(c)}\underline{\Delta}] \alpha &= 2 \underline{\kappa} [\nabla_3, {}^{(c)}\underline{\Delta}] \alpha + \text{Good}, \\
\left[\frac{1}{2} \underline{\kappa}^2, {}^{(c)}\underline{\Delta}_2 \right] \alpha &= \text{Good}.
\end{aligned}$$

Hence, using the previous commutation Lemma,

$$\begin{aligned}
[Q, {}^{(c)}\underline{\Delta}_2] \alpha &= \nabla_3 [\nabla_3, {}^{(c)}\underline{\Delta}] \alpha + [\nabla_3, {}^{(c)}\underline{\Delta}] e_3 \alpha + 2 \underline{\kappa} [\nabla_3, {}^{(c)}\underline{\Delta}] \alpha + \text{Good} \\
&= \nabla_3 \left(-\underline{\kappa} {}^{(c)}\underline{\Delta} \alpha + r^{-1} \mathfrak{d}^{\leq 2}(\Gamma_b \cdot \alpha) \right) + \left(-\underline{\kappa} {}^{(c)}\underline{\Delta} \nabla_3 \alpha + r^{-1} \mathfrak{d}^{\leq 2}(\Gamma_b \cdot \nabla_3 \alpha) \right) \\
&+ 2 \underline{\kappa} \left(-\underline{\kappa} {}^{(c)}\underline{\Delta} \alpha + r^{-1} \mathfrak{d}^{\leq 2}(\Gamma_b \cdot \alpha) \right) + \text{Good} \\
&= -\underline{\kappa} (\nabla_3 {}^{(c)}\underline{\Delta} \alpha + {}^{(c)}\underline{\Delta} \nabla_3 \alpha) - (\nabla_3 \underline{\kappa} + 2 \underline{\kappa}^2) {}^{(c)}\underline{\Delta} \alpha + \text{Good} \\
&= -\underline{\kappa} (2 \nabla_3 {}^{(c)}\underline{\Delta} \alpha - [\nabla_3, {}^{(c)}\underline{\Delta}] \alpha) - (\nabla_3 \underline{\kappa} + 2 \underline{\kappa}^2) {}^{(c)}\underline{\Delta} \alpha + \text{Good} \\
&= -2 \underline{\kappa} \nabla_3 {}^{(c)}\underline{\Delta} \alpha - (\nabla_3 \underline{\kappa} + 3 \underline{\kappa}^2) {}^{(c)}\underline{\Delta} \alpha + \text{Good}.
\end{aligned}$$

Note that

$$(\nabla_{3\underline{\kappa}} + 3\underline{\kappa}^2)^{(c)}\Delta\alpha = \left(\frac{5}{2}\underline{\kappa}^2 + r^{-1}\wp\Gamma_b\right)^{(c)}\Delta\alpha = \frac{5}{2}\underline{\kappa}^2 + r^{-2}\wp\Gamma_g \cdot \wp^{\leq 2}\alpha.$$

Hence,

$$[Q, {}^{(c)}\Delta_2]\alpha = -2\underline{\kappa}\nabla_3 {}^{(c)}\Delta\alpha - \frac{5}{2}\underline{\kappa}^2 {}^{(c)}\Delta\alpha + \text{Good}$$

as stated. \square

Lemma A.14.19. *We have*

$$Q(fg) = Q(f)g + fQ(g) + 2\nabla_3 f \nabla_3 g - \frac{1}{2}\underline{\kappa}^2 fg.$$

Also,

$$\begin{aligned} [Q, fe_4]g &= Q(f)\nabla_4 g + f[Q, e_4]g + 2\nabla_3 f \nabla_3 \nabla_4 g - \frac{1}{2}\underline{\kappa}^2 f \nabla_4 g, \\ [Q, f\nabla_3]g &= Q(f)\nabla_3 g + f[Q, \nabla_3]g + 2\nabla_3 f \nabla_3 \nabla_3 g - \frac{1}{2}\underline{\kappa}^2 f \nabla_3 g. \end{aligned}$$

Proof. Recall that,

$$Q = \nabla_3 \nabla_3 + 2\underline{\kappa}\nabla_3 + \frac{1}{2}\underline{\kappa}^2.$$

Hence,

$$\begin{aligned} Q(fg) &= \left[\nabla_3 \nabla_3 + 2\underline{\kappa}\nabla_3 + \frac{1}{2}\underline{\kappa}^2 \right] (fg) \\ &= (\nabla_3 \nabla_3 f)g + f(\nabla_3 \nabla_3 g) + 2\nabla_3 f \nabla_3 g + 2\underline{\kappa}(\nabla_3 f g + f \nabla_3 g) + \frac{1}{2}\underline{\kappa}^2 fg \\ &= \left(\nabla_3 \nabla_3 f + 2\underline{\kappa}\nabla_3 f \right) g + 2\nabla_3 f \nabla_3 g + fQ(g) \\ &= Q(f)g + fQ(g) + 2\nabla_3 f \nabla_3 g - \frac{1}{2}\underline{\kappa}^2 fg. \end{aligned}$$

Also,

$$\begin{aligned} [Q, f\nabla_4]g &= Q(f\nabla_4 g) - f\nabla_4 Q(g) = Q(f)\nabla_4 g + fQ\nabla_4(g) + 2\nabla_3 f \nabla_3 \nabla_4 g - \frac{1}{2}\underline{\kappa}^2 f \nabla_4 g \\ &\quad - f\nabla_4 Q(g) \\ &= \left(Q(f) - \frac{1}{2}\underline{\kappa}f \right) \nabla_4 g + f[Q, \nabla_4]g + 2\nabla_3 f \nabla_3 \nabla_4 g. \end{aligned}$$

Similarly,

$$[Q, f\nabla_3]g = \left(Q(f) - \frac{1}{2}\underline{\kappa}^2 f \right) \nabla_3 g + f[Q, \nabla_3]g + 2\nabla_3 f \nabla_3 \nabla_3 g$$

as stated. \square

A.14.3 Main commutation

Proposition A.14.20. *The following identity holds true.*

$$[Q, \mathcal{L}]\alpha = -2\underline{\kappa}\nabla_4 Q(\alpha) + C_Q Q(\alpha) + \text{Good}, \quad (\text{A.14.19})$$

where,

$$C_Q = -8\rho - \frac{7}{2}\underline{\kappa}\underline{\kappa}.$$

Proof. In view of Lemma A.14.13, we have

$$\mathcal{L}\alpha = -\nabla_4\nabla_3\alpha + {}^{(c)}\Delta_2\alpha - \frac{5}{2}\underline{\kappa}\nabla_3\alpha - \frac{1}{2}\underline{\kappa}\nabla_4\alpha - \left(-4\rho + \frac{1}{2}\underline{\kappa}\underline{\kappa}\right)\alpha = \text{Good}_2.$$

Hence, we infer

$$\begin{aligned} [Q, \mathcal{L}]\alpha &= -[Q, \nabla_4\nabla_3]\alpha + [Q, \Delta_2]\alpha - \frac{1}{2}[Q, \underline{\kappa}\nabla_4]\alpha - \frac{5}{2}[Q, \underline{\kappa}\nabla_3]\alpha + \left[Q, 4\rho - \frac{1}{2}\underline{\kappa}\underline{\kappa}\right]\alpha \\ &= I + J + K + L + M \end{aligned} \quad (\text{A.14.20})$$

with I, J, K, L, M denoting each of the commutators on the left of (A.14.20).

Expression for I

In view of Lemma A.14.16 we have, for $I = -[Q, \nabla_4\nabla_3]\alpha$,

$$\begin{aligned} I &= (2\rho - \underline{\kappa}\underline{\kappa})\nabla_3\nabla_3\alpha - \underline{\kappa}^2\nabla_4\nabla_3\alpha - \frac{1}{2}\underline{\kappa}^3\nabla_4\alpha - \left(-\frac{1}{2}\underline{\kappa}\underline{\kappa}^2 + 3\rho\underline{\kappa}\right)\nabla_3\alpha \\ &\quad - \frac{3}{2}\underline{\kappa}^2\left(-\frac{1}{2}\underline{\kappa}\underline{\kappa} + 2\rho\right)\alpha + \text{Good}. \end{aligned} \quad (\text{A.14.21})$$

Expression for J

Using Lemma A.14.18,

$$J = [Q, {}^{(c)}\Delta]\alpha = -2\underline{\kappa}\nabla_3 {}^{(c)}\Delta\alpha - \frac{5}{2}\underline{\kappa}^2 {}^{(c)}\Delta\alpha.$$

Recalling the definition of \mathcal{L} and the fact that $\mathcal{L}\alpha = \text{Good}_1$ we write,

$$\mathbb{A}_2\alpha = \nabla_4\nabla_3\alpha + \frac{5}{2}\kappa\nabla_3\alpha + \frac{1}{2}\underline{\kappa}\nabla_4\alpha + \left(-4\rho + \frac{1}{2}\kappa\underline{\kappa}\right)\alpha + \text{Good}_1.$$

Hence,

$$\begin{aligned} J &= -2\underline{\kappa}\nabla_3\left(\nabla_4\nabla_3\alpha + \frac{5}{2}\kappa\nabla_3\alpha + \frac{1}{2}\underline{\kappa}\nabla_4\alpha + \left(-4\rho + \frac{1}{2}\kappa\underline{\kappa}\right)\alpha\right) \\ &\quad - \frac{5}{2}\underline{\kappa}^2\left(\nabla_4\nabla_3\alpha + \frac{5}{2}\kappa\nabla_3\alpha + \frac{1}{2}\underline{\kappa}\nabla_4\alpha + \left(-4\rho + \frac{1}{2}\kappa\underline{\kappa}\right)\alpha\right) \\ &= -2\underline{\kappa}\nabla_3\nabla_4\nabla_3\alpha - 5\underline{\kappa}\nabla_3\nabla_3\alpha - \underline{\kappa}^2\nabla_3\nabla_4\alpha - 2\underline{\kappa}\left(-4\rho + \frac{1}{2}\kappa\underline{\kappa}\right)\nabla_3\alpha \\ &\quad - 2\underline{\kappa}\left(\frac{5}{2}\nabla_3\kappa\nabla_3\alpha + \frac{1}{2}\nabla_3\underline{\kappa}\nabla_4\alpha + \nabla_3\left(-4\rho + \frac{1}{2}\kappa\underline{\kappa}\right)\alpha\right) \\ &\quad - \frac{5}{2}\underline{\kappa}^2\left(\nabla_4\nabla_3\alpha + \frac{5}{2}\kappa\nabla_3\alpha + \frac{1}{2}\underline{\kappa}\nabla_4\alpha + \left(-4\rho + \frac{1}{2}\kappa\underline{\kappa}\right)\alpha\right). \end{aligned}$$

According to Lemma A.14.14

$$\begin{aligned} \nabla_3\nabla_4\nabla_3\alpha &= \nabla_4\nabla_3\nabla_3\alpha + [\nabla_3, \nabla_4]\nabla_3\alpha = \nabla_4\nabla_3\nabla_3\alpha + 2\rho\nabla_3\alpha + r^{-1}\Gamma_g\mathfrak{O}^{\leq 1}\nabla_3\alpha \\ &= \nabla_4\nabla_3\nabla_3\alpha + 2\rho\nabla_3\alpha + \text{Good}, \\ \nabla_3\nabla_4\alpha &= \nabla_4\nabla_3\alpha + 4\rho\alpha + \text{Good}_1. \end{aligned}$$

We deduce, modulo Good error terms,

$$\begin{aligned} J &= -2\underline{\kappa}(\nabla_4\nabla_3\nabla_3\alpha + 2\rho\nabla_3\alpha) - 5\underline{\kappa}\nabla_3\nabla_3\alpha - \underline{\kappa}^2(\nabla_4\nabla_3\alpha + 4\rho\alpha) - 2\underline{\kappa}\left(-4\rho + \frac{1}{2}\kappa\underline{\kappa}\right)\nabla_3\alpha \\ &\quad - 5\underline{\kappa}\nabla_3\kappa\nabla_3\alpha - \underline{\kappa}\nabla_3\underline{\kappa}\nabla_4\alpha - 2\underline{\kappa}\nabla_3\left(-4\rho + \frac{1}{2}\kappa\underline{\kappa}\right)\alpha \\ &\quad - \frac{5}{2}\underline{\kappa}^2\left(\nabla_4\nabla_3\alpha + \frac{5}{2}\kappa\nabla_3\alpha + \frac{1}{2}\underline{\kappa}\nabla_4\alpha + \left(-4\rho + \frac{1}{2}\kappa\underline{\kappa}\right)\alpha\right). \end{aligned}$$

Grouping terms we rewrite in the form,

$$J = -2\underline{\kappa}\nabla_4\nabla_3\nabla_3\alpha - 5\underline{\kappa}\nabla_3\nabla_3\alpha + J_{43}\nabla_4\nabla_3\alpha + J_4\nabla_4\alpha + J_3\nabla_3\alpha + J_0\alpha.$$

We calculate the coefficients J_{43}, J_4, J_3, J_0 as follows.

$$\begin{aligned}
J_{43} &= -\underline{\kappa}^2 - \frac{5}{2}\underline{\kappa}^2 = -\frac{7}{2}\underline{\kappa}^2, \\
J_4 &= -\underline{\kappa}\nabla_3\underline{\kappa} - \frac{5}{4}\underline{\kappa}^3 = -\underline{\kappa}\left(-\frac{1}{2}\underline{\kappa}^2 + r^{-1}\wp\Gamma_b\right) - \frac{5}{4}\underline{\kappa}^3 = -\frac{3}{4}\underline{\kappa}^3 + r^{-2}\wp\Gamma_b, \\
J_3 &= -4\underline{\kappa}\rho - 2\underline{\kappa}\left(-4\rho + \frac{1}{2}\underline{\kappa}\underline{\kappa}\right) - 5\underline{\kappa}\nabla_3\underline{\kappa} - \frac{25}{4}\underline{\kappa}^3 \\
&= 4\underline{\kappa}\rho - \frac{29}{4}\underline{\kappa}^3 - 5\underline{\kappa}\left(-\frac{1}{2}\underline{\kappa}\underline{\kappa} + 2\rho + r^{-1}\wp\Gamma_g\right) \\
&= -6\underline{\kappa}\rho - \frac{19}{4}\underline{\kappa}^3 + r^{-2}\wp\Gamma_g, \\
J_0 &= -4\rho\underline{\kappa}^2 - 2\underline{\kappa}\nabla_3\left(-4\rho + \frac{1}{2}\underline{\kappa}\underline{\kappa}\right) - \frac{5}{2}\underline{\kappa}^2\left(-4\rho + \frac{1}{2}\underline{\kappa}\underline{\kappa}\right) \\
&= 6\rho\underline{\kappa}^2 - \frac{5}{4}\underline{\kappa}\underline{\kappa}^3 + 8\underline{\kappa}\nabla_3\rho - \underline{\kappa}(\underline{\kappa}\nabla_3\underline{\kappa} + \underline{\kappa}\nabla_3\underline{\kappa}) \\
&= 6\rho\underline{\kappa}^2 - \frac{5}{4}\underline{\kappa}\underline{\kappa}^3 + 8\underline{\kappa}\left(-\frac{3}{2}\underline{\kappa}\rho + r^{-1}\wp\Gamma_g\right) \\
&\quad - \underline{\kappa}\left(\underline{\kappa}\left(-\frac{1}{2}\underline{\kappa}\underline{\kappa} + 2\rho + r^{-1}\Gamma_g\right) + \underline{\kappa}\left(-\frac{1}{2}\underline{\kappa}^2 + r^{-1}\wp\Gamma_b\right)\right) \\
&= -8\underline{\kappa}^2\rho - \frac{5}{4}\underline{\kappa}\underline{\kappa}^3 + \underline{\kappa}\underline{\kappa}^3 + r^{-3}\wp\Gamma_b + r^{-2}\Gamma_g.
\end{aligned}$$

Hence

$$\begin{aligned}
J_4\nabla_4\alpha &= -\frac{3}{4}\underline{\kappa}^3 + \text{Good}, \\
J_3\nabla_3\alpha &= \left(-6\underline{\kappa}\rho - \frac{19}{4}\underline{\kappa}^3\right)\nabla_3\alpha + \text{Good}, \\
J_0\alpha &= -8\underline{\kappa}^2\rho - \frac{1}{4}\underline{\kappa}\underline{\kappa}^3 + \text{Good}.
\end{aligned}$$

We finally derive,

$$\begin{aligned}
J &= -2\underline{\kappa}\nabla_4\nabla_3\nabla_3\alpha - 5\underline{\kappa}\underline{\kappa}\nabla_3\nabla_3\alpha - \frac{7}{2}\underline{\kappa}^2\nabla_4\nabla_3\alpha \\
&\quad - \frac{3}{4}\underline{\kappa}^3\nabla_4\alpha - \left(6\underline{\kappa}\rho + \frac{19}{4}\underline{\kappa}\underline{\kappa}^2\right)\nabla_3\alpha - \left(8\underline{\kappa}^2\rho + \frac{1}{4}\underline{\kappa}\underline{\kappa}^3\right)\alpha + \text{Good}. \tag{A.14.22}
\end{aligned}$$

Expression for K

Also, using Lemma A.14.19 and Lemma A.14.16 (according to which we have the identity $[Q, \nabla_4]\alpha = (2\rho + \kappa\underline{\kappa})\nabla_3\alpha + \frac{1}{2}\kappa\underline{\kappa}^2\alpha + \text{Good}_1$)

$$\begin{aligned} K &= -\frac{1}{2}[Q, \underline{\kappa}\nabla_4]\alpha = -\frac{1}{2}\left(Q(\underline{\kappa})\nabla_4\alpha + \underline{\kappa}[Q, \nabla_4]\alpha + 2\nabla_3\underline{\kappa}\nabla_3\nabla_4\alpha - \frac{1}{2}\underline{\kappa}^3\nabla_4\alpha\right) \\ &= -\frac{1}{2}\left(Q(\underline{\kappa}) - \frac{1}{2}\underline{\kappa}^3\right)\nabla_4\alpha - \frac{1}{2}\underline{\kappa}\left((2\rho + \kappa\underline{\kappa})\nabla_3\alpha + \frac{1}{2}\kappa\underline{\kappa}^2\alpha\right) \\ &\quad - \nabla_3\underline{\kappa}\nabla_3\nabla_4\alpha + \text{Good}. \end{aligned}$$

Hence,

$$K = -\nabla_3\underline{\kappa}\nabla_3\nabla_4\alpha - \frac{1}{2}\left(Q(\underline{\kappa}) - \frac{1}{2}\underline{\kappa}^3\right)\nabla_4\alpha - \frac{1}{2}\underline{\kappa}(2\rho + \kappa\underline{\kappa})\nabla_3\alpha - \frac{1}{4}\kappa\underline{\kappa}^3\alpha + \text{Good}.$$

We calculate the expression,

$$\begin{aligned} Q(\underline{\kappa}) - \frac{1}{2}\underline{\kappa}^3 &= \nabla_3\nabla_3\underline{\kappa} + 2\underline{\kappa}\nabla_3\underline{\kappa} = \nabla_3\left(-\frac{1}{2}\underline{\kappa}^2 + r^{-1}\wp\Gamma_b\right) + 2\underline{\kappa}\left(-\frac{1}{2}\underline{\kappa}^2 + r^{-1}\wp\Gamma_b\right) \\ &= -\underline{\kappa}(\nabla_3\underline{\kappa} + \underline{\kappa}^2) + \nabla_3(r^{-1}\wp\Gamma_b) + r^{-2}\wp\Gamma_b \\ &= -\frac{1}{2}\underline{\kappa}^3 + \nabla_3(r^{-1}\wp\Gamma_b) + r^{-2}\wp\Gamma_b. \end{aligned}$$

Hence,

$$K = -\nabla_3\underline{\kappa}\nabla_3\nabla_4\alpha + \frac{1}{4}\underline{\kappa}^3\nabla_4\alpha - \frac{1}{2}\underline{\kappa}(2\rho + \kappa\underline{\kappa})\nabla_3\alpha - \frac{1}{4}\kappa\underline{\kappa}^3\alpha + \nabla_3(r^{-1}\wp\Gamma_b)\nabla_4\alpha + \text{Good}.$$

We note that,

$$\begin{aligned} \nabla_3(r^{-1}\wp\Gamma_b)\nabla_4\alpha &= \nabla_3(r^{-1}\wp\Gamma_b\nabla_4\alpha) - r^{-1}\wp\Gamma_b\nabla_3\nabla_4\alpha \\ &= \nabla_3(r^{-1}\wp\Gamma_g\mathfrak{d}^{\leq 1}\alpha) - r^{-2}\wp\Gamma_g\mathfrak{d}^{\leq 2}\alpha = \text{Good}. \end{aligned}$$

We deduce,

$$\begin{aligned} K &= -\nabla_3\underline{\kappa}\nabla_3\nabla_4\alpha + \frac{1}{4}\underline{\kappa}^3\nabla_4\alpha - \frac{1}{2}\underline{\kappa}(2\rho + \kappa\underline{\kappa})\nabla_3\alpha - \frac{1}{4}\kappa\underline{\kappa}^3\alpha + \text{Good} \\ &= -\left(-\frac{1}{2}\underline{\kappa}^2 + r^{-1}\wp\Gamma_b\right)\nabla_3\nabla_4\alpha + \frac{1}{4}\underline{\kappa}^3\nabla_4\alpha - \frac{1}{2}\underline{\kappa}(2\rho + \kappa\underline{\kappa})\nabla_3\alpha - \frac{1}{4}\kappa\underline{\kappa}^3\alpha + \text{Good} \\ &= \frac{1}{2}\underline{\kappa}^2\nabla_3\nabla_4\alpha + \frac{1}{4}\underline{\kappa}^3\nabla_4\alpha - \frac{1}{2}\underline{\kappa}(2\rho + \kappa\underline{\kappa})\nabla_3\alpha - \frac{1}{4}\kappa\underline{\kappa}^3\alpha + \text{Good}. \end{aligned}$$

In view of Lemma A.14.14 $[\nabla_3, \nabla_4]\alpha = 4\rho\alpha + r^{-1}\Gamma_g\wp^{\leq 1}\alpha$. Hence

$$\begin{aligned} K &= \frac{1}{2}\underline{\kappa}^2(\nabla_4\nabla_3\alpha + 4\rho\alpha + r^{-1}\Gamma_g\wp^{\leq 1}\alpha) + \frac{1}{4}\underline{\kappa}^3\nabla_4\alpha - \frac{1}{2}\underline{\kappa}(2\rho + \kappa\underline{\kappa})\nabla_3\alpha - \frac{1}{4}\kappa\underline{\kappa}^3\alpha + \text{Good} \\ &= \frac{1}{2}\underline{\kappa}^2\nabla_4\nabla_3\alpha + \frac{1}{4}\underline{\kappa}^3\nabla_4\alpha - \frac{1}{2}\underline{\kappa}(2\rho + \kappa\underline{\kappa})\nabla_3\alpha + \underline{\kappa}^2\left(2\rho - \frac{1}{4}\kappa\underline{\kappa}\right)\alpha + \text{Good}. \end{aligned}$$

We have thus derived

$$K = \frac{1}{2}\underline{\kappa}^2\nabla_4\nabla_3\alpha + \frac{1}{4}\underline{\kappa}^3\nabla_4\alpha - \frac{1}{2}\underline{\kappa}(2\rho + \kappa\underline{\kappa})\nabla_3\alpha + \underline{\kappa}^2\left(2\rho - \frac{1}{4}\kappa\underline{\kappa}\right)\alpha + \text{Good(A.14.23)}$$

Expression for L

According to Lemma A.14.19 and Lemma A.14.16 (according to which we have the identity $[Q, \nabla_3]\alpha = \underline{\kappa}^2\nabla_3\alpha + \frac{1}{2}\underline{\kappa}^3\alpha + \text{Good}_1$)

$$\begin{aligned} L &= -\frac{5}{2}[Q, \kappa e_3]\alpha = -\frac{5}{2}\left(Q(\kappa)\nabla_3\alpha + \kappa[Q, \nabla_3]\alpha + 2\nabla_3\kappa\nabla_3\nabla_3\alpha - \frac{1}{2}\kappa\underline{\kappa}^2\nabla_3\alpha\right) \\ &= -\frac{5}{2}\left(Q(\kappa)\nabla_3\alpha + \kappa\left(\underline{\kappa}^2\nabla_3\alpha + \frac{1}{2}\underline{\kappa}^3\alpha\right) + 2\nabla_3\kappa\nabla_3\nabla_3\alpha - \frac{1}{2}\kappa\underline{\kappa}^2\nabla_3\alpha\right) + \text{Good} \\ &= -5\nabla_3\kappa\nabla_3\nabla_3\alpha - \frac{5}{2}\left(Q(\kappa) + \frac{1}{2}\kappa\underline{\kappa}\right)\nabla_3\alpha - \frac{5}{4}\kappa\underline{\kappa}^3\alpha + \text{Good}. \end{aligned}$$

Note that

$$\begin{aligned} Q(\kappa) &= \nabla_3\nabla_3\kappa + 2\underline{\kappa}\nabla_3\kappa + \frac{1}{2}\kappa\underline{\kappa}^2 \\ &= \nabla_3\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho + r^{-1}\wp\Gamma_g\right) + 2\underline{\kappa}\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho + r^{-1}\wp\Gamma_g\right) + \frac{1}{2}\kappa\underline{\kappa}^2 \\ &= -\frac{1}{2}(\kappa\nabla_3\underline{\kappa} + \underline{\kappa}\nabla_3\kappa) + 2\left(-\frac{3}{2}\underline{\kappa}\rho + r^{-1}\wp\Gamma_g\right) - \kappa\underline{\kappa}^2 + 4\rho\underline{\kappa} + \frac{1}{2}\kappa\underline{\kappa}^2 \\ &\quad + e_3(r^{-1}\wp\Gamma_g) + r^{-2}\wp\Gamma_g \\ &= -\frac{1}{2}(\kappa\nabla_3\underline{\kappa} + \underline{\kappa}\nabla_3\kappa) + \rho\underline{\kappa} - \frac{1}{2}\kappa\underline{\kappa}. \end{aligned}$$

Therefore,

$$\begin{aligned} Q(\kappa) &= -\frac{1}{2}\kappa\left(-\frac{1}{2}\underline{\kappa}^2 + r^{-1}\wp\Gamma_b\right) - \frac{1}{2}\underline{\kappa}\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho + r^{-1}\wp\Gamma_g\right) + \underline{\kappa}\rho - \frac{1}{2}\kappa\underline{\kappa}^2 \\ &\quad + r^{-1}\wp^{\leq 2}\Gamma_g = r^{-1}\wp^{\leq 2}\Gamma_g. \end{aligned}$$

We deduce,

$$\begin{aligned} L &= -5\nabla_3\kappa\nabla_3\nabla_3\alpha - \frac{5}{4}\kappa\underline{\kappa}^2\nabla_3\alpha + \frac{5}{4}\kappa\underline{\kappa}^3\alpha + \text{Good} \\ &= -5\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho\right)\nabla_3\nabla_3\alpha - \frac{5}{4}\kappa\underline{\kappa}^2\nabla_3\alpha + \frac{5}{4}\kappa\underline{\kappa}^3\alpha + \text{Good}. \end{aligned}$$

Therefore,

$$L = -5\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho\right)\nabla_3\nabla_3\alpha - \frac{5}{4}\kappa\underline{\kappa}^2\nabla_3\alpha + \frac{5}{4}\kappa\underline{\kappa}^3\alpha + \text{Good}. \quad (\text{A.14.24})$$

Expression for M

Similarly, according to Lemma A.14.19,

$$M = \left[Q, 4\rho - \frac{1}{2}\kappa\underline{\kappa}\right]\alpha = Q\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right)\alpha + 2\nabla_3\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right)\nabla_3\alpha - \frac{1}{2}\kappa^2\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right)\alpha$$

i.e.,

$$M = Q\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right)\alpha + 2\nabla_3\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right)\nabla_3\alpha - \frac{1}{2}\kappa^2\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right)\alpha.$$

We calculate,

$$\begin{aligned} \nabla_3\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right) &= 4\nabla_3\rho - \frac{1}{2}\kappa\nabla_3\underline{\kappa} - \frac{1}{2}\underline{\kappa}\nabla_3\kappa \\ &= 4\left(-\frac{3}{2}\underline{\kappa}\rho + r^{-1}\wp\Gamma_g\right) - \frac{1}{2}\kappa\left(-\frac{1}{2}\underline{\kappa}^2 + r^{-1}\wp\Gamma_b\right) \\ &\quad - \frac{1}{2}\underline{\kappa}\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho + r^{-1}\wp\Gamma_g\right) \\ &= -7\underline{\kappa}\rho + \frac{1}{2}\kappa\underline{\kappa}^2 + r^{-1}\wp\Gamma_g. \end{aligned}$$

We deduce,

$$M = \left(Q\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right) - \frac{1}{2}\kappa^2\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right)\right)\alpha + \left(-7\underline{\kappa}\rho + \frac{1}{2}\kappa\underline{\kappa}^2\right)\nabla_3\alpha + \text{Good}.$$

It remains to calculate

$$\begin{aligned}
M_0 &= Q\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right) - \frac{1}{2}\underline{\kappa}^2\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right) = \nabla_3\nabla_3\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right) + 2\underline{\kappa}\nabla_3\left(4\rho - \frac{1}{2}\kappa\underline{\kappa}\right) \\
&= \nabla_3\left(-7\underline{\kappa}\rho + \frac{1}{2}\kappa\underline{\kappa}^2 + r^{-1}\wp\Gamma_g\right) + 2\underline{\kappa}\left(-7\underline{\kappa}\rho + \frac{1}{2}\kappa\underline{\kappa}^2 + r^{-1}\wp\Gamma_g\right) \\
&= -7\rho\nabla_3\underline{\kappa} - 7\underline{\kappa}\nabla_3\rho + \frac{1}{2}\underline{\kappa}^2\nabla_3\underline{\kappa} + \kappa\underline{\kappa}\nabla_3\underline{\kappa} + 2\underline{\kappa}\left(-7\underline{\kappa}\rho + \frac{1}{2}\kappa\underline{\kappa}^2\right) \\
&\quad + \nabla_3(r^{-1}\wp\Gamma_g) + r^{-2}\wp\Gamma_g.
\end{aligned}$$

Hence,

$$\begin{aligned}
M_0 &= -7\rho\left(-\frac{1}{2}\underline{\kappa}^2 + r^{-1}\wp\Gamma_b\right) - 7\underline{\kappa}\left(-\frac{3}{2}\underline{\kappa}\rho + r^{-1}\wp\Gamma_g\right) + \frac{1}{2}\underline{\kappa}^2\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho + r^{-1}\wp\Gamma_g\right) \\
&\quad + \kappa\underline{\kappa}\left(-\frac{1}{2}\underline{\kappa}^2 + r^{-1}\wp\Gamma_b\right) + 2\underline{\kappa}\left(-7\underline{\kappa}\rho + \frac{1}{2}\kappa\underline{\kappa}^2\right) + \nabla_3(r^{-1}\wp\Gamma_g) + r^{-2}\wp\Gamma_g \\
&= \underline{\kappa}^2\rho + \frac{1}{4}\kappa\underline{\kappa}^3 + \nabla_3(r^{-1}\wp\Gamma_g) + r^{-2}\wp\Gamma_g.
\end{aligned}$$

We conclude,

$$M = \left(\underline{\kappa}^2\rho + \frac{1}{4}\kappa\underline{\kappa}^3\right)\alpha + 2\left(-7\underline{\kappa}\rho + \frac{1}{2}\kappa\underline{\kappa}^2\right)\nabla_3\alpha + \text{Good}. \quad (\text{A.14.25})$$

Indeed note that

$$\nabla_3(r^{-1}\wp\Gamma_g)\alpha = \nabla_3(r^{-1}\wp\Gamma_g\alpha) - r^{-1}\wp\Gamma_g\nabla_3\alpha = \text{Good}.$$

End of the proof of Proposition A.14.20

Using the equations (A.14.21)–(A.14.25) we deduce, back to (A.14.20),

$$\begin{aligned}
[Q, \mathcal{L}]\alpha &= I + J + K + L + M \\
&= (2\rho - \kappa\underline{\kappa})\nabla_3\nabla_3\alpha - \underline{\kappa}^2\nabla_4\nabla_3\alpha - \frac{1}{2}\underline{\kappa}^3\nabla_4\alpha - \left(-\frac{1}{2}\kappa\underline{\kappa}^2 + 3\rho\underline{\kappa}\right)\nabla_3\alpha \\
&\quad - \frac{3}{2}\underline{\kappa}^2\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho\right)\alpha - 2\underline{\kappa}\nabla_4\nabla_3\nabla_3\alpha - 5\kappa\underline{\kappa}\nabla_3\nabla_3\alpha - \frac{7}{2}\underline{\kappa}^2\nabla_4\nabla_3\alpha \\
&\quad - \frac{3}{4}\underline{\kappa}^3\nabla_4\alpha - \left(6\underline{\kappa}\rho + \frac{19}{4}\kappa\underline{\kappa}^2\right)\nabla_3\alpha - \left(8\underline{\kappa}^2\rho + \frac{1}{4}\kappa\underline{\kappa}^3\right)\alpha \\
&\quad + \frac{1}{2}\underline{\kappa}^2\nabla_4\nabla_3\alpha + \frac{1}{4}\underline{\kappa}^3\nabla_4\alpha - \frac{1}{2}\underline{\kappa}(2\rho + \kappa\underline{\kappa})\nabla_3\alpha + \underline{\kappa}^2\left(2\rho - \frac{1}{4}\kappa\underline{\kappa}\right)\alpha \\
&\quad - 5\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho\right)\nabla_3\nabla_3\alpha - \frac{5}{4}\kappa\underline{\kappa}^2\nabla_3\alpha - \frac{5}{4}\kappa\underline{\kappa}^3\alpha \\
&\quad + \left(\underline{\kappa}^2\rho + \frac{1}{4}\kappa\underline{\kappa}^3\right)\alpha + 2\left(-7\underline{\kappa}\rho + \frac{1}{2}\kappa\underline{\kappa}^2\right)\nabla_3\alpha + \text{Good}.
\end{aligned}$$

We deduce,

$$[Q, \mathcal{L}]\alpha = -2\underline{\kappa}\nabla_4\nabla_3\nabla_3\alpha + C'_{33}\nabla_3\nabla_3\alpha + C'_{43}\nabla_4\nabla_3\alpha + C'_4\nabla_4\alpha + C'_3\nabla_3\alpha + C'_0\alpha$$

with,

$$\begin{aligned}
C'_{33} &= (2\rho - \kappa\underline{\kappa}) - 5\kappa\underline{\kappa} - 5\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho\right) = -8\rho - \frac{7}{2}\kappa\underline{\kappa}, \\
C'_{43} &= -\underline{\kappa}^2 - \frac{7}{2}\underline{\kappa}^2 + \frac{1}{2}\underline{\kappa}^2 = -4\underline{\kappa}^2, \\
C'_4 &= -\frac{1}{2}\underline{\kappa}^3 - \frac{3}{4}\underline{\kappa}^3 + \frac{1}{4}\underline{\kappa}^3 = -\underline{\kappa}^3, \\
C'_3 &= \frac{1}{2}\kappa\underline{\kappa}^2 - 3\rho\underline{\kappa} - \left(6\underline{\kappa}\rho + \frac{19}{4}\kappa\underline{\kappa}^2\right) - \frac{1}{2}\underline{\kappa}(2\rho + \kappa\underline{\kappa}) - \frac{5}{4}\kappa\underline{\kappa}^2 + 2\left(-7\underline{\kappa}\rho + \frac{1}{2}\kappa\underline{\kappa}^2\right) \\
&= -24\underline{\kappa}\rho - 5\kappa\underline{\kappa}^2, \\
C'_0 &= -\frac{3}{2}\underline{\kappa}^2\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\rho\right) - \left(8\underline{\kappa}^2\rho + \frac{1}{4}\kappa\underline{\kappa}^3\right) + \underline{\kappa}^2\left(2\rho - \frac{1}{4}\kappa\underline{\kappa}\right) - \frac{5}{4}\kappa\underline{\kappa}^3 + \left(\underline{\kappa}^2\rho + \frac{1}{4}\kappa\underline{\kappa}^3\right) \\
&= -8\underline{\kappa}^2\rho - \frac{3}{4}\kappa\underline{\kappa}^3.
\end{aligned}$$

Finally we write, recalling the definition of $Q = \nabla_3\nabla_3 + 2\underline{\kappa}\nabla_3 + \frac{1}{2}\underline{\kappa}^2$,

$$\nabla_3\nabla_3\alpha = Q(\alpha) - 2\underline{\kappa}\nabla_3\alpha - \frac{1}{2}\underline{\kappa}^2\alpha$$

and,

$$\nabla_4 \nabla_3 \nabla_3 \alpha = \nabla_4 Q(\alpha) - 2\underline{\kappa} \nabla_4 \nabla_3 \alpha - \frac{1}{2} \underline{\kappa}^2 \nabla_4 \alpha - 2 \nabla_4 \underline{\kappa} \nabla_3 \alpha - \underline{\kappa} \nabla_4 \underline{\kappa} \alpha.$$

Hence,

$$\begin{aligned} -2\underline{\kappa} \nabla_4 \nabla_3 \nabla_3 \alpha + C'_{33} \nabla_3 \nabla_3 \alpha &= -2\underline{\kappa} \nabla_4 Q(\alpha) + 4\underline{\kappa}^2 \nabla_4 \nabla_3 \alpha + \underline{\kappa}^3 \nabla_4 \alpha + 4\underline{\kappa} \nabla_4 \underline{\kappa} \nabla_3 \alpha \\ &+ 2\underline{\kappa}^2 \nabla_4 \underline{\kappa} \alpha + C'_{33} \left(Q(\alpha) - 2\underline{\kappa} \nabla_3 \alpha - \frac{1}{2} \underline{\kappa}^2 \alpha \right). \end{aligned}$$

We deduce,

$$\begin{aligned} [Q, \mathcal{L}] \alpha &= -2\underline{\kappa} \nabla_4 Q(\alpha) + C'_{33} Q(\alpha) + 4\underline{\kappa}^2 \nabla_4 \nabla_3 \alpha + \underline{\kappa}^3 \nabla_4 \alpha + (4\underline{\kappa} \nabla_4 \underline{\kappa} - 2\underline{\kappa} C'_{33}) \nabla_3 \alpha \\ &+ \left(2\underline{\kappa}^2 \nabla_4 \underline{\kappa} - \frac{1}{2} \underline{\kappa}^2 C'_{33} \right) \alpha + C'_{43} \nabla_4 \nabla_3 \alpha + C'_4 \nabla_4 \alpha + C'_3 \nabla_3 \alpha + C'_0 \alpha. \end{aligned}$$

Thus, setting $C_Q = C'_{33}$, we deduce,

$$[Q, \mathcal{L}] \alpha = -2\underline{\kappa} \nabla_4 Q(\alpha) + C_Q Q(\alpha) + C_{43} \nabla_4 \nabla_3 \alpha + C_4 \nabla_4 \alpha + C_3 \nabla_3 \alpha + C_0 \alpha + \text{Good}$$

where,

$$\begin{aligned} C_Q &= C'_{33} = -8\rho - \frac{7}{2} \underline{\kappa} \underline{\kappa}, \\ C_{43} &= 4\underline{\kappa}^2 + C'_{43} = 4\underline{\kappa}^2 - 4\underline{\kappa}^2 = 0, \\ C_4 &= \underline{\kappa}^3 + C'_4 = \underline{\kappa}^3 - \underline{\kappa}^3 = 0. \end{aligned}$$

Also,

$$\begin{aligned} C_3 &= 2\underline{\kappa} (2 \nabla_4 \underline{\kappa} - C'_{33}) + C'_3 \\ &= 2\underline{\kappa} \left(-\underline{\kappa} \underline{\kappa} + 4\rho + r^{-1} \not\partial \Gamma_g + 8\rho + \frac{7}{2} \underline{\kappa} \underline{\kappa} \right) + C'_3 \\ &= 2\underline{\kappa} \left(12\rho + \frac{5}{2} \underline{\kappa} \underline{\kappa} \right) + (-24\underline{\kappa} \rho - 5\underline{\kappa} \underline{\kappa}^2) + r^{-2} \not\partial \Gamma_g \\ &= r^{-2} \not\partial \Gamma_g, \\ C_0 &= 2\underline{\kappa}^2 \nabla_4 \underline{\kappa} - \frac{1}{2} \underline{\kappa}^2 C'_{33} + C'_0 \\ &= 2\underline{\kappa}^2 \left(-\frac{1}{2} \underline{\kappa} \underline{\kappa} + 2\rho + r^{-1} \not\partial \Gamma_g \right) + \frac{1}{2} \underline{\kappa}^2 \left(8\rho + \frac{7}{2} \underline{\kappa} \underline{\kappa} \right) - 8\underline{\kappa}^2 \rho - \frac{3}{4} \underline{\kappa} \underline{\kappa}^3 \\ &= 8\underline{\kappa}^2 \rho + \frac{3}{4} \underline{\kappa} \underline{\kappa}^3 - 8\underline{\kappa}^2 \rho - \frac{3}{4} \underline{\kappa} \underline{\kappa}^3 + r^{-3} \not\partial \Gamma_g \\ &= r^{-3} \not\partial \Gamma_g. \end{aligned}$$

We have therefore checked that,

$$[Q, \mathcal{L}]\alpha = -2\underline{\kappa}\nabla_4 Q(\alpha) + C_Q Q(\alpha) + \text{Good}, \quad C_Q = -8\rho - \frac{7}{2}\underline{\kappa}\underline{\kappa},$$

as stated in Proposition A.14.20. □

A.14.4 Proof of Theorem 2.4.7

We start with the following,

Lemma A.14.21. *We have,*

$$\square_2(fr^4) = r^4\square_2 f - 2r^4(\underline{\kappa}e_4 f + \kappa e_3 f) + r^4(-5\underline{\kappa}\underline{\kappa} - 4\rho)f + O(r^4\mathfrak{d}^{\leq 1}\Gamma_g \cdot f).$$

We postpone the proof of the lemma to the end of the section and continue below the proof of the theorem. According to Lemma A.14.13

$$\mathcal{L}(\alpha) = \text{Good}_2$$

where \mathcal{L} is the operator

$$\mathcal{L}\alpha = -\nabla_4\nabla_3\alpha + {}^{(c)}\Delta_2\alpha - \frac{5}{2}\kappa\nabla_3\alpha - \frac{1}{2}\underline{\kappa}\nabla_4\alpha - \left(-4\rho + \frac{1}{2}\underline{\kappa}\underline{\kappa}\right)\alpha.$$

Applying Q and recalling the definition of the error terms Good we derive,

$$\mathcal{L}(Q\alpha) = -[Q, \mathcal{L}]\alpha + \text{Good}.$$

Thus, in view of Proposition A.14.20,

$$[Q, \mathcal{L}]\alpha = -2\underline{\kappa}\nabla_4 Q(\alpha) + C_Q Q(\alpha), \quad C_Q = -8\rho - \frac{7}{2}\underline{\kappa}\underline{\kappa}.$$

We deduce,

$$\mathcal{L}(Q\alpha) = 2\underline{\kappa}\nabla_4 Q(\alpha) - C_Q Q(\alpha).$$

Therefore, modulo Good terms,

$$\begin{aligned} 2\underline{\kappa}\nabla_4 Q(\alpha) - C_Q Q(\alpha) &= -\nabla_4\nabla_3(Q\alpha) + {}^{(c)}\Delta_2(Q\alpha) - \frac{5}{2}\kappa\nabla_3 Q(\alpha) - \frac{1}{2}\underline{\kappa}\nabla_4 Q(\alpha) \\ &\quad - \left(-4\rho + \frac{1}{2}\underline{\kappa}\underline{\kappa}\right) Q(\alpha). \end{aligned}$$

We deduce

$$-\nabla_4\nabla_3(Q\alpha) + {}^{(c)}\underline{\Delta}_2(Q\alpha) - \frac{5}{2}\underline{\kappa}\nabla_3Q(\alpha) - \frac{5}{2}\underline{\kappa}\nabla_4Q(\alpha) + \left(C_Q - \left(-4\rho + \frac{1}{2}\underline{\kappa}\underline{\kappa}\right)\right)Q(\alpha) + \text{Good}.$$

In view of the expression for \square_2 in the second part of the Lemma A.14.13 we rewrite in the form

$$\square_2Q(f) - 2\underline{\kappa}\nabla_3Q(\alpha) - 2\underline{\kappa}\nabla_4Q(\alpha) - (4\rho + 4\underline{\kappa}\underline{\kappa})Q(\alpha) = \text{Good} + r^{-1}\Gamma_g \cdot \not\partial Q(\alpha).$$

Finally, making use of Lemma A.14.21 and recalling that $\mathfrak{q} = r^4Q(\alpha)$,

$$\begin{aligned} \square_2\mathfrak{q} &= r^4\square_2(Q\alpha) - 2r^4(\underline{\kappa}e_4(Q\alpha) + \underline{\kappa}e_3(Q\alpha)) + r^4(-5\underline{\kappa}\underline{\kappa} - 4\rho)Qf + O(r^4\mathfrak{d}^{\leq 1}\Gamma_g \cdot Q(\alpha)) \\ &= r^4\left(2\underline{\kappa}\nabla_3Q(\alpha) + 2\underline{\kappa}\nabla_4Q(\alpha) + (4\rho + 4\underline{\kappa}\underline{\kappa})Q(\alpha) + \text{Good}\right) \\ &\quad - 2r^4(\underline{\kappa}e_4(Q\alpha) + \underline{\kappa}e_3(Q\alpha)) + r^4(-5\underline{\kappa}\underline{\kappa} - 4\rho)Qf + O(\mathfrak{d}^{\leq 1}\Gamma_g \cdot \mathfrak{q}) \\ &= -\underline{\kappa}\underline{\kappa}\mathfrak{q} + r^4\text{Good}. \end{aligned}$$

This ends the proof of Theorem 2.4.7.

Proof of Lemma A.14.21

We have,

$$\begin{aligned} \square_2(fr^4) &= \mathbf{D}^\alpha\mathbf{D}_\alpha(fr^4) = \mathbf{D}^\alpha(\mathbf{D}_\alpha fr^4 + f\mathbf{D}_\alpha r^4) \\ &= r^4\square_2f + 2\mathbf{D}_\alpha(r^4)\mathbf{D}^\alpha f + f\square(r) \\ &= r^4\square_2f - (e_3(r^4)e_4f + e_4(r^4)e_3f) + f\square(r^4) + r^4\Gamma_g\mathfrak{d}f \\ &= r^4\square_2f - 4r^3(e_3(r)e_4f + e_4(r)e_3f) + f\square(r^4) + r^4\Gamma_g\mathfrak{d}f \\ &= r^4\square_2f - 2r^4\left((\underline{\kappa} + \Gamma_b)e_4f + (\underline{\kappa} + \Gamma_g)e_3f\right) + f\square(r^4) + r^4\Gamma_g \cdot \mathfrak{d}f \\ &= r^4\square_2f - 2r^4(\underline{\kappa}e_4f + \underline{\kappa}e_3f) + f\square(r^4) + r^4\Gamma_g \cdot f. \end{aligned}$$

Also,

$$\begin{aligned} \square(r^4) &= -e_4(e_3(r^4)) - \frac{1}{2}\underline{\kappa}e_4(r^4) + \left(-\frac{1}{2}\underline{\kappa} + 2\omega\right)e_3(r^4) + \underline{\Delta}(r^4) + 2\underline{\eta}e_\theta(r^4) \\ &= -4e_4(r^3e_3(r)) - 2r^3\underline{\kappa}\frac{r}{2}(\underline{\kappa} + \Gamma_g) + 4r^3\left(-\frac{1}{2}\underline{\kappa} + 2\omega\right)\frac{r}{2}(\underline{\kappa} + \Gamma_b) + \underline{\Delta}(r^4) + 2\underline{\eta}e_\theta(r^4) \\ &= -12r^2(e_4r)(e_3r) - 4r^3e_4e_3r - r^4\underline{\kappa}\underline{\kappa} + 2r^4\left(-\frac{1}{2}\underline{\kappa} + 2\omega\right)\underline{\kappa} + O(r^3\Gamma_b) \\ &= -3r^4(\underline{\kappa} + \Gamma_b)(\underline{\kappa} + \Gamma_g) - 4r^3e_4\left(\frac{r}{2}(\underline{\kappa} + \Gamma_b)\right) - 2r^4\underline{\kappa}\underline{\kappa} + 4r^4\omega\underline{\kappa} + O(r^3\Gamma_b). \end{aligned}$$

Hence,

$$\square(r^4) = -5r^4\kappa_{\underline{\kappa}} + 4r^4\omega_{\underline{\kappa}} - 2r^3e_4(r\underline{\kappa}) + O(r^4\mathfrak{d}^{\leq 1}\Gamma_g).$$

Note that,

$$\begin{aligned} e_4(r\underline{\kappa}) &= re_4(\underline{\kappa}) + \frac{r}{2}\underline{\kappa}(\kappa + \Gamma_g) = r \left(-\frac{1}{2}\kappa_{\underline{\kappa}} + 2\omega_{\underline{\kappa}} + 2\mathfrak{d}_1\underline{\eta} + 2\rho \right) + \frac{r}{2}\underline{\kappa}(\kappa + \Gamma_g) \\ &= 2r\rho + 2r\omega_{\underline{\kappa}} + O(\mathfrak{d}^{\leq 1}\Gamma_g). \end{aligned}$$

Hence,

$$\begin{aligned} \square(r^4) &= -5r^4\kappa_{\underline{\kappa}} + 4r^4\omega_{\underline{\kappa}} - 2r^3(2r\rho + 2r\omega_{\underline{\kappa}}) + O(r^4\mathfrak{d}^{\leq 1}\Gamma_g) \\ &= r^4(-5\kappa_{\underline{\kappa}} - 4\rho) + O(r^4\mathfrak{d}^{\leq 1}\Gamma_g). \end{aligned}$$

We conclude

$$\square_2(fr^4) = r^4\square_2f - 2r^4(\underline{\kappa}e_4f + \kappa e_3f) + r^4(-5\kappa_{\underline{\kappa}} - 4\rho)f + O(r^4\mathfrak{d}^{\leq 1}\Gamma_g)$$

as stated.

Appendix B

APPENDIX TO CHAPTER 8

B.1 Proof of Proposition 8.4.1

Proposition B.1.1. *The following wave equations hold true.*

1. *The null curvature component ρ verifies the identity*

$$\square_{\mathbf{g}}\rho := \underline{\kappa}e_4\rho + \kappa e_3\rho + \frac{3}{2}(\underline{\kappa}\kappa + 2\rho)\rho + Err[\square_{\mathbf{g}}\rho],$$

where

$$\begin{aligned} Err[\square_{\mathbf{g}}\rho] &= \frac{3}{2}\rho \left(-\frac{1}{2}\underline{\vartheta}\vartheta + 2(\underline{\xi}\xi + \eta\eta) \right) + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega} \right) \left(\frac{1}{2}\underline{\vartheta}\alpha - \zeta\beta - 2(\underline{\eta}\beta + \xi\underline{\beta}) \right) \\ &\quad - \frac{1}{2}\underline{\vartheta}\phi_2^*\beta + (\zeta - \eta)e_3\beta - \eta e_3(\Phi)\beta - \underline{\xi}(e_4\beta + e_4(\Phi)\beta) - \underline{\beta}\beta \\ &\quad - e_3 \left(-\frac{1}{2}\underline{\vartheta}\alpha + \zeta\beta + 2(\underline{\eta}\beta + \xi\underline{\beta}) \right) \\ &\quad - \phi_1^*(\underline{\kappa})\beta + 2\phi_1^*(\underline{\omega})\beta + 3\eta\phi_1^*(\rho) - \phi_1 \left(-\vartheta\underline{\beta} + \underline{\xi}\alpha \right) - 2\eta e_\theta\rho. \end{aligned}$$

2. *The small curvature quantity,*

$$\tilde{\rho} := r^2 \left(\rho + \frac{2m}{r^3} \right)$$

verifies the wave equation,

$$\begin{aligned} \square_{\mathbf{g}}(\tilde{\rho}) + \frac{8m}{r^3}\tilde{\rho} &= -6m \frac{\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2}\right)}{r^2} - \frac{3m}{r} \left(\underline{\kappa} + \frac{4\underline{\Upsilon}}{r^2} \right) \\ &\quad - \frac{3m}{r} (\underline{A}\underline{\kappa} + \underline{A}\kappa) + Err[\square_{\mathbf{g}}\tilde{\rho}], \end{aligned}$$

where

$$\begin{aligned} Err[\square_{\mathbf{g}}\tilde{\rho}] &:= -\frac{6m}{r}\underline{A}\underline{A} + \frac{3}{r^2}\tilde{\rho}^2 + \frac{3}{2} \left(\frac{4}{3}\underline{A} \frac{e_3(r)}{r} + \frac{4}{3}\underline{A} \frac{e_4(r)}{r} \right) \tilde{\rho} \\ &\quad + \left(\frac{3}{2} \left(\underline{\kappa}\underline{\kappa} - \frac{8m}{r^3} + \frac{2}{3r^2}\square_{\mathbf{g}}(r^2) \right) + \frac{8m}{r^3} \right) \tilde{\rho} \\ &\quad - \underline{A}e_3(\tilde{\rho}) - \underline{A}e_4(\tilde{\rho}) + \frac{2}{r}\underline{A}e_3(m) + \frac{2}{r}\underline{A}e_4(m) \\ &\quad + 4D^a(m)D_a \left(\frac{1}{r} \right) + \frac{2}{r}\square_{\mathbf{g}}(m) + 4r \underline{\phi}_1^*(r) \underline{\phi}_1^*(\rho) + r^2 Err[\square_{\mathbf{g}}\rho]. \end{aligned}$$

Proof. We prove the result in the following steps.

Step 1. We start by deriving the wave equation for ρ . From Bianchi, ρ satisfies

$$e_4\rho + \frac{3}{2}\underline{\kappa}\rho = \underline{\phi}_1\beta - \frac{1}{2}\underline{\vartheta}\alpha + \zeta\beta + 2(\underline{\eta}\beta + \underline{\xi}\beta).$$

Differentiating with respect to e_3 , we obtain

$$e_3(e_4(\rho)) + \frac{3}{2}\underline{\kappa}e_3(\rho) + \frac{3}{2}e_3(\underline{\kappa})\rho = e_3(\underline{\phi}_1\beta) + e_3 \left(-\frac{1}{2}\underline{\vartheta}\alpha + \zeta\beta + 2(\underline{\eta}\beta + \underline{\xi}\beta) \right).$$

Also, β satisfies from Bianchi

$$e_3\beta + \underline{\kappa}\beta = -\underline{\phi}_1^*\rho + 2\underline{\omega}\beta + 3\underline{\eta}\rho - \underline{\vartheta}\beta + \underline{\xi}\alpha.$$

Differentiating with respect to $\underline{\phi}_1$, we infer

$$\begin{aligned} \underline{\phi}_1(e_3\beta) + \underline{\kappa}\underline{\phi}_1\beta - \underline{\phi}_1^*(\underline{\kappa})\beta &= -\underline{\phi}_1 \underline{\phi}_1^*\rho + 2\underline{\omega}\underline{\phi}_1\beta - 2\underline{\phi}_1^*(\underline{\omega})\beta + 3\rho \underline{\phi}_1\underline{\eta} - 3\underline{\eta} \underline{\phi}_1^*(\rho) \\ &\quad + \underline{\phi}_1 \left(-\underline{\vartheta}\beta + \underline{\xi}\alpha \right) \end{aligned}$$

and hence

$$\begin{aligned} \underline{\phi}_1 \underline{\phi}_1^*\rho &= -\underline{\phi}_1(e_3\beta) - \underline{\kappa}\underline{\phi}_1\beta + 2\underline{\omega}\underline{\phi}_1\beta + 3\rho \underline{\phi}_1\underline{\eta} \\ &\quad + \underline{\phi}_1^*(\underline{\kappa})\beta - 2\underline{\phi}_1^*(\underline{\omega})\beta - 3\underline{\eta} \underline{\phi}_1^*(\rho) + \underline{\phi}_1 \left(-\underline{\vartheta}\beta + \underline{\xi}\alpha \right). \end{aligned}$$

Next, we add the equation for $\not{d}_1 \not{d}_1^* \rho$ from the one for $e_3(e_4(\rho))$. This yields

$$\begin{aligned} & e_3(e_4(\rho)) + \not{d}_1 \not{d}_1^* \rho + \frac{3}{2} \kappa e_3(\rho) + \frac{3}{2} e_3(\kappa) \rho \\ = & [e_3, \not{d}_1] \beta - \underline{\kappa} \not{d}_1 \beta + 2 \underline{\omega} \not{d}_1 \beta + 3 \rho \not{d}_1 \eta + e_3 \left(-\frac{1}{2} \underline{\vartheta} \alpha + \zeta \beta + 2(\underline{\eta} \beta + \xi \underline{\beta}) \right) \\ & + \not{d}_1^*(\underline{\kappa}) \beta - 2 \not{d}_1^*(\underline{\omega}) \beta - 3 \eta \not{d}_1^*(\rho) + \not{d}_1 \left(-\underline{\vartheta} \beta + \underline{\xi} \alpha \right). \end{aligned}$$

Next, we recall the following commutator identity

$$[e_3, \not{d}_1] \beta = -\frac{1}{2} \underline{\kappa} \not{d}_1 \beta + \frac{1}{2} \underline{\vartheta} \not{d}_2^* \beta - (\zeta - \eta) e_3 \beta + \eta e_3(\Phi) \beta + \underline{\xi} (e_4 \beta + e_4(\Phi) \beta) + \underline{\beta} \beta.$$

We infer

$$\begin{aligned} & e_3(e_4(\rho)) + \not{d}_1 \not{d}_1^* \rho + \frac{3}{2} \kappa e_3(\rho) + \frac{3}{2} e_3(\kappa) \rho + \left(\frac{3}{2} \underline{\kappa} - 2 \underline{\omega} \right) \not{d}_1 \beta - 3 \rho \not{d}_1 \eta \\ = & \frac{1}{2} \underline{\vartheta} \not{d}_2^* \beta - (\zeta - \eta) e_3 \beta + \eta e_3(\Phi) \beta + \underline{\xi} (e_4 \beta + e_4(\Phi) \beta) + \underline{\beta} \beta \\ & + e_3 \left(-\frac{1}{2} \underline{\vartheta} \alpha + \zeta \beta + 2(\underline{\eta} \beta + \xi \underline{\beta}) \right) \\ & + \not{d}_1^*(\underline{\kappa}) \beta - 2 \not{d}_1^*(\underline{\omega}) \beta - 3 \eta \not{d}_1^*(\rho) + \not{d}_1 \left(-\underline{\vartheta} \beta + \underline{\xi} \alpha \right). \end{aligned}$$

Next, we make use of the Bianchi identities and the null structure equations to compute

$$\begin{aligned} & \frac{3}{2} e_3(\kappa) \rho + \left(\frac{3}{2} \underline{\kappa} - 2 \underline{\omega} \right) \not{d}_1 \beta - 3 \rho \not{d}_1 \eta \\ = & \frac{3}{2} \rho \left(-\frac{1}{2} \underline{\kappa} \kappa + 2 \underline{\omega} \kappa + 2 \not{d}_1 \eta + 2 \rho - \frac{1}{2} \underline{\vartheta} \vartheta + 2(\underline{\xi} \xi + \eta \eta) \right) \\ & + \left(\frac{3}{2} \underline{\kappa} - 2 \underline{\omega} \right) \left(e_4 \rho + \frac{3}{2} \kappa \rho + \frac{1}{2} \underline{\vartheta} \alpha - \zeta \beta - 2(\underline{\eta} \beta + \xi \underline{\beta}) \right) - 3 \rho \not{d}_1 \eta \\ = & \left(\frac{3}{2} \underline{\kappa} - 2 \underline{\omega} \right) e_4 \rho + \frac{3}{2} \rho (\underline{\kappa} \kappa + 2 \rho) \\ & + \frac{3}{2} \rho \left(-\frac{1}{2} \underline{\vartheta} \vartheta + 2(\underline{\xi} \xi + \eta \eta) \right) + \left(\frac{3}{2} \underline{\kappa} - 2 \underline{\omega} \right) \left(\frac{1}{2} \underline{\vartheta} \alpha - \zeta \beta - 2(\underline{\eta} \beta + \xi \underline{\beta}) \right). \end{aligned}$$

This yields

$$\begin{aligned}
& e_3(e_4(\rho)) - \underline{\Delta}\rho + \frac{3}{2}\kappa e_3(\rho) + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega}\right) e_4\rho + \frac{3}{2}\rho(\underline{\kappa}\kappa + 2\rho) \\
= & -\frac{3}{2}\rho\left(-\frac{1}{2}\underline{\vartheta}\vartheta + 2(\underline{\xi}\xi + \eta\eta)\right) - \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega}\right)\left(\frac{1}{2}\underline{\vartheta}\alpha - \zeta\beta - 2(\underline{\eta}\beta + \xi\underline{\beta})\right) \\
& + \frac{1}{2}\underline{\vartheta}\underline{\phi}_2^*\beta - (\zeta - \eta)e_3\beta + \eta e_3(\Phi)\beta + \underline{\xi}(e_4\beta + e_4(\Phi)\beta) + \underline{\beta}\beta \\
& + e_3\left(-\frac{1}{2}\underline{\vartheta}\alpha + \zeta\beta + 2(\underline{\eta}\beta + \xi\underline{\beta})\right) \\
& + \underline{\phi}_1^*(\underline{\kappa})\beta - 2\underline{\phi}_1^*(\underline{\omega})\beta - 3\eta\underline{\phi}_1^*(\rho) + \underline{\phi}_1(-\vartheta\underline{\beta} + \underline{\xi}\alpha),
\end{aligned}$$

where we used the fact that $\underline{\phi}_1\underline{\phi}_1^* = -\underline{\Delta}$.

Next, recall the formula for the wave operator acting on a scalar ψ

$$\square_{\mathbf{g}}\psi = -e_3e_4\psi + \underline{\Delta}\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi - \frac{1}{2}\kappa e_3\psi + 2\eta e_\theta\psi.$$

We infer

$$\begin{aligned}
& e_3(e_4(\rho)) - \underline{\Delta}\rho + \frac{3}{2}\kappa e_3(\rho) + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega}\right) e_4\rho + \frac{3}{2}\rho(\underline{\kappa}\kappa + 2\rho) \\
= & -\square_{\mathbf{g}}\rho + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right) e_4\rho - \frac{1}{2}\kappa e_3\rho + 2\eta e_\theta\rho \\
& + \frac{3}{2}\kappa e_3(\rho) + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega}\right) e_4\rho + \frac{3}{2}\rho(\underline{\kappa}\kappa + 2\rho)
\end{aligned}$$

and hence

$$\begin{aligned}
\square_{\mathbf{g}}\rho & = \underline{\kappa}e_4\rho + \kappa e_3\rho + \frac{3}{2}(\underline{\kappa}\kappa + 2\rho)\rho \\
& + \frac{3}{2}\rho\left(-\frac{1}{2}\underline{\vartheta}\vartheta + 2(\underline{\xi}\xi + \eta\eta)\right) + \left(\frac{3}{2}\underline{\kappa} - 2\underline{\omega}\right)\left(\frac{1}{2}\underline{\vartheta}\alpha - \zeta\beta - 2(\underline{\eta}\beta + \xi\underline{\beta})\right) \\
& - \frac{1}{2}\underline{\vartheta}\underline{\phi}_2^*\beta + (\zeta - \eta)e_3\beta - \eta e_3(\Phi)\beta - \underline{\xi}(e_4\beta + e_4(\Phi)\beta) - \underline{\beta}\beta \\
& - e_3\left(-\frac{1}{2}\underline{\vartheta}\alpha + \zeta\beta + 2(\underline{\eta}\beta + \xi\underline{\beta})\right) \\
& - \underline{\phi}_1^*(\underline{\kappa})\beta + 2\underline{\phi}_1^*(\underline{\omega})\beta + 3\eta\underline{\phi}_1^*(\rho) - \underline{\phi}_1(-\vartheta\underline{\beta} + \underline{\xi}\alpha) - 2\eta e_\theta\rho.
\end{aligned}$$

Step 2. We derive the following, identity

$$\begin{aligned} \square_{\mathbf{g}}(r^2\rho) &= -Ae_3(r^2\rho) - \underline{A}e_4(r^2\rho) \\ &\quad + \frac{3}{2} \left(\frac{4}{3}A \frac{e_3(r)}{r} + \frac{4}{3}\underline{A} \frac{e_4(r)}{r} + \kappa\underline{\kappa} + 2\rho + \frac{2}{3r^2}\square_{\mathbf{g}}(r^2) \right) r^2\rho \\ &\quad + 4r \not{d}_1^*(r) \not{d}_1^*(\rho) + r^2\text{Err}[\square_{\mathbf{g}}\rho]. \end{aligned} \tag{B.1.1}$$

Proof. $r^2\rho$ satisfies the following wave equation

$$\square_{\mathbf{g}}(r^2\rho) = r^2\square_{\mathbf{g}}\rho + 2D^a(r^2)D_a(\rho) + \rho\square_{\mathbf{g}}(r^2).$$

On the other hand, recall that we have

$$\square_{\mathbf{g}}\rho = \underline{\kappa}e_4\rho + \kappa e_3\rho + \frac{3}{2}(\underline{\kappa}\kappa + 2\rho)\rho + \text{Err}[\square_{\mathbf{g}}\rho].$$

We deduce

$$\begin{aligned} \square_{\mathbf{g}}(r^2\rho) &= (r^2\kappa - e_4(r^2))e_3\rho + (r^2\underline{\kappa} - e_3(r^2))e_4\rho \\ &\quad + \frac{3}{2} \left(\kappa\underline{\kappa} + 2\rho + \frac{2}{3r^2}\square_{\mathbf{g}}(r^2) \right) r^2\rho + 4r \not{d}_1^*(r) \not{d}_1^*(\rho) + r^2\text{Err}[\square_{\mathbf{g}}\rho] \\ &= -Ae_3(r^2\rho) - \underline{A}e_4(r^2\rho) \\ &\quad + \frac{3}{2} \left(\frac{4}{3}A \frac{e_3(r)}{r} + \frac{4}{3}\underline{A} \frac{e_4(r)}{r} + \kappa\underline{\kappa} + 2\rho + \frac{2}{3r^2}\square_{\mathbf{g}}(r^2) \right) r^2\rho + 4r \not{d}_1^*(r) \not{d}_1^*(\rho) \\ &\quad + r^2\text{Err}[\square_{\mathbf{g}}\rho] \end{aligned}$$

as desired. □

Step 3. We now derive the desired formula for $\square_{\mathbf{g}}\tilde{\rho}$. In view of the definition of $\tilde{\rho}$, we have

$$\begin{aligned} \square_{\mathbf{g}}(\tilde{\rho}) &= \square_{\mathbf{g}}(r^2\rho) + \square_{\mathbf{g}}\left(\frac{2m}{r}\right) \\ &= \square_{\mathbf{g}}(r^2\rho) + 2m\square_{\mathbf{g}}\left(\frac{1}{r}\right) + 4D^a(m)D_a\left(\frac{1}{r}\right) + \frac{2}{r}\square_{\mathbf{g}}(m). \end{aligned}$$

Together with B.1.1 we deduce,

$$\begin{aligned} \square_{\mathbf{g}}(\tilde{\rho}) &= -Ae_3(r^2\rho) - \underline{A}e_4(r^2\rho) \\ &\quad + \frac{3}{2} \left(\frac{4}{3}A \frac{e_3(r)}{r} + \frac{4}{3}\underline{A} \frac{e_4(r)}{r} + \kappa\underline{\kappa} + 2\rho + \frac{2}{3r^2}\square_{\mathbf{g}}(r^2) \right) r^2\rho + 4r \not{d}_1^*(r) \not{d}_1^*(\rho) \\ &\quad + 2m\square_{\mathbf{g}}\left(\frac{1}{r}\right) + 4D^a(m)D_a\left(\frac{1}{r}\right) + \frac{2}{r}\square_{\mathbf{g}}(m) + 4r \not{d}_1^*(r) \not{d}_1^*(\rho) + r^2\text{Err}[\square_{\mathbf{g}}\rho]. \end{aligned}$$

Next, we use $r^2\rho = \tilde{\rho} - 2mr^{-1}$. This yields

$$\begin{aligned}
& \square_{\mathbf{g}}(\tilde{\rho}) - \frac{3}{2} \left(\kappa \underline{\kappa} - \frac{8m}{r^3} + \frac{2}{3r^2} \square_{\mathbf{g}}(r^2) \right) \tilde{\rho} \\
= & 2m \square_{\mathbf{g}} \left(\frac{1}{r} \right) - \frac{3m}{r} \kappa \underline{\kappa} + \frac{12m^2}{r^4} - \frac{2m}{r^3} \square_{\mathbf{g}}(r^2) \\
& - 6m \underline{A} \frac{e_3(r)}{r^2} - 6m \underline{A} \frac{e_4(r)}{r^2} \\
& + \frac{3}{r^2} \tilde{\rho}^2 + \frac{3}{2} \left(\frac{4}{3} \underline{A} \frac{e_3(r)}{r} + \frac{4}{3} \underline{A} \frac{e_4(r)}{r} \right) \tilde{\rho} \\
& - \underline{A} e_3(\tilde{\rho}) - \underline{A} e_4(\tilde{\rho}) + \frac{2}{r} \underline{A} e_3(m) + \frac{2}{r} \underline{A} e_4(m) \\
& + 4D^a(m) D_a \left(\frac{1}{r} \right) + \frac{2}{r} \square_{\mathbf{g}}(m) + 4r \not{d}_1^*(r) \not{d}_1^*(\rho) + r^2 \text{Err}[\square_{\mathbf{g}}\rho].
\end{aligned}$$

Note that in Schwarzschild,

$$\frac{3}{2} \left(\kappa \underline{\kappa} - \frac{8m}{r^3} + \frac{2}{3r^2} \square_{\mathbf{g}}(r^2) \right) = -\frac{8m}{r^3}$$

and hence

$$\begin{aligned}
& \square_{\mathbf{g}}(\tilde{\rho}) + \frac{8m}{r^3} \tilde{\rho} \\
= & 2m \square_{\mathbf{g}} \left(\frac{1}{r} \right) - \frac{3m}{r} \kappa \underline{\kappa} + \frac{12m^2}{r^4} - \frac{2m}{r^3} \square_{\mathbf{g}}(r^2) \\
& - 6m \frac{e_3(r)}{r^2} - 6m \underline{A} \frac{e_4(r)}{r^2} \\
& + \frac{3}{r^2} \tilde{\rho}^2 + \frac{3}{2} \left(\frac{4}{3} \underline{A} \frac{e_3(r)}{r} + \frac{4}{3} \underline{A} \frac{e_4(r)}{r} \right) \tilde{\rho} + \left(\frac{3}{2} \left(\kappa \underline{\kappa} - \frac{8m}{r^3} + \frac{2}{3r^2} \square_{\mathbf{g}}(r^2) \right) + \frac{8m}{r^3} \right) \tilde{\rho} \\
& - \underline{A} e_3(\tilde{\rho}) - \underline{A} e_4(\tilde{\rho}) + \frac{2}{r} \underline{A} e_3(m) + \frac{2}{r} \underline{A} e_4(m) \\
& + 4D^a(m) D_a \left(\frac{1}{r} \right) + \frac{2}{r} \square_{\mathbf{g}}(m) + 4r \not{d}_1^*(r) \not{d}_1^*(\rho) + r^2 \text{Err}[\square_{\mathbf{g}}\rho].
\end{aligned}$$

Also, we have

$$\begin{aligned}
\square_{\mathbf{g}} \left(\frac{1}{r} \right) - \frac{1}{r^3} \square_{\mathbf{g}}(r^2) &= -\frac{\square_{\mathbf{g}}(r)}{r^2} + 2 \frac{\mathbf{D}^\alpha(r) \mathbf{D}_\alpha(r)}{r^3} - 2 \frac{\square_{\mathbf{g}}(r)}{r^2} - 2 \frac{\mathbf{D}^\alpha(r) \mathbf{D}_\alpha(r)}{r^3} \\
&= -3 \frac{\square_{\mathbf{g}}(r)}{r^2}
\end{aligned}$$

and hence

$$\begin{aligned} & \square_{\mathbf{g}}(\tilde{\rho}) + \frac{8m}{r^3}\tilde{\rho} - 6m\frac{\square_{\mathbf{g}}(r)}{r^2} - \frac{3m}{r}\kappa\underline{\kappa} + \frac{12m^2}{r^4} \\ & - 6m\frac{e_3(r)}{r^2} - 6m\underline{A}\frac{e_4(r)}{r^2} + \frac{3}{r^2}\tilde{\rho}^2 + \frac{3}{2}\left(\frac{4}{3}\underline{A}\frac{e_3(r)}{r} + \frac{4}{3}\underline{A}\frac{e_4(r)}{r}\right)\tilde{\rho} \\ & + \left(\frac{3}{2}\left(\kappa\underline{\kappa} - \frac{8m}{r^3} + \frac{2}{3r^2}\square_{\mathbf{g}}(r^2)\right) + \frac{8m}{r^3}\right)\tilde{\rho} \\ & - Ae_3(\tilde{\rho}) - \underline{A}e_4(\tilde{\rho}) + \frac{2}{r}Ae_3(m) + \frac{2}{r}\underline{A}e_4(m) \\ & + 4D^a(m)D_a\left(\frac{1}{r}\right) + \frac{2}{r}\square_{\mathbf{g}}(m) + 4r\mathcal{d}_1^*(r)\mathcal{d}_1^*(\rho) + r^2\text{Err}[\square_{\mathbf{g}}\rho]. \end{aligned}$$

Finally, since

$$-6m\frac{e_3(r)}{r^2} - 6m\underline{A}\frac{e_4(r)}{r^2} = -3m\underline{A}\frac{\kappa}{r} - 3m\underline{A}\frac{\kappa}{r} - \frac{6m}{r}\underline{A}\underline{A}$$

and

$$-6m\frac{\square_{\mathbf{g}}(r)}{r^2} - \frac{3m}{r}\kappa\underline{\kappa} + \frac{12m^2}{r^4} = -6m\frac{\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2}\right)}{r^2} - \frac{3m}{r}\left(\kappa\underline{\kappa} + \frac{4\Upsilon}{r^2}\right),$$

we obtain

$$\begin{aligned} \square_{\mathbf{g}}(\tilde{\rho}) + \frac{8m}{r^3}\tilde{\rho} &= -6m\frac{\square_{\mathbf{g}}(r) - \left(\frac{2}{r} - \frac{2m}{r^2}\right)}{r^2} - \frac{3m}{r}\left(\kappa\underline{\kappa} + \frac{4\Upsilon}{r^2}\right) \\ & - 3m\underline{A}\frac{\kappa}{r} - 3m\underline{A}\frac{\kappa}{r} - \frac{6m}{r}\underline{A}\underline{A} \\ & + \frac{3}{r^2}\tilde{\rho}^2 + \frac{3}{2}\left(\frac{4}{3}\underline{A}\frac{e_3(r)}{r} + \frac{4}{3}\underline{A}\frac{e_4(r)}{r}\right)\tilde{\rho} \\ & + \left(\frac{3}{2}\left(\kappa\underline{\kappa} - \frac{8m}{r^3} + \frac{2}{3r^2}\square_{\mathbf{g}}(r^2)\right) + \frac{8m}{r^3}\right)\tilde{\rho} \\ & - Ae_3(\tilde{\rho}) - \underline{A}e_4(\tilde{\rho}) + \frac{2}{r}Ae_3(m) + \frac{2}{r}\underline{A}e_4(m) \\ & + 4D^a(m)D_a\left(\frac{1}{r}\right) + \frac{2}{r}\square_{\mathbf{g}}(m) + 4r\mathcal{d}_1^*(r)\mathcal{d}_1^*(\rho) + r^2\text{Err}[\square_{\mathbf{g}}\rho]. \end{aligned}$$

□

Appendix C

APPENDIX TO CHAPTER 9

C.1 Proof of Lemma 9.2.6

We start with the following

Lemma C.1.1. *Let $k \geq 0$ an integer and let $f \in \mathfrak{s}_k(\mathbf{S})$. Then, we have*

$$\begin{aligned}
 (\phi_k^{\mathbf{S}} f)^{\#} &= \frac{\sqrt{\gamma}}{\sqrt{\gamma^{\mathbf{S}} \#}} \left\{ \overset{\circ}{\phi}_k(f^{\#}) + \left(\frac{k}{2} U \int_0^1 \left(\sqrt{\gamma}(e_{\theta}(\kappa) - e_{\theta}(\vartheta)) \right)^{\# \lambda} d\lambda \right. \right. \\
 &\quad + \frac{k}{4} S \int_0^1 \left(\sqrt{\gamma} e_{\theta}(\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - 2\underline{b}\gamma^{1/2} e_{\theta} \Phi) \right)^{\# \lambda} d\lambda \\
 &\quad \left. \left. + \frac{k}{4} \left(\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - 2\underline{b}\gamma^{1/2} e_{\theta} \Phi \right)^{\#} U' + \frac{k}{2} (\kappa + \vartheta)^{\#} S' \right) f^{\#} \right\}
 \end{aligned}$$

where for $0 \leq \lambda \leq 1$, $\#_{\lambda}$ denotes the pull back by

$$\psi_{\lambda}(\overset{\circ}{u}, \overset{\circ}{s}, \theta) := (\overset{\circ}{u} + \lambda U(\theta), \overset{\circ}{s} + \lambda S(\theta), \theta).$$

Proof. For $p \in \overset{\circ}{S}$ and f a \mathbf{Z} -invariant scalar function on S , we have by definition of the push forward of a vectorfield

$$[\Psi_{\#}(\partial_{\theta})f]_{\Psi(p)} = [\partial_{\theta}(f \circ \Psi)]_p.$$

We infer

$$(\phi_k^{\mathbf{S}} f)^{\#} = \frac{1}{\sqrt{\gamma^{\mathbf{S}} \#}} \left(\partial_{\theta}(f^{\#}) + k \partial_{\theta}(\Phi^{\#}) f^{\#} \right)$$

and hence

$$\begin{aligned} (\not{d}_k^{\mathbf{S}} f)^{\#} &= \frac{\sqrt{\gamma}}{\sqrt{\gamma^{\mathbf{S}} \#}} \left(e_{\theta}(f^{\#}) + k e_{\theta}(\Phi) f^{\#} + k(e_{\theta}(\Phi^{\#}) - e_{\theta}(\Phi)) f^{\#} \right) \\ &= \frac{\sqrt{\gamma}}{\sqrt{\gamma^{\mathbf{S}} \#}} \left(\overset{\circ}{d}_k(f^{\#}) + k(e_{\theta}(\Phi^{\#}) - e_{\theta}(\Phi)) f^{\#} \right). \end{aligned}$$

Next, we have

$$e_{\theta}(\Phi^{\#}) - e_{\theta}(\Phi) = \sqrt{\gamma}^{-1} \left(\partial_{\theta}(\Phi^{\#}) - \partial_{\theta}\Phi \right)$$

and

$$\begin{aligned} & \left(\partial_{\theta}(\Phi^{\#}) - \partial_{\theta}\Phi \right) (\overset{\circ}{u}, \overset{\circ}{s}, \theta) \\ &= \partial_{\theta}[\Phi(\overset{\circ}{u} + U(\theta), \overset{\circ}{s} + S(\theta), \theta)] - \partial_{\theta}\Phi(\overset{\circ}{u}, \overset{\circ}{s}, \theta) \\ &= (\partial_{\theta}\Phi)(\overset{\circ}{u} + U(\theta), \overset{\circ}{s} + S(\theta), \theta) - \partial_{\theta}\Phi(\overset{\circ}{u}, \overset{\circ}{s}, \theta) + \left[(\partial_u\Phi)^{\#}U' + (\partial_s\Phi)^{\#}S' \right] (\overset{\circ}{u}, \overset{\circ}{s}, \theta) \\ &= \int_0^1 \frac{d}{d\lambda} \left[(\partial_{\theta}\Phi)(\overset{\circ}{u} + \lambda U(\theta), \overset{\circ}{s} + \lambda S(\theta), \theta) \right] d\lambda + \left[(\partial_u\Phi)^{\#}U' + (\partial_s\Phi)^{\#}S' \right] (\overset{\circ}{u}, \overset{\circ}{s}, \theta) \\ &= U(\theta) \int_0^1 (\partial_u\partial_{\theta}\Phi)(\overset{\circ}{u} + \lambda U(\theta), \overset{\circ}{s} + \lambda S(\theta), \theta) d\lambda \\ & \quad + S(\theta) \int_0^1 (\partial_s\partial_{\theta}\Phi)(\overset{\circ}{u} + \lambda U(\theta), \overset{\circ}{s} + \lambda S(\theta), \theta) d\lambda + \left[(\partial_u\Phi)^{\#}U' + (\partial_s\Phi)^{\#}S' \right] (\overset{\circ}{u}, \overset{\circ}{s}, \theta) \end{aligned}$$

which we rewrite

$$\partial_{\theta}(\Phi^{\#}) - \partial_{\theta}\Phi = U \int_0^1 (\partial_u\partial_{\theta}\Phi)^{\#\lambda} d\lambda + S \int_0^1 (\partial_s\partial_{\theta}\Phi)^{\#\lambda} d\lambda + (\partial_u\Phi)^{\#}U' + (\partial_s\Phi)^{\#}S'$$

where $\#_{\lambda}$ denotes the pull back by the map $\psi_{\lambda}(\overset{\circ}{u}, \overset{\circ}{s}, \theta) = (\overset{\circ}{u} + \lambda U(\theta), \overset{\circ}{s} + \lambda S(\theta), \theta)$.

Next, recall that,

$$\partial_s = e_4, \quad \partial_u = \frac{1}{2} (e_3 - \underline{\Omega}e_4 - \underline{b}\gamma^{1/2}e_{\theta}), \quad \partial_{\theta} = \sqrt{\gamma}e_{\theta}.$$

Hence,

$$\begin{aligned} \partial_{\theta}\partial_s\Phi &= \sqrt{\gamma}e_{\theta}e_4(\Phi) \\ \partial_{\theta}\partial_u\Phi &= \frac{1}{2}\sqrt{\gamma}e_{\theta} (e_3\Phi - \underline{\Omega}e_4\Phi - \underline{b}\gamma^{1/2}e_{\theta}\Phi) \end{aligned}$$

which yields

$$\begin{aligned}
 \partial_\theta(\Phi^\#) - \partial_\theta\Phi &= U \int_0^1 \left(\sqrt{\gamma} e_\theta e_4(\Phi) \right)^{\#\lambda} d\lambda \\
 &\quad + S \int_0^1 \left(\frac{1}{2} \sqrt{\gamma} e_\theta (e_3\Phi - \underline{\Omega}e_4\Phi - \underline{b}\gamma^{1/2}e_\theta\Phi) \right)^{\#\lambda} d\lambda \\
 &\quad + \frac{1}{2} \left(e_3\Phi - \underline{\Omega}e_4\Phi - \underline{b}\gamma^{1/2}e_\theta\Phi \right)^\# U' + (e_4\Phi)^\# S' \\
 &= \frac{1}{2} U \int_0^1 \left(\sqrt{\gamma} (e_\theta(\kappa) - e_\theta(\vartheta)) \right)^{\#\lambda} d\lambda \\
 &\quad + \frac{1}{4} S \int_0^1 \left(\sqrt{\gamma} e_\theta (\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - 2\underline{b}\gamma^{1/2}e_\theta\Phi) \right)^{\#\lambda} d\lambda \\
 &\quad + \frac{1}{4} \left(\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - 2\underline{b}\gamma^{1/2}e_\theta\Phi \right)^\# U' + \frac{1}{2} (\kappa + \vartheta)^\# S'.
 \end{aligned}$$

We deduce

$$\begin{aligned}
 (\mathfrak{d}_k^{\mathbf{S}} f)^\# &= \frac{\sqrt{\gamma}}{\sqrt{\gamma^{\mathbf{S}}\#}} \left\{ \mathring{\mathfrak{d}}_k(f^\#) + \left(\frac{k}{2} U \int_0^1 \left(\sqrt{\gamma} (e_\theta(\kappa) - e_\theta(\vartheta)) \right)^{\#\lambda} d\lambda \right. \right. \\
 &\quad \left. \left. + \frac{k}{4} S \int_0^1 \left(\sqrt{\gamma} e_\theta (\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - 2\underline{b}\gamma^{1/2}e_\theta\Phi) \right)^{\#\lambda} d\lambda \right. \right. \\
 &\quad \left. \left. + \frac{k}{4} \left(\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - 2\underline{b}\gamma^{1/2}e_\theta\Phi \right)^\# U' + \frac{k}{2} (\kappa + \vartheta)^\# S' \right) f^\# \right\}.
 \end{aligned}$$

This concludes the proof of the lemma. □

We are ready to prove the higher derivative comparison Lemma 9.2.6 which we recall below.

Lemma C.1.2. *Let $\mathring{S} \subset \mathcal{R} = \mathcal{R}(\mathring{\epsilon}, \mathring{\delta})$ as in Definition 9.1.1 verifying the assumptions **A1-A3**. Let $\Psi : \mathring{S} \rightarrow \mathbf{S}$ be \mathbf{Z} -invariant deformation. Assume the bound*

$$\|(U', S')\|_{L^\infty(\mathring{S})} + \mathring{r}^{-1} \max_{0 \leq s \leq s_{max}} \|(U', S')\|_{\mathfrak{h}_s(\mathring{S}, \mathring{\mathfrak{g}})} \lesssim \mathring{\delta}. \tag{C.1.1}$$

Then, we have for any reduced scalar h defined on \mathcal{R}

$$\|h\|_{\mathfrak{h}_s(\mathbf{S})} \lesssim \sup_{\mathcal{R}} |\mathfrak{d}^{\leq k} h| \quad \text{for } 0 \leq s \leq s_{max}.$$

Also, if $f \in \mathfrak{h}_s(\mathbf{S})$ and $f^\#$ is its pull-back by ψ , we have

$$\|f\|_{\mathfrak{h}_s(\mathbf{S})} = \|f^\#\|_{\mathfrak{h}_s(\mathring{S}, \mathring{\mathfrak{g}}^{\mathbf{S}, \#})} = \|f^\#\|_{\mathfrak{h}_s(\mathring{S}, \mathring{\mathfrak{g}})} (1 + O(\mathring{\epsilon})) \quad \text{for } 0 \leq s \leq s_{max}.$$

Remark C.1.3. Note that the estimates of the lemma are independent of the size $\overset{\circ}{r}$ of the sphere $\overset{\circ}{S} = S(\overset{\circ}{u}, \overset{\circ}{s}) \subset \mathcal{R}$, see Definition 9.1.1. To simplify the argument below we assume $\overset{\circ}{r} \approx 1$. The general case can be easily deduced by a simple scaling argument or making obvious adjustments in the inequalities below.

Proof. We argue by iteration. We consider the following iteration assumptions

$$\text{If (9.2.14) holds, then we have } \|h\|_{\mathfrak{h}_s(\mathbf{s})} \lesssim \sup_{\mathcal{R}} |\mathfrak{d}^{\leq s} h|, \quad (\text{C.1.2})$$

and

$$\text{If (9.2.14) holds, then we have } \|f^\#\|_{\mathfrak{h}_s(\overset{\circ}{S}, \overset{\circ}{g}^{\mathbf{s}, \#})} = \|f^\#\|_{\mathfrak{h}_s(\overset{\circ}{S}, \overset{\circ}{g})} (1 + O(\delta)). \quad (\text{C.1.3})$$

First, note that (C.1.2) holds trivially for $s = 0$ and (C.1.3) holds for $s = 0$ by Lemma 9.2.3. Thus, from now on, we assume that (C.1.2) holds for some s with $0 \leq s \leq s_{max} - 1$ and that (C.1.3) holds for some s with $0 \leq s \leq s_{max} - 3$, and our goal is to prove that it also holds for s replaced by $s + 1$.

We start with (C.1.2). We have

$$\mathfrak{d}_k^{\mathbf{s}} h = e_\theta^{\mathbf{s}} h + e_\theta^{\mathbf{s}}(\Phi)h.$$

Now, recall that we have

$$e_\theta^{\mathbf{s}} = \frac{1}{\sqrt{\gamma^{\mathbf{s}}}} \partial_\theta^{\mathbf{s}}, \quad \partial_\theta^{\mathbf{s}}|_{\Psi(p)} = \left(\left(S' - \frac{1}{2} \underline{\Omega} U' \right) e_4 + \frac{1}{2} U' e_3 + \sqrt{\gamma} \left(1 - \frac{1}{2} \underline{b} U' \right) e_\theta \right) \Big|_{\Psi(p)}.$$

This yields

$$\begin{aligned} (\mathfrak{d}_k^{\mathbf{s}} h)|_{\Psi(p)} &= \left\{ \frac{1}{\sqrt{\gamma^{\mathbf{s}}}} \left(\left(S' - \frac{1}{2} \underline{\Omega} U' \right) e_4(h) + \left(S' - \frac{1}{2} \underline{\Omega} U' \right) e_4(\Phi)h \right. \right. \\ &\quad \left. \left. + \frac{1}{2} U' e_3(h) + \frac{1}{2} U' e_3(\Phi)h + \sqrt{\gamma} \left(1 - \frac{1}{2} \underline{b} U' \right) \mathfrak{d}_k(h) \right) \right\} \Big|_{\Psi(p)}. \end{aligned}$$

Together with the iteration assumption (C.1.3), we infer

$$\begin{aligned} \|\mathfrak{d}_k^{\mathbf{s}} h\|_{\mathfrak{h}_s(\mathbf{s})} &= \|(\mathfrak{d}_k^{\mathbf{s}} h)^\#\|_{\mathfrak{h}_s(\overset{\circ}{S}, \overset{\circ}{g})} (1 + O(\delta)) \\ &\lesssim \left\| \frac{1}{\sqrt{\gamma^{\mathbf{s}, \#}}} \left(\left(S' - \frac{1}{2} \underline{\Omega}^\# U' \right) (e_4(h))^\# + \left(S' - \frac{1}{2} \underline{\Omega}^\# U' \right) (e_4(\Phi)h)^\# \right. \right. \\ &\quad \left. \left. + \frac{1}{2} U' (e_3(h))^\# + \frac{1}{2} U' (e_3(\Phi)h)^\# + \sqrt{\gamma^\#} \left(1 - \frac{1}{2} \underline{b}^\# U' \right) (\mathfrak{d}_k(h))^\# \right) \right\|_{\mathfrak{h}_s(\overset{\circ}{S}, \overset{\circ}{g})} \end{aligned}$$

i.e.,

$$\begin{aligned} \|\mathcal{A}_k^{\mathbf{S}} h\|_{\mathfrak{h}_s(\mathbf{S})} &\lesssim \|(\mathcal{A}_k(h))^{\#}\|_{\mathfrak{h}_s(\dot{S}, \dot{g})} + \left\| \left(\frac{\sqrt{\gamma^{\#}}}{\sqrt{\gamma^{\mathbf{S}, \#}}} - 1 \right) (\mathcal{A}_k(h))^{\#} \right\|_{\mathfrak{h}_s(\dot{S}, \dot{g})} \\ &\quad + \left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}, \#}}} \left(\left(S' - \frac{1}{2} \underline{\Omega}^{\#} U' \right) (e_4(h))^{\#} + \left(S' - \frac{1}{2} \underline{\Omega}^{\#} U' \right) (e_4(\Phi)h)^{\#} \right. \right. \\ &\quad \left. \left. + \frac{1}{2} U'(e_3(h))^{\#} + \frac{1}{2} U'(e_3(\Phi)h)^{\#} - \frac{1}{2} \sqrt{\gamma^{\#}} \underline{b}^{\#} U'(\mathcal{A}_k(h))^{\#} \right) \right\|_{\mathfrak{h}_s(\dot{S}, \dot{g})}. \end{aligned}$$

Together with a non sharp product rule in $\mathfrak{h}_s(\dot{S}, \dot{g})$ and the repeated use of the iteration assumptions (C.1.2) (C.1.3), we can bound the right hand side of the above inequality by

$$\begin{aligned} &\lesssim \left(1 + \left(\|\sqrt{\gamma}\|_{\mathfrak{h}_s(\mathbf{S})} + \|(\sqrt{\gamma})^{\#}\|_{\mathfrak{h}_1^{\infty}(\dot{S})} \right) \left(\left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}}}} \right\|_{\mathfrak{h}_s(\mathbf{S})} + \left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}, \#}}} \right\|_{\mathfrak{h}_1^{\infty}(\dot{S})} \right) \right) \|\mathcal{A}_k(h)\|_{\mathfrak{h}_s(\mathbf{S})} \\ &\quad + \|(U', S')\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^{\infty}(\dot{S})} \left(\left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}}}} \right\|_{\mathfrak{h}_s(\mathbf{S})} + \left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}, \#}}} \right\|_{\mathfrak{h}_1^{\infty}(\dot{S})} \right) \\ &\quad \times \left(1 + \|(\underline{\Omega}, \underline{b}\sqrt{\gamma})\|_{\mathfrak{h}_s(\mathbf{S})} + \|(\underline{\Omega}, \underline{b}\sqrt{\gamma})^{\#}\|_{\mathfrak{h}_1^{\infty}(\dot{S})} \right) \left\| (e_3, e_4, \mathcal{A}_k)h, e_3(\Phi)h, e_4(\Phi)h \right\|_{\mathfrak{h}_s(\mathbf{S})} \end{aligned}$$

Therefore $\|\mathcal{A}_k^{\mathbf{S}} h\|_{\mathfrak{h}_s(\mathbf{S})}$ can be bounded by

$$\begin{aligned} &\lesssim \left(1 + \left(\|\sqrt{\gamma}\|_{\mathfrak{h}_s(\mathbf{S})} + \|(\sqrt{\gamma})^{\#}\|_{\mathfrak{h}_1^{\infty}(\dot{S})} \right) \left(\left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}}}} \right\|_{\mathfrak{h}_s(\mathbf{S})} + \left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}, \#}}} \right\|_{\mathfrak{h}_1^{\infty}(\dot{S})} \right) \right) \sup_{\mathcal{R}} |\mathfrak{d}^{\leq s} \mathcal{A}_k h| \\ &\quad + \left(\left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}}}} \right\|_{\mathfrak{h}_s(\mathbf{S})} + \left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}, \#}}} \right\|_{\mathfrak{h}_1^{\infty}(\dot{S})} \right) \left(1 + \|(\underline{\Omega}, \underline{b}\sqrt{\gamma})\|_{\mathfrak{h}_s(\mathbf{S})} + \|(\underline{\Omega}, \underline{b}\sqrt{\gamma})^{\#}\|_{\mathfrak{h}_1^{\infty}(\dot{S})} \right) \\ &\quad \times \sup_{\mathcal{R}} |\mathfrak{d}^{\leq s} (\mathfrak{d}h, e_3(\Phi)h, e_4(\Phi)h)|, \end{aligned}$$

where we used in the last inequality the assumption (9.2.14) on (U', S') . Together with

(9.1.12) and (9.1.15), we infer

$$\begin{aligned} \|\mathcal{A}_k^{\mathbf{S}} h\|_{\mathfrak{h}_s(\mathbf{S})} &\lesssim \left\{ \left(1 + \left(1 + \|(\sqrt{\gamma})^\#\|_{\mathfrak{h}_1^\infty(\dot{S})} \right) \left(\left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}}}} \right\|_{\mathfrak{h}_s(\mathbf{S})} + \left\| \frac{1}{\sqrt{\gamma^{\mathbf{S},\#}} } \right\|_{\mathfrak{h}_1^\infty(\dot{S})} \right) \right) \right. \\ &\quad \left. + \left(1 + \left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}}}} \right\|_{\mathfrak{h}_s(\mathbf{S})} + \left\| \frac{1}{\sqrt{\gamma^{\mathbf{S},\#}} } \right\|_{\mathfrak{h}_1^\infty(\dot{S})} \right) \left(1 + \|(\underline{\Omega}, \underline{b}\sqrt{\gamma})^\#\|_{\mathfrak{h}_1^\infty(\dot{S})} \right) \right\} \\ &\quad \times \sup_{\mathcal{R}} |\mathfrak{d}^{\leq s+1} h|. \end{aligned}$$

Also, for a reduced scalar v defined on \mathcal{R} , we have in view of the assumption (9.2.14) on (U', S')

$$\begin{aligned} \|v^\#\|_{\mathfrak{h}_1^\infty(\dot{S})} &= \|v \circ \psi\|_{\mathfrak{h}_1^\infty(\dot{S})} \\ &\lesssim \left(1 + \sup_{0 \leq \theta \leq \pi} |\psi'(\theta)| \right) \sup_{\mathcal{R}} |\mathfrak{d}^{\leq 1} v| \\ &\lesssim \left(1 + \|(U', S')\|_{\mathfrak{h}_1^\infty(\dot{S})} \right) \sup_{\mathcal{R}} |\mathfrak{d}^{\leq 1} v| \\ &\lesssim (1 + \delta) \sup_{\mathcal{R}} |\mathfrak{d}^{\leq 1} v|. \end{aligned} \tag{C.1.4}$$

Together with (9.1.12) and (9.1.15), we infer

$$\|\mathcal{A}_k^{\mathbf{S}} h\|_{\mathfrak{h}_s(\mathbf{S})} \lesssim \left\{ 1 + \left\| \frac{1}{\sqrt{\gamma^{\mathbf{S}}}} \right\|_{\mathfrak{h}_s(\mathbf{S})} + \left\| \frac{1}{\sqrt{\gamma^{\mathbf{S},\#}} } \right\|_{\mathfrak{h}_1^\infty(\dot{S})} \right\} \sup_{\mathcal{R}} |\mathfrak{d}^{\leq s+1} h|.$$

Now, recall that

$$\begin{aligned} \gamma^{\mathbf{S}}(\psi(\theta)) &= \gamma(\psi(\theta)) + \left(\underline{\Omega}(\psi(\theta)) + \frac{1}{4}(b(\psi(\theta)))^2 \gamma(\psi(\theta)) \right) (U'(\theta))^2 - 2U'(\theta)S'(\theta) \\ &\quad - \gamma(\psi(\theta))\underline{b}(\psi(\theta))U'(\theta). \end{aligned}$$

Together with a repeated application of the iteration assumptions and a non sharp product rule in $\mathfrak{h}_s(\dot{S}, \dot{g})$ and (C.1.4), this yields

$$\begin{aligned} &\|\gamma^{\mathbf{S}}\|_{\mathfrak{h}_s(\mathbf{S})} + \|\gamma^{\mathbf{S},\#}\|_{\mathfrak{h}_1^\infty(\dot{S})} \\ &\lesssim \left(1 + \sup_{\mathcal{R}} |\mathfrak{d}^{\leq 1}(\gamma, \underline{\Omega}, \underline{b}^2 \gamma, \underline{b}\gamma)| + \sup_{\mathcal{R}} |\mathfrak{d}^{\leq s}(\gamma, \underline{\Omega}, \underline{b}^2 \gamma, \underline{b}\gamma)| \right) \\ &\quad \times \left(1 + \|(U', S')\|_{\mathfrak{h}_1^\infty(\dot{S})} + \|(U', S')\|_{\mathfrak{h}_s(\dot{S}, \dot{g})} \right) \\ &\lesssim 1 \end{aligned}$$

where we used in the last estimate the assumption (9.2.14) on (U', S') and (9.1.15). We infer

$$\left\| \frac{1}{\sqrt{\gamma \mathbf{S}}} \right\|_{\mathfrak{h}_s(\mathbf{S})} + \left\| \frac{1}{\sqrt{\gamma \mathbf{S}, \#}} \right\|_{\mathfrak{h}_1^\infty(\mathring{S})} \lesssim 1$$

and hence

$$\| \phi_k^{\mathbf{S}} h \|_{\mathfrak{h}_s(\mathbf{S})} \lesssim \sup_{\mathcal{R}} | \mathfrak{d}^{\leq s+1} h |$$

which corresponds to the first of our iteration assumption (C.1.2) with s replaced with $s + 1$ for $s \leq s_{max} - 1$.

Next, we focus on recovering the second iteration assumption (C.1.3) with s replaced with $s + 1$ for $s \leq s_{max} - 3$. Recall from Lemma C.1.1 that we have for $f \in \mathfrak{s}_k(\mathbf{S})$

$$\begin{aligned} (\phi_k^{\mathbf{S}} f)^\# &= \frac{\sqrt{\gamma}}{\sqrt{\gamma \mathbf{S}, \#}} \left\{ \mathring{\phi}_k(f^\#) + \left(\frac{k}{2} U \int_0^1 (\sqrt{\gamma}(e_\theta(\kappa) - e_\theta(\vartheta)))^{\#\lambda} d\lambda \right. \right. \\ &\quad + \frac{k}{4} S \int_0^1 (\sqrt{\gamma} e_\theta(\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - 2\underline{b}\gamma^{1/2} e_\theta \Phi))^{\#\lambda} d\lambda \\ &\quad \left. \left. + \frac{k}{4} (\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - 2\underline{b}\gamma^{1/2} e_\theta \Phi)^\# U' + \frac{k}{2} (\kappa + \vartheta)^\# S' \right) f^\# \right\} \end{aligned}$$

where for $0 \leq \lambda \leq 1$, $\#_\lambda$ denotes the pull back by

$$\psi_\lambda(\mathring{u}, \mathring{s}, \theta) = (\mathring{u} + \lambda U(\theta), \mathring{s} + \lambda S(\theta), \theta).$$

For convenience, we rewrite some of the terms as follows

$$\begin{aligned} e_\theta(\kappa) - e_\theta(\vartheta) &= -\phi_1^*(\kappa) - \frac{1}{2}(\phi_1 \vartheta - \phi_2^* \vartheta), \\ \underline{b}\gamma^{1/2} e_\theta \Phi &= \frac{1}{2}\gamma^{1/2}(\phi_1 \underline{b} + \phi_2^* \underline{b}), \end{aligned}$$

and

$$\begin{aligned}
& e_\theta (\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\underline{\kappa} - \underline{\vartheta}) - 2\underline{b}\gamma^{1/2}e_\theta\Phi) \\
= & -\mathcal{A}_1^*(\underline{\kappa}) - \frac{1}{2}(\mathcal{A}_1\underline{\vartheta} - \mathcal{A}_2^*\underline{\vartheta}) + \mathcal{A}_1^*(\underline{\Omega}\underline{\kappa}) - \mathcal{A}_1^*(\underline{\Omega})\underline{\vartheta} + \frac{1}{2}\underline{\Omega}(\mathcal{A}_1\underline{\vartheta} - \mathcal{A}_2^*\underline{\vartheta}) \\
& + \mathcal{A}_1^*(\gamma^{1/2})(\mathcal{A}_1\underline{\vartheta} + \mathcal{A}_2^*\underline{\vartheta})\underline{b} - 2\gamma^{1/2}e_\theta(\underline{b}e_\theta\Phi) \\
= & -\mathcal{A}_1^*(\underline{\kappa}) - \frac{1}{2}(\mathcal{A}_1\underline{\vartheta} - \mathcal{A}_2^*\underline{\vartheta}) + \mathcal{A}_1^*(\underline{\Omega}\underline{\kappa}) - \mathcal{A}_1^*(\underline{\Omega})\underline{\vartheta} + \frac{1}{2}\underline{\Omega}(\mathcal{A}_1\underline{\vartheta} - \mathcal{A}_2^*\underline{\vartheta}) \\
& + \mathcal{A}_1^*(\gamma^{1/2})(\mathcal{A}_1\underline{\vartheta} + \mathcal{A}_2^*\underline{\vartheta})\underline{b} - 2\gamma^{1/2}(-e_\theta(\Phi)\mathcal{A}_2^*\underline{b} - K\underline{b}) \\
= & -\mathcal{A}_1^*(\underline{\kappa}) - \frac{1}{2}(\mathcal{A}_1\underline{\vartheta} - \mathcal{A}_2^*\underline{\vartheta}) + \mathcal{A}_1^*(\underline{\Omega}\underline{\kappa}) - \mathcal{A}_1^*(\underline{\Omega})\underline{\vartheta} + \frac{1}{2}\underline{\Omega}(\mathcal{A}_1\underline{\vartheta} - \mathcal{A}_2^*\underline{\vartheta}) \\
& + \mathcal{A}_1^*(\gamma^{1/2})(\mathcal{A}_1\underline{\vartheta} + \mathcal{A}_2^*\underline{\vartheta})\underline{b} + \frac{1}{2}\gamma^{1/2}(\mathcal{A}_2\mathcal{A}_2^*\underline{b} + \mathcal{A}_3^*\mathcal{A}_2^*\underline{b}) + 2\gamma^{1/2}K\underline{b}
\end{aligned}$$

where we used the identities

$$\begin{aligned}
e_\theta(e_\theta(\Phi)) &= -(e_\theta(\Phi))^2 - K, \\
2\gamma^{1/2}e_\theta\Phi\mathcal{A}_2^*\underline{b} &= \frac{1}{2}\gamma^{1/2}(\mathcal{A}_2\mathcal{A}_2^*\underline{b} + \mathcal{A}_3^*\mathcal{A}_2^*\underline{b}).
\end{aligned}$$

This yields

$$\begin{aligned}
(\mathcal{A}_k^{\mathbf{S}}f)^\# &= \frac{\sqrt{\gamma}}{\sqrt{\gamma^{\mathbf{S}}\#}} \left\{ \mathring{\mathcal{A}}_k(f^\#) + \left(\frac{k}{2}U \int_0^1 \left(\sqrt{\gamma} \left(-\mathcal{A}_1^*(\underline{\kappa}) - \frac{1}{2}(\mathcal{A}_1\underline{\vartheta} - \mathcal{A}_2^*\underline{\vartheta}) \right) \right)^{\#\lambda} d\lambda \right. \right. \\
& + \frac{k}{4}S \int_0^1 \left(\sqrt{\gamma} \left(-\mathcal{A}_1^*(\underline{\kappa}) - \frac{1}{2}(\mathcal{A}_1\underline{\vartheta} - \mathcal{A}_2^*\underline{\vartheta}) + \mathcal{A}_1^*(\underline{\Omega}\underline{\kappa}) - \mathcal{A}_1^*(\underline{\Omega})\underline{\vartheta} \right. \right. \\
& + \frac{1}{2}\underline{\Omega}(\mathcal{A}_1\underline{\vartheta} - \mathcal{A}_2^*\underline{\vartheta}) + \mathcal{A}_1^*(\gamma^{1/2})(\mathcal{A}_1\underline{\vartheta} + \mathcal{A}_2^*\underline{\vartheta})\underline{b} \\
& \left. \left. + \frac{1}{2}\gamma^{1/2}(\mathcal{A}_2\mathcal{A}_2^*\underline{b} + \mathcal{A}_3^*\mathcal{A}_2^*\underline{b}) + 2\gamma^{1/2}K\underline{b} \right) \right)^{\#\lambda} d\lambda \\
& \left. + \frac{k}{4} \left(\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\underline{\kappa} - \underline{\vartheta}) - \gamma^{1/2}(\mathcal{A}_1\underline{\vartheta} + \mathcal{A}_2^*\underline{\vartheta})\underline{b} \right)^\# U' + \frac{k}{2}(\underline{\kappa} + \underline{\vartheta})^\# S' \right) f^\# \}.
\end{aligned}$$

Next, we take the $\mathfrak{h}_s(\mathring{S}, \mathring{g})$ -norm of this identity, and we use the iteration assumption to

replace the norm on the left-hand side with the $\mathfrak{h}_s(\mathring{S}, \mathring{g}^{\mathbf{S}, \#})$ -norm. We infer

$$\begin{aligned} & \|(\mathring{d}_k^{\mathbf{S}} f)^\# \|_{\mathfrak{h}_s(\mathring{S}, \mathring{g}^{\mathbf{S}, \#})} (1 + O(\delta)) \\ = & \left\| \frac{\sqrt{\gamma}}{\sqrt{\gamma^{\mathbf{S}} \#}} \left\{ \mathring{d}_k(f^\#) + \left(\frac{k}{2} U \int_0^1 \left(\sqrt{\gamma} \left(-\mathring{d}_1^*(\kappa) - \frac{1}{2}(\mathring{d}_1\vartheta - \mathring{d}_2^*\vartheta) \right) \right)^{\#_\lambda} d\lambda \right. \right. \right. \\ & + \frac{k}{4} S \int_0^1 \left(\sqrt{\gamma} \left(-\mathring{d}_1^*(\underline{\kappa}) - \frac{1}{2}(\mathring{d}_1\underline{\vartheta} - \mathring{d}_2^*\underline{\vartheta}) + \mathring{d}_1^*(\underline{\Omega}\kappa) - \mathring{d}_1^*(\underline{\Omega})\vartheta \right. \right. \\ & + \frac{1}{2}\underline{\Omega}(\mathring{d}_1\underline{\vartheta} - \mathring{d}_2^*\underline{\vartheta}) + \mathring{d}_1^*(\gamma^{1/2})(\mathring{d}_1\vartheta + \mathring{d}_2^*\vartheta)\underline{b} \\ & \left. \left. \left. + \frac{1}{2}\gamma^{1/2}(\mathring{d}_2 \mathring{d}_2^*\underline{b} + \mathring{d}_3^* \mathring{d}_2^*\underline{b}) + 2\gamma^{1/2}K\underline{b} \right) \right)^{\#_\lambda} d\lambda \right. \\ & \left. \left. + \frac{k}{4} \left(\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - \gamma^{1/2}(\mathring{d}_1\vartheta + \mathring{d}_2^*\vartheta)\underline{b} \right)^\# U' + \frac{k}{2}(\kappa + \vartheta)^\# S' \right) f^\# \right\} \Big\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g}^\circ)}. \end{aligned}$$

Next, we use a non sharp product rule in $\mathfrak{h}_s(\mathring{S}, \mathring{g}^\circ)$ to infer

$$\begin{aligned} & \|(\mathring{d}_k^{\mathbf{S}} f)^\# \|_{\mathfrak{h}_s(\mathring{S}, \mathring{g}^{\mathbf{S}, \#})} (1 + O(\delta)) \\ = & \left(1 + O(1) \left\| \frac{\sqrt{\gamma}}{\sqrt{\gamma^{\mathbf{S}} \#}} - 1 \right\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g}^\circ) \cap \mathfrak{h}_1^\infty(\mathring{S})} \right) \left\{ \left\| \mathring{d}_k(f^\#) \right\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g}^\circ)} \right. \\ & + O(1) \left(\|U\|_{\mathfrak{h}_{s+1}(\mathring{S}, \mathring{g}^\circ)} \int_0^1 \left\| \left(\sqrt{\gamma} \left(-\mathring{d}_1^*(\kappa) - \frac{1}{2}(\mathring{d}_1\vartheta - \mathring{d}_2^*\vartheta) \right) \right)^{\#_\lambda} \right\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g}^\circ) \cap \mathfrak{h}_1^\infty(\mathring{S})} d\lambda \right. \\ & + \|S\|_{\mathfrak{h}_{s+1}(\mathring{S}, \mathring{g}^\circ)} \int_0^1 \left\| \left(\sqrt{\gamma} \left(-\mathring{d}_1^*(\underline{\kappa}) - \frac{1}{2}(\mathring{d}_1\underline{\vartheta} - \mathring{d}_2^*\underline{\vartheta}) + \mathring{d}_1^*(\underline{\Omega}\kappa) - \mathring{d}_1^*(\underline{\Omega})\vartheta \right. \right. \right. \\ & + \frac{1}{2}\underline{\Omega}(\mathring{d}_1\underline{\vartheta} - \mathring{d}_2^*\underline{\vartheta}) + \mathring{d}_1^*(\gamma^{1/2})(\mathring{d}_1\vartheta + \mathring{d}_2^*\vartheta)\underline{b} \\ & \left. \left. \left. + \frac{1}{2}\gamma^{1/2}(\mathring{d}_2 \mathring{d}_2^*\underline{b} + \mathring{d}_3^* \mathring{d}_2^*\underline{b}) + 2\gamma^{1/2}K\underline{b} \right) \right)^{\#_\lambda} \right\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g}^\circ) \cap \mathfrak{h}_1^\infty(\mathring{S})} d\lambda \\ & + \left\| \left(\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - \gamma^{1/2}(\mathring{d}_1\vartheta + \mathring{d}_2^*\vartheta)\underline{b} \right)^\# \right\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g}^\circ) \cap \mathfrak{h}_1^\infty(\mathring{S})} \|U'\|_{\mathfrak{h}_{s+1}(\mathring{S}, \mathring{g}^\circ)} \\ & \left. + \left\| (\kappa + \vartheta)^\# \right\|_{\mathfrak{h}_s(\mathring{S}, \mathring{g}^\circ) \cap \mathfrak{h}_1^\infty(\mathring{S})} \|S'\|_{\mathfrak{h}_{s+1}(\mathring{S}, \mathring{g}^\circ)} \right\} \|f^\# \|_{\mathfrak{h}_{s+1}(\mathring{S}, \mathring{g}^\circ)}. \end{aligned}$$

Since $s + 1 \leq s_{max} - 2$, we infer in view of (9.2.14) and the fact that $U(0) = S(0) = 0$,

$$\begin{aligned}
& \|(\not{d}_k^{\mathbf{S}} f)^\# \|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}}^{\mathbf{S}, \#})} (1 + O(\delta)) \\
&= \left(1 + O(1) \left\| \frac{\sqrt{\gamma}}{\sqrt{\gamma^{\mathbf{S}, \#}}} - 1 \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}}) \cap \mathfrak{h}_1^\infty(\dot{S})} \right) \left\{ \left\| \not{d}_k^\circ(f^\#) \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}})} \right. \\
&+ O(\delta) \left(\int_0^1 \left\| \left(\sqrt{\gamma} \left(-\not{d}_1^*(\kappa) - \frac{1}{2}(\not{d}_1 \vartheta - \not{d}_2^* \vartheta) \right) \right)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}}) \cap \mathfrak{h}_1^\infty(\dot{S})} d\lambda \right. \\
&+ \int_0^1 \left\| \left(\sqrt{\gamma} \left(-\not{d}_1^*(\underline{\kappa}) - \frac{1}{2}(\not{d}_1 \underline{\vartheta} - \not{d}_2^* \underline{\vartheta}) + \not{d}_1^*(\underline{\Omega} \kappa) - \not{d}_1^*(\underline{\Omega}) \vartheta \right. \right. \right. \\
&+ \frac{1}{2} \underline{\Omega}(\not{d}_1 \underline{\vartheta} - \not{d}_2^* \underline{\vartheta}) + \not{d}_1^*(\gamma^{1/2})(\not{d}_1 \vartheta + \not{d}_2^* \vartheta) \underline{b} \\
&+ \left. \left. \left. + \frac{1}{2} \gamma^{1/2}(\not{d}_2 \not{d}_2^* \underline{b} + \not{d}_3^* \not{d}_2^* \underline{b}) + 2\gamma^{1/2} K \underline{b} \right) \right)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}}) \cap \mathfrak{h}_1^\infty(\dot{S})} d\lambda \\
&+ \left\| \left(\underline{\kappa} - \underline{\vartheta} - \underline{\Omega}(\kappa - \vartheta) - \gamma^{1/2}(\not{d}_1 \vartheta + \not{d}_2^* \vartheta) \underline{b} \right)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}}) \cap \mathfrak{h}_1^\infty(\dot{S})} \\
&\left. + \left\| (\kappa + \vartheta)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}}) \cap \mathfrak{h}_1^\infty(\dot{S})} \right) \|f^\# \|_{\mathfrak{h}_{s+1}(\dot{S}, \dot{\mathcal{G}})} \}.
\end{aligned}$$

Next, we have by the iteration assumption (C.1.3)

$$\begin{aligned}
& \left\| \left(\not{d}^{\leq 2}(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \right)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}})} + \sup_{0 \leq \lambda \leq 1} \left\| \left(\not{d}^{\leq 2}(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \right)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}})} \\
&\lesssim \left\| \left(\not{d}^{\leq 2}(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \right)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}}^{\mathbf{S}, \#})} + \sup_{0 \leq \lambda \leq 1} \left\| \left(\not{d}^{\leq 2}(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \right)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}}^{\mathbf{S}, \# \lambda})} \\
&\lesssim \left\| \not{d}^{\leq 2}(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \right\|_{\mathfrak{h}_s(S)} + \sup_{0 \leq \lambda \leq 1} \left\| \not{d}^{\leq 2}(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \right\|_{\mathfrak{h}_s(\mathbf{S}_\lambda)}
\end{aligned}$$

where the surface \mathbf{S}_λ is the image of \dot{S} by ψ_λ . Since $s \leq s_{max} - 3$, we infer in view of our iteration assumption (C.1.2) and our assumptions (9.1.12) (9.1.15) on the (u, s) -foliation

$$\begin{aligned}
& \left\| \left(\not{d}^{\leq 2}(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \right)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}})} + \sup_{0 \leq \lambda \leq 1} \left\| \left(\not{d}^{\leq 2}(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \right)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{\mathcal{G}})} \\
&\lesssim \sup_{\mathcal{R}} \left| \not{d}^{\leq s} \not{d}^{\leq 2}(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \check{\Gamma} \right| \\
&\lesssim \sup_{\mathcal{R}} \left| \not{d}^{\leq s+2}(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \check{\Gamma} \right| \lesssim \delta. \tag{C.1.5}
\end{aligned}$$

Also, we have

$$\begin{aligned}
 & \left\| \left(\not{d}^{\leq 2} \left(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon \right) \right)^{\#} \right\|_{\mathfrak{h}_1^{\infty}(\check{S})} + \sup_{0 \leq \lambda \leq 1} \left\| \left(\not{d}^{\leq 2} \left(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon \right) \right)^{\#\lambda} \right\|_{\mathfrak{h}_1^{\infty}(\check{S})} \\
 = & \left\| \left(\not{d}^{\leq 2} \left(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon \right) \right) \circ \psi \right\|_{\mathfrak{h}_1^{\infty}(\check{S})} + \sup_{0 \leq \lambda \leq 1} \left\| \left(\not{d}^{\leq 2} \left(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon \right) \right) \circ \psi_{\lambda} \right\|_{\mathfrak{h}_1^{\infty}(\check{S})} \\
 \lesssim & \left(\sup_{\mathcal{R}} \left| \mathfrak{d}^{\leq 3} \left(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon \right) \right| \right) \left(1 + \sup_{0 \leq \theta \leq \pi} |\psi'(\theta)| \right) \\
 \lesssim & \left(\sup_{\mathcal{R}} \left| \mathfrak{d}^{\leq 3} \left(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon \right) \right| \right) \left(1 + \|(U', S')\|_{\mathfrak{h}_1^{\infty}(\check{S})} \right) \\
 \lesssim & \overset{\circ}{\epsilon}
 \end{aligned}$$

where we used our assumptions (9.1.12)–(9.1.15) on the (u, s) -foliation and our assumption (9.2.14) on (U', S') . Therefore,

$$\begin{aligned}
 & \left\| \left(\not{d}^{\leq 2} \left(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon \right) \right)^{\#} \right\|_{\mathfrak{h}_1^{\infty}(\check{S})} \lesssim \overset{\circ}{\delta} \\
 \sup_{0 \leq \lambda \leq 1} & \left\| \left(\not{d}^{\leq 2} \left(\check{\Gamma}, r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon \right) \right)^{\#\lambda} \right\|_{\mathfrak{h}_1^{\infty}(\check{S})} \lesssim \overset{\circ}{\delta}.
 \end{aligned} \tag{C.1.6}$$

We deduce

$$\begin{aligned}
 & \|(\not{d}_k^{\mathbf{S}} f)^{\#}\|_{\mathfrak{h}_s(\check{S}, \check{g}^{\mathbf{S}, \#})} (1 + O(\overset{\circ}{\delta})) \\
 = & \left(1 + O(1) \left\| \frac{\sqrt{\gamma}}{\sqrt{\gamma^{\mathbf{S}, \#}}} - 1 \right\|_{\mathfrak{h}_s(\check{S}, \check{g}^{\circ}) \cap \mathfrak{h}_1^{\infty}(\check{S})} \right) \left\{ \left\| \not{d}_k^{\circ}(f^{\#}) \right\|_{\mathfrak{h}_s(\check{S}, \check{g}^{\circ})} + O(\overset{\circ}{\delta}) \|f^{\#}\|_{\mathfrak{h}_{s+1}(\check{S}, \check{g}^{\circ})} \right\}.
 \end{aligned}$$

Next, we estimate the term in the RHS involving γ and $\gamma^{\mathbf{S}, \#}$. From the proof of Lemma 9.2.3, we have

$$\begin{aligned}
 \gamma^{\mathbf{S}, \#} - \gamma &= \frac{1}{2} U \int_0^1 ((e_3 - \underline{\Omega} e_4 - \underline{b} \gamma^{1/2} e_{\theta}) \gamma)^{\#\lambda} d\lambda + S \int_0^1 (e_4 \gamma)^{\#\lambda} d\lambda \\
 &+ \left(\underline{\Omega} + \frac{1}{4} \underline{b}^2 \gamma \right)^{\#} (U')^2 - 2U'S' - (\gamma \underline{b})^{\#} U'.
 \end{aligned}$$

Using a non sharp product rule, we infer

$$\begin{aligned}
& \|\gamma^{\mathbf{S},\#} - \gamma\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} \\
\lesssim & \|U\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} \int_0^1 \left\| \left((e_3 - \underline{\Omega}e_4 - \underline{b}\gamma^{1/2}e_\theta) \gamma \right)^{\#\lambda} \right\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} d\lambda \\
& \|S\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} \int_0^1 \left\| (e_4\gamma)^{\#\lambda} \right\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} d\lambda \\
& + \left\| \left(\underline{\Omega} + \frac{1}{4}\underline{b}^2\gamma \right)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} \|U'\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} + \|U'\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} \|S'\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} \\
& + \left\| (\gamma\underline{b})^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} \|U'\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} \\
\lesssim & \dot{\delta} \int_0^1 \left\| \left(\underline{d}^{\leq 1} (r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon) \right)^{\#\lambda} \right\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} d\lambda \\
& + \dot{\delta} \left\| (r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon)^\# \right\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} + \dot{\delta}
\end{aligned}$$

where we used our assumption (9.2.14) on (U', S') and the fact that $U(0) = S(0) = 0$. Using the estimates (C.1.5) (C.1.6) for $(r^{-2}\gamma - 1, \underline{b}, \underline{\Omega} + \Upsilon)$, we infer

$$\|\gamma^{\mathbf{S},\#} - \gamma\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} \lesssim \dot{\delta}.$$

Together with (9.1.15) for γ , we infer

$$\left\| \frac{\sqrt{\gamma}}{\sqrt{\gamma^{\mathbf{S},\#}}} - 1 \right\|_{\mathfrak{h}_s(\dot{S}, \dot{g}) \cap \mathfrak{h}_1^\infty(\dot{S})} \lesssim \dot{\delta}$$

and hence

$$\|(\underline{d}_k^{\mathbf{S}} f)^\#\|_{\mathfrak{h}_s(\dot{S}, \dot{g}^{\mathbf{S},\#})} (1 + O(\dot{\delta})) = \left(1 + O(\dot{\delta}) \right) \left\{ \left\| \dot{d}_k(f^\#) \right\|_{\mathfrak{h}_s(\dot{S}, \dot{g})} + O(\dot{\delta}) \|f^\#\|_{\mathfrak{h}_{s+1}(\dot{S}, \dot{g})} \right\}.$$

Now, we have

$$\begin{aligned}
\|f^\#\|_{\mathfrak{h}_{s+1}(\dot{S}, \dot{g}^{\mathbf{S},\#})} &= \|f^\#\|_{L^2(\dot{S}, \dot{g}^{\mathbf{S},\#})} + \|(\underline{d}_k^{\mathbf{S}} f)^\#\|_{\mathfrak{h}_s(\dot{S}, \dot{g}^{\mathbf{S},\#})}, \\
\|f^\#\|_{\mathfrak{h}_{s+1}(\dot{S}, \dot{g})} &= \|f^\#\|_{L^2(\dot{S}, \dot{g})} + \|\dot{d}_k(f^\#)\|_{\mathfrak{h}_s(\dot{S}, \dot{g})}.
\end{aligned}$$

Together with Lemma 9.2.3, this yields

$$\|f^\#\|_{\mathfrak{h}_{s+1}(\dot{S}, \dot{g}^{\mathbf{S},\#})} = \|f^\#\|_{\mathfrak{h}_{s+1}(\dot{S}, \dot{g})} (1 + O(\dot{\delta})).$$

This corresponds to our iteration assumption (C.1.3) with s replaced with $s + 1$ for $s \leq s_{max} - 3$. Thus, we have finally derived both iteration assumption (C.1.3) and (C.1.2) with s replaced with $s + 1$ respectively for $s \leq s_{max} - 1$ and $s \leq s_{max} - 3$. Hence, we deduce that they hold respectively for $0 \leq s \leq s_{max}$ and $0 \leq s \leq s_{max} - 2$, i.e.

$$\|h\|_{\mathfrak{h}_k(\mathbf{s})} \lesssim \sup_{\mathcal{R}} |\mathfrak{d}^{\leq k} h| \text{ for } 0 \leq s \leq s_{max}$$

and

$$\|f^\#\|_{\mathfrak{h}_s(\mathring{S}, \mathfrak{g}^{\mathbf{s}, \#})} = \|f^\#\|_{\mathfrak{h}_s(\mathring{S}, \mathring{\mathfrak{g}})} (1 + O(\mathring{\delta})) \text{ for } 0 \leq s \leq s_{max} - 2.$$

Together with Lemma 9.2.2, we deduce

$$\|f\|_{\mathfrak{h}_s(\mathbf{s})} = \|f^\#\|_{\mathfrak{h}_{s+1}(\mathring{S}, \mathfrak{g}^{\mathbf{s}, \#})} = \|f^\#\|_{\mathfrak{h}_{s+1}(\mathring{S}, \mathring{\mathfrak{g}})} (1 + O(\mathring{\delta})) \text{ for all } 0 \leq s \leq s_{max} - 2.$$

This concludes the proof of the lemma. □

Appendix D

APPENDIX TO CHAPTER 10

D.1 Horizontal S -tensors

Consider a null pair e_3, e_4 on (\mathbf{M}, \mathbf{g}) and, at every point $p \in \mathbf{M}$ the horizontal space $S = \{e_3, e_4\}^\perp$. Let γ the metric induced on S . By definition, for all $X, Y \in \mathbf{T}_S \mathbf{M}$, i.e. vectors in \mathbf{M} tangent to S ,

$$h(X, Y) = \mathbf{g}(X, Y)$$

For any $Y \in T(\mathbf{M})$ we define its horizontal projection,

$$Y^\perp = Y + \frac{1}{2}\mathbf{g}(Y, e_3)e_4 + \frac{1}{2}\mathbf{g}(Y, e_4)e_3 \quad (\text{D.1.1})$$

Definition D.1.1. A k -covariant tensor-field U is said to be S -horizontal, $U \in \mathbf{T}_S^k(\mathbf{M})$, if for any X_1, \dots, X_k we have,

$$U(Y_1, \dots, Y_k) = U(Y_1^\perp, \dots, Y_k^\perp)$$

We define the projection operator,

$$\Pi_\mu^\nu := \delta_\mu^\nu - \frac{1}{2}(e_3)_\mu(e_4)^\nu - \frac{1}{2}(e_4)_\mu(e_3)^\nu$$

Clearly $\Pi_\alpha^\mu \Pi_\mu^\beta = \Pi_\alpha^\beta$. An arbitrary tensor $U_{\alpha_1 \dots \alpha_m}$ is said to an S -horizontal tensor, or simply S -tensor, if

$$\Pi_{\alpha_1}^{\beta_1} \dots \Pi_{\alpha_m}^{\beta_m} U_{\beta_1 \dots \beta_m} = U_{\alpha_1 \dots \alpha_m}.$$

Definition D.1.2. Given $X \in \mathbf{T}(\mathbf{M})$ and $Y \in \mathbf{T}_S(\mathbf{M})$ we define,

$$\dot{\mathbf{D}}_X Y := (\mathbf{D}_X Y)^\perp$$

Remark D.1.3. In the particular case when S is integrable and both $X, Y \in \mathbf{T}_S \mathbf{M}$ then $\dot{\mathbf{D}}_X Y$ is the standard induced covariant differentiation on S .

Definition D.1.4. Given a general, covariant, S -horizontal tensor-field U we define its horizontal covariant derivative according to the formula,

$$\dot{\mathbf{D}}_X U(Y_1, \dots, Y_k) = X(U(Y_1, \dots, Y_k)) - U(\dot{\mathbf{D}}_X Y_1, \dots, Y_k) - \dots - U(Y_1, \dots, \dot{\mathbf{D}}_X Y_k) \quad (D.1.2)$$

where $X \in \mathbf{TM}$ and $Y_1, \dots, Y_k \in \mathbf{T}_S \mathbf{M}$.

Proposition D.1.5. For all $X \in \mathbf{TM}$ and $Y_1, Y_2 \in \mathbf{T}_S \mathbf{M}$,

$$Xh(Y_1, Y_2) = h(\dot{\mathbf{D}}_X Y_1, Y_2) + h(Y_1, \dot{\mathbf{D}}_X Y_2).$$

Proof. Indeed,

$$\begin{aligned} Xh(Y_1, Y_2) &= Xg(Y_1, Y_2) = g(\mathbf{D}_X Y_1, Y_2) + g(Y_1, \mathbf{D}_X Y_2) = g(\dot{\mathbf{D}}_X Y_1, Y_2) + g(Y_1, \dot{\mathbf{D}}_X Y_2) \\ &= h(\dot{\mathbf{D}}_X Y_1, Y_2) + h(Y_1, \dot{\mathbf{D}}_X Y_2) \end{aligned}$$

□

Given an orthonormal frame e_1, e_2 on S we have,

$$\dot{\mathbf{D}}_\mu e_A = \sum_{B=1,2} (\Lambda_\mu)_{AB} e_B \quad A, B = 1, 2$$

where,

$$(\Lambda_\mu)_{\alpha\beta} := g(\mathbf{D}_\mu e_\beta, e_\alpha)$$

D.1.1 Mixed tensors

We consider tensors $\mathbf{T}^k \mathbf{M} \otimes \mathbf{T}_S^l \mathbf{M}$, i.e. tensors of the form,

$$U_{\mu_1 \dots \mu_k, A_1 \dots A_l}$$

for which we define,

$$\begin{aligned} \dot{\mathbf{D}}_\mu U_{\nu_1 \dots \nu_k, A_1 \dots A_l} &= e_\mu U_{\nu_1 \dots \nu_k, A_1 \dots A_l} - U_{\mathbf{D}_\mu \nu_1 \dots \nu_k, A_1 \dots A_l} - \dots - U_{\nu_1 \dots \nu_k, \mathbf{D}_\mu A_1 \dots A_l} \\ &\quad - U_{\nu_1 \dots \nu_k, \dot{\mathbf{D}}_\mu A_1 \dots A_l} - U_{\nu_1 \dots \nu_k, A_1 \dots \dot{\mathbf{D}}_\mu A_l} \end{aligned}$$

We are now ready to prove the following,

Proposition D.1.6. *We have the curvature formula*

$$(\dot{\mathbf{D}}_\mu \dot{\mathbf{D}}_\nu - \dot{\mathbf{D}}_\nu \dot{\mathbf{D}}_\mu) \Psi_A = \mathbf{R}_A{}^B{}_{\mu\nu} \Psi_B$$

More generally,

$$(\dot{\mathbf{D}}_\mu \dot{\mathbf{D}}_\nu - \dot{\mathbf{D}}_\nu \dot{\mathbf{D}}_\mu) \Psi_{\lambda A} = \mathbf{R}_\lambda{}^\sigma{}_{\mu\nu} \Psi_{\sigma A} + \mathbf{R}_A{}^B{}_{\mu\nu} \Psi_{\lambda B}$$

Proof. Straightforward verification. □

D.1.2 Invariant Lagrangian

We introduce,

$$\mathcal{L} = g^{\mu\nu} h_{AB} \dot{\mathbf{D}}_\mu \Psi^A \dot{\mathbf{D}}_\nu \Psi^B + V h_{AB} \Psi^A \Psi^B$$

Proposition D.1.7. *The Euler Lagrange equations are given by:*

$$\square \Psi^A = V \Psi^A$$

where $\square \Psi^A := \mathbf{g}^{\mu\nu} \dot{\mathbf{D}}_\mu \dot{\mathbf{D}}_\nu \Psi^A$.

Proof. The variation of the action is given by,

$$\begin{aligned} 0 &= 2 \int_{\mathbf{M}} h_{AB} \left(\mathbf{g}^{\mu\nu} \dot{\mathbf{D}}_\mu \Psi^A \dot{\mathbf{D}}_\nu (\delta \Psi)^B + V \Psi^A \delta \Psi^B \right) dv_{\mathbf{g}} \\ &= 2 \int_{\mathbf{M}} \mathbf{D}_\nu \left(\mathbf{g}^{\mu\nu} h_{AB} \dot{\mathbf{D}}_\mu \Psi^A (\delta \Psi)^B \right) dv_{\mathbf{g}} - 2 \int_{\mathbf{M}} h_{AB} \left(\mathbf{g}^{\mu\nu} \dot{\mathbf{D}}_\nu \dot{\mathbf{D}}_\mu \Psi^A (\delta \Psi)^B - V \Psi^A \delta \Psi^B \right) dv_{\mathbf{g}} \\ &= -2 \int_{\mathbf{M}} h_{AB} \left(\mathbf{g}^{\mu\nu} \dot{\mathbf{D}}_\nu \dot{\mathbf{D}}_\mu \Psi^A (\delta \Psi)^B - V \Psi^A \delta \Psi^B \right) dv_{\mathbf{g}} \end{aligned}$$

from which the proposition follows. □

D.1.3 Comparison of the Lagrangians

Let $\Psi \in \mathcal{S}_2(\mathcal{M})$ and $\psi \in \mathfrak{s}_2$ its reduced form. Note that the Lagrangian of the scalar equation

$$\square_{\mathbf{g}} \psi = V \psi + 4(\epsilon_\theta \Phi)^2 \psi$$

is given by,

$$\mathcal{L}(\psi) : = \mathbf{g}^{\mu\nu} \partial_\mu \psi \partial_\nu \psi + (V + 4(e_\theta \Phi)^2) \psi^2$$

while the Lagrangian for,

$$\square_{\mathbf{g}} \Psi = V \Psi$$

is given by

$$\mathcal{L}(\Psi) = \mathbf{g}^{\mu\nu} \dot{\mathbf{D}}_\mu \Psi \cdot \dot{\mathbf{D}}_\nu \Psi + V \Psi \cdot \Psi$$

Proposition D.1.8. *We have,*

$$\mathcal{L}(\Psi) = 2\mathcal{L}(\psi) \tag{D.1.3}$$

Proof. Observe that,

$$\mathbf{g}^{\mu\nu} \dot{\mathbf{D}}_\mu \Psi \dot{\mathbf{D}}_\nu \Psi = -\dot{\mathbf{D}}_3 \Psi \cdot \dot{\mathbf{D}}_4 \Psi + \dot{\mathbf{D}}_\theta \Psi \cdot \dot{\mathbf{D}}_\theta \Psi + \dot{\mathbf{D}}_\varphi \Psi \cdot \dot{\mathbf{D}}_\varphi \Psi$$

Now, recalling that,

$$\begin{aligned} \nabla_\varphi e_\varphi &= -e_\theta \Phi e_\theta, & \nabla_\varphi e_\theta &= e_\theta(\Phi) e_\varphi \\ \nabla_\theta e_\theta &= 0 & \nabla_\theta e_\varphi &= 0 \end{aligned}$$

we deduce

$$\begin{aligned} \dot{\mathbf{D}}_3 \Psi \cdot \dot{\mathbf{D}}_4 \Psi &= e_3 \Psi \cdot e_4 \Psi = 2e_3 \psi e_4 \psi \\ \dot{\mathbf{D}}_\theta \Psi \cdot \dot{\mathbf{D}}_\theta \Psi &= \dot{\mathbf{D}}_\theta \Psi_{\theta\theta} \dot{\mathbf{D}}_\theta \Psi_{\theta\theta} + 2\dot{\mathbf{D}}_\theta \Psi_{\theta\varphi} \dot{\mathbf{D}}_\theta \Psi_{\theta\varphi} + \dot{\mathbf{D}}_\theta \Psi_{\varphi\varphi} \dot{\mathbf{D}}_\theta \Psi_{\varphi\varphi} \\ &= 2(e_\theta \psi)^2 \\ \dot{\mathbf{D}}_\varphi \Psi \cdot \dot{\mathbf{D}}_\varphi \Psi &= \dot{\mathbf{D}}_\varphi \Psi_{\theta\theta} \dot{\mathbf{D}}_\varphi \Psi_{\theta\theta} + 2\dot{\mathbf{D}}_\varphi \Psi_{\theta\varphi} \dot{\mathbf{D}}_\varphi \Psi_{\theta\varphi} + \dot{\mathbf{D}}_\varphi \Psi_{\varphi\varphi} \dot{\mathbf{D}}_\varphi \Psi_{\varphi\varphi} \\ &= 2(e_\varphi \psi)^2 + 2(-\Psi_{\dot{\mathbf{D}}_\varphi \theta\varphi} - \Psi_{\theta \dot{\mathbf{D}}_\varphi \varphi}) \cdot (-\Psi_{\dot{\mathbf{D}}_\varphi \theta\varphi} - \Psi_{\theta \dot{\mathbf{D}}_\varphi \varphi}) \\ &= 2(e_\varphi \psi)^2 + 2(-e_\theta(\Phi) \Psi_{\varphi\varphi} + e_\theta(\Phi) \Psi_{\theta\theta}) \cdot (-e_\theta(\Phi) \Psi_{\varphi\varphi} + e_\theta(\Phi) \Psi_{\theta\theta}) \\ &= 2(e_\varphi \psi)^2 + 8(e_\theta \Phi)^2 \psi^2 \end{aligned}$$

Hence,

$$\mathbf{g}^{\mu\nu} \dot{\mathbf{D}}_\mu \Psi \dot{\mathbf{D}}_\nu \Psi = -2e_3 \psi e_4 \psi + 2(e_\theta \psi)^2 + 2(e_\varphi \psi)^2 + 4(e_\theta \Phi)^2 \psi^2$$

and

$$\mathcal{L}(\Psi) = -2e_3 \Psi e_4 \psi + 2(e_\theta \psi)^2 + 2(e_\varphi \psi)^2 + 8(e_\theta \Phi)^2 \psi^2 + 2V \psi^2$$

□

D.1.4 Energy-Momentum tensor

Consider the energy-momentum tensor,

$$\mathcal{Q}_{\mu\nu} := \dot{\mathbf{D}}_\mu \Psi \cdot \dot{\mathbf{D}}_\nu \Psi - \frac{1}{2} \mathbf{g}_{\mu\nu} \left(\dot{\mathbf{D}}_\lambda \Psi \cdot \dot{\mathbf{D}}^\lambda \Psi + V \Psi \cdot \Psi \right)$$

Lemma D.1.9. *We have,*

$$\mathbf{D}^\nu \mathcal{Q}_{\mu\nu} = \dot{\mathbf{D}}_\mu \Psi \cdot (\dot{\square} \Psi - V \Psi) + \dot{\mathbf{D}}^\nu \Psi^A \mathbf{R}_{AB\nu\mu} \Psi^B - \frac{1}{2} \mathbf{D}_\mu V \Psi \cdot \Psi$$

Proof. We have,

$$\begin{aligned} \mathbf{D}^\nu \mathcal{Q}_{\mu\nu} &= \dot{\mathbf{D}}^\nu \dot{\mathbf{D}}_\nu \Psi \cdot \dot{\mathbf{D}}_\mu \Psi + \dot{\mathbf{D}}^\nu \Psi \cdot \left(\dot{\mathbf{D}}_\nu \dot{\mathbf{D}}_\mu - \dot{\mathbf{D}}_\mu \dot{\mathbf{D}}_\nu \right) \Psi - V \mathbf{D}_\mu \Psi \cdot \Psi - \frac{1}{2} \mathbf{D}_\mu V \Psi \cdot \Psi \\ &= \dot{\mathbf{D}}_\mu \Psi \cdot \dot{\mathbf{D}}^\nu \dot{\mathbf{D}}_\nu \Psi + \dot{\mathbf{D}}^\nu \Psi^A \mathbf{R}_{AB\nu\mu} \Psi^B - V \mathbf{D}_\mu \Psi \cdot \Psi - \frac{1}{2} \mathbf{D}_\mu V \Psi \cdot \Psi \\ &= \dot{\mathbf{D}}_\mu \Psi \cdot (\dot{\square} \Psi - V \Psi) + \dot{\mathbf{D}}^\nu \Psi^A \mathbf{R}_{AB\nu\mu} \Psi^B - \frac{1}{2} \mathbf{D}_\mu V \Psi \cdot \Psi \end{aligned}$$

□

Lemma D.1.10. *Relative to an arbitrary \mathbf{Z} -polarized frame $e_3, e_4, e_\theta, e_\varphi$ we have,*

$$\begin{aligned} \mathcal{Q}_{33} &= |e_3 \Psi|^2, \\ \mathcal{Q}_{44} &= |e_4 \Psi|^2, \\ \mathcal{Q}_{34} &= |\nabla \Psi|^2 + V |\Psi|^2. \end{aligned}$$

If ψ is the reduced form of Ψ ,

$$\begin{aligned} \mathcal{Q}_{33} &= 2(e_3 \psi)^2, \\ \mathcal{Q}_{44} &= 2(e_4 \psi)^2, \\ \mathcal{Q}_{34} &= 2(e_\theta \psi)^2 + 2(e_\varphi \psi)^2 + 2V |\psi|^2 + 8(e_\theta \Phi)^2 \psi^2. \end{aligned}$$

Also,

$$\mathbf{g}^{\mu\nu} \mathcal{Q}_{\mu\nu} = -\mathcal{L}(\Psi) - V |\Psi|^2,$$

$$|\mathcal{L}(\Psi)| \lesssim |e_3 \Psi| |e_4 \Psi| + |\nabla \Psi|^2 + V |\Psi|^2,$$

and

$$\begin{aligned} |\mathcal{Q}_{AB}| &\leq |e_3 \Psi| |e_4 \Psi| + |\nabla \Psi|^2 + V |\Psi|^2, \\ |\mathcal{Q}_{A3}| &\leq |e_3 \Psi| |\nabla \Psi|, \\ |\mathcal{Q}_{A4}| &\leq |e_4 \Psi| |\nabla \Psi|. \end{aligned}$$

D.2 Standard Calculation

Proposition D.2.1. *Consider an admissible spacetime \mathcal{M} and $\Psi \in \mathcal{S}_2(\mathcal{M})$ and X a vectorfield of the form,*

$$X = ae_3 + be_4,$$

1. *The 1-form $\mathcal{P}_\mu = \mathcal{Q}_{\mu\nu}X^\nu$ verifies,*

$$\mathbf{D}^\mu \mathcal{P}_\mu = X^\mu \dot{\mathbf{D}}_\mu \Psi \cdot (\dot{\square}\Psi - V\Psi) - X(V)\Psi \cdot \Psi$$

2. *Let X as above, w a scalar and \mathbf{M} a one form. Define,*

$$\mathcal{P}_\mu = \mathcal{P}_\mu[X, w, M] = \mathcal{Q}_{\mu\nu}X^\nu + \frac{1}{2}w\Psi \cdot \dot{\mathbf{D}}_\mu \Psi - \frac{1}{4}|\Psi|^2\partial_\mu w + \frac{1}{4}|\Psi|^2 M_\mu$$

Then, with $|\Psi|^2 := \Psi \cdot \Psi$,

$$\begin{aligned} \mathbf{D}^\mu \mathcal{P}_\mu[X, w, M] &= \frac{1}{2}\mathcal{Q} \cdot {}^{(X)}\pi - \frac{1}{2}X(V)\Psi \cdot \Psi + \frac{1}{2}w\mathcal{L}[\Psi] - \frac{1}{4}|\Psi|^2\Box_{\mathbf{g}}w \\ &\quad + \frac{1}{4}|\Psi|^2\mathbf{Div}M + \frac{1}{2}\Psi \cdot \dot{\mathbf{D}}_\mu \Psi M^\mu + \left(X(\Psi) + \frac{1}{2}w\Psi\right) \cdot (\dot{\square}\Psi - V\Psi) \end{aligned}$$

Proof. Let $\mathcal{P}_\mu[X, 0, 0] = \mathcal{Q}_{\mu\nu}X^\nu$, Then,

$$\begin{aligned} \mathbf{D}^\mu \mathcal{P}_\mu &= X^\mu \dot{\mathbf{D}}_\mu \Psi \cdot \left(\dot{\mathbf{D}}^\nu \dot{\mathbf{D}}_\nu \Psi - V\Psi\right) + X^\mu \dot{\mathbf{D}}^\nu \Psi^A \mathbf{R}_{AB\nu\mu} \Psi^B - \frac{1}{2}X^\mu \mathbf{D}_\mu V\Psi \cdot \Psi \\ &= X^\mu \dot{\mathbf{D}}_\mu \Psi \cdot (\dot{\square}\Psi - V\Psi) - \frac{1}{2}X(V)|\Psi|^2 \end{aligned}$$

Assume $X = ae_3 + be_4$. Then, since only the middle components of \mathbf{R} are relevant, and recalling that $\mathbf{R}_{AB43} = -{}^*\rho \in_{AB} = 0$, we derive,

$$X^\mu \dot{\mathbf{D}}^\nu \Psi^A \mathbf{R}_{AB\nu 3} \Psi^B = a\dot{\mathbf{D}}^4 \Psi^A \mathbf{R}_{AB43} \Psi^B + b\dot{\mathbf{D}}^3 \Psi^A \mathbf{R}_{AB434} \Psi^B = 0$$

To prove the second part of the proposition we write with $\mathcal{N}[\Psi] := \dot{\square}\Psi - V\Psi$,

$$\begin{aligned} \mathbf{D}^\mu \mathcal{P}_\mu[X, w, M] &= \frac{1}{2}\mathcal{Q} \cdot {}^{(X)}\pi + X(\Psi) \cdot \mathcal{N}[\Psi] - \frac{1}{2}X(V)\Psi \cdot \Psi + \frac{1}{2}\mathbf{D}^\mu w \Psi \cdot \dot{\mathbf{D}}_\mu \Psi \\ &\quad + \frac{1}{2}w \dot{\mathbf{D}}^\mu \Psi \cdot \dot{\mathbf{D}}_\mu \Psi + \frac{1}{2}w\Psi \Box_{\mathbf{g}}\Psi - \frac{1}{2}\Psi \cdot \dot{\mathbf{D}}^\mu \Psi \partial_\mu w - \frac{1}{4}|\Psi|^2\Box_{\mathbf{g}}w \\ &\quad + \frac{1}{4}|\Psi|^2\mathbf{Div}M + \frac{1}{2}\Psi \cdot \dot{\mathbf{D}}_\mu \Psi M^\mu \\ &= \frac{1}{2}\mathcal{Q} \cdot {}^{(X)}\pi - \frac{1}{2}X(V)\Psi \cdot \Psi + \frac{1}{2}w \dot{\mathbf{D}}^\mu \Psi \cdot \dot{\mathbf{D}}_\mu \Psi + \frac{1}{2}w\Psi (V\Psi + \mathcal{N}[\Psi]) \\ &\quad - \frac{1}{4}|\Psi|^2\Box_{\mathbf{g}}w + \frac{1}{4}|\Psi|^2\mathbf{Div}M + \frac{1}{2}\Psi \cdot \dot{\mathbf{D}}_\mu \Psi M^\mu + X(\Psi) \cdot \Psi \cdot \mathcal{N}[\Psi] \end{aligned}$$

Hence,

$$\begin{aligned} \mathbf{D}^\mu \mathcal{P}_\mu[X, w, M] &= \frac{1}{2} \mathcal{Q} \cdot {}^{(X)}\pi - \frac{1}{2} X(V) \Psi \cdot \Psi + \frac{1}{2} w \mathcal{L}[\Psi] - \frac{1}{4} |\Psi|^2 \square_{\mathbf{g}} w \\ &+ \frac{1}{4} |\Psi|^2 \mathbf{Div} M + \frac{1}{2} \Psi \cdot \dot{\mathbf{D}}_\mu \Psi M^\mu + \left(X(\Psi) + \frac{1}{2} w \Psi \right) \cdot \mathcal{N}[\Psi] \end{aligned}$$

as desired. \square

Remark D.2.2. *As consequence of the proposition above we deduce that every time we use vectorfields of the form $ae_3 + be_4$ as multipliers, the equation $\square\Psi - V\Psi = \mathcal{N}$ is treated exactly in the same manner as the scalar equation $\square\psi - V\psi = N$.*

Remark D.2.3. *Note that in Schwarzschild our potential $V = \kappa \underline{\kappa} = 4\Upsilon r^{-2}$ verifies,*

$$\begin{aligned} \frac{1}{4} \partial_r V &= \partial_r \left[r^{-2} \left(1 - \frac{2m}{r} \right) \right] = -2r^{-3} \left(1 - \frac{2m}{r} \right) + \frac{2m}{r^4} \\ &= -2 \frac{r - 3m}{r^4}. \end{aligned}$$

D.3 Vectorfield X_f

Lemma D.3.1. *Let $X_f := fe_4$. Then with ${}^{(X)}\Lambda = \frac{2f}{r}$ and ${}^{(X)}\tilde{\pi} = {}^{(X)}\pi - {}^{(X)}\Lambda \mathbf{g} = {}^{(X)}\pi - \frac{2f}{r} \mathbf{g}$,*

- *We have,*

$$\begin{aligned} {}^{(X)}\tilde{\pi}_{44} &= 0, \quad {}^{(X)}\pi_{4\varphi} = 0, \quad {}^{(X)}\pi_{3\varphi} = 0, \\ {}^{(X)}\tilde{\pi}_{43} &= -2e_4 f + 4f\omega + \frac{4f}{r} = -2 \left(e_4(f) - \frac{2f}{r} \right) + 4f\omega, \\ {}^{(X)}\tilde{\pi}_{4\theta} &= 2f\xi, \\ {}^{(X)}\tilde{\pi}_{AB} &= 2f {}^{(1+3)}\chi_{AB} - \frac{2f}{r} \not{g}_{AB} = 2f \left({}^{(1+3)}\chi_{AB} - \frac{1}{r} \delta_{AB} \right), \\ {}^{(X)}\tilde{\pi}_{3\theta} &= 2f(\eta + \zeta), \\ {}^{(X)}\tilde{\pi}_{33} &= -8f\underline{\omega} - 4e_3(f). \end{aligned} \tag{D.3.1}$$

- In particular, we have,

$$\begin{aligned}
{}^{(X)}\tilde{\pi}_{43} &= -2f' + \frac{4f}{r} + O(\epsilon) \min\{w_{1,1}, w_{2,1/2}\} (|f| + r|f'|), \\
{}^{(X)}\tilde{\pi}_{4A} &= \epsilon \min\{w_{2,1}, w_{3,1/2}\}, \\
{}^{(X)}\tilde{\pi}_{AB} &= O(\epsilon) \min\{w_{1,1}, w_{2,1/2}\}|f|, \\
{}^{(X)}\tilde{\pi}_{3A} &= O(\epsilon)w_{1,1}|f|, \\
{}^{(X)}\tilde{\pi}_{33} &= 4f'\Upsilon - 4\Upsilon' + O(\epsilon)w_{1,1}(|f| + r|f'|).
\end{aligned} \tag{D.3.2}$$

- We have,

$$\Box {}^{(X)}\Lambda = \frac{2}{r}f'' + O\left(\frac{m}{r^4} + \epsilon w_{3,1}\right) (|f| + r|f'| + r^2|f''|). \tag{D.3.3}$$

Proof. We calculate ${}^{(X)}\pi_{\alpha\beta} = \mathbf{g}(D_{e_\alpha}X, e_\beta) + \mathbf{g}(\mathbf{D}_{e_\beta}X, e_\alpha)$,

$$\begin{aligned}
{}^{(X)}\pi_{44} &= 0 \\
{}^{(X)}\pi_{43} &= -2e_4f + 4f\omega \\
{}^{(X)}\pi_{4\theta} &= 2f\xi \\
{}^{(X)}\pi_{AB} &= 2f{}^{(1+3)}\chi_{AB} \\
{}^{(X)}\pi_{3\theta} &= 2f(\eta + \zeta) \\
{}^{(X)}\pi_{33} &= -8f\underline{\omega} - 4e_3(f)
\end{aligned}$$

We deduce, for ${}^{(X)}\tilde{\pi} = {}^{(X)}\pi - {}^{(X)}\Lambda\mathbf{g} = {}^{(X)}\pi - \frac{2f}{r}\mathbf{g}$,

$$\begin{aligned}
{}^{(X)}\tilde{\pi}_{44} &= 0 \\
{}^{(X)}\tilde{\pi}_{43} &= -2e_4f + 4f\omega + \frac{4f}{r} = -2\left(e_4(f) - \frac{2f}{r}\right) + 4f\omega \\
{}^{(X)}\tilde{\pi}_{4\theta} &= 2f\xi \\
{}^{(X)}\tilde{\pi}_{AB} &= 2f{}^{(1+3)}\chi_{AB} - \frac{2f}{r}\mathcal{g}_{AB} = 2f\left({}^{(1+3)}\chi_{AB} - \frac{1}{r}\delta_{AB}\right) \\
{}^{(X)}\tilde{\pi}_{3\theta} &= 2f(\eta + \zeta) \\
{}^{(X)}\tilde{\pi}_{33} &= -8f\underline{\omega} - 4e_3(f)
\end{aligned}$$

Under the assumptions (10.2.8)–(10.2.9) on the Ricci coefficients (with respect to the

frame (e'_3, e'_4) , we deduce,

$$\begin{aligned}
 {}^{(X)}\tilde{\pi}_{43} &= -2e_4f + 4f\omega = -2f' + \frac{4f}{r} - 2f'(e_4(r) - 1) + 4f(\omega - 1) \\
 &= -2f' + \frac{4f}{r} + \epsilon \min\{w_{1,1}, w_{2,1/2}\} (|f| + r|f'|) \\
 {}^{(X)}\tilde{\pi}_{4A} &= \epsilon \min\{w_{2,1}, w_{3,1/2}\}, \\
 {}^{(X)}\tilde{\pi}_{AB} &= \epsilon \min\{w_{1,1}, w_{2,1/2}\}|f| \\
 {}^{(X)}\tilde{\pi}_{3A} &= \min\{w_{1,1}, w_{2,1/2}\}|f| \\
 {}^{(X)}\tilde{\pi}_{33} &= -8f\underline{\omega} - 4e_3(f) = -8f \left(\frac{m}{r^2} + \epsilon w_{1,1} \right) - 4f'(-\Upsilon + \epsilon w_{0,1}) \\
 &= 4f'\Upsilon - 4\Upsilon' + \epsilon w_{1,1} (|f| + r|f'|)
 \end{aligned}$$

To prove formula (D.3.3) we make use of the following (see also Lemma 10.1.11),

Lemma D.3.2. *If $h = h(r)$ then*

$$\square h = \Upsilon h''(r) + \left(\frac{2}{r} - \frac{2m}{r^2} \right) h' + O(\epsilon)w_{2,1} (|h| + r|h'| + r^2|h''|)$$

Proof. For a general scalar h ,

$$\square h = -\frac{1}{2}(e_3e_4 + e_4e_3)h + \underline{\Delta}h + \left({}^{(1+3)}\underline{\omega} - \frac{1}{2} {}^{(1+3)}\text{tr}\underline{\chi} \right) e_4h + ({}^{(1+3)}\omega - \frac{1}{2} {}^{(1+3)}\text{tr}\chi) e_3h$$

with $\underline{\Delta}h = e_\theta e_\theta h + (e_\theta \Phi)^2 e_\theta h = 0$ if h is radial. Thus,

$$\begin{aligned}
 \square h &= -\frac{1}{2}(e_3e_4 + e_4e_3)h + ({}^{(1+3)}\underline{\omega} - \frac{1}{2} {}^{(1+3)}\text{tr}\underline{\chi})e_4h + ({}^{(1+3)}\omega - \frac{1}{2} {}^{(1+3)}\text{tr}\chi)e_3h \\
 &= -f''(e_3r)(e_4r) - \frac{1}{2}h'(e_3e_4 + e_4e_3)r + h' \left[({}^{(1+3)}\underline{\omega} - \frac{1}{2} {}^{(1+3)}\text{tr}\underline{\chi})e_4r + ({}^{(1+3)}\omega - \frac{1}{2} {}^{(1+3)}\text{tr}\chi)e_3r \right] \\
 &= -h''(-\Upsilon + O(\epsilon)w_{0,1})(1 + O(\epsilon)w_{1,1}) + \left(\frac{m}{r^2} + O(\epsilon)w_{1,1} \right) h' \\
 &+ h' \left[\left(\frac{m}{r^2} + \frac{\Upsilon}{r} + O(\epsilon)w_{1,1} \right) (1 + O(\epsilon)w_{1,1}) + \left(-\frac{1}{r} + O(\epsilon)w_{1,1} \right) (-\Upsilon + O(\epsilon)w_{0,1}) \right] \\
 &= \Upsilon h'' + \left(\frac{2}{r} - \frac{2m}{r^2} \right) h' + O(\epsilon)w_{2,1} (|h| + r|h'| + r^2|h''|)
 \end{aligned}$$

which concludes the proof of Lemma D.3.2. □

In view of Lemma D.3.2,

$$\square^{(X)}\Lambda = \square\left(\frac{2f}{r}\right) = \Upsilon\left(\frac{2f}{r}\right)'' + \left(\frac{2}{r} - \frac{2m}{r^2}\right)\left(\frac{2f}{r}\right)' + O(\epsilon)w_{3,1}(|f| + r|f'| + r^2|f''|)$$

Note that,

$$\begin{aligned} \Upsilon\left(\frac{2f}{r}\right)'' + \left(\frac{2}{r} - \frac{2m}{r^2}\right)\left(\frac{2f}{r}\right)' &= \Upsilon\left(\frac{2f''}{r} - \frac{4f'}{r^2} + \frac{4f}{r^3}\right) + \left(\frac{2}{r} - \frac{2m}{r^2}\right)\left(\frac{2f'}{r} - \frac{2f}{r^2}\right) \\ &= \frac{2\Upsilon}{r}f'' - (\Upsilon - 1)\frac{4f'}{r^2} + (\Upsilon - 1)\frac{4f}{r^3} - \frac{2m}{r^2}\left(\frac{2f'}{r} - \frac{2f}{r^2}\right) \\ &= \frac{2}{r} + O\left(\frac{m}{r^4}\right)(|f| + r|f'| + r^2|f''|) \end{aligned}$$

Hence,

$$\square^{(X)}\Lambda = \frac{2}{r}f'' + O\left(\frac{m}{r^4}\epsilon w_{3,1}\right)(|f| + r|f'| + r^2|f''|)$$

as desired. This concludes the proof of Lemma D.3.1. \square

D.4 Proof of Proposition 10.3.1

In view of the following Leibniz rule which holds for any scalar f ,

$$\begin{aligned} -\mathbb{A}_2(f\psi) &= \mathbb{A}_2^*(f\psi) + 2Kf\psi \\ &= \mathbb{A}_2^*(f\mathbb{A}_2\psi + e_\theta(f)\psi) + 2Kf\psi \\ &= -f\mathbb{A}_2\psi - e_\theta(f)\mathbb{A}_2\psi + e_\theta(f)\mathbb{A}_3^*\psi - \mathbb{A}_0(f)\psi, \end{aligned}$$

we have the following computation

$$\begin{aligned} e_4(\square_2(r\psi)) &= e_4(r\square_2\psi) - e_4(e_3(r)e_4\psi) - e_4(e_4(r)e_3\psi) - 2e_4(e_\theta(r)\mathbb{A}_2\psi) \\ &\quad + 2e_4(e_\theta(r)\mathbb{A}_3^*\psi) + e_4(\square_0(r)\psi) \\ &= e_4(r\square_2\psi) - e_4\left(\frac{r}{2}(\underline{\kappa} + \underline{A})e_4\psi\right) - e_4\left(\frac{r}{2}(\kappa + A)e_3\psi\right) \\ &\quad + e_4(\square_0(r)\psi) + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi \\ &= e_4(r\square_2\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_4\psi) - \frac{1}{2}e_4(r\kappa e_3\psi) + e_4(\square_0(r)\psi) + r^{-1}\text{Err}, \end{aligned}$$

where we have introduced the notation, used throughout the proof of Proposition 10.3.1,

$$\text{Err} := r^2\Gamma_g e_4 e_3 \psi + r\Gamma_b e_4 \mathfrak{d} \psi + \mathfrak{d}^{\leq 1}(\Gamma_b)\mathfrak{d}^{\leq 1}\psi + r\mathfrak{d}^{\leq 1}(\Gamma_g)e_3\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^2\psi.$$

Next, recall that we have

$$\square_2\psi = -e_4e_3\psi + \underline{\Delta}_2\psi + \left(2\omega - \frac{1}{2}\kappa\right)e_3\psi - \frac{1}{2}\underline{\kappa}e_4\psi + 2\underline{\eta}e_\theta\psi.$$

We infer

$$\begin{aligned} \square_0(r) &= -e_4(e_3(r)) + \underline{\Delta}_2(r) + \left(2\omega - \frac{1}{2}\kappa\right)e_3(r) - \frac{1}{2}\underline{\kappa}e_4(r) + 2\underline{\eta}e_\theta(r) \\ &= -e_4\left(\frac{r}{2}(\underline{\kappa} + \underline{A})\right) + \left(2\omega - \frac{1}{2}\kappa\right)\frac{r}{2}(\underline{\kappa} + \underline{A}) - \frac{1}{2}\underline{\kappa}\frac{r}{2}(\kappa + A) + r^{-1}\mathfrak{d}^{\leq 1}\Gamma_g \\ &= -\frac{1}{2}e_4(r\underline{\kappa}) + \frac{1}{2}\left(2\omega - \frac{1}{2}\kappa\right)r\underline{\kappa} - \frac{1}{4}r\underline{\kappa}\kappa + r\mathfrak{d}^{\leq 1}\Gamma_g \\ &= \frac{2}{r} + O(r^{-2}) + \mathfrak{d}^{\leq 1}\Gamma_b \end{aligned}$$

and hence

$$\begin{aligned} e_4(\square_2(r\psi)) &= e_4(r\square_2\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_4\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_3\psi) + e_4(\square_0(r)\psi) + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi \\ &= e_4(r\square_2\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_4\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_3\psi) + e_4\left(\frac{2}{r}\psi\right) + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err} \end{aligned}$$

so that

$$e_4(r\square_2\psi) = e_4(\square_2(r\psi)) + \frac{1}{2}e_4(r\underline{\kappa}e_4\psi) + \frac{1}{2}e_4(r\underline{\kappa}e_3\psi) - e_4\left(\frac{2}{r}\psi\right) + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err}.$$

We infer

$$\begin{aligned} \square_2(e_4(r\psi)) - e_4(r\square_2\psi) &= [\square_2, e_4](r\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_4\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_3\psi) + e_4\left(\frac{2}{r}\psi\right) \\ &\quad + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err}. \end{aligned}$$

Next, using again

$$\square_2\psi = -e_4e_3\psi + \underline{\Delta}_2\psi + \left(2\omega - \frac{1}{2}\kappa\right)e_3\psi - \frac{1}{2}\underline{\kappa}e_4\psi + 2\underline{\eta}e_\theta\psi,$$

we infer

$$\begin{aligned} [\square_2, e_4]\psi &= -e_4[e_3, e_4]\psi + [\underline{\Delta}_2, e_4]\psi + \left(2\omega - \frac{1}{2}\kappa\right)[e_3, e_4]\psi - e_4\left(2\omega - \frac{1}{2}\kappa\right)e_3\psi \\ &\quad + \frac{1}{2}e_4(\underline{\kappa})e_4\psi + 2\underline{\eta}[e_\theta, e_4]\psi - 2e_4(\underline{\eta})e_\theta\psi \\ &= -e_4[e_3, e_4]\psi + [\underline{\Delta}_2, e_4]\psi + \left(2\omega - \frac{1}{2}\kappa\right)[e_3, e_4]\psi - \left(2e_4(\omega) - \frac{1}{2}\left(-\frac{1}{2}\kappa^2 - 2\omega\kappa\right)\right)e_3\psi \\ &\quad + \frac{1}{2}\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\omega\underline{\kappa} + 2\rho\right)e_4\psi + 2\underline{\eta}[e_\theta, e_4]\psi + r^{-2}\mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}\psi. \end{aligned}$$

Now, recall

$$[e_3, e_4] = 2\underline{\omega}e_4 - 2\underline{\omega}e_3 + 2(\underline{\eta} - \underline{\eta})e_\theta,$$

and, in view of Lemma 2.1.51, the following commutation formulae for reduced scalars

1. If $f \in \mathfrak{s}_k$,

$$\begin{aligned} [\not{d}_k, e_4] &= \frac{1}{2}\kappa \not{d}_k f + \text{Com}_k(f), \\ \text{Com}_k(f) &= -\frac{1}{2}\vartheta \not{d}_{k+1}^* f - (\zeta + \underline{\eta})e_4 f - k\underline{\eta}e_4 \Phi f - \xi(e_3 f + ke_3(\Phi)f) - k\beta f. \end{aligned}$$

2. If $f \in \mathfrak{s}_{k-1}$

$$\begin{aligned} [\not{d}_k^*, e_4]f &= \frac{1}{2}\kappa \not{d}_k f + \text{Com}_k^*(f), \\ \text{Com}_k^*(f) &= -\frac{1}{2}\vartheta \not{d}_{k-1} f + (\zeta + \underline{\eta})e_4 f - (k-1)\underline{\eta}e_4 \Phi f + \xi(e_3 f - (k-1)e_3(\Phi)f) \\ &\quad - (k-1)\beta f. \end{aligned}$$

We infer

$$\begin{aligned} [\square_2, e_4]\psi &= -e_4\left((2\underline{\omega}e_4 - 2\underline{\omega}e_3 + 2\underline{\eta}e_\theta)\psi\right) + \kappa\Delta_2\psi \\ &\quad + \left(2\underline{\omega} - \frac{1}{2}\kappa\right)\left(2\underline{\omega}e_4 - 2\underline{\omega}e_3 + 2\underline{\eta}e_\theta\right)\psi - \left(2e_4(\omega) + \frac{1}{4}\kappa^2 + \omega\kappa\right)e_3\psi \\ &\quad + \frac{1}{2}\left(-\frac{1}{2}\kappa\underline{\kappa} + 2\underline{\omega}\underline{\kappa} + 2\rho\right)e_4\psi + r^{-2}\mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi \\ &= 2e_4(\omega e_3\psi) + \kappa\Delta_2\psi - \left(2e_4(\omega) + \frac{1}{4}\kappa^2\right)e_3\psi - \frac{1}{4}\kappa\underline{\kappa}e_4\psi + O(r^{-4})\mathfrak{d}^{\leq 1}\psi + r^{-2}\text{Err} \\ &= 2\underline{\omega}e_4(e_3\psi) + \kappa\Delta_2\psi - \frac{1}{4}\kappa^2 e_3\psi - \frac{1}{4}\kappa\underline{\kappa}e_4\psi + O(r^{-4})\mathfrak{d}^{\leq 1}\psi + r^{-2}\text{Err}. \end{aligned}$$

This implies

$$\begin{aligned} [\square_2, e_4](r\psi) &= 2\underline{\omega}e_4(e_3(r\psi)) + \kappa\Delta_2(r\psi) - \frac{1}{4}\kappa^2 e_3(r\psi) - \frac{1}{4}\kappa\underline{\kappa}e_4(r\psi) + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err} \\ &= 2\underline{\omega}e_3(e_4(r\psi)) + 2\underline{\omega}[e_4, e_3]r\psi + \kappa\Delta_2(r\psi) - \frac{1}{4}\kappa^2 e_3(r\psi) - \frac{1}{4}\kappa\underline{\kappa}e_4(r\psi) \\ &\quad + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err} \\ &= 2\underline{\omega}e_3(e_4(r\psi)) + \kappa\Delta_2(r\psi) - \frac{1}{4}\kappa^2 e_3(r\psi) - \frac{1}{4}\kappa\underline{\kappa}e_4(r\psi) + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err} \end{aligned}$$

and hence

$$\begin{aligned}
& \square_2(e_4(r\psi)) - e_4(r\square_2\psi) \\
= & [\square_2, e_4](r\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_4\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_3\psi) + e_4\left(\frac{2}{r}\psi\right) + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err} \\
= & 2\omega e_3(e_4(r\psi)) - \frac{1}{2}e_4(r\underline{\kappa}e_4\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_3\psi) + \kappa\Delta_2(r\psi) - \frac{1}{4}\kappa^2 e_3(r\psi) \\
& - \frac{1}{4}\kappa\underline{\kappa}e_4(r\psi) + e_4\left(\frac{2}{r}\psi\right) + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err}.
\end{aligned}$$

Next, we compute

$$\begin{aligned}
& -\frac{1}{2}e_4(r\underline{\kappa}e_4\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_3\psi) + \kappa\Delta_2(r\psi) - \frac{1}{4}\kappa^2 e_3(r\psi) + e_4\left(\frac{2}{r}\psi\right) \\
= & -\frac{1}{2}e_4(\underline{\kappa}(e_4(r\psi) - e_4(r)\psi)) - \frac{1}{2}e_3(e_4(r\underline{\kappa}\psi)) - \frac{1}{2}[e_4, e_3](r\underline{\kappa}\psi) + \frac{1}{2}e_4(e_3(r\underline{\kappa})\psi) \\
& + r\underline{\kappa}\Delta_2\psi - \frac{1}{4}r\underline{\kappa}^2 e_3\psi - \frac{1}{4}\kappa^2 e_3(r)\psi + \frac{2}{r^2}e_4(r\psi) + e_4\left(\frac{2}{r^2}\right)r\psi + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi \\
= & -\frac{1}{2}e_4(\underline{\kappa}(e_4(r\psi))) + \frac{1}{2}e_4\left(\frac{r}{2}(\underline{\kappa} + A)\psi\right) - \frac{1}{2}e_3(\underline{\kappa}e_4(r\psi)) - \frac{1}{2}e_3(e_4(\underline{\kappa})r\psi) \\
& - \frac{1}{2}\left(-2\underline{\omega}e_4 + 2\omega e_3 - 2(\underline{\eta} - \underline{\eta})e_\theta\right)(r\underline{\kappa}\psi) + \frac{1}{2}e_4(e_3(r\underline{\kappa})\psi) \\
& + r\underline{\kappa}\Delta_2\psi - \frac{1}{4}r\underline{\kappa}^2 e_3\psi - \frac{1}{4}\kappa^2 \frac{r}{2}(\underline{\kappa} + \underline{A})\psi + \frac{2}{r^2}e_4(r\psi) - \frac{4e_4(r)}{r^2}\psi + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi
\end{aligned}$$

i.e.

$$\begin{aligned}
& -\frac{1}{2}e_4(r\underline{\kappa}e_4\psi) - \frac{1}{2}e_4(r\underline{\kappa}e_3\psi) + \kappa\Delta_2(r\psi) - \frac{1}{4}\kappa^2 e_3(r\psi) + e_4\left(\frac{2}{r}\psi\right) \\
= & -\frac{1}{2}e_4(\underline{\kappa}(e_4(r\psi))) + \frac{1}{4}e_4(r\underline{\kappa}\psi) - \frac{1}{2}e_3(\underline{\kappa}e_4(r\psi)) - \frac{1}{2}e_3\left(\left(-\frac{1}{2}\kappa^2 - 2\omega\underline{\kappa}\right)r\psi\right) \\
& - \omega e_3(r\underline{\kappa}\psi) + \frac{1}{2}e_4(e_3(r\underline{\kappa})\psi) + r\underline{\kappa}\Delta_2\psi - \frac{1}{4}r\underline{\kappa}^2 e_3\psi - \frac{1}{8}r\underline{\kappa}^2 \underline{\kappa}\psi + \frac{2}{r^2}e_4(r\psi) - \frac{2\underline{\kappa}}{r}\psi \\
& + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err} \\
= & -\frac{1}{2}e_4(\underline{\kappa}(e_4(r\psi))) + \frac{1}{4}\underline{\kappa}\underline{\kappa}e_4(r\psi) + \frac{1}{4}e_4(\underline{\kappa}\underline{\kappa})r\psi - \frac{1}{2}e_3(\underline{\kappa}e_4(r\psi)) - \frac{1}{2}e_3\left(\left(-\frac{1}{2}\kappa^2 - 2\omega\underline{\kappa}\right)r\psi\right) \\
& - \omega e_3(r\underline{\kappa}\psi) + \frac{1}{2}r^{-1}e_3(r\underline{\kappa})e_4(r\psi) + \frac{1}{2}e_4(r^{-1}e_3(r\underline{\kappa}))r\psi + r\underline{\kappa}\Delta_2\psi - \frac{1}{4}r\underline{\kappa}^2 e_3\psi - \frac{1}{8}r\underline{\kappa}^2 \underline{\kappa}\psi \\
& + \frac{2}{r^2}e_4(r\psi) - \frac{2\underline{\kappa}}{r}\psi + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err}.
\end{aligned}$$

We infer

$$\begin{aligned}
& \square_2(e_4(r\psi)) - e_4(r\square_2\psi) \\
&= 2\omega e_3(e_4(r\psi)) - \frac{1}{2}e_4(\underline{\kappa}(e_4(r\psi))) + \frac{1}{4}e_4(\underline{\kappa}\kappa)r\psi - \frac{1}{2}e_3(\kappa e_4(r\psi)) \\
&\quad - \frac{1}{2}e_3\left(\left(-\frac{1}{2}\kappa^2 - 2\omega\kappa\right)r\psi\right) - \omega e_3(r\kappa\psi) + \frac{1}{2}r^{-1}e_3(r\kappa)e_4(r\psi) + \frac{1}{2}e_4(r^{-1}e_3(r\kappa))r\psi \\
&\quad + r\kappa\Delta_2\psi - \frac{1}{4}r\kappa^2e_3\psi - \frac{1}{8}r\kappa^2\underline{\kappa}\psi + \frac{2}{r^2}e_4(r\psi) - \frac{2\kappa}{r}\psi + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err}.
\end{aligned}$$

Since $e_4(r\psi) = r\Upsilon\check{e}_4\psi$, this may be rewritten as

$$\begin{aligned}
& \square_2(r\Upsilon\check{e}_4\psi) - e_4(r\square_2\psi) \\
&= 2\omega e_3(r\Upsilon\check{e}_4\psi) - \frac{1}{2}e_4(r\Upsilon\underline{\kappa}\check{e}_4\psi) + \frac{1}{4}e_4(\underline{\kappa}\kappa)r\psi - \frac{1}{2}e_3(r\Upsilon\kappa\check{e}_4\psi) \\
&\quad - \frac{1}{2}e_3\left(\left(-\frac{1}{2}\kappa^2 - 2\omega\kappa\right)r\psi\right) - \omega e_3(r\kappa\psi) + \frac{1}{2}e_3(r\kappa)\Upsilon\check{e}_4\psi + \frac{1}{2}e_4(r^{-1}e_3(r\kappa))r\psi \\
&\quad + r\kappa\Delta_2\psi - \frac{1}{4}r\kappa^2e_3\psi - \frac{1}{8}r\kappa^2\underline{\kappa}\psi + \frac{2}{r}\Upsilon\check{e}_4\psi - \frac{2\kappa}{r}\psi + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err}.
\end{aligned}$$

Now, since

$$\square_2\psi = -e_3e_4\psi + \Delta_2\psi + \left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi - \frac{1}{2}\kappa e_3\psi + 2\eta e_\theta\psi,$$

we have

$$\begin{aligned}
r\kappa\Delta_2\psi &= r\kappa\square_2\psi + r\kappa e_3e_4\psi - r\kappa\left(2\underline{\omega} - \frac{1}{2}\underline{\kappa}\right)e_4\psi + \frac{1}{2}r\kappa^2e_3\psi + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_b)\mathfrak{d}^{\leq 1}\psi \\
&= r\kappa\square_2\psi + r\kappa e_3(r^{-1}e_4(r\psi)) - r\kappa e_3(r^{-1}e_4(r)\psi) + \frac{1}{2}\kappa\underline{\kappa}e_4(r\psi) \\
&\quad - \frac{1}{2}\kappa\underline{\kappa}e_4(r)\psi + \frac{1}{2}r\kappa^2e_3\psi + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_b)\mathfrak{d}^{\leq 1}\psi \\
&= r\kappa\square_2\psi + r\kappa e_3(\Upsilon\check{e}_4\psi) - \frac{1}{2}r\kappa e_3(\kappa)\psi + \frac{1}{2}\kappa\underline{\kappa}r\Upsilon\check{e}_4\psi - \frac{r}{4}\kappa^2\underline{\kappa}\psi + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_b)\mathfrak{d}^{\leq 1}\psi
\end{aligned}$$

and hence

$$\begin{aligned}
& \square_2(r\Upsilon\check{e}_4\psi) - e_4(r\square_2\psi) \\
&= 2\omega e_3(r\Upsilon\check{e}_4\psi) - \frac{1}{2}e_4(r\Upsilon\underline{\kappa}\check{e}_4\psi) + \frac{1}{4}e_4(\underline{\kappa}\kappa)r\psi - \frac{1}{2}e_3(r\Upsilon\kappa\check{e}_4\psi) \\
&\quad - \frac{1}{2}e_3\left(\left(-\frac{1}{2}\kappa^2 - 2\omega\kappa\right)r\psi\right) - \omega e_3(r\kappa\psi) + \frac{1}{2}e_3(r\kappa)\Upsilon\check{e}_4\psi + \frac{1}{2}e_4(r^{-1}e_3(r\kappa))r\psi \\
&\quad + r\kappa\square_2\psi + r\kappa e_3(\Upsilon\check{e}_4\psi) - \frac{1}{2}r\kappa e_3(\kappa)\psi - \frac{r}{4}\kappa^2\underline{\kappa}\psi - \frac{1}{4}r\kappa^2e_3\psi - \frac{1}{8}r\kappa^2\underline{\kappa}\psi \\
&\quad - \frac{2\kappa}{r}\psi + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err}.
\end{aligned}$$

Next, we compute

$$\begin{aligned}
& \frac{1}{4}e_4(\underline{\kappa}\kappa)r\psi - \frac{1}{2}e_3\left(\left(-\frac{1}{2}\kappa^2 - 2\omega\kappa\right)r\psi\right) - \omega e_3(r\kappa\psi) + \frac{1}{2}e_4(r^{-1}e_3(r\kappa))r\psi \\
& - \frac{1}{2}r\kappa e_3(\kappa)\psi - \frac{r}{4}\kappa^2\underline{\kappa}\psi - \frac{1}{4}r\kappa^2e_3\psi - \frac{1}{8}r\kappa^2\underline{\kappa}\psi - \frac{2\kappa}{r}\psi \\
& = \frac{r}{2}\kappa\rho\psi + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_b)\psi \\
& = O(r^{-3})\psi + r^{-1}\mathfrak{d}^{\leq 1}(\Gamma_b)\psi
\end{aligned}$$

so that

$$\begin{aligned}
\Box_2(r\Upsilon\check{e}_4\psi) &= e_4(r\Box_2\psi) + r\kappa\Box_2\psi + 2\omega e_3(r\Upsilon\check{e}_4\psi) - \frac{1}{2}e_4(r\Upsilon\underline{\kappa}\check{e}_4\psi) - \frac{1}{2}e_3(r\Upsilon\kappa\check{e}_4\psi) \\
& \quad + \frac{1}{2}e_3(r\kappa)\Upsilon\check{e}_4\psi + r\kappa e_3(\Upsilon\check{e}_4\psi) + O(r^{-3})\mathfrak{d}^{\leq 1}\psi + r^{-1}\text{Err}.
\end{aligned}$$

Since

$$\begin{aligned}
\Box_2(r^2\check{e}_4\psi) &= r\Upsilon^{-1}\Box_2(r\Upsilon\check{e}_4\psi) - e_3(r\Upsilon^{-1})e_4(r\Upsilon\check{e}_4\psi) - e_4(r\Upsilon^{-1})e_3(r\Upsilon\check{e}_4\psi) \\
& \quad + \Box_0(r\Upsilon^{-1})r\Upsilon\check{e}_4\psi + \mathfrak{d}^{\leq 1}(\Gamma_g)\mathfrak{d}^{\leq 2}\psi,
\end{aligned}$$

we infer

$$\begin{aligned}
\Box_2(r^2\check{e}_4\psi) &= r\Upsilon^{-1}e_4(r\Box_2\psi) + r^2\Upsilon^{-1}\kappa\Box_2\psi + 2r\Upsilon^{-1}\omega e_3(r\Upsilon\check{e}_4\psi) - \frac{1}{2}r\Upsilon^{-1}e_4(r\Upsilon\underline{\kappa}\check{e}_4\psi) \\
& \quad - \frac{1}{2}r\Upsilon^{-1}e_3(r\Upsilon\kappa\check{e}_4\psi) + \frac{1}{2}r e_3(r\kappa)\check{e}_4\psi + r^2\Upsilon^{-1}\kappa e_3(\Upsilon\check{e}_4\psi) \\
& \quad - e_3(r\Upsilon^{-1})e_4(r\Upsilon\check{e}_4\psi) - e_4(r\Upsilon^{-1})e_3(r\Upsilon\check{e}_4\psi) \\
& \quad + \Box_0(r\Upsilon^{-1})r\Upsilon\check{e}_4\psi + O(r^{-2})\mathfrak{d}^{\leq 1}\psi + \text{Err}.
\end{aligned}$$

Now, we have

$$\begin{aligned}
& 2r\Upsilon^{-1}\omega e_3(r\Upsilon\check{e}_4\psi) - \frac{1}{2}r\Upsilon^{-1}e_4(r\Upsilon\underline{\kappa}\check{e}_4\psi) \\
& - \frac{1}{2}r\Upsilon^{-1}e_3(r\Upsilon\kappa\check{e}_4\psi) + \frac{1}{2}r e_3(r\kappa)\check{e}_4\psi + r^2\Upsilon^{-1}\kappa e_3(\Upsilon\check{e}_4\psi) \\
& - e_3(r\Upsilon^{-1})e_4(r\Upsilon\check{e}_4\psi) - e_4(r\Upsilon^{-1})e_3(r\Upsilon\check{e}_4\psi) + \Box_0(r\Upsilon^{-1})r\Upsilon\check{e}_4\psi \\
& = 2r\frac{1-\frac{3m}{r}}{\Upsilon}e_4(\check{e}_4\psi) \\
& + \left\{ 2r\Upsilon^{-1}\omega e_3(r\Upsilon) - \frac{1}{2}r\Upsilon^{-1}e_4(r\Upsilon\underline{\kappa}) - \frac{1}{2}r\Upsilon^{-1}e_3(r\Upsilon\kappa) + \frac{1}{2}r e_3(r\kappa) + r^2\Upsilon^{-1}\kappa e_3(\Upsilon) \right. \\
& \quad \left. - e_3(r\Upsilon^{-1})e_4(r\Upsilon) - e_4(r\Upsilon^{-1})e_3(r\Upsilon) + \Box_0(r\Upsilon^{-1})r\Upsilon \right\} \check{e}_4\psi + \text{Err}.
\end{aligned}$$

Also, we have

$$\begin{aligned}
& 2r\Upsilon^{-1}\omega e_3(r\Upsilon) - \frac{1}{2}r\Upsilon^{-1}e_4(r\Upsilon\underline{\kappa}) - \frac{1}{2}r\Upsilon^{-1}e_3(r\Upsilon\kappa) + \frac{1}{2}re_3(r\kappa) + r^2\Upsilon^{-1}\kappa e_3(\Upsilon) \\
& - e_3(r\Upsilon^{-1})e_4(r\Upsilon) - e_4(r\Upsilon^{-1})e_3(r\Upsilon) + \square_0(r\Upsilon^{-1})r\Upsilon \\
= & 4 + O(r^{-1}) + r\Gamma_b \\
= & -r^2\underline{\kappa\underline{\kappa}} + O(r^{-1}) + r\Gamma_b.
\end{aligned}$$

We infer

$$\begin{aligned}
(\square_2 + \underline{\kappa\underline{\kappa}})(r^2\check{e}_4\psi) &= r\Upsilon^{-1}e_4(r\square_2\psi) + r^2\Upsilon^{-1}\kappa\square_2\psi + 2r\frac{1-\frac{3m}{r}}{\Upsilon}e_4(\check{e}_4\psi) \\
&+ O(r^{-2})\mathfrak{d}^{\leq 1}\psi + \text{Err}.
\end{aligned}$$

In view of the wee equation satisfied by ψ , i.e.

$$\square_2\psi + \underline{\kappa\underline{\kappa}}\psi = N,$$

we have

$$\begin{aligned}
& r\Upsilon^{-1}e_4(r\square_2\psi) + r^2\Upsilon^{-1}\kappa\square_2\psi + 2r\frac{1-\frac{3m}{r}}{\Upsilon}e_4(\check{e}_4\psi) \\
= & r\Upsilon^{-1}e_4(r(N - \underline{\kappa\underline{\kappa}}\psi)) + r^2\Upsilon^{-1}\kappa(N - \underline{\kappa\underline{\kappa}}\psi) + \frac{2}{r}\frac{1-\frac{3m}{r}}{\Upsilon}e_4(r^2\check{e}_4\psi) - 4\frac{1-\frac{3m}{r}}{\Upsilon}e_4(r)\check{e}_4\psi \\
= & r\Upsilon^{-1}e_4(rN) + r^2\Upsilon^{-1}\kappa N + \frac{2}{r}\frac{1-\frac{3m}{r}}{\Upsilon}e_4(r^2\check{e}_4\psi) + \frac{4m}{r}\check{e}_4\psi - 2r^2\Upsilon^{-1}\kappa\rho\psi + \mathfrak{d}^{\leq 1}(\Gamma_b)\mathfrak{d}^{\leq 1}\psi \\
= & r^2\left(\Upsilon^{-1}e_4(N) + \frac{3}{r}N\right) + \frac{2}{r}\frac{1-\frac{3m}{r}}{\Upsilon}e_4(r^2\check{e}_4\psi) + O(r^{-2})\mathfrak{d}^{\leq 1}\psi + \mathfrak{d}^{\leq 1}(\Gamma_b)\mathfrak{d}^{\leq 1}\psi,
\end{aligned}$$

from which we deduce

$$(\square_2 + \underline{\kappa\underline{\kappa}})(r^2\check{e}_4\psi) = r^2\left(\Upsilon^{-1}e_4(N) + \frac{3}{r}N\right) + \frac{2}{r}\frac{1-\frac{3m}{r}}{\Upsilon}e_4(r^2\check{e}_4\psi) + O(r^{-2})\mathfrak{d}^{\leq 1}\psi + \text{Err}.$$

Since

$$\check{\psi} = f_2\check{e}_4\psi = \frac{f_2}{r^2}r^2\check{e}_4\psi,$$

we infer

$$\begin{aligned}
(\square_2 + \underline{\kappa\underline{\kappa}})\check{\psi} &= \frac{f_2}{r^2}(\square_2 + \underline{\kappa\underline{\kappa}})(r^2\check{e}_4\psi) - e_3\left(\frac{f_2}{r^2}\right)e_4(r^2\check{e}_4\psi) - e_4\left(\frac{f_2}{r^2}\right)e_3(r^2\check{e}_4\psi) \\
&+ e_\theta\left(\frac{f_2}{r^2}\right)\not{d}_2(r^2\check{e}_4\psi) - e_\theta\left(\frac{f_2}{r^2}\right)\not{d}_3^*(r^2\check{e}_4\psi) + \square_0\left(\frac{f_2}{r^2}\right)r^2\check{e}_4\psi
\end{aligned}$$

and hence

$$\begin{aligned}
(\square_2 + \kappa \underline{\kappa}) \check{\psi} &= f_2 \left(\Upsilon^{-1} e_4(N) + \frac{3}{r} N \right) + \frac{f_2}{r^2} \left\{ \frac{2}{r} \frac{1 - \frac{3m}{r}}{\Upsilon} e_4(r^2 \check{e}_4 \psi) + O(r^{-2}) \mathfrak{d}^{\leq 1} \psi + \text{Err} \right\} \\
&\quad - e_3 \left(\frac{f_2}{r^2} \right) e_4(r^2 \check{e}_4 \psi) - e_4 \left(\frac{f_2}{r^2} \right) e_3(r^2 \check{e}_4 \psi) \\
&\quad + e_\theta \left(\frac{f_2}{r^2} \right) \not{d}_2(r^2 \check{e}_4 \psi) - e_\theta \left(\frac{f_2}{r^2} \right) \not{d}_3^*(r^2 \check{e}_4 \psi) + \square_0 \left(\frac{f_2}{r^2} \right) r^2 \check{e}_4 \psi.
\end{aligned}$$

Now, recall that Err is defined by

$$\text{Err} = r^2 \Gamma_g e_4 e_3 \psi + r \Gamma_b e_4 \mathfrak{d} \psi + \mathfrak{d}^{\leq 1}(\Gamma_b) \mathfrak{d}^{\leq 1} \psi + r \mathfrak{d}^{\leq 1}(\Gamma_g) e_3 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^2 \psi.$$

so that

$$\begin{aligned}
(\square_2 + \kappa \underline{\kappa}) \check{\psi} &= f_2 \left(\Upsilon^{-1} e_4(N) + \frac{3}{r} N \right) + \frac{f_2}{r^2} \left\{ \frac{2}{r} \frac{1 - \frac{3m}{r}}{\Upsilon} e_4(r^2 \check{e}_4 \psi) + O(r^{-2}) \mathfrak{d}^{\leq 1} \psi + r^2 \Gamma_g e_4 e_3 \psi \right. \\
&\quad \left. + r \Gamma_b e_4 \mathfrak{d} \psi + \mathfrak{d}^{\leq 1}(\Gamma_b) \mathfrak{d}^{\leq 1} \psi + r \mathfrak{d}^{\leq 1}(\Gamma_g) e_3 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^2 \psi \right\} \\
&\quad - e_3 \left(\frac{f_2}{r^2} \right) e_4(r^2 \check{e}_4 \psi) - e_4 \left(\frac{f_2}{r^2} \right) e_3(r^2 \check{e}_4 \psi) \\
&\quad + e_\theta \left(\frac{f_2}{r^2} \right) \not{d}_2(r^2 \check{e}_4 \psi) - e_\theta \left(\frac{f_2}{r^2} \right) \not{d}_3^*(r^2 \check{e}_4 \psi) + \square_0 \left(\frac{f_2}{r^2} \right) r^2 \check{e}_4 \psi.
\end{aligned}$$

In view of

$$\square_2 \psi = -e_4 e_3 \psi + \not{A}_2 \psi + \left(2\omega - \frac{1}{2} \kappa \right) e_3 \psi - \frac{1}{2} \underline{\kappa} e_4 \psi + 2 \underline{\eta} e_\theta \psi,$$

we have

$$\begin{aligned}
r^2 \Gamma_g e_4 e_3 \psi &= r^2 \Gamma_g \left(-\square_2 \psi + \not{A}_2 \psi + \left(2\omega - \frac{1}{2} \kappa \right) e_3 \psi - \frac{1}{2} \underline{\kappa} e_4 \psi + 2 \underline{\eta} e_\theta \psi \right) \\
&= -r^2 \Gamma_g N + r \Gamma_g e_3 \psi + \Gamma_g \mathfrak{d}^{\leq 2} \psi
\end{aligned}$$

and hence

$$\begin{aligned}
(\square_2 + \kappa_{\underline{\kappa}}) \check{\psi} &= f_2 \left(\Upsilon^{-1} e_4(N) + \frac{3}{r} N \right) + \frac{f_2}{r^2} \left\{ \frac{2}{r} \frac{1 - \frac{3m}{r}}{\Upsilon} e_4(r^2 \check{e}_4 \psi) + O(r^{-2}) \mathfrak{d}^{\leq 1} \psi \right. \\
&\quad \left. + r \Gamma_b e_4 \mathfrak{d} \psi + \mathfrak{d}^{\leq 1}(\Gamma_b) \mathfrak{d}^{\leq 1} \psi + r \mathfrak{d}^{\leq 1}(\Gamma_g) e_3 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^2 \psi \right\} \\
&\quad - e_3 \left(\frac{f_2}{r^2} \right) e_4(r^2 \check{e}_4 \psi) - e_4 \left(\frac{f_2}{r^2} \right) e_3(r^2 \check{e}_4 \psi) \\
&\quad + e_\theta \left(\frac{f_2}{r^2} \right) \not\phi_2(r^2 \check{e}_4 \psi) - e_\theta \left(\frac{f_2}{r^2} \right) \not\phi_3^*(r^2 \check{e}_4 \psi) + \square_0 \left(\frac{f_2}{r^2} \right) r^2 \check{e}_4 \psi.
\end{aligned}$$

In particular, we have for $r \geq 6m_0$

$$\begin{aligned}
(\square_2 + \kappa_{\underline{\kappa}}) \check{\psi} &= r^2 \left(\Upsilon^{-1} e_4(N) + \frac{3}{r} N \right) + \frac{2}{r \Upsilon} \left(1 - \frac{3m}{r} \right) e_4 \check{\psi} \\
&\quad + O(r^{-2}) \mathfrak{d}^{\leq 1} \psi + r \Gamma_b e_4 \mathfrak{d} \psi + \mathfrak{d}^{\leq 1}(\Gamma_b) \mathfrak{d}^{\leq 1} \psi + r \mathfrak{d}^{\leq 1}(\Gamma_g) e_3 \psi + \mathfrak{d}^{\leq 1}(\Gamma_g) \mathfrak{d}^2 \psi
\end{aligned}$$

and for $4m_0 \leq r \leq 6m_0$,

$$(\square_2 + \kappa_{\underline{\kappa}}) \check{\psi} = f_2 \left(\Upsilon^{-1} e_4(N) + \frac{3}{r} N \right) + O(1) \mathfrak{d}^2 \psi.$$

This concludes the proof Proposition 10.3.1.

Bibliography

- [1] S. Aksteiner, private communication.
- [2] S. Aksteiner, L. Andersson, T. Bäckdahl, A. G. Shah and B. F. Whiting, *Gauge-invariant perturbations of Schwarzschild spacetime*, arXiv:1611.08291.
- [3] L. Andersson and P. Blue, *Hidden symmetries and decay for the wave equation on the Kerr spacetime*, arXiv:0908.2265, to appear in *Annals of Math*.
- [4] Y. Angelopoulos, S. Aretakis and D. Gajic, *A vector field approach to almost-sharp decay for the wave equation on spherically symmetric, stationary spacetimes*, arXiv:1612.01565.
- [5] J. M. Bardeen and W. H. Press, *Radiation fields in the Schwarzschild background*, *J. Math. Phys.*, 14 (1973), pp. 719.
- [6] L. Bieri, *An extension of the stability theorem of the Minkowski space in general relativity*, Thesis, ETH, 2007.
- [7] P. Blue and A. Soffer, *Semilinear wave equations on the Schwarzschild manifold. I. Local decay estimates*, *Adv. Differential Equations* **8** (2003), 595–614.
- [8] P. Blue and A. Soffer, *Errata for “Global existence and scattering for the nonlinear Schrödinger equation on Schwarzschild manifolds”*, “*Semilinear wave equations on the Schwarzschild manifold I: Local Decay Estimates*”, and “*The wave equation on the Schwarzschild metric II: Local Decay for the spin 2 Regge Wheeler equation*”, gr-qc/0608073, 6 pages.
- [9] P. Blue and A. Soffer, *A space-time integral estimate for a large data semi-linear wave equation on the Schwarzschild manifold*, *Lett. Math. Phys.* **81** (2007), 227–238.
- [10] P. Blue and J. Sterbenz, *Uniform decay of local energy and the semi-linear wave equation on Schwarzschild space*, *Comm. Math. Phys.* **268** (2006), 481–504.

- [11] S. Chandrasekhar, *The mathematical theory of black holes*, 1983, Oxford Classic Texts in the Physical Sciences.
- [12] S. Chandrasekhar, *On the equations governing the perturbations of the Schwarzschild black hole*, P. Roy. Soc. Lond. A Mat., 343 (1975), pp. 289–298.
- [13] Y. Choquet-Bruhat, *Théorème d'existence pour certains systèmes d'équations aux dérivées partielles non linéaires*, Acta Math. 88, 141–225 (1952).
- [14] Y. Choquet-Bruhat and R Geroch, *Global Aspects of the Cauchy Problem in General Relativity* Commun. Math. Phys. 14, 329–335 (1969).
- [15] D. Christodoulou, *Global solutions of nonlinear hyperbolic equations for small initial data*, Comm. Pure Appl. Math. **39**, 267–282 (1986).
- [16] D. Christodoulou and S. Klainerman, *Asymptotic properties of linear field theories in Minkowski space*, Communications on Pure and Applied Mathematics, Vol. 43, 137–199 (1990).
- [17] D. Christodoulou and S. Klainerman, *The global nonlinear stability of the Minkowski space*, Princeton University Press (1993).
- [18] M. Dafermos, *The mathematical analysis of black holes in general relativity*, ICM, Seoul 2014.
- [19] M. Dafermos and I. Rodnianski, *The red-shift effect and radiation decay on black hole spacetimes*, Comm. Pure Appl. Math. **62** (2009), 859–919.
- [20] M. Dafermos and I. Rodnianski, *A new physical-space approach to decay for the wave equation with applications to black hole spacetimes*, XVIIth International Congress on Mathematical Physics, World Sci. Publ., Hackensack, NJ, 2010, pp. 421–432.
- [21] M. Dafermos and I. Rodnianski, *A proof of the uniform boundedness of solutions to the wave equation on slowly rotating Kerr backgrounds*, Invent. Math. **185** (2011), 467–559.
- [22] M. Dafermos, G. Holzegel and I. Rodnianski, *Linear stability of the Schwarzschild solution to gravitational perturbations*, arXiv:1601.06467.
- [23] M. Dafermos, G. Holzegel, and I. Rodnianski, *A scattering theory construction of dynamical black hole spacetimes*, arXiv:1306.5534, to appear in J. Diff. Geom.
- [24] M. Dafermos, I. Rodnianski, and Y. Shlapentokh-Rothman, *Decay for solutions of the wave equation on Kerr exterior spacetimes iii: The full subextremal case $|a| < m$* , arxiv:1402.7034.

- [25] E. Giorgi, *An Introduction to Coupled gravitational and electromagnetic perturbations of Reissner-Nordstrom spacetime in a polarized setting*, preprint.
- [26] P. Hintz and A. Vasy, *The global non-linear stability of the Kerr-de Sitter family of black holes*, arXiv:1606.04014.
- [27] P. Hintz, *Non-linear stability of the Kerr-Newman-de Sitter family of charged black holes*, arXiv:1612.04489.
- [28] P. Hintz and A. Vasy, *A global analysis proof of the stability of Minkowski space and the polyhomogeneity of the metric*, arXiv:1711.00195.
- [29] G. Holzegel, *Ultimately Schwarzschildian Spacetimes and the Black Hole Stability Problem*, arXiv:1010.3216.
- [30] P. K. Hung, J. Keller and M. T. Wang, *Linear stability of Schwarzschild spacetime: the Cauchy problem of metric coefficients*, arXiv:1702.02843.
- [31] A. D. Ionescu and S. Klainerman, *On the global stability of the wave-map equation in Kerr spaces with small angular momentum*, *Annals of PDE* **1**, 1–78 (2015).
- [32] S. Klainerman, *Long time behavior of solutions to nonlinear wave equations*, “Proceedings of the ICM”, (Warsaw, 1982), 1209–1215.
- [33] S. Klainerman, *Uniform decay estimates and the Lorentz invariance of the classical wave equations*, *Communications on Pure and Applied Mathematics*, **38** (1985), 321–332.
- [34] S. Klainerman, *The Null Condition and global existence to nonlinear wave equations*, *Lect. in Appl. Math.*, **23**, (1986) 293–326.
- [35] S. Klainerman, *Remarks on the global Sobolev inequalities*, *Comm. Pure and Applied Math.*, Vol. 40, 111–117 (1987).
- [36] S. Klainerman, *PDE as a unified subject*. GAFA 2000 (Tel Aviv 1999), *Geom. Funct. Anal.* **2000**, Special Volume, Part 1, 279–315.
- [37] S. Klainerman and F. Nicolò, *The evolution problem in general relativity*. *Progress in Mathematical Physics* **25**, Birkhauser, Boston, 2003, +385 pp.
- [38] S. Klainerman and F. Nicolò, *Peeling properties of asymptotic solutions to the Einstein vacuum equations*, *Classical Quantum Gravity* **20** (2003), no 14, 3215–3257.
- [39] H. Lindblad and I. Rodnianski, *The global stability of Minkowski space-time in harmonic gauge*, *Annals of Math.*, **171**, 2010, pp 1401–1477.

- [40] J. Luk, *The Null Condition and Global Existence for Nonlinear Wave Equations on Slowly Rotating Kerr Spacetimes*, *Annales Henri Poincaré*, **11** (2010), 805–880.
- [41] S. Ma, *Uniform energy bound and Morawetz estimate for extreme component of spin fields in the exterior of a slowly rotating black hole II: Linearized gravity*, arXiv:1708.07385.
- [42] M. Mars, *A spacetime characterization of the Kerr metric*, *Classical Quantum Gravity* **16** (1999), 2507–2523.
- [43] M. Mars and J. Senovilla, *Axial Symmetry and conformal Killing vectorfields*, arXiv:gr-qc/0201045.
- [44] Y. Martel and F. Merle, *Asymptotic stability of solitons for subcritical generalized KdV equations*, *Arch. Ration. Mech. Anal.* **157** (2001), no. 3, 219–254.
- [45] J. Marzuola, J. Metcalfe, D. Tataru and M. Tohaneanu, *Strichartz estimates on Schwarzschild black hole backgrounds*, *Comm. Math. Phys.* **293** (2010), 37–83.
- [46] F. Merle and P. Raphaël, *On universality of blow-up profile for L^2 critical nonlinear Schrödinger equation*, *Invent. Math.* **156** (2004), no. 3, 565–672.
- [47] V. Moncrief, *Gravitational perturbations of spherically symmetric systems. I. The exterior problem*, *Ann. Phys.*, **88** (1975), pp. 323–342.
- [48] W. Press and S. Teukolsky, *Perturbations of a rotating black hole. II. Dynamical stability of the Kerr metric* *Astrophys. J.* **185** (1973), 649–673.
- [49] T. Regge and J. A. Wheeler, *Stability of a Schwarzschild singularity*. *Phys. Rev.* (2), **108**:1063–1069, 1957.
- [50] J. Sbierski, *On the Existence of a Maximal Cauchy Development for the Einstein Equations - a Dezornification*, *Annales Henri Poincaré* February 2016, Volume 17, Issue 2, pp 301–329.
- [51] Y. Shlapentokh-Rothman, *Quantitative Mode Stability for the Wave Equation on the Kerr Spacetime*, *Annales Henri Poincaré*. **16**, 2013, 10.1007/s00023-014-0315-7.
- [52] V. Schlue, *Decay of linear waves on higher dimensional Schwarzschild black holes*, *Analysis & PDE* **6**, (2013), 515–600.
- [53] J. Stogin, *Princeton PHD thesis*, 2017.
- [54] D. Tataru and M. Tohaneanu, *A local energy estimate on Kerr black hole backgrounds*, *IMRN* (2011), 248–292.

- [55] S. A. Teukolsky, *Perturbations of a rotating black hole. I. Fundamental equations for gravitational, electromagnetic, and neutrino-field perturbations*, *Astrophys. J.*, 185 (1973), pp. 635–648.
- [56] C. V. Vishveshwara, *Stability of the Schwarzschild metric*, *Phys. Rev. D*, 1 (1970), 2870–2879.
- [57] R. M. Wald, *Construction of Solutions of Gravitational, Electromagnetic, or other Perturbation Equations from Solutions of Decoupled Equations*, *Phys. Rev. Lett.*, 41, 203–206, 1978.
- [58] G. Weinstein, *On rotating black holes in equilibrium in general relativity*, *Comm. Pure Appl. Math.* **43** (1990), 903–948.
- [59] B. Whiting, *Mode stability of the Kerr black hole*, *J. Math. Phys.* 30 (1989), 1301–1305.
- [60] F. J. Zerilli, *Effective potential for even-parity Regge-Wheeler gravitational perturbation equations*, *Phys. Rev. Lett.*, 24 (1970), pp. 737–738.