# NONABELIAN HODGE THEORY FOR KLT SPACES AND DESCENT THEOREMS FOR VECTOR BUNDLES

DANIEL GREB, STEFAN KEBEKUS, THOMAS PETERNELL, AND BEHROUZ TAJI

ABSTRACT. We generalise Simpson's nonabelian Hodge correspondence to the context of projective varieties with klt singularities. The proof relies on a descent theorem for numerically flat vector bundles along birational morphisms. In its simplest form, this theorem asserts that given any klt variety X and any resolution of singularities, then any vector bundle on the resolution that appears to come from X numerically, does indeed come from X. Furthermore and of independent interest, a new restriction theorem for semistable Higgs sheaves defined on the smooth locus of a normal, projective variety is established.

### Contents

1.	Introduction	1
2.	Notation and elementary facts	3
3.	The nonabelian Hodge correspondence for locally free Higgs sheaves	9
4.	Descent of vector bundles from klt spaces	11
5.	Descent of vector bundles to klt spaces	14
6.	The restriction theorem for semistable Higgs sheaves	16
7.	Ascent of semistable Higgs bundles	19
8.	Proof of the nonabelian Hodge correspondence for klt spaces	26
References		29

#### 1. Introduction

1.1. Nonabelian Hodge theory for singular spaces. Given a projective manifold X, a seminal result of Simpson, [Sim92], exhibits a natural equivalence between the category of local systems and the category of semistable, locally free Higgs sheaves with vanishing Chern classes. The first main result of this paper extends this correspondence to projective varieties with Kawamata log terminal (=klt) singularities.

**Theorem 1.1** (Nonabelian Hodge correspondence for klt spaces, Theorem 3.4). Let X be a projective, klt variety. Then, there exists an equivalence between the category of local systems and the category of semistable, locally free Higgs sheaves with vanishing Chern classes.  $\Box$ 

We refer the reader to [GKPT15, Sect. 5] or to the survey [GKT18, Sect. 6] for the (rather delicate) notions of Higgs sheaves, morphisms of Higgs sheaves and pull-back. Semistability is also discussed there.

1

Date: 6th November 2018.

<sup>2010</sup> Mathematics Subject Classification. 14E30, 53C07.

Key words and phrases. Nonabelian Hodge Theory, KLT Singularities.

DG was partially supported by the DFG-Collaborative Research Center SFB/TR 45. SK gratefully acknowledges support through a joint fellowship of the Freiburg Institute of Advanced Studies (FRIAS) and the University of Strasbourg Institute for Advanced Study (USIAS). BT was partially supported by the DFG-Research Training Group GK1821. Research was partially completed while SK and TP were visiting the National University of Singapore in 2017. TP was supported by the DFG project "Zur Positivität in der komplexen Geometrie".

In fact, there exists a unique way to choose the correspondences of Theorem 1.1 that makes them functorial in morphisms between klt spaces, and compatible with Simpson's construction wherever this makes sense; we refer to Section 3 on page 9 for a precise formulation. In particular, functoriality applies to resolutions of singularities, as well as morphisms whose images are entirely contained in the singular loci of the target varieties. Our results imply that the pull-back of semistable, locally free Higgs sheaves with vanishing Chern classes under any of these maps remains semistable.

1.2. **Descent of vector bundles.** The proof of Theorem 1.1 relies in part on a descent theorem for vector bundles which is of independent interest. To put the result into perspective, consider a desingularisation  $\pi:\widetilde{X}\to X$  of a normal variety. It is well-known that if X has rational singularities, then any line bundle on  $\widetilde{X}$  that comes from X topologically does in fact come from X holomorphically. If X is klt, we generalise this result to vector bundles of arbitrary rank: we show that any vector bundle on  $\widetilde{X}$  that appears to come from X numerically does indeed come from X.

**Theorem 1.2** (Descent of vector bundles to klt spaces, Theorem 5.1). Let  $f: X \to Y$  be a birational, projective morphism of normal, quasi-projective varieties. Assume that there exists a Weil  $\mathbb{Q}$ -divisor  $\Delta_Y$  such that  $(Y, \Delta_Y)$  is klt. If  $\mathscr{F}_X$  is any locally free, f-numerically flat sheaf on X, then there exists a locally free sheaf  $\mathscr{F}_Y$  on Y such that  $\mathscr{F}_X \cong f^*\mathscr{F}_Y$ .  $\square$ 

The notion of "numerical flatness for vector bundles" generalises the notion of "numerical triviality" for line bundles and is recalled in Definition 2.11 below. The importance of Theorem 1.2 in the current investigation stems from the fact that locally free Higgs sheaves coming from local systems on resolutions of klt spaces are numerically flat relative to the resolution morphism; see Proposition 7.9.

In addition to the descent result for vector bundles spelled out above, Theorem 5.1 also discusses descent of locally free Higgs sheaves. Theorem 4.1 contains a related result where X (rather than Y) is assumed to be klt.

1.3. **Optimality of results.** We expect that varieties with klt singularities form the largest natural class where our results can possibly hold in full generality. The construction of a pull-back functor for Higgs sheaves is rather delicate and hinges on the existence of functorial pull-back for reflexive differentials, for morphisms between klt spaces. For classes of varieties with singularities that are just slightly more general than klt, there are elementary examples, [Keb13, Ex. 1.9], which show that *functorial* pull-back for reflexive differentials does not exist, even though it is known that reflexive differentials still lift to resolutions of singularities in these cases, [GK14, Thm. 1.4]. In particular, it is *not possible* to define functorial pull-back of Higgs sheaves for these spaces.

For spaces with arbitrary singularities, we do not expect that a correspondence between the two categories in Theorem 1.1 holds, even at the level of objects.

1.4. **Applications.** Theorem 1.1 has applications to the quasi-étale uniformisation problem for minimal varieties of general type. Eventually, we expect that all uniformisation theorems of nonabelian Hodge theory have analogues for klt varieties. In particular we expect to generalise the uniformisation result [GKPT15, Thm. 1.2] to arbitrary klt varieties: if X is minimal, klt and of general type, and if equality holds in the  $\mathbb{Q}$ -Miyaoka-Yau inequality, [GKPT15, Thm. 1.1], i.e.,

$$(2(n+1)\cdot\widehat{c}_2(\mathscr{T}_X)-n\cdot\widehat{c}_1(\mathscr{T}_X)^2)\cdot [K_X]^{n-2}=0,$$

then the canonical model of X is a singular ball quotient.

To keep the current paper reasonably short, these applications will appear in a separate paper, cf. [GKPT18]. Please see the survey paper [GKT18] for related results in this direction.

One might expect similar uniformisation results in more general contexts, such as "pairs" or "orbifolds". In the setting of pairs, the Miyaoka-Yau inequality has already been established in [GT16]. We refer to [GKPT15, Sect. 10] for a more precise formulation.

- 1.5. **Relation to the work of Mochizuki.** In a large body of work, T. Mochizuki set up a complete theory of Higgs bundles on arbitrary smooth quasi-projective varieties; among others, see [Moc06, Moc07a, Moc07b]. Our study differs from Mochizuki's in at least two aspects: First, while our main result, Theorem 1.1 above, traces a correspondence between topological and algebro-geometric properties of X, thereby taking the singularities of X into account, the correspondence established by Mochizuki in the present setup would focus on the connection between such properties for the smooth locus  $X_{\text{reg}}$ . Second, ultimately our correspondence is induced geometrically from the nonabelian Hodge correspondence in the projective case, which in turn requires much less sophisticated analytic results than Mochizuki's approach. However, Mochizuki's theory will be used in the sequel paper [GKPT18].
- 1.6. **Structure of the paper.** Section 2 gathers notation, known results and global conventions that will be used throughout the paper. Section 3 formulates the nonabelian Hodge correspondence for klt spaces in detail, discusses functoriality and its consequence, and states a number of singular generalisations of Simpson's classical results.

The results are then proven in the remaining sections. Sections 4 and 5 prepare for the proof, establishing the descent theorems for vector bundles mentioned in 1.2 above. Section 6 establishes a restriction theorem of Mehta-Ramanathan type for Higgs sheaves that is slightly more general than the versions found in the literature. This restriction theorem is then used in Section 7 to prove that Higgs bundles with vanishing Chern classes on projective, klt varieties that are semistable with respect to an ample divisor, remain semistable with respect to *ample* divisors when pulled back to a resolution of singularities. With these preparations in place, the nonabelian Hodge correspondences can then be constructed in Section 8 without much effort.

1.7. **Acknowledgements.** We would like to thank numerous colleagues for discussions, including Daniel Barlet, Oliver Bräunling, Andreas Höring, Annette Huber, Shane Kelly, Jong-Hae Keum, Adrian Langer, and Jörg Schürmann. We thank the referee for carefully reading the paper, and for her or his very valuable and interesting remarks.

#### 2. Notation and elementary facts

- 2.1. **Global conventions.** Throughout the present paper, all varieties and schemes will be defined over the complex numbers. We will freely switch between the algebraic and analytic context if no confusion is likely to arise. Apart from that, we follow the notation used in the standard reference books [Har77, KM98], with the exception that for klt pairs  $(X, \Delta)$ , the boundary divisor  $\Delta$  is always assumed to be effective. Varieties are always assumed to be irreducible and reduced.
- 2.2. **Varieties, sets and morphisms.** Normal varieties are  $S_2$ , which implies that regular functions can be extended across sets of codimension two. The following notation will be used.

Notation 2.1 (Big and small subsets). Let X be a normal, quasi-projective variety. A Zariski-closed subset  $Z \subset X$  is called small if  $codim_X Z \geq 2$ . A Zariski-open subset  $U \subseteq X$  is called big if  $X \setminus U$  is small. A birational morphism  $\varphi: X \to Y$  of normal, projective varieties is called a small morphism if there exists a big open set  $X^{\circ} \subseteq X$  such that  $\varphi|_{X^{\circ}}$  is an open immersion.

*Notation* 2.2 (Set-theoretic fibre). Given a morphism of varieties  $\varphi: X \to Y$  and a point  $y \in Y$ , we call the reduced scheme  $\varphi^{-1}(y)_{\text{red}}$  the *set-theoretic fibre* of  $\varphi$  over y.

**Definition 2.3** (Covers and covering maps, Galois morphisms). A cover or covering map is a finite, surjective morphism  $\gamma: Y \to X$  of normal, quasi-projective varieties or complex spaces. The covering map  $\gamma$  is called Galois if there exists a finite group  $G \subseteq \operatorname{Aut}(Y)$  such that X is isomorphic to the quotient map  $Y \to Y/G$ .

**Definition 2.4** (Quasi-étale morphisms). A morphism  $f: X \to Y$  between normal varieties is called quasi-étale if f is of relative dimension zero and étale in codimension one. In other words, f is quasi-étale if  $\dim X = \dim Y$  and if there exists a closed, subset  $Z \subseteq X$  of codimension  $\operatorname{codim}_X Z \ge 2$  such that  $f|_{X \setminus Z} : X \setminus Z \to Y$  is étale.

*Note.* Combining Definitions 2.3 and 2.4, a quasi-étale cover is finite, surjective, and étale in codimension one.

2.3. **Line bundles and sheaves.** Reflexive sheaves are in many ways easier to handle than arbitrary coherent sheaves, and we will therefore frequently take reflexive hulls. The following notation will be used.

*Notation* 2.5 (Reflexive hull). Given a normal, quasi-projective variety X and a coherent sheaf  $\mathscr E$  on X of rank r, write

$$\Omega_X^{[p]} := \left(\Omega_X^p\right)^{**}, \quad \mathscr{E}^{[m]} := \left(\mathscr{E}^{\otimes m}\right)^{**}, \quad \operatorname{Sym}^{[m]}\mathscr{E} := \left(\operatorname{Sym}^m\mathscr{E}\right)^{**}$$

and det  $\mathscr{E} := (\wedge^r \mathscr{E})^{**}$ . Given any morphism  $f: Y \to X$ , write  $f^{[*]}\mathscr{E} := (f^*\mathscr{E})^{**}$ .

**Definition 2.6** (Relative Picard number). Given a projective surjection  $f: X \to Y$  of normal, quasi-projective varieties, let  $N_1(X/Y)$  be the  $\mathbb{R}$ -vector space generated by irreducible curves  $C \subseteq X$  such that f(C) is a point, modulo numerical equivalence. The dimension of  $N_1(X/Y)$  is called the relative Picard number of X/Y and is denoted by  $\rho(X/Y)$ . Let  $\overline{NE}(X/Y) \subseteq N_1(X/Y)$  be the closed cone generated by classes of effective curves which are contracted by f.

2.4. **Cycles.** The Chow variety represents a functor. The associated notion of "families of cycles" is however somewhat awkward to formulate. For the sake of simplicity, we restrict ourselves to families over normal base varieties where the definition becomes somewhat simpler. The book [BM14] discusses these matters in detail.

Notation 2.7 (Families of cycles). Let X be a quasi-projective, n-dimensional variety, not necessarily normal. Given any subscheme  $Y \subseteq X$ , we denote the associated cycle by [Y]. Let  $f: X \to Y$  be an equidimensional morphism of normal, algebraic varieties, of relative dimension d. Recall from [Kol96, I Thm. 3.17] or [BM14, Thm. 3.4.1 on p. 449] that f is then a well defined family of d-dimensional proper algebraic cycles over Y, in the sense of [Kol96, I Def. 3.10]. In particular, if  $g \in Y$  is any closed point, Kollár defines in [Kol96, I Def. 3.10.4] the associated cycle-theoretic fibre, which we denote by  $f^{[-1]}(y)$ .

Warning 2.8. In the setting of Notation 2.7, it is generally *not* true that the cycle-theoretic fibre  $f^{[-1]}(y)$  is the cycle associated with the scheme-theoretic fibre  $f^{-1}(y)$ . Using the notation introduced above, the cycles  $f^{[-1]}(y)$  and  $[f^{-1}(y)]$  do not agree in general. An example is discussed in the preprint version of this paper.

Reminder 2.9 (Pull-back of Weil divisors). For arbitrary morphisms  $f: X \to Y$  normal, projective varieties, there is generally no good notion of pull-back for Weil-divisors. If f is finite, or more generally equidimensional, then a good pull-back map exists. We refer to [CKT16, Sect. 2.4.1] for a discussion.

2.5. **Numerical classes.** We briefly fix our notation for numerical classes and intersection numbers. The following definition allows to discuss slope and stability for arbitrary sheaves on arbitrarily singular spaces. We refer to [GKP16, Sect. 4.1] for a more detailed discussion.

Notation 2.10 (Numerical classes and intersection numbers). Let X be a projective variety. If  $\mathscr{L} \in \operatorname{Pic}(X)$  is invertible and  $D \in \operatorname{Div}(X)$  a Cartier divisor, we denote the numerical classes in  $N^1(X)_{\mathbb{R}}$  by  $[\mathscr{L}]$  and [D]; see  $[\operatorname{Kol96}]$ , Sect. II.4] for a brief definition and discussion of the space  $N^1(X)_{\mathbb{R}}$  of numerical Cartier divisor classes. If A is any purely d-dimensional cycle on X, we denote the intersection number with numerical classes of invertible sheaves  $\mathscr{L}_i$  by  $[\mathscr{L}_1] \cdots [\mathscr{L}_d] \cdot A \in \mathbb{Z}$ . For brevity of notation we will often write  $[\mathscr{L}_1] \cdots [\mathscr{L}_n] \in \mathbb{Z}$  instead of the more correct  $[\mathscr{L}_1] \cdots [\mathscr{L}_n] \cdot [X]$ . Ditto for intersection with Cartier divisors.

For the reader's convenience, we recall the notion of "numerical flatness for vector bundles", which generalises the notion of "numerical triviality" for line bundles.

**Definition 2.11** (Nefness and numerical flatness, [DPS94, Def. 1.17]). Let  $\varphi: X \to Y$  be a projective morphism of quasi-projective varieties. Given a locally free sheaf  $\mathscr{F}$  on X, consider the composed morphism

$$\mathbb{P}(\mathscr{F}) \xrightarrow{\delta} X \xrightarrow{g} Y.$$

The sheaf  $\mathscr{F}$  is called  $\varphi$ -nef if  $[\mathscr{O}_{\mathbb{P}(\mathscr{F})}(1)]$  is  $\delta$ -nef, that is  $[\mathscr{O}_{\mathbb{P}(\mathscr{F})}(1)] \cdot C \geq 0$ , for every irreducible curve  $C \subset \mathbb{P}(\mathscr{F})$  such that  $\delta(C)$  is a point. The sheaf  $\mathscr{F}$  is called  $\varphi$ -numerically flat if  $\mathscr{F}$  and its dual  $\mathscr{F}^*$  are both  $\varphi$ -nef. If Y is a point, we simply say that  $\mathscr{F}$  is numerically flat.

*Remark* 2.12 (Alternate formulations of numerical flatness). Setting as in Definition 2.11. The following conditions are equivalent.

- (2.12.1) The bundle  $\mathscr{F}$  and its dual  $\mathscr{F}^*$  are both  $\varphi$ -nef.
- (2.12.2) The bundle  $\mathscr{F}$  is  $\varphi$ -nef and the invertible sheaf  $(\det \mathscr{F})^*$  is  $\varphi$ -nef.
- (2.12.3) The bundle  $\mathscr{F}$  is  $\varphi$ -nef and the invertible sheaf det  $\mathscr{F}$  is  $\varphi$ -numerically trivial.

Chern classes of numerically flat bundles vanish. The proof is based on two deep facts: the Uhlenbeck-Yau theorem asserting the existence of Hermite-Einstein metrics on stable vector bundles, and the resulting Kobayashi-Lübke flatness criterion derived from Lübke's inequality on Chern classes of Hermite-Einstein vector bundles.

**Theorem 2.13** (Chern classes of numerically flat bundles, [DPS94, Cor. 1.19]). Let X be a smooth, projective variety and  $\mathscr{F}$  a numerically flat, locally free sheaf on X. Then, all Chern classes  $c_i(\mathscr{F}) \in H^{2i}(X, \mathbb{R})$  vanish.

2.6. **KLT spaces, the Basepoint-free Theorem and contractions.** We will mostly work with klt pairs  $(X, \Delta)$ , but the boundary divisor  $\Delta$  is usually irrelevant in our discussion. We will use the following shorthand notation throughout. We refer to [KM98] for a standard reference concerning klt pairs but recall the global convention that the boundary divisor is always assumed to be effective in this paper.

**Definition 2.14** (KLT spaces). A normal, quasi-projective variety is called a klt space if there exists an effective Weil  $\mathbb{Q}$ -divisor  $\Delta$  such that  $(X, \Delta)$  is klt.

*Note.* Recall that a pair  $(X, \Delta)$  as above is called klt<sup>1</sup> if the Weil  $\mathbb{Q}$ -divisor  $K_X + \Delta$  is  $\mathbb{Q}$ -Cartier,  $\lfloor \Delta \rfloor = 0$ , and if for one (equivalently: for every) resolution of singularities,  $\pi : \widetilde{X} \to X$ , there exists a  $\mathbb{Q}$ -linear equivalence of the form

$$K_{\widetilde{X}} + \pi_*^{-1} \Delta \sim_{\mathbb{Q}} \pi^* (K_X + \Delta) + \sum_i a_i \cdot E_i,$$

where the  $E_i \subset \widetilde{X}$  are  $\pi$ -exceptional and the numbers  $a_i \in \mathbb{Q}$  satisfy  $a_i > -1$  for all i.

<sup>&</sup>lt;sup>1</sup>This is short for "Kawamata log terminal".

The results of Section 4 rely in part on the Basepoint-free Theorem and its immediate Corollary 2.16, which proves Theorem 4.1 for line bundles. We recall the statement for the reader's convenience. Full details are found in the standard references, cf. including [KMM87, Thm. 3-1-1 and Rem. 3-1-2(i)], [KM98, Thm. 3.24] and [Laz04, Thm. 2.1.27].

**Theorem 2.15** (Basepoint-free Theorem). Let  $\varphi: X \to Y$  be a projective surjection of normal, quasi-projective varieties. Assume that there exists an effective Weil  $\mathbb{Q}$ -divisor  $\Delta$  on X such that  $(X, \Delta)$  is klt. If L is any  $\varphi$ -nef Cartier divisor on X and  $m \in \mathbb{N}^+$  any number such that  $m \cdot L - (K_X + \Delta)$  is  $\varphi$ -nef and  $\varphi$ -big, then there exists a unique factorisation via a normal variety Z,

$$(2.15.1) X \xrightarrow{\varphi} Z \xrightarrow{\beta} Y,$$

such that the following holds.

- (2.15.2) The morphisms  $\alpha$  and  $\beta$  are surjective. The morphism  $\alpha$  has connected fibres.
- (2.15.3) The Cartier divisor L is the pull-back of a  $\beta$ -ample Cartier divisor  $L_Z$  on Z.
- (2.15.4) If  $Y^{\circ} \subseteq Y$  is open with preimages  $X^{\circ}$  and  $Z^{\circ}$ , and if  $L|_{X^{\circ}}$  is  $\varphi|_{X^{\circ}}$ -ample, then  $\alpha|_{X^{\circ}}: X^{\circ} \to Z^{\circ}$  is isomorphic.

*Note.* Item (2.15.3) implies that the morphism  $\alpha$  contracts exactly those irreducible curves C with  $[L] \cdot C = 0$  and dim  $\varphi(C) = 0$ .

**Corollary 2.16** (Descent of invertible sheaves). Let  $\varphi: X \to Y$  be a birational, projective morphism of normal, quasi-projective varieties, and let L be any  $\varphi$ -numerically trivial Cartier divisor on X. If there exists a Weil  $\mathbb{Q}$ -divisor  $\Delta$  on X such that  $(X, \Delta)$  is klt and such that  $-(K_X + \Delta)$  is  $\varphi$ -nef, then L is linearly equivalent to the pull-back of a Cartier divisor on Y.

*Proof.* We claim that L is  $\varphi$ -nef and that  $D := L - (K_X + \Delta)$  is  $\varphi$ -nef and  $\varphi$ -big. Relative nefness of L and D holds by assumption. The condition that D is  $\varphi$ -big is void since  $\varphi$  is assumed to be birational. Theorem 2.15 thus gives a factorisation of  $\varphi$  via a normal variety Z as in (2.15.1), and  $\beta$ -ample Cartier divisor  $L_Z$  on Z such that  $L \sim \alpha^* L_Z$ .

As a next step, we claim that  $\beta$  is finite. If not, we would find a curve  $C \subseteq X$  which is mapped to a point by  $\varphi$ , but not by  $\alpha$ . The image curve  $\alpha(C)$  would them be contained in a  $\beta$ -fibre, and would thus have positive intersection with the  $\beta$ -ample divisor  $L_Z$ . In particular,  $\deg L|_C = \deg \alpha^*(L_Z)|_C > 0$ , contradicting the assumption that L is  $\varphi$ -numerically trivial

In summary, see that  $\beta$  is both birational and finite. Zariski's main theorem, [GW10, Cor. 12.88], applies to show that  $\beta$  is isomorphic. Corollary 2.16 follows.

We list some basic properties of the contraction morphism associated with an extremal face and refer to [KMM87, Def. 3-2-3] for terminology.

**Proposition 2.17** (Relative contractions of extremal faces). Assume we are given a sequence of projective surjections between normal, quasi-projective varieties,  $f: X \to Y$  and  $Y \to Z$ . Assume also that there exists an effective Weil  $\mathbb{Q}$ -divisor  $\Delta$  on X such that  $(X, \Delta)$  is klt.

- (2.17.1) If f has only connected fibres and if  $-(K_X + \Delta)$  is f-ample, then there exists a  $(K_X + \Delta)$ -negative extremal face F of  $\overline{NE}(X/Z)$  such that f is the contraction morphism of F.
- (2.17.2) If there exists a  $(K_X + \Delta)$ -negative extremal face F of  $\overline{NE}(X/Z)$  such that f is the contraction morphism of F, then relative Picard numbers are additive,

$$\rho(X/Z) = \rho(X/Y) + \rho(Y/Z).$$

The extremal face F is an extremal ray if and only if  $\rho(X/Y) = 1$ .

*Proof.* Item (2.17.1) is [KMM87, Lem. 3-2-5(1)]. The additivity of Picard numbers in Item (2.17.2) is [KMM87, Lem. 3-2-5(3)]. It remains to consider the relation between Picard numbers and the dimension of F in Item (2.17.2).

If the dimension of F is one, it follows from the definition that  $\rho(X/Y) = 1$  once we know that there *is* a curve in X that is mapped to a point by f. This is shown in [KMM87, Lem. 3-2-4]. As for the converse direction, if  $\rho(X/Y) = 1$ , the definition implies that there must be curves that are contracted. The face F cannot be empty, and is necessarily of dimension one.

*Warning* 2.18. Picard numbers are generally not additive for compositions of arbitrary morphisms. See [KM98, Sect. 2.2] for a discussion and [KMM87, Rem. 3-2-6] for an explicit example.

2.7. **Sheaves with operators and Higgs sheaves.** Higgs sheaves and sheaves with operators on singular varieties were defined and discussed in detail in [GKPT15, Sects. 4 and 5]. We briefly recall the main definitions here and then discuss pullback of Higgs sheaves.

## 2.7.1. Fundamental definitions. -

**Definition 2.19** (Sheaf with an operator, invariant subsheaves, [GKPT15, Def. 4.1]). Let X be a normal, quasi-projective variety and  $\mathscr{W}$  be a coherent sheaf of  $\mathscr{O}_X$ -modules. A sheaf with a  $\mathscr{W}$ -valued operator is a pair  $(\mathscr{E},\theta)$  where  $\mathscr{E}$  is a coherent sheaf and  $\theta:\mathscr{E}\to\mathscr{E}\otimes\mathscr{W}$  is an  $\mathscr{O}_X$ -linear sheaf morphism.

**Definition 2.20** (Invariant subsheaf, [GKPT15, Def. 4.8]). Setting as in Definition 2.19. A coherent subsheaf  $\mathscr{F} \subseteq \mathscr{E}$  is called  $\theta$ -invariant if  $\theta(\mathscr{F})$  is contained in the image of the natural map  $\mathscr{F} \otimes \mathscr{W} \to \mathscr{E} \otimes \mathscr{W}$ . Call  $\mathscr{F}$  generically invariant if the restriction  $\mathscr{F}|_U$  is invariant with respect to  $\theta|_U$ , where  $U \subseteq X$  is the maximal, dense, open subset where  $\mathscr{W}$  is locally free.

**Definition 2.21** (Stability of sheaves with operator, [GKPT15, Def. 4.13]). Let X be a normal, projective variety and H be any nef,  $\mathbb{Q}$ -Cartier  $\mathbb{Q}$ -divisor on X. Let  $(\mathcal{E}, \theta)$  be a sheaf with an operator, as in Definition 2.19, were  $\mathcal{E}$  is torsion free. We say that  $(\mathcal{E}, \theta)$  is semistable with respect to H if the inequality  $\mu_H(\mathcal{F}) \leq \mu_H(\mathcal{E})$  holds for all generically  $\theta$ -invariant subsheaves  $\mathcal{F} \subseteq \mathcal{E}$  with  $0 < \operatorname{rank} \mathcal{F} < \operatorname{rank} \mathcal{E}$ . The pair  $(\mathcal{E}, \theta)$  is called stable with respect to H if strict inequality holds. Direct sums of stable sheaves with operator are called polystable.

On a singular variety, some attention has to be paid concerning the definition of "Higgs sheaf" at singular points. Again, we recall the relevant definitions here.

**Definition 2.22** (Higgs sheaf, stability, morphisms, [GKPT15, Defs. 5.1 and 5.2, Sect. 5.6]). Let X be a normal variety. A Higgs sheaf is a pair  $(\mathscr{E}, \theta)$  of a coherent sheaf  $\mathscr{E}$  of  $\mathscr{O}_X$ -modules, together with an  $\Omega_X^{[1]}$ -valued operator  $\theta: \mathscr{E} \to \mathscr{E} \otimes \Omega_X^{[1]}$ , called Higgs field, such that the composed morphism

$$\mathscr{E} \xrightarrow{\quad \theta \quad } \mathscr{E} \otimes \Omega_X^{[1]} \xrightarrow{\quad \theta \otimes \mathrm{Id} \quad } \mathscr{E} \otimes \Omega_X^{[1]} \otimes \Omega_X^{[1]} \xrightarrow{\quad \mathrm{Id} \otimes [\wedge] \quad } \mathscr{E} \otimes \Omega_X^{[2]}$$

vanishes. A Higgs sheaf is called stable if it is stable as a sheaf with an  $\Omega_X^{[1]}$ -valued operator, ditto for semistable and polystable. A morphism of Higgs sheaves, written  $f:(\mathscr{E}_1,\theta_1)\to (\mathscr{E}_2,\theta_2)$ , is a morphism  $f:\mathscr{E}_1\to\mathscr{E}_2$  of coherent sheaves that commutes with the Higgs fields,  $(f\otimes \operatorname{Id}_{\Omega_X^{[1]}})\circ\theta_1=\theta_2\circ f$ .

2.7.2. *Pull-back.* If  $f: Y \to X$  is a morphism of normal varieties, and if  $(\mathscr{E}, \theta)$  is a Higgs sheaf on X, there is generally no way to equip the pull-back sheaf  $f^*\mathscr{E}$  with a Higgs field, even if  $\mathscr{E}$  is locally free. It is a non-trivial fact that pull-backs do exist for klt spaces. We refer to [GKPT15, Sects. 5.3 and 5.4] for details. In brief, if X is klt, recall from [Keb13,

Thms. 1.3 and 5.2] that there exists a natural pull-back functor for reflexive differentials on klt pairs that is compatible with the usual pull-back of Kähler differentials and gives rise to a sheaf morphism

$$d_{\text{refl}}f: f^*\Omega_X^{[1]} \to \Omega_Y^{[1]}.$$

One can then define a Higgs field on  $f^*\mathscr{E}$  as the composition of the following morphisms,

$$(2.22.1) \hspace{1cm} f^{*}\mathscr{E} \xrightarrow{f^{*}\theta} f^{*}\Big(\mathscr{E} \otimes \Omega_{X}^{[1]}\Big) = f^{*}\mathscr{E} \otimes f^{*}\Omega_{X}^{[1]} \xrightarrow{\operatorname{Id}_{f^{*}\mathscr{E}} \otimes d_{\operatorname{refl}}f} f^{*}\mathscr{E} \otimes \Omega_{Y}^{[1]}.$$

2.8. **Stability for sheaves on the smooth locus.** In the situation discussed in the present paper, it makes sense to generalise the stability notions of Section 2.7 to the case where the (Higgs-) sheaves are defined on the smooth locus of a normal variety only.

**Definition 2.23** (Slope and stability for sheaves on the smooth locus). Let X be a normal, projective variety and let  $\mathscr{E}^{\circ}$  be a torsion free, coherent sheaf on  $X_{\text{reg}}$  of positive rank. If  $H \in \text{Div}(X)$  is nef, define the slope of  $\mathscr{E}^{\circ}$  with respect to H as

$$\mu_H(\mathscr{E}^{\circ}) := \frac{c_1(\iota_*\mathscr{E}^{\circ}) \cdot [H]^{\dim X - 1}}{\operatorname{rank} \mathscr{E}^{\circ}},$$

where  $\iota: X_{\text{reg}} \to X$  is the inclusion.

Remark 2.24 (Algebraicity assumption). We underline that  $\mathscr{E}^{\circ}$  is assumed to be algebraic in Definition 2.23. For coherent *analytic* sheaves on  $X_{\text{reg}}^{an}$ , the push-forward  $\iota_*\mathscr{E}^{\circ}$  need not be coherent in general.

**Definition 2.25** (Slope and stability for sheaves on the smooth locus). Setting as in Definition 2.23. If  $\mathscr{W}^{\circ}$  is coherent on  $X_{\text{reg}}$  and if  $\theta^{\circ}: \mathscr{E}^{\circ} \to \mathscr{E}^{\circ} \otimes \mathscr{W}^{\circ}$  is a  $\mathscr{W}^{\circ}$ -valued operator, say that  $(\mathscr{E}^{\circ}, \theta^{\circ})$  is stable with respect to H, if the inequality  $\mu_H(\mathscr{F}^{\circ}) < \mu_H(\mathscr{E}^{\circ})$  holds for all generically  $\theta^{\circ}$ -invariant subsheaves  $\mathscr{F}^{\circ} \subseteq \mathscr{E}^{\circ}$  with  $0 < \text{rank } \mathscr{F}^{\circ} < \text{rank } \mathscr{E}^{\circ}$ . Analogously, define notions of semistable and polystable for sheaves with operators on  $X_{\text{reg}}$ , ditto for Higgs sheaves.

The following two lemmas summarise properties of the generalised stability notions that will be used later. Proofs are elementary and therefore omitted.

**Lemma 2.26** (Restriction of stable sheaves to  $X_{\text{reg}}$ ). Let X be a normal, projective variety, and let  $(\mathcal{E}, \theta)$  be a torsion free sheaf with a  $\mathcal{W}$ -valued operator. If  $H \in \text{Div}(X)$  is nef, then  $(\mathcal{E}, \theta)$  is semistable (resp. stable) with respect to H as a sheaf with a  $\mathcal{W}$ -valued operator if and only if  $(\mathcal{E}, \theta)|_{X_{\text{reg}}}$  is semistable (resp. stable) with respect to H as a sheaf with a  $\mathcal{W}$   $|_{X_{\text{reg}}}$ -valued operator.

**Lemma 2.27** (Extensions of operators from subsheaves). Let X be a normal, projective variety and let  $(\mathcal{E}^{\circ}, \theta^{\circ})$  be a torsion free sheaf on  $X_{reg}$  with a  $\mathcal{W}^{\circ}$ -valued operator. Let  $\iota^{\circ}: \mathcal{W}^{\circ} \hookrightarrow \mathcal{V}^{\circ}$  is an inclusion of coherent sheaves on  $X_{reg}$  and consider the natural  $\mathcal{V}^{\circ}$ -valued operator  $\tau^{\circ}$  that is defined as the composition

$$\mathcal{E}^{\circ} \xrightarrow{\theta^{\circ}} \mathcal{E}^{\circ} \otimes \mathcal{W}^{\circ} \xrightarrow{\operatorname{Id} \otimes \iota^{\circ}} \mathcal{E}^{\circ} \otimes \mathcal{V}^{\circ}.$$

If  $H \in Div(X)$  is nef, then,  $(\mathcal{E}^{\circ}, \theta^{\circ})$  semistable (resp. stable) with respect to H as a sheaf with a  $\mathcal{W}^{\circ}$ -valued operator if and only if  $(\mathcal{E}^{\circ}, \tau^{\circ})$  semistable (resp. stable) with respect to H as a sheaf with a  $\mathcal{V}^{\circ}$ -valued operator.

- 3. The nonabelian Hodge correspondence for locally free Higgs sheaves
- 3.1. **Main result.** Generalising work of Simpson, [Sim92, Cor. 3.10], in this section we formulate a nonabelian Hodge correspondence for locally free Higgs sheaves on klt spaces, relating such locally free Higgs sheaves to representations of the fundamental group. The formulation uses the fact that every algebraic variety admits a distinguished, *canonical* resolution of singularities, see [BM97, Rem. 1.16ff and Sect. 13]. The following additional notation will be used.

*Notation* 3.1 (Categories in the nonabelian Hodge correspondence). Given a normal, projective variety X, consider the following categories.

 $\operatorname{Higgs}_X$  Locally free Higgs sheaves  $(\mathscr{E},\theta)$  on X having the property that there exists an ample divisor  $H\in\operatorname{Div}(X)$  such that  $(\mathscr{E},\theta)$  is semistable with respect to H and additionally satisfies  $ch_1(\mathscr{E})\cdot [H]^{n-1}=ch_2(\mathscr{E})\cdot [H]^{n-2}=0$ .

 $LSys_x$  Local systems on X.

Notation 3.2 (Higgs sheaves). There is no uniform definition of Higgs sheaves on singular spaces in the literature. Throughout the present paper, we use the definition given in Section 2.7.1. (Semi)stability of Higgs sheaves is defined and discussed in [GKPT15, Sect. 5.6]. We refer to [Del70, Sect. I.1] for a discussion of the basic properties of local systems.

Notation 3.3 (Nonabelian Hodge correspondence for manifolds). If X is smooth, Simpson's nonabelian Hodge correspondence gives an equivalence between the categories  $\operatorname{Higgs}_X$  and  $\operatorname{LSys}_X$ . We denote the relevant functors by  $\eta_X:\operatorname{LSys}_X\to\operatorname{Higgs}_X$  and  $\mu_X:\operatorname{Higgs}_X\to\operatorname{LSys}_X$ .

The following is now the main result. As we will see in Section 3.2 below, the properties spelled out in (3.4.2) and (3.4.3) immediately imply that the nonabelian Hodge correspondence presented here is in fact fully functorial in morphisms of klt spaces.

**Theorem 3.4** (Nonabelian Hodge correspondence for klt spaces). For every projective klt space X, there exists an equivalence of categories, given by functors  $\eta_X : \mathsf{LSys}_X \to \mathsf{Higgs}_X$  and  $\mu_X : \mathsf{Higgs}_X \to \mathsf{LSys}_X$  such that the following additional properties hold.

- (3.4.1) If X is smooth, then  $\eta_X$  and  $\mu_X$  equal the functors from Simpson's nonabelian Hodge correspondence.
- (3.4.2) If  $(\mathscr{E}, \theta) \in \operatorname{Higgs}_X$  and  $\pi : \widetilde{X} \to X$  is the canonical resolution of singularities, then  $\pi^*(\mathscr{E}, \theta) \in \operatorname{Higgs}_{\widetilde{X}}$ , and there exists a canonical isomorphism of local systems,

$$M_{\pi,(\mathscr{E},\theta)}: \pi^*\mu_X(\mathscr{E},\theta) \to \mu_{\widetilde{X}}(\pi^*(\mathscr{E},\theta)).$$

(3.4.3) If  $E \in LSys_X$  is any local system and  $\pi : \widetilde{X} \to X$  is the canonical resolution of singularities, then there exists a canonical isomorphism of Higgs sheaves,

$$N_{\pi,E}:\pi^*\eta_X(\mathsf{E})\to\eta_{\widetilde{X}}(\pi^*\mathsf{E}).$$

*Note* (Pull-back of Higgs sheaves). Item (3.4.2) discusses the pull-back of the Higgs sheaf  $\eta_X(E)$  from X to the resolution of singularities,  $\widetilde{X}$ , as discussed in Section 2.7.2 above.

Theorem 3.4 is shown in Section 8.1. Section 7 prepares for the proof.

3.2. **Functoriality.** Items (3.4.2) and (3.4.3) of Theorem 3.4 allow to describe the functors  $\eta_X$  and  $\mu_X$  on any given klt space X in terms of the classical nonabelian Hodge correspondence that exists on the canonical resolution of singularities. In fact, a much more general functoriality holds true.

**Theorem 3.5** (Functoriality in morphisms). The correspondence of Theorem 3.4 is functorial in morphisms. More precisely, for every morphism  $f: Y \to X$  of projective klt spaces, every  $E \in \mathsf{LSys}_X$  and every  $(\mathscr{E}, \theta) \in \mathsf{Higgs}_X$ , there exist canonical isomorphisms

$$M_{f,(\mathscr{E},\theta)}: f^*\mu_X(\mathscr{E},\theta) \to \mu_Y\big(f^*(\mathscr{E},\theta)\big) \quad and \quad N_{f,\mathsf{E}}: f^*\eta_X(\mathsf{E}) \to \eta_Y\big(f^*\mathsf{E}\big).$$

The collection of these isomorphisms satisfies the following properties.

**Functoriality:** Given morphisms  $g: Z \to Y$  and  $f: Y \to X$  between projective klt spaces and  $E \in \mathsf{LSys}_X$ , then the following diagram commutes,

$$g^*f^*\eta_X(\mathsf{E}) \xrightarrow{g^*N_{f,\mathsf{E}}} g^*\eta_Y\big(f^*\mathsf{E}\big) \xrightarrow{N_{g,f^*\mathsf{E}}} \eta_Z\big(g^*f^*\mathsf{E}\big).$$

Ditto for the functor  $\mu_{\bullet}$  and the isomorphisms  $M_{\bullet, \bullet}$ .

**Behaviour under canonical resolution:** For  $\pi: X \to X$  the canonical resolution of a projective klt space, the morphisms  $M_{\pi, \bullet}$  and  $N_{\pi, \bullet}$  equal the isomorphisms given in Items (3.4.2) and (3.4.3) of Theorem 3.4.

**Compatibility:** If  $f: Y \to X$  is a morphism between smooth projective varieties, then  $M_{f,\bullet}$  and  $N_{f,\bullet}$  are the standard isomorphisms given by functoriality of Simpson's nonabelian Hodge correspondence.

Theorem 3.5 is shown in Section 8.2 below.

*Remark* 3.6 (Uniqueness). One verifies in the blink of an eye that functoriality, behaviour under canonical resolution and compatibility determine the isomorphisms  $M_{\bullet,\bullet}$  and  $N_{\bullet,\bullet}$  uniquely.

Remark 3.7 (General resolutions). Theorem 3.5 implies in particular that the statement of Theorem 3.4 holds for any resolution of singularities, not just the canonical resolution. Taken together with [Tak03], for any klt space this establishes an equivalence of categories of Higgs $_{\widetilde{X}}$ , with Higgs $_{\widetilde{X}}$ , where  $\widetilde{X}$  is any resolution of singularities of X.

As a further consequence of functoriality, we observe that the nonabelian Hodge correspondence respects group actions and relates *G*-linearised local systems to Higgs *G*-sheaves in the sense of [GKPT15, Def. 5.1].

**Corollary 3.8** (*G*-linearised local systems and Higgs *G*-sheaves). Let *X* be a projective klt space, and let *G* be a group acting on *X* via a group morphism  $G \to \operatorname{Aut}(X)$ .

(3.8.1) If  $(\mathcal{E}, \theta) \in \operatorname{Higgs}_X$  carries the structure of a Higgs G-sheaf, given by isomorphisms  $\varphi_g : g^*(\mathcal{E}, \theta) \to (\mathcal{E}, \theta)$ , then the following composed maps endow the local system  $\mu_X(\mathcal{E}, \theta)$  with a G-linearisation,

$$g^*\mu_X(\mathcal{E},\theta) \xrightarrow{\quad M_{g,(\mathcal{E},\theta)} \quad} \mu_X\big(g^*(\mathcal{E},\theta)\big) \xrightarrow{\quad \mu_X(\varphi_g) \quad} \mu_X(\mathcal{E},\theta).$$

(3.8.2) If  $E \in LSys_X$  carries a G-linearisation given by isomorphisms  $\varphi_g : g^*E \to E$ , then the following composed maps endow  $\eta_X(E)$  with the structure of a Higgs G-sheaf,

$$g^* \eta_X(\mathsf{E}) \xrightarrow{N_{g,\mathsf{E}}} \eta_X(g^*\mathsf{E}) \xrightarrow{\eta_X(\varphi_g)} \eta_X(\mathsf{E}).$$

3.3. **Independence of polarisation.** As in Simpson's original setup, a Higgs bundle on a klt space X is in Higgs $_X$  if and only if it satisfies the conditions of Notation 3.1 with respect to any ample class. The following proposition makes this assertion precise.

**Theorem 3.9** (Independence of polarisation). Let X be a projective klt space of dimension n. Given any locally free Higgs sheaf  $(\mathcal{E}, \theta)$  on X, the following statements are equivalent.

- (3.9.1) There exists an ample divisor  $H \in \text{Div}(X)$  such that  $ch_1(\mathscr{E}) \cdot [H]^{n-1} = ch_2(\mathscr{E}) \cdot [H]^{n-2} = 0$  and such that  $(\mathscr{E}, \theta)$  is semistable with respect to H.
- (3.9.2) For all ample divisors  $H \in \text{Div}(X)$ , we have  $ch_1(\mathscr{E}) \cdot [H]^{n-1} = ch_2(\mathscr{E}) \cdot [H]^{n-2} = 0$  and  $(\mathscr{E}, \theta)$  is semistable with respect to H.
- (3.9.3) All Chern classes  $c_i(\mathcal{E}) \in H^{2i}(X, \mathbb{Q})$  vanish and  $(\mathcal{E}, \theta)$  is semistable with respect to any ample divisor on X.

- (3.9.4) There exists a resolution of singularities,  $\pi: \widetilde{X} \to X$ , and an ample divisor  $\widetilde{H} \in \operatorname{Div}(\widetilde{X})$  such that  $\operatorname{ch}_1(\pi^*\mathscr{E}) \cdot [\widetilde{H}]^{n-1} = \operatorname{ch}_2(\pi^*\mathscr{E}) \cdot [\widetilde{H}]^{n-2} = 0$  and such that  $\pi^*(\mathscr{E}, \theta)$  is semistable with respect to  $\widetilde{H}$ .
- (3.9.5) For any resolution of singularities,  $\pi: \widetilde{X} \to X$ , and any ample divisor  $\widetilde{H} \in \text{Div}(\widetilde{X})$ , we have intersection numbers  $ch_1(\pi^*\mathscr{E}) \cdot [\widetilde{H}]^{n-1} = ch_2(\pi^*\mathscr{E}) \cdot [\widetilde{H}]^{n-2} = 0$  and  $\pi^*(\mathscr{E}, \theta)$  is semistable with respect to  $\widetilde{H}$ .

The analogous equivalences hold when "semistable" is replaced by "stable" or "polystable".

Theorem 3.9 is shown in Section 8.3 below.

3.4. Harmonic bundles, differential graded categories. Simpson constructs his nonabelian Hodge correspondence first in the case of polystable Higgs bundles and semisimple local systems. In this setup, the correspondence is a consequence of existence theorems for pluri-harmonic metrics on the underlying bundles. Given their central role in the theory, we remark that the nonabelian Hodge correspondence for klt spaces, Theorem 3.4 also has a description in terms of harmonic structures, although in our case the harmonic metric exists on the smooth locus of the underlying space only. The proof of the following proposition is simple and therefore omitted: Item (3.4.3) of Theorem 3.4 allows to relate the correspondence on X to that on a resolution.

**Proposition 3.10** (Hodge correspondence for klt spaces via harmonic bundles). Let X be a projective, klt space and  $E \in LSys_X$  a semisimple local system on X, with underlying  $C^{\infty}$ -bundle E. We claim that there exists a tame and purely imaginary harmonic bundle  $(E^{\circ}, \bar{\partial}_{E^{\circ}}, \theta^{\circ}, h^{\circ})$  on  $X_{reg}$  with the following two properties.

- (3.10.1) The induced flat bundle  $(E^{\circ}, \nabla_{h^{\circ}})$  corresponds to the local system  $E|_{X_{ree}}$ .
- (3.10.2) Writing  $\mathscr{E}^{\circ}$  for the sheaf of holomorphic sections in  $(E^{\circ}, \bar{\partial}_{E^{\circ}})$ , the functor  $\eta_X$  of the nonabelian Hodge correspondence satisfies  $(\mathscr{E}^{\circ}, \theta^{\circ}) \cong \eta_X(\mathsf{E})|_{X_{reg}}$ .

Remark 3.11. We refer to [Sab13, Sect. 1] for a discussion of harmonic bundles, and to [Sim90, p. 723] and [Moc07b, Sect. 22.1] for the notions of "tame" and "purely imaginary".

As a second point, we note that Theorem 3.4 includes equivalences of differential graded categories (=DGCs) that appear in Simpson's nonabelian Hodge theory. For detailed discussion of DGCs and their relation to this theory we refer the reader to [Sim92, Sect. 3] and the references therein. Given a resolution  $\pi:\widetilde{X}\to X$ , Theorem 3.4 implies that there is an equivalence of DGCs between the category of extensions of stable Higgs bundles with vanishing Chern classes on X and the category of flat connections on  $\widetilde{X}$ . The same conclusions as in the smooth projective setting then follow.

## 4. Descent of vector bundles from KLT spaces

The proof of Theorem 5.1, our main result concerning descent of vector bundles to klt spaces, relies on the following auxiliary statement, which we prove in this section.

**Theorem 4.1** (Descent of vector bundles from klt spaces). Let  $\varphi: X \to Y$  be a projective, birational morphism of normal, quasi-projective varieties. Assume that there exists a Weil  $\mathbb{Q}$ -divisor  $\Delta_X$  on X such that the pair  $(X, \Delta_X)$  is klt and  $-(K_X + \Delta_X)$  is  $\varphi$ -nef. If  $\mathscr{F}_X$  is any locally free,  $\varphi$ -numerically flat sheaf on X, then there exists a locally free sheaf  $\mathscr{F}_Y$  on Y such that  $\mathscr{F}_X \cong \varphi^* \mathscr{F}_Y$ .

4.1. **Proof of Theorem 4.1: Setup and notation.** We maintain notation and assumptions of Theorem 4.1 throughout the present Section 4. To avoid trivial cases, we may assume throughout the proof that the rank of  $\mathscr{F}_X$  is positive. Set  $r := (\operatorname{rank} \mathscr{F}_X) - 1$ . Using Grothendieck's terminology, we consider the associated  $\mathbb{P}^r$ -bundle  $\mathbb{P}_X := \mathbb{P}_X(\mathscr{F}_X)$ .

The following diagram summarises the situation

Since  $\rho_X$  is a locally trivial  $\mathbb{P}^r$ -bundle, the variety  $\mathbb{P}_X$  is normal, and the pair  $(\mathbb{P}_X, \Delta_{\mathbb{P}_X})$  is klt, where  $\Delta_{\mathbb{P}_X} := \rho_X^* \Delta_X$ . Let  $Y^\circ \subseteq Y$  be the maximal open set over which  $\varphi$  is isomorphic, and write  $X^\circ := \varphi^{-1}(X^\circ)$ . As Y is normal, the subset  $Y^\circ$  is big.

We will also consider the invertible sheaves  $\mathscr{L}_X := \mathscr{O}_{\mathbb{P}_X(\mathscr{F}_X)}(1)$  and  $\mathscr{M}_X := \det \mathscr{F}_X$ . The assumption that  $\mathscr{F}_X$  is  $\varphi$ -numerically flat has immediate consequences for  $\mathscr{L}_X$  and  $\mathscr{M}_X$ , which we state for later reference.

Observation 4.2. -

- (4.2.1) The sheaf  $\mathcal{L}_X$  is  $\delta$ -nef.
- (4.2.2) The sheaf  $\mathcal{M}_X$  is  $\varphi$ -numerically trivial. In particular, Corollary 2.16 implies the existence of an invertible sheaf  $\mathcal{M}_Y \in \operatorname{Pic}(Y)$  such that  $\mathcal{M}_X \cong \varphi^* \mathcal{M}_Y$ .

For convenience of notation, choose Cartier divisors  $L_X$  and  $M_Y$  representing the bundles  $\mathscr{L}_X$  and  $\mathscr{M}_Y$ . The Cartier divisor  $M_X := \varphi^* M_Y$  will then represent  $\mathscr{M}_X$ .

4.2. **Proof of Theorem 4.1: Factorisation of**  $\delta$ . We aim to construct a locally free sheaf  $\mathscr{F}_Y$  on Y. Rather than doing so directly, we will first construct a factorisation of  $\delta$  via a morphism  $\rho_Y : \mathbb{P}_Y \to Y$  that agrees with  $\mathbb{P}_X$  over the big open set where  $Y^{\circ}$ . Later, we will show that  $\rho_Y$  is equidimensional, and has in fact the structure of a linear  $\mathbb{P}^r$ -bundle. The sheaf  $\mathscr{F}_Y$  will then be constructed as the push-forward of the relative hyperplane bundle on  $\mathbb{P}_Y$ .

Claim 4.3 (Construction of  $\mathbb{P}_Y$ ). There exists a commutative diagram of surjective projective morphisms with connected fibres extending Diagram (4.1.1) as follows,

$$\begin{array}{c|c}
\mathbb{P}_{X} & \xrightarrow{\Phi} \mathbb{P}_{Y} \\
\downarrow^{\rho_{X}} & & \downarrow^{\rho_{Y}} \\
X & \xrightarrow{\varphi, \text{ birational}} Y,
\end{array}$$

as well as a  $\rho_Y$ -ample sheaf  $\mathcal{L}_Y \in \operatorname{Pic}(\mathbb{P}_Y)$  such that  $\mathcal{L}_X \cong \Phi^* \mathcal{L}_Y$ . The variety  $\mathbb{P}_Y$  and the morphisms of Diagram (4.3.1) are unique up to isomorphism, and the following holds in addition.

- (4.3.2) The restricted morphism  $\Phi|_{\rho_X^{-1}(X^\circ)}: \rho_X^{-1}(X^\circ) \to \rho_Y^{-1}(Y^\circ)$  is isomorphic. In particular,  $\Phi$  is birational.
- (4.3.3) We have an isomorphism of sheaves  $\mathcal{M}_Y \cong \det((\rho_Y)_* \mathcal{L}_Y)$ .

*Proof of Claim 4.3.* The factorisation of  $\delta$  via an intermediate variety  $\mathbb{P}_Y$  will be constructed using Theorem 2.15 ("Basepoint-free Theorem"). To apply the theorem, it suffices to show that  $L_X$  is  $\delta$ -nef and that

$$D := L_X - \left( K_{\mathbb{P}_X} + \Delta_{\mathbb{P}_X} \right)$$

is  $\delta$ -nef and  $\delta$ -big. Relative nefness of  $L_X$  is clear from Observation (4.2.1). To analyse D, we use the standard formula for the canonical bundle of a projectivised vector bundle to obtain a  $\mathbb{Q}$ -linear equivalence of  $\mathbb{Q}$ -divisors,

(4.3.4) 
$$D \sim_{\mathbb{Q}} (r+2) \cdot L_X - \rho_X^* (K_X + \Delta_X + M_X).$$

The divisor  $L_X$  is  $\rho_X$ -ample and therefore ample on the general fibre of  $\delta$ . Since  $\varphi$  is birational, Equation (4.3.4) implies that D is  $\delta$ -big. Relative nefness of D also follows from

Equation (4.3.4), using Observations (4.2.1) and (4.2.2), as well as the assumption that  $-(K_X + \Delta_X)$  is  $\varphi$ -nef. Theorem 2.15 thus applies and yields a unique factorisation as in Diagram (4.3.1), as well as a  $\rho_Y$ -ample sheaf  $\mathcal{L}_Y$  such that  $\mathcal{L}_X \cong \Phi^* \mathcal{L}_Y$ .

Item (4.3.2) follows from Item (2.15.4) of Theorem 2.15, using the fact that D is  $\rho_X$ -ample over  $X^\circ$ , and therefore  $\delta$ -ample over  $Y^\circ$ . It remains to show that  $\mathcal{M}_Y \cong \det((\rho_Y)_* \mathcal{L}_Y)$ . By Item (4.3.2), an isomorphism exists at least over the big open set  $Y^\circ$ . But since both sides of the equation are reflexive, the isomorphism exists globally. This ends the proof of Claim 4.3.

4.3. **Proof of Theorem 4.1: Equidimensionality.** As indicated above, we will now show that  $\rho_Y$  is equidimensional. This is the point where the assumption that  $\mathscr{F}_X$  is  $\varphi$ -numerically flat is used in a crucial way.

Claim 4.4 (Equidimensionality of  $\rho_Y$ ). The morphism  $\rho_Y$  is equidimensional, of relative dimension r.

*Proof of Claim 4.4.* We argue by contradiction and assume that there exists a point  $y \in Y$  whose fibre  $\rho_Y^{-1}(y)$  contains a subvariety Z of dimension r+1. Recalling that  $\mathcal{L}_Y$  is  $\rho_Y$ -ample, we have a positive intersection number

$$(4.4.1) [\mathscr{L}_Y]^{r+1} \cdot [Z] > 0.$$

We will see that this is absurd. To this end, let  $F \subseteq \varphi^{-1}(y)$  be any irreducible component. Choose a desingularisation  $\pi: \widetilde{F} \to F$  and extend Diagram (4.3.1) to the left by taking fibre products as follows,

$$\mathbb{P}_{\widetilde{F}} \xrightarrow{\Pi} \mathbb{P}_{F} \xrightarrow{\Phi} \mathbb{P}_{X} \xrightarrow{\Phi} \mathbb{P}_{Y} \qquad \mathbb{P}_{\widetilde{F}} \xrightarrow{\Pi} \mathbb{P}_{F} \xrightarrow{\Phi|_{\mathbb{P}_{F}}} \rho_{Y}^{-1}(y) \xrightarrow{\Phi} \mathbb{P}_{Y} \\
\rho_{\widetilde{F}} \downarrow \qquad \rho_{F} \downarrow \qquad \rho_{X} \downarrow \qquad \rho_{Y} \downarrow \qquad \downarrow \rho_{\widetilde{F}} \qquad \downarrow \rho_{F} \qquad \downarrow \rho_{Y} \\
\widetilde{F} \xrightarrow{\pi} F \xrightarrow{\pi} F \xrightarrow{\varphi|_{F}} \{y\} \xrightarrow{\Phi} Y$$

Setting  $\mathscr{F}_{\widetilde{F}} := \pi^* \mathscr{F}_X$  and  $\mathscr{L}_{\widetilde{F}} := \Pi^* \mathscr{L}_X$ , we obtain identifications

$$\mathbb{P}_{\widetilde{F}} \cong \mathbb{P}_{\widetilde{F}}\big(\mathscr{F}_{\widetilde{F}}\big) \quad \text{and} \quad \mathscr{L}_{\widetilde{F}} \cong \mathscr{O}_{\mathbb{P}_{\widetilde{F}}(\mathscr{F}_{\widetilde{F}})}(1).$$

If  $Z_{\widetilde{F}} \subseteq \mathbb{P}_{\widetilde{F}}$  is any r+1-dimensional subvariety that dominates Z, Inequality (4.4.1) immediately implies that

$$(4.4.2) \qquad [\mathscr{L}_{\widetilde{F}}]^{r+1} \cdot [Z_{\widetilde{F}}] = [\Pi^* \Phi^* \mathscr{L}_Y]^{r+1} \cdot [Z_{\widetilde{F}}] > 0.$$

On that other hand, as the pullback of a numerically flat bundle,  $\mathscr{F}_{\widetilde{F}}$  is numerically flat. Theorem 2.13 hence implies that all Chern classes  $c_i(\mathscr{F}_{\widetilde{F}})$  of  $\mathscr{F}_{\widetilde{F}}$  vanish. A standard Chern class computation on projectivised vector bundles, [Ful98, Rem. 3.2.4 on p. 55], thus gives

$$(4.4.3) c_1(\mathscr{L}_{\widetilde{F}})^{r+1} = -\sum_{i=1}^r \underbrace{c_i(\rho_{\widetilde{F}}^*\mathscr{F}_{\widetilde{F}})} \cdot c_1(\mathscr{L}_{\widetilde{F}})^{r+1-i} = 0.$$

Items (4.4.2) and (4.4.3) are obviously in contradiction. The assumption that  $\rho_Y^{-1}(y)$  contains an (r+1)-dimensional subvariety is thus absurd. In summary, we obtain that  $\rho_Y$  is equidimensional, thus finishing the proof of Claim 4.4.

Building on work of Kollár and Höring-Novelli, it has been shown by Araujo and Druel [AD14, Prop. 4.10] that equidimensionality and the existence of the relatively ample sheaf  $\mathcal{L}_Y$  whose restriction to general  $\rho_Y$ -fibres is the hyperplane bundle implies that  $\rho_Y$  has the structure of a linear bundle. We briefly recall the argument.

Claim 4.5 (Linear bundle structure of  $\rho_Y$ ). The sheaf  $\mathscr{F}_Y := (\rho_Y)_* \mathscr{L}_Y$  is locally free on Y. The morphism  $\rho_Y : \mathbb{P}_Y \to Y$  can be identified as the projection  $\mathbb{P}_Y(\mathscr{F}_Y) \to Y$ . The invertible sheaf  $\mathscr{L}_Y$  becomes  $\mathscr{O}_{\mathbb{P}_Y(\mathscr{F}_Y)}(1)$  under this identification.

*Proof.* By (4.3.2), we know that the general fibre  $\mathbb{P}_{Y,y}$  of  $\rho_Y$  is isomorphic to  $\mathbb{P}^r$ , with  $\mathcal{L}_Y|_{\mathbb{P}_{Y,y}} \cong \mathcal{O}_{\mathbb{P}^r}(1)$ . Since  $\mathcal{L}_Y$  is  $\rho_Y$ -ample, [HN13, Prop. 3.1] applies to guarantee that in fact all fibres of  $\rho_Y$  are irreducible and generically reduced, and that the normalisation of any fibre is isomorphic to  $\mathbb{P}^r$ . In particular, the Hilbert polynomial of the normalisation of the fibres is constant, and [Kol11, Thm. 12] applies to show that the variety  $\mathbb{P}_Y$  admits a simultaneous normalisation, which is a finite, birational morphism  $\eta: \mathbb{P}_Y \to \mathbb{P}_Y$  with the property that the morphism  $\rho_Y \circ \eta$  is flat and that all its fibres are normal. But since  $\mathbb{P}_Y$  already is normal, Zariski's main theorem, [Har77, V Thm. 5.2], applies to show that  $\eta$  is isomorphic, and all fibres of  $\rho_Y$  are therefore smooth, and isomorphic to  $\mathbb{P}^r$ . Grauert's theorem, [Har77, III Cor. 12.9] thus shows that  $\mathscr{F}_Y$  is locally free, and that the natural morphism  $(\rho_Y)^*\mathscr{F}_Y = (\rho_Y)^*(\rho_Y)_*\mathscr{L}_Y \to \mathscr{L}_Y$  is surjective. The universal property of projectivisation, [Har77, II Prop. 7.12], thus gives a morphism  $\alpha: \mathbb{P}_Y \to \mathbb{P}_Y(\mathscr{F}_Y)$  that identifies  $\mathscr{L}_Y$  with the pull-back of  $\mathcal{O}_{\mathbb{P}_Y(\mathscr{F}_Y)}(1)$ . But since the identification  $\mathscr{L}_Y|_{(\rho_Y)^{-1}(y)} \cong \mathcal{O}_{\mathbb{P}^r}(1)$  now holds for every  $y \in Y$ , the morphism  $\alpha$  is clearly bijective, hence isomorphic by Zariski's main theorem.

4.4. **Proof of Theorem 4.1: End of proof.** It remains to show that  $\mathscr{F}_X \cong \varphi^* \mathscr{F}_Y$ . Recalling that

$$\mathscr{F}_Y = (\rho_Y)_* \mathscr{L}_Y$$
 and  $\mathscr{F}_X = (\rho_X)_* \mathscr{L}_X = (\rho_X)_* \Phi^* \mathscr{L}_Y$ ,

there exists a natural morphism  $\alpha: \varphi^* \mathscr{F}_Y \to \mathscr{F}_X$ , cf. [Har77, III Rem. 9.3.1]. The restriction of the morphism  $\alpha$  to the open set  $X^\circ$  is clearly isomorphic. We claim that it is isomorphic everywhere. To this end, consider its determinant  $\det \alpha: \varphi^* \mathscr{M}_Y \to \mathscr{M}_X$ . Using that  $\varphi^* \mathscr{M}_Y \cong \mathscr{M}_X$ , the determinant can be seen as a section in  $\mathscr{H}om(\mathscr{M}_X, \mathscr{M}_X) \cong \mathscr{O}_X$ , and hence as a function on X that does not vanish on  $X^\circ$ . But since the  $\varphi$ -exceptional set  $E:=X\setminus X^\circ$  is contracted to the small subvariety  $Y\setminus Y^\circ$  of Y, the function in fact cannot vanish anywhere. It follows that the morphism  $\det \alpha$  is isomorphic, and hence so is  $\alpha$ . This ends the proof of Theorem 4.1.

## 5. Descent of vector bundles to KLT spaces

In the present section, we will prove the following theorem, a simplified form of which appeared as Theorem 1.2 in the introduction.

**Theorem 5.1** (Descent of vector bundles to klt spaces). Let  $f: X \to Y$  be a birational, projective morphism of normal, quasi-projective varieties. Assume Y to be a klt space. Then, the following holds.

- (5.1.1) If  $\mathscr{F}_X$  is any locally free, f-numerically flat sheaf on X, then there exists a locally free sheaf  $\mathscr{F}_Y$  on Y such that  $\mathscr{F}_X \cong f^*\mathscr{F}_Y$ .
- (5.1.2) If  $(\mathscr{F}_X, \theta_X)$  is any locally free Higgs sheaf on X, where  $\mathscr{F}_X$  is f-numerically flat, then there exists a locally free Higgs sheaf  $(\mathscr{F}_Y, \theta_Y)$  on Y such that  $(\mathscr{F}_X, \theta_X) \cong f^*(\mathscr{F}_Y, \theta_Y)$ .
- 5.1. **Proof of Theorem 5.1: Setup and Notation.** We maintain notation and assumptions of Theorem 5.1 throughout the present Section 5. Choose an effective Weil  $\mathbb{Q}$ -divisor  $\Delta_Y$  such that  $(Y, \Delta_Y)$  is klt. We denote by  $\Delta_X := f_*^{-1} \Delta_Y$  the strict transform of  $\Delta_Y$ , and by  $E_X := \operatorname{Exc}(f)$  the divisorial part of the f-exceptional locus  $\operatorname{Exc}(f)$ .
- 5.2. **Proof of Statement (5.1.1).** Consider a resolution of singularities,  $\pi: \widetilde{X} \to X$ . If we can show that  $\pi^* \mathscr{F}_X$  is of the form  $(f \circ \pi)^* \mathscr{F}_Y$  for a suitable sheaf  $\mathscr{F}_Y$  on Y, then  $\mathscr{F}_X$  will be isomorphic to  $f^* \mathscr{F}_Y$  by the projection formula. We are therefore free to replace X by  $\widetilde{X}$  and assume without loss of generality that the following holds.

Assumption w.l.o.g. 5.2. The variety X is smooth, the f-exceptional set equals  $E_X$ , and  $\operatorname{supp}(\Delta_X + E_X)$  is an snc divisor in X.

Step 1: Factorisation via an f-relative MMP. We will factor the resolution  $f: X \to Y$  via a relative minimal model program of X over Y, cf. the discussion in [GKKP11, Sect. 23], whose organisation we will follow closely. By the definition of "klt pair" there exist effective f-exceptional divisors F and G without common components such that  $\lfloor F \rfloor = 0$  and such that the following  $\mathbb{Q}$ -linear equivalence holds:

$$K_X + \Delta_X + F \sim_{\mathbb{Q}} f^*(K_Y + \Delta_Y) + G.$$

For  $\varepsilon \in (0,1) \cap \mathbb{Q}$  we let  $\Delta_{\varepsilon} := \Delta_X + F + \varepsilon \cdot E$ . For  $0 < \varepsilon \ll 1$  small enough, the pair  $(X,\Delta)$  is klt. Fix one such  $\varepsilon$  and let  $H \in \mathbb{Q}$  Div(X) be an f-ample divisor such that  $(X,\Delta_{\varepsilon} + H)$  is still klt and  $K_X + \Delta_{\varepsilon} + H$  is f-nef. We may then run the f-relative  $(X,\Delta_{\varepsilon})$  minimal model program with scaling of H, cf. [BCHM10, Cor. 1.4.2] to obtain a diagram

$$X = X_{n} - \stackrel{\varphi_{n}}{\longrightarrow} X_{n-1} \stackrel{\varphi_{n-1}}{\longrightarrow} \cdots - \stackrel{\varphi_{1}}{\longrightarrow} X_{0}$$

$$f \downarrow \qquad \qquad f_{n} \downarrow \qquad \qquad f_{n-1} \downarrow \qquad \qquad f_{0} \downarrow$$

$$Y = Y = Y = Y = Y$$

with the following properties.

- (5.2.1) The spaces  $X_i$  are  $\mathbb{Q}$ -factorial. Writing  $\Delta_{X_i}$  for the cycle-theoretic image of  $\Delta_{\varepsilon}$ , the pairs  $(X_i, \Delta_{X_i})$  are klt.
- (5.2.2) The maps  $\varphi_i$  are either divisorial contractions of  $(K_{X_i} + \Delta_{X_i})$ -negative extremal rays in  $R_i \subseteq \overline{NE}(X_i/Y)$  or flips associated to small contractions of such rays.
- (5.2.3) The log-canonical divisor  $K_{X_0} + \Delta_{X_0}$  is  $f_0$ -nef, and it hence follows from the negativity lemma, [GKKP11, Lem. 2.16.2], that the morphism  $f_0$  is small and crepant, cf. [GKKP11, Claim 23.4]. In other words,  $\Delta_{X_0} = (f_0)_*^{-1}(\Delta_Y)$  and  $K_{X_0} + \Delta_{X_0} \sim_{\mathbb{Q}} (f_0)^*(K_Y + \Delta_Y)$ . As a consequence, also  $-(K_{X_0} + \Delta_{X_0})$  is  $f_0$ -nef.

Step 2: Construction of bundles. Next, we construct vector bundles on the  $X_i$ .


Claim 5.3. There exist locally free sheaves  $\mathscr{F}_{X_i}$  on the varieties  $X_i$  such that the following holds

- (5.3.1) The sheaf  $\mathscr{F}_{X_n}$  equals  $\mathscr{F}_{X}$ .
- (5.3.2) Given any index *i*, the sheaf  $\mathscr{F}_{X_i}$  is  $f_i$ -numerically flat.
- (5.3.3) If i > 0 is any index such that  $\mathscr{F}_{X_{i-1}}$  is isomorphic to  $(f_{i-1})^*\mathscr{G}$  for a locally free sheaf  $\mathscr{G}$  on Y, then  $\mathscr{F}_{X_i} \cong (f_i)^*\mathscr{G}$ .

*Proof of Claim 5.3.* We construct the vector bundles inductively. Start by setting  $\mathscr{F}_{X_n} := \mathscr{F}_X$ . Next, assume that we are given an index i > 0 for which vector bundles  $\mathscr{F}_{X_n}$ , ...,  $\mathscr{F}_{X_i}$  have already been constructed. We consider the cases where  $\varphi_i$  is a divisorial contraction and where it is a flip separately.

Divisorial contraction. If  $\varphi_i: X_i \to X_{i-1}$  is a divisorial contraction, then  $-(K_{X_i} + \Delta_{X_i})$  is  $\varphi_i$ -nef. Theorem 4.1 ("Descent of vector bundles from klt spaces") hence proves the existence of a locally free sheaf  $\mathscr{F}_{X_{i-1}}$  on  $X_{i-1}$  such that  $\mathscr{F}_{X_i} \cong \varphi_i^* \mathscr{F}_{X_{i-1}}$ . This isomorphism guarantees that Properties (5.3.2) and (5.3.3) both hold.

*Flip.* If  $\varphi_i: X_i \to X_{i-1}$  is a flip, consider the associated "flipping diagram",



where  $\alpha$  is obtained by contracting a  $(K_{X_i} + \Delta_{X_i})$ -extremal ray, which implies as above that  $-(K_{X_i} + \Delta_{X_i})$  is  $\alpha$ -nef. In this setting, Theorem 4.1 again proves the existence of a locally free sheaf  $\mathscr{F}_Z$  on Z such that  $\mathscr{F}_{X_i} \cong \alpha^* \mathscr{F}_Z$ . Properties (5.3.2) and (5.3.3) will hold once we set  $\mathscr{F}_{X_{i-1}} := \beta^* \mathscr{F}_Z$ . This finishes the proof of Claim 5.3.

Step 3: End of proof. To end the proof of Statement (5.1.1), recall from (5.2.3) that  $(X_0, \Delta_{X_0})$  is klt and that  $-(K_{X_0} + \Delta_{X_0})$  is  $f_0$ -nef. Since  $\mathscr{F}_{X_0}$  is  $f_0$ -numerically flat by (5.3.2), we may therefore apply Theorem 4.1 to obtain a locally free sheaf  $\mathscr{F}_Y$  on Y such that  $\mathscr{F}_{X_0} \cong (f_0)^*\mathscr{F}_Y$ . A repeated application of Property (5.3.3) then shows that the sheaves  $\mathscr{F}_{X_1}$ ,  $\mathscr{F}_{X_2}$ , ...,  $\mathscr{F}_{X_n} = \mathscr{F}$  are all pull-backs of  $\mathscr{F}_Y$ . Statement (5.1.1) follows.

## 5.3. **Proof of Statement (5.1.2).** For $\bullet \in \{X, Y\}$ , set

$$\mathscr{A}^i_{\bullet} := \mathscr{H}om(\mathscr{F}_{\bullet}, \mathscr{F}_{\bullet} \otimes \Omega^{[i]}_{\bullet}).$$

A Higgs fields on  $\mathscr{F}_Y$  is, by definition, a section of the sheaf  $\mathscr{A}_Y^1$  such that the induced section in  $\mathscr{A}_Y^2$  vanishes. In our case, the sheaf  $\mathscr{F}_Y$  is locally free, which implies that both  $\mathscr{A}_Y^1$  and  $\mathscr{A}_Y^2$  are reflexive. To give a Higgs field on  $\mathscr{F}_Y$  it is therefore equivalent to give a Higgs field on the restriction of  $\mathscr{F}_Y$  to any big open subset of Y. The Higgs field  $\theta_X$ , however clearly induces a Higgs field on the restriction of  $\mathscr{F}_Y$  to the big open set where  $f^{-1}$  is well-defined and isomorphic. The existence of  $\theta_Y$  follows.

It remains to show that  $f^*(\mathscr{F}_Y, \theta_Y) \cong (\mathscr{F}_X, \theta_X)$ . In other words, we need to show that the two sections  $\theta_X$  and  $f^*\theta_Y \in H^0(X, \mathscr{A}_X^1)$  agree. They will clearly agree over the open set  $f^{-1}(Y^\circ)$ . Since  $\mathscr{F}_X$  is locally free and  $\mathscr{A}_X^1$  therefore reflexive, this suffices to show that they are the same. This finishes the proof of Theorem 5.1.

#### 6. The restriction theorem for semistable Higgs sheaves

The proof of the nonabelian Hodge correspondence uses the following restriction theorem for (semi)stable Higgs sheaves, which generalises a number of earlier results including [GKPT15, Thm. 5.22].

**Theorem 6.1** (Restriction of (semi)stable Higgs sheaves). Let X be a normal, projective variety,  $\dim X \geq 2$ , and let  $H \in \operatorname{Div}(X)$  be big and semiample. Given any torsion free Higgs sheaf  $(\mathscr{E}^{\circ}, \theta^{\circ})$  on  $X_{\text{reg}}$  that is semistable (resp. stable) with respect to H in the sense of Definition 2.25, there exists an integer  $M \in \mathbb{N}^+$  satisfying the following conditions: If  $B \subseteq |m \cdot H|$  is any basepoint free linear system with m > M, then there exists a dense, open subset  $B^{\circ} \subseteq B$  such that the following properties hold for all  $D \in B^{\circ}$ .

- (6.1.1) The hypersurface D is irreducible and normal, and  $D_{\text{reg}} = D \cap X_{\text{reg}}$ .
- (6.1.2) The Higgs sheaf  $(\mathscr{E}^{\circ}, \theta^{\circ})|_{D_{reg}}$  is torsion free and semistable (resp. stable) with respect to  $H|_{D}$ .

Remark 6.2 (Algebraicity assumption). We stress that the Higgs sheaf  $(\mathcal{E}^{\circ}, \theta^{\circ})$  of Theorem 6.1 is assumed to be algebraic.

Remark 6.3 (Restriction theorem for sheaves on X). Recalling from Lemma 2.26 that a Higgs sheaf on X is semistable (resp. stable) if and only if its restriction to  $X_{\text{reg}}$  is semistable (resp. stable), Theorem 6.1 immediately implies a restriction theorem for torsion free Higgs sheaves on X that is more general than the results found in the literature. In practical applications, the variety X might admit a finite group action, and the linear system  $B \subseteq |m \cdot H|$  might be chosen to contain invariant divisors only.

Theorem 6.1 is shown in Section 6.2 below. The following corollary discusses the behaviour of semistability under pull-back. It complements [GKPT15, Sect. 5.6], where the (G-)stable case was discussed. Its proof, spelled out in the arXiv version of this paper, arXiv:1711.08159, applies Theorem 6.1 repeatedly to cut down to a curve, where the result is classically known.

**Corollary 6.4** (Semistability under generically finite morphisms). Let X and Y be two projective, klt spaces. Let  $H \in Div(X)$  be big and semiample, and let  $f: Y \to X$  be a surjective and generically finite morphism. Let  $(\mathcal{E}, \theta)$  be a reflexive Higgs sheaf on X.

- ullet If  $\mathscr E$  is locally free, then the following are equivalent.
  - (6.4.1) The Higgs bundle  $(\mathcal{E}, \theta)$  is semistable with respect to H.
  - (6.4.2) The Higgs bundle  $f^*(\mathcal{E}, \theta)$  is semistable with respect to  $f^*H$ .
- If Y is smooth, then the following are equivalent.
  - (6.4.3) The Higgs sheaf  $(\mathcal{E}, \theta)$  is semistable with respect to H.
  - (6.4.4) The Higgs sheaf  $f^{[*]}(\mathcal{E},\theta)$  is semistable with respect to  $f^*H$ .

Remark 6.5. One might wonder why the assumptions in Corollary 6.4 are so much more restrictive compared to Theorem 6.1. If X is not klt, then pull-back of Higgs sheaves does not exist in general. If the sheaf  $\mathscr E$  is not locally free and Y is not smooth, its pull-back is generally neither reflexive nor torsion free, and no good notion of "stability" is defined in this case. Also, we do not know whether the reflexive pull-back  $f^{[*]}\mathscr E$  carries a natural Higgs field in this case.

6.1. **Restriction theorem for sheaves with operators.** The following restriction theorem for sheaves with operators is a generalisation of [GKPT15, Thm. A.3 in the arXiv version, arXiv:1511.08822]. It serves as the main technical tool used in the proof of the restriction theorem for Higgs sheaves, Theorem 6.1.

**Theorem 6.6** (Restriction theorem for sheaves with operators). Let X be a normal, projective variety,  $\dim X \geq 2$ , let H be a big and semiample divisor on X, and  $\mathcal{W}^{\circ}$  be a reflexive sheaf on  $X_{\text{reg}}$ . Let  $(\mathcal{E}^{\circ}, \theta^{\circ})$  be a torsion free sheaf on  $X_{\text{reg}}$  with a  $\mathcal{W}^{\circ}$ -valued operator, and assume that  $(\mathcal{E}^{\circ}, \theta^{\circ})$  is semistable (resp. stable) with respect to H. Then, there exists  $M \in \mathbb{N}^+$  satisfying the following conditions: If  $B \subseteq |m \cdot H|$  is any basepoint free linear system with m > M, then there exists a dense, open subset  $B^{\circ} \subseteq B$  such that the following properties hold for all  $D \in B^{\circ}$ .

- (6.6.1) The hypersurface D is irreducible and normal, and  $D_{reg} = X_{reg} \cap D$ .
- (6.6.2) The sheaf  $\mathscr{E}^{\circ}|_{D_{reg}}$  is torsion free, and  $(\mathscr{E}^{\circ}|_{D_{reg}}, \theta^{\circ}|_{D_{reg}})$  is semistable (resp. stable) with respect to  $H|_{D}$ , as a sheaf with a  $\mathscr{W}|_{D_{reg}}$ -valued operator.

Remark 6.7 (Algebraicity and restriction theorem for sheaves on X). As before, we underline that  $(\mathscr{E}^{\circ}, \theta^{\circ})$  is assumed to be algebraic. Also as before, Theorem 6.6 implies a restriction theorem for sheaves  $\mathscr{E}$  with operators that are defined on all of X. In particular, if  $\mathscr{E}$  is a torsion free sheaf on X (=sheaf equipped with the zero operator) that is not necessarily semistable, then  $\mu_H^{\max}(\mathscr{E}) = \mu_{H|D}^{\max}(\mathscr{E}|_D)$  for all  $D \in B^{\circ}$ , and the Harder-Narasimhan filtration of  $\mathscr{E}|_D$  equals the restriction of the Harder-Narasimhan filtration of  $\mathscr{E}$ , in symbols:  $\mathrm{HN}^{\bullet}_H(\mathscr{E})|_D = \mathrm{HN}^{\bullet}_H(\mathscr{E}|_D)$ .

*Proof of Theorem 6.6.* Let  $\iota: X^\circ \to X$  be the inclusion map. Set  $\mathscr E = \iota_*(\mathscr E^\circ)$  and  $\mathscr W = \iota_*(\mathscr W^\circ)$ . The sheaf  $\mathscr E$  is torsion free; furthermore,  $\mathscr W$  is reflexive and therefore embeds into a locally free sheaf  $\mathscr V$ . Composing morphisms as in Lemma 2.27, the operator  $\theta^\circ$  induces an operator  $\tau^\circ: \mathscr E^\circ \to \mathscr E^\circ \otimes \mathscr V|_{X_{\text{reg}}}$ , and Lemma 2.27 shows that  $(\mathscr E^\circ, \tau^\circ)$ , considered as a sheaf with a  $\mathscr V|_{X_{\text{reg}}}$ -valued operator, is again semistable (resp. stable) with respect to H. Pushing forward, we can extend the operator  $\tau^\circ$ , which a priori is only defined on  $X_{\text{reg}}$ , to an operator  $\tau: \mathscr E \to \mathscr E \otimes \mathscr V$ . Lemma 2.26 then shows that  $(\mathscr E, \tau)$  is again semistable (resp. stable) with respect to H, when considered as a sheaf with a  $\mathscr V$ -valued operator. We claim the following.

(6.8.1) There exists a number  $M \in \mathbb{N}^+$  and for every m > M and for every basepoint free linear system  $B \subseteq |m \cdot B|$  an open, dense subsystem  $B^\circ$  such that (6.6.1) holds, such that  $\mathscr{E}|_D$  is torsion free, and such that  $(\mathscr{E}|_D, \theta|_D)$  is semistable (resp. stable) with respect to  $H|_D$ , as a sheaf with a  $\mathscr{V}|_D$ -valued operator.

As soon as (6.8.1) is true, Lemma 2.26 asserts that  $(\mathscr{E}|_{D_{\text{reg}}}, \tau|_{D_{\text{reg}}}) = (\mathscr{E}^{\circ}|_{D_{\text{reg}}}, \tau^{\circ}|_{D_{\text{reg}}})$  is semistable (resp. stable) with respect to H as a sheaf with an  $\mathscr{V}|_{D_{\text{reg}}}$ -valued operator. A final application of Lemma 2.27 then shows that  $(\mathscr{E}^{\circ}|_{D_{\text{reg}}}, \theta^{\circ}|_{D_{\text{reg}}})$  is semistable (resp. stable) with respect to H, which would end the proof.

It remains to show (6.8.1). To this end, observe that for any resolution of singularities,  $\pi: \widetilde{X} \to X$ , the operator  $\tau$  induces an operator on the torsion free pullback of  $\mathscr{E}$  to  $\widetilde{X}$ ,

$$\widetilde{\tau}: \pi^* \mathcal{E} \bigg/ \mathrm{tor} \to \pi^* (\mathcal{E} \otimes \mathcal{V}) \bigg/ \mathrm{tor} = \pi^* \mathcal{E} \bigg/ \mathrm{tor} \otimes \pi^* \mathcal{V}.$$

Replacing *X* by a suitable resolution and replacing  $(\mathscr{E}, \tau)$  by  $(\pi^*\mathscr{E}/\text{tor}, \widetilde{\tau})$ , we may assume without loss of generality that *X* is smooth and that  $\mathscr E$  is locally free. Now, if  $(\mathscr E, \tau)$  is stable with respect to H, the stability claim of Theorem 6.6 has been shown already for every  $D \in B$  fulfilling (6.6.1) and having the property that  $\mathscr{E}|_D$  is torsion free, see [Lan15, Thm. 9] and [GKPT15, Thm. A.3 in the arXiv version]. These two assumptions can be guaranteed for D belonging to an open subsystem B° by the classical Bertini theorem and by [Gro66, Thm. 12.2.1], respectively. In case where  $(\mathcal{E}, \tau)$  only semistable, the assumption that  $\mathcal{V}$ is a locally free guarantees the existence of a Jordan-Hölder filtration, which presents  $(\mathscr{E}, \tau)$  as a repeated extension of stable sheaves with  $\mathscr{V}$ -valued operator of equal slope. The claim then follows by induction on the length of the filtration.

6.2. **Proof of Theorem 6.1.** For convenience of notation, write  $n := \dim X$ , let  $\iota : X_{\text{reg}} \to X$ X be the inclusion map, and let  $\mathscr{E} := \iota_* \mathscr{E}^{\circ}$  be the torsion free extension of  $\mathscr{E}^{\circ}$  to X. Twisting the Higgs sheaf  $(\mathscr{E}^{\circ}, \theta^{\circ})$  with a sufficiently positive line bundle and noticing that semistability considerations are unaffected by this operation, we may also assume that the following holds.

Assumption w.l.o.g. 6.9. The slope of  $\mathscr{E}$  is positive,  $\mu_H(\mathscr{E}) > 0$ .

*Choice* 6.10. Choose  $M \gg 0$  large enough so that the following holds.

- (6.10.1) If m > M and  $B \subset |mH|$  is base point free, then Theorem 6.6 and Remark 6.7 apply to yield a dense, open set  $B' \subset B$  such that for every  $D \in B'$ , we have  $\mu_H^{\max}(\mathscr{E}) = \mu_{H|_D}^{\max}(\mathscr{E}|_D).$
- (6.10.2) The number M is larger than the number given by Theorem 6.6 for  $(\mathcal{E}^{\circ}, \theta^{\circ})$  as a sheaf with an  $\Omega^1_{X_{\mathrm{reg}}}$  -valued operator. (6.10.3) We have  $[M\cdot H]^n > nr\cdot \mu_H^{\mathrm{max}}(\mathscr{E})$ .

Step 1: Argument by contradiction. Assume for the remainder of the proof that we are given a number m > M and a basepoint free linear system  $B \subseteq |m \cdot H|$ . Let  $B^{\circ}$  be the intersection of the open subsets given by the two applications of (6.10.1) and (6.10.2) above, and let  $D \in B^{\circ}$  be any element. We aim to show that  $(\mathscr{E}^{\circ}, \theta^{\circ})|_{D_{\text{reg}}}$  is semistable (resp. stable) with respect to H. We argue by contradiction and assume that this is not the case.

Assumption 6.11. There exists a generically Higgs-invariant, saturated subsheaf 0 ⊊  $\mathscr{F}_{D_{\text{reg}}} \subsetneq \mathscr{E}^{\circ}|_{D_{\text{reg}}} \text{ with } \mu_{H|_{D}}(\mathscr{F}_{D_{\text{reg}}}) > \mu_{H}(\mathscr{E}) \text{ (resp. } \geq \text{ instead of } >).$ 

Step 2: Cutting down. Repeated applications of Theorem 6.6 (and Remark 6.7) allow to find an increasing sequence of numbers  $M < m \le m_2 \le \cdots \le m_{n-1}$  and hyperplanes  $D_i \in [m_i \cdot H]$  such that the associated intersection  $C := D \cap D_2 \cap \cdots \cap D_{n-1}$  has the following properties.

- (6.12.1) The scheme C is a smooth curve, and entirely contained in  $X_{\text{reg}}$ . The sheaves  $\mathscr{E}|_C$  and  $\mathscr{F}_{D_{\text{reg}}}|_C$  are torsion free and hence locally free. The natural morphism  $\mathscr{F}_{D_{\text{reg}}}|_{C} \to \mathscr{E}|_{C}^{\circ}$  is an injection. (6.12.2) We have  $\mu_{H}^{\max}(\mathscr{E}) = \mu_{H|_{C}}^{\max}(\mathscr{E}|_{C})$ .
- (6.12.3) Because of (6.10.2), the restriction ( $\mathscr{E}|_C$ ,  $\theta^{\circ}|_C$ ) is semistable (resp. stable) as a sheaf with an  $\Omega_X^1|_C$ -valued operator.

П

Step 3: Computation. In the following, write  $\mathscr{F}_C := \mathscr{F}_{D_{\text{reg}}}|_C$  and consider the associated sequence

$$(6.13.1) 0 \longrightarrow \mathscr{F}_C \longrightarrow \mathscr{E}|_C \xrightarrow{q} \mathscr{Q} \longrightarrow 0.$$

Let  $N_{C/X}^*$  denote the conormal bundle of C in X. If  $\mathscr{B} \subseteq \mathscr{Q} \otimes N_{C/X}^*$  is any coherent subsheaf of positive rank, then we will show in this step that

(6.13.2) 
$$\deg_{\mathcal{C}}(\mathscr{B}) \le (n-1)r \cdot \mu_H^{\max}(\mathscr{E}) - [m \cdot H]^n.$$

To prove (6.13.2), consider first any coherent subsheaf  $\mathscr{A} \subseteq \mathscr{Q}$  of positive rank. We obtain from (6.13.1) an exact sequence  $0 \to \mathscr{F}_C \to q^{-1} \mathscr{A} \xrightarrow{q} \mathscr{A} \to 0$ , which allows to estimate the degree of  $\mathscr{A}$  as follows,

$$\begin{split} \deg_C(\mathscr{A}) &= \deg_C(q^{-1}\mathscr{A}) - \deg_C(\mathscr{F}_C) \leq \deg_C(q^{-1}\mathscr{A}) & \text{ since } \mu_{H|_C}(\mathscr{F}_C) \geq \mu_H(\mathscr{E}) > 0 \\ &\leq \operatorname{rank}(q^{-1}\mathscr{A}) \cdot \mu_{H|_C}(q^{-1}\mathscr{A}) & \text{ definition of slope} \\ &\leq \operatorname{rank}(\mathscr{E}) \cdot \mu_H^{\max}(\mathscr{E}) & \text{ Item (6.12.2)} \end{split}$$

In order to apply this inequality to the problem at hand, recall that C is constructed as a complete intersection. The normal bundle of C in X is hence described as

$$N_{C/X} = \bigoplus_i N_{D_i/X}|_C$$
, where  $N_{D_i/X}|_C \cong \mathscr{O}_C(m \cdot H|_C)$  for all  $i$ .

We can therefore view  $\mathscr{A} := \mathscr{B}(m \cdot H|_C)$  as a subsheaf of  $\mathscr{Q}^{\oplus n-1}$ . An induction using the inequality obtained above then shows the following, which immediately implies (6.13.2),

$$\deg_{C}(\mathscr{B}) + \operatorname{rank}(\mathscr{B}) \cdot m \cdot \deg_{C}(H|_{C}) = \deg_{C}\mathscr{B}(m \cdot H|_{C}) \stackrel{\operatorname{Induction}}{\leq} (n-1)r \cdot \mu_{H}^{\max}(\mathscr{E}).$$

Step 4: End of proof. Consider the operator  $\theta^{\circ}|_{C}$  and its restriction  $\theta_{\mathscr{F}}: \mathscr{F}_{C} \to \mathscr{E}|_{C} \otimes \Omega^{1}_{X}|_{C}$ . The target of  $\theta_{\mathscr{F}}$  appears in the following commutative square,

$$\begin{split} \mathscr{E}|_C \otimes \Omega^1_X|_C & \longrightarrow \mathscr{E}|_C \otimes \Omega^1_C \\ q \otimes \operatorname{Id} \downarrow & \downarrow \\ 0 & \longrightarrow \mathscr{Q} \otimes N^*_{C/X} & \xrightarrow{\alpha} \mathscr{Q} \otimes \Omega^1_X|_C & \xrightarrow{\beta} \mathscr{Q} \otimes \Omega^1_C & \longrightarrow 0. \end{split}$$

Assumption 6.11 and Item (6.12.3) guarantees that  $\mathscr{F}_C$  is not invariant with respect to  $\theta^\circ|_C$ . The composed map  $(q\otimes \mathrm{Id})\circ\theta_\mathscr{F}$  is therefore *not* the zero morphism. But since  $\mathscr{F}_C$  is a Higgs-invariant subsheaf of  $(\mathscr{E}^\circ,\theta^\circ)|_C$  by assumption, the morphism  $\beta\circ(q\otimes \mathrm{Id})\circ\theta_\mathscr{F}$  is zero. In summary, we obtain a non-trivial morphism  $\tau:\mathscr{F}_C\to\mathscr{Q}\otimes N^*_{C/X}$ . But then, we compute degrees as follows,

$$\begin{split} (n-1)r \cdot \mu_H^{\max}(\mathcal{E}) - [m \cdot H]^n & \geq \deg(\operatorname{img} \tau) & \operatorname{Inequality} \ (6.13.2) \\ & = \deg_C(\mathcal{F}_C) - \deg_C(\ker \tau) \\ & \geq -\deg_C(\ker \tau) & \operatorname{Assumptions} \ 6.9, \ 6.11 \\ & \geq -\operatorname{rank}(\ker \tau) \cdot \mu_H^{\max}(\mathcal{E}) & \operatorname{Item} \ (6.12.2) \\ & \geq -r \cdot \mu_H^{\max}(\mathcal{E}) & \operatorname{Assumption} \ 6.9. \end{split}$$

We obtain a contradiction to (6.10.3), which finishes the proof of Theorem 6.1.

## 7. Ascent of semistable Higgs bundles

The proof of the nonabelian Hodge correspondence on klt spaces relies on the following result, which can be seen as an inverse to the descent results obtained in the first part of this paper. It asserts that the pull-back of a Higgs sheaf in  $\operatorname{Higgs}_X$  to any resolution of singularities is again in  $\operatorname{Higgs}_{\bullet}$ .

**Theorem 7.1** (Ascent of semistable Higgs bundles). Let X be a projective, klt space and let  $\pi: \widetilde{X} \to X$  be a resolution of singularities. If  $(\mathscr{E}, \theta) \in \text{Higgs}_{\widetilde{X}}$ , then  $\pi^*(\mathscr{E}, \theta) \in \text{Higgs}_{\widetilde{X}}$ .

The following is an almost immediate consequence of Theorem 7.1 and Theorem 5.1, Item (5.1.2); see also Fact 7.7 below.

**Corollary 7.2** (Boundedness of Higgs sheaves). Let X be a projective, klt space and let  $r \in \mathbb{N}^+$  be any number. Let F be the family of locally free Higgs sheaves  $(\mathscr{F}, \Theta) \in \text{Higgs}_X$  with rank  $\mathscr{F} = r$ . Then, the family F is bounded.

Theorem 7.1 will be shown in Sections 7.1–7.6 below. The main difficulty here is that it is not clear from the outset that a pull-back of semistable Higgs bundle via the resolution map is again semistable with respect to an *ample* divisor. It turns out that the assumption of vanishing Chern classes is sufficient to resolve this problem in dimension two. The higher-dimensional case will be reduced to the surface case by restriction techniques.

7.1. **Preparation for the proof of Theorem 7.1: Boundedness.** We will use the following iterated Bertini-type theorem for bounded families of Higgs sheaves, generalising [GKP16, Cor. 5.3], where the same result was shown for reflexive sheaves without Higgs field that are defined on a projective variety, and not just on some big open subset.

**Proposition 7.3** (Iterated Bertini-type theorem for bounded families on  $X_{\text{reg}}$ ). Let X be a normal, projective variety of dimension  $n \geq 2$ , let  $H \in \text{Div}(X)$  be ample and let  $X^{\circ} \subseteq X_{\text{reg}}$  be a big, open subset. Let  $(\mathscr{E}_{X^{\circ}}, \theta_{\mathscr{E}_{X^{\circ}}})$  be a locally free Higgs sheaf on  $X^{\circ}$ , and let  $F^{\circ}$  be a bounded family of locally free Higgs sheaves on  $X^{\circ}$ . If  $m \gg 0$  is large enough and if  $(D_1, D_2, \ldots, D_{n-1}) \in |m \cdot H|^{\times n-1}$  is a general tuple with associated complete intersection curve  $C := D_1 \cap \cdots \cap D_{n-1}$ , then  $C \subset X^{\circ}$  is smooth and the following holds for all Higgs sheaves  $(\mathscr{F}_{X^{\circ}}, \theta_{\mathscr{F}_{X^{\circ}}})$  in  $F^{\circ}$ :

$$(\mathcal{F}_{X^\circ},\theta_{\mathcal{F}_{X^\circ}})\cong(\mathcal{E}_{X^\circ},\theta_{\mathcal{E}_{X^\circ}})\quad\Leftrightarrow\quad (\mathcal{F}_{X^\circ},\theta_{\mathcal{F}_{X^\circ}})|_C\cong(\mathcal{E}_{X^\circ},\theta_{\mathcal{E}_{X^\circ}})|_C.$$

*Proof.* As a first step in the proof, we extend all relevant sheaves from  $X^{\circ}$  to X. To this end, let  $\mathscr{E}_X$  be the unique reflexive sheaf on X whose restriction to  $X^{\circ}$  is  $\mathscr{E}_{X^{\circ}}$ . Doing the same with the sheaves that appear in  $\mathscr{F}^{\circ}$ , observe that the family

 $F := \{\text{isomorphism classes of reflexive } \mathscr{F}_X \text{ on } X \text{ such that there exists} \}$ 

a member 
$$(\mathscr{F}_{X^{\circ}}, \theta_{\mathscr{F}_{X^{\circ}}})$$
 in  $\mathsf{F}^{\circ}$  with  $\mathscr{F}_{X}|_{X^{\circ}} \cong \mathscr{F}_{X^{\circ}}$ 

is likewise bounded. This fact is crucial in the proof of the following two claims.

*Claim* 7.4. Setting  $D_1^{\circ} := D_1 \cap X^{\circ}$ , the restriction maps

$$(7.4.1) \qquad \operatorname{Hom}_{X^{\circ}}(\mathscr{F}_{X^{\circ}}, \mathscr{E}_{X^{\circ}}) \to \operatorname{Hom}_{D_{\circ}^{\circ}}(\mathscr{F}_{X^{\circ}}|_{D_{\circ}^{\circ}}, \mathscr{E}_{X^{\circ}}|_{D_{\circ}^{\circ}})$$

$$(7.4.3) \qquad \operatorname{Hom}_{X^{\circ}}\left(\mathscr{F}_{X^{\circ}}, \, \mathscr{F}_{X^{\circ}} \otimes \Omega^{1}_{X^{\circ}}\right) \to \operatorname{Hom}_{C}\left(\mathscr{F}_{X^{\circ}}|_{C}, \, \mathscr{F}_{X^{\circ}}|_{C} \otimes \Omega^{1}_{X^{\circ}}|_{C}\right)$$

are isomorphic, for all  $(\mathscr{F}_{X^{\circ}}, \theta_{\mathscr{F}_{X^{\circ}}})$  in  $F^{\circ}$ .

*Proof of Claim 7.4.* For brevity, we consider (7.4.1) only and leave the rest to the reader. Since  $m \gg 0$  is assumed to be large, we have vanishing

$$h^0(X, \mathcal{H}om(\mathscr{F}_X, \mathscr{E}_X) \otimes \mathscr{J}_D) = h^1(X, \mathcal{H}om(\mathscr{F}_X, \mathscr{E}_X) \otimes \mathscr{J}_D) = 0$$

for all members  $\mathscr{F}_X$  of the bounded family F. In fact, vanishing for  $h^1$  follows from [Gro68, Exp. XII, Prop. 1.5] since X is normal, which implies by [Har80, Prop. 1.3] that the sheaves  $\mathscr{H}om(\mathscr{F}_X,\mathscr{E}_X)\otimes\mathscr{J}_D$  have depth  $\geq 2$  at every point of X. As a consequence, we obtain that the natural restriction maps,

(7.4.4) 
$$\operatorname{Hom}_X(\mathscr{F}_X, \mathscr{E}_X) \to H^0(D_1, \mathscr{H}om(\mathscr{F}_X, \mathscr{E}_X)|_{D_1})$$

are isomorphic for all  $\mathscr{F}_X$  in F. In order to relate (7.4.4) to the problem at hand, use boundedness of F again and recall from [HL10, Cor. 1.1.14] that the sheaves  $\mathscr{F}_X|_{D_1}$ ,  $\mathscr{E}_X|_{D_1}$  and  $\mathscr{H}om(\mathscr{F}_X,\mathscr{E}_X)|_{D_1}$  are reflexive on the normal variety  $D_1$ , for all  $\mathscr{F}_X$  in F. As a consequence, we find that the natural restriction morphisms,

$$\operatorname{Hom}_{X} \left( \mathscr{F}_{X}, \, \mathscr{E}_{X} \right) \to \operatorname{Hom}_{X^{\circ}} \left( \mathscr{F}_{X^{\circ}}, \, \mathscr{E}_{X^{\circ}} \right)$$

$$H^{0} \left( D_{1}, \, \mathscr{H}om(\mathscr{F}_{X}, \mathscr{E}_{X})|_{D_{1}} \right) \to \operatorname{Hom}_{D_{1}^{\circ}} \left( \mathscr{F}_{X^{\circ}}|_{D_{1}^{\circ}}, \, \mathscr{E}_{X^{\circ}}|_{D_{1}^{\circ}} \right)$$
are all isomorphic.  $\square$  (Claim 7.4)

Claim 7.5. If  $\iota_C: C \to X^\circ$  denotes the inclusion, then the composed morphisms  $\delta_{\mathscr{F}_{X^\circ}}$  of the following diagrams,

$$(7.5.1) \quad \operatorname{Hom}_{C}\left(\mathscr{F}_{X^{\circ}}|_{C}, \, \mathscr{F}_{X^{\circ}}|_{C} \otimes N_{C/X}^{*}\right) \xrightarrow{\operatorname{restriction}} \operatorname{Hom}_{C}\left(\mathscr{F}_{X^{\circ}}|_{C}, \, \mathscr{F}_{X^{\circ}}|_{C} \otimes \Omega_{X^{\circ}}^{1}|_{C}\right) \xrightarrow{\delta_{\mathscr{F}_{X^{\circ}}}} \operatorname{Hom}_{C}\left(\mathscr{F}_{X^{\circ}}|_{C}, \, \mathscr{F}_{X^{\circ}}|_{C} \otimes \Omega_{X^{\circ}}^{1}|_{C}\right) \xrightarrow{\delta_{\mathscr{F}_{X^{\circ}}}} \operatorname{Hom}_{C}\left(\mathscr{F}_{X^{\circ}}|_{C}, \, \mathscr{F}_{X^{\circ}}|_{C} \otimes \Omega_{C}^{1}\right),$$

are injective, for all  $(\mathscr{F}_{X^{\circ}}, \theta_{\mathscr{F}_{X^{\circ}}})$  in  $F^{\circ}$ .

Proof of Claim 7.5. By exactness of the vertical sequence, it suffices to show that

(7.5.2) 
$$\operatorname{Hom}_{C}(\mathscr{F}_{X^{\circ}}|_{C}, \mathscr{F}_{X^{\circ}}|_{C} \otimes N_{C/X}^{*}) = \{0\}, \text{ for all } \mathscr{F}_{X^{\circ}} \text{ in } F^{\circ}.$$

To this end, observe that the normal bundle of C in X is very positive. In fact, we have  $N_{C/X} \cong \mathcal{O}_X(m \cdot H)^{\oplus n-1}|_C$  by construction. Since  $m \gg 0$  is assumed to be large, it follows from semicontinuity of the Harder-Narasimhan filtration, [HL10, Thm. 2.3.2], that

$$\mu_X^{\max}\big(\big(\mathscr{F}_X^*\otimes\mathscr{F}_X\otimes\mathscr{O}_X(-m\cdot H)\big)^{**}\big)<0,\,\text{for all }\mathscr{F}_X\text{ in F}.$$

In particular, Flenner's version of the Mehta-Ramanathan theorem, [HL10, Thm. 7.1.1], implies that the sheaf

$$\mathscr{H}om_{C}(\mathscr{F}_{X^{\circ}}|_{C},\,\mathscr{F}_{X^{\circ}}|_{C}\otimes N_{C/X}^{*})$$

has no section.

Coming back to the proof of Proposition 7.3, we need to show the implication " $\Leftarrow$ " only. Assume that an element  $(\mathscr{F}_{X^{\circ}}, \theta_{\mathscr{F}_{X^{\circ}}})$  in  $\mathsf{F}^{\circ}$  and an isomorphism of Higgs sheaves,

$$\varphi_C: (\mathscr{F}_{X^\circ}, \theta_{\mathscr{F}_{X^\circ}})|_C \to (\mathscr{E}_{X^\circ}, \theta_{\mathscr{E}_{X^\circ}})|_C$$

are given. In other words, denoting the obvious inclusion map by  $\iota_C:C\to X^\circ$ , we are given an isomorphism of sheaves,  $\varphi_C:\mathscr{F}_{X^\circ}|_C\to\mathscr{E}_{X^\circ}|_C$  and a commutative diagram

$$(7.6.1) \begin{array}{c} \mathscr{F}_{C} \xrightarrow{(\mathrm{Id}_{\mathscr{F}_{C}} \otimes d\iota_{C}) \circ (\theta_{\mathscr{F}_{X^{\circ}}}|_{C})} \to \mathscr{F}_{C} \otimes \Omega^{1}_{C} \\ \varphi_{C} & \downarrow \varphi_{C} \otimes \mathrm{Id}_{\Omega^{1}_{C}} & \text{where} & \mathscr{F}_{C} := \mathscr{F}_{X^{\circ}}|_{C} \\ \mathscr{E}_{C} := \mathscr{E}_{X^{\circ}}|_{C}. \end{array}$$

Item (7.4.2) guarantees that the sheaf morphism  $\varphi_C$  extends in a unique manner to an isomorphism  $\varphi_{X^\circ}: \mathscr{F}_{X^\circ} \to \mathscr{E}_{X^\circ}$ . This way, we obtain two Higgs fields on  $\mathscr{F}_{X^\circ}$ , namely

$$\theta_{\mathscr{F}_{X^{\circ}}} \qquad \text{and} \qquad \theta'_{\mathscr{F}_{X^{\circ}}} \coloneqq (\varphi_{X^{\circ}} \otimes \operatorname{Id}_{\Omega^{1}_{X^{\circ}}})^{-1} \circ \theta_{\mathscr{E}_{X^{\circ}}} \circ \varphi_{X^{\circ}}.$$

Diagram (7.6.1) implies that both Higgs fields  $\theta_{\mathscr{F}_{X^{\circ}}}$  and  $\theta'_{\mathscr{F}_{X^{\circ}}}$  restrict to the same Higgs field on  $\mathscr{F}_{X^{\circ}}|_{C}$ . But then Claim 7.5 guarantees that the two Higgs fields  $\theta_{\mathscr{F}_{X^{\circ}}}$  and  $\theta'_{\mathscr{F}_{X^{\circ}}}$ 

are in fact equal. In other words, the sheaf isomorphism  $\varphi_{X^{\circ}}$  is an isomorphism of Higgs sheaves.

We use the following boundedness result for families of Higgs sheaves. Its proof is very similar to [GKPT15, Claim 8.5 and proof] and therefore omitted. We emphasise that unlike Corollary 7.2, the result formulated here does not depend on Theorem 7.1 and can therefore be used in the proof of that theorem.

**Fact 7.7** (Boundedness of Higgs sheaves). Let X be a klt space, let  $r \in \mathbb{N}^+$  be any number, and let  $\pi: \widetilde{X} \to X$  be a resolution of singularities. Let F be the family of locally free Higgs sheaves  $(\mathscr{F}, \Theta)$  on X, that satisfy rank  $\mathscr{F} = r$  and have the additional property that  $\pi^*(\mathscr{F}, \Theta) \in \text{Higgs}_{\widetilde{X}}$ . Then, the family F is bounded.

7.2. **Preparation for the proof of Theorem 7.1: Vanishing of Chern classes.** The following observation is rather standard, but used several times in this section.

**Lemma 7.8** (Description of bundles with pull-back in  $\operatorname{Higgs}_{\widetilde{X}}$ ). Let X be a projective, klt space and let  $\pi:\widetilde{X}\to X$  be a resolution of singularities. If  $(\mathscr{E},\theta)$  is any locally free Higgs sheaf on X such that  $\pi^*(\mathscr{E},\theta)\in\operatorname{Higgs}_{\widetilde{X}}$ , then all Chern classes  $c_i(\mathscr{E})\in H^{2i}(X,\mathbb{Q})$  vanish and  $(\mathscr{E},\theta)$  is semistable with respect to any nef divisor  $N\in\operatorname{Div}(X)$ .

*Proof.* To prove that  $(\mathscr{E},\theta)$  is semistable with respect to any nef divisor, it is equivalent to show that  $\pi^*(\mathscr{E},\theta)$  is semistable with respect to any nef divisor on  $\widetilde{X}$ , [GKPT15, Prop. 5.20]. Argue by contradiction and assume that  $\pi^*(\mathscr{E},\theta)$  is *not* semistable with respect some nef  $N \in \mathrm{Div}(\widetilde{X})$ . As non-semistability is an open condition, if  $A \in \mathrm{Div}(\widetilde{X})$  is ample and  $\delta \in \mathbb{Q}^+$  sufficiently small, then  $\pi^*(\mathscr{E},\theta)$  fails to be semistable with respect to the ample divisor  $N + \delta \cdot A$ . This contradicts the fact that  $\pi^*(\mathscr{E},\theta)$  is semistable with respect to *any* ample divisor on  $\widetilde{X}$ , cf. [BHR06, Thm. 1.3]. Semistability of  $(\mathscr{E},\theta)$  with respect to any nef divisor follows as desired.

Concerning Chern classes, recall from [Sim92, Rem. on p. 36] that the  $C^{\infty}$ -bundle  $\widetilde{E}$  on  $\widetilde{X}^{an}$  underlying  $\widetilde{\mathscr{E}}$  is induced by a local system. Takayama has shown in [Tak03, Prop. 2.1] that  $X^{an}$  can be covered by contractible, open subsets  $U_i$  with simply connected inverse images  $(\pi^{an})^{-1}(U_i)$ . We infer that the  $C^{\infty}$ -bundle E on  $X^{an}$  underlying  $\mathscr{E} \cong \pi_* \widetilde{\mathscr{E}}$  is again induced by a local system. But then, a classical result of Deligne-Sullivan, [DS75], implies that there exists a finite, étale cover  $\gamma:\widehat{X}\to X$  where  $(\gamma^{an})^*\mathcal{E}$  is trivial. Vanishing of Chern classes downstairs then follows from the Leray spectral sequence and from the splitting of the natural map  $\mathbb{Q}_X\to (\gamma^{an})_*\mathbb{Q}_{\widehat{Y}}$ .

7.3. **Preparation for the proof of Theorem 7.1: Numerical flatness of Higgs sheaves.** The following observation allows to apply the results obtained in Section 5 above to Higgs sheaves on resolutions of klt spaces.

**Proposition 7.9** (Numerical flatness of Higgs sheaves). Let  $\pi: \widetilde{X} \to X$  be a resolution of a klt space, and let E be any local system on  $\widetilde{X}$ . Then, the corresponding locally free Higgs sheaf  $(\mathcal{E}, \theta)$  is  $\pi$ -numerically flat.

*Proof.* Given any smooth, projective curve C and any morphism  $\gamma:C\to\widetilde{X}$  such that  $\pi\circ\gamma$  is constant, we need to show that both  $\gamma^*\mathscr{E}$  and its dual are nef. To this end, recall from [Tak03, p. 827] that every fibre of  $\pi$  admits a small, simply connected neighbourhood U, open in the Euclidean topology. The local system  $\gamma^*(\pi^*\mathsf{E})$  is therefore trivial. As Simpson's nonabelian Hodge correspondence is functorial in morphisms between manifolds, [Sim92, Rem. on p. 36], it follows that the pull-back  $\gamma^*(\mathscr{E},\theta)$  corresponds to the trivial local system, and is therefore trivial itself; in particular, the pullback  $\gamma^*\widetilde{\mathscr{E}}$  and its dual are both nef, as desired.

7.4. **Proof of Theorem 7.1 if** X **is a surface.** The following proposition immediately implies Theorem 7.1 ("Ascent of semistable Higgs bundles") in case where X is a surface.

**Proposition 7.10** (Independence of the polarisation for surfaces). Let X be a smooth projective surface. Given a locally free Higgs sheaf  $(\mathscr{E}, \theta)$  on X, the following statements are equivalent.

- (7.10.1) There exists a big and nef divisor N such that  $ch_1(\mathscr{E}) \cdot N = ch_2(\mathscr{E}) = 0$  and such that  $(\mathscr{E}, \theta)$  is semistable with respect to N.
- (7.10.2) All Chern classes of  $c_i(\mathscr{E}) \in H^{2i}(X, \mathbb{Q})$  vanish and  $(\mathscr{E}, \theta)$  is semistable with respect any ample divisor on X.

*Proof.* Using the fact that big and nef divisors are limits of ample divisors, the direction  $(7.10.2) \Rightarrow (7.10.1)$  is immediate. We will therefore consider the converse direction  $(7.10.1) \Rightarrow (7.10.2)$  for the remainder of the proof. Set  $r := \text{rank } \mathcal{E}$ .

Step 1. Proof in case where  $(\mathscr{E}, \theta)$  is stable with respect to N. Using the assumption on stability with respect to a big and nef class, recall from [Lan02, Thm. 2.1] or [GKPT15, Thm. 6.1] that  $\mathscr{E}$  satisfies the Bogomolov-Gieseker inequality,

$$(7.10.3) 2r \cdot c_2(\mathscr{E}) - (r-1) \cdot c_1(\mathscr{E})^2 \ge 0.$$

Substituting the assumption  $ch_2(\mathscr{E}) = 0$  into (7.10.3), we obtain that  $c_1(\mathscr{E})^2 \geq 0$ . On the other hand, the assumption  $ch_1(\mathscr{E}) \cdot N = 0$ , together with the Hodge index theorem implies that  $c_1(\mathscr{E})^2 \leq 0$ , with equality if and only if  $c_1(\mathscr{E}) = 0$ . We obtain that both  $c_1(\mathscr{E})$  and  $c_2(\mathscr{E})$  vanish, as desired.

Next, let  $H \subset S$  be any ample divisor. By openness of stability, [GKPT15, Prop. 4.17], there exists  $\varepsilon \in \mathbb{Q}^+$  such that  $(\mathscr{E}, \theta)$  is stable with respect to the ample class  $(N + \varepsilon H)$ . Lemma 7.8 then asserts that  $(\mathscr{E}, \theta)$  is stable with respect to any ample divisor. This finishes the proof in the stable case.

Step 2. Proof in general – setup. Since X is smooth, there exists a Jordan-Hölder filtration for the Higgs bundle  $(\mathscr{E},\theta)$  and the nef polarisation N. More precisely, we obtain a filtration

$$0 = \mathscr{E}_0 \subsetneq \mathscr{E}_1 \subsetneq \cdots \subsetneq \mathscr{E}_{t-1} \subsetneq \mathscr{E}_t = \mathscr{E},$$

with the following properties.

- (7.10.4) Each of the sheaves  $\mathcal{E}_i$  is saturated in  $\mathcal{E}_{i+1}$ , hence reflexive, hence locally free since X is a surface.
- (7.10.5) The torsion free quotients  $\mathcal{Q}_i := \mathcal{E}_i/\mathcal{E}_{i-1}$  satisfy  $c_1(\mathcal{Q}_i) \cdot N = 0$ , and inherit Higgs fields  $\tau_i$  making  $(\mathcal{Q}_i, \tau_i)$  stable with respect to N.

Moreover,  $\Omega_X^1$  being locally free, the reflexive hulls  $\mathcal{Q}_i^{**}$ , which are automatically locally free, also inherit Higgs fields, say  $\tau_i^{**}$ , which make  $(\mathcal{Q}_i^{**}, \tau_i^{**})$  stable with respect to N. Since  $\mathcal{Q}_i$  and  $\mathcal{Q}_i^{**}$  agree in codimension one,  $c_1(\mathcal{Q}_i) = c_1(\mathcal{Q}_i^{**})$ , and  $c_1(\mathcal{Q}_i^{**}) \cdot N = 0$ .

Step 3. Proof in general – the Chern character of  $\mathcal{Q}_i^{**}$ . We will prove in this step that

$$(7.10.6) ch_2(\mathcal{Q}_i^{**}) = ch_2(\mathcal{Q}_i) = 0 \text{for all } i,$$

following [Sim92, proof of Thm. 2] closely. In fact, since  $c_2(\mathcal{Q}_i^{**}) \leq c_2(\mathcal{Q}_i)$ , cf. [HL10, p. 80], we have the inverse inequality for the second Chern character,  $ch_2(\mathcal{Q}_i) \leq ch_2(\mathcal{Q}_i^{**})$ . Additivity of Chern characters then implies that

$$(7.10.7) 0 = ch_2(\mathscr{E}) = \sum ch_2(\mathscr{Q}_i) \le \sum ch_2(\mathscr{Q}_i^{**}).$$

On the other hand, the Bogomolov-Gieseker inequality, [Lan02, Thm. 2.1] or [GKPT15, Thm. 6.1], give that

$$(7.10.8) 2 \cdot ch_2(\mathcal{Q}_i^{**}) \le \frac{1}{\operatorname{rank} \mathcal{Q}_i} \cdot c_1(\mathcal{Q}_i^{**})^2.$$

As before, equality  $c_1(\mathcal{Q}_i^{**}) \cdot N = 0$  together with Hodge index theorem implies that  $c_1(\mathcal{Q}_i^{**})^2 \leq 0$ , so Inequality (7.10.8) reduces to  $ch_2(\mathcal{Q}_i^{**}) \leq 0$ . Hence (7.10.7) implies (7.10.6).

Step 4. Proof in general – end of proof. Equation (7.10.6) allows us to apply the results of Step 1 to the Higgs bundles  $(\mathcal{Q}_i^{**}, \tau_i^{**})$ . This implies in particular that all Chern classes of  $\mathcal{Q}_i^{**}$  vanish, so that the sheaves  $\mathcal{Q}_i$  and  $\mathcal{Q}_i^{**}$  agree to start with. The Higgs sheaf  $(\mathcal{E}, \theta)$  is thus (an iterated) extension of Higgs bundles with vanishing Chern classes that are stable with respect to any ample polarisation. Item (7.10.2) follows, which ends the proof of Proposition 7.10 and therefore the proof of Theorem 7.1 in dimension 2.

7.5. **Proof of Theorem 7.1 if** X **is maximally quasi-étale.** We will now prove Theorem 7.1 under the additional assumption that the natural push-forward morphism of algebraic fundamental groups,  $\widehat{\pi}_1(X_{\text{reg}}) \to \widehat{\pi}_1(X)$ , is isomorphic. Following [GKP16], we say shortly that X is maximally quasi-étale. This assumption will be maintained throughout the present Section 7.5. Choose a divisor  $\Delta$  such that  $(X, \Delta)$  is klt.

Notation 7.11. Choose an ample divisor  $H \in \text{Div}(X)$  such that  $(\mathscr{E}, \theta)$  is semistable with respect to H and satisfies  $ch_1(\mathscr{E}) \cdot [H]^{n-1} = ch_2(\mathscr{E}) \cdot [H]^{n-2} = 0$ . Set  $(\widetilde{\mathscr{E}}, \widetilde{\theta}) := \pi^*(\mathscr{E}, \theta)$  and  $\widetilde{H} := \pi^*H$ . If A is a subvariety of B, we denote the obvious inclusion morphism by  $\iota_A$ .

Finally, let F be the family of locally free Higgs sheaves  $(\mathscr{F},\Theta)$  on X, that satisfy rank  $\mathscr{F}=\operatorname{rank}\mathscr{E}$  and have the additional property that  $\pi^*(\mathscr{F},\Theta)\in\operatorname{Higgs}_{\widetilde{X}}$ . Recall from Fact 7.7 that this family is bounded.

Step 1: Choice of a complete intersection surface. Our proof relies on the choice of a sufficiently general complete intersection surface  $S \subset X$  to which we can restrict. To be precise, choosing a sufficiently increasing sequence of numbers  $0 \ll m_1 \ll m_2 \ll \cdots \ll m_{n-2}$  as well as a sufficiently general tuple of hyperplanes,

$$(D_1,\ldots,D_{n-2})\in |m_1\cdot H|\times\cdots\times |m_{n-2}\cdot H|,$$

the following will hold.


- (7.12.1) The intersection  $S := D_1 \cap \cdots \cap D_{n-2}$  is an irreducible, normal surface, not contained in any component of  $\Delta$ , and the pair  $(S, \Delta|_S)$  is klt —Seidenberg's theorem [BS95, Thm. 1.7.1] and [KM98, Lem. 5.17].
- (7.12.2) The restriction  $(\mathcal{E}, \theta)|_S$  is semistable with respect to the ample divisor  $H|_S$  Restriction theorem, Theorem 6.1.
- (7.12.3) The natural morphism  $i_*: \pi_1(S_{\text{reg}}) \to \pi_1(X_{\text{reg}})$ , induced by the inclusion  $i: S_{\text{reg}} \to X_{\text{reg}}$ , is isomorphic —Lefschetz hyperplane theorem for fundamental groups, [GM88, Thm. in Sect. II.1.2].
- (7.12.4) Given any Higgs sheaf  $(\mathscr{F}, \Theta) \in F$ , then  $(\mathscr{E}, \theta) \cong (\mathscr{F}, \Theta)$  if and only if  $(\mathscr{E}, \theta)|_S \cong (\mathscr{F}, \Theta)|_S$  —Iterated Bertini theorem, Proposition 7.3.

Let  $\widetilde{S} \subset \widetilde{X}$  be the strict transform of S in  $\widetilde{X}$ . Then  $\widetilde{S}$  is a smooth surface and  $\pi_S : \widetilde{S} \to S$  is a resolution. The following diagram summarises the situation.

(7.12.5) 
$$\widetilde{S}^{C} \xrightarrow{\iota_{\widetilde{S}}} \widetilde{X} \\
\pi_{S} \downarrow \qquad \qquad \downarrow_{\pi} \\
S \xrightarrow{\iota_{S}} X.$$

Claim 7.13. The natural morphism of étale fundamental groups,  $\widehat{\pi}_1(\widetilde{S}) \to \widehat{\pi}_1(\widetilde{X})$  is isomorphic. In particular every local system on  $\widetilde{S}$  is the restriction of a local system on  $\widetilde{X}$ .

*Proof.* Consider the following diagram of push-forward morphisms of étale fundamental groups,



The group morphisms  $\alpha_S$  and  $\alpha_X$  are isomorphisms by Takayama, [Tak03]. The morphism  $\beta_S$  is surjective by e.g. [FL81, 0.7.B on p. 33], and by the fact the profinite completion is right exact, [RZ10, Lem. 3.2.3 and Prop. 3.2.5]. The morphism  $\beta_X$  is isomorphic by assumption, and  $(\iota_{S_{\text{reg}}})_*$  is isomorphic by Item (7.12.3). The remaining arrows  $(\iota_{\widetilde{S}})_*$  and  $(\iota_S)_*$  must then also be isomorphic. The asserted extension of local systems comes from the fact that representation of  $\pi_1(S)$  comes from a representation of  $\pi_1(X)$ , cf. [Gro70, Thm. 1.2b], or [GKP16, Sect. 8.1] for a detailed pedestrian proof


Step 2: End of proof. We consider the restriction of  $(\widetilde{\mathscr{E}}, \widetilde{\theta})$  to  $\widetilde{S}$ . Commutativity of Diagram (7.12.5) and the fact that all spaces involved are klt allow us to apply [GKPT15, Lem. 5.9] to conclude that

(7.13.1) 
$$(\widetilde{\mathscr{E}}, \widetilde{\theta})|_{\widetilde{S}} \cong \pi_{S}^{*}((\mathscr{E}, \theta)|_{S}),$$

which is hence semistable with respect to  $\pi_S^*(H|_S)$ , thanks to Item (7.12.2) above and [GKPT15, Prop. 5.19]. Proposition 7.10 therefore implies that  $(\widetilde{\mathscr{E}},\widetilde{\theta})|_{\widetilde{S}}\in \operatorname{Higgs}_{\widetilde{S}}$ . The classical nonabelian Hodge correspondence applies, and gives a local system  $\mathsf{E}_{\widetilde{S}}\in \mathsf{LSys}_{\widetilde{S}}$ . According to Claim 7.13, we find a local system  $\mathsf{E}_{\widetilde{X}}\in \mathsf{LSys}_{\widetilde{X}}$  whose restriction  $\mathsf{E}_{\widetilde{X}}|_{\widetilde{S}}$  is isomorphic to  $\mathsf{E}_{\widetilde{S}}$ . The classical nonabelian Hodge correspondence on  $\widetilde{X}$  thus yields a Higgs sheaf  $(\widetilde{\mathscr{F}},\widetilde{\Theta})\in \mathsf{Higgs}_{\widetilde{X}}$  with vanishing Chern classes, whose restriction  $(\widetilde{\mathscr{F}},\widetilde{\Theta})|_{\widetilde{S}}$  is isomorphic to  $(\widetilde{\mathscr{E}},\widetilde{\theta})|_{\widetilde{S}}$  by functoriality. It follows from Proposition 7.9 that  $\widetilde{\mathscr{F}}$  is  $\pi$ -numerically flat. Item (5.1.2) of Theorem 5.1 therefore yields a locally free Higgs sheaf  $(\mathscr{F},\Theta)\in \mathsf{F}$  whose restriction  $(\mathscr{F},\Theta)|_S$  is isomorphic to  $(\mathscr{E},\theta)|_S$  owing to (7.13.1). In particular, Item (7.12.4) applies to show that  $(\mathscr{E},\theta)\cong (\mathscr{F},\Theta)$ . By construction,  $\pi^*(\mathscr{E},\theta)\cong (\widetilde{\mathscr{F}},\widetilde{\Theta})\in \mathsf{Higgs}_{\widetilde{X}}$ , which concludes the proof of Theorem 7.1 in case where X is maximally quasi-étale.

7.6. **Proof of Theorem 7.1 in the general setting.** Finally, we prove Theorem 7.1 ("Ascent of semistable Higgs bundles") without any additional assumptions. Choose an ample divisor  $H \in \text{Div}(X)$  such that  $(\mathscr{E}, \theta)$  is semistable with respect to H and satisfies  $ch_1(\mathscr{E}) \cdot [H]^{n-1} = ch_2(\mathscr{E}) \cdot [H]^{n-2} = 0$  and write  $(\widetilde{\mathscr{E}}, \widetilde{\theta}) := \pi^*(\mathscr{E}, \theta)$ .

Recall from [GKP16, Thm. 1.5] that there exists a quasi-étale cover  $f: Y \to X$  such that  $\widehat{\pi}_1(Y_{\text{reg}}) \cong \widehat{\pi}_1(Y)$ . Choose one such f, note that the corresponding Y is again klt, and let  $\widetilde{Y}$  be a desingularisation of the (unique) irreducible component of the fibre product  $Y \times_X \widetilde{X}$  that dominates both Y and  $\widetilde{X}$ . We obtain a diagram of surjections as follows,



Recall from Corollary 6.4 that the Higgs sheaf  $(\mathscr{E}_Y, \theta_Y) := f^*(\mathscr{E}, \theta)$  is semistable with respect to the ample divisor  $H_Y := f^*H$ . It clearly satisfies  $ch_1(\mathscr{E}_Y) \cdot [H_Y]^{n-1} = ch_2(\mathscr{E}_Y) \cdot [H_Y]^{n-2} = 0$ . In particular,  $(\mathscr{E}_Y, \theta_Y) \in \text{Higgs}_Y$ . We have seen in Section 7.5 that this implies that  $\widetilde{\pi}^*(\mathscr{E}_Y, \theta_Y) \in \text{Higgs}_{\widetilde{Y}}$ . Lemma 7.8 thus yields the following.

(7.13.2) All Chern classes  $c_i(\widetilde{\pi}^* \mathscr{E}_Y) = c_i(\widetilde{f}^* \widetilde{\mathscr{E}}) \in H^{2i}(\widetilde{Y}, \mathbb{Q})$  vanish.

(7.13.3) The Higgs sheaf  $\widetilde{\pi}^*(\mathscr{E}_Y, \theta_Y)$  is semistable with respect to *any* nef divisor on  $\widetilde{Y}$ .

Let  $\widetilde{H} \in \text{Div}(\widetilde{X})$  be any ample. Item (7.13.2) immediately implies that

$$ch_1(\widetilde{\mathcal{E}}) \cdot [\widetilde{H}]^{n-1} = ch_2(\widetilde{\mathcal{E}}) \cdot [\widetilde{H}]^{n-2} = 0.$$

In a similar vein, Item (7.13.3) and Corollary 6.4 imply that  $(\widetilde{\mathscr{E}}, \widetilde{\theta})$  is semistable with respect to  $\widetilde{H}$ . This concludes the proof of Theorem 7.1.

#### 8. Proof of the nonabelian Hodge correspondence for klt spaces

8.1. Proof of Theorem 3.4 ("Nonabelian Hodge correspondence for klt spaces"). Using the results on descent and ascent for Higgs bundles, Theorems 5.1 and 7.1, we construct the relevant functors between  $\operatorname{Higgs}_X$  and  $\operatorname{LSys}_X$ . Once the functors are constructed, we show that they indeed satisfy all the claims made in Theorem 3.4. We maintain assumptions and notation of Theorem 3.4 throughout Section 8.1. The canonical resolution of singularities is denoted by  $\pi:\widetilde{X}\to X$ .

Step 1: from local systems to bundles. We establish the first half of Theorem 3.4, constructing a functor that maps local systems to Higgs bundles. Given a local system  $\mathsf{E} \in \mathsf{LSys}_X$ , consider the Higgs bundle  $(\widetilde{\mathscr{E}},\widetilde{\theta}) := \eta_{\widetilde{X}}(\pi^*\mathsf{E})$  associated to  $\pi^*\mathsf{E}$  via the nonabelian Hodge correspondence on the manifold  $\widetilde{X}$ . Recall that  $(\widetilde{\mathscr{E}},\widetilde{\theta}) \in \mathsf{Higgs}_{\widetilde{X}}$  and set  $\mathscr{E} := \pi_*\widetilde{\mathscr{E}}$ . Proposition 7.9 and Item (5.1.2) of Theorem 5.1 imply that  $\mathscr{E}$  is locally free and that it carries a unique Higgs field  $\theta$  such that  $(\widetilde{\mathscr{E}},\widetilde{\theta}) \cong \pi^*(\mathscr{E},\theta)$ . Lemma 7.8 applies to show that  $(\mathscr{E},\theta) \in \mathsf{Higgs}_X$ . In summary, we have constructed a mapping  $\eta_X : \mathsf{LSys}_X \to \mathsf{Higgs}_X$ .

If  $e: E_1 \to E_2$  is a morphism of local systems, we obtain a morphism  $\widetilde{e}: \pi^*(E_1) \to \pi^*(E_2)$  and hence a morphism between the associated Higgs sheaves,  $\eta_{\widetilde{X}}(\widetilde{e}): (\widetilde{\mathcal{E}}_1, \widetilde{\theta}_1) \to (\widetilde{\mathcal{E}}_2, \widetilde{\theta}_2)$ . Denoting the associated, locally free Higgs sheaves on X by  $(\mathcal{E}_1, \theta_1)$  and  $(\mathcal{E}_2, \theta_2)$ , an elementary computation shows that  $\eta_{\widetilde{X}}(\widetilde{e})$  descends to a morphism  $\eta_X(e): (\mathcal{E}_1, \theta_1) \to (\mathcal{E}_2, \theta_2)$ . In other words, we constructed a functor  $\eta_X: \mathsf{LSys}_X \to \mathsf{Higgs}_X$ .

*Observation* 8.1. If *X* is smooth, then the canonical resolution of singularities is the identity, and  $\eta_X$  equals the functor given by the nonabelian Hodge correspondence.

*Observation* 8.2. The natural map  $\pi^*\mathscr{E} = \pi^*\pi_*\widetilde{\mathscr{E}} \to \widetilde{\mathscr{E}}$  induces an isomorphism of Higgs sheaves,  $N_{\pi,\mathsf{E}} : \pi^*\eta_X(\mathsf{E}) \to \eta_{\widetilde{X}}(\pi^*\mathsf{E})$ .

Step 2: from bundles to local systems. Given a Higgs bundle  $(\mathscr{E},\theta) \in \operatorname{Higgs}_X$ , consider the Higgs bundle  $(\widetilde{\mathscr{E}},\widetilde{\theta}) := \pi^*(\mathscr{E},\theta)$ . Theorem 7.1 asserts that  $(\widetilde{\mathscr{E}},\widetilde{\theta})$  is in  $\operatorname{Higgs}_{\widetilde{X}}$ . Apply the classical nonabelian Hodge correspondence on the manifold  $\widetilde{X}$ , in order to obtain a local system  $\operatorname{E}_{\widetilde{X}} := \mu_{\widetilde{X}}(\widetilde{\mathscr{E}},\widetilde{\theta})$ . As before, Takayama's result [Tak03, p. 827] shows that  $\operatorname{E} := \pi_*\operatorname{E}_{\widetilde{X}}$  is a local system on X. We leave it to the reader to show that this construction does indeed give a functor  $\mu_X : \operatorname{Higgs}_X \to \operatorname{LSys}_X$  and that the following observations hold.

*Observation* 8.3. If X is smooth, then the canonical resolution of singularities is the identity, and  $\mu_X$  equals the functor given by the classical nonabelian Hodge correspondence.

Observation 8.4. The natural map  $\pi^*\mathsf{E} = \pi^*\pi_*\,\mathsf{E}_{\widetilde{X}} \to \mathsf{E}_{\widetilde{X}}$  induces an isomorphism of local systems,  $M_{\pi_*(\mathscr{E},\theta)}: \pi^*\mu_X(\mathsf{E}) \to \mu_{\widetilde{X}}(\pi^*\mathsf{E})$ .

Step 3: equivalence of categories. Let X be a projective klt space, and let  $\pi:\widetilde{X}\to X$  be the canonical resolution. The functors  $\eta_{\widetilde{X}}$  and  $\mu_{\widetilde{X}}$  associated with the classical nonabelian Hodge correspondence on the manifold  $\widetilde{X}$  form an equivalence of categories: the compositions  $\eta_{\widetilde{X}}\circ \mu_{\widetilde{X}}$  and  $\mu_{\widetilde{X}}\circ \eta_{\widetilde{X}}$  is naturally isomorphic to the identities on Higgs $_{\widetilde{X}}$  and LSys $_{\widetilde{X}}$ , respectively. One checks immediately that these isomorphisms descend to X, showing that the functors  $\eta_X$  and  $\mu_X$  constructed above do indeed give an equivalence of categories.

Step 4: end of proof. It remains to show Items (3.4.1)–(3.4.3) of Theorem 3.4. These, however, follow immediately from Observations 8.1–8.4. The proof of Theorem 3.4 is thus complete.

8.2. **Proof of Theorem 3.5** ("Functoriality in morphisms"). To keep the paper reasonably short, we will only consider functoriality of the functors  $\eta_{\bullet}$ . Functoriality of  $\mu_{\bullet}$  follows along the same lines of argument. We construct the isomorphisms  $N_{\bullet,\bullet}$  in some detail, but leave the tedious and lengthy verification of the construction's properties to the reader, as none of the required arguments is in any way surprising or holds a promise of new insight.

Step 1: lifting morphisms. For morphism between manifolds, functoriality of the nonabelian Hodge correspondence is classically known, [Sim92, Rem. on p. 36]. If  $f: X \to Y$  is a morphism between klt spaces that are not smooth, the following claim allows to lift f to a morphism between spaces that are smooth, though possibly of higher dimension.

*Claim* 8.5. Given a morphism  $f: X \to Y$  of projective klt spaces, there exists a smooth, projective variety  $\widetilde{W}$ , and a commutative diagram as follows,

(8.5.1) 
$$\widetilde{W} \xrightarrow{\widetilde{f}} W \subseteq \widetilde{X}$$

$$\downarrow \pi_{W}, \text{ conn. fibres} \downarrow \pi, \text{ canon. resolution}$$

$$\widetilde{Y} \xrightarrow{\pi_{Y}, \text{ canon. resolution}} Y \xrightarrow{f} Z \subseteq X,$$

where Z = f(Y) and  $W = \widetilde{f}(\widetilde{W})$ .

*Proof.* Recall from [HM07, Cor. 1.7(1)] that there exists an irreducible variety  $W \subseteq \pi^{-1}(Z) \subseteq \widetilde{X}$  such that the induced morphism  $\pi_W : W \to Z$  is surjective, and such that its general fibre is rationally connected, and in particular irreducible. Choose one such W. Consider the fibre product  $W \times_Z \widetilde{Y}$ , choose the unique component that dominates both W and Y, and let  $\widetilde{W}$  its canonical resolution. The general fibre of  $\Pi$  is again rationally connected and irreducible, in particular connected.

Step 2: Construction of  $N_{\bullet,\bullet}$ . With the help of Claim 8.5, we construct the isomorphisms  $N_{\bullet,\bullet}$  in this section.

Construction 8.6. Assume we are given a morphism  $f: X \to Y$  of projective klt spaces and a local system  $E \in \mathsf{LSys}_X$ . Choose a diagram (D) as in Claim 8.5 and construct an isomorphism

$$\alpha_{\mathsf{D}}: \Pi^* \pi_Y^* f^* \eta_X(\mathsf{E}) \to \Pi^* \pi_Y^* \eta_Y (f^* \mathsf{E})$$

as the composition of the following canonical isomorphisms,

The following canonical isomorphisms, 
$$\Pi^* \pi_Y^* f^* \eta_X(\mathsf{E}) \cong \widetilde{f}^* \pi^* \eta_X(\mathsf{E}) \qquad \text{Commutativity of (D)}$$

$$\cong \widetilde{f}^* \eta_{\widetilde{X}}(\pi^* \mathsf{E}) \qquad \text{Item (3.4.3) of Theorem 3.4}$$

$$\cong \eta_{\widetilde{W}}(\widetilde{f}^* \pi^* \mathsf{E}) \qquad \text{Simpson's functoriality for } \widetilde{f}: \widetilde{W} \to \widetilde{X}$$

$$\cong \eta_{\widetilde{W}}(\Pi^* \pi_Y^* f^* \mathsf{E}) \qquad \text{Commutativity of (8.5.1)}$$

$$\cong \Pi^* \eta_{\widetilde{Y}} (\pi_Y^* f^* \mathsf{E}) \qquad \text{Simpson's functoriality for } \Pi : \widetilde{W} \to \widetilde{Y} \\ \cong \Pi^* \pi_Y^* \eta_Y (f^* \mathsf{E}) \qquad \text{Item (3.4.3) of Theorem 3.4.}$$

Since  $\pi_Y \circ \Pi$  has connected fibres, the isomorphism  $\alpha_D$  descends to an isomorphism

$$\beta_{\mathsf{D}}: f^*\eta_X(\mathsf{E}) \to \eta_Y(f^*\mathsf{E}).$$

Claim 8.7. In the setting of Construction 8.6, the morphism  $\beta_D$  is independent of the choices made. More precisely, given two diagrams  $D_1$  and  $D_2$  as in Claim 8.5, then  $\beta_{D_1} = \beta_{D_2}$ . We can therefore choose D arbitrarily and set  $N_{f,E} := \beta_D$ .


Step 3: end of proof. The isomorphisms  $N_{\bullet,\bullet}$  constructed in Step 2 clearly have the expected behaviour under canonical resolution and satisfy the compatibility conditions spelled out in Theorem 3.5. We leave it to the reader to verify functoriality and to write down the analogous construction of the morphisms  $M_{\bullet,\bullet}$ .

8.3. **Proof of Theorem 3.9 ("Independence of polarisation").** We establish the following sequence of implications.

$$(3.9.5) \xrightarrow[\text{trivial}]{} (3.9.4) \Longrightarrow (3.9.3) \xrightarrow[\text{trivial}]{} (3.9.2) \xrightarrow[\text{trivial}]{} (3.9.1)$$

Implication (3.9.4)  $\Rightarrow$  (3.9.3) in the semistable case. This is given by Lemma 7.8 ("Description of bundles whose pull-back is in Higgs $_{\widetilde{X}}$ ").

Implication  $(3.9.4) \Rightarrow (3.9.3)$  in the stable case. Choose  $\pi: \widetilde{X} \to X$  and  $\widetilde{H} \in \operatorname{Div}(\widetilde{X})$  as in (3.9.4). Vanishing of Chern classes has been shown in the semistable case. As for stability, let  $H \in \operatorname{Div}(X)$  be any ample divisor. Choose a sufficiently large number  $m \gg 0$  such that  $|m \cdot H|$  is basepoint free, choose sufficiently general hyperplanes  $H_1, \ldots, H_{\dim X - 1} \in |m \cdot H|$  and consider the associated complete intersection curve  $C := H_1 \cap \cdots \cap H_{\dim X - 1} \subset X$ . Observe that C is entirely contained in  $X_{\text{reg}}$ , and that  $\pi$  is isomorphic near C. Denote the preimage by  $\widetilde{C} := \pi^{-1}C$ . We obtain the following obvious morphisms between fundamental groups,



In particular, we see that the morphism  $\widetilde{i}_*$  is surjective. Apply the nonabelian Hodge correspondence for klt spaces, Theorem 3.4 to obtain local systems E on X and  $\pi^*$  E  $\cong \mu_{\widetilde{X}}(\pi^*(\mathscr{E},\theta))$  on  $\widetilde{X}$ . The following will then end the proof.

$$\pi^*\left(\mathscr{E},\theta\right) \text{ is stable w.r.t. } \widetilde{H} \Rightarrow \pi^* \text{ E is irreducible} \qquad \begin{array}{ll} [\operatorname{Sim92}, \operatorname{p. 18ff}] \\ \Rightarrow (\pi^*\operatorname{E})|_{\widetilde{C}} \text{ is irreducible} & \operatorname{surjectivity of } \widetilde{i_*} \\ \Rightarrow \operatorname{E}|_{C} \text{ is irreducible} & \operatorname{isomorphic} \\ \Rightarrow (\mathscr{E},\theta)|_{C} \text{ is stable} & [\operatorname{Sim92}, \operatorname{p. 18ff}] \\ \Rightarrow (\mathscr{E},\theta) \text{ is stable w.r.t. } H & \operatorname{genl. choice} \end{array}$$

Implication  $(3.9.4) \Rightarrow (3.9.3)$  in the polystable case. Analogous to the above, with "irreducible" replaced by "semisimple".

Implication  $(3.9.1) \Rightarrow (3.9.5)$  in the semistable case. This is Theorem 7.1 ("Ascent of semistable Higgs bundles") together with [BHR06, Thm. 1.3].

Implication  $(3.9.1) \Rightarrow (3.9.5)$  in the stable case. The pull-back bundle  $\pi^*(\mathscr{E}, \theta)$  is clearly stable with respect to the nef bundle  $\pi^*H$ , [GKPT15, Prop. 5.19]. But then openness of stability, [GKPT15, Prop. 4.17] asserts that  $\pi^*(\mathscr{E}, \theta)$  is stable with respect to one ample, hence any ample bundle.

*Implication*  $(3.9.1) \Rightarrow (3.9.5)$  *in the polystable case.* Same as the stable case.

#### REFERENCES

- [AD14] Carolina Araujo and Stéphane Druel. On codimension 1 del Pezzo foliations on varieties with mild singularities. *Math. Ann.*, 360(3-4):769−798, 2014. DOI:10.1007/s00208-014-1053-3. Preprint arXiv:1210.4013. ↑13
- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James McKernan. Existence of minimal models for varieties of log general type. J. Amer. Math. Soc., 23:405–468, 2010. DOI:10.1090/S0894-0347-09-00649-3. ↑ 15
- [BHR06] Ugo Bruzzo and Daniel Hernández Ruipérez. Semistability vs. nefness for (Higgs) vector bundles. Differential Geom. Appl., 24(4):403–416, 2006. DOI:10.1016/j.difgeo.2005.12.007. ↑ 22, 28
- [BM97] Edward Bierstone and Pierre D. Milman. Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant. *Invent. Math.*, 128(2):207–302, 1997. DOI:10.1007/s002220050141. ↑ 9
- [BM14] Daniel Barlet and Jón Magnússon. Cycles analytiques complexes. I. Théorèmes de préparation des cycles, volume 22 of Cours Spécialisés [Specialized Courses]. Société Mathématique de France, Paris, 2014. ↑ 4
- [BS95] Mauro C. Beltrametti and Andrew J. Sommese. The adjunction theory of complex projective varieties, volume 16 of De Gruyter Expositions in Mathematics. Walter de Gruyter & Co., Berlin, 1995. DOI:10.1515/9783110871746. ↑ 24
- [CKT16] Benoît Claudon, Stefan Kebekus, and Behrouz Taji. Generic positivity and applications to hyper-bolicity of moduli spaces. Preprint arXiv:1610.09832. To appear as a chapter in a forthcoming book on hyperbolicity, October 2016. ↑ 4
- [Del70] Pierre Deligne. Équations différentielles à points singuliers réguliers. Springer-Verlag, Berlin, 1970. Lecture Notes in Mathematics, Vol. 163. DOI:10.1007/BFb0061194. ↑ 9
- [DPS94] Jean-Pierre Demailly, Thomas Peternell, and Michael Schneider. Compact complex manifolds with numerically effective tangent bundles. J. Algebraic Geom., 3(2):295−345, 1994. ↑ 5
- [DS75] Pierre Deligne and Dennis Sullivan. Fibrés vectoriels complexes à groupe structural discret. C. R. Acad. Sci. Paris Sér. A-B, 281(24):Ai, A1081−A1083, 1975. ↑ 22
- [FL81] William Fulton and Robert Lazarsfeld. Connectivity and its applications in algebraic geometry. In *Algebraic geometry (Chicago, Ill., 1980)*, volume 862 of *Lecture Notes in Math.*, pages 26−92. Springer, Berlin, 1981. DOI:10.1007/BFb0090889. ↑ 25, 28
- [Ful98] William Fulton. Intersection Theory, volume 2 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, second edition, 1998. DOI:10.1007/978-1-4612-1700-8. ↑ 13
- [GK14] Patrick Graf and Sándor J. Kovács. An optimal extension theorem for 1-forms and the Lipman-Zariski conjecture. *Doc. Math.*, 19:815–830, 2014. Preprint arXiv:1301.7315. ↑ 2
- [GKKP11] Daniel Greb, Stefan Kebekus, Sándor J. Kovács, and Thomas Peternell. Differential forms on log canonical spaces. Inst. Hautes Études Sci. Publ. Math., 114(1):87–169, November 2011. DOI:10.1007/s10240-011-0036-0 An extended version with additional graphics is available as arXiv:1003.2913. ↑ 15
- [GKP16] Daniel Greb, Stefan Kebekus, and Thomas Peternell. Étale fundamental groups of Kawamata log terminal spaces, flat sheaves, and quotients of abelian varieties. *Duke Math. J.*, 165(10):1965–2004, 2016. DOI:10.1215/00127094-3450859. Preprint arXiv:1307.5718. ↑ 4, 20, 24, 25
- [GKPT15] Daniel Greb, Stefan Kebekus, Thomas Peternell, and Behrouz Taji. The Miyaoka-Yau inequality and uniformisation of canonical models. Preprint arXiv:1511.08822. To appear in the Annales scientifiques de l'École normale supérieure, November 2015. ↑ 1, 2, 3, 7, 9, 10, 16, 17, 18, 22, 23, 25, 29
- [GKPT18] Daniel Greb, Stefan Kebekus, Thomas Peternell, and Behrouz Taji. Harmonic metrics on higgs sheaves and uniformisation of varieties of general type. Preprint arXiv:1804.01266, April 2018. ↑ 2, 3
- [GKT18] Daniel Greb, Stefan Kebekus, and Behrouz Taji. Uniformisation of higher-dimensional varieties. In Tommaso de Fernex, Brendan Hassett, Mircea Mustaţă, Martin Olsson, Mihnea Popa, and Richard Thomas, editors, Algebraic Geometry: Salt Lake City 2015, volume 1 of Proceedings of Symposia in Pure Mathematics, pages 277−308. American Mathematical Society, American Mathematical Society, 2018. Preprint arXiv:1608.06644. ↑ 1, 2

- [GM88] Mark Goresky and Robert D. MacPherson. Stratified Morse theory, volume 14 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1988. DOI:10.1007/978-3-642-71714-7. ↑ 24, 28
- [Gro66] Alexandre Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. Troisiéme partie. *Inst. Hautes Études Sci. Publ. Math.*, (28):255, 1966. Revised in collaboration with Jean Dieudonné. numdam.PMIHES-1966-28-5-0. ↑ 18
- [Gro68] Alexander Grothendieck. Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2). North-Holland Publishing Co., Amsterdam, 1968. Augmenté d'un exposé par Michèle Raynaud, Séminaire de Géométrie Algébrique du Bois-Marie, 1962, Advanced Studies in Pure Mathematics, Vol. 2. Also available as arXiv:math/0511279. ↑ 20
- [Gro70] Alexander Grothendieck. Représentations linéaires et compactification profinie des groupes discrets. *Manuscripta Math.*, 2:375−396, 1970. DOI:10.1007/BF01719593. ↑ 25
- [GT16] Henri Guenancia and Behrouz Taji. Orbifold stability and Miyaoka-Yau inequality for minimal pairs. Preprint arXiv:1611.05981, November 2016. ↑ 3
- [GW10] Ulrich Görtz and Torsten Wedhorn. Algebraic geometry I, Schemes, With Examples and Exercises. Advanced Lectures in Mathematics. Vieweg + Teubner, Wiesbaden, 2010. DOI:10.1007/978-3-8348-9722-0. ↑ 6
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52. DOI:10.1007/978-1-4757-3849-0. ↑ 3, 14
- [Har80] Robin Hartshorne. Stable reflexive sheaves. *Math. Ann.*, 254(2):121−176, 1980 DOI: 10.1007/BF01467074. ↑ 20
- [HL10] Daniel Huybrechts and Manfred Lehn. The geometry of moduli spaces of sheaves. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2010. DOI:10.1017/CBO9780511711985. ↑ 21, 23
- [HM07] Christopher D. Hacon and James McKernan. On Shokurov's rational connectedness conjecture. Duke Math. J., 138(1):119–136, 2007. euclid.dmj/1178738561. Preprint arXiv:math/0504330. ↑ 27
- [HN13] Andreas Höring and Carla Novelli. Mori contractions of maximal length. Publ. Res. Inst. Math. Sci., 49(1):215–228, 2013. DOI:10.4171/PRIMS/103. Preprint arXiv:1201.4009. ↑ 14
- [Keb13] Stefan Kebekus. Pull-back morphisms for reflexive differential forms. Adv. Math., 245:78–112, 2013. DOI:10.1016/j.aim.2013.06.013. Preprint arXiv:1210.3255. ↑ 2, 8
- [KM98] János Kollár and Shigefumi Mori. Birational geometry of algebraic varieties, volume 134 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. DOI:10.1017/CBO9780511662560.
  † 3. 5. 6. 7. 24
- [KMM87] Yujiro Kawamata, Katsumi Matsuda, and Kenji Matsuki. Introduction to the minimal model problem. In Algebraic geometry, Sendai, 1985, volume 10 of Adv. Stud. Pure Math., pages 283–360. North-Holland, Amsterdam, 1987. ↑ 6, 7
- [Kol96] János Kollár. Rational curves on algebraic varieties, volume 32 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, 1996. DOI:10.1007/978-3-662-03276-3. ↑ 4, 5
- [Kol11] János Kollár. Simultaneous normalization and algebra husks. Asian J. Math., 15(3):437–449, 2011. DOI:10.4310/AJM.2011.v15.n3.a6. Preprint arXiv:0910.1076. ↑ 14
- [Lan02] Adrian Langer. A note on Bogomolov's instability and Higgs sheaves. In Algebraic geometry, pages 237–256. de Gruyter, Berlin, 2002.  $\uparrow$  23
- [Lan15] Adrian Langer. Bogomolov's inequality for Higgs sheaves in positive characteristic. Invent. Math., 199(3):889–920, 2015. DOI:10.1007/s00222-014-0534-z.  $\uparrow$  18
- [Laz04] Robert Lazarsfeld. Positivity in algebraic geometry. I. Classical setting: line bundles and linear series, volume 48 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. Springer-Verlag, Berlin, 2004. DOI:10.1007/978-3-642-18808-4. ↑ 6
- [Moc06] Takuro Mochizuki. Kobayashi-Hitchin correspondence for tame harmonic bundles and an application. *Astérisque*, (309):viii+117, 2006. ↑ 3
- [Moc07a] Takuro Mochizuki. Asymptotic behaviour of tame harmonic bundles and an application to pure twistor *D*-modules. I. *Mem. Amer. Math. Soc.*, 185(869):xii+324, 2007. DOI:10.1090/memo/0869. ↑ 3
- [Moc07b] Takuro Mochizuki. Asymptotic behaviour of tame harmonic bundles and an application to pure twistor *D*-modules. II. *Mem. Amer. Math. Soc.*, 185(870):xii+565, 2007. DOI:10.1090/memo/0870. ↑ 3,
- [RZ10] Luis Ribes and Pavel Zalesskii. Profinite groups, volume 40 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, second edition, 2010. DOI:10.1007/978-3-642-01642-4. ↑ 25

- [Sab13] Claude Sabbah. Théorie de Hodge et correspondance de Hitchin-Kobayashi sauvages (d'après T. Mochizuki). *Astérisque*, (352):Exp. No. 1050, viii, 205−241, 2013. Séminaire Bourbaki. Vol. 2011/2012. Exposés 1043−1058. ↑ 11
- [Sim90] Carlos T. Simpson. Harmonic bundles on noncompact curves. *J. Amer. Math. Soc.*, 3(3):713−770, 1990. DOI:10.2307/1990935. ↑ 11
- [Sim92] Carlos T. Simpson. Higgs bundles and local systems. *Inst. Hautes Études Sci. Publ. Math.*, 75:5–95, 1992. DOI:10.1007/BF02699491, numdam.PMIHES-1992-75-5-0. ↑ 1, 9, 11, 22, 23, 27, 28
- [Tak03] Shigeharu Takayama. Local simple connectedness of resolutions of log-terminal singularities. *Internat. J. Math.*, 14(8):825–836, 2003. DOI:10.1142/S0129167X0300196X. ↑ 10, 22, 25, 26, 28

Daniel Greb, Essener Seminar für Algebraische Geometrie und Arithmetik, Fakultät für Mathematik, Universität Duisburg–Essen, 45117 Essen, Germany

```
E-mail address: daniel.greb@uni-due.de
URL: http://www.esaga.uni-due.de/daniel.greb
```

Stefan Kebekus, Mathematisches Institut, Albert-Ludwigs-Universität Freiburg, Ernst-Zermelo-Strasse 1, 79104 Freiburg im Breisgau, Germany & Freiburg Institute for Advanced Studies (FRIAS), Freiburg im Breisgau, Germany

```
E-mail address: stefan.kebekus@math.uni-freiburg.de URL: https://cplx.vm.uni-freiburg.de
```

Thomas Peternell, Mathematisches Institut, Universität Bayreuth, 95440 Bayreuth, Germany *E-mail address*: thomas.peternell@uni-bayreuth.de

URL: http://www.komplexe-analysis.uni-bayreuth.de

BEHROUZ TAJI, UNIVERSITY OF NOTRE DAME, DEPARTMENT OF MATHEMATICS, 278 HURLEY, NOTRE DAME, IN 46556 USA

```
E-mail address: btaji@nd.edu
URL: http://sites.nd.edu/b-taji
```