

Broadband study of blazar 1ES 1959+650 during flaring state in 2016

S. R. Patel¹, A. Shukla², V. R. Chitnis¹, D. Dorner², K. Mannheim², B. S. Acharya¹, B. J. Nagare³

¹ Department of High Energy Physics, Tata Institute of Fundamental Research, Mumbai 400005, India

² Universität Würzburg, 97074 Würzburg, Germany

³ Department of Physics, University of Mumbai, Santacruz (East), Mumbai-400098, India

ABSTRACT

Aims. The nearby TeV blazar 1ES 1959+650 ($z=0.047$) was reported to be in flaring state during June - July 2016 by Fermi-LAT, FACT, MAGIC and VERITAS collaborations. We studied the spectral energy distributions (SEDs) in different states of the flare during MJD 57530 - 57589 using simultaneous multiwaveband data to understand the possible broadband emission scenario during the flare.

Methods. The UV/optical and X-ray data from UVOT and XRT respectively on board Swift and high energy γ -ray data from Fermi-LAT are used to generate multiwaveband lightcurves as well as to obtain high flux states and quiescent state SEDs. The correlation and lag between different energy bands is quantified using discrete correlation function. The synchrotron self Compton (SSC) model was used to reproduce the observed SEDs during flaring and quiescent states of the source.

Results. A decent correlation is seen between X-ray and high energy γ -ray fluxes. The spectral hardening with increase in the flux is seen in X-ray band. The powerlaw index vs flux plot in γ -ray band indicates the different emission regions for 0.1 - 3 GeV and 3-300 GeV energy photons. Two zone SSC model satisfactorily fits the observed broadband SEDs. The inner zone is mainly responsible for producing synchrotron peak and high energy γ -ray part of the SED in all states. The second zone is mainly required to produce less variable optical/UV and low energy γ -ray emission.

Conclusions. Conventional single zone SSC model does not satisfactorily explain broadband emission during observation period considered. There is an indication of two emission zones in the jet which are responsible for producing broadband emission from optical to high energy γ -rays.

Key words. Radiation mechanisms: non-thermal – Galaxies: BL Lacertae objects: individual: 1ES 1959+650 – Gamma rays: general – X-rays: galaxies

1. Introduction

Blazars are a subclass of Active Galactic Nuclei (AGN) having relativistic jets pointing close to our line of sight (Urry & Padovani 1995). Jets emit highly variable non-thermal radiation spanning wide band of frequencies from radio to γ -rays. Blazars include two types of objects, BL Lacertae (BL Lac) characterized by featureless optical spectra and flat spectrum radio quasars (FSRQ) which show prominent emission lines.

The spectral energy distributions (SEDs) of these objects show characteristic two hump structure. The first low frequency hump is attributed to the synchrotron radiation from relativistic electrons, while second high frequency hump is understood as possibly corresponding to inverse Compton scattering of these synchrotron photons (SSC i.e. Synchrotron Self-Compton) or external photons (EC i.e. External Compton) by the same population of electrons. Alternative explanation for origin of the second hump is given in terms of hadronic models including neutral pion decay, proton synchrotron (Mannheim 1998) etc. Comprehensive review of these mechanisms is given by Böttcher (2007).

Depending on the position of synchrotron peak in SED, BL Lacs are further divided into three classes (Padovani & Giommi 1995). Low frequency BL Lac (LBL) objects exhibit synchrotron peak in IR-Optical band, intermediate frequency BL Lac objects (IBL) have their synchrotron peak at optical-UV frequencies and high frequency BL Lac (HBL) objects show synchrotron peak in UV - X-ray band.

The HBL object, 1ES 1959+650 ($z = 0.047$) was first detected in radio band with NARO Green Bank 91 m telescope (Becker et al. 1991; Gregory & Condon 1991) and later in X-ray (Schachter et al. 1993) using Imaging Proportional counter (IPC) on board Einstein Observatory. TeV emission from this source was first observed by Utah Seven Telescope Array collaboration with total significance of 3.9σ above 600 GeV (Nishiyama 1999). The source was later observed during 2002 May 16 to July 8, with strong detection significance of $> 20\sigma$ by Whipple 10 m telescope (Holder et al. 2003). Since then it has shown several flaring episodes at VHE (> 100 GeV) γ -ray energies, with the most noticeable one being in 2002 when it showed enhanced TeV emission without any contemporaneous X-ray flare (Aharonian et al. 2003; Krawczynski et al. 2004; Daniel et al. 2005; Reimer et al. 2005). In 2004, the source was observed in low state by MAGIC collaboration with a flux of about 20% of the Crab and at $\sim 8\sigma$ significance level above ~ 180 GeV (Albert et al. 2006). Broadband variability of the source was studied in 2012 using strictly simultaneous observations from VERITAS and Swift and reflected emission scenario was used to explain the variability (Aliu et al. 2014). The preliminary analysis of data from Fermi-LAT and various ground based Cherenkov experiments such as FACT, MAGIC, VERITAS as reported by Buson et al. (2016), indicated flaring activity in the source 27 April 2016 onwards.

In the present paper, we have examined the multiwaveband emission from this source over 800 days during MJD 57000 to

57800 (9 December 2014 to 16 February 2017) and studied the broadband variability of this source for the period from MJD 57530 to 57589 (22 May 2016 - 20 July 2016) during which it showed increased flux in X-ray, Fermi-LAT (100 MeV - 300 GeV) and TeV bands. To have good statistics in LAT energy band (0.1 -300 GeV), we chose six periods of 10 days each to sample the complete flare and investigated its emission mechanism in different states during the flare using SSC model. We also studied SED corresponding to 10 days period when source was in low state. The paper is organized as follows. In section 2 various data sets and analysis methods are described. In section 3 timing and spectral studies are elaborated. The SED modeling is outlined in section 4 followed by discussion and conclusions in section 5.

2. Multiwaveband observations and analysis

We have studied the multiwaveband data from radio to γ -rays spanning the period of more than two years from 9 December 2014 (MJD 57000) to 16 February 2017 (MJD 57800). We analyzed UV-optical data from Swift-UVOT, X-ray data from Swift-XRT and high energy γ -ray data from Fermi-LAT. We also used publicly available data from OVRO, SPOL, MAXI and Swift-BAT. Details of these data sets and analysis procedure are given below.

2.1. High energy γ -ray observations

High energy γ -ray data covering the energy range of 100 MeV - 300 GeV was obtained from Large Area Telescope (LAT) on board Fermi spacecraft (Atwood et al. 2009). Data were analyzed using Science Tools version v10r0p5. User contributed *erico* package (Sanchez & Deil 2013) was used. The events were extracted from the circular region of interest (ROI) of 20° centered on the source. Zenith angle cut of 90° was applied to filter the background γ -rays from Earth's limb. To select good time intervals, filter with '(DATA_QUAL>0)&&(LAT_CONFIG==1)' was used. The spectral analysis was carried out using isotropic emission model (*iso_P8R2_SOURCE_V6_v06.txt*) and galactic diffuse emission component model (*gll_iem_v06.fit*) with post launch instrument response function (*P8R2_SOURCE_V6*) and using unbinned likelihood analysis. The sources lying within ROI of 15° radius around the 1ES 1959+650 from the 3FGL catalog were included in the model XML file. In likelihood fit, both spectral and normalization parameters of the sources within 5° radius around the source were left free to vary while keeping parameters for all other sources fixed at their catalog value. There are 72 point sources and 1 extended source in the model file. The parameters of extended source were also kept free in maximum likelihood analysis. Source spectrum was modeled with a power law. The lightcurves were generated with 10 days binning in energy range of 0.1 -3 GeV and 3 - 300 GeV.

2.2. X-ray observations

Publicly available hard X-ray data from Burst Alert Telescope (BAT) on board Swift are used¹. These are daily average count rates over the energy range of 15-50 keV. Soft X-ray data covering the energy range of 2-20 keV from Monitor of All sky X-ray Images (MAXI) on board International Space Station (ISS, Mat-

suoka et al. (2009)) are obtained from MAXI website². These are daily average flux values.

Also soft X-ray data covering the energy range of 0.3 - 8 keV from X-ray telescope (XRT) on board Swift have been used (Burrows et al. 2005). The data were analyzed using XRT data analysis software (XRTDAS) distributed within the HEASOFT package (v6.19). The *xrtpipeline-0.13.2* tool was used to generate the cleaned event files. Data from 34 observations during 22 April - 20 July 2016 corresponding to flare region were analyzed. The source and background data were extracted from the circular region of 20 pixels radius around the source and 40 pixels radius region away from the source respectively. The spectral data were combined into six groups of 10 days each as shown in Fig. 1 and rebinned with minimum 20 photons per bin. Six spectra were fitted with absorbed powerlaw model as well as with log parabola model. Spectral form of log parabola model is given by

$$dN/dE = K(E/E_b)^{-\alpha-\beta \log(E/E_b)} \quad (1)$$

where α is the spectral index and E_p is the point of maximum curvature given by

$$E_p = E_b 10^{(2-\alpha)/2\beta} \quad (2)$$

While fitting the spectrum, E_b is fixed at 1 keV. To correct for interstellar absorption of soft X-rays along line of sight, neutral hydrogen column density (N_H) is fixed at $1.0 \times 10^{21} \text{ cm}^{-2}$ (Kalberla et al. 2005). Swift XRT light curve spanning two years data, over the energy range of 0.3-10 keV is shown in Fig. 1. This is publicly available Swift-XRT lightcurve obtained from the website³.

2.3. UV, optical and radio observations

We have analyzed Swift-UVOT (Roming et al. 2005) data for the period of two years. Data are available in six different filters covering optical and UV band, viz. V, B, U, UVW1, UVM2 and UVW2. For each filter, images were added using tool *uvotimsum* and flux/magnitude values were obtained using tool *uvotsource*. For V, B and U filters, source counts were extracted from circular region with radius of 5" around the source location, whereas for UVW1, UVM2 and UVW2 filters, region with radius of 10" was used. The Galactic extinction correction (Schlegel et al. 1998) of $E_{B-V} = 0.177$ mag was applied to observed magnitude. Observed magnitudes were then converted into flux using zero point magnitudes (Poole et al. 2008). Host galaxy contribution (Tagliaferrri et al. 2008) of 1.1 mJy, 0.4 mJy and 0.1 mJy were subtracted for V, B and U filter respectively. No correction is applied at UV frequencies as host galaxy contribution is negligible at these frequencies. Fig. 1 shows lightcurves from optical U and ultraviolet M2 filters. Other UVOT bands show similar trend, hence only two bands are shown to avoid cluttering.

1ES 1959+650 is being monitored with SPOL CCD Imaging/Spectropolarimeter at steward observatory at University of Arizona (Smith et al. 2009) regularly as a part of the Fermi multiwavelength support programme. The publicly available optical R-band and V-band photometric and linear polarization data were obtained from the SPOL website⁴. The R-band fluxes are

² MAXI : http://134.160.243.77/star_data/J1959+651/J1959+651_00055054g_lc_1day_all.dat

³ XRT : <http://www.swift.psu.edu/monitoring/data/1ES1959+650/lightcurve2.txt>

¹ BAT : <http://swift.gsfc.nasa.gov/results/transients/weak/1ES1959p650.lc4.txt> <http://james.as.arizona.edu/~psmith/Fermi/>




Fig. 1. Two years light curves of IES 1959+650 from MJD 57000 to 57800, Panel-1 : Radio flux density at 15 GHz in Jy ; Panel-2 : Degree of polarization from CCD-SPOL observations , Panel-3 : Polarization angle from CCD-SPOL observations, Panel-4 : UVOT-U, UVOT-W2 band and CCD-SPOL-R band fluxes in mJy (Corrected for host galaxy contribution), Panel-5 : Swift-XRT count rate in counts/s, Panel-6 : MAXI flux in photons $\text{cm}^{-2}\text{s}^{-1}$ (daily average), Panel-7 : Swift-BAT flux in counts $\text{cm}^{-2}\text{s}^{-1}$ (daily average), Panel-8 : Fermi-LAT 0.1-3 GeV flux in photons $\text{cm}^{-2}\text{s}^{-1}$ (10 days average), Panel-9 : Fermi-LAT 3-300 GeV flux in photons $\text{cm}^{-2}\text{s}^{-1}$ (10 days average) ; SEDs are computed for the periods marked by dotted lines.

corrected for host galaxy contribution which is 0.84 mJy (Nilsson et al. 2007). From available data the degree of linear polarization and polarization angle were found to vary between 0.37% to 3.98% and 107° to 173° respectively.

1ES 1959+650 is also observed regularly in radio as a part of Fermi monitoring programme by Owens Valley Radio Observatory (OVRO; Richards et al. (2011)). The publicly available data at 15 GHz were used in lightcurve from OVRO website⁵.

3. Results

In this section we present results from multiwaveband temporal and spectral studies of 1ES 1959+650 collected over two years period from MJD 57000 to 57800.

3.1. Multiwaveband temporal studies

Fig. 1 shows the lightcurve of 1ES 1959+650 for the period starting from MJD 57000 to MJD 57800. Panels corresponding to various wavebands have been arranged in increasing order of frequency from top to bottom, starting with radio lightcurve from OVRO in the topmost panel to high energy γ -rays from Fermi-LAT in the bottom-most panel. Fermi-LAT flux values over the energy ranges 0.1-3 GeV and 3-300 GeV shown in last two panels, are averaged over 10 days bins. Data from Swift-BAT and MAXI are averaged over a day, whereas Swift-XRT and Swift-UVOT data points correspond to individual observations with typical duration of about hours/minutes. SPOL and OVRO have an integration time of a few seconds. The source exhibited the flare in 2016 during MJD 57530 - 57589 which is clearly seen in the figure particularly in X-ray and γ -ray bands. We studied correlated variability in various wavebands. In this work we have concentrated on detailed studies during flaring episode around MJD 57530 - 57589 (22 May - 20 July 2016). This episode is divided into six periods of ten days each. Quiescent state data during MJD 57177 - 57186 (4 -13 June 2015) is compared with this flare.

3.1.1. Variability

We have studied variability of lightcurves in various wavebands on various time scales. Variability is estimated in terms of fractional variability amplitude, F_{var} parameter (Vaughan et al. 2003; Chitnis et al. 2009). This parameter estimates variability intrinsic to the source and is given by

$$F_{var} = \sqrt{\frac{S^2 - \sigma_{err}^2}{\bar{x}^2}} \quad (3)$$

where S^2 is the sample variance, σ_{err}^2 the mean square error and \bar{x} the unweighted sample mean. The error on the F_{var} is given by

$$\sigma_{F_{var}} = \sqrt{\left(\frac{\sqrt{\frac{1}{2N} \frac{\sigma_{err}^2}{\bar{x}^2 F_{var}}}}{\bar{x}}\right)^2 + \left(\frac{\sqrt{\sigma_{err}^2} \frac{1}{\bar{x}}}{\bar{x}}\right)^2} \quad (4)$$

Variability strength is estimated on time scales of 10 and 20 days in various wavebands in present work. Results are given in Table. 1 and plotted in Fig. 2 for 10 days binning. The large error bar on BAT is due to poor sensitivity of the instrument.

⁵ OVRO : http://www.astro.caltech.edu/ovroblazars/data.php?page=data_return




Fig. 2. Fractional variability for 10 days binned lightcurve

Table 1. Fractional variability strength in various wavebands

Waveband	10 days binning	20 days binning
Radio (15 GHz)	11.16 +/- 1.19	11.46 +/- 1.59
Optical U	27.91 +/- 2.65	26.96 +/- 3.34
UV M2	25.28 +/- 2.35	23.29 +/- 2.88
XRT (0.3 - 8 keV)	43.62 +/- 3.95	38.27 +/- 4.58
MAXI (2-20 keV)	52.45 +/- 6.13	48.30 +/- 6.84
BAT (15-50 keV)	90.47 +/- 15.78	96.39 +/- 16.66
Fermi-LAT (0.1-3 GeV)	53.45 +/- 4.97	51.08 +/- 6.21
Fermi-LAT (3-300 GeV)	58.42 +/- 6.55	52.88 +/- 7.45

Variability seems to increase with frequency from radio to X-rays and decrease in high energy γ -rays compared to hard X-rays. Similar trend was seen in Mkn 421 (Sinha et al. 2016).

3.1.2. Correlations

The correlations between various lightcurves are quantified over the entire observation period (MJD 57000 to 57800) using discrete correlation function (DCF) (Edelson & Krolik 1988). For two discrete data sets a_i and b_j , the unbinned discrete correlation is defined as,

$$UDCF_{ij} = \frac{(a_i - \bar{a})(b_j - \bar{b})}{\sqrt{(\sigma_a^2 - e_a^2)(\sigma_b^2 - e_b^2)}} \quad (5)$$

for all measured pairs (a_i, b_j) having pairwise lag $\Delta t_{ij} = t_j - t_i$. σ_a and σ_b are standard deviations of each data train and e_a, e_b are the measurement errors associated with them. DCF is then given by averaging M pairs for which $(\tau - \Delta\tau/2) \leq \Delta t_{ij} < (\tau + \Delta\tau/2)$,

$$DCF(\tau) = \frac{1}{M} UDCF_{ij} \quad (6)$$

The error on DCF is given by,

$$\sigma_{DCF}(\tau) = \frac{1}{M-1} \left\{ \sum [UDCF_{ij} - DCF(\tau)] \right\}^{1/2} \quad (7)$$

Each data set is linearly detrended (Welsh 1999) before using above function. The correlation coefficients along with lags between the lightcurves is listed in Table- 2 and their plots are shown in Fig. 3. Values of correlation coefficients given in table and figure are estimated for 3 days binning in lag to improve statistics, except for 0.1-3 GeV - 3-300 GeV case where bin




Fig. 3. Discrete correlation function vs time lag between various energy bands

size of 10 days is used. These values are consistent with one day binning of lag in all the cases. It can be seen that there is a strong correlation between optical (V band) and UV (M2 band) at zero lag. X-ray data shows good correlation with low and high energy γ -rays with no visible lag. Correlation is better with higher energy γ -rays. Low and high energy γ -rays are also correlated. The mild correlation is seen between optical (V band) and γ -rays with a lag of about 38-90 days. Similar lag is seen between radio and γ -rays. On the other hand, correlation between optical and X-ray band is rather weak. This indicates X-ray and γ -rays, particularly higher energy ones, may have similar origin. Whereas optical/UV band emission may have different origin. The study of long-term data (2005-2014) carried out by Kapanadze et al. (2016) showed that source displayed slow variation in optical R-band and exhibited optical flare lasting for several months to years. They also report relatively weak correlation between XRT and UVOT band.

Table 2. Maximum DCF between various wavebands

Bands	DCF	Lag (Days)
XRT-0.1-3 GeV LAT	0.58 +/- 0.14	0.0
XRT-3-300 GeV LAT	0.92 +/- 0.19	0.0
0.1-3 GeV LAT-3-300 GeV LAT	0.72 +/- 0.17	0.0
UVOT (V)- XRT	0.38 +/- 0.08	-32.8
OVRO (15 GHz)-0.1-3 GeV LAT	0.81 +/- 0.22	-41.8
OVRO (15 GHz)-3-300 GeV LAT	0.79 +/- 0.17	-68.7
UVOT (V)- UVOT (M2)	1.00 +/- 0.11	0.0
UVOT (V)-0.1-3 GeV LAT	0.61 +/- 0.18	-89.6
UVOT (V)-3-300 GeV LAT	0.41 +/- 0.17	-38.8

3.1.3. Lognormality

Lognormality, i.e., log-normal distribution of flux and linear rms-flux relation, has been detected in several X-ray binaries (Uttley & McHardy 2001; Scaringi et al. 2012). It has also been detected in blazars including BL Lac (detected in X-ray regime, Giebels & Degrange (2009)), in multiple wavebands in PKS 2155-304 (Chevalier et al. 2015), Mkn 421 (Sinha et al. 2016), IES 1011+496 (Sinha et al. 2017), PKS 1510-089 (Kushwaha et al. 2016) etc. Flux distributions for IES 1959+650 covering two years' data for various wavebands from radio to γ -ray are shown in Fig. 4. Data are fitted with Gaussian as well as lognormal distribution and fits are shown. Reduced χ^2 values for both the distributions are listed in Table 3. Reduced χ^2 is lower for lognormal compared to Gaussian fit in all the cases, except for 0.1-3 GeV Fermi-LAT. In order to check significance of this reduction in χ^2 , the F-statistic was used. Assuming null hypothesis, i.e. no significant difference in variances from these two models, F-values were calculated ($F_{calculated}$). These were compared with F-values for 95% confidence level ($F_{95\%}$), for ν_g and ν_l degrees of freedom (corresponding to Gaussian and lognormal fit respectively). These values are listed in Table 3 for each waveband. It can be seen that $F_{calculated}$ are less than $F_{95\%}$ in all the cases, which means that we can not reject null hypothesis. In other words, even though lognormal gives lower χ^2 and better fit, Gaussian behaviour can not be ruled out at 95% confidence because of low number of degrees of freedom (dof). Fig. 5 shows the plot of excess variance $\sigma_{excess}^2 = (S^2 - \sigma_{err}^2)$ as a function of flux for various X-ray wavebands, binned over a period of 10 days. The excess variance were calculated for those bins which are having at least 5 flux points and it is plotted against mean flux. We do not see very clear linear trend in these plots as was seen in case of Mkn 421 by Sinha et al. (2016).




Fig. 4. Gaussian and Lognormal fit to flux distribution in various energy bands. The binning used is same as in Fig. 1 except for MAXI and BAT where 10 days binning is used.




Fig. 5. Excess variance vs mean flux in different X-ray energy bands : The data are binned into 10 days bins and excess variance is computed only for those bins which have at least 5 flux measurements

3.2. Spectral studies

We studied X-ray spectra using Swift-XRT data for the flare state during MJD 57530 - 57589. This flare was divided into six states, each spanning ten days period. These states are marked by ver-

tical dotted lines in Fig. 1. We also studied X-ray spectrum in the interval MJD 57177 - 57187 which corresponds to the quiescent state of the source. These seven average spectra for each bin were fitted with a powerlaw with line of sight absorption, over

Table 3. Goodness of fit for flux distributions in various energy bands

Waveband	Gaussian χ_r^2 (dof (ν_g))	Lognormal χ_r^2 (dof (ν_l))	$F_{calculated}$	$F_{95\%}$
OVRO (15GHz)	10.82 (6)	6.16 (6)	1.77	4.28
UVOT (Optical)	3.35 (6)	1.80 (6)	1.86	4.28
UVOT (UV)	5.56 (3)	2.67 (3)	2.08	9.28
Swift XRT (0.3 - 8 keV)	2.52 (5)	2.18 (5)	1.16	5.05
MAXI (0.2 - 20 keV)	6.82 (5)	4.25 (5)	1.60	5.05
Swift BAT (15 - 50 keV)	3.58 (5)	1.43 (5)	1.60	5.05
Fermi LAT (0.1 - 3 GeV)	1.08 (5)	2.20 (5)	2.05	5.05
Fermi LAT (3 - 300 GeV)	1.69 (4)	0.96 (4)	1.76	6.39

Notes. dof is number of bins minus number of model parameters which is three for both the models; $F_{calculated}$ is ratio of two variances with larger variance in numerator ; $F_{95\%}$ is F-statistic value at 95% confidence level for ν_g and ν_l degrees of freedom

the energy range of 0.3-8 keV. Alternatively spectra were also fitted with logparabola model with line of sight absorption. Best fit parameters for both the models are given in Table 4. Logparabola seems to fit data better than powerlaw, as seen from improvement in values of reduced χ^2 .




Fig. 6. Red: Powerlaw index vs X-ray flux (The correlation coefficient is -0.72), Blue: Logparabola index vs X-ray flux (The correlation coefficient is -0.61)




Fig. 7. Powerlaw index vs LAT flux. The correlation coefficient for 0.1 - 3 GeV band is 0.60

We also analyzed X-ray spectra from individual observations of XRT during the flare. Fig. 6 shows the plot of flux in 0.3-8 keV




Fig. 8. Hardness ratio in Fermi band vs 0.1 - 3 GeV flux, Points corresponding to flare is shown in blue

range as a function of X-ray photon index for both powerlaw and logparabola model. X-ray spectrum seems to harden with the increase in the flux. Similar behaviour from this source is also reported by Gutierrez et al. (2006). However no such correlation was found when source was observed in low state during 2007 - 2011 (Aliu et al. 2013). Spectral hardening with increase in flux is seen in several other blazars including Mkn 421 (Sinha et al. 2016); Mkn 501 (Krawczynski et al. 2000).

We also investigated spectral parameters in γ -ray band using Fermi-LAT data, dividing it into two energy ranges, 0.1-3 GeV and 3-300 GeV. Spectra binned over ten days were fitted with a powerlaw. Variation of flux with powerlaw index for both low and high energy bands is shown in Fig. 7. While lower energy band shows increase in the flux with increase in the photon index, no such trend is seen at higher energies. Similar behaviour at γ -ray energies was seen for Mkn 501 (Shukla et al. 2015). Such trend might be due to different emission regions for low and high energy γ -rays. Fig. 8 shows hardness ratio, defined as a ratio of 3-300 GeV to 0.1-3 GeV flux, as a function of 0.1-3 GeV flux. This also indicates that for the period of 800 days the spectrum is becoming softer at higher flux.

4. SED evolution and modeling

We have studied the evolution of SED during flare. For this purpose, multiwaveband SEDs were generated for six flux states in the flare as well as for one quiescent state. First we tried to model these SEDs using simple single zone Synchrotron Self

Table 4. Spectral fitting parameters for 10 days binned Swift-XRT data

Period (MJD)	Powerlaw Index	Reduced χ^2 (prob)	Logparabola		Reduced χ^2 (prob)
			α	β	
Q (57177-57186)	2.19 +/- 0.02	1.34 (1.55E-03)	2.04 +/- 0.00	0.49 +/- 0.07	1.05 (2.91E-01)
F1 (57530-57539)	2.09 +/- 0.01	1.44 (3.03E-07)	1.97 +/- 0.02	0.33 +/- 0.04	1.17 (2.16E-02)
F2 (57540-57549)	1.89 +/- 0.01	1.72 (2.92E-27)	1.81 +/- 0.01	0.21 +/- 0.01	1.21 (1.89E-04)
F3 (57550-57559)	1.82 +/- 0.01	1.58 (2.16E-19)	1.75 +/- 0.01	0.15 +/- 0.01	1.33 (2.56E-08)
F4 (57560-57569)	1.88 +/- 0.02	0.96 (6.81E-01)	1.81 +/- 0.02	0.17 +/- 0.04	0.91 (8.65E-01)
F5 (57570-57579)	1.74 +/- 0.01	1.28 (6.81E-05)	1.61 +/- 0.02	0.28 +/- 0.03	1.05 (2.05E-01)
F6 (57580-57589)	1.96 +/- 0.01	1.39 (1.51E-08)	1.87 +/- 0.01	0.23 +/- 0.02	1.13 (2.64E-02)

Notes. Values in the brackets in column 3 and 6 denote the value of null hypothesis probability.

Compton (SSC) model using the code developed by Krawczynski et al. (2004). This model assumes the emission zone to be a blob of radius R traveling down the jet with bulk Lorentz factor Γ towards the observer at an angle of θ . The emission zone is filled with non-thermal electron distribution and randomly oriented magnetic field. The electron population can be described by broken powerlaw, having low and high energy indices p_1 and p_2 . The radius R of the emission region can be constrained by observed doubling time scale t_{var} using relation,

$$R \sim \frac{c\delta t_{var}}{(1+z)} \quad (8)$$

The six flux states during different epochs of the flare are denoted by F1, F2, F3, F4, F5 and F6 while quiescent state is denoted by Q. The SEDs were modeled with jet parameters θ and Γ to be 3° and 9.4 respectively which corresponds to Doppler factor of ~ 15 . The value of Doppler factor is consistent with the value reported in the past (Krawczynski et al. 2004; Gutierrez et al. 2006). The blob radius of 7.84×10^{16} cm was assumed. While modeling θ is kept fixed at 3 degree and it is also assumed that emitting region has same cross section as jet.

We find that due to sharp curvature of SEDs in X-ray region, we always underpredict the optical/UV flux with one zone SSC model. Also except for F2 state, one zone SSC model does not fit γ -ray flux below ~ 3 GeV. Hence we made an attempt to explain the observed SEDs with two zone SSC model in which resultant emission is the sum of emission from two comoving blobs of different sizes. We assumed the size of the second (outer) blob to correspond to $t_{var} \sim 10$ days and having same Doppler factor as that of the inner blob. The cross plot of the flux-power law index in 0.1-3 GeV and 3-300 GeV bands suggests the different emission regions for these energy bands (Fig. 7). It is reported in long term (2005-2014) multiwavelength cross-correlation study that one zone SSC scenario was not always suitable to explain the emission from this source (Kapanadze et al. 2016).

The fit for all six high states is shown in Fig. 9 and one quiescent state is shown in Fig. 10. The Fig. 11 shows the variation of inner blob emission as the flare evolves. The SSC model parameters for two zones are listed in Table. 5. Apart from conventional single-zone SSC model, External Compton (EC), leptohadronic model and two independent SSC models were tried for fitting SEDs in the past (Backes et al. 2012). Authors report that two zone SSC model described the observed SED reasonably well for this source. The two zone SSC model was also used to reproduce observed SED during 2011 observation of Mkn 501 (Shukla et al. 2015, 2016).

In homogeneous leptonic model, one expects a break in electron energy distribution (EED) where Lorentz factor is given by

$$\gamma_c = \frac{3\pi m_e c^2}{\sigma_t B^2 R} \quad (9)$$

At this γ_c , the escape time from the source equals the synchrotron cooling time (Abdo et al. 2011; Graff et al. 2008). The Fig. 12 shows variation of fitted B with γ_{br} for flare states F1-F6. Theoretical curve based on Eq.(9) assuming $R = 7.84 \times 10^{16}$ cm is shown by solid line for inner blob. As per Eq. (9), we find that observed B is upto $\sim 45\%$ lower than the expected one (Fig. 12). This means that we observe break in electron spectrum at lower energies than the one expected in case of similar escape and cooling time scales.

5. Discussion and conclusions

The lognormal fit to the flux distribution of two years of data gives improved reduced χ^2 as compared to Gaussian fit. However we do not see this difference between models to be significant based on F-test. This could be due to smaller size of the data set. Since tail is seen in the flux distribution, possibly longer data set may show significant preference to lognormal fit. Several authors have seen lognormal behaviour of flux distribution in X-ray binaries and BL-Lac sources. The lognormal flux behaviour in BL-Lac is generally considered as indication of the accretion disk variability's imprint onto the jet (McHardy 2008). In the present data set, we do not see lognormality convincingly. Longer data set may lead to conclusive result.


The temporal analysis of two years data on 1ES 1959+650 shows that source did not exhibit significant variation in optical band. However, source showed significant flux variation in X-ray and γ -ray band. The DCFs are computed to quantify the correlations and lags between measurements from various instruments covering different energy bands over two years. Significant correlation is seen between fluxes of X-ray and both high and low energy γ -rays. No significant correlation of optical/UV fluxes with other energy bands was seen. This could be due to different emission region of optical/UV photons and these photons are up-scattered to low energy γ -rays, which might be observed by Fermi-LAT.

We found the X-ray spectrum to be curved during most of the observations and hence logparabola fit to the X-ray spectrum is found to be better than conventional power law fit. Log parabola fit to X-ray data also suggest that particle acceleration should be stochastic in nature. The spectral evolution was found to be 'harder when brighter' in X-ray during the observation period of ~ 2 months covering flare. However, 'softer when brighter' behaviour was also seen in γ -rays (Fig. 8) when source is not very active.

Table 5. SSC model parameters for flare and quiescent states for two zones with Doppler factor (δ) ~ 15 , $\theta = 3$, $R_{inner}=7.84 \times 10^{16}$ cm ($t_{var} \sim 2$ days) and $R_{outer}=3.92 \times 10^{17}$ cm ($t_{var} \sim 10$ days)

SED state	B [10^{-2} G]	U_e [10^{-3} erg/cc]	γ_{min} [10^4]	γ_{max} [10^6]	γ_{br} [10^5]	p_1	p_2	η (U_e/U_B)
Q (Inner blob)	1.10	0.50	2.46	1.55	5.52	2.40	3.40	103.85
F1 (Inner blob)	0.95	1.10	5.51	3.10	4.92	2.35	3.35	306.33
F2 (Inner blob)	1.01	1.85	0.31	3.10	9.81	2.15	3.15	451.32
F3 (Inner blob)	0.93	1.80	0.62	3.90	9.37	2.10	3.10	528.72
F4 (Inner blob)	1.05	1.50	6.18	3.32	9.81	2.15	3.15	341.92
F5 (Inner blob)	0.90	1.40	3.47	3.48	14.18	2.10	3.10	434.39
F6 (Inner blob)	0.90	1.00	1.96	3.90	6.19	2.00	3.00	341.31
Q (Outer blob)	0.70	0.13	0.20	0.98	0.62	2.85	3.85	66.07
F1 (Outer blob)	0.45	0.14	0.49	2.46	0.78	2.75	3.75	170.65
F2 (Outer blob)	0.43	0.14	0.44	1.96	1.24	2.70	3.70	194.94
F3 (Outer blob)	0.48	0.18	0.25	2.46	1.55	2.70	3.70	194.94
F4 (Outer blob)	0.45	0.23	0.31	4.92	0.74	2.70	3.70	279.25
F5 (Outer blob)	0.45	0.23	0.31	4.92	0.74	2.70	3.70	279.25
F6 (Outer blob)	0.50	0.22	0.20	4.92	0.78	2.50	3.50	221.17

Notes. U_e : Electron energy density ; γ_{min} : Minimum value of Lorentz factor of electron present in the emission region ; γ_{max} : Maximum value of Lorentz factor of electron present in the emission region ; γ_{br} : Lorentz factor at break in electron injection spectrum ; η : Equipartition coefficient


Fig. 9. SEDs fitted with two zone SSC model for six states during flare




Fig. 10. Quiescent state SED fitted with two zone SSC model

We found a significant change in electron density across the flare in both inner and outer regions by modeling of SEDs. However a maximum $\sim 20\%$ change is observed in magnetic field. Either injection of new particles or re-acceleration of particles might have increased electron energy density in the inner zone. This enhanced electron energy density might be responsible for the high state and flaring activity of the source. Also similar magnetic field in outer blob might be due to re-amplification of magnetic field by passing shock wave. Narrow EED and spectral hardening are found in the electron spectrum during the flaring period from inner blob. The narrow EED can be reconciled as a stochastic acceleration process via Fermi II order acceleration scenario, where randomly moving Alfvén waves may accelerate particles in turbulent medium.

The observed break in the particle spectrum found by SED modeling is at much lower energies than the expected from canonical jet model (Eq. 9) as seen from Fig. 12. This observed break in particle spectrum might be an outcome of effective inverse Compton cooling and this break appears at Lorentz factor where inverse Compton cooling equals adiabatic losses. In our SED modeling inverse Compton losses dominate synchrotron losses. Similar behaviour is observed in Mkn 501 (Acciari et al. 2011). The magnetic field and electron energy density in inner region was found to be an order of magnitude higher than the outer region. Moreover, electron spectra were found much softer than inner blob. The particle spectra in outer blob might be outcome of shock acceleration where particles are cooled through synchrotron and IC losses.

With increase in the flux synchrotron peak is found to shift to right and vice-versa i.e. 'bluer when brighter' behaviour of the source during this period. Similar behaviour is reported by Tagliaferri et al. (2008). Among the six states F2, F3 and F4 states sampled the peak of the flare. During F3 and F4 states the high energy γ -ray flux is comparable while there is a significant increase in the X-ray flux as the state evolved from F3 to F4. It was then followed by the decrease in the flux in both synchrotron and IC region as the flare decays in F5 and F6 state.

We are able to reproduce SEDs satisfactorily with broken powerlaw electron distribution with $(p_2 - p_1) \sim 1$, for both inner and outer regions. This corresponds to spectral index change of $\Delta \alpha = 0.5$ for synchrotron emission, which is expected in the case of canonical cooling break in homogeneous model. We could reproduce all the flaring states and one quiescent state with Doppler factor of ~ 15 for both comoving plasma blobs. Higher γ_{min} seen in inner blob compared to the outer blob suggests the




Fig. 11. Variation of inner zone of SSC emission during 6 flare and one quiescent states




Fig. 12. The temporal evolution of model parameter B with γ_{br} for inner zone ; Theoretical curve shown by black continuous line is as per Eq. (9).

narrower EED of inner blob. The narrow EED is responsible for producing the hard spectrum (Tavecchio et al. 2009; Katarzyński et al. 2006; Lefa et al. 2011). During the flaring activity inner blob is mostly responsible for synchrotron emission for all the states while outer blob is responsible for optical/UV and low energy γ -ray ($0.1 - \sim 3$ GeV). Except for F2 state, the outer blob contributes to observed SEDs significantly for all the states including quiescent one in reproducing both the humps. However in F2 state, outer blob was required to reproduce the optical/UV emission from source. We see clear plateau in this low energy γ -ray in F1, F3 and F4 states. Such plateau will be seen if the observed emission is sum of the emission from different region (Shukla et al. 2015).

Acknowledgements. In this paper, we used Enrico, a community-developed Python package to simplify Fermi-LAT analysis (Sanchez & Deil 2013). This research has made use of the XRT Data Analysis Software (XRTDAS) developed under the responsibility of the ASI Science Data Center (ASDC), Italy. Data from the Steward Observatory spectropolarimetric monitoring project were used. This program is supported by Fermi Guest Investigator grants NNX08AW56G, NNX09AU10G, NNX12AO93G, and NNX15AU81G. In this research we used data from the OVRO 40-m monitoring program (Richards et al. 2011) which is supported in part by NASA grants NNX08AW31G, NNX11A043G, and NNX14AQ89G and NSF grants AST-0808050 and AST-1109911. We thank A. R. Rao for his useful suggestions and comments.

References

- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2011, *ApJ*, 736, 131
- Acciari, V. A., Arlen, T., Aune, T., et al. 2011, *ApJ*, 729, 2
- Aharonian, F., Akhperjanian, A., Beilicke, M., et al. 2003, *A&A*, 406, L9
- Albert, J., Aliu, E., Anderhub, H., et al. 2006, *ApJ*, 639, 761
- Aliu, E., Archambault, S., Arlen, T., et al. 2014, *ApJ*, 797, 89
- Aliu, E., Archambault, S., Arlen, T., et al. 2013, *ApJ*, 775, 3
- Atwood, W. B., Abdo, A. A., Ackermann, M., et al. 2009, *ApJ*, 697, 1071
- Backes, M., Uellenbeck, M., Hayashida, M., et al. 2012, in *American Institute of Physics Conference Series*, Vol. 1505, American Institute of Physics Conference Series, 522–525
- Becker, R. H., White, R. L., & Edwards, A. L. 1991, *ApJS*, 75, 1
- Böttcher, M. 2007, *Ap&SS*, 309, 95
- Burrows, D. N., Hill, J. E., Nousek, J. A., et al. 2005, *Space Sci. Rev.*, 120, 165
- Buson, S., Magill, J. D., Dorner, D., et al. 2016, *The Astronomer's Telegram*, 9010
- Chevalier, J., Kastendieck, M. A., Rieger, F. M., et al. 2015, in *International Cosmic Ray Conference*, Vol. 34, 34th International Cosmic Ray Conference (ICRC2015), 829
- Chitnis, V. R., Pendharkar, J. K., Bose, D., et al. 2009, *ApJ*, 698, 1207
- Daniel, M. K., Badran, H. M., Bond, I. H., et al. 2005, *ApJ*, 621, 181
- Edelson, R. A. & Krolik, J. H. 1988, *ApJ*, 333, 646
- Giebels, B. & Degrange, B. 2009, *A&A*, 503, 797
- Graff, P. B., Georganopoulos, M., Perlman, E. S., & Kazanas, D. 2008, *ApJ*, 689, 68
- Gregory, P. C. & Condon, J. J. 1991, *ApJS*, 75, 1011
- Gutierrez, K., Badran, H. M., Bradbury, S. M., et al. 2006, *ApJ*, 644, 742
- Holder, J., Bond, I. H., Boyle, P. J., et al. 2003, *ApJ*, 583, L9
- Kalberla, P. M. W., Burton, W. B., Hartmann, D., et al. 2005, *A&A*, 440, 775
- Kapanadze, B., Romano, P., Vercellone, S., et al. 2016, *MNRAS*, 457, 704
- Katarzyński, K., Ghisellini, G., Tavecchio, F., Gracia, J., & Maraschi, L. 2006, *MNRAS*, 368, L52
- Krawczynski, H., Coppi, P. S., Maccarone, T., & Aharonian, F. A. 2000, *A&A*, 353, 97
- Krawczynski, H., Hughes, S. B., Horan, D., et al. 2004, *ApJ*, 601, 151
- Kushwaha, P., Chandra, S., Misra, R., et al. 2016, *ApJ*, 822, L13
- Lefa, E., Rieger, F. M., & Aharonian, F. 2011, *ApJ*, 740, 64
- Mannheim, K. 1998, *Science*, 279, 684
- Matsuoka, M., Kawasaki, K., Ueno, S., et al. 2009, *PASJ*, 61, 999
- McHardy, I. 2008, in *Blazar Variability across the Electromagnetic Spectrum*, 14
- Nilsson, K., Pasanen, M., Takalo, L. O., et al. 2007, *A&A*, 475, 199
- Nishiyama, T. 1999, *International Cosmic Ray Conference*, 3, 370
- Padovani, P. & Giommi, P. 1995, *MNRAS*, 277, 1477
- Poole, T. S., Breeveld, A. A., Page, M. J., et al. 2008, *MNRAS*, 383, 627
- Reimer, A., Böttcher, M., & Postnikov, S. 2005, *ApJ*, 630, 186
- Richards, J. L., Max-Moerbeck, W., Pavlidou, V., et al. 2011, *ApJS*, 194, 29
- Roming, P. W. A., Kennedy, T. E., Mason, K. O., et al. 2005, *Space Sci. Rev.*, 120, 95
- Sanchez, D. A. & Deil, C. 2013, *ArXiv e-prints* [[arXiv:1307.4534](https://arxiv.org/abs/1307.4534)]
- Scaringi, S., Körding, E., Uttley, P., et al. 2012, *MNRAS*, 421, 2854
- Schachter, J. F., Stocke, J. T., Perlman, E., et al. 1993, *ApJ*, 412, 541
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Shukla, A., Chitnis, V. R., Singh, B. B., et al. 2015, *ApJ*, 798, 2
- Shukla, A., Mannheim, K., Chitnis, V. R., et al. 2016, *ApJ*, 832, 177
- Sinha, A., Sahayanathan, S., Acharya, B. S., et al. 2017, *ApJ*, 836, 83
- Sinha, A., Shukla, A., Saha, L., et al. 2016, *A&A*, 591, A83
- Smith, P. S., Montiel, E., Rightley, S., et al. 2009, *ArXiv e-prints* [[arXiv:0912.3621](https://arxiv.org/abs/0912.3621)]
- Tagliaferri, G., Foschini, L., Ghisellini, G., et al. 2008, *ApJ*, 679, 1029
- Tavecchio, F., Ghisellini, G., Ghirlanda, G., Costamante, L., & Franceschini, A. 2009, *MNRAS*, 399, L59
- Urry, C. M. & Padovani, P. 1995, *PASP*, 107, 803
- Uttley, P. & McHardy, I. M. 2001, *MNRAS*, 323, L26
- Vaughan, S., Edelson, R., Warwick, R. S., & Uttley, P. 2003, *MNRAS*, 345, 1271
- Welsh, W. F. 1999, *PASP*, 111, 1347