

What is a dineutron? Exploring its manifestation and nature in two-neutron emission using a dynamical dineutron model

L.V. Grigorenko,^{1,2,3} J.S. Vaagen,⁴ and M.V. Zhukov⁵

¹*Flerov Laboratory of Nuclear Reactions, JINR, RU-141980 Dubna, Russia*

²*National Research Nuclear University “MEPhI”,
Kashirskoye shosse 31, RU-115409 Moscow, Russia*

³*National Research Centre “Kurchatov Institute”, Kurchatov sq. 1, RU-123182 Moscow, Russia*

⁴*Institute of Physics and Technology, University of Bergen, N-5007 Bergen, Norway*

⁵*Department of Physics, Chalmers University of Technology, S-41296 Göteborg, Sweden*

Emission of two neutrons or two protons in reactions and decays is often discussed in terms of “dineutron” or “diproton” emission. The discussion often leans intuitively on something described by Migdal-Watson approximation. In this work we propose a way to formalize situations of dineutron emission. It is demonstrated that properly formally defined dineutron emission may reveal properties which are drastically different from those traditionally expected, and properties which are actually observed in three-body decays.

I. INTRODUCTION

The idea of final state interaction (FSI) treatment in Migdal-Watson approximation [1, 2] is one of the basic concepts of nuclear reaction theory. In this approach the low-energy modification is predicted in the relative energy spectra of decay fragments interacting in the final state. This modification is related to the spectrum of fragments which defines pragmatic usage of the approach. The low-energy cross section in the corresponding channel (with energy E) is factorized as

$$d\sigma/dE \sim F_{FSI}(E) F_{PV}(E_T, E), \quad (1)$$

where F_{PV} is the “phase volume” contribution. In three-body decays with total decay energy E_T this term is

$$F_{PV}(E_T, E) = \sqrt{E(E_T - E)}. \quad (2)$$

The FSI term is obtained as

$$F_{FSI}(E) = \frac{1}{C_0^2} \frac{1}{2ME [\cot^2 \delta_l(E) + 1]}, \quad (3)$$

where M is the reduced mass in the channel of interest. The Coulomb penetration factor C_0 for $l = 0$ is defined via Sommerfeld parameter η

$$C_0^2(E) = \frac{2\pi\eta(E)}{\exp[2\pi\eta(E)] - 1}, \quad \eta(E) = \frac{Z_1 Z_2 \alpha}{\sqrt{2E/M}}, \quad (4)$$

and tends to unity in the case of neutral particles. For s -wave interaction of neutral particles the FSI term can be approximated in terms of the effective range approach as

$$F_{FSI}(E) = \frac{a^2}{1 + 2ME a^2}, \quad (5)$$

where a is an s -wave scattering length. Thus in the original Migdal-Watson approximation the low-energy modification of the cross section is sensitive to just one parameter: the scattering length in the channel of interest. For

emission of two neutrons such a behavior of relative energy distribution give rise to the notion of a “dineutron” particle as a specific object of research.

In spite of the fact that the “dineutron” idea is quite old there remain several aspects of theoretical importance, explored in current studies.

(a) $d(n, np)n$ reaction (and analogous reactions) as a tool to study n - n scattering length. There exists a problem of charge symmetry breaking for n - n and p - p channels (difference in the s -wave scattering lengths). Since it is very difficult to study neutron-neutron collisions directly, indirect methods have to be applied (e.g. Refs. [3–5] and Refs. therein).

(b) A Hanbury-Brown-Twiss “HBT interferometry”-like approach for high-energy collisions [6, 7]. This “femtoscscopy” approach allows to extract characteristics of the collision region, from which the emission of correlated particles is observed. It was suggested in Refs. [8–10] that an analogous “HBT interferometry”-like approach can be used for reactions with light exotic nuclei to extract the radial characteristics of a neutron halo. In this work we try to find out which information can actually be extracted in such studies.

(c) “Dineutron emission” in decays of light exotic nuclei. The even-neutron systems beyond the neutron dripline typically decay via direct emission of two neutrons. This process is sometimes discussed in terms of “dineutron emission”. The declared discovery of “dineutron emission” in decay of ^{16}Be has recently produced a heated discussion, see Refs. [11–13]. In this work we try to clarify this discussion by improved assessment of its theoretical constituents.

Some sources of current confusion in the discussion of a “dineutron” are as follows.

(i) It is important to note that we consider emission of a “dineutron” with *low* total decay energy E_T . However, it could be two physically very different sources: two-neutron *decays* of low-lying resonant states or *reactions* leading to population of low-lying three-body continuum. The formal description of these situations is very differ-

ent.

(ii) The “dineutron” is often described as a spatial correlation of two neutrons in the nuclear interior caused by the pairing interaction. It is often erroneously assumed that such a compact spatial configuration should exhibit itself as low-energy enhancement in the spectrum of two neutrons. This vision contradicts the uncertainty principle: a short-distance correlation should correspond to large relative momenta. So, the considerable large-momentum enhancement by spatial “dineutron” (caused by pairing) should be effectively overcome in the process of decay by the low-momentum enhancement (caused by the final state interaction). It appears that the issue of such an interplay defines applicability of the Migdal-Watson approximation and it is especially addressed in this work.

As an illustrating case of dineutron emission we have selected the ground state decay of ^{26}O . The latter has recently attracted considerable attention both experimental [14–17] and theoretical [18–24]. In our previous works the two-neutron emission from the ^{26}O g.s. has been studied in various theoretical approximations including sophisticated three-body decay and reaction models [19, 23, 25]. Generally, we find the dineutron approximation too poor and that complete three-body calculations (treating all pairwise final state interactions in the system on the same ground) are required to deal with three-body decays in all their complexity. However, using the limited model we explore two important tasks. (1) We attempt to clarify the question how the dineutron emission should look like if such a process takes place in reality for whatever reason. The results of these studies could be very discouraging for those who utilize this concept without sufficient theoretical background (2) We solve some methodological problems of our approach to three-body decays in fully controllable conditions. This helps to further validate our results concerning several complicated aspects of true $2n$ emission.

II. THEORETICAL APPROXIMATIONS

The discussion of “dineutron emission” is often lacking clarity because the object is loosely defined. For three models described in this Section it is defined which physical situation is considered in each case. The first two models described below are commonly used (or implied to be used), while the third model is developed in this work.

A. Trivial two-body dineutron emission

Let us consider emission of dineutron from a static source Φ with rms radius b_{nn} . The decay probability can in this case be defined via the outgoing flux

$$j_E = \frac{1}{M} \text{Im} \left[\Psi_E^{(+)\dagger}(\mathbf{r}) \nabla \Psi_E^{(+)}(\mathbf{r}) \right] \Big|_{r \rightarrow \infty},$$




FIG. 1. Neutro-neutron energy correlation for dineutron emission by a two-body source function. All curves are normalized to unity maximum value.

associated with wave function (WF)

$$\Psi_E^{(+)}(\mathbf{r}) = \int d^3r' G_E^{(+)}(\mathbf{r}; \mathbf{r}') \Phi(\mathbf{r}'). \quad (6)$$

For the source function Φ defined by first oscillator WF ϕ_{00}

$$\Phi(\mathbf{r}) = \sum_{lm} \frac{\phi_{nl}(r)}{r} Y_{lm}(\hat{r}),$$

$$\phi_{00}(r) = \frac{1}{b_{nn}^{3/2}} \left(\frac{54}{\pi} \right)^{1/4} r \exp\left(-\frac{3r^2}{4b_{nn}^2}\right), \quad (7)$$

one gets the nucleon-nucleon low-energy correlations shown in Fig. 1. The radius parameter b_{nn} in (7) is defined in such a way that it is equal to the source rms radius.

The simple nucleon-nucleon interaction is used in this work, acting only in the s -wave of two neutrons and defined by potential with Gaussian formfactor

$$V_{nn}(r) = V_{nn}^{(0)} \exp[-(r/r^{(0)})^2]. \quad (8)$$

For depth $V_{nn}^{(0)} = -31$ MeV and width $r^{(0)} = 1.8$ fm this potential produce the scattering length $a = -18.7$ fm. The Migdal-Watson expression provides the peak in the n - n energy correlation spectrum at about 115-120 keV (depending on the scattering length experimental uncertainty $a_s = 18.9 \pm 0.4$ fm). The peak produced for emission off a static source is somewhat different: it is located at somewhat lower energies of 70–100 keV for realistic “sizes” of the dineutron correlation of $b_{nn} = 3$ –7 fm. Also the shapes of the spectrum are quite sensitive to the radius parameter b_{nn} . Look for further discussion of this issue in Section III D.

This approximation provides some qualitative idea about what could be called “dineutron emission”. However, there exist unique situation in which the description of two-neutron emission by Eq. (6) becomes adequate, see the end in the next Subsection.

B. Static three-body dineutron model

Let us consider the emission of dineutron from a static source Φ consisting of two nucleon WFs occupying some orbital configurations:

$$\Phi_{JM}(\mathbf{r}_1, \mathbf{r}_2) = [\Phi(\mathbf{r}_1) \otimes \Phi(\mathbf{r}_2)]_{JM}. \quad (9)$$

This model we mention in the following as a static dineutron model (S2nM). After conversion of this source to Jacobi coordinates (so-called ‘‘T’’ system)

$$\mathbf{X} = \mathbf{r}_1 - \mathbf{r}_2, \quad \mathbf{Y} = (\mathbf{r}_1 + \mathbf{r}_2)/2, \quad (10)$$

the dineutron emission from this source can be treated exactly

$$\Psi_{E_T}^{(+)} = \frac{1}{\hat{T} + V_{nn} - E_T + i\epsilon} \Phi, \quad (11)$$

$$\Psi_{E_T, JM}^{(+)}(\mathbf{X}, \mathbf{Y}) = \int d^3X' d^3Y' G_{E_T}^{(+)}(\mathbf{X}, \mathbf{Y}; \mathbf{X}', \mathbf{Y}') \times \Phi_{JM}(\mathbf{X}', \mathbf{Y}'). \quad (12)$$

Here the three-body Green’s function can be given in a simple analytical form

$$G_{E_T}^{(+)}(\mathbf{X}, \mathbf{Y}; \mathbf{X}', \mathbf{Y}') = \frac{1}{2\pi i} \int dE_x G_{E_x}^{(+)}(\mathbf{X}; \mathbf{X}') \times G_{E_T - E_x}^{(+)}(\mathbf{Y}; \mathbf{Y}'), \quad (13)$$

where the Y variable Green’s function corresponds to plane wave propagation, while the X variable Green’s function incorporates the n - n final state interaction.

The model is called ‘‘static’’ in the sense that the properties of the source are totally decoupled from the properties of the final state interaction. The realistic scenario for such a model is sudden removal of the core from a two-nucleon halo system. This is not an improbable scenario for high-energy direct knockout reactions. For example, it was demonstrated in Ref. [26] that about 50% of the ${}^4\text{He}({}^6\text{He}, 2\alpha)$ cross section even at not very high beam energy of ~ 25 AMeV can be related to quasi-free knockout of the α core from the ${}^6\text{He}$ nucleus. In such a case the source function Φ_{JM} can be immediately related to the WF of the valence halo nucleons, paving way to studies of this WF structure.

It should be noted that there exists only one approximation in which the S2nM is reduced to the ‘‘trivial dineutron emission’’ of the previous Section. This is realized if the source function can be written in the factorized form:

$$\Phi(\mathbf{r}_1, \mathbf{r}_2) \equiv \Phi(\mathbf{X}) \Phi(\mathbf{Y}). \quad (14)$$

The one and only case when this is possible is when the $J = 0$ source is represented by two lowest s -wave oscillator WFs $\phi_{nl}(\mathbf{r})$

$$\Phi(\mathbf{r}_1, \mathbf{r}_2) \equiv \phi_{00}(\mathbf{r}_1) \phi_{00}(\mathbf{r}_2). \quad (15)$$

In this case all the information contained in the nucleon-nucleon momentum distribution is fully described by Eq. (6). This is exactly the situation considered in the applications of the HBT interferometry ideas to high-energy reactions: emission of independent particles from thermal source with Gaussian radial formfactor is formalized exactly by this model. For sources stemming from low-energy nuclear reactions this approximation is too poor because of variety of radial nucleon WFs deviating from Gaussian shapes and variety of angular momentum couplings defined by the investigated valence nucleon configurations.

C. Dynamic three-body dineutron model

In the case of the resonance state decays the S2nM can not be a reasonable approximation being associated with a certain reaction class. For resonant states in the limit of infinite lifetime the emission process should become totally insensitive to the population mechanism (as we have mentioned above the S2nM can be associated with certain reaction class). As an adequate dynamic approximation to the dineutron emission we now consider the following Dynamic Dineutron Model (D2nM). The decay of a three-body system is considered by solving a Schrödinger equation for WF $\Psi^{(+)}$ with pure outgoing wave boundary conditions and complex energy

$$(\hat{H}_3 - E_T + i\Gamma/2)\Psi_{E_T}^{(+)} = 0, \quad \hat{H}_3 = \hat{T}_3 + \hat{V}_3(\rho) + V_{nn}(r_{nn}). \quad (16)$$

The three-body Hamiltonian \hat{H}_3 contains kinetic energy term \hat{T}_3 , phenomenological three-body potential \hat{V}_3 depending on one variable, the hyperradius ρ , and nucleon nucleon potential V_{nn} .

To solve the three-body Schrödinger equation Eq. (16) we use the Hyperspherical Harmonics (HH) method and the iterative procedure developed in Ref. [27]. In the first step we use the hyperspherical harmonics method with ‘‘box’’ outgoing boundary conditions also defining the real part of the decay energy E_T :

$$(\hat{H}_3 - E_T)\Psi_{\text{box}} = 0.$$

Then the WF with outgoing asymptotic is derived solving the inhomogeneous equation

$$(\hat{H}_3 - E_T)\Psi_{E_T}^{(+)} = -(i\Gamma/2)\Psi_{\text{box}}$$

The obtained solution $\Psi_{E_T}^{(+)}$ may have problems with convergence, connected with effective ‘‘long-range’’ character of nucleon-nucleon interaction in s -wave. Near perfect work-around for such problems exists for simplified Hamiltonians, which include only one or two final state interactions and therefore there exist analytical Green’s function. In brief, we can rearrange Eq. (16) in the following way:

$$\Psi_{E_T}^{(+)} = -\frac{1}{\hat{T}_3 + V_{nn} - E_T + i\epsilon} \hat{V}_3(\rho) \Psi_{E_T}^{(+)}. \quad (17)$$

Here we have analytically known Green's function $\hat{G}_{E_T}^{(+)}$ of Eq. (13) which makes possible iterative improvement of the solution $\Psi_{E_T}^{(+)}$ providing the “corrected” WF $\Psi_{E_T, \text{corr}}^{(+)}$

$$\Psi_{E_T, \text{corr}}^{(+)} = -\hat{G}_{E_T}^{(+)} \hat{V}_3(\rho) \Psi_{E_T}^{(+)} . \quad (18)$$

Convergence of the procedure is guaranteed for the short-range potential $\hat{V}_3(\rho)$. There is also a simple criterion to check the consistency of the procedure: The resonant state widths and three-body momentum distributions obtained before and after some iteration of “correction” step should coincide.

III. WHAT AFFECTS DINEUTRON STRUCTURE?

A. Structure effects in D2nM

In this Section we try to isolate the internal nuclear structure effects on the dineutron emission. In D2nM we form the required structure by selection of the three-body potential $\hat{V}_3(\rho)$ to be different for hyperspherical components with different K values

$$\hat{V}_3(\rho) = \sum_K \frac{V_{3,K}}{1 + \exp[(\rho - a_\rho)/d_\rho]} \hat{P}_K . \quad (19)$$

Here a Woods-Saxon formfactor is selected for potential, while \hat{P}_K is projector on the states with definite K values.

We are considering presumably the lowest excitations with $J^\pi = 0^+$. In the proposed model the lowest energy three-body 0^+ WF has only one component with $L = 0$, $S = 0$, $l_x = 0$, $l_y = 0$, which corresponds to a dineutron in s -wave motion relative the core. The potential parameters used in the calculations are listed in Table I. The total decay energy E_T for each calculation is controlled just by the one running parameter V_3 .

The results are shown in Fig. 2. The upper panels illustrate the spatial correlations in the “T” Jacobi system (X is distance between two neutrons and Y is distance between n - n center of mass and heavy fragment). The selection of a structure strongly dominated by the $K = 0$, $K = 2$, or $K = 4$ component by potential in Table I, leads to corresponding population of very pure $[s^2]$, $[p^2]$, and $[d^2]$ quantum configurations. Domination of these

TABLE I. Depth parameters of the three-body potential $\hat{V}_3(\rho)$ in Eq. (19) which are used for calculations providing different dominant $[l^2]_0$ configurations. Geometry parameters $a_\rho = 4$ fm and $d_\rho = 0.8$ fm were also used.

Case	$V_{3,0}$	$V_{3,2}$	$V_{3,4}$
$[s^2]$	V_3	0	0
$[p^2]$	200	V_3	0
$[d^2]$	200	200	V_3

structures is clearly seen in Fig. 2 (a), (b), and (c) as presence of one, two, or three peaks of the WF in the internal region. Such correlation patterns are connected with Pauli principle and are often referred to as “Pauli focusing”, Ref. [28].

The energy correlations between two emitted neutrons, for WFs with corresponding internal structures, are illustrated in lower panels of Fig. 2. They are expressed in terms of fractional energy variable

$$\varepsilon = E_{nn}/E_T .$$

It is shown that for decay energies $E_T < 150$ keV the obtained correlations are relatively close to the three-body “phase volume”

$$d\Gamma/d\varepsilon \sim \sqrt{\varepsilon(1-\varepsilon)} .$$

Thus, for such decay energies the n - n FSI is not strong enough to noticeably modify the phase volume distribution. Only at about $E_T \sim 500$ keV does the correlation patterns begin to deviate considerably from the phase volume decay. At this and higher energies the structure effects are seen to play a dominant role.

For the n - n decay of the $[s^2]$ configuration the D2nM is providing expected results with explicit low-energy peak associated with n - n final state interaction. With energy increase this peak becomes sharper and sharper in the ε variable. However, if we plot the energy correlation in terms of real n - n relative energy E_{nn} , one can see in Fig. 3 that for decay energies $E_T > 150$ keV the E_{nn} peak position drifts slowly to higher energies. The peak position is stabilized at energies $E_{nn} \sim 80 - 90$ keV for $E_T \sim 500$ keV and depends only very weakly on E_T after that. To get the E_{nn} peak values above 100 keV, the decay energies E_T exceeding 5 MeV are required.

For the n - n decay of the $[p^2]$ configuration the energy evolution of the decay patterns is much more complicated. First at about $E_T \sim 500$ keV a kind of “antidineutron” is formed, providing peak at $\varepsilon > 0.5$ values. At around $E_T \sim 1$ MeV in addition to “antidineutron” an expected dineutron low- ε peak arise. At even higher energies $E_T > 2 - 3$ MeV the dineutron peak becomes the dominant feature of the spectrum, but integral intensities in the dineutron and antidineutron configurations is about equal. This evidently reflects the double-hump internal spatial configuration of the $[p^2]$ structure. So, we can conclude here that for decays with sufficiently high decay energies the n - n momentum distributions formed by n - n FSI can be used to extract information on the internal nuclear structure.

For the n - n decay of the $[d^2]$ configuration the most odd-looking results are obtained. It is clear that the correlation patterns for decay energies above $E_T \sim 1$ MeV tend to reflect the triple-hump configuration of the WF in the internal region. However, in contrast to the $[p^2]$ case no expressed low-energy n - n peak is obtained in the whole considered E_T domain.

We observe that, in contrast to common expectations, if we assume that the decay process is *totally governed* by




FIG. 2. Spatial correlations in the internal region are illustrated by the real part of the decay WF $\Psi_{E_T}^{(+)}$ for $E_T = 50$ keV. Panels (a), (b), and (c) correspond to calculations showing in the internal region dominant $[s^2]_0$, $[p^2]_0$, and $[d^2]_0$ configurations, respectively. Energy correlations between two neutrons for different total decay energies E_T are given in corresponding panels (d), (e), and (f). Gray lines show the three-body phase volume distribution. All surfaces and curves are normalized to unity maximum value.




FIG. 3. The energy distributions for relative energy between two neutrons E_{nn} are shown for different total decay energies E_T . For calculations with dominant $[s^2]_0$ configuration in nuclear interior, see Fig. 2 (a,d). All curves are normalized to unity maximum value.

n - n FSI, this does not mean that a simple picture with a single low-energy “dineutron” peak is obtained. The important prerequisite of the latter is dineutron emission from a $[s^2]$ configuration.

B. System size effect in D2nM

As we have mentioned above that an important motivation for n - n correlation studies was connected with the idea that spatial size of the emitting n - n configuration may be established here. As we have shown in Section III A the results for emission from $[p^2]$ and $[d^2]$ configurations contain a lot of information about structure and cannot be the right tool here. Hence we study this aspect of the model using example of $[s^2]$ configuration decay, demonstrating an easier way for interpretation of results.

To vary the nuclear system size we have performed calculations with three-body potential V_3 radius chosen to be strongly different from that in Table I Section III A. The real part of three-body WFs $\Psi_{E_T}^{(+)}$ obtained with $a_\rho = 3$ fm and $a_\rho = 7$ fm are shown in Fig. 4. It is clear that the radial extent of the nuclear system in the two considered cases is drastically different. The energy distributions between two neutrons associated with dineutron emission are provided in Fig. 5 for three different total decay energies E_T . We see that variation of the size of the emitting system practically do not effect the n - n correlations.




FIG. 4. Three-body WFs $\Psi^{(+)}$ (real part) calculated with different three-body potential width parameters a_ρ (equal 3 and 7 fm) and $E_T = 150$ keV. All surfaces are normalized to unity maximum value.




FIG. 5. Dineutron sensitivity to the size of the emitting 3-body system. The energy distributions between two neutrons for two different three-body potential size parameters a_ρ (equal 3 and 7 fm) and for different total decay energies E_T . All curves are normalized to unity maximum value.

C. System geometry effect in S2nM

The observation of the previous Subsection is in strong contrast to expectations. How it could be that the distance between neutrons in the emitting source do not effect the observed n - n correlations? It can, however, be understood recalling that in the method used for vari-




FIG. 6. Neutron-neutron energy correlation for static emission of dineutron from sources with different three-body T-system “geometries”. Two cases are illustrated: (a) $E_T = 0.5$ MeV, $\langle X \rangle = 3$ fm and (b) $E_T = 3$ MeV, $\langle X \rangle = 6$ fm, while the rms $\langle Y \rangle$ distance is varied. All curves are normalized to unity maximum value.

ation of the nuclear size we actually vary the ρ value for the whole system. This mean that we synchronously change both the mean sizes in X and Y coordinates. Let us consider analytical source function for static emission of dineutron which allows to vary the ratio $\langle X \rangle / \langle Y \rangle$:

$$\Phi(\mathbf{X}, \mathbf{Y}) = \Phi(\mathbf{X})\Phi(\mathbf{Y}), \quad (20)$$

where the radial functions $\Phi(\mathbf{r})$ are defined by Eq. (7). The calculation results are provided in Fig. 6 and they really demonstrate that even for emission off pure $[s^2]_0$ configuration broad variety of different energy distribution is possible. Here we have to conclude that in contrast with common expectations even for the emission off $[s^2]$ configuration the dineutron correlation is sensitive not to the mean distance $\langle X \rangle$ of the emitting source, but to the “geometry” of the source — the ratio of $\langle X \rangle$ and $\langle Y \rangle$.

D. Static emission of dineutron vs. Migdal-Watson approximation

The above calculations demonstrate broad variety of dineutron correlation patterns depending on emission conditions. A natural question here is: Why is the Migdal-Watson picture so widespread used as generic understanding of dineutron emission phenomena? The energy distributions obtained in the D2nM for $[s^2]$ case for different energies are also agree extremely well with Migdal-Watson results, see Fig. 7. What is the reason — is this type of correlation by necessity obtained for emission from a $[s^2]$ configuration? If we study systematically the correlation dependence on geometry of the source for static dineutron emission, the reason becomes clear. Fig. 6 shows examples of correlation evolution for systematic variation of the source geometry in S2nM. There is a broad variety of possible correlation pictures. We find, however, that for certain geometries, namely, for

$$\langle Y \rangle \lesssim \langle X \rangle \lesssim 2\langle Y \rangle, \quad (21)$$




FIG. 7. Energy distributions for relative energy E_{nn} between two neutrons calculated in D2nM, are given for different total decay energies E_T and compared with Migdal-Watson approximation (thin solid curves of the same color). The calculations correspond to dominant $[s^2]_0$ configuration in the nuclear interior. All curves are normalized to unity maximum value.

the correlations vary quite slowly and approach the Migdal-Watson results. It is clear that we can define a ratio of $\langle X \rangle$ and $\langle Y \rangle$ values such that S2nM results coincide with Migdal-Watson, see Fig. 8.

To interpret these results we should recall that for an independent particle model with two nucleons populating the same orbital configurations the condition $\langle X \rangle = 2\langle Y \rangle$ is satisfied (or what is the same, the average angle between two nucleons is equal to $\pi/2$). In reality the nucleon-nucleon interaction leads to formation of a closer configuration of two nucleons in the nuclear interior, which is also often referred to as “dineutron”. This leads to smaller $\langle X \rangle/\langle Y \rangle$ values compared to that in the independent particle model. The realistic values reside exactly in the range given above by Eq. (21). To illustrate this statement the calculated geometrical characteristics of some two-nucleon halo systems are provided in Table II. The geometry of continuum WFs obtained in the D2nM can be roughly estimated via the WF main peak position in the $\{X, Y\}$ plane, see Figs. 2 (a) and 4. It also satisfies the condition in Eq. (21).

We conclude that the Migdal-Watson approximation for dineutron emission (n - n FSI totally defines the decay dynamics) works nearly perfect when the $[s^2]_0$ source geometry is defined by formation of spatial “dineutron” correlation induced by pairing interactions in the internal nuclear region.

IV. DISCUSSION

A. General

The emission of two nucleons is often discussed in terms of a dominating “diproton” or “dineutron” decay




FIG. 8. Ratio between $\langle X \rangle$ and $\langle Y \rangle$ values at which the E_{nn} energy correlation for S2nM emission from a $[s^2]$ configuration coincides with one obtained in Migdal-Watson approximation. Solid and dotted curves correspond to total decay energies E_T equal 0.5 and 3 MeV respectively. The hatched region qualitatively corresponds to realistic relations between $\langle X \rangle$ and $\langle Y \rangle$ values. The nuclei mentioned in Table II are shown by blue diamonds.

mechanism. In this work we have tried to bring some clarity to the issue by constructing a model which allows to explicitly isolate effect of the nucleon-nucleon final state interaction. Based on the obtained results we can conclude that from theoretical formal point of view the common vision of “dineutron” as a low-energy enhancement in the the nucleon-nucleon energy distribution is not substantiated.

It seem that in the discussions of dinucleon emission there is some misunderstanding about relation of *necessary* and *sufficient* conditions. If we *observe* low-energy emission enhancement in the nucleon-nucleon spectrum this enhancement is evidently connected with N - N FSI. This condition can be regarded as necessary, because the huge scattering length in the N - N channel (~ 20 fm) means that in nuclear physics we do not have systems which can emit nucleons in such a way that they are outside the FSI range. This thing is unavoidable and

TABLE II. T-geometry of several bound three-cluster (e.g. two-nucleon halo) systems residing near the dripline obtained in the three-body cluster model calculations. The ${}^3\text{H}$ properties are trivially inferred from experimental data on charge radius.

Nucleus	Model	$\langle X \rangle$	$\langle Y \rangle$	$\langle X \rangle/\langle Y \rangle$	Ref.
${}^3\text{H}$	$n+n+p$	2.85	2.47	1.15	[29]
${}^6\text{He}$	${}^4\text{He}+n+n$	4.77	3.69	1.29	[30]
${}^{11}\text{Li}$	${}^9\text{Li}+n+n$	6.69	5.55	1.21	[31]
${}^{17}\text{Ne}$	${}^{15}\text{O}+p+p$	4.45	3.06	1.45	[32]
${}^{22}\text{C}$	${}^{20}\text{C}+n+n$	7.47	5.15	1.45	[33]

thus trivial. However, as we have shown in this work for various emission conditions, the presence of N - N FSI as the only factor governing two-nucleon emission does not lead to unique result (low-energy emission enhancement in the nucleon-nucleon spectrum). Even in the simplified dineutron theoretical model the major factor defining the nucleon-nucleon relative energy distributions in the final state are structure and spatial distributions in the internal region.

This result strongly discourages discussion of nucleon-nucleon correlations, observed in reactions and decays, in loosely defined terms such as a “diproton” or “dineutron” reaction mechanisms. In contrast it supports our confidence that comprehensive treatment of three-body decay mechanisms in all their complexity is a promising approach for extraction of information about nuclear interior and reaction mechanisms.

B. Lifetimes in the D2nM by example of ^{26}O

Here we consider how the lifetimes obtained in D2nM are compared with results of different decay models. This is illustrated by example of ^{26}O g.s. $2n$ decay, see Fig. 9.

The “direct decay model” estimates [25] assumes independent emission of nucleons from definite shell configurations. This model contains sensitivity to interactions in the core-nucleon channel, while the nucleon-nucleon FSI is neglected. The D2nM results provide the similar dependence of the decay width on energy in a broad energy range in the assumption about direct emission of nucleons off $[s^2]_0$ configuration. However, the decay is about one order of the magnitude faster in the case of D2nM. This is evidently connected to additional boost for $2n$ penetration due to n - n interaction in the subbarrier region. This observation is also consistent with results of $2p$ decay studies: the “diproton decay” estimates are providing the largest width values among all models, typically considerably overestimating widths relative to experiment [34].

Three-body model calculations of ^{26}O decay from Ref. [19] demonstrated strong sensitivity of width to details of core- n interactions, indicated by hatched area between red dotted curves in Fig. 9. It can be seen in the Figure that D2nM calculations with realistic assumption about $[d^2]$ structure of ^{26}O g.s. provide results consistent with complete three-body model calculations. So, application of the D2nM for lifetime estimates seems to be correct within an order of the magnitude.

C. Correlations in the decay of ^{26}O

Another issue for D2nM is how correlations between neutrons compare to results obtained in different models. This is illustrated by example of low-energy $2n$ decay of the ^{26}O ground state, see Fig. 10. This figure shows both the energy correlations for parameter $\varepsilon = E_{nn}/E_T$ and




FIG. 9. Lifetime of the ^{26}O g.s. in D2nM (black curves) for different structure assumptions are compared with direct decay model estimates of [25] (gray curves) and three-body calculations [19] (hatched area between red dashed curves).

angular correlations for angle θ_{nn} . The ε and θ_{nn} correlations are not independent and reflect the same type of correlation dynamics in different representations. For consistency with our previous works (e.g. Refs. [19, 34]) θ_{nn} is defined as the angle between momenta \mathbf{k}_{n_1} and $-\mathbf{k}_{n_2}$. It should be noted that angle θ_{nn} in Fig. 10 is defined as $\pi - \hat{\theta}_{nn}$, where $\hat{\theta}_{nn}$ is angle in [21, 35]. It can be seen that all the previous three-body model calculations [19, 21, 35] predict similar correlations behavior which can be interpreted as effective repulsion between neutrons in the final state (average angle between neutron emission directions is more than 90 degrees). In contrast the D2nM predicts small effective attraction: The peak in the energy distribution is shifted to slightly smaller ε values than for the “phase space” distribution shown for reference in Fig. 10 (a). So, for *correlations*, the “dineutron” assumption provides a qualitatively wrong trend in the case of low-energy ^{26}O g.s. decay.

D. Correlations in the decay of ^5H

D2nM calculations for decay of a $[s^2]$ configuration demonstrate nice agreement for n - n with Migdal-Watson approximation, see Fig. 7. Also we found that for low total decay energies $E_T > 150$ keV some kind of scaling behavior is obtained, see Fig. 3: The peak in the E_{nn} spectrum slowly drifts to higher energies with total decay energy E_T increase. It is interesting to note that analogous scaling behavior was observed in the studies of two-neutron decay of ^5H [36], see Fig. 11. The E_{nn} relative energy spectra were carefully reconstructed in this work for several decay energies of ^5H . However, what we see is that drift of the E_{nn} peak to higher energy in data




FIG. 10. Neutron-neutron correlations in the decay of ^{26}O g.s. in D2nM, $E_T = 150$ keV — black curve in panel (a), in three-body models of Refs. [19], $E_T = 75$ keV — red dashed curves in panels (a) and (b), [21], $E_T = 150$ keV — green dash-dotted curve in panel (b), and [35], $E_T = 150$ keV — blue dotted curve in panel (b). Energy distributions are normalized to unity maximum value; angular distributions are normalized for integration over $d\cos(\theta_{nn})$.

continues up to $E_T = 5$ MeV — the maximal energy obtained in this experiment. So, we see that E_T evolution of low-energy peak in Migdal-Watson approximation is strongly different from experimentally observed picture.

Some precaution is needed here because the majority of the mentioned spectrum is connected with the decay of excited states of ^5H expected to have $[p^2]_2$ orbital configuration. Within D2nM it is possible to study the decay specifically of this configuration. Compared to Migdal-Watson results (and data as well) the D2nM provides here the low-energy peak even at lower energies. The double-hump structure connected with decay of $[p^2]_2$ configuration and observed in the decay of ^5H is present in D2nM results. However, the calculated energy trend predicts enhancement of the large ε hump with energy E_T , while in experiment decrease was actually observed.

Thus none of the predicted “dineutron” trends is supported by the experimental data.

E. HBT-like approaches to n - n correlations

The neutron-neutron correlations in the “trivial dineutron” treatment of Section II A depend only on one parameter — the radial size of the neutron source. If we integrate the S2nM correlation spectrum (see Section II B) over the momentum connected with motion in the Y variable, we retain only the information about neutron-neutron relative distance [this is especially evident for the factorized source (14)]. This fact defines the use of intensity interferometry approach as a femtoscopy tool in high-energy physics. The idea to use the n - n correlations in the decays following reactions with exotic nuclei has two obstacles.

(i) For decay of higher shell configurations (such as $[p^2]$ and $[d^2]$) even such an integral correlation information cannot be straightforwardly related to radial character-




FIG. 11. The experimental n - n relative energy spectra for ^5H from Ref. [36] reconstructed for different total ^5H decay energies E_T . Thin solid lines of the same color in panel (a) show the Migdal-Watson approximation for the same energy. The D2nM calculations for decay of $[p^2]_2$ configuration are given in panel (b). All curves are normalized to unity maximum value.

istics of the source.

(ii) Technically the existing experimental setups are arranged in such a way that their acceptance for $2n$ events is drastically falling with total energy of two neutrons in the projectile frame. This fall is typically taking place in the energy range 1–3 MeV. For this reason we consider the two-neutron events with energy maximum E_T value of 3 MeV in this work. We have demonstrated that n - n correlations with such fixed total decay energy could be sensitive to structure, geometry, but not to radial size of the spatial n - n correlation.

These issues probably makes the interpretation of the neutron-neutron correlation data in Refs. [8–10] not quite consistent.

V. CONCLUSIONS

The dineutron emission is studied in this work in three different models, each with application to certain realistic scenario. The new development we introduce in this work is the Dynamic Dineutron Model (D2nM) which combines semirealistic internal structure for the nuclear interior with nucleon-nucleon interaction solely governing the emission process. This model is a subset of the

complete three-body problem which allows nice illustration of an efficiently isolated “dineutron emission” aspect of this problem. We argue that if we discuss the dineutron emission at all, this should be a formally correct realization of a theoretical description for such a process.

The results of this work require to critically reconsider several questions which have become a common place in discussions. In particular we have demonstrated the following:

- (i) The low-energy n - n correlation is typically a testing ground for indications of “dineutron emission”. We have to state that from a formal point of view a broad variety of “dineutron” correlation patterns is possible. A single low-energy peak, even within the simplified D2nM assumptions, should be the indication of emission from $[s^2]$ configuration strictly *with certain geometry*.
- (ii) We found that the idea to define the size of emitting region via n - n correlations, inherited from the HBT-like approach in high-energy physics, does not work for nuclear decays and reactions where sources have definite

shell structure and spin-parity. Even for emission from $[s^2]$ configuration, dineutron correlation is sensitive to the exact geometry of the internal WF (e.g. average angle between neutrons).

- (iii) We have given an illustrative explanation why the Migdal-Watson approach works well in the nuclear systems. It is shown that Migdal-Watson-like correlation patterns originate from $[s^2]$ configurations, where two neutrons are more focused in the space than in the independent particle case. This is a natural WF geometry effect of the attractive pairing interaction in a nucleus and thus such a spatial configuration is typically “pre-conditioned” for many processes of two-neutron emission.

ACKNOWLEDGMENTS

L.V.G. was partly supported by the Russian Science Foundation (grant No. 17-12-01367). We are grateful Prof. I.G. Mukha for useful comments.

-
- [1] K. Watson, Phys. Rev. C **88**, 1163 (1952).
 - [2] A. Migdal, Sov. Phys. JETP **1**, 2 (1955).
 - [3] A. Gardestig, Journal of Physics G: Nuclear and Particle Physics **36**, 053001 (2009).
 - [4] H. Witala and W. Glockle, Journal of Physics G: Nuclear and Particle Physics **37**, 064003 (2010).
 - [5] A. H. Couture, T. B. Clegg, S. Tajima, C. R. Howell, B. Fallin, J. H. Esterline, A. S. Crowell, B. J. Crowe, D. M. Markoff, L. C. Cumberbatch, R. S. Pedroni, and H. Witala, Phys. Rev. C **85**, 054004 (2012).
 - [6] Y. Sinyukov, R. Lednický, S. Akkelin, J. Pluta, and B. Erazmus, Physics Letters B **432**, 248 (1998).
 - [7] M. Lisa, S. Pratt, R. Soltz, and U. Wiedemann, Ann.Rev.Nucl.Part.Sci. **55**, 357 (2005).
 - [8] F. Marques, M. Labiche, N. Orr, J. Angélique, L. Axelsson, B. Benoit, U. Bergmann, M. Borge, W. Catford, S. Chappell, N. Clarke, G. Costa, N. Curtis, A. D’Arrigo, F. de Oliveira Santos, E. de Gies Brennand, O. Dorvaux, M. Freer, B. Fulton, G. Gardina, C. Gregori, S. Grevy, D. Guillemaud-Mueller, F. Hanappe, B. Heusch, B. Jonson, C. L. Brun, S. Leenhardt, M. Lewitowicz, M. Lopez, K. Markenroth, M. Motta, A. Mueller, T. Nilsson, A. Ninane, G. Nyman, I. Piqueras, K. Riisager, M. S. Laurent, F. Sarazin, S. Singer, O. Sorlin, and L. Stuttgé, Physics Letters B **476**, 219 (2000).
 - [9] F. M. Marqués, M. Labiche, N. A. Orr, J. C. Angélique, L. Axelsson, B. Benoit, U. C. Bergmann, M. J. G. Borge, W. N. Catford, S. P. G. Chappell, N. M. Clarke, G. Costa, N. Curtis, A. D’Arrigo, E. d. G. Brennand, F. d. O. Santos, O. Dorvaux, G. Fazio, M. Freer, B. R. Fulton, G. Gardina, S. Grévy, D. Guillemaud-Mueller, F. Hanappe, B. Heusch, B. Jonson, C. L. Brun, S. Leenhardt, M. Lewitowicz, M. J. López, K. Markenroth, A. C. Mueller, T. Nilsson, A. Ninane, G. Nyman, I. Piqueras, K. Riisager, M. G. Saint Laurent, F. Sarazin, S. M. Singer, O. Sorlin, and L. Stuttgé, Phys. Rev. C **64**, 061301 (2001).
 - [10] F.M.Marques, Few-Body Systems **31**, 145 (2002).
 - [11] A. Spyrou, Z. Kohley, T. Baumann, D. Bazin, B. A. Brown, G. Christian, P. A. DeYoung, J. E. Finck, E. Lunderberg, S. Mosby, W. A. Peters, A. Schiller, J. K. Smith, J. Snyder, M. J. Strongman, M. Thoennessen, and A. Volya, Phys. Rev. Lett. **108**, 102501 (2012).
 - [12] F. M. Marqués, N. A. Orr, N. L. Achouri, F. Delaunay, and J. Gibelin, Phys. Rev. Lett. **109**, 239201 (2012).
 - [13] A. Spyrou, Z. Kohley, T. Baumann, D. Bazin, B. A. Brown, G. Christian, P. A. DeYoung, J. E. Finck, N. Frank, E. Lunderberg, S. Mosby, W. A. Peters, A. Schiller, J. K. Smith, J. Snyder, M. J. Strongman, M. Thoennessen, and A. Volya, Phys. Rev. Lett. **109**, 239202 (2012).
 - [14] E. Lunderberg, P. A. DeYoung, Z. Kohley, H. Atanayake, T. Baumann, D. Bazin, G. Christian, D. Divaratne, S. M. Grimes, A. Haagsma, J. E. Finck, N. Frank, B. Luther, S. Mosby, T. Nagi, G. F. Peaslee, A. Schiller, J. Snyder, A. Spyrou, M. J. Strongman, and M. Thoennessen, Phys. Rev. Lett. **108**, 142503 (2012).
 - [15] C. Caesar, J. Simonis, T. Adachi, Y. Aksyutina, J. Alcantara, S. Altstadt, H. Alvarez-Pol, N. Ashwood, T. Aumann, V. Avdeichikov, M. Barr, S. Beceiro, D. Bemmerer, J. Benlliure, C. A. Bertulani, K. Boretzky, M. J. G. Borge, G. Burgunder, M. Caamano, E. Casarejos, W. Catford, J. Cederkäll, S. Chakraborty, M. Chartier, L. Chulkov, D. Cortina-Gil, U. Datta Pramanik, P. Diaz Fernandez, I. Dillmann, Z. Elekes, J. Enders, O. Ershova, A. Estrade, F. Farinon, L. M. Fraile, M. Freer, M. Freudenberger, H. O. U. Fynbo, D. Galaviz, H. Geissel, R. Gernhäuser, P. Golubev, D. Gonzalez Diaz, J. Hagdahl, T. Heftrich, M. Heil, M. Heine, A. Heinz, A. Henriques, M. Holl, J. D. Holt, G. Ickert, A. Ignatov, B. Jakobsson, H. T. Johansson, B. Jonson, N. Kalantar-Nayestanaki, R. Kanungo, A. Kelic-Heil, R. Knöbel, T. Kröll, R. Krücken, J. Kurcewicz, M. Labiche, C. Langer, T. Le Bleis, R. Lemmon,

- O. Lepyoshkina, S. Lindberg, J. Machado, J. Marganec, V. Marousov, J. Menéndez, M. Mostazo, A. Movsesyan, A. Najafi, T. Nilsson, C. Nociforo, V. Panin, A. Perea, S. Pietri, R. Plag, A. Prochazka, A. Rahaman, G. Rastrepina, R. Reifarth, G. Ribeiro, M. V. Ricciardi, C. Rigollet, K. Riisager, M. Röder, D. Rossi, J. Sanchez del Rio, D. Savran, H. Scheit, A. Schwenk, H. Simon, O. Sorlin, V. Stoica, B. Streicher, J. Taylor, O. Tengblad, S. Terashima, R. Thies, Y. Togano, E. Uberseder, J. Van de Walle, P. Velho, V. Volkov, A. Wagner, F. Wamers, H. Weick, M. Weigand, C. Wheldon, G. Wilson, C. Wimmer, J. S. Winfield, P. Woods, D. Yakorev, M. V. Zhukov, A. Zilges, M. Zoric, and K. Zuber (R3B collaboration), *Phys. Rev. C* **88**, 034313 (2013).
- [16] Z. Kohley, T. Baumann, G. Christian, P. A. DeYoung, J. E. Finck, N. Frank, B. Luther, E. Lunderberg, M. Jones, S. Mosby, J. K. Smith, A. Spyrou, and M. Thoennessen, *Phys. Rev. C* **91**, 034323 (2015).
- [17] Y. Kondo, T. Nakamura, R. Tanaka, R. Minakata, S. Ogoshi, N. A. Orr, N. L. Achouri, T. Aumann, H. Baba, F. Delaunay, P. Doornenbal, N. Fukuda, J. Gibelin, J. W. Hwang, N. Inabe, T. Isobe, D. Kameda, D. Kanno, S. Kim, N. Kobayashi, T. Kobayashi, T. Kubo, S. Leblond, J. Lee, F. M. Marqués, T. Motobayashi, D. Murai, T. Murakami, K. Muto, T. Nakashima, N. Nakatsuka, A. Navin, S. Nishi, H. Otsu, H. Sato, Y. Satou, Y. Shimizu, H. Suzuki, K. Takahashi, H. Takeda, S. Takeuchi, Y. Togano, A. G. Tuff, M. Vandebrouck, and K. Yoneda, *Phys. Rev. Lett.* **116**, 102503 (2016).
- [18] A. Volya and V. Zelevinsky, *Phys. Rev. C* **74**, 064314 (2006).
- [19] L. V. Grigorenko, I. G. Mukha, and M. V. Zhukov, *Phys. Rev. Lett.* **111**, 042501 (2013).
- [20] A. Volya and V. Zelevinsky, *Physics of Atomic Nuclei* **77**, 969 (2014).
- [21] K. Hagino and H. Sagawa, *Phys. Rev. C* **89**, 014331 (2014).
- [22] K. Hagino and H. Sagawa, *Phys. Rev. C* **90**, 027303 (2014).
- [23] L. V. Grigorenko and M. V. Zhukov, *Phys. Rev. C* **91**, 064617 (2015), [ArXiv:1503.03186].
- [24] K. Fosse, J. Rotureau, N. Michel, and W. Nazarewicz, *Phys. Rev. C* **96**, 024308 (2017).
- [25] L. V. Grigorenko, I. G. Mukha, C. Scheidenberger, and M. V. Zhukov, *Phys. Rev. C* **84**, 021303(R) (2011).
- [26] S. I. Sidorchuk, A. S. Fomichev, M. S. Golovkov, L. V. Grigorenko, V. A. Gorshkov, A. V. Gorshkov, S. A. Krupko, Y. T. Oganessian, A. M. Rodin, R. S. Slepnev, S. V. Stepantsov, G. M. Ter-Akopian, and R. Wolski, *Nucl. Phys. A* **840**, 1 (2010).
- [27] L. V. Grigorenko and M. V. Zhukov, *Phys. Rev. C* **76**, 014008 (2007).
- [28] B. Danilin, M. Zhukov, A. Korshennikov, V. Efros, and L. Chulkov, *Sov. J. Nucl. Phys.* **48**, 766 (1988), [*Yad. Fiz.* **48**, 1208 (1988)].
- [29] I. Angeli and K. Marinova, *Atomic Data and Nuclear Data Tables* **99**, 69 (2013).
- [30] M. V. Zhukov, B. Danilin, D. Fedorov, J. Bang, I. Thompson, and J.S.Vaagen, *Phys. Rep.* **231**, 151 (1993).
- [31] N. B. Shul'gina, B. Jonson, and M. V. Zhukov, *Nucl. Phys. A* **825**, 175 (2009).
- [32] L. Grigorenko, I. Mukha, and M. Zhukov, *Nuclear Physics A* **713**, 372 (2003), erratum *Nucl. Phys. A* **740** (2004) 401.
- [33] N. Shul'gina, *Phys. Rev. C*, to be submitted (2017).
- [34] M. Pfützner, M. Karny, L. V. Grigorenko, and K. Riisager, *Rev. Mod. Phys.* **84**, 567 (2012).
- [35] K. Hagino and H. Sagawa, *Few-Body Syst.* **57**, 185 (2016).
- [36] M. S. Golovkov, L. V. Grigorenko, A. S. Fomichev, S. A. Krupko, Y. T. Oganessian, A. M. Rodin, S. I. Sidorchuk, R. S. Slepnev, S. V. Stepantsov, G. M. Ter-Akopian, R. Wolski, M. G. Itkis, A. A. Bogatchev, N. A. Kondratiev, E. M. Kozulin, A. A. Korshennikov, E. Y. Nikolskii, P. Roussel-Chomaz, W. Mitig, R. Palit, V. Bouchat, V. Kinnard, T. Materna, F. Hanappe, O. Dorvaux, L. Stuttgé, A. A. Yurkhimchuk, V. V. Perevozchikov, Y. I. Vinogradov, S. K. Grishchkin, S. V. Zlatoustovskiy, V. Lapoux, R. Raabe, and L. Nalpas, *Phys. Rev. C* **72**, 064612 (2005).