

# ON BLOW-UP CONDITIONS FOR SOLUTIONS OF HIGHER ORDER DIFFERENTIAL INEQUALITIES

A.A. KON'KOV AND A.E. SHISHKOV

ABSTRACT. For differential inequalities of the form

$$\sum_{|\alpha|=m} (-1)^m \partial^\alpha a_\alpha(x, u) \geq b(x)|u|^\lambda \quad \text{in } \mathbb{R}^n, \quad n \geq 1,$$

where  $a_\alpha$  and  $b$  are some functions, we obtain conditions guaranteeing that any solution is identically equal to zero. We construct examples which show that the obtained conditions are sharp.

## 1. INTRODUCTION

We study solutions of the inequality

$$\sum_{|\alpha|=m} (-1)^m \partial^\alpha a_\alpha(x, u) \geq b(x)|u|^\lambda \quad \text{in } \mathbb{R}^n, \quad (1.1)$$

where  $n \geq 1$  and  $m \geq 1$  are some integers and  $\lambda$  is a real number. It is assumed that  $b$  is a positive measurable function and

$$|a_\alpha(x, \zeta)| \leq a(x)|\zeta| \quad (1.2)$$

with some positive measurable function  $a$  for almost all  $x = (x_1, \dots, x_n) \in \mathbb{R}^n$  and for all  $\zeta \in \mathbb{R}$  and  $|\alpha| = m$ . As is customary, by  $\alpha = (\alpha_1, \dots, \alpha_n)$  we mean a multi-index. In so doing,  $|\alpha| = \alpha_1 + \dots + \alpha_n$  and  $\partial^\alpha = \partial^{|\alpha|} / \partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}$ . Let us also denote by  $B_r$  an open ball in  $\mathbb{R}^n$  of radius  $r > 0$  centered at zero.

A function  $u$  is called a solution of inequality (1.1) if  $b(x)|u|^\lambda \in L_{1,loc}(\mathbb{R}^n)$  and  $a_\alpha(x, u) \in L_{1,loc}(\mathbb{R}^n)$  for all  $|\alpha| = m$  and, moreover,

$$\int_{\mathbb{R}^n} \sum_{|\alpha|=m} a_\alpha(x, u) \partial^\alpha \varphi \, dx \geq \int_{\mathbb{R}^n} b(x)|u|^\lambda \varphi \, dx \quad (1.3)$$

for any non-negative function  $\varphi \in C_0^\infty(\mathbb{R}^n)$ .

Our aim is to obtain conditions guaranteeing that every solution of (1.1) is trivial. Questions treated in our paper were studied earlier by a number of authors mainly for inequalities of the second order [1–11, 13–18]. Higher order inequalities have been studied much less. Until recently, the only effective method for their study was the method described in [6, 13]. However, this method loses its exactness in the case where the function  $1/b$  performs large oscillations in a neighborhood of infinity. In particular, it can not be applied to inequalities considered in Example 2.2.

---

1991 *Mathematics Subject Classification.* 35B44, 35B08, 35J30, 35J70.

*Key words and phrases.* Higher order differential inequalities, Nonlinear inequalities, Entire solutions.

The work of the second author is supported by the “RUDN University Program 5–100”.

## 2. MAIN RESULTS

**Theorem 2.1.** *Let  $\lambda > 1$  and*

$$\int_1^\infty r^{(m-n)\lambda+n-1} q(r) dr = \infty, \quad (2.1)$$

where

$$q(r) = \operatorname{ess\,inf}_{B_{\sigma r} \setminus B_{r/\sigma}} a^{-\lambda} b$$

for some real number  $\sigma > 1$ . Then any solution of (1.1) is trivial, i.e.  $u(x) = 0$  for almost all  $x \in \mathbb{R}^n$ .

**Theorem 2.2.** *Let  $\lambda > 1$  and (2.1) be valid, where*

$$q(r) = \left( \frac{1}{r^n} \int_{B_{\sigma r} \setminus B_{r/\sigma}} a^{\lambda/(\lambda-1)}(x) b^{-1/(\lambda-1)}(x) dx \right)^{1-\lambda} \quad (2.2)$$

for some real number  $\sigma > 1$ . Then any solution of (1.1) is trivial.

*Remark 2.1.* In relation (2.2), we do not exclude that

$$\int_{B_{\sigma r} \setminus B_{r/\sigma}} a^{\lambda/(\lambda-1)}(x) b^{-1/(\lambda-1)}(x) dx = \infty$$

for some real numbers  $r > 0$ . Since  $\lambda > 1$ , in this case, we have  $q(r) = 0$ .

Theorems 2.1 and 2.2 are proved in Section 3.

*Example 2.1.* Consider the inequality

$$(-1)^k \Delta^k u \geq b(x) |u|^\lambda \quad \text{in } \mathbb{R}^n, \quad (2.3)$$

where  $\lambda > 1$  and  $b$  is a positive measurable function such that

$$b(x) \geq b_0 |x|^l, \quad b_0 = \operatorname{const} > 0, \quad (2.4)$$

for almost all  $x$  from a neighborhood of infinity. According to Theorem 2.1, if

$$l \geq (n - 2k)\lambda - n, \quad (2.5)$$

then any solution of (2.3) is trivial. It can be seen that (2.5) coincides with the well-known blow-up condition given in [13, Example 5.2].

Now, assume that, in a neighborhood of infinity, the function  $b$  satisfy the inequality

$$b(x) \geq b_0 |x|^{(n-2k)\lambda-n} \log^\nu |x|, \quad b_0 = \operatorname{const} > 0. \quad (2.6)$$

In other words, we examine the critical exponent  $l = (n - 2k)\lambda - n$  in (2.4). If

$$\nu \geq -1, \quad (2.7)$$

then in accordance with Theorem 2.1 any solution of (2.3) is trivial.

In the case of  $n > 2k$ , condition (2.7) is exact. Really, let  $\nu < -1$ . We put

$$w_0(r) = (2 + r)^{-n} \log^{-(\nu+\lambda)/(\lambda-1)}(2 + r)$$

and

$$w_i(r) = \frac{1}{n-2} \int_0^r \left( \frac{\rho}{r} \right)^{n-2} \rho w_{i-1}(\rho) d\rho + \frac{1}{n-2} \int_r^\infty \rho w_{i-1}(\rho) d\rho, \quad i = 1, \dots, k.$$

For all  $1 \leq i \leq k$  we obviously have

$$-\left( \frac{d^2}{dr^2} + \frac{n-1}{r} \frac{d}{dr} \right) w_i = w_{i-1}$$

and

$$w_i(r) \sim r^{2i-n} \log^{-(1+\nu)/(\lambda-1)} r \quad \text{as } r \rightarrow \infty,$$

i.e.

$$c_1 r^{2i-n} \log^{-(1+\nu)/(\lambda-1)} r \leq w_i(r) \leq c_2 r^{2i-n} \log^{-(1+\nu)/(\lambda-1)} r$$

with some constants  $c_1 > 0$  and  $c_2 > 0$  for all  $r$  in a neighborhood of infinity. Thus, taking

$$u(x) = \varepsilon w_k(|x|), \quad (2.8)$$

where  $\varepsilon > 0$  is a sufficiently small real number, we obtain nontrivial solution of (2.3) with some positive function  $b$  for which (2.6) holds.

*Example 2.2.* In (2.3), let the positive measurable function  $b$  satisfy the inequality

$$b(x) \geq b_0 \sum_{i=1}^{\infty} \chi_{\omega_i}(x) |x|^{(n-2k)\lambda-n} \log^{\nu} |x|, \quad b_0 = \text{const} > 0, \quad (2.9)$$

for almost all  $x$  from a neighborhood of infinity, where  $\chi_{\omega_i}$  is the characteristic function of the set  $\omega_i = \{x \in \mathbb{R}^n : 2^{2i} < |x| < 2^{2i+1}\}$ , i.e.

$$\chi_{\omega_i}(x) = \begin{cases} 1, & x \in \omega_i, \\ 0, & x \in \mathbb{R}^n \setminus \omega_i. \end{cases}$$

In addition, let  $\lambda > 1$  and (2.7) be valid. Then, applying Theorem 2.1 with  $\sigma = 2^{1/4}$ , we obtain that any solution of (2.3) is trivial. As in Example 2.1, condition (2.7) is exact. Really, the right-hand side of (2.9) does not exceed the right-hand side of (2.6). Thus, in the case of  $\nu < -1$ , formula (2.8) provides us again with the desirable solution.

In general, let the positive function  $b$  satisfy the inequality

$$b(x) \geq \sum_{i=1}^{\infty} b_i \chi_{B_{2r_i} \setminus B_{r_i}}(x),$$

where  $b_i \geq 0$  and  $r_i > 0$  are some real numbers with  $r_{i+1} \geq 2r_i$ ,  $i = 1, 2, \dots$ . According to Theorem 2.1, if  $\lambda > 1$  and

$$\sum_{i=1}^{\infty} r_i^{(m-n)\lambda+n} b_i = \infty,$$

then any solution of (2.3) is trivial.

### 3. PROOF OF THEOREMS 2.1 AND 2.2

From now on we assume that  $u$  is a solution of (1.1) and, moreover,  $\lambda > 1$  and  $\sigma > 1$  are some real numbers. By  $C$  we mean various positive constants that can depend only on  $n$ ,  $m$ ,  $\lambda$ , and  $\sigma$ . Let us denote

$$J(r) = \int_{B_r} b(x) |u|^{\lambda} dx$$

and

$$h(r) = \left( \frac{1}{r^n} \int_{B_{\sqrt{\sigma}r} \setminus B_{r/\sqrt{\sigma}}} a^{\lambda/(\lambda-1)}(x) b^{-1/(\lambda-1)}(x) dx \right)^{1-\lambda},$$

where  $r$  runs over the set of positive real numbers. As in the case of the function  $q$ , it is assumed that  $h(r) = 0$  if

$$\int_{B_{\sqrt{\sigma}r} \setminus B_{r/\sqrt{\sigma}}} a^{\lambda/(\lambda-1)}(x) b^{-1/(\lambda-1)}(x) dx = \infty.$$

**Lemma 3.1.** *For arbitrary real numbers  $0 < r_1 < r_2$  such that  $r_2 \leq \sqrt{\sigma}r_1$  the following relationship holds:*

$$J(r_2) - J(r_1) \geq C(r_2 - r_1)^{\lambda m} r_1^{(1-\lambda)n} \sup_{(r_1, r_2)} h J^\lambda(r_1). \quad (3.1)$$

*Proof.* Let  $\varphi_0 \in C^\infty(\mathbb{R})$  be a non-decreasing function satisfying the conditions

$$\varphi_0|_{(-\infty, 0]} = 0 \quad \text{and} \quad \varphi_0|_{[1, \infty)} = 1.$$

We put

$$\varphi(x) = \varphi_0 \left( \frac{r_2 - |x|}{r_2 - r_1} \right).$$

It is easy to see that

$$\int_{\mathbb{R}^n} b(x) |u|^\lambda \varphi dx \geq \int_{B_{r_1}} b(x) |u|^\lambda dx$$

and

$$\left| \int_{\mathbb{R}^n} \sum_{|\alpha|=m} a_\alpha(x, u) \partial^\alpha \varphi dx \right| \leq \frac{C}{(r_2 - r_1)^m} \int_{B_{r_2} \setminus B_{r_1}} a(x) |u| dx.$$

Combining these estimates with (1.3), we obtain

$$\int_{B_{r_2} \setminus B_{r_1}} a(x) |u| dx \geq C(r_2 - r_1)^m \int_{B_{r_1}} b(x) |u|^\lambda dx. \quad (3.2)$$

If

$$\int_{B_{r_2} \setminus B_{r_1}} a^{\lambda/(\lambda-1)}(x) b^{-1/(\lambda-1)}(x) dx = \infty,$$

then (3.1) is obvious since  $h(r) = 0$  for all  $r \in (r_1, r_2)$ . Hence, we have to consider only the case where

$$\int_{B_{r_2} \setminus B_{r_1}} a^{\lambda/(\lambda-1)}(x) b^{-1/(\lambda-1)}(x) dx < \infty.$$

In this case, by the Hölder inequality, we have

$$\begin{aligned} \int_{B_{r_2} \setminus B_{r_1}} a(x) |u| dx &\leq \left( \int_{B_{r_2} \setminus B_{r_1}} a^{\lambda/(\lambda-1)}(x) b^{-1/(\lambda-1)}(x) dx \right)^{(\lambda-1)/\lambda} \\ &\quad \times \left( \int_{B_{r_2} \setminus B_{r_1}} b(x) |u|^\lambda dx \right)^{1/\lambda}. \end{aligned}$$

According to (3.2), this implies the estimate

$$\int_{B_{r_2} \setminus B_{r_1}} b(x)|u|^\lambda dx \geq C(r_2 - r_1)^{\lambda m} \left( \int_{B_{r_2} \setminus B_{r_1}} a^{\lambda/(\lambda-1)}(x)b^{-1/(\lambda-1)}(x) dx \right)^{1-\lambda} \\ \times \left( \int_{B_{r_1}} b(x)|u|^\lambda dx \right)^\lambda$$

which yields relationship (3.1) immediately.  $\square$

**Lemma 3.2.** *Let  $0 < r_1 < r_2$  be some real numbers such that  $r_2 \leq \sqrt{\sigma}r_1$  and  $2J(r_1) \leq J(r_2)$ . If  $J(r_1) > 0$ , then*

$$J^{(1/\lambda-1)/m}(r_1) - J^{(1/\lambda-1)/m}(r_2) \geq C \int_{r_1}^{r_2} r^{(1/\lambda-1)n/m} h^{1/(\lambda m)}(r) dr.$$

*Proof.* We construct a finite sequence of real numbers  $\{\rho_i\}_{i=0}^k$ . Let  $\rho_0 = r_1$ . Assume further that  $\rho_i$  is already defined. If  $4J(\rho_i) \geq J(r_2)$ , we put  $r_{i+1} = r_2$ ,  $k = i + 1$  and stop. Otherwise we take  $\rho_{i+1} \in (\rho_i, r_2)$  such that  $J(\rho_{i+1}) = 2J(\rho_i)$ . Since  $J$  is a continuous function, such a real number  $\rho_{i+1}$  obviously exists. It is also clear that this procedure must terminate at a finite step.

From the construction of the sequence  $\{\rho_i\}_{i=0}^k$ , it follows that

$$r_1 \leq \rho_i < \rho_{i+1} \leq r_2 \quad \text{and} \quad 2J(\rho_i) \leq J(\rho_{i+1}) \leq 4J(\rho_i), \quad i = 0, \dots, k-1. \quad (3.3)$$

By Lemma 3.1, for all  $0 \leq i \leq k-1$  we obtain

$$J(\rho_{i+1}) - J(\rho_i) \geq C(\rho_{i+1} - \rho_i)^{\lambda m} \rho_i^{(1-\lambda)n} \sup_{(\rho_i, \rho_{i+1})} h J^\lambda(\rho_i)$$

or, in other words,

$$\left( \frac{J(\rho_{i+1}) - J(\rho_i)}{J^\lambda(\rho_i)} \right)^{1/(\lambda m)} \geq C(\rho_{i+1} - \rho_i) \rho_i^{(1/\lambda-1)n/m} \sup_{(\rho_i, \rho_{i+1})} h^{1/(\lambda m)}. \quad (3.4)$$

Due to (3.3) the following inequalities hold:

$$J^{(1/\lambda-1)/m}(\rho_i) - J^{(1/\lambda-1)/m}(\rho_{i+1}) \\ \geq C \left( \frac{J(\rho_{i+1}) - J(\rho_i)}{J^\lambda(\rho_i)} \right)^{1/(\lambda m)}, \quad i = 0, \dots, k-1. \quad (3.5)$$

Moreover, since  $\rho \mapsto \rho^{(1/\lambda-1)/m}$  is a decreasing function on the interval  $(0, \infty)$ , it is obvious that

$$(\rho_{i+1} - \rho_i) \rho_i^{(1/\lambda-1)n/m} \sup_{(\rho_i, \rho_{i+1})} h^{1/(\lambda m)} \\ \geq \int_{\rho_i}^{\rho_{i+1}} r^{(1/\lambda-1)n/m} h^{1/(\lambda m)}(r) dr, \quad i = 0, \dots, k-1. \quad (3.6)$$

Combining (3.4), (3.5), and (3.6), we have

$$J^{(1/\lambda-1)/m}(\rho_i) - J^{(1/\lambda-1)/m}(\rho_{i+1}) \\ \geq C \int_{\rho_i}^{\rho_{i+1}} r^{(1/\lambda-1)n/m} h^{1/(\lambda m)}(r) dr, \quad i = 0, \dots, k-1.$$

Finally, summing the last inequalities over all  $0 \leq i \leq k-1$ , we complete the proof.  $\square$

**Lemma 3.3.** *Let  $0 < r_1 < r_2$  be some real numbers such that  $r_2 = \sqrt{\sigma}r_1$  and  $J(r_2) \leq 2J(r_1)$ . If  $J(r_1) > 0$ , then*

$$J^{1-\lambda}(r_1) - J^{1-\lambda}(r_2) \geq C \int_{r_1}^{r_2} r^{(m-n)\lambda+n-1} h(r) dr.$$

*Proof.* Lemma 3.1 implies the estimate

$$(J(r_2) - J(r_1))J^{-\lambda}(r_1) \geq C(r_2 - r_1)^{\lambda m} r_1^{(1-\lambda)n} \sup_{(r_1, r_2)} h. \quad (3.7)$$

By Lagrange's average value theorem, we obtain

$$J^{1-\lambda}(r_1) - J^{1-\lambda}(r_2) = (\lambda - 1)(J(r_2) - J(r_1))R^{-\lambda}.$$

for some real number  $R \in (J(r_1), J(r_2))$ . Since  $R \leq 2J(r_1)$ , this allows us to assert that

$$J^{1-\lambda}(r_1) - J^{1-\lambda}(r_2) \geq C(J(r_2) - J(r_1))J^{-\lambda}(r_1).$$

Taking into account the condition  $r_2 = \sqrt{\sigma}r_1$ , we also have

$$(r_2 - r_1)^{\lambda m} r_1^{(1-\lambda)n} \sup_{(r_1, r_2)} h \geq C \int_{r_1}^{r_2} r^{(m-n)\lambda+n-1} h(r) dr.$$

In virtue of (3.7) the last two inequalities readily yield the desirable estimate.  $\square$

**Lemma 3.4** (see [12, Lemma 2.6]). *Let  $0 < \alpha \leq 1$ ,  $\varkappa > 1$ ,  $\nu > 1$ ,  $r_1 > 0$ , and  $r_2 > 0$  be real numbers with  $r_2 \geq \nu r_1$ . Then*

$$\left( \int_{r_1}^{r_2} \psi^\alpha(r) dr \right)^{1/\alpha} \geq A \int_{r_1}^{r_2} r^{1/\alpha-1} \gamma(r) dr$$

for any measurable function  $\psi : [r_1, r_2] \rightarrow [0, \infty)$ , where

$$\gamma(r) = \operatorname{ess\,inf}_{(r/\varkappa, \varkappa r)} \psi$$

and  $A > 0$  is a constant depending only on  $\alpha$ ,  $\varkappa$ , and  $\nu$ .

*Proof of Theorem 2.2.* Assume the converse. Then there is a real number  $r_0 > 0$  such that  $J(r_0) > 0$ . Let us put  $r_i = \sigma^{i/2}r_0$ ,  $i = 1, 2, \dots$ . We denote by  $\Xi_1$  the set of non-negative integers  $i$  for which  $J(r_{i+1}) \geq 2J(r_i)$ . In so doing, let  $\Xi_2$  be the set of all other non-negative integers.

If  $i \in \Xi_1$ , then in accordance with Lemma 3.2 we obtain

$$J^{(1/\lambda-1)/m}(r_i) - J^{(1/\lambda-1)/m}(r_{i+1}) \geq C \int_{r_i}^{r_{i+1}} r^{(1/\lambda-1)n/m} h^{1/(\lambda m)}(r) dr. \quad (3.8)$$

In turn, if  $i \in \Xi_2$ , then Lemma 3.3 implies the estimate

$$J^{1-\lambda}(r_i) - J^{1-\lambda}(r_{i+1}) \geq C \int_{r_i}^{r_{i+1}} r^{(m-n)\lambda+n-1} h(r) dr. \quad (3.9)$$

Let us suppose that

$$\sum_{i \in \Xi_1} \int_{r_i}^{r_{i+1}} r^{(m-n)\lambda+n-1} q(r) dr = \infty, \quad (3.10)$$

where  $q$  is defined by (2.2). Then, summing inequalities (3.8) over all  $i \in \Xi_1$ , we obtain

$$J^{(1/\lambda-1)/m}(r_0) \geq C \sum_{i \in \Xi_1} \int_{r_i}^{r_{i+1}} r^{(1/\lambda-1)n/m} h^{1/(\lambda m)}(r) dr,$$

whence in virtue of  $\lambda m > 1$  it follows that

$$\begin{aligned} J^{1-\lambda}(r_0) &\geq C \left( \sum_{i \in \Xi_1} \int_{r_i}^{r_{i+1}} r^{(1/\lambda-1)n/m} h^{1/(\lambda m)}(r) dr \right)^{\lambda m} \\ &\geq C \sum_{i \in \Xi_1} \left( \int_{r_i}^{r_{i+1}} r^{(1/\lambda-1)n/m} h^{1/(\lambda m)}(r) dr \right)^{\lambda m}. \end{aligned} \quad (3.11)$$

Using Lemma 3.4 with  $\psi(r) = r^{(1-\lambda)n}h(r)$ ,  $\alpha = 1/(\lambda m)$ , and  $\varkappa = \nu = \sqrt{\sigma}$ , we obtain

$$\left( \int_{r_i}^{r_{i+1}} r^{(1/\lambda-1)n/m} h^{1/(\lambda m)}(r) dr \right)^{\lambda m} \geq C \int_{r_i}^{r_{i+1}} r^{(m-n)\lambda+n-1} q(r) dr$$

for all  $i \in \Xi_1$ . Hence, (3.11) implies the inequality

$$J^{1-\lambda}(r_0) \geq C \sum_{i \in \Xi_1} \int_{r_i}^{r_{i+1}} r^{(m-n)\lambda+n-1} q(r) dr$$

whose right-hand side is equal to infinity while the left-hand side is bounded. We obviously arrive at a contradiction. Therefore, (3.10) can not be valid, and due to condition (2.1) we have

$$\sum_{i \in \Xi_2} \int_{r_i}^{r_{i+1}} r^{(m-n)\lambda+n-1} q(r) dr = \infty. \quad (3.12)$$

Summing inequalities (3.9) over all  $i \in \Xi_2$ , one can conclude that

$$J^{1-\lambda}(r_0) \geq C \sum_{i \in \Xi_2} \int_{r_i}^{r_{i+1}} r^{(m-n)\lambda+n-1} h(r) dr.$$

Since  $h(r) \geq q(r)$  for all  $r \in (0, \infty)$ , this yields

$$J^{1-\lambda}(r_0) \geq C \sum_{i \in \Xi_2} \int_{r_i}^{r_{i+1}} r^{(m-n)\lambda+n-1} q(r) dr.$$

In virtue of (3.12) the last inequality again leads us to a contradiction.  $\square$

*Proof of Theorem 2.1.* We note that

$$\left( \frac{1}{r^n} \int_{B_{\sigma r} \setminus B_{r/\sigma}} a^{\lambda/(\lambda-1)}(x) b^{-1/(\lambda-1)}(x) dx \right)^{1-\lambda} \geq C \operatorname{ess\,inf}_{B_{\sigma r} \setminus B_{r/\sigma}} a^{-\lambda} b$$

for all  $r \in (0, \infty)$ . Thus, Theorem 2.1 follows immediately from Theorem 2.2.  $\square$

#### 4. GENERALIZATIONS

The condition of strict positivity of the function  $a$  can be omitted. Instead, we assume that

$$b(x) \geq a^\lambda(x) f(x)$$

with some non-negative function  $f \in L_{\infty, \text{loc}}(\mathbb{R}^n)$  for almost all  $x \in \mathbb{R}^n$ .

**Theorem 4.1.** *Let  $\lambda > 1$  and (2.1) be valid, where*

$$q(r) = \operatorname{ess\,inf}_{B_{\sigma r} \setminus B_{r/\sigma}} f$$

for some real number  $\sigma > 1$ . Then any solution of (1.1) is trivial.

**Theorem 4.2.** *Let  $\lambda > 1$  and (2.1) be valid, where*

$$q(r) = \left( \frac{1}{r^n} \int_{B_{\sigma r} \setminus B_{r/\sigma}} f^{-1/(\lambda-1)}(x) dx \right)^{1-\lambda}$$

for some real number  $\sigma > 1$ . Then any solution of (1.1) is trivial.

It can be seen that

$$\left( \frac{1}{r^n} \int_{B_{\sigma r} \setminus B_{r/\sigma}} f^{-1/(\lambda-1)}(x) dx \right)^{1-\lambda} \geq C \operatorname{ess\,inf}_{B_{\sigma r} \setminus B_{r/\sigma}} f$$

for all  $r \in (0, \infty)$ . Thus, Theorem 4.1 is a consequence of Theorem 4.2. In turn, to prove Theorem 4.2, it is sufficient to establish the validity of Lemma 3.1 with

$$h(r) = \left( \frac{1}{r^n} \int_{B_{\sqrt{\sigma}r} \setminus B_{r/\sqrt{\sigma}}} f^{-1/(\lambda-1)}(x) dx \right)^{1-\lambda}.$$

Really, in the case of

$$\int_{B_{r_2} \setminus B_{r_1}} f^{-1/(\lambda-1)}(x) dx = \infty,$$

Lemma 3.1 is evident since  $h(r) = 0$  for all  $r \in (r_1, r_2)$ . If

$$\int_{B_{r_2} \setminus B_{r_1}} f^{-1/(\lambda-1)}(x) dx < \infty,$$

then (3.2) implies the estimate

$$\int_{B_{r_2} \setminus B_{r_1}} f^{-1/\lambda}(x) b^{1/\lambda}(x) |u| dx \geq C(r_2 - r_1)^m \int_{B_{r_1}} b(x) |u|^\lambda dx. \quad (4.1)$$

By the Hölder inequality, we obtain

$$\begin{aligned} \int_{B_{r_2} \setminus B_{r_1}} f^{-1/\lambda}(x) b^{1/\lambda}(x) |u| dx &\leq \left( \int_{B_{r_2} \setminus B_{r_1}} f^{-1/(\lambda-1)}(x) dx \right)^{(\lambda-1)/\lambda} \\ &\quad \times \left( \int_{B_{r_2} \setminus B_{r_1}} b(x) |u|^\lambda dx \right)^{1/\lambda}, \end{aligned}$$

whence in accordance with (4.1) it follows that

$$\begin{aligned} \int_{B_{r_2} \setminus B_{r_1}} b(x) |u|^\lambda dx &\geq C(r_2 - r_1)^{\lambda m} \left( \int_{B_{r_2} \setminus B_{r_1}} f^{-1/(\lambda-1)}(x) dx \right)^{1-\lambda} \\ &\quad \times \left( \int_{B_{r_1}} b(x) |u|^\lambda dx \right)^\lambda. \end{aligned}$$

This obviously proves (3.1).

Finally, note that, instead of (1.2), we can consider the inequality

$$|a_\alpha(x, \zeta)| \leq a(x) |\zeta|^p,$$

where  $p > 0$  is some real number, as was done in [13]. However, this does not increase the generality since we can always use the replacement  $v = |u|^p \operatorname{sign} u$ .



## REFERENCES

- [1] A. Farina, J. Serrin, Entire solutions of completely coercive quasilinear elliptic equations, *J. Diff. Eq.* 250 (2011) 4367–4408.
- [2] A. Farina, J. Serrin, Entire solutions of completely coercive quasilinear elliptic equations, II, *J. Diff. Eq.* 250 (2011) 4409–4436.
- [3] R. Filippucci, P. Pucci, M. Rigoli, Non-existence of entire solutions of degenerate elliptic inequalities with weights, *Arch. Ration. Mech. Anal.* 188 (2008) 155–179; Erratum, 188 (2008) 181.
- [4] R. Filippucci, Nonexistence of positive entire weak solutions of elliptic inequalities, *Nonlinear Anal.* 70 (2009) 2903–2916.
- [5] R. Filippucci, P. Pucci, M. Rigoli, On entire solutions of degenerate elliptic differential inequalities with nonlinear gradient terms, *J. Math. Anal. Appl.* 356 (2009) 689–697.
- [6] V. A. Galaktionov, E. L. Mitidieri, S. I. Pohozaev, Blow-up for higher-order parabolic, hyperbolic, dispersion and Schroedinger equations, *Monographs and Research Notes in Mathematics*, CRC Press, Boca Raton, FL, 2014.
- [7] V. A. Galaktionov, J. L. Vazquez, Blow-up of a class of solutions with free boundaries for the Navier-Stokes equations, *Adv. Diff. Eq.* 4 (1997) 291–321.
- [8] V. A. Galaktionov, J. L. Vazquez, Necessary and sufficient conditions of complete blow-up and extinction for one-dimensional quasilinear heat equations, *Arch. Ration. Mech. Anal.* 129 (1996) 225–244.
- [9] J.B. Keller, On solution of  $\Delta u = f(u)$ . *Comm. Pure Appl. Math.* 10 (1957) 503–510.
- [10] V.A. Kondratiev, E.M. Landis, Qualitative properties of the solutions of a second-order nonlinear equations, *Mat. Sb.* 135 (1988) 346–360.
- [11] A.A. Kon'kov, On properties of solutions of quasilinear second-order elliptic inequalities, *Nonlinear Analysis* 123–124 (2015), 89–114.
- [12] A.A. Kon'kov, On solutions of non-autonomous ordinary differential equations, *Izv. Math.* 65 (2001) 285–327.
- [13] E. Mitidieri, S.I. Pohozaev, A priori estimates and blow-up of solutions to nonlinear partial differential equations and inequalities, *Proc. V.A. Steklov Inst. Math.* 234 (2001) 3–383.
- [14] M. Marcus, A.E. Shishkov, Fading absorption in non-linear elliptic equations, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 30 (2013) 315–336.
- [15] Y. Naito, H. Usami, Entire solutions of the inequality  $\operatorname{div}(A(|Du|)Du) = f(u)$ , *Math. Z.* 225 (1997) 167–175.
- [16] Y. Naito, H. Usami, Nonexistence results of positive entire solutions for quasilinear elliptic inequalities, *Canad. Math. Bull.* 40 (1997) 244–253.
- [17] R. Osserman, On the inequality  $\Delta u \geq f(u)$ , *Pacific J. Math.* 7 (1957) 1641–1647.
- [18] L. Veron, Comportement asymptotique des solutions d'équations elliptiques semi-linéaires dans  $\mathbb{R}^n$ , *Ann. Math. Pure Appl.* 127 (1981) 25–50.

DEPARTMENT OF DIFFERENTIAL EQUATIONS, FACULTY OF MECHANICS AND MATHEMATICS,  
MOSCOW LOMONOSOV STATE UNIVERSITY, VOROBYOVY GORY, MOSCOW, 119992 RUSSIA  
*E-mail address:* `konkov@mech.math.msu.su`

RUDN UNIVERSITY, MIKLUKHO-MAKLAYA STR. 6, MOSCOW, 117198 RUSSIA; INSTITUTE  
OF APPLIED MATHEMATICS AND MECHANICS OF NAS OF UKRAINE, DOBROVOL'SKOGO STR.  
1, SLAVYANSK, 84116 UKRAINE  
*E-mail address:* `aeshkv@yahoo.com`