

# Log-aesthetic Curves as Similarity Geometric Analogue of Euler's Elasticae

Jun-ichi INOBUCHI

Institute of Mathematics, University of Tsukuba  
Tsukuba 305-8571, Japan  
e-mail: inoguchi@math.tsukuba.ac.jp

Kenji KAJIWARA

Institute of Mathematics for Industry, Kyushu University  
744 Motoooka, Fukuoka 819-0395, Japan  
e-mail: kaji@imi.kyushu-u.ac.jp

Kenjiro T. MIURA

Graduate School of Science and Technology, Shizuoka University  
3-5-1 Johoku, Hamamatsu, Shizuoka, 432-8561, Japan  
e-mail: miura.kenjiro@shizuoka.ac.jp

Masayuki SATO

Serio Corporation  
2-21-10 Shinkawa, Chuo-ku, Tokyo 104-0033, Japan

Wolfgang K. SCHIEF

School of Mathematics and Statistics, The University of New South Wales  
Sydney, NSW 2052, Australia  
e-mail: w.schief@unsw.edu.au

Yasuhiro SHIMIZU

UEL Corporation  
1-1-1 Toyosu, Koto-ku, Tokyo 135-8560, Japan

## Abstract

In this paper we consider the log-aesthetic curves and their generalization which are used in CAGD. We consider those curves under similarity geometry and characterize them as stationary integrable flow on plane curves which is governed by the Burgers equation. We propose a variational formulation of those curves whose Euler-Lagrange equation yields the stationary Burgers equation. Our result suggests that the log-aesthetic curves and their generalization can be regarded as the similarity geometric analogue of Euler's elasticae.

## 1 Introduction

In this paper we consider a class of plane curves in CAGD called the *log-aesthetic curves* (LAC) and their generalization called the *quasi aesthetic curves* (qAC), and present their new mathematical characterization by using the theory of integrable systems. In 1744, Euler introduced the elastic energy of a plane curve which is the totality of squared signed curvature, and studied a variational problem of plane curves in Euclidean geometry [4]. The solutions to this problem are called *elasticae*. On the other hand, a standard isoperimetric deformation of arc length parametrized plane curves is governed by the *modified Korteweg de Vries (mKdV) equation* for the signed curvature [5], which is one of the most important integrable equations. Then it was seen that elasticae are

characterized as travelling wave solutions to the mKdV equation. In this paper, we consider LAC and qAC in the framework of *similarity geometry*, and characterize them as the similarity geometric analogue of elasticae. We formulate them as stationary curves with respect to the simplest integrable flow given by the *Burgers equation*. We then give a variational formulation by introducing a functional (called the fairing energy) for plane curves, from which we deduce the stationary Burgers equation as the Euler-Lagrange equation. This gives a new mathematical characterization of LAC and qAC and demonstrates the usefulness of Klein geometry and integrable systems as tools of CAGD.

## 2 Elasticae and mKdV flows

To clarify the mathematical background and motivations of this paper, we recall briefly here the basic facts on the elasticae and the Goldstein-Petrich flows. An arc length parametrized curve  $\gamma(s)$  in the Euclidean plane  $\mathbb{E}^2$  is said to be an *elastica* if it is a critical point of the *elastic energy*:

$$E(\gamma) = \int_0^\ell \frac{1}{2} \kappa(s)^2 ds, \quad (1)$$

through total length preserving variations. Here  $s$  is the arc length parameter and  $\kappa(s)$  is Euclidean (signed) curvature. We consider the smooth plane curves  $\gamma : [0, \ell] \rightarrow \mathbb{E}^2$  with two endpoints and tangent vectors at the ends being fixed. Then the Euler-Lagrange equation obtained from the variation of the curve is

$$2\kappa'' + \kappa^3 - \lambda\kappa = 0, \quad (2)$$

where  $\lambda$  is a constant and  $' = d/ds$ . For more information on the elasticae, we refer to [1, 16].

On the other hand, we consider the isoperimetric deformation of plane curves of the form (*Goldstein-Petrich flow* [5], see also [8]):

$$\dot{\gamma} = -\kappa' N^E - \frac{1}{2} \kappa^2 T^E.$$

Here  $\dot{\phantom{x}} = \partial/\partial t$ ,  $T^E = \gamma'$  and  $N^E = JT^E$ , where  $J$  is the positive  $\pi/2$ -rotation, are the unit tangent vector field and the unit normal vector field. The Frenet frame  $F^E = (T^E, N^E) \in \text{SO}(2)$  satisfies

$$(F^E)^{-1} \dot{F}^E = \kappa J, \quad (F^E)^{-1} \dot{F}^E = -\left(\kappa'' + \frac{1}{2} \kappa^3\right) J, \quad (3)$$

where the first equation is the Frenet formula. The compatibility condition of (3) yields the mKdV equation

$$\dot{\kappa} + \kappa''' + \frac{3}{2} \kappa^2 \kappa' = 0. \quad (4)$$

In the terminology of integrable systems, (3) is nothing but a *Lax pair* (for vanishing spectral parameter). The travelling wave solutions  $\kappa(s - \lambda t/2)$  to the mKdV equation satisfy  $2\kappa'' + \kappa^3 - \lambda\kappa = c_0$  for some constant  $c_0 \in \mathbb{R}$ . Comparing with (2), we see that every elastica is a solution to the Goldstein-Petrich flow.




Figure 1: Log-aesthetic curves. Left: LAC with various parameters. Right: LAC with  $\alpha = 2$  (circle involute curve).

### 3 Log-aesthetic curves and similarity geometry

According to [11], an arc length parametrized plane curve  $\gamma(s) \in \mathbb{R}^2$  is said to be an LAC of slope  $\alpha$  if its signed curvature radius  $q$  satisfies

$$q(s)^\alpha = as + b \quad (\alpha \neq 0), \quad q(s) = \exp(as + b) \quad (\alpha = 0), \quad a, b \in \mathbb{R}. \quad (5)$$

The class of LAC's includes some well known plane curves, for instance, the logarithmic spiral ( $\alpha = 1$ ), the clothoid ( $\alpha = -1$ ) and the *Nielsen spiral* ( $\alpha = 0$ ). Figure 1 illustrates some examples of LAC.

Recent advancements on the LAC have been promising as indicated by Levien and Séquin [9]. Theoretical backgrounds have been set up, such as variational formulation of LAC for free-form surface design [12], and fast computational algorithm [19]. An important development for industrial design practices has been presented in [10], where it is proved that a unique solution exists for the  $G^1$  interpolation problem using an LAC segment when  $\alpha < 0$ . For more details of the LAC we refer to [13, 14].

Those developments have been made based on the basic characterization (5) in the framework of Euclidean geometry. However, (5) is too simple to identify the underlying geometric structure. Consequently, we do not have a “good” guideline to generate a larger class of “aesthetic” geometric objects including LAC based on a sound mathematical background. In this paper, we clarify that LAC fits well the framework of the similarity geometry.

The similarity plane geometry is a Klein plane geometry associated with the group of similarity transformations, *i.e.*, isometries and scalings:  $\mathbb{R}^2 \ni \mathbf{p} \mapsto rA\mathbf{p} + \mathbf{b}$ ,  $A \in \text{SO}(2)$ ,  $r \in \mathbb{R}_+$ ,  $\mathbf{b} \in \mathbb{R}^2$ . The natural parameter of plane curves in similarity geometry is the *turning angle*  $\theta = \int \kappa(s) ds$ . Let  $\gamma(\theta)$  be a plane curve in similarity geometry parametrized by  $\theta$ . The *similarity tangent vector field*  $T$  and the *similarity normal vector field*  $N$  are defined by  $T = \gamma'$ ,  $N = J\gamma'$ , respectively, where  $' = d/d\theta$ . Then  $T$  and  $N$  satisfy the *similarity Frenet equations*:

$$T' = -uT + N, \quad N' = -T - uN, \quad (6)$$

for some function  $u(\theta)$  which is called the *similarity curvature*. One can check that plane curves in similarity geometry are uniquely determined by the similarity curvature up to similarity transformations. The similarity curvature  $u$  is related to the signed curvature radius  $q = 1/\kappa$  by the

*Cole-Hopf transformation:*

$$u = -\frac{q'}{q}. \quad (7)$$

Surprisingly, the notion of LAC is shown to be invariant under similarity transformations. For instance, the slope  $\alpha$  is expressed as  $\alpha = 1 + u'/u^2$ . LAC and qAC are reformulated in terms of similarity geometry as follows [6, 17, 18]: A plane curve  $\gamma(\theta)$  is said to be a LAC of slope  $\alpha$  if its similarity curvature satisfies the Bernoulli equation:

$$u' = (\alpha - 1)u^2. \quad (8)$$

Similarly, a plane curve is said to be a qAC of slope  $\alpha$  if its similarity curvature obeys the Riccati equation:

$$u' = (\alpha - 1)u^2 + c, \quad c \in \mathbb{R} : \text{const.} \quad (9)$$

## 4 Burgers flows on similarity plane curves

It is known that plane curves in similarity geometry admit isogonal deformations governed by the Burgers hierarchy [3]. The simplest flow is given by

$$\dot{\gamma} = (b - u)T - N, \quad b \in \mathbb{R} : \text{const.} \quad (10)$$

The similarity Frenet frame  $F = (T, N)$  satisfies the similarity Frenet equation and the deformation equation

$$F^{-1}F' = -uI + J, \quad F^{-1}\dot{F} = (-u' + u^2 + 1 - bu)I + bJ, \quad (11)$$

where  $I$  is the identity matrix. The compatibility condition of (11) yields the *Burgers equation*:

$$\dot{u} = (u' - u^2 + bu)'. \quad (12)$$

The Burgers equation is linearized in terms of the signed curvature radius via the Cole-Hopf transformation (7) as  $\dot{q} = q'' + bq'$ . We note that the parameter  $b$  corresponds to reparametrization of the curve. Imposing the *stationary ansatz*  $\dot{u} = 0$ , the Burgers equation (12) reduces to the Riccati equation:

$$u' = u^2 - bu + c, \quad c \in \mathbb{R}. \quad (13)$$

We recover the Riccati equation (9) with  $\alpha = 2$  by putting  $b = 0$ . Note that (9) is obtained formally from (13) by replacing the variable as  $u \rightarrow (\alpha - 1)u$ . In this sense, qAC is characterized as the stationary curves of the Burgers flow.

## 5 Fairing energy in similarity geometry

Let us discuss the variational formulation of LAC and qAC. We introduce the *fairing energy functional*  $\mathcal{F}^{\lambda,a}$  by

$$\mathcal{F}^{\lambda,a}(\gamma) = \int_{\theta_1}^{\theta_2} \frac{1}{2} \left\{ a^2 u(\theta)^2 + \lambda \left( \frac{q_1 q_2}{q(\theta)^2} \right)^a \right\} d\theta, \quad (14)$$

where  $a = \alpha - 1$ ,  $\lambda$  is an arbitrary constant and  $q_i = q(\theta_i)$  ( $i = 1, 2$ ). The above functional is invariant under similarity transformations and its name ‘‘fairing energy’’ is motivated from the fairing procedure in digital style design of industrial products. We write the variation of  $\gamma$  as  $\delta\gamma = \xi(\theta)T(\theta) + \eta(\theta)N(\theta)$ . Standard variational calculus yields  $\delta q/q = \phi - \psi'$ ,  $\delta u/u = -(u\psi' + \phi - \psi')$  and  $\delta\theta = \psi$ , where  $\phi(\theta) = \xi' - \xi u - \eta$ ,  $\psi(\theta) = \eta' - \eta u + \xi$ . By using those data, we obtain the *first variational formula* of the fairing energy:

$$\delta\mathcal{F}^{\lambda,a}(\gamma) = -\frac{1}{2} \left[ a^2 u(\tilde{\phi} - \tilde{\psi}') + H(\gamma)\tilde{\psi} \right]_{\theta_1}^{\theta_2} + \frac{a}{2} \int_{\theta_1}^{\theta_2} \left\{ au' - \lambda \left( \frac{q_1 q_2}{q^2} \right)^a \right\} (\tilde{\phi} - \tilde{\psi}' + u\tilde{\psi}) d\theta, \quad (15)$$

where  $\tilde{\phi} = \phi - \frac{\phi(\theta_1) + \phi(\theta_2)}{2}$ ,  $\tilde{\psi} = \psi - \frac{\psi(\theta_1) + \psi(\theta_2)}{2}$ , and  $H(\gamma) = a^2 u(\theta)^2 - \lambda \left( \frac{q_1 q_2}{q^2} \right)^a$ . The first variation formula implies that if  $\gamma$  is a critical point of the fairing energy for deformations which respect the boundary condition in (15), then  $\gamma$  satisfies

$$au' - \lambda \left( \frac{q_1 q_2}{q^2} \right)^a = 0, \quad (16)$$

which is equivalent to the Riccati equation for qAC (9) together with (7). Note that  $H(\gamma)$  is a first integral of (16). If we require preservation of the total similarity arc length  $\delta(\theta_2 - \theta_1) = 0$ , then the boundary term vanishes iff  $\phi(\theta_1) = \phi(\theta_2)$ . This implies that  $\frac{q_2}{q_1}$ , namely, the ratio of length of tangent vectors at the endpoints is preserved by the variation. Note that this condition is invariant with respect to similarity transformations.

**Theorem 5.1.** *If a plane curve  $\gamma$  is a critical point of the fairing energy  $\mathcal{F}^{\lambda,a}$  (14) under the assumption of preservation of total similarity arc length and the boundary condition that the ratio of length of tangent vectors at the endpoints is preserved, then the similarity curvature  $u$  satisfies  $u' = au^2 + c$ , where  $c$  is a constant. Therefore, quasi aesthetic curves of slope  $\alpha \neq 1$  are critical points of the fairing functional.*

**Remark 5.2.** Moreton and Séquin [15] considered the functional  $\int_{\gamma} \left( \frac{dk}{ds} \right)^2 ds \cdot \left( \int_{\gamma} ds \right)^3$ , which is invariant under similarity transformations. A curve which is a critical point of this functional is called a *scale-invariant minimum variation curve* (SI-MVC) in [2]. This is different from (14), since the above functional is rewritten as  $\left( \int \frac{u(\theta)^2}{q(\theta)^3} d\theta \right) \left( \int_{\theta_1}^{\theta_2} q(\theta)^3 d\theta \right)$ .

The mathematical characterization of LAC and qAC presented in this paper may enable us to generalize the notion of aesthetic curves. For instance, it may be possible to construct discrete LAC and qAC preserving underlying integrable structure, which may be useful for a numerical approach. Associated results will be reported in a forthcoming paper. It is noted that discrete isogonal deformations of discrete plane curves governed by the discrete Burgers equation have been studied in [7].

## Acknowledgment

This work was partially supported by JSPS KAKENHI, JP25289021, JP16H03941, JP16K13763, JP15K04834, JP26630038, JST RISTEX Service Science, Solutions and Foundation Integrated

Research Program, ImPACT Program of the Council for Science, Technology and Innovation. The authors acknowledge the support by 2016 IMI Joint Use Program Short-term Joint Research No. 20160016. Y. Shimizu acknowledges the support by 2015 IMI Joint Use Program Short-term Visiting Researcher No.20150010. The authors would like to express their sincere gratitude to Prof. Miyuki Koiso, Prof. Hiroyuki Ochiai, Prof. Nozomu Matsuura and Prof. Sampei Hirose for invaluable comments and fruitful discussions.

## References

- [1] Bryant R., Griffiths, P., 1986, Reductions for constrained variational problems and  $\int \kappa^2/2ds$ . Amer. J. Math. **108**(3), 525—570.
- [2] Brook A., Bruckstein A.M., Kimmel, R., 2005, On similarity-invariant fairness measures. in: International Conference on Scale-Space Theories in Computer Vision, Lecture Notes in Computer Sciences **3459** , pp. 456–467.
- [3] Chou, K.-S., Qu, C.-Z., 2002, Integrable equations arising from motions of plane curves. Phys. D **162** 9–33.
- [4] Euler, L., 1744, Methodus inveniendi lineas curvas maximi minimive proprietate gaudentes; Additamentum I: de curvis elasticis, Lausanne and Geneva. English translation: Oldfather, W.A., Ellis, C.A, Brown, D.M., 1933, Leonhard Euler’s elastic curves. Isis **20**(1), 72–160 (<http://www.jstor.org/stable/224885>).
- [5] Goldstein, R., Petrich, D.M., 1991, The Korteweg-de Vries hierarchy as dynamics of closed curves in the plane. Phys. Rev. Lett. **67** (23), 3203–3206.
- [6] Inoguchi, J., 2016, Attractive plane curves in differential geometry. in: Mathematical Progress in Expressive Image Synthesis III. Mathematics for Industry, vol 24, Springer, pp. 121–135.
- [7] Kajiwara, K., Kuroda, T., Matsuura, N., 2016, Isogonal deformation of discrete plane curves and discrete Burgers hierarchy. Pac. J. Math. Ind. **8**, Article Number 3, 14 pages.
- [8] Lamb Jr. G.L., 1976, Solitons and the motion of helical curves. Physical Review Letters **37**, 235–237.
- [9] Levien, R.; C H.Sequin, C.H., 2009, Interpolating splines: Which is the fairest of them all? Comput.-Aided Design and Appl., **6**(1), 91–102.
- [10] Meek, D.S., Saito, T., Walton, D.J., Yoshida, Y., 2012, Planar Two-point G1 Hermite Interpolating Log-aesthetic Spirals. J. Comput. Appl. Math. **236**(17), 4485-4493.
- [11] Miura, K.T., 2006, A general equation of aesthetic curves and its self-affinity. Comput.-Aided Design Appl. **3**(1-4) , 457–464.
- [12] Miura, K.T., Shirahata, R., Agari, S., Usuki, S., Gobithaasan, R.U., 2012, Variational Formulation of the Log-aesthetic Surface and Development of Discrete Durface Filters. Comput.-Aided Design Appl., **9**(6), 901–914.

- [13] Miura, K.T., Gobithaasan R.U., 2014, Aesthetic Curves and Surfaces in Computer Aided Geometric Design, *Int. J. of Automation Technol.* **8**(3), 304–316.
- [14] Miura K.T., Gobithaasan R.U, 2016, Aesthetic Design with Log-Aesthetic Curves and Surfaces. *Mathematical Progress in Expressive Image Synthesis III, Mathematics for Industry*, vol 24, Springer, pp.107–119.
- [15] Moreton, H.P., Séquin, C.H., 1993, Scale-invariant minimum-cost curves: fair and robust design implements. *Computer Graphics Forum* **12**(3) Eurographics '93 (R. J. Hubbard and R. Juan, eds), C473–C484.
- [16] Mumford, D., 1994, Elastica and computer vision. in: Bajaj C.L. ed., *Algebraic Geometry and its Applications*, Springer Verlag, pp. 491–506.
- [17] Sato M., Shimizu, Y., 2015, Log-aesthetic curves and Riccati equations from the viewpoint of similarity geometry. *JSIAM Letters* **7** , 21–24.
- [18] Sato, M., Shimizu, Y., 2016, Generalization of log-aesthetic curves by Hamiltonian formalism. *JSIAM Letters* **8**, 49–52.
- [19] Ziatdinov, R., Yoshida, N., Kim, T., 2012, Analytic parametric equations of log-aesthetic curves in terms of incomplete gamma functions, *Comput. Aided Geom. Design* **29**(2), 129–140.