A note for global existence of a two-dimensional chemotaxis-haptotaxis model with remodeling of non-diffusible attractant

Jiashan Zheng *

School of Mathematics and Statistics Science, Ludong University, Yantai 264025, P.R.China

Abstract

In this paper, we study the following the coupled chemotaxis-haptotaxis model with remodeling of non-diffusible attractant

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) - \xi \nabla \cdot (u \nabla w) + \mu u (1 - u - w), \\ v_t = \Delta v - v + u, \\ w_t = -vw + \eta w (1 - u - w), \end{cases}$$
(0.1)

in a bounded smooth domain $\Omega \subseteq \mathbb{R}^2$ with zero-flux boundary conditions, where χ , ξ and η are positive parameters. Under appropriate regularity assumptions on the initial data (u_0, v_0, w_0) , by developing some L^p -estimate techniques, we prove the global existence and uniqueness of classical solutions when $\mu > 0$, where μ is the logistic growth rate of cancer cells. This result removes the additional restriction of μ is sufficiently large in [32] (J. Diff. Eqns., 263(2)(2017), 1269–1292) for the global existence of solutions.

Key words: Chemotaxis-haptotaxis; Global existence; Non-diffusible attractant2010 Mathematics Subject Classification: 92C17, 35K55, 35K59, 35K20

^{*}Corresponding author. E-mail address: zhengjiashan2008@163.com (J.Zheng)

1 Introduction

The oriented movement of biological cells or organisms in response to a chemical gradient is called chemotaxis (see Calvez and Carrillo [2], Fontelos et al. [8], Hillen and Painter [13], Horstmann [15, 16], Jäger and Luckhaus [20], Kavallaris and P. Souplet [21], Nagai [30], Perthame [33], Sherratt [35], Winkler [51]). To describe chemotaxis of cell populations, the signal is produced by the cells, in 1970, Keller and Segel (see [22]) proposed an important variant of the quasilinear chemotaxis model

$$\begin{cases} u_t = \nabla \cdot (\phi(u)\nabla u) - \chi \nabla \cdot (u\nabla v), & x \in \Omega, t > 0, \\ v_t = \Delta v + u - v, & x \in \Omega, t > 0. \end{cases}$$
(1.1)

The interesting feature of quasilinear Keller–Segel types of models (1.1) is the possibility of blow-up of solutions in finite time, which strongly depends on the space dimension (see e.g. Horstmann et al. [15, 17], Rascle and Ziti [34]). In fact, solutions of (1.1) may blow up in finite time when $N \geq 2$ (Herrero and Velázquez [11], Osaki et al. [31], Winkler [52]). In the higher-dimensional case when $N \geq 3$, small total mass of cells appears to be insufficient to rule out blow-up in (Winkler et al. [51, 18]). Some recent studies show that the large nonlinear diffusion function (see Ishida et al. [19], Tao and Winkler [42], Zheng [56, 60]), the nonlinear chemotactic sensitivity function (see Fujie et al. [9]) and the (generalized) logistic growth term (see Lankeit [24], Winkler et al. [47, 50, 53], Zheng [55, 60]) may prevent the blow-up of solutions. One important extension of the classical Keller–Segel model to a more complex cell migration mechanism was proposed by Chaplain and Lolas (see Chaplain and Lolas [4]) in order to describe processes of cancer invasion. In 2006, Chaplain and Lolas ([4]) described the process of cancer invasion on the macroscopic scale by the chemotaxishaptotaxis system (with remodeling of non-diffusible attractant)

$$u_{t} = \Delta u - \chi \nabla \cdot (u \nabla v) - \xi \nabla \cdot (u \nabla w) + \mu u (1 - u - w), \quad x \in \Omega, t > 0,$$

$$\tau v_{t} = \Delta v + u - v, \quad x \in \Omega, t > 0,$$

$$w_{t} = -vw + \eta w (1 - u - w), \quad x \in \Omega, t > 0,$$

$$\frac{\partial u}{\partial \nu} - \chi u \frac{\partial v}{\partial \nu} - \xi \frac{\partial w}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0, \quad x \in \partial\Omega, t > 0,$$

$$u(x, 0) = u_{0}(x), \tau v(x, 0) = \tau v_{0}(x), w(x, 0) = w_{0}(x), \quad x \in \Omega,$$

(1.2)

where $\tau > 0, \eta \ge 0, \Omega \subseteq \mathbb{R}^N, N \ge 1$ is the physical domain which we assume to be bounded with smooth boundary, χ , ξ , μ and η measure the chemotactic sensitivities and haptotactic sensitivities, the proliferation rate of the cells and the remodeling rate of the extracellular matrix (ECM), respectively. Here the unknown quantities u = u(x,t), v = v(x,t) and w = w(x,t) denote the density of cancer cells, the concentration of enzyme and the density of healthy tissue, respectively.

The model (1.2) accounts for both chemotactic migration of cancer cells towards a diffusible matrix-degrading enzyme (MDE) secreted by themselves, and haptotactic migration towards a static tissue, also referred to as ECM (Chaplain et al. [5, 10], Liotta and T. Clair [26]). This on the one hand opens new fields of applications to modeling approaches in the style pursued by Keller and Segel (see [22]), but on the other hand it gives rise to new mathematical challenges due to more involved couplings.

If $\chi = 0$, the PDE system (1.2) becomes the haptotaxis-only system (with remodeling of non-diffusible attractant)

$$\begin{cases} u_t = \Delta u - \xi \nabla \cdot (u \nabla w) + \mu u (1 - u - w), & x \in \Omega, t > 0, \\ v_t = \Delta v + u - v, & x \in \Omega, t > 0, \\ w_t = -vw + \eta w (1 - u - w), & x \in \Omega, t > 0. \end{cases}$$
(1.3)

Global existence and asymptotic behavior of solutions to (1.3) have been investigated in [6, 7, 48, 27, 46, 29, 36] and [37] for the case $\eta = 0$ and $\eta > 0$, respectively.

When $\eta = 0$, the PDE system (1.2) is reduced to the chemotaxis-haptotaxis system

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) - \xi \nabla \cdot (u \nabla w) + \mu u (1 - u - w), & x \in \Omega, t > 0, \\ \tau v_t = \Delta v + u - v, & x \in \Omega, t > 0, \\ w_t = -vw, & x \in \Omega, t > 0. \end{cases}$$
(1.4)

When $\tau = 0$ denotes that the diffusion rate of the MDE is much greater than that of cancer cells (see Chaplain and Lolas [4], Winkler et al. [1, 43]). In [40], Tao and Wang proved that model (1.4) possesses a unique global bounded classical solution for any $\mu > 0$ in two space dimensions, and for large $\mu > 0$ in three space dimensions; Tao and Winkler ([44]) studied global boundedness for model (1.4) with the condition $\mu > \frac{(N-2)^+}{N}\chi$, furthermore, they gave

the exponential decay of w in the large time limit for the additional explicit smallness on w_0 ; While, if $\tau = 1$ in (1.2), Tao and Wang ([39]) proved that model (1.2) possesses a unique global-in-time classical solution for any $\chi>0$ in one space dimension, and for small $\frac{\chi}{\mu}>0$ in two and three space dimensions; Tao ([36]) improved the result of [39] for any $\mu > 0$ in two space dimension; Additionally, recent studies have shown that the solution behavior can be also impacted by the nonlinear diffusion (see Tao and Winkler [41], Wang et al. [28, 49, 59]) and the (generalized) logistic damping (see Cao [3], Hillen et al. [14], Zheng et al. [58, 61]). Compared with the chemotaxis-only system, chemotaxis-only system and the chemotaxis-haptotaxis system, the coupled chemotaxis-haptotaxis system with remodeling of non-diffusible attractant ($\eta > 0$ in (1.2)) is much less understood (Chaplain and Lolas [4], Pang and Wang [32], Tao and Winkler [45]). The main technical difficulty in their proof stems from the effects of the strong coupling in (1.2) on the spatial regularity of u, v and w when $\eta > 0$. When $\eta = 0$, one can build a one-sided pointwise estimate which connects Δw to v (see Lemma 2.2 of [3] or (3.10) of [49]). Relying on such a pointwise estimate, we can derive two useful energy-type inequalities that bypass $\int_{\Omega} u^{p-1} \nabla \cdot (u \nabla w)$ (see Lemma 3.2 of |59|). Using such information along with coupled estimate techniques and the boundedness of the $\|\nabla v(\cdot, t)\|_{L^2(\Omega)}$, we establish estimates on $\int_{\Omega} u^p + |\nabla v|^{2q}$ for any p and q > 1 (see Lemmata 3.3 and 3.4 of [59]), which results in the boundedness of u in $L^{\infty}(\Omega)$ by using the standard regularity theory of parabolic equation and performing the Moser iteration procedure (see Lemma 3.5 of [59]). However, for the model (1.2) with $\eta > 0$, one needs to estimate the chemotaxis-related integral term $\int_{\Omega} a^p |\nabla v|^2 dx$ (see (3.28) in [45]) or $\int_{\Omega} e^{-(p+1)(t-s)} a^p |\nabla v|^2 dx ds$ (see (3.8) of [32]) with $a := u e^{-\xi w}$, which proves to be much more technically demanding. In [32], assuming that $\mu > \xi \eta \max\{\|u_0\|_{L^{\infty}(\Omega)}, 1\} + \mu^*(\chi^2, \xi)$ (the hypothesis can not be dropped (see the proof of Lemma 3.2 of [32])), Pang and Wang showed that the problem (1.2) admits a unique global solution $(u, v, w) \in (C^{2,1}(\bar{\Omega} \times (0, \infty)))^3$. Moreover, u is bounded in $\Omega \times (0, \infty)$. However, to the best of our knowledge, it is still an open problem to determine whether or not in the case N = 2 some unbounded solutions may exist in (1.2) with small $\mu > 0$. Indeed, as pointed by [1] (see also [38]), the hypothesis on $\mu > 0$ may yield the classical global solution. So, it is natural to ask whether the solution is globally existence when $\mu > 0$. In this paper, we give a positive answer to this question.

Motivated by the aforementioned papers, the purpose of this work is to establish global solvability of (1.2). Our main result in this respect reads as follows.

Theorem 1.1. Let $\tau > 0, \chi > 0, \xi > 0$ and $\eta > 0$. Assume that $\Omega \subseteq \mathbb{R}^2$ is a bounded domain with smooth boundary and the initial data (u_0, v_0, w_0) is supposed to satisfy the following conditions

$$u_{0} \in C^{2+\vartheta}(\bar{\Omega}) \quad \text{with} \quad u_{0} \geq 0 \quad \text{in} \quad \Omega \quad \text{and} \quad \frac{\partial u_{0}}{\partial \nu} = 0 \quad \text{on} \quad \partial\Omega,$$

$$v_{0} \in C^{2+\vartheta}(\bar{\Omega}) \quad \text{with} \quad v_{0} \geq 0 \quad \text{in} \quad \Omega \quad \text{and} \quad \frac{\partial v_{0}}{\partial \nu} = 0 \quad \text{on} \quad \partial\Omega,$$

$$w_{0} \in C^{2+\vartheta}(\bar{\Omega}) \quad \text{with} \quad w_{0} \geq 0 \quad \text{in} \quad \bar{\Omega} \quad \text{and} \quad \frac{\partial w_{0}}{\partial \nu} = 0 \quad \text{on} \quad \partial\Omega$$

$$(1.5)$$

with some $\vartheta \in (0,1)$. If $\mu > 0$, then there exists a triple $(u, v, w) \in (C^{2,1}(\bar{\Omega} \times (0,\infty)))^3$ which solves (1.2) in the classical sense. Moreover, u and v are bounded in $\Omega \times (0,\infty)$.

Remark 1.1. (i) If $w \equiv 0$, (the PDE system (1.2) is reduced to the chemotaxis-only system), it is not difficult to obtain that the solutions under the conditions of Theorem 1.1 are uniformly bounded when N = 2, which coincides with the results of Osaki et al. ([31]).

(ii) From Theorem 1.1, we derive that solutions of model (1.2) are global and bounded for any $\eta = 0, \mu > 0$ and $N \leq 2$, which coincides with the result of Tao ([38]).

Without loss of generality, we may assume $\tau = 1$ in (1.2), since, for $\tau > 0$ can be proved very similarly.

The plan of this paper is as follows. In Section 2, we give some basic results and some preliminary lemmata as a preparation for the arguments in the later sections. In Section 3, firstly, by using the technical lemma (Lemma 3.2) and employing the variation-of-constants formula, we may establish the boundedness of $\int_{\Omega} a^{q_0}(q_0 > 1)$, where $a = ue^{-\xi w}$. In addition, we shall involve the variation-of-constants formula and L^p -estimate techniques to gain the boundedness of $\int_{\Omega} a^p(p > 1)$. Finally, using the Alikakos–Moser iteration, we finally established the $L^{\infty}(\Omega)$ bound of a (see the proof of Theorem 1.1).

2 Preliminaries

Before formulating our main results, we first recall some preliminary lemmas used throughout this paper. To begin with, let us collect some basic solution properties which essentially have already been used in [18] (see also Winkler [51], Zhang and Li [54]).

Lemma 2.1. ([18]) For $p \in (1, \infty)$, let $A := A_p$ denote the sectorial operator defined by

$$A_p u := -\Delta u \quad \text{for all} \quad u \in D(A_p) := \{ \varphi \in W^{2,p}(\Omega) | \frac{\partial \varphi}{\partial \nu} |_{\partial \Omega} = 0 \}.$$

$$(2.1)$$

The operator A + 1 possesses fractional powers $(A + 1)^{\alpha} (\alpha \ge 0)$, the domains of which have the embedding properties

$$D((A+1)^{\alpha}) \hookrightarrow W^{1,p}(\Omega) \quad if \quad \alpha > \frac{1}{2}.$$
 (2.2)

If $m \in \{0,1\}$, $p \in [1,\infty]$ and $q \in (1,\infty)$ with $m - \frac{N}{p} < 2\alpha - \frac{N}{q}$, then we have

$$||u||_{W^{m,p}(\Omega)} \le C ||(A+1)^{\alpha}u||_{L^q(\Omega)} \quad for \ all \ u \in D((A+1)^{\alpha}),$$
 (2.3)

where C is a positive constant. The fact that the spectrum of A is a p-independent countable set of positive real numbers $0 = \mu_0 < \mu_1 < \mu_2 < \cdots$ entails the following consequences: For all $1 \le p < q < \infty$ and $u \in L^p(\Omega)$ the general L^p - L^q estimate

$$\|(A+1)^{\alpha}e^{-tA}u\|_{L^{q}(\Omega)} \leq ct^{-\alpha-\frac{N}{2}(\frac{1}{p}-\frac{1}{q})}e^{(1-\mu)t}\|u\|_{L^{p}(\Omega)}$$
(2.4)

for any t > 0 and $\alpha \ge 0$ with some $\mu > 0$.

In deriving some preliminary estimates for v, we shall make use of following the property referred to as a variation of Maximal Sobolev Regularity (see e.g. Theorem 3.1 of [12] or [3]).

Lemma 2.2. Suppose $\gamma \in (1, +\infty)$, $g \in L^{\gamma}((0, T); L^{\gamma}(\Omega))$. Let v be a solution of the following initial boundary value

$$\begin{cases} v_t - \Delta v + v = g, \quad (x,t) \in \Omega \times (0,T), \\ \frac{\partial v}{\partial \nu} = 0, \quad (x,t) \in \partial \Omega \times (0,T), \\ v(x,0) = v_0(x), \quad (x,t) \in \Omega. \end{cases}$$
(2.5)

Then there exists a positive constant C_{γ} such that if $s_0 \in [0,T)$, $v(\cdot,s_0) \in W^{2,\gamma}(\Omega)(\gamma > N)$ with $\frac{\partial v(\cdot,s_0)}{\partial \nu} = 0$, then,

$$\int_{s_0}^T e^{\gamma s} \|v(\cdot,t)\|_{W^{2,\gamma}(\Omega)}^{\gamma} ds \le C_{\gamma} \left(\int_{s_0}^T e^{\gamma s} \|g(\cdot,s)\|_{L^{\gamma}(\Omega)}^{\gamma} ds + e^{\gamma s_0} (\|v_0(\cdot,s_0)\|_{W^{2,\gamma}(\Omega)}^{\gamma}) \right).$$
(2.6)

Proof. Letting $c(x,s) = e^s v(x,s)$. Then we derive that c satisfies

$$\begin{cases} c_s(x,s) - \Delta c(x,s) = f(x,s), & (x,s) \in \Omega \times (0,T), \\ \frac{\partial c}{\partial \nu} = 0, & (x,s) \in \partial \Omega \times (0,T), \\ c(x,0) = v_0(x), & (x,s) \in \Omega, \end{cases}$$
(2.7)

where $f(x,s) = e^s g(x,s)$. Applying the Maximal Sobolev Regularity (see e.g. Theorem 3.1 of [12]) to c, we derive that

$$\int_{0}^{T} \|\Delta c(\cdot, s)\|_{L^{\gamma}(\Omega)}^{\gamma} ds + \int_{0}^{T} \|c(\cdot, s)\|_{L^{\gamma}(\Omega)}^{\gamma} ds + \int_{0}^{T} \|c_{s}(\cdot, s)\|_{L^{\gamma}(\Omega)}^{\gamma} ds \\
\leq C_{1,\gamma} \left(\int_{0}^{T} \|f(\cdot, s)\|_{L^{\gamma}(\Omega)}^{\gamma} ds + (\|c_{0}\|_{L^{\gamma}(\Omega)}^{\gamma} + \|\Delta c_{0}\|_{L^{\gamma}(\Omega)}^{\gamma}) \right).$$
(2.8)

Substituting v into the above inequality and changing the variables imply

$$\int_{0}^{T} e^{\gamma s} (\|v(\cdot,t)\|_{L^{\gamma}(\Omega)}^{\gamma} + \|\Delta v(\cdot,t)\|_{L^{\gamma}(\Omega)}^{\gamma}) ds
\leq C_{1,\gamma} \left(\int_{0}^{T} e^{\gamma s} \|g(\cdot,s)\|_{L^{\gamma}(\Omega)}^{\gamma} ds + (\|c_{0}\|_{L^{\gamma}(\Omega)}^{\gamma} + \|\Delta c_{0}\|_{L^{\gamma}(\Omega)}^{\gamma}) \right).$$
(2.9)

On the other hand, by the elliptic L^p -estimate,

$$\|v\|_{W^{2,\gamma}(\Omega)} \le C_{2,\gamma}(\|\Delta v\|_{L^{\gamma}(\Omega)} + \|\Delta v\|_{L^{\gamma}(\Omega)}) \quad \text{for any } v \in W^{2,\gamma}(\Omega) \quad \text{with } \frac{\partial v}{\partial \nu} = 0.$$
 (2.10)

Consequently, combining (2.9) with (2.10), for any $s_0 > 0$, replacing v(t) by $v(t + s_0)$, we derive (2.6).

The Young inequality ([23]): Let $1 < p, q < +\infty$, $\frac{1}{p} + \frac{1}{q} = 1$. Then for any positive constants a and b, we have

$$ab \le \varepsilon a^p + \frac{1}{q} (\varepsilon p)^{-\frac{q}{p}} b^q.$$

The following lemma deals with local-in-time existence and uniqueness of a classical solution for the problem (1.2) (see [32]).

Lemma 2.3. ([32]) Assume that the nonnegative functions u_0, v_0 , and w_0 satisfies (1.5) for some $\vartheta \in (0, 1)$. Then there exists a maximal existence time $T_{max} \in (0, \infty]$ and a triple of nonnegative functions

$$\begin{cases} a \in C^0(\bar{\Omega} \times [0, T_{max})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max})), \\ v \in C^0(\bar{\Omega} \times [0, T_{max})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max})), \\ w \in C^{2,1}(\bar{\Omega} \times [0, T_{max})), \end{cases}$$

which solves (1.2) classically and satisfies

$$0 \le w \le \rho := \max\{1, \|w_0\|_{L^{\infty}(\Omega)}\} \quad in \quad \Omega \times (0, T_{max}).$$
(2.11)

Moreover, if $T_{max} < +\infty$, then

$$\|a(\cdot,t)\|_{L^{\infty}(\Omega)} + \|\nabla w(\cdot,t)\|_{L^{5}(\Omega)} \to \infty \quad as \quad t \nearrow T_{max}.$$
(2.12)

Firstly, by Lemma 2.3, we can pick $s_0 \in (0, T_{max})$, $s_0 \leq 1$ and $\beta > 0$ such that

$$||u(\tau)||_{L^{\infty}(\Omega)} \leq \beta, \quad ||v(\tau)||_{W^{1,\infty}(\Omega)} \leq \beta \text{ and } ||w(\tau)||_{W^{2,\infty}(\Omega)} \leq \beta \text{ for all } \tau \in [0, s_0].$$
 (2.13)

In some parts of our subsequent analysis, we introduce the variable transformation (see Tao et al. [40, 41, 45], Pang and Wang [32])

$$a = ue^{-\xi w},\tag{2.14}$$

upon which (1.2) takes the form

$$\begin{aligned} a_t &= e^{-\xi w} \nabla \cdot (e^{\xi w} \nabla a) - \chi e^{-\xi w} \nabla \cdot (e^{\xi w} a \nabla v) + \xi a v w + a(\mu - \xi \eta w)(1 - e^{\xi w} a - w), \ x \in \Omega, t > 0, \\ v_t &= \Delta v + a e^{\xi w} - v, \quad x \in \Omega, t > 0, \\ w_t &= -v w + \eta w (1 - a e^{\xi w} - w), \quad x \in \Omega, t > 0, \\ \frac{\partial a}{\partial \nu} &= \frac{\partial v}{\partial \nu} = \frac{\partial w}{\partial \nu} = 0, \quad x \in \partial \Omega, t > 0, \\ a(x, 0) &:= a_0(x) = u_0(x) e^{-\xi w_0(x)}, v(x, 0) = v_0(x), w(x, 0) = w_0(x), \quad x \in \Omega. \end{aligned}$$

$$(2.15)$$

3 Proof of the main result

In this section, we are going to establish an iteration step to develop the main ingredient of our result. Firstly, based on the ideas of Lemma 3.1 in [32] (see also Lemma 2.1 of [50]), we can derive the following properties of solutions of (1.2).

Lemma 3.1. Under the assumptions in theorem 1.1, we derive that there exists a positive constant C such that the solution of (1.2) satisfies

$$\int_{\Omega} u(x,t) + \int_{\Omega} v^2(x,t) + \int_{\Omega} |\nabla v(x,t)|^2 \le C \quad \text{for all} \quad t \in (0, T_{max}).$$
(3.1)

Moreover, for each $T \in (0, T_{max})$, one can find a constant C > 0 independent of ε such that

$$\int_{0}^{T} \int_{\Omega} [|\nabla v|^{2} + u^{2} + |\Delta v|^{2}] \le C.$$
(3.2)

Lemma 3.2. Let

$$A_{1} = \frac{1}{\delta+1} \left(\frac{\delta+1}{\delta}\right)^{-\delta} \left[\frac{\delta(\delta-1)}{2}\chi^{2}\right]^{\delta+1} C_{7} C_{\delta+1} e^{\xi(\delta-1)}$$
(3.3)

and $H(y) = y + A_1 y^{-\delta}$ for y > 0. For any fixed $\delta \ge 1, C_7, \chi, C_{\delta+1} > 0$, then

$$\min_{y>0} H(y) = \frac{\delta(\delta-1)\chi^2}{2} (C_7 C_{\delta+1})^{\frac{1}{\delta+1}}.$$

Proof. It is easy to verify that

$$H'(y) = 1 - A_1 \delta y^{-\delta - 1}.$$

Let H'(y) = 0, we have

$$y = (A_1 \delta)^{\frac{1}{\delta+1}} \,.$$

On the other hand, by $\lim_{y\to 0^+} H(y) = +\infty$ and $\lim_{y\to +\infty} H(y) = +\infty$, we have

$$\min_{y>0} H(y) = H[(A_1\delta)^{\frac{1}{\delta+1}}] = \frac{\delta(\delta-1)\chi^2}{2} (C_7 C_{\delta+1})^{\frac{1}{\delta+1}},$$

whereby the proof is completed.

Lemma 3.3. Let μ, χ, η and ξ be the positive constants. Assuming that (a, v, w) is a solution to (2.15) on $(0, T_{max})$. Then for all p > 1, there exists a positive constant $C := C(p, |\Omega|, \mu, \chi, \xi, \eta, \beta)$ such that

$$\int_{\Omega} a^p(x,t)dx \le C \quad \text{for all } t \in (0,T_{max}).$$
(3.4)

Proof. Firstly, assuming that $p \leq 2$. A straightforward differentiation, using (2.15) and two integrations by parts yields

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} e^{\xi w} a^{p} + (p+1) \int_{\Omega} e^{\xi w} a^{p} \\ &= \xi \int_{\Omega} e^{\xi w} a^{p} \cdot \{-vw + \eta w(1 - ae^{\xi w} - w)\} \\ &+ p \int_{\Omega} e^{\xi w} a^{p-1} \cdot \{e^{-\xi w} \nabla \cdot (e^{\xi w} \nabla a) - \chi e^{-\xi w} \nabla \cdot (e^{\xi w} a \nabla v)\} \\ &+ a\xi vw + a(\mu - \xi \eta w)(1 - ae^{\xi w} - w)\} + (p+1) \int_{\Omega} e^{\xi w} a^{p} \\ &= -p(p-1) \int_{\Omega} e^{\xi w} a^{p-2} |\nabla a|^{2} + p(p-1)\chi \int_{\Omega} e^{\xi w} a^{p-1} \nabla a \cdot \nabla v \\ &+ (p-1)\xi \int_{\Omega} e^{\xi w} a^{p} vw + \int_{\Omega} e^{\xi w} a^{p} \{(p+1) + (p-1)\xi \eta w(w-1) + p\mu(1-w)\} \\ &+ \int_{\Omega} e^{2\xi w} a^{p+1} [(p-1)\xi \eta w - p\mu] \\ &\coloneqq J_{1} + J_{2} + J_{3} + J_{4} + J_{5} \text{ for all } t \in (0, T_{max}). \end{aligned}$$

Now, in light of (2.11) and the Young inequality, we derive that

$$J_{3} \leq \varepsilon_{1} \int_{\Omega} e^{2\xi w} a^{p+1} + \frac{1}{p+1} (\varepsilon_{1} \times \frac{p+1}{p})^{-p} [(p-1)\xi]^{p+1} \int_{\Omega} e^{\xi w (1-p)} v^{p+1} \\ \leq \varepsilon_{1} \int_{\Omega} e^{2\xi w} a^{p+1} + \frac{1}{p+1} (\varepsilon_{1} \times \frac{p+1}{p})^{-p} [(p-1)\xi]^{p+1} \int_{\Omega} v^{p+1} \text{ for all } t \in (0, T_{max}),$$

$$(3.6)$$

$$J_{4} \leq [(p+1) + (p-1)\xi\eta\rho^{2} + p\mu] \int_{\Omega} e^{\xi w} a^{p}$$

$$\leq (p+1)[1 + \xi\eta\rho^{2} + \mu] \int_{\Omega} e^{\xi w} a^{p}$$

$$\leq \varepsilon_{2} \int_{\Omega} e^{2\xi w} a^{p+1} + \frac{1}{p+1} (\varepsilon_{2} \times \frac{p+1}{p})^{-p} (p+1)^{p+1} [1 + \xi\eta\rho^{2} + \mu]^{p+1} |\Omega| \text{ for all } t \in (0, T_{max}),$$
(3.7)

$$J_{5} \leq \int_{\Omega} e^{2\xi w} a^{p+1} [(p-1)\xi \eta \rho - p\mu] \text{ for all } t \in (0, T_{max})$$
(3.8)

and

$$J_{2} \leq \frac{p(p-1)}{2} \int_{\Omega} e^{\xi w} a^{p-2} |\nabla a|^{2} + \frac{p(p-1)}{2} \chi^{2} \int_{\Omega} e^{\xi w} a^{p} |\nabla v|^{2}$$

$$\leq \frac{p(p-1)}{2} \int_{\Omega} e^{\xi w} a^{p-2} |\nabla a|^{2} + \lambda_{0} \int_{\Omega} e^{2\xi w} a^{p+1}$$

$$+ \frac{1}{p+1} (\lambda_{0} \times \frac{p+1}{p})^{-p} [\frac{p(p-1)}{2} \chi^{2}]^{p+1} \int_{\Omega} e^{(1-p)\xi w} |\nabla v|^{2(p+1)}$$

$$\leq \frac{p(p-1)}{2} \int_{\Omega} e^{\xi w} a^{p-2} |\nabla a|^{2} + \lambda_{0} \int_{\Omega} e^{2\xi w} a^{p+1}$$

$$+ \frac{1}{p+1} (\lambda_{0} \times \frac{p+1}{p})^{-p} [\frac{p(p-1)}{2} \chi^{2}]^{p+1} \int_{\Omega} |\nabla v|^{2(p+1)} \text{ for all } t \in (0, T_{max})$$

$$(3.9)$$

and any small positive constants $\varepsilon_1, \varepsilon_2$ and λ_0 .

Inserting (3.7)–(3.9) into (3.5), we derive that

$$\frac{d}{dt} \int_{\Omega} e^{\xi w} a^{p} + (p+1) \int_{\Omega} e^{\xi w} a^{p} + \int_{\Omega} e^{2\xi w} a^{p+1} [p\mu - \varepsilon_{1} - \varepsilon_{2} - \lambda_{0} - (p-1)\xi\eta\rho] \\
\leq \frac{1}{p+1} (\lambda_{0} \times \frac{p+1}{p})^{-p} [\frac{p(p-1)}{2}\chi^{2}]^{p+1} \int_{\Omega} |\nabla v|^{2(p+1)} \\
+ C_{1}(\varepsilon_{1}, \varepsilon_{2}) \text{ for all } t \in (0, T_{max}),$$
(3.10)

where

$$C_{1}(\varepsilon_{1},\varepsilon_{2}) := \frac{1}{p+1} (\varepsilon_{2} \times \frac{p+1}{p})^{-p} (p+1)^{p+1} [1+\xi\eta\rho^{2}+\mu]^{p+1} |\Omega| + \frac{1}{p+1} (\varepsilon_{1} \times \frac{p+1}{p})^{-p} [(p-1)\xi]^{p+1} \int_{\Omega} v^{p+1}.$$
(3.11)

Next, from Lemma 3.1, N = 2 and the Gagliardo–Nirenberg inequality, it follows that

$$\|v(\cdot,t)\|_{L^p(\Omega)} \le C_2 \text{ for all } p \ge 1 \text{ and } t \in (0,T_{max}).$$
 (3.12)

This along with (3.11) entails

$$C_{1}(\varepsilon_{1},\varepsilon_{2}) \leq C_{3}(\varepsilon_{1},\varepsilon_{2})$$

$$:= \frac{1}{p+1} (\varepsilon_{2} \times \frac{p+1}{p})^{-p} (p+1)^{p+1} [1+\xi\eta\rho^{2}+\mu]^{p+1} |\Omega| \qquad (3.13)$$
$$+ C_{2} \frac{1}{p+1} (\varepsilon_{1} \times \frac{p+1}{p})^{-p} [(p-1)\xi]^{p+1}.$$

From this and (3.11) we also obtain

$$\frac{d}{dt} \int_{\Omega} e^{\xi w} a^{p} + (p+1) \int_{\Omega} e^{\xi w} a^{p} + \int_{\Omega} e^{2\xi w} a^{p+1} [p\mu - \varepsilon_{1} - \varepsilon_{2} - \lambda_{0} - (p-1)\xi\eta\rho] \\
\leq \frac{1}{p+1} (\lambda_{0} \times \frac{p+1}{p})^{-p} [\frac{p(p-1)}{2}\chi^{2}]^{p+1} \int_{\Omega} |\nabla v|^{2(p+1)} + C_{3}(\varepsilon_{1},\varepsilon_{2}) \text{ for all } t \in (0, T_{max}).$$
(3.14)

Then for any $t \in (s_0, T_{max})$, by means of the variation-of constants representation for the above inequality, we can estimate

$$\int_{\Omega} e^{\xi w} a^{p}(\cdot, t) + [p\mu - \varepsilon_{1} - \varepsilon_{2} - \lambda_{0} - (p-1)\xi\eta\rho] \int_{s_{0}}^{t} \int_{\Omega} e^{-(p-1)(t-s)} e^{2\xi w} a^{p+1} \\
\leq \int_{\Omega} u^{p}(s_{0}, t) + \frac{1}{p+1} (\lambda_{0} \times \frac{p+1}{p})^{-p} [\frac{p(p-1)}{2}\chi^{2}]^{p+1} \int_{s_{0}}^{t} \int_{\Omega} e^{-(p-1)(t-s)} |\nabla v|^{2(p+1)} \quad (3.15) \\
+ C_{3}(\varepsilon_{1}, \varepsilon_{2}) \text{ for all } t \in (0, T_{max}).$$

Next, according to the Gagliardo–Nirenberg inequality, (3.12) and Lemma 3.1, we can choose C_4 and C_5 such that

$$\begin{aligned} \|\nabla v(\cdot,s)\|_{L^{2(p+1)}(\Omega)}^{2(p+1)} &\leq C_4 \|v(\cdot,s)\|_{W^{2,p+1}(\Omega)}^{p+1} \|\nabla v(\cdot,s)\|_{L^2(\Omega)}^{p+1} \\ &\leq C_5 \|v(\cdot,s)\|_{W^{2,p+1}(\Omega)}^{p+1} \text{ for all } t \in (0,T_{max}). \end{aligned}$$
(3.16)

Therefore, due to $p \leq 2$, with the help of (3.16), applying (2.6) of Lemma 2.2 with $\gamma = p+1$, we obtain

for all $t \in (s_0, T_{max})$, where

$$C_{6} := \frac{1}{p+1} (\lambda_{0} \times \frac{p+1}{p})^{-p} [\frac{p(p-1)}{2} \chi^{2}]^{p+1} C_{5} C_{p+1} e^{\gamma s_{0}} \|v_{0}(\cdot, s_{0})\|_{W^{2,\gamma}(\Omega)}^{\gamma} \quad \text{and} \quad C_{7} := C_{5} e^{\xi}.$$
(3.18)

Substituting (3.17) into (3.15), we derive

$$\int_{\Omega} e^{\xi w} a^{p}(\cdot, t) + [p\mu - \varepsilon_{1} - \varepsilon_{2} - \lambda_{0} - (p-1)\xi\eta\rho] \int_{s_{0}}^{t} \int_{\Omega} e^{-(p-1)(t-s)} e^{2\xi w} a^{p+1} \\
\leq \frac{1}{p+1} (\lambda_{0} \times \frac{p+1}{p})^{-p} [\frac{p(p-1)}{2} \chi^{2}]^{p+1} C_{7} C_{p+1} \int_{s_{0}}^{t} \int_{\Omega} e^{-(p-1)(t-s)} e^{2\xi w} a^{p+1}(x,s) dx ds + C_{8}(\varepsilon_{1},\varepsilon_{2}) \\$$
(3.19)

for all $t \in (0, T_{max})$, where

$$C_8(\varepsilon_1, \varepsilon_2) := C_3(\varepsilon_1, \varepsilon_2) + C_6.$$

Choosing $\lambda_0 = (A_1 p)^{\frac{1}{p+1}}$ in (3.19) and using Lemma 3.2, we derive

$$\int_{\Omega} e^{\xi w} a^{p}(\cdot, t) \\
+ [p\mu - \varepsilon_{1} - \varepsilon_{2} - \frac{p(p-1)\chi^{2}}{2} (C_{7}C_{p+1})^{\frac{1}{p+1}} - (p-1)\xi\eta\rho] \int_{s_{0}}^{t} \int_{\Omega} e^{-(p-1)(t-s)} e^{2\xi w} a^{p+1} \\
\leq C_{8}(\varepsilon_{1}, \varepsilon_{2}).$$
(3.20)

Now, for any positive constants μ, χ, ξ and η , we may pick $p_0 > 1$ which is close to 1 such that

$$p_0\mu - \frac{p_0(p_0 - 1)\chi^2}{2} (C_7 C_{p_0 + 1})^{\frac{1}{p_0 + 1}} - (p_0 - 1)\xi\eta\rho > 0, \qquad (3.21)$$

thus, we can choose ε_1 and ε_2 appropriately small such that

$$0 < \varepsilon_1 + \varepsilon_2 < p_0 \mu - \frac{p_0(p_0 - 1)\chi^2}{2} (C_7 C_{p_0 + 1})^{\frac{1}{p_0 + 1}} - (p_0 - 1)\xi\eta\rho.$$
(3.22)

Collecting (3.20) and (3.22), we derive that for some $p_0 > 1$, there exists a positive constant C_9 such that

$$\int_{\Omega} u^{p_0}(x,t)dx \le C_9 \quad \text{for all} \quad t \in (s_0, T_{max}).$$
(3.23)

Next, we fix $q < \frac{2p_0}{(2-p_0)^+}$ and choose some $\alpha > \frac{1}{2}$ such that

$$q < \frac{1}{\frac{1}{p_0} - \frac{1}{2} + \frac{2}{2}(\alpha - \frac{1}{2})} \le \frac{2p_0}{(2 - p_0)^+}.$$
(3.24)

Now, involving the variation-of-constants formula for v, we have

$$v(t) = e^{-(A+1)}v(s_0) + \int_{s_0}^t e^{-(t-s)(A+1)}u(s)ds, \quad t \in (s_0, T_{max}).$$
(3.25)

Hence, it follows from (2.13) and (3.25) that

$$\| (A+1)^{\alpha} v(t) \|_{L^{q}(\Omega)}$$

$$\leq C_{10} \int_{s_{0}}^{t} (t-s)^{-\alpha-\frac{2}{2}(\frac{1}{p_{0}}-\frac{1}{q})} e^{-\mu(t-s)} \| u(s) \|_{L^{p_{0}}(\Omega)} ds + C_{10} s_{0}^{-\alpha-\frac{2}{2}(1-\frac{1}{q})} \| v(s_{0},t) \|_{L^{1}(\Omega)}$$

$$\leq C_{10} \int_{0}^{+\infty} \sigma^{-\alpha-\frac{2}{2}(\frac{1}{p_{0}}-\frac{1}{q})} e^{-\mu\sigma} d\sigma + C_{10} s_{0}^{-\alpha-\frac{2}{2}(1-\frac{1}{q})} \beta.$$

$$(3.26)$$

Hence, in light of Lemma 2.1, due to (3.24) and (3.26), we have

$$\int_{\Omega} |\nabla v(t)|^q \le C_{11} \quad \text{for all} \quad t \in (s_0, T_{max}) \tag{3.27}$$

and $q \in [1, \frac{2p_0}{(2-p_0)^+})$. Finally, in view of (2.13) and (3.27), we can get

$$\int_{\Omega} |\nabla v(t)|^q \le C_{12} \text{ for all } t \in (0, T_{max}) \text{ and } q \in [1, \frac{2p_0}{(2-p_0)^+})$$
(3.28)

with some positive constant C_{12} . Now, due to the Sobolev imbedding theorems and N = 2, we conclude that

 $||v(\cdot,t)||_{L^{\infty}(\Omega)} \le C_{13} \text{ for all } t \in (0, T_{max}).$ (3.29)

Applying the Young inequality, one obtains from (2.11), (2.15) and (3.29) that for any

Next, with the help of the Gagliardo-Nirenberg inequality (see e.g. [57]) yields that

$$C_{14} \int_{\Omega} a^{p+1} = C_{14} \|a^{\frac{p}{2}}\|_{L^{2}(\frac{p+1}{p})}^{2\frac{(p+1)}{p}}(\Omega)$$

$$\leq C_{15}(\|\nabla a^{\frac{p}{2}}\|_{L^{2}(\Omega)}^{\mu_{1}}\|a^{\frac{p}{2}}\|_{L^{\frac{2p_{0}}{p}}(\Omega)}^{1-\mu_{1}} + \|a^{\frac{p}{2}}\|_{L^{\frac{2p_{0}}{p}}(\Omega)}^{2\frac{(p+1)}{p}}$$

$$\leq C_{16}(\|\nabla a^{\frac{p}{2}}\|_{L^{2}(\Omega)}^{2(\mu-\mu)} + 1)$$

$$= C_{16}(\|\nabla u^{\frac{p}{2}}\|_{L^{2}(\Omega)}^{\frac{2(p-p_{0}+1)}{p}} + 1)$$
(3.31)

with some positive constants C_{15}, C_{16} and

$$\mu_1 = \frac{\frac{p}{p_0} - \frac{p}{p+1}}{\frac{p}{p_0}} = \frac{p+1-p_0}{p+1} \in (0,1).$$

Since, $p_0 > 1$ yields $p_0 < \frac{2p_0}{2(2-p_0)^+}$, in light of the Hölder inequality and (3.28), we derive

$$\frac{\chi^2 p(p-1)}{2} e^{\xi \rho} \int_{\Omega} a^p |\nabla v|^2 \le \frac{\chi^2 p(p-1)}{2} e^{\xi \rho} \left(\int_{\Omega} a^{\frac{p_0}{p_0-1}p} \right)^{\frac{p_0-1}{p_0}} \left(\int_{\Omega} |\nabla v|^{2p_0} \right)^{\frac{1}{p_0}} \le C_{17} \|a^{\frac{p}{2}}\|_{L^2^{\frac{p_0}{p_0-1}}(\Omega)}^2,$$
(3.32)

where C_{17} is a positive constant. Since $q_0 > 1$ and $p > q_0 - 1$, we have

$$\frac{p_0}{p} \le \frac{p_0}{p_0 - 1} < +\infty,$$

which together with the Gagliardo–Nirenberg inequality (see e.g. [57]) implies that

$$C_{17} \|a^{\frac{p}{2}}\|_{L^{2\frac{p_{0}}{p_{0}-1}}(\Omega)}^{2} \leq C_{18} (\|\nabla a^{\frac{p}{2}}\|_{L^{2}(\Omega)}^{\mu_{2}}\|a^{\frac{p}{2}}\|_{L^{\frac{2p_{0}}{p}}(\Omega)}^{1-\mu_{2}} + \|a^{\frac{p}{2}}\|_{L^{\frac{2p_{0}}{p}}(\Omega)}^{2})^{2}$$

$$\leq C_{19} (\|\nabla a^{\frac{p}{2}}\|_{L^{2}(\Omega)}^{2\mu_{2}} + 1)$$

$$= C_{19} (\|\nabla a^{\frac{p}{2}}\|_{L^{2}(\Omega)}^{\frac{2(p-p_{0}+1)}{p}} + 1)$$
(3.33)

with some positive constants C_{18}, C_{19} and

$$\mu_2 = \frac{\frac{p}{p_0} - \frac{p}{\frac{p}{p_0}}}{\frac{p}{p_0}} \in (0, 1).$$

Moreover, an application of the Young inequality shows that

$$C_{14} \int_{\Omega} a^{p+1} + \frac{\chi^2 p(p-1)}{2} e^{\xi \rho} \int_{\Omega} a^p |\nabla v|^2 \leq \frac{p(p-1)}{4} \int_{\Omega} a^{p-2} |\nabla a|^2 + C_{20}$$

$$\leq \frac{p(p-1)}{4} \int_{\Omega} e^{\xi w} a^{p-2} |\nabla a|^2 + C_{20}.$$
(3.34)

Inserting (3.34) into (3.30), we conclude that

$$\frac{d}{dt} \int_{\Omega} e^{\xi w} a^p + \frac{p(p-1)}{4} \int_{\Omega} e^{\xi w} a^{p-2} |\nabla a|^2 + p\mu \int_{\Omega} e^{2\xi w} a^{p+1} \le C_{21}.$$
 (3.35)

Therefore, integrating the above inequality with respect to t yields

$$||a(\cdot,t)||_{L^p(\Omega)} \le C_{22} \text{ for all } p \ge 1 \text{ and } t \in (0,T_{max})$$
 (3.36)

for some positive constant C_{22} . The proof of Lemma 3.3 is complete.

Remark 3.1. Since, in this paper, we only assume that $\mu > 0$ which is different from [32] (see the hypothesis of Lemma 3.2 to [32]), firstly by using the technical lemma (see Lemma 3.2), we could conclude the boundedness of $\int_{\Omega} a^{q_0}(q_0 > 1)$, then in light of the variationof-constants formula and L^q - L^p estimates for the heat semigroup, we may finally derive the boundedness of $\int_{\Omega} a^p$ (for any p > 1).

Our main result on global existence and boundedness thereby becomes a straightforward consequence of Lemma 2.3 and Lemma 3.3.

The proof of Theorem 1.1

Proof. Firstly, in light of (2.11), due to Lemma 3.3, we derive that there exist positive constants $p_0 > 2$ and C_1 such that

$$\|u(\cdot,t)\|_{L^{p_0}(\Omega)} \le C_1 \text{ for all } t \in (0, T_{max}).$$
(3.37)

Next, employing the standard estimate for Neumann semigroup provides C_2 and $C_3 > 0$ such that

$$\begin{aligned} \|\nabla v(t)\|_{L^{\infty}(\Omega)} &\leq C_{2} \int_{s_{0}+\infty}^{t} (t-s)^{-\alpha-\frac{2}{2p_{0}}} e^{-\mu(t-s)} \|u(s)\|_{L^{q_{0}}(\Omega)} ds + C_{2} s_{0}^{-\alpha} \|v(s_{0},t)\|_{L^{\infty}(\Omega)} \\ &\leq C_{2} \int_{0}^{+\infty} \sigma^{-\alpha-\frac{2}{2p_{0}}} e^{-\mu\sigma} d\sigma + C_{2} s_{0}^{-\alpha} \beta \\ &\leq C_{3} \text{ for all } t \in (0, T_{max}). \end{aligned}$$

$$(3.38)$$

Applying the Young inequality, in light of (2.11) and the first equation of (2.15), one obtains from (3.38) that for any $p \ge 4$

$$\frac{d}{dt} \int_{\Omega} e^{\xi w} a^{p} + p(p-1) \int_{\Omega} e^{\xi w} a^{p-2} |\nabla a|^{2} + \int_{\Omega} e^{\xi w} a^{p} \\
= \xi \int_{\Omega} e^{\xi w} a^{p} \cdot \{-vw + \eta w(1 - ae^{\xi w} - w)\} \\
+ p \int_{\Omega} e^{\xi w} a^{p-1} \cdot \{e^{-\xi w} \nabla \cdot (e^{\xi w} \nabla a) - \chi e^{-\xi w} \nabla \cdot (e^{\xi w} a \nabla v)\} \\
+ a\xi vw + a(\mu - \xi \eta w)(1 - ae^{\xi w} - w)\} + p \int_{\Omega} e^{\xi w} a^{p} \\
\leq \frac{p(p-1)}{4} \int_{\Omega} e^{\xi w} a^{p-2} |\nabla a|^{2} + p(p-1)\chi^{2}C_{4} \int_{\Omega} e^{\xi w} a^{p} \\
+ (p-1)\xi \int_{\Omega} e^{\xi w} a^{p} vw + \int_{\Omega} e^{\xi w} a^{p} \{(p+1) + (p-1)\xi \eta w(w-1) + p\mu(1-w)\} \\
+ \int_{\Omega} e^{2\xi w} a^{p+1} [(p-1)\xi \eta w - p\mu] \\
\leq \frac{p(p-1)}{4} \int_{\Omega} e^{\xi w} a^{p-2} |\nabla a|^{2} + C_{5}p^{2} (\int_{\Omega} a^{p+1} + 1) \text{ for all } t \in (0, T_{max}),$$
(3.39)

where C_4 and C_5 are independent of p. Here and throughout the proof of Theorem 1.1, we shall denote by $C_i (i \in \mathbb{N})$ several positive constants independent of p. Therefore, (3.39) implies that

$$\frac{d}{dt} \int_{\Omega} e^{\xi w} a^p + C_6 \int_{\Omega} |\nabla a^{\frac{p}{2}}|^2 + \int_{\Omega} e^{\xi w} a^p \le C_5 p^2 (\int_{\Omega} a^{p+1} + 1) \text{ for all } t \in (0, T_{max}).$$
(3.40)

Next, once more by means of the Gagliardo–Nirenberg inequality, we can estimate

$$C_{5}p^{2} \int_{\Omega} a^{p+1} = C_{5}p^{2} \|a^{\frac{p}{2}}\|_{L^{\frac{2(p+1)}{p}}(\Omega)}^{\frac{2(p+1)}{p}(\Omega)} \leq C_{7}p^{2}(\|\nabla a^{\frac{p}{2}}\|_{L^{2}(\Omega)}^{\frac{2(p+1)}{p}\varsigma_{1}} \|a^{\frac{p}{2}}\|_{L^{1}(\Omega)}^{\frac{2(p+1)}{p}\varsigma_{1}} + \|a^{\frac{p}{2}}\|_{L^{1}(\Omega)}^{\frac{2(p+1)}{p}}) = C_{7}p^{2}(\|\nabla a^{\frac{p}{2}}\|_{L^{2}(\Omega)}^{\frac{p+2}{p}} \|a^{\frac{p}{2}}\|_{L^{1}(\Omega)} + \|a^{\frac{p}{2}}\|_{L^{1}(\Omega)}^{\frac{2(p+1)}{p}}) \leq C_{6}\|\nabla a^{\frac{p}{2}}\|_{L^{2}(\Omega)}^{2} + C_{8}p^{\frac{4p}{p-2}} \|a^{\frac{p}{2}}\|_{L^{1}(\Omega)}^{\frac{2p}{p-2}} + C_{7}p^{2}\|a^{\frac{p}{2}}\|_{L^{1}(\Omega)}^{\frac{2(p+1)}{p}} \leq C_{6}\|\nabla a^{\frac{p}{2}}\|_{L^{2}(\Omega)}^{2} + C_{9}p^{\frac{4p}{p-2}} \|a^{\frac{p}{2}}\|_{L^{1}(\Omega)}^{\frac{2p}{p-2}},$$

$$(3.41)$$

where

$$0 < \varsigma_1 = \frac{2 - \frac{2p}{2(p+1)}}{1 - \frac{2}{2} + 2} = \frac{p+2}{2(p+1)} < 1.$$

Here we have use the fact that $\frac{4p}{p-2} \ge 2$. Therefore, inserting (3.41) into (3.40), we derive that

$$\frac{d}{dt} \int_{\Omega} e^{\xi w} a^{p} + \int_{\Omega} e^{\xi w} a^{p} \leq C_{9} p^{\frac{4p}{p-2}} \|a^{\frac{p}{2}}\|_{L^{1}(\Omega)}^{\frac{2p}{p-2}} + C_{5} p^{2} \\
\leq C_{10} p^{\frac{4p}{p-2}} \left(\max\{1, \|u^{\frac{p}{2}}\|_{L^{1}(\Omega)} \right)^{\frac{2p}{p-2}}.$$
(3.42)

Now, choosing $p_i = 2^{i+2}$ and letting $M_i = \max\{1, \sup_{t \in (0,T)} \int_{\Omega} a^{\frac{p_i}{2}}\}$ for $T \in (0, T_{max})$ and $i = 1, 2, 3, \cdots$. Then we obtain from (3.42) that

$$\frac{d}{dt} \int_{\Omega} e^{\xi w} a^{p_i} + \int_{\Omega} e^{\xi w} a^{p_i} \le C_{11} p_i^{\frac{2p_i}{p_i - 2}} M_{i-1}^{\frac{2p_i}{p_i - 2}}(T), \qquad (3.43)$$

which, together with the comparison argument entails that there exists a $\lambda > 1$ independent of *i* such that

$$M_{i}(T) \leq \max\{\lambda^{i} M_{i-1}^{\frac{2p_{i}}{p_{i}-2}}(T), e^{\xi} |\Omega| ||a_{0}||_{L^{\infty}(\Omega)}^{p_{i}}\}.$$
(3.44)

Here we use the fact that $\kappa_i := \frac{2p_i}{p_i-2} \leq 4$. Now, if $\lambda^i M_{i-1}^{\kappa_i}(T) \leq e^{\xi \rho} |\Omega| ||a_0||_{L^{\infty}(\Omega)}^{p_i}$ for infinitely many $i \geq 1$, we get

$$\left(\sup_{t\in(0,T)}\int_{\Omega}a^{p_{i-1}}(\cdot,t)\right)^{\frac{1}{p_{i-1}}} \le \left(\frac{e^{\xi\rho}|\Omega|\|a_0\|_{L^{\infty}(\Omega)}^{p_i}}{\lambda^i}\right)^{\frac{1}{p_{i-1}\kappa_i}}$$
(3.45)

for such i, which entails that

$$\sup_{t \in (0,T)} \|a(\cdot,t)\|_{L^{\infty}(\Omega)} \le \|a_0\|_{L^{\infty}(\Omega)}.$$
(3.46)

Otherwise, if $\lambda^i M_{i-1}^{\kappa_i}(T) > e^{\xi} |\Omega| ||a_0||_{L^{\infty}(\Omega)}^{p_i}$ for all sufficiently large *i*, then by (3.44), we derive that

 $M_i(T) \le \lambda^i M_{i-1}^{\kappa_i}(T)$ for all sufficiently large *i*. (3.47)

Hence, we may choose λ large enough such that

$$M_i(T) \le \lambda^i M_{i-1}^{\kappa_i}(T) \quad \text{for all} \quad i \ge 1.$$
(3.48)

Therefore, based on a straightforward induction (see e.g. Lemma 3.12 of [45]) we have

$$M_i(T) \le \lambda^{i + \sum_{j=2}^i (j-1) \cdot \prod_{k=j}^i \kappa_k} M_0^{\prod_{k=1}^i \kappa_k} \quad \text{for all } i \ge 1.$$
 (3.49)

where $\kappa_k := 2(1 + \varepsilon_k)$ satisfies $\varepsilon_k = \frac{4}{p_k - 2} \leq \frac{C_{12}}{2^k}$ for all $k \geq 1$ with some $C_{12} > 0$. Therefore, due to the fact that $\ln(1 + x) \leq x(x \geq 0)$, we derive

$$\Pi_{k=j}^{i} := 2^{i+1-j} e^{\sum_{k=j}^{i} \ln(1+\varepsilon_{j})}$$

$$\leq 2^{i+1-j} e^{\sum_{k=j}^{i} \varepsilon_{j}}$$

$$\leq 2^{i+1-j} e^{C_{12}} \text{ for all } i \geq 1 \text{ and } j \in \{1, \dots, i\},$$
(3.50)

which implies that

$$\frac{\sum_{j=2}^{i} (j-1) \cdot \prod_{k=j}^{i} \kappa_{k}}{2^{i+2}} \leq \frac{\sum_{j=2}^{i} (j-1)2^{i+1-j} e^{C_{12}}}{2^{i+2}}$$
$$\leq \frac{e^{C_{12}}}{2} \sum_{j=2}^{i} \frac{(j-1)}{2^{j}}$$
$$\leq \frac{e^{C_{12}}}{2} (\frac{1}{2} + \frac{1}{2^{2}})$$
$$= \frac{3e^{C_{12}}}{8}.$$

By the definition of p_i , we easily deduce from (3.49) that

$$M_{i}^{\frac{1}{p_{i}}}(T) \leq \lambda^{\frac{i}{2^{i+2}} + \frac{\sum_{j=2}^{i} (j-1) \cdot \Pi_{k=j}^{i} \kappa_{k}}{2^{i+2}}} M_{0}^{\frac{\Pi_{k=1}^{i} \kappa_{k}}{2^{i+2}}} \leq \lambda^{\frac{i}{2^{i+2}}} \lambda^{\frac{3e^{C_{12}}}{8}} M_{0}^{\frac{e^{C_{12}}}{4}}.$$
(3.51)

which after taking $i \to \infty$ and $T \nearrow T_{max}$ readily implies that

$$\|a(\cdot,t)\|_{L^{\infty}(\Omega)} \le \lambda^{\frac{3e^{C_{12}}}{8}} M_0^{\frac{e^{C_{12}}}{4}} \quad \text{for all} \quad t \in (0, T_{max}).$$
(3.52)

Employing almost exactly the same arguments as in the proof of Lemmata 3.5–3.6 in [32] (the minor necessary changes are left as an easy exercise to the reader), and taking advantage of (3.38) and (3.52), we conclude the estimate for any $T < T_{max}$,

$$\|\nabla w(\cdot, t)\|_{L^5(\Omega)} \le C \quad \text{for all} \quad t \in (0, T).$$

$$(3.53)$$

Now, with the above estimate in hand, using (3.46) and (3.52), employing the extendibility criterion provided by Lemma 2.3, we may prove Theorem 1.1.

Remark 3.2. If $\mu > \xi \eta \max\{||u_0||_{L^{\infty}(\Omega)}, 1\} + \mu^*(\chi^2, \xi)$ (see the proof of Lemma 3.4 to [32]), one only need to estimate $Cp^2 \int_{\Omega} a^p$ other than $Cp^2(\int_{\Omega} a^{p+1}+1)$, which is different from this paper.

Acknowledgement: This work is partially supported by the National Natural Science Foundation of China (No. 11601215), the Natural Science Foundation of Shandong Province of China (No. ZR2016AQ17) and the Doctor Start-up Funding of Ludong University (No. LA2016006).

References

- N. Bellomo, A. Belloquid, Y. Tao, M. Winkler, Toward a mathematical theory of Keller– Segel models of pattern formation in biological tissues, Math. Models Methods Appl. Sci., 25(9)(2015), 1663–1763.
- [2] V. Calvez, J. A. Carrillo, Volume effects in the Keller-Segel model: Energy estimates preventing blow-up, J. Math. Pures Appl., (9)(86)(2006), 155–175.
- [3] X. Cao, Boundedness in a three-dimensional chemotaxis-haptotaxis model, Z. Angew. Math. Phys., 67(1)(2016), 1–13.
- [4] M. A. J. Chaplain, G. Lolas, Mathematical modelling of cancer invasion of tissue: dynamic heterogeneity, Net. Hetero. Med., 1(2006), 399–439.
- [5] M. A. J. Chaplain, A. R. A. Anderson, Mathematical modelling of tissue invasion, in Cancer Modelling and Simulation, L. Preziosi, ed., Chapman Hall/CRC, Boca Raton, FL, 2003, 269–297.
- [6] L. Corrias, B. Perthame, H. Zaag, A chemotaxis model motivated by angiogenesis, C.
 R. Acad. Sci. Paris, Ser. I., 336(2003), 141–146.
- [7] L. Corrias, B. Perthame, H. Zaag, Global solutions of some chemotaxis and angiogenesis systems in high space dimensions, Milan J. Math., 72(2004), 1–28.

- [8] M. A. Fontelos, A. Friedman, B. Hu, Mathematical analysis of a model for the initiation of angiogenesis, SIAM J. Math. Anal., 33(2002), 1330–1355.
- K. Fujie, M. Winkler, T. Yokota, Boundedness of solutions to parabolic-elliptic Keller-Segel systems with signal-dependent sensitivity, Math. Methods Appl. Sci., 38(2015), 1212–1224.
- [10] A. Gerisch, M. A. J. Chaplain, Mathematical modelling of cancer cell invasion of tissue: Local and nonlocal models and the effect of adhesion, J. Theoret. Biol., 250(2008), 684– 704.
- [11] M. Herrero, J. Velázquez, A blow-up mechanism for a chemotaxis model, Ann. Scuola Norm. Super. Pisa Cl. Sci., 24(4)(1997), 633–683.
- [12] M. Hieber, J. Prüss, Heat kernels and maximal L^p-L^q estimate for parabolic evolution equations, Comm. Partial Diff. Eqns., 22(1997), 1647–1669.
- T. Hillen, K. J. Painter, A use's guide to PDE models for chemotaxis, J. Math. Biol., 58(2009), 183–217.
- [14] T. Hillen, K. J. Painter, M. Winkler, Convergence of a cancer invasion model to a logistic chemotaxis model, Math. Models Methods Appl. Sci., 23(2013), 165–198.
- [15] D. Horstmann, On the existence of radially symmetric blow-up solutions for the Keller-Segel model, J. Math. Biol., 44(2002), 463–478.
- [16] D. Horstmann, From 1970 until present: the Keller-Segel model in chemotaxis and its consequences, I. Jahresberichte der Deutschen Mathematiker-Vereinigung, 105(2003), 103–165.
- [17] D. Horstmann, G. Wang, Blow-up in a chemotaxis model without symmetry assumptions, Eur. J. Appl. Math., 12(2001), 159–177.
- [18] D. Horstmann, M. Winkler, Boundedness vs. blow-up in a chemotaxis system, J. Diff. Eqns, 215(2005), 52–107.

- [19] S. Ishida, K. Seki, T, Yokota, Boundedness in quasilinear Keller-Segel systems of parabolic-parabolic type on non-convex bounded domains, J. Diff. Eqns., 256(2014), 2993–3010.
- [20] W. Jäger, S. Luckhaus, On explosions of solutions to a system of partial differential equations modelling chemotaxis, Trans. Am. Math. Soc., 329(1992), 819–824.
- [21] N. I. Kavallaris, P. Souplet, Grow-up rate and refined asymptotics for a two-dimensional Patlak-Keller-Segel model in a disk, SIAM J. Math. Anal., 40(2009), 1852–1881.
- [22] E. Keller, L. Segel, Initiation of slime mold aggregation viewed as an instability, J. Theor. Biol., 26(1970), 399–415.
- [23] O. A. Ladyzenskaja, V. A. Solonnikov, N. N. Ural'eva, Linear and Quasi-linear Equations of Parabolic Type, Amer. Math. Soc. Transl. 23, AMS, Providence, RI, 1968.
- [24] J. Lankeit, Eventual smoothness and asymptotics in a three-dimensional chemotaxis system with logistic source, J. Diff. Eqns., 258(2015), 1158–1191.
- behaviour |25| J. Lankeit, Long-term inachemotaxis-fluid system with lo-Math. Models Methods Appl. Sci., 26(2016),2071,DOI: *qistic* source, http://dx.doi.org/10.1142/S021820251640008X.
- [26] L.A. Liotta, T. Clair, *Checkpoint for invasion*, Nature, 405(2000), 287–288.
- [27] G. Liţanu, C. Morales-Rodrigo, Asymptotic behavior of global solutions to a model of cell invasion, Math. Models Methods Appl. Sci., 20(2010), 1721–1758.
- [28] J. Liu, J. Zheng, Yifu Wang, Boundedness in a quasilinear chemotaxis-haptotaxis system with logistic source, Z. Angew. Math. Phys., 67(2)(2016), 1–33.
- [29] A. Marciniak-Czochra, M. Ptashnyk, Boundedness of solutions of a haptotaxis model, Math. Models Methods Appl. Sci., 20(2010), 449–476.
- [30] T. Nagai, Blow-up of radially symmetric solutions to a chemotaxis system, Adv. Math. Sci. Appl., 5(1995), 581–601.

- [31] K. Osaki, T. Tsujikawa, A. Yag, M. Mimura, Exponential attractor for a chemotaxis growth system of equations, Nonlinear Anal. TMA., 51(2002), 119–144.
- [32] P. Pang, Y. Wang, Global existence of a two-dimensional chemotaxis-haptotaxis model with remodeling of non-diffusible attractant, J. Diff. Eqns., 263(2)(2017), 1269–1292.
- [33] B. Perthame, Transport Equations in Biology, Birkhäuser Verlag, Basel, Switzerland, 2007.
- [34] M. Rascle, C. Ziti, Finite time blow-up in some models of chemotaxis, J. Math. Biol., 33(1995), 388–414.
- [35] J. A. Sherratt, Chemotaxis and chemokinesis in eukaryotic cells: The Keller-Segel equations as an approximation to a detailed model, Bull. Math. Biol., 56(1994), 129–146.
- [36] Y. Tao, Global existence of classical solutions to a combined chemotaxis-haptotaxis model with logistic source, J. Math. Anal. Appl., 354(2009), 60–69.
- [37] Y. Tao, Global existence for a haptotaxis model of cancer invasion with tissue remodeling, Nonlinear Anal. RWA., 12(2011), 418–435.
- [38] Y. Tao, Boundedness in a two-dimensional chemotaxis-haptotaxis system, Journal of Oceanography, 70(70)(2014), 165–174.
- [39] Y. Tao, M. Wang, Global solution for a chemotactic-haptotactic model of cancer invasion, Nonlinearity, 21(2008), 2221–2238.
- [40] Y. Tao, M. Wang, A combined chemotaxis-haptotaxis system: The role of logistic source, SIAM J. Math. Anal., 41(2009), 1533–1558.
- [41] Y. Tao, M. Winkler, A chemotaxis-haptotaxis model: the roles of porous medium diffusion and logistic source, SIAM J. Math. Anal., 43(2011), 685–704.
- [42] Y. Tao, M. Winkler, Boundedness in a quasilinear parabolic–parabolic Keller–Segel system with subcritical sensitivity, J. Diff. Eqns., 252(2012), 692–715.

- [43] Y. Tao, M. Winkler, Boundedness and stabilization in a multi-dimensional chemotaxishaptotaxis model, Proceedings of the Royal Society of Edinburgh, 144(2014), 1067–1084.
- [44] Y. Tao, M. Winkler, Dominance of chemotaxis in a chemotaxis-haptotaxis model, Nonlinearity, 27(2014), 1225–1239.
- [45] Y. Tao, M. Winkler, Energy-type estimates and global solvability in a two-dimensional chemotaxis-haptotaxis model with remodeling of non-diffusible attractant, J. Diff. Eqns., 257(2014), 784–815.
- [46] Y. Tao, M. Winkler, Large time behavior in a multidimensional chemotaxis-haptotaxis model with slow signal diffusion, SIAM J. Math. Anal., 47(6)(2015), 4229–4250.
- [47] J. I. Tello, M. Winkler, A chemotaxis system with logistic source, Comm. Partial Diff. Eqns., 32(2007), 849–877.
- [48] C. Walker, G. F. Webb, Global existence of classical solutions for a haptotaxis model, SIAM J. Math. Anal., 38(2007), 1694–1713.
- [49] Y. Wang, Boundedness in the higher-dimensional chemotaxis-haptotaxis model with nonlinear diffusion, J. Diff. Eqns., 260(2)(2016), 1975–1989.
- [50] M. Winkler, Boundedness in the higher-dimensional parabolic-parabolic chemotaxis system with logistic source, Comm. Partial Diff. Eqns., 35(2010), 1516–1537.
- [51] M. Winkler, Aggregation vs. global diffusive behavior in the higher-dimensional Keller-Segel model, J. Diff. Eqns., 248(2010), 2889–2905.
- [52] M. Winkler, Finite-time blow-up in the higher-dimensional parabolic-parabolic Keller-Segel system, J. Math. Pures Appl., 100(2013), 748–767.
- [53] M. Winkler, Global asymptotic stability of constant equilibriain a fully parabolic chemotaxis system with strong logistic dampening, J. Diff. Eqns., 257(2014), 1056–1077.
- [54] Q. Zhang, Y. Li, Global boundedness of solutions to a two-species chemotaxis system,
 Z. Angew. Math. Phys., 66(1)(2015), 83–93.

- [55] J. Zheng, Boundedness of solutions to a quasilinear parabolic-elliptic Keller-Segel system with logistic source, J. Diff. Eqns., 259(1)(2015), 120–140.
- [56] J. Zheng, Boundedness of solutions to a quasilinear parabolic-parabolic Keller-Segel system with logistic source, J. Math. Anal. Appl., 431(2)(2015), 867–888.
- [57] J. Zheng, Optimal controls of multidimensional modified Swift-Hohenberg equation, International Journal of Control, 88(10)(2015), 1–18.
- [58] J. Zheng, Boundedness of solution of a higher-dimensional parabolic-ODE-parabolic chemotaxis-haptotaxis model with generalized logistic source, Nonlinearity, 30(2017), 1987–2009.
- [59] J. Zheng, Boundedness of solutions to a quasilinear higher-dimensional chemotaxishaptotaxis model with nonlinear diffusion, Discrete and Continuous Dynamical Systems, (37)(1)(2017), 627–643.
- [60] J. Zheng, A note on boundedness of solutions to a higher-dimensional quasi-linear chemotaxis system with logistic source, Z. Angew. Math.Mech., (97)(4)(2017), 414–421.
- [61] J. Zheng, Y. Wang, Boundedness and decay behavior in a higher-dimensional quasilinearchemotaxis system with nonlinear logistic source, Comp. Math. Appl., 72(10)(2016), 2604–2619.