Asymptotic Expansions of Fundamental Solutions in Parabolic Homogenization

Jun Geng[∗] Zhongwei Shen†

Abstract

For a family of second-order parabolic systems with rapidly oscillating and timedependent periodic coefficients, we investigate the asymptotic behavior of fundamental solutions and establish sharp estimates for the remainders.

1 Introduction

In this paper we study the asymptotic behavior of fundamental solutions $\Gamma_{\varepsilon}(x, t; y, s)$ for a family of second-order parabolic operators $\partial_t + \mathcal{L}_{\varepsilon}$ with rapidly oscillating and time-dependent periodic coefficients. Specifically, we consider

$$
\mathcal{L}_{\varepsilon} = -\text{div}\left(A\big(x/\varepsilon, t/\varepsilon^2\big)\nabla\right) \tag{1.1}
$$

in $\mathbb{R}^d \times \mathbb{R}$, where $\varepsilon > 0$ and $A(y, s) = (a_{ij}^{\alpha\beta}(y, s))$ with $1 \le i, j \le d$ and $1 \le \alpha, \beta \le m$. Throughout the paper we will assume that the coefficient matrix $A = A(y, s)$ is real, bounded measurable and satisfies the ellipticity condition,

$$
||A||_{\infty} \le \mu^{-1} \quad \text{and} \quad \mu|\xi|^2 \le a_{ij}^{\alpha\beta}(y,s)\xi_i^{\alpha}\xi_j^{\beta} \tag{1.2}
$$

for any $\xi = (\xi_i^{\alpha}) \in \mathbb{R}^{m \times d}$ and a.e. $(y, s) \in \mathbb{R}^{d+1}$, where $\mu > 0$. We also assume that A is 1-periodic; i.e.,

$$
A(y + z, s + t) = A(y, s)
$$
 for $(z, t) \in \mathbb{Z}^{d+1}$ and a.e. $(y, s) \in \mathbb{R}^{d+1}$. (1.3)

Under these assumptions it is known that as $\varepsilon \to 0$, the operator $\partial_t + \mathcal{L}_{\varepsilon}$ G-converges to a parabolic operator $\partial_t + \mathcal{L}_0$ with constant coefficients [\[4\]](#page-24-0).

In the scalar case $m = 1$, it follows from a celebrated theorem of John Nash [\[17\]](#page-25-0) that local solutions of $(\partial_t + \mathcal{L}_{\varepsilon})u_{\varepsilon} = 0$ are Hölder continuous. More precisely, if $(\partial_t + \mathcal{L}_{\varepsilon})u_{\varepsilon} = 0$ in $Q_{2r} = Q_{2r}(x_0, t_0)$ for some $(x_0, t_0) \in \mathbb{R}^{d+1}$ and $0 < r < \infty$, where

$$
Q_r(x_0, t_0) = B(x_0, r) \times (t_0 - r^2, t_0), \tag{1.4}
$$

[∗]Supported in part by the NNSF of China (no. 11571152) and Fundamental Research Funds for the Central Universities (lzujbky-2017-161).

[†]Supported in part by NSF grant DMS-1600520.

then there exists some $\sigma \in (0,1)$, depending only on d and μ , such that

$$
||u_{\varepsilon}||_{C^{\sigma,\sigma/2}(Q_r)} \le Cr^{-\sigma} \left(\frac{1}{|Q_{2r}|}\int_{Q_{2r}} |u_{\varepsilon}|^2\right)^{1/2},\tag{1.5}
$$

where $C > 0$ depends only on d and μ . In particular, C and σ are independent of $\varepsilon > 0$. The periodicity assumption [\(1.3\)](#page-0-0) is not needed here. It follows that the fundamental solution $\Gamma_{\varepsilon}(x,t;y,s)$ for $\partial_t + \mathcal{L}_{\varepsilon}$ exists and satisfies the Gaussian estimate

$$
|\Gamma_{\varepsilon}(x,t;y,s)| \le \frac{C}{(t-s)^{\frac{d}{2}}} \exp\left\{-\frac{\kappa |x-y|^2}{t-s}\right\}
$$
(1.6)

for any $x, y \in \mathbb{R}^d$ and $-\infty < s < t < \infty$, where $\kappa > 0$ depends only on μ and $C > 0$ depends on d and μ (also see [\[1,](#page-24-1) [9\]](#page-24-2) for lower bounds).

If $m \geq 2$, the global Hölder estimate [\(1.5\)](#page-1-0) for $1 < r < \infty$ was established recently in [\[10\]](#page-24-3) for any $\sigma \in (0,1)$ under the assumptions that A is elliptic, periodic, and $A \in VMO_{x}$ (see [\(2.4\)](#page-4-0) for the definition of VMO_x). We mention that the local Hölder estimate for $0 < r < \varepsilon$ without the periodicity condition was obtained earlier in $[5, 16]$ $[5, 16]$. Consequently, by $[12, 6]$ $[12, 6]$, the matrix of fundamental solutions $\Gamma_{\varepsilon}(x,t;y,s) = (\Gamma_{\varepsilon}^{\alpha\beta}(x,t;y,s)),$ with $1 \leq \alpha, \beta \leq m$, exists and satisfies the estimate [\(1.6\)](#page-1-1), where $\kappa > 0$ depends only on μ . The constant $C > 0$ in [\(1.6\)](#page-1-1) depends on d, m, μ and the function $A^{\#}(r)$ in [\(2.5\)](#page-5-0), but not on $\varepsilon > 0$.

The primary purpose of this paper is to study the asymptotic behavior, as $\varepsilon \to 0$, of $\Gamma_{\varepsilon}(x,t;y,s), \nabla_x \Gamma_{\varepsilon}(x,t;y,s), \nabla_y \Gamma_{\varepsilon}(x,t;y,s)$, and $\nabla_x \nabla_y \Gamma_{\varepsilon}(x,t;y,s)$. Our main results extend the analogous estimates for elliptic operators $-\text{div}(A(x/\varepsilon)\nabla)$ in [\[3,](#page-24-7) [14\]](#page-24-8) to the parabolic setting. As demonstrated in the elliptic case [\[15\]](#page-24-9), the estimates in this paper open the doors for the use of layer potentials in solving initial-boundary value problems for the parabolic operator $\partial_t + \mathcal{L}_{\varepsilon}$ with sharp estimates that are uniform in $\varepsilon > 0$.

Let $\Gamma_0(x, t; y, s)$ denote the matrix of fundamental solutions for the homogenized operator $\partial_t + \mathcal{L}_0$. Our first result provides the sharp estimate for $\Gamma_{\varepsilon} - \Gamma_0$.

Theorem 1.1. Suppose that the coefficient matrix A satisfies conditions (1.2) and (1.3) . If $m \geq 2$, we also assume that $A \in VMO_x$. Then

$$
|\Gamma_{\varepsilon}(x,t;y,s) - \Gamma_0(x,t;y,s)| \le \frac{C\varepsilon}{(t-s)^{\frac{d+1}{2}}} \exp\left\{-\frac{\kappa|x-y|^2}{t-s}\right\}
$$
(1.7)

for any $x, y \in \mathbb{R}^d$ and $-\infty < s < t < \infty$, where $\kappa > 0$ depends only on μ . The constant C depends on d, m, μ , and $A^{\#}$ (if $m \geq 2$).

Let $\chi(y,s) = (\chi_i^{\alpha\beta})$ $j_j^{\alpha\beta}(y,s)$, where $1 \leq j \leq d$ and $1 \leq \alpha, \beta \leq m$, denote the matrix of correctors for $\partial_t + \mathcal{L}_{\varepsilon}$ (see Section 2 for its definition). The next theorem gives an asymptotic expansion for $\nabla_x \Gamma_\varepsilon(x,t;y,s)$.

Theorem 1.2. Suppose that the coefficient matrix A satisfies conditions (1.2) and (1.3) . Also assume that A is Hölder continuous,

$$
|A(x,t) - A(y,s)| \le \tau (|x - y| + |t - s|^{1/2})^{\lambda}
$$
\n(1.8)

for any $(x, t), (y, s) \in \mathbb{R}^{d+1}$, where $\tau \geq 0$ and $\lambda \in (0, 1)$. Then

$$
\left| \nabla_x \Gamma_{\varepsilon}(x, t; y, s) - \left(1 + \nabla \chi(x/\varepsilon, t/\varepsilon^2) \right) \nabla_x \Gamma_0(x, t; y, s) \right|
$$

\n
$$
\leq \frac{C\varepsilon}{(t-s)^{\frac{d+2}{2}}} \log(2 + \varepsilon^{-1}|t-s|^{1/2}) \exp\left\{-\frac{\kappa |x-y|^2}{t-s}\right\}
$$
(1.9)

for any $x, y \in \mathbb{R}^d$ and $-\infty < s < t < \infty$, where $\kappa > 0$ depends only on μ . The constant C depends on d, m, μ , and (λ, τ) in [\(1.8\)](#page-1-2).

With the summation convention this means that for $1 \leq i \leq d$ and $1 \leq \alpha, \beta \leq m$,

$$
\left| \frac{\partial \Gamma_{\varepsilon}^{\alpha\beta}}{\partial x_i}(x,t;y,s) - \frac{\partial \Gamma_0^{\alpha\beta}}{\partial x_i}(x,t;y,s) - \frac{\partial \chi_j^{\alpha\gamma}}{\partial x_i}(x/\varepsilon,t/\varepsilon^2) \frac{\partial \Gamma_0^{\gamma\beta}}{\partial x_j}(x,t;y,s) \right| \tag{1.10}
$$

is bounded by the RHS of [\(1.9\)](#page-2-0). Let $\widetilde{A}(y, s) = (\widetilde{a}_{ij}^{\alpha\beta}(y, s))$, where $\widetilde{a}_{ij}^{\alpha\beta}(y, s) = a_{ji}^{\beta\alpha}(y, -s)$. Let $\widetilde{\Gamma}_{\varepsilon}(x,t;y,s) = \left(\widetilde{\Gamma}_{\varepsilon}^{\alpha\beta}(x,t;y,s)\right)$ denote the matrix of fundamental solutions for the operator $\partial_t + \widetilde{\mathcal{L}}_{\varepsilon}$, where $\widetilde{\mathcal{L}}_{\varepsilon} = -\text{div}(\widetilde{A}(x/\varepsilon, t/\varepsilon^2)\nabla)$. Then

$$
\Gamma_{\varepsilon}^{\alpha\beta}(x,t;y,s) = \widetilde{\Gamma}_{\varepsilon}^{\beta\alpha}(y,-s;x,-t). \tag{1.11}
$$

Since \widetilde{A} satisfies the same conditions as A, it follows from [\(1.9\)](#page-2-0), [\(1.10\)](#page-2-1) and [\(1.11\)](#page-2-2) that

$$
\left| \frac{\partial \Gamma_{\varepsilon}^{\beta \alpha}}{\partial y_i}(x, t; y, s) - \frac{\partial \Gamma_0^{\beta \alpha}}{\partial y_i}(x, t; y, s) - \frac{\partial \widetilde{\chi}_j^{\alpha \gamma}}{\partial y_i}(y/\varepsilon, -s/\varepsilon^2) \frac{\partial \Gamma_0^{\beta \gamma}}{\partial y_j}(x, t; y, s) \right| \tag{1.12}
$$

is bounded by the RHS of [\(1.9\)](#page-2-0), where $\widetilde{\chi}(y, s) = (\widetilde{\chi}_j^{\alpha\beta})$ $\left(\alpha^\beta_j(y,s)\right)$ denotes the correctors for $\partial_t+\widetilde{\mathcal{L}}_\varepsilon.$ That is

$$
\left| \nabla_y \Gamma_\varepsilon^T(x, t; y, s) - \left(I + \nabla \widetilde{\chi}(y/\varepsilon, -s/\varepsilon^2) \right) \nabla_y \Gamma_0^T(x, t; y, s) \right|
$$

\$\leq \frac{C\varepsilon}{(t-s)^{\frac{d+2}{2}}} \log(2 + \varepsilon^{-1}|t-s|^{1/2}) \exp\left\{-\frac{\kappa |x-y|^2}{t-s}\right\}, \qquad (1.13)\$

where Γ_{ε}^{T} denotes the transpose of the matrix Γ_{ε} .

We also obtain an asymptotic expansion for $\nabla_x \nabla_y \Gamma_\varepsilon(x, t; y, s)$.

Theorem 1.3. Under the same assumptions on A as in Theorem [1.2,](#page-1-3) the estimate

$$
\left| \frac{\partial}{\partial x_i \partial y_j} \left\{ \Gamma_{\varepsilon}^{\alpha \beta}(x, t; y, s) \right\} - \frac{\partial}{\partial x_i} \left\{ \delta^{\alpha \gamma} x_k + \varepsilon \chi_k^{\alpha \gamma}(x/\varepsilon, t/\varepsilon^2) \right\} \frac{\partial^2}{\partial x_k \partial y_\ell} \left\{ \Gamma_0^{\gamma \sigma}(x, t; y, s) \right\} \frac{\partial}{\partial y_j} \left\{ \delta^{\beta \sigma} y_\ell + \varepsilon \widetilde{\chi}_\ell^{\beta \sigma}(y/\varepsilon, -s/\varepsilon^2) \right\} \right|
$$

$$
\leq \frac{C\varepsilon}{(t-s)^{\frac{d+3}{2}}} \log(2 + \varepsilon^{-1}|t-s|^{1/2}) \exp\left\{ -\frac{\kappa |x-y|^2}{t-s} \right\}
$$
(1.14)

holds for $x, y \in \mathbb{R}^d$ and $-\infty < s < t < \infty$, where κ depends only on μ . The constant C depends on d, m, μ , and (λ, τ) in [\(1.8\)](#page-1-2).

In the scale case $m = 1$, the estimate (1.7) , without the exponential factor, is known under the conditions that A is elliptic, periodic, symmetric, and time-independent (see $[13,$ p.77] and its references). This was proved by using the Floquet-Bloch decomposition of the fundamental solutions and by studying the spectral properties of elliptic operators

 $-(\nabla + ik) \cdot A(\nabla + ik)$ in a periodic cell, where $i = \sqrt{-1}$ and $k \in \mathbb{R}^d$. Such an approach is not available when the coefficient matrix A is time-dependent. To the best of authors' knowledge, the Gaussian bound in Theorem [1.1](#page-1-5) as well our estimates in Theorems [1.2](#page-1-3) and [1.3](#page-2-3) is new even in the case that $m = 1$ and A is time-independent.

As a corollary of Theorems [1.1](#page-1-5) and [1.2,](#page-1-3) we establish an interesting result on equistabilization for time-dependent coefficients (cf. [\[13,](#page-24-10) p.77]).

Corollary 1.4. Assume that A satisfies the same conditions as in Theorem [1.1.](#page-1-5) Let $f \in$ $L^{\infty}(\mathbb{R}^d)$ and u_{ε} be the bounded solution of the Cauchy problem,

$$
\begin{cases}\n(\partial_t + \mathcal{L}_{\varepsilon})u_{\varepsilon} = 0 & \text{in } \mathbb{R}^d \times (0, \infty), \\
u_{\varepsilon} = f & \text{on } \mathbb{R}^d \times \{t = 0\},\n\end{cases}
$$
\n(1.15)

with $\varepsilon = 1$ or 0. Then for any $x \in \mathbb{R}^d$ and $t \ge 1$,

$$
|u_1(x,t) - u_0(x,t)| \le Ct^{-1/2} \|f\|_{\infty}.
$$
\n(1.16)

Furthermore, if A is Hölder continuous,

$$
\left|\nabla u_1^{\alpha}(x,t) - \nabla u_0^{\alpha}(x,t) - \nabla \chi_j^{\alpha\beta}(x,t) \frac{\partial u_0^{\beta}}{\partial x_j}(x,t)\right| \le Ct^{-1} \log(2+t) \|f\|_{\infty}
$$
\n(1.17)

for any $x \in \mathbb{R}^d$ and $t \geq 1$.

We now describe some of the key ideas in the proof of Theorems [1.1,](#page-1-5) [1.2](#page-1-3) and [1.3.](#page-2-3) As indicated earlier, our main results extend the analogous results in [\[3,](#page-24-7) [14\]](#page-24-8) for the elliptic operators $-\text{div}(A(x/\varepsilon)\nabla)$, where $A = A(y)$ is elliptic and periodic. Our general approach is inspired by the work in [\[14\]](#page-24-8), which uses a two-scale expansion and relies on regularity estimates that are uniform in $\varepsilon > 0$. Following [\[11\]](#page-24-11), we consider the two-scale expansion

$$
w_{\varepsilon} = u_{\varepsilon}(x,t) - u_0(x,t) - \varepsilon \chi(x/\varepsilon, t/\varepsilon^2) S_{\varepsilon}(\nabla u_0) - \varepsilon^2 \phi(x/\varepsilon, t/\varepsilon^2) \nabla S_{\varepsilon}(\nabla u_0), \tag{1.18}
$$

where $\chi(y, s)$ and $\phi(y, s)$ are correctors and dual correctors respectively for $\partial_t + \mathcal{L}_{\varepsilon}$ (see Section 2 for their definitions). In [\(1.18\)](#page-3-0) the operator S_{ε} is a parabolic smoothing operator at scale ε . In comparison with the elliptic case, an extra term is added in the RHS of [\(1.18\)](#page-3-0). This modification allows us to show if $(\partial_t + \mathcal{L}_\varepsilon)u_\varepsilon = (\partial_t + \mathcal{L}_0)u_0$, then

$$
(\partial_t + \mathcal{L}_{\varepsilon})w_{\varepsilon} = \varepsilon \operatorname{div}(F_{\varepsilon})
$$
\n(1.19)

for some function F_{ε} , which depends only on u_0 . As a consequence, we may apply the uniform interior L^{∞} estimates established in [\[10\]](#page-24-3) to the function w_{ε} . To fully utilize the ideas above, we will consider the functions

$$
u_{\varepsilon}(x,t) = \int_{-\infty}^{t} \int_{\mathbb{R}^d} \Gamma_{\varepsilon}(x,t;y,s) f(y,s) e^{-\psi(y)} dy ds,
$$

$$
u_0(x,t) = \int_{-\infty}^{t} \int_{\mathbb{R}^d} \Gamma_0(x,t;y,s) f(y,s) e^{-\psi(y)} dy ds,
$$
 (1.20)

where ψ is a Lipchitz function in \mathbb{R}^d and $f \in C_0^{\infty}(Q_r(y_0, s_0); \mathbb{R}^m)$. The main technical step in proving Theorem [1.1](#page-1-5) involves bounding the L^{∞} norm

$$
||e^{\psi}(u_{\varepsilon} - u_0)||_{L^{\infty}(Q_r(x_0, t_0))}
$$
\n(1.21)

by $||f||_{L^2(Q_r(y_0,s_0))}$, where $0 < \varepsilon < r = c\sqrt{t_0-s_0}$. We remark that the use of weighted inequalities with weight e^{ψ} to generate the exponential factor in the Gaussian bound is more or less well known. Our approach may be regarded as a variation of the standard one found in $[12, 6]$ $[12, 6]$ (also see earlier work in $[9, 7, 8]$ $[9, 7, 8]$ $[9, 7, 8]$).

The proof of Theorem [1.2](#page-1-3) uses the estimate in Theorem [1.1.](#page-1-5) The stronger assumption that \vec{A} is Hölder continuos allows us to apply the uniform interior Lipschitz estimate obtained in [\[10\]](#page-24-3) to the function w_{ε} in [\(1.18\)](#page-3-0). To see Theorem [1.3,](#page-2-3) one uses the fact that as a function of (x, t) , $\nabla_y \Gamma_{\varepsilon}(x, t; y, s)$ is a solution of $(\partial_t + \mathcal{L}_{\varepsilon})u_{\varepsilon} = 0$, away from the pole (y, s) .

We end this section with some notations that will be used throughout the paper. A function $h = h(y, s)$ in \mathbb{R}^{d+1} is said to be 1-periodic if h is periodic with respect to \mathbb{Z}^{d+1} . We will use the notations

$$
\oint_E f = \frac{1}{|E|} \int_E f \quad \text{and} \quad h^{\varepsilon}(x, t) = h(x/\varepsilon, t/\varepsilon^2)
$$

for $\varepsilon > 0$, as well as the summation convention that the repeated indices are summed. Finally, we shall use κ to denote positive constants that depend only on μ , and C constants that depend at most on d, m, μ and the smoothness of A, but never on ε .

2 Preliminaries

Let $\mathcal{L}_{\varepsilon} = -\text{div}\,(A^{\varepsilon}(x,t)\nabla),$ where $A^{\varepsilon}(x,t) = A(x/\varepsilon,t/\varepsilon^2)$. Assume that $A(y,s)$ is 1-periodic in (y, s) and satisfies the ellipticity condition [\(1.2\)](#page-0-1). For $1 \leq j \leq d$ and $1 \leq \beta \leq m$, the corrector $\chi_j^{\beta} = \chi_j^{\beta}$ $j^{\beta}(y,s) \,=\, (\chi_{j}^{\alpha\beta}% ,\,\omega_{j}^{\beta}(y,s)) \label{m33}%$ $j^{\alpha\rho}(y,s)$ is defined as the weak solution of the following cell problem:

$$
\begin{cases}\n(\partial_s + \mathcal{L}_1)(\chi_j^{\beta}) = -\mathcal{L}_1(P_j^{\beta}) & \text{in } Y, \\
\chi_j^{\beta} = \chi_j^{\beta}(y, s) & \text{is 1-periodic in } (y, s), \\
\int_Y \chi_j^{\beta} = 0,\n\end{cases}
$$
\n(2.1)

where $Y = [0, 1)^{d+1}$, P_i^{β} $j_{j}^{\beta}(y) = y_{j}e^{\beta},$ and $e^{\beta} = (0, \ldots, 1, \ldots, 0)$ with 1 in the β^{th} position. Note that

$$
(\partial_s + \mathcal{L}_1)(\chi_j^{\beta} + P_j^{\beta}) = 0 \quad \text{in } \mathbb{R}^{d+1}.
$$
 (2.2)

By the rescaling property of $\partial_t + \mathcal{L}_{\varepsilon}$, one obtains that

$$
(\partial_t + \mathcal{L}_{\varepsilon}) \left\{ \varepsilon \chi_j^{\beta}(x/\varepsilon, t/\varepsilon^2) + P_j^{\beta}(x) \right\} = 0 \quad \text{in } \mathbb{R}^{d+1}.
$$
 (2.3)

We say $A \in VMO_x$ if

$$
\lim_{r \to 0} A^{\#}(r) = 0,\tag{2.4}
$$

where

$$
A^{\#}(r) = \sup_{\substack{0 < \rho < r \\ (x,t) \in \mathbb{R}^{d+1}}} \int_{t-\rho^2}^t \int_{y \in B(x,\rho)} \int_{z \in B(x,\rho)} |A(y,s) - A(z,s)| \, dz dy ds. \tag{2.5}
$$

Observe that if $A(y, s)$ is continuous in the variable y, uniformly in (y, s) , then $A \in VMO_x$.

Lemma 2.1. Assume that $A(y, s)$ is 1-periodic in (y, s) and satisfies [\(1.2\)](#page-0-1). If $m \geq 2$, we also assume $A \in VMO_x$. Then $\chi_j^{\beta} \in L^{\infty}(Y; \mathbb{R}^m)$.

Proof. In the scalar case $m = 1$, this follows from [\(2.2\)](#page-4-1) by Nash's classical estimate. Moreover, the estimate

$$
\left(\oint_{Q_r(x,t)} |\nabla \chi_j^{\beta}|^2\right)^{1/2} \le Cr^{\sigma-1} \tag{2.6}
$$

holds for any $0 < r < 1$ and $(x, t) \in \mathbb{R}^{d+1}$, where $Q_r(x, t) = B(x, r) \times (t - r^2, t)$, and $C > 0$ and $\sigma \in (0,1)$ depend on d and μ . If $m \geq 2$ and $A \in VMO_x$, the boundedness of χ_j^{β} j follows from the interior $W^{1,p}$ estimates for local solutions of $(\partial_t + \mathcal{L}_1)(u) = \text{div}(f)$ [\[5,](#page-24-4) [16\]](#page-25-1). In this case the estimate (2.6) holds for any $\sigma \in (0,1)$. case the estimate [\(2.6\)](#page-5-1) holds for any $\sigma \in (0,1)$.

Let $\widehat{A} = (\widehat{a}_{ij}^{\alpha\beta})$, where $1 \le i, j \le d, 1 \le \alpha, \beta \le m$, and

$$
\widehat{a}_{ij}^{\alpha\beta} = \int_{Y} \left[a_{ij}^{\alpha\beta} + a_{ik}^{\alpha\gamma} \frac{\partial}{\partial y_k} \chi_j^{\gamma\beta} \right];\tag{2.7}
$$

that is

$$
\widehat{A} = \int_Y \left\{ A + A \nabla \chi \right\}.
$$

It is known that the constant matrix \widehat{A} satisfies the ellipticity condition,

$$
\mu |\xi|^2 \le \widehat{a}_{ij}^{\alpha \beta} \xi_i^{\alpha} \xi_j^{\beta} \qquad \text{for any } \xi = (\xi_j^{\beta}) \in \mathbb{R}^{m \times d}, \tag{2.8}
$$

and $|\hat{a}_{ij}^{\alpha\beta}| \leq \mu_1$, where $\mu_1 > 0$ depends only on d, m and μ [\[4\]](#page-24-0). Denote

$$
\mathcal{L}_0 = -\mathrm{div}(\widehat{A}\nabla).
$$

Then $\partial_t + \mathcal{L}_0$ is the homogenized operator for the family of parabolic operators $\partial_t + \mathcal{L}_{\varepsilon}$, $\varepsilon > 0$.

To introduce the dual correctors, we consider the 1-periodic matrix-valued function

$$
B = A + A\nabla\chi - \hat{A}.\tag{2.9}
$$

More precisely, $B = B(y, s) = (b_{ij}^{\alpha\beta})$, where $1 \le i, j \le d, 1 \le \alpha, \beta \le m$, and

$$
b_{ij}^{\alpha\beta} = a_{ij}^{\alpha\beta} + a_{ik}^{\alpha\gamma} \frac{\partial \chi_j^{\gamma\beta}}{\partial y_k} - \widehat{a}_{ij}^{\alpha\beta}.
$$
 (2.10)

Lemma 2.2. Let $1 \leq j \leq d$ and $1 \leq \alpha, \beta \leq m$. Then there exist 1-periodic functions $\phi_{kij}^{\alpha\beta}(y, s)$ in \mathbb{R}^{d+1} such that $\phi_{kij}^{\alpha\beta} \in H^1(Y)$,

$$
b_{ij}^{\alpha\beta} = \frac{\partial}{\partial y_k} (\phi_{kij}^{\alpha\beta}) \quad \text{and} \quad \phi_{kij}^{\alpha\beta} = -\phi_{ikj}^{\alpha\beta}, \tag{2.11}
$$

where $1 \leq k, i \leq d+1, b_{ij}^{\alpha\beta}$ is defined by [\(2.10\)](#page-5-2) for $1 \leq i \leq d, b_{(d+1)j}^{\alpha\beta} = -\chi_j^{\alpha\beta}$ $_{j}^{\alpha \beta }$, and we have used the notation $y_{d+1} = s$.

Proof. This lemma was proved in [\[10\]](#page-24-3). We give a proof here for reader's convenience. By [\(2.1\)](#page-4-2) and [\(2.7\)](#page-5-3), $b_{ij}^{\alpha\beta} \in L^2(Y)$ and

$$
\int_{Y} b_{ij}^{\alpha\beta} = 0 \tag{2.12}
$$

for $1 \leq i \leq d+1$. It follows that there exist $f_{ij}^{\alpha\beta} \in H^2(Y)$ such that

$$
\begin{cases} \Delta_{d+1} f_{ij}^{\alpha \beta} = b_{ij}^{\alpha \beta} & \text{in } \mathbb{R}^{d+1}, \\ f_{ij}^{\alpha \beta} & \text{is 1-periodic} \end{cases}
$$
 (2.13)

where Δ_{d+1} denotes the Laplacian in \mathbb{R}^{d+1} . Write

$$
b_{ij}^{\alpha\beta} = \frac{\partial}{\partial y_k} \left\{ \frac{\partial}{\partial y_k} f_{ij}^{\alpha\beta} - \frac{\partial}{\partial y_i} f_{kj}^{\alpha\beta} \right\} + \frac{\partial}{\partial y_i} \left\{ \frac{\partial}{\partial y_k} f_{kj}^{\alpha\beta} \right\},
$$
(2.14)

where the index k is summed from 1 to $d+1$. Note that by (2.1) ,

$$
\sum_{i=1}^{d+1} \frac{\partial b_{ij}^{\alpha \beta}}{\partial y_i} = \sum_{i=1}^d \frac{\partial}{\partial y_i} b_{ij}^{\alpha \beta} - \frac{\partial}{\partial s} \chi_j^{\alpha \beta} = 0.
$$
 (2.15)

In view of [\(2.13\)](#page-6-0) this implies that

$$
\sum_{i=1}^{d+1} \frac{\partial}{\partial y_i} f_{ij}^{\alpha \beta}
$$

is harmonic in \mathbb{R}^{d+1} . Since it is 1-periodic, it must be constant. Consequently, by (2.14) , we obtain

$$
b_{ij}^{\alpha\beta} = \frac{\partial}{\partial y_k} (\phi_{kij}^{\alpha\beta}), \qquad (2.16)
$$

where

$$
\phi_{kij}^{\alpha\beta} = \frac{\partial}{\partial y_k} f_{ij}^{\alpha\beta} - \frac{\partial}{\partial y_i} f_{kj}^{\alpha\beta} \tag{2.17}
$$

is 1-periodic and belongs to $H^1(Y)$. It is easy to see that $\phi_{kij}^{\alpha\beta} = -\phi_{ikj}^{\alpha\beta}$. \Box

The 1-periodic functions $(\phi_{kij}^{\alpha\beta})$, given by Lemma [2.2,](#page-6-2) are called dual correctors for the family of parabolic operators $\partial_t + \mathcal{L}_{\varepsilon}$, $\varepsilon > 0$.

Lemma 2.3. Let $\phi = (\phi_{kij}^{\alpha\beta})$ be the dual correctors, given by Lemma [2.2.](#page-6-2) Under the same assumption as in Lemma [2.1,](#page-5-4) one has $\phi_{kij}^{\alpha\beta} \in L^{\infty}(Y)$.

Proof. It follows from [\(2.6\)](#page-5-1) that if $(x, t) \in \mathbb{R}^{d+1}$ and $0 < r < 1$,

$$
\int_{Q_r(x,t)} |b_{ij}^{\alpha\beta}|^2 \le Cr^{d+2\sigma} \tag{2.18}
$$

for some $\sigma \in (0,1)$. By covering the interval $(t-r,t)$ with intervals of length r^2 , we obtain

$$
\int_{B_r(x,t)} |b_{ij}^{\alpha\beta}|^2 \le Cr^{d-1+2\sigma},
$$

where $B_r(x,t) = B(x,r) \times (t-r,t)$. Hence, by Hölder's inequality,

$$
\int_{B_r(x,t)}|b_{ij}^{\alpha\beta}|\le Cr^{d+\sigma}.
$$

Thus, for any $(x, t) \in Y$,

$$
\int_{Y} \frac{|b_{ij}^{\alpha\beta}(y,s)|}{(|x-y|+|t-s|)^{d}} dyds \leq C \sum_{j=1}^{\infty} 2^{jd} \int_{|y-x|+|t-s|\sim 2^{-j}} |b_{ij}^{\alpha\beta}(y,s)| dyds
$$
\n
$$
\leq C.
$$
\n(2.19)

In view of [\(2.13\)](#page-6-0), by using the fundamental solution for Δ_{d+1} in \mathbb{R}^{d+1} , we may show that

$$
\|\nabla_{y,s} f_{ij}^{\alpha\beta}\|_{L^{\infty}(Y)} \leq C \|\nabla_{y,s} f_{ij}^{\alpha\beta}\|_{L^{2}(Y)} + \sup_{(x,t)\in Y} \int_Y \frac{|b_{ij}^{\alpha\beta}(y,s)|}{(|x-y|+|t-s|)^d} dyds,
$$

where $\nabla_{y,s}$ denotes the gradient in \mathbb{R}^{d+1} . This, together with [\(2.19\)](#page-7-0), shows that $|\nabla_{y,s} f_{ij}^{\alpha\beta}| \in$ $L^{\infty}(Y)$. By [\(2.17\)](#page-6-3) we obtain $\phi_{kij}^{\alpha\beta} \in L^{\infty}(Y)$. \Box

Remark 2.4. Suppose $A = A(y, s)$ is Hölder continuous in (y, s) . By [\(2.2\)](#page-4-1) and the standard regularity theory for $\partial_s + \mathcal{L}_1$, $\nabla \chi(y, s)$ is Hölder continuous in (y, s) . It follows that $b_{ij}^{\alpha\beta}(y, s)$ is Hölder continuous in (y, s) . In view of [\(2.13\)](#page-6-0) and [\(2.17\)](#page-6-3) one may deduce that $\nabla_{y,s} \phi_{kij}^{\alpha\beta}$ is Hölder continuous in (y, s) . This will be used in the proof of Theorems [1.2](#page-1-3) and [1.3.](#page-2-3)

Theorem 2.5. Suppose that A satisfies the conditions [\(1.2\)](#page-0-1) and [\(1.3\)](#page-0-0). If $m \geq 2$, we also assume $A \in VMO_x$. Let u_{ε} be a weak solution of $(\partial_t + \mathcal{L}_{\varepsilon})u_{\varepsilon} = \text{div}(f)$ in $Q_{2r} = Q_{2r}(x_0, t_0)$ for some $0 < r < \infty$, where $f = (f_i^{\alpha}) \in L^p(Q_{2r}; \mathbb{R}^{m \times d})$ for some $p > d + 2$. Then

$$
||u_{\varepsilon}||_{L^{\infty}(Q_{r})} \le C \left\{ \left(\int_{Q_{2r}} |u_{\varepsilon}|^{2} \right)^{1/2} + r \left(\int_{Q_{2r}} |f|^{p} \right)^{1/p} \right\},
$$
\n(2.20)

where C depends only on d, m, p, μ , and $A^{\#}$ in [\(2.5\)](#page-5-0) (if $m \geq 2$).

Proof. If $m = 1$, this follows from the well known Nash's estimate. The periodicity is not needed. If $m \ge 2$, [\(2.20\)](#page-7-1) follows from the uniform interior Hölder estimate in [\[10,](#page-24-3) Theorem 1.1]. 1.1].

Under the assumptions on A in Theorem [2.5,](#page-7-2) the matrix of fundamental solutions for $\partial_t + \mathcal{L}_{\varepsilon}$ in \mathbb{R}^{d+1} exists and satisfies the Gaussian estimate [\(1.6\)](#page-1-1). This follows from the L^{∞} estimate (2.20) by a general result in [\[12\]](#page-24-5) (also see [\[2,](#page-24-14) [6\]](#page-24-6)).

Theorem 2.6. Suppose that A satisfies conditions (1.2) and (1.3) . Also assume that A satisfies the Hölder condition [\(1.8\)](#page-1-2). Let u_{ε} be a weak solution of $(\partial_t + \mathcal{L}_{\varepsilon})u_{\varepsilon} = F$ in $Q_{2r} =$ $Q_{2r}(x_0, t_0)$ for some $0 < r < \infty$, where $F \in L^p(Q_{2r}; \mathbb{R}^m)$ for some $p > d+2$. Then

$$
\|\nabla u_{\varepsilon}\|_{L^{\infty}(Q_{r})} \le C \left\{ \frac{1}{r} \left(\int_{Q_{2r}} |u_{\varepsilon}|^{2} \right)^{1/2} + r \left(\int_{Q_{2r}} |F|^{p} \right)^{1/p} \right\},
$$
\n(2.21)

where C depends only on d, m, p, μ , and (λ, τ) in [\(1.8\)](#page-1-2).

Proof. This was proved in [\[10,](#page-24-3) Theorem 1.2].

The Lipschitz estimate [\(2.21\)](#page-8-0) allows us to bound $\nabla_x \Gamma_\varepsilon(x,t;y,s)$, $\nabla_y \Gamma_\varepsilon(x,t;y,s)$ and $\nabla_x \nabla_y \Gamma_\varepsilon(x,t;y,s).$

Theorem 2.7. Assume that A satisfies the same conditions as in Theorem [2.6.](#page-8-1) Then

$$
|\nabla_x \Gamma_{\varepsilon}(x, t; y, s)| + |\nabla_y \Gamma_{\varepsilon}(x, t; y, s)| \le \frac{C}{(t - s)^{\frac{d+1}{2}}} \exp\left\{-\frac{\kappa |x - y|^2}{t - s}\right\},\tag{2.22}
$$

$$
|\nabla_x \nabla_y \Gamma_{\varepsilon}(x, t; y, s)| \le \frac{C}{(t - s)^{\frac{d+2}{2}}} \exp\left\{-\frac{\kappa |x - y|^2}{t - s}\right\},\tag{2.23}
$$

for any $x, y \in \mathbb{R}^d$ and $-\infty < s < t < \infty$, where $\kappa > 0$ depends only on μ . The constant C depends on d, m, μ , and (λ, τ) in [\(1.8\)](#page-1-2).

Proof. Fix $x_0, y_0 \in \mathbb{R}^d$ and $s_0 < t_0$. Let $u_\varepsilon(x,t) = \Gamma_\varepsilon(x,t; y_0, s_0)$. Then $(\partial_t + \mathcal{L}_\varepsilon)u_\varepsilon = 0$ in $Q_{2r}(x_0, t_0)$, where $r = \sqrt{t_0 - s_0}/8$. The estimate for $|\nabla_x \Gamma(x_0, t_0; y_0, s_0)|$ now follows from [\(2.21\)](#page-8-0) and [\(1.6\)](#page-1-1) (with a different κ). In view of [\(1.11\)](#page-2-2) this also gives the estimate for $|\nabla_y \Gamma_\varepsilon(x_0, t_0; y_0, s_0)|$. Finally, to see (2.23) , we let $v_\varepsilon(x, t) = \nabla_y \Gamma_\varepsilon(x, t; y_0, s_0)$. Then $(\partial_t + \mathcal{L}_\varepsilon)v_\varepsilon = 0$ in $Q_{2r}(x_0, t_0)$. By applying [\(2.21\)](#page-8-0) to v_ε and using the estimate in [\(2.22\)](#page-8-3) for $\nabla_v \Gamma_\varepsilon(x, t; y, s)$, we obtain the desired estimate for $|\nabla_x \nabla_v \Gamma_\varepsilon(x_0, t_0; y_0, s_0)|$. $\nabla_{\mathbf{y}}\Gamma_{\varepsilon}(x,t;y,s)$, we obtain the desired estimate for $|\nabla_{x}\nabla_{\mathbf{y}}\Gamma_{\varepsilon}(x_0,t_0;y_0,s_0)|$.

3 A two-scale expansion

Suppose that

$$
(\partial_t + \mathcal{L}_\varepsilon)u_\varepsilon = (\partial_t + \mathcal{L}_0)u_0\tag{3.1}
$$

 \Box

in $\Omega \times (T_0, T_1)$, where $\Omega \subset \mathbb{R}^d$. Let S_ε be a linear operator to be chosen later. Following [\[11\]](#page-24-11), we consider the two-scale expansion $w_{\varepsilon} = (w_{\varepsilon}^{\alpha})$, where

$$
w_{\varepsilon}^{\alpha}(x,t) = u_{\varepsilon}^{\alpha}(x,t) - u_{0}^{\alpha}(x,t) - \varepsilon \chi_{j}^{\alpha\beta}(x/\varepsilon,t/\varepsilon^{2}) S_{\varepsilon} \left(\frac{\partial u_{0}^{\beta}}{\partial x_{j}}\right) - \varepsilon^{2} \phi_{(d+1)ij}^{\alpha\beta}(x/\varepsilon,t/\varepsilon^{2}) \frac{\partial}{\partial x_{i}} S_{\varepsilon} \left(\frac{\partial u_{0}^{\beta}}{\partial x_{j}}\right),
$$
\n(3.2)

and $\chi_i^{\alpha\beta}$ $\phi_j^{\alpha\beta}$, $\phi_{(d+1)ij}^{\alpha\beta}$ are the correctors and dual correctors introduced in the last section. The repeated indices i, j in [\(3.2\)](#page-9-0) are summed from 1 to d.

Proposition 3.1. Let $u_{\varepsilon} \in L^2(T_0, T_1; H^1(\Omega))$ and $u_0 \in L^2(T_0, T_1; H^2(\Omega))$. Let w_{ε} be defined by [\(3.2\)](#page-9-0). Assume [\(3.1\)](#page-8-4) holds in $\Omega \times (T_0, T_1)$. Then

$$
(\partial_t + \mathcal{L}_{\varepsilon})w_{\varepsilon} = \varepsilon \operatorname{div}(F_{\varepsilon})
$$
\n(3.3)

 \Box

in $\Omega \times (T_0, T_1)$, where $F_{\varepsilon} = (F^{\alpha}_{\varepsilon,i})$ and

$$
F_{\varepsilon,i}^{\alpha}(x,t) = \varepsilon^{-1} \left(a_{ij}^{\alpha\beta}(x/\varepsilon,t/\varepsilon^2) - \widehat{a}_{ij}^{\alpha\beta} \right) \left(\frac{\partial u_0^{\beta}}{\partial x_j} - S_{\varepsilon} \left(\frac{\partial u_0^{\beta}}{\partial x_j} \right) \right) + a_{ij}^{\alpha\beta}(x/\varepsilon,t/\varepsilon^2) \chi_k^{\beta\gamma}(x/\varepsilon,t/\varepsilon^2) \frac{\partial}{\partial x_j} S_{\varepsilon} \left(\frac{\partial u_0^{\beta}}{\partial x_k} \right) + \phi_{ikj}^{\alpha\beta}(x/\varepsilon,t/\varepsilon^2) \frac{\partial}{\partial x_k} S_{\varepsilon} \left(\frac{\partial u_0^{\beta}}{\partial x_j} \right) + \varepsilon \phi_{i(d+1)j}^{\alpha\beta}(x/\varepsilon,t/\varepsilon^2) \partial_t S_{\varepsilon} \left(\frac{\partial u_0^{\beta}}{\partial x_j} \right) - a_{ij}^{\alpha\beta}(x/\varepsilon,t/\varepsilon^2) \left(\frac{\partial}{\partial x_j} (\phi_{(d+1)\ell k}^{\beta\gamma}) \right) (x/\varepsilon,t/\varepsilon^2) \frac{\partial}{\partial x_\ell} S_{\varepsilon} \left(\frac{\partial u_0^{\gamma}}{\partial x_k} \right) - \varepsilon a_{ij}^{\alpha\beta}(x/\varepsilon,t/\varepsilon^2) \phi_{(d+1)\ell k}^{\beta\gamma}(x/\varepsilon,t/\varepsilon^2) \frac{\partial^2}{\partial x_j \partial x_\ell} S_{\varepsilon} \left(\frac{\partial u_0^{\gamma}}{\partial x_k} \right).
$$
\n(3.4)

The repeated indices i, j, k, ℓ are summed from 1 to d.

Proof. This proposition was proved in [\[11,](#page-24-11) Theorem 2.2].

We now introduce a parabolic smoothing operator. Let

$$
\mathcal{O} = \{(x, t) \in \mathbb{R}^{d+1} : |x|^2 + |t| \le 1\}.
$$

Fix a nonnegative function $\theta = \theta(x,t) \in C_0^{\infty}(\mathcal{O})$ such that $\int_{\mathbb{R}^{d+1}} \theta = 1$. Let $\theta_{\varepsilon}(x,t) =$ $\varepsilon^{-d-2}\theta(x/\varepsilon,t/\varepsilon^2)$. Define

$$
S_{\varepsilon}(f)(x,t) = f * \theta_{\varepsilon}(x,t) = \int_{\mathbb{R}^{d+1}} f(x - y, t - s) \theta_{\varepsilon}(y,s) \, dyds. \tag{3.5}
$$

Lemma 3.2. Let $g = g(x, t)$ be a 1-periodic function in (x, t) and $\psi = \psi(x)$ a bounded Lipschitz function in \mathbb{R}^d . Then

$$
||e^{\psi}g^{\varepsilon}S_{\varepsilon}(f)||_{L^{p}(\mathbb{R}^{d+1})} \leq C e^{\varepsilon ||\nabla\psi||_{\infty}} ||g||_{L^{p}(Y)} ||e^{\psi}f||_{L^{p}(\mathbb{R}^{d+1})}
$$
\n(3.6)

for any $1 \le p < \infty$, where $g^{\varepsilon}(x,t) = g(x/\varepsilon, t/\varepsilon^2)$ and C depends only on d and p.

Proof. Using $\int_{\mathbb{R}^{d+1}} \theta_{\varepsilon} = 1$ and Hölder's inequality, we obtain

$$
|S_{\varepsilon}(e^{-\psi}f)(x,t)|^{p} \leq \int_{\mathbb{R}^{d+1}} |e^{-\psi(y)}f(y,s)|^{p} \theta_{\varepsilon}(x-y,t-s) dyds.
$$

It follows that

$$
|e^{\psi(x)}S_{\varepsilon}(e^{-\psi}f)(x,t)|^{p} \leq \int_{\mathbb{R}^{d+1}} |e^{\psi(x)-\psi(y)}f(y,s)|^{p} \theta_{\varepsilon}(x-y,t-s) dyds
$$

$$
\leq e^{\varepsilon p \|\nabla \psi\|_{\infty}} \int_{\mathbb{R}^{d+1}} |f(y,s)|^{p} \theta_{\varepsilon}(x-y,t-s) dyds,
$$

where we have used the facts that $|\psi(x) - \psi(y)| \leq ||\nabla \psi||_{\infty} |x - y|$ and $\theta_{\varepsilon}(x - y, t - s) = 0$ if $|x - y| > \varepsilon$, for the last step. Hence, by Fubini's Theorem,

$$
\int_{\mathbb{R}^{d+1}} |g^{\varepsilon}(x,t)|^p |e^{\psi} S_{\varepsilon}(e^{-\psi}f)(x,t)|^p dxdt
$$
\n
$$
\leq e^{\varepsilon p \|\nabla \psi\|_{\infty}} \sup_{(y,s)\in \mathbb{R}^{d+1}} \int_{\mathbb{R}^{d+1}} |g^{\varepsilon}(x,t)|^p \theta_{\varepsilon}(x-y,t-s) dxdt \int_{\mathbb{R}^{d+1}} |f(y,s)|^p dyds
$$
\n
$$
\leq C e^{\varepsilon p \|\nabla \psi\|_{\infty}} \|g\|_{L^p(Y)}^p \|f\|_{L^p(\mathbb{R}^{d+1})}^p,
$$

where C depends only on d . This gives (3.6) .

Remark 3.3. Let $\Omega \subset \mathbb{R}^d$ and $(T_0, T_1) \subset \mathbb{R}$. Define

$$
\Omega_{\varepsilon} = \left\{ x \in \mathbb{R}^d : \text{dist}(x, \Omega) < \varepsilon \right\}. \tag{3.7}
$$

Observe that for $(x, t) \in \Omega \times (T_0, T_1)$, $S_{\varepsilon}(f)(x, t) = S_{\varepsilon}(f\eta_{\varepsilon})(x, t)$, where $\eta_{\varepsilon} = \eta_{\varepsilon}(x, t)$ is the characteristic function of $\Omega_{\varepsilon} \times (T_0 - \varepsilon^2, T_1 + \varepsilon^2)$. By applying [\(3.6\)](#page-10-0) to the function $f\eta_{\varepsilon}$, one may deduce that

$$
\int_{T_0}^{T_1} \int_{\Omega} |e^{\psi} g^{\varepsilon} S_{\varepsilon}(f)|^p \, dx dt \le C e^{\varepsilon p \|\nabla \psi\|_{\infty}} \|g\|_{L^p(Y)}^p \int_{T_0 - \varepsilon^2}^{T_1 + \varepsilon^2} \int_{\Omega_{\varepsilon}} |e^{\psi} f|^p \, dx dt. \tag{3.8}
$$

Using $\int_{\mathbb{R}^{d+1}} |\nabla \theta_{\varepsilon}| dx dt \leq C \varepsilon^{-1}$, the same argument as in the proof of Lemma [3.2](#page-10-1) also shows that

$$
\int_{T_0}^{T_1} \int_{\Omega} |e^{\psi} g^{\varepsilon} \nabla S_{\varepsilon}(f)|^p \, dx dt \le C \varepsilon^{-p} e^{\varepsilon p \|\nabla \psi\|_{\infty}} \|g\|_{L^p(Y)}^p \int_{T_0 - \varepsilon^2}^{T_1 + \varepsilon^2} \int_{\Omega_{\varepsilon}} |e^{\psi} f|^p \, dx dt \tag{3.9}
$$

for $1 \leq p < \infty$, where C depends only on d and p.

 \Box

Lemma 3.4. Let S_{ε} be defined as in [\(3.5\)](#page-9-1). Let $1 \leq p < \infty$ and ψ be a bounded Lipschitz function in \mathbb{R}^d . Then for $\Omega \subset \mathbb{R}^d$ and $(T_0, T_1) \subset \mathbb{R}$,

$$
\int_{T_0}^{T_1} \int_{\Omega} |e^{\psi} (S_{\varepsilon}(\nabla f) - \nabla f)|^p \, dx dt
$$
\n
$$
\leq C \varepsilon^p e^{\varepsilon p ||\nabla \psi||_{\infty}} \int_{T_0 - \varepsilon^2}^{T_1 + \varepsilon^2} \int_{\Omega_{\varepsilon}} |e^{\psi} (|\nabla^2 f| + |\partial_t f|)|^p \, dx dt,
$$
\n(3.10)

where Ω_{ε} is given by [\(3.7\)](#page-10-2) and C depends only on d and p.

Proof. Write

$$
S_{\varepsilon}(\nabla f)(x,t) - \nabla f(x,t) = J_1(x,t) + J_2(x,t),
$$

where

$$
J_1(x,t) = \int_{\mathbb{R}^{d+1}} \theta_{\varepsilon}(y,s) \big(\nabla f(x-y,t-s) - \nabla f(x-y,t) \big) dyds,
$$

$$
J_2(x,t) = \int_{\mathbb{R}^{d+1}} \theta_{\varepsilon}(y,s) \big(\nabla f(x-y,t) - \nabla f(x,t) \big) dyds.
$$

To estimate J_2 , we observe that by Hölder's inequality and the fact $\int_{\mathbb{R}^{d+1}} \theta_{\varepsilon} dy ds = 1$,

$$
|J_2(x,t)|^p \leq \int_{\mathbb{R}^{d+1}} \theta_{\varepsilon}(y,s) |\nabla f(x-y,t) - \nabla f(x,t)|^p dy ds,
$$

and

$$
|\nabla f(x - y, t) - \nabla f(x, t)| = \Big| \int_0^1 \frac{\partial}{\partial \tau} \nabla f(x - \tau y, t) d\tau \Big|
$$

\n
$$
\leq |y| \int_0^1 |\nabla^2 f(x - \tau y, t)| d\tau
$$

\n
$$
\leq |y| \left(\int_0^1 |\nabla^2 f(x - \tau y, t)|^p d\tau \right)^{1/p}
$$

.

It follows by Fubini's Theorem that

$$
\int_{T_0}^{T_1} \int_{\Omega} |e^{\psi(x)} J_2(x, t)|^p dx dt
$$
\n
$$
\leq \int_{T_0}^{T_1} \int_{\Omega} \int_{\mathbb{R}^{d+1}} \int_0^1 e^{p\psi(x)} \theta_{\varepsilon}(y, s) |y|^p |\nabla^2 f(x - \tau y, t)|^p d\tau dy ds dx dt
$$
\n
$$
\leq \varepsilon^p e^{\varepsilon p ||\nabla \psi||_{\infty}} \int_{T_0}^{T_1} \int_{\Omega} \int_{\mathbb{R}^{d+1}} \int_0^1 e^{p\psi(x - \tau y)} \theta_{\varepsilon}(y, s) |\nabla^2 f(x - \tau y, t)|^p d\tau dy ds dx dt
$$
\n
$$
\leq \varepsilon^p e^{\varepsilon p ||\nabla \psi||_{\infty}} \int_{T_0}^{T_1} \int_{\Omega_{\varepsilon}} |e^{\psi} \nabla^2 f|^p dx dt,
$$

where we have used the facts that $|\psi(x) - \psi(x - \tau y)| \leq |\tau||y|| |\nabla \psi||_{\infty}$ and $\theta_{\varepsilon}(y, s) = 0$ if $|y| > \varepsilon$.

Finally, to estimate J_1 , we first use integration by parts to obtain

$$
|J_1(x,t)| \leq \int_{\mathbb{R}^{d+1}} |\nabla \theta_{\varepsilon}(y,s)||f(x-y,t-s) - f(x-y,t)| dyds.
$$

By Hölder's inequality,

$$
|J_1(x,t)|^p \leq C\varepsilon^{1-p} \int_{\mathbb{R}^{d+1}} |\nabla \theta_{\varepsilon}(y,s)||f(x-y,t-s) - f(x-y,t)|^p dy ds,
$$

where we have also used the fact $\int_{\mathbb{R}^{d+1}} |\nabla \theta_{\varepsilon}| dy ds \leq C \varepsilon^{-1}$. Using

$$
|f(x-y,t-s) - f(x-y,t)| \leq \Big| \int_0^1 \frac{\partial}{\partial \tau} f(x-y,t-\tau s) d\tau \Big|
$$

$$
\leq |s| \left(\int_0^1 |\partial_t f(x-y,t-\tau s)|^p d\tau \right)^{1/p},
$$

we see that by Fubini's Theorem,

$$
\int_{T_0}^{T_1} \int_{\Omega} |e^{\psi(x)} J_1(x,t)|^p dx dt
$$
\n
$$
\leq C \varepsilon^{1-p} \int_{T_0}^{T_1} \int_{\Omega} \int_{\mathbb{R}^{d+1}} \int_{0}^{1} e^{p\psi(x)} |\nabla \theta_{\varepsilon}(y,s)| |s|^p |\partial_t f(x-y,t-\tau s)|^p d\tau dy ds dx dt
$$
\n
$$
\leq C \varepsilon^{1+p} e^{\varepsilon p ||\nabla \psi||_{\infty}} \int_{T_0}^{T_1} \int_{\Omega} \int_{\mathbb{R}^{d+1}} \int_{0}^{1} e^{p\psi(x-y)} |\nabla \theta_{\varepsilon}(y,s)| |\partial_t f(x-y,t-\tau s)|^p d\tau dy ds dx dt
$$
\n
$$
\leq C \varepsilon^{p} e^{\varepsilon p ||\nabla \psi||_{\infty}} \int_{T_0 - \varepsilon^2}^{T_1 + \varepsilon^2} \int_{\Omega_{\varepsilon}} |e^{\psi} \partial_t f|^p dx dt,
$$

where we have used the facts that $|\psi(x) - \psi(x - y)| \le ||\nabla \psi||_{\infty} |y|$ and $\theta_{\varepsilon}(y, s) = 0$ if $|y| > \varepsilon$ or $|s| > \varepsilon^2$. This, together with the estimate for J_2 , completes the proof.

Theorem 3.5. Let $F_{\varepsilon} = (F_{\varepsilon,i}^{\alpha})$ be given by [\(3.4\)](#page-9-2) and $1 \leq p < \infty$. Then for any $\Omega \subset \mathbb{R}^d$ and $(T_0, T_1) \subset \mathbb{R},$

$$
\int_{T_0}^{T_1} \int_{\Omega} |e^{\psi} F_{\varepsilon}|^p dx dt \le C e^{\varepsilon p \|\nabla \psi\|_{\infty}} \int_{T_0 - \varepsilon^2}^{T_1 + \varepsilon^2} \int_{\Omega_{\varepsilon}} \left\{ |e^{\psi} \nabla^2 u_0|^p + |e^{\psi} \partial_t u_0|^p \right\} dx dt, \tag{3.11}
$$

where Ω_{ε} is given by [\(3.7\)](#page-10-2) and C depends only on d, m, p and μ .

Proof. Observe that

$$
\int_{T_0}^{T_1} \int_{\Omega} |e^{\psi} F_{\varepsilon}|^p dx dt \leq C \varepsilon^{-p} \int_{T_0}^{T_1} \int_{\Omega} |\nabla u_0 - S_{\varepsilon} (\nabla u_0)|^p e^{p\psi} dx dt \n+ C \int_{T_0}^{T_1} \int_{\Omega} |\chi^{\varepsilon}|^p |S_{\varepsilon} (\nabla^2 u_0)|^p e^{p\psi} dx dt \n+ C \int_{T_0}^{T_1} \int_{\Omega} |\phi^{\varepsilon}|^p |S_{\varepsilon} (\nabla^2 u_0)|^p e^{p\psi} dx dt \n+ C \varepsilon^p \int_{T_0}^{T_1} \int_{\Omega} |\phi^{\varepsilon}|^p |\nabla S_{\varepsilon} (\partial_t u_0)|^p e^{p\psi} dx dt \n+ C \int_{T_0}^{T_1} \int_{\Omega} |(\nabla \phi)^{\varepsilon}|^p |S_{\varepsilon} (\nabla^2 u_0)|^p e^{p\psi} dx dt \n+ C \varepsilon^p \int_{T_0}^{T_1} \int_{\Omega} |\phi^{\varepsilon}|^p |\nabla S_{\varepsilon} (\nabla^2 u_0)|^p e^{p\psi} dx dt,
$$
\n(3.12)

where C depends only on d and μ . In [\(3.12\)](#page-13-0) we have also used the observation that $\partial_t S_{\varepsilon}(\nabla u_0) = \nabla S_{\varepsilon}(\partial_t u_0)$ and $\nabla S_{\varepsilon}(\nabla u_0) = S_{\varepsilon}(\nabla^2 u_0)$.

We now proceed to bound each term in the RHS of [\(3.12\)](#page-13-0), using Lemma [3.4](#page-11-0) and Remark [3.3.](#page-10-3) By Lemma [3.4,](#page-11-0) the first term in the RHS of [\(3.12\)](#page-13-0) is bounded by

$$
Ce^{p\varepsilon \|\nabla \psi\|_{\infty}} \int_{T_0 - \varepsilon^2}^{T_1 + \varepsilon^2} \int_{\Omega_{\varepsilon}} |e^{\psi} (|\nabla^2 u_0| + |\partial_t u_0|)|^p \, dx dt. \tag{3.13}
$$

Using [\(3.8\)](#page-10-4) we may bound the second, third, fifth terms in the RHS of [\(3.12\)](#page-13-0) by

$$
Ce^{p\varepsilon \|\nabla\psi\|_{\infty}} \int_{T_0 - e^2}^{T_1 + \varepsilon^2} \int_{\Omega_{\varepsilon}} |e^{\psi} \nabla^2 u_0|^p \, dxdt. \tag{3.14}
$$

Finally, by [\(3.9\)](#page-10-5), the fourth and sixth terms in the RHS of [\(3.12\)](#page-13-0) are bounded by [\(3.13\)](#page-13-1). This completes the proof. \Box

4 Weighted estimates for $\partial_t + \mathcal{L}_0$

Recall that $\Gamma_0(x, t; y, s)$ denotes the matrix of fundamental solutions for the homogenized operator $\partial_t + \mathcal{L}_0$ in \mathbb{R}^{d+1} . Since \mathcal{L}_0 is a second-order elliptic operator with constant coefficients, it is well known that $\Gamma_0(x, t; y, s) = \Gamma_0(x - y, t - s; 0, 0)$ and

$$
|\nabla_x^M \partial_t^N \Gamma_0(x, t; y, s)| \le \frac{C}{(t - s)^{\frac{d + M + 2N}{2}}} \exp\left\{-\frac{\kappa |x - y|^2}{t - s}\right\}
$$
(4.1)

for any $M, N \geq 0$, where $\kappa > 0$ depends only on μ , and C depends on d, m, M, N and μ .

Let $\psi : \mathbb{R}^d \to \mathbb{R}$ be a bounded Lipschitz function and

$$
u_0(x,t) = \int_{-\infty}^t \int_{\mathbb{R}^d} \Gamma_0(x,t;y,s) f(y,s) e^{-\psi(y)} dy ds,
$$
 (4.2)

where $f \in C_0^{\infty}(\mathbb{R}^{d+1}; \mathbb{R}^m)$. Then

$$
(\partial_t + \mathcal{L}_0) u_0 = e^{-\psi} f \quad \text{in } \mathbb{R}^{d+1}.
$$
 (4.3)

The goal of this section is to prove the following.

Theorem 4.1. Let u_0 be defined by [\(4.2\)](#page-13-2). Suppose that $f(x,t) = 0$ for $t \leq s_0$. Then

$$
\int_{s_0}^t \int_{\mathbb{R}^d} \left| e^{\psi} \left(|\nabla^2 u_0| + |\partial_t u_0| \right) \right|^2 dx dt \le C e^{\kappa (t - s_0) \|\nabla \psi\|_{\infty}^2} \int_{s_0}^t \int_{\mathbb{R}^d} |f|^2 dx dt \tag{4.4}
$$

for any $s_0 < t < \infty$, where $\kappa > 0$ depends only on μ and C depend only on d and μ .

We start with an estimate on a lower order term.

Lemma 4.2. Let u_0 be defined by [\(4.2\)](#page-13-2). Suppose that $f(x,t) = 0$ for $t < s_0$. Then

$$
\int_{s_0}^t \int_{\mathbb{R}^d} |e^{\psi} \nabla u_0|^2 dx dt \le C(t - s_0) e^{\kappa_1 (t - s_0) \|\nabla \psi\|_{\infty}^2} \int_{s_0}^t \int_{\mathbb{R}^d} |f|^2 dx dt \tag{4.5}
$$

for any $s_0 < t < \infty$, where $\kappa_1 > 0$ depends only on μ and C depend only on d and μ . *Proof.* It follows from [\(4.1\)](#page-13-3) that for $x, y \in \mathbb{R}^d$ and $t > s$,

$$
e^{\psi(x) - \psi(y)} |\nabla_x \Gamma_0(x, t; y, s)| \le \frac{C}{(t - s)^{\frac{d+1}{2}}} \exp \left\{ \psi(x) - \psi(y) - \frac{\kappa |x - y|^2}{t - s} \right\}
$$

$$
\le \frac{C}{(t - s)^{\frac{d+1}{2}}} \exp \left\{ ||\nabla \psi||_{\infty} |x - y| - \frac{\kappa |x - y|^2}{t - s} \right\}.
$$

This, together with the inequality,

$$
|\nabla \psi||_{\infty}|x-y| \le \frac{(t-s)\|\nabla \psi\|_{\infty}^2}{2\kappa} + \frac{\kappa|x-y|^2}{2(t-s)},\tag{4.6}
$$

yields

$$
e^{\psi(x) - \psi(y)} |\nabla_x \Gamma_0(x, t; y, s)| \le C e^{\frac{(t-s) ||\nabla \psi||_{\infty}^2}{2\kappa}} \cdot \frac{1}{(t-s)^{\frac{d+1}{2}}} e^{-\frac{\kappa |x - y|^2}{2(t-s)}}.
$$
\n(4.7)

It follows that

$$
|e^{\psi(x)}\nabla u_0(x,t)| \leq \int_{s_0}^t \int_{\mathbb{R}^d} e^{\psi(x) - \psi(y)} |\nabla_x \Gamma_0(x,t;y,s)| |f(y,s)| dy ds
$$

\n
$$
\leq C e^{\frac{(t-s_0) ||\nabla \psi||_{\infty}^2}{2\kappa}} \int_{s_0}^t \int_{\mathbb{R}^d} \frac{1}{(t-s)^{\frac{d+1}{2}}} e^{-\frac{\kappa |x-y|^2}{2(t-s)}} |f(y,s)| dy ds
$$

\n
$$
\leq C e^{\frac{(t-s_0) ||\nabla \psi||_{\infty}^2}{2\kappa}} (t-s_0)^{\frac{1}{4}} \left(\int_{s_0}^t \int_{\mathbb{R}^d} \frac{1}{(t-s)^{\frac{d+1}{2}}} e^{-\frac{\kappa |x-y|^2}{2(t-s)}} |f(y,s)|^2 dy ds \right)^{1/2},
$$

where we have used Hölder's inequality for the last step. The estimate (4.5) now follows by Fubini's Theorem. \Box **Proof of Theorem [4.1.](#page-14-1)** In view of (4.3) we have

$$
(\partial_t + \mathcal{L}_0) \frac{\partial u_0}{\partial x_k} = \frac{\partial}{\partial x_k} (e^{-\psi} f)
$$

in \mathbb{R}^{d+1} . It follows that

$$
\int_{\mathbb{R}^d} \partial_t \nabla u_0 \cdot (\nabla u_0) e^{2\psi} \, dx - \int_{\mathbb{R}^d} \widehat{a}_{ij}^{\alpha\beta} \frac{\partial^3 u_0^{\beta}}{\partial x_i \partial x_j \partial x_k} \cdot \frac{\partial u_0^{\alpha}}{\partial x_k} e^{2\psi} \, dx = \int_{\mathbb{R}^d} \frac{\partial}{\partial x_k} (e^{-\psi} f^{\alpha}) \frac{\partial u_0^{\alpha}}{\partial x_k} e^{2\psi} \, dx.
$$

Using integration by parts, we obtain

$$
\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^d}|\nabla u_0|^2 e^{2\psi} dx + \int_{\mathbb{R}^d} \widehat{a}_{ij}^{\alpha\beta} \frac{\partial^2 u_0^{\beta}}{\partial x_j \partial x_k} \cdot \frac{\partial^2 u_0^{\alpha}}{\partial x_i \partial x_k} e^{2\psi} dx \n= -\int_{\mathbb{R}^d} f \cdot (\Delta u_0) e^{\psi} dx - \int_{\mathbb{R}^d} e^{-\psi} f^{\alpha} \frac{\partial u_0^{\alpha}}{\partial x_k} \frac{\partial e^{2\psi}}{\partial x_k} dx - \int_{\mathbb{R}^d} \widehat{a}_{ij}^{\alpha\beta} \frac{\partial^2 u_0^{\beta}}{\partial x_j \partial x_k} \cdot \frac{\partial u_0^{\alpha}}{\partial x_k} \frac{\partial e^{2\psi}}{\partial x_i} dx.
$$

By the ellipticity of \mathcal{L}_0 , this yields

$$
\frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^d} |\nabla u_0|^2 e^{2\psi} dx + \mu \int_{\mathbb{R}^d} |\nabla^2 u_0|^2 e^{2\psi} dx \n\leq C \int_{\mathbb{R}^d} |f| |\nabla^2 u_0| e^{\psi} dx + C \int_{\mathbb{R}^d} |f|^2 dx + C ||\nabla \psi||_{\infty}^2 \int_{\mathbb{R}^d} |\nabla u_0|^2 e^{2\psi} dx \n+ C ||\nabla \psi||_{\infty} \int_{\mathbb{R}^d} |\nabla^2 u_0| |\nabla u_0| e^{2\psi} dx,
$$

where C depends only on d and μ . Using Cauchy inequality, we may further deduce that

$$
\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^d} |\nabla u_0|^2 e^{2\psi} \, dx + \frac{\mu}{2} \int_{\mathbb{R}^d} |\nabla^2 u_0|^2 e^{2\psi} \, dx \n\leq C \int_{\mathbb{R}^d} |f|^2 \, dx + C \|\nabla \psi\|_{\infty}^2 \int_{\mathbb{R}^d} |\nabla u_0|^2 e^{2\psi} \, dx.
$$

We now integrate the inequality above in t over the interval (s_0, s_1) . This leads to

$$
\frac{1}{2} \int_{\mathbb{R}^d} |\nabla u_0(x, s_1)|^2 e^{2\psi} dx + \frac{\mu}{2} \int_{s_0}^{s_1} \int_{\mathbb{R}^d} |\nabla^2 u_0|^2 e^{2\psi} dx dt \n\leq C \int_{s_0}^{s_1} \int_{\mathbb{R}^d} |f|^2 dx dt + C \|\nabla \psi\|_{\infty}^2 \int_{s_0}^{s_1} \int_{\mathbb{R}^d} |\nabla u_0|^2 e^{2\psi} dx dt \n\leq C e^{\kappa (s_1 - s_0) \|\nabla \psi\|_{\infty}^2} \int_{s_0}^{s_1} \int_{\mathbb{R}^d} |f|^2 dx dt,
$$
\n(4.8)

where we have used [\(4.5\)](#page-14-0) for the last inequality. Estimate [\(4.4\)](#page-14-3) follows readily from [\(4.8\)](#page-15-0). \Box

5 Proof of Theorem [1.1](#page-1-5)

We start with some weighted estimates.

Lemma 5.1. Suppose that

$$
\begin{cases}\n(\partial_t + \mathcal{L}_{\varepsilon})w_{\varepsilon} = \varepsilon \operatorname{div}(F_{\varepsilon}) & \text{in } \mathbb{R}^d \times (s_0, \infty), \\
w_{\varepsilon} = 0 & \text{on } \mathbb{R}^d \times \{t = s_0\}.\n\end{cases}
$$
\n(5.1)

Let $\psi : \mathbb{R}^d \to \mathbb{R}$ be a bounded Lipschitz function. Then for any $t > s_0$,

$$
\int_{\mathbb{R}^d} |w_{\varepsilon}(x,t)|^2 e^{2\psi(x)} dx \le C\varepsilon^2 e^{\kappa(t-s_0)\|\nabla\psi\|_{\infty}^2} \int_{s_0}^t \int_{\mathbb{R}^d} |F_{\varepsilon}(x,s)|^2 e^{2\psi(x)} dx ds, \tag{5.2}
$$

where $\kappa > 0$ and $C > 0$ depends only on μ . Proof. Let

$$
I(t) = \int_{\mathbb{R}^d} |w_{\varepsilon}(x,t)|^2 e^{2\psi(x)} dx.
$$
 (5.3)

Note that

$$
I'(t) = 2 \int_{\mathbb{R}^d} \langle \partial_t w_{\varepsilon}, e^{2\psi} w_{\varepsilon} \rangle dx
$$

\n
$$
= -2 \int_{\mathbb{R}^d} \langle \mathcal{L}_{\varepsilon}(w_{\varepsilon}), e^{2\psi} w_{\varepsilon} \rangle dx + 2\varepsilon \int_{\mathbb{R}^d} \langle \operatorname{div}(F_{\varepsilon}), e^{2\psi} w_{\varepsilon} \rangle dx
$$

\n
$$
= -2 \int_{\mathbb{R}^d} A^{\varepsilon} \nabla w_{\varepsilon} \cdot \nabla (e^{2\psi} w_{\varepsilon}) dx - 2\varepsilon \int_{\mathbb{R}^d} F_{\varepsilon} \cdot \nabla (e^{2\psi} w_{\varepsilon}) dx
$$

\n
$$
= -2 \int_{\mathbb{R}^d} A^{\varepsilon} \nabla w_{\varepsilon} \cdot (\nabla w_{\varepsilon}) e^{2\psi} dx - 2 \int_{\mathbb{R}^d} A^{\varepsilon} \nabla w_{\varepsilon} \cdot \nabla (e^{2\psi}) w_{\varepsilon} dx
$$

\n
$$
- 2\varepsilon \int_{\mathbb{R}^d} F_{\varepsilon} \cdot (\nabla w_{\varepsilon}) e^{2\psi} dx - 2\varepsilon \int_{\mathbb{R}^d} F_{\varepsilon} \cdot \nabla (e^{2\psi}) w_{\varepsilon} dx,
$$

where \langle , \rangle denotes the pairing in $H^{-1}(\mathbb{R}^d; \mathbb{R}^m) \times H^1(\mathbb{R}^d; \mathbb{R}^m)$. It follows that

$$
I'(t) \leq -2\mu \int_{\mathbb{R}^d} |\nabla w_{\varepsilon}|^2 e^{2\psi} dx + \kappa ||\nabla \psi||_{\infty} \int_{\mathbb{R}^d} |\nabla w_{\varepsilon}| |w_{\varepsilon}| e^{2\psi} dx + 2\varepsilon \int_{\mathbb{R}^d} |\nabla w_{\varepsilon}| |F_{\varepsilon}| e^{2\psi} dx + 4\varepsilon ||\nabla \psi||_{\infty} \int_{\mathbb{R}^d} |w_{\varepsilon}| |F_{\varepsilon}| e^{2\psi} dx,
$$

where $\kappa > 0$ depends only on μ . By Cauchy inequality this implies that

$$
I'(t) \leq \kappa \|\nabla \psi\|_{\infty}^2 I(t) + \kappa \varepsilon^2 \int_{\mathbb{R}^d} |F_{\varepsilon}(x, t)|^2 e^{2\psi} dx,
$$
\n(5.4)

where $\kappa > 0$ depends only on μ . Hence,

$$
\frac{d}{dt}\left\{I(t)e^{-\kappa(t-s_0)\|\nabla\psi\|_{\infty}^2}\right\} \leq C\varepsilon^2 e^{-\kappa(t-s_0)\|\nabla\psi\|_{\infty}^2} \int_{\mathbb{R}^d} |F_{\varepsilon}(x,t)|^2 e^{2\psi} dx.
$$

Since $I(s_0) = 0$, it follows that

$$
I(t) \leq C\varepsilon^2 \int_{s_0}^t \int_{\mathbb{R}^d} e^{\kappa(t-s) \|\nabla \psi\|_{\infty}^2} |F_{\varepsilon}(x,s)|^2 e^{2\psi} dx ds
$$

$$
\leq C\varepsilon^2 e^{\kappa(t-s_0) \|\nabla \psi\|_{\infty}^2} \int_{s_0}^t \int_{\mathbb{R}^d} |F_{\varepsilon}(x,s)|^2 e^{2\psi} dx ds.
$$

This completes the proof.

Lemma 5.2. Suppose that $u_{\varepsilon} \in L^2((-\infty,T); H^1(\mathbb{R}^d))$ and $u_0 \in L^2((-\infty,T); H^2(\mathbb{R}^d))$ for any $T \in \mathbb{R}$, that

$$
\begin{cases}\n(\partial_t + \mathcal{L}_\varepsilon)u_\varepsilon = (\partial_t + \mathcal{L}_0)u_0 & \text{in } \mathbb{R}^{d+1}, \\
u_\varepsilon(x,t) = u_0(x,t) = 0 & \text{for } t \le s_0.\n\end{cases}
$$

Let w_{ε} be defined by [\(3.2\)](#page-9-0), where the operator S_{ε} is given by [\(3.5\)](#page-9-1). Then for any $t > s_0$,

$$
\int_{\mathbb{R}^d} |w_{\varepsilon}(x,t)|^2 e^{2\psi(x)} dx
$$
\n
$$
\leq C \varepsilon^2 e^{2\varepsilon \|\nabla \psi\|_{\infty} + \kappa (t-s_0) \|\nabla \psi\|_{\infty}^2} \int_{s_0}^{t+\varepsilon^2} \int_{\mathbb{R}^d} \left\{ |\nabla^2 u_0(x,s)|^2 + |\partial_s u_0(x,s)|^2 \right\} e^{2\psi(x)} dx ds,
$$
\n(5.5)

where ψ is a bounded Lipschitz function in \mathbb{R}^d , κ depends only on μ , and C depends only on d, m and μ .

Proof. This follows readily from Lemma [5.1](#page-16-0) and Theorem [3.5](#page-12-0) with $p = 2$.

The next theorem gives a weighted L^{∞} estimate.

Theorem 5.3. Assume that A is 1-periodic and satisfies [\(1.2\)](#page-0-1). If $m \geq 2$, we also assume that $A \in VMO_{x}$. Suppose that $(\partial_t + \mathcal{L}_{\varepsilon})u_{\varepsilon} = (\partial_t + \mathcal{L}_0)u_0$ in $B(x_0, 3r) \times (t_0 - 5r^2, t_0 + r^2)$ for some $(x_0, t_0) \in \mathbb{R}^{d+1}$ and $\varepsilon \le r < \infty$. Then

$$
\|e^{\psi}(u_{\varepsilon} - u_0)\|_{L^{\infty}(Q_r(x_0, t_0))}\n\n\leq Ce^{3r\|\nabla\psi\|_{\infty}} \left(\int_{Q_{2r}(x_0, t_0)} |e^{\psi}(u_{\varepsilon} - u_0)|^2\right)^{1/2}\n\n+ C\varepsilon r e^{3r\|\nabla\psi\|_{\infty}} \|e^{\psi}(|\nabla^2 u_0| + |\partial_t u_0|)\|_{L^{\infty}(B(x_0, 3r)\times(t_0 - 5r^2, t_0 + r^2))}\n\n+ C\varepsilon e^{3r\|\nabla\psi\|_{\infty}} \|e^{\psi}\nabla u_0\|_{L^{\infty}(B(x_0, 3r)\times(t_0 - 5r^2, t_0 + r^2))},
$$
\n(5.6)

 \Box

where ψ is a Lipschitz function in \mathbb{R}^d and C depends only on d, m, μ and $A^{\#}$ (if $m \geq 2$).

Proof. Let w_{ε} be defined by [\(3.2\)](#page-9-0). Then $(\partial_t + \mathcal{L}_{\varepsilon})w_{\varepsilon} = \varepsilon \operatorname{div}(F_{\varepsilon})$ in $Q_{2r}(x_0, t_0)$, where F_{ε} is given by [\(3.4\)](#page-9-2). It follows by Theorem [2.5](#page-7-2) that

$$
||w_{\varepsilon}||_{L^{\infty}(Q_r(x_0,t_0))} \leq C \left\{ \left(\int_{Q_{2r}(x_0,t_0)} |w_{\varepsilon}|^2 \right)^{1/2} + \varepsilon r \left(\int_{Q_{2r}(x_0,t_0)} |F_{\varepsilon}|^p \right)^{1/p} \right\},\qquad(5.7)
$$

where $p > d + 2$. This leads to

$$
||u_{\varepsilon}-u_0||_{L^{\infty}(Q_r(x_0,t_0))} \leq C \left(\int_{Q_{2r}(x_0,t_0)} |u_{\varepsilon}-u_0|^2 \right)^{1/2} + C\varepsilon r \left(\int_{Q_{2r}(x_0,t_0)} |F_{\varepsilon}|^p \right)^{1/p} + C\varepsilon ||S_{\varepsilon}(\nabla u_0)||_{L^{\infty}(Q_{2r}(x_0,t_0))} + C\varepsilon^2 ||S_{\varepsilon}(\nabla^2 u_0)||_{L^{\infty}(Q_{2r}(x_0,t_0))},
$$

where we have used the boundedness of χ and ϕ in Lemmas [2.1](#page-5-4) and [2.3.](#page-7-3) Hence, using $|\psi(x) - \psi(y)| \leq 2r \|\nabla \psi\|_{\infty}$ for $x, y \in B(x_0, 2r)$, we obtain

$$
||e^{\psi}(u_{\varepsilon}-u_{0})||_{L^{\infty}(Q_{r}(x_{0},t_{0}))} \leq Ce^{2r||\nabla\psi||_{\infty}} \left(\int_{Q_{2r}(x_{0},t_{0})} |e^{\psi}(u_{\varepsilon}-u_{0})|^{2}\right)^{1/2} + C\varepsilon r e^{2r||\nabla\psi||_{\infty}} \left(\int_{Q_{2r}(x_{0},t_{0})} |e^{\psi}F_{\varepsilon}|^{p}\right)^{1/p} + C\varepsilon e^{2r||\nabla\psi||_{\infty}} ||e^{\psi}S_{\varepsilon}(\nabla u_{0})||_{L^{\infty}(Q_{2r}(x_{0},t_{0}))} + C\varepsilon^{2} e^{2r||\nabla\psi||_{\infty}} ||e^{\psi}S_{\varepsilon}(\nabla^{2}u_{0})||_{L^{\infty}(Q_{2r}(x_{0},t_{0}))}.
$$
\n
$$
(5.8)
$$

Finally, we use Theorem [3.5](#page-12-0) to bound the second term in the RHS of [\(5.8\)](#page-18-0). This yields

$$
\|e^{\psi}(u_{\varepsilon}-u_{0})\|_{L^{\infty}(Q_{r}(x_{0},t_{0}))}
$$
\n
$$
\leq Ce^{2r\|\nabla\psi\|_{\infty}}\left(\int_{Q_{2r}(x_{0},t_{0})}|e^{\psi}(u_{\varepsilon}-u_{0})|^{2}\right)^{1/2}
$$
\n
$$
+ C\varepsilon r e^{3r\|\nabla\psi\|_{\infty}}\left(\int_{t_{0}-5r^{2}}^{t_{0}+r^{2}}\int_{B(x_{0},3r)}\left\{|e^{\psi}\nabla^{2}u_{0}|^{p}+|e^{\psi}\partial_{t}u_{0}|^{p}\right\}\right)^{1/p}
$$
\n
$$
+ C\varepsilon e^{2r\|\nabla\psi\|_{\infty}}\|e^{\psi}S_{\varepsilon}(\nabla u_{0})\|_{L^{\infty}(Q_{2r}(x_{0},t_{0}))} + C\varepsilon^{2}e^{2r\|\nabla\psi\|_{\infty}}\|e^{\psi}S_{\varepsilon}(\nabla^{2}u_{0})\|_{L^{\infty}(Q_{2r}(x_{0},t_{0}))},
$$

 \Box

where $p > d + 2$ and we also used the assumption $\varepsilon \le r$. Estimate [\(5.6\)](#page-17-0) now follows.

We are now in a position to give the proof of Theorem [1.1.](#page-1-5)

Proof of Theorem [1.1.](#page-1-5) We begin by fixing $x_0, y_0 \in \mathbb{R}^{d+1}$ and $s_0 < t_0$. We may assume that

$$
\varepsilon < r = (t_0 - s_0)^{1/2} / 100.
$$

For otherwise the desired estimate [\(1.7\)](#page-1-4) follows directly from [\(1.6\)](#page-1-1).

For $f \in C_0^{\infty}(Q_r(y_0, s_0); \mathbb{R}^m)$, define

$$
u_{\varepsilon}(x,t) = \int_{-\infty}^{t} \int_{\mathbb{R}^d} e^{-\psi(y)} \Gamma_{\varepsilon}(x,t;y,s) f(y,s) dyds,
$$

$$
u_0(x,t) = \int_{-\infty}^{t} \int_{\mathbb{R}^d} e^{-\psi(y)} \Gamma_0(x,t;y,s) f(y,s) dyds,
$$

where ψ is a bounded Lipschitz function in \mathbb{R}^d to be chosen later. Then

$$
(\partial_t + \mathcal{L}_\varepsilon)u_\varepsilon = (\partial_t + \mathcal{L}_0)u_0 = e^{-\psi}f \quad \text{in } \mathbb{R}^{d+1}
$$

and $u_{\varepsilon}(x,t) = u_0(x,t) = 0$ if $t \leq s_0$. Let w_{ε} be defined by [\(3.2\)](#page-9-0). It follows from Lemma [5.2](#page-17-1) and Theorem [4.1](#page-14-1) that

$$
\int_{\mathbb{R}^d} |w_{\varepsilon}(x,t)|^2 e^{2\psi(x)} dx \le C\varepsilon^2 e^{2\varepsilon \|\nabla\psi\|_{\infty} + \kappa(t - s_0 + \varepsilon^2)\|\nabla\psi\|_{\infty}^2} \int_{s_0}^{t + \varepsilon^2} \int_{\mathbb{R}^d} |f|^2 dx ds \tag{5.9}
$$

for any $t > s_0$.

Next, we use [\(5.6\)](#page-17-0) to obtain

$$
\|e^{\psi}(u_{\varepsilon} - u_0)\|_{L^{\infty}(Q_r(x_0, t_0))}\n\n\leq Ce^{3r\|\nabla\psi\|_{\infty}} \left(\int_{Q_{2r}(x_0, t_0)} |e^{\psi}w_{\varepsilon}|^2\right)^{1/2}\n\n+ C\varepsilon r e^{3r\|\nabla\psi\|_{\infty}} \|e^{\psi}(|\nabla^2 u_0| + |\partial_t u_0|)\|_{L^{\infty}(B(x_0, 3r)\times(t_0 - 5r^2, t_0 + r^2))}\n\n+ C\varepsilon e^{3r\|\nabla\psi\|_{\infty}} \|e^{\psi}\nabla u_0\|_{L^{\infty}(B(x_0, 3r)\times(t_0 - 5r^2, t_0 + r^2))}.
$$
\n(5.10)

Since supp $(f) \subset Q_r(y_0, s_0)$, it follows from the estimate (4.1) for $\Gamma_0(x, t; y, s)$ that

$$
|\nabla^2 u_0(x,t)| + |\partial_t u_0(x,t)| + r^{-1} |\nabla u_0(x,t)|
$$

\n
$$
\leq C \exp\left\{-\frac{\kappa |x_0 - y_0|^2}{t_0 - s_0}\right\} \int_{Q_r(y_0, s_0)} |f e^{-\psi}| dy ds \tag{5.11}
$$

for any $x \in B(x_0, 3r)$ and $|t - t_0| \leq 5r^2$, where $\kappa > 0$ depends only on μ . Thus, by [\(5.10\)](#page-19-0), we obtain

$$
\|e^{\psi}(u_{\varepsilon} - u_0)\|_{L^{\infty}(Q_r(x_0, t_0))}\n\n\leq Ce^{3r\|\nabla\psi\|_{\infty}} \left(\int_{Q_{2r}(x_0, t_0)} |e^{\psi}w_{\varepsilon}|^2\right)^{1/2}\n\n+ \varepsilon r e^{c(|x_0 - y_0| + r)\|\nabla\psi\|_{\infty}} \exp\left\{-\frac{\kappa |x_0 - y_0|^2}{t_0 - s_0}\right\} \int_{Q_r(y_0, s_0)} |f| \, dy ds\n\n\leq C \varepsilon r e^{cr\|\nabla\psi\|_{\infty}} \left\{e^{cr^2\|\nabla\psi\|_{\infty}^2} + e^{c|x_0 - y_0|\|\nabla\psi\|_{\infty}} \exp\left\{-\frac{\kappa |x_0 - y_0|^2}{t_0 - s_0}\right\}\right\}\n\n\cdot \left(\int_{Q_r(y_0, s_0)} |f|^2\right)^{1/2},
$$
\n(5.12)

where we have used (5.9) for the last step. By duality this implies that

$$
\left(\oint_{Q_r(y_0,s_0)} \left| e^{\psi(x)-\psi(y)} \left(\Gamma_{\varepsilon}(x,t,y,s) - \Gamma_0(x,t;y,s) \right) \right|^2 dyds \right)^{1/2}
$$
\n
$$
\leq C\varepsilon r^{-d-1} e^{cr \|\nabla\psi\|_{\infty}} \left\{ e^{cr^2 \|\nabla\psi\|_{\infty}^2} + e^{c|x_0-y_0|\|\nabla\psi\|_{\infty}} \exp\left\{ -\frac{\kappa |x_0-y_0|^2}{t_0-s_0} \right\} \right\}
$$
\n(5.13)

for any $(x, t) \in Q_r(x_0, t_0)$.

To deduce the L^{∞} bound for

$$
e^{\psi(x)-\psi(y)}\big(\Gamma_{\varepsilon}(x,t;y,s)-\Gamma_0(x,t;y,s)\big)
$$

from its \mathcal{L}^2 bound in [\(5.13\)](#page-19-1), we apply Theorem [5.3](#page-17-2) (with ψ replaced by $-\psi$ and A replaced by $A = A(y, s) = A^*(y, -s)$ to the functions

$$
v_{\varepsilon}(y, s) = \Gamma_{\varepsilon}(x_0, t_0; y, -s)
$$
 and $v_0(y, s) = \Gamma_0(x_0, t_0, y, -s)$.

Note that $(\partial_t + \mathcal{L}_\varepsilon)v_\varepsilon = (\partial_t + \mathcal{L}_0)v_0 = 0$ in $B(y_0, 3r) \times (-s_0 - 5r^2, -s_0 + r^2)$. Since A satisfies the same conditions as A , we obtain

$$
\begin{split}\n&|e^{\psi(x_{0})-\psi(y_{0})}(v_{\varepsilon}(y_{0},-s_{0})-v_{0}(y_{0},-s_{0}))| \\
&\leq Ce^{3r\|\nabla\psi\|_{\infty}}\left(\int_{Q_{r}(y_{0},-s_{0})}|e^{\psi(x_{0})-\psi(y)}(v_{\varepsilon}-v_{0})|^{2}dyds\right)^{1/2} \\
&+ C\varepsilon r^{-d-1}e^{cr\|\nabla\psi\|_{\infty}}e^{\psi(x_{0})-\psi(y_{0})}\exp\left\{-\frac{\kappa|x_{0}-y_{0}|^{2}}{t_{0}-s_{0}}\right\} \\
&= Ce^{3r\|\nabla\psi\|_{\infty}}\left(\int_{Q_{r}(y_{0},s_{0}+r^{2})}|e^{\psi(x_{0})-\psi(y)}(\Gamma_{\varepsilon}(x_{0},t_{0};y,s)-\Gamma_{0}(x_{0},t_{0};y,s))|^{2}dyds\right)^{1/2} \quad(5.14) \\
&+ C\varepsilon r^{-d-1}e^{cr\|\nabla\psi\|_{\infty}}e^{\psi(x_{0})-\psi(y_{0})}\exp\left\{-\frac{\kappa|x_{0}-y_{0}|^{2}}{t_{0}-s_{0}}\right\} \\
&\leq C\varepsilon r^{-d-1}e^{cr\|\nabla\psi\|_{\infty}}\left\{e^{cr^{2}\|\nabla\psi\|_{\infty}}+e^{c|x_{0}-y_{0}|\|\nabla\psi\|_{\infty}}\exp\left\{-\frac{\kappa|x_{0}-y_{0}|^{2}}{t_{0}-s_{0}}\right\}\right\},\n\end{split}
$$

where we have used [\(5.13\)](#page-19-1) for the last inequality.

Finally, as in [\[12,](#page-24-5) [6\]](#page-24-6), we let $\psi(y) = \gamma \psi_0(|y - y_0|)$, where $\gamma \ge 0$ is to be chosen, $\psi_0(\rho) = \rho$ if $\rho \leq |x_0 - y_0|$, and $\psi_0(\rho) = |x_0 - y_0|$ if $\rho > |x_0 - y_0|$. Note that $\|\nabla \psi\|_{\infty} = \gamma$ and $\psi(x_0) - \psi(y_0) = \gamma |x_0 - y_0|$. It follows from [\(5.14\)](#page-20-0) that

$$
\begin{split} &\left|\Gamma_{\varepsilon}(x_0, t_0; y_0, s_0) - \Gamma_0(x_0, t_0; y_0, s_0)\right| \\ &\leq C \varepsilon r^{-d-1} e^{-\gamma |x_0 - y_0| + c\gamma \sqrt{t_0 - s_0}} \left\{ e^{c\gamma^2 (t_0 - s_0)} + e^{c\gamma |x_0 - y_0|} \exp\left\{ -\frac{\kappa |x_0 - y_0|^2}{t_0 - s_0} \right\} \right\}, \end{split} \tag{5.15}
$$

where $c > 0$ depends at most on μ . If $|x_0 - y_0| \leq 2c\sqrt{t_0 - s_0}$, we may simply choose $\gamma = 0$. This gives

$$
\begin{aligned} |\Gamma_{\varepsilon}(x_0, t_0; y_0, s_0) - \Gamma_0(x_0, t_0; y_0, s_0) &\leq C \varepsilon r^{-d-1} \\ &\leq C \varepsilon (t_0 - s_0)^{-\frac{d+1}{2}} \exp\left\{-\frac{\kappa |x_0 - y_0|^2}{t_0 - s_0}\right\}. \end{aligned}
$$

If $|x_0 - y_0| > 2c\sqrt{t_0 - s_0}$, we choose

$$
\gamma = \frac{\delta |x_0 - y_0|}{t_0 - s_0}.
$$

Note that

$$
-\gamma |x_0 - y_0| + c\gamma \sqrt{t_0 - s_0} + c\gamma^2 (t_0 - s_0)
$$

= $-\delta (1 - c\delta) \frac{|x_0 - y_0|^2}{t_0 - s_0} + c\delta \frac{|x_0 - y_0|}{\sqrt{t_0 - s_0}}$

$$
\leq \left\{ -\delta (1 - c\delta) + (1/2)\delta \right\} \frac{|x_0 - y_0|^2}{t_0 - s_0}
$$

$$
\leq \frac{-\delta |x_0 - y_0|^2}{4(t_0 - s_0)},
$$

if $\delta \leq (1/4)c^{-1}$. Also, observe that

$$
c\gamma\sqrt{t_0 - s_0} + c\gamma |x_0 - y_0| - \frac{\kappa |x_0 - y_0|^2}{t_0 - s_0} \le \left\{ (1/2)\delta + c\delta - \kappa \right\} \frac{|x_0 - y_0|^2}{t_0 - s_0} \le -\frac{\kappa |x_0 - y_0|^2}{2(t_0 - s_0)},
$$

if $\delta \leq (1/2)(c+1/2)^{-1}\kappa$. Recall that $r = (100)^{-1}\sqrt{t_0 - s_0}$. As a result, we have proved that there exists $\kappa_1 > 0$, depending only on μ , such that

$$
|\Gamma_{\varepsilon}(x_0,t_0;y_0,s_0)-\Gamma_0(x_0,t_0;y_0,s_0)|\leq \frac{C\varepsilon}{(t_0-s_0)^{\frac{d+1}{2}}}\exp\left\{-\frac{\kappa_1|x_0-y_0|^2}{t_0-s_0}\right\}.
$$

This completes the proof of Theorem [1.1.](#page-1-5)

6 Proof of Theorem [1.2](#page-1-3)

Define

$$
||F||_{C^{\lambda,0}(K)} = \sup \left\{ \frac{|F(x,t) - F(y,t)|}{|x - y|^{\lambda}} : (x,t), (y,t) \in K \text{ and } x \neq y \right\}.
$$

The proof of Theorem [1.2](#page-1-3) relies on the following Lipschitz estimate.

Theorem 6.1. Assume that A satisfies conditions (1.2) , (1.3) and (1.8) . Suppose that

$$
(\partial_t + \mathcal{L}_\varepsilon)u_\varepsilon = (\partial_t + \mathcal{L}_0)u_0
$$

in $Q_{2r}(x_0, t_0)$ for some $(x_0, t_0) \in \mathbb{R}^{d+1}$ and $\varepsilon \le r < \infty$. Then

$$
\|\nabla u_{\varepsilon} - \nabla u_{0} - (\nabla \chi)^{\varepsilon} \nabla u_{0}\|_{L^{\infty}(Q_{r}(x_{0},t_{0}))}
$$
\n
$$
\leq C r^{-1} \left(\int_{Q_{2r}(x_{0},t_{0})} |u_{\varepsilon} - u_{0}|^{2} \right)^{1/2} + C \varepsilon r^{-1} \|\nabla u_{0}\|_{L^{\infty}(Q_{2r}(x_{0},t_{0}))}
$$
\n
$$
+ C \varepsilon \ln \left[\varepsilon^{-1} r + 2 \right] \|\nabla^{2} u_{0}\| + \varepsilon |\partial_{t} \nabla u_{0}| + \varepsilon |\nabla^{3} u_{0}| \|_{L^{\infty}(Q_{2r}(x_{0},t_{0}))}
$$
\n
$$
+ C \varepsilon^{1+\lambda} \|\nabla^{2} u_{0}\| + \varepsilon |\partial_{t} \nabla u_{0}| + \varepsilon |\nabla^{3} u_{0}| \|_{C^{\lambda,0}(Q_{2r}(x_{0},t_{0}))},
$$
\n
$$
(6.1)
$$

where C depends only on d, m, μ and (λ, τ) in [\(1.8\)](#page-1-2).

Proof. Let

$$
w_{\varepsilon} = u_{\varepsilon} - u_0 - \varepsilon \chi_j^{\varepsilon} \frac{\partial u_0}{\partial x_j} - \varepsilon^2 \phi_{(d+1)ij}^{\varepsilon} \frac{\partial^2 u_0}{\partial x_i \partial x_j},
$$
(6.2)

where $\chi_j^{\varepsilon}(x,t) = \chi_j(x/\varepsilon,t/\varepsilon^2)$ and $\phi_{(d+1)ij}^{\varepsilon}(x,t) = \phi_{(d+1)ij}(x/\varepsilon,t/\varepsilon^2)$. It follows by Proposition 3.1 that $(\partial_t + \mathcal{L}_{\varepsilon})w_{\varepsilon} = \varepsilon \operatorname{div}(F_{\varepsilon})$ in $\widetilde{Q}_{2r}(x_0, t_0)$, where F_{ε} is given by [\(3.4\)](#page-9-2) with S_{ε} being the identity operator. Choose a cut-off function $\varphi \in C(\mathbb{R}^{d+1})$ such that

$$
\begin{cases}\n0 \le \varphi \le 1, & \varphi = 1 \text{ in } Q_{3r/2}(x_0, t_0), \\
\varphi(x, t) = 0 \text{ if } |x - x_0| \ge (7r/4) \text{ or } t < t_0 - (7r/4)^2, \\
|\nabla \varphi| \le Cr^{-1}, & |\nabla^2 \varphi| + |\partial_t \varphi| \le Cr^{-2}.\n\end{cases}
$$

 \Box

Using

$$
(\partial_t + \mathcal{L}_{\varepsilon})(\varphi w_{\varepsilon}) = (\partial_t \varphi)w_{\varepsilon} + \varepsilon \operatorname{div}(\varphi F_{\varepsilon}) - \varepsilon F_{\varepsilon}(\nabla \varphi) - \operatorname{div}(A^{\varepsilon}(\nabla \varphi)w_{\varepsilon}) - A^{\varepsilon} \nabla w_{\varepsilon}(\nabla \varphi),
$$

where $A^{\varepsilon}(x,t) = A(x/\varepsilon,t/\varepsilon^2)$, we may deduce that for any $(x,t) \in Q_r(x_0,t_0)$,

$$
w_{\varepsilon}(x,t) = \int_{-\infty}^{t} \int_{\mathbb{R}^{d}} \Gamma_{\varepsilon}(x,t;y,s) \{ (\partial_{s}\varphi)w_{\varepsilon} - \varepsilon F_{\varepsilon}(\nabla\varphi) - A^{\varepsilon}\nabla w_{\varepsilon}(\nabla\varphi) \} dy ds - \int_{-\infty}^{t} \int_{\mathbb{R}^{d}} \nabla_{y} \Gamma_{\varepsilon}(x,t;y,s) \{ \varepsilon \varphi F_{\varepsilon} + A^{\varepsilon}(\nabla\varphi)w_{\varepsilon} \} dy ds = I(x,t) + J(x,t),
$$

where

$$
J(x,t) = -\varepsilon \int_{-\infty}^{t} \int_{\mathbb{R}^d} \nabla_y \Gamma_{\varepsilon}(x,t;y,s) \varphi(y,s) F_{\varepsilon}(y,s) dyds.
$$

Since $\varphi = 1$ in $Q_{3r/2}(x_0, t_0)$, we see that for $(x, t) \in Q_r(x_0, t_0)$,

$$
\begin{split} |\nabla I(x,t)| &\leq C\int_{-\infty}^t \int_{\mathbb{R}^d} |\nabla_x \Gamma_{\varepsilon}(x,t;y,s)| \{ |\partial_s \varphi| |w_{\varepsilon}| + \varepsilon |F_{\varepsilon}| |\nabla \varphi| + |\nabla w_{\varepsilon}| |\nabla \varphi| \} dy ds \\ &\quad + C\int_{-\infty}^t \int_{R^d} |\nabla_x \nabla_y \Gamma_{\varepsilon}(x,t;y,s)| |\nabla \varphi| |w_{\varepsilon}| dy ds \\ &\leq C\left\{ \frac{1}{r} \int_{Q_{2r}(x_0,t_0)} |w_{\varepsilon}| + \varepsilon \int_{Q_{2r}(x_0,t_0)} |F_{\varepsilon}| + \int_{Q_{7r/4}(x_0,t_0)} |\nabla w_{\varepsilon}| \right\} \\ &\leq C\left\{ \frac{1}{r} \left(\int_{Q_{2r}(x_0,t_0)} |w_{\varepsilon}|^2 \right)^{1/2} + \varepsilon \left(\int_{Q_{2r}(x_0,t_0)} |F_{\varepsilon}|^2 \right)^{1/2} \right\}, \end{split}
$$

where we have used (parabolic) Cacciopoli's inequality for the last step. In view of [\(3.4\)](#page-9-2) with S_{ε} being the identity operator,

$$
|F_{\varepsilon}| \le C \{ |\nabla^2 u_0| + \varepsilon |\partial_t \nabla u_0| + \varepsilon |\nabla^3 u_0| \},
$$

where we have used the boundedness of $\nabla \phi$ (see Remark [2.4\)](#page-7-4). It follows that $\|\nabla I\|_{L^{\infty}(Q_r(x_0,t_0))}$ is bounded by the RHS of [\(6.1\)](#page-21-0).

Finally, to estimate $J(x, t)$, we write

$$
J(x,t) = -\varepsilon \int_{-\infty}^{t} \int_{\mathbb{R}^d} \nabla_y \{ \Gamma_{\varepsilon}(x,t;y,s) \varphi(y,s) \} \left(F_{\varepsilon}(y,s) - F_{\varepsilon}(x,s) \right) dy ds + \varepsilon \int_{-\infty}^{t} \int_{\mathbb{R}^d} \Gamma_{\varepsilon}(x,t;y,s) (\nabla \varphi)(y,s) F_{\varepsilon}(y,s) dy ds.
$$

It follows that for $(x, t) \in Q_r(x_0, t_0)$,

$$
|\nabla J(x,t)| \leq \varepsilon \int_{Q_{2r}(x_0,t_0)} |\nabla_x \nabla_y \{ \Gamma_{\varepsilon}(x,t;y,s)\varphi(y,s) \}||F_{\varepsilon}(y,s) - F_{\varepsilon}(x,s)| dyds
$$

+
$$
\varepsilon \int_{Q_{2r}(x_0,t_0)} |\nabla_x \Gamma_{\varepsilon}(x,t;y,s)||\nabla \varphi(y,s)||F_{\varepsilon}(y,s)| dyds
$$
(6.3)

$$
\leq C\varepsilon \int_{Q_{2r}(x_0,t_0)} \frac{|F_{\varepsilon}(y,s) - F_{\varepsilon}(x,s)|}{(|x-y| + |t-s|^{1/2})^{d+2}} dyds + C\varepsilon \int_{Q_{2r}(x_0,t_0)} |F_{\varepsilon}|
$$

To bound the first integral in the RHS of [\(6.3\)](#page-22-0), we subdivide the domain $Q_{2r}(x_0, t_0)$ into $Q_{\varepsilon}(x,t)$ and $Q_{2r}(x_0,t_0) \setminus Q_{\varepsilon}(x,t)$. On $Q_{2r}(x_0,t_0) \setminus Q_{\varepsilon}(x,t)$, we use the bound

$$
|F_{\varepsilon}(y,s)-F_{\varepsilon}(x,s)|\leq 2||F_{\varepsilon}||_{L^{\infty}(Q_{2r}(x_0,t_0))},
$$

while for $Q_{\varepsilon}(x,t)$, we use

$$
|F_{\varepsilon}(y,s) - F_{\varepsilon}(x,s)| \leq |x-y|^{\lambda} ||F||_{C^{\lambda,0}(Q_{2r}(x_0,t_0))}.
$$

This leads to

$$
|\nabla J(x,t)| \leq C\varepsilon \ln \left[\varepsilon^{-1}r + 1\right] ||F_{\varepsilon}||_{L^{\infty}(Q_{2r}(x_0,t_0))} + C\varepsilon^{1+\lambda} ||F_{\varepsilon}||_{C^{\lambda,0}(Q_{2r}(x_0,t_0))}
$$

\n
$$
\leq C\varepsilon \ln \left[\varepsilon^{-1}r + 1\right] |||\nabla^2 u_0| + \varepsilon |\partial_t \nabla u_0| + \varepsilon |\nabla^3 u_0| ||_{L^{\infty}(Q_{2r}(x_0,t_0))}
$$

\n
$$
+ C\varepsilon^{1+\lambda} |||\nabla^2 u_0| + \varepsilon |\partial_t \nabla u_0| + \varepsilon |\nabla^3 u_0|| ||_{C^{\lambda,0}(Q_{2r}(x_0,t_0))}.
$$

Thus, in view of the estimate for $\nabla I(x,t)$, we have proved that $\|\nabla w_{\varepsilon}\|_{L^{\infty}(Q_r(x_0,t_0))}$ is bounded by the RHS of [\(6.1\)](#page-21-0). Since

$$
\|\nabla w_{\varepsilon}-\{\nabla u_{\varepsilon}-\nabla u_0-(\nabla\chi)^{\varepsilon}\nabla u_0\}\|_{L^{\infty}(Q_r(x_0,t_0))}\leq C\varepsilon\||\nabla^2 u_0|+\varepsilon|\nabla^3 u_0|\|_{L^{\infty}(Q_r(x_0,t_0))},
$$

the estimate [\(6.1\)](#page-21-0) follows.

To prove Theorem [1.2,](#page-1-3) we fix $x_0, y_0 \in \mathbb{R}^d$ and $s_0 < t_0$. We may assume that $\varepsilon < (t_0 - s_0)/8$. For otherwise the estimate [\(1.9\)](#page-2-0) follows directly from [\(2.22\)](#page-8-3). We apply Theorem [6.1](#page-21-1) to the functions

$$
u_{\varepsilon}(x,t) = \Gamma_{\varepsilon}(x,t;y_0,s_0)
$$
 and $u_0(x,t) = \Gamma_0(x,t;y_0,s_0)$

in $Q_{2r}(x_0, t_0)$, where $r = (t_0 - s_0)/8$. Note that $(\partial_t + \mathcal{L}_\varepsilon)u_\varepsilon = (\partial_t + \mathcal{L}_0)u_0 = 0$ in $Q_{4r}(x_0, t_0)$. To bound the first term in the RHS of [\(6.1\)](#page-21-0), we use the estimate [\(1.7\)](#page-1-4) in Theorem [1.1.](#page-1-5) All other terms in the RHS of [\(6.1\)](#page-21-0) may be handled easily by using the estimates[\(4.1\)](#page-13-3) for $\Gamma_0(x, t; y, s)$. We leave the details to the reader.

7 Proof of Theorem [1.3](#page-2-3)

To prove Theorem [1.3,](#page-2-3) we fix $x_0, y_0 \in \mathbb{R}^d$ and $s_0 < t_0$. As before, we may assume that $\varepsilon < (t_0 - s_0)/8$. For otherwise the estimate [\(1.14\)](#page-2-4) follows directly from [\(2.23\)](#page-8-2).

Let $r = (t_0 - s_0)/8$. Fix $1 \leq j \leq d$ and $1 \leq \beta \leq m$. We apply Theorem [6.1](#page-21-1) to the functions $u_{\varepsilon} = (u_{\varepsilon}^{\alpha})$ and $u_0 = (u_0^{\alpha})$ in $Q_{2r}(x_0, t_0)$, where

$$
u_{\varepsilon}^{\alpha}(x,t) = \frac{\partial}{\partial y_{j}} \Big\{ \Gamma_{\varepsilon}^{\alpha\beta} \Big\} (x,t;y_{0},s_{0})
$$

$$
u_{0}^{\alpha}(x,t) = \frac{\partial}{\partial y_{\ell}} \Big\{ \Gamma_{0}^{\alpha\sigma} \Big\} (x,t;y_{0},s_{0}) \cdot \Big\{ \delta^{\beta\sigma} \delta_{j\ell} + \frac{\partial}{\partial y_{j}} (\widetilde{\chi}_{\ell}^{\beta\sigma}) (y_{0}/\varepsilon, -s_{0}/\varepsilon^{2}) \Big\} ,
$$

where $\tilde{\chi}$ denotes the correctors for $\partial_t + \tilde{\mathcal{L}}_{\varepsilon}$. Observe that $(\partial_t + \mathcal{L}_{\varepsilon})u_{\varepsilon} = (\partial_t + \mathcal{L}_0)u_0 = 0$ in $Q_{4r}(x_0, t_0)$. To bound the first term in the RHS of [\(6.1\)](#page-21-0), we use the estimate [\(1.13\)](#page-2-5). As in the proof of Theorem [1.1,](#page-1-5) all other terms in the RHS of [\(6.1\)](#page-21-0) may be handled readily by using estimate [\(4.1\)](#page-13-3) for $\Gamma_0(x, t; y, s)$.

 \Box

References

- [1] D. G. Aronson, Bounds for the fundamental solution of a parabolic equation, Bull. Amer. Math. Soc. **73** (1967), 890–896.
- [2] P. Auscher, Regularity theorems and heat kernel for elliptic operators, J. London Math. Soc. (2) 54 (1996), no. 2, 284–296.
- [3] M. Avellaneda and F. Lin, L^p bounds on singular integrals in homogenization, Comm. Pure Appl. Math. 44 (1991), no. 8-9, 897–910.
- [4] A. Bensoussan, J.-L. Lions, and G. Papanicolaou, Asymptotic analysis for periodic structures, AMS Chelsea Publishing, Providence, RI, 2011, Corrected reprint of the 1978 original [MR0503330].
- [5] S. Byun, *Optimal* $W^{1,p}$ regularity theory for parabolic equations in divergence form, J. Evol. Equ. 7 (2007), no. 3, 415–428.
- [6] S. Cho, H. Dong, and S. Kim, On the Green's matrices of strongly parabolic systems of second order, Indiana Univ. Math. J. 57 (2008), no. 4, 1633–1677.
- [7] E. B. Davies, Explicit constants for Gaussian upper bounds on heat kernels, Amer. J. Math. 109 (1987), no. 2, 319–333.
- [8] \Box , Heat kernel bounds for second order elliptic operators on Riemannian manifolds, Amer. J. Math. 109 (1987), no. 3, 545–569.
- [9] E. B. Fabes and D. W. Stroock, A new proof of Moser's parabolic Harnack inequality using the old ideas of Nash, Arch. Rational Mech. Anal. 96 (1986), no. 4, 327–338.
- [10] J. Geng and Z. Shen, Uniform regularity estimates in parabolic homogenization, Indiana Univ. Math. J. 64 (2015), 697–733.
- [11] , Convergence rates in parabolic homogenization with time-dependent periodic coefficients, J. Funct. Anal. 272 (2017), no. 5, 2092–2113.
- [12] S. Hofmann and S. Kim, Gaussian estimates for fundamental solutions to certain parabolic systems, Publ. Mat. 48 (2004), no. 2, 481–496.
- [13] V. V. Jikov, S. M. Kozlov, and O. A. Oleinik, Homogenization of differential operators and integral functionals, Springer-Verlag, Berlin, 1994, Translated from the Russian by G. A. Yosifian [G. A. Iosifyan].
- [14] C. Kenig, F. Lin, and Z. Shen, Periodic homogenization of Green and Neumann functions, Comm. Pure Appl. Math. 67 (2014), no. 8, 1219–1262.
- [15] C. Kenig and Z. Shen, Layer potential methods for elliptic homogenization problems, Comm. Pure Appl. Math. 64 (2011), no. 1, 1–44.
- [16] N. V. Krylov, Parabolic and elliptic equations with VMO coefficients, Comm. Partial Differential Equations 32 (2007), no. 1-3, 453–475.
- [17] J. Nash, Continuity of solutions of parabolic and elliptic equations, Amer. J. Math. 80 (1958), 931–954.

Jun Geng, School of Mathematics and Statistics, Lanzhou University, Lanzhou, P.R. China. E-mail:gengjun@lzu.edu.cn

Zhongwei Shen, Department of Mathematics, University of Kentucky, Lexington, Kentucky 40506, USA. E-mail: zshen2@uky.edu