

Weak-strong uniqueness for the general Ericksen–Leslie system in three dimensions*

Etienne Emmrich[†] Robert Lasarzik[‡]

Version October 25, 2021

Abstract

We study the Ericksen–Leslie system equipped with a quadratic free energy functional. The norm restriction of the director is incorporated by a standard relaxation technique using a double-well potential. We use the relative energy concept, often applied in the context of compressible Euler- or related systems of fluid dynamics, to prove weak-strong uniqueness of solutions. A main novelty is that the relative energy inequality is proved for a system with a nonconvex energy.

Keywords: Liquid crystal, Ericksen–Leslie equation, Existence, Weak solution, Weak-strong uniqueness,
MSC (2010): 35Q35, 35K52, 35D30, 76A15

Contents

1	Introduction	1
2	Model and main result	4
3	Properties of the strong solution	6
4	Integration-by-parts formulae	8
5	Proof of the main result	9

1 Introduction

This paper is devoted to the weak-strong uniqueness of weak solutions to the three-dimensional Ericksen–Leslie model describing liquid crystal flow. The Ericksen–Leslie model (proposed by Ericksen [8] and Leslie [16]) is a very successful model for nematic liquid crystals and agrees with experiments (see [1, Sec. 11.1, p. 463]). The particular model strongly depends on the choice of the free energy.

Recently, global existence of weak solutions for a very general class of free energies was shown in [7]. In this article, we prove weak-strong uniqueness of these solutions for a special, physically relevant free energy. The weak-strong uniqueness property says that the weak solution coincides with a weak solution admitting additional regularity as long as the latter exists. We use the concept of relative energy (see Feiereisl, Jin and Novotný [11]), which can

*This work was funded by CRC 901 *Control of self-organizing nonlinear systems: Theoretical methods and concepts of application* (Project A8).

[†]Technische Universität Berlin, Institut für Mathematik, Straße des 17. Juni 136, 10623 Berlin, Germany
emmrich@math.tu-berlin.de

[‡]Technische Universität Berlin, Institut für Mathematik, Straße des 17. Juni 136, 10623 Berlin, Germany
lasarzik@math.tu-berlin.de

also be used to consider other problems such as the stability of an equilibrium (see Feireisl [9]), singular limits for vanishing coefficients (see Breit, Feireisl and Hofmanova [2] or Feireisl [10]) or to derive a posteriori estimates for modeling errors (see Fischer [13]). In the paper at hand, we generalize the relative energy approach to a model with a nonconvex energy.

1.1 Review of known results

A simplified Ericksen–Leslie model

$$\begin{aligned} \partial_t \mathbf{d} + (\mathbf{v} \cdot \nabla) \mathbf{d} &= \Delta \mathbf{d} + \frac{1}{\varepsilon^2} (|\mathbf{d}|^2 - 1) \mathbf{d}, \\ \partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} + \nabla p - \Delta \mathbf{d} &= -\nabla \cdot (\nabla \mathbf{d}^T \nabla \mathbf{d}), \\ \nabla \cdot \mathbf{v} &= 0, \end{aligned} \tag{1.1}$$

was first considered in Lin and Liu [17], where global existence of weak solutions as well as local existence of strong solutions was shown. Later, Lin and Liu [18] showed the same result for a generalized system. Existence of weak solutions to the model considered in the article at hand (see (2.1) below) equipped with the Dirichlet energy with double-well potential

$$F(\mathbf{d}, \nabla \mathbf{d}) = \frac{k}{2} |\nabla \mathbf{d}|^2 + \frac{1}{4\varepsilon} (|\mathbf{d}|^2 - 1)^2.$$

was first proved in Cavaterra, Rocca and Wu [3]. In [7] the existence of weak solutions to the model considered in the article at hand was proved for a more general class of free-energies.

The concept of weak-strong uniqueness was first considered by Prodi in 1959 (see [21]) and Serrin in 1962 (see [23]). Both studied the Navier–Stokes equation and showed weak-strong uniqueness for a class of weak solutions fulfilling additional regularity requirements.

There is several work on the weak-strong uniqueness property for different simplifications of the Ericksen–Leslie model. Zhao and Liu [26] established weak-strong uniqueness for the simplified system (1.1) with different assumptions on the strong solution. Dai [4, 5] established weak-strong uniqueness for a simplified incompressible model and a more general incompressible Ericksen–Leslie model with additional assumptions on the weak solution, which cannot be shown to hold in general. Yang et al. [24] showed the weak-strong uniqueness for the incompressible simplified Ericksen–Leslie system with no nonlinear penalization using ideas of Feireisl et al. [12] based upon relative entropy and suitable weak solutions. In the article at hand, we use similar ideas. However, we are able to incorporate the nonlinear part of the free energy in the relative entropy, which we call relative energy, and to show the weak-strong uniqueness without further assumptions on the weak solution. This is done by adapting the relative energy to the nonconvex energy of the system. A similar weak-strong uniqueness result for measure-valued solutions to the Ericksen–Leslie system equipped with the nonconvex Oseen–Frank energy (see [14] for the existence of such solutions) was recently proved by the second author [15].

1.2 Notation

Vectors of \mathbb{R}^3 are denoted by bold small Latin letters. Matrices of $\mathbb{R}^{3 \times 3}$ are denoted by bold capital Latin letters. Moreover, numbers are denoted by small Latin or Greek letters, and capital Latin letters are reserved for potentials.

The Euclidean inner product in \mathbb{R}^3 is denoted by a dot, $\mathbf{a} \cdot \mathbf{b} := \mathbf{a}^T \mathbf{b} = \sum_{i=1}^3 \mathbf{a}_i \mathbf{b}_i$ for $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$. The Frobenius inner product in the space $\mathbb{R}^{3 \times 3}$ of matrices is denoted by a double dot, $\mathbf{A} : \mathbf{B} := \text{tr}(\mathbf{A}^T \mathbf{B}) = \sum_{i,j=1}^3 \mathbf{A}_{ij} \mathbf{B}_{ij}$ for $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{3 \times 3}$. We also employ the corresponding Euclidean norm with $|\mathbf{a}|^2 = \mathbf{a} \cdot \mathbf{a}$ for $\mathbf{a} \in \mathbb{R}^3$ and Frobenius norm with $|\mathbf{A}|^2 = \mathbf{A} : \mathbf{A}$ for $\mathbf{A} \in \mathbb{R}^{3 \times 3}$. The product of a fourth order with a second order tensor is defined by

$$\mathbf{\Gamma} : \mathbf{A} := \left[\sum_{k,l=1}^3 \mathbf{\Gamma}_{ijkl} \mathbf{A}_{kl} \right]_{i,j=1}^3, \quad \mathbf{\Gamma} \in \mathbb{R}^{3 \times 3 \times 3 \times 3}, \mathbf{A} \in \mathbb{R}^{3 \times 3}.$$

The standard matrix and matrix-vector multiplication, however, is written without an extra sign for brevity,

$$\mathbf{AB} = \left[\sum_{j=1}^3 \mathbf{A}_{ij} \mathbf{B}_{jk} \right]_{i,k=1}^3, \quad \mathbf{Aa} = \left[\sum_{j=1}^3 \mathbf{A}_{ij} \mathbf{a}_j \right]_{i=1}^3, \quad \mathbf{A} \in \mathbb{R}^{3 \times 3}, \mathbf{B} \in \mathbb{R}^{3 \times 3}, \mathbf{a} \in \mathbb{R}^3.$$

The outer product is denoted by $\mathbf{a} \otimes \mathbf{b} = \mathbf{ab}^T = [\mathbf{a}_i \mathbf{b}_j]_{i,j=1}^3$ for $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$. Note that $\text{tr}(\mathbf{a} \otimes \mathbf{b}) = \mathbf{a} \cdot \mathbf{b}$. The symmetric and skew-symmetric part of a matrix are denoted by $\mathbf{A}_{\text{sym}} := \frac{1}{2}(\mathbf{A} + \mathbf{A}^T)$ and $\mathbf{A}_{\text{skw}} := \frac{1}{2}(\mathbf{A} - \mathbf{A}^T)$ for $\mathbf{A} \in \mathbb{R}^{3 \times 3}$, respectively.

We use the Nabla symbol ∇ for real-valued functions $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ as well as vector-valued functions $\mathbf{f} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ denoting

$$\nabla f := \left[\frac{\partial f}{\partial \mathbf{x}_i} \right]_{i=1}^3, \quad \nabla \mathbf{f} := \left[\frac{\partial \mathbf{f}_i}{\partial \mathbf{x}_j} \right]_{i,j=1}^3.$$

For brevity, we write $\nabla \mathbf{f}^T$ instead of $(\nabla \mathbf{f})^T$. The divergence of a vector-valued function $\mathbf{f} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ and a matrix-valued function $\mathbf{A} : \mathbb{R}^3 \rightarrow \mathbb{R}^{3 \times 3}$ is defined by

$$\nabla \cdot \mathbf{f} := \sum_{i=1}^3 \frac{\partial \mathbf{f}_i}{\partial \mathbf{x}_i} = \text{tr}(\nabla \mathbf{f}), \quad \nabla \cdot \mathbf{A} := \left[\sum_{j=1}^3 \frac{\partial \mathbf{A}_{ij}}{\partial \mathbf{x}_j} \right]_{i=1}^3.$$

Note that $(\mathbf{v} \cdot \nabla) \mathbf{f} = (\nabla \mathbf{f}) \mathbf{v} = \nabla \mathbf{f} \mathbf{v}$ for vector-valued functions $\mathbf{v}, \mathbf{f} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$.

Throughout this paper, let $\Omega \subset \mathbb{R}^3$ be a bounded domain of class $\mathcal{C}^{3,1}$. We rely on the usual notation for spaces of continuous functions, Lebesgue and Sobolev spaces. Spaces of vector-valued functions are emphasized by bold letters, for example $\mathbf{L}^p(\Omega) := L^p(\Omega; \mathbb{R}^3)$, $\mathbf{W}^{k,p}(\Omega) := W^{k,p}(\Omega; \mathbb{R}^3)$. If it is clear from the context, we also use this bold notation for spaces of matrix-valued functions. For brevity, we often omit calling the domain Ω . The standard inner product in $L^2(\Omega; \mathbb{R}^3)$ is denoted by (\cdot, \cdot) and in $L^2(\Omega; \mathbb{R}^{3 \times 3})$ by $(\cdot; \cdot)$.

The space of smooth solenoidal functions with compact support is denoted by $\mathcal{C}_{c,\sigma}^\infty(\Omega; \mathbb{R}^3)$. By $\mathbf{L}_\sigma^p(\Omega)$, $\mathbf{H}_{0,\sigma}^1(\Omega)$, and $\mathbf{W}_{0,\sigma}^{1,p}(\Omega)$, we denote the closure of $\mathcal{C}_{c,\sigma}^\infty(\Omega; \mathbb{R}^3)$ with respect to the norm of $\mathbf{L}^p(\Omega)$, $\mathbf{H}^1(\Omega)$, and $\mathbf{W}^{1,p}(\Omega)$, respectively ($1 \leq p < \infty$).

The dual space of a Banach space V is always denoted by V^* and equipped with the standard norm; the duality pairing is denoted by $\langle \cdot, \cdot \rangle$. The duality pairing between $\mathbf{L}^p(\Omega)$ and $\mathbf{L}^q(\Omega)$ (with $1/p + 1/q = 1$), however, is denoted by (\cdot, \cdot) or $(\cdot; \cdot)$.

The Banach space of linear bounded operators mapping a Banach space V into itself is denoted by $\mathcal{L}(V)$ and equipped with the usual norm. For a given Banach space V , Bochner–Lebesgue spaces are denoted, as usual, by $L^p(0, T; V)$. Moreover, $W^{1,p}(0, T; V)$ denotes the Banach space of abstract functions in $L^p(0, T; V)$ whose weak time derivative exists and is again in $L^p(0, T; V)$ (see also Diestel and Uhl [6, Section II.2] or Roubířek [22, Section 1.5] for more details). We often omit the time interval $(0, T)$ and the domain Ω and just write, e.g., $L^p(\mathbf{W}^{k,p})$ for brevity. By $\mathcal{C}_w([0, T]; V)$, we denote the spaces of abstract functions mapping $[0, T]$ into V that are continuous with respect to the weak topology in V .

By $\mathbf{\Lambda}$, we denote a constant tensor of order 4 that is symmetric, i.e., $\mathbf{\Lambda}_{ijkl} = \mathbf{\Lambda}_{klij}$, $ijkl \in \{1, 2, 3\}$, and obeys the strong ellipticity condition (see Mc Lean [19]), i.e., there exists $\eta > 0$ such that

$$(\mathbf{a} \otimes \mathbf{b}) : \mathbf{\Lambda} : (\mathbf{a} \otimes \mathbf{b}) \geq \eta |\mathbf{a}|^2 |\mathbf{b}|^2 \quad \text{for all } \mathbf{a}, \mathbf{b} \in \mathbb{R}^3. \quad (1.2)$$

We introduce the norm $\|\cdot\|_{\mathbf{\Lambda}} := \|\cdot\|_{L^1} : \mathbf{\Lambda} : \cdot\|_{L^1}^{1/2}$. The norm $\|\nabla \cdot\|_{\mathbf{\Lambda}}$ is equivalent to the \mathbf{H}^1 -norm on \mathbf{H}_0^1 . We use the abbreviation $\Delta_{\mathbf{\Lambda}} \mathbf{d}$ for the operator $\nabla \cdot \mathbf{\Lambda} : \nabla \mathbf{d}$ for $\mathbf{d} \in \mathbf{H}^2$.

Finally, by $c > 0$, we denote a generic positive constant and by C_δ a constant depending on a given parameter $\delta > 0$.

2 Model and main result

We consider the general Ericksen–Leslie system, which was investigated in [7]. In comparison to the model in [7], we consider a particular free energy function and reformulate the stress tensor. The system is given by

$$\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} + \nabla p - \nabla \mathbf{d}^T \mathbf{q} - \nabla \cdot \mathbf{T}^L = \mathbf{g}, \quad (2.1a)$$

$$\partial_t \mathbf{d} + (\mathbf{v} \cdot \nabla) \mathbf{d} - (\nabla \mathbf{v})_{\text{skw}} \mathbf{d} + \lambda (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} + \gamma \mathbf{q} = 0, \quad (2.1b)$$

$$\nabla \cdot \mathbf{v} = 0. \quad (2.1c)$$

The vectorfields $\mathbf{v} : \overline{\Omega} \times [0, T] \rightarrow \mathbb{R}^3$ and $\mathbf{d} : \overline{\Omega} \times [0, T] \rightarrow \mathbb{R}^3$ represent the velocity field and the director field, respectively. The pressure is denoted by $p : \overline{\Omega} \times [0, T] \rightarrow \mathbb{R}$. In the article at hand, we do not address the problem of existence or uniqueness of the pressure. For the free energy potential, we choose the function

$$F(\mathbf{d}, \nabla \mathbf{d}) := \frac{1}{2} \nabla \mathbf{d} : \mathbf{\Lambda} : \nabla \mathbf{d} + \frac{1}{4\varepsilon} (|\mathbf{d}|^2 - 1)^2. \quad (2.2)$$

Here $\mathbf{\Lambda}$ is a constant symmetric fourth order tensor fulfilling the strong ellipticity condition (1.2) (see Mc Lean [19] and Section 1.2). Moreover, $\varepsilon > 0$ denotes the fixed parameter for the relaxation of the requirement $|\mathbf{d}| = 1$. We do not address the question of the limit $\varepsilon \rightarrow 0$. For such a singular limit analysis for $\varepsilon \rightarrow 0$ in the context of the Ericksen–Leslie model, we refer to [15].

The free energy is the functional induced by the free energy potential,

$$\mathcal{F}(\mathbf{d}) := \int_{\Omega} F(\mathbf{d}, \nabla \mathbf{d}) \, dx = \frac{1}{2} \|\nabla \mathbf{d}\|_{\mathbf{\Lambda}}^2 + \frac{1}{4\varepsilon} \|\|\mathbf{d}\|^2 - 1\|_{L^2}^2. \quad (2.3)$$

The vector \mathbf{q} is the variational derivative of the free energy,

$$\mathbf{q} := \frac{\delta \mathcal{F}}{\delta \mathbf{d}} = -\Delta_{\mathbf{\Lambda}} \mathbf{d} + \frac{1}{\varepsilon} (|\mathbf{d}|^2 - 1) \mathbf{d}. \quad (2.4)$$

For the definition of the operator $\Delta_{\mathbf{\Lambda}}$, see Section 1.2. In comparison to the system studied in [7], the divergence of the Ericksen stress given by $\nabla \cdot \mathbf{T}^E = \nabla \cdot (\nabla \mathbf{d}^T (\partial F / \partial \nabla \mathbf{d}))$ is replaced by $-\nabla \mathbf{d}^T \mathbf{q}$. This reformulation is valid due to the integration-by-parts formula

$$(\nabla \cdot \mathbf{T}^E, \boldsymbol{\varphi}) = -(\nabla \mathbf{d}^T \mathbf{q}, \boldsymbol{\varphi}) + (\nabla F, \boldsymbol{\varphi})$$

derived in [7, Section 3.3] that holds for every test function $\boldsymbol{\varphi} \in \mathcal{C}_c^\infty(\Omega; \mathbb{R}^3)$. Hence, via a reformulation, the term F can be incorporated in the pressure and one ends up with the formulation (2.1). The Leslie stress tensor is given by

$$\begin{aligned} \mathbf{T}^L := & \mu_1 (\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d}) \mathbf{d} \otimes \mathbf{d} + \mu_4 (\nabla \mathbf{v})_{\text{sym}} - \gamma (\mu_2 + \mu_3) (\mathbf{d} \otimes \mathbf{q})_{\text{sym}} + (\mathbf{d} \otimes \mathbf{q})_{\text{skw}} \\ & + ((\mu_5 + \mu_6) - \lambda (\mu_2 + \mu_3)) (\mathbf{d} \otimes (\nabla \mathbf{v})_{\text{sym}} \mathbf{d})_{\text{sym}}, \end{aligned} \quad (2.5)$$

Note that in view of (2.1b), the formulation (2.5) is equivalent to the formulation of the Leslie stress in [7]. In order to assure the dissipative character of the system, we assume that the parameters λ , γ , μ_1 , μ_2 , μ_3 , μ_4 , μ_5 , and μ_6 satisfy

$$\begin{aligned} \mu_1 > 0, \quad \mu_4 > 0, \quad \gamma > 0, \quad (\mu_5 + \mu_6) - \lambda (\mu_2 + \mu_3) > 0 \\ 4\gamma ((\mu_5 + \mu_6) - \lambda (\mu_2 + \mu_3)) > (\gamma (\mu_2 + \mu_3) - \lambda)^2. \end{aligned} \quad (2.6)$$

Finally, we assume that $\mathbf{g} \in L^2(0, T; (\mathbf{H}_{0,\sigma}^1)^*)$. We equip the system with initial conditions and Dirichlet boundary conditions such that

$$\mathbf{v}(\mathbf{x}, 0) = \mathbf{v}_0(\mathbf{x}) \quad \text{for } \mathbf{x} \in \Omega, \quad \mathbf{v}(\mathbf{x}, t) = \mathbf{0} \quad \text{for } (t, \mathbf{x}) \in [0, T] \times \partial\Omega, \quad (2.7a)$$

$$\mathbf{d}(\mathbf{x}, 0) = \mathbf{d}_0(\mathbf{x}) \quad \text{for } \mathbf{x} \in \Omega, \quad \mathbf{d}(\mathbf{x}, t) = \mathbf{d}_1 \quad \text{for } (t, \mathbf{x}) \in [0, T] \times \partial\Omega. \quad (2.7b)$$

We always assume that $\mathbf{d}_0 = \mathbf{d}_1$ on $\partial\Omega$, which is a compatibility condition providing regularity. For the initial and boundary values, we assume the regularity

$$\mathbf{v}_0 \in \mathbf{L}_\sigma^2, \quad \mathbf{d}_0 \in \mathbf{H}^1, \quad \text{and} \quad \mathbf{d}_1 \in \mathbf{H}^{3/2}(\partial\Omega). \quad (2.8)$$

Definition 2.1. The pair (\mathbf{v}, \mathbf{d}) is said to be a weak solution to system (2.1)–(2.8) if

$$\begin{aligned} \mathbf{v} &\in L^\infty(0, T; \mathbf{L}_\sigma^2) \cap L^2(0, T; \mathbf{H}_{0, \sigma}^1) \cap W^{1,2}(0, T; (\mathbf{W}_{0, \sigma}^{1,6})^*), \\ \mathbf{d} &\in L^\infty(0, T; \mathbf{H}^1) \cap L^2(0, T; \mathbf{H}^2) \cap W^{1,2}(0, T; \mathbf{L}^{3/2}), \end{aligned} \quad (2.9)$$

and

$$\int_0^T \langle \partial_t \mathbf{v}, \boldsymbol{\varphi} \rangle ds + \int_0^T ((\mathbf{v} \cdot \nabla) \mathbf{v}, \boldsymbol{\varphi}) ds - \int_0^T \langle \nabla \mathbf{d}^T \mathbf{q}, \boldsymbol{\varphi} \rangle dt + \int_0^T (\mathbf{T}^L : \nabla \boldsymbol{\varphi}) dt - \int_0^T \langle \mathbf{g}, \boldsymbol{\varphi} \rangle dt = 0, \quad (2.10a)$$

$$\int_0^T (\partial_t \mathbf{d}, \boldsymbol{\psi}(t)) dt + \int_0^T (((\mathbf{v} \cdot \nabla) \mathbf{d}, \boldsymbol{\psi}) - ((\nabla \mathbf{v})_{\text{skw}} \mathbf{d}, \boldsymbol{\psi}) + \lambda ((\nabla \mathbf{v})_{\text{sym}} \mathbf{d}, \boldsymbol{\psi}) + \gamma \langle \mathbf{q}, \boldsymbol{\psi} \rangle) dt = 0 \quad (2.10b)$$

for all test functions $\boldsymbol{\varphi} \in L^2(0, T; \mathbf{W}_{0, \sigma}^{1,6})$ and $\boldsymbol{\psi} \in L^2(0, T; \mathbf{L}^3)$.

The global existence of weak solutions was proved under the given assumptions (2.1)–(2.8) in [7, Theorem 3.1] for a domain of class \mathcal{C}^2 .

Definition 2.2. A weak solution (\mathbf{v}, \mathbf{d}) (see Definition 2.1) is said to be a suitable weak solution to system (2.1) if it is a weak solution and additionally satisfies the energy inequality

$$\begin{aligned} &\frac{1}{2} \|\mathbf{v}(t)\|_{\mathbf{L}^2}^2 + \mathcal{F}(\mathbf{d}(t)) + \int_0^t (\mu_1 \|\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d}\|_{\mathbf{L}^2}^2 + \mu_4 \|(\nabla \mathbf{v})_{\text{sym}}\|_{\mathbf{L}^2}^2) ds \\ &\quad + \int_0^t ((\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)) \|(\nabla \mathbf{v})_{\text{sym}} \mathbf{d}\|_{\mathbf{L}^2}^2 + \gamma \|\mathbf{q}\|_{\mathbf{L}^2}^2) ds \\ &\leq \frac{1}{2} \|\mathbf{v}_0\|_{\mathbf{L}^2}^2 + \mathcal{F}(\mathbf{d}_0) + \int_0^t (\langle \mathbf{g}, \mathbf{v} \rangle + (\gamma(\mu_2 + \mu_3) - \lambda) \langle \mathbf{q}, (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} \rangle) ds \end{aligned} \quad (2.11)$$

for almost all $t \in (0, T)$.

Definition 2.3. A weak solution $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$ (see Definition 2.1) is said to be a strong solution to (2.1) if it admits the additional regularity

$$\tilde{\mathbf{v}} \in L^2(0, T; \mathbf{W}^{1,6}), \quad \tilde{\mathbf{d}} \in L^\infty(0, T; \mathbf{L}^{12}) \cap L^2(0, T; \mathbf{W}^{2,3}), \quad \tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}} \in L^2(0, T; \mathbf{L}^6). \quad (2.12)$$

Remark 2.4. For $\mu_1 = 0$ it would be sufficient to assume the regularity $\tilde{\mathbf{v}} \in L^2(0, T; \mathbf{W}^{1,3} \cap \mathbf{L}^\infty)$ and $(\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}} \in L^2(0, T; \mathbf{L}^3)$ instead of $\tilde{\mathbf{v}} \in L^2(0, T; \mathbf{W}^{1,6})$ and $\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}} \in L^2(0, T; \mathbf{L}^6)$.

We can now state the main theorem of this paper.

Theorem 2.5. Let $\Omega \subset \mathbb{R}^3$ be a domain of class \mathcal{C}^2 . Let (\mathbf{v}, \mathbf{d}) be a suitable weak solution (see Definition 2.2) to the Ericksen–Leslie system (2.1)–(2.8) and $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$ a strong solution (see Definition 2.3) to the same initial and boundary conditions (2.7)–(2.8).

Then

$$\mathbf{v} \equiv \tilde{\mathbf{v}}, \quad \mathbf{d} \equiv \tilde{\mathbf{d}}.$$

Remark 2.6. Theorem 2.5 is a direct consequence of Lemma 5.1. In Lemma 5.1, even the continuous dependence on the initial values is shown as long as a strong solution exists.

Before we present the proof of the main result, we give an important remark on the existence of suitable weak solutions.

Remark 2.7 (Existence of suitable weak solutions). In our recent work [7], we proved global existence of weak solutions to the system (2.1) in the sense of Definition 2.1. This is done by establishing a Galerkin approximation leading to an approximate system whose solutions $(\mathbf{v}_n, \mathbf{d}_n)$ obey the energy equality as in (3.1). This allows us to show

a priori estimates for the sequence of solutions to the approximate system and extract weakly- and weakly*-converging subsequences. In the end, it is possible to identify the limit of these subsequences with the solution (\mathbf{v}, \mathbf{d}) to (2.10). It turns out that the energy inequality (2.11) cannot be shown to hold for the limit. The a priori estimates for the approximate system (see [7]) imply the following weak convergences

$$\begin{aligned} \mathbf{v}_n &\overset{*}{\rightharpoonup} \mathbf{v} && \text{in } L^\infty(0, T; \mathbf{L}_\sigma^2) \cap L^2(0, T; \mathbf{H}_{0, \sigma}^1) \cap W^{1,2}(0, T; (\mathbf{W}_{0, \sigma}^{1,6})^*), \\ \mathbf{q}_n &\rightharpoonup \mathbf{q} && \text{in } L^2(0, T; \mathbf{L}^2), \\ (\nabla \mathbf{v}_n)_{\text{sym}} \mathbf{d}_n &\rightharpoonup (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} && \text{in } L^2(0, T; \mathbf{L}^2), \\ \mathbf{d}_n \cdot (\nabla \mathbf{v}_n)_{\text{sym}} \mathbf{d}_n &\rightharpoonup \mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} && \text{in } L^2(0, T; \mathbf{L}^2), \\ \mathbf{d}_n &\overset{*}{\rightharpoonup} \mathbf{d} && \text{in } L^\infty(0, T; \mathbf{H}^1) \cap L^2(0, T; \mathbf{H}^2) \cap W^{1,2}(0, T; \mathbf{L}^{3/2}). \end{aligned}$$

Due to the weakly lower semi-continuity of the appearing norms, one can deduce that

$$\begin{aligned} &\liminf_{n \rightarrow \infty} \left(\frac{1}{2} \|\mathbf{v}_n(t)\|_{\mathbf{L}^2}^2 + \mathcal{F}(\mathbf{d}_n(t)) + \int_0^t (\mu_1 \|\mathbf{d}_n \cdot (\nabla \mathbf{v}_n)_{\text{sym}} \mathbf{d}_n\|_{\mathbf{L}^2}^2 + \mu_4 \|(\nabla \mathbf{v}_n)_{\text{sym}}\|_{\mathbf{L}^2}^2) \, ds \right. \\ &\quad \left. + \int_0^t ((\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)) \|(\nabla \mathbf{v}_n)_{\text{sym}} \mathbf{d}_n\|_{\mathbf{L}^2}^2 + \gamma \|\mathbf{q}_n\|_{\mathbf{L}^2}^2 - \langle \mathbf{g}, \mathbf{v}_n \rangle) \, ds \right) \\ &\geq \left(\frac{1}{2} \|\mathbf{v}(t)\|_{\mathbf{L}^2}^2 + \mathcal{F}(\mathbf{d}(t)) + \int_0^t (\mu_1 \|\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d}\|_{\mathbf{L}^2}^2 + \mu_4 \|(\nabla \mathbf{v})_{\text{sym}}\|_{\mathbf{L}^2}^2) \, ds \right. \\ &\quad \left. + \int_0^t ((\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)) \|(\nabla \mathbf{v})_{\text{sym}} \mathbf{d}\|_{\mathbf{L}^2}^2 + \gamma \|\mathbf{q}\|_{\mathbf{L}^2}^2 - \langle \mathbf{g}, \mathbf{v} \rangle) \, ds \right). \end{aligned} \tag{2.13}$$

Note that $\mathbf{v}_n \in \mathcal{C}_w([0, T]; \mathbf{L}_\sigma^2)$ and $\mathbf{d}_n \in \mathcal{C}_w([0, T]; \mathbf{H}^1)$. However, we are not able to identify the limit of the remaining term $(\mathbf{q}_n, (\nabla \mathbf{v}_n)_{\text{sym}} \mathbf{d}_n)$ since \mathbf{q}_n and $(\nabla \mathbf{v}_n)_{\text{sym}} \mathbf{d}_n$ only converge weakly. Thus, it is not clear whether a suitable solution in the sense of Definition 2.2 exists.

Nevertheless, the existence of a suitable solution (see Definition 2.2) can be shown when assuming Parodi's relation $\gamma(\mu_2 + \mu_3) = \lambda$. Then the last term in the energy inequality (2.11) vanishes, and with (2.13) the energy inequality also holds for the limit of the approximate solutions, which is the weak solution.

3 Properties of the strong solution

Lemma 3.1 (Energy equality). *A strong solution $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$ (see Definition 2.3) of the system (2.1) fulfills the energy equality*

$$\begin{aligned} &\frac{1}{2} \|\tilde{\mathbf{v}}(t)\|_{\mathbf{L}^2}^2 + \mathcal{F}(\tilde{\mathbf{d}}(t)) + \int_0^t (\mu_1 \|\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2 + \mu_4 \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^2}^2) \, ds \\ &\quad + \int_0^t ((\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)) \|(\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2 + \gamma \|\tilde{\mathbf{q}}\|_{\mathbf{L}^2}^2) \, ds \\ &= \frac{1}{2} \|\tilde{\mathbf{v}}_0\|_{\mathbf{L}^2}^2 + \mathcal{F}(\tilde{\mathbf{d}}_0) + \int_0^t (\langle \mathbf{g}, \tilde{\mathbf{v}} \rangle + (\gamma(\mu_2 + \mu_3) - \lambda) (\tilde{\mathbf{q}}, (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}})) \, ds \end{aligned} \tag{3.1}$$

for $t \in [0, T]$.

Proof. Due to the regularity assumptions (2.12) on the strong solution, we can take $(\tilde{\mathbf{v}}, \tilde{\mathbf{q}})$ as test functions in (2.10) and obtain the energy equality in the same way as in [7, Proposition 4.1]. \square

Lemma 3.2 (Regularity of the strong solution). *A strong solution $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$ (see Definition 2.3) admits the regularity*

$$\partial_t \tilde{\mathbf{v}} \in L^2(0, T; (\mathbf{H}_{0, \sigma}^1)^*), \quad \partial_t \tilde{\mathbf{d}} \in L^1(0, T; \mathbf{L}^3). \tag{3.2}$$

Proof. First, we estimate the time derivative of $\tilde{\mathbf{v}}$. Let $\boldsymbol{\varphi} \in L^2(0, T; \mathbf{H}_{0,\sigma}^1)$ be a test function in (2.10a). We estimate the terms individually. Because of the continuous embedding of $\mathbf{H}_{0,\sigma}^1$ into \mathbf{L}^6 , we obtain for the convection term

$$\int_0^T |(\tilde{\mathbf{v}}(t) \cdot \nabla) \tilde{\mathbf{v}}(t), \boldsymbol{\varphi}(t)| dt \leq \|\tilde{\mathbf{v}}\|_{L^\infty(\mathbf{L}^2)} \|\nabla \tilde{\mathbf{v}}\|_{L^2(\mathbf{L}^3)} \|\boldsymbol{\varphi}\|_{L^2(\mathbf{H}_{0,\sigma}^1)}.$$

Similarly, the Ericksen stress can be estimated as

$$\int_0^T |(\nabla \tilde{\mathbf{d}}(t)^T \tilde{\mathbf{q}}(t), \boldsymbol{\varphi}(t))| dt \leq \|\nabla \tilde{\mathbf{d}}\|_{L^\infty(\mathbf{L}^2)} \|\tilde{\mathbf{q}}\|_{L^2(\mathbf{L}^3)} \|\boldsymbol{\varphi}\|_{L^2(\mathbf{H}_{0,\sigma}^1)}.$$

For the right-hand side, we have that

$$\int_0^T |\langle \mathbf{g}(t), \boldsymbol{\varphi}(t) \rangle| dt \leq \|\mathbf{g}\|_{L^2((\mathbf{H}_{0,\sigma}^1)^*)} \|\boldsymbol{\varphi}\|_{L^2(\mathbf{H}_0^1)}.$$

With the definition of the Leslie stress tensor (see (2.5)), we get

$$\begin{aligned} \int_0^T |(\mathbf{T}^L(t); \nabla \boldsymbol{\varphi}(t))| dt &\leq \left(\mu_1 \|\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{L^2(\mathbf{L}^6)} \|\tilde{\mathbf{d}}\|_{L^\infty(\mathbf{L}^6)}^2 + \mu_4 \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{L^2(\mathbf{L}^2)} \right. \\ &\quad \left. + (|\gamma(\mu_2 + \mu_3)| + 1) \|\tilde{\mathbf{d}}\|_{L^\infty(\mathbf{L}^6)} \|\tilde{\mathbf{q}}\|_{L^2(\mathbf{L}^3)} \right. \\ &\quad \left. + ((\mu_5 + \mu_6) - \lambda(\mu_2 + \mu_3)) \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{L^2(\mathbf{L}^6)} \|\tilde{\mathbf{d}}\|_{L^\infty(\mathbf{L}^6)}^2 \right) \|\boldsymbol{\varphi}\|_{L^2(\mathbf{H}_{0,\sigma}^1)} \\ &\leq c \left(\left(\|\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{L^2(\mathbf{L}^6)} + \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{L^2(\mathbf{L}^6)} + \|\tilde{\mathbf{q}}\|_{L^\infty(\mathbf{L}^3)} \right) \left(\|\tilde{\mathbf{d}}\|_{L^\infty(\mathbf{L}^6)}^2 + 1 \right) \right) \|\boldsymbol{\varphi}\|_{L^2(\mathbf{H}_{0,\sigma}^1)}. \end{aligned}$$

Due to the regularity assumptions on the strong solution (see Definition 2.3), the variational derivative of the free energy can be estimated in terms of the $L^2(0, T; \mathbf{L}^3)$ -norm by standard embeddings and the Gagliardo–Nirenberg inequality [25, Section 21.19],

$$\|\tilde{\mathbf{q}}\|_{L^2(\mathbf{L}^3)} \leq |\mathbf{A}| \|\tilde{\mathbf{d}}\|_{L^2(\mathbf{W}^{2,3})} + \frac{1}{\varepsilon} \left(\|\tilde{\mathbf{d}}\|_{L^6(\mathbf{L}^9)}^3 + \|\tilde{\mathbf{d}}\|_{L^2(\mathbf{L}^3)} \right) \leq c \left(\|\tilde{\mathbf{d}}\|_{L^2(\mathbf{W}^{2,3})} + \|\tilde{\mathbf{d}}\|_{L^2(\mathbf{W}^{2,3})}^{1/3} \|\tilde{\mathbf{d}}\|_{L^\infty(\mathbf{L}^6)}^{8/3} + 1 \right). \quad (3.3)$$

Note that ε is a constant parameter. Altogether, we see that $\partial_t \tilde{\mathbf{v}} \in L^2(0, T; (\mathbf{H}_{0,\sigma}^1)^*)$ and

$$\begin{aligned} \|\partial_t \tilde{\mathbf{v}}\|_{L^2((\mathbf{H}_{0,\sigma}^1)^*)} &\leq \|\tilde{\mathbf{v}}\|_{L^\infty(\mathbf{L}^2)} \|\nabla \tilde{\mathbf{v}}\|_{L^2(\mathbf{L}^3)} + \|\nabla \tilde{\mathbf{d}}\|_{L^\infty(\mathbf{L}^2)} \|\tilde{\mathbf{q}}\|_{L^2(\mathbf{L}^3)} + \|\mathbf{g}\|_{L^2((\mathbf{H}_{0,\sigma}^1)^*)} \\ &\quad + c \left(\left(\|\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{L^2(\mathbf{L}^6)} + \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{L^2(\mathbf{L}^6)} + \|\tilde{\mathbf{q}}\|_{L^\infty(\mathbf{L}^3)} \right) \left(\|\tilde{\mathbf{d}}\|_{L^\infty(\mathbf{L}^6)}^2 + 1 \right) \right) \end{aligned}$$

with $\tilde{\mathbf{q}}$ estimated in (3.3).

Recalling equation (2.10b) and estimate (3.3), standard embeddings show that $\partial_t \tilde{\mathbf{d}} \in L^1(0, T; \mathbf{L}^3)$ with

$$\begin{aligned} \|\partial_t \tilde{\mathbf{d}}\|_{L^1(\mathbf{L}^3)} &\leq \|\tilde{\mathbf{v}}\|_{L^2(\mathbf{L}^\infty)} \|\nabla \tilde{\mathbf{d}}\|_{L^2(\mathbf{L}^3)} + \|(\nabla \tilde{\mathbf{v}})_{\text{skw}}\|_{L^2(\mathbf{L}^3)} \|\tilde{\mathbf{d}}\|_{L^2(\mathbf{L}^\infty)} + |\lambda| \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{L^2(\mathbf{L}^3)} \|\tilde{\mathbf{d}}\|_{L^2(\mathbf{L}^\infty)} + \|\tilde{\mathbf{q}}\|_{L^1(\mathbf{L}^3)} \\ &\leq c \left(\|\tilde{\mathbf{v}}\|_{L^2(\mathbf{W}^{1,6})} \|\tilde{\mathbf{d}}\|_{L^2(\mathbf{W}^{2,3})} + \|\tilde{\mathbf{q}}\|_{L^2(\mathbf{L}^3)} \right). \end{aligned}$$

□

In the course of the proof of Theorem 2.5, we shall employ the following three integration-by-parts formulae.

4 Integration-by-parts formulae

Lemma 4.1. *For functions (\mathbf{v}, \mathbf{d}) and $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$ fulfilling (2.9) and (3.2), respectively, the integration-by-parts formulae*

$$\begin{aligned} (\mathbf{v}(t), \tilde{\mathbf{v}}(t)) - (\mathbf{v}(s), \tilde{\mathbf{v}}(s)) &= \int_s^t (\langle \mathbf{v}(\tau), \partial_t \tilde{\mathbf{v}}(\tau) \rangle + \langle \partial_t \mathbf{v}(\tau), \tilde{\mathbf{v}}(\tau) \rangle) d\tau, \\ (\nabla \mathbf{d}(t); \mathbf{\Lambda} : \nabla \tilde{\mathbf{d}}(t)) - (\nabla \mathbf{d}(s); \mathbf{\Lambda} : \nabla \tilde{\mathbf{d}}(s)) &= - \int_s^t ((\partial_t \mathbf{d}(\tau), \Delta_{\mathbf{\Lambda}} \tilde{\mathbf{d}}(\tau)) + (\Delta_{\mathbf{\Lambda}} \mathbf{d}(\tau), \partial_t \tilde{\mathbf{d}}(\tau))) d\tau, \\ (|\mathbf{d}(t)|^2, |\tilde{\mathbf{d}}(t)|^2) - (|\mathbf{d}(s)|^2, |\tilde{\mathbf{d}}(s)|^2) &= 2 \int_s^t ((\partial_t \mathbf{d}(\tau) \cdot \mathbf{d}(\tau), |\tilde{\mathbf{d}}(\tau)|^2) + (|\mathbf{d}(\tau)|^2, \partial_t \tilde{\mathbf{d}}(\tau) \cdot \tilde{\mathbf{d}}(\tau))) d\tau \end{aligned} \quad (4.1)$$

hold true for every $s, t \in [0, T]$.

Proof. We choose two approximate sequences $\mathbf{v}_n \in \mathcal{C}^1([0, T]; \mathbf{H}_{0, \sigma}^1)$ and $\tilde{\mathbf{v}}_n \in \mathcal{C}^1([0, T]; \mathbf{W}_{0, \sigma}^{1,6})$ such that

$$\begin{aligned} \mathbf{v}_n &\rightarrow \mathbf{v} \in L^2(0, T; \mathbf{H}_{0, \sigma}^1) \cap W^{1,2}(0, T; (\mathbf{W}_{0, \sigma}^{1,6})^*) \\ \tilde{\mathbf{v}}_n &\rightarrow \tilde{\mathbf{v}} \in L^2(0, T; \mathbf{W}_{0, \sigma}^{1,6}) \cap W^{1,2}(0, T; (\mathbf{H}_{0, \sigma}^1)^*), \end{aligned} \quad (4.2)$$

which is possible in view of density. For the approximate sequences, the integration-by-parts formula

$$(\mathbf{v}_n(t), \tilde{\mathbf{v}}_n(t)) - (\mathbf{v}_n(s), \tilde{\mathbf{v}}_n(s)) = \int_s^t ((\mathbf{v}_n(\tau), \partial_t \tilde{\mathbf{v}}_n(\tau)) + (\partial_t \mathbf{v}_n(\tau), \tilde{\mathbf{v}}_n(\tau))) d\tau \quad (4.3)$$

obviously holds true for all $s, t \in [0, T]$. In the following, we derive estimates for the terms on the left-hand side of (4.3). Let us define a partition of the unity via a function $\phi \in \mathcal{C}^1([0, T])$ with

$$|\phi(t)| \leq 1 \quad \text{for all } t \in [0, T], \quad \phi(0) = 0 \text{ and } \phi(T) = 1.$$

Let $\mathbf{u} \in \mathcal{C}^1([0, T]; \mathbf{H}_{0, \sigma}^1)$ and $\tilde{\mathbf{u}} \in \mathcal{C}^1([0, T]; \mathbf{W}_{0, \sigma}^{1,6})$. We have that

$$\mathbf{u}(t) = \phi(t)\mathbf{u}(t) + (1 - \phi(t))\mathbf{u}(t), \quad \tilde{\mathbf{u}}(t) = \phi(t)\tilde{\mathbf{u}}(t) + (1 - \phi(t))\tilde{\mathbf{u}}(t).$$

We abbreviate $\bar{\phi}(t) := 1 - \phi(t)$ for all $t \in [0, T]$ such that $\bar{\phi}(T) = 0$. It can easily be seen that for $t \in [0, T]$

$$\begin{aligned} \phi(t)(\mathbf{u}(t), \tilde{\mathbf{u}}(t)) &= \phi(0)(\mathbf{u}(0), \tilde{\mathbf{u}}(0)) + \int_0^t \phi'(\tau)(\mathbf{u}(\tau), \tilde{\mathbf{u}}(\tau)) d\tau + \int_0^t \phi(\tau)((\partial_t \mathbf{u}(\tau), \tilde{\mathbf{u}}(\tau)) + (\mathbf{u}(\tau), \partial_t \tilde{\mathbf{u}}(\tau))) d\tau \\ \bar{\phi}(t)(\mathbf{u}(t), \tilde{\mathbf{u}}(t)) &= \bar{\phi}(T)(\mathbf{u}(T), \tilde{\mathbf{u}}(T)) + \int_t^T \phi'(\tau)(\mathbf{u}(\tau), \tilde{\mathbf{u}}(\tau)) d\tau - \int_t^T \bar{\phi}(\tau)((\partial_t \mathbf{u}(\tau), \tilde{\mathbf{u}}(\tau)) + (\mathbf{u}(\tau), \partial_t \tilde{\mathbf{u}}(\tau))) d\tau. \end{aligned}$$

Summing up the two previous equations, we find that for all $t \in [0, T]$

$$\begin{aligned} (\mathbf{u}(t), \tilde{\mathbf{u}}(t)) &= \int_0^T \phi'(\tau)(\mathbf{u}(\tau), \tilde{\mathbf{u}}(\tau)) d\tau + \int_0^T \phi(\tau)((\partial_t \mathbf{u}(\tau), \tilde{\mathbf{u}}(\tau)) + (\mathbf{u}(\tau), \partial_t \tilde{\mathbf{u}}(\tau))) d\tau \\ &\quad - \int_t^T ((\partial_t \mathbf{u}(\tau), \tilde{\mathbf{u}}(\tau)) + (\mathbf{u}(\tau), \partial_t \tilde{\mathbf{u}}(\tau))) d\tau \\ &\leq \max_{t \in [0, T]} |\phi'(t)| \|\mathbf{u}\|_{L^2(L^2)} \|\tilde{\mathbf{u}}\|_{L^2(L^2)} + 2\|\partial_t \mathbf{u}\|_{L^2((\mathbf{W}_{0, \sigma}^{1,6})^*)} \|\tilde{\mathbf{u}}\|_{L^2(\mathbf{W}_{0, \sigma}^{1,6})} + 2\|\mathbf{u}\|_{L^2(\mathbf{H}_{0, \sigma}^1)} \|\partial_t \tilde{\mathbf{u}}\|_{L^2((\mathbf{H}_{0, \sigma}^1)^*)}. \end{aligned} \quad (4.4)$$

The above estimate is now applied to the left-hand side of (4.3). Since $\{\mathbf{v}_n\}$ and $\{\tilde{\mathbf{v}}_n\}$ are Cauchy sequences in the spaces indicated in (4.2), we see with

$$(\mathbf{v}_n(t), \tilde{\mathbf{v}}_n(t)) - (\mathbf{v}_m(t), \tilde{\mathbf{v}}_m(t)) = (\mathbf{v}_n(t) - \mathbf{v}_m(t), \tilde{\mathbf{v}}_n(t)) + (\mathbf{v}_m(t), \tilde{\mathbf{v}}_n(t) - \tilde{\mathbf{v}}_m(t)) \quad \text{for } t \in [0, T] \quad \text{and } m, n \in \mathbb{N}$$

and the estimate (4.4) that $\{(\mathbf{v}_n, \tilde{\mathbf{v}}_n)\}$ is a Cauchy sequence in $\mathcal{C}([0, T])$. The continuous functions are complete and the limit is unique such that $\{(\mathbf{v}_n, \tilde{\mathbf{v}}_n)\}$ converges in $\mathcal{C}([0, T])$ to $(\mathbf{v}, \tilde{\mathbf{v}})$. For the approximation of the terms on the right-hand side of the identity (4.3), we see that the difference of the approximation and the limit can be estimated by

$$\begin{aligned} \left| \int_s^t \langle \mathbf{v}(\tau), \partial_t \tilde{\mathbf{v}}(\tau) \rangle - \langle \mathbf{v}_n(\tau), \partial_t \tilde{\mathbf{v}}_n(\tau) \rangle \, d\tau \right| &\leq \|\mathbf{v}\|_{L^2(\mathbf{H}_{0,\sigma}^1)} \|\partial_t \tilde{\mathbf{v}} - \partial_t \tilde{\mathbf{v}}_n\|_{L^2((\mathbf{H}_{0,\sigma}^1)^*)} + \|\mathbf{v} - \mathbf{v}_n\|_{L^2((\mathbf{H}_{0,\sigma}^1)^*)} \|\partial_t \tilde{\mathbf{v}}_n\|_{L^2((\mathbf{H}_{0,\sigma}^1)^*)}, \\ \left| \int_s^t \langle \partial_t \mathbf{v}(\tau), \tilde{\mathbf{v}}(\tau) \rangle - \langle \partial_t \mathbf{v}_n(\tau), \tilde{\mathbf{v}}_n(\tau) \rangle \, d\tau \right| &\leq \|\partial_t \mathbf{v} - \partial_t \mathbf{v}_n\|_{L^2((\mathbf{W}_{0,\sigma}^{1,6})^*)} \|\tilde{\mathbf{v}}\|_{L^2(\mathbf{W}_{0,\sigma}^{1,6})} + \|\partial_t \mathbf{v}_n\|_{L^2((\mathbf{W}_{0,\sigma}^{1,6})^*)} \|\tilde{\mathbf{v}} - \tilde{\mathbf{v}}_n\|_{L^2(\mathbf{W}_{0,\sigma}^{1,6})}. \end{aligned}$$

The right-hand sides of the above estimates converge to zero for $n \rightarrow \infty$ since $\{\mathbf{v}_n\}$ and $\{\tilde{\mathbf{v}}_n\}$ converge to \mathbf{v} and $\tilde{\mathbf{v}}$ in the sense of (4.2). This proves that the integration-by-parts formula (4.3) holds for \mathbf{v} and $\tilde{\mathbf{v}}$.

For proving the second and third formula in (4.1), the weak solution \mathbf{d} and the strong solution $\tilde{\mathbf{d}}$ are approximated by sequences of smooth functions \mathbf{d}_n and $\tilde{\mathbf{d}}_n$ in $L^2(0, T; \mathbf{H}^2) \cap W^{1,2}(0, T; \mathbf{L}^{3/2})$ and $L^2(0, T; \mathbf{W}^{2,3}) \cap W^{1,2}(0, T; \mathbf{L}^2)$, respectively. The approximate sequences can be chosen such that every element, i.e., \mathbf{d}_n and $\tilde{\mathbf{d}}_n$, of this sequences fulfills the same boundary conditions as \mathbf{d} and $\tilde{\mathbf{d}}$, respectively. Since these boundary values of \mathbf{d}_n and $\tilde{\mathbf{d}}_n$ are constant in time, their time derivative vanishes on the boundary. Hence, there are no boundary terms in the integration-by-parts formula

$$\begin{aligned} (\nabla \mathbf{d}_n(t); \mathbf{\Lambda} : \nabla \tilde{\mathbf{d}}_n(t)) - (\nabla \mathbf{d}_n(s); \mathbf{\Lambda} : \nabla \tilde{\mathbf{d}}_n(s)) &= \int_s^t ((\nabla \partial_t \mathbf{d}_n(\tau); \mathbf{\Lambda} : \nabla \tilde{\mathbf{d}}_n(\tau)) + (\nabla \mathbf{d}_n(\tau); \mathbf{\Lambda} : \nabla \partial_t \tilde{\mathbf{d}}_n(\tau))) \, d\tau \\ &= - \int_s^t ((\partial_t \mathbf{d}_n(\tau), \Delta_{\mathbf{\Lambda}} \tilde{\mathbf{d}}_n(\tau)) + (\Delta_{\mathbf{\Lambda}} \mathbf{d}_n(\tau), \partial_t \tilde{\mathbf{d}}_n(\tau))) \, d\tau. \end{aligned}$$

Going to the limit in n shows the second integration-by-parts formula in (4.1). The third integration-by-parts formula in (4.1) is proved by the same approximation. \square

5 Proof of the main result

We define the relative energy for two solutions to system (2.1)

$$\mathcal{E}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) := \frac{1}{2} \|\mathbf{v} - \tilde{\mathbf{v}}\|_{\mathbf{L}^2}^2 + \frac{1}{2} \|\nabla \mathbf{d} - \nabla \tilde{\mathbf{d}}\|_{\mathbf{\Lambda}}^2 + \frac{1}{4\mathcal{E}} \|(|\mathbf{d}|^2 - 1) - (|\tilde{\mathbf{d}}|^2 - 1)\|_{\mathbf{L}^2}^2, \quad (5.1)$$

and the relative dissipation by

$$\begin{aligned} \mathcal{W}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) &:= \mu_1 \|\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - \tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2 + \mu_4 \|(\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^2}^2 \\ &\quad + (\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)) \|(\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2 + \gamma \|\mathbf{q} - \tilde{\mathbf{q}}\|_{\mathbf{L}^2}^2 \end{aligned} \quad (5.2)$$

Lemma 5.1. *Let (\mathbf{v}, \mathbf{d}) be a suitable weak solution (see Definition 2.2) to system (2.1) for given initial values $(\mathbf{v}_0, \mathbf{d}_0)$. Let $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$ be a strong solution (see Definition 2.3) to system (2.1) for given initial values $(\tilde{\mathbf{v}}_0, \tilde{\mathbf{d}}_0)$. Then for almost all $t \in [0, T]$*

$$\mathcal{E}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}})(t) \leq \mathcal{E}(\mathbf{v}_0, \mathbf{d}_0 | \tilde{\mathbf{v}}_0, \tilde{\mathbf{d}}_0) e^{\int_0^t \mathcal{H}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}})(s) \, ds}, \quad (5.3)$$

where \mathcal{H} is given by

$$\begin{aligned} \mathcal{H}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) &= \\ c \left(1 + \|\mathbf{d}\|_{L^\infty(\mathbf{L}^6)}^2 + \|\tilde{\mathbf{d}}\|_{L^\infty(\mathbf{L}^6)}^2 \right) & \left(\|\tilde{\mathbf{v}}\|_{\mathbf{W}^{1,6}}^2 + \|\tilde{\mathbf{q}}\|_{\mathbf{L}^3}^2 + \|\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 + \|\partial_t \tilde{\mathbf{d}}\|_{\mathbf{L}^3} + \| |\tilde{\mathbf{d}}|^2 - 1 \|_{\mathbf{L}^6}^2 + \|\mathbf{v}\|_{\mathbf{L}^6}^2 + \|\nabla \tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2 \right) \end{aligned} \quad (5.4)$$

and c is a possibly large constant.

Remark 5.2. The functional (5.4) only depends on the two norms $\|\mathbf{v}\|_{L^2(\mathbf{L}^6)}$ and $\|\mathbf{d}\|_{L^\infty(\mathbf{L}^6)}$ of the weak solution, which are known to be finite. Additionally, it depends on several norms of the strong solution $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$. Due to the regularity assumptions (3.2) and estimate (3.3), the functional \mathcal{H} is bounded in $L^1(0, T)$. For the relative energy and the relative dissipation, note that $\mathcal{E}(\mathbf{v}, \mathbf{d}|\tilde{\mathbf{v}}, \tilde{\mathbf{d}}) \in L^\infty(0, T)$ and $\mathcal{W}(\mathbf{v}, \mathbf{d}|\tilde{\mathbf{v}}, \tilde{\mathbf{d}}) \in L^1(0, T)$ due to (2.9) and (2.11), respectively.

Proof. Consider the relative energy

$$\begin{aligned} \mathcal{E}(\mathbf{v}, \mathbf{d}|\tilde{\mathbf{v}}, \tilde{\mathbf{d}}) &= \frac{1}{2}\|\mathbf{v}\|_{L^2}^2 + \frac{1}{2}\|\nabla \mathbf{d}\|_{\Lambda}^2 + \frac{1}{4\varepsilon}\|\mathbf{d}\|^2 - 1\|_{L^2}^2 + \frac{1}{2}\|\tilde{\mathbf{v}}\|_{L^2}^2 + \frac{1}{2}\|\nabla \tilde{\mathbf{d}}\|_{\Lambda}^2 + \frac{1}{4\varepsilon}\|\tilde{\mathbf{d}}\|^2 - 1\|_{L^2}^2 \\ &\quad - (\mathbf{v}, \tilde{\mathbf{v}}) - (\nabla \mathbf{d}; \Lambda : \nabla \tilde{\mathbf{d}}) - \frac{1}{2\varepsilon}(|\mathbf{d}|^2 - 1, |\tilde{\mathbf{d}}|^2 - 1). \end{aligned}$$

We insert the energy inequality (2.11) for the weak solution (\mathbf{v}, \mathbf{d}) and the energy equality (3.1) for the smooth solution $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$. This leads to

$$\begin{aligned} \mathcal{E}(\mathbf{v}, \mathbf{d}|\tilde{\mathbf{v}}, \tilde{\mathbf{d}}) &\leq \frac{1}{2}\|\mathbf{v}_0\|_{L^2}^2 + \frac{1}{2}\|\tilde{\mathbf{v}}_0\|_{L^2}^2 + \mathcal{F}(\mathbf{d}_0) + \mathcal{F}(\tilde{\mathbf{d}}_0) \\ &\quad - \mu_1 \int_0^t (\|\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}}\|_{L^2}^2 + \|\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{L^2}^2) \, ds - \mu_4 \int_0^t (\|(\nabla \mathbf{v})_{\text{sym}}\|_{L^2}^2 + \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{L^2}^2) \, ds \\ &\quad - (\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)) \int_0^t (\|(\nabla \mathbf{v})_{\text{sym}}\mathbf{d}\|_{L^2}^2 + \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\tilde{\mathbf{d}}\|_{L^2}^2) \, ds - \gamma \int_0^t (\|\mathbf{q}\|_{L^2}^2 + \|\tilde{\mathbf{q}}\|_{L^2}^2) \, ds \\ &\quad + \int_0^t \langle \mathbf{g}, \mathbf{v} + \tilde{\mathbf{v}} \rangle \, ds + (\gamma(\mu_2 + \mu_3) - \lambda) \int_0^t ((\mathbf{q}, (\nabla \mathbf{v})_{\text{sym}}\mathbf{d}) + (\tilde{\mathbf{q}}, (\nabla \tilde{\mathbf{v}})_{\text{sym}}\tilde{\mathbf{d}})) \, ds \\ &\quad - (\mathbf{v}, \tilde{\mathbf{v}}) - (\nabla \mathbf{d}; \Lambda : \nabla \tilde{\mathbf{d}}) - \frac{1}{2\varepsilon}(|\mathbf{d}|^2 - 1, |\tilde{\mathbf{d}}|^2 - 1). \end{aligned}$$

Adding the integral over the relative dissipation gives

$$\begin{aligned} \mathcal{E}(\mathbf{v}, \mathbf{d}|\tilde{\mathbf{v}}, \tilde{\mathbf{d}}) + \int_0^t \mathcal{W}(\mathbf{v}, \mathbf{d}|\tilde{\mathbf{v}}, \tilde{\mathbf{d}}) \, ds &\leq \frac{1}{2}\|\mathbf{v}_0\|_{L^2}^2 + \frac{1}{2}\|\tilde{\mathbf{v}}_0\|_{L^2}^2 + \mathcal{F}(\mathbf{d}_0) + \mathcal{F}(\tilde{\mathbf{d}}_0) \\ &\quad - 2\mu_1 \int_0^t (\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}}\mathbf{d}, \tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}}\tilde{\mathbf{d}}) \, ds - 2\mu_4 \int_0^t ((\nabla \mathbf{v})_{\text{sym}}; (\nabla \tilde{\mathbf{v}})_{\text{sym}}) \, ds \\ &\quad - 2(\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)) \int_0^t ((\nabla \mathbf{v})_{\text{sym}}\mathbf{d}, (\nabla \tilde{\mathbf{v}})_{\text{sym}}\tilde{\mathbf{d}}) \, ds - 2\gamma \int_0^t (\mathbf{q}, \tilde{\mathbf{q}}) \, ds \\ &\quad + \int_0^t \langle \mathbf{g}, \mathbf{v} + \tilde{\mathbf{v}} \rangle \, ds + (\gamma(\mu_2 + \mu_3) - \lambda) \int_0^t ((\mathbf{q}, (\nabla \mathbf{v})_{\text{sym}}\mathbf{d}) + (\tilde{\mathbf{q}}, (\nabla \tilde{\mathbf{v}})_{\text{sym}}\tilde{\mathbf{d}})) \, ds \\ &\quad - (\mathbf{v}, \tilde{\mathbf{v}}) - (\nabla \mathbf{d}; \Lambda : \nabla \tilde{\mathbf{d}}) - \frac{1}{2\varepsilon}(|\mathbf{d}|^2 - 1, |\tilde{\mathbf{d}}|^2 - 1). \end{aligned} \tag{5.5}$$

The last term can be written via Lemma 4.1 as

$$\begin{aligned} &-\frac{1}{2\varepsilon}(|\mathbf{d}|^2 - 1, |\tilde{\mathbf{d}}|^2 - 1)(t) + \frac{1}{2\varepsilon}(|\mathbf{d}_0|^2 - 1, |\tilde{\mathbf{d}}_0|^2 - 1) = -\frac{1}{2\varepsilon} \int_0^t ((|\mathbf{d}|^2 - 1, \partial_t |\tilde{\mathbf{d}}|^2) + (\partial_t |\mathbf{d}|^2, |\tilde{\mathbf{d}}|^2 - 1)) \, ds \\ &= -\frac{1}{2\varepsilon} \int_0^t (((|\mathbf{d}|^2 - 1) - (|\tilde{\mathbf{d}}|^2 - 1), \partial_t |\tilde{\mathbf{d}}|^2) + (\partial_t |\mathbf{d}|^2 + \partial_t |\tilde{\mathbf{d}}|^2, |\tilde{\mathbf{d}}|^2 - 1)) \, ds \\ &= -\frac{1}{2\varepsilon} \int_0^t (((|\mathbf{d}|^2 - 1) - (|\tilde{\mathbf{d}}|^2 - 1), 2\partial_t \tilde{\mathbf{d}} \cdot (\tilde{\mathbf{d}} - \mathbf{d})) + (\partial_t |\mathbf{d}|^2 - \tilde{\mathbf{d}}|^2, |\tilde{\mathbf{d}}|^2 - 1)) \, ds \\ &\quad - \frac{1}{2\varepsilon} \int_0^t (((|\mathbf{d}|^2 - 1) - (|\tilde{\mathbf{d}}|^2 - 1), 2\partial_t \tilde{\mathbf{d}} \cdot \mathbf{d}) + (2\partial_t (\mathbf{d} \cdot \tilde{\mathbf{d}}), |\tilde{\mathbf{d}}|^2 - 1)) \, ds \\ &= -\frac{1}{2\varepsilon} \int_0^t (((|\mathbf{d}|^2 - 1) - (|\tilde{\mathbf{d}}|^2 - 1), 2\partial_t \tilde{\mathbf{d}} \cdot (\tilde{\mathbf{d}} - \mathbf{d})) + (\partial_t |\mathbf{d}|^2 - \tilde{\mathbf{d}}|^2, |\tilde{\mathbf{d}}|^2 - 1)) \, ds \\ &\quad - \frac{1}{\varepsilon} \int_0^t (((|\mathbf{d}|^2 - 1)\mathbf{d}, \partial_t \tilde{\mathbf{d}}) + (\partial_t \mathbf{d}, (|\tilde{\mathbf{d}}|^2 - 1)\tilde{\mathbf{d}})) \, ds \end{aligned}$$

Recall the definition of the variational derivative of the free energy (see (2.4)),

$$-\frac{1}{\varepsilon} \int_0^t (((|\mathbf{d}|^2 - 1)\mathbf{d}, \partial_t \tilde{\mathbf{d}}) + (\partial_t \mathbf{d}, (|\tilde{\mathbf{d}}|^2 - 1)\tilde{\mathbf{d}})) \, ds + \int_0^t ((\partial_t \tilde{\mathbf{d}}, \Delta_\Lambda \mathbf{d}) + (\partial_t \mathbf{d}, \Delta_\Lambda \tilde{\mathbf{d}})) \, ds = - \int_0^t ((\mathbf{q}, \partial_t \tilde{\mathbf{d}}) + (\partial_t \mathbf{d}, \tilde{\mathbf{q}})) \, ds.$$

Due to the integration-by-parts formulae (4.1) and the two previous equations, the last line in (5.5) can be reformulated as

$$\begin{aligned} & -(\mathbf{v}, \tilde{\mathbf{v}}) - (\nabla \mathbf{d}; \mathbf{\Lambda} : \nabla \tilde{\mathbf{d}}) - \frac{1}{2\varepsilon} (|\mathbf{d}|^2 - 1, |\tilde{\mathbf{d}}|^2 - 1) = -(\mathbf{v}_0, \tilde{\mathbf{v}}_0) - (\nabla \mathbf{d}_0; \mathbf{\Lambda} : \nabla \tilde{\mathbf{d}}_0) - \frac{1}{2\varepsilon} (|\mathbf{d}_0|^2 - 1, |\tilde{\mathbf{d}}_0|^2 - 1) \\ & - \frac{1}{2\varepsilon} \int_0^t (((|\mathbf{d}|^2 - 1) - (|\tilde{\mathbf{d}}|^2 - 1), 2\partial_t \tilde{\mathbf{d}} \cdot (\tilde{\mathbf{d}} - \mathbf{d})) + (\partial_t |\mathbf{d} - \tilde{\mathbf{d}}|^2, |\tilde{\mathbf{d}}|^2 - 1)) \, ds \\ & - \int_0^t (\langle \mathbf{v}, \partial_t \tilde{\mathbf{v}} \rangle + \langle \partial_t \mathbf{v}, \tilde{\mathbf{v}} \rangle + (\mathbf{q}, \partial_t \tilde{\mathbf{d}}) + (\partial_t \mathbf{d}, \tilde{\mathbf{q}})) \, ds. \end{aligned} \quad (5.6)$$

Note that the sum of terms with the initial conditions $(\mathbf{v}_0, \mathbf{d}_0)$ and $(\tilde{\mathbf{v}}_0, \tilde{\mathbf{d}}_0)$ appearing in (5.5) and (5.6) is the relative energy of the initial values,

$$\frac{1}{2} \|\mathbf{v}_0\|_{\mathbf{L}^2}^2 + \frac{1}{2} \|\tilde{\mathbf{v}}_0\|_{\mathbf{L}^2}^2 + \mathcal{F}(\mathbf{d}_0) + \mathcal{F}(\tilde{\mathbf{d}}_0) - (\mathbf{v}_0, \tilde{\mathbf{v}}_0) - (\nabla \mathbf{d}_0 \mathbf{\Lambda} \nabla \tilde{\mathbf{d}}_0) - \frac{1}{2\varepsilon} (|\mathbf{d}_0|^2 - 1, |\tilde{\mathbf{d}}_0|^2 - 1) = \mathcal{E}(\mathbf{v}_0, \mathbf{d}_0 | \tilde{\mathbf{v}}_0, \tilde{\mathbf{d}}_0).$$

We use the fact that (\mathbf{v}, \mathbf{d}) and $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$ are solutions to calculate the last line in (5.6) explicitly. In order to handle the last line in (5.6), we use the fact that (\mathbf{v}, \mathbf{d}) and $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$ are solutions to (2.1). This shows with (5.5) that

$$\begin{aligned} & \mathcal{E}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) + \int_0^t \mathcal{W}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) \, ds \leq \mathcal{E}(\mathbf{v}_0, \mathbf{d}_0 | \tilde{\mathbf{v}}_0, \tilde{\mathbf{d}}_0) + \\ & \int_0^t (((\tilde{\mathbf{v}} \cdot \nabla) \tilde{\mathbf{v}}, \mathbf{v}) + ((\mathbf{v} \cdot \nabla) \mathbf{v}, \tilde{\mathbf{v}})) \, ds + \int_0^t (((\tilde{\mathbf{v}} \cdot \nabla) \tilde{\mathbf{d}}, \mathbf{q}) + ((\mathbf{v} \cdot \nabla) \mathbf{d}, \tilde{\mathbf{q}}) - (\nabla \tilde{\mathbf{d}}^T \tilde{\mathbf{q}}, \mathbf{v}) - (\nabla \mathbf{d}^T \mathbf{q}, \tilde{\mathbf{v}})) \, ds \\ & + \mu_1 \int_0^t ((\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, (\nabla \mathbf{v})_{\text{sym}} : (\tilde{\mathbf{d}} \otimes \tilde{\mathbf{d}} - \mathbf{d} \otimes \mathbf{d})) + (\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d}, (\nabla \tilde{\mathbf{v}})_{\text{sym}} : (\mathbf{d} \otimes \mathbf{d} - \tilde{\mathbf{d}} \otimes \tilde{\mathbf{d}})) \, ds \\ & + (\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)) \int_0^t (((\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, (\nabla \mathbf{v})_{\text{sym}} (\tilde{\mathbf{d}} - \mathbf{d})) + ((\nabla \mathbf{v})_{\text{sym}} \mathbf{d}, (\nabla \tilde{\mathbf{v}})_{\text{sym}} (\mathbf{d} - \tilde{\mathbf{d}}))) \, ds \\ & - \gamma(\mu_2 + \mu_3) \int_0^t ((\tilde{\mathbf{d}} \otimes \tilde{\mathbf{q}}; (\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}) + (\mathbf{d} \otimes \mathbf{q}; (\nabla \tilde{\mathbf{v}})_{\text{sym}} - (\nabla \mathbf{v})_{\text{sym}})) \, ds \\ & - \int_0^t ((\tilde{\mathbf{d}} \otimes \tilde{\mathbf{q}}; (\nabla \mathbf{v})_{\text{skw}}) + (\mathbf{d} \otimes \mathbf{q}; (\nabla \tilde{\mathbf{v}})_{\text{skw}}) - ((\nabla \tilde{\mathbf{v}})_{\text{skw}} \tilde{\mathbf{d}}, \mathbf{q}) - ((\nabla \mathbf{v})_{\text{skw}} \mathbf{d}, \tilde{\mathbf{q}})) \, ds \\ & + \lambda \int_0^t (((\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, \mathbf{q} - \tilde{\mathbf{q}}) + ((\nabla \mathbf{v})_{\text{sym}} \mathbf{d}, \tilde{\mathbf{q}} - \mathbf{q})) \, ds \\ & - \frac{1}{2\varepsilon} \int_0^t (((|\mathbf{d}|^2 - 1) - (|\tilde{\mathbf{d}}|^2 - 1), 2\partial_t \tilde{\mathbf{d}} \cdot (\tilde{\mathbf{d}} - \mathbf{d})) + (\partial_t |\mathbf{d} - \tilde{\mathbf{d}}|^2, |\tilde{\mathbf{d}}|^2 - 1)) \, ds \\ & = \mathcal{E}(\mathbf{v}_0, \mathbf{d}_0 | \tilde{\mathbf{v}}_0, \tilde{\mathbf{d}}_0) + I_1 + I_2 + \mu_1 I_3 + (\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)) I_4 - \gamma(\mu_2 + \mu_3) I_5 + I_6 + \lambda I_7 - \frac{1}{2\varepsilon} I_8. \end{aligned}$$

In the above equation we have employed the weak formulation for the solutions (\mathbf{v}, \mathbf{d}) and $(\tilde{\mathbf{v}}, \tilde{\mathbf{d}})$ tested with $(\tilde{\mathbf{v}}, \tilde{\mathbf{q}})$ and (\mathbf{v}, \mathbf{q}) , respectively. Note that the choice of test functions is justified due to the additional regularity (see Lemma 3.2). We calculate and estimate the terms in the above equation individually. In the following let $\delta > 0$. For the first term I_1 , we observe that

$$\begin{aligned} I_1 &= \int_0^t (((\mathbf{v} \cdot \nabla (\mathbf{v} - \tilde{\mathbf{v}}), \tilde{\mathbf{v}} - \mathbf{v}) + (((\mathbf{v} - \tilde{\mathbf{v}}) \cdot \nabla) \tilde{\mathbf{v}}, \tilde{\mathbf{v}} - \mathbf{v})) \, ds = \int_0^t ((\mathbf{v} - \tilde{\mathbf{v}}) \otimes (\tilde{\mathbf{v}} - \mathbf{v}); (\nabla \tilde{\mathbf{v}})_{\text{sym}}) \, ds \\ &\leq C_\delta \int_0^t \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^3}^2 \|\mathbf{v} - \tilde{\mathbf{v}}\|_{\mathbf{L}^2}^2 \, ds + \delta \int_0^t \|\mathbf{v} - \tilde{\mathbf{v}}\|_{\mathbf{L}^6}^2 \, ds. \end{aligned}$$

We recall that \mathbf{v} and $\tilde{\mathbf{v}}$ are solenoidal such that $((\mathbf{v} \cdot \nabla) \mathbf{w}, \mathbf{w}) = ((\tilde{\mathbf{v}} \cdot \nabla) \mathbf{w}, \mathbf{w}) = 0$ for all $\mathbf{w} \in \mathbf{H}_{0,\sigma}^1$.

The term I_2 can be estimated by

$$\begin{aligned} I_2 &= \int_0^t ((\mathbf{v} \cdot \nabla)(\mathbf{d} - \tilde{\mathbf{d}}), \tilde{\mathbf{q}}) + ((\tilde{\mathbf{v}} \cdot \nabla)(\tilde{\mathbf{d}} - \mathbf{d}), \mathbf{q}) \, ds = \int_0^t (((\mathbf{v} - \tilde{\mathbf{v}}), (\nabla \mathbf{d} - \nabla \tilde{\mathbf{d}})^T \tilde{\mathbf{q}}) + ((\tilde{\mathbf{v}} \cdot \nabla)(\tilde{\mathbf{d}} - \mathbf{d}), \mathbf{q} - \tilde{\mathbf{q}})) \, ds \\ &\leq C_\delta \int_0^t (\|\tilde{\mathbf{q}}\|_{\mathbf{L}^3}^2 \|\nabla \mathbf{d} - \nabla \tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2) \, ds + \delta \int_0^t \|\mathbf{v} - \tilde{\mathbf{v}}\|_{\mathbf{L}^6}^2 \, ds + C_\delta \int_0^t \|\tilde{\mathbf{v}}\|_{\mathbf{L}^\infty}^2 \|\nabla \tilde{\mathbf{d}} - \nabla \mathbf{d}\|_{\mathbf{L}^2}^2 \, ds + \delta \int_0^t \|\mathbf{q} - \tilde{\mathbf{q}}\|_{\mathbf{L}^2}^2 \, ds. \end{aligned}$$

With the standard embedding $\mathbf{H}_{0,\sigma}^1 \hookrightarrow \mathbf{L}^6$ and Korn's inequality [19, Theorem 10.1], there is a constant c such that $\|\mathbf{v} - \tilde{\mathbf{v}}\|_{\mathbf{L}^6} \leq c \|(\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^2}$. We rearrange and estimate the term I_3 by

$$\begin{aligned} I_3 &= \int_0^t (\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, ((\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}) : (\tilde{\mathbf{d}} \otimes \tilde{\mathbf{d}} - \mathbf{d} \otimes \mathbf{d})) \, ds \\ &\quad + \int_0^t (\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - \tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, (\nabla \tilde{\mathbf{v}})_{\text{sym}} : (\mathbf{d} \otimes \mathbf{d} - \tilde{\mathbf{d}} \otimes \tilde{\mathbf{d}})) \, ds \\ &= \int_0^t (\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, ((\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}) : (\tilde{\mathbf{d}} \otimes (\tilde{\mathbf{d}} - \mathbf{d}))) \, ds \\ &\quad + \int_0^t (\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, ((\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}) \cdot (\tilde{\mathbf{d}} - \mathbf{d})) \, ds \\ &\quad + \int_0^t (\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, (\nabla \tilde{\mathbf{v}})_{\text{sym}} : (\tilde{\mathbf{d}} - \mathbf{d}) \otimes (\tilde{\mathbf{d}} - \mathbf{d})) \, ds \\ &\quad + \int_0^t (\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - \tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, (\nabla \tilde{\mathbf{v}})_{\text{sym}} : (\mathbf{d} - \tilde{\mathbf{d}}) \otimes \tilde{\mathbf{d}}) \, ds \\ &\quad + \int_0^t (\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - \tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, (\nabla \tilde{\mathbf{v}})_{\text{sym}} : \mathbf{d} \otimes (\mathbf{d} - \tilde{\mathbf{d}})) \, ds \\ &\leq C_\delta \|\tilde{\mathbf{d}}\|_{\mathbf{L}^\infty(\mathbf{L}^6)}^2 \int_0^t \|\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \, ds + \delta \int_0^t \|(\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^2}^2 \, ds \\ &\quad + C_\delta \int_0^t \|\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^3}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \, ds + \delta \int_0^t \|(\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2 \, ds \\ &\quad + \int_0^t (\|\tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^3}^2 + \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^3}^2) \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \, ds \\ &\quad + C_\delta \|\tilde{\mathbf{d}}\|_{\mathbf{L}^\infty(\mathbf{L}^6)}^2 \int_0^t \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^6}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \, ds + \delta \int_0^t \|\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - \tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2 \, ds \\ &\quad + C_\delta \|\mathbf{d}\|_{\mathbf{L}^\infty(\mathbf{L}^6)}^2 \int_0^t \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^6}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \, ds + \delta \int_0^t \|\mathbf{d} \cdot (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - \tilde{\mathbf{d}} \cdot (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2 \, ds. \end{aligned}$$

The embedding $\mathbf{H}_0^1 \hookrightarrow \mathbf{L}^6$ together with Poincaré's inequality (see Morrey [20, Thm. 6.5.6.]) assures that

$$\|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6} \leq c \|\nabla \mathbf{d} - \nabla \tilde{\mathbf{d}}\|_{\mathbf{L}^2}.$$

We continue with I_4 ,

$$\begin{aligned} I_4 &= \int_0^t ((\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, ((\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}) (\tilde{\mathbf{d}} - \mathbf{d})) \, ds + \int_0^t ((\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}, (\nabla \tilde{\mathbf{v}})_{\text{sym}} (\mathbf{d} - \tilde{\mathbf{d}})) \, ds \\ &\leq C_\delta \int_0^t \|(\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^3}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \, ds + \delta \int_0^t \|(\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^2}^2 \, ds \\ &\quad + C_\delta \int_0^t \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^3}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \, ds + \delta \int_0^t \|(\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2 \, ds. \end{aligned}$$

The term I_5 can be rearranged as

$$\begin{aligned} I_5 &= \int_0^t \left(((\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}, (\tilde{\mathbf{d}} - \mathbf{d}) \otimes \tilde{\mathbf{q}}) + ((\nabla \tilde{\mathbf{v}})_{\text{sym}} - (\nabla \mathbf{v})_{\text{sym}}, \mathbf{d} \otimes (\mathbf{q} - \tilde{\mathbf{q}})) \right) \mathrm{d}s \\ &= \int_0^t \left(((\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}, (\tilde{\mathbf{d}} - \mathbf{d}) \otimes \tilde{\mathbf{q}}) + ((\nabla \tilde{\mathbf{v}})_{\text{sym}}(\mathbf{d} - \tilde{\mathbf{d}}), \mathbf{q} - \tilde{\mathbf{q}}) \right) \mathrm{d}s \\ &\quad + \int_0^t \left((\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}} - (\nabla \mathbf{v})_{\text{sym}} \mathbf{d}, \mathbf{q} - \tilde{\mathbf{q}} \right) \mathrm{d}s, \end{aligned}$$

and thus be estimated by

$$\begin{aligned} I_5 &\leq \delta \int_0^t \|(\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^2}^2 \mathrm{d}s + \delta \int_0^t \|\mathbf{q} - \tilde{\mathbf{q}}\|_{\mathbf{L}^2}^2 \mathrm{d}s + C_\delta \int_0^t (\|\tilde{\mathbf{q}}\|_{\mathbf{L}^3}^2 + \|(\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^3}^2) \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \mathrm{d}s \\ &\quad + \int_0^t \left((\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}} - (\nabla \mathbf{v})_{\text{sym}} \mathbf{d}, \mathbf{q} - \tilde{\mathbf{q}} \right) \mathrm{d}s. \end{aligned}$$

The term I_6 is bounded by

$$\begin{aligned} I_6 &= \int_0^t \left(((\nabla \mathbf{v})_{\text{skw}}(\tilde{\mathbf{d}} - \mathbf{d}), \tilde{\mathbf{q}}) + ((\nabla \tilde{\mathbf{v}})_{\text{skw}}(\mathbf{d} - \tilde{\mathbf{d}}), \mathbf{q}) \right) \mathrm{d}s \\ &= \int_0^t \left((((\nabla \mathbf{v})_{\text{skw}} - (\nabla \tilde{\mathbf{v}})_{\text{skw}})(\tilde{\mathbf{d}} - \mathbf{d}), \tilde{\mathbf{q}}) + ((\nabla \tilde{\mathbf{v}})_{\text{skw}}(\mathbf{d} - \tilde{\mathbf{d}}), \mathbf{q} - \tilde{\mathbf{q}}) \right) \mathrm{d}s \\ &\leq C_\delta \int_0^t \|\tilde{\mathbf{q}}\|_{\mathbf{L}^3}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \mathrm{d}s + \delta \int_0^t \|(\nabla \mathbf{v})_{\text{skw}} - (\nabla \tilde{\mathbf{v}})_{\text{skw}}\|_{\mathbf{L}^2}^2 \mathrm{d}s \\ &\quad + C_\delta \int_0^t \|(\nabla \tilde{\mathbf{v}})_{\text{skw}}\|_{\mathbf{L}^3}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \mathrm{d}s + \delta \int_0^t \|\mathbf{q} - \tilde{\mathbf{q}}\|_{\mathbf{L}^2}^2 \mathrm{d}s. \end{aligned}$$

Note that due to Korn's [19, Theorem 10.1] and Poincaré's inequality, we find that

$$\|(\nabla \mathbf{v})_{\text{skw}} - (\nabla \tilde{\mathbf{v}})_{\text{skw}}\|_{\mathbf{L}^2} \leq \|\mathbf{v} - \tilde{\mathbf{v}}\|_{\mathbf{H}_0^1} \leq c \|(\nabla \mathbf{v})_{\text{sym}} - (\nabla \tilde{\mathbf{v}})_{\text{sym}}\|_{\mathbf{L}^2}.$$

The term I_7 is already in the form desired. Finally, we estimate I_8 . Starting with the first term, we observe that

$$\int_0^t \left((|\mathbf{d}|^2 - 1) - (|\tilde{\mathbf{d}}|^2 - 1) \right) (\tilde{\mathbf{d}} - \mathbf{d}), \partial_t \tilde{\mathbf{d}} \mathrm{d}s \leq \int_0^t \|\partial_t \tilde{\mathbf{d}}\|_{\mathbf{L}^3} \left(\|\mathbf{d}\|_{\mathbf{L}^2}^2 - \|\tilde{\mathbf{d}}\|_{\mathbf{L}^2}^2 + \|\mathbf{d} - \tilde{\mathbf{d}}\|_{\mathbf{L}^6}^2 \right) \mathrm{d}s.$$

Since $\tilde{\mathbf{d}}$ is a strong solution (see Definition 2.3), $\partial_t \tilde{\mathbf{d}}$ is in $L^1(0, T; \mathbf{L}^3)$ due to Lemma 3.2.

Now we reformulate the second term of I_8 . Using that equation (2.10b) is fulfilled by \mathbf{d} and $\tilde{\mathbf{d}}$, respectively, yields

$$\begin{aligned} &\frac{1}{2} \int_0^t (|\tilde{\mathbf{d}}|^2 - 1, \partial_t |\mathbf{d} - \tilde{\mathbf{d}}|^2) \mathrm{d}s \\ &= \int_0^t ((|\tilde{\mathbf{d}}|^2 - 1)(\mathbf{d} - \tilde{\mathbf{d}}), \partial_t \mathbf{d} - \partial_t \tilde{\mathbf{d}}) \mathrm{d}s \\ &= \int_0^t ((|\tilde{\mathbf{d}}|^2 - 1)(\tilde{\mathbf{d}} - \mathbf{d}), (\mathbf{v} \cdot \nabla) \mathbf{d} - (\tilde{\mathbf{v}} \cdot \nabla) \tilde{\mathbf{d}} - (\nabla \mathbf{v})_{\text{skw}} \mathbf{d} + (\nabla \tilde{\mathbf{v}})_{\text{skw}} \tilde{\mathbf{d}}) \mathrm{d}s \\ &\quad + \int_0^t ((|\tilde{\mathbf{d}}|^2 - 1)(\tilde{\mathbf{d}} - \mathbf{d}), \lambda((\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}) + \gamma(\mathbf{q} - \tilde{\mathbf{q}})) \mathrm{d}s \\ &= J_1 + J_2. \end{aligned}$$

The term J_1 can be rewritten as

$$\begin{aligned} J_1 &= \int_0^t \left((|\tilde{\mathbf{d}}|^2 - 1)(\tilde{\mathbf{d}} - \mathbf{d}), \nabla \tilde{\mathbf{d}}(\mathbf{v} - \tilde{\mathbf{v}}) \right) \mathrm{d}s + \int_0^t (|\tilde{\mathbf{d}}|^2 - 1, \mathbf{v} \cdot (\nabla \mathbf{d} - \nabla \tilde{\mathbf{d}})(\tilde{\mathbf{d}} - \mathbf{d})) \mathrm{d}s \\ &\quad - \int_0^t \left((|\tilde{\mathbf{d}}|^2 - 1)(\tilde{\mathbf{d}} - \mathbf{d}), (\nabla \mathbf{v})_{\text{skw}}(\mathbf{d} - \tilde{\mathbf{d}}) \right) \mathrm{d}s - \int_0^t \left((|\tilde{\mathbf{d}}|^2 - 1)(\tilde{\mathbf{d}} - \mathbf{d}), ((\nabla \mathbf{v})_{\text{skw}} - (\nabla \tilde{\mathbf{v}})_{\text{skw}}) \tilde{\mathbf{d}} \right) \mathrm{d}s. \end{aligned}$$

We observe that the third term, i.e., $((|\tilde{\mathbf{d}}|^2 - 1)(\tilde{\mathbf{d}} - \mathbf{d}), (\nabla \mathbf{v})_{\text{skw}}(\mathbf{d} - \tilde{\mathbf{d}}))$ vanishes since $(\nabla \mathbf{v})_{\text{skw}}$ is skew-symmetric. Hence, we can estimate J_1 by

$$\begin{aligned} J_1 &\leq C_\delta \int_0^t (\|\tilde{\mathbf{d}}\|^2 - 1)_{L^6}^2 + \|\nabla \tilde{\mathbf{d}}\|_{L^2}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{L^6}^2 ds + \delta \int_0^t \|\mathbf{v} - \tilde{\mathbf{v}}\|_{L^6}^2 ds \\ &\quad + \int_0^t (\|\tilde{\mathbf{d}}\|^2 - 1)_{L^6}^2 + \|\mathbf{v}\|_{L^6}^2 (\|\nabla \mathbf{d} - \nabla \tilde{\mathbf{d}}\|_{L^2}^2 + \|\mathbf{d} - \tilde{\mathbf{d}}\|_{L^6}^2) ds \\ &\quad + C_\delta \|\tilde{\mathbf{d}}\|_{L^\infty(L^6)}^2 \int_0^t \|\tilde{\mathbf{d}}\|^2 - 1 \|_{L^6}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{L^6}^2 ds + \delta \int_0^t \|(\nabla \mathbf{v})_{\text{skw}} - (\nabla \tilde{\mathbf{v}})_{\text{skw}}\|_{L^2}^2 ds. \end{aligned}$$

It remains to estimate the term J_2 :

$$J_2 \leq C_\delta \|\tilde{\mathbf{d}}\|_{L^\infty(L^6)}^2 \int_0^t \|\tilde{\mathbf{d}}\|^2 - 1 \|_{L^6}^2 \|\mathbf{d} - \tilde{\mathbf{d}}\|_{L^6}^2 ds + \delta \int_0^t (\lambda \|(\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{L^2}^2 + \gamma \|\mathbf{q} - \tilde{\mathbf{q}}\|_{L^2}^2) ds.$$

Inserting everything back into (5.5) yields

$$\begin{aligned} \mathcal{E}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}})(t) + \int_0^t \mathscr{W}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) ds &\leq \mathcal{E}(\mathbf{v}_0, \mathbf{d}_0 | \tilde{\mathbf{v}}_0, \tilde{\mathbf{d}}_0) \\ &\quad + (\gamma(\mu_2 + \mu_3) - \lambda) \int_0^t (\mathbf{q} - \tilde{\mathbf{q}}, (\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}) ds \\ &\quad + \delta c \int_0^t \mathscr{W}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) ds + \int_0^t \mathscr{K}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) \mathcal{E}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) ds. \end{aligned}$$

Since the constants are assumed to fulfill the dissipative relation (2.6) we can find a real number $\zeta \in (0, 1)$ such that

$$(\gamma(\mu_2 + \mu_3) - \lambda)^2 \leq \zeta^2 4\gamma(\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)). \quad (5.7)$$

The relative energy can be estimated further on with Youngs and Hölder's inequality, such that

$$\begin{aligned} \mathcal{E}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}})(t) + \int_0^t \mathscr{W}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) ds &\leq \mathcal{E}(\mathbf{v}_0, \mathbf{d}_0 | \tilde{\mathbf{v}}_0, \tilde{\mathbf{d}}_0) \\ &\quad + \zeta \int_0^t (\gamma \|\mathbf{q} - \tilde{\mathbf{q}}\|_{L^2}^2 + (\mu_5 + \mu_6 - \lambda(\mu_2 + \mu_3)) \|(\nabla \mathbf{v})_{\text{sym}} \mathbf{d} - (\nabla \tilde{\mathbf{v}})_{\text{sym}} \tilde{\mathbf{d}}\|_{L^2}^2) ds \\ &\quad + \delta c \int_0^t \mathscr{W}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) ds + \int_0^t \mathscr{K}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) \mathcal{E}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) ds \\ &\leq \mathcal{E}(\mathbf{v}_0, \mathbf{d}_0 | \tilde{\mathbf{v}}_0, \tilde{\mathbf{d}}_0) + (\zeta + \delta c) \int_0^t \mathscr{W}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) ds + \int_0^t \mathscr{K}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) \mathcal{E}(\mathbf{v}, \mathbf{d} | \tilde{\mathbf{v}}, \tilde{\mathbf{d}}) ds. \end{aligned}$$

We now choose δ sufficiently small such that $\delta \leq (1 - \zeta)/c$. Thus, the relative dissipation \mathscr{W} can be absorbed into the left-hand side. Note that c does not depend on δ , but only on the constants arising from the embeddings and Korn's inequality as well as the constants of the system (see (2.6)). The assertion (5.3) immediately follows from Gronwall's estimate (see Remark 5.2). \square

Proof of Theorem 2.5. The main Theorem 2.5 is a direct consequence of Lemma 5.1. \square

References

- [1] A. N. Beris and B. J. Edwards. *Thermodynamics of flowing systems with internal microstructure*. Oxford University Press, New York, 1994.
- [2] D. Breit, E. Feireisl, and M. Hofmanová. Incompressible limit for compressible fluids with stochastic forcing. *Arch. Rational Mech. Anal.*, 222(2):895–926, 2016.

- [3] C. Cavaterra, E. Rocca, and H. Wu. Global weak solution and blow-up criterion of the general Ericksen–Leslie system for nematic liquid crystal flows. *J. Differential Equations*, 255(1):24–57, 2013.
- [4] M. Dai. Existence of regular solutions to an Ericksen–Leslie model of the liquid crystal system. *Commun. Math. Sci.*, 13(7):1711–1740, 2015.
- [5] M. Dai, J. Qing, and M. Schonbek. Regularity of solutions to the liquid crystals systems in \mathbb{R}^2 and \mathbb{R}^3 . *Nonlinearity*, 25(2):513–532, 2012.
- [6] J. Diestel and J. J. Uhl, Jr. *Vector measures*. American Mathematical Society, Providence, Rhode Island, 1977.
- [7] E. Emmrich and R. Lasarzik. Existence of weak solutions to the Ericksen–Leslie model for a general class of free energies. *ArXiv e-prints*, 1711.10277, 2017.
- [8] J. L. Ericksen. Conservation laws for liquid crystals. *J. Rheol.*, 5:23–34, 1961.
- [9] E. Feireisl. Relative entropies in thermodynamics of complete fluid systems. *Discrete Contin. Dyn. Syst.*, 32(9):3059–3080, 2012.
- [10] E. Feireisl. Relative entropies, dissipative solutions, and singular limits of complete fluid systems. In *Hyperbolic Problems: Theory, Numerics, Applications*, volume 8 of *AIMS on Applied Mathematics*, pages 11–28. AIMS, Springfield, USA, 2014.
- [11] E. Feireisl, B. J. Jin, and A. Novotný. Relative entropies, suitable weak solutions, and weak-strong uniqueness for the compressible Navier–Stokes system. *J. Math. Fluid Mech.*, 14(4):717–730, 2012.
- [12] E. Feireisl, A. Novotný, and Y. Sun. Suitable weak solutions to the Navier–Stokes equations of compressible viscous fluids. *Indiana Univ. Math. J.*, 60(2):611–631, 2011.
- [13] J. Fischer. A posteriori modeling error estimates for the assumption of perfect incompressibility in the Navier–Stokes equation. *SIAM J. Numer. Anal.*, 53(5):2178–2205, 2015.
- [14] R. Lasarzik. Measure-valued solutions to the Ericksen–Leslie model equipped with the Oseen–Frank energy. *ArXiv e-prints*, 1711.04638, Nov. 2017.
- [15] R. Lasarzik. Weak-strong uniqueness for measure-valued solutions to the Ericksen–Leslie model equipped with the Oseen–Frank free energy. *ArXiv e-prints*, 1711.03371, Nov. 2017.
- [16] F. M. Leslie. Some constitutive equations for liquid crystals. *Arch. Rational Mech. Anal.*, 28(4):265–283, 1968.
- [17] F.-H. Lin and C. Liu. Nonparabolic dissipative systems modeling the flow of liquid crystals. *Comm. Pure Appl. Math.*, 48(5):501–537, 1995.
- [18] F.-H. Lin and C. Liu. Existence of solutions for the Ericksen–Leslie system. *Arch. Rational Mech. Anal.*, 154(2):135–156, 2000.
- [19] W. McLean. *Strongly elliptic systems and boundary integral equations*. Cambridge University Press, Cambridge, 2000.
- [20] C. B. Morrey Jr. *Multiple integrals in the calculus of variations*. Springer, Berlin, 1966.
- [21] G. Prodi. Un teorema di unicità per le equazioni di Navier–Stokes. *Ann. Mat. Pura Appl. (4)*, 48:173–182, 1959.
- [22] T. Roubíček. *Nonlinear partial differential equations with applications*. Birkhäuser, Basel, 2005.
- [23] J. Serrin. On the interior regularity of weak solutions of the Navier–Stokes equations. *Arch. Rational Mech. Anal.*, 9:187–195, 1962.
- [24] Y.-F. Yang, C. Dou, and Q. Ju. Weak-strong uniqueness property for the compressible flow of liquid crystals. *J. Differential Equations*, 255(6):1233–1253, 2013.
- [25] E. Zeidler. *Nonlinear functional analysis and its applications. II/A*. Springer-Verlag, New York, 1990.
- [26] J.-H. Zhao and Q. Liu. Weak-strong uniqueness of hydrodynamic flow of nematic liquid crystals. *Electron. J. Differential Equations*, 2012(182):1–16, 2012.