

# HARDY'S INEQUALITY FOR THE FRACTIONAL POWERS OF THE GRUSHIN OPERATOR

RAKESH BALHARA

ABSTRACT. We prove Hardy's inequality for the fractional powers of the generalized sublaplacian and the fractional powers of the Grushin operator. We also find an integral representation and a ground state representation for the fractional powers of generalized sublaplacian.

## 1. INTRODUCTION AND MAIN RESULTS

The study of various kinds of inequalities for various differential operators are important in understanding many practical problems in physics. Moreover, sharpness of the constants involved in these inequalities is directly related to the existence and nonexistence results for certain partial differential equations.

The well known Hardy's inequality for continuously differentiable functions on  $\mathbb{R}^n$  ( $n \geq 3$ ) is given by

$$(1.1) \quad \int_{\mathbb{R}^n} |\nabla f|^2 dx \geq c_n \int_{\mathbb{R}^n} \frac{|f(x)|^2}{|x|^2} dx,$$

where  $\nabla$  is the standard gradient operator on  $\mathbb{R}^n$ . Moreover, sharp value of the constant  $c_n$ , involved in the inequality is known to be equal to  $\frac{(n-2)^2}{4}$ . By sharp value of the constant, we mean that the inequality will not hold true if we take value of  $c_n > \frac{(n-2)^2}{4}$ .

We recall that the classical Laplacian  $\Delta$  on  $\mathbb{R}^n$  is defined by  $\Delta = -\sum_{j=1}^n \frac{\partial^2}{\partial x_j^2}$ . For  $f \in L^2(\mathbb{R}^n)$  such that  $\Delta f \in L^2(\mathbb{R}^n)$ , the inequality (1.1) can be shown equivalent to the inequality:

$$(1.2) \quad \langle \Delta f, f \rangle \geq \frac{(n-2)^2}{4} \int_{\mathbb{R}^n} \frac{|f(x)|^2}{|x|^2} dx.$$

Hardy's inequality has been generalized for fractional powers of Laplacian. Recall that fractional powers of Laplacian  $\Delta^s$ , for  $s > 0$  is defined via spectral decomposition (or Fourier transform) as

$$(1.3) \quad \Delta^s f(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} |\xi|^{2s} \widehat{f}(\xi) e^{ix \cdot \xi} d\xi,$$

---

2010 *Mathematics Subject Classification*. Primary: 35A23. Secondary: 26A33, 26D10, 42B37, 42C10, 47A63.

*Key words and phrases*. fractional Grushin operator, fractional generalized sublaplacian, Hardy's Inequality, ground state representation, Hecke-Bochner formula.

Author is financially supported by UGC-CSIR. Also he wants to thank his guide Prof S. Thangavelu for his continuous help and suggestions.

where  $\widehat{f}(\xi)$  is the Fourier transform defined by  $\int_{\mathbb{R}^n} f(x)e^{-ix \cdot \xi} dx$ . With this definition, we state the Hardy's inequality for the fractional powers of Laplacian. For  $f \in L^2(\mathbb{R}^n)$  such that  $\Delta^s f \in L^2(\mathbb{R}^n)$ , we have for  $0 < s < 1$

$$(1.4) \quad \langle \Delta^s f, f \rangle \geq 4^s \left( \frac{\Gamma(\frac{n+2s}{4})}{\Gamma(\frac{n-2s}{4})} \right)^2 \int_{\mathbb{R}^n} \frac{|f(x)|^2}{|x|^{2s}} dx.$$

Though the constant involved in the inequality is sharp, the equality is never achieved for any non zero function. Hardy's inequality for the fractional powers of Laplacian has been extensively studied in literature. We refer to [1][4][6][13] for more details.

On the other hand, there is another version of Hardy's inequality for fractional powers of Laplacian, where the homogeneous weight  $|x|^{2s}$  has been replaced by a non-homogeneous weight  $(\delta + |x|^2)^{2s}$ ,  $\delta > 0$ :

$$(1.5) \quad \langle \Delta^s f, f \rangle \geq (4\delta)^s \left( \frac{\Gamma(\frac{n+2s}{2})}{\Gamma(\frac{n-2s}{2})} \right)^2 \int_{\mathbb{R}^n} \frac{|f(x)|^2}{(\delta + |x|^2)^{2s}} dx.$$

The constant in the inequality is sharp and equality is achieved for  $f(x) = (\delta + |x|^2)^{-(n-2s)/2}$ . Though the inequality (1.5) is well known, we are unable to find a reference where this inequality is actually proved.

In this article, we are interested in proving a similar inequality for the fractional powers of Grushin operator. Recall that the Grushin operator  $\mathcal{G}$  on  $\mathbb{R}^{n+1}$  is defined by

$$(1.6) \quad \mathcal{G} = -\frac{1}{2} \left( \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2} + |x|^2 \frac{\partial^2}{\partial w^2} \right).$$

Using the spectral decomposition, we can define fractional powers of Grushin operator  $\mathcal{G}^s$  for any  $s > 0$  by

$$(1.7) \quad \mathcal{G}^s f(x, w) = \frac{1}{2\pi} \int_{\mathbb{R}} \sum_{k=0}^{\infty} ((k+n/2)|\lambda|)^s \mathcal{P}_k(\lambda) f^\lambda(x) e^{-i\lambda w} d\lambda,$$

where  $\mathcal{P}_k(\lambda)$  are the orthogonal projections of  $L^2(\mathbb{R}^n)$  onto the eigenspaces  $E_k^\lambda$  corresponding to the eigenvalues  $(2k+n)|\lambda|$  of the scaled Hermite operator  $\mathcal{H}(\lambda)$  defined on  $\mathbb{R}^n$  as

$$(1.8) \quad \mathcal{H}(\lambda) = -\sum_{j=1}^n \frac{\partial^2}{\partial x_j^2} + \lambda^2 |x|^2,$$

and  $f^\lambda$  is the inverse Fourier transform of  $f$  in the the last variable, that is

$$(1.9) \quad f^\lambda(x) = \int_{\mathbb{R}} f(x, w) e^{iw\lambda} dw.$$

However, it is convenient to work with the following modified fractional powers:

$$(1.10) \quad \mathcal{G}_s f(x, w) = \frac{1}{2\pi} \int_{\mathbb{R}} \sum_{k=0}^{\infty} (2|\lambda|)^s \frac{\Gamma(\frac{2k+n}{4} + \frac{1+s}{2})}{\Gamma(\frac{2k+n}{4} + \frac{1-s}{2})} \mathcal{P}_k(\lambda) f^\lambda(x) e^{-i\lambda w} d\lambda.$$

Notice that  $\mathcal{G}^s$  differs from  $\mathcal{G}_s$  by a bounded operator, that is there exists a bounded operator  $V_s$  such that  $\mathcal{G}^s = V_s \mathcal{G}_s$ , which justifies the proving of Hardy type inequality for  $\mathcal{G}_s$ .

We denote by  $W^{s,2}(\mathbb{R}^{n+1})$ , the Sobolev space consisting of all  $L^2(\mathbb{R}^{n+1})$  functions such that  $\mathcal{G}_s f \in L^2(\mathbb{R}^{n+1})$ . The main theorem that we will prove in this article is the following:

**Theorem 1.1.** *For  $f \in W^{s,2}(\mathbb{R}^{n+1})$ ,  $0 < s < 1$  and  $\delta > 0$ , we have*

$$\langle \mathcal{G}_s f, f \rangle \geq (4\delta)^s \left( \frac{\Gamma(\frac{n/2+s+1}{2})}{\Gamma(\frac{n/2-s+1}{2})} \right)^2 \int_{\mathbb{R}^{n+1}} \frac{|f(x, w)|^2}{\left( \left( \delta + \frac{|x|^2}{2} \right)^2 + w^2 \right)^s} dx dw.$$

Also the constant in the inequality is sharp and equality is achieved for

$$f(x, w) = \left( \left( \delta + \frac{|x|^2}{2} \right)^2 + w^2 \right)^{-\frac{n/2-s+1}{2}}.$$

In order to prove the above theorem, we prove an analogous theorem for the fractional powers of generalized sublaplacian. We define generalized sublaplacian  $\mathcal{L}$  on  $\mathbb{R}^+ \times \mathbb{R}$  for  $\alpha > -1/2$  by

$$(1.11) \quad \mathcal{L} = -\frac{1}{2} \left( \frac{\partial^2}{\partial x^2} + \frac{2\alpha+1}{x} \frac{\partial}{\partial x} + x^2 \frac{\partial^2}{\partial w^2} \right).$$

Using the spectral decomposition, we define fractional powers of generalized sublaplacian  $\mathcal{L}^s$  for  $s > 0$  as

$$(1.12) \quad \mathcal{L}^s f(x, w) = \frac{1}{\pi \Gamma(\alpha+1)} \int_{\mathbb{R}} \sum_{k=0}^{\infty} \left( (|\lambda|(2k+\alpha+1))^s \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x) \right) |\lambda|^{\alpha+1} e^{-i\lambda w} d\lambda,$$

where  $\widehat{f}(\lambda, k)$  is the Laguerre transform defined as

$$\widehat{f}(\lambda, k) = \frac{\Gamma(\alpha+1)\Gamma(k+1)}{\Gamma(\alpha+k+1)} \int_0^\infty \left( \int_{\mathbb{R}} f(x, w) e^{i\lambda w} dw \right) \phi_{k,\lambda}^\alpha(x) x^{2\alpha+1} dx,$$

and  $\phi_{k,\lambda}^\alpha$  defined as

$$\phi_{k,\lambda}^\alpha(x) = L_k^\alpha(|\lambda|x^2) e^{-\frac{1}{2}|\lambda|x^2},$$

with  $L_k^\alpha$  denoting the Laguerre polynomials of order  $\alpha$ . However, it is convenient to work with the following modified fractional powers of  $\mathcal{L}$ . For  $s > 0$ , we define  $\mathcal{L}_s$  by

$$(1.13) \quad \mathcal{L}_s f(x, w) = \frac{1}{\pi \Gamma(\alpha+1)} \int_{\mathbb{R}} \sum_{k=0}^{\infty} \left( (2|\lambda|)^s \frac{\Gamma(\frac{2k+\alpha+1}{2} + \frac{1+s}{2})}{\Gamma(\frac{2k+\alpha+1}{2} + \frac{1-s}{2})} \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x) \right) |\lambda|^{\alpha+1} e^{-i\lambda w} d\lambda.$$

Again, we denote by  $W^{s,2}(\mathbb{R}^+ \times \mathbb{R})$ , the space consisting of all functions  $f$  such that both  $f$  and  $\mathcal{L}_s f$  belong to  $L^2(\mathbb{R}^+ \times \mathbb{R})$ . We will prove the following Hardy type inequality for  $\mathcal{L}_s$ .

**Theorem 1.2.** *For  $f \in W^{s,2}(\mathbb{R}^+ \times \mathbb{R})$ ,  $0 < s < 1$  and  $\delta > 0$ , we have*

$$\langle \mathcal{L}_s f, f \rangle \geq (4\delta)^s \left( \frac{\Gamma(\frac{\alpha+s+2}{2})}{\Gamma(\frac{\alpha-s+2}{2})} \right)^2 \int_{\mathbb{R}^+} \int_{\mathbb{R}} \frac{|f(x, w)|^2}{\left( \left( \delta + \frac{x^2}{2} \right)^2 + w^2 \right)^s} dw dx.$$

Moreover, the constant in the inequality is sharp and equality is achieved for

$$f(x, w) = ((\delta + x^2/2)^2 + w^2)^{-\frac{\alpha-s+2}{2}}.$$

We outline the contents of this paper. In Section 2, we give preliminaries, definitions and facts concerning Laguerre expansions, fractional powers of sublaplacian, fractional powers of Grushin, spherical harmonics and Hecke Bochner formula. In Section 3, we will prove the Hardy type inequality for the fractional powers of sublaplacian. Integral representation and ground state representation for the fractional powers of sublaplacian are also calculated in this Section. In Section 4, Hardy type inequality for the fractional powers of Grushin will be proved.

## 2. PRELIMINARIES

**2.1. Laguerre expansions on  $\mathbb{R}^+$ .** Let  $\alpha > -1/2$ . We equip  $Y = \mathbb{R}^+$  with the measure  $x^{2\alpha+1} dx$ , where  $dx$  is the standard Lebesgue measure. For  $\lambda \in \mathbb{R} \setminus \{0\}$  and  $k = 0, 1, 2, \dots$ , we define *Laguerre functions*  $\phi_{k,\lambda}^\alpha$  by

$$(2.1) \quad \phi_{k,\lambda}^\alpha(x) = L_k^\alpha(|\lambda|x^2)e^{-\frac{1}{2}|\lambda|x^2},$$

where  $L_k^\alpha$  are the Laguerre polynomials of type  $\alpha$ . We also define

$$(2.2) \quad \tilde{\phi}_{k,\lambda}^\alpha(x) = \left( \frac{2\Gamma(k+1)|\lambda|^{\alpha+1}}{\Gamma(\alpha+k+1)} \right)^{\frac{1}{2}} \phi_{k,\lambda}^\alpha(x).$$

**Proposition 2.1.** *For  $\lambda \neq 0$ , the collection  $\{\tilde{\phi}_{k,\lambda}^\alpha(x)\}_{k=0}^\infty$  forms an orthonormal basis for  $L^2(Y, x^{2\alpha+1} dx)$ .*

For proof see [11] [Proposition 2.4.2].

For  $x \in Y$ , define *Laguerre translation*  $T_{\alpha,\lambda}^x$  for functions on  $Y$  by

$$(2.3) \quad T_{\alpha,\lambda}^x f(y) = \frac{\Gamma(\alpha+1)2^\alpha}{\sqrt{2\pi}} \int_0^\pi f((x^2 + y^2 + 2xy \cos(\theta))^{\frac{1}{2}}) j_{\alpha-\frac{1}{2}}(|\lambda|xy \sin \theta) \sin^{2\alpha} \theta d\theta,$$

where  $j_\alpha(t) = J_\alpha(t)t^{-\alpha}$ . Here  $J_\alpha$  is the Bessel functions of order  $\alpha$ . Though the definition of Laguerre translation is quite complicated, but its action on Laguerre functions is simple enough.

**Proposition 2.2.** *We have*

$$T_{\alpha,\lambda}^y \phi_{k,\lambda}^\alpha(x) = \frac{\Gamma(\alpha+1)\Gamma(k+1)}{\Gamma(\alpha+k+1)} \phi_{k,\lambda}^\alpha(y) \phi_{k,\lambda}^\alpha(x).$$

For proof we refer to [12][Theorem 6.1.2].

Using Laguerre translation, we define *Laguerre convolution*  $f *_\lambda g$  for functions  $f, g \in L^1(Y, x^{2\alpha+1} dx)$  as

$$(2.4) \quad f *_\lambda g(x) = \int_0^\infty T_{\alpha,\lambda}^y f(x) g(y) y^{2\alpha+1} dy.$$

We quickly recall Hilbert space theory for  $L^2(Y, x^{2\alpha+1} dx)$ . Since  $\{\tilde{\phi}_{k,\lambda}^\alpha\}_{k=0}^\infty$  forms an orthonormal basis for  $L^2(Y, x^{2\alpha+1} dx)$ , we have  $f = \sum_{k=0}^\infty \langle f, \tilde{\phi}_{k,\lambda}^\alpha \rangle \tilde{\phi}_{k,\lambda}^\alpha$  in  $L^2$  norm. So, for

$k = 0, 1, 2, \dots$ , if we also define *Laguerre coefficients*  $\hat{f}(k)$  for the function  $f \in L^2(Y, x^{2\alpha+1} dx)$  by

$$(2.5) \quad \hat{f}(k) = \frac{\Gamma(\alpha+1)\Gamma(k+1)}{\Gamma(\alpha+k+1)} \int_Y f(x) \phi_{k,\lambda}^\alpha(x) x^{2\alpha+1} dx,$$

then we have

$$(2.6) \quad f = \frac{2|\lambda|^{\alpha+1}}{\Gamma(\alpha+1)} \sum_{k=0}^{\infty} \hat{f}(k) \phi_{k,\lambda}^\alpha$$

in  $L^2(Y, x^{2\alpha+1} dx)$  norm. Moreover, using Proposition 2.2 one can check that

$$(2.7) \quad f *_{\lambda} \phi_{k,\lambda}^\alpha = \hat{f}(k) \phi_{k,\lambda}^\alpha.$$

Hence we have another representation for  $f \in L^2(Y, x^{2\alpha+1} dx)$ :

**Proposition 2.3.** *For  $f \in L^2(Y, x^{2\alpha+1} dx)$ , we have*

$$f = \frac{2|\lambda|^{\alpha+1}}{\Gamma(\alpha+1)} \sum_{k=0}^{\infty} f *_{\lambda} \phi_{k,\lambda}^\alpha$$

in  $L^2$  norm.

Again using Proposition 2.2, one can check that  $\phi_{k,\lambda}^\alpha *_{\lambda} \phi_{j,\lambda}^\alpha(x) = \frac{\Gamma(\alpha+1)}{2|\lambda|^{\alpha+1}} \phi_{k,\lambda}^\alpha(x) \delta_{k,j}$ , where  $\delta_{k,j}$  is the kronecker delta function. Using this we can easily calculate Laguerre coefficients of  $f *_{\lambda} g$  for  $f, g \in L^2(Y, x^{2\alpha+1})$ .

**Proposition 2.4.** *For  $f, g \in L^2(Y, x^{2\alpha+1})$ , Laguerre coefficients of  $f *_{\lambda} g$  are related to Laguerre coefficients of  $f$  and  $g$  by*

$$(\widehat{f *_{\lambda} g})(k) = \hat{f}(k) \hat{g}(k).$$

**2.2. Laguerre transform on  $\mathbb{R}^+ \times \mathbb{R}$ .** Let  $X = \mathbb{R}^+ \times \mathbb{R}$  and  $\alpha > -1/2$ . We will denote the elements of  $X$  by Greek letters  $\xi, \eta$ , etc, with the understanding that  $\xi = (x, w)$  means  $x \in \mathbb{R}^+$  and  $w \in \mathbb{R}$ . We equip  $X$  with measure  $d\mu(x, w) = x^{2\alpha+1} dx dw$ , where  $dx$  and  $dw$  are the standard Lesbegue measures. For  $\xi = (x, w)$  and  $\eta = (y, v)$ , and  $\theta, \phi \in \mathbb{R}$ , we define product

$$(\xi, \eta)_{\theta, \phi} = ((x^2 + y^2 - 2xy \cos(\theta))^{\frac{1}{2}}, w - v + xy \cos(\phi) \sin(\theta))$$

Define measure  $\nu$  on  $[0, \pi] \times [0, \pi]$  by:

$$d\nu(\theta, \phi) = \frac{\alpha}{\pi} (\sin(\phi))^{2\alpha-1} (\sin(\theta))^{2\alpha} d\theta d\phi,$$

where  $d\theta$  and  $d\phi$  are again standard Lesbegue measures. For  $f \in L^1(X, \mu)$ , we define *generalized translation operator*  $T^\eta$  for  $\eta \in X$  by

$$(2.8) \quad T^\eta f(\xi) = \int_0^\pi \int_0^\pi f((\xi, \eta)_{(\theta, \phi)}) d\nu(\theta, \phi).$$

Using the generalized translation operator, we define convolution  $f * g$ , for  $f, g \in L^1(X, \mu)$  by

$$(2.9) \quad f * g(\xi) = \int_X T^\eta f(\xi) g(\eta) d\mu(\eta).$$

For  $\lambda \in \mathbb{R} \setminus \{0\}$  and  $k = 0, 1, 2, \dots$ , define

$$(2.10) \quad \psi_{k,\lambda}^\alpha(x, w) = \frac{\Gamma(k+1)\Gamma(\alpha+1)}{\Gamma(\alpha+k+1)} e^{i\lambda w} \phi_{k,\lambda}^\alpha(x).$$

Though the definition of generalized translation is complicated, its action on  $\psi_{k,\lambda}^\alpha$  is simple.

**Proposition 2.5.** *We have*

$$T^{(y,-v)}\psi_{k,\lambda}^\alpha(x, w) = \psi_{k,\lambda}^\alpha(y, v)\psi_{k,\lambda}^\alpha(x, w).$$

We refer to [9][Lemma 4.2] for the proof.

For  $f \in L^1(X)$ ,  $\lambda \in \mathbb{R} \setminus \{0\}$  and  $k = 0, 1, 2, \dots$ , we define its *Laguerre transform*  $\widehat{f}(\lambda, k)$  by

$$(2.11) \quad \widehat{f}(\lambda, k) = \int_X f(x, w) \psi_{k,\lambda}^\alpha(x, w) d\mu(x, w).$$

**Proposition 2.6.** *For  $f \in L^2(X)$ , we have*

$$f(x, w) = \frac{1}{\pi\Gamma(\alpha+1)} \int_{\mathbb{R}} \sum_{k=0}^{\infty} \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x) |\lambda|^{\alpha+1} e^{-i\lambda w} d\lambda,$$

in  $L^2(X)$  norm.

For proof see [10][Lemma 3.1].

For  $f \in L^1(X, \mu)$ , we define

$$(2.12) \quad f^\lambda(x) = \int_{\mathbb{R}} f(x, w) e^{i\lambda w} dw,$$

that is inverse Fourier transform in the second variable. For  $f \in L^2(X)$ , if we calculate the Laguerre coefficients of  $f^\lambda$ , we immediately notice that

$$(2.13) \quad \widehat{f^\lambda}(k) = \widehat{f}(\lambda, k).$$

Also using the Proposition 2.5, one can check that

$$(2.14) \quad \widehat{f * g} = \widehat{f} \widehat{g}.$$

**2.3. Fractional powers of generalized sublaplacian.** For  $\alpha > -1/2$ , we define *generalized sublaplacian*  $\mathcal{L}$  on  $X$  by

$$(2.15) \quad -\frac{1}{2} \left( \frac{\partial^2}{\partial x^2} + \frac{2\alpha+1}{x} \frac{\partial}{\partial x} + x^2 \frac{\partial^2}{\partial w^2} \right).$$

This operator is positive and symmetric in  $L^2(X)$ . Using the fact that Laguerre polynomials satisfy the following identity

$$(2.16) \quad x \frac{d^2}{dx^2} L_k^\alpha(x) + (\alpha+1-x) \frac{d}{dx} L_k^\alpha(x) + k L_k^\alpha(x) = 0,$$

one can check that

$$(2.17) \quad \mathcal{L}\psi_{k,\lambda}^\alpha = |\lambda|(2k+\alpha+1)\psi_{k,\lambda}^\alpha.$$

Thus  $\psi_{k,\lambda}^\alpha$  are the eigenvectors for  $\mathcal{L}$  with  $|\lambda|(2k + \alpha + 1)$  as corresponding eigenvalues. Moreover, for  $f \in L^2(X)$  such that  $\mathcal{L}f \in L^2(X)$ , we have

$$(2.18) \quad \widehat{\mathcal{L}f}(\lambda, k) = |\lambda|(2k + \alpha + 1)\widehat{f}(\lambda, k).$$

Using Proposition 2.6 and (2.17), we obtain the following spectral decomposition of  $\mathcal{L}$ :

$$(2.19) \quad \mathcal{L}f(x, w) = \frac{1}{\pi\Gamma(\alpha + 1)} \int_{\mathbb{R}} \left( \sum_{k=0}^{\infty} |\lambda|(2k + \alpha + 1) \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x) \right) |\lambda|^{\alpha+1} e^{-i\lambda w} d\lambda.$$

Therefore, using spectral decomposition, we define fractional powers of the generalized sublaplacian  $\mathcal{L}^s$  for  $0 < s < 1$ :

$$(2.20) \quad \mathcal{L}^s f(x, w) = \frac{1}{\pi\Gamma(\alpha + 1)} \int_{\mathbb{R}} \left( \sum_{k=0}^{\infty} (|\lambda|(2k + \alpha + 1))^s \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x) \right) |\lambda|^{\alpha+1} e^{-i\lambda w} d\lambda.$$

However, it is convenient to work with the following modified fractional power of  $\mathcal{L}$ . For  $0 < s < 1$ , we define  $\mathcal{L}_s$  by

$$(2.21) \quad \mathcal{L}_s f(x, w) = \frac{1}{\pi\Gamma(\alpha + 1)} \int_{\mathbb{R}} \left( \sum_{k=0}^{\infty} (2|\lambda|)^s \frac{\Gamma(\frac{2k+\alpha+1}{2} + \frac{1+s}{2})}{\Gamma(\frac{2k+\alpha+1}{2} + \frac{1-s}{2})} \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x) \right) |\lambda|^{\alpha+1} e^{-i\lambda w} d\lambda.$$

Thus  $\mathcal{L}_s$  corresponds to the spectral multiplier

$$(2.22) \quad (2|\lambda|)^s \frac{\Gamma(\frac{2k+\alpha+1}{2} + \frac{1+s}{2})}{\Gamma(\frac{2k+\alpha+1}{2} + \frac{1-s}{2})}.$$

Finally, we define  $W^{s,2}(X)$  as the space consisting of  $f \in L^2(X)$  such that  $\mathcal{L}_s f \in L^2(X)$ .

**2.4. Heat semigroup associated with  $\mathcal{L}$ .** The *heat semigroup*  $e^{-t\mathcal{L}}$  generated by  $\mathcal{L}$  is defined by the relation

$$(2.23) \quad \widehat{e^{-t\mathcal{L}}f}(\lambda, k) = e^{-|\lambda|(2k+\alpha+1)t} \widehat{f}(\lambda, k).$$

Thus we have

$$(2.24) \quad e^{-t\mathcal{L}}f(x, w) = f * h_t(x, w),$$

where  $h_t$  is the *heat kernel* associated with  $\mathcal{L}$  given by

$$(2.25) \quad \widehat{h}_t(\lambda, k) = e^{-|\lambda|(2k+\alpha+1)t}.$$

Although the expression for  $h_t(x, w)$  is not known explicitly, we have the explicit expression for  $h_t^\lambda(x)$ .

**Proposition 2.7.** *We have*

$$h_t^\lambda(x) = \frac{2}{\Gamma(\alpha + 1)} \left( \frac{\lambda}{2 \sinh(\lambda t)} \right)^{\alpha+1} e^{-\frac{\lambda}{2} x^2 \coth(\lambda t)}.$$

*Proof.* Using Proposition 2.6, we have

$$\begin{aligned} h_t(x, w) &= \frac{1}{\pi\Gamma(\alpha + 1)} \sum_{k=0}^{\infty} \int_{-\infty}^{\infty} e^{-|\lambda|(2k+\alpha+1)t} \phi_{k,\lambda}^\alpha(x) |\lambda|^{\alpha+1} e^{-i\lambda w} d\lambda \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} h_t^\lambda(x) e^{-i\lambda w} d\lambda, \end{aligned}$$

where

$$h_t^\lambda(x) = \frac{2}{\Gamma(\alpha + 1)} \sum_{k=0}^{\infty} e^{-|\lambda|(2k+\alpha+1)t} \phi_{k,\lambda}^\alpha(x) |\lambda|^{\alpha+1}.$$

Using the generating function identity for Laguerre functions

$$\sum_{k=0}^{\infty} L_k^\alpha(x) r^k = (1-r)^{-(\alpha+1)} e^{-\frac{rx}{1-r}},$$

we simplify to get the desired expression for  $h_t^\lambda$ . □

For  $0 < s < 1$  and  $t > 0$ , we define  $K_{t,s}(x, w)$  by

$$(2.26) \quad K_{t,s}^\lambda(x) = h_t^\lambda(x) \left( \frac{\lambda t}{\sinh \lambda t} \right)^{s+1}.$$

**Lemma 2.8.** *We have the following properties of  $K_{t,s}$*

$$(2.27) \quad \int_X K_{t,s}(x, w) d\mu(x, w) = 1.$$

$$(2.28) \quad \int_X T^\eta K_{t,s}(\xi) d\mu(\xi) = \int_X T^\eta K_{t,s}(\xi) d\mu(\eta) = 1.$$

$$(2.29) \quad f * K_{t,s}(\xi) = K_{t,s} * f(\xi).$$

*Proof.* We begin with the definition

$$\int_{-\infty}^{\infty} K_{t,s}(x, w) e^{i\lambda w} dw = h_t^\lambda(x) \left( \frac{\lambda t}{\sinh \lambda t} \right)^{s+1}.$$

Making  $\lambda$  go to 0, we get

$$\int_{-\infty}^{\infty} K_{t,s}(x, w) dw = \frac{2}{\Gamma(\alpha + 1)} \frac{1}{(2t)^{\alpha+1}} e^{-\frac{x^2}{2t}}.$$

Therefore,

$$\begin{aligned} \int_X K_{t,s}(x, w) d\mu(x, w) &= \int_0^\infty \frac{2}{\Gamma(\alpha + 1)} \frac{1}{(2t)^{\alpha+1}} e^{-\frac{x^2}{2t}} x^{2\alpha+1} dx \\ &= \int_0^\infty \frac{2}{\Gamma(\alpha + 1)} e^{-x^2} x^{2\alpha+1} dx \\ &= 1. \end{aligned}$$

Next, using [9][Lemma 3.1], we have

$$\int_X T^\eta f(\xi) g(\xi) d\mu(\xi) = \int_X f(\xi) T^{\eta*} g(\xi) d\mu(\xi),$$

where  $(x, w)^* = (x, -w)$ . Take  $g = 1$  and  $f = K_{t,s}$  and use (2.27) to conclude

$$\int_X T^\eta K_{t,s}(\xi) d\mu(\xi) = 1.$$

On the other hand,

$$\int_X T^\eta K_{t,s}(\xi) d\mu(\xi) = \int_X T^{\xi*} K_{t,s}(\eta^*) d\mu(\xi)$$



is just a change of variables; combining with the fact that  $K_{t,s}$  is an even function in the second variable, we conclude second half of the (2.28).

Finally, (2.29) follows from the change of variable and the fact that  $K_{t,s}$  is an even function in the second variable.  $\square$

For  $0 < s < 1$ , we define

$$(2.30) \quad K_s(x, w) = \int_0^\infty K_{t,s}(x, w) t^{-s-1} dt.$$

We will show  $K_s$  is a positive function, more precisely:

**Proposition 2.9.** *For  $0 < s < 1$ , we have*

$$K_s(x, w) = \frac{2^{2\alpha+2s+3} \left( \Gamma\left(\frac{\alpha+s+2}{2}\right) \right)^2}{\pi \Gamma(\alpha+1)} \frac{1}{(x^4 + 4w^2)^{\frac{s+\alpha+2}{2}}}.$$

*Proof.* Calculations are borrowed from [7][Proposition 4.2]. We repeat for the sake of completeness. We start with the expression

$$\int_{-\infty}^\infty K_s(x, w) e^{i\lambda w} dw = \int_0^\infty h_t^\lambda(x) \left( \frac{t|\lambda|}{\sinh t|\lambda|} \right)^{s+1} t^{-s-1} dt.$$

Using Proposition 2.7, and since the functions involved are even in  $\lambda$ , we have

$$\int_{-\infty}^\infty K_s(x, w) e^{i\lambda w} dw = \frac{2}{\Gamma(\alpha+1)} \int_0^\infty \left( \frac{\lambda}{\sinh t\lambda} \right)^{\alpha+s+2} e^{-\frac{1}{2}\lambda(\coth t\lambda)x^2} dt.$$

As the Fourier transform of  $K_s$  in the central variable  $w$  is an even function of  $\lambda$  we have, after taking the Fourier transform in the variable  $\lambda$ ,

$$K_s(x, w) = \frac{2}{\pi \Gamma(\alpha+1)} \int_0^\infty \int_0^\infty (\cos \lambda w) \left( \frac{\lambda}{\sinh t\lambda} \right)^{\alpha+s+2} e^{-\frac{1}{2}\lambda(\coth t\lambda)x^2} d\lambda dt.$$

By the change of variables  $\lambda \rightarrow \lambda x^{-2}$ ,  $t \rightarrow tx^2$ , we obtain

$$(2.31) \quad K_s(x, wx^2) = x^{-2(\alpha+s+2)} K_s(1, w).$$

Thus

$$\begin{aligned} K_s(1, w) &= \frac{2}{\pi \Gamma(\alpha+1)} \int_0^\infty \int_0^\infty (\cos \lambda w) \left( \frac{\lambda}{\sinh t\lambda} \right)^{\alpha+s+2} e^{-\frac{\lambda}{2}(\coth t\lambda)} dt d\lambda \\ &= \frac{2}{\pi \Gamma(\alpha+1)} \int_0^\infty \left( \int_0^\infty (\cos \lambda w) \lambda^{\alpha+s+1} e^{-\frac{\lambda}{2}(\coth t\lambda)} d\lambda \right) (\sinh t)^{-(\alpha+s+2)} dt. \end{aligned}$$

The integral in  $\lambda$  can be evaluated by using [5, p. 498, 3.944.6]:

$$(2.32) \quad \int_0^\infty x^{\mu-1} e^{-\beta x} (\cos \delta x) dx = \frac{\Gamma(\mu)}{(\delta^2 + \beta^2)^{\mu/2}} \cos\left(\mu \arctan \frac{\delta}{\beta}\right),$$

valid for  $\operatorname{Re} \mu > 0$ ,  $\operatorname{Re} \beta > |\operatorname{Im} \delta|$ . Taking with  $\mu = \alpha + s + 2$ ,  $\beta = \frac{1}{2}(\coth t)$  and  $\delta = w$ . Then, we get

$$\int_0^\infty (\cos \lambda w) \lambda^{\alpha+s+1} e^{-\frac{\lambda}{2}(\coth t\lambda)} d\lambda = \frac{\Gamma(\alpha+s+2) \cos\left((\alpha+s+2) \arctan\left(\frac{2w}{\coth t}\right)\right)}{\left(w^2 + \frac{1}{4} \coth^2 t\right)^{\frac{\alpha+s+2}{2}}}.$$

Thus

$$(2.33) \quad K_s(1, w) = \frac{2\Gamma(\alpha + s + 2)}{\pi\Gamma(\alpha + 1)} \int_0^\infty \frac{\cos\left((\alpha + s + 2) \arctan\left(\frac{2w}{\coth t}\right)\right)}{\left(w^2 + \frac{1}{4} \coth^2 t\right)^{\frac{\alpha+s+2}{2}}} (\sinh t)^{-(\alpha+s+2)} dt.$$

With the change of variables  $u = \frac{2w}{\coth t}$  we have that the latter integral equals

$$\begin{aligned} & \int_0^{2w} \left(\frac{u^2}{4w^2 - u^2}\right)^{-\frac{(\alpha+s+2)}{2}} \left(w^2 + \frac{4w^2}{4u^2}\right)^{-\frac{(\alpha+s+2)}{2}} \cos[(\alpha + s + 2) \arctan u] \frac{2w}{4w^2 - u^2} du \\ &= 2w^{-(\alpha+s+1)} \int_0^{2w} (4w^2 - u^2)^{\frac{\alpha+s}{2}} (1 + u^2)^{-\frac{\alpha+s+2}{2}} \cos[(\alpha + s + 2) \arctan u] du \\ &= 2^{\alpha+s+1} w^{-1} \int_0^{2w} \left(1 - \frac{u^2}{4w^2}\right)^{\frac{\alpha+s}{2}} (1 + u^2)^{-\frac{\alpha+s+2}{2}} \cos[(\alpha + s + 2) \arctan u] du. \end{aligned}$$

Thus, with this and (2.33) we have

$$(2.34) \quad \mathcal{K}_s(1, w) = \frac{2^{\alpha+s+2}\Gamma(\alpha + s + 2)}{\pi\Gamma(\alpha + 1)} w^{-1} I,$$

where

$$I := \int_0^{2w} \left(1 - \frac{u^2}{4w^2}\right)^{\frac{\alpha+s}{2}} (1 + u^2)^{-\frac{\alpha+s+2}{2}} \cos[(\alpha + s + 2) \arctan u] du.$$

Now we will see that the above integral can be explicitly computed in terms of Legendre functions. Making a second change of variable  $\arctan u = z$ , the integral  $I$  becomes

$$I = \int_0^{\arctan 2w} \left(\cos^2 z - \frac{\sin^2 z}{4w^2}\right)^{\frac{\alpha+s}{2}} \cos[(\alpha + s + 2)z] dz.$$

We can rewrite the above integral as

$$\begin{aligned} I &= \int_0^{\arctan 2w} \left(\frac{1 + \cos 2z}{2} - \frac{1 - \cos 2z}{2 \cdot 4w^2}\right)^{\frac{\alpha+s}{2}} \cos[(\alpha + s + 2)z] dz \\ &= 2^{-\frac{\alpha+s}{2}} \int_0^{\arctan 2w} \left((\cos 2z) \left(1 + \frac{1}{4w^2}\right) - \left(\frac{1}{4w^2} - 1\right)\right)^{\frac{\alpha+s}{2}} \cos[(\alpha + s + 2)z] dz \\ &= \left(\frac{1 + 4w^2}{8w^2}\right)^{\frac{\alpha+s}{2}} \int_0^{\arctan 2w} \left(\cos 2z - \frac{1 - 4w^2}{1 + 4w^2}\right)^{\frac{\alpha+s}{2}} \cos[(\alpha + s + 2)z] dz \\ &= \frac{1}{2} \left(\frac{1 + 4w^2}{8w^2}\right)^{\frac{\alpha+s}{2}} \int_0^{2 \arctan 2w} (\cos \beta - \cos \gamma)^{\frac{\alpha+s}{2}} \cos\left[\frac{(\alpha + s + 2)}{2} \beta\right] d\beta, \end{aligned}$$

where  $\cos \gamma = \frac{1-16w^2}{1+16w^2}$ . The integral can be evaluated using ([5, p. 406, 3.663.1]):

$$(2.35) \quad \int_0^u (\cos x - \cos u)^{\nu-\frac{1}{2}} \cos ax dx = \sqrt{\frac{\pi}{2}} (\sin u)^\nu \Gamma\left(\nu + \frac{1}{2}\right) P_{a-\frac{1}{2}}^{-\nu}(\cos u),$$

valid for  $\operatorname{Re} \nu > -\frac{1}{2}$ ,  $a > 0$ ,  $0 < u < \pi$ , where  $P_{a-\frac{1}{2}}^{-\nu}$  is an associated Legendre function of the first kind (see for instance [5, Sections 8.7-8.8]). Also Recall the following representation for the associated Legendre function ([5, p. 969, 8.755])

$$(2.36) \quad P_\nu^{-\nu}(\cos \varphi) = \frac{\left(\frac{\sin \varphi}{2}\right)^\nu}{\Gamma(1 + \nu)}.$$

Taking  $\nu = \frac{\alpha+s+1}{2}$  and  $a = \frac{\alpha+s+2}{2}$  in (2.35) and using the representation for the associated Legendre function (2.36), the latter integral becomes

$$\begin{aligned} \sqrt{\frac{\pi}{2}} (\sin \gamma)^{\frac{\alpha+s+1}{2}} \Gamma\left(\frac{\alpha+s+2}{2}\right) P_{\frac{\alpha+s+1}{2}}^{-\frac{\alpha+s+1}{2}}(\cos \gamma) &= \sqrt{\frac{\pi}{2}} \Gamma\left(\frac{\alpha+s+2}{2}\right) (\sin \gamma)^{\frac{\alpha+s+1}{2}} \frac{(\sin \gamma)^{\frac{\alpha+s+1}{2}}}{2^{\frac{\alpha+s+1}{2}} \Gamma\left(\frac{\alpha+s+3}{2}\right)} \\ &= \sqrt{\frac{\pi}{2}} \frac{\Gamma\left(\frac{\alpha+s+2}{2}\right)}{2^{\frac{\alpha+s+1}{2}} \Gamma\left(\frac{\alpha+s+3}{2}\right)} (\sin^2 \gamma)^{\frac{\alpha+s+1}{2}} \\ &= \sqrt{\frac{\pi}{2}} \frac{\Gamma\left(\frac{\alpha+s+2}{2}\right)}{2^{\frac{\alpha+s+1}{2}} \Gamma\left(\frac{\alpha+s+3}{2}\right)} \left(\frac{4w}{1+4w^2}\right)^{\alpha+s+1}, \end{aligned}$$

because  $\sin^2 \gamma = \frac{16w^2}{(1+4w^2)^2}$ . This gives

$$(2.37) \quad I = \frac{1}{2} \sqrt{\frac{\pi}{2}} \frac{\Gamma\left(\frac{\alpha+s+2}{2}\right)}{2^{\frac{\alpha+s+1}{2}} \Gamma\left(\frac{\alpha+s+3}{2}\right)} \left(\frac{1+4w^2}{8w^2}\right)^{\frac{\alpha+s}{2}} \left(\frac{4w}{1+4w^2}\right)^{\alpha+s+1} = \sqrt{\pi} \frac{\Gamma\left(\frac{\alpha+s+2}{2}\right)}{\Gamma\left(\frac{\alpha+s+3}{2}\right)} w (1+4w^2)^{-\frac{\alpha+s+2}{2}}.$$

Finally, plugging (2.37) into (2.34), we have

$$K_s(1, w) = \frac{2^{\alpha+s+2} \Gamma(\alpha+s+2)}{\pi \Gamma(\alpha+1)} \sqrt{\pi} \frac{\Gamma\left(\frac{\alpha+s+2}{2}\right)}{\Gamma\left(\frac{\alpha+s+3}{2}\right)} (1+4w^2)^{-\frac{\alpha+s+2}{2}},$$

or, by (2.31)

$$K_s(x, w) = x^{-2(\alpha+s+2)} K_s\left(1, \frac{w}{x^2}\right) = c_{\alpha,s} (x^4 + 4w^2)^{-\frac{\alpha+s+2}{2}}.$$

where the constant  $c_{\alpha,s}$  is given by

$$c_{\alpha,s} = \frac{2^{\alpha+s+2} \Gamma(\alpha+s+2)}{\sqrt{\pi} \Gamma(\alpha+1)} \frac{\Gamma\left(\frac{\alpha+s+2}{2}\right)}{\Gamma\left(\frac{\alpha+s+3}{2}\right)}.$$

By using Legendre's duplication formula

$$(2.38) \quad \sqrt{\pi} \Gamma(2z) = 2^{2z-1} \Gamma(z) \Gamma\left(z + \frac{1}{2}\right)$$

with  $z = \frac{\alpha+s+2}{2}$ , and after simplification, we get

$$c_{\alpha,s} = \frac{2^{2\alpha+2s+3} \left(\Gamma\left(\frac{\alpha+s+2}{2}\right)\right)^2}{\pi \Gamma(\alpha+1)}.$$

This completes the proof of the Proposition.  $\square$

We have shown that  $K_s$  is a positive function. Moreover, generalized translation is a positive operator, see [9][Proposition 3.2]. Therefore,  $T^\eta K_s \geq 0$ . In other words,

$$(2.39) \quad \int_X T^\eta K_{t,s}(\xi) t^{-s-1} dt \geq 0, \quad \forall \xi, \eta \in X.$$

**2.5. Fractional powers of Grushin operator.** Let  $H = \mathbb{R}^n \times \mathbb{R}$  with the understanding that  $(x, w) \in H$  means  $x \in \mathbb{R}^n$  and  $w \in \mathbb{R}$ . We equip  $H$  with the measure  $d\mu(x, w) = dx dw$ , where  $dx$  and  $dw$  are the usual Lebesgue measures on  $\mathbb{R}^n$  and  $\mathbb{R}$ . We define *Grushin operator*  $\mathcal{G}$  on  $H$  by

$$(2.40) \quad \mathcal{G} = -\frac{1}{2} \left( \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2} + |x|^2 \frac{\partial^2}{\partial w^2} \right).$$

For  $\lambda \in \mathbb{R} \setminus \{0\}$ , we define the *scaled Hermite operator*  $\mathcal{H}(\lambda)$  on  $\mathbb{R}^n$  by

$$(2.41) \quad \mathcal{H}(\lambda) = - \left( \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2} - \lambda^2 |x|^2 \right)$$

For multi-index  $\beta \in \mathbb{N}^n$ , define  $\Phi_\beta(x) = h_{\beta_1}(x_1)h_{\beta_2}(x_2) \cdots h_{\beta_n}(x_n)$ , where  $\beta = (\beta_1, \dots, \beta_n)$  and  $h_{\beta_i}$  are normalized Hermite functions. Further, for  $\lambda \in \mathbb{R} \setminus \{0\}$  define

$$\Phi_\beta^\lambda(x) = |\lambda|^{\frac{n}{4}} \Phi_\beta(\sqrt{|\lambda|}x).$$

The collection  $\{\Phi_\beta^\lambda\}_{\beta \in \mathbb{N}^n}$  forms an orthonormal basis for  $L^2(\mathbb{R}^n)$ , see [11][Theorem 1.2.2]. Also they are eigenfunctions for the scaled Hermite operator, that is

$$(2.42) \quad \mathcal{H}(\lambda)\Phi_\beta^\lambda = (2|\beta| + n)|\lambda|\Phi_\beta^\lambda.$$

For  $\lambda \in \mathbb{R} \setminus \{0\}$  and  $k = 0, 1, 2, \dots$ , define  $\mathcal{P}_k(\lambda)$  as projections of  $L^2(\mathbb{R}^n)$  onto  $E_k^\lambda$ , the eigenspace corresponding to eigenvalue  $(2k + n)|\lambda|$ . In other words,

$$(2.43) \quad \mathcal{P}_k(\lambda)f = \sum_{|\beta|=k} \langle f, \Phi_\beta^\lambda \rangle \Phi_\beta^\lambda.$$

Thus we have

$$(2.44) \quad \mathcal{H}(\lambda) = \sum_{k=0}^{\infty} (2k + n)|\lambda|\mathcal{P}_k(\lambda).$$

Finally, using the Fourier transform and the above spectral decomposition of scaled Hermite operator (Equation (2.44)), we have the following spectral decomposition of  $\mathcal{G}$ :

$$(2.45) \quad \mathcal{G}f(x, w) = \frac{1}{2\pi} \int_{\mathbb{R}} \sum_{k=0}^{\infty} (k + n/2)|\lambda|\mathcal{P}_k(\lambda)f^\lambda(x)e^{-i\lambda w} d\lambda.$$

Therefore, a natural way to define fractional powers of Grushin operator is via spectral decomposition:

$$(2.46) \quad \mathcal{G}^s f(x, w) = \frac{1}{2\pi} \int_{\mathbb{R}} \sum_{k=0}^{\infty} ((k + n/2)|\lambda|)^s \mathcal{P}_k(\lambda)f^\lambda(x)e^{-i\lambda w} d\lambda.$$

However, it is convenient to with following modified fractional powers of  $\mathcal{G}$ . For  $0 < s < 1$ , we define  $\mathcal{G}_s$  by

$$(2.47) \quad \mathcal{G}_s f(x, w) = \frac{1}{2\pi} \int_{\mathbb{R}} \sum_{k=0}^{\infty} (2|\lambda|)^s \frac{\Gamma(\frac{2k+n}{4} + \frac{1+s}{2})}{\Gamma(\frac{2k+n}{4} + \frac{1-s}{2})} \mathcal{P}_k(\lambda)f^\lambda(x)e^{-i\lambda w} d\lambda.$$

Also we define  $W^{s,2}(H)$  as the space consists of all those functions  $f$  in  $L^2(H)$  such that  $\mathcal{G}_s f \in L^2(H)$  too.

**2.6. Spherical harmonics and Hecke-Bochner formula.** Let us quickly recall some facts about spherical harmonics and solid harmonics. We refer to [8][Chapter 4] for missing details. Let  $\mathfrak{H}_m$  denote the space of spherical harmonics of degree  $m$ . Let  $\{Y_{m,j}\}_{j=1}^{a_m}$  denotes the orthogonal basis of  $\mathfrak{H}_m$ , where  $a_m$  denotes the dimension of  $\mathfrak{H}_m$ . We know that  $L^2(\mathbb{S}^{n-1}) = \bigoplus_{m=0}^{\infty} \mathfrak{H}_m$  and the collection  $\{Y_{m,j}\}$ , for  $j = 1, 2, \dots, a_m$  and  $m = 0, 1, 2, \dots$ , forms orthonormal basis for  $L^2(\mathbb{S}^{n-1})$ . Note that  $\mathbb{S}^{n-1}$  denotes the unit sphere in  $\mathbb{R}^n$ . Define solid harmonics  $P_{m,j}(x) = |x|^m Y_{m,j}(x/|x|)$  for  $j = 1, 2, \dots, a_m$  and  $m = 0, 1, \dots$ . Let  $\mathfrak{h}_m$  denotes the space consisting of linear combination of functions of the form  $f(|x|)P(x)$ , where  $f$  varies over radial functions and  $P \in \mathfrak{H}_m$ , with the stipulation that each  $f(|x|)P(x) \in L^2(\mathbb{R}^n)$ . With these definitions, we have  $L^2(\mathbb{R}^n) = \bigoplus_{m=0}^{\infty} \mathfrak{h}_m$ . So for  $f \in L^2(\mathbb{R}^{n+1})$ , we have

$$(2.48) \quad f(x, w) = \sum_{m=0}^{\infty} \sum_{j=1}^{a_m} f_{m,j}(|x|, w) P_{m,j}(x),$$

where  $f_{m,j}(|x|, w) = \int_{\mathbb{S}^{n-1}} f(|x|\omega, w) P_{m,j}(|x|\omega) d\omega$ .

Finally, we recall Hecke-Bochner formula, which describes how Hermite projections act on solid harmonics, proof of which can be found in the book [12][Theorem 3.4.1].

**Proposition 2.10.** *Suppose  $f \in L^2(\mathbb{R}^n)$  is such that  $f = gP$ , where  $g$  is radial and  $P$  is a solid harmonics of degree  $m$ . Then we have*

$$\mathcal{P}_{2k+m}(\lambda)f(x) = R_{k,m}^{\lambda}(g)\phi_{k,\lambda}^{\alpha}(|x|)P(x)$$

where

$$R_{k,m}^{\lambda}(g) = \frac{2|\lambda|^{\alpha+1}\Gamma(k+1)}{\Gamma(\alpha+k+1)} \int_0^{\infty} g(s)\phi_{k,\lambda}^{\alpha}(s)s^{2\alpha+1} ds$$

and  $\alpha = \frac{n}{2} + m - 1$ . For other values of  $j$ ,  $\mathcal{P}_j(\lambda)f = 0$ .

### 3. HARDY'S INEQUALITY FOR GENERALIZED SUBAPLACIAN

For  $-1 < s < 1$  and  $\delta > 0$ , we define

$$(3.1) \quad u_{s,\delta}(x, w) = \left( \left( \delta + \frac{x^2}{2} \right)^2 + w^2 \right)^{-\frac{s+\alpha+2}{2}}.$$

**Proposition 3.1.** *For  $0 < s < 1$ , we have*

$$\mathcal{L}_s u_{-s,\delta}(\xi) = (4\delta)^s \left( \frac{\Gamma(\frac{\alpha+s+2}{2})}{\Gamma(\frac{\alpha-s+2}{2})} \right)^2 u_{s,\delta}(\xi).$$

*Proof.* We prove the result by calculating Laguerre transform on both sides. This has been already done in Ciaurri et al [2], but we repeat the calculations for the convenience of the readers. Define

$$L(a, b, c) = \int_0^{\infty} e^{-a(2x+1)} x^{b-1} (1+x)^{-c} dx.$$

We start with the generating function identity for the Laguerre functions:

$$\sum_{k=0}^{\infty} z^k L_k^{\alpha}(x^2) e^{-\frac{1}{2}x^2} = (1-z)^{-\alpha-1} e^{-\frac{1}{2}\frac{1+z}{1-z}x^2}.$$

Therefore, we have

$$(3.2) \quad \sum_{k=0}^{\infty} \left( \frac{y}{y+|\lambda|} \right)^k L_k^\alpha(|\lambda|x^2) e^{-\frac{1}{2}|\lambda|x^2} = |\lambda|^{-\alpha-1} (y+|\lambda|)^{\alpha+1} e^{-\frac{1}{2}(2y+|\lambda|x^2)}.$$

For functions  $f, g$  defined on  $(0, \infty)$ , let  $F, G$  be their Laplace transforms defined by

$$F(a+ib) = \int_0^\infty e^{-(a+ib)y} f(y) dy, \quad G(a, ib) = \int_0^\infty e^{-(a+ib)y} g(y) dy, \quad a > 0, b \in \mathbb{R}$$

Let  $\beta = \frac{1}{2}(\alpha + s + 2)$ . Then with  $f(y) = g(y) = \Gamma^{-1}(\beta) y^{\beta-1} e^{-\delta y}$ , we have

$$F(a+ib) = G(a+ib) = (\delta + a + ib)^{-\beta}.$$

On the other hand, it can be checked, see [[3], Lemma 3.4], that

$$\int_{-\infty}^{\infty} F(a+ib) \overline{G(a+ib)} e^{-i|\lambda|b} db = 2\pi \int_0^\infty f(y)g(y+|\lambda|) e^{-a(2y+|\lambda|)} dy.$$

Taking  $a = \frac{1}{2}x^2$ , we have

$$\int_{-\infty}^{\infty} \left( \left( \delta + \frac{1}{2}x^2 \right)^2 + b^2 \right)^{-\frac{1}{2}(s+\alpha+2)} e^{-i|\lambda|b} db = 2\pi \int_0^\infty f(y)g(y+|\lambda|) e^{-\frac{1}{2}(2y+|\lambda|x^2)} dy.$$

Since  $u_{s,\delta}(x, w)$  is symmetric in  $w$  variable, therefore we have

$$u_{s,\delta}^\lambda(x) = 2\pi \int_0^\infty f(y)g(y+|\lambda|) e^{-\frac{1}{2}(2y+|\lambda|x^2)} dy.$$

Using (3.2), we have

$$u_{s,\delta}^\lambda(x) = \frac{2|\lambda|^{\alpha+1}}{\Gamma(\alpha+1)} \sum_{k=0}^{\infty} c_{k,\delta}^\lambda(s) \phi_{k,\lambda}^\alpha(x),$$

where the coefficients are given by

$$\begin{aligned} c_{k,\delta}^\lambda(s) &= \pi \Gamma(\alpha+1) \int_0^\infty f(y)g(y+|\lambda|) (y+|\lambda|)^{-(k+\alpha+1)} y^k dy \\ &= \frac{\pi \Gamma(\alpha+1) |\lambda|^s}{(\Gamma(\beta))^2} \int_0^\infty e^{-\delta(2y+|\lambda|)} y^{\beta+k-1} (y+|\lambda|)^{\beta-k-\alpha-2} dy \\ &= \frac{\pi \Gamma(\alpha+1) |\lambda|^s}{(\Gamma(\beta))^2} L \left( \delta|\lambda|, \frac{2k+\alpha+2+s}{2}, \frac{2k+\alpha+2-s}{2} \right). \end{aligned}$$

Notice that  $\widehat{u_{s,\delta}}(\lambda, k) = \widehat{u_{s,\delta}^\lambda}(k) = c_{k,\delta}^\lambda(s)$ . Also, according to [[3], Proposition 3.6] the function  $L$  satisfies the following identity

$$\frac{(2\lambda)^a}{\Gamma(a)} L(\lambda, a, b) = \frac{(2\lambda)^b}{\Gamma(b)} L(\lambda, b, a)$$

for all  $(a, b \in \mathbb{C})$  and  $\lambda > 0$ . Using this we get

$$(3.3) \quad c_{k,\delta}^\lambda(-s) = (2\delta)^s |\lambda|^{-s} \left( \frac{\Gamma(\frac{\alpha+s+2}{2})}{\Gamma(\frac{\alpha-s+2}{2})} \right)^2 \frac{\Gamma(\frac{2k+\alpha+1}{2} + \frac{1-s}{2})}{\Gamma(\frac{2k+\alpha+1}{2} + \frac{1+s}{2})} c_{k,\delta}^\lambda(s).$$

Rearranging the terms,

$$(2|\lambda|)^s \frac{\Gamma\left(\frac{2k+\alpha+1}{2} + \frac{1+s}{2}\right)}{\Gamma\left(\frac{2k+\alpha+1}{2} + \frac{1-s}{2}\right)} c_{k,\delta}^\lambda(-s) = (4\delta)^s \left(\frac{\Gamma\left(\frac{\alpha+s+2}{2}\right)}{\Gamma\left(\frac{\alpha-s+2}{2}\right)}\right)^2 c_{k,\delta}^\lambda(s),$$

which is nothing but

$$\widehat{\mathcal{L}_s u_{-s,\delta}}(\lambda, k) = (4\delta)^s \left(\frac{\Gamma\left(\frac{\alpha+s+2}{2}\right)}{\Gamma\left(\frac{\alpha-s+2}{2}\right)}\right)^2 \widehat{u_{s,\delta}}(\lambda, k),$$

and hence the result.  $\square$

Next, we find an integral representation for  $\mathcal{L}_s$  in an analogous way as it has been found for the fractional powers of sublaplacian on Heisenberg group by Roncal et al [7][Section 4].

**Theorem 3.2.** *For  $0 < s < 1$  and  $f \in W^{s,2}(X)$ , we have*

$$\mathcal{L}_s f(\xi) = \frac{1}{|\Gamma(-s)|} \int_0^\infty (f(\xi) - f * K_{t,s}(\xi)) t^{-s-1} dt,$$

where  $K_{t,s}$  is defined in (2.26).

*Proof.* We begin with the identity (see [5]. p.382, 3.541.1)

$$2^{1-s} \int_0^\infty e^{-(\mu+1)t} (\sinh t)^{-s} dt = \frac{\Gamma(1-s)\Gamma\left(\frac{\mu}{2} + \frac{1+s}{2}\right)}{\Gamma\left(\frac{\mu}{2} + \frac{1-s}{2} + 1\right)},$$

which gives

$$(3.4) \quad (\mu+1-s) \int_0^\infty e^{-(\mu+1)t} (\sinh t)^{-s} dt = \frac{2^s \Gamma(1-s)\Gamma\left(\frac{\mu}{2} + \frac{1+s}{2}\right)}{\Gamma\left(\frac{\mu}{2} + \frac{1-s}{2}\right)}.$$

Also we have

$$\begin{aligned} (\mu+1) \int_0^\infty e^{-(\mu+1)t} (\sinh t)^{-s} dt &= \int_0^\infty \frac{d}{dt} (1 - e^{-(\mu+1)t}) (\sinh t)^{-s} dt \\ &= s \int_0^\infty (1 - e^{-(\mu+1)t}) (\sinh t)^{-s-1} (\cosh t) dt. \end{aligned}$$

Therefore, plugging the latter into (3.4), we get

$$\begin{aligned} \frac{2^s \Gamma(1-s)\Gamma\left(\frac{\mu}{2} + \frac{1+s}{2}\right)}{\Gamma\left(\frac{\mu}{2} + \frac{1-s}{2}\right)} &= s \int_0^\infty \left( \cosh t - e^{-(\mu+1)t} (\cosh t + \sinh t) \right) (\sinh t)^{-s-1} dt \\ &= s \int_0^\infty (\cosh t - e^{-\mu t}) (\sinh t)^{-s-1} dt \\ &= s \int_0^\infty (\cosh t - 1) (\sinh t)^{-s-1} dt + s \int_0^\infty (1 - e^{-\mu t}) (\sinh t)^{-s-1} dt \\ &= c_1 s + s \int_0^\infty (1 - e^{-\mu t}) (\sinh t)^{-s-1} dt, \end{aligned}$$

where  $c_1$  is the constant given by

$$c_1 := \int_0^\infty (\cosh t - 1)(\sinh t)^{-s-1} dt.$$

Thus, by taking  $\mu = 2k + \alpha + 1$  and changing  $t$  into  $|\lambda|t$ , we have

$$\frac{2^s \Gamma(1-s)}{s} \frac{\Gamma\left(\frac{2k+\alpha+1}{2} + \frac{1+s}{2}\right)}{\Gamma\left(\frac{2k+\alpha+1}{2} + \frac{1-s}{2}\right)} = c_1 + |\lambda| \int_0^\infty (1 - e^{-(2k+\alpha+1)|\lambda|t})(\sinh t|\lambda|)^{-s-1} dt.$$

Multiplying both sides by  $\frac{2|\lambda|^{s+\alpha+1}}{\Gamma(\alpha+1)} \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x)$ , we have

$$\begin{aligned} \frac{\Gamma(1-s)}{s} \frac{2|\lambda|^{\alpha+1}}{\Gamma(\alpha+1)} (2|\lambda|)^s \frac{\Gamma\left(\frac{2k+\alpha+1}{2} + \frac{1+s}{2}\right)}{\Gamma\left(\frac{2k+\alpha+1}{2} + \frac{1-s}{2}\right)} \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x) &= c_1 \frac{2|\lambda|^{\alpha+1}}{\Gamma(\alpha+1)} |\lambda|^s \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x) \\ &+ \frac{2|\lambda|^{\alpha+1}}{\Gamma(\alpha+1)} \int_0^\infty (1 - e^{-(2k+\alpha+1)|\lambda|t}) \left(\frac{t|\lambda|}{\sinh t\lambda}\right)^{s+1} \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x) t^{-s-1} dt. \end{aligned}$$

Using (2.7), (2.4) and summing over  $k$ , we obtain

$$\begin{aligned} (3.5) \quad \frac{\Gamma(1-s)}{s} \frac{2|\lambda|^{\alpha+1}}{\Gamma(\alpha+1)} (2|\lambda|)^s \sum_{k=0}^\infty \frac{\Gamma\left(\frac{2k+\alpha+1}{2} + \frac{1+s}{2}\right)}{\Gamma\left(\frac{2k+\alpha+1}{2} + \frac{1-s}{2}\right)} \widehat{f}(\lambda, k) \phi_{k,\lambda}^\alpha(x) \\ = c_1 |\lambda|^s f^\lambda(x) + \int_0^\infty (f^\lambda(x) - f^\lambda *_\lambda h_t^\lambda(x)) \left(\frac{t\lambda}{\sinh t\lambda}\right)^{s+1} t^{-s-1} dt. \end{aligned}$$

We now rewrite the last integral as a sum of the following two integrals:

$$A = f^\lambda(x) \int_0^\infty \left( \left(\frac{t\lambda}{\sinh t\lambda}\right)^{s+1} - 1 \right) t^{-s-1} dt,$$

$$B = \int_0^\infty \left( f^\lambda(x) - \left(\frac{t\lambda}{\sinh t\lambda}\right)^{s+1} f^\lambda *_\lambda h_t^\lambda(x) \right) t^{-s-1} dt.$$

Note that the first integral  $A$  is equal to

$$|\lambda|^s f^\lambda(x) \int_0^\infty \left( \left(\frac{t}{\sinh t}\right)^{s+1} - 1 \right) t^{-s-1} dt =: -c_2 |\lambda|^s f^\lambda(x).$$

It happens that  $c_1 = c_2$ . Indeed,

$$\begin{aligned} c_1 - c_2 &= \int_0^\infty (\cosh t - 1)(\sinh t)^{-s-1} dt + \int_0^\infty \left( \left(\frac{t}{\sinh t}\right)^{s+1} - 1 \right) t^{-s-1} dt \\ &= \int_0^\infty ((\cosh t)(\sinh t)^{-s-1} - t^{-s-1}) dt. \end{aligned}$$

Consider the integral

$$\int_\delta^\infty (\cosh t)(\sinh t)^{-s-1} dt = \int_{\sinh \delta}^\infty t^{-s-1} dt = \int_\delta^\infty t^{-s-1} dt - \int_\delta^{\sinh \delta} t^{-s-1} dt.$$

This gives

$$\int_\delta^\infty ((\cosh t)(\sinh t)^{-s-1} - t^{-s-1}) dt = - \int_\delta^{\sinh \delta} t^{-s-1} dt,$$



which converges to 0 as  $\delta \rightarrow 0$ . Finally, using the expression of  $B$ , we multiply (3.5) by  $e^{-i\lambda w}$  and integrate over  $\lambda$  variable to get

$$\mathcal{L}_s f(x, w) = \frac{s}{\Gamma(1-s)} \int_0^\infty (f(x, w) - f * K_{t,s}(x, w)) t^{-s-1} dt.$$

Since  $\frac{s}{\Gamma(1-s)} = \frac{1}{|\Gamma(-s)|}$ , we obtain the desired representation of  $\mathcal{L}_s$ .  $\square$

We can modify the integral representation using the properties of  $K_{t,s}$  (Proposition 2.8):

**Proposition 3.3.** *For  $0 < s < 1$ ,  $f \in W^{s,2}(X)$ , we have*

$$\mathcal{L}_s f(\xi) = \frac{1}{\Gamma(-s)} \int_0^\infty \int_X (f(\xi) - f(\eta)) T^\eta K_{t,s}(\xi) t^{-s-1} d\mu(\eta) dt.$$

*Proof.* From last theorem, we have

$$\begin{aligned} \mathcal{L}_s f(\xi) &= \frac{1}{\Gamma(-s)} \int_0^\infty (f(\xi) - f * K_{t,s}(\xi)) t^{-s-1} dt \\ &= \frac{1}{\Gamma(-s)} \int_0^\infty \left( \int_X f(\xi) T^\eta K_{t,s}(\xi) d\mu(\eta) - \int_X T^\eta K_{t,s}(\xi) f(\eta) d\mu(\eta) \right) t^{-s-1} dt \\ &= \frac{1}{\Gamma(-s)} \int_0^\infty \int_X (f(\xi) - f(\eta)) T^\eta K_{t,s}(\xi) d\mu(\eta) t^{-s-1} dt. \end{aligned}$$

$\square$

**Proposition 3.4.** *For  $0 < s < 1$  and  $f, g \in W^{s,2}(X)$ , we have*

$$\langle \mathcal{L}_s f, g \rangle = \frac{1}{2\Gamma(-s)} \int_0^\infty \int_X \int_X (f(\xi) - f(\eta)) \overline{(g(\xi) - g(\eta))} T^\eta K_{t,s}(\xi) d\mu(\eta) d\mu(\xi) \frac{dt}{t^{s+1}}.$$

*Proof.* We have

$$\begin{aligned} \langle \mathcal{L}_s f, g \rangle &= \frac{1}{\Gamma(-s)} \int_X \left( \int_0^\infty \int_X (f(\xi) - f(\eta)) \overline{g(\xi)} T^\eta K_{t,s}(\xi) d\mu(\eta) \frac{dt}{t^{s+1}} \right) d\mu(\xi) \\ &= \frac{1}{\Gamma(-s)} \int_X \left( \int_0^\infty \int_X (f(\eta) - f(\xi)) \overline{g(\eta)} T^\xi K_{t,s}(\eta) d\mu(\xi) \frac{dt}{t^{s+1}} \right) d\mu(\eta) \\ &= -\frac{1}{\Gamma(-s)} \int_X \left( \int_0^\infty \int_X (f(\xi) - f(\eta)) \overline{g(\eta)} T^{\eta*} K_{t,s}(\xi^*) d\mu(\xi) \frac{dt}{t^{s+1}} \right) d\mu(\eta), \end{aligned}$$

where  $(x, w)^* = (x, -w)$ . Using,  $T^{\eta*} K_{t,s}(\xi^*) = T^\eta K_{t,s}(\xi)$  and Fubini's Theorem, we get:

$$\langle \mathcal{L}_s f, g \rangle = -\frac{1}{\Gamma(-s)} \int_X \left( \int_0^\infty \int_X (f(\xi) - f(\eta)) \overline{g(\eta)} T^\eta K_{t,s}(\xi) d\mu(\eta) \frac{dt}{t^{s+1}} \right) d\mu(\xi).$$

Hence, we have

$$\begin{aligned} \langle \mathcal{L}_s f, g \rangle &= \frac{1}{2\Gamma(-s)} \int_X \left( \int_0^\infty \int_X (f(\xi) - f(\eta)) \overline{g(\xi) - g(\eta)} T^\eta K_{t,s}(\xi) d\mu(\eta) \frac{dt}{t^{s+1}} \right) d\mu(\xi). \\ &= \frac{1}{2\Gamma(-s)} \int_0^\infty \left( \int_X \int_X (f(\xi) - f(\eta)) \overline{g(\xi) - g(\eta)} T^\eta K_{t,s}(\xi) d\mu(\eta) d\mu(\xi) \right) \frac{dt}{t^{s+1}}. \end{aligned}$$

$\square$

Finally, for  $0 < s < 1$  and  $\delta > 0$ , we define *ground state representation*  $\mathcal{H}[f]$  for  $f \in W^{s,2}(X)$  by:

$$\mathcal{H}[f] = \langle \mathcal{L}_s f, f \rangle - A_{\alpha,s} \int_X \frac{|f(x, w)|^2}{\left( (\delta + \frac{x^2}{2})^2 + w^2 \right)^s},$$

where

$$A_{\alpha,s} = (4\delta)^s \left( \frac{\Gamma(\frac{\alpha+s+2}{2})}{\Gamma(\frac{\alpha-s+2}{2})} \right)^2.$$

**Proposition 3.5.** *Let  $0 < s < 1$ ,  $\delta > 0$  and  $F \in C_c^\infty(X)$ , that is  $F$  is an infinitely differentiable function defined on  $X$  with compact support. If we define  $G(\xi) = \frac{F(\xi)}{u_{-s,\delta}(\xi)}$ , then*

$$\mathcal{H}_s[F] = \frac{1}{2\Gamma(-s)} \int_0^\infty \int_X \int_X |G(\xi) - G(\eta)|^2 T^\eta K_{t,s}(\xi) u_{-s,\delta}(\xi) u_{-s,\delta}(\eta) d\mu(\eta) d\mu(\xi) \frac{dt}{t^{s+1}},$$

where  $u_{s,\delta}$  is defined in (3.1).

*Proof.* In the previous proposition, if we take  $g(\xi) = u_{-s,\delta}(\xi)$  and  $f(\xi) = \frac{|F(\xi)|^2}{u_{-s,\delta}(\xi)}$ , then we have  $\langle \mathcal{L}_s f, g \rangle$  equals to

$$\frac{1}{2\Gamma(-s)} \int_0^\infty \int_X \int_X (g(\xi) - g(\eta)) \left( \frac{F^2(\xi)}{g(\xi)} - \frac{F^2(\eta)}{g(\eta)} \right) T^\eta K_{t,s}(\xi) d\mu(\eta) d\mu(\xi) \frac{dt}{t^{s+1}}.$$

On simplification, we get

$$\frac{1}{2\Gamma(-s)} \int_0^\infty \int_X \int_X \left( |F(\xi) - F(\eta)|^2 - \left| \frac{F(\xi)}{g(\xi)} - \frac{F(\eta)}{g(\eta)} \right|^2 g(\xi)g(\eta) \right) T^\eta K_{t,s}(\xi) d\mu(\eta) d\mu(\xi) \frac{dt}{t^{s+1}}.$$

On the other hand, using Proposition 3.1 and the fact that  $\mathcal{L}_s$  is self-adjoint, we have  $\langle \mathcal{L}_s f, g \rangle$  equals to

$$(4\delta)^s \left( \frac{\Gamma(\frac{\alpha+s+2}{2})}{\Gamma(\frac{\alpha-s+2}{2})} \right)^2 \int_X \frac{|F(\xi)|^2}{u_{-s,\delta}(\xi)} u_{s,\delta}(\xi) d\mu(\xi).$$

Equating both, and noting that  $u_{-s,\delta}(x, w)/u_{s,\delta}(x, w) = ((\delta + x^2/2)^2 + w^2)^s$ , we get the desired result.  $\square$

Finally we prove Hardy inequality for fractional powers of generalized sublaplacian.

*Proof of Theorem 1.2.* Let  $f \in C_c^\infty(X)$ . Define  $g(\xi) = f(\xi)/u_{-s,\delta}(\xi)$ , where  $u_{s,\delta}$  is defined in (3.1). Using the last Proposition and Fubini's Theorem, we have

$$\mathcal{H}_s[f] = \frac{1}{2\Gamma(-s)} \int_X \int_X \left( \int_0^\infty T^\eta K_{t,s}(\xi) \frac{dt}{t^{s+1}} \right) |g(\xi) - g(\eta)|^2 u_{-s,\delta}(\xi) u_{-s,\delta}(\eta) d\mu(\eta) d\mu(\xi).$$

Since the generalized translation operator is a positive operator (see [9][Proposition 3.2]) and  $\int_0^\infty K_{t,s}(\xi) t^{-s-1} dt \geq 0$  (Proposition 2.9), we have for all  $\xi, \eta \in X$

$$\int_0^\infty T^\eta K_{t,s}(\xi) \frac{dt}{t^{s+1}} \geq 0.$$

Also the remaining terms in the expression of  $\mathcal{H}_s[f]$  are positive for all  $\xi$  and  $\eta$ . Therefore, we conclude  $\mathcal{H}_s[f] \geq 0$  for all  $f \in C_c^\infty(X)$ . Hence, for  $f \in C_c^\infty(X)$ , we have

$$\langle \mathcal{L}_s f, f \rangle \geq (4\delta)^s \left( \frac{\Gamma(\frac{\alpha+s+2}{2})}{\Gamma(\frac{\alpha-s+2}{2})} \right)^2 \int_X \frac{|f(\xi)|^2}{\left( (\delta + \frac{x^2}{2})^2 + w^2 \right)^s} d\mu(\xi).$$

Next, let  $f \in W^{s,2}(X)$ . Since  $C_c^\infty(X)$  dense in  $W^{s,2}(X)$ , therefore there exists a sequence  $\{f_j\}$  with each  $f_j \in C_c^\infty(X)$  such that  $f_j \rightarrow f$  in  $W^{s,2}(X)$ . Passing to a sub-sequence, we can assume  $f_j \rightarrow f$  pointwise a.e.. The continuity of inner-product on  $W^{s,2}(X)$  implies  $\langle \mathcal{L}_s f_j, f_j \rangle \rightarrow \langle \mathcal{L}_s f, f \rangle$ . On the other hand, the inequality  $|f(x, w)|^2 / \left( (\delta + x^2)^2 + w^2 \right)^s \leq \delta^{-2s} |f(x, w)|^2$  together with Dominated convergence theorem, implies

$$\int_X \frac{|f_j(\xi)|^2}{\left( (\delta + \frac{x^2}{2})^2 + w^2 \right)^s} d\mu(\xi) \rightarrow \int_X \frac{|f(\xi)|^2}{\left( (\delta + \frac{x^2}{2})^2 + w^2 \right)^s} d\mu(\xi).$$

From this we conclude that the inequality holds for all  $f \in W^{s,2}(X)$ .

Finally, we note that the both sides of the inequality are equal for  $f = u_{-s,\delta}$ . Hence the constant involved in the inequality is sharp.  $\square$

#### 4. PROOF OF THE MAIN THEOREM

We recall from Section 2.6 that for  $j = 1, 2, \dots, a_m$  and  $m = 0, 1, 2, \dots$ ,  $\{Y_{m,j}\}$  forms orthonormal basis for  $L^2(\mathbb{S}^{n-1})$ . Corresponding to each spherical harmonic  $Y_{m,j}$ , we define solid harmonics  $P_{m,j}$  on  $\mathbb{R}^n$  by

$$P_{m,j}(x) = |x|^m Y_{m,j}(x/|x|).$$

Moreover, for  $f \in L^2(\mathbb{R}^{n+1})$ , we have

$$f(x, w) = \sum_{m=0}^{\infty} \sum_{j=1}^{a_m} f_{m,j}(|x|, w) P_{m,j}(x),$$

where  $f_{m,j}(|x|, w) = \int_{\mathbb{S}^{n-1}} f(|x|\omega, w) P_{m,j}(|x|\omega) d\omega$ .

Suppose  $f \in W^{s,2}(H)$ , such that  $f(x, w) = g(|x|, w)P(x)$ , where  $P$  is a solid harmonics of degree  $m$ . Then using spectral decomposition and the Hecke Bockner formula (Proposition 2.10), we have

$$\begin{aligned} \mathcal{G}_s f(x, w) &= \frac{1}{2\pi} \int_{\mathbb{R}} \sum_{k=0}^{\infty} (2|\lambda|)^s \frac{\Gamma(\frac{2k+n}{4} + \frac{1+s}{2})}{\Gamma(\frac{2k+n}{4} + \frac{1-s}{2})} \mathcal{P}_k(\lambda) f^\lambda(x) e^{-i\lambda w} d\lambda \\ &= \int_{\mathbb{R}} \sum_{k=0}^{\infty} \left( d_{m,k}^n(s) A_{k,m}^\lambda(g) \phi_{k,\lambda}^{n/2+m-1}(|x|) \right) P(x) |\lambda|^{n/2+m+s} e^{-i\lambda w} d\lambda, \end{aligned}$$

where

$$d_{m,k}^n(s) = \frac{2^{s+1} \Gamma(\frac{2k+n/2+m+1+s}{2})}{2\pi \Gamma(\frac{2k+n/2+m+1-s}{2})} \frac{\Gamma(k+1)}{\Gamma(k+n/2+m)},$$

and

$$A_{k,m}^\lambda(g) = \int_0^\infty g^\lambda(r) \phi_{k,\lambda}^{n/2+m-1}(r) r^{n+2m-1} dr.$$

Therefore, using the orthogonality of solid harmonics (with respect to inner product inherited from  $L^2(\mathbb{S}^{n-1})$ ), we have

$$\langle \mathcal{G}_s f, f \rangle = \int_{\mathbb{R}} \sum_{k=0}^{\infty} d_{m,k}^n(s) |A_{k,m}^\lambda(g)|^2 |\lambda|^{\frac{n}{2}+m+s} d\lambda.$$

Moreover, treating  $g$  as a function on  $X$ , we have

$$\langle \mathcal{L}_s g, g \rangle = \frac{2^{s+1}}{2\pi} \int_{\mathbb{R}} \sum_{k=0}^{\infty} \frac{\Gamma(\frac{2k+\alpha+2+s}{2})}{\Gamma(\frac{2k+\alpha+2-s}{2})} \frac{\Gamma(k+1)}{\Gamma(k+\alpha+1)} |B_{k,\alpha}^\lambda(g)|^2 |\lambda|^{\alpha+1+s} d\lambda,$$

where  $B_{k,\alpha}^\lambda(g) = \int_0^\infty g^\lambda(x) \phi_{k,\lambda}^\alpha(x) x^{2\alpha+1} dx$ . So for  $\alpha = \frac{n}{2} + m - 1$ , we have

$$(4.1) \quad \langle \mathcal{G}_s f, f \rangle = \langle \mathcal{L}_s g, g \rangle.$$

Now let  $f \in C_c^\infty(H)$ . Since  $C_c^\infty(H)$  dense in  $W^{s,2}(H)$ , we have

$$f(x, w) = \sum_{m=0}^{\infty} \sum_{j=1}^{a_m} f_{m,j}(|x|, w) P_{m,j}(x),$$

where  $f_{m,j}(|x|, w) = \int_{\mathbb{S}^{n-1}} f(|x|\omega, w) P_{m,j}(|x|\omega) d\omega$ . Using (4.1), we have

$$\begin{aligned} \langle \mathcal{G}_s f, f \rangle &= \sum_{m=0}^{\infty} \sum_{j=1}^{a_m} \langle \mathcal{G}_s(f_{m,j} P_{m,j}), f_{m,j} P_{m,j} \rangle \\ &= \sum_{m=0}^{\infty} \sum_{j=1}^{a_m} \langle \mathcal{L}_s f_{m,j}, f_{m,j} \rangle. \end{aligned}$$

Using the Theorem 1.2, we have

$$\begin{aligned} \langle \mathcal{G}_s f, f \rangle &\geq \sum_{m=0}^{\infty} \sum_{j=1}^{a_m} (4\delta)^s \left( \frac{\Gamma(\frac{n/2+m+s+1}{2})}{\Gamma(\frac{n/2+m-s+1}{2})} \right)^2 \int_{\mathbb{R}} \int_0^\infty \frac{|f_{m,j}(x, w)|^2 x^{n+2m-1}}{\left( (\delta + \frac{x^2}{2})^2 + w^2 \right)^s} dx dw \\ &\geq \inf_{m \geq 0} \left\{ (4\delta)^s \left( \frac{\Gamma(\frac{n/2+m+s+1}{2})}{\Gamma(\frac{n/2+m-s+1}{2})} \right)^2 \right\} \sum_{m=0}^{\infty} \sum_{j=1}^{a_m} \int_{\mathbb{R}} \int_0^\infty \frac{|f_{m,j}(x, w)|^2 x^{n+2m-1}}{\left( (\delta + \frac{|x|^2}{2})^2 + w^2 \right)^s} dx dw. \end{aligned}$$

Moreover, we have

$$\int_{\mathbb{R}^n} |f(x, w)|^2 dx = \sum_{m=0}^{\infty} \sum_{j=1}^{a_m} \int_0^\infty |f_{m,j}(r, w)|^2 r^{n+2m-1} dr,$$

and

$$\inf_{m \geq 0} \left\{ (4\delta)^s \left( \frac{\Gamma(\frac{n/2+m+s+1}{2})}{\Gamma(\frac{n/2+m-s+1}{2})} \right)^2 \right\} = (4\delta)^s \left( \frac{\Gamma(\frac{n/2+s+1}{2})}{\Gamma(\frac{n/2-s+1}{2})} \right)^2.$$

Therefore,

$$\langle \mathcal{G}_s f, f \rangle \geq (4\delta)^s \left( \frac{\Gamma(\frac{n/2+s+1}{2})}{\Gamma(\frac{n/2-s+1}{2})} \right)^2 \int_{\mathbb{R}} \int_{\mathbb{R}^n} \frac{|f(x, w)|^2}{\left( (\delta + \frac{|x|^2}{2})^2 + w^2 \right)^s} dx dw.$$

Once we proved the inequality for  $f \in C_c^\infty(H)$ , we argue as in the proof of Theorem 3 to conclude that the inequality holds true for  $f \in W^{s,2}(H)$ .

Finally, we show that the constants involved in the inequality are sharp. For  $-1 < s < 1$  and  $\delta > 0$ , define  $v_{s,\delta}$  on  $H$  by

$$v_{s,\delta}(x, w) = ((\delta + |x|^2/2)^2 + w^2)^{-\frac{n/2+1+s}{2}}.$$

Using Proposition 3.1, we have

$$\begin{aligned} \langle \mathcal{G}_s v_{-s,\delta}, v_{-s,\delta} \rangle &= \langle \mathcal{L}_s v_{-s,\delta}, v_{-s,\delta} \rangle \\ &= (4\delta)^s \frac{\Gamma^2(\frac{n/2+s+1}{2})}{\Gamma^2(\frac{n/2-s+1}{2})} \langle v_{s,\delta}, v_{-s,\delta} \rangle \\ &= (4\delta)^s \frac{\Gamma^2(\frac{n/2+s+1}{2})}{\Gamma^2(\frac{n/2-s+1}{2})} \int_{\mathbb{R}} \int_0^\infty \frac{x^{n-1} dx dw}{\left( (\delta + \frac{x^2}{2})^2 + w^2 \right)^{n/2+1}} \\ &= (4\delta)^s \frac{\Gamma^2(\frac{n/2+s+1}{2})}{\Gamma^2(\frac{n/2-s+1}{2})} \int_{\mathbb{R}} \int_{\mathbb{R}^n} \frac{|u_{-s,\delta}(x, w)|^2 dx dw}{\left( (\delta + \frac{x^2}{2})^2 + w^2 \right)^s}. \end{aligned}$$

Therefore, equality is achieved for  $f = u_{-s,\delta}$ . Hence the constants involved in the inequality are sharp.

## REFERENCES

- [1] Beckner W, Pitts inequality and the fractional Laplacian: sharp error estimates, *Forum Math.* **24** (2012), 177209
- [2] Ciaurri O, Roncal L, Thangavelu S, Hardy-type inequalities for fractional powers of the Dunkl-Hermite operator, arXiv preprint arXiv:1602.04997 (2016)
- [3] Cowling M and Haagerup U, Completely bounded multipliers of the Fourier algebra of a simple Lie group of real rank one, *Invent. Math.* **96** (1989), 507549
- [4] Frank R L, Lieb E H and Seiringer R, Hardy-Lieb-Thirring inequalities for fractional Schrödinger operators, *J. Amer. Math. Soc.* **21** (2008), 925950
- [5] Gradshteyn I S and Ryzhik I M, Table of Integrals, Series and Products, Seventh edition, Elsevier Academic Press, Amsterdam, 2007
- [6] Herbst I W, Spectral theory of the operator  $(p^2 + m^2)^{1/2} Z e^2/r$ , *Commun. Math. Phys.* **53** (1977), 285294
- [7] Roncal L and Thangavelu S, Hardy's inequality for fractional powers of the sublaplacian on the Heisenberg group, *Advances in Mathematics* **302** (2016): 106-158.
- [8] Stein E M and Weiss G, Introduction to Fourier analysis on Euclidean spaces(PMS-32), Vol. 32, Princeton university press, 2016
- [9] Stempak K, An algebra associated with the generalized sublaplacian, *Studia Math.* **88** (1988), 245-256
- [10] Stempak K, Mean summability methods for Laguerre series, *Trans. Amer. Math. Soc.* **322**(1990), 671-690
- [11] Thangavelu S, An introduction to the uncertainty principle. Hardys theorem on Lie groups. With a foreword by Gerald B. Folland, Progress in Mathematics **217**. Birkhäuser, Boston, MA, 2004
- [12] Thangavelu S, Lectures on Hermite and Laguerre expansions, Math. Notes. **42**, Princeton University Press, Princeton, NJ, 1993
- [13] Yafaev D, Sharp constants in the Hardy-Rellich inequalities, *J. Funct. Anal.* **168** (1999), 121144

(R. Balhara) DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, 560 012 BANGALORE,  
INDIA

*E-mail address:* `rakeshbalhara@gmail.com`