# MULTIPLICITY AND CONCENTRATION RESULTS FOR A FRACTIONAL CHOQUARD EQUATION VIA PENALIZATION METHOD

#### VINCENZO AMBROSIO

ABSTRACT. This paper is devoted to the study of the following fractional Choquard equation

$$\varepsilon^{2s}(-\Delta)^s u + V(x)u = \varepsilon^{\mu-N} \left(\frac{1}{|x|^{\mu}} * F(u)\right) f(u) \text{ in } \mathbb{R}^N,$$

where  $\varepsilon > 0$  is a parameter,  $s \in (0,1), N > 2s, (-\Delta)^s$  is the fractional Laplacian, V is a positive continuous potential with local minimum,  $0 < \mu < 2s$ , and f is a superlinear continuous function with subcritical growth. By using the penalization method and the Ljusternik-Schnirelmann theory, we investigate the multiplicity and concentration of positive solutions for the above problem.

#### 1. Introduction

In this paper we deal with the following nonlinear fractional Choquard equation

$$\varepsilon^{2s}(-\Delta)^s u + V(x)u = \varepsilon^{\mu - N} \left(\frac{1}{|x|^{\mu}} * F(u)\right) f(u) \text{ in } \mathbb{R}^N,$$
(1.1)

where  $\varepsilon > 0$  is a parameter,  $s \in (0,1)$ , N > 2s and  $0 < \mu < 2s$ . The potential  $V : \mathbb{R}^N \to \mathbb{R}$  is a continuous function verifying the following hypotheses:

- $(V_1) \inf_{x \in \mathbb{R}^N} V(x) = V_0 > 0;$
- $(V_2)$  there exists a bounded open set  $\Lambda \subset \mathbb{R}^N$  such that

$$V_0 < \min_{x \in \partial \Lambda} V(x).$$

Concerning the nonlinearity  $f: \mathbb{R} \to \mathbb{R}$ , we assume that f is a continuous function such that f(t) = 0 for t < 0, and satisfies the following conditions:

- $(f_1) \lim_{t \to 0} \frac{f(t)}{t} = 0;$
- $(f_2)$  there exists  $q \in (2, \frac{2_s^*}{2}(2 \frac{\mu}{N}))$ , where  $2_s^* = \frac{2N}{N-2s}$ , such that  $\lim_{t \to \infty} \frac{f(t)}{t^{q-1}} = 0$ ;
- $(f_3)$  f verifies the following Ambrosetti-Rabinowitz type condition [5]:

$$0 < 4F(t) = 4 \int_0^t f(\tau) d\tau \le 2f(t) t$$
 for all  $t > 0$ ;

 $(f_4)$  The map  $t \mapsto \frac{f(t)}{t}$  is increasing for every t > 0.

The nonlocal operator  $(-\Delta)^s$  is the fractional Laplacian which may be defined for any  $u: \mathbb{R}^N \to \mathbb{R}$  sufficiently smooth by

$$(-\Delta)^{s} u(x) = -\frac{C(N,s)}{2} \int_{\mathbb{R}^{N}} \frac{u(x+y) + u(x-y) - 2u(x)}{|y|^{N+2s}} dy \quad (x \in \mathbb{R}^{N}),$$

where C(N, s) is a suitable normalization constant; see for instance [16, 28].

<sup>2010</sup> Mathematics Subject Classification. 35A15, 35B09, 35R11, 45G05.

Key words and phrases. Fractional Choquard equation; penalization method; multiplicity of solutions.

We recall that the problem (1.1) is motivated by the search of standing wave solutions for the following fractional Schrödinger equation

$$i\frac{\partial \psi}{\partial t} = (-\Delta)^s \psi + V(x)\psi - \left(\frac{1}{|x|^{\mu}} * |\Psi|^q\right) |\Psi|^{q-2} \Psi \quad (t, x) \in \mathbb{R} \times \mathbb{R}^N,$$

which naturally models many physical problems, such as phase transition, conservation laws, especially in fractional quantum mechanics. For physical motivations we refer to [6, 7, 14, 18, 19, 20, 21, 22, 36].

When  $s=1, V(x)\equiv 1, \varepsilon=1$  and  $F(u)=\frac{|u|^2}{2}$ , (1.1) boils down to the Choquard-Pekar equation

$$-\Delta u + u = \left(\frac{1}{|x|^{\mu}} * |u|^2\right) u \text{ in } \mathbb{R}^N$$
 (1.2)

introduced by Pekar [33] to describe the quantum mechanics of a polaron. Subsequently, Choquard used (1.2) to describe an electron trapped in its own hole as approximation to Hartree-Fock Theory of one component plasma; see [24, 34].

The early existence and symmetry results are due to Lieb [23] and Lions [25]. Later, Ma and Zhao [26] obtained some qualitative properties of positive solutions considering powers like  $|u|^q$ . Moroz and Van Shaftingen [30] investigated regularity, radial symmetry and asymptotic behavior at infinity of positive solutions for a generalized Choquard equation. Alves and Yang [4] studied multiplicity and concentration of positive solutions for a Choquard equation. Further results on Choquard equations can be found in [1, 3, 31, 35, 40, 41].

In the case  $s \in (0,1)$ , only few recent papers considered fractional Choquard equations like (1.1). In [13] d'Avenia et al. considered the following fractional Choquard equation

$$(-\Delta)^s u + u = \left(\frac{1}{|x|^{\mu}} * |u|^p\right) |u|^{p-2} u \text{ in } \mathbb{R}^N,$$

obtaining regularity, existence and non existence, symmetry and decay properties of solutions. Frank and Lenzman [20] established uniqueness of nonnegative ground states for the  $L^2$  critical boson star equation

$$(-\Delta)^{\frac{1}{2}}u + u = \left(\frac{1}{|x|^{\mu}} * |u|^2\right)u \text{ in } \mathbb{R}^3,$$

by using variational methods and the extension technique [10]. Coti Zelati and Nolasco [12] obtained existence of ground state solutions for a pseudo-relativistic Hartree-equation via critical point theory. Shen et al. [37] investigated the existence of ground state solutions for a fractional Choquard equation involving a nonlinearity satisfying Berestycki-Lions type assumptions. Chen and Liu [11] studied an autonomous fractional Choquard equation via Nehari manifold and concentration-compactness arguments. Belchior et al. [8] dealt with existence, regularity and polynomial decay for a fractional Choquard equation involving the fractional p-Laplacian.

Motivated by the above papers, in this work we focus our attention on the multiplicity and the concentration of positive solutions of (1.1), involving a potential and a continuous nonlinearity satisfying the assumptions  $(V_1)$ - $(V_2)$  and  $(f_1)$ - $(f_4)$  respectively. In particular, we are interested in relating the number of positive solutions of (1.1) with the topology of the set  $M = \{x \in \Lambda : V(x) = V_0\}$ . In order to state precisely our result, we recall that if Y is a given closed set of a topological space X, we denote by  $cat_X(Y)$  the Ljusternik-Schnirelmann category of Y in X, that is the least number of closed and contractible sets in X which cover Y.

The main result of this paper is the following:

**Theorem 1.1.** Suppose that V verifies  $(V_1)$ - $(V_2)$ ,  $0 < \mu < 2s$  and f satisfies  $(f_1)$ - $(f_4)$  with  $2 < q < \frac{2(N-\mu)}{N-2s}$ . Then, for any  $\delta > 0$  such that  $M_{\delta} = \{x \in \mathbb{R}^N : dist(x,M) \leq \delta\} \subset \Lambda$ , there exists  $\varepsilon_{\delta} > 0$ 

such that, for any  $\varepsilon \in (0, \varepsilon_{\delta})$ , the problem (1.1) has at least  $cat_{M_{\delta}}(M)$  positive solutions. Moreover, if  $u_{\varepsilon}$  denotes one of these positive solutions and  $x_{\varepsilon} \in \mathbb{R}^{N}$  its global maximum, then

$$\lim_{\varepsilon \to 0} V(x_{\varepsilon}) = V_0.$$

Firstly, we note that the restriction on the exponent q is justified by the Hardy-Littlewood-Sobolev inequality (see Theorem 2.2 in Section 2). Indeed, if  $F(u) = |u|^q$ , then the term

$$\int_{\mathbb{R}^N} \left( \frac{1}{|x|^{\mu}} * F(u) \right) F(u) \, dx$$

is well-defined if  $F(u) \in L^t(\mathbb{R}^N)$  for t > 1 such that  $\frac{2}{t} + \frac{\mu}{N} = 2$ . Hence, recalling that  $H^s(\mathbb{R}^N)$  is continuously embedded into  $L^r(\mathbb{R}^N)$  for any  $r \in [2, 2_s^*]$ , we need to require that  $tq \in [2, 2_s^*]$ , which leads to assume that

$$2 - \frac{\mu}{N} \le q \le \frac{2_s^*}{2} \left( 2 - \frac{\mu}{N} \right).$$

Now, we give a sketch of the proof of Theorem 1.1. Inspired by [4], we adapt the del Pino-Felmer penalization technique [15] considering an auxiliary problem. It consists in making a suitable modification on the nonlinearity f, solving a modified problem and then check that, for  $\varepsilon$  sufficiently small, the solutions of the modified problem are indeed solutions of the original one. Differently from the case s=1, in our setting a more accurate investigation is needed due to the presence of two non-local terms. Moreover, the nonlinearity appearing in (1.1) is only continuous (while  $f \in C^1$ in [4]), so to overcome the non-differentiability of the associated Nehari manifold, we will use some abstract critical point results due to Szulkin and Weth [39]. Concerning the multiplicity result for the modified problem, we resemble some ideas due to Benci and Cerami in [9], based on the comparison between the category of some sublevel sets of the modified functional and the category of the set M. Finally, in order to prove that the solutions  $u_{\varepsilon}$  of the modified problem are solutions of the problem (1.1), we adapt a Moser iteration argument [29] to establish  $L^{\infty}$ -estimates, and after showed that the convolution term remains bounded, we exploit some useful properties of the Bessel kernel [2, 19] to obtain the desired result. To our knowledge, this is the first result in which the concentration and the multiplicity of solutions to (1.1) are considered by using penalization argument and Ljusternik-Schnirelmann category theory.

The plan of the paper is the following: In Section 2 we introduce the functional setting and the modified problem. The Section 3 is devoted to the existence of positive solutions to the autonomous problem associated to (1.1). In Section 4, we obtain a multiplicity result using Ljusternik-Schnirelmann theory. Finally, exploiting a Moser iteration scheme, we are able to prove that for  $\varepsilon$  small enough, the solutions of the modified problem are indeed solutions of (1.1).

### 2. Functional setting

For any  $s \in (0,1)$ , we denote by  $\mathcal{D}^{s,2}(\mathbb{R}^N)$  the completion of  $C_0^{\infty}(\mathbb{R}^N)$  with respect to

$$[u]^{2} = \iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{2}}{|x - y|^{N+2s}} dx dy = \|(-\Delta)^{\frac{s}{2}} u\|_{L^{2}(\mathbb{R}^{N})}^{2},$$

that is

$$\mathcal{D}^{s,2}(\mathbb{R}^N) = \left\{ u \in L^{2^*_s}(\mathbb{R}^N) : [u]_{H^s(\mathbb{R}^N)} < \infty \right\}.$$

Let us introduce the fractional Sobolev space

$$H^{s}(\mathbb{R}^{N}) = \left\{ u \in L^{2}(\mathbb{R}^{N}) : \frac{|u(x) - u(y)|}{|x - y|^{\frac{N+2s}{2}}} \in L^{2}(\mathbb{R}^{2N}) \right\}$$

endowed with the natural norm

$$||u|| = \sqrt{[u]^2 + ||u||_{L^2(\mathbb{R}^N)}^2}.$$

We collect the following useful results.

**Theorem 2.1.** [16] Let  $s \in (0,1)$  and N > 2s. Then there exists a sharp constant  $S_* = S(N,s) > 0$  such that for any  $u \in H^s(\mathbb{R}^N)$ 

$$||u||_{L^{2_*^*}(\mathbb{R}^N)}^2 \le S_*^{-1}[u]^2. \tag{2.1}$$

Moreover  $H^s(\mathbb{R}^N)$  is continuously embedded in  $L^q(\mathbb{R}^N)$  for any  $q \in [2, 2_s^*]$  and compactly in  $L^q_{loc}(\mathbb{R}^N)$  for any  $q \in [2, 2_s^*]$ .

**Lemma 2.1.** [19] Let N > 2s. If  $(u_n)$  is a bounded sequence in  $H^s(\mathbb{R}^N)$  and if

$$\lim_{n \to \infty} \sup_{y \in \mathbb{R}^N} \int_{B_R(y)} |u_n|^2 dx = 0$$

where R > 0, then  $u_n \to 0$  in  $L^t(\mathbb{R}^N)$  for all  $t \in (2, 2_s^*)$ .

**Theorem 2.2.** [24] Let r, t > 1 and  $0 < \mu < N$  such that  $\frac{1}{r} + \frac{\mu}{N} + \frac{1}{t} = 2$ . Let  $f \in L^r(\mathbb{R}^N)$  and  $h \in L^t(\mathbb{R}^N)$ . Then there exists a sharp constant  $C(r, N, \mu, t) > 0$  independent of f and h such that

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{f(x)h(y)}{|x-y|^{\mu}} \, dx dy \le C(r, N, \mu, t) \|f\|_{L^r(\mathbb{R}^N)} \|h\|_{L^t(\mathbb{R}^N)}.$$

For any  $\varepsilon > 0$ , we denote by  $H^s_{\varepsilon}$  the completion of  $C_0^{\infty}(\mathbb{R}^N)$  with respect to the norm

$$||u||_{\varepsilon}^2 = \iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dxdy + \int_{\mathbb{R}^N} V(\varepsilon x)u^2(x) dx.$$

It is clear that  $H^s_{\varepsilon}$  is a Hilbert space with respect to the inner product

$$(u,v)_{\varepsilon} = \iint_{\mathbb{R}^{2N}} \frac{(u(x) - u(y))}{|x - y|^{N+2s}} (v(x) - v(y)) \, dx dy + \int_{\mathbb{R}^N} V(\varepsilon x) uv \, dx.$$

By using the change of variable  $u(x) \mapsto u(\varepsilon x)$  we can see that the problem (1.1) is equivalent to the following one

$$(-\Delta)^{s} u + V(\varepsilon x) u = \left(\frac{1}{|x|^{\mu}} * F(u)\right) f(u) \text{ in } \mathbb{R}^{N}.$$
 (2.2)

Fix  $\ell > 2$  and a > 0 such that  $\frac{f(a)}{a} = \frac{V_0}{\ell}$ , and we introduce the functions

$$\tilde{f}(t) := \begin{cases} f(t) & \text{if } t \le a \\ \frac{V_0}{\ell} t & \text{if } t > a, \end{cases}$$

and

$$g(x,t) = \chi_{\Lambda}(x)f(t) + (1 - \chi_{\Lambda}(x))\tilde{f}(t),$$

where  $\chi_{\Lambda}$  is the characteristic function on  $\Lambda$ , and we write  $G(x,t) = \int_0^t g(x,\tau) d\tau$ . Let us note that from the assumptions  $(f_1)$ - $(f_4)$ , g satisfies the following properties:

- $(g_1) \lim_{t\to 0} \frac{g(x,t)}{t} = 0$  uniformly in  $x \in \mathbb{R}^N$ ;
- $(g_2)$   $\lim_{t\to\infty} \frac{g(x,t)}{t^{q-1}} = 0$  uniformly in  $x \in \mathbb{R}^N$ ;
- $(g_3) \ 0 < 4G(x,t) \le 2g(x,t)t \text{ for any } x \in \Lambda \text{ and } t > 0, \text{ and } 0 \le 2G(x,t) \le g(x,t)t \le \frac{V_0}{\ell}t^2 \text{ for any } x \in \mathbb{R}^N \setminus \Lambda \text{ and } t > 0,$
- $(g_4)$   $t \mapsto g(x,t)$  and  $t \mapsto \frac{G(x,t)}{t}$  are increasing for all  $x \in \mathbb{R}^N$  and t > 0. Thus we consider the following auxiliary problem

$$(-\Delta)^{s} u + V(\varepsilon x) u = \left(\frac{1}{|x|^{\mu}} * G(\varepsilon x, u)\right) g(\varepsilon x, u) \text{ in } \mathbb{R}^{N}$$
(2.3)

and we note that if u is a solution of (2.3) such that

$$u(x) < a \text{ for all } x \in \mathbb{R}^N \setminus \Lambda_{\varepsilon},$$
 (2.4)

where  $\Lambda_{\varepsilon} := \{x \in \mathbb{R}^N : \varepsilon x \in \Lambda\}$ , then u solves (2.2), in view of the definition of g. It is clear that, weak solutions to (2.3) are critical points of the  $C^1$ -functional  $J_{\varepsilon} : H_{\varepsilon}^s \to \mathbb{R}$  defined by

$$J_{\varepsilon}(u) = \frac{1}{2} \|u\|_{\varepsilon}^{2} - \Sigma_{\varepsilon}(u)$$

where

$$\Sigma_{\varepsilon}(u) = \frac{1}{2} \int_{\mathbb{R}^N} \left( \frac{1}{|x|^{\mu}} * G(\varepsilon x, u) \right) G(\varepsilon x, u) dx.$$

We begin proving that  $J_{\varepsilon}$  satisfies the assumptions of the mountain pass theorem [5].

**Lemma 2.2.**  $J_{\varepsilon}$  has a mountain pass geometry, that is

- (i) there exist  $\alpha, \rho > 0$  such that  $J_{\varepsilon}(u) \geq \alpha$  for any  $u \in H^s_{\varepsilon}$  such that  $||u||_{\varepsilon} = \rho$ ;
- (ii) there exists  $e \in H^s_{\varepsilon}$  with  $||e||_{\varepsilon} > \rho$  such that  $J_{\varepsilon}(e) < 0$ .

*Proof.* From  $(g_1)$  and  $(g_2)$ , it follows that that for any  $\eta > 0$  there exists  $C_{\eta} > 0$  such that

$$|g(\varepsilon x, t)| \le \eta |t| + C_{\eta} |t|^{q-1}. \tag{2.5}$$

By using Theorem 2.2 and (2.5), we get

$$\left| \int_{\mathbb{R}^N} \left( \frac{1}{|x|^{\mu}} * G(\varepsilon x, u) \right) G(\varepsilon x, u) dx \right| \le C \|G(\varepsilon x, u)\|_{L^t(\mathbb{R}^N)} \|G(\varepsilon x, u)\|_{L^t(\mathbb{R}^N)} \le C \left( \int_{\mathbb{R}^N} (|u|^2 + |u|^q dx)^t \right)^{\frac{2}{t}},$$
(2.6)

where  $\frac{1}{t} = \frac{1}{2}(2 - \frac{\mu}{N})$ . Since  $2 < q < \frac{2_s^*}{2}(2 - \frac{\mu}{N})$ , we can see that  $tq \in (2, 2_s^*)$ , and from Theorem 2.1, we have

$$\left(\int_{\mathbb{R}^N} (|u|^2 + |u|^q \, dx)^t\right)^{\frac{2}{t}} \le C(\|u\|_{\varepsilon}^2 + \|u\|_{\varepsilon}^q)^2. \tag{2.7}$$

Taking into account (2.6) and (2.7) we can deduce that

$$\left| \int_{\mathbb{R}^N} \left( \frac{1}{|x|^{\mu}} * G(\varepsilon x, u) \right) G(\varepsilon x, u) \, dx \right| \le C(\|u\|_{\varepsilon}^2 + \|u\|_{\varepsilon}^q)^2 \le C(\|u\|_{\varepsilon}^4 + \|u\|_{\varepsilon}^{2q}).$$

As a consequence

$$J(u) \ge \frac{1}{2} ||u||_{\varepsilon}^{2} - C(||u||_{\varepsilon}^{4} + ||u||_{\varepsilon}^{2q}),$$

and being q>2 we can see that (i) holds. Fix a positive function  $u_0\in H^s(\mathbb{R}^N)\setminus\{0\}$  with  $supp(u_0)\subset\Lambda_{\varepsilon}$ , and we set

$$h(t) = \Sigma_{\varepsilon} \left( \frac{tu_0}{\|u_0\|_{\varepsilon}} \right) \text{ for } t > 0.$$

Since  $G(\varepsilon x, u_0) = F(u_0)$  and by using  $(f_3)$ , we deduce that

$$h'(t) = \Sigma_{\varepsilon}' \left(\frac{tu_0}{\|u_0\|_{\varepsilon}}\right) \frac{u_0}{\|u_0\|_{\varepsilon}}$$

$$= \int_{\mathbb{R}^N} \left(\frac{1}{|x|^{\mu}} * F\left(\frac{tu_0}{\|u_0\|_{\varepsilon}}\right)\right) f\left(\frac{tu_0}{\|u_0\|_{\varepsilon}}\right) \frac{u_0}{\|u_0\|_{\varepsilon}} dx$$

$$= \frac{4}{t} \int_{\mathbb{R}^N} \frac{1}{2} \left(\frac{1}{|x|^{\mu}} * F\left(\frac{tu_0}{\|u_0\|_{\varepsilon}}\right)\right) \frac{1}{2} f\left(\frac{tu_0}{\|u_0\|_{\varepsilon}}\right) \frac{tu_0}{\|u_0\|_{\varepsilon}} dx$$

$$> \frac{4}{t} h(t). \tag{2.8}$$

Integrating (2.8) on  $[1, t||u_0||_{\varepsilon}]$  with  $t > \frac{1}{||u_0||_{\varepsilon}}$ , we find

$$h(t||u_0||_{\varepsilon}) \ge h(1)(t||u_0||_{\varepsilon})^4$$

which gives

$$\Sigma_{\varepsilon}(tu_0) \ge \Sigma_{\varepsilon} \left(\frac{u_0}{\|u_0\|_{\varepsilon}}\right) \|u_0\|_{\varepsilon}^4 t^4.$$

Therefore, we have

$$J_{\varepsilon}(tu_0) = \frac{t^2}{2} \|u_0\|_{\varepsilon}^2 - \Sigma_{\varepsilon}(tu_0) \le C_1 t^2 - C_2 t^4 \text{ for } t > \frac{1}{\|u_0\|_{\varepsilon}}.$$

Taking  $e = tu_0$  with t sufficiently large, we can see that (ii) holds.

Since  $supp(u_0) \subset \Lambda_{\varepsilon}$ , there exists  $\kappa > 0$  independent of  $\varepsilon, l, a$  such that

$$c_{\varepsilon} = \inf_{u \in H_{\varepsilon}^{s} \setminus \{0\}} \max_{t \ge 0} J_{\varepsilon}(tu) < \kappa.$$

Now, let us define

$$\mathcal{B} = \{ u \in H^s(\mathbb{R}^N) : ||u||_{\varepsilon}^2 \le 4(\kappa + 1) \}$$

and we set

$$\tilde{K}_{\varepsilon}(u)(x) = \frac{1}{|x|^{\mu}} * G(\varepsilon x, u).$$

We prove the following useful lemma.

**Lemma 2.3.** Assume that  $(f_1)$ - $(f_3)$  hold and  $2 < q < \frac{2(N-\mu)}{N-2s}$ . Then there exists  $\ell_0 > 0$  such that

$$\frac{\sup_{u\in\mathcal{B}}\|\tilde{K}_{\varepsilon}(u)(x)\|_{L^{\infty}(\mathbb{R}^{N})}}{\ell_{0}}<\frac{1}{2} \text{ for any } \varepsilon>0.$$

*Proof.* Let us prove that there exists  $C_0 > 0$  such that

$$\sup_{u \in \mathcal{B}} \|\tilde{K}_{\varepsilon}(u)(x)\|_{L^{\infty}(\mathbb{R}^{N})} \le C_{0}. \tag{2.9}$$

We observe that

$$|G(\varepsilon x, u)| \le C(|u|^2 + |u|^q) \text{ for all } \varepsilon > 0.$$
 (2.10)

By using (2.10), we can see that

$$|\tilde{K}_{\varepsilon}(u)(x)| = \left| \int_{\mathbb{R}^{N}} \frac{G(\varepsilon x, u)}{|x - y|^{\mu}} \, dy \right|$$

$$\leq \left| \int_{|x - y| \leq 1} \frac{G(\varepsilon x, u)}{|x - y|^{\mu}} \, dy \right| + \left| \int_{|x - y| > 1} \frac{G(\varepsilon x, u)}{|x - y|^{\mu}} \, dy \right|$$

$$\leq C \int_{|x - y| \leq 1} \frac{|u(y)|^{2} + |u(y)|^{q}}{|x - y|^{\mu}} \, dy + C \int_{\mathbb{R}^{N}} (|u|^{2} + |u|^{q}) \, dy$$

$$\leq C \int_{|x - y| \leq 1} \frac{|u(y)|^{2} + |u(y)|^{q}}{|x - y|^{\mu}} \, dy + C$$

$$(2.11)$$

where in the last line we used Theorem 2.1 and  $||u||_{\varepsilon}^2 \leq 4(\kappa+1)$ . Now, we take

$$t \in \left(\frac{N}{N-\mu}, \frac{N}{N-2s}\right] \text{ and } r \in \left(\frac{N}{N-\mu}, \frac{2N}{q(N-2s)}\right].$$

By applying Hölder inequality and by using Theorem 2.1 and  $||u||_{\varepsilon}^2 \leq 4(\kappa+1)$ , we can see that

$$\int_{|x-y| \le 1} \frac{|u(y)|^2}{|x-y|^{\mu}} dy \le \left( \int_{|x-y| \le 1} |u|^{2t} dy \right)^{\frac{1}{t}} \left( \int_{|x-y| \le 1} \frac{1}{|x-y|^{\frac{t\mu}{t-1}}} dy \right)^{\frac{t-1}{t}} \\
\le C_* (4(\kappa+1))^2 \left( \int_{\rho < 1} \rho^{N-1-\frac{t\mu}{t-1}} d\rho \right)^{\frac{t-1}{t}} < \infty.$$
(2.12)

because of  $N-1-\frac{t\mu}{t-1}>-1$ . Similarly, we get

$$\int_{|x-y| \le 1} \frac{|u(y)|^q}{|x-y|^{\mu}} dy \le \left( \int_{|x-y| \le 1} |u|^{rq} dy \right)^{\frac{1}{r}} \left( \int_{|x-y| \le 1} \frac{1}{|x-y|^{\frac{r-1}{r-1}}} dy \right)^{\frac{r-1}{r}} \\
\le C_* (4(\kappa+1))^q \left( \int_{\rho \le 1} \rho^{N-1-\frac{r\mu}{r-1}} d\rho \right)^{\frac{r-1}{r}} < \infty$$
(2.13)

in view of  $N-1-\frac{r\mu}{r-1}>-1$ . Putting together (2.12) and (2.13) we can see that

$$\int_{|x-y|<1} \frac{|u(y)|^2 + |u(y)|^q}{|x-y|^{\mu}} \, dy \le C \text{ for all } x \in \mathbb{R}^N$$

which in view of (2.11) yields (2.9). Then there exists  $\ell_0 > 0$  such that

$$\frac{\sup_{u\in\mathcal{B}}\|\tilde{K}_{\varepsilon}(u)(x)\|_{L^{\infty}(\mathbb{R}^{N})}}{\ell_{0}}\leq \frac{C_{0}}{\ell_{0}}<\frac{1}{2}.$$

Then, we take a > 0 the unique number such that

$$\frac{f(a)}{a} = \frac{V_0}{\ell_0}$$

and we consider the penalized problem (2.3) with these choices. Now, let us introduce the Nehari manifold associated to (2.3), that is

$$\mathcal{N}_{\varepsilon} := \{ u \in H_{\varepsilon}^{s} \setminus \{0\} : \langle J_{\varepsilon}'(u), u \rangle = 0 \}.$$

By using Theorem 2.2 and  $(g_1)$ - $(g_2)$ , we can see that for all  $u \in \mathcal{N}_{\varepsilon}$ 

$$||u||_{\varepsilon}^2 \le C(||u||_{\varepsilon}^4 + ||u||_{\varepsilon}^{2q}),$$

so there exists r > 0 such that

$$||u||_{\varepsilon} \ge r \text{ for all } u \in \mathcal{N}_{\varepsilon}, \varepsilon > 0.$$
 (2.14)

Let us denote by  $S_{\varepsilon}$  the unitary sphere in  $H_{\varepsilon}^{s}$ . Since f is only continuous, the next two results will play a fundamental role to overcome the non-differentiability of  $\mathcal{N}_{\varepsilon}$ .

**Lemma 2.4.** Suppose that V satisfies  $(V_1)$ - $(V_2)$  and f verifies  $(f_1)$ - $(f_4)$ . Then, the following facts hold true:

- (a) For any  $u \in H_{\varepsilon}^s \setminus \{0\}$ , let  $h_u : \mathbb{R}_+ \to \mathbb{R}$  be defined by  $h_u(t) := J_{\varepsilon}(tu)$ . Then, there is a unique  $t_u > 0$  such that  $h'_u(t) > 0$  in  $(0, t_u)$  and  $h'_u(t) < 0$  in  $(t_u, \infty)$ .
- (b) There is  $\tau > 0$ , independent on u, such that  $t_u \geq \tau$  for every  $u \in \mathbb{S}_{\varepsilon}$ . Moreover, for each compact set  $W \subset \mathbb{S}_{\varepsilon}$ , there is  $C_W > 0$  such that  $t_u \leq C_W$  for every  $u \in W$ .
- (c) The map  $\hat{m}_{\varepsilon}: H_{\varepsilon}^{s} \setminus \{0\} \to \mathcal{N}_{\varepsilon}$  given by  $\hat{m}_{\varepsilon}(u) := t_{u}u$  is continuous and  $m_{\varepsilon} := \hat{m}|_{\mathbb{S}_{\varepsilon}}$  is a homeomorphism between  $\mathbb{S}_{\varepsilon}$  and  $\mathcal{N}_{\varepsilon}$ . Moreover,  $m_{\varepsilon}^{-1}(u) = \frac{u}{\|u\|_{\varepsilon}}$ .

*Proof.* (a) From the proof of Lemma 2.2 we can see that  $h_u(0) = 0$ ,  $h_u(t) > 0$  for t small and  $h_u(t) < 0$  for t large. Then, by the continuity of  $h_u$ , it is easy to see that there exists  $t_u > 0$  such that  $\max_{t\geq 0} h_u(t) = h_u(t_u)$ ,  $t_u u \in \mathcal{N}_{\varepsilon}$  and  $h'_u(t_u) = 0$ . Now, we note that

$$tu \in \mathcal{N}_{\varepsilon} \iff ||u||_{\varepsilon}^{2} = \int_{\mathbb{R}^{N}} \left( \frac{1}{|x|^{\mu}} * \frac{G(\varepsilon x, tu)}{t} \right) g(\varepsilon x, tu) u \, dx,$$

so, by using  $(g_4)$ , we get the uniqueness of a such  $t_u$ .

(b) Let  $u \in \mathbb{S}_{\varepsilon}$ . Recalling that  $h'_u(t_u) = 0$ , and using  $(g_1)$ - $(g_2)$ , Theorem 2.2 (see estimates in Lemma ??), and Theorem 2.1, we get for any  $\xi > 0$  small

$$t_u^2 = \int_{\mathbb{R}^N} \tilde{K}_{\varepsilon}(t_u u) g(\varepsilon x, t_u u) t_u u \, dx \le \xi C_1 t_u^4 + C_2 C_{\xi} t_u^{2q}.$$

Being q > 2, there exists  $\tau > 0$  independent of u, such that  $t_u \geq \tau$ . Now, by using  $(g_3)$ , we can observe that

$$J_{\varepsilon}(v) = J_{\varepsilon}(v) - \frac{1}{4} \langle J_{\varepsilon}'(v), v \rangle$$

$$= \frac{1}{4} ||v||_{\varepsilon}^{2} - \frac{1}{4} \int_{\mathbb{R}^{N}} \left( \frac{1}{|x|^{\mu}} * G(\varepsilon x, u) \right) \left[ 2G(\varepsilon x, u) - g(\varepsilon x, u)u \right] dx$$

$$\geq \frac{1}{4} ||v||_{\varepsilon}^{2} \quad \text{for any } v \in \mathcal{N}_{\varepsilon}.$$
(2.15)

Hence, if  $W \subset \mathbb{S}_{\varepsilon}$  is a compact set, and  $(u_n) \subset W$  is such that  $t_{u_n} \to \infty$ , it follows that  $u_n \to u$  in  $H_{\varepsilon}^s$ , and  $J_{\varepsilon}(t_{u_n}u_n) \to -\infty$ . Taking  $v_n = t_{u_n}u_n \in \mathcal{N}_{\varepsilon}$  in (2.15), we can see that

$$0 < \frac{1}{4} \le \frac{J_{\varepsilon}(t_{u_n} u_n)}{t_{u_n}^2} \le 0 \text{ as } n \to \infty$$

which gives a contradiction.

(c) Since (a) and (b) hold, we can apply Proposition 8 in [39] to deduce the thesis.  $\Box$ 

**Remark 2.1.** From the estimates in (b), we can deduce that  $J_{\varepsilon}$  is coercive on  $\mathcal{N}_{\varepsilon}$ , because for all  $u \in \mathcal{N}_{\varepsilon}$ 

$$J_{\varepsilon}(u) \ge \frac{1}{4} \|u\|_{\varepsilon}^2 \to \infty \text{ as } \|u\|_{\varepsilon} \to \infty.$$

Taking into account the above estimate and (2.14), we can also see that  $J_{\varepsilon}|_{\mathcal{N}_{\varepsilon}}$  is bounded below by some positive constant.

Let us define the maps  $\hat{\psi}_{\varepsilon}: H_{\varepsilon}^{s} \setminus \{0\} \to \mathbb{R}$  by  $\hat{\psi}_{\varepsilon}(u) := J_{\varepsilon}(\hat{m}_{\varepsilon}(u))$ , and  $\psi_{\varepsilon} := \hat{\psi}|_{\mathbb{S}_{\varepsilon}}$ . The next result is a consequence of Lemma 2.4. For more details, see Proposition 9 and Corollary 10 in [39].

**Proposition 2.1.** Suppose that V satisfies  $(V_1)$ - $(V_2)$  and f verifies  $(f_1)$ - $(f_4)$ . Then, one has:

(a)  $\hat{\psi}_{\varepsilon} \in C^1(H^s_{\varepsilon} \setminus \{0\}, \mathbb{R})$  and

$$\langle \hat{\psi}'_{\varepsilon}(u), v \rangle = \frac{\|\hat{m}_{\varepsilon}(u)\|_{\varepsilon}}{\|u\|_{\varepsilon}} \langle J'_{\varepsilon}(\hat{m}_{\varepsilon}(u)), v \rangle,$$

for every  $u \in H^s_{\varepsilon} \setminus \{0\}$  and  $v \in H^s_{\varepsilon}$ ;

- (b)  $\psi_{\varepsilon} \in C^1(\mathbb{S}_{\varepsilon}, \mathbb{R})$  and  $\langle \psi'_{\varepsilon}(u), v \rangle = \|m_{\varepsilon}(u)\|_{\varepsilon} \langle J'_{\varepsilon}(m_{\varepsilon}(u)), v \rangle$ , for every  $v \in T_u \mathbb{S}_{\varepsilon}$ .
- (c) If  $(u_n)$  is a  $(PS)_d$  sequence for  $\psi_{\varepsilon}$ , then  $(m_{\varepsilon}(u_n))$  is a  $(PS)_d$  sequence for  $J_{\varepsilon}$ . Moreover, if  $(u_n) \subset \mathcal{N}_{\varepsilon}$  is a bounded  $(PS)_d$  sequence for  $J_{\varepsilon}$ , then  $(m_{\varepsilon}^{-1}(u_n))$  is a  $(PS)_d$  sequence for the functional  $\psi_{\varepsilon}$ ;
- (d) u is a critical point of  $\psi_{\varepsilon}$  if and only if  $m_{\varepsilon}(u)$  is a nontrivial critical point for  $J_{\varepsilon}$ . Moreover, the corresponding critical values coincide and

$$\inf_{u \in \mathbb{S}_{\varepsilon}} \psi_{\varepsilon}(u) = \inf_{u \in \mathcal{N}_{\varepsilon}} J_{\varepsilon}(u).$$

**Remark 2.2.** As in [39], we have the following characterization of the infimum of  $J_{\varepsilon}$  on  $\mathcal{N}_{\varepsilon}$ :

$$c_{\varepsilon} = \inf_{u \in \mathcal{N}_{\varepsilon}} J_{\varepsilon}(u) = \inf_{u \in H_{\varepsilon}^{s} \setminus \{0\}} \max_{t > 0} J_{\varepsilon}(tu) = \inf_{u \in \mathbb{S}_{\varepsilon}} \max_{t > 0} J_{\varepsilon}(tu).$$

In the next result we show that  $J_{\varepsilon}$  verifies a local compactness condition.

**Lemma 2.5.**  $J_{\varepsilon}$  satisfies the  $(PS)_c$  condition for all  $c \in [c_{\varepsilon}, \kappa]$ .

*Proof.* Let  $(u_n)$  be a Palais-Smale sequence at the level c, that is  $J_{\varepsilon}(u_n) \to c$  and  $J'_{\varepsilon}(u_n) \to 0$ . We divide the proof in two main steps.

**Step** 1: For any  $\eta > 0$  there exists  $R = R_{\eta} > 0$  such that

$$\limsup_{n \to \infty} \int_{\mathbb{R}^N \setminus B_R} |(-\Delta)^{\frac{s}{2}} u_n|^2 + V(\varepsilon x) u_n^2 \, dx < \eta. \tag{2.16}$$

By using  $(g_3)$ , we can see that

$$J_{\varepsilon}(u_n) - \frac{1}{4} \langle J_{\varepsilon}'(u_n), u_n \rangle = \left(\frac{1}{2} - \frac{1}{4}\right) \|u_n\|_{H_{\varepsilon}^s}^2 + \frac{1}{4} \int_{\mathbb{R}^N} \left(\frac{1}{|x|^{\mu}} * G(\varepsilon x, u_n)\right) g(\varepsilon x, u_n) u_n dx$$
$$- \frac{1}{2} \int_{\mathbb{R}^N} \left(\frac{1}{|x|^{\mu}} * G(\varepsilon x, u_n)\right) G(\varepsilon x, u_n) dx$$
$$\geq \frac{1}{4} \|u_n\|_{\varepsilon}^2,$$

so there exists  $n_0 \in \mathbb{N}$  such that

$$||u_n||_{\varepsilon}^2 \leq 4(\kappa+1)$$
 for all  $n \geq n_0$ .

Therefore, we may assume that  $u_n \rightharpoonup u$  in  $H^s(\mathbb{R}^N)$  and  $u_n \rightarrow u$  in  $L^r_{loc}(\mathbb{R}^N)$  for any  $r \in [2, 2_s^*)$ . Moreover, by Lemma 2.3, we can deduce that

$$\frac{\sup_{n\geq n_0} \|K_{\varepsilon}(u_n)(x)\|_{L^{\infty}(\mathbb{R}^N)}}{\ell_0} \leq \frac{1}{2}.$$
(2.17)

Fix R > 0 and let  $\psi_R \in C^{\infty}(\mathbb{R}^N)$  be a function such that  $\psi_R = 0$  in  $B_{R/2}$ ,  $\psi_R = 1$  in  $B_R^c$ ,  $\psi_R \in [0,1]$  and  $|\nabla \eta_R| \leq C/R$ . Since  $(u_n \psi_R)$  is bounded we can see that

$$\int_{\mathbb{R}^{N}} (-\Delta)^{\frac{s}{2}} u_{n}(-\Delta)^{\frac{s}{2}} (u_{n}\psi_{R}) + V(\varepsilon x) \psi_{R} u_{n}^{2} dx = \langle J_{\varepsilon}'(u_{n}), u_{n}\psi_{R} \rangle + \int_{\mathbb{R}^{N}} \left( \frac{1}{|x|^{\mu}} * G(\varepsilon x, u_{n}) \right) g(\varepsilon x, u_{n}) u_{n}\psi_{R} dx$$

$$= o_{n}(1) + \int_{\mathbb{R}^{N}} \left( \frac{1}{|x|^{\mu}} * G(\varepsilon x, u_{n}) \right) g(\varepsilon x, u_{n}) u_{n}\psi_{R} dx$$
(2.18)

For  $n \ge n_0$  and  $\varepsilon > 0$  fixed, take R > 0 big enough such that  $\Lambda_{\varepsilon} \subset B_{R/2}$ . Then, by using  $(g_3)$  with  $\ell_0$  as in Lemma 2.3, we deduce that

$$\begin{split} \int_{\mathbb{R}^{N}\backslash B_{R/2}} |(-\Delta)^{\frac{s}{2}} u_{n}|^{2} + V(\varepsilon x) u_{n}^{2} \, dx &\leq \int_{\mathbb{R}^{N}\backslash B_{R/2}} \left(\frac{1}{|x|^{\mu}} * G(\varepsilon x, u_{n})\right) g(\varepsilon x, u_{n}) u_{n} \, dx + o_{n}(1) \\ &- \iint_{\mathbb{R}^{2N}} \frac{(u_{n}(x) - u_{n}(u))(\psi_{R}(x) - \psi_{R}(y))}{|x - y|^{N + 2s}} u_{n}(y) \, dx dy \\ &\leq \int_{\mathbb{R}^{N}\backslash B_{R/2}} \frac{\sup_{n \geq n_{0}} \|\tilde{K}_{\varepsilon}(u_{n})(x)\|_{L^{\infty}(\mathbb{R}^{N})}}{\ell_{0}} V_{0} u_{n}^{2} \, dx + o_{n}(1) \\ &- \iint_{\mathbb{R}^{2N}} \frac{(u_{n}(x) - u_{n}(u))(\psi_{R}(x) - \psi_{R}(y))}{|x - y|^{N + 2s}} u_{n}(y) \, dx dy, \end{split}$$

which together with (2.17) yields

$$\frac{1}{2} \int_{\mathbb{R}^{N} \setminus B_{R/2}} |(-\Delta)^{\frac{s}{2}} u_{n}|^{2} + V(\varepsilon x) u_{n}^{2} dx \le o_{n}(1) - \iint_{\mathbb{R}^{2N}} \frac{(u_{n}(x) - u_{n}(u))(\psi_{R}(x) - \psi_{R}(y))}{|x - y|^{N + 2s}} u_{n}(y) dx dy.$$

Now, we note that the Hölder inequality and the boundedness of  $(u_n)$  imply that

$$\left| \iint_{\mathbb{R}^{2N}} \frac{(u_n(x) - u_n(u))(\psi_R(x) - \psi_R(y))}{|x - y|^{N+2s}} u_n(y) \, dx dy \right|$$

$$\leq \left( \iint_{\mathbb{R}^{2N}} \frac{|u_n(x) - u_n(y)|^2}{|x - y|^{N+2s}} \, dx dy \right)^{\frac{1}{2}} \left( \iint_{\mathbb{R}^{2N}} \frac{|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} u_n^2(y) \, dx dy \right)^{\frac{1}{2}}$$

$$\leq C \left( \iint_{\mathbb{R}^{2N}} \frac{|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} u_n^2(y) \, dx dy \right)^{\frac{1}{2}}.$$

Therefore, it is enough to prove that

$$\lim_{R \to \infty} \limsup_{n \to \infty} \iint_{\mathbb{R}^{2N}} \frac{|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N + 2s}} u_n^2(y) \, dx dy = 0$$

to conclude our first claim.

Let us note that  $\mathbb{R}^{2N}$  can be written as

$$\mathbb{R}^{2N} = ((\mathbb{R}^N \setminus B_{2R}) \times (\mathbb{R}^N \setminus B_{2R})) \cup ((\mathbb{R}^N \setminus B_{2R}) \times B_{2R}) \cup (B_{2R} \times \mathbb{R}^N) =: X_R^1 \cup X_R^2 \cup X_R^3.$$

Then

$$\iint_{\mathbb{R}^{2N}} \frac{|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} u_n^2(x) dx dy = \iint_{X_R^1} \frac{|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} u_n^2(x) dx dy 
+ \iint_{X_R^2} \frac{|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} u_n^2(x) dx dy + \iint_{X_R^3} \frac{|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} u_n^2(x) dx dy.$$
(2.19)

Now, we estimate each integral in (2.19). Since  $\psi_R = 1$  in  $\mathbb{R}^N \setminus B_{2R}$ , we have

$$\iint_{X_R^1} \frac{u_n^2(x)|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} dx dy = 0.$$
 (2.20)

Let k > 4. Clearly, we have

$$X_R^2 = (\mathbb{R}^N \setminus B_{2R}) \times B_{2R} = ((\mathbb{R}^{2N} \setminus B_{kR}) \times B_{2R}) \cup ((B_{kR} \setminus B_{2R}) \times B_{2R})$$

Let us observe that, if  $(x,y) \in (\mathbb{R}^{2N} \setminus B_{kR}) \times B_{2R}$ , then

$$|x - y| \ge |x| - |y| \ge |x| - 2R > \frac{|x|}{2}.$$

Therefore, taking into account  $0 \le \psi_R \le 1$ ,  $|\nabla \psi_R| \le \frac{C}{R}$  and applying Hölder inequality, we can see

$$\iint_{X_R^2} \frac{u_n^2(x)|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} dxdy 
= \int_{\mathbb{R}^{2N} \backslash B_{kR}} \int_{B_{2R}} \frac{u_n^2(x)|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} dxdy + \int_{B_{kR} \backslash B_{2R}} \int_{B_{2R}} \frac{u_n^2(x)|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} dxdy 
\leq 2^{2+N+2s} \int_{\mathbb{R}^N \backslash B_{kR}} \int_{B_{2R}} \frac{u_n^2(x)}{|x|^{N+2s}} dxdy + \frac{C}{R^2} \int_{B_{kR} \backslash B_{2R}} \int_{B_{2R}} \frac{u_n^2(x)}{|x - y|^{N+2(s-1)}} dxdy 
\leq CR^N \int_{\mathbb{R}^N \backslash B_{kR}} \frac{u_n^2(x)}{|x|^{N+2s}} dx + \frac{C}{R^2} (kR)^{2(1-s)} \int_{B_{kR} \backslash B_{2R}} u_n^2(x)dx 
\leq CR^N \left( \int_{\mathbb{R}^N \backslash B_{kR}} |u_n(x)|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} \left( \int_{\mathbb{R}^N \backslash B_{kR}} \frac{1}{|x|^{\frac{N^2}{2s} + N}} dx \right)^{\frac{2s}{N}} + \frac{Ck^{2(1-s)}}{R^{2s}} \int_{B_{kR} \backslash B_{2R}} u_n^2(x)dx 
\leq \frac{C}{k^N} \left( \int_{\mathbb{R}^N \backslash B_{kR}} |u_n(x)|^{2_s^*} dx \right)^{\frac{2}{s^*}} + \frac{Ck^{2(1-s)}}{R^{2s}} \int_{B_{kR} \backslash B_{2R}} u_n^2(x)dx 
\leq \frac{C}{k^N} + \frac{Ck^{2(1-s)}}{R^{2s}} \int_{B_{kR} \backslash B_{2R}} u_n^2(x)dx. \tag{2.21}$$

Now, fix  $\varepsilon \in (0,1)$ , and we note that

$$\iint_{X_R^3} \frac{u_n^2(x)|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} dxdy$$

$$\leq \int_{B_{2R} \setminus B_{\varepsilon R}} \int_{\mathbb{R}^N} \frac{u_n^2(x)|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} dxdy + \int_{B_{\varepsilon R}} \int_{\mathbb{R}^N} \frac{u_n^2(x)|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} dxdy. \quad (2.22)$$

Let us estimate the first integral in (2.22). Then.

$$\int_{B_{2R}\setminus B_{\varepsilon R}} \int_{\mathbb{R}^N \cap \{y: |x-y| < R\}} \frac{u_n^2(x) |\psi_R(x) - \psi_R(y)|^2}{|x-y|^{N+2s}} \, dx dy \le \frac{C}{R^{2s}} \int_{B_{2R}\setminus B_{\varepsilon R}} u_n^2(x) dx$$

and

$$\int_{B_{2R} \setminus B_{\varepsilon R}} \int_{\mathbb{R}^N \cap \{y: |x-y| \ge R\}} \frac{u_n^2(x) |\psi_R(x) - \psi_R(y)|^2}{|x-y|^{N+2s}} \, dx dy \le \frac{C}{R^{2s}} \int_{B_{2R} \setminus B_{\varepsilon R}} u_n^2(x) dx$$

from which we have

$$\int_{B_{2R}\setminus B_{\varepsilon R}} \int_{\mathbb{R}^N} \frac{u_n^2(x)|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} dx dy \le \frac{C}{R^{2s}} \int_{B_{2R}\setminus B_{\varepsilon R}} u_n^2(x) dx. \tag{2.23}$$

By using the definition of  $\psi_R$ ,  $\varepsilon \in (0,1)$ , and  $\psi_R \leq 1$ , we have

$$\int_{B_{\varepsilon R}} \int_{\mathbb{R}^{N}} \frac{u_{n}^{2}(x)|\psi_{R}(x) - \psi_{R}(y)|^{2}}{|x - y|^{N + 2s}} dxdy = \int_{B_{\varepsilon R}} \int_{\mathbb{R}^{N} \setminus B_{R}} \frac{|u_{n}(x)|^{2}|\psi_{R}(x) - \psi_{R}(y)|^{2}}{|x - y|^{N + 2s}} dxdy$$

$$\leq 4 \int_{B_{\varepsilon R}} \int_{\mathbb{R}^{N} \setminus B_{R}} \frac{u_{n}^{2}(x)}{|x - y|^{N + 2s}} dxdy$$

$$\leq C \int_{B_{\varepsilon R}} u_{n}^{2}(x) dx \int_{(1 - \varepsilon)R}^{\infty} \frac{1}{r^{1 + 2s}} dr$$

$$= \frac{C}{[(1 - \varepsilon)R]^{2s}} \int_{B_{\varepsilon R}} u_{n}^{2}(x) dx \tag{2.24}$$

where we use the fact that if  $(x, y) \in B_{\varepsilon R} \times (\mathbb{R}^N \setminus B_R)$ , then  $|x - y| > (1 - \varepsilon)R$ . Taking into account (2.22), (2.23) and (2.24) we deduce

$$\iint_{X_R^3} \frac{u_n^2(x)|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} dxdy$$

$$\leq \frac{C}{R^{2s}} \int_{B_{2R} \setminus B_{\varepsilon R}} |u_n(x)|^2 dx + \frac{C}{[(1 - \varepsilon)R]^{2s}} \int_{B_{\varepsilon R}} u_n^2(x) dx. \tag{2.25}$$

Putting together (2.19), (2.20), (2.21) and (2.25), we can infer

$$\iint_{\mathbb{R}^{2N}} \frac{u_n^2(x)|\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N+2s}} dxdy$$

$$\leq \frac{C}{k^N} + \frac{Ck^{2(1-s)}}{R^{2s}} \int_{B_{kR} \setminus B_{2R}} u_n^2(x)dx + \frac{C}{R^{2s}} \int_{B_{2R} \setminus B_{\varepsilon R}} |u_n(x)|^2 dx + \frac{C}{[(1-\varepsilon)R]^{2s}} \int_{B_{\varepsilon R}} u_n^2(x)dx. \tag{2.26}$$

Since  $(u_n)$  is bounded in  $H^s(\mathbb{R}^N)$ , we may assume that  $u_n \to u$  in  $L^2_{loc}(\mathbb{R}^N)$  for some  $u \in H^s(\mathbb{R}^N)$ . Then, taking the limit as  $n \to \infty$  in (2.26), we have

$$\begin{split} & \limsup_{n \to \infty} \iint_{\mathbb{R}^{2N}} \frac{|u_n(x)|^2 |\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N + 2s}} \, dx dy \\ & \leq \frac{C}{k^N} + \frac{Ck^{2(1 - s)}}{R^{2s}} \int_{B_{kR} \backslash B_{2R}} |u(x)|^2 dx + \frac{C}{R^{2s}} \int_{B_{2R} \backslash B_{\varepsilon R}} |u(x)|^2 dx + \frac{C}{[(1 - \varepsilon)R]^{2s}} \int_{B_{\varepsilon R}} |u(x)|^2 dx \\ & \leq \frac{C}{k^N} + Ck^2 \left( \int_{B_{kR} \backslash B_{2R}} |u(x)|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} + C \left( \int_{B_{2R} \backslash B_{\varepsilon R}} |u(x)|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} + C \left( \frac{\varepsilon}{1 - \varepsilon} \right)^{2s} \left( \int_{B_{\varepsilon R}} |u(x)|^{2_s^*} dx \right)^{\frac{2}{2_s^*}}, \end{split}$$

where in the last passage we use Hölder inequality. Since  $u \in L^{2_s^*}(\mathbb{R}^N)$ , k > 4 and  $\varepsilon \in (0,1)$ , we obtain

$$\limsup_{R \to \infty} \int_{B_{kR} \setminus B_{2R}} |u(x)|^{2_s^*} dx = \limsup_{R \to \infty} \int_{B_{2R} \setminus B_{\varepsilon R}} |u(x)|^{2_s^*} dx = 0.$$

Choosing  $\varepsilon = \frac{1}{k}$ , we get

$$\begin{split} & \limsup_{R \to \infty} \limsup_{n \to \infty} \iint_{\mathbb{R}^{2N}} \frac{u_n^2(x) |\psi_R(x) - \psi_R(y)|^2}{|x - y|^{N + 2s}} \, dx dy \\ & \leq \lim_{k \to \infty} \limsup_{R \to \infty} \Big[ \frac{C}{k^N} + C k^2 \left( \int_{B_{kR} \backslash B_{2R}} |u(x)|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} + C \left( \int_{B_{2R} \backslash B_{\frac{1}{k}R}} |u(x)|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} \\ & + C \left( \frac{1}{k - 1} \right)^{2s} \left( \int_{B_{\frac{1}{k}R}} |u(x)|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} \Big] \\ & \leq \lim_{k \to \infty} \frac{C}{k^N} + C \left( \frac{1}{k - 1} \right)^{2s} \left( \int_{\mathbb{R}^N} |u(x)|^{2_s^*} dx \right)^{\frac{2}{2_s^*}} = 0. \end{split}$$

**Step** 2: Let us prove that  $u_n \to u$  in  $H^s_{\varepsilon}$  as  $n \to \infty$ . Set  $\Psi_n = ||u_n - u||^2_{\varepsilon}$  and we observe that

$$\Psi_n = \langle J_{\varepsilon}'(u_n), u_n \rangle - \langle J_{\varepsilon}'(u_n), u \rangle + \int_{\mathbb{R}^N} \left( \frac{1}{|x|^{\mu}} * G(\varepsilon x, u_n) \right) g(\varepsilon x, u_n) (u_n - u) \, dx + o_n(1). \tag{2.27}$$

Let us note that  $\langle J'_{\varepsilon}(u_n), u_n \rangle = \langle J'_{\varepsilon}(u_n), u \rangle = o_n(1)$ , so in view of (2.27), we need to show that

$$\int_{\mathbb{R}^N} \left( \frac{1}{|x|^{\mu}} * G(\varepsilon x, u_n) \right) g(\varepsilon x, u_n) (u_n - u) dx = o_n(1),$$

to infer that  $\Psi_n \to 0$  as  $n \to \infty$ .

We observe that  $G(\varepsilon x, u_n)$  is bounded in  $L^{\frac{2N}{2N-\mu}}(\mathbb{R}^N)$  (since  $q < \frac{2^*}{2}(2-\frac{\mu}{N})$ ),  $u_n \to u$  a.e. in  $\mathbb{R}^N$ , and G is continuous, so we deduce that

$$G(\varepsilon x, u_n) \rightharpoonup G(\varepsilon x, u) \text{ in } L^{\frac{2N}{2N-\mu}}(\mathbb{R}^N).$$
 (2.28)

In virtue of Theorem 2.2, we know that the convolution term

$$\frac{1}{|x|^{\mu}} * h(x) \in L^{\frac{2N}{\mu}}(\mathbb{R}^N) \text{ for all } h \in L^{\frac{2N}{2N-\mu}}(\mathbb{R}^N)$$

is a linear bounded operator from  $L^{\frac{2N}{2N-\mu}}(\mathbb{R}^N)$  to  $L^{\frac{2N}{\mu}}(\mathbb{R}^N)$ , so we can see that

$$\tilde{K}_{\varepsilon}(u_n) = \frac{1}{|x|^{\mu}} * G(\varepsilon x, u_n) \rightharpoonup \frac{1}{|x|^{\mu}} * G(\varepsilon x, u) = \tilde{K}_{\varepsilon}(u) \text{ in } L^{\frac{2N}{\mu}}(\mathbb{R}^N).$$
 (2.29)

Since g has a subcritical growth, by using Theorem 2.1 and (2.29), we obtain

$$\lim_{n \to \infty} \int_{B_R} \tilde{K}_{\varepsilon}(u_n) g(\varepsilon x, u_n) (u_n - u) \, dx = 0.$$
 (2.30)

From the growth assumption and the boundedness of  $K_{\varepsilon}(u_n)$  we obtain

$$\int_{\mathbb{R}^N \setminus B_R} \tilde{K}_{\varepsilon}(u_n) |g(\varepsilon x, u_n) u_n| \, dx \le C_1 \int_{\mathbb{R}^N \setminus B_R} u_n^2 \, dx.$$

By the Step 1 and Theorem 2.1, for any  $\eta > 0$  there exists  $R_{\eta} > 0$  such that

$$\limsup_{n \to \infty} \int_{\mathbb{R}^N \setminus B_R} \tilde{K}_{\varepsilon}(u_n) |g(\varepsilon x, u_n) u_n| \, dx \le C_2 \eta.$$

In similar way, from Hölder inequality, we can see that

$$\limsup_{n\to\infty} \int_{\mathbb{R}^N\setminus B_R} \tilde{K}_{\varepsilon}(u_n) |g(\varepsilon x, u_n)u| \, dx \le C_3 \eta.$$

Taking into account the above limits we can infer that

$$\lim_{n \to \infty} \int_{\mathbb{R}^N} \tilde{K}_{\varepsilon}(u_n) g(\varepsilon x, u_n) (u_n - u) \, dx = 0.$$

Finally, we prove the following result:

**Lemma 2.6.** The functional  $\psi_{\varepsilon}$  satisfies the  $(PS)_c$  on  $\mathbb{S}_{\varepsilon}$  for any  $c \in [c_{\varepsilon}, \kappa]$ .

Proof. Let  $(u_n) \subset \mathbb{S}_{\varepsilon}$  be a  $(PS)_c$  sequence for  $\psi_{\varepsilon}$ . Then  $\psi_{\varepsilon}(u_n) \to c$  and  $\|\psi'_{\varepsilon}(u_n)\|_* \to 0$ , where  $\|\cdot\|_*$  denotes the norm in the dual space of  $(T_{u_n}\mathbb{S}_{\varepsilon})^*$ . By using Proposition 2.1-(c), we can infer that  $(m_{\varepsilon}(u_n))$  is a  $(PS)_c$  sequence for  $J_{\varepsilon}$ . In view of Lemma 2.5, we can see that, up to a subsequence, there exists  $u \in \mathbb{S}_{\varepsilon}$  such that  $m_{\varepsilon}(u_n) \to m_{\varepsilon}(u)$  in  $H^s_{\varepsilon}$ . From Lemma 2.4-(c), we conclude that  $u_n \to u$  in  $\mathbb{S}_{\varepsilon}$ .

## 3. The autonomous problem

In this section we deal with the limit problem associated to (2.2), namely

$$(-\Delta)^s u + V_0 u = \left(\frac{1}{|x|^{\mu}} * F(u)\right) f(u) \text{ in } \mathbb{R}^N.$$
(3.1)

In what follows, we denote the above problem with  $(P_{V_0})$ .

The functional  $J_{V_0}: H_0^s \to \mathbb{R}$  associated to the above problem is given by

$$J_{V_0}(u) = \frac{1}{2} ||u||_{V_0}^2 - \Sigma_0(u),$$

where  $H_0^s$  is the space  $H^s(\mathbb{R}^N)$  endowed with the norm

$$||u||_{V_0}^2 = [u]^2 + \int_{\mathbb{R}^N} V_0 u^2 dx,$$

and

$$\Sigma_0(u) = \frac{1}{2} \int_{\mathbb{R}^N} \left( \frac{1}{|x|^{\mu}} * F(u) \right) F(u) dx.$$

Let us consider the following Nehari manifold

$$\mathcal{N}_{V_0} = \{ u \in H_0^s \setminus \{0\} : \langle J'_{V_0}(u), u \rangle = 0 \}$$

and let us denote by  $\mathbb{S}_0$  the unit sphere in  $H_0^s$ . Arguing as in the proofs of Lemma 2.4 and Proposition 2.1, we can see that the following results hold.

**Lemma 3.1.** Suppose that f verifies  $(f_1)$ - $(f_4)$ .

Then, the following facts hold true:

- (a) For any  $u \in H_0^s \setminus \{0\}$ , let  $h_u : \mathbb{R}_+ \to \mathbb{R}$  be defined by  $h_u(t) := J_{V_0}(tu)$ . Then, there is a unique  $t_u > 0$  such that  $h'_u(t) > 0$  in  $(0, t_u)$  and  $h'_u(t) < 0$  in  $(t_u, +\infty)$ .
- (b) There is  $\tau > 0$ , independent on u, such that  $t_u \geq \tau$  for every  $u \in \mathbb{S}_0$ . Moreover, for each compact set  $\mathcal{W} \subset \mathbb{S}_0$ , there is  $C_{\mathcal{W}} > 0$  such that  $t_u \leq C_{\mathcal{W}}$  for every  $u \in \mathcal{W}$ .
- (c) The map  $\hat{m}_0: H_0^s \setminus \{0\} \to \mathcal{N}_{V_0}$  given by  $\hat{m}_0(u) := t_u u$  is continuous and  $m_0 := \hat{m}|_{\mathbb{S}_0}$  is a homeomorphism between  $\mathbb{S}_0$  and  $\mathcal{N}_{V_0}$ . Moreover,  $m_0^{-1}(u) = \frac{u}{\|u\|_{V_0}}$ .

Let us define the maps  $\hat{\psi}_0: H_0^s \setminus \{0\} \to \mathbb{R}$  by  $\hat{\psi}_0(u) := J_{V_0}(\hat{m}_0(u))$ , and  $\psi := \hat{\psi}_0|_{\mathbb{S}_0}$ . Then we have

**Proposition 3.1.** Suppose that f verifies  $(f_1)$ - $(f_4)$ . Then, one has:

(a)  $\hat{\psi}_0 \in C^1(H_0^s \setminus \{0\}, \mathbb{R})$  and

$$\langle \hat{\psi}_0'(u), v \rangle = \frac{\|\hat{m}_0(u)\|_{V_0}}{\|u\|_{V_0}} \langle J_{\varepsilon}'(\hat{m}_0(u)), v \rangle,$$

for every  $u \in H_0^s \setminus \{0\}$  and  $v \in H_0^s$ ;

- (b)  $\psi_0 \in C^1(\mathbb{S}_0, \mathbb{R}) \text{ and } \langle \psi_0'(u), v \rangle = \|m_0(u)\|_{V_0} \langle J_{V_0}'(m_0(u)), v \rangle, \text{ for every } v \in T_u \mathbb{S}_0.$
- (c) If  $(u_n)$  is a  $(PS)_d$  sequence for  $\psi_0$ , then  $(m_0(u_n))$  is a  $(PS)_d$  sequence for  $J_{V_0}$ . Moreover, if  $(u_n) \subset \mathcal{N}_{V_0}$  is a bounded  $(PS)_d$  sequence for  $J_{V_0}$ , then  $(m_0^{-1}(u_n))$  is a  $(PS)_d$  sequence for the functional  $\psi_0$ ;
- (d) u is a critical point of  $\psi_0$  if and only if  $m_0(u)$  is a nontrivial critical point for  $J_{V_0}$ . Moreover, the corresponding critical values coincide and

$$\inf_{u \in \mathbb{S}_0} \psi_0(u) = \inf_{u \in \mathcal{N}_{V_0}} J_{V_0}(u).$$

Moreover, we have the following characterization of the infimum of  $J_0$  on  $\mathcal{N}_{V_0}$ 

$$c_{V_0} = \inf_{u \in \mathcal{N}_{V_0}} J_{V_0}(u) = \inf_{u \in H_0^s \setminus \{0\}} \max_{t > 0} J_{V_0}(tu) = \inf_{u \in \mathbb{S}_0} \max_{t > 0} J_{V_0}(tu). \tag{3.2}$$

The next Lemma allows us to assume that the weak limit of a  $(PS)_c$  sequence is nontrivial.

**Lemma 3.2.** Let  $(u_n) \subset H_0^s$  be a  $(PS)_c$  sequence for  $J_{V_0}$  and such that  $u_n \rightharpoonup 0$ . Then, only one of the following alternatives holds.

- (a)  $u_n \to 0$  in  $H_0^s$ , or
- (b) there exists a sequence  $(\tilde{y}_n) \subset \mathbb{R}^N$ , and constants R > 0 and  $\gamma > 0$  such that

$$\liminf_{n \to \infty} \int_{B_R(\tilde{y}_n)} |u_n|^2 \, dx \ge \gamma > 0.$$

*Proof.* Suppose that (b) does not hold. Then, for all R > 0, we have

$$\lim_{n \to \infty} \sup_{y \in \mathbb{R}^N} \int_{B_R(y)} |u_n|^2 dx = 0.$$

Since we know that  $(u_n)$  is bounded in  $H_0^s$ , we can use Lemma 2.1 to deduce that  $u_n \to 0$  in  $L^q(\mathbb{R}^N)$  for any  $q \in (2, 2_s^*)$ . By using  $(f_1)$ - $(f_2)$ , we know that for all  $\eta > 0$  there exists  $C_{\eta} > 0$  such that

$$|F(t)| \le \eta |t|^2 + C_{\eta} |t|^q,$$

so, applying Hardy-Littlewood-Sobolev inequality, we get

$$\int_{\mathbb{R}^N} \left( \frac{1}{|x|^{\mu}} * F(u_n) \right) f(u_n) u_n \, dx = o_n(1).$$

Taking into account  $\langle J_0'(u_n), u_n \rangle = o_n(1)$ , we can infer that  $||u_n||_{V_0} \to 0$  as  $n \to \infty$ .

Now, we prove the following result for the autonomous problem.

**Lemma 3.3.** Let  $(w_n) \subset H_0^s$  be a  $(PS)_{c_{V_0}}$  sequence for  $J_{V_0}$ . Then the problem  $(P_{V_0})$  has a positive ground state.

*Proof.* Arguing as in Lemma 2.2, we can see that  $J_0$  has a mountain pass geometry. As a consequence of the mountain pass theorem without the (PS) condition (see [42]), there exists a Palais-Smale sequence  $(u_n) \subset H_0^s$  such that

$$J_{V_0}(u_n) \to c_{V_0}$$
 and  $J'_{V_0}(u_n) \to 0$ .

Since

$$J_{V_0}(u_n) - \frac{1}{4} \langle J'_{V_0}(u_n), u_n \rangle \ge \frac{1}{4} ||u_n||_{V_0}^2,$$

it is easy to deduce that  $(u_n)$  is bounded in  $H_0^s$ . By using  $(f_1)$ - $(f_2)$ , we know that  $||u||_{V_0} \ge r$  for all  $u \in \mathcal{N}_{V_0}$ . Then, arguing as in the proof of Lemma 3.2, we can see that there exists a sequence  $(y_n) \subset \mathbb{R}^N$ , and constants R > 0 and  $\gamma > 0$  such that

$$\liminf_{n \to \infty} \int_{B_R(y_n)} |u_n|^2 \, dx \ge \gamma > 0.$$

Set  $v_n = u_n(\cdot - y_n)$ . Since  $J_{V_0}$  and  $J'_{V_0}$  are both invariant by translation, it holds that

$$J'_{V_0}(v_n) \to 0 \text{ and } J_{V_0}(v_n) \to c_{V_0}.$$

We observe that  $(v_n)$  is also bounded in  $H_0^s$ , so we may assume that  $v_n \to v$  in  $H_0^s$ , for some  $v \neq 0$ . Now, we show that v is a weak solution to  $(P_{V_0})$ . Fix  $\varphi \in C_0^{\infty}(\mathbb{R}^N)$ . Recalling that  $(v_n)$  is bounded, we can argue as in the proof of Lemma 2.3 to deduce that  $\|\frac{1}{|x|^{\mu}} * F(v_n)\|_{L^{\infty}(\mathbb{R}^N)} \leq C$  for any  $n \in \mathbb{N}$ .

Then, using the fact that f has subcritical growth and  $v_n \to v$  in  $L^r_{loc}(\mathbb{R}^N)$  for any  $r \in [1, 2_s^*)$ , we can see that the Dominated Convergence Theorem gives

$$\int_{\mathbb{R}^N} \left( \frac{1}{|x|^\mu} * F(v_n) \right) f(v_n) \varphi \, dx \to \int_{\mathbb{R}^N} \left( \frac{1}{|x|^\mu} * F(v) \right) f(v) \varphi \, dx.$$

This combined with the weak convergence of  $(v_n)$  yields

$$o_n(1) = \langle J'_{V_0}(v_n), \varphi \rangle \to \langle J'_{V_0}(v), \varphi \rangle.$$

From the density of  $C_0^{\infty}(\mathbb{R}^N)$  in  $H_0^s$ , we get  $\langle J'_{V_0}(v), \varphi \rangle = 0$  for all  $\varphi \in H_0^s$ . In particular,  $v \in \mathcal{N}_{V_0}$ . Using the definition of  $c_{V_0}$  together with Fatou's Lemma, we also deduce that  $J_{V_0}(v) = c_{V_0}$ .

Now, recalling that f(t) = 0 for  $t \le 0$  and  $(x - y)(x^- - y^-) \ge |x^- - y^-|^2$  for all  $x, y \in \mathbb{R}$ , it is easy to deduce that  $\langle J'_{V_0}(v), v^- \rangle = 0$  implies that  $v \ge 0$  in  $\mathbb{R}^N$ .

Proceeding as in the proof of Lemma 2.3, we can see that  $K(x) := \frac{1}{|x|^{\mu}} * F(v)$  is bounded in  $\mathbb{R}^N$ , so similar arguments developed in Lemma 5.1 below, allow us to deduce that  $v \in L^{\infty}(\mathbb{R}^N)$ . Since f has subcritical growth and K(x) is bounded, we can see that  $K(x)f(v) \in L^{\infty}(\mathbb{R}^N)$ , so we can apply Proposition 2.9 in [38] to infer that  $v \in C^{0,\alpha}(\mathbb{R}^N)$  for some  $\alpha \in (0,1)$ . Using the Harnack inequality [38], we can conclude that v > 0 in  $\mathbb{R}^N$ .

The next result is a compactness result on autonomous problem which we will use later.

**Lemma 3.4.** Let  $(\tilde{v}_n) \subset \mathcal{N}_{V_0}$  be such that  $J_0(\tilde{v}_n) \to c_{V_0}$ . Then  $(\tilde{v}_n)$  has a convergent subsequence in  $H_0^s$ .

*Proof.* Since  $(\tilde{v}_n) \subset \mathcal{N}_{V_0}$  and  $J_{V_0}(\tilde{v}_n) \to c_{V_0}$ , we can apply Lemma 3.1-(c) and Proposition 3.1-(d) to infer that

$$w_n = m_0^{-1}(\tilde{v}_n) = \frac{\tilde{v}_n}{\|\tilde{v}_n\|_{V_0}} \in \mathbb{S}_0$$

and

$$\psi_0(w_n) = J_{V_0}(\tilde{v}_n) \to c_{V_0} = \inf_{v \in S_0} \psi_0(v).$$

Hence, by using the Ekeland's variational principle [17], we can find  $(\tilde{w}_n) \subset \mathbb{S}_0$  such that  $(\tilde{w}_n)$  is a  $(PS)_{c_{V_0}}$  sequence for  $\psi_0$  on  $\mathbb{S}_0$  and  $\|\tilde{w}_n - w_n\|_{V_0} = o_n(1)$ . From Proposition 3.1-(c), we can deduce that  $m_0(\tilde{w}_n)$  is a  $(PS)_{c_{V_0}}$  sequence of  $J_0$ . By applying Lemma 3.3, it follows that there exists  $\tilde{w} \in \mathbb{S}_0$  such that  $m_0(\tilde{w}_n) \to m_0(\tilde{w})$  in  $H_0^s$ . This fact, together with Lemma 3.1-(c), and  $\|\tilde{w}_n - w_n\|_{V_0} = o_n(1)$ , allow us to conclude that  $\tilde{v}_n \to \tilde{v}$  in  $H_0^s$ .

## 4. Multiplicity results

In order to study the multiplicity of solutions to (1.1), we need introduce some useful tools. Let us consider  $\delta > 0$  such that  $M_{\delta} \subset \Lambda$ , where

$$M_{\delta} = \{ x \in \mathbb{R}^N : dist(x, M) \le \delta \}.$$

and  $\eta \in C_0^{\infty}(\mathbb{R}_+, [0, 1])$  satisfying  $\eta(t) = 1$  if  $0 \le t \le \frac{\delta}{2}$  and  $\eta(t) = 0$  if  $t \ge \delta$ . For any  $y \in M$ , we define

$$\Psi_{\varepsilon,y}(x) = \eta(|\varepsilon x - y|)w\left(\frac{\varepsilon x - y}{\varepsilon}\right)$$

where w is a positive ground state solution for  $J_{V_0}$  (by Lemma 3.3). Let us denote by  $t_{\varepsilon} > 0$  the unique positive number verifying

$$\max_{t>0} J_{\varepsilon}(t\Psi_{\varepsilon,y}) = J_{\varepsilon}(t_{\varepsilon}\Psi_{\varepsilon,y}).$$

Finally, we consider  $\Phi_{\varepsilon}(y) = t_{\varepsilon} \Psi_{\varepsilon,y}$ .

In next lemma we prove an important relationship between  $\Phi_{\varepsilon}$  and the set M.

**Lemma 4.1.** The functional  $\Phi_{\varepsilon}$  satisfies the following limit

$$\lim_{\varepsilon \to 0} J_{\varepsilon}(\Phi_{\varepsilon}(y)) = c_{V_0} \text{ uniformly in } y \in M.$$

*Proof.* Assume by contradiction that there exist  $\delta_0 > 0$ ,  $(y_n) \subset M$  and  $\varepsilon_n \to 0$  such that

$$|J_{\varepsilon_n}(\Phi_{\varepsilon_n}(y_n)) - c_{V_0}| \ge \delta_0. \tag{4.1}$$

We first show that  $\lim_{n\to\infty} t_{\varepsilon_n} < \infty$ . Let us observe that by using the change of variable  $z = \frac{\varepsilon_n x - y_n}{\varepsilon_n}$ , if  $z \in B_{\frac{\delta}{\varepsilon_n}}(0)$ , it follows that  $\varepsilon_n z \in B_{\delta}(0)$  and  $\varepsilon_n z + y_n \in B_{\delta}(y_n) \subset M_{\delta} \subset \Lambda$ .

Since G = F on  $\Lambda$ , we can see that

$$J_{\varepsilon_n}(\Phi_{\varepsilon_n}(z_n)) = \frac{t_{\varepsilon_n}^2}{2} \int_{\mathbb{R}^N} |(-\Delta)^{\frac{s}{2}} (\eta(|\varepsilon_n z|) w(z))|^2 dz + \frac{t_{\varepsilon_n}^2}{2} \int_{\mathbb{R}^N} V(\varepsilon_n z + y_n) (\eta(|\varepsilon_n z|) w(z))^2 dz - \sum_0 (t_{\varepsilon_n} \eta(|\varepsilon_n z|) w(z)).$$

$$(4.2)$$

In view of the Dominated Convergence Theorem and Lemma 5 in [32], we can see that

$$\lim_{n \to \infty} \|\Psi_{\varepsilon_n, y_n}\|_{\varepsilon_n} = \|w\|_{V_0}$$

and

$$\lim_{n\to\infty} \Sigma_0(\Psi_{\varepsilon_n,y_n}) = \Sigma_0(w).$$

By using  $t_{\varepsilon_n} \Psi_{\varepsilon_n, y_n} \in \mathcal{N}_{\varepsilon_n}$  and the assumptions on f, it is easy to prove that  $t_{\varepsilon_n} \to t_0 > 0$ . Moreover, being

$$t_{\varepsilon_n}^2 \|\Psi_{\varepsilon_n, y_n}\|_{\varepsilon_n}^2 = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{F(t_{\varepsilon_n} \Psi_{\varepsilon_n, y_n}) f(t_{\varepsilon_n} \Psi_{\varepsilon_n, y_n}) t_{\varepsilon_n} \Psi_{\varepsilon_n, y_n}}{|x - y|^{\mu}}$$
(4.3)

we can deduce that

$$||w||_{V_0}^2 = \lim_{n \to \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{F(t_{\varepsilon_n} \Psi_{\varepsilon_n, y_n}) f(t_{\varepsilon_n} \Psi_{\varepsilon_n, y_n}) t_{\varepsilon_n} \Psi_{\varepsilon_n, y_n}}{t_{\varepsilon_n}^2 |x - y|^{\mu}}.$$

Taking into account that w is a ground state to  $(P_{V_0})$  and using  $(f_4)$ , we can conclude that  $t_{\varepsilon_n} \to 1$ . As a consequence

$$\lim_{n \to \infty} \Sigma_0(t_{\varepsilon_n} \eta(|\varepsilon_n z|) w(z)) = \Sigma_0(w)$$

and this yields

$$\lim_{n \to \infty} J_{\varepsilon_n}(\Phi_{\varepsilon_n}(y_n)) = J_0(w) = c_{V_0},$$

which contradicts (4.1).

Now, we consider  $\delta > 0$  such that  $M_{\delta} \subset \Lambda$ , and choose  $\rho = \rho(\delta) > 0$  such that  $M_{\delta} \subset B_{\delta}(0)$ . We define  $\Upsilon : \mathbb{R}^N \to \mathbb{R}^N$  by setting  $\Upsilon(x) = x$  for  $|x| \leq \rho$  and  $\Upsilon(x) = \frac{\rho x}{|x|}$  for  $|x| \geq \rho$ . Then we define the barycenter map  $\beta_{\varepsilon} : \mathcal{N}_{\varepsilon} \to \mathbb{R}^N$  given by

$$\beta_{\varepsilon}(u) = \frac{\int_{\mathbb{R}^N} \Upsilon(\varepsilon x) u^2(x) \, dx}{\int_{\mathbb{R}^N} u^2(x) \, dx}.$$

**Lemma 4.2.** The function  $\beta_{\varepsilon}$  verifies the following limit

$$\lim_{\varepsilon \to 0} \beta_{\varepsilon}(\Phi_{\varepsilon}(y)) = y \text{ uniformly in } y \in M.$$

*Proof.* Suppose by contradiction that there exists  $\delta_0 > 0$ ,  $(y_n) \subset M$  and  $\varepsilon_n \to 0$  such that

$$|\beta_{\varepsilon_n}(\Phi_{\varepsilon_n}(y_n)) - y_n| \ge \delta_0. \tag{4.4}$$

By using the definitions of  $\Phi_{\varepsilon_n}(y_n)$ ,  $\beta_{\varepsilon_n}$  and  $\eta$ , and using a change of variable, we can see that

$$\beta_{\varepsilon_n}(\Psi_{\varepsilon_n}(y_n)) = y_n + \frac{\int_{\mathbb{R}^N} [\Upsilon(\varepsilon_n z + y_n) - y_n] |\eta(|\varepsilon_n z|) w(z)|^2 dz}{\int_{\mathbb{R}^N} |\eta(|\varepsilon_n z|) w(z)|^2 dz}.$$

Taking into account  $(y_n) \subset M \subset B_\rho$  and the Dominated Convergence Theorem, we can infer that

$$|\beta_{\varepsilon_n}(\Phi_{\varepsilon_n}(y_n)) - y_n| = o_n(1)$$

which contradicts (4.4).

The next compactness result will be fundamental to show that the solutions of the modified problem are solutions of the original problem.

**Lemma 4.3.** Let  $\varepsilon_n \to 0$  and  $(u_n) \subset \mathcal{N}_{\varepsilon_n}$  be such that  $J_{\varepsilon_n}(u_n) \to c_{V_0}$ . Then there exists  $(\tilde{y}_n) \subset \mathbb{R}^N$  such that  $v_n(x) = u_n(x + \tilde{y}_n)$  has a convergent subsequence in  $H^s(\mathbb{R}^N)$ . Moreover, up to a subsequence,  $y_n = \varepsilon_n \tilde{y}_n \to y_0 \in M$ .

Proof. Since  $\langle J'_{\varepsilon_n}(u_n), u_n \rangle = 0$  and  $J_{\varepsilon_n}(u_n) \to c_{V_0}$ , we can see that  $(u_n)$  is bounded in  $H^s_{\varepsilon_n}$ . Note that  $c_{V_0} > 0$ , and since  $||u_n||_{\varepsilon_n} \to 0$  would imply  $J_{\varepsilon_n}(u_n) \to 0$ , we can argue as in Lemma 3.2 to obtain a sequence  $(\tilde{y}_n) \subset \mathbb{R}^N$ , and constants R > 0 and  $\gamma > 0$  such that

$$\liminf_{n \to \infty} \int_{B_R(\tilde{y}_n)} |u_n|^2 dx \ge \gamma > 0.$$
(4.5)

Now, we set  $v_n(x) = u_n(x + \tilde{y}_n)$ . Then,  $(v_n)$  is bounded in  $H_0^s$ , and we may assume that  $v_n \rightharpoonup v \not\equiv 0$  in  $H_0^s$  as  $n \to \infty$ . Fix  $t_n > 0$  such that  $\tilde{v}_n = t_n v_n \in \mathcal{N}_{V_0}$ . Since  $u_n \in \mathcal{N}_{\varepsilon_n}$ , we can see that

$$c_{V_0} \le J_{V_0}(\tilde{v}_n) = J_{V_0}(t_n u_n) \le J_{\varepsilon_n}(t_n u_n) \le J_{\varepsilon_n}(u_n) = c_{V_0} + o_n(1)$$

which gives  $J_{V_0}(\tilde{v}_n) \to c_{V_0}$ . In particular, we get  $\tilde{v}_n \rightharpoonup \tilde{v}$  in  $H_0^s$  and  $t_n \to t^* > 0$ . Then, from the uniqueness of the weak limit, we have  $\tilde{v} = t^*v \not\equiv 0$ .

By using Lemma 3.4, we can see that

$$\tilde{v}_n \to \tilde{v} \text{ in } H_0^s.$$
 (4.6)

In order to complete the proof of the lemma, we consider  $y_n = \varepsilon_n \tilde{y}_n$ . Our claim is to show that  $(y_n)$  admits a subsequence, still denoted by  $y_n$ , such that  $y_n \to y_0$ , for some  $y_0 \in M$ . Firstly, we prove that  $(y_n)$  is bounded. We argue by contradiction, and we assume that, up to a subsequence,  $|y_n| \to \infty$  as  $n \to \infty$ . Since

$$||u_n||_{\varepsilon_n}^2 = \int_{\mathbb{R}^N} \left( \frac{1}{|x|^{\mu}} * G(\varepsilon x, u_n) \right) g(\varepsilon x, u_n) u_n,$$

and  $J_{\varepsilon_n}(u_n) \to c_{V_0}$ , we can see that  $u_n \in \mathcal{B}$  for all n big enough. Then, in view of Lemma 2.3, there exists  $C_0 > 0$  such that

$$\sup_{n \in \mathbb{N}} \left\| \frac{1}{|x|^{\mu}} * G(\varepsilon x, u_n) \right\|_{L^{\infty}(\mathbb{R}^N)} < C_0.$$

Fixed R>0 such that  $\Lambda\subset B_R(0)$ , and assume that  $|y_n|>2R$ . Then, for all  $z\in B_{\frac{R}{\varepsilon_n}}(0)$ ,

$$|\varepsilon_n z + y_n| \ge |y_n| - |\varepsilon_n z| > R. \tag{4.7}$$

By using the change of variable  $x \mapsto z + \tilde{y}_n$  and (4.7), we deduce that

$$[v_n]^2 + \int_{\mathbb{R}^N} V_0 v_n^2 dx \le C_0 \int_{\mathbb{R}^N} g(\varepsilon_n z + y_n, v_n) v_n dx$$

$$\le C_0 \int_{B_{\frac{R}{\varepsilon_n}}(0)} \tilde{f}(v_n) v_n dx + C_0 \int_{\mathbb{R}^N \setminus B_{\frac{R}{\varepsilon_n}}(0)} f(v_n) v_n dx. \tag{4.8}$$

Then, by using the fact that  $v_n \to v$  in  $H_0^s$  as  $n \to \infty$  and that  $\tilde{f}(t) \leq \frac{V_0}{\ell_0}t$ , we can see that (4.8) implies that

$$[v_n]^2 + \int_{\mathbb{D}^N} V_0 v_n^2 \, dx = o_n(1),$$

that is  $v_n \to 0$  in  $H_0^s$ , which is a contradiction. Therefore,  $(y_n)$  is bounded, and we may assume that  $y_n \to y_0 \in \mathbb{R}^N$ . Clearly, if  $y_0 \notin \overline{\Lambda}$ , then we can argue as before and we deduce that  $v_n \to 0$  in  $H_0^s$ , which is impossible. Hence  $y_0 \in \overline{\Lambda}$ . Now, we note that if  $V(y_0) = V_0$ , then we can infer that  $y_0 \notin \partial \Lambda$  in view of  $(V_2)$ , and then  $y_0 \in M$ . Therefore, in the next step, we show that  $V(y_0) = V_0$ .

Suppose by contradiction that  $V(y_0) > V_0$ . Then, by using  $\tilde{v}_n \to \tilde{v}$  in  $H_0^s$  and Fatou's Lemma, we get

$$c_{V_0} = J_{V_0}(\tilde{v}) < \frac{1}{2} \left( [\tilde{v}]^2 + \int_{\mathbb{R}^N} V(y_0) \tilde{v}^2 \right) - \Sigma_0(\tilde{v})$$

$$\leq \liminf_{n \to \infty} \left[ \frac{1}{2} [\tilde{v}_n]^2 + \frac{1}{2} \int_{\mathbb{R}^N} V(\varepsilon_n z + y_n) \tilde{v}_n^2 dx - \Sigma_0(\tilde{v}_n) \right]$$

$$\leq \liminf_{n \to \infty} J_{\varepsilon_n}(t_n u_n) \leq \liminf_{n \to \infty} J_{\varepsilon_n}(u_n) = c_{V_0}$$

which gives a contradiction.

Now, we introduce a subset  $\tilde{\mathcal{N}}_{\varepsilon}$  of  $\mathcal{N}_{\varepsilon}$  by setting

$$\tilde{\mathcal{N}}_{\varepsilon} = \{ u \in \mathcal{N}_{\varepsilon} : J_{\varepsilon}(u) \le c_{V_0} + h(\varepsilon) \},$$

where  $h: \mathbb{R}_+ \to \mathbb{R}_+$  is such that  $h(\varepsilon) \to 0$  as  $\varepsilon \to 0$ . Given  $y \in M$ , we can use Lemma 4.1 to conclude that  $h(\varepsilon) = |J_{\varepsilon}(\Phi_{\varepsilon}(y)) - c_{V_0}| \to 0$  as  $\varepsilon \to 0$ . Hence  $\Phi_{\varepsilon}(y) \in \tilde{\mathcal{N}}_{\varepsilon}$ , and  $\tilde{\mathcal{N}}_{\varepsilon} \neq \emptyset$  for any  $\varepsilon > 0$ . Moreover, we have the following lemma.

### Lemma 4.4.

$$\lim_{\varepsilon \to 0} \sup_{u \in \tilde{\mathcal{N}}_{\varepsilon}} dist(\beta_{\varepsilon}(u), M_{\delta}) = 0.$$

*Proof.* Let  $\varepsilon_n \to 0$  as  $n \to \infty$ . For any  $n \in \mathbb{N}$ , there exists  $u_n \in \tilde{\mathcal{N}}_{\varepsilon_n}$  such that

$$\sup_{u \in \tilde{\mathcal{N}}_{\varepsilon_n}} \inf_{y \in M_{\delta}} |\beta_{\varepsilon_n}(u) - y| = \inf_{y \in M_{\delta}} |\beta_{\varepsilon_n}(u_n) - y| + o_n(1).$$

Therefore, it is suffices to prove that there exists  $(y_n) \subset M_\delta$  such that

$$\lim_{n \to \infty} |\beta_{\varepsilon_n}(u_n) - y_n| = 0. \tag{4.9}$$

We note that  $(u_n) \subset \tilde{\mathcal{N}}_{\varepsilon_n} \subset \mathcal{N}_{\varepsilon_n}$ , from which we deuce that

$$c_{V_0} \le c_{\varepsilon_n} \le J_{\varepsilon_n}(u_n) \le c_{V_0} + h(\varepsilon_n).$$

This yields  $J_{\varepsilon_n}(u_n) \to c_{V_0}$ . By using Lemma 4.3, there exists  $(\tilde{y}_n) \subset \mathbb{R}^N$  such that  $y_n = \varepsilon_n \tilde{y}_n \in M_\delta$  for n sufficiently large. By setting  $v_n = u_n(\cdot + \tilde{y}_n)$  and using a change of variable, we can see that

$$\beta_{\varepsilon_n}(u_n) = y_n + \frac{\int_{\mathbb{R}^N} [\Upsilon(\varepsilon_n z + y_n) - y_n] v_n^2(z) \, dz}{\int_{\mathbb{R}^N} v_n^2(z) \, dz}.$$

Since  $\varepsilon_n z + y_n \to y \in M$ , we deduce that  $\beta_{\varepsilon_n}(u_n) = y_n + o_n(1)$ , that is (4.9) holds.

#### 5. Proof of Theorem 1.1

This last section is devoted to the proof of the main result of this work. Firstly, we show that (2.3) admits at least  $cat_{M_{\delta}}(M)$  positive solutions. In order to achieve our aim, we recall the following result for critical points involving Ljusternik-Schnirelmann category. For the details of the proof one can see [27].

**Theorem 5.1.** Let U be a  $C^{1,1}$  complete Riemannian manifold (modelled on a Hilbert space). Assume that  $h \in C^1(U, \mathbb{R})$  bounded from below and satisfies  $-\infty < \inf_U h < d < k < \infty$ . Moreover, suppose that h satisfies Palais-Smale condition on the sublevel  $\{u \in U : h(u) \le k\}$  and that d is not a critical level for h. Then

$$card\{u \in h^d : \nabla h(u) = 0\} \ge cat_{h^d}(h^d).$$

Since  $\mathcal{N}_{\varepsilon}$  is not a  $C^1$  submanifold of  $H^s_{\varepsilon}$ , we cannot apply Theorem 5.1 directly. Fortunately, from Lemma 2.4, we know that the mapping  $m_{\varepsilon}$  is a homeomorphism between  $\mathcal{N}_{\varepsilon}$  and  $\mathbb{S}_{\varepsilon}$ , and  $\mathbb{S}_{\varepsilon}$  is a  $C^1$  submanifold of  $H^s_{\varepsilon}$ . So we can apply Theorem 5.1 to  $\psi_{\varepsilon}(u) = J_{\varepsilon}(\hat{m}_{\varepsilon}(u))|_{\mathbb{S}_{\varepsilon}} = J_{\varepsilon}(m_{\varepsilon}(u))$ , where  $\psi_{\varepsilon}$  is given in Proposition 2.1.

**Theorem 5.2.** Assume that  $(V_1)$ - $(V_2)$  and  $(f_1)$ - $(f_4)$  hold. Then, for any  $\delta > 0$  there exists  $\bar{\varepsilon}_{\delta} > 0$  such that, for any  $\varepsilon \in (0, \bar{\varepsilon}_{\delta})$ , problem (2.3) has at least  $cat_{M_{\delta}}(M)$  positive solutions.

*Proof.* For any  $\varepsilon > 0$ , we define  $\alpha_{\varepsilon} : M \to \mathbb{S}_{\varepsilon}$  by setting  $\alpha_{\varepsilon}(y) = m_{\varepsilon}^{-1}(\Phi_{\varepsilon}(y))$ . By using Lemma 4.1 and the definition of  $\psi_{\varepsilon}$ , we can see that

$$\lim_{\varepsilon \to 0} \psi_{\varepsilon}(\alpha_{\varepsilon}(y)) = \lim_{\varepsilon \to 0} J_{\varepsilon}(\Phi_{\varepsilon}(y)) = c_{V_0} \text{ uniformly in } y \in M.$$

Then, there exists  $\bar{\varepsilon} > 0$  such that  $\tilde{\mathbb{S}}_{\varepsilon} := \{ w \in \mathbb{S}_{\varepsilon} : \psi_{\varepsilon}(w) \leq c_{V_0} + h(\varepsilon) \} \neq \emptyset$  for all  $\varepsilon \in (0, \bar{\varepsilon})$ . Taking into account Lemma 4.1, Lemma 2.4-(c), Lemma 4.2 and Lemma 4.4, we can find  $\bar{\varepsilon} = \bar{\varepsilon}_{\delta} > 0$  such that the following diagram

$$M \stackrel{\Phi_{\varepsilon}}{\to} \tilde{\mathcal{N}}_{\varepsilon} \stackrel{m_{\varepsilon}^{-1}}{\to} \tilde{\mathbb{S}}_{\varepsilon} \stackrel{m_{\varepsilon}}{\to} \tilde{\mathcal{N}}_{\varepsilon} \stackrel{\beta_{\varepsilon}}{\to} M_{\delta}$$

is well defined for any  $\varepsilon \in (0, \bar{\varepsilon})$ .

By using Lemma 4.2, there exists a function  $\theta(\varepsilon,y)$  with  $|\theta(\varepsilon,y)| < \frac{\delta}{2}$  uniformly in  $y \in M$  for all  $\varepsilon \in (0,\bar{\varepsilon})$  such that  $\beta_{\varepsilon}(\Phi_{\varepsilon}(y)) = y + \theta(\varepsilon,y)$  for all  $y \in M$ . Then, we can see that  $H(t,y) = y + (1-t)\theta(\varepsilon,y)$  with  $(t,y) \in [0,1] \times M$  is a homotopy between  $\beta_{\varepsilon} \circ \Phi_{\varepsilon} = (\beta_{\varepsilon} \circ m_{\varepsilon}) \circ \alpha_{\varepsilon}$  and the inclusion map  $id: M \to M_{\delta}$ . This fact and Lemma 4.3 in [9] implies that  $\cot_{\tilde{\mathbb{S}}_{\varepsilon}}(\tilde{\mathbb{S}}_{\varepsilon}) \geq \cot_{M_{\delta}}(M)$ . On the other hand, let us choose a function  $h(\varepsilon) > 0$  such that  $h(\varepsilon) \to 0$  as  $\varepsilon \to 0$  and such that  $c_{V_0} + h(\varepsilon)$  is not a critical level for  $J_{\varepsilon}$ . For  $\varepsilon > 0$  small enough, we deduce from Lemma 2.6 that  $\psi_{\varepsilon}$  satisfies the Palais-Smale condition in  $\tilde{\mathbb{S}}_{\varepsilon}$ . So, it follows from Theorem 5.1 that  $\psi_{\varepsilon}$  has at least  $\cot_{\tilde{\mathbb{S}}_{\varepsilon}}(\tilde{\mathbb{S}}_{\varepsilon})$  critical points on  $\tilde{\mathbb{S}}_{\varepsilon}$ . By Proposition 2.1-(d) we conclude that  $J_{\varepsilon}$  admits at least  $\cot_{M_{\delta}}(M)$  critical points.

Now, we use a Moser iteration argument [29] which will be fundamental to study of behavior of the maximum points of the solutions.

**Lemma 5.1.** Let  $\varepsilon_n \to 0$  and  $u_n \in \widetilde{\mathcal{N}}_{\varepsilon_n}$  be a solution to (2.3). Then  $v_n = u_n(\cdot + \tilde{y}_n)$  satisfies the following problem

$$\begin{cases}
(-\Delta)^s v_n + V_n(x) v_n = \left(\frac{1}{|x|^{\mu}} * G_n(v_n)\right) g_n(v_n) & \text{in } \mathbb{R}^N \\
v_n \in H^s(\mathbb{R}^N) & \text{in } \mathbb{R}^N, \\
v_n > 0 & \text{in } \mathbb{R}^N,
\end{cases}$$
(5.1)

where  $V_n(x) = V(\varepsilon_n x + \varepsilon_n \tilde{y}_n)$ ,  $g_n(v_n) = g(\varepsilon_n x + \varepsilon_n \tilde{y}_n, v_n)$ ,  $\varepsilon_n \tilde{y}_n \to y \in M$ , and there exists C > 0 such that  $||v_n||_{L^{\infty}(\mathbb{R}^N)} \leq C$  for all  $n \in \mathbb{N}$ .

*Proof.* For any L > 0 and  $\beta > 1$ , let us define the function

$$\gamma(v_n) = \gamma_{L,\beta}(v_n) = v_n v_{L,n}^{2(\beta-1)} \in H^s(\mathbb{R}^N)$$

where  $v_{L,n} = \min\{u_n, L\}$ . Since  $\gamma$  is an increasing function, we have

$$(a-b)(\gamma(a)-\gamma(b)) \ge 0$$
 for any  $a,b \in \mathbb{R}$ .

Let us consider

$$\mathcal{E}(t) = \frac{|t|^2}{2}$$
 and  $\Gamma(t) = \int_0^t (\gamma'(\tau))^{\frac{1}{2}} d\tau$ .

Then, by applying Jensen inequality we get for all  $a, b \in \mathbb{R}$  such that a > b,

$$\mathcal{E}'(a-b)(\gamma(a)-\gamma(b)) = (a-b)(\gamma(a)-\gamma(b)) = (a-b)\int_a^b \gamma'(t)dt$$
$$= (a-b)\int_a^b (\Gamma'(t))^2 dt \ge \left(\int_a^b (\Gamma'(t)) dt\right)^2.$$

The same argument works when  $a \leq b$ . Therefore

$$\mathcal{E}'(a-b)(\gamma(a)-\gamma(b)) \ge |\Gamma(a)-\Gamma(b)|^2 \text{ for any } a,b \in \mathbb{R}.$$
(5.2)

By using (5.2), we can see that

$$|\Gamma(v_n)(x) - \Gamma(v_n)(y)|^2 \le (v_n(x) - v_n(y))((v_n v_{L,n}^{2(\beta-1)})(x) - (v_n v_{L,n}^{2(\beta-1)})(y)). \tag{5.3}$$

Choosing  $\gamma(v_n) = v_n v_{L,n}^{2(\beta-1)}$  as test-function in (5.1), and using (5.3), we obtain

$$[\Gamma(v_n)]^2 + \int_{\mathbb{R}^N} V_n(x)|v_n|^2 v_{L,n}^{2(\beta-1)} dx$$

$$\leq \iint_{\mathbb{R}^{2N}} \frac{(v_n(x) - v_n(y))}{|x - y|^{N+2s}} ((v_n v_{L,n}^{2(\beta-1)})(x) - (v_n v_{L,n}^{2(\beta-1)})(y)) dx dy + \int_{\mathbb{R}^N} V_n(x)|v_n|^2 v_{L,n}^{2(\beta-1)} dx$$

$$= \int_{\mathbb{R}^N} \left(\frac{1}{|x|^{\mu}} * G_n(v_n)\right) g_n(v_n) v_n v_{L,n}^{2(\beta-1)} dx. \tag{5.4}$$

Since

$$\Gamma(v_n) \ge \frac{1}{\beta} v_n v_{L,n}^{\beta - 1},$$

and using Theorem 2.1, we have

$$[\Gamma(v_n)]^2 \ge S_* \|\Gamma(v_n)\|_{L^{2_s^*}(\mathbb{R}^N)}^2 \ge \left(\frac{1}{\beta}\right)^2 S_* \|v_n v_{L,n}^{\beta-1}\|_{L^{2_s^*}(\mathbb{R}^N)}^2. \tag{5.5}$$

On the other hand, from the boundedness of  $(v_n)$ , it follows that there exists  $C_0 > 0$  such that

$$\sup_{n \in \mathbb{N}} \left\| \frac{1}{|x|^{\mu}} * G(\varepsilon x, v_n) \right\|_{L^{\infty}(\mathbb{R}^N)} < C_0.$$
 (5.6)

By the assumption  $(g_1)$  and  $(g_2)$ , for any  $\xi > 0$  there exists  $C_{\xi} > 0$  such that

$$|g_n(v_n)| \le \xi |v_n| + C_\xi |v_n|^{q-1}.$$
 (5.7)

Taking  $\xi \in (0, V_0)$ , and using (5.5), (5.6) and (5.7), we can see that (5.4) yields

$$||v_n v_{L,n}^{\beta-1}||_{L^{2_s^*}(\mathbb{R}^N)}^2 \le C\beta^2 \int_{\mathbb{R}^N} |v_n|^q v_{L,n}^{2(\beta-1)} dx.$$

Set  $w_{L,n} := v_n v_{L,n}^{\beta-1}$ . By applying Hölder inequality, we get

$$||w_{L,n}||_{L^{2_s^*}(\mathbb{R}^N)}^2 \le C\beta^2 \left( \int_{\mathbb{R}^N} v_n^{2_s^*} dx \right)^{\frac{q-2}{2_s^*}} \left( \int_{\mathbb{R}^N} w_{L,n}^{\alpha_s^*} dx \right)^{\frac{2}{\alpha_s^*}}$$

where  $\alpha_s^* := \frac{22_s^*}{2_s^* - (q-2)} \in (2, 2_s^*)$ .

Since  $(v_n)$  is bounded in  $H^s(\mathbb{R}^N)$ , we deduce that

$$||w_{L,n}||_{L^{2_s^*}(\mathbb{R}^N)}^2 \le C\beta^2 ||w_{L,n}||_{L^{\alpha_s^*}(\mathbb{R}^N)}^2.$$
(5.8)

Now, we observe that if  $v_n^{\beta} \in L^{\alpha_s^*}(\mathbb{R}^N)$ , from the definition of  $w_{L,n}$ , and by using the fact that  $v_{L,n} \leq v_n$  and (5.8), we obtain

$$||w_{L,n}||_{L^{2_s^*}(\mathbb{R}^N)}^2 \le C\beta^2 \left( \int_{\mathbb{R}^N} v_n^{\beta \alpha_s^*} dx \right)^{\frac{2}{\alpha_s^*}} < \infty.$$
 (5.9)

By passing to the limit in (5.9) as  $L \to +\infty$ , the Fatou's Lemma yields

$$||v_n||_{L^{\beta 2_s^*}(\mathbb{R}^N)} \le C^{\frac{1}{\beta}} \beta^{\frac{1}{\beta}} ||v_n||_{L^{\beta \alpha_s^*}(\mathbb{R}^N)}$$
(5.10)

whenever  $v_n^{\beta \alpha_s^*} \in L^1(\mathbb{R}^N)$ .

Now, we set  $\beta := \frac{2_s^*}{\alpha_s^*} > 1$ , and we observe that, being  $v_n \in L^{2_s^*}(\mathbb{R}^N)$ , the above inequality holds for this choice of  $\beta$ . Then, observing that  $\beta^2 \alpha_s^* = \beta 2_s^*$ , it follows that (5.10) holds with  $\beta$  replaced by  $\beta^2$ . Therefore, we can see that

$$\|v_n\|_{L^{\beta^2 2_s^*}(\mathbb{R}^N)} \leq C^{\frac{1}{\beta^2}} \beta^{\frac{2}{\beta^2}} \|v_n\|_{L^{\beta^2 \alpha_s^*}(\mathbb{R}^N)} \leq C^{\left(\frac{1}{\beta} + \frac{1}{\beta^2}\right)} \beta^{\frac{1}{\beta} + \frac{2}{\beta^2}} \|v_n\|_{L^{\beta \alpha_s^*}(\mathbb{R}^N)}.$$

Iterating this process, and recalling that  $\beta \alpha^* := 2_s^*$ , we can infer that for every  $m \in \mathbb{N}$ 

$$||v_n||_{L^{\beta^m 2_s^*}(\mathbb{R}^N)} \le C^{\sum_{j=1}^m \frac{1}{\beta^j}} \beta^{\sum_{j=1}^m j\beta^{-j}} ||v_n||_{L^{2_s^*}(\mathbb{R}^N)}.$$
(5.11)

Taking the limit in (5.11) as  $m \to +\infty$  and recalling that  $||v_n||_{L^{2_s^*}(\mathbb{R}^N)} \leq K$ , we get

$$||v_n||_{L^{\infty}(\mathbb{R}^N)} \le C^{\sigma_1} \beta^{\sigma_2} K,$$

where

$$\sigma_1 := \sum_{j=1}^{\infty} \frac{1}{\beta^j} < \infty$$
 and  $\sigma_2 := \sum_{j=1}^{\infty} \frac{j}{\beta^j} < \infty$ .

At this point, we are ready to give the proof of our main result.

Proof of Theorem 1.1. Take  $\delta > 0$  such that  $M_{\delta} \subset \Lambda$ . We begin proving that there exists  $\tilde{\varepsilon}_{\delta} > 0$  such that for any  $\varepsilon \in (0, \tilde{\varepsilon}_{\delta})$  and any solution  $u_{\varepsilon} \in \tilde{\mathcal{N}}_{\varepsilon}$  of (2.3), it holds

$$||u_{\varepsilon}||_{L^{\infty}(\mathbb{R}^{N}\setminus\Lambda_{\varepsilon})} < a. \tag{5.12}$$

Assume by contradiction that there exist  $\varepsilon_n \to 0$ ,  $u_{\varepsilon_n} \in \tilde{\mathcal{N}}_{\varepsilon_n}$  such that  $J'_{\varepsilon_n}(u_{\varepsilon_n}) = 0$  and  $\|u_{\varepsilon_n}\|_{L^{\infty}(\mathbb{R}^N \setminus \Lambda_{\varepsilon_n})} \ge a$ . Since  $J_{\varepsilon_n}(u_{\varepsilon_n}) \le c_{V_0} + h(\varepsilon_n)$  and  $h(\varepsilon_n) \to 0$ , we can argue as in the first part of the proof of Lemma 4.3, to deduce that  $J_{\varepsilon_n}(u_{\varepsilon_n}) \to c_{V_0}$ . Then, by using Lemma 4.3, we can find  $(\tilde{y}_n) \subset \mathbb{R}^N$  such that  $y_n := \varepsilon_n \tilde{y}_n \to y_0 \in M$ .

Now, if we choose r > 0 such that  $B_r(y_0) \subset B_{2r}(y_0) \subset \Lambda$ , we can see  $B_{\frac{r}{\varepsilon_n}}(\frac{y_0}{\varepsilon_n}) \subset \Lambda_{\varepsilon_n}$ . In particular, for any  $y \in B_{\frac{r}{\varepsilon_n}}(\tilde{y}_n)$  there holds

$$\left| y - \frac{y_0}{\varepsilon_n} \right| \le |y - \tilde{y}_n| + \left| \tilde{y}_n - \frac{y_0}{\varepsilon_n} \right| < \frac{2r}{\varepsilon_n}$$
 for  $n$  sufficiently large.

Therefore

$$\mathbb{R}^N \setminus \Lambda_{\varepsilon_n} \subset \mathbb{R}^N \setminus B_{\frac{r}{\varepsilon_n}}(\tilde{y}_n)$$
 for any  $n$  big enough.

Now, denoting by  $v_n(x) = u_{\varepsilon_n}(x + \tilde{y}_n)$ , we can see that

$$(-\Delta)^s v_n + v_n = h_n \text{ in } \mathbb{R}^N,$$

where

$$h_n := v_n - V_n(x)v_n + \left(\frac{1}{|x|^{\mu}} * G_n(v_n)\right) g_n(v_n),$$

and  $v_n \to v$  converges strongly in  $L^p(\mathbb{R}^N)$  for any  $p \in [2, \infty)$ , in view of Lemma 5.1.

Since  $\langle J'_{\varepsilon_n}(u_{\varepsilon_n}), u_{\varepsilon_n} \rangle = 0$  and  $J_{\varepsilon_n}(u_{\varepsilon_n}) \to c_{V_0}$ , we may assume  $u_{\varepsilon_n} \in \mathcal{B}$  for all n big enough, so that

$$\left\| \frac{1}{|x|^{\mu}} * G(\varepsilon x, u_{\varepsilon_n}) \right\|_{L^{\infty}(\mathbb{R}^N)} < C_0.$$

As a consequence, recalling that  $\varepsilon_n \tilde{y}_n \to y_0 \in M$ , we get

$$||h_n||_{L^{\infty}(\mathbb{R}^N)} \le C$$
 and  $h_n \to v - V(y_0)v + \left(\frac{1}{|x|^{\mu}} * F(v)\right)f(v)$  in  $L^p(\mathbb{R}^N)$   $\forall p \in [2, \infty)$ .

Hence,  $v_n = \mathcal{K} * h_n$ , where  $\mathcal{K}$  is the Bessel kernel [19], and we can argue as in [2] to prove that

$$\lim_{|x| \to \infty} \sup_{n \in \mathbb{N}} |v_n(x)| = 0,$$

which implies that there exists R > 0 such that  $v_n(x) < a$  for  $|x| \ge R$  and  $n \in \mathbb{N}$ . Thus

$$u_{\varepsilon_n}(x) < a \text{ for any } x \in \mathbb{R}^N \setminus B_R(\tilde{y}_n), n \in \mathbb{N}.$$

As a consequence, there exists  $\nu \in \mathbb{N}$  such that for any  $n \geq \nu$  and  $\frac{r}{\varepsilon_n} > R$ , it holds

$$\mathbb{R}^N \setminus \Lambda_{\varepsilon_n} \subset \mathbb{R}^N \setminus B_{\frac{r}{\varepsilon_n}}(\tilde{y}_n) \subset \mathbb{R}^N \setminus B_R(\tilde{y}_n),$$

which gives  $u_{\varepsilon_n}(x) < a$  for any  $x \in \mathbb{R}^N \setminus \Lambda_{\varepsilon_n}$ , and this is impossible.

Now, let  $\bar{\varepsilon}_{\delta}$  given in Theorem 5.2 and take  $\varepsilon_{\delta} = \min\{\tilde{\varepsilon}_{\delta}, \bar{\varepsilon}_{\delta}\}$ . Fix  $\varepsilon \in (0, \varepsilon_{\delta})$ . By Theorem 5.2, we know that problem (2.3) admits  $cat_{M_{\delta}}(M)$  nontrivial solutions  $u_{\varepsilon}$ . Since  $u_{\varepsilon} \in \tilde{\mathcal{N}}_{\varepsilon}$  satisfies (5.12), from the definition of g it follows that  $u_{\varepsilon}$  is a solution of (2.2).

Now, we study the behavior of the maximum points of  $u_n \in H^s_{\varepsilon_n}$  solutions to the problem (2.3). Let us observe that  $(g_1)$  implies that we can find  $\gamma \in (0, a)$  such that

$$g(\varepsilon x, t)t \le \frac{V_0}{\ell_0} t^2 \text{ for any } x \in \mathbb{R}^N, t \le \gamma.$$
 (5.13)

Arguing as before, we can find R > 0 such that

$$||u_n||_{L^{\infty}(B_R^c(\tilde{y}_n))} < \gamma. \tag{5.14}$$

Moreover, up to extract a subsequence, we may assume that

$$||u_n||_{L^{\infty}(B_R(\tilde{y}_n))} \ge \gamma. \tag{5.15}$$

Indeed, if (5.15) does not hold, in view of (5.14) we can see that  $||u_n||_{L^{\infty}(\mathbb{R}^N)} < \gamma$ . Then, by using  $\langle J'_{\varepsilon_n}(u_n), u_n \rangle = 0$ , (5.13) and

$$\left\| \frac{1}{|x|^{\mu}} * G(\varepsilon x, u_n) \right\|_{L^{\infty}(\mathbb{R}^N)} < C_0,$$

we can infer

$$||u_n||_{\varepsilon_n}^2 \le C_0 \int_{\mathbb{R}^N} g(\varepsilon_n x, u_n) u_n \, dx \le \frac{C_0}{\ell_0} \int_{\mathbb{R}^N} V_0 u_n^2 \, dx$$

which together with  $\frac{C_0}{\ell_0} < \frac{1}{2}$  yields  $||u_n||_{\varepsilon_n} = 0$ , and this gives a contradiction. As a consequence, (5.15) holds. Taking into account (5.14) and (5.15) we can deduce that the maximum points  $p_n \in \mathbb{R}^N$  of  $u_n$  belong to  $B_R(\tilde{y}_n)$ . Therefore,  $p_n = \tilde{y}_n + q_n$  for some  $q_n \in B_R(0)$ . Hence,  $\eta_n = \varepsilon_n \tilde{y}_n + \varepsilon_n q_n$  is the maximum point of  $\hat{u}_n(x) = u_n(x/\varepsilon_n)$ . Since  $|q_n| < R$  for any  $n \in \mathbb{N}$  and  $\varepsilon_n \tilde{y}_n \to y_0 \in M$ , from the continuity of V we can infer that

$$\lim_{n \to \infty} V(\eta_{\varepsilon_n}) = V(y_0) = V_0,$$

which ends the proof of the Theorem 1.1.

**Acknowledgements.** The author would like to express his sincere thanks to the anonymous referee for his/her careful reading of the manuscript and valuable comments and suggestions.

The paper has been carried out under the auspices of the INdAM - GNAMPA Project 2017 titled: Teoria e modelli per problemi non locali.

### References

- [1] N. Ackermann, On a periodic Schrödinger equation with nonlocal superlinear part, Math. Z. 248 (2004) 423-443.
- [2] C. O. Alves and O. H. Miyagaki, Existence and concentration of solution for a class of fractional elliptic equation in  $\mathbb{R}^N$  via penalization method, Calc. Var. Partial Differential Equations 55 (2016), no. 3, Art. 47, 19 pp.
- [3] C. A. Alves and M. Yang, Existence of semiclassical ground state solutions for a generalized Choquard equation,
   J. Differ. Equ. 257 (2014), 4133-4164.
- [4] C. A. Alves and M. Yang, Investigating the multiplicity and concentration behaviour of solutions for a quasilinear Choquard equation via penalization method, Proc. Roy. Soc. Edinburgh Sect. A 146 (2016), 23–58.
- [5] A. Ambrosetti P. H. and Rabinowitz, Dual variational methods in critical point theory and applications, J. Funct. Anal. 14 (1973), 349–381.
- [6] V. Ambrosio, Multiplicity of positive solutions for a class of fractional Schrödinger equations via penalization method, Ann. Mat. Pura Appl. (4) 196 (2017), no. 6, 2043–2062.
- [7] V. Ambrosio, Ground states for superlinear fractional Schrödinger equations in R<sup>N</sup>, Ann. Acad. Sci. Fenn. Math. 41 (2016), no. 2, 745–756.
- [8] P. Belchior, H. Bueno, O. H. Miyagaki and G. A. Pereira, Remarks about a fractional Choquard equation: ground state, regularity and polynomial decay, Nonlinear Analysis 164 (2017), 38–53.
- [9] V. Benci and G. Cerami, Multiple positive solutions of some elliptic problems via the Morse theory and the domain topology, Calc. Var. Partial Differential Equations 2 (1994), no. 1, 29–48.
- [10] L.A. Caffarelli and L.Silvestre, An extension problem related to the fractional Laplacian, Comm. Partial Differential Equations 32 (2007),1245–1260.
- [11] Y. H. Chen and C. Liu, Ground state solutions for non-autonomous fractional Choquard equations, Nonlinearity 29 (2016), no. 6, 1827–1842.
- [12] V. Coti Zelati and M. Nolasco, Ground states for pseudo-relativistic Hartree equations of critical type, Rev. Mat. Iberoam. 29 (2013), no. 4, 1421–1436.
- [13] P. d'Avenia, G. Siciliano, and M. Squassina, On fractional Choquard equations, Math. Models Methods Appl. Sci. 25 (2015), no. 8, 1447–1476.
- [14] J. Dávila, M. del Pino and J. Wei, Concentrating standing waves for the fractional nonlinear Schrödinger equation, J. Differential Equations 256 (2014), no. 2, 858–892.
- [15] M. Del Pino and P.L. Felmer, Local Mountain Pass for semilinear elliptic problems in unbounded domains, Calc. Var. Partial Differential Equations, 4 (1996), 121–137.
- [16] E. Di Nezza, G. Palatucci and E. Valdinoci, Hitchhiker's guide to the fractional Sobolev spaces, Bull. Sci. math. 136 (2012), 521–573.
- [17] I. Ekeland, On the variational principle, J. Math. Anal. Appl. 47 (1974), 324–353.
- [18] M. M. Fall, M. Fethi and E. Valdinoci, Ground states and concentration phenomena for the fractional Schrödinger equation, Nonlinearity 28 (2015), no. 6, 1937–1961.
- [19] P. Felmer, A. Quaas and J. Tan, Positive solutions of the nonlinear Schrödinger equation with the fractional Laplacian, Proc. Roy. Soc. Edinburgh Sect. A 142 (2012), 1237–1262.
- [20] R.L. Frank and E. Lenzmann, On ground states for the  $L^2$ -critical boson star equation, preprint arXiv:0910.2721.
- [21] X. He and W. Zou, Existence and concentration result for the fractional Schrödinger equations with critical nonlinearities, Calc. Var. Partial Differential Equations 55 (2016), no. 4, Paper No. 91, 39 pp.
- [22] N. Laskin, Fractional Schrödinger equation, Phys. Rev. E (3) 66 (2002), no. 5, 056108, 7 pp. 81Q05.
- [23] E. H. Lieb Existence and uniqueness of the minimizing solution of Choquards nonlinear equation, Stud. Appl. Math. 57 (1977), 93–105.
- [24] E. H. Lieb and M. Loss, Analysis, Graduate Studies in Mathematics, 14. American Mathematical Society, Providence, RI, 1997. xviii+278 pp.
- [25] P.L. Lions, The Choquard equation and related questions, Nonlinear Anal. 4 (1980), 1063–1072.
- [26] L. Ma and L.Zhao Classification of positive solitary solutions of the nonlinear Choquard equation, Archive for Rational Mechanics and Analysis, 195 (2010), 455–467.
- [27] J. Mawhin and M. Willen, Critical Point Theory and Hamiltonian Systems, Springer-Verlag, 1989.
- [28] G. Molica Bisci, V. Rădulescu and R. Servadei, Variational Methods for Nonlocal Fractional Problems, Cambridge University Press, 162 Cambridge, 2016.
- [29] J. Moser, A new proof of De Giorgi's theorem concerning the regularity problem for elliptic differential equations, Comm. Pure Appl. Math. 13 (1960), 457–468.

- [30] V. Moroz and J. Van Schaftingen, Groundstates of nonlinear Choquard equations: existence, qualitative properties and decay asymptotics, J. Funct. Anal. 265 (2013), 153–84.
- [31] V. Moroz and J. Van Schaftingen, A guide to the Choquard equation, J. Fixed Point Theory Appl. 19 (2017), no. 1, 773–813.
- [32] G. Palatucci and A. Pisante, Improved Sobolev embeddings, profile decomposition, and concentration-compactness for fractional Sobolev spaces, Calc. Var. Partial Differential Equations 50 (2014), no. 3-4, 799–829.
- [33] S. Pekar, Untersuchung über die Elektronentheorie der Kristalle (Akademie Verlag, 1954).
- [34] R. Penrose, Quantum computation, entanglement and state reduction, Philos. Trans. Roy. Soc. 356 (1998) 1–13.
- [35] S. Secchi, A note on Schrödinger-Newton systems with decaying electric potential, Nonlinear Anal. 72 (2010), no. 9-10, 3842–3856.
- [36] S. Secchi, Ground state solutions for nonlinear fractional Schrödinger equations in  $\mathbb{R}^N$ , J. Math. Phys. **54** (2013), 031501.
- [37] Z. Shen, F. Gao, and M. Yang, Ground states for nonlinear fractional Choquard equations with general nonlinearities, Math. Methods Appl. Sci. 39 (2016), no. 14, 4082–4098.
- [38] L. Silvestre, Regularity of the obstacle problem for a fractional power of the Laplace operator, Comm. Pure Appl. Math., 60 (2007), no.1, 67–112.
- [39] A. Szulkin and T. Weth, The method of Nehari manifold, in Handbook of Nonconvex Analysis and Applications, edited by D. Y. Gao and D. Montreanu (International Press, Boston, 2010), pp. 597–632.
- [40] M. Yang, J. Zhang, and Y. Zhang, Multi-peak solutions for nonlinear Choquard equation with a general nonlinearity, Commun. Pure Appl. Anal. 16 (2017), no. 2, 493–512.
- [41] J. Wei and M. Winter, Strongly interacting bumps for the Schrödinger-Newton equation, J. Math. Phys., **50** (2009).
- [42] M. Willem, *Minimax theorems*, Progress in Nonlinear Differential Equations and their Applications, 24. Birkhäuser Boston, Inc., Boston, MA, 1996. x+162 pp.

DIPARTIMENTO DI SCIENZE PURE E APPLICATE (DISPEA), UNIVERSITÀ DEGLI STUDI DI URBINO 'CARLO BO' PIAZZA DELLA REPUBBLICA, 13 61029 URBINO (PESARO E URBINO, ITALY)

E-mail address: vincenzo.ambrosio@uniurb.it