

# Arrangements of Pseudocircles: On Circularizability\*

Stefan Felsner<sup>[0000-0002-6150-1998]</sup> and Manfred Scheucher<sup>[0000-0002-1657-9796]</sup>

Institut für Mathematik, Technische Universität Berlin, Germany  
{felsner,scheucher}@math.tu-berlin.de

**Abstract.** An arrangement of pseudocircles is a collection of simple closed curves on the sphere or in the plane such that every pair is either disjoint or intersects in exactly two crossing points. We call an arrangement intersecting if every pair of pseudocircles intersects twice. An arrangement is circularizable if there is a combinatorially equivalent arrangement of circles.

In this paper we present the results of the first thorough study of circularizability. We show that there are exactly four non-circularizable arrangements of 5 pseudocircles (one of them was known before). We also show that exactly three out of the 2131 digon-free intersecting arrangement of 6 pseudocircles are non-circularizable.

Most of our non-circularizability proofs depend on incidence theorems like Miquel's. In other cases we contradict circularizability by considering a continuous deformation where the circles of an assumed circle representation grow or shrink in a controlled way.

The claims that we have all non-circularizable arrangements with the given properties are based on a program that generated all arrangements up to a certain size. Given the complete lists of arrangements, we used heuristics to find circle representations. Examples where the heuristics failed were examined by hand.


**Keywords:** circularizability · incidence theorems · great-(pseudo)circles

## 1 Introduction

Arrangements of pseudocircles generalize arrangements of circles in the same vein as arrangements of pseudolines generalize arrangements of lines. The study of arrangements of pseudolines was initiated by Levi [12] in 1918. Since then arrangements of pseudolines were intensively studied. The handbook article on the topic [5] lists more than 100 references. To the best of our knowledge the study of arrangements of pseudocircles was initiated by Grünbaum [8].

---

\*Partially supported by the DFG Grants FE 340/11-1 and FE 340/12-1. Manfred Scheucher was partially supported by the ERC Advanced Research Grant no. 267165 (DISCONV). We gratefully acknowledge the computing time granted by TBK Automatisierung und Messtechnik GmbH and by the Institute of Software Technology, Graz University of Technology. We also thank the anonymous reviewers for valuable comments.



**Fig. 1:** The 3 connected arrangements of  $n = 3$  pseudocircles. (a) *Krupp*, (b) *NonKrupp*, (c) *3-Chain*.

A *pseudocircle* is a simple closed curve in the plane or on the sphere. An *arrangement of pseudocircles* is a collection of pseudocircles with the property that the intersection of any two of the pseudocircles is either empty or consists of two points where the curves cross. Other authors also allow touching pseudocircles, e.g. [1]. A cell of the arrangement with  $k$  crossings on its boundary is a  $k$ -*cell*. A 2-cell is also called a *digon* (some authors call it a *lens*), and a 3-cell is also called a *triangle*. An arrangement  $\mathcal{A}$  of pseudocircles is

***simple***, if no three pseudocircles of  $\mathcal{A}$  intersect in a common point;  
***connected***, if the graph of the arrangement is connected;  
***intersecting***, if any two pseudocircles of  $\mathcal{A}$  intersect.

In this paper we assume that arrangements are simple and connected.

Two arrangements  $\mathcal{A}$  and  $\mathcal{B}$  are *isomorphic* if they induce homeomorphic cell decompositions of the compactified plane, i.e., on the sphere. In particular, the isomorphism class of an arrangement of pseudocircles in the plane is closed under changes of the unbounded cell.

Figure 1 shows the three connected arrangements of three pseudocircles. We call the unique digon-free intersecting arrangement the *Krupp*<sup>1</sup>. The second intersecting arrangement is the *NonKrupp*; this arrangement has digons. The non-intersecting arrangement is the *3-Chain*.

Every triple of great-circles on the sphere induces a Krupp arrangement, hence, we call an arrangement of pseudocircles an *arrangement of great-pseudocircles* if every subarrangement induced by three pseudocircles is a Krupp.

Some authors think of arrangements of great-pseudocircles when they speak about arrangements of pseudocircles, this is e.g. common practice in the theory of oriented matroids. In fact, arrangements of great-pseudocircles serve to represent rank 3 oriented matroids, cf. [2].

**Definition 1.** *An arrangement of pseudocircles is circularizable if there is an isomorphic arrangement of circles.*

Preceding our work there have been only few results about circularizability of arrangements of pseudocircles. Edelsbrunner and Ramos [4] presented an intersecting arrangement of 6 pseudocircles (with digons) which has no realization

<sup>1</sup>This name refers to the logo of the Krupp AG, a German steel company. Krupp was the largest company in Europe at the beginning of the 20th century.

with circles, i.e., it is not circularizable. Linhart and Ortner [13] found a non-circularizable non-intersecting arrangement of 5 pseudocircles with digons (see Figure 2(b)). They also proved that every intersecting arrangement of at most 4 pseudocircles is circularizable. Kang and Müller [9] extended the result by showing that all arrangements with at most 4 pseudocircles are circularizable. They also proved that deciding circularizability for connected arrangements is NP-hard.

## 2 Overview

In Section 3 we present some background on arrangements of pseudocircles and provide tools that will be useful for non-circularizability proofs.

In Section 4 we study arrangements of great-pseudocircles – this class of arrangements of pseudocircles is in bijection with projective arrangements of pseudolines. Our main theorem in this section is the Great-Circle Theorem which allows to transfer knowledge regarding arrangements of pseudolines to arrangements of pseudocircles.

**Theorem 1 (Great-Circle Theorem).** *An arrangement of great-pseudocircles is circularizable (i.e., has a circle representation) if and only if it has a great-circle representation.*


In the last two sections we present the full classification of circularizable and non-circularizable arrangements among all connected arrangements of 5 pseudocircles and all digon-free intersecting arrangements of 6 pseudocircles. With the aid of computers we generated the complete lists of connected arrangements of  $n \leq 6$  pseudocircles and of intersecting arrangements of  $n \leq 7$  pseudocircles. The respective numbers are shown in Table 1. Given the complete lists of arrangements, we used automatized heuristics to find circle representations. Examples where the heuristics failed had to be examined by hand.

Computational issues and algorithmic ideas are omitted here – we refer the interested reader to the full version of this paper [7]. The encoded lists of arrangements of up to  $n = 6$  pseudocircles and circle representations are available on our webpage [6].

The list of circle representations together with the non-circularizability proofs given in Sections 5 yields the following theorem.

**Theorem 2.** *The four equivalence classes of arrangements  $\mathcal{N}_5^1$ ,  $\mathcal{N}_5^2$ ,  $\mathcal{N}_5^3$ , and  $\mathcal{N}_5^4$  (shown in Figure 2) are the only non-circularizable ones among the 984 equivalence classes of connected arrangements of  $n = 5$  pseudocircles.*

Note that  $\mathcal{N}_5^1$  is the only non-circularizable intersecting arrangement on 5 pseudocircles. Non-circularizability of  $\mathcal{N}_5^2$  was previously shown by Linhart and Ortner [13]. We give an alternative proof which also shows the non-circularizability of  $\mathcal{N}_5^3$ . Jonathan Wild and Christopher Jones, contributed sequences A250001 and A288567 to the On-Line Encyclopedia of Integer Sequences (OEIS). These



**Fig. 2:** The four non-circularizable arrangements on  $n = 5$  pseudocircles: (a)  $\mathcal{N}_5^1$ , (b)  $\mathcal{N}_5^2$ , (c)  $\mathcal{N}_5^3$ , and (d)  $\mathcal{N}_5^4$ .

$n$	3	4	5	6	7
<b>connected</b>	3	21	984	609 423	?
+digon-free	1	3	30	4 509	?
<b>intersecting</b>	2	8	278	145 058	447 905 202
+digon-free	1	2	14	2 131	3 012 972
<b>great-p.c.s</b>	1	1	1	4	11

**Table 1:** Number of combinatorially different arrangements of  $n$  pseudocircles.

sequences count certain classes of arrangements of circles and pseudocircles. Wild and Jones also looked at circularizability and independently found Theorem 2 (personal communication).


Concerning arrangements of 6 pseudocircles, we were able to fully classify digon-free intersecting arrangements.

**Theorem 3.** *The three equivalence classes of arrangements  $\mathcal{N}_6^\Delta$ ,  $\mathcal{N}_6^2$ , and  $\mathcal{N}_6^3$  (shown in Figure 3) are the only non-circularizable ones among the 2131 equivalence classes of digon-free intersecting arrangements of  $n = 6$  pseudocircles.*

In Section 6, we give non-circularizability proofs for  $\mathcal{N}_6^\Delta$ ,  $\mathcal{N}_6^2$ , and  $\mathcal{N}_6^3$ . In fact, for the non-circularizability of  $\mathcal{N}_6^\Delta$  and  $\mathcal{N}_6^2$ , respectively, we have two proofs of different flavors: One proof (see Section 6) uses continuous deformations similar to the proof of the Great-Circle Theorem (Theorem 1) and the other proof (omitted in this version) is based on an incidence theorem. The incidence theorem used for  $\mathcal{N}_6^\Delta$  might be new and of independent interest.

By enumerating and realizing all arrangement of  $n \leq 4$  pseudocircles, we have an alternative proof of Kang and Müller’s result, that all arrangements of  $n \leq 4$  are circularizable [9].

In the full version [7] we have further results, for example, non-circularizability proofs for some intersecting arrangements on  $n = 6$  pseudocircles with digons.



**Fig. 3:** The three non-circularizable digon-free intersecting arrangements for  $n = 6$ : (a)  $\mathcal{N}_6^A$ , (b)  $\mathcal{N}_6^2$ , and (c)  $\mathcal{N}_6^3$ . Inner triangles are colored gray. Note that in (b) and (c) the outer face is also a triangle.

### 3 Preliminaries: Basic Properties and Tools

Stereographic projections map circles to circles (if we consider a line to be a circle containing the point at infinity), therefore, circularizability on the sphere and in the plane is the same concept. Arrangements of circles can be mapped to isomorphic arrangements of circles via Möbius transformations. In this context, the sphere is identified with the extended complex plane  $\mathbb{C} \cup \{\infty\}$ .

Let  $\mathcal{C}$  be an arrangement of circles represented on the sphere. Each circle of  $\mathcal{C}$  spans a plane in 3-space, hence, we obtain an arrangement  $\mathcal{E}(\mathcal{C})$  of planes in  $\mathbb{R}^3$ . In fact, a fixed sphere  $S$  conveys a bijection between (not necessarily connected) circle arrangements on  $S$  and arrangements of planes with the property that each plane of the arrangement intersects  $S$ .

Consider two circles  $C_1, C_2$  of a circle arrangement  $\mathcal{C}$  on  $S$  and the corresponding planes  $E_1, E_2$  of  $\mathcal{E}(\mathcal{C})$ . The intersection of  $E_1$  and  $E_2$  is either empty (i.e.,  $E_1$  and  $E_2$  are parallel) or a line  $\ell$ . The line  $\ell$  intersects  $S$  if and only if  $C_1$  and  $C_2$  intersect, in fact,  $\ell \cap S = C_1 \cap C_2$ .


With three pairwise intersecting circles  $C_1, C_2, C_3$  we obtain three planes  $E_1, E_2, E_3$  intersecting in a vertex  $v$  of  $\mathcal{E}(\mathcal{C})$ . It is notable that  $v$  is in the interior of the ball bounded by  $S$  if and only if the three circles form a Krupp in  $\mathcal{C}$ .

**Lemma 1.** *Let  $\mathcal{C}$  be an arrangement of circles represented on the sphere. Three circles  $C_1, C_2, C_3$  of  $\mathcal{C}$  form a Krupp if and only if the three corresponding planes  $E_1, E_2, E_3$  intersect in a single point in the interior of the ball bounded by  $S$ .*

Digons are also nicely characterized: A pair  $C_1, C_2$  of circles forms a digon of  $\mathcal{C}$  if and only if the segment of  $\ell$  in the interior of  $S$  contains no vertex of  $\mathcal{E}(\mathcal{C})$ .

#### 3.1 Incidence Theorems

The smallest non-stretchable arrangements of pseudolines are closely related to the incidence theorems of Pappos and Desargues. A construction already



**Fig. 4:** (a) A non-stretchable arrangement of pseudolines from Pappos's Theorem. (b) A non-circularizable arrangement of pseudocircles from Miquel's Theorem.

described by Levi [12] is depicted in Figure 4(a). Pappos's Theorem states that, in a configuration of 8 lines as shown in the figure in black, the 3 white points are collinear, i.e., a line containing two of them also contains the third. Therefore, the arrangement including the red pseudoline has no corresponding arrangement of straight lines, i.e., it is not stretchable.

Miquel's Theorem asserts that, in a configuration of 5 circles as shown in Figure 4(b) in black, the 4 white points are cocircular, i.e., a circle containing three of them also contains the fourth. Therefore, the arrangement including the red pseudocircle cannot be circularized.

Next we state an incidence theorem that will be used in later proofs of non-circularizability. (For a proof, see Appendix A.)

**Lemma 2 (First Four-Circles Incidence Lemma).** *Let  $\mathcal{C}$  be an arrangement of four circles  $C_1, C_2, C_3, C_4$  such that none of them is contained in the interior of another one, and such that  $(C_1, C_2)$ ,  $(C_2, C_3)$ ,  $(C_3, C_4)$ , and  $(C_4, C_1)$  are touching. Then there is a circle  $C^*$  passing through these four touching points in the given cyclic order.*

In the course of the paper we will meet further incidence theorems, e.g. Lemma 4 and Lemma 5.

### 3.2 Flips and Deformations of Pseudocircles

Let  $\mathcal{C}$  be an arrangement of circles. Imagine that the circles of  $\mathcal{C}$  start moving independently, i.e., the position of their centers and their radii depend on a time parameter  $t$  in a continuous way. This yields a family  $\mathcal{C}(t)$  of arrangements with  $\mathcal{C}(0) = \mathcal{C}$ . Let us assume that the set  $T$  of all  $t$  for which  $\mathcal{C}(t)$  is not simple or contains touching circles is discrete and for each  $t \in T$  the arrangement  $\mathcal{C}(t)$  contains either a single point where 3 circles intersect or a single touching. If  $t_1 < t_2$  are consecutive in  $T$ , then all arrangements  $\mathcal{C}(t)$  with  $t \in (t_1, t_2)$  are isomorphic. Selecting one representative from each such class, we get a list  $\mathcal{C}_0, \mathcal{C}_1, \dots$  of simple arrangements such that two consecutive (non-isomorphic) arrangements  $\mathcal{C}_i, \mathcal{C}_{i+1}$  are either related by a triangle flip or by a digon flip.

We will make use of controlled changes in circle arrangements, in particular, we grow or shrink specified circles of an arrangement to produce touchings or points where 3 circles intersect. The following lemma will be of use frequently. (For a proof, see Appendix B.)

**Lemma 3 (Digon Collapse Lemma).** *Let  $\mathcal{C}$  be an intersecting arrangement of  $n \geq 3$  circles in the plane and let  $C$  be a circles from  $\mathcal{C}$ . If  $C$  has no incident triangle in its interior, then we can shrink  $C$  into its interior such that the combinatorics of the arrangement remain the same except that two digons collapse to touchings. Moreover, the two corresponding circles touch  $C$  from the outside.*

In the following we will sometimes use the dual version of the lemma, whose statement is obtained from the Digon Collapse Lemma by changing interior to exterior and outside to inside. The validity of the dual lemma is seen by applying a Möbius transformation which exchanges interior and exterior of  $C$ .


Triangle flips and digon flips are also central to the work of Snoeyink and Hershberger [17]. They have shown that an arrangement  $\mathcal{C}$  of pseudocircles can be swept with a sweepfront  $\gamma$  starting at any pseudocircle  $C \in \mathcal{C}$ , i.e.,  $\gamma_0 = C$ . The sweep consists of two stages, one for sweeping the interior of  $C$ , the other for sweeping the exterior. At any fixed time  $t$  the sweepfront  $\gamma_t$  is a closed curve such that  $\mathcal{C} \cup \{\gamma_t\}$  is an arrangement of pseudocircles. Moreover, this arrangement is simple except for a discrete set  $T$  of times where sweep events happen. The sweep events are triangle flips or digon flips involving  $\gamma_t$ .

### 4 Arrangements of Great-Pseudocircles

Central projections map between arrangements of great-circles on a sphere  $S$  and arrangements of lines on a plane. Changes of the plane preserve the isomorphism class of the projective arrangement of lines. In fact, arrangements of lines in the projective plane are in one-to-one correspondence to arrangements of great-circles.

In this section we generalize this concept to arrangements of pseudolines and show that there is a one-to-one correspondence to arrangements of great-pseudocircles. As already mentioned, this correspondence is not new (see e.g. [2]).

An Euclidean arrangement of  $n$  pseudolines can be represented by  $x$ -monotone pseudolines, see e.g [5]. As illustrated in Figure 5, an  $x$ -monotone representation can be glued with a mirrored copy of itself to form an arrangement of  $n$  pseudo-circles. The resulting arrangement is intersecting and has no NonKrupp subarrangement, hence, it is an arrangement of great-pseudocircles.



**Fig. 5:** Obtaining an arrangement of great-pseudocircles from an Euclidean arrangement  $\mathcal{L}$  of pseudolines and its mirrored copy. The gray boxes highlight the arrangement  $\mathcal{L}$  and its mirrored copy.

For a pseudocircle  $C$  of an arrangement of  $n$  great-pseudocircles the cyclic order of crossings on  $C$  is *antipodal*, i.e., the infinite sequence corresponding to the cyclic order crossings of  $C$  with the other pseudocircles is periodic of order  $n - 1$ . If we consider projections of projective arrangements of  $n$  pseudolines, then this order does not depend on the choice of the projection. In fact, projective arrangements of  $n$  pseudolines are in bijection with arrangements of  $n$  great-pseudocircles.

Projective arrangements of pseudolines are also known as projective abstract order types or oriented matroids of rank 3. The precise numbers of such arrangements are known for  $n \leq 11$ , see [10,11]. Hence the numbers of great-pseudocircle arrangements given in Table 1 are not new.

#### 4.1 The Great-Circle Theorem and its Applications


Let  $\mathcal{A}$  be an arrangement of great-pseudocircles and let  $\mathcal{L}$  be the corresponding projective arrangement of pseudolines. Central projections show that, if  $\mathcal{L}$  is realizable with straight lines, then  $\mathcal{A}$  is realizable with great-circles, and conversely. In fact, due to Theorem 1, it is sufficient that  $\mathcal{A}$  is circularizable to conclude that  $\mathcal{A}$  is realizable with great-circles and  $\mathcal{L}$  is realizable with straight lines.

*Proof (of Theorem 1).* Consider an arrangement of circles  $\mathcal{C}$  on the unit sphere  $S$  that realizes an arrangement of great-pseudocircles. Let  $\mathcal{E}(\mathcal{C})$  be the arrangement of planes spanned by the circles of  $\mathcal{C}$ . Since  $\mathcal{C}$  realizes an arrangement of great-pseudocircles, every triple of circles forms a Krupp, hence, the point of intersection of any three planes of  $\mathcal{E}(\mathcal{C})$  is in the interior of  $S$ .

Imagine the planes of  $\mathcal{E}(\mathcal{C})$  moving towards the origin. To be precise, for time  $t \geq 1$  let  $\mathcal{E}_t := \{1/t \cdot E : E \in \mathcal{E}(\mathcal{C})\}$ . Since all intersection points of the initial arrangement  $\mathcal{E}_1 = \mathcal{E}(\mathcal{C})$  are in the interior of the sphere  $S$ , the circle arrangement obtained by intersecting the moving planes  $\mathcal{E}_t$  with the sphere  $S$  remains the same (isomorphic). Moreover, every circle in this arrangement converges to a great-circle as  $t \rightarrow +\infty$ , and the statement follows.  $\square$

From the theorem it follows that an arrangement of pseudolines is stretchable if and only if the corresponding arrangement of great-pseudocircles is circularizable. Since deciding stretchability is known to be  $\exists\mathbb{R}$ -complete (see e.g. [14,15]), the hardness of stretchability directly carries over to hardness of circularizability.

It is known that all (not necessarily simple) arrangements  $n \leq 8$  pseudolines are stretchable and that the simple non-Pappos arrangement is the unique non-stretchable simple projective arrangements of 9 pseudoline, see e.g. [5]. This again carries over to arrangements of great-pseudocircles. Bokowski and Sturmfels [3] have shown that infinite families of minimal non-stretchable arrangements of pseudolines exist, i.e., non-stretchable arrangements where every proper subarrangement is stretchable. Again, this carries over to arrangement of pseudocircles.



**Fig. 6:** An illustration of the non-circularizability proof of  $\mathcal{N}_5^1$ . The auxiliary circle  $C^*$  is drawn dashed.

## 5 Arrangements of 5 Pseudocircles


We have realizations<sup>2</sup> as circle arrangements of all 984 connected arrangements of 5 pseudocircles except for the four arrangements mentioned in Theorem 2. Since all arrangements with  $n \leq 4$  pseudocircles have representations as arrangements of circles, there are no disconnected non-circularizable examples with  $n \leq 5$ . Hence, the four arrangements  $\mathcal{N}_5^1$ ,  $\mathcal{N}_5^2$ ,  $\mathcal{N}_5^3$ , and  $\mathcal{N}_5^4$  are the only candidates for non-circularizability. In this section, we show that they are indeed not circularizable. This proves Theorem 2.

For the non-circularizability proof of  $\mathcal{N}_5^1$  we need the following additional incidence lemma. (A proof is given in Appendix C.)

**Lemma 4 (Second Four-Circles Incidence Lemma).** *Let  $\mathcal{C}$  be an arrangement of four circles  $C_1, C_2, C_3, C_4$  such that every pair of them is touching or forms a digon, and every circle is involved in at least two touchings. Then there is a circle  $C^*$  passing through the digon or touching point of each of the following pairs of circles  $(C_1, C_2)$ ,  $(C_2, C_3)$ ,  $(C_3, C_4)$ , and  $(C_4, C_1)$  in this cyclic order.*

*Proof (non-circularizability of  $\mathcal{N}_5^1$ ).* Suppose for a contradiction that there is an equivalent arrangement  $\mathcal{C}$  of circles. We apply the Digon Collapse Lemma (Lemma 3) to shrink  $C_2$ ,  $C_3$ , and  $C_4$  into their respective interiors. We also use the dual of the Digon Collapse Lemma for  $C_1$ . In the resulting subarrangement  $\mathcal{C}'$  formed by these four transformed circles  $C'_1, C'_2, C'_3, C'_4$ , each of the four circles is involved in at least two touchings. Moreover, since the intersection of  $C'_i$  and  $C'_j$  in  $\mathcal{C}'$  is contained in the intersection of  $C_i$  and  $C_j$  in  $\mathcal{C}$ , each of the four points  $p_{12}$ ,  $p_{23}$ ,  $p_{34}$ , and  $p_{41}$  lies in the original digons of  $\mathcal{C}$  which respectively are touching points or points from the digons of  $(C'_1, C'_2)$ ,  $(C'_2, C'_3)$ ,  $(C'_3, C'_4)$ , and  $(C'_4, C'_1)$ . It follows that the circle  $C_5$  has  $p_{12}$  and  $p_{34}$  in its interior but  $p_{23}$  and  $p_{41}$  in its exterior. Figure 6 gives an illustration.

<sup>2</sup> Representations of these arrangements are available at our website [6].



**Fig. 7:** An illustration of the non-circularizability proofs of (a)  $\mathcal{N}_5^2$ , (b)  $\mathcal{N}_5^3$ , and (c)  $\mathcal{N}_5^4$ . The auxiliary circle  $C^*$  is drawn dashed.

By applying Lemma 4 to  $\mathcal{C}'$  we obtain a circle  $C^*$  which passes through the points  $p_{12}$ ,  $p_{23}$ ,  $p_{34}$ , and  $p_{41}$  (in this order). Now the two circles  $C_5$  and  $C^*$  intersect in four points. This is impossible, and hence  $\mathcal{N}_5^4$  is not circularizable.  $\square$

*Proof (non-circularizability of  $\mathcal{N}_5^2$  and  $\mathcal{N}_5^3$ ).* Suppose for a contradiction that there is an equivalent arrangement  $\mathcal{C}$  of circles. We label the circles as illustrated in Figures 7(a) and 7(b). Since  $C_5$  is not incident to any digon, applying the Digon Collapse Lemma (Lemma 3) to  $C_1$  and  $C_3$  yields an arrangement  $\mathcal{C}'$  with four touching points  $p_{12}, p_{23}, p_{34}, p_{41}$ , where  $p_{ij}$  is the touching of  $C'_i$  and  $C'_j$ . To be precise, since  $\mathcal{C}$  is not intersecting, we first shrink  $C_1$  in the arrangement  $\mathcal{C} \setminus \{C_3\}$  and then  $C_3$  in the arrangement  $\mathcal{C} \setminus \{C_1\}$ . Since the respective interiors of  $C_1$  and  $C_3$  are disjoint, we obtain the desired arrangement.

From Lemma 2 it follows that there is a circle  $C^*$  which passes through the points  $p_{12}$ ,  $p_{23}$ ,  $p_{34}$ , and  $p_{41}$  in this cyclic order. Since the point  $p_{ij}$  lies inside the digon formed by  $C_i$  and  $C_j$  in the arrangement  $\mathcal{C}$ , it follows that the circle  $C_5$  has  $p_{12}, p_{34}$  in its interior and  $p_{23}, p_{41}$  in its exterior. Therefore, the two circles  $C_5$  and  $C^*$  intersect in four points. This is impossible and, therefore,  $\mathcal{N}_5^2$  and  $\mathcal{N}_5^3$  are not circularizable.  $\square$

To prove the non-circularizability of  $\mathcal{N}_5^4$ , we make use of the following incidence lemma. (A proof of this lemma can be found in Appendix D.)

**Lemma 5 (Third Four-Circles Incidence Lemma).** *Let  $\mathcal{C}$  be an arrangement of four circles  $C_1, C_2, C_3, C_4$  such that  $(C_1, C_2)$ ,  $(C_2, C_3)$ ,  $(C_3, C_4)$ , and  $(C_4, C_1)$  are touching, moreover,  $C_4$  is in the interior of  $C_1$  and the exterior of  $C_3$ , and  $C_2$  is in the interior of  $C_3$  and the exterior of  $C_1$ , see Figure 12. Then there is a circle  $C^*$  passing through the four touching points in the given cyclic order.*

*Proof (non-circularizability of  $\mathcal{N}_5^4$ ).* Suppose there is an isomorphic arrangement of circles  $\mathcal{C}$ . Referring to the labeling as shown in Figure 7(c) we shrink the circles  $C_2$  and  $C_4$  such that the pairs  $(C_1, C_2)$ ,  $(C_2, C_3)$ ,  $(C_3, C_4)$ ,  $(C_4, C_1)$  (which form digons in  $\mathcal{C}$ ) touch. With these touchings the four circles  $C_1, C_2, C_3, C_4$  form the configuration of Lemma 5. Hence there is a circle  $C^*$  containing the four touching points in the given cyclic order. Now the two circles  $C^*$  and  $C_5$  intersect in four points. This is impossible and, therefore,  $\mathcal{N}_5^4$  is not circularizable.  $\square$

## 6 Arrangements of 6 Pseudocircles

We have realizations<sup>2</sup> as circle arrangements of all 2131 intersecting digon-free arrangements of 6 pseudocircles except for the three arrangements mentioned in Theorem 3. In the following, we present one of our two non-circularizability proofs for  $\mathcal{N}_6^{\Delta}$  and  $\mathcal{N}_6^2$ , respectively, and our non-circularizability proof for  $\mathcal{N}_6^3$ . The other non-circularizability proofs can be found in the full version [7].

Since  $\mathcal{N}_6^{\Delta}$  (shown in Figure 3(a)) is intersecting, digon-free, and each of the eight triangles of  $\mathcal{N}_6^{\Delta}$  is a NonKrupp, the non-circularizability of  $\mathcal{N}_6^{\Delta}$  is an immediate consequence of the following theorem:

**Theorem 4.** *Let  $\mathcal{A}$  be a connected digon-free arrangement of pseudocircles. If every triple of pseudocircles which forms a triangle is NonKrupp, then  $\mathcal{A}$  is not circularizable.*

*Proof.* Assume for a contradiction that there exists an isomorphic arrangement of circles  $\mathcal{C}$  on the unit sphere  $S$ . Let  $\mathcal{E}(\mathcal{C})$  be the arrangements of planes spanned by the circles of  $\mathcal{C}$ . Imagine the planes of  $\mathcal{E}(\mathcal{C})$  moving away from the origin. To be precise, for time  $t \geq 1$  let  $\mathcal{E}_t := \{t \cdot E : E \in \mathcal{E}(\mathcal{C})\}$ . Consider the arrangement induced by intersecting the moving planes  $\mathcal{E}_t$  with the sphere  $S$ . Since  $\mathcal{C}$  has Non-Krupp triangles, it is not a great-circle arrangement and some planes of  $\mathcal{E}(\mathcal{C})$  do not contain the origin. All planes from  $\mathcal{E}(\mathcal{C})$ , which do not contain the origin, will eventually lose the intersection with  $S$ , hence some event has to happen.

When the isomorphism class of the intersection of  $\mathcal{E}_t$  with  $S$  changes, we see a triangle flip, or a digon flip, or some isolated circle disappears. Since initially there is no digon and no isolated circle, the first event is a triangle flip. Triangles of  $\mathcal{C}$  correspond to NonKrupp subarrangements, hence, the intersection point of their planes is outside of  $S$  (Lemma 1). This shows that a triangle flip event is also impossible. This contradiction implies that  $\mathcal{A}$  is non-circularizable.  $\square$


The following theorem is of the same flavor and directly implies the non-circularizability of  $\mathcal{N}_6^2$  (shown in Figure 3(b)), since  $\mathcal{N}_6^2$  is intersecting, not an arrangement of great-pseudocircles, and since each triangle in  $\mathcal{N}_6^2$  is Krupp.

**Theorem 5.** *Let  $\mathcal{A}$  be an intersecting arrangement of pseudocircles which is not an arrangement of great-pseudocircles. If every triple of pseudocircles which forms a triangle is Krupp, then  $\mathcal{A}$  is not circularizable.*

We outline the proof: Suppose a realization of  $\mathcal{A}$  exists on the sphere. If we grow the sphere over time, the arrangement will eventually become (isomorphic to) an arrangement of great-circles. We consider the first event that occurs. As the sphere grows, no digon is collapsed. Since  $\mathcal{A}$  is intersecting, no digon is created, and, since all triangles are Krupp, the corresponding intersection points of their planes lie inside  $S$ . Therefore, no event can occur – a contradiction.

We remark that there is one additional arrangement of 6 pseudocircles where non-circularizability can be proved by using Theorem 5.

The arrangement  $\mathcal{N}_6^3$  is shown in Figure 3(c) and Figure 8(b). To prove its non-circularizability, we again use an incidence lemma.



**Fig. 8:** (a) An illustration for Lemma 6. (b) The non-circularizable arrangement  $\mathcal{N}_6^2$  with 3 dashed pseudolines illustrating the proof.

**Lemma 6 (Richter-Gebert [16]).** *Let  $\ell_1, \ell_2, \ell_3$  be lines,  $C_1, C_2, C_3$  be circles, and  $p_1, p_2, p_3, q_1, q_2, q_3$  be points, such that for  $\{i, j, k\} = \{1, 2, 3\}$  point  $p_i$  is incident to line  $\ell_i$ , circle  $C_j$ , and circle  $C_k$ , while point  $q_i$  is incident to circle  $C_i$ , line  $\ell_j$ , and line  $\ell_k$ . Then  $C_1, C_2$ , and  $C_3$  have a common point of intersection.*

This lemma was mentioned by Richter-Gebert as a relative of Pappos's Theorem, cf. [16, page 26]. Figure 8(a) gives an illustration.

*Proof (non-circularizability of  $\mathcal{N}_6^3$ ).* Suppose that  $\mathcal{N}_6^3$  has a representation  $\mathcal{C}$  as a circle arrangement in the plane. We refer to circles and intersection points via the label of the corresponding object in Figure 8(b). As in the figure, we assume without loss of generality that the triangular cell spanned by  $C_4, C_5$ , and  $C_6$  is the outer cell of the arrangement.

Consider the region  $R := R_{24} \cup R_{35}$  where  $R_{ij}$  denotes the intersection of the respective interiors of  $C_i$  and  $C_j$ . The two straight-line segments  $p_1p'_1$  and  $p_3p'_3$  are fully contained in  $R_{35}$  and  $R_{24}$ , respectively, and have alternating end points along the boundary of  $R$ , hence they cross inside the region  $R_{24} \cap R_{35}$ .

From rotational symmetry we obtain that the three straight-line segments  $p_1p'_1$ ,  $p_2p'_2$ , and  $p_3p'_3$  intersect pairwise.

For  $i = 1, 2, 3$ , let  $\ell_i$  denote the line spanned by  $p_i$  and  $p'_i$ , let  $q_i$  denote the intersection-point of  $\ell_{i+1}$  and  $\ell_{i+2}$ , and let  $C'_i$  denote the circle spanned by  $q_i, p_{i+1}, p_{i+2}$  (indices modulo 3). Note that  $\ell_i$  contains  $p_i, q_{i+1}, q_{i+2}$ . These are precisely the conditions for the incidences of points, lines, and circles in Lemma 6. Hence, the three circles  $C'_1, C'_2$ , and  $C'_3$  intersect in a common point.

Let  $T$  be the triangle with corners  $p_1, p_2, p_3$ . Since  $p_2$  and  $p_3$  are on  $C_1$ , and  $q_1$  lies inside of  $C_1$ , we find that the intersection of the interior of  $C'_1$  with  $T$  is a subset of the intersection of the interior of  $C_1$  with  $T$ . The respective containments also hold for  $C'_2$  and  $C_2$  and for  $C'_3$  and  $C_3$ . Moreover, since  $C'_1, C'_2$ , and  $C'_3$  intersect in a common point, the union of the interiors of  $C'_1, C'_2$ , and  $C'_3$  contains  $T$ . Hence, the union of interiors of the  $C_1, C_2$ , and  $C_3$  also contains  $T$ . This shows that in  $\mathcal{C}$  there is no face corresponding to the gray triangle; see Figure 8(b). This contradicts the assumption that  $\mathcal{C}$  is a realization of  $\mathcal{N}_6^3$ , whence the arrangement is not circularizable.  $\square$

## References

1. Agarwal, P.K., Nevo, E., Pach, J., Pinchasi, R., Sharir, M., Smorodinsky, S.: Lenses in Arrangements of Pseudo-circles and Their Applications. *Journal of the ACM* **51**(2), 139–186 (2004). <https://doi.org/10.1145/972639.972641>
2. Björner, A., Las Vergnas, M., White, N., Sturmfels, B., Ziegler, G.M.: *Oriented Matroids*, Encyclopedia of Mathematics and its Applications, vol. 46. Cambridge University Press, 2 edn. (1999). <https://doi.org/10.1017/CBO9780511586507>
3. Bokowski, J., Sturmfels, B.: An Infinite Family of Minor-Minimal Non-realizable 3-chirotopes. *Mathematische Zeitschrift* **200**(4), 583–589 (1989). <https://doi.org/10.1007/BF01160956>
4. Edelsbrunner, H., Ramos, E.A.: Inclusion-Exclusion Complexes for Pseudodisk Collections. *Discrete & Computational Geometry* **17**, 287–306 (1997). <https://doi.org/10.1007/PL00009295>
5. Felsner, S., Goodman, J.E.: Pseudoline Arrangements. In: Toth, O'Rourke, Goodman (eds.) *Handbook of Discrete and Computational Geometry*. CRC Press, 3 edn. (2018). <https://doi.org/10.1201/9781315119601>
6. Felsner, S., Scheucher, M.: Webpage: Homepage of Pseudocircles, <http://www3.math.tu-berlin.de/pseudocircles>
7. Felsner, S., Scheucher, M.: Arrangements of Pseudocircles: On Circularizability. <http://arXiv.org/abs/1712.02149> (2017)
8. Grünbaum, B.: Arrangements and Spreads, CBMS Regional Conference Series in Mathematics, vol. 10. AMS (1972 (reprinted 1980)). <https://doi.org/10.1090/cbms/010>
9. Kang, R.J., Müller, T.: Arrangements of Pseudocircles and Circles. *Discrete & Computational Geometry* **51**, 896–925 (2014). <https://doi.org/10.1007/s00454-014-9583-8>
10. Knuth, D.E.: *Axioms and Hulls*, LNCS, vol. 606. Springer (1992). [https://doi.org/10.1007/3-540-55611-7\\_1](https://doi.org/10.1007/3-540-55611-7_1)
11. Krasser, H.: Order Types of Point Sets in the Plane. Ph.D. thesis, Institute for Theoretical Computer Science, Graz University of Technology, Austria (2003)
12. Levi, F.: Die Teilung der projektiven Ebene durch Gerade oder Pseudogerade. *Berichte über die Verhandlungen der sächsischen Akademie der Wissenschaften, Leipzig, Mathematisch-physische Klasse* **78**, 256–267 (1926)
13. Linhart, J., Ortner, R.: An Arrangement of Pseudocircles Not Realizable with Circles. *Beiträge zur Algebra und Geometrie* **46**(2), 351–356 (2005). <https://doi.org/10.1007/s00454-014-9583-8>
14. Matoušek, J.: Intersection graphs of segments and  $\exists\mathbb{R}$  (2014), <http://arXiv.org/abs/1406.2636>
15. Mnëv, N.E.: The universality theorems on the classification problem of configuration varieties and convex polytopes varieties. In: Viro, O.Y., Vershik, A.M. (eds.) *Topology and Geometry — Rohlin Seminar*. LNM, vol. 1346, pp. 527–543. Springer (1988). <https://doi.org/10.1007/bfb0082792>
16. Richter-Gebert, J.: *Perspectives on Projective Geometry – A Guided Tour through Real and Complex Geometry*. Springer (2011). <https://doi.org/10.1007/978-3-642-17286-1>
17. Snoeyink, J., Hershberger, J.: Sweeping Arrangements of Curves. In: Goodman, Pollack, Steiger (eds.) *Discrete & Computational Geometry*, DIMACS Series in Discrete Mathematics and Theoretical Computer Science, vol. 6, pp. 309–349. AMS (1991). <https://doi.org/10.1145/73833.73872>

### A Proof of Lemma 2




Fig. 9: An illustration for the proof of Lemma 2.

*Proof (of Lemma 2).* Apply a Möbius transformation  $\Gamma$  that maps the touching point of  $C_1$  and  $C_2$  to the point  $\infty$  of the extended complex plane. This maps  $C_1$  and  $C_2$  to a pair  $L_1, L_2$  of parallel lines and the discs of  $C_1$  and  $C_2$  are mapped to disjoint halfplanes. We may assume that  $L_1$  and  $L_2$  are horizontal and that  $L_1$  is above  $L_2$ . Circles  $C_3$  and  $C_4$  are mapped to touching circles  $C'_3$  and  $C'_4$ . Moreover,  $C'_3$  is touching  $L_2$  from above and  $C'_4$  is touching  $L_1$  from below. Figure 9 shows a sketch of the situation.

Let  $\ell'$  be the line, which is tangent to  $C'_3$  and  $C'_4$  at their touching point  $p$ . Consider the two segments from  $p$  to  $C'_3 \cap L_2$  and from  $p$  to  $C'_4 \cap L_1$ . Elementary considerations show the following equalities of angles:  $\alpha_3 = \alpha_4$ ,  $\beta_3 = \beta_4$ ,  $\gamma_3 = \gamma_4$ , and  $\gamma_3 = \gamma_4$ . Hence, there is a line  $\ell$  containing the images of the four touchings points. Consequently, the circle  $C^* = \Gamma^{-1}(\ell)$  contains the four touching points of  $\mathcal{C}$ , i.e., they are cocircular.

### B Proof of Lemma 3




Fig. 10: An illustration of the Digon Collapse Lemma.

*Proof.* As illustrated in Figure 10, we shrink the radius of  $C$  until the first flip occurs. Since  $C$  has no incident triangles in the interior, the flip must be a digon flip. Since  $\mathcal{C}$  is intersecting, it is impossible that a new digon is created, i.e.,  $C$  would lose the intersection to the touching circle from further shrinking, whence the circle is touching  $C$  from the outside. (Note that several digons might collapse at the same time.)

If  $C$  has only one touching point  $p$ , we shrink the radius and simultaneously move the center towards  $p$  such that  $p$  stays a touching until a second digon becomes a touching. Again the touching is from the outside of  $C$ .

### C Proof of Lemma 4





Fig. 11: Illustration for the proof of Lemma 4.

*Proof.* We first deal with the case where  $C_1$  and  $C_3$  form a digon. The assumptions imply that there is at most one further digon which might then be formed by  $C_2$  and  $C_4$ . In particular, the four pairs mentioned in the statement of the lemma form touchings and, as illustrated in the first row of Figure 11, we find a circle  $C^*$  that is incident to those four touching points. In the following let  $p_{ij}$  denote the touching point of  $C_i$  and  $C_j$ .

Think of the circles as being in the extended complex plane. Apply a Möbius transformation  $\Gamma$  that maps one of the points of intersection of  $C_1$  and  $C_3$  to the point  $\infty$ . This maps  $C_1$  and  $C_3$  to a pair of lines. The images of  $C_2$  and  $C_4$  are circles which touch the two lines corresponding to  $C_1$  and  $C_3$  and mutually either touch or form a digon. The first row of Figure 11 gives an illustration. Since the centers of  $C_2$  and  $C_4$  lie on the bisector  $\ell$  of the lines  $\Gamma(C_1)$  and  $\Gamma(C_3)$ , the touchings of  $C_2$  and  $C_4$  are symmetric with respect to  $\ell$ . Therefore, there is a circle  $C$  with center on  $\ell$  that contains the images of the four points  $p_{12}$ ,



**Fig. 12:** An illustration for the proof of Lemma 5.

$p_{23}$ ,  $p_{34}$ , and  $p_{41}$ . The circle  $C^* = \Gamma^{-1}(C)$  contains the four points, i.e., they are cocircular.

If  $C_1$  and  $C_2$  form a digon then there is at most one further digon formed by  $C_3$  and  $C_4$ . Again apply a Möbius transformation  $\Gamma$  that sends  $p_{13}$  to  $\infty$ . This maps  $C_1$  and  $C_3$  to parallel lines, each touched by one of  $C_2$  and  $C_4$ . The second row of Figure 11 shows that there is a circle  $C$  such that  $C^* = \Gamma^{-1}(C)$  has the claimed property.

## D Proof of Lemma 5

*Proof.* Since  $C_1$  is touching  $C_2$  and  $C_4$  which are respectively inside and outside  $C_3$  the two circles  $C_1$  and  $C_3$  intersect. Apply a Möbius transformation  $\Gamma$  that maps a crossing point of  $C_1$  and  $C_3$  to the point  $\infty$  of the extended complex plane. This maps  $C_1$  and  $C_3$  to a pair  $L_1, L_3$  of lines. The images  $C'_2, C'_4$  of  $C_2$  and  $C_4$  are separated by the lines  $L_1, L_3$  and each of them is touching both lines. Figure 12 illustrates the situation. The figure also shows that a circle  $C'$  through the four touching points exists. The circle  $C^* = \Gamma^{-1}(C')$  has the claimed properties.