

A new 24+24 matter multiplet of N=2 conformal supergravity

Subramanya Hegde,^a Ivano Lodato^b and Bindusar Sahoo^a

^a*Indian Institute of Science Education and Research, Vithura, Thiruvananthapuram - 695551, India*

^b*Department of Physics and Center for Field Theory and Particle Physics, Fudan University, 220 Handan Road, 200433 Shanghai, China*

E-mail: smhegde14@iisertvm.ac.in, ilodato@fudan.edu.cn,
bsahoo@iisertvm.ac.in

ABSTRACT: We write down the 48+48 component multiplet of supercurrents for a rigid N=2 tensor multiplet in four spacetime dimensions along with its supersymmetry transformation law. This gives us the transformation of the linearized supergravity multiplet which couples to this supercurrent multiplet. At the linearized level, this 48+48 component supergravity multiplet decouples into the 24+24 component linearized standard Weyl multiplet known in the literature and a new 24+24 component irreducible multiplet that contains a dilaton and dilatino. By a consistent application of the supersymmetry algebra with field dependent structure constants appropriate to N=2 conformal supergravity, we find the full transformation law for this newly found multiplet. We present a set of constraints which can be consistently imposed on the new multiplet to obtain a restricted minimal 8+8 off-shell matter multiplet. We also show as an example the precise embedding of the tensor multiplet inside the new multiplet.

ARXIV EPRINT: [1712.xxxxx](https://arxiv.org/abs/1712.xxxxx)

Contents

1	Introduction	1
2	N=2 conformal supergravity in four spacetime dimensions	3
3	The dilaton matter multiplet	5
4	Tensor multiplet and the multiplet of supercurrents	8
5	Linearized supergravity multiplet	10
6	Constraining the dilaton matter multiplet	15
7	The embedding of the tensor multiplet in the dilaton matter multiplet	18
8	Conclusions and future directions	19

1 Introduction

The use of conformal symmetries plays a crucial role in the construction of theories of supergravity. While the physical Poincaré theory can be obtained via a simple gauge-fixing procedure, the higher degree of symmetry in the conformal theories allows for a rigorous rearrangement of the off-shell degrees of freedom within multiplets shorter than the one that contains Poincaré supergravity degrees of freedom. Furthermore, techniques generally referred as “multiplet calculus” have been developed over the years to simplify the often massive task of constructing invariant couplings of supergravity, especially higher derivative ones.

Off-shell conformal supergravities exist for dimensions $D \leq 6$. The higher dimensional theories are typically related to the lower dimensional theory via dimensional reduction. For a connection between the $N = (1,0)$ chiral theory in 6 dimensions as well as $N = 2$ theory in four dimensions with the $N = 1$ theory in five dimensions¹ via dimensional reduction (uplift) see [4] and [5] respectively. Hence, one can expect that given a result in higher dimensional theories, a similar result would exist in the lower dimensional theory from dimensional reduction arguments. A relevant example of such connection is the existence of the two versions of Weyl multiplets: the standard and dilaton Weyl, which were constructed in six dimensions [6] and later on in five dimensions [1, 2], also exists in the four dimensional case [7–10]. While the precise reduction procedure and identification of off-shell degrees of freedom

¹We refer to the minimal supergravity in five dimensions formulated in terms of eight supercharges as N=1. However some people, for example [1–3], refer to it as N=2.

is being currently worked out, it is easy to realize that the reduction of the 32+32 off-shell five dimensional dilaton Weyl multiplet would give the 24+24 four dimensional dilaton-Weyl multiplet plus a short 8+8 vector matter multiplet in four dimensions .

It is worth pointing out that the dilaton-Weyl multiplet could be obtained in two different yet equivalent ways in five dimensions. The first procedure involves analyzing the coupling of a standard Weyl multiplet with an on-shell vector multiplet [1]. The vanishing of the vector multiplet equations of motion furnishes a set of constraints, which can be used to fix the auxiliary fields of the standard Weyl multiplet in terms of the auxiliary fields of the dilaton Weyl multiplet. The same procedure is used in a forthcoming paper [10] to obtain the $N = 2$ dilaton-Weyl multiplet in four dimensions.

The second method in five dimensions was based on the construction of the current multiplet for a non-conformal vector multiplet. The standard action for a non conformal vector multiplet in five dimensions would not be scale invariant and hence the trace of the energy momentum tensor $\sigma = \theta^\mu_\mu$ would be a non trivial component of the current multiplet. When coupling the current multiplet to linearized gravity (Weyl) multiplet, the scalar current σ will couple to a scalar field of dimension 1: the dilaton. From the linearized gravity multiplet it is in principle a simple, yet computationally challenging, task to derive the full non-linear transformations of the dilaton version of the Weyl multiplet. It is worth emphasizing that, in this method, the non-conformal nature of the matter multiplet chosen was crucial to obtain the Weyl multiplet with a dilaton.

One naïvely expects the same method to produce the dilaton Weyl multiplet in four dimensions, the only caveat being that the vector multiplet in four dimensions is conformally invariant. Hence a different off-shell non-conformal multiplet needs to be chosen for the construction of the current multiplet, such as the tensor multiplet. Nevertheless, the naïve expectations turns out to be incorrect.

Specifically we find that the linearized multiplet of gravity that couples to the current multiplet of the tensor multiplet has 48+48 components, which can be arranged into the standard Weyl multiplet and a new irreducible and unconstrained off-shell 24+24 matter multiplet. We will henceforth refer to it as the “dilaton matter multiplet” for reasons that will be clear in section-3. To the best of our knowledge this is the first time such a “large” matter multiplet is presented. We also find that this multiplet can be consistently restricted to a 8+8 component multiplet and show the precise identifications encoding the embedding of the tensor multiplet inside the dilaton matter multiplet.

The paper is organized as follows. In section 2, we will describe the features of four dimensional conformal supergravity relevant to our paper (see [11] for all the details and [12] for an extensive review on the subject). In section 3, we will present the dilaton matter multiplet along with its full supersymmetry transformation. This is the main result of our paper. In sections 4 and 5 we will present the details of the construction of the dilaton matter multiplet starting from the current multiplet of a rigid tensor multiplet. In section 6, we present the set of consistent constraints that restricts the multiplet to 8+8 components. Finally in section 7, we find the embedding of the tensor multiplet within the dilaton matter

Table 1.

Field	SU(2) Ireps	Weyl weight (w)	Chiral weight (c)	Chirality fermions
e_μ^a	1	-1	0	–
$V_\mu^i{}_j$	3	0	0	–
A_μ	1	0	0	–
b_μ	1	0	0	–
T_{ab}^-	1	1	-1	–
D	1	2	0	–
ψ_μ^i	2	-1/2	-1/2	+1
χ^i	2	3/2	-1/2	+1

multiplet. We will end with conclusions and future directions.

2 N=2 conformal supergravity in four spacetime dimensions

Conformal supergravity with N-extended supersymmetry is a gauge theory based on $SU(2, 2|N)$ superconformal algebra, the supersymmetric generalization of the conformal algebra. It is invariant under two different types of supersymmetry generators, the Q-supersymmetry and the special S-supersymmetry. The multiplet of fields that contains the gauge fields of the superconformal algebra is known as Weyl multiplet. For $N = 2$, the Weyl multiplet has 24+24 (bosonic+fermionic) off-shell degrees of freedom. In order to describe the multiplet, we fix our notations as follows. We denote the spacetime indices by greek letters $\mu, \nu \dots$, the local Lorentz indices by latin letters $a, b, \dots$, SU(2) indices by $i, j \dots$. As we mentioned before, two different versions of the Weyl multiplet exists in four dimensions: here we will focus on the standard formulation. The field content of the standard Weyl multiplet contains the gauge fields of the $SU(2, 2|2)$ algebra e_μ^a (vielbein), an SU(2) doublet of Majorana spinors whose positive and negative chiral projections are denoted by ψ_μ^i and $\psi_{\mu i}$ respectively (gravitinos), an SU(2) triplet of gauge fields $V_\mu^i{}_j$ corresponding to SU(2) R-symmetry, gauge field A_μ corresponding to U(1) R-symmetry, gauge field b_μ corresponding to the dilatation symmetry. Apart from the above mentioned gauge fields, the Weyl multiplet also contains several auxiliary fields that are required to balance bosonic and fermionic degrees of freedom. They include an SU(2) doublet of Majorana spinors whose positive and negative chiral projections are denoted by χ^i and χ_i respectively, a real tensor T_{ab} anti-symmetric in its Lorentz indices and a real scalar field D . The above mentioned field content of the Weyl multiplet can be tabulated in Table-1 along with their chiral and Weyl weights. Here $T_{ab}^\mp$ refers to the (anti)self-dual components of T_{ab} . Complex conjugation is used to raise/lower SU(2) indices and change the duality properties of a tensor. It hence flips the chiral weight as

well as chirality (for fermions) but not the Weyl weight. The gauge fields ω_μ^{ab} corresponding to local Lorentz transformations, f_μ^a corresponding to special conformal transformation and $\phi_{\mu i}$ corresponding to S-supersymmetry are dependent and are determined by the following set of conventional constraints:

$$\begin{aligned} R(P)_{\mu\nu}{}^a &= 0 , \\ \gamma^\mu \left(R(Q)_{\mu\nu}{}^i + \frac{1}{2} \gamma_{\mu\nu} \chi^i \right) &= 0 , \\ e^\nu{}_b R(M)_{\mu\nu a}{}^b - i \tilde{R}(A)_{\mu a} + \frac{1}{4} T^+{}_{ab} T^-{}_\mu{}^b - \frac{3}{2} D e_{\mu a} &= 0 . \end{aligned} \quad (2.1)$$

The super-covariant curvature $R(P)$, $R(Q)$ and $R(M)$ appearing above are associated with local translation, Q-supersymmetry and local Lorentz transformations respectively and $\tilde{R}(A)$ is the dual of the curvature associated with the U(1) R-symmetry. The supersymmetry commutators, in particular $[\delta_Q, \delta_Q]$, gets modified by field dependent structure constants as mentioned below:

$$\begin{aligned} [\delta_Q(\epsilon_1), \delta_Q(\epsilon_2)] &= \delta^{(cov)}(\xi) + \delta_M(\varepsilon) + \delta_K(\Lambda_K) + \delta_S(\eta) + \delta_{gauge} , \\ [\delta_S(\eta), \delta_Q(\epsilon)] &= \delta_M(\bar{\eta}_i \gamma^{ab} \epsilon^i + \text{h.c.}) + \delta_D(\bar{\eta}^i \epsilon_i + \text{h.c.}) + \delta_A(i \bar{\eta}_i \epsilon^i + \text{h.c.}) \\ &\quad + \delta_V(-2 \bar{\eta}^i \epsilon_j - (\text{h.c. ; traceless})) , \\ [\delta_S(\eta_1), \delta_S(\eta_2)] &= \delta_K(\bar{\eta}_{2i} \gamma^a \eta_1^i + \text{h.c.}) . \end{aligned} \quad (2.2)$$

The infinitesimal transformations, δ_Q , δ_S , δ_M , δ_K , δ_D , δ_A and δ_V , corresponds to Q-supersymmetry, S-supersymmetry, local Lorentz, special conformal, dilatation, U(1) and SU(2) R-symmetry respectively. The infinitesimal covariant transformation, $\delta^{(cov)}$ is defined as:

$$\delta^{(cov)}(\xi) = \delta_{gct}(\xi) + \sum_T \delta^T(-\xi^\mu h_\mu(T)) , \quad (2.3)$$

where δ_{gct} is general coordinate transformation. The sum is over all superconformal transformation (and any other gauge transformation present in the multiplet) except local translation and the field $h_\mu(T)$ is the corresponding gauge field, except for Q and S-supersymmetry, where $h_\mu(T)$ is $\frac{1}{2}$ times the corresponding gauge fields and for SU(2) R-symmetry where $h_\mu(T)$ is $-\frac{1}{2}$ times the corresponding gauge field. Finally the effect of any other gauge transformation (for example in vector multiplet) is encoded in δ_{gauge} that appears on the RHS of $[\delta_Q, \delta_Q]$ algebra in (2.2). The parameters that appear above in (2.2) read:

$$\begin{aligned} \xi^\mu &= 2 \bar{\epsilon}_2^i \gamma^\mu \epsilon_{1i} + \text{h.c.} , \\ \varepsilon_{ab} &= \varepsilon^{ij} \bar{\epsilon}_{1i} \epsilon_{2j} T_{ab}^- + \text{h.c.} , \\ \Lambda_K^a &= \varepsilon^{ij} \bar{\epsilon}_{1i} \epsilon_{2j} D_b T^{-ba} - \frac{3}{2} \bar{\epsilon}_2^i \gamma^a \epsilon_{1i} D + \text{h.c.} , \\ \eta^i &= 6 \bar{\epsilon}_1^i \epsilon_2^j \chi_j . \end{aligned} \quad (2.4)$$

Apart from the Weyl multiplet(s), there exists a large variety of $N = 2$ superconformal matter multiplets in four dimensions [13–19]. These multiplets all contain 8+8 off-shell degrees of freedom, with the exception of the hypermultiplet which is an on-shell multiplet².

Below, we will present a new $N = 2$ off-shell matter multiplet, which contains instead 24+24 degrees of freedom and forms an irreducible representation of the $SU(2, 2|2)$ superconformal algebra.

3 The dilaton matter multiplet

The new multiplet we present in this section contains a real scalar ϕ of Weyl weight +1 that can be interpreted as the dilaton. For this reason we will refer to this multiplet as the “dilaton matter multiplet”. Its bosonic content encompasses a traceless anti-hermitian $SU(2)$ triplet of Lorentz vector $S_a^i{}_j$,

$$(S_a^i{}_j)^* \equiv S_{ai}{}^j = -S_a^j{}_i, \quad S_a^i{}_i = 0, \quad (3.1)$$

an $SU(2)$ triplet of complex scalars E_{ij} (symmetric in the $SU(2)$ indices), and an $SU(2)$ quintuplet of pseudo-real scalars C_{ijkl} (completely symmetric in the $SU(2)$ indices) satisfying the reality condition

$$(C_{ijkl})^* \equiv C^{ijkl} = \varepsilon^{im}\varepsilon^{jn}\varepsilon^{kp}\varepsilon^{lq}C_{mnpq}. \quad (3.2)$$

The fermionic field content is given to be an $SU(2)$ doublet of Majorana fermions whose positive chirality components are denoted by Λ_i and an $SU(2)$ quadruplet of Majorana fermions whose positive chirality component are denoted by Ξ_{ijk} (completely symmetric in the $SU(2)$ indices). We present a schematic Table-2 with the full field content along with the Weyl and chiral weights and chirality.

All the field components are invariant under special conformal transformation³, and they transform under Q- and S-supersymmetry as shown below⁴.

$$\begin{aligned} \delta\phi &= -\frac{\phi}{2}\bar{\epsilon}^i\Lambda_i + \text{h.c.}, \\ \delta\Lambda^i &= -2\not{P}\epsilon^i - \left(\not{S}^i{}_j\epsilon^j + 2\varepsilon^{ik}\varepsilon^{jl}\epsilon_j E_{lk}\right) - \frac{1}{2}\bar{\Lambda}^i\Lambda^j\epsilon_j - \frac{1}{4}\bar{\Lambda}^j\gamma_a\Lambda_j\gamma^a\epsilon^i + \frac{1}{8}\bar{\Lambda}^i\gamma_{ab}\Lambda^j\gamma^{ab}\epsilon_j \\ &\quad - 2\eta^i, \\ \delta S_a^i{}_j &= \bar{\epsilon}_j\gamma_a\chi^i + \frac{2}{3}\bar{\epsilon}_j\gamma_a\not{D}\Lambda^i - 2\bar{\epsilon}_j D_a\Lambda^i - \frac{1}{3}\varepsilon^{li}\varepsilon^{nk}\bar{\epsilon}_n\gamma_a\Xi_{ljk} + \frac{1}{24}\bar{\epsilon}_j\gamma_a\gamma.T^-\Lambda_k\varepsilon^{ik} \end{aligned}$$

²The existence of off-shell hypermultiplet is subtle. One can formulate it in the presence of an off-shell central charge. Alternatively one can also formulate it in harmonic superspace [20], without the need for central charge. The latter formulation requires infinite degrees of freedom.

³This follows in a straightforward way from the $[\delta_S, \delta_S]$ commutators (2.2).

⁴We use the usual conventions for symmetrized non-consecutive indices, i.e. the indices that are not symmetrized are surrounded by vertical lines. For instance, we will write a multi-indices tensor product $A_i B_{jkl}$, symmetrized in (ijl) as $A_{(i} B_{j|k|l)}$.

Table 2.

Field	SU(2) Irreps	Weyl weight (w)	Chiral weight (c)	Chirality (Fermions)
ϕ	1	1	0	–
$S_a^i{}_j$	3	1	0	–
E_{ij}	3	1	-1	–
C_{ijkl}	5	2	0	–
Λ_i	2	1/2	1/2	+1
Ξ_{ijk}	4	3/2	-1/2	+1

$$\begin{aligned}
& -\frac{1}{3}\bar{\epsilon}_j\gamma_a\Lambda_k E_{lm}\epsilon^{il}\epsilon^{km} - \frac{2}{3}\bar{\epsilon}^i\gamma_a\mathcal{S}^k{}_j\Lambda_k - \frac{1}{2}\bar{\epsilon}^i\mathcal{S}^k{}_j\gamma_a\Lambda_k + \frac{1}{2}\bar{\epsilon}^k\gamma_a\mathcal{S}^i{}_j\Lambda_k - \frac{2}{3}\bar{\epsilon}^i\gamma_a\not{P}\Lambda_j \\
& - \bar{\epsilon}^i\not{P}\gamma_a\Lambda_j - \frac{1}{24}\bar{\Lambda}^i\Lambda^k\bar{\epsilon}_j\gamma_a\Lambda_k - \frac{1}{32}\bar{\Lambda}^i\gamma_{bc}\Lambda^k\bar{\epsilon}_j\gamma^{bc}\gamma_a\Lambda_k - (\text{h.c.;traceless}) , \\
\delta E_{ij} = & 2\bar{\epsilon}^{(l}\chi^k)\epsilon_{ik}\epsilon_{jl} - \frac{2}{3}\bar{\epsilon}^{(l}\not{D}\Lambda^k)\epsilon_{ik}\epsilon_{jl} + \frac{1}{3}\bar{\epsilon}^k\Xi_{ijk} - \frac{1}{12}\bar{\epsilon}^k\gamma.T^-\Lambda_{(i}\epsilon_{j)k} + \frac{2}{3}\bar{\epsilon}^k\Lambda_{(i}E_{j)k} \\
& - 2\bar{\epsilon}_{(i}\Lambda^k E_{j)k} - \frac{2}{3}\bar{\epsilon}^k\Lambda_k E_{ij} + \bar{\epsilon}_k\Lambda^k E_{ij} - \frac{1}{3}\bar{\epsilon}^{(l}\mathcal{S}^m{}_k\Lambda^k\epsilon_{il}\epsilon_{jm} - \frac{2}{3}\bar{\epsilon}^{(k}\not{P}\Lambda^l)\epsilon_{ik}\epsilon_{jl} \\
& - \frac{1}{12}\bar{\epsilon}^{(l}\gamma_a\Lambda^k)\bar{\Lambda}^m\gamma^a\Lambda_m\epsilon_{il}\epsilon_{jk} , \\
\delta\Xi_{ijk} = & \frac{3}{2}\epsilon_{mn}\epsilon_{lp}\left[D_a S^{al}{}_{(i}\delta_j^n\delta_k^p) - 2\gamma^{ab}D_a S_b{}^l{}_{(i}\delta_j^n\delta_k^p) - \gamma.R(V)^l{}_{(i}\delta_j^n\delta_k^p)\right]\epsilon^m + 6\not{D}E_{(ij}\epsilon_{k)} \\
& - C_{ijkl}\epsilon^l - 6E^{ln}E_{m(i}\epsilon_{j|l|}\epsilon_{k)n}\epsilon^m - 6\not{P}E_{(ij}\epsilon_{k)} + 3\mathcal{S}^m{}_{(i}E_{j)k}\epsilon_m - 6\mathcal{S}^m{}_{(i}E_{j|m|}\epsilon_{k)} \\
& + 3S^{am}{}_{(i}S^{bn}{}_{j}\epsilon_{k)m}\epsilon_{ln}\gamma_{ab}\epsilon^l + 3P^a S_a{}^l{}_{(i}\epsilon_{j|l|}\epsilon_{k)m}\epsilon^m - \frac{3}{4}\gamma.T^-E^{lm}\epsilon^n\epsilon_{(i|l|}\epsilon_{j|m|}\epsilon_{k)n} \\
& + \bar{\Lambda}^l\gamma_a\not{D}\Lambda^m\gamma^a\epsilon_{(i}\epsilon_{j|l|}\epsilon_{k)m} + \frac{1}{4}\bar{\Lambda}^l\not{D}\Lambda_{(i}\epsilon^m\epsilon_{j|l|}\epsilon_{k)m} - \frac{3}{8}\bar{\Lambda}^l\not{D}\gamma_{ab}\Lambda_{(i}\gamma^{ab}\epsilon^m\epsilon_{j|l|}\epsilon_{k)m} \\
& - \frac{1}{4}\bar{\Lambda}_{(i}\not{D}\Lambda^l\epsilon^m\epsilon_{j|l|}\epsilon_{k)m} + \frac{1}{8}\bar{\Lambda}_{(i}\gamma_{ab}\not{D}\Lambda^l\gamma^{ab}\epsilon^m\epsilon_{j|l|}\epsilon_{k)m} - \frac{3}{2}\bar{\Lambda}_{(i}R(Q)_{ab}{}^l\gamma^{ab}\epsilon^m\epsilon_{j|l|}\epsilon_{k)m} \\
& + \frac{1}{2}\bar{\Lambda}^p\Xi^{lmn}\epsilon^q\epsilon_{il}\epsilon_{jm}\epsilon_{kn}\epsilon_{pq} - \frac{1}{2}\bar{\Lambda}^{(m}\Xi^{np)l}\epsilon^q\epsilon_{im}\epsilon_{jn}\epsilon_{kp}\epsilon_{lq} - \frac{1}{8}\bar{\Lambda}_l\gamma_{ab}\Xi_{ijk}\gamma^{ab}\epsilon^l \\
& - \bar{\Lambda}_{(i}\Xi_{jk)l}\epsilon^l + \frac{1}{8}\bar{\Lambda}_{(i}\gamma_{ab}\Xi_{jk)l}\gamma^{ab}\epsilon^l - \frac{1}{2}\bar{\Lambda}^l\gamma_a\Xi_{ijk}\gamma^a\epsilon_l + \bar{\Lambda}^l\gamma_a\Xi_{l(ij}\gamma^a\epsilon_{k)} \\
& - \frac{3}{2}\bar{\Lambda}_{(i}\chi^l\epsilon^m\epsilon_{j|l|}\epsilon_{k)m} - \frac{3}{2}\bar{\Lambda}_{(i}\gamma_{ab}\chi^l\gamma^{ab}\epsilon^m\epsilon_{j|l|}\epsilon_{k)m} + \frac{3}{2}\bar{\Lambda}^l\chi_{(i}\epsilon^m\epsilon_{j|l|}\epsilon_{k)m} \\
& - 3\bar{\Lambda}^l\gamma_a\chi^m\gamma^a\epsilon_{(i}\epsilon_{j|l|}\epsilon_{k)m} + \frac{1}{4}\epsilon_{l(k}\bar{\Lambda}_i\Lambda_j)\gamma.T^-\epsilon^l - \frac{1}{8}\epsilon_{l(k}\bar{\Lambda}^l\gamma_a\Lambda_i\gamma.T^-\gamma^a\epsilon_{j)} \\
& + \frac{1}{2}\bar{\Lambda}^{(l}\Lambda^m E^{n)p}\epsilon^q\epsilon_{il}\epsilon_{jm}\epsilon_{kn}\epsilon_{pq} - \frac{1}{2}\bar{\Lambda}^p\Lambda^{(l}E^{mn)}\epsilon^q\epsilon_{il}\epsilon_{jm}\epsilon_{kn}\epsilon_{pq} + \frac{1}{2}\bar{\Lambda}_l\Lambda_{(i}E_{j)k}\epsilon^l \\
& - \frac{1}{2}\bar{\Lambda}_{(i}\Lambda_j E_{k)l}\epsilon^l + \frac{1}{4}\bar{\Lambda}_l\gamma_{ab}\Lambda_{(i}E_{j)k})\gamma^{ab}\epsilon^l + \frac{1}{4}\bar{\Lambda}^l\gamma_a\Lambda_l\gamma^a\epsilon_{(i}E_{j)k)} - \bar{\Lambda}^l\gamma_a\Lambda_{(i}\gamma^a\epsilon_{j)}E_{k)l}
\end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2}\bar{\Lambda}^l \mathcal{S}^n_l \Lambda_{(i} \epsilon^m \varepsilon_{j|n|} \varepsilon_{k)m} + \frac{1}{4} \varepsilon_{(i|n|} \varepsilon_{j|m|} \bar{\Lambda}^l \mathcal{S}^n_k \Lambda_l \epsilon^m - \frac{1}{8} \bar{\Lambda}^l \mathcal{S}^n_l \gamma_{ab} \Lambda_{(i} \gamma^{ab} \epsilon^m \varepsilon_{j|n|} \varepsilon_{k)m} \\
& + \frac{3}{16} \varepsilon_{(i|n|} \varepsilon_{j|m|} \bar{\Lambda}^l \mathcal{S}^n_k \gamma^{ab} \Lambda_l \gamma_{ab} \epsilon^m + \frac{1}{2} \bar{\Lambda}^n \Lambda^m \mathcal{S}^l_{(i} \varepsilon_{j\varepsilon k)m} \varepsilon_{nl} + \frac{1}{2} \bar{\Lambda}^l \mathcal{P} \Lambda_{(i} \epsilon^m \varepsilon_{j|l|} \varepsilon_{k)m} \\
& + \frac{1}{16} \bar{\Lambda}^l \gamma_{ab} \Lambda^m \mathcal{S}^n_{(i} \gamma^{ab} \epsilon_{j\varepsilon k)n} \varepsilon_{lm} + \frac{1}{2} \bar{\Lambda}^l \mathcal{P} \gamma_{ab} \Lambda_{(i} \gamma^{ab} \epsilon^m \varepsilon_{j|l|} \varepsilon_{k)m} + \bar{\Lambda}^l \Lambda^m \mathcal{P} \epsilon_{(i} \varepsilon_{j|l|} \varepsilon_{k)m} \\
& + \frac{1}{8} \bar{\Lambda}^m \Lambda^n \bar{\Lambda}^l \gamma_a \Lambda_l \gamma^a \epsilon_{(i} \varepsilon_{j|m|} \varepsilon_{k)n} - \frac{1}{16} \bar{\Lambda}^l \Lambda^m \bar{\Lambda}_n \gamma_{ab} \Lambda_{(i} \gamma^{ab} \epsilon^n \varepsilon_{j|l|} \varepsilon_{k)m} , \\
\delta C_{ijkl} &= \bar{\epsilon}_{(i} \Gamma_{jkl)} + \varepsilon_{ip} \varepsilon_{jq} \varepsilon_{kr} \varepsilon_{ls} \bar{\epsilon}^{(p} \Gamma^{qrs)} + (\bar{\epsilon}_m \Lambda^m + \bar{\epsilon}^m \Lambda_m) C_{ijkl} . \tag{3.3}
\end{aligned}$$

In the above, the composite field $P_a \equiv \phi^{-1} D_a \phi$ and Γ_{ijk} is defined as:

$$\begin{aligned}
\Gamma_{ijk} &= -2\mathcal{D}\Xi_{ijk} - 3D_a S^{an} {}_{(i} \Lambda^m \varepsilon_{j|n|} \varepsilon_{k)m} + 6\mathcal{D}E_{(ij} \Lambda_{k)} - 2C_{ijkl} \Lambda^l + 2\mathcal{D}\Lambda_{(i} E_{j k)} \\
& + 2D_a \Lambda^l S^{an} {}_{(i} \varepsilon_{j|n|} \varepsilon_{k)l} - 2\gamma^{ab} D_a \Lambda^l S_b^n {}_{(i} \varepsilon_{j|n|} \varepsilon_{k)l} - 4\Xi^{lmn} E_{l(i} \varepsilon_{j|m|} \varepsilon_{k)n} + 2\mathcal{S}^l_{(i} \Xi_{jk)l} \\
& + 12\chi_{(i} E_{jk)} - 6\mathcal{S}^l_{(i} \chi^m{}_{j|l|} \varepsilon_{k)m} - 4\mathcal{P}E_{(ij} \Lambda_{k)} - 4P_a S^{an} {}_{(i} \varepsilon_{j|n|} \varepsilon_{k)l} \Lambda^l \\
& - 2P_a \gamma^{ab} S_b^n {}_{(i} \varepsilon_{j|n|} \varepsilon_{k)l} \Lambda^l - 2\mathcal{S}^l_{(i} E_{jk)} \Lambda_l - 2\mathcal{S}^l_{(i} E_{j|l|} \Lambda_{k)} - 4E_{(ij} \varepsilon_{k)m} \varepsilon_{ln} E^{mn} \Lambda^l \\
& + 12\varepsilon_{(i|m|} \varepsilon_{j|n|} E_{k)l} E^{mn} \Lambda^l + S_a^m {}_{(i} S^{an} {}_{j\varepsilon k)m} \varepsilon_{ln} \Lambda^l + \gamma_{ab} S^{am} {}_{(i} S^{bn} {}_{j\varepsilon k)m} \varepsilon_{ln} \Lambda^l \\
& + \frac{1}{2} \gamma \cdot T^+ E_{(ij} \varepsilon_{k)l} \Lambda^l + \frac{1}{4} \mathcal{S}^l_{(i} \gamma \cdot T^- \Lambda_{j\varepsilon k)l} + \frac{1}{2} \bar{\Lambda}^l \Lambda^m \mathcal{D}\Lambda_{(i} \varepsilon_{j|l|} \varepsilon_{k)m} \\
& + \frac{5}{4} \bar{\Lambda}^l \gamma^a \Lambda_{(i} \varepsilon_{j|l|} \varepsilon_{k)m} D_a \Lambda^m + \frac{5}{16} \bar{\Lambda}^l \gamma^{bc} \gamma^a \Lambda_{(i} \varepsilon_{j|l|} \varepsilon_{k)m} \gamma_{bc} D_a \Lambda^m \\
& + \frac{3}{2} \bar{\Lambda}^l \Lambda^m \chi_{(i} \varepsilon_{j|l|} \varepsilon_{k)m} - \frac{3}{2} \bar{\Lambda}^l \gamma^a \Lambda_{(i} \varepsilon_{j|l|} \varepsilon_{k)m} \gamma_a \chi^m + \frac{1}{2} \bar{\Lambda}^q \Lambda^{(p} \Xi^{ml)n} \varepsilon_{ip} \varepsilon_{jm} \varepsilon_{kl} \varepsilon_{qn} \\
& + \frac{1}{2} \bar{\Lambda}^l \gamma_a \Lambda_{(i} \gamma^a \Xi_{jk)l} + \frac{3}{2} \bar{\Lambda}^l \Lambda^m \mathcal{P} \Lambda_{(i} \varepsilon_{j|l|} \varepsilon_{k)m} - \frac{1}{4} \bar{\Lambda}^l \Lambda^m \mathcal{S}^n_{(i} \varepsilon_{j|n|} \varepsilon_{k)m} \Lambda_l \\
& - \frac{1}{32} \varepsilon_{mn} \bar{\Lambda}^m \gamma^{ab} \Lambda^n \gamma_{ab} \mathcal{S}^l_{(i} \Lambda_{j\varepsilon k)l} - \frac{1}{2} \bar{\Lambda}^l \Lambda^{(m} E^{pq)} \Lambda^n \varepsilon_{ip} \varepsilon_{jq} \varepsilon_{km} \varepsilon_{ln} \\
& - \bar{\Lambda}_{(i} \Lambda_{j} E_{k)l} \Lambda^l - \bar{\Lambda}_l \Lambda_{(i} E_{jk)} \Lambda^l + \frac{1}{8} \bar{\Lambda}^l \Lambda^m \bar{\Lambda}^n \gamma_a \Lambda_n \gamma^a \Lambda_{(i} \varepsilon_{j|l|} \varepsilon_{k)m} . \tag{3.4}
\end{aligned}$$

To obtain the above result, we first construct the current multiplet of a rigid N=2 tensor multiplet in four spacetime dimensions. This multiplet will couple to a linearized supergravity multiplet containing the graviton and gravitino. Since the tensor multiplet is not conformal i.e its action in flat spacetime is not invariant under a scale transformation, the energy momentum tensor $\theta_{\mu\nu}$ will not be traceless and the supersymmetry current $J_{\mu i}$ will not be gamma-traceless. Hence, the trace of the energy momentum tensor, $\sigma = \theta^\mu{}_\mu$ and the gamma-trace of the supersymmetry current, $\lambda_i = \gamma^\mu J_{\mu i}$ will be non trivial components of the current multiplet. These components will couple to a dilaton and a dilatino respectively in the linearized supegravity multiplet. In this way one expects to find a different version of the Weyl multiplet with a dilaton and a dilatino. This is typically referred to as dilaton Weyl multiplet in the literature for five dimensional and six dimensional conformal supergravity [1, 6]. In five dimensions [1], this was found exactly in the above mentioned way, by investigating the current multiplet for a five dimensional N=2 vector multiplet. However, following the identical procedure in four dimensions, and building the current multiplet for a non-conformal tensor

multiplet in four spacetime dimensions, we find that the corresponding linearized supergravity multiplet that couples to this current multiplet gets decomposed into a 24+24 linearized version of the four dimensional standard Weyl multiplet and a new 24+24 component matter multiplet, which is a linearized version of the dilaton matter multiplet presented in (3.3). The full non linear transformations are obtained by a standard application of the supersymmetry algebra (2.2).

4 Tensor multiplet and the multiplet of supercurrents

An off-shell rigid N=2 tensor multiplet in four spacetime dimensions [19] encompasses 8+8 degrees of freedom, organized in an SU(2) triplet of pseudo real scalars L_{ij} satisfying the reality constraint $(L_{ij})^* \equiv L^{ij} = \varepsilon^{ik}\varepsilon^{jl}L_{kl}$; an SU(2) doublet of Majorana spinors, whose positive components are denoted by ϕ^i ; a real tensor gauge field $E_{\mu\nu}$ with vector gauge transformation $\delta E_{\mu\nu} = 2\partial_{[\mu}\Lambda_{\nu]}$. To close the algebra off-shell, an auxiliary complex scalar field G is required. When G is set to zero through its algebraic equations of motion, the supersymmetry transformations of the tensor multiplet components are given by :

$$\begin{aligned}\delta E_{\mu\nu} &= i\bar{\epsilon}^i\gamma_{\mu\nu}\phi^j\epsilon_{ij} + \text{h.c.} , \\ \delta\phi^i &= \not{\partial}L^{ij}\epsilon_j + \epsilon^{ij}\not{H}\epsilon_j , \\ \delta L^{ij} &= 2\bar{\epsilon}^{(i}\phi^{j)} + 2\epsilon^{ik}\epsilon^{j\ell}\bar{\epsilon}_{(k}\phi_{\ell)} .\end{aligned}\tag{4.1}$$

The action that is invariant under the above susy transformation reads:

$$S = \int d^4x \left[H_\mu H^\mu - \bar{\phi}^i \overleftrightarrow{\not{\partial}} \phi_i - \frac{1}{2} \partial_\mu L^{ij} \partial^\mu L_{ij} \right] ,\tag{4.2}$$

where we defined

$$H^\mu = \frac{i}{6} \varepsilon^{\mu\nu\rho\sigma} H_{\nu\rho\sigma} .\tag{4.3}$$

The three form $H_{\mu\nu\rho}$ is the gauge-invariant field strength of the tensor gauge field $E_{\mu\nu}$ defined as $H_{\mu\nu\rho} \equiv 3\partial_{[\mu}E_{\nu\rho]}$. The equations of motion for the fields are given by:

$$\not{\partial}\phi^i = 0 , \quad \partial^2 L_{ij} = 0 , \quad \partial_{[\mu}H_{\nu]} = 0 .\tag{4.4}$$

The tensor gauge field also satisfies the Bianchi identity $\partial^\mu H_\mu = 0$. Apart from the above mentioned supersymmetry transformation (4.1), the action (4.2) is also invariant under an SU(2) $\times$ U(1) R-symmetry.

The current multiplet associated to the tensor multiplet action contains the following components: the energy momentum tensor $\theta_{\mu\nu}$ arising from translation invariance, a supersymmetry current $J_{\mu i}$ arising from the supersymmetry invariance, an SU(2) current $v_\mu{}^i{}_j$ and a U(1) current a_μ arising from the SU(2) $\times$ U(1) R-symmetry. Since the multiplet is non conformal, the

trace σ of the energy momentum tensor and the gamma-trace λ_i of the supersymmetry current will also be a part of the current multiplet. The above mentioned components of the current multiplet in terms of the tensor multiplet fields are given as below:

$$\begin{aligned}
\theta_{\mu\nu} &= H_\mu H_\nu - \frac{1}{2}\eta_{\mu\nu}H^2 + \frac{1}{2}\bar{\phi}^i\gamma_{(\mu}\overleftrightarrow{\partial}_{\nu)}\phi_i + \frac{1}{2}\partial_\mu L^{ij}\partial_\nu L_{ij} - \frac{1}{4}\eta_{\mu\nu}\partial_\rho L^{ij}\partial^\rho L_{ij} , \\
\sigma &\equiv \theta_\mu^\mu = -H^2 - \frac{1}{2}\partial^\mu L^{ij}\partial_\mu L_{ij} , \\
v_\mu{}^i{}_j &= \frac{1}{8}L^{ik}\overleftrightarrow{\partial}_\mu L_{jk} - \frac{1}{4}\left(\bar{\phi}^i\gamma_\mu\phi_j - \frac{1}{2}\delta_j^i\bar{\phi}^k\gamma_\mu\phi_k\right) , \\
a_\mu &= -\frac{i}{8}\bar{\phi}^l\gamma_\mu\phi_l , \\
J_{\mu i} &= \frac{1}{2}\gamma_\mu\mathbb{H}\phi^j\varepsilon_{ij} - H_\mu\phi^j\varepsilon_{ij} + \phi^j\partial_\mu L_{ij} - \frac{1}{2}\gamma_\mu\mathbb{D}L_{ij}\phi^j , \\
\lambda_i &\equiv \gamma^\mu J_{\mu i} = \mathbb{H}\phi^j\varepsilon_{ij} - \mathbb{D}L_{ij}\phi^j .
\end{aligned} \tag{4.5}$$

The other components of the current multiplet are obtained by supersymmetry and read

$$\begin{aligned}
t_\mu{}^i{}_j &= \frac{1}{2}H_\mu L^{ik}\varepsilon_{kj} + \frac{1}{4}\left(\bar{\phi}^i\gamma_\mu\phi_j - \frac{1}{2}\delta_j^i\bar{\phi}^k\gamma_\mu\phi_k\right) , \\
e^{ij} &= \bar{\phi}^i\phi^j , \quad \tilde{a}_\mu = \varepsilon_{ij}\bar{\phi}^i\partial_\mu\phi^j , \quad d = \frac{1}{2}L^{ij}L_{ij} , \\
b_{\mu\nu}^- &= \frac{1}{2}\bar{\phi}^i\gamma_{\mu\nu}\phi^j\varepsilon_{ij} , \quad c^{ijkl} = L^{(ij}L^{kl)} , \\
\Sigma^{ijk} &= L^{(ij}\phi^k) , \quad \xi_j = L_{jk}\phi^k .
\end{aligned} \tag{4.6}$$

Together, eqs. (4.5) and (4.6) form the 48+48 component current multiplet of the rigid N=2 tensor multiplet in four spacetime dimensions. They satisfy the following conservation, reality and trace relations.

$$\begin{aligned}
\partial^\mu\theta_{\mu\nu} &= 0 , \quad \sigma = \theta_\mu^\mu , \quad \partial^\mu J_{\mu i} = 0 , \quad \lambda_i = \gamma^\mu J_{\mu i} , \\
(v_\mu{}^i{}_j)^* &= -v_\mu{}^j{}_i , \quad v_\mu{}^i{}_i = 0 , \quad \partial_\mu v^{\mu i}{}_j = 0 , \\
(t_\mu{}^i{}_j)^* &= -t_\mu{}^j{}_i , \quad t_\mu{}^i{}_i = 0 , \\
\partial_\mu a^\mu &= 0 , \quad (a_\mu)^* = a_\mu , \quad \partial_\mu \tilde{a}^\mu = 0 , \\
\partial^\mu b_{\mu\nu}^- &= \tilde{a}_\nu , \quad \frac{1}{2}\varepsilon_{\mu\nu\rho\sigma}b^{-\rho\sigma} = -b_{\mu\nu}^- ,
\end{aligned} \tag{4.7}$$

and transform under supersymmetry as :

$$\begin{aligned}
\delta\theta_{\mu\nu} &= \bar{\epsilon}^i\gamma_{\lambda(\mu}\partial^\lambda J_{\nu)i} + \text{h.c.} , \\
\delta\sigma &= \bar{\epsilon}^i\mathbb{D}\lambda_i + \text{h.c.} , \\
\delta t_\mu{}^i{}_j &= \frac{1}{2}\bar{\epsilon}^i J_{\mu j} - \frac{1}{2}\bar{\epsilon}^i\gamma_\mu\lambda_j + \frac{1}{4}\varepsilon_{lm}\varepsilon_{jk}\bar{\epsilon}^l\mathbb{D}\left(\gamma_\mu\Sigma^{imk}\right) - \frac{1}{3}\bar{\epsilon}^i\gamma_\mu\mathbb{D}\xi_j - \frac{1}{3}\bar{\epsilon}^i\partial_\mu\xi_j \\
&\quad - (\text{h.c.;traceless}) ,
\end{aligned}$$

$$\begin{aligned}
\delta v_\mu{}^i{}_j &= \frac{1}{2}\bar{\epsilon}^i J_{\mu j} - \frac{1}{4}\epsilon_{kj}\epsilon_{lm}\bar{\epsilon}^l \gamma_\mu \not{\partial} \Sigma^{ikm} - \frac{1}{3}\bar{\epsilon}^i \partial_\mu \xi_j + \frac{1}{4}\epsilon_{kj}\epsilon_{lm}\bar{\epsilon}^l \partial_\mu \Sigma^{ikm} + \frac{1}{3}\bar{\epsilon}^i \gamma_\mu \not{\partial} \xi_j \\
&\quad - (\text{h.c.; traceless}) , \\
\delta a_\mu &= \frac{i}{4}\bar{\epsilon}_k J_\mu^k + \text{h.c.} , \\
\delta b_{\mu\nu}^- &= -\epsilon^{ik}\bar{\epsilon}_i \gamma^\alpha \gamma_{\mu\nu} J_{\alpha k} , \\
\delta \tilde{a}_\mu &= 2\epsilon^{ik}\bar{\epsilon}_i \not{\partial} J_{\mu k} + \epsilon^{ik}\bar{\epsilon}_i \gamma_\mu \not{\partial} \lambda_k - \epsilon^{ik}\bar{\epsilon}_i \partial_\mu \lambda_k , \\
\delta e^{ij} &= -2\bar{\epsilon}_k \not{\partial} \Sigma^{ijk} - 2\epsilon^{k(i}\epsilon^{j)l}\bar{\epsilon}_k \lambda_l - \frac{8}{3}\epsilon^{k(i}\epsilon^{j)l}\bar{\epsilon}_k \not{\partial} \xi_l , \\
\delta d &= 2\bar{\epsilon}_i \xi^i + \text{h.c.} , \\
\delta c^{ijkl} &= 4\bar{\epsilon}^{(i}\Sigma^{jkl)} + 4\epsilon^{im}\epsilon^{jn}\epsilon^{kp}\epsilon^{lq}\bar{\epsilon}_{(m}\Sigma_{npq)} , \\
\delta J_{\mu i} &= \theta_{\mu\nu}\gamma^\nu \epsilon_i + (\gamma_\rho \gamma_{\mu\sigma} + \gamma_{\mu\sigma} \gamma_\rho) \partial^\sigma v^{\rho k}{}_i \epsilon_k + (\gamma_\rho \gamma_{\mu\sigma} + \gamma_{\mu\sigma} \gamma_\rho) \partial^\sigma t^{\rho k}{}_i \epsilon_k \\
&\quad - \frac{1}{4}\partial_\mu b_{\nu\rho}^- \gamma^{\nu\rho} \epsilon^j \epsilon_{ij} - \gamma \partial^\rho e^{lj} \gamma_{\mu\rho} \epsilon^k \epsilon_{ij} \epsilon_{kl} + \gamma_{\mu\rho} \tilde{a}^\rho \epsilon^j \epsilon_{ij} + i(\gamma_\rho \gamma_{\mu\sigma} - 3\gamma_{\mu\sigma} \gamma_\rho) \partial^\sigma a^\rho \epsilon_i , \\
\delta \lambda_i &= \sigma \epsilon_i + 4\not{\partial} \psi_i^k \epsilon_k + 4\not{\partial} t_i^{\mu k} \epsilon_k - 4\partial_\mu t_i^{\mu k} \epsilon_k - 8i\not{\partial} \phi \epsilon_i + 3\not{\partial} \epsilon^j \epsilon_{ij} - \frac{1}{4}\not{\partial} (\gamma \cdot b^-) \epsilon^j \epsilon_{ij} , \\
\delta \xi_j &= -2t_j^l \epsilon_l - 4\phi_j^l \epsilon_l + 6i\phi \epsilon_j - \frac{3}{8}b^- \cdot \gamma \epsilon^m \epsilon_{jm} - \frac{1}{2}\epsilon_{jl}\epsilon_{km} e^{lk} \epsilon^m + \frac{1}{2}\not{\partial} d \epsilon_j , \\
\delta \Sigma^{ijk} &= -2\epsilon^{l(i}\not{\partial}^{jk)} \epsilon_l + 2\epsilon^{l(i}\not{\partial}^{jk)} \epsilon_l + \frac{1}{2}\not{\partial} c^{ijkl} \epsilon_l - e^{(ij} \epsilon^{k)} . \tag{4.8}
\end{aligned}$$

5 Linearized supergravity multiplet

The components of the current multiplet will couple to the linearized components of a supergravity multiplet as shown below in (5.1). This will fix the basis and normalization of the fields belonging to our linearized gravity multiplet⁵. The invariant action reads:

$$\begin{aligned}
S = \int d^4x \left[\frac{1}{2}\theta^{\mu\nu} h_{\mu\nu} + \sigma\varphi + d\mathcal{D} + \frac{1}{24}c^{ijkl}C_{ijkl} + \frac{1}{4}b_{\mu\nu}^- T^{-\mu\nu} - 2v_\mu{}^i{}_j V^{\mu j}{}_i + e^{ij}E_{ij} \right. \\
\left. + 4a_\mu A^\mu + (t_\mu{}^i{}_j - v_\mu{}^i{}_j) S^{\mu j}{}_i + 2\bar{J}_{\mu i} \psi^{\mu i} + \bar{\lambda}_i \Lambda^i + \bar{\xi}_i \zeta^i + \frac{1}{3}\bar{\Sigma}_{ijk}\Xi^{ijk} + \tilde{a}_\mu \tilde{A}^\mu + \text{h.c.} \right] . \tag{5.1}
\end{aligned}$$

Supersymmetry invariance of the above action (5.1) gives us the linearized transformation of the fields. Due to the constraints (4.7) satisfied by the currents, the fields coupling to them will have gauge symmetries. For instance, the field $V_\mu{}^i{}_j$ will have an SU(2) gauge symmetry $\delta V_\mu{}^i{}_j \sim \partial_\mu \Lambda^i{}_j$ because it couples to a conserved current (second line of 4.7). As a consequence, we can neglect some pure gauge terms $\sim \bar{\epsilon}^i \partial_\mu \Lambda_j$ that will appear in the supersymmetry variation of $V_\mu{}^i{}_j$. Alternatively one can add terms proportional to $\partial^\mu v_\mu{}^j{}_i \epsilon_j$ (which is zero due to the relations (4.7)) in the supersymmetry variation of λ_i in (4.8), which

⁵If we choose different numerical factors in the coupling or take different linear combination of fields coupling to the currents, we will have a different choice of basis and normalization for the fields which will be related to our choice via field redefinition.

cancels against any pure gauge-like terms of the form shown above in the supersymmetry variation of $V_\mu^i{}_j$.

Analogously, due to the last line of (4.7), the fields $T_{\mu\nu}^-$ and $\tilde{A}_\mu$ that couple to the current $b_{\mu\nu}^-$ and $\tilde{a}_\mu$ have a complex vector gauge transformation as shown below.

$$\begin{aligned}\delta T_{\mu\nu}^- &= \partial_{[\mu}\Lambda_{\nu]} - \frac{1}{2}\varepsilon_{\mu\nu\rho\sigma}\partial^\rho\Lambda^\sigma, \\ \delta\tilde{A}_\mu &= \Lambda_\mu.\end{aligned}\tag{5.2}$$

From the above equation, it is obvious that the field $\tilde{A}_\mu$ is a pure gauge under this complex vector gauge transformation and can be gauged away by using the gauge fixing condition $\tilde{A}_\mu = 0$. However, we need to compensate the supersymmetry transformation of $\tilde{A}_\mu$ by a field dependent complex vector gauge transformation. This will only affect the supersymmetry transformation of $T_{\mu\nu}^-$. After doing the above set of exercises, we obtain the linearized transformation of the fields belonging to the linearized multiplet:

$$\begin{aligned}\delta C^{ijkl} &= -2\bar{\epsilon}^{(l}\delta\Xi^{ijk)} - 2\epsilon^{im}\epsilon^{jn}\epsilon^{kp}\epsilon^{lq}\bar{\epsilon}_{(m}\delta\Xi_{npq)}, \\ \delta\mathcal{D} &= -\frac{1}{4}\bar{\epsilon}_i\delta\chi^i + \text{h.c.}, \\ \delta h^{\mu\nu} &= 2\bar{\epsilon}_i\gamma^{(\mu}\psi^{\nu)i} + \text{h.c.}, \\ \delta\varphi &= -\frac{1}{2}\bar{\epsilon}_i\Lambda^i + \text{h.c.}, \\ \delta E_{ij} &= \frac{1}{3}\bar{\epsilon}^k\Xi_{ijk} - \frac{1}{2}\bar{\epsilon}^{(l}\zeta^{k)}\varepsilon_{ik}\varepsilon_{jl}, \\ \delta V_\mu^j{}_i &= -\bar{\epsilon}_i(\gamma_{\rho\sigma}\gamma_\mu + \gamma_\mu\gamma_{\rho\sigma})\partial^\sigma\psi^{\rho j} - 2\bar{\epsilon}_i\gamma_\mu\delta\Lambda^j + \frac{3}{2}\bar{\epsilon}_i\gamma_\mu\zeta^j - (\text{h.c.; traceless}), \\ \delta S_\mu^j{}_i &= \bar{\epsilon}_i(\gamma_{\rho\sigma}\gamma_\mu + \gamma_\mu\gamma_{\rho\sigma})\partial^\sigma\psi^{\rho j} + 2\bar{\epsilon}_i\gamma_\mu\delta\Lambda^j - 2\bar{\epsilon}_i\partial_\mu\Lambda^j - \bar{\epsilon}_i\gamma_\mu\zeta^j - \frac{1}{3}\varepsilon^{lj}\varepsilon^{nk}\bar{\epsilon}_n\gamma^\mu\Xi_{ilk} \\ &\quad - (\text{h.c.; traceless}), \\ \delta T_{\mu\nu}^- &= \frac{3}{2}\bar{\epsilon}^i\gamma_{\mu\nu}\zeta^j\varepsilon_{ij} - 2\bar{\epsilon}^i\gamma_{\mu\nu}\delta\Lambda^j\varepsilon_{ij} + 8\varepsilon_{ij}\bar{\epsilon}^i\left(\partial_{[\mu}\psi_{\nu]}^j\right)_{|A.S.D.} + 2\bar{\epsilon}^i\gamma_{\mu\nu}\gamma_{\rho\sigma}\partial^\rho\psi^{\sigma j}\varepsilon_{ij}, \\ \delta A_\mu &= +\frac{i}{4}\bar{\epsilon}_i(\gamma_{\rho\sigma}\gamma_\mu + 3\gamma_\mu\gamma_{\rho\sigma})\partial^\sigma\psi^{\rho i} + \frac{3i}{4}\bar{\epsilon}_j\gamma_\mu\zeta^j - i\bar{\epsilon}_j\gamma_\mu\delta\Lambda^j + \text{h.c.}, \\ \delta\psi^{\mu i} &= -\frac{1}{2}\gamma_{\lambda\nu}\bar{\epsilon}^i\partial^\lambda h^{\mu\nu} + V^{\mu i}{}_j\epsilon^j + iA^\mu\bar{\epsilon}^i - \frac{1}{8}\varepsilon^{ij}(\gamma \cdot T^-)\gamma^\mu\epsilon_j, \\ \delta\Lambda^i &= -2\delta\varphi\bar{\epsilon}^i - \delta^i{}_j\epsilon^j - 2\varepsilon^{ik}\varepsilon^{jl}\epsilon_j E_{lk}, \\ \delta\zeta^i &= -\frac{4}{3}\delta\mathcal{S}^i{}_j\epsilon^j - \frac{4}{3}\gamma^{\mu\nu}\partial_\mu V_\nu^i{}_j\epsilon^j + \frac{8}{3}\varepsilon^{ik}\varepsilon^{lj}\delta E_{lk}\epsilon_j - 4\mathcal{D}\bar{\epsilon}^i, \\ \delta\Xi_{ijk} &= \frac{3}{2}\varepsilon_{mn}\varepsilon_{lp}\left[\partial_\mu S^{\mu l}{}_{(i}\delta_j^n\delta_k^p) - 2\gamma^{\mu\nu}\partial_\mu S_\nu^l{}_{(i}\delta_j^n\delta_k^p) - 2\gamma^{\mu\nu}\partial_\mu V_\nu^l{}_{(i}\delta_j^n\delta_k^p)\right]\bar{\epsilon}^m + 6\delta E_{(ij}\epsilon_{k)} \\ &\quad - C_{ijkl}\bar{\epsilon}^l.\end{aligned}\tag{5.3}$$

In the above variations, we observe that the derivative of the gravitino ψ_μ^i appears in antisymmetric form $\sim \partial_{[\mu}\psi_{\nu]}^i$, in the supersymmetry variation of some of the fields like $V_\mu^i{}_j$,

$S_\mu^i{}_j$, $T_{\mu\nu}^-$ and A_μ . This is important because derivatives of gauge fields should be traded off for the corresponding curvature $R(Q)$ in this case) and it can only happen if the derivatives appear in an anti-symmetric form. But before trading off $\partial\psi$ in terms of $R(Q)$ in the above variations, we recall that the curvature $R(Q)$ will contain a term proportional to the S-gauge field ϕ_μ^i at the linearized level. This happens because the gravitino ψ_μ^i transforms under S-supersymmetry which is known to be $\delta_S \psi_\mu^i = -\gamma_\mu \eta^i$ from the superconformal algebra. Hence, the linearized form of the curvature $R(Q)_{\mu\nu}^i$, is given as

$$R(Q)_{\mu\nu}^i = 2\partial_{[\mu}\psi_{\nu]}^i - \gamma_{[\mu}\phi_{\nu]}^i, \quad (5.4)$$

where ϕ_μ^i is the gauge field corresponding to S-supersymmetry. Using (5.4) one can trade off the antisymmetric derivative on the gravitino for the curvature $R(Q)$ in the supersymmetry variation of the above mentioned fields. However, we also need to impose the curvature constraint on $R(Q)$. The conventional constraint for $R(Q)$ (2.1) involves the standard Weyl multiplet field χ^i . However, at this point since we have not yet identified the field χ^i , which would be some linear combination of the fields appearing in (5.3), it is difficult to implement the conventional constraint for the curvature $R(Q)$ from (2.1). We will instead impose the following unconventional constraint on $R(Q)$ ⁶

$$\gamma^\mu R(Q)_{\mu\nu}^i = 0, \quad (5.5)$$

which, at the linearized level, gives:

$$\gamma^{\mu\nu}\partial_\mu\psi_\nu^i = -\frac{3}{2}\gamma^\nu\phi_\nu^i. \quad (5.6)$$

We can see that, when we try to re-write $\partial_{[\mu}\psi_{\nu]}^i$ in terms of $R(Q)$ and use the above constraint in the linearized transformation (5.3), bare S-gauge field ϕ_μ^i will pop out as shown below.

$$\begin{aligned} \delta V_\mu^i{}_j &= 2\bar{\epsilon}_j\phi_\mu^i - 2\bar{\epsilon}_j\gamma_\mu\gamma\cdot\phi^i - 2\bar{\epsilon}_j\not{\partial}\Lambda^i + \frac{3}{2}\bar{\epsilon}_j\gamma_\mu\zeta^i - (\text{h.c.;traceless}), \\ \delta A_\mu &= \frac{i}{2}\bar{\epsilon}_i\phi_\mu^i - i\bar{\epsilon}_j\gamma_\mu\gamma\cdot\phi^j + \frac{3i}{4}\bar{\epsilon}_j\gamma_\mu\zeta^j - i\bar{\epsilon}_j\gamma_\mu\not{\partial}\Lambda^j + \text{h.c.}, \\ \delta T_{\mu\nu}^- &= 4\epsilon_{ij}\bar{\epsilon}^i R(Q)_{\mu\nu}^j - 2\bar{\epsilon}^i\gamma_{\mu\nu}\not{\partial}\Lambda^j\epsilon_{ij} - 2\bar{\epsilon}^i\gamma_{\mu\nu}\gamma\cdot\phi^j\epsilon_{ij} + \frac{3}{2}\bar{\epsilon}^i\gamma_{\mu\nu}\zeta^j\epsilon_{ij}, \\ \delta S_\mu^i{}_j &= 2\bar{\epsilon}_i\gamma_\mu\not{\partial}\Lambda^j - 2\bar{\epsilon}_i\partial_\mu\Lambda^j + 2\bar{\epsilon}_i\gamma_\mu\gamma\cdot\phi^j - 2\bar{\epsilon}_i\gamma_\mu\phi^j - \bar{\epsilon}_i\gamma_\mu\zeta^j - \frac{1}{3}\epsilon^{lj}\epsilon^{nk}\bar{\epsilon}_n\gamma^\mu\Xi_{ilk} \\ &\quad - (\text{h.c.;traceless}). \end{aligned} \quad (5.7)$$

Among the fields that appear above in (5.7), the fields like $S_\mu^i{}_j$ and $T_{\mu\nu}^-$ do not couple to any conserved current and as a result they will not correspond to any gauge fields of the multiplet.

⁶The conventional and the unconventional constraints are related by a shift in the S-gauge field by a term proportional to the auxiliary fermion χ^i . Once we correctly identify the χ^i , we can go back to the conventional constraint by making an appropriate shift in the S-gauge field.

Rather they will be the auxiliary fields. Such fields should not have any bare S-gauge field in their transformation rules, which should be either be inside a covariant derivative or a curvature. Hence, terms involving bare S-gauge field that appears in (5.7) for $T_{\mu\nu}^-$ and S_μ^{ij} , should be absorbed into the covariant derivative of Λ^i . This can be done, if we let Λ^i to transform under S-supersymmetry as

$$\delta_S \Lambda^i = -2\eta^i. \quad (5.8)$$

As a consequence, the covariant derivative of Λ^i at the linearized level is as shown below:

$$D_\mu \Lambda^i = \partial_\mu \Lambda^i + \phi_\mu^i. \quad (5.9)$$

In terms of the covariant derivative on Λ^i , the supersymmetry variation (5.7) takes the following form

$$\begin{aligned} \delta V_\mu^{ij} &= 2\bar{\epsilon}_j \phi_\mu^i - 2\bar{\epsilon}_j \not{D} \Lambda^i + \frac{3}{2} \bar{\epsilon}_j \gamma_\mu \zeta^i - (\text{h.c.; traceless}), \\ \delta A_\mu &= \frac{i}{2} \bar{\epsilon}_i \phi_\mu^i + \frac{3i}{4} \bar{\epsilon}_j \gamma_\mu \zeta^j - i\bar{\epsilon}_j \gamma_\mu \not{D} \Lambda^j + \text{h.c.}, \\ \delta T_{\mu\nu}^- &= 4\varepsilon_{ij} \bar{\epsilon}^i R(Q)_{\mu\nu}^j - 2\bar{\epsilon}^i \gamma_{\mu\nu} \not{D} \Lambda^j \varepsilon_{ij} + \frac{3}{2} \bar{\epsilon}^i \gamma_{\mu\nu} \zeta^j \varepsilon_{ij}, \\ \delta S_\mu^{ij} &= 2\bar{\epsilon}_i \gamma_\mu \not{D} \Lambda^j - 2\bar{\epsilon}_i D_\mu \Lambda^j - \bar{\epsilon}_i \gamma_\mu \zeta^j - \frac{1}{3} \varepsilon^{lj} \varepsilon^{nk} \bar{\epsilon}_n \gamma^\mu \Xi_{ilk} - (\text{h.c.; traceless}). \end{aligned} \quad (5.10)$$

By looking at the transformation of the R-symmetry gauge fields V_μ^{ij} and A_μ , we find that the standard Weyl multiplet χ^i can be identified with the following linear combination at the linearized level⁷.

$$\chi^i = -\frac{3}{4} \zeta^i + \not{D} \Lambda^i. \quad (5.11)$$

By making a supersymmetry transformation on the above, we find that the following linear combination is identified with the standard-Weyl multiplet field “ D ” at the linearized level.

$$D = 3\mathcal{D} - 2\partial^2 \varphi. \quad (5.12)$$

Finally, we also redefine the scalar field φ as⁸

$$\phi = e^\varphi. \quad (5.13)$$

Having identified the standard-Weyl multiplet field χ^i , we would prefer going back to the conventional constraint by making a field redefinition on the S-gauge field: $\phi_\mu^i \rightarrow \phi_\mu^i - \frac{1}{2} \gamma_\mu \chi^i$. In terms of the above redefined fields (5.11-5.13 as well as the redefined S-gauge field), we see that the linearized components $h_{\mu\nu}$, V_μ^{ij} , A_μ , $T_{\mu\nu}^-$, D , ψ_μ^i , χ^i decouples from the rest and

⁷Here, we need to keep in mind that we are working with the unconventional constraint and hence the variation of V_μ^{ij} and A_μ should be compared with the results known in the literature after shifting the S-gauge field so that we can correctly identify the χ field.

⁸The redefinitions are redundant at the linearized level, but really important at the non-linear level.

they transform, under Q-supersymmetry, exactly like the linearization of the standard Weyl multiplet known in the literature in the conventional constraints. The full supersymmetry transformation of the above fields including the S-supersymmetry transformation follows from the superconformal algebra (2.2) and is already known. The remaining 24+24 components ϕ , E_{ij} , S_{μ}^i , C_{ijkl} , Λ^i , Ξ_{ijk} transform as given below.

$$\begin{aligned}
\delta\phi &= -\frac{\phi}{2}\bar{\epsilon}^i\Lambda_i + \text{h.c.} , \\
\delta\Lambda^i &= -2\mathcal{P}\epsilon^i - \left(\mathcal{S}_j^i\epsilon^j + 2\epsilon^{ik}\epsilon^{jl}\epsilon_j E_{lk} \right) - 2\eta^i , \\
\delta S_a^i{}_j &= \bar{\epsilon}_j\gamma_a\chi^i + \frac{2}{3}\bar{\epsilon}_j\gamma_a\mathcal{D}\Lambda^i - 2\bar{\epsilon}_j D_a\Lambda^i - \frac{1}{3}\epsilon^{li}\epsilon^{nk}\bar{\epsilon}_n\gamma_a\Xi_{ljk} - (\text{h.c.; traceless}) , \\
\delta E_{ij} &= 2\bar{\epsilon}^{(l}\chi^{k)}\epsilon_{ik}\epsilon_{jl} - \frac{2}{3}\bar{\epsilon}^{(l}\mathcal{D}\Lambda^{k)}\epsilon_{ik}\epsilon_{jl} + \frac{1}{3}\bar{\epsilon}^k\Xi_{ijk} , \\
\delta\Xi_{ijk} &= \frac{3}{2}\epsilon_{mn}\epsilon_{lp} \left[\partial_\mu S^{\mu l}{}_{(i}\delta_j^n\delta_k^p - 2\gamma^{\mu\nu}\partial_\mu S_{\nu}{}^l{}_{(i}\delta_j^n\delta_k^p - \gamma \cdot R(V)^l{}_{(i}\delta_j^n\delta_k^p \right] \epsilon^m + 6\mathcal{D}E_{(ij}\epsilon_{k)} \\
&\quad - C_{ijkl}\epsilon^l , \\
\delta C^{ijkl} &= -2\bar{\epsilon}^{(l}\mathcal{D}\Xi^{ijk)} - 2\epsilon^{im}\epsilon^{jn}\epsilon^{kp}\epsilon^{lq}\bar{\epsilon}_{(m}\mathcal{D}\Xi_{npq)} .
\end{aligned} \tag{5.14}$$

Here, we have defined $P_a = \phi^{-1}\partial_a\phi$ and $R(V)_{\mu\nu}{}^i{}_j = 2\partial_{[\mu}V_{\nu]}{}^i{}_j$. The above result (5.14) is a linearization of our final result (3.3)⁹. The above linearized transformation gives us the Weyl and chiral weights of the fields as mentioned in Table 2. Because of the S-susy transformation of Λ_i , it is clear that Λ_i will have the same chiral and Weyl weight as η_i which are $-1/2$ and $+1/2$ respectively. Further, we obtain the weights of E_{ij} , $S_a^i{}_j$, C_{ijkl} and Ξ_{ijk} by looking at the above susy transformation (5.14) and the knowledge of the weights of the Q-susy parameter ϵ^i . The chiral weight of ϕ has to be zero because it is a real scalar field. The Weyl weight of ϕ is found by acting the commutator $[\delta_S, \delta_Q]$ on it.

Thereafter, we apply the algebra (2.2) consistently to obtain the final result (3.3). In the process, we assume that the S-transformation of $S_a^i{}_j$, E_{ij} , C_{ijkl} and Ξ_{ijk} is zero. This is because any S-transformation of the above mentioned fields can be removed by a field redefinition with a term containing Λ^i . This is possible simply because the S-transformation of Λ^i is proportional to the S-susy parameter η^i . For example, suppose the S-transformation of E_{ij} was:

$$\delta_S E_{ij} = \alpha\bar{\eta}^{(l}\Lambda^{k)}\epsilon_{il}\epsilon_{jk} , \tag{5.15}$$

then we could redefine E_{ij} as a primary (S-invariant) field

$$\tilde{E}_{ij} = E_{ij} + \frac{\alpha}{4}\bar{\Lambda}^{(l}\Lambda^{k)}\epsilon_{il}\epsilon_{jk} . \tag{5.16}$$

⁹We have partially non-linearized our results (see $\delta\phi$ and first term of $\delta\Lambda^i$) by working with the field ϕ instead of φ which are related by an exponentiation as given in (5.13).

6 Constraining the dilaton matter multiplet

In this section, we will show how to consistently constrain the new multiplet to obtain a restricted multiplet with 8+8 components. This discussion is in line with the one on N=2 chiral multiplet in the literature [21]. The chiral multiplet of N=2 supergravity is a large multiplet with 16+16 components. It is known that a chiral multiplet of Weyl weight $w = 1$ for the lowest component can be consistently reduced by imposing a reality constraint on an SU(2) triplet of field B_{ij} belonging to the chiral multiplet. Analogously, the 24+24 components dilaton matter multiplet presented in this paper can be reduced to a 8+8 matter multiplet, by imposing a reality constraint on the field E_{ij} . However, since E_{ij} has a non trivial chiral weight $c = -1$ (as opposed to B_{ij} of chiral multiplet which has $c = 0$ if $w = 1$ for the lowest component), we can only constrain it to a “real” field up to an overall phase factor, which carries the chiral weight, i.e we can constrain E_{ij} to take the following form:

$$E_{ij} = e^{-i\sigma/2} \mathcal{L}_{ij} , \quad (6.1)$$

where $\mathcal{L}_{ij}$ satisfies the reality constraint:

$$\mathcal{L}^{ij} = \varepsilon^{ik} \varepsilon^{jl} \mathcal{L}_{kl} . \quad (6.2)$$

Trivially, under chiral transformation, the phase σ transforms as:

$$\delta_{U(1)}(\Lambda)\sigma = 2\Lambda . \quad (6.3)$$

As a consequence of (6.1), E_{ij} and its complex conjugate $\bar{E}_{ij} \equiv \varepsilon_{ik} \varepsilon_{jl} E^{kl}$ are related by the constraint:

$$\mathcal{R}_{ij} := \bar{E}_{ij} - e^{i\sigma} E_{ij} = 0 . \quad (6.4)$$

The above constraint can be taken as the starting point of a procedure to obtain the full set of constraints by applying supersymmetry. For the supersymmetry transformation of $\mathcal{R}_{ij}$, we get:

$$\delta \mathcal{R}_{ij} = \bar{\epsilon}_{(i} \Theta_{j)} + \bar{\epsilon}_k \Upsilon^{lmk} \varepsilon_{il} \varepsilon_{jm} - e^{-i\sigma} \varepsilon_{ik} \varepsilon_{jl} \left(\bar{\epsilon}^{(k} \Theta^{l)} + \bar{\epsilon}^m \Upsilon_{mnp} \varepsilon^{kn} \varepsilon^{lp} \right) , \quad (6.5)$$

with

$$\begin{aligned} \Theta_j &= 2\chi_j - \frac{2}{3} \not{D} \Lambda_j + \frac{1}{12} \gamma \cdot T^+ \Lambda^l \varepsilon_{jl} + \frac{2}{3} \Lambda^l \bar{E}_{jl} + \frac{1}{3} \not{S}^k{}_j \Lambda_k - \frac{2}{3} \not{P} \Lambda_j \\ &\quad + \frac{1}{12} \bar{\Lambda}^m \gamma_a \Lambda_m \gamma^a \Lambda_j - \frac{2i}{3} \zeta^l \bar{E}_{jl} , \\ \Upsilon^{lmk} &= \frac{1}{3} \Xi^{lmk} - \Lambda^{(l} E^{mk)} - i \zeta^{(l} E^{mk)} . \end{aligned} \quad (6.6)$$

Here, we have defined ζ_l as the supersymmetry variation of the phase σ :

$$\delta \sigma = \bar{\epsilon}_k \zeta^k + \text{h.c.} . \quad (6.7)$$

Since $\mathcal{R}_{ij} = 0$ is a supersymmetric invariant constraint, we need to set its supersymmetry variation to zero, i.e. $\Theta_i = 0 = \Upsilon_{ijk}$, to get the other set of constraints. From the first identity we derive the relation between ζ^k and the components of the dilaton matter multiplet. Specifically:

$$i\zeta^k = \Lambda^k + 3E^{-2}\alpha_j \bar{E}^{jk} , \quad (6.8)$$

with

$$\begin{aligned} \alpha_j &\equiv 2\chi_j - \frac{2}{3}\not{D}\Lambda_j + \frac{1}{12}\gamma \cdot T^+ \Lambda^l \varepsilon_{jl} + \frac{1}{3}\not{\mathcal{S}}^k{}_j \Lambda_k - \frac{2}{3}\not{P}\Lambda_j + \frac{1}{12}\bar{\Lambda}^m \gamma_a \Lambda_m \gamma^a \Lambda_j , \\ E^2 &\equiv E^{ij} E_{ij} . \end{aligned} \quad (6.9)$$

The constraint $\Upsilon^{lmk} = 0$, together with (6.8), gives an expression for Ξ^{ijk} completely determined in terms of the lower weight fields:

$$\Xi^{ijk} = 9E^{-2}\alpha_l E^{(ij} \bar{E}^{kl)} + 6\Lambda^{(i} E^{jk)} . \quad (6.10)$$

At this point, one can check that $\delta_S \Xi_{ijk} = 0$ by using the S-transformation of α_j :

$$\delta_S \alpha_j = \frac{8}{3} \bar{E}_{jk} \eta^k .$$

Also from (6.8), one finds that $\delta_S \zeta^i = -2i\eta^i$, and hence the $[\delta_Q, \delta_S]$ commutator acts on σ as:

$$[\delta_S(\eta), \delta_Q(\epsilon)] \sigma = -2i\bar{\epsilon}_k \eta^k + 2i\bar{\epsilon}^k \eta_k . \quad (6.11)$$

This is consistent with the U(1) transformation of σ given in (6.3).

Now, since Ξ_{ijk} is completely determined in terms of the lower weight fields, the number of independent fermionic components of the multiplet is reduced to 8 (Λ_i).

We obtain further constraints on the bosonic fields by setting the supersymmetry variation of Υ_{ijk} to zero. For simplicity, we present only the bosonic terms:

$$\delta\Upsilon_{ijk} = A_{ijkl}\epsilon^l + B_{(ij}\varepsilon_{k)l}\epsilon^l + D^{ab}{}_{ijkl}\gamma_{ab}\epsilon^l + F^{ab}{}_{(ij}\varepsilon_{k)l}\gamma_{ab}\epsilon^l + \not{G}_{(ij}\epsilon_{k)} + \not{J}_{ijkl}\epsilon_m \varepsilon^{lm} , \quad (6.12)$$

with

$$\begin{aligned} A_{ijkl} &= \frac{1}{3} \left[-C_{ijkl} - 18E^{-2} D E_{(ij} \bar{E}_{kl)} - 12E^{-2} D_a P^a E_{(ij} \bar{E}_{kl)} - 6E^{-2} D_a S^{am}{}_{(i} E_{jk} \bar{E}_{l)m} \right. \\ &\quad \left. - 12E^{-2} P_a S^{am}{}_{(i} E_{jk} \bar{E}_{l)m} - \frac{3}{2} E^{-2} S^n{}_{m \cdot} S^m{}_{n} E_{(ij} \bar{E}_{kl)} - 12E^{-2} P_a P^a E_{(ij} \bar{E}_{kl)} \right. \\ &\quad \left. + 6E_{(ij} \bar{E}_{kl)} \right] + \text{fermions} , \\ B_{ij} &= \mathcal{L}^{-2} \mathcal{L}_{ij} \mathcal{L}_{kl} \mathcal{G}^{kl} + \text{fermions} , \\ F^{ab}{}_{ij} &= \mathcal{H}_{abij} - \mathcal{L}^{-2} \mathcal{L}_{ij} \mathcal{L}^{kl} \mathcal{H}_{abkl} + \text{fermions} , \\ D^{ab}{}_{ijkl} &= \frac{4}{3} \mathcal{L}^{-2} \mathcal{H}^{ab}{}_{m(i} \mathcal{L}_{jk} \mathcal{L}_{l)n} \varepsilon^{mn} + \text{fermions} , \end{aligned}$$

$$\begin{aligned}
J_{ijkl}^a &= \frac{4}{3} \mathcal{L}^{-2} G_{n(i}^a \mathcal{L}_{jk} \mathcal{L}_{l)m} \varepsilon^{mn} + \text{fermions} , \\
G_{ij}^a &= 3e^{-i\sigma/2} \left(D^a \mathcal{L}_{ij} - (\mathcal{L}^{-1} D^a \mathcal{L}) \mathcal{L}_{ij} - S^{ak} ({}_i \mathcal{L}_j)_k \right) + \text{fermions} .
\end{aligned} \tag{6.13}$$

In the above expressions, we have defined¹⁰:

$$\begin{aligned}
\mathcal{G}^{ij} &= \frac{3}{2} [D_a S^{aij} + 2P_a S^{aij}] , \\
\mathcal{H}_{abij} &= -\frac{3}{2} \left[D_{[a} S_{b]ij} + \frac{1}{2} S_{am} ({}_i S_b^m{}_j) + \frac{1}{2} R(V)_{abij} \right] .
\end{aligned} \tag{6.14}$$

By setting (6.12) to zero, we obtain the following set of constraints:

$$\begin{aligned}
A_{ijkl} &= 0 , \\
\mathcal{G}^{ij} \mathcal{L}_{ij} &= 0 , \\
\mathcal{H}_{abij} &= \mathcal{L}^{-2} \mathcal{L}_{ij} \mathcal{L}^{mn} \mathcal{H}_{abmn} , \\
G_{ij}^a &= 0 .
\end{aligned} \tag{6.15}$$

The first constraint fixes C_{ijkl} in terms of the lower weight field. The third constraint in (6.15) is not an independent one, since it is equivalent to $\mathcal{L}^{lm} \varepsilon_{jm} D_{[b} G_{a]il} = 0$. Hence, we are left with a scalar (1) and a vector-tensor (12) constraints, which read explicitly:

$$\begin{aligned}
(D_a S^{aij} + 2P_a S^{aij}) \mathcal{L}_{ij} &= 0 , \\
\left(D^a \mathcal{L}_{ij} - (\mathcal{L}^{-1} D^a \mathcal{L}) \mathcal{L}_{ij} - S^{ak} ({}_i \mathcal{L}_j)_k \right) &= 0 .
\end{aligned} \tag{6.16}$$

At this point, it is trivial to check that the second constraint above encompasses only 12-4=8 independent components, since a contraction with $\mathcal{L}^{ij}$ gives zero identically. Together, (6.16) constitute a set of 9 independent constraints on $S_a^i{}_j$. Hence, $S_a^i{}_j$ has only 3 independent components. Furthermore, the second identity in (6.16) can be solved for $S_a^i{}_j$ in terms of a 4-vector $\mathcal{H}_a$. To see this, consider the set of equalities:

$$\begin{aligned}
S_a^k ({}_i \mathcal{L}_j)_k &= S_a^k {}_i \mathcal{L}_{jk} - S_a^k {}_{[i} \mathcal{L}_{j]k} \\
&= S_a^k {}_i \mathcal{L}_{jk} - \frac{1}{2} S_a^{km} \mathcal{L}_{mk} \varepsilon_{ij} \\
&= S_a^k {}_i \mathcal{L}_{jk} - \frac{1}{2} \phi^{-2} \mathcal{L} \mathcal{H}_a \varepsilon_{ij} ,
\end{aligned} \tag{6.17}$$

where, we defined $S_a^{mk} \mathcal{L}_{mk} = \phi^{-2} \mathcal{L} \mathcal{H}_a$. The pre-factor $\phi^{-2} \mathcal{L}$ is chosen conveniently so that $\mathcal{H}_a$ has Weyl weight +3 and it satisfies a simple Bianchi identity as we will see below. Substituting (6.17) into the second identity in (6.16) and contracting with $\mathcal{L}^{lj}$, we can solve explicitly for $S_a^i{}_j$:

$$S_a^i{}_j = \mathcal{L}^{-2} [\mathcal{L}^{im} D_a \mathcal{L}_{jm} - \mathcal{L}_{jm} D_a \mathcal{L}^{im}] + \phi^{-2} \mathcal{L}^{-1} \mathcal{H}_a \mathcal{L}^{ik} \varepsilon_{kj} . \tag{6.18}$$

¹⁰The indices of $S_a^i{}_j$ and $R(V)_{ab}{}^i{}_j$ are raised/lowered by using the rule $A_{ij} = \varepsilon_{ik} A^k{}_j$ and $A^{ij} = A^i{}_k \varepsilon^{kj}$.

Plugging this solution in the first identity in (6.16), we obtain a Bianchi identity on $\mathcal{H}_a$, i.e.:

$$D_a \mathcal{H}^a = 0 . \quad (6.19)$$

Thus, $\mathcal{H}_a$ can be interpreted as the dual of a 3-form field strength corresponding to a 2-form gauge field, up to possible fermionic terms.

To sum up, we have seen that, $S_a^i{}_j$ has 3 independent components which can be traded off for the dual of a three form field strength $\mathcal{H}_a$ satisfying a Bianchi identity. Together with the real scalar field ϕ (1), the phase σ (1) and $\mathcal{L}_{ij}$ (3), they make up the total 8 bosonic components expected for a minimal matter multiplet of N=2 supergravity.

7 The embedding of the tensor multiplet in the dilaton matter multiplet

In this section we want to show the precise embedding of the N=2 tensor multiplet in the dilaton matter multiplet, which satisfies the constraint discussed in the previous section. This will show the equivalence between the tensor multiplet and the restricted dilaton matter multiplet. For completeness we write the full supersymmetry transformation of the tensor multiplet in a conformal supergravity background [19, 22]:

$$\begin{aligned} \delta L_{ij} &= 2\bar{\epsilon}_{(i}\varphi_{j)} + 2\varepsilon_{ik}\varepsilon_{jl}\bar{\epsilon}^{(k}\varphi^{l)} , \\ \delta\varphi^i &= \not{D}L^{ij}\epsilon_j + \not{H}\varepsilon^{ij}\epsilon_j - G\epsilon^i + 2L^{ij}\eta_j , \\ \delta G &= -2\bar{\epsilon}_i\not{D}\varphi^i - 6\bar{\epsilon}_i\chi_j L^{ij} + \frac{1}{4}\varepsilon^{ij}\bar{\epsilon}_i\gamma \cdot T^+\varphi_j + 2\bar{\eta}_i\varphi^i , \\ \delta E_{\mu\nu} &= i\bar{\epsilon}^i\gamma_{\mu\nu}\varphi^j\varepsilon_{ij} + 2iL_{ij}\varepsilon^{jk}\bar{\epsilon}^i\gamma_{[\mu}\psi_{\nu]k} + \text{h.c.} . \end{aligned} \quad (7.1)$$

Here we have defined H_a as the dual of the 3-form field strength H_{abc} as shown below:

$$H_a = \frac{i}{6}\varepsilon_{abcd}H^{bcd} , \quad (7.2)$$

where $H_{\mu\nu\rho}$ is the super-covariant field strength associated with tensor gauge field $E_{\mu\nu}$ given as:

$$H_{\mu\nu\rho} = 3\partial_{[\mu}E_{\nu\rho]} - \frac{3i}{2}\bar{\psi}_{[\mu}{}^i\gamma_{\nu\rho]}\varphi^j\varepsilon_{ij} + \frac{3i}{2}\bar{\psi}_{[\mu}{}^i\gamma_{\nu\rho]}\varphi_j\varepsilon^{ij} - \frac{3i}{2}L_{ij}\varepsilon^{jk}\bar{\psi}_{[\mu}{}^i\gamma_{\nu}\psi_{\rho]k} . \quad (7.3)$$

Now, it is possible to verify that the following combinations of the tensor multiplet fields,

$$\begin{aligned} \phi^4 &= L^2 , \\ \Lambda^i &= -2L^{-2}L^{ij}\varphi_j , \\ E_{ij} &= L^{-4}L_{ij}L^{kl}\bar{\varphi}_k\varphi_l - L^{-2}\bar{G}L_{ij} , \\ S_a^i{}_j &= 2L^{-2}H_aL^{ik}\varepsilon_{kj} + 4L^{-4}L^{ik}L_{jm}\bar{\varphi}^m\gamma_a\varphi_k - L^{-2}\bar{\varphi}^i\gamma_a\varphi_j - \frac{1}{2}L^{-2}\delta_j^i\bar{\varphi}^m\gamma_a\varphi_m , \\ &\quad + L^{-2}\left(L^{ik}D_aL_{jk} - L_{jk}D_aL^{ik}\right) , \end{aligned}$$

$$\begin{aligned}
\Xi_{ijk} = & -24L^{-6}L^{mn}\bar{\varphi}_m\varphi_nL_{(ij}L_{k)l}\varphi^l + 6L^{-4}\bar{\varphi}_l\varphi_{(i}L_{jk)}\varphi^l - 6L^{-4}L^{lm}\not{D}L_{l(i}L_{jk)}\varphi_m \\
& + 6L^{-4}L^{mn}\not{H}\varphi_nL_{(ij}\varepsilon_{k)m} + 12L^{-4}\bar{G}L_{(ij}L_{k)l}\varphi^l + 6L^{-2}\not{D}\varphi_{(i}L_{jk)} \\
& + 18L^{-2}L_{(ij}L_{k)l}\chi^l - \frac{3}{4}L^{-2}\gamma \cdot T^- \varphi^l L_{(ij}\varepsilon_{k)l} \\
C_{ijkl} = & 6L^{-4}G\bar{G}L_{(ij}L_{kl)} - 18L^{-2}DL_{(ij}L_{kl)} + 6L^{-4}H^aH_aL_{(ij}L_{kl)} \\
& - 12L^{-4}H^aD_aL^{mn}\varepsilon_{m(i}L_{jk}L_{l)n} - 6L^{-4}L^{mn}D_aL_{mn}D^aL_{(ij}L_{kl)} - 6L^{-2}D_aD^aL_{(ij}L_{kl)} \\
& + 12L^{-4}L^{mn}D_aL_{m(i}L_{jk}D^aL_{l)n} + 3L^{-4}D_aL_{mn}D^aL^{mn}L_{(ij}L_{kl)} - 9L^{-2}\bar{\chi}^m\varphi^nL_{(ij}\varepsilon_{k|m|}\varepsilon_{l)n} \\
& - 36L^{-4}\bar{\chi}^m\varphi^nL_{(ij}L_{k|m|}L_{l)n} + 6L^{-6}\bar{G}L_{mn}\bar{\varphi}^m\varphi^nL_{(ij}L_{kl)} + 6L^{-4}G\bar{\varphi}_{(i}\varphi_{j}L_{kl)} \\
& - 18L^{-6}GL^{mn}\bar{\varphi}_m\varphi_nL_{(ij}L_{kl)} - 6L^{-4}\bar{G}\bar{\varphi}^m\varphi^nL_{(ij}\varepsilon_{k|m|}\varepsilon_{l)n} - 24L^{-6}\bar{G}\bar{\varphi}^m\varphi^nL_{(ij}L_{k|m|}L_{l)n} \\
& + 36L^{-6}\bar{\varphi}^m\gamma^a\varphi_nL^{sn}D_aL_{s(i}L_{jk}L_{l)m} - 36L^{-6}\bar{\varphi}^m\gamma^a\varphi_nL^{ns}H_aL_{(ij}L_{k|m|}\varepsilon_{l)s} \\
& - 6L^{-4}\bar{\varphi}^m\gamma^a\varphi_{(i}L_{jk}D_aL_{l)m} + 6L^{-4}\bar{\varphi}^m\gamma^a\varphi_{(i}L_{jk}\varepsilon_{l)m}H_a - 6L^{-4}\bar{\varphi}^m\gamma^a\varphi_mD_aL_{(ij}L_{kl)} \\
& - 12L^{-4}L^{st}\varphi_s\not{D}\varphi^mL_{(ij}\varepsilon_{k|t|}\varepsilon_{l)m} - \frac{3}{2}L^{-4}\bar{\varphi}^m\gamma \cdot T^- \varphi^nL_{m(i}L_{jk}\varepsilon_{l)n} \\
& + 18L^{-4}L^{mn}\bar{\chi}_m\varphi_nL_{(ij}L_{kl)} - \frac{27}{2}L^{-2}\bar{\chi}_{(i}\varphi_{j}L_{kl)} - 6L^{-8}L^{mn}\bar{\varphi}_m\varphi_nL_{st}\bar{\varphi}^s\varphi^tL_{(ij}L_{kl)} \\
& + 12L^{-6}\bar{\varphi}^m\varphi^n\bar{\varphi}_m\varphi_nL_{(ij}L_{kl)} + 12L^{-6}L_{mn}\bar{\varphi}^m\varphi^n\bar{\varphi}_{(i}\varphi_{j}L_{kl)} - 60L^{-6}\bar{\varphi}^m\varphi^n\bar{\varphi}_m\varphi_{(i}L_{jk}L_{l)n} \\
& - 48L^{-8}L^{st}\bar{\varphi}_s\varphi_t\bar{\varphi}^m\varphi^nL_{m(i}L_{jk}L_{l)n} + 144L^{-8}L^{st}\bar{\varphi}_s\varphi_t\bar{\varphi}^m\varphi^nL_{(ij}L_{k|m|}L_{l)n} \tag{7.4}
\end{aligned}$$

transform as the components of the dilaton matter multiplet (3.3). One can check that the above combinations satisfy the constraints presented in the previous section. The relation between the tensor multiplet and the independent components of the restricted dilaton matter multiplet can be obtained from the identification (7.4) and are given as:

$$\begin{aligned}
\phi^4 &= L^2, \\
\Lambda^i &= -2L^{-2}L^{ij}\varphi_j, \\
e^{-i\sigma/2} &= \left(\frac{\mathcal{Z}}{\bar{\mathcal{Z}}}\right)^{1/2}, \\
\mathcal{L}_{ij} &= |\mathcal{Z}|L_{ij}, \\
\mathcal{H}_a &\sim H_a, \quad \text{Up to fermion bilinears.} \tag{7.5}
\end{aligned}$$

Here $\mathcal{Z}$ is defined as the following combination of tensor multiplet fields:

$$\mathcal{Z} = L^{-4}L^{kl}\bar{\varphi}_k\varphi_l - L^{-2}\bar{G} . \tag{7.6}$$

Thus, we see that the off-shell N=2 tensor multiplet is equivalent to the restricted version of the dilaton matter multiplet presented in this paper¹¹.

8 Conclusions and future directions

Superconformal couplings of the matter multiplets and the gauge multiplet are crucial in our understanding of the structure of supergravity that can arise from them. Apart from

¹¹We thank Daniel Butter for pointing out this possibility.

the structure of the higher derivative terms that one can obtain, they also tell us about the geometric structure of the matter couplings. For example, it is well-known that the scalar manifold corresponding to vector multiplet coupled to $N=2$ conformal supergravity in four dimensions, upon gauge fixing and eliminating the auxiliary fields, corresponds to a special Kähler manifold (see [23–25] and [12] for a review). Similar geometric structure exists for other matter couplings in four dimensional as well as five dimensional supergravity. Hence, the discovery of new matter multiplets opens up the possibilities of gaining new insights into supergravity structures arising from them. In this paper, we have presented one such multiplet, the dilaton matter multiplet, which contains $24+24$ off-shell degrees of freedom. We have also shown that this multiplet can be consistently restricted to $8+8$ components, which can be identified with the components of a tensor multiplet. One possible future direction is the construction of a density formula in terms of this new multiplet. By using the composite expressions (7.4) in terms of the tensor multiplet fields, one would then obtain the coupling of tensor multiplet to conformal supergravity. This would perhaps be different from the coupling of the improved tensor multiplet to conformal supergravity obtained in [22] which was based on the chiral density formula. We leave this study for a future work. It would also be interesting to see if results analogous to the ones presented in section 7 can be obtained for vector and Weyl multiplets, and the possibility of new formulations of conformal and Poincaré supergravity.

A complete knowledge of the structure of supergravity arising from various matter and gauge couplings, in an off-shell formulation, is crucial to our understanding of black holes, in particular the effective quantum corrections to their entropy. In fact, using the standard Weyl multiplet and other matter multiplets known so far, several higher derivative invariants have been constructed in four spacetime dimensions [26–29] and their effect on black hole entropy have been studied [28, 30–32]. It has also been shown that the above results are able to correctly reproduce the microscopic entropy of BPS black holes [28, 30, 32] as well as some class of non BPS black holes [31]. However, the result presented in this paper practically allows for the existence of new, unknown couplings in supergravity, and their connection to higher dimensional theories and string theory in general. As an inevitable consequence, the need will arise to revisit and put under scrutiny every result mentioned above (and more), based on the analysis of invariant couplings of supergravity.

Acknowledgments

The authors would like to thank Daniel Butter and Bernard de Wit for their helpful comments on this work. We would also like to thank the organizers of the Indian National Strings Meeting (NSM) 2015 (IISER Mohali) for giving us a platform for illuminating discussions which led to the idea behind this work.

References

- [1] E. Bergshoeff, T. de Wit, R. Halbersma, S. Cucu, M. Derix and A. Van Proeyen, *Weyl multiplets of $N=2$ conformal supergravity in five-dimensions*, *JHEP* **06** (2001) 051, [[hep-th/0104113](#)].
- [2] T. Fujita and K. Ohashi, *Superconformal tensor calculus in five-dimensions*, *Prog. Theor. Phys.* **106** (2001) 221–247, [[hep-th/0104130](#)].
- [3] K. Hanaki, K. Ohashi and Y. Tachikawa, *Supersymmetric Completion of an R^{**2} term in Five-dimensional Supergravity*, *Prog. Theor. Phys.* **117** (2007) 533, [[hep-th/0611329](#)].
- [4] T. Kugo and K. Ohashi, *Supergravity tensor calculus in 5-D from 6-D*, *Prog. Theor. Phys.* **104** (2000) 835–865, [[hep-ph/0006231](#)].
- [5] N. Banerjee, B. de Wit and S. Katmadas, *The Off-Shell 4D/5D Connection*, *JHEP* **03** (2012) 061, [[1112.5371](#)].
- [6] E. Bergshoeff, E. Sezgin and A. Van Proeyen, *Superconformal Tensor Calculus and Matter Couplings in Six-dimensions*, *Nucl. Phys.* **B264** (1986) 653.
- [7] M. Muller, *Minimal $N=2$ off-shell supergravity*, *Phys. Lett.* **B172** (1986) 353–356.
- [8] M. Muller, *Minimal $N = 2$ Supergravity in Superspace*, *Nucl. Phys.* **B282** (1987) 329–348.
- [9] W. Siegel, *Curved extended superspace from Yang-Mills theory a la strings*, *Phys. Rev.* **D53** (1996) 3324–3336, [[hep-th/9510150](#)].
- [10] D. Butter, S. Hegde, I. Lodato and B. Sahoo, *$N = 2$ dilaton-Weyl multiplet in 4D Supergravity*, **1712.XXXX**.
- [11] E. Bergshoeff, M. de Roo and B. de Wit, *Extended Conformal Supergravity*, *Nucl. Phys.* **B182** (1981) 173–204.
- [12] T. Mohaupt, *Black hole entropy, special geometry and strings*, *Fortsch. Phys.* **49** (2001) 3–161, [[hep-th/0007195](#)].
- [13] P. Fayet, *Fermi-Bose Hypersymmetry*, *Nucl. Phys.* **B113** (1976) 135.
- [14] R. Grimm, M. Sohnius and J. Wess, *Extended Supersymmetry and Gauge Theories*, *Nucl. Phys.* **B133** (1978) 275–284.
- [15] B. de Wit and J. W. van Holten, *Multiplets of Linearized $SO(2)$ Supergravity*, *Nucl. Phys.* **B155** (1979) 530–542.
- [16] B. de Wit, J. W. van Holten and A. Van Proeyen, *Structure of $N=2$ Supergravity*, *Nucl. Phys.* **B184** (1981) 77.
- [17] M. Sohnius, K. S. Stelle and P. C. West, *Off Mass Shell Formulation of Extended Supersymmetric Gauge Theories*, *Phys. Lett.* **92B** (1980) 123–127.
- [18] F. Brandt, *New $N=2$ supersymmetric gauge theories: The Double tensor multiplet and its interactions*, *Nucl. Phys.* **B587** (2000) 543–567, [[hep-th/0005086](#)].
- [19] B. de Wit and F. Saueressig, *Off-shell $N=2$ tensor supermultiplets*, *JHEP* **09** (2006) 062, [[hep-th/0606148](#)].

- [20] A. Galperin, E. Ivanov, S. Kalitsyn, V. Ogievetsky and E. Sokatchev, *Unconstrained $N=2$ Matter, Yang-Mills and Supergravity Theories in Harmonic Superspace*, *Class. Quant. Grav.* **1** (1984) 469–498.
- [21] M. de Roo, J. W. van Holten, B. de Wit and A. Van Proeyen, *Chiral Superfields in $N = 2$ Supergravity*, *Nucl. Phys.* **B173** (1980) 175–188.
- [22] B. de Wit, R. Philippe and A. Van Proeyen, *The Improved Tensor Multiplet in $N = 2$ Supergravity*, *Nucl. Phys.* **B219** (1983) 143–166.
- [23] B. de Wit and A. Van Proeyen, *Potentials and Symmetries of General Gauged $N=2$ Supergravity: Yang-Mills Models*, *Nucl. Phys.* **B245** (1984) 89–117.
- [24] B. de Wit, P. G. Lauwers and A. Van Proeyen, *Lagrangians of $N=2$ Supergravity - Matter Systems*, *Nucl. Phys.* **B255** (1985) 569–608.
- [25] E. Cremmer, C. Kounnas, A. Van Proeyen, J. P. Derendinger, S. Ferrara, B. de Wit et al., *Vector Multiplets Coupled to $N=2$ Supergravity: SuperHiggs Effect, Flat Potentials and Geometric Structure*, *Nucl. Phys.* **B250** (1985) 385–426.
- [26] B. de Wit, *$N=2$ electric - magnetic duality in a chiral background*, *Nucl. Phys. Proc. Suppl.* **49** (1996) 191–200, [[hep-th/9602060](#)].
- [27] B. de Wit, *$N=2$ symplectic reparametrizations in a chiral background*, *Fortsch. Phys.* **44** (1996) 529–538, [[hep-th/9603191](#)].
- [28] B. de Wit, S. Katmadas and M. van Zalk, *New supersymmetric higher-derivative couplings: Full $N=2$ superspace does not count!*, *JHEP* **01** (2011) 007, [[1010.2150](#)].
- [29] D. Butter, B. de Wit, S. M. Kuzenko and I. Lodato, *New higher-derivative invariants in $N=2$ supergravity and the Gauss-Bonnet term*, *JHEP* **12** (2013) 062, [[1307.6546](#)].
- [30] B. Sahoo and A. Sen, *Higher derivative corrections to non-supersymmetric extremal black holes in $N=2$ supergravity*, *JHEP* **09** (2006) 029, [[hep-th/0603149](#)].
- [31] N. Banerjee, S. Bansal and I. Lodato, *The Resolution of an Entropy Puzzle for 4D non-BPS Black Holes*, *JHEP* **05** (2016) 142, [[1602.05326](#)].
- [32] D. Butter, B. de Wit and I. Lodato, *Non-renormalization theorems and $N=2$ supersymmetric backgrounds*, *JHEP* **03** (2014) 131, [[1401.6591](#)].