Ergodic pairs for singular or degenerate fully nonlinear operators

I. Birindelli, F. Demengel, F. Leoni

Abstract

We study the ergodic problem for fully nonlinear operators which may be singular or degenerate when the gradient of solutions vanishes. We prove the convergence of both explosive solutions and solutions of Dirichlet problems for approximating equations. We further characterize the ergodic constant as the infimum of constants for which there exist bounded sub solutions. As intermediate results of independent interest, we prove a priori Lipschitz estimates depending only on the norm of the zeroth order term, and a comparison principle for equations having no zero order terms.

1 Introduction

In 1989, in a fundamental paper [\[18\]](#page-25-0), Lasry and Lions study solutions of

$$
-\Delta u + |\nabla u|^q + \lambda u = f(x) \text{ in } \Omega
$$

that blow up on the boundary of Ω . Here $q > 1$ and Ω is a \mathcal{C}^2 bounded domain in \mathbb{R}^N . In particular they introduce the concept of ergodic pair. Among other things, they prove that in the subquadratic case $q \leq 2$ there exists a unique constant c_{Ω} , called ergodic constant, and there exists a unique, up to a constant, solution of

$$
-\Delta \varphi + |\nabla \varphi|^q - c_{\Omega} = f(x) \text{ in } \Omega, \ \varphi = +\infty \text{ on } \partial \Omega.
$$

It is well known that for $q = 2, -c_{\Omega}$ is just the principal eigenvalue of $(-\Delta+f)(\cdot)$. This important paper has generated a huge and interesting literature, also in connection with the stochastic interpretation of the problem. Interestingly, while the concept of principal eigenvalue has been extended to fully nonlinear operators of different types (see e.g. [\[10\]](#page-25-1), [\[6\]](#page-24-0)), the notion of ergodic constant has not been much investigated in fully nonlinear settings. The scope of this paper is to give a thoroughly picture of the ergodic pairs and the related blowing up solutions and solutions with Dirichlet boundary condition for approximating equations.

We now detail the main theorems. In the whole paper Ω denotes a \mathcal{C}^2 bounded domain of \mathbb{R}^N ; \mathcal{S} denotes the space of symmetric matrices in \mathbb{R}^N . We consider a uniformly elliptic homogenous operator F, i.e. a continuous function $F : \mathcal{S} \to \mathbb{R}$ satisfying:

there exist
$$
0 < a < A
$$
 such that for all $M, N \in S$, with $N \ge 0$, and for all $t > 0$,
\n
$$
a \operatorname{tr}(N) \le F(M + N) - F(M) \le A \operatorname{tr}(N), \quad F(tM) = tF(M).
$$
\n(1.1)

We will always consider the differential operator

$$
G(\nabla u, D^2 u) = -|\nabla u|^\alpha F(D^2 u) + |\nabla u|^\beta
$$

with $\alpha > -1$ and $\alpha + 1 < \beta \leq \alpha + 2$. This reduces to the Lasry Lions case for $\alpha = 0$ and $a = A = 1$.

Theorem 1.1. Suppose that f is bounded and locally Lipschitz continuous in Ω , and that F satisfies [\(1.1\)](#page-0-0). Consider the Dirichlet problems

$$
\begin{cases}\n-|\nabla u|^{\alpha}F(D^2u) + |\nabla u|^{\beta} = f & \text{in } \Omega \\
u = 0 & \text{on } \partial\Omega,\n\end{cases}
$$
\n(1.2)

and, for $\lambda > 0$,

$$
\begin{cases}\n-|\nabla u|^{\alpha}F(D^2u) + |\nabla u|^{\beta} + \lambda |u|^{\alpha}u = f & \text{in } \Omega\\
u = 0 & \text{on } \partial\Omega.\n\end{cases}
$$
\n(1.3)

The following alternative holds.

- 1. Suppose that there exists a bounded sub solution of [\(1.2\)](#page-1-0). Then the solution u_{λ} of [\(1.3\)](#page-1-1) satisfies: (u_λ) is bounded and uniformly converging up to a sequence $\lambda_n \to 0$ to a solution of [\(1.2\)](#page-1-0).
- 2. Suppose that there is no solution for the Dirichlet problem (1.2) . Then, (u_λ) satisfies, up to a sequence $\lambda_n \to 0$ and locally uniformly in Ω ,
	- (a) $u_{\lambda} \rightarrow -\infty$;
	- (b) there exists a constant $c_{\Omega} \geq 0$ such that $\lambda |u_{\lambda}|^{\alpha} u_{\lambda} \to -c_{\Omega}$;
	- (c) c_{Ω} is an ergodic constant and $v_{\lambda} = u_{\lambda} + |u_{\lambda}|_{\infty}$ converges to a solution of the ergodic problem

$$
\begin{cases}\n-|\nabla v|^{\alpha} F(D^2 v) + |\nabla v|^{\beta} = f + c_{\Omega} & \text{in } \Omega \\
v = +\infty & \text{on } \partial\Omega\n\end{cases}
$$
\n(1.4)

whose minimum is zero.

The standard notion of viscosity solution fails when the operator is singular, i.e., in this paper, when α < 0, hence we will consider viscosity solutions as defined in [\[6\]](#page-24-0).

Theorem [1.1](#page-1-2) gives a construction of an ergodic pair (c_{Ω}, v) as a limit of solutions of Dirichlet problems when problem [\(1.2\)](#page-1-0) does not admit any solution. More in general, we will prove the existence of ergodic pairs under the regularity condition

$$
F(\nabla d(x) \otimes \nabla d(x)) \text{ is } C^2 \text{ in a neighborhood of } \partial \Omega, \qquad (1.5)
$$

where $d(x)$ denotes here the distance function from $\partial\Omega$. We observe that [\(1.5\)](#page-1-3) is certainly satisfied if the domain Ω is of class \mathcal{C}^3 and the operator F is \mathcal{C}^2 , but there can be also cases with non smooth F satisfying [\(1.5\)](#page-1-3). For instance, for all operators $F(M)$ which depend only on the eigenvalues of M, such as Pucci's operators, $F(\nabla d(x) \otimes \nabla d(x))$ is a constant function as long as $|\nabla d(x)| = 1$.

Under condition [\(1.5\)](#page-1-3), we will show that there exists a unique ergodic constant c_{Ω} , which shares some properties with the principal eigenvalue of the operator $|\nabla u|^\alpha F(D^2u)$, even when it is not the principal eigenvalue.

Indeed, let us define, as e.g. in [\[5\]](#page-24-1), [\[21\]](#page-25-2),

$$
\mu^* = \inf \{ \mu : \exists \varphi \in C(\overline{\Omega}), -|\nabla \varphi|^\alpha F(D^2 \varphi) + |\nabla \varphi|^\beta \le f + \mu \}.
$$

Note that μ^* depends on f and Ω , but if there is no ambiguity we will not precise this dependence.

Theorem 1.2. Suppose that f is bounded and locally Lipschitz continuous in Ω , and that F satisfies [\(1.1\)](#page-0-0) and [\(1.5\)](#page-1-3). Then, there exists an ergodic constant c_{Ω} and it satisfies

- 1. c_{Ω} is unique;
- 2. $c_{\Omega} = \mu^*$;
- 3. the map $\Omega \mapsto c_{\Omega}$ is nondecreasing with respect to the domain, and continuous;
- 4. if either $\alpha = 0$ or $\alpha \neq 0$ and $\sup_{\Omega} f + c_{\Omega} < 0$, then μ^* is not achieved. Moreover, if $\Omega' \subset\subset \Omega$, then $c_{\Omega'} < c_{\Omega}$.

In order to prove these results many questions need to be addressed. Clearly the first one is the existence of solutions for (1.3) when $\lambda > 0$, but even though it is fundamental, this is extraneous to the spirit of this note and it can be found in [\[8\]](#page-24-2). The interested reader will see that it is done through a Perron's procedure i.e. constructing sub and super solutions of [\(1.3\)](#page-1-1) together with a comparison principle and some Lipschitz estimates depending on the L^{∞} norm of the solution.

Theorems [1.1](#page-1-2) and [1.2](#page-2-0) are obtained by means of several intermediate results, most of which are of independent interest. A first fundamental tool is an interior Lipschitz estimate for solutions of equation [\(1.3\)](#page-1-1) that does not depend directly on the L^{∞} norm of the solution but only on the norm of the zero order term. In the linear case, these kind of estimates were obtained by Capuzzo Dolcetta, Leoni, Porretta in [\[11\]](#page-25-3), and the proof we use is inspired by theirs. In order to extend the result to the present fully nonlinear singular case, we have to address several non trivial technical difficulties, see Section [2.](#page-3-0) After that, we give the proof of Theorem [1.1](#page-1-2) in Section [3.](#page-10-0)

In Section [4](#page-12-0) we focus on existence and estimates for explosive solutions of the approximating λ equation, i.e. solutions u of

$$
\begin{cases}\n-|\nabla u|^{\alpha}F(D^2u) + |\nabla u|^{\beta} + \lambda |u|^{\alpha}u = f & \text{in } \Omega \\
u = +\infty & \text{on } \partial\Omega.\n\end{cases}
$$

Here, the function f is assumed to be continuous in Ω , but it is allowed to be unbounded on the boundary, as long as its growth is controlled. This is an important feature that will be needed in the proof of Theorem [1.2.](#page-2-0) Construction of explosive solutions in the fully nonlinear setting includes the works by Alarcón, Quaas $[1]$, by Esteban, Felmer and Quaas $[15]$ and by Demengel, Goubet $[14]$, where only suitable zero order terms are considered. Capuzzo Dolcetta, Leoni and Vitolo in [\[12,](#page-25-6) [13\]](#page-25-7) construct radial explosive solutions in some degenerate cases. The construction we do in order to give existence and estimates of blowing up solutions slightly differs from the standard proof for linear operators (see Remark [4.2\)](#page-16-0), and we obtain solutions satisfying non constant boundary asymptotics. Moreover, our proof can be carried on for other classes of operators, as e.g. the p–Laplacian or some generalizations such as

$$
F(p, M) = |p|^{\alpha} (q_1 tr M + q_2 \frac{Mp \cdot p}{|p|^2}),
$$

with $q_1 > 0$ and $q_1 + q_2 > 0$. When $p > 2$, using the variational form of the p-Laplacian, and its linearity with respect to the Hessian, Leonori and Porretta proved in [\[19\]](#page-25-8) such estimates and existence results. So our result for $\alpha < 0$ covers the case $p < 2$ that was not considered there.

Uniqueness of solution is a sensitive matter for degenerate elliptic equations, nonetheless, when $\alpha > 0$ i.e. in the degenerate case, uniqueness of explosive solutions is proved for any β . Instead, in the singular case, i.e. when $\alpha < 0$, we have some restriction on β and on the behavior of f near the boundary. In a forthcoming paper, [\[9\]](#page-25-9), we prove some $W^{2,p}$ estimates for any p when f is continuous in the case $\alpha \leq 0$, which give $\mathcal{C}^{1,\eta}$ estimates. This provides the uniqueness of blowing up solutions when $\alpha \leq 0$, without any restriction on β and f.

As we have seen when comparing the ergodic constant with the principal eigenvalue, in some sense the forcing term f in (1.2) plays the role of the zero order term for linear problems, hence it is not surprising that for $f < 0$ the comparison principle holds even if the operator is degenerate and the equation is not proper. This is the spirit of the comparison principle that we prove in Section [5.](#page-17-0) The change of equation that allows to prove the comparison principle of Theorem [5.1](#page-17-1) is sort of standard, but the computation which follows is original and ad hoc for our setting. The work of Leonori, Porretta and Riey [\[20\]](#page-25-10) has been a source of inspiration.

Finally, in Section [6,](#page-20-0) after proving the existence of ergodic constants and estimating the ergodic functions near the boundary, we complete the proof of Theorem [1.2.](#page-2-0)

Notations

- We use $d(x)$ to denote a \mathcal{C}^2 positive function in Ω with coincides with the distance function from the boundary in a neighborhood of $\partial\Omega$
- For $\delta > 0$, we set $\Omega_{\delta} = \{x \in \Omega : d(x) > \delta\}$
- We denote by \mathcal{M}^+ , \mathcal{M}^- the Pucci's operators with ellipticity constants a, A, namely, for all $M \in \mathcal{S}$,

$$
\mathcal{M}^+(M) = A \operatorname{tr}(M^+) - a \operatorname{tr}(M^-)
$$

$$
\mathcal{M}^-(M) = a \operatorname{tr}(M^+) - A \operatorname{tr}(M^-)
$$

and we often use that, as a consequence of (1.1) , for all $M, N \in \mathcal{S}$ one has

$$
\mathcal{M}^-(N) \le F(M+N) - F(M) \le \mathcal{M}^+(N)
$$

2 A priori Lipschitz-type estimates

In the note [\[8\]](#page-24-2), we prove the following result

Theorem 2.1. Assume that f is bounded and continuous in Ω . Then, for any $\lambda > 0$, there exists a unique solution $u_{\lambda} \in C(\overline{\Omega})$ of [\(1.3\)](#page-1-1), which is Lipschitz continuous up to the boundary, and satisfies

$$
|u_{\lambda}|_{W^{1,\infty}(\Omega)} \leq C(|u_{\lambda}|_{\infty}, |f-\lambda|u_{\lambda}|^{\alpha}u_{\lambda}|_{\infty}, a, A, \alpha, \beta).
$$

This is obtained, through Perron's method, by constructing sub and super solutions and using the following general comparison principle.

Theorem 2.2. Suppose that b is Lipschitz continuous in Ω , $\zeta : \mathbb{R} \to \mathbb{R}$ is a non decreasing function and f and g are continuous in Ω . Let u be a bounded by above viscosity sub solution of

$$
-|\nabla u|^{\alpha} F(D^2 u) + b(x)|\nabla u|^{\beta} + \zeta(u) \le g
$$

and let v be a bounded by below viscosity super solution of

$$
-|\nabla v|^{\alpha} F(D^2 v) + b(x)|\nabla v|^{\beta} + \zeta(v) \ge f.
$$

If either $g \leq f$ and ζ is increasing or $g < f$ then, $u \leq v$ on $\partial\Omega$ implies that $u \leq v$ in Ω .

For the proofs of the above results we refer to [\[8\]](#page-24-2).

The rest of this section is devoted to prove a priori Lipschitz estimates for solutions of the equation

$$
-|\nabla u|^{\alpha} F(D^2 u) + |\nabla u|^{\beta} + \lambda |u|^{\alpha} u = f,
$$
\n(2.1)

that depend on $\lambda |u|_{\infty}^{\alpha+1}$, but not on $|u|_{\infty}$. Our estimates will be a consequence of the following result, in which we denote by B the unit ball centred at the origin in \mathbb{R}^N .

Proposition 2.3. Let F satisfy [\(1.1\)](#page-0-0) and, for $\lambda \geq 0$, $\alpha > -1$ and $\beta > \alpha + 1$, let u and v be respectively a bounded sub solution and a bounded from below super solution of equation [\(2.1\)](#page-4-0) in B, with f Lipschitz continuous in B. Then, for any positive $p \geq \frac{(2+\alpha-\beta)^+}{\beta-\alpha-1}$ $\frac{(\mathcal{A} + \alpha - \beta)}{\beta - \alpha - 1}$, there exists a positive constant M, depending only on $p, \alpha, \beta, a, A, N, ||f - \lambda|u|^\alpha u||_\infty$ and on the Lipschitz constant of f, such that, for all $x, y \in B$ one has

$$
u(x) - v(y) \le \sup_B (u - v)^+ + M \frac{|x - y|}{(1 - |y|)^{\frac{\beta + \alpha^{-}}{\beta - \alpha - 1}}} \left[1 + \left(\frac{|x - y|}{(1 - |x|)} \right)^p \right]
$$

Proof. We argue as in the proof of Theorem 3.1 in [\[11\]](#page-25-3).

Let us define a "distance" function d which equals $1 - |x|$ near the boundary and it is extended in B as a \mathcal{C}^2 function satisfying, for some constant $c_1 > 0$,

$$
\begin{cases}\n d(x) = 1 - |x| & \text{if } |x| > \frac{1}{2} \\
 \frac{1-|x|}{2} \le d(x) \le 1 - |x| & \text{for all } x \in \bar{B} \\
 |Dd(x)| \le 1, \ -c_1 I_N \le D^2 d(x) \le 0 & \text{for all } x \in \bar{B}\n\end{cases}
$$

For $\xi = \frac{|x-y|}{d(x)}$ $\frac{x-y_1}{d(x)}$, we consider the function

$$
\phi(x, y) = \frac{k}{d(y)^{\tau}} |x - y| (L + \xi^{p}) + \sup_{B} (u - v)^{+}
$$

where $p > 0$ is a fixed exponent satisfying $p \geq \frac{(2+\alpha-\beta)^+}{\beta-\alpha-1}$ $\frac{(\beta+\alpha-\beta)^+}{\beta-\alpha-1}, \ \tau = \frac{\beta+\alpha^-}{\beta-\alpha^-}$ $\frac{\beta+\alpha}{\beta-\alpha-1}$ and L, k are suitably large positive constants to be chosen later.

The statement is proved if we show that for all $(x, y) \in B^2$ one has

$$
u(x) - u(y) \le \phi(x, y).
$$

By contradiction, let us assume that $u(x) - u(y) - \phi(x, y) > 0$ somewhere. Then, necessarily the supremum is achieved on a pair (x, y) with $x \neq y$ and $d(x), d(y) > 0$. Using Ishii's Lemma of [\[16\]](#page-25-11), one gets that on such a point (x, y) , for all $\epsilon > 0$, there exist symmetric matrices X_{ϵ} and Y_{ϵ} such that

$$
(\nabla_x \phi, X_{\epsilon}) \in J^{2,+}u(x), \; (-\nabla_y \phi, -Y_{\epsilon}) \in J^{2,-}v(y)
$$

$$
-\left(\frac{1}{\epsilon} + |D^2 \phi|\right)I_{2N} \le \left(\begin{array}{cc} X_{\epsilon} & O \\ O & Y_{\epsilon} \end{array}\right) \le D^2 \phi + \epsilon (D^2 \phi)^2.
$$
 (2.2)

We proceed in the proof by considering separately the cases $\alpha \geq 0$ and $\alpha < 0$.

The case $\alpha \geq 0$. Since u is a sub solution and v a super solution, by the positive 1–homogeneity of ${\cal F}$ we have in this case

$$
\begin{cases}\n-F(|\nabla_x \phi|^\alpha X_\epsilon) + |\nabla_x \phi|^\beta + \lambda |u|^\alpha u(x) \le f(x) \\
-F(-|\nabla_y \phi|^\alpha Y_\epsilon) + |\nabla_y \phi|^\beta + \lambda |v|^\alpha v(y) \ge f(y)\n\end{cases}
$$

Subtracting the above inequalities and using also that $u(x) - v(y) > \phi(x, y) \geq 0$, for any $t > 0$ we can write

$$
t F\left(|\nabla_x \phi|^\alpha X_\epsilon \right) - \left[F\left((1+t) |\nabla_x \phi|^\alpha X_\epsilon \right) - F\left(-|\nabla_y \phi|^\alpha Y_\epsilon \right) \right]
$$

$$
\leq |\nabla_y \phi|^\beta - |\nabla_x \phi|^\beta + f(x) - f(y),
$$

and therefore

$$
t\left|\nabla_x\phi\right|^\beta \leq F\left((1+t)|\nabla_x\phi|^\alpha X_\epsilon\right) - F\left(-|\nabla_y\phi|^\alpha Y_\epsilon\right) +|\nabla_y\phi|^\beta - |\nabla_x\phi|^\beta + t\left(f-\lambda|u|^\alpha u\right)^+ + f(x) - f(y).
$$

By the uniform ellipticity of F , it then follows that

$$
t \left| \nabla_x \phi \right|^\beta \leq \mathcal{M}^+ \left((1+t) \left| \nabla_x \phi \right|^\alpha X_\epsilon + \left| \nabla_y \phi \right|^\alpha Y_\epsilon \right) + \left| \nabla_y \phi \right|^\beta - \left| \nabla_x \phi \right|^\beta + t \left(f - \lambda |u|^\alpha u \right)^+ + f(x) - f(y) .
$$

By multiplying the right inequality of [\(2.2\)](#page-4-1) on the left and on the right by

$$
\left(\begin{array}{cc} \sqrt{1+t}|\nabla_x \phi|^{\alpha/2}I_N & O \\ O & |\nabla_y \phi|^{\alpha/2}I_N \end{array} \right)
$$

and testing the resulting inequality on vectors of the form (v, v) with $v \in \mathbb{R}^N$, we further obtain

$$
(1+t)|\nabla_x \phi|^\alpha X_\epsilon+|\nabla_y \phi|^\alpha Y_\epsilon\leq Z_{\alpha,t}+O(\epsilon)\,,
$$

with

$$
Z_{\alpha,t} = (1+t)|\nabla_x \phi|^{\alpha} D_{xx}^2 \phi + \sqrt{1+t}|\nabla_x \phi|^{\alpha/2} |\nabla_y \phi|^{\alpha/2} \left[D_{xy}^2 \phi + \left(D_{xy}^2 \phi \right)^t \right] + |\nabla_y \phi|^{\alpha} D_{yy}^2 \phi. \tag{2.3}
$$

Hence, after letting $\epsilon \to 0$, we get

$$
t\left|\nabla_x\phi\right|^{\beta} \le \mathcal{M}^+\left(Z_{\alpha,t}\right) + \left|\nabla_y\phi\right|^{\beta} - \left|\nabla_x\phi\right|^{\beta} + t\left(f - \lambda|u|^\alpha u\right)^+ + f(x) - f(y). \tag{2.4}
$$

We now proceed by evaluating the right hand side terms of [\(2.4\)](#page-5-0). An explicit computation shows that, setting $\eta = \frac{|x-y|}{d(y)}$ $\frac{x-y}{d(y)}$ and $\zeta = \frac{x-y}{|x-y|}$, one has

$$
\nabla_x \phi(x, y) = \frac{k}{d(y)^\tau} \left[(L + (1+p)\xi^p)\zeta - p\xi^{p+1}\nabla d(x) \right],
$$

as well as

$$
\nabla_y \phi(x, y) = -\frac{k}{d(y)^\tau} \left[(L + (1+p)\xi^p)\zeta + \tau \eta \left(L + \xi^p \right) \nabla d(y) \right].
$$

From now on we denote with c possibly different positive constants which depend only on p , N , a , A , α and β. As discussed in [\[11\]](#page-25-3), for $L > 1$ fixed suitably large depending only on p, one has

$$
|\nabla_x \phi| \ge ck \frac{1 + \xi^{p+1}}{d(y)^\tau}
$$
\n(2.5)

and

$$
|\nabla_x \phi|, \ |\nabla_y \phi| \le ck \frac{1 + \xi^{p+1}}{d(y)^{\tau+1}}.
$$
\n(2.6)

Moreover, we notice that one has also

$$
|\nabla_y \phi| \ge \frac{k}{d(y)^\tau} \left[L + (1+p)\xi^p - \tau \eta (L+\xi^p) |\nabla d(y)| \right] \ge ck \frac{1+\xi^p}{d(y)^\tau} \quad \text{if } \tau \eta \le \frac{1}{2}.
$$
 (2.7)

On the other hand, the second order derivatives of ϕ may be written as follows

$$
D_{xx}^{2}\phi = \frac{k}{d(y)^{\tau}} \left\{ \frac{[L+(1+p)\xi^{p}]}{[x-y]}B + p(1+p)\frac{\xi^{p-1}}{d(x)}T - p(1+p)\frac{\xi^{p}}{d(x)}(C+C^{t}) + p(1+p)\frac{\xi^{p+1}}{d(x)}\nabla d(x) \otimes \nabla d(x) - p\xi^{p+1}D^{2}d(x) \right\}
$$

$$
D_{xy}^{2}\phi = -\frac{k}{d(y)^{\tau}} \left\{ \frac{[L+(1+p)\xi^{p}]}{[x-y]}B + p(1+p)\frac{\xi^{p-1}}{d(x)}T - p(1+p)\frac{\xi^{p}}{d(x)}C^{t} + \frac{\tau [L+(1+p)\xi^{p}]}{d(y)}E - \frac{\tau p\xi^{p+1}}{d(y)}\nabla d(x) \otimes \nabla d(y) \right\}
$$

$$
D_{yy}^{2}\phi = \frac{k}{d(y)^{\tau}} \left\{ \frac{[L+(1+p)\xi^{p}]}{[x-y]}B + p(1+p)\frac{\xi^{p-1}}{d(x)}T + \frac{\tau [L+(1+p)\xi^{p}]}{d(y)}(E+E^{t}) + \frac{\tau (\tau+1)\eta (L+\xi^{p})}{d(y)}\nabla d(y) \otimes \nabla d(y) - \tau \eta (L+\xi^{p})D^{2}d(y) \right\}
$$

with $B = I_N - \zeta \otimes \zeta$, $T = \zeta \otimes \zeta$, $C = \zeta \otimes \nabla d(x)$ and $E = \zeta \otimes \nabla d(y)$. Therefore, the matrix $Z_{\alpha,t}$ defined in [\(2.3\)](#page-5-1) is given by

$$
Z_{\alpha,t} = \frac{k}{d(y)^{\tau}} \left\{ \left(\sqrt{1+t} |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right)^2 \left[\frac{(L+(1+p)\xi^p)}{|x-y|} B + p(1+p) \frac{\xi^{p-1}}{d(x)} T \right] \right.- \sqrt{1+t} |\nabla_x \phi|^{\alpha/2} \left(\sqrt{1+t} |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right) \frac{p(1+p)\xi^p}{d(x)} (C + C^t) - |\nabla_y \phi|^{\alpha/2} \left(\sqrt{1+t} |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right) \frac{\tau(L+(1+p)\xi^p)}{d(y)} (E + E^t) + p(1+t) |\nabla_x \phi|^{\alpha} \left[\frac{(1+p)\xi^{p+1}}{d(x)} \nabla d(x) \otimes \nabla d(x) - \xi^{p+1} D^2 d(x) \right] + \sqrt{1+t} |\nabla_x \phi|^{\alpha/2} |\nabla_y \phi|^{\alpha/2} \frac{\tau p \xi^{(p+1)}}{d(y)} [\nabla d(x) \otimes \nabla d(y) + \nabla d(y) \otimes \nabla d(x)] + \tau |\nabla_y \phi|^{\alpha} \left[\frac{(1+\tau)\eta(L+\xi^p)}{d(y)} \nabla d(y) \otimes \nabla d(y) - \eta(L+\xi^p) D^2 d(y) \right] \right\},
$$

and, recalling that $\xi = |x - y|/d(x)$ and that $d < 1$ in B, this yields the estimate

$$
\mathcal{M}^{+}(Z_{\alpha,t}) \leq \frac{ck}{d(y)^{\tau}} \left\{ \left(\sqrt{1+t} |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right)^2 \frac{1+\xi^p}{|x-y|} \right. \\ \left. + \sqrt{1+t} |\nabla_x \phi|^{\alpha/2} | \sqrt{1+t} |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} | \frac{\xi^p}{d(x)} \right. \\ \left. + |\nabla_y \phi|^{\alpha/2} | \sqrt{1+t} |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} | \frac{1+\xi^p}{d(y)} \right. \\ \left. + (1+t) |\nabla_x \phi|^{\alpha} \frac{\xi^{p+1}}{d(x)} + \sqrt{1+t} |\nabla_x \phi|^{\alpha/2} |\nabla_y \phi|^{\alpha/2} \frac{\xi^{p+1}}{d(y)} + |\nabla_y \phi|^{\alpha} \eta \frac{1+\xi^p}{d(y)} \right\}.
$$

By observing that

$$
\left| \sqrt{1+t} |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right| \leq (\sqrt{t+1}-1) |\nabla_x \phi|^{\alpha/2} + \left| |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right|
$$

and by applying the trivial inequalities $\sqrt{1+t} - 1 \le t$, $\sqrt{1+t} (\sqrt{t+1} - 1) \le t$, $\sqrt{1+t} \le 1+t$, after rearranging terms we then deduce

$$
\mathcal{M}^{+}(Z_{\alpha,t}) \leq \frac{ck}{d(y)^{\tau}} \left\{ t^{2} |\nabla_{x}\phi|^{\alpha} \frac{1+\xi^{p}}{|x-y|} \right.
$$

+
$$
t \left[|\nabla_{x}\phi|^{\alpha} \frac{\xi^{p} + \xi^{p+1}}{d(x)} + |\nabla_{x}\phi|^{\alpha/2} |\nabla_{y}\phi|^{\alpha/2} \left(\frac{1+\xi^{p} + \xi^{p+1}}{d(y)} + \frac{\xi^{p}}{d(x)} \right) \right]
$$

+
$$
\left(|\nabla_{x}\phi|^{\alpha/2} - |\nabla_{y}\phi|^{\alpha/2} \right)^{2} \frac{1+\xi^{p}}{|x-y|}
$$

+
$$
||\nabla_{x}\phi|^{\alpha/2} - |\nabla_{y}\phi|^{\alpha/2} |\left(|\nabla_{x}\phi|^{\alpha/2} \frac{\xi^{p}}{d(x)} + |\nabla_{y}\phi|^{\alpha/2} \frac{1+\xi^{p}}{d(y)} \right)
$$

+
$$
|\nabla_{x}\phi|^{\alpha} \frac{\xi^{p+1}}{d(x)} + |\nabla_{y}\phi|^{\alpha} \frac{(1+\xi^{p})\eta}{d(y)} + |\nabla_{x}\phi|^{\alpha/2} |\nabla_{y}\phi|^{\alpha/2} \frac{\xi^{p+1}}{d(y)} \right\}
$$
(2.8)

We now recall that, as proved in [\[11\]](#page-25-3), for all $q, \gamma > 0$ one has

$$
\frac{\xi^q}{d(x)^\gamma} \le 2^\gamma \frac{1 + \xi^{q+\gamma}}{d(y)^\gamma} \,. \tag{2.9}
$$

Moreover, if $\alpha \geq 2$, the mean value theorem, the bounds [\(2.6\)](#page-6-0), [\(2.9\)](#page-7-0) and the explicit expression of $\nabla_x \phi + \nabla_y \phi$ imply that

$$
\left| |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right| \leq c \max \left\{ |\nabla_x \phi|^{\alpha/2 - 1}, |\nabla_y \phi|^{\alpha/2 - 1} \right\} |\nabla_x \phi + \nabla_y \phi|
$$

$$
\leq c k^{\alpha/2} \frac{(1 + \xi^{p+1})^{\alpha/2 - 1}}{d(y)^{\alpha/2(\tau + 1) - 1}} \left(\frac{\xi^p}{d(x)} + \frac{1 + \xi^p}{d(y)} \right) |x - y|
$$

$$
\leq c \left[k \frac{(1 + \xi^{p+1})}{d(y)^{\tau + 1}} \right]^{\alpha/2} |x - y|
$$

Analogously, if $\alpha < 2$ but $\tau \eta \leq 1/2$, from [\(2.5\)](#page-6-1), [\(2.7\)](#page-6-2) and again [\(2.9\)](#page-7-0) we deduce

$$
\left| |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right| \le c \left[k \frac{(1 + \xi^p)}{d(y)^\tau} \right]^{\alpha/2 - 1} k \frac{(1 + \xi^{p+1})}{d(y)^{\tau+1}} |x - y|
$$

and therefore, since $\xi \leq \frac{1}{2\tau - 1}$ for $\eta \leq \frac{1}{2\tau}$, we obtain in this case

$$
\left| |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right| \leq c \left[k \frac{(1 + \xi^{p+1})}{d(y)^{\tau + 2/\alpha}} \right]^{\alpha/2} |x - y|.
$$

Finally, if $\alpha < 2$ and $\tau\, \eta > 1/2,$ that is $|x-y| > d(y)/2\tau,$ we have

$$
\left| |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right|
$$

\n
$$
\leq |\nabla_x \phi + \nabla_y \phi|^{\alpha/2} \leq c \left[\frac{k(1 + \xi^{p+1})|x - y|}{d(y)^{\tau+1}} \right]^{\alpha/2}
$$

\n
$$
\leq c \left[\frac{k(1 + \xi^{p+1})}{d(y)^{\tau+1}} \right]^{\alpha/2} \left(\frac{2\tau}{d(y)} \right)^{1 - \alpha/2} |x - y| = c \left[k \frac{(1 + \xi^{p+1})}{d(y)^{\tau+2/\alpha}} \right]^{\alpha/2} |x - y|.
$$

Thus, in all cases we obtain

$$
\left| |\nabla_x \phi|^{\alpha/2} - |\nabla_y \phi|^{\alpha/2} \right| \le c \left[k \frac{(1 + \xi^{p+1})}{d(y)^{\tau + \max\{1, 2/\alpha\}}} \right]^{\alpha/2} |x - y| \le c \frac{|\nabla_x \phi|^{\alpha/2}}{d(y)^{\max\{\alpha/2, 1\}}} |x - y|.
$$
 (2.10)

By using inequalities (2.6) , (2.5) , (2.9) and (2.10) , from estimate (2.8) it then follows

$$
\mathcal{M}^+(Z_{\alpha,t}) \le \frac{ck|\nabla_x \phi|^\alpha}{d(y)^\tau} \left\{ t^2 \frac{1+\xi^p}{|x-y|} + t \frac{1+\xi^{p+2}}{d(y)^{\alpha/2+1}} + |x-y| \frac{1+\xi^{p+2}}{d(y)^{\alpha+2}} \right\}.
$$
 (2.11)

Moreover, since $p \ge \frac{2+\alpha-\beta}{\beta-\alpha-1}$ and $\tau \ge \frac{\alpha+2}{2(\beta-\alpha-1)}$, by using again (2.5) , we further deduce

$$
\mathcal{M}^+(Z_{\alpha,t}) \le ck|\nabla_x \phi|^\alpha \left\{ t^2 \frac{1+\xi^p}{d(y)^\tau |x-y|} + t \frac{|\nabla_x \phi|^{\beta-\alpha}}{k^{\beta-\alpha}} + \frac{|x-y|(1+\xi^{p+2})}{d(y)^\tau + \alpha+2} \right\}.
$$

Using the above inequality jointly with [\(2.4\)](#page-5-0) yields

$$
t \left| \nabla_x \phi \right| ^{\beta - \alpha} \leq c k \left\{ t^2 \frac{1 + \xi^p}{d(y)^\tau |x - y|} + t \frac{\left| \nabla_x \phi \right| ^{\beta - \alpha}}{k^{\beta - \alpha}} + \frac{\left| x - y \right| (1 + \xi^{p+2})}{d(y)^\tau + \alpha + 2} \right\}
$$

$$
+ \left| \nabla_x \phi \right| ^{-\alpha} \left(\left| \nabla_y \phi \right| ^{\beta} - \left| \nabla_x \phi \right| ^{\beta} \right) + t \left| \nabla_x \phi \right| ^{-\alpha} \left(f - \lambda |u|^\alpha u \right)^+
$$

$$
+ \left| \nabla_x \phi \right| ^{-\alpha} \left(f(x) - f(y) \right) ,
$$

and therefore, being $\beta > \alpha + 1,$ for k sufficiently large one has

$$
\frac{t}{2} |\nabla_x \phi|^{\beta-\alpha} - t^2 \frac{ck(1+\xi^p)}{d(y)^\tau |x-y|} \leq \frac{ck|x-y|(1+\xi^{p+2})}{d(y)^\tau + \alpha+2} + |\nabla_x \phi|^{-\alpha} \left(|\nabla_y \phi|^{\beta} - |\nabla_x \phi|^{\beta} \right) + t |\nabla_x \phi|^{-\alpha} \left(f - \lambda |u|^\alpha u \right)^+ + |\nabla_x \phi|^{-\alpha} \left(f(x) - f(y) \right).
$$

We now choose $t > 0$ in order to maximize the left hand side, namely

$$
t = \frac{|\nabla_x \phi|^{\beta-\alpha} d(y)^\tau |x-y|}{4ck(1+\xi^p)}.
$$

We then obtain

$$
|\nabla_x \phi|^{2(\beta-\alpha)} \leq c \left\{ \frac{k^2 (1+\xi^{2(p+1)})}{d(y)^{2\tau+\alpha+2}} + k|\nabla_x \phi|^{-\alpha} \frac{(1+\xi^p) \left(|\nabla_y \phi|^\beta - |\nabla_x \phi|^\beta \right)}{|x-y| d(y)^\tau} + |\nabla_x \phi|^{\beta-2\alpha} (f-\lambda|u|^\alpha u)^+ + k|\nabla_x \phi|^{-\alpha} \frac{(1+\xi^p) (f(x)-f(y))}{|x-y| d(y)^\tau} \right\}.
$$

Moreover, arguing as for [\(2.10\)](#page-8-0) in the case $\alpha \geq 2$, we also have

$$
\left| |\nabla_y \phi|^{\beta} - |\nabla_x \phi|^{\beta} \right| \leq c k^{\beta} |x - y| \frac{(1 + \xi^{p+1})^{\beta}}{d(y)^{(r+1)\beta}},
$$

so that

$$
\begin{aligned} |\nabla_x \phi|^{2(\beta-\alpha)} &\leq C\left\{\frac{k^2(1+\xi^{(p+1)})^2}{d(y)^{2\tau+\alpha+2}}+|\nabla_x \phi|^{-\alpha}\frac{k^{\beta+1}(1+\xi^{p+1})^{\beta+1}}{d(y)^{\tau(\beta+1)+\beta}}\right.\\&\left.+\left|\nabla_x \phi\right|^{\beta-2\alpha}+|\nabla_x \phi|^{-\alpha}\frac{k(1+\xi^p)}{d(y)^{\tau}}\right\}\,, \end{aligned}
$$

for some constant $C > 0$ depending now also on $||(f - \lambda |u|^\alpha u)^+||_\infty$ and on the Lipschitz constant of f.

By inequality [\(2.5\)](#page-6-1) it then follows

$$
|\nabla_x \phi|^{2(\beta-\alpha)} \leq C \left\{ \frac{|\nabla_x \phi|^2}{d(y)^{\alpha+2}} + \frac{|\nabla_x \phi|^{\beta-\alpha+1}}{d(y)^{\beta}} + |\nabla_x \phi|^{\beta-2\alpha} + |\nabla_x \phi|^{1-\alpha} \right\}
$$

$$
\leq C \left\{ \frac{|\nabla_x \phi|^{2+\frac{\alpha+2}{\tau}}}{k^{\frac{\alpha+2}{\tau}}} + \frac{|\nabla_x \phi|^{\beta-\alpha+1+\frac{\beta}{\tau}}}{k^{\frac{\beta}{\tau}}} + |\nabla_x \phi|^{\beta-2\alpha} \right\},
$$

Recalling that $\alpha > 0$, $\beta > \alpha + 1$ and $\tau = \frac{\beta}{\beta - \alpha - 1}$, the last inequality gives a contradiction for k large enough.

The case $\alpha < 0$. As proved in [\[7\]](#page-24-4), if $\alpha < 0$ a sub solution u and super solution v of equation [\(2.1\)](#page-4-0) satisfy respectively in the viscosity sense

$$
\begin{cases}\n-F(D^2u) + |\nabla u|^{\beta-\alpha} + \lambda |\nabla u|^{-\alpha} |u|^\alpha u \leq |\nabla u|^{-\alpha} f \\
-F(D^2v) + |\nabla v|^{\beta-\alpha} + \lambda |\nabla v|^{-\alpha} |v|^\alpha v \geq |\nabla v|^{-\alpha} f\n\end{cases}
$$

From [\(2.2\)](#page-4-1) in this case it then follows that

$$
\begin{cases}\n-F(X_{\epsilon}) + |\nabla_x \phi|^{\beta-\alpha} + \lambda |\nabla_x \phi|^{-\alpha} |u|^{\alpha} u(x) \leq |\nabla_x \phi|^{-\alpha} f(x) \\
-F(-Y_{\epsilon}) + |\nabla_y \phi|^{\beta-\alpha} + \lambda |\nabla_y \phi|^{-\alpha} |v|^{\alpha} v(y) \geq |\nabla_y \phi|^{-\alpha} f(y)\n\end{cases}
$$

and, arguing as in the previous case, we now obtain for any $t > 0$

$$
t\left|\nabla_x\phi\right|^{\beta-\alpha} \leq \mathcal{M}^+\left(Z_{0,t}\right) + \left|\nabla_y\phi\right|^{\beta-\alpha} - \left|\nabla_x\phi\right|^{\beta-\alpha} + t\left|\nabla_x\phi\right|^{-\alpha}\left(f-\lambda|u|^\alpha u\right) - \left(f-\lambda|u|^\alpha u\right)\left(\left|\nabla_y\phi\right|^{-\alpha} - \left|\nabla_x\phi\right|^{-\alpha}\right) + \left|\nabla_y\phi\right|^{-\alpha}\left(f(x)-f(y)\right),
$$

where $Z_{0,t}$ is defined by [\(2.3\)](#page-5-1) (with $\alpha = 0$).

By applying inequalities [\(2.11\)](#page-8-1) (with $\alpha = 0$), [\(2.5\)](#page-6-1), [\(2.6\)](#page-6-0), [\(2.10\)](#page-8-0), in the present case, taking into account that $\beta - \alpha > 1$ and $0 < -\alpha < 1$, we deduce that

$$
|\nabla_x \phi|^{2(\beta-\alpha)} \le C \left\{ \frac{|\nabla_x \phi|^2}{d(y)^2} + \frac{|\nabla_x \phi|^{\beta-\alpha+1}}{d(y)^{\beta-\alpha}} + |\nabla_x \phi|^{\beta-2\alpha} \right\},
$$

for some constant $C > 0$ depending on $p, \alpha, \beta, a, A, N, ||f - \lambda|u|^\alpha u||_\infty$ and on the Lipschitz constant of f. Since now $\tau = \frac{\beta - \alpha}{\beta - \alpha - 1}$, we reach a contradiction for k sufficiently large as before.

As in [\[11\]](#page-25-3), the above Proposition and a scaling argument for solutions of equation [\(2.1\)](#page-4-0) give the following result.

Theorem 2.4. Let F satisfy [\(1.1\)](#page-0-0) and, for $\lambda \geq 0, \alpha > -1$ and $\beta > \alpha + 1$, let u be a continuous solution of equation [\(2.1\)](#page-4-0) in $\Omega \subset \mathbb{R}^N$, with f Lipschitz continuous in Ω . Then, u is locally Lipschitz continuous in Ω and there exists a positive constant M, depending only on $\alpha, \beta, a, A, N, ||f - \lambda|u|^\alpha u||_\infty$ and on the Lipschitz constant of f, such that at any differentiability point $x \in \Omega$ one has

$$
|\nabla u(x)| \le \frac{M}{\operatorname{dist}_{\partial\Omega}(x)^{\frac{1}{\beta-\alpha-1}}}.
$$

3 Proof of Theorem [1.1.](#page-1-2)

By using the Lipschitz estimates obtained in the previous section, we can now prove Theorem [1.1.](#page-1-2)

Proof of Theorem [1.1.](#page-1-2) Let u_{λ} be a solution of [\(1.3\)](#page-1-1). We begin by giving a bound that will be useful in the whole proof. Observe that u^+_λ is a sub solution of

$$
-|\nabla u_\lambda^+|^{\alpha} F(D^2 u_\lambda^+) \le |f|_\infty;
$$

from known estimates, see [\[6\]](#page-24-0), this implies that

$$
|u_{\lambda}^{+}|_{\infty} \le c_1 |f|_{\infty}^{\frac{1}{1+\alpha}}.
$$
\n(3.1)

Let us consider first the case when there exists a sub solution φ of [\(1.2\)](#page-1-0). Then, $\varphi - |\varphi|_{\infty}$ is a sub solution of equation [\(1.3\)](#page-1-1), and by the comparison principle we deduce $u_\lambda \geq \varphi - |\varphi|_\infty$. Thus, in this case (u_λ) is uniformly bounded in Ω . The Lipschitz estimates in Theorem [2.1](#page-3-1) then yield that u_λ is uniformly converging up to a sequence to a Lipschitz solution of problem [\(1.2\)](#page-1-0).

We now treat the second case, i.e. we suppose that [\(1.2\)](#page-1-0) has no solutions. In particular $|u_\lambda|_\infty$ diverges, since otherwise we could extract from (u_λ) a subsequence converging to a solution of [\(1.2\)](#page-1-0).

On the other hand, since $-\left(\frac{|f|_{\infty}}{\lambda}\right)$ $\frac{d}{\lambda}$ $\frac{1}{\lambda}$ is a sub solution of [\(1.3\)](#page-1-1), by the comparison principle we obtain $u_\lambda^- \leq \left(\frac{|f|_\infty}{\lambda}\right)$ $\frac{d}{\lambda}$ $\frac{1}{1+\alpha}$, which, jointly with [\(3.1\)](#page-10-1), yields $\lambda |u_{\lambda}|_{\infty}^{1+\alpha} \leq c_1 |f|_{\infty}$. Hence, there exists $(x_{\lambda}) \subset \Omega$ such that $u_\lambda(x_\lambda) = -|u_\lambda|_\infty \to -\infty$ and there exists a constant $c_\Omega \geq 0$ such that, up to a subsequence, $\lambda |u_{\lambda}|_{\infty}^{1+\alpha} \to c_{\Omega}.$

We will show, as in [\[18\]](#page-25-0) (see also [\[19\]](#page-25-8) and [\[21\]](#page-25-2)), that $v_\lambda = u_\lambda + |u_\lambda|_\infty = u_\lambda - u_\lambda(x_\lambda)$ converges up to a subsequence to a function v such that the pair (c_{Ω}, v) solves [\(1.4\)](#page-1-4).

Clearly, v_λ satisfies in Ω

$$
-|\nabla v_\lambda|^{\alpha} F(D^2 v_\lambda) + |\nabla v_\lambda|^{\beta} + \lambda (v_\lambda)^{1+\alpha} = f + \lambda (v_\lambda^{\alpha+1} - |u_\lambda|^{\alpha} u_\lambda) \ge f.
$$

Next, we set

$$
\gamma = \frac{2+\alpha-\beta}{\beta-1-\alpha},
$$

and for $s, \delta_o > 0$ to be chosen sufficiently small, let us consider the function

$$
\begin{aligned}\n\phi(x) &= \frac{\sigma}{(d(x) + s)^{\gamma}} - \frac{\sigma}{(\delta_o + s)^{\gamma}} \quad \text{if } \gamma > 0, \\
\phi(x) &= -\sigma \log(d(x) + s) + \sigma \log(\delta_o + s) \quad \text{if } \gamma = 0,\n\end{aligned} \tag{3.2}
$$

where $\sigma = ((\gamma + 1)\frac{a}{2})^{\frac{1}{\beta - \alpha - 1}} \gamma^{-1}$ if $\gamma > 0$, $\sigma = \frac{a}{2}$ if $\gamma = 0$. A direct computation shows that, for $d(x) \leq \delta_0$ with δ_0 small enough, in the case $\gamma > 0$ one has

$$
-|\nabla\phi|^{\alpha}\mathcal{M}^-(D^2\phi)+|\nabla\phi|^{\beta}+\lambda\phi^{1+\alpha}\leq -\frac{a(\sigma\gamma)^{\alpha+1}}{2(d+s)^{(\gamma+1)\beta}}+\frac{A(\sigma\gamma)^{\alpha+1}|D^2d|_{\infty}}{(d+s)^{(\gamma+1)(\alpha+1)}}+\frac{\lambda\sigma^{\alpha+1}}{(d+s)^{\gamma(\alpha+1)}},
$$

and, in the case $\gamma = 0$,

$$
-|\nabla \phi|^{\alpha} \mathcal{M}^-(D^2 \phi) + |\nabla \phi|^{\beta} + \lambda \phi^{1+\alpha} \leq -\frac{\sigma^{\alpha+2}}{(d+s)^{\alpha+2}} + \frac{A \sigma^{\alpha+1} |D^2 d|_{\infty}}{(d+s)^{(\alpha+1)}} + \lambda \left(-\sigma \log(d+s)\right)^{\alpha+1}.
$$

In both cases, by the ellipticity of F and for δ_o and s sufficiently small, we obtain

$$
-|\nabla \phi|^{\alpha} F(D^2 \phi) + |\nabla \phi|^{\beta} + \lambda \phi^{1+\alpha} \le -|f|_{\infty} \le f(x) \quad \text{in } \Omega \setminus \overline{\Omega}_{\delta_0}.
$$

Moreover, one has $\phi = 0 \le v_\lambda$ on $\partial \Omega_{\delta_\alpha}$ and $\phi \le |u_\lambda|_\infty = v_\lambda$ on $\partial \Omega$ for λ sufficiently small in dependence of s. The comparison principle then yields

$$
v_{\lambda} \ge \phi > 0 \quad \text{in } \Omega \setminus \overline{\Omega}_{\delta_0} \,. \tag{3.3}
$$

Since $v_{\lambda}(x_{\lambda}) = 0$, from [\(3.3\)](#page-11-0) we deduce that $(x_{\lambda}) \subset \Omega_{\delta_o}$. The interior Lipschitz estimate of Theorem [2.4](#page-10-2) then yields that $v_{\lambda} = u_{\lambda} - u_{\lambda}(x_{\lambda})$ is locally uniformly bounded and locally uniformly Lipschitz continuous. This proves both statement [2a](#page-1-5) of the theorem and that (v_λ) is locally uniformly converging up to a subsequence to a Lipschitz continuous function $v_o \geq 0$ in Ω . Moreover, since also (x_λ) converges up to a subsequence to some point $x_o \in \Omega_{\delta_0}$, we obtain $v_o(x_o) = 0$. We observe further that, locally uniformly in Ω , one has

$$
\lim_{\lambda \to 0} \lambda |u_{\lambda}|^{\alpha} u_{\lambda} = \lim_{\lambda \to 0} \lambda |u_{\lambda}|_{\infty}^{\alpha+1} \frac{|v_{\lambda} - |u_{\lambda}|_{\infty}|^{\alpha} (v_{\lambda} - |u_{\lambda}|_{\infty})}{|u_{\lambda}|_{\infty}^{\alpha+1}} = -c_{\Omega}.
$$

This yields statement [2b](#page-1-6) and, letting $\lambda \to 0$ in the equation satisfied by v_λ , also that v_o is a viscosity solution of

$$
-|\nabla v_o|^{\alpha} F(D^2 v_o) + |\nabla v_o|^{\beta} = f + c_{\Omega}.
$$

Finally, letting $\lambda \to 0$ in inequality [\(3.3\)](#page-11-0), we obtain $v_o \ge \phi$ in $\Omega \setminus \Omega_{\delta_0}$, which in turn implies, by letting $s \to 0$ and $x \to \partial \Omega$, that $v_o(x) \to +\infty$ as $d(x) \to 0$. This completely proves statement [2c](#page-1-7) and concludes the proof of the theorem.

 \Box

4 Existence of explosive solutions.

In this section we prove the existence of solutions of [\(2.1\)](#page-4-0) blowing up at the boundary. The results obtained here will be used in the proof of the existence of ergodic pairs.

Throughout the present section, on the one hand we assume the regularity condition [\(1.5\)](#page-1-3), but, on the other hand, we drop the assumption on the boundedness of the right hand side f , and we consider continuous functions in Ω , possibly unbounded as $d(x) \to 0$.

For what follows, we set

$$
\gamma = \frac{2+\alpha-\beta}{\beta-1-\alpha},
$$

and we denote by $C(x)$ a non negative function of class C^2 in Ω satisfying in a neighborhood of $\partial\Omega$

$$
C(x) = ((\gamma + 1)F(\nabla d(x) \otimes \nabla d(x)))^{\frac{1}{\beta - \alpha - 1}} \gamma^{-1} \quad \text{if} \quad \gamma > 0,
$$

\n
$$
C(x) = F(\nabla d(x) \otimes \nabla d(x)) \quad \text{if} \quad \gamma = 0.
$$
\n(4.1)

Theorem 4.1. Let $\beta \in (\alpha + 1, \alpha + 2], \lambda > 0$ and let F satisfy [\(1.1\)](#page-0-0) and [\(1.5\)](#page-1-3). Let further $f \in \mathcal{C}(\Omega)$ be bounded from below and such that

$$
\lim_{d(x)\to 0} f(x)d(x)^{\frac{\beta}{\beta-1-\alpha}-\gamma_0} = 0, \qquad (4.2)
$$

for some $\gamma_0 \geq 0$. Then, there exists a solution u of

$$
\begin{cases}\n-|\nabla u|^{\alpha}F(D^2u) + |\nabla u|^{\beta} + \lambda |u|^{\alpha}u = f & \text{in } \Omega, \\
u = +\infty & \text{on } \partial\Omega.\n\end{cases}
$$
\n(4.3)

Moreover, any solution u of [\(4.3\)](#page-12-1) satisfies: for any $\nu > 0$ and for any $0 \le \gamma_1 \le \gamma_0$, with $\gamma_1 < 1$, and $\gamma_1 < \gamma$ when $\gamma > 0$, there exists $D = \frac{D_1}{\lambda^{1/(\alpha+1)}}$, with $D_1 > 0$ depending on $\nu, \gamma_1, \alpha, \beta, a, A, |d|_{\mathcal{C}^2(\Omega)}, |C|_{\mathcal{C}^2(\Omega)}$ and on f, such that, for all $x \in \Omega$,

$$
\frac{C(x)}{d(x)^{\gamma}} - \frac{\nu}{d(x)^{\gamma - \gamma_1}} - D \le u(x) \le \frac{C(x)}{d(x)^{\gamma}} + \frac{\nu}{d(x)^{\gamma - \gamma_1}} + D \quad \text{if } \gamma > 0,
$$
\n
$$
|\log d(x)| \left(C(x) - \nu d(x)^{\gamma_1} \right) - D \le u(x) \le |\log d(x)| \left(C(x) + \nu d(x)^{\gamma_1} \right) + D \quad \text{if } \gamma = 0.
$$
\n
$$
(4.4)
$$

Furthermore, the solution u is unique

- for $\alpha > 0$ and any β ,
- for $\alpha < 0$ and any $\beta > \frac{1-\alpha-\alpha^2}{1-\alpha}$ $\frac{1-\alpha-\alpha^2}{1-\alpha}$, provided that f satisfies [\(4.2\)](#page-12-2) with $\gamma_0 > -\alpha \gamma$.

Proof. We give the proof in the case $\gamma > 0$, the reader can easily see the changes to be made when $\gamma = 0.$

1. Estimates and existence.

We will get the conclusion by showing that, for every $\nu > 0$ and for any $0 \le \gamma_1 \le \gamma_0$, with $\gamma_1 <$ $\min\{1,\gamma\}$, there exist $D=\frac{D_1}{\lambda^{1/\alpha+1}}>0$, a super solution \overline{w} and a sub solution \underline{w}^s satisfying

$$
C(x)(d+s)^{-\gamma} - \nu(d+s)^{-\gamma+\gamma_1} - D \leq \underline{w}^s(x) \leq \overline{w}(x) \leq C(x)d^{-\gamma} + \nu d^{-\gamma+\gamma_1} + D,
$$

for any $s > 0$ sufficiently small.

Assume for a while that this is proved. Then, problem (4.3) does have solutions and any solution u of [\(4.3\)](#page-12-1) satisfies $\underline{w}^o \le u \le \overline{w}$. Indeed, for any $R > 0$, we can consider the solution u_R of

$$
\begin{cases}\n-|\nabla u_R|^{\alpha} F(D^2 u_R) + |\nabla u_R|^{\beta} + \lambda |u_R|^{\alpha} u_R = f_R & \text{in } \Omega \\
u_R = R & \text{on } \partial\Omega,\n\end{cases}
$$

with $f_R = \min\{f, R\}$. By Theorem [2.2,](#page-3-2) u_R is monotone increasing with respect to R and satisfies $\underline{w}^s \leq u_R \leq \overline{w}$, provided that $R > \max_{\partial \Omega} \underline{w}^s(x)$. Moreover, u_R is locally uniformly Lipschitz continuous by Theorem [2.1.](#page-3-1) Thus, u_R is locally uniformly convergent as $R \to +\infty$ to a solution <u>u</u> of [\(4.3\)](#page-12-1) such that $\underline{w}^0 \le \underline{u} \le \overline{w}$. By definition, \underline{u} is the so called minimal explosive solution. The maximal explosive solution \overline{u} is then obtained as the limit for $\delta \to 0$ of the minimal explosive solutions in Ω_{δ} . Thus, it follows that for any solution u of problem (4.3) one has

$$
C(x)d^{-\gamma}-\nu d^{-\gamma+\gamma_1}-D\leq \underline{u}\leq u\leq \overline{u}\leq C(x)d^{-\gamma}+\nu d^{-\gamma+\gamma_1}+D\,.
$$

Let us now proceed to the construction of \overline{w} and \underline{w}^s .

Let $\delta > 0$ be so small that in the set $\Omega \setminus \Omega_{2\delta} = \{d(x) < 2\delta\}$ the function d satisfies $|\nabla d| = 1$ and C satisfies [\(4.1\)](#page-12-3). For $x \in \Omega \setminus \Omega_{2\delta}$, let us consider the function

$$
\varphi(x) = C(x)d(x)^{-\gamma} + \nu d(x)^{\gamma_1 - \gamma}.
$$

One has

$$
\nabla \varphi(x) = -\gamma C(x) d^{-\gamma - 1} \left(1 + \nu \frac{(\gamma - \gamma_1)}{\gamma C(x)} d^{\gamma_1} \right) \nabla d + d^{-\gamma} \nabla C
$$

and

$$
D^2 \varphi(x) = \gamma (\gamma + 1) C(x) d^{-\gamma - 2} \left(1 + \nu \frac{(\gamma - \gamma_1)(\gamma - \gamma_1 + 1)}{\gamma (\gamma + 1) C(x)} d^{\gamma_1} \right) \nabla d \otimes \nabla d
$$

$$
- \gamma C(x) d^{-\gamma - 1} \left(1 + \nu \frac{(\gamma - \gamma_1)}{\gamma C(x)} d^{\gamma_1} \right) D^2 d
$$

$$
- \gamma d^{-\gamma - 1} \left(\nabla d \otimes \nabla C + \nabla C \otimes \nabla d \right) + d^{-\gamma} D^2 C.
$$

By ellipticity of F and by definition of $C(x)$ it then follows

$$
F(D^2\varphi) \leq \frac{\gamma(\gamma+1)C(x)}{d^{\gamma+2}} \left(1 + \nu \frac{(\gamma-\gamma_1)(\gamma-\gamma_1+1)}{\gamma(\gamma+1)C(x)} d^{\gamma_1}\right) F(\nabla d \otimes \nabla d) + \frac{K_1}{d^{\gamma+1}}
$$

=
$$
\frac{(\gamma C(x))^{\beta-\alpha}}{d^{\gamma+2}} \left(1 + \nu \frac{(\gamma-\gamma_1)(\gamma-\gamma_1+1)}{\gamma(\gamma+1)C(x)} d^{\gamma_1}\right) + \frac{K_1}{d^{\gamma+1}},
$$

for a constant $K_1 > 0$ depending on $\nu, \alpha, \beta, \gamma_1, a, A, |D^2 d|_{\infty}$ and $|C|_{\mathcal{C}^2(\Omega)}$. In what follows we denote by K_i , $i = 1, 2, \ldots$, different constants depending on these quantities.

Hence, we obtain

$$
-|\nabla\varphi|^{\alpha}F(D^{2}\varphi)+|\nabla\varphi|^{\beta}
$$
\n
$$
\geq |\nabla\varphi|^{\alpha}\left[-\frac{(\gamma C(x))^{\beta-\alpha}}{d^{\gamma+2}}\left(1+\nu\frac{(\gamma-\gamma_{1})(\gamma-\gamma_{1}+1)}{\gamma(\gamma+1)C(x)}d^{\gamma_{1}}\right)-\frac{K_{1}}{d^{\gamma+1}}+|\nabla\varphi|^{\beta-\alpha}\right]
$$
\n
$$
\geq |\nabla\varphi|^{\alpha}\left[-\frac{(\gamma C(x))^{\beta-\alpha}}{d^{\gamma+2}}\left(1+\nu\frac{(\gamma-\gamma_{1})(\gamma-\gamma_{1}+1)}{\gamma(\gamma+1)C(x)}d^{\gamma_{1}}\right)-\frac{K_{1}}{d^{\gamma+1}} + \frac{(\gamma C(x))^{\beta-\alpha}}{d^{\gamma+1}(\beta-\alpha)}\left(1+\nu\frac{(\gamma-\gamma_{1})}{\gamma C(x)}d^{\gamma_{1}}\right)^{\beta-\alpha}-\frac{K_{2}}{d^{\gamma+1}(\beta-\alpha)-1}\right]
$$
\n
$$
\geq |\nabla\varphi|^{\alpha}\left[-\frac{(\gamma C(x))^{\beta-\alpha}}{d^{\gamma+2}}\left(1+\nu\frac{(\gamma-\gamma_{1})(\gamma-\gamma_{1}+1)}{\gamma(\gamma+1)C(x)}d^{\gamma_{1}}\right)-\frac{K_{1}}{d^{\gamma+1}} + \frac{(\gamma C(x))^{\beta-\alpha}}{d^{\gamma+1}(\beta-\alpha)}\left(1+\nu\frac{(\beta-\alpha)(\gamma-\gamma_{1})}{\gamma C(x)}d^{\gamma_{1}}\right)-\frac{K_{2}}{d^{\gamma+1}(\beta-\alpha)-1}\right]
$$

Recalling that $\gamma = \frac{\alpha+2-\beta}{\beta-\alpha-1}$, we finally deduce

$$
-|\nabla \varphi|^{\alpha} F(D^2 \varphi) + |\nabla \varphi|^{\beta}
$$

\n
$$
\geq |\nabla \varphi|^{\alpha} \left[\nu \frac{(\gamma - \gamma_1)(1 + \gamma_1)}{(\gamma + 1)} \frac{(\gamma C(x))^{\beta - \alpha - 1}}{d^{\gamma + 2 - \gamma_1}} - \frac{K_3}{d^{\gamma + 1}} \right]
$$

\n
$$
= \frac{|(\gamma C(x) + \nu (\gamma - \gamma_1) d^{\gamma_1}) \nabla d - d \nabla C|^{\alpha}}{d^{\beta - \alpha - 1} - \gamma_1} \left[\nu \frac{(\gamma - \gamma_1)(1 + \gamma_1)}{(\gamma + 1)} (\gamma C(x))^{\beta - \alpha - 1} - K_3 d^{1 - \gamma_1} \right]
$$

Since $\gamma_1 \leq \gamma_0$, by assumption [\(4.2\)](#page-12-2) on f the last inequality implies that, for δ sufficiently small,

$$
-|\nabla \varphi|^\alpha F(D^2 \varphi) + |\nabla \varphi|^\beta \ge f(x) \quad \text{in } \Omega \setminus \Omega_{2\delta},
$$

and therefore also that

$$
-|\nabla \varphi|^\alpha F(D^2 \varphi) + |\nabla \varphi|^\beta + \lambda \varphi^{\alpha+1} \ge f(x) \quad \text{in } \Omega \setminus \Omega_{2\delta}.
$$

Clearly, the same inequality holds also for $\varphi_1(x) = \varphi(x) + D$, for any $D > 0$. Next, for $x \in {\delta \le d(x) \le 2\delta}$, we consider the function

$$
\varphi_2(x) = \delta^{-\gamma} C(x) e^{\frac{1}{d(x) - 2\delta} + \frac{1}{\delta}} + E,
$$

with $E > 0$ to be conveniently fixed. A direct computation shows that, for δ sufficiently small in dependence of $|\nabla C|_{\infty}$, one has

$$
|\nabla \varphi_2|^{\alpha} |F(D^2 \varphi_2)| + |\nabla \varphi_2|^{\beta} \leq K_4 \quad \text{in } \overline{\Omega}_\delta \setminus \Omega_{2\delta}.
$$

Thus, if E is chosen satisfying $E \geq \left(\frac{|f|_{L^{\infty}(\Omega_{\delta})} + K_4}{\lambda} \right)$ $\frac{\alpha_{\delta}+K_4}{\lambda}$ $\frac{1}{\alpha+1}$, then we obtain

$$
-|\nabla \varphi_2|^{\alpha} F(D^2 \varphi_2) + |\nabla \varphi_2|^{\beta} + \lambda \varphi_2^{\alpha+1} \ge f(x) \quad \text{in } \overline{\Omega}_\delta \setminus \Omega_{2\delta}.
$$

Let us further restrict the smallness of δ by requiring that

$$
A\gamma \delta^{-\gamma - 1} + (\gamma - \gamma_1)\delta^{\gamma_1 - \gamma - 1} < a\delta^{-2 - \gamma} \,. \tag{4.5}
$$

Then, setting $D = \left(\frac{|f|_{L^{\infty}(\Omega_{\delta})} + K_4}{\lambda}\right)$ $\left(\frac{\Omega_{\delta}+K_4}{\lambda}\right)^{\frac{1}{\alpha+1}}$ and $E=\nu\delta^{\gamma_1-\gamma}+D$, the function

$$
\overline{w}(x) = \begin{cases}\nC(x)d(x)^{-\gamma} + \nu d^{-\gamma + \gamma_1} + D & \text{for } d < \delta \\
\delta^{-\gamma} C(x)e^{\frac{1}{d(x) - 2\delta} + \frac{1}{\delta}} + E & \text{for } \delta \le d \le 2\delta \\
E & \text{for } d > 2\delta\n\end{cases}
$$

is the required super solution in Ω . Indeed, in the set Ω_{δ} , \overline{w} is of class \mathcal{C}^2 and it is a super solution by the properties of φ_2 and by the fact that locally constant functions satisfy $|\nabla u|^{\alpha} F(D^2 u) = 0$. On the other hand, \overline{w} is a super solution in $\Omega \setminus \Omega_{2\delta}$ by the properties of φ_1 and φ_2 and the fact that $\varphi_2(x) < \varphi_1(x)$ for $d(x) > \delta$, as it follows by evaluating at points x such that $d(x) = \delta$ the derivatives of φ_1 and φ_2 along the direction $\nabla d(x)$ and by using [\(4.5\)](#page-14-0).

As far as the sub solution is concerned, for $s > 0$, $\nu > 0$ and $x \in \Omega \setminus \Omega_{2\delta}$, let us consider the function

$$
\varphi^{s}(x) = C(x)(d(x) + s)^{-\gamma} - \nu(d(x) + s)^{-\gamma + \gamma_1}.
$$

Analogous computations as above give that

$$
-|\nabla\varphi^s|^{\alpha} F(D^2\varphi^s) + |\nabla\varphi^s|^{\beta} + \lambda|\varphi^s|^{\alpha}\varphi^s
$$

$$
\leq |\nabla\varphi^s|^{\alpha} \left[-\frac{(\gamma C(x))^{\beta-\alpha-1}}{(d+s)^{\gamma+2-\gamma_1}} \nu \frac{(\gamma-\gamma_1)(1+\gamma_1)}{\gamma+1} + \frac{K_5}{(d+s)^{\gamma+1}} \right] + \lambda \frac{A^{\alpha+1}}{(d+s)^{\gamma(\alpha+1)}} \leq f(x),
$$

for δ and s sufficiently small, since f is bounded from below.

Moreover, setting $C_{\delta} = \max_{\{d(x) = \delta\}} C(x)$ and choosing $D \geq C_{\delta} \delta^{-\gamma} + \left(\frac{|f^{-}|_{\infty}}{\lambda}\right)$ $\frac{1}{\lambda}$ $\frac{1}{\alpha+1}$, the constant function $C_{\delta}(\delta + s)^{-\gamma} - \nu(\delta + s)^{-\gamma + \gamma_1} - D$ is also a sub solution in Ω . Therefore, the function

$$
\underline{w}^s(x) = \begin{cases}\n\max \left\{ C(x)(d+s)^{-\gamma} - \nu(d+s)^{-\gamma+\gamma_1} - D, \\
C_\delta(\delta+s)^{-\gamma} - \nu(\delta+s)^{-\gamma+\gamma_1} - D\right\} & \text{in } \Omega \setminus \Omega_\delta \\
C_\delta(\delta+s)^{-\gamma} - \nu(\delta+s)^{-\gamma+\gamma_1} - D & \text{in } \Omega_\delta\n\end{cases}
$$

is the wanted sub solution.

2. Uniqueness.

We prove that $u = \overline{u}$.

Remark that, by estimates [\(4.4\)](#page-12-4), for any $\theta < 1$ and for any $c \in \mathbb{R}$, there exists δ such that $\theta \overline{u}(x) - c \leq$ $\underline{u}(x)$ for $d(x) \leq \delta$.

The case $\alpha \geq 0$. Observe that, for all $t \in \mathbb{R}$ and $c > 0$, one has

$$
|t-c|^{\alpha}(t-c) - |t|^{\alpha}t \le -2^{-\alpha}c^{\alpha+1}.
$$

From this, we deduce that

$$
-|\nabla(\theta \overline{u}-c)|^{\alpha} F(D^2(\theta \overline{u}-c)) + |\nabla(\theta \overline{u}-c)|^{\beta} + \lambda |\theta \overline{u}-c|^{\alpha}(\theta \overline{u}-c) \leq \theta^{\alpha+1} f(x) - \lambda 2^{-\alpha} c^{\alpha+1},
$$

and the choice

$$
c = c_{\theta} = \left(\frac{2^{\alpha}(1 - \theta^{\alpha+1})|f^{-}|\infty}{\lambda}\right)^{\frac{1}{\alpha+1}}
$$

then yields

$$
-|\nabla(\theta \overline{u} - c)|^{\alpha} F(D^2(\theta \overline{u} - c)) + |\nabla(\theta \overline{u} - c)|^{\beta} + \lambda |\theta \overline{u} - c|^{\alpha}(\theta \overline{u} - c)
$$

$$
\leq \theta^{\alpha+1} f(x) - (1 - \theta^{\alpha+1}) ||f^-||_{\infty} \leq f(x).
$$

By applying Theorem [2.2,](#page-3-2) it then follows that $\theta \overline{u} - c_{\theta} \leq \underline{u}$ in Ω , and letting $\theta \to 1$ we obtain the uniqueness of the explosive solution in the case $\alpha \geq 0$.

The case $\alpha < 0$. In this case we use the inequality

$$
|t-c|^{\alpha}(t-c)-|t|^{\alpha}t\leq -2^{\alpha}(\alpha+1)K^{\alpha}c\,,
$$

which holds true for all $0 < c < K$ and $t \in \mathbb{R}$ such that $|t| \leq K$. Let $C_1 = \sup_{\Omega} |\overline{u}(x)| d(x)^{\gamma}$, which is finite by [\(4.4\)](#page-12-4). Then, for any $\delta > 0$, one has $|\overline{u}| < \frac{C_1}{\delta^{\gamma}}$ in Ω_{δ} . Therefore, for any $0 < \theta < 1$ and $0 < c < \frac{C_1}{\delta^{\gamma}}$, and for $x \in \Omega_{\delta}$, we have

$$
-|\nabla(\theta \overline{u} - c)|^{\alpha} F(D^2(\theta \overline{u} - c)) + |\nabla(\theta \overline{u} - c)|^{\beta} + \lambda |\theta \overline{u} - c|^{\alpha}(\theta \overline{u} - c)
$$

$$
\leq \theta^{\alpha+1} f(x) - \lambda (2C_1)^{\alpha} (\alpha+1) \delta^{-\alpha \gamma} c.
$$

We choose, as before,

$$
c = c_{\theta,\delta} = \frac{|f^-|_{\infty}(1 - \theta^{\alpha+1})}{\lambda (2C_1)^{\alpha}(\alpha+1)\delta^{-\alpha\gamma}},
$$

which is admissible for δ sufficiently small, since $\alpha > -1$. This yields

$$
-|\nabla(\theta\overline{u}-c_{\theta,\delta})|^{\alpha}F(D^2(\theta\overline{u}-c_{\theta,\delta})) + |\nabla(\theta\overline{u}-c_{\theta,\delta})|^{\beta} + \lambda|\theta\overline{u}-c_{\theta,\delta}|^{\alpha}(\theta\overline{u}-c_{\theta,\delta}) \leq f \quad \text{in } \Omega_{\delta}.
$$

On the other hand, by estimates [\(4.4\)](#page-12-4) with $\nu = 1$, we have

$$
\theta \overline{u} - c_{\theta, \delta} \le \underline{u} \quad \text{ on } \partial \Omega_{\delta}
$$

provided that

$$
\delta = \delta_{\theta} = \left(\frac{a(1-\theta)}{2(1+D)}\right)^{\frac{1}{\gamma_1}}
$$

.

With this choice of δ , we then deduce from Theorem [2.2](#page-3-2) that $\theta \overline{u} - c_{\theta, \delta_{\theta}} \leq \underline{u}$ in $\Omega_{\delta_{\theta}}$. Finally, we let θ → 1. We observe that, by the restrictions assumed on β and f in the case α < 0, we can choose γ_1 satisfying $\gamma_1 > -\alpha \gamma$. Therefore, $c_{\theta, \delta_{\theta}} \to 0$ as θ → 1, and we conclude that $\overline{u} \leq \underline{u}$ in Ω . satisfying $\gamma_1 > -\alpha \gamma$. Therefore, $c_{\theta, \delta_{\theta}} \to 0$ as $\theta \to 1$, and we conclude that $\overline{u} \leq \underline{u}$ in Ω .

Remark 4.2. Let us put in evidence that estimates (4.4) imply that any solution u of (4.3) satisfies

$$
\lim_{d(x)\to 0} \frac{u(x) d(x)^\gamma}{C(x)} = 1 \text{ if } \gamma > 0, \quad \lim_{d(x)\to 0} \frac{u(x)}{|\log d(x)| C(x)} = 1 \text{ if } \gamma = 0.
$$

Moreover, if f satisfies [\(4.2\)](#page-12-2) with $\gamma_0 = 0$, then necessarily $\gamma_1 = 0$ and [\(4.4\)](#page-12-4) reduce to

$$
(C(x) - \nu)d(x)^{-\gamma} - D \le u(x) \le (C(x) + \nu)d(x)^{-\gamma} + D \text{ if } \gamma > 0,
$$

$$
(C(x) - \nu) |\log d(x)| - D \le u(x) \le (C(x) + \nu) |\log d(x)| + D \text{ if } \gamma = 0,
$$

for any $\nu > 0$, with $D > 0$ depending in particular on ν and λ . The above estimates are the classical ones for explosive solutions firstly obtained in the semilinear case in [\[18\]](#page-25-0), where $C(x)$ is a constant function. Also the case $\gamma_0 = 1$ has been considered in [\[18\]](#page-25-0), and in this case more refined estimates have been obtained. In the nonlinear case, analogous estimates for $\gamma_0 \geq 1$ would require further regularity assumptions on the non constant function $C(x)$. Estimates [\(4.4\)](#page-12-4) are interesting in the intermediate cases $0 \leq \gamma_0 < 1$, in which they are new also for linear operators and yield a uniqueness result in the non linear singular case $\alpha < 0$.

5 A comparison principle for non linear degenerate/singular equations without zero order terms

This section is devoted to some comparison principle for fully non linear equations without zero order terms. For analogous results concerning non singular operators, see [\[2\]](#page-24-5), [\[3\]](#page-24-6).

Theorem 5.1. Let b be a continuous and bounded function in Ω and, when $\alpha \neq 0$, let f be a bounded continuous function such that $f \leq -m < 0$. Let u and v be respectively sub and super solution of

$$
-|\nabla u|^{\alpha} F(D^2 u) + b(x)|\nabla u|^{\beta} = f \quad \text{in } \Omega.
$$
 (5.1)

If u and v are bounded and at least one of the two is Lipschitz continuous then the comparison principle holds i.e.

$$
u \le v \quad on \quad \partial\Omega \Rightarrow u \le v \quad in \quad \Omega.
$$

Proof. Without loss of generality, we will suppose that u is Lipschitz continuous.

The case $\alpha = 0$ is quite standard, it is enough to construct strict sub solutions that converge uniformly to u. For $\kappa > 0$ to be chosen, let $u_{\epsilon} = u + \epsilon e^{-\kappa x_1} - \epsilon$, with e.g. $\Omega \subset \{x_1 > 0\}$. By the mean value theorem:

$$
|\nabla u_{\epsilon}|^{\beta} \leq |\nabla u|^{\beta} + \beta \kappa \epsilon e^{-\kappa x_1} (|\nabla u| + \epsilon k)^{\beta - 1}.
$$

Since $\beta \leq 2$ in the case $\alpha = 0$, we can choose κ such that $a\kappa > 2\beta(|\nabla u|_{\infty} + \epsilon k)^{\beta - 1}$. Then, u_{ϵ} is a strict sub solution of [\(5.1\)](#page-17-2), being

$$
F(D^2u_{\epsilon}) \ge F(D^2u) + a\kappa^2 \epsilon e^{-\kappa x_1}
$$

\n
$$
\ge F(D^2u) + \beta k \epsilon e^{-kx_1} (|\nabla u|_{\infty} + \epsilon k)^{\beta - 1} + \frac{a\kappa^2 \epsilon}{2} e^{-\kappa x_1}
$$

\n
$$
\ge |\nabla u_{\epsilon}|^{\beta} - f + \frac{a\kappa^2 \epsilon}{2} e^{-\kappa x_1}.
$$

Furthermore $u_{\epsilon} \leq u \leq v$ on $\partial \Omega$. We are now in a position to apply the comparison principle in Theorem [2.2,](#page-3-2) and we obtain that $u_{\epsilon} \leq v$ in Ω . To conclude, let $\epsilon \to 0$. This computation has been done for a classical solution u , but, with obvious changes, it can be made rigorous for viscosity solutions.

For the case $\alpha \neq 0$ and $f < 0$, we use the change of function $u = \varphi(z)$, $v = \varphi(w)$ with

$$
\varphi(s) = -\gamma(\alpha + 1)\log\left(\delta + e^{-\frac{s}{\alpha + 1}}\right).
$$

This function is used in [\[2\]](#page-24-5), [\[3\]](#page-24-6), [\[4\]](#page-24-7), [\[19\]](#page-25-8), [\[20\]](#page-25-10).

We choose δ small enough in order that the range of φ covers the ranges of u and v. The constant γ will be chosen small enough depending only on a, α , β , inf $\Omega(-f)$ and $|b|_{\infty}$; in this proof, any constant of this type will be called universal. Observe that $\varphi' > 0$ while $\varphi'' < 0$.

In the viscosity sense, z and w are respectively sub and super solution of

$$
-|\nabla z|^{\alpha} F(D^2 z + \frac{\varphi''(z)}{\varphi'(z)} \nabla z \otimes \nabla z) + b(x)\varphi'(z)^{\beta - \alpha - 1} |\nabla z|^{\beta} + \frac{-f}{(\varphi'(z))^{\alpha + 1}} = 0.
$$
 (5.2)

We define

$$
H(x,s,p)=\frac{-a\varphi^{\prime\prime}(s)}{\varphi^\prime(s)}|p|^{2+\alpha}+b(x)\varphi^\prime(s)^{\beta-\alpha-1}|p|^\beta+\frac{-f(x)}{\varphi^\prime(s)^{\alpha+1}}.
$$

The point is to prove that at \bar{x} , a maximum point of $z - w$, $\frac{\partial H(\bar{x},s,p)}{\partial s} > 0$ for all p. This will be sufficient to get a contradiction. A simple computation gives

$$
\varphi' = \frac{\gamma e^{-\frac{s}{\alpha+1}}}{\delta + e^{-\frac{s}{\alpha+1}}}, \ \varphi'' = \frac{-\gamma \delta e^{-\frac{s}{\alpha+1}}}{(\alpha+1)(\delta + e^{-\frac{s}{\alpha+1}})^2}.
$$

Hence

$$
\left(\frac{-\varphi''}{\varphi'}\right)' = \frac{\delta}{(\alpha+1)^2} \frac{e^{-\frac{s}{\alpha+1}}}{(\delta + e^{-\frac{s}{\alpha+1}})^2} \text{ i.e. } \left(\frac{-\varphi''}{\varphi'}\right)' = -\frac{\varphi''}{(\alpha+1)\gamma} > 0.
$$

Differentiating H with respect to s gives:

$$
\partial_s H = a|p|^{\alpha+2} \frac{-\varphi''}{(\alpha+1)\gamma} + (-f) \frac{-(\alpha+1)\varphi''}{(\varphi')^{\alpha+2}} + b(x)|p|^{\beta}(\beta-\alpha-1)(\varphi')^{\beta-\alpha-2}\varphi''.
$$

Since $-\varphi''$ is positive, we need to prove that

$$
K := \frac{a|p|^{\alpha+2}}{(\alpha+1)\gamma} + (-f)\frac{\alpha+1}{(\varphi')^{\alpha+2}} - |b|_{\infty}|p|^{\beta}(\beta-\alpha-1)(\varphi')^{\beta-\alpha-2} > 0.
$$

We start by treating the case $\beta < \alpha + 2$.

Observe first that the boundedness of u and v , implies that there exists universal positive constants c_o and $c₁$ such that

$$
c_o \gamma \le \varphi' \le c_1 \gamma.
$$

Hence, it is easy to see that there exist three positive universal constants C_i , $i = 1, 2, 3$ such that

$$
K > \frac{C_1|p|^{\alpha+2}}{\gamma} + \frac{C_2}{\gamma^{\alpha+2}} - \frac{C_3|p|^\beta}{\gamma^{\alpha+2-\beta}}.
$$

We choose $\gamma = \min\left\{1, (\frac{C_3}{C_2})^{\beta}, (\frac{C_3}{C_1})^{\frac{1}{\alpha+1-\beta}}\right\}$. With this choice of γ , for $|p| \leq 1$,

$$
\frac{C_1|p|^{\alpha+2}}{\gamma}+\frac{C_2}{\gamma^{\alpha+2}}-\frac{C_3|p|^\beta}{\gamma^{\alpha+2-\beta}}\geq \frac{C_2}{\gamma^{\alpha+2}}-\frac{C_3}{\gamma^{\alpha+2-\beta}}>0;
$$

while for $|p| \geq 1$,

$$
\frac{C_1|p|^{\alpha+2}}{\gamma} + \frac{C_2}{\gamma^{\alpha+2}} - \frac{C_3|p|^\beta}{\gamma^{\alpha+2-\beta}} \ge \frac{(C_1)|p|^{\alpha+2}}{\gamma} - \frac{C_3|p|^\beta}{\gamma^{\alpha+2-\beta}} > 0.
$$

If $\beta = \alpha + 2$, just take $\gamma < \frac{a}{(\alpha+1)|b|_{\infty}}$.

This gives that for γ small enough depending only on min($-f$), α , $|b|_{\infty}$ and β one has, for some universal constant C,

$$
\partial_s H(x, s, p) \ge C > 0. \tag{5.3}
$$

We now conclude the proof of the comparison principle.

We will distinguish the case $\alpha > 0$ and $\alpha < 0$. In the first case we introduce $\psi_j(x, y) = z(x) - w(y) - \frac{j}{2}|x - y|^2$ where $q > \frac{\alpha+2}{\alpha+1}$. We detail the case $\alpha > 0$.

Suppose by contradiction that $u > v$ somewhere in Ω , then $z > w$ somewhere, since φ is increasing, while on the boundary $z \leq w$. Then the supremum of $z - w$ is positive and it is achieved inside Ω . Hence ψ_j reaches a positive maximum in $(x_j, y_j) \in \Omega \times \Omega$.

By Ishii's lemma [\[16\]](#page-25-11), there exists $(X_j, Y_j) \in S \times S$ such that

$$
(p_j, X_j) \in \overline{J}^{2,+}z(x_j), (p_j, -Y_j) \in \overline{J}^{2,-}w(y_j), \text{ with } p_j = j(x_j - y_j)
$$

and

$$
\left(\begin{array}{cc} X_j & 0 \\ 0 & Y_j \end{array}\right) \leq j \left(\begin{array}{cc} I & -I \\ -I & I \end{array}\right).
$$

On (x_j, y_j) , by a continuity argument, for j large enough one has

$$
z(x_j) > w(y_j) + \frac{\sup(z - w)}{2}
$$

.

Note for later purposes that since z or w are Lipschitz, $p_j = j(x_j - y_j)$ is bounded. Observe that, the monotonicity of $\frac{\varphi''}{\varphi'}$ $\frac{\rho}{\varphi'}$ implies that

$$
N = p_j \otimes p_j \left(\frac{\varphi''(z(x_j))}{\varphi'(z(x_j))} - \frac{\varphi''(w(y_j))}{\varphi'(w(y_j))} \right) \leq 0.
$$

Using the fact that z and w are respectively sub and super solutions of the equation (5.2) , the estimate (5.3) and that H is decreasing in the second variable, one obtains:

$$
0 \geq \frac{-f(x_j)}{(\varphi')^{\alpha+1}(z(x_j))} - |p_j|^{\alpha} F(X_j + \frac{\varphi''(z(x_j))}{\varphi'(z(x_j))} p_j \otimes p_j) + b(x_j) |p_j|^{\beta} \varphi'(z(x_j))^{\beta-\alpha-1}
$$

\n
$$
\geq \frac{-f(x_j)}{(\varphi')^{\alpha+1}(z(x_j))} - |p_j|^{\alpha} F(-Y_j + \frac{\varphi''(w(y_j))}{\varphi'(w(y_j))} p_j \otimes p_j) +
$$

\n
$$
+ a|p_j|^{2+\alpha} \left(\frac{\varphi''(w(y_j))}{\varphi'(w(y_j))} - \frac{\varphi''(z(x_j))}{\varphi'(z(x_j))}\right) + |p_j|^{\beta} b(x_j) (\varphi'(z(x_j))^{\beta-\alpha-1})
$$

\n
$$
\geq \frac{f(y_j) - f(x_j)}{(\varphi'(w(y_j))^{\alpha+1}} + (b(x_j) - b(y_j)) |p_j|^{\beta} (\varphi'(w(y_j)))^{\beta-\alpha-1}
$$

\n
$$
+ H(x_j, z(x_j), p_j) - H(x_j, w(y_j), p_j)
$$

\n
$$
\geq C(z(x_j) - w(y_j)) + \frac{o(1)}{\gamma^{\alpha+1}}.
$$

Here we have used the continuity of f and b, the boundedness of p_j and that

$$
\psi(x_j, y_j) \ge \sup(\psi(x_j, x_j), \psi(y_j, y_j)).
$$

Passing to the limit one gets a contradiction, since (x_j, y_j) converges to (\bar{x}, \bar{x}) such that $z(\bar{x}) > w(\bar{z})$. This ends the case $\alpha \geq 0$.

In the case $\alpha < 0$, the proof is similar but we need to make sure that one can choose $x_j \neq y_j$. This can be done proceeding as in [8]. can be done proceeding as in [\[8\]](#page-24-2).

6 Ergodic pairs

In this section we consider, for $c \in \mathbb{R}$, the equation

$$
-|\nabla u|^{\alpha} F(D^2 u) + |\nabla u|^{\beta} = f + c \text{ in } \Omega.
$$
\n(6.1)

Definition 6.1. Suppose that c is some constant (depending on f, Ω , β , α , and F) such that there exists $\varphi \in \mathcal{C}(\Omega)$, solution of [\(6.1\)](#page-20-1), such that $\varphi \to +\infty$ at $\partial \Omega$. We will say that c is an ergodic constant, φ is an ergodic function and (c, φ) is an ergodic pair.

We suppose, as usual, that $\alpha > -1$, $\beta \in (\alpha + 1, \alpha + 2]$ and recall that $\gamma = \frac{2+\alpha-\beta}{\beta-\alpha-1}$ and $C(x)$ satisfies [\(4.1\)](#page-12-3). In the following subsections, we prove the existence and show several properties of ergodic pairs.

6.1 Existence of ergodic constants and boundary behavior of ergodic functions

Theorem [1.1](#page-1-2) provides the existence of a nonnegative ergodic constant under the assumption that problem [\(1.2\)](#page-1-0) does not have a solution. In the next result we obtain the existence of ergodic constants using approximating explosive solutions.

Theorem 6.2. Let F and f be as in Theorem [4.1,](#page-12-5) and assume further that f is locally Lipschitz continuous in Ω . Then, there exists an ergodic constant $c \in \mathbb{R}$.

Proof. By Theorem [4.1,](#page-12-5) for $\lambda > 0$ there exists a solution U_{λ} of problem [\(4.3\)](#page-12-1), which satisfies estimates [\(4.4\)](#page-12-4). Recalling the dependence on λ of the constant D which appears in [\(4.4\)](#page-12-4), we see that $\lambda |U_{\lambda}|^{\alpha}U_{\lambda}$ is locally bounded in Ω , uniformly with respect to $0 < \lambda < 1$. Let us fix an arbitrary point $x_0 \in \Omega$. Then, there exists $c \in \mathbb{R}$ such that, up to a sequence $\lambda_n \to 0$,

$$
\lambda |U_{\lambda}(x_0)|^{\alpha}U_{\lambda}(x_0) \to -c.
$$

On the other hand, Theorem [2.4](#page-10-2) yields that U_{λ} is locally uniformly Lipschitz continuous. Therefore, for x in a compact subset of Ω , one has

$$
\lambda ||U_{\lambda}(x)|^{\alpha}U_{\lambda}(x) - |U_{\lambda}(x_0)|^{\alpha}U_{\lambda}(x_0)| \leq \lambda |U_{\lambda}(x) - U_{\lambda}(x_0)|^{\alpha+1} \to 0 \quad \text{if } \alpha \leq 0,
$$

as well as, using again estimates [\(4.4\)](#page-12-4),

$$
\lambda ||U_{\lambda}(x)|^{\alpha}U_{\lambda}(x) - |U_{\lambda}(x_0)|^{\alpha}U_{\lambda}(x_0) || \leq \lambda \frac{K}{\lambda^{\frac{\alpha}{\alpha+1}}} |U_{\lambda}(x) - U_{\lambda}(x_0) || \to 0 \quad \text{ if } \alpha > 0.
$$

It then follows that c does not depend on the choice of x_0 and, up to a sequence and locally uniformly in Ω, one has

$$
\lambda |U_{\lambda}|^{\alpha}U_{\lambda} \to -c \, .
$$

Moreover, the function $V_\lambda(x) = U_\lambda(x) - U_\lambda(x_0)$ is locally uniformly bounded, locally uniformly Lipschitz continuous and satisfies

$$
-|\nabla V_{\lambda}|^{\alpha} F(D^2 V_{\lambda}) + |\nabla V_{\lambda}|^{\beta} = f - \lambda |U_{\lambda}|^{\alpha} U_{\lambda} \text{ in } \Omega.
$$

If V denotes the local uniform limit of V_{λ} for a sequence $\lambda_n \to 0$, then one has

$$
-|\nabla V|^{\alpha} F(D^2 V) + |\nabla V|^{\beta} = f + c \text{ in } \Omega.
$$

Finally, arguing as in the proof of Theorem [1.1](#page-1-2) and using Theorem [2.2,](#page-3-2) we have that, for some $\delta_0 > 0$ sufficiently small,

$$
V_{\lambda} \ge \phi + \min_{d(x) = \delta_0} V_{\lambda} \quad \text{in } \Omega \setminus \Omega_{\delta_0},
$$

with ϕ defined in [\(3.2\)](#page-11-1) for arbitrary $s > 0$. Letting λ , $s \to 0$ we deduce that $V(x) \to +\infty$ as $d(x) \to 0$. This shows that (c, V) is an ergodic pair and concludes the proof.

We now prove that ergodic functions satisfy on the boundary the same asymptotic identities as the explosive solutions of [\(4.3\)](#page-12-1).

Theorem 6.3. Let F and f be as in Theorem [4.1.](#page-12-5) Then, any ergodic function u satisfies

$$
\lim_{d(x)\to 0} \frac{u(x) d(x)^\gamma}{C(x)} = 1 \text{ if } \gamma > 0, \quad \lim_{d(x)\to 0} \frac{u(x)}{|\log d(x)| C(x)} = 1 \text{ if } \gamma = 0. \tag{6.2}
$$

 \Box

Proof. As in the proof of Theorem [4.1,](#page-12-5) we consider only the case $\gamma > 0$. The computations made in the proof of Theorem [4.1](#page-12-5) (for $\gamma_1 = 0$) show that, for all $\nu > 0$ and for $\delta_0 > 0$ sufficiently small, the function $\overline{w}_{\nu,\delta}(x) := \frac{C(x)+\nu}{(d(x)-\delta)^{\gamma}}$ satisfies for $\delta < d(x) < \delta_0$

$$
-|\nabla \overline{w}_{\nu,\delta}|^{\alpha} F(D^2 \overline{w}_{\nu,\delta}) + |\nabla \overline{w}_{\nu,\delta}|^{\beta} \ge c_1 \nu(d(x)-\delta)^{-\frac{\beta}{\beta-\alpha-1}}
$$

where $c_1 > 0$ is a constant depending on $\alpha, \beta, a, A, |D^2d|_{\infty}$ and $|C|_{\mathcal{C}^2(\Omega)}$. By assumption (4.2) on f, this implies that

$$
-|\nabla \overline{w}_{\nu,\delta}|^{\alpha} F(D^2 \overline{w}_{\nu,\delta}) + |\nabla \overline{w}_{\nu,\delta}|^{\beta} > f(x) + c = -|\nabla u|^{\alpha} F(D^2 u) + |\nabla u|^{\beta} \text{ in } \Omega_{\delta} \setminus \Omega_{\delta_0}
$$

for $\delta_0 = \delta_0(\nu)$ small enough. Hence, we are in the hypothesis of Theorem [2.2](#page-3-2) and we deduce that

$$
u \le M_{\nu} + \overline{w}_{\nu,\delta}
$$
 in $\Omega_{\delta} \setminus \Omega_{\delta_0}$,

with $M_{\nu} = \sup_{d(x) = \delta_0} u(x)$. Letting $\delta \to 0$ we obtain that

$$
u \le M_{\nu} + (C(x) + \nu)d(x)^{-\gamma}
$$
 in $\Omega \setminus \Omega_{\delta_0}$.

This in turn implies that

$$
\lim_{d(x)\to 0} u(x)^{\alpha+1} d(x)^{\frac{\beta}{\beta-\alpha-1}-\gamma_0} = 0
$$

for all γ_0 such that

$$
\gamma_0 < \frac{\beta}{\beta - \alpha - 1} - (\alpha + 1)\gamma = \alpha + 2.
$$

Since $\alpha + 2 > 1$, we obtain in particular that the function $|u|^{\alpha}u$ satisfies condition [\(4.2\)](#page-12-2) with $\gamma_0 = 1$. Note also that $|u|^{\alpha}u$ is bounded from below in Ω since it is continuous and blows up on the boundary. Finally, we observe that u satisfies

$$
-|\nabla u|^{\alpha} F(D^2 u) + |\nabla u|^{\beta} + |u|^{\alpha} u = f + c + |u|^{\alpha} u,
$$

where the right hand side $f + c + |u|^\alpha u$ satisfies condition [\(4.2\)](#page-12-2) with an exponent $\gamma_0 = \min{\gamma_0(f), 1}$, $\gamma_0(f)$ being the exponent appearing in the condition [\(4.2\)](#page-12-2) satisfied by f. Hence, by applying Theorem [4.1,](#page-12-5) we obtain that u satisfies the boundary estimates [\(4.4\)](#page-12-4) with $\lambda = 1$ and the constant D depending also on u itself. Estimates [\(4.4\)](#page-12-4) in turn imply relations [\(6.2\)](#page-21-0). \Box

6.2 Uniqueness and further properties of the ergodic constant: proof of Theorem [1.2.](#page-2-0)

Throughout this section we assume that f is bounded and locally Lipschitz continuous. In the introduction we have defined $\mu^* \in \mathbb{R} \cup \{-\infty\}$ as

$$
\mu^* = \inf \{ \mu : \exists \varphi \in C(\overline{\Omega}), -|\nabla \varphi|^\alpha F(D^2 \varphi) + |\nabla \varphi|^\beta \le f + \mu \}.
$$

It is easy to see that $\mu^* \le -\inf_{\Omega} f$. A better upper bound on μ^* depending on the domain Ω is given by the following result.

Proposition 6.4. If $\Omega \subset [0, R] \times \mathbb{R}^{N-1}$, then

$$
\mu^* \le -\inf_{\Omega} f - \frac{K_1}{R^{\frac{\beta}{\beta - \alpha - 1}}}
$$

for a positive constant $K_1 = K_1(a, \alpha, \beta)$.

Proof. The function $\varphi(x) = Cx_1^{\frac{\alpha+2}{\alpha+1}}$ with $C = \left[\frac{a}{2(\alpha+1)}\right]^{\frac{1}{\beta-\alpha-1}} \left(\frac{\alpha+1}{\alpha+2}\right) R^{-\frac{\beta}{(\alpha+1)(\beta-\alpha-1)}}$ satisfies, for some constant $K_1 = K_1(a, \alpha, \beta)$:

$$
-|\nabla\varphi|^{\alpha}F(D^2\varphi)+|\nabla\varphi|^{\beta}\leq -\frac{a(\alpha+2)^{1+\alpha}}{(\alpha+1)^{2+\alpha}}C^{1+\alpha}+\frac{(\alpha+2)^{\beta}}{(\alpha+1)^{\beta}}C^{\beta}x_1^{\frac{\beta}{\alpha+1}}\leq -K_1R^{-\frac{\beta}{\beta-\alpha-1}}.
$$

Hence, by its definition, $\mu^* \le -\inf_{\Omega} f - K_1 R^{-\frac{\beta}{\beta - \alpha - 1}}$.

We are now ready to give the proof of Theorem [1.2.](#page-2-0)

Proof of Theorem [1.2.](#page-2-0) Here we set $c_{\Omega} = c$. The existence of c is given by Theorem [6.2.](#page-20-2)

Proof of [1.](#page-2-1) Suppose that c and c' are two ergodic constants, and let φ and φ' be respectively corresponding ergodic functions. By Theorem [6.3](#page-21-1) the ratio of φ and φ' goes to 1 as $d(x) \to 0$; hence, for any $\theta < 1$, the supremum of $\theta \varphi - \varphi'$ is achieved in the interior of Ω since $\theta \varphi - \varphi'$ blows down to $-\infty$ as $d(x) \to 0$.

 \Box

We observe that

$$
-|\nabla(\theta\varphi)|^{\alpha} F(D^2(\theta\varphi)) + |\theta\nabla\varphi|^{\beta} \leq \theta^{1+\alpha}(f+c)
$$

$$
-|\nabla\varphi'|^{\alpha} F(D^2\varphi') + |\nabla\varphi'|^{\beta} = f+c
$$

From standard comparison arguments in viscosity solutions theory, see [\[17\]](#page-25-12), it follows that at a maximum point \bar{x} of $\theta\varphi - \varphi'$ one has $f(\bar{x}) + c' \leq \theta^{1+\alpha}(f(\bar{x}) + c)$. Letting $\theta \to 1$, we get $c' \leq c$. Exchanging the roles of c and c' we conclude that $c = c'$.

Proof of [2.](#page-2-2) Let $\mu < c$ and suppose by contradiction that there exists $\varphi \in C(\overline{\Omega})$ satisfying

$$
-|\nabla \varphi|^\alpha F(D^2\varphi) + |\nabla \varphi|^\beta \le f + \mu.
$$

Let u be an ergodic function corresponding to c. Clearly, $\sup_{\Omega}(\varphi - u)$ is attained in Ω . Again by standard viscosity arguments, we obtain that at a maximum point \bar{x} of $\varphi - u$ one has

$$
f(\bar{x}) + \mu \ge f(\bar{x}) + c \,,
$$

which is a contradiction. Hence, we deduce

$$
\{\mu : \exists \varphi \in C(\overline{\Omega}), -|\nabla \varphi|^\alpha F(D^2 \varphi) + |\nabla \varphi|^\beta \le f + \mu\} \subset [c, +\infty),
$$

which implies that μ^* is finite and $\mu^* \geq c$. On the other hand, by definition of μ^* , for any $\mu < \mu^*$ the problem

$$
\begin{cases}\n-|\nabla u|^{\alpha}F(D^2u) + |\nabla u|^{\beta} = f + \mu & \text{in } \Omega \\
u = 0 & \text{on } \partial\Omega\n\end{cases}
$$

does not have solution. Theorem [1.1](#page-1-2) then implies that there exists an ergodic constant $c_{f+\mu} \geq 0$ for the right hand side $f + \mu$. On the other hand, by the uniqueness proved in [1.](#page-2-1) above, one has $c = \mu + c_{f+\mu}$. Hence, $c \ge \mu$ and, therefore, $c \ge \mu^*$.

Proof of [3.](#page-2-3) The nondecreasing monotonicity of c with respect to the domain Ω is an immediate consequence of point [2.](#page-2-2) above and the definition of μ^* .

Let us now prove the "continuity" of $\Omega \mapsto c_{\Omega}$, in the following weak sense. For $\delta > 0$ small, let c_{δ} denote the ergodic constant in Ω_{δ} . Then, c_{δ} is nondecreasing as δ decreases to zero, and $c_{\delta} \leq c = c_{\Omega}$. Let u_{δ} be an ergodic function in Ω_{δ} , $x_0 \in \Omega$ be a fixed point and let us set $v_{\delta} = u_{\delta} - u_{\delta}(x_0)$.

By Theorem [2.4,](#page-10-2) v_{δ} is locally uniformly bounded and locally uniformly Lipschitz continuous in Ω_{δ} . Thus, up to a sequence $\delta_n \to 0$, v_δ converges locally uniformly in Ω to a solution v of the equation with right hand side $f + \lim_{\delta \to 0} c_{\delta}$. Moreover, arguing as in the proof of Theorem [6.2,](#page-20-2) we have that $v_\delta(x) \ge C_0 (d(x) - \delta)^{-\gamma}$ if $\gamma > 0$, and $v_\delta(x) \ge -C_0 \log (d(x) - \delta)$ if $\gamma = 0$, for some constant $C_0 > 0$ and for $\delta < d(x) \leq \delta_0$. Letting $\delta \to 0$, we get that $v(x) \to +\infty$ as $d(x) \to 0$. Hence, v is an ergodic function in Ω and, by point [1.](#page-2-1), $\lim_{\delta \to 0} c_{\delta}$ is the ergodic constant c.

Proof of [4.](#page-2-4)

We prove that the constant μ^* is not achieved. Suppose by contradiction that there exists $\varphi \in C(\overline{\Omega})$ such that

$$
-|\nabla \varphi|^\alpha F(D^2\varphi) + |\nabla \varphi|^\beta \le f + \mu^\star = f + c \, .
$$

On the other hand, let u be an ergodic function in Ω .

We observe that for all constants $M, \varphi + M$ is still a bounded sub solution, whereas u is a solution satisfying $u = +\infty$ on $\partial\Omega$. Theorem [5.1](#page-17-1) applied in a smaller domain Ω_{δ} then yields $u \geq \varphi + M$ for arbitrarily large M, which clearly is a contradiction.

A similar argument proves the strict increasing behavior of the ergodic constant. Let $\Omega' \subset\subset \Omega$ and suppose by contradiction that $c_{\Omega'} = c_{\Omega}$. Let $u_{\Omega'}$ and u_{Ω} be ergodic functions respectively in Ω' and Ω . For every constant M, both $u_{\Omega} + M$ and $u_{\Omega'}$ satisfy [\(6.1\)](#page-20-1) in Ω' , with $u_{\Omega} + M$ bounded and $u_{\Omega'} = +\infty$ on $\partial\Omega'$. Hence, Theorem [5.1](#page-17-1) yields the contradiction $u_{\Omega'} \ge u_{\Omega} + M$ for every M.

Remark 6.5. We remark that, thanks to Proposition [6.4,](#page-22-0) the condition sup_{Ω} f + c < 0 appearing in Theorem [1.2](#page-2-0)[-4.](#page-2-4) is satisfied in one of the following cases:

- f is constant in Ω ;
- the oscillation sup_Ω f –inf_Ω f of f is suitably small, in dependence of the length of the projections of Ω on the coordinated axes;
- in at least one direction Ω is suitably narrow, in dependence of the oscillation of f in Ω .

Acknowledgment. Part of this work has been done while the first and third authors were visiting the UMR 80-88, University of Cergy Pontoise, and the second one was visiting Sapienza University of Rome supported by GNAMPA- INDAM .

References

- [1] S. Alarc´on, A. Quaas, Large viscosity solutions for some fully nonlinear equations, Nonlinear Differ. Equ. Appl. 20 (2013), 1453–1472
- [2] G. Barles, J. Busca, Existence and comparison results for fully non linear degenerate elliptic equations without zeroth order terms, Communications in Partial Differential Equations , 26(11- 12), (2001), 2323–2337.
- [3] G. Barles, F. Murat, Uniqueness and the maximum principle for quasilinear elliptic equations with quadratic growth conditions. Archive for Rational mechanics and analysis 133.1 (1995), 77–101.
- [4] G. Barles, A. Porretta, Uniqueness for unbounded solutions to stationary viscous Hamilton-Jacobi $equation$ Ann. Scuola Norm. Sup Pisa, Cl Sci (5) , Bol V (2006) , 107-136.
- [5] G. Barles, A. Porretta, T. Tabet Tchamba, On the large time behavior of solutions of the Dirichlet problem for subquadratic viscous Hamilton-Jacobi equations Journal de Mathématiques Pures et Appliquées, 94, (2010), 497–519.
- [6] I. Birindelli, F. Demengel First eigenvalue and Maximum principle for fully nonlinear singular operators, Advances in Differential equations, Vol (2006) 11 (1), 91–119.
- [7] I. Birindelli, F. Demengel, $C^{1,\beta}$ regularity for Dirichlet problems associated to fully nonlinear degenerate elliptic equations, ESAIM COCV, (2014), 20 (40), 1009–1024.
- [8] I. Birindelli, F. Demengel, F. Leoni, Existence, comparison and regularity results for a class of fully non linear singular or degenerate equations, in progress.
- [9] I. Birindelli, F. Demengel, F. Leoni, On the $W^{2,p}$ and $\mathcal{C}^{1,\gamma}$ regularity for fully non linear singular or degenerate equations, in preparation.
- [10] J. Busca, M. Esteban, A. Quaas, Nonlinear eigenvalues and bifurcation problems for Pucci's operators, Ann. Inst. H. Poincaré Anal. Non Linéaire, 22 (2005), no. 2, 187–206.
- [11] I. Capuzzo Dolcetta, F. Leoni, A. Porretta, Hölder's estimates for degenerate elliptic equations with coercive Hamiltonian, Transactions of the American Mathematical Society, Vol 362, n.9, (2010), 4511–4536.
- [12] I. Capuzzo Dolcetta, F. Leoni, A. Vitolo, On the inequality $F(x, D^2u) \ge f(u) + g(u)|Du|^q$, Math. Ann., Vol. 365, n.1-2 (2016), 423–448.
- [13] I. Capuzzo Dolcetta, F. Leoni, A.Vitolo, Entire subsolutions of fully nonlinear degenerate elliptic equations, Bull. Inst. Math. Acad. Sinica (New Series) 9(2) (2014), 147–161.
- [14] F. Demengel, O. Goubet, Existence of boundary blow up solutions for singular or degenerate fully nonlinear equations, Comm. Pure and Applied Anal., 12 (2) (2013), 621–645.
- [15] M. Esteban, P. Felmer, A. Quaas, Super-linear elliptic equation for fully nonlinear operators without growth restrictions for the data, Proc. Roy. Soc. Edinburgh, 53, (2010), 125–141
- [16] H. Ishii, Viscosity solutions of Nonlinear fully nonlinear equations Sugaku Expositions , Vol 9, number 2, December 1996.
- [17] M.G. Crandall,H. Ishii, P.L. Lions, User's guide to viscosity solutions of second order partial differential equations, Bull. Amer. Math. Soc. (N.S.) 27 (1992), no. 1, 1–67.
- [18] J.M. Lasry, P. L. Lions, Nonlinear Elliptic Equations with Singular Boundary Conditions and Stochastic Control with state Constraints, Math. Ann. 283, (1989), 583–630.
- [19] T. Leonori, A. Porretta, Large solutions and gradient bounds for quasilinear elliptic equations, Comm. in Partial Differential Equations 41, 6 (2016), 952–998.
- [20] T. Leonori, A. Porretta, G. Riey, Comparison principles for p-Laplace equations with lower order terms, Annali di Matematica Pura ed Applicata 196(3), (2017), 877–903.
- [21] A. Porretta, The ergodic limit for a viscous Hamilton–Jacobi equation with Dirichlet conditions, Rend. Lincei Mat. Appl. 21 (2010), 59–78.