

Minimax Estimation of Large Precision Matrices with Bandable Cholesky Factor

Yu Liu and Zhao Ren
University of Pittsburgh¹

Abstract

This paper considers the estimation of large precision matrices. We focus on two commonly used types of parameter spaces with a bandable structure on the Cholesky factor of precision matrices. Last decade witnesses significant methodological advances and wide scientific applications in meteorology and spectroscopy. However, the minimax theory has still been largely unknown, as opposed to the well established minimax results over the corresponding two types of bandable covariance matrices. In this paper, we develop the optimal rates of convergence under both the operator norm and the Frobenius norm. A striking phenomenon is found: two types of parameter spaces are fundamentally different under the operator norm but enjoy the same rate optimality under the Frobenius norm, which is in sharp contrast to the equivalence of two types of bandable covariance matrices under both norms. This fundamental difference is largely established by carefully constructing the corresponding minimax lower bounds using the Le Cam and Assouad methods. Two new estimation procedures are developed: for the operator norm, our optimal procedure is based on a novel local cropping estimator targeting on all principle submatrices of the precision matrix while for the Frobenius norm, our optimal procedure relies on a delicate regression-based block-thresholding rule. We further establish rate optimality in the nonparanormal model by applying our local cropping procedure to the rank-based estimators. In the end, we carry out numerical studies to demonstrate the performance of our approach as well as illustrate the fundamental difference between two types of parameter spaces.

Keywords: optimal rate of convergence, precision matrix, local cropping, tapering, Cholesky factor, minimax lower bound, operator norm, Frobenius norm, block-thresholding
AMS 2000 Subject Classification: Primary 62H12; secondary 62F12, 62C20, 62G09.

¹Department of Statistics, University of Pittsburgh, Pittsburgh, PA 15260. e-mail: yul125@pitt.edu, zren@pitt.edu

1 Introduction

Covariance matrix plays a fundamental role in many important multivariate statistical problems. They include the principal component analysis, linear and quadratic discriminant analysis, clustering analysis, regression analysis and conditional dependence relationship studies in graphical models. During the last two decades, with the advances of technology, it is very common that the datasets are high-dimensional (the dimension p can be much larger than the sample size n) in many applications such as genomics, fMRI data, astrophysics, spectroscopic imaging, risk management, portfolio allocation and numerical weather forecasting [26, 20, 25, 33, 45, 39]. It has been well-known that the sample covariance matrix performs poorly and can yield to invalid conclusions in the high-dimensional settings. For example, see [48, 49, 29, 21, 40, 30] for details on the limiting behaviors of the spectra of sample covariance matrices when both n and p increase.

To avoid the curse of dimensionality, structural assumptions are imposed in literature in order to estimate the covariance matrix or its inverse, the precision matrix, consistently under various loss functions. While mostly driven by the specific scientific applications, popular structures include *ordered sparsity* (bandable covariance matrices, precision matrices with bandable Cholesky factor), *unordered sparsity* (sparse covariance matrices, sparse precision matrices) and other more complicated ones such as certain combination of sparsity and low-rankness (spike covariance matrices, covariance with tensor product, latent graphical models). Many estimation procedures have been proposed accordingly to estimate high-dimensional covariance/precision matrices via taking advantages of these specific structures. For example, banding ([51, 7, 52, 8]) and tapering methods ([24, 13]) were developed to estimate bandable covariance matrices or precision matrices with bandable Cholesky factor; thresholding procedures were used in [6, 22, 15] to estimate sparse covariance matrices; penalized likelihood estimation ([28, 55, 18, 2, 43, 32, 41]) and penalized regression methods ([37, 54, 16, 47, 42]) are designed for sparse precision matrix estimation. In addition, loss functions critically determine the intrinsic estimation difficulty and corresponding efficient estimation procedures. For matrix estimation, important loss functions include operator norm loss ($\|S\|_{\text{op}} = \sup_{\|x\|_2=1} \|Sx\|_2$ for a matrix $S \equiv [s_{ij}]_{p \times q} \in \mathbb{R}^{p \times q}$), Frobenius norm loss ($\|S\|_{\text{F}} = (\sum_{i,j} s_{ij}^2)^{\frac{1}{2}}$) and its equivalent forms such as Bregman divergence loss. Among all losses, the operator loss is arguably the most important one which is regarded as a truly “two-directional” problem because it cannot be essentially reduced to a problem of estimating a single or multiple vectors.

During the last decade, fundamental difficulty of various covariance/precision matrices estimation problems have been carefully investigated in terms of the minimax risks under the operator norm loss among other losses, especially for those *ordered and unordered sparsity structures*. Specifically, for unordered structures, [14] considered the problems of optimal estimation of sparse covariance while [10] (see [42] as well) established the optimality results for estimating sparse precision matrices. For ordered structures, [13] established the optimal rates of convergence over two types of bandable covariance ma-

trices. In addition, with an extra Toeplitz structure, [11] studied optimal estimation of two types of bandable Toeplitz covariance matrices. However, it is still largely unknown about the optimality results on estimating precision matrices with bandable Cholesky factor. See an exposure paper with discussion [12] and references therein on minimax results of covariance/precision matrix estimation under some other losses.

In this paper, we provide a solution to this open problem by establishing the optimal rates of convergence over two types of precision matrices with bandable Cholesky factor. Thus, this paper completes the minimaxity results of all four sparsity structures commonly considered in literature. Both the operator norm and Frobenius norm are considered. We begin with introducing the bandable Cholesky factor of the precision matrix. Assume that $\mathbf{X} = (X_1, \dots, X_p)^T$ is a centered p -variate random vector with covariance matrix Σ . Let $\hat{X}_i = \sum_{t=1}^{i-1} a_{it} X_t = \mathbf{X}_{1,i-1}^T \mathbf{a}_i$ be the regression of X_i on its previous variables $\mathbf{X}_{1,i-1} = (X_1, X_2, \dots, X_{i-1})^T$ in population. (Define $\hat{X}_1 = 0$.) Set A as the lower triangular matrix with zeros on the diagonal and coefficients \mathbf{a}_i^T arranged in the rows. Denote the residual $\epsilon = \mathbf{X} - \hat{\mathbf{X}} = (I - A)\mathbf{X}$ and $D = \text{Var}(\epsilon)$. The regression theory implies the residuals are uncorrelated, and thus the matrix D is diagonal. The modified Cholesky decomposition of Ω is

$$\Omega = \Sigma^{-1} = (I - A)^T D^{-1} (I - A), \quad (1)$$

where $I - A$ is the Cholesky factor of Ω . There is a natural order on the variables based on the above Cholesky decomposition. Indeed, the well-known AR(k) model can be characterized by the k -banded Cholesky factor $A \equiv [a_{ij}]_{p \times p}$ of the precision matrix in which $a_{ij} = 0$ if $i - j > k$. Inspired by the auto-regression model, we consider the bandable structures imposed on the Cholesky factor. More specifically, for $M > 0$, $\eta > 1$ we define the parameter space $\mathcal{P}_\alpha(\eta, M)$ of precision matrices by

$$\mathcal{P}_\alpha(\eta, M) = \left\{ \Omega : \eta^{-1} \leq \lambda_{\min}(\Omega) \leq \lambda_{\max}(\Omega) < \eta, \right. \\ \left. \max_i \sum_{j < i-k} |a_{ij}| < M k^{-\alpha}, \quad k \in [i-1] \right\}. \quad (2)$$

Here, $\lambda_{\max}(\Omega)$, $\lambda_{\min}(\Omega)$ are the maximum and minimum eigenvalues of Ω and the index set $[p] = \{1, 2, \dots, p\}$. This parameter space was first proposed in [7]. The parameter α specifies how fast the sequence a_{ij} decays to zero as j goes away from i . Unlike the order in the bandable covariance matrices [6, 13], in which large distance $|i - j|$ implies nearly independence, the order in bandable Cholesky factor encodes nearly conditional independence in the sense that the coefficients a_{ij} is close to zero when $i - j > 0$ is large.

Although several approaches have been developed to estimate the precision matrix with bandable Cholesky factor, the optimality question remains mostly open, partially due to the following two reasons. (i) Intuitively, one would expect the minimax rate of convergence over $\mathcal{P}_\alpha(\eta, M)$ under the operator norm to be the same as that over the class of bandable covariance matrices with the same decay parameter α . Under sub-Gaussian assumptions, [13] established the optimal rate of convergence $\mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \asymp n^{\frac{-2\alpha+1}{2\alpha}} + \frac{\log p}{n}$

uniformly for all bandable covariance matrices $\Sigma = \Omega^{-1} = [\sigma_{ij}]_{p \times p}$ with bounded spectra such that $\max_i \sum_{|j-i|>k} |\sigma_{ij}| < Mk^{-\alpha}$, $k \in [p]$. To establish such a rate of convergence for $\mathcal{P}_\alpha(\eta, M)$, [34] provided a lower bound with the matching rate. However, we show a surprising result in this paper that estimation over $\mathcal{P}_\alpha(\eta, M)$ is a much harder task than that over bandable covariance matrices. Therefore, the lower bound in [34] is sub-optimal, and all attempts on showing the same rate of convergence $n^{\frac{-2\alpha+1}{2\alpha}} + \frac{\log p}{n}$ intrinsically cannot succeed. (ii) From the methodological aspect, due to the regression interpretation of the Cholesky decomposition (1), almost all existing methods rely on an intermediate estimator of A obtained by running regularized regression of each variable against its previous variables $X_i \sim \sum_{j=1}^{i-1} a_{ij}X_j$. For instance, [7] estimated each row of A by fitting the banded regression model $X_i \sim \sum_{j=\max\{1, i-k\}}^{i-1} a_{ij}X_j$ with some bandwidth k . [50] used an AIC or BIC penalty to pick the best bandwidth k . In addition, [28] proposed adding a Lasso or Ridge penalty while [35] proposed using a nested Lasso penalty to the regression. See, for instance, [3, 34] for Bayesian approaches following the similar idea. While the typical analysis for those estimation procedures in a row-wise fashion is to bound the operator norm by its matrix ℓ_1/ℓ_∞ norm, which may provide optimal rates of convergence under the operator norm loss for unordered sparsity structure (see, i.e., [14, 10] for sparse covariance and precision matrices estimation), it is well-received that such an analysis would be sub-optimal for ordered sparsity structure as seen in bandable covariance matrix estimation [13, 7]. Therefore, in order to obtain rate-optimality over $\mathcal{P}_\alpha(\eta, M)$, a novel analysis or even a new estimation approach is expected.

In regard to the above two issues, we provide satisfactory solutions in this paper. We at the first time show that the rate of convergence under the operator norm over $\mathcal{P}_\alpha(\eta, M)$ is intrinsically slower than that over the counterpart class of bandable covariance matrices. This is achieved via a novel minimax lower bound construction. Moreover, in order to obtain a rate-optimal estimator, we propose a novel local cropping estimator which does not rely on any estimator of A , and thus requires a new analysis. Our local cropping approach targets on accurate estimation of principal submatrices of the precision matrix under the operator norm, which results in a tradeoff between one variance term and two bias terms. The name comes after the idea of estimating each principal submatrix of the precision matrix, which is to crop the center k by k submatrix of the inverse of $3k$ by $3k$ sample covariance matrix using their neighbors in two directions of the same size. (During the finalizing process of this paper, we realized that a similar estimator is independently proposed to estimate precision matrices with a different structure [27].) Since our procedure does not directly explore the structure on each row of A , the analysis of bias terms is much involved, requiring a block-wise partition strategy. More details are discussed in Sections 2.1 and 3.1. In the end, besides $\mathcal{P}_\alpha(\eta, M)$, a similar type of classes of parameter spaces with bandable Cholesky factor is considered as well,

$$\mathcal{Q}_\alpha(\eta, M) = \left\{ \Omega : \eta^{-1} \leq \lambda_{\min}(\Omega) \leq \lambda_{\max}(\Omega) < \eta, \right. \\ \left. |a_{ij}| < M(i-j)^{-\alpha-1}, \quad j \in [i-1] \right\}. \quad (3)$$

We further establish another surprising result: the optimal rates of convergence of two spaces, namely $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$, are different under the operator norm. This remarkable distinction is significantly different from the comparison of two similar types of parameter spaces for bandable covariance matrices in [13] and bandable Toeplitz covariance matrices in [11]. The contrast of minimax results on $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$ is summarized in Theorem 1 below. We mainly focus on the high-dimensional setting, assuming that $\log p = O(n)$ and $n = O(p)$. Otherwise, one can always obtain trivial constant minimax rate (i.e., inconsistency) or the minimax rate as the smaller of p/n and the one in Theorem 1.

Theorem 1 *Under normality assumption, the minimax risk of estimating the precision matrix Ω over the parameter space $\mathcal{P}_\alpha(\eta, M)$ with $\alpha > \frac{1}{2}$ given in (2) under the operator norm satisfies*

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{P}_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \asymp n^{\frac{-2\alpha+1}{2\alpha}} + \frac{\log p}{n}. \quad (4)$$

The minimax risk of estimating the precision matrix Ω over the parameter space $\mathcal{Q}_\alpha(\eta, M)$ given in (3) under the operator norm satisfies

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{Q}_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \asymp n^{\frac{-2\alpha}{2\alpha+1}} + \frac{\log p}{n}. \quad (5)$$

Moreover, we also consider the minimax rates of convergence of precision matrix estimation under the Frobenius norm loss over $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$. This time, we prove that two types of spaces enjoy the same optimal rate of convergence. Together with the different rates of convergence under the operator norm loss, we demonstrate the intrinsic difference between operator norm and Frobenius norm. The Frobenius norm of a p by p matrix is defined as the ℓ_2 vector norm of all entries. Driven by this fact, our estimation approach is naturally obtained by optimally estimating A and D in (1) separately. Due to the decay structure in $\mathcal{P}_\alpha(\eta, M)$, which is defined in terms of nested ℓ_1 norm of each row of A , our estimator is based on regression with a delicate block-thresholding rule. The minimax procedure is motivated by wavelet nonparametric function estimation, although the space $\mathcal{P}_\alpha(\eta, M)$ cannot be exactly described by any Besov ball ([9, 19]). We summarize the optimality result under the Frobenius norm in Theorem 2 below.

Theorem 2 *Under normality assumption, the minimax risk of estimating the precision matrix Ω over $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$ given in (2) and (3) satisfies*

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{P}_\alpha(\eta, M)} \frac{1}{p} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{F}}^2 \asymp \inf_{\tilde{\Omega}} \sup_{\mathcal{Q}_\alpha(\eta, M)} \frac{1}{p} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{F}}^2 \asymp n^{-\frac{2\alpha+1}{2\alpha+2}}. \quad (6)$$

The rest of the paper is organized as follows. First, we propose our estimation procedures for precision matrix estimation in Section 2. In particular, local cropping estimators and regression-based block-thresholding estimators are designed for estimating precision

matrices under the operator norm and the Frobenius norm respectively. Section 3 establishes the optimal rates of convergence under the operator norm for two commonly used types of parameter spaces $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$. The upper bounds are obtained by studying the bias-variance tradeoff of the local cropping estimators. The minimax lower bounds are obtained by testing arguments, which reveal a fundamental and striking difference between two spaces when considering operator norm loss. Section 4 considers rate-optimal estimation under the Frobenius norm via carefully studying the properties of regression-based block-thresholding estimators. The results reveal that the fundamental difficulty of estimation for two parameter spaces are the same when considering Frobenius norm loss. In Section 5, we extend the results to nonparanormal models for inverse correlation matrix estimation by applying local cropping procedure to rank-based estimators. Section 6 presents the numerical performance of our local cropping procedure to illustrate the difference between two parameter spaces by simulation studies. We also demonstrate the sub-optimality of banding estimators, compared to our optimal procedures. All technical lemmas used in proofs of main results are relegated to the supplement.

Notation. we introduce some basic notations that will be used in the rest of the paper. $\mathbf{1}(\cdot)$ indicates the indicator function while $\mathbf{1}$ indicates the all-ones vector. $\text{sgn}(\cdot)$ indicates the sign function. $\lfloor s \rfloor$ represents the largest integer which is no more than s . $\lceil s \rceil$ represents the smallest integer which is no less than s . Define $a_n \asymp b_n$ if there is a constant $C > 0$ independent of n such that $C^{-1} \leq a_n/b_n \leq C$. For any vector x , $\|x\|_p$ indicates its ℓ_p norm. For any p by q matrix $S = [s_{ij}]_{p \times q} \in \mathbb{R}^{p \times q}$, we use S^T to denote its transpose. The ℓ_p matrix norm is define as $\|S\|_p = \sup_{\|x\|_p=1} \|Sx\|_p$. The ℓ_2 matrix norm is also called the operator norm or the spectral norm. The Frobenius norm is defined as $\|S\|_F = (\sum_{i,j} s_{ij}^2)^{\frac{1}{2}}$. $\lambda_{\max}(S)$ and $\lambda_{\min}(S)$ are the largest and smallest singular value of S . When S is a real symmetric matrix, $\lambda_{\max}(S)$ and $\lambda_{\min}(S)$ also denote the largest and smallest magnitudes of its eigenvalue values. $\text{row}_i(S)$ and $\text{col}_i(S)$ indicate the i -th row and column of matrix S . $a : b$ denotes the index set $\{a, a+1, \dots, b\}$. $[p]$ is short for the set $1 : p$. For the random vector $\mathbf{X} \in \mathbb{R}^{p \times 1}$ and the data matrix $\mathbf{Z} \in \mathbb{R}^{n \times p}$, $\mathbf{X}_{a:b}$ and $\mathbf{Z}_{a:b}$ indicates the $(a : b)$ -th columns of \mathbf{X}^T and \mathbf{Z} . For any p by p matrix S , $\text{diag}(S)$ denotes the diagonal matrix with diagonal entries being those on the main diagonal of S while for any vector v , $\text{diag}(v)$ denotes the diagonal matrix with diagonal entries being v . In the estimation procedure under the operator norm, we use the matrix notation in the form of $S_m^{(k)}$ to facilitate the proof, where S is always a square matrix, m indicates the location information, and (k) indicates that the size of $S_m^{(k)}$ is k . Throughout the paper we denote by C a generic positive constant which may vary from place to place but only depends on α , η , M and possibly some sub-Gaussian distribution constant ρ in (17).

2 Methodologies

In this section, we introduce our methodologies for estimating precision matrices over $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$ under both the operator norm and the Frobenius norm. Assume

that $\mathbf{X} = (X_1, \dots, X_p)^T$, a p -variate random vector with mean zero and precision matrix Ω_p . Our estimation procedures are based on its n i.i.d. copies $\mathbf{Z} \in \mathbb{R}^{n \times p}$. We write $\mathbf{Z} = (\mathbf{Z}_1, \dots, \mathbf{Z}_p)$, where each \mathbf{Z}_i consists of n i.i.d. copies of X_i . Our estimation procedures are different under the operator norm and the Frobenius norm.

2.1 Estimation Procedure under the Operator Norm

We focus on the estimation problem under the operator norm first. As we discussed in the introduction, almost all existing methodologies ([50, 28, 7]) directly appeal the Cholesky decomposition of the precision matrix. They first estimate the Cholesky factor A and D by auto-regression and then estimate the precision matrix according to $\Omega = (I - A)^T D^{-1} (I - A)$. The corresponding analysis in the row-wise fashion may not be suitable for the operator norm loss. In this paper, we propose a novel local cropping estimator, which focuses on the estimation of Ω directly.

To facilitate the illustration of the estimation procedure, we define two matrix operators. The **cropping operator** is designed to crop the center block out of the matrix. For a p by p matrix $E \equiv [e_{ij}]_{p \times p}$, we define the $k \times k$ matrix $\mathbf{C}_m^k(E) \equiv [c_{ij}]_{k \times k}$, where $1 \leq m \leq p - k + 1$, with

$$c_{ij} = e_{i+m-1, j+m-1}, \text{ when } 1 \leq i, j \leq k. \quad (7)$$

The parameter m indicates the location and k indicates the dimension. It is clear that $\mathbf{C}_m^k(E)$ is a principal submatrix of E . The **expanding operator** is designed to put a small matrix onto a large zero matrix. For a k by k matrix, $C \equiv [c_{ij}]_{k \times k}$, define the $p \times p$ matrix $\mathbf{E}_m^p(C) \equiv [e_{ij}]_{p \times p}$, where $1 \leq m \leq p - k + 1$, with

$$e_{ij} = c_{i-m+1, j-m+1}, \text{ when } m \leq i, j \leq m + k - 1, \text{ otherwise } e_{ij} = 0. \quad (8)$$

The parameter m indicates the location and p indicates the dimension. Note that for a k by k matrix C , we have $\mathbf{C}_m^k(\mathbf{E}_m^p(C)) = C$.

In addition, for technical reasons (of obtaining rates of convergence in expectation rather than in probability), we introduce a **projection operator**. For a real matrix S , let the singular value decomposition of S be $S = U\Lambda V^T$ with $UU^T = I$, $VV^T = I$ and $\Lambda = \text{diag}(\lambda_i)$. Let $\Lambda^* = \text{diag}(\lambda_i^*)$, where $\lambda_i^* = \min\{\max\{\lambda_i, \eta^{-1}\}, \eta\}$, then define

$$\mathbf{P}_\eta(S) = U\Lambda^*V^T. \quad (9)$$

For a symmetric matrix S , we modify $\mathbf{P}_\eta(\cdot)$ a little bit and define $\mathbf{P}_\eta(S) = U\Lambda^*U^T$, where $S = U\Lambda U^T$ is its eigen-decomposition. Since all eigenvalues of $\mathbf{P}_\eta(\cdot)$ are in the interval $[\eta^{-1}, \eta]$, $\mathbf{P}_\eta(S)$ is always invertible and positive definite.

We are ready to construct the local cropping estimator $\tilde{\Omega}_k^{\text{op}}$ with bandwidth $k < p$. At a high level, we first propose an estimator of each principal submatrix of size k and $2k$ in Ω using cropping and extending operators. Then we arrange over those local estimators

to estimate Ω . Since the core idea of estimating those local estimators in our procedure is to crop the inverse of sample covariance matrix with a relatively larger size, we call $\tilde{\Omega}_k^{\text{op}}$ in (12) **the local cropping estimator**.

Specifically, we first define an estimator $\tilde{\Omega}_m^{(k)}$ of the principal submatrix $\mathbf{C}_m^k(\Omega)$ at each location m . To this end, we select the sample covariance matrix with a relative larger size, in this case, $3k$. Let the modified local sample covariance matrix be

$$\tilde{\Sigma}_{m-k}^{(3k)} = \mathbf{P}_\eta(\mathbf{C}_{m-k}^{3k}(\frac{1}{n}\mathbf{Z}^T\mathbf{Z})). \quad (10)$$

Note that the operator $\mathbf{P}_\eta(\cdot)$ guarantees $\tilde{\Sigma}_{m-k}^{(3k)}$ to be invertible. Then we use the center part of its inverse to estimate $\mathbf{C}_m^k(\Omega)$, i.e.,

$$\tilde{\Omega}_m^{(k)} = \mathbf{C}_{k+1}^k((\tilde{\Sigma}_{m-k}^{(3k)})^{-1}). \quad (11)$$

Similarly, we can define local estimators of $\tilde{\Omega}_m^{(2k)}$ via replacing k by $2k$. Arranging over these estimators in the form of weighted sum, we obtain the estimator of Ω , that is,

$$\tilde{\Omega}_k^{\text{op}} = \mathbf{P}_\eta\left(\frac{1}{k}\left(\sum_{m=2-2k}^p \mathbf{E}_m^p(\tilde{\Omega}_m^{(2k)}) - \sum_{m=2-k}^p \mathbf{E}_m^p(\tilde{\Omega}_m^{(k)})\right)\right). \quad (12)$$

The operator $\mathbf{E}_m^p(\cdot)$ makes these local estimators in the correct places. The final step (12) is motivated by the analysis of optimal bandable covariance matrix estimation procedure proposed in [13]. Indeed, the optimal tapering estimator in [13] can be rewritten as a sum of many principal submatrices of the sample covariance matrix in a similar way as (12). In contrast, our estimator is not in a form of tapering the sample covariance matrix. However, in the analysis of our local cropping estimator in Section 3, the direct target of $\tilde{\Omega}_k^{\text{op}}$ is certain tapered population precision matrix with bandwidth k . There are natural bias and variance terms involved in the distance of $\tilde{\Omega}_k^{\text{op}}$ and its direct target. Together with the bias of the tapered population precision matrix, our analysis involves two bias terms and one variance term, which critically determine the optimal choice of bandwidth.

In Section 3, we show that the local cropping estimator with an optimal choice of bandwidth would achieve the minimax risk under the operator norm over parameter spaces $\mathcal{P}_\alpha(\eta, M)$ in (2) and $\mathcal{Q}_\alpha(\eta, M)$ in (3). However, the optimal choices of bandwidth are fundamentally distinct between $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$. Specifically, we show that the optimal bandwidth over $\mathcal{P}_\alpha(\eta, M)$ is $k \asymp n^{\frac{1}{2\alpha}}$ while that one over $\mathcal{Q}_\alpha(\eta, M)$ is $k \asymp n^{\frac{1}{2\alpha+1}}$.

Remark 1 *Of note, the estimator $\tilde{\Omega}_k^{\text{op}}$ depends on $\mathbf{Z}_{2-4k}, \dots, \mathbf{Z}_{p+4k-1}$. The index of variable is clear most of the time, while we need to be careful when it is close to the boundary. When the index is beyond the index set $[p]$, we shrink the size of the corresponding block by discarding the data with meaningless indexes.*

2.2 Estimation Procedure under the Frobenius Norm

Under the Frobenius norm, our estimation procedure is based on the Cholesky decomposition of the precision matrix (1). More specifically, we estimate the matrix A and D respectively by auto-regression, and then combine them together to construct the estimator of Ω . The following estimation procedure applies to both the parameter space $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$ as we will show that they enjoy the same optimal rate of convergence in Section 4.

Our estimator of the i -th row of A is based on the regression of X_i against its previous variables. Unlike those existing methods ([50, 28, 7]) which rely on certain banding or penalized approaches for such a regression problem, we apply a block-thresholding procedure due to the decay structure in $\mathcal{P}_\alpha(\eta, M)$ which is defined in terms of nested ℓ_1 norm. To this end, we first regress X_i against $\mathbf{X}_{i-k_1:i-1} = (X_{i-k_1}, \dots, X_{i-1})^T$ with a slightly larger bandwidth $k_1 = \lceil n^{\frac{2\alpha+1}{(2\alpha+2)2\alpha}} \rceil$. Recall that the $n \times 1$ matrix \mathbf{Z}_i consists of n observations of X_i , and the $n \times k_1$ matrix $\mathbf{Z}_{i-k_1:i-1}$ represents n observations of $\mathbf{X}_{i-k_1:i-1}$. The empirical regression coefficients are

$$\hat{\mathbf{a}}_i = (\hat{a}_{i(i-k_1)}, \dots, \hat{a}_{i(i-1)})^T = (\mathbf{Z}_{i-k_1:i-1}^T \mathbf{Z}_{i-k_1:i-1})^{-1} \mathbf{Z}_{i-k_1:i-1}^T \mathbf{Z}_i. \quad (13)$$

We then further threshold the coefficients of $\hat{\mathbf{a}}_i$ by taking advantages of the bandable structure of the Cholesky factor A . Specifically,

$$\hat{a}_{i(i-j)}^* = \hat{a}_{i(i-j)} \mathbf{1}(|\hat{a}_{i(i-j)}| > \lambda_j), j \in [k_1], \quad (14)$$

where $\lambda_j = (\lceil \log_2^j - \log_2^k \rceil R)^{\frac{1}{2}}$, $k = n^{\frac{1}{2\alpha+2}}$ and $R = \eta \|(\mathbf{Z}_{i-k_1:i-1}^T \mathbf{Z}_{i-k_1:i-1})^{-1}\|_{\text{op}}$. Note that we apply a block-thresholding rule in which the size of each block doubles sequentially in (14). Our procedure is inspired by the optimal estimation procedure over Besov balls for many nonparametric function estimation problems, or equivalently, the corresponding Gaussian sequence models (See [9] the reference therein). We emphasize that any linear estimator of the coefficients $\hat{\mathbf{a}}_i$ cannot yield to the optimal rates of convergence in our setting under the Frobenius norm.

Our estimator of $I - A$ can be constructed by arranging $\hat{\mathbf{a}}_i^*$, $i \in [p]$ accordingly with an identity matrix. Set the ij -th entry of \hat{A}^* as \hat{a}_{ij}^* when $i \in [p]$, $(i-j) \in [k_1]$, otherwise as zero. We also need to bound the singular values of $(I - \hat{A}^*)$. To this end, we define

$$\widetilde{I - A} = \mathbf{P}_\eta(I - \hat{A}^*),$$

as our estimator of $(I - A)$, where $\mathbf{P}_\eta(\cdot)$ is defined in Equation (9).

The estimation of D is based on the sample variances of those empirical residuals in the regression between X_i and $\mathbf{X}_{i-k_1:i-1} = (X_{i-k_1}, \dots, X_{i-1})^T$. For each i , the sample variance of the empirical residual is

$$\hat{d}_i = \frac{1}{n - k_1} \mathbf{Z}_i^T (I - \mathbf{M}_i)^T (I - \mathbf{M}_i) \mathbf{Z}_i, \quad (15)$$

where $\mathbf{M}_i = \mathbf{Z}_{i-k_1:i-1}(\mathbf{Z}_{i-k_1:i-1}^T \mathbf{Z}_{i-k_1:i-1})^{-1} \mathbf{Z}_{i-k_1:i-1}^T$. Let $\hat{D} = \text{diag}(\hat{d}_1, \hat{d}_2, \dots, \hat{d}_p)$. We define $\tilde{D} = \mathbf{P}_\eta(\hat{D})$ as our estimator of D .

Finally, define our estimator of Ω as

$$\tilde{\Omega}_k^F = (\widetilde{I - A})^T \tilde{D}^{-1} (\widetilde{I - A}). \quad (16)$$

Remark 2 For the parameter space $\mathcal{Q}_\alpha(\eta, M)$, a much simpler banding estimation scheme on the empirical regression coefficients is able to achieve the minimax risk. Set $k = \lceil n^{\frac{1}{2\alpha+2}} \rceil$. We use the empirical residuals and coefficients obtained by regressing each X_i against $\mathbf{X}_{i-k:i-1}$ to directly construct the estimators of A and D . It can be proved that this estimator achieves the minimax risk over the parameter space $\mathcal{Q}_\alpha(\eta, M)$.

3 Rate Optimality under the Operator Norm

In this section, we establish the optimal rates of convergence for estimating the precision matrix over the parameter spaces $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$ given in (2) and (3) under the operator norm. We first derive the risk upper bound of the local cropping estimator in Section 3.1 over parameter space $\mathcal{P}_\alpha(\eta, M)$. Then we provide a matching risk lower bound by applying the Assouad's method and the Le Cam's method in Section 3.2 over $\mathcal{P}_\alpha(\eta, M)$. The establishment of the rate optimality over the parameter space $\mathcal{Q}_\alpha(\eta, M)$ is similar to the one over $\mathcal{P}_\alpha(\eta, M)$, which is provided in Section 3.3.

Recall that for $1 \leq q < \infty$, the matrix ℓ_q norm of a matrix S is defined by $\|S\|_q = \max_{\|x\|_q=1} \|Sx\|_q$. It is well known that $\frac{\|S\|_F}{\sqrt{p}} \leq \|S\|_{\text{op}} \leq \sqrt{\|S\|_1 \|S\|_\infty}$. In addition, for a real symmetric matrix S , $\|S\|_{\text{op}} \leq \|S\|_1$. In this section, we assume that $\mathbf{X} = (X_1, \dots, X_p)^T$ follows certain sub-Gaussian distribution with constant $\rho > 0$, that is,

$$\mathbb{P}\{|v^T(\mathbf{X} - \mathbb{E}\mathbf{X})| > t\} \leq 2 \exp(-t^2/(2\rho)), \quad (17)$$

for all $t > 0$ and all unit vectors $\|v\|_2 = 1$.

3.1 Minimax Upper Bound under the Operator Norm over $\mathcal{P}_\alpha(\eta, M)$

In this section, we develop the following upper bound of our estimation procedure proposed in Section 2.1.

Theorem 3 When $\lceil n^{\frac{1}{2\alpha}} \rceil \leq p$, the local cropping estimator defined in (12) of the precision matrix Ω over $\mathcal{P}_\alpha(\eta, M)$ with $\alpha > \frac{1}{2}$ given in (2) satisfies

$$\sup_{\mathcal{P}_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega}_k - \Omega\|_{\text{op}}^2 \leq C k^{-2\alpha+1} + C \frac{\log p + k}{n}.$$

When $k = \lceil n^{\frac{1}{2\alpha}} \rceil$, we have

$$\sup_{\mathcal{P}_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega}_k - \Omega\|_{\text{op}}^2 \leq C n^{\frac{-2\alpha+1}{2\alpha}} + C \frac{\log p}{n}.$$

The optimal choice of $k \asymp \lceil n^{\frac{1}{2\alpha}} \rceil$ is due to the bias-variance tradeoff. Combining Theorem 3 with the minimax lower bound derived in Section 3.2, we immediately obtain that the local cropping estimator is rate optimal among all estimators.

Proof. As we discussed in Section 2.1, the direct target of our local cropping estimator is certain tapered population precision matrix with bandwidth k , which can be written as a weighted sum of many principal submatrices of the population precision matrix. We construct this corresponding tapered population precision matrix Ω_k^* as follows. Denote the precision matrix $\Omega \equiv [\omega_{ij}]_{p \times p}$. We define $\Omega_k^* \equiv [\omega_{ij}^*]_{p \times p}$ such that for $i, j \in [p]$,

$$\omega_{ij}^* = m_{ij} \omega_{ij}, \text{ where } m_{ij} = \max\{0, 2 - \frac{1}{k}|i - j|\} - \max\{0, 1 - \frac{1}{k}|i - j|\}. \quad (18)$$

The following lemma elucidates the decomposition of this tapered precision matrix Ω_k^* .

Lemma 1 *The Ω_k^* defined in (18) can be written as*

$$\Omega_k^* = \frac{1}{k} \left(\sum_{m=2}^{2k+1} \left(\sum_{j=-1}^{\lfloor p/2k \rfloor} \mathbf{E}_{m+2kj}^p (\mathbf{C}_{m+2kj}^{2k}(\Omega)) \right) - \sum_{m=2}^{k+1} \left(\sum_{j=-1}^{\lfloor p/k \rfloor} \mathbf{E}_{m+kj}^p (\mathbf{C}_{m+2kj}^k(\Omega)) \right) \right).$$

The proof of Lemma 1 can be found in [13] (refer to the proof of Lemma 1 with covariance matrix therein replaced by the precision matrix), and thus omitted. Define

$$\tilde{\Omega}_k^* = \frac{1}{k} \left(\sum_{m=2}^{2k+1} \left(\sum_{j=-1}^{\lfloor p/2k \rfloor} \mathbf{E}_{m+2kj}^p (\tilde{\Omega}_{m+2kj}^{(2k)}) \right) - \sum_{m=2}^{k+1} \left(\sum_{j=-1}^{\lfloor p/k \rfloor} \mathbf{E}_{m+kj}^p (\tilde{\Omega}_{m+2kj}^{(k)}) \right) \right).$$

It is easy to check $\tilde{\Omega}_k^{\text{op}} = \mathbf{P}_\eta(\tilde{\Omega}_k^*)$. Since the eigenvalues of Ω are in the interval $[\eta^{-1}, \eta]$, according to Equation (54) in Lemma 21, we have

$$\mathbb{E} \|\tilde{\Omega}_k^{\text{op}} - \Omega\|_{\text{op}} \leq 2\mathbb{E} \|\tilde{\Omega}_k^* - \Omega\|_{\text{op}}.$$

The minimax risk can be decomposed into

$$\begin{aligned} \mathbb{E} \|\tilde{\Omega}_k^{\text{op}} - \Omega\|_{\text{op}}^2 &\leq 4\mathbb{E} \|\tilde{\Omega}_k^* - \Omega\|_{\text{op}}^2 \\ &\leq 8\mathbb{E} \|\tilde{\Omega}_k^* - \Omega_k^*\|_{\text{op}}^2 + 8\|\Omega_k^* - \Omega\|_{\text{op}}^2. \end{aligned} \quad (19)$$

The following lemma bounds the bias between our direct target Ω_k^* and the population precision matrix.

Lemma 2 *For Ω in the parameter space $\mathcal{P}_\alpha(\eta, M)$ defined in (2) with $\alpha > \frac{1}{2}$, Ω_k^* defined in (18), we have*

$$\|\Omega_k^* - \Omega\|_{\text{op}}^2 \leq Ck^{-2\alpha+1}.$$

Remark 3 *Unlike existing methods, our procedure does not directly utilize the decay structure of Cholesky factor. Consequently, the proof of Lemma 2 is involved and requires a block-wise partition strategy.*

Then we turn to the analysis of $\mathbb{E}\|\tilde{\Omega}_k^* - \Omega_k^*\|_{\text{op}}^2$.

$$\begin{aligned} \mathbb{E}\|\tilde{\Omega}_k^* - \Omega_k^*\|_{\text{op}}^2 &\leq 2\mathbb{E}\left(\frac{1}{k} \sum_{m=2}^{2k+1} \left\| \sum_{j=-1}^{\lfloor p/2k \rfloor} \mathbf{E}_{m+2kj}^p(\tilde{\Omega}_{m+2kj}^{(2k)}) - \sum_{j=-1}^{\lfloor p/2k \rfloor} \mathbf{E}_{m+2kj}^p(\mathbf{C}_{m+2kj}^{2k}(\Omega)) \right\|_{\text{op}}\right)^2 \\ &\quad + 2\mathbb{E}\left(\frac{1}{k} \sum_{m=2}^{k+1} \left\| \sum_{j=-1}^{\lfloor p/k \rfloor} \mathbf{E}_{m+kj}^p(\tilde{\Omega}_{m+kj}^{(k)}) - \sum_{j=-1}^{\lfloor p/k \rfloor} \mathbf{E}_{m+kj}^p(\mathbf{C}_{m+kj}^k(\Omega)) \right\|_{\text{op}}\right)^2. \end{aligned} \quad (20)$$

These two terms can be bounded in the same way, we only focus on the second term.

$$\begin{aligned} &\mathbb{E}\left(\frac{1}{k} \sum_{m=2}^{k+1} \left\| \sum_{j=-1}^{\lfloor p/k \rfloor} \mathbf{E}_{m+kj}^p(\tilde{\Omega}_{m+kj}^{(k)}) - \sum_{j=-1}^{\lfloor p/k \rfloor} \mathbf{E}_{m+kj}^p(\mathbf{C}_{m+kj}^k(\Omega)) \right\|_{\text{op}}\right)^2 \\ &\leq \mathbb{E}\left(\max_m \left\| \sum_{j=-1}^{\lfloor p/k \rfloor} (\mathbf{E}_{m+kj}^p(\tilde{\Omega}_{m+kj}^{(k)}) - \mathbf{E}_{m+kj}^p(\mathbf{C}_{m+kj}^k(\Omega))) \right\|_{\text{op}}^2\right) \\ &\leq \mathbb{E}\left(\max_{m,j} \|\tilde{\Omega}_{m+kj}^{(k)} - \mathbf{C}_{m+kj}^k(\Omega)\|_{\text{op}}^2\right) \\ &\leq 2\mathbb{E}\left(\max_{m \in [p]} \|\tilde{\Omega}_m^{(k)} - \mathbf{C}_{k+1}^k((\mathbf{C}_m^{3k}(\Omega^{-1})^{-1}))\|_{\text{op}}^2\right) + 2\left(\max_{m \in [p]} \|\mathbf{C}_{k+1}^k((\mathbf{C}_m^{3k}(\Omega^{-1})^{-1})) - \mathbf{C}_m^k(\Omega)\|_{\text{op}}^2\right), \end{aligned} \quad (21)$$

where we further have variance term and bias term of local estimators. For the variance term in (21), we further have

$$\|\tilde{\Omega}_m^{(k)} - \mathbf{C}_{k+1}^k((\mathbf{C}_m^{3k}(\Omega^{-1})^{-1}))\|_{\text{op}} \quad (22)$$

$$\begin{aligned} &\leq \|(\tilde{\Sigma}_{m-k}^{(3k)})^{-1} - (\mathbf{C}_m^{3k}(\Omega^{-1})^{-1})\|_{\text{op}} \\ &\leq \eta^2 \|\tilde{\Sigma}_{m-k}^{(3k)} - \mathbf{C}_m^{3k}(\Omega^{-1})\|_{\text{op}} \\ &\leq 2\eta^2 \|\mathbf{C}_m^{3k}(\frac{1}{n}\mathbf{Z}\mathbf{Z}^T) - \mathbf{C}_m^{3k}(\Omega^{-1})\|_{\text{op}}. \end{aligned} \quad (23)$$

The last two inequalities hold because of the fact that the eigenvalues of $\tilde{\Sigma}_{m-k}^{(3k)}$ and $\mathbf{C}_m^{3k}(\Omega^{-1})$ are in the interval $[\eta^{-1}, \eta]$, and Lemma 21. The following concentration inequality of sample covariance matrix facilitates our proof.

Lemma 3 *For the observations \mathbf{Z} following certain sub-Gaussian distribution with constant ρ with precision matrix Ω , we have*

$$\mathbb{E}\left(\max_{m \in [p]} \|(\mathbf{C}_m^{3k}(\frac{1}{n}\mathbf{Z}\mathbf{Z}^T) - \mathbf{C}_m^{3k}(\Omega^{-1}))\|_{\text{op}}^2\right) \leq C \frac{\log p + k}{n}.$$

Lemma 3 is an extension of the result in Chapter 2 of [44]. Its proof can be found in Lemma 3 of [13].

Combining Lemma 3, (21) and (23), we have

$$\mathbb{E} \left(\max_{m \in [p]} \|\tilde{\Omega}_m^{(k)} - \mathbf{C}_{k+1}^k((\mathbf{C}_m^{3k}(\Omega^{-1})^{-1}))\|_{\text{op}}^2 \right) \leq C \frac{\log p + k}{n}. \quad (24)$$

We turn to bounding the bias term in local estimator in (21).

Lemma 4 *Assume that $\Omega \in \mathcal{P}_\alpha(\eta, M)$ defined in (2) with $\alpha > \frac{1}{2}$. Then we have*

$$\|\mathbf{C}_{k+1}^k((\mathbf{C}_m^{3k}(\Omega^{-1})^{-1})) - \mathbf{C}_m^k(\Omega)\|_{\text{op}}^2 \leq C k^{-2\alpha+1}.$$

The specific constant C is given in the proof of this lemma. Lemma 4, together with (24), (21) and (20), implies that

$$\mathbb{E} \|\tilde{\Omega}_k^* - \Omega_k^*\|_{\text{op}}^2 \leq C \frac{\log p + k}{n} + C k^{-2\alpha+1}. \quad (25)$$

Plugging Lemma 2 and (25) into (19), we finish the proof of Theorem 3. ■

3.2 Minimax Lower Bound under the Operator Norm over $\mathcal{P}_\alpha(\eta, M)$

Theorem 3 in Section 3.1 proves that the local cropping estimator defined in (12) attains the convergence rate of $n^{-\frac{2\alpha+1}{2\alpha}} + \frac{\log p}{n}$. In this section, we establish the following matching lower bound, which proves the rate optimality of the local cropping estimator.

Theorem 4 *The minimax risk of estimating the precision matrix Ω over $\mathcal{P}_\alpha(\eta, M)$ defined in (2) under the operator norm with $\alpha \geq \frac{1}{2}$ satisfies*

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{P}_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \geq \frac{\tau^2}{32} \left(n^{-\frac{2\alpha+1}{2\alpha}} + \frac{\log p}{n} \right), \quad (26)$$

where $0 < \tau < \min\{M, \frac{1}{4}\eta^{-1}, \eta^{\frac{1}{2}} - 1\}$.

Remark 4 *Theorems 3 and 4 together show that the minimax risk for estimating the precision matrices over $\mathcal{P}_\alpha(\eta, M)$ stated in (4) of Theorem 1. It is worthwhile to notice that there is no consistent estimator over $\mathcal{P}_\alpha(\eta, M)$ under the operator norm, when $\alpha \leq \frac{1}{2}$.*

Proof. The lower bound of parameter space $\mathcal{P}_\alpha(\eta, M)$ can be established by the lower bounds over its subsets. We construct two subsets \mathcal{P}_1 and \mathcal{P}_2 and calculate the lower bound over those two subsets separately. Let τ be a positive constant which is less than $\min\{M, \frac{1}{4}\eta^{-1}, \eta^{\frac{1}{2}} - 1\}$.

First, we construct \mathcal{P}_1 . Set $k = \min\{\lceil n^{\frac{1}{2\alpha}} \rceil, \frac{p}{2}\}$. Set the index set $\Theta = \{0, 1\}^k$, i.e., for any $\theta \equiv \{\theta_i\}_{1 \leq i \leq k} \in \Theta$, each θ_i is either 0 or 1. Then we define the $k \times k$ matrix $A_k^*(\theta) \equiv [a_{ij}]_{k \times k}$ with $a_{ij} = \tau n^{-\frac{1}{2}} \theta_i \mathbf{1}(j = k)$ and

$$A(\theta) = \begin{bmatrix} 0_{k \times k} & 0_{k \times k} & 0_{k \times (p-2k)} \\ A_k^*(\theta) & 0_{k \times k} & 0_{k \times (p-2k)} \\ 0_{(p-2k) \times k} & 0_{(p-2k) \times k} & 0_{(p-2k) \times (p-2k)} \end{bmatrix}.$$

We then define \mathcal{P}_1 as the collection of 2^k matrices indexed by Θ ,

$$\mathcal{P}_1 = \{\Omega(\theta) : \Omega(\theta) = (I_p - A(\theta))^T (I_p - A(\theta)), \theta \in \Theta\}. \quad (27)$$

Next, we construct \mathcal{P}_2 as the collection of the diagonal matrices in the following equation,

$$\mathcal{P}_2 = \left\{ \Omega(m) \equiv [w_{ij}(m)]_{p \times p} : w_{ij}(m) = \mathbf{1}(i = j) - \frac{\tau a^{\frac{1}{2}}}{1 + \tau a^{\frac{1}{2}}} \mathbf{1}(i = j = m), 0 \leq m \leq p \right\}, \quad (28)$$

where $a = \min\{\frac{\log p}{n}, 1\}$.

Lemma 5 \mathcal{P}_1 and \mathcal{P}_2 are subsets of $\mathcal{P}_\alpha(\eta, M)$.

Note that we assume $\log p = O(n)$ and $n = O(p)$. Without loss of generality, we further assume $\log p < n < p$. For any estimator $\tilde{\Omega}$ based on n i.i.d. observations, we establish the lower bounds over those two subsets in Sections 3.2.1 and 3.2.2 respectively,

$$\sup_{\mathcal{P}_1} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \geq \frac{\tau^2}{16} n^{-1} \min\{n^{\frac{1}{2\alpha}}, \frac{p}{2}\} \geq \frac{\tau^2}{16} n^{-\frac{2\alpha+1}{2\alpha}}, \quad (29)$$

$$\sup_{\mathcal{P}_2} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \geq \frac{\tau^2}{16} n^{-1} \min\{\log p, n\} \geq \frac{\tau^2 \log p}{16 n}. \quad (30)$$

According to Lemma 5, $(\mathcal{P}_1 \cup \mathcal{P}_2) \subset \mathcal{P}_\alpha(\eta, M)$. Therefore, we obtain

$$\begin{aligned} \sup_{\mathcal{P}_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 &\geq \max\{\sup_{\mathcal{P}_1} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2, \sup_{\mathcal{P}_2} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2\} \\ &\geq \frac{\tau^2}{32} \left(n^{-\frac{2\alpha+1}{2\alpha}} + \frac{\log p}{n} \right), \end{aligned}$$

which completes the proof of Theorem 4. ■

We introduce some further notation before establishing (29) using Assouad's method in Section 3.2.1 and (30) using Le Cam's method in Section 3.2.2. Let $H(\theta, \theta') = \sum_{i=1}^k |\theta_i - \theta'_i|$ be the Hamming distance on $\{0, 1\}^k$, which is the number of different elements between θ and θ' . The total variation affinity $\|P \wedge Q\| = \int p \wedge q \, d\mu$, where p and q are the density functions of two probability measure P and Q with respect to any common dominating measure μ .

3.2.1 Assouad's method in proof of (29)

Assouad's method [1] is a powerful tool to provide the lower bound over distributions indexed by the hypercube $\Theta = \{0, 1\}^k$. Let P_θ be the distribution generated from observations indexed by $\Omega(\theta)$. The proof of Lemma 6 can be found in [53], and thus omitted.

Lemma 6 (Assouad) *Let $\tilde{\Omega}$ be an estimator based on observations from a distribution in the collection $\{P_\theta, \theta \in \Theta\}$, where $\Theta = \{0, 1\}^k$. Then*

$$\sup_{\theta \in \Theta} 2^2 \mathbb{E}_\theta \|\tilde{\Omega} - \Omega(\theta)\|_2^2 \geq \min_{H(\theta, \theta') \geq 1} \frac{\|\Omega(\theta) - \Omega(\theta')\|_2^2 k}{H(\theta, \theta')} \frac{1}{2} \min_{H(\theta, \theta')=1} \|P_\theta \wedge P_{\theta'}\|.$$

Applying the Assouad's method to the subset \mathcal{P}_1 , we have the following results.

Lemma 7 *Let P_θ be the joint distribution of n i.i.d. observations from $N(0, \Omega(\theta)^{-1})$, where $\Omega(\theta) \in \mathcal{P}_1$ defined in (27). Then*

$$\min_{H(\theta, \theta')=1} \|P_\theta \wedge P_{\theta'}\| \geq 0.5.$$

Lemma 8 *Consider all $\Omega(\theta) \in \mathcal{P}_1$ defined in (27). Then*

$$\min_{H(\theta, \theta') \geq 1} \frac{\|\Omega(\theta) - \Omega(\theta')\|_2^2}{H(\theta, \theta')} \geq (\tau n^{-\frac{1}{2}})^2.$$

Lemmas 6, 7 and 8 together imply the desired (29), with the choice $k = \lceil n^{\frac{1}{2\alpha}} \rceil$. The proofs of the above lemmas can be found in the supplement.

3.2.2 Le Cam's method in proof of (30)

Le Cam's method can be used to establish the lower bound via testing a single distribution against a convex hull of distributions. Set $r = \inf_{m \in [p]} \|\Omega(0) - \Omega(m)\|_{\text{op}}^2$. Let P_i be the distribution generated from observations indexed by $\Omega(i)$, where $0 \leq i \leq p$. Define $\bar{P} = \sum_{m=1}^p P_m$. The proof of the following lemma can be found in [53], and thus omitted.

Lemma 9 (Le Cam) *Let $\tilde{\Omega}$ be an estimator based on observations from a distribution in the collection $\{P_i, 0 \leq i \leq p\}$. Then*

$$\sup_{0 \leq m \leq p} \mathbb{E} \|\tilde{\Omega} - \Omega(m)\|_{\text{op}}^2 \geq \frac{1}{2} r \|P_0 \wedge \bar{P}\|.$$

Applying Le Cam's method to the subset \mathcal{P}_2 , we obtain that $r = \left(\frac{\tau a^{\frac{1}{2}}}{1 + \tau a^{\frac{1}{2}}} \right)^2 \geq \frac{1}{4} \tau^2 a$ and the following results.

Lemma 10 *Let P_m be the joint distribution of n i.i.d. observations from $N(0, \Omega(m)^{-1})$, where $\Omega(m) \in \mathcal{P}_2$ defined in (28). Then*

$$\|P_0 \wedge \bar{P}\| > \frac{7}{8}.$$

Combining the above results in Theorems 9 and 10, we obtain the desired (30), i.e.,

$$\sup_{0 \leq m \leq p} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \geq \frac{7}{64} \tau^2 a \geq \frac{\tau^2}{16} \min\left\{\frac{\log p}{n}, 1\right\}.$$

3.3 Rate Optimality under the Operator Norm over $\mathcal{Q}_\alpha(\eta, M)$

3.3.1 Minimax upper bound

In this section, we establish the optimal rate of convergence over $\mathcal{Q}_\alpha(\eta, M)$ under the operator norm. Our estimation procedure remains the local cropping estimator, except that the bandwidth is $k \asymp n^{\frac{1}{2\alpha+1}}$, which is due to smaller bias terms.

Theorem 5 *When $\lceil n^{\frac{1}{2\alpha+1}} \rceil \leq p$, the local cropping estimator defined in (12) of the precision matrix Ω over $\mathcal{Q}_\alpha(\eta, M)$ given in (3) satisfies*

$$\sup_{\mathcal{Q}_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega}_k^{\text{op}} - \Omega\|_{\text{op}}^2 \leq Ck^{-2\alpha} + C \frac{\log p + k}{n},$$

when $k = \lceil n^{\frac{1}{2\alpha+1}} \rceil$, we have

$$\sup_{\mathcal{Q}_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega}_k^{\text{op}} - \Omega\|_{\text{op}}^2 \leq Cn^{\frac{-2\alpha}{2\alpha+1}} + C \frac{\log p}{n}.$$

Proof. We employ the same proof strategy as that of Theorem 3. Only two lemmas bounding bias terms need to be replaced. We only emphasize the differences here.

We replace Lemma 2 in the proof by Lemma 11, which bounds the distance of the population precision matrix and its tapered one.

Lemma 11 *For Ω in the parameter space $\mathcal{Q}_\alpha(\eta, M)$ defined in (3), Ω_k^* is defined in (18), we have*

$$\|\Omega_k^* - \Omega\|_{\text{op}}^2 \leq Ck^{-2\alpha}.$$

In addition, we replace Lemma 4 by Lemma 12, which bounds the bias term of each local estimator.

Lemma 12 *For $\Omega \in \mathcal{Q}_\alpha(\eta, M)$ defined in (2) with $\alpha > 0$, we have*

$$\|\mathbf{C}_{k+1}^k((\mathbf{C}_m^{3k}(\Omega^{-1})^{-1}) - \mathbf{C}_m^k(\Omega))\|_{\text{op}}^2 \leq Ck^{-2\alpha}.$$

The remaining part of the proof remains the same, including a similar upper bound for the variance term stated in Lemma 3. Therefore, we complete our proof. ■

3.3.2 Minimax lower bound

Theorem 6 *The minimax risk for estimating the precision matrix Ω over $\mathcal{Q}_\alpha(\eta, M)$ defined in (3) under the operator norm with $\alpha > 0$ satisfies*

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{Q}_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \geq \frac{\tau^2}{32} \left(n^{\frac{-2\alpha}{2\alpha+1}} + \frac{\log p}{n} \right). \quad (31)$$

Remark 5 Theorems 5 and 6 together show that the minimax risk for estimating the precision matrices over $\mathcal{Q}_\alpha(\eta, M)$ stated in (5) of Theorem 1. In contrast to $\mathcal{P}_\alpha(\eta, M)$, the optimal rate of convergence over $\mathcal{Q}_\alpha(\eta, M)$ is faster. In particular, rate-optimal local cropping estimators are always consistent as long as $\alpha > 0$.

Proof. To establish the lower bound for $\mathcal{Q}_\alpha(\eta, M)$ in which the decay of a_{ij} is in the entry-wise fashion, we repeat the proof scheme in Section 3.2 with a few changes. Let τ be a positive constant which is less than $\min\{M, \frac{1}{4}\eta^{-1}, \eta^{\frac{1}{2}} - 1\}$.

Set $k = \min\{\lceil n^{\frac{1}{2\alpha+1}} \rceil, \frac{p}{2}\}$ and the index set $\Theta = \{0, 1\}^k$, i.e., for any $\theta \in \Theta$, $\theta \equiv \{\theta_i\}_{1 \leq i \leq k}$, each θ_i is either 0 or 1. Define the $k \times k$ matrix $B_k^*(\theta) \equiv [b_{ij}]_{k \times k}$ with $b_{ij} = \tau(nk)^{-\frac{1}{2}}\theta_i$. Define

$$B(\theta) = \begin{bmatrix} 0_{k \times k} & 0_{k \times k} & 0_{k \times (p-2k)} \\ B_k^*(\theta) & 0_{k \times k} & 0_{k \times (p-2k)} \\ 0_{(p-2k) \times k} & 0_{(p-2k) \times k} & 0_{(p-2k) \times (p-2k)} \end{bmatrix}.$$

We construct the collection of 2^k matrices as

$$\mathcal{P}_3 = \{\Omega(\theta) : \Omega(\theta) = (I_p - B(\theta))^T(I_p - B(\theta)), \theta \in \Theta\}. \quad (32)$$

Lemma 13 \mathcal{P}_3 is a subset of $\mathcal{Q}_\alpha(\eta, M)$.

Let P_θ be the joint distribution of n i.i.d. observations from $N(0, \Omega(\theta)^{-1})$, where $\Omega(\theta) \in \mathcal{P}_3$ defined in (32). Parallel to Lemmas 7 and 8 and the lower bound (29) for \mathcal{P}_1 , we establish the following lower bound for \mathcal{P}_3 .

Lemma 14 Consider all $\Omega(\theta) \in \mathcal{P}_3$ defined in (32). Then

$$\min_{H(\theta, \theta')=1} \|P_\theta \wedge P_{\theta'}\| \geq 0.5, \quad (33)$$

$$\min_{H(\theta, \theta') \geq 1} \frac{\|\Omega(\theta) - \Omega(\theta')\|_2^2}{H(\theta, \theta')} \geq (\tau n^{-\frac{1}{2}})^2. \quad (34)$$

According to Assouad's method, for any estimator $\tilde{\Omega}$ based on n i.i.d. observations, we have

$$\sup_{\mathcal{P}_3} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \geq \frac{\tau^2}{16} n^{-1} \min\{n^{\frac{1}{2\alpha+1}}, \frac{p}{2}\}. \quad (35)$$

It is easy to show $(\mathcal{P}_3 \cup \mathcal{P}_2) \subset \mathcal{Q}_\alpha(\eta, M)$, where \mathcal{P}_2 is defined in (28). Therefore, combining (35) and (30), we complete the proof of Theorem 6. ■

Remark 6 The estimation of the covariance matrix Σ is of significant importance as well. We propose the estimator of Σ by inverting our estimator $\tilde{\Omega}_k^{\text{op}}$ given in (12). The results and the analysis given in Section 3 can be used to establish the minimax optimality

of our estimator under the operator norm. According to the inequality $\|(\tilde{\Omega}_k^{\text{op}})^{-1} - \Sigma\|_{\text{op}} \leq \|(\tilde{\Omega}_k^{\text{op}})^{-1}\|_{\text{op}} \|\tilde{\Omega}_k^{\text{op}} - \Omega\|_{\text{op}} \|\Omega^{-1}\|_{\text{op}}$ and the fact that both $\|(\tilde{\Omega}_k^{\text{op}})^{-1}\|_{\text{op}}$ and $\|\Omega^{-1}\|_{\text{op}}$ are bounded by η , we establish the upper bound of our estimator $(\tilde{\Omega}_k^{\text{op}})^{-1}$. Furthermore, considering the analog between the covariance matrix and the precision matrix in the subset \mathcal{P}_1 and \mathcal{P}_2 defined in (27) and (28), the matching lower bound can be proved by a similar argument in Section 3.2. Therefore, we have the following rate optimality of estimating the covariance matrix under the operator norm, which can be achieved by estimator $(\tilde{\Omega}_k^{\text{op}})^{-1}$,

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{P}_{\alpha}(\eta, M)} \mathbb{E} \|\tilde{\Omega}^{-1} - \Omega^{-1}\|_{\text{op}}^2 \asymp n^{\frac{-2\alpha+1}{2\alpha}} + \frac{\log p}{n},$$

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{Q}_{\alpha}(\eta, M)} \mathbb{E} \|\tilde{\Omega}^{-1} - \Omega^{-1}\|_{\text{op}}^2 \asymp n^{\frac{-2\alpha}{2\alpha+1}} + \frac{\log p}{n}.$$

4 Rate Optimality under the Frobenius Norm

In this section, we establish that the optimal rates of convergence for estimating the precision matrix over the parameter spaces $\mathcal{P}_{\alpha}(\eta, M)$ and $\mathcal{Q}_{\alpha}(\eta, M)$ are identical under the Frobenius norm. Since $\mathcal{Q}_{\alpha}(\eta, \alpha M) \subset \mathcal{P}_{\alpha}(\eta, M)$, one immediately obtain that

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{P}_{\alpha}(\eta, M)} \frac{1}{p} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{F}}^2 \geq \inf_{\tilde{\Omega}} \sup_{\mathcal{Q}_{\alpha}(\eta, \alpha M)} \frac{1}{p} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{F}}^2. \quad (36)$$

In order to show (6) in Theorem 2, it suffices to establish the upper bound over the parameter space $\mathcal{P}_{\alpha}(\eta, M)$ and the matching lower bound over the parameter space $\mathcal{Q}_{\alpha}(\eta, M)$. We assume that \mathbf{X} follows the p -variate Gaussian distribution, with mean zero and precision matrix Ω in this section.

4.1 Minimax Upper Bound under the Frobenius Norm

In this section, we establish the following risk upper bound of the regression-based block-thresholding estimation procedure we proposed in Section 2.2 under the Frobenius norm over $\mathcal{P}_{\alpha}(\eta, M)$.

Theorem 7 Assume $\lceil n^{\frac{1}{2\alpha+2}} \rceil \leq p$. The estimator defined in (16) of the precision matrix Ω over $\mathcal{P}_{\alpha}(\eta, M)$ and $\mathcal{Q}_{\alpha}(\eta, \alpha M)$ given in (2) and (3) with $k = \lceil n^{\frac{1}{2\alpha+2}} \rceil$ satisfies

$$\sup_{\mathcal{Q}_{\alpha}(\eta, \alpha M)} \frac{1}{p} \mathbb{E} \|\tilde{\Omega}_k^{\text{F}} - \Omega\|_{\text{F}}^2 \leq \sup_{\mathcal{P}_{\alpha}(\eta, M)} \frac{1}{p} \mathbb{E} \|\tilde{\Omega}_k^{\text{F}} - \Omega\|_{\text{F}}^2 \leq C n^{-\frac{2\alpha+1}{2\alpha+2}}. \quad (37)$$

Proof. We focus on the second inequality since the first one is trivial. Note that $\tilde{\Omega}_k^{\text{F}} = (\widetilde{I - A})^T \widetilde{D}^{-1} (\widetilde{I - A})$ according to (16) while $\Omega = (I - A)^T D^{-1} (I - A)$. The risk upper bound can be controlled by bounding $\widetilde{I - A} - (I - A)$ and $\widetilde{D} - D$. To this end, we first provide some properties of our estimator.

Lemma 15 Assume that \mathbf{X} follows the p -variate Gaussian distribution with mean zero and precision matrix $\Omega = (I - A)^T D^{-1} (I - A)$, which belongs to parameter space $\mathcal{P}_\alpha(\eta, M)$ defined in (2). For any fixed i , d_i is the i -th diagonal of D , \mathbf{a}_i is the i -th row of A . \hat{d}_i is defined in (15), and $\hat{\mathbf{a}}_i^*$ is the i -th row of \hat{A}^* defined in (14). Then we have

$$\begin{aligned}\mathbb{E}|\hat{d}_i - d_i|^2 &\leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}, \\ \mathbb{E}\|\hat{\mathbf{a}}_i^* - \mathbf{a}_i\|_{\text{op}}^2 &\leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}.\end{aligned}$$

We are ready to establish the upper bounds of $\widetilde{I - A} - (I - A)$ and $\tilde{D} - D$ separately. Note that $\|\tilde{D}^{-1}\|_{\text{op}} \leq \eta$ and $\|D^{-1}\|_{\text{op}} \leq \eta$, which is due to Lemma 22. Therefore, Lemma 21 yields $\mathbb{E}\|\tilde{D} - D\|_{\text{F}}^2 \leq 4\mathbb{E}\|\hat{D} - D\|_{\text{F}}^2$, which further implies that

$$\begin{aligned}\frac{1}{p}\mathbb{E}\|D^{-1} - \tilde{D}^{-1}\|_{\text{F}}^2 &\leq \frac{1}{p}\mathbb{E}\|\tilde{D}^{-1}\|_{\text{op}}^2 \|\tilde{D} - D\|_{\text{F}}^2 \|D^{-1}\|_{\text{op}}^2 \\ &\leq 4\eta^4 \frac{1}{p}\mathbb{E}\|\hat{D} - D\|_{\text{F}}^2 \\ &\leq 4\eta^4 \frac{1}{p} \sum_i \mathbb{E}|\hat{d}_i - d_i|^2.\end{aligned}$$

Together with Lemma 15, it follows that

$$\frac{1}{p}\mathbb{E}\|D^{-1} - \tilde{D}^{-1}\|_{\text{F}}^2 \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}. \quad (38)$$

Next, we turn to prove that $\frac{1}{p}\mathbb{E}\|\widetilde{I - A} - (I - A)\|_{\text{F}}^2 \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}$. Lemma 21 implies

$$\frac{1}{p}\mathbb{E}\|\widetilde{I - A} - (I - A)\|_{\text{F}}^2 \leq \frac{4}{p}\mathbb{E}\|\hat{A}^* - A\|_{\text{F}}^2 \leq \frac{4}{p} \sum_i \mathbb{E}\|\hat{\mathbf{a}}_i^* - \mathbf{a}_i\|_{\text{op}}^2.$$

Combining above equation with Lemma 15, we have

$$\frac{1}{p}\mathbb{E}\|\widetilde{I - A} - (I - A)\|_{\text{F}}^2 \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}. \quad (39)$$

At last, we derive the risk upper bound of our estimator. It is clear that $\|\widetilde{I - A}\|_{\text{op}} \leq \eta$, $\|\tilde{D}^{-1}\|_{\text{op}} \leq \eta$. According to Lemma 22, $\|I - A\|_{\text{op}} \leq \eta$, $\|D^{-1}\|_{\text{op}} \leq \eta$. Combining these facts with (38) and (39), we have

$$\begin{aligned}\frac{1}{p}\mathbb{E}\|\tilde{\Omega} - \Omega\|_{\text{F}}^2 &\leq \frac{3}{p}\mathbb{E}(\|I - A\|_{\text{op}}^2 \|D^{-1}\|_{\text{op}}^2 \|\widetilde{I - A} - (I - A)\|_{\text{F}}^2 \\ &\quad + \|I - A\|_{\text{op}}^2 \|D^{-1} - \tilde{D}^{-1}\|_{\text{F}}^2 \|\widetilde{I - A}\|_{\text{op}}^2 \\ &\quad + \|\widetilde{I - A} - (I - A)\|_{\text{F}}^2 \|\tilde{D}^{-1}\|_{\text{op}}^2 \|\widetilde{I - A}\|_{\text{op}}^2) \\ &\leq 6\eta^4 \frac{1}{p}\mathbb{E}\|\widetilde{I - A} - (I - A)\|_{\text{F}}^2 + 3\eta^4 \frac{1}{p}\mathbb{E}\|D^{-1} - \tilde{D}^{-1}\|_{\text{F}}^2 \\ &\leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}.\end{aligned}$$

Therefore, we finish the proof of Theorem 7. ■

4.2 Minimax Lower Bound under the Frobenius Norm

In this section, we establish the matching lower bound $n^{-\frac{2\alpha+1}{2\alpha+2}}$ over parameter spaces $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$.

Theorem 8 *The minimax risk for estimating the precision matrix Ω over $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, \alpha M)$ under the Frobenius norm satisfies*

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{P}_\alpha(\eta, M)} \frac{1}{p} \mathbb{E} \|\tilde{\Omega} - \Omega\|_F^2 \geq \inf_{\tilde{\Omega}} \sup_{\mathcal{Q}_\alpha(\eta, \alpha M)} \frac{1}{p} \mathbb{E} \|\tilde{\Omega} - \Omega\|_F^2 \geq \frac{\tau^2}{32} n^{-\frac{2\alpha+1}{2\alpha+2}}.$$

Remark 7 *The minimax risk for estimating the precision matrices over $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$ under the Frobenius norm in Theorem 2 immediately follows from Theorems 7 and 8.*

Proof. It is sufficient to establish the lower bound over $\mathcal{Q}_\alpha(\eta, M)$ since the first inequality immediately follows from (36). We construct a least favorable subset in $\mathcal{Q}_\alpha(\eta, M)$. Without loss of generality, we assume $\frac{p}{2k}$ is an integer where $k = \min\{\lceil n^{\frac{1}{2\alpha+2}} \rceil, \frac{p}{2}\}$. Define the index set $\Theta' = \{0, 1\}^{\frac{kp}{2}}$. For each $\theta \in \Theta'$, we further denote it as $\frac{p}{2k}$ many k^2 dimensional vectors, i.e., $\theta = \{\theta(s)\}_{1 \leq s \leq \lceil \frac{p}{2k} \rceil}$, where $\theta(s)_{ij}$ is equal to 0 or 1. For such an index θ , there is a corresponding $p \times p$ block diagonal matrix $C(\theta)$ such that each $k \times k$ block $C_s(\theta(s)) \equiv [c(s)_{ij}]_{k \times k}$, where $c(s)_{ij} = \tau n^{-\frac{1}{2}} \theta(s)_{ij}$, $s \in \lceil \frac{p}{2k} \rceil$. We set τ as a positive constant which is less than $\min\{M, \frac{1}{4}\eta^{-1}, \eta^{\frac{1}{2}} - 1\}$.

$$C(\theta) = \begin{bmatrix} \boxed{\begin{matrix} 0_k & 0_k \\ C_1(\theta(1)) & 0_k \end{matrix}} & 0_{2k} & \cdots & 0_{2k} \\ 0_{2k} & \boxed{\begin{matrix} 0_k & 0_k \\ C_2(\theta(2)) & 0_k \end{matrix}} & \cdots & 0_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ 0_{2k} & 0_{2k} & \cdots & \boxed{\begin{matrix} 0_k & 0_k \\ C_{\lceil \frac{p}{2k} \rceil}(\theta(\lceil \frac{p}{2k} \rceil)) & 0_k \end{matrix}} \end{bmatrix}.$$

Finally, we define the subset of $\mathcal{Q}_\alpha(\eta, M)$ indexed by Θ' as follows

$$\mathcal{P}_4 = \{\Omega(\theta) : \Omega(\theta) = (I_p - C(\theta))^T (I_p - C(\theta)), \theta \in \Theta'\}. \quad (40)$$

Lemma 16 \mathcal{P}_4 is a subset of $\mathcal{Q}_\alpha(\eta, M)$.

Applying Lemma 6 to \mathcal{P}_4 , we obtain that,

$$\inf_{\tilde{\Omega}} \max_{\theta \in \Omega(\Theta')} 2^2 \mathbb{E}_\theta \|\tilde{\Omega} - \Omega(\theta)\|_F^2 \geq \min_{H(\theta, \theta') \geq 1} \frac{\|\Omega(\theta) - \Omega(\theta')\|_F^2}{H(\theta, \theta')} \frac{kp}{4} \min_{H(\theta, \theta')=1} \|P_\theta \wedge P_{\theta'}\| \quad (41)$$

Lemma 17 *Let P_θ be the joint distribution of n i.i.d. observations from $N(0, \Omega(\theta)^{-1})$, where $\Omega(\theta) \in \mathcal{P}_4$ defined in (40). Then*

$$\min_{H(\theta, \theta')=1} \|P_\theta \wedge P_{\theta'}\| \geq 0.5, \quad (42)$$

and

$$\min_{H(\theta, \theta') \geq 1} \frac{\|\Omega(\theta) - \Omega(\theta')\|_F^2}{H(\theta, \theta')} \geq \tau^2 n^{-1}. \quad (43)$$

Applying Lemma 17 into (41), we obtain

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{Q}_\alpha(\eta, M)} \frac{1}{p} \mathbb{E} \|\tilde{\Omega} - \Omega\|_F^2 \geq \inf_{\tilde{\Omega}} \sup_{\mathcal{P}_4} \frac{1}{p} \mathbb{E} \|\tilde{\Omega} - \Omega\|_F^2 \geq \frac{\tau^2}{32} n^{-1} \min \left\{ n^{\frac{1}{2\alpha+2}}, \frac{p}{2} \right\},$$

which completes the proof of Theorem 8, noting that $n < p$. ■

5 Extension to Nonparanormal Distributions

In this section, we extend the minimax framework to the nonparanormal model. Assume that $\mathbf{X} = (X_1, X_2, \dots, X_p)^T$ follows the p -variate Gaussian distribution with covariance matrix Σ . Instead of n i.i.d. copies $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n$ of \mathbf{X} , we only observe their transformations. Specifically, let $\mathbf{Y} = (f_1(X_1), f_2(X_2), \dots, f_p(X_p))^T$ be the transformed variables of \mathbf{X} , where each f_i is certain unknown strictly increasing function. Then our observation is $\mathbf{Z} = (\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_n)^T \in \mathbb{R}^{n \times p}$, where each \mathbf{Y}_i is the transformed \mathbf{X}_i . This is a form of the Gaussian copula model [5], or the nonparanormal model [36]. To avoid the identifiability issue, we set $\text{diag}(\Sigma) = I$, which makes Σ the correlation matrix. Here we consider the same structural assumption as in previous sections on the inverse of the correlation matrix, which is denoted by Ω . Based on $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$ defined in (2) and (3), the following two types of parameter spaces are of interest,

$$\mathcal{P}'_\alpha(\eta, M) = \left\{ \{\Omega, \{f_i\}\} : \begin{array}{l} \text{diag}(\Omega^{-1}) = I, \quad \Omega \in \mathcal{P}_\alpha(\eta, M); \\ f_i \text{ is strictly increasing, } i \in [p]. \end{array} \right\}, \quad (44)$$

and

$$\mathcal{Q}'_\alpha(\eta, M) = \left\{ \{\Omega, \{f_i\}\} : \begin{array}{l} \text{diag}(\Omega^{-1}) = I, \quad \Omega \in \mathcal{Q}_\alpha(\eta, M); \\ f_i \text{ is strictly increasing, } i \in [p]. \end{array} \right\}. \quad (45)$$

Our goal is to estimate the latent correlation structure, the inverse of the correlation matrix Ω , using the observation \mathbf{Z} . We establish the minimax risk of estimating Ω over the parameter spaces $\mathcal{P}'_\alpha(\eta, M)$ and $\mathcal{Q}'_\alpha(\eta, M)$ under the operator norm in the following theorem.

Theorem 9 *Assume $\log p = O(n)$, $n = O(p)$. Then for the nonparanormal model, the minimax risk of estimating Ω under the operator norm over $\mathcal{P}'_\alpha(\eta, M)$ with $\alpha > \frac{1}{2}$ satisfies*

$$\inf_{\tilde{\Omega}} \sup_{\{\Omega, \{f_i\}\} \in \mathcal{P}'_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \asymp n^{\frac{1-2\alpha}{2\alpha}} + \frac{\log p}{n}. \quad (46)$$

The minimax risk of estimating Ω under the operator norm over $\mathcal{Q}'_\alpha(\eta, M)$ satisfies

$$\inf_{\tilde{\Omega}} \sup_{\{\Omega, \{f_i\}\} \in \mathcal{Q}'_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \asymp n^{\frac{-2\alpha}{1+2\alpha}} + \frac{\log p}{n}. \quad (47)$$

Rank-based estimator are widely applied in the nonparanormal model. Progress has been made in this field during the last decade especially for high-dimensional statistics. For instance, see [38] for bandable correlation matrix estimation, [4] for Gaussian graphical models, and [23] for multi-task regression via Cholesky decomposition.

5.1 Estimation Procedure

We introduce our rate-optimal estimation procedure over the parameter spaces $\mathcal{P}'_\alpha(\eta, M)$ and $\mathcal{Q}'_\alpha(\eta, M)$ under the operator norm. The approach to estimate the inverse of the correlation matrix in nonparanormal model is almost the same as the estimation scheme of the precision matrix under the operator norm in Section 2.1, except that the sample covariance matrix needs to be replaced by its rank-based nonparametric variant via Kendall's tau (τ) [31] or Spearman's correlation coefficient rho (ρ) [46].

Kendall's tau is defined as

$$\hat{\tau}_{ij} = \frac{2}{n(n-1)} \sum_{1 \leq k_1 < k_2 \leq n} \text{sgn}(Z_{k_1 i} - Z_{k_2 i}) \text{sgn}(Z_{k_1 j} - Z_{k_2 j}).$$

Then define

$$\hat{\Sigma}^\tau = [\sin(\frac{\pi}{2} \hat{\tau}_{ij})]_{p \times p}. \quad (48)$$

Spearman's rho is defined as

$$\hat{\rho}_{ij} = \frac{\sum_{k=1}^n (r_{ki} - (n+1)/2)(r_{kj} - (n+1)/2)}{\sqrt{\sum_{k=1}^n (r_{ki} - (n+1)/2)^2 \sum_{k=1}^n (r_{kj} - (n+1)/2)^2}},$$

where r_{ij} is the rank of Z_{ij} among $Z_{1j}, Z_{2j}, \dots, Z_{nj}$. Define

$$\hat{\Sigma}^\rho = [2 \sin(\frac{\pi}{6} \hat{\rho}_{ij})]_{p \times p}. \quad (49)$$

It is well-known that both $\hat{\Sigma}^\tau$ and $\hat{\Sigma}^\rho$ are unbiased estimators of the population correlation matrix Σ . We adopt almost the same estimation procedure proposed in Section 2.1, but replacing $\frac{1}{n} \mathbf{Z}^T \mathbf{Z}$ in (10) with either $\hat{\Sigma}^\tau$ or $\hat{\Sigma}^\rho$. By this way, we construct the nonparametric local cropping estimators $\tilde{\Omega}_k^\tau$ and $\tilde{\Omega}_k^\rho$ in replace of $\tilde{\Omega}_k^{\text{op}}$ in (12). Note that the optimal choices of the bandwidth k are picked differently over two types of parameter spaces $\mathcal{P}'_\alpha(\eta, M)$ and $\mathcal{Q}'_\alpha(\eta, M)$ as we did over $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$ in Section 2.1.

5.2 Minimax Optimality

In this section, we prove that both $\tilde{\Omega}_k^\tau$ and $\tilde{\Omega}_k^\rho$ achieve the minimax optimality under the operator norm over the parameter space $\mathcal{P}'_\alpha(\eta, M)$. The minimax optimality over $\mathcal{Q}'_\alpha(\eta, M)$ can be established in the same way.

First, we derive the risk lower bound over $\mathcal{P}'_\alpha(\eta, M)$. Following the same strategy in Section 3.2, we construct the two subset \mathcal{P}'_1 and \mathcal{P}'_2 based on \mathcal{P}_1 and \mathcal{P}_2 given in (27) and (28). Define the subset

$$\mathcal{P}'_1 = \left\{ \{\Omega, \{f_i\}\} : \begin{array}{l} \Omega = \text{diag}(\Omega'^{-1})^{\frac{1}{2}} \Omega' \text{diag}(\Omega'^{-1})^{\frac{1}{2}}, \quad \Omega' \in \mathcal{P}_1; \\ f_i(x) = \text{diag}(\Omega'^{-1})^{\frac{1}{2}}_i x, \quad i \in [p]. \end{array} \right\}, \quad (50)$$

Lemma 18 \mathcal{P}'_1 is a subset of $\mathcal{P}'_\alpha(\eta^2, M\eta)$.

One can easily check that the probability measure of the Gaussian distribution with precision matrix Ω and transformation $\{f_i\}$, where $\{\Omega, \{f_i\}\} \in \mathcal{P}'_1$ is equivalent to the the probability measure of the Gaussian distribution with precision matrix Ω' , where $\Omega' \in \mathcal{P}_1$. Therefore, by this one-to-one correspondence of probability measure between index sets \mathcal{P}'_1 and \mathcal{P}_1 , we immediately have

$$\sup_{\{\Omega, \{f_i\}\} \in \mathcal{P}'_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \geq \sup_{\{\Omega, \{f_i\}\} \in \mathcal{P}'_1} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 = \sup_{\Omega' \in \mathcal{P}_1} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2. \quad (51)$$

Applying the same proof strategy in Section 3.2, we have the following result.

Lemma 19 Let $\Omega'(\theta) \in \mathcal{P}_1$ defined in (27), $\Omega = \text{diag}(\Omega'^{-1})^{\frac{1}{2}} \Omega' \text{diag}(\Omega'^{-1})^{\frac{1}{2}}$, we have

$$\min_{H(\theta, \theta') \geq 1} \frac{\|\Omega(\theta) - \Omega(\theta')\|_2^2}{H(\theta, \theta')} \geq (\tau n^{-\frac{1}{2}})^2,$$

where $0 < \tau < \min\{M\eta, \frac{1}{4}\eta^{-2}, \eta - 1\}$.

Then by Assoaud's lemma, we obtain that

$$\inf_{\tilde{\Omega}} \sup_{\Omega' \in \mathcal{P}_1} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \geq \frac{\tau^2}{32} n^{\frac{1-2\alpha}{2\alpha}}.$$

We can use a similar strategy to construct \mathcal{P}'_2 . Note that in this case we need to put $\tau a^{\frac{1}{2}}$ on the first sub-diagonal in $I - A$, instead of the diagonal of Σ . Combined with the result from the Le Cam's lemma on the subset \mathcal{P}'_2 , we have

$$\sup_{\{\Omega, \{f_i\}\} \in \mathcal{P}'_\alpha(\eta, M)} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \geq C(n^{\frac{1-2\alpha}{2\alpha}} + \frac{\log p}{n}). \quad (52)$$

Next, we turn to the risk upper bound of our rank-based local cropping estimators. The risk can be decomposed into the bias terms and the variance term in the same fashion in Section 3.1. Since the bias terms are deterministic and only due to the bandable structure of the Cholesky factor of the inverse correlation matrix, the upper bounds of two bias terms we derived in Lemma 2 and Lemma 4 still hold. For the variance term, a simple extension of Theorem 1 in [38] provides the following result.

Lemma 20 *For any Ω such that $\text{diag}(\Omega^{-1}) = I$ and $\Omega \in \mathcal{P}_\alpha(\eta, M) \cup \mathcal{Q}_\alpha(\eta, M)$, we have*

$$\mathbb{E}\left(\max_{m \in [p]} \|(\mathbf{C}_m^{3k}(\hat{\Sigma}^\tau) - (\mathbf{C}_m^{3k}(\Omega^{-1}))\|_{\text{op}}^2\right) \leq C \frac{\log p + k}{n},$$

$$\mathbb{E}\left(\max_{m \in [p]} \|(\mathbf{C}_m^{3k}(\hat{\Sigma}^\rho) - (\mathbf{C}_m^{3k}(\Omega^{-1}))\|_{\text{op}}^2\right) \leq C \frac{\log p + k}{n}.$$

Replacing Lemma 3 by Lemma 20 and following the rest of the proof in Theorem 3, we finally obtain that

$$\sup_{\{\Omega, \{f_i\}\} \in \mathcal{P}'_\alpha(\eta, M)} \mathbb{E}\|\tilde{\Omega}_k^\tau - \Omega\|_{\text{op}}^2 \asymp \sup_{\{\Omega, \{f_i\}\} \in \mathcal{P}'_\alpha(\eta, M)} \mathbb{E}\|\tilde{\Omega}_k^\rho - \Omega\|_{\text{op}}^2 \leq Cn^{\frac{1-2\alpha}{2\alpha}} + C \frac{\log p}{n}. \quad (53)$$

The lower bound (52) and upper bound (53) together give the optimal rate of convergence (46) in Theorem 9. The optimal rate of convergence (47) in Theorem 9 can be proved similarly. Therefore, we establish the minimax framework for nonparanormal distributions.

6 Numerical Studies

In this section, we turn to the numerical performance of the proposed rate-optimal estimators under the operator norm for $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$ defined in (2) and (3) to further illustrate the fundamental difference of $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$. In addition, we compare them with the banding estimator proposed in [6], which is based on the auto-regression between variables. Specifically, for a given bandwidth $k < n$, the banding estimator is defined as $\tilde{\Omega}^{BL} = (I - \tilde{A}^{BL})^T (\tilde{D}^{BL})^{-1} (I - \tilde{A}^{BL})$. Here the i -th row of the lower triangular matrix \tilde{A}^{BL} is the vector $\hat{\mathbf{a}}_i$ in (13), i.e., the least square estimates of the coefficients for the regression of X_i against $\mathbf{X}_{i-k:i-1}$. The i -th entry of the diagonal matrix \tilde{D}^{BL} is the estimate of the residual variance for the regression of X_i against $\mathbf{X}_{i-k:i-1}$.

6.1 Simulation in $\mathcal{Q}_\alpha(\eta, M)$ under the operator norm

We first focus on the parameter space $\mathcal{Q}_\alpha(\eta, M)$ and compare the performance of local cropping estimator and the banding estimator. Specifically, we generate the precision matrix in the following form:

$$\Omega = (I - A)^T D^{-1} (I - A), \quad A = [a_{ij}]_{p \times p}, \quad D = I_p,$$

where $a_{ij} = -(i - j)^{-\alpha-1}$ when $i > j$; otherwise $a_{ij} = 0$. It is easy to check that $\Omega \in \mathcal{Q}_\alpha(\eta, 1)$ with some large $\eta > 0$. The simulation is done with a range of parameter values for p, n, α . Specifically, the decay rate α ranges from 0.5 to 2 with a step of 0.5, the sample size n ranges from 500 to 4000, the dimension p ranges from 500 to 2000.

In this setting, we compare our local cropping estimator (denoted as cropping.Q.) with the banding estimator (denoted as BL) proposed in [6]. According to [6], the bandwidth of banding estimator is chosen as $k \asymp (n/\log p)^{1/(2\alpha+2)}$. The optimal bandwidth

over $\mathcal{Q}_\alpha(\eta, M)$ is $k \asymp n^{1/(2\alpha+1)}$. In the simulation, the bandwidth of BL estimator is $\lfloor (n/\log p)^{1/(2\alpha+2)} \rfloor$ and the bandwidth of crop.Q is $\lfloor n^{1/(2\alpha+1)} \rfloor$.

Table 1 reports the average errors of the banding estimator (BL) and local cropping estimator (crop.Q) under the operator norm over 100 replications. The smaller errors in each experiment are highlighted in boldface. Fig. 6.2 displays the boxplots of the errors of BL and crop.Q.

It can be seen from Table 1 that crop.Q outperforms BL in most cases with a few exceptions when n is small. As the sample size increases, the average errors of both methods decrease, which matches our intuition. In addition, the dimension p has minor effect on the errors of both estimators, which is partially reflected by the optimal rates (dominating term $n^{\frac{-2\alpha}{2\alpha+1}}$) obtained in Theorem 1. For each fixed dimension p , the superiority crop.Q over BL becomes more significant as the sample size n increases, which implies that BL estimator is indeed sub-optimal.

6.2 Simulation in $\mathcal{P}_\alpha(\eta, M)$ under the operator norm

We demonstrate the fundamental difference between two types of parameter space $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$ by numerical studies in this section. Of note, although local cropping estimators proposed in (12) are rate-optimal over both $\mathcal{P}_\alpha(\eta, M)$ and $\mathcal{Q}_\alpha(\eta, M)$, the corresponding optimal choices of bandwidth are distinct. We generate precision matrices in the following way to guarantee that Ω is always in $\mathcal{P}_\alpha(\eta, M)$ but not in $\mathcal{Q}_\alpha(\eta, M)$ with some fixed η and M . Considering

$$\Omega = (I - A)^T D^{-1} (I - A), \quad A = [a_{ij}]_{p \times p}, \quad D = I_p,$$

where the first column of A is $a_{i1} = -2(i-1)^{-\alpha}$, $2 \leq i \leq p$. The remaining entries are all zeros. It is easy to check that $\Omega \in \mathcal{P}_\alpha(\eta, 2)$ with some large $\eta > 0$. The simulation is carried out with a similar range of values for p , n , α as in Section 6.1. Note that the consistent estimator exists only if $\alpha > 0.5$. Therefore, in this setting, the decay rate α varies among 1, 1.5 and 2.

The optimal choice of bandwidth of our local cropping estimator in $\mathcal{P}_\alpha(\eta, M)$ is $k \asymp n^{1/2\alpha}$, which is different from the one of crop.Q. We denote this rate-optimal estimator in $\mathcal{P}_\alpha(\eta, M)$ by crop.P. In the simulation, the bandwidth of crop.P is $\lfloor n^{1/2\alpha} \rfloor$. We also include BL estimator as a reference.

Table 1 reports the average errors of the three procedures, crop.P, crop.Q and BL, under the operator norm over 100 replications. The smallest errors in each experiment are highlighted in boldface. Fig. 6.2 plots the boxplots of their errors for $p = 500, 1000, 2000$.

Since Ω always belongs to $\mathcal{P}_\alpha(\eta, M)$ but not $\mathcal{Q}_\alpha(\eta, M)$, the estimator crop.Q is sub-optimal and thus expected to have an inferior performance. Table 2 shows this point, i.e., for fixed p and α , the advantage of crop.P is more obvious as n increases. Especially, crop.P outperforms the other two estimators when $n = 4000$. We also see a similar pattern as in Table 1 that p has minor effect on the errors of all the estimators.

Table 1: The average errors under the operator norm of the banding estimator (BL) and the local cropping estimator (crop.Q) over 100 replications.

p	n	$\alpha = 0.5$		$\alpha = 1$		$\alpha = 1.5$		$\alpha = 2$	
		crop.Q	BL	crop.Q	BL	crop.Q	BL	crop.Q	BL
500	500	4.68	5.44	1.64	2.38	1.18	1.16	0.93	0.81
	1000	3.29	4.89	1.17	1.72	0.82	1.08	0.66	0.69
	2000	2.47	4.45	0.89	1.33	0.59	0.69	0.48	0.59
	4000	1.84	3.80	0.62	1.07	0.41	0.64	0.34	0.53
1000	500	4.96	5.74	1.75	2.40	1.30	1.19	0.99	0.84
	1000	3.43	5.19	1.24	1.74	0.86	1.10	0.68	0.70
	2000	2.58	4.75	0.93	1.35	0.62	0.71	0.51	0.60
	4000	1.93	4.10	0.66	1.33	0.44	0.65	0.36	0.55
2000	500	5.14	5.97	1.85	2.41	1.33	1.21	1.06	0.89
	1000	3.58	5.41	1.30	1.76	0.90	1.12	0.72	0.71
	2000	2.69	4.97	0.98	1.37	0.65	0.73	0.54	0.62
	4000	2.01	4.32	0.69	1.34	0.45	0.66	0.38	0.55

Table 2: The average errors under the operator norm of the banding estimator (BL) and the local cropping estimator (crop.P & crop.Q) with two different bandwidths over 100 replications.

p	n	$\alpha = 1$			$\alpha = 1.5$			$\alpha = 2$		
		crop.P	crop.Q	BL	crop.P	crop.Q	BL	crop.P	crop.Q	BL
500	500	1.50	1.18	2.32	0.66	0.73	0.86	0.52	0.65	0.53
	1000	1.09	0.96	1.80	0.47	0.56	0.83	0.38	0.56	0.45
	2000	0.83	0.80	1.53	0.35	0.43	0.55	0.27	0.32	0.41
	4000	0.64	0.68	1.33	0.26	0.35	0.54	0.19	0.24	0.38
1000	500	1.50	1.20	2.36	0.68	0.74	0.91	0.57	0.68	0.59
	1000	1.12	0.98	1.82	0.49	0.58	0.81	0.39	0.55	0.46
	2000	0.84	0.81	1.54	0.37	0.44	0.55	0.27	0.32	0.41
	4000	0.65	0.68	1.52	0.26	0.35	0.53	0.19	0.24	0.38
2000	500	1.51	1.21	2.39	0.69	0.75	0.96	0.62	0.71	0.63
	1000	1.16	1.00	1.81	0.51	0.60	0.84	0.39	0.56	0.46
	2000	0.85	0.81	1.55	0.39	0.44	0.56	0.27	0.33	0.41
	4000	0.65	0.69	1.70	0.26	0.35	0.53	0.19	0.24	0.38




Figure 1: The boxplot of the errors from the local cropping estimator with the optimal bandwidth in $\mathcal{Q}_\alpha(\eta, M)$ (cropping.Q) and the banding estimator (BL) over 100 replications.

7 Appendix

Theorem 1 immediately follows from Theorems 3, 4, 5 and 6 while Theorem 2 immediately follows from Theorems 7 and 8. We provide key lemmas in this section in the proof of those main theorems.

7.1 Preliminary Lemmas: Lemma 21, Lemma 22, and Lemma 23

In this section, we provide three preliminary lemmas and their proofs, which are important in the proof of the other lemmas.




Figure 2: The boxplot of the errors from the local cropping estimator with the optimal bandwidth in $\mathcal{P}_\alpha(\eta, M)$ (cropping.P), the local cropping estimator with the optimal bandwidth in $\mathcal{Q}_\alpha(\eta, M)$ (cropping.Q) and the banding estimator (BL) over 100 replications.

7.1.1 Lemma 21 and the Proof

Lemma 21 *For any matrix A , such that $\eta^{-1} \leq \lambda_{\min}(A) \leq \lambda_{\max}(A) \leq \eta$. Operator $\mathbf{P}_\eta(\cdot)$ is defined in (9). we have,*

$$\|A - \mathbf{P}_\eta(S)\|_{\text{op}} \leq 2\|A - S\|_{\text{op}}, \quad (54)$$

$$\|A - \mathbf{P}_\eta(S)\|_{\text{F}} \leq 2\|A - S\|_{\text{F}}. \quad (55)$$

Proof. Since

$$\|A - \mathbf{P}_\eta(S)\|_* \leq \|S - \mathbf{P}_\eta(S)\|_* + \|A - S\|_*,$$

we only need to prove that

$$\|S - \mathbf{P}_\eta(S)\|_* \leq \|A - S\|_*,$$

where $\|\cdot\|_*$ is either the operator norm or Frobenius norm. Assume that the singular value decomposition of S is $U^T S V = W$. Since operator norm and Frobenius norm are invariant to the orthogonal transformation, it is sufficient to prove

$$\|U^T S V - U^T \mathbf{P}_\eta(S) V\|_* \leq \|U^T A V - U^T S V\|_*.$$

Here, $U^T S V$ and $U^T \mathbf{P}_\eta(S) V$ are the diagonal matrices. $\eta^{-1} \leq \lambda_{\min}(U^T A V) \leq \lambda_{\max}(U^T A V) \leq \eta$. Without loss of generality, we only need to prove that for any diagonal matrix $W \equiv \text{diag}(w_i)$,

$$\|W - W_\eta\|_* \leq \|A - W\|_*,$$

where $W_\eta = \text{diag}(\max\{\min\{w_i, \eta\}, \eta^{-1}\})$ and $\eta^{-1} \leq \lambda_{\min}(A) \leq \lambda_{\max}(A) \leq \eta$. The same result can be derived easily in the case of symmetric matrix S using similar argument on the eigen-decomposition. For the operator norm, Note the fact that

$$\|A - W\|_{\text{op}} \geq \|W\|_{\text{op}} - \|A\|_{\text{op}},$$

$$\|A - W\|_{\text{op}} \geq \frac{\|A^{-1} - W^{-1}\|_{\text{op}}}{\|W^{-1}\|_{\text{op}} \|A^{-1}\|_{\text{op}}} \geq \frac{\|W^{-1}\|_{\text{op}} - \|A^{-1}\|_{\text{op}}}{\|W^{-1}\|_{\text{op}} \|A^{-1}\|_{\text{op}}} \geq \|A^{-1}\|_{\text{op}}^{-1} - \|W^{-1}\|_{\text{op}}^{-1}.$$

Therefore, we have

$$\begin{aligned} \|A - W\|_{\text{op}} &\geq \max\{\|W\|_{\text{op}} - \|A\|_{\text{op}}, \|A^{-1}\|_{\text{op}}^{-1} - \|W^{-1}\|_{\text{op}}^{-1}, 0\} \\ &\geq \max\{w_i - \|A\|_{\text{op}}, \|A^{-1}\|_{\text{op}}^{-1} - w_i, 0\} \\ &\geq \max\{w_i - \eta, \eta^{-1} - w_i, 0\} \\ &= \max\{|w_i - \max\{\min\{w_i, \eta\}, \eta^{-1}\}|\} \\ &= \|W_\eta - W\|_{\text{op}}. \end{aligned}$$

The third inequality is because of the fact that $\|A\|_{\text{op}} \leq \eta$, $\|A^{-1}\|_{\text{op}}^{-1} \geq \eta^{-1}$.

For the Frobenius norm, with $A \equiv [a_{ij}]_{p \times p}$, we have

$$\begin{aligned} \|A - W\|_{\text{F}}^2 &= \sum_{i \neq j} a_{ij}^2 + \sum_i (a_{ii} - w_i)^2 \\ &= \sum_i (\sum_j a_{ij}^2 + w_i^2 - 2a_{ii}w_i) \\ &\geq \sum_i (\sum_j a_{ij}^2 + w_i^2 - 2\sqrt{\sum_j a_{ij}^2} w_i) \\ &= \sum_i (w_i - \sqrt{\sum_j a_{ij}^2})^2. \end{aligned}$$

Since $\eta^{-1} \leq \lambda_{\min}(A) \leq \lambda_{\max}(A) \leq \eta$, we obtain $\eta^{-1} \leq \sqrt{\sum_j a_{ij}^2} \leq \eta$. It is easy to check that for any i , $(\sqrt{\sum_j a_{ij}^2} - w_i)^2 \geq (\max\{\min\{w_i, \eta\}, \eta^{-1}\} - w_i)^2$. Then we have

$$\|A - W\|_F^2 \geq \|W_\eta - W\|_F^2,$$

that yields $\|A - W\|_F \geq \|W_\eta - W\|_F$. ■

7.1.2 Lemma 22 and the Proof

Lemma 22 *Assume that Ω satisfies that its magnitudes of all the eigenvalues are in the interval $[\eta^{-1}, \eta]$, let $\Sigma = \Omega^{-1}$ and $\Omega = (I - A)^T D^{-1} (I - A)$ (see Equation (1)), then,*

$$\|\Sigma\|_{\text{op}} \leq \eta, \text{ and } \|\Sigma^{-1}\|_{\text{op}} \leq \eta, \quad (56)$$

$$\|D\|_{\text{op}} \leq \eta, \text{ and } \|D^{-1}\|_{\text{op}} \leq \eta, \quad (57)$$

$$\|I - A\|_{\text{op}} \leq \eta, \text{ and } \|(I - A)^{-1}\|_{\text{op}} \leq \eta. \quad (58)$$

Furthermore, any sub-matrix of Ω , Σ , D and $I - A$ also share this property.

Proof. Since $\Sigma = \Omega^{-1}$, all the eigenvalues of Σ are in the interval $[\eta^{-1}, \eta]$. Let $\Omega \equiv [\omega_{ij}]_{p \times p}$, $\Sigma \equiv [\sigma_{ij}]_{p \times p}$, then $\forall i$, $\omega_{ii} \geq \lambda_{\min}(\Omega) \geq \eta^{-1}$; $\sigma_{ii} \leq \lambda_{\max}(\Sigma) \leq \eta$. ω_{ii} can be interpreted as the variance of the residual of the regression X_i and $\mathbf{X}_{-i} = (X_1 \dots X_{i-1}, X_{i+1} \dots X_p)^T$. σ_{ii} can be interpreted as the variance of the residual of the regression X_i and $\mathbf{0}$. The i -th entry of D , the variance of the residual of the regression X_i and $\mathbf{X}_{1:i-1} = (X_1 \dots X_{i-1})^T$, has the bound $\omega_{ii} \leq d_i \leq \sigma_{ii}$, then $\|D\|_{\text{op}} \leq \eta$ and $\|D^{-1}\|_{\text{op}} \leq \eta$.

And for any unit vector u, v such that $(I - A)u = \lambda v$, we have

$$u^T \Omega u = u^T (I - A)^T D^{-1} (I - A) u = \lambda^2 v^T D^{-1} v,$$

that is $\lambda^2 = (u^T \Omega u) / (v^T D^{-1} v) \in [\eta^{-2}, \eta^2]$. So,

$$\eta^{-1} \leq \lambda_{\min}(I - A) \leq \lambda_{\max}(I - A) \leq \eta.$$

For any sub-matrix Z^* of Z , Z^* can be represented by PZQ , where matrix P, Q are the diagonal matrices whose diagonals are 0 or 1. Since the singular values of matrix P and Q are 0 or 1, it is easy to show $\lambda_{\min}(Z) \leq \lambda_{\min}(Z^*) \leq \lambda_{\max}(Z^*) \leq \lambda_{\max}(Z)$. It implies that any sub-matrix of Ω , Σ and $(I - A)$ share that property. ■

7.1.3 Lemma 23 and the Proof

Lemma 23 *Let the projection of X_i on the linear span of $\mathbf{X}_{1:i-1}$ be*

$$\hat{X}_i = \mathbf{X}_{1:i-1}^T \boldsymbol{\beta}_i, \quad \epsilon_i = \text{Var}(X_i - \hat{X}_i).$$

Let the projection of X_i on the linear span of $\mathbf{X}_{i-k:i-1}$ be

$$\hat{X}_i^{[k]} = \mathbf{X}_{i-k:i-1}^T \boldsymbol{\gamma}'_i = \mathbf{X}_{1:i-1}^T \boldsymbol{\gamma}_i, \quad \delta_i = \text{Var}(X_i - \hat{X}_i^{[k]}),$$

where $\boldsymbol{\gamma}_i = (0, \boldsymbol{\gamma}'_i)^T$. If $\boldsymbol{\beta}_i = (\beta_{i1}, \dots, \beta_{i(i-1)})^T$ satisfies

$$\sum_{j < i-k} |\beta_{ij}| < Mk^{-\alpha}, \quad \forall 1 \leq k \leq i-1, \quad (59)$$

then,

$$\|\boldsymbol{\beta}_i - \boldsymbol{\gamma}_i\|_{\text{op}} \leq 2\eta^2 Mk^{-\alpha}, \quad (60)$$

$$|\epsilon_i - \delta_i| \leq 4\eta^4 Mk^{-\alpha}. \quad (61)$$

If $\boldsymbol{\beta}_i = (\beta_{i1}, \dots, \beta_{i(i-1)})^T$ satisfies

$$|\beta_{ij}| < M(i-j)^{-\alpha-1}, \quad \forall 1 \leq j \leq i-1 \quad (62)$$

then,

$$\|\boldsymbol{\beta}_i - \boldsymbol{\gamma}_i\|_{\text{op}} \leq 2\eta^2 M(2\alpha+2)^{\frac{1}{2}} k^{-\alpha-1/2}, \quad (63)$$

$$|\epsilon_i - \delta_i| \leq 4\eta^4 M(2\alpha+2)^{\frac{1}{2}} k^{-\alpha-1/2}. \quad (64)$$

Proof. To facilitate the proof, we divide Σ_i into several block matrix,

$$\Sigma_i = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} & \Sigma_{13} \\ \Sigma_{21} & \Sigma_{22} & \Sigma_{23} \\ \Sigma_{31} & \Sigma_{32} & \Sigma_{33} \end{bmatrix},$$

where $\Sigma_{11}, \Sigma_{22}, \Sigma_{33}$ are the covariance matrices of $\mathbf{X}_{1:i-k-1}, \mathbf{X}_{i-k:i-1}, X_i$.

The following statement can be proved by simple linear algebra.

In regression $X_i \sim \mathbf{X}_{1:i-1}$,

$$\hat{X}_i = \mathbf{X}_{1:i-1}^T \boldsymbol{\beta}_i = \mathbf{X}_{1:i-k-1}^T \boldsymbol{\beta}_{i[-k]} + \mathbf{X}_{i-k:i-1}^T \boldsymbol{\beta}_{i[k]}, \quad \epsilon_i = \text{Var}(X_i - \hat{X}_i).$$

We have,

$$\begin{aligned} \Sigma_{23} &= \Sigma_{21} \boldsymbol{\beta}_{i[-k]} + \Sigma_{22} \boldsymbol{\beta}_{i[k]}, \\ \epsilon_i &= (-\boldsymbol{\beta}_{i[-k]}^T, -\boldsymbol{\beta}_{i[k]}^T, 1) \Sigma_i (-\boldsymbol{\beta}_{i[-k]}, -\boldsymbol{\beta}_{i[k]}, 1)^T. \end{aligned}$$

In regression $X_i \sim \mathbf{X}_{i-k:i-1}$,

$$\hat{X}_i^{[k]} = \mathbf{X}_{i-k:i-1}^T \boldsymbol{\gamma}_i = \mathbf{X}_{i-k:i-1}^T \boldsymbol{\gamma}'_i, \quad \delta_i = \text{Var}(X_i - \hat{X}_i^{[k]}).$$

So,

$$\begin{aligned} \Sigma_{23} &= \Sigma_{22} \boldsymbol{\gamma}'_i, \\ \delta_i &= (\mathbf{0}^T, -\boldsymbol{\gamma}'_i{}^T, 1) \Sigma_i (\mathbf{0}, -\boldsymbol{\gamma}'_i, 1)^T. \end{aligned}$$

Equation (59) implies

$$\|\beta_{i[-k]}\|_{\text{op}} \leq \|\beta_{i[-k]}\|_1 \leq Mk^{-\alpha}.$$

Then we have,

$$\begin{aligned} \|\gamma_i - \beta_i\|_{\text{op}} &\leq \|\mathbf{0} - \beta_{i[-k]}\|_{\text{op}} + \|\gamma'_i - \beta_{i[k]}\|_{\text{op}} \\ &\leq \|\beta_{i[-k]}\|_{\text{op}} + \|\Sigma_{22}^{-1}\Sigma_{21}\beta_{i[-k]}\|_{\text{op}} \\ &\leq Mk^{-\alpha} + \eta^2 Mk^{-\alpha} \\ &\leq 2\eta^2 Mk^{-\alpha}. \end{aligned}$$

We finish the proof of Equation (60).

Σ_i is the sub-matrix of Σ , Lemma 22 implies that $\|\Sigma_i\|_{\text{op}} \leq \eta$. It is easy to verify that $\|(-\beta_i, 1)\|_{\text{op}} \leq \|I - A\|_{\text{op}} \leq \eta$. Let Σ' be the covariance matrix of $\mathbf{X}_{i-k:i-1}$, Σ' is the sub-matrix of Σ , then $\|\Sigma'^{-1}\|_{\text{op}} \leq \eta$. Let $\Omega' = (\Sigma')^{-1} = (I - B)^T E^{-1} (I - B)$, so $\|I - B\|_{\text{op}} \leq \eta$ by Lemma 22, then $\|(-\gamma_i, 1)\|_{\text{op}} \leq \|I - B\|_{\text{op}} \leq \eta$.

$$\begin{aligned} |\epsilon_i - \delta_i| &\leq |(-\beta_i^T, 1)\Sigma_i(-\beta_i, 1)^T - (-\gamma_i^T, 1)\Sigma_i(-\gamma_i, 1)^T| \\ &\leq |(-\beta_i^T + \gamma_i^T, 0)\Sigma_i(-\beta_i, 1)^T| + |(-\gamma_i^T, 1)\Sigma_i(\beta_i - \gamma_i, 0)^T| \\ &\leq 4\eta^4 Mk^{-\alpha}. \end{aligned}$$

We finish the proof of Equation (61). Similarly, Equation (62) implies

$$\|\beta_{i[-k]}\|_{\text{op}} \leq M(2\alpha + 2)^{1/2} k^{-\alpha-1/2}.$$

Then,

$$\begin{aligned} \|\gamma_i - \beta_i\|_{\text{op}} &\leq \|\beta_{i[-k]}\|_{\text{op}} + \|\Sigma_{22}^{-1}\Sigma_{21}\|_{\text{op}} \|\beta_{i[-k]}\|_{\text{op}} \\ &\leq 2\eta^2 M(2\alpha + 2)^{1/2} k^{-\alpha-1/2}. \end{aligned}$$

Again, we can show $\|\Sigma_i\|_{\text{op}} \leq \eta$, $\|(-\beta_i, 1)\|_{\text{op}} \leq \eta$ and $\|(-\gamma_i, 1)\|_{\text{op}} \leq \eta$.

$$\begin{aligned} |\epsilon_i - \delta_i| &\leq |(-\beta_i^T, 1)\Sigma_i(-\beta_i, 1)^T - (-\gamma_i^T, 1)\Sigma_i(-\gamma_i, 1)^T| \\ &\leq |(-\beta_i^T + \gamma_i^T, 0)\Sigma_i(-\beta_i, 1)^T| + |(-\gamma_i^T, 1)\Sigma_i(\beta_i - \gamma_i, 0)^T| \\ &\leq 4\eta^4 M(2\alpha + 2)^{1/2} k^{-\alpha-1/2}. \end{aligned}$$

We finish the proof of Equation (63) and (64).

7.2 Proof of Lemma 2 and Lemma 4 in analysis of Theorem 3

In this section, we prove Lemma 2 and Lemma 4 to establish Theorem 3.

7.2.1 Proof of Lemma 2

For $\Omega \equiv [\omega_{ij}]_{p \times p}$, define

$$bd_k(\Omega) = [\omega_{ij} \mathbf{1}(|i - j| \leq k)]_{p \times p}. \quad (65)$$

We have

$$\begin{aligned} \Omega_k^* &= \frac{1}{k} \sum_{i=k}^{2k} bd_i(\Omega), \\ \|\Omega - \Omega_k^*\|_{\text{op}} &\leq \frac{1}{k} \sum_{i=k}^{2k} (\|\Omega - bd_i(\Omega)\|_{\text{op}}). \end{aligned}$$

First, we prove

$$\|\Omega - bd_k(\Omega)\|_{\text{op}} \leq Ck^{\frac{1}{2}-\alpha}.$$

Define $D^{-\frac{1}{2}}(I - A)$ as $B = [b_{ij}]$, then $\Omega = B^T B$, and

$$\max_i \sum_{j < i-k} |b_{ij}| \leq M\eta^{\frac{1}{2}} k^{-\alpha}.$$

Set $bd_k(B)$ as B_k , and $B - bd_k(B)$ as B_{-k} . Then

$$\begin{aligned} \Omega &= B^T B \\ &= (B_k + B_{-k})^T (B_k + B_{-k}) \\ &= B_k^T B_k + B_{-k}^T B_k + B_k^T B_{-k} + B_{-k}^T B_{-k}. \end{aligned}$$

One can check each entry of $(B_k)^T B_k$ to prove

$$bd_k((B_k)^T B_k) = (B_k)^T B_k.$$

$$\begin{aligned} \|\Omega - bd_k(\Omega)\|_{\text{op}} &= \|B^T B - bd_k(B^T B)\|_{\text{op}} \\ &= \|B_{-k}^T B_k + B_k^T B_{-k} + B_{-k}^T B_{-k} - bd_k(B_{-k}^T B_k + B_k^T B_{-k} + B_{-k}^T B_{-k})\|_{\text{op}} \\ &\leq \|B_{-k}^T B_k + B_k^T B_{-k} + B_{-k}^T B_{-k}\|_{\text{op}} + \|bd_k(B_{-k}^T B_k + B_k^T B_{-k} + B_{-k}^T B_{-k})\|_{\text{op}} \\ &= \|B_{-k}^T B + B^T B_{-k} - B_{-k}^T B_{-k}\|_{\text{op}} + \|bd_k(B_{-k}^T B_k + B_k^T B_{-k} + B_{-k}^T B_{-k})\|_{\text{op}} \\ &\leq \|B_{-k}^T B\|_{\text{op}} + \|B^T B_{-k}\|_{\text{op}} + \|B_{-k}^T B_{-k}\|_{\text{op}} \\ &\quad + \|bd_k(B_{-k}^T B_k)\|_{\text{op}} + \|bd_k(B_k^T B_{-k})\|_{\text{op}} + \|bd_k(B_{-k}^T B_{-k})\|_{\text{op}}. \end{aligned}$$

We need to control $\|B_{-k}\|_{\text{op}}$. Here we can even control the operator norm of the matrix whose entry is the absolute value of the entry in B_{-k} .

Lemma 24

$$\|B_{-k}^+\|_{\text{op}} \leq Ck^{\frac{1}{2}-\alpha}$$

where X^+ is the matrix whose entry is the magnitude of the corresponding entry in X .

Proof. Assume D_i is the matrix composed by the $(2^{i-1}k + 1)th$ to $(2^i k)th$ sub-diagonals in matrix B^+ . Let D_0 be the the matrix composed by the $1st$ to kth sub-diagonals in matrix B^+ . Then

$$B_k^+ = D_0, B_{-k}^+ = \sum_{i=1}^{\log_2(p/k)} (D_i).$$

For any $i \geq 1$,

$$\|D_i\|_\infty \leq M\eta^{\frac{1}{2}}(2^{i-1}k)^{-\alpha}.$$

Since there are $2^{i-1}k$ entries in one column of D_i , then

$$\|D_i\|_1 \leq M\eta^{\frac{1}{2}}(2^{i-1}k)^{1-\alpha}.$$

$$\begin{aligned} \|D_i\|_{\text{op}} &\leq (\|D_i\|_\infty \|D_i\|_1)^{\frac{1}{2}} \\ &\leq M\eta^{\frac{1}{2}}(2^{i-1}k)^{\frac{1}{2}-\alpha} \\ &= M\eta^{\frac{1}{2}}k^{\frac{1}{2}-\alpha} \times 2^{(i-1)(\frac{1}{2}-\alpha)} \\ &\leq M\eta^{\frac{1}{2}}k^{\frac{1}{2}-\alpha} \times (2^{\frac{1}{2}-\alpha})^{i-1}. \end{aligned}$$

Then,

$$\begin{aligned} \|B_{-k}^+\|_{\text{op}} &\leq \sum_{i=1}^{\log_2(p/k)} \|D_i\|_{\text{op}} \\ &\leq \sum_{i=1}^{\log_2(p/k)} (M\eta^{\frac{1}{2}}k^{\frac{1}{2}-\alpha} \times (2^{\frac{1}{2}-\alpha})^{i-1}) \\ &= M\eta^{\frac{1}{2}}k^{\frac{1}{2}-\alpha} \times \sum_{i=1}^{\log_2(p/k)} (2^{\frac{1}{2}-\alpha})^{i-1} \\ &\leq CM\eta^{\frac{1}{2}}k^{\frac{1}{2}-\alpha}. \end{aligned}$$

■

Lemma 25 *Let X be a matrix, then*

$$\|X\|_{\text{op}} \leq \|X^+\|_{\text{op}},$$

and

$$\|bd_k(X)\|_{\text{op}} \leq \|X^+\|_{\text{op}}.$$

Proof. Assume $\|X\|_{\text{op}} = u^T X v$, then

$$\|X\|_{\text{op}} = u^T X v = \sum_{i,j} u_i x_{ij} v_j \leq \sum_{i,j} |u_i| |x_{ij}| |v_j| = (u^+)^T X^+ v^+ \leq \|X^+\|_{\text{op}}.$$

Assume $bd_k(X) = [x_{ij}^*]$, $\|bd_k(X)\|_{\text{op}} = u^T (bd_k(X)) v$, then

$$\|bd_k(X)\|_{\text{op}} = u^T (bd_k(X)) v = \sum_{i,j} u_i x_{ij}^* v_j \leq \sum_{i,j} |u_i| |x_{ij}| |v_j| = (u^+)^T X^+ v^+ \leq \|X^+\|_{\text{op}}.$$

■

So,

$$\|B_{-k}\|_{\text{op}} \leq \|B_{-k}^+\|_{\text{op}} \leq Ck^{\frac{1}{2}-\alpha}.$$

We have

$$\begin{aligned} \|B^T B_{-k}\|_{\text{op}} &\leq \|B\|_{\text{op}} \|B_{-k}\|_{\text{op}} \leq Ck^{\frac{1}{2}-\alpha}, \\ \|B_{-k}^T B\|_{\text{op}} &\leq \|B\|_{\text{op}} \|B_{-k}\|_{\text{op}} \leq Ck^{\frac{1}{2}-\alpha}, \\ \|B_{-k}^T B_{-k}\|_{\text{op}} &\leq \|B_{-k}\|_{\text{op}} \|B_{-k}\|_{\text{op}} \leq Ck^{1-2\alpha}, \end{aligned}$$

$$\|bd_k(B_{-k}^T B_{-k})\|_{\text{op}} \leq \|(B_{-k}^T B_{-k})^+\|_{\text{op}} \leq \|(B_{-k}^+)^T B_{-k}^+\|_{\text{op}} \leq \|B_{-k}\|_{\text{op}} \|B_{-k}\|_{\text{op}} \leq Ck^{1-2\alpha}.$$

Then we turn to bound $\|bd_k(B_{-k}^T B_k)\|_{\text{op}} + \|bd_k(B_k^T B_{-k})\|_{\text{op}}$. We need to control $\|bd_k(B_{-k}^T B_k)\|_{\infty}$ and $\|bd_k(B_{-k}^T B_k)\|_1$.

For the h -th row of $bd_k(B_{-k}^T B_k)$,

$$\begin{aligned} \|\text{row}_h(bd_k(B_{-k}^T B_k))\|_1 &\leq \sum_{i=h+1}^{h+k} \left(\sum_{j=h+k+1}^{h+2k} |b_{jh}| |b_{ij}| \right) \\ &\leq \sum_{j=h+k+1}^{h+2k} \left(\sum_{i=h+1}^{h+k} |b_{jh}| |b_{ij}| \right) \\ &\leq \sum_{j=h+k+1}^{h+2k} M^2 \eta ((j-h)^{-\alpha} (j-h-k)^{-\alpha}) \\ &\leq \sum_{j=1}^k M^2 \eta ((j+k)^{-\alpha} j^{-\alpha}) \\ &\leq M^2 \eta k^{-\alpha} \sum_{j=1}^k j^{-\alpha} \\ &\leq Ck^{1-2\alpha}. \end{aligned}$$

So $\|bd_k(B_{-k}^T B_k)\|_{\infty} \leq Ck^{1-2\alpha}$.

For the h -th column of $bd_k(B_{-k}^T B_k)$,

$$\begin{aligned}
\|col_h(bd_k(B_{-k}^T B_k))\|_1 &\leq \sum_{i=h-k}^{h-1} \left(\sum_{j=h+1}^{h+k} |b_{ji}| |b_{jh}| \right) \\
&\leq \sum_{j=h+1}^{h+k} \left(\sum_{i=h-k}^{h-1} |b_{ji}| |b_{jh}| \right) \\
&\leq \sum_{j=h+1}^{h+k} M^2 \eta(k^{-\alpha} (j-h)^{-\alpha}) \\
&\leq \sum_{j=1}^k M^2 \eta(k^{-\alpha} j^{-\alpha}) \\
&\leq Ck^{1-2\alpha}.
\end{aligned}$$

So $\|bd_k(B_{-k}^T B_k)\|_1 \leq Ck^{1-2\alpha}$.

It is clear that

$$\begin{aligned}
\|bd_k(B_{-k}^T B_k)\|_{\text{op}} &\leq (\|bd_k(B_{-k}^T B_k)\|_{\infty} \|bd_k(B_{-k}^T B_k)\|_1) \leq Ck^{1-2\alpha}, \\
\|bd_k(B_k^T B_{-k})\|_{\text{op}} &\leq (\|bd_k(B_{-k}^T B_k)\|_{\infty} \|bd_k(B_{-k}^T B_k)\|_1) \leq Ck^{1-2\alpha}.
\end{aligned}$$

Combining the equations above, we prove that

$$\|\Omega - bd_k(\Omega)\|_{\text{op}} \leq Ck^{\frac{1}{2}-\alpha}.$$

So,

$$\begin{aligned}
\|\Omega - \Omega_k^*\|_{\text{op}} &\leq Ck^{\frac{1}{2}-\alpha}, \\
\|\Omega - \Omega_k^*\|_{\text{op}}^2 &\leq Ck^{1-2\alpha}.
\end{aligned}$$

7.2.2 Proof of Lemma 4

Define the $p \times q$ matrix

$$R_{p \times q}^{a,b,k} = [r_{ij}]_{p \times q}, \quad r_{ij} = \mathbf{1} (i - a = j - b, a \leq i < a + k, b \leq j < b + k). \quad (66)$$

Note that for a $m \times n$ matrix T , $R_{p \times m}^{a,b,c} T R_{n \times q}^{c,d,f}$ means that we select the sub-matrix of T with its $b : (b + c - 1)$ rows and $d : (d + f - 1)$ columns, then put it onto the entry with the index (a, c) of a $p \times q$ zero matrix. Let $\Phi_m^{(k)} = \mathbf{E}_m^p(\mathbf{C}_m^k(\Omega))$. Note that $\Phi_m^{(k)}$ is a p by p matrix. Then one can check

$$\begin{aligned}
\Phi_m^{(k)} &= R_{p \times p}^{m,m,k} \Omega R_{p \times p}^{m,m,k} \\
&= R_{p \times p}^{m,m,k} (I - A)^T D^{-1} (I - A) R_{p \times p}^{m,m,k}.
\end{aligned}$$

Define

$$\begin{aligned}
\Phi_m^{(k)'} &= R_{p \times p}^{m,m,k} (I - A)^T R_{p \times p}^{m,m,2k} D^{-1} R_{p \times p}^{m,m,2k} (I - A) R_{p \times p}^{m,m,k} \\
&= \underbrace{R_{p \times p}^{m,m,k} (I - A)^T R_{p \times p}^{m,m,2k}}_{H^T} \underbrace{R_{p \times p}^{m,m,2k} D^{-1} R_{p \times p}^{m,m,2k}}_{D'^{-1}} \underbrace{R_{p \times p}^{m,m,2k} (I - A) R_{p \times p}^{m,m,k}}_H \\
&= H^T D'^{-1} H.
\end{aligned}$$

Define t_i is the i -th row of $(I - A)R_{p \times p}^{m,m,k}$.

Since $I - A$ is the lower triangular matrix, then

$$(I - A)R_{p \times p}^{m,m,k} = (0, 0, \dots, 0, t_m, \dots, t_p)^T.$$

$R_{p \times p}^{m,m,2k} (I - A)R_{p \times p}^{m,m,k}$ keeps the m -th to $(m + 2k - 1)$ -th rows in $(I - A)R_{p \times p}^{m,m,k}$. Then,

$$(I - A)R_{p \times p}^{m,m,k} - R_{p \times p}^{m,m,2k} (I - A)R_{p \times p}^{m,m,k} = (0, 0, \dots, 0, t_{m+2k}, \dots, t_p)^T.$$

and for $m + 2k \leq i \leq p$,

$$\|t_i\|_{\text{op}} \leq \|t_i\|_1 \leq M(i - m - k + 1)^{-\alpha}. \quad (67)$$

So,

$$\begin{aligned}
&\|(I - A)R_{p \times p}^{m,m,k} - R_{p \times p}^{m,m,2k} (I - A)R_{p \times p}^{m,m,k}\|_{\text{op}}^2 \\
&\leq \|(\|t_{m+2k}\|_{\text{op}}, \dots, \|t_p\|_{\text{op}})^T\|_{\text{op}}^2 \\
&\leq \|(M(k + 1)^{-\alpha}, \dots, M(p - m - k)^{-\alpha})^T\|_{\text{op}}^2 \\
&\leq 2\alpha M^2 k^{-2\alpha+1}.
\end{aligned}$$

Since both $(I - A)R_{p \times p}^{m,m,k}$ and $R_{p \times p}^{m,m,2k} (I - A)R_{p \times p}^{m,m,k}$ are the sub-matrices of $I - A$, their operator norm can be bounded by η , then

$$\begin{aligned}
\|\Phi_m^{(k)} - \Phi_m^{(k)'}\|_{\text{op}}^2 &\leq \|R_{p \times p}^{m,m,k} (I - A)^T D^{-1} (I - A) R_{p \times p}^{m,m,k} \\
&\quad - R_{p \times p}^{m,m,k} (I - A)^T R_{p \times p}^{m,m,2k} D^{-1} R_{p \times p}^{m,m,2k} (I - A) R_{p \times p}^{m,m,k}\|_2^2 \\
&\leq \|(I - A)R_{p \times p}^{m,m,k} - R_{p \times p}^{m,m,2k} (I - A)R_{p \times p}^{m,m,k}\|_{\text{op}}^2 \|D^{-1}\|_{\text{op}}^2 \\
&\quad \times (\|R_{p \times p}^{m,m,k} (I - A)^T R_{p \times p}^{m,m,2k}\|_{\text{op}}^2 + \|(I - A)R_{p \times p}^{m,m,k}\|_{\text{op}}^2) \\
&\leq 2\eta^2 \|(I - A)R_{p \times p}^{m,m,k} - R_{p \times p}^{m,m,2k} (I - A)R_{p \times p}^{m,m,k}\|_{\text{op}}^2 \\
&\leq 4\alpha\eta^2 M^2 k^{-2\alpha+1}. \quad (68)
\end{aligned}$$

Define $\Psi_m^{(k)} = \mathbf{E}_m^p \left(\mathbf{C}_{k+1}^k (\mathbf{C}_{3k}^m (\Omega^{-1})^{-1}) \right)$. Note that $\Psi_m^{(k)}$ is a p by p matrix. One can check that

$$\begin{aligned}
\Psi_m^{(k)} &= R_{p \times 3k}^{m,k+1,k} \mathbf{C}_{3k}^m (\Omega^{-1})^{-1} R_{3k \times p}^{k+1,m,k} \\
&= R_{p \times 3k}^{m,k+1,k} (I - G)^T F^{-1} (I - G) R_{3k \times p}^{k+1,m,k} \\
&= \underbrace{R_{p \times 3k}^{m,k+1,k} (I - G)^T R_{3k \times p}^{k+1,m,2k}}_{K^T} \underbrace{R_{p \times 3k}^{m,k+1,2k} F^{-1} R_{3k \times p}^{k+1,m,2k}}_{F'^{-1}} \underbrace{R_{p \times 3k}^{m,k+1,2k} (I - G) R_{3k \times p}^{k+1,m,k}}_K \\
&= K^T F'^{-1} K.
\end{aligned}$$

For any fixed i , the i -th row of H is part of the coefficients of the regression $X_{m+i} \sim \mathbf{X}_{1:m+i-1}$ and the i -th row of K is part of the coefficients of the regression $X_{m+i} \sim \mathbf{X}_{m-k:k+i}$. According to Lemma 23, the operator norm of any row of $(H - K)$ can be bounded by $2M\eta^2(k+i)^{-\alpha}$. The dimension of the non-zero part of $H - K$ is $k \times 2k$. So we have

$$\|H - K\|_{\text{op}}^2 \leq 8\eta^4 M^2 k^{-2\alpha+1}.$$

Similarly, the i -th elements of D' and F' is the residual of the above two regressions. According to Lemma 23, $\|D' - F'\|_{\text{op}}^2 \leq 16\eta^8 M^2 k^{-2\alpha}$. So,

$$\|D'^{-1} - F'^{-1}\|_{\text{op}}^2 \leq \|D'^{-1}\|_{\text{op}}^2 \|D' - F'\|_{\text{op}}^2 \|F'^{-1}\|_{\text{op}}^2 \leq 16M^2 \eta^{12} k^{-2\alpha}.$$

Then by Lemma 22, the operator norms of $I - A$, D^{-1} , $I - G$ and F^{-1} can be bounded by η . So the operator norm of their sub-matrices H , D'^{-1} , K and F'^{-1} can also be bounded by η . Then we have

$$\begin{aligned} & \|\Phi_m^{(k)'} - \Psi_m^{(k)}\|_{\text{op}}^2 \\ & \leq \|H^T D'^{-1} H - K^T F'^{-1} K\|_{\text{op}}^2 \\ & \leq 3(\|D'^{-1} H\|_{\text{op}}^2 + \|F'^{-1} K\|_{\text{op}}^2) \|H - K\|_{\text{op}}^2 + 3\|K\|_{\text{op}}^2 \|H\|_{\text{op}}^2 \|D'^{-1} - F'^{-1}\|_{\text{op}}^2 \quad (69) \\ & \leq 6\eta^4 \times 8\eta^4 M^2 k^{-2\alpha+1} + 3\eta^4 \times 16M^2 \eta^{12} k^{-2\alpha} \\ & \leq 96M^2 \eta^{16} k^{-2\alpha+1}. \end{aligned}$$

Based on the Equation (68) and (69), we have

$$\begin{aligned} & \|\Phi_m^{(k)} - \Psi_m^{(k)}\|_{\text{op}}^2 \\ & \leq 2\|\Phi_m^{(k)} - \Phi_m^{(k)'}\|_{\text{op}}^2 + 2\|\Phi_m^{(k)'} - \Psi_m^{(k)}\|_{\text{op}}^2 \\ & \leq 2 \times (4\alpha\eta^2 M^2 k^{-2\alpha+1} + 96M^2 \eta^{16} k^{-2\alpha+1}) \\ & \leq 400\alpha M^2 \eta^{16} k^{-2\alpha+1}. \end{aligned}$$

Remember that $\Phi_m^{(k)} = \mathbf{E}_m^p(\mathbf{C}_m^k(\Omega))$ and $\Psi_m^{(k)} = \mathbf{E}_m^p(\mathbf{C}_{k+1}^k(\mathbf{C}_{3k}^m(\Omega^{-1})^{-1}))$. Then

$$\|\mathbf{C}_{k+1}^k(\mathbf{C}_{3k}^m(\Omega^{-1})^{-1}) - \mathbf{C}_m^k(\Omega)\|_{\text{op}} \leq 400\alpha M^2 \eta^{16} k^{-2\alpha+1}.$$

So we finish the proof of Lemma 4. ■

7.3 Proof of Lemma 5, Lemma 7, Lemma 8, and Lemma 10 in the analysis of Theorem 4

In this section, we prove Lemma 5, Lemma 7, Lemma 8, and Lemma 10 to establish Theorem 4.

7.3.1 Proof of Lemma 5

First we prove that $\mathcal{P}_1 \in \mathcal{P}_\alpha(\eta, M)$. Let $A(\theta) \equiv a_{ij}$. We know the exact value of each entry. It is easy to check $\sum_{i-j > k} |a_{ij}| \leq Mk^{-\alpha}$. One can check $\Omega(\theta) \in \mathcal{P}_1$ has the specific form of

$$\Omega(\theta) = \begin{bmatrix} I_k + (A_k^*(\theta))^T A_k^*(\theta) & -(A_k^*(\theta))^T & 0_{k \times (p-2k)} \\ -A_k^*(\theta) & I_k & 0_{k \times (p-2k)} \\ 0_{(p-2k) \times k} & 0_{(p-2k) \times k} & I_{p-2k} \end{bmatrix}.$$

Let $\mathbf{1}$ denote the vector with all 1's in Θ . We find

$$\begin{aligned} \lambda_{\max}(\Omega(\theta)) &\leq \lambda_{\max}(\Omega(\mathbf{1})) = (1 + \lambda_{\max}(A_k^*(\mathbf{1})))^2 = (1 + \sqrt{k/n}\tau)^2 \leq (1 + \tau)^2 \leq \eta; \\ \lambda_{\min}(\Omega(\theta)) &\geq \lambda_{\min}(\Omega(\mathbf{1})) = (1 + \lambda_{\max}(A_k^*(\mathbf{1})))^{-2} = (1 + \sqrt{k/n}\tau)^{-2} \geq (1 + \tau)^{-2} \geq \eta^{-1}. \end{aligned}$$

The eigenvalues of $\Omega(\theta)$ are in the interval $[\eta^{-1}, \eta]$. So $\mathcal{P}_1 \in \mathcal{P}_\alpha(\eta, M)$.

Then we turn to prove that $\mathcal{P}_2 \in \mathcal{P}_\alpha(\eta, M)$. The Cholesky factor A of $\Omega(m)$ is the identity matrix. The minimum eigenvalue of $\Omega(m)$ is $(1 + \tau a^{\frac{1}{2}})^{-1}$, which is greater than η^{-1} and maximum one is 1, which is less than η . So $\mathcal{P}_2 \in \mathcal{P}_\alpha(\eta, M)$.

7.3.2 Proof of Lemma 7

Since $\|P_\theta \wedge P_{\theta'}\| = 1 - \frac{1}{2}\|P_\theta - P_{\theta'}\|_1$, we turn to control $\max_{H(\theta, \theta')=1} \|P_\theta - P_{\theta'}\|_1$. According to First Pinsker's inequality in [17],

Lemma 26 (First Pinsker's Inequality)

$$\begin{aligned} \|P_\theta - P_{\theta'}\|_1^2 &\leq \frac{1}{2}K(P_{\theta'}|P_\theta) \\ &= \frac{n}{2} \left[\frac{1}{2} \text{tr}(\Sigma(\theta')\Sigma(\theta)^{-1}) - \frac{1}{2} \log \det(\Sigma(\theta')\Sigma(\theta)^{-1}) - \frac{p}{2} \right] \end{aligned}$$

where $K(\cdot|\cdot)$ is the Kullback-Leibler divergence.

One can check $\Sigma(\theta) \in \mathcal{P}_1$ has the form of

$$\begin{aligned} \Sigma(\theta) &= (I + A(\theta))(I + A(\theta)^T) \\ &= \begin{bmatrix} I_k & (A_k^*(\theta))^T & 0_{k \times (p-2k)} \\ (A_k^*(\theta)) & I_k + (A_k^*(\theta))(A_k^*(\theta))^T & 0_{k \times (p-2k)} \\ 0_{(p-2k) \times k} & 0_{(p-2k) \times k} & I_{p-2k} \end{bmatrix}. \end{aligned}$$

Assume that $\Sigma(\theta') = D + \Sigma(\theta)$, and let $d = \theta' - \theta$. Note that $\max H(\theta, \theta') = 1$, then D has the form of

$$\begin{aligned} D &= (A(\theta') - A(\theta)) + (A(\theta') - A(\theta))^T + (A(\theta')A(\theta')^T - A(\theta)A(\theta)^T) \\ &= A(d) + A(d)^T + A(\theta')A(d)^T + A(d)A(\theta)^T. \end{aligned}$$

Let $\mathbf{1}$ denote the all-ones vector, then we have

$$\|D\|_{\text{op}} \leq 2(1 + \|A_k^*(\mathbf{1})\|_{\text{op}})\|A_k^*(d)\|_{\text{op}} \leq 4\tau n^{-\frac{1}{2}}, \quad (70)$$

$$\|D\|_{\text{F}} \leq 2(1 + \|A_k^*(\mathbf{1})\|_{\text{F}})\|A_k^*(d)\|_{\text{F}} \leq 4\tau n^{-\frac{1}{2}}. \quad (71)$$

Furthermore, we have $\|D\Sigma(\theta)^{-1}\|_{\text{op}}^2 \leq n^{-\frac{1}{2}}$, $\|D\Sigma(\theta)^{-1}\|_{\text{F}}^2 \leq \|\Sigma(\theta)^{-1}\|_{\text{op}}^2\|D\|_{\text{F}}^2 \leq n^{-\frac{1}{2}}$. Since $\Sigma(\theta')\Sigma(\theta)^{-1} = I + D\Sigma(\theta)^{-1}$, $\frac{1}{2}\text{tr}(I + D\Sigma(\theta)^{-1}) = \frac{p}{2} + \frac{1}{2}\text{tr}(D\Sigma(\theta)^{-1})$, $\log \det(I + D\Sigma(\theta)^{-1}) = \text{tr}(D\Sigma(\theta)^{-1}) + \sum_i (\log(1 + \lambda_i) - \lambda_i)$, where λ_i 's are the eigenvalues of $D\Sigma(\theta)^{-1}$. Then applying the First Pinsker's inequality in this case, we have

$$\begin{aligned} \|P_\theta - P_{\theta'}\|_1^2 &\leq \frac{n}{2} \left[\frac{1}{2}\text{tr}(\Sigma(\theta')\Sigma(\theta)^{-1}) - \frac{1}{2}\log \det(\Sigma(\theta')\Sigma(\theta)^{-1}) - \frac{p}{2} \right] \\ &= \frac{n}{2} \left[\frac{1}{2}\text{tr}(I + D\Sigma(\theta)^{-1}) - \frac{1}{2}\log \det(I + D\Sigma(\theta)^{-1}) - \frac{p}{2} \right] \\ &= \frac{n}{2} \sum_i (\lambda_i - \log(1 + \lambda_i)). \end{aligned}$$

$|\lambda_i| \leq \|D\Sigma(\theta)^{-1}\|_{\text{op}}$. So all the λ_i are bounded by $\pm n^{-\frac{1}{2}}$. By Taylor expansion, we have $\sum_i (\lambda_i - \log(1 + \lambda_i)) \leq 2 \sum_i \lambda_i^2 = 2\|D\Sigma(\theta)^{-1}\|_{\text{F}}^2$. We have

$$\|P_\theta - P_{\theta'}\|_1^2 \leq n\|D\Sigma(\theta)^{-1}\|_{\text{F}}^2 \leq 1,$$

then $\|P_\theta \wedge P_{\theta'}\| \geq 0.5$.

7.3.3 Proof of Lemma 8

The proof is as follows,

$$\begin{aligned} &\|\Omega(\theta') - \Omega(\theta)\|_{\text{op}}^2 \\ &\geq \sup_{\|u\|_2=1, \|v\|_2=1} (u^T, 0, 0) \begin{bmatrix} A_k^*(\theta)A_k^*(\theta)^T - A_k^*(\theta')A_k^*(\theta')^T & A_k^*(\theta')^T - A_k^*(\theta)^T & 0_{k \times (p-2k)} \\ A_k^*(\theta') - A_k^*(\theta) & 0_{k \times k} & 0_{k \times (p-2k)} \\ 0_{(p-2k) \times k} & 0_{(p-2k) \times k} & 0_{p-2k} \end{bmatrix} (0, v, 0)^T \\ &\geq \sup_{\|u\|_2=1, \|v\|_2=1} u^T (A_k^*(\theta') - A_k^*(\theta))v \\ &\geq \|A_k^*(\theta' - \theta)\|_{\text{op}}^2 \\ &= H(\theta', \theta)(\tau n^{-\frac{1}{2}})^2. \end{aligned}$$

So,

$$\min_{H(\theta, \theta') \geq 1} \frac{\|\Omega(\theta) - \Omega(\theta')\|_2^2}{H(\theta, \theta')} \geq (\tau n^{-\frac{1}{2}})^2.$$

7.3.4 Proof of Lemma 10

Since

$$\|P_0 \wedge \bar{P}\| \geq 1 - \frac{1}{2} \left(\int \frac{f_m^2}{f_0} du - 1 \right)^{\frac{1}{2}},$$

we turn to bound $\int \frac{f_m^2}{f_0} du - 1$. We have

$$\begin{aligned}
\int \frac{f_m^2}{f_0} du - 1 &= \frac{1}{p^2} \left(\sum_k \int \frac{f_k^2}{f_0} du + \sum_{i \neq j} \int \frac{f_i f_j}{f_0} du \right) - 1 \\
&= \frac{1}{p^2} (p(1 - \tau^2 k n^{-1})^{-\frac{n}{2}} + p^2 - p) - 1 \\
&\leq \frac{1}{2p} \tau^2 k \\
&\leq \frac{1}{16},
\end{aligned}$$

then

$$\|P_0 \wedge \bar{P}\| \geq 1 - \frac{1}{2} \times \frac{1}{4} = \frac{7}{8}.$$

7.4 Proof of Lemma 11, Lemma 12 in the analysis of Theorem 5

In this section, we prove Lemma 11, Lemma 12 to establish Theorem 5.

7.4.1 Proof of Lemma 11

It is easy to prove that $tp_k(\Omega) = \frac{1}{k} \sum_{i=k}^{2k} bd_i(\Omega)$, where $bd_i(\cdot)$ is defined in Equation (65). Then,

$$\|tp_k(\Omega) - \Omega\|_{\text{op}} \leq \frac{1}{k} \sum_{i=k}^{2k} \|bd_i(\Omega) - \Omega\|_{\text{op}}$$

The matrix $bd_k(\Omega) - \Omega$ is composed by a lower triangular matrix U and its conjugate U^T . Note that $\Omega = (I - A)^T D^{-1} (I - A)$, where $A \equiv [a_{ij}]_{p \times p}$. Let $\text{row}_i(S)$ and $\text{col}_i(S)$ be the i -th row and column of matrix S . Since $\Omega \in \mathcal{Q}_\alpha(\eta, M)$, for any fixed i , $\|U\|_\infty = \max_i (\|\text{row}_i(U)\|_1)$. Let $a_{ss} = 1, \forall s$. we have

$$\begin{aligned}
\|U\|_\infty &\leq \max_i \|\text{row}_i(U)\|_1 \\
&\leq \max_i \eta \sum_{s=i}^p |a_{si}| \sum_{j=1}^{i-k} |a_{sj}| \\
&\leq \max_i \eta M \sum_{s=i}^p (s-i)^{-\alpha-1} (s-i+k)^{-\alpha} \\
&\leq C_\eta M k^{-\alpha}.
\end{aligned}$$

For any fixed j ,

$$\begin{aligned}
\|U\|_1 &\leq \max_j \|\text{col}_j(U)\|_1 \\
&\leq \max_j \eta M \sum_{s=j+k}^p |a_{sj}| \\
&\leq \max_j \eta M \sum_{s=j+k}^p (s-j)^{-\alpha-1} \\
&\leq C\eta M k^{-\alpha}.
\end{aligned}$$

Since $\|U\|_{\text{op}} \leq \sqrt{\|U\|_{\infty}\|U\|_1}$ and $\|bd_k(\Omega) - \Omega\|_{\text{op}} \leq \|U\|_{\text{op}} + \|U^T\|_{\text{op}}$, so

$$\|tp_k(\Omega) - \Omega\|_{\text{op}} \leq \frac{1}{k} \sum_{i=k}^{2k} \|bd_i(\Omega) - \Omega\|_{\text{op}} \quad (72)$$

$$\begin{aligned}
&\leq \|bd_k(\Omega) - \Omega\|_{\text{op}} \\
&\leq 2C\eta M k^{-\alpha}.
\end{aligned} \quad (73)$$

7.4.2 Proof of Lemma 12

The proof of Lemma 12 is basically the same with the one of Lemma 4, except for several steps.

Since $\Omega \in \mathcal{Q}_{\alpha}(\eta, M)$, Equation (67) should be updated by

$$\|t_i\|_{\text{op}} \leq M(i - m - k + 1)^{-\alpha-1/2},$$

and then Equation (68) is replaced by

$$\|\Phi_{m-k}^{(3k)} - \Phi_{m-k}^{(3k)'}\|_{\text{op}}^2 \leq (4\alpha + 2)M^2 k^{-2\alpha}.$$

According to Lemma 23, the operator norm of any row of $H' - K$ can be bounded by $2M\eta^2(2\alpha + 2)^{1/2}k^{-\alpha-1/2}$, so $\|H' - K\|_{\text{op}}^2 \leq 8\eta^4 M^2(2\alpha + 2)k^{-2\alpha}$ and $\|D' - F'\|_{\text{op}}^2 \leq 16\eta^8 M^2(2\alpha + 2)k^{-2\alpha}$. Equation (69) is replaced by

$$\|\Phi_{m-k}^{(3k)'} - \Psi_{m-k}^{(3k)}\|_{\text{op}}^2 \leq 96(2\alpha + 2)M^2\eta^{16}k^{-2\alpha}.$$

So, we have

$$\|\Phi_{m-k}^{(3k)} - \Psi_{m-k}^{(3k)}\|_{\text{op}}^2 \leq 1200\alpha M^2\eta^{16}k^{-2\alpha}.$$

7.5 Proof of Lemma 13, Lemma 14 in the analysis of Theorem 6

In this section, we prove Lemma 13, Lemma 14 to establish Theorem 6.

7.5.1 Proof of Lemma 13

Let $B(\theta) \equiv [b_{ij}]$, then we have the specific value of each entry. It is easy to check $b_{ij} \leq M(i-j)^{-\alpha-1}$. One can check $\Omega(\theta) \in \mathcal{P}_3$ has the specific form of

$$\Omega(\theta) = \begin{bmatrix} I_k + B_k^*(\theta)^T B_k^*(\theta) & -B_k^*(\theta)^T & 0_{k \times (p-2k)} \\ -B_k^*(\theta) & I_k & 0_{k \times (p-2k)} \\ 0_{(p-2k) \times k} & 0_{(p-2k) \times k} & I_{p-2k} \end{bmatrix} \quad (74)$$

Let $\mathbf{1}$ denote the all-ones vector. We find

$$\begin{aligned} \lambda_{\max}(\Omega(\theta)) &\leq \lambda_{\max}(\Omega(\mathbf{1})) = (1 + \lambda_{\max}(B_k^*(\mathbf{1})))^2 = (1 + \sqrt{k/n}\tau)^2 \leq (1 + \tau)^2 \leq \eta; \\ \lambda_{\min}(\Omega(\theta)) &\geq \lambda_{\min}(\Omega(\mathbf{1})) = (1 + \lambda_{\max}(B_k^8(\mathbf{1})))^{-2} = (1 + \sqrt{k/n}\tau)^{-2} \geq (1 + \tau)^{-2} \geq \eta^{-1}. \end{aligned}$$

The eigenvalues of $\Omega(\theta)$ are in the interval $[\eta^{-1}, \eta]$. So $\mathcal{P}_3 \in \mathcal{P}_\alpha(\eta, M)$.

7.5.2 Proof of Lemma 14

The proof follows most part of the one of Lemma 7. Some inequalities in Lemma 7 needs to be rechecked. Assume that $\Sigma(\theta) = \{\Omega(\theta)^{-1} : \Omega(\theta) \in \mathcal{P}_3\}$ and $D = \Sigma(\theta') - \Sigma(\theta)$, one can verify that D has the same form except that A_k^* is replaced by B_k^* . We can show

$$\|D\Sigma(\theta)^{-1}\|_{\text{op}} \leq \eta\|D\|_{\infty} \leq \eta(\tau(nk)^{-\frac{1}{2}}k + (\tau(nk)^{-\frac{1}{2}})^2k^2) \leq \frac{1}{2}.$$

$$\|D\Sigma(\theta)^{-1}\|_{\text{F}}^2 \leq \eta^2(2k(\tau(nk)^{-\frac{1}{2}})^2 + (2k-1)((\tau(nk)^{-\frac{1}{2}})^2k)^2) \leq 4\eta^2(\tau n^{-\frac{1}{2}})^2.$$

So Equation (33) still holds in this case. As for the Equation (34), similarly,

$$\begin{aligned} \|\Omega(\theta') - \Omega(\theta)\|_{\text{op}}^2 &\geq \|(\theta' - \theta)B_k^*\|_{\text{op}}^2 = H(\theta, \theta')(\tau(nk)^{-\frac{1}{2}})^2k = H(\theta, \theta')(\tau n^{-\frac{1}{2}})^2, \\ \min_{H(\theta, \theta') \geq 1} \frac{\|\Omega'(\theta) - \Omega'(\theta')\|_2^2}{H(\theta, \theta')} &\geq (\tau n^{-\frac{1}{2}})^2. \end{aligned}$$

Plugging the result of Equation (33) and (34) in Lemma 6, we have

$$\inf_{\tilde{\Omega}} \sup_{\mathcal{P}_3} \mathbb{E} \|\tilde{\Omega} - \Omega\|_{\text{op}}^2 \geq \frac{\tau^2}{16} n^{-1} k = \frac{\tau^2}{16} n^{-1} \min\{n^{\frac{1}{2\alpha+1}}, \frac{p}{2}\}.$$

7.6 Proof of Lemma 15 in the analysis of Theorem 7

In this section, we prove Lemma 15 to establish Theorem 7.

7.6.1 Proof of Lemma 15

Let the projection of X_i on the linear span of $\mathbf{X}_{1:i-1}$ be

$$\hat{X}_i = \mathbf{X}_{1,i-1}^T \mathbf{a}_i, \quad d_i = \text{Var}(X_i - \hat{X}_i).$$

Let the projection of X_i on the linear span of $\mathbf{X}_{i-k:i-1}$ be

$$\hat{X}_i^{[k_1]} = \mathbf{X}_{i-k:i-1}^T \mathbf{b}'_i = \mathbf{X}_{1,i-1}^T \mathbf{b}_i, \quad e_i = \text{Var}(X_i - \hat{X}_i^{[k_1]}),$$

where $\mathbf{b}_i = (0, \mathbf{b}'_i)^T$, $k_1 = \lceil n^{\frac{2\alpha+1}{(2\alpha+2)2\alpha}} \rceil$. Note that $\hat{\mathbf{a}}_i$ and \hat{d}_i is the empirical coefficient and residual of the regression $X_i \sim \mathbf{X}_{i-k:i-1}$. First, we prove the $\mathbb{E}|\hat{d}_i - d_i|^2 \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}$.

$$\mathbb{E}|\hat{d}_i - d_i|^2 \leq 2\mathbb{E}|\hat{d}_i - e_i|^2 + 2|e_i - d_i|^2.$$

According to Lemma 23 we know that $|e_i - d_i|^2 \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}$. The regression theory implies that, $(n-k)\hat{d}_i/e_i \sim \chi^2(n-k)$. So we have

$$\mathbb{P}(|\hat{d}_i/e_i - 1| > t) \leq 2\exp(-(n-k)t^2/8), \quad \forall t \in (0, 1).$$

Then

$$\mathbb{E}|\hat{d}_i - e_i|^2 \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}.$$

So, we prove that $\mathbb{E}|\hat{d}_i - d_i|^2 \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}$.

Then we turn to prove $\mathbb{E}(\|\hat{\mathbf{a}}_i^* - \mathbf{a}_i\|_{\text{op}}^2) \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}$. For any fixed i , based on the conditional normal distribution, condition on $\mathbf{Z}_{i-k:i-1}$,

$$\hat{\mathbf{a}}_i|\mathbf{Z}_{i-k:i-1} = (0, \dots, 0, \hat{a}_{i(i-k)}, \dots, \hat{a}_{i(i-1)})^T|\mathbf{Z}_{i-k:i-1} \sim N(\mathbf{b}_i, (\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1} \text{Var}(X_i|\mathbf{Z}_{i-k:i-1})).$$

So that for any $0 < j < i$, $\hat{a}_{i(i-j)}|\mathbf{Z}_{i-k:i-1} = b_{i(i-j)} + \sigma_j$, where σ_j follows normal distribution with bounded variance $\text{Var}(\sigma_j) \leq \|(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}} \text{Var}(X_i|\mathbf{Z}_{i-k:i-1}) \leq \eta\|(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}}$. Since $\hat{a}_{i(i-j)}^* = \hat{a}_{i(i-j)} \mathbf{1}(|\hat{a}_{i(i-j)}| > \lambda_j)$, we can show, condition on $\mathbf{Z}_{i-k:i-1}$

$$|\hat{a}_{i(i-j)}^* - b_{i(i-j)}| \leq \min\{|b_{i(i-j)}|, 3/2\lambda_j\} + |3\sigma_j| \mathbf{1}(|\sigma_j| > \lambda_j/2). \quad (75)$$

We need to prove Equation (75) first. One can check that

$$|\hat{a}_{i(i-j)}^* - b_{i(i-j)}| = \max\{|\sigma_j| \mathbf{1}(|b_{i(i-j)} + \sigma_j| > \lambda_j), |b_{i(i-j)}| \mathbf{1}(|b_{i(i-j)} + \sigma_j| \leq \lambda_j)\}.$$

For the first term, we have

$$\begin{aligned} |\sigma_j| \mathbf{1}(|b_{i(i-j)} + \sigma_j| > \lambda_j) &\leq |\sigma_j| \mathbf{1}(|\sigma_j| > \lambda_j/2) + |\sigma_j| \mathbf{1}(|\sigma_j| \leq \lambda_j/2 \cap |b_{i(i-j)}| > \lambda_j/2) \\ &\leq |\sigma_j| \mathbf{1}(|\sigma_j| > \lambda_j/2) + \min\{|b_{i(i-j)}|, \lambda_j/2\}. \end{aligned}$$

Similarly, for the second term, we have

$$\begin{aligned} |b_{i(i-j)}| \mathbf{1}(|b_{i(i-j)} + \sigma_j| \leq \lambda_j) &\leq |b_{i(i-j)}| \mathbf{1}(|b_{i(i-j)} + \sigma_j| \leq \lambda_j \cap |\sigma_j| > \lambda_j/2) \\ &\quad + |b_{i(i-j)}| \mathbf{1}(|b_{i(i-j)} + \sigma_j| \leq \lambda_j \cap |\sigma_j| \leq \lambda_j/2) \\ &\leq |3\sigma_j| \mathbf{1}(|\sigma_j| > \lambda_j/2) + \min\{|b_{i(i-j)}|, 3\lambda_j/2\}. \end{aligned}$$

We finish the proof of Equation (75).

Then,

$$\begin{aligned}
& \mathbb{E}(\|\hat{\mathbf{a}}_i^* - \mathbf{b}_i\|_{\text{op}}^2 | \mathbf{Z}_{i-k:i-1}) \\
& \leq 2 \sum_{j=1}^{i-1} \min\{|b_{i(i-j)}|, (3/2\lambda_j)\}^2 + 2 \sum_{j=1}^{i-1} \mathbb{E}((3\sigma_j)^2 \mathbf{1}(|\sigma_j| > \lambda_j/2) | \mathbf{Z}_{i-k:i-1}) \\
& \leq 4 \sum_{j=1}^{i-1} \min\{|a_{i(i-j)}|, (3/2\lambda_j)\}^2 + 18 \sum_{j=1}^{i-1} \mathbb{E}(\sigma_j^2 \mathbf{1}(|\sigma_j| > \lambda_j/2) | \mathbf{Z}_{i-k:i-1}) + 2\|\mathbf{a}_i - \mathbf{b}_i\|_{\text{op}}^2 \\
& \leq 6 \sum_{j=1}^{i-1} ((|a_{i(i-j)}|\lambda_j)) + 18 \sum_{j=1}^{i-1} \mathbb{E}(\sigma_j^2 \mathbf{1}(|\sigma_j| > \lambda_j/2) | \mathbf{Z}_{i-k:i-1}) + Cn^{-\frac{2\alpha+1}{2\alpha+2}}
\end{aligned} \tag{76}$$

Form Lemma 23, we know that $\|\mathbf{a}_i - \mathbf{b}_i\|_{\text{op}}^2 \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}$.

We can show

$$\begin{aligned}
\sum_{j=1}^{i-1} \lambda_j |a_{i(i-j)}| & \leq \sum_{k=J_0}^{J_1+1} \sqrt{(k-J_0)R} \times M(2^{-(k-J_0)\alpha} - 2^{-(k+1-J_0)\alpha})2^{-J_0\alpha} \\
& \leq n^{-1/2}2^{-J_0\alpha} M \sqrt{\|n(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}}} \left(\sum_{k=0}^{J_1-J_0+1} 2^{-k\alpha} \sqrt{k} \right) \\
& \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}} \sqrt{\|n(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}}} \\
& \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}} \max\{\|n(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}}, 1\},
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{j=1}^{i-1} \mathbb{E}(\sigma_j^2 \mathbf{1}(|\sigma_j| > \lambda_j/2) | \mathbf{Z}_{i-k:i-1}) \\
& \leq \sum_{j=1}^{i-1} \mathbb{E}(\eta \|(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}} \mathbf{1}(|\sigma_j| > \lambda_j/2) | \mathbf{Z}_{i-k:i-1}) \\
& \leq \eta \|(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}} \mathbb{P}(|\sigma_j| > \lambda_j/2 | \mathbf{Z}_{i-k:i-1}) \\
& \leq n^{-1} C \|n(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}} \sum_{j=1}^{i-1} \exp\left(-\frac{\lambda_j^2}{8\eta \|(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}}}\right) \\
& \leq n^{-1} C \|n(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}} \sum_{k=J_0}^{J_1+1} \exp\left(-\frac{k-J_0}{8}\right) (2^k) \\
& \leq Cn^{-1} \|n(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}}.
\end{aligned}$$

Since $\mathbb{E}\|n(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}} \leq \mathbb{E}\|n(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1} - \Omega_{i-k,k}\|_{\text{op}} + \|\Omega_{i-k,k}\|_{\text{op}} \leq$

$1 + \eta$ can be bounded by constant. So,

$$\begin{aligned}
& \mathbb{E}(\|\hat{\mathbf{a}}_i^* - \mathbf{b}_i\|_{\text{op}}^2) \\
&= \mathbb{E}(\mathbb{E}(\|\hat{\mathbf{a}}_i^* - \mathbf{b}_i\|_{\text{op}}^2 | \mathbf{Z}_{i-k:i-1})) \\
&\leq Cn^{-\frac{2\alpha+1}{2\alpha+2}} \max\{\mathbb{E}\|n(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}}, 1\} + Cn^{-1} \mathbb{E}\|n(\mathbf{Z}_{i-k:i-1}^T \mathbf{Z}_{i-k:i-1})^{-1}\|_{\text{op}} + Cn^{-\frac{2\alpha+1}{2\alpha+2}} \\
&\leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}.
\end{aligned}$$

Then,

$$\mathbb{E}(\|\hat{\mathbf{a}}_i^* - \mathbf{a}_i\|_{\text{op}}^2) \leq 2\mathbb{E}(\|\hat{\mathbf{a}}_i^* - \mathbf{b}_i\|_{\text{op}}^2) + 2\mathbb{E}(\|\mathbf{b}_i - \mathbf{a}_i\|_{\text{op}}^2) \leq Cn^{-\frac{2\alpha+1}{2\alpha+2}}.$$

7.7 Proof of Lemma 16, Lemma 17 in the analysis of Theorem 8

In this section, we prove Lemma 16, Lemma 17 to establish Theorem 8.

7.7.1 Proof of Lemma 16

Let $C(\theta) \equiv [c_{ij}]_{k \times k}$, it is easy to check $\sum_{i-j > k} |c_{ij}| \leq Mk^{-\alpha-1}$. One can check $\Omega(\theta)$ in \mathcal{P}_4 has the specific form of

$$\Omega(\theta) = \begin{bmatrix} E_1(\theta(1)) & 0_{2k} & \dots & 0_{2k} \\ 0_{2k} & E_2(\theta(2)) & \dots & 0_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ 0_{2k} & 0_{2k} & \dots & E_{\lceil \frac{p}{2k} \rceil}(\theta(\lceil \frac{p}{2k} \rceil)) \end{bmatrix},$$

where,

$$E_s = \begin{bmatrix} I_k + C_s(\theta(s))^T C_s(\theta(s)) & -C_s(\theta(s))^T \\ -C_s(\theta(s)) & I_k \end{bmatrix}.$$

Let $\mathbf{1}$ be the all-ones vector, then

$$\lambda_{\max}(\Omega(\theta)) \leq \max_s \lambda_{\max}(E_{2k}^s(\theta(s))) \leq \lambda_{\max}(E_{2k}^s(\mathbf{1})) = (1 + \lambda_{\max} C_k^s(\mathbf{1}))^2 \leq (1 + \tau)^2 \leq \eta,$$

$$\lambda_{\min}(\Omega(\theta)) \geq \min_s \lambda_{\min}(E_{2k}^s(\theta(s))) \geq \lambda_{\min}(E_{2k}^s(\mathbf{1})) = (1 + \lambda_{\max} C_k^s(\mathbf{1}))^{-2} \geq (1 + \tau)^{-2} \geq \eta^{-1}.$$

The eigenvalues of $\Omega(\theta)$ are in the interval $[\eta^{-1}, \eta]$. So $\mathcal{P}_4 \in \mathcal{P}_\alpha(\eta, M)$.

7.7.2 Proof of Lemma 17

For equation (42), the proof is almost the same as that of Lemma 7. Since Lemma 17 is about the distributions in the subset \mathcal{P}_4 , we need to recheck the inequalities (70) and (71). Assume that $\Sigma(\theta) = \{\Omega(\theta)^{-1} : \Omega(\theta) \in \mathcal{P}_4\}$ and $D = \Sigma(\theta') - \Sigma(\theta)$, note that $H(\theta, \theta') = 1$, in this case, one can verify that we only need to replace A_k^* by C_k^s . Then we still have

$$\begin{aligned}
\|D\|_{\text{op}} &\leq 2(1 + \|C_k^s(\mathbf{1})\|_{\text{op}}) \|C_k^s(d)\|_{\text{op}} \leq 4\tau n^{-\frac{1}{2}}, \\
\|D\|_{\text{F}} &\leq 2(1 + \|C_k^s(\mathbf{1})\|_{\text{F}}) \|C_k^s(d)\|_{\text{F}} \leq 4\tau n^{-\frac{1}{2}}.
\end{aligned}$$

So Equation (42) holds in this case. As for the Equation (43), similarly,

$$\begin{aligned}\|\Omega(\theta') - \Omega(\theta)\|_F^2 &\geq \sum_s \|C_k^*(\theta'(s) - \theta(s))\|_F^2 = H(\theta, \theta')(\tau n^{-\frac{1}{2}})^2, \\ \min_{H(\theta, \theta') \geq 1} \frac{\|\Omega'(\theta) - \Omega'(\theta')\|_2^2}{H(\theta, \theta')} &\geq \tau^2 n^{-1}.\end{aligned}$$

7.8 Proof of Lemma 18, Lemma 19 in the analysis of Theorem 9

In this section, we prove Lemma 18 and Lemma 19 to establish Theorem 9.

7.8.1 Proof of Lemma 18

It is easy to show $\text{diag}(\Omega) = I$. Now we need to verify $\Omega \in \mathcal{P}_\alpha(\eta^2, M\eta)$. Let S denote $(\text{diag}(\Omega'^{-1}))^{\frac{1}{2}}$. The Cholesky decomposition of Ω is:

$$\begin{aligned}\Omega &= S\Omega'S \\ &= S(I - A')^T D'^{-1}(I - A')S \\ &= (S - A'S)^T D'^{-1}(S - A'S) \\ &= (I - S^{-1}A'S)^T S D'^{-1}S (I - S^{-1}A'S).\end{aligned}$$

According to Lemma 22, $\eta^{-\frac{1}{2}} \leq \lambda_{\min}(S) \leq \lambda_{\max}(S) \leq \eta^{\frac{1}{2}}$ and $\eta^{-1} \leq \lambda_{\min}(\Omega') \leq \lambda_{\max}(\Omega') \leq \eta$. Therefore, we obtain $\eta^{-2} \leq \lambda_{\min}(\Omega) \leq \lambda_{\max}(\Omega) \leq \eta^2$. Let $A \equiv [a_{ij}]_{p \times p} = S^{-1}A'S$. The desired result $\Omega \in \mathcal{P}_\alpha(\eta^2, M\eta)$ then immediately follows from that $\max_i \sum_{j < i-k} |a_{ij}| < M\eta k^{-\alpha}$ for $k \in [i-1]$.

Since $\text{diag}(\Omega'^{-1})_i^{\frac{1}{2}} > 0$, f_i is a strictly increasing function. Therefore, $\{\Omega, \{f_i\}\} \in \mathcal{P}'_\alpha(\eta^2, M\eta)$.

7.8.2 Proof of Lemma 19

One can check

$$\Omega(\theta) = \begin{bmatrix} I_k + A_k^*(\theta)^T A_k^*(\theta) & -A_k^*(\theta)^T \text{diag}(I_k + A_k^*(\theta) A_k^*(\theta)^T)^{\frac{1}{2}} & 0_{k \times (p-2k)} \\ -\text{diag}(I_k + A_k^*(\theta) A_k^*(\theta)^T)^{\frac{1}{2}} A_k^*(\theta) & \text{diag}(I_k + A_k^*(\theta) A_k^*(\theta)^T) & 0_{k \times (p-2k)} \\ 0_{(p-2k) \times k} & 0_{(p-2k) \times k} & I_{p-2k} \end{bmatrix}.$$

Then,

$$\begin{aligned}\|\Omega(\theta') - \Omega(\theta)\|_{\text{op}}^2 &\geq \|\text{diag}(I_k + A_k^*(\theta) A_k^*(\theta)^T)^{\frac{1}{2}} A_k^*(\theta) - \text{diag}(I_k + A_k^*(\theta') A_k^*(\theta')^T)^{\frac{1}{2}} A_k^*(\theta')\|_{\text{op}}^2 \\ &= H(\theta, \theta')(1 + (\tau n^{-\frac{1}{2}})^2)(\tau n^{-\frac{1}{2}})^2,\end{aligned}$$

which further implies

$$\min_{H(\theta, \theta') \geq 1} \frac{\|\Omega(\theta) - \Omega(\theta')\|_2^2}{H(\theta, \theta')} \geq (\tau n^{-\frac{1}{2}})^2.$$

References

- [1] Patrice Assouad. Deux remarques sur l'estimation. *Comptes rendus des séances de l'Académie des sciences. Série 1, Mathématique*, 296(23):1021–1024, 1983.
- [2] Onureena Banerjee, Laurent El Ghaoui, and Alexandre dAspremont. Model selection through sparse maximum likelihood estimation for multivariate Gaussian or binary data. *Journal of Machine learning research*, 9(Mar):485–516, 2008.
- [3] Sayantan Banerjee and Subhashis Ghosal. Posterior convergence rates for estimating large precision matrices using graphical models. *Electronic Journal of Statistics*, 8(2):2111–2137, 2014.
- [4] Rina Foygel Barber and Mladen Kolar. Rocket: Robust confidence intervals via Kendall's tau for transelliptical graphical models. *arXiv preprint arXiv:1502.07641*, 2015.
- [5] Peter J Bickel, Chris AJ Klaassen, Peter J Bickel, Y Ritov, J Klaassen, Jon A Wellner, and YA'Acov Ritov. *Efficient and adaptive estimation for semiparametric models*. Johns Hopkins University Press Baltimore, 1993.
- [6] Peter J Bickel and Elizaveta Levina. Covariance regularization by thresholding. *The Annals of Statistics*, 36(6):2577–2604, 2008.
- [7] Peter J Bickel and Elizaveta Levina. Regularized estimation of large covariance matrices. *The Annals of Statistics*, 36(1):199–227, 2008.
- [8] Jacob Bien, Florentina Bunea, and Luo Xiao. Convex banding of the covariance matrix. *Journal of the American Statistical Association*, 111(514):834–845, 2016.
- [9] T Tony Cai. Minimax and adaptive inference in nonparametric function estimation. *Statistical Science*, 27(1):31–50, 2012.
- [10] T Tony Cai, Weidong Liu, and Harrison H Zhou. Estimating sparse precision matrix: Optimal rates of convergence and adaptive estimation. *The Annals of Statistics*, 44(2):455–488, 2016.
- [11] T Tony Cai, Zhao Ren, and Harrison H Zhou. Optimal rates of convergence for estimating Toeplitz covariance matrices. *Probability Theory and Related Fields*, 156(1-2):101–143, 2013.
- [12] T Tony Cai, Zhao Ren, and Harrison H Zhou. Estimating structured high-dimensional covariance and precision matrices: Optimal rates and adaptive estimation. *Electronic Journal of Statistics*, 10(1):1–59, 2016.
- [13] T Tony Cai, Cun-Hui Zhang, and Harrison H Zhou. Optimal rates of convergence for covariance matrix estimation. *The Annals of Statistics*, 38(4):2118–2144, 2010.

- [14] T Tony Cai and Harrison H Zhou. Optimal rates of convergence for sparse covariance matrix estimation. *The Annals of Statistics*, 40(5):2389–2420, 2012.
- [15] Tony Cai and Weidong Liu. Adaptive thresholding for sparse covariance matrix estimation. *Journal of the American Statistical Association*, 106(494):672–684, 2011.
- [16] Tony Cai, Weidong Liu, and Xi Luo. A constrained ℓ_1 minimization approach to sparse precision matrix estimation. *Journal of the American Statistical Association*, 106(494):594–607, 2011.
- [17] I Csizsár. Information-type indices of the divergence of distributions. i, ii. *Magyar Tud. Akad. Mat. Fiz. Oszt. Közl.* 17 (1967), 123–149; *ibid*, 17:267–291, 1967.
- [18] Alexandre d’Aspremont, Onureena Banerjee, and Laurent El Ghaoui. First-order methods for sparse covariance selection. *SIAM Journal on Matrix Analysis and Applications*, 30(1):56–66, 2008.
- [19] Bernard Delyon and Anatoli Juditsky. On minimax wavelet estimators. *Applied and Computational Harmonic Analysis*, 3(3):215–228, 1996.
- [20] Michael B Eisen, Paul T Spellman, Patrick O Brown, and David Botstein. Cluster analysis and display of genome-wide expression patterns. *Proceedings of the National Academy of Sciences*, 95(25):14863–14868, 1998.
- [21] Nouredine El Karoui. On the largest eigenvalue of Wishart matrices with identity covariance when n , p and p/n tend to infinity. *arXiv preprint math/0309355*, 2003.
- [22] Nouredine El Karoui. Operator norm consistent estimation of large-dimensional sparse covariance matrices. *The Annals of Statistics*, 36(6):2717–2756, 2008.
- [23] Jianqing Fan, Lingzhou Xue, and Hui Zou. Multitask quantile regression under the transnormal model. *Journal of the American Statistical Association*, 111(516):1726–1735, 2016.
- [24] Reinhard Furrer and Thomas Bengtsson. Estimation of high-dimensional prior and posterior covariance matrices in Kalman filter variants. *Journal of Multivariate Analysis*, 98(2):227–255, 2007.
- [25] Thomas M Hamill, Jeffrey S Whitaker, and Chris Snyder. Distance-dependent filtering of background error covariance estimates in an ensemble kalman filter. *Monthly Weather Review*, 129(11):2776–2790, 2001.
- [26] Mark H Heyer and F Peter Schloerb. Application of principal component analysis to large-scale spectral line imaging studies of the interstellar medium. *The Astrophysical Journal*, 475(1):173, 1997.

- [27] Addison Hu and Sahand Negahban. Minimax estimation of bandable precision matrices. In *Advances in Neural Information Processing Systems*, pages 4893–4901, 2017.
- [28] Jianhua Z Huang, Naiping Liu, Mohsen Pourahmadi, and Linxu Liu. Covariance matrix selection and estimation via penalised normal likelihood. *Biometrika*, 93(1):85–98, 2006.
- [29] Iain M Johnstone. On the distribution of the largest eigenvalue in principal components analysis. *The Annals of Statistics*, pages 295–327, 2001.
- [30] Iain M Johnstone and Arthur Yu Lu. On consistency and sparsity for principal components analysis in high dimensions. *Journal of the American Statistical Association*, 104(486):682–693, 2009.
- [31] Maurice G Kendall. A new measure of rank correlation. *Biometrika*, 30(1/2):81–93, 1938.
- [32] Clifford Lam and Jianqing Fan. Sparsistency and rates of convergence in large covariance matrix estimation. *The Annals of Statistics*, 37(6B):4254, 2009.
- [33] Olivier Ledoit and Michael Wolf. Improved estimation of the covariance matrix of stock returns with an application to portfolio selection. *Journal of Empirical Finance*, 10(5):603–621, 2003.
- [34] Kyoungjae Lee and Jaeyong Lee. Estimating large precision matrices via modified Cholesky decomposition. *arXiv preprint arXiv:1707.01143*, 2017.
- [35] Elizaveta Levina, Adam Rothman, and Ji Zhu. Sparse estimation of large covariance matrices via a nested lasso penalty. *The Annals of Applied Statistics*, 2(1):245–263, 2008.
- [36] Han Liu, John Lafferty, and Larry Wasserman. The nonparanormal: Semiparametric estimation of high dimensional undirected graphs. *Journal of Machine Learning Research*, 10(Oct):2295–2328, 2009.
- [37] Nicolai Meinshausen and Peter Bühlmann. High-dimensional graphs and variable selection with the lasso. *The Annals of Statistics*, pages 1436–1462, 2006.
- [38] Ritwik Mitra and Cun-Hui Zhang. Multivariate analysis of nonparametric estimates of large correlation matrices. *arXiv preprint arXiv:1403.6195*, 2014.
- [39] Nikhil Padmanabhan, Martin White, Harrison H Zhou, and Ross O’Connell. Estimating sparse precision matrices. *Monthly Notices of the Royal Astronomical Society*, 460(2):1567–1576, 2016.

- [40] Debashis Paul. Asymptotics of sample eigenstructure for a large dimensional spiked covariance model. *Statistica Sinica*, 17(4):1617, 2007.
- [41] Pradeep Ravikumar, Martin J Wainwright, Garvesh Raskutti, and Bin Yu. High-dimensional covariance estimation by minimizing ℓ_1 -penalized log-determinant divergence. *Electronic Journal of Statistics*, 5:935–980, 2011.
- [42] Zhao Ren, Tingni Sun, Cun-Hui Zhang, and Harrison H Zhou. Asymptotic normality and optimalities in estimation of large Gaussian graphical models. *The Annals of Statistics*, 43(3):991–1026, 2015.
- [43] Adam J Rothman, Peter J Bickel, Elizaveta Levina, and Ji Zhu. Sparse permutation invariant covariance estimation. *Electronic Journal of Statistics*, 2:494–515, 2008.
- [44] Leonas Saulis and VA Statulevicius. *Limit theorems for large deviations*, volume 73. Springer Science & Business Media, 2012.
- [45] Juliane Schäfer and Korbinian Strimmer. A shrinkage approach to large-scale covariance matrix estimation and implications for functional genomics. *Statistical applications in genetics and molecular biology*, 4(1), 2005.
- [46] Charles Spearman. Spearmans rank correlation coefficient. *Amer J Psychol*, 15:72–101, 1904.
- [47] Tingni Sun and Cun-Hui Zhang. Sparse matrix inversion with scaled lasso. *The Journal of Machine Learning Research*, 14(1):3385–3418, 2013.
- [48] Kenneth W Wachter. Probability plotting points for principal components. In *Ninth Interface Symposium Computer Science and Statistics*, pages 299–308. Prindle, Weber and Schmidt, Boston, 1976.
- [49] Kenneth W Wachter. The strong limits of random matrix spectra for sample matrices of independent elements. *The Annals of Probability*, 6(1):1–18, 1978.
- [50] Wei Biao Wu and Mohsen Pourahmadi. Nonparametric estimation of large covariance matrices of longitudinal data. *Biometrika*, 90(4):831–844, 2003.
- [51] Wei Biao Wu and Mohsen Pourahmadi. Banding sample autocovariance matrices of stationary processes. *Statistica Sinica*, pages 1755–1768, 2009.
- [52] Luo Xiao and Florentina Bunea. On the theoretic and practical merits of the banding estimator for large covariance matrices. *arXiv preprint arXiv:1402.0844*, 2014.
- [53] Bin Yu. Assouad, Fano, and Le Cam. *Festschrift for Lucien Le Cam*, 423:435, 1997.
- [54] Ming Yuan. High dimensional inverse covariance matrix estimation via linear programming. *Journal of Machine Learning Research*, 11(Aug):2261–2286, 2010.

- [55] Ming Yuan and Yi Lin. Model selection and estimation in the Gaussian graphical model. *Biometrika*, 94(1):19–35, 2007.