

WEIGHTED COMPLETE CONTINUITY FOR THE COMMUTATOR OF MARCINKIEWICZ INTEGRAL

GUOEN HU

ABSTRACT. Let Ω be homogeneous of degree zero, have mean value zero and integrable on the unit sphere, and \mathcal{M}_Ω be the higher-dimensional Marcinkiewicz integral associated with Ω . In this paper, the author considers the complete continuity on weighted $L^p(\mathbb{R}^n)$ spaces with $A_p(\mathbb{R}^n)$ weights, weighted Morrey spaces with $A_p(\mathbb{R}^n)$ weights, for the commutator generated by $\text{CMO}(\mathbb{R}^n)$ functions and \mathcal{M}_Ω when Ω satisfies certain size conditions.

1. INTRODUCTION

As an analogy of the classical Littlewood-Paley g -function, Marcinkiewicz [31] introduced the operator

$$\mathcal{M}(f)(x) = \left(\int_0^\pi \frac{|F(x+t) - F(x-t) - 2F(x)|^2}{t^3} dt \right)^{\frac{1}{2}},$$

where $F(x) = \int_0^x f(t)dt$. This operator is now called Marcinkiewicz integral. Zygmund [40] proved that \mathcal{M} is bounded on $L^p([0, 2\pi])$ for $p \in (1, \infty)$. Stein [34] generalized the Marcinkiewicz operator to the case of higher dimension. Let Ω be homogeneous of degree zero, integrable and have mean value zero on the unit sphere S^{n-1} . Define the Marcinkiewicz integral operator \mathcal{M}_Ω by

$$(1.1) \quad \mathcal{M}_\Omega(f)(x) = \left(\int_0^\infty |F_{\Omega,t}f(x)|^2 \frac{dt}{t^3} \right)^{\frac{1}{2}},$$

where

$$F_{\Omega,t}f(x) = \int_{|x-y| \leq t} \frac{\Omega(x-y)}{|x-y|^{n-1}} f(y) dy$$

for $f \in \mathcal{S}(\mathbb{R}^n)$. Stein [34] proved that if $\Omega \in \text{Lip}_\alpha(S^{n-1})$ with $\alpha \in (0, 1]$, then \mathcal{M}_Ω is bounded on $L^p(\mathbb{R}^n)$ for $p \in (1, 2]$. Benedek, Calderón and Panzon showed that the $L^p(\mathbb{R}^n)$ boundedness ($p \in (1, \infty)$) of \mathcal{M}_Ω holds true under the condition that $\Omega \in C^1(S^{n-1})$. Using the one-dimensional result and Riesz transforms similarly as in the case of singular integrals (see [7]) and interpolation, Walsh [38] proved that for each $p \in (1, \infty)$, $\Omega \in L(\ln L)^{1/r}(\ln \ln L)^{2(1-2/r')}(S^{n-1})$ is a sufficient condition such that \mathcal{M}_Ω is bounded on $L^p(\mathbb{R}^n)$, where $r = \min\{p, p'\}$ and $p' = p/(p-1)$. Ding, Fan and Pan [17] proved that if $\Omega \in H^1(S^{n-1})$ (the Hardy space on S^{n-1}), then \mathcal{M}_Ω is bounded on $L^p(\mathbb{R}^n)$ for all $p \in (1, \infty)$; Al-Salmam, Al-Qassem, Cheng and Pan [3] proved that $\Omega \in L(\ln L)^{1/2}(S^{n-1})$ is a sufficient condition such that \mathcal{M}_Ω is bounded on $L^p(\mathbb{R}^n)$ for all $p \in (1, \infty)$. Ding, Fan and Pan [18] considered the boundedness on weighted $L^p(\mathbb{R}^n)$ with $A_p(\mathbb{R}^n)$ when $\Omega \in L^q(S^{n-1})$ for some

1991 *Mathematics Subject Classification.* Primary 42B25, Secondary 47B07.

Key words and phrases. commutator, Marcinkiewicz integral, completely continuous operator, Fourier transform, weighted Morrey space.

The research was supported by the NNSF of China under grant #11371370.

$q \in (1, \infty]$, where and in the following, for $p \in [1, \infty)$, $A_p(\mathbb{R}^n)$ denotes the weight function class of Muckenhoupt, see [24] for the definitions and properties of $A_p(\mathbb{R}^n)$. For other works about the operator defined by (1.1), see [2, 3, 9, 17, 19, 21] and the related references therein.

The commutator of \mathcal{M}_Ω is also of interest and has been considered by many authors (see [36, 27, 20, 8, 26]). Let $b \in \text{BMO}(\mathbb{R}^n)$, the commutator generated by \mathcal{M}_Ω and b is defined by

$$(1.2) \quad \mathcal{M}_{\Omega, b} f(x) = \left(\int_0^\infty \left| \int_{|x-y| \leq t} (b(x) - b(y)) \frac{\Omega(x-y)}{|x-y|^{n-1}} f(y) dy \right|^2 \frac{dt}{t^3} \right)^{\frac{1}{2}}.$$

Torchinsky and Wang [36] showed that if $\Omega \in \text{Lip}_\alpha(S^{n-1})$ ($\alpha \in (0, 1]$), then $\mathcal{M}_{\Omega, b}$ is bounded on $L^p(\mathbb{R}^n)$ with bound $C\|b\|_{\text{BMO}(\mathbb{R}^n)}$ for all $p \in (1, \infty)$. Hu and Yan [27] proved the $\Omega \in L(\ln L)^{3/2}(S^{n-1})$ is a sufficient condition such that $\mathcal{M}_{\Omega, b}$ is bounded on L^2 . Ding, Lu and Yabuta [20] considered the weighted estimates for $\mathcal{M}_{\Omega, b}$, and proved that if $\Omega \in L^q(S^{n-1})$ for some $q \in (1, \infty]$, then for $p \in (q', \infty)$ and $w \in A_{p/q'}(\mathbb{R}^n)$, or $p \in (1, q)$ and $w^{-1/(p-1)} \in A_{p'/q'}(\mathbb{R}^n)$, $\mathcal{M}_{\Omega, b}$ is bounded on $L^p(\mathbb{R}^n, w)$. Chen and Lu [8] improved the result in [27] and showed that if $\Omega \in L(\ln L)^{3/2}(S^{n-1})$, then $\mathcal{M}_{\Omega, b}$ is bounded on $L^p(\mathbb{R}^n)$ with bound $C\|b\|_{\text{BMO}(\mathbb{R}^n)}$ for all $p \in (1, \infty)$.

Let $\text{CMO}(\mathbb{R}^n)$ be the closure of $C_0^\infty(\mathbb{R}^n)$ in the $\text{BMO}(\mathbb{R}^n)$ topology, which coincide with $\text{VMO}(\mathbb{R}^n)$, the space of functions of vanishing mean oscillation introduced by Coifman and Weiss [16], see also [6]. Uchiyama [37] proved that if T is a Calderón-Zygmund operator, and $b \in \text{BMO}(\mathbb{R}^n)$, then the commutator of T defined by

$$[b, T]f(x) = b(x)Tf(x) - T(bf)(x),$$

is a compact operator on $L^p(\mathbb{R}^n)$ ($p \in (1, \infty)$) if and only if $b \in \text{CMO}(\mathbb{R}^n)$. Chen and Ding [12] considered the compactness of $\mathcal{M}_{\Omega, b}$ on $L^p(\mathbb{R}^n)$, and proved that if Ω satisfies certain regularity condition of Dini type, then for $p \in (1, \infty)$, $\mathcal{M}_{\Omega, b}$ is compact on $L^p(\mathbb{R}^n)$ if and only if $b \in \text{CMO}(\mathbb{R}^n)$. Using the ideas from [10], Mao, Sawano and Wu [30] consider the compactness of $\mathcal{M}_{\Omega, b}$ when Ω satisfies the size condition that

$$(1.3) \quad \sup_{\zeta \in S^{n-1}} \int_{S^{n-1}} |\Omega(\eta)| \left(\ln \frac{1}{|\eta \cdot \zeta|} \right)^\theta d\eta < \infty,$$

and proved that if Ω satisfies (1.3) for some $\theta \in (3/2, \infty)$, then for $b \in \text{CMO}(\mathbb{R}^n)$ and $p \in (4\theta/(4\theta-3), 4\theta/3)$, $\mathcal{M}_{\Omega, b}$ is compact on $L^p(\mathbb{R}^n)$. Recently, Chen and Hu [11] proved that if $\Omega \in L(\log L)^{\frac{1}{2}}(S^{n-1})$, then for $b \in \text{CMO}(\mathbb{R}^n)$ and $p \in (1, \infty)$, $\mathcal{M}_{\Omega, b}$ is completely continuous on $L^p(\mathbb{R}^n)$. Our first purpose of this paper is to consider the complete continuity on weighted $L^p(\mathbb{R}^n)$ spaces $\mathcal{M}_{\Omega, b}$ when $\Omega \in L^q(S^{n-1})$ for some $q \in (1, \infty]$. To formulate our main result, we first recall some definitions.

Definition 1.1. Let \mathcal{X} be a normed linear spaces and \mathcal{X}^* be its dual space, $\{x_k\} \subset \mathcal{X}$ and $x \in \mathcal{X}$, If for all $f \in \mathcal{X}^*$,

$$\lim_{k \rightarrow \infty} |f(x_k) - f(x)| = 0,$$

then $\{x_k\}$ is said to converge to x weakly, or $x_k \rightharpoonup x$.

Definition 1.2. Let \mathcal{X} , \mathcal{Y} be two Banach spaces and S be a bounded operator from \mathcal{X} to \mathcal{Y} .

- (i) If for each bounded set $\mathcal{G} \subset \mathcal{X}$, $S\mathcal{G} = \{Sx : x \in \mathcal{G}\}$ is a strongly pre-compact set in \mathcal{Y} , then S is called a compact operator from \mathcal{X} to \mathcal{Y} ;
- (ii) if for $\{x_k\} \subset \mathcal{X}$ and $x \in \mathcal{X}$,

$$x_k \rightharpoonup x \text{ in } \mathcal{X} \Rightarrow \|Sx_k - Sx\|_{\mathcal{Y}} \rightarrow 0,$$

then S is said to be a completely continuous operator.

It is well known that, if \mathcal{X} is a reflexive space, and S is completely continuous from \mathcal{X} to \mathcal{Y} , then S is also compact from \mathcal{X} to \mathcal{Y} . On the other hand, if S is a linear compact operator from \mathcal{X} to \mathcal{Y} , then S is also a completely continuous operator. However, if S is not linear, then S is compact do not imply that S is completely continuous. For example, the operator

$$Sx = \|x\|_{l^2}$$

is compact from l^2 to \mathbb{R} , but not completely continuous.

Our first result in this paper can be stated as follows.

Theorem 1.3. *Let Ω be homogeneous of degree zero, have mean value zero on S^{n-1} and $\Omega \in L^q(S^{n-1})$ for some $q \in (1, \infty]$. Suppose that p and w satisfy one of the following conditions*

- (i) $p \in (q', \infty)$ and $w \in A_{p/q'}(\mathbb{R}^n)$;
- (ii) $p \in (1, q)$ and $w^{-1/(p-1)} \in A_{p'/q'}(\mathbb{R}^n)$;
- (iii) $p \in (1, \infty)$ and $w^{q'} \in A_p(\mathbb{R}^n)$.

Then for $b \in \text{CMO}(\mathbb{R}^n)$, $\mathcal{M}_{\Omega, b}$ is completely continuous on $L^p(\mathbb{R}^n, w)$.

Our argument used in the proof of Theorem 1.3 also leads to the complete continuity of $\mathcal{M}_{\Omega, b}$ on weighted Morrey spaces.

Definition 1.4. Let $p \in (0, \infty)$, w be a weight and $\lambda \in (0, 1)$. The weighted Morrey space $L^{p, \lambda}(\mathbb{R}^n, w)$ is defined as

$$L^{p, \lambda}(\mathbb{R}^n, w) = \{f \in L^p_{\text{loc}}(\mathbb{R}^n) : \|f\|_{L^{p, \lambda}(\mathbb{R}^n, w)} < \infty\},$$

with

$$\|f\|_{L^{p, \lambda}(\mathbb{R}^n, w)} = \sup_{y \in \mathbb{R}^n, r > 0} \left(\frac{1}{\{w(B(y, r))\}^\lambda} \int_{B(y, r)} |f(x)|^p w(x) dx \right)^{1/p},$$

here $B(y, r)$ denotes the ball in \mathbb{R}^n centered at y and having radius r , and $w(B(y, r)) = \int_{B(y, r)} w(z) dz$. For simplicity, we use $L^{p, \lambda}(\mathbb{R}^n)$ to denote $L^{p, \lambda}(\mathbb{R}^n, 1)$.

The Morrey space $L^{p, \lambda}(\mathbb{R}^n)$ was introduced by Morrey [17]. It is well-known that this space is closely related to some problems in PED (see [32, 33]), and has interest in harmonic analysis (see [1] and the references therein). Komori and Shirai [28] introduced the weighted Morrey spaces and considered the properties on weighted Morrey spaces for some classical operators. Chen, Ding and Wang [13] considered the compactness of $\mathcal{M}_{\Omega, b}$ on Morrey spaces. They proved that if $\lambda \in (0, 1)$, $\Omega \in L^q(S^{n-1})$ for $q \in (1/(1-\lambda), \infty]$ and satisfies a regularity condition of L^q -Dini type, then $\mathcal{M}_{\Omega, b}$ is compact on $L^{p, \lambda}(\mathbb{R}^n)$. Our second purpose of this paper is to prove the complete continuity of $\mathcal{M}_{\Omega, b}$ on weighted Morrey spaces with $A_p(\mathbb{R}^n)$ weights.

Theorem 1.5. *Let Ω be homogeneous of degree zero, have mean value zero on S^{n-1} and $\Omega \in L^q(S^{n-1})$ for some $q \in (1, \infty]$. Suppose that $p \in (q', \infty)$, $\lambda \in (0, 1)$ and $w \in A_{p/q'}(\mathbb{R}^n)$; or $p \in (1, q')$, $w^r \in A_1(\mathbb{R}^n)$ for some $r \in (q', \infty)$ and $\lambda \in (0, 1 - r'/q)$. Then for $b \in \text{CMO}(\mathbb{R}^n)$, $\mathcal{M}_{\Omega, b}$ is completely continuous on $L^{p, \lambda}(\mathbb{R}^n, w)$.*

Remark 1.6. The proofs of Theorems 1.3 involve some ideas used in [10] and a sufficient condition of strongly pre-compact set in $L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)$ with $w \in A_p(\mathbb{R}^n)$. To prove Theorem 1.5, we will establish a lemma which clarify the relationship of the bounds on $L^p(\mathbb{R}^n, w)$ and the bounds on $L^{p, \lambda}(\mathbb{R}^n, w)$ for a class of sublinear operators, see Lemma 4.1 below.

We make some conventions. In what follows, C always denotes a positive constant that is independent of the main parameters involved but whose value may differ from line to line. We use the symbol $A \lesssim B$ to denote that there exists a positive constant C such that $A \leq CB$. For a set $E \subset \mathbb{R}^n$, χ_E denotes its characteristic function. Let M be the Hardy-Littlewood maximal operator. For $r \in (0, \infty)$, we use M_r to denote the operator $M_r f(x) = (M(|f|^r)(x))^{1/r}$. For a locally integrable function f , the sharp maximal function $M^\sharp f$ is defined by

$$M^\sharp f(x) = \sup_{Q \ni x} \inf_{c \in \mathbb{C}} \frac{1}{|Q|} \int_Q |f(y) - c| dy.$$

2. APPROXIMATION

Let Ω be homogeneous of degree zero, integrable on S^{n-1} . For $t \in [1, 2]$ and $j \in \mathbb{Z}$, set

$$(2.1) \quad K_t^j(x) = \frac{1}{2^j} \frac{\Omega(x)}{|x|^{n-1}} \chi_{\{2^{j-1}t < |x| \leq 2^j t\}}(x).$$

As it was proved in [23], if $\Omega \in L^q(S^{n-1})$ for some $q \in (1, \infty]$, then there exists a constant $\alpha \in (0, 1)$ such that for $t \in [1, 2]$ and $\xi \in \mathbb{R}^n \setminus \{0\}$,

$$(2.2) \quad |\widehat{K}_t^j(\xi)| \lesssim \|\Omega\|_{L^q(S^{n-1})} \min\{1, |2^j \xi|^{-\alpha}\}.$$

Here and in the following for $h \in \mathcal{S}'(\mathbb{R}^n)$, \widehat{h} denotes the Fourier transform of h . Moreover, if $\int_{S^{n-1}} \Omega(x') dx' = 0$, then

$$(2.3) \quad |\widehat{K}_t^j(\xi)| \lesssim \|\Omega\|_{L^1(S^{n-1})} \min\{1, |2^j \xi|\}.$$

Let

$$\widetilde{\mathcal{M}}_\Omega f(x) = \left(\int_1^2 \sum_{j \in \mathbb{Z}} |F_j f(x, t)|^2 dt \right)^{\frac{1}{2}},$$

with

$$F_j f(x, t) = \int_{\mathbb{R}^n} K_t^j(x - y) f(y) dy.$$

For $b \in \text{BMO}(\mathbb{R}^n)$, let $\widetilde{\mathcal{M}}_{\Omega, b}$ be the commutator of $\widetilde{\mathcal{M}}_\Omega$ defined by

$$\widetilde{\mathcal{M}}_{\Omega, b} f(x) = \left(\int_1^2 \sum_{j \in \mathbb{Z}} |F_{j, b} f(x, t)|^2 dt \right)^{1/2},$$

with

$$F_{j,b}f(x,t) = \int_{\mathbb{R}^n} (b(x) - b(y))K_t^j(x-y)f(y)dy.$$

A trivial computation leads to that

$$(2.4) \quad \mathcal{M}_\Omega f(x) \approx \widetilde{\mathcal{M}}_\Omega f(x), \quad \widetilde{\mathcal{M}}_{\Omega,b}f(x) \approx \widetilde{\mathcal{M}}_{\Omega,b}f(x).$$

Let $\phi \in C_0^\infty(\mathbb{R}^n)$ be a nonnegative function such that $\int_{\mathbb{R}^n} \phi(x)dx = 1$, $\text{supp } \phi \subset \{x : |x| \leq 1/4\}$. For $l \in \mathbb{Z}$, let $\phi_l(y) = 2^{-nl}\phi(2^{-l}y)$. It is easy to verify that for any $\varsigma \in (0, 1)$,

$$(2.5) \quad |\widehat{\phi}_l(\xi) - 1| \lesssim \min\{1, |2^l \xi|^\varsigma\}.$$

Let

$$F_j^l f(x,t) = \int_{\mathbb{R}^n} K_t^j * \phi_{j-l}(x-y)f(y)dy.$$

Define the operator $\widetilde{\mathcal{M}}_\Omega^l$ by

$$(2.6) \quad \widetilde{\mathcal{M}}_\Omega^l f(x) = \left(\int_1^2 \sum_{j \in \mathbb{Z}} |F_j^l f(x,t)|^2 dt \right)^{\frac{1}{2}}.$$

This section is devoted to the approximation of $\widetilde{\mathcal{M}}_\Omega$ by $\widetilde{\mathcal{M}}_\Omega^l$. We will prove following theorem.

Theorem 2.1. *Let Ω be homogeneous of degree zero and have mean value zero. Suppose that $\Omega \in L^q(S^{n-1})$ for some $q \in (1, \infty]$, p and w are the same as in Theorem 1.3, then for $l \in \mathbb{N}$,*

$$\|\widetilde{\mathcal{M}}_\Omega f - \widetilde{\mathcal{M}}_\Omega^l f\|_{L^p(\mathbb{R}^n, w)} \lesssim 2^{-\varrho_p l} \|f\|_{L^p(\mathbb{R}^n, w)},$$

with $\varrho_p \in (0, 1)$ a constant depending only on p, n and w .

To prove Theorem 2.1, we will use some lemmas.

Lemma 2.2. *Let Ω be homogeneous of degree zero and belong to $L^q(S^{n-1})$ for some $q \in (1, \infty]$, K_t^j be defined as in (2.1). Then for $t \in [1, 2]$, $l \in \mathbb{N}$, $R > 0$ and $y \in \mathbb{R}^n$ with $|y| < R/4$,*

$$\sum_{j \in \mathbb{Z}} \sum_{k=1}^{\infty} (2^k R)^{\frac{n}{q'}} \left(\int_{2^k R < |x| \leq 2^{k+1} R} |K_t^j * \phi_{j-l}(x+y) - K_t^j * \phi_{j-l}(x)|^q dx \right)^{\frac{1}{q}} \lesssim l.$$

For the proof of Lemma 2.2, see [39].

Lemma 2.3. *Let Ω be homogeneous of degree zero and $\Omega \in L^q(S^{n-1})$ for some $q \in (1, \infty]$, $p \in (1, q)$ and $w^{-1/(p-1)} \in A_{p'/q'}(\mathbb{R}^n)$. Then*

$$(2.7) \quad \left\| \left(\sum_{j \in \mathbb{Z}} |K_t^j * \phi_{j-l} * f_j|^2 \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim \left\| \left(\sum_{j \in \mathbb{Z}} |f_j|^2 \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)}.$$

Proof. Let M_Ω be the maximal operator defined by

$$(2.8) \quad M_\Omega h(x) = \sup_{r>0} \frac{1}{|B(x,r)|} \int_{B(x,r)} |\Omega(x-y)h(y)|dy.$$

We know from the proof of Lemma 1 in [22] that for $p \in (1, 2]$,

$$(2.9) \quad \left\| \left(\sum_{j \in \mathbb{Z}} |M_\Omega f_j|^2 \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim \left\| \left(\sum_{j \in \mathbb{Z}} |f_j|^2 \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)},$$

provided that $p \in (q', \infty)$ and $w \in A_{p/q'}(\mathbb{R}^n)$, or $p \in (1, q)$ and $w^{-1/(p-1)} \in A_{p'/q'}(\mathbb{R}^n)$. On the other hand, it is easy to verify that

$$|K_t^j * \phi_{j-l} * f_j(x)| \lesssim M_\Omega M f_j(x).$$

The inequality (2.9), together with the weighted vector-valued inequality of M (see Theorem 3.1 in [4]), proves that (2.7) hold when $p \in (1, 2]$, $p \in (q', \infty)$ and $w \in A_{p/q'}(\mathbb{R}^n)$, or $p \in (1, q)$ and $w^{-1/(p-1)} \in A_{p'/q'}(\mathbb{R}^n)$. This, via a standard duality argument, shows that (2.7) holds when $p \in (2, \infty)$, $p \in (1, q)$ and $w^{-1/(p-1)} \in A_{p'/q'}(\mathbb{R}^n)$. \square

Proof of Theorem 2.1. We employ the ideas used in [39]. By Fourier transform estimates (2.2) and (2.5), and the Plancherel theorem, we know that

$$\begin{aligned} \|\widetilde{\mathcal{M}}_\Omega f - \widetilde{\mathcal{M}}_\Omega^l f\|_{L^2(\mathbb{R}^n)}^2 &= \int_1^2 \left\| \left(\sum_{j \in \mathbb{Z}} |F_l f(\cdot, t) - F_j^l f(\cdot, t)|^2 \right)^{\frac{1}{2}} \right\|_{L^2(\mathbb{R}^n)}^2 dt \\ &= \int_1^2 \sum_{j \in \mathbb{Z}} \int_{\mathbb{R}^n} |\widehat{K}_t^j(\xi)|^2 |1 - \widehat{\phi}_{j-l}(\xi)|^2 |\widehat{f}(\xi)|^2 d\xi dt \\ &\lesssim 2^{-\alpha l} \|f\|_{L^2(\mathbb{R}^n)}^2. \end{aligned}$$

Now let p and w be the same as in Theorem 1.3. Recall that \mathcal{M}_Ω is bounded on $L^p(\mathbb{R}^n, w)$ and so is $\widetilde{\mathcal{M}}_\Omega$. Thus, by interpolation with changes of measures of Stein and Weiss [35], it suffices to prove that

$$(2.10) \quad \|\widetilde{\mathcal{M}}_\Omega^l f\|_{L^p(\mathbb{R}^n, w)} \lesssim l \|f\|_{L^p(\mathbb{R}^n, w)}.$$

We now prove (2.10) for the case $p \in (1, q)$ and $w^{-1/(p-1)} \in A_{p'/q'}(\mathbb{R}^n)$. Let $\psi \in C_0^\infty(\mathbb{R}^n)$ be a radial function such that $\text{supp } \psi \subset \{1/4 \leq |\xi| \leq 4\}$ and

$$\sum_{i \in \mathbb{Z}} \psi(2^{-i}\xi) = 1, \quad |\xi| \neq 0.$$

Define the multiplier operator S_i by

$$\widehat{S_i f}(\xi) = \psi(2^{-i}\xi) \widehat{f}(\xi).$$

Set

$$\begin{aligned} E_1 f(x) &= \sum_{m=-\infty}^0 \left(\int_1^2 \sum_j |K_t^j * \phi_{j-l} * (S_{m-j} f)(x)|^2 dt \right)^{\frac{1}{2}}, \\ E_2 f(x) &= \sum_{m=1}^{\infty} \left(\int_1^2 \sum_j |K_t^j * \phi_{j-l} * (S_{m-j} f)(x)|^2 dt \right)^{\frac{1}{2}}. \end{aligned}$$

It then follows that for $f \in \mathcal{S}(\mathbb{R}^n)$,

$$\left\| \left(\int_1^2 \sum_j |K_t^j * \phi_{j-l} * f(x)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n)} \leq \sum_{i=1}^2 \|E_i f\|_{L^p(\mathbb{R}^n)}.$$

We now estimate the term E_1 . By Fourier transform estimate (2.3), we know that

$$\begin{aligned}
 (2.11) \quad & \left\| \left(\int_1^2 \sum_j \left| K_t^j * \phi_{j-l} * (S_{m-j}f)(x) \right|^2 dt \right)^{\frac{1}{2}} \right\|_{L^2(\mathbb{R}^n)}^2 \\
 &= \int_1^2 \int_{\mathbb{R}^n} \sum_{j \in \mathbb{Z}} \left| K_t^j * \phi_{j-l} * (S_{m-j}f)(x) \right|^2 dx dt \\
 &\lesssim \sum_{j \in \mathbb{Z}} \int_{\mathbb{R}^n} |2^j \xi| |\psi(2^{-m+j} \xi)|^2 |\widehat{f}(\xi)|^2 d\xi \\
 &\leq 2^{2m} \|f\|_{L^2(\mathbb{R}^n)}^2.
 \end{aligned}$$

On the other hand, applying the Minkowski inequality, Lemma 2.3 and the weighted Littlewood-Paley theory, we have that

$$\begin{aligned}
 (2.12) \quad & \left\| \left(\int_1^2 \sum_j \left| K_t^j * \phi_{j-l} * (S_{m-j}f)(x) \right|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)}^2 \\
 &\leq \int_1^2 \left(\int_{\mathbb{R}^n} \left(\sum_{j \in \mathbb{Z}} \left| K_t^j * \phi_{j-l} * (S_{m-j}f)(x) \right|^2 \right)^{p/2} w(x) dx \right)^{2/p} dt \\
 &\leq \|f\|_{L^p(\mathbb{R}^n, w)}^2, \quad p \in [2, \infty).
 \end{aligned}$$

To estimate

$$\left\| \left(\int_1^2 \sum_j \left| K_t^j * \phi_{j-l} * (S_{m-j}f)(x) \right|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)}$$

for $p \in (1, 2)$, we consider the mapping \mathcal{F} defined by

$$\mathcal{F} : \{h_j(x)\}_{j \in \mathbb{Z}} \longrightarrow \{K_t^j * \phi_{j-l} * h_j(x)\}.$$

Note that for any $t \in (1, 2)$,

$$|K_t^j * \phi_{j-l} * h_j(x)| \lesssim M M_\Omega h_j(x).$$

We choose $p_0 \in (1, p)$ such that $w^{-1/(p_0-1)} \in A_{p'_0/q'}(\mathbb{R}^n)$. Then by the weighted estimates for M_Ω (see [22]), we have that

$$(2.13) \quad \int_{\mathbb{R}^n} \int_1^2 \sum_{j \in \mathbb{Z}} |K_t^j * \phi_{j-l} * h_j(x)|^{p_0} dt w(x) dx \lesssim \int_{\mathbb{R}^n} \sum_{j \in \mathbb{Z}} |h_j(x)|^{p_0} w(x) dx.$$

Also, we have that

$$\sup_{j \in \mathbb{Z}} \sup_{t \in [1, 2]} |K_t^j * \phi_{j-l} * h_j(x)| \lesssim \sup_{j \in \mathbb{Z}} |h_j(x)|.$$

which implies that for $p_1 \in (1, \infty)$,

$$(2.14) \quad \left\| \sup_{j \in \mathbb{Z}} \sup_{t \in [1, 2]} |K_t^j * \phi_{j-l} * h_j| \right\|_{L^{p_1}(\mathbb{R}^n, w)} \lesssim \left\| \sup_{j \in \mathbb{Z}} |h_j| \right\|_{L^{p_1}(\mathbb{R}^n, w)}.$$

By interpolation, we deduce from the inequalities (2.13) and (2.14) (with $p_0 \in (1, 2)$, $p_1 \in (2, \infty)$ and $1/p = 1/2 + (2 - p_0)/(2p_1)$) that

$$\left\| \left(\int_1^2 \sum_{j \in \mathbb{Z}} |K_t^j * \phi_{j-l} * h_j|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim \left\| \left(\sum_{j \in \mathbb{Z}} |h_j|^2 \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)},$$

and so

$$\begin{aligned} & \left\| \left(\int_1^2 \sum_j \left| K_t^j * \phi_{j-l} * (S_{m-j}f) \right|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \\ & \lesssim \left\| \left(\sum_{j \in \mathbb{Z}} |S_{m-j}f|^2 \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim \|f\|_{L^p(\mathbb{R}^n, w)}, \quad p \in (1, 2). \end{aligned}$$

This, along with (2.12), states that for $p \in (1, q)$,

$$(2.15) \quad \left\| \left(\int_1^2 \sum_j \left| K_t^j * \phi_{j-l} * (S_{m-j}f) \right|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim \|f\|_{L^p(\mathbb{R}^n, w)}.$$

Again by interpolating, the inequalities (2.11) and (2.15) give us that for $p \in (1, q)$,

$$\left\| \left(\int_1^2 \sum_j \left| K_t^j * \phi_{j-l} * (S_{m-j}f)(x) \right|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim 2^{t_p m} \|f\|_{L^p(\mathbb{R}^n, w)}.$$

with $t_p \in (0, 1)$ a constant depending only on p . Therefore,

$$\|E_1 f\|_{L^p(\mathbb{R}^n, w)} \lesssim \|f\|_{L^p(\mathbb{R}^n, w)}.$$

We consider the term E_2 . Again by the Plancherel theorem and the Fourier transform estimates (2.2) and (2.5), we have that

$$\begin{aligned} (2.16) \quad & \left\| \left(\int_1^2 \sum_{j \in \mathbb{Z}} \left| K_t^j * \phi_{j-l} * (S_{m-j}f)(x) \right|^2 dt \right)^{\frac{1}{2}} \right\|_{L^2(\mathbb{R}^n)}^2 \\ & = \int_1^2 \sum_{j \in \mathbb{Z}} \int_{\mathbb{R}^n} |\widehat{K}_t^j(\xi)|^2 |\psi(2^{-m+j}\xi)|^2 |\widehat{f}(\xi)|^2 d\xi dt \\ & \lesssim \sum_{j \in \mathbb{Z}} \int_{\mathbb{R}^n} |2^j \xi|^{-2\alpha} |2^{j-l} \xi|^\alpha |\psi(2^{-m+j}\xi)|^2 |\widehat{f}(\xi)|^2 d\xi \\ & \lesssim 2^{-m\alpha} \|f\|_{L^2(\mathbb{R}^n)}^2. \end{aligned}$$

As in the inequality (2.15), we have that

$$(2.17) \quad \left\| \left(\int_1^2 \sum_{j \in \mathbb{Z}} \left| K_t^j * \phi_{j-l} * (S_{m-j}f)(x) \right|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim \|f\|_{L^p(\mathbb{R}^n, w)}.$$

Interpolating the inequalities (2.16) and (2.17) then shows that

$$\left\| \left(\int_1^2 \sum_{j \in \mathbb{Z}} \left| K_t^j * \phi_{j-l} * (S_{m-j}f)(x) \right|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim 2^{-t_p m} \|f\|_{L^p(\mathbb{R}^n, w)}.$$

This gives the desired estimate for E_2 . Combining the estimates for E_1 and E_2 then yields (2.10) for the case $p \in (1, q)$ and $w^{-1/(p-1)} \in A_{p'/q'}(\mathbb{R}^n)$.

We now prove (2.10) for the case of $p \in (q', \infty)$ and $w \in A_{p/q'}(\mathbb{R}^n)$. By a standard argument, it suffices to prove that

$$(2.18) \quad M^\sharp(\widetilde{\mathcal{M}}_\Omega^l f)(x) \lesssim l M_{q'} f(x),$$

To prove (2.18), let $x \in \mathbb{R}^n$ and Q be a cube containing x . Decompose f as

$$f(y) = f(y)\chi_{4nQ}(y) + f(y)\chi_{\mathbb{R}^n \setminus 4nQ}(y) =: f_1(y) + f_2(y).$$

It is obvious that $\widetilde{\mathcal{M}}_\Omega^l$ is bounded on $L^{q'}(\mathbb{R}^n)$. Thus,

$$(2.19) \quad \frac{1}{|Q|} \int_Q \widetilde{\mathcal{M}}_\Omega^l f_1(y) dy \lesssim \left(\frac{1}{|Q|} \int_Q \{ \widetilde{\mathcal{M}}_\Omega^l f_1(y) \}^{q'} dy \right)^{1/q'} \lesssim M_{q'} f(x).$$

Let $x_0 \in Q$ such that $\widetilde{\mathcal{M}}_\Omega^l f_2(x_0) < \infty$. For $y \in Q$, it follows from Lemma 2.2 that

$$(2.20) \quad \begin{aligned} & \left| \widetilde{\mathcal{M}}_\Omega^l f_2(y) - \widetilde{\mathcal{M}}_\Omega^l f_2(x_0) \right| \\ & \lesssim \left(\int_1^2 \sum_{j \in \mathbb{Z}} |K_t^j * \phi_{j-l} * f_2(y) - K_t^j * \phi_{j-l} * f_2(x_0)|^2 dt \right)^{\frac{1}{2}} \\ & \lesssim \left(\int_1^2 \left(\sum_{j \in \mathbb{Z}} |K_t^j * \phi_{j-l} * f_2(y) - K_t^j * \phi_{j-l} * f_2(x_0)| \right)^2 dt \right)^{\frac{1}{2}} \\ & \lesssim l M_{q'} f(x). \end{aligned}$$

Combining the estimates (2.19) and (2.20) leads to that

$$\inf_{c \in \mathbb{C}} \frac{1}{|Q|} \int_Q |\widetilde{\mathcal{M}}_\Omega^l f(y) - c| dy \lesssim l M_{q'} f(x)$$

and then establish (2.18).

Finally, we see that (2.10) holds for the case of $p \in (1, \infty)$ and $w^{q'} \in A_p(\mathbb{R}^n)$, if we invoke the interpolation argument used in the proof of Theorem 2 in [29]. This completes the proof of Theorem 2.1. \square

3. PROOF OF THEOREM 1.3

We begin with some preliminary lemmas.

Lemma 3.1. *Let Ω be homogeneous of degree zero and belong to $L^1(S^{n-1})$, K_t^j be defined as in (2.1). Then for $l \in \mathbb{N}$, $t \in [1, 2]$, $s \in (1, \infty]$, $j_0 \in \mathbb{Z}_-$ and $y \in \mathbb{R}^n$ with $|y| < 2^{j_0-4}$,*

$$\begin{aligned} & \sum_{j > j_0} \sum_{k \in \mathbb{Z}} 2^{kn/s} \left(\int_{2^k < |x| \leq 2^{k+1}} |K_t^j * \phi_{j-l}(x+y) - K_t^j * \phi_{j-l}(x)|^{s'} dx \right)^{\frac{1}{s'}} \\ & \lesssim 2^{l(n+1)} 2^{-j_0} |y|. \end{aligned}$$

For the proof of Lemma 3.1, see [11].

For $t \in [1, 2]$ and $j \in \mathbb{Z}$, let K_t^j be defined as in (2.1), ϕ and ϕ_l (with $l \in \mathbb{N}$) be the same as in Section 2. For $b \in \text{BMO}(\mathbb{R}^n)$, let $\widetilde{\mathcal{M}}_{\Omega, b}^l$ be the commutator of $\widetilde{\mathcal{M}}_\Omega^l$ defined by

$$\widetilde{\mathcal{M}}_{\Omega, b}^l f(x) = \left(\int_1^2 \sum_{j \in \mathbb{Z}} |F_{j, b}^l f(x, t)|^2 dt \right)^{\frac{1}{2}},$$

with

$$F_{j, b}^l f(x, t) = \int_{\mathbb{R}^n} (b(x) - b(y)) K_t^j * \phi_{j-l}(x-y) f(y) dy.$$

For $j_0 \in \mathbb{Z}$, define the operator $\widetilde{\mathcal{M}}_\Omega^{l, j_0}$ by

$$\widetilde{\mathcal{M}}_\Omega^{l, j_0} f(x) = \left(\int_1^2 \sum_{j \in \mathbb{Z}: j > j_0} |F_{j, b}^l f(x, t)|^2 dt \right)^{\frac{1}{2}},$$

and the commutator $\widetilde{\mathcal{M}}_{\Omega, b}^{l, j_0}$

$$\widetilde{\mathcal{M}}_{\Omega, b}^{l, j_0} f(x) = \left(\int_1^2 \sum_{j \in \mathbb{Z}: j > j_0} |F_{j, b}^l f(x, t)|^2 dt \right)^{\frac{1}{2}},$$

with $b \in \text{BMO}(\mathbb{R}^n)$.

Lemma 3.2. *Let Ω be homogeneous of degree zero and integrable on S^{n-1} . Then for $b \in C_0^\infty(\mathbb{R}^n)$, $l \in \mathbb{N}$, $j_0 \in \mathbb{Z}_-$,*

$$|\widetilde{\mathcal{M}}_{\Omega, b}^{l, j_0} f(x) - \widetilde{\mathcal{M}}_{\Omega, b}^l f(x)| \lesssim 2^{j_0} M M_\Omega f(x).$$

Proof. Let $b \in C_0^\infty(\mathbb{R}^n)$ with $\|\nabla b\|_{L^\infty(\mathbb{R}^n)} = 1$. For $t \in [1, 2]$, by the fact that $\text{supp } K_t^j * \phi_{j-l} \subset \{x : 2^{j-2} \leq |x| \leq 2^{j+2}\}$, it is easy to verify that

$$\begin{aligned} & \sum_{j \leq j_0} \int_{\mathbb{R}^n} |K_t^j * \phi_{j-l}(x-y)| |x-y| |f(y)| dy \\ & \lesssim \sum_{j \leq j_0} \sum_{k \in \mathbb{Z}} 2^k \int_{2^k < |x-y| \leq 2^{k+1}} |K_t^j * \phi_{j-l}(x-y)| |f(y)| dy \\ & \lesssim \sum_{j \leq j_0} \sum_{|k-j| \leq 3} 2^k \int_{2^k < |x-y| \leq 2^{k+1}} |K_t^j * \phi_{j-l}(x-y)| |f(y)| dy \\ & \lesssim 2^{j_0} M_\Omega M f(x). \end{aligned}$$

Thus,

$$\begin{aligned} & \left| \widetilde{\mathcal{M}}_{\Omega, b}^{l, j_0} f(x) - \widetilde{\mathcal{M}}_{\Omega, b}^l f(x) \right|^2 \\ & \leq \sum_{j < j_0} \int_1^2 \left| \int_{\mathbb{R}^n} (b(x) - b(y)) K_t^j * \phi_{j-l}(x-y) f(y) \right|^2 dt \\ & \lesssim \int_1^2 \left(\sum_{j \leq j_0} \int_{\mathbb{R}^n} |x-y| |K_t^j * \phi_{j-l}(x-y) f(y)| dy \right)^2 dt \\ & \lesssim \{2^{j_0} M_\Omega M f(x)\}^2. \end{aligned}$$

The desired conclusion now follows immediately. \square

Let $p, r \in [1, \infty)$, $q \in [1, \infty]$ and w be a weight, $L^p(L^q([1, 2]), l^r; \mathbb{R}^n, w)$ be the space of sequences of functions defined by

$$L^p(L^q([1, 2]), l^r; \mathbb{R}^n, w) = \{ \vec{f} = \{f_k\}_{k \in \mathbb{Z}} : \|\vec{f}\|_{L^p(L^q([1, 2]), l^r; \mathbb{R}^n, w)} < \infty \},$$

with

$$\|\vec{f}\|_{L^p(L^q([1, 2]), l^r; \mathbb{R}^n, w)} = \left\| \left(\int_1^2 \left(\sum_{k \in \mathbb{Z}} |f_k(x, t)|^r \right)^{\frac{q}{r}} dt \right)^{1/q} \right\|_{L^p(\mathbb{R}^n, w)}.$$

With usual addition and scalar multiplication, $L^p(L^q([1, 2]), l^r; \mathbb{R}^n, w)$ is a Banach space.

Lemma 3.3. *Let $p \in (1, \infty)$ and $w \in A_p(\mathbb{R}^n)$, $\mathcal{G} \subset L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)$. Suppose that \mathcal{G} satisfies the following five conditions:*

- (a) \mathcal{G} is bounded, that is, there exists a constant C such that for all $\{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,
- $$\|\vec{f}\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)} \leq C;$$

- (b) for each fixed $\epsilon > 0$, there exists a constant $A > 0$, such that for all $\{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$\left\| \left(\int_1^2 \sum_{k \in \mathbb{Z}} |f_k(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \chi_{\{|\cdot| > A\}}(\cdot) \right\|_{L^p(\mathbb{R}^n, w)} < \epsilon;$$

- (c) for each fixed $\epsilon > 0$ and $N \in \mathbb{N}$, there exists a constant $\varrho > 0$, such that for all $\{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$\left\| \sup_{|h| \leq \varrho} \left(\int_1^2 \sum_{|k| \leq N} |f_k(x, t) - f_k(x+h, t)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} < \epsilon;$$

- (d) for each fixed $\epsilon > 0$ and $N \in \mathbb{N}$, there exists a constant $\sigma \in (0, 1/2)$ such that for all $\{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$\left\| \sup_{|s| \leq \sigma} \left(\int_1^2 \sum_{|k| \leq N} |f_k(\cdot, t+s) - f_k(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} < \epsilon,$$

- (e) for each fixed $D > 0$ and $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for all $\{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$\left\| \left(\int_1^2 \sum_{|k| > N} |f_k(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \chi_{B(0, D)} \right\|_{L^p(\mathbb{R}^n, w)} < \epsilon.$$

Then \mathcal{G} is a strongly pre-compact set in $L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)$.

Proof. We employ the argument used in the proof of [14, Theorem 5], with some refined modifications. Our goal is to prove that, for each fixed $\epsilon > 0$, there exists a $\delta = \delta_\epsilon > 0$ and a mapping Φ_ϵ on $L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)$, such that $\Phi_\epsilon(\mathcal{G}) = \{\Phi_\epsilon(\vec{f}) : \vec{f} \in \mathcal{G}\}$ is a strong pre-compact set in the space $L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)$, and for any $\vec{f}, \vec{g} \in \mathcal{G}$,

$$(3.1) \quad \begin{aligned} \|\Phi_\epsilon(\vec{f}) - \Phi_\epsilon(\vec{g})\|_{L^p(L^2([1,2]), l^2; \mathbb{R}^n, w)} &< \delta \\ \Rightarrow \|\vec{f} - \vec{g}\|_{L^p(L^2([1,2]), l^2; \mathbb{R}^n, w)} &< 8\epsilon. \end{aligned}$$

If we can prove this, then by Lemma 6 in [14], we see that \mathcal{G} is a strongly pre-compact set in $L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)$.

Now let $\epsilon > 0$. We choose $A > 1$ large enough as in assumption (b), $N \in \mathbb{N}$ such that for all $\{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$\left\| \left(\int_1^2 \sum_{|k| > N} |f_k(\cdot, t)|^2 dt \right)^{1/2} \chi_{B(0, 2A)} \right\|_{L^p(\mathbb{R}^n, w)} < \epsilon.$$

Let $\varrho \in (0, 1/2)$ small enough as in assumption (c) and $\sigma \in (0, 1/2)$ small enough such that (d) holds true. Let Q be the largest cube centered at the origin such that $2Q \subset B(0, \varrho)$, Q_1, \dots, Q_J be J copies of Q such that they are non-overlapping, and $\overline{B(0, A)} \subset \cup_{j=1}^J Q_j \subset B(0, 2A)$. Let $I_1, \dots, I_L \subset [1, 2]$ be non-overlapping intervals with same length $|I|$, such that $|s - t| \leq \sigma$ for all $s, t \in I_j$ ($j = 1, \dots, L$)

and $\cup_{j=1}^N I_j = [1, 2]$. Define the mapping Φ_ϵ on $L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)$ by

$$\begin{aligned} \Phi_\epsilon(\vec{f})(x, t) = & \left\{ \dots, 0, \dots, 0, \sum_{i=1}^J \sum_{j=1}^L m_{Q_i \times I_j}(f_{-N}) \chi_{Q_i \times I_j}(x, t), \right. \\ & \sum_{i=1}^J \sum_{j=1}^L m_{Q_i \times I_j}(f_{-N+1}) \chi_{Q_i \times I_j}(x, t), \dots, \\ & \left. \sum_{i=1}^J \sum_{j=1}^L m_{Q_i \times I_j}(f_N) \chi_{Q_i \times I_j}(x, t), 0, \dots \right\}, \end{aligned}$$

where and in the following,

$$m_{Q_i \times I_j}(f_k) = \frac{1}{|Q_i|} \frac{1}{|I_j|} \int_{Q_i \times I_j} f_k(x, t) dx dt.$$

We claim that Φ_ϵ is bounded on $L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)$. In fact, if $p \in [2, \infty)$, we have by the Hölder inequality that

$$\begin{aligned} |m_{Q_i \times I_j}(f_k)| & \leq \left(\frac{1}{|Q_i||I_j|} \int_{I_j \times Q_i} |f_k(y, t)|^p w(y) dy dt \right)^{\frac{1}{p}} \\ & \quad \times \left(\frac{1}{|Q_i|} \int_{Q_i} w^{-\frac{1}{p-1}}(y) dy \right)^{\frac{1}{p'}}, \end{aligned}$$

and

$$\begin{aligned} & \sum_{|k| \leq N} \left(\frac{1}{|Q_i||I_j|} \int_{I_j} \int_{Q_i} |f_k(y, t)|^p w(y) dy dt \right)^{2/p} \\ & \lesssim N^{1-2/p} \left(\sum_{|k| \leq N} \frac{1}{|Q_i||I_j|} \int_{I_j \times Q_i} |f_k(y, t)|^p w(y) dy dt \right)^{2/p}. \end{aligned}$$

Therefore,

$$\begin{aligned} \|\Phi_\epsilon(\vec{f})\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)}^p & = \sum_{i=1}^J \sum_{j=1}^L \int_{I_j} \int_{Q_i} \left(\sum_{|k| \leq N} |m_{Q_i \times I_j}(f_k)|^2 \right)^{p/2} w(x) dx dt \\ & \lesssim N^{p/2-1} \sum_{i=1}^J \sum_{j=1}^L \int_{I_j} \int_{Q_i} \sum_{|k| \leq N} |f_k(y, t)|^p w(y) dy dt \\ & \leq N^{p/2} \sum_{i=1}^J \sum_{j=1}^L \int_{I_j} \int_{Q_i} \left\{ \sum_{|k| \leq N} |f_k(y, t)|^2 \right\}^{\frac{p}{2}} w(y) dy dt \\ & \leq N^{p/2} \|\vec{f}\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)}^p. \end{aligned}$$

On the other hand, for $p \in (1, 2)$ and $w \in A_p(\mathbb{R}^n)$, we choose $\gamma \in (0, 1)$ such that $w \in A_{p-\gamma}(\mathbb{R}^n)$. Note that

$$\sup_{-N \leq k \leq N} \sup_{t \in [1, 2]} \left| \sum_{i=1}^J \sum_{j=1}^L m_{Q_i \times I_j}(f_k) \chi_{Q_i \times I_j}(x, t) \right| \lesssim \sup_{k \in \mathbb{Z}} \sup_{t \in [1, 2]} |f_k(x, t)|,$$

which implies that for $p_1 \in (1, \infty)$,

$$(3.2) \quad \|\Phi_\epsilon(\vec{f})\|_{L^{p_1}(L^\infty([1, 2]), l^\infty; \mathbb{R}^n, w)} \lesssim \|\vec{f}\|_{L^{p_1}(L^\infty([1, 2]), l^\infty; \mathbb{R}^n, w)}.$$

We also have that for $p_0 = p - \gamma$,

$$|m_{Q_i \times I_j}(f_k)| \leq \left(\frac{1}{|Q_i||I_j|} \int_{I_j} \int_{Q_i} |f_k(y, t)|^{p_0} w(y) dy dt \right)^{\frac{1}{p_0}} \left(\frac{1}{|Q_i|} \int_{Q_i} w^{-\frac{1}{p_0-1}}(y) dy \right)^{\frac{1}{p_0}},$$

and so

$$(3.3) \quad \|\Phi_\epsilon(\vec{f})\|_{L^{p_0}(L^{p_0}([1, 2]), l^{p_0}; \mathbb{R}^n, w)} \lesssim \|\vec{f}\|_{L^{p_0}(L^{p_0}([1, 2]), l^{p_0}; \mathbb{R}^n, w)}.$$

By interpolation, we can deduce from (3.2) and (3.3) that in this case

$$\|\Phi_\epsilon(\vec{f})\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)} \lesssim \|\vec{f}\|_{L^p(L^2([0, 1]), l^2, \mathbb{R}^n, w)}^p.$$

Our claim then follows directly, and so $\Phi_\epsilon(\mathcal{G}) = \{\Phi_\epsilon(\vec{f}) : \vec{f} \in \mathcal{G}\}$ is strongly pre-compact in $L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)$.

We now verify (3.1). Denote $\mathcal{D} = \cup_{i=1}^J Q_i$ and write

$$\begin{aligned} & \|\vec{f}\chi_{\mathcal{D}} - \Phi_\epsilon(\vec{f})\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)} \\ & \leq \left\| \left(\int_1^2 \sum_{|k| \leq N} \left| f_k(\cdot, t)\chi_{\mathcal{D}} - \sum_{i=1}^J \sum_{j=1}^L m_{Q_i \times I_j}(f_k)\chi_{Q_i \times I_j}(x, t) \right|^2 dt \right)^{1/2} \right\|_{L^p(\mathbb{R}^n, w)} \\ & \quad + \left\| \left(\int_1^2 \sum_{|k| > N} |f_k(\cdot, t)|^2 \right)^{\frac{1}{2}} \chi_{B(0, 2A)} \right\|_{L^p(\mathbb{R}^n, w)}. \end{aligned}$$

Noting that for $x \in Q_i$ with $1 \leq i \leq J$,

$$\begin{aligned} & \left\{ \int_1^2 \sum_{|k| \leq N} \left| f_k(x, t)\chi_{\mathcal{D}}(x) - \sum_{u=1}^J \sum_{v=1}^L m_{Q_u \times I_v}(f_k)\chi_{Q_u \times I_v}(x, t) \right|^2 dt \right\}^{\frac{1}{2}} \\ & \lesssim |Q|^{-1/2} |I|^{-1/2} \left\{ \sum_{j=1}^L \int_{I_j} \int_{Q_i} \int_{I_j} \sum_{|k| \leq N} |f_k(x, t) - f_k(y, s)|^2 dy ds dt \right\}^{\frac{1}{2}} \\ & \lesssim |Q|^{-1/2} \left\{ \int_{2Q} \int_1^2 \sum_{|k| \leq N} |f_k(x, s) - f_k(x+h, s)|^2 ds dh \right\}^{\frac{1}{2}} \\ & \quad + |I|^{-1/2} \left\{ \sum_{j=1}^L \int_{I_j} \int_{I_j} \sum_{|k| \leq N} |f_k(x, t) - f_k(x, s)|^2 dt ds \right\}^{\frac{1}{2}} \\ & \lesssim \sup_{|h| \leq \varrho} \left(\int_1^2 \sum_{|k| \leq N} |f_k(x, t) - f_k(x+h, t)|^2 dt \right)^{\frac{1}{2}} \\ & \quad + \sup_{|s| \leq \sigma} \left(\int_1^2 \sum_{|k| \leq N} |f_k(x, t+s) - f_k(x, t)|^2 dt \right)^{\frac{1}{2}}, \end{aligned}$$

we then get that

$$\sum_{i=1}^J \int_{Q_i} \left\{ \int_1^2 \sum_{|k| \leq N} \left| f_k(x, t) - \sum_{l=1}^J m_{Q_l}(f_k)\chi_{Q_l}(x) \right|^2 dt \right\}^{p/2} w(x) dx \lesssim 2\epsilon.$$

It then follows from the assumption (b) that for all $\vec{f} \in \mathcal{G}$,

$$\begin{aligned} \|\vec{f} - \Phi_\epsilon(\vec{f})\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)} &\leq \|\vec{f}\chi_{\mathcal{D}} - \Phi_\epsilon(\vec{f})\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)} \\ &+ \left\| \left(\int_1^2 \sum_{k \in \mathbb{Z}} |f_k(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \chi_{\{|\cdot| > A\}}(\cdot) \right\|_{L^p(\mathbb{R}^n, w)} \\ &< 3\epsilon. \end{aligned}$$

Noting that

$$\begin{aligned} \|\vec{f} - \vec{g}\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)} &\leq \|\vec{f} - \Phi_\epsilon(\vec{f})\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)} \\ &+ \|\Phi_\epsilon(\vec{f}) - \Phi_\epsilon(\vec{g})\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)} \\ &+ \|\vec{g} - \Phi_\epsilon(\vec{g})\|_{L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)}, \end{aligned}$$

we then get (3.1) and finish the proof of Lemma 3.3. \square

Proof of Theorem 1.3. Let $j_0 \in \mathbb{Z}_-$, $b \in C_0^\infty(\mathbb{R}^n)$ with $\text{supp } b \subset B(0, R)$, p and w be the same as in Theorem 1.3. Without loss of generality, we may assume that $\|b\|_{L^\infty(\mathbb{R}^n)} + \|\nabla b\|_{L^\infty(\mathbb{R}^n)} = 1$. We claim that

(i) for each fixed $\epsilon > 0$, there exists a constant $A > 0$ such that

$$\left\| \left(\int_1^2 \sum_{j \in \mathbb{Z}} |F_{j,b}^l f(x, t)|^2 dt \right)^{1/2} \chi_{\{|\cdot| > A\}}(\cdot) \right\|_{L^p(\mathbb{R}^n, w)} < \epsilon \|f\|_{L^p(\mathbb{R}^n, w)};$$

(ii) for $s \in (1, \infty)$,

$$\begin{aligned} (3.4) \quad &\left(\int_1^2 \sum_{j > j_0} |F_{j,b}^l f(x, t) - F_{j,b}^l f(x+h, t)|^2 dt \right)^{1/2} \\ &\lesssim 2^{-j_0} |h| \left(\widetilde{\mathcal{M}}_\Omega^{l, j_0} f(x) + 2^{l(n+1)} M_s f(x) \right); \end{aligned}$$

(iii) for each $\epsilon > 0$ and $N \in \mathbb{N}$, there exists a constant $\sigma \in (0, 1/2)$ such that

$$\begin{aligned} (3.5) \quad &\left\| \sup_{|s| \leq \sigma} \left(\int_1^2 \sum_{|j| \leq N} |F_{j,b}^l f(x, s+t) - F_{j,b}^l f(x, t)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \\ &< \epsilon \|f\|_{L^p(\mathbb{R}^n, w)}; \end{aligned}$$

(iv) for each fixed $D > 0$ and $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$(3.6) \quad \left\| \left(\int_1^2 \sum_{j > N} |F_{j,b}^l(\cdot, t)|^2 dt \right)^{1/2} \chi_{B(0, D)} \right\|_{L^p(\mathbb{R}^n, w)} < \epsilon \|f\|_{L^p(\mathbb{R}^n, w)}.$$

We now prove claim (i). Let $t \in [1, 2]$. For each fixed $x \in \mathbb{R}^n$ with $|x| > 4R$, observe that $\text{supp } K_t^j * \phi_{j-l} \subset \{2^{j-2} \leq |y| \leq 2^{j+2}\}$, and $\int_{|z| < R} |K_t^j * \phi_{j-l}(x-z)| dz \neq 0$ only if $2^j \approx |x|$. A trivial computation shows that

$$\begin{aligned} \int_{|z| < R} |K_t^j * \phi_{j-l}(x-z)| dz &\lesssim \left(\int_{|z| < R} |K_t^j * \phi_{j-l}(x-z)|^2 dz \right)^{\frac{1}{2}} R^{\frac{n}{2}} \\ &\lesssim \left(\int_{\frac{|x|}{2} \leq |z| < 2|x|} |K_t^j * \phi_{j-l}(z)|^2 dz \right)^{\frac{1}{2}} R^{\frac{n}{2}} \\ &\lesssim \|K_t^j\|_{L^1(S^{n-1})} \|\phi_{j-l}\|_{L^2(\mathbb{R}^n)} R^{\frac{n}{2}} \\ &\lesssim 2^{nl/2} |x|^{-\frac{n}{2}} R^{\frac{n}{2}}. \end{aligned}$$

On the other hand, we have that

$$\begin{aligned} & \sum_{j \in \mathbb{Z}} \left(\int_{|y| < R} |K_t^j * \phi_{j-l}(x-y)| |f(y)|^s dy \right)^{\frac{1}{s}} \\ &= \sum_{j \in \mathbb{Z}: 2^j \approx |x|} \left(\int_{|x|/2 \leq |y-x| \leq 2|x|} |K_t^j * \phi_{j-l}(x-y)| |f(y)|^s dy \right)^{\frac{1}{s}} \\ &\lesssim \left(M_\Omega M(|f|^s)(x) \right)^{1/s}. \end{aligned}$$

Another application of the Hölder inequality then yields

$$\begin{aligned} (3.7) \quad \sum_{j \in \mathbb{Z}} |F_{j,b}^l f(x,t)|^2 &\lesssim \sum_{j \in \mathbb{Z}} \left(\int_{|y| < R} |K_t^j * \phi_{j-l}(x-y)| |f(y)|^s dy \right)^{2/s} \\ &\quad \times \left(\int_{|y| < R} |K_t^j * \phi_{j-l}(x-y)| dy \right)^{2/s'} \\ &\lesssim 2^{\frac{2l}{s'}} |x|^{-\frac{n}{s'}} R^{\frac{n}{s'}} \left(M_\Omega M(|f|^s)(x) \right)^{2/s}. \end{aligned}$$

This, in turn leads to our claim (i).

We turn our attention to claim (ii). Write

$$|F_{j,b}^l f(x,t) - F_{j,b}^l f(x+h,t)| \leq |b(x) - b(x+h)| |F_{j,b}^l f(x,t)| + J_j^l f(x,t),$$

with

$$J_j^l f(x,t) = \left| \int_{\mathbb{R}^n} (K_t^j * \phi_{j-l}(x-y) - K_t^j * \phi_{j-l}(x+h-y)) (b(x+h) - b(y)) f(y) dy \right|.$$

It follows from Lemma 3.1 that

$$\begin{aligned} \left(\sum_{j > j_0} |J_j^l f(x,t)|^2 \right)^{\frac{1}{2}} &\lesssim \sum_{j > j_0} \int_{\mathbb{R}^n} |K_t^j * \phi_{j-l}(x-y) - K_t^j * \phi_{j-l}(x+h-y)| |f(y)| dy \\ &\lesssim 2^{l(n+1)} |h| 2^{-j_0} M_s f(x). \end{aligned}$$

Therefore,

$$\begin{aligned} &\left(\int_1^2 \sum_{j > j_0} |F_{j,b}^l f(x,t) - F_{j,b}^l f(x+h,t)|^2 dt \right)^{\frac{1}{2}} \\ &\lesssim |h| \widetilde{\mathcal{M}}_\Omega^{l,j_0} f(x) + 2^{l(n+1)} 2^{-j_0} |h| M_s f(x). \end{aligned}$$

We now verify claim (iii). For each fixed $\sigma \in (0, 1/2)$ and $t \in [1, 2]$, let

$$U_{t,\sigma}^j(z) = \frac{1}{2^j} \frac{|\Omega(z)|}{|z|^{n-1}} \chi_{\{2^j(t-\sigma) \leq |z| \leq 2^j t\}} + \frac{1}{2^j} \frac{|\Omega(z)|}{|z|^{n-1}} \chi_{\{2^{j+1}t \leq |z| \leq 2^{j+1}(t+\sigma)\}},$$

and

$$G_{l,t,\sigma}^j f(x) = \int_{\mathbb{R}^n} (U_{t,\sigma}^j * |\phi_{j-l}|)(x-y) |f(y)| dy.$$

Note that for $t \in [1, 2]$,

$$\|U_{t,\sigma}^j * |\phi_{j-l}|\|_{L^1(\mathbb{R}^n)} \lesssim \sigma, \quad \sup_{|j| \leq N} \sup_{t \in [1, 2]} |G_{l,t,\sigma}^j f(x)| \lesssim M M_\Omega f(x).$$

Thus,

$$(3.8) \quad \left\| \sup_{|j| \leq N} \sup_{t \in [1, 2]} |G_{l,t,\sigma}^j f| \right\|_{L^\infty(\mathbb{R}^n)} \lesssim \sigma \|f\|_{L^\infty(\mathbb{R}^n)},$$

and

$$(3.9) \quad \left\| \sup_{|j| \leq N} \sup_{t \in [1, 2]} |G_{l, t, \sigma}^j f| \right\|_{L^p(\mathbb{R}^n, w)} \lesssim \|MM_\Omega\|_{L^p(\mathbb{R}^n, w)} \lesssim \|f\|_{L^p(\mathbb{R}^n, w)}.$$

Interpolating the estimates (3.8) and (3.9) shows that if $p_1 \in (p, \infty)$,

$$(3.10) \quad \left\| \sup_{|j| \leq N} \sup_{t \in [1, 2]} |G_{l, t, \sigma}^j f| \right\|_{L^{p_1}(\mathbb{R}^n, w)} \lesssim \sigma^{1-p/p_1} \|f\|_{L^{p_1}(\mathbb{R}^n, w)}.$$

On the other hand, if $p_0 \in (1, p)$, it then follows from the weighted estimates M and M_Ω that

$$(3.11) \quad \int_{\mathbb{R}^n} \int_1^2 \sum_{|j| \leq N} |G_{l, t, \sigma}^j f(x)|^{p_0} dt w(x) dx \lesssim N \|f\|_{L^{p_0}(\mathbb{R}^n, w)}^{p_0}.$$

Choosing $p_1 \in (2, \infty)$ such that $1/p = 1/2 + (2 - p_0)/(2p_1)$ in (3.10), we get from (3.10) and (3.11) that for $p \in (1, 2)$,

$$(3.12) \quad \left\| \left(\int_1^2 \sum_{|j| \leq N} |G_{l, t, \sigma}^j f(x)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim N \sigma^{\tau_1} \|f\|_{L^p(\mathbb{R}^n, w)},$$

with $\tau_1 \in (0, 1)$ a constant. If $p \in [2, \infty)$, we obtain from the Minkowski inequality and the Young inequality that

$$(3.13) \quad \begin{aligned} & \left\| \left(\int_1^2 \sum_{|j| \leq N} |G_{l, t, \sigma}^j f(x)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)}^2 \\ & \lesssim \left\{ \int_{\mathbb{R}^n} \left(\int_1^2 \left(\sum_{|j| \leq N} \int_{\mathbb{R}^n} (U_{l, t, \sigma}^j * |\phi_{j-l}|)(x-y) |f(y)| dy \right)^2 dt \right)^{\frac{p}{2}} w(x) dx \right\}^{\frac{2}{p}} \\ & \lesssim \int_1^2 \left\{ \sum_{|j| \leq N} \left(\int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} (U_{l, t, \sigma}^j * |\phi_{j-l}|)(x-y) |f(y)| dy \right)^p w(x) dx \right)^{\frac{1}{p}} \right\}^2 dt \\ & \lesssim N^2 \|f\|_{L^p(\mathbb{R}^n, w)}^2. \end{aligned}$$

Also, we have that

$$(3.14) \quad \begin{aligned} & \left\{ \int_{\mathbb{R}^n} \left(\int_1^2 \sum_{|j| \leq N} \left(\int_{\mathbb{R}^n} (U_{l, t, \sigma}^j * |\phi_{j-l}|)(x-y) |f(y)| dy \right)^2 dt \right)^{\frac{p}{2}} dx \right\}^{\frac{2}{p}} \\ & \lesssim \int_1^2 \left\{ \sum_{|j| \leq N} \left\| (U_{l, t, \sigma}^j * |\phi_{j-l}| * |f|) \right\|_{L^p(\mathbb{R}^n)} \right\}^2 dt \\ & \lesssim (2N\sigma)^2 \|f\|_{L^p(\mathbb{R}^n)}^2, \quad p \in [2, \infty). \end{aligned}$$

The inequalities (3.13) and (3.14), via interpolation with changes of measures, give us that for $p \in [2, \infty)$,

$$(3.15) \quad \left\| \left(\int_1^2 \sum_{|j| \leq N} |G_{l, t, \sigma}^j f(x)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim N \sigma^{\tau_2} \|f\|_{L^p(\mathbb{R}^n, w)},$$

with $\tau_2 \in (0, 1)$ a constant. Since

$$\sup_{|s| \leq \sigma} |F_{j, b}^l f(x, t) - F_{j, b}^l f(x, t+s)| \leq G_{l, t, \sigma}^j f(x),$$

our claim (iii) now follow from (3.12) and (3.15) immediately if we choose $\sigma = \epsilon/(2N)$.

It remains to prove (iv). Let $D > 0$ and $N \in \mathbb{N}$ such that $2^{N-2} > D$. Then for $j > N$ and $x \in \mathbb{R}^n$ with $|x| \leq D$,

$$\begin{aligned} \int_{\mathbb{R}^n} |K_t^j * \phi_{j-l}(x-y)f(y)| dy &\leq \int_{\mathbb{R}^n} |K_t^j * \phi_{j-l}(x-y)f(y)| \chi_{\{|y| \leq 2^{j+3}\}} dy \\ &\lesssim \int_{|y| \leq 2^{j+3}} |f(y)| dy \|K_t^j\|_{L^1(\mathbb{R}^n)} \|\phi_{j-l}\|_{L^\infty(\mathbb{R}^n)} \\ &\lesssim 2^{nl} 2^{-nj/p} \|f\|_{L^p(\mathbb{R}^n)}. \end{aligned}$$

Therefore,

$$(3.16) \quad \left\| \left(\int_1^2 \sum_{j>N} |F_{j,b}^l f(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \chi_{B(0,D)} \right\|_{L^p(\mathbb{R}^n)} \lesssim 2^{nl} \left(\frac{D}{2N}\right)^{n/p} \|f\|_{L^p(\mathbb{R}^n)}.$$

It is obvious that

$$(3.17) \quad \left\| \left(\int_1^2 \sum_{j>N} |F_{j,b}^l f(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \chi_{B(0,D)} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim l \|f\|_{L^p(\mathbb{R}^n, w)}.$$

Interpolating the inequalities (3.16) and (3.17) yields

$$\left\| \left(\int_1^2 \sum_{j>N} |F_{j,b}^l f(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \chi_{B(0,D)} \right\|_{L^p(\mathbb{R}^n, w)} \lesssim 2^{\tau_3 nl} \left(\frac{D}{2N}\right)^{\frac{\tau_3 n}{p}} \|f\|_{L^p(\mathbb{R}^n, w)},$$

with $\tau_3 \in (0, 1)$ a constant depending only on w . The claim (iv) now follows directly.

We can now conclude the proof of Theorem 1.3. Let $p \in (1, \infty)$. Note that

$$\widetilde{\mathcal{M}}_{\Omega, b}^{l, j_0} f(x) \leq \widetilde{\mathcal{M}}_{\Omega, b}^l f(x),$$

and so $\widetilde{\mathcal{M}}_{\Omega, b}^{l, j_0}$ is bounded on $L^p(\mathbb{R}^n, w)$. Our claims (i)-(iv), via Lemma 3.3, prove that for $b \in C_0^\infty(\mathbb{R}^n)$, $l \in \mathbb{N}$ and $j_0 \in \mathbb{Z}_-$, the operator $\mathcal{F}_{j_0}^l$ defined by

$$(3.18) \quad \mathcal{F}_{j_0}^l : f(x) \rightarrow \{\dots, 0, \dots, F_{j_0, b}^l f(x, t), F_{j_0+1, b}^l f(x, t), \dots\}$$

is compact from $L^p(\mathbb{R}^n, w)$ to $L^p(L^2([1, 2]), l^2; \mathbb{R}^n, w)$. Thus, $\widetilde{\mathcal{M}}_{\Omega, b}^{l, j_0}$ is completely continuous on $L^p(\mathbb{R}^n, w)$. This, via Lemma 3.2 and Theorem 2.1, shows that for $b \in C_0^\infty(\mathbb{R}^n)$, $\widetilde{\mathcal{M}}_{\Omega, b}$ is completely continuous on $L^p(\mathbb{R}^n, w)$. Note that

$$|\mathcal{M}_{\Omega, b} f_k(x) - \mathcal{M}_{\Omega, b} f(x)| \lesssim \mathcal{M}_{\Omega, b}(f_k - f)(x) \lesssim \widetilde{\mathcal{M}}_{\Omega, b}(f_k - f)(x).$$

Thus, for $b \in C_0^\infty(\mathbb{R}^n)$, $\mathcal{M}_{\Omega, b}$ is completely continuous on $L^p(\mathbb{R}^n, w)$. Recalling that $\mathcal{M}_{\Omega, b}$ is bounded on $L^p(\mathbb{R}^n, w)$ with bound $C\|b\|_{\text{BMO}(\mathbb{R}^n)}$, we obtain that for $b \in \text{CMO}(\mathbb{R}^n)$, $\mathcal{M}_{\Omega, b}$ is completely continuous on $L^p(\mathbb{R}^n, w)$. \square

4. PROOF OF THEOREM 1.5

The following lemma will be useful in the proof of Theorem 1.5, and is of independent interest.

Lemma 4.1. *Let $u \in (1, \infty)$, $m \in \mathbb{N} \cup \{0\}$, S be a sublinear operator which satisfies that*

$$|Sf(x)| \leq \int_{\mathbb{R}^n} |b(x) - b(y)|^m |W(x-y)f(y)| dy,$$

with $b \in \text{BMO}(\mathbb{R}^n)$, and

$$(4.1) \quad \sup_{R>0} R^{n/u} \left(\int_{R \leq |x| \leq 2R} |W(x)|^{u'} dx \right)^{1/u'} \lesssim 1.$$

(a) *Let $p \in (u, \infty)$, $\lambda \in (0, 1)$ and $w \in A_{p/u}(\mathbb{R}^n)$. If S is bounded on $L^p(\mathbb{R}^n, w)$ with bound $D \|b\|_{\text{BMO}(\mathbb{R}^n)}^m$, then for some $\varepsilon \in (0, 1)$,*

$$\|Sf\|_{L^{p,\lambda}(\mathbb{R}^n, w)} \lesssim (D + D^\varepsilon) \|b\|_{\text{BMO}(\mathbb{R}^n)}^m \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)}.$$

(b) *Let $p \in (1, u)$, $w^r \in A_1(\mathbb{R}^n)$ for some $r \in (u, \infty)$ and $\lambda \in (0, 1 - r'/u')$. If S is bounded on $L^p(\mathbb{R}^n, w)$ with bound D , then for some $\varepsilon \in (0, 1)$,*

$$\|Sf\|_{L^{p,\lambda}(\mathbb{R}^n)} \lesssim (D + D^\varepsilon) \|b\|_{\text{BMO}(\mathbb{R}^n)}^m \|f\|_{L^{p,\lambda}(\mathbb{R}^n)}.$$

Proof. For simplicity, we only consider the case of $m = 1$ and $\|b\|_{\text{BMO}(\mathbb{R}^n)} = 1$. For fixed ball B and $f \in L^{p,\lambda}(\mathbb{R}^n, w)$, decompose f as

$$f(y) = f(y)\chi_{2B}(y) + \sum_{k=1}^{\infty} f(y)\chi_{2^{k+1}B \setminus 2^k B}(y) = \sum_{k=0}^{\infty} f_k(y).$$

It is obvious that

$$\int_B |Sf_0(y)|^p w(y) dy \lesssim D^p \int_{2B} |f(y)|^p w(y) dy \lesssim D^p \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)}^p \{w(B)\}^\lambda.$$

Let $\theta \in (1, p/u)$ such that $w \in A_{p/(\theta u)}(\mathbb{R}^n)$. For each $k \in \mathbb{N}$, let $S_k f(x) = S(f\chi_{2^{k+1}B \setminus 2^k B})(x)$. Then S_k is also sublinear. We have by the Hölder inequality that for each $x \in B$,

$$\begin{aligned} |S_k f(x)| &\lesssim |b(x) - m_B(b)| \|f_k\|_{L^u(\mathbb{R}^n)} \left(\int_{2^k B} |W(x-y)|^{u'} dy \right)^{1/u'} \\ &\quad + \|(b - m_B(b))f_k\|_{L^u(\mathbb{R}^n)} \left(\int_{2^k B} |W(x-y)|^{u'} dy \right)^{1/u'} \\ &\lesssim |b(x) - m_B(b)| \|f_k\|_{L^p(\mathbb{R}^n, w)} \left(\int_{2^k B} w^{-\frac{1}{p/u-1}}(y) dy \right)^{\frac{1}{u(p/u)'}} |2^k B|^{-\frac{1}{u}} \\ &\quad + \left(\int_{2^{k+1}B} |b(y) - m_B(b)|^{p\theta'} dy \right)^{1/(p\theta')} \|f_k\|_{L^p(\mathbb{R}^n, w)} \\ &\quad \times \left(\int_{2^k B} w^{-\frac{1}{p/(\theta u)-1}}(y) dy \right)^{\frac{1}{u(p/(\theta u))'}} |2^k B|^{-\frac{1}{u}}, \end{aligned}$$

here, $m_B(b)$ denotes the mean value of b on B . It follows from the John-Nirenberg inequality that

$$\left(\int_{2^{k+1}B} |b(y) - m_B(b)|^{p\theta'} dy \right)^{\frac{1}{p\theta'}} \lesssim k |2^k B|^{\frac{1}{p\theta'}}.$$

Therefore, for $q \in (1, \infty)$ and $k \in \mathbb{N}$, we have

$$(4.2) \quad \|S_k f\|_{L^q(B, w)} \lesssim k \{w(B)\}^{\frac{1}{q} - \frac{1}{p}} \left(\frac{w(B)}{w(2^k B)} \right)^{1/p} \|f_k\|_{L^p(\mathbb{R}^n, w)}.$$

On the other hand, we deduce from the $L^p(\mathbb{R}^n, w)$ boundedness of S that

$$(4.3) \quad \int_B |S_k f(y)|^p w(y) dy \lesssim D^p \int_{2^k B} |f(x)|^p w(x) dx$$

We then get from (4.2) (with $q = p$) and (4.3) that for $\sigma \in (0, 1)$,

$$(4.4) \quad \int_B |S_k f(y)|^p w(y) dy \lesssim k^p D^{p(1-\sigma)} \left(\frac{w(B)}{w(2^k B)} \right)^\sigma \int_{2^k B} |f(x)|^p w(x) dx$$

Recall that $w \in A_{p/u}(\mathbb{R}^n)$. Thus, there exists a constant $\tau \in (0, 1)$,

$$\frac{w(B)}{w(2^k B)} \lesssim \left(\frac{|B|}{|2^k B|} \right)^\tau,$$

see [24]. For fixed $\lambda \in (0, 1)$, we choose σ sufficiently close to 1 such that $0 < \lambda < \sigma$. It then follows from (4.4) that

$$\begin{aligned} \sum_{k=0}^{\infty} \left(\int_B |S_k f(y)|^p dy \right)^{\frac{1}{p}} &\lesssim D^{1-\sigma} \{w(B)\}^{\frac{\lambda}{p}} \sum_{k=0}^{\infty} (k+1) 2^{-kn\tau(\sigma-\lambda)/p} \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)} \\ &\lesssim D^{1-\sigma} \{w(B)\}^{\lambda/p} \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)}. \end{aligned}$$

This leads to the conclusion (a).

Now we turn our attention to conclusion (b). From (4.1), it is obvious that for $y \in 2^{k+1}B \setminus 2^k B$,

$$\int_B |W(x-y)| |b(x) - m_B(b)| w(x) dx \lesssim |2^k B|^{-1/u} |B|^{\frac{1}{u\vartheta}} \left(\int_B w^{u\vartheta}(x) dx \right)^{\frac{1}{u\vartheta}},$$

with $\vartheta \in (1, \infty)$ small enough such that $w^{u\vartheta} \in A_1(\mathbb{R}^n)$. This, in turn implies that

$$\begin{aligned} &\int_B \int_{2^{k+1}B \setminus 2^k B} |W(x-y)h(y)| dy |b(x) - m_B(b)| w(x) dx \\ &\lesssim 2^{kn/u'} \frac{w(B)}{w(2^k B)} \int_{2^k B} h(y) w(y) dy. \end{aligned}$$

Therefore, for $s \in (1, \infty)$,

$$\begin{aligned} (4.5) \quad \int_B |S_k f(x)| w(x) dx &\lesssim 2^{kn/u'} \frac{w(B)}{w(2^k B)} \int_{2^k B} |f(x)| w(x) dx \\ &\quad + 2^{kn/u'} \frac{w(B)}{w(2^k B)} \int_{2^k B} |f(x)| |b(x) - m_B(b)| w(x) dx \\ &\lesssim k 2^{\frac{kn}{u'}} \frac{w(B)}{w(2^{k+1}B)} \left(\int_{2^k B} |f(x)|^s w(x) dx \right)^{\frac{1}{s}} \{w(2^k B)\}^{\frac{1}{s'}}. \end{aligned}$$

Also, we get by (4.2) that for $q \in (u, \infty)$ and $\theta \in (0, 1)$ with $\theta q \in (u, \infty)$,

$$(4.6) \quad \|S_k f\|_{L^q(B, w)} \lesssim k \{w(B)\}^{\frac{1}{q} - \frac{1}{\theta q}} \left(\frac{w(B)}{w(2^k B)} \right)^{\frac{1}{\theta q}} \|f\|_{L^{\theta q}(2^{k+1}B, w)}.$$

For $p \in (1, \infty)$, we choose $q \in (u, \infty)$ and $\theta \in (0, 1)$, $s \in (1, \infty)$ which is close to 1 sufficiently such that $1/p = t + (1-t)/q$ and $1/p = t/s + (1-t)/(\theta q)$, with $t \in (0, 1/p)$. By interpolating, we obtain from the inequalities (4.5) and (4.6) that

$$\|S_k f\|_{L^p(\mathbb{R}^n, w)} \lesssim k 2^{\frac{kn}{p'u'}} \left(\frac{w(B)}{w(2^k B)} \right)^{1/p} \|f\|_{L^p(2^k B, w)}.$$

The fact that $w^r \in A_1(\mathbb{R}^n)$ tells us that

$$\frac{w(B)}{w(2^k B)} \lesssim 2^{-kn(r-1)/r},$$

see [24, p. 306]. This, together with the fact that S is bounded on $L^p(\mathbb{R}^n, w)$ with bound D , gives us that for any $\omega \in (0, 1)$,

$$\begin{aligned} \left(\int_B |S_k f(x)|^p w(x) dx \right)^{1/p} &\lesssim D^{1-\omega} k 2^{\frac{\omega kn}{p u'}} \left(\frac{w(B)}{w(2^k B)} \right)^{\omega/p} \|f\|_{L^p(2^k B, w)} \\ &\lesssim \{w(B)\}^{\lambda/p} D^{1-\omega} k 2^{\frac{kn}{p} \left(\frac{\omega}{u'} - \frac{\omega-\lambda}{r'} \right)} \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)}. \end{aligned}$$

For fixed $\lambda \in (0, 1 - r'/u')$, we choose $\omega \in (\lambda, 1)$ sufficiently close to 1 such that $\omega/u' - (\omega - \lambda)/r' < 0$. Summing over the last inequality yields conclusion (b). \square

Let $p, r \in [1, \infty)$, $\lambda \in (0, 1)$, $q \in [1, \infty]$ and w be a weight. Define the space $L^{p,\lambda}(L^q([1, 2]), l^r; \mathbb{R}^n, w)$ by

$$L^{p,\lambda}(L^q([1, 2]), l^r; \mathbb{R}^n, w) = \{ \vec{f} = \{f_k\}_{k \in \mathbb{Z}} : \|\vec{f}\|_{L^{p,\lambda}(L^q([1, 2]), l^r; \mathbb{R}^n, w)} < \infty \},$$

with

$$\|\vec{f}\|_{L^{p,\lambda}(L^q([1, 2]), l^r; \mathbb{R}^n, w)} = \left\| \left(\int_1^2 \left(\sum_{k \in \mathbb{Z}} |f_k(x, t)|^r \right)^{\frac{q}{r}} dt \right)^{\frac{1}{q}} \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)}.$$

With usual addition and scalar multiplication, $L^{p,\lambda}(L^q([1, 2]), l^r; \mathbb{R}^n, w)$ is a Banach space.

Lemma 4.2. *Let $p \in (1, \infty)$, $\lambda \in (0, 1)$ and $w \in A_p(\mathbb{R}^n)$, \mathcal{G} be a subset in $L^{p,\lambda}(L^2([1, 2]), l^2; \mathbb{R}^n, w)$. Suppose that \mathcal{G} satisfies the following five conditions:*

- (a) \mathcal{G} is a bounded set in $L^{p,\lambda}(L^2([1, 2]), l^2; \mathbb{R}^n, w)$;
- (b) for each fixed $\epsilon > 0$, there exists a constant $A > 0$, such that for all $\{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$\left\| \left(\int_1^2 \sum_{k \in \mathbb{Z}} |f_k(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \chi_{\{|\cdot| > A\}}(\cdot) \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} < \epsilon;$$

- (c) for each fixed $\epsilon > 0$ and $N \in \mathbb{N}$, there exists a constant $\varrho > 0$, such that for all $\vec{f} = \{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$\left\| \sup_{|h| \leq \varrho} \left(\int_1^2 \sum_{|k| \leq N} |f_k(\cdot, t) - f_k(\cdot + h, t)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} < \epsilon;$$

- (d) for each fixed $\epsilon > 0$ and $N \in \mathbb{N}$, there exists a constant $\sigma \in (0, 1/2)$ such that for all $\vec{f} = \{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$\left\| \sup_{|s| \leq \sigma} \left(\int_1^2 \sum_{|k| \leq N} |f_k(\cdot, t+s) - f_k(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} < \epsilon,$$

- (e) for each fixed $D > 0$ and $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for all $\{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$\left\| \left(\int_1^2 \sum_{|k| > N} |f_k(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \chi_{B(0, D)} \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} < \epsilon.$$

Then \mathcal{G} is strongly pre-compact in $L^{p,\lambda}(L^2([1, 2]), l^2; \mathbb{R}^n, w)$.

Proof. The proof is similar to the proof of Lemma 3.3, and so we only give the outline here. It suffices to prove that, for each fixed $\epsilon > 0$, there exists a $\delta = \delta_\epsilon > 0$ and a mapping Φ_ϵ on $L^{p,\lambda}(L^2([1, 2]), l^2; \mathbb{R}^n, w)$, such that $\Phi_\epsilon(\mathcal{G}) = \{\Phi_\epsilon(\vec{f}) : \vec{f} \in \mathcal{G}\}$ is a strongly pre-compact set in $L^{p,\lambda}(L^2([1, 2]), l^2; \mathbb{R}^n, w)$, and for any $\vec{f}, \vec{g} \in \mathcal{G}$,

$$\|\Phi_\epsilon(\vec{f}) - \Phi_\epsilon(\vec{g})\|_{L^{p,\lambda}(L^2([1,2]), l^2; \mathbb{R}^n, w)} < \delta \Rightarrow \|\vec{f} - \vec{g}\|_{L^{p,\lambda}(L^2([1,2]), l^2; \mathbb{R}^n, w)} < 8\epsilon.$$

For fixed $\epsilon > 0$, we choose $A > 1$ large enough as in assumption (b), and $N \in \mathbb{N}$ such that for all $\{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$\left\| \left(\int_1^2 \sum_{|k| > N} |f_k(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \chi_{B(0, 2A)} \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} < \epsilon.$$

Let $Q, Q_1, \dots, Q_J, \mathcal{D}, I_1, \dots, I_L \subset [1, 2]$, and Φ_ϵ be the same as in the proof of Lemma 3.2. For such fixed N , let ϱ and $\sigma \in (0, 1/2)$ small enough such that for all $\vec{f} = \{f_k\}_{k \in \mathbb{Z}} \in \mathcal{G}$,

$$(4.7) \quad \left\| \sup_{|h| \leq \varrho} \left(\int_1^2 \sum_{|k| \leq N} |f_k(\cdot, t) - f_k(\cdot + h, t)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} < \frac{\epsilon}{2J};$$

$$(4.8) \quad \left\| \sup_{|s| \leq \sigma} \left(\int_1^2 \sum_{|k| \leq N} |f_k(\cdot, t + s) - f_k(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} < \frac{\epsilon}{2J},$$

We can verify that Φ_ϵ is bounded on $L^{p,\lambda}(L^2([1, 2]), l^2; \mathbb{R}^n, w)$, and consequently, $\Phi_\epsilon(\mathcal{G}) = \{\Phi_\epsilon(\vec{f}) : \vec{f} \in \mathcal{G}\}$ is a strongly pre-compact set in $L^{p,\lambda}(L^2([1, 2]), l^2; \mathbb{R}^n, w)$. Recall that for $x \in Q_i$ with $1 \leq i \leq J$,

$$\begin{aligned} & \left\{ \int_1^2 \sum_{|k| \leq N} \left| f_k(x, t) - \sum_{v=1}^L m_{Q_i \times I_v}(f_k) \chi_{I_v}(t) \right|^2 dt \right\}^{\frac{1}{2}} \\ & \lesssim \sup_{|h| \leq \varrho} \left(\int_1^2 \sum_{|k| \leq N} |f_k(x, t) - f_k(x + h, t)|^2 dt \right)^{\frac{1}{2}} \\ & \quad + \sup_{|s| \leq \sigma} \left(\int_1^2 \sum_{|k| \leq N} |f_k(x, t + s) - f_k(x, t)|^2 dt \right)^{\frac{1}{2}}. \end{aligned}$$

For a ball $B(y, r)$, a trivial computation involving (4.7) and (4.8), leads to that

$$\begin{aligned} & \int_{B(y, r)} \left(\int_1^2 \sum_{|k| \leq N} \left| f_k(x, t) \chi_{\mathcal{D}} - \sum_{i=1}^J \sum_{j=1}^L m_{Q_i \times I_j}(f_k) \chi_{Q_i \times I_j}(x, t) \right|^2 dt \right)^{\frac{p}{2}} w(x) dx \\ & = \sum_{i=1}^J \int_{B(y, r) \cap Q_i} \left(\int_1^2 \sum_{|k| \leq N} \left| f_k(x, t) - \sum_{i=1}^J m_{Q_i \times I_j}(f_k) \chi_{I_j}(t) \right|^2 dt \right)^{\frac{p}{2}} w(x) dx \\ & \lesssim \epsilon \{w(B(y, r))\}^\lambda. \end{aligned}$$

Therefore,

$$\begin{aligned}
& \int_{B(y,r)} \|\vec{f}\chi_{\mathcal{D}} - \Phi_{\epsilon}(\vec{f})\|_{L^2([1,2],l^2)}^p w(x) dx \\
& \lesssim \int_{B(y,r)} \left(\int_1^2 \sum_{|k|\leq N} |f_k(x,t)\chi_{\mathcal{D}} - \sum_{i=1}^J \sum_{j=1}^L m_{Q_i \times I_j}(f_k)\chi_{Q_i \times I_j}(x,t)|^2 dt \right)^{\frac{p}{2}} w(x) dx \\
& \quad + \int_{B(y,r)} \left(\int_1^2 \sum_{|k|>N} |f_k(x,t)|^2 \right)^{p/2} \chi_{B(0,2A)}(x) w(x) dx \\
& \lesssim 2\epsilon \{w(B(y,r))\}^{\lambda}.
\end{aligned}$$

It then follows from the assumption (b) that for all $\vec{f} \in \mathcal{G}$,

$$\begin{aligned}
\|\vec{f} - \Phi_{\epsilon}(\vec{f})\|_{L^{p,\lambda}(L^2([1,2],l^2;\mathbb{R}^n),w)} & \leq \|\vec{f}\chi_{\mathcal{D}} - \Phi_{\epsilon}(\vec{f})\|_{L^p(L^2([1,2],l^2;\mathbb{R}^n),w)} + \epsilon \\
& < 3\epsilon,
\end{aligned}$$

and

$$\|\vec{f} - \vec{g}\|_{L^{p,\lambda}(\mathbb{R}^n)} < 6\epsilon + \|\Phi_{\epsilon}(f) - \Phi_{\epsilon}(\vec{g})\|_{L^{p,\lambda}(\mathbb{R}^n)}.$$

This completes the proof of Lemma 4.2. \square

Proof of Theorem 1.5. We only consider the case of $p \in (q', \infty)$, $w \in A_{p/q'}(\mathbb{R}^n)$ and $\lambda \in (0, 1)$. Recall that $\mathcal{M}_{\Omega,b}$ is bounded on $L^p(\mathbb{R}^n, w)$. Thus, by Lemma 4.2, we know that $\mathcal{M}_{\Omega,b}$ is bounded on $L^{p,\lambda}(\mathbb{R}^n, w)$. Thus, it suffices to prove that for $b \in C_0^{\infty}(\mathbb{R}^n)$, $\mathcal{M}_{\Omega,b}$ is completely continuous on $L^{p,\lambda}(\mathbb{R}^n, w)$.

Let $j_0 \in \mathbb{Z}_-$, $b \in C_0^{\infty}(\mathbb{R}^n)$ with $\text{supp } b \subset B(0, R)$ and $\|b\|_{L^{\infty}(\mathbb{R}^n)} + \|\nabla b\|_{L^{\infty}(\mathbb{R}^n)} =$

1. Let $\widetilde{K}^j(z) = \frac{|\Omega(z)|}{|z|^n} \chi_{\{2^{j-1} \leq |z| \leq 2^{j+2}\}}(z)$. By the Minkowski inequality,

$$\begin{aligned}
\left(\int_1^2 \sum_{j \in \mathbb{Z}} |F_{j,b}^l f(x,t)|^2 dt \right)^{\frac{1}{2}} & \leq \left(\sum_{j \in \mathbb{Z}} \int_1^2 |F_{j,b}^l f(x,t)|^2 dt \right)^{\frac{1}{2}} \\
& \lesssim \sum_{j \in \mathbb{Z}} \int_{\mathbb{R}^n} \widetilde{K}^j * \phi_{j-l}(x-y) |f(y)| dy.
\end{aligned}$$

It is obvious that $\text{supp } \widetilde{K}^j * \phi_{j-l} \subset \{x : 2^{j-3} \leq |x| \leq 2^{j+3}\}$, and for any $R > 0$,

$$\int_{R \leq |x| \leq 2R} \left| \sum_{j \in \mathbb{Z}} \widetilde{K}^j * \phi_{j-l}(x) \right|^q dx \leq \sum_{j: 2^j \approx R} \|\widetilde{K}^j * \phi_{j-l}\|_{L^q(\mathbb{R}^n)}^q \lesssim R^{-nq+n}.$$

Let $\epsilon > 0$. We deduce from Lemma 4.1 and the inequality (3.7) that, there exists a constant $A > 0$, such that for all $f \in L^p(\mathbb{R}^n)$,

$$(4.9) \quad \left\| \left(\int_1^2 \sum_{j \in \mathbb{Z}} |F_{j,b}^l f(x,t)|^2 dt \right)^{\frac{1}{2}} \chi_{\{|\cdot| > A\}}(\cdot) \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} < \epsilon \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)}.$$

Recall that $\widetilde{\mathcal{M}}_{\Omega}^{l,j_0}$ is bounded on $L^{p,\lambda}(\mathbb{R}^n, w)$. For $r > 1$ small enough, M_r is also bounded on $L^{p,\lambda}(\mathbb{R}^n, w)$ (see [28]). Thus by (3.4), we know that there exists a constant $\rho > 0$, such that

$$\begin{aligned}
(4.10) \quad & \left\| \sup_{|h| \leq \rho} \left(\int_1^2 \sum_{j > j_0} |F_{j,b}^l f(\cdot, t) - F_{j,b}^l f(\cdot + h, t)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} \\
& \lesssim \epsilon \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)}.
\end{aligned}$$

It follows from Lemma 4.1, estimate (3.5) that for each $N \in \mathbb{N}$, there exists a constant $\sigma \in (0, 1/2)$ such that

$$(4.11) \quad \left\| \sup_{|s| \leq \sigma} \left(\int_1^2 \sum_{|j| \leq N} |F_{j,b}^l f(\cdot, s+t) - F_{j,b}^l f(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} < \epsilon \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)}.$$

We also obtain by Lemma 4.1 and (3.6) that for each fixed $D > 0$, there exists $N \in \mathbb{N}$ such that

$$(4.12) \quad \left\| \left(\int_1^2 \sum_{j > N} |F_{j,b}^l f(\cdot, t)|^2 dt \right)^{\frac{1}{2}} \chi_{B(0,D)} \right\|_{L^{p,\lambda}(\mathbb{R}^n, w)} < \epsilon \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)}.$$

The inequalities (4.9)-(4.12), via Lemma 4.2, tell us for any $j_0 \in \mathbb{Z}_-$, the operator $\mathcal{F}_{j_0}^l$ defined by (3.18) is compact from $L^{p,\lambda}(\mathbb{R}^n, w)$ to $L^{p,\lambda}(L^2([1, 2]), l^2; \mathbb{R}^n, w)$. On the other hand, by Lemma 4.1, Theorem 2.1 and Lemma 3.2, we know that

$$\|\widetilde{\mathcal{M}}_{\Omega} f - \widetilde{\mathcal{M}}_{\Omega}^l f\|_{L^{p,\lambda}(\mathbb{R}^n, w)} \lesssim 2^{-\varepsilon \varrho_p l} \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)},$$

and

$$\|\widetilde{\mathcal{M}}_{\Omega, b}^{l, j_0} f - \widetilde{\mathcal{M}}_{\Omega, b}^l f\|_{L^{p,\lambda}(\mathbb{R}^n, w)} \lesssim 2^{\varepsilon j_0} \|f\|_{L^{p,\lambda}(\mathbb{R}^n, w)}.$$

As it was shown in the proof of Theorem 1.3, we can deduce from the last facts that $\mathcal{M}_{\Omega, b}$ is completely continuous on $L^{p,\lambda}(\mathbb{R}^n, w)$ when $b \in C_0^\infty(\mathbb{R}^n)$. This completes the proof of Theorem 1.5. \square

REFERENCES

- [1] D. R. Aamas and J. Xiao, Morrey potential and harmonic analysis, *Ark. Mat.* **50** (2012), 201-230.
- [2] A. Al-Salman, On the L^2 boundedness of parametric Marcinkiewicz integral operator, *J. Math. Anal. Appl.* **375** (2011) 745-752.
- [3] A. Al-Salman, H. Al-Qassem, L. Cheng and Y. Pan, L^p bounds for the function of Marcinkiewicz, *Math. Research Letter* **9** (2002), 697-700.
- [4] K. F. Anderson and R. T. John, Weighted inequality for vector-valued maximal functions and singular integrals, *Studia Math.* **69** (1980), 19-31.
- [5] J. Alvarez, R. Bagby, D. Kurtz and C. Pérez, Weighted estimates for commutators of linear operators, *Studia Math.* **104** (1993), 195-209.
- [6] G. Bourdaud, M. Lanze de Cristoforis and W. Sickel, Functional calculus on BMO and related spaces, *J. Funct. Anal.* **189** (2002), 515-538.
- [7] A. P. Calderón and A. Zygmund, On singular integrals, *Amer. J. Math.* **78** (1956), 289-309.
- [8] D. Chen and S. Lu, L^p boundedness for commutators of parabolic Littlewood-Paley operators with rough kernels, *Math. Nachr.* **284** (2011), 973-986.
- [9] J. Chen, D. Fan and Y. Pan, A note on a Marcinkiewicz integral operator, *Math. Nachr.* **227** (2001), 33-42.
- [10] J. Chen and G. Hu, Compact commutators of rough singular integral operators, *Canad. Math. Bull.*, **64** (2012), 257-281.
- [11] Completely continuous commutator of Marcinkiewicz integral, *Publ. Math. Debrecen*, **90**(2017), 11-32.
- [12] Y. Chen and Y. Ding, Compactness characterization of commutators for Littlewood-Paley operators, *Kodai Math. J.* **32** (2009), 256-323.
- [13] Y. Chen, Y. Ding and X. Wang, Compactness for commutators of Marcinkiewicz integral on Morrey spaces, *Taiwanese. J. Math.* **15** (2011), 633-658.
- [14] A. Clop and V. Cruz, Weighted estimates for Beltrami equations, *Ann. Acad. Sci. Fenn. Math.* **38** (2013), 91-113.
- [15] R. Coifman, R. Rochberg and G. Weiss, Factorization theorems for Hardy spaces in several variables. *Ann. Math.* **103** (1976), 611-635.

- [16] R. Coifman and G. Weiss, Extension of Hardy spaces and their use in analysis, *Bull. Amer. Math. Soc.* **83** (1977), 569-645.
- [17] Y. Ding, D. Fan and Y. Pan, L^p -boundedness of Marcinkiewicz integrals with Hardy space function kernel, *Acta Math. Sinica (English Series)* **16** (2000), 593-600.
- [18] Y. Ding, D. Fan and Y. Pan, Weighted boundedness for a class of rough Marcinkiewicz integral, *Indiana Univ. Math. J.* **48** (1999), 1037-1055.
- [19] Y. Ding, S. Lu and K. Yabuta, A problem on rough Marcinkiewicz functions, *J. Austral. Math. Soc.* **71** (2001), 1-9.
- [20] Y. Ding, S. Lu and K. Yabuta, On commutators of Marcinkiewicz integrals with rough kernel, *J. Math. Anal. Appl.* **275** (2002), 60-68.
- [21] Y. Ding, Q. Xue and K. Yabuta, A remark to the L^2 boundedness of parametric Marcinkiewicz integral, *J. Math. Anal. Appl.* **387** (2012), 691-697.
- [22] J. Duoandikoetxea, Weighted norm inequalities for homogeneous singular integrals, *Trans. Amer. Math. Soc.* **336** (1993), 869-880.
- [23] J. Duoandikoetxea and J. L. Rubio de Francia, Maximal and singular integrals via Fourier transform estimates. *Invent. Math.* **84** (1986), 541-561.
- [24] L. Grafakos, *Modern Fourier analysis*, GTM250, 2nd Edition, Springer, New York, 2008.
- [25] X. Guo and G. Hu, Compactness of the commutators of homogeneous singular integral operators *Sci. China Math.* **58** (2015), 2347-2362.
- [26] G. Hu, $L^p(\mathbb{R}^n)$ boundedness for a class of g -functions and its applications, *J. Hokkaido Math.* **32** (2003), 497-521.
- [27] G. Hu and D. Yan, On commutator of the Marcinkiewicz integral, *J. Math. Anal. Appl.* **283** (2003), 351-361.
- [28] Y. Komori and S. Shirai, Weighted Morrey spaces and a singular integral operator, *Math. Nachr.* **282** (2009), 219-232.
- [29] D. S. Kurtz and R. Wheeden, Results on weighted norm inequalities for multipliers. *Trans. Amer. Math. Soc.* **255**(1979), 343-362.
- [30] S. Mao, Y. Sawano and H. Wu, On the compactness of commutators for rough Marcinkiewicz integral operators, *Taiwanese J. Math.* **19** (2015), 1777-1793.
- [31] J. Marcinkiewicz, Sur quelques intégrales du type de Dini, *Ann. Soc. Polon. Math.* **17** (1938), 42-50.
- [32] A. Ruiz and L. Vega, Unique continuation for Schrödinger operators with potential in Morrey space, *Publ. Mat.* **35** (1991), 291-298.
- [33] Z. Shen, Boundary value problems in Morrey spaces for elliptic systems on Lipschitz domains, *Amer. J. Math.* **125** (2003), 1079-1115.
- [34] E. M. Stein, On the function of Littlewood-Paley, Lusin and Marcinkiewicz, *Trans. Amer. Math. Soc.* **88** (1958), 430-466.
- [35] E. M. Stein and G. Weiss, Interpolation of operators with changes of measures. *Trans. Amer. Math. Soc.* **87** (1958), 159-172.
- [36] A. Torchinsky and S. Wang, A note on the Marcinkiewicz ntegral, *Colloq. Math.* **47** (1990), 235-243.
- [37] A. Uchiyama, On the compactness of operators of Hankel type. *Tohoku Math. J.* **30** (1978), 163-171.
- [38] T. Walsh, On the function of Marcinkiewicz, *Studia Math.* **44** (1972), 203-217.
- [39] D. K. Watson, Weighted estimates for singular integrals via Fourier transform estimates. *Duke Math. J.* **60** (1990), 389-399.
- [40] A. Zygmund, *Trigonometric series*, 3rd Edition, Cambridge Univ. Press, Cambridge, 2002.

GUOEN HU, DEPARTMENT OF APPLIED MATHEMATICS, ZHENGZHOU INFORMATION SCIENCE AND TECHNOLOGY INSTITUTE, ZHENGZHOU 450001, P. R. CHINA

E-mail address: guoenxx@163.com