

# Eligibility Traces and Plasticity on Behavioral Time Scales: Experimental Support of neoHebbian Three-Factor Learning Rules

Wulfram Gerstner\*, Marco Lehmann, Vasiliki Liakoni, Dane Corneil, and Johanni Brea

School of Computer Science & School of Life Sciences

École Polytechnique Fédérale de Lausanne,

1015 Lausanne EPFL, Switzerland

(\*) corresponding author

January 17, 2018

## Abstract

Most elementary behaviors such as moving the arm to grasp an object or walking into the next room to explore a museum evolve on the time scale of seconds; in contrast, neuronal action potentials occur on the time scale of a few milliseconds. Learning rules of the brain must therefore bridge the gap between these two different time scales. Modern theories of synaptic plasticity have postulated that the co-activation of pre- and postsynaptic neurons sets a flag at the synapse, called an eligibility trace, that leads to a weight change only if an additional factor is present while the flag is set. This third factor, signaling reward, punishment, surprise, or novelty, could be implemented by the phasic activity of neuromodulators or specific neuronal inputs signaling special events. While the theoretical framework has been developed over the last decades, experimental evidence in support of eligibility traces on the time scale of seconds has been collected only during the last few years. Here we review, in the context of three-factor rules of synaptic plasticity, four key experiments that support the role of synaptic eligibility traces in combination with a third factor as a biological implementation of neoHebbian three-factor learning rules.

**Keywords** Eligibility trace, Hebb Rule, Reinforcement Learning, Neuromodulators, Surprise, synaptic tagging, Synaptic Plasticity, Behavioral Learning

## 1 Introduction

Humans are able to learn novel behaviors such as pressing a button, swinging a tennis racket, or breaking at a red traffic light; they are also able to form memories of salient events, learn to distinguish flowers, and to establish a mental map when exploring a novel environment. Memory formation and behavioral learning is linked to changes of synaptic connections (Martin et al.,

2000). Long-lasting synaptic changes, necessary for memory, can be induced by Hebbian protocols that combine the activation of presynaptic terminals with a manipulation of the voltage or the firing state of the postsynaptic neuron (Lisman, 2003). Traditional experimental protocols of long-term potentiation (LTP) (Bliss and Lømo, 1973; Bliss and Collingridge, 1993), long-term depression (LTD) (Levy and Stewart, 1983; Artola and Singer, 1993) and spike-timing dependent plasticity (STDP) (Markram et al., 1997; Zhang et al., 1998; Sjöström et al., 2001) neglect that additional factors such as neuromodulators or other gating signals might be necessary to permit synaptic changes (Gu, 2002; Hasselmo, 2006; Reynolds and Wickens, 2002). Early STDP experiments that involved neuromodulators mainly focused on tonic bath application of modulatory factors (Pawlak et al., 2010). However, from the perspective of formal learning theories, to be reviewed below, the timing of modulatory factors is just as crucial (Schultz and Dickinson, 2000; Schultz, 2002). From the theoretical perspective, STDP under the control of neuromodulators leads to the framework of three-factor learning rules (Xie and Seung, 2004; Legenstein et al., 2008; Vasilaki et al., 2009) where an eligibility trace represents the Hebbian idea of co-activation of pre- and postsynaptic neurons (Hebb, 1949) while modulation of plasticity by additional gating signals is represented generically by a 'third factor' (Crow, 1968; Barto, 1985; Legenstein et al., 2008). Such a third factor could represent variables such as 'reward minus expected reward' (Williams, 1992; Schultz, 1998; Sutton and Barto, 1998) or the saliency of an unexpected event (Ljunberg and W. Schultz, 1992; Redgrave and Gurney, 2006).

In an earlier paper (Frémaux and Gerstner, 2016) we reviewed the theoretical literature of, and experimental support for, three-factor rules available by the end of 2013. During recent years, however, the experimental procedures advanced significantly and provided direct physiological evidence of eligibility traces and three-factor learning rules for the first time, making an updated review of three-factor rules necessary. In the following, we – a group of theoreticians – review five experimental papers indicating support of eligibility traces in striatum (Yagishita et al., 2014), cortex (He et al., 2015), and hippocampus (Brzosko et al., 2015, 2017; Bittner et al., 2017). We will close with a few remarks on the paradoxical nature of theoretical predictions in the field of computational neuroscience.

## 2 Hebbian rules versus three-factor rules

Learning rules describe the change of the strength of a synaptic contact between a presynaptic neuron  $j$  and a postsynaptic neuron  $i$ . The strength of an excitatory synaptic contact can be defined by the amplitude of the postsynaptic potential which is closely related to the spine volume and the number of AMPA receptors (Matsuzaki et al., 2001). Synapses contain complex molecular machineries (Lisman, 2003; Redondo and Morris, 2011; Hugarir and Nicoll, 2013; Lisman, 2017), but for the sake of transparency of the arguments, we will keep the mathematical notation as simple as possible and characterize the synapse by two variables only: the first one is the synaptic strength  $w_{ij}$ , measured as spine volume or amplitude of postsynaptic potential, and the second one is a synapse-internal variable  $e_{ij}$  which is not directly visible in standard electrophysiological experiments. In our view, the internal variable  $e_{ij}$  represents a metastable

transient state of interacting molecules in the spine head or a multi-molecular substructure in the postsynaptic density which serves as a synaptic flag indicating that the synapse is ready for an increase or decrease of its spine volume (Bosch et al., 2014). The precise biological nature of  $e_{ij}$  is not important to understand the theories and experiments that are reviewed below. We refer to  $e_{ij}$  as the 'synaptic flag' or the 'eligibility trace' and to  $w_{ij}$  as the 'synaptic weight' or 'strength' of the synaptic contact. A change of the synaptic flag indicates a 'candidate weight change' (Frémaux et al., 2010) whereas a change of  $w_{ij}$  indicates an actual, measurable, change of the synaptic weight. Before we turn to three-factor rules, let us discuss conventional models of Hebbian learning.

## 2.1 Hebbian learning rules

Hebbian learning rules are the mathematical summary of the outcome of experimental protocols inducing long-term potentiation (LTP) or long-term depression (LTD) of synapses. Suitable experimental protocols include strong extracellular stimulation of presynaptic fibers (Bliss and Lømo, 1973; Levy and Stewart, 1983), manipulation of postsynaptic voltage in the presence of presynaptic spike arrivals (Artola and Singer, 1993), or spike-timing dependent plasticity (STDP) (Markram et al., 1997; Sjöström et al., 2001). In all mathematical formulations of Hebbian learning, the synaptic flag variable  $e_{ij}$  is sensitive to the *combination* of presynaptic spike arrival and a postsynaptic variable, such as the voltage at the location of the synapse. Under a Hebbian learning rule, repeated presynaptic spike arrivals at a synapse of a neuron at rest do not cause a change of the synaptic variable. Similarly, an elevated postsynaptic potential in the absence of a presynaptic spike does not cause a change of the synaptic variable. Thus Hebbian learning always needs two factors for a synaptic change: a factor caused by a presynaptic signal such as glutamate; and a factor that depends on the state of the postsynaptic neuron.

What are these factors? We can think of the presynaptic factor as the time course of glutamate available in the synaptic cleft or bound to the postsynaptic membrane. Note that the term 'presynaptic factor' that we will use in the following *does not imply* that the physical location of the presynaptic factor is inside the presynaptic terminal - the factor could very well be located in the postsynaptic membrane as long as it *only* depends on the amount of available neurotransmitters. The postsynaptic factor might be the calcium in the synaptic spine (Shouval et al., 2002; Rubin et al., 2005), a calcium-related second messenger molecule (Graupner and Brunel, 2007), or simply the voltage at the site of the synapse (Brader et al., 2007; Clopath et al., 2010).

We remind the reader that we always use the index  $j$  to refer to the presynaptic neuron and the index  $i$  to refer to the postsynaptic one. For the sake of simplicity, let us call the presynaptic factor  $x_j$  (representing the activity of the presynaptic neuron or the amount of glutamate in the synaptic cleft) and the postsynaptic factor  $y_i$  (representing the state of the postsynaptic neuron). In a Hebbian learning rule, changes of the synaptic flag  $e_{ij}$  need both  $x_j$  and  $y_i$

$$\frac{d}{dt}e_{ij} = \eta x_j g(y_i) - e_{ij}/\tau_e \quad (1)$$

where  $\eta$  is the constant learning rate,  $\tau_e$  is a decay time constant and  $g(y_j)$  is some arbitrary, potentially nonlinear, function of the postsynaptic variable  $y_i$ . Thus, the synaptic flag  $e_{ij}$  acts as a *correlation detector* between presynaptic activity  $x_j$  and the state of the postsynaptic neuron  $y_i$ . In some models, there is no decay or the decay is considered negligible on the time scale of one experiment ( $\tau_e \rightarrow \infty$ ).

Let us discuss two examples. In the Bienenstock-Cooper Munro (BCM) model of developmental cortical plasticity (Bienenstock et al., 1982) the presynaptic factor  $x_j$  is the firing rate of the presynaptic neuron and  $g(y_i) = (y_i - \theta) y_i$  is a quadratic function with  $y_i$  the postsynaptic firing rate and  $\theta$  a threshold rate. Thus, if both pre- and postsynaptic neurons fire together at a high rate  $x_j = y_i > \theta$  then the synaptic flag  $e_{ij}$  increases. In the BCM model, just like in most other conventional models, a change of the synaptic flag (i.e., an internal state of the synapse) leads instantaneously to a change of the weight  $e_{ij} \rightarrow w_{ij}$  so that an experimental protocol results immediately in a measurable weight change. With the BCM rule and other similar rules (Oja, 1982; Miller and MacKay, 1994), the synaptic weight increases if both presynaptic and postsynaptic neuron are highly active, implementing the slogan 'fire together, wire together' (Lowel and Singer, 1992; Shatz, 1992); cf. Fig. 1A(i).

As a second example, we consider the Clopath model (Clopath et al., 2010). In this model, there are two correlation detectors implemented as synaptic flags  $e_{ij}^+$  and  $e_{ij}^-$  for LTP and LTD, respectively. The synaptic flag  $e_{ij}^+$  for LTP uses a presynaptic factor  $x_j^+$  (related to the amount of glutamate available in the synaptic cleft) which increases with each presynaptic spike and decays back to zero over the time of a few milliseconds (Clopath et al., 2010). The postsynaptic factor for LTP depends on the postsynaptic voltage  $y_i$  via a function  $g(y_i) = a_+[y_i - \theta_+] \bar{y}_i$  where  $a_+$  is a positive constant,  $\theta_+$  a voltage threshold, square brackets denote the rectifying piecewise linear function, and  $\bar{y}_i$  a running average of the voltage with a time constant of tens of milliseconds. An analogous, but simpler, combination of presynaptic spikes and postsynaptic voltage defines the second synaptic flag  $e_{ij}^-$  for LTD (Clopath et al., 2010). The total change of the synaptic weight is the combination of the two synaptic flags for LTP and LTD:  $dw_{ij}/dt = de_{ij}^+/dt - de_{ij}^-/dt$ . Note that, since both synaptic flags  $e_{ij}^+$  and  $e_{ij}^-$  depend on the postsynaptic voltage, postsynaptic spikes are not a necessary condition for changes, in agreement with voltage-dependent protocols (Artola and Singer, 1993; Ngezahayo et al., 2000). Thus, in voltage-dependent protocols, and similarly in voltage-dependent models, 'wiring together' is possible without 'firing together' - indicating that the theoretical framework sketched above goes beyond a narrow view of Hebbian learning; cf. Fig. 1A(ii).

If we restrict the discussion of the postsynaptic variable to super-threshold spikes, then the Clopath model becomes identical to the triplet STDP model (Pfister et al., 2006) which is in turn closely related to other nonlinear STDP models (Senn et al., 2001; Froemke and Dan, 2002; Izhikevich and Desai, 2003) as well as to the BCM model discussed above (Pfister et al., 2006; Gjorjjeva et al., 2011). Classic pair-based STDP models (Gerstner et al., 1996; Kempster et al., 1999; Song et al., 2000; van Rossum et al., 2000; Rubin et al., 2001) are further examples of the general theoretical framework of Eq. (1) and so are some models of structural plasticity (Helias et al., 2008; Deger et al., 2012; Fauth et al., 2015). Hebbian models of synaptic consolidation




Figure 1: **A.** Two Hebbian protocols and one three-factor learning protocol. (i) Hebbian STDP protocol with presynaptic spikes (presynaptic factor) followed by a burst of postsynaptic spikes (postsynaptic factor). Synapses in the stimulated pathway (green) will typically show LTP while an unstimulated synapse (red) will not change its weight (Markram et al., 1997). (ii) Hebbian voltage pairing protocol of presynaptic spikes (presynaptic factor) with a depolarization of the postsynaptic neuron (postsynaptic factor). Depending on the amount of depolarization the stimulated pathway (green) will show LTP or LTD while an unstimulated synapse (red) does not change its weight (Artola and Singer, 1993; Ngezahayo et al., 2000). (iii) Results of a Hebbian induction protocol are influenced by a third factor (blue) even if it is given after a delay  $d$ . The third factor could be a neuromodulator such as dopamine, acetylcholine, norepinephrine, or serotonin (Pawlak et al., 2010; Yagishita et al., 2014; He et al., 2015; Brzosko et al., 2015, 2017; Bittner et al., 2017). **B.** Specificity of three-factor learning rules. (i) Presynaptic input spikes (green) arrive at two different neurons, but only one of these also shows postsynaptic activity (orange spikes). (ii) A synaptic flag is set only at the synapse with a Hebbian co-activation of pre- and postsynaptic factors; the synapse becomes then eligible to interact with the third factor (blue). Spontaneous spikes of other neurons do not interfere. (iii) The interaction of the synaptic flag with the third factor leads to a strengthening of the synapse (green).

have several hidden flag variables (Fusi et al., 2005; Barrett et al., 2009; Benna and Fusi, 2016) but can also be situated as examples within the general framework of Hebbian rules. Note that in most of the examples so far the measured synaptic weight is a linear function of the synaptic flag variable(s). However, this does not need to be the case. For example, in some voltage-based (Brader et al., 2007) or calcium-based models (Shouval et al., 2002; Rubin et al., 2005), the synaptic flag is transformed into a weight change only if  $e_{ij}$  is above or below some threshold, or only after some further filtering.

To summarize, in the theoretical literature the class of Hebbian models is a rather general framework encompassing all those models that are driven by a combination of presynaptic activity and the state of the postsynaptic neuron. In this view, Hebbian models depend on two factors related to the activity of the presynaptic and the state of the postsynaptic neuron. The correlations between the two factors can be extracted on different time scales using one or, if necessary, several flag variables. The flag variables trigger a change of the measured synaptic weight. In the following we build on Hebbian learning, but extend the theoretical framework to include a third factor.

## 2.2 Three-factor learning rules

We are interested in a framework where a Hebbian co-activation of two neurons leaves one or several flags (eligibility trace) at the synapse connecting these neurons. The flag is not directly visible and does not automatically trigger a change of the synaptic weight. An actual weight change is implemented only if a third signal, e.g., a phasic increase of neuromodulator activity or an additional input (signaling the occurrence of a special event) is present at the same time or in the near future. Theoreticians refer to such a plasticity model as a three-factor learning rule (Xie and Seung, 2004; Legenstein et al., 2008; Vasilaki et al., 2009; Frémaux et al., 2013; Frémaux and Gerstner, 2016). Three-factor rules have also been called 'neoHebbian' (Lisman et al., 2011; Lisman, 2017) or 'heterosynaptic (modulatory-input dependent)' (Bailey et al., 2000) and can be traced back to the 1960s (Crow, 1968), if not earlier. To our knowledge the wording 'three factors' was first used by (Barto, 1985). The terms eligibility and eligibility traces have been used in (Klopf, 1972; Sutton and Barto, 1981; Barto et al., 1983; Barto, 1985; Williams, 1992; Schultz, 1998; Sutton and Barto, 1998) but in some of the early studies it remained unclear whether eligibility traces can be set by presynaptic activity alone (Klopf, 1972; Sutton and Barto, 1981) or only by Hebbian co-activation of pre- and postsynaptic neurons (Barto et al., 1983; Barto, 1985; Williams, 1992; Schultz, 1998; Sutton and Barto, 1998).

The basic idea of a modern eligibility trace is that a synaptic flag variable  $e_{ij}$  is set according to Eq. (1) by coincidences between presynaptic activity  $x_j$  and a postsynaptic factor  $y_i$ . The update of the synaptic weight  $w_{ij}$ , as measured via the spine volume or the amplitude of the excitatory postsynaptic potential (EPSP), is given by

$$\frac{d}{dt}w_{ij} = e_{ij} M_{3rd}(t) \quad (2)$$

where  $M_{3rd}(t)$  refers to the third factor (Izhikevich, 2007; Legenstein et al., 2008; Frémaux et al., 2013). Thus, a third factor is needed to transform the eligibility trace into a weight change; cf. Fig. 1A(iii). Note that the weight change is proportional to  $M_{3rd}(t)$ . Thus, the third factor influences the speed of learning. In the absence of the third factor ( $M_{3rd}(t) = 0$ ), the synaptic weight is not changed. We emphasize that a positive value of the synaptic flag in combination with a negative value of  $M_{3rd}(t)$  leads to a decrease of the weight. Therefore, the third factor also influences the *direction* of change.

What could be the source of such a third factor? The third factor could be triggered by attentional processes, surprising events, or reward. Phasic signals of neuromodulators such as dopamine, serotonin, acetylcholine, or noradrenaline are obvious candidates of a third factor, but potentially not the only ones. Note that axonal branches of most dopaminergic, serotonergic, cholinergic, or adrenergic neurons project broadly onto large regions of cortex so that a phasic neuromodulator signal arrives at many neurons and synapses in parallel (Schultz, 1998). Since neuromodulatory information is shared by many neurons, the variable  $M_{3rd}(t)$  of the third factor has no neuron-specific index (neither  $i$  nor  $j$ ) in our mathematical formulation. Because of its unspecific nature, the theory literature sometimes refers to the third factor as a 'global'

broadcasting signal, even though in practice not every brain region and every synapse is reached by each neuromodulator.

Note that we mathematically define a phasic signal as the deviation from the running average so that  $M_{3rd}(t)$  in Eq. (2) can take positive and negative values. However, the third factor could also be biologically implemented by *positive* excursions of the activity using two different neuromodulators with very low baseline activity. The activity of the first modulator could indicate positive values of the third factor and that of the second modulator negative ones - similar to ON and OFF cells in the retina. Similarly, the framework of neoHebbian three-factor rules is general enough to enable biological implementations with separate eligibility traces for LTP and LTD as discussed above in the context of the Clopath model (Clopath et al., 2010).

### 2.3 Examples and theoretical predictions

There are several known examples in the theoretical literature of neoHebbian three-factor rules. We briefly present three of these and formulate expectations derived from the theoretical framework which we would like to compare to experimental results in the next section.

As a first example, we consider the relation of neoHebbian three-factor rules to reward-based learning. Temporal Difference (TD) algorithms such as SARSA( $\lambda$ ) or TD( $\lambda$ ) from the theory of reinforcement learning (Sutton and Barto, 1998) as well as learning rules derived from policy gradient theories (Williams, 1992) can be interpreted in neuronal networks in terms of neoHebbian three-factor learning rules. The resulting plasticity rules are applied to synapses connecting 'state-neurons' (e.g., place cells coding for the current location of an animal) to 'action neurons' (e.g., cells initiating an action program such as 'turn left') (Brown and Sharp, 1995; Suri and Schultz, 1999; Arleo and Gerstner, 2000; Foster et al., 2000; Xie and Seung, 2004; Loewenstein and Seung, 2006; Florian, 2007; Izhikevich, 2007; Legenstein et al., 2008; Vasilaki et al., 2009; Frémaux et al., 2013); for a review, see (Frémaux and Gerstner, 2016). The eligibility trace is increased during the joint activation of 'state-neurons' and 'action-neurons' and decays exponentially thereafter consistent with the framework of Eq. (1). The third factor is defined as reward minus expected reward where the exact definition of expected reward depends on the implementation details. A long line of research by Wolfram Schultz and colleagues (Schultz et al., 1997; Schultz, 1998; Schultz and Dickinson, 2000; Schultz, 2002) indicates that phasic increases of the neuromodulator dopamine have the necessary properties required for a third factor in the theoretical framework of reinforcement learning.

However, despite the rich literature on dopamine and reward-based learning accumulated during the last 25 years, measurements of the decay time constant  $\tau_e$  of the eligibility trace  $e_{ij}$  in Eq. (1) have, to our knowledge, not become available before 2015. From the mathematical framework of neoHebbian three-factor rules it is clear that, in the context of action learning, the time constant of the eligibility trace (i.e., the duration of the synaptic flag) should roughly match the time span from the initiation of an action to the delivery of reward. As an illustration, let us imagine a baby that attempts to grasp her bottle of milk. The typical duration of one grasping movement is in the range of a second, but potentially only the third grasping attempt might be

successful. Let us suppose that each grasping movement corresponds to the co-activation of some neurons in the brain. If the duration of the synaptic flag is much less than a second, the co-activation of pre- and postsynaptic neurons that sets the synaptic flag (eligibility trace) cannot be linked to the reward one second later and synapses do not change. If the duration of the synaptic flag is much longer than a second, then the two 'wrong' grasping attempts are reinforced nearly as strongly as the third, successful one which mixes learning of 'wrong' co-activations with the correct ones. Hence, *the existing theory of three-factor learning rules predicts that the synaptic flag (eligibility trace for action learning) should be in the range of a typical elementary action, about 200ms to 2s*; see, for example, p. 15 of (Schultz, 1998)<sup>1</sup>, p.3 of (Izhikevich, 2007), <sup>2</sup>, p.3 of (Legenstein et al., 2008)<sup>3</sup>, p. 13327 of (Frémaux et al., 2010)<sup>4</sup>, or p. 13 of (Frémaux et al., 2013)<sup>5</sup>. An eligibility trace of 100ms or 20 seconds would be less useful for learning a typical elementary action or delayed reward task than an eligibility trace in the range of 200ms to 2s. The expected time scale of the synaptic eligibility trace should roughly match the maximal delay of reinforcers in conditioning experiments (Thorndike, 1911; Pavlov, 1927; J.Blac et al., 1985), linking synaptic processes to behavior. For human behavior, delaying a reinforcer by 10 seconds during ongoing actions decreases learning compared to immediate reinforcement (Okouchi, 2009).

As a second example, we consider situations that go beyond standard reward-based learning. Even in the absence of reward, a surprising event might trigger a combination of neuromodulators such as noradrenaline, acetylcholine and dopamine that may act as third factor for synaptic plasticity. Imagine a small baby lying in the cradle with an attractive colorful object swinging above him. He spontaneously makes several arm movements until finally he succeeds, by chance, to grasp the object. There is no food reward for this action. However, the fact that he can now turn the object, look at it from different sides, or put it in his mouth is satisfying because it leads to many novel (and exciting!) stimuli. The basic idea is that, in such situations, novelty or surprise acts as a reinforcer even in complete absence of food rewards (Schmidhuber, 1991; Singh et al., 2004; Oudeyer et al., 2007). Theoreticians have studied these ideas in the context of curiosity (Schmidhuber, 2010), information gain during active exploration (Storck et al., 1995; Sun et al., 2011; Schmidhuber, 2006; Sun et al., 2011; Little and Sommer, 2013; Friston et al., 2016), and via formal definitions of surprise (Storck et al., 1995; Itti and Baldi, 2009; Schmidhuber, 2010; Shannon, 1948; Friston, 2010; Faraji et al., 2018). Note that surprise is not always linked to active exploration but can also occur in a passive situation, e.g. listening to tone beeps or viewing simple stimuli (Squires et al., 1976; Kolossa et al., 2013, 2015; Meyniel et al., 2016). Measurable physiological responses to surprise include pupil dilation (Hess and Polt, 1960) and the P300 component of the electroencephalogram (Squires et al., 1976).

---

<sup>1</sup>'Learning ... (with dopamine) on striatal synapses ... requires hypothetical traces of synaptic activity that last until reinforcement occurs and makes those synapses eligible for modification ...'

<sup>2</sup>'(the eligibility trace  $c$ ) decays to  $c = 0$  exponentially with the time constant  $\tau_c = 1s$ '

<sup>3</sup>'the time scale of the eligibility trace is assumed in this article to be on the order of seconds'

<sup>4</sup>'candidate weight changes  $e_{ij}$  decay to zero with a time constant  $\tau_e = 500ms$ . The candidate weight changes  $e_{ij}$  are known as the eligibility trace in reinforcement learning'

<sup>5</sup>'the time scales of the eligibility traces we propose, (are) on the order of hundreds of milliseconds, .. Direct experimental evidence of eligibility traces still lacks, ...'



If surprise can play a role similar to reward, then surprise-transmitting broadcast signals should speed-up plasticity. Indeed, theories of surprise as well as hierarchical Bayesian models predict a faster change of model parameters for surprising stimuli than for known ones (Yu and Dayan, 2005; Nassar et al., 2010; Mathys et al., 2011, 2014; Faraji et al., 2018) similar to, but more general than, the well-known Kalman filters (Kalman, 1960). Since the translation of these abstract models into networks of spiking neurons is still missing, precise predictions for surprise modulation of plasticity in the form of three-factor rules are not yet available. However, if we consider noradrenaline, acetylcholine, and/or dopamine as candidate neuromodulators signaling novelty and surprise, we expect that these neuromodulators should have a strong effect on plasticity so as to boost learning of surprising stimuli. The influence of tonic applications of various neuromodulators on synaptic plasticity has been shown in numerous studies (Gu, 2002; Hasselmo, 2006; Reynolds and Wickens, 2002; Pawlak et al., 2010). However, in the context of the above examples, we are interested in phasic neuromodulatory signals. Phasic signals conveying moments of surprise are most useful for learning if they are either synchronous with the stimulus to be learned (e.g., passive listening or viewing) or arise with a delay corresponding to one exploratory movement (e.g. grasping). Hence, we predict from these considerations a decay constant  $\tau_e$  of the synaptic flag in the range of 1 second, but with a pronounced effect for synchronous or near-synchronous events.

As our final example, we would like to comment on synaptic consolidation. The synaptic tagging-and-capture hypothesis (Frey and Morris, 1997; Reymann and Frey, 2007; Redondo and Morris, 2011) perfectly fits in the framework of three-factor learning rules: The joint pre- and postsynaptic activity sets the synaptic flag (called 'tag' in the context of consolidation) which decays back to zero over the time of one hour. To stabilize synaptic weights beyond one hour an additional factor is needed to trigger protein synthesis required for long-term maintenance of synaptic weights (Redondo and Morris, 2011; Reymann and Frey, 2007). Neuromodulators such as dopamine have been identified as the necessary third factor for consolidation (Bailey et al., 2000; Reymann and Frey, 2007; Redondo and Morris, 2011; Lisman, 2017). Indeed, modern computational models of synaptic consolidation take into account the effect of neuromodulators (Clopath et al., 2008; Ziegler et al., 2015) in a framework reminiscent of the three-factor rule defined by Eqs. (1) and (2) above. However, there are two noteworthy differences. First, in contrast to reward-based learning, the decay time  $\tau_e$  of the synaptic tag  $e_{ij}$  is in the range of one hour rather than one second, consistent with slice experiments (Frey and Morris, 1997) as well as with behavioral experiments (Moncada and Viola, 2007). Second, in slices, the measured synaptic weights  $w_{ij}$  are increased a few minutes after the end of the induction protocol and decay back with the time course of the synaptic tag whereas in the simplest implementation of the three-factor rule framework as formulated in Eqs. (1) and (2) the visible weight is only updated at the moment when the third factor is present. However, slightly more involved models where the visible weight depends on both the synaptic tag variable and the long-term stable weight (Clopath et al., 2008; Ziegler et al., 2015) correctly account for the time course of the measured synaptic weights in consolidation experiments (Frey and Morris, 1997; Reymann and Frey, 2007; Redondo and Morris, 2011).

In summary, the neoHebbian three-factor rule framework has a wide range of applicability. It is well established in the context of synaptic consolidation where the duration of the flag ('synaptic tag') extracted from slice experiments (Frey and Morris, 1997) is in the range of one hour, consistent with fear conditioning experiments (Moncada and Viola, 2007). This time scale is significantly longer than what is needed for behavioral learning of elementary actions or for memorizing surprising events. Theoreticians therefore hypothesized that a process analogous to setting a tag ('eligibility trace') must also exist on the time scale of one second. The next section discusses some recent experimental evidence supporting this theoretical prediction.

### 3 Experimental evidence for eligibility traces

Recent experimental evidence for eligibility traces in striatum (Yagishita et al., 2014), cortex (He et al., 2015), and hippocampus (Brzosko et al., 2015, 2017; Bittner et al., 2017) is reviewed in the following three subsections.

#### 3.1 Eligibility traces in dendritic spines of medial spiny striatal neurons in nucleus accumbens

In their elegant imaging experiment of dendritic spines of nucleus accumbens neurons, Yagishita et al. (2014) mimicked presynaptic spike arrival by glutamate uncaging (presynaptic factor), paired it with three postsynaptic spikes immediately afterward (postsynaptic factor), repeated this STDP-like pre-before-post sequence ten times, and combined it with optogenetic stimulation of dopamine fibers (3rd factor) at various delays (Yagishita et al., 2014). The ten repetitions of the pre-before-post sequence at 10 Hz took about one second while stimulation of dopaminergic fibers (10 dopamine pulses at 30Hz) projecting from the ventral tegmental area (VTA) to nucleus accumbens took about 0.3s. In their paper, dopamine was counted as delayed by one second if the dopamine stimulation started immediately after the end of the one-second long induction period (delay = difference in switch-on time of STDP and dopamine), but for consistency with other data we define the delay  $d$  here as the time passed since the end of the STDP protocol. After 15 complete trials the spine volume, an indicator of synaptic strength (Matsuzaki et al., 2001), was measured and compared with the spine volume before the induction protocol. The authors found that dopamine promoted spine enlargement only if phasic dopamine was given in a narrow time window during or immediately after the 1s-long STDP protocol; cf. Fig. 2A.

The maximum enlargement of spines occurred if the dopamine signal started during the STDP protocol ( $d = -0.4s$ ), but even at a delay of  $d = 1s$  LTP was still visible. Giving dopamine too early ( $d = -2s$ ) or too late ( $d = +4s$ ) had no effect. Spine enlargement corresponded to an increase in the amplitude of excitatory postsynaptic currents indicating that the synaptic weight was indeed strengthened after the protocol (Yagishita et al., 2014). Thus, we can summarize that we have in the *striatum* a three-factor learning rule for the induction of LTP where the decay of the eligibility trace occurs on a time scale of 1s; cf. Fig. 2A.




Figure 2: Experimental support for synaptic eligibility traces. Fractional weight change (vertical axis) as a function of delay  $d$  of third factor (horizontal axis) for various protocols (schematically indicated at the bottom of each panel). **A**. In striatum medial spiny cells, stimulation of presynaptic glutamatergic fibers (green) followed by three postsynaptic action potentials (STDP with pre-post-post-post at +10ms) repeated 10 times at 10Hz yields LTP if dopamine fibers are stimulated during the presentation ( $d < 0$ ) or shortly afterward ( $d = 0$ s or  $d = 1$ s) but not if dopamine is given with a delay  $d = 4$ s; redrawn after Fig. 1 of (Yagishita et al., 2014), with delay  $d$  defined as time since end of STDP protocol. **B**. In cortical layer 2/3 pyramidal cells, stimulation of two independent presynaptic pathways (green and red) from layer 4 to layer 2/3 by a single pulse combined with a burst of four postsynaptic spikes (orange). If the pre-before-post stimulation was combined with a pulse of norepinephrine (NE) receptor agonist isoproterenol with a delay of 0 or 5s, the protocol gave LTP (blue trace). If the post-before-pre stimulation was combined with a pulse of serotonin (5-HT) of a delay of 0 or 2.5s, the protocol gave LTD (red trace); redrawn after Fig. 6 of (He et al., 2015). **C**. In hippocampus CA1, a post-before-pre ( $\Delta t = -20$ ms) induction protocol yields LTP if dopamine is present during induction or given with a delay  $d$  of zero or one minute, but yields LTD if dopamine is absent or given with a delay of 30min; redrawn after Figs 1F, 3C and 2B (square data point at delay of 1min) of (Brzosko et al., 2015). **D**. In hippocampus CA1, 10 extracellular stimuli of presynaptic fibers at 20Hz cause depolarization of the postsynaptic potential. The timing of a complex spike (calcium plateau potential) triggered by current injection (during 300ms) after a delay  $d$ , is crucial for the amount of LTP. If we interpret presynaptic spike arrival as the first, and postsynaptic depolarization as the second factor, the complex spike could be associated with a third factor; redrawn after Fig.3 of (Bittner et al., 2017). Height of boxes gives a very rough estimate of standard deviation - see original papers and figures for details.

To arrive at these results, Yagishita et al. (2014) concentrated on medial spinal neurons in the nucleus accumbens core, a part of the ventral striatum of the basal ganglia. Functionally, striatum is a particularly interesting candidate for reinforcement learning (Brown and Sharp, 1995; Schultz, 1998; Doya, 2000a; Arleo and Gerstner, 2000; Daw et al., 2005) for several reasons. First, striatum receives highly processed sensory information from neocortex and hippocampus through glutamatergic synapses (Mink, 1996; Middleton and Strick, 2000; Haber et al., 2006). Second, striatum also receives dopamine input associated with reward processing (Schultz, 1998). Third, striatum is, together with frontal cortex, involved in selection of motor action programs (Mink, 1996; Seo et al., 2012).

On the molecular level, the striatal three-factor plasticity depended on NMDA, CaMKII, protein synthesis, and dopamine D1 receptors (Yagishita et al., 2014). CaMKII increases were found to be localized in the spine and to have roughly the same time course as the critical window for phasic dopamine suggesting that CaMKII could be involved in the 'synaptic flag' triggered by the STDP-like induction protocol while protein kinase A (PKA) was found to have an unspecific cell-wide distribution suggesting an interpretation of PKA as a molecule linked to the dopamine-triggered third factor (Yagishita et al., 2014).

### **3.2 Two distinct eligibility traces for LTP and LTD in cortical synapses**

In a recent experiment of He et al. (2015), layer 2/3 pyramidal cells in slices from prefrontal or visual cortex were stimulated by an STDP protocol, either pre-before-post for LTP induction or post-before-pre for LTD induction. A neuromodulator was applied with a delay after a single STDP sequence before the whole protocol was repeated; cf. Fig. 2B. Neuromodulators, either norepinephrine (NE), serotonin (5-HT), dopamine (DA), or acetylcholine (ACh) were ejected from a pipette for 10 seconds or from endogenous fibers (using optogenetics) for one second (He et al., 2015). It was found that NE was necessary for LTP whereas 5-HT was necessary for LTD. DA or ACh agonists had no effect in visual cortex but DA had a positive effect on LTP induction in frontal cortex (He et al., 2015).

For the STDP protocol, He et al. (2015) used extracellular stimulation of two presynaptic pathways from layer 4 to layer 2/3 (presynaptic factor) combined with a burst of 4 postsynaptic action potentials (postsynaptic factor), either pre-before-post or post-before-pre. In a first variant of the experiment, the STDP stimulation was repeated 200 times at 10 Hz corresponding to a total stimulation time of 20s before the NE or 5-HT was given. In a second variant, instead of an STDP protocol, they paired presynaptic stimulation (first factor) with postsynaptic depolarization (second factor) to -10mV to induce LTP, or to -40mV to induce LTD. With both protocols it was found that LTP can be induced if the neuromodulator NE (third factor) arrived with a delay of 5 seconds or less after the LTP protocol, but not 10 seconds. LTD could be induced if 5-HT (third factor) arrives with a delay of 2.5 seconds or less after the LTD protocol, but not 5 seconds (He et al., 2015).

A third variant of the experiment involved optogenetic stimulation of the noradrenaline, dopamine, or serotonin pathway by repeated light pulses during 1s applied immediately, or a

few seconds, after a minimal STDP protocol consisting of a single presynaptic and four postsynaptic pulses (either pre-before-post or post-before-pre), a protocol that is physiologically more plausible. The minimal sequence of STDP pairing and neuromodulation was repeated 40 times at intervals of 20s. Results with optogenetic stimulation were consistent with those mentioned above and showed in addition that application of NE or 5-HT immediately before the STDP stimulus did not induce LTP or LTD. *Overall these results indicate that in visual and frontal cortex, pre-before-post pairing leaves an eligibility trace that decays over 5-10 seconds and that can be converted into LTP by the neuromodulator noradrenaline. Similarly, post-before-pre pairing leaves a shorter eligibility trace that decays over 3 seconds and can be converted into LTD by the neuromodulator serotonin; cf. Fig. 2B.*

Functionally, a theoretical model in the same paper (He et al., 2015) showed that the measured three-factor learning rules with two separate eligibility traces stabilized and prolonged network activity so as to allow 'event prediction'. The authors hypothesized that these three-factor rules were related to reward-based learning in cortex such as perceptual learning in monkey (Schoups et al., 2001) or mice (Poort et al., 2015) or reward prediction (Shuler and Bear, 2006). The relation to surprise was not discussed but might be a direction for further explorations.

Molecularly, the transformation of the Hebbian pre-before-post eligibility trace into LTP involves beta adrenergic receptors and intracellular cyclic AMP whereas the transformation of the post-pre eligibility trace into LTD involves the 5-HT<sub>2c</sub> receptor (He et al., 2015). Both receptors are anchored at the postsynaptic density consistent with a role in the transformation of an eligibility trace into actual weight changes (He et al., 2015).

### **3.3 Eligibility traces in hippocampus**

Two experimental groups studied eligibility traces in CA1 hippocampal neurons using complementary approaches. In the studies of Brzosko et al. (2015,2017), CA1 neurons in hippocampal slices were stimulated during about 8 minutes in an STDP protocol involving 100 repetitions (at 0.2Hz) of pairs of one extracellularly delivered presynaptic stimulation pulse (presynaptic factor) and one postsynaptic action potential (postsynaptic factor) (Brzosko et al., 2015). Repeated pre-before-post with a relative timing +10 ms gave LTP (in the presence of natural endogenous dopamine) whereas post-before-pre (-20ms) gave LTD. However, with additional dopamine (third factor) in the bathing solution, post-before-pre at -20ms gave LTP (Zhang et al., 2009). Similarly, an STDP protocol with post-before-pre at -10ms resulted in LTP when endogenous dopamine was present, but in LTD when dopamine was blocked (Brzosko et al., 2015). Thus dopamine broadens the STDP window for LTP into the post-before-pre regime (Zhang et al., 2009; Pawlak et al., 2010). Moreover, in the presence of ACh during the STDP stimulation protocol, pre-before-post at +10ms also gave LTD (Brzosko et al., 2017). Thus ACh broadens the LTD window.

The crucial experiment of Brzosko et al. (2015) involved a delay in the dopamine (Brzosko et al., 2015). Brzosko et al. started to perfuse dopamine either immediately after the end of the post-before-pre (-20ms) induction protocol or with a delay. Since the dopamine was given

for about 10 minutes, it cannot be considered as a phasic signal – but at least the *start* of the dopamine perfusion was delayed. Brzosko et al. found that the stimulus that would normally have given LTD turned into LTP if the delay of dopamine was in the range of one minute or less, but not if dopamine started 10 minutes after the end of the STDP protocol (Brzosko et al., 2015). Note that for the conversion of LTD into LTP, it was important that the synapses were weakly stimulated at low rate while dopamine was present. Similarly, a prolonged pre-before-post protocol at +10ms in the presence of ACh gave rise to LTD, but with dopamine given with a delay of less than 1 minute the same protocol gave LTP (Brzosko et al., 2017). To summarize, *in the hippocampus a prolonged post-before-pre protocol (or a pre-before-post protocol in the presence of ACh) yields visible LTD, but also sets an invisible synaptic flag for LTP. If dopamine is applied with a delay of less than one minute, the synaptic flag is converted into a positive weight change under continued weak presynaptic stimulation; cf. Fig. 2C.*

Molecularly, the conversion of LTD into LTP after repeated stimulation of pre-before-post pulse pairings depended on NMDA receptors and on the cyclic adenosine monophosphate (cAMP) - PKA signaling cascade (Brzosko et al., 2015). The source of dopamine could be in the Locus Coeruleus which would make a link to arousal and novelty (Takeuchi et al., 2016) or from other dopamine nuclei linked to reward (Schultz, 1998). Since the time scale of the synaptic flag reported in (Brzosko et al., 2015, 2017) was in the range of minutes, the process studied by Brzosko et al. could be related to synaptic consolidation (Frey and Morris, 1997; Reymann and Frey, 2007; Redondo and Morris, 2011; Lisman, 2017) rather than eligibility traces in reinforcement learning where shorter time constants are needed (Izhikevich, 2007; Legenstein et al., 2008; Frémaux et al., 2010, 2013). The computational study in Brzosko et al. (2017) used an eligibility trace with a time constant of 2s and showed that dopamine as a reward signal induced learning of reward location while ACh during exploration enabled a fast relearning after a shift of the reward location (Brzosko et al., 2017).

The second study combined in vivo with in vitro data (Bittner et al., 2017). From in vivo studies it has been known that CA1 neurons in mouse hippocampus can develop a novel, reliable, and rather broadly tuned, place field in a single trial under the influence of a 'calcium plateau potential' (Bittner et al., 2015), visible as a complex spike at the soma. Moreover, an artificially induced complex spike was sufficient to induce such a novel place field in vivo (Bittner et al., 2015, 2017).

In additional slice experiments, several input fibers from CA3 to CA1 neurons were stimulated by 10 pulses from an extracellular electrode during one second. The resulting nearly synchronous inputs at, probably, multiple synapses caused a total EPSP that was about 10mV above baseline at the soma, and potentially somewhat larger in the dendrite, but did not cause somatic spiking of the CA1 neuron. The stimulated synapses showed LTP if the presynaptic stimulation was paired with a calcium plateau potential (complex spike) in the postsynaptic neuron. LTP occurred, even if the presynaptic stimulation stopped one or two seconds before the start of the plateau potential or if the plateau potential started before the presynaptic stimulation (Bittner et al., 2017). The protocol has a remarkable efficiency since potentiation was around 200% after only 5 pairings. Thus, *the joint activation of many synapses sets a flag at the*

*activated synapses which is translated into LTP if a calcium plateau potential (complex spike) occurs a few seconds before or after the synaptic activation; cf. Fig. 2D. Molecularly, the plasticity processes implied NMDA receptors and calcium channels (Bittner et al., 2017).*

Functionally, synaptic plasticity in hippocampus is particularly important because of the role of hippocampus in spatial memory (O'keefe and Nadel, 1978). CA1 neurons get input from CA3 neurons which have a narrow place field. The emergence of a broad place field in CA1 has therefore been interpreted as linking several CA3 neurons (that cover for example the 50 cm of the spatial trajectory traversed by the rat before the current location) to a single CA1 cell that codes for the current location (Bittner et al., 2017). Note that at the typical running speed of rodents, 50 cm correspond to several seconds of running. The broad activity of CA1 cells has therefore been interpreted as a predictive representation of upcoming events or places (Bittner et al., 2017). What could such an upcoming event be? For a rodent exploring a T-maze it might for example be important to develop a more precise spatial representation at the T-junction than inside one of the long corridors. With a broad CA1 place field located at the T-junction, information about the upcoming bifurcation could become available several seconds before the animal reaches the junction.

Bittner et al. interpreted their findings as the signature of an unusual form of STDP with a particularly long coincidence window on the behavioral time scale (Bittner et al., 2017). Given that the time span of several seconds between presynaptic stimulation and postsynaptic complex spike is outside the range of a potential causal relation between input and output, they classified the plasticity rule as non-Hebbian because the presynaptic neurons do not participate in firing the postsynaptic one (Bittner et al., 2017). As an alternative view, we propose to classify the findings of Bittner et al. as the signature of an eligibility trace that was left by the joint occurrence of a presynaptic spike arriving from CA3 (presynaptic factor) and a subthreshold depolarization at the location of the synapse in the postsynaptic CA1 neuron (postsynaptic factor); cf. Fig. 2D. In this view, the setting of the synaptic flag is caused by a 'Hebbian'-type induction, except that on the postsynaptic side there are no spikes but just depolarization, consistent with the role of depolarization as a postsynaptic factor (Artola and Singer, 1993; Ngezahayo et al., 2000; Sjöström et al., 2001; Clopath et al., 2010). In this view, the findings of Bittner et al. suggest that the synaptic flag set by the induction protocol leaves an eligibility trace which decays over 2s. If a plateau potential (related to the third factor) is generated during these two seconds, the eligibility trace caused by the induction protocol is transformed into a measurable change of the synaptic weight. The third factor  $M_{3rd}(t)$  in Eq. (2) could correspond to the complex spike, filtered with a time constant of about one second. Importantly, plateau potentials can be considered as neuron-wide signals (Bittner et al., 2015) triggered by surprising novel or rewarding events (Bittner et al., 2017). In this view, the results of Bittner et al. are consistent with the framework of neoHebbian three-factor learning rules. If the plateau potentials are indeed linked to surprising events, the three-factor rule framework predicts that in vivo many neurons in CA1 receive such a third input as a broadcast-like signal. However, only those neurons that also get, at the same time, sufficiently strong input from CA3 might develop the visible plateau potential (Bittner et al., 2015).

The main difference between the two alternative views is that, in the model discussed in Bittner et al. (2017), *each activated synapse* is marked by an eligibility trace (which is independent of the state of the postsynaptic neuron) whereas in the view of the three-factor rule, the eligibility trace is set only if the presynaptic activation coincides with a strong depolarization of the postsynaptic membrane. Thus, in the model of Bittner et al. the eligibility trace is set by the presynaptic factor alone whereas in the three-factor rule description it is set by the combination of pre- and postsynaptic factors. The two models can be distinguished in future experiments where either the postsynaptic voltage is controlled during presynaptic stimulation or where the number of simultaneously stimulated input fibers is minimized. The prediction of the three-factor rule is that spike arrival at a single synapse, or spike arrival in conjunction with a very small depolarization of less than 2 mV above rest, is not sufficient to set an eligibility trace. Therefore, LTP will not occur in these cases even if a calcium plateau potential occurs one second later.

## 4 Discussion and Conclusion

### 4.1 Policy gradient versus TD-learning

Algorithmic models of TD-learning with discrete states and in discrete time do not need eligibility traces that extend beyond one time step (Sutton and Barto, 1998). In a scenario where the only reward is given in a target state that is several action steps away from the initial state, reward information shifts, over multiple trials, from the target state backwards, even if the one-step eligibility trace connects only one state to the next (Sutton and Barto, 1998). Nevertheless, extended eligibility traces across multiple time steps are considered convenient heuristic tools to speed up learning in temporal difference algorithms such as  $TD(\lambda)$  or  $SARSA(\lambda)$  (Singh and Sutton, 1996; Sutton and Barto, 1998).

In policy gradient methods (Williams, 1992) as well as in continuous space-time TD-learning (Doya, 2000b; Frémaux et al., 2013) eligibility traces appear naturally in the formulation of the problem of reward maximization. Importantly, a large class of TD-learning and policy gradient methods can be formulated as three-factor rules for spiking neurons where the third factor is defined as reward minus expected reward (Frémaux and Gerstner, 2016). In policy gradient methods and related three-factor rules, expected reward is calculated as a running average of the reward (Frémaux et al., 2010) or fixed to zero by choice of reward schedule (Florian, 2007; Legenstein et al., 2008). In TD-learning the expected reward in a given time step is defined as the difference of the value of the current state and that of the next state (Sutton and Barto, 1998). In the most recent large-scale applications of reinforcement learning the expected immediate reward in policy gradient is calculated by a TD-algorithm for state-dependent value estimation (Greensmith et al., 2004; Mnih et al., 2016). An excellent modern summary of Reinforcement Learning Algorithms and their historical predecessors can be found in (Sutton and Barto, 2018).



## 4.2 Specificity

If phasic neuromodulator signals are broadcasted over large areas of the brain, the question arises whether synaptic plasticity can still be selective. In the framework of three-factor rules, specificity is inherited from the synaptic flags which are set by the combination of presynaptic spike arrival *and* an elevated postsynaptic voltage at the location of the synapses. The requirement is met only for a small subset of synapses, because presynaptic alone or postsynaptic activity alone are not sufficient; cf. Fig. 1B. Furthermore, among all the flagged synapses only those that show, over many trials, a correlation with the reward signal will be consistently reinforced (Legenstein et al., 2008; Loewenstein and Seung, 2006). Specificity can further be enhanced by an attentional feedback mechanism (Roelfsema and van Ooyen, 2005; Roelfsema et al., 2010) that restricts the number of eligible synapses to the 'interesting' ones, likely to be involved in the task. Such an attentional gating signal acts as an additional factor and turns the three-factor into a four-factor learning rule (Rombouts et al., 2015).

## 4.3 Mapping to Neuromodulators

The third factor is likely to be related to neuromodulators, but from the perspective of a theoretician there is no need to assign one neuromodulator to surprise and another one to reward. Indeed, the theoretical framework also works if each neuromodulator codes for a different combination of variables such as surprise, novelty or reward, just as we can use different coordinate systems to describe the same physical system (Frémaux and Gerstner, 2016). Thus, whether dopamine is purely reward related or also novelty related (Ljunberg and W. Schultz, 1992; Schultz, 1998; Redgrave and Gurney, 2006) is not critical for the development of three-factor learning rules as long as dimensions relating to novelty, surprise, and reward are all covered by the set of neuromodulators.

Complexity in biology is increased by the fact that dopamine neurons projecting from the VTA to the striatum can have separate circuits and functions changing from reward in ventral striatum to novelty in the tail of striatum (Menegas et al., 2017). Similarly, dopaminergic fibers starting in the VTA can have a different function than those starting in Locus Coeruleus (Takeuchi et al., 2016). The framework of three-factor rules is general enough to allow for these, and many other, variations.

## 4.4 Alternatives to eligibility traces for bridging the gap between the behavioral and neuronal timescales

From a theoretical point of view, there is nothing – apart conceptual elegance – to favor eligibility traces over alternative neuronal mechanisms to associate events that are separated by a second or more. For example, memory traces hidden in the rich firing activity patterns of a recurrent network (Maass et al., 2002; Jaeger and Haas, 2004; Buonomano and Maass, 2009; Sussillo, 2009) or short-term synaptic plasticity in recurrent networks (Mongillo et al., 2008) could be involved in learning behavioral tasks with delayed feedback. In some models, neuronal,

rather than synaptic, activity traces have been involved in learning a delayed paired-associate task (Brea et al., 2016) and a combination of synaptic eligibility traces with prolonged single-neuron activity has been used for learning on behavioral time scales (Rombouts et al., 2015). The empirical studies reviewed here support the idea that the brain makes use of the elegant solution with synaptic eligibility traces and three-factor learning rules, but do not exclude that other mechanisms work in parallel.

#### 4.5 The paradoxical nature of predictions in computational neuroscience

If a neuroscientist thinks of a theoretical model, he often imagines a couple of assumptions at the beginning, a set of results derived from simulations or mathematical analysis, and ideally a few novel predictions - but is this the way modeling works? There are at least two types of predictions in computational neuroscience, detailed predictions and conceptual predictions. Well-known examples of detailed predictions have been generated from variants of multi-channel biophysical Hodgkin-Huxley type (Hodgkin and Huxley, 1952) models such as: 'if channel X is blocked then we predict that ...' where X is a channel with known dynamics and predictions include depolarization, hyperpolarization, action potential firing, action potential backpropagation or failure thereof. All of these are useful predictions readily translated to and tested in experiments.

Conceptual predictions derived from abstract conceptual models are potentially more interesting, but more difficult to formulate. Conceptual models develop ideas and form our thinking of how a specific neuronal system could work to solve a behavioral task such as working memory (Mongillo et al., 2008), action selection and decision making (Sutton and Barto, 1998), long-term stability of memories (Lisman, 1985; Crick, 1984; Fusi et al., 2005), memory formation and memory recall (Willshaw et al., 1969; Hopfield, 1982). Paradoxically these models often make no detailed predictions in the sense indicated above. Rather, in these and other conceptual theories, the most relevant model features are formulated as *assumptions* which may be considered, in a loose sense, as playing the role of *conceptual predictions*. To formulate it as a short slogan: Assumptions are predictions. Let us return to the conceptual framework of three-factor rules: the purification of rough ideas into the role of three factors is the important conceptual work - and part of the assumptions. Moreover, the specific choice of time constant in the range of one second for the eligibility trace has been formulated by theoreticians as one of the model assumptions, rather than as a prediction; cf. the footnotes in section 'Examples and theoretical predictions'. Why is this the case?

Most theoreticians shy away from calling their conceptual modeling work a 'prediction', because there is no logical necessity that the brain must work the way they assume in their model - the brain could have found a less elegant, different, but nevertheless functional solution to the problem under consideration; see the examples in the previous subsection. What a good conceptual model in computational neuroscience shows is that there *exists* a (nice) solution that should ideally not be in obvious contradiction with too many known facts. Importantly, conceptual models necessarily rely on assumptions which in many cases have not (yet) been

shown to be true. The response of referees to modeling work in experimental journals therefore often is: 'but this has never been shown'. Indeed, some assumptions may look far-fetched or even in contradiction with known facts: for example, to come back to eligibility traces, experiments on synaptic tagging-and-capture have shown in the 1990s that the time scale of a synaptic flag is in the range of one *hour* (Frey and Morris, 1997; Reymann and Frey, 2007; Redondo and Morris, 2011; Lisman, 2017), whereas the theory of eligibility traces for action learning needs a synaptic flag on the time scale of one *second*. Did synaptic tagging results imply that three-factor rules for action learning were wrong, because they used the wrong time scale? Or, on the contrary, did these experimental results rather imply that a biological machinery for three-factor rules was indeed in place which could therefore, for other neuron types and brain areas, be used and re-tuned to a different time scale (Frémaux et al., 2013)?

As mentioned earlier, the concepts of eligibility traces and three-factor rules can be traced back to the 1960ies, from models formulated in words (Crow, 1968), to firing rate models formulated in discrete time and discrete states (Klopf, 1972; Sutton and Barto, 1981; Barto et al., 1983; Barto, 1985; Williams, 1992; Schultz, 1998; Sutton and Barto, 1998; Bartlett and Baxter, 1999), to models with spikes in a continuous state space and an explicit time scale for eligibility traces (Xie and Seung, 2004; Loewenstein and Seung, 2006; Florian, 2007; Izhikevich, 2007; Legenstein et al., 2008; Vasilaki et al., 2009; Frémaux et al., 2013). Despite the mismatch with the known time scale of synaptic tagging in hippocampus (and lack of experimental support in other brain areas), theoreticians persisted, polished their theories, talked at conferences about these models, until eventually the experimental techniques and the scientific interests of experimentalists were aligned to directly test the assumptions of these theories. In view of the long history of three-factor learning rules, the recent elegant experiments (Yagishita et al., 2014; He et al., 2015; Brzosko et al., 2015, 2017; Bittner et al., 2017) provide an instructive example of how conceptual theories can influence experimental neuroscience.

## Funding and Acknowledgments

This project has been funded by the European Research Council (grant agreement no. 268 689, “MultiRules”), by the European Union Horizon 2020 Framework Program under grant agreement no. 720270 (Human Brain Project), and by the Swiss National Science Foundation (no. 200020 165538).

## References

- Arleo, A. and W. Gerstner (2000). Spatial cognition and neuro-mimetic navigation: a model of hippocampal place cell activity. *Biological Cybernetics* 83(3), 287–299.
- Artola, A. and W. Singer (1993). Long-term depression of excitatory synaptic transmission and its relationship to long-term potentiation. *Trends Neurosci.* 16(11), 480–487.
- Bailey, C., M. Giustetto, Y.-Y. Huang, R. Hawkins, and E. Kandel (2000). Is heterosynaptic modulation essential for stabilizing hebbian plasticity and memory. *Nat. Rev. Neurosci.* 1, 11–20.

- Barrett, A., G. Billings, R. Morris, and M. van Rossum (2009). State based model of long-term potentiation and synaptic tagging and capture. *PLOS Comput. Biol.* 5, e1000259.
- Bartlett, P. L. and J. Baxter (1999). Hebbian synaptic modification in spiking neurons that learn. Technical report, Australian National University.
- Barto, A., R. Sutton, and C. Anderson (1983). Neuronlike adaptive elements that can solve difficult learning and control problems. *IEEE transactions on systems, man, and cybernetics* 13, 835–846.
- Barto, A.G. (1985). Learning by statistical cooperation of self-interested neuron-like computing elements. *Human Neurobiol.* 4, 229–256.
- Benna, M. and S. Fusi (2016). Computational principles of synaptic memory consolidation. *Nat. Neurosci.* 19, 1697–1706.
- Bienenstock, E., L. Cooper, and P. Munroe (1982). Theory of the development of neuron selectivity: orientation specificity and binocular interaction in visual cortex. *J. Neurosci.* 2, 32–48.
- Bittner, K., C. Grienberger, S. Vaidya, A. Milstein, J. Macklin, J. Suh, S. Tonegawa, and J. Magee (2015). Conjunctive input processing drives feature selectivity in hippocampal ca1 neurons. *Nat. Neurosci.* 357, 1133–1142. doi:10.1038/nn.4062.
- Bittner, K., A. Milstein, C. Grienberger, S. Romani, and J. Magee (2017). Behavioral time scale synaptic plasticity underlies ca1 place fields. *Nature* 357, 1033–1036.
- Bliss, T. and T. Lømo (1973). Long-lasting potentiation of synaptic transmission in the dentate area of anaesthetized rabbit following stimulation of the perforant path. *J. Physiol.* 232, 351–356.
- Bliss, T. V. P. and G. L. Collingridge (1993). A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 361, 31–39.
- Bosch, M., J. Castro, T. Saneyoshi, H. Matsuno, M. Sur, and Y. Hayashi (2014). Structural and molecular remodeling of dendritic spine substructures during long-term-potentiation. *Neuron* 82, 444–459.
- Brader, J., W. Senn, and S. Fusi (2007). Learning real-world stimuli in a neural network with spike-driven synaptic dynamics. *Neural Computation* 19, 2881–2912.
- Brea, J., A. Gaal, R. Urbancik, and W. Senn (2016). Prospective coding by spiking neurons. *PLOS Comput. Biol.* 12, e100500. doi:10.1371/journal.pcbi.100500.
- Brown, M. and P. Sharp (1995). Simulation of spatial-learning in the Morris water maze by a neural network model of the hippocampal-formation and nucleus accumbens. *Hippocampus* 5, 171–188.
- Brzosko, Z., W. Schultz, and O. Paulsen (2015). Retroactive modulation of spike timingdependent plasticity by dopamine. *eLife* 4, e09685. DOI: 10.7554/eLife.09685.
- Brzosko, Z., S. Zannone, W. Schultz, , C. Clopath, and O. Paulsen (2017). Sequential neuromodulation of hebbian plasticity offers mechanism for effective reward-based navigation. *eLife* 6, e27756. DOI: 10.7554/eLife.27756.
- Buonomano, D. and W. Maass (2009). State-dependent computations: spatiotemporal processing in cortical networks. *Nat. Rev. Neuroscience* 10, 113–125.
- Clopath, C., L. Busing, E. Vasilaki, and W. Gerstner (2010). Connectivity reflects coding: A model of voltage-based spike-timing-dependent-plasticity with homeostasis. *Nature Neuroscience* 13, 344–352.
- Clopath, C., L. Ziegler, E. Vasilaki, L. Busing, and W. Gerstner (2008). Tag-trigger-consolidation: A model of early and late long-term-potentiation and depression. *PLOS Comput. Biol.* 4, e1000248.

- Crick, F. (1984). Neurobiology - memory and molecular turnover. *Nature* 312, 101.
- Crow, T. (1968). Cortical synapses and reinforcement: a hypothesis. *Nature* 219, 736–737.
- Daw, N., Y. Niv, and P. Dayan (2005). Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control. *Nature Neuroscience* 8, 1704–1711.
- Deger, M., M. Helias, S. Rotter, and M. Diesmann (2012). Spike-timing dependence of structural plasticity explains cooperative synapse formation in the neocortex. *PLOS Comput. Biol.* 8, e1002689.
- Doya, K. (2000a). Complementary roles of basal ganglia and cerebellum in learning and motor control. *Current opinion in neurobiology* 10(6), 732–739.
- Doya, K. (2000b). Temporal difference learning in continuous time and space. *Neural Comput.* 12, 219–245.
- Faraji, M., K. Preuschoff, and W. Gerstner (2018). Balancing new against old information: The role of puzzlement surprise in learning. *Neural Computation* 30, xx–xx.
- Fauth, M., F. Worgotter, and C. Tetzlaff (2015). The formation of multi-synaptic connections by the interaction of synaptic and structural plasticity and their functional consequences. *PLOS Comput. Biol.* 11, e1004031.
- Florian, R. V. (2007). Reinforcement learning through modulation of spike-timing-dependent synaptic plasticity. *Neural Computation* 19, 1468–1502.
- Foster, D., R. Morris, and P. Dayan (2000). Models of hippocampally dependent navigation using the temporal difference learning rule. *Hippocampus* 10, 1–16.
- Frémaux, N. and W. Gerstner (2016). Neuromodulated spike-timing dependent plasticity and theory of three-factor learning rules. *Front. Neural Circuits* 9, 85.
- Frémaux, N., H. Sprekeler, and W. Gerstner (2010). Functional requirements for reward-modulated spike-timing-dependent plasticity. *J. Neurosci.* 40, 13326–13337.
- Frémaux, N., H. Sprekeler, and W. Gerstner (2013). Reinforcement learning using continuous time actor-critic framework with spiking neurons. *PLOS Comput. Biol.* 9, 4.
- Frey, U. and R. Morris (1997). Synaptic tagging and long-term potentiation. *Nature* 385, 533 – 536.
- Friston, K. (2010). The free-energy principle: a unified brain theory? *Nat. Rev. Neurosci.* 11, 127–138.
- Friston, K., T. Fitzgerald, F. Rigoli, P. Schwartenbeck, J. O’Doherty, and G. Pezzulo (2016). Active inference and learning. *Neurosci. and Behav. Rev.* 68, 862–879.
- Froemke, R. and Y. Dan (2002). Spike-timing dependent plasticity induced by natural spike trains. *Nature* 416, 433–438.
- Fusi, S., P. Drew, and L. Abbott (2005). Cascade models of synaptically stored memories. *Neuron* 45, 599–611.
- Gerstner, W., R. Kempter, J. van Hemmen, and H. Wagner (1996). A neuronal learning rule for sub-millisecond temporal coding. *Nature* 383(6595), 76–78.
- Gjorjjeva, J., C. Clopath, J. Audet, and J. Pfister (2011). A triplet spike-timingdependent plasticity model generalizes the bienenstockcoopermunro rule to higher-order spatiotemporal correlations. *Proc. Natl. Sci. Acad. USA* 108, 19383–19388.
- Graupner, M. and N. Brunel (2007). Stdp in a bistable synapse model based on CaMKII and associate signaling pathways. *PLOS Comput. Biol.* 3, e221. doi:10.1371/journal.pcbi.0030221.

- Greensmith, E., P. Bartlett, and J. Baxter (2004). Variance reduction techniques for gradient estimates in reinforcement learning. *J. Machine Learn. Res.* 5, 1471–1530.
- Gu, Q. (2002). Neuromodulatory transmitter systems in the cortex and their role in cortical plasticity. *Neuroscience* 111, 815–835.
- Haber, S., K.-S. Kim, P. Mailly, and R. Calzavara (2006). Reward-related cortical inputs define a large striatal region in primates that interface with associative cortical connections, providing a substrate for incentive-based learning. *J. Neurosci.* 26, 8368–8376.
- Hasselmo, M. (2006). The role of acetylcholine in learning and memory. *Curr. Opinion Neurobiol.* 16, 710–715.
- He, K., M. Huertas, S. Z. Hong, X. T. abnd Johannes W Hell, H. Shouval, and A. Kirkwood (2015). Distinct eligibility traces for ltp and ltd in cortical synapses. *Neuron* 88, 528–538. <http://dx.doi.org/10.1016/j.neuron.2015.09.037>.
- Hebb, D. O. (1949). *The Organization of Behavior*. New York: Wiley.
- Helias, M., S. Rotter, M.-O. Gewaltig, and M. Diesmann (2008). Structural plasticity controlled by calcium based correlation detection. *Front. Comput. Neurosci.* 2, 7.
- Hess, E. and J. Polt (1960). Pupil size as related to interest value of visual stimuli. *Science* 132, 349–350.
- Hodgkin, A. L. and A. F. Huxley (1952). A quantitative description of membrane current and its application to conduction and excitation in nerve. *J Physiol* 117(4), 500–544.
- Hopfield, J. J. (1982). Neural networks and physical systems with emergent collective computational abilities. *Proc. Natl. Acad. Sci. USA* 79, 2554–2558.
- Huganir, R. and R. Nicoll (2013). AMPARs and synaptic plasticity: the last 25 years. *Neuron* 80, 704–717.
- Itti, L. and P. Baldi (2009). Bayesian surprise attracts human attention. *Vision Research* 49, 1295–1306.
- Izhikevich, E. (2007). Solving the distal reward problem through linkage of STDP and dopamine signaling. *Cerebral Cortex* 17, 2443–2452.
- Izhikevich, E. and N. Desai (2003). Relating stdp to bcm. *Neural Computation* 15, 1511–1523.
- Jaeger, H. and H. Haas (2004). Harnessing nonlinearity: Predicting chaotic systems and saving energy in wireless communication. *Science* 304, 78–80.
- J.Blac, J. Belluzzi, and L. Stein (1985). Reinforcement delay of one second severley impairs acquisition of brain self-stimulation. *Brain Res.* 359, 113–119.
- Kalman, R. (1960). A new approach to linear filtering and prediciton problems. *J. Basic Eng.* 82, 35–45.
- Kempter, R., W. Gerstner, and J. L. van Hemmen (1999). Hebbian learning and spiking neurons. *Phys. Rev. E* 59, 4498–4514.
- Klopf, A. (1972). Brain function and adaptive systems - a heterostatic theory. *Air Force Cambridge research laboratories, special reports* 133, 1–70.
- Kolossa, A., T. Fingscheidt, K. Wessel, and B. Kopp (2013). A model-based approach to trial-by-trial p300 amplitude fluctuations. *Front. Hum. Neurosci.* 6, 359.
- Kolossa, A., B. Kopp, and T. Fingscheidt (2015). A computational analysis of the neural bases of bayesian inference. *NeuroImage* 106(222-237).

- Legenstein, R., D. Pecevski, and W. Maass (2008). A learning theory for reward-modulated spike-timing-dependent plasticity with application to biofeedback. *PLoS Comput. Biol.* 4, e1000180.
- Levy, W. B. and D. Stewart (1983). Temporal contiguity requirements for long-term associative potentiation/depression in hippocampus. *Neurosci*, 8, 791–797.
- Lisman, J. (1985). A mechanism for memory storage insensitive to molecular turnover: a bistable autophosphorylating kinase. *Proc. Natl. Acad. Sci. USA* 82, 3055–3057.
- Lisman, J. (2003). Long-term potentiation: outstanding questions and attempted synthesis. *Phil. Trans. R. Soc. Lond B: Biological Sciences* 358, 829 – 842.
- Lisman, J. (2017). Glutamatergic synapses are structurally and biochemically complex because of multiple plasticity processes: long-term potentiation, long-term depression, short-term potentiation and scaling. *Phil. Trans. Roy. Soc. B* 372, 20160260.
- Lisman, J., A. Grace, and E. Duzel (2011). A neoHebbian framework for episodic memory; role of dopamine-dependent late ltp. *Trends Neurosci.* 34, 536–547.
- Little, D. and F. Sommer (2013). Learning and exploration in action-perception loops. *Front. Neural Circuits* 7, 37.
- Ljunberg, T. and P. A. and W. Schultz (1992). Responses of monkey dopamine neurons during learning of behavioral interactions. *J. Neurophysiol.* 67, 145–163.
- Loewenstein, Y. and H. Seung (2006). Operant matching is a generic outcome of synaptic plasticity based on the covariance between reward and neural activity. *Proc. Natl. Acad. Sci. USA* 103, 15224–15229.
- Lowel, S. and W. Singer (1992). Selection of intrinsic horizontal connections in the visual cortex by correlated neuronal activity. *Science* 255, 209–212.
- Maass, W., T. Natschläger, and H. Markram (2002). Real-time computing without stable states: a new framework for neural computation based on perturbations. *Neural Computation*, 2531–2560.
- Markram, H., J. Lübke, M. Frotscher, and B. Sakmann (1997). Regulation of synaptic efficacy by coincidence of postsynaptic AP and EPSP. *Science* 275, 213–215.
- Martin, S., P. Grimwood, and R. Morris (2000). Synaptic plasticity and memory: an evaluation of the hypothesis. *Ann. Rev. Neurosci.* 23, 649–711.
- Mathys, C., J. Daunizeau, K. Friston, and K. Stephan (2011). A bayesian foundation for individual learning under uncertainty. *Front. Hum. Neurosci.* 5, 39.
- Mathys, C. D., E. I. Lomakina, K. Daunizeau, S. Iglesias, K. Brodersen, K. Friston, and K. Stephan (2014). Uncertainty in perception and the hierarchical gaussian filter. *Front. Hum. Neurosci.* 8, 825.
- Matsuzaki, M., G. Ellis-Davies, T. Nemoto, M. Iione, Y. Miyashita, and H. Kasai (2001). Dendritic spine geometry is critical for ampa receptor expression in hippocampal ca1 pyramidal neurons. *Nat. Neurosci.* 4, 1086–1092.
- Menegas, W., B. Babayan, N. Uchida, and M. Watabe-Uchida (2017). Opposite initialization to novel cues in dopamine signaling in ventral and posterior striatum in mice. *eLife* 6, e21886. doi: 10.7554/eLife.21886.
- Meyniel, F., M. Maheu, and S. Dehaene (2016). Human inferences about sequences: A minimal transition probability model. *PLoS Comput. Biol.* 12, e1005260.
- Middleton, F. A. and P. Strick (2000). Basal ganglia and cerebellar loops: motor and cognitive circuitry. *Brain Res. Rev.* 31, 236–250.

- Miller, K. D. and D. J. C. MacKay (1994). The role of constraints in hebbian learning. *Neural Computation* 6, 100–126.
- Mink, J. (1996). The basal ganglia: focused selection and inhibition of competing motor programs. *Progr. Neurobiol.* 50, 381–425.
- Mnih, V., A. Badia, M. Mirza, A. Graves, T. Harley, T. Lillicrap, D. Silver, and K. Kavukcuoglu (2016). Asynchronous methods for deep reinforcement learning. *arXiv 1602.01783v2*, 1–19.
- Moncada, D. and H. Viola (2007). Induction of long-term memory by exposure to novelty requires protein synthesis: Evidence for a behavioral tagging. *J. Neurosci* 27, 7476–7481.
- Mongillo, G., O. Barak, and M. Tsodyks (2008). Synaptic theory of working memory. *Science* 319, 1543–1546.
- Nassar, M., R. Wilson, B. Heasley, and J. Gold (2010). An approximately bayesian delta-rule model explains the dynamics of belief updating in a changing environment. *J. Neurosci.* 30, 12366–12378.
- Ngezahayo, A., M. Schachner, and A. Artola (2000). Synaptic activation modulates the induction of bidirectional synaptic changes in adult mouse hippocampus. *J. Neuroscience* 20, 2451–2458.
- Oja, E. (1982). A simplified neuron model as a principal component analyzer. *J. Mathematical Biology* 15, 267–273.
- O’keefe, J. and L. Nadel (1978). *The hippocampus as a cognitive map*, Volume 3. Clarendon Press Oxford.
- Okouchi, H. (2009). Response acquisition by humans with delayed reinforcement. *J. Exp. Anal. of Beh.* 91, 377–390.
- Oudeyer, P., F. Kaplan, and V. Hafner (2007). Intrinsic motivation systems for autonomous mental development. *IEEE Trans. Evol. Comput.* 11, 265–286.
- Pavlov, I. P. (1927). *Conditioned reflexes*. Oxford Univ. Press.
- Pawlak, V., J. Wickens, A. Kirkwood, and J. Kerr (2010). Timing is not everything: neuromodulation opens the STDP gate. *Front. Synaptic Neurosci.* 2, 146.
- Pfister, J.-P., T. Toyozumi, D. Barber, and W. Gerstner (2006). Optimal spike-timing dependent plasticity for precise action potential firing in supervised learning. *Neural Computation* 18, 1318–1348.
- Poort, H., A. Khan, M. Pachitariu, A. Nemri, I. Orsolich, J. Krupic, M. Bauza, M. Sahani, G. Keller, T. Mrsic-Flogel, and S. Hofer (2015). Learning enhances sensory and multiple non-sensory representations in primary visual cortex. *Neuron* 86, 1478–1490.
- Redgrave, P. and K. Gurney (2006). The short-latency dopamine signal: a role in discovering novel actions? *Nat. Rev. Neurosci.* 7, 967–975.
- Redondo, R. L. and R. G. M. Morris (2011). Making memories last: the synaptic tagging and capture hypothesis. *Nat Rev Neurosci* 12, 17–30.
- Reymann, K. and J. Frey (2007). The late maintenance of hippocampal ltp: requirements, phases, synaptic tagging, late associativity and implications. *Neuropharmacology* 52, 24–40.
- Reynolds, J. and J. Wickens (2002). Dopamine-dependent plasticity of corticostriatal synapses. *Neural Networks* 15, 507–521.
- Roelfsema, P. and A. van Ooyen (2005). Attention-gated reinforcement learning of internal representations for classification. *Neural Computation* 17, 2176–2214.



- Roelfsema, P., A. van Ooyen, and T. Watanabe (2010). Perceptual learning rules based on reinforcers and attention. *Trends in Cogn. Sci.* 14, 64–71. doi:10.1016/j.tics.2009.11.00.
- Rombouts, J., S. Bothe, and R. Roelfsema (2015). How attention can create synaptic tags for the learning of working memories in sequential tasks. *PLoS Comput. Biol.* 11, e1004060.
- Rubin, J., R. Gerkin, G.-Q. Bi, and C. Chow (2005). Calcium time course as a signal for spike-timing-dependent plasticity. *J. Neurophysiology* 93, 2600–2613.
- Rubin, J., D. D. Lee, and H. Sompolinsky (2001). Equilibrium properties of temporally asymmetric Hebbian plasticity. *Physical Review Letters* 86, 364–367.
- Schmidhuber, J. (1991). Curious model-building control systems. In *Proceedings of the International Joint Conference on Neural Networks, Singapore*, Volume 2, pp. 1458–1463. IEEE press.
- Schmidhuber, J. (2006). Developmental robotics, optimal artificial curiosity, creativity, music, and the fine arts. *Connection Science* 18, 173–187.
- Schmidhuber, J. (2010). Formal theory of creativity, fun, and intrinsic motivation (1990–2010). *IEEE Trans. Autom. Mental Developm.* 2, 230–247.
- Schoups, A., R. Vogels, N. Quian, and G. A. Orban (2001). Practising orientation identification improves orientation coding in v1 neurons. *Nature* 412, 549–553.
- Schultz, W. (1998). Predictive reward signal of dopamine neurons. *J. Neurophysiooogy* 80, 1–27.
- Schultz, W. (2002, 10). Getting formal with dopamine and reward. *Neuron* 36(2), 241–263.
- Schultz, W., P. Dayan, and R. Montague (1997). A neural substrate for prediction and reward. *Science* 275, 1593–1599.
- Schultz, W. and A. Dickinson (2000). Neuronal coding of prediction errors. *Annual Reviews of Neuroscience* 23, 472–500.
- Senn, W., M. Tsodyks, and H. Markram (2001). An algorithm for modifying neurotransmitter release probability based on pre- and postsynaptic spike timing. *Neural Computation* 13, 35–67.
- Seo, M., E. Lee, and B. Averbeck (2012). Action selection and action value in frontal-striatal circuits. *Neuron* 74, 947–960.
- Shannon, C. (1948). A mathematical theory of communication. *Bell System Technical Journal* 27, 37–423.
- Shatz, C. (1992). The developing brain. *Sci. Am.* 267, 60–67.
- Shouval, H. Z., M. F. Bear, and L. N. Cooper (2002). A unified model of nmda receptor dependent bidirectional synaptic plasticity. *Proc. Natl. Acad. Sci. USA* 99, 10831–10836.
- Shuler, M. and M. Bear (2006). Reward timing in the primary visual cortex. *Science* 311, 1606–1609.
- Singh, S., A. Barto, and N. Chentanez (2004). Intrinsically motivated reinforcement learning. *Advances of Neural Information Processing Systems, NIPS 17*, 1281–1288.
- Singh, S. and R. Sutton (1996). Reinforcement learning with replacing eligibility traces. *Machine Learning* 22, 123–158.
- Sjöström, P., G. Turrigiano, and S. Nelson (2001). Rate, timing, and cooperativity jointly determine cortical synaptic plasticity. *Neuron* 32, 1149–1164.

- Song, S., K. Miller, and L. Abbott (2000). Competitive Hebbian learning through spike-time-dependent synaptic plasticity. *Nature Neuroscience* 3, 919–926.
- Squires, K., C. Wickens, N. Squires, and E. Donchin (1976). The effect of stimulus sequence on the waveform of the cortical event-related potential. *Science* 193, 1141–1146.
- Storck, J., S. Hochreiter, and J. Schmidhuber (1995). Reinforcement-driven information acquisition in non-deterministic environments. In *Proc. ICANN'95, vol.2*, pp. 159–164. EC2-CIE, Paris.
- Sun, Y., F. Gomez, and J. Schmidhuber (2011). Planning to be surprised: Optimal bayesian exploration in dynamic environments. In *Artificial General Intelligence*, pp. 41–51. Springer.
- Suri, R. E. and W. Schultz (1999). A neural network with dopamine-like reinforcement signal that learns a spatial delayed response task. *Neuroscience* 91, 871890.
- Sussillo, D. (2009). Generating coherent patterns of activity from chaotic neural networks. *Neuron* 63, 544–557.
- Sutton, R.S and A.G. Barto (1998). *Reinforcement learning*. MIT Press, Cambridge.
- Sutton, R.S. and A.G. Barto (2018). *Reinforcement learning: an introduction*. MIT Press, Cambridge, 2nd edition
- Sutton, R. S. and A. G. Barto (1981). Towards a modern theory of adaptive networks: expectation and prediction. *Psychol. Review* 88, 135–171.
- Takeuchi, T., A. Duzskiewicz, A. Sonneborn, P. Spooner, M. Yamasaki, M. Watanabe, C. Smith, G. Fernandez, K. Deisseroth, R. Greene, and R. Morris (2016). Locus coeruleus and dopaminergic consolidation of everyday memory. *Nature* 537, 357362.
- Thorndike, E. (1911). *Animal Intelligence*. Darien, CT: Hafner.
- van Rossum, M. C. W., G. Q. Bi, and G. G. Turrigiano (2000). Stable Hebbian learning from spike timing-dependent plasticity. *J. Neuroscience* 20, 8812–8821.
- Vasilaki, E., N. Frémaux, R. Urbanczik, W. Senn, and W. Gerstner (2009). Spike-based reinforcement learning in continuous state and action space: When policy gradient methods fail. *PLOS Comput.Biol.* 5, e1000586.
- Williams, R. (1992). Simple statistical gradient-following methods for connectionist reinforcement learning. *Machine Learning* 8, 229–256.
- Willshaw, D. J., O. P. Bunemann, and H. C. Longuet-Higgins (1969). Non-holographic associative memory. *Nature* 222, 960–962.
- Xie, X. and S. Seung (2004). Learning in neural networks by reinforcement of irregular spiking. *Phys. Rev. E* 69, 41909.
- Yagishita, S., A. Hayashi-Takagi, G. Ellis-Davies, H. Urakubo, S. Ishii, and H. Kasai (2014). A critical time window for dopamine actions on the structural plasticity of dendritic spines. *Science* 345, 1616–1620.
- Yu, A. and P. Dayan (2005). Uncertainty, neuromodulation, and attention. *Neuron* 46, 681–692.
- Zhang, J., P. Lau, and G. Bi (2009). Gain in sensitivity and loss in temporal contrast of stdp by dopaminergic modulation at hippocampal synapses. *Proc. Natl. Aca. Sci. USA* 106, 13028–13033.
- Zhang, L., H. Tao, C. Holt, W.A.Harris, and M.-M. Poo (1998). A critical window for cooperation and competition among developing retinotectal synapses. *Nature* 395, 37–44.
- Ziegler, L., F. Zenke, D. Kastner, and W. Gerstner (2015). Synaptic consolidation: from synapses to behavioral modeling. *J. Neuroscience* 35, 1319–1334.