On the complexity of the outer-connected bondage and the outer-connected reinforcement problems

M. Hashemipour¹, M. R. Hooshmandasl², A. Shakiba³

^{1,2}Department of Computer Science, Yazd University, Yazd, Iran.

^{1,2,3}The Laboratory of Quantum Information Processing, Yazd University, Yazd, Iran.

³Department of Computer Science, Vali-e-Asr University of Rafsanjan, Rafsanjan, Iran e-mail: ¹mhashemi@stu.yazd.ac.ir, ²hooshmandasl@yazd.ac.ir, ³ali.shakiba@vru.ac.ir.

Abstract

Let G=(V,E) be a graph. A subset $S\subseteq V$ is a dominating set of G if every vertex not in S is adjacent to a vertex in S. A set $\tilde{D}\subseteq V$ of a graph G=(V,E) is called an outer-connected dominating set for G if (1) \tilde{D} is a dominating set for G, and (2) $G[V\setminus \tilde{D}]$, the induced subgraph of G by $V\setminus \tilde{D}$, is connected. The minimum size among all outer-connected dominating sets of G is called the outer-connected domination number of G and is denoted by $\tilde{\gamma}_c(G)$. We define the outer-connected bondage number of a graph G as the minimum number of edges whose removal from G results in a graph with an outer-connected domination number larger than the one for G. Also, the outer-connected reinforcement number of a graph G is defined as the minimum number of edges whose addition to G results in a graph with an outer-connected domination number, which is smaller than the one for G. This paper shows that the decision problems for the outer-connected bondage and the outer-connected reinforcement numbers are NP-hard. Also, the exact values of the bondage number are determined for several classes of graphs.

1 Introduction

The terminology and notation on graph theory in this paper in general follows the reference [9]. Let G = (V, E) be a graph with vertex set V and edge set E. The graph G is called to be of order |V| and size |E|. Also, we use V(G) and E(G) to denote the vertex set and the edge set for the graph G, respectively. Let V be a vertex in V. The open neighborhood of V is denoted by $N_G(V)$ and is defined as $\{u \in V : \{u, v\} \in E(G)\}$. Similarly, the closed neighborhood of V is denoted by $V_G(V)$ and is defined as $V_G(V)$ whenever the graph $V_G(V)$ is a vertex, we simply write $V_G(V)$ and $V_G(V)$ to denote $V_G(V)$ and $V_G(V)$, respectively. A leaf vertex in $V_G(V)$ is a vertex of degree one. We denote the path of order $V_G(V)$ be a vertex of order $V_G(V)$ is a subset $V_G(V)$ and the star of order $V_G(V)$ is a dominating set of $V_G(V)$ is called a $V_G(V)$ is called a

Domination is one of the most widely studied topics in graph theory, e.g. [9,18] and references therein. This paper studies some issues in a particular variation of domination, namely the *outer-connected domination*. The concept of outer-connected domination number is introduced by Cyman in [3] and is further studied by others in [1,13]. The outer-connected domination problem is shown to be an **NP**-complete problem for arbitrary graphs in [3]. A set $\tilde{D} \subseteq V$ of a graph G = (V, E) is called an *outer-connected dominating set* for G if (1) \tilde{D} is a dominating set for G, and (2) $G[V \setminus \tilde{D}]$, the induced subgraph of G by $V \setminus \tilde{D}$, is connected. The minimum size among all outer-connected dominating sets of G is called the *outer-connected domination number* of G and is denoted by $\tilde{\gamma}_c(G)$ [3]. An outer-connected dominating set \tilde{D} is called a $\tilde{\gamma}$ - set of G if $|\tilde{D}| = \tilde{\gamma}_c(G)$.

In this paper, we focus on two graph alterations and their effects on the outer-connected domination number, (1) the removal of edges from a graph and (2) the addition of edges to a graph. The bondage and the reinforcement numbers are two important parameters for measuring the vulnerability and the stability of the network domination under link failure and link addition. The bondage number of G, denoted by b(G), is the minimum number of edges whose removal from G results in a graph with a domination number larger than the one for G. The reinforcement number of G, denoted by r(G), is the smallest number of edges whose addition to G results in a graph with a domination number smaller than the one for G. The bondage and the reinforcement numbers in graphs are very interesting research problems and were introduced by Fink et al. in [4] and Kok, Mynhardt in [14], respectively. Hattingh et al. in [8] showed that the problem of the restrained bondage is NP-complete, even for bipartite graphs. Also, he has determined the exact values of the bondage number for several classes of graphs. Moreover, the reinforcement number for digraphs has been studied by Huang, Wang and Xu in [12]. Hu and Xu in [11] showed that the problems of the bondage, the total bondage, the reinforcement and the total reinforcement numbers for an arbitrary graph are all NPhard, in general. Recently, Xu in [19] gave a review article on the bondage numbers. Moreover, Hu and Sohn in [10] showed that these problems remain NP-hard, even for bipartite graphs. Xu, Hu and Lu in [15] studied the complexity of p-reinforcement and paired bondage problems in general graphs. Jafari Rad in [16] showed that the problems of the p-reinforcement, the p-total reinforcement, the total restrained reinforcement and the k-rainbow reinforcement are all NP-hard for bipartite graphs. In addition, he also in [17] showed that the problems of the paired bondage, the total restrained bondage, the independent bondage and the k-rainbow bondage numbers are all NP-hard, even if they are restricted to bipartite graphs. From the algorithmic point of view, Hartnell et al. in [6] designed a linear time algorithm to compute the bondage number of a

The outer-connected bondage number of a graph G, where G does not have any isolated vertex, is denoted by $b_{OCD}(G)$ and is equal to the minimum number of edges whose removal from G results in a graph with an outer-connected domination number larger than the one for G. The outer-connected reinforcement number of a graph G which does not have any isolated vertices is denoted by $r_{OCD}(G)$ and is equal to the smallest number of edges whose addition from G results in a graph with an outer-connected domination number smaller than the one for G.

The rest of the paper is organized as follows: In Section 2, we describe some necessary preliminaries. In Sections 3, we show that the decision problem for the outer-connected reinforcement number in general graphs is **NP**-hard. In sections 4, we show that the outer-connected bondage number is also **NP**-hard in general graphs. In the other words, we show that there are no polynomial time algorithms to compute these values for graphs, unless P = NP. Finally, in Section 5, we determined the exact value of bondage number for several classes of graphs.

2 Preliminaries

In order to show the **NP**-hardness of the aforementioned problems, we do a polynomial time reduction from 3-satisfiability problem, 3-SAT, which is known to be an **NP**-complete problem [5]. For concreteness, Let \mathcal{U} be a set of Boolean variables. A truth assignment for \mathcal{U} is a mapping $f: \mathcal{U} \to \{T, F\}$. If f(u) = T, then u is said to be "true" with respect to f. In the case that f(u) = F, then u is said to be "false" with respect to f. If f(u) = T is a variable in f(u) = T, then f(u) = T is true if and only if the variable f(u) = T is true with respect to f(u) = T and f(u) = T is true if and only if the variable f(u) = T is true with respect to f(u) = T.

A clause over \mathcal{U} is a set of literals over \mathcal{U} which represents the disjunction of these literals. It is said to be satisfied by a truth assignment if and only if at least one of its members is true with respect to that assignment. Similarly, a collection $\mathcal{C} = \{C_1, C_2, \dots, C_m\}$ of clauses over \mathcal{U} is satisfiable if and only if there exists some truth assignment for \mathcal{U} , which simultaneously satisfies all the clauses C_i in \mathcal{C} for $i = 1, 2, \dots, m$. Such a truth assignment is called a satisfying truth assignment for \mathcal{C} . Given these notations, the 3-SAT problem is specified as follows:

3-SAT problem:




Figure 1: An instance of the outer-connected reinforcement problem. Bold points are the dominator vertices, k = 1 and $\tilde{\gamma}_c = 6$.

Instance: A collection $C = \{C_1, C_2, \dots, C_m\}$ of clauses over a finite set of variables \mathcal{U} such that $|C_j| = 3$ for $j = 1, 2, \dots, m$.

Question: Is there a truth assignment for \mathcal{U} which satisfies all the clauses in \mathcal{C} ?

Theorem 2.1. (See Theorem 3.1 in [2].) The 3-SAT problem is NP-complete.

3 NP-hardness for the outer-connected reinforcement problem

In this section, we show that the outer-connected reinforcement problem for general graphs, is an NP-hard problem. The outer-connected reinforcement problem is defined as follows:

Outer-connected reinforcement problem:

Instance: A graph G with no isolated vertices and a positive integer k.

Question: Is $r_{OCD}(G) \leq k$?

Theorem 3.1. The outer-connected reinforcement problem is NP-hard.

Proof. We show the NP-hardness of the outer-connected reinforcement problem by a polynomial reduction from the 3-SAT.

Let $I = \{U = \{u_1, u_2, \dots, u_n\}$, $C = \{C_1, C_2, \dots, C_m\}$ be an arbitrary instance of the 3-SAT problem. Without loss of generality, consider that k = 1. We construct a graph G such that this instance of 3-SAT will be satisfiable if and only if G has an outer-connected reinforcement of cardinality equal to 1, i.e. $r_{OCD}(G) = 1$. Next, we describe the construction of G.

To each $u_i \in \mathcal{U}$, we associate a triangle $\mathcal{S}_i = \{u_i, v_i, \bar{u}_i\}$. Note that $v_i \notin \mathcal{U}$. For each clause $C_j \in \mathcal{C}$, we associate a single vertex c_j and add edges $\{c_j, u_i\}(\{c_j, \bar{u}_i\})$ if the literal $u_i(\bar{u}_i)$ appears in clause C_j , for $j = 1, 2, \dots, m$, respectively. Finally, we add vertices x and y and join them to every vertex c_j for $j = 1, 2, \dots, m$ and add edges $\{x, y\}$, $\{y, u_i\}$ and $\{y, \bar{u}_i\}$ for $i = 1, 2, \dots, n$.

For example, consider a 3-SAT instance $(\mathcal{U} = \{u_1, u_2, u_3, u_4, u_5\}$, $\mathcal{C} = \{C_1, C_2, C_3, C_4\}$, where $C_1 = \{\bar{u_1}, \bar{u_2}, u_3\}$, $C_2 = \{u_1, u_3, u_5\}$, $C_3 = \{\bar{u_3}, \bar{u_4}, u_5\}$ and $C_4 = \{\bar{u_1}, \bar{u_3}, u_4\}$. Figure 1 illustrates the constructed graph corresponding to this instance.

It can be easily seen that the construction can be accomplished in polynomial time, since the graph G contains 3n + m + 2 vertices and 5n + 5m + 1 edges. All that remains to be shown is that the $I = (\mathcal{U}, \mathcal{C})$ is satisfiable if and only if $r_{OCD}(G) = 1$. To this end, we will first show the following three claims.

Claim 3.1.1. For any graph G constructed as is described above, we have $\tilde{\gamma}_c(G) = n + 1$.

Proof. Let \tilde{D} be a $\tilde{\gamma}$ -set of G. Then $\tilde{\gamma}_c(G) = |\tilde{D}| \geq n+1$ since it is necessary that $|\tilde{D} \cap V(S_i)| \geq 1$ for $i = 1, 2, \ldots, n$ and also $|\tilde{D} \cap N[x]| \geq 1$. On the other hand, the set $\tilde{D}' = \{x, u_1, u_2, \ldots, u_n\}$ is an outer-connected dominating set for G, which implies that $\tilde{\gamma}_c(G) \leq |\tilde{D}'| = n+1$. Thus, we obtain $\tilde{\gamma}_c(G) = n+1$.

Claim 3.1.2. Let \tilde{D}_e denotes a $\tilde{\gamma}$ -set of G+e for an arbitrary edge $e \in E(\bar{G})$. If there exists an edge $e \in E(\bar{G})$ such that $\tilde{\gamma}_c(G+e) = n$, then for i = 1, 2, ..., n, we have $|\tilde{D}_e \cap V(S_i)| = 1$ such that $c_j \notin \tilde{D}_e$ for j = 1, 2, ..., m and $y \notin \tilde{D}_e$.

Proof. Since the connection between the vertices in $V \setminus \tilde{D}_e$ is due to vertex y, then $y \notin \tilde{D}_e$. On the contrary, suppose that $|\tilde{D}_e \cap V(\mathcal{S}_\ell)| = 0$ for some $\ell = 1, 2, ..., n$. Since v_ℓ needs to be dominated by vertices in \tilde{D}_e and $v_\ell, u_\ell, \bar{u}_\ell \notin \tilde{D}_e$, then one of the end-vertices of the edge e should be v_ℓ , otherwise \tilde{D}_e dominates it via the edge e in G + e and for every $i \neq \ell$, we have $|\tilde{D}_e \cap V(\mathcal{S}_i)| \geq 1$, since \tilde{D}_e dominates all the vertices v_i .

It is clear that the vertices u_{ℓ} and \bar{u}_{ℓ} do not simultaneously appear in the same clause in \mathcal{C} , so, there is no j such that the vertex c_j is adjacent to both of them. Since u_{ℓ} and \bar{u}_{ℓ} should be dominated by \tilde{D}_e , then there exists two distinct vertices $c_j, c_{\ell} \in \tilde{D}_e$ such that c_j and c_{ℓ} dominate u_{ℓ} and \bar{u}_{ℓ} , respectively.

Hence, $|\tilde{D}_e| \geq n+1$, which is a contradiction. Therefore, we have $|\tilde{D}_e \cap V(S_i)| = 1$ for all $i = 1, 2, \ldots, n$ and $c_i \notin \tilde{D}_e$ for every j, since $|\tilde{D}_e| = n$.

Claim 3.1.3. The 3-SAT instance $(\mathcal{U}, \mathcal{C})$ is satisfiable if and only if $r_{OCD}(G) = 1$.

Proof. Suppose that $r_{OCD}=1$, which means that there exists an edge e in \bar{G} such that $\tilde{\gamma}_c(G+e)=n$. Let \tilde{D}_e be a $\tilde{\gamma}$ -set of G+e. Then, by Claim 3.1.2, for all $i=1,2,\ldots,n$, we have $|\tilde{D}_e\cap V(\mathcal{S}_i)|=1$. To be precise, we have either $\tilde{D}_e\cap V(\mathcal{S}_i)=\{v_i\}$, $\tilde{D}_e\cap V(\mathcal{S}_i)=\{u_i\}$ or $\tilde{D}_e\cap V(\mathcal{S}_i)=\{\bar{u}_i\}$ for all $i=1,2,\ldots,n$.

Assume that the mapping $f: \mathcal{U} \to \{T, F\}$ is defined as

$$f(u_i) = \begin{cases} T, & \text{if } u_i \in \tilde{D}_e \text{ or } v \in \tilde{D}_e, \\ F, & \text{if } \bar{u}_i \in \tilde{D}_e. \end{cases}$$
 (1)

We want to show that the mapping f is a satisfying truth assignment for $I = (\mathcal{U}, \mathcal{C})$. So, it is sufficient to show that f satisfies every clause in \mathcal{C} . We choose an arbitrary clause $C_j \in \mathcal{C}$. Since the corresponding vertex c_j to clause C_j is not adjacent to any vertices in correspondence with the set $\{v_i : 1 \leq i \leq n\}$, there exists an index i such that c_j is dominated by $u_i \in \tilde{D}_e$ or $\bar{u}_i \in \tilde{D}_e$. Assume that c_j is dominated by $u_i \in \tilde{D}_e$, then u_i is adjacent to vertex c_j in G, namely, the literal u_i is in the clause C_j . Since $u_i \in \tilde{D}_e$, we have $f(u_i) = T$ by Equation 1. So, f satisfies the clause C_j .

Now, suppose that the vertex c_j is dominated by vertex $\bar{u}_i \in \tilde{D}_e$. So, \bar{u}_i is adjacent to c_j in G, namely, the literal \bar{u}_i is in the clause C_j . Since $\bar{u}_i \in \tilde{D}_e$, we have $f(u_i) = F$ by Equation 1, which implies that \bar{u}_i is assigned the truth value T by f. So, the clause C_j is satisfied by f. Since clause C_j is chosen arbitrarily, all the clauses in C are satisfied by f, which implies that I = (U, C) is satisfiable.

Conversly, suppose that $f: \mathcal{U} \to \{T, F\}$ is a satisfying truth assignment for \mathcal{C} and \tilde{D}' is a subset of V(G) that is constructed as follows.



Figure 2: The graphs H_i and S

If $f(u_i) = T$, then we put the vertex u_i in \tilde{D}' and if $f(u_i) = F$, we put the vertex \bar{u}_i in \tilde{D}' . Therefore, we have $|\tilde{D}'| = n$. For j = 1, 2, ..., m, at least one of the literals in clause C_j is true under the assignment of f, given that f is a satisfying truth assignment for $I = (\mathcal{U}, \mathcal{C})$. So, by the construction of G, the corresponding vertex c_j in G is adjacent to at least one vertex in \tilde{D}' . Without loss of generality, let $f(u_1) = T$. Then, \tilde{D}' is a dominating set for $G + \{x, u_1\}$. On the other hand, the induced graph $G[V \setminus \tilde{D}']$ is connected. Hence, \tilde{D}' is an outer-connected dominating set for $G + \{x, u_1\}$ and $\tilde{\gamma}_c(G + \{x, u_1\}) \leq |\tilde{D}'| = n$.

By Claim 3.1.1, we have $\tilde{\gamma}_c(G) = n+1$. Therefore, we obtain $\tilde{\gamma}_c(G + \{x, u_1\}) \leq n < n+1 = \tilde{\gamma}_c(G)$, which means that $r_{OCD} = 1$.

Claims 3.1.1, 3.1.2 and 3.1.3 conclude the proof.

4 The NP-hardness of the outer-connected bondage

In this section, we show that the outer-connected bondage problem for general graphs is an \mathbf{NP} -hard problem. Consider the following decision problem.

Outer-connected bondage problem:

Instance: A graph G with no isolated vertices and a positive integer k.

Question: Is $b_{OCD}(G) \leq k$?

Theorem 4.1. The outer-connected bondage problem is NP-hard.

Proof. Let $I = \{u_1, u_2, \dots, u_n\}$, $C = \{C_1, C_2, \dots, C_m\}$ be an arbitrary instance of the 3-SAT problem. For an arbitrary positive integer k, we will construct a graph G such that this instance of 3-SAT will be satisfiable if and only if G has an outer-connected bondage of cardinality of at most k, i.e. $b_{OCD}(G) \leq k$. The graph G is constructed as follows.

To each $u_i \in \mathcal{U}$, we associate a vertex set $\mathcal{H}_i = \{u_i, v_i, \bar{u}_i, x_i, y_i\}$ and add edges $\{x_i, u_i\}, \{y_i, \bar{u}_i\}, \{u_i, v_i\}$, and $\{\bar{u}_i, v_i\}$ for i = 1, 2, ..., n. For each clause $C_j \in \mathcal{C}$, we associate a single vertex c_j and add edge $\{c_j, u_i\}(\{c_j, \bar{u}_i\})$ if the literal $u_i(\bar{u}_i)$ is present in the clause C_j , where j = 1, 2, ..., m. Then, we add a set of vertices $S = \{s_1, s_2, s_3, s_4\}$ and join s_1, s_3 and s_4 to vertices c_j and s_2 . Finally, we add a vertex t to the graph G, edges $\{t, s_1\}, \{t, s_3\}, \{t, s_4\}, \{t, u_i\}$ and $\{t, \bar{u}_i\}$ for i = 1, 2, ..., n and set k = 1.

It can be easily seen that the construction can be accomplished in polynomial time, since the graph G contains 5n + m + 5 vertices and 6m + 6n + 6 edges. All that remains to be shown is that $I = (\mathcal{U}, \mathcal{C})$ is satisfiable if and only if $b_{OCD}(G) \leq k$. Without loss of generality, let k = 1. To this end, we will first prove the following five claims.

Claim 4.1.1. For any graph G constructed as above, we have $\tilde{\gamma}_c(G) \geq 3n + 1$.

Proof. Let \tilde{D} be a $\tilde{\gamma}$ -set of G. Then, $\tilde{\gamma}_c(G) = |\tilde{D}| \geq 3n+1$, since $|\tilde{D} \cap V(\mathcal{H}_i)| \geq 3$ for $i = 1, 2, \ldots, n$. Note that to dominate a vertex v_i , we need at least one vertex and the leaf vertices x_i and y_i to be in \tilde{D} . Moreover, $|\tilde{D} \cap N[s_2]| \geq 1$.

Claim 4.1.2. If $\tilde{\gamma}_c(G) = 3n + 1$, then $c_j, t \notin \tilde{D}$ for j = 1, 2, ..., m, $\tilde{D} \cap V(S) = \{s_2\}$ and $|\tilde{D} \cap V(\mathcal{H}_i)| = 3$ for i = 1, 2, ..., n.

Proof. Since the connection between \mathcal{H}_i and S is due to the vertex t, then $t \notin \tilde{D}$. Suppose that $\tilde{\gamma}_c(G) = 3n+1$. Then, $|\tilde{D} \cap V(\mathcal{H}_i)| = 3$ for $i=1,2,\ldots,n$, while $|\tilde{D} \cap V(S)| = 1$. Consequently, $c_j \notin \tilde{D}$ for $j=1,2,\ldots,m$. Simultaneously, if $\tilde{D} \cap V(S) = \{s_1\}$, then, s_3 and s_4 are not dominated. Hence, $s_1 \notin \tilde{D}$ and similarly, $s_3, s_4 \notin \tilde{D}$. So, $\tilde{D} \cap V(S) = \{s_2\}$.

Claim 4.1.3. The 3-SAT instance $I = (\mathcal{U}, \mathcal{C})$ is satisfiable if and only if $\tilde{\gamma}_c(G) = 3n + 1$.

Proof. Suppose that $\tilde{\gamma}_c(G) = 3n + 1$ and c_j is an arbitrary vertex. By Claim 4.1.2, this vertex is adjacent to either $u_i \in \tilde{D}$ or $\bar{u}_i \in \tilde{D}$, since $s_1, s_3, s_4 \notin \tilde{D}$. As $|\tilde{D} \cap V(\mathcal{H}_i)| = 3$ for i = 1, 2, ..., n, it follows that either $\tilde{D} \cap V(\mathcal{H}_i) = \{x_i, y_i, u_i\}$, $\tilde{D} \cap V(\mathcal{H}_i) = \{x_i, y_i, \bar{u}_i\}$ or $\tilde{D} \cap V(\mathcal{H}_i) = \{x_i, y_i, v_i\}$. Let the mapping $f : \mathcal{U} \to \{T, F\}$ be defined as

$$f(u_i) = \begin{cases} T, & \text{if } u_i \in \tilde{D} \text{ or } v_i \in \tilde{D}, \\ F, & \text{if } \bar{u_i} \in \tilde{D}. \end{cases}$$
 (2)

To prove that the values assigned by the mapping f is a satisfying truth assignment for $I=(\mathcal{U},\mathcal{C})$, it is sufficient to show that f satisfies every clause in \mathcal{C} . Let $C_j \in \mathcal{C}$ be an arbitrarily clause. Since the corresponding vertex to the clause C_j is not adjacent to any vertex in correspondence with the set $\{v_i, x_i, y_i : 1 \le i \le n\}$, there exists an i such that c_j is dominated by either $u_i \in \tilde{D}$ or $\bar{u}_i \in \tilde{D}$. Without loss of generality, assume that c_j is dominated by $u_i \in \tilde{D}$. So, u_i is adjacent to c_j in G, namely the literal u_i is in the clause C_j . Since $u_i \in \tilde{D}$, we have $f(u_i) = T$ by Equation 2. So, the values assigned by the mapping f satisfies the clause C_j .

Now, suppose that the vertex c_j is dominated by vertex $\bar{u}_i \in \tilde{D}$. So, the vertex \bar{u}_i is adjacent to the vertex c_j in G, namely the literal \bar{u}_i is in the clause C_j . Since $\bar{u}_i \in \tilde{D}$, we have $f(u_i) = F$ by Equation 2, which implies that \bar{u}_i is assigned the truth value T by f and the clause C_j is satisfied by f. Since C_j was chosen arbitrarily, all the clauses in C are satisfied by f, which implies that $I = (\mathcal{U}, C)$ is satisfiable.

Conversly, suppose that $f: \mathcal{U} \to \{T, F\}$ is a satisfying truth assignment for \mathcal{C} and \tilde{D}' is a subset of V(G) which is constructed as follows. If $f(u_i) = T$, then we put the vertex u_i in \tilde{D}' and if $f(u_i) = F$, then we put the vertex \bar{u}_i in \tilde{D}' . Therefore, we have $|\tilde{D}'| = n$. For $j = 1, 2, \ldots, m$, at least one of the literals in C_j is true under the assignment f, because the mapping f is a satisfying truth assignment for $I = (\mathcal{U}, \mathcal{C})$. So, by the construction of G, the vertex is in correspondence to C_j in G is adjacent to at least one vertex in \tilde{D}' . Then, $D = \tilde{D}' \cup (\bigcup_{i=1}^n \{x_i, y_i\}) \cup \{s_2\}$ is a dominating set for G. On the other hand, the induced graph $G[V \setminus D]$ is connected. Hence, D is an outer-connected dominating set for G and $\tilde{\gamma}_c(G) \leq |D| = 3n + 1$. By Claim 4.1.1, we have $\tilde{\gamma}_c(G) \geq 3n + 1$. Therefore, we obtain $\tilde{\gamma}_c(G) = 3n + 1$.

Claim 4.1.4. For every $e \in E(G)$, we have $\tilde{\gamma}_c(G-e) \leq 3n+2$.

Proof. Suppose that $E' = \{\{s_2, s_3\}, \{s_2, s_4\}, \{s_1, c_j\}, \{u_i, v_i\}, \{y_i, \bar{u}_i\}, \{t, s_1\}, \{v_i, \bar{u}_i\}, \{t, \bar{u}_i\}\}$ and $E'' = E \setminus E'$. Let $e \in E''$ be an edge. It is clear that the set $D' = (\bigcup_{i=1}^n \{x_i, y_i, u_i\}) \cup \{s_1, s_2\}$ is an outer-connected dominating set for G - e, since every vertex in $V \setminus D'$ is adjacent to a vertex

in D' due to an edge in E', and the induced graph $(G-e)[V \setminus D']$ is connected. This connection is established by vertices t and s_i for $i \neq 1, 2$. Given that |D'| = 3n + 2, then $\tilde{\gamma}_c(G-e) \leq 3n + 2$. We have four cases to consider:

Case 1: If either $e = \{s_2, s_3\}$, $e = \{s_1, c_j\}$ or $e = \{t, s_1\}$, then $D' = (\bigcup_{i=1}^n \{x_i, y_i, u_i\}) \cup \{s_3, s_2\}$ is an outer-connected dominating set for G - e and $\tilde{\gamma}_c(G - e) \leq |D'| = 3n + 2$.

Case 2: If $e = \{s_2, s_4\}$, then $D' = (\bigcup_{i=1}^n \{x_i, y_i, u_i\}) \cup \{s_4, s_2\}$ is an outer-connected dominating set for G - e and $\tilde{\gamma}_c(G - e) \leq |D'| = 3n + 2$.

Case 3: If either $e = \{y_i, \bar{u}_i\}$, $e = \{\bar{u}_i, v_i\}$ or $e = \{u_i, v_i\}$, then $D' = (\bigcup_{i=1}^n \{x_i, y_i, v_i\}) \cup \{s_1, s_2\}$ is an outer-connected dominating set for G - e and $\tilde{\gamma}_c(G - e) \leq |D'| = 3n + 2$.

Case 4: If $e = \{t, \bar{u}_i\}$, then $D' = (\bigcup_{i=1}^n \{x_i, y_i, \bar{u}_i\}) \cup \{s_1, s_2\}$ is an outer-connected dominating set for G - e and $\tilde{\gamma}_c(G - e) \leq |D'| = 3n + 2$.

Claim 4.1.5. $\tilde{\gamma}_c(G) = 3n + 1$ if and only if $b_{OCD}(G) = 1$.

Proof. First, suppose that $\tilde{\gamma}_c(G) = 3n + 1$. Let $e = \{s_1, s_2\}$ and $\tilde{\gamma}_c(G) = \tilde{\gamma}_c(G - e)$. If \tilde{D} is a $\tilde{\gamma}$ -set of G - e, then \tilde{D} is a $\tilde{\gamma}$ -set for G of cardinality 3n + 1. By Claim 4.1.2, we have $c_j, t \notin \tilde{D}$ for $j = 1, 2, \ldots, m$ and $\tilde{D} \cap V(S) = \{s_2\}$. So, the vertex s_1 is not dominated by \tilde{D} , which is a contradiction. Hence, $\tilde{\gamma}_c(G) < \tilde{\gamma}_c(G - e)$. So, $b_{OCD}(G) = 1$.

Next, assume that $b_{OCD}(G) = 1$. By Claim 4.1.1, it follows that $\tilde{\gamma}_c(G) \geq 3n+1$. Suppose that e is an edge such that $\tilde{\gamma}_c(G) < \tilde{\gamma}_c(G-e)$. By Claim 4.1.4, we have $3n+1 \leq \tilde{\gamma}_c(G) < \tilde{\gamma}_c(G-e) \leq 3n+2$ which implies that $\tilde{\gamma}_c(G) = 3n+1$.

Therefore, by Claims 4.1.3 and 4.1.5, we have $b_{OCD}(G) = 1$ if and only if $I = (\mathcal{U}, \mathcal{C})$ is satisfiable.

5 Exact values for $b_{OCD}(G)$

In this section, we establish several theorems on the exact values of $b_{OCD}(G)$.

Lemma 5.1. Let $G = K_n$ be a complete graph with $n \ge 3$ vertices. If $\lceil \frac{n+1}{2} \rceil - 1$ edges are removed from G, then G is still connected.

Proof. Two cases are considered:

1. If n is odd, then $\lceil \frac{n+1}{2} \rceil = \frac{n+1}{2}$ and the number of remaining edges by removing $\lceil \frac{n+1}{2} \rceil - 1$ edges from G is equal to

$$\frac{n(n-1)}{2} - (\frac{n+1}{2} - 1) = \frac{(n-1)^2}{2}.$$
 (3)

On the other hand, a graph of order n cannot be disconnected if it is simple and the number of its edges is greater than $\frac{(n-1)(n-2)}{2}$. This is because the complete graph on n-1 vertices has $\frac{(n-1)(n-2)}{2}$ edges. Consequently, since $\frac{(n-1)^2}{2} > \frac{(n-1)(n-2)}{2}$ and $n \geq 3$, the graph G remains connected even after the removal of $\lceil \frac{n+1}{2} \rceil - 1$ edges.

2. If n is even, then $\lceil \frac{n+1}{2} \rceil = \frac{n+2}{2}$ and the number of remaining edges by removing $\lceil \frac{n+1}{2} \rceil - 1$ edges from G is equal

$$\frac{n(n-1)}{2} - (\frac{n+2}{2} - 1) = \frac{n(n-2)}{2}.$$
 (4)

On the other hand, a graph of order n cannot be disconnected if it is simple and the number of its edges is greater than $\frac{(n-1)(n-2)}{2}$. This is because the complete graph with n-1 vertices has $\frac{(n-1)(n-2)}{2}$ edges. Consequently, since $\frac{n(n-2)}{2} > \frac{(n-1)(n-2)}{2}$ and $n \geq 3$, the graph G remains connected even after the removal of $\lceil \frac{n+1}{2} \rceil - 1$ edges.

Theorem 5.2. Let K_n be a complete graph with $n \geq 3$ vertices. Then, we have

$$b_{OCD}(K_n) = \begin{cases} 1, & \text{if } n = 3, \\ \lceil \frac{n}{2} \rceil, & \text{otherwise.} \end{cases}$$
 (5)

Proof. If n = 3, then we have $\tilde{\gamma}_c(K_3) = 1$. By removing any edges from G, it turns into a P_3 . So, we have $b_{OCD}(K_3) = 1$ since $\tilde{\gamma}_c(P_3) = 2$.

Now, suppose that n>3. Let the graph G' be obtained by removing fewer than $\lceil \frac{n}{2} \rceil$ edges from G. Then, G' contains at least a vertex of degree n-1. Let the vertex v be of degree n-1. In the other hand, according to the Lemma 5.1 we have the induced graph $G'[V\setminus v]$ is connected. Then, $\tilde{\gamma}_c(G')=1$. So, we have

$$b_{OCD}(G) \ge \lceil \frac{n}{2} \rceil. \tag{6}$$

Now, we need to consider the following two cases:

- 1. If n is even. Let H be the graph obtained by removing $\lceil \frac{n}{2} \rceil$ independent edges from G. Then the degree of every vertex $v \in V(H)$ is n-2. So, we have $\tilde{\gamma}_c(H) \geq 2$.
- 2. If n is odd. Let H be the graph obtained by removing $\frac{n-1}{2}$ independent edges from G. Then, there is exactly one vertex $v \in V(H)$ such that the degree of v is n-1. If we remove one edge incident with v, then we have $\tilde{\gamma}_c(H) \geq 2$.

In either cases, by removing $\lceil \frac{n}{2} \rceil$ we have $\tilde{\gamma}_c(G) < \tilde{\gamma}_c(H)$. So,

$$b_{OCD}(G) \le \lceil \frac{n}{2} \rceil. \tag{7}$$

Therefore, by Equations 6 and 7, we have $b_{OCD}(G) = \lceil \frac{n}{2} \rceil$.

Theorem 5.3. Let C_n be a cycle graph with $n \geq 3$ vertices. Then, we have

$$b_{OCD}(C_n) = \begin{cases} 1, & if \quad n = 3, \\ \lceil \frac{n}{3} \rceil, & otherwise. \end{cases}$$
 (8)

Proof. If n=3, then we have $\tilde{\gamma}_c(C_3)=1$. By removing any edges from G, it turns into a P_3 . So, $b_{OCD}(C_3)=1$ since $\tilde{\gamma}_c(P_3)=2$.

Cyman in [3] has shown that if $n \geq 4$, then $\tilde{\gamma}_c(C_n) = \tilde{\gamma}_c(P_n) = n-2$. So, for $n \geq 4$, we have $b_{OCD}(C_3) \geq 1$. By removing some of the edges from P_n , a set of components $\{H_1, H_2, \cdots, H_m\}$ is obtained such that every component is a path. If $H = \bigcup_{i=1}^m H_i$ and $\tilde{D}(H)$ is an outer-connected dominating set for H, then $\tilde{D}(H) = \bigcup_{i=1}^{m-1} H_i \cup \tilde{D}(H_m)$ (See Lemma 3.1 in [7]). Therefore, if there exists at least one component such as H_i with four vertices, then $\tilde{\gamma}_c(C_n) = \tilde{\gamma}_c(P_n) = \tilde{\gamma}_c(H) = n-2$. Otherwise, we have $\tilde{\gamma}_c(C_n) = \tilde{\gamma}_c(P_n) < \tilde{\gamma}_c(H) = n-1$. So, we need to break the path P_n in such a way that there exist no more than three vertices in any component H_i . To this end, we have to remove $\lceil \frac{n}{3} \rceil - 1$ edges from P_n , which with the deleted edge from C_n are counted to $\lceil \frac{n}{3} \rceil$.

Theorem 5.4. Let P_n be a path with $n \geq 3$ vertices. Then, we have

$$b_{OCD}(P_n) = \begin{cases} 1, & \text{if } n = 2, \\ 2, & \text{if } n = 3, \\ \lceil \frac{n}{3} \rceil - 1, & \text{otherwise.} \end{cases}$$
 (9)

Proof. The cases for n=2 and n=3 are quite clear. The proof of the case $n \ge 4$ is the same as in Theorem 5.3.

Theorem 5.5. The graph G is a galaxy of order $n \ge 4$ if and only if $b_{OCD}(G) = |E(G)|$.

Proof. According to the Observation 2 in [3] and Lemma 3.1 in [7], it is clear that if G is a galexy, then $\tilde{\gamma}_c(G) = n-1$. Moreover, the only graph with outer-connected domination number equal to n is \bar{K}_n . So, $b_{OCD}(G) = |E(G)|$. Converesly, suppose that $b_{OCD}(G) = |E(G)|$. If a component H of G is not a star, then H either contains a cycle or a P_4 , which means that $\tilde{\gamma}_c(G) \leq n-2$. Let $e = \{v_1, v_2\}$ be an edge in the cycle or the P_4 . If H is a graph obtained by removing all edges from G expect e, then we have $\tilde{\gamma}_c(H) = n-1 > \tilde{\gamma}_c(G)$. This implies $b_{OCD}(G) \leq |E(G)| - 1$, which is a contradiction.

References

- [1] AKHBARI, M., HASNI, R., FAVARON, O., KARAMI, H., AND SHEIKHOLESLAMI, S. M. On the outer-connected domination in graphs. *Journal of Combinatorial Optimization* (2013), 1–9.
- [2] COOK, S. The P versus NP problem. The millennium prize problems (2000), 87–104.
- [3] CYMAN, J. The outer-connected domination number of a graph. Australasian Journal of Combinatorics 38 (2007), 35–46.
- [4] Fink, J. F., Jacobson, M. S., Kinch, L. F., and Roberts, J. The bondage number of a graph. Discrete Mathematics 86, 1-3 (1990), 47–57.
- [5] GAREY, M. R., AND JOHNSON, D. S. Computers and intractability, vol. 29. wh freeman New York, 2002.
- [6] HARTNELL, B., JØRGENSEN, L. K., VESTERGAARD, P. D., AND WHITEHEAD, C. Edge stability of the k-domination number of trees. Bulletin of the Institute of Combinatorics and its Applications 22 (1998), 31–40.
- [7] Hashemipour, M., Hooshmandasl, M., and Shakiba, A. On outer-connected domination for graph products. arXiv preprint arXiv:1708.00188 (2017).
- [8] HATTINGH, J. H., AND PLUMMER, A. R. Restrained bondage in graphs. *Discrete Mathematics* 308, 23 (2008), 5446–5453.
- [9] HAYNES, T. W., HEDETNIEMI, S., AND SLATER, P. Fundamentals of domination in graphs. CRC Press, 1998.
- [10] Hu, F.-T., and Sohn, M. Y. The algorithmic complexity of bondage and reinforcement problems in bipartite graphs. *Theoretical Computer Science* 535 (2014), 46–53.
- [11] Hu, F.-T., AND Xu, J.-M. On the complexity of the bondage and reinforcement problems. *Journal of Complexity* 28, 2 (2012), 192–201.
- [12] HUANG, J., WANG, J.-W., AND XU, J.-M. Reinforcement numbers of digraphs. Discrete Applied Mathematics 157, 8 (2009), 1938–1946.

- [13] Keil, J. M., and Pradhan, D. Computing a minimum outer-connected dominating set for the class of chordal graphs. *Information Processing Letters* 113, 14 (2013), 552–561.
- [14] Kok, J., and Mynhardt, C. Reinforcement in graphs. Congr. Numer 79 (1990), 225–231.
- [15] Lu, Y., Hu, F.-T., AND Xu, J.-M. On the p-reinforcement and the complexity. *Journal of Combinatorial Optimization* 29, 2 (2015), 389–405.
- [16] RAD, N. J. On the complexity of reinforcement in graphs. Discussiones Mathematicae Graph Theory 36, 4 (2016), 877–887.
- [17] RAD, N. J., AND KAMARULHAILI, H. On the complexity of some bondage problems in graphs. australasian journal of combinatorics 68, 2 (2017), 265–275.
- [18] TERESA, W. H., STEPHEN, T. H., AND PETER, J. S. Domination in graph, advanced topics. *Marcel Dekkar, New York* (1998).
- [19] Xu, J.-M. On bondage numbers of graphs: a survey with some comments. *International Journal of Combinatorics* 2013 (2013).