

ORIGINAL ARTICLE

Hybrid Sterility Can Only be Primary When Acting as a Reproductive Barrier for Sympatric Speciation

Donald R. Forsdyke¹

Department of Biomedical and Molecular Sciences, Queen's University, Kingston, ON, Canada

Running title: The Primary Role of Hybrid Sterility

¹E-mail: forsdyke@queensu.ca

Abstract Parental gametes unite to form a zygote that develops into an adult with gonads that, in turn, produce gametes. Interruption of this germinal cycle by prezygotic or postzygotic reproductive barriers can result in two independent cycles, each with the potential to evolve into a new species. When the speciation process is complete, members of each species are fully reproductively isolated from those of the other. During speciation a primary barrier may be supported and eventually superceded by a later appearing secondary barrier. For those holding certain cases of prezygotic isolation to be primary (e.g. elephant cannot copulate with mouse), the onus is to show that they had not been preceded over evolutionary time by periods of postzygotic hybrid inviability (genically determined) or sterility (genically or chromosomally determined). Likewise, the onus is upon those holding cases of hybrid inviability to be primary (e.g. Dobzhansky-Muller epistatic incompatibilities), to show that they had not been preceded by periods, however brief, of hybrid sterility. The latter, when acting as a sympatric barrier that initiates reproductive isolation, can only be primary. In many cases, hybrid sterility results from incompatibilities between parental chromosomes that attempt to pair during meiosis in the gonad of their offspring (Winge-Crowther-Bateson incompatibilities).

Keywords Chromosomal speciation . Germinal cycle . Haldane's rule . Heirarchical barriers . Meiosis . Reproductive isolation

Introduction

Successful sexual crossing results in offspring that are fertile and so able to continue the line. Barriers that impede crossing of individuals within a species can facilitate the branching of that species into two species. This review considers three broad reproductive barriers in the context of successive generational life cycles, with an emphasis on the mechanism and importance of the hybrid sterility barrier which, if operative in sympatry (Foote 2018), can only be primary. When another barrier is proposed to be primary, there is an onus on the proposer to show that that barrier had not been preceded by a period, however brief, of some degree of hybrid sterility (Forsdyke 2017a, b, c; 2018).

The Advantage of Sterility

Since he construed natural selection as working positively when organisms survive and produce offspring, Charles Darwin found it difficult to understand how something so negative as sterility in an otherwise normal offspring (hybrid) could be favoured by natural selection (Darwin 1862). Today, the paradox of an “inherent difficulty in genetically dissecting the phenotype that prevents its own transfer to progeny,” is acknowledged (Gregorova et al. 2017). However, hybrid sterility in offspring can be dissected when it is regarded as a phenotypic expression, not of an offspring character, but of a character of the parents of that offspring.

While the sterility character is disadvantageous *for offspring* – for whom it would be a barrier preventing continuation of the line – sterile offspring can be the manifestation of an emergent *parental character* that, indeed, could be favoured by natural selection. The phenotype manifested in their hybrid would involve a generational cycle immediately following that of the parental generation. As with other characters that natural selection might target, first there would have to have been internal genomic variations within parental or ancestral generations. When the character emerged, natural selection would act, either positively if the character was advantageous, or negatively if the character was disadvantageous.

While an offspring may inherit various overt characters, such as tallness or hair colour, an inherited sterility character is more subtle. When the offspring crosses with *certain members* of its species, there is a failure to produce further offspring (the potential grandchildren of its parents). However, when it crosses with *other members*, fertile offspring are produced. Likewise,

those certain members with whom it fails to cross may themselves find other members with whom they may successfully cross. Thus, an individual can cross with some members of its species, but not with others. The problem is not with the individuals, but with their pairing relationships. They are reproductively compatible with some, but reproductively incompatible with others. The subtle character they manifest is *selective reproductive isolation*. How could this be favoured by natural selection?

When there exists some degree of hybrid sterility between sectors of a species, then members of that species can progressively split into two groups. *Within* each group mutual fertility exists among its members, but *between* members of different groups fertility is decreased. As proposed long ago by Romanes (1886), isolation may often have *preceded* character differentiation, not the converse (Forsdyke 2001, pp. 47–63; Fuller et al. 2017). When there is reproductive isolation there is then the possibility of the eventual division of a species into two species that evolve different sets of characters. Members of these two species may, when viewed as a collective, meet the challenges of natural selection with a wider range of phenotypic options than the members of the one species from which they originated. In this circumstance, possession of a sterility/fertility character that facilitated speciation could be favoured by natural selection. This would be especially apparent with members of a species that had arrived at an evolutionary dead end in that they were having difficulties meeting new challenges. There might be more flexibility if there were division into two species.

Non-Mendelian Inheritance of the Residue

Aware of Romanes' work, in the first decade of the twentieth century William Bateson recognized that, unlike the transfer to offspring of more conventional characters that were distributed according to Mendel's laws, the *selective sterility* character was often distributed to *all* the offspring of a *particular* couple. Likewise, for another couple, the *selective fertility* character would be distributed to *all* offspring (Forsdyke 2010). While Bateson sometimes found that the transmission of sterility did indeed show a Mendelian distribution, thus indicating a relationship with what we now call genes (genic sterility), this only explained hybrid sterility in a minority of cases. There was something else, a factor, apparently of a non-genic nature, that could also be manifest as hybrid sterility.

Having already coined many of the terms now familiar to geneticists – e.g. homozygote, heterozygote, epistasis – he pondered a name for that factor. The philosopher William James had written of the “unclassified residuum” that floats around the “orderly facts” of a science and “proves less easy to attend to than to ignore” (Forsdyke 2016, p. 397). Bateson chose the name “residue” for the new factor. Furthermore, he held this likely to be fundamental to Darwin’s great question: how do species originate?

Winge-Crowther-Bateson Incompatibility

The chromosomal basis of the Bateson’s residue emerged in the 1920s (Cock and Forsdyke 2008). Initiation of a speciation process requires mechanisms for achieving some degree of reproductive isolation. This decreases the probability that the process would be subverted by recombination between the genomes of diverging types. Such recombination would tend to blend (homogenize), rather than retain, the differences that were responsible for the initiation of, and/or were sustaining, the speciation process (see later). Be it of genic, or non-genic, origin, hybrid sterility is one of the three broad barriers to reproduction that can arise in sympatry among certain members of a species. An outcome of the blockage can be branching into two species (see later).

With the assistance of a physician (Crowther 1922), Bateson tentatively equated non-genic hybrid sterility with an incompatibility between parental chromosomes when they pair at meiosis within the gonad of their offspring (Bateson 1922). This failed homology search would be the basis of his mysterious “residue.” Since a similar view had been advanced by the yeast geneticist, Öjvind Winge (1917), we can refer to this eponymously as a Winge-Crowther-Bateson (WCB) incompatibility. By contrast, the well known Dobzhansky-Muller (DM) incompatibility refers to a form of genic incompatibility involving adverse epistatic interactions between the products of parental genes (Forsdyke 2011).

While WCB incompatibilities may involve chromosomes on a microscopically observable scale (White 1978; King 1993; Fuller et al. 2017), there has been a growing focus on finer sequence changes (Forsdyke 1996, 2001, 2016, 2017a, b; Reese and Forsdyke 2016; see later). Furthermore, there has been growing support for the idea that meiotic “homology search and recognition can occur independently from strand invasion and genetic exchange” (Chapman et

transmission (prezygotic isolation), or development (hybrid inviability), or gamete formation (hybrid sterility). When speciation is complete (bottom), two independent germinal cycles maintain the continuity of species B and C through the generations. To simplify, male and females here share a common cycle

The Germinal Cycle

For sexual organisms that are considered high on the evolutionary scale there is a clearly demarcated reproductive (germinal) cycle. Figure 1 shows a simplified form of the reproductive cycle in a species (A), with the cycles of the two sexes merged and their bodies, having acted as hosts to their gonads, being discarded in each generation (grey boxes). The cycling process continues within the confines of species A until one of the three fundamental barriers to cycle operation arises as a primary barrier. This primary barrier may affect many crosses until secondary, and perhaps tertiary, barriers come into play. There is a progressively irreversible divergence into new lines with independent cycles that are seen today as species B and C.

When viewed abstractly, cycles operate continuously and there may be no obvious start or end-points. A blockage at any one of three main points (1, 2 or 3) either halts or slows a cycle (Fig. 2a). When halted, subsequent downstream points of potential blockage cannot contribute to cycle arrest. When slowed, these subsequent points of potential blockage may contribute to, and eventually replace, the point of initial blockage that occurred in an earlier generational cycle. They become *secondary* blockage points that, likewise, may eventually be replaced by *tertiary* blockage points in later generational cycles.

The actual reproductive cycle differs from the abstract and is shown in Fig. 2b with separate cycles for the two sexes (α and α that have closely related DNA sequences). A cycle operates in an individual of one sex and interacts, through its generation of gametes, with the reproductive cycle that operates in an individual of another sex. Both cycles are naturally interrupted at one point (1) when gamete transmission occurs, and a new cycle emerges in the form of an individual (α) with some of the characteristics of the parenting individuals (upper part of Fig. 2b). This process can recur, so establishing a line of α organisms.




Fig. 2 Roles of hierarchically related barriers in interruption of germinal cycles. Since the germinal cycle is recursive, *any point*, be it before or after union of gametes to form a zygote, can mediate primary arrest of the cycle (points 1, 2 and 3; red lines in Figure 2A). The arrow indicates clockwise progression, but numbering is anticlockwise. When 3 is primary the cycle arrests and downstream events (2 and 1) cannot occur in that individual. When 2 is primary, then downstream event (1) cannot occur in that individual. However, when 1 is primary it cannot affect 3 and 2 *in the same individual* because sexual collaboration with a partner (α with α) generates a new cycle (upper part of Fig. 2B). Thus, it is normal for the cycle to interrupt when a hybrid sterility barrier (1) is absent. If prezygotic (transmission) barriers (3) are absent, gametes from pairing partners can meet (dashed black lines) as a new individual (another α with a new cycle). If hybrid inviability barriers (2) are absent in that new individual, then gonadal gametogenesis can follow, provided hybrid sterility barriers (1) are still absent. The cycle can then resume in another α individual, and so on (not shown). However, collaboration with a partner with a slightly divergent DNA sequence (α with β^1) may generate a new cycle (α/β^1) that

arrests at the hybrid sterility barrier (middle thick red line). In this circumstance, to continue their respective lines, α/β^1 must meet another α/β^1 (not shown), and β^1 must meet another β^1 (not shown). Should there be further DNA divergence (β^1 to β^2) then a collaboration with an α (not shown) or with the few α/β^1 offspring who chance to evade their cycle's hybrid sterility barrier (shown as dashed grey line), could be stopped by the hybrid inviability barrier (2; bottom thick red line). This new cycle's hybrid sterility barrier 1 is then excluded. To continue their lines, α/β^1 must find another α/β^1 (not shown), and β^2 must find another β^2 (not shown). With further DNA divergence (β^2 to β^3 ; not shown), the transmission barrier (3) can come into play, so excluding 2 and 1

With each crossing there is some assortment of parental characters to constitute a new offspring and, while there may be no loss of character-forming *potential*, in classical Mendelian terms there is no blending of characters. This is seen most obviously in the inheritance of sex. Although her father was male, a daughter does not inherit her father's sex. Although his mother was female, a son does not inherit his mother's sex. Multiple genes are involved in sex determination, but various features of the sex chromosomes tend to impede their recombinational re-assortment, so the genes act as a unit and there is no blending.

However, when multiple genes contribute to a character (such as height in humans), then, as Mendel recognized, blending is usual (Forsdyke 2016, pp. 145-146). In such circumstances, a tendency for divergence into different forms (say tall and short in humans) will be countered by the blending that occurs (e.g. when a tall person crosses with a short person). For different lines to emerge, tall must cross with tall and short must cross with short. To achieve such *selective* crossing, some externally or internally imposed form of reproductive isolation (reproductive selection) must occur. There must be barriers.

We are here concerned with natural, internal, non-geographic, barriers to crossing that can be hierarchically related (primary, secondary, tertiary). Among these, hybrid sterility is special in that, when manifest, it is primary. Within one organism, the sterility barrier cannot be replaced by another barrier, yet, within a subsequent organism in the line, another barrier (secondary) may arise to pre-empt the emergence of hybrid sterility in that organism. Likewise, within a subsequent organism, a tertiary barrier may arise to pre-empt the re-emergence of the secondary barrier within the line (Fig. 2b lower).

Defective gamete transmission always trumps inviability, since without transmission there can be no zygote to develop into an embryo. And if there is embryo there can be no adult for

gametogenesis. Even if transmission required far fewer genes than development, over long evolutionary time scales mutations in transmission-related genes would be bound to occur. Accordingly, when genic differences underlie cycle interruption, those defects are most likely to *finalize* among transmission-related genes. As far as reproductive isolation is concerned, any pre-existing barrier-creating differences in genes acting post-zygotically then become irrelevant.

Indeed, many modern species have ended up controlled by barrier 3 where multiple genes have failed to coadapt (e.g. elephant cannot copulate with a mouse). However, in others 2 remains the identified barrier. Neither identification excludes the possibility of an earlier role for 1 as a fundamental barrier acting at the time of an initial divergence into two species. Although unlikely, even barrier 1 may sometimes have persisted (see later). When there are successive barriers, the temporal sequence may sometimes be 1, 2, 3. Once pre-empted, early acting barriers are liberated from their reproductive isolating role.

Thus, regarding the speciation process, there is a “multi-dimensional ... continuum” (Mérot et al. 2017). Although an early member of a hierarchical sequence may sometimes be forestalled (preempted) by a later, there should be evidence that the early member had not acted before concluding that it was not primary. A clear determination that a hybrid sterility barrier was absent at initiation may prove difficult since, once liberated from its reproductive isolation role, it may respond to other pressures (Forsdyke 2001, p. 22). However, new approaches to the sequencing of “natural DNA archives” (Olajos et al. 2017), may now permit a finer analysis of sequence changes during speciation.

Sex Chromosomes and Haldane’s Rule

A primary role for hybrid sterility was suggested from the generalization that, prior to the development of full sterility among offspring (F1 hybrids), the sterility may be partial in that only one sex (male in mice) is affected (Haldane 1922). Furthermore, experimental exchanges (“introgressions”) of individual chromosomes between “closely related mouse subspecies” have shown that “heterospecific autosomal pairs in sterile hybrids are more prone to asynapsis than the homospecific pairs in which both homologs came from the same species” (Bhattacharyya et al., 2013). Even among heterospecific pairs, some chromosomes appear more disposed to

asynapsis than others. Thus, Bhattacharyya et al. (2014) note that “the number of unsynapsed autosomes per cell varies, indicating the same type of cis-acting mechanism operates on individual autosomes.” Their observations are attributed, not to some mobile (“trans-acting”) genic factor with the ability to single out individual chromosome pairs, but to “their fast evolving non-genic divergence,” which could have affected some chromosome pairs more than others (Bhattacharyya et al. 2013).

Being *already* disparate, the sex chromosomes in male mice (XY) regularly fail to pair along most of their lengths. Relative to females with homologous sex chromosomes (XX), this gives males (the “heterogametic” sex) a head-start along the path to full sterility (Haldane’s rule for hybrid sterility; Forsdyke 2000). While conceding an important role of genes in this process (*prdm9* and a hemizygous gene on the male X chromosome), Bhattacharyya et al. (2013) conclude that “variation in pairing failure is under genic control,” but the sterility itself “is chromosomal, caused by heterospecific pairing incompatibilities.” They deem this supportive of similar suggestions regarding sterility in both fruit fly (Naviera and Maside 1998; Moehring, 2011), and yeast (“simple sequence divergence acted upon by the mismatch repair system;” Louis 2009).

Whether sex chromosomes are the same size (homomorphic) or of different size (heteromorphic), these considerations also apply to plants with independent sexes (Ironsides and Filatov 2005). Indeed, Delph and Demuth (2016), claiming to have “the best explanation for male rarity in some *Silene* hybrids,” consider that “although the original chromosomal mechanism ... largely fell out of favor, recent work has argued for its importance on theoretical grounds.” Furthermore, demonstrating the progressive involvement of autosomes *after* a primary involvement of sex chromosomes, Hu and Filatov (2016) find an “increased species divergence and reduced gene flow on the *Silene* X-chromosome,” but gene flow involving autosomal loci is still “sufficient to homogenize the gene pools of the two species.”

The Pairing Mechanism

Disparities in the DNA sequences that parents have contributed to their offspring can suffice to impair the chromosome pairing needed for error-correction (Bernstein et al. 2017). When the

disparity within their offspring's gonad is not correctable, then only within the bounds of an emerging new species will each parent be likely to find a non-disparate partner (Forsdyke 2018).

For successful homologous recombination two DNA duplexes must pair and exchange segments. This requires both recognition of some degree of sequence similarity and strand breakage. The temporal order of these events is contentious. A popular model postulates cutting to produce a single strand that *then* seeks a pairing partner in the homologous duplex (Szostak et al. 1983). However, a growing view, summarized by Zickler and Kleckner (2015), suggests otherwise:

A prominent, but still mysterious, feature of chromosome biology is the ability of homologous chromosomes, or chromosomal regions, to specifically recognize and pair with one another in the apparent absence of DNA lesions (DSBs) or recombination. ... Recombination-independent pairing ... plays prominent roles for premeiotic and meiotic programs, where it is defined as pairing that occurs before and/or without ... DSBs.

A “cut first” model implies a *localized commitment* prior to pairing. A “pair first” model should more reliably afford reversible genome-wide homology *testing*, without commitment (McGavin, 1977; Wilson, 1979; Boeteng et al. 2013). An initial alignment through “kissing” interactions between the loops of extruded DNA stem-loop structures requires only loop-loop base-pair complementation (Forsdyke 2007). Thus, initial *breakage-independent* homology recognition might not lead to strand breakage and segment exchange. This would require that dispersed loop homologies be interspersed with stem sequences that were also homologous.

Indeed, from studies of homology-directed DNA changes in fungi (repeat-induced point mutation; RIP), Gladyshev and Kleckner (2014) provide “a new perspective ... for ... the breakage-independent recognition of homology that underlies RIP and, potentially, other processes where sequence-specific pairing of intact chromosomes is involved.” Thus, “the nucleotide composition of participating DNA molecules is identified as an important factor,” and “homology recognition is modulated by the underlying sequence” (Gladyshev and Kleckner 2016). Accordingly, “sequence information can be compared directly between double-stranded DNA molecules during RIP,” and there is the potential for application to “other processes where homologous pairing of intact DNA molecules is observed.” This view is supported by later studies (Gladyshev and Kleckner 2017; Chapman et al. 2017).

However, the “pair first” and “cut first” views on the process by which genomes of paternal and maternal origin exchange information in the gonad of their child, should not necessarily be mutually exclusive. A limited number of pair-first sites might suffice (i.e. provide anchor points) to assist the close apposition of homologous chromosomes. Once this alignment was achieved, the homology-search task of single-strands liberated by a “cut first” mechanism should be easier.

Ideal evidence on the mechanisms through which sympatric homology searches might fail would involve species where primary sympatric hybrid sterility barriers had not been superseded by later acting barriers. As indicated earlier, retroviruses can help here.

Base Composition of Retroviruses

When we compare two viral species that have a *common* host cell, with two viral species that, even within a common host, do not share a common *cell*, we would expect to observe a fundamental difference related to their reproductive isolation mechanisms. If that difference is found to apply to other viral pairs that occupy a common host cell, then a fundamental isolation mechanism may have been identified. Retroviruses provided an example (Forsdyke 1996).

An important measure of base composition is the proportion of the individual bases G and C, among the four bases (A, C, G, T). This is expressed as GC%. The AIDS virus, HIV1, has a low GC% value. Thus, in its DNA form it is AT-rich. This happens to mainly reflect an increase in A, which is a purine (R), so the “R-loading” index is very high (Fig. 3). In contrast, the T cell leukaemia virus, HTLV1, has a high GC% value. Its GC-richness largely reflects an increase in C, which is not a purine, so the index of R-loading is very low.

The host of both these viruses is the T-lymphocyte. Assuming their evolution from a common ancestor and a continuity in their need to frequent a common host cell, then initial small difference in base composition could have played a role in preventing recombination between them within that host cell (analogous to a hybrid sterility barrier). This weak primary barrier to their recombination seems not to have been followed by the emergence of an effective secondary barrier. Their gene products had not adversely interacted (no ‘hybrid inviability’). Furthermore, although there are mechanisms to prevent ‘superinfection’ of an already infected cell (Forsdyke 2016, p184-186), their co-entry into a host cell was not absolutely denied (no ‘prezygotic isolation’). Rather, the primary barrier would have been progressively strengthened. This is a likely explanation for the extensive base-composition difference that we see between today’s co-

infecting viruses (insect viruses, herpes viruses, retroviruses; Forsdyke 1996). It is tempting to extend this explanation to the wide difference in GC% of certain protozoal parasites that undergo meiosis within a common host (Lee et al. 2004).




Fig. 3 Extreme differences in base compositions of two modern viruses that are presumed to have arisen from a common ancestor and are still capable of co-occupying a cell where mutual recombination could threaten their integrities. Base composition values (GC%) are from Bronson and Anderson (1994). Purine-loading (R-loading) indices are from Cristillo et al. (2001). Base compositions of the more flexible third positions of codons are shown in parentheses. HIV1 preservation correlates with greater AT-richness. HTLV1 preservation correlates with greater GC-richness. These base compositions correlate with specific oligonucleotide frequencies. A mismatch between oligonucleotides would impede the DNA ‘kissing’ interactions that may precede homologous recombination

Base composition differences are displayed not only with individual bases, but also with groups of bases (oligonucleotides). Thus, in HIV1 we find higher frequencies of AT-rich oligonucleotides (e.g. AAT, AAA, TAA), while in HTLV1 there are higher frequencies of GC-rich oligonucleotides (e.g. CCG, CCC, GCC). Species differences in these oligonucleotide frequencies, rather than in the individual bases (Brbić et al. 2015), should be critical in preventing recombination and thus maintaining the integrity of virus species. These oligonucleotides seem the best molecular candidates for the Bateson’s “residue” role.

Conclusions

Whether in sympatry or allopatry, any barrier in the germinal cycle can be primary in its potential to lead to divergence into species. However, in sympatry, hybrid sterility when operative can only be primary. It should be formally excluded before concluding that another barrier is primary.

Acknowledgements

Virgil Reese gave valuable advice. Queen's University hosts my biohistory web pages (<http://post.queensu.ca/~forsdyke/speciat5.htm> or <https://archive-it.org/collections/7641>).

The *arXiv* preprint server hosts an early version of this review.

References

- Bateson W (1922) Interspecific sterility. *Nature* 110:76
- Bernstein H, Bernstein C, Michod RE (2017) Sex in microbial pathogens. *Inf Genet Evol* 57:8–25
- Bhattacharyya T, Gregorova S, Mihola O et al (2013) Mechanistic basis of infertility of mouse intersubspecific hybrids. *Proc Natl Acad Sci USA* 110:E468–E477
- Bhattacharyya T, Reifova R, Gregorova S et al (2014) X chromosome control of meiotic chromosome synapsis in mouse inter-subspecific hybrids. *PLoS Genet* 10:e1004088
- Boateng KA, Bellani MA, Gregoretta IV et al (2013) Homologous pairing preceding SPO11-mediated double-strand breaks in mice. *Dev Cell* 24:196–1205
- Brbić M, Warnecke T, Kriško A et al (2015) Global shifts in genome and proteome composition are very tightly coupled. *Genome Biol Evol* 7:1519–1532
- Bronson EC, Anderson JN (1994) Nucleotide composition as a driving force in the evolution of retroviruses. *J Mol Evol* 38:506–532
- Chapman KM, Wilkey MM, Potter KE et al (2017) High homology is not required at the site of strand invasion during recombinational double-strand break repair in mammalian chromosomes. *DNA Repair* 60:1–8
- Cristillo AD, Mortimer JR, Barrette IH, et al. (2001) Double-stranded RNA as a not-self alarm signal: to evade, most viruses purine-load their RNAs, but some (HTLV-1, Epstein-Barr) pyrimidine-load. *J Theor Biol* 208:475–491

- Cock AG, Forsdyke DR (2008) Treasure your exceptions. The science and life of William Bateson. Springer, New York.
- Crowther CR (1922) Evolutionary faith and modern doubts. *Nature* 109:777
- Darwin CR (1862) Notes on the causes of cross and hybrid sterility. In: Burkhardt F, Porter DM, Harvey J, Topham JR (eds) *Correspondence of Charles Darwin* 10:700–711. Cambridge University Press, Cambridge
- Delph LF, Demuth JP (2016) Haldane's rule: genetic bases and their empirical support. *J Hered* 107:383–391
- Foot AD (2018) Sympatric speciation in the genomic era. *Trends Ecol Evol* 33: 86–95
- Forsdyke DR (1996) Different biological species "broadcast" their DNAs at different (G+C)% "wavelengths". *J Theor Biol* 178:405–417
- Forsdyke DR (2000) Haldane's rule: hybrid sterility affects the heterogametic sex first because sexual differentiation is on the path to species differentiation. *J Theor Biol* 204:443–452
- Forsdyke DR (2001) *The origin of species, revisited*. McGill-Queen's University Press, Montreal
- Forsdyke DR (2007) Molecular sex: the importance of base composition rather than homology when nucleic acids hybridize. *J Theor Biol* 249:325–330
- Forsdyke DR (2010) George Romanes, William Bateson, and Darwin's "weak point." *Notes Rec RS* 64:139–154
- Forsdyke DR (2011) The B in BDM. William Bateson did not advocate a genic speciation theory. *Heredity* 106:202
- Forsdyke DR (2016) *Evolutionary bioinformatics*, 3rd edn. Springer, New York
- Forsdyke DR (2017a) Speciation: Goldschmidt's chromosomal heresy, once supported by Gould and Dawkins, is again reinstated. *Biol Theor* 12:4–12
- Forsdyke DR (2017b) Base composition, speciation, and why the mitochondrial barcode precisely classifies. *Biol Theor* 12:157–168

- Forsdyke DR (2017c) The chromosomal basis of species initiation: Prdm9 as an anti-speciation gene. bioRxiv 170860. doi: <https://doi.org/10.1101/170860>
- Forsdyke DR (2018) The chromosomal basis of species initiation: *Prdm9* as an anti-speciation gene. Biol J Linn Soc (submitted)
- Fuller ZL, Leonard CJ, Young RE et al. (2017) The role of chromosomal inversions in speciation. bioRxiv 211771. doi: <https://doi.org/10.1101/211771>
- Gladyshev E, Kleckner N (2014) Direct recognition of homology between double helices of DNA in *Neurospora crassa*. Nat Commun 5:3509
- Gladyshev E, Kleckner N (2016) Recombination-independent recognition of DNA homology for repeat-induced point mutation (RIP) is modulated by the underlying nucleotide sequence. PLoS Genet 12:e1006015
- Gladyshev E, Kleckner N (2017) Recombination-independent recognition of DNA homology for repeat-induced point mutation. Curr Genet 63:389–400
- Gregorova S, Gergelits V, Chvatalova I et al. (2017) Modulation of Prdm9-controlled meiotic chromosome asynapsis overrides hybrid sterility in mice. bioRxiv 203505. doi: <https://doi.org/10.1101/203505>
- Haldane JBS (1922) Sex ratio and unisexual sterility in hybrid animals. J Genet 12:101–109
- Hu X-S, Filatov DA (2016) The large-X effect in plants: increased species divergence and reduced gene flow on the *Silene* X-chromosome. Mol Ecol 25:2609–2619
- Ironside JE, Filatov DA (2005) Extreme population structure and high interspecific divergence of the *Silene* Y chromosome. Genetics 171:705–713
- King M (1993) Species evolution. The role of chromosome change. Cambridge University Press, Cambridge
- Lee S-J, Mortimer JR, Forsdyke DR (2004) Genomic conflict settled in favour of the species rather than of the gene at extreme GC% values. Appl Bioinf 3:219–228
- Louis EC (2009) Origins of reproductive isolation. Nature 457:549–560

- McGavin S (1977) A model for the specific pairing of homologous double-stranded nucleic acid molecules during genetic recombination. *Heredity* (Edinb) 39:15–25
- Mérot C, Salazar C, Merrill RM et al (2017) What shapes the continuum of reproductive isolation? Lessons from *Heliconius* butterflies. *Proc R Soc B* 284:20170335
- Moehring AJ (2011) Heterozygosity and its unexpected correlations with hybrid sterility. *Evolution* 65:2621–2630
- Naviera HF, Maside XR (1998) The genetics of hybrid male sterility in *Drosophila*. In: Howard DJ, Berlocher SH (eds) *Endless forms: species and speciation*. Oxford University Press, Oxford, pp. 330–338
- Olajos F, Bokma F, Bartels P et al. (2018) Estimation species colonization dates using DNA in lake sediment. *Methods Ecol Evol* (in press)
- Reese VR, Forsdyke DR (2016) Meiotic pairing inadequacies at the levels of X chromosome, gene, or base: Epigenetic tagging for transgenerational error-correction guided by a future homologous duplex. *Biol Theor* 11:150–157
- Romanes GJ (1886) Physiological selection: an additional suggestion on the origin of species. *J Linn Soc (Zool)* 19:337–411
- Szostak JW, Orr-Weaver TL, Rothstein RJ et al (1983) The double-strand-break repair model for recombination. *Cell* 33:25–35
- White MJD (1978) *Modes of speciation*. Freeman, San Francisco, pp. 323–349 .
- Wilson JH (1979) Nick-free formation of reciprocal heteroduplexes: a simple solution to the topological problem. *Proc Natl Acad Sci USA* 76:3641–3645
- Winge Ö (1917) The chromosomes, their number and general importance. *Compt Rend Trav Lab Carlsberg* 13:131–275
- Zickler D, Kleckner N (2015) Recombination, pairing, and synapsis of homologs during meiosis. *Cold Spring Harb Pers Biol* 7(6). pii: a016626.