DIMENSION FREE PROPERTIES OF STRONG MUCKENHOUPT AND REVERSE HÖLDER WEIGHTS FOR RADON MEASURES

O. BEZNOSOVA AND A. REZNIKOV

ABSTRACT. In this paper we prove self-improvement properties of strong Muckenhoupt and Reverse Hölder weights with respect to a general Radon measure on \mathbb{R}^n . We derive our result via a Bellman function argument. An important feature of our proof is that it uses only the Bellman function for the one-dimensional problem for Lebesgue measure; with this function in hand, we derive dimension free results for general measures and dimensions.

Keywords: Bellman function, Dimension free estimates, Muckenhoupt weights, Reverse Hölder weights.

Mathematics Subject Classification: Primary: 42B35; Secondary: 43A85

1. Introduction

It is well known that Muckenhoupt weights on a real line with respect to the Lebesgue measure satisfy self-improvement properties in the following sense: for p > q we always have $A_q \subset A_p$; but also for any function $w \in A_p$, there is an $\varepsilon > 0$ such that $w \in A_{p-\varepsilon}$ (we refer to Definition 1 for precise definitions). Besides that, there always exists a q such that $w \in RH_q$. These self-improvement properties allow one to prove many important results in harmonic analysis, see, e.g., [4] or a more recent paper [5]. In [7], the authors considered strong Muckenhoupt classes; in particular, it was proven that for a Radon measure μ on \mathbb{R}^n which is absolutely continuous with respect to the Lebesgue measure dx, any weight $w \in A_p^*$ satisfies a Reverse Hölder property with an exponent that does not depend on the dimension n.

For p > 1, we say that w belongs to the strong Muckenhoupt class with respect to μ , $w \in A_p^*$, if there exists a number Q > 1 such that for any rectangular box $R \subset \mathbb{R}^n$ with edges parallel to axis, we have

$$\langle w \rangle_R \langle w^{-1/(p-1)} \rangle_R^{p-1} \leqslant Q,$$

where $\langle \phi \rangle_R$ denotes the average of the function ϕ over R:

$$\langle \varphi \rangle_R := \frac{1}{\mu(R)} \int\limits_R \varphi(x) d\mu(x).$$

For p > 1, we say that w belongs to the strong Reverse Hölder class with respect to μ , $w \in RH_p^*$, if there exists a constant Q > 1 such that for any rectangular box R with edges parallel to axis, we have

$$\langle w^p \rangle_R^{1/p} \leqslant Q \langle w \rangle_R.$$

We proceed with the following definition.

Date: August 25, 2021.

Definition 1. Let μ be a Radon measure on \mathbb{R}^n and w be a function which is positive μ -a.e. For p > 1, we denote the A_n^* -characteristic of w by

$$[w]_p := \sup_{R} \langle w \rangle_R \langle w^{-1/(p-1)} \rangle_R^{p-1}$$

and the Reverse Hölder characteristic of w by

$$[w]_{RH_p} := \sup_{R} \langle w^p \rangle_R^{1/p} \langle w \rangle_R^{-1},$$

where both suprema are taken over rectangular boxes R with edges parallel to axis. If $[w]_p < \infty$, we have $w \in A_p^*$ and if $[w]_{RH_p} < \infty$, we have $w \in RH_p^*$.

In [7] it was proved that if μ is an absolutely continuous Radon measure on \mathbb{R}^n and $[w]_p < \infty$, then for some q > 1 we have $[w]_{RH_q} < C < \infty$ with an explicit dimension free estimates on q and C. It is of a particular importance that we can take

$$q = 1 + \frac{1}{2^{p+2}[w]_p}.$$

To prove this result, the authors used a clever version of the Calderón–Zygmund decomposition from [6]. The aim of this paper is to derive a sharp result from the one-dimensional case for Lebesgue measure (i.e., for the classical A_p and RH_p classes on \mathbb{R}). In this case the result from [7] can be obtained, for example, by means of a so-called Bellman function; i.e., a function of two variables that satisfies certain boundary and concavity conditions in its domain. In the one-dimensional case this function is known explicitly, see [10]. It has been understood for some time that, for classes of functions like A_p , RH_p or BMO_p , when we work with their strong multi-dimensional analogs (e.g., A_p^* and RH_p^*), the one-dimensional Bellman function should prove the higher-dimensional results with dimension free constants. For the Lebesgue measure and the inclusion $RH_p^* \subset A_q^*$, this was done in [1]. The trick of using the Bellman function for one-dimensional problems was also used in [3], [2] and [9] (in a slightly different setting, the same trick was also used in [8]). In this paper, we present a simple version of this trick for general measures; we prove the result from [7] as well as all other results of self-improving type for strong Muckenhoupt and Reverse Hölder weights.

Acknowledgements: we are very grateful to Vasiliy Vasyunin for helpful discussions and suggestions on the presentation of this paper.

2. STATEMENT OF THE MAIN RESULT

2.1. **Properties of Muckenhoupt weights** A_p^* . For $p_1 := -1/(p-1)$ and every $t \in [0,1]$ define $u_{p_1}^{\pm}(t)$ to be solutions of the equation

$$(1-u)(1-p_1u)^{-1/p_1}=t.$$

The function $u_{p_1}^+$ is decreasing and maps [0,1] onto [0,1]; the function $u_{p_1}^-$ is increasing and maps [0,1] onto $[1/p_1,0]$. For a fixed Q > 1, define

(1)
$$s_{p_1}^{\pm} = s_{p_1}^{\pm}(Q) := u_{p_1}^{\pm}(1/Q).$$

Our first main result is the following.

Theorem 2.1. Let μ be a Radon measure on \mathbb{R}^n with $\mu(H) = 0$ for every hyperplane H orthogonal to one of the coordinate axis. Fix numbers p > 1 and Q > 1 and set $p_1 := -1/(p-1)$. Then for every weight w with $[w]_p = Q$ we have

$$w \in A_q^*, \quad 1 - s_{p_1}^-(Q) < q < \infty,$$

and

$$w \in RH_a^*$$
, $1 \le q < 1/s_{p_1}^+(Q)$,

where $s_{p_1}^{\pm}(Q)$ are defined in (1). These ranges for q are sharp for n=1 and $\mu=dx$.

2.2. **Properties of Reverse Hölder weights.** For p > 1 and every $t \in [0,1]$ we define $v_p^{\pm}(t)$ to be solutions of the equation

$$(1 - pv)^{1/p} (1 - v)^{-1} = t.$$

In this case, v_p^+ is a decreasing function that maps [0,1] onto [0,1/p] and v_p^- is an increasing function that maps [0,1] onto $[-\infty,0]$. As before for a fixed Q>1, we define

(2)
$$s_p^{\pm} = s_p^{\pm}(Q) := v_p^{\pm}(1/Q).$$

Our second main result concerning Reverse Hölder weights is the following.

Theorem 2.2. Let μ be a Radon measure on \mathbb{R}^n with $\mu(H) = 0$ for every hyperplane H orthogonal to one of the coordinate axis. Fix numbers p > 1 and Q > 1. Then for every weight w with $[w]_{RH_p} = Q$ we have

$$w \in A_q^*$$
, $1 - s_p^-(Q) < q < \infty$,

and

$$w \in RH_q^*, \quad 1 \leqslant q < 1/s_p^+(Q),$$

where $s_p^{\pm}(Q)$ are defined in (2). These ranges for q are sharp for n=1 and $\mu=dx$.

3. PROOF OF THE MAIN RESULTS

We begin with the following Theorem from [10]. This theorem ensures the existence of a certain Bellman function for a one-dimensional problem. In what follows, by letters without sub-indices (e.g., x, x^{\pm}) we denote points in \mathbb{R}^2 and by letters with sub-indices we denote the corresponding coordinates (e.g., x_1^+ denotes the first coordinate of x^+).

Theorem 3.1 (Theorem 1 in [10]). Fix p > 1 and set $p_1 := -1/(p-1)$. Also fix an $r \in (1/s_{p_1}^-, p_1] \cup [1, 1/s_{p_1}^+)$ for $s_{p_1}^\pm(Q)$ defined in (1). For every Q > 1 there exists a nonnegative function $B_Q(x)$ defined in the domain $\Omega_Q := \{x = (x_1, x_2) \in \mathbb{R}^2 : 1 \le x_1 x_2^{-1/p_1} \le Q\}$ with the following property: $B_Q(x)$ is continuous in x and Q, and for any line segment $[x^-, x^+] \subset \Omega_Q$ and $x = \lambda x^- + (1 - \lambda)x^+$, $\lambda \in [0, 1]$, we have

$$B_O(x) \geqslant \lambda B_O(x^-) + (1 - \lambda) B_O(x^+).$$

Moreover, $B(x_1, x_1^{p_1}) = x_1^r$ and $B_Q(x) \leq c(r, Q)x_1^r$ for some positive constant c(r, Q) and every $x \in \Omega_Q$.

To use the concavity property of the function B_Q for our proof, we need the following lemma. Its proof is given in [10, Lemma4] with an interval instead of the rectangle; however, the proof remains the same in our case.

Lemma 3.2. Let the measure μ be as before. Fix two numbers $Q_1 > Q > 1$ and a rectangular box $R \subset \mathbb{R}^n$ with edges parallel to the axis. For every coordinate vector \mathbf{e} , there exists a hyperplane H normal to \mathbf{e} that splits R into two rectangular boxes R^1 and R^2 with the following properties:

- (i) For i = 1, 2 we have $\mu(R^i)/\mu(R) \in (c, 1-c)$ for some constant $c \in (0,1)$;
- (ii) For every weight w with $[w]_p \leq Q$, we have $[x^1, x^2] \subset \Omega_{O_1}$ and, therefore,

$$B_{Q_1}(x_1,x_2) \geqslant \frac{\mu(R^1)}{\mu(R)} B_{Q_1}(x_1^1,x_2^1) + \frac{\mu(R^2)}{\mu(R)} B_{Q_1}(x_2^i,x_2^2),$$

where

$$x_1 = \langle w \rangle_R, \qquad x_2 = \langle w^{p_1} \rangle_R$$

 $x_1^i = \langle w \rangle_{R^i}, \qquad x_2^i = \langle w^{p_1} \rangle_{R^i}.$

We are ready to prove our main result.

Proof of Theorem 2.1. Fix a rectangular box R with edges parallel to the axis, and take any $Q_1 > Q$. We first explain how we split R into two rectangular boxes. Take one of the (n-1)-dimensional faces of R, call it R_{n-1} , that has the largest (n-1)-area. Among all (n-2)-dimensional faces of R_{n-1} , take one of those (call it R_{n-2}) that have the largest (n-2)-area. We proceed like this to get R_{n-1}, \ldots, R_1 . Now take a vector \mathbf{e} that is orthogonal to every R_i , $i=1,\ldots,n-1$.* We now split R according to Lemma 3.2. Notice that all the corresponding i-dimensional faces of R^1 and R^2 have smaller i-areas than the corresponding i-dimensional faces of R. We now take the boxes R^1 and R^2 and repeat the same procedure. If we repeat this M times, we get a family of rectangular boxes $\mathcal{R} = \{R^{i,M}\}_{i=1,\ldots,2^M}$. Denote

$$\begin{aligned} x_1 &= \left< w \right>_R, & x_2 &= \left< w^{p_1} \right>_R \\ x_1^{i,M} &= \left< w \right>_{R^{i,M}}, & x_2^{i,M} &= \left< w^{p_1} \right>_{R^{i,M}}. \end{aligned}$$

Abusing the notation, we also define step-functions

$$x_1^M(t) := \sum_{i=1}^{2^M} x_1^{i,M} \mathbb{1}_{R^{i,M}}(t), \qquad x_2^M(t) := \sum_{i=1}^{2^M} x_2^{i,M} \mathbb{1}_{R^{i,M}}(t).$$

From the construction of rectangular boxes, we notice that $x_1^M(t) \to w(t)$ and $x_2^M(t) \to w^{p_1}(t)$ as $M \to \infty$ for μ -a.e. $t \in R$. Indeed, our splitting procedure (and the fact that we have $\mu(R^i)/\mu(R) \in (1-c,c)$ at every step) guarantees that

$$\max_{i=1,\dots,2^M} \mathrm{diam}(R^{i,M}) \to 0, \ M \to \infty,$$

and we obtain the convergence of $x_1^M(t)$ and $x_2^M(t)$ from the Lebegue differentiation theorem for Radon measures. Therefore,

$$B_{Q_1}(x_1, x_2) \geqslant \sum_{i=1}^{2^M} \frac{\mu(R^{i,M})}{\mu(R)} B_{Q_1}(x_1^{i,M}, x_2^{i,M}) = \frac{1}{\mu(R)} \int_R B_{Q_1}(x_1^M(t), x_2^M(t)) d\mu(t).$$

^{*}In the case n=2, we just take **e** orthogonal to the longest side of R.

By the Fatou lemma,

$$(3) \quad B_{Q_1}(x_1, x_2) \geqslant \frac{1}{\mu(R)} \int_R \lim_{M \to \infty} B_{Q_1}(x_1^M(t), x_2^M(t)) d\mu(t) = \\ \frac{1}{\mu(R)} \int_R B_{Q_1}(w(t), w^{p_1}(t)) d\mu(t) = \frac{1}{\mu(R)} \int_R w^r(t) dt = \langle w^r \rangle_R.$$

Since $B_{Q_1}(x_1,x_2)$ is continuous in Q_1 and the above estimate holds for any $Q_1 > Q$, we get

$$c(r,Q)\langle w\rangle_R^r = c(r,Q)x_1^r \leqslant \langle w^r\rangle_R.$$

If we use this estimate for $q=r\in [1,1/s_{p_1}^+)$, we obtain $w\in RH_q^*$. If we use this estimate for $-1/(q-1)=r\in (1/s_{p_1}^-,p_1]$, we obtain $w\in A_q^*$ for $1-s_{p_1}^-(Q)< q<\infty$.

To prove Theorem 2.2 we need to use a different Bellman function B_Q . Namely, the following result holds.

Theorem 3.3 (Theorem 1 in [10]). Fix p > 1 and $r \in (1/s_p^-, 1] \cup [p, 1/s_p^+)$ for s_p^\pm defined in (2). For every Q > 1 there exists a non-negative function $B_Q(x)$ defined in the domain $\Omega_Q := \{x = (x_1, x_2) \in \mathbb{R}^2 : 1 \le x_1 x_2^{-1/p} \le Q\}$ with the following property: $B_Q(x)$ is continuous in x and Q, and for any line segment $[x^-, x^+] \subset \Omega_Q$ and $x = \lambda x^- + (1 - \lambda)x^+$, $\lambda \in [0, 1]$, we have

$$B_{Q}(x) \geqslant \lambda B_{Q}(x^{-}) + (1 - \lambda)B_{Q}(x^{+}).$$

Moreover, $B(x_1, x_1^p) = x_1^r$ and $B_Q(x) \leq c(r, Q)x_1^r$ for some positive constant c(r, Q) and every $x \in \Omega_O$.

We also notice that the analog of Lemma 3.2 reads the same, and with this in hand, the proof of Theorem 2.1 is analogous to the proof of Theorem 2.2; we leave the details to the reader.

REFERENCES

- [1] M. Dindoš and T. Wall. The sharp A_p constant for weights in a reverse-Hölder class. *Rev. Mat. Iberoam.*, 25(2):559–594, 2009.
- [2] K. Domelevo, S. Petermichl, and J. Wittwer. A linear dimensionless bound for the weighted Riesz vector. *Bull. Sci. Math.*, 141(5):385–407, 2017.
- [3] O. Dragičević and A. Volberg. Bellman functions and dimensionless estimates of Littlewood-Paley type. *J. Operator Theory*, 56(1):167–198, 2006.
- [4] J. García-Cuerva and J. L. Rubio de Francia. *Weighted norm inequalities and related topics*, volume 116 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1985. Notas de Matemática [Mathematical Notes], 104.
- [5] T. Hytönen and C. Pérez. Sharp weighted bounds involving A_{∞} . Anal. PDE, 6(4):777–818, 2013.
- [6] A. A. Korenovskyy, A. K. Lerner, and A. M. Stokolos. On a multidimensional form of F. Riesz's "rising sun" lemma. *Proc. Amer. Math. Soc.*, 133(5):1437–1440, 2005.
- [7] T. Luque, C. Pérez, and E. Rela. Reverse Hölder property for strong weights and general measures. *J. Geom. Anal.*, 27(1):162–182, 2017.
- [8] F. Nazarov, A. Reznikov, S. Treil, and A. Volberg. A Bellman function proof of the L^2 bump conjecture. *J. Anal. Math.*, 121:255–277, 2013.
- [9] S. Treil. Sharp A_2 estimates of Haar shifts via Bellman function. In *Recent trends in analysis*, volume 16 of *Theta Ser. Adv. Math.*, pages 187–208. Theta, Bucharest, 2013.
- [10] V. I. Vasyunin. Mutual estimates for L^p-norms and the Bellman function. Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI), 355(Issledovaniya po Line<u>inym Operatoram i Teorii Funktsii.</u> 36):81–138, 237–238, 2008.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALABAMA *E-mail address*: ovbeznosova@ua.edu

Department of Mathematics, Florida State University

E-mail address: reznikov@math.fsu.edu