


# REALIZATION OF SHIFT GRAPHS AS DISJOINTNESS GRAPHS OF 1-INTERSECTING CURVES IN THE PLANE

TORSTEN MÜTZE, BARTOSZ WALCZAK, AND VEIT WIECHERT

The *disjointness graph* of a family  $\mathcal{F}$  of curves in the plane is the graph with vertex set  $\mathcal{F}$  and with edges between the pairs of disjoint curves. In the *shift graph*  $H_m$ , the vertices are the ordered pairs  $(i, j)$  satisfying  $1 \leq i < j \leq m$ , and two such pairs  $(i, j)$  and  $(k, \ell)$  form an edge if and only if  $j = k$  or  $\ell = i$ . The graphs  $H_m$  are triangle-free and have unbounded chromatic number [1, Theorem 6] (in fact, it is well known that  $\chi(H_m) = \lceil \log_2 m \rceil$ ). We strengthen a result of Pach, Tardos, and Tóth [5, Theorem 5] showing that  $H_m$  can be realized as the disjointness graph of a family  $\mathcal{F}_m$  of polygonal curves in the plane with the following properties: (a) every curve in  $\mathcal{F}_m$  is made of 3 straight-line segments, and (b) any two curves in  $\mathcal{F}_m$  intersect in at most one point.


Let  $\prec$  denote the lexicographic order on the vertices of  $H_m$ :  $(i, j) \prec (k, \ell)$  when  $i < k$  or  $i = k$  and  $j < \ell$ . First, we construct a realization in which the curve  $c(i, j)$  representing the vertex  $(i, j)$  of  $H_m$  consists of 4 straight-line segments as illustrated, where

- the  $A$ -parts occur in the order  $\prec$  from right to left,
- the  $B$ -parts occur in the order  $\prec$  from bottom to top,
- the  $C$ -parts occur in the order  $\prec$  from top-right to bottom-left,
- the  $D$ -parts occur in the order  $\prec$  from left to right,
- the  $A$ -part of  $c(i, j)$  intersects the  $B$ -parts of  $c(k, \ell)$  up to  $c(j - 1, m)$ ,
- the  $D$ -part of  $c(i, j)$  intersects the  $B$ -parts of  $c(k, \ell)$  down to  $c(j + 1, j + 2)$  (if  $j \leq m - 2$ ).



If  $(i, j) \prec (k, \ell)$ , then  $c(i, j)$  and  $c(k, \ell)$  intersect if and only if  $j \neq k$ , as required. By a standard stretching argument (see e.g. [4, Theorem 3]), the  $AB$ -parts of the curves can be replaced by single straight-line segments with the same intersection pattern.

The curves in the non-stretched realization can be arranged to touch (but not cross) a common straight line. By contrast, triangle-free intersection graphs of curves touching (but not crossing) a common straight line have bounded chromatic number [3]. An analogous contrast but in the opposite way holds for graphs realized by straight-line segments: while there exist triangle-free segment intersection graphs with arbitrarily large chromatic number [6], the proof technique of Larman et al. [2, Theorem 1] shows that disjointness graphs of straight-line segments (or, more generally,  $x$ -monotone curves) in the plane satisfy  $\chi = O(\omega^4)$ .



## REFERENCES

- [1] Paul Erdős and András Hajnal, Some remarks on set theory. IX. Combinatorial problems in measure theory and set theory, *Michigan Math. J.* 11 (2), 107–127, 1964.
- [2] David Larman, Jiří Matoušek, János Pach, and Jenő Törőcsik, A Ramsey-type result for convex sets, *Bull. London Math. Soc.* 26 (2), 132–136, 1994.
- [3] Sean McGuinness, Colouring arcwise connected sets in the plane I, *Graphs Comb.* 16 (4), 429–439, 2000.
- [4] Matthias Middendorf and Frank Pfeiffer, The max clique problem in classes of string-graphs, *Discrete Math.* 108 (1–3), 365–372, 1992.
- [5] János Pach, Gábor Tardos, and Géza Tóth, Disjointness graphs of segments, in: Boris Aronov and Matthew J. Katz (eds.), *33rd International Symposium on Computational Geometry (SoCG 2017)*, vol. 77 of *Leibniz International Proceedings in Informatics (LIPIcs)*, 59:1–15, Leibniz-Zentrum für Informatik, Dagstuhl, 2017.
- [6] Arkadiusz Pawlik, Jakub Kozik, Tomasz Krawczyk, Michał Lasoń, Piotr Micek, William T. Trotter, and Bartosz Walczak, Triangle-free intersection graphs of line segments with large chromatic number, *J. Comb. Theory Ser. B* 105, 6–10, 2014.