

# DIVERGENT ON AVERAGE DIRECTIONS OF TEICHMÜLLER GEODESIC FLOW

PAUL APISA AND HOWARD MASUR

ABSTRACT. The set of directions from a quadratic differential that diverge on average under Teichmüller geodesic flow has Hausdorff dimension exactly equal to one-half.

## 1. INTRODUCTION

Suppose  $g_t$  is a flow on a topological space  $S$ . We say that  $p \in S$  is *divergent* if  $\{g_t p\}_{t \geq 0}$  eventually leaves every compact set of  $S$ . We say that  $p$  is *divergent on average* if the proportion of time that  $\{g_t p\}_{0 \leq t \leq T}$  spends in any compact subset of  $S$  tends to zero as  $T$  tends to  $\infty$ .

Let  $\mathcal{M}_{g,n}$  be the moduli space of closed genus  $g$  Riemann surfaces with  $n$  punctures. The cotangent space of  $\mathcal{M}_{g,n}$  coincides with the moduli space  $Q_{g,n}$  of holomorphic quadratic differentials on surfaces in  $\mathcal{M}_{g,n}$  and admits an  $\mathrm{SL}(2, \mathbb{R})$  action generated by complex scalar multiplication  $r_\theta(q) = e^{i\theta}q$  and Teichmüller geodesic flow  $g_t$ . (see the survey paper of [Zor06] for the definition of the flat structure defined by a quadratic or Abelian differential and the Teichmüller geodesic flow). The space  $Q_{g,n}$  admits an  $\mathrm{SL}(2, \mathbb{R})$ -invariant stratification by specifying the number of zeros and poles and their orders of vanishing.

For any holomorphic quadratic differential  $q$  it is a consequence of Chaika-Eskin [CE15, Theorem 1.1] that the set of directions  $\theta$  such that  $r_\theta q$  diverges on average under the Teichmüller flow in its stratum has measure zero. We prove the following result.

**Theorem 1.** *For a quadratic or Abelian differential  $q$  the set of directions  $\theta$  such that  $r_\theta q$  diverges on average (either in its stratum or in  $Q_{g,n}$ ) has Hausdorff dimension exactly equal to  $\frac{1}{2}$ .*

As a corollary, we have

**Corollary 1.** *Let  $\mathcal{H}$  be any stratum of quadratic or Abelian differentials. Let  $f : \mathcal{H} \rightarrow \mathbb{R}$  be any compactly supported continuous function. Then for any  $q \in \mathcal{H}$  the set of  $\theta \in [0, 2\pi)$  such that*

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(g_t r_\theta q) dt = 0$$

has Hausdorff dimension at least  $\frac{1}{2}$ .

**Remark 1.** For any quadratic differential the set of directions that diverge on average in  $Q_{g,n}$  is contained in the set of directions that diverge on average in the stratum. In al-Saqban-Apisa-Erchenko-Khalil-Mirzadeh-Uyanik [ASAE<sup>+</sup>17], the authors adapted the techniques of Kadyrov-Kleinbock-Lindenstrauss-Margulis [KKLM14] to show that the latter set has Hausdorff dimension at most  $\frac{1}{2}$  (this result improves on results of Masur [Mas91, Mas92]). Therefore, the novelty of the current work is establishing the lower bound of Hausdorff dimension  $\frac{1}{2}$  for the set of directions that diverge on average in  $Q_{g,n}$ .

**Remark 2.** The methods of [ASAE<sup>+</sup>17] in fact show that the Hausdorff dimension of the set of directions that diverge on average in any open invariant subset of a stratum is at most  $\frac{1}{2}$ . Therefore, Theorem 1 remains true when divergence on average is considered in a stratum with finitely many affine invariant submanifolds deleted.

**Remark 3.** In the classical case of  $SL(2, \mathbb{R})/SL(2, \mathbb{Z})$  it is known that the set of directions that diverge on average has Hausdorff dimension  $\frac{1}{2}$ . In fact, the behavior of a geodesic is determined by the continued fraction expansion of its endpoint  $x = [a_0, a_1, a_2, \dots]$  (see for instance Dani [Dan85]). The geodesic diverges on average if and only if  $(\prod_{i=1}^n a_i)^{\frac{1}{n}}$  goes to  $\infty$  as  $n \rightarrow \infty$  (see Choudhuri [Cho17, Theorem 1.2]) and this set has Hausdorff dimension  $\frac{1}{2}$  by [FLWW09, Theorem 1.2]. In Cheung [Che11], it is shown that the set of real numbers for which  $a_n$  tends to  $\infty$  at a certain prescribed rate has Hausdorff dimension  $\frac{1}{2}$ . Our construction in higher genus Teichmüller space is modeled on this construction.

**Connection with Previous Results.** For any holomorphic quadratic differential  $q$  on a Riemann surface  $X$ , every direction gives a foliation of the underlying Riemann surface. By Masur [Mas92, Theorem 1.1], if the Teichmüller geodesic in a given direction is recurrent, then the foliation is uniquely ergodic. In particular, the non-uniquely ergodic directions -  $NUE(q)$  - are divergent directions.

By results of Strebel [Str84] and Katok-Zemlyakov [ZK75], the collection of directions with non-minimal flow -  $NM(q)$  - is countable. In [MS91], the main theorem is that outside finitely many exceptional strata of quadratic differentials (the exceptions being the ones where every flat structure induced by a holomorphic quadratic differential has a holonomy double cover that is a translation covering of a flat torus), there is a constant  $\delta > 0$  depending on the stratum so that for almost-every quadratic differential  $q$  in the stratum the set of directions with

non-ergodic flow -  $\text{NE}(q)$  - has Hausdorff dimension at least  $\delta$ . The sequence of inclusions is then

$$\text{NM}(q) \subseteq \text{NE}(q) \subseteq \text{NUE}(q) \subseteq D(q) \subseteq \text{DA}(q)$$

where  $D(q)$  and  $\text{DA}(q)$  are the set of directions that diverge (resp. diverge on average). The set  $D(q)$  was shown to have measure zero in Kerckhoff-Masur-Smillie [KMS86, Theorem 4].

The set  $\text{NUE}(q)$  was shown to have Hausdorff dimension at most  $\frac{1}{2}$  by the main theorem of Masur [Mas92]. Recently, Chaika-Masur [CM18] showed that for hyperelliptic components of strata of Abelian differentials this inequality is actually an equality for almost every Abelian differential.

**Problem 1.** Is it the case that the Hausdorff dimension of  $\text{NUE}(q)$  is either 0 or  $\frac{1}{2}$  for all quadratic differentials  $q$ .

For all known examples, the dimension is either 0 or  $\frac{1}{2}$ . Despite the fact that  $\text{NE}(q)$  has positive Hausdorff dimension for a full measure set of quadratic differentials (outside of finitely many exceptional strata), in each stratum there is a dense set of Veech surfaces, for which  $D(q) = \text{NM}(q)$  and hence is countable. The fact that  $D(q)$  is positive dimensional for a full measure set of  $q$  and zero-dimensional for a dense set of  $q$  shows that an analogue of Theorem 1 for divergent directions does not exist in general.

**Acknowledgements.** The authors thank Jon Chaika for suggesting this problem. They also thank Alex Eskin and Kasra Rafi for helpful conversations. This material is based upon work supported by the National Science Foundation under Grant No. DMS-1607512. HM gratefully acknowledges their support. This material is also based upon work supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE-1144082. PA gratefully acknowledges their support.

## 2. CYLINDERS AND THE THICK-THIN DECOMPOSITION

Given a Riemann surface  $X$  with a holomorphic quadratic differential there are two natural metrics on  $X$  - the hyperbolic metric and the flat metric induced by the quadratic differential. Assume that the quadratic differential has unit area. Fix  $\delta > 0$  small enough so that two curves of hyperbolic-length less than  $\delta$  are disjoint.

The thick-thin decomposition of the flat surface is defined as follows (for simplicity we only state the definition for an Abelian differential -  $(X, \omega)$  - which will be sufficient for our purpose). Let  $\gamma_k$  be the simple

closed curves on  $X$  of hyperbolic-length less than  $\delta$ . There is a geodesic representative of  $\gamma_k$  in the flat metric on  $(X, \omega)$ . Either the flat-geodesic is unique or it is contained in a flat cylinder. In the first case, cut out the unique flat-geodesic from  $(X, \omega)$  and in the second excise the entire cylinder. The resulting components are called  $\delta$ -thick-pieces. The size of a thick-piece is defined as the smallest flat length of a simple closed curve in the thick piece that is not homotopic to a boundary curve. These definitions are due to Rafi [Raf07] who showed that in a thick piece the hyperbolic length of any closed curve is comparable to its flat length divided by the size of the piece.

In Eskin-Kontsevich-Zorich [EKZ, Geometric Compactification Theorem (Theorem 10)], the following is established (we only state a version for Abelian differentials)

**Theorem 2.** *Let  $(X_n, \omega_n)$  be any sequence of unit-area translation surfaces that are not contained in a compact subset of a stratum of Abelian differentials. By passing to a subsequence assume that  $(X_n, \omega_n)$  converges to a stable differential  $\omega$  on a nodal Riemann surface  $X$ . Let  $\delta_0$  be less than half the injectivity radius of  $X$  in the hyperbolic metric on the desingularized surface. Then there is a subsequence of  $(X_n, \omega_n)$  so that each thick component converges to a nonzero meromorphic quadratic differential when the flat metric on the thick component is renormalized so that its size is one.*

**Lemma 3.** *Under the hypotheses of Theorem 2, if a thick piece has area bounded away from 0 on  $(X_n, \omega_n)$  for all  $n$  then its size is also bounded away from zero.*

*Proof.* By Eskin-Kontsevich-Zorich [EKZ, Lemma 4.2], each thick piece can be triangulated in the flat metric with saddle connections of length comparable to the size of the thick piece. Any such triangulation has a bounded number of triangles (depending only on the stratum) and so if the size of a thick piece tends to zero its area must as well.  $\square$

**Definition 1.** Suppose that  $A$  is an annulus around a curve  $\gamma$  on  $X$ . The annulus  $A$  is called regular if it is of the form  $\{p : d(p, \gamma) < r\}$  for some  $r$  where  $d(\cdot, \cdot)$  denotes distance in the flat metric. The annulus is primitive if additionally it contains no singularities in its interior. If  $A$  is a primitive regular annulus that is not a flat cylinder, then define  $\mu(A) := \log \left( \frac{|\gamma_o|}{|\gamma_i|} \right)$  where  $|\cdot|$  denotes flat length and  $\gamma_o$  (resp  $\gamma_i$ ) is the longer (resp. shorter) boundary curve of  $A$  and is called the outer (resp. inner) curve of  $A$ . This definition agrees with the one made in Minsky [Min92] up to a multiplicative constant that only depends on the stratum containing  $(X, \omega)$ .

**Lemma 4.** *Under the hypotheses of Theorem 2, every flat cylinder in  $(X_n, \omega_n)$  around a  $\delta_0$ -hyperbolically short curve has the length of its core curve tend to 0 as  $n \rightarrow \infty$ .*

*Proof.* By Maskit [Mas85, Corollary 2], since the  $\delta_0$ -hyperbolically short curves have lengths tending to zero in the hyperbolic metric, their extremal length also tends to zero as  $n \rightarrow \infty$ . The extremal length of a curve  $\gamma$  is defined to be the reciprocal of the modulus of the largest topological annulus embedded in the hyperbolic surface whose waist curve is freely homotopic to  $\gamma$ .

Therefore, each  $\delta_0$ -hyperbolically short curve  $\gamma$  is contained in a topological annulus whose modulus tends to  $\infty$  as  $n \rightarrow \infty$ . By Minsky [Min92, Theorem 4.5 and 4.6] (note that the inequality  $\leq m_0$  should be  $\geq m_0$  in the statement of the Theorem 4.6), either  $\gamma$  is contained in a flat cylinder whose modulus is unbounded in  $n$  or there is a primitive regular annulus  $A_n \subseteq (X_n, \omega_n)$  contained in a thick piece whose core curve is homotopic to  $\gamma$  and so that  $\mu(A_n)$  tends to  $\infty$  as  $n$  increases.

Notice that if the modulus of the flat cylinder containing  $\gamma$  tends to  $\infty$  then the flat length of  $\gamma$  tends to zero since each  $(X_n, \omega_n)$  is unit-area. Therefore, suppose that for each  $n$  there is a primitive regular annulus  $A_n$  whose core curve is homotopic to  $\gamma$  and so that  $\mu(A_n)$  is unbounded in  $n$ .

Let  $\ell_n$  be the flat length of  $\gamma$  on  $(X_n, \omega_n)$  and let  $a_n$  be the area of the thick piece containing  $A_n$ . The flat distance across  $A_n$  is at most  $h_n := \frac{a_n}{\ell_n}$ . The flat length of the outer curve  $A_n$  in the flat metric is at most  $2\ell_n + 2\pi M h_n$  where  $M$  is some integer only depending on the stratum. Therefore,  $\mu(A_n) \leq \log\left(2 + \frac{2\pi M}{\ell_n}\right)$ . Since  $\mu(A_n)$  is unbounded in  $n$ ,  $\ell_n \rightarrow 0$  as  $n \rightarrow \infty$ .  $\square$

A similar argument to the one above is given in Choi-Rafi-Series [CRS08, Corollary 5.4]. The rest of this section is devoted to the proof of the following proposition.

**Proposition 1.** *Given a stratum of quadratic or Abelian differentials, there are constants  $N_0, c$  so that for any unit area  $(X, \omega)$  there is a cylinder  $C_\beta$  with*

- $|\beta| < N_0$
- $\text{area}(C_\beta) > c$

*Proof.* By taking a holonomy double cover in the quadratic differential case, we will assume we are in the Abelian case. We will prove the proposition in this case by finding a cylinder and then projecting to

find a cylinder for the quadratic differential. We begin with the proof of the first conclusion. The following argument is essentially found in Masur [Mas86].

**Step 1: There is a constant  $N_0$  so that every translation surface in the stratum contains a cylinder whose core curve has length less than  $N_0$ .**

Proceed by induction on the dimension of the stratum. The base case of genus one translation surfaces is trivial. Suppose now that we are in a stratum of Abelian differentials on genus  $g > 1$  surfaces. Suppose to a contradiction that  $(X_n, \omega_n)$  is a sequence of unit area translation surfaces in the stratum so that the length of the shortest cylinder tends to  $\infty$ . By passing to a subsequence we may suppose that  $(X_n, \omega_n)$  converges to a stable differential  $(X, \omega)$ . Letting  $\delta_0 > 0$  be half the injectivity radius on the desingularized hyperbolic surface  $X$ , Theorem 2 implies that after passing to another subsequence we may suppose that each  $\delta_0$ -thick piece converges.

**Step 1a: There is a thick piece that has definite area on each  $(X_n, \omega_n)$ .** In the thick-thin decomposition, the only positive-area sub-surfaces that are not contained in a thick piece are flat cylinders around  $\delta_0$ -hyperbolically short curves. By Lemma 4, the lengths of the core curves of these cylinders tend to zero along the sequence  $(X_n, \omega_n)$ , which would contradict our hypotheses. Therefore, there are no such cylinders and some thick piece therefore has definite area,

**Step 1b: There is a cylinder of bounded length in the definite-area thick piece**

By Lemma 3, any thick piece that has definite area on each  $(X_n, \omega_n)$  also has definite size for each  $(X_n, \omega_n)$ . Therefore, the thick piece must converge to a holomorphic (not just meromorphic) Abelian differential on  $(X, \omega)$  by Theorem 2. By the induction hypothesis, this Abelian differential contains a cylinder of bounded length (where the bound only depends on the genus of the limiting component of the desingularized Riemann surface  $X$ ). Since the thick pieces of definite area converge in the flat metric, it follows that for sufficiently large  $n$  they contain a cylinder of bounded length, which is a contradiction.

**Step 2: There is a constant  $c > 0$  so that the cylinder found in the previous step may be assumed to have area at least  $c$**

For the second conclusion, again proceed by induction on the dimension of the stratum. The base case of genus one translation surfaces is straightforward. Suppose now that we are in a stratum of Abelian differentials on genus  $g > 1$  surfaces. Suppose to a contradiction that  $(X_n, \omega_n)$  is a sequence of unit area translation surfaces in the stratum so that the area of the largest-area cylinder with core curve of length

less than  $N_0$  tends to 0. Suppose without loss of generality that the value  $N_0$  is taken to be larger than the value of  $N_0 + 1$  for any lower dimensional stratum of Abelian differentials. As above, pass to a subsequence  $(X_n, \omega_n)$  that converges to a stable Abelian differential  $(X, \omega)$ , let  $\delta_0$  be less than half the injectivity radius of the desingularized hyperbolic metric on  $X$ , and suppose that the  $\delta_0$  thick pieces converge as in Theorem 2.

Again we claim there is a thick piece that has definite area on each  $(X_n, \omega_n)$ . In the thick-thin decomposition, the only positive-area sub-surfaces that are not contained in a thick piece are flat cylinders around  $\delta_0$ -hyperbolically short curves. However, by Lemma 4 these cylinders have the length of their core curves tend to zero along the sequence  $(X_n, \omega_n)$ . If some such cylinder had definite area then we would contradict our hypotheses on the sequence  $(X_n, \omega_n)$ . Therefore, all such cylinders have area tending to zero and some thick piece therefore has definite area.

Then just as before, every thick piece that has definite area on each  $(X_n, \omega_n)$  contains a cylinder of length smaller than  $N_0$  and definite area for all sufficiently large  $n$ , which is a contradiction.  $\square$

**Remark 4.** In every stratum of quadratic and Abelian differentials there is a unit-area flat surface that contains a unit-area cylinder whose circumference is length one. On such a surface every cylinder has circumference of length at least one and so  $N_0 \geq 1$ .

### 3. PROOF OF THEOREM 1

In this section we give a proof of Theorem 1 based on Lemma 5 - proved in this section - and Proposition 2 - proved in the next. The strategy is essentially found in Cheung [Che11, pages 23-24], in the context of continued fractions.

**Definition 2.** Suppose  $(X, \omega)$  is either a quadratic or Abelian differential. Given a cylinder  $C_\beta$  with core curve  $\beta$  and area at least  $c$  on  $(X, \omega)$ , let  $|\beta|$  denote the length of  $\beta$  and  $0 \leq \theta_\beta \leq \pi/2$  the angle it makes with the horizontal direction. Define  $I_\beta$  to be the interval of angles centered at  $\theta_\beta$  with radius  $\frac{1}{|\beta|^2 \log |\beta|}$ . Fix an integer  $M$  to be determined later. Say that a cylinder of area at least  $c$  with core curve  $\beta'$  is a *child* of  $\beta$  if

$$|\beta| \log |\beta| \leq |\beta'| \leq M |\beta| \log |\beta|$$

and  $I_{\beta'} \subseteq I_\beta$ .

**Lemma 5.** *Let  $0 < \epsilon < 1$ . Suppose  $\beta'$  is a child of  $\beta$  and  $\theta \in I_{\beta'} \subset I_{\beta}$ . Suppose too that  $\frac{4}{\epsilon^2} < \log |\beta|$ . Then for all*

$$t \in [\log |\beta|, \log |\beta'|],$$

*except for a subset of size at most  $\log \frac{4M}{\epsilon^2}$ ,  $\beta$  has flat length at most  $\epsilon$  on  $g_t r_{\theta}(X, \omega)$ .*

*Proof.* Suppose without loss of generality that we have rotated  $(X, \omega)$  so that  $\theta$  is the vertical direction. Let  $h(t)$  and  $v(t)$  be the horizontal (resp. vertical) component of the period of  $\beta$  on  $g_t r_{\theta}(X, \omega)$ . When  $t = \log |\beta| + \log \left(\frac{2}{\epsilon}\right)$  we see that

$$v(t) \leq \frac{\epsilon}{2} \quad \text{and} \quad h(t) \leq \left(\frac{2|\beta|}{\epsilon}\right) |\beta| \sin \left(\frac{1}{|\beta|^2 \log |\beta|}\right) \leq \frac{2\epsilon^{-1}}{\log |\beta|} \leq \frac{\epsilon}{2}.$$

Similarly when  $t = \log |\beta'| - \log \left(\frac{2M}{\epsilon}\right)$  we see that

$$v(t) \leq \frac{\epsilon}{2} \quad \text{and} \quad h(t) \leq (M|\beta| \log |\beta|) \left(\frac{\epsilon}{2M}\right) |\beta| \sin \left(\frac{1}{|\beta|^2 \log |\beta|}\right) \leq \frac{\epsilon}{2}.$$

Therefore, for all times

$$t \in \left[ \log |\beta| + \log \left(\frac{2}{\epsilon}\right), \log |\beta'| - \log \left(\frac{2M}{\epsilon}\right) \right]$$

the curve  $\beta$  has length at most  $\epsilon$  on  $g_t r_{\theta}(X, \omega)$  □

Let  $D$  be the collection of angles  $\theta$  that can be written as

$$\{\theta\} = \bigcap_n I_{\beta_n}$$

where  $(\beta_n)$  is a sequence of curves so that  $\beta_n$  is a child of  $\beta_{n-1}$  for all  $n$ .

**Corollary 6.** *Any angle  $\theta \in D$  is a divergent on average direction in the moduli space of Riemann surfaces  $\mathcal{M}_{g,n}$  (not just in the stratum of quadratic differentials).*

*Proof.* Let  $\epsilon > 0$ . By the Mumford compactness theorem,  $\mathcal{M}_{g,n}$  has a compact exhaustion by sets  $K_{\epsilon}$  of Riemann surfaces on which all simple closed curves have hyperbolic length at least  $\epsilon$ . By Maskit [Mas85], for sufficiently small  $\epsilon$ , there is an  $\epsilon' > 0$  so that  $K_{\epsilon}$  is contained in the set of Riemann surfaces on which all simple closed curves have extremal length at least  $\epsilon'$ .

Let  $\epsilon'' := \sqrt{c\epsilon'}$  and let  $t_n := \log |\beta_n|$ . Since  $(|\beta_n|)_n$  is an increasing sequence that tends to  $\infty$  let  $N$  be an integer such that  $\frac{4}{(\epsilon'')^2} < \log |\beta_n|$  for  $n > N$ . By Lemma 5, for all  $n > N$  and for all but at most  $\log \frac{4M}{(\epsilon'')^2}$



times in  $[t_n, t_{n+1}]$  the translation surfaces  $\{g_t r_\theta(X, \omega)\}_{t=t_n}^{t_{n+1}}$  contain a cylinder with core curve  $\beta_n$  of length less than  $\epsilon''$  and of area at least  $c$ . For these times  $\beta_n$  has extremal length at most  $\frac{(\epsilon'')^2}{c} = \epsilon'$  and hence the underlying Riemann surface lies outside of  $K_\epsilon$ . Since  $t_{n+1} - t_n$  tends to  $\infty$  as  $n \rightarrow \infty$  whereas the amount of time spent in  $K_\epsilon$  for times in  $[t_n, t_{n+1}]$  is at most  $\log \frac{4M}{(\epsilon'')^2}$ , we see that  $\{g_t r_\theta(X, \omega)\}$  spends asymptotically zero percent of its time in  $K_\epsilon$  as desired.  $\square$

**Proposition 2.** *There are constants  $C, \delta$ , and  $\nu$  depending on  $N_0$  and  $c$  so that for every integer  $M \geq 2$  the following holds. Suppose that  $(X, \omega)$  is a quadratic or Abelian differential and let  $\beta$  be the core curve of a cylinder with area at least  $c$ . Define  $N_\beta := \log |\beta|$  and  $\rho_\beta := \frac{C}{N_\beta}$ .*

*For  $|\beta|$  sufficiently large there is a subset  $J_\beta \subset \left[\frac{1}{\delta}N_\beta, \frac{M}{2N_0}N_\beta\right] \cap \mathbb{Z}$  so that for each  $j \in J_\beta$  there is a child  $\beta_j$  of  $\beta$  so that*

- (1) *Successive integers  $j, k \in J_\beta$  satisfy  $|k - j| \leq 2\nu$ .*
- (2) *The distance between  $I_{\beta_j}$  and  $I_{\beta_k}$  for any two children  $\beta_j, \beta_k$  is at least  $\rho_\beta |I_\beta| = \frac{C}{N_\beta^2 |\beta|^2}$ .*
- (3) *For  $j \in J_\beta$  and letting  $N_{\beta_j} := \log |\beta_j|$ , we have -*

$$\frac{N_\beta |\beta|}{N_{\beta_j} |\beta_j|} \geq \frac{1}{2N_0 j}.$$

Assuming Proposition 2 we will now prove Theorem 1.

*Proof of Theorem 1:* Setting  $s = \frac{1}{2}$  we see that

$$\sum_{\beta' \in D_\beta} \frac{\rho_{\beta'}^s |I_{\beta'}|^s}{\rho_\beta^s |I_\beta|^s} = \sum_{\beta' \in D_\beta} \left(\frac{CN_\beta}{CN_{\beta'}}\right)^{1/2} \left(\frac{N_\beta |\beta|^2}{N_{\beta'} |\beta'|^2}\right)^{1/2} = \sum_{\beta' \in D_\beta} \frac{N_\beta |\beta|}{N_{\beta'} |\beta'|}$$

where  $D_\beta$  is the set of children of  $\beta$  constructed in Proposition 2. We can assume  $|\beta|$  is large enough so that the proposition holds. By assumption (3) the sum on the right is greater than

$$\sum_{j \in J_\beta} \frac{1}{2N_0 j} \geq \frac{1}{4N_0 \nu} \log \left(\frac{\delta M}{2N_0}\right)$$

where the inequality follows from assumption (1). Now the right side is greater than or equal to one whenever  $M > \frac{2N_0}{\delta} e^{4N_0 \nu}$ . This shows - by Cheung [Che11, Theorem 3.3] - that the Hausdorff dimension of  $D$  is at least  $\frac{1}{2}$ . Therefore, the Hausdorff dimension of the set of directions that diverge on average is exactly equal to  $\frac{1}{2}$  by [ASAE<sup>+</sup>17].  $\square$

## 4. PROOF OF PROPOSITION 2

Fix a unit area quadratic or Abelian differential  $(X, \omega)$ . Let  $\beta$  be the core curve of a cylinder  $\mathcal{C}$ , which has area at least  $c$ . Recall that  $N_\beta := \log |\beta|$ , which, for simplicity of notation, we will just denote  $N$  in the sequel. Finally, given two straight line segments  $s$  and  $s'$  in  $(X, \omega)$  we will let  $\theta(s, s')$  denote the angle between them. In the sequel, this angle will always be taken to be positive.

**Step 1: Defining a set of proto-children**

Let  $\sigma$  be at the shortest saddle connection crossing  $\mathcal{C}$ . Let  $\sigma_j$  be  $\sigma$  after  $j$  right Dehn twists have been applied in  $\mathcal{C}$ . We will only consider  $\delta^{-1}N \leq j \leq \frac{M}{2N_0}N$  where  $\delta \in (0, 1)$  is a constant depending only on  $N_0$  and  $c$  that will be determined later. These saddle connections will be called proto-children. We have

$$\frac{c}{2j|\beta|^2} \leq \tan \theta(\sigma_j, \beta) \leq \frac{1}{j|\beta|^2} \quad \text{and} \quad \frac{c}{2j^2|\beta|^2} \leq \sin \theta(\sigma_j, \sigma_{j+1}) \leq \frac{1}{j^2|\beta|^2}.$$

The estimates on the left come from the the area assumption on the cylinder which says that the component of  $\sigma_j$  perpendicular to  $\beta$  is between  $c/|\beta|$  and  $1/|\beta|$  and the fact that for  $|\beta|$  large enough (for example  $|\beta| > 3$ ),

$$j|\beta| \leq |\sigma_j| \leq \sqrt{2}j|\beta|.$$

The estimate on the right comes from the identity.

$$c \leq \text{Area}(\mathcal{C}) = |\sigma_j \times \sigma_{j+1}| \leq 1.$$

**Step 2: There is a collection of cylinders close to the proto-children that are children of  $\beta$** 

We would like to find a cylinder with core curve  $\beta_j$  close in angle to  $\sigma_j$  and satisfying the properties of Lemma 2. To produce  $\beta_j$ , begin by applying a rotation -  $r_{\theta_j}$  - so that  $\sigma_j$  is vertical, and then apply  $g_{t_j}$  for

$$t_j := \log |\sigma_j|$$

so that  $\sigma_j$  has length 1 on  $g_{t_j}r_{\theta_j}(X, \omega)$ .

By Proposition 1, on  $g_{t_j}r_{\theta_j}(X, \omega)$  there is a cylinder whose core curve is  $\beta_j$  with circumference of length at most  $N_0$ , and area at least  $c$ . Denote by  $v_j$  the vertical component of its holonomy and  $h_j$  the horizontal component on  $g_{t_j}r_{\theta_j}(X, \omega)$ .

On  $r_{\theta_j}(X, \omega)$  the vertical component of  $\beta_j$  is  $e^{t_j}v_j$  and the horizontal component is  $e^{-t_j}h_j$ . Thus

$$(4.0.1) \quad \tan \theta(\beta_j, \sigma_j) = e^{-2t_j} \frac{h_j}{v_j}.$$

**Lemma 7.** *For any  $\epsilon > 0$  there is an  $L > 0$  so that if  $|\beta| > L$  then,*

- (1)  $\sin \theta(\beta_j, \sigma_j) \leq \tan \theta(\beta_j, \sigma_j) < \epsilon$  and  
 (2)  $|\frac{|\beta_j|}{e^{t_j} v_j} - 1| < \epsilon$ .

*Proof.* Since  $h_j \leq N_0$ , as  $|\beta| \rightarrow \infty$  we have  $h_j e^{-t_j} \rightarrow 0$ . Now on  $(X, \omega)$  there is a lower bound on the length of any closed curve so  $v_j e^{t_j}$  is bounded away from 0. Together with (4.0.1) this gives (1). Conclusion (2) follows from the fact that  $|\beta_j| = \sqrt{e^{2t_j} v_j^2 + e^{-2t_j} h_j^2}$ .  $\square$

Using Lemma 7 we will suppose that  $|\beta|$  is sufficiently large so that

$$\frac{1}{2} \leq \frac{\sin \theta(\beta_j, \sigma_j)}{\theta(\beta_j, \sigma_j)} \leq \frac{\tan \theta(\beta_j, \sigma_j)}{\theta(\beta_j, \sigma_j)} \leq 2$$

and so that  $|\frac{|\beta_j|}{e^{t_j} v_j} - 1| < \frac{1}{2}$ .

**Lemma 8.** *There exists  $\delta = \delta(c, N_0) > 0$  such that*

- (1) *If  $\max\{h_j, v_j\} < \delta$ , then  $\beta_j$  is disjoint from  $\beta_k$  for  $|k - j| \leq 3g - 3$ .*  
 (2) *If  $\max(v_j, v_k) < \delta$ , then  $\beta_j$  is disjoint from  $\beta_k$  for  $|k - j| \leq 3g - 3$ .*

*Proof.* If  $|k - j| \leq 3g - 3$  and  $|\beta| > 9g - 9$  then,

$$(4.0.2) \quad \frac{1}{2} \leq \frac{e^{2t_k}}{e^{2t_j}} = \frac{|\sigma_k|^2}{|\sigma_j|^2} \leq 2.$$

Therefore,

$$\begin{aligned} \sin \theta(\beta_j, \beta_k) &\leq \sin \theta(\beta_j, \sigma_j) + \sin \theta(\sigma_j, \sigma_k) + \sin \theta(\sigma_k, \beta_k) \\ &\leq e^{-2t_j} \frac{h_j}{v_j} + \frac{|k - j|}{|\sigma_j| |\sigma_k|} + e^{-2t_k} \frac{h_k}{v_k}. \end{aligned}$$

The quantity on the right hand side is therefore bounded above by

$$(4.0.3) \quad 2e^{-2t_j} \left( (3g - 3) + \frac{h_j}{v_j} + \frac{h_k}{v_k} \right).$$

On the other hand suppose  $\beta_j$  and  $\beta_k$  intersect. For simplicity assume that  $\beta_j$  is vertical. Since the area of the cylinder with core curve  $\beta_j$  is at least  $c$ , we have that the distance across the cylinder is bounded below by  $\frac{c}{|\beta_j|} \geq \frac{c}{2v_j e^{t_j}}$ . Thus the horizontal component of  $\beta_k$  is bounded below by  $\frac{ce^{-t_j}}{2v_j}$  and so

$$\sin \theta(\beta_j, \beta_k) \geq \frac{ce^{-t_j}}{2v_j |\beta_k|} \geq \frac{c}{4} \frac{e^{-2t_j}}{v_j v_k}.$$

Comparing this inequality to (4.0.3) we get

$$c \leq 8((3g - 3)v_j v_k + h_j v_k + h_k v_j).$$

By assumption  $h_j, h_k, v_j$ , and  $v_k$  are bounded above by  $N_0$ . Therefore, if  $\max(h_j, v_j)$  or  $\max(v_j, v_k)$  is less than or equal to  $\delta := \frac{c}{32N_0(3g-3)}$ , then we have  $c < c$ , which is a contradiction; and so  $\beta_j$  and  $\beta_k$  must be disjoint.  $\square$

We continue with proof of Proposition 2. On a genus  $g$  translation surface there are at most  $3g - 3$  pairwise disjoint cylinders. Therefore, for any  $j$ , there is some  $\ell \in \{j, j + 1, \dots, j + 3g - 4\}$  with  $v_\ell \geq \delta$ . We ignore the  $j$  with  $v_j \leq \delta$ , which leaves at least proportion  $1/(3g - 3)$  of the original  $\beta_j$  each of which satisfies  $v_j > \delta$ . Considering only these  $j$ , we will now verify that they satisfy the definition of children given in Definition 2.

By construction all of the  $\beta_j$  are core curves of cylinders of area at least  $c$ . The length of each  $\beta_j$  satisfies the following:

$$j\delta|\beta| \leq \delta|\sigma_j| = \delta e^{t_j} \leq |\beta_j| \leq N_0 e^{t_j} = N_0 |\sigma_j| \leq \sqrt{2} N_0 j |\beta|.$$

Since  $j$  lies between  $\delta^{-1}N$  and  $\frac{M}{\sqrt{2}N_0}N$  we see that

$$|\beta| \log |\beta| \leq |\beta_j| \leq M |\beta| \log |\beta|.$$

By (4.0.1), we have

$$\tan \theta(\sigma_j, \beta_j) \leq e^{-2t_j} \frac{N_0}{\delta} = \frac{N_0}{|\sigma_j|^2 \delta} \leq \frac{\delta^{-1} N_0}{(j|\beta|)^2}.$$

This estimate implies that

$$\theta(\beta, \beta_j) \leq \frac{2(\delta^{-1} N_0 + 1)}{(j|\beta|)^2}.$$

The farthest point of  $I_{\beta_j}$  from  $\theta_\beta$  is therefore,

$$\frac{2(\delta^{-1} N_0 + 1)}{(j|\beta|)^2} + \frac{1}{|\beta_j|^2 \log |\beta_j|} \leq \frac{1}{|\beta|^2 \log |\beta|} \left( \frac{2\delta^{-1} N_0 + 3}{\log |\beta|} \right)$$

which shows that  $I_{\beta_j} \subseteq I_\beta$  whenever  $|\beta| > \exp(2\delta^{-1} N_0 + 3)$ . This shows that the  $\beta_j$  are children of  $\beta$  in the sense of Definition 2.

### Step 3: Verifying property (1) of Proposition 2

Set

$$\nu = \frac{4N_0(3g-3)}{c\delta}.$$

Notice that our choice of  $\delta$  implies that  $\nu > 3g - 3$ . Delete more  $\beta_j$  so that any two successive indices differ by at least  $\nu$  and at most  $2\nu$ .

Then for any two successive remaining  $\beta_j, \beta_k$ , with  $k > j$

$$\begin{aligned}
2\theta(\beta_j, \beta_k) &\geq \sum_{i=j}^{k-1} \sin \theta(\sigma_i, \sigma_{i+1}) - \tan \theta(\sigma_j, \beta_j) - \tan \theta(\sigma_k, \beta_k) \\
&\geq \sum_{i=j}^{k-1} \frac{c}{2i^2|\beta|^2} - \frac{2\delta^{-1}N_0}{(j|\beta|)^2} \\
&\geq \frac{1}{(j|\beta|)^2} \left( \frac{c(3g-3)(4N_0)}{2c\delta} - 2\delta^{-1}N_0 \right) \\
&= \frac{2\delta^{-1}N_0(3g-4)}{(j|\beta|)^2} \geq \frac{2C_0}{|\beta_j|^2}
\end{aligned}$$

for some constant  $C_0 = C_0(N_0, c)$ . In particular, these remaining  $\beta_j$  are now distinct. We define  $D_\beta$  to be the set of these remaining children  $\beta_j$ . Notice that property (1) of Proposition 2 - which states that the successive indices are at most  $2\nu$  apart - is automatic.

**Step 4: Verifying property (2) of Proposition 2**

By definition the interval  $I_{\beta_j}$  with center  $\theta_{\beta_j}$  has radius  $\frac{1}{|\beta_j|^2 \log |\beta_j|}$  whereas  $\theta(\beta_j, \beta_k) \geq \frac{C_0}{|\beta_j|^2}$ . The distance between two distinct intervals  $I_{\beta_j}$  and  $I_{\beta_k}$  is at least

$$\frac{1}{|\beta|^2 \log^2 |\beta|} \left( \frac{C_0}{M^2} - \frac{1}{\log |\beta|} \right)$$

Letting  $C := \frac{C_0}{2M^2}$  we see that this distance is at least  $\frac{C}{N_\beta^2 |\beta|^2}$  whenever  $|\beta| > e^C$ , which is property (2).

**Step 5: Verifying property (3) of Proposition 2**

We have

$$|\beta_j| \leq N_0 |\sigma_j| \leq \sqrt{2} N_0 j |\beta|$$

so

$$\frac{|\beta|}{|\beta_j|} \geq \frac{1}{\sqrt{2} N_0 j}$$

which implies since  $j \leq M \log |\beta|$

$$N_{\beta_j} = \log |\beta_j| \leq \log N_0 + \log M + \log \sqrt{2} + \log \log |\beta| + \log |\beta|$$

and this gives

$$\begin{aligned}
\frac{N_\beta}{N_{\beta_j}} &\geq \frac{\log |\beta|}{\log N_0 + \log M + \log \sqrt{2} + \log \log |\beta| + \log |\beta|} \\
&= \frac{1}{1 + \frac{\log M + \log N_0 + \log \sqrt{2} + \log \log |\beta|}{\log |\beta|}}.
\end{aligned}$$

For sufficiently large  $N_\beta = \log |\beta|$  this lower bound is at least  $\frac{1}{\sqrt{2}}$ . Together with the above lower bound on  $\frac{|\beta|}{|\beta_j|}$  gives property (3) of Proposition 2.

## REFERENCES

- [ASAE<sup>+</sup>17] Hamid Al-Saqban, Paul Apisa, Alena Erchenko, Osama Khalil, Shahriar Mirzadeh, and Caglar Uyanik, *Exceptional directions for the Teichmüller geodesic flow and Hausdorff dimension*, 2017, arXiv:1711.10542.
- [CE15] Jon Chaika and Alex Eskin, *Every flat surface is Birkhoff and Osledelets generic in almost every direction*, J. Mod. Dyn. **9** (2015), 1–23. MR 3395258
- [Che11] Yitwah Cheung, *Hausdorff dimension of the set of singular pairs*, Ann. of Math. (2) **173** (2011), no. 1, 127–167. MR 2753601
- [Cho17] Manoj Choudhuri, *On cusp excursions of geodesics and Diophantine approximation*, 2017, arXiv:1703.09002.
- [CM18] Jon Chaika and Howard Masur, *The set of non-uniquely ergodic d-IETs has Hausdorff codimension 1/2*, 2018, arXiv:1801.00770.
- [CRS08] Young-Eun Choi, Kasra Rafi, and Caroline Series, *Lines of minima and Teichmüller geodesics*, Geom. Funct. Anal. **18** (2008), no. 3, 698–754. MR 2438996
- [Dan85] S. G. Dani, *Divergent trajectories of flows on homogeneous spaces and Diophantine approximation*, J. Reine Angew. Math. **359** (1985), 55–89. MR 794799
- [EKZ] Alex Eskin, Maxim Kontsevich, and Anton Zorich, *Sum of Lyapunov exponents of the Hodge bundle with respect to the Teichmüller geodesic flow*, preprint, arXiv 1112.5872 (2012).
- [FLWW09] Ai-Hua Fan, Ling-Min Liao, Bao-Wei Wang, and Jun Wu, *On Khintchine exponents and Lyapunov exponents of continued fractions*, Ergodic Theory Dynam. Systems **29** (2009), no. 1, 73–109. MR 2470627
- [KKLM14] S. Kadyrov, D. Y. Kleinbock, E. Lindenstrauss, and G. A. Margulis, *Singular systems of linear forms and non-escape of mass in the space of lattices*, arXiv:1407.5310.
- [KMS86] Steven Kerckhoff, Howard Masur, and John Smillie, *Ergodicity of billiard flows and quadratic differentials*, Ann. of Math. (2) **124** (1986), no. 2, 293–311. MR 855297 (88f:58122)
- [Mas85] Bernard Maskit, *Comparison of hyperbolic and extremal lengths*, Ann. Acad. Sci. Fenn. Ser. A I Math. **10** (1985), 381–386. MR 802500 (87c:30062)
- [Mas86] Howard Masur, *Closed trajectories for quadratic differentials with an application to billiards*, Duke Math. J. **53** (1986), no. 2, 307–314. MR 850537
- [Mas91] ———, *Hausdorff dimension of divergent Teichmüller geodesics*, Trans. Amer. Math. Soc. **324** (1991), no. 1, 235–254. MR 984857
- [Mas92] ———, *Hausdorff dimension of the set of nonergodic foliations of a quadratic differential*, Duke Math. J. **66** (1992), no. 3, 387–442.

- [Min92] Yair N. Minsky, *Harmonic maps, length, and energy in Teichmüller space*, J. Differential Geom. **35** (1992), no. 1, 151–217. MR 1152229
- [MS91] Howard Masur and John Smillie, *Hausdorff dimension of sets of non-ergodic measured foliations*, Ann. of Math. (2) **134** (1991), no. 3, 455–543. MR 1135877
- [Raf07] Kasra Rafi, *Thick-thin decomposition for quadratic differentials*, Math. Res. Lett. **14** (2007), no. 2, 333–341. MR 2318629
- [Str84] Kurt Strebel, *Quadratic differentials*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 5, Springer-Verlag, Berlin, 1984. MR 743423
- [ZK75] A. N. Zemljakov and A. B. Katok, *Topological transitivity of billiards in polygons*, Mat. Zametki **18** (1975), no. 2, 291–300. MR 0399423
- [Zor06] Anton Zorich, *Flat surfaces*, Frontiers in number theory, physics, and geometry. I, Springer, Berlin, 2006, pp. 437–583.