AN INDECOMPOSABLE CONTINUUM AS SUBPOWER HIGSON CORONA

YUTAKA IWAMOTO

ABSTRACT. In this paper, we study topological properties of the subpower Higson coronas of proper metric spaces and show that the subpower Higson corona of the half open interval with the usual metric is an indecomposable continuum. Continuous surjections from Higson-type coronas onto a Higson-type compactifications of the half open interval are also constructed.

1. Introduction

Let (X, d) be a metric space with a metric d and let $B_d(x, r)$ denote the closed ball of radius r centered at $x \in X$. A metric d on X is called *proper* if all balls $B_d(x, r)$ are compact. The Higson compactification is a compactification defined by the coarse structure of a proper metric space and plays an important role in the large-scale geometry [14]. Also the sublinear Higson compactification is known as a compactification defined by the sublinear coarse structure of a proper metric space [3], [5]. There are several ways to define a Higson type compactification, by a coarse structure [4], [5], [14], by a large scale structure [6], and by a closed subring of the algebra of all continuous bounded real-valued functions [10], [11]. The subpower Higson compactification was introduced in [11] as a compactification defined by a closed subring of the algebra of all continuous bounded real-valued functions. And the asymptotic power dimension was studied in [12] related to the subpower Higson corona.

In this paper, we adopt the definition associated with a closed subrings of the algebra of all continuous bounded real-valued functions and study topological properties of the subpower Higson compactifications and their coronas. It is known that a Higson-type corona νX can be realized by a discrete subspace of a proper metric space X. The Higson corona and the subpower Higson corona contain a copy of the Stone-Čech remainder $\beta \mathbb{N} \setminus \mathbb{N}$ of the natural numbers \mathbb{N} [10], [11]. Also it was shown that the Higson corona of the half open interval $[0, \infty)$ with the usual metric is, like the Stone-Čech remainder $\beta[0,\infty)\setminus[0,\infty)$ [2], an indecomposable continuum [9]. We show that the subpower Higson corona of the half open interval with the usual metric is also an indecomposable continuum. Then we construct continuous surjections from Higson-type coronas onto Higson-type compactifications of the half open interval.

²⁰¹⁰ Mathematics Subject Classification. Primary 54D40; Secondary 54C45, 54D05, 54E40. Keywords: Higson corona; indecomposable continuum; Stone-Čech compactification

2. Basic properties of the subpower Higson compactification

A (not necessarily continuous) function $s: \mathbb{R}_+ \to \mathbb{R}_+$ between the set of positive real numbers is called asymptotically subpower (resp. asymptotically sublinear) if for every $\alpha > 0$ there exists $t_0 > 0$ such that $s(t) < t^{\alpha}$ (resp. $s(t) < \alpha t$) whenever $t > t_0$. The set of all asymptotically subpower functions (resp. all asymptotically sublinear functions, all positive constant functions) is denoted by \mathcal{P} (resp. \mathcal{L}, \mathcal{H}). Since every constant functions are asymptotically subpower and every asymptotically subpower functions are asymptotically sublinear, it follows that $\mathcal{H} \subset \mathcal{P} \subset \mathcal{L}$.

In what follows, a metric space (X, d) is assumed to have a base point x_0 , and the distance $d(x_0, x)$ of $x \in X$ from x_0 is denoted by |x|. For a subset A of X, the diameter of A is denoted by diam_d A, that is, diam_d $A = \sup\{d(x, y) \mid x, y \in A\}$.

Let (X, d) and (Y, ρ) be proper metric spaces. A map $f: X \to Y$ is called *Higson* subpower (resp. Higson sublinear, Higson) provided that

$$\lim_{|x| \to \infty} \operatorname{diam}_{\rho} f(B_d(x, s(|x|))) = 0$$

for each $s \in \mathcal{P}$ (resp. $s \in \mathcal{L}$, $s \in \mathcal{H}$). Hence, a map $f : X \to Y$ is Higson subpower if and only if, given asymptotically subpower function s and $\varepsilon > 0$, there exists a compact subset $K \subset X$ such that $\operatorname{diam}_{\rho} f(B_d(x, s(|x|)) < \varepsilon$ whenever $x \in X \setminus K$.

Let $C^*(X)$ be the set of all bounded real-valued continuous functions on X. For a subset F of $C^*(X)$, let $e_F: X \to \prod_{f \in F} I_f$, where $I_f = [\inf f, \sup f]$, $f \in F$, be the evaluation map of F defined by $(e_F(x))_f = f(x)$ for every $x \in X$. It is known that if $F \subset C^*(X)$ separates points from closed sets, then e_F is an embedding [15, 8.16]. Identifying X with $e_F(X)$, the closure $\overline{e_F(X)}$ of $e_F(X)$ in $\prod_{f \in F} I_f$ gives a compactification of X.

For a proper metric space X, we consider the following three subsets of $C^*(X)$:

$$C_H(X) = \{ f \in C^*(X) \mid f \text{ satisfies } (*)_s^f \text{ for every } s \in \mathcal{H} \},$$

$$C_P(X) = \{ f \in C^*(X) \mid f \text{ satisfies } (*)_s^f \text{ for every } s \in \mathcal{P} \},$$

$$C_L(X) = \{ f \in C^*(X) \mid f \text{ satisfies } (*)_s^f \text{ for every } s \in \mathcal{L} \}.$$

Then they are closed subrings of $C^*(X)$ with respect to the sup-metric. Also, they contain all constant maps and separate points from closed sets. Hence the subrings $C_H(X)$, $C_P(X)$ and $C_L(X)$ determine compactifications of X. Since $\mathcal{H} \subset \mathcal{P} \subset \mathcal{L}$, it follows that $C_L(X) \subset C_P(X) \subset C_H(X)$ (cf. [11]).

Let c_1X and c_2X be compactifications of X. We say $c_1X \succeq c_2X$ provided that there is a continuous map $f: c_1X \to c_2X$ such that $f|_X = \mathrm{id}_X$. We note that a continuous map $f: c_1X \to c_2X$ with $f|_X = \mathrm{id}_X$ is unique and surjective. If $c_1X \preceq c_2X$ and $c_1X \succeq c_2X$, then we say that c_1X and c_2X are equivalent.

The Higson compactification $h_H(X)$, the subpower Higson compactification $h_P(X)$ and the sublinear Higson compactification $h_L(X)$ are compactifications defined by closed subrings $C_H(X)$, $C_P(X)$ and $C_L(X)$ respectively, that is, $h_A(X)$ is equivalent to $\overline{e_{C_A}(X)}$, where $A \in \{H, P, L\}$ and $C_A = C_A(X)$. Their coronas are defined by $\nu_A X = h_A(X) \setminus X$ for each $A \in \{H, P, L\}$. These three compactifications are referred to as the Higson type compactifications. We note that $h_L(X) \leq h_P(X) \leq h_H(X)$ since $C_L(X) \subset C_P(X) \subset C_H(X)$.

One of the basic property of the Higson type compactifications is the following:

(\sharp) A bounded continuous map $f: X \to \mathbb{R}$ has an extension $\hat{f}: h_A(X) \to \mathbb{R}$ if and only if $f \in C_A(X)$,

where $A \in \{H, P, L\}$. This condition holds for any closed subring of $C^*(X)$ which contains all constant maps and separates points from closed sets [7, Problem 3.12.22(e)]. See [1] for a comprehensive and detailed description of this property.

In this section, we shall derive some basic properties concerning the Higson subpower compactifications using the basic ideas of [4], [5] and [10].

The following proposition is a fundamental property of the Higson type compactification which is derived from the property (\natural) and can be shown by a similar argument to that in [10, Proposition 1].

Proposition 2.1 (cf. [10]). Let X be a proper metric space. Then the subpower (resp. sublinear) Higson compactification is the unique compactification of X such that if Y is any compact metric space and $f: X \to Y$ is continuous, then f has a continuous extension $\hat{f}: h_P(X) \to Y$ (resp. $\hat{f}: h_L(X) \to Y$) if and only if f satisfies $(*)_s^f$ for any $s \in \mathcal{P}$ (resp. for any $s \in \mathcal{L}$).

Let (X,d) be a metric space. We call a finite system E_1, \ldots, E_n of closed subsets of X diverges as a power function if there exist $\alpha > 0$ and $r_0 > 0$ such that $\max\{d(x, E_i) \mid 1 \leq i \leq n\} \geq |x|^{\alpha}$ whenever $|x| > r_0$. If a finite system E_1, \ldots, E_n diverge as a power function then $(\bigcap_{i=1}^n E_i) \cap (X \setminus B_d(x_0, r)) = \emptyset$ for some r > 0. As in the case of Higson compactification [4, Lemma 2.2], we have the following:

Lemma 2.2 (cf. [4]). Let (X,d) be a non-compact proper metric space. Let E_1, \ldots, E_n be a finite system of closed subsets of X such that $\bigcap_{i=1}^n E_i = \emptyset$. Let $f_i: X \to \mathbb{R}_+$ be the map defined by $f_i(x) = d(x, E_i)$, $1 \le i \le n$ and let $F = \sum_{i=1}^n f_i$. If the system E_1, \ldots, E_n diverges as a power function then the map $g_i = f_i/F : X \to [0,1]$ is Higson subpower.

Proof. First we note that the well-definedness of g_i follows from the assumption $\bigcap_{i=1}^n E_i = \emptyset$. Let $\alpha, r_1 > 0$ be positive numbers such that $\max\{d(x, E_i) \mid 1 \le i \le n\} \ge |x|^{2\alpha}$ whenever $|x| > r_1$. Then $F(x) \ge |x|^{2\alpha}$ whenever $|x| > r_1$. Let $s: \mathbb{R}_+ \to \mathbb{R}_+$ be an asymptotically subpower function. Then there exists $r_2 > 0$ such that $s(t) < t^{\alpha}$ for every $t > r_2$. Given $\varepsilon > 0$, we can take $r_3 > 0$ so that $(n+1)/t^{\alpha} < \varepsilon/2$ for every $t > r_3$.

Put $r_0 = \max\{r_1, r_2, r_3\}$. Let x be a point with $|x| > r_0$ and let $y \in B_d(x, s(|x|))$. Then we have

$$|g_{i}(x) - g_{i}(y)| = \left| \frac{f_{i}(x)}{F(x)} - \frac{f_{i}(y)}{F(y)} \right| = \left| \frac{f_{i}(x) - f_{i}(y)}{F(x)} + \frac{f_{i}(y)}{F(y)} \cdot \frac{F(y) - F(x)}{F(x)} \right|$$

$$\leq \left| \frac{d(x,y)}{F(x)} \right| + \left| \frac{F(y) - F(x)}{F(x)} \right| \leq \frac{(n+1)s(|x|)}{F(x)}$$

$$\leq \frac{(n+1)s(|x|)}{|x|^{2\alpha}} < \frac{(n+1)|x|^{\alpha}}{|x|^{2\alpha}} = \frac{(n+1)}{|x|^{\alpha}} < \frac{\varepsilon}{2}.$$

Hence, diam $g_i(B_d(x, s(|x|))) \le \varepsilon$ whenever $|x| > r_0$. Thus g_i is Higson subpower for every $1 \le i \le n$.

For a subset A of a proper metric space X, \overline{A} denotes the closure of A in the subpower Higson compactification h_PX of X. As in case of the Higson corona [4, Proposition 2.3] and of the sublinear Higson corona [5, Lemma 2.3], we have the following.

Proposition 2.3 (cf. [4], [5]). Let (X, d) be a non-compact proper metric space. For a system E_1, \ldots, E_n of closed subsets of X, the following conditions are equivalent:

- $(1) \ \nu_P X \cap \left(\cap_{k=1}^n \overline{E_k} \right) = \emptyset,$
- (2) the system E_1, \ldots, E_n diverges as a power function.

Proof. Suppose that the condition (2) does not hold. Then there exists a sequence $(x_i)_{i=1}^{\infty}$ in X such that $\lim_{i\to\infty}|x_i|=\infty$, $|x_i|<|x_{i+1}|$ and $d(x_i,E_k)<|x_i|^{1/i}$ for every $i\in\mathbb{N}$ and $1\leq k\leq n$. Let p be a cluster point of the sequence $(x_i)_{i=1}^{\infty}$. Then $p\in\nu_P X$. Put $t_i=|x_i|$ for each $i\in\mathbb{N}$. Let $s:\mathbb{R}_+\to\mathbb{R}_+$ be the (non-continuous) function defined by $s(t)=|x_i|^{1/i}$ when $t\in[t_i,t_{i+1})$ and s(t)=1 when $t\in[0,t_1)$. It is easy to see that s is an asymptotically subpower function and $d(x_i,E_k)< s(|x_i|)$ for every $i\in\mathbb{N}$ and $1\leq k\leq n$. Hence, if f is any element of $C_P(X)$, the condition $(*)_s^f$ implies that the distance between $f(x_i)$ and $f(E_k)$ tends to zero as i tends to infinity. Considering the evaluation map $e_{C_P(X)}: X\to\prod_{f\in C_P(X)}I_f$, it follows from this fact that $p\in\overline{E_k}$ for every $1\leq k\leq n$. Thus $p\in\nu_P X\cap(\cap_{k=1}^n\overline{E_k})\neq\emptyset$.

Suppose that the system E_1,\ldots,E_n diverges as a power function. Then we have $(\cap_{k=1}^n E_k) \cap (X \setminus B_d(x_0,r)) = \emptyset$ for some r>0. Put $F_k = \operatorname{cl}_X(E_k \setminus B_d(x_0,r+1))$ for each $k=1,\ldots,n$. Then the system F_1,\ldots,F_n diverges as a power function and $\cap_{k=1}^n F_k = \emptyset$. Note that $\overline{F_k} \cap \nu_P X = \overline{E_k} \cap \nu_P X$ for every $1 \le k \le n$. By Lemma 2.2 there exist Higson subpower maps $g_k : X \to [0,1], 1 \le k \le n$, such that $F_k \subset g_k^{-1}(0)$ and $\Sigma_{k=1}^n g_k = 1$. By Proposition 2.1, there exists an extension $G_k : h_P X \to [0,1]$ of g_k . Then $\Sigma_{k=1}^n G_k = 1$ and $\overline{F_k} \subset G_k^{-1}(0)$. However, the condition $\Sigma_{k=1}^n G_k = 1$ leads that $\nu_P X \cap (\cap_{k=1}^n \overline{E_k}) \subset \cap_{k=1}^n \overline{F_k} \subset \cap_{k=1}^n G_k^{-1}(0) = \emptyset$.

We note that the implication $(1) \Rightarrow (2)$ for the case n = 2 of the above proposition was in fact shown by Kucab and Zarichnyi in [13].

Proposition 2.4 (cf. [10]). Let X be a proper metric space and let Y be a closed subset of X with the induced metric from X. Then the closure \overline{Y} of Y in $h_P(X)$ is equivalent to the subpower Higson compactification $h_P(Y)$ of Y.

Proof. By [7, Theorem 3.5.5], it suffices to show $\operatorname{cl}_{h_P(X)}A \cap \operatorname{cl}_{h_P(X)}B = \emptyset$ if and only if $\operatorname{cl}_{h_P(Y)}A \cap \operatorname{cl}_{h_P(Y)}B = \emptyset$ for every pair A, B of closed subsets of Y. It follows from Proposition 2.3 that the condition $\operatorname{cl}_{h_P(X)}A \cap \operatorname{cl}_{h_P(X)}B = \emptyset$ if and only if the system A, B diverges as a power function in X and $A \cap B = \emptyset$. This is equivalent to the condition that the system A, B diverges as a power function in Y and $A \cap B = \emptyset$ since the metric on Y is inherited from X.

Let (X,d) be a metric space. For any R>0, a subset Y of X is called R-dense in X if $B_d(x,R)\cap Y\neq\emptyset$ for every $x\in X$.

Corollary 2.5 (cf. [10]). Let (X, d) be a proper metric space and let Y be a subset of X. If Y is R-dense in X for some R > 0 then $\overline{Y} \setminus Y = \nu_P X$. In particular, $\nu_P Y$ is homeomorphic to $\nu_P X$.

Proof. Let $x \in \nu_P X$. Let U and V be open neighborhoods of x in $h_P(X)$ such that $x \in V \subset \overline{V} \subset U$. Put $E = X \cap (h_P(X) \setminus U)$ and $F = X \cap \overline{V}$. By Proposition 2.3, the system E, F diverges as a power function since $\nu_P X \cap \overline{E} \cap \overline{F} = \emptyset$. Thus, there are $\alpha, r > 0$ such that $d(y, E) > |y|^{\alpha}$ whenever $y \in F$ and |y| > r. Since $V \cap \nu_P X \neq \emptyset$ and X is a proper metric space, we can take $z \in F$ so that |z| > r.

We may assume that $R < r^{\alpha}$. Then $B_d(z,R) \cap E = \emptyset$, i.e., $B_d(z,R) \subset U$. Since Y is R-dense in $X, \emptyset \neq Y \cap B_d(z,R) \subset Y \cap U$. This fact leads that $x \in \overline{Y}$ since we can take U as an arbitrarily small neighborhood of x. Hence $\nu_P X$ is contained in $\overline{Y} \setminus Y$, i.e., $\overline{Y} \setminus Y = \nu_P X$. By Proposition 2.4, $\overline{Y} \setminus Y$ is homeomorphic to $\nu_P Y$. Thus $\nu_P Y$ is homeomorphic to $\nu_P X$.

3. Continua as subpower Higson coronas of $[0, \infty)$

Throughout this section, 0 is assumed to be the base point of the half-open interval $[0,\infty)$, that is, $x_0=0\in[0,\infty)$. A metric d on $[0,\infty)$ defined by d(x,y)= $\sqrt{(x-y)^2}$ is called the usual metric on $[0,\infty)$. For notational simplicity, we identify t with $|t| = d(x_0, t)$ for every $t \in [0, \infty)$ when we consider the usual metric on $[0,\infty)$. A continuum is a nonempty, compact and connected Hausdorff space. A subcontinuum is a continuum which is a subset of a continuum. A continuum is called *decomposable* if it can be represented as the union of two of its proper subcontinua. A continuum which is not decomposable is said to be *indecomposable*.

Let $X = [0, \infty)$ be the half open interval with the usual metric. The indecomposability of the Stone-Čech remainder $\beta X \setminus X$ and the Higson corona $\nu_H X$ was proved in [2] and in [9] respectively. We show here that the subpower Higson corona $\nu_P X$ is also an indecomposable continuum. Let K be a proper closed subset of $\beta X \setminus X$ with non-empty interior in $\beta X \setminus X$. To see the indecomposability of $\beta X \setminus X$, Bellamy [2] constructed a continuous map $f: X \to [0,1]$ such that $f^*(K) = \{0,1\}$, where $f^*: \beta X \to [0,1]$ is the extension of f. Hence K is disconnected. We note that such the extension f^* to the Stone-Čech compactification always exists since f is a bounded continuous map. And then he constructed a continuous surjection from a given non-degenerate subcontinuum of $\beta X \setminus X$ onto a given metric continuum. Our strategy is essentially the same as the Bellamy's. To see the indecomposability of $\nu_P X$, we will construct a map $\psi: X \to [0,1]$ such that $\psi^*(K') = \{0,1\}$ for a given proper closed subset K' of $\nu_P X$ with non-empty interior in $\nu_P X$, where $\psi^*: h_P X \to [0,1]$ is the extension of ψ . In order to ensure that a map $\psi: X \to [0,1]$ has the extension $\psi^*: h_P X \to [0,1]$, we will carefully construct ψ to be Higson subpower (cf. Proposition 2.1). Then we will construct a continuous surjection from $\nu_P X$ onto the Higson type compactification $h_A(X)$ for each $A \in \{H, P, L\}$.

The following lemma plays an essential role to analyze the subpower Higson corona of the half-open interval $[0, \infty)$.

Lemma 3.1. Let $X = [0, \infty)$ be the half open interval with the usual metric d. Let U and V be non-empty disjoint open subsets of $h_P(X)$ such that $U \cap \nu_P X \neq \emptyset$ and $V \cap \nu_P X \neq \emptyset$. Then there exist a natural number $k \geq 3$ and sequences $(a_n)_{n=1}^{\infty}$ and $(b_n)_{n=1}^{\infty}$ with $0 = b_0 < a_1 < b_1 < \cdots < b_{n-1} < a_n < b_n < \cdots$ satisfying the following conditions:

- (1) $(a_n)^{1/k} > 2^n$,
- (2) $b_{n-1} + (b_{n-1})^{1/k} < a_n$,
- (3) $[b_{n-1}, a_n] \cap U \neq \emptyset$,
- $(4) b_n = a_n + (a_n)^{2/k},$
- (5) $[a_n, b_n] \subset V \cap X$,
- (6) $a_n b_{n-1} > 2^{n-1} (a_n)^{1/k},$ (7) $b_n a_n > 2^n (b_n)^{1/k}.$

Proof. Let z be a point in $V \cap \nu_P X$ and let W be an open neighborhood of z in $h_P X$ such that $z \in W \subset \overline{W} \subset V$. Then

$$\nu_P X \cap \overline{(X \setminus (V \cap X))} \cap \overline{(\overline{W} \cap X)} \subset \nu_P X \cap (h_P(X) \setminus V) \cap \overline{W} = \emptyset.$$

By Proposition 2.3, the system $X \setminus (V \cap X)$, $\overline{W} \cap X$ diverges as a power function. Hence, there exist $\alpha > 0$ and $r_0 > 0$ such that

$$d(t, X \setminus (V \cap X)) > t^{\alpha}$$
 whenever $t \in \overline{W} \cap X$ and $t > r_0$.

Take a natural number $k \geq 3$ so that $\alpha > 2/k$. Then the condition above implies that

(A)
$$B_d(t, t^{2/k}) \subset V \cap X$$
 whenever $t \in \overline{W} \cap X$ and $t \geq r_0$.

Then we consider the following two inequalities:

(B)
$$\begin{cases} x^k > x, \\ x^2 > 2^k (x + x^{2/k}). \end{cases}$$

Since $k \geq 3$, there is $t_1 > 2$ such that x satisfies (B) whenever $x > t_1$. Then we take $a_1 \in W \cap X$ so that $a_1 > \max\{r_0, (t_1)^k\}$. This is possible since the set $W \cap X$ is cofinal in X by the condition $W \cap \nu_P X \neq \emptyset$. Hence, a_1 satisfies the conditions (1) and (2). Also, the condition $U \cap \nu_P X \neq \emptyset$ allows us to make a_1 satisfy the condition (3). Then we define b_1 by (4). Since $a_1 \in \overline{W} \cap X$ and $d(a_1, b_1) = (a_1)^{2/k}$, we have $b_1 \in B_d(a_1, (a_1)^{2/k}) \subset V \cap X$ by (A). Then the condition (5) follows from the fact that the metric d is geodesic. To see that a_1 and b_1 satisfy the conditions (6) and (7), put $a_1 = x$. Then $b_1 = x + x^{2/k}$ by (4). Since $a_1 > t_1$, we have $(a_1 - b_0)^k = x^k > x = ((a_1)^{1/k})^k$ and $(b_1 - a_1)^k = x^2 > 2^k(x + x^{2/k}) = (2(b_1)^{1/k})^k$ by (B). Thus the conditions (6) and (7) are satisfied.

Suppose that $b_{i-1} < a_i < b_i$ have been constructed for i < n. Then we consider the following two inequalities:

(C)
$$\begin{cases} (x - b_{n-1})^k > 2^{(n-1)k}x, \\ x^2 > 2^{nk}(x + x^{2/k}). \end{cases}$$

Since $k \geq 3$, there exists $t_n > 2^n$ such that x satisfies (C) whenever $x > t_n$. As in the first step, we can take $a_n \in W \cap X$ with $a_n > \max\{r_0, (t_n)^k\}$ so that the conditions (1)–(3) are satisfied. Then we define b_n by (4). The condition (5) follows from (A) and the fact that the metric d is geodesic. Finally, to see the conditions (6) and (7), we put $a_n = x$. Then $b_n = x + x^{2/k}$ by (4). Since $a_n > t_n$, it follows from (C) that $(a_n - b_{n-1})^k = (x - b_{n-1})^k > 2^{(n-1)k}x = (2^{n-1}(a_n)^{1/k})^k$ and $(b_n - a_n)^k = x^2 > 2^{nk}(x + x^{2/k}) = (2^n(b_n)^{1/k})^k$. The conditions (6) and (7) are satisfied.

Proposition 3.2. Let $X = [0, \infty)$ be the half open interval with the usual metric d. If K is a proper closed subset of $\nu_P X$ with non-empty interior in $\nu_P X$ then K is disconnected.

Proof. Let $x \in \operatorname{Int}_{\nu_P X} K$ and $y \in \nu_P X \setminus K$. Let U be an open neighborhood of x in $h_P X$ such that $\overline{U} \cap \nu_P X \subset K$. Since $y \notin \overline{U}$, we can take an open neighborhood V of y in $h_P X$ such that $y \in V \subset \overline{V} \subset h_P X \setminus (K \cup \overline{U})$. By Lemma 3.1, there exist a natural number $k \geq 3$ and sequences $(a_n)_{n=1}^{\infty}$ and $(b_n)_{n=1}^{\infty}$ with $0 = b_0 < a_1 < b_1 < \dots < b_{n-1} < a_n < b_n < \dots$ satisfying the following conditions:

- (1) $(a_n)^{1/k} > 2^n$,
- (2) $b_{n-1} + (b_{n-1})^{1/k} < a_n$,
- $(3) [b_{n-1}, a_n] \cap U \neq \emptyset,$
- (4) $b_n = a_n + (a_n)^{2/k}$
- (5) $[a_n, b_n] \subset V \cap X$,
- (6) $a_n b_{n-1} > 2^{n-1} (a_n)^{1/k},$ (7) $b_n a_n > 2^n (b_n)^{1/k}.$

Claim 3.2.1. Let $u \in B_d(t, t^{1/k})$. Then we have the following:

- (i) If $t \in [a_n, b_n)$ then $u \in (b_{n-1}, a_{n+1})$.
- (ii) If $t \in [b_n, a_{n+1})$ then $u \in (a_n, b_{n+1})$.

Proof of Claim 3.2.1. Suppose that $t \in [a_n, b_n)$. If u < t then $u \ge t - t^{1/k} \ge a_n - (a_n)^{1/k} \ge a_n - 2^{n-1}(a_n)^{1/k} > b_{n-1}$ by the monotone increasing-property of the correspondence $s \mapsto s - s^{1/k}$ on $(1, \infty)$ and (6). If t < u then $u \le t + t^{1/k} < t$ $b_n + (b_n)^{1/k} < a_{n+1}$ by (2). Thus the condition (i) is satisfied.

Next suppose that $t \in [b_n, a_{n+1})$. If u < t then, as above, using (7), $u \ge t - t^{1/k} \ge b_n - (b_n)^{1/k} > b_n - 2^n (b_n)^{1/k} > a_n$. If u > t then $u \le t + t^{1/k} < a_{n+1} + (a_{n+1})^{1/k} < a_{n+1} + (a_{n+1})^{2/k} = b_{n+1}$ by (4). Thus the condition (ii) is satisfied. \diamondsuit

We define $\psi: X \to [0,1]$ by

$$\psi(t) = \begin{cases} 0, & \text{if } b_{2n-2} \le t < a_{2n-1}, \\ \frac{t - a_{2n-1}}{b_{2n-1} - a_{2n-1}}, & \text{if } a_{2n-1} \le t < b_{2n-1}, \\ 1, & \text{if } b_{2n-1} \le t < a_{2n}, \\ \frac{b_{2n} - t}{b_{2n} - a_{2n}}, & \text{if } a_{2n} \le t < b_{2n}. \end{cases}$$

Then it is easy to see that the function ψ is well-defined and continuous. Now we shall evaluate the diameter of $\psi(B_d(t,t^{1/k}))$.

Claim 3.2.2. If $a_n \le t < a_{n+1}$ then $|\psi(t) - \psi(u)| < 1/2^n$ for every $u \in B_d(t, t^{1/k})$.

Proof of Claim 3.2.2. Let $u \in B_d(t, t^{1/k})$. By the definition of ψ , we check the claim by dividing it into four cases, $t \in [a_{2n-1}, b_{2n-1}), t \in [b_{2n-1}, a_{2n}), t \in [a_{2n}, b_{2n})$ and $t \in [b_{2n}, a_{2n+1})$. We show here the first two cases. The other two cases can be shown in a similar fashion.

Suppose that $t \in [a_{2n-1}, b_{2n-1})$. By Claim 3.2.1, it suffices to consider the three cases that $u \in (b_{2n-2}, a_{2n-1}), u \in [a_{2n-1}, b_{2n-1})$ and $u \in [b_{2n-1}, a_{2n})$. If $u \in (b_{2n-2}, a_{2n-1})$ then, using (7),

$$\begin{split} |\psi(t)-\psi(u)| &= \psi(t) = \frac{t-a_{2n-1}}{b_{2n-1}-a_{2n-1}} \leq \frac{t-u}{b_{2n-1}-a_{2n-1}} \\ &\leq \frac{t^{1/k}}{b_{2n-1}-a_{2n-1}} < \frac{(b_{2n-1})^{1/k}}{b_{2n-1}-a_{2n-1}} \\ &< \frac{(b_{2n-1})^{1/k}}{2^{2n-1}(b_{2n-1})^{1/k}} < \frac{1}{2^{2n-1}}. \end{split}$$

If $u \in [a_{2n-1}, b_{2n-1})$ then we have

$$|\psi(t) - \psi(u)| = \frac{|t - u|}{b_{2n-1} - a_{2n-1}} \le \frac{t^{1/k}}{b_{2n-1} - a_{2n-1}} < \frac{1}{2^{2n-1}}.$$

If $u \in [b_{2n-1}, a_{2n})$ then we have

$$|\psi(t) - \psi(u)| = 1 - \psi(t) = 1 - \frac{t - a_{2n-1}}{b_{2n-1} - a_{2n-1}} = \frac{b_{2n-1} - t}{b_{2n-1} - a_{2n-1}}$$

$$\leq \frac{u - t}{b_{2n-1} - a_{2n-1}} \leq \frac{t^{1/k}}{b_{2n-1} - a_{2n-1}} < \frac{1}{2^{2n-1}}.$$

Thus $|\psi(t) - \psi(u)| < 1/2^{2n-1}$ whenever $t \in [a_{2n-1}, b_{2n-1})$.

Next suppose $t \in [b_{2n-1}, a_{2n})$. By Claim 3.2.1, it suffices to consider the three cases that $u \in (a_{2n-1}, b_{2n-1}), \ u \in [b_{2n-1}, a_{2n}) \ \text{and} \ u \in [a_{2n}, b_{2n})$. Let $u \in (a_{2n-1}, b_{2n-1})$. Since $u \ge t - t^{1/k} \ge b_{2n-1} - (b_{2n-1})^{1/k}$, it follows that

$$(\dagger) b_{2n-1} - u \le (b_{2n-1})^{1/k}.$$

So, using (7) and (\dagger) , we have

$$|\psi(t) - \psi(u)| = 1 - \psi(u) = 1 - \frac{u - a_{2n-1}}{b_{2n-1} - a_{2n-1}} = \frac{b_{2n-1} - u}{b_{2n-1} - a_{2n-1}}$$

$$\leq \frac{(b_{2n-1})^{1/k}}{b_{2n-1} - a_{2n-1}} < \frac{(b_{2n-1})^{1/k}}{2^{2n-1}(b_{2n-1})^{1/k}} < \frac{1}{2^{2n-1}}.$$

If $u \in [b_{2n-1}, a_{2n})$ then $|\psi(t) - \psi(u)| = 1 - 1 = 0$. Finally, let $u \in [a_{2n}, b_{2n})$. Then, using (4) and (1), we have

$$|\psi(t) - \psi(u)| = 1 - \psi(u) = 1 - \frac{b_{2n} - u}{b_{2n} - a_{2n}} = \frac{u - a_{2n}}{b_{2n} - a_{2n}}$$

$$< \frac{u - t}{b_{2n} - a_{2n}} \le \frac{t^{1/k}}{b_{2n} - a_{2n}} = \frac{t^{1/k}}{(a_{2n})^{2/k}}$$

$$< \frac{(a_{2n})^{1/k}}{(a_{2n})^{2/k}} = \frac{1}{(a_{2n})^{1/k}} < \frac{1}{2^{2n}} < \frac{1}{2^{2n-1}}.$$

 \Diamond

Thus $|\psi(t) - \psi(u)| < 1/2^{2n-1}$ whenever $t \in [b_{2n-1}, a_{2n})$.

Claim 3.2.3. The map $\psi: X \to [0,1]$ is Higson subpower.

Proof of Claim 3.2.3. Let $\varepsilon > 0$ and let $s : X \to \mathbb{R}_+$ be an asymptotically subpower function. Then there exists r > 0 such that $s(t) < t^{1/k}$ for every t > r. Let m be a natural number such that $1/2^m < \varepsilon/2$. Put $t_0 = \max\{r, a_m\}$. Then, for every $t > t_0$ and $u \in B_d(t, t^{1/k})$, we have $B_d(t, s(t)) \subset B_d(t, t^{1/k})$ and $|\psi(t) - \psi(u)| < 1/2^m$ by Claim 3.2.2. Thus diam $\psi(B_d(t, s(t))) \le \dim \psi(B_d(t, t^{1/k})) \le 2/2^m < \varepsilon$ for every $t > t_0$. Hence ψ is Higson subpower.

By Claim 3.2.3 and Proposition 2.1, there exists an extension $\Psi: h_P X \to [0,1]$ of ψ . By (3), we obtain a sequence $(c_n)_{n=1}^{\infty}$ such that $b_n < c_n < a_{n+1}$ and $c_n \in U$. Then $\psi(c_{2n}) = 0$ and $\psi(c_{2n-1}) = 1$ for every n. Let z_0 and z_1 be cluster points of the sequences $(c_{2n})_{n=1}^{\infty}$ and $(c_{2n-1})_{n=1}^{\infty}$ respectively. Then both z_0 and z_1 are contained in $\overline{U} \cap \nu_P X \subset K$. It follows from the continuity of Ψ that $\Psi(z_0) = 0$ and $\Psi(z_1) = 1$. Hence $z_0 \in \Psi^{-1}(0) \cap K \neq \emptyset$ and $z_1 \in \Psi^{-1}(1) \cap K \neq \emptyset$. However, $\Psi^{-1}((0,1)) \cap K = \emptyset$. Indeed, if there exists $z \in K$ such that $\Psi(z) = t \in (0,1)$ then z has a neighborhood $W \subset h_P X$ such that $\Psi(W) \subset (0,1)$. Since $K \cap \overline{V} = \emptyset$, we may assume $W \cap V = \emptyset$. Then we have $\Psi(W \cap X) = \psi(W \cap X) \subset (0,1)$, i.e., $W \cap X \subset \psi^{-1}((0,1))$. On the other hand, $\psi^{-1}((0,1)) \subset V \cap X$ by (5) and our

construction of ψ . So we have $\emptyset \neq W \cap X \subset V \cap X$ which implies that $W \cap V \neq \emptyset$, a contradiction. Thus $\Psi(K) = \{0,1\}$, i.e., K is disconnected.

As we have seen, the function $\psi: X \to [0,1]$ constructed above is Higson subpower. However it is not Higson sublinear. Indeed, take sequences $(a_n)_{n=1}^{\infty}$, $(b_n)_{n=1}^{\infty}$ and $k \geq 3$ as in Proposition 3.2. Let $\xi: \mathbb{R}_+ \to \mathbb{R}_+$ be the function defined by $\xi(t) = t^{2/k}$. Since $k \geq 3$, it follows that ξ is an asymptotically sublinear function. Since $b_n = a_n + (a_n)^{2/k}$, the closed ball $B_d(a_n, \xi(a_n))$ contains at least two points a_n and b_n . By our construction of ψ , we have $(\psi(a_n), \psi(b_n)) \in \{(0,1), (1,0)\}$, i.e., diam $\psi(B_d(a_n, \xi(a_n))) = 1$ for every n. Hence, ψ is not Higson sublinear.

Theorem 3.3. Let $X = [0, \infty)$ be the half open interval with the usual metric. Then the subpower Higson corona $\nu_P X$ is a non-metrizable indecomposable continuum

Proof. Let $A_n = [n, \infty) \subset X$. Then $\overline{A_n}$ is a continuum and $\overline{A_n} \supset \overline{A_{n+1}}$ for every $n \in \mathbb{N}$. Note that $\nu_P X = \bigcap_{n=1}^{\infty} \overline{A_n}$. Thus $\nu_P X$ is a continuum as the intersection of the decreasing sequence $(\overline{A_n})_{n=1}^{\infty}$ of continua. Non-metrizability of $\nu_P X$ follows from the fact that $\nu_P X$ contains a copy of $\beta \mathbb{N} \setminus \mathbb{N}$ [11] which has the cardinality $2^{\mathfrak{c}}$ [8]. Finally, if $\nu_P X$ is expressed as the union of two non-degenerate proper closed subsets K and L of $\nu_P X$ then both of K and L must have non-empty interiors in $\nu_P X$. Thus both K and L are disconnected by Proposition 3.2.

The following is an example, given in [9], of a proper metric on $[0, \infty)$ which derives a decomposable Higson corona. We show that the same example also derives a decomposable subpower Higson corona.

Example 3.4. There exists a proper metric ρ on the half open interval $[0, \infty)$ such that the Higson subpower corona is a decomposable continuum. Indeed, let $f:[0,\infty)\to\mathbb{R}^2$ be the embedding given by $f(t)=(t,t\sin t)$ and let $X=f([0,\infty))$. Let ρ be the metric on X inherited from \mathbb{R}^2 . We shall show that the Higson subpower corona $\nu_P X$ of (X,ρ) is a decomposable continuum. Put

$$\begin{split} Y &= \{(x,y) \in X \mid y \ge 0\}, \\ Z &= \{(x,y) \in X \mid y \le 0\}, \\ A &= \{(x,y) \in \mathbb{R}^2 \mid x \ge 0, \ |y| \le x\}, \\ B &= \{(x,y) \in A \mid y \ge 0\}, \\ C &= \{(x,y) \in A \mid y \le 0\}. \end{split}$$

Then $X = Y \cup Z$, $A = B \cup C$, $Y \subset B$ and $Z \subset C$. For each $A \subset X$, \overline{A} denotes the closure of A in the subpower Higson compactification $h_P(X)$ of X.

First we note that

$$\nu_P X = \overline{X} \setminus X = (\overline{Y} \setminus Y) \cup (\overline{Z} \setminus Z).$$

Indeed, $\overline{X} \setminus X \supset (\overline{Y} \cup \overline{Z}) \setminus (Y \cup Z) = (\overline{Y} \setminus Y) \cup (\overline{Z} \setminus Z)$. If $z \in \overline{X} \setminus X$ has an open neighborhood U with $U \cap Y = \emptyset$ then every neighborhood of z must intersect with Z by the density of X in \overline{X} , i.e., $z \in \overline{Z} \setminus Z$.

Next we shall show that both $\overline{Y} \setminus Y$ and $\overline{Z} \setminus Z$ are subcontinua of $\nu_P X$. By Proposition 2.4, $\overline{Y} \setminus Y$ is homeomorphic to $\nu_P Y$ (notation: $\overline{Y} \setminus Y \approx \nu_P Y$) and $\overline{Z} \setminus Z \approx \nu_P Z$. It is easy to see that Y is R-dense in B for some R > 0. Hence $\nu_P Y$ is homeomorphic to $\nu_P B$ by Corollary 2.5. Thus, we have $\overline{Y} \setminus Y \approx \nu_P B$.

Similarly, we have $\overline{Z} \setminus Z \approx \nu_P C$. For each $n \in \mathbb{N}$, let $B_n = \{(x,y) \in B \mid x \geq n\}$. Then $\nu_P B = \bigcap_{i=1}^{\infty} \operatorname{cl}_{\nu_P B}(B_n)$. Hence $\nu_P B$ is a continuum as the intersection of the decreasing sequence $(\operatorname{cl}_{\nu_P B}(B_n))_{n=1}^{\infty}$ of continua. Similarly, it follows that $\nu_P C$ is a continuum. Thus, both $\overline{Y} \setminus Y$ and $\overline{Z} \setminus Z$ are subcontinua of $\nu_P X$.

Finally, we shall show that both $\overline{Y} \setminus Y$ and $\overline{Z} \setminus Z$ are proper subcontinua of $\nu_P X$. Let $D = \{ f(t) \mid t = \pi/2 + 2n\pi, n \in \mathbb{N} \}$ and $E = \{ f(t) \mid t = 3\pi/2 + 2n\pi, n \in \mathbb{N} \}.$ Then $\overline{Y} \setminus Y \supset \overline{D} \setminus D \neq \emptyset$. It is easy to see that the systems D, Z diverges as a power function. Hence we have $\overline{D} \cap \overline{Z} \cap \nu_P X = \emptyset$ by Proposition 2.3, i.e., $(\overline{Y}\backslash Y)\backslash(\overline{Z}\backslash Z)\supset\overline{D}\backslash D\neq\emptyset$. Similarly, we have $(\overline{Z}\backslash Z)\backslash(\overline{Y}\backslash Y)\supset\overline{E}\backslash E\neq\emptyset$. Thus, both $\overline{Y} \setminus Y$ and $\overline{Z} \setminus Z$ are proper subcontinua of $\nu_P X$ with $\nu_P X = (\overline{Y} \setminus Y) \cup (\overline{Z} \setminus Z)$. Hence $\nu_P X$ is a decomposable continuum.

Let $K \subset X$ be a continuum and let $a, b \in K$ be distinct two points. If there is no proper subcontinuum of K containing both a and b then K is said to be *irreducible* between a and b.

The following lemma was proved by Bellamy [2, Lemma 1] for the Stone-Čech compactification of $[0,\infty)$. The proof is valid for any compactification of $[0,\infty)$. Here we give a proof for the reader's sake.

Lemma 3.5. Every compactification $\alpha([0,\infty))$ of the half-open interval $[0,\infty)$ is an irreducible continuum between 0 and every point z of $\alpha([0,\infty)) \setminus [0,\infty)$.

Proof. The connectivity of $\alpha([0,\infty))$ is obvious. Let $z \in \alpha([0,\infty)) \setminus [0,\infty)$. Suppose that there exists a proper closed subset K of $\alpha([0,\infty))$ such that $0,z\in K$. Since $\alpha([0,\infty))\setminus K$ is a non-empty open subset of $\alpha([0,\infty))$, there exists a point $t\in$ $[0,\infty)\cap(\alpha([0,\infty))\setminus K)$. Then $U=K\cap[0,t)=K\cap[0,t]$ is a non-empty closed and open subset of K since $0 \in U$. On the other hand, $z \in K \setminus U \neq \emptyset$, that is, K is not connected.

Theorem 3.6. Let $X = [0, \infty)$ be the half open interval with the usual metric d. For each $A \in \{H, P, L\}$, there exists a continuous surjection $\xi_A : \nu_P X \to h_A(X)$ from the subpower Higson corona $\nu_P X$ onto the Higson type compactification $h_A(X)$.

Proof. We shall construct a continuous surjection $\xi_H: \nu_P X \to h_H(X)$ from the subpower Higson corona $\nu_P X$ onto the Higson compactification $h_H(X)$. Then, for each $A \in \{P, L\}$, a required surjection is obtained by the composition $\alpha_A^H \circ \xi_H$: $\nu_P X \to h_A(X)$ where $\alpha_A^H : h_H(X) \to h_A(X)$ is the canonical surjection assured by the relation $h_H(X) \succeq h_A(X)$.

Let x and y be distinct two points in $\nu_P X$. Let U and V be open subsets of $h_P(X)$ such that $x \in U$, $y \in V$ and $\overline{U} \cap \overline{V} = \emptyset$. Then, by Lemma 3.1, there exist a natural number $k \geq 3$ and sequences $(a_n)_{n=1}^{\infty}$ and $(b_n)_{n=1}^{\infty}$ with $0 = b_0 < a_1 < b_1 < a_2 < a_2 < a_3 < a_2 < a_4 < a_4 < a_4 < a_5 < a_$ $\cdots < b_{n-1} < a_n < b_n < \cdots$ satisfying the following conditions:

- $(1) (a_n)^{1/k} > 2^n$
- (2) $b_{n-1} + (b_{n-1})^{1/k} < a_n$,
- (3) $[b_{n-1}, a_n] \cap U \neq \emptyset$,
- $(4) b_n = a_n + (a_n)^{2/k},$
- (5) $[a_n, b_n] \subset V \cap X$,
- (6) $a_n b_{n-1} > 2^{n-1} (a_n)^{1/k},$ (7) $b_n a_n > 2^n (b_n)^{1/k}.$

Then we define $\theta: X \to X$ by

$$\theta(t) = \begin{cases} 0, & \text{if } b_{2n-2} \le t < a_{2n-1}, \\ \frac{(2n-1)(t-a_{2n-1})}{b_{2n-1} - a_{2n-1}}, & \text{if } a_{2n-1} \le t < b_{2n-1}, \\ 2n-1, & \text{if } b_{2n-1} \le t < a_{2n}, \\ \frac{(2n-1)(b_{2n}-t)}{b_{2n} - a_{2n}}, & \text{if } a_{2n} \le t < b_{2n}. \end{cases}$$

Let $\psi: X \to [0,1]$ be the map constructed in Proposition 3.2, regarded as a map of X into X, i.e., $\psi: X \to X$. Then it follows from our construction that $\theta(t) = (2n-1)\psi(t)$ whenever $b_{2n-2} \le t < b_{2n}$.

Let $\ell: \mathbb{R}_+ \to \mathbb{R}_+$ be a function defined by $\ell(t) = t/2^t$. On should note that ℓ is monotonically decreasing on $(1/\log 2, \infty)$. In particular, $\lim_{t\to\infty} \ell(t) = 0$. Then we have the following.

Claim 3.6.1. If $n \geq 3$ and $t \in [a_{2n-1}, a_{2n+1})$ then $|\theta(t) - \theta(u)| < \ell(2n-1)$ for every $u \in B_d(t, t^{1/k})$.

Proof of Claim 3.6.1. Let $u \in B_d(t, t^{1/k})$. If $t \in [a_{2n-1}, a_{2n})$ then $u \in (b_{2n-2}, b_{2n})$ by Claim 3.2.1. Then $\theta(t) = (2n-1)\psi(t)$ and $\theta(u) = (2n-1)\psi(u)$. By Claim 3.2.2, we have $|\theta(t) - \theta(u)| = (2n-1)|\psi(t) - \psi(u)| < (2n-1)/2^{2n-1} = \ell(2n-1)$.

Suppose that $t \in [a_{2n}, b_{2n})$. Then $\theta(t) = (2n-1)\psi(t)$. By Claim 3.2.1, $u \in (b_{2n-1}, a_{2n+1})$. If $u \in (b_{2n-1}, b_{2n})$ then $\theta(u) = (2n-1)\psi(u)$. If $u \in [b_{2n}, a_{2n+1})$ then $\theta(u) = (2n-1)\psi(u)$ since $\theta(u) = \psi(u) = 0$. Hence, we obtain $|\theta(t) - \theta(u)| < \ell(2n-1)$ as above.

Now suppose that $t \in [b_{2n}, a_{2n+1})$. In this case, $\theta(t) = \psi(t) = 0$. So we have $\theta(t) = (2n-1)\psi(t)$. By Claim 3.2.1, $u \in (a_{2n}, b_{2n+1})$. If $u \in (a_{2n}, b_{2n})$ then $\theta(u) = (2n-1)\psi(u)$. If $u \in [b_{2n}, a_{2n+1})$ then $\theta(u) = (2n-1)\psi(u)$ since $\theta(u) = \psi(u) = 0$. Hence, we have $|\theta(t) - \theta(u)| < \ell(2n-1)$ as above. Finally, if $u \in [a_{2n+1}, b_{2n+1})$ then, by the conditions (4) and (1), we have

$$|\theta(t) - \theta(u)| = \theta(u) = \frac{(2n+1)(u - a_{2n+1})}{b_{2n+1} - a_{2n+1}} = \frac{(2n+1)(u - a_{2n+1})}{(a_{2n+1})^{2/k}}$$

$$< \frac{(2n+1)t^{1/k}}{(a_{2n+1})^{2/k}} < \frac{(2n+1)(a_{2n+1})^{1/k}}{(a_{2n+1})^{2/k}} = \frac{2n+1}{(a_{2n+1})^{1/k}}$$

$$< \frac{2n+1}{2^{2n+1}} = \ell(2n+1).$$

We note that $\ell(2n+1) < \ell(2n-1)$ since ℓ is decreasing on $(1/\log 2, \infty)$ and $2n-1 \ge 5 > 1/\log 2 = 3.3$. Thus we have $|\theta(t) - \theta(u)| < \ell(2n-1)$.

Claim 3.6.2. If $n \geq 3$ and $t \geq a_{2n-1}$ then $|\theta(t) - \theta(u)| < \ell(2n-1)$ for every $u \in B_d(t, t^{1/k})$.

Proof of Claim 3.6.2. Since ℓ is monotonically decreasing on $(1/\log 2, \infty)$ and $2n-1 \ge 5 > 1/\log 2 = 3.3$, this is an easy consequence of Claim 3.6.1.

Claim 3.6.3. For every $f \in C_H(X)$, the composition $f \circ \theta : X \to \mathbb{R}$ is Higson subpower.

Proof of Claim 3.6.3. Let $f: X \to \mathbb{R}$ be a continuous Higson map and let $\varepsilon > 0$ be given. If $t \ge a_5$ then $|\theta(t) - \theta(u)| < \ell(5)$ for every $u \in B_d(t, t^{1/k})$ by Claim 3.6.2. Since f is Higson, there exists K > 0 such that

(a)
$$\operatorname{diam} f(B_d(t, \ell(5))) < \varepsilon \text{ for every } t > K.$$

By the compactness of the interval [0,K] and the continuity of f, we can take $\delta>0$ so that

(b)
$$|f(t) - f(u)| < \varepsilon/3$$
 whenever $t \in [0, K]$ and $|t - u| < 3\delta$.

Without loss of generality, we may assume that $\delta < \ell(5)$. Since $\lim_{t\to\infty} \ell(t) = 0$, we can take an integer $n_0 \geq 3$ so that $\ell(2n_0 - 1) < \delta/2$.

Let $s: X \to \mathbb{R}_+$ be an asymptotically subpower function. We take $t_1 > 0$ so that $s(t) < t^{1/k}$ for every $t > t_1$. Then we put $t_0 = \max\{t_1, a_{2n_0-1}\}$.

Let $t > t_0$. It follows from $t > t_1$ that $\theta(B_d(t, s(t))) \subset \theta(B_d(t, t^{1/k}))$. Since $t > a_{2n_0-1}$, it follows from Claim 3.6.2 that $|\theta(t) - \theta(u)| < \ell(2n_0 - 1)$ for every $u \in B_d(t, t^{1/k})$, i.e., $\theta(B_d(t, t^{1/k})) \subset B_d(\theta(t), 2\ell(2n_0 - 1))$. Since $\ell(2n_0 - 1) < \delta/2$, we have $B_d(\theta(t), 2\ell(2n_0 - 1)) \subset B_d(\theta(t), \delta)$. As a summary, we have

$$\theta(B_d(t,s(t))) \subset B_d(\theta(t),\delta).$$

Suppose that $\theta(t) \leq K$. It follows from the condition (b) that

$$\operatorname{diam} f \circ \theta(B_d(t, s(t))) \leq \operatorname{diam} f(B_d(\theta(t), \delta)) < \varepsilon.$$

Now suppose that $\theta(t) > K$. Since $\delta < \ell(5)$, we have $B_d(\theta(t), \delta) \subset B_d(\theta(t), \ell(5))$. By the condition (a), we have

diam
$$f \circ \theta(B_d(t, s(t))) < \text{diam } f(B_d(\theta(t), \ell(5))) < \varepsilon$$
.

 \Diamond

Hence diam $f \circ \theta(B_d(t, s(t))) < \varepsilon$ for every $t > t_0$.

Claim 3.6.4. The map $\theta: X \to X$ has an extension $\Theta: h_P(X) \to h_H(X)$.

Proof of Claim 3.6.4. Let $e: X \to \prod_{f \in C_H(X)} I_f$ be the evaluation map. Since $h_H(X)$ is equivalent to $\overline{e(X)}$, it suffices to see that $e \circ \theta: X \to \prod_{f \in C_H(X)} I_f$ can be extended to $h_P(X) \to \overline{e(X)}$. For every $f \in C_H(X)$, the composition $f \circ \theta: X \to I_f$ is Higson subpower by Claim 3.6.3, so it can be extended to the map $\widehat{f \circ \theta}: h_P(X) \to I_f$ by Proposition 2.1. Hence $e \circ \theta$ can be extended to the map $\widehat{e \circ \theta}: h_P(X) \to \prod_{f \in C_P(X)} I_f$ so that $\widehat{e \circ \theta}(h_P(X)) \subset \overline{e(X)}$.

Let $(c_n)_{n=1}^{\infty}$ and $(d_n)_{n=1}^{\infty}$ be sequences such that $c_n \in [b_{2n-2}, a_{2n-1}]$ and $d_n \in [b_{2n-1}, a_{2n}]$ and let p and q be cluster points of $(c_n)_{n=1}^{\infty}$ and $(d_n)_{n=1}^{\infty}$ respectively. Put $C = \{c_n \mid n \in \mathbb{N}\}$ and $D = \{d_n \mid n \in \mathbb{N}\}$. Then the system C, D diverges as a power function by (1) and (7). Hence $p \in \overline{C} \cap \nu_P X$ and $q \in \overline{D} \cap \nu_P X$ are distinct two points in $\nu_P X$. By Claim 3.6.4, there exists an extension $\Theta : h_P(X) \to h_H(X)$ of θ . Then $\Theta(p) = 0$ and $\Theta(q) \in \nu_H X$. Thus $\Theta(\nu_P X)$ is a continuum containing 0 and $\Theta(q) \in \nu_H X$. So, $\Theta(\nu_P X)$ must coincide with $h_H(X)$ by Lemma 3.5. Hence the map $\xi_H = \Theta|\nu_P X : \nu_P X \to h_H(X)$ is a required surjection.

Corollary 3.7. Let $X = [0, \infty)$ be the half open interval with the usual metric d. For each $A \in \{H, P, L\}$, there exists a continuous surjection $\eta_A : \nu_H X \to h_A(X)$ from the Higson corona $\nu_H X$ onto the Higson type compactification $h_A(X)$.

Proof. Let $A \in \{H, P, L\}$ and let $\xi_A : \nu_P X \to h_A(X)$ be a continuous surjection given by Theorem 3.6. By the relation $h_H(X) \succeq h_P(X)$, there is the canonical surjection $\alpha : h_H(X) \to h_P(X)$, such that $\alpha|_X = \mathrm{id}_X$. Since X is dense in $h_H(X)$ and $\alpha|_X = \mathrm{id}_X$, it follows that $\alpha(\nu_H X) = \nu_P X$. Thus the composition $\eta_A = \xi_A \circ \alpha|_{\nu_H X} : \nu_H X \to h_A(X)$ is a required surjection.

Question 1. Does Proposition 3.2 hold for sublinear Higson corona? In particular, is the sublinear Higson corona of the half open interval with the usual metric an indecomposable continuum?

Let W be a non-degenerate subcontinuum of $\beta[0,\infty) \setminus [0,\infty)$ and let M be a metric continuum. It is known [2] that there exist continuous surjections $f:W\to\beta[0,\infty)$ and $g:W\to M$.

Question 2. Let $X = [0, \infty)$ be the half open interval with the usual metric d. Let W be a non-degenerate subcontinuum of $\nu_P(X)$ and let M be a metric continuum. Do there exist continuous surjections $f: W \to h_P(X)$ and $g: W \to M$?

Acknowledgement

I would like to express my sincere gratitude to the referee for his/her comments and valuable suggestions which helped to improve the manuscript.

References

- [1] B.J. Ball and S. Yokura Compactifications determined by subsets of $C^*(X)$, Topology Appl. 13 (1982), 1–13.
- [2] D. Bellamy, A non-metric indecomposable continuum, Duke Math. J., 38 (1971), 15–20.
- [3] M. Cencelj, J. Dydak, J. Smrekar and A. Vavpetič, Sublinear Higson corona and Lipschitz extensions, Houston J. Math., 37 (2011), No. 4, 1307–1322.
- [4] A. N. Dranishnikov, J. Keesling and V. V. Uspenskij, On the Higson corona of uniformly contractible spaces, Topology 37(4) (1999), 791–803.
- [5] A. N. Dranishnikov and J. Smith, On asymptotic Assouad-Nagata dimension, Topology Appl., 154 (2007), 934-952.
- [6] J. Dydak and C. S. Hoffland, An alternative definition of coarse structures, Topology Appl., 155 (2008), 1013–1021.
- [7] R. Engelking, General topology, Heldermann Verlag, Berlin, Revised and completed ed., 1989.
- [8] L. Gillman and M. Jerison, Rings of continuous functions, D. van Nostrand Company, Inc., Princeton, 1960.
- [9] Y. Iwamoto and K. Tomoyasu, Higson compactifications obtained by expanding and contracting the half-open interval, Tsukuba J. Math., 25 (2001), No. 1, 179–186.
- [10] J. Keesling, The one-dimensional Čech cohomology of the Higson compactification and its corona, Topology Proc., 19 (1994), 129–148.
- [11] J. Kucab and M. Zarichnyi, Subpower Higson corona of a metric space, Algebra and Discrete Mathematics, 17 (2014), No. 2, 280–287.
- [12] J. Kucab and M. Zarichnyi, On asymptotic power dimension, Topology Appl., 201 (2016), 124–130.
- [13] J. Kucab and M. Zarichnyi, On convergence in the subpower Higson corona of metric spaces, arXiv:1701.00721v2 [math.GN], 2017.
- [14] J. Roe, Lectures on Coarse Geometry, University Lecture Series, Vol. 31, AMS, 2003.
- [15] S. Willard, General Topology, Addison-Wesley, 1970.

Department of Engineering Science,

National Institute of Technology,

Niihama College, Niihama, 792-8580, Japan

E-mail address: iwamoto@sci.niihama-nct.ac.jp