

ON PERIODIC SOLUTIONS TO LAGRANGIAN SYSTEM WITH CONSTRAINTS

OLEG ZUBELEVICH

DEPT. OF THEORETICAL MECHANICS,
MECHANICS AND MATHEMATICS FACULTY,
M. V. LOMONOSOV MOSCOW STATE UNIVERSITY
RUSSIA, 119899, MOSCOW, MGU

ABSTRACT. A Lagrangian system is considered. The configuration space is a non-compact manifold that depends on time. A set of periodic solutions has been found.

1. INTRODUCTION

Existence problems for periodic solutions to Lagrangian systems have intensively been studied since the beginning of the 20th century and even earlier. There is an immense number of different results and methods developed in this field. We mention only a few of them which are mach closely related to this article.

In [2] periodic solutions have been obtained for the Lagrangian system with Lagrangian

$$L(t, x, \dot{x}) = \frac{1}{2}g_{ij}(x)\dot{x}^i\dot{x}^j - W(t, x), \quad x = (x^1, \dots, x^m) \in \mathbb{R}^m$$

here and in the sequel we use the Einstein summation convention. The form g_{ij} is symmetric and positive definite:

$$g_{ij}\xi^i\xi^j \geq \text{const}_1 \cdot |\xi|^2.$$

The potential is as follows $W(t, x) = V(x) + g(t) \sum_{i=1}^m x^i$, where V is a bounded function $|V(x)| \leq \text{const}_2$ and g is an ω -periodic function. The functions V, g_{ij} are even.

Under these assumptions the authors prove that there exists a non-trivial ω -periodic solution.

2000 *Mathematics Subject Classification.* 34C25, 70F20 .

Key words and phrases. Lagrangian systems, periodic solutions.

Partially supported by grant RFBR 18-01-00887.

Our main tool to obtain periodic solutions is variation technique. Variational problems and Hamiltonian systems have been studied extensively. Classic references of these subjects are [6], [7], [4].

2. THE MAIN THEOREM

Introduce some notations. Let $x = (x^1, \dots, x^m)$ and $\varphi = (\varphi^1, \dots, \varphi^n)$ be points of the standard \mathbb{R}^m and \mathbb{R}^n respectively. Then let z stand for the point $(x, \varphi) \in \mathbb{R}^{m+n}$. By $|\cdot|$ denote the standard Euclidean norm of \mathbb{R}^k , $k = m, m+n$, that is $|x|^2 = \sum_{i=1}^k (x^i)^2$.

The variable z can consist only of x without φ or conversely.

The main object of our study is the following Lagrangian system with Lagrangian

$$L(t, z, \dot{z}) = \frac{1}{2}g_{ij}\dot{z}^i\dot{z}^j + a_i\dot{z}^i - V, \quad z = (z^1, \dots, z^{m+n}). \quad (2.1)$$

Remark 1. *The term $a_i\dot{z}^i$ in the Lagrangian corresponds to the so called gyroscopic forces. For example, the Coriolis force and the Lorentz force are gyroscopic.*

The functions g_{ij}, a_i, V depend on (t, z) and belong to $C^2(\mathbb{R}^{m+n+1})$; moreover all these functions are 2π -periodic in each variable φ^j and ω -periodic in the variable t , $\omega > 0$. For all $(t, z) \in \mathbb{R}^{m+n+1}$ it follows that $g_{ij} = g_{ji}$.

We also assume that there are positive constants C, M, A, K such that for all (t, z) and $\xi \in \mathbb{R}^{m+n}$ we have

$$|a_i(t, z)| \leq C + M|z|, \quad V(t, z) \leq A|z|^2, \quad \frac{1}{2}g_{ij}(t, z)\xi^i\xi^j \geq K|\xi|^2. \quad (2.2)$$

System (2.1) obeys the following ideal constraints:

$$f_j(t, z) = 0, \quad j = 1, \dots, l < m+n, \quad f_j \in C^2(\mathbb{R}^{m+n+1}). \quad (2.3)$$

The functions f_j are also 2π -periodic in each variable φ^j and ω -periodic in the variable t .

Introduce a set

$$F = \{(t, z) \in \mathbb{R}^{m+n+1} \mid f_j(t, z) = 0, \quad j = 1, \dots, l\}.$$

Assume that

$$\text{rank} \left(\frac{\partial f_j}{\partial z^k}(t, z) \right) = l \quad (2.4)$$

for all $(t, z) \in F$. So that F is a smooth manifold.

Assume also that all the functions f_j are either odd:

$$f_j(-t, -z) = -f_j(t, z) \quad (2.5)$$

or even

$$f_j(-t, -z) = f_j(t, z). \quad (2.6)$$

Remark 2. *Actually, it is sufficient to say that all the functions are defined and have formulated above properties in some open symmetric vicinity of the manifold F . We believe that this generalization is unimportant and keep referring to the whole space \mathbb{R}^{m+n+1} just for simplicity of exposition.*

Definition 1. *We shall say that a function $z(t) \in C^2(\mathbb{R}, \mathbb{R}^{n+m})$ is a solution to system (2.1), (2.3) if there exists a set of functions*

$$\{\alpha^1, \dots, \alpha^l\} \subset C(\mathbb{R})$$

such that

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{z}^i}(t, z(t), \dot{z}(t)) - \frac{\partial L}{\partial z^i}(t, z(t), \dot{z}(t)) = \alpha^j(t) \frac{\partial f_j}{\partial z^i}(t, z(t)) \quad (2.7)$$

and

$$(t, z(t)) \in F \quad (2.8)$$

for all real t .

In the absence of constraints (2.3) right side of (2.7) is equal to zero and the condition (2.8) is dropped.

The functions α^i are defined from equations (2.7), (2.3) uniquely.

Theorem 2.1. *Assume that*

1) *all the functions are even:*

$$g_{ij}(-t, -z) = g_{ij}(t, z), \quad a_i(-t, -z) = a_i(t, z), \quad V(-t, -z) = V(t, z);$$

2) *the following inequality holds*

$$K - \frac{M\omega}{\sqrt{2}} - \frac{A\omega^2}{2} > 0; \quad (2.9)$$

3) *Assume that for some $\nu = (\nu_1, \dots, \nu_n) \in \mathbb{Z}^n$ there is a function $\tilde{z}(t) = (\tilde{x}(t), \tilde{\varphi}(t)) \in C^1(\mathbb{R}, \mathbb{R}^{m+n})$ such that*

$$\tilde{z}(-t) = -\tilde{z}(t), \quad \tilde{x}(t + \omega) = \tilde{x}(t), \quad \tilde{\varphi}(t + \omega) = \tilde{\varphi}(t) + 2\pi\nu$$

and

$$(t, \tilde{z}(t)) \in F, \quad t \in \mathbb{R}. \quad (2.10)$$

Then system (2.1), (2.3) has a solution $z(t) = (x(t), \varphi(t))$ such that

1) *the function z is odd: $z(-t) = -z(t)$;*

2) *$x(t + \omega) = x(t)$, $\varphi(t + \omega) = \varphi(t) + 2\pi\nu$;*

3) *the functions $\{\alpha^i\}$ are ω -periodic and odd provided the constraints are odd; and α^j are even for the even constraints.*

This assertion remains valid in the absence of constraints (2.3).

Actually the solution stated in this theorem is as smooth as it is allowed by smoothness of the Lagrangian L and the functions f_j up to C^∞ .

Remark 3. 1) If all the functions do not depend on t then we can choose ω to be arbitrary small and inequality (2.9) is satisfied. Taking a vanishing sequence of ω , we obtain infinitely many periodic solutions of the same homotopic type;

2) Suppose that $M = 0$ and $V(t, z) \leq A_1|x|^\alpha$, $\alpha < 2$. Choose a small constant $A > 0$ such that inequality (2.9) is satisfied then choose a constant $c_1 > 0$ such that for all $|x|$ one has $A_1|x|^\alpha \leq A|x|^2 + c_1$. Now the second condition of the theorem is satisfied for the new potential

$$V_1 = V - c_1 \leq A|x|^2.$$

Remark 4. The condition 2) of the Theorem is essential for non-compact manifold F or when the constraints are absent. Indeed, system

$$L(t, x, \dot{x}) = \frac{1}{2}\dot{x}^2 - \frac{1}{2}(x - \sin t)^2$$

obeys all the conditions except inequality (2.9). It is easy to see that the corresponding equation $\ddot{x} + x = \sin t$ does not have periodic solutions.

When F is compact then condition 2) can be dropped and the theorem follows from the standard results.

2.1. Examples. Our first example is as follows.




FIGURE 1. the tube and the ball

A thin tube can rotate freely in the vertical plane about a fixed horizontal axis passing through its centre O . A moment of inertia of the tube about this axis is equal to J . The mass of the tube is distributed symmetrically such that tube's centre of mass is placed at the point O .

Inside the tube there is a small ball which can slide without friction. The mass of the ball is m . The ball can pass by the point O and fall out from the ends of the tube.

The system undergoes the standard gravity field \mathbf{g} .

It seems to be evident that for typical motion the ball reaches an end of the tube and falls down out the tube. It is surprisingly, at least for the first glance, that this system has very many periodic solutions such that the tube turns around several times during the period.

The sense of generalized coordinates ϕ, x is clear from Figure 1.

The Lagrangian of this system is as follows

$$L(x, \phi, \dot{x}, \dot{\phi}) = \frac{1}{2} \left(m\dot{x}^2 + J \right) \dot{\phi}^2 + \frac{1}{2} m\dot{x}^2 - mgx \sin \phi. \quad (2.11)$$

Form theorem 2.1 it follows that for any constant $\omega > 0$ system (2.11) has a solution $\phi(t), x(t)$, $t \in \mathbb{R}$ such that

- 1) $x(t) = -x(-t)$, $\phi(t) = -\phi(-t)$;
- 2) $x(t + \omega) = x(t)$, $\phi(t + \omega) = \phi(t) + 2\pi$.

This result shows that for any $\omega > 0$ the system has an ω -periodic motion such that the tube turns around once during the period. The length of the tube should be chosen properly.

Our second example is a counterexample. Let us show that the first condition of the theorem 2.1 can not be omitted.

Consider a mass point m that slides on a right circular cylinder of radius r . The surface of the cylinder is perfectly smooth. The axis x of the cylinder is parallel to the gravity \mathbf{g} and directed upwards.

The Lagrangian of this system is

$$L(x, \varphi, \dot{x}, \dot{\varphi}) = \frac{m}{2} \left(r^2 \dot{\varphi}^2 + \dot{x}^2 \right) - mgx. \quad (2.12)$$

All the conditions except the evenness are satisfied but it is clear this system does not have periodic solutions.

3. PROOF OF THEOREM 2.1

In this section we use several standard facts from functional analysis and the Sobolev spaces theory [3], [1].

1. Recall that the Sobolev space $H_{\text{loc}}^1(\mathbb{R})$ consists of functions $u(t)$, $t \in \mathbb{R}$ such that $u, \dot{u} \in L_{\text{loc}}^2(\mathbb{R})$. The following embedding holds $H_{\text{loc}}^1(\mathbb{R}) \subset C(\mathbb{R})$.

Lemma 3.1. *Let $u \in H_{\text{loc}}^1(\mathbb{R})$ and $u(0) = 0$. Then for any $a > 0$ we have*

$$\|u\|_{L^2(0,a)}^2 \leq \frac{a^2}{2} \|\dot{u}\|_{L^2(0,a)}^2, \quad \|u\|_{C[0,a]}^2 \leq a \|\dot{u}\|_{L^2(0,a)}^2.$$

Here and below the notation $\|\dot{u}\|_{L^2(0,a)}$ implies that

$$\left\| \dot{u} \Big|_{(0,a)} \right\|_{L^2(0,a)}$$

the same is concerned to $\|u\|_{C[0,a]}$ etc.

This Lemma is absolutely standard, nevertheless just for completeness of exposition sake we bring a sketch of its proof.

Proof of Lemma 3.1. From formula

$$u(t) = \int_0^t \dot{u}(s) ds \quad (3.1)$$

it follows that

$$\int_0^a u^2(s) ds = \int_0^a \left(\int_0^t \dot{u}(s) ds \right)^2 dt.$$

It remains to observe that by the Cauchy inequality

$$\left| \int_0^t \dot{u}(s) ds \right| \leq \int_0^t |\dot{u}(s)| ds \leq \|\dot{u}\|_{L^2(0,a)} \left(\int_0^t ds \right)^{1/2}, \quad t \in [0, a].$$

This implies the first inequality of the lemma. The second inequality also follows from formula (3.1) and the Cauchy inequality.

The Lemma is proved.

2. Here we collect several spaces which are needed in the sequel.

Definition 2. By X denote a space of functions $u \in H_{\text{loc}}^1(\mathbb{R})$ such that for all $t \in \mathbb{R}$ the following conditions hold

$$u(-t) = -u(t), \quad u(t + \omega) = u(t).$$

By virtue of lemma 3.1, the mapping $u \mapsto \|\dot{u}\|_{L^2(0,\omega)}$ determines a norm in X . This norm is denoted by $\|u\|$. The norm $\|\cdot\|$ is equivalent to the standard norm of $H^1[0,\omega]$. The space $(X, \|\cdot\|)$ is a Banach space. Since the norm $\|\cdot\|$ is generated by an inner product

$$(u, v)_X = \int_0^\omega \dot{u}(t)\dot{v}(t) dt$$

the space X is also a real Hilbert space, particularly this implies that X is a reflexive Banach space.

Definition 3. Let Φ stand for the space $\{ct + u(t) \mid c \in \mathbb{R}, \quad u \in X\}$.

By the same argument, $(\Phi, \|\cdot\|)$ is a reflexive Banach space. Observe also that $\Phi = \mathbb{R} \oplus X$ and by direct calculation we get

$$\|\psi\|^2 = \omega c^2 + \|u\|^2, \quad \psi(t) = ct + u(t) \in \Phi.$$

Observe that $X \subset \Phi$.

Definition 4. Let E stand for the space

$$X^m \times \Phi^n = \{z(t) = (x^1, \dots, x^m, \varphi^1, \dots, \varphi^n)(t) \mid x^i \in X, \quad \varphi^j \in \Phi\}.$$

The space E is also a real Hilbert space with an inner product defined as follows

$$(z, y)_E = \int_0^\omega \sum_{i=1}^{m+n} \dot{x}^i(t) \dot{y}^i(t) dt,$$

where $z = (z^k), y = (y^k) \in E, \quad k = 1, \dots, m+n$.

We denote the corresponding norm in E by the same symbol and write

$$\|z\|^2 = \|\|z\|\|^2 = \sum_{k=1}^{m+n} \|z^k\|^2.$$

The space E is also a reflexive Banach space.

Introduce the following set

$$E_0 = \left\{ (x, \varphi) \in E \mid \varphi^j = \frac{2\pi\nu_j}{\omega} t + u_j, \quad \forall x \in X, \quad \forall u_j \in X, \quad j = 1, \dots, n \right\}.$$

This set is a closed plane of codimension n in E .

If $(x, \varphi) \in E_0$ then $\varphi(t + \omega) = \varphi(t) + 2\pi\nu$.

Definition 5. Let Y stand for the space

$$\{u \in L_{\text{loc}}^2(\mathbb{R}) \mid u(t) = u(-t), \quad u(t+\omega) = u(t) \quad \text{almost everywhere in } \mathbb{R}\}.$$

The space Y^{m+n} is a Hilbert space with respect to the inner product

$$(z, y)_Y = \int_0^\omega \sum_{i=1}^{m+n} \dot{x}^i(t) \dot{y}^i(t) dt,$$

where $z = (z^k), y = (y^k) \in Y^{m+n}, \quad k = 1, \dots, m+n$.

Definition 6. Let W stand for the set

$$\{z(\cdot) \in E_0 \mid (t, z(t)) \in F, \quad t \in \mathbb{R}\}.$$

3. By formula (2.10) the set W is non-void.

Introduce the Action Functional $S : W \rightarrow \mathbb{R}$,

$$S(z) = \int_0^\omega L(t, z, \dot{z}) dt.$$

Our next goal is to prove that this functional attains its minimum.

Observe that $|x| \leq |z|$ then by using estimates (2.2) we get

$$S(z) \geq \int_0^\omega (K|\dot{z}|^2 - |\dot{z}|(C + M|z|) - A|z|^2) dt.$$

From the Cauchy inequality and Lemma 3.1 it follows that

$$\begin{aligned}\int_0^\omega |\dot{z}| |z| dt &\leq \frac{\omega}{\sqrt{2}} \|z\|^2, \\ \int_0^\omega |z|^2 dt &\leq \frac{\omega^2}{2} \|z\|^2, \\ \int_0^\omega |\dot{z}| dt &\leq \sqrt{\omega} \|z\|.\end{aligned}$$

We finally obtain

$$S(z) \geq \left(K - \frac{M\omega}{\sqrt{2}} - \frac{A\omega^2}{2} \right) \|z\|^2 - C\sqrt{\omega} \|z\|. \quad (3.2)$$

Taking into account Remark 4 we proceed with the assumption that the manifold F is not compact.

By formula (2.9) the functional S is coercive:

$$S(z) \rightarrow \infty \quad (3.3)$$

as $\|z\| \rightarrow \infty$.

Note that the Action Functional which corresponds to system (2.12) is also coercive but, as we see above, property (3.3) by itself does not imply existence results.

4. Let $\{z_k\} \subset W$ be a minimizing sequence:

$$S(z_k) \rightarrow \inf_{z \in W} S(z)$$

as $k \rightarrow \infty$.

By formula (3.3) the sequence $\{z_k\}$ is bounded: $\sup_k \|z_k\| < \infty$. Since the space E is reflexive, this sequence contains a weakly convergent subsequence. Denote this subsequence in the same way: $z_k \rightarrow z_*$ weakly in E .

Moreover, the space $H^1[0, \omega]$ is compactly embedded in $C[0, \omega]$. Thus extracting a subsequence from the subsequence and keeping the same notation we also have

$$\max_{t \in [0, \omega]} |z_k(t) - z_*(t)| \rightarrow 0, \quad (3.4)$$

as $k \rightarrow \infty$.

The set E_0 is convex and strongly closed therefore it is weakly closed: $z_* \in E_0$. By continuity (3.4) one also gets $z_* \in W$.

5. Let us show that $\inf_{z \in W} S(z) = S(z_*)$.

Lemma 3.2. *Let a sequence $\{u_k\} \subset \Phi$ weakly converge to $u \in \Phi$ (or $u_k, u \in X$ and $u_k \rightarrow u$ weakly in X); and also $\max_{t \in [0, \omega]} |u_k(t) - u(t)| \rightarrow 0$ as $k \rightarrow \infty$.*

Then for any $f \in C(\mathbb{R})$ and for any $v \in L^2(0, \omega)$ it follows that

$$\int_0^\omega f(u_k) \dot{u}_k v dt \rightarrow \int_0^\omega f(u) \dot{u} v dt, \quad \text{as } k \rightarrow \infty.$$

Indeed,

$$\begin{aligned} & \int_0^\omega f(u_k) \dot{u}_k v dt \\ &= \int_0^\omega (f(u_k) - f(u)) \dot{u}_k v dt + \int_0^\omega f(u) \dot{u}_k v dt. \end{aligned}$$

The function f is uniformly continuous in a compact set

$$\left[\min_{t \in [0, \omega]} \{u(t)\} - c, \max_{t \in [0, \omega]} \{u(t)\} + c \right]$$

with some constant $c > 0$. Consequently we obtain

$$\max_{t \in [0, \omega]} |f(u_k(t)) - f(u(t))| \rightarrow 0.$$

Since the sequence $\{u_k\}$ is weakly convergent it is bounded:

$$\sup_k \|u_k\| < \infty$$

particularly, we get

$$\|\dot{u}_k\|_{L^2(0, \omega)} < \infty.$$

So that as $k \rightarrow \infty$

$$\begin{aligned} & \left| \int_0^\omega (f(u_k) - f(u)) \dot{u}_k v dt \right| \\ & \leq \|v(f(u_k) - f(u))\|_{L^2(0, \omega)} \cdot \|\dot{u}_k\|_{L^2(0, \omega)} \rightarrow 0. \end{aligned}$$

To finish the proof it remains to observe that a function

$$w \mapsto \int_0^\omega f(u) \dot{w} v dt$$

belongs to Φ' (or to X'). Indeed,

$$\left| \int_0^\omega f(u) \dot{w} v dt \right| \leq \max_{t \in [0, \omega]} |f(u(t))| \cdot \|v\|_{L^2(0, \omega)} \|w\|.$$

6. The following lemma is proved similarly.

Lemma 3.3. *Let a sequence $\{u_k\} \subset \Phi$ (or $\{u_k\} \subset X$) be such that*

$$\max_{t \in [0, \omega]} |u_k(t) - u(t)| \rightarrow 0$$

as $k \rightarrow \infty$.

Then for any $f \in C(\mathbb{R})$ and for any $v \in L^1(0, \omega)$ it follows that

$$\int_0^\omega f(u_k)v dt \rightarrow \int_0^\omega f(u)v dt$$

as $k \rightarrow \infty$.

7. Introduce a function $p_k(t, \xi) = L(t, z_k, \dot{z}_* + \xi)$. The function p_k is a quadratic polynomial of $\xi \in \mathbb{R}^{m+n}$, so that

$$p_k(t, \xi) = L(t, z_k, \dot{z}_*) + \frac{\partial L}{\partial \dot{z}^i}(t, z_k, \dot{z}_*)\xi^i + \frac{1}{2} \frac{\partial^2 L}{\partial \dot{z}^j \partial \dot{z}^i}(t, z_k, \dot{z}_*)\xi^i \xi^j.$$

The last term in this formula is non-negative:

$$\frac{\partial^2 L}{\partial \dot{z}^j \partial \dot{z}^i}(t, z_k, \dot{z}_*)\xi^i \xi^j = g_{ij}(t, z_k)\xi^i \xi^j \geq 0.$$

We consequently obtain

$$p_k(t, \xi) \geq L(t, z_k, \dot{z}_*) + \frac{\partial L}{\partial \dot{z}^i}(t, z_k, \dot{z}_*)\xi^i.$$

It follows that

$$\begin{aligned} S(z_k) &= \int_0^\omega p_k(t, \dot{z}_k - \dot{z}_*) dt \geq \int_0^\omega L(t, z_k, \dot{z}_*) dt \\ &\quad + \int_0^\omega \frac{\partial L}{\partial \dot{z}^i}(t, z_k, \dot{z}_*)(\dot{z}_k^i - \dot{z}_*^i) dt. \end{aligned} \quad (3.5)$$

From Lemma 3.2 and Lemma 3.3 it follows that

$$\int_0^\omega L(t, z_k, \dot{z}_*) dt \rightarrow \int_0^\omega L(t, z_*, \dot{z}_*) dt \quad \text{as } k \rightarrow \infty,$$

and

$$\int_0^\omega \frac{\partial L}{\partial \dot{z}^i}(t, z_k, \dot{z}_*)(\dot{z}_k^i - \dot{z}_*^i) dt \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Passing to the limit as $k \rightarrow \infty$ in (3.5) we finally yield

$$\inf_{z \in W} S(z) \geq S(z_*) \implies \inf_{z \in W} S(z) = S(z_*).$$

Remark 5. *Basing upon these formulas one can estimate the norm $\|z_*\|$. Indeed, take a function $\hat{z} \in W$ then due to formula (3.2) one obtains*

$$S(\hat{z}) \geq S(z_*) \geq \left(K - \frac{M\omega}{\sqrt{2}} - \frac{A\omega^2}{2} \right) \|z_*\|^2 - C\sqrt{\omega}\|z_*\|,$$

here $S(\hat{z})$ is an explicitly calculable number.

8. Now from this point we begin proving the theorem under the assumption that the constraints are odd (2.5).

Thus for any $v \in X^{m+n}$ such that

$$\frac{\partial f_j}{\partial z^k}(t, z_*) v^k(t) = 0$$

it follows that

$$\left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} S(z_* + \varepsilon v) = 0.$$

Introduce a linear functional

$$b : X^{m+n} \rightarrow \mathbb{R}, \quad b(v) = \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} S(z_* + \varepsilon v),$$

and a linear operator

$$A : X^{m+n} \rightarrow X^l, \quad (Av)_j = \frac{\partial f_j}{\partial z^k}(t, z_*) v^k.$$

It is clear, both these mappings are bounded and

$$\ker A \subset \ker b.$$

Lemma 3.4. *The operator A maps X^{m+n} onto X^l that is*

$$A(X^{m+n}) = X^l.$$

Proof. By $\tilde{A}(t)$ denote the matrix

$$\frac{\partial f_j}{\partial z^k}(t, z_*(t)).$$

It is convenient to consider our functions to be defined on the circle $t \in \mathbb{S} = \mathbb{R}/(\omega\mathbb{Z})$.

Fix an element $w \in X^l$. Let us cover the circle \mathbb{S} with open intervals U_i , $i = 1, \dots, N$ such that there exists a set of functions

$$v_i \in H^1(U_i), \quad \tilde{A}(t)v_i(t) = w(t), \quad t \in U_i, \quad i = 1, \dots, N.$$

And let ψ_i be a smooth partition of unity subordinated to the covering $\{U_i\}$. A function $\tilde{v}(t) = \sum_{i=1}^N \psi_i(t)v_i(t)$ belongs to $H^1(\mathbb{S})$ and for each t it follows that $\tilde{A}(t)\tilde{v}(t) = w(t)$. But the function \tilde{v} is not obliged to be odd.

Since $\tilde{A}(-t) = \tilde{A}(t)$ we have

$$\tilde{A}(t)v(t) = w(t), \quad v(t) = \frac{\tilde{v}(t) - \tilde{v}(-t)}{2} \in X^{m+n}.$$

The Lemma is proved.

9. Recall a lemma from functional analysis [5].

Lemma 3.5. *Let E, H, G be Banach spaces and*

$$A : E \rightarrow H, \quad B : E \rightarrow G$$

be bounded linear operators; $\ker A \subseteq \ker B$.

If the operator A is onto then there exists a bounded operator $\Gamma : H \rightarrow G$ such that $B = \Gamma A$.

Thus there is a linear function $\Gamma \in (X^l)'$ such that

$$b(v) = \Gamma A(v), \quad v \in X^{m+n}.$$

Or by virtue of the Riesz representation theorem, there exists a set of functions $\{\gamma^1, \dots, \gamma^l\} \subset X$ such that

$$\left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} S(z_* + \varepsilon v) = \int_0^\omega \dot{\gamma}^j(t) \frac{d}{dt} \left(\frac{\partial f_j}{\partial z^k}(t, z_*) v^k(t) \right) dt$$

for all $v \in X^{m+n}$.

10. Every element $v \in X^{m+n}$ is presented as follows

$$v(t) = \int_0^t y(s) ds,$$

where $y \in Y^{m+n}$ is such that

$$\int_0^\omega y(s) ds = 0.$$

Introduce a linear operator $h : Y^{m+n} \rightarrow \mathbb{R}^{m+n}$ by the formula

$$h(y) = \int_0^\omega y(s) ds.$$

Define a linear functional $q : Y^{m+n} \rightarrow \mathbb{R}$ by the formula

$$q(y) = (b - \Gamma A)v, \quad v(t) = \int_0^t y(s) ds.$$

Now all our observations lead to

$$\ker h \subseteq \ker q.$$

Therefore, there exists a linear functional $\lambda : \mathbb{R}^{m+n} \rightarrow \mathbb{R}$ such that

$$q = \lambda h$$

Let us rewrite the last formula explicitly. There are real constants λ_k such that for any $y^k \in Y$ one has

$$\begin{aligned} & \int_0^\omega \left(\frac{\partial L}{\partial z^k}(t, z_*, \dot{z}_*) y^k(t) + \frac{\partial L}{\partial z^k}(t, z_*, \dot{z}_*) \int_0^t y^k(s) ds \right) dt \\ &= \int_0^\omega \dot{\gamma}^j(t) \frac{\partial f_j}{\partial z^k}(t, z_*) y^k(t) dt + \int_0^\omega \dot{\gamma}^j(t) \frac{d}{dt} \left(\frac{\partial f_j}{\partial z^k}(t, z_*) \right) \int_0^t y^k(s) ds dt \\ &+ \lambda_k \int_0^\omega y^k(s) ds. \end{aligned}$$

11. By the Fubini theorem we obtain

$$\begin{aligned} & \int_0^\omega \frac{\partial L}{\partial z^k}(t, z_*, \dot{z}_*) y^k(t) dt + \int_0^\omega y^k(s) \int_s^\omega \frac{\partial L}{\partial z^k}(t, z_*, \dot{z}_*) dt ds \\ &= \int_0^\omega \dot{\gamma}^j(t) \frac{\partial f_j}{\partial z^k}(t, z_*) y^k(t) dt + \int_0^\omega y^k(s) \int_s^\omega \dot{\gamma}^j(t) \frac{d}{dt} \left(\frac{\partial f_j}{\partial z^k}(t, z_*) \right) dt ds \\ &+ \lambda_k \int_0^\omega y^k(s) ds. \end{aligned} \tag{3.6}$$

In this formula the functions

$$\frac{\partial L}{\partial z^k}(t, z_*, \dot{z}_*), \quad \dot{\gamma}^j(t) \frac{\partial f_j}{\partial z^k}(t, z_*)$$

are even and ω -periodic functions of t .

The functions

$$\dot{\gamma}^j(t) \frac{d}{dt} \left(\frac{\partial f_j}{\partial z^k}(t, z_*) \right), \quad \frac{\partial L}{\partial z^k}(t, z_*, \dot{z}_*)$$

are odd and ω -periodic. So that the functions

$$\int_s^\omega \dot{\gamma}^j(t) \frac{d}{dt} \left(\frac{\partial f_j}{\partial z^k}(t, z_*) \right) ds, \quad \int_s^\omega \frac{\partial L}{\partial z^k}(t, z_*, \dot{z}_*) ds$$

are even and ω -periodic in s .

Therefore, equation (3.6) is rewritten as $(y, q)_Y = 0$ for any $y = (y^1, \dots, y^{m+n}) \in Y^{m+n}$ and $q = (q_1, \dots, q_{m+n})$ stands for

$$\begin{aligned} q_k &= \frac{\partial L}{\partial z^k}(t, z_*, \dot{z}_*) + \int_t^\omega \frac{\partial L}{\partial z^k}(s, z_*, \dot{z}_*) ds \\ &- \dot{\gamma}^j(t) \frac{\partial f_j}{\partial z^k}(t, z_*) - \int_t^\omega \dot{\gamma}^j(s) \frac{d}{ds} \left(\frac{\partial f_j}{\partial z^k}(s, z_*) \right) ds - \lambda_k. \end{aligned}$$

Consequently we obtain the following system

$$\begin{aligned} & \frac{\partial L}{\partial \dot{z}^k}(t, z_*, \dot{z}_*) + \int_t^\omega \frac{\partial L}{\partial z^k}(s, z_*, \dot{z}_*) ds \\ &= \dot{\gamma}^j(t) \frac{\partial f_j}{\partial z^k}(t, z_*) + \int_t^\omega \dot{\gamma}^j(s) \frac{d}{ds} \left(\frac{\partial f_j}{\partial z^k}(s, z_*) \right) ds + \lambda_k. \end{aligned} \quad (3.7)$$

here $k = 1, \dots, m + n$.

If we formally differentiate both sides of equations (3.7) in t then we obtain the Lagrange equations (2.7) with $\alpha^j = \dot{\gamma}^j$.

Equations (3.7) hold for almost all $t \in (0, \omega)$ but all the functions contained in (3.7) are defined for all $t \in \mathbb{R}$.

Now we employ the following trivial observation: if $w \in L^1_{\text{loc}}(\mathbb{R})$ is an ω -periodic and odd function then for any constant $a \in \mathbb{R}$ a function

$$t \mapsto \int_a^t w(s) ds$$

is also ω -periodic.

Hence the functions

$$\int_t^\omega \dot{\gamma}^j(s) \frac{d}{ds} \left(\frac{\partial f_j}{\partial z^k}(s, z_*) \right) ds, \quad \int_t^\omega \frac{\partial L}{\partial z^k}(s, z_*, \dot{z}_*) ds$$

are ω -periodic and equation (3.7) holds for almost all $t \in \mathbb{R}$.

12. Let g^{ij} stand for the components of the matrix inverse to (g_{ij}) : $g^{ij}g_{ik} = \delta_k^j$. Present equation (3.7) in the form

$$\begin{aligned} & \dot{z}_*^j(t) = g^{kj}(t, z_*(t)) \\ & \cdot \left(\lambda_k + \dot{\gamma}^i(t) \frac{\partial f_i}{\partial z^k}(t, z_*(t)) + \int_t^\omega \dot{\gamma}^i(s) \frac{d}{ds} \left(\frac{\partial f_i}{\partial z^k}(s, z_*(s)) \right) ds \right. \\ & \left. - \int_t^\omega \frac{\partial L}{\partial z^k}(s, z_*(s), \dot{z}_*(s)) ds - a_k(t, z_*(t)) \right). \end{aligned} \quad (3.8)$$

Together with equation (3.8) consider equations

$$\frac{\partial f_j}{\partial t}(t, z_*(t)) + \frac{\partial f_j}{\partial z^k}(t, z_*(t)) \dot{z}_*^k(t) = 0. \quad (3.9)$$

These equations follow from (2.3).

Recall that by the Sobolev embedding theorem, $z_* \in X \subset C(\mathbb{R})$.

Due to (2.4) we have

$$\det B(t, z_*) \neq 0, \quad B(t, z_*) = \left(g^{kj}(t, z_*) \frac{\partial f_i}{\partial z^k}(t, z_*) \frac{\partial f_l}{\partial z^j}(t, z_*) \right)$$

for all t . Substituting \dot{z}_* from (3.8) to (3.9) we can express $\dot{\gamma}^j$ and see $\dot{\gamma}^j \in C(\mathbb{R})$. Thus from (3.8) it follows that $\dot{z}_* \in C(\mathbb{R})$.

Applying this argument again we obtain $\ddot{\gamma}^j, \ddot{z}_* \in C(\mathbb{R})$.

This proves the theorem for the case of odd constraints.

13. Let us discuss the proof of the theorem under the assumption that the constraints are even (2.6).

Definition 7. By Z denote a space of functions $u \in H_{\text{loc}}^1(\mathbb{R})$ such that for all $t \in \mathbb{R}$ the following conditions hold

$$u(-t) = u(t), \quad u(t + \omega) = u(t).$$

The space Z^l is a real Hilbert space with respect to an inner product

$$(u, v)_Z = \sum_{i=1}^l \int_0^\omega (u_i(t)v_i(t) + \dot{u}_i(t)\dot{v}_i(t)) dt.$$

This is the standard inner product in $H^1[0, \omega]$.

So what changed now? The operator A takes the space X^{m+n} onto the space Z^l . The proof of this fact is the same as in Lemma 3.4.

By the Riesz representation theorem, there exists a set of functions $\{\gamma^1, \dots, \gamma^l\} \subset Z$ such that

$$\begin{aligned} b(v) &= \int_0^\omega \dot{\gamma}^j(t) \frac{d}{dt} \left(\frac{\partial f_j}{\partial z^k}(t, z_*) v^k(t) \right) dt \\ &\quad + \int_0^\omega \gamma^j(t) \frac{\partial f_j}{\partial z^k}(t, z_*) v^k(t) dt \end{aligned}$$

for all

$$v = \int_0^t y(s) ds, \quad y \in Y^{m+n}, \quad \int_0^\omega y(s) ds = 0.$$

So that equation (3.7) is replaced with the following one

$$\begin{aligned} &\frac{\partial L}{\partial z^k}(t, z_*, \dot{z}_*) + \int_t^\omega \frac{\partial L}{\partial z^k}(s, z_*, \dot{z}_*) ds \\ &= \dot{\gamma}^j(t) \frac{\partial f_j}{\partial z^k}(t, z_*) + \int_t^\omega \dot{\gamma}^j(s) \frac{d}{ds} \left(\frac{\partial f_j}{\partial z^k}(s, z_*) \right) ds \\ &\quad + \int_t^\omega \gamma^j(s) \frac{\partial f_j}{\partial z^k}(s, z_*) ds + \lambda_k. \end{aligned}$$

Here $k = 1, \dots, m + n$. By the same argument the functions

$$\int_t^\omega \gamma^j(s) \frac{\partial f_j}{\partial z^k}(s, z_*) ds, \quad \int_t^\omega \dot{\gamma}^j(s) \frac{d}{ds} \left(\frac{\partial f_j}{\partial z^k}(s, z_*) \right) ds$$

are ω -periodic and one can put $\alpha^j = \ddot{\gamma}^j - \gamma^j$.

Other argument is the same as above. The theorem is proved.

REFERENCES

- [1] R. A. Adams J.J.F. Fournier: Sobolev Spaces, Elsevier, 2nd Edn. 2003.
- [2] A. Capozzi, D. Fortunato, A. Salvatore: Periodic Solutions of Lagrangian Systems with Bounded Potential, Journal of Mathematical Analysis and Applications, Vol.124, p.482-494 (1987)
- [3] R. Edwards: Functional Analysis. New York, 1965.
- [4] I. Ekeland, R. Tmam, Convex Analysis and Variational Problems, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1999.
- [5] A. Kolmogorov, S. Fomin Elements of the Theory of Functions and Functional Analysis. Ukraine, 1999.
- [6] J. Mawhin, M. Willem, Critical Point Theory and Hamiltonian Systems, Springer-Verlag, New York, 1989.
- [7] M. Struwe, Variational Methods, Springer-Verlag, Berlin, 2008.