

# A computational perspective of the role of Thalamus in cognition

Nima Dehghani<sup>1,2, a)</sup> and Ralf D. Wimmer<sup>3</sup>

<sup>1)</sup>*Department of Physics, Massachusetts Institute of Technology*

<sup>2)</sup>*Center for Brains, Minds and Machines (CBMM), Massachusetts Institute of Technology*

<sup>3)</sup>*Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology*

(Dated: October 18, 2018)

Thalamus has traditionally been considered as only a relay source of cortical inputs, with hierarchically organized cortical circuits serially transforming thalamic signals to cognitively-relevant representations. Given the absence of local excitatory connections within the thalamus, the notion of thalamic ‘relay’ seemed like a reasonable description over the last several decades. Recent advances in experimental approaches and theory provide a broader perspective on the role of the thalamus in cognitively-relevant cortical computations, and suggest that only a subset of thalamic circuit motifs fit the relay description. Here, we discuss this perspective and highlight the potential role for the thalamus – and specifically mediodorsal (MD) nucleus – in dynamic selection of cortical representations through a combination of intrinsic thalamic computations and output signals that change cortical network functional parameters. We suggest that through the contextual modulation of cortical computation, thalamus and cortex jointly optimize the information/cost trade-off in an emergent fashion. We emphasize that coordinated experimental and theoretical efforts will provide a path to understanding the role of the thalamus in cognition, along with an understanding to augment cognitive capacity in health and disease.

Keywords: Thalamo-cortical system, Recurrent Neural Network, Reservoir Computing, Multi-objective Optimization, Cognitive Computing, Artificial Intelligence

## CORTICO-CENTRIC VIEW OF PERCEPTUAL AND COGNITIVE PROCESSING

Until recently, cognition [in mammalian, bird and reptilian nervous system] has been viewed as a cortico-centric process, with thalamus considered to only play the mere role of a relay system. This classic view, much driven by the visual hierarchical model of the mammalian cortex<sup>35</sup>, puts thalamus at the beginning of a feedforward hierarchy. The transmission of information from thalamus to early sensory cortex (V1 in the visual system for example), and the gradual increasing complex representations from V2 to MT/IT and eventually prefrontal cortex (PFC), constitute the core of the perceptual representation under the hierarchical model. A recent comparative study of biologically-plausible Convolutional Neural Networks (CNN) and the visual ventral stream, emphasizes on the feature enrichment through the network of hierarchically interconnected computational modules (layers in the neural network and areas in the ventral stream)<sup>157</sup>.

The strictly static feedforward model has since morphed to a dynamic hierarchical model due to the discoveries of the role of feedback from higher cortical areas to lower cortical areas<sup>52</sup>. These dynamic hierarchies are even considered to be favorable for recursive optimizations, where the overall optimization can be achieved by breaking the problem into smaller ones to find the optimum for each of these smaller problems. Such recursive optimization does not need to be confined to just one

cortical area and each of such distributed optimizations may even be solved differently.<sup>90</sup>. This view of cortical computation is also paralleled with the growing use of recurrent neural networks (RNN) that can capture the dynamics of single neurons or neural population in a variety of tasks. As such, RNNs can mimic (a) context-dependent prefrontal response<sup>89</sup> or (b) can reproduce the temporal scaling of neural responses in medial frontal cortex and caudate nucleus<sup>151</sup>. Although it is clear that higher level cortical feedback reaches all the way down to thalamus (Wimmer et al, other refs), the main attribute of perception/cognition remains cortico-centric under the umbrella of dynamic hierarchical models or RNN embodiment of cortical cognitive functions. Since the computation that is carried by the system should match the computing elements at the appropriate scale<sup>25</sup> a mismatch between these presumed computational systems and the underlying circuitry becomes vividly apparent. First, associative cortex (and not just sensory cortex) receives thalamic input. Second, certain thalamic territories receive their primary (driving) input from the cortex, rather than sensory periphery, some of which are likely to be highly convergent on a single thalamic cell level, suggesting at least a cortical modulation of sensory input. Third, thalamic projections can be broad and diffuse, suggesting a modulatory, rather than a relay function. These points are indicative that thalamus may play a central part in cognitive processes. But what does including thalamus add that could not be achieved in cortical loops? One advantage that the thalamus could bring into the cortical equation is flexibility. For example, the same sensory cue may have different meanings according to a particular context a subject is in. A recent study<sup>118</sup> shows that thalamus may be well suited to

<sup>a)</sup>Electronic mail: [nima.dehghani@mit.edu](mailto:nima.dehghani@mit.edu)

reorganize functional connectivity in frontal cortices in response to such contextual changes, allowing for a more flexible switching of rule to action mapping. We suggest that the unique cognitive capability of the thalamo-cortical system is tightly bound to parallel processing and contextual modulation that are enabled by the diver-

sity of computing nodes (including thalamic and cortical structures) and complexity of the computing architecture (Fig. 1). We will start with a brief overview of the thalamic architecture, followed by experimental evidence and a computational perspective of the thalamic role in contextual cognitive computation.




Figure 1. **Cognitive computing morphospace:** Morphospace of a few example biological and synthetic computing engines in a multidimensional layout. The thalamo-cortical system stands out as a unique system with high cognitive capacity, massive parallel processing and extreme diversity of the computing nodes. Other computing systems occupy less desirable domains of this morphospace. *Logic Gates*: NAND, NOR; *Genetic Logic Gate*: Synthetic biology adaptation of logic gate; *FPGA*: field-programmable gate array (configurable integrated circuit); *CNN*: Convolutional Neural Network; *Physarium Machine*: programmable amorphous biological computer experimentally implemented in slime mould. *Reservoir Computing*: A reservoir of recurrent neurons that dynamically change their activity to nonlinearly map the input to a new space; *DNA computing*: A computing paradigm where many different DNA molecules are used to perform large number of logical computations in parallel; *Connection Machine*: the first commercial supercomputer designed for problems of Artificial Intelligence (AI) using hardware enabled parallel processing. For a multi-dimensional representation of global/local computation, serial/parallel processing and complex/simple computation, see<sup>137</sup>. For an idealized landscape of computation involving the degree of relevance of space, agent (cell) diversity and distributed computing see<sup>138</sup>.

## THALAMIC ARCHITECTURE: ANATOMICAL AND FUNCTIONAL FEATURES

Traditionally, thalamic nuclei (see Fig. 2) are defined as collections of neurons that are segregated by gross fea-

tures such as white matter tracts and various types of tissue staining procedures<sup>59</sup>. This gross anatomical classification has been equated with a functional one, where individual thalamic nuclei giving rise to a set of defined functions<sup>59,60</sup>. More recent fine anatomical studies chal-

lunge this notion, showing that within individual nuclei, single cell input/output connectivity patterns are quite variable.

Thalamus lacks lateral excitatory connections and rather receives inputs from other subcortical structures and/or cortex. In fact, a major feature of forebrain expansion across evolution is the invasion of the thalamus by cortical inputs<sup>45,121</sup>. Most (90-95%) afferents to the relay nuclei are not from the sensory organs<sup>60,133</sup>. Recent anatomical studies have shown a great diversity of cortical input type, strength and inferred degree of convergence, even within individual thalamic nuclei<sup>17</sup> (see Fig. 3 for an example view of cell and network architecture diversity).

Excitatory inputs, mostly, arrive as feedbacks from layer 6 of the cortex as well as from brainstem reticular formation. In addition to the diversity of excitatory inputs, thalamic circuits receive a diverse set of inhibitory inputs (note that local GABAergic thalamic interneurons are mostly absent in non-LGN relay nuclei). The two major systems of inhibitory control are the thalamic reticular nucleus (TRN), a shell of inhibitory nucleus surrounding thalamic excitatory nuclei, and the extra-thalamic inhibitory system; a group of inhibitory projects across the fore-, mid- and hindbrain (see<sup>47</sup> for a review on thalamic inhibition). Perhaps a major differentiating feature of these two systems (TRN and ETI) is temporal precision. One of the key characteristics of thalamus is lack of direct local loops. Only a very small group of inhibitory neurons with local connections exists in thalamus<sup>142</sup>. A mechanistic consequence of this architecture is the differential control of thalamic response gain and selectivity (Fig. 3 F and G), with the TRN controlling the first as observed in sensory systems<sup>111</sup>, and ETI controlling the latter as observed in motor systems<sup>147</sup>. For example, basal ganglia control of thalamic responses, a form of ETI control, would be implemented through thalamic disinhibition, which is not only dependent on ETI input, but also a special type of thalamic conductance that enables high frequency ‘bursting’ upon release from inhibition<sup>29,43</sup>.

Overall, the variety of thalamic inputs (both excitatory and inhibitory), combined with intrinsic thalamic features such as excitability and morphology will determine the type of intrinsic computations that thalamus performs. This view appears to be consistent with recent observations of confidence encoding in both sensory<sup>69</sup> and motor systems<sup>58,84</sup>.

Thalamic relay nuclei mostly project to the cortical middle layers in a topographic fashion. However, the majority of thalamic structures also project more diffusely to the cortical superficial layers, such as mediodorsal (MD), posteriodmedial complex (POm) and pulvinar for example (see Fig. 3 for an example of thalamic cell and circuit diversity). These diffuse projections seem poorly suited to relay information in a precise manner. Rather, they might have a modulatory role of cortical function. Further, a great degree of diversity can be observed at the level of thalamic axonal terminals within the cortex.




Figure 2. **Schematic layout of thalamic nuclei:** Three cross sections of monkey thalamus. *AD*: anterodorsal nucleus; *AM*: anteromedial nucleus; *AV*: anteroventral nucleus; *CM*: centromedian nucleus; *CN*: caudate nucleus; *H*: habenular nucleus; *IL*: intralaminar nuclei; *LD*: lateral dorsal nucleus; *LGN*: lateral geniculate nucleus; *MD*: mediodorsal nucleus; *MGN(D)*: medial geniculate nucleus (dorsal); *MGN(V)*: medial geniculate nucleus (ventral); *PU*: pulvinar; *TRN*: thalamic reticular nucleus (not a relay nucleus); *VA*: ventral anterior nucleus; *VL*: ventral lateral nucleus; *VPI*: ventral posterior nucleus (inferior); *VPL*: ventral posterior nucleus (lateral); *VPM*: ventral posterior nucleus medial. Redrawn from<sup>133</sup>.

While the idea of a thalamic relay was consolidated by observing that the main LGN neurons thought to be associated with form vision (M and P pathways) exhibit spatially compact cortical terminals, recent anatomical studies of individual neurons across the thalamus show a variety of terminal sizes and degree of spatial spread and intricate computational architecture (Fig. 3). This complexity of the architecture and diversity of the computing nodes are among the key factors that set apart the thalamo-cortical system from other conventional and unconventional computing engines (Fig. 1). Part of the complication in understanding how these anatomical types give rise to different functions is their potential for contacting different sets of excitatory and inhibitory cortical neurons.

Specifically, among thalamic nuclei, mediodorsal thalamus (MD) seems to have a connectivity pattern that is distinctively different from the classic sensory nuclei. Cortico-thalamic projections to MD originate from both layers V and VI of PFC<sup>41,44,94</sup>. But, in contrast to relay nuclei, cortical input to MD terminates both extraglomerularly and within the synaptic glomeruli, suggesting that cortex plays a different role in MD activity (in comparison to LGN for example)<sup>127</sup>. Additionally, optogenetic and *in vitro* electrophysiology techniques have

revealed that MD not only projects to the layer I but also has additional terminations in layer III<sup>20</sup>. MD projections to PFC synapse with both excitatory and inhibitory cells<sup>20</sup> where the triggered triggered feedforward inhibition could play a variety of roles from regulating dendritic action potentials<sup>67</sup> to imposing a narrow temporal window within which the excitation can reach the target<sup>21</sup>. These elaborate input and output connectivities of the thalamocortical architecture point to the non-unitary computational role of thalamus. Among thalamic nuclei, specifically MD (and likely other MD-like nuclei) has the architecture necessary to be involved in modulatory computational roles rather than the well-known relay functions. In the next sections, we provide the experimental evidence and algorithmic designs pointing to this modulatory role. This idea of the thalamus controlling cortical state parameters is highlighted in Figs. 4, 5 and the next section.

## MANY FACETS OF THALAMIC COMPUTATION

It is commonly thought that processes like attention, decision making and working memory are implemented through distributed computations across multiple cortical regions<sup>19,93,128</sup>. However, it is unclear how these computations are coordinated to drive relevant behavioral outputs. From an anatomical standpoint, the thalamus is strategically positioned to perform this function, but relatively little is known about its broad functional engagement in cognition. The thalamic cellular composition and network structure constrain how cortex receives and processes information. The thalamus is the major input to the cortex and interactions between the two structures are critical for sensation, action and cognition<sup>61,99,131</sup>. Despite recent studies showing that the mammalian thalamus contains several circuit motifs, each with specific input/output characteristics, the thalamus is traditionally viewed as a relay to or between cortical regions<sup>133</sup>. That the active role of thalamus in cognition is beyond relay stems from a) the fact that sensory input to thalamus is much limited in comparison to input from other structures, such as cortex and basal ganglia, b) the experimental evidence showing number of nuclei modulating cortical neural processing according to behavioral context and c) lesions of certain nuclei such as pulvinar and MD cause severe attention and memory deficits<sup>125</sup>. Below we will discuss how the distinctive anatomical architecture and computational role of pulvinar and MD differ from relay nuclei such as LGN.

It is worth mentioning that this view of bona fide thalamic computations is quite distinct from the one in which thalamic responses reflect their inputs, with only linear changes in response size. This property of reflecting an input (with only slight modification of amplitude) was initially observed in the lateral geniculate nucleus (LGN), which receives inputs from the retina. LGN responses to specific sensory inputs (their receptive fields,


RF) are very similar to those in the retina itself, arguing that there is little intrinsic computation happening in the LGN itself outside of gain control. Success in early vision studies<sup>53,54</sup> might have inadvertently given rise to the LGN relay function being generalized across the thalamus. The strictly feedforward thalamic role in cognition requires reconsideration<sup>48</sup>; only a few thalamic territories receive peripheral sensory inputs and project cortically in a localized manner, as the LGN does<sup>36,59,64,115,131</sup>.

The largest thalamic structures in mammals, the MD and pulvinar contain many neurons that receive convergent cortical inputs and project diffusely across multiple cortical layers and regions<sup>17,122</sup>. For example, the primate pulvinar has both territories that receive topographical, non-convergent inputs from the striate cortex<sup>122</sup> and others that receive convergent inputs from non-striate visual cortical (and frontal) areas<sup>92</sup>. This same thalamic nucleus also receives inputs from the superior colliculus<sup>109</sup>, a subcortical region receiving retinal inputs. This suggests that the pulvinar contains multiple input ‘motifs’ solely based on the diversity of excitatory input. Such input diversity is not limited to the pulvinar, but is seen within many thalamic nuclei across the mammalian forebrain<sup>11</sup>. Local inactivation of pulvinar neurons results in reduced neural activity in primary visual cortex<sup>114</sup> suggesting a feedforward role. Recent studies, however, indicate that pulvinar may have additional roles. For example, pulvinar inactivation was shown to increase low-frequency cortical oscillations<sup>158</sup>. Given that such activity is often associated with inattention and sleep, this study suggested that pulvinar may keep cortical regions in an activated state that would allow responsiveness to top-down input from other areas to modulate ongoing activity according to attentional demands. A different study showed that during perceptual decision making pulvinar neurons encode choice confidence, rather than stimulus category<sup>69</sup>. Together, these recent findings strongly argue for more pulvinar functions beyond relaying information.

In the case of MD, direct sensory input is limited<sup>94</sup> and the diffuse, distributed projections to cortex<sup>73</sup> are poorly suited for information relay; this input/output connectivity suggests different functions. Recent studies<sup>12,118,126</sup> have begun to shed light on the type of computation that MD performs. Taking advantage of the genetic accessibility of the mouse for neuronal manipulations, these studies have revealed that MD coordinates task-relevant activity in the prefrontal cortex (PFC) in a manner analogous to a contextual signal regulating distinct attractor states within a computing reservoir. Specifically, in a task where animals had to keep a rule in mind over a brief delay period (Fig. 4A), PFC neurons show population-level persistent activity following the rule presentation, a sensory cue that instructs the animal to direct its attention to either vision or audition (Fig. 4B,C). MD neurons show responses that are devoid of categorical selectivity (Fig. 4D), yet are critical for selective PFC activity; optogenetic MD inhibition diminishes this activity, while MD

activation augments it. The conclusion is that MD inputs enhance synaptic connectivity among PFC neurons or may adjust the activity of PFC neurons through selective

filtering of the thalamic inputs. In other words, delay-period MD activity maintains rule-selective PFC representations by augmenting local excitatory recurrence<sup>126</sup>.



**Figure 3. Diversity and complexity of thalamo-cortical architectures:** (A) Comparative size of a single RGC (retinal ganglion cell) terminal (red) and an LGN neuron (black); Redrawn from<sup>36</sup>. (B) Complete terminal arbor of a single LGN neuron projecting to V1; Redrawn from<sup>115</sup>. (C) Projection of a TC (thalamo-cortical) neuron to cat motor cortex (with only 23 terminals in TRN versus 1632 terminals in VA/VL); Redrawn from<sup>64,104</sup>. (D,E) Axonal arborization of a single MD neuron (D) and a single POM neuron (E). Principal target layers are: layer I (green), layer II-IV (green-yellow), layer V (blue), layer VI (purple); Redrawn from<sup>73</sup>. Note the comparative size of panels A-E (scale bar at 100  $\mu$ m). (F,G) Synaptic network of thalamo-cortical (Th-cx), Reticular thalamic cell (RE) and local circuit (L-circ) thalamic interneuron in rodents (F) and feline/primates (G). Note that rodents do not have L-circ. Afferent axon excites Th-cx, which in return sends the signal to cortex. RE inhibitory effect on Th-cx cells varies depending on the excitatory drive to each Th-cx cell (F: compare the two neurons on the right versus the one on the left). Axonal collaterals of an RE cell could inhibit another RE cell (G: top RE inhibits the bottom RE), which releases the activity of L-circ leading to inhibition of weakly excited Th-cx (bottom) adjacent to the active Th-cx (top). Panels F and G are redrawn from<sup>143</sup> based on experiments from<sup>141?</sup>.

In a related study, a delayed nonmatching-to-sample T-maze working memory task<sup>12</sup>, it was shown that MD amplification and maintenance of higher PFC activity indicated correct performance during the subsequent choice phase of the task. Interestingly, MD-dependent increased PFC activity was much more pronounced during the later (in delay) rather than earlier part of the task. These findings indicate that PFC might have to recursively pull in MD to sustain cortical representations as working memory weakens with time. Together these studies indicate that PFC cognitive computation can not be dissociated from MD activity. Further evidence for the critical role of the MD-PFC interaction for cognition is the disrupted fronto-thalamic anatomical and functional connectivity seen in neurodevelopmental disorders<sup>91,95,98,107,155</sup>.

### Can MD select cortical subnetworks based on contextual modulation?

Why would a recurrent network (PFC) computation depend on its interaction with a non-recurrent (MD) non-relay network? What computational advantage such system would have? Using a chemogenetic approach, a recent study suggested that information flow in the MD-PFC network can be unidirectional. While both inactivating PFC-to-thalamus and MD-to-cortex pathways impaired recognition of a change in reward value in rats performing a decision making task, only the inactivation of MD-to-cortex pathway had an impact on the behavioral response to a change in action-reward relationship<sup>1</sup>. Given that a sensory stimulus may require a different action depending on the context in which it occurs, the ability to flexibly re-route the active PFC subnetwork to a different output may be crucial. In an architecture like the PFC-MD network, where MD can modulate PFC functional connectivity, MD might well be suited to re-route the ongoing activity in a context dependent manner. In fact, in the mouse cognitive task described above (Fig. 4A), a subset of MD neurons showed substantial spike rate modulation during task engagement compared to when the animals is in its home in cage (see Fig. 4E)<sup>126</sup>. In contrast, PFC neurons show very little difference in spike rates when the animal gets engaged in the task. This suggests that perhaps different subsets of MD neurons are capable of encoding task ‘contexts’, which has been shown experimentally<sup>118</sup>. Subsequently, each given subset could unlock a distinct cortical association; this hypothesis is now experimentally verified<sup>118</sup>. These MD subsets have to be able to shift the cortical states dynamically while maintaining the selectivity based on the subset of cortical connections they target. This idea would also fit with the paradigm shift indicating that thalamic neurons exert dynamical control over information relay to cortex<sup>8,105</sup>.

Overall, the anatomical and neurophysiological data show that the thalamic structure and cortico-thalamic network circuitry, and the interplay between thalamus

and cortex, shape the frame within which thalamus plays the dual role of relay and modulator. Under this framework, different thalamic nuclei carry out multitude of functions including but not limited to information relay. A suggestion of this comparative computational role of LGN, pulvinar and MD is depicted in Fig 8. The importance of (non-relay) thalamic nuclei’s regulatory influence on cortical function is also reflected by the disorders that emerge due to thalamic dysfunctions. Specifically, lesions to pulvinar and MD lead to severe attention and memory deficits<sup>9,124</sup>. The disruption of MD-PFC communication is the likely cause of these cognitive impairment. As mentioned earlier, the back and forth interaction between MD and PFC is necessary during the task acquisition period and is reflected by an increase in (beta frequency) MD-PFC synchrony<sup>106</sup>. In addition, MD also regulates the neural synchrony of PFC neurons<sup>123</sup>. Decreasing MD spiking activity (by hyperpolarizing MD neurons) leads to disrupted MD-PFC synchrony and impaired performance in the delayed non-match to sample task<sup>106</sup>. Moreover, schizophrenics show both a reduced beta and gamma frequency deficit<sup>146</sup>, significant reduction of MD volume<sup>2,112</sup> and total number of MD neurons<sup>112</sup>. While it remains unclear whether the MD loss of neuron is primary or secondary to PFC pathology<sup>112</sup>, it is evident that MD is regulating PFC plasticity, cognitive flexibility<sup>9</sup> and contextual processing<sup>118,126</sup>.

### IS THALAMUS A READ-WRITE MEDIUM FOR CORTICAL PARALLEL PROCESSING?

The connectivity pattern of relay and non-relay point to the dichotomy of algorithmic constrains that are imposed by these specific thalamic structures. There are stark contrasts between sensory (LGN-like) versus non-sensory thalamic nuclei thalamocortical and cortico-thalamic connectivity profiles. Anatomical tracing and radiographic studies have shown that while relay nuclei have preserved topographic focal projections the to middle cortical layer, major thalamic structures (MD, Pulvinar, POm) have a more diffuse projection to the superficial layers of cortex<sup>41,70,94</sup>. Among non-relay nuclei, MD shows interesting characteristics, projecting not only to the layer I but also to the outer banks of layer III<sup>20</sup>. Only a small fraction of MD to PFC projections end in middle cortical layers and more than 90% have modulatory and project diffusely to superficial layers of PFC<sup>149,150,159</sup>. In addition, cortico-thalamic axons to non-relay nuclei show a dual role for cortical influence on thalamus<sup>41,94</sup>. For example, not only layer VI/V PFC project huge number of axons directly to MD but also send collaterals to the reticular thalamic nucleus and indirectly influence MD activity<sup>41,127,132</sup>. These massive reciprocal connectivity is metabolically very costly for the brain. If brain were to operate as a simple pattern matching system, a far less costly feed-forward network (similar to the structure suggested by<sup>157</sup>) would have been more economical. The

complex connectivity of non-relay thalamus and cortex point to computations that are beyond pattern matching. This view also matches what we know of cortical computation, namely that it involves multiple sources of expertise (each decoding certain aspect of the incoming stimuli), along with the fact that these expertise modules operate in a highly parallel mode. To coordinate these parallel yet convergent cortical processing modules, there is a need for a system that is commonly visible to each

module. In addition, collectively, these modules should keep track of the changes in the stimuli in the environment and integrate the information in time. This contextual processing would require small modifications of processed information in individual modules. Or for handling the local constraints, it may need repetition of certain algorithms until the satisfactory results are reached. The connectivity profile that we discussed provides the platform to run such computations.




Figure 4. **MD-PFC interactions during sustained rule representations.** (A) attentional control task design. (B) Example peri-stimulus time histogram (PSTH) for a neuron tuned to attend to vision rule signaled through low pass noise cue (C) Examples showing that rule-specificity is maintained across distinct PFC rule-tuned populations. (D) PSTHs of four MD neurons showing consistent lack of rule specificity. (E) Example rasters and PSTHs of an MD and PFC neuron when the animal is engaged in the task and outside of the behavioral arena. In contrast to PFC, MD neurons show the contextual difference in a change in firing rate. Figure is redrawn from<sup>7</sup>.

If the brain were to function as a simple pattern matching system without wiring and metabolic constraints, evolution would just expand the size and depth of the network to the point that it could potentially memorize a large number of possible patterns. Possibly, evolution would have achieved this approximation of arbitrary patterns by evolving a deep network. This would be a desirable solution since any system can be defined as a polynomial Hamiltonians of low order, which can be accurately approximated by neural networks<sup>82</sup>. But cognition is much more than template matching and classification achieved by a neural network. The limits of template matching methods in dealing with (rotation, translation and scale) invariance in object recognition quickly became known to neuroscientists and in early works on computer vision. One of the early pioneers of AI, Oliver Selfridge, proposed Pandemonium architecture to overcome this issue<sup>129</sup>. Selfridge envisioned serially connected distinct demons (an image demon, followed by a set of parallel feature demons, followed by a set of parallel cognitive demons and eventually a decision demon), that independently perceive parts of the input before reaching a consensus together through a mixture of serial culmination of evidence from parallel processing. This simple feedforward computational pattern recognition model is (in some ways) a predecessor to modern day connectionist feedforward neural networks, much like what we discussed earlier in the text. However, despite its simplicity, Pandemonium was a leap forward in understanding that the intensity of (independent parallel) activity along with a need to a summation inference are the keys to move from simple template matching to a system that has a concept about the processed input. A later extension of this idea was proposed by Allen Newell as the Blackboard model: *“Metaphorically, we can think of a set of workers, all looking at the same blackboard: each is able to read everything that is on it and to judge when he has something worthwhile to add to it. This conception is just that of Selfridge’s Pandemonium: a set of demons independently looking at the total situation and shrieking in proportion to what they see that fits their natures”*<sup>101</sup>. Blackboard AI systems, adapted based on this model, have a common knowledge base (blackboard) that is iteratively updated (written to and read from) by a group of knowledge sources (specialist modules), and a control shell (organizing the updates by knowledge sources)<sup>102</sup>.

Interestingly, this computational metaphor can also be extended to the interaction between thalamus and cortex, though thalamic blackboard is not a passive one as in the blackboard systems<sup>50,96,97</sup>. Although, initially the active blackboard was used as an analogy for LGN computation, the nature of MD connectivity and its communication with cortex seem much more suitable to the type of computations that is enabled by an active blackboard. Starting with an input, thalamus as the common blackboard visible to processing (cortical) modules, initially presents the problem (input) for parallel processing by modules. Here by module, we refer to a group of cortical

neurons that form a functional assembly which may or may not be clustered together (in a column for example). By iteratively reading (via thalamo-cortical projections) from and writing (via cortico-thalamic projections) to this active blackboard, expert pattern recognition modules, gradually refine their initial guess based on their internal processing and the updates of the common knowledge. This process continues until the problem is solved (Fig. 5).

This iterative communication between non-relay thalamus and cortex suggests that cortico-thalamic projections return the results of computations (that was carried in parallel cortical modules) back to the thalamus. The integration of these revisions in the next thalamic output happens via the synaptic input and dendritic arbors of the non-relay thalamic neurons. One of the major differences in synaptic organization of MD from that of the sensory nuclei is that cortical axons target MD neurons both extraglomerularly and within the synaptic glomeruli<sup>127</sup>. In sensory nuclei, these within-glomeruli synaptic sites are particularly designated for receiving major ascending sensory afferents<sup>139</sup>. In MD, such large terminals may even engulf multiple synaptic contacts<sup>110</sup> and are positioned on proximal dendrites of thalamic neurons<sup>127</sup>. These within-glomeruli multi-synaptic structures exhibit fast kinetics, large postsynaptic currents and strong short-term depression<sup>110</sup>. Interestingly, the short-term depression is combined with fast recovery after repetitive stimulation<sup>110</sup>, enabling generation of synaptic activity patterns that can match the frequency of cortico-thalamic inputs arriving via these large terminals<sup>140</sup>. As a result, these potent synaptic structures provide the platform for PFC inputs to play a much more active role in shaping the thalamic response in relay (MD) nuclei in comparison to the role that cortical feedback to sensory thalamic nuclei may play<sup>127</sup>.

Distinctive biophysical characteristics of non-relay thalamocortical projections play a complementary role in the computational scheme echoing an active blackboard. Interestingly, activating MD does not generate spikes across a population of prefrontal cortical neurons they project to, while activating LGN generates spikes in primary visual cortex<sup>118,126</sup>. Instead, MD activation results in overall enhancement of inhibitory tone, coupled with enhanced local recurrent connectivity within the PFC. Although thalamocortical feed-forward inhibition is also observed in somatosensory barrel cortex in response to thalamic stimulation<sup>40</sup>, MD-evoked inhibition in PFC exerts a more powerful inhibitory gain control<sup>27</sup>. This difference must be rooted in the particular cortical target pattern of MD projections. Specifically, MD directly targets parvalbumin-positive PV (and not somatostatin-SOM) interneurons in layer I and III<sup>18,27,74,75,120</sup>, while (for example) VM only “weakly” activates variety of layer I interneurons<sup>20</sup>. In fact, when SOM interneurons are silenced, MD-evoked feedforward inhibition is enhanced<sup>27</sup>. As a result, while VM plays a more important role in excitation/inhibition balance, MD plays the modulator



role via varying the integration time window and temporal precision of cortical responses<sup>18</sup>. While individual pyramidal neurons harbor a broad response dynamics, the feedforward inhibition regulates the population dynamic via graded recruitment of individual neurons<sup>66</sup>. Increased conductance, noisy voltage fluctuations, and depolarization are the not-necessarily exclusive factors that define how the changes in the background input, i.e. stimulus changes and their contextual relevance, affect the gain<sup>16,113</sup>. In addition, while sensory thalamocortical synapses (i.e. LGN to visual cortex) onto fast-spiking inhibitory neurons manifest much higher release probability than those onto pyramidal cells, MD projections show similar presynaptic release probability among the two inhibitory and excitatory cortical neurons<sup>27</sup>. The co-variation of excitatory and feed-forward inhibitory response sets the control for graded recruitment of pyramidal neurons into population response<sup>66</sup>. This control in itself is evoked by prior cortical excitation of MD, and as a result, the altered activity of PV interneurons in PFC can bias the response towards passive vs flexible contextual processing in a manner that is distinctively different from the observed response of the sensory cortices<sup>27</sup>. These mechanisms show us why the chemogenetic inhibition of MD leads to impaired working memory and flexible goal-directed behaviors<sup>106,107</sup>. Similarly, schizophrenics show reduced MD-PFC functional coupling<sup>95</sup> and deficits in prefrontal PV interneurons<sup>81</sup>, highlighting the importance of the modulatory effect of MD on PFC function<sup>27</sup>.

## I. COMPUTATIONAL AND METABOLIC CONSTRAINTS

To process the changing stimuli and altering contextual cues, and in order to achieve cognitive flexibility, the thalamocortical system needs to harbor temporal buffering. The mechanisms that we described here point to the ways in which such buffer may take place, namely: a) changes in the stimuli/context are constantly reprocessed by the cortex and the outputs are rewritten to thalamus, b) thalamus constantly reshapes the cortical population dynamics. MD is changing the mode by which PFC neurons interact with one another, initiating and updating different attractor dynamics underlying distinct cognitive inputs. As a result, the thalamocortical system, collectively and at any instant, keeps an updated description of the stimulus/context over some computational cycles up to the present (Fig. 5). Perhaps the upper bound of these computational cycles is tightly bound to the particulars of the cortico-thalamic and thalamocortical connectivity and biophysical constraints. However, since we are dealing with a biological system with finite resources, this back and forth communication needs to have certain characteristics to provide a viable computational solution. First and foremost, the control of interaction and its scheduling has to have a plausible biological component and should bind solutions as time

evolves. Second, to avoid turning into an NP-hard (non-deterministic polynomial-time hardness) problem, there must exist a mechanism that stops this iterative computation once an approximation has been reached (Fig. 5). Here, we propose a specific solution to the first problem and a plausible one for the later issue. We suggest that phase-dependent contextual modulation serves to deal with the first issue and a multi-objective optimization of efficiency (computational information gain) and economy (computational cost, i.e. metabolic needs and the required time for computation) handles the second issue (Fig. 6). In both cases, we suggest that thalamus plays an integral role in conjunction with cortex.

### Computational constraints and the role of thalamus in phase-dependent contextual modulation

As mentioned earlier, we know that hierarchical convolutional neural networks (HCNN), which can recapitulate certain properties of static hierarchical forward models, can not capture any processes that need to store prior states<sup>156</sup>. As a result, context-dependent processing can be extremely hard to implement in neural network models<sup>117</sup>. The most widely used ANNs (Feedforward nets, i.e. multilayer perceptrons/Deep Learning algorithms) face fundamental deficiencies: the ubiquitous training algorithms (such as back-propagation), i) have no biophysical plausibility, ii) have high computational cost (number of operations and speed), and iii) require millions of examples for proper adjustment of the connections' weights. These features render feedforward NNs not suitable for temporal information processing. In contrast, recurrent neural networks (RNNs) can universally approximate the state of dynamical systems<sup>38</sup>, and because of their dynamical memory are well suited for contextual computation. If the higher cortical areas were to show some features of RNN-like networks, as manifested by the dynamical response of single neurons<sup>89</sup>, then we anticipate that the local computation (interaction between neighboring neurons) to be mostly driven by external biases. The thalamic projections could then play the role of bias where they seed the state of the network. From both anatomical studies and electrophysiological investigations<sup>46</sup>, we know that thalamus is at a prime position to modify the signal based on the cognitive processing that is happening in the cortex<sup>12,126</sup>. This thalamic-driven regulation entails "binding in time" since MD-like thalamus modifies its output cortex at a given time and is itself influenced by what is perceived by the cortex in time prior. But how can the "binding in time" avoid locking-in the thalamic function to a set of inputs at a given time? How can thalamus constantly be both ahead of cortex and yet keep track of the past information? The secret may be embedded in the non-recurrent intrinsic structure of thalamus, the recurrent structure of

the higher cortical areas, , and the phase-sensitive detec-


To expand the idea further, let's revisit some core attributes of cognitive processing. Based on the observations of behavior, higher cognition requires "efficient computation", "time delay feedback", the capacity to "retain information" and "contextual" computational properties. Such computational cognitive process surpass the computational capacity of simple RNN-like networks. The essential required properties of a complex cognitive system of such kind are: 1) input should be nonlinearly mapped onto the high-dimensional state, while different inputs map onto different states, 2) slightly different states should map onto identical targets, 3) only recent past should influence the state and network is essentially unaware of remote past, 4) a phase-locked loop should decode information that is already encoded in time and 5) the combination of 1-4, should optimize sensory processing based on the context. The first three attributes of such system have close relevance to constrains and computational properties of higher cortical areas (prefrontal). The same three are also the main features of reservoir computing, namely "separation property", "approximation property" and "fading memory"<sup>56,57,85,86</sup>. Interestingly, and RC system can "non-linearly" map a lower dimensional system to a high-dimensional space facilitating classification of the elements of the low-dimensional space. The last two properties match the structure and computational constraints of non-relay thalamic system as a contextual modulator that is phasically changing the input to the RC system. In fact, in an RC model of prefrontal cortex, addition of a phase neuron significantly improved the networks performance in complex cognitive tasks. The phase neuron improves the performance by generating input driven attractor dynamics that best matched the input<sup>33</sup>. This advantageous phase-based bias effect is not limited to the simulation or physiological RC-like neural circuitry. In a recent study, electronic implementation and numerical studies of a limited RC system of a single nonlinear node with delayed feedback has shown efficient information processing<sup>3</sup>. Such reservoir's transient dynamical response follows delay-dynamical systems, and only a limited set of parameters are required to set the rich dynamical properties of delay systems<sup>55</sup>. This system was able to effectively process time-dependent signals.

The phase neuron<sup>33</sup> and delayed dynamical RC<sup>3</sup> both show properties that resemble the structure-function of MD-like thalamus as discussed here. Specifically, the phasic recruitment of cortical neurons is invoked due to the combination of cortical influence on non-relay thalamic neurons through direct and indirect corticothalamic projections (see reticular nucleus inhibitory influence on thalamic neurons, Fig.??). In a given cycle

of computation (Fig. 5), cortical feedback leads to the release of GABA (via reticular nucleus) in MD<sup>68</sup>. It has been shown that the increased GABA alters the opening of T-type Ca<sup>2+</sup> channels<sup>22</sup>, which in the case of MD, results in enhanced MD-PFC interaction; yielding mutual drive of the corticothalamic and thalamocortical activity together<sup>68</sup>. Through this calcium-based low-threshold spiking, the gradual synchronization of MD and PFC ensues<sup>62,68</sup>. As a result, MD units and PFC show strong phase-locked synchrony<sup>106</sup>. This gradual phase-locking mechanism forms the basis of the temporal dynamics that can nonlinearly (through consecutive cycles of computation) change the cortical activity as a result of novel stimuli or unexpected contextual changes as observed experimentally<sup>12,126</sup>. In fact, abnormal activity of T-type Ca<sup>2+</sup> channels in MD, leads to hypersynchrony in PFC neurons and frontal lobe-specific seizures<sup>68</sup>. The interactions between non-relay thalamus and cortex, collectively, is neither feedforward, nor locally recurrent, but it has a mixture of non-recurrent phase encoder that keeps copies of the past processing and modulates the sensory input relay and its next step processing (Fig. 5). The distinctive short-term dynamics are well matched with the divergent structure-function relationship of sensory and non-relay MD-like thalamic nuclei. These features further emphasize that the perceptual and cognitive processing can not be solely cortico-centric operations.

## BIOLOGICAL CONSTRAINS AND THE ROLE OF THALAMUS IN COMPUTATIONAL OPTIMIZATION


Computation and optimization are two sides of the same coin. But how does the brain optimize the computations that would match its required objective, i.e. cognitive processing? There is a current trend of thinking that brain optimizes some arbitrary functions, with the hopes that the future discovery of these unknown functions may guide us to establish a link between brain's operations and deep learning<sup>90</sup>. This line of approach to optimizational (and computational) operations of the brain has few flaws. First, it avoids specifying what function the brain is supposed to optimize (and as a result it remains vague). Second, it refrains from addressing certain limitations that brain has to cope with due to biological constrains. First of these limitations is the importance of using just enough resources to solve the current perceptual problem. Second is the necessity to come up with a solution just in (the needed) time. The importance of "just-enough" and "just-in-time" computation in



**Figure 5. Schematic representation of thalamic cognitive contextual computation.** (A): In the case of static data, a set of function/weight modules can yield good classification. Function represents a polynomial (since any system that is known to be a polynomial Hamiltonians of low order can be accurately approximated by neural networks<sup>82</sup>) and the weights exemplify the connection matrix of an artificial neural network encapsulating this polynomial. Stacking multiple of such module can increase the accuracy of polynomial approximation (such as in the case of CNN). (B): Thalamo-cortical computation for contextual processing of dynamic data. Each dataframe is processed by a weight/pointer module (thalamus MD-like structure) which like a blackboard is writable by different sets of neuronal assemblies in cortex. Thalamic pointers assign the assemblies; modules' weights adjust the influence of each assembly in further computational step [inset C shows a non-recurrent thalamic nuclei (MD-like) modulating the weights in the PFC (reservoir and readout). Here, depending on the context (blue or red), the interactions between MD and Reservoir, between Reservoir and Readout, and between Readout and MD could pursue one of the two possible outcomes. Specifically, MD changes the weights in the Reservoir to differentially set assemblies that produce two different attractor states, each leading to one of the two possible network outputs]. In (B, C), each operation of the thalamic module is itself influenced, not only by the current frame ( $t$ ), but also by the computation carried by cortex module on the prior frame ( $t - 1$ ). Cortical module is composed of multiple assemblies where each operate similar to the function/weight module of the static case. These assemblies are locally recurrent and each cell may be recruited to a different assembly during each operation. This mechanism could explain why prefrontal cells show mixed selectivity in their responses to stimuli (as reported in<sup>39,116</sup>). Through this recursive interaction between thalamus and cortex, cognition emerges not as just a pattern matching computation, but through contextual computation of dynamic data (bottom right schematic drawing).

cortical computation should not be overlooked<sup>30</sup>. If the first condition is not met, the organism can not sustain

continued activity since the metabolic demand surpasses the dedicated energetic expenditure and the animal can



**Figure 6. Dynamic role of thalamo-cortical system in the information/cost optimization.** (A) Iso-maps of information (A1) and cost (A2) in the domain specified by  $\omega$  and  $\lambda$  (functions of cortical and thalamic activity). Information across each Iso-quant curve ( $\eta_1$  for example) is constant and is achieved at a certain mixture of  $\omega$  and  $\lambda$ . Optimal information can be obtained by moving outward (arrows, A1). Cost optimization can be achieved by moving inward (A2). (B) since information and cost are both defined in the domain of  $\omega$  and  $\lambda$ , thalamus and cortex jointly contribute to information and cost optimization. The points where the iso-quant curves' tangents are equal (black dashed line), provide the optimal combination of information/cost (green curve). In any cycle of cognitive operation, depending on the prior state of the system ( $\omega$  and  $\lambda$ ), the nearest points on the green curve are the optimal solutions for ending that cycle. (C) mapping of the optima curve to information/cost domain shows all pareto efficient allocations (cyan curve). The slope of the pareto frontier shows how the system trades cost versus information: along the pareto curve, efficiency is constant but the exchange between information and cost is not. All allocations inside of this curve could be improved as thalamus and cortex interact. The grey zone shows the biophysically not-permissible allocation of computation and resource.

not survive. In fact, the communication in neural networks are highly constrained by number of factors, specifically the energetic demands of the network operations<sup>78</sup>.

From estimates of the cost of cortical computation<sup>79</sup>, we know that the high cost of spiking forces the brain to rely on sparse communication and using only a small fraction of the available neurons<sup>5,134</sup>. While, theoretically, cortex can dedicate a large number of neurons (and very high dynamical space) to solve any cognitive task, metabolic demand of such high-energetic neural activity renders such mechanism highly inefficient. As a result, the “law of diminishing returns” dictates that increased energetic cost causing excessive pooling of active neurons to an assembly would be penalized<sup>103</sup>. The penalization for unnecessary high-energetic neural activity, in itself, should be driven by the nature of computation rather than being formulated as a fixed arbitrary threshold imposed by an external observer. On the other hand, a system can resort to low-cost computation at any given time but dedicate long enough time to solve the task on hand. Naturally, such system would not be very relevant to the biological systems since time is of essence. If an animal dedicates a long instance of its computational capacity to solve a problem, the environment has changed before it reaches a solution and the solution becomes obsolete. A deer would never have an advantage for its brain to have fully analyzed the visual scene instead of spotting the approaching wolf and shifting resources to the most-needed task, i.e. escape. As a result, many of the optimization techniques and concepts that may be relevant to artificial neural networks are irrelevant to embodied computational cognition of the brain. The optimization that the brain requires is not aiming for the best possible performance, but rather needs to reach a good mixture of economy and efficiency.

Not surprisingly, these constrains, i.e. efficiency and economy, are cornerstones of homeostasis and are observed across many scales in living systems<sup>145</sup>. The simple “Integral feedback” acts as the mainstay of control feedback in such homeostatic systems (such as E Coli heat-shock or DNA repair after exposure to gamma radiation)<sup>26,31,32,71</sup>. Change in input leads to change in the output and the proportional change in the controller aiming to reset the output to the desired regime. When the integral feedback is disrupted, the system can no longer reach proper homeostasis and either efficiency or economy (or even both) will be sub-optimal<sup>31,32,145</sup>. Many different etiologies could be behind the integral feedback disruption, but the outcome is loss of robust response in uncertain environments. The presence of feed-forward and feedback loops provide the means for robust and fast operation in processing fluctuating incoming inputs. This feedback regulation and operational robustness has an energetic and computational cost for the system. Although for simple systems it is feasible to associate the exact cost of an operation to the overall computational cost of the system, scaling the metabolic cost of feedback regulations to large networks remains a challenge since it will involve multiple feedback loops, non-linear dynamics and numerous uncertain parameters<sup>23</sup>. Specifically in the case of a single neurons, branching

architecture, non-uniform ion channel distributions and conduction states of action potentials affect the rate of energy consumption<sup>63</sup>. However, this electrochemical energy of single neuron operation does not linearly scale to the spent energy at networks level<sup>152,153</sup>. The total energy function of neural populations will depend not only on the energy function of single neurons and their coupling in a given neural population, but also on the flow of information between different populations<sup>152,154</sup>. When a large pool of neurons is recruited to form multiple assemblies to perform a certain computation, it is the interactions between the assemblies that will define the collective behavior of the discrete components. Since the coupled processes show additive entropy productions<sup>28</sup>, the total energetic optimality of the desired function would depend on the feedback loops between the assemblies and how these feedbacks control the intrinsic energy expenditure of a given assembly. These attributes are inline with the general principle of modular composition of biological systems<sup>24,25,51</sup>. From the dynamical systems' perspective, to understand the operational principles (here, of large assembly of neurons), we do not need to strip down the assembly to its individual component level (here, individual neurons)<sup>25</sup>. As a result the optimal control at the functional scale of modules where the interaction between the system's modules take place<sup>23,25</sup>.

The constraints that we discussed above, directly translate to the computational operations of thalamocortical system as we discussed. Instead of just trying to deal with one fitness function at a time (where the minima of the landscape would be deemed as "the" optima), the brain has to perform a multi-objective optimization, finding solutions to both metabolic cost (economy) and just-in-time (efficiency) computation. Thus we can infer that a unique solution does not exist for such a problem. Rather, any optimization for computational efficiency will cost us economy and any optimization for economy will cost us efficiency. In such case, a multi-objective optimization pareto frontier is desirable. Pareto frontier of information/cost will be the set of solutions where any other point in the space is objectively worse for both of the objectives<sup>42,72,145</sup>. As a result, the optimization mechanism should push the system to this frontier. The iterative dynamical interaction between thalamus and cortex seems to provide an elegant solution for this problem (We discuss this in more details below).

In addition to these theoretical rationals, we also wish to point to some observations that support the emergent optimization in the thalamo-cortical system. For example, metabolic studies have shown that following thalamic injuries, a misbalance in cortical metabolism ensues<sup>6,7,77,80</sup>. Moreover, in healthy humans (and not in mood disorder patients), the metabolic rate of thalamus directly relates to the power of cortical oscillations<sup>83</sup>. The misbalance in cortical metabolism have been observed in variety of nuclei damages, but are specially pronounced in mediodorsal, centre median or pulvinar injuries<sup>6,7</sup>. In addition, low-frequency and high-

frequency stimulation of MD can induce long-term depression/potential or in mPFC (medial PFC); however, the exact sign and magnitude of the differential modulation of thalamo-prefrontal functions under low and high input drive depends on the lack or presence of Muscarinic and Nicotinic modulation<sup>14</sup>. Just as thalamic injury/modulation can change the cortical activity and metabolism, cortical injuries (due to stroke for example) can cause an attenuation of the excitatory feedback to thalamus and lead to thalamo-cortical dysrhythmia<sup>148</sup>. Regardless of where the initial injury has occurred, the disrupted thalamo-cortical interaction is conjoined with a misbalance in metabolism. The resultant out of balance activity leads to cognitive disorders that can happen in form of disrupted information processing due to cortical hypersynchrony as a results of excessive thalamic spiking<sup>68</sup> or faulty modulation of sensory signals and loss of the normal correlation between glucose metabolism in the thalamus and PFC<sup>15,65</sup>. The exact cellular/subcellular mechanism that lies beneath the joint fluctuations of firing and metabolic of cortex and thalamus is not very well understood and number of mechanisms may act (not necessarily exclusively). For example, reduced cortical feedback may lead to thalamic hyperpolarization, and the resultant de-inactivation of voltage-gated T-type Ca channels may cause the neurons to switch from tonic spiking to a pathological bursting<sup>148</sup>. Or it could be that the thalamic drive of the inhibitory neurons in the cortex not only directly affect the cortical mode of firing<sup>34</sup> but also change the glycogenolysis in astrocytes through Vasoactive intestinal peptide (VIP) interneurons<sup>87,88</sup>. Interestingly, and in contrast to the noradrenergic afferent fibers that span horizontally across cortical domains, VIP neurons have a bipolar architecture and therefore their effect is spatially limited<sup>88</sup>, likely correlated to the size of the functional assemblies that are recruited to perform a computational task. Whichever the exact mechanism at the cellular level is, the collective activity of modulatory thalamus and cortex drives the optimization that inherently can not be controlled by the information available at the scale of single neurons and solely in cortex.

To formalize multiobjective optimization, consider a set of functions,  $\omega$  and  $\lambda$  of  $f_{Th}$  (firing rate of thalamic cell) and  $f_{Cx}$  (firing rate of cortical cells). Uncertainty (or its opposite, information) and computational cost (a mixture of time and metabolic expense) can both be mapped to this functional space of  $\omega(f_{Th}, f_{Cx})$  and  $\lambda(f_{Th}, f_{Cx})$  (Fig. 6A1,2). Let's define computational cost and information as product and linear sum of cortical and thalamic activity ( $\alpha f_{Th}^n \cdot \beta f_{Cx}^n, \theta \frac{d_{f_{Th}}^n}{dt} + \psi \frac{d_{f_{Cx}}^n}{dt}$ ; with  $\alpha, \beta, \theta, \psi$  as coefficients) to reflect the logarithmic nature of information (entropy) and the fact that biological cost is an accelerating function of the cost-inducing variables<sup>26</sup>. The hypothetical space of cost/information is depicted in Fig. 6, where top panels show indifference maps of information (A1) and cost (A2). The example simulations and parametric plots of the cost and infor-

mation functions defined as above are shown in Fig. 7. In each indifference map, along each iso-quant curve, the total functional attribute is the same. For example, anywhere on the  $\eta_1$  curve, the uncertainty (or information) in our computational engine is the same. However, different iso-quant curves represent different levels of the functional attribute. For example, moving outward increases information (reduces uncertainty) as  $\eta_1 < \eta_2 < \eta_3 < \eta_4$  and thus if computational cost was not a constrain, the optimal solution would have existed on  $\eta_4$  or further away (Fig.6 A1). In contrast, moving inward would preserve the cost ( $\zeta_1 < \zeta_2 < \zeta_3 < \zeta_4$ ) if the computational engine did not have the objective of reducing uncertainty (Fig. 6A2). Since information and cost are interdependent and both depend on the interaction between thalamus and cortex, we suggest that information/cost optimization happens through an iterative interaction between thalamus and cortex (note the blackboard analogy and contextual modulation discussed above). Since we defined both information and cost as a set of iso-quant curves in the functional space of  $\omega(f_{Th}, f_{Cx})$  and  $\lambda(f_{Th}, f_{Cx})$ , they can be co-represented in the same space (Fig. 6B). Optimal solutions for information/cost optimizations are simply the solutions to where the tangents of the iso-quant curves are equal (see the tangents [black dashed lines] and points A, B, C and D, in Fig. 6B). These points create a set of optimal solutions for the tradeoff between information and cost (green curve). Mapping of the optimal solutions to the computational efficiency space  $E$ , gives us the pareto efficient curve (cyan curve, Fig. 6C). Anywhere inside the curve is not pareto efficient (i.e. information gain and computational cost can change in such a way that, collectively, the system can be in a better state (on the pareto curve). Points outside of the pareto efficient curve are not available to the current state of the system due to the coefficients of  $\omega$  and  $\lambda$ . A change in these coefficients can potentially shape a different co-representation of information and cost (see Fig 7, top row for 3 different instances of  $\omega$  and  $\lambda$  based on different  $\alpha, \beta, \theta, \psi$  values), and thus a different pareto efficient curve (see Fig 7, bottom row). These different possible pareto frontiers can be set based on the prior state of the system and the complexity of the computational problem on hand. For example, the modulatory MD-like thalamic triggering of feedforward inhibition of layer I interneurons and layers II/III pyramidal<sup>20</sup> may tune the cortical activity to a sustained profile of arousal during wakefulness<sup>49</sup>. Or in contrast to relay thalamus where maximum responsiveness to transient signals (such as sensory stimuli onset/offset) is needed<sup>13,119</sup>, the MD-like modulatory drive may be invoked for tasks where working memory and contextual processing are needed<sup>27,106</sup>. Nonetheless, the computational efficiency of the system can not be infinitely pushed outward because of the system's intrinsic biophysical constrains (neurons and their wiring). The shaded region in Fig 7, bottom row, shows this non-permissible zone.

In the defined computational efficiency space  $E$ ,




Figure 7. **Dynamic parameter space of thalamo-cortical joint optimization of information/cost.** Three different realization of information/cost interaction as a function of thalamic and cortical activity ( $\omega, \lambda$ ) and the corresponding pareto curves (see Fig. 6 for details of this optimization construct). Pareto curve shows the optimal set of both cost and information that can be obtained given the biophysical constrains of neurons and networks connecting them. Every point on the pareto frontier shows technically efficient levels for a given parameter set of  $\omega, \lambda$  (see text for more details). All the points inside the curve are feasible but are not maximally efficient. The slope (marginal rate of transformation between cost and information) shows how in order to increase information, cost has to change. The dynamic nature of interaction between thalamus and cortex enables an emergent optimization of information/cost depending on the computational problem on hand and the prior state of the system.

composed of the two variables information and cost (as the objective functions, shown in bottom panels of Fig. 6 and Fig. 7), solving a computational problem is represented by a decrease in uncertainty. However, any change in uncertainty has an associated cost. First derivative of the pareto frontier shows “marginal rate of substitution” as  $\frac{\Delta_{info}}{\Delta_{cost}}$ . This ratio varies among different points on the pareto efficient curve. If we take two points on the pareto curve in the computational efficiency space, such as A and C for example, computational efficiency of these two points are equal  $E_A(\eta_1, \lambda_4) = E_C(\eta_3, \lambda_2)$ . The change in efficiency of point A with respect to information and cost, are the partial derivatives  $\frac{\partial E_A}{\partial info}$  and  $\frac{\partial E_A}{\partial cost}$ , respectively. As a result,  $\frac{\partial E_A}{\partial info} d_{info} + \frac{\partial E_A}{\partial cost} d_{cost} = 0$ , meaning that there is constant efficiency along the pareto curve, the tradeoff between information and cost is not constant. The optimization in this space is not based on some fixed built-in algorithm or arbitrary thresholds by an external observer. Rather, information/cost optimization is the result of back and forth interaction between thalamus and cortex. Based on the computational perspective

that we have portrayed, thalamus seems to be poised to operate as an optimizer. Thalamus receives a copy of (sensory) input while relaying it, and receives an efferent copy from the processor (cortex), while trying to efficiently bind the information from past and present and sending it back to cortex. The outcome of such emergent optimization, is a pareto front in the economy-efficiency landscape (Fig. 6,7). If the cortex were to be the sole conductor of cognitive processing, the dynamics of the relay and cortical processing would meander in the parameter space and not yielding any optimization that can provide a feasible solution to economic and just-in-time computation. Such system is doomed to fail, either due to metabolic costs or due to computational freeze over time ; thus more or less be a useless cognitive engine. In contrast, with the help of an optimizer that acts as a contextual modulator, the acceptable parameters will be confined to a manifold within the parameter space. Such regime would be a sustainable and favorable domain for cognitive computing. This property shows another important facet of a thalamo-cortical computational cognitive system and the need to move passed the cortico-centric view of cognition. An important consequence of this formalization is that it provides us testable hypotheses for objectively evaluating information and cost optimization. By careful simultaneous measurements of thalamic and cortical collective activity, during different states and under different neurotransmitter modulatory effects, one should be able to examine the distinctive interaction of cortex and MD-like versus relay thalamic nuclei. Although we wish to emphasize that while information processing is a fundamentally energy-consuming process<sup>10,108</sup> and one can drive theoretical estimates of the energetic cost of the activity of a population of neurons, the exact translation of bit to watts in adaptive information processing systems (such as thalamocortical) can only be verified experimentally<sup>37</sup>. Without proper and careful measurements, it is impossible to predict how much more reliable the collective computation could get at the expense of energy<sup>4</sup>. Likewise, the degree to which the energy is traded for accuracy/speed (or their combination) will be a hard challenge for the experimentalists measuring the collective activity<sup>76</sup>.

### CONCLUDING REMARKS: REFRAMING THALAMIC FUNCTION ABOVE AND BEYOND INFORMATION RELAY

Lately, new evidence about the possible role of thalamus has started to challenged the cortico-centric view of perception/cognition. Anatomical studies and physiological measurements have begun to unravel the importance of the Cortico-Thalamo-Cortical loops in cognitive

processes<sup>8?</sup>. Under this emerging paradigm, thalamus plays two distinctive roles: a) information relay, b) modulation of cortical function<sup>133</sup>, where the neocortex does not work in isolation but is largely dependent on thalamus. In contrast to cortical networks which operate as specialized memory devices via their local recurrent excitatory connections, the thalamus is devoid of local connections, and is instead optimized for capturing state information that is distributed across multiple cortical nodes while animals are engaged in context-dependent task switching<sup>126</sup>. This allows the thalamus to explicitly represent task context (corresponding to different combinations of cortical states), and through its unique projection patterns to the cortex, different thalamic inputs modify the effective connections between cortical neurons<sup>12,126</sup>.

Here, we started with a brief overview of the architecture of thalamus, the back and forth communication between thalamus and cortex, then we provided the electrophysiological evidence of thalamic modulatory function, and concluded with a computational frame that encapsulates the architectural and functional attributes of the thalamic role in cognition. In such frame, the computational efficiency of the cognitive computing machinery is achieved through iterative interactions between thalamus and cortex embedded in the hierarchical organization (Fig. 4, 5). Under this emergent view, thalamus serves not only as relay, but also as a read/write medium for cortical processing , playing a crucial role in contextual modulation of cognition (Fig. 8). Such multiscale organization of computational processes is a necessary requirement for design of the intelligent systems<sup>24,135,136</sup>. Distributed computing in biological systems in most cases operates without central control<sup>100</sup>. This is well reflected in the computational perspective that we discussed here. We suggest that through the continuous contextual modulation of cortical activity, thalamus (along with cortex) plays a significant role in emergent optimization of computational efficiency and computational cost. This phenomenon has a deep relation with phase transitions in complex networks. Different states (phases) of the network are associated with the connectivity of the computing elements (see thalamic weight/pointer and cortical function/weight modules in Fig. 5). Interestingly, intrinsic properties of the complex networks do not define the phase transitions in system. Rather, the interplay of the system with its external environment shapes the landscape where phase transitions occur<sup>130</sup>. This parallel in well-studied physical systems and neuronal networks of thalamo-cortical system show the importance of the interplay between thalamus and cortex in cognitive computation and optimization. The proposed frame for contextual cognitive computation and the emergent information/cost optimization in thalamo-cortical system can guide us in designing novel AI architecture.




Figure 8. **The emergent view of thalamic role in cognition.** (Top) In the traditional view, serial processing of information confines the role of thalamus to only a relay station. (Bottom) the view that is discussed in this manuscript considers thalamus as a key player in cognition, above and beyond relay to sensory cortices. Through combining the efferent readout from cortex with sensory afferent, MD-like thalamic nuclei modulate further activity of the higher cortex. The contextual modulation enabled by MD is composed of distinctively parallel operations (individual circles represent the non-recurrent nature of these processes due to lack of local excitatory connections). Under this view, and the computational operatives discussed here, the thalamo-cortical system (and not just cortex) is in charge of contextual cognitive computing. The computation enabled by Pulvinar/PO like nuclei is different from LGN and also from MD-like nuclei.

## ACKNOWLEDGMENTS

We wish to thank Michael Halassa for helpful discussions.

## REFERENCES

- <sup>1</sup>Alcaraz, F., Fresno, V., Marchand, A. R., Kremer, E. J., Couverture, E., and Wolff, M. (2018). Thalamocortical and corticothalamic pathways differentially contribute to goal-directed behaviors in the rat. *eLife*, 7.
- <sup>2</sup>Alelú-Paz, R. and Giménez-Amaya, J. M. (2008). The mediodorsal thalamic nucleus and schizophrenia. *Journal of psychiatry & neuroscience: JPN*, 33(6):489.
- <sup>3</sup>Appeltant, L., Soriano, M. C., Van der Sande, G., Danckaert, J., Massar, S., Dambre, j., Schrauwen, B., Mirasso, C. R., and Fischer, I. (2011). Information processing using a single dynamical node as complex system. *Nature Communications*, 2:468.
- <sup>4</sup>Ay, N., Flack, J., and Krakauer, D. C. (2007). Robustness and complexity co-constructed in multimodal signalling networks. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 362(1479):441–447.
- <sup>5</sup>Baddeley, R., Abbott, L. F., Booth, M. C. A., Sengpiel, F., Freeman, T., Wakeman, E. A., and Rolls, E. T. (1997). Responses of neurons in primary and inferior temporal visual cortices to natural scenes. *Proceedings of the Royal Society B: Biological Sciences*, 264(1389):1775–1783.
- <sup>6</sup>Baron, J., D’antona, R., Pantano, P., Serdaru, M., Samson, Y., and Bousser, M. (1986). Effects of thalamic stroke on energy metabolism of the cerebral cortex: a positron tomography study in man. *Brain*, 109(6):1243–1259.
- <sup>7</sup>Baron, J., Levasseur, M., Mazoyer, B., Legault-Demare, F., Mauguiere, F., Pappata, S., Jedynak, P., Derome, P., Cambier, J., and Tran-Dinh, S. (1992). Thalamocortical diaschisis: positron emission tomography in humans. *Journal of Neurology, Neurosurgery & Psychiatry*, 55(10):935–942.
- <sup>8</sup>Basso, M. A., Uhlrich, D., and Bickford, M. E. (2005). Cortical Function: A View from the Thalamus. *Neuron*, 45(4):485–488.
- <sup>9</sup>Baxter, M. (2013). Mediodorsal thalamus and cognition in non-human primates. *Frontiers in Systems Neuroscience*, 7:38.
- <sup>10</sup>Bennett, C. H. (2003). Notes on landauer’s principle, reversible computation, and maxwell’s demon. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics*, 34(3):501 – 510. Quantum Information and Computation.
- <sup>11</sup>Bickford, M. E. (2016). Thalamic Circuit Diversity: Modulation of the Driver/Modulator Framework. *Frontiers in Neural Circuits*, 9.
- <sup>12</sup>Bolkan, S. S., Stujenske, J. M., Parnaudeau, S., Spellman, T. J.,



- Rauffenbart, C., Abbas, A. I., Harris, A. Z., Gordon, J. A., and Kellendonk, C. (2017). Thalamic projections sustain prefrontal activity during working memory maintenance. *Nature Neuroscience*, 20(7):987–996.
- <sup>13</sup>Bruno, R. M. and Sakmann, B. (2006). Cortex is driven by weak but synchronously active thalamocortical synapses. *Science*, 312(5780):1622–1627.
- <sup>14</sup>Bueno-Junior, L. S., Lopes-Aguiar, C., Ruggiero, R. N., Romcy-Pereira, R. N., and Leite, J. P. (2012). Muscarinic and nicotinic modulation of thalamo-prefrontal cortex synaptic plasticity in vivo. *PLOS ONE*, 7:1–11.
- <sup>15</sup>Byne, W., Buchsbaum, M. S., Kemether, E., Hazlett, v., Shinwari, A., Mitropoulou, V., and Siever, L. J. (2001). Magnetic resonance imaging of the thalamic mediodorsal nucleus and pulvinar in schizophrenia and schizotypal personality disorder. *Archives of General Psychiatry*, 58(2):133–140.
- <sup>16</sup>Cardin, J. A., Palmer, L. A., and Contreras, D. (2008). Cellular mechanisms underlying stimulus-dependent gain modulation in primary visual cortex neurons in vivo. *Neuron*, 59:150–160.
- <sup>17</sup>Clasca, F., Rubio-Garrido, P., and Jabaudon, D. (2012). Unveiling the diversity of thalamocortical neuron subtypes. *Eur J Neurosci*, 35:1524–32.
- <sup>18</sup>Collins, D. P., Anastasiades, P. G., Marlin, J. J., and Carter, A. G. (2018). Reciprocal circuits linking the prefrontal cortex with dorsal and ventral thalamic nuclei. *Neuron*, 98(2):366–379.e4.
- <sup>19</sup>Corbetta, M. (1998). Frontoparietal cortical networks for directing attention and the eye to visual locations: identical, independent, or overlapping neural systems? *Proceedings of National Academy of Science*, 95:831–8.
- <sup>20</sup>Cruikshank, S. J., Ahmed, O. J., Stevens, T. R., Patrick, S. L., Gonzalez, A. N., Elmaleh, M., and Connors, B. W. (2012). Thalamic control of layer 1 circuits in prefrontal cortex. *Journal of Neuroscience*, 32(49):17813–17823.
- <sup>21</sup>Cruikshank, S. J., Lewis, T. J., and Connors, B. W. (2007). Synaptic basis for intense thalamocortical activation of feedforward inhibitory cells in neocortex. *Nature Neuroscience*, 10.
- <sup>22</sup>Crunelli, V. and Leresche, N. (1991). A role for gabab receptors in excitation and inhibition of thalamocortical cells. *Trends in Neurosciences*, 14(1):16 – 21.
- <sup>23</sup>Csete, M. E. and Doyle, J. C. (2002). Reverse engineering of biological complexity. *Science*, 295(5560):1664–1669.
- <sup>24</sup>Dehghani, N. (2017). Design of the Artificial: lessons from the biological roots of general intelligence. *ArXiv*.
- <sup>25</sup>Dehghani, N. (2018). Theoretical principles of multiscale spatiotemporal control of neuronal networks: A complex systems perspective. *Frontiers in Computational Neuroscience*, 12:81.
- <sup>26</sup>Dekel, E. and Alon, U. (2005). Optimality and evolutionary tuning of the expression level of a protein. *Nature*, 436(7050):588–592.
- <sup>27</sup>Delevich, K., Tucciarone, J., Huang, Z. J., and Li, B. (2015). The mediodorsal thalamus drives feedforward inhibition in the anterior cingulate cortex via parvalbumin interneurons. *Journal of Neuroscience*, 35(14):5743–5753.
- <sup>28</sup>Demirel, Y. (2011). *Energy Coupling*, pages 419–440. MIT Press.
- <sup>29</sup>Deniau, J. M. and Chevalier, G. (1985). Disinhibition as a basic process in the expression of striatal functions. II. The striatonigral influence on thalamocortical cells of the ventromedial thalamic nucleus. *Brain Research*, 334(2):227–233.
- <sup>30</sup>Douglas, R. J. and Martin, K. A. (2007). Mapping the Matrix: The Ways of Neocortex. *Neuron*, 56(2):226–238.
- <sup>31</sup>El-Samad, H. J., Goff, J. P., and Khamash, M. H. (2002). Calcium Homeostasis and Parturient Hypocalcemia: An Integral Feedback Perspective. *Journal of Theoretical Biology*, 214(1):17–29.
- <sup>32</sup>El-Samad, H. J., Kurata, H., Doyle, J. C., Gross, C. A., and Khamash, M. H. (2005). Surviving heat shock: Control strategies for robustness and performance. *Proceedings of the National Academy of Sciences*, 102(8):2736–2741.
- <sup>33</sup>Enel, P., Procyk, E., Quilodran, R., and Dominey, P. F. (2016). Reservoir Computing Properties of Neural Dynamics in Prefrontal Cortex. *PLOS Computational Biology*, 12(6):e1004967.
- <sup>34</sup>Fan, D., Duan, L., Wang, Q., and Luan, G. (2017). Combined effects of feedforward inhibition and excitation in thalamocortical circuit on the transitions of epileptic seizures. *Frontiers in Computational Neuroscience*, 11:59.
- <sup>35</sup>Felleman, D. J. and Essen, D. C. V. (1991). Distributed Hierarchical Processing in the Primate Cerebral Cortex. *Cerebral Cortex*, 1(1):1–47.
- <sup>36</sup>FitzGibbon, T., Eriköz, B., Grünert, U., and Martin, P. R. (2015). Analysis of the lateral geniculate nucleus in dichromatic and trichromatic marmosets. *Journal of Comparative Neurology*, 523(13):1948–1966.
- <sup>37</sup>Flack, J. (2017). *Life's Information Hierarchy*, page 283–302. Cambridge University Press.
- <sup>38</sup>Funahashi, K. and Nakamura, Y. (1993). Approximation of dynamical systems by continuous time recurrent neural networks. *Neural Networks*, 6(6):801–806.
- <sup>39</sup>Fusi, S., Miller, E. K., and Rigotti, M. (2016). Why neurons mix: high dimensionality for higher cognition. *Current Opinion in Neurobiology*, 37:66–74.
- <sup>40</sup>Gabernet, L., Jadhav, S. P., Feldman, D. E., Carandini, M., and Scanziani, M. (2005). Somatosensory integration controlled by dynamic thalamocortical feed-forward inhibition. *Neuron*, 48(2):315–327.
- <sup>41</sup>Giguere, M. and Goldman-Rakic, P. S. (1988). Mediodorsal nucleus: Areal, laminar, and tangential distribution of afferents and efferents in the frontal lobe of rhesus monkeys. *Journal of Comparative Neurology*, 277(2):195–213.
- <sup>42</sup>Godfrey, P., Shipley, R., and Gryz, J. (2006). Algorithms and analyses for maximal vector computation. *The VLDB Journal*, 16(1):5–28.
- <sup>43</sup>Goldberg, J. H., Farries, M. A., and Fee, M. S. (2013). Basal ganglia output to the thalamus: still a paradox. *Trends in Neurosciences*, 36(12):695–705.
- <sup>44</sup>Goldman-Rakic, P. S. and Porrino, L. J. (1985). The primate mediodorsal (md) nucleus and its projection to the frontal lobe. *Journal of Comparative Neurology*, 242(4):535–560.
- <sup>45</sup>Grant, E., Hoerder-Suabedissen, A., and Molnár, Z. (2012). Development of the Corticothalamic Projections. *Frontiers in Neuroscience*, 6.
- <sup>46</sup>Groh, A., Bokor, H., Mease, R. A., Plattner, V. M., Hangya, B., Stroh, A., Deschenes, M., and Acsády, L. (2014). Convergence of Cortical and Sensory Driver Inputs on Single Thalamocortical Cells. *Cerebral Cortex*, 24(12):3167–3179.
- <sup>47</sup>Halassa, M. M. and Acsády, L. (2016). Thalamic Inhibition: Diverse Sources Diverse Scales. *Trends in Neurosciences*, 39(10):680–693.
- <sup>48</sup>Halassa, M. M. and Kastner, S. (2017). Thalamic functions in distributed cognitive control. *Nature Neuroscience*, 20(12):1669–1679.
- <sup>49</sup>Harris, K. D. and Thiele, A. (2011). Cortical state and attention. *Nature Reviews Neuroscience*, 12:509 EP –.
- <sup>50</sup>Harth, E. M., Unnikrishnan, K. P., and Pandya, A. S. (1987). The inversion of sensory processing by feedback pathways: a model of visual cognitive functions. *Science*, 237(4811):184–187.
- <sup>51</sup>Hartwell, L. H., Hopfield, J. J., Leibler, S., and Murray, A. W. (1999). From molecular to modular cell biology. *Nature*, 402:C47 EP –.
- <sup>52</sup>Heeger, D. J. (2017). Theory of cortical function. *Proceedings of the National Academy of Sciences*, 114(8):1773–1782.
- <sup>53</sup>Hubel, D. H. and Wiesel, T. N. (1959). Receptive fields of single neurones in the cat's striate cortex. *The Journal of Physiology*, 148(3):574–591.
- <sup>54</sup>Hubel, D. H. and Wiesel, T. N. (1962). Receptive fields binocular interaction and functional architecture in the cat's visual cortex. *The Journal of Physiology*, 160(1):106–154.
- <sup>55</sup>Ikeda, K. and Matsumoto, K. (1987). High-dimensional chaotic behavior in systems with time-delayed feedback. *Physica D:*

- Nonlinear Phenomena*, 29(1-2):223–235.
- <sup>56</sup>Jaeger, H. (2001). The echo state approach to analysing and training recurrent neural networks. Technical report.
- <sup>57</sup>Jaeger, H. (2007). Echo state network. *Scholarpedia*, 2(9):2330.
- <sup>58</sup>Jazayeri, M. and Shadlen, M. N. (2015). A Neural Mechanism for Sensing and Reproducing a Time Interval. *Current Biology*, 25(20):2599–2609.
- <sup>59</sup>Jones, E. G. (1981). Functional subdivision and synaptic organization of the mammalian thalamus. *Int Rev Physiol*, 25:173–245.
- <sup>60</sup>Jones, E. G. (1985). Principles of Thalamic Organization. In *The Thalamus*, pages 85–149. Springer US.
- <sup>61</sup>Jones, E. G. (1998). Viewpoint: the core and matrix of thalamic organization. *Neuroscience*, 85(2):331–345.
- <sup>62</sup>Jones, E. G. (2002). Thalamic circuitry and thalamocortical synchrony. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 357(1428):1659–1673.
- <sup>63</sup>Ju, H., Hines, M. L., and Yu, Y. (2016). Cable energy function of cortical axons. *Scientific Reports*, 6:29686 EP –.
- <sup>64</sup>Kakei, S., Na, J., and Shinoda, Y. (2001). Thalamic terminal morphology and distribution of single corticothalamic axons originating from layers 5 and 6 of the cat motor cortex. *The Journal of Comparative Neurology*, 437(2):170–185.
- <sup>65</sup>Katz, M., Buchsbaum, M. S., Siegel Jr, B. V., Wu, J., Haier, R. J., and Bunney Jr, W. E. (1996). Correlational patterns of cerebral glucose metabolism in never-medicated schizophrenics. *Neuropsychobiology*, 33(1):1–11.
- <sup>66</sup>Khubieh, A., Ratté, S., Lankarany, M., and Prescott, S. A. (2016). Regulation of cortical dynamic range by background synaptic noise and feedforward inhibition. *Cerebral Cortex*, 26(8):3357–3369.
- <sup>67</sup>Kim, H. G., Beierlein, M., and Connors, B. W. (1995). Inhibitory control of excitable dendrites in neocortex. *Journal of Neurophysiology*, 74(4):1810–1814.
- <sup>68</sup>Kim, J., Woo, J., Park, Y.-G., Chae, S., Jo, S., Choi, J. W., Jun, H. Y., Yeom, Y. I., Park, S. H., Kim, K. H., Shin, H.-S., and Kim, D. (2011). Thalamic t-type  $ca_{2+}$  channels mediate frontal lobe dysfunctions caused by a hypoxia-like damage in the prefrontal cortex. *Journal of Neuroscience*, 31(11):4063–4073.
- <sup>69</sup>Komura, Y., Nikkuni, A., Hirashima, N., Uetake, T., and Miyamoto, A. (2013). Responses of pulvinar neurons reflect a subject's confidence in visual categorization. *Nature Neuroscience*, 16(6):749–755.
- <sup>70</sup>Krettek, J. E. and Price, J. L. (1977). The cortical projections of the mediodorsal nucleus and adjacent thalamic nuclei in the rat. *Journal of Comparative Neurology*, 171(2):157–191.
- <sup>71</sup>Krishna, S., Maslov, S., and Sneppen, K. (2007). UV-Induced Mutagenesis in Escherichia coli SOS Response: A Quantitative Model. *PLoS Computational Biology*, 3(3):e41.
- <sup>72</sup>Kung, H.-T., Luccio, F., and Preparata, F. P. (1975). On Finding the Maxima of a Set of Vectors. *Journal of the ACM*, 22(4):469–476.
- <sup>73</sup>Kuramoto, E., Pan, S., Furuta, T., Tanaka, Y. R., Iwai, H., Yamanaka, A., Ohno, S., Kaneko, T., Goto, T., and Hioki, H. (2017). Individual mediodorsal thalamic neurons project to multiple areas of the rat prefrontal cortex: A single neuron-tracing study using virus vectors. *Journal of Comparative Neurology*, 525(1):166–185.
- <sup>74</sup>Kuroda, M., Yokofujita, J., and Murakami, K. (1998). An ultrastructural study of the neural circuit between the prefrontal cortex and the mediodorsal nucleus of the thalamus. *Progress in Neurobiology*, 54(4):417–458.
- <sup>75</sup>Kuroda, M., Yokofujita, J., Oda, S., and Price, J. L. (2004). Synaptic relationships between axon terminals from the mediodorsal thalamic nucleus and  $\gamma$ -aminobutyric acidergic cortical cells in the prelimbic cortex of the rat. *Journal of Comparative Neurology*, 477(2):220–234.
- <sup>76</sup>Lan, G., Sartori, P., Neumann, S., Sourjik, V., and Tu, Y. (2012). The energy–speed–accuracy trade-off in sensory adaptation. *Nature Physics*, 8:422 EP –.
- <sup>77</sup>Larson, C. L., Davidson, R. J., Abercrombie, H. C., Ward, R. T., Schaefer, S. M., Jackson, D. C., Holden, J. E., and Perlman, S. B. (1998). Relations between pet-derived measures of thalamic glucose metabolism and eeg alpha power. *Psychophysiology*, 35(2):162–169.
- <sup>78</sup>Laughlin, S. B. and Sejnowski, T. J. (2003). Communication in neuronal networks. *Science*, 301:1870–4.
- <sup>79</sup>Lennie, P. (2003). The Cost of Cortical Computation. *Current Biology*, 13(6):493–497.
- <sup>80</sup>Levasseur, M., Baron, J., Sette, G., Legault-Demare, F., Pappata, S., Manguiere, F., Benoit, N., Dinh, S. T., Degos, J., Laplane, D., et al. (1992). Brain energy metabolism in bilateral paramedian thalamic infarcts: a positron emission tomography study. *Brain*, 115(3):795–807.
- <sup>81</sup>Lewis, D. A., Curley, A. A., Glausier, J. R., and Volk, D. W. (2012). Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends in Neurosciences*, 35(1):57–67.
- <sup>82</sup>Lin, H. W., Tegmark, M., and Rolnick, D. (2017). Why Does Deep and Cheap Learning Work So Well? *Journal of Statistical Physics*, 168(6):1223–1247.
- <sup>83</sup>Lindgren, K. A., Larson, C. L., Schaefer, S. M., Abercrombie, H. C., Ward, R. T., Oakes, T. R., Holden, J. E., Perlman, S. B., Benca, R. M., and Davidson, R. J. (1999). Thalamic metabolic rate predicts eeg alpha power in healthy control subjects but not in depressed patients. *Biological Psychiatry*, 45(8):943–952.
- <sup>84</sup>Ma, W. J. and Jazayeri, M. (2014). Neural Coding of Uncertainty and Probability. *Annual Review of Neuroscience*, 37(1):205–220.
- <sup>85</sup>Maass, W., Natschlaeger, T., and Markram, H. (2003). Computational Models for Generic Cortical Microcircuits. In *Computational Neuroscience*. Chapman and Hall/CRC.
- <sup>86</sup>Maass, W., Natschläger, T., and Markram, H. (2002). Real-Time Computing Without Stable States: A New Framework for Neural Computation Based on Perturbations. *Neural Computation*, 14(11):2531–2560.
- <sup>87</sup>Magistretti, P. J. (2006). Neuron–glia metabolic coupling and plasticity. *Journal of Experimental Biology*, 209(12):2304–2311.
- <sup>88</sup>Magistretti, P. J. and Allaman, I. (2015). A cellular perspective on brain energy metabolism and functional imaging. *Neuron*, 86(4):883–901.
- <sup>89</sup>Mante, V., Sussillo, D., Shenoy, K. V., and Newsome, W. T. (2013). Context-dependent computation by recurrent dynamics in prefrontal cortex. *Nature*, 503(7474):78–84.
- <sup>90</sup>Marblestone, A. H., Wayne, G., and Kording, K. P. (2016). Toward an Integration of Deep Learning and Neuroscience. *Frontiers in Computational Neuroscience*, 10.
- <sup>91</sup>Marenco, S., Stein, J. L., Savostyanova, A. A., Sambataro, F., Tan, H.-Y., Goldman, A. L., Verchinski, B. A., Barnett, A. S., Dickinson, D., Apud, J. A., Callicott, J. H., Meyer-Lindenberg, A., and Weinberger, D. R. (2012). Investigation of Anatomical Thalamo-Cortical Connectivity and fMRI Activation in Schizophrenia. *Neuropsychopharmacology*, 37(2):499–507.
- <sup>92</sup>Mathers, L. H. (1972). The synaptic organization of the cortical projection to the pulvinar of the squirrel monkey. *The Journal of Comparative Neurology*, 146(1):43–59.
- <sup>93</sup>Mesulam, M. M. (1990). Large-scale neurocognitive networks and distributed processing for attention, language, and memory. *Ann Neurol*, 28:597–613.
- <sup>94</sup>Mitchell, A. S. (2015). The mediodorsal thalamus as a higher order thalamic relay nucleus important for learning and decision-making. *Neurosci Biobehav Rev*, 54:76–88.
- <sup>95</sup>Mitelman, S. A., Byrne, W., Kemether, E. M., Hazlett, E. A., and Buchsbaum, M. S. (2005). Metabolic Disconnection Between the Mediodorsal Nucleus of the Thalamus and Cortical Brodmann's Areas of the Left Hemisphere in Schizophrenia. *American Journal of Psychiatry*, 162(9):1733–1735.
- <sup>96</sup>Mumford, D. (1991). On the computational architecture of the neocortex. I: The role of the thalamo-cortical loop. *Biological*

- Cybernetics*, 65(2):135–145.
- <sup>97</sup>Mumford, D. (1992). On the computational architecture of the neocortex. II The role of cortico-cortical loops. *Biological Cybernetics*, 66(3):241–251.
- <sup>98</sup>Nair, A., Treiber, J. M., Shukla, D. K., Shih, P., and Müller, R.-A. (2013). Impaired thalamocortical connectivity in autism spectrum disorder: a study of functional and anatomical connectivity. *Brain*, 136(6):1942–1955.
- <sup>99</sup>Nakajima, M. and Halassa, M. M. (2017). Thalamic control of functional cortical connectivity. *Current Opinion in Neurobiology*, 44:127–131.
- <sup>100</sup>Navlakha, S. and Bar-Joseph, Z. (2014). Distributed information processing in biological and computational systems. *Communications of the ACM*, 58(1):94–102.
- <sup>101</sup>Newell, A. (1962). Some problems of basic organization in problem solving programs.
- <sup>102</sup>Nii, P. (1986). The Blackboard Model of Problem Solving and the Evolution of Blackboard Architectures. *The AI Magazine*, 7(2):38–53.
- <sup>103</sup>Niven, J. E. and Laughlin, S. B. (2008). Energy limitation as a selective pressure on the evolution of sensory systems. *Journal of Experimental Biology*, 211(11):1792–1804.
- <sup>104</sup>Ohno, S., Kuramoto, E., Furuta, T., Hioki, H., Tanaka, Y., Fujiyama, F., Sonomura, T., Uemura, M., Sugiyama, K., and Kaneko, T. (2012). A morphological analysis of thalamocortical axon fibers of rat posterior thalamic nuclei: a single neuron tracing study with viral vectors. *Cereb Cortex*, 22:2840–57.
- <sup>105</sup>Parnaudeau, S., Bolkan, S. S., and Kellendonk, C. (2017). The Mediodorsal Thalamus: An Essential Partner of the Prefrontal Cortex for Cognition. *Biological Psychiatry*.
- <sup>106</sup>Parnaudeau, S., O’Neill, P.-K., Bolkan, S. S., Ward, R. D., Abbas, A. I., Roth, B. L., Balsam, P. D., Gordon, J. A., and Kellendonk, C. (2013). Inhibition of mediodorsal thalamus disrupts thalamofrontal connectivity and cognition. *Neuron*, 77(6):1151–1162.
- <sup>107</sup>Parnaudeau, S., Taylor, K., Bolkan, S. S., Ward, R. D., Balsam, P. D., and Kellendonk, C. (2015). Mediodorsal Thalamus Hypofunction Impairs Flexible Goal-Directed Behavior. *Biological Psychiatry*, 77(5):445–453.
- <sup>108</sup>Parrondo, J. M. R., Horowitz, J. M., and Sagawa, T. (2015). Thermodynamics of information. *Nature Physics*, 11:131 EP –.
- <sup>109</sup>Partlow, G. D., Colonnier, M., and Szabo, J. (1977). Thalamic projections of the superior colliculus in the rhesus monkey, *Macaca mulatta*. A light and electron microscopic study. *The Journal of Comparative Neurology*, 171(3):285–317.
- <sup>110</sup>Pelzer, P., Horstmann, H., and Kuner, T. (2017). Ultrastructural and functional properties of a giant synapse driving the piriform cortex to mediodorsal thalamus projection. *Frontiers in Synaptic Neuroscience*, 9:3.
- <sup>111</sup>Pinault, D. (2004). The thalamic reticular nucleus: structure function and concept. *Brain Research Reviews*, 46(1):1–31.
- <sup>112</sup>Popken, G. J., Bunney, W. E., Potkin, S. G., and Jones, E. G. (2000). Subnucleus-specific loss of neurons in medial thalamus of schizophrenics. *Proceedings of the National Academy of Sciences*, 97(16):9276–9280.
- <sup>113</sup>Prescott, S. A. and De Koninck, Y. (2003). Gain control of firing rate by shunting inhibition: Roles of synaptic noise and dendritic saturation. *Proceedings of the National Academy of Sciences*, 100(4):2076–2081.
- <sup>114</sup>Purushothaman, G., Marion, R., Li, K., and Casagrande, V. (2012). Gating and control of primary visual cortex by pulvinar. *Nat Neurosci*, 15:905–12.
- <sup>115</sup>Raczkowski, D. and Fitzpatrick, D. (1990). Terminal arbors of individual physiologically identified geniculocortical axons in the tree shrew’s striate cortex. *The Journal of Comparative Neurology*, 302(3):500–514.
- <sup>116</sup>Rigotti, M., Barak, O., Warden, M. R., Wang, X.-J., Daw, N. D., Miller, E. K., and Fusi, S. (2013). The importance of mixed selectivity in complex cognitive tasks. *Nature*, 497(7451):585–590.
- <sup>117</sup>Rigotti, M., Rubin, D. B. D., Wang, X.-J., and Fusi, S. (2010). Internal representation of task rules by recurrent dynamics: the importance of the diversity of neural responses. *Frontiers in Computational Neuroscience*, 4.
- <sup>118</sup>Rikhye, R. V., Wimmer, R. D., and Halassa, M. M. (2018). Towards an integrative theory of thalamic function. *Nature Neuroscience (In press)*, xxx(xxx):xxx–xxx.
- <sup>119</sup>Rose, H. J. and Metherate, R. (2005). Auditory thalamocortical transmission is reliable and temporally precise. *Journal of Neurophysiology*, 94(3):2019–2030.
- <sup>120</sup>Rotaru, D. C., Barrionuevo, G., and Sesack, S. R. (2005). Mediodorsal thalamic afferents to layer iii of the rat prefrontal cortex: Synaptic relationships to subclasses of interneurons. *Journal of Comparative Neurology*, 490(3):220–238.
- <sup>121</sup>Rouiller, E. M. and Welker, E. (2000). A comparative analysis of the morphology of corticothalamic projections in mammals. *Brain Research Bulletin*, 53(6):727–741.
- <sup>122</sup>Rovo, Z., Ulbert, I., and Acsády, L. (2012). Drivers of the primate thalamus. *J Neurosci*, 32:17894–908.
- <sup>123</sup>Saalmann, Y. B. (2014). Intralaminar and medial thalamic influence on cortical synchrony, information transmission and cognition. *Frontiers in Systems Neuroscience*, 8:83.
- <sup>124</sup>Saalmann, Y. B. and Kastner, S. (2011). Cognitive and perceptual functions of the visual thalamus. *Neuron*, 71(2):209 – 223.
- <sup>125</sup>Saalmann, Y. B. and Kastner, S. (2015). The cognitive thalamus. *Frontiers in Systems Neuroscience*, 9:39.
- <sup>126</sup>Schmitt, L. I., Wimmer, R. D., Nakajima, M., Happ, M., Mofakham, S., and Halassa, M. M. (2017). Thalamic amplification of cortical connectivity sustains attentional control. *Nature*, 545(7653):219–223.
- <sup>127</sup>Schwartz, M. L., Dekker, J. J., and Goldman-Rakic, P. S. (1991). Dual mode of corticothalamic synaptic termination in the mediodorsal nucleus of the rhesus monkey. *Journal of Comparative Neurology*, 309(3):289–304.
- <sup>128</sup>Scott, B. B., Constantinople, C., Akrami, A., Hanks, T. D., Brody, C. D., and Tank, D. W. (2017). Fronto-parietal Cortical Circuits Encode Accumulated Evidence with a Diversity of Timescales. *Neuron*, 95:385–398.e5.
- <sup>129</sup>Selfridge, O. G. (1959). Pandemonium: a paradigm for learning in . In Blake, D. V. and Uttley, A. M., editors, *Proceedings of the Symposium on Mechanisation of Thought Processes*, pages 513–526, London. National Physical Laboratory.
- <sup>130</sup>Seoane, L. F. and Solé, R. (2015). Phase transitions in Pareto optimal complex networks. *Physical Review E*, 92(3).
- <sup>131</sup>Sherman, S. M. (2016). Thalamus plays a central role in ongoing cortical functioning. *Nature Neuroscience*, 16(4):533–541.
- <sup>132</sup>Sherman, S. M. and Guillery, R. W. (2002). The role of the thalamus in the flow of information to the cortex. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 357(1428):1695–1708.
- <sup>133</sup>Sherman, S. M. and Guillery, R. W. (2013). *Functional Connections of Cortical Areas*. The MIT Press.
- <sup>134</sup>Shoham, S., O’Connor, D. H., and Segev, R. (2006). How silent is the brain: is there a “dark matter” problem in neuroscience? *Journal of Comparative Physiology A*, 192(8):777–784.
- <sup>135</sup>Simon, H. A. (1962). The Architecture of Complexity. *Proceedings of the American Philosophical Society*, 106(6):467–482.
- <sup>136</sup>Simon, H. A. (1969). *The Sciences of the Artificial*, chapter The Architecture of Complexity. MIT Press.
- <sup>137</sup>Sipper, M. (1999). The emergence of cellular computing. *Computer*, 32(7):18–26.
- <sup>138</sup>Solé, R. V. and Macia, J. (2013). Expanding the landscape of biological computation with synthetic multicellular consortia. *Natural Computing*, 12(4):485–497.
- <sup>139</sup>Spacek, J. and Lieberman, A. (1974). Ultrastructure and three-dimensional organization of synaptic glomeruli in rat somatosensory thalamus. *Journal of anatomy*, 117(Pt 3):487.
- <sup>140</sup>Steriade, M. and Deschenes, M. (1984). The thalamus as a neuronal oscillator. *Brain Research Reviews*, 8(1):1–63.

- <sup>141</sup>Steriade, M., Domich, L., and Oakson, G. (1986). Reticular thalamic neurons revisited: activity changes during shifts in states of vigilance. *J Neurosci*, 6:68–81.
- <sup>142</sup>Steriade, M. and Llinás, R. R. (1988). The functional states of the thalamus and the associated neuronal interplay. *Physiological Reviews*, 68(3):649–742.
- <sup>143</sup>Steriade, M. and Pare, D. (2007). Morphology and electrophysiological properties of thalamic neurons. In *Gating in Cerebral Networks*, pages 1–26. Cambridge University Press.
- <sup>144</sup>Steriade, M., Parent, A., and Hada, J. (1984). Thalamic projections of nucleus reticularis thalami of cat: A study using retrograde transport of horseradish peroxidase and fluorescent tracers. *The Journal of Comparative Neurology*, 229(4):531–547.
- <sup>145</sup>Szekely, P., Sheftel, H., Mayo, A., and Alon, U. (2013). Evolutionary Tradeoffs between Economy and Effectiveness in Biological Homeostasis Systems. *PLoS Computational Biology*, 9(8):e1003163.
- <sup>146</sup>Uhlhaas, P. J. and Singer, W. (2010). Abnormal neural oscillations and synchrony in schizophrenia. *Nature Reviews Neuroscience*, 11:100 EP –.
- <sup>147</sup>Urbain, N. and Deschênes, M. (2007). Motor Cortex Gates Vibrissal Responses in a Thalamocortical Projection Pathway. *Neuron*, 56(4):714–725.
- <sup>148</sup>van Wijngaarden, J. B. G., Zucca, R., Finnigan, S., and Verschure, P. F. M. J. (2016). The impact of cortical lesions on thalamo-cortical network dynamics after acute ischaemic stroke: A combined experimental and theoretical study. *PLoS Computational Biology*, 12(8):1–16.
- <sup>149</sup>Viaene, A. N., Petrof, I., and Sherman, S. M. (2011a). Properties of the thalamic projection from the posterior medial nucleus to primary and secondary somatosensory cortices in the mouse. *Proceedings of the National Academy of Sciences*, 108(44):18156–18161.
- <sup>150</sup>Viaene, A. N., Petrof, I., and Sherman, S. M. (2011b). Synaptic properties of thalamic input to the subgranular layers of primary somatosensory and auditory cortices in the mouse. *Journal of Neuroscience*, 31(36):12738–12747.
- <sup>151</sup>Wang, J., Narain, D., Hosseini, E. A., and Jazayeri, M. (2018). Flexible timing by temporal scaling of cortical responses. *Nature Neuroscience*.
- <sup>152</sup>Wang, R. and Wang, Z. (2014). Energy distribution property and energy coding of a structural neural network. *Frontiers in Computational Neuroscience*, 8:14.
- <sup>153</sup>Wang, R., Zhang, Z., and Chen, G. (2009). Energy coding and energy functions for local activities of the brain. *Neurocomputing*, 73(1):139 – 150.
- <sup>154</sup>Wang, R. and Zhu, Y. (2016). Can the activities of the large scale cortical network be expressed by neural energy? a brief review. *Cognitive neurodynamics*, 10(1):1–5.
- <sup>155</sup>Woodward, N. D., Giraldo-Chica, M., Rogers, B., and Cascio, C. J. (2017). Thalamocortical Dysconnectivity in Autism Spectrum Disorder: An Analysis of the Autism Brain Imaging Data Exchange. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 2(1):76–84.
- <sup>156</sup>Yamins, D. L. K. and DiCarlo, J. J. (2016). Using goal-driven deep learning models to understand sensory cortex. *Nature Neuroscience*, 19(3):356–365.
- <sup>157</sup>Yamins, D. L. K., Hong, H., Cadieu, C. F., Solomon, E. A., Seibert, D., and DiCarlo, J. J. (2014). Performance-optimized hierarchical models predict neural responses in higher visual cortex. *Proceedings of the National Academy of Sciences*, 111(23):8619–8624.
- <sup>158</sup>Zhou, H., Schafer, R. J., and Desimone, R. (2016). Pulvinar-cortex interactions in vision and attention. *Neuron*, 89(1):209–220.
- <sup>159</sup>Zikopoulos, B. and Barbas, H. (2007). Parallel driving and modulatory pathways link the prefrontal cortex and thalamus. *PLoS ONE*, 2(9):1–19.