An extension of Azzalini's method

Filippo Domma

Department of Economics, Statistics and Finance, University of Calabria, Italy

Božidar V. Popović

University of Montenegro, Faculty of Science and Mathematics, Podgorica, Montenegro

Saralees Nadarajah

School of Mathematics, University of Manchester, Manchester, UK

Abstract

The aim of this paper is to extend Azzalini's method. This extension is done in two stages: consider two dependent and non-identically distributed random variables say X_1 and X_2 ; model the dependence between X_1 and X_2 by a copula. To illustrate the new method, we assume X_1 and X_2 are exponential random variables. This assumption leads to a new distribution called the Generalized Weighted Exponential Distribution (GWED), a generalization of Gupta and Kundu (2009)'s Weighted Exponential Distribution (WED). Some mathematical properties of the GWED are derived, and its parameters estimated by maximum likelihood. The GWED is applied to biochemical data sets showing its good performance compared to the WED.

Keywords and phrases: Azzalini's method; Copula; Hidden truncation; Weighted distribution.

MSC 2010: 60E05, 62P10, 62G30, 62F10

1 Introduction

In real life, there are many data sets that are asymmetric, multimodal and heavy tailed. This has motivated many researchers to develop non-normal and/or skewed distributions. In the statistical literature, there are various techniques to build non-normal and/or skewed distributions. Nowadays, the most widely used technique for introducing asymmetry in a symmetric distribution is that due to Azzalini (1985). With reference to the normal distribution, this technique can be described as follows: a random variable Z is said to have the skew-normal distribution with parameter α, written as $Z \sim SN(\alpha)$, if its probability density function (pdf) is $f(z; \alpha) = 2\Phi(\alpha z) \phi(z)$, where $\Phi(\cdot)$ denotes the standard normal cumulative distribution function (cdf), $\phi(\cdot)$ denotes the standard normal pdf, and z and α are real numbers (Azzalini, 1985). An enormous literature exists on the study of the skew-normal distribution and its extension to the multivariate case.

Azzalini's method can also be described in terms of conditional distributions: let X_i , $i = 1, 2$ be independent random variables with pdfs $f_i(x_i)$ and cdfs $F_i(x_i)$. Then, the conditional pdf of $X = X_1$ given $\alpha X_1 > X_2$ is

$$
f_X(x) = \frac{f_1(x)F_2(\alpha x)}{P(\alpha X_1 > X_2)}.
$$
\n(1)

Observe that $P(\alpha X_1 > X_2) = \int^{+\infty}$ θ $\int \frac{\alpha x_1}{2}$ θ $f_2(x_2) dx_2$ 1 $f_1(x_1) dx_1 = E_{X_1} [F_2(\alpha X)], \text{ where}$ $E_{X_1}(\cdot)$ denotes the expectation with respect to X_1 . Equation [\(1\)](#page-1-0) can be interpreted as a weighted distribution with weight function $w(x) = F_2(\alpha x)$. If we set $f_1(x_1) = f_2(x_2) = \phi(x)$, i.e., X_1 and X_2 are standard normal random variables, we obtain the skew-normal distribution.

In the literature, [\(1\)](#page-1-0) has been used to construct new skewed distributions from a given symmetric distribution, for example, skew-t (Gupta et al., 2002), skew-Cauchy (Arnold and Beaver, 2000b; Gupta et al., 2002), skew-Laplace (Gupta et al., 2002; Aryal and Nadarajah, 2005) and skew-logistic (Gupta et al., 2002; Nadarajah, 2009). However, there is little work on the use of Azzalini's method for non-symmetric distributions. Gupta and Kundu (2009) took X_1 and X_2 in [\(1\)](#page-1-0) to be independent and identical exponential random variables with scale parameter λ , giving

$$
f_X(x) = \frac{\alpha + 1}{\alpha} \lambda e^{-\lambda x} \left(1 - e^{-\alpha \lambda x} \right).
$$

Gupta and Kundu (2009) called this the Weighted Exponential Distribution (WED). Shakhatreh (2012) studied a two-parameter version of the WED. Mahdy (2011, 2013) proposed weighted gamma and weighted Weibull distributions.

The aim of this paper is to extend Azzalini's method in two stages: take X_1 and X_2 to be dependent and non-identically distributed random variables; model their dependence using a copula. After a general discussion on the potentials, we illustrate this method by assuming X_1 and X_2 are exponential random variables. This assumption leads us to a new distribution called the Generalized Weighted Exponential Distribution (GWED), a generalization of the WED. Although the GWED is defined using Azzalini's method in terms of weighted distribution, we will show that it can also be interpreted as a hidden truncated distribution (Arnold and Beaver, 2000a). Moreover, the GWED can be interpreted as a finite mixture. The mixture representation enables us to derive mathematical properties of the GWED like its cdf, moments, and the moment generating function. The skewness of the WED due to Gupta and Kundu (2009) takes values in $[\sqrt{2},2]$ while that of the two-parameter WED due to Shakhatreh (2012) takes values in $\left[\frac{2}{\sqrt{2}}\right]$ $\left[\frac{1}{3},2\right]$. The GWED allows for wider values for skewness. Another importance feature is that the GWED allows for non-monotonic hazard rate functions (hrfs). The hrfs of the WED are always monotonic.

The paper is organized as follows. The extension of Azzalini's method is described in Section 2. Details (including mathematical properties, estimation issues and applications) of an important special case are given in Section 3.

2 The extension of Azzalini's method

Unlike Azzalini's method, we consider two dependent and non-identically distributed random variables. This extension can be used to construct any distribution.

First note that the denominator in [\(1\)](#page-1-0) in the case of dependence of X_1 and X_2 can be

expressed as

$$
P(\alpha X_1 > X_2) = \int_0^{+\infty} \int_0^{\alpha x_1} f(x_1, x_2) dx_1 dx_2
$$

=
$$
\int_0^{+\infty} \left\{ \int_0^{\alpha x_1} c(F_1(x_1), F_2(x_2)) f_2(x_2) dx_2 \right\} f_1(x_1) dx_1
$$

=
$$
E_{X_1} \left[\int_0^{\alpha X_1} c(F_1(X_1), F_2(x_2)) f_2(x_2) dx_2 \right],
$$
 (2)

where $c(F_1(x_1), F_2(x_2)) = \frac{\partial^2 C(F_1(x_1), F_2(x_2))}{\partial F_1(x_1)\partial F_2(x_2)}$ $\frac{C(F_1(x_1), F_2(x_2))}{\partial F_1(x_1)\partial F_2(x_2)}$ is the copula pdf. Accordingly, the weight function is $w(x) = \int^{\alpha x_1}$ $\dot{0}$ $c(F_1(x_1), F_2(x_2)) f_2(x_2) dx_2$. If $\alpha = 1, (2)$ $\alpha = 1, (2)$ reduces to the stress-strength model widely studied in the literature, see, for example, Kotz et al. (2003); using a copula, Domma and Giordano (2013) have recently highlighted the role of dependence between stress and strength on a reliability measure defined by [\(2\)](#page-2-0).

We are now able to provide

Definition 1 Let X_i , $i = 1, 2$ be dependent random variables with pdfs $f_i(x_i; \xi_i)$, cdfs $F_i(x_i; \xi_i)$ and joint pdf $f(x_1, x_2; \xi_1, \xi_2, \theta) = c(F_1(x_1; \xi_1), F_2(x_2; \xi_2); \theta) f_1(x_1; \xi_1) f_2(x_2; \xi_2),$ where $c(\cdot, \cdot)$ is a copula pdf. Then, the random variable $X = X_1 | \alpha X_1 > X_2$ is said to have a Generalized Weighted Distribution if its pdf is

$$
f^{w}(x; \xi_{1}, \xi_{2}, \alpha, \theta) = \frac{f_{1}(x; \xi_{1}) \int_{-\infty}^{\alpha x} c(F_{1}(x; \xi_{1}), F_{2}(x_{2}; \xi_{2}); \theta) f_{2}(x_{2}; \xi_{2}) dx_{2}}{E_{X_{1}} \left[\int_{-\infty}^{\alpha X_{1}} c(F_{1}(X_{1}), F_{2}(x_{2}); \xi_{1}) f_{2}(x_{2}; \xi_{2}) dx_{2} \right]}.
$$
(3)

The use of copula is motivated by the fact that it allows for the dependence structure to be treated separately from the marginal components of a joint distribution. Various forms of dependence structures (linear, non-linear, tail dependence, etc) can be used. Also X_1 and X_2 need not necessarily belong to the same parametric family.

To better understand the role of [\(3\)](#page-2-1) in defining new distributions, we now consider a recent expansion applicable to a wide range of bivariate copulas (Nadarajah, 2014):

$$
C\left(F_1\left(x_1;\xi_1\right), F_2\left(x_2;\xi_2\right); \theta\right) = \sum_{i=1}^p \gamma_i \left[F_1\left(x_1;\xi_1\right)\right]^{a_i} \left[F_2\left(x_2;\xi_2\right)\right]^{b_i},\tag{4}
$$

where $p \ge 1$ and $\{(\gamma_i, a_i, b_i) : i \ge 1\}$ are real numbers. The corresponding joint pdf is

$$
c(F_1(x_1;\xi_1), F_2(x_2;\xi_2); \theta) = \sum_{i=1}^p \gamma_i a_i b_i [F_1(x_1;\xi_1)]^{a_i-1} [F_2(x_2;\xi_2)]^{b_i-1}.
$$

Then, the denominator in [\(3\)](#page-2-1) reduces to

$$
P(\alpha X_1 > X_2) = \sum_{i=1}^p \gamma_i a_i E_{X_1} \left[(F_1(X_1; \xi_1))^{a_i-1} (F_2(\alpha X_1; \xi_2))^{b_i} \right],
$$

while weight function is

$$
w(x) = \sum_{i=1}^{p} \gamma_i a_i (F_1(x; \xi_1))^{a_i-1} (F_2(\alpha x; \xi_2))^{b_i}.
$$

Consequently, the new pdf is

$$
f^{w}(x; \xi_{1}, \xi_{2}, \alpha, \theta) = \overline{K} f_{1}(x; \xi_{1}) \sum_{i=1}^{p} \gamma_{i} a_{i} (F_{1}(x; \xi_{1}))^{a_{i}-1} (F_{2}(\alpha x; \xi_{2}))^{b_{i}}, \qquad (5)
$$

where

$$
\overline{K} = \left\{ \sum_{i=1}^{p} \gamma_i a_i E_{X_1} \left[\left(F_1 \left(X_1; \xi_1 \right) \right)^{a_i-1} \left(F_2 \left(\alpha X_1; \xi_2 \right) \right)^{b_i} \right] \right\}^{-1}.
$$

The properties of the new pdf [\(5\)](#page-3-0) depend on the specified forms of the cdfs $F_1(x_1;\xi_1)$ and $F_2(x_2;\xi_2)$. In the next section, we study a simple but an important special case that generalizes a distribution proposed by Gupta and Kundu (2009).

3 A special case: Generalized weighted exponentiated distribution

Here, we illustrate an application of the extension of Azzalini's method. We take X_i , $i = 1, 2$ to be exponential random variables with cdfs $F_i(x) = \lambda_i e^{-\lambda_i x}$, and C to be the Farlie-Gumbel-Morgenstern copula (Morgenstern, 1956) defined by

$$
C(u_1, u_2) = u_1 u_2 \{1 + \theta [1 - u_1] [1 - u_2] \}.
$$

These choices yield a new distribution called the Generalized Weighted Exponential Distribution (GWED). This distribution generalizes the WED due to Gupta and Kundu (2009). We will show that the GWED can be interpreted as a hidden truncated distribution (Arnold and Beaver, 2000a) and as a finite mixture. We will also discuss the behavior of the pdf and the hrf of the GWED.

The Farlie-Gumbel-Morgenstern copula is a special case of [\(4\)](#page-2-2) for $p = 4$, $\gamma_1 = 1 + \theta$, $a_1 = 1$, $b_1 = 1, \ \gamma_2 = -\theta, \ a_2 = 1, \ b_2 = 2, \ \gamma_3 = -\theta, \ a_3 = 2, \ b_3 = 1, \ \gamma_4 = \theta, \ a_4 = 2 \text{ and } b_4 = 2.$ Consequently, after simple algebra, we find that $\overline{K} = \left\{ \frac{\lambda_2 \alpha}{\lambda_1 + \lambda_2 \alpha} + \theta \lambda_1 \left[\frac{2}{2\lambda_1 + \lambda_2 \alpha} + \frac{1}{\lambda_1 + 2\lambda_2 \alpha} - \frac{2}{\lambda_1 + \lambda_2 \alpha} \right] \right\}$ $\lambda_1+\lambda_2\alpha$ $\big] \big\}^{-1}$. Now, we can provide the definition of the GWED.

Definition 2 Let X_i , $i = 1,2$ be exponential random variables with pdfs $f_i(x_i; \lambda_i) = \lambda_i e^{-\lambda_i x_i}$, cdfs $F_i(x_i; \lambda_i) = 1 - e^{-\lambda_i x_i}$ and the joint pdf

$$
f(x_1, x_2; \lambda_1, \lambda_2, \theta) = \left\{ 1 + \theta \left[1 - 2e^{-\lambda_1 x_1} \right] \left[1 - 2e^{-\lambda_2 x_2} \right] \right\} f_1(x_1; \lambda_1) f_2(x_2; \lambda_2).
$$

Then, the random variable $X = X_1 | \alpha X_1 > X_2$ is said to have the GWED if its pdf is

$$
f^{w}(x; \lambda_{1}, \lambda_{2}, \alpha, \theta) = Ke^{-\lambda_{1}x} \left(1 - e^{-\lambda_{2}\alpha x}\right) \left[1 - \theta e^{-\lambda_{2}\alpha x} \left(1 - 2e^{-\lambda_{1}x}\right)\right],
$$
\n(6)

where $K = \lambda_1 \overline{K}$.

It is easy to see that the WED due to Gupta and Kundu (2009) is the special case of the GWED for $\lambda_1 = \lambda_2 = \lambda$ and $\theta = 0$.

Proposition 3 shows that $f^w(x; \lambda_1, \lambda_2, \alpha, \theta)$ in [\(6\)](#page-3-1) can be interpreted as a hidden truncated pdf (Arnold and Beaver, 2000a).

Proposition 3 Suppose Z and Y are two dependent positive random variables with joint pdf

$$
f_{Z,Y}(z,y;\lambda_1,\lambda_2,\theta) = \lambda_1 \lambda_2 z e^{-(\lambda_1+\lambda_2 y)z} \left\{ 1 + \theta \left[1 - 2e^{-\lambda_1 z} \right] \left[1 - 2e^{-\lambda_2 y z} \right] \right\}.
$$

Then, the pdf of $X = Z|Y \leq \alpha$ is equal to $f^w(x; \lambda_1, \lambda_2, \alpha, \theta)$ in [\(6\)](#page-3-1).

Proof. First, observe that the conditional cdf of $X = Z|Y \leq \alpha$ is

$$
F_{X=Z|Y<\alpha}(x)=\frac{\int_0^x\int_0^\alpha f_{Z,Y}(z,y)\mathrm{d}y\mathrm{d}z}{\int_0^\alpha f_Y(y)\mathrm{d}y}.
$$

After simple algebra, we have

$$
\int_0^\alpha f_{Z,Y}(z,y;\lambda_1,\lambda_2,\theta) dy = \lambda_1 \left\{ (1+\theta)e^{-\lambda_1 z} \left(1 - e^{-\lambda_2 \alpha z} \right) - \theta e^{-\lambda_1 z} \left(1 - e^{-2\lambda_2 \alpha z} \right) - 2\theta e^{-2\lambda_1 z} \left(1 - e^{-\lambda_2 \alpha z} \right) + 2\theta e^{-2\lambda_1 z} \left(1 - e^{-2\lambda_2 \alpha z} \right) \right\}
$$

and

$$
\int_0^x \int_0^\alpha f_{Z,Y}(z,y;\lambda_1,\lambda_2,\theta) dy = \lambda_1 \left\{ (1+\theta) \left[\frac{1 - e^{-\lambda_1 x}}{\lambda_1} - \frac{1 - e^{-(\lambda_1 + \lambda_2 \alpha)x}}{\lambda_1 + \lambda_2 \alpha} \right] - \theta \left[\frac{1 - e^{-\lambda_1 x}}{\lambda_1} - \frac{1 - e^{-(\lambda_1 + 2\lambda_2 \alpha)x}}{\lambda_1 + 2\lambda_2 \alpha} \right] - 2\theta \left[\frac{1 - e^{-2\lambda_1 x}}{2\lambda_1} - \frac{1 - e^{-(2\lambda_1 + \lambda_2 \alpha)x}}{2\lambda_1 + \lambda_2 \alpha} \right] + 2\theta \left[\frac{1 - e^{-2\lambda_1 x}}{2\lambda_1} - \frac{1 - e^{-2(\lambda_1 + \lambda_2 \alpha)x}}{2(\lambda_1 + \lambda_2 \alpha)} \right] \right\}.
$$
\n(7)

Moreover, the marginal pdf of Y is

$$
f_Y(y; \lambda_1, \lambda_2, \theta) = \lambda_1 \lambda_2 \left\{ \frac{1 + 2\theta}{\left(\lambda_1 + \lambda_2 y\right)^2} - \frac{2\theta}{\left(\lambda_1 + 2\lambda_2 y\right)^2} - \frac{2\theta}{\left(2\lambda_1 + \lambda_2 y\right)^2} \right\}.
$$

Consequently,

$$
P(Y \le \alpha) = \frac{\lambda_2 \alpha}{\lambda_1 + \lambda_2 \alpha} + \lambda_1 \theta \left[\frac{2}{2\lambda_1 + \lambda_2 \alpha} + \frac{1}{\lambda_1 + 2\lambda_2 \alpha} - \frac{2}{\lambda_1 + \lambda_2 \alpha} \right].
$$
 (8)

By combining [\(7\)](#page-4-0) with [\(8\)](#page-4-1), we obtain the cdf of $X|Y \leq \alpha$, i.e., $F_{X=Z|Y \leq \alpha}(x; \lambda_1, \lambda_2, \alpha, \theta)$. By differentiating with respect to x, it can be verified that the conditional pdf $f_{X=Z|Y\leq\alpha}(x;\lambda_1,\lambda_2,\alpha,\theta)$ is equal to [\(6\)](#page-3-1). The proof is complete. \Box

It is easy to see that $\lim_{x\to 0^+} f^w(x; \lambda_1, \lambda_2, \alpha, \theta) = \lim_{x\to +\infty} f^w(x; \lambda_1, \lambda_2, \alpha, \theta) = 0$. Therefore, there exists at least one $x > 0$, say x_0 , such that $\frac{\partial f^w(x; \lambda_1, \lambda_2, \alpha, \theta)}{\partial x}$ $\Big|_{x=x_0} = 0$. Figure [1](#page-5-0) plots [\(6\)](#page-3-1) for different parameter values.

Proposition 4 interprets [\(6\)](#page-3-1) as a finite mixture of exponential pdfs $f_j(x; \tau) = \tau e^{-\tau x}$.

Proposition 4 The pdf $f^w(x; \lambda_1, \lambda_2, \alpha, \theta)$ can be expressed as

$$
f^{w}(x; \lambda_{1}, \lambda_{2}, \alpha, \theta) = p_{1} f_{1}(x; \lambda_{1}) + p_{2} f_{2}(x; \lambda_{1} + \lambda_{2} \alpha) + p_{3} f_{3}(x; 2\lambda_{1} + \lambda_{2} \alpha) + p_{4} f_{4}(x; \lambda_{1} + 2\lambda_{2} \alpha) + p_{5} f_{5}(x; 2(\lambda_{1} + \lambda_{2} \alpha)),
$$
\n(9)

where $p_1 = \frac{K}{\lambda_1}$, $p_2 = -\frac{K(1+\theta)}{\lambda_1 + \lambda_2 \alpha}$ $\frac{K(1+\theta)}{\lambda_1+\lambda_2\alpha}$, $p_3=\frac{2K\theta}{2\lambda_1+\lambda_2\alpha}$, $p_4=\frac{K\theta}{\lambda_1+2\lambda_2\alpha}$ and $p_5=-\frac{K\theta}{\lambda_1+\lambda_2\alpha}$.

Proof. Follows by simple algebra. \Box

For details about mixtures, we refer readers to Titterington et al. (1985, page 50). A mixture of distributions is a valid distribution if the sum of weights is equal to 1. In the case of [\(9\)](#page-4-2), it is straightforward to verify that $\sum_{n=1}^{5}$ $j=1$ $p_j = 1$. The mixture representation [\(9\)](#page-4-2) enables us to derive

Figure 1: Examples of the GWED pdf for $(\lambda_1, \lambda_2, \alpha, \theta) = (1, 1, 1, -0.99)$ (solid line), $(\lambda_1, \lambda_2, \alpha, \theta) = (1, 1, 1, 0.99)$ (dashed line), $(\lambda_1, \lambda_2, \alpha, \theta) = (1, 1, 0.5, -0.99)$ (dotted line), $(\lambda_1, \lambda_2, \alpha, \theta) = (1, 1, 0.5, 0.99)$ (dot-dashed line).

mathematical properties of the GWED like its cdf, moments, and moment generating function. For example, the cdf of the GWED can be expressed as

$$
F^{w}(x; \lambda_{1}, \lambda_{2}, \alpha, \theta) = p_{1}F_{1}(x; \lambda_{1}) + p_{2}F_{2}(x; \lambda_{1} + \lambda_{2}\alpha) + p_{3}F_{3}(x; 2\lambda_{1} + \lambda_{2}\alpha) + p_{4}F_{4}(x; \lambda_{1} + 2\lambda_{2}\alpha) + p_{5}F_{5}(x; 2(\lambda_{1} + \lambda_{2}\alpha)).
$$

An alternative expression for the cdf can be obtained from [\(9\)](#page-4-2) by using the Taylor series for exponential functions:

$$
F^{w}(x; \lambda_{1}, \lambda_{2}, \alpha, \theta) = \sum_{i \geq 1} a_{i} x^{i} = \sum_{i \geq 0} a_{i+1} x^{i+1} = x \sum_{i \geq 0} d_{i} x^{i},
$$
\n(10)

where

$$
d_{i} = a_{i+1} = \frac{(-1)^{i+1}}{(i+1)!} \left[p_{1} \lambda_{1}^{i+1} + p_{2} \left(\lambda_{1} + \lambda_{2} \alpha \right)^{i+1} + p_{3} \left(2\lambda_{1} + \lambda_{2} \alpha \right)^{i+1} + p_{4} \left(\lambda_{1} + 2\lambda_{2} \alpha \right)^{i+1} + p_{5} \left(\lambda_{1} + \lambda_{2} \alpha \right)^{i+1} \right].
$$

The hrf of the GWED given by $h^{w}(x; \lambda_1, \lambda_2, \alpha, \theta) = \frac{f^{w}(x; \lambda_1, \lambda_2, \alpha, \theta)}{1 - F^{w}(x; \lambda_1, \lambda_2, \alpha, \theta)}$ $\frac{1-F^w(x;\lambda_1,\lambda_2,\alpha,\theta)}{1-F^w(x;\lambda_1,\lambda_2,\alpha,\theta)}$ can also be expressed using the mixture representation [\(9\)](#page-4-2). It will take a complicated expression. However, the behavior of the hrf can be easily assessed: we can verify that $\lim_{x\to 0^+} h^w(x; \lambda_1, \lambda_2, \alpha, \theta) = 0$ and $\lim_{x\to+\infty} h^w(x; \lambda_1, \lambda_2, \alpha, \theta) = \lambda_1$, where the latter follows by L'Hospital rule.

Figures [2](#page-6-0) and [3](#page-6-1) illustrate the behavior of the hrf for different parameter values. We see that $h^{w}(x; \lambda_1, \lambda_2, \alpha, \theta)$ of the GWED is more flexible than the hrf of the WED. The hrfs of the GWED appear monotonic for negative dependence and non-monotonic for positive dependence.

3.1 Some mathematical properties

Let X be a GWE random variable. Proposition 5 derives the nth moment and the moment generating function of X.

Figure 2: (a) The hrf of the GWED for $\theta < 0$, $\alpha = 0.5$ (red line), $\theta < 0$, $\alpha = 1$ (green line), $\theta < 0$, $\alpha = 2$ (blue line) and $\theta < 0$, $\alpha = 5$ (black line); (b) The hrf of the GWED for $\theta > 0$, $\alpha = 0.5$ (red line), $\theta > 0$, $\alpha = 1$ (green line), $\theta > 0$, $\alpha = 2$ (blue line) and $\theta > 0$, $\alpha = 5$ (black line).

Figure 3: (a) The hrf of the WED for $\alpha = 0.5$ (red line), $\alpha = 1$ (green line), $\alpha = 2$ (blue line) and $\alpha = 5$ (black line); (b) The hrf of the GWED for $\theta = 0$, $\alpha = 0.5$ (red line), $\theta = 0$, $\alpha = 1$ (green line), $\theta = 0$, $\alpha = 2$ (blue line) and $\theta = 0$, $\alpha = 5$ (black line).

Proposition 5 The nth moment and the moment generating function of X can be expressed as

$$
\mu'_n = n! \left\{ \frac{p_1}{\lambda_1^n} + \frac{p_2}{\left(\lambda_1 + \lambda_2\alpha\right)^n} + \frac{p_3}{\left(2\lambda_1 + \lambda_2\alpha\right)^n} + \frac{p_4}{\left(\lambda_1 + 2\lambda_2\alpha\right)^n} + \frac{p_5}{2^n \left(\lambda_1 + \lambda_2\alpha\right)^n} \right\}
$$

and

$$
\mathcal{M}(t) = \frac{p_1 \lambda_1}{\lambda_1 - t} + \frac{p_2 (\lambda_1 + \lambda_2 \alpha)}{\lambda_1 + \lambda_2 \alpha - t} + \frac{p_3 (2 \lambda_1 + \lambda_2 \alpha)}{2 \lambda_1 + \lambda_2 \alpha - t} + \frac{p_4 (\lambda_1 + 2 \lambda_2 \alpha)}{\lambda_1 + 2 \lambda_2 \alpha - t} + \frac{2p_5 (\lambda_1 + \lambda_2 \alpha)}{2 \lambda_1 + 2 \lambda_2 \alpha - t},
$$

respectively, for $\lambda_1 > t$.

Proof. Follows from Proposition 4. \Box

The central moments (μ_r) and cumulants (κ_r) of X can be calculated from

$$
\mu_r = \sum_{k=0}^r (-1)^k {r \choose k} \mu_1'^k \mu_{r-k}' \quad \text{and} \quad \kappa_r = \mu_r' - \sum_{k=1}^{r-1} {r-1 \choose k-1} \kappa_k \mu_{r-k}',
$$

respectively, where $\kappa_1 = \mu'_1$. Note that $\kappa_2 = \mu'_2 - \mu'^2_{1, \kappa_3} = \mu'_3 - 3\mu'_2\mu'_1 + 2\mu'^3_1, \kappa_4 = \mu'_4 - 4\mu'_3\mu'_1 3\mu'^2_2 + 12\mu'_2\mu'^2_1 - 6\mu'^4_1$, etc. The skewness $\gamma_1 = \kappa_3/\kappa_2^{3/2}$ and kurtosis $\gamma_2 = \kappa_4/\kappa_2^2$ follow from the second, third and fourth cumulants.

Figures [4](#page-7-0) and [5](#page-7-1) plot skewness and kurtosis as functions of λ_2 and α when λ_1 and θ are fixed. Skewness assumes values greater than 2, implying that the GWED possesses a wider range of skewness values than the two-parameter WED due to Shakhatreh (2012).

 (a) (b)

Figure 4: Skewness for different values of λ_2 and α when $\lambda_1 = 5$, $\theta = 0.99$. Skewness for different values of λ_2 and α when (a) $\lambda_1 = 5$, $\theta = 0.99$, (b) $\lambda_1 = 5$, $\theta = -0.99$.

 (a) (b)

Figure 5: Kurtosis for different values of λ_2 and α when $\lambda_1 = 5$, $\theta = 0.99$.

The shape of many distributions can be usefully described by conditional moments. These moments play an important role in measuring inequality, for example, income quantiles, Lorenz curve and Bonferroni curve. Proposition 6 derives the nth conditional moment and the conditional moment generating function of X.

Proposition 6 The nth conditional moment and the conditional moment generating function of X can be expressed as

$$
m_n(x_0) = p_1 \frac{\gamma (n+1, \lambda_1 x_0)}{\lambda_1^n} + p_2 \frac{\gamma (n+1, (\lambda_1 + \lambda_2 \alpha) x_0)}{(\lambda_1 + \lambda_2 \alpha)^n} + p_3 \frac{\gamma (n+1, (2\lambda_1 + \lambda_2 \alpha) x_0)}{(2\lambda_1 + \lambda_2 \alpha)^n} + p_4 \frac{\gamma (n+1, (\lambda_1 + 2\lambda_2 \alpha) x_0)}{(\lambda_1 + 2\lambda_2 \alpha)^n} + p_5 \frac{\gamma (n+1, 2(\lambda_1 + \lambda_2 \alpha) x_0)}{2^n (\lambda_1 + \lambda_2 \alpha)^n}
$$

and

$$
\mathcal{M}^{\star}(t) = \frac{p_1\lambda_1 \left[1 - e^{-(\lambda_1 - t)x_0}\right]}{\lambda_1 - t} + \frac{p_2(\lambda_1 + \lambda_2\alpha) \left[1 - e^{-(\lambda_1 + \lambda_2\alpha - t)x_0}\right]}{\lambda_1 + \lambda_2\alpha - t} + \frac{p_3(2\lambda_1 + \lambda_2\alpha) \left[1 - e^{-(2\lambda_1 + \lambda_2\alpha - t)x_0}\right]}{2\lambda_1 + \lambda_2\alpha - t} + \frac{p_4(\lambda_1 + 2\lambda_2\alpha) \left[1 - e^{-(\lambda_1 + 2\lambda_2\alpha - t)x_0}\right]}{\lambda_1 + 2\lambda_2\alpha - t} + \frac{2p_5(\lambda_1 + \lambda_2\alpha) \left[1 - e^{-(2\lambda_1 + 2\lambda_2\alpha - t)x_0}\right]}{2\lambda_1 + 2\lambda_2\alpha - t},
$$

respectively, for $\lambda_1 > t$, where $\gamma(a,x) = \int^x$ 0 $t^{a-1}e^{-t}$ dt denotes the lower incomplete gamma function.

Proof. Follows from Proposition 4. \Box

Probability weighted moments (PWMs) formally defined as

$$
\tau_{s,r} = E\left\{X^s \left[F^w \left(X\right)\right]^r\right\} = \int_0^{+\infty} x^s F^w(x)^r f^w(x) \mathrm{d}x
$$

are used to summarize distributions. They are also used for estimation of parameters especially when the inverse cdf cannot be expressed explicitly. Proposition 7 derives an expression for the PWMs of X.

Proposition 7 The PWMs of X can be expressed as

$$
\tau_{s,r} = \sum_{i\geq 0} c_{r,i}(s+r+i)! \left[\frac{p_1}{\lambda_1^{s+r+i}} + \frac{p_2}{(\lambda_1 + \lambda_2 \alpha)^{s+r+i}} + \frac{p_3}{(2\lambda_1 + \lambda_2 \alpha)^{s+r+i}} + \frac{p_4}{(\lambda_1 + 2\lambda_2 \alpha)^{s+r+i}} + \frac{p_5}{2^{s+r+i}(\lambda_1 + \lambda_2 \alpha)^{s+r+i}} \right],
$$

where $c_{r,0} = d_0^r$ and $c_{r,m} = (a_0 m)^{-1} \sum_{r=0}^{m}$ $k=1$ $[k(r + 1) - m] d_k c_{r,m-k}$ for $m \ge 1$.

Proof. By definition,

$$
\tau_{s,r} = K \int_0^{+\infty} x^{s+r} \left[\sum_{i \ge 0} d_i x^i \right]^r f^w(x; \lambda_1, \lambda_2, \alpha, \theta) dx.
$$

By equation (0.314) in Gradshteyn and Ryzhik (2000),

$$
\left[\sum_{i\geq 0} d_i x^i\right]^r = \sum_{i\geq 0} c_{r,i} x^i,
$$

where $c_{r,0} = d_0^r$ and $c_{r,m} = (a_0 m)^{-1} \sum_{r=0}^{m}$ $k=1$ $[k(r+1) - m] d_k c_{r,m-k}$ for $m \geq 1$. The result follows by using [\(9\)](#page-4-2). \square

We now invert [\(10\)](#page-5-1) to obtain a power series expansion for the quantile function say $x = Q(u)$ of the GWED. We shall use the Lagrange theorem. We assume that the power series expansion

$$
g = F^{w}(x) = g_0 + \sum_{n=1}^{+\infty} f_n (x - x_0)^n
$$

holds, where $F^w(x)$ is analytic at a point x_0 . Then, the inverse function $x = Q(u) = (F^w(u))^{-1}$ exists in the neighborhood of some point $u = u_0$.

Proposition 8 The quantile function of X can be expressed as

$$
Q(u) = \sum_{n=1}^{+\infty} b_n u^n,
$$
\n(11)

where $b_n = q_{n,n-1}/(nd_0^n)$, $q_{n,i} = i^{-1} \sum_{i=1}^{i}$ $m=1$ $[m(n+1) - i] p_m q_{n,i-m}, q_{n,0} = p_0^n = 1$ and $p_i =$ $-m_0^{-1}\sum^i$ $_{k=1}$ $d_{k+1}p_{i-k}, i \geq 1$ with $p_0 = 1$.

Proof. According to Markushevich (1965, volume 2, page 88), a power series of $x = Q(u)$ is

$$
x = Q(u) = x_0 + \sum_{n=1}^{+\infty} h_n (u - u_0)^n,
$$

where

$$
h_n = \frac{1}{n!} \frac{d^{n-1}}{dz^{n-1}} \left\{ [\psi(x)]^n \right\} \Big|_{x=x_0}
$$
 and $\psi(x) = \frac{x - x_0}{F^w(x) - F^w(x_0)}$.

Setting $x_0 = 0$ and $u_0 = 0$, and using [\(10\)](#page-5-1), we have

$$
\psi(x) = \frac{x}{F^w(x) - 0} = \frac{1}{\sum_{i \ge 0} d_i x^i}.
$$

The inverse of the power series \sum $i \geq 0$ $d_i x^i$ follows from Gradshteyn and Ryzhik (2000, equation (0.313) :

$$
\psi(x) = \frac{1}{\sum_{i \ge 0} d_i x^i} = \frac{1}{d_0} \sum_{i \ge 0} p_i x^i,
$$

where the coefficients p_i can be determined from $p_i = -m_0^{-1} \sum_{i=1}^{i}$ $_{k=1}$ $d_{k+1}p_{i-k}, i \geq 1$ with $p_0 = 1$. Using Gradshteyn and Ryzhik (2000, equation (0.314)), we can write $\psi(x)^n = \frac{1}{d!}$ $\frac{1}{d_0^n}$ $\overline{}$ $i \geq 0$ $q_{n,i}x^i,$

where $q_{n,i}$ for $i = 1, 2, \ldots$ are given by $q_{n,i} = i^{-1} \sum_{i=1}^{i}$ $m=1$ $[m(n+1) - i] p_m q_{n,i-m}$, and $q_{n,0} = p_0^n = 1$. The $q_{n,i}$ can be determined from $q_{n,0}, \ldots, q_{n,i-1}$ and therefore from p_0, \ldots, p_i . The derivative of order $(n-1)$ of $\psi(x)^n$ can be expressed as

$$
h_n = \frac{1}{n!} \frac{d^{n-1}}{dx^{n-1}} \{ [\psi(x)]^n \} \Big|_{x=0} = \frac{q_{n,n-1}}{n d_0^n}.
$$

Hence, the power series for the quantile function reduces to [\(11\)](#page-9-0). \Box

3.2 Maximum likelihood estimation

Here, we consider estimation of the parameters $\eta = (\lambda_1, \lambda_2, \alpha, \theta)$ of the GWED by the method of maximum likelihood. We suppose x_1, x_2, \ldots, x_n is a random sample from the GWED. Then, the log-likelihood function is

$$
\ell(\eta) = n \left\{ \log \lambda_1 - \log \left[\frac{\lambda_2 \alpha}{\lambda_1 + \lambda_2 \alpha} + \theta \lambda_1 \left(\frac{2}{2\lambda_1 + \lambda_2 \alpha} + \frac{1}{\lambda_1 + 2\lambda_2 \alpha} - \frac{2}{\lambda_1 + \lambda_2 \alpha} \right) \right] \right\}
$$

$$
- \lambda_1 \sum_{i=1}^n x_i + \sum_{i=1}^n \log \left(1 - e^{-\lambda_2 \alpha x_i} \right) + \sum_{i=1}^n \log \left[1 - \theta e^{-\lambda_2 \alpha x_i} \left(1 - 2e^{-\lambda_1 x_i} \right) \right].
$$

Differentiating $\ell(\eta)$ with respect to λ_1 , λ_2 , α and θ , we obtain the normal equations

$$
\frac{\partial \ell(\eta)}{\partial \lambda_1} = \frac{n}{\lambda_1} - n\lambda_2 \alpha \frac{A}{B} - \sum_{i=1}^n x_i - 2\theta \sum_{i=1}^n x_i \frac{e^{-(\lambda_1 + \lambda_2 \alpha)x_i}}{1 - \theta e^{-\lambda_2 \alpha x_i} \left(1 - 2e^{-\lambda_1 x_i}\right)} = 0,
$$
\n
$$
\frac{\partial \ell(\eta)}{\partial \lambda_2} = n\lambda_1 \alpha \frac{A}{B} + \alpha \sum_{i=1}^n \frac{x_i e^{-\lambda_2 \alpha x_i}}{1 - e^{-\lambda_2 \alpha x_i}} + \theta \alpha \sum_{i=1}^n x_i \frac{e^{-\lambda_2 \alpha x_i} - 2e^{-(\lambda_1 + \lambda_2)x_i}}{1 - \theta e^{-\lambda_2 \alpha x_i} \left(1 - 2e^{-\lambda_1 x_i}\right)} = 0,
$$
\n
$$
\frac{\partial \ell(\eta)}{\partial \alpha} = -n\lambda_1 \lambda_2 \frac{A}{B} - \lambda_2 \sum_{i=1}^n \frac{x_i e^{-\lambda_2 \alpha x_i}}{1 - e^{-\lambda_2 \alpha x_i}} + \theta \lambda_2 \sum_{i=1}^n x_i \frac{e^{-\lambda_2 \alpha x_i} - 2e^{-(\lambda_1 + \lambda_2)x_i}}{1 - \theta e^{-\lambda_2 \alpha x_i} \left(1 - 2e^{-\lambda_1 x_i}\right)} = 0,
$$
\n
$$
\frac{\partial \ell(\eta)}{\partial \theta} = -\frac{n\lambda_1 \left(\frac{2}{2\lambda_1 + \lambda_2 \alpha} + \frac{1}{\lambda_1 + 2\lambda_2 \alpha} - \frac{2}{\lambda_1 + \lambda_2 \alpha}\right)}{B} - \sum_{i=1}^n \frac{e^{-\lambda_2 \alpha x_i} \left(1 - 2e^{-\lambda_1 x_i}\right)}{1 - \theta e^{-\lambda_2 \alpha x_i} \left(1 - 2e^{-\lambda_1 x_i}\right)} = 0,
$$

where

$$
A = -\frac{1}{(\lambda_1 + \lambda_2 \alpha)^2} + 2\theta \left[\frac{1}{(2\lambda_1 + \lambda_2 \alpha)^2} + \frac{1}{(\lambda_1 + 2\lambda_2 \alpha)^2} - \frac{1}{(\lambda_1 + \lambda_2 \alpha)^2} \right]
$$

and

$$
B = \frac{\lambda_2 \alpha}{\lambda_1 + \lambda_2 \alpha} + \theta \lambda_1 \left(\frac{2}{2\lambda_1 + \lambda_2 \alpha} + \frac{1}{\lambda_1 + 2\lambda_2 \alpha} - \frac{2}{\lambda_1 + \lambda_2 \alpha} \right).
$$

The maximum likelihood estimates say $\hat{\eta} = (\hat{\lambda}_1, \hat{\lambda}_2, \hat{\alpha}, \hat{\theta})$ are the simultaneous solutions of the normal equations. These equations do not yield explicit solutions. Hence, the maximum likelihood estimates must be obtained numerically.

Table 1: Descriptive statistics of the CRP data.

	Minimum Maximum Mean Median SD			Skewness Kurtosis	
0.15	7.85	1.8867 1.485		1.5450 1.1777	4.1425

According to Cox and Hinkley (1979), the asymptotic distribution of $\hat{\eta}$ can be approximated by the multivariate normal distribution, $N_4\left(0,[\mathbf{J}(\hat{\boldsymbol{\eta}})]^{-1}\right)$, where $\mathbf{J}(\hat{\boldsymbol{\eta}})$ denotes the inverse of the observed information matrix evaluated at $\hat{\eta}$. Due to algebraic complexity and in order to save space, we have not reported the expression for $J(\hat{\eta})$. This approximation can be used to construct approximate confidence intervals and hypothesis tests for λ_1 , λ_2 , α and θ .

Numerical calculations not reported here showed that the surface of the $\ell(\eta)$ was smooth. Numerical routines for maximization of $\ell(\eta)$ were able to locate the maximum for a wide range of starting values. However, to ease computations it is useful to have reasonable starting values. These can be obtained, for example, by the method of moments. For $r = 1, 2, 3, 4$ let $m_r =$ 1 n $\sum_{n=1}^{\infty}$ $\frac{i=1}{i}$ x_i^r denote the first four sample moments. Equating these moments with the theoretical versions given in Section 3.1, we have $m_r = E(X^r)$ for $r = 1, 2, 3, 4$. These equations can be solved simultaneously to obtain the moments estimates.

3.3 Application

Here, we illustrate the flexibility of the GWED using two biochemical real data sets: C-reactive protein (CRP) data and insulin data. These data were collected through a study whose aim was to examine the correlation of the RBP4 (retinol-binding protein-4) with anthropometric measurements (BMI-body mass index and WC-waist circumference), insulin resistance, metabolic and kidney parameters and inflammation. The participants of the study were 128 obese diabetic patients. This research was carried out in Primary Health Care Center (Center of Laboratory Diagnostics) and Center of Clinical-Laboratory Diagnostics, Clinical Center of Montenegro. These data sets are original in that they have not been analyzed in the statistics literature before.

We fitted the GWED to both data sets by the method of maximum likelihood. The fit of the GWED was compared with that of the WED due to Gupta and Kundu (2009). As criteria for comparison, we used the Akaike information criterion (AIC) and the p-value of the Kolmogorov-Smirnov test.

Both distributions were fitted by executing the R function optimx (Nash and Varadhan, 2011) for a wide range of starting values. This sometimes resulted in more than one maximum, but at least one maximum was identified each time. In cases of more than one maximum, we took the maximum likelihood estimates to correspond to the largest of the maxima.

3.3.1 Application 1: CRP data

CRP is a protein found in the blood, the levels of which rise in response to inflammation (i.e., CRP is an acute-phase protein). The CRP gene is located on the first chromosome (1q21-q23). Table [1](#page-11-0) gives descriptive statistics of the CRP data set. We see that the data has positive skewness and kurtosis greater than that of the normal distribution.

The MLEs of the model parameters, their standard errors, the AIC values, and the p-values of the Kolmogorov-Smirnov test for the CRP data are reported in Table [2.](#page-12-0) Based on the AIC values and the p-values, we see that the GWED provides a better fit than the WED for the CRP data.

Figure [6\(](#page-12-1)a) plots the survival function for the fitted GWED and the empirical survival function for the CRP data. Figure [6\(](#page-12-1)b) plots the pdfs for the fitted GWED and WED and the empirical pdf for the CRP data. Figure $7(a)$ plots the cdfs for the fitted GWED and WED and

Table 2: MLEs of the model parameters, AIC values and p-values for the CRP data.

Model λ_1	λ 2	α		λ	AIC	<i>p</i> -value
GWED 0.7280	0.8316	0.8121 (0.0874) (0.0792) (0.1010) (0.0936)	0.7991		574.84 0.819	
WED		8.5271 (1.0024)		0.5857 (0.08715)	777.76 0	

the empirical cdf for the CRP data. All these figures suggest that the GWED provides a better fit to the CRP data.

 (a) (b)

Figure 6: (a) Fitted GWE survival function and the empirical survival function for the CRP data; (b) Fitted GWE and WE pdfs and the empirical pdf for the CRP data.

Figure 7: (a) Fitted GWE and WE cdfs and the empirical cdf for the CRP data; (b) Fitted GWE and WE cdfs and the empirical cdf for the insulin data.

3.3.2 Application 2: Insulin data

Insulin is a peptide hormone, produced by beta cells of the pancreas, and is central to regulating carbohydrate and fat metabolism in the body. Insulin causes cells in the liver, skeletal muscles, and fat tissue to absorb glucose from the blood. In the liver and skeletal muscles, glucose is stored as glycogen, and in fat cells (adipocytes) it is stored as triglycerides. The descriptive statistics of the insulin data are given in Table [3.](#page-13-0) This data is also positively skewed and has kurtosis greater than that of the normal distribution.

The MLEs of the model parameters, their standard errors, the AIC values, and the p-values of the Kolmogorov-Smirnov test for the insulin data are reported in Table [3.](#page-13-0) Based on the AIC

Table 3: Descriptive statistics of the insuline data.						
	Minimum Maximum Mean Median SD				Skewness Kurtosis	
2.4	72.4	9.616 6.8		9.1306 1.7050		24.158

Table 4: MLEs of the model parameters, AIC values and p-values for the insulin data.

values and the p-values, we see again that the GWED provides a better fit than the WED for the insulin data.

Figure [8\(](#page-13-1)a) plots the survival function for the fitted GWED and the empirical survival function for the insulin data. Figure [8\(](#page-13-1)b) plots the pdfs for the fitted GWED and WED and the empirical pdf for the insulin data. Figure [7\(](#page-12-2)b) plots the cdfs for the fitted GWED and WED and the empirical cdf for the insulin data. All these figures suggest that the GWED again provides a better fit to the insulin data.

$$
\rm (a)\qquad \qquad (b)
$$

0.0 0.4 0.8

 0.4

 $\overline{0}$

0 10 20 30 40 50 60 70

95% CI Kaplan−−Meier GWED model Frequency 0 20 40 60 PDF of the GWED PDF of the WED

Figure 8: (a) Fitted GWE survival function and the empirical survival function for the insulin data; (b) Fitted GWE and WE pdfs and the empirical pdf for the insulin data.

x

0 20 40 60 80

4 Concluding remarks and future research

In this paper, we have proposed an extension of Azzalini's method. As an illustration of this extension, we have introduced a four-parameter Generalized Weighted Exponential distribution (GWED). This new distribution generalizes of the weighted exponential distribution due to Gupta and Kundu (2009). The method that we have proposed can be used to generalize any distribution.

We have studied mathematical properties of the GWED. We have shown that the GWED can be interpreted as a hidden truncated distribution (Arnold and Beaver, 2000a). The dependence parameter rather than the marginal parameters plays an important role in making the GWED flexible in terms of skewness, kurtosis, shape of the hazard rate function and other characteristics.

We have discussed maximum likelihood estimation of the parameters of the GWED and provided two real data applications. The applications show that the GWED can be used quite effectively to give better fits than the WED. We hope that the GWED may attract wider

applications in statistics.

In conclusion, it is important to highlight that although this paper has mainly concentrated on the GWED, our method can be used for any distribution. For instance, one can consider a general class of cdfs defined by

$$
F(x) = F(x; a, b, c) = [a\xi(x) + b]^{c},
$$

where $a \neq 0$, $b \neq 0$ and $c \neq 0$ are parameters, and $\xi(x)$ is a monotonic and differentiable function on $(F^{-1}(0), F^{-1}(1))$. This class for proper choices of a, b, c and $\xi(\cdot)$ can contain various distributions studied in the literature like the power function, Dagum (Burr type III), Pareto, inverse Weibull, reflected exponential and rectangular distributions.

Acknowledgement

We are grateful to Dr. Aleksandra Klisić, a biochemist in Primary Health Unit in Podgorica, Montenegro, for allowing us to use the CRP and insulin data.

We would like to thank to the Editor and to anonymous referees whose comments greatly improved quality of the paper.

References

Arnold, B.C. and Beaver, R.J. (2000a). Hidden truncation models. Sankhya, 62, pp. 23-35.

Arnold, B.C. and Beaver, R.J. (2000b). The skew-Cauchy distribution. Statistics and Probability Letters, 49, pp. 285-290.

Aryal, G. and Nadarajah, S. (2005). On the skew-Laplace distribution. Journal of Information and Optimization Sciences, 26, pp. 205-217.

Azzalini, A. (1985). A class of distributions which includes the normal ones. Scandinavian Journal of Statistics, 12, pp. 171-178.

Cox, D.R. and Hinley, D.V. (1979). Theoretical Statistics. Chapman and Hall, London.

Domma, F. and Giordano, S. (2013). A copula-based approach to account for dependence in stress-strength models. Statistical Papers, 54, pp. 807-826.

Gradshteyn, I.S. and Ryzhik, I.M. (2000). Table of Integrals, Series, and Products. Academic Press, San Diego.

Gupta, A.K., Chang, F.C. and Haung, W.J. (2002). Some skew-symmetric models. Random Operators and Stochastic Equations, 10, pp. 133-140.

Gupta, R.D. and Kundu, D. (2009). A new class of weighted exponential distributions. Statistics, 43, pp. 621-634.

Kotz, S., Lumelskii, Y. and Pensky, M. (2003). The Stress-Strength Model and Its Generalizations: Theory and Applications. World Scientific Publishing, Singapore.

Mahdy, M. (2011). A class of weighted gamma distributions and its properties. Economic Quality Control, 26, pp. 133-144.

Mahdy, M. (2013). A class of weighted Weibull distributions and its properties. Studies in Mathematical Sciences, 6, pp. 35-45.

Markushevich, A.I. (1965). Theory of Functions of a Complex Variable. Chelsea Publication Company.

Morgenstern, D. (1956). Einfache Beispiele zweidimensionaler Verteilungen. Mitteilingsblatt für Mathematische Statistik, 8, pp. 234-235.

Nadarajah, S. (2009). On the skew-logistic distribution. AStA Advances in Statistical Analysis, 93, pp. 187-203.

Nadarajah, S. (2014). Expansions for bivariate copulas. Submitted.

Nash, J.C. and Varadhan, R. (2011). Unifying optimization algorithms to aid software system users: optimx for R. Journal of Statistical Software, 43, pp. 1-14.

Shakhatreh, M.K. (2012). A two-parameter of weighted exponential distributions. Statistics and Probability Letters, 82, pp. 252-261.

Titterington, D.M., Smith, A.F.M. and Makov, U.E. (1985). Statistical Analysis of Finite Mixture Distributions. John Wiley and Sons, New York.