

# Dagger and Dilation

in the Category of Von Neumann Algebras

Bastiaan Eelco WESTERBAAN

Third corrected version  
(april 26th, 2019)

### Identifiers

hdl: 2066/201785

arXiv: 1803.01911

ISBN: 978-94-6332-485-4

### Persistent links

<https://arxiv.org/abs/1803.01911>

<https://doi.org/2066/201785>

<https://hdl.handle.net/2066/201785>

### Source code

$\LaTeX$  <https://github.com/westerbaan/theses>

cover <https://github.com/westerbaan/ndpt>

Printed by GVO drukkers & vormgevers B.V., Ede, <https://proefschriften.nl>.

Where applicable, © 2019 B.E. Westerbaan,  available under CC BY, [1].

# Dagger and Dilation

in the Category of Von Neumann Algebras

## Proefschrift

ter verkrijging van de graad van doctor  
aan de Radboud Universiteit Nijmegen  
op gezag van de rector magnificus prof. dr. J.H.J.M. VAN KRIEKEN,  
volgens besluit van het college van decanen  
in het openbaar te verdedigen

op

**dinsdag 14 mei 2019**

om

**12.30 uur precies**

door

**Bastiaan Eelco WESTERBAAN**

geboren op 30 augustus 1988  
te Nijmegen

**Promotor:**

Prof. dr. B.P.F. JACOBS

**Manuscriptcommissie:**

Prof. dr. J.D.M. MAASSEN

Prof. dr. P. PANANGADEN  
(McGill University, Canada)

Prof. dr. P. SELINGER  
(Dalhousie University, Canada)

Dr. C.J.M. HEUNEN  
(University of Edinburgh, Verenigd Koninkrijk)

Dr. S. STATON  
(University of Oxford, Verenigd Koninkrijk)

# Dagger and Dilation

in the Category of Von Neumann Algebras

## Doctoral Thesis

to obtain the degree of doctor  
from Radboud University Nijmegen  
on the authority of the Rector Magnificus prof. dr. J.H.J.M. VAN KRIEKEN,  
according to the decision of the Council of Deans  
to be defended in public

on

**Tuesday, May 14, 2019**

at

**12.30 hours**

by

**Bastiaan Eelco WESTERBAAN**

born on August 30, 1988  
in Nijmegen (the Netherlands)

## **Supervisor:**

Prof. dr. B.P.F. JACOBS

## **Doctoral Thesis Committee:**

Prof. dr. J.D.M. MAASSEN

Prof. dr. P. PANANGADEN  
(McGill University, Canada)

Prof. dr. P. SELINGER  
(Dalhousie University, Canada)

Dr. C.J.M. HEUNEN  
(University of Edinburgh, United Kingdom)

Dr. S. STATON  
(University of Oxford, United Kingdom)



<b>1</b>	<b>Introduction</b>	<b>134</b>
<b>2</b>	<b>Dilations</b>	<b>135</b>
2.1	Stinespring's theorem . . . . .	136
2.2	Hilbert $C^*$ -modules . . . . .	141
2.2.1	The basics of Hilbert $C^*$ -modules . . . . .	142
2.2.2	Ultranorm uniformity . . . . .	146
2.2.3	Self-dual completion . . . . .	150
2.3	Paschke dilations . . . . .	154
2.4	Kaplansky density theorem for Hilbert $C^*$ -modules . . . . .	158
2.5	More on self-dual Hilbert $C^*$ -modules . . . . .	159
2.5.1	Exterior tensor product . . . . .	164
2.6	Pure maps . . . . .	168
<b>3</b>	<b>Diamond, and then, dagger</b>	<b>173</b>
3.1	Effect algebras and related structures . . . . .	174
3.1.1	Effect monoids . . . . .	178
3.1.2	Effect modules . . . . .	179
3.2	Effectuses . . . . .	180
3.2.1	From partial to total . . . . .	181
3.2.2	From total to partial . . . . .	182
3.2.3	Predicates, states and scalars . . . . .	190
3.2.4	Effectus of abstract $M$ -convex sets . . . . .	193
3.3	Effectuses with quotients . . . . .	197
3.4	Effectuses with comprehension and images . . . . .	199
3.4.1	Sharp predicates . . . . .	203
3.5	$\diamond$ -effectuses . . . . .	206
3.6	$\&$ -effectuses . . . . .	211
3.7	$\dagger$ -effectuses . . . . .	214
3.7.1	Dilations . . . . .	221
3.8	Comparisons . . . . .	224
3.8.1	Dagger kernel categories . . . . .	224
3.8.2	Sequential product . . . . .	225
3.8.3	Homological categories . . . . .	226
<b>4</b>	<b>Outlook</b>	<b>229</b>

# Chapter 1

## Introduction

This thesis is a mathematical study of quantum computing, concentrating on two related, but independent topics. First up are *dilations*, covered in chapter 2. In chapter 3 “diamond, and then, dagger” we turn to the second topic: *effectus theory*. Both chapters, or rather parts, can be read separately and feature a comprehensive introduction of their own. For now, we focus on what they have in common.

134

Both parts study the category of von Neumann algebras (with ncpsu-maps\* between them). A von Neumann algebra is a model for a (possibly infinite-dimensional) quantum datatype, for instance:

II

$M_2$	a qubit	$M_3 \otimes M_3$	an (ordered) pair of qutrits
$M_3$	a qutrit	$\mathbb{C}^2$	a bit
$M_2 \oplus M_3$	a qubit or a qutrit	$\mathcal{B}(\ell^2)$	quantum integers

The ncpsu-maps between these algebras represent physical (but not necessarily computable) operations in the opposite direction (as in the ‘Heisenberg picture’):

1. *Measure a qubit in the computational basis* (qubit  $\rightarrow$  bit)  
 $\mathbb{C}_2 \rightarrow M_2, \quad (\lambda, \mu) \mapsto \lambda |0\rangle\langle 0| + \mu |1\rangle\langle 1|,$  using bracket notation<sup>†</sup>
2. *Discard the second qubit* (qubit  $\times$  qubit  $\rightarrow$  qubit)  
 $M_2 \rightarrow M_2 \otimes M_2, \quad a \mapsto a \otimes 1$
3. *Apply a Hadamard-gate to a qubit* (qubit  $\rightarrow$  qubit)  
 $M_2 \rightarrow M_2, \quad a \mapsto H^* a H,$  where  $H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$

---

\*An abbreviation for normal completely positive contractive (i.e. subunital) linear maps.

<sup>†</sup>See e.g. [91, fig. 2.1], so here  $|0\rangle \equiv \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ,  $\langle 0| \equiv (1 \ 0)$ ,  $|1\rangle \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  and  $|0\rangle\langle 0| = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ .

4. *Initialize a qutrit as 0* (empty  $\rightarrow$  qutrit)  
 $M_3 \rightarrow \mathbb{C}, \quad a \mapsto \langle 0|a|0\rangle$

- III The first part of this thesis starts with a concrete question: does the famous Stinespring dilation theorem [106], a cornerstone of quantum theory, extend to every ncpsu-map between von Neumann algebras? Roughly, this means that every ncpsu-map splits in two maps of a particularly nice form. We will see that indeed it does. The insights attained pondering this question lead naturally, as they so often do in mathematics, to other new results: for instance, we learn more about the tensor product of dilations (167I) and the pure maps associated to them (171VII). But perhaps the mathematically most interesting by-catch of pondering the question are the contributions (see 135VI) to the theory of self-dual Hilbert  $C^*$ -modules over a von Neumann algebra, which underlie these dilations.
- IV In contrast to the first part of this thesis, the second part starts with the abstract (or rather: vague) question: what characterizes the category of von Neumann algebras? The goal is to find a set of reasonable axioms which pick out the category of von Neumann algebras among all other categories. In other words: a reconstruction of quantum theory with categorical axioms (presupposing von Neumann algebras are the right model of quantum theory). Unfortunately, we do not reach this goal here.<sup>‡</sup> However, by studying von Neumann algebras with this self-imposed restriction, one is led to, as again is often the case in mathematics, upon several new notions and results, which are of independent interest, such as, for instance,  $\diamond$ -adjointness, pure maps and the existence of their dagger.
- V **Twin theses** A substantial part of the research presented here has been performed in close collaboration with my twin brother Abraham Westerbaan. Our results are interwoven to such a degree that separating them is difficult and, more importantly, detrimental to the clarity of presentation. Therefore we decided to write our theses as two consecutive volumes (with consecutive numbering, more on that later). Abraham's thesis [124] is the first volume and covers the preliminaries on von Neumann algebras and, among other results, an axiomatization of the sequential product  $b \mapsto \sqrt{ab}\sqrt{a}$ . The present thesis is the second volume and builds upon the results of Abraham's thesis, in its study of dilations and effectus theory. Because of this clear division, a significant amount of Abraham's original work appears in my thesis and, vice versa, my work appears in his. However, the large majority of the new results in each of our theses is our own.
- VI **Preliminaries** For the first part of this thesis, only knowledge is presumed of, what is considered to be, the basic theory of von Neumann algebras. Unfor-

---

<sup>‡</sup>Building on this thesis, van de Wetering recently found a reconstruction of finite-dimensional quantum theory [117, 118].

tunately, this ‘basic theory’ takes most introductory texts close to a thousand pages to develop. This situation has led my twin brother to ambitiously redevelop the basic theory of von Neumann algebras tailored to the needs of our theses (and hopefully others in the field). In his thesis, Abraham presents [124] a clear, comprehensive and self-contained development of the basic theory of von Neumann algebras in less than 200 pages. For a traditional treatment, I can wholeheartedly recommend the books of Kadison and Ringrose [73]. For the second part of this thesis, only elementary category theory is required as covered in any introductory text, like e.g. [9].

**Writing style** This thesis is divided into points, each a few paragraphs long. Points are numbered with Roman numerals in the margin: this is point VII. Points are organized into groups, which are assigned Arabic numerals, that appear at the start of the group in the margin. The group of this point is 134. Group numbers below 134 appear in my twin brother’s thesis [124]. Page numbers have been replaced by the group numbers that appear on the spread.<sup>§</sup> This unusual system allows for very short references: instead of writing ‘see Theorem 2.3.4 on page 123’, we simply write ‘see 154 III’. Even better: ‘see the middle of the proof of Theorem 2.3.4 on page 123’ becomes the more precise ‘see 154 VIII’.

Somewhat unconventionally, this thesis includes exercises. Most exercises replace straight-forward, but repetitive, proofs. Other exercises, marked with an asterisk \*, are harder and explore tangents. In both cases, these exercises are not meant to force the reader into the role of a pupil, but rather to aid her, by reducing clutter. Solutions to these exercises (page 197) and errata to the printed version of this text (page 239), can be found at the end of this PDF. Possibly further errata will be made available on the arXiv under 1803.01911.

**Advertisements** This thesis is based on the publications [22,69,126,127]. During my doctoral research, I also published the following papers, which although somewhat related, would’ve made this thesis too broad in scope if included.

1. In *A Kochen–Specker System has at least 22 Vectors* [116], Uijlen and I justify the title building upon the work of Arends, Ouaknine and Wampler.
2. In *Unordered Tuples in Quantum Computation* [35] Furber and I argue that  $M_3 \oplus \mathbb{C}$  is the right algebra modeling an unordered pair of qubits. Then we compute the algebra for an unordered  $n$ -tuple of  $d$ -level systems.
3. In *Picture-perfect QKD* [76] Kissinger, Tull and I show how to graphically derive a security bound on quantum key distribution using the continuity of Stinespring’s dilation.
4. In [100], Schwabe and I work out the details of a quantum circuit to break binary  $\mathcal{MQ}$  using Grover’s algorithm.

---

<sup>§</sup>A spread are two pages that are visible at the same time.

In the second part of this thesis, we use the notion of effectus (introduced by Jacobs [57]), solely for studying quantum theory. Effectus theory, however, has more breadth to it, see e.g. [3, 20, 21, 58–64, 67, 68, 70]. For a long, but comprehensive introduction, see [23].

IX Next, I would like to mention some other recent approaches to the mathematical study of quantum theory which are related in spirit and of which I am aware. First, there are colleagues who also study particular categories related to quantum theory. Spearheaded by Coecke and Abramsky [2], dozens of authors around Oxford have developed insightful graphical calculi by studying the category of Hilbert spaces categorically [25]. Not explicitly categorically (but certainly in spirit) is the work around operational probabilistic theories, see e.g. [27]. Besides these approaches (and that of effectus theory), there are many other varied categorical forages into quantum theory, e.g. [34, 78, 96, 105]. Then there are authors who pick out different structures from quantum theory to study. Most impressive is the work of Alfsen and Shultz, who studied and axiomatized the state spaces of many operator algebras, including von Neumann algebras [4]. Recently, the poset of commutative subalgebras of a fixed operator algebra has received attention, see e.g. [51, 52, 85]. Finally, tracing back to von Neumann himself, the lattice of projections (representing sharp predicates) and the poset of effects (fuzzy predicates), has have been studied by many, see e.g. [29].

IXa **About the cover** One of the most familiar structures in operator algebras (the mathematical field of this thesis) is perhaps the infinite-dimensional Hilbert space  $\ell^2(\mathbb{N})$ . What would its unit sphere look like? The cover depicts an approximation: a 12-dimensional spherical shell split into two. On the outside the pieces are completely reflective — on the inside they are dark blue and only mostly reflective. The pieces are sat between two (11-dimensional) hyperplanes with a somewhat reflective checkerboard pattern. The planes are cropped at a certain radius from the origin turning them into *hyperdisks*. The picture is rendered using a homebrew<sup>¶</sup> *path tracer*, which simulates how rays of light would bounce to end up in the camera. The calculations involved (such as the one to determine how a ray bounces from an  $n$ -dimensional sphere) are all conveniently expressed and performed using inner products. The primary rays cast by the camera have the biggest presence in the first three dimensions, only ten percent in the fourth, five percent in the fifth et cetera. This causes more higher-dimensional oddities to appear in the image where rays have travelled further or bounced farther in the higher dimensions.

X **Funding** My research has been supported by the European Research Council under grant agreement № 320571.

XI **Acknowledgments** Progress in theoretical fields is often dramatized as the fruit

---

<sup>¶</sup>Source code is available at <https://github.com/westerbaan/ndpt>.

of personal struggles of *einzelgängers* grown in deep focus and isolation. But what is missing in this depiction is that often the seeds were sown in casual conversation among colleagues. I would like to acknowledge two seeds I'm aware of. First, it was Sam Staton who suggested looking at a universal property for instruments, which led to quotients and comprehension. Secondly, Chris Heunen insisted, against my initial skepticism, that Stinespring's dilation must have a universal property. I've also had the immense pleasure of discussing research with Robin Adams, Guillaume Allais, Henk Barendregt, Kenta Cho, Joan Daemen, Robert Furber, Bart Jacobs, Bas Joosten, Martti Karvonen, Aleks Kissinger, Mike Koeman, Hans Maassen, Joshua Moerman, Teun van Nuland, Dusko Pavlovic, Arnoud van Rooij, Frank Roumen, Marc Schoolderman, Peter Schwabe, Peter Selinger, Alexandra Silva, Michael Skeide, Pawel Sobocinski, Sean Tull, Sander Uijlen, Marloes Venema, Abraham Westerbaan, John van de Wetering and Fabio Zanasi. I'm particularly grateful to Arnoud, Abraham and John for proofreading this thesis. I'm honored to have been received by Dusko for a two-month research visit, on which he, Jason, Jon, Mariam, Liang-Ting, Marine, Muzamil, Pawel and Peter-Michael made me feel at home on the other side of the planet. On this side of the planet, it is hard to imagine a friendlier group of colleagues. I'm much obliged to Marko van Eekelen, Wouter Geraedts, Engelbert Hubbers, Carlo Meijer and Joeri de Ruiter with whom I've performed security audits as a side-gig, which allowed for an extension of my position. I'm grateful to my supervisor, Bart Jacobs, who showed me a realistic optimism, which is in such short supply these days. I'm touched by the patience of my friends, (extended) family and Annelies, during the final stretch of writing this thesis.

Last, but not least, I would like to share my deep appreciation for the (under)graduate education I've been fortunate enough to receive at the Radboud Universiteit. Only now, a few years later, I can see how rare it is, not only for its uncompromised treatment of the topics at hand, but also for the personal attention to each student. Of all the great professors, Henk Barendregt, Mai Gehrke, Arnoud van Rooij and Wim Veldman have been of particular personal influence to me — I owe them a lot.



# Chapter 2

## Dilations

In this chapter we will study dilations. The common theme is that a complicated map is actually the composition of a simpler map after a representation into a larger algebra. The famous Gel'fand–Naimark–Segal construction (see 30 VI in [124]) is a dilation theorem in disguise: it shows that every state is a vector state on a larger algebra: 135

**Theorem (GNS')** For each pu-map\*  $\omega: \mathcal{A} \rightarrow \mathbb{C}$  from a C\*-algebra  $\mathcal{A}$ , there is a Hilbert space  $\mathcal{H}$ , an miu-map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  and a vector  $x \in \mathcal{H}$  such that  $\omega = h \circ \varrho$  where  $h: \mathcal{B}(\mathcal{H}) \rightarrow \mathbb{C}$  is given by  $h(T) \equiv \langle x, Tx \rangle$ . II

Probably the most famous dilation theorem is that of Stinespring [106, thm. 1]. (We will see a detailed proof later, in 137 II.) III

**Theorem (Stinespring)** For every cp-map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  between a C\*-algebra  $\mathcal{A}$  and the C\*-algebra of bounded operators on a Hilbert space  $\mathcal{H}$ , there is a Hilbert space  $\mathcal{K}$ , an miu-map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{K})$  and a bounded operator  $V: \mathcal{H} \rightarrow \mathcal{K}$  such that  $\varphi = \text{ad}_V \circ \varrho$ , where  $\text{ad}_V: \mathcal{B}(\mathcal{K}) \rightarrow \mathcal{B}(\mathcal{H})$  is given by  $\text{ad}_V(T) \equiv V^*TV$ . Furthermore IV

1. if  $\varphi$  is unital, then  $V$  is an isometry *and*
2. if  $\mathcal{A}$  is a von Neumann algebra and  $\varphi$  is normal, then  $\varrho$  is normal as well.

Stinespring's theorem is fundamental in the study of quantum information and quantum computing; it is used to prove entropy inequalities (e.g. [84]), bounds on optimal cloners (e.g. [123]), full completeness of quantum programming languages (e.g. [105]), security of quantum key distribution (e.g. [76,80]), to analyze V

---

\*pu-map is an abbreviation for positive unital linear map. See 10 II and 44 XV for other abbreviations, like nmiu-map and npsu-map.

quantum alternation (e.g. [13]), to categorify quantum processes (e.g. [102]) and as an axiom to single out quantum theory among information processing theories [18]. These are only a few examples of the many thousands building on Stinespring’s discovery.

Stinespring’s theorem only applies to maps of the form  $\mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  and so do most of its useful consequences. One wonders: is there an extension of Stinespring’s theorem to arbitrary ncp-maps  $\mathcal{A} \rightarrow \mathcal{B}$ ? A different and perhaps less frequently asked question is whether Stinespring’s dilation is categorical in some way? That is: does it have a defining universal property? Both questions turn out to have an affirmative answer: Paschke’s generalization of GNS for Hilbert  $C^*$ -modules [93] turns out to have the same universal property as Stinespring’s dilation and so it extends Stinespring to arbitrary ncp-maps.

Dilation	Maps	LHS	RHS
(normal) GNS [36], 30 VI, II	(n)p-maps $\mathcal{A} \xrightarrow{\varphi} \mathbb{C}$ (Hilbert space $\mathcal{H}$ ; von Neumann algebra $\mathcal{A}$ ; $x \in \mathcal{H}$ )	(n)miu $\mathcal{A} \xrightarrow{\varrho} \mathcal{B}(\mathcal{H})$	ncp vector state $\mathcal{B}(\mathcal{H}) \xrightarrow{\langle x, (\cdot)x \rangle} \mathbb{C}$
(normal) Stinespring [106], IV	(n)cp-maps $\mathcal{A} \xrightarrow{\varphi} \mathcal{B}(\mathcal{H})$ (Hilbert spaces $\mathcal{H}, \mathcal{K}$ ; vN alg. $\mathcal{A}$ ; $V \in \mathcal{B}(\mathcal{H}, \mathcal{K})$ )	(n)miu $\mathcal{A} \xrightarrow{\varrho} \mathcal{B}(\mathcal{H})$	pure ncp-map $\mathcal{B}(\mathcal{H}) \xrightarrow{\text{ad}_V} \mathcal{B}(\mathcal{H})$
Paschke [93, 127], 154 III	ncp-maps $\mathcal{A} \xrightarrow{\varphi} \mathcal{B}$ (Von Neumann algebras $\mathcal{A}, \mathcal{B}, \mathcal{P}$ )	nmiu $\mathcal{A} \xrightarrow{\varrho} \mathcal{P}$	pure ncp-map $\mathcal{P} \xrightarrow{h} \mathcal{B}$
KSGNS [75], cf. [81]	cp-maps $\mathcal{A} \xrightarrow{\varphi} \mathcal{B}^a(E)$ ( $C^*$ -algs $\mathcal{A}, \mathcal{B}$ ; Hilb. $\mathcal{B}$ -modules $E, F$ ; $T \in \mathcal{B}^a(F, E)$ )	miu $\mathcal{A} \xrightarrow{\varrho} \mathcal{B}^a(F)$	cp-map $\mathcal{B}^a(F) \xrightarrow{\text{ad}_T} \mathcal{B}^a(E)$

Table 2.1: Various dilation theorems. Maps  $\varphi$  of the given type split as  $\varphi = h \circ \varrho$ , where  $\varrho$  is as in LHS and  $h$  is as in RHS. The first three appear prominently in this thesis; the last, KSGNS, is only briefly touched upon in 155.

VI We start this chapter with a detailed proof of Stinespring’s theorem. Then we show that it obeys a universal property. Before we move on to Paschke’s GNS, we need to develop Paschke’s theory of self-dual Hilbert  $C^*$ -modules. Paschke’s work builds on Sakai’s characterization of von Neumann algebras, which would take considerable effort to develop in detail. Thus to be as self-contained as possible, we avoid Sakai’s characterization (in contrast to our article [127]) and give new proofs of Paschke’s results where required. One major difference is that we will use a completion (see 150 II) of a uniform space instead of considering the dual space of a Hilbert  $C^*$ -module. This uniformity (which we call the *ultranorm uniformity*) plays a central rôle in our development of the theory and many of the new results. We finish the first part of this chapter by constructing Paschke’s

dilation and establishing that it indeed extends Stinespring’s dilation. We start the second part of this chapter by characterizing when the representation in the Paschke dilation of an ncp-map is injective (see 156 II), which is a generalization of our answer [125] to the same question about the Stinespring dilation. Then we allow for an intermezzo to prove several new results on (self-dual) Hilbert  $C^*$ -modules, notably:

1. we generalize the Kaplansky density theorem to Hilbert  $C^*$ -modules, see 158 II;
2. we prove that every self-dual Hilbert  $C^*$ -module over a von Neumann algebra factor is of a particularly nice form, see 162 IV, and
3. we greatly simplify the construction of the exterior tensor product for Hilbert  $C^*$ -modules over von Neumann algebras in 164 II and show that its self-dual completion is particularly well-behaved, see e.g. 165 VI.

We use this new insight into the exterior tensor product to compute the tensor product of dilations, see 167 I. In the final sections of this chapter we turn our attention to the maps that occur on the right-hand side of the Paschke dilation. In Stinespring’s dilation these maps are always of the form  $\text{ad}_V$ , but in Paschke’s dilation different maps may appear — in any case, they are pure in the sense of 100 I. The main result of these last sections is an equivalence between this and another seemingly unrelated notion of purity: we show that an ncp-map  $\varphi$  is pure in the sense of 100 I if and only if the map on the left-hand side of the Paschke dilation of  $\varphi$  is surjective. This is a bridge to the next chapter in which we prove in greater generality that these pure maps have a  $\dagger$ -structure, see 215 III.

## 2.1 Stinespring’s theorem

In the proof of Stinespring’s theorem (137 II), it will be necessary to “complete”<sup>†</sup> a (complex) vector space with inner product into a Hilbert space. The details of this classical result were only sketched 30 I, where it’s used in the GNS-construction (30 VI). Here we will work through the details as the corresponding completion (150 II) required in Paschke’s dilation is more complex and its exposition will benefit from this familiar analog.

136

---

<sup>†</sup>Note that we do not require an inner product to be definite. The inner product on a Hilbert space *is* definite. Thus the completion will quotient out those vectors with zero norm with respect to the inner product.

- II **Proposition** Let  $V$  be a (complex) vector space with inner product  $[\cdot, \cdot]$ . There is a Hilbert space  $\mathcal{H}$  together with a bounded linear map  $\eta: V \rightarrow \mathcal{H}$  such that 1.  $[v, w] = \langle \eta(v), \eta(w) \rangle$  for all  $v, w \in V$  and 2. the image of  $\eta$  is dense in  $\mathcal{H}$ .
- III **Proof** We will form  $\mathcal{H}$  from the set of Cauchy sequences from  $V$  with a little twist. Recall that two Cauchy sequences  $(v_n)_n$  and  $(w_n)_n$  in  $V$  are said to be equivalent if for every  $\varepsilon > 0$  there is a  $n_0$  such that  $\|v_n - w_n\| \leq \varepsilon$  for all  $n \geq n_0$ , where as usual  $\|v\| \equiv \sqrt{[v, v]}$ . Call a Cauchy sequence  $(v_n)_n$  **fast** if for each  $n_0$  we have  $\|v_n - v_m\| \leq \frac{1}{2^{n_0}}$  for all  $n, m \geq n_0$ . Clearly every Cauchy sequence has a fast subsequence which is (as all subsequences are) equivalent with it.
- IV Define  $\mathcal{H}$  to be the set of fast Cauchy sequences modulo equivalence.<sup>‡</sup> For brevity we will denote an element of  $\mathcal{H}$ , which is an equivalence class of fast Cauchy sequences, simply by a single representative. Also, we often tersely write  $v$  for the Cauchy sequence  $(v_n)_n$ . The set  $\mathcal{H}$  is a metric space with the standard distance  $d(v, w) \equiv \lim_{n \rightarrow \infty} \|v_n - w_n\|$ . It is independent of representatives as by the reverse and regular triangle inequality combined, we get

$$\| \|v_n - w_n\| - \|v'_n - w'_n\| \| \leq \|v_n - v'_n\| + \|w_n - w'_n\| \rightarrow 0$$

for  $v, w$  equivalent with a Cauchy sequence  $v'$ , respectively  $w'$ . To show  $\mathcal{H}$  is a complete metric space, assume  $v^1, v^2, \dots$  is a fast Cauchy sequence of fast Cauchy sequences. We will show  $(v_n^n)_n$  is a Cauchy sequence. Here the restriction to fast sequences bears fruit: without it, the limit sequence is (notationally) harder to construct. Assume  $n, m, k \geq N$  for some  $N$ . We have

$$\|v_n^n - v_m^m\| \leq \|v_n^n - v_k^k\| + \|v_k^k - v_m^m\| + \|v_k^m - v_m^m\| \leq \|v_k^n - v_k^m\| + \frac{2}{2^N}.$$

Because  $\lim_{k \rightarrow \infty} \|v_k^n - v_k^m\| = d(v^n, v^m) \leq 2^{-N}$  we can find a  $k \geq N$  such that  $\|v_k^n - v_k^m\| \leq \frac{2}{2^N}$  and so  $\|v_n^n - v_m^m\| \leq \frac{4}{2^N}$ . Thus  $(v_n^n)_n$  is a Cauchy sequence. It's easily checked  $v^1, v^2, \dots$  converges to  $(v_n^n)_n$  with respect to  $d$ , with one caveat: the sequence  $(v_n^n)_n$  might not be fast and so, might not be in  $\mathcal{H}$ . Like any other Cauchy sequence,  $(v_n^n)_n$  has a fast subsequence. The equivalence class of this subsequence is the limit of  $v^1, v^2, \dots$  in  $\mathcal{H}$ . We have shown  $\mathcal{H}$  is a complete metric space.

- V Define  $\eta: V \rightarrow \mathcal{H}$  to be the map which sends  $v$  to the equivalence class of the constant sequence  $(v)_n$ . By construction of  $\eta$  and  $\mathcal{H}$ , the image of  $\eta$  is dense in  $\mathcal{H}$ . Let  $f: V \rightarrow X$  be any uniformly continuous map to a complete metric space  $X$ . We will show there is a unique continuous  $g: \mathcal{H} \rightarrow X$  such that  $g \circ \eta = f$ . From the uniform continuity it easily follows that  $f$  maps Cauchy sequences to Cauchy sequences and preserves equivalence between them. Together with the completeness of  $X$  there is a unique  $g: \mathcal{H} \rightarrow X$  fixed

---

<sup>‡</sup>We use fast Cauchy sequences in the definition of  $\mathcal{H}$  only to shorten the text: any Cauchy sequence is equivalent to a fast one anyway.

by  $g((v_n)_n) = \lim_{n \rightarrow \infty} f(v_n)$ . Clearly  $g \circ \eta = f$ . Finally, uniqueness of  $g$  follows readily from the fact  $g$  is fixed on the image of  $\eta$ , which is dense in  $\mathcal{H}$ .

Thus we directly get a scalar multiplication on  $\mathcal{H}$  by extending  $\eta \circ r_z: V \rightarrow \mathcal{H}$  where  $r_z(v) = zv$  is scalar multiplication by  $z \in \mathbb{C}$ , which is uniformly continuous. In fact  $z(v_n)_n \equiv (zv_n)_n$ . Extending addition is more involved, but straightforward: given representatives  $v, w \in \mathcal{H}$  we know  $(v_n + w_n)_n$  is Cauchy in  $V$  by uniform continuity of addition and so picking a fast subsequence  $v + w$  of  $(v_n + w_n)_n$  fixes an addition on  $\mathcal{H}$ . With this expression for addition, and the similar one for scalar multiplication, it is easy to see they turn  $\mathcal{H}$  into a complex vector space with zero  $\eta(0)$  for which  $\eta$  is linear. Extending the inner product is bit trickier. VI

To define the inner product on  $\mathcal{H}$ , first note that for Cauchy sequences  $v$  and  $w$  in  $V$  we have (using Cauchy–Schwarz, 32 VI, on the second line): VII

$$\begin{aligned} |[v_n, w_n] - [v_m, w_m]| &= |[v_n, w_n - w_m] + [v_n - v_m, w_m]| \\ &\leq \|v_n\| \|w_n - w_m\| + \|v_n - v_m\| \|w_m\|. \end{aligned}$$

Thus as  $(\|v_n\|)_n$  and  $(\|w_n\|)_n$  are bounded we see  $([v_n, w_n])_n$  is Cauchy. In a similar fashion we see  $|[v_n, w_n] - [v'_n, w'_n]| \leq \|v_n\| \|w_n - w'_n\| + \|v_n - v'_n\| \|w'_n\| \rightarrow 0$  as  $n \rightarrow \infty$  for  $v'$  and  $w'$  Cauchy sequences equivalent to  $v$  respectively  $w$ . Thus  $([v_n, w_n])_n$  is equivalent to  $([v'_n, w'_n])_n$  and so we define  $\langle v, w \rangle \equiv \lim_{n \rightarrow \infty} [v_n, w_n]$  on  $\mathcal{H}$ . With vector space structure from VI we easily see this is an inner product  $\mathcal{H}$ . The metric induced by the inner product coincides with  $d$ :

$$\langle v - w, v - w \rangle = \lim_n [v_n - w_n, v_n - w_n] = \lim_n \|v_n - w_n\|^2 = d(v, w)^2.$$

And so  $\mathcal{H}$  is a Hilbert space. Finally,  $\langle \eta(v), \eta(w) \rangle = [v, w]$  is direct. □

We are ready to prove Stinespring's dilation theorem. 137

**Proof of Stinespring's theorem, 135 IV** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  be any cp-map. II  
Write  $\mathcal{A} \odot \mathcal{H}$  for the tensor product of  $\mathcal{A}$  and  $\mathcal{H}$  as vector spaces. We will define  $\mathcal{K}$  as the Hilbert space completion of  $\mathcal{A} \odot \mathcal{H}$  with respect to the inner product  $[\cdot, \cdot]$  fixed by

$$[a \otimes x, b \otimes y] \equiv \langle x, \varphi(a^*b)y \rangle_{\mathcal{H}}. \quad (2.1)$$

To proceed, we have to show that  $[\cdot, \cdot]$  is indeed an inner product. By linear extension, (2.1) fixes a sesquilinear form  $[\cdot, \cdot]$ . As  $\varphi$  preserves involution as a positive map, see 10 IV, the sesquilinear form  $[\cdot, \cdot]$  is also symmetric, i.e.  $[t, s] = \overline{[s, t]}$ . Assume  $\sum_{i=1}^n a_i \otimes x_i$  is an arbitrary element of  $\mathcal{A} \odot \mathcal{H}$ . To see  $[\cdot, \cdot]$  is positive, we need to show

$$0 \leq \left[ \sum_i a_i \otimes x_i, \sum_j a_j \otimes x_j \right] \equiv \sum_{i,j} \langle x_i, \varphi(a_i^* a_j) x_j \rangle.$$

The trick is to consider the matrix algebra  $M_n\mathcal{B}(\mathcal{H})$  acting on  $\mathcal{H}^{\oplus n}$  in the obvious way:  $(A(x_1, \dots, x_n)^t)_i \equiv \sum_j A_{ij}x_j$ . (In fact, this yields an isomorphism  $M_n\mathcal{B}(\mathcal{H}) \cong \mathcal{B}(\mathcal{H}^{\oplus n})$ .) Writing  $x$  for  $(x_1, \dots, x_n)^t$  in  $\mathcal{H}^{\oplus n}$ , we get

$$\sum_{i,j} \langle x_i, \varphi(a_i^* a_j) x_j \rangle = \langle x, (M_n\varphi)((a_i^* a_j)_{ij}) x \rangle_{\mathcal{H}^{\oplus n}}. \quad (2.2)$$

The matrix  $(a_i^* a_j)_{ij}$  is positive as  $(C^*C)_{ij} = a_i^* a_j$  with  $C_{ij} \equiv \frac{1}{\sqrt{n}} a_j$ . By complete positivity  $M_n\varphi((a_i^* a_j)_{ij})$  is positive and so is (2.2), hence  $[\cdot, \cdot]$  is an inner product. Write  $\eta: \mathcal{A} \odot \mathcal{H} \rightarrow \mathcal{K}$  for the Hilbert space completion of  $\mathcal{A} \odot \mathcal{H}$  with the inner product described in 136II.

III We need a lemma before we continue. Let  $T: \mathcal{A} \odot \mathcal{H} \rightarrow \mathcal{A} \odot \mathcal{H}$  be a bounded linear map. We show there is an extension  $\hat{T}: \mathcal{K} \rightarrow \mathcal{K}$ . The operator  $T$  is uniformly continuous and so is  $\eta \circ T: \mathcal{A} \odot \mathcal{H} \rightarrow \mathcal{K}$ . Thus by 136V there is a unique continuous extension  $\hat{T}: \mathcal{K} \rightarrow \mathcal{K}$  with  $\hat{T}(\eta(t)) = \eta(T(t))$  for all  $t \in \mathcal{A} \odot \mathcal{H}$ . Clearly  $\hat{T}$  is linear on the image of  $\eta$ , which is dense and so  $\hat{T}$  is linear everywhere. As  $\hat{T}$  is linear and continuous, it must be bounded, so  $\hat{T} \in \mathcal{B}(\mathcal{K})$ . It is easy to see  $\widehat{T+S} = \hat{T} + \hat{S}$  and  $\widehat{\lambda T} = \lambda \hat{T}$  for operators  $S, T$  on  $\mathcal{A} \odot \mathcal{H}$  and  $\lambda \in \mathbb{C}$ . Also  $\widehat{TS} = \hat{T}\hat{S}$ : indeed  $\widehat{T\hat{S}}\eta(t) = \hat{T}\hat{S}\eta(t) = \hat{T}\eta(St) = \eta(TSt)$  and so by uniqueness  $\hat{T}\hat{S} = \widehat{TS}$ .

IV Assume  $b \in \mathcal{A}$ . Let  $\varrho_0(b)$  be the operator on  $\mathcal{A} \odot \mathcal{H}$  fixed by

$$\varrho_0(b) a \otimes x \equiv (ba) \otimes x.$$

Clearly  $\varrho_0$  is linear, unital and multiplicative. We want to show  $\varrho_0(b)$  is bounded for fixed  $b \in \mathcal{A}$ , so that we can define  $\varrho(b) \equiv \widehat{\varrho_0(b)} \in \mathcal{B}(\mathcal{K})$  using III. To show  $\varrho_0(b)$  is bounded, we claim that in  $M_n\mathcal{B}(\mathcal{H})$  we have

$$(a_i^* b^* b a_j)_{ij} \leq \|b\|^2 (a_i^* a_j)_{ij}. \quad (2.3)$$

Indeed, as  $b^*b \leq \|b\|^2$ , there is some  $c$  with  $c^*c = \|b\|^2 - b^*b$ . Define  $C_{ij} \equiv \frac{1}{\sqrt{n}} c a_j$ . We compute:  $(C^*C)_{ij} = a_i^* c^* c a_j = a_i^* (\|b\|^2 - b^*b) a_j$  and so (2.3) holds. Hence

$$\begin{aligned} \|\varrho_0(b) \sum_i a_i \otimes x_i\|^2 &= \|\sum_i (b a_i) \otimes x_i\|^2 \\ &= \langle x, (M_n\varphi)((a_i^* b^* b a_j)_{ij}) x \rangle \\ &\leq \|b\|^2 \langle x, (M_n\varphi)((a_i^* a_j)_{ij}) x \rangle \quad \text{by (2.3)} \\ &= \|b\|^2 \|\sum_i a_i \otimes x_i\|^2 \end{aligned}$$

and so  $\varrho_0(b)$  is bounded. Define  $\varrho(b): \mathcal{A} \rightarrow \mathcal{B}(\mathcal{K})$  using III by  $\varrho(b) \equiv \widehat{\varrho_0(b)}$ .

We already know  $\varrho$  is a mu-map. We want to show it preserves involution:  $\varrho(c^*) = \varrho(c)^*$  for all  $c \in \mathcal{A}$ . Indeed, for every  $a, b \in \mathcal{A}$  and  $x, y \in \mathcal{H}$  we have

$$[\varrho_0(c^*)a \otimes x, b \otimes y] = [(c^*a) \otimes x, b \otimes y] = \langle x, \varphi(a^*cb)y \rangle = [a \otimes x, \varrho_0(c)b \otimes y].$$

Hence  $\langle \varrho(c^*)\eta(a \otimes x), \eta(b \otimes y) \rangle = \langle \eta(a \otimes x), \varrho(c)\eta(b \otimes y) \rangle$ . As the linear span of vectors of the form  $\eta(a \otimes x)$  is dense in  $\mathcal{H}$ , we see  $\varrho(c^*)$  is adjoint to  $\varrho(c)$  and so  $\varrho(c^*) = \varrho(c)^*$ , as desired.

Let  $V_0: \mathcal{H} \rightarrow \mathcal{A} \odot \mathcal{H}$  be given by  $V_0x = 1 \otimes x$ . This operator  $V_0$  is bounded: VI

$$\|V_0x\| = \langle x, \varphi(1^*1)x \rangle^{\frac{1}{2}} = \|\sqrt{\varphi(1)}x\| \leq \|\sqrt{\varphi(1)}\| \|x\|.$$

Define  $V \equiv \eta \circ V_0$ . For all  $a \in \mathcal{A}$  and  $x, y \in \mathcal{H}$ , we have

$$[a \otimes x, V_0y] = \langle x, \varphi(a^*)y \rangle = \langle x, \varphi(a)^*y \rangle = \langle \varphi(a)x, y \rangle$$

and so  $V^*$  satisfies  $V^*\eta(a \otimes x) = \varphi(a)x$ . Hence for all  $a \in \mathcal{A}$  and  $x \in \mathcal{H}$

$$V^*\varrho(a)Vx = V^*\varrho(a)\eta(1 \otimes x) = V^*\eta(a \otimes x) = \varphi(a)x.$$

We have shown  $\text{ad}_V \circ \varrho = \varphi$ . If  $\varphi$  is unital, then  $V^*Vx = V^*(1 \otimes x) = \varphi(1)x = x$  for all  $x \in \mathcal{H}$  and so  $V$  is an isometry. (Or equivalently  $\text{ad}_V$  is unital.)

Assume  $\mathcal{A}$  is a von Neumann algebra. It remains to be shown that  $\varrho$  is normal when  $\varphi$  is. So assume  $\varphi$  is normal. To start, note that the vector functionals  $\langle a \otimes x, (\cdot)a \otimes x \rangle$  are together a faithful set of np-functionals. Thus, by 48 II, it is sufficient to show that VII

$$b \mapsto \langle a \otimes x, \varrho(b)a \otimes x \rangle \equiv \langle x, \varphi(a^*ba)x \rangle$$

is normal for every  $a \in \mathcal{A}$  and  $x \in \mathcal{H}$ . This is indeed the case, as normal maps are closed under composition and  $\langle x, (\cdot)x \rangle$  is normal by 37 IX,  $a^*(\cdot)a$  by 44 VIII and  $\varphi$  by assumption. □

**Remark** Stinespring's theorem generalizes GNS. It seems tempting to directly prove Stinespring instead of GNS — however, for our version with ncp-maps, we used GNS itself in the proof of Stinespring's theorem to show that  $\varrho$  is normal. VIII

In the following series of exercises, the reader will develop a few consequences of Stinespring's theorem which are in some fields better-known than Stinespring's theorem itself. As preparation we study the nmiu-maps between type I von Neumann algebras. The following result can be derived from the more general [110, thm. 5.5], but we will give a direct proof, which seems to be new. [130] 138

**Proposition** Let  $\varrho: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  be any non-zero<sup>§</sup> nmiu-map for Hilbert spaces  $\mathcal{H}$  and  $\mathcal{K}$ . There is a Hilbert space  $\mathcal{H}'$  and a unitary  $U: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}'$  (see 34a IV) such that  $\varrho(a) = U^*(a \otimes 1)U$  for all  $a \in \mathcal{B}(\mathcal{H})$ . II

<sup>§</sup> If  $\mathcal{H}$  is 0-dimensional, then there zero-map  $\mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  is an nmiu-map.

- III **Proof** To start, we will show that  $\varrho$  is injective by proving  $\ker \varrho = \{0\}$ . Clearly  $\ker \varrho$  is a two-sided ideal — indeed, if  $\varrho(a) = 0$  for some  $a \in \mathcal{B}(\mathcal{H})$ , then  $\varrho(ab) = \varrho(a)\varrho(b) = 0$  and  $\varrho(ba) = 0$  for any  $b \in \mathcal{B}(\mathcal{H})$ . As  $\varrho$  is normal, we have  $\varrho(\sup D) = \sup_{d \in D} \varrho(d) = 0$  for any bounded directed  $D \subseteq \ker \varrho$  of self-adjoint elements. So by 69II, we know  $\ker \varrho = z\mathcal{B}(\mathcal{H})$  for some central projection  $z \in \mathcal{B}(\mathcal{H})$ . Either  $z = 0$  or  $z = 1$ . Clearly  $z \neq 1$ , for otherwise  $0 = \varrho \neq 0$ , which is absurd. So  $z = 0$  and  $\ker \varrho = \{0\}$ , thus  $\varrho$  is injective.
- IV Pick an orthonormal basis  $(e_i)_{i \in I}$  of  $\mathcal{H}$ . Write  $p_i$  for the projection onto  $\mathbb{C}e_i$ , i.e.  $p_i \equiv |e_i\rangle\langle e_i|$ . Choose any  $i_0 \in I$  and write  $\mathcal{H}' \equiv \varrho(p_{i_0})\mathcal{H}$ . Pick any orthonormal basis  $(d_j)_{j \in J}$  of  $\mathcal{H}'$  and write  $q_j$  for the projection onto  $\mathbb{C}d_j$ , i.e.  $q_j \equiv |d_j\rangle\langle d_j|$ . Let  $u_i$  denote the unique partial isometry fixed by  $u_i e_i = e_{i_0}$  and  $u_i e_j = 0$  for  $j \neq i$  — viz.  $u_i \equiv |e_{i_0}\rangle\langle e_i|$ . Define  $r_{ij} \equiv \varrho(u_i^*)d_j$ . We will show  $(r_{ij})_{(i,j) \in I \times J}$  is an orthonormal basis of  $\mathcal{H}$ . For this, it is sufficient to show  $r_{ij}$  are unit vectors and  $\sum_{i,j} |r_{ij}\rangle\langle r_{ij}| = 1$  (where the sum is defined as supremum over the partial sums), using that then  $|r_{ij}\rangle\langle r_{ij}|$  are pairwise orthogonal projections by 55XIII. Clearly these  $r_{ij}$  are unit vectors, as by  $u_i u_i^* = p_{i_0}$  and  $\varrho(p_{i_0})d_j = d_j$ , we have  $\|r_{ij}\|^2 = \langle \varrho(u_i u_i^*)d_j, d_j \rangle = 1$ . To show  $1 = \sum_{i,j} |r_{ij}\rangle\langle r_{ij}|$ , we compute

$$\begin{aligned}
1 &= \varrho(1) && \text{as } \varrho \text{ is unital} \\
&= \varrho\left(\sum_i p_i\right) && \text{as } e_i \text{ is an orthon. basis} \\
&= \sum_i \varrho(p_i) && \text{as } \varrho \text{ is normal} \\
&= \sum_i \varrho(u_i^* p_{i_0} u_i) && \text{by defn. } u_i \\
&= \sum_i \varrho(u_i)^* \varrho(p_{i_0}) \varrho(u_i) && \text{as } \varrho \text{ is mi-map} \\
&= \sum_i \varrho(u_i)^* \left(\sum_j q_j\right) \varrho(u_i) && \text{as } d_j \text{ is an orthon. basis of } \mathcal{H}' \\
&= \sum_{i,j} \varrho(u_i)^* q_j \varrho(u_i) && \text{by 44VIII} \\
&= \sum_{i,j} |r_{ij}\rangle\langle r_{ij}|.
\end{aligned}$$

- V Let  $U: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}'$  denote the unitary fixed by  $U r_{ij} = e_i \otimes d_j$ . For any  $i \in I, j \in J$  and  $a \in \mathcal{B}(\mathcal{H})$ , it is straightforward to show that  $|ae_i\rangle\langle e_{i_0}| = \sum_k \langle e_k, ae_i \rangle |e_k\rangle\langle e_{i_0}|$ , where the sum converges ultrastrongly and consequently

$$\begin{aligned}
\varrho(a) r_{ij} &= \varrho(a)\varrho(|e_i\rangle\langle e_{i_0}|) d_j && \text{by defn. } r_{ij} \text{ and } u_i \\
&= \varrho(|ae_i\rangle\langle e_{i_0}|) d_j
\end{aligned}$$

$$\begin{aligned}
&= \varrho\left(\sum_k \langle e_k, ae_i \rangle |e_k\rangle \langle e_{i_0}| \right) d_j \\
&= \left(\sum_k \langle e_k, ae_i \rangle \varrho(|e_k\rangle \langle e_{i_0}|)\right) d_j && \text{as } \varrho \text{ is us-cont. by 44 XV} \\
&= \sum_k \langle e_k, ae_i \rangle \varrho(|e_k\rangle \langle e_{i_0}|) d_j \\
&= \sum_k \langle e_k, ae_i \rangle U^* e_k \otimes d_j && \text{by dfn. } U \text{ and } u_k \\
&= U^* \left(\sum_k e_k \langle e_k, ae_i \rangle\right) \otimes d_j \\
&= U^* (ae_i) \otimes d_j && \text{as } e_k \text{ is an orthn. basis} \\
&= U^* (a \otimes 1) U r_{ij} && \text{by dfn. } r_{ij}.
\end{aligned}$$

We have shown  $\varrho(a) = U^*(a \otimes 1)U$ . □

**Corollary** The nmiu-isomorphisms  $\mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$  are precisely of the form  $\text{ad}U$  VI  
for some unitary  $U: \mathcal{H} \rightarrow \mathcal{K}$ .

**Exercise\*** Use II and 135 IV to show that for every ncp-map  $\varphi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$  VII  
there are a Hilbert space  $\mathcal{K}'$  and a bounded operator  $V: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{K}'$  such  
that  $\varphi(a) = V^*(a \otimes 1)V$ .

Conclude that any quantum channel  $\Phi$  from  $\mathcal{H}$  to itself (i.e. completely positive trace-preserving linear map between trace-class operators over  $\mathcal{H}$ , see e.g. [112, §2.6.2]) is of the form  $\Phi(\varrho) = \text{tr}_{\mathcal{K}'}[U^* \varrho \otimes |v_0\rangle \langle v_0| U]$  for some unitary  $U$ .

**Exercise\* (Kraus' decomposition)** Let  $\varphi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$  be any ncp-map. VIII  
By VII there is a Hilbert space  $\mathcal{K}'$  and bounded operator  $V: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{K}'$   
such that  $\varphi(a) = V^*(a \otimes 1)V$ . Choose an orthonormal basis  $(e_i)_{i \in I}$  of  $\mathcal{K}'$ .  
Show  $\varphi(a) = \sum_{i \in I} V^*(a \otimes |e_i\rangle \langle e_i|)V$ , where the sum converges ultraweakly. De-  
duce there are projections  $P_i: \mathcal{H} \otimes \mathcal{K}' \rightarrow \mathcal{H}$  with  $\varphi(a) = \sum_{i \in I} V^* P_i^* a P_i V$ .  
Conclude all ncp-maps  $\varphi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$  are of the form  $\varphi(a) = \sum_{i \in I} V_i^* a V_i$   
for some bounded operators  $V_i: \mathcal{H} \rightarrow \mathcal{H}$ , for which the partial sums of  $\sum_i V_i^* V_i$   
are bounded. These  $V_i$  are called **Kraus operators** for  $\varphi$ . (For a different ap-  
proach, see e.g. [28, thm. 2.3].)

Now assume  $\mathcal{H}$  and  $\mathcal{K}$  are finite dimensional.<sup>¶</sup> Show that in this case  
the total number of Kraus operators can be chosen to be less than or equal  
to  $(\dim \mathcal{H}) \cdot (\dim \mathcal{K})$ . (There is a similar fact for infinite-dimensional  $\mathcal{H}$   
and  $\mathcal{K}$ , where the product of numbers is replaced by a maximum of cardinals.)

**Definition** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  be any ncp-map for some Hilbert space  $\mathcal{H}$  139

<sup>¶</sup>For arbitrary  $\mathcal{H}$  and  $\mathcal{K}$ , the cardinality of the set of Kraus operators can be chosen to be less than or equal to  $(\dim \mathcal{H}) \cdot (\dim \mathcal{K})$ , with dimension understood as the cardinality of an orthonormal basis and the usual product on cardinals.

and von Neumann algebra  $\mathcal{A}$ . A **normal Stinespring dilation** (cf. [94, ch. 4]) is a triple  $(\mathcal{H}, \varrho, V)$  with  $\mathcal{H}$  a Hilbert space,  $\varrho: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  an nmiu-map (i.e. normal representation) and  $V: \mathcal{H} \rightarrow \mathcal{H}$  a bounded operator such that  $\varphi$  decomposes as  $\varphi = \text{ad}_V \circ \varrho$ . We just saw every  $\varphi$  has (at least one) normal Stinespring dilation. A (normal) Stinespring dilation  $(\mathcal{H}, \varrho, V)$  is said to be **minimal** if the linear span of  $\varrho(\mathcal{A})V\mathcal{H} \equiv \{ \varrho(a)Vx; a \in \mathcal{A}, x \in \mathcal{H} \}$  is dense in  $\mathcal{H}$ . By construction the (normal) Stinespring dilation from 137 II is minimal. Every (normal) Stinespring dilation  $(\mathcal{H}, \varrho, V)$  of  $\varphi$  restricts to a minimal dilation  $(\mathcal{H}', \varrho, V)$  for  $\varphi$ , where  $\mathcal{H}' \subseteq \mathcal{H}$  is the norm-closure of the linear span of  $\varrho(\mathcal{A})V\mathcal{H}$ .

- II It is a well-known fact that all minimal normal Stinespring dilations for a fixed  $\varphi$  are unitarily equivalent, see for instance [94, prop. 4.2]. We will adapt its proof to show that a minimal normal Stinespring dilation admits a universal property. Later we will use this to prove a second universal property 140 III that allows us to generalize Stinespring's dilation to arbitrary ncp-maps. The modification of the familiar argument is mostly straightforward, except for the following lemma, which we published earlier as [127, lem. 11].

- III **Lemma** Let  $\varrho: \mathcal{A} \rightarrow \mathcal{B}$  and  $\varrho': \mathcal{A} \rightarrow \mathcal{C}$  be nmiu-maps between von Neumann algebras, and let  $\sigma: \mathcal{C} \rightarrow \mathcal{B}$  be an ncp-map such that  $\sigma \circ \varrho' = \varrho$ . Then

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{\varrho} & \mathcal{B} \\ & \searrow \varrho' & \uparrow \sigma \\ & & \mathcal{C} \end{array}$$

$$\sigma(\varrho'(a_1)c\varrho'(a_2)) = \varrho(a_1)\sigma(c)\varrho(a_2)$$

for all  $a_1, a_2 \in \mathcal{A}$  and  $c \in \mathcal{C}$ .

- IV **Proof** By Theorem 3.1 of [24] (see 34 XVIII), we know that for all  $c, d \in \mathcal{C}$ :

$$\sigma(d^*d) = \sigma(d)^*\sigma(d) \implies \sigma(cd) = \sigma(c)\sigma(d). \quad (2.4)$$

We can apply this to  $\varrho'(a)^*\varrho'(a)$  with  $a \in \mathcal{A}$  — indeed we have

$$\sigma(\varrho'(a)^*\varrho'(a)) = \sigma(\varrho'(a^*a)) = \varrho(a^*a) = \varrho(a)^*\varrho(a) = \sigma(\varrho'(a))^*\sigma(\varrho'(a))$$

and so by (2.4), we have  $\sigma(c\varrho'(a)) = \sigma(c)\sigma(\varrho'(a)) = \sigma(c)\varrho(a)$  for all  $c \in \mathcal{C}$ . Thus also (taking adjoints):  $\sigma(\varrho'(a)c) = \varrho(a)\sigma(c)$  for all  $c \in \mathcal{C}$ . Hence

$$\sigma(\varrho'(a_1)c\varrho'(a_2)) = \varrho(a_1)\sigma(c\varrho'(a_2)) = \varrho(a_1)\sigma(c)\varrho(a_2)$$

for all  $a_1, a_2 \in \mathcal{A}$  and  $c \in \mathcal{C}$  as desired.  $\square$

- V **Proposition** Assume  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  is any ncp-map with normal Stinespring dilations  $(\mathcal{H}, \varrho, V)$  and  $(\mathcal{H}', \varrho', V')$ . If  $(\mathcal{H}, \varrho, V)$  is minimal, then there is a unique isometry  $S: \mathcal{H} \rightarrow \mathcal{H}'$  such that  $SV = V'$  and  $\varrho = \text{ad}_S \circ \varrho'$ .

**Proof** (This is the same proof we gave in [127].) First we will deal with a pathological case. If  $V = 0$ , then  $\varphi = 0$ ,  $V' = 0$ ,  $\mathcal{H} = \{0\}$  and  $\varrho = 0$ . Thus the unique linear map  $S: \{0\} \rightarrow \mathcal{H}'$  satisfies the requirements. Assume  $V \neq 0$ . VI

(Uniqueness) Let  $S_1, S_2: \mathcal{H} \rightarrow \mathcal{H}'$  be any isometries for which  $S_k V = V'$  and  $\text{ad}_{S_k} \circ \varrho' = \varrho$  for  $k = 1, 2$ . We want to show  $S_1 = S_2$ . First we will prove that  $\text{ad}_{S_1} = \text{ad}_{S_2}$ . For  $k = 1, 2$ ,  $n, m \in \mathbb{N}$ ,  $a_1, \dots, a_n, \alpha_1, \dots, \alpha_m \in \mathcal{A}$ ,  $x_1, \dots, x_n, y_1, \dots, y_m \in \mathcal{H}$  and  $c \in \mathcal{B}(\mathcal{H}')$ , we have VII

$$\begin{aligned} \langle \text{ad}_{S_k}(c) \sum_i \varrho(a_i) V x_i, \sum_j \varrho(\alpha_j) V y_j \rangle &= \sum_{i,j} \langle V^* \varrho(\alpha_j^*) \text{ad}_{S_k}(c) \varrho(a_i) V x_i, y_j \rangle \\ &\stackrel{\text{III}}{=} \sum_{i,j} \langle V^* \text{ad}_{S_k}(\varrho'(\alpha_j^*) c \varrho'(a_i)) V x_i, y_j \rangle \\ &= \sum_{i,j} \langle (V')^* \varrho'(\alpha_j^*) c \varrho'(a_i) V' x_i, y_j \rangle. \end{aligned}$$

As the linear span of  $\varrho(\mathcal{A})V\mathcal{H}$  is dense in  $\mathcal{H}$ , we see  $\text{ad}_{S_1}(c) = \text{ad}_{S_2}(c)$  for all  $c \in \mathcal{B}(\mathcal{H}')$ . So  $\text{ad}_{S_1} = \text{ad}_{S_2}$  and thus  $\lambda S_1 = S_2$  for some  $\lambda \in \mathbb{C}$ , cf. [126, lem. 9]. As  $V \neq 0$ , there is an  $x \in \mathcal{H}$  with  $Vx \neq 0$ . Also  $S_1 Vx \neq 0$  as  $S_1$  is an isometry and thus injective. From this and  $S_1 Vx = V'x = S_2 Vx = \lambda S_1 Vx$ , we get  $\lambda = 1$ . Hence  $S_1 = S_2$ .

(Existence) For any  $n \in \mathbb{N}$ ,  $a_1, \dots, a_n \in \mathcal{A}$  and  $x_1, \dots, x_n \in \mathcal{H}$ , we have VIII

$$\begin{aligned} \left\| \sum_i \varrho(a_i) V x_i \right\|^2 &= \sum_{i,j} \langle V^* \varrho(a_j^* a_i) V x_i, x_j \rangle \\ &= \sum_{i,j} \langle \varphi(a_j^* a_i) x_i, x_j \rangle \\ &= \left\| \sum_i \varrho'(a_i) V' x_i \right\|^2. \end{aligned}$$

It follows (again using denseness of the linear span of  $\varrho(\mathcal{A})V\mathcal{H}$ ), that there is a unique isometry  $S: \mathcal{H} \rightarrow \mathcal{H}'$  such that  $S\varrho(a)Vx = \varrho'(a)V'x$  for all  $a \in \mathcal{A}$  and  $x \in \mathcal{H}$ . As  $SVx = S\varrho(1)Vx = \varrho'(1)V'x = V'x$  for all  $x \in \mathcal{H}$ , we have  $SV = V'$ . Furthermore we have

$$\begin{aligned} S\varrho(a) \sum_i \varrho(a_i) V x_i &= \sum_i S\varrho(aa_i) V x_i \\ &= \sum_i \varrho'(aa_i) V' x_i \\ &= \varrho'(a) \sum_i \varrho'(a_i) V' x_i \\ &= \varrho'(a) S \sum_i \varrho(a_i) V x_i. \end{aligned}$$

for all  $n \in \mathbb{N}$ ,  $a, a_1, \dots, a_n \in \mathcal{A}$  and  $x_1, \dots, x_n \in \mathcal{H}$ . So  $S\varrho(a) = \varrho'(a)S$ . Thus  $S^* \varrho'(a)S = S^* S \varrho(a) = \varrho(a)$  for all  $a \in \mathcal{A}$ , whence  $\text{ad}_S \circ \varrho' = \varrho$ .  $\square$

- IX Exercise\*** The statement of the universal property **V** for Stinespring was proposed by Chris Heunen in an unpublished note. In this note he shows it is equivalent to the existence of a left-adjoint, as we will see in this exercise.

Let  $\text{Rep}_{\text{cp}}$  denote the category with as objects ncpu-maps  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  and as arrows between  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  and  $\varphi': \mathcal{A}' \rightarrow \mathcal{B}(\mathcal{H}')$  pairs  $(m, S)$  where  $m: \mathcal{A} \rightarrow \mathcal{A}'$  is an nmiu-map and  $S: \mathcal{H} \rightarrow \mathcal{H}'$  is an isometry such that  $\varphi = \text{ad}_S \circ \varphi' \circ m$ . Write  $\text{Rep}$  for the full subcategory of  $\text{Rep}_{\text{cp}}$  consisting of objects  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  that are nmiu.

Show using **V** that the inclusion functor  $U: \text{Rep} \rightarrow \text{Rep}_{\text{cp}}$  has a left adjoint.

- IXa Remarks** Recently there have been two interesting categorical discoveries concerning Stinespring's dilation theorem, which deserve a mention. Parzygnat found a rather different characterization of Stinespring's dilation theorem as an adjunction [92]. Secondly, Huot and Staton discovered that the category of completely positive maps between matrix algebras is the universal monoidal category with terminal unit with a functor from the category of matrix algebras with isometries between them. [54] This fact is surprisingly closely related to Stinespring's dilation theorem for matrix algebras.


- X** In the special case of ncp-maps of the form  $\varphi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$ , there is a different and noteworthy property concerning dilations 'of the same dimension' called *essential uniqueness of purification*. This property is the cornerstone of several publications, notably [18], which popularized it.

- XI Exercise\* (Essential uniqueness of purification)** Let  $\varphi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$  be any ncp-map. By 138 VII here is a Hilbert space  $\mathcal{K}'$  and bounded operator  $V: \mathcal{K} \rightarrow \mathcal{H} \otimes \mathcal{K}'$  such that  $V^*(a \otimes 1)V = \varphi(a)$ . Assume there is a bounded operator  $W: \mathcal{K} \rightarrow \mathcal{H} \otimes \mathcal{K}'$  with  $W^*(a \otimes 1)W = \varphi(a)$  as well. We will show there is a unitary  $U: \mathcal{K}' \rightarrow \mathcal{K}'$  such that  $V = (1 \otimes U)W$ .

First show that in the special case the dilations are minimal (i.e. the linear span of  $\{(a \otimes 1)Vx; a \in \mathcal{B}(\mathcal{H}), x \in \mathcal{K}\}$  is dense in  $\mathcal{H} \otimes \mathcal{K}'$ ), there is a (unique) unitary  $U_0$  on  $\mathcal{H} \otimes \mathcal{K}'$  with  $V = U_0W$  and  $(a \otimes 1)U_0 = U_0(a \otimes 1)$  for all  $a \in \mathcal{B}(\mathcal{H})$ . Derive from the latter that for each  $y \in \mathcal{K}'$  and unit-vector  $e \in \mathcal{H}$ , there is a  $y' \in \mathcal{K}'$  with  $U_0(e \otimes y) = e \otimes y'$ . Conclude  $U_0 = 1 \otimes U$  for some unitary  $U$  on  $\mathcal{K}'$ . Show that the general case follows from the special case in which the dilations are minimal.

- 140** We are ready to state the more general universal property that Stinespring's dilation admits, which will allow us to generalize it to arbitrary ncp-maps.

**Definition** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  be any ncp-map between von Neumann algebras. A **Paschke dilation** of  $\varphi$  is a triple  $(\mathcal{P}, \varrho, h)$  with a von Neumann algebra  $\mathcal{P}$ ; an nmiu-map  $\varrho: \mathcal{A} \rightarrow \mathcal{P}$  and ncp-map  $h: \mathcal{P} \rightarrow \mathcal{B}$  with  $\varphi = h \circ \varrho$  and the following universal property holds.




For every triple  $(\mathcal{P}', \varrho', h')$  with a von Neumann algebra  $\mathcal{P}'$ ; an nmiu-map  $\varrho': \mathcal{A} \rightarrow \mathcal{P}'$  and ncp-map  $h': \mathcal{P}' \rightarrow \mathcal{B}$  with  $\varphi = h' \circ \varrho'$ , there is a unique ncp-map  $\sigma: \mathcal{P}' \rightarrow \mathcal{P}$  (the “mediating map”) with  $\sigma \circ \varrho' = \varrho$  and  $h \circ \sigma = h'$ .

**Theorem** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  be any ncp-map together with a minimal normal Stinespring dilation  $(\mathcal{K}, \varrho, V)$ . Then  $(\mathcal{B}(\mathcal{K}), \varrho, \text{ad}_V)$  is a Paschke dilation of  $\varphi$ .

**Proof** (This is essentially the same proof we gave in [127].)

Let  $\mathcal{P}'$  be any von Neumann algebra together with nmiu-map  $\varrho': \mathcal{A} \rightarrow \mathcal{P}'$  and ncp-map  $h': \mathcal{P}' \rightarrow \mathcal{B}(\mathcal{H})$  for which it holds that  $h' \circ \varrho' = \varphi$ . We must show that there is a unique ncp-map  $\sigma: \mathcal{P}' \rightarrow \mathcal{B}(\mathcal{K})$  such that  $\sigma \circ \varrho' = \varrho$  and  $\text{ad}_V \circ \sigma = h'$ . First we will prove uniqueness of this map  $\sigma$ . Then we will show that such a  $\sigma$  exists by appealing to the universal property 139V.



(Uniqueness) Let  $\sigma_1, \sigma_2: \mathcal{P}' \rightarrow \mathcal{B}(\mathcal{K})$  be two such ncp-maps. Similarly to the uniqueness proof in 139V, we compute that for all  $n, m \in \mathbb{N}$ ,  $a_1, \dots, a_n, \alpha_1, \dots, \alpha_m \in \mathcal{A}$ ,  $x_1, \dots, x_n, y_1, \dots, y_m \in \mathcal{H}$  and  $c \in \mathcal{P}'$ , we have

$$\begin{aligned} \langle \sigma_k(c) \sum_i \varrho(a_i) V x_i, \sum_j \varrho(a_j) V x_j \rangle &= \sum_{i,j} \langle V^* \varrho(\alpha_j^*) \sigma_k(c) \varrho(a_i) V x_i, y_j \rangle \\ &\stackrel{139\text{III}}{=} \sum_{i,j} \langle V^* \sigma_k(\varrho'(\alpha_j^*) c \varrho'(a_i)) V x_i, y_j \rangle \\ &= \sum_{i,j} \langle h'(\varrho'(\alpha_j^*) c \varrho'(a_i)) x_i, y_j \rangle. \end{aligned}$$

By definition of minimality of a Stinespring dilation, the set of vectors of the form  $\sum_i \varrho(a_i) V x_i$  is norm dense in  $\mathcal{K}$  and so  $\sigma_1(c) = \sigma_2(c)$  as desired.

(Existence) Let  $(\tilde{\mathcal{K}}, \tilde{\varrho}, \tilde{V})$  be a minimal normal Stinespring dilation of  $h'$ . The triple  $(\tilde{\mathcal{K}}, \tilde{\varrho} \circ \varrho', \tilde{V})$  is a normal Stinespring dilation of  $\varphi$ . Thus by 139V there is a (unique) isometry  $S: \mathcal{K} \rightarrow \tilde{\mathcal{K}}$  with  $SV = \tilde{V}$  and  $\text{ad}_S \circ \tilde{\varrho} \circ \varrho' = \varrho$ . Define  $\sigma = \text{ad}_S \circ \tilde{\varrho}$ . Clearly  $\sigma \circ \varrho' = \text{ad}_S \circ \tilde{\varrho} \circ \varrho' = \varrho$  and  $\text{ad}_V \circ \sigma = \text{ad}_V \circ \text{ad}_S \circ \tilde{\varrho} = \text{ad}_{SV} \circ \tilde{\varrho} = \text{ad}_{\tilde{V}} \circ \tilde{\varrho} = h'$  as desired.  $\square$

VII The converse holds as well: by VIII every Paschke dilation of a map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  is up to a unique nmiu-isomorphism a minimal Stinespring dilation. Later we will show that every ncp-map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  has a Paschke dilation: it turns out that Paschke's generalization of GNS to Hilbert  $C^*$ -modules fits the bill. Before we show this, we look at some other examples and basic properties of Paschke dilations. First, as Paschke dilations are defined by a simple universal property, they are unique up to a unique nmiu-isomorphism:

VIII **Lemma** If  $(\mathcal{P}_1, \varrho_1, h_1)$  and  $(\mathcal{P}_2, \varrho_2, h_2)$  are two Paschke dilations for the same ncp-map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$ , then there is a unique (nmiu) isomorphism  $\vartheta: \mathcal{P}_1 \rightarrow \mathcal{P}_2$  with  $\vartheta \circ \varrho_1 = \varrho_2$  and  $h_2 \circ \vartheta = h_1$ .

IX **Proof** There are unique mediating maps  $\sigma_1: \mathcal{P}_1 \rightarrow \mathcal{P}_2$  and  $\sigma_2: \mathcal{P}_2 \rightarrow \mathcal{P}_1$ . It is easy to see  $\sigma_1 \circ \sigma_2$  satisfies the same defining property as the unique mediating map  $\text{id}: \mathcal{P}_2 \rightarrow \mathcal{P}_2$  and so  $\sigma_1 \circ \sigma_2 = \text{id}$ . Similarly  $\sigma_2 \circ \sigma_1 = \text{id}$ . Define  $\vartheta = \sigma_1$ . We just saw  $\vartheta$  is an ncp-isomorphism. Note  $\vartheta(1) = \vartheta(\varrho_1(1)) = \varrho_2(1) = 1$  and so  $\vartheta$  is unital. But then  $\vartheta$  is an nmiu-isomorphism by 99IX.  $\square$

X **Exercise** The following can be shown using only the defining universal property.

1. Show  $(\mathcal{A}, \varrho, \text{id})$  is a Paschke dilation of an nmiu-map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}$ .
2. Let  $(\mathcal{P}, \varrho, h)$  be any Paschke dilation. Prove  $(\mathcal{P}, \text{id}, h)$  is a Paschke dilation for  $h$ . (Hint: use  $\text{id}: \mathcal{P} \rightarrow \mathcal{P}$  is the unique ncp-map with  $\text{id} \circ \varrho = \varrho$  and  $h \circ \text{id} = h$ .)
3. Let  $(\varphi_2^1): \mathcal{A} \rightarrow \mathcal{B}_1 \oplus \mathcal{B}_2$  be an ncp-map. Show  $(\mathcal{P}_1 \oplus \mathcal{P}_2, (\varrho_2^1), h_1 \oplus h_2)$  is a Paschke dilation of  $(\varphi_2^1)$  if  $(\mathcal{P}_i, \varrho_i, h_i)$  is a Paschke dilation of  $\varphi_i$  for  $i = 1, 2$ .
4. Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  be any ncp-map with Paschke dilation  $(\mathcal{P}, \varrho, h)$  and  $\lambda \in \mathbb{R}$  with  $\lambda > 0$ . Prove  $(\mathcal{P}, \varrho, \lambda h)$  is a Paschke dilation of  $\lambda \varphi$ .

XI **Examples** Later we will be able to compute more dilations.

1. In 167I, we will see that if  $(\mathcal{P}_i, \varrho_i, h_i)$  is a Paschke dilation of an ncp-map  $\varphi_i: \mathcal{A}_i \rightarrow \mathcal{B}_i$  for  $i = 1, 2$ , then  $(\mathcal{P}_1 \otimes \mathcal{P}_2, \varrho_1 \otimes \varrho_2, h_1 \otimes h_2)$  is a Paschke dilation of  $\varphi_1 \otimes \varphi_2$ .
2. In 171II, we will show that  $(\llbracket p \rrbracket \mathcal{A}, h_{\llbracket p \rrbracket}, h_p)$  is a Paschke dilation of the ncp-map  $h: \mathcal{A} \rightarrow p\mathcal{A}p$  given by  $h(a) = pap$  for some projection  $p \in \mathcal{A}$ .
3. In 171VII, it's proven that a map  $\varphi$  with Paschke dilation  $(\mathcal{P}, \varrho, h)$  is pure (in the sense of 100I) if and only if  $\varrho$  is surjective.
4. In 169XI the Paschke dilations of  $\varphi$  and  $c \circ \varphi$  for a filter  $c$  are related.

## 2.2 Hilbert $C^*$ -modules

Recall that by GNS every state (pu-map)  $\omega: \mathcal{A} \rightarrow \mathbb{C}$  on a von Neumann algebra  $\mathcal{A}$  splits as  $h \circ \varrho$  for some nmiu-map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ , Hilbert space  $\mathcal{H}$  and vector state  $h(T) = \langle x, Tx \rangle$  for some  $x \in \mathcal{H}$ . It is easy to see  $(\mathcal{H}, \varrho, h)$  is a Paschke dilation of  $\omega$ . 141

Paschke showed that GNS can be generalized by replacing the scalars  $\mathbb{C}$  with an arbitrary von Neumann algebra  $\mathcal{B}$ : every ‘state’ ncp-map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  splits as  $h \circ \varrho$  for some nmiu-map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}^a(X)$ , and ‘vector state’  $h(T) = \langle x, Tx \rangle$ , where  $X$  is a self-dual Hilbert  $C^*$ -module over  $\mathcal{B}$ ,  $x \in X$  and  $\mathcal{B}^a(X)$  are the  $\mathcal{B}$ -linear bounded operators on  $X$ . It turns out  $(\mathcal{B}^a(X), \varrho, h)$  is a Paschke dilation of  $\varphi$ . Before we will prove this, we first develop the theory of (self-dual) Hilbert  $C^*$ -modules.

We already encountered Hilbert  $C^*$ -modules in the development of the basics of completely positive maps in my twin brother’s thesis [124]. (They appear in 321.) As Hilbert  $C^*$ -modules are central to the present topic, we will prove some basic results again, which were already covered by my twin.

**Definition** Let  $\mathcal{B}$  be a  $C^*$ -algebra and  $X$  be a right  $\mathcal{B}$ -module. A  $\mathcal{B}$ -valued inner product on  $X$  is a map  $\langle \cdot, \cdot \rangle: X \times X \rightarrow \mathcal{B}$  such that II

1.  $\langle x, \cdot \rangle: X \rightarrow \mathcal{B}$  is  $\mathcal{B}$ -linear for all  $x \in X$  — viz.  $\langle x, yb \rangle = \langle x, y \rangle b$ ;
2.  $\langle x, y \rangle^* = \langle y, x \rangle$  for all  $x, y \in X$  and
3.  $\langle x, x \rangle \geq 0$  for all  $x \in X$ .

A  $\mathcal{B}$ -valued inner product is called **definite** provided  $x = 0$  whenever  $\langle x, x \rangle = 0$ . A **pre-Hilbert  $\mathcal{B}$ -module**  $X$  is a right  $\mathcal{B}$ -module  $X$  together with a definite  $\mathcal{B}$ -valued inner product.

It will turn out  $\|x\| = \|\langle x, x \rangle\|^{\frac{1}{2}}$  defines a norm on a pre-Hilbert  $\mathcal{B}$ -module  $X$ . A **Hilbert  $\mathcal{B}$ -module** (also called a **Hilbert  $C^*$ -module over  $\mathcal{B}$** ) is a pre-Hilbert  $\mathcal{B}$ -module which is complete in this norm.

A pre-Hilbert  $\mathcal{B}$ -module is called **self dual** if for each bounded  $\mathcal{B}$ -linear  $\tau: X \rightarrow \mathcal{B}$ , there is a  $t \in X$  such that  $\tau(x) = \langle t, x \rangle$  for all  $x \in X$ . IIa

**Beware** In his original definition [93, §2.1], Paschke chose to make his  $\mathcal{B}$ -valued inner product  $\mathcal{B}$ -linear on the left instead of on the right. He also assumes his inner products to be definite. IIb

**Examples** Several examples [93] of (pre-)Hilbert  $C^*$ -modules follow, to each of which we will return in the remainder. III

- Hilbert  $C$ -modules are the same as Hilbert spaces. They are all self dual.

- Any  $C^*$ -algebra  $\mathcal{B}$  is a self-dual Hilbert  $C^*$ -module over itself with  $\mathcal{B}$ -valued inner product  $\langle a, b \rangle = a^*b$ . By the  $C^*$ -identity, the Hilbert  $\mathcal{B}$ -module norm on  $\mathcal{B}$  coincides with the norm of the  $C^*$ -algebra.
- More generally: if  $J \subseteq \mathcal{B}$  is any right ideal of  $\mathcal{B}$ , then  $J$  is a pre-Hilbert  $\mathcal{B}$ -module with  $\langle a, b \rangle = a^*b$ . If  $J$  is closed with respect to the norm, then  $J$  is a Hilbert  $\mathcal{B}$ -module. It might not be self dual.
- Principal right-ideals are self dual: the right ideal  $e\mathcal{B}$  for a projection  $e \in \mathcal{B}$  is a self-dual Hilbert  $\mathcal{B}$ -module with  $\langle ea, eb \rangle = a^*eb$ . Later we will see that every self-dual Hilbert  $\mathcal{B}$ -module over a von Neumann algebra  $\mathcal{B}$  is built up from such  $e\mathcal{B}$ .
- If  $X$  and  $Y$  are (pre-)Hilbert  $\mathcal{B}$ -modules, then their direct sum  $X \oplus Y$  (as right  $\mathcal{B}$ -modules) is a (pre-)Hilbert  $\mathcal{B}$ -module with the familiar inner product  $\langle (x, y), (x', y') \rangle = \langle x, x' \rangle + \langle y, y' \rangle$ . If  $X$  and  $Y$  are self dual, then so is  $X \oplus Y$ .

## 2.2.1 The basics of Hilbert $C^*$ -modules

**142** Perhaps the most important stones in the foundation of the theory of operator algebras are the Cauchy–Schwarz inequality  $|\langle x, y \rangle|^2 \leq \|x\|^2 \|y\|^2$  and the polarization identity  $\langle x, y \rangle = \frac{1}{4} \sum_{k=0}^3 i^k \|i^k x + y\|^2$ , see 4XV. Also in the development of Hilbert  $C^*$ -modules, their generalizations support the bulk of the structure.

**II Definition** Let  $X$  be a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product  $\langle \cdot, \cdot \rangle$  for some  $C^*$ -algebra  $\mathcal{B}$ . For any positive functional  $f: \mathcal{B} \rightarrow \mathbb{C}$ , write  $\langle x, y \rangle_f = f(\langle x, y \rangle)$  and  $\|x\|_f = \langle x, x \rangle_f^{1/2}$ . This  $\langle \cdot, \cdot \rangle_f$  is a complex-valued inner product on  $X$ . Hence, by 4XV, we know that  $\|\cdot\|_f$  is a seminorm (i.e. a not-necessarily-positive-definite norm.)

**IIa** We start with the Cauchy–Schwarz inequality for  $\mathcal{B}$ -valued inner products. The proof is the same as Paschke gives in [93, prop. 2.3 (ii)], but we use it to prove a slightly more general result.

**III Proposition (Cauchy–Schwarz)** Let  $X$  be a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product  $\langle \cdot, \cdot \rangle$  for some  $C^*$ -algebra  $\mathcal{B}$ . Then  $\langle x, y \rangle \langle y, x \rangle \leq \| \langle y, y \rangle \| \langle x, x \rangle$ .

**IV Proof** Let  $f: \mathcal{B} \rightarrow \mathbb{C}$  be any state (pu-map). Since the states on  $\mathcal{B}$  are order separating, see 22VIII, it suffices to show  $f(\langle x, y \rangle \langle y, x \rangle) \leq \| \langle y, y \rangle \| f(\langle x, x \rangle)$ . If  $f(\langle x, y \rangle \langle y, x \rangle) = 0$  the inequality holds trivially, so assume  $f(\langle x, y \rangle \langle y, x \rangle) \neq 0$ .

Using Cauchy–Schwarz for  $\langle \cdot, \cdot \rangle_f$ , we derive

$$\begin{aligned} f(\langle x, y \rangle \langle y, x \rangle)^2 &= \langle x, y \langle y, x \rangle \rangle_f^2 \\ &\leq \|x\|_f^2 \|y \langle y, x \rangle\|_f^2 \\ &= \|x\|_f^2 f(\langle x, y \rangle \langle y, y \rangle \langle y, x \rangle) \\ &\leq \|x\|_f^2 \| \langle y, y \rangle \| f(\langle x, y \rangle \langle y, x \rangle), \end{aligned}$$

which yields the inequality by dividing by  $f(\langle x, y \rangle \langle y, x \rangle)$ .  $\square$

**Exercise** Let  $X$  be a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product  $\langle \cdot, \cdot \rangle$  for some  $C^*$ -algebra  $\mathcal{B}$ . First use Cauchy–Schwarz (III) to show  $\| \langle x, y \rangle \| \leq \|x\| \|y\|$ . With this, derive  $\|x\| = \| \langle x, x \rangle \|^{1/2}$  is a seminorm on  $X$  with  $\|x \cdot b\| \leq \|x\| \|b\|$  for all  $b \in \mathcal{B}$ . (See [93, prop. 2.3].) V

From this, it follows easily that that the addition and the module action are jointly uniformly continuous and that the inner product is uniformly continuous in each argument separately. (The inner product is also jointly continuous, but not uniformly.) VI

**Definition** Let  $V$  be a right  $\mathcal{B}$ -module for some  $C^*$ -algebra  $\mathcal{B}$ . A  $\mathcal{B}$ -sesquilinear form on  $V$  is a map  $B: V \times V \rightarrow \mathcal{B}$  such that for each  $y \in V$  the maps  $x \mapsto B(y, x)$  and  $x \mapsto B(x, y)^*$  are  $\mathcal{B}$ -linear — that is: VII

$$\beta^* \langle x, y \rangle b = \langle x\beta, yb \rangle \quad \text{for all } x, y \in X \text{ and } b, \beta \in \mathcal{B}.$$

**Remark** It seems that  $\mathcal{B}$ -sesquilinear forms do not explicitly appear in the established literature, even though its associated polarization identity is often used implicitly. VIIa

**Example** Let  $X$  be a pre-Hilbert  $\mathcal{B}$ -module. For every  $\mathcal{B}$ -linear  $T: X \rightarrow X$  the map  $\langle (\cdot), T(\cdot) \rangle$  is a  $\mathcal{B}$ -sesquilinear form. In 152V we will see that on a self-dual Hilbert  $C^*$ -module all bounded  $\mathcal{B}$ -sesquilinear forms are of this form. VIII

**Exercise** Let  $B$  be a  $\mathcal{B}$ -sesquilinear form. Prove the polarization identity: IX

$$B(x, y) = \frac{1}{4} \sum_{k=0}^3 i^k B(i^k x + y, i^k x + y).$$

**Definition** Let  $X, Y$  be pre-Hilbert  $\mathcal{B}$ -modules for some  $C^*$ -algebra  $\mathcal{B}$ . A  $(C-)$ linear map  $T: X \rightarrow Y$  is said to be adjointable if there is a linear map  $S: Y \rightarrow X$  such that  $\langle y, Tx \rangle_Y = \langle Sy, x \rangle_X$  for all  $x \in X$  and  $y \in Y$ . Adjoints are unique: if  $S$  and  $S'$  are both adjoints of  $T$  an easy computation shows  $\langle (S - S')y, (S - S')'y \rangle = 0$  for all  $y \in Y$ . We write  $T^*$  for the adjoint of  $T$  (if it exists) and  $\mathcal{B}^a(X)$  for the subset of (norm-)bounded operators on  $X$  which are adjointable. (See [93, §2].) 143

- la The second part of the following lemma is well-known, but is proven in a new way using the equivalence which is inspired by [93, thm. 2.8 (ii)].
- ll **Lemma** For a linear map  $T: X \rightarrow Y$  between pre-Hilbert  $\mathcal{B}$ -modules  $X, Y$  and real number  $B > 0$ , the following are equivalent.
1.  $\|Tx\| \leq B\|x\|$  for all  $x \in X$ .
  2.  $\|\langle y, Tx \rangle\| \leq B\|y\|\|x\|$  for all  $x \in X$  and  $y \in Y$ .

If  $T$  is bounded and adjointable, then  $\|T^*\| = \|T\|$  and  $\|T^*T\| = \|T\|^2$ .

- liii **Proof** Assume  $\|Tx\| \leq B\|x\|$  for all  $x \in X$ . Then by 142V we find that for all  $y \in X$  we have  $\|\langle y, Tx \rangle\| \leq \|y\|\|Tx\| \leq B\|y\|\|x\|$ , as desired.

For the converse, pick  $x \in X$  and assume  $\|\langle y, Tx \rangle\| \leq B\|y\|\|x\|$  for all  $y \in Y$ . Then  $\|Tx\|^2 = \|\langle Tx, Tx \rangle\| \leq B\|Tx\|\|x\|$ . So  $\|Tx\| \leq B\|x\|$  by dividing  $\|Tx\|$  if  $\|Tx\| \neq 0$  and trivially otherwise.

Now assume  $T$  is adjointable and bounded. Then  $\|\langle x, T^*y \rangle\| = \|\langle y, Tx \rangle\| \leq \|T\|\|y\|\|x\|$  for all  $x \in X, y \in Y$  and so by the previous  $\|T^*\| \leq \|T\|$ . As adjoints are unique and  $T$  is an adjoint of  $T^*$  we get  $T^{**} = T$  and so  $\|T\| = \|T^{**}\| \leq \|T^*\| \leq \|T\|$ , as desired.

For the final identity, note that for any  $x \in X$  we have

$$\|Tx\|^2 = \|\langle Tx, Tx \rangle\| = \|\langle x, T^*Tx \rangle\| \leq \|x\|\|T^*Tx\| \leq \|x\|^2\|T^*T\|$$

and so  $\|T\| \leq \|T^*T\|^{\frac{1}{2}}$ . Hence  $\|T\|^2 \leq \|T^*T\| \leq \|T^*\|\|T\| = \|T\|^2$ .  $\square$

- lV **Proposition** Let  $X$  be a Hilbert  $\mathcal{B}$ -module for some  $C^*$ -algebra  $\mathcal{B}$ . With composition as multiplication, adjoint as involution and operator norm, the set of adjointable operators  $\mathcal{B}^a(X)$  is a  $C^*$ -algebra. [93, §2]

- lV **Proof** It is easy to see  $T^* + S^*$  is an adjoint of  $T + S$  for  $T, S \in \mathcal{B}^a(X)$  and so  $T + S$  is adjointable with  $(T + S)^* = T^* + S^*$ . Similarly it is easy to see  $\mathcal{B}^a(X)$  is closed under scalar multiplication, composition and involution with  $(\lambda T)^* = \bar{\lambda}T^*, (TS)^* = S^*T^*$  and  $T^{**} = T$ . Also the identity 1 is clearly bounded and self-adjoint. So  $\mathcal{B}^a(X)$  is a unital  $*$ -algebra. By ll the  $C^*$ -identity holds. It only remains to be shown that  $\mathcal{B}^a(X)$  is complete. Let  $T_1, T_2, \dots$  be a Cauchy sequence in  $\mathcal{B}^a(X)$ . As  $X$  is a Banach space  $\mathcal{B}(X)$  is complete and so  $T_n \rightarrow T$  for some  $T \in \mathcal{B}(X)$ . We have to show  $T$  is adjointable. For any  $n, m \in \mathbb{N}$  we have  $\|T_n^* - T_m^*\| = \|(T_n - T_m)^*\| = \|T_n - T_m\|$  and so  $T_1^*, T_2^*, \dots$  is also Cauchy and converges to, say,  $S \in \mathcal{B}(X)$ . So

$$\langle Sx, y \rangle = \langle \lim_n T_n^*x, y \rangle = \lim_n \langle T_n^*x, y \rangle = \lim_n \langle x, T_n y \rangle = \langle x, Ty \rangle,$$

for any  $x \in X$  and  $y \in Y$ . Thus  $S = T^*$  and  $T \in \mathcal{B}^a(X)$  as desired.  $\square$

- 144 **Proposition** Let  $X$  be a Hilbert  $\mathcal{B}$ -module for some  $C^*$ -algebra  $\mathcal{B}$ . The vector states on  $\mathcal{B}^a(X)$  are order separating — that is:  $T \geq 0$  if and only if  $\langle x, Tx \rangle \geq 0$  for all  $x \in X$ .

**Proof** (Proof from [81, lem. 4.1].) Assume  $T \in \mathcal{B}^a(X)$  with  $T \geq 0$ . Then  $T = S^*S$  for some  $S \in \mathcal{B}^a(X)$  and so  $\langle x, Tx \rangle = \langle Sx, Sx \rangle \geq 0$  for all  $x \in X$ . ||

For the converse, assume  $\langle x, Tx \rangle \geq 0$  for all  $x \in X$ . We claim  $T$  is self-adjoint. By the polarization identity  $\langle x, Ty \rangle = \frac{1}{4} \sum_{k=0}^3 i^k \langle i^k x + y, T(i^k x + y) \rangle$ , see 142IX, we have  $\langle x, Ty \rangle = \langle Tx, y \rangle$  for all  $x, y \in X$  and so  $T^* = T$ . There are positive  $T_+, T_- \in \mathcal{B}^a(X)$  with  $T = T_+ - T_-$  and  $T_+ T_- = 0$ , see 24II. By assumption  $0 \leq \langle T_- x, T T_- x \rangle$  from which  $-\langle x, T_-^3 x \rangle \geq 0$ , but  $T_-^3 \geq 0$  so  $T_-^3 = 0$ . Thus  $T_- = 0$  (by the functional calculus, 28II) and so  $T \geq 0$ . □

**Lemma** Let  $T: X \rightarrow Y$  be an adjointable linear map between pre-Hilbert  $\mathcal{B}$ -modules  $X, Y$ . Then  $T$  is  $\mathcal{B}$ -linear. [93, §2] |||

**Proof** We have  $\langle y, (Tx)b \rangle = \langle T^*y, x \rangle b = \langle y, T(xb) \rangle$  for any  $x \in X, y \in Y$  and  $b \in \mathcal{B}$ . In particular we get  $\langle (Tx)b - T(xb), (Tx)b - T(xb) \rangle = 0$  taking  $y = (Tx)b - T(xb)$  and so  $T(x)b = T(xb)$ . □ IV

**Proposition** Let  $X$  and  $Y$  be right  $\mathcal{B}$ -modules with  $\mathcal{B}$ -valued inner products for some  $C^*$ -algebra  $\mathcal{B}$ . If  $T: X \rightarrow Y$  is a bounded  $\mathcal{B}$ -linear map, then we have the inequality  $\langle Tx, Tx \rangle \leq \|T\|^2 \langle x, x \rangle$  for any  $x \in X$ . (cf. [93, rem. 2.9].) V

**Proof** (The proof is a slight variation on the first part of [93, thm. 2.8].) Pick  $\varepsilon > 0$ . By 17V we know  $\langle x, x \rangle + \varepsilon$  is invertible. Define  $h = (\langle x, x \rangle + \varepsilon)^{-\frac{1}{2}}$ . Clearly VI

$$0 \leq \langle xh, xh \rangle = h(\langle x, x \rangle + \varepsilon)h - h^2\varepsilon = 1 - h^2\varepsilon \leq 1.$$

Thus  $\|xh\| \leq 1$  and so  $\|h \langle Tx, Tx \rangle h\| = \|T(xh)\|^2 \leq \|T\|^2$ . From this and  $0 \leq h \langle Tx, Tx \rangle h$  we get  $h \langle Tx, Tx \rangle h \leq \|T\|^2$ . Hence, by dividing by  $h$  on both sides,

$$\langle Tx, Tx \rangle \leq \|T\| h^{-2} = \|T\|^2 (\langle x, x \rangle + \varepsilon)$$

for all  $\varepsilon > 0$  and so  $\langle Tx, Tx \rangle \leq \|T\|^2 \langle x, x \rangle$  as desired. □

**Proposition** Let  $\mathcal{B}$  be a von Neumann algebra,  $X$  be a Hilbert  $\mathcal{B}$ -module and  $x \in X$ . Then the map  $h: \mathcal{B}^a(X) \rightarrow \mathcal{B}$  defined by  $h(T) = \langle x, Tx \rangle$  is completely positive. 145

**Proof** Pick any  $n \in \mathbb{N}$ ,  $T_1, \dots, T_n \in \mathcal{B}^a(X)$  and  $b_1, \dots, b_n \in \mathcal{B}$  and compute ||

$$\begin{aligned} \sum_{i,j} b_i^* h(T_i^* T_j) b_j &= \sum_{i,j} b_i^* \langle x, T_i^* T_j x \rangle b_j \\ &= \sum_{i,j} \langle T_i x b_i, T_j x b_j \rangle \\ &= \left\langle \sum_i T_i x b_i, \sum_i T_i x b_i \right\rangle \geq 0, \end{aligned}$$

which shows  $h$  is indeed completely positive. □

## 2.2.2 Ultranorm uniformity

**146** In 136 II we saw how to complete a complex vector space with inner product to a Hilbert space. Now we will see how to complete a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product to a *self-dual* Hilbert  $\mathcal{B}$ -module under the assumption  $\mathcal{B}$  is a von Neumann algebra. We will use a different construction than Paschke used [93]. Paschke showed that for a pre-Hilbert  $\mathcal{B}$ -module  $X$  the set of functionals  $X'$  (i.e.  $\mathcal{B}$ -module homomorphisms into  $\mathcal{B}$ ) turns out to be a self-dual Hilbert  $\mathcal{B}$ -module, which he uses as the completion of  $X$ . A considerable part of his paper [93] is devoted to this construction. It requires Sakai's characterization of von Neumann algebras, which we have not covered. To avoid developing Sakai's theory, we give a different construction. Instead of embedding  $X$  into a dual space, we will stay closer to the similar fact for Hilbert spaces and use a topological completion. A simple metric completion will not do, we will need to complete  $X$  as a *uniform space*. Uniformities are structures that sit between topological spaces and metric spaces: every metric space is a uniformity (see V) and every uniformity is a topological space. We will refresh the basics of uniformities — for a thorough treatment, see for instance [133, ch. 9].

**II Definition** A **uniform space** is a set  $X$  together with a family of relations  $\Phi \subseteq \wp(X \times X)$  called **entourages** satisfying the following conditions.

1. The set of entourages  $\Phi$  is a filter. That is: (a) if  $\varepsilon, \delta \in \Phi$ , then  $\varepsilon \cap \delta \in \Phi$  and (b) if  $\varepsilon \subseteq \delta$  and  $\varepsilon \in \Phi$ , then  $\delta \in \Phi$ .
2. For every  $\varepsilon \in \Phi$  and  $x \in X$  we have  $x \varepsilon x$ .
3. For each  $\varepsilon \in \Phi$ , there is a  $\delta \in \Phi$  such that  $\delta^2 \subseteq \varepsilon$ . So, if  $x \delta y$  and  $y \delta z$  then  $x \varepsilon z$  for any  $x, y, z \in X$ .
4. For every  $\varepsilon \in \Phi$ , there is a  $\delta \in \Phi$  with  $\delta^{-1} \subseteq \varepsilon$ . So, if  $x \delta y$  then  $y \varepsilon x$  for any  $x, y \in X$ .

A uniform space is **Hausdorff** whenever  $x \varepsilon y$  for all  $\varepsilon \in \Phi$  implies that  $x = y$ .

**III** The elements of  $X$  are the points of the uniform space. The entourages  $\varepsilon \in \Phi$  are generalized distances: one reads  $x \varepsilon y$  as ‘the point  $x$  is at most  $\varepsilon$  far from to  $y$ ’. With this in mind, the second axiom states every point is arbitrarily close to itself. The third axiom requires that for every entourage  $\varepsilon$  there is an entourage which acts like  $\varepsilon/2$ . There might be several  $\varepsilon/2$  which fit the bill. Whenever we write  $\varepsilon/2$  we implicitly pick some entourage that satisfies  $(\varepsilon/2)^2 \subseteq \varepsilon$ . Also we will use the obvious shorthand  $\varepsilon/4 = (\varepsilon/2)/2$ .

**IIIa Definition** Let  $X$  be a set together with a family of relations  $B \subseteq \wp(X \times X)$ . The set  $B$  is said to be a **subbase (for a uniformity on  $X$ )** if it satisfies axioms 2, 3 and 4 of II. (Cf. [133, dfn. 5.5].)

**Exercise** Let  $X$  be a set together with a subbase  $B$ . Write  $\Phi$  for the filter generated by  $B$  (that is:  $\delta \in \Phi$  iff  $\varepsilon_1 \cap \dots \cap \varepsilon_n \subseteq \delta$  for some  $\varepsilon_1, \dots, \varepsilon_n \in B$ ). Show that  $(X, \Phi)$  is a uniform space. IV

**Examples** Using the definition of a subbase, it is easy to describe the entourages of some common uniformities. V

1. Let  $(X, d)$  be a metric space. Define  $\hat{\varepsilon} \equiv \{(x, y); d(x, y) \leq \varepsilon\}$  for any  $\varepsilon > 0$ . The set  $B \equiv \{\hat{\varepsilon}; \varepsilon > 0\}$  is a subbase and so fixes a uniformity  $\Phi$  on  $X$ . In this sense every metric space is a uniformity.
2. Let  $X$  be a set together with a (infinite) family  $(d_\alpha)_{\alpha \in I}$  of pseudometrics.<sup>||</sup> Define  $E_{\alpha, \varepsilon} = \{(x, y); d_\alpha(x, y) \leq \varepsilon\}$ . Then  $B \equiv \{E_{\alpha, \varepsilon}; \varepsilon > 0, \alpha \in I\}$  is a subbase and fixes a uniformity  $\Phi$  on  $X$ . (Every uniform space is of this form, see e.g. [133, thm. 39.11].)
3. Let  $V$  be a vector space with a family of seminorms  $(\|\cdot\|_\alpha)_{\alpha \in I}$ . As a special case of the previous,  $V$  has a uniformity fixed by pseudometrics  $d_\alpha(x, y) = \|x - y\|_\alpha$ . (If additionally  $x = 0$  provided  $\|x\|_\alpha = 0$  for all  $\alpha$ , then  $V$  is called a locally convex space.)
4. Assume  $\mathcal{B}$  is a von Neumann algebra. There are several uniformities on  $\mathcal{B}$  — two of which are of particular interest to us. The **ultrastrong uniformity** on  $\mathcal{B}$  is fixed by the seminorms  $\|b\|_f = f(b^*b)^{\frac{1}{2}}$  where  $f: \mathcal{B} \rightarrow \mathbb{C}$  is an np-map. The **ultraweak uniformity** on  $\mathcal{B}$  is given by the seminorms  $|f(b)|$  for np-maps  $f: \mathcal{B} \rightarrow \mathbb{C}$ .

We are ready to define a uniformity for Hilbert  $\mathcal{B}$ -modules. Later, we will complete  $X$  with respect to this uniformity. VI

**Definition** Let  $\mathcal{B}$  be a von Neumann algebra. Assume  $X$  is a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product  $[\cdot, \cdot]$ . We call the uniformity on  $X$  given by seminorms  $\|x\|_f = f([x, x])^{\frac{1}{2}}$  for np-maps  $f: \mathcal{B} \rightarrow \mathbb{C}$  the **ultranorm uniformity**. VII

The ultranorm uniformity will play a very similar rôle to the norm for Hilbert spaces. If  $\mathcal{B} = \mathbb{C}$ , then the ultranorm uniformity is the same as the uniformity induced by the norm. If  $X = \mathcal{B}$  with  $[a, b] = a^*b$ , then the ultranorm uniformity coincides with the ultrastrong uniformity. VIII

**Beware** The ultranorm uniformity is (in general) not given by a single norm. Furthermore the ultranorm uniformity is weaker than the norm uniformity (that is: norm convergence implies ultranorm convergence, but not necessarily the other way around), even though on von Neumann algebras the ultraweak uniformity is stronger than the weak uniformity. IX

---

<sup>||</sup>Like a metric, a pseudometric  $d$  is symmetric, obeys the triangle inequality and  $d(x, x) = 0$ , but unlike a metric  $d(x, y) = 0$  is not required to entail  $x = y$ .

× **Remarks** The ultranorm uniformity has appeared earlier in the literature under various different names; for instance as the  $\tau_1$ -topology in [88, thm. 3.5.1] and as the s-topology in [38, 2.9]. In previous work the ultranorm uniformity only plays a minor rôle, often amongst other topologies. In this thesis we put the ultranorm uniformity center-stage as the right generalization of the norm and give it the ‘ultranorm’ name to match this view.

**147 Definition** Let  $X$  be a uniform space with entourages  $\Phi$ . It is easy to translate the familiar notions for metric spaces to uniform spaces.

1. A net  $(x_\alpha)_\alpha$  is said to **converge** to  $x$  if for each  $\varepsilon \in \Phi$  there is an  $\alpha_0$  such that for all  $\alpha > \alpha_0$  we have  $x \varepsilon x_\alpha$ . Note that a net can converge to two different points. Indeed, this is the case if we start off with a pseudometric that is not a metric.
2. A net  $(x_\alpha)_\alpha$  is called **Cauchy** if for each  $\varepsilon \in \Phi$  there is an  $\alpha_0$  such that for all  $\alpha, \beta > \alpha_0$  we have  $x_\alpha \varepsilon x_\beta$ . The uniform space  $X$  is **complete** when every Cauchy net converges.
3. Let  $(Y, \Psi)$  be a second uniform space. A map  $f: X \rightarrow Y$  is said to be **uniformly continuous** if for each  $\varepsilon \in \Psi$  there is a  $\delta \in \Phi$  such that for all  $x \delta y$  we have  $f(x) \varepsilon f(y)$ . The map  $f$  is merely **continuous** if for each  $x$  and  $\varepsilon \in \Psi$ , there is a  $\delta \in \Phi$  such that for all  $x \delta y$  we know  $f(x) \varepsilon f(y)$ .
4. We say two Cauchy nets  $(x_\alpha)_{\alpha \in I}$  and  $(y_\beta)_{\beta \in J}$  are **equivalent**, in symbols:  $(x_\alpha)_\alpha \sim (y_\beta)_\beta$ , when for every  $\varepsilon \in \Phi$  there are  $\alpha_0 \in I$  and  $\beta_0 \in J$  such that for all  $\alpha \geq \alpha_0$  and  $\beta \geq \beta_0$  we have  $x_\alpha \varepsilon y_\beta$ .
5. A subset  $D \subseteq X$  is said to be **dense** if for each  $\varepsilon \in \Phi$  and  $x \in X$ , there is a  $y \in D$  with  $x \varepsilon y$ .

II **Exercise** In the same setting as the previous definition.

1. Show that equivalence of Cauchy nets is an equivalence relation. Show that if  $(x_\alpha)_\alpha$  is a subnet of a Cauchy net  $(y_\alpha)_\alpha$ , that then  $(x_\alpha)_\alpha$  is equivalent to  $(y_\alpha)_\alpha$ .
2. Prove that if  $(x_\alpha)_\alpha$  and  $(y_\alpha)_\alpha$  are equivalent Cauchy nets and  $x_\alpha \rightarrow x$ , then also  $y_\alpha \rightarrow x$ .
3. Assume  $(x_\alpha)_\alpha$  is a net with  $x_\alpha \rightarrow x$  and  $x_\alpha \rightarrow y$ . Prove  $x = y$  whenever  $X$  is Hausdorff.
4. Show that if  $f: X \rightarrow Y$  is a continuous map between uniform spaces and we have  $x_\alpha \rightarrow x$  in  $X$ , then  $f(x_\alpha) \rightarrow f(x)$  in  $Y$ .

5. Assume  $f$  is a uniformly continuous map between uniform spaces. Show that  $f$  maps Cauchy nets to Cauchy nets and furthermore that  $f$  maps equivalent Cauchy nets to equivalent Cauchy nets.
6. Suppose  $D \subseteq X$  is a dense subset. Show that for each  $x \in X$  there is a Cauchy net  $(d_\alpha)_{\alpha \in \Phi}$  in  $D$  with  $d_\alpha \rightarrow x$  in  $X$ .
7. Assume  $f, g: X \rightarrow Y$  are continuous maps between uniform spaces where  $Y$  is Hausdorff. Conclude from 4 and 6 that  $f$  and  $g$  are equal whenever they agree on a dense subset of  $X$ .

**Exercise** Let  $(X_i)_{i \in I}$  be a family of sets with uniformities  $(\Phi_i)_{i \in I}$ . For each  $i_0 \in I$  and  $\varepsilon \in \Phi_{i_0}$ , define a relation on  $\prod_{i \in I} X_i$  by  $(x_i)_{i \in I} \hat{\varepsilon} (y_i)_{i \in I} \iff x_{i_0} \varepsilon y_{i_0}$ . Show  $\{\hat{\varepsilon}; \varepsilon \in \Phi_i, i \in I\}$  is a subbase for  $\prod_{i \in I} X_i$ ; that the projections  $\pi_i: \prod_{i \in I} X_i \rightarrow X_i$  are uniformly continuous with respect to them and that they make  $\prod_{i \in I} X_i$  into the product of  $(X_i)_{i \in I}$  in the category of uniform spaces with uniformly continuous maps. |||

**Proposition** Let  $X$  and  $Y$  be right  $\mathcal{B}$ -modules with  $\mathcal{B}$ -valued inner products. A bounded  $\mathcal{B}$ -linear map  $T: X \rightarrow Y$  is uniformly ultranorm continuous. 148

**Proof** Let  $f: \mathcal{B} \rightarrow \mathbb{C}$  be an np-map and  $\varepsilon > 0$ . Assume  $\|x - y\|_f \leq \frac{\varepsilon}{\|T\|}$ . By 144V we have  $\langle T(x - y), T(x - y) \rangle \leq \|T\|^2 \langle x - y, x - y \rangle$  and so ||

$$\|Tx - Ty\|_f = f(\langle T(x - y), T(x - y) \rangle)^{\frac{1}{2}} \leq \|T\| \|x - y\|_f \leq \varepsilon.$$

Thus  $T$  is uniformly continuous w.r.t. the ultranorm uniformity. □

**Corollary** Let  $\mathcal{B}$  be a von Neumann algebra and  $X$  a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product. For  $x_0 \in X$  the maps |||

$$\begin{array}{lll} (x, y) \mapsto x + y & x \mapsto \langle x_0, x \rangle & b \mapsto x_0 b \\ X \times X \rightarrow X & X \rightarrow \mathcal{B} & \mathcal{B} \rightarrow X \end{array}$$

are all uniformly continuous w.r.t. the ultranorm uniformity.

**Exercise** Let  $\mathcal{B}$  be a von Neumann algebra and  $X$  a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product. Show that  $x \mapsto xb$  is ultranorm continuous for any  $b \in \mathcal{B}$ . IV

**Proposition** Let  $X$  be a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product  $[\cdot, \cdot]$  for some  $C^*$ -algebra  $\mathcal{B}$ . If  $x_\alpha \rightarrow x$  and  $y_\alpha \rightarrow y$  in the ultranorm uniformity, for some nets  $(x_\alpha)_{\alpha \in I}, (y_\alpha)_{\alpha \in I}$ , then  $[x_\alpha, y_\alpha] \rightarrow [x, y]$  ultraweakly. V

**Proof** Let  $\varepsilon > 0$  and np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$  be given. Note that  $\|y_\alpha\|_f \rightarrow \|y\|_f$ ; indeed by the reverse triangle inequality  $|\|y_\alpha\|_f - \|y\|_f| \leq \|y_\alpha - y\|_f \rightarrow 0$ . Thus we can find  $\alpha_1$  such that  $\|y_\alpha\|_f \leq \|y\|_f + 1$  for  $\alpha \geq \alpha_1$ . VI

We can find  $\alpha_2$  such that for all  $\alpha \geq \alpha_2$  we have  $\|x - x_\alpha\|_f \leq \frac{1}{2}\varepsilon(\|y\|_f + 1)^{-1}$  and  $\|y - y_\alpha\|_f \leq \frac{1}{2}\varepsilon(\|x\|_f + 1)^{-1}$ . Pick  $\alpha_0 \geq \alpha_1, \alpha_2$ . Then for  $\alpha \geq \alpha_0$  we have

$$\begin{aligned} |f([x_\alpha, y_\alpha] - [x, y])| &= |[x_\alpha - x, y_\alpha]_f + [x, y_\alpha - y]_f| \\ &\leq \|x_\alpha - x\|_f \|y_\alpha\|_f + \|x\|_f \|y_\alpha - y\|_f && \text{by 142 III} \\ &\leq \|x_\alpha - x\|_f (\|y\|_f + 1) + (\|x\|_f + 1) \|y_\alpha - y\|_f \\ &\leq \varepsilon. \end{aligned}$$

Thus  $[x_\alpha, y_\alpha] \rightarrow [x, y]$  ultraweakly.  $\square$

**VII Corollary** Let  $X$  be a Hilbert  $\mathcal{B}$ -module together with an ultranorm-dense subset  $D \subseteq X$ . The vector states from  $D$  are order separating: for each  $T \in \mathcal{B}^a(X)$  we have  $T \geq 0$  if and only if  $\langle x, Tx \rangle \geq 0$  for all  $x \in D$ . Cf. 144 I.

**149 Definition** Let  $X$  be pre-Hilbert  $\mathcal{B}$ -module for a von Neumann algebra  $\mathcal{B}$ . Let  $E \subseteq X$  be some subset.

1.  $E$  is **orthogonal** if  $\langle e, d \rangle = 0$  for  $e, d \in E$  with  $e \neq d$ .
2.  $E$  is **orthonormal** if additionally  $\langle e, e \rangle$  is a non-zero projection for  $e \in E$ .
3. A family  $(b_e)_{e \in E}$  in  $\mathcal{B}$  is said to be  **$\ell^2$ -summable** if the partial sums of  $\sum_{e \in E} b_e^* b_e$  are bounded.
4.  $E$  is an **(orthonormal) basis** if it is orthonormal and additionally
  - (a) for each  $x \in X$  we have

$$x = \sum_{e \in E} e \langle e, x \rangle,$$

where the sum converges in the ultranorm uniformity *and*

- (b) the sum  $\sum_{e \in E} e b_e$  converges in the ultranorm uniformity for any  $\ell^2$ -summable  $(b_e)_{e \in E}$  in  $\mathcal{B}$ .

**II Notation** As with Hilbert spaces, it will sometimes be more convenient to view an orthonormal basis as a sequence/family  $(e_i)_{i \in I}$ , instead of a subset. This should cause no confusion.

**IIa Remark** The first assumption ‘(a)’ in the definition of an orthonormal basis  $E$  requires that every element  $x$  can be reconstructed from its coefficients  $(\langle e, x \rangle)_{e \in E}$ . This assumption fails, for instance, if there is a non-zero element orthogonal to all basis elements. The second assumption ‘(b)’ requires that behind any a reasonable family of coefficients, there is a corresponding element. This second assumption fails if the module is not ultranorm complete, see V.

**Beware** In the literature, there are several different definitions of an orthonormal basis for a Hilbert  $\mathcal{B}$ -module  $X$ . IIb

1. Some [82, 88] define an orthonormal basis as an orthonormal subset  $E \subseteq X$  such that  $x = \sum_{e \in E} e \langle e, x \rangle$  for all  $x \in X$  where the sum converges in the *norm*. This notion of basis is independent of ours: there is a basis in our sense that is not a basis in their sense ( $\{|0\rangle\langle 0|, |1\rangle\langle 1|, \dots\}$  in  $\mathcal{B}(\ell^2)$ ) and vice versa (the standard Hilbert module  $H_{\mathcal{B}(\ell^2)}$  defined in [82, §3] has a basis in their sense, but not a basis in ours as it is not self dual, cf. V.)
2. Others such as [12, lemma 8.5.23] define an orthonormal basis only when  $X$  is a self-dual (V) Hilbert  $C^*$ -module over a von Neumann algebra. In this case an orthonormal basis is defined as an orthonormal subset  $E \subseteq X$  such that  $x = \sum_{e \in E} e \langle e, x \rangle$  for all  $x \in X$  where the sum converges in the  $w^*$ -topology of  $X$  (with respect to the unique Banach space predual of  $X$  for which the inner product is separately  $w^*$ -continuous, see [12, lemma 8.5.4].) This notion is equivalent to ours, but only defined for self-dual modules over von Neumann algebras.

**Exercise** Let  $X$  be a pre-Hilbert  $\mathcal{B}$ -module for a  $C^*$ -algebra  $\mathcal{B}$ . Show that we have  $e \langle e, e \rangle = e$  for any  $e \in X$  where  $\langle e, e \rangle$  is a projection. III

(Hint: consider  $\|e(1 - \langle e, e \rangle)\|^2$ ; cf. [93, thm. 3.12].)

**Exercise** Let  $X$  be a pre-Hilbert  $\mathcal{B}$ -module for a von Neumann algebra  $\mathcal{B}$  with an orthonormal basis  $E \subseteq X$ . Show **Parseval's identity** holds, that is: IV

$$\langle x, x \rangle = \sum_{e \in E} \langle x, e \rangle \langle e, x \rangle,$$

where the sum converges ultraweakly. [93] (Hint: use 148V.)

**Theorem** Assume  $\mathcal{B}$  is a von Neumann algebra. For any pre-Hilbert  $\mathcal{B}$ -module  $X$ , the following conditions are equivalent. V

1.  $X$  is self dual.
2.  $X$  is ultranorm complete.
3. Every norm-bounded ultranorm-Cauchy net in  $X$  converges.
4.  $X$  has an orthonormal basis.

**Proof** We will prove  $1 \Rightarrow 3 \Rightarrow 4 \Rightarrow 2 \Rightarrow 4 \Rightarrow 1$ . The equivalence  $1 \Leftrightarrow 4$  was already shown by Paschke in [93, thm. 3.12], but we give a different proof. The equivalence  $1 \Leftrightarrow 3$  turns out proven by Frank already [32, thm. 3.2] in a different way. The final equivalence  $1 \Leftrightarrow 2$  is new, as far as we know. VI

VII (1  $\Rightarrow$  3, self-dual  $X$  are bounded complete) Let  $X$  be a self-dual pre-Hilbert  $\mathcal{B}$ -module. To show norm-bounded ultranorm completeness, assume  $(x_\alpha)_\alpha$  is an ultranorm-Cauchy net in  $X$  with  $\|x_\alpha\| \leq B$  for some  $B \geq 0$ . For the moment pick an  $y \in X$ . By 142III we have  $f(\langle x_\alpha, y \rangle \langle y, x_\alpha \rangle) \leq \|y\|^2 f(\langle x_\alpha, x_\alpha \rangle)$  for any np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$  and so  $\langle y, x_\alpha \rangle$  is an ultrastrong-Cauchy net in  $\mathcal{B}$ . Von Neumann algebras are ultrastrongly complete, see 77I, so we can define  $\tau(y) = (\text{uslim}_\alpha \langle y, x_\alpha \rangle)^*$ , where **uslim** denotes the ultrastrong limit. As addition and multiplication by a fixed element are ultrastrongly continuous (see 45IV)  $\tau$  is  $\mathcal{B}$ -linear. We will show  $\tau$  is bounded. For each  $\alpha$  we have  $\langle x_\alpha, y \rangle \langle y, x_\alpha \rangle \leq \|y\|^2 \langle x_\alpha, x_\alpha \rangle \leq \|y\|^2 B^2$  and so

$$\tau(y)\tau(y)^* \stackrel{46II}{=} \text{uwlim}_\alpha \langle x_\alpha, y \rangle \langle y, x_\alpha \rangle \leq \|y\|^2 B^2,$$

where **uwlim** denotes the ultraweak limit. Thus  $\|\tau(y)\|^2 = \|\tau(y)\tau(y)^*\| \leq \|y\|^2 B^2$ . So  $\tau$  is bounded. By self-duality, there is a  $t \in X$  such that  $\tau(y) = \langle t, y \rangle$  for all  $y \in X$ . And so for each np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$  we have

$$\|\langle y, t - x_\alpha \rangle\|_f = \|\langle y, t \rangle - \langle y, x_\alpha \rangle\|_f = \|\tau(y)^* - \langle y, x_\alpha \rangle\|_f \rightarrow 0. \quad (2.5)$$

In 142II we introduced the inner product  $\langle \cdot, \cdot \rangle_f$  defined by  $\langle x, y \rangle_f \equiv f(\langle x, y \rangle)$ . From Kadison's inequality 30IV it follows  $|f(a)|^2 \leq f(a^*a)f(1)$  and so

$$|\langle y, t - x_\alpha \rangle_f|^2 \leq f(\langle t - x_\alpha, y \rangle \langle y, t - x_\alpha \rangle) f(1) = \|\langle y, t - x_\alpha \rangle\|_f^2 f(1). \quad (2.6)$$

Combining (2.5) and (2.6), we see  $|\langle y, t - x_\alpha \rangle_f| \rightarrow 0$ . Let  $\varepsilon > 0$  be given. As  $\|x_\alpha\|_f$  is norm bounded we can find  $B > 0$  such that  $\|x_\alpha\|_f \leq B$  for all  $\alpha$ . As  $(x_\alpha)_\alpha$  is ultranorm Cauchy, we can pick  $\alpha$  such that  $\|x_\beta - x_\alpha\|_f \leq (\|t\|_f + B)^{-1} \frac{\varepsilon}{2}$  for all  $\beta \geq \alpha$ . Find  $\beta$  such that  $|\langle t - x_\alpha, t - x_\beta \rangle_f| \leq \frac{\varepsilon}{2}$ . Putting it all together:

$$\begin{aligned} \langle t - x_\alpha, t - x_\alpha \rangle_f &\leq |\langle t - x_\alpha, t - x_\beta \rangle_f| + |\langle t - x_\alpha, x_\beta - x_\alpha \rangle_f| \\ &\leq \frac{\varepsilon}{2} + \|t - x_\alpha\|_f \|x_\beta - x_\alpha\|_f \\ &\leq \frac{\varepsilon}{2} + (\|t\|_f + B) \|x_\beta - x_\alpha\|_f \leq \varepsilon. \end{aligned}$$

Thus  $x_\alpha$  converges ultranorm to  $t$ . So  $X$  is ultranorm bounded complete.

VIII (3  $\Rightarrow$  4, bounded complete  $X$  has an orthonormal basis) Assume norm-bounded ultranorm-Cauchy nets converge in  $X$ . By Zorn's lemma, there is a maximal orthonormal subset  $E \subseteq X$ . We will show  $E$  is a basis.

Let  $(b_e)_{e \in E}$  be an  $E$ -tuple in  $\mathcal{B}$  such that the partial sums of  $\sum_{e \in E} b_e^* b_e$  are bounded. We want to show  $\sum_{e \in E} e b_e$  converges in the ultranorm uniformity. As the summands of  $\sum_{e \in E} b_e^* b_e$  are positive, the partial sums form a directed bounded net that converges ultrastrongly to its supremum  $\sum_{e \in E} b_e^* b_e$ . By III,

we have  $e \langle e, e \rangle = e$  for each  $e \in E$  and so  $\langle x, e \rangle \langle e, e \rangle = \langle x, e \rangle$  for every  $x \in X$ . Thus for any finite subset  $S \subseteq E$

$$\left\langle \sum_{e \in S} eb_e, \sum_{e \in S} eb_e \right\rangle = \sum_{e \in S} b_e^* \langle e, e \rangle b_e \leq \sum_{e \in S} b_e^* b_e.$$

To show  $\sum_e eb_e$  converges ultranorm, pick any np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$ . As  $\sum_{e \in E} b_e^* b_e$  converges ultraweakly, we have  $\sum_{e \in E} f(b_e^* b_e) < \infty$ . Thus the tails of the series  $\sum_{e \in E-S} f(b_e^* b_e)$  tend to zero, hence for finite  $S, T \subseteq E$  we know

$$\left\| \sum_{e \in S} eb_e - \sum_{e \in T} eb_e \right\|_f^2 = \left\| \sum_{e \in S \Delta T} eb_e \right\|_f^2 \leq \sum_{e \in S \Delta T} f(b_e^* b_e).$$

The latter vanishes as  $S \cap T$  grows. Thus the partial sums of  $\sum_{e \in E} eb_e$  are ultranorm Cauchy and norm-bounded by  $\left\| \sum_{e \in E} b_e^* b_e \right\|$ , so  $\sum_{e \in E} eb_e$  converges in the ultranorm uniformity.

Pick  $x \in X$ . We have to show  $\sum_{e \in E} x \langle e, x \rangle$  converges in the ultranorm uniformity to  $x$ . For any finite subset  $S \subseteq E$  we have

$$0 \leq \left\langle x - \sum_{e \in S} e \langle e, x \rangle, x - \sum_{e \in S} e \langle e, x \rangle \right\rangle = \langle x, x \rangle - \sum_{e \in S} \langle x, e \rangle \langle e, x \rangle.$$

Rearranging we find [Bessel's inequality](#):

$$\sum_{e \in S} \langle x, e \rangle \langle e, x \rangle \leq \langle x, x \rangle.$$

Hence the  $E$ -tuple  $(\langle e, x \rangle)_e$  is  $\ell^2$ -summable and so  $\sum_{e \in E} e \langle e, x \rangle$  converges in the ultranorm uniformity. Consider  $x' = x - \sum_{e \in E} e \langle e, x \rangle$ . We are done if we can show  $x' = 0$ .

To see  $x' = 0$ , we first show that  $X$  has [polar decomposition](#): for each  $y \in X$ , there is an  $u \in X$  with  $\langle u, u \rangle = \lceil \langle y, y \rangle \rceil$  and  $y = u \langle y, y \rangle^{\frac{1}{2}}$ . By 80IV the positive element  $\langle y, y \rangle^{\frac{1}{2}}$  has an approximate pseudoinverse  $h_1, h_2, \dots$  — that is:  $h_n \langle y, y \rangle^{\frac{1}{2}} = \langle y, y \rangle^{\frac{1}{2}} h_n = \lceil h_n \rceil$  and  $\sum_n h_n \langle y, y \rangle^{\frac{1}{2}} = \sum_n \lceil h_n \rceil = \lceil \langle y, y \rangle \rceil$ . Note

$$\langle y h_n, y h_m \rangle = (h_n \langle y, y \rangle^{\frac{1}{2}}) (\langle y, y \rangle^{\frac{1}{2}} h_m) = \lceil h_n \rceil \lceil h_m \rceil$$

and so using that  $\lceil h_1 \rceil, \lceil h_2 \rceil, \dots$  are pairwise orthogonal, we see

$$\left\langle \sum_{n=1}^N y h_n, \sum_{n=1}^N y h_n \right\rangle = \sum_{n=1}^N \sum_{m=1}^N \lceil h_n \rceil \lceil h_m \rceil = \sum_{n=1}^N \lceil h_n \rceil \leq \lceil \langle y, y \rangle \rceil.$$

Thus  $(\sum_{n=1}^N y h_n)_N$  forms an ultranorm-Cauchy net norm bounded by 1. By assumption this bounded net converges: write  $u \equiv \sum_n y h_n$ . (So in the special

case  $X = \mathcal{B}$ , we have  $u = y/\langle y, y \rangle^{\frac{1}{2}}$ , see 81 III.) We derive

$$\begin{aligned}
u \langle y, y \rangle^{\frac{1}{2}} &= \sum_n y h_n \langle y, y \rangle^{\frac{1}{2}} && \text{by dfn. } u \text{ and 148 IV} \\
&= \sum_n y [h_n] && \text{by dfn. } h_n \\
&= y \sum_n [h_n] && \text{by 148 III} \\
&= y [\langle y, y \rangle] && \text{by dfn. } h_n \\
&= y,
\end{aligned}$$

where the last equality follows from  $y(1 - [\langle y, y \rangle]) = 0$ , which is justified by

$$\langle y(1 - [\langle y, y \rangle]), y(1 - [\langle y, y \rangle]) \rangle = (1 - [\langle y, y \rangle]) \langle y, y \rangle (1 - [\langle y, y \rangle]) = 0.$$

This finishes the proof of polar decomposition.

Recall we want to show  $x' \equiv x - \sum_{e \in E} e \langle e, x \rangle = 0$ . Reasoning towards contradiction, assume  $x' \neq 0$ . By polar decomposition  $x' = u \langle x', x' \rangle^{\frac{1}{2}}$  for some  $u \in X$  with  $\langle u, u \rangle = [\langle x', x' \rangle]$ . Note  $\langle u, u \rangle \neq 0$  for otherwise  $x' = 0$ , quod non. For any  $e_0 \in E$ , we have

$$\langle e_0, x' \rangle = \langle e_0, x - \sum_{e \in E} e \langle e, x \rangle \rangle \stackrel{148 \text{ III}}{=} \langle e_0, x \rangle - \sum_{e \in E} \langle e_0, e \rangle \langle e, x \rangle = 0$$

and so  $0 = \langle e, x' \rangle = \langle e, u \rangle \langle x', x' \rangle^{\frac{1}{2}}$  for any  $e \in E$ . Thus

$$\langle e, u \rangle = \langle e, u \rangle \langle u, u \rangle = \langle e, u \rangle [\langle x', x' \rangle] \stackrel{60 \text{ VIII}}{=} 0.$$

Thus  $E \cup \{u\}$  contradicts the maximality of  $E$ . Hence  $x' = 0$  and so  $x = \sum_{e \in E} e \langle e, x \rangle$  as desired.

IX (4  $\Rightarrow$  2, basis implies completeness) Assume  $X$  has an orthonormal basis  $E$ . Let  $(x_\alpha)_\alpha$  be an ultranorm-Cauchy net. Pick  $e \in E$ . By 148 III  $(\langle e, x_\alpha \rangle)_\alpha$  is an ultrastrong-Cauchy net. As von Neumann algebras are ultrastrongly complete (see 77 I) there is a  $b_e \in \mathcal{B}$  to which  $\langle e, x_\alpha \rangle$  converges ultrastrongly. We will show  $x_\alpha$  converges in the ultranorm uniformity to  $\sum_{e \in E} e b_e$ .

We will first show that  $\sum_{e \in E} e b_e$  converges. As before, it is sufficient to show  $\sum_{e \in E} b_e^* b_e$  is bounded. For this, it suffices to show  $\sum_{e \in E} f(b_e^* b_e)$  is bounded for each np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$ , as ultraweakly-bounded nets are norm bounded, see 87 VIII. For any  $e \in E$  we have  $\|b_e\|_f = \lim_\alpha \|\langle e, x_\alpha \rangle\|_f$  and

so  $f(b_e^* b_e) = \lim_\alpha f(\langle x_\alpha, e \rangle \langle e, x_\alpha \rangle)$ . Pick a finite subset  $S \subseteq E$ . Then we have

$$\begin{aligned} f\left(\sum_{e \in S} b_e^* b_e\right) &= \sum_{e \in S} \lim_\alpha f(\langle x_\alpha, e \rangle \langle e, x_\alpha \rangle) \\ &= \lim_\alpha f\left(\sum_{e \in S} \langle x_\alpha, e \rangle \langle e, x_\alpha \rangle\right) \\ &\leq f(\langle x_\alpha, x_\alpha \rangle) \qquad \text{by Bessel's inequality.} \end{aligned}$$

As  $x_\alpha$  is ultranorm Cauchy, the net  $f(\langle x_\alpha, x_\alpha \rangle)$  must be Cauchy and so bounded. Hence  $\sum_{e \in E} e b_e$  converges in the ultranorm uniformity.

To prove  $x_\alpha$  converges to  $\sum_{e \in E} e b_e$ , pick any  $\varepsilon > 0$  and np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$ . We want to find  $\alpha_0$  such that  $\|x_\alpha - \sum_{e \in E} e b_e\|_f \leq \varepsilon$  for all  $\alpha \geq \alpha_0$ . As  $(x_\alpha)_\alpha$  is ultranorm Cauchy, we can pick an  $\alpha_0$  such that  $\|x_\alpha - x_\beta\|_f \leq \frac{1}{2\sqrt{2}}\varepsilon$  for all  $\alpha, \beta \geq \alpha_0$ . As  $x_\alpha = \sum_{e \in E} e \langle e, x_\alpha \rangle$  we find with Parseval's identity (IV) that

$$\|x_\alpha - \sum_{e \in E} e b_e\|_f^2 = \left\| \sum_{e \in E} e(\langle e, x_\alpha \rangle - b_e) \right\|_f^2 = \sum_{e \in E} \|\langle e, x_\alpha \rangle - b_e\|_f^2.$$

Take a finite subset  $S \subseteq E$  such that  $\sum_{e \in E-S} \|\langle e, x_\alpha \rangle - b_e\|_f^2 \leq \frac{1}{2}\varepsilon^2$ . If we can also show  $\sum_{e \in S} \|\langle e, x_\alpha \rangle - b_e\|_f^2 \leq \frac{1}{2}\varepsilon^2$ , then we are done. For any  $\beta \geq \alpha$  we have using the triangle inequality of  $\|\cdot\|_f$  on  $X$

$$\begin{aligned} \left(\sum_{e \in S} \|\langle e, x_\alpha \rangle - b_e\|_f^2\right)^{\frac{1}{2}} &= \left\| \sum_{e \in S} e \langle e, x_\alpha \rangle - e b_e \right\|_f \\ &\leq \left\| \sum_{e \in S} e \langle e, x_\alpha - x_\beta \rangle \right\|_f + \left\| \sum_{e \in S} e \langle e, x_\beta \rangle - e b_e \right\|_f \\ &= \left(\sum_{e \in S} \|\langle e, x_\alpha - x_\beta \rangle\|_f^2\right)^{\frac{1}{2}} + \left(\sum_{e \in S} \|\langle e, x_\beta \rangle - b_e\|_f^2\right)^{\frac{1}{2}}. \end{aligned}$$

We want to show the previous is bounded by  $\frac{1}{\sqrt{2}}\varepsilon$ . We bound the two terms separately. For the first term using Bessel's inequality and our choice of  $\alpha_0$  we have  $\sum_{e \in S} \|\langle e, x_\alpha - x_\beta \rangle\|_f^2 \leq \|x_\alpha - x_\beta\|_f^2 \leq \left(\frac{1}{2\sqrt{2}}\varepsilon\right)^2$ . As  $\|\langle e, x_\beta \rangle - b_e\|_f$  vanishes for each  $e \in S$  and  $S$  is finite, we can find sufficiently large  $\beta$  such that the right term is also bounded by  $\frac{1}{2\sqrt{2}}\varepsilon$ . Thus  $x_\alpha$  converges in the ultranorm uniformity to  $\sum_{e \in E} e b_e$ , as desired.

(2  $\Rightarrow$  4, completeness implies basis) Follows from 3  $\Rightarrow$  4 as 2  $\Rightarrow$  3 is trivial. X  
(4  $\Rightarrow$  1, basis implies self-duality) Assume  $X$  has an orthonormal basis  $E$ . XI  
Suppose  $\tau: X \rightarrow \mathcal{B}$  is a bounded  $\mathcal{B}$ -linear map. We want to show there is some  $t \in X$  such that  $\tau(x) = \langle t, x \rangle$  for all  $x \in X$ . Pick any  $x \in X$ . By assumption  $\sum_{e \in E} e \langle e, x \rangle$  converges in the ultranorm uniformity to  $x$ . As by 148  $\tau$  is

ultranorm continuous we see

$$\tau(x) = \sum_{e \in E} \tau(e) \langle e, x \rangle = \sum_{e \in E} \langle e\tau(e)^*, x \rangle.$$

Thus if we can show  $\sum_e e\tau(e)^*$  converges in the ultranorm, we are done. By assumption this is the case if  $(\tau(e)^*)_{e \in E}$  is  $\ell^2$ -summable — that is: the partial sums of  $\sum_{e \in E} \tau(e)\tau(e)^*$  must be bounded. We will show that for finite  $S \subseteq E$  we have  $\|\sum_{e \in S} e\tau(e)^*\| \leq \|\tau\|$ , which is sufficient as then  $\|\sum_{e \in S} \tau(e)\tau(e)^*\| = \|\sum_{e \in S} e\tau(e)^*\|^2 \leq \|\tau\|^2$ . For the moment pick an arbitrary  $x \in X$ . By Bessel's inequality we have  $\|\sum_{e \in S} e \langle e, x \rangle\|^2 = \sum_{e \in S} \|\langle x, e \rangle \langle e, x \rangle\| \leq \|x\|^2$ . Thus

$$\|\langle \sum_{e \in S} e\tau(e)^*, x \rangle\| = \|\tau(\sum_{e \in S} e \langle e, x \rangle)\| \leq \|\tau\| \|x\|.$$

Substituting  $\sum_{e \in S} e\tau(e)^*$  for  $x$ , we find

$$\|\sum_{e \in S} e\tau(e)^*\|^2 = \|\langle \sum_{e \in S} e\tau(e)^*, \sum_{e \in S} e\tau(e)^* \rangle\| \leq \|\tau\| \|\sum_{e \in S} e\tau(e)^*\|.$$

If  $\|\sum_{e \in S} e\tau(e)^*\| = 0$ , then we are done. Otherwise, divide both sides of the previous equation by  $\|\sum_{e \in S} e\tau(e)^*\|$  to find  $\|\sum_{e \in S} e\tau(e)^*\| \leq \|\tau\|$  as desired.  $\square$

### 2.2.3 Self-dual completion

**150** In 136II we saw that every vector space with inner product can be completed into a Hilbert space. We continue with the generalization to Hilbert  $C^*$ -modules. Paschke proves a similar result [93, thm. 3.2] in a completely different way.

**II Theorem** Let  $\mathcal{B}$  be a von Neumann algebra and  $V$  a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product  $[\cdot, \cdot]$ . There is a self-dual Hilbert  $\mathcal{B}$ -module  $X$  together with a bounded  $\mathcal{B}$ -linear  $\eta: V \rightarrow X$  such that 1.  $[v, w] = \langle \eta(v), \eta(w) \rangle$  and 2. the image of  $\eta$  is ultranorm dense in  $X$ .

**III Proof** Before we start the proof proper, we give an overview.

**IV (Overview)** By 149V and 148V one would expect that one can simply extend the inner product of  $V$  to its ultranorm completion  $\overline{V}$  by

$$[(x_\alpha)_\alpha, (y_\alpha)_\alpha] = \text{uwlim}_\alpha [x_\alpha, y_\alpha], \quad (2.7)$$

which will turn  $\overline{V}$  into a self-dual Hilbert  $\mathcal{B}$ -module. It will turn out that  $\overline{V}$  is indeed a self-dual Hilbert  $\mathcal{B}$ -module with this inner product, but it isn't clear at all how to show directly that the ultraweak limit in (2.7) converges.

We will sketch an indirect construction, before delving into the details. As with a metric completion,  $\overline{V}$  consists of equivalence classes of ultranorm Cauchy nets in  $V$ . Let  $V_0 \subseteq \overline{V}$  denote the equivalence classes of constant Cauchy nets.  $V_0$  is a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product using the operations of  $V$ . Let  $V_1 \subseteq \overline{V}$  denote the limits of norm-bounded ultranorm Cauchy nets over  $V_0$ . Using the norm boundedness we can extend the module structure and inner product of  $V_0$  to  $V_1$ . If all norm-bounded ultranorm Cauchy-nets over  $V_1$  would converge in  $V_1$ , then  $V_1$  would be a self-dual Hilbert  $\mathcal{B}$ -module and  $V_1$  would even be ultranorm complete. Later, when we have established that  $\overline{V}$  is a Hilbert  $\mathcal{B}$ -module, we will find out that in fact  $\overline{V} \subseteq V_1$  due to 158II — i.e.  $V_1$  is already ultranorm complete. However, in a most frustrating state of affairs, we do not know how to prove this directly without first proving that  $\overline{V}$  is a Hilbert  $\mathcal{B}$ -module. Instead, we repeat: let  $V_2 \subseteq \overline{V}$  denote the limits of norm-bounded ultranorm Cauchy nets over  $V_1$ . We have to go further: for any  $n \in \mathbb{N}$  define  $V_{n+1} = \sigma(V_n)$  where  $\sigma(U) \subseteq \overline{V}$  denotes the limits of norm-bounded ultranorm Cauchy over  $U$ .

$$V_0 \subseteq V_1 \subseteq V_2 \subseteq V_3 \subseteq \cdots \subseteq \bigcup_{n \in \mathbb{N}} V_n.$$

Will we be able to show that  $\bigcup_n V_n$  is finally ultranorm complete? Not yet. We have to go even further. Define  $V_\omega = \sigma(\bigcup_{n \in \mathbb{N}} V_n)$ ,  $V_{\omega+1} = \sigma(V_\omega)$  and so on: for any ordinal number  $\alpha > 0$ , define  $V_\alpha = \sigma(\bigcup_{\beta < \alpha} V_\beta)$ . Now  $V_\alpha$  is an ascending chain of subsets of  $\overline{V}$ . Thus  $V_{\alpha_0} = V_{\alpha_0+1}$  for some  $\alpha_0 \leq |2^{\overline{V}}|$  (for otherwise  $|2^{\overline{V}}| \leq |\overline{V}|$ ). Thus norm-bounded ultranorm Cauchy nets in  $V_{\alpha_0}$  must converge. We will see that at each step we can extend the module structure and inner product and with those  $V_{\alpha_0}$  is self-dual. Thus  $V_{\alpha_0}$  is ultranorm complete and actually  $V_{\alpha_0} = \overline{V}$ .

$$V_0 \subseteq V_1 \subseteq \cdots \subseteq V_\omega \subseteq V_{\omega+1} \subseteq \cdots \subseteq V_{2\omega} \subseteq \cdots \subseteq V_{\alpha_0} = \overline{V}$$

To avoid requiring familiarity with transfinite induction, we will phrase the proof using Zorn's lemma instead.

( $\overline{V}$ : fast nets) As we need some details in its construction, we will explicitly define  $\overline{V}$ , the completion of  $V$  in the ultranorm uniformity using Cauchy nets. There are other ways to construct a completion of a uniform space, see for instance [133, thm. 39.12].

Let  $\Phi$  denote the set of entourages of the ultranorm uniformity on  $V$ ; for its definition see 146 VII.  $\Phi$  is a filter and thus can be used as index set for a net using reverse inclusion as order. (Thus  $\varepsilon \geq \delta \Leftrightarrow \varepsilon \subseteq \delta$ .) We say a net  $(x_\alpha)_{\alpha \in \Phi}$  indexed by entourages is **fast** if for every  $\varepsilon \in \Phi$  and  $\alpha, \beta \geq \varepsilon$  we have  $x_\alpha \varepsilon^2 x_\beta$ .

Every Cauchy net is equivalent to a fast one, but this is not as evident as in the metric case. Let  $(x_\alpha)_{\alpha \in I}$  be an arbitrary Cauchy net in  $V$ . By definition

we can find  $\alpha_\varepsilon \in I$  such that for each  $\varepsilon \in \Phi$  and  $\alpha, \beta \geq \alpha_\varepsilon$  we have  $x_\alpha \varepsilon x_\beta$ . Unfortunately  $\alpha_{(\cdot)}$  need not be order preserving and so  $(x_{\alpha_\varepsilon})_{\varepsilon \in \Phi}$  need not be a subnet of  $(x_\alpha)_{\alpha \in I}$ . However, we claim the net  $(x_{\alpha_\varepsilon})_{\varepsilon \in \Phi}$  is Cauchy, fast and equivalent to  $(x_\alpha)_{\alpha \in I}$ . To show it's Cauchy and fast, pick any  $\varepsilon \in \Phi$  and  $\zeta, \xi \geq \varepsilon$  (that is:  $\zeta, \xi \subseteq \varepsilon$ ). As  $I$  is directed, we can find  $\beta \in I$  with  $\beta \geq \alpha_\zeta, \alpha_\xi$ . By definition of  $\alpha_{(\cdot)}$  we have  $x_{\alpha_\zeta} \zeta x_\beta \xi x_{\alpha_\xi}$  and so  $x_{\alpha_\zeta} \varepsilon^2 x_{\alpha_\xi}$ , as desired. To show equivalence, assume  $\varepsilon \in \Phi$  is given. Assume  $\delta \geq \varepsilon/2$  and  $\beta \geq \alpha_{\varepsilon/2}$ . There is a  $\gamma \in I$  with  $\gamma \geq \alpha_\delta, \beta$ . Then  $x_\beta \varepsilon/2 x_\gamma \delta x_{\alpha_\delta}$  and so as  $\delta \subseteq \varepsilon/2$  we get  $x_\beta \varepsilon x_{\alpha_\delta}$ . We have shown equivalence.

A different fact about fast nets will be useful later on: if fast Cauchy nets  $(x_\alpha)_{\alpha \in \Phi}$  and  $(y_\alpha)_{\alpha \in \Phi}$  are equivalent, then we can find for every  $\varepsilon \in \Phi$  some  $\beta \in \Phi$  such that for all  $\gamma \geq \beta$ , we have  $x_\gamma \varepsilon y_\gamma$ .

VI ( $\bar{V}$ : the uniform space  $N$ ) Write  $N$  for the set of fast Cauchy nets over  $V$ . Later we will define  $\bar{V}$  as  $N$  modulo equivalence. Because of a subtlety with the definition of the uniformity on  $\bar{V}$  later, it is helpful to consider  $N$  separately. Let  $\varepsilon \in \Phi$  be given. For nets  $(x_\alpha)_\alpha$  and  $(y_\alpha)_\alpha$  in  $N$ , define

$$(x_\alpha)_\alpha \hat{\varepsilon} (y_\alpha)_\alpha \Leftrightarrow \exists \beta \in \Phi \forall \gamma \geq \beta. x_\gamma \varepsilon y_\gamma.$$

If  $\varepsilon \subseteq \delta$ , then  $\hat{\varepsilon} \subseteq \hat{\delta}$  and  $\hat{\varepsilon}_1 \circ \hat{\varepsilon}_2 \subseteq \widehat{\varepsilon_1 \circ \varepsilon_2}$ . So  $\widehat{\varepsilon/2} \circ \widehat{\varepsilon/2} \subseteq \widehat{\varepsilon/2 \circ \varepsilon/2} \subseteq \hat{\varepsilon}$ , which is one requirement for  $\{\hat{\varepsilon}; \varepsilon \in \Phi\}$  to be a subbase for  $N$ . The others are easy as well. Also  $\widehat{\varepsilon_1 \cap \varepsilon_2} = \hat{\varepsilon}_1 \cap \hat{\varepsilon}_2$  and so each entourage of  $N$  has some  $\hat{\varepsilon}$  as subset. For the following, it is helpful to remark now that Cauchy nets  $(x_\alpha)_\alpha$  and  $(y_\alpha)_\alpha$  over  $V$  are equivalent if and only if  $(x_\alpha)_\alpha \hat{\varepsilon} (y_\alpha)_\alpha$  for all  $\varepsilon \in \Phi$ .

VII ( $\bar{V}$ :  $N$  is complete) As every Cauchy net is equivalent to a fast one, it is sufficient to show convergence of fast Cauchy nets. Thus let  $((x_\alpha^\gamma)_\alpha)_\gamma$  be a fast Cauchy net in  $N$ . First we will show  $(x_\alpha^{\hat{\alpha}})_\alpha$  is a Cauchy net. It might not be fast, so formally  $((x_\alpha^\gamma)_\alpha)_\gamma$  cannot converge to it, but it will converge to an equivalent fast Cauchy net. As  $((x_\alpha^\gamma)_\alpha)_\gamma$  is fast, we know that for all  $\gamma_1, \gamma_2 \geq \hat{\varepsilon}$  we have  $(x_\alpha^{\gamma_1})_\alpha \hat{\varepsilon}^2 (x_\alpha^{\gamma_2})_\alpha$ . So there is a  $\zeta_{\gamma_1, \gamma_2} \in \Phi$  such that for  $\alpha \geq \zeta_{\gamma_1, \gamma_2}$  we have  $x_\alpha^{\gamma_1} \varepsilon^2 x_\alpha^{\gamma_2}$ . Suppose  $\alpha, \beta \geq \varepsilon$ . Pick  $\xi \geq \alpha, \beta, \zeta_{\hat{\alpha}, \hat{\beta}}$ . Then we have  $x_\alpha^{\hat{\alpha}} \varepsilon^2 x_\xi^{\hat{\alpha}} \varepsilon^2 x_\xi^{\hat{\beta}} \varepsilon^2 x_\beta^{\hat{\beta}}$  and so  $(x_\alpha^{\hat{\alpha}})_\alpha$  is Cauchy. Let  $(y_\alpha)_\alpha$  be a fast Cauchy net equivalent to  $(x_\alpha^{\hat{\alpha}})_\alpha$ . We want to prove  $(x_\alpha^\gamma)_\alpha \rightarrow (y_\alpha)_\alpha$ . As  $(y_\alpha)_\alpha$  is equivalent to  $(x_\alpha^{\hat{\alpha}})_\alpha$  we can find  $\alpha_0$  such that  $y_\alpha \varepsilon/4 x_\alpha^{\hat{\alpha}}$  whenever  $\alpha \geq \alpha_0$ . As  $((x_\alpha^\gamma)_\alpha)_\gamma$  is fast Cauchy, we know that  $(x_\beta^{\hat{\alpha}})_\beta \widehat{\varepsilon/4} (x_\beta^\gamma)_\beta$  if  $\hat{\alpha}, \gamma \geq \widehat{\varepsilon/8}$ . Thus there is some  $\beta_0$  such that  $x_\beta^{\hat{\alpha}} \varepsilon/4 x_\beta^\gamma$  for  $\beta \geq \beta_0$  and  $\hat{\alpha}, \gamma \geq \widehat{\varepsilon/8}$ . Thus for  $\beta \geq \beta_0 \cap \varepsilon/8$ ,  $\alpha \geq \alpha_0 \cap \varepsilon/8$  and  $\gamma \geq \widehat{\varepsilon/8}$  we get  $y_\alpha \varepsilon/4 x_\alpha^{\hat{\alpha}} \varepsilon/4 x_\beta^{\hat{\alpha}} \varepsilon/4 x_\beta^\gamma \varepsilon/4 x_\alpha^\gamma$ . Thus  $(y_\alpha)_\alpha \hat{\varepsilon} (x_\alpha^\gamma)_\alpha$  whenever  $\gamma \geq \widehat{\varepsilon/8}$ , so  $(x_\alpha^\gamma)_\alpha \rightarrow (y_\alpha)_\alpha$ . We have shown  $N$  is complete.

VIII ( $\bar{V}$ : uniformity) Let  $\bar{V}$  denote the set of fast Cauchy nets over  $V$  modulo equivalence. Write  $\eta: V \rightarrow \bar{V}$  for the map that sends  $x \in V$  to the equivalence class of the constant Cauchy net  $(x)_{\alpha \in \Phi}$ . For brevity, we may write  $\hat{x} = \eta(x)$ . As

announced before, define  $V_0 \equiv \eta(V)$ .

What uniformity to put on  $\overline{V}$ ? Unfortunately the relations  $\hat{\varepsilon}$  do not necessarily preserve equivalence ( $\sim$ ) of Cauchy nets: there might be  $(x_\alpha)_\alpha \sim (x'_\alpha)_\alpha$  and  $(y_\alpha)_\alpha \sim (y'_\alpha)_\alpha$  in  $N$  such that  $(x_\alpha)_\alpha \hat{\varepsilon} (y_\alpha)_\alpha$ , but not  $(x'_\alpha)_\alpha \hat{\varepsilon} (y'_\alpha)_\alpha$  for some  $\varepsilon \in \Phi$ . Instead we define for any  $\varepsilon \in \Phi$  the following entourages on  $\overline{V}$ :

$$(x_\alpha)_\alpha \tilde{\varepsilon} (y_\alpha)_\alpha \iff (x'_\alpha)_\alpha \hat{\varepsilon} (y'_\alpha)_\alpha \text{ for all } (x'_\alpha)_\alpha \sim (x_\alpha)_\alpha \text{ and } (y'_\alpha)_\alpha \sim (y_\alpha)_\alpha.$$

By definition  $\tilde{\varepsilon}$  respects equivalence and can be considered as a relation on  $\overline{V}$ . It is not hard to verify  $\{\tilde{\varepsilon}; \varepsilon \in \Phi\}$  is a subbase for  $\overline{V}$  and  $\varepsilon_1 \cap \varepsilon_2 = \tilde{\varepsilon}_1 \cap \tilde{\varepsilon}_2$ . It is easy to see  $\overline{V}$  is a Hausdorff uniform space. Write  $\tilde{\Phi}$  for the generated set of entourages on  $\overline{V}$ . The entourages of  $\overline{V}$  are exactly those relations which have some  $\tilde{\varepsilon}$  as a subset.

Furthermore, if  $(x_\alpha)_\alpha \hat{\varepsilon} (y_\alpha)_\alpha$ , then  $(x_\alpha)_\alpha \tilde{\varepsilon}^3 (y_\alpha)_\alpha$ . Thus if a net  $(x^\gamma_\alpha)_\alpha$  converges to  $(y_\alpha)_\alpha$  in  $N$ , then so do their equivalence classes in  $\overline{V}$ . Hence  $\overline{V}$  is complete. (The inclusion  $\hat{\varepsilon} \subseteq \tilde{\varepsilon}^3 \subseteq \tilde{\varepsilon}^3$  implies that the  $\hat{\varepsilon}$  and  $\tilde{\varepsilon}$  generate the same uniformity on  $N$ .)

( $\overline{V}$  is a right  $\mathcal{B}$ -module) First we will define an addition on  $\overline{V}$ . The addition on  $V$  is uniformly continuous (as the uniformity on  $V$  is given by seminorms, see 147 III for the uniformity on  $V^2$ .) Thus for any  $(x_\alpha)_\alpha$  and  $(y_\alpha)_\alpha$  in  $N$ , the net  $(x_\alpha + y_\alpha)_\alpha$  is again Cauchy, see 147 II. Also because of the uniform continuity of addition: if  $(x_\alpha)_\alpha \sim (x'_\alpha)_\alpha$  and  $(y_\alpha)_\alpha \sim (y'_\alpha)_\alpha$ , then  $(x_\alpha + y_\alpha)_\alpha \sim (x'_\alpha + y'_\alpha)_\alpha$ . There is a net in  $N$  equivalent to  $(x_\alpha + y_\alpha)_\alpha$ . This fixes an addition on  $\overline{V}$ , which turns it into an Abelian group. By construction  $\eta(x + y) = \eta(x) + \eta(y)$ . IX

Pick any  $b \in \mathcal{B}$ . We show the map  $r_b: V \rightarrow V$  given by  $r_b(x) = xb$  is also uniformly continuous, which requires us to unfold the definitions further than with addition. Assume we are given an entourage in  $V$ , that is: np-maps  $f_1, \dots, f_n: \mathcal{B} \rightarrow \mathbb{C}$  and  $\varepsilon > 0$ . For any np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$ , the map  $b * f$  given by  $(b * f)(x) \equiv f(b * xb)$  is also an np-map, see 72 III. Clearly  $\|xb\|_f = f([xb, xb])^{\frac{1}{2}} = f(b * [x, x]b)^{\frac{1}{2}} = \|x\|_{b * f}$  for any  $x \in V$ . Thus if  $\|x - y\|_{b * f_i} \leq \varepsilon$  for  $1 \leq i \leq n$ , then  $\|xb - yb\|_{f_i} = \|x - y\|_{b * f_i} \leq \varepsilon$  as well. Hence  $r_b$  is uniformly continuous. As before, this uniform continuity allows us to define a right  $\mathcal{B}$ -action on  $\overline{V}$  by sending the equivalence class of  $(x_\alpha)_\alpha$  to  $(x_\alpha b)_\alpha$ . By definition  $\eta(x)b = \eta(xb)$ . It is straightforward to check this turns  $\overline{V}$  into a right  $\mathcal{B}$ -module.

(Extending  $\|\cdot\|_f$  to  $\overline{V}$ ) Let  $(x_\alpha)_\alpha$  be a Cauchy net over  $V$  and  $f: \mathcal{B} \rightarrow \mathbb{C}$  be any np-map. From the reverse triangle inequality  $|\|x_\alpha\|_f - \|x_\beta\|_f| \leq \|x_\alpha - x_\beta\|_f$ , it follows  $(\|x_\alpha\|_f)_\alpha$  is Cauchy. Define  $\|(x_\alpha)_\alpha\|_f = \lim_\alpha \|x_\alpha\|_f$ . Again using the reverse triangle inequality  $|\|x_\alpha\|_f - \|x'_\alpha\|_f| \leq \|x_\alpha - x'_\alpha\|_f$  we see  $\|(x_\alpha)_\alpha\|_f = \|(x'_\alpha)_\alpha\|_f$  whenever  $(x_\alpha)_\alpha \sim (x'_\alpha)_\alpha$ . Thus  $\|\cdot\|_f$  lifts to  $\overline{V}$ . X

These extended norms  $\|\cdot\|_f$  also induce a uniformity on  $\overline{V}$ . We will now show it is the same uniformity. Let  $((x^\gamma_\alpha)_\alpha)_\gamma$  be a net in  $\overline{V}$ . It is sufficient to show that  $(x^\gamma_\alpha)_\alpha \rightarrow 0$  in the original uniformity if and only if  $\|(x^\gamma_\alpha)_\alpha\|_f \rightarrow 0$  for all np-maps  $f: \mathcal{B} \rightarrow \mathbb{C}$ .

Assume  $(x_\alpha^\gamma)_\alpha \rightarrow 0$  in  $\overline{V}$ . That is: for each  $f: \mathcal{B} \rightarrow \mathbb{C}$  and  $\varepsilon > 0$  there is a  $\gamma_0$  such that for each  $\gamma \geq \gamma_0$  there is an  $\alpha_0$  such that  $\|x_\alpha^\gamma\|_f \leq \varepsilon$  when  $\alpha \geq \alpha_0$ . Let any  $f: \mathcal{B} \rightarrow \mathbb{C}$  and  $\varepsilon > 0$  be given. Find such  $\gamma_0$  for  $f$  and  $\varepsilon$ . Then  $\|(x_\alpha^\gamma)_\alpha\|_f = \lim_\alpha \|x_\alpha^\gamma\|_f \leq \varepsilon$  whenever  $\gamma \geq \gamma_0$  and so  $\|(x_\alpha^\gamma)_\alpha\|_f \rightarrow 0$  as desired.

For the converse, assume  $\|(x_\alpha^\gamma)_\alpha\|_f \rightarrow 0$  for all np-maps  $f: \mathcal{B} \rightarrow \mathbb{C}$ . Let  $\varepsilon > 0$  and np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$  be given. There is some  $\gamma_0$  such that  $\lim_\alpha \|x_\alpha^\gamma\|_f = \|(x_\alpha^\gamma)_\alpha\|_f \leq \frac{1}{2}\varepsilon$  for  $\gamma \geq \gamma_0$ . Hence  $\|x_\alpha^\gamma\|_f \leq \varepsilon$  for sufficiently large  $\alpha$ , which is to say  $(x_\alpha^\gamma)_\alpha \rightarrow 0$  in  $\overline{V}$ .

XI (Induction set-up) The majority of the remaining work is to show we can define an inner product on  $\overline{V}$ . As sketched earlier, we will extend the inner product from  $V_0$  step-by-step. Call a subset  $W \subseteq \overline{V}$  together with a  $\mathcal{B}$ -valued inner product  $[\cdot, \cdot]$  a **compatible extension** if

1.  $V_0 \subseteq W$ ;
2.  $W$  is a  $\mathcal{B}$ -submodule of  $\overline{V}$ ;
3. the inner product turns  $W$  into a pre-Hilbert  $\mathcal{B}$ -module *and*
4.  $\|x\|_f^{\overline{V}} = \|x\|_f^W$  for all  $x \in W$  and np-maps  $f: \mathcal{B} \rightarrow \mathbb{C}$ , where  $\|x\|_f^W = f([x, x]_W)^{\frac{1}{2}}$  and  $\|\cdot\|_f^{\overline{V}}$  is the extension of  $\|\cdot\|_f$  on  $V$  to  $\overline{V}$  as in X.

As  $\|\cdot\|_f^{\overline{V}}$  also generate the uniformity on  $\overline{V}$ , the last requirement implies that on a compatible extension  $W$  the ultranorm uniformity and the uniformity induced by  $\overline{V}$  coincide.

For compatible extensions  $W_1, W_2 \subseteq \mathcal{V}$  we say  $W_1 \leq W_2$  iff  $W_1 \subseteq W_2$  and  $[v, w]_{W_1} = [v, w]_{W_2}$  for all  $v, w \in W_1$ . Later, we will apply Zorn's lemma to the compatible extensions ordered in this way.

XII (Induction base case) The inner product of  $V$  lifts to  $V_0$ . Indeed,  $\eta(x) = 0$  if and only if  $\|x\| = 0$ . So by Cauchy-Schwarz  $[x, y] = 0$  whenever  $\eta(x) = 0$ . Thus  $[x, y] = [x', y']$  if  $\eta(x) = \eta(x')$  and  $\eta(y) = \eta(y')$ . Define  $[\eta(x), \eta(y)] = [x, y]$ . It's easy to see  $V_0$  is a pre-Hilbert  $\mathcal{B}$ -submodule of  $\overline{V}$ . Finally, for any np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$  we have  $\|\eta(x)\|_f^{\overline{V}} = \|x\|_f^V = \|\eta(x)\|_f^{V_0}$ , as desired.

XIII (Induction step) Assume  $W \subseteq \overline{V}$  with  $[\cdot, \cdot]$  is a compatible extension. Let  $\sigma(W)$  denote the limits of norm-bounded Cauchy nets over  $W$ . We will define a  $\mathcal{B}$ -valued inner product on  $\sigma(W)$  that turns it into a compatible extension. In fact  $W \leq \sigma(W)$ . By assumption the induced uniformity on  $W$  is the ultranorm uniformity and so addition and the  $\mathcal{B}$ -action are uniformly continuous. Hence  $\sigma(W)$  is again a submodule.

To define the inner product, assume  $x, y \in \sigma(W)$ . There are  $\Phi$ -indexed norm-bounded Cauchy nets  $(x_\alpha)_\alpha$  and  $(y_\alpha)_\alpha$  over  $W$  such that  $x_\alpha \rightarrow x$  and  $y_\alpha \rightarrow y$ .

For any np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$  we have by Cauchy–Schwarz

$$\begin{aligned} |f([x_\alpha, y_\alpha] - [x_\beta, y_\beta])| &= |[x_\alpha, y_\alpha - y_\beta]_f + [x_\alpha - x_\beta, y_\beta]_f| \\ &\leq \|x_\alpha\|_f \|y_\alpha - y_\beta\|_f + \|x_\alpha - x_\beta\|_f \|y_\beta\|_f. \end{aligned} \quad (2.8)$$

(If not otherwise specified  $\|\cdot\|_f = \|\cdot\|_f^W$ .) As  $(x_\alpha)_\alpha$  is norm bounded and  $\|x_\alpha\|_f^2 = f([x_\alpha, x_\alpha]) \leq \|f\| \|x_\alpha\|^2$ , we see  $(\|x_\alpha\|_f)_\alpha$  is bounded. Similarly  $(\|y_\beta\|_f)_\beta$  is bounded. Thus from (2.8) it follows  $([x_\alpha, y_\alpha])_\alpha$  is a Cauchy net in the ultraweak uniformity of  $\mathcal{B}$ , see 146 V. As  $\|[x_\alpha, y_\alpha]\| \leq \|x_\alpha\| \|y_\alpha\|$ , see 142 III, it is also a norm-bounded net. By 77 I norm-bounded Cauchy nets in the ultraweak uniformity converge, so  $\text{uwlim}_\alpha [x_\alpha, y_\alpha]$  exists.

Assume we are given any  $\Phi$ -indexed norm-bounded Cauchy nets  $(x')_\alpha$  and  $(y')_\alpha$  over  $W$  with  $(x'_\alpha)_\alpha \rightarrow x$  and  $(y'_\alpha)_\alpha \rightarrow y$ . Then, like (2.8):

$$\begin{aligned} |f([x_\alpha, y_\alpha] - [x'_\alpha, y'_\alpha])| &= |[x_\alpha, y_\alpha - y'_\alpha]_f + [x_\alpha - x'_\alpha, y'_\alpha]_f| \\ &\leq \|x_\alpha\|_f \|y_\alpha - y'_\alpha\|_f + \|x_\alpha - x'_\alpha\|_f \|y'_\alpha\|_f. \end{aligned}$$

From this it follows  $\text{uwlim}_\alpha [x_\alpha, y_\alpha] = \text{uwlim}_\alpha [x'_\alpha, y'_\alpha]$ . We are justified to define

$$[x, y] \equiv \text{uwlim}_\alpha [x_\alpha, y_\alpha].$$

We will show  $[\cdot, \cdot]$  is a  $\mathcal{B}$ -valued inner product. Pick any  $x, y, z \in \sigma(W)$  and  $\Phi$ -indexed norm-bounded Cauchy nets  $(x_\alpha)_\alpha$ ,  $(y_\alpha)_\alpha$  and  $(z_\alpha)_\alpha$  over  $W$  with  $x_\alpha \rightarrow x$ ,  $y_\alpha \rightarrow y$  and  $z_\alpha \rightarrow z$  in  $\overline{V}$ . So  $x_\alpha + y_\alpha \rightarrow x + y$ . With ultraweak continuity of addition

$$[x + y, z] = \text{uwlim}_\alpha [x_\alpha + y_\alpha, z_\alpha] = \text{uwlim}_\alpha [x_\alpha, z_\alpha] + \text{uwlim}_\alpha [y_\alpha, z_\alpha] = [x, z] + [y, z].$$

In a similar fashion one proves the other axioms of a  $\mathcal{B}$ -valued inner product:  $[x, y] = [y, x]^*$  follows from the ultraweak continuity of  $(\cdot)^*$ ;  $\mathcal{B}$ -homogeneity follows from ultraweak continuity of  $c \mapsto cb$  (see 45 IV) and  $[x, x] \geq 0$  follows from ultraweak-closedness of the positive cone of  $\mathcal{B}$ .

To prove definiteness of  $[\cdot, \cdot]$ , assume  $[x, x] = 0$  for some  $x \in \sigma(W)$ . Pick some  $\Phi$ -indexed norm-bounded Cauchy net  $(x_\alpha)_\alpha$  over  $W$  with  $x_\alpha \rightarrow x$ . For any np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$  we have  $\lim_\alpha f([x_\alpha, x_\alpha]) = f(\text{uwlim}_\alpha [x_\alpha, x_\alpha]) = f([x, x]) = 0$ . Thus  $x_\alpha$  converges in the ultranorm uniformity on  $W$  to 0. Hence  $x = 0$ .

Thus  $\sigma(W)$  is a pre-Hilbert  $\mathcal{B}$ -module. It remains to be shown that  $\|\cdot\|_f^{\sigma(W)}$  and  $\|\cdot\|_f^{\overline{V}}$  agree. Pick any  $x \in \sigma(W)$  and  $f: \mathcal{B} \rightarrow \mathbb{C}$ . By definition there is a norm-bounded Cauchy net  $x_\alpha$  over  $W$  for which  $x_\alpha \rightarrow x$  in  $\overline{V}$ . Then

$$\|x\|_f^{\sigma(W)} = f(\text{uwlim}_\alpha [x_\alpha, x_\alpha])^{\frac{1}{2}} = \lim_\alpha \|x_\alpha\|_f^W = \lim_\alpha \|x_\alpha\|_f^{\overline{V}} = \|x\|_f^{\overline{V}},$$

as desired. We have shown that  $\sigma(W)$  is a compatible extension.

XIV (Induction limit step) Assume  $\mathcal{V}$  is a non-empty chain of compatible extensions. See XI for the order on compatible extensions. Write  $W = \bigcup \mathcal{V}$ . We will turn  $W$  into a compatible extension. Clearly  $W$  is a submodule of  $\overline{V}$ . As  $\mathcal{V}$  is not empty,  $V_0 \subseteq W$ . For any  $v, w \in W$  there is some  $W' \in \mathcal{V}$  with  $v, w \in W'$ . Define  $[v, w]_W = [v, w]_{W'}$ . This turns  $W$  into a pre-Hilbert  $\mathcal{B}$ -module as all its axioms only involve finitely many elements, which are contained in some single compatible extension far enough up in the chain. Finally, to see the seminorms agree, pick any np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$  and  $x \in W$ . Then  $x \in W'$  for some  $W' \in \mathcal{V}$ . Consequently

$$\|x\|_f^{\overline{V}} = \|x\|_f^{W'} = f([x, x]_{W'})^{\frac{1}{2}} = f([x, x]_W)^{\frac{1}{2}} = \|x\|_f^W,$$

as desired. We have shown  $W$  is a compatible extension.

XV (Self-duality) We have seen that every non-empty chain of compatible extensions has an upper bound. Thus by Zorn's lemma there is a maximal compatible extension  $W \subseteq \overline{V}$ . By maximality  $\sigma(W) = W$  and so in  $W$  norm-bounded ultranorm-Cauchy nets converge. Thus by 149V  $W$  must be self-dual and ultranorm complete. By completeness  $W = \overline{V}$ . Define  $X = \overline{V}$  and  $\langle x, y \rangle = [x, y]_W$ . Clearly  $[v, w] = \langle \eta(v), \eta(w) \rangle$  and the image of  $\eta$  is norm dense.  $\square$

**151** The completion 150II we just constructed has a universal property, which can be seen as a generalization of [93, prop. 3.6].

la **Lemma** Assume  $\mathcal{B}$  is a von Neumann algebra and  $V$  is a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product. Let  $X$  be any self-dual Hilbert  $\mathcal{B}$ -module together with inner product preserving  $\mathcal{B}$ -linear map  $\eta: V \rightarrow X$  such that the image of  $\eta$  is ultranorm dense in  $X$ . (E.g. the completion into a self-dual Hilbert  $\mathcal{B}$ -module from 150II.) Then  $\eta$  has the following universal property.

Let  $Y$  be a self-dual Hilbert  $\mathcal{B}$ -module. For each bounded  $\mathcal{B}$ -linear map  $T: V \rightarrow Y$ , there is a unique bounded  $\mathcal{B}$ -linear map  $\hat{T}: X \rightarrow Y$  such that  $\hat{T} \circ \eta = T$ .

II **Proof** By 148I bounded  $\mathcal{B}$ -linear maps are ultranorm uniformly continuous. As  $\eta(V)$  is ultranorm dense in  $X$ , the extension  $\hat{T}$  is unique if it exists. Pick any  $x \in X$ . There is a net  $x_\alpha$  in  $V$  such that  $\eta(x_\alpha) \rightarrow x$  ultranorm. So  $\eta(x_\alpha)$  is ultranorm Cauchy. As  $\eta$  preserves inner product,  $x_\alpha$  is ultranorm Cauchy as well. As  $T$  is ultranorm uniformly continuous  $T(x_\alpha)$  is ultranorm Cauchy. With the same motions we see  $T(x_\alpha) \sim T(x'_\alpha)$  when  $x_\alpha \sim x'_\alpha$ . Using this and that  $Y$  is ultranorm complete, see 149V, we may define  $\hat{T}x = \text{unlim}_\alpha T(x_\alpha)$ , where **unlim** denotes the limit in the ultranorm uniformity. Clearly  $\hat{T} \circ \eta = T$ . It is straightforward to check that  $\mathcal{B}$ -linearity of  $\hat{T}$  follows from  $\mathcal{B}$ -linearity of  $T$  and ultranorm continuity of addition and the  $\mathcal{B}$ -module action.

It remains to be shown  $\hat{T}$  is bounded. By 144V we have

$$\langle \hat{T}x_\alpha, \hat{T}x_\alpha \rangle \leq \|T\|^2 [x_\alpha, x_\alpha] = \|T\|^2 \langle \eta(x_\alpha), \eta(x_\alpha) \rangle$$

and so using 148V we see that

$$\langle \hat{T}x, \hat{T}x \rangle = \text{uwlim}_\alpha \langle Tx_\alpha, Tx_\alpha \rangle \leq \|T\|^2 \text{uwlim}_\alpha \langle \eta(x_\alpha), \eta(x_\alpha) \rangle = \|T\|^2 \langle x, x \rangle,$$

which implies that  $\|\hat{T}\| \leq \|T\|$  as desired. (In fact  $\|\hat{T}\| = \|T\|$ .) □

Before we show that every ncp-map has a Paschke dilation, we study  $\mathcal{B}^a(X)$  a bit more for self-dual  $X$ . We start with  $\mathcal{B}$ -sesquilinear forms. 152

**Definition** Let  $V$  be a normed right  $\mathcal{B}$ -module. A sesquilinear form  $B: V \times V \rightarrow \mathcal{B}$  is said to be **bounded** if there is an  $r \geq 0$  such that  $\|B(x, y)\| \leq r\|x\|\|y\|$  for all  $x, y \in V$ . II

**Example** Let  $X$  be a pre-Hilbert  $\mathcal{B}$ -module. For every  $T \in \mathcal{B}^a(X)$  the map  $\langle (\cdot), T(\cdot) \rangle$  is a bounded  $\mathcal{B}$ -sesquilinear form. III

For self-dual  $X$  every bounded  $\mathcal{B}$ -sesquilinear form arises in this way. IV

**Proposition** Let  $X$  be a self-dual Hilbert  $\mathcal{B}$ -module. For every bounded  $\mathcal{B}$ -sesquilinear form  $B$  on  $X$ , there is a unique  $T \in \mathcal{B}^a(X)$  with  $B(x, y) = \langle x, Ty \rangle$  for all  $x, y \in X$ . V

**Proof** For each  $y \in X$  the map  $B(\cdot, y)^*$  is  $\mathcal{B}$ -linear and bounded. Thus by self-duality of  $X$  there is a unique  $t_y$  for each  $y \in X$  such that  $\langle t_y, x \rangle = B(x, y)^*$  for all  $x \in X$ . It is easy to see  $\langle t_{y+y'}, x \rangle = \langle t_y + t_{y'}, x \rangle$  and  $\langle t_{yb}, x \rangle = \langle t_y b, x \rangle$ , and so by uniqueness  $y \mapsto t_y$  is  $\mathcal{B}$ -linear. Define  $Ty = t_y$ . Clearly  $\langle x, Ty \rangle = \langle t_y, x \rangle^* = B(x, y)$  and  $T$  must be the unique such  $\mathcal{B}$ -linear map. Reasoning in the same way, we see there is a unique  $S$  with  $B(y, x)^* = \langle x, Sy \rangle$ . Hence  $\langle x, Ty \rangle = B(x, y) = \langle y, Sx \rangle^* = \langle Sx, y \rangle$ , so  $T$  is adjointable. It remains to be shown  $T$  is bounded. There is an  $r \geq 0$  such that for all  $x \in X$ : VI

$$\|Tx\|^2 = \|\langle Tx, Tx \rangle\| = \|B(Tx, x)\| \leq r\|Tx\|\|x\|.$$

So  $r$  is a bound by dividing out  $\|Tx\|$  if  $\|Tx\| \neq 0$  and trivially otherwise. □

**Remark** In 36V a more general result was shown. VII

**Exercise** Suppose  $T: X \rightarrow Y$  is a bounded  $\mathcal{B}$ -linear map between Hilbert  $\mathcal{B}$ -modules. Show that if  $X$  is self dual, then  $T$  is adjointable. [93, prop. 3.4] VIII

**Exercise** Let  $V$  be a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product for some von Neumann algebra  $\mathcal{B}$  and  $\eta: V \rightarrow X$  be the ultranorm completion of  $V$  from 150II. Show that IX

$$\{ \langle \hat{x}, (\cdot) \hat{x} \rangle: X \rightarrow \mathcal{B}; x \in V \} \quad (\text{where } \hat{x} \equiv \eta(x))$$

is an order separating set of ncp-maps, see 21II. (That is:  $T \geq 0$  iff  $\langle \hat{x}, T\hat{x} \rangle \geq 0$  for all  $x \in V$ . Consequently  $S = T$  iff  $\langle \hat{x}, T\hat{x} \rangle = \langle \hat{x}, S\hat{x} \rangle$  for all  $x \in V$ .)

- X **Theorem** Suppose  $X$  is a self-dual Hilbert  $\mathcal{B}$ -module for a von Neumann algebra  $\mathcal{B}$ . Then  $\mathcal{B}^a(X)$  is a von Neumann algebra.
- XI **Proof** The theorem is due to Paschke [93, prop. 3.10]; we give a new proof, which also appeared as 49 II. We already know that  $\mathcal{B}^a(X)$  is a  $C^*$ -algebra, see 143 IV.
- XII (bounded order completeness) Let  $(T_\alpha)_\alpha$  be a norm-bounded net of self-adjoint elements of  $\mathcal{B}^a(X)$ . We have to show it has a supremum. Pick  $x \in X$  and  $r \in \mathbb{R}$ ,  $r \geq 0$  such that  $\|T_\alpha\| \leq r$  for all  $\alpha$ . By 144 I  $(\langle x, T_\alpha x \rangle)_\alpha$  is a norm-bounded net of self-adjoint elements of  $\mathcal{B}$  and so it has a supremum to which it converges ultrastrongly by 44 XIV. In particular

$$\frac{1}{4} \sum_{k=0}^3 i^k \langle i^k x + y, T_\alpha(i^k x + y) \rangle = \langle x, T_\alpha y \rangle$$

converges ultrastrongly for all  $x, y \in X$ . Define  $B(x, y) = \text{uslim}_\alpha \langle x, T_\alpha y \rangle$ . As addition and multiplication by a fixed element are ultrastrongly continuous (see 45 IV), we see that  $B$  is a  $\mathcal{B}$ -sesquilinear form. By 142 III and 144 V we have  $\langle T_\alpha y, x \rangle \langle x, T_\alpha y \rangle \leq r^2 \|x\|^2 \langle y, y \rangle$  for each  $\alpha$ , hence

$$\begin{aligned} \|B(x, y)\|^2 &= \|B(x, y)^* B(x, y)\| \\ &\stackrel{46 \text{ II}}{=} \|\text{uwlim}_\alpha \langle T_\alpha y, x \rangle \langle x, T_\alpha y \rangle\| \\ &\leq r^2 \|x\|^2 \|y\|^2. \end{aligned}$$

So  $B$  is a bounded  $\mathcal{B}$ -sesquilinear form. By V there is a unique  $T \in \mathcal{B}^a(X)$  with  $\langle x, Ty \rangle = B(x, y) = \text{uslim}_\alpha \langle x, T_\alpha y \rangle$ . Noting  $B(x, y) = \text{uwlim}_\alpha \langle x, T_\alpha y \rangle$ , it is easy to see  $T$  is self-adjoint. Using 144 I and the fact that ultraweak limits respect the order (44 XI), we see  $\langle x, T_\alpha x \rangle \leq \langle x, Tx \rangle$  for all  $x \in X$  and so  $T_\alpha \leq T$ . Now suppose we are given a self-adjoint  $S \in \mathcal{B}^a(X)$  with  $T_\alpha \leq S$  for all  $\alpha$ . With a similar argument we see  $T \leq S$  and so  $S$  is the supremum of  $(T_\alpha)_\alpha$ .

- XIII (separating normal states) By 144 I, the states  $\langle x, (\cdot)x \rangle$  are separating. We are done if we can show that  $\langle x, (\cdot)x \rangle$  is normal, i.e. preserves suprema of bounded directed sets of self-adjoint elements. To this end, pick  $x \in X$  and let  $(T_\alpha)_\alpha$  be a net with supremum  $T$ . As before  $(\langle x, T_\alpha x \rangle)_\alpha$  is a norm bounded net which converges ultrastrongly to its supremum. Now, we just saw

$$\langle x, Tx \rangle = \text{uwlim}_\alpha \langle x, T_\alpha x \rangle = \sup_\alpha \langle x, T_\alpha x \rangle$$

and so indeed  $\langle x, (\cdot)x \rangle$  is normal; the normal states are separating and consequently  $\mathcal{B}^a(X)$  is a von Neumann algebra.  $\square$

- 153 **Proposition** Assume  $T: X \rightarrow Y$  is an adjointable bounded module map between Hilbert  $\mathcal{B}$ -modules. Define  $\text{ad}_T: \mathcal{B}^a(Y) \rightarrow \mathcal{B}^a(X)$  by  $\text{ad}_T(B) = T^*BT$ . The map  $\text{ad}_T$  is completely positive. If  $X$  and  $Y$  are self-dual, then  $\text{ad}_T$  is normal.

**Proof** For any  $n \in \mathbb{N}$ ,  $B_1, \dots, B_n \in \mathcal{B}^a(X)$  and  $A_1, \dots, A_n \in \mathcal{B}^a(Y)$  we have ||

$$\sum_{i,j} B_j^* T^* A_j^* A_i T B_i = \left( \sum_i A_i T B_i \right)^* \left( \sum_j A_j T B_j \right) \geq 0$$

and so  $\text{ad}_T$  is completely positive.

Now we show  $\text{ad}_S$  is normal. By 48|| and 144|, it suffices to show that for every  $x \in X$ , the map  $T \mapsto \langle x, \text{ad}_S(T)x \rangle = \langle Sx, T Sx \rangle$  is normal, which it indeed is by 152XIII. ||| □

**Exercise** Let  $\mathcal{A}$  be a C\*-algebra. Assume  $n \in \mathbb{N}$  and  $a_1, \dots, a_n \in \mathcal{A}$ . Use | to show  $\varphi: \mathcal{A} \rightarrow M_n \mathcal{A}$  given by  $\varphi(d) = (a_i^* d a_j)_{ij}$  is an ncp-map. IV

## 2.3 Paschke dilations

We are ready to show that every ncp-map has a Paschke dilation. That is: 154  
for every ncp-map  $\varphi$  there is a triple  $(\mathcal{P}, \varrho, h)$  with  $\varphi = h \circ \varrho$  via  $\mathcal{P}$  for some ncp-map  $h$ , nmiu-map  $\varrho$  and von Neumann algebra  $\mathcal{P}$ , which is minimal in the sense that for any such triple  $(\mathcal{P}', \varrho', h')$  there is a unique ncp-map  $\sigma: \mathcal{P}' \rightarrow \mathcal{P}$  with  $h' = h \circ \sigma$  and  $\varrho = \sigma \circ \varrho'$ . See 140||.

**Definition** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  be any ncp-map between von Neumann algebras. A complex bilinear map  $B: \mathcal{A} \times \mathcal{B} \rightarrow X$ , where  $X$  is a self-dual Hilbert  $\mathcal{B}$ -module is called  $\varphi$ -compatible if there is an  $r > 0$  such that for all  $n \in \mathbb{N}$ ,  $a_1, \dots, a_n \in \mathcal{A}$  and  $b_1, \dots, b_n \in \mathcal{B}$  we have ||

$$\left\| \sum_i B(a_i, b_i) \right\|^2 \leq r \cdot \left\| \sum_{i,j} b_i^* \varphi(a_i^* a_j) b_j \right\| \quad (2.9)$$

and  $B(a, b_1) b_2 = B(a, b_1 b_2)$  for all  $a \in \mathcal{A}$  and  $b_1, b_2 \in \mathcal{B}$ .

**Theorem** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  be any ncp-map between von Neumann algebras. |||

1. There is a self-dual Hilbert  $\mathcal{B}$ -module  $\mathcal{A} \otimes_\varphi \mathcal{B}$  and  $\varphi$ -compatible bilinear

$$\otimes: \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{A} \otimes_\varphi \mathcal{B}$$

such that for every  $\varphi$ -compatible bilinear map  $B: \mathcal{A} \times \mathcal{B} \rightarrow Y$  there is a unique bounded module map  $T: \mathcal{A} \otimes_\varphi \mathcal{B} \rightarrow Y$  such that  $T(a \otimes b) = B(a, b)$  for all  $a \in \mathcal{A}$  and  $b \in \mathcal{B}$ .

2. For  $a_0 \in \mathcal{A}$ , there is a unique  $\varrho(a_0) \in \mathcal{B}^a(\mathcal{A} \otimes_\varphi \mathcal{B})$  fixed by

$$\varrho(a_0)(a \otimes b) = (a_0 a) \otimes b. \quad (a \in \mathcal{A}, b \in \mathcal{B})$$

Furthermore  $a \mapsto \varrho(a)$  yields a nmiu-map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}^a(\mathcal{A} \otimes_\varphi \mathcal{B})$ .

3. The map  $h: \mathcal{B}^a(\mathcal{A} \otimes_\varphi \mathcal{B}) \rightarrow \mathcal{B}$  given by  $h(T) = \langle 1 \otimes 1, T(1 \otimes 1) \rangle$  is ncp.
4. The map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}^a(\mathcal{A} \otimes_\varphi \mathcal{B})$  together with  $1 \otimes 1 \in \mathcal{A} \otimes_\varphi \mathcal{B}$  has the following universal property.

Let  $\varrho': \mathcal{A} \rightarrow \mathcal{B}^a(X)$  be an nmiu-map for a self-dual Hilbert  $\mathcal{B}$ -module  $X$  together with an element  $e \in X$  such that  $\varphi = h' \circ \varrho'$  for  $h'(T) \equiv \langle e, Te \rangle$ . Then: there is a unique inner product preserving  $\mathcal{B}$ -linear map  $S: \mathcal{A} \otimes_\varphi \mathcal{B} \rightarrow X$  such that  $\text{ad}_S \circ \varrho' = \varrho$  and  $S(1 \otimes 1) = e$ .

5.  $(\mathcal{B}^a(\mathcal{A} \otimes_\varphi \mathcal{B}), \varrho, h)$  is a Paschke dilation of  $\varphi$ , see 140 II.

- IV Proof** (This is a simplified version of the proof we published earlier in [127]. The construction of  $\mathcal{A} \otimes_\varphi \mathcal{B}$  is essentially due to Paschke [93, thm. 5.2]: the self-dual completion of  $X$  in [93, thm. 5.2] is isomorphic to  $\mathcal{A} \otimes_\varphi \mathcal{B}$ .)
- V** (1:  $\mathcal{A} \odot_\varphi \mathcal{B}$ ) The algebraic tensor product  $\mathcal{A} \odot \mathcal{B}$  is a right  $\mathcal{B}$ -module with the action  $(\sum_i a_i \otimes b_i)\beta = \sum_i a_i \otimes (b_i\beta)$ . On  $\mathcal{A} \odot \mathcal{B}$ , define

$$\left[ \sum_i a_i \otimes b_i, \sum_j \alpha_j \otimes \beta_j \right] \equiv \sum_{i,j} b_i^* \varphi(a_i^* \alpha_j) \beta_j.$$

By complete positivity of  $\varphi$  this is a  $\mathcal{B}$ -valued inner product on  $\mathcal{A} \odot \mathcal{B}$ . By 150 II, there is a self-dual Hilbert  $\mathcal{B}$ -module  $\mathcal{A} \otimes_\varphi \mathcal{B}$  and  $\mathcal{B}$ -linear inner product-preserving  $\eta: \mathcal{A} \odot \mathcal{B} \rightarrow \mathcal{A} \otimes_\varphi \mathcal{B}$  with ultranorm-dense range. Define a bilinear map  $\otimes: \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{A} \otimes_\varphi \mathcal{B}$  by  $a \otimes b = \eta(a \odot b)$ . By definition we have

$$\left\| \sum_i a_i \otimes b_i \right\|^2 = \left\| \left[ \sum_i a_i \otimes b_i, \sum_j a_j \otimes b_j \right] \right\| = \left\| \sum_{i,j} b_i^* \varphi(a_i^* a_j) b_j \right\|$$

and so  $\otimes$  is a  $\varphi$ -compatible bilinear map.

Let  $B: \mathcal{A} \times \mathcal{B} \rightarrow Y$  be a  $\varphi$ -compatible bilinear map to some self-dual Hilbert  $\mathcal{B}$ -module  $Y$ . We must show that there is a unique bounded module map  $T: \mathcal{A} \otimes_\varphi \mathcal{B} \rightarrow Y$  such that  $T(a \otimes b) = B(a, b)$  for all  $a \in \mathcal{A}$  and  $b \in \mathcal{B}$ . By the defining property of the algebraic tensor product, there is a unique linear map  $T_0: \mathcal{A} \odot \mathcal{B} \rightarrow Y$  such that  $T_0(a \odot b) = B(a, b)$  for all  $a \in \mathcal{A}$  and  $b \in \mathcal{B}$ . By definition of  $\varphi$ -compatibility and the inner product on  $\mathcal{A} \odot \mathcal{B}$ , the map  $T_0$  is in fact bounded and  $\mathcal{B}$ -linear. By 151 Ia this map extends uniquely to a bounded module map  $T: \mathcal{A} \otimes_\varphi \mathcal{B} \rightarrow Y$  with  $T(a \otimes b) = B(a, b)$ , as desired.

- VI** (2:  $\varrho: \mathcal{A} \rightarrow \mathcal{B}^a(\mathcal{A} \otimes_\varphi \mathcal{B})$ ) Let  $a_0 \in \mathcal{A}$  be given. To show the module map  $\varrho(a_0)$  exists, it suffices to show that the bilinear map  $B: \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{A} \otimes_\varphi \mathcal{B}$  given by  $B(a, b) = (a_0 a) \otimes b$  is  $\varphi$ -compatible. Clearly  $B(a, b\beta) = B(a, b)\beta$ . To prove (2.9), let  $n \in \mathbb{N}$ ,  $a_1, \dots, a_n \in \mathcal{A}$  and  $b_1, \dots, b_n \in \mathcal{B}$  be given. The row vector  $(a_1 \ \cdots \ a_n)$  is an  $\mathcal{A}$ -linear map  $a: \mathcal{A}^{\oplus n} \rightarrow \mathcal{A}$  in the usual

way:  $a(\alpha_1, \dots, \alpha_n) = a_1\alpha_1 + \dots + a_n\alpha_n$ . Similarly the column vector  $(b_1 \dots b_n)^t$  is a  $\mathcal{B}$ -linear map  $b: \mathcal{B} \rightarrow \mathcal{B}^{\oplus n}$  with  $b(\beta) = (b_1\beta, \dots, b_n\beta)$ . We compute

$$\begin{aligned} \left\| \sum_i B(a_i, b_i) \right\|^2 &= \left\| \sum_i (a_0 a_i) \otimes b_i \right\|^2 \\ &= \left\| \sum_{i,j} b_i^* \varphi(a_i^* a_0^* a_0 a_j) b_j \right\| \\ &= \|b^*(M_n \varphi)(a^* a_0^* a_0 a) b\| \\ &\leq \|a_0^* a_0\| \|b^*(M_n \varphi)(a^* a) b\| \\ &= \|a_0\|^2 \left\| \sum_{i,j} b_i^* \varphi(a_i^* a_j) b_j \right\|. \end{aligned}$$

Thus  $B$  is  $\varphi$ -compatible and so there is a unique  $\mathcal{B}$ -linear bounded module map  $\varrho(a_0): \mathcal{A} \otimes_{\varphi} \mathcal{B} \rightarrow \mathcal{A} \otimes_{\varphi} \mathcal{B}$  with  $\varrho(a_0)(a \otimes b) = (a_0 a) \otimes b$ . Clearly  $\varrho$  is a unital, multiplicative and involution preserving map. It remains to be shown  $\varrho$  is normal. By 48II and 152IX, it suffices to show that  $d \mapsto \langle \hat{x}, \varrho(d)\hat{x} \rangle$  is normal for every  $x \in \mathcal{A} \odot \mathcal{B}$ . Find  $n \in \mathbb{N}$ , row vector  $a \equiv (a_1 \dots a_n)$  and column vector  $b \equiv (b_1 \dots b_n)$  such that  $x = \sum_{i=1}^n a_i \otimes b_i$ . We compute

$$\langle \hat{x}, \varrho(d)\hat{x} \rangle = \sum_{i,j} b_i^* \varphi(a_i^*(d) a_j) b_j = b^*(M_n \varphi)(a^* d a) b.$$

So  $d \mapsto \langle \hat{x}, \varrho(d)\hat{x} \rangle$  is normal by 153I and 49IV, whence  $\varrho$  is normal.

(3:  $h: \mathcal{B}^a(\mathcal{A} \otimes_{\varphi} \mathcal{B}) \rightarrow \mathcal{B}$ ) Define  $h(T) \equiv \langle 1 \otimes 1, T 1 \otimes 1 \rangle$ . It is completely VII  
positive by 145I and normal by 152XIII.

(Uniqueness  $\sigma$ ) Before we continue with 4, we will already prove the uniqueness VIII  
property for point 5. Note that  $(h \circ \varrho)(a) = \langle 1 \otimes 1, a \otimes 1 \rangle = \varphi(a)$  for all  $a \in \mathcal{A}$  and so  $\varphi = h \circ \varrho$ . Assume  $\varphi = h' \circ \varrho'$  for some nmiu-map  $\varrho': \mathcal{A} \rightarrow \mathcal{P}'$ , ncp-map  $h: \mathcal{P}' \rightarrow \mathcal{B}$  and von Neumann algebra  $\mathcal{P}'$ . For point 5 we must show there is a unique ncp-map  $\sigma: \mathcal{P}' \rightarrow \mathcal{B}^a(\mathcal{A} \otimes_{\varphi} \mathcal{B})$  such that  $h \circ \sigma = h'$  and  $\sigma \circ \varrho' = \varrho$ . Let  $\sigma_1, \sigma_2: \mathcal{P}' \rightarrow \mathcal{B}^a(\mathcal{A} \otimes_{\varphi} \mathcal{B})$  be ncp-maps with  $h \circ \sigma_k = h'$  and  $\sigma_k \circ \varrho' = \varrho$ ,  $k = 1, 2$ . We must show  $\sigma_1 = \sigma_2$ . Let  $c \in \mathcal{P}'$  and  $x \in \mathcal{A} \odot \mathcal{B}$  be given. By 152IX it suffices to prove that  $\langle \hat{x}, \sigma_1(c)\hat{x} \rangle = \langle \hat{x}, \sigma_2(c)\hat{x} \rangle$ . Find  $n \in \mathbb{N}$ ,  $a_1, \dots, a_n \in \mathcal{A}$  and  $b_1, \dots, b_n \in \mathcal{B}$  such that  $x = \sum_i a_i \otimes b_i$ . Note  $a_i \otimes b_i = \varrho(a_i)(1 \otimes 1)b_i$ . So for any  $k = 1, 2$ , we find

$$\begin{aligned} \langle \hat{x}, \sigma_k(c)\hat{x} \rangle &= \sum_{i,j} b_i^* h(\varrho(a_i^*) \sigma_k(c) \varrho(a_j)) b_j \\ &= \sum_{i,j} b_i^* h(\sigma_k(\varrho'(a_i^*) c \varrho'(a_j))) b_j \quad \text{by 139III} \\ &= \sum_{i,j} b_i^* h'(\varrho'(a_i^*) c \varrho'(a_j)) b_j. \end{aligned} \tag{2.10}$$

Thus  $\sigma_1 = \sigma_2$  as desired. One can show such  $\sigma$  exists by using (2.10) as defining formula. We will use, however, an indirect but shorter approach by first considering the ‘spatial case’, which was suggested by Michael Skeide.

- IX (4: spatial case) Let  $X$  be a self-dual Hilbert  $\mathcal{B}$ -module together with  $e \in X$  and nmiu-map  $\mathcal{A} \rightarrow \mathcal{B}^a(X)$  such that  $\varphi = h' \circ \varrho'$  with  $h'(T) = \langle e, Te \rangle$ . To prove uniqueness, assume that for  $k = 1, 2$ , we have  $\mathcal{B}$ -linear inner product preserving  $S_k: \mathcal{A} \otimes_{\varphi} \mathcal{B} \rightarrow X$  with  $\text{ad}_{S_k} \circ \varrho' = \varrho$  and  $S_k 1 \otimes 1 = e$ . Then  $h \circ \text{ad}_{S_k} = h'$  and so (as  $\text{ad}_{S_k}$  is an ncp-map by 153I), we know  $\text{ad}_{S_1} = \text{ad}_{S_2}$  by VIII. Hence there is a  $\lambda \in \mathbb{C}, \lambda \neq 0$  with  $S_1 = \lambda S_2$ , cf. [126, lemma 9]. If  $e \neq 0$ , then from  $e = S_1 1 \otimes 1 = \lambda S_2 1 \otimes 1 = \lambda e$  it follows  $\lambda = 1$  and so  $S_1 = S_2$ . In the other case, if  $e = 0$ , then  $h' = 0$  and so  $\varphi = 0$ , whence  $\mathcal{A} \otimes_{\varphi} \mathcal{B} = \{0\}$  and indeed  $S_1 = S_2 = 0$  trivially. We continue with existence. There is a unique linear  $S_0: \mathcal{A} \odot \mathcal{B} \rightarrow X$  fixed by  $S_0(a \otimes b) = \varrho'(a)eb$ . Clearly  $S_0$  is also  $\mathcal{B}$ -linear. For any  $x, y \in \mathcal{A} \odot \mathcal{B}$ , say  $x = \sum_i a_i \otimes b_i$  and  $y = \sum_j \alpha_j \otimes \beta_j$ , we have

$$\begin{aligned} \langle S_0 x, S_0 y \rangle &= \sum_{i,j} \langle \varrho'(a_i)eb_i, \varrho'(\alpha_j)e\beta_j \rangle \\ &= \sum_{i,j} b_i^* \langle e, \varrho'(a_i^* \alpha_j)e \rangle \beta_j \\ &= \sum_{i,j} b_i^* \varphi(a_i^* \alpha_j) \beta_j \\ &= [x, y]. \end{aligned}$$

Thus  $S_0$  preserves the inner product. In particular  $S_0$  is bounded and so there is a unique bounded  $\mathcal{B}$ -linear  $S: \mathcal{A} \otimes_{\varphi} \mathcal{B} \rightarrow X$  with  $S \circ \eta = S_0$ . For all  $x \in \mathcal{A} \odot \mathcal{B}$  we have  $\langle \hat{x}, S^* S \hat{x} \rangle = \langle \hat{x}, \hat{x} \rangle$  and so  $S^* S = 1$  by 152IX. Thus  $S$  preserves the inner product. By definition  $S(1 \otimes 1) = \varrho'(1)e1 = e$ . Pick any  $a \in \mathcal{A}$ . A straightforward computation shows  $S\varrho(a) = \varrho'(a)S$  hence  $S^* \varrho'(a) = \varrho(a)S^*$  and  $S^* \varrho'(a)S = \varrho(a)S^* S = \varrho(a)$ . We have shown  $\text{ad}_S \circ \varrho' = \text{ad}_S$ .

- X (5,  $\sigma$  existence) Assume  $\varphi = h' \circ \varrho'$  for some nmiu-map  $\varrho': \mathcal{A} \rightarrow \mathcal{P}'$ , ncp-map  $h: \mathcal{P}' \rightarrow \mathcal{B}$  and von Neumann algebra  $\mathcal{P}'$ . It remains to be shown there is a  $\sigma: \mathcal{P}' \rightarrow \mathcal{B}^a(\mathcal{A} \otimes_{\varphi} \mathcal{B})$  with  $\sigma \circ \varrho' = \varrho$  and  $h \circ \sigma = h'$ . To apply the previous point, we perform the whole construction for  $h'$  instead of  $\varphi$  yielding  $h' = h_{h'} \circ \varrho_{h'}$  with  $\varrho_{h'}: \mathcal{P}' \rightarrow \mathcal{B}^a(\mathcal{P}' \otimes_{h'} \mathcal{B})$  and  $h_{h'}: \mathcal{B}^a(\mathcal{P}' \otimes_{h'} \mathcal{B}) \rightarrow \mathcal{B}$ . By IX there is a unique  $\mathcal{B}$ -linear map  $S: \mathcal{A} \otimes_{\varphi} \mathcal{B} \rightarrow \mathcal{P}' \otimes_{h'} \mathcal{B}$  with  $S^* S = 1$ ,  $S1 \otimes 1 = 1 \otimes 1$  and  $\text{ad}_S \circ \varrho_{h'} \circ \varrho' = \varrho$ . Define  $\sigma = \text{ad}_S \circ \varrho_{h'}$ . This  $\sigma$  fits the bill:  $\sigma \circ \varrho' = \text{ad}_S \circ \varrho_{h'} \circ \varrho' = \varrho$  and  $h \circ \sigma = h \circ \text{ad}_S \circ \varrho_{h'} = h_{h'} \circ \varrho_{h'} = h'$ .  $\square$

**155** We have shown that any ncp-map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  admits a dilation using Paschke’s generalization of GNS to Hilbert  $C^*$ -modules. There is also a generalization of Stinespring’s theorem to Hilbert  $C^*$ -modules due to Kasparov [75]:

**Theorem (KSGNS)** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}^a(X)$  be a cp-map for some  $C^*$ -algebras  $\mathcal{A}$ ,  $\mathcal{B}$  and Hilbert  $\mathcal{B}$ -module  $X$ . There exists a Hilbert  $\mathcal{B}$ -module  $Y$ , miu-map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}^a(Y)$  and adjointable  $\mathcal{B}$ -linear  $T: Y \rightarrow X$  with  $\varphi = \text{ad}_T \circ \varrho$ . II

For our purposes (the dilation of an arbitrary ncp-map  $\mathcal{A} \rightarrow \mathcal{B}$ ), Paschke's GNS construction sufficed. We leave open whether KSGNS admits a universal property like 139V and whether it is a Paschke dilation. III

We have all the tools to characterize the ncp-maps  $\varphi$  for which the Paschke representation is injective. This is a generalization of our answer [125] to the same question for the Stinespring embedding. 156

**Theorem** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  be any ncp-map with Paschke dilation  $(\mathcal{P}, \varrho, h)$ . Then  $[\varrho] = \llbracket \varphi \rrbracket$ . ( $\llbracket \varphi \rrbracket$  is the central carrier of  $\varphi$ , see 69I.) Thus  $\varrho$  is injective ( $[\varrho] = 1$ ) if and only if  $\varphi$  maps no non-zero central projection to zero ( $\llbracket \varphi \rrbracket = 1$ ). II

**Proof** (This is a simplified version of the proof we published earlier in [93, thm. 30].) By 140VIII, it is sufficient to prove the equivalence for the dilation constructed in 154III. Let  $p \in \mathcal{A}$  be any projection. We will show  $p \leq [\varrho]^\perp$  iff  $p \leq \llbracket \varphi \rrbracket^\perp$ . Assume  $p \leq [\varrho]$ , viz.  $\varrho(p) = 0$ . Pick any  $a \in \mathcal{A}$  and  $b \in \mathcal{B}$ . Then  $\varrho(p)a \otimes b = 0$ . By definition of  $\varrho$  and the inner product, this is the case iff  $b^* \varphi(a^* p^* p a) b = 0$ . In particular  $\varphi(a^* p a) = 0$ . In other words  $a^* p a \leq [\varphi]^\perp$ . This happens if and only if  $a [\varphi] a^* \leq p^\perp$  by 55V. Equivalently  $[a [\varphi] a^*] \leq p^\perp$ . As  $a$  was arbitrary, we see (using 68I), that  $\llbracket \varphi \rrbracket = \bigcup_{a \in \mathcal{A}} [a [\varphi] a^*] \leq p^\perp$ , as desired. For the converse, note that the two implications we just used are, in fact, equivalences: if  $\varphi(a^* p a) = 0$ , then  $b^* \varphi(a^* p a) b = 0$  for any  $b \in \mathcal{B}$  and if  $\varrho(p)a \otimes b = 0$  for all  $a \in \mathcal{A}$  and  $b \in \mathcal{B}$ , then  $\varrho(p) = 0$ , by 152IX. □

We will now prove a useful connection between the ncp-maps *nep-below*  $\varphi$  (as defined in a moment) and the commutant of the image of a Paschke representation (“ $\varrho$ ”) of  $\varphi$ . We will use it to give an alternative proof of the fact that pure maps and nmiu-maps are extreme among all ncpu-maps. 157

**Definition** For linear maps  $\varphi, \psi: \mathcal{A} \rightarrow \mathcal{B}$ , we say  $\varphi$  is nep-below  $\psi$  (in symbols:  $\varphi \leq_{\text{nep}} \psi$ ) whenever  $\psi - \varphi$  is an ncp-map. Furthermore, write II

$$[0, \varphi]_{\text{nep}} = \{\psi; \psi: \mathcal{A} \rightarrow \mathcal{B}; 0 \leq_{\text{nep}} \psi \leq_{\text{nep}} \varphi\}.$$

**Definition** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  be any ncp-map with Paschke dilation  $(\mathcal{P}, \varrho, h)$ . For  $t \in \varrho(\mathcal{A})^\square$ , the commutant of  $\varrho(\mathcal{A})$ , define  $\varphi_t = h(t\varrho(a))$ . III

The following is a generalization [93, prop. 5.4] to arbitrary Paschke dilations. IIIa

**Theorem** Assume  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  is an ncp-map with Paschke dilation  $(\mathcal{P}, \varrho, h)$ . The map  $t \mapsto \varphi_t$  is a linear order isomorphism  $[0, 1]_{\varrho(\mathcal{A})^\square} \rightarrow [0, \varphi]_{\text{nep}}$ . IV

- V **Proof** First we show the correspondence holds for the Paschke dilation constructed in 154 III. This proof is a slight variation on [93, prop. 5.4]. Then we show the correspondence carries over to arbitrary Paschke dilations.
- VI (Set-up) Let  $(\mathcal{B}^a(\mathcal{A} \otimes_{\varphi} \mathcal{B}), \varrho, h)$  denote the Paschke dilation constructed in 154 III. Clearly  $T \mapsto \varphi_T$  is linear for  $T \in \varrho(\mathcal{A})^{\square}$ . Pick  $T \in \varrho(\mathcal{A})^{\square}$ . As  $\sqrt{T} \in \varrho(\mathcal{A})^{\square}$  we have  $\varphi_T(a) = h(\sqrt{T}\varrho(a)\sqrt{T})$ . Thus  $\varphi_T$  is ncp. In particular, if  $T \leq S$  for some  $S \in \varrho(\mathcal{A})^{\square}$ , then  $\varphi_{S-T} = \varphi_S - \varphi_T$  is ncp and so  $\varphi_T \leq_{\text{ncp}} \varphi_S$ . Consequently, if  $T \leq 1$ , then  $\varphi_T \leq \varphi_1 = \varphi$ .
- VII (Order embedding) Let  $T \in \varrho(\mathcal{A})^{\square}$  be given such that  $\varphi_T$  is ncp. We must show  $T \geq 0$ . Pick any  $x \in \mathcal{A} \odot \mathcal{B}$ . By 152 IX and the construction of  $\mathcal{A} \otimes_{\varphi} \mathcal{B}$ , it is sufficient to show  $\langle \hat{x}, T\hat{x} \rangle \geq 0$ . Say  $\hat{x} \equiv \sum_i a_i \otimes b_i$ . Then

$$\langle \hat{x}, T\hat{x} \rangle = \sum_{i,j} b_i^* \langle 1 \otimes 1, T\varrho(a_i^* a_j) 1 \otimes 1 \rangle b_j = \sum_{i,j} b_i^* \varphi_T(a_i^* a_j) b_j \geq 0.$$

Thus  $T \mapsto \varphi_T$  is an order embedding and in particular an injection.

- VIII (Surjectivity) Pick any  $\psi \in [0, \varphi]_{\text{ncp}}$ . We will show there is a  $T \in \varrho(\mathcal{A})^{\square}$  with  $\varphi_T = \psi$  and  $0 \leq T \leq 1$ . Let  $(\mathcal{B}^a(\mathcal{A} \otimes_{\psi} \mathcal{B}), \varrho_{\psi}, h_{\psi})$  denote the Paschke dilation of  $\psi$  constructed in 154 III. Define  $B: \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{A} \otimes_{\psi} \mathcal{B}$  by  $B(a, b) = a \otimes b$ . This  $B$  is  $\varphi$ -compatible — indeed, for any  $n \in \mathbb{N}$ ,  $a_1, \dots, a_n \in \mathcal{A}$  and  $b_1, \dots, b_n \in \mathcal{B}$ , it follows from  $\psi \leq_{\text{ncp}} \varphi$  that

$$\left\| \sum_i B(a_i, b_i) \right\|^2 = \left\| \sum_{i,j} b_i^* \psi(a_i^* a_j) b_j \right\| \leq \left\| \sum_{i,j} b_i^* \varphi(a_i^* a_j) b_j \right\|.$$

So by 154 III, there is a unique bounded module map  $W: \mathcal{A} \otimes_{\varphi} \mathcal{B} \rightarrow \mathcal{A} \otimes_{\psi} \mathcal{B}$  fixed by  $W a \otimes b = a \otimes b$ . As  $\mathcal{A} \otimes_{\psi} \mathcal{B}$  is self-dual, the map  $W$  has an adjoint  $W^*$  by 152 VIII. For any  $a \in \mathcal{A}$ , we have  $W\varrho(a) = \varrho_{\psi}(a)W$  and so  $\varrho(a)W^* = W^*\varrho_{\psi}(a)$ . Thus  $W^*W\varrho(a) = W^*\varrho_{\psi}(a)W = \varrho(a)W^*W$ . Apparently  $W^*W \in \varrho(\mathcal{A})^{\square}$ . Define  $T \equiv W^*W$ . Clearly  $0 \leq T$ . As  $\psi \leq_{\text{ncp}} \varphi$ , we have for each  $x \equiv \sum_i a_i \otimes b_i$ , the inequality  $\langle x, x \rangle_{\psi} \leq \langle x, x \rangle_{\varphi}$  and so

$$\langle x, Tx \rangle_{\varphi} = \langle Wx, Wx \rangle_{\psi} = \langle x, x \rangle_{\psi} \leq \langle x, x \rangle_{\varphi}.$$

Thus  $T \leq 1$  by 152 IX. For any  $a \in \mathcal{A}$  we have

$$\begin{aligned} \varphi_T(a) &= \langle 1 \otimes 1, W^*W a \otimes 1 \rangle_{\varphi} \\ &= \langle W 1 \otimes 1, W a \otimes 1 \rangle_{\psi} \\ &= \langle 1 \otimes 1, a \otimes 1 \rangle_{\psi} \\ &= \psi(a), \end{aligned}$$

as desired. We have shown the correspondence holds for the Paschke dilation  $(\mathcal{B}^a(\mathcal{A} \otimes_{\varphi} \mathcal{B}), \varrho, h)$  constructed in 154 III.

(Arbitrary dilation) Let  $(\mathcal{P}', \varrho', h')$  be any Paschke dilation of  $\varphi$ . For brevity IX  
write  $\mathcal{P} \equiv \mathcal{A} \otimes_{\varphi} \mathcal{B}$  and for clarity write  $\varphi_T^{\mathcal{P}}$  and  $\varphi_t^{\mathcal{P}'}$  to distinguish between  $\varphi_T$   
and  $\varphi_t$  for  $T \in \varrho(\mathcal{A})^{\square}$  and  $t \in \varrho'(\mathcal{A})^{\square}$ . By 140VIII there is a unique nmiu-  
isomorphism  $\vartheta: \mathcal{P}' \rightarrow \mathcal{P}$  such that  $\vartheta \circ \varrho' = \varrho$  and  $h \circ \vartheta = h'$ . It is easy to  
see  $\vartheta$  restricts to a linear order isomorphism  $[0, 1]_{\varrho'(\mathcal{A})^{\square}} \rightarrow [0, 1]_{\varrho(\mathcal{A})^{\square}}$ . For  
any  $t \in \varrho'(\mathcal{A})^{\square}$  and  $a \in \mathcal{A}$  we have

$$\varphi_t^{\mathcal{P}'}(a) = h'(t\varrho'(a)) = h(\vartheta(t\varrho'(a))) = h(\vartheta(t)\varrho(a)) = \varphi_{\vartheta(t)}^{\mathcal{P}}(a)$$

and so  $t \mapsto \varphi_t^{\mathcal{P}'}$  is a linear order isomorphism as it is the composition of the  
linear order isomorphisms  $\vartheta^{-1}$  and  $t \mapsto \varphi_t^{\mathcal{P}}$ . □

We return to  $t \mapsto \varphi_t$  in 172, where we connect it to extremeness. X

## 2.4 Kaplansky density theorem for Hilbert $C^*$ -modules

Before we continue our study of Paschke dilations, we need to develop some more 158  
theory on (self-dual) Hilbert  $C^*$ -modules. First we will prove a generalization  
of the Kaplansky density theorem for Hilbert  $C^*$ -modules, which we will need  
to compute the tensor product of Paschke dilations.

The original density theorem (74IV) is an important foundational result in  
the theory of operator algebras; quoting Pedersen: “The density theorem is  
Kaplansky’s great gift to mankind. It can be used every day, and twice on  
Sundays.” [95, §2.3.4]. We used the original theorem already several times;  
notably to show that a von Neumann algebra is ultrastrongly complete (77I  
via 75VIII). This ultrastrong completeness was used to define operator divi-  
sion (81I) (and in turn filters 96I), to give the spatial tensor product a universal  
property (112XI) and in this thesis to show ultranorm completeness of self-dual  
Hilbert  $C^*$ -modules (149V).

So what does is this density theorem again? In essence, it is a remedy for  
the fact that strongly converging nets might not be bounded. The theorem  
is usually stated as follows: the unit ball of the SOT-closure of a self-adjoint  
algebra of operators  $\mathcal{A}$  is contained in the SOT-closure of the unit ball of  $\mathcal{A}$ .  
For our generalization, it is more convenient to consider the following variation.

**Kaplansky density theorem** Let  $\mathcal{B}$  be a von Neumann algebra with ultrastrongly- 1a  
dense  $C^*$ -subalgebra  $\mathcal{A} \subseteq \mathcal{B}$ . Then: for every element  $a \in \mathcal{A}$ , there is a net  $b_{\alpha}$   
in  $\mathcal{B}$  with  $b_{\alpha} \rightarrow a$  ultranorm and  $\|b_{\alpha}\| \leq \|b\|$  for all  $\alpha$ .

Now we are ready to state and prove our generalization, which is inspired by 1b  
the proof of the regular Kaplansky density theorem given in [7, thm. 1.2.2].  
Cf. 74IV.

- II **Kaplansky density theorem for Hilbert  $C^*$ -modules** Let  $X$  be a Hilbert  $\mathcal{B}$ -module for a von Neumann algebra  $\mathcal{B}$  with an ultranorm-dense  $\mathcal{A}$ -submodule  $D \subseteq X$ , where  $\mathcal{A} \subseteq \mathcal{B}$  is some  $C^*$ -subalgebra with  $\langle y, y \rangle \in \mathcal{A}$  for all  $y \in D$ .

Then: for every element  $x \in X$ , there is a net  $x_\alpha$  in  $D$  with  $x_\alpha \rightarrow x$  ultranorm and  $\|x_\alpha\| \leq \|x\|$  for all  $\alpha$ .

- III **Proof** Let  $x \in X$  be given. The case  $x = 0$  is trivial. Assume  $x \neq 0$ . Without loss of generality, we may assume  $\|x\| = 1$ . For this  $x$  and any  $y \in X$ , define

$$h(y) \equiv y \cdot \frac{2}{1 + \langle y, y \rangle} \quad g(x) \equiv x \cdot \frac{1}{1 + \sqrt{1 - \langle x, x \rangle}}.$$

We claim  $h$  is ultranorm continuous,  $h(g(x)) = x$  and  $\|h(y)\| \leq 1$  for all  $y \in X$ . From this, the promised result follows. Indeed, let  $x_\alpha$  be a net in  $D$  that converges ultranorm to  $g(x)$ . Then  $h(x_\alpha)$  is still in  $D$  and furthermore  $h(x_\alpha) \rightarrow h(g(x)) = x$  ultranorm and  $\|h(x_\alpha)\| \leq 1 = \|x\|$ , as desired.

- IV We start with  $\|h(y)\| \leq 1$ . The real map  $h_r: \lambda \mapsto 4\lambda(1 + \lambda)^{-2}$  is bounded by 1 and so by the functional calculus we see  $\|h(y)\|^2 = \|4\langle y, y \rangle(1 + \langle y, y \rangle)^{-2}\| = \|\overline{h_r(\langle y, y \rangle)}\| \leq 1$ . We continue with the proof of  $h(g(x)) = x$ . Note  $\langle g(x), g(x) \rangle = (1 + \sqrt{1 - \langle x, x \rangle})^{-2} \langle x, x \rangle$  and so with basic algebra

$$\begin{aligned} h(g(x)) &= x \cdot \frac{1}{1 + \sqrt{1 - \langle x, x \rangle}} \frac{2}{1 + (1 + \sqrt{1 - \langle x, x \rangle})^{-2} \langle x, x \rangle} \\ &= x \cdot 2 \left( \frac{(1 + \sqrt{1 - \langle x, x \rangle})^2 + \langle x, x \rangle}{1 + \sqrt{1 - \langle x, x \rangle}} \right)^{-1} \\ &= x. \end{aligned}$$

- V (Ultrnorm continuity  $h$ ) The real work is in the proof of the ultranorm continuity of  $h$ . Suppose  $y_\alpha \rightarrow y$  ultranorm in  $X$ . Define

$$\begin{aligned} A_1 &\equiv \frac{\langle y, y \rangle}{(1 + \langle y, y \rangle)^2} - \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} \frac{\langle y, y \rangle}{1 + \langle y, y \rangle} \\ A'_1 &\equiv \frac{\langle y_\alpha, y_\alpha \rangle}{(1 + \langle y_\alpha, y_\alpha \rangle)^2} - \frac{1}{1 + \langle y, y \rangle} \frac{\langle y_\alpha, y_\alpha \rangle}{1 + \langle y_\alpha, y_\alpha \rangle} \\ A_2 &\equiv \frac{1}{1 + \langle y, y \rangle} \langle y_\alpha - y, y_\alpha \rangle \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} \\ A'_2 &\equiv \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} \langle y - y_\alpha, y \rangle \frac{1}{1 + \langle y, y \rangle}. \end{aligned}$$

A straightforward computations shows

$$\langle h(y) - h(y_\alpha), h(y) - h(y_\alpha) \rangle = A_1 + A'_1 + A_2 + A'_2. \quad (2.11)$$

We will show that each of these terms converges ultraweakly to 0 as  $\alpha \rightarrow \infty$ . We start with  $A_1$ . We need two facts. First, for any positive  $b \in \mathcal{B}$ , we have  $0 \leq \frac{1}{1+b} \leq 1$  and  $0 \leq \frac{b}{1+b} \leq 1$ . Secondly

$$\begin{aligned} \frac{1}{1 + \langle y, y \rangle} - \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} &= \frac{1}{1 + \langle y, y \rangle} ((1 + \langle y_\alpha, y_\alpha \rangle) - (1 + \langle y, y \rangle)) \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} \\ &= \frac{1}{1 + \langle y, y \rangle} (\langle y_\alpha, y_\alpha \rangle - \langle y, y \rangle) \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle}. \end{aligned}$$

Putting these facts and the definition of  $A_1$  together, we get

$$\begin{aligned} A_1 &= \left( \frac{1}{1 + \langle y, y \rangle} - \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} \right) \frac{\langle y, y \rangle}{1 + \langle y, y \rangle} \\ &= \frac{1}{1 + \langle y, y \rangle} (\langle y_\alpha, y_\alpha \rangle - \langle y, y \rangle) \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} \frac{\langle y, y \rangle}{1 + \langle y, y \rangle} \\ &= \frac{1}{1 + \langle y, y \rangle} (\langle y_\alpha - y, y_\alpha \rangle + \langle y, y_\alpha - y \rangle) \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} \frac{\langle y, y \rangle}{1 + \langle y, y \rangle}. \end{aligned}$$

As multiplying with constants is ultraweakly continuous (see 45 IV), it is sufficient to show that ultraweakly, as  $\alpha \rightarrow \infty$ , we have

$$\langle y_\alpha - y, y_\alpha \rangle \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} \rightarrow 0 \quad \langle y, y_\alpha - y \rangle \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} \rightarrow 0. \quad (2.12)$$

For any np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$ , we have

$$\begin{aligned} \left| f \left( \langle y_\alpha - y, y_\alpha \rangle \frac{1}{1 + \langle y_\alpha, y_\alpha \rangle} \right) \right|^2 &= |\langle y_\alpha - y, y_\alpha (1 + \langle y_\alpha, y_\alpha \rangle)^{-1} \rangle_f|^2 \\ &\leq \|y_\alpha - y\|_f^2 \cdot \|y_\alpha (1 + \langle y_\alpha, y_\alpha \rangle)^{-1}\|_f^2 \\ &\leq \|y_\alpha - y\|_f^2 \cdot \|y_\alpha (1 + \langle y_\alpha, y_\alpha \rangle)^{-1}\|^2 \cdot \|f\| \\ &\leq \|y_\alpha - y\|_f^2 \cdot \|f\| \rightarrow 0. \end{aligned}$$

We have shown the LHS of (2.12). The proof for the RHS is different, but simpler. So  $A_1$  vanishes ultraweakly. In a similar way one sees  $A'_1 \rightarrow 0$  ultraweakly. The proofs for  $A_2, A'_2 \rightarrow 0$  are very similar.  $\square$

## 2.5 More on self-dual Hilbert $C^*$ -modules

We have a second look at self-dual Hilbert  $C^*$ -modules over von Neumann algebras and their orthonormal bases. We start with some good news. In  $\mathcal{B}(\mathcal{H})$  the linear span of  $|e_i\rangle\langle e_j|$  is ultraweakly dense for any orthonormal basis  $(e_i)_{i \in I}$  of  $\mathcal{H}$ . For self-dual Hilbert  $\mathcal{B}$ -modules, we have a similar result.

II **Definition** Let  $X$  be a Hilbert  $\mathcal{B}$ -module. For any  $x, y \in X$ , define the bounded operator  $|x\rangle\langle y| \in \mathcal{B}^a(X)$  by  $|x\rangle\langle y|z = x\langle y, z\rangle$ .

IIa **Remark** The operator  $|x\rangle\langle y|$  is usually denoted by  $\theta_{x,y}$  in the literature, see for instance [88, §2.2].

III The following rules are easy to check. For  $x, y, v, w \in X$  and  $b \in \mathcal{B}$  we have

$$|x\rangle\langle y|^* = |y\rangle\langle x| \quad |xb\rangle\langle y| = |x\rangle\langle yb^*| \quad |x\rangle\langle y||v\rangle\langle w| = |x\langle y, v\rangle\rangle\langle w|.$$

If  $\langle e, e\rangle$  is a projection, then  $|e\rangle\langle e|$  is a projection (using 149 III). For any  $T \in \mathcal{B}^a(X)$  we have  $T|x\rangle\langle y| = |Tx\rangle\langle y|$  and  $|x\rangle\langle y|T^* = |x\rangle\langle Ty|$ .

IV **Proposition** Let  $X$  be a self-dual Hilbert  $\mathcal{B}$ -module for a von Neumann algebra  $\mathcal{B}$ . If  $(e_i)_{i \in I}$  is an orthonormal basis of  $X$ , then the linear span of the operators  $\{|e_i b\rangle\langle e_j|; i, j \in I, b \in \mathcal{B}\}$  is ultraweakly dense in  $\mathcal{B}^a(X)$ .

V **Proof** We start with some preparation.

VI For a finite subset  $S \subseteq I$ , write  $p_S = \sum_{i \in S} |e_i\rangle\langle e_i|$ . The  $(|e_i\rangle\langle e_i|)_{i \in I}$  are pairwise orthogonal projections. Thus  $p_S$  is a projection. Also  $p_S \leq p_{S'}$  when  $S \subseteq S'$ . Pick any  $x \in X$ . As  $p_S x = \sum_{i \in S} e_i \langle e_i, x\rangle$ , we have

$$\langle x, p_S x\rangle = \langle p_S x, p_S x\rangle = \sum_{i, j \in S} \langle x, e_i\rangle \langle e_i, e_j\rangle \langle e_j, x\rangle = \sum_{i \in S} \langle x, e_i\rangle \langle e_i, x\rangle,$$

and so by Parseval  $\sup_S \langle x, p_S x\rangle = \langle x, x\rangle$ . Hence  $\sup_S p_S = 1$  by 22 VIII and so  $p_S$  converges ultraweakly to 1 by 44 XIV.

VII Pick any  $T \in \mathcal{B}^a(X)$ . For  $i, j \in I$  we have

$$|e_i\rangle\langle e_i|T|e_j\rangle\langle e_j| = |e_i\rangle\langle e_i||Te_j\rangle\langle e_j| = |e_i\rangle\langle e_i, Te_j\rangle\rangle\langle e_j|.$$

and so  $p_S T p_S$  is in the linear span of  $\{|e_i b\rangle\langle e_j|\}$ . Thus it is sufficient to prove  $p_S T p_S$  converges ultraweakly to  $T$ .

VIII Pick any np-map  $f: \mathcal{B} \rightarrow \mathbb{C}$ . We want to show  $|f(T - p_S T p_S)| \rightarrow 0$ . Clearly

$$|f(T - p_S T p_S)| \leq |f((1 - p_S)T)| + |f(p_S T (1 - p_S))|.$$

Using Cauchy-Schwarz and VI we see the second term vanishes

$$|f(p_S T (1 - p_S))|^2 \leq f(p_S T T^* p_S) f(1 - p_S) \leq \|T\|^2 f(1) f(1 - p_S) \rightarrow 0.$$

Similarly the first term vanishes and so indeed  $p_S T p_S \rightarrow T$  ultraweakly.  $\square$

IX **Proposition** Let  $X$  be a self-dual Hilbert  $\mathcal{B}$ -module for a von Neumann algebra  $\mathcal{B}$ . If  $x_\alpha \rightarrow x$  ultranorm for a norm-bounded net  $x_\alpha$  in  $X$ , then  $|x_\alpha\rangle\langle y| \rightarrow |x\rangle\langle y|$  ultraweakly for any  $y \in Y$ .

X **Proof** We need some preparation. Define

$$\Omega \equiv \{f(\langle x, (\cdot)x\rangle); f: \mathcal{B} \rightarrow \mathbb{C} \text{ np-map, } x \in X\}.$$

From 144 I it follows  $\Omega$  is order separating. We will use this fact twice. First we will prove  $\| |z\rangle\langle y| \| \leq \|z\| \|y\|$ . For any  $\omega \in \Omega$ , say  $\omega \equiv f(\langle x, (\cdot)x \rangle)$ , we have

$$\omega(|y\rangle\langle y|) = f(\langle x, y \rangle \langle y, x \rangle) \leq f(\langle x, x \rangle) \|y\|^2 = \|\omega\| \|y\|^2$$

and so from 21 VII it follows  $\| |y\rangle\langle y| \| \leq \|y\|^2$ . Whence

$$\begin{aligned} \| |z\rangle\langle y| \|^2 &= \| |y\rangle\langle z| |z\rangle\langle y| \| \\ &= \| |y\rangle\langle z, z|^{\frac{1}{2}} \langle y, z|^{\frac{1}{2}} \| \\ &\leq \| |y\rangle\langle z, z|^{\frac{1}{2}} \|^2. \end{aligned}$$

Hence  $\| |z\rangle\langle y| \| \leq \|y\rangle\langle z, z|^{\frac{1}{2}} \| \leq \|y\| \|z\|$ , as promised.

Now we start the proof proper. As  $|\cdot\rangle\langle y|$  is linear and  $x_\alpha - x$  is norm-bounded, we may assume without loss of generality that  $x_\alpha \rightarrow 0$  ultranorm. We have to prove  $|x_\alpha\rangle\langle y| \rightarrow 0$  ultraweakly. Let  $g: \mathcal{B}^a(X) \rightarrow \mathbb{C}$  be an arbitrary np-map. Using the order separation of  $\Omega$  a second time and the fact that  $\omega(T^*(\cdot)T) \in \Omega$  for all  $T \in \mathcal{B}^a(X)$  and  $\omega \in \Omega$ , we see that the linear span of  $\Omega$  is operator norm dense among all np-maps  $\mathcal{B}^a(X) \rightarrow \mathbb{C}$  by 90 II. So there are  $n \in \mathbb{N}$  and  $\omega_1, \dots, \omega_n \in \Omega$  with  $\|g - g'\| \leq \varepsilon$ , where  $g' \equiv \sum_{i=1}^n \omega_i$ . By unfolding definitions and 148 V, it follows easily that  $\omega_i(|\cdot\rangle\langle y|)$  is ultranorm continuous. So  $g'(|\cdot\rangle\langle y|)$  is ultranorm continuous as well. Pick any  $\varepsilon > 0$ . By the previous, there is an  $\alpha_0$  such that for all  $\alpha \geq \alpha_0$  we have  $|g'(|x_\alpha\rangle\langle y|)| \leq \varepsilon$ , hence

$$\begin{aligned} |g(|x_\alpha\rangle\langle y|)| &\leq |(g - g')(|x_\alpha\rangle\langle y|)| + |g'(|x_\alpha\rangle\langle y|)| \\ &\leq \|g - g'\| \cdot \| |x_\alpha\rangle\langle y| \| + \varepsilon \\ &\leq \varepsilon (\|x_\alpha\| \cdot \|y\| + 1). \end{aligned}$$

As  $x_\alpha$  is norm-bounded and  $\varepsilon$  was arbitrary, we conclude  $g(|x_\alpha\rangle\langle y|) \rightarrow 0$ . Thus  $|x_\alpha\rangle\langle y| \rightarrow 0$  ultraweakly, as desired.  $\square$

For a Hilbert space  $\mathcal{H}$  with linear subspace  $V \subseteq \mathcal{H}$  we have  $V \subseteq V^{\perp\perp} = \overline{V}$  and  $\mathcal{H} \cong V^{\perp\perp} \oplus V^\perp$ . We generalize this to self-dual Hilbert  $C^*$ -modules over von Neumann algebras. Let's introduce the cast: 160

**Exercise** Let  $\mathcal{B}$  be a von Neumann algebra. Assume  $X$  and  $Y$  are self-dual Hilbert  $\mathcal{B}$ -modules with orthonormal bases  $E \subseteq X$  and  $F \subseteq Y$ . Write  $\kappa_1: X \rightarrow X \oplus Y$  and  $\kappa_2: Y \rightarrow X \oplus Y$  for the maps  $\kappa_1(x) = (x, 0)$  and  $\kappa_2(y) = (0, y)$ . Show that  $\kappa_1(E) \cup \kappa_2(F)$  is an orthonormal basis of  $X \oplus Y$ . Conclude that  $X \oplus Y$  is self-dual. II

**Definition** Let  $X$  be a Hilbert  $C^*$ -module with subset  $V \subseteq X$ . Write  $V^\perp$  for the orthocomplement of  $V$ , defined by III

$$V^\perp = \{x; x \in X; \langle x, v \rangle = 0 \text{ for all } v \in V\}.$$

**IV Proposition** Assume  $X$  is a self-dual Hilbert  $\mathcal{B}$ -module for a von Neumann algebra  $\mathcal{B}$ . Let  $V \subseteq X$  be any subset. Then

1.  $V^\perp$  is a ultranorm-closed Hilbert  $\mathcal{B}$ -submodule of  $X$  (and so is  $V^{\perp\perp}$ );
2.  $V^{\perp\perp}$  is the ultranorm closure of the  $\mathcal{B}$ -linear span of  $V$  and
3.  $V^{\perp\perp} \oplus V^\perp \cong X$  as Hilbert  $C^*$ -modules via  $(x, y) \mapsto x + y$ .

**V Proof** It is easy to see  $V^\perp$  is a submodule of  $X$  and  $V \subseteq V^{\perp\perp}$ . To show  $V^\perp$  is ultranorm closed in  $X$ , assume  $x_\alpha$  is a net in  $V^\perp$  converging ultranorm to  $x \in X$ . For each  $v \in V$  we have  $\langle v, \text{unlim}_\alpha x_\alpha \rangle \stackrel{148\text{III}}{=} \text{uslim}_\alpha \langle v, x_\alpha \rangle = 0$  and so  $x \in V^\perp$  as well. Thus  $V^\perp$  is ultranorm complete (and so self dual by 149V).

**VI** Write  $W$  for the ultranorm closure of the  $\mathcal{B}$ -linear span of  $V$  in  $X$ . It follows from 148III, that  $W$  is a submodule. Clearly the induced uniformity of  $X$  on  $W$  is the same as its own ultranorm uniformity: thus  $W$  is ultranorm complete and  $W \subseteq V^{\perp\perp}$ . By 149V there is an orthonormal basis  $(e_i)_{i \in I}$  of  $W$ . Going back to the construction of  $(e_i)_{i \in I}$  (in 149VIII), we see that we can extend it to a maximal orthonormal subset of  $X$ , which must be a basis for the whole of  $X$ . Pick any  $(d_j)_{j \in J}$  in  $X$  such that  $\{e_i\}_{i \in I} \cup \{d_j\}_{j \in J}$  is an orthonormal basis of  $X$ .

**VII** By construction  $d_j \in W^\perp \subseteq V^\perp$  for every  $j \in J$ . For any  $x \in X$  we have  $x = \sum_i e_i \langle e_i, x \rangle + \sum_j d_j \langle d_j, x \rangle$ . If  $x \in V^{\perp\perp}$  then  $x = \sum_i e_i \langle e_i, x \rangle$  as  $\langle x, d_j \rangle = 0$  for each  $j \in J$ . Thus  $V^{\perp\perp} \subseteq W$ , so  $V^{\perp\perp} = W$ . In particular, as  $V^\perp$  is already an ultranorm closed submodule, we find  $W^\perp = (V^\perp)^{\perp\perp} = V^\perp$ .

**VIII** So  $(e_i)_i$  is a basis for  $W = V^{\perp\perp}$  and  $(d_j)_j$  is a basis for  $W^\perp = V^\perp$ . For brevity, let  $\vartheta: V^{\perp\perp} \oplus V^\perp \rightarrow X$  denote the map  $(y, y') \mapsto y + y'$ . Clearly  $\vartheta$  is bounded  $\mathcal{B}$ -linear. As  $\langle y, y' \rangle = 0$  for  $(y, y') \in V^{\perp\perp} \oplus V^\perp$  we easily see  $\vartheta$  is inner product preserving and thus injective. As  $\{e_i\}_{i \in I} \cup \{d_j\}_{j \in J}$  is an orthonormal basis of  $X$ ,  $\vartheta$  is surjective. As  $\vartheta$  is inner product preserving, its inverse is bounded — hence  $\vartheta$  is an isomorphism of Hilbert  $C^*$ -modules, as promised.  $\square$

**IX Exercise** Assume  $X$  is a self-dual Hilbert  $\mathcal{B}$ -module for a von Neumann algebra  $\mathcal{B}$ . Suppose  $E \subseteq X$  is an orthonormal set. Show

1.  $E$  is a basis of  $E^{\perp\perp}$  and
2. for any  $x \in X$ , we have  $x \in E^{\perp\perp}$  if and only if  $\langle x, x \rangle = \sum_{e \in E} \langle x, e \rangle \langle e, x \rangle$ .

**X Exercise** Let  $X$  be a self-dual Hilbert  $\mathcal{B}$ -module for some von Neumann algebra  $\mathcal{B}$ . Show that for any  $x_1, \dots, x_n \in X$ , there is a finite orthonormal basis of  $\{x_1, \dots, x_n\}^{\perp\perp}$  consisting of at most  $n$  elements. (Use the orthonormalization in the last part of 149VIII.)

Every Hilbert space  $\mathcal{H}$  is isomorphic to some  $\ell^2(I)$ . In fact, the cardinality of  $I$  is the only thing that matters, in the following sense: if  $\ell^2(I) \cong \ell^2(J)$ , then  $I$  and  $J$  have the same cardinality. What about Hilbert  $C^*$ -modules? 161

**Exercise** In this exercise we prove [93, thm. 3.12]. Let  $\mathcal{B}$  be a von Neumann algebra and  $(p_i)_{i \in I}$  a family of projections from  $\mathcal{B}$ . Write  $\ell^2((p_i)_{i \in I})$  for the set of  $I$ -tuples  $(b_i)_{i \in I}$  from  $\mathcal{B}$  that are  $\ell^2$ -summable (149) and furthermore  $[b_i b_i^*] \leq p_i$  for every  $i \in I$ . Show that  $\ell^2((p_i)_{i \in I})$  is a right  $\mathcal{B}$ -module with coordinatewise operations. Prove that for any  $(b_i)_i$  and  $(c_i)_i$  in  $\ell^2((p_i)_{i \in I})$  the sum  $\sum_{i \in I} b_i^* c_i =: \langle (b_i)_i, (c_i)_i \rangle$  converges ultraweakly and turns  $\ell^2((p_i)_{i \in I})$  into a pre-Hilbert  $\mathcal{B}$ -module. Conclude  $\ell^2((p_i)_{i \in I})$  is self-dual and, in fact, that for every self-dual Hilbert  $\mathcal{B}$ -module  $X$  with any orthonormal basis  $(e_i)_{i \in I}$ , we have  $X \cong \ell^2(\langle e_i, e_i \rangle)_{i \in I}$ . II

Let  $\mathcal{B}$  be a von Neumann algebra and  $(p_i)_{i \in I}, (q_j)_{j \in J}$  families of projections from  $\mathcal{B}$ . What can we say if  $\ell^2((p_i)_{i \in I}) \cong \ell^2((q_j)_{j \in J})$ ? ( $\ell^2$  as defined in II.) Let's start with some counterexamples. III

1. Do all orthonormal bases have the same cardinality?  $\mathcal{B}$  is a self-dual Hilbert  $\mathcal{B}$ -module over itself. Clearly, the element 1 by itself is an orthonormal basis of  $\mathcal{B}$ . However, for every projection  $p \in \mathcal{B}$  the pair  $\{p, 1-p\}$  is also an orthonormal basis. Worse still, for  $\mathcal{B} \equiv \mathcal{B}(\ell^2(\mathbb{N}))$ , the set

$$\{|0\rangle\langle 0|, |1\rangle\langle 1|, |2\rangle\langle 2|, \dots\}.$$

is an orthonormal basis.

2. Perhaps any two orthonormal bases have a common coarsening in some sense? Take  $\mathcal{B} = M_3$ . It might not be obvious at this moment, but both

$$\{|0\rangle\langle 0| + |2\rangle\langle 2|, |1\rangle\langle 1|\} \quad \text{and} \quad \{|0\rangle\langle 0|, |2\rangle\langle 2| + |1\rangle\langle 1|\}$$

are orthonormal bases of  $M_3$  over itself. (Combine IV with V.) There does not seem to be an obvious way in which these two bases have a common coarsening of sorts.

3. Maybe  $\sum_{i \in I} \langle e_i, e_i \rangle = \sum_{j \in J} \langle d_j, d_j \rangle$  for finite orthonormal bases  $(e_i)_{i \in I}$  and  $(d_j)_{j \in J}$ ? For commutative  $\mathcal{B}$  this indeed holds using Parseval's identity. However, in general it fails: both  $\{|0\rangle\langle 0|, |1\rangle\langle 1|\}$  and  $\{|0\rangle\langle 0|, |1\rangle\langle 0|\}$  are orthonormal bases of  $M_2$  over itself, but  $|0\rangle\langle 0| + |1\rangle\langle 1| \neq 2|0\rangle\langle 0|$ .

**Exercise** Let  $(e_i)_{i \in I}$  be some orthonormal basis of a Hilbert  $\mathcal{B}$ -module  $X$ . Show that if  $(u_i)_{i \in I}$  is a family of partial isometries from  $\mathcal{B}$  (see 79 IV) such that  $u_i u_i^* = \langle e_i, e_i \rangle$  for all  $i \in I$ , then  $(e_i u_i)_{i \in I}$  is an orthonormal basis of  $X$ . IV

Use this and II to conclude that  $\ell^2((p_i)_{i \in I}) \cong \ell^2((q_i)_{i \in I})$  for any projections  $(p_i)_{i \in I}, (q_i)_{i \in I}$  from  $\mathcal{B}$  with  $p_i \sim q_i$  for  $i \in I$ . (Here,  $\sim$  denotes the [Murray–von Neumann equivalence](#). That is:  $p \sim q$  iff there is a partial isometry  $u$  with  $u^*u = p$  and  $uu^* = q$ . Eg.  $|0\rangle\langle 0| \sim |+\rangle\langle +|$  via  $|0\rangle\langle +|$ .)

- V **Exercise** Let  $(e_i)_{i \in I}$  be some orthonormal basis of a Hilbert  $\mathcal{B}$ -module  $X$  with distinguished  $1, 2 \in I$ . Show that if  $\langle e_1, e_1 \rangle + \langle e_2, e_2 \rangle \leq 1$ , then we can make a new orthonormal basis of  $X$  by removing  $e_1$  and  $e_2$  and inserting  $e_1 + e_2$ .

Conclude  $p\mathcal{B} \oplus q\mathcal{B} \cong (p + q)\mathcal{B}$  for projections  $p, q \in \mathcal{B}$  with  $p + q \leq 1$ .

- 162 Before we can prove our normal form (in IV), we need some more *comparison theory of projections*, which was touched upon in 83 II. Recall that for projections  $p, q$  in a von Neumann algebra  $\mathcal{A}$ , we write  $p \lesssim q$  if there is a partial isometry  $u$  with  $u^*u = p$  and  $uu^* \leq q$ . In 83 IV we saw  $\lesssim$  preorders the projections.

- II **Proposition** Let  $\mathcal{B}$  be a von Neumann algebra that is a factor (i.e. with center  $\mathbb{C}1$ ). For any two projections  $p, q \in \mathcal{B}$ , either  $p \lesssim q$  or  $q \lesssim p$  (see 83 II).

- III **Proof** (For a more traditional proof, see e.g. [73, prop. 6.2.4].) Let  $p, q \in \mathcal{B}$  be any projections. By Zorn's lemma, there is a maximal set  $U \subseteq \mathcal{B}$  of partial isometries with  $\sum_{u \in U} u^*u \leq p$  and  $\sum_{u \in U} uu^* \leq q$ . Define  $p_0 \equiv p - \sum_{u \in U} u^*u$  and  $q_0 \equiv q - \sum_{u \in U} uu^*$ . Note that if  $p_0 = 0$ , then  $p \lesssim q$  via  $\sum_{u \in U} u$ , c.f. 89 III. Similarly, if  $q_0 = 0$ , then  $q \lesssim p$ . So, reasoning towards contradiction, suppose  $p_0 \neq 0$  and  $q_0 \neq 0$ . As  $\mathcal{B}$  is a factor  $\llbracket q_0 \rrbracket = 1$  and so  $0 < p_0 = p_0 \llbracket q_0 \rrbracket p_0 = p_0 (\bigcup_{a \in \mathcal{B}} [a^* q_0 a]) p_0 = \bigcup_{a \in \mathcal{B}} [p_0 a^* q_0 a p_0]$  by 68 I and 60 IX. Pick an  $a \in \mathcal{B}$  with  $[p_0 a^* q_0 a p_0] \neq 0$ . Then  $u_0 \equiv [q_0 a p_0]$ , see 82 I, is a non-zero partial isometry satisfying  $u_0^* u_0 = [p_0 a^* q_0 a p_0] \leq p_0$  and  $u_0 u_0^* = [q_0 a p_0 a^* q_0] \leq q_0$ , contradicting maximality of  $U$ .  $\square$

- IV **Theorem** Let  $\mathcal{B}$  be a von Neumann algebra that is a factor (i.e. with center  $\mathbb{C}1$ ). Suppose  $X$  is a self-dual Hilbert  $\mathcal{B}$ -module. Either there is an infinite cardinal  $\kappa$  such that  $X \cong \ell^2((1)_{\alpha \in \kappa})$ , with  $\ell^2$  as in 161 II, or there is a natural number  $n \in \mathbb{N}$  and a projection  $p \in \mathcal{B}$  such that  $X \cong \ell^2((1, \dots, 1, p))$ , where 1 occurs  $n$  times.

- V **Proof** The case  $X = \{0\}$  is covered by  $n = 0$  and  $p = 0$ . Assume  $X \neq \{0\}$ .

- VI As a first step we will prove that either

1.  $X$  has an orthonormal basis  $E$  such that  $\langle e, e \rangle = 1$  for some  $e \in E$  or
2. there is a single vector  $e \in X$  such that  $\{e\}$  is an orthonormal basis.

Pick any orthonormal basis  $E_0$  of  $X$ . Let  $P$  denote the poset of subsets  $U \subseteq E_0 \times \mathcal{B}$  satisfying

1.  $u$  is a partial isometry with  $uu^* = \langle e, e \rangle$  for every  $(e, u) \in U$  and

2. the domains of these  $u$  are orthogonal:  $\sum_{(e,u) \in U} u^*u \leq 1$ .

For any non-empty chain  $C \subseteq P$  it is easy to see  $\bigcup C \in P$  and so by Zorn's lemma, there is a maximal element  $U_0 \in P$ . Define

$$e_0 = \sum_{(e,u) \in U_0} eu \quad E_1 = E_0 \setminus \{e; (e,u) \in U_0\}.$$

The set  $\{e_0\} \cup E_1$  is an orthonormal basis of  $X$  — indeed, it is clearly orthonormal and if  $x = \sum_{e \in E_0} e \langle e, x \rangle$ , then

$$x = e_0 \left( \sum_{(e,u) \in U_0} u^* \langle e, x \rangle \right) + \sum_{e \in E_1} e \langle e, x \rangle.$$

If  $E_1 = \emptyset$ , then  $\{e_0\}$  is an orthonormal basis of  $X$  and we have shown point 2.

For the other case, assume  $E_1 \neq \emptyset$ . Pick  $e_1 \in E_1$ . Write  $p_0 = \langle e_0, e_0 \rangle$  and  $p_1 = \langle e_1, e_1 \rangle$ . Suppose  $p_1 \lesssim 1 - p_0$ . Then  $vv^* = p_1$  and  $v^*v \leq 1 - p_0$  for some  $v \in \mathcal{B}$ . Hence  $v^*v + p_0 \leq 1$ . So  $U_0 \cup \{(e_1, v)\} \in P$  contradicting maximality of  $U_0$ . Apparently  $p_1 \not\lesssim 1 - p_0$ . So by II, we must have  $1 - p_0 \lesssim p_1$ . Let  $v \in \mathcal{B}$  be such that  $vv^* = 1 - p_0$  and  $v^*v \leq p_1$ . Write  $p'_0 = 1 - v^*v$ . As  $1 - p'_0 \leq p_1 = 1 - (1 - p_1)$  we have  $(1 - p'_0) + (1 - p_1) \leq 1$ . Define  $q = 1 - ((1 - p'_0) + (1 - p_1))$ . We now have the following splittings of 1.

$$\begin{array}{|c|c|c|} \hline & \overbrace{p'_0} & \\ \hline 1 - p_1 & q & v^*v \\ \hline & \underbrace{p_1} & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline p_0 & vv^* \\ \hline \end{array}$$

Note  $q = p'_0 + p_1 - 1$  and  $p'_0, p_1 \geq q$ . Define

$$D = \{d_0, d_1\} \quad \text{where} \quad d_0 = e_1q \quad d_1 = e_0 + e_1v^*.$$

From  $p_1q = q$  it follows  $D$  is orthogonal. Clearly  $\langle d_0, d_0 \rangle = q$  and

$$\langle d_1, d_1 \rangle = \langle e_0, e_0 \rangle + vp_1v^* = p_0 + vv^* = 1.$$

So  $D$  is an orthonormal set. Using  $0 = (1 - p_0)p_0 = vv^*p_0$  and  $q + v^*v = p_1$ , we see that for any  $x \equiv e_0b_0 + e_1b_1$  with  $b_0 \in p_0\mathcal{B}$  and  $b_1 \in p_1\mathcal{B}$ , we have

$$\begin{aligned} d_0 \langle d_0, x \rangle + d_1 \langle d_1, x \rangle &= d_0b_1 + d_1(vb_1 + b_0) \\ &= e_1qb_1 + e_0vb_1 + e_1v^*vb_1 + e_0b_0 + e_1v^*b_0 \\ &= e_1qb_1 + e_0p_0vv^*vb_1 + e_1v^*vb_1 + e_0b_0 + e_1v^*vv^*p_0b_0 \\ &= e_1qb_1 + e_1v^*vb_1 + e_0b_0 \\ &= e_1(q + v^*v)b_1 + e_0b_0 \\ &= e_1b_1 + e_0b_0 = x. \end{aligned}$$

Thus  $D$  is an orthonormal basis of  $\{e_0, e_1\}^{\perp\perp}$ . Hence  $E \equiv D \cup (E_1 - \{e_1\})$  is an orthonormal basis of  $X$  with  $d_1 \in E$  such that  $\langle d_1, d_1 \rangle = 1$ .

- VII For brevity, call  $X$  **1-dim** if there is a one-element orthonormal basis of  $X$ . In VI we saw how to create an orthonormal basis  $D$  of a non-1-dim  $X$  with an  $e_0 \in D$  such that  $\langle e_0, e_0 \rangle = 1$ . If  $\{e_0\}^\perp$  is not 1-dim, we can apply VI on  $\{e_0\}^\perp$  (instead of  $X$ ) to find an orthonormal basis  $D'$  of  $\{e_0\}^\perp$  with  $e_1 \in D'$  such that  $\langle e_1, e_1 \rangle = 1$ . This procedure can be continued using Zorn's lemma as follows. Let  $P$  denote the poset of orthogonal subsets  $E \subseteq X$  satisfying  $\langle e, e \rangle = 1$  for all  $e \in E$ . Clearly  $\emptyset \in P$  and for any non-empty chain  $C \subseteq P$  we have  $\bigcup C \in P$ . By Zorn's lemma there is a maximal  $E \in P$ . Suppose  $E^\perp \neq \{0\}$  and  $E^\perp$  is not 1-dim. Then by VI we can find an orthonormal basis  $E'$  of  $E^\perp$  together with  $e \in E'$  such that  $\langle e, e \rangle = 1$ . Now  $E' \cup \{e\} \in P$ , contradicting maximality of  $E$ . Apparently  $E^\perp = \{0\}$  or  $E^\perp$  is 1-dim. If  $E^\perp = \{0\}$  then we are done taking for  $\kappa$  the cardinality of  $E$ . For the other case, assume  $E^\perp = \{e\}^{\perp\perp}$  for some  $e \in X$  with  $\langle e, e \rangle$  a non-zero projection. If  $E$  is finite, then we are done as well. So, assume  $E$  is infinite. Pick a sequence  $e_1, e_2, \dots \in E$  of distinct elements. Write  $p = \langle e, e \rangle$  and  $E_0 = \{e_n; n \in \mathbb{N}\}$ . Define

$$E_1 = \{e + e_1(1 - p), e_1p + e_2(1 - p), e_2p + e_3(1 - p), \dots\}.$$

It is easy to see  $E_1$  is an orthogonal set with  $\langle d, d \rangle = 1$  for all  $d \in E_1$ . Furthermore  $E_1$  is an orthonormal basis for  $E_0 \cup \{e\}$ , hence  $(E - E_0) \cup E_1$  is the desired orthonormal basis of  $X$ .  $\square$

- VIII Does this settle the issue raised in 161I? One might hope that  $\ell^2((1_{\mathcal{B}})_{\alpha \in \kappa}) \cong \ell^2((1_{\mathcal{B}})_{\beta \in \lambda})$  implies  $\kappa = \lambda$  for all cardinals  $\kappa, \lambda$ . This is not the case. Indeed, if  $\mathcal{B}$  is a type III factor of operators on a separable Hilbert space, then for any non-zero projection  $p$ , we have  $p \sim 1 - p$  (by combining [73, dfn. 6.5.1] and [73, cor. 6.3.5]) and so  $\mathcal{B} \cong p\mathcal{B} \oplus (1 - p)\mathcal{B} \cong \mathcal{B} \oplus \mathcal{B}$  as Hilbert  $C^*$ -modules.

163 Before we continue to the next topic, the exterior tensor product of Hilbert  $C^*$ -modules, we will show that the ultranorm completion 150II is determined by its universal property. We will need this fact in our treatment of the exterior tensor product.

- II **Proposition** Let  $\mathcal{B}$  be a von Neumann algebra and  $V$  a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product  $[\cdot, \cdot]$ . There is an up-to-isomorphism unique self-dual Hilbert  $\mathcal{B}$ -module  $X$  together with inner product preserving  $\mathcal{B}$ -linear  $\eta: V \rightarrow X$  with the following universal property.

For every bounded  $\mathcal{B}$ -linear map  $T: V \rightarrow Y$  to some self-dual Hilbert  $\mathcal{B}$ -module  $Y$ , there is a unique bounded  $\mathcal{B}$ -linear map  $\hat{T}: X \rightarrow Y$  with  $\hat{T} \circ \eta = T$ .

Moreover, for such  $\eta: V \rightarrow X$ , the image of  $V$  under  $\eta$  is ultranorm dense.

**Proof** We already know such a  $X$  and  $\eta: V \rightarrow X$  exist by 150II and 151 Ia. III  
 For this one  $\eta(V)$  is ultranorm dense in  $X$ . Assume there is another self-dual Hilbert  $\mathcal{B}$ -module  $X_2$  with inner product preserving  $\mathcal{B}$ -linear  $\eta_2: V \rightarrow X_2$  satisfying the universal property. By the universal property of  $\eta$  applied to  $\eta_2$ , there must exist a unique  $U: X \rightarrow X_2$  with  $U \circ \eta = \eta_2$ . Similarly, there is a  $V: X_2 \rightarrow X$  with  $V \circ \eta_2 = \eta$ . Clearly  $\text{id}_X \circ \eta = V \circ \eta_2 = V \circ U \circ \eta$  and so by the uniqueness  $\text{id} = V \circ U$ . Similarly  $U \circ V = \text{id}$ . It remains to be shown  $U$  preserves the inner product. As

$$\langle \eta(x), \eta(y) \rangle = [x, y] = \langle \eta_2(x), \eta_2(y) \rangle = \langle U\eta(x), U\eta(y) \rangle$$

for all  $x, y \in X$ , we know  $\eta(X)$  is ultranorm dense. So by ultranorm continuity and bijectivity of  $U$ , we know  $U(\eta(X)) = \eta_2(X)$  is ultranorm dense in  $X_2$ . Hence  $U$  preserves the inner product by 148V □

## 2.5.1 Exterior tensor product

Suppose  $X$  is a Hilbert  $\mathcal{A}$ -module and  $Y$  is a Hilbert  $\mathcal{B}$ -module for some  $C^*$ -algebras  $\mathcal{A}$  and  $\mathcal{B}$ . The algebraic tensor product  $X \odot Y$  is a right  $\mathcal{A} \odot \mathcal{B}$ -module via  $(x \otimes y) \cdot (a \otimes b) = (xa) \otimes (yb)$  and has an  $\mathcal{A} \odot \mathcal{B}$ -valued inner product fixed by  $[x \otimes y, x' \otimes y'] = \langle x, x' \rangle \otimes \langle y, y' \rangle$ . Write  $\mathcal{A} \otimes_\sigma \mathcal{B}$  for the spatial  $C^*$ -tensor product [73, §11.3] of  $\mathcal{A}$  and  $\mathcal{B}$ . The inner product on  $X \odot Y$  can be extended to an  $\mathcal{A} \otimes_\sigma \mathcal{B}$ -valued inner product on the norm completion of  $X \odot Y$  on which it is definite [81]. This is a Hilbert  $C^*$ -module, which is called the *exterior tensor product* of  $X$  and  $Y$ . It is quite difficult to perform this construction: Lance discusses the difficulties in [81, ch. 4]. But if we assume  $\mathcal{A}$  and  $\mathcal{B}$  are von Neumann algebras *and* that  $X$  and  $Y$  are self-dual, it is rather easier. In fact, we will directly construct the ultranorm completion of the exterior tensor product, which is a self-dual Hilbert  $\mathcal{A} \otimes \mathcal{B}$ -module, which we call the *self-dual exterior tensor product*. This self-dual exterior tensor product behaves much better than the plain exterior tensor product. As a first example, we will see (in 165VI) that  $\mathcal{B}^a(X) \otimes \mathcal{B}^a(Y) \cong \mathcal{B}^a(X \otimes Y)$  for the self-dual exterior tensor product, which is false in general for the plain exterior tensor product. The self-dual exterior tensor product also appears naturally when computing the Paschke dilation of a tensor product  $\varphi \otimes \psi$ , see 167I. 164

**Theorem** Suppose  $\mathcal{A}$  and  $\mathcal{B}$  are von Neumann algebras. For any self-dual Hilbert  $\mathcal{A}$ -module  $X$  and self-dual Hilbert  $\mathcal{B}$ -module  $Y$ , there is an up-to-isomorphism unique self-dual Hilbert  $\mathcal{A} \otimes \mathcal{B}$ -module  $X \otimes Y$ , called the (self-dual exterior) tensor product, together with a  $\mathcal{A} \odot \mathcal{B}$ -linear injective inner product preserving map  $\eta: X \odot Y \rightarrow X \otimes Y$  with the following universal property. II

For every self-dual Hilbert  $\mathcal{A} \otimes \mathcal{B}$ -module  $Z$  with bounded  $\mathcal{A} \odot \mathcal{B}$ -linear  $T: X \odot Y \rightarrow Z$ , there is a unique bounded  $\mathcal{A} \otimes \mathcal{B}$ -linear

map  $\hat{T}: X \otimes Y \rightarrow Z$  with  $\hat{T} \circ \eta = T$ ,

where we norm  $X \odot Y$  with  $\|t\| = \|[t, t]\|_{\mathcal{A} \otimes \mathcal{B}}^{\frac{1}{2}}$ . Write  $x \hat{\otimes} y = \eta(x \otimes y)$ . For any such  $X \otimes Y$  we have the following.

1. The image of  $X \odot Y$  under  $\eta$  is ultranorm dense in  $X \otimes Y$ .
2. If  $(e_i)_{i \in I}$  is an orthonormal basis of  $X$  and  $(d_j)_{j \in J}$  of  $Y$ , then

(a)  $(e_i \hat{\otimes} d_j)_{i,j \in I \times J}$  is an orthonormal basis of  $X \otimes Y$  and

(b) the linear span of

$$\{ |(e_i a) \hat{\otimes} (d_j b)\rangle \langle e_k \otimes d_l|; a \in \mathcal{A}, b \in \mathcal{B}, i, k \in I, j, l \in J \}$$

is ultraweakly dense in  $\mathcal{B}^a(X \otimes Y)$ .

**III Proof** Pick orthonormal bases  $(e_i)_{i \in I}$  and  $(d_j)_{j \in J}$  of  $X$  and  $Y$  respectively. Write  $p_{ij} \equiv \langle e_i, e_i \rangle \otimes \langle d_j, d_j \rangle \in \mathcal{A} \otimes \mathcal{B}$ . Define  $X \otimes Y \equiv \ell^2((p_{ij})_{i,j \in I \times J})$ , with  $\ell^2$  as in 161 II. This definition of  $X \otimes Y$  seems to depend on the choice of bases  $(e_i)_i$  and  $(d_j)_j$ . We show it does not (up-to-isomorphism), in IX.

**IV (Definition  $\eta$ )** Pick  $x \in X$  and  $y \in Y$ . Write  $x_i \equiv \langle e_i, x \rangle$  and  $y_j \equiv \langle d_j, y \rangle$ . Then

$$\begin{aligned} \langle x, x \rangle \otimes \langle y, y \rangle &= \left( \sum_i x_i^* x_i \right) \otimes \left( \sum_j y_j^* y_j \right) && \text{by Parseval} \\ &= \sum_{i,j} x_i^* x_i \otimes y_j^* y_j \\ &= \sum_{i,j} (x_i \otimes y_j)^* (x_i \otimes y_j). \end{aligned}$$

Hence  $(x_i \otimes y_j)_{i,j}$  is  $\ell^2$ -summable, and so there is a unique  $\mathcal{A} \odot \mathcal{B}$ -linear map  $\eta: X \odot Y \rightarrow X \otimes Y$  fixed by  $\eta(x \otimes y) = (x_i \otimes y_j)_{i,j}$ .

**V ( $\eta$  preserves inner product)** Pick additional  $x' \in X$  and  $y' \in Y$ . Writing  $x'_i = \langle e_i, x' \rangle$  and  $y'_j = \langle d_j, y' \rangle$ , we have

$$\langle \eta(x \otimes y), \eta(x' \otimes y') \rangle = \sum_{i,j} (x_i \otimes y_j)^* \otimes (x'_i \otimes y'_j) = \sum_{i,j} x_i^* x'_i \otimes y_j^* y'_j$$

and

$$\left( \sum_i x_i^* x'_i \right) \otimes \left( \sum_j y_j^* y'_j \right) = \langle x, x' \rangle \otimes \langle y, y' \rangle = \langle x \otimes y, x' \otimes y' \rangle.$$

Thus it is sufficient to show  $\sum_{i,j} x_i^* x'_i \otimes y_j^* y'_j = \left( \sum_i x_i^* x'_i \right) \otimes \left( \sum_j y_j^* y'_j \right)$ , which holds as the product np-functionals are separating by definition, see 108 II. Thus indeed  $\eta$  preserves the inner product.

( $\eta$  is injective) Let  $n \in \mathbb{N}$ ,  $x_1, \dots, x_n \in X$  and  $y_1, \dots, y_n \in Y$  be given such that  $\eta(\sum_l x_l \otimes y_l) = 0$ . By 160X there are orthonormal  $e'_1, \dots, e'_m \in X$  such that  $x_l = \sum_i e'_i \langle e'_i, x_l \rangle$ . Similarly  $y_l = \sum_j d'_j \langle d'_j, y_l \rangle$  for some orthonormal  $d'_1, \dots, d'_m \in Y$ . As  $\eta$  preserves the inner product, the elements  $e'_i \hat{\otimes} d'_j$  are again orthonormal and so from

$$0 = \sum_l x_l \hat{\otimes} y_l = \sum_{i,j} (e'_i \hat{\otimes} d'_j) \left( \sum_l \langle e'_i, x_l \rangle \otimes \langle d'_j, y_l \rangle \right)$$

it follows  $\sum_l \langle e'_i, x_l \rangle \otimes \langle d'_j, y_l \rangle = 0$  for all  $i, j$ . Consequently

$$\sum_l x_l \otimes y_l = \sum_{i,j} (e'_i \otimes d'_j) \left( \sum_l \langle e'_i, x_l \rangle \otimes \langle d'_j, y_l \rangle \right) = 0,$$

which shows  $\eta$  is injective (and that the inner product on  $X \odot Y$  is definite).

(Image  $\eta$  ultranorm dense) For brevity, write  $E \equiv \{e_i \hat{\otimes} d_j; i, j \in I \times J\}$ . By 108II (and 73VIII)  $\mathcal{A} \odot \mathcal{B}$  is ultrastrongly dense in  $\mathcal{A} \otimes \mathcal{B}$ . So  $E(\mathcal{A} \odot \mathcal{B})$  is ultranorm dense in  $E(\mathcal{A} \otimes \mathcal{B})$ , see 148III. In turn the linear span of  $E(\mathcal{A} \otimes \mathcal{B})$  is (by construction) ultranorm dense in  $X \otimes Y$ . Thus the linear span of  $E(\mathcal{A} \odot \mathcal{B})$  is ultranorm dense in  $X \otimes Y$ . Hence the image of  $\eta$ , which contains the linear span of  $E(\mathcal{A} \odot \mathcal{B})$ , is ultranorm dense in  $X \otimes Y$ .

(Universal property) Let  $Z$  be a self-dual Hilbert  $\mathcal{A} \otimes \mathcal{B}$ -module with some bounded  $\mathcal{A} \odot \mathcal{B}$ -linear  $T: X \odot Y \rightarrow Z$ . If  $X \odot Y$  were an  $\mathcal{A} \otimes \mathcal{B}$ -module and both  $\eta$  and  $T$  were  $\mathcal{A} \otimes \mathcal{B}$ -linear, we could simply apply 151Ia to get the desired  $\hat{T}$ . Instead, we will retrace the steps of its proof and modify it for the present situation. Uniqueness of  $\hat{T}$  is the same:  $\hat{T}$  is fixed by the ultranorm-dense image of  $\eta$  as it is bounded  $\mathcal{A} \otimes \mathcal{B}$ -linear and so ultranorm continuous by 148I. The argument for existence of  $\hat{T}$  is more subtle. We will show

$$\langle Tt, Tt \rangle \leq \|T\|^2 [t, t] \quad \text{for all } t \in X \odot Y \quad (2.13)$$

as a replacement for the ultranorm continuity of  $T$ . We cannot apply 144V directly: as  $\mathcal{A} \odot \mathcal{B}$  is not a  $C^*$ -algebra, we cannot construct  $(\langle t, t \rangle + \varepsilon)^{-\frac{1}{2}}$  required for its proof. To prove (2.13), we will work in the norm completions. Write  $\overline{\mathcal{A} \odot \mathcal{B}}$  for the norm-closure of  $\mathcal{A} \odot \mathcal{B}$  in  $\mathcal{A} \otimes \mathcal{B}$ . Clearly  $\overline{\mathcal{A} \odot \mathcal{B}}$  is a  $C^*$ -algebra (in fact it is the spatial  $C^*$ -tensor product of  $\mathcal{A}$  and  $\mathcal{B}$ ). Write  $\overline{X \odot Y}$  for the norm-closure of the image of  $\eta$  in  $X \otimes Y$ . By norm continuity (see 142V) the operations of  $X \otimes Y$  restrict to turn  $\overline{X \odot Y}$  into a Hilbert  $\overline{\mathcal{A} \odot \mathcal{B}}$ -module. As  $T$  is uniformly continuous, there is a unique bounded  $\overline{\mathcal{A} \odot \mathcal{B}}$ -linear  $\overline{T}: \overline{X \odot Y} \rightarrow Z$  with  $\overline{T} \circ \eta = T$  and  $\|T\| = \|\overline{T}\|$ . Now we can apply the proof of 144V to  $\overline{T}$  to find  $\langle \overline{T}s, \overline{T}s \rangle \leq \|\overline{T}\|^2 \langle s, s \rangle$  for  $s \in \overline{X \odot Y}$ . Substituting  $s = \eta(t)$ , we get (2.13).

To define  $\hat{T}$ , pick any  $t \in X \otimes Y$ . As the image of  $\eta$  is ultranorm dense, there

is a net  $(\eta(t_\alpha))_\alpha$  with  $\eta(t_\alpha) \rightarrow t$  ultranorm. For any np-map  $f: \mathcal{A} \otimes \mathcal{B} \rightarrow \mathbb{C}$

$$\begin{aligned} \|T(t_\alpha) - T(t_\beta)\|_f^2 &= f(\langle T(t_\alpha - t_\beta), T(t_\alpha - t_\beta) \rangle) \\ &\leq \|T\|^2 f([t_\alpha - t_\beta, t_\alpha - t_\beta]) \\ &= \|T\|^2 f(\langle \eta(t_\alpha) - \eta(t_\beta), \eta(t_\alpha) - \eta(t_\beta) \rangle) \rightarrow 0. \end{aligned}$$

So  $(Tt_\alpha)_\alpha$  is ultranorm Cauchy in  $Z$ . With a similar argument we see  $(Tt_\alpha)_\alpha$  is equivalent to  $(Tt'_\alpha)_\alpha$  whenever  $(\eta t_\alpha)_\alpha$  is equivalent to  $(\eta t'_\alpha)_\alpha$ . So we may define

$$\hat{T}t = \operatorname{unlim}_\alpha Tt_\alpha.$$

Clearly  $\hat{T} \circ \eta = T$ . With the same argument as for 151 la, we see  $\hat{T}$  is bounded and additive. To show  $\hat{T}$  is  $\mathcal{A} \otimes \mathcal{B}$ -linear, pick any  $b \in \mathcal{A} \otimes \mathcal{B}$ . There is a net  $b_\beta$  in  $\mathcal{A} \odot \mathcal{B}$  with  $b_\beta \rightarrow b$  ultrastrongly. If  $\hat{T}$  is ultranorm continuous, we are done:

$$(\hat{T}t)b = \operatorname{unlim}_\beta (\operatorname{unlim}_\alpha Tt_\alpha)b_\beta = \operatorname{unlim}_\beta \operatorname{unlim}_\alpha T(t_\alpha b_\beta) = \operatorname{unlim}_\beta \hat{T}(tb_\beta) = \hat{T}(tb).$$

To show  $\hat{T}$  is ultranorm continuous, assume  $t^\alpha \rightarrow 0$  ultranorm in  $X \otimes Y$ . It is sufficient to show  $\hat{T}t^\alpha \rightarrow 0$  ultranorm. Let  $f: \mathcal{A} \otimes \mathcal{B} \rightarrow \mathbb{C}$  be any np-map. Write  $\Phi$  for the set of entourages for ultranorm uniformity on  $X \otimes Y$ . As the image of  $\eta$  is ultranorm dense, there is a  $\Phi$ -indexed Cauchy net  $(t_E^\alpha)_{E \in \Phi}$  in  $\eta(X \odot Y)$  with  $t_E^\alpha \in E$  for each  $E \in \Phi$ . Thus  $\hat{T}t^\alpha = \operatorname{unlim}_E Tt_E^\alpha$ . As  $t^\alpha \rightarrow 0$  there is some  $\alpha_0$  such that for all  $\alpha \geq \alpha_0$  we have  $\|t^\alpha\|_f \leq \varepsilon$ . We can find  $E_0 \geq E_{f, \varepsilon}$  such that for all  $E \geq E_0$  we have  $\|\hat{T}t^{\alpha_0} - Tt_E^{\alpha_0}\|_f \leq \varepsilon$ . For  $E \geq E_0$  we have  $E \geq E_{f, \varepsilon}$  and so

$$\|Tt_{E_0}^{\alpha_0}\|_f \stackrel{(2.13)}{\leq} \|T\| \|t_{E_0}^{\alpha_0}\|_f \leq \|T\| (\|t_{E_0}^{\alpha_0} - t^{\alpha_0}\|_f + \|t^{\alpha_0}\|_f) \leq 2\|T\|\varepsilon.$$

Consequently

$$\|\hat{T}t^{\alpha_0}\|_f \leq \|\hat{T}t^{\alpha_0} - Tt_{E_0}^{\alpha_0}\|_f + \|Tt_{E_0}^{\alpha_0}\|_f \leq \varepsilon + 2\|T\|\varepsilon.$$

So  $\hat{T}t^\alpha \rightarrow 0$  ultranorm, as desired.

- IX (Uniqueness) Assume there is a self-dual Hilbert  $\mathcal{A} \otimes \mathcal{B}$ -module  $X \otimes_2 Y$  together with a injective inner product-preserving  $\mathcal{A} \odot \mathcal{B}$ -linear map  $\eta_2: X \odot Y \rightarrow X \otimes_2 Y$  also satisfying the universal property. With the same reasoning as in 163II there is an invertible bounded  $\mathcal{A} \otimes \mathcal{B}$ -linear  $U: X \otimes Y \rightarrow X \otimes_2 Y$  with  $U \circ \eta = \eta_2$ . Clearly  $\langle Ux, Uy \rangle = \langle x, y \rangle$  for  $x, y \in E(\mathcal{A} \odot \mathcal{B})$ , where we defined  $E = \{e_i \hat{\otimes} d_j; i, j \in I \times J\}$ . As we saw the linear span of  $E(\mathcal{A} \odot \mathcal{B})$  is ultranorm dense in  $X \otimes Y$ , we see  $U$  must preserve the inner product by 148V.

Property 1 from the statement of the Theorem follows immediately from the fact that the exterior tensor product is unique up to an isomorphism which respects the embeddings. Assume  $(e'_i)_{i \in I'}$  is an orthonormal basis of  $X$  and  $(d'_j)_{j \in J'}$  is an orthonormal basis of  $Y$ . We will show  $E_2 = \{e'_i \hat{\otimes} d'_j; i, j \in I' \times J'\}$  is an orthonormal basis of  $X \hat{\otimes} Y$ . Clearly  $E_2$  is orthonormal. By 160 IX and 160 IV it is sufficient to show  $e_{i_0} \hat{\otimes} d_{j_0} \in E_2^{\perp\perp}$  for all  $i_0 \in I$  and  $j_0 \in J$ . Write  $a_i = \langle e'_i, e_{i_0} \rangle$  and  $b_j = \langle d'_j, d_{j_0} \rangle$ . By 160 IX and V it suffices to show the latter equality in X

$$\left( \sum_i a_i^* a_i \right) \otimes \left( \sum_j b_j^* b_j \right) = \langle e_{i_0} \hat{\otimes} d_{j_0}, e_{i_0} \hat{\otimes} d_{j_0} \rangle = \sum_{i,j} a_i^* a_i \otimes b_j^* b_j,$$

which holds as product np-functionals are separating by definition, see 108 II.

Finally we prove that the linear span of XI

$$D \equiv \{ |(e'_i a) \hat{\otimes} (d'_j b) \langle e_k \otimes d_l |; a \in \mathcal{A}, b \in \mathcal{B}, i, k \in I', j, l \in J' \}$$

is ultraweakly dense in  $\mathcal{B}^a(X \otimes Y)$ . By 159 IV, the linear span of (the set of operators of the form)  $|(e'_i \hat{\otimes} d'_j) t \langle e'_k \hat{\otimes} d'_l |$  with  $t \in \mathcal{A} \otimes \mathcal{B}$  is ultraweakly dense in  $\mathcal{B}^a(X \otimes Y)$ . We will show that  $t \in \mathcal{A} \odot \mathcal{B}$  suffices. By 108 II, 73 VIII and 74 IV, each  $t \in \mathcal{A} \otimes \mathcal{B}$  is the ultrastrong limit of a norm bounded net  $t_\alpha$  in  $\mathcal{A} \odot \mathcal{B}$ . So by 159 IX (and 148 III), we see  $|(e'_i \hat{\otimes} d'_j) t_\alpha \langle e'_k \hat{\otimes} d'_l |$  converges ultraweakly to  $|(e'_i \hat{\otimes} d'_j) t \langle e'_k \hat{\otimes} d'_l |$ . So indeed, the linear span of  $D$  is ultraweakly dense in  $\mathcal{B}^a(X \otimes Y)$ . □

**Examples** We give a few examples of the self-dual exterior tensor product and relate it to other notions of tensor product. To avoid confusion, we will denote the self-dual exterior tensor product by  $\otimes_{\text{ext}}$  (instead instead of the plain  $\otimes$ ). XII

1. Any Hilbert space is a self-dual Hilbert  $\mathbb{C}$  module. For Hilbert spaces  $\mathcal{H}$  and  $\mathcal{K}$  we have  $\mathcal{H} \otimes_{\text{Hilb}} \mathcal{K} \cong \mathcal{H} \otimes_{\text{ext}} \mathcal{K}$ , where  $\otimes_{\text{Hilb}}$  denotes the regular tensor product of Hilbert spaces.
2. Every von Neumann algebra is a self-dual Hilbert  $\mathbb{C}^*$ -module over itself. For any von Neumann algebras  $\mathcal{A}$  and  $\mathcal{B}$  we have  $\mathcal{A} \otimes_{\text{vN}} \mathcal{B} \cong \mathcal{A} \otimes_{\text{ext}} \mathcal{B}$  as Hilbert  $\mathcal{A} \otimes_{\text{vN}} \mathcal{B}$ -modules, where  $\mathcal{A} \otimes_{\text{vN}} \mathcal{B}$  denotes the (spatial) tensor product of von Neumann algebras.
3. Let  $X$  be a self-dual Hilbert  $\mathcal{A}$ -module and  $Y$  be a self-dual Hilbert  $\mathcal{B}$ -module for von Neumann algebras  $\mathcal{A}, \mathcal{B}$ . The norm closure of  $X \odot Y$  in  $X \otimes_{\text{ext}} Y$  is the plain exterior tensor product of  $X$  and  $Y$ . In turn, the ultranorm completion of the exterior tensor product of  $X$  and  $Y$  is isomorphic to  $X \otimes_{\text{ext}} Y$ , the self-dual exterior tensor product.

4. In 167I it will be shown that

$$(\mathcal{A}_1 \otimes_{\varphi_1} \mathcal{B}_1) \otimes_{\text{ext}} (\mathcal{A}_2 \otimes_{\varphi_2} \mathcal{B}_2) \cong (\mathcal{A}_1 \otimes \mathcal{A}_2) \otimes_{\varphi_1 \otimes \varphi_2} (\mathcal{B}_1 \otimes \mathcal{B}_2)$$

for all ncp-maps  $\varphi_i: \mathcal{A}_i \rightarrow \mathcal{B}_i$  ( $i = 1, 2$ ) between von Neumann algebras.

**165** For Hilbert spaces  $\mathcal{H}$  and  $\mathcal{K}$ , we have  $\mathcal{B}(\mathcal{H} \otimes \mathcal{K}) \cong \mathcal{B}(\mathcal{H}) \otimes \mathcal{B}(\mathcal{K})$ . However, for Hilbert modules  $X$  and  $Y$  and the regular (not necessarily self-dual) exterior tensor product it is not true in general that  $\mathcal{B}^a(X) \otimes \mathcal{B}^a(Y) \cong \mathcal{B}^a(X \otimes Y)$ . However, if  $X$  and  $Y$  are self-dual Hilbert modules over von Neumann algebras, then we do have the familiar formula for the self-dual exterior tensor product. We need some preparation.

**II Setting** Let  $X$  be a self-dual Hilbert  $\mathcal{A}$ -module and  $Y$  be a self-dual Hilbert  $\mathcal{B}$ -module for von Neumann algebras  $\mathcal{A}$  and  $\mathcal{B}$ .

**III Proposition** For any operators  $S \in \mathcal{B}^a(X)$  and  $T \in \mathcal{B}^a(Y)$ , there is a unique operator  $S \otimes T \in \mathcal{B}^a(X \otimes Y)$  fixed by  $(S \otimes T)(x \otimes y) = (Sx) \otimes (Ty)$  for any  $x \in X$  and  $y \in Y$ .

**IV Proof** We will define  $S \otimes T$  using the universal property proven in 164II. Write  $\Theta: X \odot Y \rightarrow X \otimes Y$  for the map  $\Theta(x \otimes y) = (Sx) \otimes (Ty)$ . It is easy to see  $\Theta$  is  $\mathcal{A} \odot \mathcal{B}$ -linear. The hard part is to show that  $\Theta$  is bounded. Pick any  $t \in X \odot Y$  — say  $t \equiv \sum_{i=1}^n x_i \otimes y_i$ . To start, note  $(\langle Sx_i, Sx_j \rangle)_{ij}$  is positive in  $M_n \mathcal{A}$  — indeed, for any  $a_1, \dots, a_n \in \mathcal{A}$  we have  $\sum_{i,j} a_i^* \langle Sx_i, Sx_j \rangle a_j = \langle Sx, Sx \rangle \geq 0$  where  $x \equiv \sum_i x_i a_i$  and so it is positive by 33II. By reasoning in the same way for  $\sqrt{\|S\|^2 - S^*S}$  instead of  $S$ , it follows  $(\langle Sx_i, Sy_i \rangle)_{ij} \leq (\|S\|^2 \langle x_i, x_j \rangle)_{ij}$ . Similarly  $0 \leq (\langle Ty_i, Ty_j \rangle)_{ij} \leq (\|T\|^2 \langle y_i, y_j \rangle)_{ij}$  in  $M_n \mathcal{B}$ . Write  $s: (\mathcal{A} \otimes \mathcal{B})^{\oplus n} \rightarrow \mathcal{A} \otimes \mathcal{B}$  for the  $\mathcal{A} \otimes \mathcal{B}$ -linear map given by  $s(t_1, \dots, t_n) = t_1 + \dots + t_n$ . ( $s$  is the row vector filled with 1s). It is easy to see  $sAs^*1 = \sum_{i,j} A_{ij}$  for any matrix  $A$  over  $\mathcal{A} \otimes \mathcal{B}$ . As a final ingredient, recall that the bilinear map  $M_n(\otimes): M_n \mathcal{A} \times M_n \mathcal{B} \rightarrow M_n(\mathcal{A} \otimes \mathcal{B})$  defined in the obvious way is positive by 113IV and 108II. Putting everything together:

$$\begin{aligned} \langle \Theta t, \Theta t \rangle &= \sum_{i,j} \langle (Sx_i) \otimes (Ty_i), (Sx_j) \otimes (Ty_j) \rangle \\ &= \sum_{i,j} \langle Sx_i, Sx_j \rangle \otimes \langle Ty_i, Ty_j \rangle \\ &= s (\langle Sx_i, Sx_j \rangle \otimes \langle Ty_i, Ty_j \rangle)_{ij} s^* 1 \\ &= s M_n(\otimes) ((\langle Sx_i, Sx_j \rangle)_{ij}, (\langle Ty_i, Ty_j \rangle)_{ij}) s^* 1 \\ &\leq s M_n(\otimes) (\|S\|^2 \langle x_i, x_j \rangle)_{ij}, (\|T\|^2 \langle y_i, y_j \rangle)_{ij}) s^* 1 \\ &= \|S\|^2 \|T\|^2 \langle t, t \rangle. \end{aligned}$$

We have proven that  $\Theta$  is bounded (by  $\|S\|\|T\|$ ). Thus by 164 II, there exists a unique map  $S \otimes T: X \otimes Y \rightarrow X \otimes Y$  with  $(S \otimes T)(x \otimes y) = \Theta(x \otimes y) \equiv (Sx) \otimes (Ty)$ , which is what we desired to prove.  $\square$

**Exercise** Prove that for the operation  $\otimes$  defined in III, we have V

1.  $|x_1\rangle\langle x_2| \otimes |y_1\rangle\langle y_2| = |x_1 \otimes y_1\rangle\langle x_2 \otimes y_2|$ ,
2.  $1 \otimes 1 = 1$ ;
3.  $(S \otimes T)(S' \otimes T') = (SS') \otimes (TT')$  and
4.  $(S \otimes T)^* = S^* \otimes T^*$ .

(Hint: use that by the reasoning of 164 VII the linear span of

$$\{(ea) \otimes (db); (a, b, e, d) \in \mathcal{A} \times \mathcal{B} \times E \times D\}$$

is ultranorm dense in  $X \otimes Y$  for any bases  $E$  of  $X$  and  $D$  of  $Y$ .)

**Theorem** There is an nmiu-isomorphism  $\vartheta: \mathcal{B}^a(X) \otimes \mathcal{B}^a(Y) \cong \mathcal{B}^a(X \otimes Y)$  VI  
fixed by  $\vartheta(S \otimes T) = S \otimes T$ .

**Proof** We will show that the bilinear map  $\Theta: \mathcal{B}^a(X) \times \mathcal{B}^a(Y) \rightarrow \mathcal{B}^a(X \otimes Y)$  VII  
given by  $\Theta(S, T) = S \otimes T$  is a tensor product in the sense of 108 II using 116 VII.  
From this it follows by 114 II, that there is a nmiu-isomorphism

$$\vartheta: \mathcal{B}^a(X) \otimes \mathcal{B}^a(Y) \cong \mathcal{B}^a(X \otimes Y)$$

with  $\vartheta(S \otimes T) = \Theta(S, T) = S \otimes T$ , as promised.

From V, it follows  $\Theta$  is an miu-bilinear map. By 164 II the linear span of operators of the form  $|e_i a\rangle\langle e_k| \otimes |d_j b\rangle\langle d_l| = |(e_i a) \otimes (d_j b)\rangle\langle e_k \otimes d_l|$  is ultraweakly dense in  $\mathcal{B}^a(X \otimes Y)$  for an orthonormal basis  $e_i \otimes d_j$  of  $X \otimes Y$ . So the image of  $\Theta$  generates  $\mathcal{B}^a(X \otimes Y)$ . VIII

It remains to be shown that there are sufficiently many product functionals (see 108 II for the definition of product functional and their sufficiency). Write IX

$$\begin{aligned}\Omega_X &\equiv \{f(\langle x, (\cdot)x \rangle); f: \mathcal{A} \rightarrow \mathbb{C} \text{ np-map}, x \in X\} \\ \Omega_Y &\equiv \{g(\langle y, (\cdot)y \rangle); g: \mathcal{B} \rightarrow \mathbb{C} \text{ np-map}, y \in Y\}.\end{aligned}$$

From 144 I it follows  $\Omega_X$  and  $\Omega_Y$  are order separating. For  $\sigma \in \Omega_X$  and  $\tau \in \Omega_Y$  — say  $\sigma \equiv f(\langle x, (\cdot)x \rangle)$ ,  $\tau \equiv g(\langle y, (\cdot)y \rangle)$  — define  $\sigma \otimes \tau \equiv (f \otimes g)(x \otimes y, (\cdot)x \otimes y)$  and  $\Omega \equiv \{\sigma \otimes \tau; \sigma \in \Omega_X, \tau \in \Omega_Y\}$ . We will show that  $\Omega$  is faithful and that  $(\sigma \otimes \tau)(\Theta(S, T)) = \sigma(S)\tau(T)$  for all  $S \in \mathcal{B}^a(X)$  and  $T \in \mathcal{B}^a(Y)$ , which is sufficient to show  $\Theta$  is a tensor product. Indeed for any  $S \in \mathcal{B}^a(X)$  and

$T \in \mathcal{B}^a(Y)$ , we have

$$\begin{aligned}
 (\sigma \otimes \tau)(\Theta(S, T)) &= (f \otimes g)(\langle x \otimes y, (S \otimes T)(x \otimes y) \rangle) \\
 &= (f \otimes g)(\langle x \otimes y, (Sx) \otimes (Ty) \rangle) \\
 &= (f \otimes g)(\langle x, Sx \rangle \otimes \langle y, Ty \rangle) \\
 &= f(\langle x, Sx \rangle) \cdot g(\langle y, Ty \rangle) \\
 &= \sigma(S)\tau(T).
 \end{aligned}$$

- × Finally, to show that  $\Omega$  is faithful, assume  $A \in \mathcal{B}^a(X \otimes Y)$ ,  $A \geq 0$  and  $(\sigma \otimes \tau)(A) = 0$  for all  $\sigma \otimes \tau \in \Omega$ . Then for all  $x \in X$  and  $y \in Y$ , we have  $\langle x \otimes y, A(x \otimes y) \rangle = 0$ , thus  $\|\sqrt{A}(x \otimes y)\|^2 = 0$ , hence  $\sqrt{A}(x \otimes y) = 0$ . So  $\sqrt{A}$  vanishes on  $X \otimes Y$ , which is ultranorm dense in  $X \otimes Y$ , whence  $\sqrt{A} = 0$ . So  $A = 0$ .  $\square$

**166** To compute the tensor product of Paschke dilations, we will need to know a bit more about the ultranorm continuity of the exterior tensor product.

- II **Lemma** Let  $X$  be a self-dual Hilbert  $\mathcal{A}$ -module and  $Y$  be a self-dual Hilbert  $\mathcal{B}$ -module for von Neumann algebras  $\mathcal{A}$  and  $\mathcal{B}$ . If  $x_\alpha \rightarrow x$  and  $y_\alpha \rightarrow y$  ultranorm for norm-bounded nets  $x_\alpha$  in  $X$  and  $y_\alpha$  in  $Y$ , then  $x_\alpha \otimes y_\alpha \rightarrow x \otimes y$  ultranorm.
- III **Proof** As  $x_\alpha \otimes y_\alpha - x \otimes y = (x_\alpha - x) \otimes y_\alpha + x \otimes (y_\alpha - y)$ , it is sufficient to show that both  $(x_\alpha - x) \otimes y_\alpha \rightarrow 0$  and  $x \otimes (y_\alpha - y) \rightarrow 0$  ultranorm. Clearly

$$\begin{aligned}
 \langle (x_\alpha - x) \otimes y_\alpha, (x_\alpha - x) \otimes y_\alpha \rangle &= \langle x_\alpha - x, x_\alpha - x \rangle \otimes \langle y_\alpha, y_\alpha \rangle \\
 &= (\langle x_\alpha - x, x_\alpha - x \rangle \otimes 1) \cdot (1 \otimes \langle y_\alpha, y_\alpha \rangle).
 \end{aligned}$$

The maps  $b \mapsto 1 \otimes b$  and  $a \mapsto a \otimes 1$  are ncp so both  $1 \otimes \langle y_\alpha, y_\alpha \rangle$  and  $\langle x_\alpha - x, x_\alpha - x \rangle \otimes 1$  are bounded nets of positive elements. Also  $\langle x_\alpha - x, x_\alpha - x \rangle \otimes 1 \rightarrow 0$  as  $x_\alpha \rightarrow x$  ultranorm. Thus 44 III applies and we see  $(\langle x_\alpha - x, x_\alpha - x \rangle \otimes 1) \cdot (1 \otimes \langle y_\alpha, y_\alpha \rangle) \rightarrow 0$  ultraweakly, hence  $(x_\alpha - x) \otimes y_\alpha \rightarrow 0$  ultranorm. With a similar argument one shows  $x \otimes (y_\alpha - y) \rightarrow 0$  ultranorm.  $\square$

- IV **Lemma** Let  $X$  be a self-dual Hilbert  $\mathcal{A}$ -module and  $Y$  be a self-dual Hilbert  $\mathcal{B}$ -module for von Neumann algebras  $\mathcal{A}$  and  $\mathcal{B}$ . Suppose  $U \subseteq X$  and  $V \subseteq Y$  are ultranorm-dense submodules. Then the linear span of

$$U \otimes V \equiv \{u \otimes v; u \in U, v \in V\}$$

is ultranorm dense in  $X \otimes Y$ .

- V **Proof** It is sufficient to show that every  $x \otimes y \in X \otimes Y$  is the ultranorm limit of elements from  $U \otimes V$  as the linear span of elements of the form  $x \otimes y$  is ultranorm dense in  $X \otimes Y$ . So pick any element of the form  $x \otimes y \in X \otimes Y$ . By the Kaplansky density theorem for Hilbert  $C^*$ -modules, see 158 II, there are norm-bounded nets  $x_\alpha$  in  $U$  and  $y_\alpha$  in  $V$  with  $x_\alpha \rightarrow x$  and  $y_\alpha \rightarrow y$  ultranorm. So by II, we see  $x_\alpha \otimes y_\alpha \rightarrow x \otimes y$ .  $\square$

**Lemma** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  be an ncp-map between von Neumann algebras together with ultrastrongly dense  $*$ -subalgebras  $\mathcal{A}' \subseteq \mathcal{A}$  and  $\mathcal{B}' \subseteq \mathcal{B}$ . Then: the linear span of  $\mathcal{T} \equiv \{a \otimes b; a \in \mathcal{A}', b \in \mathcal{B}'\}$  is ultranorm dense in  $\mathcal{A} \otimes_{\varphi} \mathcal{B}$ , which was defined in 154 III. VI

**Proof** It is sufficient to show that any  $a \otimes b \in \mathcal{A} \otimes_{\varphi} \mathcal{B}$  is the ultranorm limit of elements from  $\mathcal{T}$ . Pick any  $a \otimes b \in \mathcal{A} \otimes_{\varphi} \mathcal{B}$ . By 74 VI there are norm-bounded nets  $a_{\alpha}$  in  $\mathcal{A}'$  and  $b_{\alpha}$  in  $\mathcal{B}'$  with  $a_{\alpha} \rightarrow a$  and  $b_{\alpha} \rightarrow b$  ultrastrongly. Say  $\|b_{\alpha}\|, \|a_{\alpha}\| \leq B$  for some  $B > 0$ . We will show  $a_{\alpha} \otimes b_{\alpha} - a \otimes b \rightarrow 0$  ultranorm. As usual, we split it into two: VII

$$a_{\alpha} \otimes b_{\alpha} - a \otimes b = a_{\alpha} \otimes (b_{\alpha} - b) + (a_{\alpha} - a) \otimes b.$$

For the first term, note

$$\begin{aligned} \langle a_{\alpha} \otimes (b_{\alpha} - b), a_{\alpha} \otimes (b_{\alpha} - b) \rangle &= (b_{\alpha} - b)^* \varphi(a_{\alpha}^* a_{\alpha}) (b_{\alpha} - b) \\ &\leq B^2 \|\varphi\| (b_{\alpha} - b)^* (b_{\alpha} - b), \end{aligned}$$

which converges to 0 ultraweakly as  $b_{\alpha} \rightarrow b$  ultrastrongly. The second

$$\langle (a_{\alpha} - a) \otimes b, (a_{\alpha} - a) \otimes b \rangle = b^* \varphi((a_{\alpha} - a)^* (a_{\alpha} - a)) b,$$

converges ultraweakly to 0 as  $a_{\alpha} \rightarrow a$  ultrastrongly and  $b^* \varphi(\cdot) b$  is normal and  $(a_{\alpha} - a)^* (a_{\alpha} - a)$  converges ultraweakly to 0 as  $a_{\alpha} \rightarrow a$  ultrastrongly. □

**Theorem** Suppose  $\varphi_i: \mathcal{A}_i \rightarrow \mathcal{B}_i$  is an ncp-map between von Neumann algebras with Paschke dilation  $(\mathcal{P}_i, \varrho_i, h_i)$  for  $i = 1, 2$ . Then  $(\mathcal{P}_1 \otimes \mathcal{P}_2, \varrho_1 \otimes \varrho_2, h_1 \otimes h_2)$  is a Paschke dilation of  $\varphi_1 \otimes \varphi_2$ . Furthermore 167

$$(\mathcal{A}_1 \otimes_{\varphi_1} \mathcal{B}_1) \otimes (\mathcal{A}_2 \otimes_{\varphi_2} \mathcal{B}_2) \cong (\mathcal{A}_1 \otimes \mathcal{A}_2) \otimes_{\varphi_1 \otimes \varphi_2} (\mathcal{B}_1 \otimes \mathcal{B}_2). \quad (2.14)$$

**Proof** We will prove (2.14) first. We proceed as follows. To start, we prove that  $(\mathcal{A}_1 \odot \mathcal{B}_1) \odot (\mathcal{A}_2 \odot \mathcal{B}_2)$  (with the obvious inclusion) is ultranorm dense in  $(\mathcal{A}_1 \otimes_{\varphi_1} \mathcal{B}_1) \otimes (\mathcal{A}_2 \otimes_{\varphi_2} \mathcal{B}_2)$ . Then we show that  $(\mathcal{A}_1 \odot \mathcal{A}_2) \odot (\mathcal{B}_1 \odot \mathcal{B}_2)$  is ultranorm dense in  $(\mathcal{A}_1 \otimes \mathcal{A}_2) \otimes_{\varphi_1 \otimes \varphi_2} (\mathcal{B}_1 \otimes \mathcal{B}_2)$ . We continue and show that the natural bijection  $U_0: (\mathcal{A}_1 \odot \mathcal{B}_1) \odot (\mathcal{A}_2 \odot \mathcal{B}_2) \cong (\mathcal{A}_1 \odot \mathcal{A}_2) \odot (\mathcal{B}_1 \odot \mathcal{B}_2)$  preserves the induced inner products and so extends uniquely to an isomorphism (2.14). Using this isomorphism, we prove that the tensor product of the standard Paschke dilations is a dilation of the tensor product. From this, we finally derive the stated result. II

(Density) By construction,  $\mathcal{A}_1 \odot \mathcal{B}_1$  is an ultranorm-dense  $\mathcal{B}_1$ -submodule of  $\mathcal{A}_1 \otimes_{\varphi_1} \mathcal{B}_1$ . Similarly  $\mathcal{A}_2 \odot \mathcal{B}_2$  is ultranorm dense in  $\mathcal{A}_2 \otimes_{\varphi_2} \mathcal{B}_2$ . Thus by 166 IV  $(\mathcal{A}_1 \odot \mathcal{B}_1) \odot (\mathcal{A}_2 \odot \mathcal{B}_2)$  is ultranorm dense in  $(\mathcal{A}_1 \otimes_{\varphi_1} \mathcal{B}_1) \otimes (\mathcal{A}_2 \otimes_{\varphi_2} \mathcal{B}_2)$ . By 108 II (with 73 VIII and 74 VI), we know  $\mathcal{A}_1 \odot \mathcal{A}_2$  is an ultrastrongly-dense III

subalgebra of  $\mathcal{A}_1 \otimes \mathcal{A}_2$ . Similarly  $\mathcal{B}_1 \odot \mathcal{B}_2$  is ultrastrongly dense in  $\mathcal{B}_1 \otimes \mathcal{B}_2$ . Hence  $(\mathcal{A}_1 \odot \mathcal{A}_2) \odot (\mathcal{B}_1 \odot \mathcal{B}_2)$  is ultranorm dense in  $(\mathcal{A}_1 \otimes \mathcal{A}_2) \otimes_{\varphi_1 \otimes \varphi_2} (\mathcal{B}_1 \otimes \mathcal{B}_2)$  by 166 VI,

IV (Inner product preservation) Clearly

$$\begin{aligned} & \langle (a_1 \otimes b_1) \otimes (a_2 \otimes b_2), (\alpha_1 \otimes \beta_1) \otimes (\alpha_2 \otimes \beta_2) \rangle \\ &= \langle a_1 \otimes b_1, \alpha_1 \otimes \beta_1 \rangle \otimes \langle a_2 \otimes b_2, \alpha_2 \otimes \beta_2 \rangle \\ &= (b_1^* \varphi_1(a_1^* \alpha_1) \beta_1) \otimes (b_2^* \varphi_2(a_2^* \alpha_2) \beta_2) \\ &= (b_1 \otimes b_2)^* (\varphi_1 \otimes \varphi_2)((a_1 \otimes a_2)^* (\alpha_1 \otimes \alpha_2)) (\beta_1 \otimes \beta_2) \\ &= \langle (a_1 \otimes a_2) \otimes (b_1 \otimes b_2), (\alpha_1 \otimes \alpha_2) \otimes (\beta_1 \otimes \beta_2) \rangle; \end{aligned}$$

so  $U_0: (\mathcal{A}_1 \odot \mathcal{B}_1) \odot (\mathcal{A}_2 \odot \mathcal{B}_2) \cong (\mathcal{A}_1 \odot \mathcal{A}_2) \odot (\mathcal{B}_1 \odot \mathcal{B}_2)$  preserves the inner product by linearity.

V As  $U_0$  preserves the inner product, it is ultranorm continuous. So with the usual reasoning, there is a unique bounded  $\mathcal{B}_1 \odot \mathcal{B}_2$ -linear extension

$$U_1: (\mathcal{A}_1 \otimes_{\varphi_1} \mathcal{B}_1) \odot (\mathcal{A}_2 \otimes_{\varphi_2} \mathcal{B}_2) \rightarrow (\mathcal{A}_1 \otimes \mathcal{A}_2) \otimes_{\varphi_1 \otimes \varphi_2} (\mathcal{B}_1 \otimes \mathcal{B}_2).$$

In turn, by the universal property of the self-dual exterior tensor product, there exists a unique bounded  $\mathcal{B}_1 \otimes \mathcal{B}_2$ -linear extension

$$U: (\mathcal{A}_1 \otimes_{\varphi_1} \mathcal{B}_1) \otimes (\mathcal{A}_2 \otimes_{\varphi_2} \mathcal{B}_2) \rightarrow (\mathcal{A}_1 \otimes \mathcal{A}_2) \otimes_{\varphi_1 \otimes \varphi_2} (\mathcal{B}_1 \otimes \mathcal{B}_2).$$

By the ultranorm density III and 148 V we know  $U$  also preserves the inner product. Also from III, it follows that  $U$  is surjective. Thus  $U$  is an isomorphism with  $U^{-1} = U^*$ . We have shown (2.14). By 165 VI there is a unique nmiu-isomorphism

$$\vartheta: \mathcal{B}^a(\mathcal{A}_1 \otimes_{\varphi_1} \mathcal{B}_1) \otimes \mathcal{B}^a(\mathcal{A}_2 \otimes_{\varphi_2} \mathcal{B}_2) \rightarrow \mathcal{B}^a((\mathcal{A}_1 \otimes_{\varphi_1} \mathcal{B}_1) \otimes (\mathcal{A}_2 \otimes_{\varphi_2} \mathcal{B}_2))$$

fixed by  $\vartheta(S \otimes T) = S \otimes T$ . Let  $(\mathcal{B}^a((\mathcal{A}_1 \otimes \mathcal{A}_2) \otimes_{\varphi_1 \otimes \varphi_2} (\mathcal{B}_1 \otimes \mathcal{B}_2)), \varrho, h)$  and  $(\mathcal{B}^a(\mathcal{A}_i \otimes_{\varphi_i} \mathcal{B}_i), \varrho_i, h_i)$  (for  $i = 1, 2$ ) denote the Paschke dilations of  $\varphi_1 \otimes \varphi_2$  and  $\varphi_i$ , respectively, constructed in 154 III.

VI Our next goal is to show

$$\text{ad}_{U^*} \circ \vartheta \circ (\varrho_1 \otimes \varrho_2) = \varrho \quad h \circ \text{ad}_{U^*} \circ \vartheta = h_1 \otimes h_2, \quad (2.15)$$

which is sufficient to show that  $(\mathcal{B}^a(\mathcal{A}_1 \otimes_{\varphi_1} \mathcal{B}_1) \otimes \mathcal{B}^a(\mathcal{A}_2 \otimes_{\varphi_2} \mathcal{B}_2), \varrho_1 \otimes \varrho_2, h_1 \otimes h_2)$

is a Paschke dilation of  $\varphi_1 \otimes \varphi_2$ . We start with the equation on the left.

$$\begin{aligned}
& ((\text{ad}_{U^*} \circ \vartheta \circ (\varrho_1 \otimes \varrho_2))(\alpha_1 \otimes \alpha_2)) (a_1 \otimes a_2) \otimes (b_1 \otimes b_2) \\
&= ((\text{ad}_{U^*} \circ \vartheta)(\varrho_1(\alpha_1) \otimes \varrho_2(\alpha_2))) (a_1 \otimes a_2) \otimes (b_1 \otimes b_2) \\
&= U(\varrho_1(\alpha_1) \otimes \varrho_2(\alpha_2)) U^* (a_1 \otimes a_2) \otimes (b_1 \otimes b_2) \\
&= U(\varrho_1(\alpha_1) \otimes \varrho_2(\alpha_2)) (a_1 \otimes b_1) \otimes (a_2 \otimes b_2) \\
&= U((\alpha_1 a_1) \otimes b_1) \otimes ((\alpha_2 a_2) \otimes b_2) \\
&= ((\alpha_1 \otimes \alpha_2) \cdot (a_1 \otimes a_2)) \otimes (b_1 \otimes b_2) \\
&= \varrho(\alpha_1 \otimes \alpha_2) (a_1 \otimes a_2) \otimes (b_1 \otimes b_2).
\end{aligned}$$

As the linear span of elements of the form  $(a_1 \otimes a_2) \otimes (b_1 \otimes b_2)$  is ultranorm dense in  $(\mathcal{A}_1 \otimes \mathcal{A}_2) \otimes_{\varphi_1 \otimes \varphi_2} (\mathcal{B}_1 \otimes \mathcal{B}_2)$ , we see  $(\text{ad}_{U^*} \circ \vartheta \circ (\varrho_1 \otimes \varrho_2))(\alpha_1 \otimes \alpha_2) = \varrho(\alpha_1 \otimes \alpha_2)$  for all  $\alpha_1 \in \mathcal{A}_1$  and  $\alpha_2 \in \mathcal{A}_2$ . In turn, as the linear span of elements of the form  $\alpha_1 \otimes \alpha_2$  is ultrastrongly dense in  $\mathcal{A}_1 \otimes \mathcal{A}_2$  and ncp-maps are ultrastrongly continuous, we see  $\text{ad}_{U^*} \circ \vartheta \circ (\varrho_1 \otimes \varrho_2) = \varrho$ . We continue with the equation on the right of (2.15).

$$\begin{aligned}
(h \circ \text{ad}_{U^*} \circ \vartheta)(T \otimes S) &= \langle (1 \otimes 1) \otimes (1 \otimes 1), U(T \otimes S) U^*(1 \otimes 1) \otimes (1 \otimes 1) \rangle \\
&= \langle U^*(1 \otimes 1) \otimes (1 \otimes 1), (T \otimes S) U^*(1 \otimes 1) \otimes (1 \otimes 1) \rangle \\
&= \langle (1 \otimes 1) \otimes (1 \otimes 1), (T \otimes S)(1 \otimes 1) \otimes (1 \otimes 1) \rangle \\
&= \langle 1 \otimes 1, T 1 \otimes 1 \rangle \otimes \langle 1 \otimes 1, S 1 \otimes 1 \rangle \\
&= h_1(T) \otimes h_2(S) \\
&= (h_1 \otimes h_2)(T \otimes S)
\end{aligned}$$

From ultrastrong density, again, it follows  $h \circ \text{ad}_{U^*} \circ \vartheta = h_1 \otimes h_2$ .  $\square$

## 2.6 Pure maps

Schrödinger's equation is invariant under the reversal of time and so the isolated quantum mechanical processes (ncp-maps) described by it are invertible. In stark contrast, the ncp-maps corresponding to measurement and discarding are rarely invertible. There are, however, many processes in between which might not be invertible, but are still pure enough to be 'reversed' in some consistent and useful fashion. For instance, the ncp-map  $\text{ad}_V$  has the obvious reversal  $(\text{ad}_V)^\dagger = \text{ad}_{V^*}$  (which is in general not the inverse.) Stinespring's theorem (135 IV) implies that every ncp-map into  $\mathcal{B}(\mathcal{H})$  splits as a reversible  $\text{ad}_V$  after a (possibly) non-reversible nmiu-map  $\varrho$ . Using Paschke's dilation, we see that every ncp-map factors as an nmiu-map  $\varrho$  before some particular ncp-map  $h$ . Is this ncp-map  $h$  reversible in some sense? In 171 VII we will see that these  $h$  are pure (in the

168

sense of 100I) and in 215III we will see that these pure maps are (in a sense) reversible.

- II The reversibility of pure maps (in a much more general setting) is the main result of the next chapter. In the remainder of this chapter, we study purity of ncp-maps a bit more. We consider a seemingly unrelated alternative notion of purity, which is based on Paschke's dilation, and show (in 171VII) that this alternative notion is in fact equivalent to the original. We end this chapter by giving a different proof of the ncp-extremeness of pure and nmiu-maps.
- III Before we continue, let's rule out generalizations of more familiar notions of purity to arbitrary ncp-maps.

1. A state  $\varphi: \mathcal{A} \rightarrow \mathbb{C}$  is called pure if it is an extreme point among all states. It is unreasonable to define an arbitrary ncp-map to be pure if it is extreme, as every nmiu-map (including von Neumann measurements and discarding) is extreme among the unital ncp-maps. This follows from 102III and 102V, but will be proven again in 172VIII. In fact, in 172XII we show that every ncp-map is the composition of two extreme maps.
2. Inspired by the GNS-correspondence between pure states and irreducible representations, Størmer defines a map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  to be pure if the only maps below  $\varphi$  in the completely positive ordering are scalar multiples of  $\varphi$ . For every central element  $z$  of  $\mathcal{A}$ , the map  $a \mapsto za$  is below  $\text{id}_{\mathcal{A}}$  and so the identity on non-factors is not pure in the sense of Størmer, even though the identity map is clearly reversible.

- IV Let's recall how we came to the notion of purity in 100I, by generalizing  $\text{ad}_V$ . Pick  $V: \mathcal{H} \rightarrow \mathcal{K}$ . By polar decomposition, see 82I, we know that  $V = UA$  for  $A \equiv \sqrt{V^*V}: \mathcal{H} \rightarrow [A]\mathcal{H}$  and some isometry  $U: [A]\mathcal{H} \rightarrow \mathcal{K}$ . Thus we have  $\text{ad}_V = \text{ad}_A \circ \text{ad}_U$ . These two maps admit dual universal properties:  $\text{ad}_U$  is a *corner* and  $\text{ad}_A$  is a *filter*. An ncp-map  $\varphi$  is pure if it is the composition of a filter after a corner. It turns out that  $\varphi$  is pure in this sense if and only if the map on the left-hand side of its Paschke dilation is surjective.

**169** We recall the definition of corner from [126, dfn. 2] and 95I.

- II **Definition** An ncp-map  $h: \mathcal{A} \rightarrow \mathcal{B}$  is a **corner** for  $a \in [0, 1]_{\mathcal{A}}$  if  $h(a) = h(1)$  and

for every ncp-map  $f: \mathcal{A} \rightarrow \mathcal{C}$  with  $f(1) = f(a)$ , there is a unique ncp-map  $f': \mathcal{B} \rightarrow \mathcal{C}$  with  $f = f' \circ h$ .

- III **Remark** In the next chapter we will study maps with a similar universal property as corners in a more general setting, but with all arrows in the reverse direction. To help alleviate some confusion caused by the change of direction, we call those maps *comprehensions*, see 199II. See also [23].

**Example** The map  $h_a: \mathcal{A} \rightarrow [a] \mathcal{A} [a]$  given by  $b \mapsto [a] b [a]$  is a corner of  $a \in [0, 1]_{\mathcal{A}}$ . We call  $h_a$  the **standard corner** of  $a$ . See 98 I, 95 II or [126, prop. 5]. IV

**Lemma** If  $(\mathcal{P}, \varrho, h)$  is a Paschke dilation of a unital ncp-map, then  $h$  is a corner. V

**Proof** (This is a different proof of our similar result [127, cor. 27].) Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  be any unital ncp-map. It is sufficient to prove  $h$  is a corner for the standard dilation  $(\mathcal{P}, \varrho, h)$  constructed in 154 III, so  $\mathcal{P} \equiv \mathcal{B}^a(\mathcal{A} \otimes_{\varphi} \mathcal{B})$  and  $h(T) = \langle e, Te \rangle$ , where  $e \equiv 1 \otimes 1$ . Write  $p \equiv |e\rangle\langle e|$ . From  $\langle e, e \rangle = \varphi(1) = 1$  it follows  $p$  is a projection. Define  $\vartheta: \mathcal{B} \rightarrow p\mathcal{P}p$  by  $\vartheta(b) = |e \cdot b\rangle\langle e|$ . Some straightforward computations (using 159 III) show  $\vartheta$  is an miu-map with inverse  $pTp \mapsto h(pTp)$ . So  $\vartheta$  is an miu-isomorphism and thus also normal. Furthermore  $h = \vartheta^{-1}(p(\cdot)p)$  and so  $h$  is indeed a corner (of  $p$ ). VI  $\square$

We recall the definition of filter, see 96 I. VII

**Definition** An ncp-map  $c: \mathcal{A} \rightarrow \mathcal{B}$  is a **filter** for  $b \in \mathcal{B}$ ,  $b \geq 0$  if  $c(1) \leq b$  and VIII

for every ncp-map  $f: \mathcal{C} \rightarrow \mathcal{B}$  with  $f(1) \leq b$ , there is a unique ncp-map  $f': \mathcal{C} \rightarrow \mathcal{A}$  with  $f = c \circ f'$ .

**Remark** The direction-reversed counterpart of filters are called *quotients*, see 197 II IX and [23]. Filters were introduced by us in [126, dfn. 2] under the name *compressions*. To avoid confusion with [4] and be closer to e.g. [132], we changed the name from compression to filter. Filters are named after polarization filters: the action of a polarization filter on the polarization of a photon is the same as a filter for the projection of the space of polarizations the polarization filter allows to pass without disturbance. Combining two polarization filters corresponds to composing the corresponding filters. This composed filter is in general a filter for an effect instead of a projection. This corresponds with the fact that there might not be a polarization of the photon anymore, with which it can pass both filters undisturbed.

**Example** The map  $c_b: [b] \mathcal{B} [b] \rightarrow \mathcal{B}$  given by  $a \mapsto \sqrt{b} a \sqrt{b}$  is a filter of  $b \in \mathcal{B}$ ,  $b \geq 0$ . We call  $c_b$  the **standard filter** of  $b$ . See 96 V, 98 I or [126, prop. 6]. X

**Exercise** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  be any ncp-map between von Neumann algebras. XI

1. Let  $c: \mathcal{B} \rightarrow \mathcal{C}$  be any filter. Show that if  $(\mathcal{P}, \varrho, h)$  is a Paschke dilation of  $\varphi$ , that then  $(\mathcal{P}, \varrho, c \circ h)$  is a Paschke dilation of  $c \circ \varphi$ .
2. Assume  $c': \mathcal{C}' \rightarrow \mathcal{B}$  is a filter of  $\varphi(1)$ . Prove that there is a unique unital ncp-map  $\varphi'$  with  $\varphi = c \circ \varphi'$ . Conclude that if  $(\mathcal{P}, \varrho, h)$  is a Paschke dilation of  $\varphi'$ , that then  $(\mathcal{P}, \varrho, c' \circ h)$  is a Paschke dilation of  $\varphi$ .

(See 221 IV and 197 VII for the proofs in the more general case.)

**Exercise** Derive (perhaps from X and 60 VIII), that filters are injective. XII

**170 Definition** Filters, corners and their compositions are called **pure**.

**Ia Remark** This definition is equivalent to our previous definition of purity given in [127, dfn. 21]; see 100 III.

**II Examples** Isomorphisms are pure. Less trivial examples follow.

1. The pure maps  $\mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  are precisely of the form  $\text{ad}_T$  for some bounded operator  $T: \mathcal{H} \rightarrow \mathcal{H}$ .
2. Let  $(\mathcal{P}, \varrho, h)$  be a Paschke dilation of some ncp-map  $\varphi$ . In 169 V we saw that if  $\varphi$  is unital, that then  $h$  is a corner (and thus pure). In general,  $h = c \circ h'$  for some filter  $c$  and corner  $h'$  by 169 XI, and so  $h$  is pure.

**III Remark** By definition, the set of pure maps is closed under composition. In the next chapter we will see (215 III) that there is a unique dagger  $\dagger$  which turns the subcategory of von Neumann algebras with pure maps into a  $\dagger$ -category such that  $f^\dagger$  is contraposed to  $f$ ;  $\diamond$ -positive maps are  $\dagger$ -positive and  $\dagger$ -positive maps have unique roots. For this unique  $\dagger$ , we have  $(\text{ad}_V)^\dagger = \text{ad}_{V^*}$ .

**IV Exercise** Show that any surjective nmiu-map between von Neumann algebras is a corner of a central projection, hence pure — and conversely, that any corner of a central projection is a surjective nmiu-map.

**171** We will prove that a map is pure if and only if the left-hand side map of its Paschke dilation is surjective. We need the following result.

**II Theorem** Let  $\mathcal{A}$  be a von Neumann algebra together with a projection  $p \in \mathcal{A}$ . Then a Paschke dilation of the standard corner  $h_p: \mathcal{A} \rightarrow p\mathcal{A}p$ ,  $a \mapsto pap$  is given by  $(\llbracket p \rrbracket \mathcal{A}, h_{\llbracket p \rrbracket}, h'_p)$ , where  $\llbracket p \rrbracket$  is the central carrier of  $p$ , see 68 I;  $h_{\llbracket p \rrbracket}$  is the standard corner for  $\llbracket p \rrbracket$  and  $h'_p: \llbracket p \rrbracket \mathcal{A} \rightarrow p\mathcal{A}p$  is the restriction of  $h_p$ .

**III Proof** (This is a simplified proof of the result we published earlier. [127, thm. 28]) Let  $(\mathcal{A} \otimes_{h_p} p\mathcal{A}p, \varrho, h)$  denote the Paschke dilation constructed in 154 III. We proceed in three steps: first we show  $\mathcal{A}p$  is a self-dual Hilbert  $p\mathcal{A}p$ -module, then we prove  $\mathcal{A} \otimes_{h_p} p\mathcal{A}p \cong \mathcal{A}p$  and finally we show  $\mathcal{B}^a(\mathcal{A}p) \cong \llbracket p \rrbracket \mathcal{A}$ .

**IV** Note  $\mathcal{A}p$  is a pre-Hilbert  $p\mathcal{A}p$ -module with scalar multiplication  $(\alpha p) \cdot (pap) = \alpha pap$  and inner product  $\langle \alpha p, ap \rangle = p\alpha^*ap$ . Since by the C\*-identity the norm on  $\mathcal{A}p$  as a Hilbert module coincides with that as a subset of  $p\mathcal{A}p$ ;  $\mathcal{A}p$  is norm closed in  $\mathcal{A}$  and  $\mathcal{A}$  is norm complete, we see  $\mathcal{A}p$  is a Hilbert  $p\mathcal{A}p$ -module.

Recall  $\mathcal{A}$  itself is a self-dual Hilbert  $\mathcal{A}$ -module. The uniformity induced on  $\mathcal{A}p$  by the ultranorm uniformity of  $\mathcal{A}$  coincides with the ultranorm uniformity of  $\mathcal{A}p$  itself. The module  $\mathcal{A}p$  is ultranorm closed in  $\mathcal{A}$  as  $x_\alpha \rightarrow x$  ultranorm implies  $x_\alpha p \rightarrow xp$  ultranorm. As self-duality is equivalent to ultranorm completeness, we see  $\mathcal{A}p$  is self dual as well.

Concerning  $\mathcal{A} \otimes_{h_p} p\mathcal{A}p \cong \mathcal{A}p$ , define  $B: \mathcal{A} \times p\mathcal{A}p \rightarrow \mathcal{A}p$  by  $B(\alpha, pap) = \alpha pap$ . A straightforward computation shows  $B$  is a  $h_p$ -compatible complex bilinear map (see 154 II) and so by 154 III, there is a unique bounded module map  $U: \mathcal{A} \otimes_{h_p} p\mathcal{A}p \rightarrow \mathcal{A}p$  fixed by  $U(\alpha \otimes pap) = B(\alpha, pap) \equiv \alpha pap$ . We will show  $U$  is an isomorphism. Clearly  $U$  is surjective. V

An easy computation shows  $a \otimes pap - ap\alpha p \otimes p = 0$  for all  $a, \alpha \in \mathcal{A}$ . So the set  $\{a \otimes p; a \in \mathcal{A}\}$  is ultranorm dense in  $\mathcal{A} \otimes_{h_p} p\mathcal{A}p$ . From this and  $\langle \alpha \otimes p, a \otimes p \rangle = p\alpha^*ap = \langle U\alpha \otimes p, Ua \otimes p \rangle$  we see  $U$  preserves the inner product. Hence  $U$  is an isomorphism with  $U^* = U^{-1}$  given by  $U^*a = a \otimes p$ .

To show  $\mathcal{B}^a(\mathcal{A}p) \cong \llbracket p \rrbracket \mathcal{A}$ , we will first find a convenient orthonormal basis of  $\mathcal{A}p$ . By 83 V, there are partial isometries  $(u_i)_i$  with  $\sum_i u_i u_i^* = \llbracket p \rrbracket$  and  $u_i^* u_i \leq p$ . Write  $E = \{u_i; i \in I\}$ . As  $u_i = u_i p$  and  $u_i = u_i u_i^* u_i$  we see  $E$  is an orthonormal subset of  $\mathcal{A}p$ . For any  $ap \in \mathcal{A}p$ , we have VI

$$ap = a \llbracket p \rrbracket p = \llbracket p \rrbracket ap = \left( \sum_i u_i u_i^* \right) ap = \sum_i u_i u_i^* ap = \sum_i u_i \langle u_i, ap \rangle,$$

where the sums converge ultrastrongly in  $\mathcal{A}$ . Thus the last sum converges ultranorm in  $\mathcal{A}p$  as well. Hence  $E$  is an orthonormal basis of  $\mathcal{A}p$ .

Define  $\varrho_0: \mathcal{A} \rightarrow \mathcal{B}^a(\mathcal{A}p)$  by  $\varrho_0 = \text{ad}_{U^*} \circ \varrho$ . Note  $\varrho_0$  is an nmiu-map and  $\varrho_0(\alpha)ap = \alpha ap$ . The map  $\varrho_0$  is surjective: indeed for any  $T \in \mathcal{B}^a(\mathcal{A}p)$

$$T(ap) = \sum_i T(u_i)u_i^* ap = \left( \sum_i T(u_i)u_i^* \right) ap = \varrho_0 \left( \sum_i T(u_i)u_i^* \right) ap.$$

Next, we show  $\llbracket \varrho_0 \rrbracket = \llbracket p \rrbracket$ . It suffices to prove for all  $\alpha \in \mathcal{A}, \alpha \geq 0$ , that

$$\alpha ap = 0 \text{ for all } a \in \mathcal{A} \iff \alpha \llbracket p \rrbracket = 0.$$

Clearly,  $\alpha \llbracket p \rrbracket = 0$  implies  $\alpha ap = \alpha \llbracket p \rrbracket ap = 0$ . For the converse, assume that we have  $\alpha ap = 0$  for all  $a \in \mathcal{A}$ . Then in particular  $\alpha u_i p = 0$ , hence  $\alpha \llbracket p \rrbracket = \alpha \sum_i u_i u_i^* = \sum_i \alpha u_i p u_i^* = 0$ . Indeed  $\llbracket \varrho_0 \rrbracket = \llbracket p \rrbracket$ .

By 170 IV there is an nmiu-isomorphism  $\varphi: \llbracket p \rrbracket \mathcal{A} \rightarrow \mathcal{B}^a(\mathcal{A}p)$  with  $\varrho_0 = \varphi \circ h_{\llbracket p \rrbracket}$ . Hence  $h_{\llbracket p \rrbracket} = (\varphi^{-1} \circ \text{ad}_{U^*}) \circ \varrho$ . We compute

$$h'_p \circ h_{\llbracket p \rrbracket} = h_p = h \circ \varrho = h \circ \text{ad}_U \circ \varrho_0 = h \circ \text{ad}_U \circ \varphi \circ h_{\llbracket p \rrbracket}.$$

Using surjectivity of  $h_{\llbracket p \rrbracket}$  and rearranging, we find  $h'_p \circ (\varphi^{-1} \circ \text{ad}_{U^*}) = h$ . Thus indeed  $(\llbracket p \rrbracket \mathcal{A}, h_{\llbracket p \rrbracket}, h'_p)$  is a Paschke dilation of  $h_p$ . □

**Theorem** Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  be an ncp-map between von Neumann algebras with Paschke dilation  $(\mathcal{P}, \varrho, h)$ . The map  $\varphi$  is pure if and only if  $\varrho$  is surjective. VII

**Proof** Assume  $\varrho$  is surjective. The kernel of  $\varrho$  is an ultraweakly closed two-sided ideal and so  $\ker \varrho = z\mathcal{A}$  for some central projection  $z$ , see 69 II. The standard corner  $h_{z^\perp}: \mathcal{A} \rightarrow z^\perp \mathcal{A}$  is the quotient-map of  $z\mathcal{A}$  and so by the isomorphism VIII

theorem  $\varrho$  is a corner as well. In 170 II we saw  $h$  is pure. So  $\varphi$  is the composition of pure maps, hence pure itself.

For the converse, assume  $\varphi$  is pure. For brevity, write  $p \equiv [\varphi]$ . There is a unique ncp-map  $c$  with  $\varphi = c \circ h_p$ , where  $h_{[\varphi]}$  is the standard corner of  $p$ . The map  $c$  is a filter, see 98 IX. By II, the triple  $(\llbracket p \rrbracket \mathcal{A}, h_{\llbracket p \rrbracket}, h'_p)$  is a Paschke dilation of  $h_p$ , where  $h'_p$  denotes the restriction of  $h_p$  to  $\llbracket p \rrbracket \mathcal{A}$ . So by 169 XI we see  $(\llbracket p \rrbracket \mathcal{A}, h_{\llbracket p \rrbracket}, c \circ h'_p)$  is a Paschke dilation of  $\varphi$ . Clearly  $h_{\llbracket p \rrbracket}$  is surjective. If  $(\mathcal{P}', \varrho', h')$  is any other Paschke dilation of  $\varphi$ , then  $\varrho' = \vartheta \circ h_{\llbracket p \rrbracket}$  for some isomorphism  $\vartheta$  and so  $\varrho'$  is surjective as well.  $\square$

**172** The order correspondence 157 IV has several corollaries.

**II Definition** Assume  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  is an ncp-map between von Neumann algebras. We say  $\varphi$  is **ncp-extreme** if  $\varphi$  is an extreme point among the ncp-maps with the same value on 1 [108] — that is: for any  $0 < \lambda < 1$ , ncp-maps  $\varphi_1, \varphi_2: \mathcal{A} \rightarrow \mathcal{B}$  with  $\varphi_1(1) = \varphi_2(1) = \varphi(1)$  we have

$$\lambda\varphi_1 + (1 - \lambda)\varphi_2 = \varphi \implies \varphi_1 = \varphi_2 = \varphi.$$

**III Theorem** Assume  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  is an ncp-map with Paschke dilation  $(\mathcal{P}, \varrho, h)$ . Then the following are equivalent.

1. The map  $h$  is injective on  $\varrho(\mathcal{A})^\square$ , the commutant of  $\varrho(\mathcal{A})$ .
2. The map  $h$  is injective on  $[0, 1]_{\varrho(\mathcal{A})^\square}$ .
3. The map  $\varphi$  is ncp-extreme.

**IV Proof** (Based on [8, prop. 1.4.6] and [93, thm. 5.4].) We prove  $1 \Rightarrow 2 \Rightarrow 3 \Rightarrow 1$ .

**V**  $(1 \Rightarrow 2)$  Trivial.

**VI**  $(2 \Rightarrow 3)$  Assume  $h$  is injective on  $[0, 1]_{\varrho(\mathcal{A})^\square}$ . Suppose  $\varphi = \lambda\psi + (1 - \lambda)\psi'$  for some  $0 < \lambda < 1$  and ncp  $\psi, \psi': \mathcal{A} \rightarrow \mathcal{B}$  with  $\psi(1) = \psi'(1) = \varphi(1)$ . Note  $0 \leq_{\text{ncp}} \lambda\psi \leq_{\text{ncp}} \varphi$  hence  $\lambda\psi = \varphi_t$  for some  $t \in [0, 1]_{\varrho(\mathcal{A})^\square}$  by 157 IV. So

$$h(\lambda 1) = \varphi_{\lambda 1}(1) = \lambda\varphi(1) = \lambda\psi(1) = \varphi_t(1) = h(t).$$

By the injectivity of  $h$ , we find  $t = \lambda 1$ , thus  $\lambda\psi = \lambda\varphi$  and so  $\psi = \varphi$ . The proof of  $\psi' = \varphi$  is the same. Hence  $\varphi$  is ncp-extreme.

**VII**  $(3 \Rightarrow 1)$  Suppose  $\varphi$  is ncp-extreme. Pick any  $a \in \varrho(\mathcal{A})^\square$  with  $h(a) = 0$ . We want to show  $a = 0$ . Clearly  $h(a)_{\mathbb{R}} = h(a_{\mathbb{R}}) = 0$  and  $h(a)_{\mathbb{I}} = h(a_{\mathbb{I}}) = 0$ , so it is sufficient to prove  $a_{\mathbb{R}} = a_{\mathbb{I}} = 0$ . Thus we may assume without loss of generality that  $a$  is self adjoint.

By 170 II  $h$  is pure and thus splits as  $h = c \circ h_p$  for some filter  $c: p\mathcal{P}p \rightarrow \mathcal{B}$  and the standard corner  $h_p: \mathcal{P} \rightarrow p\mathcal{P}p$  for  $p = [\mathcal{A}]$ . By 169 XII filters are

injective and so from the assumption  $0 = h(a) = c(pap)$  it follows  $pap = 0$ . Using 9X and some easy algebra, we can find  $0 < \mu, \lambda$  such that  $\frac{1}{4} \leq \mu a + \lambda \leq \frac{3}{4}$ . Define  $t = \mu a + \lambda$ . For the moment, note  $\varphi_t$  is ncp. As  $pap = 0$ , we see  $ptp = \lambda p$  and so  $\frac{1}{4}p \leq ptp = \lambda p \leq \frac{3}{4}p$ . Thus either  $0 < \lambda < 1$  or  $p = 0$ . If  $p = 0$  then  $\varphi = 0$  hence  $\mathcal{P} = \{0\}$  and  $a = 0$  as desired. For the non-trivial case, assume  $0 < \lambda < 1$ . Note  $\varphi_t(1) = h(t) = c(ptp) = \lambda c(p1p) = \lambda\varphi(1)$  and similarly  $\varphi_{1-t}(1) = (1 - \lambda)\varphi(1)$ . Define  $\psi_1 = \lambda^{-1}\varphi_t$  and  $\psi_2 = (1 - \lambda)^{-1}\varphi_{1-t}$ . By the previous,  $\psi_1$  and  $\psi_2$  are ncp-maps with  $\psi_1(1) = \psi_2(1) = \varphi(1)$ . Also  $\lambda\psi_1 + (1 - \lambda)\psi_2 = \varphi$ . As  $\varphi$  is ncp-extreme, we must have  $\psi_1 = \psi_2 = \varphi$ . So  $\varphi_t = \lambda\varphi = \varphi_{\lambda 1}$ , thus  $\lambda = t \equiv \mu a + \lambda$ . Rearranging:  $\mu a = 0$ . As  $\mu > 0$ , we find  $a = 0$ , as desired.  $\square$

**Corollary** Any nmiu-map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}$  is ncp-extreme. (For a different proof, see VIII e.g. [108, thm. 3.5].) VIII

**Proof** Easy:  $(\mathcal{B}, \varrho, \text{id})$  is a Paschke dilation of  $\varrho$  and  $\text{id}$  is injective.  $\square$  IX

**Theorem** Any pure ncp-map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  is ncp-extreme. X

**Proof** By 100III  $\varphi = c \circ h_p$  for a filter  $c: p\mathcal{A}p \rightarrow \mathcal{B}$  and projection  $p \equiv \lceil \varphi \rceil$ . Combining 169XI and 171II we see  $(\llbracket p \rrbracket \mathcal{A}, h_{\llbracket p \rrbracket}, c \circ h_p)$  is a Paschke dilation of  $\varphi$ . Appealing to III, we are done if we can show  $c \circ h_p$  is injective on  $[0, 1]_{h_{\llbracket p \rrbracket}(\mathcal{A})}$ . As  $h_{\llbracket p \rrbracket}$  is surjective,  $h_{\llbracket p \rrbracket}(\mathcal{A})^\square = Z(\llbracket p \rrbracket \mathcal{A})$ . Assume  $t \in [0, 1]_{Z(\llbracket p \rrbracket \mathcal{A})}$  with  $c(ptp) = 0$ . We want to show  $t = 0$ . As  $c$  is injective by 169XII, we must have  $ptp = 0$ . So  $p \leq 1 - \lceil t \rceil$  and  $\llbracket p \rrbracket \leq 1 - \lceil t \rceil$ . Hence  $\llbracket p \rrbracket \leq \llbracket p \rrbracket - \lceil t \rceil \leq \llbracket p \rrbracket$ . So  $t \leq \lceil t \rceil = 0$ , as desired.  $\square$

**Corollary** Any ncp-map is the composition of two ncp-extreme maps. XII



## Chapter 3

# Diamond, and then, dagger

In the previous chapter and in [124] we have studied categorical properties of von Neumann algebras. In this chapter we change pace: we study categories that resemble the category of von Neumann algebras. The goal of this line of study is to identify axioms which uniquely pick out the category of von Neumann algebras. We do not reach this ambitious goal — instead we will build up to a characterization of the existence of a unique  $\dagger$ -structure on the pure maps of a von Neumann algebras-like category. 173

As a basic axiom we will require the categories we consider to be effectuses (which will be defined in 180!). For us the definition of an effectus will play a similar role to that of a topological space for a geometer. In many applications, general topological spaces on their own are of little interest: the axioms are so weak that there are many (for the application) pathological examples. These weak axioms, however, are very expressive in the sense that they allow for the definition of many useful notions. Herein lies the strength of topological spaces — as a stepping stone: many important classes of mathematical spaces are just plain topological spaces with a few additional axioms. Similarly, there will be many pathological effectuses. Their use for us lies in their expressiveness. II

Before we dive into effectuses, we need to learn about effect algebras (and related structures), which were introduced by mathematical physicists as a generalization of Boolean algebra to study fuzzy predicates in quantum mechanics. For us, effectuses are the categorical counterpart of effect algebras.\* After this, we continue with a brief, but thorough development of the basic theory of effectuses. There is a lot more to say about effectuses (see [23]), which has been omitted to avoid straining the reader. We will indulge, however, in one tangent, which III

---

\*Effect algebroïds [97, 98] are a different categorification with applications in cohomology.

can be skipped: the study of abstract convex sets in §3.2.4. For the moment, think of an effectus as a generalization of the opposite category of von Neumann algebras: the objects represent the physical systems (or data types, if you like) and the maps represent the physical processes (or properly typed programs).

- IV The first two axioms we add on top of those of an effectus are the existence of quotients and comprehension. It's helpful to discuss the origin of these axioms. It started with the desire to axiomatize categorically the sequential product on a von Neumann algebra  $\mathcal{A}$ , that is the operation  $a \& b \equiv \sqrt{ab}\sqrt{a}$ , which represents sequential measurement of first  $a$  *and then*  $b$ . The sequential product is a quantum mechanical generalization of classical conjunction (logical 'and'). In contrast to the tame  $\wedge$  in a Boolean algebras, the sequential product does not obey a lot of insightful formulas. The sequential product as a binary operation has received quite some attention [6, 37, 44–47, 71, 72, 83, 104, 111, 122]. We approach the sequential product in a rather different way: we take a step back and consider the map  $\text{asrt}_a(b) \equiv a \& b$  for fixed  $a$ . ( $\text{asrt}_a$  is named after the **assert** statement used in many programming languages, see 211 III.) This map  $\text{asrt}_a$  factors in two:

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{\pi} & [a] \mathcal{A} [a] & \xrightarrow{\xi} & \mathcal{A} \\ b \vdash & \xrightarrow{\quad} & [a] b [a] & \xrightarrow{\quad} & \sqrt{ab}\sqrt{a}. \end{array}$$

It turned out that both  $\pi$  and  $\xi$  have nice and dual (in a sense) defining universal properties which can be expressed in an effectus:  $\pi$  is a comprehension and  $\xi$  is a quotient.<sup>†</sup>

- V The existence of quotients and comprehension in some effectus  $C$  is interesting on its own, but with two additional axioms (the existence of images and preservation of images under orthocomplementation), we get a surprisingly firm handle on the *possibilistic* side of  $C$ , by means of the existence of a certain functor  $(\ )_{\diamond}: C \rightarrow \text{OMLat}$  to the category of orthomodular lattices. In the guiding example of von Neumann algebras, we have  $f_{\diamond} = g_{\diamond}$  if and only there are no post-measurement  $a$  and initial state  $\omega$  that can determine with certainty whether  $f$  or  $g$  has been performed.<sup>‡</sup> With this functor we can introduce several possibilistic notions of which  $\diamond$ -adjoint and  $\diamond$ -positive are the most important. Because of the central role of  $\diamond$ , we call an effectus with these axioms a  $\diamond$ -effectus.
- VI The axioms of a  $\diamond$ -effectus do not force any coherence between the quotients and comprehension. This has been a roadblock to the axiomatization of the sequential product for quite a while. The key insight was the following: the map  $\text{asrt}_a$  (i.e.  $b \mapsto a \& b$ ) is the unique  $\diamond$ -positive map on  $\mathcal{A}$  with  $\text{asrt}_a(1) = a$ . We turn this theorem into an axiom: an  $\&$ -effectus is a  $\diamond$ -effectus where there are

<sup>†</sup>Beware: an effectus quotient is not the same thing as a von Neumann-algebra quotient.

<sup>‡</sup>In symbols:  $f_{\diamond} = g_{\diamond} \iff (\forall a, \omega. \omega(f(a)) = 0 \iff \omega(g(a)) = 0)$ .

such unique  $\diamond$ -positive maps  $\text{asrt}_a$  and where an additional polar decomposition axiom holds. In an  $\&$ -effectus, the predicates on an object carry a canonical binary operation  $\&$  which is the intended sequential product in the case of von Neumann algebras.

In an effectus, we call a map pure if it can be written as the composition of quotients and comprehensions. In 171 VII we saw that in the case of von Neumann algebras, a map is pure (in this sense) if and only if the corresponding Paschke embedding is surjective. In particular, the pure maps  $\mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  are exactly of the form  $\text{ad}_v$  (for bounded operators  $v: \mathcal{H} \rightarrow \mathcal{H}$ ) which carry a  $\dagger$ -structure, namely  $(\text{ad}_v)^\dagger = \text{ad}_{v^*}$ . This begs the question: can we define a  $\dagger$  on all pure maps in an  $\&$ -effectus? The main result of this chapter is Theorem 215 III, which gives necessary and sufficient conditions for an  $\&$ -effectus to have a well-behaved  $\dagger$ -structure on its pure maps. After this apogee, we tie up some loose ends and discuss how these structures compare to those already known in the literature. VII

### 3.1 Effect algebras and related structures

Before we turn to effectuses, it is convenient to introduce some lesser-known algebraic structures of which the effect algebra is the most important. It will turn out that the set of predicates associated to an object in an effectus can be arranged into an effect algebra. In this way, effect algebras play the same role for effectuses as Heyting algebras for toposes. First, we will recall the definition of partial commutative monoid (PCM) — as later in 180 X we will see that the partial maps between two objects in an effectus can be arranged into a PCM. 174

**Definition** A **partial commutative monoid (PCM)**  $M$  is a set  $M$  together with distinguished element  $0 \in M$  and a partial binary operation  $\odot$  such that for all  $a, b, c \in M$  — writing  $a \perp b$  whenever  $a \odot b$  is defined — we have II

1. (*partial commutativity*) if  $a \perp b$ , then  $b \perp a$  and  $a \odot b = b \odot a$ ;
2. (*partial associativity*) if  $a \perp b$  and  $a \odot b \perp c$ , then  $b \perp c$ ,  $a \perp b \odot c$  and  $(a \odot b) \odot c = a \odot (b \odot c)$  and
3. (*zero*)  $0 \perp a$  and  $0 \odot a = a$ .

A map  $f: M \rightarrow N$  between PCMs  $M$  and  $N$  is called an **PCM homomorphism** if for all  $a, b \in E$  with  $a \perp b$ , we have  $f(a) \perp f(b)$  and  $f(a) \odot f(b) = f(a \odot b)$ . Write **PCM** for the category of PCMs with these homomorphisms. (Cf. [65, 121])

For  $a, b$  in a PCM  $M$ , we say  $a \leq b$  iff  $a \odot c = b$  for some  $c \in M$ .

**Exercise** Show a PCM is preordered by  $\leq$ . III

- IV The partial commutativity and associativity of a PCM ensure that a sum only depends on which elements occur (and how often), instead of their order. Formally: if  $(\cdots(x_1 \otimes x_2) \otimes \cdots) \otimes x_n$  exists some  $x_1, \dots, x_n$  in some PCM, then so does  $(\cdots(x_{\pi(1)} \otimes x_{\pi(2)}) \otimes \cdots) \otimes x_{\pi(n)}$  for any permutation  $\pi$  of  $\{1, \dots, n\}$  and

$$(\cdots(x_1 \otimes x_2) \otimes \cdots) \otimes x_n = (\cdots(x_{\pi(1)} \otimes x_{\pi(2)}) \otimes \cdots) \otimes x_{\pi(n)}.$$

Thus parantheses are superfluous and we will often leave them out:

$$\bigotimes_{i=1}^n x_i \equiv x_1 \otimes \cdots \otimes x_n \equiv (\cdots(x_1 \otimes x_2) \otimes \cdots) \otimes x_n.$$

**175 Definition** A PCM  $E$  together with distinguished element  $1 \in E$  is called an **effect algebra (EA)** [31] provided that

1. (*orthocomplement*) for every  $a$  there is a unique  $a^\perp$  with  $a \otimes a^\perp = 1$  and
2. (*zero-one*) if  $a \perp 1$ , then  $a = 0$ .

A map  $f: E \rightarrow F$  between effect algebras  $E$  and  $F$  is called an **effect algebra homomorphism** if it is a PCM homomorphism that preserves 1; concretely:

1. (*additive*)  $a \perp b$  implies  $f(a) \perp f(b)$  and  $f(a) \otimes f(b) = f(a \otimes b)$  and
2. (*unital*)  $f(1) = 1$ .

(It follows that  $f(0) = 0$  and  $f(a^\perp) = f(a)^\perp$ . See 176V.) Write **EA** for the category of effect algebras with these homomorphisms. A subset  $D \subseteq E$  is a **sub-effect algebra** of  $E$  if  $0, 1 \in D$  and for any  $a, b \in D$  with  $a \perp b$ , we have  $a \otimes b \in D$  and  $a^\perp \in D$ .

II **Examples** There are many examples of effect algebras — we only give a few.

1. The unit interval  $[0, 1]$  with partial addition is an effect algebra — i.e.:  $x \perp y$  whenever  $x + y \leq 1$  and then  $x \otimes y = x + y$ ,  $x^\perp = 1 - x$ .
2. Generalizing the previous: if  $G$  is an ordered group with distinguished element 1, then the order interval  $[0, 1]_G \equiv \{x; x \in G; 0 \leq x \leq 1\}$  is an effect algebra with  $x \perp y$  whenever  $x + y \leq 1$ ;  $x \otimes y = x + y$  and  $x^\perp = 1 - x$ .
3. In particular, if  $\mathcal{A}$  is any von Neumann algebra, then the set of **effects**  $[0, 1]_{\mathcal{A}}$  forms an effect algebra with  $a \perp b$  whenever  $a + b \leq 1$ ;  $a \otimes b = a + b$  and  $a^\perp = 1 - a$ . The ‘effect’ in effect algebra originates from this example.
4. Any orthomodular lattice  $L$  (defined in 177IV) is an effect algebra with the same orthocomplement,  $x \perp y$  whenever  $x \leq y^\perp$  and  $x \otimes y = x \vee y$ . (See e.g. [129, prop. 27].)

5. In particular, any Boolean algebra  $L$  is an effect algebra with complement as orthocomplement,  $x \perp y$  whenever  $x \wedge y = 0$  and  $x \otimes y = x \vee y$ .
6. The one-element Boolean algebra  $1 \equiv \{0 = 1\}$  is the final object in EA and the two-element Boolean algebra  $2 \equiv \{0, 1\}$  is the initial object in EA.

**Exercise** Let  $E$  and  $F$  be effect algebras. First show that the cartesian product  $E \times F$  is an effect algebra with componentwise operations. Show that this is in fact the categorical product of  $E$  and  $F$  in EA. (The category EA is in fact complete and cocomplete; for this and more categorical properties, see [65].) III

**Exercise** There is a small redundancy in our definition of effect algebra: show that the zero axiom ( $x \otimes 0 = x$ ) follows from the remaining axioms (partial commutativity, partial associativity, orthocomplement and zero-one.) IV

**Proposition** In any effect algebra  $E$  with  $a, b, c \in E$ , we have V

1. (*involution*)  $a^{\perp\perp} = a$ ;
2.  $1^\perp = 0$  and  $0^\perp = 1$ ;
3. (*positivity*) if  $a \otimes b = 0$ , then  $a = b = 0$ ;
4. (*cancellation*) if  $a \otimes c = b \otimes c$ , then  $a = b$ ;
5. the relation  $\leq$  (from 174II) partially orders  $E$ ;
6.  $a \leq b$  if and only if  $b^\perp \leq a^\perp$ ;
7. if  $a \leq b$  and  $b \perp c$ , then  $a \perp c$  and  $a \otimes c \leq b \otimes c$  and
8.  $a \perp b$  if and only if  $a \leq b^\perp$ .

**Proof** (These proofs are well-known, see for instance [29].) By partial commutativity and definition of orthocomplement both  $a^\perp \otimes a = 1$  and  $a^\perp \otimes a^{\perp\perp} = 1$ . So by uniqueness of orthocomplement, we must have  $a = a^{\perp\perp}$ , which is point 1. Clearly  $0 \otimes 1 = 1$ , so  $1^\perp = 0$  and  $0^\perp = 1$ , which is point 2. For point 3, assume  $a \otimes b = 0$ . Then  $a \otimes b \perp 1$  and so by partial associativity  $b \perp 1$ . By zero-one, we get  $b = 0$ . Similarly  $a = 0$ , which shows point 3. For point 4, assume  $a \otimes c = b \otimes c$ . From partial associativity and commutativity, we get  $((a \otimes b)^\perp \otimes a) \otimes c = ((a \otimes b)^\perp \otimes a) \otimes b = 1$  and so by uniqueness of orthocomplement  $c = ((a \otimes b)^\perp \otimes a)^\perp = b$ , which is point 4. By 174III, we only need to show that  $\leq$  is antisymmetric for point 5. So assume  $a \leq b$  and  $b \leq a$  for some  $a, b \in E$ . Pick  $c, d \in E$  with  $a \otimes c = b$  and  $b \otimes d = a$ . Then  $a = (a \otimes c) \otimes d = a \otimes (c \otimes d)$ . By cancellation  $c \otimes d = 0$ . So by positivity  $c = d = 0$ . Hence  $a = b$ . For point 6, assume  $a \leq b$ . Pick  $c \in E$

with  $a \otimes c = b$ . Clearly  $a \otimes a^\perp = 1 = b \otimes b^\perp = a \otimes c \otimes b^\perp$ , so by cancellation  $a^\perp = c \otimes b^\perp$ , which is to say  $b^\perp \leq a^\perp$ . For point 7, assume  $a \leq b$  and  $b \perp c$ . Pick  $d$  with  $a \otimes d = b$ . By partial associativity and commutativity we have  $b \otimes c = (a \otimes d) \otimes c = (a \otimes c) \otimes d$ , so  $a \perp c$  and  $a \otimes c \leq b \otimes c$ . For point 8, first assume  $a \perp b$ . Then  $a \otimes b \otimes (a \otimes b)^\perp = 1 = b \otimes b^\perp$ . So by cancellation  $b^\perp = a \otimes (a \otimes b)^\perp$ , hence  $a \leq b^\perp$ . For the converse, assume  $a \leq b^\perp$ . Then  $a \otimes c = b^\perp$  for some  $c$ . Hence  $a \otimes c \perp b$  and so by partial associativity and commutativity, we get  $a \perp b$ , as desired.  $\square$

**176 Definition** Suppose  $E$  is an effect algebra. Write  $b \ominus a$  for the (by cancellation) unique element (if it exists) with  $a \otimes (b \ominus a) = b$ .

**II Exercise\*** Show that for any effect algebra  $E$ , we have

(D1)  $a \ominus b$  is defined if and only if  $b \leq a$ ;

(D2)  $a \ominus b \leq a$  (if defined);

(D3)  $a \ominus (a \ominus b) = b$  (if defined) and

(D4) if  $a \leq b \leq c$ , then  $c \ominus b \leq c \ominus a$  and  $(c \ominus a) \ominus (c \ominus b) = b \ominus a$ .

**III** Let  $E$  be a poset with maximum element 1 and partial binary operation  $\ominus$  satisfying (D1)–(D4). Define  $a \otimes b = c \Leftrightarrow c \ominus b = a$  and  $a^\perp = 1 \ominus a$ . Show that this turns  $E$  into an effect algebra with compatible order and  $\ominus$ .

**IV Remark** Such a structure  $(E, \ominus, 1)$  is called a **difference-poset (D-poset)** [77] and is, as we have just seen, an alternative way to axiomatize effect algebras.

**V Exercise** Show that for an effect algebra homomorphism  $f: E \rightarrow F$ , we have

1. (*preserves zero*)  $f(0) = 0$ ;
2. (*order preserving*) if  $a \leq b$ , then  $f(a) \leq f(b)$ ;
3. if  $a \ominus b$  is defined, then  $f(a \ominus b) = f(a) \ominus f(b)$  and
4. consequently  $f(a^\perp) = f(a)^\perp$ .

**177** For real numbers we have  $a + b = \min\{a, b\} + \max\{a, b\}$ . The following is a generalization to effect algebras.

**1a Proposition** Suppose  $E$  is an effect algebra. If the infimum  $a \wedge b$  exists for some  $a, b \in E$  with  $a \perp b$ , then their supremum  $a \vee b$  exists as well and

$$a \otimes b = (a \wedge b) \otimes (a \vee b).$$

**Proof** (This result appeared in my master's thesis [129, prop. 15]. It turns out that it was already proven before (in D-poset form) [29, prop. 1.8.2].) The result follows from this lemma: if  $(x \ominus c) \wedge (x \ominus d)$  exists, then  $c \vee d$  exists as well and  $(x \ominus c) \wedge (x \ominus d) = x \ominus (c \vee d)$ . Indeed: II

$$a \wedge b = ((a \otimes b) \ominus a) \wedge ((a \otimes b) \ominus b) = (a \otimes b) \ominus (a \vee b)$$

and so  $(a \wedge b) \otimes (a \vee b) = a \otimes b$ , as desired.

Now, we prove the lemma. Assume  $(x \ominus c) \wedge (x \ominus d)$  exists. Note  $x \ominus c = (x^\perp \otimes c)^\perp$ . III  
As  $z \mapsto z^\perp$  is an order anti-isomorphism, we see  $(x^\perp \otimes c) \vee (x^\perp \otimes d)$  exists and  $(x^\perp \otimes c) \vee (x^\perp \otimes d) = ((x \ominus c) \wedge (x \ominus d))^\perp$ . Clearly  $(x^\perp \otimes c) \vee (x^\perp \otimes d) \geq x^\perp$ . Write  $r := ((x^\perp \otimes c) \vee (x^\perp \otimes d)) \ominus x^\perp$ . We will show  $r = c \vee d$ . It is easy to see  $r \geq c$  and  $r \geq d$ . Assume  $s$  is any element with  $s \geq c$  and  $s \geq d$ . Note  $x^\perp \otimes s \geq x^\perp \otimes c$  and  $x^\perp \otimes s \geq x^\perp \otimes d$ , hence  $x^\perp \otimes s \geq (x^\perp \otimes c) \vee (x^\perp \otimes d)$  and so  $s \geq r$ . We have shown  $r = c \vee d$ . Consequently  $(x \ominus c) \wedge (x \ominus d) = ((x^\perp \otimes c) \vee (x^\perp \otimes d))^\perp = (r \otimes x^\perp)^\perp = x \ominus (c \vee d)$ , as promised. □

**Definition** An **ortholattice**  $L$  is a bounded lattice with orthocomplement — that is: it has a minimum 0, a maximum 1 and a unary operation  $( )^\perp$  satisfying IV

1.  $a \wedge a^\perp = 0$ ;
2.  $a \vee a^\perp = 1$ ;
3.  $a \leq b \Rightarrow b^\perp \leq a^\perp$  and
4.  $a^{\perp\perp} = a$ .

An ortholattice is **orthomodular** provided

$$a \leq b \quad \Longrightarrow \quad a \vee (a^\perp \wedge b) = b.$$

See, for instance [11, 29, 74].

**Example** The lattice of projections in a von Neumann algebra is an orthomodular lattice with orthocomplement  $p^\perp \equiv 1 - p$ . V

**Proposition** An effect algebra  $E$  that is an ortholattice is also orthomodular. VI

**Proof** (For a different proof, see [29, prop. 1.5.8].) Assume  $a \leq b$ . We have to show  $a \vee (a^\perp \wedge b) = b$ . Note  $a \wedge (a^\perp \wedge b) \leq a \wedge a^\perp = 0$  and so by **la** we have  $a \vee (a^\perp \wedge b) = a \otimes (a^\perp \wedge b)$ . Thus it is sufficient to prove  $a^\perp \wedge b = b \ominus a$ . Note  $a^\perp = (b \ominus a) \otimes b^\perp$  and  $b = (b \ominus a) \otimes a$ . Similar to **II** one can show that VII

$$(b \ominus a) \otimes (b^\perp \wedge a) = ((b \ominus a) \otimes b^\perp) \wedge ((b \ominus a) \otimes a) \equiv a^\perp \otimes b,$$

but  $b^\perp \wedge a \leq b^\perp \wedge b = 0$  and so  $a^\perp \otimes b = b \ominus a$ , as desired. □

### 3.1.1 Effect monoids

In 190II we will see that the scalars of an effectus form an effect monoid: 178

II **Definition** An **effect monoid**  $M$  [56] is an effect algebra together with a binary operation  $\odot$  such that for all  $a, b, c, d \in M$ , we have

1. (*unit*)  $1 \odot a = a \odot 1 = a$ ;
2. (*associativity*)  $(a \odot b) \odot c = a \odot (b \odot c)$  and
3. (*distributivity*) if  $a \perp b$  and  $c \perp d$ , then the following sum exists and furthermore  $(a \odot c) \vee (b \odot c) \vee (a \odot d) \vee (b \odot d) = (a \vee b) \odot (c \vee d)$ .

(Phrased categorically: an effect monoid is a monoid in EA with the obvious tensor product that relates bimorphisms to morphisms, see [56, 65].) An effect monoid  $M$  is said to be **commutative** if we have  $a \odot b = b \odot a$  for all  $a, b \in M$ . A map  $f: M \rightarrow N$  between effect monoids is called an **effect monoid homomorphism** if it is an effect algebra homomorphism and furthermore  $f(a \odot b) = f(a) \odot f(b)$  for all  $a, b \in M$ .

III **Examples** Effect monoids are less abundant than effect algebras.

1. The effect algebra  $[0, 1]$  is a commutative effect monoid with the usual product. (This is the only way to turn  $[0, 1]$  into an effect monoid [129, prop. 41].)
2. We saw earlier that every Boolean algebra is an effect algebra with  $x \vee y = x \vee y$  defined iff  $x \wedge y = 0$ . The Boolean algebra is turned into an effect monoid with  $x \odot y \equiv x \wedge y$ . In fact, every finite (not necessarily commutative) effect monoid is of this form [129, prop. 40] and thus commutative.
3. In particular: the two-element Boolean algebra  $2 = 0, 1$  from 175 II is an effect monoid with  $x \odot y = x \wedge y$ .
4. There is a non-commutative effect monoid based on the lexicographically ordered vector space  $\mathbb{R}^5$ , see [129, cor. 51].
5. Let  $M$  be an effect monoid. Write  $M^{\text{op}}$  for the effect monoid  $M$  with the opposite multiplication — that is:  $a \odot_M b = b \odot_{M^{\text{op}}} a$ .

IIIa **Exercise** Show that  $a \odot 0 = a = 0 \odot a$  for any  $a$  in an effect monoid  $M$ .

IV Later, in a tangent, we will need the following specific fact about effect monoids.

V **Exercise** Assume  $M$  is an effect monoid with  $a_1, \dots, a_n, b_1, \dots, b_n \in M$  such that  $\bigvee_i a_i = 1$  and  $\bigvee_i a_i \odot b_i = 1$ . Prove  $a_i \odot b_i = a_i$  for every  $1 \leq i \leq n$ .

### 3.1.2 Effect modules

179 We will see that in an effectus, the effect monoid of scalars will act on every effect algebra of predicates, turning them into effect modules (see 190 II):

**Definition** Suppose  $M$  is an effect monoid. An **effect module**  $E$  over  $M$  [65] is an effect algebra together with an operation  $M \times E \rightarrow E$  denoted by  $(\lambda, a) \mapsto \lambda \cdot a$  such that for all  $a, b \in E$  and  $\lambda, \mu \in M$ , we have II

1.  $(\lambda \odot \mu) \cdot a = \lambda \cdot (\mu \cdot a)$ ;
2. if  $a \perp b$ , then  $\lambda \odot a \perp \lambda \odot b$  and  $(\lambda \odot a) \vee (\lambda \odot b) = \lambda \odot (a \vee b)$ ;
3. if  $\lambda \perp \mu$ , then  $\lambda \odot a \perp \mu \odot a$  and  $(\lambda \odot a) \vee (\mu \odot a) = (\lambda \vee \mu) \odot a$  and
4.  $1 \odot a = a$ .

(Categorically: an effect module over  $M$  is an  $M$ -action.) An effect algebra homomorphism  $f: E \rightarrow F$  between effect modules over  $M$  is an  $M$ -**effect module homomorphism** provided  $\lambda \cdot f(a) = f(\lambda \cdot a)$  for all  $\lambda \in M$  and  $a \in E$ . Write  $\mathbf{EMod}_M$  for the category of effect modules over  $M$  with effect module homomorphisms between them.

**Examples** There are many effect modules. III

1. Every effect algebra is an effect module over the two-element effect monoid  $2$ . (In fact  $\mathbf{EA} \cong \mathbf{EMod}_2$ .) The only effect module up-to-isomorphism over the one-element effect monoid  $1$  is the one-element effect algebra  $1$  itself.
2. Effect modules over  $[0, 1]$  are the same thing as **convex effect algebras**, see [43, 48]. If  $V$  is an ordered real vector space with order unit  $u$ , then  $[0, u]$  is an effect module over  $[0, 1]$ . In fact, every effect module over  $[0, 1]$  is of this form [48]. See also [66, thm. 3] for the stronger categorical equivalence.

## 3.2 Effectuses

An effectus comes in two guises: axiomatizing either a category of total maps or a category of partial maps. We will start off with the total form as it has the simplest axioms. Later we will prefer to work with the partial form.

**Definition** A category  $C$  is said to be an **effectus in total form** [23, 57, 69] if 180

1.  $C$  has finite coproducts (hence an initial object  $0$ ) and a final object  $1$ ;

2. all diagrams of the following form<sup>§</sup> are pullbacks

$$\begin{array}{ccc}
 X + Y & \xrightarrow{\text{id}+!} & X + 1 \\
 \text{!+id} \downarrow & & \downarrow \text{!+id} \\
 1 + Y & \xrightarrow{\text{id}+!} & 1 + 1
 \end{array}
 \qquad
 \begin{array}{ccc}
 X & \xrightarrow{!} & 1 \\
 \kappa_1 \downarrow & & \downarrow \kappa_1 \\
 X + Y & \xrightarrow{\text{!+!}} & 1 + 1
 \end{array}
 \quad (3.1)$$

3. and the following two arrows are jointly monic.

$$\begin{array}{ccc}
 1 + 1 + 1 & \xrightarrow{[\kappa_1, \kappa_2, \kappa_2]} & 1 + 1 \\
 & \xrightarrow{[\kappa_2, \kappa_1, \kappa_2]} & 1 + 1
 \end{array}$$

An arrow  $f: X \rightarrow Y + 1$  is called a **partial map** and written  $f: X \dashrightarrow Y$ .

- II One with an interest in physics might think of the objects of an effectus as physical systems and its arrows as the physical operations between them. The final object  $1$  is the physical system with a single state. The coproduct  $X + Y$  is the system that can be prepared as either  $X$  or as  $Y$ . An arrow  $1 \rightarrow X$  corresponds to a physical preparation of the system  $X$  and an arrow  $X \rightarrow 1 + 1$  is a yes–no measurement.
- III In this sense an effectus can be seen as a generalized probabilistic theory — not unlike the operational-probabilistic theories (OPT) of Chiribella et al [17]. There are differences: for instance, OPTs are equipped with a parallel composition of systems, effectuses are not and OPTs always have the unit interval as scalars, effectuses might not. In [113] Tull compares effectuses and OPTs. There are other approaches to categorical probabilistic theories; see, for instance [40, 131].
- IV Studying programming languages, one would do better thinking of the objects of an effectus as data types and its arrows as the allowed operations between them (semantics of programs). The final object  $1$  is the unit data type. The coproduct  $X + Y$  is the union data type of  $X$  and  $Y$ . An arrow  $1 \rightarrow X$  is a value of  $X$  and an arrow  $X \rightarrow 1 + 1$  is a predicate on  $X$ .
- V Our main example of an effectus in total form is the category  $\text{vN}^{\text{op}}$  of von Neumann algebras with completely positive normal unital maps in the opposite direction. (To see  $\text{vN}^{\text{op}}$  is an effectus in total form, adapt the proof of 191 II.) The partial maps correspond to ncp-maps  $f$  with  $f(1) \leq 1$ . (Equivalently: contractive ncp-maps.) Some other examples appear later on in 191 III, 192 III, 196 II and 189a I. For a comprehensive list of examples, see [23].

---

<sup>§</sup>We write  $\kappa_i$  for coproduct coprojections; square brackets  $[f, g]$  for coproduct cotupling;  $h + k = [\kappa_1 \circ h, \kappa_2 \circ k]$  and  $!$  for the unique maps associated to either the final object  $1$  or initial object  $0$ .

Let  $C$  be an effectus in total form. Given two arrows  $f: X \rightarrow Y + 1$  and  $g: Y \rightarrow Z + 1$  (i.e. partial maps  $X \rightarrow Y$  and  $Y \rightarrow Z$ ) their composition as partial maps is defined as  $g \hat{\circ} f \equiv [g, \kappa_2] \circ f$ . Write  $\text{Par } C$  for the **category of partial maps**, which has the same objects as  $C$ , but as arrows  $X \rightarrow Y$  in  $\text{Par } C$  we take arrows of the form  $X \rightarrow Y + 1$  in  $C$ , which we compose using  $\hat{\circ}$  and with identity on  $X$  in  $\text{Par } C$  given by  $\kappa_1: X \rightarrow X + 1$ .<sup>¶</sup> The category  $\text{Par } C$  is not an effectus in total form — instead it is an effectus *in partial form*. VI

**Definition** A category  $C$  is called an **effectus in partial form** [19, 23] if VII

1.  $C$  is a finPAC<sup>||</sup> [19] (cf. [5]) — that is:
  - (a)  $C$  has finite coproducts  $(+, 0)$ ;
  - (b)  $C$  is PCM-enriched — that is:
    - i. every homset  $C(X, Y)$  has a partial binary operation  $\otimes$  and distinguished element  $0 \in C(X, Y)$  that turns  $C(X, Y)$  into a partial commutative monoid, see 174 II;
    - ii. if  $f \perp g$  then both  $(h \circ f) \perp (h \circ g)$  and  $(f \circ k) \perp (g \circ k)$  and
 
$$h \circ (f \otimes g) = (h \circ f) \otimes (h \circ g) \quad (f \otimes g) \circ k = (f \circ k) \otimes (g \circ k)$$
 for any  $f, g: X \rightarrow Y$ ,  $h: Y \rightarrow Y'$  and  $k: X' \rightarrow X$  and
    - iii. (zero)  $0 \circ f = 0$  and  $f \circ 0 = 0$  for any  $f: X \rightarrow Y$  and
  - (c) (compatible sum) for any  $b: X \rightarrow Y + Y$  we have  $\triangleright_1 \circ b \perp \triangleright_2 \circ b$ , where  $\triangleright_i: Y + Y \rightarrow Y$  are **partial projectors**\*\* defined by  $\triangleright_1 \equiv [\text{id}, 0]$  and  $\triangleright_2 \equiv [0, \text{id}]$  and
  - (d) (untying) if  $f \perp g$ , then  $\kappa_1 \circ f \perp \kappa_2 \circ g$ , where  $\kappa_1$  and  $\kappa_2$  are coprojections on the same coproduct and
2. it has effects — that is: there is a distinguished object  $I$  such that
  - (a) for each object  $X$ , the PCM  $C(X, I) \equiv \text{Pred } X$  is an effect algebra, see 175 I — in particular  $C(X, I)$  has a maximum element  $1$ ;
  - (b) if  $1 \circ f \perp 1 \circ g$ , then  $f \perp g$  and
  - (c) if  $1 \circ f = 0$ , then  $f = 0$ .

In this setting, a map  $f: X \rightarrow Y$  is called **total** if  $1 \circ f = 1$ .

**Remark** The untying and zero axioms are redundant: they follows from the others. We include them, as they are part of the definition of finPAC. VIII

<sup>¶</sup>In categorial parlance:  $\text{Par } C$  is the Kleisli category of the monad  $(\ ) + 1: C \rightarrow C$ .

<sup>||</sup>Finitarily partially additive category.

\*\*Later, in 181 VII, we will use the more slightly more general  $\triangleright_1: X_1 + X_2 \rightarrow X_1$  and  $\triangleright_2: X_1 + X_2 \rightarrow X_2$  defined by  $\triangleright_1 = [\text{id}, 0]$  and  $\triangleright_2 = [0, \text{id}]$ .

- IX At first glance an effectus in partial form seems to have a much richer structure than an effectus in total form. This is not the case — effectuses in total and partial form are two views on the same thing [19, 23]:
- X **Theorem (Cho)** Let  $C$  be an effectus in total form and  $D$  be an effectus in partial form.
  1. The category  $\text{Par } C$  is an effectus in partial form with  $I = 1$ .
  2. The total maps of  $D$  form an effectus in total form  $\text{Tot } D$ .
  3. Nothing is lost:  $\text{Par}(\text{Tot } D) \cong D$  and  $\text{Tot}(\text{Par } C) \cong C$ .
- XI (The result can be rephrased categorically as a 2-equivalence of the 2-category of effectuses in partial form and the 2-category of effectuses in total form. For the details, see [19, §5].) To prove the Theorem (in 188II), we need some preparation.

### 3.2.1 From partial to total

**181** We will first show that the subcategory of total maps of an effectus in partial form is an effectus in total form. This proof and especially the demonstration that the squares in (3.1) are pullbacks, will elucidate the axioms of an effectus in total form and will make the proof in the opposite direction more palatable.

II **Lemma** In an effectus in partial form, coprojections are total.

III **Proof** (For the original and different proof see [19, lem. 4.7(5)].) Let  $\kappa_1: X \rightarrow X + Y$  be any coprojection. We have to show  $1 \circ \kappa_1 = 1$ . Note that the first 1 denotes the maximum of  $\text{Pred } X + Y$  and the second the maximum of  $\text{Pred } X$ . Hence  $1_X = 1_X \circ \text{id}_X = (1_X \circ [\text{id}_X, 0]) \circ \kappa_1 \leq 1_{X+Y} \circ \kappa_1 \leq 1_X$ . Indeed  $1 \circ \kappa_1 = 1$ .  $\square$

IV **Proposition** In an effectus in partial form, the cotupling bijection  $(f, g) \mapsto [f, g]$  is a PCM-isomorphism [23]— that is:

1.  $[f, g] \perp [f', g']$  if and only if  $f \perp f'$  and  $g \perp g'$ ;
2. if  $[f, g] \perp [f', g']$ , then  $[f, g] \otimes [f', g'] = [f \otimes f', g \otimes g']$  and
3.  $[0, 0] = 0$ .

Furthermore  $[1, 1] = 1$  for maps into  $I$ , so the cotupling map is an effect algebra isomorphism  $\text{Pred}(X) \times \text{Pred}(Y) \cong \text{Pred}(X + Y)$ .

V **Proof** First we show  $[h, l] = [h, 0] \otimes [0, l]$ . By the compatible sum axiom

$$[h, 0] = \triangleright_1 \circ (h + l) \perp \triangleright_2 \circ (h + l) = [0, l].$$

By PCM-enrichment  $([h, 0] \otimes [0, l]) \circ \kappa_1 = ([h, 0] \circ \kappa_1) \otimes ([0, l] \circ \kappa_1) = h$ . Similarly  $([h, 0] \otimes [0, l]) \circ \kappa_2 = l$ . Thus indeed  $[h, l] = [h, 0] \otimes [0, l]$ .

Assume  $[f, g] \perp [f', g']$ . By PCM-enrichment we have  $f = [f, g] \circ \kappa_1 \perp [f', g'] \circ \kappa_1 = f'$ . Similarly  $g \perp g'$ . Conversely, assume  $f \perp f'$  and  $g \perp g'$ . Again, by PCM-enrichment  $[f, 0] = f \circ \triangleright_1 \perp f' \circ \triangleright_1 = [f', 0]$  and  $[f \otimes f', 0] = [f, 0] \otimes [f', 0]$ . Similarly  $[0, g \otimes g'] = [0, g] \otimes [0, g']$ . Putting it all together:

$$\begin{aligned} [f, g] \otimes [f', g'] &= [f, 0] \otimes [0, g] \otimes [f', 0] \otimes [0, g'] \\ &= [f \otimes f', 0] \otimes [0, g \otimes g'] \\ &= [f \otimes f', g \otimes g']. \end{aligned}$$

To show cotupling is a PCM-isomorphism, it only remains to be shown that  $[0, 0] = 0$ . As  $0 \circ \kappa_1 = 0$  and  $0 \circ \kappa_2 = 0$ , we indeed have  $0 = [0, 0]$ . Similarly by II we have  $1 \circ \kappa_1 = 1$  and  $1 \circ \kappa_2 = 1$ , so  $1 = [1, 1]$ .  $\square$

The coproduct in an effectus in partial form is almost a (bi)product: VI

**Proposition** In an effectus in partial form, we have a bijective correspondence VII

$$\frac{h: Z \rightarrow X + Y}{\frac{f: Z \rightarrow X \quad g: Z \rightarrow Y \quad 1 \circ f \perp 1 \circ g}}$$

as follows [19, lem. 4.8]:

for every  $f: Z \rightarrow X$  and  $g: Z \rightarrow Y$  with  $1 \circ f \perp 1 \circ g$ , there is a unique map  $\langle f, g \rangle: Z \rightarrow X + Y$  such that  $\triangleright_1 \circ \langle f, g \rangle = f$  and  $\triangleright_2 \circ \langle f, g \rangle = g$ , where  $\triangleright_1 = [\text{id}, 0]$  and  $\triangleright_2 = [0, \text{id}]$ . In fact  $\langle f, g \rangle = (\kappa_1 \circ f) \otimes (\kappa_2 \circ g)$ .

Conversely  $h = \langle \triangleright_1 \circ h, \triangleright_2 \circ h \rangle$  with  $1 \circ \triangleright_1 \circ h \perp 1 \circ \triangleright_2 \circ h$  for  $h: Z \rightarrow X + Y$ .

**Proof** Let  $f: Z \rightarrow X$  and  $g: Z \rightarrow Y$  be given such that  $1 \circ f \perp 1 \circ g$ . By II, we have  $1 \circ \kappa_1 \circ f = 1 \circ f \perp 1 \circ g = 1 \circ \kappa_2 \circ g$ . Thus  $\kappa_1 \circ f \perp \kappa_2 \circ g$ . Define  $\langle f, g \rangle = (\kappa_1 \circ f) \otimes (\kappa_2 \circ g)$ . By the PCM-enrichedness, we have VIII

$$\begin{aligned} \triangleright_1 \circ \langle f, g \rangle &= [\text{id}, 0] \circ ((\kappa_1 \circ f) \otimes (\kappa_2 \circ g)) \\ &= ([\text{id}, 0] \circ \kappa_1 \circ f) \otimes ([\text{id}, 0] \circ \kappa_2 \circ g) \\ &= f \otimes (0 \circ g) = f. \end{aligned}$$

Similarly  $\triangleright_2 \circ \langle f, g \rangle = g$ . To show uniqueness, assume  $f = \triangleright_1 \circ h$  and  $g = \triangleright_2 \circ h$  for some  $h: Z \rightarrow X + Y$ . Note  $\kappa_1 \circ \triangleright_1 = [\kappa_1, 0]$  and  $\kappa_2 \circ \triangleright_2 = [0, \kappa_2]$ , so by IV we have  $(\kappa_1 \circ \triangleright_1) \otimes (\kappa_2 \circ \triangleright_2) = [\kappa_1, \kappa_2] = \text{id}$ , hence

$$\begin{aligned} \langle f, g \rangle &= \langle \triangleright_1 \circ h, \triangleright_2 \circ h \rangle \\ &= (\kappa_1 \circ \triangleright_1 \circ h) \otimes (\kappa_2 \circ \triangleright_2 \circ h) \\ &= ((\kappa_1 \circ \triangleright_1) \otimes (\kappa_2 \circ \triangleright_2)) \circ h \\ &= \text{id} \circ h = h, \end{aligned}$$

which demonstrates uniqueness.

Finally, let  $h: Z \rightarrow X+Y$  be any map. We must show that  $h = \langle \triangleright_1 \circ h, \triangleright_2 \circ h \rangle$ . Note  $1 \circ \triangleright_1 = [1, 0]$  and  $1 \circ \triangleright_2 = [0, 1]$ . So by PCM-enrichment  $1 \circ \triangleright_1 \circ h \perp 1 \circ \triangleright_2 \circ h$  and so  $\triangleright_1 \circ h \perp \triangleright_2 \circ h$ . By the previous  $\langle \triangleright_1 \circ h, \triangleright_2 \circ h \rangle = (\kappa_1 \circ \triangleright_1 \circ h) \otimes (\kappa_2 \circ \triangleright_2 \circ h) = h$  as desired.  $\square$

**IX Exercise** Show that in an effectus in partial form, we have

1.  $[a, b] \circ \langle f, g \rangle = (a \circ f) \otimes (b \circ g)$ ;
2.  $1 \circ \langle f, g \rangle = (1 \circ f) \otimes (1 \circ g)$ ;
3.  $(k + l) \circ \langle f, g \rangle = \langle k \circ f, l \circ g \rangle$  and
4.  $\langle f, g \rangle \circ k = \langle f \circ k, g \circ k \rangle$

assuming  $1 \circ f \perp 1 \circ g$ . [23]

**X Remark** In an effectus in partial form, there is a straightforward generalization of the bijective correspondence VII to

$$\frac{h: Z \rightarrow X_1 + \cdots + X_n}{f_1: Z \rightarrow X_1 \quad \cdots \quad f_n: Z \rightarrow X_n \quad \perp_n 1 \circ f_n}$$

with  $f_i = \triangleright_i \circ h$  and  $h = \langle f_1, \dots, f_n \rangle \equiv (\kappa_1 \circ f_1) \otimes \cdots \otimes (\kappa_n \circ f_n)$ , where  $\triangleright_i$  is defined in the obvious way. The expected generalizations of the rules in IX also hold.

**XI Theorem** Let  $C$  be an effectus in partial form. The category of total maps of  $C$  is an effectus in total form. [19, thm. 4.10]

**XII Proof** The total maps indeed form a subcategory:  $1 \circ \text{id} = 1$  and  $1 \circ f \circ g = 1 \circ g = 1$  for composable total  $f, g$ . In II we saw coprojections are total. By IV we have  $1 \circ [f, g] = [1 \circ f, 1 \circ g] = [1, 1] = 1$  for total  $f: X \rightarrow Z$  and  $g: Y \rightarrow Z$ , so  $\text{Tot } C$  has binary coproducts. The unique map  $!: 0 \rightarrow X$  must be total as  $1 \circ ! = 1$  is the unique map  $0 \rightarrow I$ , so  $\text{Tot } C$  has initial object  $0$ , hence all finite coproducts.

**XIII** To show  $I$  is the final object of  $\text{Tot } C$ , we need  $\text{id}_I = 1$ . As  $C(I, I)$  is an effect algebra  $1 = \text{id} \otimes \text{id}^\perp$  for some  $\text{id}^\perp$ . So by PCM-enrichment  $1 = 1 \circ 1 = (1 \circ \text{id}) \otimes (1 \circ \text{id}^\perp) = 1 \otimes (1 \circ \text{id}^\perp)$ . By the zero-one axiom  $1 \circ \text{id}^\perp = 0$ . So  $\text{id}^\perp = 0$  and indeed  $\text{id} = 1$ . To show  $I$  is final in  $\text{Tot } C$ , pick any object  $X$  in  $\text{Tot } C$ . We claim  $1: X \rightarrow I$  is the unique total map. Indeed, by the previous  $1 = \text{id}_I \circ 1 = 1 \circ 1$ , so  $1$  is total and if  $h: X \rightarrow I$  is total, then  $1 = 1 \circ h = \text{id}_I \circ h = h$ .

To show the square on the left of (3.1) is a pullback in  $\text{Tot } C$ , let  $f: Z \rightarrow X + I$  XIV  
and  $g: Z \rightarrow I + Y$  be total maps with  $(\text{id} + 1) \circ g = (1 + \text{id}) \circ f$ . By VII,  $f = \langle \alpha, a \rangle$   
and  $g = \langle b, \beta \rangle$  for some maps  $\alpha: Z \rightarrow X$ ,  $\beta: Z \rightarrow Y$  and  $a, b: Z \rightarrow I$ .

$$\begin{array}{ccc}
Z & \xrightarrow{\langle \alpha, a \rangle \equiv f} & X + I \\
\downarrow \langle \alpha, \beta \rangle & \searrow \text{id} + 1 & \downarrow 1 + \text{id} \\
X + Y & \xrightarrow{\text{id} + 1} & X + I \\
\downarrow 1 + \text{id} & & \downarrow 1 + \text{id} \\
I + Y & \xrightarrow{\text{id} + 1} & I + I
\end{array}$$

$g \equiv \langle b, \beta \rangle$

By IX, we have  $1 = 1 \circ f = 1 \circ \langle \alpha, a \rangle = (1 \circ \alpha) \otimes a$  and so  $a^\perp = 1 \circ \alpha$ . Similarly  $b^\perp = 1 \circ \beta$ . Again, using IX, we see that

$$\langle b, 1 \circ \beta \rangle = (\text{id} + 1) \circ \langle b, \beta \rangle = (\text{id} + 1) \circ g = (1 + \text{id}) \circ f = \langle 1 \circ \alpha, a \rangle,$$

so  $1 \circ \beta = a = (1 \circ \alpha)^\perp$ , hence  $1 \circ \beta \perp 1 \circ \alpha$ , so  $\langle \alpha, \beta \rangle: Z \rightarrow X + Y$  exists and is total as  $1 \circ \langle \alpha, \beta \rangle = (1 \circ \alpha) \otimes (1 \circ \beta) = 1$ . We compute  $(\text{id} + 1) \circ \langle \alpha, \beta \rangle = \langle \alpha, 1 \circ \beta \rangle = \langle \alpha, a \rangle = f$ . Similarly  $(1 + \text{id}) \circ \langle \alpha, \beta \rangle = g$ . Assume  $h: Z \rightarrow X + Y$  is any map with  $(1 + \text{id}) \circ h = g$  and  $(\text{id} + 1) \circ h = f$ . Say  $h = \langle h_1, h_2 \rangle$ . Then  $\langle \alpha, a \rangle = f = (\text{id} + 1) \circ h = \langle h_1, 1 \circ h_2 \rangle$ , so  $\alpha = h_1$ . Similarly  $\beta = h_2$ . Thus  $h = \langle \alpha, \beta \rangle$ , which shows our square is indeed a pullback.

To show the square on the right of (3.1) is a pullback in  $\text{Tot } C$ , assume (using XV  
VII)  $\langle \alpha, \beta \rangle: Z \rightarrow X + Y$  is some total map with  $(1 + 1) \circ \langle \alpha, \beta \rangle = \kappa_1 \circ 1$ .

$$\begin{array}{ccc}
Z & \xrightarrow{\dots \kappa_1} & X & \xrightarrow{1} & I \\
\downarrow \langle \alpha, \beta \rangle & & \downarrow \kappa_1 & & \downarrow \kappa_1 \\
X + Y & \xrightarrow{1 + 1} & X + Y & \xrightarrow{1 + 1} & I + I
\end{array}$$

With IX, we see  $\langle 1 \circ \alpha, 1 \circ \beta \rangle = (1 + 1) \circ \langle \alpha, \beta \rangle = \kappa_1 \circ 1 = \langle 1, 0 \rangle$ . So  $\alpha$  is total and  $\beta = 0$ . Hence  $\langle \alpha, \beta \rangle = \langle \alpha, 0 \rangle = \kappa_1 \circ \alpha$  as desired.

Finally, to show  $m_1 \equiv [\kappa_1, \kappa_2, \kappa_2], m_2 \equiv [\kappa_2, \kappa_1, \kappa_2]: I + I + I \rightarrow I + I$  are jointly XVI  
monic, let  $f_1 \equiv \langle a_1, b_1, c_1 \rangle, f_2 \equiv \langle a_2, b_2, c_2 \rangle: X \rightarrow I + I + I$  be any total maps with  $m_1 \circ f_1 = m_1 \circ f_2$  and  $m_2 \circ f_1 = m_2 \circ f_2$ . Then

$$a_1 = [\text{id}, 0, 0] \circ f_1 = \triangleright_1 \circ m_1 \circ f_1 = \triangleright_1 \circ m_1 \circ f_2 = a_2.$$

and similarly from the equality involving  $m_2$ , we get  $b_1 = b_2$ . As  $f_1$  is total, we have  $1 = 1 \circ f_1 = a_1 \otimes b_1 \otimes c_1$ , so  $c_1 = (a_1 \otimes b_1)^\perp$ . With the same reasoning  $c_2 = (a_2 \otimes b_2)^\perp$ . Thus  $c_1 = c_2$  and so  $f_1 = f_2$ , as desired. □

### 3.2.2 From total to partial

**182** Let  $C$  be an effectus in total form. In this section we will show that  $\text{Par } C$  is an effectus in partial form. Before we get to work, it is helpful to discuss the axioms of an effectus in total form, now we have some experience with an effectus in partial form.

1. In 181 VII we saw that the coproduct in an effectus (in partial form) is almost a biproduct. This structure is hidden (for the most part) in the left pullback square of (3.1), which allows the formation of  $\langle \alpha, \beta \rangle$  given partial maps  $\alpha, \beta$ .
2. The right pullback square of (3.1) is used to extract a total map in  $C$  from a partial map  $f$  in  $\text{Par } C$  that is total (i.e.  $1 \hat{\circ} f = 1$ ). See 186 VIII.
3. The joint-monicity of  $[\kappa_1, \kappa_2, \kappa_2]$  and  $[\kappa_2, \kappa_1, \kappa_2]$  will imply the joint monicity of  $\triangleright_1$  and  $\triangleright_2$ , which is required for uniqueness of the partial sum of partial maps. See 186 X.

To prove the theorem, we need to study pullbacks: first in any category, then in an effectus  $C$  in total form and finally in  $\text{Par } C$ . Our proof is different from the original proof of Cho by using several more general facts about pullbacks, which we will prove next.

**183** We start with two classic facts about pullbacks.

II **Exercise** Show that if we have any pullback square — say

$$\begin{array}{ccc} P & \xrightarrow{m_1} & B \\ m_2 \downarrow \lrcorner & & \downarrow f \\ A & \xrightarrow{g} & X \end{array},$$

then  $m_1$  and  $m_2$  are jointly monic.

III **Exercise** Prove the [pullback lemma](#) — that is: if we have a commuting diagram

$$\begin{array}{ccccc} A & \xrightarrow{f} & B & \xrightarrow{g} & C \\ k \downarrow & & l \downarrow & & \downarrow m \\ X & \xrightarrow{f'} & Y & \xrightarrow{g'} & Z \end{array},$$

then we have the following two implications.

1. If the left and right inner squares are pullbacks, then so is the outer square.

2. If the outer square is a pullback and  $l$  and  $g$  are jointly monic, then the left inner square is a pullback.

**Remark** In the literature (e.g. [87, III.5 exc. 8]), one often finds a weaker second IV implication, which assumes that the right square is a pullback.

It is well-known that monos are stable under pullbacks — that is:

184

$$\text{if } \begin{array}{ccc} P & \xrightarrow{n} & X \\ g \downarrow \lrcorner & & \downarrow f \\ A & \xrightarrow{m} & B \end{array}, \quad \text{then } \begin{array}{ccc} P & \xrightarrow{\triangleright n} & X \\ g \downarrow \lrcorner & & \downarrow f \\ A & \xrightarrow{m} & B \end{array},$$

where we used  $\triangleright$  to denote a monic map. We will need an analogous result for jointly monic maps.

**Lemma** If the pairs  $(m_1, m_2)$ ,  $(n_1, g_1)$ ,  $(n_2, g_2)$  and  $(h_1, h_2)$  in the following commuting diagram are jointly monic (for instance: if they span pullback squares), II then  $(n_1 \circ h_1, n_2 \circ h_2)$  is jointly monic as well.

$$\begin{array}{ccccc} P & \xrightarrow{h_1} & P_1 & \xrightarrow{n_1} & X_1 \\ h_2 \downarrow & & \downarrow g_1 & & \downarrow f_1 \\ P_2 & \xrightarrow{g_2} & A & \xrightarrow{m_1} & B_1 \\ n_2 \downarrow & & \downarrow m_2 & & \\ X_2 & \xrightarrow{f_2} & B_2 & & \end{array}$$

**Proof** To show joint monicity of  $n_1 \circ h_1$  and  $n_2 \circ h_2$ , assume  $\alpha_1, \alpha_2: Z \rightarrow P$  are III maps with  $n_1 \circ h_1 \circ \alpha_1 = n_1 \circ h_1 \circ \alpha_2$  and  $n_2 \circ h_2 \circ \alpha_1 = n_2 \circ h_2 \circ \alpha_2$ . To start

$$m_1 \circ g_1 \circ h_1 \circ \alpha_2 = f_1 \circ n_1 \circ h_1 \circ \alpha_2 = f_1 \circ n_1 \circ h_1 \circ \alpha_1 = m_1 \circ g_1 \circ h_1 \circ \alpha_1.$$

Reasoning on the other side of the diagram, we find

$$\begin{aligned} m_2 \circ g_1 \circ h_1 \circ \alpha_2 &= m_2 \circ g_2 \circ h_2 \circ \alpha_2 \\ &= f_2 \circ n_2 \circ h_2 \circ \alpha_2 \\ &= f_2 \circ n_2 \circ h_2 \circ \alpha_1 \\ &= m_2 \circ g_2 \circ h_2 \circ \alpha_1 \\ &= m_2 \circ g_1 \circ h_1 \circ \alpha_1. \end{aligned}$$

So by joint monicity of  $m_1$  and  $m_2$ , we conclude  $g_1 \circ h_1 \circ \alpha_2 = g_1 \circ h_1 \circ \alpha_1$ . So by the joint monicity of  $n_1$  and  $g_1$  (and by assumption  $n_1 \circ h_1 \circ \alpha_2 = n_1 \circ h_1 \circ \alpha_1$ ),

we conclude  $h_1 \circ \alpha_1 = h_1 \circ \alpha_2$ . Reasoning in the same way mirrored over the diagonal, we find  $h_2 \circ \alpha_1 = h_2 \circ \alpha_2$ . Thus, by joint monicity of  $h_1$  and  $h_2$ , we conclude that  $\alpha_1 = \alpha_2$ , as desired.  $\square$

**185 Proposition** In an effectus in total form, any square of one of the two following forms is a pullback.

$$\begin{array}{ccc} X + A & \xrightarrow{\text{id}+f} & X + B \\ \downarrow \lrcorner \scriptstyle{g+\text{id}} & & \downarrow \scriptstyle{g+\text{id}} \\ Y + A & \xrightarrow{\text{id}+f} & Y + B \end{array} \qquad \begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow \lrcorner \scriptstyle{\kappa_1} & & \downarrow \scriptstyle{\kappa_1} \\ X + A & \xrightarrow{f+g} & Y + B \end{array}$$

|| **Proof** To start with the left square, consider the following commuting diagram.

$$\begin{array}{ccccc} X + A & \xrightarrow{\text{id}+f} & X + B & \xrightarrow{\text{id}+!} & X + 1 \\ \downarrow \scriptstyle{g+\text{id}} & & \downarrow \scriptstyle{g+\text{id}} & & \downarrow \scriptstyle{g+\text{id}} \\ Y + A & \xrightarrow{\text{id}+f} & Y + B & \xrightarrow{\text{id}+!} & Y + 1 \\ \downarrow \scriptstyle{!+\text{id}} & & \downarrow \scriptstyle{!+\text{id}} & & \downarrow \scriptstyle{!+\text{id}} \\ 1 + A & \xrightarrow{\text{id}+f} & 1 + B & \xrightarrow{\text{id}+!} & 1 + 1 \end{array}$$

We want to show (1) is a pullback. The inner square (4) and right rectangle (2,4) are pullbacks by axiom. Thus the inner square (2) is also a pullback by the pullback lemma, see 183 III. With the same reasoning, we see square (3) is a pullback. Thus by the pullback lemma, it is sufficient to show that left rectangle (1,3) is a pullback. The left rectangle is indeed a pullback as both the outer square (1,2,3,4) and the right rectangle (2,4) are pullbacks.

For the right square, we consider the following diagram.

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & \xrightarrow{!} & 1 \\ \downarrow \scriptstyle{\kappa_1} & & \downarrow \scriptstyle{\kappa_1} & & \downarrow \scriptstyle{\kappa_1} \\ X + A & \xrightarrow{f+g} & Y + B & \xrightarrow{!+!} & 1 + 1 \end{array}$$

The inner right and outer square are pullbacks by axiom. So the inner left square is also a pullback by the pullback lemma, as desired.  $\square$

**186 Definition** Assume  $C$  is an effectus in total form. For a map  $f: X \rightarrow Y$ , write  $\hat{f} = \kappa_1 \circ f: X \rightarrow Y + 1$ . (This is the Kleisli embedding  $C \rightarrow \text{Par } C$ .) [19]

(Recall that we use  $f: X \rightarrow Y$  to denote a map  $f: X \rightarrow Y + 1$ , see 180I.) Ia

**Exercise** Assume  $C$  is an effectus in total form. Show that if II

$$\kappa_1: X \rightarrow X + Y \leftarrow Y: \kappa_2$$

is a coproduct in  $C$ , then

$$\hat{\kappa}_1: X \rightarrow X + Y \leftarrow Y: \hat{\kappa}_2$$

is a coproduct in  $\text{Par } C$ . Show  $0$  is also the initial object of  $\text{Par } C$ .

**Beware** While cotupling in  $C$  and  $\text{Par } C$  coincide:  $[f, g]_C = [f, g]_{\text{Par } C}$  for any III  
(properly typed) partial maps  $f, g$  and  $\widehat{f +_C g} = \hat{f} +_{\text{Par } C} \hat{g}$ , however, in general

$$[\kappa_1 \circ f, \kappa_2 \circ g] = f +_C g \neq f +_{\text{Par } C} g = [[\kappa_1, \kappa_3] \circ f, [\kappa_2, \kappa_3] \circ g].$$

**Proposition** Let  $C$  be an effectus in total form. Squares of the following form IV  
are pullbacks in  $\text{Par } C$ .

$$\begin{array}{ccc} X + A & \xrightarrow{\text{id} + \hat{f}} & X + B \\ \hat{g} + \text{id} \downarrow \lrcorner & & \downarrow \hat{g} + \text{id} \\ Y + A & \xrightarrow{\text{id} + \hat{f}} & Y + B \end{array} \quad \begin{array}{ccc} A + X & \xrightarrow{\hat{f} + \text{id}} & B + X \\ \triangleright_1 \downarrow \lrcorner & & \downarrow \triangleright_1 \\ A & \xrightarrow{\hat{f}} & B \end{array}$$

**Proof** (A different proof for the pullback on the right is given in [19, lem. 4.1(5)].) V  
It is sufficient to show that the following squares are pullbacks in  $C$ .

$$\begin{array}{ccc} X + (A + 1) & \xrightarrow{\text{id} + (f + \text{id})} & X + (B + 1) \\ g + \text{id} \downarrow & & \downarrow g + \text{id} \\ Y + (A + 1) & \xrightarrow{\text{id} + (f + \text{id})} & Y + (B + 1) \end{array} \quad \begin{array}{ccc} A + (X + 1) & \xrightarrow{f + \text{id}} & B + (X + 1) \\ \text{id} + ! \downarrow & & \downarrow \text{id} + ! \\ A + 1 & \xrightarrow{f + \text{id}} & B + 1 \end{array}$$

These are indeed pullbacks by 185I. □


**Definition** Assume  $C$  is an effectus in total form. For arbitrary objects  $X, Y$  VI  
in  $C$ , write  $0 \equiv \kappa_2 \circ !: X \rightarrow Y + 1$  and  $1 \equiv \kappa_1 \circ ! \equiv \hat{!}: X \rightarrow 1 + 1$ . [19]

**Exercise** Show  $0$  is a zero object in  $\text{Par } C$  with unique zero map as in VI. VII

**Proposition** Assume  $C$  is an effectus in total form and  $f$  is a map in  $\text{Par } C$ . VIII


1. If  $1 \hat{\circ} f = 1$ , then  $f = \hat{g}$  for a unique  $g$  in  $C$ .
2. If  $1 \hat{\circ} f = 0$ , then  $f = 0$ .

- IX **Proof** (This proof is essentially the same as [19, lem. 4.7(1) & prop. 4.6].) Both points follow from the right pullback square of (3.1) as follows.



If  $1 \hat{\circ} f = 1$ , then  $(!+) \circ f = \kappa_1 \circ !$  and so there is a unique  $g$  with  $f = \kappa_1 \circ g$  as shown above on the left. For the other point, if  $1 \hat{\circ} f = 0$ , then  $(!+) \circ f = \kappa_2 \circ !$  and so  $f = \kappa_2 \circ !$  as shown above on the right.  $\square$

- X **Proposition** Assume  $C$  is an effectus in total form. The partial projectors  $\triangleright_1 \equiv [\text{id}, 0]$  and  $\triangleright_2 \equiv [0, \text{id}]$  are jointly monic in  $\text{Par } C$ . [19, lem. 4.1(4)]
- XI **Proof** (This is a new proof.) Consider the following diagram in  $\text{Par } C$ .



This diagram commutes and each of the inner squares is a pullback by IV. The maps  $\triangleright_1, \triangleright_2: 1+1 \rightarrow 1$  are jointly monic, which is a reformulation of the joint monicity axiom of an effectus in total form. Thus by 184II the outer  $\triangleright_1$  and  $\triangleright_2$  are jointly monic.  $\square$

187 **Theorem** If  $C$  is an effectus in total form, then  $\text{Par } C$  is an effectus in partial form. [19, thm. 4.2]

- II **Proof** In 186II, we saw that  $\text{Par } C$  has finite coproducts.
- III (PCM-enrichment, I) Assume  $f, g: X \rightarrow Y$ . We define that  $f \perp g$  iff there is a bound  $b: X \rightarrow Y+Y$  with  $\triangleright_1 \hat{\circ} b = f$  and  $\triangleright_2 \hat{\circ} b = g$ . (By 186X this  $b$  is unique if it exists.) In this case, define  $f \circledast g = \nabla \hat{\circ} b$ , where  $\nabla \equiv [\text{id}, \text{id}]$ . We will show  $\circledast$  with 0 as defined in 186VI forms a PCM. Partial commutativity is obvious. The map  $\kappa_1 \hat{\circ} f: X \rightarrow Y+Y$  is a bound for  $f \perp 0$  and so  $f \circledast 0 = \nabla \hat{\circ} \kappa_1 \hat{\circ} f = f$ , which

shows 0 is indeed a zero for  $\otimes$ . Only partial associativity remains. Assume  $f \perp g$  via bound  $b$  and  $f \otimes g \perp h$  via bound  $c$ . Note  $\nabla \hat{\circ} b = f \otimes g = \triangleright_1 \hat{\circ} c$  and thus by the right pullback square of 186 IV (see diagram below), there is a unique  $d: X \rightarrow (Y + Y) + Y$  with  $\triangleright_1 \hat{\circ} d = b$  and  $(\nabla + \text{id}) \hat{\circ} d = c$ .

$$\begin{array}{ccc}
 X & \xrightarrow{c} & (Y + Y) + Y \\
 \text{\scriptsize \dots} \searrow d & & \xrightarrow{\nabla + \text{id}} Y + Y \\
 \text{\scriptsize } b \searrow & & \downarrow \triangleright_1 \\
 & & Y + Y \xrightarrow{\nabla} Y \\
 & & \text{\scriptsize } \downarrow \triangleright_1
 \end{array}$$

The map  $(\triangleright_2 + \text{id}) \hat{\circ} d$  is a bound for  $g \perp h$ , indeed:

$$\begin{aligned}
 \triangleright_1 \hat{\circ} (\triangleright_2 + \text{id}) \hat{\circ} d &= \triangleright_2 \hat{\circ} \triangleright_1 \hat{\circ} d & \triangleright_2 \hat{\circ} (\triangleright_2 + \text{id}) \hat{\circ} d &= \triangleright_2 \hat{\circ} d \\
 &= \triangleright_2 \hat{\circ} b & &= \triangleright_2 \hat{\circ} (\nabla + \text{id}) \hat{\circ} d \\
 &= g & &= \triangleright_2 \hat{\circ} c \\
 & & &= h.
 \end{aligned}$$

We need one more:  $[\text{id}, \kappa_2] \hat{\circ} d$  is a bound for  $f \perp g \otimes h$ , indeed

$$\begin{aligned}
 \triangleright_1 \hat{\circ} [\text{id}, \kappa_2] \hat{\circ} d &= \triangleright_1 \hat{\circ} \triangleright_1 \hat{\circ} d & \triangleright_2 \hat{\circ} [\text{id}, \kappa_2] \hat{\circ} d &= \nabla \hat{\circ} (\triangleright_2 + \text{id}) \hat{\circ} d \\
 &= \triangleright_1 \hat{\circ} b & &= g \otimes h \\
 &= f.
 \end{aligned}$$

Finally, we compute

$$f \otimes (g \otimes h) = \nabla \hat{\circ} [\text{id}, \kappa_2] \hat{\circ} d = \nabla \hat{\circ} (\nabla + \text{id}) \hat{\circ} d = \nabla \hat{\circ} c = (f \otimes g) \otimes h,$$

which shows the partial associativity. Homsets of  $\text{Par } C$  are indeed PCMs.

(PCM-enrichment, II) Assume  $f \perp g$  with bound  $b$ . It is easy to see  $b \hat{\circ} k$  is a bound for  $f \hat{\circ} k \perp g \hat{\circ} k$  and consequently  $(f \hat{\circ} k) \otimes (g \hat{\circ} k) = \nabla \hat{\circ} b \hat{\circ} k = (f \otimes g) \hat{\circ} k$ . IV

For the other side:  $(h + h) \hat{\circ} b$  is a bound for  $h \hat{\circ} f \perp h \hat{\circ} g$ , indeed

$$\begin{aligned}
 \triangleright_1 \hat{\circ} (h + h) \hat{\circ} b &= h \hat{\circ} \triangleright_1 \hat{\circ} b = h \hat{\circ} f \\
 \triangleright_2 \hat{\circ} (h + h) \hat{\circ} b &= h \hat{\circ} \triangleright_2 \hat{\circ} b = h \hat{\circ} g.
 \end{aligned}$$

And so  $(h \hat{\circ} f) \otimes (h \hat{\circ} g) = \nabla \hat{\circ} (h + h) \hat{\circ} b = h \hat{\circ} \nabla \hat{\circ} b = h \hat{\circ} (f \otimes g)$ . Finally, the zero axiom holds by 186 VII.

(FinPAC) We just saw  $\text{Par } C$  is PCM-enriched. We already know  $\text{Par } C$  has coproducts. The compatible sum axiom holds by definition of  $\perp$ . To show  $\text{Par } C$  is a finPAC, only the untying axiom remains to be proven. If  $f \perp g$  with bound  $b$ , then  $(\kappa_1 + \kappa_2) \hat{\circ} b$  is a bound for  $\kappa_1 \hat{\circ} f \perp \kappa_2 \hat{\circ} g$ , which proves the untying axiom. V

VI (Effect algebra of predicates) Pick any predicate  $p: X \rightarrow 1 + 1$ . We define  $p^\perp = [\kappa_2, \kappa_1] \circ p$ . ( $p^\perp$  is  $p$  with swapped outcomes.) We compute

$$\begin{aligned}\triangleright_1 \hat{\circ} \hat{p} &= [\text{id}, \kappa_2] \circ \kappa_1 \circ p = p \\ \triangleright_2 \hat{\circ} \hat{p} &= [[\kappa_2, \kappa_1], \kappa_2] \circ \kappa_1 \circ p = [\kappa_2, \kappa_1] \circ p \\ \nabla \hat{\circ} \hat{p} &= [[\kappa_1, \kappa_1], \kappa_2] \circ \kappa_1 \circ p = \kappa_1 \circ ! = 1.\end{aligned}$$

So  $\hat{p}$  is a bound for  $p \perp p^\perp$  and  $p \otimes p^\perp = 1$ . To show  $p^\perp$  is the unique orthocomplement, assume  $p \otimes q = 1$  for some  $q$  via a bound  $b: X \rightarrow 1 + 1$ . Note  $1 \hat{\circ} b = \nabla \hat{\circ} b = 1$  and so by 186 VIII we know  $b = \hat{c}$  for some  $c: X \rightarrow 1 + 1$ . And so  $p = \triangleright_1 \hat{\circ} b = [\text{id}, \kappa_2] \circ \kappa_1 \circ c = c$ . Hence

$$q = \triangleright_2 \hat{\circ} b = [[\kappa_2, \kappa_1], \kappa_2] \circ \kappa_1 \circ c = [\kappa_2, \kappa_1] \circ p = p^\perp.$$

To show  $\text{Pred } X$  is an effect algebra, it only remains to be proven that the zero-one axiom holds. So assume  $1 \perp p$  via some bound  $b$ . We must show  $p = 0$ . By the following instance of the right pullback square of (3.1), we see  $b = \kappa_1 \circ \kappa_1 \circ !$ .

$$\begin{array}{ccc} X & \xrightarrow{!} & 1 \\ \text{dotted arrow} \searrow & & \downarrow \kappa_1 \\ & & 1 \\ \text{solid arrow } b \searrow & & \downarrow \kappa_1 \\ (1+1)+1 & \xrightarrow{[\text{id}, \kappa_2]} & 1+1 \\ & & \downarrow \kappa_1 \circ \kappa_1 \\ & & 1 \end{array}$$

Hence  $p = \triangleright_2 \hat{\circ} b = [[\kappa_2, \kappa_1], \kappa_1] \circ \kappa_1 \circ \kappa_1 \circ ! = \kappa_2 \circ ! = 0$ .

VII (Final axioms) In 186 VIII we proved that  $f = 0$  whenever  $1 \hat{\circ} f = 0$ . It only remains to be shown that  $f \perp g$  provided  $1 \hat{\circ} f \perp 1 \hat{\circ} g$ . Let  $b$  be a bound for  $1 \hat{\circ} f \perp 1 \hat{\circ} g$ . We apply the right pullback square of 186 IV twice in succession as follows (using  $\triangleright_2 \hat{\circ} c = \triangleright_2 \hat{\circ} (! + \text{id}) \hat{\circ} c = \triangleright_2 \hat{\circ} b = 1 \hat{\circ} g$  for the right one.)

$$\begin{array}{ccc} X & \xrightarrow{b} & 1+1 \\ \text{dotted arrow } c \searrow & & \downarrow \hat{\triangleright}_1 \\ & & Y+1 \xrightarrow{\hat{\text{id}}+\text{id}} 1+1 \\ \text{solid arrow } f \searrow & & \downarrow \hat{\triangleright}_1 \\ & & Y \xrightarrow{\hat{\text{id}}} 1 \end{array} \quad \begin{array}{ccc} X & \xrightarrow{c} & Y+1 \\ \text{dotted arrow } d \searrow & & \downarrow \hat{\triangleright}_2 \\ & & Y+Y \xrightarrow{\text{id}+\hat{\text{id}}} Y+1 \\ \text{solid arrow } g \searrow & & \downarrow \hat{\triangleright}_2 \\ & & Y \xrightarrow{\hat{\text{id}}} 1 \end{array}$$

Clearly  $\triangleright_2 \hat{\circ} d = g$  and  $\triangleright_1 \hat{\circ} d = \triangleright_1 \hat{\circ} (\text{id} + \hat{\text{id}}) \hat{\circ} d = \triangleright_1 \hat{\circ} c = f$ , so  $d$  is a bound for  $f \perp g$ , as desired.  $\square$

**188** We are ready to show the equivalence of effectuses in partial and total form.

**Proof of 180X** Let  $C$  and  $D$  be effectuses in total and respectively partial form. ||  
 The first point, that  $\text{Par } C$  is an effectus in partial form, is shown in 187I. The second point, that  $\text{Tot } D$  is an effectus in total form, is shown in 181XI. It remains to be shown that nothing is lost.

( $\text{Par Tot } D \cong D$ ) Let  $D$  be an effectus in partial form. A map  $f: X \rightarrow Y$  |||  
 in  $\text{Par Tot } D$  is by definition a map  $f: X \rightarrow Y + 1$  in  $D$  with  $1 \circ f = 1$ . Let  $P: \text{Par Tot } D \rightarrow D$  denote the identity-on-objects map  $f \mapsto \triangleright_1 \circ f$ . This is a functor: clearly  $P\hat{\text{id}} = P\kappa_1 = \text{id}$  and if  $g: Y \rightarrow Z$  in  $\text{Par Tot } D$ , then

$$\begin{aligned} \triangleright_1 \circ (g \hat{\circ} f) &= \triangleright_1 \circ [g, \kappa_2] \circ f \\ &= [\triangleright_1 \circ g, \triangleright_1 \circ \kappa_2] \circ f \\ &= [\triangleright_1 \circ g, 0] \circ (\triangleright_1 \circ f, \triangleright_2 \circ f) \\ &= \triangleright_1 \circ g \circ \triangleright_1 \circ f. \end{aligned}$$

The functor  $P$  is an isomorphism with inverse  $P'f \equiv \langle f, (1 \circ f)^\perp \rangle$ . Indeed, we have  $PP'f = \triangleright_1 \circ \langle f, (1 \circ f)^\perp \rangle = f$  and

$$P'Pf = \langle \triangleright_1 \circ f, (1 \circ \triangleright_1 \circ f)^\perp \rangle = (\kappa_1 \circ \triangleright_1 \circ f) \otimes (\kappa_2 \circ \triangleright_1 \circ f)^\perp = f.$$

( $\text{Tot Par } C \cong C$ ) Assume  $C$  is an effectus in total form. Recall  $1 \hat{\circ} \hat{g} = 1$  IV  
 and  $\widehat{f \circ g} = \hat{f} \hat{\circ} \hat{g}$ , so  $Q: C \rightarrow \text{Tot Par } C$  given by  $Qg = \hat{g}$  is an identity-on-objects functor. It is an isomorphism by the first part of 186VIII: for every  $f$  in  $\text{Tot Par } C$ , there is a unique  $g$  in  $C$  with  $f = \hat{g}$ . □

**Exercise** In this exercise we will distinguish effectuses in total and partial form. 189

1. To start, show that the initial object of an effectus in total form is strict — that is: show that any map into 0 is an isomorphism. (Hint: use the right pullback square of 185I with  $X = Y = 0$ .)
2. Conclude that if  $C$  is both an effectus in total and partial form, then every object in  $C$  is isomorphic to 0.

**Convention** For the remainder of this text, we will work with effectuses in partial form and write 1 instead of  $I$ . To spare ink, a phrase like “let  $C$  be an effectus” should be read as “let  $C$  be an effectus in partial form” and simply “ $C$  is an effectus” means either “ $C$  is an effectus in total form” or “ $C$  is an effectus in partial form”. (In non-trivial cases, this is unambiguous, as seen in I.) As before, when we took the effectus in partial form as base category, we will denote partial maps by  $f: X \rightarrow Y$  (instead of  $f: X \dashrightarrow Y$ ) and their composition as  $f \circ g$  (instead of  $f \hat{\circ} g$ ). ||

**189a Examples** Our main example of an effectus (in total form) is the category  $\text{vN}^{\text{op}}$  of von Neumann algebras with  $\text{ncpu}$ -maps between them in the opposite direction, see 180V. Its full subcategory  $\text{CvN}^{\text{op}}$  of commutative von Neumann algebras is an effectus as well.

- II In [23] several other examples of effectuses are discussed in detail:
  1. The category  $\text{OUS}^{\text{op}}$  of order unit spaces (i.e. ordered real vector spaces with distinguished order unit) with positive unit-preserving linear maps in the opposite direction is an effectus (in total form).
  2. A related example is the category  $\text{OUG}^{\text{op}}$  of order unit groups (i.e. ordered abelian groups with distinguished order unit) with positive unit-preserving homomorphisms in the opposite direction, which is also an effectus.
  3. Every extensive category [15] with final object is an effectus in total form. Examples of extensive categories with final object include
    - (a)  $\text{Set}$ , the category of sets with maps between them;
    - (b)  $\text{CRng}^{\text{op}}$ , the category of commutative unit rings with unit-preserving homomorphisms between them in the opposite direction;
    - (c)  $\text{CH}$ , the category of compact Hausdorff spaces with continuous maps between them; *and*
- III Together with van de Wetering, we recently investigated [128] the category  $\text{EJA}^{\text{op}}$  of Euclidean Jordan algebras with positive unit-preserving linear maps in the opposite direction, which is also an effectus (in total form).

### 3.2.3 Predicates, states and scalars

**190** We turn to the internal structures of an effectus. [23]

- II **Definition** Let  $C$  be an effectus (in partial form).
  1. As alluded to in the definition of an effectus in partial form, a **predicate** on  $X$  is a map  $X \rightarrow 1$ . The set of predicates  $\text{Pred } X$  on  $X$  is (by definition of an effectus in partial form) an effect algebra.
  2. A **scalar** is a predicate on  $1$ ; that is, a map  $1 \rightarrow 1$ . We write  $\text{Scal } C \equiv M \equiv \text{Pred } 1$  for the set of scalars. We multiply two scalars  $\lambda, \mu: 1 \rightarrow 1$  simply by composing  $\lambda \odot \mu \equiv \lambda \circ \mu$ . As  $1_M = \text{id}_M$  (see 181XIII) and  $C$  is PCM-enriched, the set of scalars  $M$  with  $\odot$  is an effect monoid, see 178II.
  3. A **real effectus** is an effectus where the set of scalars  $M$  is isomorphic (as effect monoid) to  $[0, 1]$ .

4. For a scalar  $\lambda: 1 \rightarrow 1$  and a predicate  $p: X \rightarrow 1$ , we write  $\lambda \cdot p \equiv \lambda \circ p$ . Again due to PCM-enrichment and  $1_M = \text{id}_M$ , this scalar multiplication turns  $\text{Pred } X$  into an  $M$ -effect module, see 179 II.
5. For  $f: X \rightarrow Y$  in  $C$ , define  $\text{Pred}(f): \text{Pred } Y \rightarrow \text{Pred } X$  by  $\text{Pred}(f)(p) = p \circ f$ . It is easy to see  $\text{Pred}(f)$  is an  $M$ -effect module homomorphism if  $f$  is total and that, in fact,  $\text{Pred}: \text{Tot } C \rightarrow \mathbf{EMod}_M^{\text{op}}$  is a functor.
6. A **substate** on  $X$  is a map  $\omega: 1 \rightarrow X$ . A **state** is a total substate. We denote the set of states on  $X$  by  $\text{Stat } X$ .
7. An effectus  $C$  has **separating predicates** if  $\text{Pred } X$  (as a set of maps  $X \rightarrow 1$ ) is jointly monic for every  $X$  in  $C$ . Similarly, an effectus has **separating states** if  $\text{Stat } X$  (as a set of maps  $1 \rightarrow X$ ) is jointly epic for every  $X$  in  $C$ .

**Examples** The category  $\mathbf{vN}^{\text{op}}$  (see 180 V) is a real effectus ( $M \cong [0, 1]$ ) with separating states and predicates. [23] The predicates on a von Neumann algebra  $\mathcal{A}$  correspond to the set of effects  $[0, 1]_{\mathcal{A}}$  and  $\text{Stat } \mathcal{A}$  can be identified with the convex set of normal states. III

We briefly cover the examples from [23] mentioned in 189a I. IV

1. The category  $\mathbf{OUS}^{\text{op}}$  (see 189a I) is a real effectus with separating predicates, but without separating states. (The states on a single order unit space are separating if and only if the order unit space is archimedean.) A predicate on an order unit space  $X$  corresponds to a point  $x \in X$  with  $0 \leq x \leq 1$ , where 1 is the distinguished order unit. The states are exactly what are called states for order unit spaces in the literature.
2. The category  $\mathbf{OUG}^{\text{op}}$  (see 189a I) has the two-element effect monoid  $2$  as scalars and separating predicates. The predicates of an order unit group  $G$  correspond to the elements  $x \in G$  with  $0 \leq x \leq 1$ , where 1 is the distinguished order unit. States on  $G$  correspond to unit-preserving positive homomorphisms  $G \rightarrow \mathbb{Z}$ , which are not separating.
3. Any extensive category with final object (such as  $\mathbf{Set}$ ,  $\mathbf{CRng}^{\text{op}}$  and  $\mathbf{CH}$ ) has as scalars the two-element effect monoid  $2$ .
  - (a) In  $\mathbf{Set}$  the predicates on a set  $X$  correspond to subsets  $U \subseteq X$  and are thus separating. States on  $X$  correspond to elements  $x \in X$ , which are also separating.
  - (b) In  $\mathbf{CRng}^{\text{op}}$  the predicates on  $R$  correspond to idempotents of  $R$ , which are not separating. States correspond to unit-preserving homomorphisms  $R \rightarrow \mathbb{Z}$ , which are not separating either.

(c) In CH the predicates on  $X$  correspond to clopen subsets  $U \subseteq X$ , which are not separating. States correspond on  $X$  correspond to its points  $x \in X$ , which are separating.

✓ The category  $\text{EJA}^{\text{op}}$  is a real effectus with separating states and predicates. The predicates on an Euclidean Jordan algebra (EJA)  $E$  correspond to its effects; i.e. the elements  $0 \leq a \leq 1$ . The states are exactly what are usually considered states for EJAs.

**191** Does every effect monoid  $M$  occur as the effect monoid of scalars of some effectus? Does every  $M$ -effect module occur as effect module of predicates? We will show they do, in the following strong sense.


II **Theorem** Let  $M$  be any effect monoid. The category  $\text{EMod}_M^{\text{op}}$  is an effectus in total form with scalars  $M$  and separating predicates. [23] In fact: every effectus  $C$  in total form with scalars  $M$  and separating predicates is equivalent to a subcategory of  $\text{EMod}_M^{\text{op}}$ .

III **Proof** To show  $\text{EMod}_M^{\text{op}}$  is an effectus, we have to show the dual axioms for  $\text{EMod}_M$ .

IV (Finite products) For every  $M$ -effect module  $E$ , there is a unique  $M$ -effect module map  $!: M \rightarrow E$ , which is given by  $\lambda \mapsto \lambda \cdot 1$ . So  $M$  is the initial object of  $\text{EMod}_M$ . The one-element  $M$ -effect module  $\{0 = 1\}$  is the final object.

If  $E$  and  $F$  are  $M$ -effect modules, then the set of pairs  $E \times F$  with componentwise operations is again an  $M$ -effect module. The effect module  $E \times F$  with obvious projections  $E \xleftarrow{\pi_1} E \times F \xrightarrow{\pi_2} F$  is the categorical product of  $E$  and  $F$  in  $\text{EMod}_M$  (as is the case with effect algebras, see 175 III).

V (Pushout diagrams) To show the pushout diagrams corresponding to (3.1) hold, assume we are given  $M$ -effect modules  $E, F, G$  with module maps  $\alpha, \beta, \delta$  that make the outer squares of the following diagrams commute.



We have to show that there are unique dashed arrows (as shown) that make these diagrams commute. We start with the left diagram. By assumption  $\alpha \circ (! \times \text{id}) = \beta \circ (\text{id} \times !)$ , so in particular  $\alpha(1, 0) = \beta(1, 0)$ . For any  $(x, y) \in E \times F$ , we have  $\alpha(x, 0) \leq \alpha(1, 0) = \beta(1, 0) \perp \beta(0, 1) \geq \beta(0, y)$ , so  $\alpha(x, 0) \perp \beta(0, y)$ . Define  $f: E \times F \rightarrow G$  by  $f(x, y) = \alpha(x, 0) \odot \beta(0, y)$ . It is easy to see  $f$  is (partially) additive and  $f(1, 1) = \alpha(1, 0) \odot \beta(0, 1) = \beta(1, 1) = 1$ , so  $f$  is a

module map. We compute

$$f(x, \lambda \cdot 1) = \alpha(x, 0) \otimes \beta(0, \lambda \cdot 1) = \alpha(x, 0) \otimes \lambda \cdot \alpha(0, 1) = \alpha(x, \lambda)$$

and so  $f \circ (\text{id} \times !) = \alpha$ . Similarly  $\beta = f \circ (! \times \text{id})$ . It is easy to see  $f$  is the unique map with  $f \circ (\text{id} \times !) = \alpha$  and  $\beta = f \circ (! \times \text{id})$  and so the left square of (3.1) is a pullback in  $\mathbf{EMod}_M^{\text{op}}$ .

Concerning the right diagram: as  $\delta \circ (! \times !) = ! \circ \pi_1$ , we have  $\delta(\lambda \cdot 1, \mu \cdot 1) = \lambda \cdot 1$  and so  $\delta(0, y) \leq \delta(0, 1) = 0$ . Hence  $\delta(x, y) = \delta(x, 0)$ . Define  $g: E \rightarrow G$  by  $g(x) = \delta(x, 0)$ . Clearly  $g$  is additive,  $g(1) = \delta(1, 0) = \delta(1, 1) = 1$  and  $\delta = g \circ \pi_1$  by definition. Obviously  $g$  is the unique such map. Thus the right square of (3.1) is a pullback in  $\mathbf{EMod}_M^{\text{op}}$ .

(Joint epicity) To show  $\mathbf{EMod}_M^{\text{op}}$  is an effectus, it only remains to be shown VI that  $\langle \pi_1, \pi_2, \pi_2 \rangle, \langle \pi_2, \pi_1, \pi_2 \rangle: M \times M \rightarrow M \times M \times M$  are jointly epic. So assume  $f, g: M \times M \times M \rightarrow E$  are two  $M$ -effect module maps with  $f \circ \langle \pi_1, \pi_2, \pi_2 \rangle = g \circ \langle \pi_1, \pi_2, \pi_2 \rangle$  and  $f \circ \langle \pi_2, \pi_1, \pi_2 \rangle = g \circ \langle \pi_2, \pi_1, \pi_2 \rangle$ . From the first equality, it follows that  $f(1, 0, 0) = g(1, 0, 0)$ . The other one implies  $f(0, 1, 0) = g(0, 1, 0)$ . Thus  $f(0, 0, 1) = f(1, 1, 0)^\perp = g(1, 1, 0)^\perp = g(0, 0, 1)$ . As for  $(\lambda_1, \lambda_2, \lambda_3) \in M^3$ , we have  $(\lambda_1, \lambda_2, \lambda_3) = \lambda_1 \cdot (1, 0, 0) \otimes \lambda_2 \cdot (0, 1, 0) \otimes \lambda_3 \cdot (0, 0, 1)$  we get  $f = g$ .

(Representation) Let  $C$  be an effectus in total form. Clearly  $\text{Pred } f = \text{Pred } g$  VII (for  $f, g: X \rightarrow Y$  in  $C$ ) if and only if  $p \circ f = (\text{Pred } f)(p) = (\text{Pred } g)(p) = p \circ g$  for every  $p \in \text{Pred } X$ . Thus the functor  $\text{Pred}: C \rightarrow \mathbf{EMod}_M^{\text{op}}$  is faithful if and only if  $C$  has separating predicates. So if  $C$  has separating predicates,  $C$  is equivalent to the subcategory  $\text{Pred } C$  of  $\mathbf{EMod}_M^{\text{op}}$ .  $\square$

**Exercise** Show that the category  $\mathbf{Rng}^{\text{op}}$  of unit rings with unit-preserving homomorphisms in the opposite direction is an effectus in total form. VIII

1. Show that the predicates on a ring  $R$  correspond to its idempotents; that  $p \perp q$  iff  $pq = qp = 0$ ,  $p^\perp = 1 - p$  and  $p \otimes q = p + q$ . (So 2 is its effect monoid of scalars.) Conclude  $\mathbf{Rng}^{\text{op}}$  does not have separating predicates.
2. Show that there is no unit-preserving ring-homomorphism  $\mathbb{Z}_2 \rightarrow \mathbb{Z}$ . Conclude  $\mathbf{Rng}^{\text{op}}$  does not have separating states.

We studied the structure of predicates in an effectus. What about the states? 192 They will turn out to form an abstract  $M$ -convex set. Before we define these, we introduce a generalized distribution monad.

**Definition** Let  $M$  be an effect monoid and  $X$  be any set. A formal  $M$ -convex combination over  $X$  is a function  $p: X \rightarrow M$  with finite support such II that  $\bigotimes_{x \in X} p(x) = 1$ . We use  $\lambda_1 |x_1\rangle \otimes \cdots \otimes \lambda_n |x_n\rangle$  as a shorthand for the formal  $M$ -convex combination  $p: X \rightarrow M$  given by  $p(x) = \bigotimes_{x_i=x} \lambda_i$ . (So  $p(x_i) = \lambda_i$  if the  $x_i$  are distinct.) We write  $\mathcal{D}_M X$  for the set of all formal  $M$ -convex combinations over  $X$ . [56]

III **Exercise\*** In this exercise you study  $\mathcal{D}_M$  as a monad. See [56, 129].

1. For  $f: X \rightarrow Y$ , define  $\mathcal{D}_M f: \mathcal{D}_M X \rightarrow \mathcal{D}_M Y$  by

$$(\mathcal{D}_M f)(p)(y) = \bigvee_{x: f(x)=y} p(x).$$

(That is:  $(\mathcal{D}_M f)(\lambda_1 |x_1\rangle \otimes \cdots \otimes \lambda_n |x_n\rangle) = \lambda_1 |f(x_1)\rangle \otimes \cdots \otimes \lambda_n |f(x_n)\rangle$ .)  
Show that this turns  $\mathcal{D}_M$  into a functor  $\mathbf{Set} \rightarrow \mathbf{Set}$ .

2. Define  $\eta: X \rightarrow \mathcal{D}_M X$  by  $\eta(x) = 1 |x\rangle$  and  $\mu: \mathcal{D}_M \mathcal{D}_M X \rightarrow \mathcal{D}_M X$  by

$$\mu(\Phi)(x) = \bigvee_{\varphi} \Phi(\varphi) \odot \varphi(x).$$

(That is:  $\mu(\bigvee_i \sigma_i | \bigvee_j \lambda_{ij} |x_{ij}\rangle) = \bigvee_{i,j} (\sigma_i \odot \lambda_{ij}) |x_{ij}\rangle$ .)  
Show that  $(\mathcal{D}_M, \eta, \mu)$  is a monad.

3. Show that  $\mathcal{Kl} \mathcal{D}_M$ , the Kleisli category of  $\mathcal{D}_M$  (see e.g. [129, §2.6]), is an effectus with  $M$  as scalars.

IV **Definition** Let  $X$  be a set together with a map  $h: \mathcal{D}_M X \rightarrow X$ . We say  $(X, h)$  is an **abstract  $M$ -convex set** provided  $h(|x\rangle) = x$  and

$$h\left(\bigvee_{i,j} \sigma_i \odot \lambda_{ij} |x_{ij}\rangle\right) = h\left(\bigvee_i \sigma_i | h\left(\bigvee_j \lambda_{ij} |x_{ij}\rangle\right)\right) \quad (3.2)$$

for every  $\bigvee_i \sigma_i | \bigvee_j \lambda_{ij} |x_{ij}\rangle$  in  $\mathcal{D}_M \mathcal{D}_M X$ . (Equivalently:  $h \circ \mu = h \circ \mathcal{D}_M h$  and  $h \circ \eta = \text{id}$ .) We call  $h$  the convex structure on  $X$ . A map  $f: X \rightarrow Y$  between abstract  $M$ -convex sets is  **$M$ -affine** if

$$f(h_X(\bigvee_i \lambda_i |x_i\rangle)) = h_Y(\bigvee_i \lambda_i |f(x_i)\rangle)$$

for every  $\bigvee_i \lambda_i |x_i\rangle$  in  $\mathcal{D}_M X$ . (Equivalently:  $h_X \circ \mathcal{D}_M f = f \circ h_Y$ .) Write  **$\mathbf{AConv}_M$**  for the category of abstract  $M$ -convex sets with  $M$ -affine maps between them. (Equivalently:  **$\mathbf{AConv}_M$**  is the Eilenberg–Moore category of the monad  $\mathcal{D}_M$ .) We call an abstract  $M$ -convex set  $(X, h)$  **cancellative** provided

$$h(\lambda |y\rangle \otimes \lambda^\perp |x_1\rangle) = h(\lambda |y\rangle \otimes \lambda^\perp |x_2\rangle) \quad \text{implies} \quad x_1 = x_2$$

for any  $x_1, x_2, y \in X$  and  $\lambda \in M, \lambda \neq 1$ .

IVa **Remark** For an abstract  $[0, 1]$ -convex set  $(X, h)$ , binary convex combinations are sufficient: for instance, for any  $\lambda \otimes \mu \neq 1$  and  $x, y, z \in X$  we have

$$h(\lambda |x\rangle \otimes \mu |y\rangle \otimes (\lambda \otimes \mu)^\perp |z\rangle) = h\left(\lambda |x\rangle \otimes \lambda^\perp |h(\mu/\lambda^\perp |y\rangle \otimes (\lambda \otimes \mu)^\perp / \lambda^\perp |z\rangle)\right).$$

This reduction to binary convex combinations uses the division on  $[0, 1]$ . Some effect monoids have a suitable division as well, but many do not as we will see later on in 195II and 195VI. It seems plausible that it is impossible to write an arbitrary three-element  $M$ -convex combination over  $\mathcal{D}_M\{x, y, z\}$  using only binary convex combinations in the case that  $M$  is the effect monoid on the unit interval of  $C[0, 1]$ . I do not have a proof to back this claim.

**Examples** We give some examples of abstract convex sets. V

1. Every convex subset  $X \subseteq V$  of some real vector space  $V$  is a cancellative abstract  $[0, 1]$ -convex set with  $h(\lambda_1 |x_1\rangle \otimes \cdots \otimes \lambda_n |x_n\rangle) = \lambda_1 x_1 + \cdots + \lambda_n x_n$ .
2. Not every abstract  $[0, 1]$ -convex set is a convex subset of a real vector space. Consider  $T \subseteq [0, 1]^2$  defined by  $T \equiv \{(x, y); x + y \leq 1\}$ . Say  $(x, y) \sim (x', y')$  whenever  $x = x'$  and either  $x \neq 0$  or  $y = y'$ . This equivalence relation  $\sim$  is a congruence (we will cover these in 193II) for the convex structure on  $T$  (inherited by  $\mathbb{R}^2$ ) and so  $T/\sim$  is an abstract  $[0, 1]$ -convex set. That is: we start off with a filled triangle  $T \equiv \triangle$ , identify all vertical lines except for the y-axis and are left with  $T/\sim \equiv \perp$ . The abstract  $[0, 1]$ -convex set  $T/\sim$  is not cancellative:

$$h\left(\frac{1}{2} |\perp\rangle \otimes \frac{1}{2} |\perp\rangle\right) = \perp = h\left(\frac{1}{2} |\perp\rangle \otimes \frac{1}{2} |\ulcorner\rangle\right), \quad \text{but } \perp \neq \ulcorner.$$

3. Semilattices are exactly abstract 2-convex sets. (2 is the two-element Effect Monoid, see 178III.) Furthermore, every semilattice  $(L, \vee)$  is an abstract  $[0, 1]$ -convex set with

$$h(\lambda_1 |x_1\rangle \otimes \cdots \otimes \lambda_n |x_n\rangle) = \bigvee_{i; \lambda_i \neq 0} x_i.$$

A semilattice  $L$  is cancellative as abstract  $[0, 1]$ -convex set if and only if  $x = y$  for all  $x, y \in L$ . See [90].

4. Every cancellative  $[0, 1]$ -convex set is isomorphic to a convex subset of a real vector space. See e.g. [69, thm. 8].

**Remarks** The study of convex subsets (of real vector spaces) as algebraic structures goes back a long time, see e.g. [30, 33, 42, 90, 107]. The more pathological abstract  $[0, 1]$ -convex sets have been studied before under different names: they are semiconvex sets in [30, 109] and convex spaces in [33]. The description of convex sets using Eilenberg–Moore algebras is probably due to Świrszcz [109]. For the even more exotic abstract  $M$ -convex sets, see [23] or [56]. VI

**Proposition** Let  $C$  be an effectus with scalars  $M$ . For any object  $X$ , the set of states  $\text{Stat } X$  is an abstract  $M^{\text{op}}$ -convex set with  $h: \mathcal{D}_{M^{\text{op}}} \text{Stat } X \rightarrow \text{Stat } X$  VII

defined by  $h(\lambda_1 |\varphi_1\rangle \otimes \cdots \otimes \lambda_n |\varphi_n\rangle) = [\varphi_1, \dots, \varphi_n] \circ \langle \lambda_1, \dots, \lambda_n \rangle$ . Furthermore, for any total map  $f: X \rightarrow Y$ , the map  $(\text{Stat } f)(\varphi) = f \circ \varphi$  is an  $M^{\text{op}}$ -affine map  $\text{Stat } X \rightarrow \text{Stat } Y$  yielding a functor  $\text{Stat}: \text{Tot } C \rightarrow \mathbf{AConv}_{M^{\text{op}}}$ . [23]

**VIII Proof** Clearly  $h(|\varphi\rangle) = \varphi$ . To show the other axiom (3.2), assume we have any  $\bigotimes_{i=1}^n \sigma_i | \bigotimes_{j=1}^{m_i} \lambda_{ij} |\varphi_{ij}\rangle \rangle$  in  $\mathcal{D}_{M^{\text{op}}} \mathcal{D}_{M^{\text{op}}} \text{Stat } X$ . Write  $\lambda_i \equiv \langle \lambda_{i1}, \dots, \lambda_{im_i} \rangle$  and  $\varphi_i \equiv [\varphi_{i1}, \dots, \varphi_{im_i}]$ . Then

$$\begin{aligned}
h\left(\bigotimes_i \sigma_i | h\left(\bigotimes_j \lambda_{ij} |\varphi_{ij}\rangle\right)\right) &= [\varphi_1 \circ \lambda_1, \dots, \varphi_n \circ \lambda_n] \circ \langle \sigma_1, \dots, \sigma_n \rangle \\
&= \bigotimes_i \varphi_i \circ \lambda_i \circ \sigma_i \\
&= \bigotimes_i \left(\bigotimes_j \varphi_{ij} \circ \lambda_{ij}\right) \circ \sigma_i \\
&= \bigotimes_{i,j} \varphi_{ij} \circ (\lambda_{ij} \circ \sigma_i) \\
&= \bigotimes_{i,j} \varphi_{ij} \circ (\sigma_i \circ_{M^{\text{op}}} \lambda_{ij}) \\
&= h\left(\bigotimes_{i,j} \sigma_i \circ_{M^{\text{op}}} \lambda_{ij} |\varphi_{ij}\rangle\right).
\end{aligned}$$

So  $\text{Stat } X$  is indeed an abstract  $M^{\text{op}}$ -convex set. Let  $f: X \rightarrow Y$  be any total map. To show  $\text{Stat } f$  is affine, it is sufficient to show that for every  $\bigotimes_i \lambda_i |\varphi_i\rangle$  in  $\mathcal{D}_{M^{\text{op}}}$ , we have  $f \circ \bigotimes_i \varphi_i \circ \lambda_i = \bigotimes_i f \circ \varphi_i \circ \lambda_i$ , which is clearly true. Functoriality of  $\text{Stat}$  is obvious.  $\square$

### 3.2.4 Effectus of abstract $M$ -convex sets

**193** For our next project, we indulge ourselves in a tangent: we investigate whether the category  $\mathbf{AConv}_M$  is an effectus. We can show it is, if  $M$  has a certain partial division. Unfortunately it will remain unclear if  $\mathbf{AConv}_M$  is an effectus in general. The case  $M = [0, 1]$  is much easier and has already been dealt with in [69]. The first step is to study the coproduct in  $\mathbf{AConv}_M$ . To construct the coproduct, we will need to know about quotients of abstract  $M$ -convex sets.

**II Exercise** In this exercise you construct the quotient of an abstract  $M$ -convex set by a congruence. Assume  $(X, h)$  is an abstract  $M$ -convex set and  $\sim \subseteq X^2$  is an equivalence relation. Write  $X/\sim$  for the set of equivalence classes of  $\sim$  and  $q: X \rightarrow X/\sim$  for  $q(x) = [x]_\sim$ , where  $[x]_\sim$  is the equivalence class of  $x$ . For  $\varphi, \psi \in \mathcal{D}_M X$ , we write  $\varphi \sim \psi$  provided  $(\mathcal{D}_M q)(\varphi) = (\mathcal{D}_M q)(\psi)$ . We say  $\sim$  is a **congruence** for  $(X, h)$  if  $\varphi \sim \psi$  implies  $h(\varphi) \sim h(\psi)$ .

1. Prove that  $q$ ,  $\mathcal{D}_M q$  and  $\mathcal{D}_M \mathcal{D}_M q$  are all surjective.

2. Show that the following are equivalent.

(a)  $\sim$  is a congruence.

(b) There is a map  $h_\sim: \mathcal{D}_M X/\sim \rightarrow X/\sim$  such that  $h_\sim \circ \mathcal{D}_M q = q \circ h$ .

By the previous point  $h_\sim$  is unique, if it exists.

3. Assume  $\sim$  is a congruence. Use the previous two points to show

$$h_\sim \circ \mathcal{D}_M h_\sim \circ \mathcal{D}_M \mathcal{D}_M q = h_\sim \circ \mu \circ \mathcal{D}_M \mathcal{D}_M q.$$

Conclude that  $(X/\sim, h_\sim)$  is an abstract  $M$ -convex set and that the quotient map  $q: X \rightarrow X/\sim$  is  $M$ -affine.

**Exercise** Assume  $f: X \rightarrow Y$  is an affine map between abstract  $M$ -convex sets. III  
Show that the kernel  $\{(x, y); f(x) = f(y)\}$  of  $f$  is a congruence.

**Exercise** Assume  $(X, h)$  is an abstract  $M$ -convex set and  $R \subseteq X^2$  is some IV  
relation. It is easy to see there is a least congruence containing  $R$ . We need to know a bit more than mere existence. Write  $R^*$  for the reflexive symmetric transitive closure of  $R$  (i.e. the least equivalence relation containing  $R$ ).

For  $\psi_1, \psi_2 \in \mathcal{D}_M X$ , we write  $\psi_1 \approx \psi_2$  if there is a *derivation*: a tuple  $\varphi_1, \dots, \varphi_n \in \mathcal{D}_M X$  (say  $\varphi_i \equiv \bigotimes_{j=1}^{n_i} \lambda_{ij} |x_{ij}\rangle$ ) such that  $\varphi_1 = \psi_1$ ,  $\varphi_n = \psi_2$  and for each  $1 \leq i < n$  one of the following two conditions holds.

1.  $n_i = n_{i+1}$  and for all  $1 \leq j \leq n_i$  we have  $x_{ij} R^* x_{(i+1)j}$  and  $\lambda_{ij} = \lambda_{(i+1)j}$ .

2.  $h(\varphi_i) = h(\varphi_{i+1})$ .

We will show that  $\sim$  given by  $x \sim y \iff \eta(x) \approx \eta(y)$  is the least congruence containing  $R$ :

1. To start, show that  $\eta(h(\psi)) \approx \psi$  and so  $\varphi \approx \psi$  implies  $h(\varphi) \sim h(\psi)$ .

2. Show that if  $\varphi \approx \psi$ , then

$$\mu\left(\lambda_0 |\psi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle\right) \approx \mu\left(\lambda_0 |\varphi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle\right).$$

for any  $\bigotimes_j \lambda_j = 1$  in  $M$  and  $\chi_1, \dots, \chi_n \in \mathcal{D}_M X$ .

3. Use the previous point to show that  $\varphi \sim \psi$  implies  $\varphi \approx \psi$ . Conclude  $\sim$  is the smallest congruence containing  $R$ .

- V **Proposition** Assume  $(X, h_X)$  and  $(Y, h_Y)$  are abstract  $M$ -convex sets. We will construct their coproduct. Note that  $\mu$  turns  $\mathcal{D}_M(X + Y)$  into an abstract  $M$ -convex set. Let  $\sim$  denote the least congruence on  $\mathcal{D}_M(X + Y)$  with

$$(\mathcal{D}_M\kappa_1)(\varphi) \sim \eta(\kappa_1(h_X(\varphi))) \quad \text{and} \quad (\mathcal{D}_M\kappa_2)(\psi) \sim \eta(\kappa_2(h_Y(\psi))) \quad (3.3)$$

for all  $\varphi \in \mathcal{D}_M X$  and  $\psi \in \mathcal{D}_M Y$ . (Alternatively:  $\sim$  is the least congruence that makes  $\eta \circ \kappa_1$  and  $\eta \circ \kappa_2$  affine.) Write  $q: \mathcal{D}_M(X + Y) \rightarrow C$  for the quotient of  $\mathcal{D}_M(X + Y)$  by  $\sim$ , see II. Denote the convex structure on  $C$  by  $h$ . Define  $c_1: X \rightarrow C$  and  $c_2: Y \rightarrow C$  by  $c_1 \equiv q \circ \eta \circ \kappa_1$  and  $c_2 \equiv q \circ \eta \circ \kappa_2$ .

Then:  $c_1: X \rightarrow C \leftarrow Y: c_2$  is a coproduct of  $X$  and  $Y$  in  $\text{AConv}_M$ .

- VI **Proof** We start with the affinity of the coprojections:

$$\begin{aligned} c_1 \circ h_X &= q \circ \eta \circ \kappa_1 \circ h_X \\ &= q \circ \mathcal{D}_M\kappa_1 && \text{by dfn. of } \sim \\ &= q \circ \mu \circ \mathcal{D}_M\eta \circ \mathcal{D}_M\kappa_1. \\ &= \mu \circ \mathcal{D}_M q \circ \mathcal{D}_M\eta \circ \mathcal{D}_M\kappa_1. \\ &= \mu \circ \mathcal{D}_M c_1, \end{aligned}$$

so  $c_1$  is affine. With similar reasoning, we see  $c_2$  is affine.

- VII Assume  $f: X \rightarrow Z$  and  $g: Y \rightarrow Z$  are two affine maps to some abstract  $M$ -convex set  $(Z, h_Z)$ . We want to show  $h_Z \circ \mathcal{D}_M[f, g]$  lifts to  $C$ . With an easy calculation, one sees  $h_Z \circ \mathcal{D}_M[f, g]$  is affine. We compute

$$\begin{aligned} h_Z \circ \mathcal{D}_M[f, g] \circ \mathcal{D}_M\kappa_1 &= h_Z \circ \mathcal{D}_M f \\ &= f \circ h_X \\ &= h_Z \circ \eta \circ [f, g] \circ \kappa_1 \circ h_X \\ &= h_Z \circ \mathcal{D}_M[f, g] \circ \eta \circ \kappa_1 \circ h_X. \end{aligned}$$

So  $((\mathcal{D}_M\kappa_1)(\varphi), \eta(\kappa_1(h_X(\varphi))))$  is in the kernel (see III) of  $h_Z \circ \mathcal{D}_M[f, g]$  for  $\varphi \in \mathcal{D}_M$ . Similarly  $((\mathcal{D}_M\kappa_2)(\psi), \eta(\kappa_2(h_Y(\psi))))$  is also in the kernel of  $h_Z \circ \mathcal{D}_M[f, g]$  for all  $\psi \in \mathcal{D}_M$ .

As kernels of affine maps are congruences and  $\sim$  is the least congruence that contains these pairs, we conclude  $\sim$  is a subset of the kernel of  $h_Z \circ \mathcal{D}_M[f, g]$ . Thus there is a unique affine  $k: C \rightarrow Z$  fixed by  $k \circ q = h_Z \circ \mathcal{D}_M[f, g]$ . Now

$$\begin{aligned} k \circ c_1 &= k \circ q \circ \eta \circ \kappa_1 \\ &= h_Z \circ \mathcal{D}_M[f, g] \circ \eta \circ \kappa_1 \\ &= h_Z \circ \eta \circ [f, g] \circ \kappa_1 \\ &= f \end{aligned}$$

and similarly  $k \circ c_2 = g$ .

Only uniqueness of  $k$  remains. So assume  $k': C \rightarrow Z$  is an affine map such VIII  
that  $k' \circ c_1 = f$  and  $k' \circ c_2 = g$ . We turn the wheel:

$$\begin{aligned}
k' \circ q &= k' \circ q \circ \mu \circ \mathcal{D}_M \eta \\
&= k' \circ h \circ \mathcal{D}_M(q \circ \eta) \\
&= h_Z \circ \mathcal{D}_M(k' \circ q \circ \eta) \\
&= h_Z \circ \mathcal{D}_M[k' \circ q \circ \eta \circ \kappa_1, k' \circ q \circ \eta \circ \kappa_2] \\
&= h_Z \circ \mathcal{D}_M[k' \circ c_1, k' \circ c_2] \\
&= h_Z \circ \mathcal{D}_M[f, g] \\
&= k \circ q.
\end{aligned}$$

So  $k = k'$  by surjectivity of  $q$ . □

**Remark** The existence of coproducts of abstract  $M$ -convex sets already follows IX  
from the more general theorem of Linton [86]. (Cf. [10].) However, for the results in the remainder of this section we need to know more about the coproduct than Linton's construction provides.

By our construction, we know that every  $z \in X + Y$  is of the form

$$z = h\left(\bigvee_{i=1}^n \lambda_i |c_1(x_i)\rangle \otimes \bigvee_{i=1}^m \sigma_i |c_2(y_i)\rangle\right)$$

for some  $\lambda_i, \sigma_i \in M$ ,  $x_i \in X$  and  $y_i \in Y$ . In general these are not unique. Indeed using IV, we see that

$$h\left(\bigvee_{i=1}^n \lambda_i |c_1(x_i)\rangle \otimes \bigvee_{i=1}^m \sigma_i |c_2(y_i)\rangle\right) = h\left(\bigvee_{i=1}^{n'} \lambda'_i |c_1(x'_i)\rangle \otimes \bigvee_{i=1}^{m'} \sigma'_i |c_2(y'_i)\rangle\right)$$

iff there is a *derivation*  $\Phi_1, \dots, \Phi_l \in \mathcal{D}_M^2(X + Y)$ , say  $\Phi_i \equiv \bigvee_j \zeta_{ij} |\varphi_{ij}\rangle$ , with

$$\Phi_1 = \left| \bigvee_{i=1}^n \lambda_i |\kappa_1(x_i)\rangle \otimes \bigvee_{i=1}^m \sigma_i |\kappa_2(y_i)\rangle \right\rangle, \quad \Phi_l = \left| \bigvee_{i=1}^{n'} \lambda'_i |\kappa_1(x_i)\rangle \otimes \bigvee_{i=1}^{m'} \sigma'_i |\kappa_2(y_i)\rangle \right\rangle$$

and for each  $1 \leq i < l$  either

1.  $\mu(\Phi_i) = \mu(\Phi_{i+1})$  or
2. for each  $j$  we have  $\zeta_{ij} = \zeta_{(i+1)j}$  and one of the following.
  - (a)  $\varphi_{ij} = \varphi_{(i+1)j}$ .
  - (b)  $\varphi_{ij} = (\mathcal{D}_M \kappa_1)(\chi)$  and  $\varphi_{(i+1)j} = (\mathcal{D}_M \kappa_1)(\chi')$  for some  $\chi, \chi' \in \mathcal{D}_M X$  with  $h_X(\chi) = h_X(\chi')$ .
  - (c)  $\varphi_{ij} = (\mathcal{D}_M \kappa_2)(\chi)$  and  $\varphi_{(i+1)j} = (\mathcal{D}_M \kappa_2)(\chi')$  for some  $\chi, \chi' \in \mathcal{D}_M Y$  with  $h_Y(\chi) = h_Y(\chi')$ .

- X **Exercise** Show that the one-element set is an abstract  $M$ -convex set and, in fact, the final object of  $\mathbf{AConv}_M$ . Moreover, show that in  $\mathbf{AConv}_M$

$$n \cdot 1 \equiv \underbrace{1 + \cdots + 1}_{n \text{ times}} \cong \mathcal{D}_M\{1, \dots, n\}.$$

(You might want to use that  $\mathcal{D}_M$ , as a left-adjoint, preserves coproducts.)

- 194 **Proposition** The category  $\mathbf{AConv}_M$  obeys all the axioms of an effectus in total form except perhaps for the left pullback square of (3.1).

- II **Proof** In 193V we proved  $\mathbf{AConv}_M$  has binary coproducts and in 193X that the one-element set  $1$  is its final object. As  $\mathcal{D}_M\emptyset = \emptyset$ , the empty set is trivially also an abstract  $M$ -convex set and in fact the initial object of  $\mathbf{AConv}_M$ .

- III We continue with joint monicity of  $[\kappa_1, \kappa_2, \kappa_2], [\kappa_2, \kappa_1, \kappa_2]: 1 + 1 + 1 \rightarrow 1 + 1$ . We can represent the convex set  $1 + 1 + 1$  as the set of triples  $(a, b, c) \in M^3$  with  $a \otimes b \otimes c = 1$  with the obvious convex structure, cf. 193X. Similarly, we identify  $1 + 1$  with pairs  $(a, a^\perp) \in M^2$ . The maps at hand are given by

$$(a, b, c) \mapsto (a, b \otimes c) \quad \text{and} \quad (a, b, c) \mapsto (b, a \otimes c). \quad (3.4)$$

Assume  $(a, b \otimes c) = (a', b' \otimes c')$  and  $(b, a \otimes c) = (b', a' \otimes c')$  for  $a, a', b, b', c, c' \in M$  with  $a \otimes b \otimes c = 1$  and  $a' \otimes b' \otimes c' = 1$ . Then clearly  $a = a'$ ,  $b = b'$  and so  $c = (a \otimes b)^\perp = (a' \otimes b')^\perp = c'$ . Thus the maps (3.4) are jointly injective, hence jointly monic.

- IV At last, we are ready to prove that the square on the right of (3.1) is a pullback diagram in  $\mathbf{AConv}_M$ . So assume  $\alpha: Z \rightarrow X + Y$  is a map in  $\mathbf{AConv}_M$  with  $(!+) \circ \alpha = \kappa_1 \circ !$ .

We have to show  $\alpha = \kappa_1 \circ \gamma$ , for a unique  $\gamma: Z \rightarrow X + Y$  in  $\mathbf{AConv}_M$ . For the moment, assume  $\kappa_1$  is injective. Then  $\gamma$ , if it exists, is unique. Pick  $z \in Z$ . We know

$$\alpha(z) = h\left(\bigvee_{i=1}^n \lambda_i |\kappa_1(x_i)\rangle \otimes \bigvee_{i=1}^m \sigma_i |\kappa_2(y_i)\rangle\right)$$

for some  $\lambda_i, \sigma_i \in M$ ,  $x_i \in X$  and  $y_i \in Y$ , where  $h$  is the convex structure on  $X + Y$ . As  $(!+) \circ \alpha = ! \circ \kappa_1$ , we must have  $\bigvee_i \sigma_i = 0$  and so  $\alpha(z) = \kappa_1(x_z)$

for some  $x_z \in X$ . This  $x_z$  is unique by supposed injectivity of  $\kappa_1$ . Define  $\gamma(z) = x_z$ . By definition  $\kappa_1 \circ \gamma = \alpha$ . To see  $\gamma$  is affine, pick any  $\varphi \in \mathcal{D}_M(Z)$ . We have  $\kappa_1(\gamma(h_Z(\varphi))) = \alpha(h_Z(\varphi)) = h(\mathcal{D}_M\alpha(\varphi)) = h(\mathcal{D}_M\kappa_1(\mathcal{D}_M\gamma(\varphi))) = \kappa_1(h_X(\mathcal{D}_M\gamma(\varphi)))$  and so indeed  $\gamma(h_Z(\varphi)) = h_X(\mathcal{D}_M\gamma(\varphi))$  — i.e.  $\gamma$  is affine.

Finally, we will show that  $\kappa_1$  is indeed injective. Assume  $\kappa_1(x_0) = \kappa_1(x'_0)$  for some  $x_0, x'_0 \in X$ . Let  $\Phi_1, \dots, \Phi_l \in \mathcal{D}_M^2(X + Y)$  be a derivation of  $\kappa_1(x_0) = \kappa_1(x'_0)$  as in 193 IX. The remainder of the proof is, in essence, a simple induction over  $1 \leq i < l$ , which has been complicated by a technicality. Define

$$\text{IH}(i) \equiv \text{“} \bigotimes_y \mu(\Phi_i)(\kappa_2(y)) = 0 \text{ and } h_X\left(\bigotimes_x \mu(\Phi_i)(\kappa_1(x)) |x\rangle\right) = x_0\text{”}.$$

Clearly IH(1) holds and if IH( $l$ ) should hold, then  $x_0 = x'_0$  as desired. So, to prove the inductive step, assume IH( $i$ ) holds. There are two possible derivation steps, see 193 IX. In the first case (i.e.  $\mu(\Phi_i) = \mu(\Phi_{i+1})$ ) it is obvious IH( $i + 1$ ) holds. So we continue with the other case, where (among others) we have  $\zeta_{ij} = \zeta_{(i+1)j}$ , where  $\Phi_i \equiv \bigotimes_j \zeta_{ij} |\varphi_{ij}\rangle$ . Without loss of generality, we may assume  $\zeta_{ij} \neq 0$ . For each  $j$  there are three possibilities, see 193 IX. The third does not occur: if  $\varphi_{ij_0} = (\mathcal{D}_M\kappa_2)(\chi)$  for some  $\chi \in \mathcal{D}_MY$  and  $j_0$ , then we reach a contradiction:

$$0 = \bigotimes_y \mu(\Phi_i)(\kappa_2(y)) = \bigotimes_{y,j} \zeta_{ij} \odot \varphi_{ij}(\kappa_2(y)) \geq \bigotimes_y \zeta_{ij_0} \odot \chi(y) \neq 0.$$

We are left with two possibilities — to distinguish those, write  $j \in J$  if  $\varphi_{ij} = \varphi_{(i+1)j}$  and  $j \notin J$  if  $\varphi_{ij} = (\mathcal{D}_M\kappa_1)(\chi)$  and  $\varphi_{(i+1)j} = (\mathcal{D}_M\kappa_1)(\chi')$  for some  $\chi, \chi' \in \mathcal{D}_MX$  with  $h_X(\chi) = h_X(\chi')$ .

We will show the first part of IH( $i + 1$ ). Pick any  $y \in Y$ . If  $j \in J$ , then  $\zeta_{(i+1)j} \odot \varphi_{(i+1)j}(\kappa_2(y)) = \zeta_{ij} \odot \varphi_{ij}(\kappa_2(y)) = 0$ . In the other case, if  $j \notin J$ , then clearly  $\zeta_{(i+1)j} \odot \varphi_{(i+1)j}(\kappa_2(y)) = 0$  as  $\varphi_{(i+1)j} = (\mathcal{D}_M\kappa_1)(\chi')$ . So, we have  $\bigotimes_y \mu(\Phi_{i+1})(\kappa_2(y)) = \bigotimes_{y,j} \zeta_{(i+1)j} \odot \varphi_{(i+1)j}(\kappa_2(y)) = 0$ .

Write  $r_{ij} = (\bigotimes_x \varphi_{ij}(\kappa_1(x)))^\perp$ . The remainder of the proof is complicated by the fact that in general  $r_{ij} \neq 0$ . We do have  $\bigotimes_j \zeta_{ij} \odot r_{ij}^\perp = 1$  and  $\bigotimes_j \zeta_{ij} = 1$  so by 178 V we see  $\zeta_{ij} \odot r_{ij}^\perp = \zeta_{ij}$ , which forces  $\zeta_{ij} \odot r_{ij} = 0$ . We compute

$$\begin{aligned} h_X\left(\bigotimes_x \mu(\Phi_i)(\kappa_1(x)) |x\rangle\right) &= h_X\left(\bigotimes_{x,j} \zeta_{ij} \odot \varphi_{ij}(\kappa_1(x)) |x\rangle\right) \\ &= h_X\left(\bigotimes_j \zeta_{ij} \odot r_{ij} |x_0\rangle \otimes \bigotimes_x \zeta_{ij} \odot \varphi_{ij}(\kappa_1(x)) |x\rangle\right) \\ &= h_X\left(\mu\left(\bigotimes_j \zeta_{ij} |r_{ij} |x_0\rangle \otimes \bigotimes_x \varphi_{ij}(\kappa_1(x)) |x\rangle\right)\right) \\ &= h_X\left(\bigotimes_j \zeta_{ij} |h_X(r_{ij} |x_0\rangle \otimes \bigotimes_x \varphi_{ij}(\kappa_1(x)) |x\rangle)\right). \end{aligned}$$

So, if we show that for each  $j$ , we have

$$\begin{aligned} h_X \left( r_{ij} |x_0\rangle \otimes \bigotimes_x \varphi_{ij}(\kappa_1(x)) |x\rangle \right) \\ = h_X \left( r_{(i+1)j} |x_0\rangle \otimes \bigotimes_x \varphi_{(i+1)j}(\kappa_1(x)) |x\rangle \right), \end{aligned} \quad (3.5)$$

then

$$x_0 = h_X \left( \bigotimes_x \mu(\Phi_i)(\kappa_1(x)) |x\rangle \right) = h_X \left( \bigotimes_x \mu(\Phi_{i+1})(\kappa_1(x)) |x\rangle \right),$$

as desired. If  $j \in J$ , then  $\varphi_{ij} = \varphi_{(i+1)j}$  and so  $r_{ij} = r_{(i+1)j}$  and obviously (3.5). So assume  $j \notin J$ . Then clearly  $r_{ij} = 0 = r_{(i+1)j}$  as  $\varphi_{ij} = (\mathcal{D}_M \kappa_1)(\chi)$  and  $\varphi_{(i+1)j} = (\mathcal{D}_M \kappa_1)(\chi')$ . Now (3.5) follows from  $h_X(\chi) = h_X(\chi')$ .  $\square$

**195** It is unclear whether the left square of (3.1) is a pullback in  $\mathbf{AConv}_M$ . We will see that it is, if we assume that  $M$  has a partial division in the following sense.

**II Definition** An **effect divisoid** is an effect monoid  $M$  with partial binary operation  $(a, b) \mapsto b \setminus a$  that is defined iff  $a \leq b$  and satisfies the following axioms.

1.  $b \setminus a$  is the unique element of  $M$  with  $b \setminus a \leq b$  and  $b \odot b \setminus a = a$
2.  $a \leq a \setminus a$
3.  $(a \setminus a) \setminus (a \setminus a) = a \setminus a$

**IIa Beware** The first axiom might be read as “ $b \setminus a$  is defined to be the unique element such that (...)” in which case the definition is cyclical and thus problematic. What is meant instead is that  $b \setminus a$  is part of the structure and satisfies

- $b \setminus a \leq b$  and  $b \odot b \setminus a = a$  and
- if  $c \leq b$  and  $b \odot c = a$ , then  $c = b \setminus a$

for all  $a, b, c \in M$  with  $a \leq b$ . In particular, it is not ruled out that there is an effect monoid which is an effect divisoid in two different ways.

**III Remark** The element  $a \setminus a$  behaves like a support-projection for  $a$ .

**IV Exercise** Show that for an effect divisoid  $M$ , the following holds.

1.  $0 \setminus 0 = 0$ ,  $1 \setminus 1 = 1$ ,  $1 \setminus a = a$ ,  $a \setminus a \odot a \setminus a = a \setminus a$  and  $a \setminus a \odot b = a \setminus a \odot b$ .
2. For any  $a \leq b \leq c$ , we have  $c \setminus b \odot b \setminus a = c \setminus a$ .

**Examples** Almost all effect monoids we encountered are effect divisoids. V

1. The effect monoid  $[0, 1]$  is an effect divisoid with  $b \setminus^a = \frac{a}{b}$  if  $b \neq 0$  and  $0 \setminus^0 = 0$ . With the same partial division, we see the two-element effect monoid  $2$  is also an effect divisoid.
2. More interestingly, if  $M_1$  and  $M_2$  are effect divisoids, then  $M_1 \times M_2$  is an effect divisoid with  $(b_1, b_2) \setminus^{(a_1, a_2)} = (b_1 \setminus^{a_1}, b_2 \setminus^{a_2})$ . In particular  $[0, 1]^n$  is an effect divisoid with component-wise partial division.
3. Later, in 2131, we will see that if  $C$  is an  $\&$ -effectus, then  $(\text{Scal } C)^{\text{op}}$  (the scalars with multiplication in the opposite direction) is an effect divisoid.
4. If  $M$  is a division effect monoid as in [19, dfn. 6.3], then  $M^{\text{op}}$  is an effect divisoid in the obvious way.
5. The effect monoid on the unit interval of  $C[0, 1]$  is *not* an effect divisoid, but the effect monoid on the unit interval of  $L^\infty[0, 1]$  is, see VI.

**Exercise\*** Let  $X$  be a compact Hausdorff space. In this exercise you will show that the unit interval of  $C(X)$  is an effect divisoid if and only if  $X$  is **basically disconnected** [39, 1H] — that is: if  $\overline{\text{supp } f}$  is open for every  $f \in C(X)$ . Equivalently:  $C(X)$  is  $\sigma$ -Dedekind complete [39, 3N.5], which is in turn equivalent to bounded  $\omega$ -completeness (i.e.  $C(X)$  has suprema, resp. infima, of bounded ascending, resp. descending, sequences). VI

1. Suppose the unit interval of  $C(X)$  is an effect divisoid. Let  $f \in C(X)$  with  $0 \leq f \leq 1$ . If  $\overline{\text{supp } f} = X$ , then you're done. Otherwise pick any  $y \notin \overline{\text{supp } f}$ . Show, using Urysohn's lemma, that there is a  $g \in C(X)$ ,  $0 \leq g \leq f \setminus^f$  with  $g(y) = 0$  and  $g(x) = 1$  for all  $x \in \overline{\text{supp } f}$ . Show that this implies  $(f \setminus^f)(y) = 0$  and so  $f \setminus^f$  is the characteristic function of  $\overline{\text{supp } f}$ . Conclude  $X$  is basically disconnected.
2. Assume  $X$  is basically disconnected. Let  $f, g \in C(X)$  with  $0 \leq f \leq g \leq 1$ . For  $n > 0$ , show that  $U_n \equiv \{x; g(x) > \frac{1}{n}\}$  has open closure and so

$$h_n \equiv \begin{cases} \frac{f(x)}{g(x)} & x \in \overline{U_n} \\ 0 & \text{otherwise.} \end{cases}$$

is continuous. Note  $h_1 \leq h_2 \leq \dots \leq 1$  and define  $g \setminus^f = \sup_n h_n$ . Show  $f \setminus^f$  is the characteristic function of  $\overline{\text{supp } f}$  and  $g \setminus^f \leq g \setminus^g$ . Prove that this partial division turns the unit interval of  $C(X)$  into an effect divisoid.

**VII Proposition** If  $a \perp b$  and  $a \otimes b \leq c$  in an effect divisoid, then  $c \setminus a \otimes b = c \setminus a \otimes c \setminus b$ .

**VIII Proof** We will first show that if  $c \odot a \perp c \odot b$ , then  $c \setminus c \odot a \perp c \setminus c \odot b$ . To start, note  $c \odot b^\perp \leq c \odot b^\perp \otimes c^\perp = (c \odot b)^\perp \leq c \odot a$  and so

$$c \setminus c \odot a = c \setminus c \odot b^\perp \odot_{c \odot b^\perp} c \odot a = c \setminus c \odot b^\perp \odot_{c \odot b^\perp} c \odot a \leq c \setminus c \odot b^\perp.$$

Hence  $(c \setminus c \odot b)^\perp \geq c \setminus c \odot b^\perp \geq c \setminus c \odot a = c \setminus c \odot a$ . Indeed:  $c \setminus c \odot a \perp c \setminus c \odot b$ .

**IX** Assume  $a \perp b$  and  $a \otimes b \leq c$ . Clearly  $c \odot c \setminus a = a \perp b = c \odot c \setminus b$ . By our initial lemma, we get  $c \setminus a = c \setminus c \odot c \setminus a \perp c \setminus c \odot c \setminus b = c \setminus b$ . Clearly  $c \odot (c \setminus a \otimes c \setminus b) = a \otimes b$ , so

$$c \setminus a \otimes b = c \setminus c \odot (c \setminus a \otimes c \setminus b) = c \setminus c \odot (c \setminus a \otimes c \setminus b) = c \setminus a \otimes c \setminus b,$$

as desired.  $\square$

**196** The following is a generalization of [19, prop. C.3]. See also [69, prop. 15].

**II Theorem** If  $M$  is an effect divisoid, then  $\mathbf{AConv}_M$  is an effectus.

**III Proof** We only have to show that the square on the left of (3.1) is a pullback in  $\mathbf{AConv}_M$  — the other axioms were proven in 194I.

$$\begin{array}{ccc}
 Z & \xrightarrow{\beta} & X + Y \xrightarrow{\text{id}+!} X + 1 \\
 \alpha \searrow & \dashrightarrow & \downarrow \text{!+id} \quad \downarrow \text{!+id} \\
 & & 1 + Y \xrightarrow{\text{id}+!} 1 + 1
 \end{array}$$

To this end, assume  $\alpha: Z \rightarrow X + 1$  and  $\beta: Z \rightarrow Y + 1$  are maps in  $\mathbf{AConv}_M$  with  $(\text{!+id}) \circ \beta = (\text{id}+!) \circ \alpha$ . We have to show there is a unique  $\gamma: Z \rightarrow X + Y$  in  $\mathbf{AConv}_M$  with  $(\text{id}+!) \circ \gamma = \beta$  and  $(\text{!+id}) \circ \gamma = \alpha$ . Pick any  $z \in Z$ . Write  $\cdot$  for the unique element of 1. By construction of the coproduct, see 193V, we have

$$\begin{aligned}
 \alpha(z) &= h_1 \left( \lambda_0 | \kappa_1(\cdot) \rangle \otimes \bigotimes_{i=1}^n \lambda_i | \kappa_2(y_i) \rangle \right) \\
 \beta(z) &= h_2 \left( \sigma_0 | \kappa_2(\cdot) \rangle \otimes \bigotimes_{i=1}^m \sigma_i | \kappa_1(x_i) \rangle \right)
 \end{aligned}$$

for some  $\lambda_i, \sigma_i \in M$ ,  $x_i \in X$  and  $y_i \in Y$ , where  $h_1$  and  $h_2$  are the convex structures on  $1 + Y$  and  $X + 1$ , respectively. Unfolding definitions (and identifying  $1 + 1 \cong M$  via  $\kappa_1(\cdot) = 1$ ), it is easy to see that by assumption

$$\bigotimes_{i=1}^m \sigma_i = (\text{!+id})(\beta(z)) = (\text{id}+!)(\alpha(z)) = \lambda_0 = \left( \bigotimes_{i=1}^n \lambda_i \right)^\perp. \quad (3.6)$$

This suggests the following definition

$$\gamma(z) = h_3\left(\bigotimes_{i=1}^n \lambda_i |\kappa_2(y_i)\rangle \otimes \bigotimes_{i=1}^m \sigma_i |\kappa_1(x_i)\rangle\right), \quad (3.7)$$

where  $h_3$  is the convex structure on  $X + Y$ . To show this a proper definition, let  $n', m' \in \mathbb{N}$ ,  $\lambda'_i, \sigma'_i \in M$ ,  $x'_i \in X$  and  $y'_i \in Y$  be any elements such that

$$\begin{aligned} \alpha(z) &= h_1\left(\lambda'_0 |\kappa_1(\cdot)\rangle \otimes \bigotimes_{i=1}^{n'} \lambda'_i |\kappa_2(y'_i)\rangle\right) \\ \beta(z) &= h_2\left(\sigma'_0 |\kappa_2(\cdot)\rangle \otimes \bigotimes_{i=1}^{m'} \sigma'_i |\kappa_1(x'_i)\rangle\right). \end{aligned}$$

We have to show

IV

$$\begin{aligned} h_3\left(\bigotimes_{i=1}^{n'} \lambda'_i |\kappa_2(y'_i)\rangle \otimes \bigotimes_{i=1}^{m'} \sigma'_i |\kappa_1(x'_i)\rangle\right) \\ = h_3\left(\bigotimes_{i=1}^n \lambda_i |\kappa_2(y_i)\rangle \otimes \bigotimes_{i=1}^m \sigma_i |\kappa_1(x_i)\rangle\right). \end{aligned} \quad (3.8)$$

Without loss of generality, we may assume  $n, n', m, m' \geq 1$  (as we allow, for instance,  $\lambda_1 = 0$ ). Due to (3.6), it is clear  $\lambda_0 = \lambda'_0$  and  $\sigma_0 = \sigma'_0$ . By 193IX we know there are derivations  $\Phi_1, \dots, \Phi_l \in \mathcal{D}_M^2(1 + Y)$  and  $\Psi_1, \dots, \Psi_k \in \mathcal{D}_M^2(X + 1)$ , say  $\Phi_i \equiv \bigotimes_j \zeta_{ij} |\varphi_{ij}\rangle$ ,  $\Psi_i \equiv \bigotimes_j \xi_{ij} |\psi_{ij}\rangle$  with

$$\begin{aligned} \Phi_1 &= |\lambda_0 |\kappa_1(\cdot)\rangle \otimes \bigotimes_{i=1}^n \lambda_i |\kappa_2(y_i)\rangle\rangle & \Phi_l &= |\lambda_0 |\kappa_1(\cdot)\rangle \otimes \bigotimes_{i=1}^{n'} \lambda'_i |\kappa_2(y'_i)\rangle\rangle \\ \Psi_1 &= |\sigma_0 |\kappa_2(\cdot)\rangle \otimes \bigotimes_{i=1}^m \sigma_i |\kappa_1(x_i)\rangle\rangle & \Psi_k &= |\sigma_0 |\kappa_2(\cdot)\rangle \otimes \bigotimes_{i=1}^{m'} \sigma'_i |\kappa_1(x'_i)\rangle\rangle. \end{aligned}$$

We are going to combine the  $\Phi_i$  and  $\Psi_i$  into a single derivation of (3.8) by first ‘applying’ the  $\Phi_i$  and then the  $\Psi_i$ . Define  $\Omega_1, \dots, \Omega_{l+k} \in \mathcal{D}_M^2(X + Y)$  by

$$\begin{aligned} \Omega_i &= \bigotimes_j \zeta_{ij} |\omega_{ij}\rangle & \text{for } 1 \leq i \leq l \\ \Omega_{i+l} &= \bigotimes_j \xi_{ij} |\omega_{(i+l)j}\rangle & \text{for } 1 \leq i \leq k, \end{aligned}$$

where  $\omega_{ij} \in \mathcal{D}_M(X + Y)$  are given by

$$\begin{aligned}
\omega_{ij}(\kappa_2(y)) &= \varphi_{ij}(\kappa_2(y)) \\
\omega_{ij}(\kappa_1(x)) &= \varphi_{ij}(\kappa_1(\cdot)) \odot (\lambda_0 \setminus \psi_{11}(\kappa_1(x)) \otimes r(\kappa_1(x))) \\
\omega_{(i+l)j}(\kappa_1(x)) &= \psi_{ij}(\kappa_1(x)) \\
\omega_{(i+l)j}(\kappa_2(y)) &= \psi_{ij}(\kappa_2(\cdot)) \odot (\sigma_0 \setminus \varphi_{11}(\kappa_2(y)) \otimes r(\kappa_2(y))) \\
r(z) &= \begin{cases} (\lambda_0 \setminus \lambda_0)^\perp & z = \kappa_1(x_1) \\ (\sigma_0 \setminus \sigma_0)^\perp & z = \kappa_2(y_1) \\ 0 & \text{otherwise,} \end{cases}
\end{aligned}$$

where on the first two lines  $1 \leq i \leq l$  and on the second two  $1 \leq i \leq k$ . Before we continue, we check whether the  $\omega_{ij}$  are distributions, as claimed. For  $1 \leq i \leq l$  this amounts to  $\bigvee_x \omega_{ij}(\kappa_1(x)) = \varphi_{ij}(\kappa_1(\cdot))$ . Indeed

$$\begin{aligned}
\bigvee_x \omega_{ij}(\kappa_1(x)) &= \varphi_{ij}(\kappa_1(\cdot)) \odot r(\kappa_1(x_1)) \otimes \bigvee_x \varphi_{ij}(\kappa_1(\cdot)) \odot \lambda_0 \setminus \psi_{11}(\kappa_1(x)) \\
&= \varphi_{ij}(\kappa_1(\cdot)) \odot (\lambda_0 \setminus \lambda_0)^\perp \otimes \varphi_{ij}(\kappa_1(\cdot)) \odot \lambda_0 \setminus \bigvee_x \psi_{11}(\kappa_1(x)) \\
&= \varphi_{ij}(\kappa_1(\cdot)) \odot (\lambda_0 \setminus \lambda_0)^\perp \otimes \varphi_{ij}(\kappa_1(\cdot)) \odot \lambda_0 \setminus \lambda_0 \\
&= \varphi_{ij}(\kappa_1(\cdot)).
\end{aligned}$$

With an analogous argument, one checks  $\omega_{(i+l)j}$  is a distribution for  $1 \leq i \leq k$ .

To show  $\Omega_i$  is a derivation (in the sense of 193IX), first pick any  $1 \leq i < l$ . We distinguish between the two allowable derivation steps. If  $\mu(\Phi_i) = \mu(\Phi_{i+1})$ , then  $\mu(\Omega_i) = \mu(\Omega_{i+1})$  by a straightforward computation and so  $\Omega_i$  has a valid step. In the other case, we have  $\zeta_{ij} = \zeta_{(i+1)j}$  and for each  $j$  we have three possibilities. In the first case, if  $\varphi_{ij} = \varphi_{(i+1)j}$ , then clearly  $\omega_{ij} = \omega_{(i+1)j}$  as well. For the second case, assume  $\varphi_{ij} = (\mathcal{D}_M \kappa_2)(\chi)$  and  $\varphi_{(i+1)j} = (\mathcal{D}_M \kappa_2)(\chi')$  for some  $\chi, \chi' \in \mathcal{D}_M Y$  with  $h_Y(\chi) = h_Y(\chi')$ . Clearly  $\varphi_{ij}(\kappa_1(\cdot)) = 0$  and so  $\omega_{ij} = (\mathcal{D}_M \kappa_2)(\chi)$ . Similarly  $\omega_{(i+1)j} = (\mathcal{D}_M \kappa_2)(\chi')$ . The third case is trivial. So  $\Omega_i$  also makes a valid step of the second kind.

With the same kind of argument, we cover  $\Omega_{i+l}$  for  $1 \leq i < k$ . The only step left to check is the one from  $\Omega_l$  to  $\Omega_{l+1}$ . Note  $\zeta_{l1} = 1$ ,  $\varphi_{l1}(\kappa_1(\cdot)) = \lambda_0$  and  $\lambda_0 \odot (\lambda_0 \setminus \lambda_0)^\perp = 0$ , so with some easy manipulation, we find

$$\Omega_l = \left| \bigvee_{i=1}^{n'} \lambda'_i \mid \kappa_2(y'_i) \right\rangle \otimes \left| \bigvee_{i=1}^m \sigma_i \mid \kappa_1(x_i) \right\rangle.$$

and in a similar fashion

$$\begin{aligned}\Omega_{l+1} &= \left| \bigotimes_{i=1}^{n'} \lambda'_i | \kappa_2(y'_i) \rangle \right\rangle \otimes \left| \bigotimes_{i=1}^m \sigma_i | \kappa_1(x_i) \rangle \right\rangle \\ \Omega_1 &= \left| \bigotimes_{i=1}^n \lambda_i | \kappa_2(y_i) \rangle \right\rangle \otimes \left| \bigotimes_{i=1}^m \sigma_i | \kappa_1(x_i) \rangle \right\rangle \\ \Omega_{l+k} &= \left| \bigotimes_{i=1}^{n'} \lambda'_i | \kappa_2(y'_i) \rangle \right\rangle \otimes \left| \bigotimes_{i=1}^{m'} \sigma'_i | \kappa_1(x'_i) \rangle \right\rangle.\end{aligned}$$

So  $\Omega_i$  is a valid derivation of (3.8).

We are in the home stretch now. Define  $\gamma(z)$  as in (3.7). It is easy to see  $\gamma$  is the unique map with  $(\text{id}+!) \circ \gamma = \beta$  and  $(! + \text{id}) \circ \gamma = \alpha$ . It remains to be shown  $\gamma$  is affine. Write V

$$q_1: \mathcal{D}_M(X+Y) \rightarrow X+Y \quad q_2: \mathcal{D}_M(1+Y) \rightarrow 1+Y \quad q_3: \mathcal{D}_M(X+1) \rightarrow X+1$$

for the quotient maps used in the construction of the coproducts. We worked hard to show that for all  $z \in Z$ ,  $\varphi \in \mathcal{D}_M Y$ ,  $\psi \in \mathcal{D}_M X$ , and  $\chi \in \mathcal{D}_M(X+Y)$  with  $\chi(\kappa_1(y)) = \varphi(\kappa_1(y))$  and  $\chi(\kappa_2(x)) = \psi(\kappa_2(x))$ , we have

$$\left. \begin{aligned} \alpha(z) &= q_1(\varphi) \\ \beta(z) &= q_2(\psi) \end{aligned} \right] \implies \gamma(z) = q_3(\chi).$$

Assume  $z = h_Z(\bigotimes_i \lambda_i | z_i \rangle)$ . We want to show  $\gamma(z) = h_3(\bigotimes_i \lambda_i | \gamma(z_i) \rangle)$ . To this end, pick  $\varphi_i, \psi_i, \chi_i$  with  $\alpha(z_i) = q_1(\varphi_i)$ ,  $\beta(z_i) = q_2(\psi_i)$ ,  $\chi_i(\kappa_1(y)) = \varphi_i(\kappa_1(y))$  and  $\chi_i(\kappa_2(x)) = \psi_i(\kappa_2(x))$ . So  $\gamma(z_i) = q_3(\chi_i)$ . Define  $\varphi, \psi, \chi$  as follows.

$$\varphi \equiv \mu\left(\bigotimes_i \lambda_i | \varphi_i \rangle\right) \quad \psi \equiv \mu\left(\bigotimes_i \lambda_i | \psi_i \rangle\right) \quad \chi \equiv \mu\left(\bigotimes_i \lambda_i | \chi_i \rangle\right)$$

Note  $q_1(\varphi) = h_1(\bigotimes_i \lambda_i | q_1(\varphi_i) \rangle) = h_1(\bigotimes_i \lambda_i | \alpha(z_i) \rangle) = \alpha(\bigotimes_i \lambda_i | z_i \rangle) = \alpha(z)$ . Similarly  $q_2(\psi) = \beta(z)$ . For any  $y \in Y$ , we have  $\chi(\kappa_1(y)) = \bigotimes_i \lambda_i \odot \chi_i(\kappa_1(y)) = \bigotimes_i \lambda_i \odot \varphi_i(\kappa_1(y)) = \varphi(\kappa_1(y))$ . Similarly  $\chi(\kappa_2(x)) = \psi(\kappa_2(x))$  for any  $x \in X$ . So  $\gamma(z) = q_3(\chi)$  and consequently

$$\gamma(z) = q_3(\chi) = h_3\left(\bigotimes_i \lambda_i | q_3(\chi_i) \rangle\right) = h_3\left(\bigotimes_i \lambda_i | \gamma(z_i) \rangle\right),$$

as desired. □

### 3.3 Effectuses with quotients

We return to the main program of this chapter: the attempted axiomatisation of the effectus  $\mathbf{vN}^{\text{op}}$ . 197

**II Definition** Let  $C$  be an effectus. We say  $C$  is an **effectus with quotients** if for each predicate  $p: X \rightarrow 1$ , there exists a map  $\xi_p: X \rightarrow X/p$  with  $1 \circ \xi_p \leq p^\perp$  satisfying the following universal property.

For any map  $f: X \rightarrow Y$  with  $1 \circ f \leq p^\perp$ , there is a unique map  $f': X/p \rightarrow Y$  such that  $f' \circ \xi_p = f$ .

Any map with this universal property is called a **quotient** for  $p$ .

**IIa Remarks** The universal property of quotients is essentially the same as the universal property of (contractive) *compressions*, which we proposed in [126]. Effectuses with quotients also appear in [23].

**III Notation** Unless otherwise specified  $\xi_p$  will denote some quotient for  $p$ .

**IV Examples** In  $\mathbf{vN}^{\text{op}}$  quotients are exactly the same thing as contractive filters, see 169 VIII, 96 I and 96 V. An example of a quotient map for  $1 - a \in \mathcal{A}$  is given by  $\xi: [a] \mathcal{A} [a] \rightarrow \mathcal{A}$  with  $\xi(b) = \sqrt{ab}\sqrt{a}$ .

**IVa** The effectus  $\mathbf{EJA}^{\text{op}}$  has quotients, which are in essence very similar to those of  $\mathbf{vN}^{\text{op}}$ , when ignoring the technical complications Jordan algebras impose. [128, prop. 25]

**IVb** The effectus  $\mathbf{OUS}^{\text{op}}$  has quotients, which are rather different from the previous two examples. An example of a quotient for a predicate  $v$  on an order unit space  $V$  is given by the inclusion of the order ideal generated by  $1 - v$

$$\langle 1 - v \rangle \equiv \{a; a \in V; \exists n. -n(1 - v) \leq a \leq n(1 - v)\} \subseteq V,$$

where the order ideal is considered as an order unit space with order-unit  $1 - v$ . The effectus  $\mathbf{OUG}^{\text{op}}$  has very similar quotients. See [23].

**V Exercise** Verify the following basic properties of quotients. (See [23].)

1. If  $\xi: X \rightarrow Y$  is a quotient for  $p$  and  $\vartheta: Y \rightarrow Z$  is an isomorphism, then  $\vartheta \circ \xi$  is a quotient for  $p$  as well.
2. Conversely, if  $\xi_1$  and  $\xi_2$  are both quotients for  $p$ , then there is a unique isomorphism  $\vartheta$  with  $\xi_1 = \vartheta \circ \xi_2$ .
3. Isomorphisms are quotients (for 0).
4. Maps into 0 are quotients (for 1).
5. If  $\xi$  is a quotient for  $p$ , then  $1 \circ \xi = p^\perp$  (Hint: apply the universal property to  $p^\perp$ ).
6. Quotients are epic.

**VI** The following proposition is easy to prove, but shows an important property of quotients: any map  $f$  factors as a total map after a quotient for  $(1 \circ f)^\perp$ .

**Proposition** Assume  $\xi_{p^\perp} : X \rightarrow X/p^\perp$  is a quotient for  $p^\perp$ . For any  $f : X \rightarrow Z$  VII  
with  $1 \circ f = p$ , there is a unique *total*  $g : X/p^\perp \rightarrow Z$  with  $f = g \circ \xi_{p^\perp}$ .

**Proof** (This is a simplified version of our previous proof in [23].) By definition VIII  
of quotient, there is a unique  $g : X/p^\perp \rightarrow Z$  with  $g \circ \xi_{p^\perp} = f$ . Note  $1 \circ g \circ \xi_{p^\perp} = 1 \circ f = p = 1 \circ \xi_{p^\perp}$ . Thus, as  $\xi_{p^\perp}$  is epi (V), we conclude  $1 \circ g = 1$ . That is:  $g$  is total.  $\square$

**Proposition** In an effectus with quotients, quotients are closed under composition. IX

**Proof** (This is essentially the same proof as we gave in [23].) Assume  $\xi_1 : X \rightarrow Y$  X  
is a quotient for  $p^\perp$  and  $\xi_2 : Y \rightarrow Z$  is a quotient for  $q^\perp$ . We will prove  $\xi_2 \circ \xi_1$  is a quotient for  $(q \circ \xi_1)^\perp$ . As our effectus has quotients, we can pick a quotient  $\xi : X \rightarrow X/(q \circ \xi_1)^\perp$  of  $(q \circ \xi_1)^\perp$ . First, some preparation. Note  $1 \circ \xi = q \circ \xi_1 \leq 1 \circ \xi_1 = p$ . Thus by the universal property of  $\xi_1$ , there is a unique map  $h_1 : Y \rightarrow X/(q \circ \xi_1)^\perp$  with  $h_1 \circ \xi_1 = \xi$ . As  $1 \circ h_1 \circ \xi_1 = 1 \circ \xi = q \circ \xi_1$  and  $\xi_1$  is epi, we see  $1 \circ h_1 = q$ . Thus by VII, there is a unique total map  $h_2 : Z \rightarrow X/(q \circ \xi_1)^\perp$  with  $h_2 \circ \xi_2 = h_1$ . Let  $g : X/(q \circ \xi_1)^\perp \rightarrow Z$  be the unique map such that  $g \circ \xi = \xi_2 \circ \xi_1$ . We are in the following situation.

$$\begin{array}{ccccc}
 X & \xrightarrow{\xi_1} & Y & \xrightarrow{\xi_2} & Z \\
 & \searrow \xi & & \swarrow h_1 & \uparrow g \\
 & & & & X/(q \circ \xi_1)^\perp \\
 & & & & \downarrow h_2
 \end{array}$$

We claim  $g$  and  $h_2$  are each other's inverse. Indeed: from  $g \circ h_2 \circ \xi_2 \circ \xi_1 = g \circ \xi = \xi_2 \circ \xi_1$  we get  $g \circ h_2 = \text{id}$  and from  $h_2 \circ g \circ \xi = h_2 \circ \xi_2 \circ \xi_1 = \xi$  we find  $h_2 \circ g = \text{id}$ . Thus  $\xi_2 \circ \xi_1 = g \circ \xi$  for an isomorphism  $g$ , which shows  $\xi_2 \circ \xi_1$  is a quotient, see V.  $\square$

**Exercise** By VII we know that each map  $f$  in an effectus with quotients factors XI  
as  $t \circ \xi$  for some total map  $t$  and quotient  $\xi$ . Show that this forms an orthogonal factorization system (cf. [79]) — that is: prove that if  $t' \circ \xi' = t \circ \xi$  for some quotient  $\xi'$  and total map  $t'$ , then there is a unique isomorphism  $\vartheta$  with  $\xi' = \vartheta \circ \xi$  and  $t = t' \circ \vartheta$ .

In 197XI we saw that every map  $f$  factors uniquely via a quotient for  $(1 \circ f)^\perp$ . 198  
In this sense quotients are the universal way to restrict to the support on the domain — so why do we call them quotients instead of, say, initial-support-maps? We answer this question in IV, but need some preparation first.

**Definition** For an effectus  $C$ , write  $\int \text{Pred} \square$  for the category with II

1. as objects pairs  $(X, p)$ , where  $X$  is an object of  $C$  and  $p \in \text{Pred } X$  and

2. an morphism between  $(X, p)$  and  $(Y, q)$  corresponds to a partial map  $f: X \rightarrow Y$  in  $C$  for which we have  $p \leq (q^\perp \circ f)^\perp$ .

Write  $U: \int \text{Pred}_\square \rightarrow C$  for the forgetful functor — that is:  $U(X, p) = X$  and  $Uf = f$  for arrows  $f$ . The functor  $U$  has left- and right-adjoint  $0 \dashv U \dashv 1$  with  $0, 1: C \rightarrow \int \text{Pred}_\square$  given by  $0X = (X, 0)$ ,  $1X = (X, 1)$  and  $0f = 1f = f$ .

III **Exercise\*** Show that for an effectus  $C$ , the following are equivalent.

1.  $C$  has quotients.
2. The functor  $0: C \rightarrow \int \text{Pred}_\square$ , has a left-adjoint  $Q: \int \text{Pred}_\square \rightarrow C$ .

IV If the existence of a left adjoint  $Q$  to  $0$  gives quotients, what does the existence of a right adjoint  $K$  to  $1$  amount to? It will turn out (199 VI) this is equivalent to the existence of so-called *comprehensions*, the topic of the next section. With both quotients and comprehension, we have a chain of four adjunctions:

$$\begin{array}{ccc}
 & \int \text{Pred}_\square & \\
 Q \swarrow & \left( \begin{array}{c} \dashv \dashv \dashv \dashv \\ \downarrow \\ \dashv \dashv \dashv \dashv \end{array} \right) & \searrow K \\
 & C & 
 \end{array}$$

There are numerous other examples of categories which also have such a chain of four adjunctions between it and a natural category of object-with-predicates. See [22] for several examples. In most of these examples, the predicates correspond to subspaces and the left-most functor to quotienting by this subspace. This is the reason for the name ‘effectus with quotients’.

V Originally, we considered the universal properties of quotients and comprehensions separately and under different names. It was Jacobs who recognized that both universal properties appear together in a chain of adjunctions.

### 3.4 Effectuses with comprehension and images

199 We continue with the formal introduction of our second axiom: the existence of *comprehension*.

II **Definition** Let  $C$  be an effectus. We say  $C$  has *comprehension* if for each predicate  $p: X \rightarrow 1$ , there exists a map  $\pi_p: \{X|p\} \rightarrow X$  with  $p \circ \pi_p = 1 \circ \pi_p$  satisfying the following universal property.

For any map  $g: Z \rightarrow X$  with  $p \circ g = 1 \circ g$ , there is a unique map  $g': Z \rightarrow \{X|p\}$  such that  $\pi_p \circ g' = g$ .

Any map with this universal property is called a **comprehension** for  $p$ .

**Beware** We do not assume comprehensions are total (in contrast to [23].) III  
 In 202 VIII we will see that in an effectus with quotients, comprehensions must be total.

**Remarks** The universal property of comprehensions is essentially the same as IIIa  
 the universal property of (contractive) *corners*, which we proposed in [126].  
 Effectuses with compression also appear in [23]. In partial form, comprehensions  
 are essentially the same as categorical kernels, see 200 V later on.

**Notation** Unless otherwise specified  $\pi_p$  will denote some comprehension of  $p$ . IV

**Examples** In  $\mathbf{vN}^{\text{op}}$  comprehensions are exactly the same thing as corners, see V  
 169 II or 95 I. An example of a comprehension for  $a \in \mathcal{A}$  is given by  $\pi: \mathcal{A} \rightarrow$   
 $[a] \mathcal{A} [a]$  with  $\pi(b) = [a] b [a]$  as proven in 95 II.

The effectus  $\mathbf{EJA}^{\text{op}}$  also has comprehension, which is very similar to those Va  
 of  $\mathbf{vN}^{\text{op}}$ . [128, prop. 24]

The comprehension maps of  $\mathbf{OUS}^{\text{op}}$  are again quite different from those  
 of  $\mathbf{vN}^{\text{op}}$  and  $\mathbf{EJA}^{\text{op}}$ . Confusingly, a comprehension for a predicate  $v$  of an order  
 unit space  $V$  is given by the vector-space-quotient map  $q: V \rightarrow V/\langle 1-v \rangle$ ,  
 where  $\langle 1-v \rangle$  is the order ideal generated by  $1-v$  defined in 197 IVb. The  
 effectus  $\mathbf{OUG}^{\text{op}}$  has similar comprehension. See [23].

Any extensive category with final object has comprehension, which includes **Set**,  
**CH** and **CRng**, see [23].

**Exercise\*** Show that an effectus  $C$  has comprehension if and only if the func- VI  
 tor  $1: C \rightarrow \int \text{Pred}_{\square}$  from 198 II has a right adjoint  $K$ .

**Exercise** Show the following basic properties of comprehensions. (See [23].) VII

1. If  $\pi: X \rightarrow Y$  is a comprehension for  $p$  and  $\vartheta: Z \rightarrow X$  is an isomorphism,  
 then  $\pi \circ \vartheta$  is a comprehension for  $p$  as well.
2. Conversely, if  $\pi_1$  and  $\pi_2$  are both comprehensions for  $p$ , then there is a  
 unique isomorphism  $\vartheta$  with  $\pi_1 = \pi_2 \circ \vartheta$ .
3. Isomorphisms are comprehensions (for 1).
4. Zero maps are comprehensions (for 0).
5. Comprehensions are monic.
6.  $p^{\perp} \circ \pi = 0$  if  $\pi$  is a comprehension for  $p$ .

Comprehensions and categorical kernels are very similar.

200

- II **Definition** Let  $C$  be a category with a zero object. A (categorical) kernel of an arrow  $f$  (in symbols:  $\ker f$ ) is an equalizer of  $f$  with the parallel zero map. (That is:  $f \circ (\ker f) = 0$  and for every  $g$  with  $f \circ g = 0$ , there exists a unique  $g'$  with  $(\ker f) \circ g' = g$ .) We will use  $\text{cok } f$  to denote a cokernel of  $f$  — that is: a kernel of  $f$  in  $C^{\text{op}}$ .
- III **Proposition** An effectus with comprehension has all kernels. The kernel of a map  $f$  is given by a comprehension  $\pi_{(1 \circ f)^\perp}$  for  $(1 \circ f)^\perp$ .
- IV **Proof** Clearly  $1 \circ f \circ \pi_{(1 \circ f)^\perp} = 0$  and so  $f \circ \pi_{(1 \circ f)^\perp} = 0$ . Assume  $g$  is any map with  $f \circ g = 0$ . Then  $1 \circ f \circ g = 0$  and so  $(1 \circ f)^\perp \circ g = 1 \circ g$ . Thus there exists a unique  $g'$  with  $\pi_{(1 \circ f)^\perp} \circ g' = g$ .  $\square$
- V **Exercise** Show that in an effectus, a map  $f$  is a comprehension for  $p$  if and only if it is a kernel of  $p^\perp$ .

201 We are ready to define purity in effectuses.

- II **Definition** In an effectus a map  $f$  is called pure if  $f = \pi \circ \xi$  for some comprehension  $\pi$  and quotient  $\xi$ .
- III **Example** Due to 100 III pure maps in  $\text{vN}^{\text{op}}$  are precisely the pure maps as defined in 170 I and 100 I.  
The pure maps  $\mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  are exactly the maps of the form  $\text{ad}_T$  where  $T$  is a contractive map  $\mathcal{H} \rightarrow \mathcal{H}$ .
- IV **Remark** In the previous chapter we discussed (in 168 III) why some more familiar notions of purity for states, like extremality, do not generalize properly to arbitrary maps. Recently, three other definitions of purity have been proposed in different contexts. [16, 26, 101] All three are inspired by the essential uniqueness of purification (see 139 XI) and require a monoidal structure to state. The identity map might not be pure in the sense of [16], which is too restrictive for our taste. Hazarding a guess, it seems likely that [26] considers maps pure which we don't and that [101] does not contain all our pure maps. The exact relation between our definition of pure and [26, 101] is left open.
- V We do not have a good handle on the structure of pure maps in an arbitrary effectus. We will need a few additional assumptions.

202 **Definition** Let  $C$  be an effectus.

1. We say  $C$  has images if for each map  $f: X \rightarrow Y$ , there is a least predicate  $\text{im } f$  on  $Y$  with the property  $(\text{im } f) \circ f = 1 \circ f$  — that is, for every predicate  $p$  on  $Y$  with  $p \circ f = 1 \circ f$ , we must have  $\text{im } f \leq p$ . For brevity,

write  $\text{im}^\perp f \equiv (\text{im } f)^\perp$ . Note  $\text{im}^\perp f$  is the greatest predicate with the property  $(\text{im}^\perp f) \circ f = 0$ .

2. We say a map  $f: X \rightarrow Y$  is **faithful** if  $\text{im } f = 1$ . That is:  $f$  is faithful if and only if  $p \circ f = 0$  implies  $p = 0$  for every predicate  $p$ .

**Notation** The expression  $\text{im } f \circ g$  is read as  $\text{im}(f \circ g)$ . II

**Remarks** The predicate  $\text{im } f$  will play for comprehension the same role as  $1 \circ f$  plays for quotients. Similarly faithful is the analogue of total. Images in effectuses are also studied in [23]. III

**Examples** In  $\text{vN}^{\text{op}}$  the image of a map  $f: \mathcal{A} \rightarrow \mathcal{B}$  is given by the least projection  $e \in \mathcal{A}$  with the property  $f(1 - e) = 0$ . The map  $f$  is faithful if and only if  $f(a^*a) = 0$  implies  $a^*a = 0$  for all  $a \in \mathcal{A}$ . IV

The effectus  $\text{EJA}^{\text{op}}$  also has all images, which are defined similarly, see [128, prop. 29]. In contrast, the effectuses  $\text{OUG}^{\text{op}}$  and  $\text{OUS}^{\text{op}}$  do not have all images, see [23]. IVa

**Exercise** Show  $\text{im } f \circ g \leq \text{im } f$ . Conclude  $\text{im } f \circ \alpha = \text{im } f$  for any iso  $\alpha$ . V

**Exercise** Show that in an effectus, any quotient is faithful. VI

In contrast to quotients, it is not clear at all whether (without additional assumptions) comprehensions are total; they are part of a factorization system or are closed under composition. VII

**Lemma** In an effectus with quotients, comprehensions are total. VIII

**Proof** Assume  $\pi$  is some comprehension for  $p$ . Let  $\xi$  be a quotient for  $(1 \circ \pi)^\perp$ . There is a total  $\pi_t$  with  $\pi = \pi_t \circ \xi$ . As  $1 \circ \pi_t \circ \xi = 1 \circ \pi = p \circ \pi = p \circ \pi_t \circ \xi$  and  $\xi$  is epi, we see  $1 \circ \pi_t = p \circ \pi_t$ . Thus, as  $\pi$  is a comprehension for  $p$ , there exists an  $f$  with  $\pi_t = \pi \circ f$ . Now  $\pi = \pi_t \circ \xi = \pi \circ f \circ \xi$  and so  $\text{id} = f \circ \xi$ , since  $\pi$  is mono. Hence  $1 = 1 \circ \text{id} = 1 \circ f \circ \xi \leq 1 \circ \xi = 1 \circ \pi$  and so  $\pi$  is total. IX  $\square$

### 3.4.1 Sharp predicates

**Definition** Let  $C$  be an effectus with comprehension and images. 203

1. We say a predicate  $p$  is (image) **sharp** if  $p = \text{im } f$  for some map  $f$ . Write  $\text{SPred } X$  for the set of all sharp predicates on  $X$ .
2. We define  $[p] = \text{im } \pi_p$ , where  $\pi_p$  is some comprehension for  $p$ . (Comprehensions for the same predicate have the same image by 202V.) Also write  $[p] = [p^\perp]^\perp$ .

- II **Beware** In [23]  $p$  is called sharp whenever  $p \wedge p^\perp = 0$ . In general this is weaker than image-sharpness, which we use in this text. In IV we will see  $[p]$  is sharp. It is unclear whether  $[p]$  is sharp (without additional assumptions).
- III **Example** In  $\mathbf{vN}^{\text{op}}$  the sharp predicates are the projections. For a predicate  $a \in \mathcal{A}$ , the sharp predicate  $[a]$  is the least projection above  $a$ .
- IV **Lemma** In an effectus with comprehension and images, we have

1.  $[p] \leq p$
2.  $\pi_p = \pi_{[p]} \circ \alpha$  for some iso  $\alpha$ ;
3.  $[[p]] = [p]$ ;
4.  $p \leq q \implies [p] \leq [q]$ ;
5.  $[p] \circ f \leq [p \circ f]$  and
6.  $[p] \circ f = 0$  iff  $p \circ f = 0$ ,

for any map  $f: X \rightarrow Y$ , predicates  $p, q$  on  $X$  and  $\pi_p$  and  $\pi_{[p]}$  comprehensions for  $p$  and  $[p]$  respectively.

- V **Proof** We will prove the statements in listed order.
- VI (Ad 1) Let  $\pi$  be a comprehension for  $p$ . By definition  $p \circ \pi = 1 \circ \pi$ . Thus  $[p] = \text{im } \pi \leq p$ , as desired.
- VII (Ad 2) It is sufficient to show  $\pi_p$  is a comprehension for  $[p]$ . First, note  $[p] \circ \pi_p = (\text{im } \pi_p) \circ \pi_p = 1 \circ \pi_p$ . To show the universal property, assume  $g: Z \rightarrow X$  is some map with  $[p] \circ g = 1 \circ g$ . Then  $1 \circ g = [p] \circ g \leq p \circ g \leq 1 \circ g$  and so  $1 \circ g = p \circ g$ . As  $\pi_p$  is a comprehension for  $p$ , there is a unique  $g'$  with  $\pi_p \circ g' = g$  and so  $\pi_p$  is indeed a comprehension for  $[p]$  as well.
- VIII (Ad 3) Follows from the previous point and 202V.
- IX (Ad 4) Pick a comprehension  $\pi_p$  for  $p$  and  $\pi_q$  for  $q$ . Note  $1 \circ \pi_p = p \circ \pi_p \leq q \circ \pi_p \leq 1 \circ \pi_p$  so  $q \circ \pi_p = 1 \circ \pi_p$  and thus  $\pi_p = \pi_q \circ f$  for some  $f$ . By 202V we see  $[p] = \text{im } \pi_p = \text{im } \pi_q \circ f \leq \text{im } \pi_q = [q]$ .
- X (Ad 5) Clearly  $p \circ f \circ \pi_{(p \circ f)^\perp} = 0$ . (Recall our convention 199IV for  $\pi$ 's.) Thus there is some  $h$  with  $f \circ \pi_{(p \circ f)^\perp} = \pi_{p^\perp} \circ h$ . By point 2 there is some (isomorphism)  $\alpha$  with  $\pi_{p^\perp} = \pi_{[p^\perp]} \circ \alpha$ . We compute

$$[p] \circ f \circ \pi_{(p \circ f)^\perp} = [p] \circ \pi_{p^\perp} \circ h = [p] \circ \pi_{[p^\perp]} \circ \alpha \circ h = [p] \circ \pi_{[p]^\perp} \circ \alpha \circ h = 0.$$

Thus  $[p] \circ f \leq \text{im}^\perp \pi_{(p \circ f)^\perp} = [(p \circ f)^\perp]^\perp = [p \circ f]$ , as promised.

- XI (Ad 6) Assume  $[p] \circ f = 0$ . From  $p \leq [p]$  it follows  $p \circ f \leq [p] \circ f = 0$ . Thus  $p \circ f = 0$ . For the converse, assume  $p \circ f = 0$ . Then  $[p \circ f] = [0] = (\text{im id})^\perp = 0$  as  $\text{id}$  is a comprehension for  $1$ . Thus  $[p] \circ f \leq [p \circ f] = 0$ .  $\square$
- XII **Exercise** Let  $C$  be an effectus with comprehension and images. Show that  $p$  is sharp if and only if  $[p] = p$ . Conclude  $\text{im } \pi_s = s$  for sharp  $s$ .
- XIII **Exercise** Show that in an effectus with comprehension and images we have the equality  $[[p] \circ f] = [p \circ f]$  for any map  $f: X \rightarrow Y$  and predicate  $p$  on  $X$ .

**Exercise** Show that in an effectus with images  $\text{im} \langle f, g \rangle = [\text{im } f, \text{im } g]$ . Conclude that a predicate  $[p, q]$  is sharp if and only if  $p$  and  $q$  are sharp. XIV

**Lemma** In an effectus with comprehension and images we have 204

$$s \leq t \iff \pi_s = \pi_t \circ h \text{ for some } h,$$

for all sharp predicates  $s, t$  on the same object.

**Proof** (This simpler proof than ours in [23].) Assume  $\pi_s = \pi_t \circ h$ . Then II

$$s \stackrel{203\text{XII}}{=} [s] = \text{im } \pi_s = \text{im } \pi_t \circ h \stackrel{202\text{V}}{\leq} \text{im } \pi_t = [t] \stackrel{203\text{XII}}{=} t.$$

as desired. Conversely assume  $s \leq t$ . Then  $t^\perp \circ \pi_s \leq s^\perp \circ \pi_s = 0$  and so  $\pi_s = \pi_t \circ h$  for some  $h$  by the universal property of  $\pi_t$ . □

**Lemma** In an effectus with images, we have  $\text{im}[f, g] = (\text{im } f) \vee (\text{im } g)$ . III

**Proof** We get  $\text{im}[f, g] \geq \text{im } f$  and  $\text{im}[f, g] \geq \text{im } g$  from IV

$$\begin{aligned} [1 \circ f, 1 \circ g] &= 1 \circ [f, g] \\ &= (\text{im}[f, g]) \circ [f, g] \\ &= [(\text{im}[f, g]) \circ f, (\text{im}[f, g]) \circ g]. \end{aligned}$$

To show  $\text{im}[f, g]$  is the least upper-bound, assume  $p \geq \text{im } f$  and  $p \geq \text{im } g$ . Then

$$1 \circ [f, g] \geq p \circ [f, g] = [p \circ f, p \circ g] \geq [(\text{im } f) \circ f, (\text{im } g) \circ g] = 1 \circ [f, g],$$

hence  $1 \circ [f, g] = p \circ [f, g]$  and so  $p \geq \text{im}[f, g]$ , as desired. □

**Corollary** In an effectus with comprehension and images, we have for any sharp  $s, t$  a supremum (among all predicates) given by  $s \vee t = \text{im}[\pi_s, \pi_t]$ . In particular:  $s \vee t$  is sharp. V

In 200V we saw that comprehensions are precisely kernels of predicates. What about cokernels and quotients? 205

**Proposition** An effectus with quotients and images has all cokernels. A cokernel of a map  $f$  is given by a quotient  $\xi_{\text{im } f}$  of  $\text{im } f$ . II

**Proof** As  $0 = (\text{im}^\perp f) \circ f = 1 \circ \xi_{\text{im } f} \circ f$ , we have  $\xi_{\text{im } f} \circ f = 0$ . Assume  $g$  is a map with  $g \circ f = 0$ . Then  $1 \circ g \leq \text{im}^\perp f$  and so  $g = g' \circ \xi_{\text{im } f}$  for a unique  $g'$ . Thus indeed,  $\xi_{\text{im } f}$  is a cokernel of  $f$ . □ III

**Exercise** Show that in an effectus with comprehension and images, a map  $f$  is a quotient of sharp  $s$  if and only if  $f$  is a cokernel of a comprehension  $\pi_s$  of  $s$ . IV

### 3.5 $\diamond$ -effectuses

**206** In quantum physics it is not uncommon to restrict one’s attention to sharp predicates (i.e. projections). Does this restriction hurt the expressivity? It does not: every (normal) state on a von Neumann algebra is determined by its values on projections. In fact, on  $\mathcal{B}(\mathcal{H})$  with  $\dim \mathcal{H} \geq 3$ , Gleason’s famous theorem states that every measure on the projections extends uniquely to a state. We take the idea of restricting oneself to projections one step further: we also want to restrict to projections for our outcomes. It is rare for quantum processes to send projections to projections (see 210 III), so instead we consider the least projection above the outcome. The simple idea of taking the restriction to sharp predicates seriously leads to a host of interesting new notions. We will give these right off the bat and study their relevance later on.

**II Definition** A  $\diamond$ -effectus (“diamond effectus”) is an effectus with quotients, comprehension and images such that  $s^\perp$  is sharp for every sharp predicate  $s$ . In a  $\diamond$ -effectus, define for  $f: X \rightarrow Y$  the following restrictions to sharp predicates.

$$\text{SPred } X \begin{array}{c} \xrightarrow{f_\diamond} \\ \xleftarrow{f^\diamond} \end{array} \text{SPred } Y \quad \text{by} \quad f^\diamond(s) = \lceil s \circ f \rceil \quad \text{and} \quad f_\diamond(s) = \text{im } f \circ \pi_s$$

1. We say maps  $f: X \rightleftarrows Y: g$  are  $\diamond$ -adjoint if  $f^\diamond = g_\diamond$ .
2. An endomap  $f: X \rightarrow X$  is  $\diamond$ -self-adjoint if  $f$  is  $\diamond$ -adjoint to itself.
3. Two maps  $f, g: X \rightarrow Y$  are  $\diamond$ -equivalent if  $f^\diamond = g^\diamond$  (or equivalently whenever  $f_\diamond = g_\diamond$ , see 207 VIIa.)
4. A pure endomap  $f$  is  $\diamond$ -positive if  $f = g \circ g$  for some  $\diamond$ -self-adjoint  $g$ .

For brevity, write  $f^\square(s) = f^\diamond(s^\perp)^\perp$  and  $f_\square(s) = f_\diamond(s^\perp)^\perp$

**III Examples** The categories  $\mathbf{vN}^{\text{op}}$ ,  $\mathbf{CvN}^{\text{op}}$ ,  $\mathbf{EJA}^{\text{op}}$  and  $\mathbf{Set}$  are all  $\diamond$ -effectuses.

**207** Let’s investigate the basic properties of  $( )^\diamond$ ,  $( )_\diamond$  and  $( )^\square$ .

**II Exercise** Show that in a  $\diamond$ -effectus both  $f^\diamond$  and  $f^\square$  are order preserving maps.

**III Proposition** For  $f: X \rightarrow Y$  in a  $\diamond$ -effectus we have

$$f^\diamond(s) \leq t^\perp \quad \iff \quad f_\diamond(t) \leq s^\perp \tag{3.9}$$

for all sharp  $s, t$ . In other words:  $f_\diamond$  is the left order-adjoint of  $f^\square$ .

**IV Proof** To start, let’s prove the order-adjunction reformulation

$$f_\diamond(s) \leq t \quad \stackrel{(3.9)}{\iff} \quad f^\diamond(t^\perp) \equiv f^\square(t)^\perp \leq s^\perp \quad \iff \quad s \leq f^\square(t).$$

To prove (3.9), first assume  $f^\diamond(s) \leq t^\perp$ . Then  $s \circ f \leq \lceil s \circ f \rceil = f^\diamond(s) \leq t^\perp = \text{im}^\perp \pi_t$ , where the last equality is due to 203XII. Thus  $s \circ f \circ \pi_t = 0$  which is to say  $s \leq \text{im}^\perp f \circ \pi_t$ , so  $f_\diamond(t) = \text{im} f \circ \pi_t \leq s^\perp$ .

For the converse, assume  $f_\diamond(t) \leq s^\perp$ . Then as before (but in the other direction) we find  $s \circ f \circ \pi_t = 0$  and so  $s \circ f \leq t^\perp$ . Hence  $f^\diamond(s) = \lceil s \circ f \rceil \leq \lceil t^\perp \rceil = \lfloor t \rfloor^\perp = t^\perp$ , as desired.  $\square$

**Exercise** Use the fact that there is an order adjunction between  $f_\diamond$  and  $f^\square$  to show that in a  $\diamond$ -effectus V

- |                                      |   |
|--------------------------------------|---|
| 1. $f_\diamond$ is order preserving; | 4. $f^\diamond$ preserves suprema;                                |
| 2. $f_\diamond$ preserves suprema;   | 5. $f_\diamond \circ f^\square \circ f_\diamond = f_\diamond$ and |
| 3. $f^\square$ preserves infima;     | 6. $f^\square \circ f_\diamond \circ f^\square = f^\square$ .     |

**Lemma** In a  $\diamond$ -effectus  $(\ )^\diamond$ ,  $(\ )^\square$  and  $(\ )_\diamond$  are functorial — that is VI

- |   |  |   |
|---|--|---|
| 1. $(\text{id})^\diamond = \text{id}$ ,                   | 3. $(\text{id})^\square = \text{id}$ ,                 | 5. $(\text{id})_\diamond = \text{id}$ and                 |
| 2. $(f \circ g)^\diamond = g^\diamond \circ f^\diamond$ , | 4. $(f \circ g)^\square = g^\square \circ f^\square$ , | 6. $(f \circ g)_\diamond = f_\diamond \circ g_\diamond$ . |

**Proof** We get  $(\text{id})^\diamond = \text{id}$  directly from 203XII. For 2 we only need a single line: VII

$$(f \circ g)^\diamond(s) = \lceil s \circ f \circ g \rceil \stackrel{203\text{XIII}}{=} \lceil \lceil s \circ f \rceil \circ g \rceil = g^\diamond(f^\diamond(s)).$$

Note that we used that ceilings are sharp. Point 3 and 4 follow easily from 1 and 2 respectively. The identity  $(\text{id})_\diamond = \text{id}$  is again 203XII. We claim  $f_\diamond \circ g_\diamond$  is left order-adjoint to  $(f \circ g)^\square$ , indeed

$$f_\diamond(g_\diamond(s)) \leq t \iff g_\diamond(s) \leq f^\square(t) \iff s \leq g^\square(f^\square(t)) = (f \circ g)^\square(t).$$

Thus by uniqueness of order adjoints, we find  $(f \circ g)_\diamond = f_\diamond \circ g_\diamond$ .  $\square$

**Exercise** Derive from III that  $f^\diamond = g^\diamond$  if and only if  $f_\diamond = g_\diamond$ . VIIa

**Lemma** In a  $\diamond$ -effectus, sharp predicates are **order sharp** — that is: for any predicate  $p$  and sharp predicate  $s$  with  $p \leq s$  and  $p \leq s^\perp$ , we must have  $p = 0$ . 208

**Proof** Note  $\lceil p \rceil \leq \lceil s \rceil = s$  and  $\lceil p \rceil \leq \lceil s^\perp \rceil = s^\perp$ . So by 204I, there is an  $h$  with  $\pi_{\lceil p \rceil} = \pi_s \circ h$ . We compute II

$$1 \circ \pi_{\lceil p \rceil} = \lceil p \rceil \circ \pi_{\lceil p \rceil} = \lceil p \rceil \circ \pi_s \circ h \leq s^\perp \circ \pi_s \circ h = 0.$$

Thus  $\pi_{\lceil p \rceil} = 0$  and so  $p \leq \lceil p \rceil = \text{im} \pi_{\lceil p \rceil} = 0$ , as desired.  $\square$

- III **Proposition (Cho)** In a  $\diamond$ -effectus, the poset  $\text{SPred } X$  of sharp predicates on any object  $X$ , is a sub-effect algebra of  $\text{Pred } X$  and an orthomodular lattice.
- IV **Proof** Let  $C$  be a  $\diamond$ -effectus with some object  $X$ . We will show  $\text{SPred } X$  is a sub-effect algebra of  $\text{Pred } X$ , which is additionally an ortholattice. By 177 VI this is sufficient to show  $\text{SPred } X$  is orthomodular.
- V (Ortholattice) In 204 V we already saw that sharp  $s, t$  have a sharp supremum  $s \vee t = \text{im}[\pi_s, \pi_t]$  in  $\text{Pred } X$ . So  $s \vee t$  is also the supremum of  $s$  and  $t$  in  $\text{SPred } X$ . As  $(\ )^\perp$  is an order anti-automorphism of both  $\text{Pred } X$  and  $\text{SPred } X$ , we know  $(s^\perp \vee t^\perp)^\perp$  is the infimum of  $s$  and  $t$  in  $\text{Pred } X$  and  $\text{SPred } X$ . As any sharp predicate  $s$  is order sharp by I, we find  $s \wedge s^\perp = 0$  (and consequently  $s^\perp \vee s = 1$ ). We have shown  $\text{SPred } X$  is an ortholattice.
- VI (Sub-EA) Clearly  $0, 1$  are sharp and  $s^\perp$  is sharp for sharp  $s$  by definition of  $\diamond$ -effectus. To prove  $\text{SPred } X$  is a sub-effect algebra of  $\text{Pred } X$ , it only remains to be shown  $s \otimes t$  is sharp for sharp and summable  $s, t$ . So, assume  $s, t$  are sharp and  $s \perp t$ . Note  $s \wedge t \leq s \wedge s^\perp = 0$  as  $\text{SPred } X$  is an ortholattice and so by 177 Ia we find  $s \otimes t = s \vee t$ , which is indeed sharp.  $\square$
- VII **Corollary** In an  $\diamond$ -effectus  $C$  the assignment  $X \mapsto \text{SPred } X$ ,  $f \mapsto (f_\diamond, f^\square)$  yields a functor from  $C$  to  $\text{OMLatGal}$ , the category of orthomodular lattices with Galois connection between them, as defined in [55].
- VIII There is a rather different formula for the infima of sharp predicates, which will be useful later on.
- IX **Lemma** In a  $\diamond$ -effectus, we have  $s \wedge t = (\pi_s)_\diamond(\pi_s^\square(t))$  for sharp predicates  $s, t$ .
- X **Proof** It is easy to see  $(\pi_s)_\diamond(\pi_s^\square(t))$  is a lower bound: indeed  $(\pi_s)_\diamond(\pi_s^\square(t)) \leq t$  and  $(\pi_s)_\diamond(\pi_s^\square(t)) \leq (\pi_s)_\diamond(1) = \text{im } \pi_s = s$ . We have to show  $(\pi_s)_\diamond(\pi_s^\square(t))$  is the greatest lower bound. Let  $r$  be any sharp predicate with  $r \leq s$  and  $r \leq t$ . By 204 I we have  $\pi_r = \pi_s \circ h$  for some  $h$ . Thus

$$(\pi_r)_\diamond = (\pi_s)_\diamond \circ h_\diamond = (\pi_s)_\diamond \circ \pi_s^\square \circ (\pi_s)_\diamond \circ h_\diamond = (\pi_s)_\diamond \circ \pi_s^\square \circ (\pi_r)_\diamond.$$

Hence  $r = (\pi_r)_\diamond(1) = ((\pi_s)_\diamond \circ \pi_s^\square \circ (\pi_r)_\diamond)(1) = (\pi_s)_\diamond(\pi_s^\square(r)) \leq (\pi_s)_\diamond(\pi_s^\square(t))$ .  $\square$

- XI **Remark** In [23] there appears a similar result due to Jacobs, which is based on subtly different assumptions.
- XII **Exercise** Show that in a  $\diamond$ -effectus, we have  $(\xi^\square \circ \xi_\diamond)(t) = s \vee t$  for sharp  $s, t$  and quotient  $\xi$  of  $s$ . (Hint: mimic IX.)

209 We turn to  $\diamond$ -adjointness.

II **Exercise** Show the following basic properties of  $\diamond$ -adjointness

1.  $f^\diamond = g_\diamond$  ( $f$  is  $\diamond$ -adjoint to  $g$ ) if and only if  $f_\diamond = g^\diamond$ .
2. If  $f$  and  $g$  are  $\diamond$ -adjoint, then  $\text{im } f = [1 \circ g]$ .

**Exercise** Show in order: III

1. If  $f$  is  $\diamond$ -self-adjoint, then  $f \circ f$  is  $\diamond$ -self-adjoint.
2. If  $f$  is  $\diamond$ -positive, then  $f$  is  $\diamond$ -self-adjoint.
3. If  $f$  is  $\diamond$ -positive and  $f \circ f$  is pure, then  $f \circ f$  is  $\diamond$ -positive.

**Lemma** Let  $\alpha$  be an isomorphism in a  $\diamond$ -effectus. Then IV

1.  $s \circ \alpha$  is sharp for sharp predicates  $s$  and
2.  $\alpha^\diamond(s) = s \circ \alpha$  and  $\alpha_\diamond(s) = s \circ \alpha^{-1}$  (so  $\alpha$  and  $\alpha^{-1}$  are  $\diamond$ -adjoint).

**Proof** Let  $s$  be a sharp predicate. Then  $s = \text{im } \pi_s$ . Note  $\text{im } \alpha^{-1} \circ \pi_s = s \circ \alpha$  — V  
indeed,  $s \circ \alpha \circ \alpha^{-1} \circ \pi_s = 1$  and when  $p \circ \alpha^{-1} \circ \pi_s = 1$ , we must have  $p \circ \alpha^{-1} \geq s$ ,  
which gives  $p \geq s \circ \alpha$  as desired. So  $s \circ \alpha$  is indeed sharp.

So  $\alpha^\diamond(s) = [s \circ \alpha] = s \circ \alpha$  and  $\alpha_\diamond(s) = \text{im } \alpha \circ \pi_s = s \circ \alpha^{-1}$  as promised. □

**Definition** A map  $f$  in a  $\diamond$ -effectus is a **sharp map** provided  $s \circ f$  is sharp for 210  
all sharp predicates  $s$ .

**Exercise** Show that the following are equivalent. II

1.  $f$  is a sharp map.
2.  $[p \circ f] = [p] \circ f$  for every predicate  $p$ .

**Example** In  $\mathbf{vN}^{\text{op}}$  the sharp maps are exactly the mini-maps (i.e. the normal III  
\*-homomorphisms). See 99 XII.

## 3.6 &-effectuses

In a  $\diamond$ -effectus quotient and comprehension are not tied together by its axioms. 211  
With two additional axioms, we will see quotient and comprehension become  
tightly interwoven.

**Definition** An **&-effectus** (“andthen effectus”) is a  $\diamond$ -effectus such that II

1. for each object  $X$  and each predicate  $p$  on  $X$ , there is a unique  $\diamond$ -positive  
map  $\text{asrt}_p: X \rightarrow X$  (“assert  $p$ ”) with  $1 \circ \text{asrt}_p = p$  (see 206 II) and
2. for every quotient  $\xi: Y \rightarrow Z$  and comprehension  $\pi: X \rightarrow Y$  the compos-  
ite  $\xi \circ \pi$  is pure.

In an  $\&$ -effectus, we define  $p \& q \equiv q \circ \text{asrt}_p$ , pronounced “ $p$  andthen  $q$ ”. For brevity, we will write  $p^2 \equiv p \& p$ .

- III **Remarks** The  $\text{asrt}_p$  maps are named after the **assert** statement found in many programming languages. The **assert** statement checks whether the provided Boolean expression (i.e. predicate) is true (at the time of execution) — and if it is, it will continue the program without further action; if it isn’t, it will halt execution immediately. Our  $\text{asrt}_p$  maps can be thought of in the same way. Alternatively,  $\text{asrt}_p$  can be viewed as a filter that blocks states for which  $p^\perp$  holds. With this intuition, the predicate  $p \& q$  corresponds to whether  $p$  and  $q$  are true, by first checking  $p$  and then checking  $q$ .

The  $\text{asrt}_p$  maps can be used to construct more complicated measurements. For instance, let  $p, q, r$  be three predicates (measurement outcomes) on an object  $X$  with  $p \otimes q \otimes r = 1$ . Then  $\langle \text{asrt}_p, \text{asrt}_q, \text{asrt}_r \rangle : X \rightarrow X + X + X$  models measuring whether  $p, q, r$  hold without discarding the state of the (possibly affected) system after measurement (in contrast to  $\langle p, q, r \rangle : X \rightarrow 1 + 1 + 1$ ). If we discard the measurement outcome, we find the map  $\text{asrt}_p \otimes \text{asrt}_q \otimes \text{asrt}_r : X \rightarrow X$ , which is the side-effect of measuring  $p, q, r$ .

- IIIa The second axiom is closely related to polar decomposition: in the case of  $\text{vN}^{\text{op}}$  (flipping direction of arrows now), the unique isomorphism  $\varphi$  such that  $h \circ c = c' \circ \varphi \circ h'$  for the suitable standard filter  $c'$  and corner  $h'$  is given by  $\text{ad}_u$  where  $u$  is the isomtry from the polar decomposition  $u\sqrt{|h|}c(1)|h| \equiv \sqrt{c(1)}|h|$  restricted to a unitary.

- IV **Examples** The category  $\text{vN}^{\text{op}}$  is an  $\&$ -effectuses with  $\text{asrt}_a : b \mapsto \sqrt{ab}\sqrt{a}$ . The first axiom is proven in 105V and the second in 100III. The full subcategory  $\text{CvN}^{\text{op}}$  of commutative von Neumann algebras is a  $\&$ -effectus as well.

The only other known example of an  $\&$ -effectus is the category  $\text{EJA}^{\text{op}}$  of Euclidean Jordan algebras with positive unital maps in the opposite direction, see [128].

- V **Proposition** For a predicate  $p$  in an  $\&$ -effectus the following are equivalent.

1.  $p$  is sharp,
2.  $p \& p = p$  and
3.  $\text{asrt}_p \circ \text{asrt}_p = \text{asrt}_p$ .

- VI **Proof** First we prove that  $p$  is sharp if and only if  $p \& p = p$ . So, assume  $p$  is sharp. As  $\diamond$ -positive maps are  $\diamond$ -self-adjoint we have  $\text{im asrt}_p = [1 \circ \text{asrt}_p] = [p] = p$ . Thus  $p \& p = p \circ \text{asrt}_p = 1 \circ \text{asrt}_p = p$ . For the converse, assume  $p \& p = p$ . From  $\text{im asrt}_p = [1 \circ \text{asrt}_p] = [p]$ , we get  $[p] \circ \text{asrt}_p = p \& [p] = 1 \circ \text{asrt}_p = p$ . By assumption  $p \& p = p$ . So  $p \& ([p] \ominus p) = 0$ . Hence  $[p] \ominus p \leq \text{im}^\perp \text{asrt}_p = [p]^\perp$ . However  $[p] \ominus p \leq [p]$ . Thus by 208I get  $[p] \ominus p = 0$ . So  $p$  is indeed sharp.

Clearly, if  $\text{asrt}_p \circ \text{asrt}_p = \text{asrt}_p$ , then  $p = 1 \circ \text{asrt}_p = 1 \circ \text{asrt}_p \circ \text{asrt}_p = p \& p$ . It only remains to be shown  $\text{asrt}_p \circ \text{asrt}_p = \text{asrt}_p$  whenever  $p$  is sharp. So assume  $p$  is sharp. By definition  $\text{asrt}_p$  is pure:  $\text{asrt}_p = \pi \circ \xi$  for some quotient  $\xi$  and comprehension  $\pi$ . By assumption  $\xi \circ \pi$  is pure as well, so there is a quotient  $\xi'$  and comprehension  $\pi'$  with  $\xi \circ \pi = \pi' \circ \xi'$ . Note  $1 \circ \xi = 1 \circ \text{asrt}_p = p$  and  $1 \circ \text{asrt}_p \circ \text{asrt}_p = p \& p = p = 1 \circ \xi$ , hence

$$1 \circ \xi = 1 \circ \text{asrt}_p \circ \text{asrt}_p = 1 \circ \pi \circ \xi \circ \pi \circ \xi = 1 \circ \pi \circ \pi' \circ \xi' \circ \xi = 1 \circ \xi' \circ \xi,$$

so  $1 \circ \xi' = 1$ . Thus  $\xi'$  is an iso. Also  $(\text{im } \pi') \circ \xi \circ \pi = (\text{im } \pi') \circ \pi' \circ \xi' = 1 \circ \xi' = 1$  from which it follows  $p = \text{im } \pi \leq (\text{im } \pi') \circ \xi \leq 1 \circ \xi = p$ . Thus  $(\text{im } \pi') \circ \xi = p = 1 \circ \xi$ . Hence  $\text{im } \pi' = 1$  and so  $\pi'$  is an isomorphism. Now we know  $\pi \circ \pi'$  is a comprehension and  $\xi' \circ \xi$  is a quotient, we see  $\text{asrt}_p \circ \text{asrt}_p$  is pure. As  $\text{asrt}_p$  is  $\diamond$ -self-adjoint, we see  $\text{asrt}_p \circ \text{asrt}_p$  is  $\diamond$ -positive. By uniqueness of positive maps  $\text{asrt}_p \circ \text{asrt}_p = \text{asrt}_p$ , as desired.  $\square$

**Proposition** Let  $C$  be an  $\&$ -effectus and  $s$  be any sharp predicate. There exist **VII** comprehension  $\pi_s$  of  $s$  and quotient  $\zeta_s$  of  $s^\perp$  such that

$$\zeta_s \circ \pi_s = \text{id} \quad \text{and} \quad \pi_s \circ \zeta_s = \text{asrt}_s.$$

In fact, for every comprehension  $\pi$  of  $s$ , there is a quotient  $\xi$  of  $s^\perp$  with  $\xi \circ \pi = \text{id}$  and  $\pi \circ \xi = \text{asrt}_s$  and conversely for every quotient  $\xi$  of  $s^\perp$  there exists a comprehension  $\pi$  of  $s$  with  $\xi \circ \pi = \text{id}$  and  $\pi \circ \xi = \text{asrt}_s$ .

**Proof** Let  $s$  be any sharp predicate. By definition  $\diamond$ -positive maps are pure and **VIII** so there is a quotient  $\xi$  and comprehension  $\pi$  with  $\pi \circ \xi = \text{asrt}_s$ . So

$$\pi \circ \xi = \text{asrt}_s \stackrel{\vee}{=} \text{asrt}_s \circ \text{asrt}_s = \pi \circ \xi \circ \pi \circ \xi.$$

Thus  $\xi \circ \pi = \text{id}$ . We compute  $s = 1 \circ \text{asrt}_s = 1 \circ \xi$  and so

$$\begin{aligned} \text{im } \pi &= \text{im } \pi \circ \xi \circ \pi && \text{as } \xi \circ \pi = \text{id} \\ &= \text{im } \text{asrt}_s \circ \pi && \text{by dfn. } \xi \text{ and } \pi \\ &= (\text{asrt}_s)_\diamond(\text{im } \pi) && \text{by dfn. } ( )_\diamond \\ &= (\text{asrt}_s)^\diamond(\text{im } \pi) && \text{by } \diamond\text{-s.a. } \text{asrt}_s \\ &= [(\text{im } \pi) \circ \pi \circ \xi] && \text{by dfn. } ( )^\diamond \\ &= [1 \circ \xi] \\ &= s. \end{aligned}$$

Thus  $\pi$  is a comprehension of  $s$  and  $\xi$  is a quotient of  $s^\perp$ . We have proven the first part.

For the second part, let  $\pi'$  be any comprehension of  $s$ . Then  $\pi' = \pi \circ \alpha$  for some iso  $\alpha$ . Define  $\xi' = \alpha^{-1} \circ \xi$ . It is easy to see  $\pi' \circ \xi' = \text{asrt}_s$  and  $\xi' \circ \pi' = \text{id}$ . The other statement is proven in a similar way.  $\square$

IX **Notation** In an  $\&$ -effectus together with chosen comprehension  $\pi_s$  for  $s$ , we will write  $\zeta_s$  for the unique quotient for  $s^\perp$  satisfying  $\zeta_s \circ \pi_s = \text{id}$  and  $\pi_s \circ \zeta_s = \text{asrt}_s$ . We call this  $\zeta_s$  the **corresponding quotient** of  $\pi_s$  and vice versa  $\pi_s$  the **corresponding comprehension** of  $\zeta_s$ .

X **Warning**  $\xi_{s^\perp} = \zeta_s!$

XI **Proposition** In an  $\&$ -effectus both comprehensions and pure maps are closed under composition.

XII **Proof** We will first prove that comprehensions are closed under composition. Assume  $\pi_1: X \rightarrow Y$  and  $\pi_2: Y \rightarrow Z$  are comprehensions with  $s = \text{im } \pi_1$  and  $t = \text{im } \pi_2$ . We will show  $\pi_2 \circ \pi_1$  is a comprehension for  $\text{im } \pi_2 \circ \pi_1$ . To this end, let  $f: V \rightarrow Z$  be any map with  $(\text{im } \pi_2 \circ \pi_1)^\perp \circ f = 0$ . As  $\text{im } \pi_2 \circ \pi_1 \leq \text{im } \pi_2 = t$  we get  $t^\perp \circ f = 0$ , so  $f = \pi_2 \circ g_2$  for a unique  $g_2: V \rightarrow Y$ . Let  $\zeta_2$  be a quotient for  $t^\perp$  such that  $\zeta_2 \circ \pi_2 = \text{id}$ , which exists by VII. Then  $s \circ \zeta_2 \circ \pi_2 \circ \pi_1 = s \circ \pi_1 = 1$  so  $s \circ \zeta_2 \geq \text{im } \pi_2 \circ \pi_1$ . Consequently

$$s \circ g_2 = s \circ \zeta_2 \circ \pi_2 \circ g_2 = s \circ \zeta_2 \circ f \geq (\text{im } \pi_2 \circ \pi_1) \circ f = 1 \circ f = 1 \circ g_2.$$

Hence there exists a unique  $g_1: V \rightarrow X$  with  $\pi_1 \circ g_1 = g_2$  and so  $\pi_2 \circ \pi_1 \circ g_1 = f$ . By monicity of  $\pi_2 \circ \pi_1$ , this  $g_1$  is unique and so  $\pi_2 \circ \pi_1$  is indeed a comprehension.

XIII Now we will prove that pure maps are closed under composition. Assume  $g: X \rightarrow Y$  and  $f: Y \rightarrow Z$  are pure maps. Say  $f = \pi_1 \circ \xi_1$  and  $g = \pi_2 \circ \xi_2$  for some comprehensions  $\pi_1, \pi_2$  and quotients  $\xi_1, \xi_2$ . By definition of  $\&$ -effectus there is a comprehension  $\pi'$  and quotient  $\xi'$  such that  $\xi_1 \circ \pi_2 = \pi' \circ \xi'$ . By the previous point  $\pi_1 \circ \pi'$  is a comprehension and  $\xi' \circ \xi_2$  is a quotient by 197IX. Thus  $f \circ g = \pi_1 \circ \pi' \circ \xi' \circ \xi_2$  is pure.  $\square$

XIV **Exercise** Show that in an  $\&$ -effectus, we have  $\text{asrt}_p \circ \text{asrt}_p = \text{asrt}_{p \& p}$ .

XV **Exercise** Show that in an  $\&$ -effectus, we have

$$\begin{aligned} \text{im } f \leq s &\iff \text{asrt}_s \circ f = f \\ 1 \circ f \leq t &\iff f \circ \text{asrt}_t = f \end{aligned}$$

for any sharp predicates  $s$  and  $t$ .

XVI **Definition** Let  $C$  be an  $\&$ -effectus. In XI we saw pure maps are closed under composition. Write **Pure**  $C$  for the subcategory of pure maps.

212 **Lemma** In an  $\&$ -effectus,  $\zeta_{[p]} \circ \text{asrt}_p$  is a quotient for  $p^\perp$ .

II **Proof** Recall  $\text{im } \text{asrt}_p = [1 \circ \text{asrt}_p] = [p]$  and so

$$\text{im } \zeta_{[p]} \circ \text{asrt}_p = (\zeta_{[p]}) \circ ([p]) = \text{im } \zeta_{[p]} \circ \pi_{[p]} = \text{im id} = 1,$$

where  $\pi_{[p]}$  is the comprehension corresponding to  $\zeta_{[p]}$ . By 211XI  $\zeta_{[p]} \circ \text{asrt}_p = \pi \circ \xi$  for some comprehension  $\pi$  and quotient  $\xi$ . Putting it together:  $1 =$

$\text{im } \zeta_{[p]} \circ \text{asrt}_p = \text{im } \pi \circ \xi \leq \text{im } \pi$ , so  $\text{im } \pi = 1$  and thus  $\pi$  is an isomorphism, hence  $\zeta_{[p]} \circ \text{asrt}_p$  is a quotient for the orthocomplement of  $[p] \circ \text{asrt}_p = p$ .  $\square$

**Proposition** Every map  $f$  in an  $\&$ -effectus factors as

III

$$f = \pi_{\text{im } f} \circ g \circ \zeta_{[1 \circ f]} \circ \text{asrt}_{1 \circ f} \quad (3.10)$$

for a unique total and faithful map  $g$ . Furthermore the following holds.

1. If  $f$  is pure, then  $g$  is an isomorphism.
2. If  $f$  is pure and  $\text{im } f = 1$ , then  $f$  is a quotient.
3. If  $f$  is pure and  $1 \circ f = 1$ , then  $f$  is a comprehension.

**Proof** By the universal property of  $\pi_{\text{im } f}$ , there is a unique  $g'$  with  $f = \pi_{\text{im } f} \circ g'$ . IV To show  $g'$  is faithful, assume  $p \circ g' = 0$  for some predicate  $p$ . Then  $0 = p \circ g' = p \circ \zeta_{\text{im } f} \circ \pi_{\text{im } f} \circ g' = p \circ \zeta_{\text{im } f} \circ f$  and so  $p \circ \zeta_{\text{im } f} \leq \text{im}^\perp f$ , hence  $p = p \circ \zeta_{\text{im } f} \circ \pi_{\text{im } f} \leq (\text{im}^\perp f) \circ \pi_{\text{im } f} = 0$ , which shows  $g'$  is indeed faithful.

Note  $1 \circ g' = 1 \circ \pi_{\text{im } f} \circ g' = 1 \circ f$ . By 197 VII and I there is a unique total  $g$  with  $g' = g \circ \zeta_{[1 \circ f]} \circ \text{asrt}_{1 \circ f}$ . Clearly (3.10) holds and  $g$  is the unique map for which (3.10) holds as comprehensions are mono and quotients are epi. As  $1 = \text{im } g' = \text{im } g \circ \zeta_{[1 \circ f]} \circ \text{asrt}_{1 \circ f} \leq \text{im } g$ , we see  $\text{im } g = 1$  and so  $g$  is faithful.

To prove point 1, assume  $f$  is pure. That is:  $f = \pi \circ \xi$  for some comprehension  $\pi$  and quotient  $\xi$ . As  $1 \circ \pi = 1$  and  $\text{im } \xi = 1$ , we see  $\text{im } \pi = \text{im } f$  and  $1 \circ \xi = 1 \circ f$ . Thus  $\pi = \pi_{\text{im } f} \circ \alpha$  and  $\xi = \beta \circ \zeta_{[1 \circ f]} \circ \text{asrt}_{1 \circ f}$  for some isomorphisms  $\alpha$  and  $\beta$ . Thus  $f = \pi_{\text{im } f} \circ \alpha \circ \beta \circ \zeta_{[1 \circ f]} \circ \text{asrt}_{1 \circ f}$  and so by uniqueness of  $g$ , we see  $g = \alpha \circ \beta$  is an isomorphism.

To prove point 2, additionally assume  $\text{im } f = 1$ . Then  $\pi_{\text{im } f}$  is an iso and so using the previous, we see  $f$  is indeed a quotient. Point 3 is just as easy.  $\square$

**Proposition** If  $C$  is an  $\&$ -effectus, then  $(\text{Scal } C)^{\text{op}}$  is an effect divisoid, see 195 II. 213

**Proof** Let  $\lambda, \mu$  be scalars with  $\lambda \leq \mu$ . Recall that the scalar  $1 = \text{id}$  and so  $\mu = 1 \circ \text{asrt}_\mu = \text{asrt}_\mu$ . By 212 I, there is a unique  $\lambda'$  with  $\lambda' \circ \zeta_{[\mu]} \circ \mu = \lambda$ . Define  $\mu \setminus \lambda = \lambda' \circ \zeta_{[\mu]}$ . II

For the moment, assume  $\lambda = \mu$ . Then  $\lambda' \circ \zeta_{[\mu]} \circ \mu = \mu = 1 \circ \zeta_{[\mu]} \circ \mu$  as  $\text{im } \mu = \text{im } \text{asrt}_\mu = [\mu]$ . By epicity of  $\zeta_{[\mu]} \circ \mu$ , we see  $\lambda' = 1$ , hence  $\mu \setminus \mu = [\mu]$ . As we have  $\mu \leq [\mu]$  and  $[[\mu]] = [\mu]$ , we see axioms 2 and 3 hold.

We return to the general case to prove axiom 1. Clearly  $\mu \setminus \lambda = \lambda' \circ \zeta_{[\mu]} \leq 1 \circ \zeta_{[\mu]} = [\mu] = \mu \setminus \mu$  and  $\mu \circ_{\text{op}} \mu \setminus \lambda = \lambda' \circ \zeta_{[\mu]} \circ \mu = \lambda$  as required. Assume  $\sigma$  is an arbitrary scalar with  $\mu \circ_{\text{op}} \sigma = \lambda$  and  $\sigma \leq \mu \setminus \mu \equiv [\mu]$ . Then  $\sigma = \sigma' \circ \zeta_{[\mu]}$  for a unique  $\sigma'$ . As  $\lambda' \circ \zeta_{[\mu]} \circ \mu = \lambda = \sigma' \circ \zeta_{[\mu]} \circ \mu$  we must have  $\lambda' = \sigma'$ , whence  $\mu \setminus \lambda = \sigma$ .  $\square$

III **Lemma** In an  $\&$ -effectus, if  $s \perp t$  for sharp  $s, t$ , then  $s \& t = 0 = t \& s$ .

IV **Proof** Write  $r \equiv (s \otimes t)^\perp$ . As  $1 = s \otimes t \otimes r$  and  $s \& s = s$ , we have

$$s = s \& 1 = s \& s \otimes s \& t \otimes s \& r = s \otimes s \& t \otimes s \& r$$

and so  $s \& t \leq s \& t \otimes s \& r = 0$ . □

V **Exercise** Show that in an  $\&$ -effectus, we have

$$p \leq s \iff s \& p = p$$

for sharp  $s$  and any predicate  $p$ .

VI **Exercise** Show that in a  $\&$ -effectus, a map  $\langle f, g \rangle$  is sharp if and only if both  $f$  and  $g$  are sharp. (Hint: use 208 III and 203 XIV.)

### 3.7 †-effectuses

214 **Definition** A †-category (“dagger category”) is a category  $C$  together with an involutive identity-on-objects functor  $(\ )^\dagger: C \rightarrow C^{\text{op}}$  — that is, for all objects  $X$  and maps  $f, g$  in  $C$ , we have

- |  |                                 |
|--|---------------------------------|
| 1. $(f \circ g)^\dagger = g^\dagger \circ f^\dagger$ | 3. $f^{\dagger\dagger} = f$ and |
| 2. $\text{id}^\dagger = \text{id}$                   | 4. $X^\dagger = X$ .            |

Cf. [14, 49, 50, 102]. In any †-category we may define the following.

1. An endomap  $f$  is called †-self-adjoint if  $f^\dagger = f$ .
2. An endomap  $f$  is †-positive if  $f = g^\dagger \circ g$  for some other map  $g$ .
3. An isomorphism  $\alpha$  is called †-unitary whenever  $\alpha^{-1} = \alpha^\dagger$ .

II **Example** The category  $\text{Hilb}$  of Hilbert spaces with bounded linear maps is a †-category with the familiar adjoint as †.

215 **Definition** We call an  $\&$ -effectus  $C$  a †-effectus (“dagger effectus”) provided

1.  $\text{Pure } C$  is a †-category satisfying  $\text{asrt}_p^\dagger = \text{asrt}_p$  and  $f$  is  $\diamond$ -adjoint to  $f^\dagger$ ;
2. for every †-positive  $f$ , there is a unique †-positive  $g$  with  $g \circ g = f$  and
3.  $\diamond$ -positive maps are †-positive.

**Examples** In VI we will see that the category  $\mathbf{vN}^{\text{op}}$  is a  $\dagger$ -effectus. Recently we have shown with van de Wetering that the category  $\mathbf{EJA}^{\text{op}}$  of Euclidean Jordan algebras with positive unital maps in the opposite direction, is a  $\dagger$ -effectus as well. [128] II

**Theorem** An  $\&$ -effectus is a  $\dagger$ -effectus if and only if III

1. for every predicate  $p$ , there is a unique predicate  $q$  with  $q \& q = p$ ;
2.  $\text{asrt}_{p\&q}^2 = \text{asrt}_p \circ \text{asrt}_q^2 \circ \text{asrt}_p$  for all predicates  $p, q$  and
3. a quotient for a sharp predicate (e.g.  $\zeta_s$ ) is a sharp map, see 210I.

**Proof** Necessity will be proven in 216XI and sufficiency in 220II. □ IV

Especially the sufficiency requires quite some preparation. For convenience, call  $C$  a  $\dagger'$ -effectus if  $C$  is an  $\&$ -effectus satisfying axioms 1, 2 and 3 from the Theorem above. V

**Corollary** The category  $\mathbf{vN}^{\text{op}}$  of von Neumann algebras with ncpu-maps in the opposite direction, which is an  $\&$ -effectus by 211IV, is also a  $\dagger$ -effectus. VI

**Remarks** The dagger on the pure maps of  $\mathbf{vN}$  (that exists by the previous corollary) is fixed by the following two rules. VIIa

1. For any (nmiu-)isomorphism  $\vartheta$  we have  $\vartheta^\dagger = \vartheta^{-1}$  as  $\vartheta$  is  $\diamond$ -adjoint to  $\vartheta^{-1}$  and pure maps are rigid, see 102IX.
2. The standard filter  $c: \mathcal{A} \rightarrow [b] \mathcal{A} [b]$  for  $b \in \mathcal{A}$  (ie.  $c(a) = \sqrt{ba}\sqrt{b}$ , see 169VIII) has as dagger  $c^\dagger: [b] \mathcal{A} [b] \rightarrow \mathcal{A}$  given by  $c^\dagger(a) = \sqrt{ba}\sqrt{b}$ . (This follows from  $\text{asrt}_p^\dagger = \text{asrt}_p$  and 216VII.)

More concretely, for any pure ncp-map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  we can find a unique isomorphism  $\vartheta: [\varphi] \mathcal{A} [\varphi] \rightarrow [\varphi(1)] \mathcal{B} [\varphi(1)]$  such that

$$\varphi(a) = \sqrt{\varphi(1)} \vartheta([\varphi] a [\varphi]) \sqrt{\varphi(1)} \quad (\text{for all } a \in \mathcal{A})$$

(this follows from 100III and 98IX) and so by the previous two rules we get

$$\varphi^\dagger(b) = [\varphi] \vartheta^{-1} \left( \sqrt{\varphi(1)} b \sqrt{\varphi(1)} \right) [\varphi] \quad (\text{for all } b \in \mathcal{B}).$$

In special cases, we can give a simpler definition of the dagger. For instance, VIIIb  
if the pure map is of the form  $\varphi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ , then  $\varphi = \text{ad}_T$  for some operator  $T: \mathcal{H} \rightarrow \mathcal{H}$  and in that case  $\varphi^\dagger = \text{ad}_{T^*}$ .

Another example is a pure map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  between finite-dimensional von Neumann algebras. The Hilbert–Schmidt inner product  $[a, b] \equiv \text{tr } a^* b$  turns a finite-dimensional von Neumann algebra into a Hilbert space. The adjoint of  $\varphi$  with respect to this inner product equals  $\varphi^\dagger$ . This can be seen by considering the special cases of nmiu-isomorphisms and standard filters (noting nmiu-isomorphisms are trace-preserving).

VII **Remarks** In an  $\&$ -effectus with separating predicates (see 190II), the second axiom of a  $\dagger'$ -effectus is equivalent to

$$(p \& q)^2 \& r = p \& (q^2 \& (p \& r))$$

for all predicates  $p, q, r$ . This is essentially the *fundamental formula of quadratic Jordan algebras* [89, §4.2] (with  $U_{xy} \equiv x^2 \& y$ ).

The second axiom of a  $\dagger$ -effectus has been considered before in  $\dagger$ -categories by Selinger [103] who calls it the *unique square root axiom*.

216 **Lemma** In a  $\dagger$ -effectus, a map is  $\dagger$ -positive if and only if it is  $\diamond$ -positive.

II **Proof** Assume  $f$  is  $\dagger$ -positive. By assumption 2, there is a (unique)  $\dagger$ -positive  $g$  with  $f = g \circ g$ . By  $\dagger$ -positivity of  $g$ , there is an  $h$  with  $g = h^\dagger \circ h$ . Using the fact  $h$  is  $\diamond$ -adjoint to  $h^\dagger$ , we see  $g$  is  $\diamond$ -self-adjoint:

$$g^\diamond = (h^\dagger \circ h)^\diamond = h^\diamond \circ (h^\dagger)^\diamond = (h^\dagger)^\diamond \circ h_\diamond = (h^\dagger \circ h)_\diamond = g_\diamond.$$

Thus  $f$  is the square of the  $\diamond$ -self-adjoint map  $g$ , hence  $f$  is  $\diamond$ -positive.  $\square$

III **Lemma** Predicates in a  $\dagger$ -effectus have a unique square root: for every predicate  $p$ , there is a unique predicate  $q$  with  $q \& q = p$ .

IV **Proof** Let  $p: X \rightarrow 1$  be any predicate. By assumption 3, the  $\diamond$ -positive map  $\text{asrt}_p$  is also  $\dagger$ -positive. So by assumption 2, there is (a unique)  $\dagger$ -positive map  $f$  with  $f \circ f = \text{asrt}_p$ . Define  $q = 1 \circ f$ . By I  $f$  is  $\diamond$ -positive and so by uniqueness of  $\diamond$ -positive maps, we get  $f = \text{asrt}_{1 \circ f} \equiv \text{asrt}_q$ . We compute

$$q \& q = q \circ \text{asrt}_q \equiv 1 \circ f \circ \text{asrt}_{1 \circ f} = 1 \circ f \circ f = 1 \circ \text{asrt}_p = p,$$

which shows  $p$  has as square root  $q$ .

To show uniqueness, assume  $p = r \& r$  for some predicate  $r$ . Note

$$\text{asrt}_p = \text{asrt}_{r \& r} \stackrel{211\text{XIV}}{=} \text{asrt}_r \circ \text{asrt}_r.$$

As  $\text{asrt}_r$  is  $\diamond$ -positive, it is also  $\dagger$ -positive by the third axiom. So by the second axiom  $\text{asrt}_r = \text{asrt}_q$ . Thus  $r = q$ , which shows uniqueness of the square root.  $\square$

V **Proposition** In an  $\&$ -effectus with square roots (e.g.  $\dagger$ - or  $\dagger'$ -effectus) we have

$$\text{asrt}_p \circ \alpha = \alpha \circ \text{asrt}_{p \circ \alpha}$$

for every isomorphism  $\alpha$  and predicate  $p$ .

VI **Proof** There is some  $q$  with  $q \& q = p$ . The map  $\alpha^{-1} \circ \text{asrt}_q \circ \alpha$  is  $\diamond$ -self-adjoint:

$$\begin{aligned} (\alpha^{-1} \circ \text{asrt}_q \circ \alpha)_\diamond &= \alpha_\diamond^{-1} \circ (\text{asrt}_q)_\diamond \circ \alpha_\diamond \\ &\stackrel{209\text{IV}}{=} \alpha_\diamond^\diamond \circ (\text{asrt}_q)^\diamond \circ (\alpha^{-1})^\diamond \\ &= (\alpha^{-1} \circ \text{asrt}_q \circ \alpha)^\diamond. \end{aligned}$$

Thus  $\alpha^{-1} \circ \text{asrt}_q \circ \alpha \circ \alpha^{-1} \circ \text{asrt}_q \circ \alpha = \alpha^{-1} \circ \text{asrt}_p \circ \alpha$  is  $\diamond$ -positive. By uniqueness of  $\diamond$ -positive maps, we get  $\alpha^{-1} \circ \text{asrt}_p \circ \alpha = \text{asrt}_{1 \circ \alpha^{-1} \circ \text{asrt}_p \circ \alpha} = \text{asrt}_{p \circ \alpha}$ . Postcomposing  $\alpha$ , we find  $\text{asrt}_p \circ \alpha = \alpha \circ \text{asrt}_{p \circ \alpha}$ , as desired.  $\square$

**Proposition** In a  $\dagger$ -effectus:  $\zeta_s^\dagger = \pi_s$  (recall convention 211 IX for  $\pi_s$  and  $\zeta_s$ ). VII

**Proof** As  $\pi_s$  is  $\diamond$ -adjoint to  $\pi_s^\dagger$ , we have VIII

$$\text{im } \pi_s^\dagger = (\pi_s^\dagger)_\diamond(1) = (\pi_s)^\diamond(1) = [1 \circ \pi_s] = 1$$

and similarly  $[1 \circ \pi_s^\dagger] = \text{im } \pi_s = s$ . As  $1 \circ \pi_s^\dagger \leq [1 \circ \pi_s^\dagger] = s$ , there is some pure  $h$  with  $\pi_s^\dagger = h \circ \zeta_s$ . By 212 III, there is also some pure and faithful  $g$  with  $\zeta_s^\dagger = \pi_s \circ g$ . Using  $\zeta_s \circ \pi_s = \text{id}$  twice we find

$$\text{id} = \text{id}^\dagger = \pi_s^\dagger \circ \zeta_s^\dagger = h \circ \zeta_s \circ \pi_s \circ g = h \circ g. \quad (3.11)$$

Now  $1 = 1 \circ h \circ g \leq 1 \circ g$  and so  $g$  is total. Clearly  $\zeta_s^\dagger \circ \zeta_s$  is  $\dagger$ -positive, hence  $\diamond$ -positive and so by uniqueness of  $\diamond$ -positive maps:

$$\zeta_s^\dagger \circ \zeta_s = \text{asrt}_{1 \circ \zeta_s^\dagger \circ \zeta_s} = \text{asrt}_{1 \circ \pi_s \circ g \circ \zeta_s} = \text{asrt}_s = \pi_s \circ \zeta_s.$$

So by epicity of  $\zeta_s$ , we find  $\zeta_s^\dagger = \pi_s$ , as desired.  $\square$

**Corollary** In a  $\dagger$ -effectus,  $\pi_s$  is  $\diamond$ -adjoint to  $\zeta_s$ . Also  $\alpha^\dagger = \alpha^{-1}$  for any iso  $\alpha$ . IX

**Exercise** Show that in an  $\&$ -effectus where every predicate has a square root and where  $\pi_s$  is  $\diamond$ -adjoint to  $\zeta_s$  (e.g. a  $\dagger$ -effectus) we have  $\text{asrt}_p \circ \zeta_s = \zeta_s \circ \text{asrt}_{p \circ \zeta_s}$ . (Hint: mimic the proof of V.) X

**Theorem** A  $\dagger$ -effectus is a  $\dagger'$ -effectus. XI

**Proof** Assume  $C$  is a  $\dagger$ -effectus. Axiom 1 is already proven in III. XII

(Ax. 2) Let  $p, q$  be predicates. To start, note  $1 \circ \text{asrt}_q \circ \text{asrt}_p = p \& q$  and XIII

$$\text{im } \text{asrt}_q \circ \text{asrt}_p = (\text{asrt}_q)_\diamond(p) = (\text{asrt}_q)^\diamond(p) = [q \& p].$$

So using 212 III we know

$$\text{asrt}_q \circ \text{asrt}_p = \pi_{[q \& p]} \circ \alpha \circ \zeta_{[p \& q]} \circ \text{asrt}_{p \& q}$$

for some iso  $\alpha$ . Applying the dagger to both sides we get, using VII and IX:

$$\begin{aligned} \text{asrt}_p \circ \text{asrt}_q &= (\text{asrt}_q \circ \text{asrt}_p)^\dagger \\ &= \text{asrt}_{p \& q} \circ \zeta_{[p \& q]}^\dagger \circ \alpha^\dagger \circ \pi_{[q \& p]}^\dagger \\ &= \text{asrt}_{p \& q} \circ \pi_{[p \& q]} \circ \alpha^{-1} \circ \zeta_{[q \& p]}. \end{aligned}$$

Combining both:

$$\begin{aligned}
& \text{asrt}_p \circ \text{asrt}_q^2 \circ \text{asrt}_p \\
&= \text{asrt}_{p\&q} \circ \pi_{[p\&q]} \circ \alpha^{-1} \circ \zeta_{[q\&p]} \circ \pi_{[q\&p]} \circ \alpha \circ \zeta_{[p\&q]} \circ \text{asrt}_{p\&q} \\
&= \text{asrt}_{p\&q} \circ \pi_{[p\&q]} \circ \zeta_{[p\&q]} \circ \text{asrt}_{p\&q} \\
&= \text{asrt}_{p\&q} \circ \text{asrt}_{[p\&q]} \circ \text{asrt}_{p\&q} \\
&\stackrel{211\text{XV}}{=} \text{asrt}_{p\&q}^2,
\end{aligned}$$

as desired.

- XIV (Ax. 3) Pick any sharp predicates  $s, t$ . We want to show  $t \circ \zeta_s$  is sharp. To this end, we will show  $t \circ \zeta_s$  is the image of  $\pi_s \circ \pi_t$ . Clearly  $t \circ \zeta_s \circ \pi_s \circ \pi_t = 1$ . Let  $p$  be any sharp predicate with  $p \circ \pi_s \circ \pi_t = 1$ . Then  $p \circ \pi_s \geq \text{im } \pi_t = t$ . So  $p^\perp \circ \pi_s \leq t^\perp$  and hence  $[p^\perp \circ \pi_s] \leq t^\perp$ . As  $\pi_s$  is  $\diamond$ -adjoint to  $\zeta_s$  by IX, we get  $t \circ \zeta_s \leq [t \circ \zeta_s] \leq p$ , which shows  $t \circ \zeta_s$  is the image of  $\pi_s \circ \pi_t$  and consequently sharp.  $\square$

- 217 Let  $f$  be a pure map in a  $\dagger$ -effectus. We will work towards the definition of  $f^\dagger$ . By 212III we know there is an iso  $\alpha$  with

$$f = \pi_{\text{im } f} \circ \alpha \circ \zeta_{[1 \circ f]} \circ \text{asrt}_{1 \circ f}.$$

In a  $\dagger$ -effectus we have  $\text{asrt}_p^\dagger = \text{asrt}_p$ ,  $\zeta_s^\dagger = \pi_s$ ,  $\alpha^\dagger = \alpha^{-1}$  and  $\pi_s^\dagger = \zeta_s$  (for corresponding  $\zeta_s$  and  $\pi_s$ ) — so we are forced to define

$$f^\dagger = \text{asrt}_{1 \circ f} \circ \pi_{[1 \circ f]} \circ \alpha^{-1} \circ \zeta_{\text{im } f}, \quad (3.12)$$

where  $\zeta_{\text{im } f}$  is the unique corresponding quotient of  $\pi_{\text{im } f}$  and  $\pi_{[1 \circ f]}$  the unique corresponding comprehension of and  $\zeta_{[1 \circ f]}$ , see 211IX. Before we declare (3.12) a definition, we have to check whether it is independent of choice of  $\pi$  (and corresponding  $\zeta$ ). So suppose  $f = \pi' \circ \alpha' \circ \zeta' \circ \text{asrt}_{1 \circ f}$  for some iso  $\alpha'$ , comprehension  $\pi'$  of  $\text{im } f$  and quotient  $\zeta'$  of  $[1 \circ f]^\perp$ . There are isos  $\beta$  and  $\gamma$  such that  $\pi' = \pi_{\text{im } f} \circ \beta$  and  $\zeta' = \gamma \circ \zeta_{[1 \circ f]}$ . We will take a moment to relate  $\alpha$  and  $\alpha'$ : as  $\pi_{\text{im } f}$  is mono and  $\zeta_{1 \circ f} \circ \text{asrt}_{1 \circ f}$  is epic, we have  $\beta \circ \alpha' \circ \gamma = \alpha$  and so  $(\alpha')^{-1} = \gamma \circ \alpha^{-1} \circ \beta$ . To continue, it is easy to see  $\beta^{-1} \circ \zeta_{\text{im } f}$  is the unique corresponding quotient to  $\pi'$  and  $\pi_{[1 \circ f]} \circ \gamma^{-1}$  is the unique corresponding comprehension to  $\zeta'$ . So with this choice of quotient and comprehension, we are forced to define

$$\begin{aligned}
f^\dagger &= \text{asrt}_{1 \circ f} \circ \pi_{[1 \circ f]} \circ \gamma^{-1} \circ \alpha'^{-1} \circ \beta^{-1} \circ \zeta_{\text{im } f} \\
&= \text{asrt}_{1 \circ f} \circ \pi_{[1 \circ f]} \circ \gamma^{-1} \circ \gamma \circ \alpha^{-1} \circ \beta \circ \beta^{-1} \circ \zeta_{\text{im } f} \\
&= \text{asrt}_{1 \circ f} \circ \pi_{[1 \circ f]} \circ \alpha^{-1} \circ \zeta_{\text{im } f},
\end{aligned}$$

which is indeed consistent with (3.12). So we are justified to declare:

**Definition** In a  $\dagger'$ -effectus, for a pure map  $f$ , define

$$f^\dagger = \text{asrt}_{1 \circ f} \circ \pi_{[1 \circ f]} \circ \alpha^{-1} \circ \zeta_{\text{im } f},$$

where  $\alpha$  is the unique iso such that  $f = \pi_{\text{im } f} \circ \alpha \circ \zeta_{[1 \circ f]} \circ \text{asrt}_{1 \circ f}$ .

**Exercise** Show that in a  $\dagger'$ -effectus, we have

$$\text{asrt}_p^\dagger = \text{asrt}_p \quad \pi_s^\dagger = \zeta_s \quad \zeta_s^\dagger = \pi_s \quad \alpha^\dagger = \alpha^{-1}$$

for a quotient  $\zeta_s$  corresponding to  $\pi_s$  and iso  $\alpha$ .

To compute  $f^{\dagger\dagger}$ , we need to put  $f^\dagger$  in the standard form of 212III. To do this, we need to pull  $\text{asrt}_p$  from one side to the other. In this section we will work towards a general result for this.

**Lemma** In a  $\dagger'$ -effectus,  $\pi_s$  is  $\diamond$ -adjoint to  $\zeta_s$ .

**Proof** Pick any sharp  $t, u$ . We have to show  $t \circ \zeta_s \leq u^\perp$  if and only if  $u \circ \pi_s \leq t^\perp$ . So assume  $t \circ \zeta_s \leq u^\perp$ . Then  $t = t \circ \zeta_s \circ \pi_s \leq u^\perp \circ \pi_s = (u \circ \pi_s)^\perp$  so  $u \circ \pi_s \leq t^\perp$ . For the converse, assume  $u \circ \pi_s \leq t^\perp$ . Then  $u \circ \text{asrt}_s = u \circ \pi_s \circ \zeta_s \leq t^\perp \circ \zeta_s$ . From this, the  $\diamond$ -self-adjointness of  $\text{asrt}_s$  and the fact that  $t^\perp \circ \zeta_s$  is sharp, we find  $(t^\perp \circ \zeta_s)^\perp \circ \pi_s \circ \zeta_s \leq u^\perp$  and so

$$\begin{aligned} t \circ \zeta_s &= (t^\perp \circ \text{id})^\perp \circ \zeta_s \\ &= (t^\perp \circ \zeta_s \circ \pi_s)^\perp \circ \zeta_s \\ &= (t^\perp \circ \zeta_s)^\perp \circ \pi_s \circ \zeta_s \\ &\leq u^\perp, \end{aligned}$$

as desired.  $\square$

**Definition** In an  $\&$ -effectus, a map  $f$  is **pristine** if it is pure and  $1 \circ f$  is sharp.

**Remarks** In general, pristine maps are not closed under composition: in  $\mathbf{vN}^{\text{op}}$ , the maps  $\text{asrt}_{|0\rangle\langle 0|}$  and  $\text{asrt}_{|+\rangle\langle +|}$  on  $M_2$  are both pristine, but their composite is not. However, with a non-standard composition of pristine maps (namely  $f \cdot g \equiv \text{asrt}_{[1 \circ f \circ g]} \circ f \circ g$ ), the pristine maps are closed under composition and even form a  $\dagger$ -category [23]. Essentially the same construction has been found independently by [53] for partial isometries between Hilbert spaces.

**Exercise** Show that in an  $\&$ -effectus, every pristine map  $h$  is of the form

$$h = \pi_{\text{im } h} \circ \alpha \circ \zeta_{1 \circ h}$$

for some iso  $\alpha$ .

**VII Proposition** In a  $\dagger'$ -effectus, with pristine map  $h$ , we have

$$\text{asrt}_p \circ h = h \circ \text{asrt}_{p \circ h}$$

for any predicate  $p$  with  $p \leq \text{im } h$ .

**VIII Proof** For brevity, write  $t = 1 \circ h$  and  $s = \text{im } h$ . By VI there is some iso  $\alpha$  such that  $h = \pi_s \circ \alpha \circ \zeta_t$ . Note  $\text{im } \text{asrt}_p = [p] \leq [s] = s$  and so  $\text{asrt}_p = \text{asrt}_s \circ \text{asrt}_p$  by the first rule of 211 XV. By the second rule of 211 XV and  $1 \circ p = p \leq s$ , we see  $p = p \circ \text{asrt}_s$ . Thus

$$\begin{aligned} \text{asrt}_p \circ \pi_s &= \text{asrt}_s \circ \text{asrt}_p \circ \text{asrt}_s \circ \pi_s \\ &= \pi_s \circ \zeta_s \circ \text{asrt}_{p \circ \pi_s \circ \zeta_s} \circ \pi_s \\ &= \pi_s \circ \text{asrt}_{p \circ \pi_s} \circ \zeta_s \circ \pi_s && \text{by 216 X and II} \\ &= \pi_s \circ \text{asrt}_{p \circ \pi_s}. \end{aligned}$$

Putting everything together

$$\begin{aligned} \text{asrt}_p \circ h &= \text{asrt}_p \circ \pi_s \circ \alpha \circ \zeta_t \\ &= \pi_s \circ \text{asrt}_{p \circ \pi_s} \circ \alpha \circ \zeta_t \\ &= \pi_s \circ \alpha \circ \text{asrt}_{p \circ \pi_s \circ \alpha} \circ \zeta_t && \text{by 216 V} \\ &= \pi_s \circ \alpha \circ \zeta_t \circ \text{asrt}_{p \circ \pi_s \circ \alpha} \circ \zeta_t && \text{by 216 X and II} \\ &= h \circ \text{asrt}_{p \circ h}, \end{aligned}$$

as desired. □

**IX Exercise** Working in a  $\dagger'$ -effectus, show in order

1. if  $h \equiv \pi_{\text{im } h} \circ \alpha \circ \zeta_{1 \circ h}$  is some pristine map, then  $h^\dagger = \pi_{1 \circ h} \circ \alpha^{-1} \circ \zeta_{\text{im } h}$ ;
2.  $h^{\dagger\dagger} = h$  for any pristine  $h$ ;
3.  $h^\dagger \circ h = \text{asrt}_{1 \circ h}$  for any pristine map  $h$ ;
4.  $p \circ h^\dagger \leq \text{im } h$  for any predicate  $p$  and pristine map  $h$  and
5. if  $p \leq 1 \circ h$ , then  $\text{asrt}_{p \circ h^\dagger} \circ h = h \circ \text{asrt}_p$  for any pristine map  $h$ .

**X Proposition** In a  $\dagger'$ -effectus, for every pure map  $f$ , there exists a unique pristine map  $h$  with  $1 \circ h = [1 \circ f]$  and  $f = h \circ \text{asrt}_{1 \circ f}$ . Furthermore  $f^\dagger = \text{asrt}_{1 \circ f} \circ h^\dagger$ .

**XI Proof** By 212 III we know  $f = \pi_{\text{im } f} \circ \alpha \circ \zeta_{[1 \circ f]} \circ \text{asrt}_{1 \circ f}$  for some iso  $\alpha$ . Define  $h = \pi_{\text{im } f} \circ \alpha \circ \zeta_{[1 \circ f]}$ . Clearly  $1 \circ h = [1 \circ f]$  and  $f = h \circ \text{asrt}_{1 \circ f}$ . Also

$$\begin{aligned} f^\dagger &\equiv \text{asrt}_{1 \circ f} \circ \pi_{[1 \circ f]} \circ \alpha^{-1} \circ \zeta_{\text{im } f} \\ &= \text{asrt}_{1 \circ f} \circ \text{asrt}_{[1 \circ f]} \circ \pi_{[1 \circ f]} \circ \alpha^{-1} \circ \zeta_{\text{im } f} && \text{by 211 XV} \\ &\equiv \text{asrt}_{1 \circ f} \circ h^\dagger. \end{aligned}$$

Only uniqueness remains. Assume  $f = h' \circ \text{asrt}_{1 \circ f}$  for some pristine map  $h'$  with  $[1 \circ h'] = [1 \circ f]$ . By 212 III and 211 XV, there is an iso  $\alpha'$  with  $h' = \pi_{\text{im } h} \circ \alpha' \circ \zeta_{[1 \circ f]}$ . As  $\zeta_{[1 \circ f]} \circ \text{asrt}_{1 \circ f}$  is a quotient (by 212 I), quotients are faithful and  $f = h' \circ \text{asrt}_{1 \circ f}$ , we see  $\text{im } h' = \text{im } f$  and so  $\alpha = \alpha'$  by 212 III.  $\square$

**Proposition** In a  $\dagger$ '-effectus, we have  $f^{\dagger\dagger} = f$  for any pure map  $f$ . XII

**Proof** By X we have  $f = h \circ \text{asrt}_{1 \circ f}$  and  $f^\dagger = \text{asrt}_{1 \circ f} \circ h^\dagger$  for some pristine  $h$  XIII  
with  $1 \circ h = [1 \circ f]$ . Clearly  $1 \circ f \leq [1 \circ f] = 1 \circ h = \text{im } h^\dagger$ , so by VII we get  $f^\dagger = \text{asrt}_{1 \circ f} \circ h^\dagger = h^\dagger \circ \text{asrt}_{1 \circ f \circ h^\dagger}$ . Consequently

$$f^{\dagger\dagger} = \text{asrt}_{1 \circ f \circ h^\dagger} \circ h^{\dagger\dagger} \stackrel{\text{IX}}{=} \text{asrt}_{1 \circ f \circ h^\dagger} \circ h \stackrel{\text{IX}}{=} h \circ \text{asrt}_{1 \circ f} = f,$$

as desired.  $\square$

Now we tackle the most tedious part of 215 III: we will show  $(f \circ g)^\dagger = g^\dagger \circ f^\dagger$  219  
in a  $\dagger$ '-effectus. To avoid too much repetition, let us fix the setting.

**Setting** Let  $f, g$  be two composable pure maps in a  $\dagger$ '-effectus. For brevity, II  
write  $p = 1 \circ f$ ,  $q = 1 \circ g$ ,  $s = \text{im } f$  and  $t = \text{im } g$ . Let  $\varphi$  and  $\psi$  be the unique isomorphisms (see 212 III) such that

$$f = \pi_s \circ \varphi \circ \zeta_{[p]} \circ \text{asrt}_p \quad \text{and} \quad g = \pi_t \circ \psi \circ \zeta_{[q]} \circ \text{asrt}_q.$$

Define  $h = \pi_s \circ \varphi \circ \zeta_{[p]}$  and  $k = \pi_t \circ \psi \circ \zeta_{[q]}$ . To compute  $(f \circ g)^\dagger$ , we have to put  $f \circ g$  in the standard form of 212 III. We will do this step-by-step, first we put  $\zeta_{[p]} \circ \text{asrt}_p \circ \pi_t$  in standard form.

$$\begin{aligned} 1 \circ \zeta_{[p]} \circ \text{asrt}_p \circ \pi_t &= [p] \circ \text{asrt}_p \circ \pi_t \stackrel{211\text{XV}}{=} p \circ \pi_t \\ \text{im } \zeta_{[p]} \circ \text{asrt}_p \circ \pi_t &= (\zeta_{[p]} \circ \text{asrt}_p)_\diamond(t) \stackrel{218\text{II}}{=} [t \circ \text{asrt}_p \circ \pi_{[p]}]. \end{aligned}$$

Thus there is a unique isomorphism  $\chi$  with

$$\zeta_{[p]} \circ \text{asrt}_p \circ \pi_t = \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \chi \circ \zeta_{[p \circ \pi_t]} \circ \text{asrt}_{p \circ \pi_t}. \quad (3.13)$$

Next, we consider  $\text{asrt}_{p \circ k} \circ \text{asrt}_q$ , clearly

$$1 \circ \text{asrt}_{p \circ k} \circ \text{asrt}_q = p \circ k \circ \text{asrt}_q = p \circ g.$$

Concerning the image, first note  $p \circ k = p \circ \pi_t \circ \psi \circ \zeta_{[q]} \leq 1 \circ \zeta_{[q]} = [q]$  and so we must have  $\text{im } \text{asrt}_{p \circ k} \leq [q]$ , which implies

$$\text{im } \text{asrt}_{p \circ k} \circ \text{asrt}_q = (\text{asrt}_{p \circ k})_\diamond(q) = [[q] \circ \text{asrt}_{p \circ k}] = [p \circ k].$$

So there is a unique isomorphism  $\omega$  such that

$$\text{asrt}_{p \circ k} \circ \text{asrt}_q = \pi_{[p \circ k]} \circ \omega \circ \zeta_{[p \circ g]} \circ \text{asrt}_{p \circ g}. \quad (3.14)$$

Next, we consider  $\zeta_{[p \circ \pi_t]} \circ \psi \circ \zeta_{[q]}$ . Note  $\psi \circ \zeta_{[q]}$  is a quotient for a sharp predicate, hence sharp and so  $[p \circ \pi_t] \circ \psi \circ \zeta_{[q]} = [p \circ \pi_t \circ \psi \circ \zeta_{[q]}] = [p \circ k]$  by 210II. As quotients are closed under composition (197IX), there is an iso  $\beta$  with

$$\zeta_{[p \circ \pi_t]} \circ \psi \circ \zeta_{[q]} = \beta \circ \zeta_{[p \circ k]}. \quad (3.15)$$

Finally, we deal with  $\pi_s \circ \varphi \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]}$ . By 211XI this is again a comprehension. We compute


$$\begin{aligned} \text{im } \pi_s \circ \varphi \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} &= (\pi_s \circ \varphi \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]}) \circ (1) \\ &\stackrel{218\text{II}}{=} \zeta_s^\diamond(\varphi \circ ([t \circ \text{asrt}_p \circ \pi_{[p]}])) \\ &= [t \circ \text{asrt}_p \circ \pi_{[p]}] \circ \varphi^{-1} \circ \zeta_s \\ &\stackrel{210\text{II}}{=} [t \circ \text{asrt}_p \circ \pi_{[p]} \circ \varphi^{-1} \circ \zeta_s] \\ &= [t \circ f^\dagger] \end{aligned}$$

and so there must be a unique iso  $\alpha$  with

$$\pi_s \circ \varphi \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} = \pi_{[t \circ f^\dagger]} \circ \alpha. \quad (3.16)$$

**III Lemma** In setting II, we have  $f \circ g = \pi_{[t \circ f^\dagger]} \circ \alpha \circ \chi \circ \beta \circ \omega \circ \zeta_{[p \circ g]} \circ \text{asrt}_{p \circ g}$ .

**IV Proof** It's a long, but easy verification, either with a diagram



or with equational reasoning

$$\begin{aligned} f \circ g &= \pi_s \circ \varphi \circ \zeta_{[p]} \circ \text{asrt}_p \circ \pi_t \circ \psi \circ \zeta_{[q]} \circ \text{asrt}_q \\ &\stackrel{(3.13)}{=} \pi_s \circ \varphi \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \chi \circ \zeta_{[p \circ \pi_t]} \circ \text{asrt}_{p \circ \pi_t} \circ \psi \circ \zeta_{[q]} \circ \text{asrt}_q \\ &\stackrel{218\text{VII}}{=} \pi_s \circ \varphi \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \chi \circ \zeta_{[p \circ \pi_t]} \circ \psi \circ \zeta_{[q]} \circ \text{asrt}_{p \circ k} \circ \text{asrt}_q \\ &\stackrel{(3.15)}{=} \pi_s \circ \varphi \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \chi \circ \beta \circ \zeta_{[p \circ k]} \circ \text{asrt}_{p \circ k} \circ \text{asrt}_q \\ &\stackrel{(3.16)}{=} \pi_{[t \circ f^\dagger]} \circ \alpha \circ \chi \circ \beta \circ \zeta_{[p \circ k]} \circ \text{asrt}_{p \circ k} \circ \text{asrt}_q \\ &\stackrel{(3.14)}{=} \pi_{[t \circ f^\dagger]} \circ \alpha \circ \chi \circ \beta \circ \zeta_{[p \circ k]} \circ \pi_{[p \circ k]} \circ \omega \circ \zeta_{[p \circ g]} \circ \text{asrt}_{p \circ g} \\ &= \pi_{[t \circ f^\dagger]} \circ \alpha \circ \chi \circ \beta \circ \omega \circ \zeta_{[p \circ g]} \circ \text{asrt}_{p \circ g}, \end{aligned}$$

whichever the Reader might prefer.  $\square$

**Corollary**  $(f \circ g)^\dagger = \text{asrt}_{p \circ g} \circ \pi_{[p \circ g]} \circ \omega^{-1} \circ \beta^{-1} \circ \chi^{-1} \circ \alpha^{-1} \circ \zeta_{[t \circ f^\dagger]}$ . V

To show  $(f \circ g)^\dagger = g^\dagger \circ f^\dagger$ , it is sufficient to proof that the ‘daggered’ version of each of the subdiagrams (3.16), (3.15), (3.13), (3.14) and 218 VII of the above diagram holds. We start with the simple ones. VI

**Lemma** In setting II, the daggered version of (3.15) holds — that is: VII

$$\pi_{[p \circ k]} \circ \beta^{-1} = \pi_{[q]} \circ \psi^{-1} \circ \pi_{[p \circ \pi_t]}.$$

**Proof** The map  $\pi_{[q]} \circ \psi^{-1} \circ \pi_{[p \circ \pi_t]} \circ \beta$  is a comprehension for  $[p \circ k]$  — indeed VIII

$$\begin{aligned} (\pi_{[q]} \circ \psi^{-1} \circ \pi_{[p \circ \pi_t]} \circ \beta) \circ (1) &= (\zeta_{[q]}^\circ \circ \psi^\circ)([p \circ \pi_t]) \\ &= [p \circ \pi_t] \circ \zeta_{[q]} \circ \psi \\ &= [p \circ \pi_t \circ \zeta_{[q]} \circ \psi] \\ &= [p \circ k]. \end{aligned}$$

Furthermore

$$\begin{aligned} \zeta_{[p \circ k]} \circ \pi_{[q]} \circ \psi^{-1} \circ \pi_{[p \circ \pi_t]} \circ \beta \\ &\stackrel{(3.15)}{=} \beta^{-1} \circ \zeta_{[p \circ \pi_t]} \circ \psi \circ \zeta_{[q]} \circ \pi_{[q]} \circ \psi^{-1} \circ \pi_{[p \circ \pi_t]} \circ \beta \\ &= \text{id}. \end{aligned}$$

So  $\pi_{[q]} \circ \psi^{-1} \circ \pi_{[p \circ \pi_t]} \circ \beta$  is the unique comprehension corresponding to  $\zeta_{[p \circ k]}$  — that is:  $\pi_{[q]} \circ \psi^{-1} \circ \pi_{[p \circ \pi_t]} \circ \beta = \pi_{[p \circ k]}$ , as desired.  $\square$

**Exercise** Show that in the setting II, the daggered version of (3.16) holds — i.e. IX

$$\alpha^{-1} \circ \zeta_{[t \circ f^\dagger]} = \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \varphi^{-1} \circ \zeta_s.$$

(Hint: mimic the proof of VII.)

**Exercise** Show that in the setting II, we have X

$$\pi_{[q]} \circ \psi^{-1} \circ \text{asrt}_{p \circ \pi_t} = \text{asrt}_{p \circ k} \circ \pi_{[q]} \circ \psi^{-1}.$$

(This is the daggered version of the subdiagram marked 218 VII.)

**Proposition** If in a  $\dagger'$ -effectus,  $\nu$  is the unique iso (cf. 216 XIII) such that XI

$$\text{asrt}_a \circ \text{asrt}_b = \pi_{[a \& b]} \circ \nu \circ \zeta_{[b \& a]} \circ \text{asrt}_{b \& a},$$

then  $\text{asrt}_b \circ \text{asrt}_a = \text{asrt}_{b \& a} \circ \pi_{[b \& a]} \circ \nu^{-1} \circ \zeta_{[a \& b]}$ .

XII **Proof** Let  $\mu$  be the unique iso with  $\text{asrt}_b \circ \text{asrt}_a = \pi_{[b \& a]} \circ \mu \circ \zeta_{[a \& b]} \circ \text{asrt}_{a \& b}$ . We will see  $\mu = \nu^{-1}$ . For brevity, write

$$\overline{a \& b} = (a \& b) \circ \pi_{[a \& b]} \quad \overline{b \& a} = (b \& a) \circ \pi_{[b \& a]}$$

By 218 IX and 211 XV, we have

$$\pi_{[a \& b]} \circ \text{asrt}_{\overline{a \& b}} \circ \zeta_{[a \& b]} = \pi_{[a \& b]} \circ \zeta_{[a \& b]} \circ \text{asrt}_{a \& b} = \text{asrt}_{a \& b}.$$

Now the second axiom of a  $\dagger$ -effectus comes into play

$$\begin{aligned} \pi_{[a \& b]} \circ \text{asrt}_{\overline{a \& b}}^2 \circ \zeta_{[a \& b]} &= \pi_{[a \& b]} \circ \text{asrt}_{\overline{a \& b}} \circ \zeta_{[a \& b]} \circ \pi_{[a \& b]} \circ \text{asrt}_{\overline{a \& b}} \circ \zeta_{[a \& b]} \\ &= \text{asrt}_{\overline{a \& b}}^2 \\ &= \text{asrt}_a \circ \text{asrt}_b^2 \circ \text{asrt}_a \\ &= \pi_{[a \& b]} \circ \nu \circ \text{asrt}_{\overline{b \& a}} \circ \zeta_{[b \& a]} \circ \pi_{[b \& a]} \circ \mu \circ \text{asrt}_{\overline{a \& b}} \circ \zeta_{[a \& b]} \\ &= \pi_{[a \& b]} \circ \nu \circ \text{asrt}_{\overline{b \& a}} \circ \mu \circ \text{asrt}_{\overline{a \& b}} \circ \zeta_{[a \& b]}. \end{aligned}$$

Thus as quotients are epis and comprehensions are monos:

$$\text{asrt}_{\overline{a \& b}}^2 = \nu \circ \text{asrt}_{\overline{b \& a}} \circ \mu \circ \text{asrt}_{\overline{a \& b}}. \quad (3.17)$$

We want to show  $\text{asrt}_{\overline{a \& b}}$  is an epi. First note

$$\overline{[a \& b]} \circ \zeta_{[a \& b]} \stackrel{210 \text{II}}{=} \overline{[a \& b] \circ \zeta_{[a \& b]}} \stackrel{211 \text{XV}}{=} [a \& b] = 1 \circ \zeta_{[a \& b]}$$

and so  $\text{im } \text{asrt}_{\overline{a \& b}} = \overline{[a \& b]} = 1$ , which tells us  $\text{asrt}_{\overline{a \& b}}$  is a quotient and therefore an epi. So, from (3.17) we get  $\text{asrt}_{\overline{a \& b}} = \nu \circ \text{asrt}_{\overline{b \& a}} \circ \mu$  and so

$$\overline{a \& b} = 1 \circ \text{asrt}_{\overline{a \& b}} = 1 \circ \nu \circ \text{asrt}_{\overline{b \& a}} \circ \mu = \overline{b \& a} \circ \mu. \quad (3.18)$$

Using this equation again in the previous, we find

$$\nu \circ \mu \circ \text{asrt}_{\overline{a \& b}} = \nu \circ \mu \circ \text{asrt}_{\overline{b \& a}} \circ \mu = \nu \circ \text{asrt}_{\overline{b \& a}} \circ \mu = \text{asrt}_{\overline{a \& b}}.$$

Thus  $\nu \circ \mu = \text{id}$  and so  $\mu = \nu^{-1}$ . Write  $l = \pi_{[b \& a]} \circ \nu^{-1} \circ \zeta_{[a \& b]}$ . Then

$$\begin{aligned} (a \& b) \circ l^\dagger &= (a \& b) \circ \pi_{[a \& b]} \circ \nu \circ \zeta_{[b \& a]} && \text{by 218 IX} \\ &= \overline{a \& b} \circ \nu \circ \zeta_{[b \& a]} \\ &= \overline{b \& a} \circ \zeta_{[b \& a]} && \text{by (3.18) and } \mu = \nu^{-1} \\ &= b \& a. \end{aligned}$$

And so, keeping in mind  $a \& b \leq [a \& b] = 1 \circ l$ , we have

$$\begin{aligned}
\text{asrt}_b \circ \text{asrt}_a &= l \circ \text{asrt}_{a\&b} \\
&= \text{asrt}_{(a\&b) \circ l^\dagger} \circ l && \text{by 218 IX} \\
&= \text{asrt}_{b\&a} \circ l \\
&= \text{asrt}_{b\&a} \circ \pi_{[b\&a]} \circ \nu^{-1} \circ \zeta_{[a\&b]},
\end{aligned}$$

as promised. □

**Corollary** In setting II, the daggered version of (3.14) holds — that is: XIII

$$\text{asrt}_q \circ \text{asrt}_{p \circ k} = \text{asrt}_{p \circ g} \circ \pi_{[p \circ g]} \circ \omega^{-1} \circ \zeta_{[p \circ k]}.$$

**Lemma** In setting II, the daggered version of (3.13) holds — that is: XIV

$$\zeta_t \circ \text{asrt}_p \circ \pi_{[p]} = \text{asrt}_{p \circ \pi_t} \circ \pi_{[p \circ \pi_t]} \circ \chi^{-1} \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]}.$$

**Proof** The heavy lifting has been done in XI already. To start, note XV

$$\begin{aligned}
\text{asrt}_p \circ \text{asrt}_t &\stackrel{211\text{XV}}{=} \text{asrt}_{[p]} \circ \text{asrt}_p \circ \text{asrt}_t \\
&= \pi_{[p]} \circ \zeta_{[p]} \circ \text{asrt}_p \circ \pi_t \circ \zeta_t \\
&\stackrel{(3.13)}{=} \pi_{[p]} \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \chi \circ \zeta_{[p \circ \pi_t]} \circ \text{asrt}_{p \circ \pi_t} \circ \zeta_t \\
&\stackrel{218\text{VII}}{=} \pi_{[p]} \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \chi \circ \zeta_{[p \circ \pi_t]} \circ \zeta_t \circ \text{asrt}_{p \circ \pi_t} \circ \zeta_t \\
&= \pi_{[p]} \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \chi \circ \zeta_{[p \circ \pi_t]} \circ \zeta_t \circ \text{asrt}_{t\&p} \\
&= \pi_{[p\&t]} \circ \alpha_2 \circ \chi \circ \beta_2 \circ \zeta_{[t\&p]} \circ \text{asrt}_{t\&p},
\end{aligned}$$

where  $\alpha_2$  and  $\beta_2$  are the unique isomorphisms such that

$$\pi_{[p]} \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} = \pi_{[p\&t]} \circ \alpha_2 \quad \zeta_{[p \circ \pi_t]} \circ \zeta_t = \beta_2 \circ \zeta_{[t\&p]}.$$

With the same reasoning as in IX and VII, we see

$$\zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \zeta_{[p]} = \alpha_2^{-1} \circ \zeta_{[p\&t]} \quad \pi_t \circ \pi_{[p \circ \pi_t]} = \pi_{[t\&p]} \circ \beta_2^{-1}.$$


Now we can apply XI:

$$\begin{aligned}
\zeta_t \circ \text{asrt}_p \circ \pi_{[p]} &\stackrel{211\text{XV}}{=} \zeta_t \circ \text{asrt}_t \circ \text{asrt}_p \circ \pi_{[p]} \\
&\stackrel{\text{XI}}{=} \zeta_t \circ \text{asrt}_{t\&p} \circ \pi_{[t\&p]} \circ \beta_2^{-1} \circ \chi^{-1} \circ \alpha_2^{-1} \circ \zeta_{[p\&t]} \circ \pi_{[p]} \\
&= \zeta_t \circ \text{asrt}_{t\&p} \circ \pi_t \circ \pi_{[p \circ \pi_t]} \circ \chi^{-1} \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \zeta_{[p]} \circ \pi_{[p]} \\
&= \zeta_t \circ \text{asrt}_{p \circ \pi_t} \circ \zeta_t \circ \pi_t \circ \pi_{[p \circ \pi_t]} \circ \chi^{-1} \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \\
&= \text{asrt}_{p \circ \pi_t} \circ \pi_{[p \circ \pi_t]} \circ \chi^{-1} \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]},
\end{aligned}$$

as desired. □

XVI **Proposition** In a  $\dagger'$ -effectus  $(f \circ g)^\dagger = g^\dagger \circ f^\dagger$  holds.

XVII **Proof** We work in setting II. The equality  $(f \circ g)^\dagger = g^\dagger \circ f^\dagger$  follows from V and the commutativity of the following diagram



or alternatively by

$$\begin{aligned}
g^\dagger \circ f^\dagger &= \text{asrt}_q \circ \pi_{[q]} \circ \psi^{-1} \circ \zeta_t \circ \text{asrt}_p \circ \pi_{[p]} \circ \varphi^{-1} \circ \zeta_s \\
&\stackrel{\text{XIV}}{=} \text{asrt}_q \circ \pi_{[q]} \circ \psi^{-1} \circ \text{asrt}_{p \circ \pi_t} \circ \pi_{[p \circ \pi_t]} \circ \chi^{-1} \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \varphi^{-1} \circ \zeta_s \\
&\stackrel{\text{X}}{=} \text{asrt}_q \circ \text{asrt}_{p \circ k} \circ \pi_{[q]} \circ \psi^{-1} \circ \pi_{[p \circ \pi_t]} \circ \chi^{-1} \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \varphi^{-1} \circ \zeta_s \\
&\stackrel{\text{VII}}{=} \text{asrt}_q \circ \text{asrt}_{p \circ k} \circ \pi_{[p \circ k]} \circ \beta^{-1} \circ \chi^{-1} \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \varphi^{-1} \circ \zeta_s \\
&\stackrel{\text{IX}}{=} \text{asrt}_q \circ \text{asrt}_{p \circ k} \circ \pi_{[p \circ k]} \circ \beta^{-1} \circ \chi^{-1} \circ \alpha^{-1} \circ \zeta_{[t \circ f^\dagger]} \\
&\stackrel{\text{XIII}}{=} \text{asrt}_{p \circ g} \circ \pi_{[p \circ g]} \circ \omega^{-1} \circ \zeta_{[p \circ k]} \circ \pi_{[p \circ k]} \circ \beta^{-1} \circ \chi^{-1} \circ \alpha^{-1} \circ \zeta_{[t \circ f^\dagger]} \\
&= \text{asrt}_{p \circ g} \circ \pi_{[p \circ g]} \circ \omega^{-1} \circ \beta^{-1} \circ \chi^{-1} \circ \alpha^{-1} \circ \zeta_{[t \circ f^\dagger]} \\
&\stackrel{\text{V}}{=} (f \circ g)^\dagger,
\end{aligned}$$

whichever the Reader might prefer.  $\square$

220 We are ready to finish the proof of 215 III.

II **Theorem** A  $\dagger'$ -effectus is a  $\dagger$ -effectus with  $\dagger$  as defined in 217 II.

III **Proof** Let  $C$  be a  $\dagger'$ -effectus.

IV (Ax. 1) By 219 XVI, 218 XII and 217 III the  $\dagger$  defined in 217 II turns Pure  $C$  into a  $\dagger$ -category. Also by 217 III, we have  $\text{asrt}_p^\dagger = \text{asrt}_p$  for any predicate  $p$ . Pick any pure  $f$ . We have to show  $f$  is  $\diamond$ -adjoint to  $f^\dagger$ . By 212 III we have  $f =$

$\pi_{\text{im } f} \circ \varphi \circ \zeta_{[1 \circ f]} \circ \text{asrt}_{1 \circ f}$  for some iso  $\alpha$ . We compute

$$\begin{aligned}
f_{\diamond} &= (\pi_{\text{im } f})_{\diamond} \circ \varphi_{\diamond} \circ (\zeta_{[1 \circ f]})_{\diamond} \circ (\text{asrt}_{1 \circ f})_{\diamond} \\
&\stackrel{218\text{II}}{=} (\zeta_{\text{im } f})_{\diamond} \circ \varphi_{\diamond} \circ (\pi_{[1 \circ f]})_{\diamond} \circ (\text{asrt}_{1 \circ f})_{\diamond} \\
&\stackrel{209\text{IV}}{=} (\zeta_{\text{im } f})_{\diamond} \circ (\varphi^{-1})_{\diamond} \circ (\pi_{[1 \circ f]})_{\diamond} \circ (\text{asrt}_{1 \circ f})_{\diamond} \\
&= (\text{asrt}_{1 \circ f} \circ \pi_{[1 \circ f]} \circ \varphi^{-1} \circ \zeta_{\text{im } f})_{\diamond} \\
&= (f^{\dagger})_{\diamond},
\end{aligned}$$

so  $f$  is indeed  $\diamond$ -adjoint to  $f^{\dagger}$ .

(Ax. 2) Let  $f$  be a  $\dagger$ -positive map. That is:  $f = h^{\dagger} \circ h$  for some pure map  $h$ . V  
By 212III we have  $h = \pi_{\text{im } h} \circ \alpha \circ \zeta_{[1 \circ h]} \circ \text{asrt}_{1 \circ h}$  for some iso  $\alpha$ . We compute

$$\begin{aligned}
f &= h^{\dagger} \circ h \\
&= \text{asrt}_{1 \circ h} \circ \pi_{[1 \circ h]} \circ \alpha^{-1} \circ \zeta_{\text{im } h} \circ \pi_{\text{im } h} \circ \alpha \circ \zeta_{[1 \circ h]} \circ \text{asrt}_{1 \circ h} \\
&= \text{asrt}_{1 \circ h} \circ \text{asrt}_{[1 \circ h]} \circ \text{asrt}_{1 \circ h} \\
&\stackrel{211\text{XV}}{=} \text{asrt}_{1 \circ h} \circ \text{asrt}_{1 \circ h}.
\end{aligned}$$

Form this it follows that  $\dagger$ -positive maps are  $\diamond$ -positive. Let  $q$  be the predicate such that  $q \& q = 1 \circ h$ . Then

$$\text{asrt}_q^{\dagger} \circ \text{asrt}_q = \text{asrt}_q \circ \text{asrt}_q \stackrel{211\text{XIV}}{=} \text{asrt}_{q \& q} = \text{asrt}_{1 \circ h}.$$

Thus  $\text{asrt}_{1 \circ h}$  is a  $\dagger$ -positive with  $\text{asrt}_{1 \circ h} \circ \text{asrt}_{1 \circ h} = f$ . We have to show  $\text{asrt}_{1 \circ h}$  is the unique map with this property. Let  $g$  be any  $\dagger$ -positive map with  $g \circ g = f$ . Recall both  $g$  and  $f$  are  $\diamond$ -positive, hence

$$\text{asrt}_{1 \circ f} = f = g \circ g = \text{asrt}_{1 \circ g} \circ \text{asrt}_{1 \circ g} \stackrel{211\text{XIV}}{=} \text{asrt}_{(1 \circ g) \& (1 \circ g)}.$$

So  $(1 \circ g) \& (1 \circ g) = 1 \circ f = (1 \circ h) \& (1 \circ h)$ . Hence  $1 \circ g = 1 \circ h$  by uniqueness of the square root. So  $g = \text{asrt}_{1 \circ g} = \text{asrt}_{1 \circ h} = h$ , as desired.

(Ax. 3) Let  $\text{asrt}_p$  be any  $\diamond$ -positive map. Write  $q$  for the unique predicate VI  
with  $q \& q = p$ . Then

$$\text{asrt}_p = \text{asrt}_{q \& q} \stackrel{211\text{XIV}}{=} \text{asrt}_q \circ \text{asrt}_q = \text{asrt}_q^{\dagger} \circ \text{asrt}_q,$$

so  $\text{asrt}_p$  is  $\dagger$ -positive, as desired. □

### 3.7.1 Dilations

Being the main topic of the first part of this thesis, we cannot finish our discussion of  $\dagger$ -effectuses without mentioning dilations. The presented abstract theory of dilations in a  $\dagger$ -effectus is preliminary and as of yet rather unimpressive.

**221**

II **Definition** Let  $f: X \rightarrow Y$  be any map in an effectus. A **dilation** of  $f$  is a triple  $(P, \varrho, h)$  of a sharp total map  $\varrho: P \rightarrow Y$  and a pure map  $h: X \rightarrow P$  such that  $\varrho \circ h = f$  and the following universal property holds.

For every triple  $(P', \varrho', h')$  with  $\varrho': P' \rightarrow Y$  total sharp,  $h': X \rightarrow P'$  arbitrary and  $f = \varrho' \circ h'$ , there is a unique  $\sigma: P \rightarrow P'$  with  $\sigma \circ h = h'$  and  $\varrho' \circ \sigma = \varrho$ .

We say an effectus **has dilations** if there exists a dilation for every map.

III **Example** The effectus  $\mathbf{vN}^{\text{op}}$  (180V) has dilations, as shown in 154III, but the full subcategory  $\mathbf{CvN}^{\text{op}}$  of commutative von Neumann algebras does not as shown in the next exercise.

IIIa **Exercise** In this exercise you will show that  $\mathbf{CvN}^{\text{op}}$  does not have dilations. First, show that a corner between commutative von Neumann algebras is multiplicative. Conclude that if  $\mathbf{CvN}^{\text{op}}$  were to have dilations, then any ncpu-map would be multiplicative, quod non.

IV **Proposition** Let  $C$  be an effectus with dilations.

1. If  $(P_1, \varrho_1, h_1)$  and  $(P_2, \varrho_2, h_2)$  are both dilations of  $f$ , then there is a unique isomorphism  $\alpha: P_1 \rightarrow P_2$  with  $\alpha \circ h_1 = h_2$  and  $\varrho_2 \circ \alpha = \varrho_1$ .
2. Conversely, if  $(P, \varrho, h)$  is a dilation of  $f$  and  $\alpha: P \rightarrow P'$  is some isomorphism, then  $(P', \varrho \circ \alpha^{-1}, \alpha \circ h)$  is also a dilation of  $f$ .
3.  $(X, \varrho, \text{id})$  is the dilation of a sharp total map  $\varrho$ .
4. If  $(P, \varrho, h)$  is some dilation, then  $(P, \text{id}, h)$  is the dilation of  $h$ .
5. For any quotient  $\xi: X \rightarrow Q$  and map  $f: Q \rightarrow Y$  with dilation  $(P, \varrho, h)$ , the triple  $(P, \varrho, h \circ \xi)$  is a dilation of  $f \circ \xi$ .
6. Conversely, if  $(P, \varrho, h)$  is a dilation of  $f \circ \xi$  for a map  $f: Q \rightarrow Y$  and quotient  $\xi: X \rightarrow Q$ , then  $(P, \varrho, h'')$  is a dilation of  $f$ , where  $h'': Q \rightarrow P$  is the unique map with  $h'' \circ \xi = h$ .
7. If for  $i = 1, 2$  the triple  $(P_i, \varrho_i, h_i)$  is a dilation of  $f_i: X_i \rightarrow Y$ , then  $(P_1 + P_2, [\varrho_1, \varrho_2], h_1 + h_2)$  is a dilation of  $[f_1, f_2]: X_1 + X_2 \rightarrow Y$ .

V **Proof** Point 1 has been proven in 140VIII. Point 2 is easily verified. To prove point 3, assume  $\varrho$  is a sharp total map. Assume  $\varrho = \varrho' \circ h'$  for some  $h'$  and sharp total  $\varrho'$ . Then  $h'$  is clearly the unique map satisfying  $h' = h' \circ \text{id}$  (and  $\varrho = \varrho' \circ h'$ ), which demonstrates point 3.

For point 4, assume  $(P, \varrho, h)$  is some dilation. Assume  $h = \varrho' \circ h'$  for some  $h'$  and sharp total  $\varrho'$ . There is a unique  $\sigma$  with  $h' = \sigma \circ h$  and  $\varrho \circ \varrho' \circ \sigma = \varrho$ . Now  $\varrho' \circ \sigma \circ h = \varrho' \circ h' = h$ . So  $\varrho' \circ \sigma$  satisfies the properties of the unique

mediating map  $\text{id}: P \rightarrow P$ , hence  $\varrho' \circ \sigma = \text{id}$ . Only uniqueness of  $\sigma$  remains, so assume  $\sigma'$  is a ncp-map with  $h' = \sigma' \circ h$  and  $\varrho' \circ \sigma' = \text{id}$ . Then  $\varrho \circ \varrho' \circ \sigma' = \varrho$  and so  $\sigma = \sigma'$ .

To demonstrate point 5, assume  $\varrho' \circ h' = f \circ \xi$  for some  $h'$  and total sharp  $\varrho'$ . Then  $1 \circ h' = 1 \circ \varrho' \circ h' = 1 \circ f \circ \xi \leq 1 \circ \xi$ , so there is a unique  $h''$  with  $h'' \circ \xi = h'$ . By epicity of  $\xi$ , we get  $\varrho' \circ h'' = f$  from  $\varrho' \circ h'' \circ \xi = \varrho' \circ h' = f \circ \xi$ . Thus there is a unique  $\sigma$  with  $\sigma \circ h = h''$  and  $\varrho' \circ \sigma = \varrho$  and so  $\sigma \circ h \circ \xi = h'' \circ \xi = h'$ . Only uniqueness of  $\sigma$  remains, so suppose  $\sigma'$  is a map with  $\sigma' \circ h \circ \xi = h'$  and  $\varrho' \circ \sigma = \varrho$ . Then  $\sigma' \circ h = h''$  by epicity of  $\xi$ , hence  $\sigma' = \sigma$  as desired.

For point 6, assume  $f = \varrho' \circ h'$  for some  $h'$  and sharp total  $\varrho'$ . Then  $f \circ \xi = \varrho' \circ h' \circ \xi$  and so there is a unique  $\sigma$  with  $\varrho \circ \sigma = \varrho'$  and  $\sigma \circ h' \circ \xi = h \equiv h'' \circ \xi$ . Thus  $\sigma \circ h' = h''$  by epicity of  $\xi$ . To show uniqueness, assume  $\sigma'$  is such that  $\sigma' \circ h' = h''$  and  $\varrho \circ \sigma' = \varrho'$ . Then  $\sigma' \circ h' \circ \xi = h'' \circ \xi = h$  and indeed  $\sigma' = \sigma$ .

To show point 7, assume  $[f_1, f_2] = \varrho' \circ h'$  for some  $h'$  and total sharp  $\varrho'$ . For any  $i = 1, 2$  we have  $\varrho' \circ h' \circ \kappa_i = f_i$ , so there is a unique  $\sigma_i$  with  $\sigma_i \circ h_i = h' \circ \kappa_i$  and  $\varrho_i = \varrho' \circ \sigma_i$ . Define  $\sigma = [\sigma_1, \sigma_2]$ . Clearly  $\sigma \circ (h_1 + h_2) = [\sigma_1 \circ h_1, \sigma_2 \circ h_2] = [h' \circ \kappa_1, h' \circ \kappa_2] = h'$  and  $\varrho' \circ \sigma = [\varrho' \circ \sigma_1, \varrho' \circ \sigma_2] = [\varrho_1, \varrho_2]$ . Assume  $\sigma'$  is any ncp-map with  $\sigma' \circ (h_1 + h_2) = h'$  and  $\varrho' \circ \sigma' = [\varrho_1, \varrho_2]$ . For  $i = 1, 2$  we have  $\varrho' \circ \sigma' \circ \kappa_i = \varrho_i$  and  $h' \circ \kappa_i = \sigma' \circ (h_1 + h_2) \circ \kappa_i = \sigma' \circ h_i$ , so  $\sigma' \circ \kappa_i = \sigma_i$ . Hence  $\sigma' = [\sigma' \circ \kappa_1, \sigma' \circ \kappa_2] = [\sigma_1, \sigma_2] = \sigma$ .  $\square$

With some additional assumptions, the existence of dilations forces the existence of some familiar quantum gates. Assume  $C$  is a  $\dagger$ -effectus with dilation with a scalar  $\lambda$  such that  $\lambda^\perp = \lambda$ . (For instance:  $\lambda = \frac{1}{2}$  in  $\mathbf{vN}^{\text{op}}$ .) Let  $(P, \langle s, s^\perp \rangle, h)$  be a dilation of the map  $\langle h, h^\perp \rangle: 1 \rightarrow 1 + 1$ . We think of  $P$  as an abstract qubit — indeed in  $\mathbf{vN}^{\text{op}}$  any dilation of  $\langle \frac{1}{2}, \frac{1}{2} \rangle$  is of the form  $(M_2, \langle e, e^\perp \rangle, h)$  where  $h$  is the vector state for some  $v \in \mathbb{C}^2$  and  $e$  is a projector on  $w\mathbb{C}$ , with  $\langle v, w \rangle = \frac{1}{\sqrt{2}}$  (for instance  $v = |0\rangle$  and  $w = |+\rangle$ ).

By assumption  $\langle s, s^\perp \rangle \circ h = \langle \lambda, \lambda \rangle$ , so  $s \circ h = s^\perp \circ h = \lambda$ . and  $\langle s^\perp, s \rangle \circ h = \langle \lambda, \lambda \rangle$ . Hence by the definition of dilation, there is a unique map  $X: P \rightarrow P$  with  $X \circ h = h$  and  $\langle s^\perp, s \rangle \circ X = \langle s, s^\perp \rangle$ , whence  $s \circ X = s^\perp \circ X$  and  $s^\perp \circ X = s \circ X$ . Clearly  $X \circ X \circ h = h$  and  $\langle s, s^\perp \rangle \circ X \circ X = \langle s, s^\perp \rangle$ , so by uniqueness of the mediating map  $\text{id}: P \rightarrow P$ , we must have  $X \circ X = \text{id}$ . Thus the isomorphism  $X$  behaves like the  $X$ -gate — indeed, in  $\mathbf{vN}^{\text{op}}$ , we have  $X = \text{ad}_X$  if  $e = |0\rangle\langle 0|$ .  $\square$

For the next gate, assume  $s$  is pure and  $\text{im } s = 1$ . Note  $s^\dagger$  is total. As  $\text{asrt}_\lambda = \lambda$ , we have  $h^\dagger \circ s^\dagger = \lambda^\dagger = \lambda$  and so  $(h^\dagger)^\perp \circ s^\dagger = \lambda^\perp = \lambda$ . Combined:  $\langle h^\dagger, (h^\dagger)^\perp \rangle \circ s^\dagger = \langle \lambda, \lambda \rangle$ . Thus using the universal property of the dilation, there is a unique map  $H: P \rightarrow P$  with  $H \circ h = s^\dagger$  and  $\langle h^\dagger, (h^\dagger)^\perp \rangle \circ H = \langle s, s^\perp \rangle$ . So  $h^\dagger \circ H = s$ . In general, we do not know a lot more about  $H$ . Assume  $H$  is an comprehen-  $\square$

sion (as it is in  $\mathbf{vN}^{\text{op}}$  in this situation). Then  $s^\dagger = H^\dagger \circ h$  and  $s = h^\dagger \circ H^\dagger$ , so  $s^\perp = (h^\dagger)^\perp \circ H^\dagger$ , whence  $\langle s, s^\perp \rangle = \langle h^\dagger, (h^\dagger)^\perp \rangle \circ H^\dagger$ . Thus, by uniqueness of  $H$ , we must have  $H^\dagger = H$ . Thus both  $H^\dagger$  and  $H$  are total, which forces  $H \circ H^\dagger = H^\dagger \circ H = \text{asrt}_1 = \text{id}$ , by uniqueness of  $\diamond$ -positive maps. Hence  $H \circ H = \text{id}$ ,  $s \circ H = h^\dagger$  and  $h^\dagger \circ H = s$ . So  $H$  is similar to the Hadamard-gate — indeed, in  $\mathbf{vN}^{\text{op}}$ , we have  $H = \text{ad}_H$  if  $e = |0\rangle\langle 0|$ .

- IV Do our additional assumptions (purity of  $s$ ,  $\text{im } s = 1$ , etc.) follow from more general principles? Can we define the other gates, like the CNOT? Do these gates interact in the usual way? Unfortunately, we do not have an answer to any of these questions.

**223** In 157 IV we saw that for any ncp-map  $\varphi$  with Paschke dilation  $(\mathcal{P}, \varrho, h)$ , there is a linear order isomorphism  $[0, 1]_{\varrho(\mathcal{A})} \square \cong [0, \varphi]_{\text{nCP}}$ . To finish the discussion of dilations, we mention how to state this correspondence in an  $\&$ -effectus.

- II **Definition** In an  $\&$ -effectus, for any predicate  $p$ , define  $\text{seff}_p \equiv \text{asrt}_p \otimes \text{asrt}_{p^\perp}$ . This map models the side-effect of measuring  $p$ . For any map  $f: X \rightarrow Y$ , define

$$\text{Inv } f \equiv \{p; p \in \text{Pred } X; f \circ \text{seff}_p = f\}.$$

This is the set of predicates whose measurement does not disturb  $f$ .

- III **Lemma** In  $\mathbf{vN}^{\text{op}}$ , for any nmiu-map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}$ , we have  $\text{Inv } \varrho = [0, 1]_{\varrho(\mathcal{A})} \square$  — that is:  $\text{Inv } \varrho$  is the set of effects of the commutator of the image of  $\varrho$ .
- IV **Proof** Assume  $a \in [0, 1]_{f(\mathcal{A})} \square$ . Simply unfolding definitions, we find  $\text{seff}_a(\varrho(b)) = \sqrt{a}\varrho(b)\sqrt{a} + \sqrt{1-a}\varrho(b)\sqrt{1-a} = \varrho(b)$  for any  $b \in \mathcal{A}$ , so  $a \in \text{Inv } \varrho$ . For the converse, assume  $a \in \text{Inv } \varrho$ . Recall  $(\mathcal{A}, \varrho, \text{id})$  is a Paschke dilation of  $\varrho$ . As

$$\varrho = \text{seff}_a \circ \varrho = (\text{asrt}_a \circ \varrho) \otimes (\text{asrt}_{1-a} \circ \varrho) \geq_{\text{nCP}} \text{asrt}_a \circ \varrho,$$

we know by 157 IV that there is a  $t \in [0, 1]_{\varrho(\mathcal{A})} \square$  with  $\text{asrt}_a(\varrho(b)) \equiv \sqrt{a}\varrho(b)\sqrt{a} = t\varrho(b)$  for all  $b \in \mathcal{A}$ , but then clearly  $a = \sqrt{a}\varrho(1)\sqrt{a} = t\varrho(1) = t$ .  $\square$

- V **Definition** Let  $C$  be an  $\&$ -effectus. For a map  $f: X \rightarrow Y$ , write

$$\downarrow f \equiv \{g; g: X \rightarrow Y; g \leq f\}.$$

We say a dilation  $(P, \varrho, h)$  of  $f$  has **the order correspondence** if there is an order isomorphism  $\Theta: \downarrow f \rightarrow \text{Inv } \varrho$  such that for every  $g \in \downarrow f$ , we have

$$g = \varrho \circ \text{asrt}_{\Theta(g)} \circ h.$$

- VI **Example** By 157 IV every dilation in  $\mathbf{vN}^{\text{op}}$  has the order correspondence.

## 3.8 Comparisons

### 3.8.1 Dagger kernel categories

To finish this chapter, we relate the structures in a  $\dagger$ -effectus with existing structures in the literature. Just like we try to axiomatize  $\mathbf{vN}^{\text{op}}$ , Heunen in the third chapter of his Ph.D-thesis attempted to axiomatize  $\mathbf{Hilb}$ , the category of Hilbert spaces with bounded operators. Heunen came very close: he proves that a  $\dagger$ -category obeying certain additional axioms must embed into  $\mathbf{Hilb}$ . [49, 3.7.18] We will discover that some of his axioms for  $\mathbf{Hilb}$  hold in  $\text{Pure}(C)$  for a  $\dagger$ -effectus  $C$  and others do not in general. (We do not cover all his axioms.) 224

**Definition** Let  $C$  be a  $\dagger$ -category. We say that an arrow  $f$  is  $\dagger$ -mono iff  $f^\dagger \circ f = \text{id}$  and dually  $f$  is  $\dagger$ -epi iff  $f \circ f^\dagger = \text{id}$ . An arrow  $f$  is a  $\dagger$ -partial isometry iff  $f = m \circ e$  for some  $\dagger$ -mono  $m$  and  $\dagger$ -epi  $e$ . II

A  $\dagger$ -kernel of  $f$  is an equalizer of  $f$  with  $0$ , which is  $\dagger$ -mono. A  $\dagger$ -kernel category is a  $\dagger$ -category with zero object and where each arrow has a  $\dagger$ -kernel [49, 3.2.20].

**Proposition** In  $\text{Pure}(C)$  for some  $\dagger$ -effectus  $C$ , the following holds. III

1.  $f$  is  $\dagger$ -mono iff  $f$  is a comprehension.  
Dually:  $f$  is  $\dagger$ -epi iff  $f$  is a quotient of a sharp predicate.

2. The  $\dagger$ -partial isometries are exactly the pristine maps, see 218 IV.

Furthermore:  $\text{Pure } C$  is a  $\dagger$ -kernel category: the  $\dagger$ -kernel of  $f$  is given by  $\pi_{(1 \circ f)^\perp}$ .

**Proof** Let  $f$  be any  $\dagger$ -mono pure map, say  $f \equiv \pi_s \circ \alpha \circ \zeta_{[p]} \circ \text{asrt}_p$  for some iso  $\alpha$ ,  $s \equiv \text{im } f$  and  $p \equiv 1 \circ f$ . As we saw before (using 211 XV), we see  $\text{id} = f^\dagger \circ f = \text{asrt}_{p^2}$ . Thus  $p \& p = p^2 = 1 = 1 \& 1$  and so by uniqueness of the square root we get  $p = 1$ . Hence  $f$  is a comprehension. IV

Dualizing, we see  $\dagger$ -epi are exactly quotients of sharp predicates. Now it is clear that the  $\dagger$ -partial isometries are exactly the pristine maps.

By the previous  $\pi_{(1 \circ f)^\perp}$  is  $\dagger$ -mono. To see it is the  $\dagger$ -kernel of  $f$ , we have to show it is the equalizer of  $f$  with  $0$ . Clearly  $(1 \circ f)^\perp \circ \pi_{(1 \circ f)^\perp} = 1$  and so  $1 \circ f \circ \pi_{(1 \circ f)^\perp} = 0$ , whence  $f \circ \pi_{(1 \circ f)^\perp} = 0$ . Assume  $f \circ g = 0$  for some pure map  $g$ . Then  $1 \circ f \circ g = 0$ , so  $1 \circ f \leq \text{im}^\perp g$ , hence  $(1 \circ f)^\perp \geq \text{im } g$ , so  $g = \pi_{(1 \circ f)^\perp} \circ g'$  for a unique  $g'$ . V  $\square$

**Exercise\*** Let  $C$  be a  $\dagger$ -effectus. In this exercise, you will show that in general  $\text{Pure } C$  does not have finite coproducts. In particular,  $\text{Pure } C$  does not have  $\dagger$ -biproducts, see [49, 3.2.15]. Consider  $C \equiv \mathbf{vN}^{\text{op}}$ . Reasoning towards contradiction, assume  $(\text{Pure } \mathbf{vN}^{\text{op}})^{\text{op}}$  has a product  $\mathbb{C} \xleftarrow{\pi_1} \mathcal{A} \xrightarrow{\pi_2} \mathbb{C}$  of  $\mathbb{C}$  and  $\mathbb{C}$ . First show that for any non-zero pure map  $f: \mathcal{A} \rightarrow \mathbb{C}$ , there exists a Hilbert space  $\mathcal{H}$ , an element  $x \in \mathcal{H}$ , a von Neumann algebra  $\mathcal{C}$  and VI

isomorphism  $\varphi: \mathcal{B}(\mathcal{H}) \oplus \mathcal{C} \rightarrow \mathcal{A}$  with  $f(\text{varphi}(T, c)) = \langle x, Tx \rangle$ . Use this on  $\pi_1, \pi_2$  and  $\langle \text{id}_{\mathbb{C}}, \text{id}_{\mathbb{C}} \rangle$  to show there is an isomorphism  $\varphi: \mathcal{B}(\mathcal{H}) \oplus \mathcal{C} \cong \mathcal{A}$  with  $\pi_1 \circ \varphi = \langle x, (\cdot)x \rangle$  and  $\pi_2 \circ \varphi = \langle y, (\cdot)y \rangle$  for some von Neumann algebra  $\mathcal{C}$ , Hilbert space  $\mathcal{H}$  and  $x, y \in \mathcal{H}$ . Press on: show  $\mathcal{C}$  is trivial. Then prove  $\dim \mathcal{H} \leq 2$ . Almost there: show  $\dim \mathcal{H} \geq 2$ . Derive a contradiction.

- VII Exercise\*** In this exercise you will show that  $\text{Pure}(\text{vN}^{\text{op}})$  does not have all coequalizers. It is helpful to first consider two concrete coequalizers in  $\text{Hilb}$  and  $\text{vN}^{\text{op}}$ . (These coequalizers capture unordered pairs of qubits.)

Write  $\sigma: \mathbb{C}^2 \otimes \mathbb{C}^2 \rightarrow \mathbb{C}^2 \otimes \mathbb{C}^2$  for the unitary fixed by  $\sigma|ij\rangle = |ji\rangle$ . The equalizer of  $\sigma$  with  $\text{id}$  exists in  $\text{Hilb}$  and is given by  $e_{\mathcal{S}}$ , the inclusion of the subspace  $\mathcal{S}$  spanned by  $\{|00\rangle, |11\rangle, |01\rangle + |10\rangle\}$  into  $\mathbb{C}^2 \otimes \mathbb{C}^2$ . ( $\mathcal{S}$  is called the symmetric tensor of  $\mathbb{C}^2$  with itself.) The coequalizer of  $\sigma$  with  $\text{id}$  is simply  $e_{\mathcal{S}}^{\dagger}$ . There is also an equalizer of  $\text{ad}_{\sigma}: M_4 \rightarrow M_4$  with  $\text{id}$  in  $\text{vN}$ , but curiously enough, it is given by  $e: M_3 \oplus \mathbb{C} \rightarrow M_4$ , where  $e(a, \lambda) = \text{ad}_{e_{\mathcal{S}}^{\dagger}}(a) + \text{ad}_{e_{\mathcal{S}}^{\dagger}}(\lambda)$  and  $\mathcal{A}$  is the subspace spanned by  $|01\rangle - |10\rangle$ . For the proof and similar results, see [35].

Assume  $\xi: \mathcal{C} \rightarrow M_4$  is some filter (i.e. quotient in the opposite category) with  $\text{ad}_{\sigma} \circ \xi = \xi$ . Show that for each  $b \in \mathcal{C}$ , we have  $\xi(b)\sigma = \sigma\xi(b)$ . From this, derive (perhaps in the same way as [35]), that  $\xi(b) = p_{\mathcal{A}}\xi(b)p_{\mathcal{A}} + p_{\mathcal{S}}\xi(b)p_{\mathcal{S}}$  for all  $b \in \mathcal{C}$ , where  $p_{\mathcal{A}} = e_{\mathcal{A}}e_{\mathcal{A}}^{\dagger}$  and  $p_{\mathcal{S}} = e_{\mathcal{S}}e_{\mathcal{S}}^{\dagger}$ . Prove that either  $\xi(b) \leq p_{\mathcal{A}}$  for all  $b \in \mathcal{C}$  or  $\xi(b) \leq p_{\mathcal{S}}$  for all  $b \in \mathcal{C}$ . Conclude  $\text{Pure}(\text{vN}^{\text{op}})$  does not have all coequalizers.

- VIII** To summarize: the category  $\text{Pure}(C)$  of a  $\dagger$ -effectus  $C$  is a  $\dagger$ -category, has  $\dagger$ -kernels and each  $\dagger$ -mono is a  $\dagger$ -kernel, but in general it does not have  $\dagger$ -biproducts or  $\dagger$ -equalizers, which shows it is not, in general, a pre-Hilbert category as in [49, 3.7.1].
- VIIIa** Recently Tull axiomatized [115] the  $\dagger$ -category  $\text{Hilb}_{\sim}$  of Hilbert spaces with bounded operators between them modulo global phase (i.e.  $T = S$  in  $\text{Hilb}_{\sim}$  iff  $T = \lambda S$  for some  $\lambda \in \mathbb{C}, |\lambda| = 1$ .) Write  $\text{Hilb}_{\sim}^c$  for the restriction of  $\text{Hilb}_{\sim}$  to contractive operators. This category  $\text{Hilb}_{\sim}^c$  is a full  $\dagger$ -subcategory of  $\text{Pure}(\text{vN}^{\text{op}})$  via the functor

$$\mathcal{H} \mapsto \mathcal{B}(\mathcal{H}), \quad (T: \mathcal{H} \rightarrow \mathcal{H}) \mapsto (\text{ad}_T: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})).$$

The category  $\text{Hilb}_{\sim}^c$  (and in fact  $\text{Hilb}_{\sim}$ ) do not have all (co)products. Tull shows that  $\text{Hilb}_{\sim}$  does have *phased biproducts*. [114, 115] This raises the question: do the phased biproducts of  $\text{Hilb}_{\sim}$  extend to  $\text{Pure}(\text{vN}^{\text{op}})$ . If this is the case, then a phased product of  $\mathbb{C}$  with itself in  $\text{Pure}(\text{vN}^{\text{op}})$  is given by:  $|0\rangle\langle 0|, |1\rangle\langle 1|: \mathbb{C} \rightarrow M_2$ . Unfortunately this leads to a contradiction. It hasn't been determined yet whether  $\text{Pure}(\text{vN}^{\text{op}})$  has phased products at all.

### 3.8.2 Sequential product

The sequential product — the operation  $a \& b \equiv \sqrt{ab}\sqrt{a}$  on the effects of a von Neumann algebra — has been studied before. In [37,47] Gudder and coauthors establish some basic algebraic properties of  $\&$  on  $\mathcal{B}(\mathcal{H})$ . Most of them are surprisingly hard to prove — for instance, the proof that  $a\&b = b\&a$  implies  $ab = ba$ , requires the Fuglede–Rosenblum–Putnam Theorem. 225

Then, in [46], Gudder and Latrémolière (G&L) characterize  $\&$  on  $\mathcal{B}(\mathcal{H})$  as the unique operation satisfying (for effects  $a, b$  and density matrix  $\varrho$ ) II

1.  $\text{tr}[(a \& \varrho)b] = \text{tr}[\varrho(a \& b)]$
2.  $a \& 1 = 1 \& a = a$
3.  $a \& (a \& b) = (a \& a) \& b = a^2 \& b$
4.  $a \mapsto a \& b$  is strongly continuous

In 105V we characterized  $\text{asrt}_a: b \mapsto a \& b$  as the unique  $\diamond$ -positive map with  $a \& 1 = a$ . How do these characterizations compare? G&L's first axiom plays a very similar role as the  $\diamond$ -self adjointness of  $\diamond$ -positive maps for us. Their third axiom is somewhat related to  $\diamond$ -positivity. Their fourth axiom seems unrelated to our characterization and conversely the purity of our  $\diamond$ -positive maps has no counterpart in their axioms.

A few years earlier, Gudder and Greechie started [47] the abstract study of some of the algebraic properties of the sequential product on arbitrary effect algebras, which has been picked up by several other authors [6, 44, 45, 71, 72, 83, 104, 111, 119, 120, 122]. III

**Definition** A **sequential effect algebra** (SEA) is an effect algebra  $E$  together with a binary operation  $\&$  satisfying IV

- (S1) if  $a \perp b$ , then  $c \& a \perp c \& b$  and  $(c \& a) \oslash (c \& b) = c \& (a \oslash b)$ ;
- (S2)  $1 \& a = a$ ;
- (S3) if  $a \& b = 0$ , then  $a \& b = b \& a$ ;
- (S4) if  $a \& b = b \& a$ , then  $a \& b^\perp = b^\perp \& a$  and  $(a \& b) \& c = a \& (b \& c)$  and
- (S5) if  $c \& a = a \& c$ ,  $c \& b = b \& c$  and  $a \perp b$ ,  
then  $c \& (a \& b) = (a \& b) \& c$  and  $c \& (a \oslash b) = (a \oslash b) \& c$ .

**Examples** The effect algebra  $[0, 1]_{\mathcal{A}}$  of effects on a von Neumann algebra  $\mathcal{A}$  is a sequential effect algebra with  $a \& b = \sqrt{ab}\sqrt{a}$ . Any commutative effect monoid is a sequential effect algebra with  $a \& b = a \odot b$ . V

**Proposition** In a  $\dagger$ -effectus, the set of predicates  $\text{Pred } X$  on any object  $X$  with  $p \& q = q \circ \text{asrt}_p$  satisfies axioms (S1), (S2) and (S3) of a SEA. VI

- VII **Proof** The proofs of (S1) and (S2) are obvious. To show (S3), assume  $p \& q = 0$  for some predicates  $p, q$ . Then  $1 \circ \text{asrt}_q \circ \text{asrt}_p = p \& q = 0$  and so  $\text{asrt}_q \circ \text{asrt}_p = 0 = \text{asrt}_0$ . Applying the dagger, we find  $0 = \text{asrt}_0^\dagger = \text{asrt}_p^\dagger \circ \text{asrt}_q^\dagger = \text{asrt}_p \circ \text{asrt}_q$ . Thus  $q \& p = 0 = p \& q$ , as desired.  $\square$
- VIII **Remark** It's unclear whether the set of predicates in a  $\dagger$ -effectus forms a sequential effect algebra. A seemingly related open problem is whether  $p \& q = q \& p$  implies that  $\text{asrt}_p \circ \text{asrt}_q = \text{asrt}_q \circ \text{asrt}_p$ .

### 3.8.3 Homological categories

226 Next we compare our effectuses to Grandis' homological categories [41], which generalize abelian categories used in homological algebra. First, we need a lemma.

- II **Lemma** Suppose  $s, t$  are any sharp predicates on the same object in a  $\dagger$ -effectus. If  $s^\perp \leq t$ , then  $s \& t$  is sharp.
- III **Proof** By 208 III, it is sufficient to show  $s \& t^\perp$  is sharp as  $(s \& t)^\perp = s \& t^\perp \otimes s^\perp$ . Note  $s^\perp \perp t^\perp$ , so by 213 III, we find  $t^\perp \& s^\perp = 0$ , hence  $t^\perp \& s = t^\perp$  is sharp. By the same reasoning as in the final part of the proof of 219 XI, there exists a (unique) pristine map  $l$  with  $\text{asrt}_s \circ \text{asrt}_{t^\perp} = l \circ \text{asrt}_{t^\perp \& s}$ ,  $l \circ l^\dagger = \text{asrt}_{[t^\perp \& s]}$  and  $l^\dagger \circ l = \text{asrt}_{[s \& t^\perp]}$ . It follows  $l \circ \text{asrt}_{t^\perp \& s} \circ l^\dagger = \text{asrt}_{s \& t^\perp}$  and so

$$\begin{aligned}
 \text{asrt}_{s \& t^\perp}^2 &= l \circ \text{asrt}_{t^\perp \& s} \circ l^\dagger \circ l \circ \text{asrt}_{t^\perp \& s} \circ l^\dagger \\
 &= l \circ \text{asrt}_{t^\perp \& s} \circ \text{asrt}_{[s \& t^\perp]} \circ \text{asrt}_{t^\perp \& s} \circ l^\dagger \\
 &\stackrel{211 \text{XV}}{=} l \circ \text{asrt}_{t^\perp \& s} \circ \text{asrt}_{t^\perp \& s} \circ l^\dagger \\
 &\stackrel{211 \text{V}}{=} l \circ \text{asrt}_{t^\perp \& s} \circ l^\dagger \\
 &= \text{asrt}_{s \& t^\perp},
 \end{aligned}$$

which shows  $s \& t^\perp$  is sharp.  $\square$

- IV **Definition** A category  $C$  is a **pointed semiexact category** [41, §1.1] if  $C$  has a zero object and all kernels and cokernels. In a pointed semiexact category:

1. We put a partial order on kernels in the usual way: we write  $n \leq m$  for kernels  $n: A \rightarrow X$  and  $m: B \rightarrow X$ , if there is an  $f: A \rightarrow B$  with  $n = m \circ f$ . If both  $n \leq m$  and  $m \leq n$ , then we write  $n \approx m$ . We denote the poset of kernels on  $A$  modulo  $\approx$  by  $\text{Nsb } A$ . (**N**ormal **s**ub**o**bjects, [41, §1.5].)
2. For every map  $f$  there is a unique map  $g$  with  $f = (\ker \text{cok } f) \circ g \circ (\text{cok } \ker f)$ . If this  $g$  is an iso, then  $f$  is called **exact**.

A pointed semiexact category  $C$  is a **pointed homological category** if

1. kernels are closed under composition;
2. cokernels are closed under composition *and*
3. (*homology axiom*) for any kernel  $m: M \rightarrow A$  and cokernel  $q: A \rightarrow Q$  with  $\ker q \leq m$ , the composition  $q \circ m$  is exact.

**Theorem** Any  $\dagger$ -effectus (in partial form) is a (pointed) homological category: V

1. a map is a kernel iff it is a comprehension;
2. a map is a cokernel iff it is a quotient of a sharp predicate *and*
3. a map is exact iff it is pristine.

**Proof** Let  $C$  be a  $\dagger$ -effectus. By 200 III and 205 II, we know  $C$  has all kernels and cokernels: a kernel of  $f$  is exactly a comprehension of  $(1 \circ f)^\perp$  and a cokernel of  $f$  is exactly a quotient of  $\text{im } f$ . By 211 XI, comprehensions are closed under composition, and so kernels are closed under composition as well. As for the cokernels, assume  $\zeta_1, \zeta_2$  are two composable cokernels. Both  $\zeta_1$  and  $\zeta_2$  are quotients of a sharp predicate. By 197 IX the composition  $\zeta_1 \circ \zeta_2$  is a quotient as well. The map  $\zeta_2$  is sharp by 197 IX and so the predicate  $1 \circ \zeta_1 \circ \zeta_2$  is sharp. Hence  $\zeta_1 \circ \zeta_2$  is a cokernel too. Unfolding definitions, it is easy to see exact maps correspond precisely to pristine maps. VI

(*Homology axiom*) We have to show  $q \circ m$  is exact for any cokernel  $q$  and kernel  $m$  with  $\ker q \leq m$ . As  $q \circ m$  is pure, it is sufficient to show  $1 \circ q \circ m$  is sharp. By 204 I, the assumption  $\ker q \leq m$  is equivalent to  $(1 \circ q)^\perp \equiv \text{im } \ker q \leq \text{im } m$  and so  $\text{im}^\perp m \leq 1 \circ q$ . Our lemma II shows  $(\text{im } m) \& (1 \circ q) \equiv 1 \circ q \circ m \circ m^\dagger$  is sharp and so VII

$$\begin{aligned}
 [1 \circ q \circ m] &= [1 \circ q \circ m] \circ m^\dagger \circ m && \text{as comprehensions are } \dagger\text{-mono} \\
 &= [1 \circ q \circ m \circ m^\dagger] \circ m && \text{as } m^\dagger \text{ is sharp} \\
 &= 1 \circ q \circ m \circ m^\dagger \circ m \\
 &= 1 \circ q \circ m,
 \end{aligned}$$

which show  $q \circ m$  is sharp, as desired. □

In any homological category, a generalization of the famous *Snake Lemma* holds. 227  
 Before we can discuss it, we need some more homological category theory.

**Definition** Let a (pointed) semiexact category be given. II

1. We say the diagram  $A \xrightarrow{f} B \xrightarrow{g} C$  is **exact at  $B$** , if  $\ker \text{cok}(f) \approx \ker(g)$ .  
 We say  $A_1 \xrightarrow{f_1} A_2 \rightarrow \cdots \rightarrow A_n$  is a **(long) exact sequence** if it is exact at  $A_2, \dots, A_{n-1}$ .

2. The poset  $\text{Nsb } A$  of kernels modulo  $\approx$  is a bounded lattice with minimum  $0$  and maximum  $1 \equiv \text{id}$ . [41, §1.5]
3. For any  $f: A \rightarrow B$ , define  $f_*: \text{Nsb } A \rightleftharpoons \text{Nsb } B: f^*$  by  $f_*(k) = \ker \text{cok}(f \circ k)$  and  $f^*(k) = \ker((\text{cok } k) \circ f)$ .
4. We say  $f$  is **left-modular at  $k$**  if  $f^*(f_*(k)) = k \vee f^*(0)$  and  $f$  is **right-modular at  $k$**  if  $f_*(f^*(k)) = k \wedge f_*(1)$ . We call  $f$  simply **left-modular** (resp. **right-modular**) if it is left-modular (resp. left-modular) at every  $k$ . We say  $f$  is **modular** if it is both left- and right-modular.

**III Example** In a  $\dagger$ -effectus, the previous concepts are related to our familiar notions as follows.

1. A diagram  $A \xrightarrow{f} B \xrightarrow{g} C$  is exact at  $B$  if and only if  $\text{im}^\perp f = [1 \circ f]$ .
2. The lattice  $\text{Nsb } A$  is isomorphic to the lattice  $\text{SPred } A$  via  $k \mapsto \text{im } k$ .
3. For any  $f$ , we have  $\text{im } f_*(k) = f_\diamond(\text{im } k)$  and  $\text{im } f^*(k) = f^\square(\text{im } k)$ .
4. A map  $f$  is left-modular at  $k$  iff  $f^\square(f_\diamond(\text{im } k)) = (\text{im } k) \vee [1 \circ f]$  and right-modular at  $k$  iff  $f_\diamond(f^\square(\text{im } k)) = (\text{im } k) \wedge \text{im } f$ .

**IV Notation** As  $\text{Nsb } A$  and  $\text{SPred } A$  are isomorphic, we will abbreviate “ $f$  is left-modular at  $\pi_s$ ” by “ $f$  is left-modular at  $s$ ” and similarly for right-modularity.

**V Lemma** In a  $\dagger$ -effectus, we have  $\pi^\square \circ \pi_\diamond = \text{id}$  and  $\zeta_\diamond \circ \zeta^\square = \text{id}$  for any comprehension  $\pi$  and sharp quotient  $\zeta$ .

**VI Proof** We start with  $\pi$ . The trick is to use  $\pi^\dagger$  is sharp:  $\pi^\square \circ \pi_\diamond(s) = \pi^\square \circ (\pi^\dagger)^\diamond(s) = [[s \circ \pi^\dagger]^\perp \circ \pi]^\perp = [(s \circ \pi^\dagger)^\perp \circ \pi]^\perp = [(s \circ \pi^\dagger \circ \pi)^\perp]^\perp = [s^\perp]^\perp = s$ . As for the other equality:  $\zeta_\diamond \circ \zeta^\square(s) = (\zeta^\dagger)^\diamond \circ \zeta^\square(s) = [[s^\perp \circ \zeta]^\perp \circ \zeta^\dagger] = [([s^\perp \circ \zeta] \circ \zeta^\dagger)^\perp] = [([s^\perp] \circ \zeta \circ \zeta^\dagger)^\perp] = [[s^\perp]^\perp] = s$ .  $\square$

**228** The following is a translation of (a part of) Grandis’ Snake Lemma [41, §3.4] into  $\dagger$ -effectus jargon. We include a proof as it highlights how the reasoning in a homological category is subtly different from the reasoning we already saw in a  $\dagger$ -effectus: maps  $l^\square$  appear, where we would have expected  $l^\diamond$ .

**II Snake Lemma** Suppose we have a commuting diagram in  $\dagger$ -effectus as follows.

$$\begin{array}{ccccccc}
 & & A & \xrightarrow{f} & B & \xrightarrow{g} & C & \xrightarrow{0} & 0 \\
 & & \downarrow a & & \downarrow b & & \downarrow c & & \\
 0 & \xrightarrow{0} & A' & \xrightarrow{h} & B' & \xrightarrow{k} & C' & & 
 \end{array}$$

Furthermore, assume

1. the diagram has exact rows;
2.  $b$  is left-modular over  $\text{im } f$ ;
3.  $b$  is right-modular over  $\text{im } h$ ;
4.  $f$  is right-modular over  $[1 \circ b]^\perp$  and
5.  $k$  is left-modular over  $\text{im } b$ .

These additional assumptions are equivalent to the following conditions.

1.  $b^\square \circ b_\circ(\text{im } f) = [1 \circ b]^\perp \vee \text{im } f$ ,
2.  $b_\circ \circ b^\square(\text{im } h) = (\text{im } h) \wedge \text{im } b$ ,
3.  $k^\square \circ k_\circ(\text{im } b) = (\text{im } h) \vee \text{im } b$ ,
4.  $f_\circ \circ f^\square \circ b^\square(0) = [1 \circ b]^\perp \wedge \text{im } f$ ,
5.  $\text{im}^\perp f = [1 \circ g]$ ,
6.  $\text{im}^\perp h = [1 \circ k]$ ,
7.  $g$  is a sharp quotient and
8.  $h$  is a comprehension.

Completing the diagram, write  $a_\pi, b_\pi, c_\pi$  for kernels of  $a, b$  and  $c$ , respectively;  $a_\zeta, b_\zeta, c_\zeta$  for the cokernels and  $\bar{f}, \bar{g}, \bar{h}, \bar{k}$  for the induced maps between the kernels and cokernels. It is easy to see

$$\bar{f} = b_\pi^\dagger \circ f \circ a_\pi \quad \bar{g} = c_\pi^\dagger \circ g \circ b_\pi \quad \bar{h} = b_\zeta \circ h \circ a_\zeta^\dagger \quad \bar{k} = c_\zeta \circ k \circ b_\zeta^\dagger.$$

Then: there is a map  $d: \ker c \rightarrow \text{cok } a$  that turns<sup>††</sup>

$$\ker a \xrightarrow{\bar{f}} \ker b \xrightarrow{\bar{g}} \ker c \xrightarrow{d} \text{cok } a \xrightarrow{\bar{h}} \text{cok } b \xrightarrow{\bar{k}} \text{cok } c \quad (3.19)$$

into a (long) exact sequence. This reveals the snake:

$$\begin{array}{ccccccc}
 \ker a & \xrightarrow{\bar{f}} & \ker b & \xrightarrow{\bar{g}} & \ker c & & \\
 \downarrow a_\pi & & \downarrow b_\pi & & \downarrow c_\pi & & \\
 A & \xrightarrow{f} & B & \xrightarrow{g} & C & \xrightarrow{0} & 0 \\
 \downarrow a & & \downarrow b & & \downarrow c & & \\
 0 & \xrightarrow{0} & A' & \xrightarrow{h} & B' & \xrightarrow{k} & C' \\
 \downarrow a_\zeta & & \downarrow b_\zeta & & \downarrow c_\zeta & & \\
 \text{cok } a & \xrightarrow{\bar{h}} & \text{cok } b & \xrightarrow{\bar{k}} & \text{cok } c & & 
 \end{array}$$

$\left. \begin{array}{l} \text{---} \\ \text{---} \\ \text{---} \end{array} \right\} d$

<sup>††</sup>Beware: we use  $\ker a$  (and  $\text{cok}$ ) inconsistently — earlier in the text  $\ker f$  refers to a kernel map — here it refers to a kernel object instead.

- III **Proof** Before we start, we give an overview of the proof. We will first show there are comprehensions  $m, h'$  and sharp quotients  $g', v$  such that the left and right faces of the following cube commute.

$$\begin{array}{ccccc}
 & & B & \xrightarrow{b} & B' \\
 & \nearrow m & \downarrow g & & \nearrow h \\
 Z & & & & A' \\
 \downarrow g' & & \downarrow c_\pi & & \downarrow a_\zeta \\
 \ker c & \xrightarrow{d} & \text{cok } a & & Q \\
 & & \downarrow h' & & \\
 & & & & 
 \end{array}$$

(These two faces are, what Grandis calls, subquotients.) Then we will prove using exactness of the rows that there is a unique lifting of  $b$  to  $b'$ :  $Z \rightarrow A'$  along the comprehensions  $m, h$ . In turn,  $d$  is defined as the unique lifting of  $b$  along the sharp quotients  $h', a_\zeta$ . ( $d$  is the map regularly induced by  $b$  along the two subquotients.) Before demonstrating the exactness of the snake, we need some non-trivial identities for  $d_\diamond$  and  $d^\square$ . (These also follow from Grandis' calculus of direct and inverse images along an induced morphism.)

- IV Define  $v \equiv \xi_{h_\diamond(\text{im } a)}$ . As  $(v \circ h)^\diamond(1) = h^\diamond(h_\diamond(\text{im } a)) = \text{im } a = 1 \circ a_\zeta$ , there must be a total pure map  $h'$  such that  $v \circ h = h' \circ a_\zeta$ . Note  $[1 \circ v]^\perp = h_\diamond(\text{im } a) \leq h_\diamond(1) = \text{im } h$ . So by the homology axiom, we know  $v \circ h$  is pristine and so  $h'$  must be a comprehension. Moving to the other side, define  $m \equiv \pi_{g^\square([1 \circ c]^\perp)}$ . Reasoning in a similar way, we see there is a unique sharp quotient  $g'$  with  $g \circ m = c_\pi \circ g'$ .
- V To show  $b$  lifts along  $m$  and  $h$ , it suffices (by the universal property of comprehensions) to show  $(b \circ m)_\diamond(1) \equiv \text{im } b \circ m \leq \text{im } h \equiv h_\diamond(1)$ .

$$\begin{aligned}
 (b \circ m)_\diamond(1) &= (b_\diamond \circ g^\square \circ c^\square)(0) && \text{by dfn. } m \\
 &= (b_\diamond \circ b^\square \circ k^\square)(0) \\
 &= (b_\diamond \circ b^\square \circ h_\diamond)(1) && \text{by exactness at } B' \\
 &\leq h_\diamond(1) && \text{as } b_\diamond \dashv b^\square.
 \end{aligned}$$

So there is a unique  $b': Z \rightarrow A'$  with  $h \circ b' = b \circ m$ . Before we continue, we note  $b' = h^\dagger \circ b \circ m$  (as  $h^\dagger \circ h = \text{id}$ ) and so surprisingly:

$$m^\square \circ b^\square \circ h_\square = b'^{\square 227} \equiv b'^\square \circ h^\square \circ h_\diamond = m^\square \circ b^\square \circ h_\diamond. \quad (3.20)$$

Next, to show  $b'$  lifts along  $g'$  and  $a_\zeta$ , it is sufficient to prove  $(b'^\square \circ a_\zeta^\square)(0) = [1 \circ b' \circ a_\zeta]^\perp \geq [1 \circ g']^\perp = g'^\square(0)$ .

$$\begin{aligned}
(b'^\square \circ a_\zeta^\square)(0) &= (b'^\square \circ a_\diamond)(1) && \text{by dfn. } a_\zeta \\
&= (m^\square \circ b^\square \circ h_\diamond \circ a_\diamond)(1) && \text{by (3.20)} \\
&= (m^\square \circ b^\square \circ b_\diamond \circ f_\diamond)(1) \\
&\geq (m^\square \circ f_\diamond)(1) && \text{as } b_\diamond \dashv b^\square \\
&= (m^\square \circ g_\square)(0) && \text{by exactness at } B \\
&= (g'^\square \circ (c_\pi)_\square)(0) && \text{by dfn. } g' \\
&\geq g'^\square(0).
\end{aligned}$$

Thus there is a unique  $d: \ker c \rightarrow \text{cok } a$  with  $d \circ g' = a_\zeta \circ b'$ . More concretely, using  $g' \circ g'^\dagger = \text{id}$ , we see  $d = a_\zeta \circ h^\dagger \circ b \circ m \circ g'^\dagger$ .

Our next goal is to show

VI

$$\begin{aligned}
d_\diamond &= (a_\zeta)_\diamond \circ h^\square \circ b_\diamond \circ m_\diamond \circ g'^\square = h'^\square \circ v_\diamond \circ b_\diamond \circ m_\diamond \circ g'^\square && (3.21) \\
&= (a_\zeta)_\diamond \circ h^\square \circ b_\diamond \circ g'^\square \circ (c_\pi)_\diamond = h'^\square \circ v_\diamond \circ b_\diamond \circ g'^\square \circ (c_\pi)_\diamond.
\end{aligned}$$

As a first step, note  $\text{im } m = (g^\square \circ c_\diamond)(1) \geq (g^\square \circ (c_\pi)_\diamond)(s)$  and so

$$\begin{aligned}
(g^\square \circ (c_\pi)_\diamond)(s) &= (g^\square \circ (c_\pi)_\diamond(s)) \wedge (\text{im } m) \\
&= (m_\diamond \circ m^\square \circ g^\square \circ (c_\pi)_\diamond)(s) && \text{by 208 IX} \\
&= (m_\diamond \circ g'^\square \circ c_\pi^\square \circ (c_\pi)_\diamond)(s) && \text{by dfn. } g' \\
&= (m_\diamond \circ g'^\square)(s) && \text{by 227 V.}
\end{aligned}$$

and so we only have to show the first two equalities of (3.21). The first is easy:

$$\begin{aligned}
d_\diamond &= d_\diamond \circ g'_\diamond \circ g'^\square && \text{by 227 V} \\
&= (a_\zeta)_\diamond \circ b'_\diamond \circ g'^\square && \text{by dfn. } d \\
&= (a_\zeta)_\diamond \circ h^\square \circ h_\diamond \circ b'_\diamond \circ g'^\square && \text{by 227 V} \\
&= (a_\zeta)_\diamond \circ h^\square \circ b_\diamond \circ m_\diamond \circ g'^\square && \text{by dfn. } b'.
\end{aligned}$$

The second is trickier. We need some preparation. Recall  $v^\square(0) \leq h_\diamond(1)$  and so

$$h_\diamond(1) = v^\square(0) \vee h_\diamond(1) \stackrel{208 \text{ XII}}{=} (v^\square \circ v_\diamond \circ h_\diamond)(1) = (v^\square \circ h'_\diamond)(1). \quad (3.22)$$

As  $h_\diamond \circ a_\zeta^\square$  is injective, the last equality follows from

$$\begin{aligned}
& (h_\diamond \circ a_\zeta^\square \circ (a_\zeta)_\diamond \circ h^\square \circ b_\diamond \circ m_\diamond \circ g'^\square)(s) \\
&= (h_\diamond(h^\square \circ b_\diamond \circ m_\diamond \circ g'^\square)(s) \vee a_\zeta^\square(0)) && \text{by 208 XII} \\
&= (h_\diamond \circ h^\square \circ b_\diamond \circ m_\diamond \circ g'^\square)(s) \vee (h_\diamond \circ a_\zeta^\square)(0) && \text{as } h_\diamond \dashv h^\square \\
&= (h_\diamond \circ h^\square \circ b_\diamond \circ m_\diamond \circ g'^\square)(s) \vee v^\square(0) && \text{by dfns. } v, a_\zeta \\
&= ((b_\diamond \circ m_\diamond \circ g'^\square)(s) \wedge h_\diamond(1)) \vee v^\square(0) && \text{by 208 IX} \\
&= (b_\diamond \circ m_\diamond \circ g'^\square)(s) \vee v^\square(0) && \text{as } b_\diamond \circ m_\diamond(1) \leq h_\diamond(1) \\
&= ((b_\diamond \circ m_\diamond \circ g'^\square)(s) \vee v^\square(0)) \wedge h_\diamond(1) && \text{as } v^\square(0) \leq h_\diamond(1) \\
&= (v^\square \circ v_\diamond \circ b_\diamond \circ m_\diamond \circ g'^\square)(s) \wedge h_\diamond(1) && \text{by 208 XII} \\
&= v^\square \circ v_\diamond \circ b_\diamond \circ m_\diamond \circ g'^\square(s) \wedge v^\square \circ h'_\diamond(1) && \text{by (3.22)} \\
&= v^\square((v_\diamond \circ b_\diamond \circ m_\diamond \circ g'^\square)(s) \wedge h'_\diamond(1)) && \text{as } v^\square \vdash v_\diamond \\
&= (v^\square \circ h'_\diamond \circ h'^\square \circ v_\diamond \circ b_\diamond \circ m_\diamond \circ g'^\square)(s) && \text{by 208 IX} \\
&= (h_\diamond \circ a_\zeta^\square \circ h'^\square \circ v_\diamond \circ b_\diamond \circ m_\diamond \circ g'^\square)(s),
\end{aligned}$$

where  $h_\diamond \circ a_\zeta^\square \circ v^\square \circ h'_\diamond$  is proven in the same way as  $g^\square \circ (c_\pi)_\diamond = m_\diamond \circ g'^\square$ .

VII We are ready to show exactness of (3.19). We will first derive exactness in  $\text{cok } a$  from the right-modularity of  $b$  over  $\text{im } h$ .

$$\begin{aligned}
d_\diamond(1) &= ((a_\zeta)_\diamond \circ h^\square \circ b_\diamond \circ g^\square \circ (c_\pi)_\diamond)(1) && \text{by (3.21)} \\
&= ((a_\zeta)_\diamond \circ h^\square \circ b_\diamond \circ g^\square \circ c^\square)(0) && \text{by dfn. } c_\pi \\
&= ((a_\zeta)_\diamond \circ h^\square \circ b_\diamond \circ b^\square \circ k^\square)(0) \\
&= ((a_\zeta)_\diamond \circ h^\square \circ b_\diamond \circ b^\square \circ h_\diamond)(1) && \text{by exactness in } B' \\
&= ((a_\zeta)_\diamond \circ h^\square)(b_\diamond(1) \wedge h_\diamond(1)) && \text{by right-modularity} \\
&= ((a_\zeta)_\diamond \circ h^\square \circ h_\diamond \circ h^\square \circ b_\diamond)(1) && \text{by 208 IX} \\
&= ((a_\zeta)_\diamond \circ h^\square \circ b_\diamond)(1) && \text{as } h^\square \vdash h_\diamond \\
&= ((a_\zeta)_\diamond \circ h^\square \circ b_\zeta^\square)(0) && \text{by dfn. } b_\zeta \\
&= ((a_\zeta)_\diamond \circ a_\zeta^\square \circ \overline{h}^\square)(0) && \text{by dfn. } \overline{h} \\
&= \overline{h}^\square(0) && \text{by 227 V.}
\end{aligned}$$

By a dual argument one derives exactness of (3.19) in  $\text{ker } c$  from the left-modularity of  $b$  over  $\text{im } f$ .

To show exactness of (3.19) in  $\text{cok } b$ , we will use left-modularity of  $k$  in  $\text{im } b$ : VIII

$$\begin{aligned}
\bar{h}_\diamond(1) &= (\bar{h}_\diamond \circ (a_\zeta)_\diamond \circ a_\zeta^\square)(1) && \text{by 227 V} \\
&= (\bar{h}_\diamond \circ (a_\zeta)_\diamond)(1) && \text{as } a_\zeta^\square \dashv (a_\zeta)_\diamond \\
&= ((b_\zeta)_\diamond \circ h_\diamond)(1) && \text{by dfn. } \bar{h} \\
&= ((b_\zeta)_\diamond \circ b_\zeta^\square \circ (b_\zeta)_\diamond \circ h_\diamond)(1) && \text{as } (b_\zeta)_\diamond \dashv b_\zeta^\square \\
&= (b_\zeta)_\diamond(h_\diamond(1) \vee b_\zeta^\square(0)) && \text{by 208 XII} \\
&= (b_\zeta)_\diamond(h_\diamond(1) \vee b_\diamond(1)) && \text{by dfn. } b_\zeta \\
&= (b_\zeta)_\diamond(k^\square(0) \vee b_\diamond(1)) && \text{exactness in } B' \\
&= ((b_\zeta)_\diamond \circ k^\square \circ k_\diamond \circ b_\diamond)(1) && \text{by left-modularity} \\
&= ((b_\zeta)_\diamond \circ k^\square \circ c_\diamond \circ g_\diamond)(1) \\
&= ((b_\zeta)_\diamond \circ k^\square \circ c_\diamond)(1) && \text{as quotients are faithful} \\
&= ((b_\zeta)_\diamond \circ k^\square \circ c_\zeta^\square)(0) && \text{by dfn. } c_\zeta \\
&= ((b_\zeta)_\diamond \circ b_\zeta^\square \circ \bar{k}^\square)(0) && \text{by dfn. } \bar{k} \\
&= \bar{k}^\square(0) && \text{by 227 V.}
\end{aligned}$$

Finally, the exactness of (3.19) in  $\text{ker } b$  is shown with a dual argument using the right-modularity of  $f$  over  $[1 \circ b]^\perp$ . □

**Remark** As  $\text{vN}^{\text{op}}$  is a  $\dagger$ -effectus, Grandis' Snake Lemma also holds for von Neumann algebras. This is somewhat puzzling for the following reason. To study spaces, one associates in algebraic geometry to each space an object (or even a diagram of objects) of an abelian category representing some algebraic invariant. The key point is that it is easier to compute and understand these algebraic invariants than it is to work with the spaces themselves. One normally thinks of a category of spaces as being quite different from a category of algebraic invariants, hence it's surprising that  $\text{vN}^{\text{op}}$ , which is a rather complicated category of (non-commutative) spaces, behaves in this way, similar to an abelian category. IX



# Chapter 4

## Outlook

For my closing remarks, I would like to speculate on possible applications and suggest possible directions for future research. 229

We started this thesis by asking the question whether the incredibly useful Stinespring dilation theorem extends to any ncp-map between von Neumann algebras. Now that we have seen it does, the next question is obvious: do the applications that make the Stinespring dilation so useful also extend to the Paschke dilation? I want to draw attention to one application in particular: proving security bounds on quantum protocols. Here one uses a continuous version of the Stinespring dilation theorem [80]. Is there also such a continuous version for Paschke dilation?

Next, we have seen that self-dual Hilbert  $C^*$ -modules over von Neumann algebras are very well-behaved compared to arbitrary Hilbert  $C^*$ -modules. Frankly, I'm surprised they haven't been studied more extensively before: the way results about Hilbert spaces generalize elegantly to self-dual Hilbert  $C^*$ -modules over von Neumann algebras seems hard to ignore. Only one immediate open question remains here: does the normal form of 162 IV extend in some way for to arbitrary von Neumann algebras? II

As announced, in the second part of this thesis, we did not reach our goal of axiomatizing the category of von Neumann algebras categorically. In our attempt, we did find several new concepts such as  $\diamond$ -adjointness,  $\diamond$ -positivity, purity defined with quotient and comprehension and the  $\dagger$  on these pure maps. Building on this work, van de Wetering recently announced [117, 118] a reconstruction of finite-dimensional quantum theory. To discuss it, we need a definition: call an effectus *operational* if its scalars are isomorphic to the real interval; both the states and the predicates are order separating; all predicate effect modules are embeddable in finite-dimensional order unit spaces *and* each state space is a closed subset of the base norm space of all unital positive functionals on the III

corresponding order unit space of predicates. It follows from [118] that the state space of any operational  $\&$ -effectus is a spectral convex set in the sense of Alfsen and Shultz and that any operational  $\dagger$ -effectus is equivalent to a subcategory of the category of Euclidean Jordan Algebras with positive maps between them in the opposite direction. Are there infinite-dimensional generalizations of these results? Are the state spaces of a real  $\&$ -effectuses perhaps spectral convex sets? Is any real  $\dagger$ -effectus equivalent to a subcategory of, say, the category of JBW-algebras with positive maps between them in the opposite direction?

# Index

- $\&$ , andthen, 211 II
  - $[0, 1]_{\mathcal{A}}$ , 175 II
  - $\{X|p\}$ , 199 II
  - !, initial/final map, 180 I
  - $X/p$ , 197 II
  - $\dagger$ , dagger
    - category, 214 I
    - effectus, 215 I
    - epi, 224 II
    - in a  $\dagger$ -effectus, 217 II
    - kernel, 224 II
      - category, 224 II
    - mono, 224 II
    - partial isometry, 224 II
    - positive, 214 I
    - self-adjoint, 214 I
    - unitary, 214 I
  - $\diamond$ , diamond
    - adjoint, 206 II
    - effectus, 206 II
    - equivalent, 206 II
    - positive, 206 II
    - self-adjoint, 206 II
  - $\hat{\varepsilon}$ , 150 VI
  - $\tilde{\varepsilon}$ , 150 VIII
  - $\lfloor \cdot \rfloor$ , floor, 203 I
  - $\lceil \cdot \rceil$ , ceiling, 203 I
  - $\hat{f}$ 
    - in an effectus, 186 I
  - $|x\rangle\langle y|$  for Hilbert  $\mathcal{B}$ -modules, 159 II
  - $\lambda \cdot p$ , 190 II
  - $\langle \cdot, \cdot \rangle$ 
    - in an effectus, 181 VII
    - $\mathcal{B}$ -valued, 141 II
  - $\leq$ 
    - among kernels, 226 IV
    - in PCM/EA, 174 II
  - $\lesssim$ , 162 I
  - $\nabla$ , 187 III
  - $\odot$ 
    - effect monoid, 178 II
    - vector spaces, 137 II
  - $\ominus$ , 176 I
  - $( )_{\diamond}, ( )^{\diamond}$ , 206 II
  - $( )_{\square}, ( )^{\square}$ , 206 II
  - $( )_{*}, ( )^{*}$ , 227 II
  - $( )^{\perp}$ 
    - element effect algebra, 175 I
    - subset Hilbert  $\mathcal{B}$ -module, 160 III
  - $\pi_p$ , 199 IV, 211 IX
  - $\triangleright_i$ , 180 VII
  - $\rightarrow$ , 180 I
  - $\otimes$ 
    - Hilbert  $\mathcal{B}$ -modules, 164 II
    - module maps, 165 III
  - $\otimes_{\varphi}$ , 154 III
  - $\xi_p$ , 197 III
  - $\zeta_s$ , 211 IX
  - $\langle \cdot, \cdot \rangle_f$ 
    - $\mathcal{B}$ -module, 142 II
  - $[\cdot, \cdot]$ 
    - cotupling, 180 I
    - inner product, 136 II, 137 II
    - $\mathcal{B}$ -valued, 146 VII, 148 V, 150 II, 150 XI, 150 XIII, 163 II
  - $[0, \varphi]_{\text{nep}}$ , 157 II
  - $\ell^2((p_i)_{i \in I})$ , 161 II
  - $\mathcal{B}^a(X)$ , 143 I
  - $\leq_{\text{nep}}$ , 157 II, 198 II
  - $\varphi_t$ , 157 III
  - $\text{ad}_V$ 
    - Hilbert  $\mathcal{B}$ -modules, 153 I
- A**
- $\text{AConv}_M$ , 192 IV
  - $\text{ad}_V$ 
    - Hilbert spaces, 135 IV
  - adjointable, 143 I

$M$ -affine, 192 IV  
asrt <sub>$p$</sub> , assert, 211 II

## B

basically disconnected, 195 VI  
(orthonormal) basis, 149 I  
Bessel's inequality, 149 VIII  
bound, 187 III

## C

cancellative  
     $M$ -convex set, 192 IV  
Cauchy net, 147 I  
    equivalent, 147 I  
    fast, 150 V  
Cauchy sequence  
    fast, 136 III  
CH, 189a II  
 $\varphi$ -compatible, 154 II  
comprehension, 199 II  
    corresponding, 211 IX  
congruence  
     $M$ -convex set, 193 II  
(abstract)  $M$ -convex set, 192 IV  
(formal)  $M$ -convex combination, 192 II  
corner, 169 II  
    standard, 169 IV  
CRng, 189a II  
CvN, 189a I

## D

$\mathcal{D}_M$ , 192 II  
dilation  
    in an effectus, 221 II  
D-poset, 176 IV

## E

EA, 175 I  
effect algebra, 175 I  
    convex, 179 III  
    homomorphism, 175 I  
    sequential-, 225 IV  
    sub-, 175 I  
effect divisoid, 195 II

effect module, 179 II  
    homomorphism, 179 II  
effect monoid, 178 II  
    commutative, 178 II  
    homomorphism, 178 II  
effects, 175 II  
effectus, 189 II  
     $\dagger'$ -, 215 V  
     $\&$ -, 211 II  
     $\diamond$ -, 206 II, 215 I  
    in partial form, 180 VII  
    in total form, 180 I  
    operational, 229 III  
    real, 190 II  
    with comprehension, 199 II  
    with dilations, 221 II  
        and order correspondence, 223 V  
    with images, 202 I  
    with quotients, 197 II  
EMod <sub>$M$</sub> , 179 II  
entourages, 146 II  
exact, 226 IV  
    at, 227 II  
    sequence, 227 II

## F

cok  $f$ , 200 II  
faithful, 202 I  
filter, 169 VIII  
    standard, 169 X

## H

Hilbert  $\mathcal{B}$ -module, 141 II  
    pre-, 141 II  
    self-dual, 141 IIa  
homological category  
    pointed, 226 IV

## I

im, im<sup>⊥</sup>, 202 I  
inner product  
     $\mathcal{B}$ -valued, 141 II  
    definite, 141 II

Inv  $f$ , 223 II

## K

$\kappa_i$ , coprojection, 180 I

ker  $f$ , 200 II

kernel

†-, 224 II

category theoretic, 200 II

Kraus operators, 138 VIII

## M

$M$ , 190 II

map

ncp(u)-, 135 II

nmiu-, 135 II

modular, 227 II

## N

ncp-extreme, 172 II

Nsb  $A$ , 226 IV

## O

orthocomplement

in an effect algebra, 175 I

w.r.t. a Hilbert  $\mathcal{B}$ -module, 160 III

orthogonal, 149 I

ortholattice, 177 IV

orthomodular

lattice, 177 IV

orthonormal, 149 I

OUG, 189a II

OUS, 189a II

## P

$\wp$ , power set, 146 II

Par  $C$ , 180 VI

Parseval's identity, 149 IV

partial map, 180 I

partial projectors, 180 VII

Paschke dilation, 140 II

PCM, 174 II

homomorphism, 174 II

polar decomposition, 149 VIII

polarization identity, 142 IX

Pred( $f$ ), 190 II

predicate, 190 II

Pred  $X$ , 180 VII

pristine, 218 IV

pullback lemma, 183 III

pure

in an effectus, 201 II

ncp-map, 170 I

Pure  $C$ , 211 XVI

## Q

quotient, 197 II

corresponding, 211 IX

## R

Rng, 191 VIII

## S

scalar, 190 II

Scal  $C$ , 190 II

SEA, 225 IV

seff $_p$ , 223 II

semiexact category

pointed, 226 IV

separating

predicates, 190 II

states, 190 II

sesquilinear form

$\mathcal{B}$ -valued, 142 VII

bounded, 152 II

Set, 189a II

sharp, 203 I

map, 210 I

order-, 208 I

SPred  $X$ , 203 I

state, 190 II

sub-, 190 II

Stat  $X$ , 190 II

Stinespring dilation, 139 I

minimal, 139 I

## T

tensor product

of Hilbert  $\mathcal{B}$ -modules, 164 II

of module maps, 165 III  
total map, 180 VII

## U

ultranorm, 146 VII  
ultrastrong, 146 V  
ultraweak, 146 V  
uniform space, 146 II  
    complete, 147 I  
    continuous, 147 I  
    convergence, 147 I  
    dense, 147 I  
    Hausdorff, 146 II  
    uniformly continuous, 147 I  
unlim, 151 II  
uslim, 149 VII  
uwlim, 149 VII

## V

$vN$ , 180 V

# Bibliography

- [1] Creative commons attribution 4.0 international public license. <https://creativecommons.org/licenses/by/4.0/legalcode>.
- [2] Samson Abramsky and Bob Coecke. A categorical semantics of quantum protocols. In *LICS 2004*, pages 415–425. IEEE. arXiv:quant-ph/0402130.
- [3] Robin Adams and Bart Jacobs. A type theory for probabilistic and Bayesian reasoning. 2015. arXiv:1511.09230.
- [4] Erik M. Alfsen and Frederic W. Shultz. *State spaces of operator algebras*. 2001. doi:10.1007/978-1-4612-0147-2.
- [5] Michael A. Arbib and Ernest G. Manes. Partially additive categories and flow-diagram semantics. *Journal of Algebra*, 62(1):203–227, 1980. doi:10.1016/0021-8693(80)90212-4.
- [6] Alvaro Arias and Stanley P. Gudder. Almost sharp quantum effects. *Journal of Mathematical Physics*, 45(11):4196–4206, 2004. doi:10.1063/1.1806532.
- [7] William Arveson. *An invitation to  $C^*$ -algebras*, volume 39. Springer, 2012. doi:10.1007/978-1-4612-6371-5.
- [8] William B. Arveson. Subalgebras of  $C^*$ -algebras. *Acta Mathematica*, 123(1):141–224, 1969. doi:10.1007/BF02392388.
- [9] Steve Awodey. *Category theory*. Oxford University Press, 2010.
- [10] Michael Barr and Charles Wells. *Toposes, triples and theories*, volume 278. Springer-Verlag New York, 1985.
- [11] Garrett Birkhoff and John von Neumann. The logic of quantum mechanics. *Annals of Mathematics*, pages 823–843, 1936. doi:10.2307/1968621.
- [12] David P Blecher, Christian Le Merdy, et al. *Operator algebras and their modules: an operator space approach*. Number 30. Oxford University Press, 2004.
- [13] Costin Bădescu and Prakash Panangaden. Quantum alternation: Prospects and problems. In *QPL 2015*, volume 195 of *EPTCS*, pages 33–42. Open Publishing Association, 2015. doi:10.4204/EPTCS.195.3.

- [14] Mark S. Burgin. Categories with involution, and correspondences in  $\gamma$ -categories. *Trudy Moskovskogo Matematicheskogo Obshchestva*, 22:161–228, 1970.
- [15] Aurelio Carboni, Stephen Lack, and Robert F.C. Walters. Introduction to extensive and distributive categories. *Journal of Pure and Applied Algebra*, 84(2):145–158, 1993. doi:10.1016/0022-4049(93)90035-R.
- [16] Giulio Chiribella. Distinguishability and copiability of programs in general process theories. *Int J. Software Informatics*, 1(2), 2014. arXiv:1411.3035.
- [17] Giulio Chiribella, Giacomo Mauro D’Ariano, and Paolo Perinotti. Probabilistic theories with purification. *Physical Review A*, 81(6):062348, 2010. doi:10.1103/PhysRevA.81.062348.
- [18] Giulio Chiribella, Giacomo Mauro D’Ariano, and Paolo Perinotti. Informational derivation of quantum theory. *Physical Review A*, 84(1):012311, 2011. doi:10.1103/PhysRevA.84.012311.
- [19] Kenta Cho. Total and partial computation in categorical quantum foundations. In *QPL 2015*, volume 195 of *EPTCS*, pages 116–135, 2015. doi:10.4204/EPTCS.195.9 arXiv:1511.01569v1.
- [20] Kenta Cho and Bart P.F. Jacobs. The EfProb library for probabilistic calculations. In *CALCO 2017*, volume 72 of *LIPICs*, pages 25:1–25:8, 2017. doi:10.4230/LIPICs.CALCO.2017.25.
- [21] Kenta Cho and Bart P.F. Jacobs. Disintegration and Bayesian inversion via string diagrams. *MSCS*, 2019. doi:10.1017/S0960129518000488 arXiv:1709.00322v3.
- [22] Kenta Cho, Bart P.F. Jacobs, Abraham A. Westerbaan, and Bas E. Westerbaan. Quotient-comprehension chains. In *QPL 2015*, volume 195 of *EPTCS*, pages 136–147, 2015. doi:10.4204/EPTCS.195.10 arXiv:1511.01570v1.
- [23] Kenta Cho, Bart P.F. Jacobs, Bas E. Westerbaan, and Abraham A. Westerbaan. An introduction to effectus theory. arXiv:1512.05813v1, 2015.
- [24] Man-Duen Choi. A Schwarz inequality for positive linear maps on  $C^*$ -algebras. *Illinois Journal of Mathematics*, 18(4):565–574, 1974. doi:10.1215/ijm/1256051007.
- [25] Bob Coecke and Aleks R. Kissinger. *Picturing quantum processes*. 2017. doi:10.1017/9781316219317.

- [26] Oscar Cunningham and Chris J.M. Heunen. Purity through factorisation. In *QPL 2017*, volume 266 of *EPTCS*, pages 315–328, 2017. doi:10.4204/EPTCS.266.20 arXiv:1705.07652.
- [27] Giacomo Mauro D’Ariano, Giulio Chiribella, and Paolo Perinotti. *Quantum Theory from First Principles*. Cambridge University Press, 2016. doi:10.1017/9781107338340.
- [28] Edward Brian Davies. *Quantum theory of open systems*. Academic Press London, 1976.
- [29] Anatolij Dvurecenskij and Sylvia Pulmannová. *New trends in quantum structures*, volume 516. Springer, 2013. doi:10.1007/978-94-017-2422-7.
- [30] Joe Flood. Semiconvex geometry. *Journal of the Australian Mathematical Society*, 30(4):496–510, 1981. doi:10.1017/S1446788700017973.
- [31] David J. Foulis and Mary K. Bennett. Effect algebras and unsharp quantum logics. *Foundations of Physics*, 24(10):1331–1352, 1994. doi:10.1007/BF02283036.
- [32] Michael Frank. Self-duality and  $C^*$ -reflexivity of Hilbert  $C^*$ -moduli. *Zeitschrift für Analysis und ihre Anwendungen*, 9(2):165–176, 1990. doi:10.4171/ZAA/390.
- [33] Tobias Fritz. Convex spaces I: Definition and examples. 2009. arXiv:0903.5522.
- [34] Robert W.J. Furber and Bart P.F. Jacobs. From Kleisli categories to commutative  $C^*$ -algebras: probabilistic Gelfand duality. In *CALCO 2013*, volume 8089 of *LNCS*, pages 141–157, 2013. doi:10.1007/978-3-642-40206-7\_12.
- [35] Robert W.J. Furber and Bas E. Westerbaan. Unordered tuples in quantum computation. In *QPL 2015*, *EPTCS*, pages 196–207, 2015. doi:10.4204/EPTCS.195.15.
- [36] Israel Gelfand and Mark Neumark. On the imbedding of normed rings into the ring of operators in Hilbert space. *Rec. Math. [Mat. Sbornik] N.S.*, 12(2):197–217, 1943. doi:10.1090/conm/167/16.
- [37] Aurelian Gheondea and Stanley P. Gudder. Sequential product of quantum effects. *Proceedings of the American Mathematical Society*, 132(2):503–512, 2004. doi:10.2307/1194082.
- [38] Paul Ghez, Ricardo Lima, and John Roberts.  $W^*$ -categories. *Pacific Journal of Mathematics*, 120(1):79–109, 1985.

- [39] Leonard Gillman and Meyer Jerison. *Rings of continuous functions*. Springer, 2013. doi:10.1007/978-1-4615-7819-2.
- [40] Stefano Gogioso and Carlo Maria Scandolo. Categorical probabilistic theories. In *QPL 2017*, volume 266 of *EPTCS*, pages 367–385, 2017. doi:10.4204/EPTCS.266.23 arXiv:1701.08075.
- [41] Marco Grandis. On the categorical foundations of homological and homotopical algebra. *Cahiers de Topologie et Géométrie Différentielle Catégoriques*, 33(2):135–175, 1992.
- [42] Stanley P. Gudder. A general theory of convexity. *Milan Journal of Mathematics*, 49(1):89–96, 1979. doi:10.1007/BF02925185.
- [43] Stanley P. Gudder. Convex structures and effect algebras. *International Journal of Theoretical Physics*, 38(12):3179–3187, 1999. doi:10.1023/A:1026678114856.
- [44] Stanley P. Gudder. Open problems for sequential effect algebras. *International Journal of Theoretical Physics*, 44(12):2199–2206, 2005. doi:10.1007/s10773-005-8015-1.
- [45] Stanley P. Gudder and Richard J. Greechie. Uniqueness and order in sequential effect algebras. *International Journal of Theoretical Physics*, 44(7):755–770, 2005. doi:10.1007/s10773-005-7054-y.
- [46] Stanley P. Gudder and Frédéric Latrémolière. Characterization of the sequential product on quantum effects. *Journal of Mathematical Physics*, 49(5):052106, 2008. doi:10.1063/1.2904475.
- [47] Stanley P. Gudder and Gabriel Nagy. Sequential quantum measurements. *Journal of Mathematical Physics*, 42(11):5212–5222, 2001. doi:10.1063/1.1407837.
- [48] Stanley P. Gudder and Sylvia Pulmannová. Representation theorem for convex effect algebras. *Commentationes Mathematicae Universitatis Carolinae*, 39(4):645–660, 1998.
- [49] Chris J.M. Heunen. *Categorical quantum models and logics*. PhD thesis, Radboud University, 2009. <https://www.math.ru.nl/~landsman/PhDChris.pdf>.
- [50] Chris J.M. Heunen and Martti Karvonen. Limits in dagger categories. arXiv:1803.06651.
- [51] Chris J.M. Heunen, Nicolaas P. Landsman, and Bas Spitters. Bohrification of operator algebras and quantum logic. *Synthese*, 186(3):719–752, 2012. doi:10.1007/s11229-011-9918-4.

- [52] Chris J.M. Heunen and Bert J. Lindenhovius. Domains of commutative  $C^*$ -subalgebras. In *LICS 2015*, pages 450–461, 2015. doi:10.1109/LICS.2015.49.
- [53] Peter Hines and Samuel L. Braunstein. The structure of partial isometries. *Semantic Techniques in Quantum Computation*, pages 361–389, 2010.
- [54] Mathieu Huot and Sam Staton. Universal properties in quantum theory. In *QPL 2018*, volume 287 of *EPTCS*, pages 213–223, 2019. doi:10.4204/EPTCS.287.12.
- [55] Bart P.F. Jacobs. Orthomodular lattices, Foulis semigroups and dagger kernel categories. *LMCS*, 6, June 2010. doi:10.2168/LMCS-6(2:1)2010 arXiv:0912.0931.
- [56] Bart P.F. Jacobs. Probabilities, distribution monads, and convex categories. *Theoretical Computer Science*, 412(28):3323–3336, 2011. doi:10.1016/j.tcs.2011.04.005.
- [57] Bart P.F. Jacobs. New directions in categorical logic, for classical, probabilistic and quantum logic. *LMCS*, 11(3), 2015. doi:10.2168/LMCS-11(3:24)2015 arXiv:1205.3940v5.
- [58] Bart P.F. Jacobs. Affine monads and side-effect-freeness. In *CMCS 2016*, volume 9608 of *LNCS*, pages 53–72, 2016. doi:10.1007/978-3-319-40370-0\_5.
- [59] Bart P.F. Jacobs. Effectuses from monads. *ENTCS*, 325:169–183, 2016. doi:10.1016/j.entcs.2016.09.037.
- [60] Bart P.F. Jacobs. Hyper normalisation and conditioning for discrete probability distributions. *LMCS*, 13(3), 2017. doi:10.23638/LMCS-13(3:17)2017 arXiv:1607.02790v3.
- [61] Bart P.F. Jacobs. Quantum effect logic in cognition. *Journal of Mathematical Psychology*, 81:1–10, 2017. doi:10.1016/j.jmp.2017.08.004.
- [62] Bart P.F. Jacobs. A recipe for state-and-effect triangles. *LMCS*, 13(2), 2017. doi:10.23638/LMCS-13(2:6)2017 arXiv:1703.09034v3.
- [63] Bart P.F. Jacobs. A channel-based perspective on conjugate priors. arXiv:1707.00269v2, 2018.
- [64] Bart P.F. Jacobs. From probability monads to commutative effectuses. *Journal of Logical and Algebraic Methods in Programming*, 94:200–237, 2018. doi:10.1016/j.jlamp.2016.11.006.

- [65] Bart P.F. Jacobs and Jorik Mandemaker. Coreflections in algebraic quantum logic. *Foundations of physics*, 42(7):932–958, 2012. doi:10.1007/s10701-012-9654-8.
- [66] Bart P.F. Jacobs, Jorik Mandemaker, and Robert W.J. Furber. The expectation monad in quantum foundations. *Information and Computation*, 250:87–114, 2016. doi:10.1016/j.ic.2016.02.009.
- [67] Bart P.F. Jacobs and Abraham A. Westerbaan. An effect-theoretic account of lebesgue integration. *ENTCS*, 319:239–253, 2015. doi:10.1016/j.entcs.2015.12.015.
- [68] Bart P.F. Jacobs and Abraham A. Westerbaan. Distances between states and between predicates. arXiv:1711.09740v2, 2018.
- [69] Bart P.F. Jacobs, Bas E. Westerbaan, and Abraham A. Westerbaan. States of convex sets. In *FoSSaCS 2015*, volume 9034 of *LNCS*, pages 87–101, 2015. doi:10.1007/978-3-662-46678-0\_6.
- [70] Bart P.F. Jacobs and Fabio Zanasi. A formal semantics of influence in Bayesian reasoning. In *MFCS 2017*, volume 83 of *LIPICs*, 2017. doi:10.4230/LIPICs.MFCS.2017.21.
- [71] Wang Jia-Mei, Wu Jun-De, and Cho Minhyung. Entropy of partitions on sequential effect algebras. *Communications in Theoretical Physics*, 53(3):399, 2010. doi:10.1088/0253-6102/53/3/01.
- [72] Shen Jun and Junde Wu. Remarks on the sequential effect algebras. *Reports on Mathematical Physics*, 63(3):441–446, 2009. doi:10.1016/S0034-4877(09)90015-5.
- [73] Richard V. Kadison and John R. Ringrose. *Fundamentals of the Theory of Operator Algebras, I & II*. 1983. doi:10.1090/gsm/015 & doi:10.1090/gsm/016.
- [74] Gudrun Kalmbach. *Orthomodular lattices*, volume 18. Academic Pr, 1983. doi:10.1016/S0304-0208(08)73817-9.
- [75] Gennadi G. Kasparov. Hilbert  $C^*$ -modules: theorems of Stinespring and Voiculescu. *Journal of Operator Theory*, pages 133–150, 1980.
- [76] Aleks R. Kissinger, Sean Tull, and Bas E. Westerbaan. Picture-perfect quantum key distribution. 2017. arXiv:1704.08668.
- [77] František Kôpka and Ferdinand Chovanec. D-posets. *Mathematica Slovaca*, 44(1):21–34, 1994.

- [78] Andre Kornell. Quantum collections. *International Journal of Mathematics*, 28(12):1750085, 2017. doi:10.1142/S0129167X17500859 arXiv:1202.2994v2.
- [79] Mareli Korostenski and Walter Tholen. Factorization systems as eilenberg-moore algebras. *Journal of Pure and Applied Algebra*, 85(1):57–72, 1993. doi:10.1016/0022-4049(93)90171-O.
- [80] Dennis Kretschmann, Dirk Schlingemann, and Reinhard F. Werner. The information-disturbance tradeoff and the continuity of Stinespring’s representation. *Information Theory, IEEE Transactions on*, 54(4):1708–1717, 2008. doi:10.1109/TIT.2008.917696.
- [81] E. Christopher Lance. *Hilbert  $C^*$ -modules: a toolkit for operator algebraists*, volume 210. Cambridge University Press, 1995. doi:10.1017/CBO9780511526206.
- [82] Giovanni Landi and Alexander Pavlov. On orthogonal systems in Hilbert  $C^*$ -modules. *Journal of Operator Theory*, pages 487–500, 2012.
- [83] Yuan Li and Xiu-Hong Sun. Sequential product and Jordan product of quantum effects. *International Journal of Theoretical Physics*, 50(4):1206–1213, 2011. doi:10.1007/s10773-010-0615-8.
- [84] Göran Lindblad. Completely positive maps and entropy inequalities. *Communications in Mathematical Physics*, 40(2):147–151, 1975.
- [85] Bert J. Lindenhovius.  $\mathcal{C}(A)$ . PhD thesis, Radboud University, 2016. doi:2066/158429.
- [86] Fred E.J. Linton. Coequalizers in categories of algebras. In *Seminar on triples and categorical homology theory*, volume 80, pages 75–90. Springer, 1969. doi:10.1007/BFb0083082.
- [87] Saunders Mac Lane. *Categories for the working mathematician*. 1971. doi:10.1007/978-1-4757-4721-8.
- [88] VM Manuilov and EV Troitsky. Hilbert  $C^*$ -and  $W^*$ -modules and their morphisms. *Journal of Mathematical Sciences*, 98(2):137–201, 2000.
- [89] Kevin McCrimmon. *A taste of Jordan algebras*. Springer, 2006. doi:10.1007/b97489.
- [90] Walter D. Neumann. On the quasivariety of convex subsets of affine spaces. *Archiv der Mathematik*, 21(1):11–16, 1970. doi:10.1007/BF01220869.
- [91] Michael A. Nielsen and Isaac L. Chuang. Quantum computation and quantum information, 2010. doi:10.1017/CBO9780511976667.

- [92] Arthur J. Parzygnat. Stinespring’s construction as an adjunction. 2018. arXiv:arXiv:1807.02533.
- [93] William L. Paschke. Inner product modules over  $B^*$ -algebras. *Transactions of the American Mathematical Society*, 182:443–468, 1973. doi:10.1090/S0002-9947-1973-0355613-0.
- [94] Vern I. Paulsen. *Completely bounded maps and operator algebras*, volume 78 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2002.
- [95] Gert K. Pedersen.  $C^*$ -algebras and their automorphism groups. *London Mathematical Society Monographs*, 1979.
- [96] Mathys P.A. Rennela, Sam Staton, and Robert W.J. Furber. Infinite-dimensionality in quantum foundations:  $W^*$ -algebras as presheaves over matrix algebras. In *QPL 2016*, volume 236 of *EPTCS*, 2017. doi:10.4204/EPTCS.236.11 arXiv:1701.00662v1.
- [97] Frank Roumen. Cohomology of effect algebras. In *QPL 2016*, volume 236 of *EPTCS*, pages 174–202, 2016. doi: arXiv:1602.00567.
- [98] Frank Roumen. *Effect algebroids*. PhD thesis, Radboud Universiteit Nijmegen, 2017. doi:2066/181438.
- [99] Shôichirô Sakai.  *$C^*$ -algebras and  $W^*$ -algebras*. 1998. doi:10.1007/978-3-642-61993-9.
- [100] Peter Schwabe and Bas E. Westerbaan. Solving binary- $\mathcal{MQ}$  with Grover’s algorithm. In *International Conference on Security, Privacy, and Applied Cryptography Engineering*, pages 303–322. Springer, 2016. doi:10.1007/978-3-319-49445-6\_17 eprint:2019/151.
- [101] John Selby and Bob Coecke. Leaks: quantum, classical, intermediate and more. *Entropy*, 19(4):174, 2017. doi:10.3390/e19040174.
- [102] Peter Selinger. Dagger compact closed categories and completely positive maps. *ENTCS*, 170:139–163, 2007. doi:10.1016/j.entcs.2006.12.018.
- [103] Peter Selinger. Idempotents in dagger categories. *ENTCS*, 210:107–122, 2008. doi:10.1016/j.entcs.2008.04.021.
- [104] Jun Shen and Junde Wu. Not each sequential effect algebra is sharply dominating. *Physics Letters A*, 373(20):1708–1712, 2009. doi:10.1016/j.physleta.2009.02.073.

- [105] Sam Staton. Algebraic effects, linearity, and quantum programming languages. In *ACM SIGPLAN Notices*, volume 50, pages 395–406. ACM, 2015. doi:10.1145/2676726.2676999.
- [106] W. Forrest Stinespring. Positive functions on C\*-algebras. *Proceedings of the American Mathematical Society*, 6(2):211–216, 1955. doi:10.2307/2032342.
- [107] Marshall H. Stone. Postulates for the barycentric calculus. *Annali di Matematica Pura ed Applicata*, 29(1):25–30, 1949. doi:10.1007/BF02413910.
- [108] Erling Størmer. *Positive linear maps of operator algebras*. Springer, 2012. doi:10.1007/978-3-642-34369-8.
- [109] Tadeusz Świrszcz. *Monadic functors and categories of convex sets*. Polish Academy of Sciences. Institute of Mathematics, 1975.
- [110] Masamichi Takesaki. *Theory of operator algebras I*. 1979. doi:10.1007/978-1-4612-6188-9.
- [111] Josef Tkadlec. Atomic sequential effect algebras. *International Journal of Theoretical Physics*, 47(1):185–192, 2008. doi:10.1007/s10773-007-9492-1.
- [112] Marco Tomamichel. *Quantum Information Processing with Finite Resources: Mathematical Foundations*, volume 5. Springer, 2015. doi:10.1007/978-3-319-21891-5.
- [113] Sean Tull. Operational theories of physics as categories. 2016. arXiv:1602.06284.
- [114] Sean Tull. Quotient categories and phases. 2018. arXiv:1801.09532.
- [115] Sean Tull. *Categorical Operational Physics*. PhD thesis, University of Oxford, 2019. arXiv:1902.00343.
- [116] Sander Uijlen and Bas E. Westerbaan. A Kochen-Specker system has at least 22 vectors. *New Generation Computing*, 34(1-2):3–23, 2016. arXiv:1412.8544 doi:10.1007/s00354-016-0202-5.
- [117] John van de Wetering. An effect-theoretic reconstruction of quantum theory. arXiv:1801.05798.
- [118] John van de Wetering. Reconstruction of quantum theory from universal filters. arXiv:1801.05798.
- [119] John van de Wetering. Sequential product spaces are jordan algebras. arXiv:1803.11139.

- [120] John van de Wetering. Three characterisations of the sequential product. *Journal of Mathematical Physics*, 59(8), 2018. doi:10.1063/1.5031089 arXiv:1803.08453.
- [121] Friedrich Wehrung. *Partial Commutative Monoids*, pages 23–69. Springer International Publishing, Cham, 2017. doi:10.1007/978-3-319-61599-8\_2.
- [122] Liu Weihua and Wu Junde. A uniqueness problem of the sequence product on operator effect algebra. *Journal of Physics A: Mathematical and Theoretical*, 42(18):185206, 2009. doi:10.1088/1751-8113/42/18/185206.
- [123] Reinhard F. Werner. Optimal cloning of pure states. *Physical Review A*, 58(3):1827, 1998. doi:10.1103/PhysRevA.58.1827.
- [124] Abraham A. Westerbaan. *The Category of Von Neumann Algebras*. PhD thesis, Radboud University, 2019. arXiv:1804.02203 doi:2066/201611.
- [125] Abraham A. Westerbaan and Bas E. Westerbaan. When does Stinespring dilation yield a faithful representation? Mathematics Stack Exchange. <http://math.stackexchange.com/q/1701574> (version: 2016-03-17).
- [126] Abraham A. Westerbaan and Bas E. Westerbaan. A universal property for sequential measurement. *Journal of Mathematical Physics*, 57(9):092203, 2016. doi:10.1063/1.4961526 arXiv:1603.00410v1.
- [127] Abraham A. Westerbaan and Bas E. Westerbaan. Paschke dilations. In *QPL 2016*, volume 236, pages 229–244, 2017. doi:10.4204/EPTCS.236.15 arXiv:1603.04353v2.
- [128] Abraham A. Westerbaan, Bas E. Westerbaan, and John van de Wetering. Pure maps between Euclidean Jordan algebras. In Peter Selinger and Giulio Chiribella, editors, *QPL 2018*, volume 287 of *EPTCS*, pages 345–364. Open Publishing Association, 2019. doi:10.4204/EPTCS.287.19.
- [129] Bas E. Westerbaan. Sequential product on effect logics. 2013. Master’s thesis, Radboud University. [https://www.ru.nl/publish/pages/813276/masterscriptie\\_bas\\_westerbaan.pdf](https://www.ru.nl/publish/pages/813276/masterscriptie_bas_westerbaan.pdf).
- [130] Bas E. Westerbaan, Abraham A. and Westerbaan. Reference request unital normal  $*$ -homomorphisms  $B(H) \rightarrow B(K)$ . Mathematics Stack Exchange. <https://math.stackexchange.com/q/3140021> (version: 2019-03-11).
- [131] Alexander Wilce. A shortcut from categorical quantum theory to convex operational theories. In *QPL 2017*, volume 266 of *EPTCS*, pages 222–236, 2017. doi:10.4204/EPTCS.266 arXiv:1802.09737.

- [132] Alexander Wilce. A royal road to quantum theory (or thereabouts). *Entropy*, 20(4):227, 2018. doi:10.3390/e20040227.
- [133] Stephen Willard. *General topology*. Addison–Wesley, 1970.



# Lekensamenvatting

De volgende twee technische vraagstukken in de wiskundige theorie van quantum computers liggen ten grondslag aan dit proefschrift.

1. Breidt de stelling van Stinespring uit naar alle von Neumann algebra's?
2. Is de categorie van von Neumann algebra's te axiomatiseren?

Het eerste vraagstuk wordt beantwoord: ja, de stelling breidt uit. Het tweede vraagstuk blijft helaas voor een groot deel onbeantwoord. De belangrijkste bijdragen van dit proefschrift aan de wetenschap zijn niet per se deze directe resultaten, maar de bijvangst die het onderzoek erna heeft opgeleverd: er worden verscheidene nieuwe stellingen bewezen en begrippen ingevoerd die toepassing hebben buiten deze vraagstukken (zoals bijvoorbeeld een uitbreiding van de stelling van Kaplansky en het begrip  $\diamond$ -geadjungeerdheid).

Maar waar gaan deze twee originele vraagstukken eigenlijk over? Wat is die stelling van Stinespring? Wat zijn von Neumann algebra's? En wat is een quantum computer überhaupt?

## Quantum computers

Een quantum computer gebruikt bepaalde quantum mechanische eigenaardigheden om sommige berekeningen veel efficiënter uit te voeren dan een traditionele computer. Een zorgelijk voorbeeld is dat quantum computer erg goed is in het ontbinden van getallen in priemfactoren. Het grootste deel van de moderne cryptografie die ons beveiligt is gestoeld op de aanname dat het vinden van zulke ontbindingen lastig is en deze wordt dan ook makkelijk door een voldoende grote quantum computer gebroken. Gelukkig zijn de huidige quantum computers nog erg klein en hebben wij de tijd om onze cryptografie aan te passen. Aan de positieve kant zijn quantum computer erg geschikt voor het simuleren van het gedrag van moleculen, waaronder het lokale effect van potentiële medicijnen.

In functie is een quantum computer te vergelijken met een grafische kaart van een computer: voor de meeste berekeningen is een normale CPU het best,

maar voor sommige berekeningen is een grafische kaart veel efficiënter. Dit komt doordat een normale CPU gemaakt is om zo snel mogelijk losse rekenstappen op afzonderlijke getallen achter elkaar uit te voeren, terwijl een grafische kaart rekenstappen uitvoert op miljoenen getallen tegelijkertijd. Niet elke berekening heeft baat bij dat grote parallelisme. Voor diegene die dat wel hebben (zoals het weergeven van een 3D-wereld in een computerspel) is het lastig om het originele algoritme aan te passen om optimaal gebruik te maken van de grafische kaart. Programmeren voor een grafische kaart vraagt om een heel andere *mindset*. Ook de theoretische analyse van algoritmes voor grafische kaarten is van een andere aard dan die voor een traditionele computer.

De situatie bij een quantum computer is vergelijkbaar, maar extremer: de efficiëntiewinst tussen een quantum computer en PC is groter dan die tussen een CPU en grafische kaart. Helaas is het vinden en analyseren van algoritmes voor quantum computers ook vele malen ingewikkelder. Er is ook geen methode of zelfs vuistregel bekend om te bepalen of een berekening efficiënter uit te voeren is op een quantum computer of niet.

## Von Neumann algebra's

Het gedrag van een algoritme voor een traditionele computer, zoals diegene die het kwadraat van een telgetal berekent, wordt vaak beschreven met wiskundige functie, zoals in dit geval de functie  $f: \mathbb{N} \rightarrow \mathbb{N}$  met  $f(n) = n^2$ . De notatie  $f: \mathbb{N} \rightarrow \mathbb{N}$  betekent dat de functie  $f$  als invoer een telgetal  $\mathbb{N} \equiv \{0, 1, 2, \dots\}$  neemt en ook een telgetal als uitvoer geeft. Een echt programma draait op een computer dat maar een beperkte hoeveelheid geheugen heeft: er is dus een grootste telgetal dat in de computer past. Dat we het algoritme beschrijven met willekeurig grote telgetallen is een idealisering van de situatie. Het zou echter onhandig zijn om alle generiek toepasbare algoritme te beschrijven als wiskundige functies tussen eindige verzamelingen (zoals  $\{0, 1, 2, 3\}$ ) in plaats van tussen oneindige verzamelingen zoals  $\mathbb{N}$ .

Dit is helaas precies de situatie voor de meeste beschrijvingen van algoritmes voor quantum computers. Het probleem is dat het analoog van  $\mathbb{N}$  en de wiskundige functies daartussen vele malen complexer van aard zijn dan de wiskundige functies die het gedrag beschrijven voor het eindige geval. Ook al dit eindige geval op zichzelf is stukken ingewikkelder dan de simpele functies tussen eindige verzamelingen voor de normale algoritmes.

*Von Neumann algebra's* zijn de wiskundige structuren die gebruikt worden om oneindige quantumdatatypes te beschrijven zoals het analoog van  $\mathbb{N}$ . Zogenaemde *ncp-functies* tussen von Neumann algebras zijn het analoog van de normale wiskundige functies tussen verzamelingen. Het zijn deze von Neumann algebra's met bijbehorende ncp-functies, die in dit proefschrift onderzocht worden om ze beter te begrijpen en eenvoudiger toe te kunnen passen.

## De stelling van Stinespring

Het is lastig om grip te krijgen op een willekeurige ncp-functie tussen von Neumann algebra's. De stelling van Stinespring helpt hierbij: deze zegt dat elke ncp-functie  $f: \mathcal{A} \rightarrow \mathcal{B}_I$  tussen von Neumann algebra's  $\mathcal{A}$  en  $\mathcal{B}_I$  (waar  $\mathcal{B}_I$  van een speciale soort is) in feite de samenstelling is van twee veel simpelere ncp-functies: eerst  $f_1: \mathcal{A} \rightarrow \mathcal{P}$  en dan  $f_2: \mathcal{P} \rightarrow \mathcal{B}_I$ . Deze bijzondere opsplitsing wordt ook wel een *dilatatie* genoemd en wordt gebruikt als cruciale stap in het bewijs van vele stellingen, niet alleen over ncp-functies zelf, maar ook over bijvoorbeeld quantum protocollen en quantum informatie. Het probleem is dat de stelling van Stinespring alleen toepasbaar is als  $\mathcal{B}_I$  van de speciale soort *type I* is. Daarmee zijn de stellingen die ermee worden bewezen vaak ook maar toepasbaar op zulke speciale von Neumann algebra's. In het eerste hoofdstuk van het proefschrift wordt een generalisatie van de stelling van Stinespring bewezen, die toepasbaar is op alle ncp-functies.

## Axiomatisatie

Waarom zijn juist von Neumann algebra's en ncp-functies geschikt voor oneindige quantum datatypes? In feite is het een wiskundige gok: von Neumann algebra's en ncp-functies lijken de eenvoudigste structuren die de voorbeelden die we kennen goed beschrijven. Eenvoud hier is relatief: de ingewikkelde definitie van een von Neumann algebra staat ver van de intuïtie van een informaticus en natuurkundige. Het valt ook te betwijfelen of elke von Neumann algebra een realistisch quantum mechanisch systeem beschrijft, laat staan een quantum datatype.

Een mogelijke manier om deze kwestie te beslechten is de volgende: wij zoeken een lijst van kenmerkende (en voor onze toepassing relevante) eigenschappen waaraan von Neumann algebra's samen met ncp-functies aan voldoen. Als er bewezen kan worden dat alleen von Neumann algebra's met ncp-functies aan deze lijst van eigenschappen kan voldoen (en geen ander soort wiskundige structuur) dan hebben wij met deze lijst van eigenschappen (nu verheven tot axioma's) een nieuw inzicht in de aard van von Neumann algebra's.

Een dergelijke aanpak is eerder al succesvol gebruikt om de regels van quantum theorie voor eindige systemen beter te begrijpen. Een voorbeeld hiervan zijn de axioma's van Chiribella et al, die quantum theorie karakteriseren door de wijze waarop informatie verwerkt wordt.

In het tweede hoofdstuk van dit proefschrift wordt gezocht naar een dergelijke axiomatisatie. Een van de kernaxioma's is het bestaan van een soort *omkering*, een zogenaemde *dagger*, op de speciale ncp-functies die aan de rechterkant voorkomen van een dilatatie, wat de twee hoofdstuken met elkaar verbindt. Helaas wordt een volledige axiomatisatie niet bereikt. Het verhaal eindigt hier niet:

mijn college van de Wetering heeft recent bewezen dat met een paar toevoegingen, de axioma's uit dit proefschrift zogeheten Euclidische Jordan algebra's (EJA's) karakteriseren. Hiermee is de kous nog niet af: EJA's zijn niet geschikt voor oneindige quantum datatypes. Dit resultaat suggereert wel de mogelijkheid dat JBW-algebra's, welke een kruising zijn van von Neumann en Jordan algebra's, uiteindelijk de juiste algebra's zijn voor onze toepassing.

Hoewel ons tweede vraagstuk niet beantwoord is, heeft de zoektocht naar geschikte axioma's een aantal nieuwe begrippen aan het licht gebracht (zoals  $\diamond$ -geadjungeerdheid, hoeken en filters), die een nieuw inzicht verschaffen in von Neumann algebra's en daarmee quantum computers in het algemeen.

# About the author

Bas Westerbaan, born in 1988, enrolled to the Radboud Universiteit as a physics student in 2007. He received his bachelor's degree in mathematics with a thesis on computability theory supervised by dr. Wim Veldman. He continued his study of logic and foundational computer science during his master's education and developed a keen interest for, what he thought to be, an unrelated mathematical field: functional analysis. These interests however, are conveniently combined in the present thesis which continues the research of his master's thesis, which was supervised by prof. Bart Jacobs as well. During his undergraduate studies, Bas also served as secretary of the board of the 150 member student club Karpe Noktem and he was elected to the student board of the faculty of science. His master's degree in mathematics was awarded cum laude in 2013.

While performing the research as a *promovendus* at the Digital Security group that led to this thesis, Bas coauthored several security audits including those of the *DigiD*- and *BerichtenBox*-app; the Dutch digital identity and mail system. In the year between the submission and defense of his thesis, Bas was employed as a post-doctoral researcher where he assisted with the development of a cryptographic system for polymorphic pseudonymisation and encryption. During that same year he also taught a quarter-year course on computer networking and was kindly invited by dr. Heunen to speak about his research at the University of Edinburgh.



# Solutions to exercises

**Solution to 138 VII.** To prove the first statement, let  $\varphi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$  be any ncp-map. If  $\varphi = 0$  then  $\mathcal{K}' = 0$  and  $V = 0$  does the job, so for the other case, assume  $\varphi \neq 0$ . By 135 IV there is a Hilbert space  $\mathcal{H}'$ , a bounded operator  $W: \mathcal{H} \rightarrow \mathcal{H}'$  and an ncpu-map  $\varrho: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H}')$  such that  $\varphi = \text{ad}_W \circ \varrho$ . Clearly  $\varrho \neq 0$ . By 138 II there is a Hilbert space  $\mathcal{K}'$  and a unitary  $U: \mathcal{H}' \rightarrow \mathcal{H} \otimes \mathcal{K}'$  with  $\varrho(A) = U^*(A \otimes 1)U$  for all  $A \in \mathcal{B}(\mathcal{H})$ . Define  $V \equiv UW$ . Then  $\varphi(A) = W^*\varrho(A)W = W^*U^*(A \otimes 1)UW = V^*(A \otimes 1)V$ , as desired.

Before we can continue with the second statement, we need to understand the relationship between quantum channels and ncpu-maps. This relationship is best understood with predual characterization of von Neumann algebras due to Sakai [99], which we have been avoiding. The characterization is as thus: a  $C^*$ -algebra  $\mathcal{A}$  is a von Neumann algebra if and only if it is isomorphic to the dual of a Banach space. Then this Banach space is unique up-to-isomorphism as it must be isomorphic to the space of normal functionals on  $\mathcal{A}$  (denoted by  $\mathcal{A}_*$ ) and is appropriately called the *predual* of  $\mathcal{A}$ . Any normal linear map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  between von Neumann algebras  $\mathcal{A}$  and  $\mathcal{B}$  yields a linear map  $\varphi_*: \mathcal{B}_* \rightarrow \mathcal{A}_*$  via  $\varphi_*(\omega) = \omega \circ \varphi$ . In the other direction, any linear map  $\varphi_*: \mathcal{B}_* \rightarrow \mathcal{A}_*$  gives rise to a normal linear map  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  by defining  $\varphi(a)(\omega) = \varphi_*(\omega)(a)$  where we identified  $\mathcal{A} \equiv (\mathcal{A}_*)^*$ . Clearly  $\varphi$  is positive precisely if  $\varphi_*$  maps positive functionals to positive functionals.

A normal state  $\omega: \mathcal{B}(\mathcal{H}) \rightarrow \mathbb{C}$  is precisely of the form  $\omega(a) = \text{tr}[\rho a]$  for some density matrix  $\rho$  over  $\mathcal{H}$ . Thus the predual of  $\mathcal{B}(\mathcal{H})$  can be identified with the set of trace-class operators over  $\mathcal{H}$ . Let  $\varphi_*$  be a linear map from the density operators on  $\mathcal{H}$  to those on  $\mathcal{K}$ . The map  $\varphi_*$  is completely positive in its usual sense if the corresponding map  $\varphi$  is completely positive. Furthermore  $\varphi$  is unital if and only if  $\varphi_*$  is trace-preserving.

To prove the second statement, let  $\Phi$  be any quantum channel mapping density matrices over  $\mathcal{H}$  to those of  $\mathcal{H}$  again. (Note: in the printed version of the thesis the exercise incorrectly assumes  $\Phi$  to map density matrices over  $\mathcal{H}$  to those over some other Hilbert space  $\mathcal{K}$ .) It follows from the previous, that

there is a unique ncpu-map  $\varphi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  with

$$\mathrm{tr}[\Phi(\rho)A] = \mathrm{tr}[\rho\varphi(A)] \quad \text{for any density matrix } \rho \in \mathcal{B}(\mathcal{H}).$$

See also [112] for a more direct approach. By the first part of the exercise, we know that there is a Hilbert space  $\mathcal{K}'$  and a bounded operator  $V: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{K}'$  with  $\varphi(A) = V^*(A \otimes 1)V$ . Tracing back the definition of  $V$ , we see that  $V$  is an isometry because  $\varphi$  is unital. Pick any orthonormal bases  $E$  and  $F$  of  $\mathcal{H}$  and  $\mathcal{K}'$  respectively. We may assume, without loss of generality, that  $\mathcal{K}'$  is not zero-dimensional by setting  $\mathcal{K}' = \mathbb{C}$  and  $V = 0$  in the case that  $\Phi = 0$ . Pick any  $f_0 \in F$  and any unitary  $U: \mathcal{H} \otimes \mathcal{K}' \rightarrow \mathcal{H} \otimes \mathcal{K}'$  with  $U^*x \otimes f_0 = Vx$ , which exists as  $V$  is an isometry. Now we compute

$$\begin{aligned} \mathrm{tr}[\varphi(A)\rho] &= \mathrm{tr}[\rho V^*(A \otimes 1)V] \\ &= \sum_{e \in E} \langle V\rho e, (A \otimes 1)V e \rangle \\ &= \sum_{e \in E} \langle U^*(\rho \otimes 1)e \otimes f_0, (A \otimes 1)U^*e \otimes f_0 \rangle. \end{aligned}$$

Inserting  $|f_0\rangle\langle f_0|$  in the previous, we may sum over all  $f \in F$  and get

$$\begin{aligned} \mathrm{tr}[\varphi(A)\rho] &= \sum_{\substack{e \in E \\ f \in F}} \langle U^*(\rho \otimes |f_0\rangle\langle f_0|)e \otimes f, (A \otimes 1)U^*e \otimes f \rangle \\ &= \sum_{\substack{e \in E \\ f \in F}} \langle (U^*(e \otimes f)), U^*(\rho \otimes |f_0\rangle\langle f_0|)U(A \otimes 1)(U(e \otimes f)) \rangle \\ &= \mathrm{tr}[U^*(\rho \otimes |f_0\rangle\langle f_0|)U(A \otimes 1)] \\ &= \mathrm{tr}[A \mathrm{tr}_{\mathcal{K}'}[U^*(\rho \otimes |f_0\rangle\langle f_0|)U]]. \end{aligned}$$

This shows that indeed  $\Phi(\rho) = \mathrm{tr}_{\mathcal{K}'}[U^*(\rho \otimes |v_0\rangle\langle v_0|)U]$  as desired with  $v_0 \equiv f_0$ .

**Solution to 138 VIII.** Let  $\varphi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  be any ncpu-map. By 138 VII there is a Hilbert space  $\mathcal{K}'$  and a bounded operator  $V: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{K}'$  with  $\varphi(A) = V^*(A \otimes 1)V$ . Let  $E$  be any orthonormal basis of  $\mathcal{K}'$ . Then  $1 = \sum_{e \in E} |e\rangle\langle e|$  where the sum converges ultraweakly and so by ultraweak continuity of  $\mathrm{ad}_V$  (44 VIII) and  $B \mapsto A \otimes B$  (116 III), we see

$$\varphi(A) = V^* \left( A \otimes \sum_{e \in E} |e\rangle\langle e| \right) V = \sum_{e \in E} V^*(A \otimes |e\rangle\langle e|)V. \quad (4.1)$$

For  $e \in E$ , define  $P_e: \mathcal{H} \otimes \mathcal{K}' \rightarrow \mathcal{H}$  by  $P_e \equiv 1 \otimes \langle e|$ , i.e.  $P_e(x \otimes y) = x\langle e, y \rangle$ .

Define  $V_e \equiv P_e V$ . Note that  $P_e^* A P_e = A \otimes |e\rangle\langle e|$  and so

$$\begin{aligned}\varphi(A) &= \sum_{e \in E} V^*(A \otimes |e\rangle\langle e|)V && \text{by (4.1)} \\ &= \sum_{e \in E} V^* P_e^* A P_e V \\ &= \sum_{e \in E} V_e^* A V_e,\end{aligned}$$

as desired. From the special case  $A = 1$ , we see that  $\sum_{e \in E} V_e^* V_e = \varphi(1)$  and so the partial sums of  $\sum_{e \in E} V_e^* V_e$  are bounded.

For the final part, assume  $\mathcal{H}$  and  $\mathcal{K}$  are finite dimensional. Recall that the standard Stinespring dilation space (say  $\mathcal{K}''$ ) for  $\varphi$  is constructed using a completion and quotient of  $\mathcal{B}(\mathcal{H}) \odot \mathcal{K}$ . As  $\mathcal{B}(\mathcal{H}) \odot \mathcal{K}$  is finite dimensional it is already complete. Hence  $\mathcal{K}''$  has dimension at most  $(\dim \mathcal{H})^2(\dim \mathcal{K})$ . By construction  $\mathcal{H} \otimes \mathcal{K}' \cong \mathcal{K}''$ , hence  $\dim \mathcal{K}' \leq (\dim \mathcal{H})(\dim \mathcal{K})$ . Recall  $E$  is a basis of  $\mathcal{K}'$  and so there are indeed at most  $(\dim \mathcal{H})(\dim \mathcal{K})$  Kraus operators.

**Solution to 139 IX.** We will show that  $U: \text{Rep} \rightarrow \text{Rep}_{\text{cp}}$  has a left adjoint by demonstrating the universal mapping property. Let  $\varphi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  be any object of  $\text{Rep}_{\text{cp}}$ . Pick any minimal Stinespring dilation  $(\mathcal{K}, \varrho, V)$  of  $\varphi$ . The map  $\varrho: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  is an object of  $\text{Rep}$ . Clearly  $\text{ad}_V \circ \varrho \circ \text{id} = \varphi$  and so  $\eta_\varphi \equiv (\text{id}, V): \varphi \rightarrow U\varrho$  is a morphism in  $\text{Rep}_{\text{cp}}$ . We will show that for each  $f: \varphi \rightarrow U\varrho'$  in  $\text{Rep}_{\text{cp}}$ , there is a unique  $f': \varrho \rightarrow \varrho'$  in  $\text{Rep}$  with  $Uf' \circ \eta_\varphi = f$ . This is sufficient to show that  $U$  has a left adjoint.

So let  $f: \varphi \rightarrow U\varrho'$  be any morphism in  $\text{Rep}_{\text{cp}}$ . Say  $\varrho': \mathcal{A}' \rightarrow \mathcal{B}(\mathcal{K}')$ . Then  $f \equiv (m', V')$  consists of a nmiu-map  $m': \mathcal{A} \rightarrow \mathcal{A}'$  and bounded operator  $V': \mathcal{H} \rightarrow \mathcal{K}'$  with  $\text{ad}_{V'} \circ \varrho' \circ m' = \varphi$ . By 139 V there is a unique bounded operator  $S: \mathcal{H} \rightarrow \mathcal{K}'$  with  $SV = V'$  and  $\varrho = \text{ad}_S \circ \varrho' \circ m'$ . This turns  $f' \equiv (m', S)$  into a morphism  $\varrho \rightarrow \varrho'$  in  $\text{Rep}$ . Furthermore  $Uf' \circ \eta_\varphi = (m' \circ \text{id}, SV) = (m', V') = f$ . To show uniqueness, assume there is some  $f'': \varrho \rightarrow \varrho'$  in  $\text{Rep}$  with  $Uf'' \circ \eta_\varphi = f$ . Say  $f'' = (m'', S'')$ . Then  $(m', V') = f = Uf'' \circ \eta_\varphi = (m'', S''V)$ . So  $m'' = m'$  and  $V' = S''V$ . The fact that  $f''$  is a morphism in  $\text{Rep}$  is equivalent to  $\text{ad}_{S''} \circ \varrho' \circ m'' = \varrho$ . Thus  $\text{ad}_{S''} \circ \varrho' \circ m' = \varrho$ . By uniqueness of  $S$ , we get  $S'' = S$ . Hence  $f'' = (m'', S'') = (m', S) = f'$ , as desired.

**Solution to 139 XI.** Let  $\varphi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$  be any ncp-map. As in the description of the exercise, let  $\mathcal{K}$  be a Hilbert space and  $V, W: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{K}'$  be bounded operators with  $V^*(a \otimes 1)V = \varphi(a) = W^*(a \otimes 1)W$ . Write  $\mathcal{V}$  for the closed linear span of  $\{(a \otimes 1)Vx; a \in \mathcal{B}(\mathcal{H}), x \in \mathcal{K}\}$  in  $\mathcal{H} \otimes \mathcal{K}'$  and similarly  $\mathcal{W}$  for that of  $\{(a \otimes 1)Wx; a \in \mathcal{B}(\mathcal{H}), x \in \mathcal{K}\}$ . Note that for any  $n \in \mathbb{N}$ ,

$x_1, \dots, x_n \in \mathcal{K}$  and  $a_1, \dots, a_n \in \mathcal{B}(\mathcal{H})$  we have

$$\begin{aligned} \left\| \sum_i (a_i \otimes 1) V x_i \right\|^2 &= \sum_{i,j} \langle x_i, V^* ((a_i^* a_j) \otimes 1) V x_j \rangle \\ &= \sum_{i,j} \langle x_i, W^* ((a_i^* a_j) \otimes 1) W x_j \rangle \\ &= \left\| \sum_i (a_i \otimes 1) W x_i \right\|^2. \end{aligned}$$

Thus there is a unique unitary  $U_0: \mathcal{W} \rightarrow \mathcal{V}$  fixed by  $U_0(a \otimes 1)Wx = U_0(a \otimes 1)Vx$ . We see  $U_0W = V$  by setting  $a = 1$ . Furthermore

$$(\alpha \otimes 1)U_0(a \otimes 1)Wx = ((\alpha a) \otimes 1)Vx = U_0(\alpha \otimes 1)(a \otimes 1)Wx$$

for any  $\alpha, a \in \mathcal{B}(\mathcal{H})$  and  $x \in \mathcal{K}$ , hence  $(\alpha \otimes 1)U_0 = U_0(\alpha \otimes 1)$ .

For any  $a \in \mathcal{B}(\mathcal{H})$ , the operator  $a \otimes 1 \in \mathcal{B}(\mathcal{H} \otimes \mathcal{K}')$  restricts to  $\mathcal{B}(\mathcal{W})$ . Pick an orthonormal basis  $E$  of  $\mathcal{H}$  and some  $e_0 \in E$ . Note that  $(|e_0\rangle\langle e_0| \otimes 1)\mathcal{W} = e_0 \otimes \mathcal{W}'$  for some closed subspace  $\mathcal{W}' \subseteq \mathcal{K}'$ . In fact, for any  $e \in E$  we have  $e \otimes \mathcal{W}' = (|e\rangle\langle e_0| \otimes 1)(e_0 \otimes \mathcal{W}') = (|e\rangle\langle e_0| \otimes 1)(1 \otimes |e_0\rangle\langle e_0|)\mathcal{W} = (|e\rangle\langle e| T \otimes 1)\mathcal{W} = (|e\rangle\langle e| \otimes 1)\mathcal{W}$ , where  $T$  is the unitary on  $\mathcal{H}$  that only swaps  $e$  and  $e_0$ . Hence  $\mathcal{W} = \mathcal{H} \otimes \mathcal{W}'$ . Similarly  $\mathcal{V} = \mathcal{H} \otimes \mathcal{V}'$  for some closed subspace  $\mathcal{V}' \subseteq \mathcal{K}'$ .

For any non-zero  $w \in \mathcal{W}'$  and unit-vector  $x \in \mathcal{H}$ , we have  $U_0(x \otimes w) = U_0(|x\rangle\langle x| \otimes 1)(x \otimes w) = (|x\rangle\langle x| \otimes 1)U_0(x \otimes w)$  so  $U_0(x \otimes w) = x \otimes y$  for some  $y \in \mathcal{V}'$ . Clearly  $\|w\| = \|x \otimes w\| = \|U_0(x \otimes w)\| = \|x \otimes y\| = \|y\|$ , so there is a unique unitary  $U_1: \mathcal{W}' \rightarrow \mathcal{V}'$  with  $U_0(x \otimes w) = x \otimes U_1w$ . It follows  $U_0 = 1 \otimes U_1$ .

As  $\mathcal{V}$  and  $\mathcal{W}$  are isomorphic, they have the same dimension and so do  $\mathcal{V}'$  and  $\mathcal{W}'$ . Consequently, there is an unitary  $U: \mathcal{K}' \rightarrow \mathcal{K}'$  extending  $U_1$ . We have  $V = (1 \otimes U)W = (1 \otimes U_1)W = U_0W = V$  as desired.

**Solution to 140 X.** We cover the points in order.

1. Let  $\varrho: \mathcal{A} \rightarrow \mathcal{B}$  be a mniw-map. Assume there are mniw  $\varrho': \mathcal{A} \rightarrow \mathcal{P}'$  and ncp  $h': \mathcal{P}' \rightarrow \mathcal{B}$  with  $h' \circ \varrho' = \varrho$ . We have to show there is a unique map  $\sigma: \mathcal{P} \rightarrow \mathcal{B}$  with  $\text{id} \circ \sigma = h'$  and  $h' \circ \varrho' = \varrho$ . Clearly  $\sigma \equiv h'$  fits the bill.
2. Let  $(\mathcal{P}, \varrho, h)$  be any Paschke dilation. We will show that  $(\mathcal{P}, \text{id}, h)$  is a Paschke dilation of  $h$ . To this end, let  $\varrho': \mathcal{A} \rightarrow \mathcal{P}'$  be any mniw-map and  $h': \mathcal{P}' \rightarrow \mathcal{B}$  be any ncp-map with  $h' \circ \varrho' = h$ . Consider  $\varrho' \circ \varrho$  and  $h'$ . By the universal property of the original dilation, there is a unique ncp-map  $\sigma: \mathcal{P}' \rightarrow \mathcal{P}$  with  $\sigma \circ \varrho' \circ \varrho = \varrho$  and  $h \circ \sigma = h'$ . Furthermore  $\text{id}: \mathcal{P} \rightarrow \mathcal{P}$  is the unique ncp-map with  $\text{id} \circ \varrho = \varrho$  and  $h \circ \text{id} = h$ . Now  $(\sigma \circ \varrho') \circ \varrho = \varrho$  and  $h \circ (\sigma \circ \varrho') = h' \circ \varrho' = h$ , so  $\sigma \circ \varrho' = \text{id}$ . We are halfway demonstrating that  $\sigma$  is also the mediating map for our dilation

of  $h$ . It remains to be shown that  $\sigma$  is the unique ncp-map with  $\sigma \circ \varrho' = \text{id}$  and  $h \circ \sigma = h'$ . So assume there is a ncp-map  $\sigma' : \mathcal{P}' \rightarrow \mathcal{P}$  with  $h \circ \sigma' = h'$  and  $\sigma' \circ \varrho' = \text{id}$ . Clearly  $\sigma' \circ \varrho' \circ \varrho = \varrho$  and so by uniqueness of  $\sigma$  as the mediating map for the original dilation, we see  $\sigma' = \sigma$ , as desired.

3. Let  $(\varphi_2^1) : \mathcal{A} \rightarrow \mathcal{B}_1 \oplus \mathcal{B}_2$  be any ncp-map. Pick Paschke dilations  $(\mathcal{P}_i, \varrho_i, h_i)$  of  $\varphi_i$  for  $i = 1, 2$ . We will show that  $(\mathcal{P}_1 \oplus \mathcal{P}_2, (\frac{\varrho_1}{\varrho_2}), h_1 \oplus h_2)$  is a Paschke dilation of  $(\varphi_2^1)$ . Clearly  $h_1 \oplus h_2 \circ (\frac{\varrho_1}{\varrho_2}) = (\frac{h_1 \circ \varrho_1}{h_2 \circ \varrho_2}) = (\varphi_2^1)$ . Let  $\varrho' : \mathcal{A} \rightarrow \mathcal{P}'$  be any nmiu-map and  $(\frac{h_1}{h_2}) : \mathcal{P}' \rightarrow \mathcal{B}_1 \oplus \mathcal{B}_2$  any ncp-map with  $(\frac{h_1}{h_2}) \circ \varrho' = (\varphi_2^1)$ . Note  $h_i \circ \varrho' = \varphi_i$  (for  $i = 1, 2$ ) and so there is a unique  $\sigma_i : \mathcal{P}' \rightarrow \mathcal{P}_i$  with  $\sigma_i \circ \varrho' = \varrho_i$  and  $h_i \circ \sigma_i = h'_i$ . We will show that  $(\frac{\sigma_1}{\sigma_2}) : \mathcal{P}' \rightarrow \mathcal{P}_1 \oplus \mathcal{P}_2$  is the unique mediating map. Clearly  $(\frac{\sigma_1}{\sigma_2}) \circ \varrho' = (\frac{\sigma_1 \circ \varrho'}{\sigma_2 \circ \varrho'}) = (\frac{\varrho_1}{\varrho_2})$  and  $(h_1 \oplus h_2) \circ (\frac{\sigma_1}{\sigma_2}) = (\frac{h'_1 \circ \sigma_1}{h'_2 \circ \sigma_2}) = (\frac{h_1}{h_2})$ . To show uniqueness of  $(\frac{\sigma_1}{\sigma_2})$ , assume there is ncp-map  $(\frac{\sigma'_1}{\sigma'_2}) : \mathcal{P}' \rightarrow \mathcal{P}_1 \oplus \mathcal{P}_2$  such that  $(\frac{\sigma'_1}{\sigma'_2}) \circ \varrho' = (\frac{\varrho_1}{\varrho_2})$  and  $(h_1 \oplus h_2) \circ (\frac{\sigma'_1}{\sigma'_2}) = (\frac{h_1}{h_2})$ . Then  $h_i \circ \sigma'_i = h_i$  and  $\sigma'_i \circ \varrho' = \varrho_i$  for  $i = 1, 2$  and so  $\sigma_i = \sigma'_i$  by the uniqueness of the separate  $\sigma_i$ . Thus indeed  $(\frac{\sigma_1}{\sigma_2}) = (\frac{\sigma'_1}{\sigma'_2})$ .

4. Let  $\varphi : \mathcal{A} \rightarrow \mathcal{B}$  be any ncp-map with Paschke dilation  $(\mathcal{P}, \varrho, h)$ . Assume  $\lambda \in \mathbb{R}, \lambda > 0$ . We will show  $(\mathcal{P}, \varrho, \lambda h)$  is a Paschke dilation of  $\lambda \varphi$ . Clearly  $\lambda h \circ \varrho = \lambda \varphi$ . To this end, assume  $\varrho' : \mathcal{A} \rightarrow \mathcal{P}'$  is a nmiu-map and  $h' : \mathcal{P}' \rightarrow \mathcal{B}$  is an ncp-map with  $h' \circ \varrho' = \lambda \varphi$ . Then  $\lambda^{-1} h' \circ \varrho' = \varphi$ . Thus there is a unique  $\sigma : \mathcal{P}' \rightarrow \mathcal{P}$  with  $\sigma \circ \varrho' = \varrho$  and  $h \circ \sigma = \lambda^{-1} h'$ . Clearly  $\lambda h \circ \sigma = h'$  and so  $\sigma$  also serves as the unique mediating map for the dilation of  $\lambda \varphi$ .

**Solution to 142V.** Let  $X$  be a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product  $\langle \cdot, \cdot \rangle$  for some C\*-algebra  $\mathcal{B}$ . Using the C\*-identity, 142III and the definition of  $\| \cdot \|$  on  $X$ , we get  $\|\langle x, y \rangle\|^2 = \|\langle x, y \rangle^* \langle x, y \rangle\| = \|\langle y, x \rangle \langle x, y \rangle\| \leq \| \langle x, x \rangle \| \langle y, y \rangle \| = \|x\|^2 \|y\|^2$ , so indeed  $\|\langle x, y \rangle\| \leq \|x\| \|y\|$ .

Now we will show  $\|x\| \equiv \|\langle x, x \rangle\|^{\frac{1}{2}}$  is a seminorm on  $X$ . Clearly  $\|x\| \geq 0$  for any  $x \in X$ . For any  $\lambda \in \mathbb{C}$  and  $x \in X$  we have  $\langle \lambda x, \lambda x \rangle = \overline{\lambda} \langle x, x \rangle \lambda = |\lambda|^2 \langle x, x \rangle$ , hence  $\|\lambda x\| = \|\lambda^2 \langle x, x \rangle\|^{\frac{1}{2}} = |\lambda| \|x\|$ . Next, for any  $x, y \in X$  we have

$$\begin{aligned} \|x + y\|^2 &= \|\langle x + y, x + y \rangle\| \\ &\leq \|\langle x, x \rangle\| + \|\langle y, y \rangle\| + \|\langle x, y \rangle\| + \|\langle x, y \rangle^*\| \\ &= \|x\|^2 + \|y\|^2 + 2\|\langle x, y \rangle\| \\ &\leq \|x\|^2 + \|y\|^2 + 2\|x\| \|y\| \\ &= (\|x\| + \|y\|)^2 \end{aligned}$$

and thus  $\|\cdot\|$  is indeed a seminorm.

Finally, we will show  $\|x \cdot b\| \leq \|x\|\|b\|$  for any  $b \in \mathcal{B}$  and  $x \in X$ . As  $\langle x, x \rangle$  is positive, we have  $\langle x, x \rangle \leq \|\langle x, x \rangle\| = \|x\|^2$ . Also recall  $a \mapsto b^*ab$  is clearly positive. Thus  $\|x \cdot b\|^2 \equiv \|\langle xb, xb \rangle\| = \|b^*\langle x, x \rangle b\| \leq \|\|x\|^2 b^*b\| = \|b\|^2\|x\|^2$  as desired.

**Solution to 142 IX.** Let  $B$  be any  $\mathcal{B}$ -sesquilinear form on a pre-Hilbert  $\mathcal{B}$ -module  $X$  for some  $C^*$ -algebra  $\mathcal{B}$ . Distributing  $B$  in  $\sum_{k=0}^3 i^k B(i^k x + y, i^k x + y)$  we get 16 terms consisting of four of each  $B(x, x)$ ,  $B(y, y)$ ,  $B(x, y)$  and  $B(y, x)$  with the following coefficients.

	$B(x, x)$	$B(y, y)$	$B(x, y)$	$B(y, x)$
$k = 0$	1	1	1	1
$k = 1$	$i \cdot (-i) \cdot i = i$	$i$	$i \cdot (-i) = 1$	$i^2 = -1$
$k = 2$	$(-1)^3 = -1$	$-1$	$(-1)^2 = 1$	$(-1)^2 = 1$
$k = 3$	$(-i) \cdot i \cdot (-i) = -i$	$-i$	$(-i) \cdot i = 1$	$(-i)^2 = -1$

Note that the coefficients in every column sum to 0, except for the coefficients for  $B(x, y)$  which sum to 4. Hence  $\sum_{k=0}^3 i^k B(i^k x + y, i^k x + y) = 4B(x, y)$ .

**Solution to 146 IV.** Let  $X$  be a set together with a subbase  $B$ . Write  $\Phi$  for the filter generated by  $B$ . Note  $B \subseteq \Phi$ . We will show  $(X, \Phi)$  is a uniform space, by proving its axioms in order.

1. By construction  $\Phi$  is a filter.
2. Pick any  $\varepsilon \in \Phi$ . We have to show  $\Delta \equiv \{(x, x); x \in X\} \subseteq \varepsilon$ . By definition of  $\Phi$ , there are  $\varepsilon_1, \dots, \varepsilon_n \in B$  with  $\varepsilon_1 \cap \dots \cap \varepsilon_n \subseteq \varepsilon$ . By definition of a base, we have  $\Delta \subseteq \varepsilon_i$  for each  $1 \leq i \leq n$  and so  $\Delta \subseteq \varepsilon_1 \cap \dots \cap \varepsilon_n \subseteq \varepsilon$  as well.
3. Pick any  $\varepsilon \in \Phi$ . By definition of  $\Phi$ , there are  $\varepsilon_1, \dots, \varepsilon_n \in B$  with  $\varepsilon_1 \cap \dots \cap \varepsilon_n \subseteq \varepsilon$ . As  $B$  is a base, there are  $\delta_1, \dots, \delta_n$  with  $\delta_i \circ \delta_i \subseteq \varepsilon_i$  for  $1 \leq i \leq n$ . Define  $\delta = \delta_1 \cap \dots \cap \delta_n$ . Then  $\delta \circ \delta \subseteq \bigcap_{i,j} \delta_i \circ \delta_j \subseteq \bigcap_i \delta_i \subseteq \bigcap_i \varepsilon_i \subseteq \varepsilon$ .
4. Pick any  $\varepsilon \in \Phi$ . By definition of  $\Phi$ , there are  $\varepsilon_1, \dots, \varepsilon_n \in B$  with  $\varepsilon_1 \cap \dots \cap \varepsilon_n \subseteq \varepsilon$ . As  $B$  is a base, there are  $\delta_1, \dots, \delta_n$  with  $\delta_i^{-1} \subseteq \varepsilon_i$ . Define  $\delta = \delta_1 \cap \dots \cap \delta_n$ . Then  $\delta^{-1} = \bigcap_i \delta_i^{-1} \subseteq \bigcap_i \varepsilon_i \subseteq \varepsilon$ .

Thus indeed,  $(X, \Phi)$  is a uniform space.

**Solution to 147 II.** We prove the subexercises in order.

1. First we will show that equivalence of Cauchy nets is an equivalence relation. As for every entourage  $\varepsilon$  we have  $x \varepsilon x$ , we see that every Cauchy net is equivalent to itself. Assume  $(x_\alpha)_\alpha \sim (y_\beta)_\beta$ . We will show  $(y_\beta)_\beta \sim (y_\alpha)_\alpha$ . Let  $\varepsilon$  be some entourage. There is some entourage  $\delta$  with  $\delta^{-1} \subseteq \varepsilon$ . By assumption there are  $\alpha_0$  and  $\beta_0$  such that  $x_\alpha \delta y_\beta$  for all  $\alpha \geq \alpha_0$  and  $\beta \geq \beta_0$ .

But then  $y_\beta \varepsilon x_\alpha$  for  $\alpha \geq \alpha_0$  and  $\beta \geq \beta_0$ . Hence  $(y_\beta)_\beta \sim (x_\alpha)_\alpha$ . To prove transitivity, assume we are given Cauchy nets  $(x_\alpha)_\alpha \sim (y_\beta)_\beta \sim (z_\gamma)_\gamma$ . Let  $\varepsilon$  be some entourage. There is an entourage  $\delta$  with  $\delta \circ \delta \subseteq \varepsilon$ . There are  $\alpha_0, \beta_0, \gamma_0$  such that  $x_\alpha \delta y_\beta$  and  $y_\beta \delta z_\gamma$  for  $\alpha \geq \alpha_0, \beta \geq \beta_0$  and  $\gamma \geq \gamma_0$ . Hence  $x_\alpha \varepsilon z_\gamma$  for such  $\alpha$  and  $\gamma$ , which shows  $(x_\alpha)_\alpha \sim (z_\gamma)_\gamma$ .

Next, assume that  $(x_\alpha)_\alpha$  is a subnet of a Cauchy net  $(y_\alpha)_\alpha$ . To show  $(x_\alpha)_\alpha \sim (y_\alpha)_\alpha$ , assume  $\varepsilon$  is some entourage. By the definition of Cauchy net, there is some  $\alpha_0$  such that  $x_\alpha \varepsilon x_\beta$  for all  $\alpha, \beta \geq \alpha_0$ . In particular  $x_\alpha \varepsilon y_\beta$  for all  $\alpha, \beta \geq \alpha_0$  which shows  $(x_\alpha)_\alpha \sim (y_\beta)_\beta$ .

2. Assume  $(x_\alpha)_\alpha \sim (y_\beta)_\beta$  and  $x_\alpha \rightarrow x$ . To prove  $y_\alpha \rightarrow x$ , pick any entourage  $\varepsilon$ . Pick  $\delta$  such that  $\delta^2 \subseteq \varepsilon$ . There are  $\alpha_0$  and  $\beta_0$  such that  $y_\beta \delta x_\alpha$  and  $x_\alpha \delta x$  for all  $\alpha \geq \alpha_0$  and  $\beta \geq \beta_0$ . Then  $y_\beta \varepsilon x$  for all  $\beta \geq \beta_0$ , whence  $y_\beta \rightarrow x$ .
3. Assume  $(x_\alpha)_\alpha \rightarrow x$  and  $(x_\alpha)_\alpha \rightarrow y$  in some Hausdorff uniform space. Let  $\varepsilon$  be any entourage. Pick  $\delta$  and  $\delta'$  with  $\delta^2 \subseteq \varepsilon$  and  $\delta' \subseteq \delta^{-1}$ . There is an  $\alpha_0$  such that  $x_\alpha \delta x$  and  $x_\alpha \delta' y$  for all  $\alpha \geq \alpha_0$ . Thus  $x \varepsilon y$ . As our space is Hausdorff the previous implies  $x = y$ .
4. Let  $f: X \rightarrow Y$  be a continuous map between uniform spaces. Assume  $x_\alpha \rightarrow x$  in  $X$ . Let  $\varepsilon$  be any entourage of  $Y$ . By continuity there is a  $\delta$  such that  $x \delta y$  implies  $f(x) \varepsilon f(y)$  for any  $y \in Y$ . There is an  $\alpha_0$  such that  $x \delta x_\alpha$  for all  $\alpha \geq \alpha_0$ . For those  $\alpha$  we also have  $f(x) \varepsilon f(x_\alpha)$ , which shows  $f(x_\alpha) \rightarrow f(x)$ .
5. Let  $f: X \rightarrow Y$  be a uniformly continuous map between uniform spaces. Let  $(x_\alpha)_\alpha$  and  $(y_\beta)_\beta$  be nets of  $X$  such that for each entourage  $\varepsilon$  of  $X$  there are  $\alpha_0, \beta_0$  with  $x_\alpha \varepsilon y_\beta$  for all  $\alpha \geq \alpha_0$  and  $\beta \geq \beta_0$ . The map  $f$  preserves this relation between the nets in the following way. Let  $\varepsilon$  be any entourage of  $Y$ . By uniform continuity there is a  $\delta$  such that  $x \delta y$  implies  $f(x) \varepsilon f(y)$ . There are  $\alpha_0$  and  $\beta_0$  with  $x_\alpha \delta y_\beta$  for all  $\alpha \geq \alpha_0$  and  $\beta \geq \beta_0$ . Hence  $f(x_\alpha) \varepsilon f(y_\beta)$  for such  $\alpha, \beta$ .

From the previous it follows that  $f$  preserves Cauchy nets (by setting  $x_\alpha = y_\alpha$ ) and that it preserves equivalence between Cauchy nets.

6. Let  $D \subseteq X$  be a dense subset of a uniform space  $X$ . Let  $x \in X$  be any point. Pick for every  $\varepsilon \in \Phi$  an element  $d_\varepsilon \in D$  with  $x \varepsilon d_\varepsilon$ . Clearly  $(d_\varepsilon)_{\varepsilon \in \Phi}$  is a net with inverse inclusion. We have  $d_\varepsilon \rightarrow x$  as  $d_\delta \varepsilon x$  whenever  $\delta \subseteq \varepsilon$ .
7. Assume  $f, g: X \rightarrow Y$  are continuous maps between uniform spaces that agree on a dense subset  $D \subseteq X$ . Let  $x \in X$  be any point. Pick a net  $x_\alpha$  from  $D$  with  $x_\alpha \rightarrow x$ . Then  $f(x) = f(\lim_\alpha x_\alpha) = \lim_\alpha f(x_\alpha) = \lim_\alpha g(x_\alpha) = g(\lim_\alpha x_\alpha) = g(x)$ . Hence  $f = g$ .

**Solution to 147 III.** Write  $B \equiv \{\hat{\varepsilon}; \varepsilon \in \Phi_i; i \in I\}$ . First we show  $B$  is a subbase, i.e. that it satisfies conditions 2, 3 and 4 of 146 II. Let  $\hat{\varepsilon}$  be an arbitrary element of  $B$  and  $i \in I$  denote the index element with  $\varepsilon \in \Phi_i$ . Assume  $(x_i)_{i \in I} \in \prod_i X_i$ . Clearly  $x_i \varepsilon x_i$  and so  $(x_i)_i \hat{\varepsilon} (x_i)_i$ . Thus  $B$  satisfies condition 2 of 146 II. Pick a  $\delta \in \Phi_i$  with  $\delta^2 \subseteq \varepsilon$ . Then  $\hat{\delta}^2 = \widehat{\delta^2} \subseteq \hat{\varepsilon}$  and so  $B$  satisfies condition 3 of 146 II. Now let  $\delta$  denote an entourage of  $X_i$  with  $\delta^{-1} \subseteq \varepsilon$ . Then  $\hat{\delta}^{-1} = \widehat{\delta^{-1}} \subseteq \hat{\varepsilon}$  and so  $B$  also satisfies condition 4 of 146 II.

Next we show that the projectors  $\pi_i: \prod_i X_i \rightarrow X_i$  are uniformly continuous. Assume  $i_0 \in I$  and  $\varepsilon$  is an entourage of  $X_{i_0}$ . Define  $\delta \equiv \hat{\varepsilon}$ . Let  $(x_i)_i$  and  $(y_i)_i$  from  $\prod_i X_i$  be given with  $(x_i)_i \delta (y_i)_i$ . Then  $x_{i_0} \varepsilon y_{i_0}$ . Thus  $\pi_{i_0}$  is indeed uniformly continuous.

To show  $(\pi_i)_i$  is a categorical product, assume we are given a uniform space  $Y$  together with uniformly continuous maps  $f_i: Y \rightarrow X_i$  for each  $i \in I$ . We have to show that there is a unique uniformly continuous map  $f: Y \rightarrow \prod_i X_i$  with  $\pi_i \circ f = f_i$  for all  $i \in I$ . Define  $f$  by  $(f(y))_i \equiv f_i(y)$ . Clearly  $\pi_i \circ f = f_i$ .

To show  $f$  is uniformly continuous, pick any entourage  $\varepsilon$  of  $\prod_i X_i$ . By definition, there are  $i_1, \dots, i_n$  and  $\varepsilon_1, \dots, \varepsilon_n$  with  $\varepsilon_j \in \Phi_{i_j}$  and  $\bigcap_j \hat{\varepsilon}_j \subseteq \varepsilon$ . For each  $1 \leq j \leq n$  pick an entourage  $\delta_j$  of  $Y$  such that  $x \delta_j y$  implies  $f_{i_j}(x) \varepsilon_j f_{i_j}(y)$ . Define  $\delta \equiv \bigcap_j \delta_j$ . Assume  $x \delta y$ . Then for each  $1 \leq j \leq n$  we have  $f_{i_j}(x) \varepsilon_j f_{i_j}(y)$  and so  $f(x) \hat{\varepsilon}_j f(y)$ , hence  $f(x) \varepsilon f(y)$ . Thus  $f$  is uniformly continuous.

To show uniqueness of  $f$ , assume there is a uniformly continuous map  $f': Y \rightarrow \prod_i X_i$  with  $\pi_i \circ f' = f_i$ . Then  $(f'(y))_i = f_i(y) = (f(y))_i$  for every  $y \in Y$  and so  $f' = f$ .

**Solution to 148 IV.** Let  $\mathcal{B}$  be a von Neumann algebra and  $X$  a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product. We will show that  $x \mapsto xb$  is ultranorm continuous for any  $b \in \mathcal{B}$ . It is sufficient to show it is ultranorm continuous at 0, so assume  $x_\alpha \rightarrow 0$  ultranorm for some net  $x_\alpha$  in  $X$ . Let  $f: \mathcal{B} \rightarrow \mathbb{C}$  be any np-map. Then  $\|x_\alpha b\|_f^2 = f([x_\alpha b, x_\alpha b]) = f(b^*[x_\alpha, x_\alpha]b) \leq \|b^*b\|f([x_\alpha, x_\alpha]) = \|b\|^2\|x_\alpha\|_f^2 \rightarrow 0$  thus  $x_\alpha b \rightarrow 0$  ultranorm as well.

**Solution to 149 III.** For brevity, write  $p \equiv \langle e, e \rangle$ . Note  $\|ep - e\|^2 = \|e(1-p)\|^2 = \langle e(1-p), e(1-p) \rangle = (1-p)\langle e, e \rangle(1-p) = (1-p)p(1-p) = 0$ . Thus  $ep - e = 0$  and so  $ep = e$  as desired.

**Solution to 149 IV.** Let  $X$  be a pre-Hilbert  $\mathcal{B}$ -module for a von Neumann algebra  $\mathcal{B}$  with orthonormal basis  $E \subseteq X$ . Assume  $x \in X$ . By definition of orthonormal basis, we know  $x = \sum_{e \in E} e \langle e, x \rangle$  where the sum converges ultranorm. That is:  $\sum_{e \in S} e \langle e, x \rangle \rightarrow x$  as  $S$  ranges over the finite subsets of  $E$ . By 148 V we see  $\langle \sum_{e \in S} e \langle e, x \rangle, \sum_{e \in S} e \langle e, x \rangle \rangle \rightarrow \langle x, x \rangle$  ultraweakly. Thus  $\sum_{e \in S} \langle x, e \rangle \langle e, x \rangle = \sum_{e, d \in S} \langle x, e \rangle \langle e, d \rangle \langle d, x \rangle = \langle \sum_{e \in S} e \langle e, x \rangle, \sum_{e \in S} e \langle e, x \rangle \rangle \rightarrow \langle x, x \rangle$  ultraweakly, as desired.

**Solution to 152 VIII.** Let  $T: X \rightarrow Y$  be a bounded  $\mathcal{B}$ -linear map between Hilbert  $\mathcal{B}$ -modules. Assume  $X$  is self dual. For any  $y \in Y$ , the map  $x \mapsto \langle y, Tx \rangle$  is  $\mathcal{B}$ -linear and bounded and so by self-duality of  $X$ , there is a  $t_y \in X$  with  $\langle t_y, x \rangle = \langle y, Tx \rangle$  for all  $x \in X$ . For any  $z, y \in Y$  and  $x \in X$ , we have  $\langle t_z + t_y, x \rangle = \langle t_z, x \rangle + \langle t_y, x \rangle = \langle z, Tx \rangle + \langle y, Tx \rangle = \langle z + y, Tx \rangle = \langle t_{z+y}, x \rangle$ . Thus  $t_z + t_y = t_{z+y}$ . For any  $\lambda \in \mathbb{C}$ ,  $y \in Y$  and  $x \in X$  we have  $\langle \lambda t_y, x \rangle = \langle \lambda y, Tx \rangle = \langle t_{\lambda y}, x \rangle$  and so  $\lambda t_y = t_{\lambda y}$ . Hence  $T^*y \equiv t_y$  defines a linear map from  $Y$  to  $X$  with  $\langle y, Tx \rangle = \langle t_y, x \rangle \equiv \langle T^*y, x \rangle$  for all  $x \in X$  and  $y \in Y$ . So  $T^*$  is the adjoint of  $T$ .

**Solution to 152 IX.** Let  $V$  be a right  $\mathcal{B}$ -module with  $\mathcal{B}$ -valued inner product for some von Neumann algebra  $\mathcal{B}$ . Write  $\eta: V \rightarrow X$  for the ultranorm completion of  $V$  from 150 II. Let  $T \in \mathcal{B}^a(X)$  be given with  $\langle \hat{x}, T\hat{x} \rangle \geq 0$  for all  $x \in V$ . We have to show  $T \geq 0$ . Let  $x \in X$  be an arbitrary vector. As all vector states on  $\mathcal{B}^a(X)$  are order separating by 148 VII, it is sufficient to show  $\langle x, Tx \rangle \geq 0$ . As the image of  $V$  under  $\eta$  is ultranorm dense in  $X$ , we can find a net  $x_\alpha$  with  $\widehat{x_\alpha} \rightarrow x$ . Then by 148 V we get  $\langle x, Tx \rangle = \text{uwl}\lim_\alpha \langle \widehat{x_\alpha}, T\widehat{x_\alpha} \rangle \geq 0$ . So indeed  $T \geq 0$ , as desired.

**Solution to 153 IV.** Let  $\mathcal{A}$  be a  $C^*$ -algebra with  $a_1, \dots, a_n \in \mathcal{A}$ . Define  $\varphi: \mathcal{A} \rightarrow M_n \mathcal{A}$  by  $\varphi(d) \equiv (a_i^* da_j)_{ij}$ . We have to show  $\varphi$  is an ncp-map. Recall  $\mathcal{A}$  is a self-dual Hilbert  $\mathcal{A}$ -module with  $\langle a, b \rangle \equiv a^*b$  and so its  $n$ -fold direct product  $\mathcal{A}^n$  is also self dual (see also 160 II.) Define  $T: \mathcal{A}^n \rightarrow \mathcal{A}$  by  $T((b_i)_i) \equiv \sum_i b_i a_i$  (i.e.  $T$  is the row-vector  $(a_i)_i$ .) Clearly  $T$  is  $\mathcal{A}$ -linear. It is also bounded:  $\|T(b_i)_i\|^2 = \sum_i \|b_i a_i\|^2 \leq \sum_i A \|b_i\|^2 = A \|(b_i)_i\|^2$ , where  $A \equiv \max_i \|a_i\|^2$ . It's easy that  $T^*(b) \equiv (a_i^* b)_i$  is the adjoint of  $T$ . We may identify  $\mathcal{B}^a(\mathcal{A}^n) = M_n$  and then  $\text{ad}_T(d) ((b_i)_i) = T^* d T (b_i)_i = T^* d \sum_i a_i b_i = (\sum_i (a_j^* da_i) b_i)_j = \varphi(d) ((b_i)_i)$ . Thus  $\text{ad}_T = \varphi$ . By 153 I the map  $\text{ad}_T$  and thus  $\varphi$  is as well.

**Solution to 160 II.** Assume  $X$  and  $Y$  are self-dual Hilbert  $\mathcal{B}$ -modules over a von Neumann algebra  $\mathcal{B}$  with orthonormal bases  $E \subseteq X$  and  $F \subseteq Y$ . Write  $G \equiv \{(e, 0); e \in E\} \cup \{(0, d); d \in D\}$ . Clearly  $G$  is orthonormal. To show  $G$  is an orthonormal basis, two conditions remain. For the first, let  $(x, y) \in X \oplus Y$  be given. As  $E$  and  $F$  are orthonormal bases we know  $x = \sum_{e \in E} e \langle e, x \rangle$  and  $y = \sum_{f \in F} f \langle f, y \rangle$ , where the sums converge ultranorm. The inclusions  $x \mapsto (x, 0)$

and  $y \mapsto (0, y)$  are bounded  $\mathcal{B}$ -linear and thus ultranorm continuous, hence

$$\begin{aligned}
\sum_{g \in G} g \langle g, (x, y) \rangle &= \left( \sum_{e \in E} (e, 0) \langle (e, 0), (x, y) \rangle \right) + \left( \sum_{f \in F} (0, f) \langle (0, f), (x, y) \rangle \right) \\
&= \left( \sum_{e \in E} (e, 0) \langle e, x \rangle \right) + \left( \sum_{f \in F} (0, f) \langle f, y \rangle \right) \\
&= \left( \sum_{e \in E} e \langle e, x \rangle, \sum_{f \in F} f \langle f, y \rangle \right) \\
&= (x, y),
\end{aligned}$$

which proves the first condition. For the second condition, let  $(b_g)_{g \in G}$  be some  $\ell^2$ -summable family from  $\mathcal{B}$ . The subfamilies  $(b_{e_0})_{e \in E}$  and  $(b_{0,f})_{f \in F}$  are  $\ell^2$ -summable as well. Hence the sums  $\sum_{e \in E} e b_{(e,0)}$  and  $\sum_{f \in F} f b_{(0,f)}$  converge ultranorm. Thus  $\sum_{e \in E} (e, 0) b_{(e,0)} + \sum_{f \in F} (0, f) b_{(0,f)} = \sum_{g \in G} g b_g$  converges ultranorm as well. We have shown  $G$  is an orthonormal basis of  $X \oplus Y$ . Consequently  $X \oplus Y$  is self dual by 149V.

**Solution to 160IX.** Assume  $X$  is a self-dual Hilbert  $\mathcal{B}$ -module for a von Neumann algebra  $\mathcal{B}$ . Suppose  $E \subseteq X$  is an orthonormal set. We will show  $E$  is an orthonormal basis of  $E^{\perp\perp}$ . Clearly  $E$  is orthonormal. Because of this, and the fact that  $E^{\perp\perp}$  is ultranorm closed (by 160IV) we know  $\sum_e b b_e$  converges ultranorm in  $E^{\perp\perp}$  for any  $\ell^2$ -family  $(b_e)_e$ . Assume  $x \in E^{\perp\perp}$ . To show  $E$  is an orthonormal base, it only remains to be shown that  $x = \sum_e e \langle e, x \rangle$ . Define  $x' \equiv x - \sum_e e \langle e, x \rangle$ . By 160IV we know  $E^{\perp\perp}$  is ultranorm closed and so  $x' \in E^{\perp\perp}$ . For any  $e_0 \in E$  we also have  $\langle e_0, x' \rangle = \langle e_0, x \rangle - \sum_{e \in E} \langle e_0, e \rangle \langle e, x \rangle = 0$  and so  $x' \in E^\perp$ . Hence  $\langle x', x' \rangle = 0$ , so indeed  $x = \sum_e e \langle e, x \rangle$  and  $E$  is an orthonormal basis of  $E^{\perp\perp}$ .

For the second part, assume  $x \in X$ . By Parseval's identity (see 149IV) we have  $\langle x, x \rangle = \sum_e \langle x, e \rangle \langle e, x \rangle$  for any  $x \in E^{\perp\perp}$ . To prove the converse, assume  $\langle x, x \rangle = \sum_e \langle x, e \rangle \langle e, x \rangle$ . By 160IV we know  $x = x' + x''$  for  $x' \in E^{\perp\perp}$  and  $x'' \in E^\perp$ . Note  $\langle e, x' \rangle = \langle e, x \rangle$  for any  $e \in E$  and so by Parseval's identity for  $E^{\perp\perp}$  we see  $\langle x', x' \rangle = \sum_e \langle x, e \rangle \langle e, x \rangle = \langle x, x \rangle$ . Now, using  $\langle x'', x' \rangle = 0$  we see  $\langle x, x \rangle = \langle x'', x'' \rangle + \langle x', x' \rangle = \langle x'', x'' \rangle + \langle x, x \rangle$  and so  $\langle x'', x'' \rangle = 0$ , whence  $x \in E^{\perp\perp}$ .

**Solution to 160X.** Let  $X$  be a self-dual Hilbert  $\mathcal{B}$ -module for some von Neumann algebra  $\mathcal{B}$ . Assume  $x_1, \dots, x_n \in X$ . We will show that there is a finite orthonormal set  $E$  of  $n$  or fewer elements such that  $E$  is a basis of  $\{x_1, \dots, x_n\}^{\perp\perp}$ . We do this by induction over  $n$ . For  $n = 0$ , the set  $E = \emptyset$  suffices. For the induction step, assume  $n > 0$  and  $E'$  is an orthonormal basis of  $\{x_1, \dots, x_{n-1}\}^{\perp\perp}$ . Write  $x' \equiv x_n - \sum_{e \in E'} e \langle e, x_n \rangle$ . If  $x' = 0$ , then  $E \equiv E'$  suffices. For the other case, assume  $x' \neq 0$ . By polar decomposition (see the end of 149VIII), there is an  $u \in X$  with  $x' = u \langle x', x' \rangle^{\frac{1}{2}}$  and  $\langle u, u \rangle = \lceil \langle x', x' \rangle \rceil$ . Define  $E \equiv E' \cup \{u\}$ .

Clearly  $E$  is an orthonormal set of  $n$  or fewer elements. By the induction assumption and 160 IX, we know  $x_i \in E'^{\perp\perp} \subseteq E^{\perp\perp}$  for  $i \leq 1 \neq n-1$ . For any  $d \in E^\perp$  we have  $\langle d, x' \rangle = \langle d, u \rangle \langle x', x' \rangle^{\frac{1}{2}} = 0$  and so  $x' \in E^{\perp\perp}$ . Clearly  $\sum_{e \in E'} e \langle e, x_n \rangle \in E'^{\perp\perp} \subseteq E^{\perp\perp}$  and so  $x_n = x' + \sum_{e \in E'} e \langle e, x_n \rangle \in E^{\perp\perp}$ . Together with the previous, we see  $\{x_1, \dots, x_n\}^{\perp\perp} \subseteq E^{\perp\perp\perp\perp} = E^{\perp\perp} \subseteq \{x_1, \dots, x_n\}^{\perp\perp}$ , thus  $E^{\perp\perp} = \{x_1, \dots, x_n\}^{\perp\perp}$ . By 160 IX we know  $E$  is an orthonormal basis of  $E^{\perp\perp}$ , which completes the proof by induction.

**Solution to 161 II.** For brevity write  $\ell^2 \equiv \ell^2((p_i)_{i \in I})$ . We will first proof that  $\ell^2$  is a right  $\mathcal{B}$ -module. Assume  $(a_i)_i, (b_i)_i \in \ell^2$ . We want to show  $(a_i + b_i)_i \in \ell^2$ . First, we to show  $\sum_i (a_i + b_i)^*(a_i + b_i)$  is bounded. Pick  $A, B \in \mathbb{R}^+$  with  $\sum_i a_i^* a_i \leq A$  and  $\sum_i b_i^* b_i \leq B$  which exist as  $(a_i)_i$  and  $(b_i)_i$  are  $\ell^2$ . Let  $f$  be any normal state on  $\mathcal{B}$  and  $S \subseteq I$  some finite subset. Then

$$\begin{aligned} f\left(\sum_{i \in S} (a_i + b_i)^*(a_i + b_i)\right) &= \sum_{i \in S} \|a_i + b_i\|_f^2 \\ &\leq \sum_{i \in S} \|a_i\|_f^2 + \|b_i\|_f^2 + 2\|a_i\|_f \|b_i\|_f \\ &\leq A + B + 2 \sum_{i \in S} \|a_i\|_f \|b_i\|_f. \end{aligned}$$

By Cauchy–Schwarz  $\sum_{i \in S} \|a_i\|_f \|b_i\|_f \leq (\sum_{i \in S} \|a_i\|_f^2)^{\frac{1}{2}} (\sum_{i \in S} \|b_i\|_f^2)^{\frac{1}{2}} \leq (AB)^{\frac{1}{2}}$ . As normal states are order separating, we see that we have a bounded and thus converging sum of positive elements  $\sum_i (a_i + b_i)^*(a_i + b_i) \leq A + B + (AB)^{\frac{1}{2}}$ . Suppose  $i \in I$ . It remains to be shown  $\lceil (a_i + b_i)(a_i + b_i)^* \rceil \leq p_i$ . Recall from 59 VI that  $\lceil xx^* \rceil \leq p_i$  if and only if  $p_i x = x$ . Clearly  $p_i(a_i + b_i) = p_i a_i + p_i b_i = a_i + b_i$  as  $(a_i)_i, (b_i)_i \in \ell^2$  and so indeed  $(a_i + b_i)_i \in \ell^2$ .

Suppose  $b \in \mathcal{B}$  and  $(a_i)_i \in \ell^2$ . Then  $b^*(\sum_i a_i^* a_i)b = \sum_i (a_i b)^* a_i b$  as  $x \mapsto b^* x b$  is normal by 44 VIII and so  $(a_i b)_i$  is  $\ell^2$ . Furthermore  $p_i a_i b = a_i b$  for any  $i \in I$ , so  $(a_i b)_i \in \ell^2$ . We have shown  $\ell^2$  is a right  $\mathcal{B}$ -module with coordinatewise operations.

Next we will show  $\langle (a_i)_i, (b_i)_i \rangle \equiv \sum_i a_i^* b_i$  defines a  $\mathcal{B}$ -valued inner product on  $\ell^2$ . First we have to show the sum converges. Suppose  $f$  is any normal state on  $\mathcal{B}$ . Pick  $A, B \in \mathbb{R}^+$  with  $\sum_i a_i^* a_i \leq A$  and  $\sum_i b_i^* b_i \leq B$ . For any finite subset  $S \subseteq I$  we have

$$\left| f\left(\sum_{i \in S} a_i^* b_i\right) \right| \leq \sum_{i \in S} |[a_i, b_i]_f| \leq \sum_{i \in S} \|a_i\|_f \|b_i\|_f \leq (AB)^{\frac{1}{2}},$$

where we used Cauchy–Schwarz for  $\mathcal{B}$ -valued inner products in the second inequality and classic Cauchy–Schwarz in the final inequality. We have shown that  $\sum_{i \in S} a_i^* b_i$  is a norm-bounded net in  $S$ . We claim it is ultraweakly Cauchy as well. For now, pick any finite sets  $S, T \subseteq I$ . Assume  $f$  is any normal state

on  $\mathcal{B}$ . We want to show that the following quantity vanishes for sufficiently large  $S \cap T$ .

$$\left| f\left(\sum_{i \in S} a_i^* b_i - \sum_{i \in T} a_i^* b_i\right) \right| \leq \left| \sum_{i \in S-T} f(a_i^* b_i) \right| + \left| \sum_{i \in T-S} f(a_i^* b_i) \right|. \quad (4.2)$$

Note that  $[(a_i)_i, (b_i)_i] \equiv \sum_{i \in S-T} f(a_i^* b_i)$  is an inner product and so

$$\left| \sum_{i \in S-T} f(a_i^* b_i) \right| \leq \left( \sum_{i \in S-T} f(a_i^* a_i) \right)^{\frac{1}{2}} \left( \sum_{i \in S-T} f(b_i^* b_i) \right)^{\frac{1}{2}}. \quad (4.3)$$

The sum  $\sum_i f(a_i^* a_i)$  converges and so  $\sum_{i \in S-T} f(a_i^* a_i)$  can be made arbitrarily small by picking sufficiently large  $S \cap T$ . And so (with a similar argument for the other factor), we see that (4.3) vanishes, which is the left term of (4.2). The argument for the other term of (4.2) is the same. Thus  $\sum_{i \in S} a_i^* b_i$  is ultraweakly Cauchy and converges by 76 III. From the fact that  $a \mapsto a^*$ ,  $a \mapsto ab$  and  $(a, b) \mapsto a + b$  are all ultraweakly continuous, it follows readily that  $\langle (a_i)_i, (b_i)_i \rangle \equiv \sum_i a_i^* b_i$  is an inner product on  $\ell^2$ . We claim this inner product is definite. Assume  $0 = \langle (a_i)_i, (a_i)_i \rangle \equiv \sum_i a_i^* a_i$ . Then  $a_i^* a_i = 0$  for each  $i \in I$  and so  $a_i = 0$ , which is to say  $(a_i)_i = 0$ . Indeed our inner product is definite. Thus  $\ell^2$  is a pre-Hilbert  $\mathcal{B}$ -module.

Write  $E \equiv \{\delta_i; i \in I\}$ , where  $(\delta_i)_j = 0$  for  $i \neq j$  and  $(\delta_i)_i = p_i$ . Clearly  $E$  is an orthonormal set. We claim it's an orthonormal basis of  $\ell^2$ . Assume  $(a_i)_i \in \ell^2$ . It is easy to see  $\sum_{i \in I} \delta_i a_i = (a_i)_i$  and so  $\sum_{\delta_i \in E} \delta_i \langle \delta_i, (a_i)_i \rangle = \sum_{i \in I} \delta_i a_i = (a_i)_i$ . It remains to be shown that  $\sum_{\delta_i \in E} \delta_i b_{\delta_i}$  converges ultranorm for any  $\ell^2$  summable  $(b_i)_{\delta_i \in E}$ , which indeed it does to  $(b_{\delta_i})_{i \in I}$  as we already saw. Thus  $\ell^2$  is self dual.

For the final part of the exercise, assume  $X$  is a self-dual Hilbert  $\mathcal{B}$ -module over some von Neumann algebra  $\mathcal{B}$  with orthonormal basis  $E \subseteq X$ . Define  $\vartheta: X \rightarrow \ell^2(((e, e))_{e \in E})$  by  $\vartheta(x) \equiv ((e, x))_e$ . Clearly  $\vartheta$  is  $\mathcal{B}$ -linear. It also preserves the inner product:  $\langle x, y \rangle = \langle x, \sum_e e \langle e, y \rangle \rangle = \sum_e \langle x, e \langle e, y \rangle \rangle = \sum_e \langle x, e \rangle \langle e, y \rangle = \langle \vartheta(x), \vartheta(y) \rangle$ . which entails it's injective. To show  $\vartheta$  is surjective, let  $(x_e)_e \in \ell^2(((e, e))_{e \in E})$  be given. The family  $(x_e)_e$  is  $\ell^2$  so  $\sum_e e b_e$  converges ultranorm. Clearly  $\vartheta(\sum_e e b_e) = ((e, \sum_e e b_e))_e = (b_e)_e$ , so  $\vartheta$  is indeed surjective. It follows  $\vartheta$  is an isomorphism  $X \cong \ell^2(((e, e))_{e \in E})$ .

**Solution to 161 IV.** Let  $X$  be a self-dual Hilbert  $\mathcal{B}$ -module  $X$  for some von Neumann algebra  $\mathcal{B}$  with orthonormal basis  $(e_i)_{i \in I}$ . Assume  $(u_i)_{i \in I}$  is a family partial isometries with  $u_i u_i^* = \langle e_i, e_i \rangle$ . We will show  $(e_i u_i)_{i \in I}$  is an orthonormal basis of  $X$ . For brevity, write  $d_i \equiv e_i u_i$ . To start, note  $\langle d_i, d_j \rangle = u_i^* \langle e_i, e_j \rangle u_j$ , which is zero if  $i \neq j$ . If  $i = j$ , then  $\langle d_i, d_i \rangle = u_i^* \langle e_i, e_i \rangle u_i = u_i^* u_i$  which is a projection. So  $(d_i)_i$  is orthonormal. Hence  $\sum_i d_i b_i$  converges for any  $\ell^2$ -family  $(d_i)_i$ . It remains to be shown that  $x = \sum_i d_i \langle d_i, x \rangle$ . Note  $\sum_i d_i \langle d_i, x \rangle =$

$\sum_i e_i \langle e_i u_i u_i^*, x \rangle = \sum_i e_i \langle e_i \langle e_i, e_i \rangle, x \rangle = \sum_i e_i \langle e_i, x \rangle = x$ , so indeed  $(d_i)_i$  is an orthonormal basis of  $X$ .

For the next part, assume  $(p_i)_{i \in I}$  and  $(q_i)_{i \in I}$  are projections with  $p_i \sim q_i$ . Let  $p_i$  denote the partial projection with  $u_i^* u_i = q_i$  and  $u_i u_i^* = p_i$ . Consider  $\ell^2((p_i)_{i \in I})$ . Define  $((\delta_i)_i) \in \ell^2$  by  $(\delta_i)_j = 0$  if  $i \neq j$  and  $(\delta_i)_i = p_i$ . In 161 II we saw  $(\delta_i)_i$  is an orthonormal basis of  $\ell^2((p_i)_{i \in I})$ . Note  $u_i u_i^* = p_i = \langle \delta_i, \delta_i \rangle$ . By the previous  $\delta_i u_i$  is another orthonormal basis of  $\ell^2((p_i)_{i \in I})$ . By the second part of 161 II we see  $\ell^2((p_i)_{i \in I}) \cong \ell^2((\langle \delta_i u_i, \delta_i u_i \rangle)_{i \in I}) = \ell^2((u_i^* p_i u_i)_{i \in I}) = \ell^2((q_i)_{i \in I})$ , as promised.

**Solution to 161 V.** Let  $X$  be a self-dual Hilbert  $\mathcal{B}$ -module. Assume  $E \subseteq X$  and  $e_1, e_2 \in X$  such that  $E \cup \{e_1, e_2\}$  is an orthonormal basis and  $\langle e_1, e_1 \rangle + \langle e_2, e_2 \rangle \leq 1$ . We will show that  $E' \equiv E \cup \{e_1 + e_2\}$  is an orthonormal basis as well. For brevity, write  $p_1 \equiv \langle e_1, e_1 \rangle$  and  $p_2 \equiv \langle e_2, e_2 \rangle$ . Clearly  $E$  itself is an orthonormal set. For any  $e \in E$  we have  $\langle e, e_1 + e_2 \rangle = \langle e, e_1 \rangle + \langle e, e_2 \rangle = 0$  and so  $E'$  is an orthogonal set. By assumption  $p_1$  and  $p_2$  are projections with  $p_1 + p_2 \leq 1$  and so by 55 XIII they are orthogonal and thus in particular  $p_1 + p_2$  is again a projection. Hence  $\langle e_1 + e_2, e_1 + e_2 \rangle = p_1 + p_2$  is a projection. Thus  $E'$  is an orthonormal set.

Let  $x \in X$  be given. By 149 III we have  $e_2 = e_2 p_2$  and so  $e_2 \langle e_1 + e_2, x \rangle = e_2 \langle (e_1 + e_2) p_2, x \rangle = e_2 \langle e_2, x \rangle$ . Similarly  $e_1 \langle e_1 + e_2, x \rangle = e_1 \langle e_1, x \rangle$ . Thus  $x = e_1 \langle e_1, x \rangle + e_2 \langle e_2, x \rangle + \sum_{e \in E} e \langle e, x \rangle = (e_1 + e_2) \langle e_1 + e_2, x \rangle + \sum_{e \in E} e \langle e, x \rangle$ , which shows the first condition on an orthonormal set to be an orthonormal basis. The second (and final) condition holds automatically as  $E'$  is an orthonormal set and  $X$  is ultranorm complete. Thus  $E'$  is indeed an orthonormal basis.

The the last part of the exercise, assume  $p, q \in \mathcal{B}$  are projections with  $p + q \leq 1$ . Clearly  $\{p + q\}$  is an orthonormal basis of  $(p + q)\mathcal{B}$  and so is  $\{p, q\}$  by the previous. Hence by 161 II we see  $(p + q)\mathcal{B} \cong \ell^2(\{p, q\}) = p\mathcal{B} \oplus q\mathcal{B}$ , as promised.

**Solution to 165 V.** Assume  $X$  is a self-dual Hilbert  $\mathcal{A}$ -module and  $Y$  is a self-dual Hilbert  $\mathcal{B}$ -module for von Neumann algebras  $\mathcal{A}$  and  $\mathcal{B}$ .

1. Assume  $x_1, x_2, x \in X$  and  $y_1, y_2, y \in Y$ . Then

$$\begin{aligned} |x_1\rangle\langle x_2| \otimes |y_1\rangle\langle y_2| x \otimes y &= (|x_1\rangle\langle x_2|x) \otimes (|y_1\rangle\langle y_2|y) \\ &= (x_1 \langle x_2, x \rangle) \otimes (y_1 \langle y_2, y \rangle) \\ &= (x_1 \otimes y_1)(\langle x_2, x \rangle \otimes \langle y_2, y \rangle) \\ &= (x_1 \otimes y_1)\langle x_2 \otimes y_2, x \otimes y \rangle \\ &= |x_1 \otimes y_1\rangle\langle x_2 \otimes y_2|x \otimes y. \end{aligned}$$

This is sufficient to show that  $|x_1\rangle\langle x_2| \otimes |y_1\rangle\langle y_2| = |x_1 \otimes y_1\rangle\langle x_2 \otimes y_2|$ , either by appealing to the defining universal property of  $X \otimes Y$  or by the related property that the  $\mathcal{A} \odot \mathcal{B}$ -linear span of  $\{x \otimes y; x \in X, y \in Y\}$  is

ultranorm dense in  $X \otimes Y$ . (In fact the  $\mathbb{C}$ -linear span is already ultranorm dense.)

2. Clearly, for any  $x \in X$  and  $y \in Y$  we have  $(1 \otimes 1)(x \otimes y) = 1x \otimes y$  and so by same token (as the conclusion of the previous point) we see  $1 = 1 \otimes 1$ .
3. Assume  $x \in X, y \in Y, S, S' \in \mathcal{B}^a(X)$  and  $T, T' \in \mathcal{B}^a(Y)$ . Then

$$(S \otimes T)(S' \otimes T')x \otimes y = (SS'x) \otimes (TT'y) = (SS') \otimes (TT')x \otimes y,$$

which is sufficient to show  $(SS') \otimes (TT') = (S \otimes T) \otimes (S' \otimes T')$  (see the conclusion of the first point.)

4. Assume  $S \in \mathcal{B}^a(X), T \in \mathcal{B}^a(Y), x_1, x_2 \in X$  and  $y_1, y_2 \in Y$ . Then

$$\begin{aligned} \langle (S \otimes T)^* x_1 \otimes y_1, x_2 \otimes y_2 \rangle &= \langle x_1 \otimes y_1, (S \otimes T)x_2 \otimes y_2 \rangle \\ &= \langle x_1 \otimes y_1, (Sx_2) \otimes (Ty_2) \rangle \\ &= \langle x_1, Sx_2 \rangle \otimes \langle y_1, Ty_2 \rangle \\ &= \langle S^*x_1, x_2 \rangle \otimes \langle T^*y_1, y_2 \rangle \\ &= \langle (S^* \otimes T^*)x_1 \otimes y_1, x_2 \otimes y_2 \rangle. \end{aligned}$$

This is sufficient (see the conclusion of the first point) to show that the vector functionals for  $(S \otimes T)^*x_1 \otimes y_1$  and  $(S^* \otimes T^*)x_1 \otimes y_1$  agree. Hence  $(S \otimes T)^*x_1 \otimes y_1 = (S^* \otimes T^*)x_1 \otimes y_1$ . In turn this is sufficient to show  $(S \otimes T)^* = S^* \otimes T^*$ , as desired.

**Solution to 169 XI.** Assume  $\varphi: \mathcal{A} \rightarrow \mathcal{B}$  is an ncp-map between von Neumann algebras.

1. Assume  $c: \mathcal{B} \rightarrow \mathcal{C}$  is a filter and that  $(\mathcal{P}, \varrho, h)$  is a Paschke dilation of  $\varphi$ . We will show that  $(\mathcal{P}, \varrho, c \circ h)$  is a Paschke dilation of  $c \circ \varphi$ . To this end, assume  $\varrho: \mathcal{A} \rightarrow \mathcal{P}'$  is some nmiu-map and  $h': \mathcal{P}' \rightarrow \mathcal{C}$  is any ncp-map with  $h' \circ \varrho' = c \circ \varphi$ . Assume  $\varphi(1) \leq 1$ . (We will reduce the general to this one later on.) Note  $h'(1) = h'(\varrho'(1)) = c(\varphi(1)) \leq c(1)$  and so by the universal property of  $c$ , there is a unique ncp-map  $h'': \mathcal{P}' \rightarrow \mathcal{B}$  with  $c \circ h'' = h'$ . Then  $c \circ \varphi = h' \circ \varrho' = c \circ h'' \circ \varrho'$ . And so  $\varphi = h'' \circ \varrho'$  as  $c$  is injective, see 169 XII. There is a unique ncp-map  $\sigma: \mathcal{P}' \rightarrow \mathcal{P}$  with  $h \circ \sigma = h''$  and  $\sigma \circ \varrho' = \varrho$ . Then  $h' = c \circ h'' = c \circ h \circ \sigma$ . Assume  $\sigma': \mathcal{P}' \rightarrow \mathcal{P}$  is any ncp-map with  $\sigma' \circ \varrho = \varrho$  and  $c \circ h \circ \sigma' = h'$ . Then  $c \circ h \circ \sigma' = h' = c \circ h''$  and so  $h \circ \sigma' = h''$ . Hence  $\sigma = \sigma'$  by the uniqueness of  $\sigma$ . Thus  $(\mathcal{P}, \varrho, c \circ h)$  is a Paschke dilation of  $c \circ \varphi$ .

Now, in the general case, assume  $\varphi(1) \not\leq 1$ . Then  $\varphi(1) \neq 0$ . Define  $\varphi' \equiv \|\varphi(1)\|^{-1}\varphi$ . Then  $\varphi'(1) = \|\varphi(1)\|^{-1}\varphi(1) \leq 1$ . By 140 X we know  $(\mathcal{P}, \varrho, \|\varphi(1)\|^{-1}h)$  is a Paschke dilation of  $\varphi'$ . Now we are in our

previous case and so  $(\mathcal{P}, \varrho, \|\varphi(1)\|^{-1}c \circ h)$  is a Paschke dilation of  $c \circ \varphi'$ . Applying 140X once again, we conclude  $(\mathcal{P}, \varrho, c \circ h)$  is a Paschke dilation of  $c \circ \varphi$  as desired.

2. Assume  $c': \mathcal{C}' \rightarrow \mathcal{B}$  is a filter of  $\varphi(1)$ . By the universal property of  $c'$ , there is a unique ncp-map  $\varphi': \mathcal{A} \rightarrow \mathcal{C}'$  with  $c' \circ \varphi' = \varphi$ . Then  $c'(\varphi'(1)) = \varphi(1) = c'(1)$  and so by injectivity of  $c'$  (see 169XII) we get  $\varphi'(1) = 1$  — that is to say:  $\varphi'$  is unital.

For the final part of this exercise, assume  $(\mathcal{P}, \varrho, h)$  is a Paschke dilation of  $\varphi'$ . By the previous point, we know  $(\mathcal{P}, \varrho, c' \circ h)$  is a Paschke dilation of  $c' \circ \varphi' = \varphi$ , as desired.

**Solution to 169XII.** Let  $c: \mathcal{C} \rightarrow \mathcal{B}$  be some filter for  $b \in \mathcal{B}$ . By 169X we know  $c_b: [b] \mathcal{B} [b] \rightarrow \mathcal{B}$  given by  $a \mapsto \sqrt{ba}\sqrt{b}$  is also a filter for  $b$ . By the universal property of  $c$ , there is a unique ncp-map  $\vartheta_1$  with  $c \circ \vartheta_1 = c_b$ . In the other direction, by the universal property of  $c_b$ , there is a unique ncp-map  $\vartheta_2$  with  $c_b \circ \vartheta_2 = c$ . Hence  $c \circ (\vartheta_1 \circ \vartheta_2) = c_b \circ \vartheta_2 = c \circ \text{id}$ . Clearly  $c(1) \leq c(1)$  and so by the universal property of  $c$  the map  $\text{id}$  is the unique ncp-map with  $c \circ \text{id} = c$ , hence  $\vartheta_1 \circ \vartheta_2 = \text{id}$ . Reasoning similarly on the other side, we see  $\vartheta_2 \circ \vartheta_1 = \text{id}$  and so  $\vartheta_2$  is an isomorphism. Recall  $c = c_b \circ \vartheta_2$  and so it is sufficient to show  $c_b$  is injective. To this end, let  $a_1, a_2 \in [b] \mathcal{B} [b]$  be given with  $c_b(a_1) = c_b(a_2)$ ; i.e.  $\sqrt{ba_1}\sqrt{b} = \sqrt{ba_2}\sqrt{b}$ . Then  $a_1 = a_2$  by 60VIII and so  $c_b$  is indeed injective.

**Solution to 170IV.** Assume  $\varrho: \mathcal{A} \rightarrow \mathcal{B}$  is a surjective nmiu-map between von Neumann algebras. We already saw in 138III that the kernel of an nmiu-map is an ultraweakly-closed two-sided ideal and hence a principal ideal of a central projection by 69II. Write  $z$  for the central projection of  $\mathcal{A}$  with  $\ker \varrho = (1-z)\mathcal{A}$ ;  $h_z: \mathcal{A} \rightarrow z\mathcal{A}$  for the map  $h_z(a) = za$  and  $\varrho': z\mathcal{A} \rightarrow \mathcal{B}$  for the restriction of  $\varrho$  to  $z\mathcal{A}$ . Clearly  $\varrho' \circ h_z = \varrho$  and so  $\varrho'$  is surjective. Furthermore  $\varrho'$  is injective, because for any  $a \in z\mathcal{A}$  with  $\varrho(a) = 0$  we have  $\varrho(a) = 0$  and so  $a \in \ker \varrho = (1-z)\mathcal{A}$ , hence  $a = 0$ . It is easy to see that  $\varrho'$  as a bijective miu-map has an miu inverse and so it is an miu-isomorphism. Consequently, it is an nmiu-isomorphism (as miu-maps are order-preserving) and in particular an ncp-isomorphism. The map  $h_z$  is a standard corner for  $z$  (see 169IV) and so  $\varrho = \varrho' \circ h_z$  is also a corner for  $z$  as  $\varrho'$  is an ncp-isomorphism.

Conversely, assume  $h: \mathcal{A} \rightarrow \mathcal{C}$  is a corner for a central projection  $z \in \mathcal{A}$ . The map  $h_z: \mathcal{A} \rightarrow z\mathcal{A}$  given by  $a \mapsto za$  is another corner for  $z$ , see 169IV. By the universal property of  $h$ , there is a unique ncp-map  $\vartheta_1: z\mathcal{A} \rightarrow \mathcal{C}$  with  $h \circ \vartheta_1 = h_z$ . On the other side, by the universal property of  $h_z$ , there is a unique ncp-map  $\vartheta_2: \mathcal{C} \rightarrow z\mathcal{A}$  with  $h_z \circ \vartheta_2 = \vartheta_1$ . Again, by the universal property of  $h$ , the map  $\text{id}: \mathcal{C} \rightarrow \mathcal{C}$  is the unique ncp-map  $\mathcal{C} \rightarrow \mathcal{C}$  with  $h = h \circ \text{id}$ . Note  $h = h_z \circ \vartheta_2 = h \circ (\vartheta_1 \circ \vartheta_2)$  and so  $\text{id} = \vartheta_1 \circ \vartheta_2$ . Similarly  $\vartheta_2 \circ \vartheta_1 = \text{id}$ . Thus  $\vartheta_1$  is an ncp-isomorphism and consequently an nmiu-

isomorphism by 99 IX. Clearly  $h_z$  is a surjective nmiu-map and thus  $h = h_z \circ \vartheta_2$  is a surjective nmiu-map as well.

**Solution to 174 III.** Reflexivity is easy:  $a \otimes 0 = 0 \otimes a = a$  by the partial commutativity and zero axiom, so  $a \leq a$ . For transitivity, assume  $a \leq b$  and  $b \leq c$ . Pick  $d, e \in M$  with  $a \otimes d = b$  and  $b \otimes e = c$ . Then by partial associativity  $c = b \otimes e = (a \otimes d) \otimes e = a \otimes (d \otimes e)$  so  $a \leq c$ .

**Solution to 175 III.** Assume  $E$  and  $F$  are effect algebras. We define  $a \otimes b$  for  $a, b \in E \times F$  whenever both  $a_1 \perp b_1$  and  $a_2 \perp b_2$  and in that case  $a \otimes b \equiv (a_1 \otimes b_1, a_2 \otimes b_2)$ . Furthermore, define  $1 \equiv (1, 1)$  and  $0 \equiv (0, 0)$ . We claim that these operations turn  $E \times F$  into an effect algebra.

We start with partial commutativity. Assume  $a, b \in E \times F$  with  $a \perp b$ . Then  $a_1 \perp b_1$  and  $a_2 \perp b_2$ , hence  $b_1 \perp a_1, b_2 \perp a_2$  and  $a \otimes b = (a_1 \otimes b_1, a_2 \otimes b_2) = (b_1 \otimes a_1, b_2 \otimes a_2) = b \otimes a$ .

We continue with partial associativity. Assume  $a, b, c \in E \times F$  with  $a \perp b$  and  $a \otimes b \perp c$ . Then  $a_i \perp b_i, a_i \otimes b_i \perp c_i$  for  $i = 1, 2$  and so  $b_i \perp c_i$  and  $(a_i \otimes b_i) \otimes c_i = a_i \otimes (b_i \otimes c_i)$ . Consequently  $(a \otimes b) \otimes c = a \otimes (b \otimes c)$ .

Next the zero axiom: for any  $a \in E \times F$  we have  $a_1 \perp 0, a_2 \perp 0, 0 \otimes a_1 = a_1$  and  $0 \otimes a_2 = a_2$ . Thus  $0 \equiv (0, 0) \perp (a_1, a_2)$  and  $0 \otimes a = a$ .

To prove the orthocomplement law, assume  $a \in E \times F$ . Then for  $a \equiv (a_1^\perp, a_2^\perp)$  we have  $a \perp a^\perp$  and  $a \otimes a^\perp = (1, 1) \equiv 1$ . Furthermore, if  $a \vee b = 1$  for some  $b \in E \times F$ , then  $b_1 = a_1^\perp$  and  $b_2 = a_2^\perp$ , hence  $b = a^\perp$ .

Only the zero-one law remains. Assume  $a \perp 1$  for some  $a \in E \times F$ . Then  $a_1 \perp 1$  and  $a_2 \perp 1$ , hence  $a_1 = 0$  and  $a_2 = 0$ . Thus  $a = 0$ , as desired. We have shown  $E \times F$  is an effect algebra with componentwise operations.

Write  $\pi_1: E \times F \rightarrow E$  and  $\pi_2: E \times F \rightarrow F$  for the maps given by  $\pi_1(a_1, a_2) = a_1$  and  $\pi_2(a_1, a_2) = a_2$ . Clearly  $\pi_1$  and  $\pi_2$  are additive and unital, hence effect algebra homomorphisms. We will show that  $E \times F$  with projections  $\pi_1$  and  $\pi_2$  forms a categorical product of  $E$  and  $F$  in EA. To this end, assume  $f_1: G \rightarrow E$  and  $f_2: G \rightarrow F$  are effect algebra homomorphisms for some effect algebra  $G$ . Define  $g: G \rightarrow E \times F$  by  $g(a) = (f_1(a), f_2(a))$ . Clearly  $g$  is the unique map with  $\pi_i \circ g = f_i$  for  $i = 1, 2$ . It is also an effect algebra homomorphism and so  $E \times F$  is indeed the product of  $E$  and  $F$ .

**Solution to 175 IV.** Let  $E$  be an effect algebra with any  $a \in E$ . We will first show that  $1^\perp = 0$  and  $a^{\perp\perp} = a$ . By the orthocomplement law  $1 \otimes 1^\perp = 1$ , so  $1^\perp \perp 1$ , hence  $1^\perp = 0$  by the zero-one law. By the orthocomplement law  $a \otimes a^\perp = 1$ . Thus  $a^\perp \otimes a = a \otimes a^\perp = 1$  by partial commutativity. Hence  $a = a^{\perp\perp}$  by uniqueness of the orthocomplement. Now, to prove the zero law, note  $a^\perp \otimes a^{\perp\perp} = 1 = (a^\perp \otimes a^{\perp\perp}) \otimes (a^\perp \otimes a^{\perp\perp})^\perp = (a^\perp \otimes a^{\perp\perp}) \otimes 0 = a^\perp \otimes (a^{\perp\perp} \otimes 0)$  by the orthocomplement law and partial associativity. Thus by uniqueness of the orthocomplement  $a^{\perp\perp} = a^{\perp\perp} \otimes 0$ , hence  $0 \otimes a = a$  by partial commutativity and  $a^{\perp\perp} = a$ .

**Solution to 176II.** We start with (D1). Assume  $b \leq a$ . Then  $b \otimes c = a$  for some  $c$ . By definition  $c = a \ominus b$ . Conversely, assume  $a \ominus b$  is defined. Then  $b \otimes (b \ominus a) = a$ , hence  $b \leq a$ . To prove (D2), assume  $a \ominus b$  is defined. Then  $a \ominus b \leq b \otimes (a \ominus b) = a$  as desired. For (D3) assume  $a \ominus b$  and  $a \ominus (a \ominus b)$  are defined. Note  $(a \ominus b) \otimes b = b \otimes (a \ominus b) = a$  and so  $b = a \ominus (a \ominus b)$ .

(D4) is next. Assume  $a \leq b \leq c$ . Then  $(c \ominus b) \otimes (b \ominus a) \otimes a = c = (c \ominus a) \otimes a$  and so by cancellation  $c \ominus b \leq (c \ominus b) \otimes (b \ominus a) = c \ominus a$ . Note  $((c \ominus a) \ominus (c \ominus b)) \otimes (c \ominus b) \otimes a = c = (b \ominus a) \otimes a \otimes (c \ominus b)$  and so by cancellation  $(c \ominus a) \ominus (c \ominus b) = b \ominus a$ .

For the final part of the exercise, assume  $E$  is a poset with maximum element 1 and a partial operation  $\ominus$  satisfying (D1), (D2), (D3) and (D4). We will show  $E$  is an effect algebra with  $a \otimes b = c \Leftrightarrow a = c \ominus b$  and  $0 = 1 \ominus 1$ .

First we have to show that if  $c_1 \ominus b = c_2 \ominus b$  for some  $b \leq c_1, c_2$  in  $E$ , then  $c_1 = c_2$  so that  $\otimes$  is at most single-valued. By (D4) we have  $(1 \ominus b) \ominus (1 \ominus c_1) = c_1 \ominus b$  and so  $(1 \ominus b) \ominus a = (1 \ominus b) \ominus (c_1 \ominus b) = (1 \ominus b) \ominus ((1 \ominus b) \ominus (1 \ominus c_1)) = 1 \ominus c_1$  by (D3). Similarly  $1 \ominus c_2 = (1 \ominus b) \ominus a$ , so with another application of (D3) we see  $c_1 = 1 \ominus (1 \ominus c_1) = 1 \ominus (1 \ominus c_2) = c_2$ .

The first axiom we will prove is partial commutativity. Assume  $a, b \in E$  with  $a \perp b$ . By definition, there is some  $c \geq b$  with  $c \ominus b = a$ . By (D3) we have  $c \ominus a = c \ominus (c \ominus b) = b$  and so  $b \perp a$  with  $b \otimes a = c = a \otimes b$  as desired. Note that we have also shown that  $a \leq a \otimes b$  and  $(a \otimes b) \ominus a = b$  for  $a \perp b$ .

Next, to prove partial associativity, assume  $a, b, c \in E$  with  $a \perp b$  and  $a \otimes b \perp c$ . Then  $a \leq a \otimes b \leq (a \otimes b) \otimes c$  and thus

$$\begin{aligned} b &= (a \otimes b) \ominus a \\ &\stackrel{(D4)}{=} (((a \otimes b) \otimes c) \ominus a) \ominus (((a \otimes b) \otimes c) \ominus (a \otimes b)) \\ &= (((a \otimes b) \otimes c) \ominus a) \ominus c, \end{aligned}$$

which shows  $b \perp c$  with  $b \otimes c = ((a \otimes b) \otimes c) \ominus a$ . From that it immediately follows that  $b \otimes c \perp a$  with  $(b \otimes c) \otimes a = (a \otimes b) \otimes c$  and so  $a \perp b \otimes c$  with  $(a \otimes b) \otimes c = a \otimes (b \otimes c)$  by partial commutativity.

We continue with the zero axiom. Assume  $a \in E$ . Clearly  $a \leq 1 \leq 1$  and so  $(1 \ominus a) \ominus (1 \ominus 1) = 1 \ominus a$  by (D4), which shows  $1 \ominus a \perp 1 \ominus 1 \equiv 0$  with  $(1 \ominus a) \otimes 0 = 1 \ominus a$ . The zero law follows by partial commutativity.

Now we prove the orthocomplement law. Assume  $a \in E$ . By (D3) we have  $1 \ominus (1 \ominus a) = a$ , which demonstrates  $a \perp 1 \ominus a \equiv a^\perp$  with  $a \otimes a^\perp = 1$ . Assume  $b \in E$  with  $a \otimes b = 1$ . By definition, we have  $a = 1 \ominus b$  and so by (D3) we see  $b = 1 \ominus (1 \ominus b) = 1 \ominus a \equiv a^\perp$ .

To show the last axiom (zero-one), assume  $a \in E$  with  $a \perp 1$ . Then  $1 \leq a \otimes 1 \leq 1$  and so  $a \otimes 1 = 1$ . Hence  $a = (a \otimes 1) \ominus 1 = 1 \ominus 1 \equiv 0$ . We have shown  $E$  is an effect algebra.

To conclude this exercise, we show that the original order and  $\ominus$  coincide with those defined for  $E$  as an effect algebra. For clarity, write  $\leq_{EA}$  and  $\ominus_{EA}$  for the latter.

To start, assume  $a \leq b$  for some  $a, b \in E$  (as a D-poset). Then  $b = a \circledast (b \ominus a)$  as trivially  $b \ominus a = b \ominus a$ . Hence  $a \leq_{\text{EA}} b$ . To prove the converse, assume  $a \leq_{\text{EA}} b$ . Then  $a \circledast c = b$  for some  $c \in E$ . Hence  $a \leq a \circledast c \leq b$ . Indeed  $\leq = \leq_{\text{EA}}$ .

Assume  $a, b \in E$ . The element  $a \ominus_{\text{EA}} b$  is defined if and only if  $b \leq a$ , which is precisely when  $a \ominus b$  is defined. Furthermore  $b \circledast (a \ominus_{\text{EA}} b) = a$  and so  $a \ominus_{\text{EA}} b = (b \circledast (a \ominus_{\text{EA}} b)) \ominus b = a \ominus b$ .

**Solution to 176 V.** Assume  $f: E \rightarrow F$  is an homomorphism between effect algebras.

1. We have  $0 \perp 1$  and so  $f(0) \perp f(1) = 1$ , hence  $f(0) = 0$  by the zero-one law.
2. Assume  $a \leq b$  in  $E$ . Then  $a \circledast c = b$  for some  $c \in E$ . Hence  $f(b) = f(a \circledast c) = f(a) \circledast f(c)$ , which shows  $f(a) \leq f(b)$ .
3. Assume  $a \ominus b$  is defined (i.e.  $a \geq b$ ). Recall  $(a \ominus b) \circledast b = a$  and so  $f(a \ominus b) \circledast f(b) = f(a)$ . Hence  $f(a \ominus b) = f(a) \ominus f(b)$ .
4. Assume  $a \in E$ . In 176 II we saw  $1 \ominus a = a^\perp$  and so by the previous point  $f(a^\perp) = f(1 \ominus a) = f(1) \ominus f(a) = 1 \ominus f(a) = f(a)^\perp$ .

**Solution to 178 IIIa.** Let  $M$  be an effect monoid with any  $a \in M$ . Then  $a \circledast 1 = a \circledast (0 \circledast 1) = (a \circledast 0) \circledast (a \circledast 1)$ . Hence by cancellation  $a \circledast 0 = 0$ . Similarly  $0 \circledast a = 0$ .

**Solution to 178 V.** Assume  $M$  is an effect monoid with  $a_1, \dots, a_n, b_1, \dots, b_n \in M$  such that  $\bigcircledast_i a_i = 1$  and  $\bigcircledast_i a_i \circledast b_i = 1$ . Note that for any  $1 \leq i \leq n$  we have  $a_i \circledast b_i \leq a_i$  as  $(a_i \circledast b_i) \circledast a_i \circledast b_i^\perp = a_i \circledast 1 = a_i$ . Thus

$$a_i^\perp = \bigcircledast_{j \neq i} a_j \geq \bigcircledast_{j \neq i} a_j \circledast b_j = (a_i \circledast b_i)^\perp \geq a_i^\perp$$

and so  $a_i^\perp = (a_i \circledast b_i)^\perp$ , hence  $a_i = a_i \circledast b_i$ .

**Solution to 181 IX.** Let  $C$  be an effectus in partial form. Assume  $f: X \rightarrow A$  and  $g: X \rightarrow B$  are any arrows in  $C$  with  $1 \circ f \perp 1 \circ g$ .

1. Assume  $[a, b]: A + B \rightarrow Y$  is any map. Then  $1 \circ a \circ f \leq 1 \circ f \perp 1 \circ g \geq 1 \circ b \circ g$  and so  $1 \circ a \circ f \perp 1 \circ b \circ g$ . Hence  $a \circ f \perp b \circ g$  and so  $[a, b] \circ \langle f, g \rangle = [a, b] \circ ((\kappa_1 \circ f) \circledast (\kappa_2 \circ g)) = ([a, b] \circ (\kappa_1 \circ f)) \circledast ([a, b] \circ (\kappa_2 \circ g)) = (a \circ f) \circledast (b \circ g)$ .
2. By the previous point and 181 IV, we have  $1 \circ \langle f, g \rangle = [1, 1] \circ \langle f, g \rangle = (1 \circ f) \circledast (1 \circ g)$ .
3. Assume  $k: A \rightarrow A'$  and  $l: B \rightarrow B'$  are two maps in  $C$ . Then by the first point of this exercise, we have  $(k + l) \circ \langle f, g \rangle \equiv [\kappa_1 \circ k, \kappa_2 \circ l] \circ \langle f, g \rangle = (\kappa_1 \circ k \circ f) \circledast (\kappa_2 \circ l \circ g) \equiv \langle k \circ f, l \circ g \rangle$ .

4. Assume  $k: X' \rightarrow X$  is some map in  $C$ . Then by PCM-enrichment of  $C$ , we get  $\langle f, g \rangle \circ k \equiv ((\kappa_1 \circ f) \oplus (\kappa_2 \circ g)) \circ k = (\kappa_1 \circ f \circ k) \oplus (\kappa_2 \circ g \circ k) \equiv \langle f \circ k, g \circ k \rangle$ .

**Solution to 183 II.** Assume  $h_1, h_2: Q \rightarrow P$  are two arrows with  $m_1 \circ h_1 = m_1 \circ h_2$  and  $m_2 \circ h_1 = m_2 \circ h_2$ . Note  $(f \circ m_1) \circ h_1 = g \circ m_2 \circ h_1 = (g \circ m_2) \circ h_2$  and so by the universal property of a pullback there is a unique map  $h: Q \rightarrow P$  with  $m_1 \circ h = m_1 \circ h_1$  and  $m_2 \circ h = m_2 \circ h_2$  and so  $h_1 = h_2 = h$ .

**Solution to 183 III.** For the first point, assume the left and right inner squares are pullbacks. To prove the outer square is a pullback, assume  $\alpha: S \rightarrow C$  and  $\beta: S \rightarrow X$  are any maps with  $g' \circ f' \circ \beta = m \circ \alpha$ . As the right inner square is a pullback, there is a unique map  $\gamma: S \rightarrow B$  with  $g \circ \gamma = \alpha$  and  $l \circ \gamma = f' \circ \beta$ . By this last equality and the fact that the left inner square is a pullback, there is a unique map  $\delta: S \rightarrow A$  with  $f \circ \delta = \gamma$  and  $k \circ \delta = \beta$ . We will prove that this  $\delta$  is also the mediating map for the outer square. Clearly  $g \circ f \circ \delta \circ g \circ \gamma = \alpha$ . Assume  $\delta': S \rightarrow A$  is some other map with  $g \circ f \circ \delta' = \alpha$  and  $k \circ \delta' \circ \beta$ . Note that  $g \circ f \circ \delta' = \alpha = g \circ \gamma$  and  $l \circ f \circ \delta' = f' \circ k \circ \delta' = f' \circ \beta = l \circ \gamma$ . Thus  $f \circ \delta' = \gamma$  as  $g$  and  $l$  are jointly monic by 183 II. Consequently  $\delta = \delta'$  and so the outer square is indeed a pullback.

For the second point, assume the outer square is a pullback and that  $l$  and  $g$  are jointly monic. We will prove that the left inner square is a pullback. To this end, assume  $\alpha: S \rightarrow B$  and  $\beta: S \rightarrow X$  are maps with  $l \circ \alpha = f' \circ \beta$ . Note  $m \circ g \circ \alpha = g' \circ l \circ \alpha = g' \circ f' \circ \beta$  and thus by the fact that the outer square is a pullback, there is a unique  $\gamma: S \rightarrow A$  with  $g \circ f \circ \gamma = g \circ \alpha$  and  $k \circ \gamma = \beta$ . This will also be the unique mediating map demonstrating that the left inner square is a pullback. We have  $l \circ f \circ \gamma = f' \circ k \circ \gamma = f' \circ \beta = l \circ \alpha$  and so by the joint monicity of  $l$  and  $g$ , we conclude  $f \circ \gamma = \alpha$ . Assume there is some map  $\gamma': S \rightarrow A$  with  $f \circ \gamma' = \alpha$  and  $k \circ \gamma' = \beta$ . Then clearly  $g \circ f \circ \gamma' = g \circ \alpha$  and so by the uniqueness of  $\gamma$  as a mediating map for the outer square, we find  $\gamma' = \gamma$ . Thus the left inner square is a pullback as well.

**Solution to 186 II.** Let  $f: X \rightarrow Z$  and  $g: Y \rightarrow Z$  be two partial maps in  $C$ ; that is  $f: X \rightarrow Z + 1$  and  $g: Y \rightarrow Z + 1$ . Then the cotuple  $[f, g]: X + Y \rightarrow Z + 1$  (with respect to the coproduct that we assumed to exist) is also a partial map  $X + Y \rightarrow Z$ . Note  $[f, g] \hat{\circ} \kappa_1 = [[f, g], \kappa_2] \circ \kappa_1 \circ \kappa_1 = f$  and  $[f, g] \hat{\circ} \kappa_2 = [[f, g], \kappa_2] \circ \kappa_1 \circ \kappa_2 = g$ . Suppose  $h: X + Y \rightarrow Z$  is a partial map with  $h \hat{\circ} \kappa_1 = f$  and  $h \hat{\circ} \kappa_2 = g$ . Then  $h \circ \kappa_1 = h \hat{\circ} \kappa_1 = f$  and  $h \circ \kappa_2 = h \hat{\circ} \kappa_2 = g$ , so  $h = [f, g]$ . This shows  $X + Y$  is also a coproduct in  $\text{Par } C$  with coprojectors  $\kappa_1$  and  $\kappa_2$ .

To show  $0$  is an initial object of  $\text{Par } C$ , assume  $X$  is some object of  $\text{Par } C$ , i.e. of  $C$ . As  $0$  is initial in  $C$ , there is a unique map  $!: 0 \rightarrow X + 1$  in  $C$  and so there is a unique map  $!: 0 \rightarrow X$  in  $\text{Par } C$ .

**Solution to 186 VII.** We already know that the initial object  $0$  of  $C$  is also initial

in  $\text{Par } C$ . We will show that it is also final in  $\text{Par } C$ . To this end, assume  $X$  is some object in  $\text{Par } C$ , i.e. of  $C$ . There is a unique map  $! : X \rightarrow 1$  in  $C$  and so there is a unique map  $! : X \rightarrow 0$  in  $\text{Par } C$ . Thus  $0$  is final in  $\text{Par } C$ .

For any objects  $X$  and  $Y$  from  $\text{Par } C$ , the zero map  $0 : X \rightarrow Y$  in  $\text{Par } C$  is by definition equal to  $i \hat{\circ} f$ , where  $f : X \rightarrow 0$  is the unique final map and  $i : 0 \rightarrow Y$  is the unique initial map. Unfolding definitions, we see  $0 = i \hat{\circ} f = [i, \kappa_2] \circ f = \kappa_2 \circ !$  as desired.

**Solution to 189I.** For the first point, assume  $C$  is an effectus in total form together with a map  $f : A \rightarrow 0$ . We will show that  $f$  is an isomorphism. By initiality of  $0$ , we know  $f \circ !_A = !_0 = \text{id}$ . By 185I, the following square is a pullback.

$$\begin{array}{ccc} 0 & \xlongequal{\quad} & 0 \\ \downarrow ! & \lrcorner & \parallel \\ A & \xrightarrow{f} & 0 \end{array}$$

As  $\text{id}_0 \circ f = f \circ \text{id}_A$ , there is a unique map  $h : A \rightarrow 0$  with  $\text{id} \circ h = f$  and  $! \circ h = \text{id}$ . So  $h = f$  and  $! \circ f = ! \circ h = \text{id}$ . Thus  $f$  is indeed an isomorphism with inverse  $!$ .

For the second point, assume  $C$  is an effectus in total and partial form. Let  $X$  and  $Y$  be any two objects. As  $0$  is final in an effectus in total form, there are maps  $!_X : X \rightarrow 0$  and  $!_Y : Y \rightarrow 0$ . As  $0$  is a strict initial object in an effectus in partial form, both  $!_X$  and  $!_Y$  are isomorphisms. Hence  $!_Y \circ !_X^{-1}$  is an isomorphism between  $X$  and  $Y$ .

**Solution to 191 VIII.** To show  $\text{Rng}^{\text{op}}$  is an effectus in total form, we shown the dual axioms for  $\text{Rng}$ . Clearly, for any two unit rings  $R$  and  $S$ , their cartesian product  $R \times S$  is a categorical product with projectors  $\pi_1(r, s) = r$  and  $\pi_2(r, s) = s$ . The integer ring  $\mathbb{Z}$  is the initial object of  $\text{Rng}$ . The zero ring (the unique unit ring with a single element  $0 = 1$ ) is the final object of  $\text{Rng}$ .

To show the pushout diagrams corresponding to (3.1) hold, assume we are given unit rings  $R, S, T$  with unit-preserving homomorphisms  $\alpha, \beta, \delta$  that make the outer squares of the following diagrams commute.

$$\begin{array}{ccc} T & \xleftarrow{\alpha} & R \times Z \\ \beta \swarrow & \text{dotted} & \nwarrow \\ R \times S & \xleftarrow{\text{id} \times !} & R \times Z \\ \uparrow ! \times \text{id} & & \uparrow ! \times \text{id} \\ \mathbb{Z} \times S & \xleftarrow{\text{id} \times !} & \mathbb{Z} \times \mathbb{Z} \end{array} \qquad \begin{array}{ccc} T & \xleftarrow{!} & Z \\ \delta \swarrow & \text{dotted} & \nwarrow \\ R & \xleftarrow{!} & Z \\ \uparrow \pi_1 & & \uparrow \pi_1 \\ R \times S & \xleftarrow{! \times !} & \mathbb{Z} \times \mathbb{Z} \end{array}$$

We have to show that there are unique dashed arrows (as shown) that make

these diagrams commute. We start with the left diagram. Define  $f: R \times S \rightarrow T$  by  $f(r, s) \equiv \alpha(r, 0) + \beta(0, s)$ . Clearly  $f$  is additive and  $f(0, 0) = \alpha(0, 0) + \beta(0, 0) = 0$ . By assumption  $\alpha(n, m) = \beta(n, m)$  for any  $n, m \in \mathbb{Z}$  and so  $\alpha(1, 0) = \beta(1, 0)$  in particular, hence  $f(1, 1) = \alpha(1, 0) + \beta(0, 1) = \beta(1, 1) = 1$ . It remains to be shown that  $f$  is multiplicative. First note that  $\alpha(1, 0)\beta(0, 1) = \beta(1, 0)\beta(0, 1) = \beta(0, 0) = 0$ . Thus  $\alpha(r, 0)\beta(0, s) = \alpha(r, 0)\alpha(1, 0)\beta(0, 1)\beta(0, s) = 0$  for any  $r \in R$  and  $s \in S$  and similarly  $\beta(0, s)\alpha(r, 0) = 0$ . Hence  $f(r, s)f(r', s') = \alpha(r, 0)\alpha(r', 0) + \beta(0, s)\beta(0, s') = \alpha(rr', 0) + \beta(0, ss') = f(rr', ss')$ . We have shown  $f$  is a unit-preserving ring homomorphism. Clearly  $f(r, m) = \alpha(r, 0) + \beta(0, m) = \alpha(r, 0) + \alpha(0, m) = \alpha(r, m)$  for any  $r \in R$  and  $m \in \mathbb{Z}$  hence  $\alpha = f \circ (\text{id} \times !)$ . Similarly  $\beta = f \circ (! \circ \text{id})$ . Assume  $f': R \times S \rightarrow T$  is any unit-preserving ring homomorphism with  $\alpha = f' \circ (\text{id} \times !)$  and  $\beta = f' \circ (! \circ \text{id})$ . Then clearly  $f'(r, 0) = \alpha(r, 0)$  and  $f'(0, s) = \beta(0, s)$ , so  $f'(r, s) = f'(r, 0) + f'(0, s) = \alpha(r, 0) + \beta(0, s) = f(r, s)$ . This shows the left square above is indeed a pushout.

We continue with the diagram on the right. By assumption  $\delta(n, m) = n$  for any  $n, m \in \mathbb{Z}$  hence  $\delta(0, s) = \delta(0, s)\delta(0, 1) = 0$  for any  $s \in S$ . Thus  $\delta(r, s) = \delta(r, 0) + \delta(0, s) = \delta(r, 0)$  for any  $r \in R$  and  $s \in S$ . Define  $g: R \rightarrow T$  by  $g(r) = \delta(r, 0)$ . Clearly  $g$  is additive, multiplicative and  $g(0, 0) = 0$ . Furthermore  $g(1) = \delta(1, 0) = \delta(1, 1) = 1$ , so it is a unit-preserving ring homomorphism. It is easy to see that  $g \circ \pi_1 = \delta$  and that  $g$  is the unique such unit-preserving ring homomorphism. We have shown the right diagram is a pushout too.

To show  $\text{Rng}^{\text{op}}$  is an effectus in total form, it only remains to be shown that  $\langle \pi_1, \pi_2, \pi_2 \rangle, \langle \pi_2, \pi_1, \pi_2 \rangle: \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}$  are jointly epic. So assume  $f, g: \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z} \rightarrow R$  are two unit-preserving ring homomorphisms for which we have  $f \circ \langle \pi_1, \pi_2, \pi_2 \rangle = g \circ \langle \pi_1, \pi_2, \pi_2 \rangle$  and  $f \circ \langle \pi_2, \pi_1, \pi_2 \rangle = g \circ \langle \pi_2, \pi_1, \pi_2 \rangle$ . By the first equality  $f(k, 0, 0) = g(k, 0, 0)$  and by the second  $f(0, l, 0) = g(0, l, 0)$  for any  $k, l \in \mathbb{Z}$ . In particular  $f(0, 0, 1) = 1 - f(1, 0, 0) - f(0, 1, 0) = 1 - g(1, 0, 0) - g(0, 1, 0) = g(0, 0, 1)$  and so  $f(0, 0, m) = mf(0, 0, 1) = mg(0, 0, 1) = g(0, 0, m)$  for any  $m \in \mathbb{Z}$ . Putting it all together:  $f(k, l, m) = f(k, 0, 0) = f(0, l, m) + f(0, 0, m) = g(k, 0, 0) = g(0, l, m) + g(0, 0, m) = g(k, l, m)$ . So  $f = g$  and so we have shown the required joint epicity. We have shown  $\text{Rng}^{\text{op}}$  is indeed an effectus in total form.

We continue with the two additional points. Let  $p: \mathbb{Z} \times \mathbb{Z} \rightarrow R$  be any predicate on  $R$ . Then  $p(1, 0) \in R$  with  $p(1, 0)^2 = p(1, 0^2) = p(1, 0)$  and so  $p(1, 0)$  is an idempotent. Furthermore  $p(0, 1) = p(1, 1) - p(1, 0) = 1 - p(1, 0)$ , so  $p$  is fixed by the idempotent  $p(1, 0)$ . Let  $e \in R$  be any idempotent. It is easy to see that  $p(n, m) \equiv ne + m(1 - e)$  is a unit-preserving ring homomorphism with  $p(1, 0) = e$ . Thus the predicates on  $R$  correspond to its idempotents. This also shows that the set of scalars of  $\text{Rng}^{\text{op}}$  is the two-element effect monoid  $\mathbf{2}$  as  $\mathbb{Z}$  has exactly two idempotents: 0 and 1.

Assume  $p \perp q$  for two predicates  $p, q: \mathbb{Z} \times \mathbb{Z} \rightarrow R$  on  $R$ . Then, by definition,  $p \perp q$  if there is a  $b: \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z} \rightarrow R$  with  $b(n, m, m) = p(n, m)$  and  $b(m, n, m) = q(n, m)$  for all  $n, m \in \mathbb{Z}$ . In particular  $p(1, 0)q(1, 0) = b(1, 0, 0)b(0, 1, 0) = 0$ ,

thus the idempotents corresponding to  $p$  and  $q$  are orthogonal. Conversely, if  $p(1, 0)$  and  $q(1, 0)$  are orthogonal, then  $b(k, l, m) = kp(1, 0) + lq(1, 0) + m(1 - p(1, 0) - q(1, 0))$  defines a unit-preserving ring homomorphism that shows  $p \perp q$  with  $(p \oplus q)(1, 0) = b(1, 1, 0) = p(1, 0) + q(1, 0)$ . From this it is also clear that  $p^\perp(1, 0) = 1 - p(1, 0)$ .

There are many examples to show unit-preserving ring homomorphisms need not be fixed on their value on idempotents (and thus that  $\mathbf{Rng}^{\text{op}}$  does not have separating predicates.) For instance, let  $R$  denote the unit ring of continuous real-valued functions on the unit interval  $[0, 1]$ . This ring has only two idempotents: the functions that are constant 0 and 1, which are also the zero and unit element (respectively) of the ring. For every  $x \in [0, 1]$  the map  $\delta_x: R \rightarrow \mathbb{R}$  given by  $\delta_x(f) = f(x)$  is a unit-preserving ring homomorphism. Clearly  $\delta_x(0) = 0 = \delta_y(0)$  and  $\delta_x(1) = 1 = \delta_y(1)$  for any  $x, y \in [0, 1]$  with  $x \neq y$ , but not  $\delta_x = \delta_y$ .

Finally, we treat the second and last point of the exercise. Reasoning towards contradiction, let  $f: \mathbb{Z}_2 \rightarrow \mathbb{Z}$  be any unit-preserving ring homomorphism. Then  $0 = f(0) = f(1 + 1) = f(1) + f(1) = 1 + 1 = 2$ , which is absurd. Thus there is no such homomorphism. However, there are two unit-preserving ring homomorphisms  $\mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}_2$  corresponding to the idempotents 0 and 1 in  $\mathbb{Z}_2$ . Hence the states cannot be separating (for otherwise there could at most be a single unit-preserving ring homomorphism  $\mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}_2$ .)

**Solution to 192 III.** We will first show that  $\mathcal{D}_M$  is a functor. To start, clearly

$$\mathcal{D}_M(\text{id})(p)(x) \equiv \bigvee_{y; \text{id}(y)=x} p(y) = p(x)$$

and so  $\mathcal{D}_M(\text{id}) = \text{id}$ . Let  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  be given. Then

$$\begin{aligned} \mathcal{D}_M(g \circ f)(p)(z) &\equiv \bigvee_{x; g(f(x))=z} p(x) \\ &= \bigvee_{y; g(y)=z} \bigvee_{x; f(x)=y} p(x) \\ &= \bigvee_{y; g(y)=z} ((\mathcal{D}_M f)(p))(y) \\ &= ((\mathcal{D}_M g)((\mathcal{D}_M f)(p)))(z) \end{aligned}$$

and so  $\mathcal{D}_M(g \circ f) = (\mathcal{D}_M g) \circ (\mathcal{D}_M f)$ .

Next, we show that  $\eta$  is a natural transformation. Let  $f: X \rightarrow Y$  be any

map. Then for any  $x_0 \in X$ , we have

$$\begin{aligned}
((\mathcal{D}_M f) \circ \eta_X)(x_0) &= \bigvee_{x; f(x)=y} (\eta_X(x_0))(x) \\
&= \bigvee_{x; \substack{f(x)=y \\ x=x_0}} 1 \\
&= (\eta_Y(f(x_0)))
\end{aligned}$$

and so  $\mathcal{D}_M f \circ \eta_X = \eta_Y \circ f$ , i.e.  $\eta$  is a natural transformation  $\text{id} \Rightarrow \mathcal{D}_M$ .

To show  $\mu$  is a natural transformation, assume  $f: X \rightarrow Y$  is some map,  $y \in Y$  and  $\Phi \in \mathcal{D}_M \mathcal{D}_M X$ . Then

$$\begin{aligned}
(\mathcal{D}_M f)(\mu_X(\Phi))(y) &= \bigvee_{x; f(x)=y} \mu_X(\Phi)(x) \\
&= \bigvee_{x; f(x)=y} \bigvee_p \Phi(p) \odot p(x) \\
&= \bigvee_p \bigvee_{x; f(x)=y} \Phi(p) \odot p(x) \\
&= \bigvee_p \Phi(p) \odot \bigvee_{x; f(x)=y} p(x) \\
&= \bigvee_p \Phi(p) \odot (\mathcal{D}_M f)(p)(y) \\
&= \bigvee_q \bigvee_{p; (\mathcal{D}_M f)(p)=q} \Phi(p) \odot q(y) \\
&= \bigvee_q \left( \bigvee_{p; (\mathcal{D}_M f)(p)=q} \Phi(p) \right) \odot q(y) \\
&= \bigvee_q (\mathcal{D}_M \mathcal{D}_M f)(\Phi)(q) \odot q(y) \\
&= \mu_Y((\mathcal{D}_M \mathcal{D}_M f)(\Phi))(y)
\end{aligned}$$

and so  $(\mathcal{D}_M f) \circ \mu_X = \mu_Y \circ (\mathcal{D}_M \mathcal{D}_M f)$ , hence  $\mu$  is a natural transformation  $\mathcal{D}_M \mathcal{D}_M \Rightarrow \mathcal{D}_M$ . We continue with the monad laws. Assume  $X$  is a set,  $x \in X$  and  $\aleph \in$

$\mathcal{D}_M \mathcal{D}_M \mathcal{D}_M X$ . Then

$$\begin{aligned}
\mu_X(\mu_{\mathcal{D}_M X}(\aleph))(x) &= \bigvee_p (\mu_{\mathcal{D}_M X})(\aleph)(p) \odot p(x) \\
&= \bigvee_p \bigvee_\Phi \aleph(\Phi) \odot \Phi(p) \odot p(x) \\
&= \bigvee_\Phi \aleph(\Phi) \odot \bigvee_p \Phi(p) \odot p(x) \\
&= \bigvee_\Phi \aleph(\Phi) \odot \mu_X(\Phi)(x) \\
&= \bigvee_p \bigvee_{\Phi; \mu_X(\Phi)=p} \aleph(\Phi) \odot p(x) \\
&= \bigvee_p (\mathcal{D}_M \mu_X)(\aleph)(p) \odot p(x) \\
&= \mu_X((\mathcal{D}_M \mu_X)(\aleph))(x)
\end{aligned}$$

and so  $\mu \circ \mu_{\mathcal{D}_M} = \mu \circ (\mathcal{D}_M)$ . For any  $x \in X$  and  $p \in \mathcal{D}_M X$ , we have

$$\mu_X(\eta_{\mathcal{D}_M X}(p))(x) = \bigvee_q \eta_{\mathcal{D}_M X}(p)(q) \odot q(x) = p(x)$$

and so  $\mu \circ \eta_{\mathcal{D}_M} = \text{id}$ . For the final monad law, assume  $x \in X$  and  $p \in \mathcal{D}_M X$ . Then

$$\begin{aligned}
\mu_X((\mathcal{D}_M \eta_X)(p))(x) &= \bigvee_q (\mathcal{D}_M \eta_X)(p)(q) \odot q(x) \\
&= \bigvee_q \bigvee_{y; \eta_X(y)=q} p(y) \odot q(x) \\
&= \bigvee_y p(y) \odot \eta_X(y)(x) \\
&= p(x)
\end{aligned}$$

and so  $\mu \circ \mathcal{D}_M \eta = \text{id}$ . We have shown  $(\mathcal{D}_M, \eta, \mu)$  is a monad.

Our next project is to show that  $\mathcal{Kl} \mathcal{D}_M$  is an effectus in total form. It is easy to see that the coproduct  $X + Y$  from  $\mathbf{Set}$  is also a coproduct in  $\mathcal{D}_M$  with coprojectors  $\eta \circ \kappa_i$ , where  $\kappa_i$  are the coprojectors for  $X + Y$  in  $\mathbf{Set}$ . (In fact, for any category  $C$  and monad  $T$ , the inclusion functor  $K: C \rightarrow \mathcal{Kl} T$  given by  $KX = X$  and  $Kf = \eta \circ f$  is a left adjoint and so  $K$  preserves colimits.) The

empty set is also the initial object of  $\mathcal{Kl} \mathcal{D}_M$ . As  $\mathcal{D}_M 1 \cong 1$ , the category  $\mathcal{Kl} \mathcal{D}_M$  has as final object the one-element set 1.

Now we will show that a square such as that on the left of (3.1) is a pullback. So assume  $\alpha: Z \rightarrow \mathcal{D}_M(X+1)$  and  $\beta: Z \rightarrow \mathcal{D}_M(1+Y)$  are maps with  $(\hat{!}+\hat{\text{id}}) \hat{\circ} \alpha = (\hat{\text{id}}+\hat{!}) \hat{\circ} \beta$ , where  $\hat{\circ}$  denotes the composition in  $\mathcal{Kl} \mathcal{D}_M$  and  $\hat{f} \equiv \eta \circ f$  the Kleisli embedding. Assume  $z \in Z$ . By assumption  $\alpha(z)(\kappa_2^*) = ((\hat{!}+\hat{\text{id}}) \hat{\circ} \alpha)(z)(\kappa_2^*) = ((\hat{\text{id}}+\hat{!}) \hat{\circ} \beta)(z)(\kappa_2^*) = \bigcirc_y \beta(z)(\kappa_2 y)$ , where  $*$  is the unique element of 1. We want to define  $\delta: Z \rightarrow \mathcal{D}_M(X+Y)$  by  $\delta(z)(\kappa_1 x) = \alpha(z)(\kappa_1 x)$  and  $\delta(z)(\kappa_2 y) = \beta(z)(\kappa_2 y)$ , but first need to check the image of  $\delta(z)$  sums to 1. Indeed  $\bigcirc_w \delta(z)(w) = \bigcirc_y \beta(z)(\kappa_2 y) \circ \bigcirc_x \alpha(z)(\kappa_1 x) = \alpha(z)(\kappa_2^*) \circ \bigcirc_x \alpha(z)(\kappa_1 x) = 1$ . Clearly  $\alpha = (\hat{\text{id}}+\hat{!}) \hat{\circ} \delta$  and  $\beta = (\hat{!}+\hat{\text{id}}) \hat{\circ} \delta$ . Suppose  $\delta': Z \rightarrow \mathcal{D}_M(X+Y)$  is some map with  $\alpha = (\hat{\text{id}}+\hat{!}) \hat{\circ} \delta'$  and  $\beta = (\hat{!}+\hat{\text{id}}) \hat{\circ} \delta'$ . Then  $\delta'(\kappa_1 x) = ((\hat{\text{id}}+\hat{!}) \hat{\circ} \delta')(\kappa_1 x) = \alpha(\kappa_1 x)$  for any  $x \in X$  and similarly  $\delta'(\kappa_2 y) = \beta(\kappa_2 y)$  for any  $y \in Y$ , so  $\delta' = \delta$ . We have shown that the square on the left of (3.1) is indeed a pullback in  $\mathcal{Kl} \mathcal{D}_M$ .

To show that the square on the right of (3.1) is also a pullback, assume  $\alpha: Z \rightarrow \mathcal{D}_M 1$  and  $\beta: Z \rightarrow \mathcal{D}_M(X+Y)$  are maps with  $(\hat{!}+\hat{!}) \hat{\circ} \beta = \hat{\kappa}_1 \hat{\circ} \alpha$ . Assume  $z \in Z$  and  $y \in Y$ . Then  $\beta(z)(\kappa_2 y) \leq ((\hat{!}+\hat{!}) \hat{\circ} \beta)(z)(\kappa_2^*) = (\hat{\kappa}_1 \hat{\circ} \alpha)(z)(\kappa_2^*) = 0$ . This allows us to define  $\delta: Z \rightarrow \mathcal{D}_M X$  by  $\delta(z)(x) = \beta(\kappa_1 x)$  as  $\bigcirc_x \delta(z)(x) = \bigcirc_x \beta(\kappa_1 x) = \bigcirc_x \beta(\kappa_1 x) \circ \bigcirc_y \beta(\kappa_2 y) = 1$ . Clearly  $\hat{\kappa}_1 \hat{\circ} \delta = \beta$  and  $\hat{!} \hat{\circ} \delta = \alpha$ . Suppose  $\delta': Z \rightarrow \mathcal{D}_M X$  is some map with  $\hat{\kappa}_1 \hat{\circ} \delta' = \beta$  and  $\hat{!} \hat{\circ} \delta' = \alpha$ . Then  $\delta'(z)(x) = (\hat{\kappa}_1 \hat{\circ} \delta')(\kappa_1 x) = \beta(z)(\kappa_1 x) = \delta(z)(x)$  and so  $\delta = \delta'$ . We have shown that the square on the right of (3.1) is a pullback in  $\mathcal{Kl} \mathcal{D}_M$ .

It only remains to be shown that the maps  $[\hat{\kappa}_1, \hat{\kappa}_2, \hat{\kappa}_2], [\hat{\kappa}_2, \hat{\kappa}_1, \hat{\kappa}_2]: 1+1+1 \rightarrow \mathcal{D}_M(1+1)$  are jointly monic in  $\mathcal{Kl} \mathcal{D}_M$ . To this end, assume  $f_1, f_2: Z \rightarrow \mathcal{D}_M(1+1+1)$  are given with  $[\hat{\kappa}_1, \hat{\kappa}_2, \hat{\kappa}_2] \hat{\circ} f_1 = [\hat{\kappa}_1, \hat{\kappa}_2, \hat{\kappa}_2] \hat{\circ} f_2$  and  $[\hat{\kappa}_2, \hat{\kappa}_1, \hat{\kappa}_2] \hat{\circ} f_1 = [\hat{\kappa}_2, \hat{\kappa}_1, \hat{\kappa}_2] \hat{\circ} f_2$ . Assume  $z \in Z$ . Clearly

$$\begin{aligned} f_1(z)(\kappa_1^*) &= ([\hat{\kappa}_1, \hat{\kappa}_2, \hat{\kappa}_2] \hat{\circ} f_1)(z)(\kappa_1^*) \\ &= ([\hat{\kappa}_1, \hat{\kappa}_2, \hat{\kappa}_2] \hat{\circ} f_2)(z)(\kappa_1^*) \\ &= f_2(z)(\kappa_1^*) \end{aligned}$$

and

$$\begin{aligned} f_1(z)(\kappa_2^*) &= ([\hat{\kappa}_2, \hat{\kappa}_1, \hat{\kappa}_2] \hat{\circ} f_1)(z)(\kappa_1^*) \\ &= ([\hat{\kappa}_2, \hat{\kappa}_1, \hat{\kappa}_2] \hat{\circ} f_2)(z)(\kappa_1^*) \\ &= f_2(z)(\kappa_2^*). \end{aligned}$$

So

$$\begin{aligned}
f_1(z)(\kappa_{3*}) &= (f_1(z)(\kappa_{1*}) \otimes f_1(z)(\kappa_{2*}))^\perp \\
&= (f_2(z)(\kappa_{1*}) \otimes f_2(z)(\kappa_{2*}))^\perp \\
&= f_2(z)(\kappa_{3*}),
\end{aligned}$$

hence  $f_1 = f_2$ . We have shown that  $\mathcal{Kl} \mathcal{D}_M$  is an effectus.

A scalar in  $\mathcal{Kl} \mathcal{D}_M$  corresponds to a map  $\lambda: 1 \rightarrow \mathcal{D}_M(1+1)$ , which corresponds in turn to an element of  $M$ . Unfolding definitions, it is straight-forward to see that the effect monoid structure defined on the scalars is the same as that of  $M$ .

**Solution to 193 II.** We start with the surjectivity of  $q$ ,  $\mathcal{D}_M q$  and  $\mathcal{D}_M \mathcal{D}_M q$ . By definition  $q$  is clearly surjective. Pick a section  $r: X/\sim \rightarrow X$  of  $q$ ; i.e.  $q \circ r = \text{id}$ . Then  $\mathcal{D}_M r$  and  $\mathcal{D}_M \mathcal{D}_M r$  are sections of  $\mathcal{D}_M q$  and  $\mathcal{D}_M \mathcal{D}_M q$  respectively, hence  $\mathcal{D}_M q$  and  $\mathcal{D}_M \mathcal{D}_M q$  are surjective.

To prove the second point, assume  $\sim$  is a congruence. Assume  $\varphi, \psi \in \mathcal{D}_M X$  with  $(\mathcal{D}_M q)(\varphi) = (\mathcal{D}_M q)(\psi)$ . By definition  $\varphi \sim \psi$  and  $h(\varphi) \sim h(\psi)$  by assumption that  $\sim$  is a congruence. Thus  $q \circ h(\varphi) = q \circ h(\psi)$ . Thus together with the surjectivity of  $\mathcal{D}_M q$ , there is a unique map  $h_\sim: \mathcal{D}_M X/\sim \rightarrow X/\sim$  fixed by  $h_\sim \circ (\mathcal{D}_M q) = q \circ h$ . To prove the converse, assume there is a map  $h_\sim: \mathcal{D}_M X/\sim \rightarrow X/\sim$  with  $h_\sim \circ (\mathcal{D}_M q) = q \circ h$  and assume  $\varphi \sim \psi$  for some  $\varphi, \psi \in \mathcal{D}_M X$ . Then  $q(h(\varphi)) = h_\sim(\mathcal{D}_M q(\varphi)) = h_\sim(\mathcal{D}_M q(\psi)) = q(h(\psi))$  and so  $h(\varphi) \sim h(\psi)$ , which shows that  $\sim$  is a congruence.

For the third point, assume  $\sim$  is a congruence. Then

$$\begin{aligned}
h_\sim \circ \mathcal{D}_M h_\sim \circ \mathcal{D}_M \mathcal{D}_M h_\sim &= h_\sim \circ \mathcal{D}_M (h_\sim \circ \mathcal{D}_M q) \\
&= h_\sim \circ \mathcal{D}_M (q \circ h) && \text{by the second point} \\
&= q \circ h \circ \mathcal{D}_M h && \text{idem} \\
&= q \circ h \circ \mu && \text{as } (X, h) \text{ is a.conv.} \\
&= h_\sim \circ \mathcal{D}_M q \circ \mu && \text{by the second point} \\
&= h_\sim \circ \mu \circ \mathcal{D}_M \mathcal{D}_M q && \text{by naturality } \mu.
\end{aligned}$$

Hence by surjectivity of  $\mathcal{D}_M \mathcal{D}_M q$ , we find  $h_\sim \circ \mathcal{D}_M h_\sim = h_\sim \circ \mu$ . Using naturality of  $\eta$ , the second point and the fact that  $(X, h)$  is an abstract convex set (in that order), we find  $h_\sim \circ \eta \circ q = h_\sim \circ \mathcal{D}_M q \circ \eta = q \circ h \circ \eta = q$  and so  $h_\sim \circ \eta = \text{id}$  by surjectivity of  $q$ . We have shown  $(X/\sim, h_\sim)$  is an abstract  $M$ -convex set. The equality  $q \circ h = h_\sim \circ \mathcal{D}_M q$  from the second point shows that  $q$  is an  $M$ -affine map.

**Solution to 193 III.** To show  $\sim$  is a congruence, assume  $\varphi \sim \psi$  for  $\varphi, \psi \in \mathcal{D}_M X$ . Write  $q: X \rightarrow X/\sim$  for the quotient map for  $\sim$  and  $f_\sim: X/\sim \rightarrow X$  for the unique map with  $f_\sim \circ q = f$ . Note that  $\mathcal{D}_M q(\varphi) = \mathcal{D}_M q(\psi)$  and so  $f(h(\varphi)) =$

$h(\mathcal{D}_M f(\varphi)) = h(\mathcal{D}_M f_{\sim}(\mathcal{D}_M q(\varphi))) = h(\mathcal{D}_M f_{\sim}(\mathcal{D}_M q(\psi))) = f(h(\psi))$  and so  $h(\varphi) \sim h(\psi)$ , which shows  $\sim$  is a congruence.

**Solution to 193 IV.** Concerning the first point: clearly  $h(\eta(h(\psi))) = h(\psi)$  and so  $(\eta(h(\psi)), \psi)$  is a derivation of  $\eta(h(\psi)) \approx \psi$ . Assume  $\varphi \approx \psi$ . Then  $\eta(h(\varphi)) \approx \varphi \approx \psi \approx \eta(h(\psi))$  and so  $h(\varphi) \sim h(\psi)$ .

We continue with the second point. Suppose we are given  $\chi_1, \dots, \chi_n, \varphi, \psi \in \mathcal{D}_M X$  and  $\lambda_0, \dots, \lambda_n \in M$  with  $\bigotimes_i \lambda_i = 1$ . To cover the first base case in the definition of a derivation, assume  $h(\varphi) = h(\psi)$ . Then

$$\begin{aligned} (h \circ \mu) \left( \lambda_0 |\psi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle \right) &= (h \circ \mathcal{D}_M h) \left( \lambda_0 |\psi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle \right) \\ &= h \left( \lambda_0 |h(\psi)\rangle \otimes \bigotimes_{j=1}^n \lambda_j |h(\chi_j)\rangle \right) \\ &= h \left( \lambda_0 |h(\varphi)\rangle \otimes \bigotimes_{j=1}^n \lambda_j |h(\chi_j)\rangle \right) \\ &= (h \circ \mu) \left( \lambda_0 |\varphi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle \right) \end{aligned}$$

and so  $\mu(\lambda_0 |\psi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle) \approx \mu(\lambda_0 |\varphi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle)$ . Next, for the other base case in the definition of a derivation, assume  $\varphi \equiv \bigotimes_{i=1}^m \mu_i |x_i\rangle$  and  $\psi \equiv \bigotimes_{i=1}^m \mu_i |y_i\rangle$  with  $x_i R^* y_i$  for  $1 \leq i \leq m$ . Then

$$\begin{aligned} \mu \left( \lambda_0 |\psi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle \right) &= \bigotimes_{i=1}^m \lambda_0 \circ \mu_i |y_i\rangle \otimes \bigotimes_{j=1}^n \bigotimes_x \lambda_j \circ \chi_j(x) |x\rangle \\ \mu \left( \lambda_0 |\varphi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle \right) &= \bigotimes_{i=1}^m \lambda_0 \circ \mu_i |x_i\rangle \otimes \bigotimes_{j=1}^n \bigotimes_x \lambda_j \circ \chi_j(x) |x\rangle. \end{aligned}$$

This shows that  $\mu(\lambda_0 |\psi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle) \approx \mu(\lambda_0 |\varphi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle)$  by the second base case of a derivation. Thus by induction  $\mu(\lambda_0 |\psi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle) \approx \mu(\lambda_0 |\varphi\rangle \otimes \bigotimes_{j=1}^n \lambda_j |\chi_j\rangle)$  whenever merely  $\varphi \approx \psi$ .

For the third point, we need some preparation. Pick representatives  $R \subseteq X$  of  $\sim$ : for every  $x \in X$  there is a unique  $r_x \in R$  with  $r_x \sim x$ . By defini-

tion  $\eta(r_x) \approx \eta(x)$ . Let  $\varphi \equiv \bigotimes_{i=1}^n \lambda_i |x_i\rangle \in \mathcal{D}_M X$  given. By the previous point

$$\begin{aligned}
\varphi &= \mu\left(\lambda_1 |\eta(x_1)\rangle \otimes \bigotimes_{i=2}^n \lambda_i |\eta(x_i)\rangle\right) \\
&\approx \mu\left(\lambda_1 |\eta(r_{x_1})\rangle \otimes \bigotimes_{i=2}^n \lambda_i |\eta(x_i)\rangle\right) \\
&= \mu\left(\lambda_2 |\eta(x_2)\rangle \otimes \lambda_1 |r_{x_1}\rangle \otimes \bigotimes_{i=3}^n \lambda_i |\eta(x_i)\rangle\right) \\
&\approx \mu\left(\lambda_2 |\eta(r_{x_2})\rangle \otimes \lambda_1 |r_{x_1}\rangle \otimes \bigotimes_{i=3}^n \lambda_i |\eta(x_i)\rangle\right) \\
&\quad \vdots \\
&\approx \bigotimes_{i=1}^n \lambda_i |r_{x_i}\rangle \\
&\approx \bigotimes_{r \in R} \left( \bigotimes_{x \sim r} \varphi(x) \right) |r\rangle.
\end{aligned}$$

Now, assume  $\varphi \sim \psi$ . Then  $\bigotimes_{x \sim x_0} \varphi(x) = \bigotimes_{x \sim x_0} \psi(x)$  for any  $x_0 \in X$  and so

$$\varphi \approx \bigotimes_{r \in R} \left( \bigotimes_{x \sim r} \varphi(x) \right) |r\rangle = \bigotimes_{r \in R} \left( \bigotimes_{x \sim r} \psi(x) \right) |r\rangle \approx \psi.$$

Clearly  $\sim$  is an equivalence relation on  $X$ . To show it is a congruence, assume  $\varphi \sim \psi$ . By the previous, we know  $\varphi \approx \psi$  and so  $h(\varphi) \sim h(\psi)$  by the first point, which shows  $\sim$  is indeed a congruence. Furthermore  $R$  is contained in  $\sim$ ; indeed, if  $x R y$ , then  $(\eta(x), \eta(y))$  is a derivation of  $\eta(x) \approx \eta(y)$  and so  $x \sim y$ .

To show  $\sim$  is the smallest congruence containing  $R$ , assume  $S \subseteq X^2$  is some congruence of  $X$  with  $R \subseteq S$ . It is sufficient to show that  $\varphi \approx \psi$  implies  $h(\varphi) S h(\psi)$ . Indeed if this is the case and  $x \sim y$ , then by definition  $\eta(x) \approx \eta(y)$  and so  $x = h(\eta(x)) S h(\eta(y)) = y$ .

We will prove that  $\varphi \approx \psi$  implies  $h(\varphi) S h(\psi)$  by induction over the definition of  $\approx$ . For the first base case in the definition of  $\approx$ , assume  $\varphi \equiv \bigotimes_{i=1}^n \lambda_i |x_i\rangle$  and  $\psi \equiv \bigotimes_{i=1}^n \lambda_i |y_i\rangle$  for some  $(x_1, y_1), \dots, (x_n, y_n) \in R^*$ . Then  $x_i S y_i$  and so  $\varphi S \psi$ , hence  $h(\varphi) S h(\psi)$ . For the other base case, suppose  $h(\varphi) = h(\psi)$  for some  $\varphi, \psi \in \mathcal{D}_M X$ . Then clearly  $h(\varphi) S h(\psi)$ . Thus, by induction  $h(\varphi) S h(\psi)$  whenever  $\varphi \approx \psi$ .

**Solution to 193 X.** As the one element set is final in  $\mathbf{Set}$ , there is a unique map  $! : \mathcal{D}_M 1 \rightarrow 1$ . This turns  $\mathcal{D}_M 1$  into an abstract  $M$ -convex set:  $! \circ \mathcal{D}_M ! =$

$! = ! \circ \mu$  and  $! \circ \eta = ! = \text{id}$ . Note that there is a single formal  $M$ -convex combination over  $1$  and so  $\mathcal{D}_M 1 \cong 1$ . As  $\text{AConv}_M$  is the Eilenberg–Moore category of the monad  $\mathcal{D}_M$ , it follows that the functor  $F: \mathbf{Set} \rightarrow \text{AConv}_M$  given by  $F(X) = (\mathcal{D}_M X, \mu)$  and  $Ff = \mathcal{D}_M f$  is a left adjoint and so preserves coproducts. Hence  $\mathcal{D}_M\{1, \dots, n\} \cong F(n \cdot 1) \cong n \cdot F(1) \cong n \cdot 1$ .

**Solution to 195 IV.** Clearly  $0 \leq 0 \setminus 0$  and  $0 \odot 0 = 0$ . Thus, by uniqueness of  $0 \setminus 0$ , we have  $0 \setminus 0 = 0$ . For any  $a \in M$ , we have  $1 \setminus a = 1 \odot 1 \setminus a = a$  and so in particular  $1 \setminus 1 = 1$ . Next, note  $a \setminus a \odot a \setminus a \leq a \setminus a$  and  $a \odot a \setminus a \odot a \setminus a = a \odot a \setminus a = a$  and so by uniqueness of  $a \setminus a$ , we have  $a \setminus a \odot a \setminus a = a \setminus a$ . For the final equation of the first point, assume  $b \in M$ . As  $a \odot b \leq a$  we know  $a \setminus a \odot b$  is defined. Note  $a \setminus a \odot b \leq a \setminus a$  and  $a \odot a \setminus a \odot b = a \odot b$ . So by uniqueness of  $a \setminus a \odot b$  we see  $a \setminus a \odot b = a \setminus a \odot b$ .

For the second point, suppose  $a, b, c \in M$  with  $a \leq b \leq c$ . Note  $c \setminus b \odot b \setminus a \leq c \setminus b \leq c \setminus c$  and  $c \odot c \setminus b \odot b \setminus a = b \odot b \setminus a = a$ . So by uniqueness of  $c \setminus a$ , we get  $c \setminus b \odot b \setminus a = c \setminus a$ , as desired.

**Solution to 195 VI.** Let  $X$  be a compact Hausdorff space. Assume that the unit interval of  $C(X)$  is an effect divisoid. To show  $X$  is basically disconnected, assume  $f \in C(X)$ . We have to show that  $\overline{\text{supp } f}$  is open. Note  $\text{supp } |f| = \text{supp } f$  and so, without loss of generality, we may assume that  $f \geq 0$ . By compactness  $f$  is bounded. Pick  $B > 0$  with  $B \geq f$ . Then  $\text{supp } B^{-1}f = \text{supp } f$  and  $\overline{B^{-1}f} \leq 1$ , so we may also assume without loss of generality, that  $0 \leq f \leq 1$ . If  $\text{supp } f = X$ , then we are done, so assume  $\overline{\text{supp } f} \neq X$ . Pick any  $y \notin \overline{\text{supp } f}$ . By Urysohn's lemma, there is a  $g \in C(X)$  with  $g(y) = 0$  and  $g(x) = 1$  for every  $x \in \overline{\text{supp } f}$ . Define  $h \equiv f \setminus f \wedge (0 \vee g)$ . Clearly  $0 \leq h \leq f \setminus f \leq 1$  and  $h(y) \leq (0 \vee g)(y) = 0$ . Note that  $f \setminus f \setminus f \setminus f \equiv f \setminus f \odot f \setminus f = f \setminus f$  and so  $f \setminus f$  is zero–one valued: a characteristic function. Pick any  $x \in \overline{\text{supp } f}$ . Then  $0 < \frac{f(x)}{f(x)} \leq f \setminus f(x) \in \{0, 1\}$ , hence  $f \setminus f(x) = 1$ . By continuity  $f \setminus f(x) = 1$  for every  $x \in \overline{\text{supp } f}$  and so  $h(x) = f \setminus f(x) \wedge g(x) = 1 \wedge 1 = 1$ . Hence  $(f \odot h)(x) = f(x)h(x) = f(x)$  for  $x \in \overline{\text{supp } f}$  and  $(f \odot h)(x) = f(x)h(x) = 0 = f(x)$  whenever  $x \notin \overline{\text{supp } f}$ . Thus  $f \odot h = f$ . By uniqueness of  $f \setminus f$ , we have  $f \setminus f = h$ . Consequently  $f \setminus f(y) = h(y) = 0$ . Recall that  $y$  was an arbitrary element  $y \notin \overline{\text{supp } f}$  and thus  $f \setminus f$  is the characteristic function of  $\overline{\text{supp } f}$ , which is thus open. Hence  $X$  is basically disconnected.

To prove the converse, assume  $X$  is basically disconnected. Let  $f, g \in C(X)$  be given with  $0 \leq f \leq g \leq 1$ . Define  $U_n \equiv \{x; g(x) > \frac{1}{n}\}$ . Note  $U_n = \text{supp}((g - \frac{1}{n}) \vee 0)$  and so  $\overline{U_n}$  is open as  $X$  is basically disconnected. Define

$$h_n \equiv \begin{cases} \frac{f(x)}{g(x)} & x \in \overline{U_n} \\ 0 & \text{otherwise.} \end{cases}$$

To show  $h_n$  is continuous, assume there is a net  $(x_\alpha)_\alpha$  with  $x_\alpha \rightarrow x$  for some  $x \in X$ . Suppose  $x \in \overline{U_n}$ . As  $\overline{U_n}$  is open, we know  $x_\alpha \in U_n$  for sufficiently large  $\alpha$  and so  $h_n(x_\alpha) \rightarrow h_n(x)$  as  $\frac{f(x)}{g(x)}$  is continuous for  $x \in \overline{U_n}$  as then  $g(x) \geq \frac{1}{n} > 0$ .

For the other case, suppose  $x \notin \overline{U_n}$ . The set  $X - \overline{U_n}$  is open and so  $x_\alpha \notin \overline{U_n}$  for sufficiently large  $\alpha$  and then  $h_n(x_\alpha) = 0 = h_n(x)$ . Thus  $h_n$  is continuous.

Clearly  $0 \leq h_1 \leq h_2 \leq \dots \leq 1$  and so we may define  $g \setminus f \equiv \sup_n h_n$  as  $C(X)$  is  $\omega$ -complete by basic disconnectedness of  $X$ . Note  $0 \leq g \setminus f \leq 1$ .

Suppose  $f = g$  and  $x \in \text{supp } f$ . Then  $h_n(x) = 1$  for all  $n > f(x)^{-1}$  and so  $f \setminus f(x) = 1$ . Thus  $f \setminus f(x) = 1$  for all  $x \in \text{supp } f$ . Write  $\chi$  for the characteristic function of  $\overline{\text{supp } f}$ . We just saw  $\chi \leq f \setminus f(x)$ . As  $\overline{U_n} \subseteq \overline{\text{supp } f}$  and  $h_n \leq 1$ , we have  $h_n \leq \chi$  for all  $n$  and so  $f \setminus f \leq \chi$ . Thus  $f \setminus f = \chi$ , the characteristic function of  $\overline{\text{supp } f}$ . In particular  $f \leq f \setminus f$  and  $f \setminus f \odot f \setminus f \equiv f \setminus f f \setminus f = f \setminus f$ .

Let  $f, g \in C(X)$  be given with  $0 \leq f \leq g \leq 1$  and  $h_1 \leq h_2 \leq \dots$  as above. We will show  $g \setminus f \leq g \setminus g$ . Assume  $x \in X$ . Suppose  $x \in \text{supp } g$ . Then  $h_n(x) = \frac{f(x)}{g(x)} \leq 1 = g \setminus g(x)$  for all  $n > g(x)^{-1}$  and so  $g \setminus f(x) \leq g \setminus g(x)$ . In particular  $g \setminus f(x) \leq g \setminus g(x)$  for all  $x \in \text{supp } g$ . For the other case, assume  $x \notin \text{supp } g$ . Then  $x \notin \overline{U_n}$  and so  $h_n(x) = 0$  for all  $n \in \mathbb{N}$ , whence  $g \setminus f(x) = 0 \leq g \setminus g(x)$ . Thus indeed  $g \setminus f \leq g \setminus g$ .

In a similar way, it is easy to see that  $g \setminus f(x) \leq \frac{f(x)}{g(x)}$  for  $x \in \text{supp } g$ . To show equality, we define

$$k_n \equiv \begin{cases} \frac{f(x)}{g(x)} & x \in \overline{U_n} \\ 1 & \text{otherwise.} \end{cases}$$

With similar reasoning as for  $h_n$ , we see that these  $k_n$  are continuous. Furthermore  $h_m \leq k_n$  for all  $m$  and so  $g \setminus f \leq k_n$  for any  $n$ . Pick any  $x \in \text{supp } g$ . Then  $g \setminus f(x) \leq k_n(x) = \frac{f(x)}{g(x)}$  for  $n > g(x)^{-1}$  and so  $g \setminus f(x) = \frac{f(x)}{g(x)}$  for any  $x \in \text{supp } g$ . Thus  $(g \odot g \setminus f)(x) = f(x)$  for  $x \in \text{supp } g$  and so  $(g \odot g \setminus f)(x) = f(x)$  for  $x \in \overline{\text{supp } g}$  by continuity. For the other case, assume  $x \notin \overline{\text{supp } g}$ . Then  $g \odot g \setminus f(x) \leq g \setminus f(x) \leq g \setminus g(x) = 0 = g(0) \geq f(0)$ . We have shown  $g \odot g \setminus f = f$ .

Only uniqueness of  $g \setminus f$  remains. So assume  $h \in C(X)$  with  $0 \leq h \leq g \setminus g$  and  $g \odot h = f$ . Assume  $x \in \text{supp } g$ . Then  $g(x)h(x) = f(x)$  and so  $h(x) = \frac{f(x)}{g(x)} = g \setminus f(x)$ . By continuity  $h(x) = g \setminus f(x)$  for all  $x \in \overline{\text{supp } g}$ . For the other case, assume  $x \notin \overline{\text{supp } g}$ . As both  $h, g \setminus f \leq g \setminus g$ , we have  $h(x) = 0 = g \setminus f(x)$ . We have shown  $h = g \setminus f$  and thus the unit interval of  $C(X)$  is an effect divisoid.

**Solution to 197 V.** Assume  $C$  is an effectus.

1. Assume  $\xi: X \rightarrow Y$  is a quotient for  $p$  and  $\vartheta: Y \rightarrow Z$  is an isomorphism. Note  $1 \circ \vartheta \circ \xi = 1 \circ \xi \leq p^\perp$ . To prove  $\vartheta \circ \xi$  is a quotient for  $p$ , assume  $f: X \rightarrow Y'$  is some map with  $1 \circ f \leq p^\perp$ . As  $\xi$  is a quotient, there exists a unique map  $f': Y \rightarrow Y'$  with  $f' \circ \xi = f$ . Clearly  $f = f' \circ \xi = (f' \circ \vartheta^{-1}) \circ (\vartheta \circ \xi)$ . Assume  $h: Z \rightarrow Y'$  is any map with  $f = h \circ (\vartheta \circ \xi)$ . Then  $h \circ \vartheta = f'$  by uniqueness of  $f'$  and so  $h = f' \circ \vartheta^{-1}$ . We have shown that  $\vartheta \circ \xi$  is a quotient of  $p$  as well.
2. Assume  $\xi_1: X \rightarrow Y_1$  and  $\xi_2: X \rightarrow Y_2$  are quotients for  $p$ . By the universal property of  $\xi_1$ , there is a unique map  $\vartheta_1: Y_1 \rightarrow Y_2$  with  $\vartheta_1 \circ \xi_1 = \xi_2$ .

Similarly, there is a unique map  $\vartheta_2: Y_2 \rightarrow Y_1$  with  $\vartheta_2 \circ \xi_2 = \xi_1$ . Note that  $\xi_1 = \vartheta_2 \circ \xi_2 = \vartheta_2 \circ \vartheta_1 \circ \xi_1$ . By the universal property of  $\xi_1$  again, the map  $\text{id}: Y_1 \rightarrow Y_1$  is the unique map with  $\text{id} \circ \xi_1 = \xi_1$  and so  $\vartheta_2 \circ \vartheta_1 = \text{id}$ . Similarly  $\vartheta_1 \circ \vartheta_2 = \text{id}$ . Thus  $\vartheta_2$  is an isomorphism with  $\vartheta_2 \circ \xi_2 = \xi_1$ . It is the unique such isomorphism by the universal property of  $\xi_2$ .

3. Assume  $f: X \rightarrow Y$  is some map. Clearly  $1 \circ f \leq 1 = 0^\perp$ . Furthermore  $f$  is itself the unique map  $f'$  with  $f' \circ \text{id} = f$ . Thus  $\text{id}$  is a quotient for  $0$ . Consequently any isomorphism is a quotient for  $0$  by the first point.
4. Assume  $X$  is any object. As  $0$  is a zero object, there is a unique map  $0_X: X \rightarrow 0$ . Clearly  $1 \circ 0 = 0 = 1^\perp$ . Assume  $f: X \rightarrow Y$  is any map with  $1 \circ f \leq 1^\perp = 0$ . Then  $f = 0$  and so  $f = 0 = 0 \circ 0_X$ . If  $f = h \circ 0_X$  for some  $h: 0 \rightarrow Y$ , then  $h = 0$  and so any map into zero is indeed a quotient for  $1$ .
5. Assume  $\xi: X \rightarrow Y$  is a quotient for  $p$ . Clearly  $1 \circ p^\perp = p^\perp \leq p^\perp$  and so there is a unique map  $f: Y: 1$  with  $f \circ \xi = p^\perp$ . Then  $p^\perp = 1 \circ p^\perp = 1 \circ f \circ \xi \leq 1 \circ \xi = p^\perp$  and so  $1 \circ \xi = p^\perp$ .
6. Assume  $f_1 \circ \xi = f_2 \circ \xi$  for some quotient  $\xi$  of  $p$ . Then  $1 \circ f_1 \circ \xi \leq 1 \circ \xi = p^\perp$ . Hence there is a unique  $f$  with  $f \circ \xi = f_1 \circ \xi$ . Both  $f_1$  and  $f_2$  fit the bill, hence  $f_1 = f = f_2$ . Thus  $\xi$  is epic.

**Solution to 197 XI.** Let  $C$  be an effectus with quotients. Assume  $t' \circ \xi' = t \circ \xi$  for some quotient  $\xi, \xi'$  and total maps  $t, t'$ . Then  $1 \circ \xi = 1 \circ t \circ \xi = 1 \circ t' \circ \xi' = 1 \circ \xi'$ . Thus by 197V there is a unique isomorphism  $\vartheta$  with  $\xi' = \vartheta \circ \xi$ . Note  $t \circ \xi = t' \circ \xi' = t' \circ \vartheta \circ \xi$  and so by epicity of  $\xi$  (see 197V), we see  $t = t' \circ \vartheta$ , as desired.

**Solution to 198 III.** Assume  $C$  is an effectus with quotients. We will show that  $0$  has a left adjoint by demonstrating the universal mapping property. Pick for every object  $(X, p)$  in  $\int \text{Pred}_\square$  a quotient  $\xi_p: X \rightarrow X/p$  of  $p$ . Note  $(0^\perp \circ \xi_p)^\perp = p$  and so  $\xi_p$  is a map  $(X, p) \rightarrow (X/p, 0)$  in  $\int \text{Pred}_\square$ . We will use these maps  $\xi_p$  as the components of the unit of the adjunction. Let  $f: (X, p) \rightarrow (Y, 0)$  be some map in  $\int \text{Pred}_\square$ . By definition of map  $1 \circ f \leq p^\perp$  and so by the universal property of  $\xi_p$ , there exists a unique map  $f': X/p \rightarrow Y$  with  $f = f' \circ \xi_p$  in  $C$ . Clearly also  $(0f) \circ \xi_p = f$  in  $\int \text{Pred}_\square$ . Let  $g: X/p \rightarrow Y$  be some map with  $(0g) \circ \xi_p = f$  as well. Then  $g \circ \xi_p = f$  and so  $g' = f'$ . This shows that  $0$  has a left adjoint.

To prove the converse, assume  $C$  is an effectus where  $0: C \rightarrow \int \text{Pred}_\square$  has a left adjoint  $Q: \int \text{Pred}_\square \rightarrow C$ . Let  $X$  be some object in  $C$  with a predicate  $p$ . Write  $\eta: (X, p) \rightarrow 0Q(X, p)$  for the  $(X, p)$  component of the unit of the adjunction  $Q \dashv 0$ . By definition of maps in  $\int \text{Pred}_\square$ , we know  $1 \circ \eta \leq p^\perp$ . We will show that  $\eta$  is a quotient for  $p$ . To this end, assume  $f: X \rightarrow Y$  is some map in  $C$  with  $1 \circ f \leq p^\perp$ . Then  $f: (X, p) \rightarrow (Y, 0)$  in  $\int \text{Pred}_\square$ . By the universal mapping property, there is a unique map  $f': Q(X, p) \rightarrow Y$  in  $C$  with  $(0f') \circ \eta = f$ ,

i.e.  $f' \circ \eta = f$ . This shows that  $\eta$  is indeed a quotient of  $p$ . Hence  $C$  has quotients.

**Solution to 199 VI.** Assume  $C$  is an effectus with comprehension. We will show that  $1$  has a right adjoint by demonstrating the dual of the universal mapping property. Pick for every object  $(X, p)$  in  $\int \text{Pred}_{\square}$  a comprehension  $\pi_p: \{X|p\} \rightarrow X$ . Note that for any map  $f$ , we have  $1 \circ f = p \circ f$  if and only if  $p^{\perp} \circ f = 0$ . Thus  $(p^{\perp} \circ \pi_p)^{\perp} = 1$ , which shows that  $\pi_p$  is a map  $(\{X|p\}, 1) \rightarrow (X, p)$  in  $\int \text{Pred}_{\square}$ . We will use these maps as components for the counit of the adjunction. Let  $f: (Y, 1) \rightarrow (X, p)$  be some map in  $C$ . By definition of maps in  $\int \text{Pred}_{\square}$ , we have  $1 \leq (p^{\perp} \circ f)^{\perp}$ , viz.  $1 \circ f = p \circ f$ . Thus, by the universal property of  $\pi_p$ , there is a unique map  $f': Y \rightarrow \{X|p\}$  in  $C$  with  $\pi_p \circ f' = f$ . Hence  $1f'$  is the unique map in  $\int \text{Pred}_{\square}$  with  $\pi_p \circ (1f') = f$  in  $\int \text{Pred}_{\square}$ . We have shown that  $1$  has a right adjoint.

To prove the converse, assume  $C$  is an effectus where  $1: C \rightarrow \int \text{Pred}_{\square}$  has a right adjoint  $K: \int \text{Pred}_{\square} \rightarrow C$ . Let  $X$  be some object of  $C$  with a predicate  $p$ . Write  $\varepsilon: 1K(X, p) \rightarrow (X, p)$  for the  $(X, p)$  component of the counit of the adjunction  $1 \dashv K$ . By definition of maps in  $\int \text{Pred}_{\square}$ , we know  $1 \leq (p^{\perp} \circ \varepsilon)^{\perp}$ , hence  $1 \circ \varepsilon = 1 \circ \varepsilon$ . We will show that  $\varepsilon$  is a comprehension for  $p$ . To this end, assume  $f: Y \rightarrow X$  is some map in  $C$  with  $1 \circ f = p \circ f$ . Then  $1 = (p^{\perp} \circ f)^{\perp}$  and so  $f$  is a map  $(Y, 1) \rightarrow (X, p)$  in  $\int \text{Pred}_{\square}$ . By the dual of the universal mapping property, there is a unique map  $f': Y \rightarrow K(X, p)$  in  $\int \text{Pred}_{\square}$  with  $\varepsilon \circ (1f') = f$ , i.e.  $\varepsilon \circ f' = f$ . This shows that  $\varepsilon$  is indeed a comprehension for  $p$ . Hence  $C$  has comprehension.

**Solution to 199 VII.** Let  $C$  be an effectus.

1. Assume  $\pi: X \rightarrow Y$  is a comprehension for  $p$  and  $\vartheta: Z \rightarrow X$  is an isomorphism. We will show  $\pi \circ \vartheta$  is a comprehension for  $p$  as well. To start, note  $1 \circ \pi \circ \vartheta = p \circ \pi \circ \vartheta$ . Assume  $f: X' \rightarrow Y$  is any map with  $1 \circ f = p \circ f$ . Then, by the universal property of  $\pi$ , there is a unique map  $f': X' \rightarrow X$  with  $\pi \circ f' = f$ . Then  $f = (\pi \circ \vartheta) \circ (\vartheta^{-1} \circ f')$ . Assume  $g: X' \rightarrow X$  is a map with  $g = (\pi \circ \vartheta) \circ g$  as well. Then  $\vartheta \circ g = f'$  by uniqueness of  $f'$  and so  $g = \vartheta^{-1} \circ f'$ . We have shown that  $\pi \circ \vartheta$  is a comprehension for  $p$  as well.
2. Assume  $\pi_1: X_1 \rightarrow Y$  and  $\pi_2: X_2 \rightarrow Y$  are comprehensions of  $p$ . By the universal property of  $\pi_1$ , there is a unique map  $\vartheta_1: X_2 \rightarrow X_1$  with  $\pi_2 = \pi_1 \circ \vartheta_1$ . On the other side, by the universal property of  $\pi_2$ , there is a unique map  $\vartheta_2: X_1 \rightarrow X_2$  with  $\pi_1 = \pi_2 \circ \vartheta_2$ . Then  $\pi_1 = \pi_2 \circ \vartheta_2 = \pi_1 \circ \vartheta_1 \circ \vartheta_2$ . By the universal property of  $\pi_1$  again, the map  $\text{id}: X_1 \rightarrow X_1$  is the unique map with  $\pi_1 \circ \text{id} = \pi_1$  and so  $\vartheta_1 \circ \vartheta_2 = \text{id}$ . Similarly  $\vartheta_2 \circ \vartheta_1 = \text{id}$ . Thus  $\vartheta_2$  is an isomorphism with  $\vartheta_1 = \pi_2 \circ \vartheta_2$ . It is the unique such isomorphism by the universal property of  $\pi_2$ .

3. Assume  $f: X \rightarrow Y$  is some map. Trivially  $1 \circ f = 1 \circ f$  and  $1 \circ \text{id} = 1 \circ \text{id}$ . The map  $f$  itself is the unique map  $g$  with  $g \circ \text{id} = f$  and so  $\text{id}$  is a comprehension for 1. Consequently any isomorphism is a comprehension for 1 by the first point.
4. Assume  $Y$  is any object. As 0 is a zero object, there is a unique map  $0_Y: 0 \rightarrow Y$ . Clearly  $1 \circ 0_Y = 0 = 0 \circ 0_Y$ . Assume  $f: X \rightarrow Y$  is any map with  $1 \circ f = 0 \circ f$ . Then  $1 \circ f = 0 \circ f = 0$  and so  $f = 0$ . The zero map is the unique map  $X \rightarrow 0$  and for that map, we have  $0_Y \circ 0 = 0 = f$  and so  $0_Y$  is a comprehension for 0.
5. Assume  $\pi \circ f_1 = \pi \circ f_2$  for some comprehension  $\pi$  of  $p$ . Then  $1 \circ \pi \circ f_1 = p \circ \pi \circ f_1$  and so by the universal property of  $\pi$ , there is a unique map  $f$  with  $\pi \circ f = \pi \circ f_1$ . Both  $f_1$  and  $f_2$  fit the bill, hence  $f_1 = f = f_2$ . Thus  $\pi$  is monic.
6. Assume  $\pi$  is a comprehension of  $p$ . Then  $(p^\perp \circ \pi) \circ (p \circ \pi) = (p^\perp \circ p) \circ \pi = 1 \circ \pi = p \circ \pi$ . Thus  $p^\perp \circ \pi = 0$  by cancellation.

**Solution to 200 V.** Let  $C$  be any effectus. Assume  $\pi: X \rightarrow Y$  is a comprehension of  $p$ . We will show  $\pi$  is a categorical kernel  $p^\perp$ . To start, note  $p^\perp \circ \pi = 0$  by 199 VII. Now assume  $f: X' \rightarrow Y$  is some map with  $p^\perp \circ f = 0$ . Then  $p \circ f = (p \circ f) \circ (p^\perp \circ f) = 1 \circ f$  and so by the universal property of  $\pi$ , there is a unique map  $f': X' \rightarrow X$  with  $f = \pi \circ f'$ . Hence  $\pi$  is a kernel of  $p^\perp$ .

To prove the converse, assume  $k: X \rightarrow Y$  is a kernel of  $p^\perp: Y \rightarrow 1$ . We have  $1 \circ k = (p \circ p^\perp) \circ k = (p \circ k) \circ (p^\perp \circ k) = p \circ k$ . Assume  $f: X' \rightarrow Y$  is some map with  $p \circ f = 1 \circ f$ . Then  $(p^\perp \circ f) \circ (p \circ f) = 1 \circ f = p \circ f$  and so by cancellation  $p^\perp \circ f = 0$ . Hence by the universal property of  $k$  there is a unique map  $f': X' \rightarrow X$  with  $k \circ f' = f$ . This shows that  $k$  is a comprehension of  $p$ .

**Solution to 202 V.** We have  $(\text{im } f) \circ f \circ g = 1 \circ f \circ g \leq 1 \circ g$  and so  $\text{im } f \circ g \leq \text{im } f$ . Consequently  $\text{im } f \circ \alpha \leq \text{im } f = \text{im } f \circ \alpha \circ \alpha^{-1} \leq \text{im } f \circ \alpha$  and so  $\text{im } f \circ \alpha = \text{im } f$ .

**Solution to 202 VI.** Assume  $\xi: X \rightarrow Y$  is some quotient and  $p$  is a predicate on  $Y$  with  $p \circ \xi = 0$ . Then  $p \circ \xi = 0 = 0 \circ \xi$  and so  $p = 0$  by epicity of  $\xi$ . Hence  $\xi$  is faithful.

**Solution to 203 XII.** Assume  $[p] = p$ . Then  $p = [p] \equiv \text{im } \pi$  for some comprehension  $\pi$  of  $p$  and so  $p$  is sharp. To prove the converse, assume  $p$  is sharp; that is:  $p = \text{im } f$  for some  $f$ . Pick any comprehension  $\pi$  of  $p$ . Clearly  $p \circ f = (\text{im } f) \circ f = 1 \circ f$  and so there is a unique  $g$  with  $f = \pi \circ g$ . By 202 V we find  $p = \text{im } f = \text{im } \pi \circ g \leq \text{im } \pi \equiv [p] \leq p$  and so  $p = [p]$ .

**Solution to 203 XIII.** Note  $[p] \leq [q]$  whenever  $p \leq q$  and so  $[p \circ f] \leq [[p] \circ f] \leq [[p \circ f]] = [p \circ f]$  as  $p \leq [p]$  and  $[p] \circ f \leq [p \circ f]$  by 203 XIII. Thus  $[p \circ f] =$

$[[p] \circ f]$ .

**Solution to 203 XIV.** To start, note  $[\text{im } f, \text{im } g] \langle f, g \rangle = ((\text{im } f) \circ f) \circledast ((\text{im } g) \circ g) = (1 \circ f) \circledast (1 \circ g) = 1 \circ \langle f, g \rangle$  by 181 IX. Assume  $p \equiv [p_1, p_2]$  is some predicate with  $p \circ \langle f, g \rangle = 1 \circ \langle f, g \rangle$ . Then  $(1 \circ f) \circledast (1 \circ g) = 1 \circ \langle f, g \rangle = p \circ \langle f, g \rangle = (p_1 \circ f) \circledast (p_2 \circ g) \leq (p_1 \circ f) \circledast (1 \circ g)$  and so by cancellation  $1 \circ f \leq p_1 \circ f \leq 1 \circ f$ , hence  $1 \circ f = p_1 \circ f$ . Thus  $\text{im } f \leq p_1$ . Similarly  $\text{im } g \leq p_2$ . Thus  $[\text{im } f, \text{im } g] \leq [p_1, p_2] = p$ . We have shown  $\text{im } \langle f, g \rangle = [\text{im } f, \text{im } g]$ .

Assume  $[p, q]$  is sharp. That is, there is some  $f$  with  $[p, q] = \text{im } f$ . Then  $[p, q] = \text{im } f = \text{im } \langle \triangleright_1 \circ f, \triangleright_2 \circ f \rangle = [\text{im } \triangleright_1 \circ f, \text{im } \triangleright_2 \circ f]$  and so  $p = \text{im } \triangleright_1 \circ f$  and  $q = \text{im } \triangleright_2 \circ f$ , which shows  $p$  and  $q$  are sharp.

To prove the converse, assume  $p$  and  $q$  are sharp. Pick  $f$  and  $g$  with  $p = \text{im } f$  and  $q = \text{im } g$ . Note  $f + g = \langle f \circ \triangleright_1, g \circ \triangleright_2 \rangle$ . We have  $\text{im } f = \text{im } f \circ \triangleright_1 \circ \kappa_1 \leq \text{im } f \circ \triangleright_1 \leq \text{im } f$  and so  $\text{im } f \circ \triangleright_1 = \text{im } f$ . Similarly  $\text{im } g \circ \triangleright_2 = \text{im } g$ . Thus  $\text{im } f + g = \text{im } \langle f \circ \triangleright_1, g \circ \triangleright_2 \rangle = [\text{im } f \circ \triangleright_1, \text{im } g \circ \triangleright_2] = [\text{im } f, \text{im } g] = [p, q]$ , which shows  $[p, q]$  is sharp.

**Solution to 205 IV.** Let  $C$  be an effectus with comprehension and images. Assume  $f: X \rightarrow Y$  is a quotient of sharp  $s$ . Let  $\pi_s$  denote a comprehension of  $s$ . Clearly  $1 \circ f \circ \pi_s = s^\perp \circ \pi_s = 0$ . Assume  $g: X \rightarrow Y'$  is some map with  $g \circ \pi_s = 0$ . Then  $1 \circ g \circ \pi_s = 0$  and so  $1 \circ g \leq \text{im}^\perp \pi_s = s^\perp$ . Thus by the universal property of  $f$  as a quotient, there is a unique map  $g': Y \rightarrow Y'$  with  $g = g' \circ f$ . This shows  $f$  is a cokernel of  $\pi_s$ .

To prove the converse, assume  $f$  is a cokernel of  $\pi_s$ . By assumption  $f \circ \pi_s = 0$ . Thus  $1 \circ f \circ \pi_s = 0$ . Hence  $1 \circ f \leq \text{im}^\perp \pi_s = s^\perp$ . Let  $g: X \rightarrow Y'$  be a map with  $1 \circ g \leq s^\perp$ . Then  $1 \circ g \circ \pi_s \leq s^\perp \circ \pi_s = 0$  and so  $g \circ \pi_s = 0$ . Thus by the universal property of  $f$  as a cokernel, there is a unique map  $g': Y \rightarrow Y'$  with  $g = g' \circ f$ . This shows  $f$  is a quotient of  $s$ .

**Solution to 207 II.** Assume  $s \leq t$ . Then  $s \circ f \leq t \circ f$ , hence  $(t \circ f)^\perp \leq (s \circ f)^\perp$  so  $[(t \circ f)^\perp] \leq [(s \circ f)^\perp]$ . Thus  $f^\circ(s) \equiv [s \circ f] \equiv [(s \circ f)^\perp]^\perp \leq [(t \circ f)^\perp]^\perp \equiv f^\circ(t)$ . This shows  $f^\circ$  is order preserving. Furthermore  $t^\perp \leq s^\perp$ , so  $f^\circ(t^\perp) \leq f^\circ(s^\perp)$ , hence  $f^\square(s) \equiv f^\circ(s^\perp)^\perp \leq f^\circ(t^\perp)^\perp \equiv f^\square(t)$ , which shows that  $f^\square$  is order preserving.

**Solution to 207 V.** Let  $f: X \rightarrow Y$  be some map in a  $\diamond$ -effectus.

1. Assume  $s \leq t$ . Clearly  $f_\diamond(s) \leq f_\diamond(t)$  so  $s \leq t \leq f^\square(f_\diamond(t))$ , hence  $f_\diamond(s) \leq f_\diamond(t)$ , which shows that  $f_\diamond$  is order preserving.
2. Assume  $D \subseteq \text{SPred } X$  is a set of sharp predicates with a supremum  $\text{sup } D$ . We will show  $f_\diamond(\text{sup } D)$  is the supremum of  $f_\diamond(D)$ . For any  $d \in D$ , we have  $d \leq \text{sup } D$  and so  $f_\diamond(d) \leq f_\diamond(\text{sup } D)$  by the previous point. Assume  $x$  is some sharp predicate of  $Y$  with  $f_\diamond(d) \leq x$  for all  $d \in D$  as well. Then  $d \leq f^\square(x)$  for all  $d \in D$  and so  $\text{sup } D \leq f^\square(x)$ . Hence  $f_\diamond(\text{sup } D) \leq x$ , which shows that  $\text{sup } f_\diamond(D) = f_\diamond(\text{sup } D)$ .

3. Assume  $D \subseteq \text{SPred } Y$  is a set of sharp predicates with an infimum  $\inf D$ . We will show  $f^\square(\inf D)$  is the infimum of  $f^\square(D)$ . For any  $d \in D$ , we have  $\inf D \leq d$  and so  $f^\square(\inf D) \leq f^\square(d)$  by 207 II. Assume  $x$  is some sharp predicate of  $X$  with  $x \leq f^\square(d)$  for all  $d \in D$  as well. Then  $f_\diamond(x) \leq d$  for all  $d \in D$  and so  $f_\diamond(x) \leq \inf D$ , hence  $x \leq f^\square(\inf D)$ . This shows that  $\inf f^\square(D) = f^\square(\inf D)$ .

4. For for any  $D \subseteq \text{SPred } Y$  with supremum, we have

$$\begin{aligned} f^\diamond(\sup D) &\equiv f^\square((\sup D)^\perp)^\perp \\ &= f^\square(\inf_{d \in D} (d^\perp))^\perp \\ &= (\inf_{d \in D} f^\square(d^\perp))^\perp \\ &= \sup_{d \in D} f^\square(d^\perp)^\perp \\ &= \sup_{d \in D} f^\diamond(d) \end{aligned}$$

and so  $f^\diamond$  preserves suprema.

5. For any  $s \in \text{SPred } X$  we have  $f_\diamond(s) \leq f_\diamond(s)$  and so  $s \leq f^\square(f_\diamond(s))$ . Similarly, for any  $t \in \text{SPred } Y$  we have  $f_\diamond(f^\square(t)) \leq t$ . Hence  $f_\diamond(f^\square(f_\diamond(s))) \leq f_\diamond(s) \leq f_\diamond(f^\square(f_\diamond(s)))$ , so  $f_\diamond = f_\diamond \circ f^\square \circ f_\diamond$ .

6. We have  $f^\square(f_\diamond(f^\square(t))) \leq f^\square(t) \leq f^\square(f_\diamond(f^\square(t)))$  and so  $f^\square = f^\square \circ f_\diamond \circ f^\square$ .

**Solution to 207 VIIa.** Let  $f, g: X \rightarrow Y$  be some maps in a  $\diamond$ -effectus. Assume  $f^\diamond = g^\diamond$  and  $t \in \text{SPred } X$ . Trivially  $g_\diamond(t) \leq (g_\diamond(t)^\perp)^\perp$  and so  $f^\diamond(g_\diamond(t)^\perp) = g^\diamond(g_\diamond(t)^\perp) \leq t^\perp$ . Thus  $f_\diamond(t) \leq g_\diamond(t)^\perp = g_\diamond(t)$ . Reasoning in the same way with  $f$  and  $g$  swapped, we see  $g_\diamond(t) \leq f_\diamond(t)$  and so  $f_\diamond = g_\diamond$ .

To prove the converse, assume  $f_\diamond = g_\diamond$  and  $s \in \text{SPred } Y$ . Trivially  $g^\diamond(s) \leq (g^\diamond(s)^\perp)^\perp$  and so  $f_\diamond(g^\diamond(s)^\perp) = g_\diamond(g^\diamond(s)^\perp) \leq s^\perp$ . Thus  $f^\diamond(t) \leq g^\diamond(t)^\perp = g^\diamond(t)$ . Reasoning in the same way with  $f$  and  $g$  swapped, we see  $g^\diamond(t) \leq f^\diamond(t)$  and so  $f^\diamond = g^\diamond$ .

**Solution to 208 XII.** As  $\xi_\diamond$  is left adjoint to  $\xi^\square$ , we have  $t \leq \xi^\square(\xi_\diamond(t))$ . Furthermore  $s = \lceil s^\perp \rceil^\perp = \lceil 1 \circ \xi \rceil^\perp = \xi^\square(0) \leq \xi^\square(\xi_\diamond(t))$ . Thus  $\xi^\square(\xi_\diamond(t))$  is an upper bound of  $s$  and  $t$ . We have to show it is the least sharp upper bound. So assume  $r$  is any sharp predicate with  $r \geq s$  and  $r \geq t$ . Then  $1 \circ \xi_r = r^\perp \leq s^\perp$  and so there is a (unique) map  $h$  with  $\xi_r = h \circ \xi$ . We have

$$(\xi_r)^\square = \xi^\square \circ h^\square = \xi^\square \circ \xi_\diamond \circ \xi^\square \circ h^\square = \xi^\square \circ \xi_\diamond \circ (\xi_r)^\square$$

and so  $r = (\xi_r)^\square(0) = \xi^\square(\xi_\diamond((\xi_r)^\square(0))) = \xi^\square(\xi_\diamond(r)) \geq \xi^\square(\xi_\diamond(t))$ .

**Solution to 209 II.** For the first point, assume  $f^\diamond = g_\diamond$  for some  $f : X \rightleftarrows Y : g$ . Trivially  $g^\diamond(s) \leq (g^\diamond(s)^\perp)^\perp$  for any  $s \in \text{SPred } X$  and so  $f^\diamond(g^\diamond(s)^\perp) = g_\diamond(g^\diamond(s)^\perp) \leq s^\perp$ . Hence  $f_\diamond(s) \leq g^\diamond(s)^{\perp\perp} = g^\diamond(s)$ . With essentially the same argument (swapping both  $f$  and  $g$  and the up/down position of the diamonds), we see  $g^\diamond(s) \leq f_\diamond(s)$ , hence  $g^\diamond = f_\diamond$ . Swapping  $f$  and  $g$  again, gives the argument for the converse.

For the second point, assume  $f$  and  $g$  are  $\diamond$ -adjoint. Then  $\text{im } f = f_\diamond(1) = g^\diamond(1) = [1 \circ g]$ , as desired.

**Solution to 209 III.** For the first point, assume  $f$  is  $\diamond$ -self-adjoint. Then  $(f \circ f)_\diamond = f_\diamond \circ f_\diamond = f^\diamond \circ f^\diamond = (f \circ f)^\diamond$  and so  $f \circ f$  is  $\diamond$ -self-adjoint as well.

For the second point, assume  $f$  is  $\diamond$ -positive. Then  $f = g \circ g$  for some  $\diamond$ -self-adjoint  $g$ . Hence  $f$  is  $\diamond$ -self-adjoint by the previous point.

For the third point, assume  $f$  is  $\diamond$ -positive and  $f \circ f$  is pure. By the previous point  $f$  is  $\diamond$ -self-adjoint and so  $f \circ f$  is  $\diamond$ -self-adjoint by the first point. Thus  $f \circ f$  is  $\diamond$ -positive.

**Solution to 210 II.** Assume  $f : X \rightarrow Y$  is a sharp map and  $p$  is a predicate on  $Y$ . Using 203 XIII and 203 XII we see  $[p \circ f] = [[p] \circ f] = [p] \circ f$ . For the converse assume  $[p] \circ f = [p \circ f]$  for every predicate  $p$  on  $Y$ . Assume  $s$  is any sharp predicate on  $Y$ . Then  $[s \circ f] = [s] \circ f = s \circ f$  and so  $s \circ f$  is sharp. Thus  $f$  is sharp.

**Solution to 211 XIV.** Note  $1 \circ \text{asrt}_p \circ \text{asrt}_p = p \circ \text{asrt}_p \equiv p \& p$ . By 209 III the map  $\text{asrt}_p \circ \text{asrt}_p$  is  $\diamond$ -positive and thus  $\text{asrt}_p \circ \text{asrt}_p = \text{asrt}_{p \& p}$  by uniqueness of  $\diamond$ -positive maps.

**Solution to 211 XV.** Assume  $f : X \rightarrow Y$  is a map in a  $\&$ -effectus with  $s \in \text{SPred } Y$  and  $t \in \text{SPred } X$ .

Assume  $\text{im } f \leq s$ . Then  $1 \circ f = s \circ f$ . Thus there is a  $g$  with  $f = \pi_s \circ g$ . Then  $\text{asrt}_s \circ f = \pi_s \circ \zeta_s \circ \pi_s \circ g = \pi_s \circ g = f$ . For the converse, assume  $\text{asrt}_s \circ f = f$ . Then  $1 \circ f = 1 \circ \text{asrt}_s \circ f = s \circ f$  and so  $\text{im } f \leq s$ . We have shown the first equivalence.

For the second equivalence, assume  $1 \circ f \leq t$ . Then  $f = h \circ \zeta_t$  for a (unique) map  $h$ . So  $f \circ \text{asrt}_t = h \circ \zeta_t \circ \pi_t \circ \zeta_t = h \circ \zeta_t = f$ . For the converse, assume  $f = f \circ \text{asrt}_t$ . Then  $1 \circ f = 1 \circ f \circ \text{asrt}_t \leq 1 \circ \text{asrt}_t = t$ .

**Solution to 213 V.** Clearly  $p \leq s$  if and only if  $1 \circ p \leq s$ . Even more trivially,  $s \& p = p$  if and only if  $p \circ \text{asrt}_s \equiv s \& p = p$ . Thus the result follows by applying the second equivalence of 211 XV with  $f = p$  and  $t = s$ .

**Solution to 213 VI.** Let  $f : X \rightarrow A$  and  $g : X \rightarrow B$  be maps in a  $\&$ -effectus. Assume  $\langle f, g \rangle$  is sharp. Pick any sharp predicate  $s$  on  $A$ . By 203 XIV we know  $[s, 0]$  is sharp and so  $[s, 0] \circ \langle f, g \rangle = (s \circ f) \vee (0 \circ g) = s \circ f$  is sharp as well. This shows  $f$  is sharp. With a similar argument we see  $g$  is sharp. For the converse, assume  $f$  and  $g$  are sharp. Assume  $[s, t]$  is some sharp predicate on  $A + B$ .

By 203XIV we know  $s$  and  $t$  are sharp. Thus  $s \circ f$  and  $t \circ g$  are sharp. Consequently  $[s, t] \circ \langle f, g \rangle = (s \circ f) \circ (t \circ g)$  is sharp as well by 208III. Thus  $\langle f, g \rangle$  is sharp, as desired.

**Solution to 216 X.** Assume  $C$  is a  $\&$ -effectus where every predicate has a square root and where  $\pi_s$  is  $\diamond$ -adjoint to  $\zeta_s$ . Let  $p$  be a predicate and  $s$  be another predicate on the same object that is sharp. Pick a predicate  $q$  with  $q \& q = p$ . The map  $f \equiv \pi_s \circ \text{asrt}_q \circ \zeta_s$  is pure and  $\diamond$ -self-adjoint:

$$f^\diamond = (\zeta_s)^\diamond \circ (\text{asrt}_q)^\diamond \circ (\pi_s)^\diamond = (\pi_s)_\diamond \circ (\text{asrt}_q)_\diamond \circ (\zeta_s)_\diamond = f_\diamond.$$

Thus  $f \circ f$  is  $\diamond$ -positive. By 211XIV, we get  $f \circ f = \pi_s \circ \text{asrt}_q \circ \zeta_s \circ \pi_s \circ \text{asrt}_q \circ \zeta_s = \pi_s \circ \text{asrt}_{q \& q} \circ \zeta_s = \pi_s \circ \text{asrt}_p \circ \zeta_s$  and so  $1 \circ f \circ f = 1 \circ \pi_s \circ \text{asrt}_p \circ \zeta_s = p \circ \zeta_s$ . By uniqueness of  $\diamond$ -positive maps, we get  $\pi_s \circ \text{asrt}_p \circ \zeta_s = f \circ f = \text{asrt}_{1 \circ f \circ f} = \text{asrt}_{p \circ \zeta_s}$ . Thus  $\text{asrt}_p \circ \zeta_s = \zeta_s \circ \pi_s \circ \text{asrt}_p \circ \zeta_s = \zeta_s \circ \text{asrt}_{p \circ \zeta_s}$ .

**Solution to 217 III.** By 211XV we have

$$\text{asrt}_p = \text{asrt}_p \circ \text{asrt}_{[p]} = \pi_{[p]} \circ \text{id} \circ \zeta_{[p]} \circ \text{asrt}_p$$

and so

$$\text{asrt}_p^\dagger = \text{asrt}_p \circ \pi_{[p]} \circ \text{id}^{-1} \circ \zeta_{[p]} = \text{asrt}_p \circ \text{asrt}_{[p]} = \text{asrt}_p.$$

Next, note  $\zeta_1 = \pi_1 = \text{asrt}_1 = \text{id}$  and so  $\pi_s = \pi_s \circ \text{id} \circ \zeta_1 \circ \text{asrt}_1$ , hence

$$\pi_s^\dagger = \text{asrt}_1 \circ \pi_1 \circ \text{id}^{-1} \circ \zeta_s = \zeta_s.$$

Also  $\zeta_s = \pi_1 \circ \text{id} \circ \zeta_s \circ \text{asrt}_1$  and so

$$\zeta_s^\dagger = \text{asrt}_1 \circ \pi_s \circ \text{id}^{-1} \circ \zeta_1 = \pi_s.$$

Finally, note  $\alpha = \text{asrt}_1 = \pi_1 \circ \alpha \circ \zeta_1$  for any isomorphism  $\alpha$  and so

$$\alpha^\dagger = \text{asrt}_1 \circ \pi_1 \circ \alpha^{-1} \circ \zeta_1 = \alpha^{-1}.$$

**Solution to 218 VI.** Assume  $h$  is a pristine map in a  $\&$ -effectus. By 212III we know there is an isomorphism  $\alpha$  with  $h = \pi_{\text{im } h} \circ \alpha \circ \zeta_{[1 \circ h]} \circ \text{asrt}_{1 \circ h}$ . As  $1 \circ h$  is sharp, we have  $\zeta_{[1 \circ h]} \circ \text{asrt}_{1 \circ h} = \zeta_{1 \circ h} \circ \pi_{1 \circ h} \circ \zeta_{1 \circ h} = \zeta_{1 \circ h}$  and so  $h = \pi_{\text{im } h} \circ \alpha \circ \zeta_{1 \circ h}$ .

**Solution to 218 IX.** Assume  $h \equiv \pi_{\text{im } h} \circ \alpha \circ \zeta_{1 \circ h}$  is a pristine map in a  $\dagger'$ -effectus, cf. 218VI.

1. Clearly  $h = \pi_{\text{im } h} \circ \alpha \circ \zeta_{[1 \circ h]} \circ \text{asrt}_{1 \circ h}$  as  $1 \circ h$  is sharp and by 211XV. Thus

$$h^\dagger = \text{asrt}_{1 \circ h} \circ \pi_{[1 \circ h]} \circ \alpha^{-1} \circ \zeta_{\text{im } h} = \pi_{1 \circ h} \circ \alpha^{-1} \circ \zeta_{\text{im } h}$$

using other equivalence of 211XV.

2. Note that by the previous point  $1 \circ h^\dagger = \text{im } h$  and  $\text{im } h^\dagger = 1 \circ h$ , hence by a second appeal to the previous point:

$$h^{\dagger\dagger} = \pi_{1 \circ h^\dagger} \circ (\alpha^{-1})^{-1} \circ \zeta_{\text{im } h^\dagger} = \pi_{\text{im } h} \circ \alpha \circ \zeta_{1 \circ h} = h.$$

3. By the first point  $h^\dagger \circ h = \pi_{1 \circ h} \circ \alpha^{-1} \circ \zeta_{\text{im } h} \circ \pi_{\text{im } h} \circ \alpha \circ \zeta_{1 \circ h} = \pi_{1 \circ h} \circ \zeta_{1 \circ h} = \text{asrt}_{1 \circ h}$  as promised.

4. Clearly  $p \circ h^\dagger \leq 1 \circ h^\dagger = \text{im } h$ .

5. Assume  $p \leq 1 \circ h$ . Then  $p \circ \text{asrt}_{1 \circ h} = p$  by 211XV and so

$$\text{asrt}_{p \circ h^\dagger} \circ h \stackrel{218\text{VII}}{=} h \circ \text{asrt}_{p \circ h^\dagger \circ h} \stackrel{\text{pt. 3}}{=} h \circ \text{asrt}_{p \circ \text{asrt}_{1 \circ h}} = h \circ \text{asrt}_p,$$

as desired.

**Solution to 219 IX.** The map  $\alpha \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \varphi^{-1} \circ \zeta_s$  is a quotient for  $[t \circ f^\dagger]$ :

$$\begin{aligned} 1 \circ \alpha \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \varphi^{-1} \circ \zeta_s &= [t \circ \text{asrt}_p \circ \pi_{[p]}] \circ \varphi^{-1} \circ \zeta_s \\ &= [t \circ \text{asrt}_p \circ \pi_{[p]} \circ \varphi^{-1} \circ \zeta_s] \\ &= [t \circ f^\dagger]. \end{aligned}$$

Furthermore

$$\begin{aligned} \alpha \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \varphi^{-1} \circ \zeta_s \circ \pi_{[t \circ f^\dagger]} \\ \stackrel{(3.16)}{=} \alpha \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \varphi^{-1} \circ \zeta_s \circ \pi_s \circ \varphi \circ \pi_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \alpha^{-1} \\ = \text{id}. \end{aligned}$$

Thus  $\alpha \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \varphi^{-1} \circ \zeta_s$  is the unique quotient corresponding to the comprehension  $\pi_{[t \circ f^\dagger]}$ . Hence  $\alpha \circ \zeta_{[t \circ \text{asrt}_p \circ \pi_{[p]}]} \circ \varphi^{-1} \circ \zeta_s = \zeta_{[t \circ f^\dagger]}$ , as desired.

**Solution to 219 X.** Note that  $\pi_{[q]} \circ \psi^{-1} = \pi_{[q]} \circ \psi^{-1} \circ \zeta_1 \circ \text{asrt}_1$  and so by definition of the dagger, we have  $(\pi_{[q]} \circ \psi^{-1})^\dagger = \text{asrt}_1 \circ \pi_1 \circ (\psi^{-1})^{-1} \circ \zeta_{[q]} = \psi \circ \zeta_{[q]}$ . Furthermore  $p \circ \pi_t \leq 1 = 1 \circ \pi_{[q]} \circ \psi^{-1}$ . Thus we can apply the fourth point of 218 IX:

$$\begin{aligned} \pi_{[q]} \circ \psi^{-1} \circ \text{asrt}_{p \circ \pi_t} &= \text{asrt}_{p \circ \pi_t \circ (\pi_{[q]} \circ \psi^{-1})^{-1}} \circ \pi_{[q]} \circ \psi^{-1} \\ &= \text{asrt}_{p \circ \pi_t \circ \psi \circ \zeta_{[q]}} \circ \pi_{[q]} \circ \psi^{-1} \\ &= \text{asrt}_{p \circ k} \circ \pi_{[q]} \circ \psi^{-1}, \end{aligned}$$

which is the desired equality.

**Solution to 221 IIIa.** Assume  $\mathcal{A}$  is a commutative von Neumann algebra with effect  $a \in \mathcal{A}$ . The standard corner  $h_a: \mathcal{A} \rightarrow [a]\mathcal{A}[a]$  for  $a$  (see 169 IV) is given by  $b \mapsto [a]b[a]$ . If  $\mathcal{A}$  is commutative, then  $h_a(bc) = [a]bc[a] = [a]b[a][a]c[a] = h_a(b)h_a(c)$  and so  $h_a$  is an nmiu-map. Every corner is a standard corner up-to an isomorphism, thus every corner on a commutative von Neumann algebra is an nmiu-map as well. If  $\text{CvN}^{\text{op}}$  were to have dilations, then any ncpu-map would be the composition of a corner and an nmiu-map. We just saw that such corners are nmiu and so any ncp-map would be an nmiu-map as well, which is absurd. Thus  $\text{CvN}^{\text{op}}$  does not have dilations in the sense of 221 II.

**Solution to 224 VI.** We will first show that for any non-zero pure map  $f: \mathcal{A} \rightarrow \mathbb{C}$ , there is a Hilbert space  $\mathcal{H}$ , an element  $x \in \mathcal{H}$ , von Neumann algebra  $\mathcal{C}$  and isomorphism  $\varphi: \mathcal{B}(\mathcal{H}) \oplus \mathcal{C} \rightarrow \mathcal{A}$  with  $f(\varphi(T, c)) = \langle x, Tx \rangle$ . By definition of purity, we know  $f = c \circ h$  for some corner  $h$  and filter  $c$  for  $f(1)$ . Clearly  $c(1) = f(1) \neq 0$  as  $f \neq 0$  and so  $c: \mathbb{C} \rightarrow \mathbb{C}$  is simply given by  $c(1)\text{id}$ , hence  $f = f(1)h$  for a corner  $h: \mathcal{A} \rightarrow \mathbb{C}$  of  $p \in \mathcal{A}$ . As  $p\mathcal{A}p \cong \mathbb{C}$  both have only two projections and so  $p$  is a minimal projection. In particular 0 is the only central element strictly below  $p$ , thus  $\llbracket p \rrbracket \mathcal{A}$  is a factor. Let  $\varrho: \llbracket p \rrbracket \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ ,  $y \in \mathcal{H}$  be a GNS-representation of the restriction of  $h$  to  $\llbracket p \rrbracket \mathcal{A}$  with  $y \in \mathcal{H}$  such that  $h(\llbracket p \rrbracket a) = \langle y, \varrho(\llbracket p \rrbracket a)y \rangle$ . This restriction of  $h$  is still a corner, hence pure and so by 171 VII we know  $\varrho$  is surjective. As  $\llbracket p \rrbracket \mathcal{A}$  is a factor,  $\varrho$  must be injective as well. Thus  $\varrho$  is an nmiu-isomorphism. Hence  $\varphi: \mathcal{B}(\mathcal{H}) \oplus \mathcal{B} \rightarrow \mathcal{A}$  defined by  $\varphi(T, c) \equiv \varrho^{-1}(T) + c$  is an nmiu-isomorphism as well. Define  $x \equiv \sqrt{f(1)}$ . Then  $f(\varphi(T, c)) = f(1)h(\varrho^{-1}(T) + c) = f(1)h(\varrho^{-1}(T)) = f(1)\langle y, \varrho(\varrho^{-1}(T))y \rangle = f(1)\langle y, Ty \rangle = \langle x, Tx \rangle$  as desired.

We continue with the main task of this exercise. Assume  $(\text{Pure vN}^{\text{op}})^{\text{op}}$  has a product  $\mathbb{C} \xleftarrow{\pi_1} \mathcal{A} \xrightarrow{\pi_2} \mathbb{C}$  of  $\mathbb{C}$  and  $\mathbb{C}$ . Note  $\pi_1 \neq 0$  for otherwise  $\text{id}_{\mathbb{C}} = \pi_1 \circ h = 0$ , where  $h: \mathbb{C} \rightarrow \mathcal{A}$  is the unique map given by the universal property of the product for  $\text{id}_{\mathbb{C}}$  and  $\text{id}_{\mathbb{C}}$ . Similarly  $\pi_2 \neq 0$ .

By assumption, there is a unique pure map  $h: \mathbb{C} \rightarrow \mathcal{A}$  with  $\pi_1 \circ h = \text{id}$  and  $\pi_2 \circ h = \text{id}$ . With the reasoning at the start, we know there is a Hilbert space  $\mathcal{H}$ , an element  $z \in \mathcal{H}$ , a von Neumann algebras  $\mathcal{C}$  and an isomorphism  $\varphi: \mathcal{B}(\mathcal{H}) \oplus \mathcal{C}$  with  $h^\dagger(\varphi(T, c)) = \langle z, Tz \rangle$ . Note that  $T \mapsto \langle z, Tz \rangle$  is the dagger of the standard filter  $c_{|z\rangle\langle z|}$ . Write  $p: \mathcal{B}(\mathcal{H}) \oplus \mathbb{C} \rightarrow \mathcal{B}(\mathcal{H})$  for the regular coprojection  $(T, c) \mapsto T$ , which is corner with  $p^\dagger(T) = (T, 0)$ . Then  $h^\dagger \circ \varphi = c_{|z\rangle\langle z|}^\dagger \circ p$  and so  $h = \varphi \circ p^\dagger \circ c_{|z\rangle\langle z|}$ . By definition of  $h$ , we have

$$\text{id} = \pi_1 \circ h = \pi_1 \circ \varphi \circ p^\dagger \circ c_{|z\rangle\langle z|}. \quad (4.4)$$

Reasoning the same way as we did for  $f$  at the start of this exercise, we know that  $\llbracket \pi_1 \circ \varphi \rrbracket$  is a minimal projection. If  $\llbracket \pi_1 \circ \varphi \rrbracket \leq (0, 1)$ , then  $\pi_1 \circ \varphi \circ p^\dagger = 0$  and so  $\text{id} = 0$  by (4.4), which is absurd. Thus  $\llbracket \pi_1 \circ \varphi \rrbracket = (p_1, 0)$  for some minimal projection  $p_1$  of  $\mathcal{B}(\mathcal{H})$ . Thus  $\pi_1(\varphi(T, c)) = \langle x, Tx \rangle$  for some  $x \in \mathcal{H}$ . Similarly  $\pi_2(\varphi(T, c)) = \langle y, Ty \rangle$  for some  $y \in \mathcal{H}$ .

Note that  $\pi_i \circ \varphi \circ (\text{id} \oplus \vartheta) = \pi_i \circ \varphi$  for any map  $\vartheta: \mathcal{C} \rightarrow \mathcal{C}$  and  $i = 1, 2$ . Hence  $\text{id} \circ \pi_i \circ \varphi \circ (\text{id} \oplus \vartheta) \circ \varphi^{-1} \circ h$  for  $i = 1, 2$  and so  $(\text{id} \oplus \vartheta) \circ \varphi^{-1} \circ h = \varphi^{-1} \circ h$  for every  $\vartheta$  by uniqueness of  $h$ , which is absurd unless  $\mathcal{C} = 0$ .

Suppose  $\dim \mathcal{H} \geq 3$ . Then there is some non-zero  $z_0 \in \mathcal{H}$  with  $z_0 \perp x$  and  $x_0 \perp y$ . So  $\pi_i \circ \varphi \circ c_{|z_0\rangle\langle z_0|} = 0$  for  $i = 1, 2$ . By the presumed universal property of  $\mathcal{A}$ , there is a unique map  $h: \mathbb{C} \rightarrow \mathcal{A}$  with  $\pi_1 \circ h = \pi_1 \circ \varphi \circ c_{|z_0\rangle\langle z_0|}$  and  $\pi_2 \circ h = \pi_2 \circ \varphi \circ c_{|z_0\rangle\langle z_0|}$ . Both  $\varphi \circ c_{|z_0\rangle\langle z_0|}$  and  $0$  fit the bill, so  $\varphi \circ c_{|z_0\rangle\langle z_0|} = 0$ , which is absurd. Thus  $\dim \mathcal{H} \leq 2$ .

Clearly  $\dim \mathcal{H} \neq 0$ , for otherwise there would be a unique pure map  $\mathbb{C} \rightarrow \mathbb{C}$ , which is absurd. If  $\dim \mathcal{H} = 1$ , then  $x = \lambda y$  for some  $\lambda \in \mathbb{C}, \lambda \neq 0$  and so  $\pi_1 = \lambda \pi_2$ , hence for any two pure maps  $f_1, f_2: \mathcal{B} \rightarrow \mathbb{C}$  we would have  $f_1 = \lambda f_2$ , which is absurd.

Reducing clutter a bit, we may assume without loss of generality  $\mathcal{A} = M_2$ ,  $\pi_1(T) = \langle x, Tx \rangle$  and  $\pi_2(T) = \langle y, Ty \rangle$  for some  $x, y \in \mathbb{C}^2$ . Write  $f_1, f_2: \mathbb{C}^2 \rightarrow \mathbb{C}$  for the maps  $f_1(\lambda, \mu) = \lambda$  and  $f_2(\lambda, \mu) = \mu$ . By the presumed universal property of  $M_2$  as product, there is a unique pure map  $f: \mathbb{C}^2 \rightarrow \mathcal{A}$  with  $\pi_1 \circ f = f_1$  and  $\pi_2 \circ f = f_2$ . From the definition of purity, it follows that  $f = c \circ h$  for some corner  $h: \mathbb{C}^2 \rightarrow [h(1)] M_2 [h(1)]$  and the standard filter  $c: [h(1)] M_2 [h(1)] \rightarrow M_2$ . Note  $[h] \mathbb{C}^2 [h] \cong [h(1)] M_2 [h(1)]$ . The former is commutative, so  $[h(1)] \neq 1$ . If  $[h(1)] = 0$ , then  $h = 0$  and so  $f_1 = 0$ , quod non. Thus  $[h(1)] M_2 [h(1)] \cong \mathbb{C}$  and so either  $[h] = (1, 0)$  or  $[h] = (0, 1)$ . In the first case  $1 = f_2(0, 1) = \pi_2(h(0, 1)) = \pi_2(0) = 0$ , which is a contradiction. The second case leads to a similar contradiction. So there is no product of  $\mathbb{C}$  and  $\mathbb{C}$  in  $(\text{Pure vN}^{\text{op}})^{\text{op}}$ .

**Solution to 224 VII.** Assume  $\xi: \mathcal{C} \rightarrow M_4$  is some filter with  $\text{ad}_\sigma \circ \xi = \xi$ . Note  $\sigma^* = \sigma$ . Assume  $c \in \mathcal{C}$ . Then  $\xi(c) = \text{ad}_\sigma(\xi(c)) = \sigma \xi(c) \sigma$ . Let  $b \in M_4$  be any element with  $\sigma b \sigma = b$ . Then  $(p_{\mathcal{J}} + p_{\mathcal{A}})b(p_{\mathcal{J}} + p_{\mathcal{A}}) = b = \sigma b \sigma = (\sigma p_{\mathcal{J}} + \sigma p_{\mathcal{A}})b(p_{\mathcal{J}} \sigma + p_{\mathcal{A}} \sigma) = (p_{\mathcal{J}} - p_{\mathcal{A}})b(p_{\mathcal{J}} - p_{\mathcal{A}})$  and so  $p_{\mathcal{A}} b p_{\mathcal{J}} + p_{\mathcal{J}} b p_{\mathcal{A}} = 0$ . As  $p_{\mathcal{A}}$  and  $p_{\mathcal{J}}$  have orthogonal ranges, we see  $p_{\mathcal{A}} b p_{\mathcal{J}} = 0 = p_{\mathcal{J}} b p_{\mathcal{A}}$ . Hence  $b = p_{\mathcal{J}} b p_{\mathcal{J}} + p_{\mathcal{A}} b p_{\mathcal{A}}$ . Furthermore  $b p_{\mathcal{J}} = p_{\mathcal{J}} b p_{\mathcal{J}}$ . If  $b$  is additionally self adjoint, then  $p_{\mathcal{J}} b = p_{\mathcal{J}} b^* p_{\mathcal{J}} = p_{\mathcal{J}} b p_{\mathcal{J}} = p_{\mathcal{J}} b$ . Thus  $p_{\mathcal{J}} \xi(c) = \xi(c) p_{\mathcal{J}}$  for any  $c \in \mathcal{C}$ . There is some isomorphism  $\vartheta: \mathcal{C} \rightarrow [\xi(1)] M_4 [\xi(1)]$  with  $\xi(c) = \sqrt{\xi(1)} \vartheta(c) \sqrt{\xi(1)}$  and so  $p_{\mathcal{J}} \sqrt{\xi(1)} b \sqrt{\xi(1)} = \sqrt{\xi(1)} b \sqrt{\xi(1)} p_{\mathcal{J}}$  for any  $b \in M_4$ . From this it follows (using the pseudoinverse of  $\sqrt{\xi(1)}$ ) that  $p_{\mathcal{J}} [\xi(1)] b [\xi(1)] = [\xi(1)] b [\xi(1)] p_{\mathcal{J}}$ . Thus  $[\xi(1)] p_{\mathcal{J}} [\xi(1)]$  is central in  $[\xi(1)] M_4 [\xi(1)]$ . So either  $[\xi(1)] p_{\mathcal{J}} [\xi(1)] = 0$  or  $[\xi(1)] p_{\mathcal{J}} [\xi(1)] = [\xi(1)]$ . In the first case, we have  $[\xi(1)] p_{\mathcal{J}} = 0$  hence  $\xi(c) = p_{\mathcal{A}} \xi(c) p_{\mathcal{A}} + p_{\mathcal{J}} \xi(c) [\xi(1)] p_{\mathcal{J}} = p_{\mathcal{A}} \xi(c) p_{\mathcal{A}} \leq p_{\mathcal{A}}$  for all  $c \in \mathcal{C}$ . With similar reasoning we see that  $\xi(c) \leq p_{\mathcal{J}}$  for all  $c \in \mathcal{C}$  in the second case.

Assume there is an equalizer  $e: E \rightarrow M_4$  of  $\text{id}_{M_4}$  and  $\text{ad}_\sigma$  in  $(\text{Pure vN}^{\text{op}})^{\text{op}}$ . By definition of purity  $e = c \circ h$  for some filter  $c$  and comprehension  $h$ . As  $e$  is supposed to be an equalizer of  $\text{id}$  and  $\text{ad}_\sigma$ , we have  $\text{ad}_\sigma \circ c \circ h = \text{ad}_\sigma \circ e = e = c \circ h$ , hence  $\text{ad}_\sigma \circ c = c$  by the surjectivity of the corner  $h$ . We have

just shown that then either  $c(b) \leq p_{\mathcal{A}}$  for all  $b$  or  $c(b) \leq p_{\mathcal{S}}$  for all  $b$ . Hence  $h(e) \leq p_{\mathcal{A}}$  for all  $e \in E$  or  $h(e) \leq p_{\mathcal{S}}$  for all  $e \in E$ . In the first case, consider the map  $\text{ad}_{e_{\mathcal{S}}^\dagger} : M_3 \rightarrow M_4$ . Clearly  $\text{ad}_\sigma \circ \text{ad}_{e_{\mathcal{S}}^\dagger} = \text{ad}_{e_{\mathcal{S}}^\dagger}$  and so there must be a (unique) map  $h : M_3 \rightarrow E$  with  $e \circ h = \text{ad}_{e_{\mathcal{S}}^\dagger}$ . However  $p_{\mathcal{A}} \geq e(h(T)) = \text{ad}_{e_{\mathcal{S}}^\dagger}(T) \leq p_{\mathcal{S}}$  for all  $T \in M_3$  and so  $\text{ad}_{e_{\mathcal{S}}^\dagger} = 0$ , quod non. We reach a contradiction in the other case in a similar fashion. Hence  $\text{Pure}\mathbf{vN}^{\text{op}}$  has no coequalizers.



# Errata

These are errata to the printed version of this thesis, which have been corrected in this PDF. Additional errata will be published on the arXiv under 1803.01911.

**Erratum to 138 VII.** The second part of the exercise starts with “Conclude that any quantum channel  $\Phi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H}) \dots$ ”. This should have read “Conclude that any quantum channel  $\Phi$  from  $\mathcal{H}$  to itself ...”

**Erratum to 139 XI.** The third-to-last sentence reads “Derive from the latter that for each  $y \in \mathcal{H}'$  and rank-one projector  $e \in \mathcal{H}$ , there is a  $y' \in \mathcal{H}'$  with  $U_0(e \otimes y) = e \otimes y'$ .” This should have been “Derive from the latter that for each  $y \in \mathcal{H}'$  and unit vector  $e \in \mathcal{H}$ , there is a  $y' \in \mathcal{H}'$  with  $U_0(e \otimes y) = e \otimes y'$ .”

**Erratum to 141 IIa.** A pre-Hilbert  $\mathcal{B}$ -module  $X$  is **self dual** if every bounded  $\mathcal{B}$ -linear  $\tau: X \rightarrow \mathcal{B}$  is of the form  $\tau(x) = \langle t, x \rangle$  for some  $t \in X$ . In print the condition that  $\tau$  should be bounded was accidentally lost when it was moved from 149 V.

**Erratum to 147 II.** The assumption that the net  $x_\alpha$  is Cauchy in point 3 is superfluous.

**Erratum to 152 VIII.** The map  $T$  should be presumed to be bounded.

**Erratum to 152 IX.** The  $C^*$ -algebra  $\mathcal{B}$  isn't assumed to be a von Neumann algebra, which it should have been (for otherwise 150 II wouldn't be applicable.)

**Erratum to 159 VIII.** The displayed inequality

$$|f(T - p_S T p_S)| \leq |f((1 - p_S)T) + |p_S T(1 - p_S)|$$

should have read  $|f(T - p_S T p_S)| \leq |f((1 - p_S)T)| + |f(p_S T(1 - p_S))|$ .

**Erratum to 169 X.** The standard filter  $c_b$  for  $b \in \mathcal{B}$  is a map “[ $b$ ]  $\mathcal{B}$  [ $b$ ]  $\rightarrow \mathcal{B}$ ” instead of a map “[ $\mathcal{B} \rightarrow [b] \mathcal{B} [b]$ ”.

**Erratum to 197 V.** Not every zero map is a quotient; only those into 0.

**Erratum to 199 VII.** Not every zero map is a comprehension; only those from 0.

**Erratum to 218 IX.** Also in the last two points of the exercise, the map  $h$  should be presumed to be pristine.

**Erratum to 224 VI.** The condition “ $f \circ \varphi = \langle x, (\cdot)x \rangle$ ” is a typo. What was meant is “ $f(\varphi(T, c)) = \langle x, Tx \rangle$  for all  $c \in \mathcal{C}$  and  $T \in \mathcal{B}(\mathcal{H})$ ”. The same mistake was made writing “ $\pi_1 \circ \varphi = \langle x, (\cdot)x \rangle$ ” and “ $\pi_2 \circ \varphi = \langle y, (\cdot)y \rangle$ ”.