

Hypoxia-inducible factor 1 α protects peripheral sensory neurons from diabetic peripheral neuropathy by suppressing accumulation of reactive oxygen species.

Daniel Rangel Rojas¹, Irmgard Tegeder², Rohini Kuner¹, Nitin Agarwal^{1*}

¹Institute of Pharmacology, Heidelberg University, Im Neuenheimer Feld 366, D-69120 Heidelberg, Germany.

²Institute for Clinical Pharmacology, Goethe-University Hospital, Frankfurt, Germany.

*Address correspondence to N.A, Institute of Pharmacology, Heidelberg University, Im Neuenheimer Feld 366, D-69120 Heidelberg, Germany. E mail: nitin.agarwal@pharma.uni-heidelberg.de

ABSTRACT

Diabetic peripheral neuropathy (DPN) is one of the most common diabetic complications. Mechanisms underlying nerve damage and sensory loss following metabolic dysfunction remain large unclear. Recently, hyperglycemia-induced mitochondrial dysfunction and the generation of ROS have gained attention as possible mechanisms of organ damage in diabetes. Hypoxia-inducible factor 1 (HIF1 α) is a key transcription factor activated by hypoxia, hyperglycemia, nitric oxide as well as ROS, suggesting a fundamental role in DPN susceptibility. Genetically-modified mutant mice, which conditionally lack HIF1 α in peripheral sensory neurons (SNS-HIF1 $\alpha^{-/-}$), were analyzed longitudinally up to 6 months in the streptozotocin (STZ) model of type1 diabetes. Behavioral measurements of sensitivity to thermal and mechanical stimuli, quantitative morphological analyses of intraepidermal nerve fiber density and measurements of reactive oxygen species (ROS) in sensory neurons in vivo were undertaken over several months post-STZ injections to delineate the role of HIF1 α .

Abbreviations: HIF1 α - Hypoxia-inducible factor 1 α ; ROS - reactive oxygen species; STZ - Streptozotocin; DPN - Diabetic peripheral neuropathy; DRG - dorsal root ganglia; SNS – sensory neuron-specific

in DPN. Longitudinal behavioral and morphological analyses at 5, 13 and 24 weeks post-STZ treatment revealed that SNS-HIF1 α ^{-/-} developed stronger hyperglycemia-evoked losses of peripheral nociceptive sensory axons associated with stronger losses of mechano- and heat sensation with a faster onset than HIF1 α ^{fl/fl} mice. Mechanistically, these histomorphologic and behavioral differences were associated with significantly higher level of STZ-induced production of ROS in sensory neurons of SNS-HIF1 α ^{-/-} mice as compared with HIF1 α ^{fl/fl}. Our results indicate that HIF1 α is as an upstream modulator of ROS in peripheral sensory neurons and exerts a protective function in suppressing hyperglycemia-induced nerve damage by limiting ROS levels. HIF1 α stabilization may be thus a new strategy target for limiting sensory loss, a debilitating late complication of diabetes.

KEY MESSAGES

- Impaired Hypoxia-inducible factor 1 α (HIF1 α) signaling leads to early onset of STZ-induced loss of sensation in mice
- STZ-induced loss of sensation in HIF1 α is associated with loss of sensory nerve fiber in skin.
- Activation of HIF1 α signaling in diabetic mice protects the sensory neurons by limiting ROS formation generated due to mitochondrial dysfunction.

KEYWORDS: Hyperglycemia, DPN, ROS, Streptozotocin, sensory neurons

Abbreviations: HIF1 α - Hypoxia-inducible factor 1 α ; ROS - reactive oxygen species; STZ - streptozotocin; DPN - Diabetic peripheral neuropathy; DRG - dorsal root ganglia; SNS – sensory neuron-specific

INTRODUCTION

Diabetes is a chronic progressive metabolic disease characterized by elevated levels of blood glucose. It is one of the largest health care problems worldwide, with more than 422 million diabetic patients [1]. Diabetic peripheral neuropathy (DPN) is one of the most common diabetic complications. Patients with DPN experience pain, tingling, prickling sensations and numbness, which can progress to complete loss of sensation and motor dysfunction. These late diabetic complications in combination with vascular damage are the leading cause of lower extremity amputations [2, 3]. Clinical research from last two decades indicates that rigorous control of glucose reduces the incidence of DPN, suggesting a major role for hyperglycemia [4].

Uncovering molecular mechanisms underlying DPN remains a major challenge and a lack of mechanistic understanding has long hampered effective therapy of DPN. Only recently, hyperglycemia-induced mitochondrial dysfunction and the subsequent generation of ROS have gained attention as possible mechanisms [5]. Prolonged hyperglycemia energizes activation of the polyol [6], and hexosamine pathways [7], activation of PKC isoforms [8] and production of advanced glycation end products (AGE) [9, 10], thereby cumulatively increasing the intracellular ratio of NADH/NAD⁺. It has been hypothesized that this depletion of cellular antioxidant potential leads to cellular damage, which is marked by the progression of DPN [11]. Hyperglycemia-induced ROS formation causes the development of a pseudohypoxia state, which modulates intracellular signaling pathways such as mitogen activation protein kinases (MAPKs), nuclear factor kappa B (NF- κ B), activator protein I (AP-1) and importantly, hypoxia-inducible factor-1 (HIF-1) [12, 13].

HIF1 is a heterodimeric transcriptional factor composed of α and β subunits (HIF1 α and HIF1 β respectively) [14, 15]. HIF1 α is an oxygen-sensitive subunit and its activity is tightly regulated by cellular oxygen concentration [15], while HIF1 β is constitutively expressed. In conditions of normoxia, HIF1 α is subjected to oxygen-dependent hydroxylation at two prolyl residues, which enables its binding to Hippiel-Lindau protein (VHL) leading to polyubiquitination and subsequent proteasomal degradation [16, 17]. In contrast, under

hypoxic conditions, hydroxylation of HIF1 α is inhibited, resulting in accumulation of HIF1 α and its heteromerization with HIF1 β . In consequence, stabilized transcription factor HIF1 binds to hypoxia response elements (HREs) and regulates the expression of target genes, leading to biological consequences.

In contrast to the clarity on hypoxia-mediated modulation of HIF1 α , the role of hyperglycemia in the modulation of HIF1 α is subject to much debate. Previous studies report contradictory observations, e.g. hyperglycemia-induced stabilization of HIF1 α in pancreatic cancer cells versus its inhibition in primary dermal fibroblasts [33, 45]. Moreover, there is no knowledge on the role of HIF1 α in hyperglycemia-induced dysfunction in peripheral sensory neurons in the course of DPN. Recently, it was reported that HIF1 α deletion results in exaggerated acute thermal and cold nociception [18] in naïve mice, suggesting that HIF1 α activity is important in peripheral nociceptive neurons. However, to date, the role of HIF1 α in pathological DPN-related structural and functional changes in peripheral sensory neurons has not been studied.

Here, we investigated the role of HIF1 α in peripheral pain-sensing (nociceptive) neurons over the chronic progression of DPN in mice genetically lacking HIF1 α conditionally in the dorsal root ganglia (DRG) by undertaking detailed longitudinal analyses spanning 6 months following induction of type1 diabetes in the streptozotocin (STZ) model. Our results uncover a previously undescribed protective role of HIF1 α in nerve damage and sensory loss associated with chronic stages of DPN and lead to the surprising observation that HIF1 α is an upstream modulator of ROS levels and oxidative stress in peripheral sensory neurons under diabetic conditions.

MATERIAL AND METHODS:

Animal experiments

Age-matched 7-9 wks old SNS-HIF1 α ^{-/-} and HIF1 α ^{fl/fl} mice were used for experiments. Animals were maintained in a temperature controlled room with a light/dark cycle (12/12). Four to five littermates were housed in individually ventilated cage-rack system (Techniplast, Italy) with free access to water and food. All animal experiments were done in accordance with ethical guidelines and approved by Regierungspräsidium Karlsruhe, Germany.

Sensory-neuron-specific knockout mice

Mice lacking HIF-1 α (SNS-HIF1 α ^{-/-}) in a sensory neuron-specific manner were described previously [18]. Briefly, mice carrying homozygous alleles of Hif1 α gene (HIF1 α ^{fl/fl}) were mated with mice expressing the Cre recombinase selectively in nociceptors (SNSCre) [19] to obtain homozygous SNS-HIF1 α ^{-/-} and HIF1 α ^{fl/fl} (control littermates). We have previously described the basic characterization of these mice in the study by Kanngiesser et al [18].

Rodent model of type 1 diabetic model

Diabetes was induced by low dose multiple injections of STZ (60 mg/kg/d) in citrate buffer, intraperitoneally (i.p), in 8 wks-old SNS-HIF1 α ^{-/-} or HIF1 α ^{fl/fl} mice for 5 consecutive days. Blood glucose levels were measured once weekly using a glucometer (Accu-Chek, Roche Diagnostics) for the entire course of the experiment. Mice above > 350 mg/dl were considered to be diabetic. Mice were analyzed over a period of 5 wks to 24 wks post-STZ injection. At 2 weeks (wks) post-STZ injection, we achieved blood glucose between 380 and 500 mg/dl. The blood glucose levels were maintained in a range between 380 and 480 mg/dl by administering insulin sub-cutaneously as required over the entire course of an experiment to ensure uniformity and consistency.

Behavioral measurements

All behavioral experiments were done in awake, unrestrained, acclimatized, age-matched adult mice. Mice were habituated to the experimental setup twice a day for 4 days prior to

the behavioral testing. The experimenter was blinded for the genotype of the groups tested in the experiment. Thermal sensitivity was measured by recording paw withdrawal latency on application of infrared (IR) heat source on the plantar surface of hindpaw. A cut off of 20 sec was set to avoid burning of tissue. Two consecutive heat applications are performed at a time interval of 5 minutes. Von Frey measurements were done to assess mechanical sensitivity as described previously [20]. Briefly, mice were placed on an elevated grid and von Frey monofilaments exerting a specific force of 0.07, 0.16, 0.4, 0.6, 1.0, 1.4, 2.0 and 4.0 g were tested on the plantar hindpaw. Each monofilament was applied for 5 times at time interval of 5 min on the plantar hindpaw. 60% response frequency was as “thresholds” as described previously at basal, 14 and 24 wks post-STZ injection.

ROS measurements in DRG

To detect production of ROS and superoxide in DRG of SNSHIF1 $\alpha^{-/-}$ and HIF1 $\alpha^{fl/fl}$ diabetic and nondiabetic mice, we intrathecally (i.t) injected MitoTrackerRedCM-H₂XROS dye (100 nm, 10 μ l, Life Technologies). Twenty four h later, mice were perfused with 1x PBS and 4% PFA and DRG were extracted and 16 μ m sections were made on cryotome [21]. The dye is in reduced form and non-fluorescent in the basal state and becomes fluorescent upon oxidation [22]. Presence of MitoTrackerRedCM-H₂XROS fluorescence at different point of time point's post-STZ injection was analyzed using confocal microscopy.

Immunohistochemistry

Immunohistochemistry was performed on punch biopsies of the plantar surface of the hindpaws in diabetic and non-diabetic HIF1 $\alpha^{fl/fl}$ an SNS-HIF1 $\alpha^{-/-}$ mice prior to and different time points post-STZ. The mice were perfused with 4% paraformaldehyde (PFA) and plantar skin was post-fixed in 4 % PFA for 24 hrs at 4°C. After overnight incubation in 30% sucrose, 16 μ m cryo sections were made. The sections were permeabilized in 0.5% PBST, washed

and blocked with 7% Horse serum (HS). The sections were incubated overnight with anti-CGRP (1:1000, sigma) primary antibody in 7% HS in PBS at 4°C [23]. Subsequently, sections were washed and incubated in Alexa Fluor-594-conjugated secondary antibody. Further, the sections were washed and mounted in Mowiol. Fluorescence images were obtained using a laser-scanning spectral confocal microscope and maximal projections were created using Leica SP8 software (Leica TCS SP8 AOBS, Bensheim, Germany). Same conditions were used for imaging of sections from control and test groups. The acquired images were analyzed using ImageJ software. The fluorescence intensity from the epidermal area was measured and the background staining from IgG control was deducted from each test sample.

Statistical analysis

All the data were calculated and are presented as mean \pm SEM. ANOVA for repeated measures followed by Bonferroni's test for multiple comparisons was employed to determine statistically significant differences. Changes with $p \leq 0.05$ were considered to be significant.

RESULTS

In vivo modulation of nociception by HIF1 α in STZ model of type 1 diabetes

This study investigated the role of HIF1 α in the modulation of levels of ROS in dorsal root ganglia in diabetic and non-diabetic mice. In order to address a potential contribution of HIF1 α to DPN, we employed a mouse line with conditional genetic loss of HIF-1 α selectively in a sensory neuron-specific manner (SNS-HIF1 α ^{-/-} mice [18, 19]. In this line, we have previously demonstrated the deletion of HIF1 α in peripheral nociceptive neurons via quantitative RT-PCR and *in situ* hybridization [18]. The STZ model of type 1 is characterized by slowly progressive β cell death and subsequently lymphocytic infiltration of pancreatic

islets [26]. To investigate the role of HIF-1 α in diabetes-induced ROS modulation, we employed a protocol of STZ injections, which gradually induces hyperglycemia and precludes acute neurotoxic effects that may result from high doses of STZ [24, 25]. Using this protocol, hyperglycemia developed to comparable levels in SNS-HIF1 $\alpha^{-/-}$ and HIF1 $\alpha^{fl/fl}$ mice (Fig 1a). Behavior measurements in STZ-injected mice are an important and direct measure of the progression of DPN. The STZ protocol reproduces sensory phenomena of human DPN, i.e. hyperalgesia, which is evident over 5-8 weeks post-STZ, and progressive loss of sensory functions, which is evident starting 20 wks post-STZ injection [27, 28]. We performed long-term studies in SNS-HIF1 $\alpha^{-/-}$ and corresponding control mice (HIF1 $\alpha^{fl/fl}$) in the STZ model to determine the course of behavioral changes, setting a focus on 5, 13 and 24 wks post-STZ injection. Thus, the time points we chose are temporally separated from potential acute toxic effects of STZ.

At 5 wks post-STZ, we observed that both SNS-HIF1 $\alpha^{-/-}$ and HIF1 $\alpha^{fl/fl}$ develop similar levels of hypersensitivity to mechanical stimuli applied to plantar surface of paw in form of graded strength von Frey hairs, evident as a decrease in mechanical nociceptive thresholds (figure 1b, c). Similarly, the development of thermal hyperalgesia, which is evident as a decrease in latency to an infrared heat lamp applied to the hindpaw plantar surface (figure 1b, c). In contrast, at 13 and 24 wks post-STZ injection, deletion of HIF1 α in sensory neurons of the DRG had a marked impact on the development of sensory losses. Upon measuring mechanical response thresholds at 13 and 24 wks post-STZ, we observed that SNS-HIF1 $\alpha^{-/-}$ showed hypoalgesia as compared to HIF1 $\alpha^{fl/fl}$ mice, indicating an increase in the magnitude of STZ-induced sensory losses (figure 1b,*, # $p < 0.0001$, ANOVA, Bonferroni's test). Similarly, in the Hargreaves test, the time required for paw withdrawal upon stimulation with radiant heat was significantly increased in STZ-treated SNS-HIF1 $\alpha^{-/-}$ mice as compared to STZ-treated HIF1 $\alpha^{fl/fl}$ mice (figure 1c, * $p < 0.05$, *** $p < 0.0001$, # $p < 0.01$, ANOVA, Bonferroni's test). Hence, SNS-HIF1 $\alpha^{-/-}$ mice developed faster onset of hypoalgesia and thus showed a stronger loss of sensory function as compared to control mice, which is an indicative of an accelerated course of DPN upon loss of HIF1 α specifically in peripheral sensory neurons.

Manifestation of DPN in SNS-HIF1 α ^{-/-} mice post-STZ injection

In order to complement behavioral abnormalities with actual changes in nerve morphology, we conducted a detailed neuropathological analysis of changes in epidermal nerve fiber density. We performed immunostaining of markers for nociceptive nerves, such as Calcitonin Gene-Related Peptide (CGRP), on skin sections in a longitudinal analysis up to 24 wks post-STZ. We analyzed DPN-associated reductions in the density of epidermal nociceptive innervation, measured by the fluorescence intensity of CGRP-positive fibers in epidermal region of the glabrous skin derived from the paw. The magnitude of this loss was higher and the progression faster in SNS-HIF1 α ^{-/-} mice as compared to HIF1 α ^{fl/fl} mice (Fig. 2a, b, * p<0.0001, # p<0.01, ANOVA, Bonferroni's test), at 13 and 24 wks post-STZ injection, indicating an aggravation of DPN manifestations in the absence of HIF1 α .

Modulation of ROS levels in DRGs of diabetic mice by HIF1 α

Previous studies have established that hyperglycemia modulates the respiratory chain of the mitochondria and augments oxidative stress [29]. Overproduction of ROS has been shown to modulate HIF1 function [30], with both positive and negative modulation being reported, and several mechanisms have been proposed [31, 32, 16]. Despite these insights on the regulation of HIF1 by ROS, none of the studies investigated the impact of HIF1 function on modulation of ROS levels itself. Therefore, we measured the levels of ROS in vivo in DRG neurons of SNS-HIF1 α ^{-/-} and control HIF1 α ^{fl/fl} mice in the basal state (i.e. naïve mice) and at 13 or 24 wks post-STZ treatment. We intrathecally injected MitoTrackerRedCM-H₂XROS (100 nm, 10 μ l), a reporter for ROS generation [22], and observed that STZ-induced DPN was linked to a significant increase in MitoTrackerRedCM-H₂XROS fluorescence in DRG neurons, suggesting ROS and superoxide build-up (Fig. 3a, b, *, # p<0.0001, ANOVA, Bonferroni's test). Importantly, the extent of ROS and superoxide accumulation was

significantly increased in SNS-HIF1 α ^{-/-} mice as compared to HIF1 α ^{fl/fl} mice at 13 and 24 wks of STZ injection (Fig. 3a, b). Thus, a selective deletion of HIF1 α resulted in inflated levels of ROS formation in peripheral sensory neurons in vivo at time points, which temporally matched the exacerbation of DPN in SNS-HIF1 α ^{-/-} mice.

DISCUSSION

The main insights revealed by this study are: (i) the transcription factor HIF1 α plays a protective role against axonal pathology and loss of peripheral nerve fibers in DPN, (ii) a loss of HIF1 α in sensory neurons of diabetic mice leads to a faster progression of DPN-associated loss of sensation at the level of organismal behavior; (iii) a loss of HIF1 α results in an enhanced build-up of ROS in sensory neurons, thereby placing ROS downstream of HIF1 α .

HIF1 α has only recently been suggested to modulate pain following direct mechanical injury to peripheral nerves [18]. However, to date, nothing was known about the role of HIF1 α in pain and nerve damage caused by metabolic dysfunction. Indeed, there is a large mechanistic diversity between diverse forms of pathological pain, such as that arising from direct mechanical injury to nerves that is focal and localized versus metabolically-induced nerve dysfunction in the course of diabetes, which is widespread and diffuse. Very little is known about how nerve damage develops under metabolic stress. We hypothesized that HIF1 α may be a vital link because it has been reported to be directly regulated by hyperglycemia in some tissues, such as endothelial cells and cardiac myocytes, with both positive and negative modulation being reported [33]. Indeed, using highly selective genetic perturbations, we observed that both structural remodeling of nerves and their functionality (via behavioral analysis) were highly affected upon a loss of HIF1 α in sensory neurons. Interestingly, only chronic changes, such as structural damage and the ensuing sensory loss at about 4-6 months post-diabetes induction were affected in SNS-HIF1 α ^{-/-} mice,

emphasizing the strength of longitudinal analyses. Particularly this late component of diabetic changes is key in the clinical situation, frequently resulting in small and large fiber neuropathies, diabetic feet and amputations [34]. In contrast, early STZ-evoked hypersensitivity was not changed upon a loss of HIF1 α in sensory neurons. Several recent studies have addressed mechanisms of early hyperalgesia in diabetic models and there is converging evidence that this involves acute modulation of ion-channels, such as members of the Transient Receptor Potential (TRP) family, such as TRPV1 and TRPA1 by oxidative or lipid metabolites [35, 36].

The nature of modulation of DPN by HIF1 α is of much interest. In terms of injury induced by hypoxia and/or hyperglycemia, both protective and destructive roles have been observed for HIF1 α depending upon the type of tissue and the context [33, 37]. Here, we observed a protective function for HIF1 α against nerve damage induced by metabolic dysfunction in a model of type 1 diabetes. These observations were further strengthened by measurement of ROS levels in DRG, which revealed that the deletion of HIF1 α results in increased accumulation of ROS in sensory neurons as compared to controls over chronic phases of diabetes. This is unexpected, because previously published studies in other tissues reported the modulation of HIF1 α activity by ROS, but there is little evidence for modulation of ROS levels by HIF1 α . In kidney tissue under hyperglycemia conditions, ROS has been shown to impair HIF1 α activation as well as expression via inhibition of NO and Rac1 [33]. Moreover, ROS can further inhibit HIF1 α by increasing ubiquitin-proteasome activity [38]. In contrast, Bonello *et. al.* demonstrate that ROS leads to HIF α expression and activation in a NF κ B-dependent manner in pulmonary artery smooth muscle cells [39]. In sensory neurons of chronically hyperglycemic mice, we observed that ROS is downstream of HIF α , which suggests that HIF α exerts its neuroprotective functions in this context by suppressing ROS. The precise mechanisms how HIF α affects ROS levels are not clarified in this study. Contributing mechanisms may include the role of HIF1 α in modulating the expression of

metabolic enzymes [40], which may decrease the metabolic overload in sensory nerves and suppress the generation of harmful ROS and free radical species.

Recent studies have implicated ROS in the modulation of acute pain [41] and the administration of ROS scavengers peripherally or spinally has antinociceptive effects in models of inflammatory and neuropathic pain [42]. It is believed that ROS production via mitochondrial respiratory chain is a causal link between hyperglycemia and pathways involved in hyperglycemia-induced tissue damage. Under hyperglycemic conditions, the increased electron flux in the respiratory chain leads to an increase in the ATP/ADP ratio and in the mitochondrial membrane potential, which result in partial inhibition of complex III and thereby induce an accumulation of electrons to generate potentially harmful complex-I ROS [29, 43]. It is postulated that accumulation of ROS may result in a bioenergetic failure, depletion of antioxidant defenses and neuroinflammation [44]. These mechanisms likely contribute to nerve damage and the ensuing sensory loss observed upon the excessive accumulation of ROS in diabetic mice lacking HIF1 α .

CONCLUSIONS

In summary, this study identifies HIF1 α as an important protective molecule against DPN-associated long-term nerve damage and sensory loss. Moreover, we propose that HIF1 α functions as an upstream suppressor of ROS levels in sensory neurons. These observations have direct relevance to a highly debilitating and intractable long-term complication of diabetes and hold significance for novel therapeutic measures designed to strengthen the protective role of HIF1 α in sensory neurons over chronic diabetes.

AUTHORS CONTRIBUTIONS

NA and RK designed the study; IT mated HIF1 α ^{fl/fl} mice and SNS-Cre mice and gave general conceptual input; DRR performed and analyzed in vivo experiments; NA and DRR

performed and analyzed in vitro experiments; NA and RK generated the SNS-Cre line and wrote the manuscript. All authors approved the final version.

ACKNOWLEDGMENTS

The authors thank Rose LeFaucheur for secretarial help, Dunja Baumgartl-Ahlert and Hans-Joseph Wrede for technical assistance and Dr. Manuela Simonetti for help with i.t injections. This work was supported by a grant from the Deutsche Forschungsgemeinschaft (DFG) in the Collaborative Research Center 1118 (SFB1118 Project B06) to N.A. and R.K. We acknowledge support from the Interdisciplinary Neurobehavioral Core (INBC) for behavioral experiments.

COMPLIANCE WITH ETHICAL STANDARDS

All animal experiments were done in accordance with ethical guidelines and approved by Regierungspräsidium Karlsruhe, Germany. The manuscript does not contain clinical studies or patient data.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

REFERENCES

- 1 Mathers CD, Loncar D (2006) Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 3(11):e442.
- 2 Duby JJ, Campbell RK, Setter SM, White JR, Rasmussen KA (2004) Diabetic neuropathy: an intensive review. *Am J Health Syst Pharm.* 61: 160-73
- 3 Colloca L, Ludman T, Bouhassira D, Baron R, Dickenson AH, Yarnitsky D, et al (2017) Neuropathic pain. *Nat Rev Dis Primers* 3: 17002
- 4 Callaghan BC, Cheng HT, Stables CL, Smith AL, Feldman EL (2012) Diabetic neuropathy: clinical manifestations and current treatments. *Lancet Neurol* 11(6):521-34
- 5 Vincent AM, Callaghan BC, Smith AL, Feldman EL (2011) Diabetic neuropathy: cellular mechanisms as therapeutic targets. *Nat Rev Neurol.* 7(10):573-83.

- 6 Oates PJ (2002) Polyol pathway and diabetic peripheral neuropathy. *Int Rev Neurobiol.* 50:325-92
- 7 Du XL, Edelstein D, Rossetti L, Fantus IG, Goldberg H, Ziyadeh F et al (2000) Hyperglycemia-induced mitochondrial superoxide overproduction activates the hexosamine pathway and induces plasminogen activator inhibitor-1 expression by increasing Sp1 glycosylation. *Proc Natl Acad Sci U S A* 97(22):12222-6
- 8 Eichberg J (2002) Protein kinase C changes in diabetes: is the concept relevant to neuropathy? *Int Rev Neurobiol* 50:61-82.
- 9 Misur I, Zarković K, Barada A, Batelja L, Milicević Z, Turk Z (2004) Advanced glycation endproducts in peripheral nerve in type 2 diabetes with neuropathy. *Acta Diabetol* 41(4):158-66.
- 10 Miyauchi Y, Shikama H, Takasu T, Okamiya H, Umeda M, Hirasaki E, et al (1996) Slowing of peripheral motor nerve conduction was ameliorated by aminoguanidine in streptozocin-induced diabetic rats. *Eur J Endocrinol* 134(4):467-73
- 11 van der Vlies D, Makkinje M, Jansens A, Braakman I, Verkleij AJ, Wirtz KW, et al (2003) Oxidation of ER resident proteins upon oxidative stress: effects of altering cellular redox/antioxidant status and implications for protein maturation. *Antioxid Redox Signal* 5(4):381-7
- 12 Fiorentino TV, Prioleta A, Zuo P, Folli F (2013) Hyperglycemia-induced oxidative stress and its role in diabetes mellitus related cardiovascular diseases. *Curr Pharm Des* 19(32):5695-703.
- 13 Catrina SB, Okamoto K, Pereira T, Brismar K, Poellinger L (2004) Hyperglycemia regulates hypoxia-inducible factor-1alpha protein stability and function. *Diabetes.* 53(12):3226-32.
- 14 Masoud GN, Li W (2015) HIF-1 α pathway: role, regulation and intervention for cancer therapy. *Acta Pharm Sin B* 5(5):378-89.
- 15 Wang GL, Jiang BH, Rue EA, Semenza GL (1995) Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proc Natl Acad Sci U S A* 92(12):5510-4
- 16 Salceda S, Caro J (1997) Hypoxia-inducible factor 1alpha (HIF-1alpha) protein is rapidly degraded by the ubiquitin-proteasome system under normoxic conditions. Its stabilization by hypoxia depends on redox-induced changes. *J Biol Chem* 272(36):22642-7.
- 17 Iwai K, Yamanaka K, Kamura T, Minato N, Conaway RC, Conaway JW, et al (1999) Identification of the von Hippel-lindau tumor-suppressor protein as part of an active E3 ubiquitin ligase complex. *Proc Natl Acad Sci U S A* 96(22):12436-41.
- 18 Kanngiesser M, Mair N, Lim HY, Zschiebsch K, Blees J, Häussler A, et al (2014) Hypoxia-inducible factor 1 regulates heat and cold pain sensitivity and persistence. *Antioxid Redox Signal* 20(16):2555-71.
- 19 Agarwal N, Offermanns S, Kuner R (2004) Conditional gene deletion in primary nociceptive neurons of trigeminal ganglia and dorsal root ganglia. *Genesis* 38(3):122-9.

- 20 Simonetti M, Agarwal N, Stösser S, Bali KK, Karaulanov E, Kamble R, et al (2014) Wnt-Fzd signaling sensitizes peripheral sensory neurons via distinct noncanonical pathways. *Neuron* 83(1):104-21.
- 21 Guo BL, Sui BD, Wang XY, Wei YY, Huang J, Chen J, et al (2013) Significant changes in mitochondrial distribution in different pain models of mice. *Mitochondrion* 13(4):292-7.
- 22 Cottet-Rousselle C, Ronot X, Leverage X, Mayol JF (2011) Cytometric assessment of mitochondria using fluorescent probes. *Cytometry A* 79(6):405-25.
- 23 Hsieh YL, Lin CL, Chiang H, Fu YS, Lue JH, Hsieh ST (2012) Role of peptidergic nerve terminals in the skin: reversal of thermal sensation by calcitonin gene-related peptide in TRPV1-depleted neuropathy. *PLoS One*. 7(11):e50805.
- 24 Koulmanda M, Qipo A, Chebrolu S, O'Neil J, Auchincloss H, Smith RN (2003) The effect of low versus high dose of streptozotocin in cynomolgus monkeys (*Macaca fascicularis*). *Am J Transplant* 3: 267 - 72.
- 25 Yorek MA (2016) Alternatives to the Streptozotocin-Diabetic Rodent. *Int Rev Neurobiol*. 127:89-112.
- 26 Lenzen S (2008) The mechanisms of alloxan- and streptozotocin-induced diabetes. *Diabetologia* 51: 216-26
- 27 Korngut L, Ma CH, Martinez JA, Toth CC, Guo GF, Singh V, et al (2012) Overexpression of human HSP27 protects sensory neurons from diabetes. *Neurobiol Dis* 47(3):436-43.
- 28 Murakami T, Iwanaga T, Ogawa Y, Fujita Y, Sato E, Yoshitomi H, et al (2013) Development of sensory neuropathy in streptozotocin-induced diabetic mice. *Brain Behav*. 3:35-41
- 29 Brownlee M (2001) Biochemistry and molecular cell biology of diabetic complications. *Nature* 414(6865):813-20
- 30 Catrina SB, Okamoto K, Pereira T, Brismar K, Poellinger L (2004) Hyperglycemia regulates hypoxia-inducible factor-1 α protein stability and function. *Diabetes* 53(12):3226-32.
- 31 Bullock JJ, Mehta SL, Lin Y, Lolla P, Li PA (2009) Hyperglycemia-enhanced ischemic brain damage in mutant manganese SOD mice is associated with suppression of HIF-1 α . *Neurosci Lett* 456(2):89-92.
- 32 Hirota K, Semenza GL (2001) Rac1 activity is required for the activation of hypoxia-inducible factor 1. *J Biol Chem* 276(24):21166-72.
- 33 Xiao H, Gu Z, Wang G, Zhao T (2013) The possible mechanisms underlying the impairment of HIF-1 α pathway signaling in hyperglycemia and the beneficial effects of certain therapies. *Int J Med Sci*. 10(10):1412-21.
- 34 Yagihashi S, Mizukami H, Sugimoto K (2011) Mechanism of diabetic neuropathy: Where are we now and where to go? *J Diabetes Investig*. 2(1):18-32.
- 35 Pabbidi RM, Yu SQ, Peng S, Khardori R, Pauza ME, Premkumar LS (2008) Influence of TRPV1 on diabetes-induced alterations in thermal pain sensitivity. *Mol Pain*. 1;4:9.
- 36 Andersson DA, Gentry C, Light E, Vastani N, Vallortigara J, Bierhaus A, et al (2013) Methylglyoxal evokes pain by stimulating TRPA1. *PLoS One*. 8(10):e77986.

- 37 Bohuslavova R, Cerychova R, Nepomucka K, Pavlinkova G (2017) Renal injury is accelerated by global hypoxia-inducible factor 1 alpha deficiency in a mouse model of STZ-induced diabetes. *BMC Endocr Disord.* 17(1):48.
- 38 Botusan IR, Sunkari VG, Savu O, Catrina AI, Grünler J, Lindberg S, et al (2008) Stabilization of HIF-1alpha is critical to improve wound healing in diabetic mice. *Proc Natl Acad Sci U S A* 105(49):19426-31.
- 39 Bonello S, Zähringer C, BelAiba RS, Djordjevic T, Hess J, Michiels C, et al (2007) A. Reactive oxygen species activate the HIF-1alpha promoter via a functional NFkappaB site. *Arterioscler Thromb Vasc Biol* 27(4):755-61.
- 40 Dengler VL, Galbraith M, Espinosa JM (2014) Transcriptional regulation by hypoxia inducible factors. *Crit Rev Biochem Mol Biol* 49(1):1-15.
- 41 Kim HK, Par, SK, Zhou JL, Tagliatela G, Chung K, Coggeshall RE, et al (2004) Reactive oxygen species (ROS) play an important role in a rat model of neuropathic pain. *Pain* 111(1-2):116-24
- 42 Yowtak J, Lee KY, Kim HY, Wang J, Kim HK, Chung K, et al (2011) Reactive oxygen species contribute to neuropathic pain by reducing spinal GABA release. *Pain* 152(4):844-52.
- 43 Murphy MP (2009) How mitochondria produce reactive oxygen species. *Biochem J* 417(1):1-13
- 44 Shah MS, Brownlee M (2016) Molecular and Cellular Mechanisms of Cardiovascular Disorders in Diabetes. *Circ Res.* 118(11):1808-29.
- 45 Liu Z, Jia X, Duan Y, Xiao H, Sundqvist KG, Permert J, et al (2013) Excess glucose induces hypoxia-inducible factor-1 α in pancreatic cancer cells and stimulates glucose metabolism and cell migration. *Cancer Biol Ther.* 14(5):428-35.




Figure legends:

Figure 1: Analysis of hyperglycemia and sensory dysfunction in mice lacking HIF1 α selectively in peripheral nociceptive neurons (SNS-HIF1 $\alpha^{-/-}$) in the Streptozotocin (STZ) model of type 1 diabetes in a longitudinal study spanning 6 months. **(A)** Blood glucose levels in SNS-HIF1 $\alpha^{-/-}$ and HIF1 $\alpha^{fl/fl}$ mice after STZ injection **(B & C)** Course of DPN-associated sensory changes upon loss of HIF1 α in peripheral nociceptive neurons. **(B)** Summary of response thresholds to mechanical stimuli (defined as a force eliciting a response of paw withdrawal at least 40% response rate) and **(C)** thermal latency before and at 5, 13 and 24 days after STZ treatment (60 mg/kg for 5 days) in SNS-HIF1 $\alpha^{-/-}$ or HIF1 $\alpha^{fl/fl}$ (n = 10 mice per group). Data are represented as mean \pm S.E.M. * $p < 0.05$ compared to baseline, # $p < 0.05$

as compared to control group two-way ANOVA for repeated measurements with Bonferroni multiple comparison test.

Figure 2: Course and quantitative morphological analysis of DPN-associated pathology of peripheral nociceptive nerves in SNS-HIF1 α ^{-/-} mice and control littermates. **(A)** Identification of intra-epidermal nerve fibers via immunostaining for CGRP in the paw skin of SNS-HIF1 α ^{-/-} and HIF1 α ^{fl/fl} mice after STZ injection. The dotted white lines represent the epidermal layer in the skin. **(B)** Quantification of reduction in epidermal nerve fiber density post-STZ injection in SNS-HIF1 α ^{-/-} and HIF1 α ^{fl/fl} mice. n=6 - 8 mice per group. Data are represented as mean \pm S.E.M. * $p < 0.05$ compared to baseline, # $p < 0.05$ as compared to control group two-way ANOVA for repeated measurements with Bonferroni multiple comparison test. Scale bar represents 30 μ m.


Figure 3: Quantitative analysis of ROS measurement in vivo in peripheral sensory neurons at stages corresponding to chronic nerve damage and sensory dysfunction in diabetic mice. **(A)** Examples and **(B)** quantitative analysis of MitoTrackerRed staining in DRGs of STZ-injected SNS-HIF1 α ^{-/-} and HIF1 α ^{fl/fl} mice after intrathecal delivery of the ROS reporter dye. Arrows indicate ROS-positive neurons; n=6 DRGs per group. * $p < 0.05$ compared to baseline, # $p < 0.05$ as compared to control group two-way ANOVA for repeated measurements with Bonferroni multiple comparison test. Scale bar represents 30 μ m.




■ SNS-HIF1 $\alpha^{-/-}$ mice

■ HIF1 $\alpha^{fl/fl}$ mice


A Immunostaining using α -CGRP antibody



B



A Measurement of ROS in DRG using MitoTrackerRedCM-H₂XROS



B

