

# CROSSED MODULES OF MONOIDS II.

## RELATIVE CROSSED MODULES

GABRIELLA BÖHM

ABSTRACT. This is the second part of a series of three strongly related papers in which three equivalent structures are studied:

- internal categories in categories of monoids; defined in terms of pullbacks relative to a chosen class of spans
- crossed modules of monoids relative to this class of spans
- simplicial monoids of so-called Moore length 1 relative to this class of spans.

The most important examples of monoids that are covered are small categories (treated as monoids in categories of spans) and bimonoids in symmetric monoidal categories (regarded as monoids in categories of comonoids). In this second part we define relative crossed modules of monoids and prove their equivalence with the relative categories of Part I.

### INTRODUCTION

Since their appearance in [17], crossed modules of groups have been intensively studied and applied in various contexts; see e.g. the reviews [13, 14, 12] and the references in them. They admit several different descriptions: a *simplicial group whose Moore complex is concentrated in degrees 1 and 2* turns out to be the internal nerve of a *strict 2-group* and the Moore complex yields a *crossed module*. These constructions establish, in fact, equivalences between these three notions.

The first (to our knowledge) proof of the equivalence between crossed modules and strict 2-groups — that is, category objects in the category of groups — can be found in [5], where it is referred also to the unpublished proof [6]. Based on the fact that groups constitute a semi-Abelian category, another short and deeply conceptual proof is due to George Janelidze [10].

More recently, however, some results on, and certain applications of crossed modules of groups were extended to crossed modules of groupoids [4] and of Hopf algebras [1, 16, 11, 8, 7]. To these generalizations Janelidze's proof can not be applied directly. Our aim is therefore to develop a wider theory of crossed modules of monoids in more general monoidal categories which are not expected to have all pullbacks (not even along split epimorphisms). We have the above two main examples in mind:

- Categories of spans whose monoids are small categories, including groupoids in particular.
- Categories of comonoids in symmetric monoidal categories whose monoids are bimonoids including Hopf monoids in particular.

In the first part [2] of this series of papers we discussed classes of spans satisfying appropriate conditions; and relative pullbacks with respect to them. Assuming that such pullbacks exist — as they do in our key examples — we introduced a monoidal

---

*Date:* March 2018.

category with monoidal product provided by these pullbacks. We defined a relative (to the chosen class of spans) category as a monoid in this monoidal category. It is given by the usual data

$$B \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{i} \\ \xleftarrow{t} \end{array} A \xleftarrow{d} A \square_B A \quad (*)$$

where  $\square_B$  is now a relative pullback.

In the current article we make the next step and prove the equivalence of the following categories for a fixed class of suitable spans in a monoidal category:

- the category of relative categories in the category of monoids,
- the category of relative crossed modules of monoids.

Our methodology is inspired by Janelidze's paper [10]: In Section 1 we investigate first some category of split epimorphisms of monoids. We obtain an equivalent description of a split epimorphism of monoids  $B \begin{array}{c} \xrightarrow{i} \\ \xleftarrow{s} \end{array} A$  in terms of a distributive law which allows for handy characterizations of possible morphisms  $t$  and  $d$  in (\*). This is used in Section 2 and Section 3, respectively, to present equivalent descriptions of some reflexive graphs of monoids in terms of relative pre-crossed modules of monoids; and of relative category objects (\*) in categories of monoids in terms of relative crossed modules of monoids. Applying our results to categories of spans and to categories of comonoids, respectively, we re-obtain the definitions of crossed modules of groupoids in [4] and of crossed modules of Hopf monoids in [16], respectively.

Our next aim is to extend to our setting the equivalence of strict 2-groups and the category of crossed modules of groups to the further category of simplicial groups whose Moore complex has length 1. This will be achieved in Part III of this series [3].

**Acknowledgement.** The author's interest in the subject was triggered by the excellent workshop 'Modelling Topological Phases of Matter – TQFT, HQFT, premodular and higher categories, Yetter-Drinfeld and crossed modules in disguise' in Leeds UK, 5-8 July 2016. It is a pleasure to thank the organizers, Zoltán Kádár, João Faria Martins, Marcos Calçada and Paul Martin for the experience and a generous invitation. Financial support by the Hungarian National Research, Development and Innovation Office NKFIH (grant K124138) is gratefully acknowledged.

## 1. SPLIT EPIMORPHISMS OF MONOIDS VERSUS DISTRIBUTIVE LAWS

We freely use definitions, notation and results from [2]. Throughout, the composition of some morphisms  $A \xrightarrow{g} B$  and  $B \xrightarrow{f} C$  in an arbitrary category will be denoted by  $A \xrightarrow{f.g} C$ . Identity morphisms will be denoted by 1 (without any reference to the (co)domain object if it causes no confusion). In any monoidal category  $\mathbf{C}$  the monoidal product will be denoted by juxtaposition and the monoidal unit will be  $I$ . For the monoidal product of  $n$  copies of the same object  $A$  also the power notation  $A^n$  will be used. For any monoid  $A$  in  $\mathbf{C}$ , the multiplication and the unit morphisms will be denoted by  $A^2 \xrightarrow{m} A$  and  $I \xrightarrow{u} A$ , respectively. If  $\mathbf{C}$  is also braided, then for the braiding the symbol  $c$  will be used.

Recall that an *admissible class*  $\mathcal{S}$  of spans in an arbitrary category was defined in [2, Definition 2.1]. The *pullback*

$$\begin{array}{ccc} A \square_B C & \xrightarrow{p_C} & C \\ p_A \downarrow & & \downarrow g \\ A & \xrightarrow{f} & B \end{array}$$

of the cospan  $A \xrightarrow{f} B \xleftarrow{g} C$  relative to such a class  $\mathcal{S}$  was introduced in [2, Definition 3.1]. [2, Assumption 4.1] asserts that there exist the relative pullbacks of those cospans whose legs are in  $\mathcal{S}$  in the sense of [2, Definition 2.9]. Under this assumption it was proven in [2, Corollary 4.6] that the spans whose legs are in  $\mathcal{S}$  (again in the sense of [2, Definition 2.9]) constitute a monoidal category. An  $\mathcal{S}$ -relative category is defined as a monoid therein, see [2, Definition 4.9].

A class of spans in a monoidal category, which is compatible with the monoidal structure — meaning *multiplicativity* and *unitality* in a natural sense — was termed *monoidal* in [2, Definition 2.5]. It is discussed in [2, Example 2.8] that a monoidal admissible class  $\mathcal{S}$  of spans in a braided monoidal category  $\mathbf{C}$  induces a monoidal admissible class of spans in the category of monoids in  $\mathbf{C}$ ; and it is shown in [2, Example 4.4] that if  $\mathcal{S}$  satisfies [2, Assumption 4.1] then so does the induced class in the category of monoids. This allows for the discussion of relative categories in the category of monoids.

In this paper we will be interested mainly in these relative categories of monoids. They contain, in particular, a split epimorphism of monoids (consisting of the morphisms  $i$  and  $s$  of  $(*)$  in the Introduction). So we start with the analysis of the following category of split epimorphisms of monoids.

**Theorem 1.1.** *Consider a monoidal admissible class  $\mathcal{S}$  of spans in a monoidal category  $\mathbf{C}$  for which [2, Assumption 4.1] holds. The following categories are equivalent.*

$\text{SplitEpiMon}_{\mathcal{S}}(\mathbf{C})$  whose

objects are split epimorphisms  $B \begin{smallmatrix} \xrightarrow{s} \\ \xleftarrow{i} \end{smallmatrix} A$  of monoids in  $\mathbf{C}$  subject to the following conditions.

- (a)  $A \xrightarrow{s} B \in \mathcal{S}$ ; so that by the unitality of  $\mathcal{S}$  and [2, Assumption 4.1], there exists the  $\mathcal{S}$ -relative pullback

$$\begin{array}{ccc} A \square_B I & \xrightarrow{p_I} & I \\ p_A \downarrow & & \downarrow u \\ A & \xrightarrow{s} & B. \end{array}$$

- (b)  $q := (A \square_B I) B \xrightarrow{p_A i} A^2 \xrightarrow{m} A$  is invertible.

morphisms are pairs of monoid morphisms  $(B \xrightarrow{b} B', A \xrightarrow{a} A')$  such that  $s'.a = b.s$  and  $i'.b = a.i$ .

$\text{DistLaw}_{\mathcal{S}}(\mathcal{C})$  whose

objects consist of monoids  $B$  and  $Y$ , a monoid morphism  $Y \xrightarrow{e} I$  and a distributive law  $BY \xrightarrow{x} YB$  subject to the following conditions.

(a')  $Y = Y \xrightarrow{e} I \in \mathcal{S}$  and  $B = B = B \in \mathcal{S}$ . Then by the monoidality of  $\mathcal{S}$  also  $YB = Y \xrightarrow{e1} B \in \mathcal{S}$  so by [2, Assumption 4.1] there exists the  $\mathcal{S}$ -relative pullback  $YB \square_B I$  in the diagram below.

(b')  $e1.x = 1e$ .

(c') The morphism  $f$  occurring in the diagram below is invertible. (It is well-defined since by (a') and condition (POST) in [2, Definition 2.1],

$$YB \xleftarrow{1u} Y \xrightarrow{e} I \in \mathcal{S}.)$$

$$\begin{array}{ccccc} Y & & & & \\ & \searrow f & & & \\ & & YB \square_B I & \xrightarrow{pI} & I \\ & & \downarrow p_{YB} & & \downarrow u \\ & & YB & \xrightarrow{e1} & B. \end{array}$$

morphisms are pairs of monoid morphisms  $(B \xrightarrow{b} B', Y \xrightarrow{y} Y')$  such that  $e'.y = e$  and  $x'.by = yb.x$ .

*Proof.* We prove the theorem by constructing mutually inverse equivalence functors. The first one  $\text{SplitEpiMon}_{\mathcal{S}}(\mathcal{C}) \rightarrow \text{DistLaw}_{\mathcal{S}}(\mathcal{C})$  sends

$$\begin{array}{ccc} \begin{array}{ccc} B & \xrightleftharpoons[s]{i} & A \\ b \downarrow & & \downarrow a \\ B' & \xrightleftharpoons[s']{i'} & A' \end{array} & \mapsto & \begin{array}{ccc} (A \square_B I, & B, & A \square_B I \xrightarrow{pI} I, B(A \square_B I) \xrightarrow{ip_A} A^2 \xrightarrow{m} A \xrightarrow{q^{-1}} (A \square_B I)B \\ a \square 1 \downarrow & \downarrow b & \\ (A' \square_{B'} I, & B', & A' \square_{B'} I \xrightarrow{pI} I, B'(A' \square_{B'} I) \xrightarrow{i'p_{A'}} A'^2 \xrightarrow{m'} A' \xrightarrow{q'^{-1}} (A' \square_{B'} I)B'. \end{array} \end{array}$$

Let us see that the object map is meaningful. By construction  $B$  is a monoid and  $B \xrightarrow{b} B'$  is a monoid morphism. By [2, Proposition 3.7 (1)]  $A \square_B I$  is a monoid and  $A \square_B I \xrightarrow{pI} I$  is a monoid morphism. By [2, Lemma 1.5]  $B(A \square_B I) \xrightarrow{ip_A} A^2 \xrightarrow{m} A \xrightarrow{q^{-1}} (A \square_B I)B$  is a distributive law. Concerning property (a'),  $I = I = I \in \mathcal{S}$  by the unitality of  $\mathcal{S}$ ; hence by [2, Lemma 3.4 (2)]  $A \square_B I = A \square_B I \xrightarrow{pI} I \in \mathcal{S}$ . By [2, Lemma 2.4 (1)] also  $B = B = B$  belongs to  $\mathcal{S}$ . Condition (b') holds since commutativity of the first diagram of

$$\begin{array}{ccc} \begin{array}{ccc} & \xrightarrow{q} & \\ (A \square_B I)B & \xrightarrow{p_A i} & A^2 \xrightarrow{m} A \\ pI1 \downarrow & & \downarrow ss \\ B & \xrightarrow{u1} & B^2 \xrightarrow{m} B \end{array} & & \begin{array}{ccc} B(A \square_B I) & \xrightarrow{ip_A} & A^2 \xrightarrow{m} A \xrightarrow{q^{-1}} (A \square_B I)B \\ 1pI \downarrow & & \downarrow ss \\ B & \xrightarrow{1u} & B^2 \xrightarrow{m} B \end{array} \end{array} \quad (1.1)$$

implies the commutativity of the second diagram. For condition (c') observe that by the unitality of the monoid morphism  $i$  the equality  $q.1u = p_A$  holds, equivalently,  $q^{-1}.p_A = 1u$ . With this identity in mind we see that the morphism  $f$  of condition (c') is equal to  $q^{-1} \square 1$  in the first diagram of

$$\begin{array}{ccc}
 \begin{array}{ccccc}
 A \square_I & & & & \\
 \downarrow p_A & \searrow^{q^{-1} \square 1} & & \searrow^{p_I} & \\
 (A \square_I) B \square_I & & & & I \\
 \downarrow p_{(A \square_I) B} & & & & \downarrow u \\
 A & \xrightarrow{q^{-1}} & (A \square_I) B & \xrightarrow{p_I 1} & B
 \end{array} & & 
 \begin{array}{ccccc}
 (A \square_I) B \square_I & & & & \\
 \downarrow p_{(A \square_I) B} & \searrow^{q \square 1} & & \searrow^{p_I} & \\
 (A \square_I) B & & & & I \\
 \downarrow p_A & & & & \downarrow u \\
 (A \square_I) B & \xrightarrow{q} & A & \xrightarrow{s} & B
 \end{array}
 \end{array}$$

Then by [2, Proposition 3.5 (2)] it is invertible with the inverse  $q \square 1$  in the second diagram. Both morphisms  $q^{-1} \square 1$  and  $q \square 1$  are well-defined by the commutativity of the first diagram of (1.1); see [2, Proposition 3.5 (1)]. This proves that the object map of our candidate functor is meaningful.

Concerning the morphism map,  $a \square 1$  is a well-defined morphism in  $\mathcal{C}$  by the assumption that  $b.s = s'.a$  (see [2, Proposition 3.5 (1)]) and it is a monoid morphism by [2, Proposition 3.7 (2)]. Condition  $p_I.(a \square 1) = p_I$  holds by construction and the other equality holds since the commutativity of the first diagram of

$$\begin{array}{ccc}
 \begin{array}{ccc}
 (A \square_I) B & \xrightarrow{p_A i} & A^2 \\
 \downarrow (a \square 1) b & & \downarrow aa \\
 (A' \square_I) B' & \xrightarrow{p_{A'} i'} & A'^2 \\
 & & \downarrow m' \\
 & & A'
 \end{array} & & 
 \begin{array}{ccc}
 B(A \square_I) & \xrightarrow{ip_A} & A^2 \\
 \downarrow b(a \square 1) & & \downarrow aa \\
 B'(A' \square_I) & \xrightarrow{i' p_{A'}} & A'^2 \\
 & & \downarrow m' \\
 & & A'
 \end{array}
 \end{array} \quad (1.2)$$

implies the commutativity of the second diagram.

In the opposite direction  $\text{DistLaw}_{\mathcal{S}}(\mathcal{C}) \rightarrow \text{SplitEpiMon}_{\mathcal{S}}(\mathcal{C})$  we propose a functor sending

$$\begin{array}{ccc}
 (Y, B, Y \xrightarrow{e} I, BY \xrightarrow{x} YB) & & B \xrightleftharpoons[u1]{e1} YB \\
 y \downarrow & \downarrow b & \downarrow b \\
 (Y', B', Y' \xrightarrow{e'} I, B'Y' \xrightarrow{x'} Y'B') & \mapsto & B' \xrightleftharpoons[u'1]{e'1} A.
 \end{array}$$

Here  $YB$  is considered with the monoid structure induced by the distributive law  $x$ , see [2, Lemma 1.4]. Then  $B \xrightarrow{u1} YB$  is a monoid morphism by [2, Lemma 1.4] again. By [2, Lemma 1.6] condition (b') implies that  $YB \xrightarrow{e1} B$  is a monoid morphism too. The rows are split epimorphisms (of monoids) by the unitality of the monoid morphism  $e$ . By (a') and the multiplicativity of  $\mathcal{S}$ ,  $YB = YB \xrightarrow{e1} B \in \mathcal{S}$  so that condition (a)

holds. For condition (b) note that the commutativity of

$$\begin{array}{ccccc}
 YB & \xlongequal{\quad} & & \xlongequal{\quad} & YB \\
 \downarrow f1 & \searrow 1u1 & & \nearrow 1m & \parallel \\
 & YB^2 & \xlongequal{\quad} & YB^2 & \\
 & \downarrow 11u1 & & \downarrow 1u11 & \\
 (YB \square_I B)_B & \xrightarrow{p_{YB}u1} & (YB)^2 & \xrightarrow{1x1} & Y^2B^2 & \xrightarrow{mm} & YB
 \end{array}$$

implies that the bottom row is the inverse of the isomorphism  $f1$  in the left column hence it is invertible. This proves that the object map is well defined.

Concerning the morphism map, it follows by the assumption  $yb.x = x'.by$  that  $yb$  is a monoid morphism, see [2, Lemma 1.6]. The monoid morphisms  $(b, yb)$  are compatible with the monomorphisms  $B \xrightarrow{u1} YB$  and  $B' \xrightarrow{u'1} Y'B'$  by the unitality of  $y$  and they are compatible with the epimorphisms  $YB \xrightarrow{e1} B$  and  $Y'B' \xrightarrow{e'1} B'$  by the assumption that  $e'.y = e$ .

So we have well-defined functors in both directions, it remains to see that their composites are naturally isomorphic to the identity functors. The composite

$$\text{SplitEpiMon}_{\mathcal{S}}(\mathcal{C}) \rightarrow \text{DistLaw}_{\mathcal{S}}(\mathcal{C}) \rightarrow \text{SplitEpiMon}_{\mathcal{S}}(\mathcal{C})$$

acts as

$$\begin{array}{ccc}
 \begin{array}{ccc} B & \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{i} \end{array} & A \\ b \downarrow & & \downarrow a \\ B' & \begin{array}{c} \xleftarrow{s'} \\ \xrightarrow{i'} \end{array} & A \end{array} & \mapsto & \begin{array}{ccc} B & \begin{array}{c} \xleftarrow{p_I 1} \\ \xrightarrow{u1} \end{array} & (A \square_I B)_B \\ b \downarrow & & \downarrow a \\ B' & \begin{array}{c} \xleftarrow{p_{I'} 1} \\ \xrightarrow{u'1} \end{array} & (A' \square_I B')_{B'} \end{array}
 \end{array}$$

We claim that a natural isomorphism from this to the identity functor has the components  $(B \xrightarrow{q} (A \square_I B)_B, (A \square_I B)_B \xrightarrow{q} A)$ . Since  $p_A$  is a monoid morphism by [2, Proposition 3.7 (1)], so is  $q$  by [2, Lemma 1.5]. The stated pair  $(1, q)$  is a morphism in  $\text{SplitEpiMon}_{\mathcal{S}}(\mathcal{C})$  by the first diagram of (1.1) and by the fact that the unitality of  $p_A$  implies  $q.1u = i$ . Naturality with respect to any morphism  $(B \xrightarrow{b} B', A \xrightarrow{a} A')$  in  $\text{SplitEpiMon}_{\mathcal{S}}(\mathcal{C})$  follows by the commutativity of the first diagram of (1.2).

Composing our functors in the opposite order

$$\text{DistLaw}_{\mathcal{S}}(\mathcal{C}) \rightarrow \text{SplitEpiMon}_{\mathcal{S}}(\mathcal{C}) \rightarrow \text{DistLaw}_{\mathcal{S}}(\mathcal{C})$$

we obtain the functor sending

$$\begin{array}{ccc}
 (Y, & B, & Y \xrightarrow{e} I, BY \xrightarrow{x} YB) \\
 y \downarrow & \downarrow b & \\
 (Y', & B', & Y' \xrightarrow{e'} I, B'Y' \xrightarrow{x'} Y'B')
 \end{array}$$

to

$$\begin{array}{c}
 (YB \square_B I, B, YB \square_B I \xrightarrow{p_I} I, B(YB \square_B I) \xrightarrow{u^1 p_{YB}} (YB)^2 \xrightarrow{1x^1} Y^2 B^2 \xrightarrow{m'm} YB \xrightarrow{f^1} (YB \square_B I)B) \\
 \begin{array}{c}
 \downarrow y^{b \square 1} \\
 \downarrow b
 \end{array} \\
 (Y' B' \square_{B'} I, B', Y' B' \square_{B'} I \xrightarrow{p_I} I, B'(Y' B' \square_{B'} I) \xrightarrow{u^1 p_{Y'B'}} (Y' B')^2 \xrightarrow{1x'^1} Y'^2 B'^2 \xrightarrow{m'm'} Y' B' \xrightarrow{f'^1} (Y' B' \square_{B'} I)B').
 \end{array}$$

We claim that a natural isomorphism from this to the identity functor has the invertible components  $(B = B, Y \xrightarrow{f} YB \square_B I)$ . By construction  $f$  is a monoid morphism, see [2, Proposition 3.7 (2)]. The compatibility of the monoid morphisms  $(1, f)$  with  $Y \xrightarrow{e} I$  and  $YB \square_B I \xrightarrow{p_I} I$  holds by the definition of  $f$  and the compatibility with the distributive laws  $BY \rightarrow YB$  and  $B(YB \square_B I) \rightarrow (YB \square_B I)B$  holds by the commutativity of

$$\begin{array}{c}
 \begin{array}{ccccccc}
 BY & \xlongequal{\quad} & BY & \xrightarrow{x} & YB & & \\
 \downarrow 1f & & \downarrow 11u & & \swarrow 11u & \parallel & \searrow f1 \\
 B(YB \square_B I) & \xrightarrow{1p_{YB}} & BYB & \xrightarrow{x1} & YB^2 & \xrightarrow{1m} & YB \xrightarrow{f1} (YB \square_B I)B.
 \end{array}
 \end{array}$$

Finally, the naturality with respect to an arbitrary morphism  $(B \xrightarrow{b} B', Y \xrightarrow{y} Y')$  in  $\text{DistLaws}_{\mathcal{S}}(\mathcal{C})$  follows by the commutativity of the diagrams

$$\begin{array}{ccc}
 \begin{array}{ccc}
 YB \square_B I & \xrightarrow{y^{b \square 1}} & Y' B' \square_{B'} I \\
 \downarrow p_{YB} & & \downarrow p_{Y'B'} \\
 Y & \xrightarrow{1u} & YB \xrightarrow{yb} & Y' B' \\
 \downarrow y & \nearrow 1u' & \uparrow p_{Y'B'} \\
 Y' & \xrightarrow{f'} & Y' B' \square_{B'} I
 \end{array} & & 
 \begin{array}{ccc}
 YB \square_B I & \xrightarrow{y^{b \square 1}} & Y' B' \square_{B'} I \\
 \downarrow p_I & & \downarrow p_I \\
 Y & \xrightarrow{e} & I \\
 \downarrow y & \nearrow e' & \uparrow p_I \\
 Y' & \xrightarrow{f'} & Y' B' \square_{B'} I
 \end{array}
 \end{array}$$

using again that  $YB \xleftarrow{p_{YB}} YB \square_B I \xrightarrow{p_I} I$  are joint monomorphisms in  $\mathcal{C}$ .  $\square$

**Example 1.2.** For any fixed set  $X$ , the category  $\mathcal{C}$  of spans over  $X$  is monoidal via the pullback over  $X$ . A monoid in  $\mathcal{C}$  is a small category with the object set  $X$  and a monoid morphism is a functor acting on the objects as the identity map. Moreover,  $\mathcal{C}$  has all pullbacks (computed in the underlying category of sets). So taking as  $\mathcal{S}$  the class of all spans in  $\mathcal{C}$ , from Theorem 1.1 we obtain the equivalence of the following categories. (Throughout  $s$  denotes the source map in any category and  $t$  denotes the target map.)

$\text{SplitEpiMon}(\mathcal{C})$  whose

objects are pairs of identity-on-objects functors  $B \xrightleftharpoons[\iota]{\sigma} A$  between categories of the common object set  $X$  such that the composite  $\sigma \iota$  is the identity functor, and the map

$$q : (A \square_B X) \square_X B = \{(a, x, b) \mid \sigma(a) = 1_x, x = t(b)\} \rightarrow A \quad (a, x, b) \mapsto a.\iota(b) \quad (1.3)$$

is invertible. (The morphism of (1.3) is invertible e.g. if  $B$  is a groupoid; then its inverse takes a morphism  $a$  to  $(a.\iota(\sigma(a)^{-1}), t(a), \sigma(a))$ .)

morphisms are pairs of identity-on-objects functors  $(A \xrightarrow{\alpha} A', B \xrightarrow{\beta} B')$  for which  $\alpha\iota = \iota'\beta$  and  $\beta\sigma = \sigma'\alpha$ .

**DistLaw(C)** whose

objects consist of categories  $B$  and  $Y$  with the common object set  $X$  such that  $\overline{Y}$  has no morphisms between non-equal objects (that is, its source map  $s$  and target map  $t$  coincide); and an *action*  $B \square_X Y = \{(b, y) | s(b) = t(y)\} \xrightarrow{\triangleright} Y$  in the sense of [4, Definition 1.1]; meaning the following axioms for all morphisms  $b, b'$  in  $B$  and  $y, y'$  in  $Y$  for which  $s(b') = t(b)$  and  $s(b) = t(y) = s(y) = t(y') = s(y')$ .

- (i)  $t(b \triangleright y) = t(b)$
- (ii)  $b \triangleright (y.y') = (b \triangleright y).(b \triangleright y')$  and  $b \triangleright 1_{s(b)} = 1_{t(b)}$
- (iii)  $(b'.b) \triangleright y = b' \triangleright (b \triangleright y)$  and  $1_{t(y)} \triangleright y = y$ .

morphisms are pairs of identity-on-objects functors  $(Y \xrightarrow{\nu} Y', B \xrightarrow{\beta} B')$  for which  $\nu(b \triangleright y) = \beta(b) \triangleright \nu(y)$  for all morphisms  $b$  in  $B$  and  $y$  in  $Y$  for which  $s(b) = t(y)$ .

Only the above description of an object in **DistLaw(C)** requires some explanation.

The monoidal unit of  $\mathbf{C}$  is the trivial span  $X = X = X$ . Its trivial monoid structure yields the discrete category  $\mathbf{D}(X)$ . An identity-on-objects functor  $Y \xrightarrow{e} \mathbf{D}(X)$  as in Theorem 1.1 exists if and only if the source and target maps of  $Y$  coincide. Then there is precisely one such functor sending any morphism to the identity morphism on its equal source and target objects. For this functor  $e$ , precisely those maps  $B \square_X Y \xrightarrow{x} Y \square_X B$  satisfy  $(e \square 1).x = 1 \square e$  which are of the form  $(b, y) \mapsto (b \triangleright y, b)$  in terms of some map  $\triangleright$  obeying condition (i). It is straightforward to see that then  $x$  is a distributive law if and only if conditions (ii) and (iii) hold.

The morphism  $f$  of Theorem 1.1 (c') is invertible because

$$\begin{array}{ccc} Y \square_X C & \xrightarrow{e \square 1} & C \\ 1 \square g \downarrow & & \downarrow g \\ Y \square_X B & \xrightarrow{e \square 1} & B \end{array} \quad (1.4)$$

is clearly a pullback of  $X$ -spans for any span morphism  $g$ .

**Example 1.3.** Let  $\mathbf{M}$  be a symmetric monoidal category in which equalizers exist and are preserved by taking the monoidal product with any object.

Take  $\mathbf{C}$  to be the category of comonoids in  $\mathbf{M}$  with the monoidal admissible class  $\mathcal{S}$  in [2, Example 2.3] of spans in  $\mathbf{C}$ . Thanks to the symmetry of  $\mathbf{M}$ , its monoidal structure is inherited by  $\mathbf{C}$ . A monoid  $A$  in  $\mathbf{C}$  is known as a *bimonoid* in  $\mathbf{M}$ . Recall that the monoidal structure of  $\mathbf{M}$  is lifted to the category of (left or right) modules over the monoid  $A$  in  $\mathbf{M}$ . A monoid (respectively, a comonoid) in the category of  $A$ -modules is known as an *A-module monoid* (respectively, *A-module comonoid*).



Recall from [2, Example 3.3] that for a cospan  $A \xrightarrow{f} B \xleftarrow{g} C$  of comonoids whose legs are in  $\mathcal{S}$ , the  $\mathcal{S}$ -relative pullback is given by the so-called *cotensor product*, defined as the equalizer

$$A \square_B C \xrightarrow{j} AC \xrightleftharpoons[1g1.1\delta]{1f1.\delta1} ABC \quad (1.5)$$

in  $\mathbf{M}$  (where  $\delta$  denotes both comultiplications of the comonoids  $A$  and  $C$ .)

Below we describe the equivalent categories of Theorem 1.1 in this context.

**SplitEpiMon $_{\mathcal{S}}$ (C)** whose

objects are split epimorphisms  $B \xrightleftharpoons[i]{s} A$  of bimonoids in  $\mathbf{M}$  subject to the following conditions.

(a) The comultiplication  $\delta$  of  $A$  satisfies  $c.s1.\delta = 1s.\delta$ .

(b) In terms of the morphism  $j$  of (1.5),  $q := (A \square_B I)B \xrightarrow{ji} A^2 \xrightarrow{m} A$  is invertible.

morphisms are pairs of bimonoid morphisms which are compatible with the epimorphisms  $s$  as well as their sections  $i$ .

**DistLaw $_{\mathcal{S}}$ (C)** whose

objects consist of a cocommutative bimonoid  $B$  and a bimonoid  $Y$  in  $\mathbf{M}$ , together with a left  $B$ -action on  $Y$  which makes  $Y$  both a left  $B$ -module monoid and a left  $B$ -module comonoid.

morphisms are pairs of bimonoid morphisms  $(B \xrightarrow{b} B', Y \xrightarrow{y} Y')$  which are compatible with the actions  $BY \xrightarrow{l} Y$  and  $B'Y' \xrightarrow{l'} Y'$  in the sense that  $l'.by = y.l$ .

This concise description of  $\text{DistLaw}_{\mathcal{S}}(\mathbf{C})$  requires a proof. Note that the monoidal unit  $I$  is now a terminal object in  $\mathbf{C}$ ; the unique morphism  $Y \rightarrow I$  is the counit  $\varepsilon$ . It obviously satisfies  $Y = Y \xrightarrow{\varepsilon} I \in \mathcal{S}$ . The other condition  $B = B = B \in \mathcal{S}$  in (a') of Theorem 1.1 reduces to the requirement that the comonoid  $B$  is cocommutative.

Next we establish a bijective correspondence between distributive laws  $BY \rightarrow YB$  satisfying property (b') of Theorem 1.1 and left actions  $BY \rightarrow Y$  as in the description above. Starting with a distributive law  $BY \xrightarrow{x} YB$ , put  $l := 1\varepsilon.x$ . It is a unital action by the left unitality of  $x$  and it is associative by the left multiplicativity of  $x$ :

$$\begin{array}{ccc}
 Y & \xlongequal{\quad} & Y \\
 \downarrow u1 & & \downarrow 1u \\
 BY & \xrightarrow{x} & YB \xrightarrow{1\varepsilon} Y \\
 & \searrow l & \nearrow
 \end{array}
 \qquad
 \begin{array}{ccccc}
 & & 1l & & \\
 & & \curvearrowright & & \\
 B^2Y & \xrightarrow{1x} & BYB & \xrightarrow{11\varepsilon} & BY \\
 \downarrow m1 & & \downarrow x1 & & \downarrow x \\
 & & YB^2 & \xrightarrow{11\varepsilon} & YB \\
 & & \downarrow 1m & & \downarrow 1\varepsilon \\
 BY & \xrightarrow{x} & YB & \xrightarrow{1\varepsilon} & Y \\
 & \searrow l & \nearrow & & 
 \end{array}$$

By the right unitality of  $x$  the unit  $I \xrightarrow{u} Y$  is a morphism of  $B$ -modules and by the right multiplicativity of  $x$  the multiplication  $Y^2 \xrightarrow{m} Y$  is a morphism of  $B$ -modules:

$$\begin{array}{ccc}
 \begin{array}{ccc}
 B & \xrightarrow{1u} & BY \\
 \parallel & & \parallel \\
 B & \xrightarrow{u1} & YB \\
 \parallel & & \parallel \\
 I & \xrightarrow{u} & Y
 \end{array} & & \begin{array}{ccc}
 BY^2 & \xrightarrow{\quad} & BY^2 & \xrightarrow{1m} & BY \\
 \delta_{11} \downarrow & & \downarrow & & \downarrow \\
 B^2Y^2 & \xrightarrow{11\delta_1} & B^2Y^3 & & BY \\
 \parallel & \swarrow & \downarrow & & \downarrow \\
 B^2Y^2 & & B^2Y^2 & & YBY \\
 \downarrow & \swarrow & \downarrow & & \downarrow \\
 (BY)^2 & \xleftarrow{111\varepsilon_1} & (BY)^2Y & & YBY \\
 \downarrow & \swarrow & \downarrow & & \downarrow \\
 YB^2Y & \xleftarrow{11\varepsilon_{11}} & (YB)^2Y & & Y^2B^2Y \\
 \downarrow & \swarrow & \downarrow & & \downarrow \\
 (YB)^2 & & (YB)^2 & & Y^2B \\
 \downarrow & & \downarrow & & \downarrow \\
 Y^2 & \xrightarrow{\quad} & Y^2 & \xrightarrow{m} & Y
 \end{array}
 \end{array}$$


(note that here we also used the comultiplicativity of  $x$ ). The condition that the counit  $Y \xrightarrow{\varepsilon} I$  is a morphism of  $B$ -modules coincides with the counitality of  $l$  and also with the counitality of  $x$ . The comultiplication  $Y \xrightarrow{\delta} Y^2$  is a morphism of  $B$ -modules, equivalently,  $l$  is comultiplicative by the comultiplicativity of  $x$ :

$$\begin{array}{ccccccc}
 BY & \xrightarrow{1\delta} & BY^2 & \xrightarrow{\delta_1} & B^2Y^2 & \xrightarrow{1c_1} & (BY)^2 \\
 \downarrow x & & \downarrow & & \downarrow & & \downarrow \\
 YB & \xrightarrow{\delta_1} & Y^2B & \xrightarrow{11\delta} & Y^2B^2 & \xrightarrow{1c_1} & (YB)^2 \\
 \downarrow 1\varepsilon & & \downarrow 11\varepsilon & & \downarrow 11\varepsilon\varepsilon & & \downarrow 1\varepsilon_1\varepsilon \\
 Y & \xrightarrow{\delta} & Y^2 & \xrightarrow{\quad} & Y^2 & \xrightarrow{\quad} & Y^2
 \end{array}$$


Conversely, in terms of an action  $l$  as above, put  $x := BY \xrightarrow{\delta_1} B^2Y \xrightarrow{1c} BYB \xrightarrow{l_1} YB$ . It clearly satisfies (b') by the counitality of  $l$  hence it is counital. It is comultiplicative by the comultiplicativity of  $l$ :

$$\begin{array}{ccccccc}
 BY & \xrightarrow{\quad} & B^2Y & \xrightarrow{\quad} & BYB & \xrightarrow{l_1} & YB \\
 \delta\delta \downarrow & & \downarrow \delta\delta\delta & & \downarrow \delta\delta\delta & & \downarrow \delta\delta \\
 B^2Y^2 & \xrightarrow{\delta\delta_{11}} & B^4Y^2 & \xrightarrow{1c_{111}} & B^4Y^2 & \xrightarrow{11c_{B^2,Y^2}} & B^2Y^2B^2 \\
 \downarrow 1c_1 & & \downarrow 11c_{B^2,Y^1} & & \downarrow 1c_{111} & & \downarrow \\
 (BY)^2 & \xrightarrow{\delta_1\delta_1} & (B^2Y)^2 & \xrightarrow{1c_1c} & (BYB)^2 & \xrightarrow{l_1l_1} & (YB)^2
 \end{array}$$

where the top-left region commutes by the coassociativity and cocommutativity of the comonoid  $B$ . This morphism  $x$  is a distributive law. Indeed, the left unitality and the left multiplicativity follow by the unitality and the associativity of the action  $l$ , respectively:



and the right unitality and the right multiplicativity of  $x$  follow using that the unit and the multiplication of  $Y$  are  $B$ -module morphisms:



The above correspondences between  $l$  and  $x$  are bijective by the commutativity of

$$\begin{array}{ccc}
 & B^2Y & \xrightarrow{1c} & BYB \\
 \delta_1 \curvearrowright & \uparrow 111\varepsilon & \nearrow 111\varepsilon & \downarrow x_1 \\
 BY & \xrightarrow{\delta\delta} & B^2Y^2 \xrightarrow{1c1} & (BY)^2 \\
 \downarrow x & & \downarrow xx & \downarrow 1\varepsilon 1 \\
 YB & \xrightarrow{\delta\delta} & Y^2B^2 \xrightarrow{1c1} & (YB)^2 \\
 & \downarrow 1\varepsilon\varepsilon 1 & \searrow 1\varepsilon\varepsilon 1 & \downarrow 1\varepsilon 1 \\
 & YB & \xlongequal{\quad} & YB
 \end{array}
 \qquad
 \begin{array}{ccc}
 BY & \xrightarrow{\delta_1} & B^2Y \\
 \downarrow 1c & & \downarrow 1\varepsilon 1 \\
 BYB & \xrightarrow{11\varepsilon} & BY \\
 \downarrow l_1 & & \downarrow l \\
 YB & \xrightarrow{1\varepsilon} & Y
 \end{array}$$

for a comultiplicative morphism  $x$  satisfying (b') and any morphism  $l$ .

Finally, we show that the morphism  $Y \xrightarrow{f} YB \square_B I$  in part (c') of Theorem 1.1 is invertible without any further assumption; its inverse is constructed as  $f^{-1} := YB \square_B I \xrightarrow{p_{YB}} YB \xrightarrow{1\varepsilon} Y$ . In order to see that it is the inverse, indeed, recall that by [2, Example 3.3] the morphism  $p_{YB}$  is the equalizer of  $YB \xrightarrow{1\delta} YB^2$  and  $YB \xrightarrow{11u} YB^2$ . Hence the following diagrams commute.

$$\begin{array}{ccc}
 Y & \xlongequal{\quad} & Y \\
 \downarrow f & & \downarrow 1u \\
 YB \square_B I & \xrightarrow{p_{YB}} & YB \xrightarrow{1\varepsilon} Y \\
 & \searrow f^{-1} & \nearrow f^{-1}
 \end{array}
 \qquad
 \begin{array}{ccccc}
 & & f^{-1} & & \\
 & & \curvearrowright & & \\
 YB \square_B I & \xrightarrow{p_{YB}} & YB & \xrightarrow{1\varepsilon} & Y \xrightarrow{f} & YB \square_B I \\
 p_{YB} \downarrow & & \downarrow 11u & & \downarrow 1u & \downarrow p_{YB} \\
 YB & \xrightarrow{1\delta} & YB^2 & \xrightarrow{1\varepsilon 1} & YB \xlongequal{\quad} & YB
 \end{array}$$

This completes the characterization of the objects of  $\text{DistLaw}_{\mathcal{S}}(\mathcal{C})$ . Concerning the morphisms  $(B \xrightarrow{b} B', Y \xrightarrow{y} Y')$ , the first condition in Theorem 1.1 is the counitality of the bimonoid morphism  $y$  hence it identically holds. The second condition in Theorem 1.1 is equivalent to  $y.l = l'.by$  by the commutativity of

$$\begin{array}{ccc}
 & \xrightarrow{x} & \\
 BY & \xrightarrow{\delta_1} & B^2Y \xrightarrow{1c} & BYB \xrightarrow{l_1} & YB \\
 by \downarrow & & \downarrow bby & \downarrow byb & \downarrow yb \\
 B'Y' & \xrightarrow{\delta'_1} & B'^2Y' \xrightarrow{1c} & B'Y'B' \xrightarrow{l'_1} & Y'B' \\
 & \searrow x' & \nearrow x' & & \\
 & & & & l'
 \end{array}
 \qquad
 \begin{array}{ccc}
 & \xrightarrow{l} & \\
 BY & \xrightarrow{x} & YB \xrightarrow{1\varepsilon} & Y \\
 by \downarrow & & \downarrow yb & \downarrow y \\
 B'Y' & \xrightarrow{x'} & Y'B' \xrightarrow{1\varepsilon'} & Y' \\
 & \searrow l' & \nearrow l' & \\
 & & & & l'
 \end{array}$$

We can apply the current example to the particular case of a finitely complete category  $\mathcal{M}$  regarded with the Cartesian monoidal structure. Then the category  $\mathcal{C}$  of comonoids in  $\mathcal{M}$  is isomorphic to  $\mathcal{M}$  and the equivalent categories of Theorem 1.1 reduce to the following ones.

$\text{SplitEpiMon}_{\mathcal{S}}(\mathbf{M})$  whose

objects are split epimorphisms  $B \begin{smallmatrix} \xrightarrow{s} \\ \xleftarrow{i} \end{smallmatrix} A$  of monoids in  $\mathbf{M}$  such that in terms

of the morphism  $j$  of (1.5),  $q := (A \square_B I)B \xrightarrow{j^i} A^2 \xrightarrow{m} A$  is invertible.

morphisms are pairs of monoid morphisms which are compatible with the epimorphisms  $s$  as well as their sections  $i$ .

$\text{DistLaw}_{\mathcal{S}}(\mathbf{M})$  whose

objects consist of monoids  $B$  and  $Y$  in  $\mathbf{M}$ , together with a left  $B$ -action on  $Y$  which makes the multiplication and the unit of the monoid  $Y$  left  $B$ -linear.

morphisms are pairs of monoid morphisms  $(B \xrightarrow{b} B', Y \xrightarrow{y} Y')$  which are compatible with the actions  $BY \xrightarrow{l} Y$  and  $B'Y' \xrightarrow{l'} Y'$  in the sense that  $l'.by = y.l$ .

Recall that a bimonoid  $B$  — with monoid structure  $(m, u)$  and comonoid structure  $(\delta, \varepsilon)$  — is a *Hopf monoid* provided that there exists a morphism  $B \xrightarrow{z} B$  — the so-called *antipode* — which renders commutative

$$\begin{array}{ccccc} B & \xrightarrow{\delta} & B^2 & \xrightarrow{z1} & B^2 \\ & \searrow \varepsilon & & & \downarrow m \\ & & I & \xrightarrow{u} & B \\ \delta \downarrow & & & & \\ B^2 & \xrightarrow{1z} & B^2 & \xrightarrow{m} & B. \end{array}$$

If the antipode exists then it is unique. It is a monoid morphism from  $B$  to the monoid with the opposite multiplication  $m.c$  and comonoid morphism from  $B$  to the comonoid with the opposite comultiplication  $c.\delta$ .

**Proposition 1.4.** (1) *The equivalent categories of Example 1.3 have equivalent full subcategories as follows.*

- *The category whose*

objects are split epimorphisms  $B \begin{smallmatrix} \xrightarrow{s} \\ \xleftarrow{i} \end{smallmatrix} A$  of bimonoids in  $\mathbf{M}$  subject to the following conditions.

- The comultiplication  $\delta$  of  $A$  satisfies  $c.s1.\delta = 1s.\delta$ .*
- $B$  is a Hopf monoid.*

morphisms are pairs of bimonoid morphisms which are compatible with the epimorphisms  $s$  as well as their sections  $i$ .

- *The category whose*

objects consist of a cocommutative Hopf monoid  $B$  and a bimonoid  $Y$  in  $\mathbf{M}$ , together with a left  $B$ -action on  $Y$  which makes  $Y$  both a left  $B$ -module monoid and a left  $B$ -module comonoid.

morphisms are pairs of bimonoid morphisms  $(B \xrightarrow{b} B', Y \xrightarrow{y} Y')$  which are compatible with the actions  $BY \xrightarrow{l} Y$  and  $B'Y' \xrightarrow{l'} Y'$  in the sense that  $l'.by = y.l$ .

- (2) *The equivalent categories of part (1) have equivalent full subcategories as follows.*

- *The category whose*  
objects are split epimorphisms  $B \begin{smallmatrix} \xrightarrow{s} \\ \xleftarrow{i} \end{smallmatrix} A$  of cocommutative Hopf monoids.  
morphisms are pairs of bimonoid morphisms which are compatible with the epimorphisms  $s$  as well as their sections  $i$ .
- *The category whose*  
objects consist of cocommutative Hopf monoids  $B$  and  $Y$  in  $\mathbf{M}$ , together with a left  $B$ -action on  $Y$  which makes  $Y$  both a left  $B$ -module monoid and a left  $B$ -module comonoid.  
morphisms are pairs of bimonoid morphisms  $( B \xrightarrow{b} B' , Y \xrightarrow{y} Y' )$  which are compatible with the actions  $BY \xrightarrow{l} Y$  and  $B'Y' \xrightarrow{l'} Y'$  in the sense that  $l'.by = y.l$ .

*Proof.* (1) The second listed category is obviously a full subcategory of  $\text{DistLaw}_S(\mathbf{C})$  of Example 1.3; thus via the equivalence of Theorem 1.1 it is equivalent to some full subcategory of  $\text{SplitEpiMon}_S(\mathbf{C})$  of Example 1.3. Our task is to show that it is the first listed category above. For that we only need to show that it is a subcategory of  $\text{SplitEpiMon}_S(\mathbf{C})$ ; that is, that for any object  $B \begin{smallmatrix} \xrightarrow{s} \\ \xleftarrow{i} \end{smallmatrix} A$  of it, the morphism  $q$  in part (b) of Example 1.3 is invertible. Following ideas in [15], we use the antipode  $z$  of  $B$  and the image of the equalizer (1.5) under the functor  $-B$  to construct the inverse:

$$\begin{array}{ccccccccccc}
 & & & & & & & & & & \xrightarrow{q^{-1}} & (A \square_B I)B \\
 & & & & & & & & & & & \downarrow j_1 \\
 A & \xrightarrow{\delta} & A^2 & \xrightarrow{1s} & AB & \xrightarrow{1\delta} & AB^2 & \xrightarrow{1z1} & AB^2 & \xrightarrow{1i1} & A^2B & \xrightarrow{m1} & AB \\
 & & & & & & & & & & & \downarrow \text{1s1.}\delta\text{1} & \downarrow \text{1u1} \\
 & & & & & & & & & & & & AB^2
 \end{array}$$

This definition works because the horizontal morphism equalizes the parallel morphisms of the fork on the right; see Figure 1. The so constructed morphism  $q^{-1}$  is the inverse of  $q$  by the commutativity of the diagrams of Figure 2 (in the second case we also need to use that the columns are equal monomorphisms).

(2) If both  $Y$  and  $B$  are cocommutative comonoids then clearly so is  $YB$ ; and if both  $Y$  and  $B$  have antipodes  $z$  then  $YB \xrightarrow{zz} YB$  is the antipode of the Hopf monoid  $YB$ .

Conversely, if  $A$  is cocommutative then evidently so is its sub-comonoid  $A \square_B I$ . If furthermore  $A$  has an antipode  $z$  then it restricts to  $A \square_B I$  by the commutativity of





FIGURE 1. Construction of  $q^{-1}$

FIGURE 2. Invertibility of  $q$ 



the following diagram.

$$\begin{array}{ccccc}
 A \square_B I & \xrightarrow{j} & A & \xrightarrow{z} & A \\
 \downarrow j & & \downarrow \delta & & \downarrow \delta \\
 & & A^2 & \xrightarrow{zz} & A^2 \\
 & & \downarrow 1s & & \downarrow 1s \\
 A & \xrightarrow{1u} & AB & & AB \\
 \downarrow z & & \searrow zz & & \downarrow 1s \\
 A & \xrightarrow{1u} & AB & & AB
 \end{array}$$

The top right region commutes by the Hopf monoid identity  $\delta.z = zz.c.\delta$  and the assumed cocommutativity of  $A$ . The bottom right region commutes since any bimonoid morphism  $s$  commutes with the antipodes.  $\square$

**Example 1.5.** Proposition 1.4 can be applied in particular to a finitely complete category  $\mathbf{M}$ , regarded as a Cartesian monoidal category. From Proposition 1.4 we obtain equivalences between the following pairs of categories.

- (1)
  - The category whose
    - objects are split epimorphisms  $B \begin{smallmatrix} \xrightarrow{s} \\ \xleftarrow{i} \end{smallmatrix} A$  of monoids in  $\mathbf{M}$  such that  $B$  is a group object.
    - morphisms are pairs of monoid morphisms which are compatible with the epimorphisms  $s$  as well as their sections  $i$ .
  - The category whose
    - objects consist of a group object  $B$  and a monoid  $Y$  in  $\mathbf{M}$ , together with a left  $B$ -action on  $Y$  which makes  $Y$  a left  $B$ -module monoid.
    - morphisms are pairs of monoid morphisms  $( B \xrightarrow{b} B' , Y \xrightarrow{y} Y' )$  which are compatible with the actions  $BY \xrightarrow{l} Y$  and  $B'Y' \xrightarrow{l'} Y'$  in the sense that  $l'.by = y.l$ .
- (2)
  - The category whose
    - objects are split epimorphisms  $B \begin{smallmatrix} \xrightarrow{s} \\ \xleftarrow{i} \end{smallmatrix} A$  of group objects.
    - morphisms are pairs of monoid morphisms which are compatible with the epimorphisms  $s$  as well as their sections  $i$ .
  - The category whose
    - objects consist of group objects  $B$  and  $Y$  in  $\mathbf{M}$ , together with a left  $B$ -action on  $Y$  which makes  $Y$  a left  $B$ -module group.
    - morphisms are pairs of bimonoid morphisms  $( B \xrightarrow{b} B' , Y \xrightarrow{y} Y' )$  which are compatible with the actions  $BY \xrightarrow{l} Y$  and  $B'Y' \xrightarrow{l'} Y'$  in the sense that  $l'.by = y.l$ .

*Remark 1.6.* There are particular symmetric monoidal categories  $\mathbf{M}$  whose cocommutative Hopf monoids constitute semi-abelian categories  $\mathbf{Hopf}(\mathbf{M})$ ; e.g. the category of sets (which is Cartesian monoidal hence the Hopf monoids are the groups, all of them cocommutative) or the category of vector spaces over an algebraically closed field (see [9]). In such cases the equivalence of Proposition 1.4 (2) is in fact the equivalence  $\mathbf{SplitEpi}(\mathbf{Hopf}(\mathbf{M})) \cong \mathbf{Act}(\mathbf{Hopf}(\mathbf{M}))$  discussed in [10, Section 1], see [10, Example 3.10].

## 2. REFLEXIVE GRAPHS OF MONOIDS VERSUS PRE-CROSSED MODULES

Consider a monoidal admissible class  $\mathcal{S}$  of spans in a monoidal category  $\mathbf{C}$  for which [2, Assumption 4.1] holds. Take an object  $B \overset{s}{\underset{i}{\rightrightarrows}} A$  in the category  $\mathbf{SplitEpiMon}_{\mathcal{S}}(\mathbf{C})$  of Theorem 1.1. Then by property (b) in Theorem 1.1, the induced morphism  $q := (A \square_B I)B \xrightarrow{p_A^i} A^2 \xrightarrow{m} A$  is invertible. Therefore by [2, Corollary 1.7] there is a bijective correspondence between the retractions  $t$  of the monoid morphism  $i$  and the monoid morphisms  $A \square_B I \xrightarrow{k} B$  rendering commutative

$$\begin{array}{ccccccc} B(A \square_B I) & \xrightarrow{ip_A} & A^2 & \xrightarrow{m} & A & \xrightarrow{q^{-1}} & (A \square_B I)B \\ \downarrow 1k & & & & & & \downarrow k1 \\ B^2 & \xrightarrow{m} & B & \xleftarrow{m} & B^2 & & \end{array}$$

The correspondence is given by

$$t \mapsto k := A \square_B I \xrightarrow{p_A} A \xrightarrow{t} B \quad k \mapsto t := A \xrightarrow{q^{-1}} (A \square_B I)B \xrightarrow{k1} B^2 \xrightarrow{m} B.$$

Combining this observation with the equivalence of Theorem 1.1, next we present an equivalent description of a suitable category of reflexive graphs of monoids. This leads to the notion of pre-crossed module over a monoid.

**Theorem 2.1.** *Consider a monoidal admissible class  $\mathcal{S}$  of spans in a monoidal category  $\mathbf{C}$  for which [2, Assumption 4.1] holds. The following categories are equivalent.*

$\mathbf{ReflGraphMon}_{\mathcal{S}}(\mathbf{C})$  whose

objects are reflexive graphs  $B \overset{s}{\underset{i}{\rightrightarrows}} A$  of monoids in  $\mathbf{C}$  subject to the following conditions.

(a)  $A = A \xrightarrow{s} B \in \mathcal{S}$  (hence the  $\mathcal{S}$ -relative pullback  $A \square_B I$  in Theorem 1.1 exists).

(b)  $q := (A \square_B I)B \xrightarrow{p_A^i} A^2 \xrightarrow{m} A$  is invertible.

morphisms are pairs of monoid morphisms  $(B \xrightarrow{b} B', A \xrightarrow{a} A')$  such that  $s'.a = b.s$ ,  $t'.a = b.t$  and  $i'.b = a.i$ .

$\mathbf{PreX}_{\mathcal{S}}(\mathbf{C})$  whose

objects consist of monoids  $B$  and  $Y$ , monoid morphisms  $Y \xrightarrow{e} I$  and  $Y \xrightarrow{k} B$  and a distributive law  $BY \xrightarrow{x} YB$  subject to the following conditions.

(a')  $Y = Y \xrightarrow{e} I \in \mathcal{S}$  and  $B = B = B \in \mathcal{S}$ .

(b')  $e1.x = 1e$  and  $m.k1.x = m.1k$ .

(c') The morphism  $f$  of Theorem 1.1 (c') is invertible.

morphisms are pairs of monoid morphisms  $(B \xrightarrow{b} B', Y \xrightarrow{y} Y')$  such that  $e'.y = e$ ,  $k'.y = b.k$  and  $x'.by = yb.x$ .

*Proof.* We show that the equivalence functors of Theorem 1.1 lift to the equivalence of the claim. In the direction  $\text{ReflGraphMon}_S(\mathcal{C}) \rightarrow \text{PreX}_S(\mathcal{C})$  we send

$$\begin{array}{ccc} B & \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{i} \\ \xleftarrow{t} \end{array} & A \\ b \downarrow & & \downarrow a \\ B' & \begin{array}{c} \xleftarrow{s'} \\ \xrightarrow{i'} \\ \xleftarrow{t'} \end{array} & A' \end{array}$$

to

$$\begin{array}{c} (A \square_B I, \quad B, \quad A \square_B I \xrightarrow{p_I} I, \quad A \square_B I \xrightarrow{p_A} A \xrightarrow{t} B, \quad B(A \square_B I) \xrightarrow{ip_A} A^2 \xrightarrow{m} A \xrightarrow{q^{-1}} (A \square_B I)B) \\ \downarrow a \square 1 \quad \downarrow b \\ (A' \square_{B'} I, \quad B', \quad A' \square_{B'} I \xrightarrow{p_I} I, \quad A' \square_{B'} I \xrightarrow{p_{A'}} A' \xrightarrow{t'} B', \quad B'(A' \square_{B'} I) \xrightarrow{i' p_{A'}} A'^2 \xrightarrow{m'} A' \xrightarrow{q'^{-1}} (A' \square_{B'} I)B') \end{array}$$

By [2, Proposition 3.7 (1)],  $p_A$  is a monoid morphism hence so is  $t.p_A$ . The second condition in (b') holds by the considerations preceding the theorem. Hence in light of the proof of Theorem 1.1 the object map is well-defined. Concerning the morphisms, the second condition holds by the commutativity of

$$\begin{array}{ccccc} A \square_B I & \xrightarrow{p_A} & A & \xrightarrow{t} & B \\ a \square 1 \downarrow & & \downarrow a & & \downarrow b \\ A' \square_{B'} I & \xrightarrow{p_{A'}} & A' & \xrightarrow{t'} & B' \end{array}$$

Thus using again the proof of Theorem 1.1 we conclude that this functor is well-defined.

In the opposite direction  $\text{PreX}_S(\mathcal{C}) \rightarrow \text{ReflGraphMon}_S(\mathcal{C})$  we put

$$\begin{array}{ccc} (Y, \quad B, \quad Y \xrightarrow{e} I, \quad Y \xrightarrow{k} B, \quad BY \xrightarrow{x} YB) & \mapsto & \begin{array}{ccc} B & \begin{array}{c} \xleftarrow{e1} \\ \xrightarrow{u1} \\ \xleftarrow{m.k1} \end{array} & YB \\ b \downarrow & & \downarrow yb \\ B' & \begin{array}{c} \xleftarrow{e'1} \\ \xrightarrow{u'1} \\ \xleftarrow{m'.k'1} \end{array} & Y'B' \end{array} \\ (Y', \quad B', \quad Y' \xrightarrow{e'} I, \quad Y' \xrightarrow{k'} B', \quad B'Y' \xrightarrow{x'} Y'B') & & \end{array}$$

By the considerations preceding the theorem  $m.k1$  is a monoid morphism. It is a retraction of  $B \xrightarrow{u1} AB$  by the unitality of  $k$ . The monoid morphisms  $(b, yb)$  are compatible with  $m.k1$  by the compatibility of  $(b, y)$  with  $k$  and the multiplicativity of  $b$ . So using again the proof of Theorem 1.1 we conclude that this functor is well-defined too.

By the commutativity of

$$\begin{array}{ccc} (A \square_B I)B \xrightarrow{p_{A^1}} AB \xrightarrow{t1} B^2 \xrightarrow{m} B & & Y \xrightarrow{=} Y \xrightarrow{k} B \\ \downarrow p_{Ai} \quad \searrow tt & & \downarrow f \quad \downarrow 1u \quad \downarrow 1u \\ A^2 & & YB \square_B I \xrightarrow{p_{YB}} YB \xrightarrow{k1} B^2 \xrightarrow{m} B \\ \downarrow m & & \downarrow m \\ A \xrightarrow{t} B & & \end{array}$$

the components  $(1, q)$  and  $(1, f)$  of the natural isomorphisms in the proof of Theorem 1.1 are morphisms in the appropriate category. This proves that the stated functors are mutually inverse equivalences.  $\square$

**Lemma 2.2.** *Consider a monoidal admissible class  $\mathcal{S}$  of spans in a monoidal category  $\mathbf{C}$  for which [2, Assumption 4.1] holds. For any object  $B \overset{s}{\underset{t}{\rightleftarrows}} A$  of the category  $\text{ReflGraphMon}_{\mathcal{S}}(\mathbf{C})$  of Theorem 2.1, the following assertions are equivalent.*

- (i)  $B \xleftarrow{t} A = A \in \mathcal{S}$ .
- (ii)  $B \xleftarrow{k:=t.p_A} A \square_B I = A \square_B I \in \mathcal{S}$ .

*Proof.* Assertion (i) implies (ii) by [2, Lemma 3.4]. Conversely, since  $B = B = B \in \mathcal{S}$  by assumption, (ii) implies  $B^2 \xleftarrow{k^1} (A \square_B I) B = (A \square_B I) B \in \mathcal{S}$  by the multiplicativity of  $\mathcal{S}$ . Hence by (PRE) also  $B^2 \xleftarrow{k^1} (A \square_B I) B \xleftarrow{q^{-1}} A \xrightarrow{q^{-1}} (A \square_B I) B \in \mathcal{S}$ . Then using the identity  $t = m.k^1.q^{-1}$  from the proof of Theorem 2.1, (i) follows by (POST) (composing by  $m$  on the left and by  $q$  on the right).  $\square$

**Example 2.3.** As in Example 1.2, take the (evidently admissible and monoidal) class of all spans in the monoidal category  $\mathbf{C}$  of spans over a fixed set  $X$ . Then the equivalent categories of Theorem 2.1 take the following forms.

$\text{ReflGraphMon}(\mathbf{C})$  whose

objects are reflective graphs  $B \overset{\sigma}{\underset{\tau}{\rightleftarrows}} A$  of categories with the common object set  $X$  and identity-on-objects functors between them, such that the map (1.3) in Example 1.2 is invertible (recall that this holds e.g. if  $B$  is a groupoid).  
morphisms are pairs of compatible identity-on-objects functors.

$\text{PreX}(\mathbf{C})$  whose

objects consist of categories  $B$  and  $Y$  of the common object set  $X$  such that in  $Y$  there are no morphisms between non-equal objects; an action (cf. Example 1.2)  $B \square_X Y \xrightarrow{\triangleright} Y$  and an identity-on-objects functor  $Y \xrightarrow{\kappa} B$  such that

$$\kappa(b \triangleright y).b = b.\kappa(y) \quad (2.1)$$

for all morphisms  $b$  in  $B$  and  $y$  in  $Y$  for which  $s(b) = t(y)$ . (If  $B$  is a groupoid then (2.1) has the equivalent form  $\kappa(b \triangleright y) = b.\kappa(y).b^{-1}$ ; so when both  $B$  and  $Y$  are groupoids we recover the notion of *pre-crossed module* of groupoids in [4, Definition 1.2].)

morphisms are pairs of identity-on-objects functors  $(B \xrightarrow{\beta} B', Y \xrightarrow{\nu} Y')$  such that  $\kappa'\nu = \beta\kappa$  and  $\nu(b \triangleright y) = \beta(b) \triangleright \nu(y)$  for all morphisms  $b$  in  $B$  and  $y$  in  $Y$  for which  $s(b) = t(y)$ .

**Example 2.4.** In the setting of Example 1.3, the equivalent categories of Theorem 2.1 take the following explicit forms.

$\text{ReflGraphMon}_{\mathcal{S}}(\mathcal{C})$  whose

objects are reflective graphs  $B \begin{smallmatrix} \xrightarrow{s} \\ \xleftrightarrow{i} \\ \xleftarrow{t} \end{smallmatrix} A$  of bimonoids in  $\mathbf{M}$  subject to the following conditions.

(a) The comultiplication  $\delta$  of  $A$  satisfies  $c.s1.\delta = 1s.\delta$ .

(b) In terms of the morphism  $j$  of (1.5),  $q := (A \square_B I)B \xrightarrow{ji} A^2 \xrightarrow{m} A$  is invertible.

morphisms are pairs of bimonoid morphisms  $(B \xrightarrow{b} B', A \xrightarrow{a} A')$  such that  $s'.a = b.s$ ,  $t'.a = b.t$  and  $i'.b = a.i$ .

$\text{PreX}_{\mathcal{S}}(\mathcal{C})$  whose

objects consist of a cocommutative bimonoid  $B$  and a bimonoid  $Y$  in  $\mathbf{M}$ , together with a left  $B$ -action on  $Y$  which makes  $Y$  both a left  $B$ -module monoid and a left  $B$ -module comonoid, and a bimonoid morphism  $Y \xrightarrow{k} B$  for which the following diagram commutes.

$$\begin{array}{ccccc} BY & \xrightarrow{\delta 1} & B^2 Y & \xrightarrow{1c} & BYB & \xrightarrow{l1} & YB \\ \downarrow 1k & & & & & & \downarrow k1 \\ B^2 & \xrightarrow{m} & B & \xleftarrow{m} & B^2 & & \end{array} \quad (2.2)$$

morphisms are pairs of bimonoid morphisms  $(B \xrightarrow{b} B', Y \xrightarrow{y} Y')$  which are compatible with the actions  $BY \xrightarrow{l} Y$  and  $B'Y' \xrightarrow{l'} Y'$  in the sense that  $l'.by = y.l$  and which satisfy  $k'.y = b.k$ .



*Remark 2.5.* Clearly, the equivalent categories of Example 2.4 have equivalent full subcategories for whose objects the bimonoid  $B$  is a cocommutative Hopf monoid (then condition (b) becomes redundant by Example 1.4). Note that whenever  $B$  has an antipode  $z$ , the commutative diagram (2.2) has an equivalent form

$$\begin{array}{ccccc} BY & \xrightarrow{\delta k} & B^3 & \xrightarrow{1c} & B^3 & \xrightarrow{mz} & B^2 \\ \downarrow l & & & & & & \downarrow m \\ Y & \xrightarrow{k} & B & & B & & \end{array} \quad (2.3)$$

occurring in [16, Definition 12 (iv)]. Their equivalence follows by the commutativity of the diagrams of Figure 3.

### 3. RELATIVE CATEGORIES OF MONOIDS VERSUS CROSSED MODULES

Consider again a monoidal admissible class  $\mathcal{S}$  of spans in a monoidal category  $\mathcal{C}$  for which [2, Assumption 4.1] holds. Take an object  $B \begin{smallmatrix} \xrightarrow{s} \\ \xleftrightarrow{i} \\ \xleftarrow{t} \end{smallmatrix} A$  of the category  $\text{ReflGraphMon}_{\mathcal{S}}(\mathcal{C})$  of Theorem 2.1 such that also  $B \xleftarrow{t} A = A \in \mathcal{S}$ ; that is, the legs of the cospan  $A \xrightarrow{s} B \xleftarrow{t} A$  are in  $\mathcal{S}$  (hence there exists its  $\mathcal{S}$ -relative pullback

FIGURE 3. Equivalence of (2.2) and (2.3) for Hopf monoids  $B$ 

$A \xleftarrow{p_1} A \square_B A \xrightarrow{p_2} A$ ). Whenever the morphism

$$q_2 := (A \square_B I)A \xrightarrow{p_{A^1}} A^2 \xrightarrow{(1 \square i)(i \square 1)} (A \square_B A)^2 \xrightarrow{m} A \square_B A \quad (3.1)$$

is invertible, we infer from [2, Corollary 1.7] that there exists at most one monoid morphism  $d$  rendering commutative

$$\begin{array}{ccccc} A \square_B I & \xrightarrow{p_A} & A & \xrightarrow{1 \square i} & A \square_B A & \xleftarrow{i \square 1} & A \\ & \searrow^{p_A} & & & \downarrow d & & \\ & & & & A & & \end{array} \quad (3.2)$$

which is our candidate to serve as the composition morphism of a relative category. By this motivation, in this section we investigate first the condition that (3.1) is invertible. Assuming so, next we show that whenever the morphism  $d$  of (3.2) exists, it makes the object  $B \overset{s}{\underset{t}{\rightleftarrows}} A$  of  $\text{ReflGraphMon}_{\mathcal{S}}(\mathcal{C})$  to an  $\mathcal{S}$ -relative category. Finally, based on Theorem 2.1, we give an equivalent description of the category of  $\mathcal{S}$ -relative categories in the category of monoids in  $\mathcal{C}$ , in terms of crossed modules introduced hereby.

### 3.1. Invertibility of some canonical morphisms.

**Lemma 3.1.** *Consider a monoidal admissible class  $\mathcal{S}$  of spans in a monoidal category  $\mathcal{C}$  for which [2, Assumption 4.1] holds. For any monoid  $B$  in  $\mathcal{C}$  for which  $B = B = B$  is in  $\mathcal{S}$ , for any span of monoids  $B \overset{t}{\leftarrow} A \overset{s}{\rightarrow} B$  with legs in  $\mathcal{S}$ , and for any natural number  $n$ , the following assertions hold. (Recall the convention  $A \square_B^0 := B$  from [2, Corollary 4.6].)*

(1) *There exists the  $\mathcal{S}$ -relative pullback*

$$\begin{array}{ccc} (A \square_B I)B \square_B A \square_B^{\square n} & \xrightarrow{p_{A \square_B^{\square n}}} & A \square_B^{\square n} \\ p_{(A \square_B I)B} \downarrow & & \downarrow t.p_1 \\ (A \square_B I)B & \xrightarrow{p_I 1} & B. \end{array}$$

(2) *There is a unique morphism  $h_n$  rendering commutative*

$$\begin{array}{ccccc} (A \square_B I)A \square_B^{\square n} & \xrightarrow{p_I 1} & & & A \square_B^{\square n} \\ \downarrow 1p_1 & \dashrightarrow^{h_n} & (A \square_B I)B \square_B A \square_B^{\square n} & \xrightarrow{p_{A \square_B^{\square n}}} & A \square_B^{\square n} \\ & & p_{(A \square_B I)B} \downarrow & & \downarrow t.p_1 \\ (A \square_B I)A & \xrightarrow{1t} & (A \square_B I)B & \xrightarrow{p_I 1} & B. \end{array}$$

(3) For a common section  $i$  of  $s$  and  $t$ , consider the morphism

$$q_{n+1} := (A \square_B I) A_B^{\square n} \xrightarrow{p_{A^1}} A A_B^{\square n} \xrightarrow{(1 \square i \square \dots \square i)(i \square 1 \square \dots \square 1)} (A_B^{\square n+1})^2 \xrightarrow{m} A_B^{\square n+1} \quad (3.3)$$

(it is well-defined by [2, Proposition 3.5] and  $q_1$  is equal to  $q$  in Theorem 1.1 (b)). If  $q_{n+1}$  is invertible for some  $n$ , then  $q_k$  is invertible for all  $0 < k \leq n$ .

(4) For a common section  $i$  of  $s$  and  $t$  the following are equivalent.

- (i)  $h_n$  in part (2) and  $q_1$  in part (3) are invertible.
- (ii)  $q_{n+1}$  in part (3) is invertible.

*Proof.* (1) By assumption  $B \xleftarrow{t} A = A \in \mathcal{S}$  and by the unitality of  $\mathcal{S}$ ,  $I = I = I \in \mathcal{S}$ . Then by [2, Lemma 3.4],

$$B \xleftarrow{t} A \xleftarrow{p_1} A_B^{\square n} = A_B^{\square n} \in \mathcal{S} \quad \text{and} \quad A \square_B I = A \square_B I \xrightarrow{p_I} I \in \mathcal{S}. \quad (3.4)$$

By assumption also  $B = B = B \in \mathcal{S}$  hence by the second assertion in (3.4) and the multiplicativity of  $\mathcal{S}$

$$(A \square_B I) B = (A \square_B I) B \xrightarrow{p_I^1} B \in \mathcal{S}. \quad (3.5)$$

The first assertion of (3.4) and (3.5) say that the legs of  $(A \square_B I) B \xrightarrow{p_I^1} B \xleftarrow{t, p_1} A_B^{\square n}$  are in  $\mathcal{S}$  hence their  $\mathcal{S}$ -relative pullback exists by assumption.

(2) By (3.4) and the multiplicativity of  $\mathcal{S}$ ,

$$(A \square_B I) B \xleftarrow{1t} (A \square_B I) A \xleftarrow{1p_1} (A \square_B I) A_B^{\square n} \xrightarrow{p_I^1} A_B^{\square n} \in \mathcal{S}.$$

Hence by the evident commutativity of the exterior of the diagram in part (2), universality of the  $\mathcal{S}$ -relative pullback in its codomain implies the existence of the unique morphism  $h_n$ .

(3) For some positive integer  $n$  assume that  $q_{n+1}$  is invertible. Then so is  $q_n$  with the inverse

$$A_B^{\square n} \xrightarrow{1 \square i} A_B^{\square n+1} \xrightarrow{q_{n+1}^{-1}} (A \square_B I) A_B^{\square n} \xrightarrow{1p_{1\dots n-1}} (A \square_B I) A_B^{\square n-1}. \quad (3.6)$$

Indeed, (3.6) renders commutative both diagrams

$$\begin{array}{ccccc} (A \square_B I) A_B^{\square n-1} & \xlongequal{\quad} & (A \square_B I) A_B^{\square n-1} & A_B^{\square n+1} & \xrightarrow{q_{n+1}^{-1}} & (A \square_B I) A_B^{\square n} & \xrightarrow{1p_{1\dots n-1}} & (A \square_B I) A_B^{\square n-1} \\ \downarrow q_n & \nearrow 1(1 \square i) & \downarrow q_{n+1} & \uparrow 1 \square i & \searrow q_{n+1} & \downarrow q_n & \searrow p_{1\dots n} & \downarrow q_n \\ A_B^{\square n} & \xrightarrow{1 \square i} & A_B^{\square n+1} & \xrightarrow{q_{n+1}^{-1}} & (A \square_B I) A_B^{\square n} & \xrightarrow{1 \square i} & A_B^{\square n} & \xrightarrow{p_{1\dots n}} & A_B^{\square n} \end{array}$$

The leftmost region of the first diagram commutes by the explicit expression (3.3) of  $q_n$  and  $q_{n+1}$ , multiplicativity of  $1 \square i$  and the functoriality of  $\square$ , see [2, Proposition 3.5 (2)]. The rightmost region of the second diagram commutes again by the explicit expression (3.3) of  $q_n$  and  $q_{n+1}$  and the multiplicativity of  $p_{1\dots n}$ .



(4) Our strategy is to prove that  $q_{n+1}$  can be rewritten as

$$(A \square_B I) A_B^{\square n} \xrightarrow{h_n} (A \square_B I) B \square_B A_B^{\square n} \xrightarrow{q \square 1} A_B^{\square n+1}. \quad (3.7)$$

Then (i) obviously implies (ii) and in view of part (3) also the opposite implication holds.

The occurring morphism  $q \square 1$  is defined as the unique morphism rendering commutative

$$\begin{array}{ccccc} (A \square_B I) B \square_B A_B^{\square n} & \xrightarrow{p_{A \square_B n}} & & & \\ \downarrow p_{(A \square_B I) B} & \searrow q \square 1 & & & \\ (A \square_B I) B & \xrightarrow{q} & A & \xrightarrow{s} & B \end{array}$$

It is well-defined by the commutativity of the first diagram of (1.1); see [2, Proposition 3.5 (2)]. The morphism of (3.7) is equal to  $q_{n+1}$  by the commutativity of both diagrams

$$\begin{array}{ccccccc} & \xrightarrow{h_n} & (A \square_B I) B \square_B A_B^{\square n} & \xrightarrow{q \square 1} & A_B^{\square n+1} & & \\ & & \downarrow p_{(A \square_B I) B} & & \downarrow p_1 & & \\ (A \square_B I) A_B^{\square n} & \xrightarrow{1p_1} & (A \square_B I) A & \xrightarrow{1t} & (A \square_B I) B & \xrightarrow{p_{A^1}} & AB & \xrightarrow{1i} & A^2 & \xrightarrow{m} & A \\ & & & & \uparrow 1t & & \uparrow 1p_1 & & \uparrow p_1 p_1 & & \uparrow p_1 \\ & & & & A^2 & & & & & & \\ & & & & \uparrow 1p_1 & & & & & & \\ & & & & AA_B^{\square n} & \xrightarrow{(1 \square i \square \dots \square i)(i \square 1)} & (A_B^{\square n+1})^2 & \xrightarrow{m} & A_B^{\square n+1} & & \end{array} \quad (3.8)$$

$$\begin{array}{ccccccc} & \xrightarrow{h_n} & (A \square_B I) B \square_B A_B^{\square n} & \xrightarrow{q \square 1} & A_B^{\square n+1} & & \\ & & \downarrow p_{A \square_B n} & & \downarrow p_{2 \dots n} & & \\ (A \square_B I) A_B^{\square n} & \xrightarrow{p_I 1} & A_B^{\square n} & \xrightarrow{u 1} & BA_B^{\square n} & \xrightarrow{(i \square \dots \square i) 1} & (A_B^{\square n})^2 & \xrightarrow{m} & A_B^{\square n} \\ & & & & \uparrow s 1 & & \uparrow p_{2 \dots n} p_{2 \dots n} & & \uparrow p_{2 \dots n} \\ & & & & AA_B^{\square n} & \xrightarrow{(1 \square i \square \dots \square i)(i \square 1)} & (A_B^{\square n+1})^2 & \xrightarrow{m} & A_B^{\square n+1} & & \end{array} \quad (3.9)$$

whose right verticals are joint monomorphisms.  $\square$

**Example 3.2.** In the category  $\mathbf{C}$  of spans over a given set  $X$  from Example 1.2, the morphisms  $h_n$  of Lemma 3.1 (2) are isomorphisms, see the pullback (1.4). Hence for any reflexive graph  $B \begin{array}{c} \xrightarrow{\sigma} \\ \xleftrightarrow{\tau} \\ \xleftarrow{\tau} \end{array} A$  of categories with common object set  $X$  and identity-on-objects functors between them, all morphisms  $\{q_n\}_{n>0}$  in Lemma 3.1 (3) are invertible if and only if  $q_1$  is so; see Lemma 3.1 (3). The latter condition holds e.g. if  $B$  is a groupoid, see Example 1.2.



**Lemma 3.4.** *Let  $\mathcal{S}$  be a monoidal admissible class of spans in a monoidal category  $\mathcal{C}$  for which [2, Assumption 4.1] holds and let  $(B, Y, Y \xrightarrow{e} I, Y \xrightarrow{k} B, BY \xrightarrow{x} YB)$  be an object of the category  $\text{PreX}_{\mathcal{S}}(\mathcal{C})$  in Theorem 2.1 such that  $B \xleftarrow{k} Y = Y \in \mathcal{S}$ . For any natural number  $n$  denote by  $B^{n+1} \xrightarrow{m^{(n)}} B$  the  $n$ -times iterated multiplication (unique by the associativity of  $m$ ; by definition the identity morphism for  $n = 0$ ) and consider the span*

$$B \xleftarrow{m^{(n)}} B^{n+1} \xleftarrow{k \dots k1} Y^n B \xrightarrow{e \dots e1} B. \quad (3.10)$$

For any natural number  $n$  the following assertions hold.

- (1) The cospan  $YB \xrightarrow{e1} B \xleftarrow{m^{(n)}} B^{n+1} \xleftarrow{k \dots k1} Y^n B$  has its legs in  $\mathcal{S}$  (hence there exists its  $\mathcal{S}$ -relative pullback  $YB \square_B Y^n B$ ).
- (2) There exists a unique morphism  $b_{n+1}$  of spans (for the spans (3.10)) rendering commutative

$$\begin{array}{ccccc}
 Y^{n+1}B & & \xrightarrow{e1 \dots 11} & & Y^n B \\
 \downarrow 1k \dots k1 & \searrow b_{n+1} & & \xrightarrow{p_{Y^n B}} & \downarrow k \dots k1 \\
 & YB \square_B Y^n B & & & B^{n+1} \\
 & \downarrow p_{YB} & & & \downarrow m^{(n)} \\
 YB^{n+1} & \xrightarrow{1m^{(n)}} & YB & \xrightarrow{e1} & B.
 \end{array}$$

- (3) If  $b_{n+1}$  in part (2) is an isomorphism then also  $b_k$  is an isomorphism for all  $0 < k \leq n$ .
- (4) For the morphism

$$q_{n+1} := (YB \square_B I)(YB) \square_B^{p_{YB}1} \xrightarrow{p_{YB}1} YB(YB) \square_B^{(1 \square u1 \square \dots \square u1)(u1 \square 1)} \xrightarrow{((YB) \square_B^{n+1})^2} \xrightarrow{m} (YB) \square_B^{n+1}$$

the following diagram commutes

$$\begin{array}{ccccccc}
 Y^{n+1}B & \xrightarrow{b_{n+1}} & YB \square_B Y^n B & \xrightarrow{1 \square b_n} & \dots & \xrightarrow{1 \square b_1} & (YB) \square_B^{n+1} \square_B \\
 \downarrow f \dots f1 & & & & & & \downarrow p_{(YB) \square_B^{n+1}} \\
 (YB \square_B I)^{n+1} B & \xrightarrow{1 \dots 1q_1} & (YB \square_B I)^n YB & \xrightarrow{1 \dots 1q_2} & \dots & \xrightarrow{q_{n+1}} & (YB) \square_B^{n+1}
 \end{array}$$

where  $f$  is the isomorphism in Theorem 1.1 (c').

- (5)  $b_{n+1}$  in part (2) is an isomorphism if and only if  $q_{n+1}$  in part (4) is an isomorphism.

*Proof.* (1) By definition the first two spans in

$$Y = Y \xrightarrow{e} I \quad B = B = B \quad B \xleftarrow{k} Y = Y \quad YB = YB \xrightarrow{e1} B \quad (3.11)$$

belong to  $\mathcal{S}$  hence so does the last one by the multiplicativity of  $\mathcal{S}$ . Again, by definition the second and the third spans of (3.11) belong to  $\mathcal{S}$  hence by the multiplicativity of

$\mathcal{S}$  so does the first one in

$$B^{n+1} \xleftarrow{k\dots k1} Y^n B = Y^n B \quad B \xleftarrow{m^{(n)}} B^{n+1} \xleftarrow{k\dots k1} Y^n B = Y^n B. \quad (3.12)$$

Then the second span of (3.12) is in  $\mathcal{S}$  by (POST).

(2) Since the first span of (3.11) and the second span of (3.12) are in  $\mathcal{S}$ , the multiplicativity of  $\mathcal{S}$  implies that so is

$$YB \xleftarrow{1m^{(n)}} YB^{n+1} \xleftarrow{1k\dots k1} Y^{n+1}B \xrightarrow{e1\dots 11} Y^n B.$$

So by the evident commutativity of the exterior of the diagram of part (2) the stated morphism  $b_{n+1}$  exists. It is a morphism of spans (for the spans (3.10)) by the commutativity of the following diagrams.

$$\begin{array}{ccc} Y^{n+1}B & \xrightarrow{e\dots e1} & B \\ b_{n+1} \downarrow & \searrow e1\dots 1 & \parallel \\ YB \square_B Y^n B & \xrightarrow{p_{Y^n B}} & Y^n B \xrightarrow{e\dots e1} B \end{array} \quad \begin{array}{ccccc} Y^{n+1}B & \xrightarrow{1k\dots k1} & YB^{n+1} & \xrightarrow{k1\dots 11} & B^{n+2} \xrightarrow{m^{(n+1)}} B \\ b_{n+1} \downarrow & & \downarrow 1m^{(n)} & & \downarrow 1m^{(n)} \\ YB \square_B Y^n B & \xrightarrow{p_{YB}} & YB & \xrightarrow{k1} & B^2 \xrightarrow{m} B \end{array}$$

(3) Since for a positive integer  $n$ ,  $Y^{n-1}B \xrightarrow{1\dots 1u1} Y^n B$  is a morphism between the spans of (3.10), the morphism in the top row of the following diagram is well-defined by [2, Proposition 3.5].

$$\begin{array}{ccc} & \xrightarrow{1\Box 1\dots 1u1} & YB \square_B Y^{n-1}B \rightarrow YB \square_B Y^n B \\ b_n \curvearrowright & & \downarrow p_{YB} \\ Y^n B = Y^n B & \xrightarrow{1k\dots k1} & YB^n \xrightarrow{1m^{(n-1)}} YB \\ \uparrow 1\dots 1u1 & \nearrow 1\dots 1m1 & \nearrow 1m^{(n)} \\ Y^{n+1}B & \xrightarrow{1k\dots k1} & YB^{n+1} \xrightarrow{1m^{(n)}} YB \\ \downarrow 1\dots 1u1 & \searrow b_{n+1} & \downarrow p_{YB} \\ & \xrightarrow{b_{n+1}} & YB \square_B Y^n B \end{array} \quad \begin{array}{ccc} & \xrightarrow{1\Box 1\dots 1u1} & YB \square_B Y^{n-1}B \rightarrow YB \square_B Y^n B \\ b_n \curvearrowright & & \downarrow p_{Y^{n-1}B} \\ Y^n B & \xrightarrow{e1\dots 11} & Y^{n-1}B \xrightarrow{1\dots 1u1} Y^n B \\ \uparrow 1\dots 1u1 & \nearrow e1\dots 1 & \nearrow p_{Y^n B} \\ Y^{n+1}B & \xrightarrow{b_{n+1}} & YB \square_B Y^n B \end{array}$$

By their commutativity we infer  $b_{n+1}.1\dots 1u1 = (1\Box 1\dots 1u1).b_n$ . Similarly, since for  $n > 0$  also  $Y^n B \xrightarrow{1\dots 1m1} Y^{n-1}B$  is a morphism between the spans of (3.10), the morphism in the top row of the following diagram is well-defined by [2, Proposition 3.5].

$$\begin{array}{ccc} b_{n+1} \curvearrowright & YB \square_B Y^n B \xrightarrow{1\Box 1\dots 1m1} & YB \square_B Y^{n-1}B \\ & \searrow p_{YB} & \downarrow p_{YB} \\ Y^{n+1}B & \xrightarrow{1k\dots k1} & YB^{n+1} \xrightarrow{1m^{(n)}} YB \\ \uparrow 1\dots 1m1 & \nearrow 1\dots 1m1 & \nearrow 1m^{(n-1)} \\ & \xrightarrow{1\dots 1m1} & YB^n \xrightarrow{1m^{(n-1)}} YB \\ \downarrow 1\dots 1m1 & \searrow b_n & \downarrow p_{YB} \\ & \xrightarrow{b_n} & YB \square_B Y^{n-1}B \end{array} \quad \begin{array}{ccc} b_{n+1} \curvearrowright & YB \square_B Y^n B \xrightarrow{1\Box 1\dots 1m1} & YB \square_B Y^{n-1}B \\ & \searrow p_{Y^n B} & \downarrow p_{Y^{n-1}B} \\ Y^{n+1}B & \xrightarrow{e1\dots 11} & Y^n B \xrightarrow{1\dots 1m1} Y^{n-1}B \\ \uparrow 1\dots 1m1 & \nearrow e1\dots 1 & \nearrow p_{Y^{n-1}B} \\ & \xrightarrow{1\dots 1m1} & YB^n \xrightarrow{1m^{(n-1)}} YB \\ \downarrow 1\dots 1m1 & \searrow b_n & \downarrow p_{Y^{n-1}B} \\ & \xrightarrow{b_n} & YB \square_B Y^{n-1}B \end{array}$$

By their commutativity,  $b_n.1\dots 1m1 = (1\Box 1\dots 1m1).b_{n+1}$ . It follows from these identities and the unitality of the monoid  $Y$  that whenever  $b_{n+1}$  is invertible then so

is  $b_n$  with the inverse

$$YB \square_B Y^{n-1}B \xrightarrow{1 \square 1 \dots 1 u 1} YB \square_B Y^n B \xrightarrow{b_{n+1}^{-1}} Y^{n+1}B \xrightarrow{1 \dots 1 m 1} Y^n B.$$

(4) We proceed by induction in  $n$ . For  $n = 0$  the diagram in the claim reduces to

$$\begin{array}{ccc} YB & \xrightarrow{b_1} & YB \square_B B \\ f_1 \downarrow & \searrow & \downarrow p_{YB} \\ (YB \square_B I)B & \xrightarrow{q_1} & YB \end{array}$$

whose upper half commutes by construction (see part (2)) and the lower half commutes since  $f_1$  and  $q_1$  are mutual inverses (see the proof of Theorem 1.1).

For any positive value of  $n$ , denote the top-right path in the diagram of the claim by  $\tilde{b}_{n+1}$  and the bottom row by  $\tilde{q}_{n+1}$ . Then the diagram takes the form


$$\begin{array}{ccc} Y^{n+1}B & \xrightarrow{\tilde{b}_{n+1}} & (YB) \square_B^{n+1} \\ f_1 \dots 1 1 \downarrow & & \parallel \\ (YB \square_B I)Y^n B & \searrow \tilde{b}_n & \\ 1 f \dots f 1 \downarrow & \xrightarrow{1 \tilde{q}_n} & (YB \square_B I)(YB) \square_B^n \xrightarrow{q_{n+1}} (YB) \square_B^{n+1} \\ & \xrightarrow{\tilde{q}_{n+1}} & \end{array}$$

The region at the bottom left corner commutes if the claim holds for  $n - 1$ ; and the commutativity of the large region is proven in Figure 4.

(5) By Theorem 1.1  $q_1$  is an isomorphism without any further assumption; it is the inverse of the isomorphism  $YB \xrightarrow{f_1} (YB \square_B I)B$ . Also  $b_1$  is an isomorphism; the inverse of the isomorphism  $YB \square_B \xrightarrow{p_{YB}} YB$  in [2, Proposition 3.6 (1)].

Assume that  $b_l$  is iso for some  $l > 1$ . Take the diagram of part (4) for  $n = 1$ ; it says  $b_2 = q_2.f_1 1$ . Since  $f$  is an isomorphism by definition and  $b_2$  is an isomorphism by part (3), also  $q_2$  is an isomorphism. If  $l = 2$  then this completes the proof. If  $l > 2$  then take next the diagram of part (4) for  $n = 2$ ; it says  $(1 \square b_2).b_3 = q_3.1 q_2.f f 1 1$ . All of the occurring morphisms but  $q_3$  are known to be isomorphisms proving that so is  $q_3$ . Repeating this reasoning for all  $n \leq l$  we conclude that  $q_n$  is an isomorphism for all  $0 < n \leq l$ .

The opposite implication is proven by the same steps. Assume that  $q_l$  is iso for some  $l > 1$ . Take the diagram of part (4) for  $n = 1$ ; it says  $b_2 = q_2.f_1 1$ . Since  $f$  is an isomorphism by definition and  $q_2$  is an isomorphism by Lemma 3.1 (3), also  $b_2$  is an isomorphism. If  $l = 2$  then this completes the proof. If  $l > 2$  then take next the diagram of part (4) for  $n = 2$ ; it says  $(1 \square b_2).b_3 = q_3.1 q_2.f f 1 1$ . All of the occurring morphisms but  $b_3$  are known to be isomorphisms proving that so is  $b_3$ . Repeating this reasoning for all  $n \leq l$  we conclude that  $b_n$  is an isomorphism for all  $0 < n \leq l$ .  $\square$

FIGURE 4. Proof of  $\tilde{b}_{n+1} = q_{n+1} \cdot \tilde{b}_n$ 

**Example 3.5.** Take  $\mathcal{S}$  to be the (monoidal and admissible) class of all spans in the monoidal category  $\mathbf{C}$  of spans over a given set. For any object of the category  $\text{ReflGraphMon}(\mathbf{C})$  of Example 2.3 and for any positive integer  $n$ , the morphism  $b_n$  in Lemma 3.4 (2) is invertible, see the pullback (1.4).

**Example 3.6.** In the setting of Example 1.3 we know from Example 3.3 that the morphism  $q_n$  of Lemma 3.1 (3) is invertible for any positive integer  $n$  and for any object of  $\text{ReflGraphMon}_{\mathcal{S}}(\mathbf{C})$ . By the isomorphism of Theorem 2.1 this means that the morphism  $q_n$  of Lemma 3.4 (4) is invertible for any object of  $\text{PreX}_{\mathcal{S}}(\mathbf{C})$ . Then also the morphism  $b_n$  of Lemma 3.4 (2) is invertible by Lemma 3.4 (5). Since the diagram

$$\begin{array}{ccccc}
 Y^n B & \xrightarrow{\quad\quad\quad} & Y^n B & & \\
 \parallel & \searrow^{\delta_Y \delta_{Y^{n-1} B}} & \delta_{Y^1 \dots 11} & \xrightarrow{\quad\quad\quad} & Y^n B \\
 & & Y^2 (Y^{n-1} B)^2 & \xrightarrow{11 \varepsilon_{Y^{n-1} B} 1 \dots 11} & Y^{n+1} B \\
 & \searrow^{1c_{Y, Y^{n-1} B} 1 \dots 11} & \downarrow & \xrightarrow{1 \varepsilon_{Y^{n-1} B} 1 \dots 11} & \uparrow & \xrightarrow{1 \varepsilon_B 1 \dots 11} \\
 Y^n B & \xrightarrow{\delta_{Y^n B}} & (Y^n B)^2 & \xrightarrow{1k \dots k1 1 \dots 11} & Y B^n Y^n B & \xrightarrow{1m^{(n-1)} 1 \dots 1} & Y B Y^n B \\
 \downarrow b_n & & \downarrow b_n b_n & & \uparrow & \xrightarrow{11 \varepsilon_Y 1 \dots 11} & \uparrow & \xrightarrow{1 \varepsilon_B 1 \dots 11} \\
 Y B \square_B Y^{n-1} B & \xrightarrow{\delta} & (Y B \square_B Y^{n-1} B)^2 & \xrightarrow{p_Y B p_{Y^{n-1} B}} & Y B Y^{n-1} B & & \\
 & & & \searrow^j & & & \\
 & & & & & & 
 \end{array}$$

commutes, we conclude that the morphism in its bottom-right path — involving the equalizer  $j$  as in (1.5) — is the inverse of  $b_n$ .

**Lemma 3.7.** Let  $\mathcal{S}$  be a monoidal admissible class of spans in a monoidal category  $\mathbf{C}$  for which [2, Assumption 4.1] holds and let  $(B, Y, Y \xrightarrow{e} I, Y \xrightarrow{k} B, BY \xrightarrow{x} YB)$  be an object of the category  $\text{PreX}_{\mathcal{S}}(\mathbf{C})$  in Theorem 2.1 such that  $B \xleftarrow{k} Y = Y \in \mathcal{S}$ . For any positive integer  $n$  the morphism  $b_n$  in Lemma 3.4 (2) satisfies the following identities.

- (1)  $b_2 \cdot u11 = u1 \square 1$
- (2)  $b_2 \cdot 1u1 = 1 \square u1$

*Proof.* Assertion (1) follows by the commutativity of the diagrams

$$\begin{array}{ccc}
 & Y^2 B \xrightarrow{b_2} Y B \square_B Y B & \\
 u11 \curvearrowright & \downarrow 1k1 & \downarrow p_1 \\
 & Y B^2 & Y B \\
 Y B \xrightarrow{k1} B^2 \xrightarrow{m} B \xrightarrow{u1} Y B & & \uparrow p_1 \\
 & & Y B \square_B Y B \\
 & \searrow^{u1 \square 1} & \\
 & & 
 \end{array}
 \qquad
 \begin{array}{ccc}
 & Y^2 B \xrightarrow{b_2} Y B \square_B Y B & \\
 u11 \curvearrowright & \searrow^{e11} & \downarrow p_2 \\
 & & Y B \\
 Y B \xrightarrow{\quad\quad\quad} Y B & & \uparrow p_2 \\
 & \searrow^{u1 \square 1} & \\
 & & Y B \square_B Y B \\
 & & 
 \end{array}$$

and part (2) follows by the commutativity of

$$\begin{array}{ccc}
 & Y^2B & \xrightarrow{b_2} & YB \square_B YB \\
 1u1 \curvearrowright & \downarrow 1k1 & & \downarrow p_1 \\
 YB & \xrightarrow{1u1} & YB^2 & \xrightarrow{1m} & YB \\
 & \searrow 1 \square u1 & & \uparrow p_1 & \\
 & & & & YB \square_B YB
 \end{array}
 \qquad
 \begin{array}{ccc}
 & Y^2B & \xrightarrow{b_2} & YB \square_B YB \\
 1u1 \curvearrowright & \searrow e11 & & \downarrow p_2 \\
 YB & \xrightarrow{e1} & B & \xrightarrow{u1} & YB \\
 & \searrow 1 \square u1 & & \uparrow p_2 & \\
 & & & & YB \square_B YB
 \end{array}$$

□

### 3.2. The composition morphism of a relative category of monoids.

**Proposition 3.8.** *Consider a monoidal admissible class  $\mathcal{S}$  of spans in a monoidal category  $\mathcal{C}$  such that [2, Assumption 4.1] holds. Take an object  $B \overset{s}{\underset{t}{\rightleftarrows}} A$  of the category  $\text{ReflGraphMon}_{\mathcal{S}}(\mathcal{C})$  of Theorem 2.1 such that the following properties hold.*

- $B \overset{t}{\leftarrow} A = A$  belongs to  $\mathcal{S}$
- the morphism  $q_3$  of Lemma 3.1 (3) is invertible.

The following assertions hold.

- (1) There is at most one monoid morphism  $d$  rendering commutative

$$\begin{array}{ccc}
 A & \xrightarrow{1 \square i} & A \square_B A & \xleftarrow{i \square 1} & A \\
 & \searrow & \downarrow d & \swarrow & \\
 & & A & & 
 \end{array}$$

- (2) The monoid morphism  $d$  of part (1) exists if and only if the following diagram commutes (recall that  $q_2$  is invertible by Lemma 3.1 (3)).

$$\begin{array}{ccccccc}
 A(A \square_B I) & \xrightarrow{1p_A} & A^2 \xrightarrow{(i \square 1)(1 \square i)} & (A \square_B A)^2 & \xrightarrow{m} & A \square_B A & \xrightarrow{q_2^{-1}} & (A \square_B I)A \\
 1p_A \downarrow & & & & & & & \downarrow p_{A^1} \\
 A^2 & \xrightarrow{m} & & A & \xleftarrow{m} & A^2 & & 
 \end{array}$$

Moreover, in this case  $d$  is equal to  $A \square_B A \xrightarrow{q_2^{-1}} (A \square_B I)A \xrightarrow{p_{A^1}} A^2 \xrightarrow{m} A$ .

- (3) Whenever the monoid morphism  $d$  of part (1) exists,  $B \overset{s}{\underset{t}{\rightleftarrows}} A \overset{d}{\longleftarrow} A \square_B A$  is an  $\mathcal{S}$ -relative category in the category of monoids in  $\mathcal{C}$ .

*Proof.* The proof is built on [2, Corollary 1.7].



(1) Since the morphism  $q_2$  in Lemma 3.1 (3) is invertible, we know from [2, Corollary 1.7] that there is at most one monoid morphism rendering commutative

$$\begin{array}{ccccc}
 A \square_B I & \xrightarrow{p_A} & A & \xrightarrow{1 \square i} & A \square_B A & \xleftarrow{i \square 1} & A \\
 & \searrow & & & \downarrow & & \swarrow \\
 & & & & A & & \\
 & \swarrow & & & \downarrow & & \searrow \\
 & & & & A & & 
 \end{array}
 \quad (3.13)$$

Since a monoid morphism  $d$  as in part (1) obviously renders commutative (3.13), this proves its uniqueness.

(2) By [2, Corollary 1.7] commutativity of the diagram of part (2) is equivalent to the existence of a (unique) monoid morphism making (3.13) commute. Since a monoid morphism  $d$  in part (1) provides such a morphism, its existence implies commutativity of the diagram of part (2).

In order to prove the converse implication, we show that any monoid morphism  $d$  making (3.13) commute renders commutative also the diagram of part (1). Recall from [2, Lemma 1.2] that the invertibility of  $q$  in Theorem 2.1 (b) implies that  $p_A$  and  $i$  are joint epimorphisms of monoids. Hence if  $d$  makes (3.13) commute then it does so the left hand side of the diagram of part (1) by  $d.(1 \square i).i = d.(i \square 1).i = i$ .

The stated expression of  $d$  immediately follows from [2, Corollary 1.7].

(3) In order to see that the monoid morphism  $d$  in part (1) is a morphism of spans, we use that by the invertibility of  $q_2$  there are unique morphisms rendering commutative the respective diagrams

$$\begin{array}{ccc}
 A \square_B I & \xrightarrow{p_A} & A & \xrightarrow{1 \square i} & A \square_B A & \xleftarrow{i \square 1} & A \\
 & \searrow & & & \downarrow & & \swarrow \\
 & & & & B & & \\
 & \swarrow & & & \downarrow & & \searrow \\
 & & & & B & & 
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 A \square_B I & \xrightarrow{p_A} & A & \xrightarrow{1 \square i} & A \square_B A & \xleftarrow{i \square 1} & A \\
 & \searrow & & & \downarrow & & \swarrow \\
 & & & & B & & \\
 & \swarrow & & & \downarrow & & \searrow \\
 & & & & B & & 
 \end{array}$$

see [2, Corollary 1.7]. Now  $s.d$  obviously makes the first diagram commute and so does  $A \square_B A \xrightarrow{p_2} A \xrightarrow{s} B$  by the commutativity of

$$\begin{array}{ccc}
 A \square_B I & \xrightarrow{p_A} & A & \xrightarrow{1 \square i} & A \square_B A \\
 \downarrow s & & \downarrow p_2 & & \downarrow p_2 \\
 B & \xrightarrow{i} & A & & A \\
 \downarrow s & & \downarrow s & & \downarrow s \\
 B & & B & & B
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 A & \xrightarrow{i \square 1} & A \square_B A \\
 \downarrow p_2 & & \downarrow p_2 \\
 A & & A \\
 \downarrow s & & \downarrow s \\
 B & & B
 \end{array}$$

Thus they are equal. Similarly, both  $t.d$  and  $A \square_B A \xrightarrow{p_1} A \xrightarrow{t} B$  render commutative the second diagram proving that they are equal.

The to-be composition morphism  $d$  in part (1) admits the unit  $i$  by construction. Its associativity follows again by [2, Corollary 1.7] since by the invertibility of  $q_3$  there is at most one morphism rendering commutative

$$\begin{array}{ccccc}
 A \square_B I & \xrightarrow{p_A} & A & \xrightarrow{1 \square i \square i} & A \square_B A \square_B A & \xleftarrow{i \square 1} & A \square_B A \\
 & \searrow & & & \downarrow & & \swarrow \\
 & & & & A & & \\
 & \swarrow & & & \downarrow & & \searrow \\
 & & & & A & & 
 \end{array}$$

Since both  $d.(d \square 1)$  and  $d.(1 \square d)$  do so by the commutativity of

$$\begin{array}{c}
 \begin{array}{ccc}
 A \square_B I & \xrightarrow{p_A} & A \xrightarrow{1 \square i \square i} A \square_B^3 \\
 & & \downarrow d \square 1 \\
 & & A \square_B A \\
 & & \downarrow d \\
 & & A
 \end{array}
 &
 \begin{array}{ccc}
 A \square_B I & \xrightarrow{p_A} & A \xrightarrow{1 \square i \square i} A \square_B^3 \\
 & & \downarrow 1 \square d \\
 & & A \square_B A \\
 & & \downarrow d \\
 & & A
 \end{array}
 &
 \begin{array}{ccc}
 A \square_B A & \xrightarrow{i \square 1 \square 1} & A \square_B^3 \\
 & & \downarrow d \square 1 \\
 & & A \square_B A \\
 & & \downarrow d \\
 & & A
 \end{array}
 &
 \begin{array}{ccc}
 A \square_B A & \xrightarrow{i \square 1 \square 1} & A \square_B^3 \\
 \downarrow d & & \downarrow 1 \square d \\
 A & \xrightarrow{i \square 1} & A \square_B A \\
 & & \downarrow d \\
 & & A
 \end{array}
 \end{array}$$

this proves their equality (modulo the omitted associativity isomorphism in [2, Proposition 3.6]).  $\square$

**Proposition 3.9.** *Consider a monoidal admissible class  $\mathcal{S}$  of spans in a monoidal category  $\mathbf{C}$  such that [2, Assumption 4.1] holds. Between  $\mathcal{S}$ -relative categories in the category of monoids in  $\mathbf{C}$  for which the morphisms  $q_2$  in Lemma 3.1 (3) are invertible, any morphism of reflexive graphs of monoids is in fact an  $\mathcal{S}$ -relative functor.*

*Proof.* Take  $\mathcal{S}$ -relative categories  $B \xrightleftharpoons[t]{s} A \xleftarrow{d} A \square_B A$  and  $B' \xrightleftharpoons[t']{s'} A' \xleftarrow{d'} A' \square_{B'} A'$  as in the claim. We need to check the compatibility of any morphism of reflexive graphs  $(B \xrightarrow{b} B', A \xrightarrow{a} A')$  with the composition morphisms  $d$  and  $d'$ . The first diagram of

$$\begin{array}{ccc}
 \begin{array}{ccccc}
 (A \square_B I)A & \xrightarrow{p_{A^1}} & A^2 & \xrightarrow{(1 \square i)(i \square 1)} & (A \square_B A)^2 & \xrightarrow{m} & A \square_B A \\
 \downarrow (a \square 1)a & & \downarrow aa & & \downarrow (a \square a)(a \square a) & & \downarrow a \square a \\
 (A' \square_{B'} I)A' & \xrightarrow{p_{A'^1}} & A'^2 & \xrightarrow{(1 \square i')(i' \square 1)} & (A' \square_{B'} A')^2 & \xrightarrow{m'} & A' \square_{B'} A' \\
 & & \downarrow q'_2 & & & & \downarrow q'_2
 \end{array}
 &
 \begin{array}{ccccc}
 A \square_B A & \xrightarrow{q_2^{-1}} & (A \square_B I)A & \xrightarrow{p_{A^1}} & A^2 & \xrightarrow{m} & A \\
 \downarrow a \square a & & \downarrow (a \square 1)a & & \downarrow aa & & \downarrow a \\
 A' \square_{B'} A' & \xrightarrow{q_2'^{-1}} & (A' \square_{B'} I)A' & \xrightarrow{p_{A'^1}} & A'^2 & \xrightarrow{m'} & A' \\
 & & \downarrow d' & & & & \downarrow d'
 \end{array}
 \end{array}$$

commutes since  $a \square a$  is multiplicative by [2, Proposition 3.7 (2)] and by the functoriality of  $\square$ ; see [2, Proposition 3.5 (2)]. It is used to prove the commutativity of the second diagram.  $\square$

### 3.3. The equivalence between relative categories and crossed modules of monoids.

**Theorem 3.10.** *Consider a monoidal admissible class  $\mathcal{S}$  of spans in a monoidal category  $\mathbf{C}$  such that [2, Assumption 4.1] holds. Use the same notation  $\mathcal{S}$  for the induced admissible class of spans in the category of monoids in  $\mathbf{C}$  from [2, Example 2.8] (also satisfying [2, Assumption 4.1] by [2, Example 4.4]). The following categories are equivalent.*

$\text{CatMon}_{\mathcal{S}}(\mathbf{C})$  whose

objects are  $\mathcal{S}$ -relative categories  $B \xrightleftharpoons[t]{s} A \xleftarrow{d} A \square_B A$  in the category of monoids

in  $\mathbf{C}$  such that the morphisms  $q_n$  of (3.3) are invertible for any positive integer  $n$ .

morphisms are  $\mathcal{S}$ -relative functors in the category of monoids in  $\mathbf{C}$ .

$\mathbf{Xmod}_{\mathcal{S}}(\mathbf{C})$  whose

objects consist of monoids  $B$  and  $Y$ , monoid morphisms  $Y \xrightarrow{e} I$  and  $Y \xrightarrow{k} B$  and a distributive law  $BY \xrightarrow{x} YB$  subject to the following conditions.

- (a')  $B \xleftarrow{k} Y = Y \in \mathcal{S}$ ,  $Y = Y \xrightarrow{e} I \in \mathcal{S}$  and  $B = B = B \in \mathcal{S}$ .
- (b')  $e1.x = 1e$  and  $m.k1.x = m.1k$ .
- (c') The morphism  $f$  of Theorem 1.1 (c') is invertible and the morphisms  $b_n$  of Lemma 3.4 (2) are invertible for all positive integers  $n$ .
- (d') Regarding  $YB$  as a monoid via the structure induced by the distributive law  $x$ , the following diagram commutes.

$$\begin{array}{ccccc}
 YBY & \xrightarrow{u111uu} & (Y^2B)^2 & \xrightarrow{b_2b_2} & (YB \square_B YB)^2 & \xrightarrow{m} & YB \square_B YB \\
 \downarrow 1x & & & & & & \downarrow b_2^{-1} \\
 Y^2B & \xrightarrow{m1} & YB & \xleftarrow{m1} & Y^2B & & 
 \end{array}$$

morphisms are pairs of monoid morphisms  $(B \xrightarrow{b} B', Y \xrightarrow{y} Y')$  such that  $e'.y = e$ ,  $k'.y = b.k$  and  $x'.by = yb.x$ .

*Proof.* It follows by Proposition 3.8 and Proposition 3.9 that  $\mathbf{CatMon}_{\mathcal{S}}(\mathbf{C})$  is a full subcategory of  $\mathbf{ReflGraphMon}_{\mathcal{S}}(\mathbf{C})$  and obviously  $\mathbf{Xmod}_{\mathcal{S}}(\mathbf{C})$  is a full subcategory of  $\mathbf{PreX}_{\mathcal{S}}(\mathbf{C})$ . Below we show that the mutually inverse functors of Theorem 2.1 restrict to functors between these subcategories thus establishing the stated equivalence.

Regarding an object  $B \xrightleftharpoons[t]{s} A \xleftarrow{d} A \square_B A$  of  $\mathbf{CatMon}_{\mathcal{S}}(\mathbf{C})$  as an object  $B \xrightleftharpoons[t]{s} A$  of  $\mathbf{ReflGraphMon}_{\mathcal{S}}(\mathbf{C})$ , the functor in the proof of Theorem 2.1 takes it to the object  $(B, A \square_B I, A \square_B I \xrightarrow{pI} I, A \square_B I \xrightarrow{pA} A \xrightarrow{t} B, B(A \square_B I) \xrightarrow{ipA} A^2 \xrightarrow{m} A \xrightarrow{q^{-1}} (A \square_B I)B)$  of the category  $\mathbf{PreX}_{\mathcal{S}}(\mathbf{C})$ ; we claim that it is in fact an object of  $\mathbf{Xmod}_{\mathcal{S}}(\mathbf{C})$ .

It satisfies the condition  $B \xleftarrow{t} A \xleftarrow{pA} A \square_B I = A \in \mathcal{S}$  by Lemma 2.2.

From Lemma 3.4 (5) we know that the morphism  $b_n$  of Lemma 3.4 (2) is invertible if and only if the left column of the commutative diagram

$$\begin{array}{ccc}
 ((A \square_B I)B \square_B I)((A \square_B I)B) \square_B^{n-1} & \xrightarrow{(q \square 1)q \square^{n-1}} & (A \square_B I)A \square_B^{n-1} \\
 \downarrow p(A \square_B I)B^1 & & \downarrow pA^1 \\
 (A \square_B I)B((A \square_B I)B) \square_B^{n-1} & \xrightarrow{qq \square^{n-1}} & AA \square_B^{n-1} \\
 \downarrow (1 \square u 1 \square \dots \square u 1)(u 1 \square 1) & & \downarrow (1 \square i \square \dots \square i)(i \square 1) \\
 (((A \square_B I)B) \square_B^n)^2 & \xrightarrow{q \square^n q \square^n} & (A \square_B^n)^2 \\
 \downarrow m & & \downarrow m \\
 ((A \square_B I)B) \square_B^n & \xrightarrow{q \square^n} & A \square_B^n
 \end{array}$$

$\curvearrowright q_n$

is invertible. Recognize the isomorphism  $q_n$  in the right column. Since also the rows are isomorphisms by assumption, so is the left column and hence  $b_n$ .

The proof of the commutativity of the diagram in part (d') requires some preparation. The commutativity of

$$\begin{array}{c}
 \begin{array}{ccccc}
 & & & & (A \square_B I) B \square_B (A \square_B I) B \xrightarrow{q \square q} A \square_B A \\
 & & & & \downarrow p_1 \\
 (A \square_B I)^2 B & \xrightarrow{1 p_A 1} & (A \square_B I) A B = (A \square_B I) A B \xrightarrow{1 t 1} & (A \square_B I) B^2 \xrightarrow{1 m} & (A \square_B I) B \xrightarrow{q} & A \\
 & & \searrow 1 1 i & \uparrow 1 1 t & \uparrow 1 t & \downarrow p_1 \\
 & & (A \square_B I) A^2 & & & A \\
 & & \searrow 1 m & & & \uparrow p_1 \\
 & & & & (A \square_B I) A \xrightarrow{q_2} & A \square_B A \\
 & & & & \uparrow 1 q & \\
 & & & & (A \square_B I) A & \\
 & & & & \downarrow 1 q & \\
 & & & & (A \square_B I) B \square_B (A \square_B I) B & \xrightarrow{q \square q} & A \square_B A \\
 & & & & \downarrow p_2 & & \downarrow p_2 \\
 (A \square_B I)^2 B & \xrightarrow{p_I 1 1} & (A \square_B I) B & \xrightarrow{q} & A & & A \\
 & & \searrow 1 q & \nearrow p_I 1 & \uparrow p_2 & & \\
 & & (A \square_B I) A & \xrightarrow{q_2} & A \square_B A & & 
 \end{array}
 \end{array}$$

proves  $(q \square q).b_2 = q_2.1q$ . (Here the bottom-right region of the first diagram commutes since the lower half of the diagram of (3.8) commutes and the bottom-right region of the second diagram commutes since the lower half of the diagram of (3.9) commutes.)

By the associativity of  $A$  and the multiplicativity of  $A \square_B I \xrightarrow{p_A} A$  also the following diagram commutes.

$$\begin{array}{ccccc}
 & & & & 1q \\
 (A \square_B I)^2 B & \xrightarrow{1 p_A i} & (A \square_B I) A^2 & \xrightarrow{1 m} & (A \square_B I) A \\
 \downarrow m 1 & & \downarrow p_A 1 1 & & \downarrow p_A 1 \\
 & & A^3 & \xrightarrow{1 m} & A^2 \\
 & & \downarrow m 1 & & \downarrow m \\
 (A \square_B I) B & \xrightarrow{p_A i} & A^2 & \xrightarrow{m} & A \\
 & & \searrow q & & \\
 & & & & 
 \end{array}$$

With the help of these identities and Lemma 3.7, and using that the region marked by (\*) commutes by Proposition 3.8 (2), the diagram of Figure 5 is seen to commute. This proves that the stated object belongs to  $\mathbf{Xmod}_S(\mathbf{C})$  indeed.

In the opposite direction, consider an object  $(B, Y, Y \xrightarrow{e} I, Y \xrightarrow{k} B, BY \xrightarrow{x} YB)$  of  $\mathbf{Xmod}_S(\mathbf{C})$  as an object of  $\mathbf{PreX}_S(\mathbf{C})$ . The functor in the proof of Theorem 2.1 takes it to the object  $B \begin{array}{c} \xleftarrow{e 1} \\ \xrightarrow{u 1} \\ \xleftarrow{m, k 1} \end{array} A$  of  $\mathbf{ReflGraphMon}_S(\mathbf{C})$ ; we claim that it can be seen as an object of  $\mathbf{CatMon}_S(\mathbf{C})$ .

By Lemma 3.4 (1) the span  $B \xleftarrow{m} B^2 \xleftarrow{k 1} YB = YB$  belongs to  $\mathcal{S}$ .

The morphism  $q_n$  of Lemma 3.4 (4) is invertible for all positive integers  $n$  by Lemma 3.4 (5).




FIGURE 5. Commutativity of the diagram in part (d')

By Proposition 3.8 (2) and (3), the reflexive graph of monoids  $B \begin{smallmatrix} \xrightarrow{e1} \\ \xleftrightarrow[u1]{m.k1} \\ \xrightarrow{m.k1} \end{smallmatrix} A$  extends to an  $\mathcal{S}$ -relative category in the category of monoids in  $\mathbf{C}$  by the commutativity of

$$\begin{array}{c}
 \begin{array}{ccccccc}
 YB(YB \square_B I) & \xrightarrow{11p_{YB}} & (YB)^2 & \xrightarrow{(u1 \square 1)(1 \square u1)} & (YB \square_B YB)^2 & \xrightarrow{m} & YB \square_B YB & \xrightarrow{q_2^{-1}} & (YB \square_B I) YB \\
 \downarrow 11p_{YB} & \searrow 11f^{-1} & \uparrow 111u & \searrow u111u1 & \nearrow b_2 b_2 & & \downarrow b_2^{-1} & & \downarrow f^{-1} 11 \\
 & & YBY & & (Y^2 B)^2 & & & & \\
 & \nearrow 111u & \downarrow 1x & & & & & & \\
 & & Y^2 B & & & & & & \\
 & & \uparrow 11m & \searrow m1 & & & & & \\
 (YB)^2 & \xrightarrow{1x1} & Y^2 B^2 & \xrightarrow{mm} & YB & \xleftarrow{mm} & Y^2 B^2 & \xleftarrow{1x1} & (YB)^2.
 \end{array} \\
 \text{(d')}
 \end{array}$$

The region at the top-right corner is the commutative diagram of Lemma 3.4 (4) for  $n = 1$ . The region bounded from below by the curved arrows commutes by Lemma 3.7. The region marked by (d') coincides with the diagram of part (d') hence it commutes.  $\square$

**Example 3.11.** As in Example 1.2, take the (evidently admissible and monoidal) class of all spans in the category  $\mathbf{C}$  of spans over a given set  $X$ . Then the equivalent categories of Theorem 3.10 take the following forms.

$\text{CatMon}(\mathbf{C})$  whose

objects are the double categories with the object set  $X$  and only identity horizontal morphisms and such that the morphism (1.3) is invertible. (This last condition holds e.g. if the vertical edge category is a groupoid.)

morphisms are the double functors which are identities on the objects (and hence on the horizontal morphisms).

$\text{Xmod}(\mathbf{C})$  whose

objects consist of categories  $B$  and  $Y$  with the common object set  $X$  such that in  $Y$  there are no morphisms between different objects; an action (see Example 1.2)  $B \square_X Y \xrightarrow{b} Y$  and an identity-on-objects functor  $Y \xrightarrow{\kappa} B$  such that

$$\kappa(b \triangleright y).b = b.\kappa(y) \quad \text{and} \quad (\kappa(y) \triangleright y').y = y.y'$$

for all morphisms  $b$  in  $B$  and  $y, y'$  in  $Y$  for which  $s(b) = t(y) = t(y')$ .

morphisms are the same as the morphisms in  $\text{PreXMon}(\mathbf{C})$ , see Example 2.3.

These equivalent categories have equivalent full subcategories in whose objects the occurring category  $B$  is a groupoid; and other equivalent full subcategories in whose objects both occurring categories are groupoids. In the latter case these are the category of categories in the category of groupoids; and the category of *crossed modules of groupoids* in [4, Definition 1.2], respectively.

**Example 3.12.** In the setting of Example 1.3, the equivalent categories of Theorem 3.10 take the following explicit forms.

$\text{CatMon}_{\mathcal{S}}(\mathbf{C})$  whose

objects are  $\mathcal{S}$ -relative categories  $B \begin{smallmatrix} \xleftarrow{s} \\ \xleftrightarrow{i} \\ \xrightarrow{t} \end{smallmatrix} A \xleftarrow{d} A \square_B A$  in the category of monoids in  $\mathbf{C}$  — that is, in the category of bimonoids in  $\mathbf{M}$  — such that the morphism  $q$  of Theorem 1.1 (b) is invertible.

morphisms are  $\mathcal{S}$ -relative functors in the category of monoids in  $\mathbf{C}$  — that is, in the category of bimonoids in  $\mathbf{M}$ .

$\text{Xmod}_{\mathcal{S}}(\mathbf{C})$  whose

objects consist of a bimonoid  $Y$  and a cocommutative bimonoid  $B$  together with a left action  $BY \xrightarrow{l} Y$  which makes  $Y$  both a  $B$ -module monoid and a  $B$ -module comonoid and a bimonoid morphism  $Y \xrightarrow{k} B$  for which the following diagrams commute.

$$\begin{array}{ccccc} Y & \xrightarrow{\delta} & Y^2 & \xrightarrow{c} & Y^2 & & BY & \xrightarrow{\delta_1} & B^2Y & \xrightarrow{1c} & BYB & \xrightarrow{l_1} & YB & & Y^2 & \xrightarrow{\delta_1} & Y^3 & \xrightarrow{1c} & Y^3 & \xrightarrow{k_{11}} & BY^2 \\ \delta \downarrow & & & & \downarrow k_1 & & 1k \downarrow & & & & & & \downarrow k_1 & & \parallel & & & & & & \downarrow l_1 \\ Y^2 & \xrightarrow{k_1} & BY & & & & B^2 & \xrightarrow{m} & B & \xleftarrow{m} & B^2 & & & & Y^2 & \xrightarrow{m} & Y & \xleftarrow{m} & Y^2 & & \end{array}$$

The third condition appears in [16, Definition 12 (v)] under the name *Peiffer condition* (motivated by the terminology for groups).

morphisms are pairs of monoid morphisms  $(B \xrightarrow{b} B', Y \xrightarrow{y} Y')$  such that  $e'.y = e, k'.y = b.k$  and  $x'.by = yb.x$ .

These equivalent categories are equivalent furthermore to the full subcategory of  $\text{RefGraphMon}_{\mathcal{S}}(\mathbf{C})$  of Example 2.3 for whose objects  $B \begin{smallmatrix} \xleftarrow{s} \\ \xleftrightarrow{i} \\ \xrightarrow{t} \end{smallmatrix} A$  the following diagrams commute.

$$\begin{array}{ccc} A & \xrightarrow{\delta} & A^2 & \xrightarrow{c} & A^2 & & A^2 & \xrightarrow{t_1} & BA \\ \delta \downarrow & & & & & & \downarrow t_1 \\ A^2 & \xrightarrow{t_1} & & & & & & & \end{array} \quad (3.14)$$

$$\begin{array}{ccccccccccc} A(A \square_B I) & \xrightarrow{\delta_1} & A^2(A \square_B I) & \xrightarrow{tc} & B(A \square_B I)A & \xrightarrow{ip_{A^1}} & A^3 & \xrightarrow{m_1} & A^2 & \xrightarrow{q^{-1}_1} & (A \square_B I)BA \\ 1p_A \downarrow & & & & & & & & & & \downarrow p_{A \in B^1} \\ A^2 & \xrightarrow{m} & & & A & \xleftarrow{m} & & & A^2 & & \end{array} \quad (3.15)$$

The above description of  $\text{CatMon}_{\mathcal{S}}(\mathbf{C})$  requires no further explanation. In the description of  $\text{Xmod}_{\mathcal{S}}(\mathbf{C})$  we need to show that the third diagram (the Peiffer condition) is equivalent to the diagram of Theorem 3.10 (d') in the current setting. The path on the right hand side of the diagram of Theorem 3.10 (d') appears as the left bottom

path of the commutative diagram

$$\begin{array}{ccccc}
YBY & \xrightarrow{\delta_{YB}1} & (YB)^2Y & \xrightarrow{k1111} & B^2YBY \\
u111uu \downarrow & & \downarrow u11u111uu & & \swarrow u11111uu \\
(Y^2B)^2 & \xrightarrow{\delta_{Y^2B}111} & (Y^2B)^3 & \xrightarrow{1k1\varepsilon_Y11111} & YB^2YBY^2B \\
b_2b_2 \downarrow & & \downarrow jb_2^{-1} & & \downarrow m111 \\
(YB \square_B YB)^2 & \xrightarrow{jj} & (YB)^4 & \xrightarrow{11111\varepsilon_B11} & (YB)^2Y^2B \xleftarrow{u1111uu} (BY)^2 \\
\downarrow m & & \downarrow 11c_{YB,YB}11 & & \downarrow 1c_{YB,Y} \\
& & (YB)^4 & \xrightarrow{1\varepsilon_B1111} & Y^3BYB \xleftarrow{u111uu} BY^2B \\
& & \downarrow 1x11x1 & & \downarrow l11 \\
& & (Y^2B^2)^2 & \xrightarrow{11\varepsilon_B\varepsilon_B1111} & Y^4B^2 \xleftarrow{u11u1u} Y^2B \\
& & \downarrow mmmm & & \downarrow mmm \\
YB \square_B YB & \xrightarrow{j} & (YB)^2 & \xrightarrow{1\varepsilon_B11} & Y^2B \xrightarrow{m1} YB \\
& & & & \downarrow b_2^{-1}
\end{array}$$

(in which  $x$  stands for the distributive law  $BY \xrightarrow{\delta_1} B^2Y \xrightarrow{1c} BYB \xrightarrow{l1} YB$  of Example 1.3). Hence it can be replaced by the top right path yielding the equivalent form

$$\begin{array}{ccc}
YBY & \xrightarrow{\delta_{YB}1} & (YB)^2Y & \xrightarrow{k1111} & B^2YBY & \xrightarrow{m111} & (BY)^2 & (3.16) \\
1\delta_B1 \downarrow & & & & & & \downarrow 1c_{YB,Y} \\
YB^2Y & & & & & & BY^2B \\
11c \downarrow & & & & & & \downarrow l11 \\
(YB)^2 & & & & & & Y^2B \\
1l1 \downarrow & & & & & & \\
Y^2B & \xrightarrow{m1} & YB & \xleftarrow{m1} & Y^2B
\end{array}$$

of the diagram of Theorem 3.10 (d'). The first diagram of Figure 6 shows that if the diagram of (3.16) commutes then the Peiffer condition in the above presentation of  $\mathbf{Xmod}_S(\mathcal{C})$  holds. The opposite implication is proven by the second diagram of Figure 6.

In order to justify the further equivalent characterization of these categories as a full subcategory of  $\mathbf{RefGraphMon}_S(\mathcal{C})$ , we need to see the equivalence of the diagram of Proposition 3.8 (2) in the current setting and the diagram of (3.15). This follows by noting that the top row of the diagram of Proposition 3.8 (2) in the current setting







FIGURE 6. Derivation of the Peiffer condition for crossed modules of binonoids

appears in the left-bottom path of the commutative diagram

$$\begin{array}{ccccc}
A(A \square_B I) & \xrightarrow{\delta 1} & A^2(A \square_B I) & \xrightarrow{1c} & A(A \square_B I)A \\
\downarrow 1p_A & \searrow \delta\delta & \downarrow 11p_A & \searrow 11p_A & \downarrow t11 \\
A^2 & \xrightarrow{\delta\delta} & A^2(A \square_B I)^2 & \xrightarrow{11p_{API}} & A^3 & \xrightarrow{1c} & A^3 & \xrightarrow{1p_{A1}} & B(A \square_B I)A \\
\downarrow (i \square 1)(1 \square i) & & \downarrow 11p_{APA} & & \downarrow t11 & & \downarrow t11 & & \downarrow 1p_{A1} \\
(A \square_B A)^2 & \xrightarrow{jj} & A^4 & \xrightarrow{111u} & BA^2 & \xrightarrow{1c} & A^3 & \xrightarrow{111u} & A^3 \\
\downarrow m & & \downarrow t11s & & \downarrow i11 & & \downarrow i11 & & \downarrow m1 \\
A \square_B A & \xrightarrow{q^{-1} \square 1} & (A \square_B I)B \square_B A & \xrightarrow{j} & (A \square_B I)BA & \xrightarrow{j} & (A \square_B I)BA & \xrightarrow{1\varepsilon_B 1} & (A \square_B I)A \\
& \searrow q_2^{-1} & \searrow h_1^{-1} & & \searrow h_1^{-1} & & \searrow h_1^{-1} & & \searrow h_1^{-1} \\
& & & & & & & & (A \square_B I)A
\end{array}$$

hence it can be replaced by the top-right path. (The expression of  $h_1^{-1}$  in the bottom-right corner was computed in Example 3.3.)

**Proposition 3.13.** *The equivalent categories of Example 3.12 have equivalent full subcategories as follows.*

- The full subcategory of  $\text{CatMon}_S(\mathcal{C})$  for whose objects  $B \begin{smallmatrix} \xleftarrow{s} \\ \xrightarrow{i} \\ \xrightarrow{t} \end{smallmatrix} A \xleftarrow{d} A \square_B A$  the bimonoid  $B$  in  $\mathcal{M}$  is a Hopf monoid.
- The full subcategory of  $\text{Xmod}_S(\mathcal{C})$  for whose objects  $(B, Y, BY \xrightarrow{l} Y, Y \xrightarrow{k} B)$  the bimonoid  $B$  in  $\mathcal{M}$  is a Hopf monoid.
- The full subcategory of  $\text{ReflGraphMon}_S(\mathcal{C})$  for whose objects  $B \begin{smallmatrix} \xleftarrow{s} \\ \xrightarrow{i} \\ \xrightarrow{t} \end{smallmatrix} A$  the following conditions hold.
  - $B$  is a Hopf monoid (with antipode  $z$ )
  - $t1.\delta = t1.c.\delta$
  - for the morphisms

$$\overrightarrow{s} := A \xrightarrow{\delta} A^2 \xrightarrow{1s} AB \xrightarrow{1z} AB \xrightarrow{1i} A^2 \xrightarrow{m} A, \quad \overleftarrow{t} := A \xrightarrow{\delta} A^2 \xrightarrow{t1} BA \xrightarrow{z1} BA \xrightarrow{i1} A^2 \xrightarrow{m} A$$

the following diagram commutes.

$$\begin{array}{ccc}
A^2 & \xrightarrow{\overrightarrow{s} \overleftarrow{t}} & A^2 & \xrightarrow{c} & A^2 \\
\overrightarrow{s} \overleftarrow{t} \downarrow & & & & \downarrow m \\
A^2 & \xrightarrow{m} & A & & 
\end{array} \tag{3.17}$$

*Proof.* The only ingredient that requires a proof is the equivalence of diagrams (3.15) and (3.17) in the case when  $B$  has an antipode  $z$ . The proof will repeatedly use the identity on  $\vec{s}$  encoded in the following commutative diagram.

$$\begin{array}{ccccccc}
 A^2 & \xrightarrow{m} & & & & & A \\
 \downarrow 1\vec{s} & \searrow \delta 1 & & & & & \downarrow \delta \\
 A^2 & \xrightarrow{11\delta} & A^3 & \xrightarrow{1c1} & A^4 & \xrightarrow{mm} & A^2 \\
 \downarrow \delta 1 & \searrow 11\vec{s} & \downarrow 1c & & \downarrow 11c & & \downarrow 1s \\
 A^3 & \xrightarrow{1c} & A^3 & \xrightarrow{1\delta 1} & A^4 & \xrightarrow{mm} & AB \\
 & & \downarrow 1\vec{s}1 & & \downarrow 11c & & \downarrow 1z \\
 & & & & A^2B^2 & \xrightarrow{mm} & AB \\
 & & & & \downarrow 11zz & & \downarrow 1i \\
 & & & & A^2B^2 & \xrightarrow{mm} & AB \\
 & & & & \downarrow 11ii & & \downarrow 1i \\
 & & & & A^4 & \xrightarrow{mm} & A^2 \\
 & & & & \downarrow 1m1 & & \downarrow m \\
 A^3 & \xrightarrow{1c} & A^3 & \xrightarrow{11s} & A^2B & \xrightarrow{11z} & A^2B & \xrightarrow{11i} & A^3 & \xrightarrow{m1} & A^2 & \xrightarrow{m} & A
 \end{array} \tag{3.18}$$

Recall from [15] that if  $B$  has an antipode  $z$  then  $A \square_B I \xrightarrow{p_A} A$  is a split monomorphism in  $\mathbf{M}$ ; a retraction is provided by  $g_A := A \xrightarrow{q^{-1}} (A \square_B I)B \xrightarrow{1\varepsilon} A \square_B I$ . Indeed,

$$g_A \cdot p_A = 1\varepsilon \cdot q^{-1} \cdot q \cdot 1u = 1\varepsilon \cdot 1u = 1.$$

On the other hand, since in Proposition 1.4  $q^{-1}$  was constructed as the unique solution of  $p_A 1 \cdot q^{-1} = \vec{s} s \cdot \delta$ , also the equality

$$p_A \cdot g_A = p_A \cdot 1\varepsilon \cdot q^{-1} = 1\varepsilon \cdot p_A 1 \cdot q^{-1} = 1\varepsilon \cdot \vec{s} s \cdot \delta = \vec{s}$$

holds, proving that  $\vec{s}$  is idempotent.

Pre-composing both paths around (3.15) with the split epimorphism  $1g_A$ , we obtain the equivalent diagram

$$\begin{array}{ccccccccccc}
 A^2 & \xrightarrow{\delta 1} & A^3 & \xrightarrow{1c} & A^3 & \xrightarrow{t11} & BA^2 & \xrightarrow{i11} & A^3 & \xrightarrow{m1} & A^2 \\
 \downarrow 1g_A & & \downarrow 11g_A & & & & \downarrow 1g_A 1 & & \downarrow 1\vec{s}1 & & \downarrow \vec{s}1 \\
 A(A \square_B I) & \xrightarrow{\delta 1} & A^2(A \square_B I) & \xrightarrow{1c} & A(A \square_B I)A & \xrightarrow{t11} & B(A \square_B I)A & \xrightarrow{ip_A 1} & A^3 & & A^2 \\
 \downarrow 1\vec{s} & & \downarrow 1p_A & & & & & & \downarrow m1 & & \downarrow \vec{s}1 \\
 A^2 & \xrightarrow{m} & & & & & & & A^2 & & A^2
 \end{array} \tag{3.19}$$

Its rightmost region commutes by (3.18) and the fact that  $\vec{s}$  is idempotent.

The morphism around the right hand side of (3.19) occurs as the left-bottom path of the commutative diagram

$$\begin{array}{ccccccc}
 A^2 & \xrightarrow{\delta 1} & A^3 & \xrightarrow{1c} & A^3 & \xrightarrow{t11} & BA^2 \\
 \delta 1 \downarrow & & \downarrow 1\delta 1 & & \downarrow 11\delta & & \downarrow i11 \\
 A^3 & \xrightarrow{\delta 11} & A^4 & & & & \\
 1c \downarrow & & \downarrow 11c & & & & \\
 A^3 & \xrightarrow{\delta 11} & A^4 & \xrightarrow{1c1} & A^4 & & \\
 t11 \downarrow & & \downarrow tt11 & & \downarrow t1t1 & & \\
 BA^2 & & B^2A^2 & & (BA)^2 & & \\
 i11 \downarrow & & \downarrow ii11 & & \downarrow i1i1 & & \\
 A^3 & \xrightarrow{\delta 11} & A^4 & \xrightarrow{1c1} & A^4 & & \\
 m1 \downarrow & & & & \downarrow 11s1 & & \\
 A^2 & & & & A^2BA & \xleftarrow{11t1} & A^4 \xleftarrow{11\delta} A^3 \\
 \downarrow \overrightarrow{s}1 & & & & \downarrow 11z1 & & \downarrow 11\overleftarrow{t} \\
 A^2 & & & & A^2BA & \xrightarrow{11i1} & A^4 \xrightarrow{11m} A^3 \\
 & & & & \downarrow 1\overrightarrow{s}11 & & \downarrow 1\overrightarrow{s}1 \\
 & & & & A^4 & \xrightarrow{11m} & A^3 \\
 & & & & \swarrow m11 & & \downarrow 1m \\
 & & & & A^3 & & A^2 \\
 & & & & \swarrow m1 & & \downarrow m \\
 A^2 & \xrightarrow{m} & A & \xleftarrow{m} & A^2 & & 
 \end{array}
 \quad (3.18)$$

Hence it can be replaced by the top-right path yielding the equivalent form

$$\begin{array}{ccccccc}
 A^2 & \xrightarrow{\delta 1} & A^3 & \xrightarrow{t11} & BA^2 & \xrightarrow{i11} & A^3 \xrightarrow{1c} A^3 \xrightarrow{1\overrightarrow{s}\overleftarrow{t}} A^3 \\
 1\overrightarrow{s} \downarrow & & & & & & \downarrow 1m \\
 A^2 & \xrightarrow{m} & A & \xleftarrow{m} & A^2 & & 
 \end{array}
 \quad (3.20)$$


of (3.19).

Finally, observe that for any morphisms  $A^2 \xrightarrow{\phi, \psi} A$  the following diagrams are equivalent:


$$\begin{array}{ccc}
 A^2 \xrightarrow{\delta 1} A^3 \xrightarrow{t11} BA^2 \xrightarrow{i11} A^3 & & A^2 \xrightarrow{\delta 1} A^3 \xrightarrow{t11} BA^2 \xrightarrow{z11} BA^2 \xrightarrow{i11} A^3 \\
 \psi \downarrow & & \phi \downarrow \\
 A \xleftarrow{m} A^4 & & A \xleftarrow{m} A^4 \\
 & & \downarrow 1\psi \\
 & & A^4
 \end{array}
 \quad (3.21)$$

Indeed, the first diagram below shows that if the first diagram of (3.21) commutes then so does the second one; and the opposite implication follows by the second diagram

below.



Applying the equivalence of the diagrams of (3.21) to  $\phi := A^2 \xrightarrow{c} A^2 \xrightarrow{\overrightarrow{s} \overleftarrow{t}} A^2 \xrightarrow{m} A$  and  $\psi := A^2 \xrightarrow{1 \overrightarrow{s}} A^2 \xrightarrow{m} A$ , we obtain from (3.20) the equivalent form



which is equivalent to (3.17) by the naturality of the symmetry  $c$ . □

The equivalent categories of Proposition 3.13 have equivalent full subcategories in whose objects both occurring bimonoids are Hopf monoids, and other equivalent full subcategories in whose objects they are both cocommutative Hopf monoids. In this way, Proposition 3.13 includes [16, Proposition 11] and [16, Theorem 14] about the equivalence between the category of so-called  $\text{Cat}^1$ -Hopf algebras and the category of crossed modules over Hopf algebras; hence in particular the equivalence between the category of  $\text{Cat}^1$ -groups and the category of crossed modules over groups in [10, Section 3.9].

## REFERENCES

- [1] Marcelo Aguiar, *Internal Categories and Quantum Groups*, Ph.D. thesis, Cornell University 1997. [1](#)
- [2] Gabriella Böhm, *Crossed modules of monoids I. Relative categories*, preprint available at [arXiv:1803.03418](#) [1](#), [2](#), [3](#), [4](#), [5](#), [6](#), [7](#), [8](#), [9](#), [12](#), [18](#), [19](#), [20](#), [21](#), [23](#), [24](#), [25](#), [26](#), [27](#), [28](#), [29](#), [31](#), [32](#), [33](#), [34](#)
- [3] Gabriella Böhm, *Crossed modules of monoids III. Simplicial monoids of Moore length 1*, in preparation. [2](#)
- [4] Ronald Brown and İlhan İçen, *Homotopies and Automorphisms of Crossed Modules of Groupoids*, Appl. Categ. Structures 11 no. 2 (2003) 185-206. [1](#), [2](#), [8](#), [20](#), [38](#)
- [5] Ronald Brown and Christopher B. Spencer,  *$\mathcal{G}$ -groupoids, crossed modules and the fundamental groupoid of a topological group*, Indagationes Mathematicae (Proceedings) 79 no. 4 (1976) 296-302. [1](#), [46](#)
- [6] John Williford Duskin, *Preliminary remarks on groups*, as quoted in [5]: Unpublished notes, Tulane University, (1969). [1](#)
- [7] Kadir Emir, *2-Crossed Modules of Hopf Algebras*, talk given at “Category Theory 2016” Halifax, Canada. slides available at <http://mysite.science.uottawa.ca/phofstra/CT2016/slides/Emir.pdf>. [1](#)
- [8] João Faria Martins, *Crossed modules of Hopf algebras and of associative algebras and two-dimensional holonomy*, J. Geom. Phys. 99 (2016), 68-110. [1](#)
- [9] Marino Gran, Gabriel Kadjo and Joost Vercruyssen, *A torsion theory in the category of cocommutative Hopf algebras*, Appl. Categ. Structures 24 no. 3 (2016), 269-282. [17](#)
- [10] George Janelidze, *Internal crossed modules*, Georgian Math. J. 10 (2003), 99-114. [1](#), [2](#), [17](#), [45](#)
- [11] Shahn Majid, *Strict quantum 2-groups*, preprint available at [arXiv:1208.6265](#). [1](#)
- [12] Simona Paoli, *Internal categorical structures in homotopical algebra*, in: “Towards Higher Categories” J. C. Baez and J. P. May (eds.), IMA volumes in Mathematics and its applications pp 85-103, Springer 2009. [1](#)
- [13] Timothy Porter, *The Crossed Menagerie: an introduction to crossed gadgetry and cohomology in algebra and topology*, Notes initially prepared for the XVI Encuentro Rioplatense de Álgebra y Geometría Algebraica, in Buenos Aires, 12-15 December 2006, extended for an MSc course (Summer 2007) at Ottawa. available at <https://ncatlab.org/timporter/show/crossed+menagerie>. [1](#)
- [14] Timothy Porter, *Homotopy Quantum Field Theories meets the Crossed Menagerie: an introduction to HQFTs and their relationship with things simplicial and with lots of crossed gadgetry*, Notes prepared for the Workshop and School on Higher Gauge Theory, TQFT and Quantum Gravity Lisbon, February, 2011. available at <https://ncatlab.org/timporter/show/HQFTs+meet+the+Crossed+Menagerie>. [1](#)
- [15] David E. Radford, *The structure of Hopf algebras with a projection*, J. Algebra 92 no. 2 (1985), 322-347. [14](#), [43](#)
- [16] José Manuel Fernández Vilaboa, María Purificación López López and Emilo Villanueva Novoa, *Cat<sup>1</sup>-Hopf Algebras and Crossed Modules*, Commun. Algebra 35 no. 1 (2006) 181-191. [1](#), [2](#), [21](#), [39](#), [45](#)
- [17] John Henry Constantine Whitehead, *On adding relations to homotopy groups*, Ann. of Math. 42 no. 2 (1941) 409-428. [1](#)

WIGNER RESEARCH CENTRE FOR PHYSICS, H-1525 BUDAPEST 114, P.O.B. 49, HUNGARY  
*E-mail address:* bohm.gabriella@wigner.mta.hu