A system-wide network reconstruction of gene regulation and metabolism in *Escherichia coli*

Anne Grimbs¹, David F. Klosik², Stefan Bornholdt², and Marc-Thorsten $\rm H\ddot{u}tt^{*1}$

¹Computational Systems Biology, Department of Life Sciences & Chemistry, Jacobs University, Bremen, 28759, Germany ²Institute for Theoretical Physics, University of Bremen, Bremen, 28359, Germany

*m.huett@jacobs-university.de

Genome-scale metabolic models have become a fundamental tool for examining metabolic principles. However, metabolism is not solely characterized by the underlying biochemical reactions and catalyzing enzymes, but also affected by regulatory events. Since the pioneering work of Covert and co-workers as well as Shlomi and co-workers it is debated, how regulation and metabolism synergistically characterize a coherent cellular state. The first approaches started from metabolic models which were extended by the regulation of the encoding genes of the catalyzing enzymes. By now, bioinformatics databases in principle allow addressing the challenge of integrating regulation and metabolism on a systemwide level. Collecting information from several databases we provide a network representation of the integrated gene regulatory and metabolic system for *Escherichia coli*, including major cellular processes, from metabolic processes via protein modification to a variety of regulatory events. Besides transcriptional regulation, we also take into account regulation of translation, enzyme activities and reactions. Our network model provides novel topological characterizations of system components based on their positions in the network. We show that network characteristics suggest a representation of the integrated system as three network domains (regulatory, metabolic and interface networks) instead of two. This new three-domain representation reveals the structural centrality of components with known high functional relevance. This integrated network can serve as a platform for understanding coherent cellular states as active subnetworks and to elucidate crossover effects between metabolism and gene regulation.

Introduction

So far, metabolic processes and gene regulatory events are typically considered individually in systemlevel investigations. However, ample evidence exists that the majority of cellular processes involves both, metabolism and gene regulation, and thus requires their joint examination [1]. One of the bestinvestigated individual examples in *Escherichia coli* (*E. coli*) is the phosphoenolpyruvate–carbohydrate phosphotransferase system (PTS) which is responsible for import and phosphorylation of sugars [2]. Additionally, the PTS is involved in the regulation of the import process depending on the available carbohydrate mixtures in the growth medium. By carbon catabolite repression and inducer exclusion, primarily the uptake of a preferred carbon source to be metabolized, such as glucose, is selected from other carbohydrates present in the growth medium. In order to understand the underlying principles, not only the effects of both 'layers', metabolism and regulation, need to be taken into account, but also their interface [3].

On a more qualitative level, the importance of the interface of metabolism and gene regulation can be illustrated by having a closer look at their most prominent representatives, namely, enzymes and metabolic transcriptional regulators. Both examples are proteins and can be thought of as a component type organizing the interplay of genes and metabolic reactions (Figure 1). For enzymes the connection is straightforward: The majority of metabolic reactions can only take place if the corresponding genes of the catalyzing enzymes are expressed. These genes, in turn, are often involved in regulatory processes, especially if they are associated with central biochemical reactions. In contrast, metabolic transcriptional regulators can be illustrated by looking at transcription factors, the probably best-investigated transcriptional regulators. Some of them require the binding of a metabolite to be active and are therefore called metabolic transcriptional regulators. In the context of the integrative view discussed here, it is noteworthy that only the interaction with a metabolic component enables their functionality as gene expression regulators.

Conventional reconstructions of E. coli's metabolism as well as of its gene regulation thoroughly describe the process itself but usually lack information on interacting elements of the other biological system. While there are numerous genome-scale metabolic reconstructions available [4–9], only a few large-scale transcriptional regulatory networks exist that are mainly based on the information from RegulonDB [10]. First attempts to integrate both cellular processes started from metabolic reconstructions which were expanded by regulatory genes and stimuli of the associated encoding metabolic genes [11, 12]. Both studies started from the metabolic model of Reed *et al.* [5] and include 104 regulatory genes and 583 regulatory rules regulating approximately 50 % of the metabolic genes. In this manner, the close proximity of regulatory events was captured but more far-reaching and global effects, *e.g.*, self-contained regulatory dynamics among genes, could not be considered. Further approaches examine the regulatory processes of the metabolic network based on the aforementioned pioneering attempts [13, 14]. For this purpose, the information about regulatory events was assembled in terms of Boolean rules as a variant of Boolean network models.

More recently, Chandrasekaran and Price [15] introduced a method called probabilistic regulation of metabolism, a new variant of regulatory flux balance analysis, i.e., the class of approaches behind some of the pioneering integrative models discussed above [11, 12]. The state of the many variants of integrating regulatory information into flux-balance analysis models has been reviewed in [16] and [17]. The necessity of achieving such data integration, even on the network level, has recently been discussed in [18]. To a certain degree all these studies consider the regulation of metabolism but only cover the proximity rather than a genome scale.

Understanding the interplay of metabolism and gene regulation will help to gain insight in cellular, system-wide responses such as to changing environmental conditions. Here, we present the database-assisted reconstruction of an integrative $E.\ coli$ network capturing metabolic as well as regulatory processes. The attribution of network components (in terms of individual vertices) to the metabolic and regulatory domains, as well as the protein interface enables the further characterization of the network in terms of its modular organization, its path statistics and the vertex centrality.

In particular, we formulate a new measure by evaluating *domain-traversing paths*, in order to quantitatively assess the role of components in the interface domain and thus identify cross-systemic key



Figure 1: Schematic representation of the involved processes and biological elements in the integrative metabolic-regulatory *E. coli* network. Gene regulatory processes primarily comprise genes (○) and several proteins (monomers ▲ as well as complexes ▲), mainly transcription factors. In contrast, metabolic processes are predominantly defined by small molecules (■) and the catalyzing biochemical reactions (●). The interactions between regulatory and metabolic processes can be mainly characterized by proteins (also modified proteins ▲) serving as enzymes and regulators, respectively. While regulatory links are represented as dashed lines, the encoding and reaction-associated links are shown as solid lines.

elements contributing to both regulatory and metabolic processes. In all cases, these topological assessments highlight system components and functional subsystems, which are well known for their biological relevance, thus emphasizing the predictive power of network topology. Employing observations on the topological (structural, network-architectural) level, in order to identify components in the system of particular functional relevance has a long tradition in network biology (and in network science in general).

The main results of our investigation are: We present an integrated network representation of gene regulation and metabolism of $E.\ coli$ and illustrate how it is a promising starting point for the structural investigation of system-wide phenomena. In particular, the network perspective suggests the explicit consideration of a protein interface between the genetic and metabolic realms of the cell. Employing network metrics we argue (1) that a three-domain partitioning is architecturally and functionally plausible, and (2) show that prominent components of the network according to the structural investigation tend to be of evident biological importance. Especially, the evaluation of possible paths through the interface domain of the network reconstruction yields well-known functional subsystems. The overlap of structural and biological investigations by focusing on a limited number of structurally outstanding components. This network model can also serve as a starting point for a range of topological analyses with methods developed in statistical physics (see, *e.g.*, [19] for a recent review).

Summarizing, in contrast to the separate analyses of (e.g., the metabolic or gene regulatory) subsystems, we expect that the integrative network model shown here will draw the attention to system-wide feedback loops not contained in the individual subsystems and to different roles of individual components, which become only visible from the perspective of interdependent networks.

Results

Database-assisted network reconstruction

By now, the dramatic growth of bioinformatics databases [20], both in content and in diversity, allows addressing the challenge of integrating regulation and metabolism on a system-wide level. We devised a semi-automated framework to integrate information from EcoCyc database [21] and RegulonDB [10] into a network for *E. coli* including major cellular processes, from metabolic processes via protein modifications to a variety of regulatory events (see Methods). Networks are an efficient data structure for integrating this wealth of information [22–24]. In this way, the vast amount of information contained in the bioinformatics databases provide an 'architectural embedding' for metabolic-regulatory networks and guides subsequent steps of model refinement and validation. We augmented and validated the resulting network based on existing reconstructions of metabolic [6, 8, 25–27] as well as of gene regulatory processes [10].

The integrative *E. coli* network constructed here comprises the three major biological components, genes, proteins, and metabolites, as well as the metabolizing reactions summing up to more than 12,000 components. Represented as a graph the network has seven types of vertices (Figure 2, Table S1) and seven different types of edges including two types of encoding associations, four reaction-associated relations, and regulatory links (Table S2). The graph representation facilitates the mapping of reactions and their catalyzing enzymes, as both are depicted as vertices. In contrast, metabolic systems are often represented as hypergraphs to illustrate the Boolean 'AND' association of reaction educts and the fixed stoichiometric ratio of the involved metabolites. Those aspects are assigned explicitely as edge properties in the graph representation. Besides the associations of reaction educts, the encoding relations of protein complexes are of Boolean 'AND' type, termed *conjunct* links. On the contrary, associations representing isoforms of protein subunits, isoenzymes as well as reaction products are implemented by Boolean 'OR' links, called *disjunct*. The third linkage type, *regulation*, covers approximately 7,300 regulatory associations, *i.e.*, transcriptional, translational as well as metabolic ones (Table S3).



Figure 2: Spring-block graph representation (using a scalable force directed placement algorithm) and vertex composition of the integrative *E. coli* network. The coverage of the pioneer model from Covert *et al.* [11] is provided in column *i*MC1010.

The metabolic and regulatory processes

The comparison with existing models reveals that the presented integrative network is a comprehensive representation of the metabolic and regulatory processes in *E. coli*. The very first approach of embedding metabolic processes in the regulatory context of Covert *et al.* [11], the *i*MC1010 model, started from a metabolic model which was extended by the regulation of the encoding genes of the catalyzing enzymes. For the purpose of determining the overlap of the integrative metabolic-regulatory network and the *i*MC1010 model, transport reactions as well as the artificial biomass reaction have been disregarded and, moreover, only unique metabolites (neglecting compartmentation) have been taken into account. Else, the different levels of details of the transport systems such as PTS as well as of the compound compartmentation would render a correct mapping impossible. Overall, the *i*MC1010 model is covered by our model to more than 89 % (Figure 2, see Table S4, column 3).

To assess the coverage of E. coli's metabolic processes, the embedded metabolic processes of the integrative E. coli network have been associated to the ones of an established E. coli metabolic reconstruction, namely the *i*AF1260 model from Feist *et al.* [6]. About 67 % of the involved biochemical reactions, compounds and genes could be mapped directly (see Table S4, column 4). Particularly, these two thirds capture almost all biologically relevant components in terms of *in silico* viability. Using flux balance analysis for simulating the biomass production capacity of the *i*AF1260 model and taking the overlap with mapped components of the integrative E. coli network revealed that for the default medium setup approximately 75 % of the essential reactions (to yield 1 % biomass) are covered by the integrative E. coli network.

Analogous to the metabolic processes, the coverage of $E.\ coli$'s gene regulation has been determined using the transcriptional regulatory network from RegulonDB [10]. This model has been assembled in a similar fashion but is accounting only for transcription factors and their regulated genes. With a coverage of more than 98 %, the transcription-related regulatory processes are considered as completely recorded in the integrative $E.\ coli$ network (see Table S4, column 5). Apart from that, for this assessment of overlap a comparison of regulatory processes associated with RNA translation as well as metabolic regulatory events is not possible since the RegulonDB transcriptional regulatory network does not consider protein and metabolic interaction processes.

The interface of metabolic and regulatory processes

The most conspicuous links between metabolic and gene regulatory processes are metabolic transcription factors, *i.e.*, gene expression regulators binding metabolites, and metabolic genes, *i.e.*, genes with significant and coordinated response on the metabolic level such as encoding enzymes. Intuitively, the interface is considered so far as the direct interactions of metabolic elements and gene regulatory domain (MD – RD).

However, by examining those interactions in more detail the topological role of proteins becomes apparent. Regarding the metabolic transcription factors, the respective metabolite binds to a protein and this metabolite-protein complex then subsequently regulates the gene expression. In the case of metabolic genes, ultimately the respective gene encodes a protein which either by itself or as a complex serves as an enzyme. In line with this, the interface of metabolic and gene regulatory processes should be considered as the series of interactions of metabolites and genes, respectively, with proteins and subsequent protein modifications. Thus, the interface does not only comprise interactions (edges) but also components (vertices), and the integrative *E. coli* network will in the following be divided into a metabolic domain, a protein interface and a regulatory domain (MD – PI – RD).

In the next section, the plausibility of the three-domain partition (and the set of biologically motivated rules devised to create it) will be assessed in comparison to the likewise proposed two-domain (MD - RD) representation.

The interface structure – a matter of network partition

In order to assess the large-scale structure of the reconstructed network we apply a set of rules that assign each vertex of the network to one of two and three domains, respectively, by considering the biological types of the vertices themselves as well as those of their neighbors (as outlined in the Methods section). Since these rules have been designed to group together vertices connected to the same biological processes we expect them to result in biologically plausible network partitions.

To complement the two functional partitions, MD - RD and MD - PI - RD, two partitions that solely take into account the vertex types have been analyzed, also representing a metabolic-regulatory division into two and three domains, respectively. For the vertex-driven two-domain partition, the sets of gene and protein vertices denote the regulatory processes while in the three-domain partition regulation is given by the set of genes and the interface domain only consists of the protein vertices. In both cases, metabolism is represented by the sets of reactions and compounds. In the three-domain case, the vertex-driven three-domain partition, the vertex set of proteins form an interface similar to the MD - PI - RD partition (Figure 3). The functional and vertex-driven three-domain partitions are of roughly similar size in terms of vertex count, while the respective two-domain partitions have a metabolic-regulatory vertex ratio of 5:1 and 4:3, respectively (see Table 1).

First, the two three-domain partitions will be compared, *i.e.*, the functional partition, MD - PI - RD, and the vertex-driven partition. In the following, we will argue that the additional third domain acts as an interface between the regulatory and metabolic domains in the functional partition, while we will see that the vertex-driven partition fails to give a coherent picture of the domain-level organization of the biological system.

Especially, it will become clear, also in later sections, that the interface domain in the functional partition contains processes that are known to play prominent roles in system-scale communication within the cell, and may therefore be considered an important component of the large-scale organizational structure of the combined regulation and metabolism of $E. \ coli$.

A simple quantity to illustrate the domain-level picture is the fraction of inter-module edges (linking to a vertex of a different domain) over all edges connected to vertices of a specific domain (*i.e.*, external and internal edges). Of course, there is no objectively 'correct' partition the result of our procedure could be measured against, but there are a number of fundamental properties that a biologically plausible partition in the given context should possess. On the one hand, a proper interface provides the main means of communication between the regulatory and the metabolic processes, *i.e.*, the majority of paths between the outer two domains should run through the interface. Indeed, the interface of the functional partition shows a considerably larger inter-module edge fraction than the remaining domains (0.7 compared to 0.5 and 0.1, Table 1), stressing its special character as a bridging module. A high inter-module edge fraction of the interface is also found in the vertex-driven partition, however, its regulatory domain shows an even higher inter-module edge fraction which indicates an entanglement between the two groups rather than one domain acting as a bridging module to another domain. This exactly give rise for the second criteria, that the domains should capture actual processes (here, structures on the level of several vertices). Unambiguously, regulatory or metabolic processes should be contained within the respective domain so that system-wide interaction takes place between processes. In the following chapter, Interface characterization, it will be shown that this actually is also the case for the interface in the functional partition. In contrast, in the vertex-driven partition already the regulatory domain show deficiencies with respect to that criterion. Since this regulatory domain solely contains gene-gene interactions the intermediate transcription factor steps are not within the domain which become visible in the almost exclusively inter-module edges, linking it to the interface domain.

Next, we compare the three-domain partitions with the two-domain partitions. While the introduction of a third domain allows to study the system in terms of an explicit interface, the partitions into two domains is much closer to common biological intuition. The question which needs to be answered is whether metabolism and gene regulation are solely interfaced by the linking processes such as gene expression, and activation or inhibition of transcription factors and genes, so that the system can appropriately be described with two domains. Or whether there is an actual interface that preferably comprises entire processes additionally including protein modifications and suchlike. Here, this question will be assessed from a topological perspective.

A relevant topological quantity is the network modularity [28] of a given network partition. For a biologically meaningful classification, one would expect on the network level that the regulatory



Functional three-domain partition

Functional two-domain partition



Vertex type-driven three-domain partition

Vertex type-driven two-domain partition

Figure 3: Graph snapshots of the four partitions: the functional three-domain partition into metabolic and regulatory domains and protein interface (MD – PI – RD) (A), the functional two-domain partition into metabolic and regulatory domains (MD – RD) (B), vertex-driven three-domain partition into compounds/reactions, proteins and genes (C), vertex-driven two-domain partition into compounds/reactions, and proteins/genes (D). Vertices are colored according to their domain-affiliation: yellow – (pseudo) regulatory and gene-focused domain, respectively, and blue – (pseudo) metabolic and compound-focused domain, respectively. The interface domain in the three-domain partitions are drawn in red. The diagrams in the top right corners of each panel show the edge composition of the system in terms of intra-domain and inter-domain edges.

Table 1: Topological properties of the functional and vertex type-driven network partitions. The functional partitions are denoted by the respective modules, metabolic domains (MD), regulatory domain (RD) and protein interface (PI). The vertex type-driven partitions are represented by the comprising vertex types, reaction (\bigcirc), compound (\blacksquare), gene (\bigcirc), and protein (\blacktriangle). For each property, the module-specific coefficients and contributions (I, II, III) are presented, respectively. For the modularity, M, the overall network coefficient (Total) is shown as well as the best coefficient is underlined, the module-specific values correspond to the terms in the sum of equation (1).

		Functional parti	tions	Vertex-driven pa	artitions
		MD – PI – RD I – II – III	$egin{array}{c} \mathrm{MD}-\mathrm{RD} \ \mathrm{I} &-\mathrm{III} \end{array}$		
Vertices	I II	8369 2286	10655	7374 2949	7374
	III	2213	2213	2545	5494
Inter-module edge fraction	I II	$0.086 \\ 0.701$	0.106	$0.319 \\ 0.915$	0.319
	III	0.485	0.485	0.969	0.49
Modularity, M	Total I II	$\frac{0.287}{0.166}\\0.042$	$0.157 \\ 0.079$	0.081 0.113 -0.027	$0.226 \\ 0.113$
	III	0.078	0.079	-0.005	0.113

and the metabolic domains show high intra-module connectivity (a large number of links are within a domain) and sparse inter-module linkages (a small number of links are between domains). Accordingly, the network modularity should be high for a successful partition. The results for the modularity are listed in Table 1. The functional partitions clearly outperform the vertex type-driven partitions. Also, when going from MD – RD to MD – PI – RD there is a notable increase in the modularity of the network ($M_2 = 0.157$, $M_3 = 0.287$). Note that here we consider specific candidates for biologically plausible partitions, while a purely topological analysis of the module structure of this large network yields a much larger set of significant modules. Here, a detailed biological interpretation is still missing and will be discussed elsewhere.

Altogether, the functional partition into metabolic domain, protein interface and regulatory domain reflects a biologically reliable classification in two delimited domains linked by a bridging module. Reinforced by the topological properties, the interface structure including full protein modification processes will be used subsequently.

Interface characterization

The interface of metabolic and gene regulatory processes of the integrative *E. coli* network comprises, as expected, predominantly proteins, *i.e.*, monomers and complexes (Table S1), and mainly protein modification processes such as protein translation, protein complex formation and biochemical protein conversion (Table S2). On closer examination, the covered processes can be divided in internal and peripheral ones. According to the bridging role of the interface, the majority of these are peripheral processes (Figure 3, Table S2). The peripheral processes, in turn, can be subdivided according to their directionality meaning from regulatory to metabolic domain (subsequently termed 'downwards') and from metabolic to regulatory domain ('upwards'), respectively. To enumerate the portion of peripheral processes forming complete paths across the interface, direct downwards and upwards links and the new topological concept of *domain-traversing paths* (or short: *traversing paths*) have to be considered. A traversing path connects regulatory and metabolic domain via the protein interface, whereby only starting and end vertex are not affiliated to the bridging domain and the path direction is considered

carefully (see Methods).

Examination of the downwards-upwards subdivision, especially the traversing paths, reveals a considerable (though biologically expected) asymmetry of the interface (Figure 4): The downwards interface is much more pronounced comprising predominantly the transcription of enzymes, *i.e.*, metabolic genes, and the formation of enzymatic protein complexes. On the contrary, the upwards interface is comparably sparse with roughly half the direct (102/283) and quarter the traversing paths (4,070/18,904) connections of the downwards interface. These few upwards processes mainly include the formation of metabolic regulators, especially transcription factors, and the corresponding regulatory events.



Figure 4: Schematic overview of the components and connections of the integrative *E. coli* network, especially those involved in the protein interface. The information about edges are presented in gray and about traversing paths are shown in dark goldenrod while the number of vertices are shown in dark blue and the traversing paths-related ones are given in dark brown, in addition. The solid lines denote direct link connections while the dashed lines the traversing paths connections.

In addition to confirming the interface asymmetry, the traversing paths reveal the bottleneck characteristic of the interface. First indications for this special property are (1) the low number of involved vertices and (2) the distribution of traversing path lengths. For both, downwards and upwards traversing paths, the number of distinct interface vertices in the traversing paths is low compared to the total number, *i.e.*, 1,393 and 449 interface vertices of 2,286 in total, respectively (Figure 4). On the other hand, for both, downwards and upwards traversing paths, emerges a remarkable clustering of paths of length 8–10 and four, six, and 9–11, respectively (Figure 5). This is in contrast to a smooth distribution one would expect in random graphs. By enumerating the involved vertices it is striking that more than 44 % of traversing paths contain one of five three-vertex-combinations, respectively. The respective combinations of downwards and upwards traversing paths pertain to three functional systems, the phosphoenolpyruvate-dependent sugar phosphotransferase system, PTS, the ribonucleotide reducing system, RNR system, as well as the nitrogen regulation two-component signal transduction system, NtrBC system (Table S5).



Figure 5: Distribution of the path lengths for the downwards (RD \rightarrow MD) and upwards traversing paths (MD \rightarrow RD), respectively (dark blue). The golden bars represent the fraction of downwards and upwards traversing paths comprising the PTS and RNR, and the NtrBC system associated vertices.

All three biological subsystems, the PTS [2, 29], the RNR [30–33] as well as the NtrBC system [34–36] are well-studied with respect to their functionality and their cellular context. A schematic representation of the three subsystems is provided in Figure 6. The PTS is an enzymatically active protein complex involved in the transport and phosphorylation of several sugars, so-called PTS-sugars [2]. In the integrative *E. coli* network more than 18 different sugars serve as potential substrates which are imported from peroxisome to cytosol at the same time (Table S6). The substrate variety together with the manifold usage of the associatively produced pyruvate point out the key role of the PTS in *E. coli*'s metabolism and, moreover, suggest that the PTS acts as a bottleneck in the interface.

The RNR system, the second system dominating the downwards traversing paths, provides the major DNA building blocks [32]. Each of the different core enzyme classes, ribonucleotide reductase class I–III, are capable of catalyzing the reduction of all four nucleotides. Its transcriptional and metabolic regulation ensures the balanced supply and, thus, avoid the increase of mutation rates and the loss of DNA replication fidelity [37]. The central cellular role which is reflected in its regulatory embedding, together with its alternate substrates point to its special position in the interface.

The NtrBC system is a two-component signal transduction system initiating the nitrogen starvation response regulation. More precisely, depending on the nitrogen availability NtrB can autophosphorylate and the transfer of the NtrB phosphate group activates the global transduction regulator, NtrC. In *E. coli*, more than 40 genes known to be activated are involved in the nitrogen-response reaction such as active transport and mobilization of nitrogen in terms of N-containing compounds (for integrative *E. coli* network see Table S7). The extensive regulatory function and the linkage to metabolism due to the allocation of ATP for NtrB autophosphorylation indicate that also the NtrBC system acts as a bottleneck in the interface, in the opposite direction to the PTS and RNR system.

The three central traversing paths systems and their biological relevance suggest that a topologically prominent position can be indicative of a biologically important functional entity. To corroborate the general validity of this indication, in the following section different topological properties have been analyzed and the prominent elements have been further characterized from a functional perspective.



Figure 6: Classical representation of the three major interface systems of the integrative *E. coli* network, the phosphoenolpyruvate-dependent sugar phosphotransferase system (PTS, A), the ribonucleotide reducing system (RNR system, B) and the nitrogen regulation two-component signal transduction system (NtrBC system, C). The edges represent biochemical reactions and the vertices denote the involved compounds and proteins. The reactions and proteins highlighted in dark blue are the most abundant vertices determining nearly half of the traversing paths (Table S5).

Cross-systemic key elements of E. coli

The integration of metabolic and regulatory events allows us to determine the key elements of $E. \ coli$, especially those beyond the individual processes. In particular, the functional three-domain partition facilitates to recover network components (in terms of individual vertices) of evident biological relevance, *e.g.*, by means of simple centrality measures. In the following, two different aspects of centrality have been examined [38]: degree centrality depicting the direct linkage of a vertex, and betweenness centrality which can be thought of as the participation of a vertex in the network flow [39].

Starting with the prominent local vertex structure, the so-called hubs (here, vertices with a total degree larger than 50), it is noticable that they are primarily compounds and proteins, in particular protein complexes and appear in all three domains (see Table S8, columns 3–5). In the metabolic domain, hubs include trivial compounds such as H^+ and H_2O and, so-called, currency metabolites, *e.g.*, ATP, NAD(P)H and coenzyme A, while hubs of regulatory processes are obviously global regulators which characteristically exhibit a remarkably strong asymmetry of in-degree and out-degree. Particularly, well-known transcriptions factors top this list such as FNR (fumarate and nitrate reduction) [40], Fis (factor for inversion stimulation) and H-NS (histone-like nucleoid structuring protein) [41]. As stated above, hubs predominantly occur in metabolic and gene regulatory domain while only a few are affiliated to the protein interface. However, it was not to be expected to identify cross-systemic elements solely based on their degree.

To assess/detect cross-systemic key elements an extended approach of degree centrality has been used that additionally accounts for the domain boundaries. The intra-domain degree fraction ξ , also termed embeddedness [42], denotes the ratio of the internal degree of a vertex, within a domain, and the total degree in the network. This measure very clearly distinguishes between, on the one hand, metabolic and regulatory hubs which show intra-domain degree fractions $\xi > 0.87$ (except one single compound with $\xi = 0.185$) and hubs in the interface which in contrast have $\xi \leq 0.06$ (see Table S8, last column). Thus, while metabolic and regulatory hubs are embedded in their respective domains, hubs in the protein interface are mainly connected to vertices in the neighboring domains. In total, seven hubs show a significant low intra-domain degree fraction pointing to their prevalent interactions with the other two domains (Figure S1 and Table S11, column 5). Six of them are affiliated to the protein interface exhibiting numerous interactions with the regulatory domain. Their linkages to the metabolic domain become visible when considering their composition, in case of the protein complexes, and their modes of action, respectively. The former involve the four protein-compound complexes CrpcAMP (cyclic-AMP receptor protein binding cyclic-AMP) [29, 43, 44], DksA-ppGpp (dnaK suppressor binding guanosine 3'-diphosphate 5'-diphosphate) [45–47], NsrR-NO (nitrite-sensitive repressor binding nitric oxide) [48–50] and Lrp-Leu (leucine-responsive regulatory protein binding leucine) [51–53] whose naming schemes already indicate the metabolic link. The latter, namely, protein complex Cra (catabolite repressor activator) and protein monomer Lrp (leucine-responsive regulatory protein) form in the presence of appropriate metabolites, *i.e.*, fructose 1,6-bisphosphate/fructose 1-phosphate and leucine, complexes affecting their regulatory effect. The remaining hub is the metabolic-domain vertex representing guanosine 5'-diphosphate 3'-diphosphate (ppGpp). Besides its special domain-affiliation among the low intra-domain degree hubs, ppGpp acts as an important regulator of both, metabolism and transcriptional processes. More precisely, it regulates several enzyme activities as well as numerous transcription initiations by allosterically binding to RNA polymerase.

So far, we demonstrated that the protein interface of the E. coli network reconstruction acts as a bridging module between regulatory and metabolic domain enabling their interaction and communication. Therefore, we expect the betweenness centrality to directly highlight vertices from the interface. Indeed, ten out of the top-25-ranked (still including currency metabolites) vertices are from the interface (see Table S9, column 5), while overall the interface only accounts for about 18 % of the vertices of the network. Especially, the already mentioned protein-compound complexes CrpcAMP and DksA-ppGpp are among these compounds. In general, currency metabolites and trivial compounds (see above) as well as global regulators are among the central components with respect to betweenness. Overall, the compliance of the most central components regarding degree and betweenness accounts approximately 50 %. Apart from that, biochemical reactions building up and/or breaking down these metabolites and proteins as well as the other involved reactants pertain to the most betweenness-central components. Component association to functional systems allows to assess the systemic feature and by considering the corresponding network affiliation to depict the candidates for cross-systemic key elements. In this manner the network analysis allows us to detect the central role of Crp-cAMP, Lrp-Leu and ppGpp on purely topological grounds, as each component is the focus of such a functional system with high betweenness. Additionally among the top-ranked vertices with respect to betweenness centrality are five further cross-systemic components which are assigned to the protein interface, namely, phosphorylated PhoB (PhoB-P), Fur-Fe²⁺, and three outer membrane proteins (Omp), OmpC, OmpE and OmpF (Table S9). The former two components are transcription factors and therefore acting in the gene regulatory domain, while at the same time they are protein complexes binding a metabolic small molecule depicting the connection to the metabolic processes. The latter three, the outer membran porins, form hydrophilic channels, enabling non-specific diffusion of small molecules across the outer membrane [54-56]. In this role these proteins represent the most obvious connections of gene regulatory and metabolic domain – their encoding genes are highly regulated while the porins enable numerous metabolic transport reactions.

By focusing on the connecting domain of gene regulation and metabolism, the two centrality measures reinforce the key role of further cross-systemic elements. Considering the protein interfaceinduced subgraph both centralities point out the vertices that top the list of the above-discussed downwards traversing paths (Table S10). In more detail, both major systems contributing to the downwards traversing paths are represented each by three vertices, namely, PTS and RNR system (Figure 6, panels A and B). Having a look at the intra-domain degree fraction, which put the focus on protein interface vertices as described above, additionally highlights a representative of the upwards traversing path system NtrBC (Figure 6, panel C), as the second non-hub (Table S11). This corroborates the predictions from the traversing paths and, thus, shows that our new topological measure reveals cross-systemic elements which otherwise only stand out under detailed scrutiny of a large amount of biological information.

Discussion

Here, we present an integrative network covering metabolic processes as well as regulatory events of $E.\ coli$ but, especially, the interaction between both systems. With more than 10,000 vertices, it comprises around two third of the metabolic processes currently integrated in metabolic reconstructions [6] and concerning regulatory events, the presented network incorporates more than 95 % of the established transcription-related processes [10]. Both, metabolic and gene regulatory processes are integrated on a genome scale rather than one of the two providing the network basis which is then expanded by closely related processes in the other subsystem, as it has been done, for example, in conventional metabolic reconstructions which solely involve the encoding genes indirectly. Hitherto, integration of transcriptomics data could only be achieved using the so-called gene-protein-reaction (GPR) associations. On the one hand, this procedure limits the applicable data set to metabolic genes and, on the other hand, it acts on the assumption that all expressed enzymes are present in their active form. Starting from the integrative $E.\ coli$ network, integrating transcriptomics data is much more straightforward and, more importantly, the complete data set can be applied. In this way, multi-domain variants of the frequently employed network-based interpretation of 'omics' data [57–61] can be formulated and indirect and regulatory impacts on metabolism can be examined.

The novelty of the reconstruction, the connection of metabolism and gene regulation, allows us not only to investigate the separate systems but also to assess their interactions. The most relevant connecting links are proteins, on the one hand, those acting as enzymes and, on the other hand, metabolic transcription factors. The functional classification, together with the topological analysis, suggests a network division into three domains: metabolic domain, protein interface and regulatory domain. This partition was corroborated by different connectivity measures and reflects a biologically reliable categorization in two delimited modules linked by a bridging module.

The principal structural feature of the network model, the three-domain organization, is reminicient of the 'bow-tie' architectures frequently discussed in the theory of complex systems, where an input and an output layer are connected via a (typically much smaller) intermediate network [62–64]. Such a bow-tie structure (or, rather, the presence of several nested bow-tie architectures) has for example been discussed for metabolic networks [65], where the diversity of inputs (nutrients) and outputs (biomass components) is much larger than the intermediate processing layer. It has been hypothesized that such a bow-tie organization is a prerequisite for the robust operation of a complex system [62, 63]. Here we observe a bow-tie organization in a system consisting of a rich 'material flow' system (metabolism) and a similarly rich 'control' system (gene regulation) connected via a protein interface.

As our topological assessment shows, the bridging character of the protein interface entails a bottleneck functionality. The analysis of the new topological measure, termed *traversing paths*, highlighted three major biological systems represented by 12 vertices forming more than 40 % of these paths (comprising in total 1465 distinct vertices). These traversing path systems, namely phosphotransferase system (PTS), ribonucleotide reducing (RNR) and nitrogen regulation two-component signal transduction (NtrBC) system, are well-investigated ones with key biological relevance for *E. coli*'s metabolism as well as its gene regulation suggesting that a topologically prominent position points to an important biologically functional entity.

Further detection of cross-systemic key elements in the network was accomplished using additional topological measures. In particular, two centrality measures were studied to account for different

aspects of importance in terms of direct linkage and participation in network flow. Apart from conspicuous components, such as trivial compounds, currency metabolites and global regulators, a group of seven hubs were revealed by degree centrality whose characteristic is a significant low intra-domain degree fraction what numerically reflects the bridging feature of the protein interface. As expected, these components are located in the interface except for one, the vertex representing guanosine 5'diphosphate 3'-diphosphate (ppGpp) which is affiliated to the metabolic domain. On the other hand, the inspection of betweenness centrality highlights rather biological systems than single components and as such point to key components detected before in their functional context. Besides trivial compounds and currency metabolites, this includes Crp-cAMP (cyclic-AMP receptor protein binding cyclic-AMP), Lrp-Leu (leucine-responsive regulatory protein binding leucine) and ppGpp which stand out due to their intra-domain degree fraction as well as seven further components already revealed as hubs.

Intriguingly, the interface-specific key elements of the network could be corroborated by exactly these two centrality measures. The assessment of the interface-induced subgraph using both centralities emphasizes altogether eight vertices of the downwards traversing paths discussed above contributing to the two major systems PTS and RNR. Taking into account the intra-domain degree fraction point out a representative of the upwards traversing path system NtrBC. In conclusion, the importance of vertices revealed by the here presented traversing paths could be reinforced by well-established topological measures showing the predictive power of the new measure.

Eventually, the key elements of the integrative E. coli network according to both centralities illustrate the importance of the different domains and their combined consideration (Table 2). Unsurprisingly, the majority of key elements are affiliated to the metabolic domain and represent trivial compounds and currency metabolites, e.g., H^+ , H_2O , ATP and $NAD(P)^+$. Moreover, predominantly cross-systemic components top this combined list of central elements. First of all, the vertices emphasized also by their low intra-domain degree fraction attract attention, namely, Crp-cAMP, Lrp-Leu and ppGpp. These vertices demonstrate the value of the integrative approach: Only when embedded in domain context their vertex importance emerged. In case of the former two components, additionally, the composition unveils the cross-systemic role, *i.e.*, a transcriptional factor protein binding a metabolic small molecule affecting its regulatory activity. Likewise the two regulatory key elements, $Fur-Fe^{2+}$ and PhoB-P, exhibit this conspicuous linkage to the metabolic domain illustrating their cross-systemic property. In other words, they belong to the so-called metabolic transcription factors and, thus, are related to the upwards interface. The opposite is the case for the three metabolic Omp (outer membran porin) transporters that are among the key elements. While their metabolic linkage is more than obvious, the relation to the regulatory domain appears when the encoding genes are examined. These are highly regulated amongst others by the global regulators Crp-cAMP, Fur-Fe²⁺, Lrp-Leu and PhoB-P. In this manner, the Omp's are classical representatives of proteins related to the downwards interface, even though they are not affiliated with it. The remaining key elements are three metabolic small molecules which are counter-intuitively also related to the interface and the cross-systemic elements detected by the traversing paths. While in case of pyruvate the connection to PTS is apparent at first glance (Figure 6, panel A), the link of glutamate and ammonium and the NtrBC system is less perceptible. The actual connecting element is glutamine which is the ligase product of glutamate and ammonium. It activates the (de)uridylylation of the regulatory protein PII which, in turn, inhibits NtrB autophosphorylation [34, 66]. Altogether, the links to the three major traversing path systems are certainly not the only important processes these elements are involved in but they reinforce their biologically central roles. Remarkably, these connecting elements show up when considering the entire network while to acknowledge their importance the interface-specific analysis is needed.

Beyond the detection of key elements, the integrative approach will allow to examine the interplay and distribution of short-term and long-term regulation in *E. coli*'s metabolism. While metabolic regulation of, for instance, enzyme activities occurs on a short time-scale, regulation of gene expression is a long-term control process. Both types of regulation have been incorporated in the network even though only on a qualitative level, *i.e.*, as activator or inhibitor. Like this, the different effective ranges Table 2: Key elements of the integrative *E. coli* network with respect to degree (DC) and betweenness centrality (BC) rank as well as their functional characteristic and cross-systemic property, respectively. Squares denote trivial compounds (□) and currency metabolites (□) while the colored arrows depict the cross-systemic contribution – ▼ downwards interface-related,
▲ upwards interface-related. The orange arrows emphasize the cross-systemic components with significant low intra-domain degree fraction and the golden ones point out elements indirectly linking to one of the major traversing paths systems.

Vertex name	DC	BC	Property
Proton	1	1	
H_2O	2	2	
ATP	5	3	
Phosphate (P)	4	4	
Proton (periplasmic)	6	5	
Crp-cAMP, transcriptional dual regulator	3	9	
ADP	10	6	
outer membran porin F	7	24	$\mathbf{\nabla}$
outer membran porin C	7	29	$\mathbf{\nabla}$
H_2O (periplasmic)	12	31	
outer membran porin E	9	34	
Fur-Fe ²⁺ , transcriptional dual regulator	25	21	
Pyrophosphate	18	30	
$\rm NAD^+$	13	35	
Phosphate (periplasmic)	19	36	
PhoB-P, transcriptional dual regulator	45	12	
Lrp-Leucine, transcriptional dual regulator	43	17	
Guanosine 5'-diphosphate 3'-diphosphate	35	26	
$NADP^+$	21	41	
Glutamate	24	40	
Pyruvate	30	39	•
Coenzyme A	23	48	
CO_2	32	41	
NH_4^+	31	46	A

in metabolism can be assessed and, thus, its covering by one or both regulation types where central metabolism is said to be highly controlled.

From the perspective of recent advances in network theory

With their balance of structural detail and functional simplicity, network models are capable of revealing organizational principles, which are hard to recognize on a smaller systemic scale (*e.g.*, by analyzing individual pathways) or in functionally richer system representations (*e.g.*, in dynamical models). One purpose of the network provided here is to enable work at the interface of statistical physics and systems biology, where the rich toolbox of complex network analysis is employed to identify functionally relevant non-random features of such biological networks.

The recent work of Jensen *et al.* [67], for example, showed that network structure can reveal, whether an enzyme is susceptible rather to genetic knockdown or pharmacologic inhibition. While in the present study, the network measures do not distinguish between different kinds of vertices or links, the rich biological meta data concerning the different biological roles of the components could be translated into distinct vertex and edge classes. In our own investigation [68] we used this fact to study, in a further example of such an interdisciplinary effort, the balance of robustness and sensitivity in the interdependent network of gene regulation and metabolism, based on the reconstructed network

provided here.

In general, we expect that our network reconstruction can serve as a relevant data resource for the application of methods from the analysis of multiplex [69] and other multilayer networks [19, 70]. Recently, there has been a growing interest in the properties of these systems, especially in the presence of explicit interdependencies between vertices [69, 71]. In contrast to monoplex networks interdependent networks can show a qualitatively different robustness against failures, *i.e.*, cascading failures leading to a sudden system breakdown at a critical initial attack size [72, 73]. The case of different vertex types (as opposed to different edge types) has been considered, for example, in the context of secure communication in a network where eavesdroppers control sets of vertices [74].

On a general level, analyzing statistics of paths with respect to the network's large-scale structure, like the domain-traversing paths used here, might prove useful for the evaluation of other networks that show (possibly more than one) interface-like features.

Concluding remarks

In summary, the analysis of network topology allows to determine key system components in the integrative *E. coli* network. In line with expectations, trivial compounds as well as currency metabolites showed up regardless of the measure that has been applied. In addition, further obvious components including several global regulators were identified. More striking is the detection of components and systems which solely emerge when analyzing specifically the interface. These hidden elements are associated to two of the biologically well-investigated functional subsystems, PTS and NtrBC. Both well-established and newly designed measures of the interface point out the same subsystems, and even the analysis of the entire network discloses components indirectly related to these hidden subsystems.

Apart from trivial and currency metabolites, every detected key element of the entire network contributes to some extent to the downwards and/or upwards interface. This unlooked-for cross-systemic property is reflected either in the complex composition, the intra-domain degree fraction, the proximity to key systems, and/or the interplay with regulatory and metabolic processes. The biological relevance of these components supports their detection and reinforces the predictive power of the novel traversing path measure. In general, we believe that the presented integrative *E. coli* network allows further investigations of the interplay of metabolism and gene regulation which will provide insights into cellular, system-wide responses.

Methods

The interconnected *E. coli* network is based on the EcoCyc database [21], release 20.0, which includes verified information of metabolic and regulatory processes (corresponds to RegulonDB 8.6 [10]) for *E. coli* K-12 substr. MG1655. The network is represented as a graph comprising five different types of vertices, encoding genes, protein monomers and complexes (including enzymes), small compounds, and (bio)chemical reactions (Table S1), as well as three types of edges, encoding and catalyzing associations, reaction connections to educts and products, and regulatory links to sources and targets (Table S2).

Extraction of database information

First, relevant information of the database has been extracted and arranged (Algorithm 1). For each regulatory process, the respective source and target were specified and converted to match one of the vertex types ('regulation.dat', file name of the EcoCyc-archive). To this end, the transcript units were separated into promoter, genes and terminator (if applicable), and the regulatory processes were multiplied per comprising gene. Moreover, each regulating RNA has been translated into its encoding gene to meet the vertex types. In case of the metabolic processes, the reaction educts and products as well as the catalyzing enzymes have been assembled and converted to match one of the vertex groups, the respective educt and product stoichiometry have been assigned and the reaction compartmentation and reversibility have been assessed ('reactions.dat'). Thereby, as cell compartments the periplasmic space, the inner membrane, and the cytosol have been taken into account and reversible reactions have been split up.

Second, vertex candidates have been validated ('reactions.dat', 'compounds.dat', 'proteins.dat', 'genes.dat', 'rnas.dat') and divided into reaction, compound, protein monomer, protein-protein complex, protein-compound

complex, protein-RNA complex, and gene. In doing so, generic terms such as DIPEPTIDES have been substituted ('classes.dat') and double annotations, *e.g.*, CPD-15709 and FRUCTOSE-6P have been decoded. Thereupon, the compositions and the encoding genes of the assembled proteins have been gathered and matched to the vertex groups and the respective logical operation and stoichiometry have been annotated ('protcplxs.col'). Based on the validated vertex lists, the regulatory and metabolic processes have been updated whereby each process was removed with at least one unidentified vertex resulting in the final edge lists.

Network implementation

With the validated vertex and edge lists the graph has been assembled and its largest weakly connected component has been extracted. The three domain partition MD - PI - RD (Tables 3 and S1) as well as the two-domain partition are implemented as vertex properties *affiliation* and *metabolic*. Algorithms 2A and 2B show how the domain affiliation of a vertex is determined by its type and its neighbors' types and affiliations.

Moreover, the mapping to the *E. coli* model of Covert *et al.* [11] has been annotated which integrates the metabolic network *i*JR904 published by Reed *et al.* [5] and the transcription regulatory events related to the encoding genes of the catalzying enzymes. To this end, genes, proteins, metabolites as well as biochemical reactions of the metabolic model have been mapped to the EcoCyc database (release 20.0), in a first step automatically based on their identifier and the resulting dictionaries have been manually curated. As the EcoCyc database does not account for compartmentation of compounds and reaction as well as for exchange reactions, unique metabolites and internal reactions have been considered resulting in a coverage of more than 93 %. By additionally disregarding internal transport reactions a coverage of 96.5 % can be achieved (Table 3).

Integrating the manually curated Covert dictionaries, each vertex has attributed (1) a unique identifier, according to the EcoCyc identifier but also indicating the compartment, (2) a unique type reference, (3) a unique assignment of the model components from Covert *et al.* [11], if applicable, and (4) the affiliations of the two- and three-domain partition. Furthermore, vertices of types gene and reaction have (5a) a name assigned, the blattner ID and the EC number, if applicable. The remaining vertices have additionally (5b) a compartment assigned, where cytosol (c), extracellular space (e), periplasmic space (p), inner membrane (i), outer membrane (o) and membrane in general (m) were taken into account. Similarly, each edge of the network has the attribute (1) type, specifying the connected vertices, and the corresponding (2) stoichiometry, where zero is assigned if not applicable or ambiguous. For edges depicting regulatory processes the stoichiometry actually denotes the mode of regulation, namely activation (+,1) inhibition (-,-1) or combined (0). These edges additionally have assigned (3) an identifier, according to the EcoCyc identifier and (4) a name, specifying the regulation type. All other edge types can be classified as either representing conjunct or disjunct links in the sense that all or solely one incoming link is required for functionality (Table S2).

The fully annotated integrative reconstruction of *E. coli*'s metabolic and regulatory processes is provided as a graph representation in Supplementary File 1.

Graph properties concerning intra- and inter-module connectivity

The following measures have been used in the assessment of the graph partitioning scheme.

Inter-module edge fraction *c*: Given the set of vertices with the domain label *D*, edges connecting these vertices to a vertex of a different label are considered *external*, while edges between vertices of the same label are *internal*. We call

$$c_D = \frac{(\#\text{external edges of D})}{\#(\text{external + internal edges of D})}$$

the inter-module edge fraction of domain D.

Network modularity M denotes the degree to which a given partition divides the network in highly connected groups, modules, which are comparably sparsely connected among each other. Therefore, the intra-module links are counted against the total degree of the module vertices (Equation 1),

$$M = \sum_{j=1}^{N_M} \left(\frac{L(v_{M_j}, w_{M_j})}{L_G} - \left(\frac{\deg(v_{M_j})}{2L_G} \right)^2 \right)$$
(1)

with $N_M - \#$ of modules, $L_G - \#$ of links of graph G, $L(v_M, w_N) = \sum_{v \in M} \sum_{w \in N} \operatorname{link}(v, w)$, $\operatorname{deg}(v_M) = \sum_{v \in M} \operatorname{deg}(v)$.

Domain-traversing paths

A traversing path connects the regulatory and the metabolic domains via the protein interface, specifically, a traversing path of length k is of the form

$$[(u, v_1), (v_1, v_2), \dots (v_{k-1}, w)]$$
⁽²⁾

where the vertices u and w are from the regulatory and the metabolic domain (and vice versa) and the vertices v_i are distinct and part of the protein interface. Starting from the set of edges directly at the intersection of two domains iteratively the vertex successors of the interface domain as well as the final, first successor in the third domain have been determined (Algorithm 3).

Vertex centrality

The key elements of the integrative E. coli network have been determined based on two graph properties.

Degree Centrality DC is a local centrality measure and denotes the total number of in- and out-going edges of a vertex, (Equation 3),

$$DC(v) = k_v = k_v^{\text{in}} + k_v^{\text{out}}.$$
(3)

Here, the vertices with a total degree greater than 50 are termed hubs.

By additionally accounting for the domain boundaries, the intra-domain degree fraction ξ (also termed embeddedness [42]) have been defined as ratio of internal degree, within domain D, and total degree of a vertex, (Equation 4),

$$\xi_D(v) = \frac{k_v^{\text{int}}}{k_v} = \frac{1}{k_v} \sum_{w \in D} \left(A_{vw} + A_{wv} \right) \tag{4}$$

where A denotes the adjacency matrix of the graph.

Betweenness Centrality BC describes the impact on the flux through the network, under the assumption that the transfer follows the shortest paths. In particular, it quantifies the fraction of shortest paths between all pairs of vertices which involve the designated vertex (Equation 5),

$$BC(v) = \sum_{s \neq v \neq t \in V} \frac{\sigma_{st}(v)}{\sigma_{st}}$$
(5)

where σ_{st} is the number of all shortest-paths between the vertices s and t while $\sigma_{st}(v)$ yields the number of these paths that run through v [39].

Algorithm 1: Extraction of database information (EcoCyc, release 20.0) on regulatory and metabolic processes.

```
regprocs = AssembleRegulatoryProcesses()
   Extract information on regulatory processes ('regulation.dat')
   REG type, REG ID, REG source, REG target, REG mode = ParseRegulation()
   for rT in REG_type do
      for rID in REG_ID[rT] do
          Convert regulation source and target to match one of the vertex groups
          if REG source[iT][iID] == RNA then
           Translate RNA into corresponding genes
          if REG_target[iT][iID] == transcript unit then
             Split up transcription units into promoters, genes and terminators
             Translate promoters and terminators into corresponding genes
          s,t = Match2Vertex(REG source[rT][rID],REG target[rT][rID])
          regprocs.append(rT,rID,s,t,REG_mode[rT][rID])
metprocs = AssembleCompartmentedMetabolicProcesses()
   Extract information on metabolic processes ('reaction.dat')
   Rxn_ID,Rxn_enz,Rxn_dir,Rxn_1,Rxn_r,Rxn_loc = ParseReactions()
   for mID in Rxn_ID do
      Match catalyzing enzyme, educts and products to one of the vertex groups
      enz,left,right = Match2Vertex(Rxn_enz[mID],Rxn_1[mID],Rxn_r[mID]])
      Annotate reaction compartmentation
      left_comp,right_comp = AnnotateCompartment(Rxn_loc[mID],left,right)
      Split up reversible reactions and reverse 'right-to-left' reactions
      if Rxn_dir == 'reversible' then
          metprocs.append(mID+'_f',enz,left_comp,right_comp)
          metprocs.append(mID+'_r',enz,right_comp,left_comp)
      else if Rxn_dir == 'right-to-left' then
         metprocs.append(mID,enz,right_comp,left_comp)
      else
       metprocs.append(mID,enz,left_comp,right_comp)
valReg, valMet, valVertices = ValidateProcessesVertices(regprocs, metprocs)
   Compile vertex candidates from regulation sources and targets as well as metabolic reactions, the
    corresponding educts and products, and encoding enzymes
   vertCands = AssembleCandidates(regprocs,metprocs)
   Assign vertex candidates to the seven types: reaction, compound, gene, protein monomer,
    protein-protein-complex, protein-compound-complex, protein-rna-complex
   rxns,cmps,gns,prts,ppc,pcc,prc = ComposeVertexLists(vertCands)
   Decode generic terms ('classes.dat') and double annotations
   curReg,curMet,curRxns,curCmps = Curation(regprocs,metprocs,rxns,cmps)
   Prune vertex lists regarding unmapped candidates
   valVertices = PruneVertices(curRxns,curCmps,gns,prts,ppc,pcc,prc)
   Prune regulatory and metabolic processes with respect to validated vertex lists
   valReg,valMet = PruneRegMetProcesses(curReg,curMet,valVertices)
   Export lists of validated vertices and links, i.e., processes
```

Algorithm 2A: Network affilition compilation based on vertex type, and the vertex neighbors types and affiliations. Affiliation assignment for non-ambiguous reactions, compounds and proteins. continued in Algorithm 2B

```
aff = AssignNonAmbiguousVertices()
   Metabolic regulatory processes will be interpreted as metabolic links
   MetRegs = [Regulation-of-Enzyme-Activity, Regulation-of-Reactions]
   for v in Vtypes(network.vertices) == reaction do
      if Vtypes(Educts(v) \land Products(v)) == compound then aff(v) = metabolic
      else if Vtypes(Educts(v) \land Products(v)) == (compound \lor protein) then
       aff(v) = interface
      else aff(v) = tba
   for v in Vtypes(network.vertices) == compound do
      Compounds which participate in at least one reaction
      if # InvolvedRxns(v) > 0 then
          if # aff(InvolvedRxns(v)) == 1 then aff(v) = aff(InvolvedRxns(v))
          else if metabolic in aff(InvolvedRxns(v)) then aff(v) = metabolic
          else if interface in aff(InvolvedRxns(v)) then aff(v) = interface
          else aff(v) = ambiguous
      Compounds which adjacent vertices are either compounds or proteins
      else if Vtypes(Neighbors(v)) == (compound v protein) then
          if regulation in Etypes(OutEdges(v)) then
             if REGtypes(OutEdges(v)) in MetRegs then aff(v) = metabolic
             else aff(v) = ambiguous
          else aff(v) = ambiguous
      else aff(v) = tba
   for v in Vtypes(network.vertices) == protein do
      Proteins with enzymatic function
      if enzyme - reaction in Etypes(OutEdges(v)) then
          if Vtypes(OutNeighbors(v)) == reaction then
             if # aff(OutNeighbors(v)) == 1 then aff(v) = aff(OutNeighbors(v))
             else if metabolic in aff(OutNeighbors(v))) then aff(v) = metabolic
             else if interface in aff(OutNeighbors(v))) then aff(v) = interface
             else aff(v) = tba
          else if regulation in Etypes(OutEdges(v)) then
             if REGtypes(OutEdges(v)) in MetRegs then aff(v) = metabolic
             else aff(v) = regulatory
          else if # aff(InvolvedRxns(OutNeighbors(v))) == 1 then aff(v) =
           aff(InvolvedRxns(OutNeighbors(v)))
          else if metabolic in aff(InvolvedRxns(OutNeighbors(v))) then aff(v) = metabolic
          else if interface in aff(InvolvedRxns(OutNeighbors(v))) then aff(v) = interface
          else aff(v) = tba
       Proteins involved in metabolic reactions
      else if educt - reaction in Etypes(OutEdges(v)) then aff(v) = interface
      Proteins involved in regulatory processes
      else if regulation in Etypes(OutEdges(v)) then
          if REGtypes(OutEdges(v)) in MetRegs then aff(v) = metabolic
          else aff(v) = regulatory
       Proteins involved in protein complex formation
      else if protein - complex in Etypes(OutEdges(v)) then
          if reaction - product in Etypes(InEdges(v)) then aff(v) = interface
          else if regulation in Etypes(InEdges(v)) then
             if REGtypes(InEdges(v)) in MetRegs then aff(v) = metabolic
             else aff(v) = regulatory
          else aff(v) = interface
      else aff(v) = ambiguous
```

Algorithm 2B: Network affilition compilation based on vertex type, and the vertex neighbors types and affiliations. Affiliation assignment for non-ambiguous genes and vertices assigned as ambiguous.



Table 3: Comparison of vertex composition and the coverage to the model from Covert *et al.* [11] of the integrative *E. coli* network (Largest WCC), the underlying full graph and the EcoCyc database (release 20.0).

	Largest V	WCC	Full grap	h	Database	y**
Vertices	EcoCyc	<i>i</i> MC1010*	EcoCyc	<i>i</i> MC1010*	EcoCyc	<i>i</i> MC1010*
Reaction	4693	569/ 767	7251	601/ 767	2617	717/ 767
Compound	2681	557/ 615	2785	558/ 615	2678	614/ 615
Gene	2545	971/1010	2801	981/1010	4506	1010/1010
Protein monomer	1917		2012			
Protein-protein complex	929	771 / 917	986	775 / 917	5709	015/017
Protein-compound complex	100	//1/ 01/	103	115/ 011	5708	015/ 017
Protein-RNA complex	3		4			
	12868	2868/3209	15942	2915/3209	15509	3151/3209

* accounted only for enzymatic reactions and unique metabolites (1076 and 762 in total); ** reactions in EcoCyc database are case insensitive for compartmentation and reversibility

Algorithm 3: Recursive algorithm for the determination of the, so-termed, domain-traversing paths from regulatory to metabolic domain and *vice versa* truly passing the interface domain.

```
input : graph G = \{V, E\},\
        map aff : V \rightarrow \{regulatory, interface, metabolic\}
downTP,upTP = TraversingPaths( G, aff )
   downTP,upTP = list()
   for (source,target) in E do
      if aff(target) == interface then
         if aff(source) == regulatory then
          InterfaceSuccessors( G, [source,target], metabolic, downTP )
         else if aff(source) == metabolic then
          InterfaceSuccessors( G, [source,target], regulatory, upTP )
recursively determine all successors
InterfaceSuccessors( G, vList, aimAff, sucList)
   for suc in outneighbours( vList[-1] ) do
      vL = list(vList)
      if suc not in vL then
         vL.append( suc )
         if aff(suc) == interface then InterfaceSuccessors( G, vL, aimAff, sucList )
          else if aff(suc) == aimAff then
          sucList.append( vL )
```

References

- Karl Kochanowski, Uwe Sauer, and Elad Noor. Posttranslational regulation of microbial metabolism. *Current Opinion in Microbiology*, 27:10–17, Oct 2015. ISSN 1369-5274. doi: 10.1016/j.mib.2015.05.007. URL http://dx.doi.org/10.1016/j.mib.2015.05.007.
- [2] Adelfo Escalante, Ania Salinas Cervantes, Guillermo Gosset, and Francisco Bolivar. Current knowledge of the Escherichia coli phosphoenolpyruvate-carbohydrate phosphotransferase system: peculiarities of regulation and impact on growth and product formation. Applied Microbiology and Biotechnology, 94(6):1483-1494, May 2012. ISSN 1432-0614. doi: 10.1007/s00253-012-4101-5. URL http: //dx.doi.org/10.1007/s00253-012-4101-5.
- [3] E Goncalves, J Bucher, A Ryll, J Niklas, K Mauch, S Klamt, M Rocha, and J Saez-Rodriguez. Bridging the layers: towards integration of signal transduction, regulation and metabolism into mathematical models. *Mol Biosyst*, 9(7):1576-1583, 2013. doi: 10.1039/C3MB25489E. URL http://pubs.rsc.org/en/ content/articlehtml/2013/mb/c3mb25489e.
- [4] J S Edwards and B Palsson. The Escherichia coli MG1655 in silico metabolic genotype: Its definition, characteristics, and capabilities. Proceedings of the National Academy of Sciences of the United States of America, 97(10):5528-5533, March 2000. ISSN 1091-6490. URL http://www.ncbi.nlm.nih. gov/pmc/articles/PMC25862/.
- [5] Jennifer L Reed, Thuy D Vo, Christophe H Schilling, and Bernhard Palsson. An expanded genome-scale model of Escherichia coli K-12 (iJR904 GSM/GPR). *Genome Biology*, 4(9):R54, 2003. ISSN 1465-6906. doi: 10.1186/gb-2003-4-9-r54. URL http://dx.doi.org/10.1186/gb-2003-4-9-r54.
- [6] Adam M Feist, Christopher S Henry, Jennifer L Reed, Markus Krummenacker, Andrew R Joyce, Peter D Karp, Linda J Broadbelt, Vassily Hatzimanikatis, and Bernhard Palsson. A genome-scale metabolic reconstruction for Escherichia coli K-12 MG1655 that accounts for 1260 ORFs and thermodynamic information. *Molecular Systems Biology*, 3:121, Jun 2007. ISSN 1744-4292. doi: 10.1038/msb4100155. URL http://dx.doi.org/10.1038/msb4100155.
- [7] Jeffrey D. Orth, Bernhard. Palsson, and R. M. T. Fleming. Reconstruction and use of microbial metabolic networks: the core Escherichia coli metabolic model as an educational guide. *EcoSal Plus*, 4(1), Sep 2010. ISSN 2324-6200. doi: 10.1128/ecosalplus.10.2.1. URL http://dx.doi.org/10.1128/ecosalplus.10.2.1.
- [8] J. D. Orth, T. M. Conrad, J. Na, J. A. Lerman, H. Nam, A. M. Feist, and B. Palsson. A comprehensive genome-scale reconstruction of Escherichia coli metabolism. *Molecular Systems Biology*, 7(1):535, Apr 2011. ISSN 1744-4292. doi: 10.1038/msb.2011.65. URL http://dx.doi.org/10.1038/msb.2011.65.
- [9] Jonathan M Monk, Colton J Lloyd, Elizabeth Brunk, Nathan Mih, Anand Sastry, Zachary King, Rikiya Takeuchi, Wataru Nomura, Zhen Zhang, Hirotada Mori, et al. iML1515, a knowledgebase that computes Escherichia coli traits. Nature biotechnology, 35(10):904, 2017.
- [10] Socorro Gama-Castro, Heladia Salgado, Alberto Santos-Zavaleta, Daniela Ledezma-Tejeida, Luis Muñiz-Rascado, Jair Santiago García-Sotelo, Kevin Alquicira-Hernández, Irma Martínez-Flores, Lucia Pannier, Jaime Abraham Castro-Mondragón, et al. RegulonDB version 9.0: high-level integration of gene regulation, coexpression, motif clustering and beyond. Nucleic Acids Research, 44(D1):D133–D143, 2015. ISSN 1362-4962. doi: 10.1093/nar/gkv1156. URL http://dx.doi.org/10.1093/nar/gkv1156.
- [11] Markus W. Covert, Eric M. Knight, Jennifer L. Reed, Markus J. Herrgard, and Bernhard Palsson. Integrating high-throughput and computational data elucidates bacterial networks. *Nature*, 429(6987):92–96, 2004. ISSN 1476-4679. doi: 10.1038/nature02456. URL http://dx.doi.org/10.1038/nature02456.
- [12] Tomer Shlomi, Yariv Eisenberg, Roded Sharan, and Eytan Ruppin. A genome-scale computational study of the interplay between transcriptional regulation and metabolism. *Molecular Systems Biology*, 3:101, 2007. ISSN 1744-4292. doi: 10.1038/msb4100141. URL http://dx.doi.org/10.1038/msb4100141.
- [13] Areejit Samal and Sanjay Jain. The regulatory network of E. coli metabolism as a Boolean dynamical system exhibits both homeostasis and flexibility of response. BMC Systems Biology, 2(1):21, 2008. ISSN 1752-0509. doi: 10.1186/1752-0509-2-21. URL http://dx.doi.org/10.1186/1752-0509-2-21.

- [14] Erwin P Gianchandani, Andrew R Joyce, Bernhard Palsson, and Jason A Papin. Functional states of the genome-scale Escherichia coli transcriptional regulatory system. *PLoS Computational Biol*ogy, 5(6):e1000403, May 2009. ISSN 1553-7358. URL http://www.ncbi.nlm.nih.gov/pmc/articles/ PMC2685017/.
- [15] Sriram Chandrasekaran and Nathan D Price. Probabilistic integrative modeling of genome-scale metabolic and regulatory networks in Escherichia coli and Mycobacterium tuberculosis. Proceedings of the National Academy of Sciences of the United States of America, 107(41):17845-17850, 2010. ISSN 1091-6490. doi: 10.1073/pnas.1005139107. URL http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2955152/.
- [16] Saheed Imam, Sascha Schäuble, Aaron N. Brooks, Nitin S. Baliga, and Nathan D. Price. Data-driven integration of genome-scale regulatory and metabolic network models. *Frontiers in Microbiology*, 6:409, 2015. ISSN 1664-302X. URL https://www.frontiersin.org/article/10.3389/fmicb.2015.00409.
- [17] RP Vivek-Ananth and Areejit Samal. Advances in the integration of transcriptional regulatory information into genome-scale metabolic models. *Biosystems*, 147:1–10, 2016.
- [18] Tong Hao, Dan Wu, Lingxuan Zhao, Qian Wang, Edwin Wang, and Jinsheng Sun. The genome-scale integrated networks in microorganisms. *Frontiers in Microbiology*, 9:296, 2018.
- [19] Nicole E. Radde and Marc-Thorsten Hütt. The Physics behind Systems Biology. EPJ Nonlinear Biomedical Physics, 4(1):7-, 2016. ISSN 2195-0008. doi: 10.1140/epjnbp/s40366-016-0034-8. URL https://doi.org/10.1140/epjnbp/s40366-016-0034-8.
- [20] Michael Y. Galperin, Xose M. Fernandez-Suarez, and Daniel J. Rigden. The 24th annual Nucleic Acids Research database issue: a look back and upcoming changes. *Nucleic Acids Research*, 45(D1):D1–D11, 2017. doi: 10.1093/nar/gkw1188. URL +http://dx.doi.org/10.1093/nar/gkw1188.
- [21] I. M. Keseler, A. Mackie, M. Peralta-Gil, A. Santos-Zavaleta, S. Gama-Castro, C. Bonavides-Martinez, C. Fulcher, A. M. Huerta, A. Kothari, M. Krummenacker, and et al. EcoCyc: fusing model organism databases with systems biology. *Nucleic Acids Research*, 41(D1):D605–D612, 2012. ISSN 1362-4962. doi: 10.1093/nar/gks1027. URL http://dx.doi.org/10.1093/nar/gks1027.
- [22] Trey Ideker and Nevan J. Krogan. Differential network biology. Molecular Systems Biology, 8:565, 2012. doi: 10.1038/msb.2011.99. URL http://dx.doi.org/10.1038/msb.2011.99.
- [23] Dexter Pratt, Jing Chen, David Welker, Ricardo Rivas, Rudolf Pillich, Vladimir Rynkov, Keiichiro Ono, Carol Miello, Lyndon Hicks, Sandor Szalma, Aleksandar Stojmirovic, Radu Dobrin, Michael Braxenthaler, Jan Kuentzer, Barry Demchak, and Trey Ideker. NDEx, the Network Data Exchange. *Cell Systems*, 1 (4):302–305, 2015. ISSN 2405-4712. doi: 10.1016/j.cels.2015.10.001. URL http://dx.doi.org/10.1016/ j.cels.2015.10.001.
- [24] Michael Ku Yu, Michael Kramer, Janusz Dutkowski, Rohith Srivas, Katherine Licon, Jason F. Kreisberg, Cherie T. Ng, Nevan Krogan, Roded Sharan, and Trey Ideker. Translation of genotype to phenotype by a hierarchy of cell subsystems. *Cell Systems*, 2(2):77–88, 2016. doi: 10.1016/j.cels.2016.02.003. URL http://dx.doi.org/10.1016/j.cels.2016.02.003.
- [25] E. J. O'Brien, J. A. Lerman, R. L. Chang, D. R. Hyduke, and B. Palsson. Genome-scale models of metabolism and gene expression extend and refine growth phenotype prediction. *Molecular Systems Biology*, 9(1):693, 2013. ISSN 1744-4292. doi: 10.1038/msb.2013.52. URL http://dx.doi.org/10.1038/ msb.2013.52.
- [26] Kieran Smallbone. Standardized network reconstruction of E. coli metabolism. 2013. URL https:// arxiv.org/abs/1304.2960.
- [27] Joanne K Liu, Edward J O'Brien, Joshua A Lerman, Karsten Zengler, Bernhard Palsson, and Adam M Feist. Reconstruction and modeling protein translocation and compartmentalization in Escherichia coli at the genome-scale. BMC Systems Biology, 8(1):110, Sep 2014. ISSN 1752-0509. doi: 10.1186/s12918-014-0110-6. URL http://dx.doi.org/10.1186/s12918-014-0110-6.
- [28] Roger Guimera and Luis A. Nunes Amaral. Functional cartography of complex metabolic networks. Nature, 433(7028):895-900, Feb 2005. ISSN 1476-4679. doi: 10.1038/nature03288. URL http://dx.doi.org/10. 1038/nature03288.

- [29] Josef Deutscher. The mechanisms of carbon catabolite repression in bacteria. Current Opinion in Microbiology, 11(2):87-93, Apr 2008. ISSN 1369-5274. doi: 10.1016/j.mib.2008.02.007. URL http: //dx.doi.org/10.1016/j.mib.2008.02.007.
- [30] L Thelander and P Reichard. Reduction of ribonucleotides. Annual Review of Biochemistry, 48(1): 133-158, Jun 1979. ISSN 1545-4509. doi: 10.1146/annurev.bi.48.070179.001025. URL http://dx.doi.org/10.1146/annurev.bi.48.070179.001025.
- [31] Marc Fontecave, Per Nordlund, Hans Eklund, and Peter Reichard. The Redox Centers of Ribonucleotide Reductase of Escherichia coli. In F.F. Nord, editor, Advances in Enzymology and Related Areas of Molecular Biology, volume 65, chapter 4, pages 147–183. Wiley-Blackwell, Nov 1992. ISBN http://id.crossref.org/isbn/9780471527602. doi: 10.1002/9780470123119.ch4. URL http://dx.doi.org/ 10.1002/9780470123119.ch4.
- [32] A. Jordan and P. Reichard. Ribonucleotide reductases. Annual Review of Biochemistry, 67(1):71-98, Jun 1998. ISSN 1545-4509. doi: 10.1146/annurev.biochem.67.1.71. URL http://dx.doi.org/10.1146/ annurev.biochem.67.1.71.
- [33] Eduard Torrents. Ribonucleotide reductases: essential enzymes for bacterial life. Frontiers in Cellular and Infection Microbiology, 4:52, Apr 2014. ISSN 2235-2988. doi: 10.3389/fcimb.2014.00052. URL http://dx.doi.org/10.3389/fcimb.2014.00052.
- [34] Peng Jiang and Alexander J Ninfa. Regulation of autophosphorylation of Escherichia coli nitrogen regulator II by the PII signal transduction protein. J Bacteriol, 181(6):1906-1911, Mar 1999. URL http://jb. asm.org/content/181/6/1906.full.
- [35] Larry Reitzer. Nitrogen assimilation and global regulation in Escherichia coli. Annual Review of Microbiology, 57(1):155-176, Oct 2003. ISSN 1545-3251. doi: 10.1146/annurev.micro.57.030502.090820. URL http://dx.doi.org/10.1146/annurev.micro.57.030502.090820.
- [36] Daniel R. Brown, Geraint Barton, Zhensheng Pan, Martin Buck, and Sivaramesh Wigneshweraraj. Nitrogen stress response and stringent response are coupled in Escherichia coli. *Nature Communications*, 5:4115, Jun 2014. ISSN 2041-1723. doi: 10.1038/ncomms5115. URL http://dx.doi.org/10.1038/ncomms5115.
- [37] C. K. Mathews. DNA precursor metabolism and genomic stability. The FASEB Journal, 20(9):1300-1314, Jul 2006. ISSN 1530-6860. doi: 10.1096/fj.06-5730rev. URL http://dx.doi.org/10.1096/fj. 06-5730rev.
- [38] Tore Opsahl, Filip Agneessens, and John Skvoretz. Node centrality in weighted networks: Generalizing degree and shortest paths. *Social Networks*, 32(3):245-251, 2010. ISSN 0378-8733. doi: http:// dx.doi.org/10.1016/j.socnet.2010.03.006. URL http://www.sciencedirect.com/science/article/pii/ S0378873310000183.
- [39] Mark Newman. Networks: An Introduction. Oxford University Press, Inc., New York, NY, USA, 2010. ISBN 0199206651, 9780199206650.
- [40] Patricia J. Kiley and Helmut Beinert. Oxygen sensing by the global regulator, FNR: the role of the iron-sulfur cluster. FEMS Microbiology Reviews, 22(5):341-352, 1998. doi: 10.1111/j.1574-6976.1998. tb00375.x. URL http://dx.doi.org/10.1111/j.1574-6976.1998.tb00375.x.
- [41] Andrew Travers and Georgi Muskhelishvili. Dna supercoiling a global transcriptional regulator for enterobacterial growth? Nat Rev Micro, 3(2):157–169, 2005. ISSN 1740-1526. doi: 10.1038/nrmicro1088. URL http://dx.doi.org/10.1038/nrmicro1088.
- [42] Santo Fortunato and Darko Hric. Community detection in networks: A user guide. Physics Reports, 659:1-44, November 2016. ISSN 0370-1573. doi: 10.1016/j.physrep.2016.09.002. URL http: //www.sciencedirect.com/science/article/pii/S0370157316302964.
- [43] A. Kolb, S. Busby, H. Buc, S. Garges, and S. Adhya. Transcriptional regulation by cAMP and its receptor protein. Annual Review of Biochemistry, 62(1):749-797, Jun 1993. ISSN 1545-4509. doi: 10.1146/ annurev.bi.62.070193.003533. URL http://dx.doi.org/10.1146/annurev.bi.62.070193.003533.

- [44] E. Fic, P. Bonarek, A. Gorecki, S. Kedracka-Krok, J. Mikolajczak, A. Polit, M. Tworzydlo, M. Dziedzicka-Wasylewska, and Z. Wasylewski. cAMP Receptor Protein from Escherichia coli as a Model of Signal Transduction in Proteins – A Review. Journal of Molecular Microbiology and Biotechnology, 17(1): 1–11, 2009. ISSN 1464-1801. doi: 10.1159/000178014. URL http://dx.doi.org/10.1159/000178014.
- [45] Lisa U. Magnusson, Anne Farewell, and Thomas Nyström. ppGpp: a global regulator in Escherichia coli. Trends in Microbiology, 13(5):236-242, May 2005. ISSN 0966-842X. doi: 10.1016/j.tim.2005.03.008. URL http://dx.doi.org/10.1016/j.tim.2005.03.008.
- [46] Katarzyna Potrykus and Michael Cashel. (p)ppGpp: still magical? Annual Review of Microbiology, 62 (1):35-51, Oct 2008. ISSN 1545-3251. doi: 10.1146/annurev.micro.62.081307.162903. URL http://dx. doi.org/10.1146/annurev.micro.62.081307.162903.
- [47] Anjana Srivatsan and Jue D Wang. Control of bacterial transcription, translation and replication by (p)ppGpp. Current Opinion in Microbiology, 11(2):100–105, Apr 2008. ISSN 1369-5274. doi: 10.1016/ j.mib.2008.02.001. URL http://dx.doi.org/10.1016/j.mib.2008.02.001.
- [48] Stephen Spiro. Regulators of bacterial responses to nitric oxide. FEMS Microbiology Reviews, 31(2): 193-211, Mar 2007. ISSN 1574-6976. doi: 10.1111/j.1574-6976.2006.00061.x. URL http://dx.doi.org/10.1111/j.1574-6976.2006.00061.x.
- [49] Jonathan D. Partridge, Diane M. Bodenmiller, Michael S. Humphrys, and Stephen Spiro. NsrR targets in the Escherichia coli genome: new insights into DNA sequence requirements for binding and a role for NsrR in the regulation of motility. *Molecular Microbiology*, 73(4):680–694, Aug 2009. ISSN 1365-2958. doi: 10.1111/j.1365-2958.2009.06799.x. URL http://dx.doi.org/10.1111/j.1365-2958.2009.06799.x.
- [50] Nicholas P. Tucker, Nick E. Le Brun, Ray Dixon, and Matthew I. Hutchings. There's NO stopping NsrR, a global regulator of the bacterial NO stress response. *Trends in Microbiology*, 18(4):149–156, Apr 2010. ISSN 0966-842X. doi: 10.1016/j.tim.2009.12.009. URL http://dx.doi.org/10.1016/j.tim.2009.12.009.
- B R Ernsting, M R Atkinson, A J Ninfa, and R G Matthews. Characterization of the regulon controlled by the leucine-responsive regulatory protein in Escherichia coli. *Journal of Bacteriology*, 174(4):1109–1118, Feb 1992. ISSN 1098-5530. doi: 10.1128/jb.174.4.1109-1118.1992. URL http://dx.doi.org/10.1128/ jb.174.4.1109-1118.1992.
- [52] J M Calvo and R G Matthews. The leucine-responsive regulatory protein, a global regulator of metabolism in Escherichia coli. *Microbiological Reviews*, 58(3):466-490, Sep 1994. ISSN 0146-0749. URL http: //www.ncbi.nlm.nih.gov/pmc/articles/PMC372976/.
- [53] Arie B. Brinkman, Thijs J. G. Ettema, Willem M. De Vos, and John Van Der Oost. The Lrp family of transcriptional regulators. *Molecular Microbiology*, 48(2):287–294, Apr 2003. ISSN 1365-2958. doi: 10.1046/j.1365-2958.2003.03442.x. URL http://dx.doi.org/10.1046/j.1365-2958.2003.03442.x.
- [54] Georg E. Schulz. Bacterial porins: structure and function. Current Opinion in Cell Biology, 5(4): 701-707, Aug 1993. ISSN 0955-0674. doi: 10.1016/0955-0674(93)90143-e. URL http://dx.doi.org/10. 1016/0955-0674(93)90143-E.
- [55] B. K. Jap and P. J. Walian. Structure and functional mechanism of porins. *Physiological Reviews*, 76 (4):1073-1088, 1996. URL http://physrev.physiology.org/content/76/4/1073.
- [56] Tilman Schirmer. General and specific porins from bacterial outer membranes. Journal of Structural Biology, 121(2):101-109, 1998. ISSN 1047-8477. doi: 10.1006/jsbi.1997.3946. URL http://dx.doi.org/ 10.1006/jsbi.1997.3946.
- [57] Carsten Marr, Marcel Geertz, Marc-Thorsten Hütt, and Georgi Muskhelishvili. Dissecting the logical types of network control in gene expression profiles. BMC Syst Biol, 2(1):18, Jan 2008. doi: 10.1186/ 1752-0509-2-18. URL https://bmcsystbiol.biomedcentral.com/articles/10.1186/1752-0509-2-18.
- [58] Nikolaus Sonnenschein, Marcel Geertz, Georgi Muskhelishvili, and Marc-Thorsten Hütt. Analog regulation of metabolic demand. BMC Syst Biol, 5(1):40, Jan 2011. doi: 10.1186/1752-0509-5-40. URL http: //www.biomedcentral.com/1752-0509/5/40.

- [59] Nikolaus Sonnenschein, José Felipe Golib Dzib, Annick Lesne, Sebastian Eilebrecht, Sheerazed Boulkroun, Maria-Christina Zennaro, Arndt Benecke, and MT Hütt. A network perspective on metabolic inconsistency. BMC Systems Biology, 6(1):41, May 2012. doi: 10.1186/1752-0509-6-41. URL https://bmcsystbiol. biomedcentral.com/articles/10.1186/1752-0509-6-41.
- [60] Carolin Knecht, Christoph Fretter, Philipp Rosenstiel, Michael Krawczak, and Marc-Thorsten Hütt. Distinct metabolic network states manifest in the gene expression profiles of pediatric inflammatory bowel disease patients and controls. *Scientific Reports*, 6:32584, 2016. doi: 10.1038/srep32584. URL https://www.nature.com/articles/srep32584.
- [61] Moritz E Beber, Patrick Sobetzko, Georgi Muskhelishvili, and Marc-Thorsten Hütt. Interplay of digital and analog control in time-resolved gene expression profiles. *EPJ Nonlinear Biomedical Physics*, 4(1):8, 2016. doi: 10.1140/epjnbp/s40366-016-0035-7. URL https://epjnonlinearbiomedphys.springeropen. com/articles/10.1140/epjnbp/s40366-016-0035-7.
- [62] Hiroaki Kitano. Biological robustness. Nature Reviews Genetics, 5:826-837, November 2004. doi: 10.1038/nrg1471. URL http://dx.doi.org/10.1038/nrg1471.
- [63] Hiroaki Kitano. Towards a theory of biological robustness. Molecular Systems Biology, 3:137, September 2007. ISSN 1744-4292. doi: 10.1038/msb4100179. URL http://www.ncbi.nlm.nih.gov/pmc/articles/ PMC2013924/.
- [64] Tamar Friedlander, Avraham E Mayo, Tsvi Tlusty, and Uri Alon. Evolution of bow-tie architectures in biology. PLoS Computational Biology, 11(3):e1004055, November 2014. ISSN 1553-7358. doi: 10.1371/ journal.pcbi.1004055. URL http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4370773/.
- [65] Marie Csete and John Doyle. Bow ties, metabolism and disease. Trends in Biotechnology, 22(9):446–450, 2004. ISSN 0167-7799. doi: 10.1016/j.tibtech.2004.07.007. URL http://dx.doi.org/10.1016/j.tibtech.2004.07.007.
- [66] W. C. van Heeswijk, H. V. Westerhoff, and F. C. Boogerd. Nitrogen assimilation in Escherichia coli: putting molecular data into a systems perspective. *Microbiology and Molecular Biology Reviews*, 77 (4):628-695, Dec 2013. ISSN 1092-2172. doi: 10.1128/mmbr.00025-13. URL http://dx.doi.org/10. 1128/MMBR.00025-13.
- [67] Karin J Jensen, Christian B Moyer, and Kevin A Janes. Network architecture predisposes an enzyme to either pharmacologic or genetic targeting. *Cell systems*, 2(2):112–121, February 2016. ISSN 2405-4720. doi: 10.1016/j.cels.2016.01.012. URL http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4770809/.
- [68] David F. Klosik, Anne Grimbs, Stefan Bornholdt, and Marc-Thorsten Hütt. The interdependent network of gene regulation and metabolism is robust where it needs to be. *Nature Communications*, 8:534, 2017. doi: 10.1038/s41467-017-00587-4. URL https://www.nature.com/articles/s41467-017-00587-4.
- [69] Filippo Radicchi and Ginestra Bianconi. Redundant interdependencies boost the robustness of multiplex networks. *Physical Review X*, 7:011013, 2017. doi: 10.1103/PhysRevX.7.011013. URL https://link. aps.org/doi/10.1103/PhysRevX.7.011013.
- [70] Mikko Kivelä, Alex Arenas, Marc Barthelemy, James P. Gleeson, Yamir Moreno, and Mason A. Porter. Multilayer networks. *Journal of Complex Networks*, 2(3):203–271, 2014. doi: 10.1093/comnet/cnu016. URL https://doi.org/10.1093/comnet/cnu016.
- [71] Jianxi Gao, Sergey V. Buldyrev, H. Eugene Stanley, and Shlomo Havlin. Networks formed from interdependent networks. *Nature Physics*, 8(1):40–48, January 2012. ISSN 1745-2473. doi: 10.1038/nphys2180. URL http://dx.doi.org/10.1038/nphys2180.
- [72] Sergey V. Buldyrev, Roni Parshani, Gerald Paul, H. Eugene Stanley, and Shlomo Havlin. Catastrophic cascade of failures in interdependent networks. *Nature*, 464(7291):1025–1028, 2010. ISSN 0028-0836. doi: 10.1038/nature08932. URL http://dx.doi.org/10.1038/nature08932.
- Seung-Woo Son, Golnoosh Bizhani, Claire Christensen, Peter Grassberger, and Maya Paczuski. Percolation theory on interdependent networks based on epidemic spreading. *Europhysics Letters*, 97:16006, 2012. ISSN 0295-5075. doi: 10.1209/0295-5075/97/16006. URL http://stacks.iop.org/0295-5075/97/i=1/a=16006.

[74] Sebastian M. Krause, Michael M. Danziger, and Vinko Zlatić. Hidden connectivity in networks with vulnerable classes of nodes. *Physical Review X*, 6:041022, 2016. doi: 10.1103/PhysRevX.6.041022. URL https://link.aps.org/doi/10.1103/PhysRevX.6.041022.

Supplementary information

Vertices	Total	RD	PI	MD
reaction	4693	0	477	4216
	2681	0	26	2655
⊖ gene	2545	1949	311	285
protein monomer	1917	198	1129	590
\blacktriangle protein-protein complex	929	65	243	621
protein-compound complex	100	0	100	0
protein-rna complex	3	1	0	2
	12868	2213	2286	8369

Table S1: Vertex composition of the integrative E. coli network in total (Total) and for the partitionregulatory domain – protein interface – metabolic domain (RD,PI,MD).

Table S2: Edge composition of the integrative *E. coli* network in total (Total), for the partition regulatory domain – protein interface – metabolic domain (RD,PI,MD) and for the peripheral edges between the three domains.

Edges		Link	Total	MD	MD/PI	PI	PI/RD	RD	RD/MD
	gene - protein	D	1916	312	0	325	803	198	278
	protein - complex	c C	1182	6	809	291	68	5	3
			4	0	0	4	0	0	0
			80	8	42	22	7	1	0
			6	0	0	0	6	0	0
	enzyme - reaction	ı D	1272	1225	5	42	0	0	0
			2775	2657	6	109	1	0	2
	educt - reaction	\mathbf{C}	7707	7374	298	35	0	0	0
			246	0	1	245	0	0	0
			181	0	3	178	0	0	0
			100	0	0	100	0	0	0
	reaction - produc	t D	8303	7892	398	13	0	0	0
			171	0	1	170	0	0	0
			252	0	7	210	35	0	0
			102	0	0	102	0	0	0
\rightarrow	transport	\mathbf{C}	291	281	8	2	0	0	0
►	regulation 3,8	R	207	0	0	0	0	207	0
▲►	1,2,9		1274	0	0	0	376	898	0
	12,14		9	4	1	2	1	1	0
	12		5	2	2	1	0	0	0
▲►●	13		4	0	2	2	0	0	0
►	1,4,9		2082	0	0	0	185	1897	0
	12		2	0	1	1	0	0	0
	12		11	6	1	1	0	1	2
	13		1	0	1	0	0	0	0
▲►	1,2		1160	0	0	0	1160	0	0
▲►○	9		2	0	0	0	0	2	0
	12		2	2	0	0	0	0	0
- *	2,5,6,7	7,10,11	98	0	0	0	0	0	98
	12		701	650	50	1	0	0	0
	12		1728	1667	59	0	0	0	2
	13		10	0	8	2	0	0	0
			31880	22086	1703	1854	2642	3210	385

C – Conjunct encoding; D – Disjunct encoding; R – Regulation

Table S3: List of the 14 different kinds of regulatory processes subsumed in the edge type *regulation* of the integrative *E. coli* network (EcoCyc, release 20.0). Each of the 7296 regulatory processes comprises the regulator source ('Regulator') and target ('Regulated entity') as well as the regulatory mode, namely activation (+) and inhibition (-).

Re	gulation type	#	Regulator	Regulated entity
1	Transcription-Factor-Binding	4302	Protein	Transunit, Promoter
2	Allosteric-Regulation-of-RNAP	219	Protein	Promoter
3	Ribosome-Mediated-Attenuation	12	RNA	Terminator
4	Protein-Mediated-Attenuation	5	Protein	Transunit, Terminator
5	Transcriptional-Attenuation	3	Compound	Transunit, Terminator
6	Rho-Blocking-Antitermination	3	Compound	Terminator
7	${\it Small-Molecule-Mediated-Attenuation}$	2	Compound	Transunit, Terminator
8	RNA-Mediated-Translation-Regulation	195	RNA	Transunit, Gene
9	Protein-Mediated-Translation-Regulation	56	Protein	Transunit, Gene
10	Compound-Mediated-Translation-Regulation	22	Protein	Transunit, Gene
11	Regulation-of-Translation	4	Compound	Transunit, Gene
$\frac{12}{13}$	Regulation-of-Enzyme-Activity Regulation-of-Reactions	$2456 \\ 15$	Compound, Protein Compound, Protein	Enzyme Reaction
14	Regulation	2	Protein	Protein

Table S4: Comparison of vertex composition of the integrative *E. coli* network and the coverage to the integrative model from Covert *et al.* [11], the metabolic model from Feist *et al.* [6], and the transcriptional regulatory network based on the RegulonDB [10].

Vertices of the integrative $E.$ col	<i>i</i> network	<i>i</i> MC1010*	<i>i</i> AF1260**	RegulonDB
Reaction	4693	569/ 767	665/1436	0/ 0
Compound	2681	557/ 615	607/ 963	0/ 0
– Gene	2545	971/1010	1168/1260	1764/1788
Protein monomer	1917			185/ 190
Protein-protein complex	929	771 / 017	0/ 0	11/ 13
Protein-compound complex	x 100	//1/ 01/	0/ 0	0/ 0
Protein-RNA complex	3			0/ 0
Total (model coverage)	12868	2868/3209	2440/3636	1960/1991
		89.4%	67.1%	98.4
Total (EcoCyc coverage)		3156/3209	3636/3636	1991/1991
		98.3%	100.0%	100.0%

 \ast accounted only for intracellular reactions and unique metabolites, in total 1076 and 762

** accounted only for intracellular reactions and unique metabolites, in total 2382 and 1668

LTaversing paths Vertex RD 7304 PTSH-1 \uparrow 7276 PTSH-1 Λ 3642 RXNO-6 Λ 3642 RXNO-6 γ 3642 RXNO-6 γ 2878 γ	K IID	1 1	f	Ē
RD 7304 PTSH-1 7276 PTSH-1 7276 PTSH-1 7304 PTSH-1 7305 PTSH-1 7304 PTSH-1 7305 PTSH-1 7306 RX00-* 2496 RX00-* 2430 4164 4151 RED-T1 4050 0X-TH1			Degree	VITPU
↓ 7276 PTSH-1 MID 3642 RXN0-6 3642 RXN0-6 2878 7304 PTSH-1 7304 PTSH-1 7276 7305 PTSH-1 7276 2430 2430 1464 4164 RIBON 4151 RED-TN 4050 OX-TH	MONOMER	HPr (histidine protein)	37	74
MID 3642 RXNO 2878 2878 7304 PTSH-1 7304 PTSH-1 7276 PTSH-1 2496 RXNO 2430 4464 RIBONI 4151 RED-T1 4050 0X-TH	PHOSPHORYLATED	HPr-P (phosphorylated HPr)	35	79
2878 7304 PTSH-1 7276 PTSH-1 2496 RXNO- 2430 4464 RIBON 4151 RED-TH 4050 OX-TH	6718	$EI-P + HPr \rightarrow HPr-P + EI$	4	4651
7304 PTSH-1 7276 PTSH-1 2496 RXNO- 2430 4464 RIBON 4151 RED-TH 4050 OX-TH				
7276 PTSH-1 2496 RXNO- 2430 4464 RIBON 4151 RED-T1 4050 OX-TH	MONOMER	HPr (histidine protein)	37	74
2496 RXNO- 2430 4464 RIBON 4151 RED-TI 4050 0X-TH	PHOSPHORYLATED	HPr-P (phosphorylated HPr)	35	79
2430 4464 RIBON 4151 RED-T1 4050 OX-TH	7166	$PEP + HPr \leftrightarrow HPr-P + Pyr$	4	4651
4464 RIBON 4151 RED-T 4050 0X-TH				
4151 RED-TI 4050 OX-TH .	UCLEOSIDE-DIP-REDUCTI-CPLX	RDPR1 (ribonucleoside-diphosphate reductase)	29	114
4050 DX-TH	HIOREDOXIN-MONOMER	(reduced) thioredoxin 1	34	82
	IOREDOXIN-MONOMER	oxidized thioredoxin 1	33	89
2046				
4464 RIBON	UCLEOSIDE-DIP-REDUCTI-CPLX	RDPR1 (ribonucleoside-diphosphate reductase)	29	114
4151 RED-TI	HIOREDOXIN2-MONOMER	(reduced) thioredoxin 2	34	82
4050 DX-TH	IOREDOXIN2-MONOMER	oxidized thioredoxin 2	33	89
2046				
8952 of over	rall 18892			
MD 1822 PROTE	IN-NRIP	NtrC-P (phosphorylated NtrC)	47	57
↓ 1572 PROTE	IN-NRIIP	NtrB-P (phosphorylated NtrB)	5 C	2879
RD 1349 NRIPH	OS-RXN	$NtrB-P + NtrC \rightarrow NtrB + NtrC-P$	4	4651
<u>1212</u> of over	rall 4070			

Table S6: Compounds that serve as reactants of the phosphoenolpyruvate–carbohydrate phosphotransferase system, so-called PTS-sugars.

Vertex ID	Vertex name
N-ACETYL-D-GLUCOSAMINE_p	<i>N</i> -acetylglucosamine
N-ACETYL-D-MANNOSAMINE_p	N-acetylmannosamine
NACMUR_p	<i>N</i> -acetylmuramate
ASCORBATE_p	Ascorbate
CELLOBIOSE_p	Cellobiose
DIHYDROXYACETONE	Dihydroxyacetone
GALACTITOL_p	Galactitol
CPD-12538_p	Glucosamine
2-DEOXY-D-GLUCOSE_p	2-Deoxyglucose
CPD-15382_p	keto-Fructose
GLC_p	Glucose
CPD-3570_p	Methylglucoside
HYDROQUINONE-O-BETA-D-GLUCOPYRANOSIDE_P	Hydroquinone-O-glucopyranoside (arbutin)
MANNITOL_P	Mannitol
CPD-12601_p	Mannose
2-0-ALPHA-MANNOSYL-D-GLYCERATE_p	2-O-Mannosylglycerate
CPD-1142_p	Salicin
SORBITOL_p	Sorbitol
TREHALOSE_p	Trehalose

	Vertex	
Vertex ID	name	Function of the encoded protein
EG10385	glnG	NtrC (inhibition)
EG10387	glnL	NtrB (inhibition)
EG10383	glnA	Glutamine synthetase (as 12-fold oligomer; inhibition)
EG12191	glnK	PII-2 (as trimer) can activate the adenylylation of glutamine synthetase
EG10386	$_{\rm glnH}$	glutamine ABC transporter - periplasmic binding protein
EG10388	glnP	glutamine ABC transporter - membrane subunit
EG10389	glnQ	glutamine ABC transporter - ATP binding subunit
EG11629	potF	putrescine ABC transporter - periplasmic binding protein
EG11630	potG	putrescine ABC transporter - ATP binding subunit
EG11631	potH	putrescine ABC transporter - membrane subunit
EG11632	potI	putrescine ABC transporter - membrane subunit
EG12124	hisJ	histidine ABC transporter - periplasmic binding protein
EG10007	hisM	arginine/histidine/lysine/ornithine ABC transporter - membrane subunit
EG10452	hisP	arginine/histidine/lysine/ornithine ABC transporter - ATP binding subunit
EG12125	hisQ	arginine/histidine/lysine/ornithine ABC transporter - membrane subunit
EG10072	argT	arginine/lysine/ornithine ABC transporter - periplasmic binding protein
EG11821	amtB	member of NH_3/NH_4^+ transporters, necessary for growth only at low NH_3 levels
G7071	cbl	Cbl DNA-binding transcriptional activator
G7072	nac	Nac DNA-binding transcriptional dual regulator
G6943	astA	Arginine succinyltransferase - 1 st step in arginine degradation II (AST pathway)
G6941	astB	Succinylarginine dihydrolase (as dimer) - 2 nd step in AST pathway
G6944	astC	Succinylornithine transaminase - 3 rd step in AST pathway
G6942	astD	Succinylglutamate semialdehyde dehydrogenase - 4 th step in AST pathway
G6940	astE	Succinylglutamate desuccinylase - 5^{th} and final reaction in AST pathway
G6523	rutA	Uracil oxygenase - 1 st step in uracil degradation III
G6522	rutB	peroxyureidoacrylate/ureidoacrylate amido hydrolase - 2 nd step in uracil degra- dation III
G6521	$\operatorname{rut}\mathbf{C}$	(predicted aminoacrylate peracid reductase - 3 rd step in uracil degradation III)
G6520	rutD	predicted aminoacrylate hydrolase - 4 th step in uracil degradation III
G6519	rutE	predicted malonic semialdehyde reductase - 5 th step in uracil degradation III
G6518	rutF	(flavin reductase - activity required for 1 st step in uracil degradation (ButA))
G6517	rutG	member of the nucleobase: cation symporter-2 (NCS2) family of transporters
	1 40 0	(probably for uracil)
G6782	ddpX	D-Ala-D-Ala dipeptidase required for wild-type peptidoglycan biosynthesis
G6781	ddpA	(predicted peptide ABC transporter - periplasmic binding component)
G6780	ddpB	(predicted peptide ABC transporter - membrane component)
G6779	ddpC	(predicted peptide ABC transporter - membrane component)
G6778	ddpD	(predicted peptide ABC transporter - ATP-binding component)
G6777	ddpF	(predicted peptide ABC transporter - ATP-binding component)
G6969	yeaG	(impact in adaptation to sustained N starvation, member of Ser protein kinases)
G6970	veaH	(impact in adaptation to sustained N starvation)
EG12834	yhdW	(predicted amino acid ABC transporter - membrane component)
EG12835	yhdX	(predicted amino acid ABC transporter - ATP-binding component)
EG12836	yhdY	(predicted amino acid ABC transporter - membrane component)
EG12837	yhdZ	(predicted amino acid ABC transporter - periplasmic binding component)

 Table S7: Regulated entities of the global transcriptional response regulator of the NtrBC system, the phosphorylated NtrC.

Table S8: Hubs of the integrative *E. coli* network with a total degree of at least 50 (DC), their module affiliation, and the differentiation in in-degree (In) and out-degree (Out) including the affiliation and linkage assignments. The last column denotes the intra-domain degree fraction, ξ , here given as percental fraction.

											I
		Affiliation		in-affilia	tion in-	linkage	out-af	filiation	out-link;	age _{E [0}	1
DC Vertex ID	Vertex name	RD PI MI	_ In	RD PI	MD C	D R	Out RD P	I MD	C D	R R	5
1412 PROTON	Proton	>	1027	15	1012	1027	385 1 9) 375	384	1 98	5
930 WATER	H_2O	>	226	74	152	226	704 2:	3 681	704	89	.6
515 CPLX0-226	Crp-cAMP, transcriptional dual regulator	>	1			1	514 513	_	1	513 0	4.
489 Pi	Phosphate (P)	>	410	×	402 1	409	; 62	2 77	40	39 98	0.
439 ATP	ATP	>	17	က	14	17	422 47	7 375	383	39 88	9.
419 PROTON_P	Proton (periplasmic)	>	96		96	96	323	323	323	100	0.
401 CPLX0-7534_0	[OmpF] ₃ , outer membran porin F complex	>		1	1		400	400	400	66	ø
401 CPLX0-7533_0	[OmpC] ₃ , outer membran porin C complex	>	, ,	Π	1		400	400	400	66	ò
399 CPLX0-7530_0	[OmpE] ₃ , outer membran porin E complex	>			1		398	398	398	66	٢.
350 ADP	ADP	>	299	35	264	299	51 9) 42	22	29 87	4.
293 CPLX0-7797	FNR, transcriptional dual regulator	>	1				$292 \ 292$			292 100	o.
247 WATER_p	H_2O (periplasmic)	>	21		21	21	226	226	226	100	0.
241 NAD	NAD ⁺	>	112		111 1	111	129	l 128	118	11 99	ci
235 NADH	NADH/H ⁺	>	115		115	115	120	l 119	110	10 99	.6
222 PC00027	IHF, transcriptional dual regulator	>	2	2	2		$220 \ 220$			220 99	Ŀ.
221 CPLX0-7705	Fis, transcriptional dual regulator	>	1		1		$220\ 220$			$220 \ 100$	0.
187 PD00288	H-NS, transcriptional dual regulator	>				1	$186 \ 186$			$186 \ 100$	o.
178 PPI	Pyrophosphate	>	147	8	139	147	31	31	6	22 95	v
177 Pi_p	Phosphate (periplasmic)	>	168		168	168	6	6	33	$6\ 100$	0.
176 PHOSPHO-ARCA	ArcA-P, transcriptional dual regulator	>	°.	က	1	2	$173 \ 173$			173 98	сi
159 NADP	NADP ⁺	>	114	4	110	114	45	l 44	34	11 96	6.
151 NADPH	NADPH/H ⁺	>	33		33	33	118	5 113	114	4 96	٢.
144 CO-A	Coenzyme A	>	50	2	$48 \ 1$	49	94 (88	77	17 94	4
138 GLT	Glutamate	>	72		72	72	66	99	61	$5\ 100$	0.
131 CPLX0-7639	$[Fur-Fe^{2+}]_2$, transcriptional dual regulator	>	1	1	1		$130 \ 130$			130 99	2
125 2-KETOGLUTARATE	2-Ketoglutarate	>	58		58	58	; 29	2 65	59	8 98	4.
124 PHOSPHO-NARL	NarL-P, transcriptional dual regulator	>	e.	3 S	1	2	$121 \ 121$			121 97	9.
121 CPLX0-8070	DksA-ppGpp	>	1	1		1	120 119	_	1	119 1	۲.
115 AMP	AMP	>	80	x	72	80	35	l 34	10	25 92	2
111 PYRUVATE	Pyruvate	>	59	ъ	54	59	52	5 47	36	$16 \ 91$	0.
98 AMMONIUM	NH_4^+	>	20		70 1	69	28	28	13	$15\ 100$	o.
87 CARBON-DIOXIDE	CO_2	>	81	3	78	81	9	9	9	96	.6
								Conti	nued on	next pag	e

Continued		
Continue		
Continue	1)	
Continu	-	
Contin	-	
Contin	_	
Contir	1	
Conti	<u> </u>	
Cont		
Cont	1.3	
Con	+	
Cor		
Co	H	
ŭ	-	
Õ	0	
\circ		
\sim	()	
-	\sim	
	-	

$\begin{array}{cccccccccccccccccccccccccccccccccccc$	5 7 7 7	25 2 27 2	49 1	49 49 26 27	1 25 25 23	$ \frac{1}{23} $	ດຕາມດາ		>	NarP-P, transcriptional dual regulator Alanine Sucrose	52 PHOSPHO-NARP 51 D-ALANINE 50 SUC
100.0 98.1	45	45		53 7 45		54 1 7	4 x	~ < 5		Acyltransferase Alkaline nhosnhatase (nerinlasmic)	4 AAS-MONOMER 3 AI.KAPHOSPHA-CPI.X D
$55 \ 96.5$			55	55	1 1	2	5		>	PhoP-P, transcriptional dual regulator	7 PHOSPHOPPHOP
98.3	49	49		8 49	1	1 8	6	>		Cyclopropane fatty acyl phospholipid synthase	S CFA-CPLX
55 3.4	1		55 1	56	2	1	2 1		>	Lrp, transcriptional dual regulator	; PD00353
58 98.3			58	58	1	1	1		>	LexA, transcriptional repressor	PC00010
59 95.2			59	59	1 2	33	с С		>	PhoB-P, transcriptional dual regulator	PHOSPHO-PHOB
61 3.2	1 ($61 \ 1$	62	1	1	1		>	Lrp-Leucine, transcriptional dual regulator	MONOMERO-155
20 96.8	, Ii	59 4	2	61	2	2	2	>		GTP	GTP (GTP)
66 98.5	1		67	67	1	1	1		>	CpxR, transcriptional dual regulator	3 PHOSPHO-CPXR
4 98.5	17	60 5	1	61	7	7	7	>		0_2	3 OXYGEN-MOLECULE
8 94.2	8	33 2	റ	36	33	1 32	с С	33		acetyl-CoA) ACETYL-COA
59 88.7	2	61	5	66	Ω	3 2	5 L	>		$\operatorname{Zinc}\left(\operatorname{Zn}^{2+} ight)$	ZN+2
$10 \ 90.8$	1	x	က	11	65	4 61	5	< 6		Adenosylhomocysteine	ADENOSYL-HOMO-CYS
76 2.6	1		$76 \ 1$	77	1	1	1		>	NsrR-nitric oxide	CPLX0-8047
75 18.5	2	12	$64 \ 1$	27	4	1 3	4	>		NTE Guanosine 5'-diphosphate 3'-diphosphate (ppGpp)	GUANOSINE_TETRAPHOSPHA
3 88.9	3	2 02	1 7	78	e S	1 2	3	>		Adenosylmethionine	S-ADENOSYLMETHIONINE
80 97.6	~		80	80	2	2	2		>	FlhDC, transcriptional dual regulator	CPLX0-3930
79 6.0	5		79 2	81	1 2	3	3		>	Cra, transcriptional dual regulator	l PC00061
R	C D	MD	RD PI	R Out	C D	I MD	n RD P	I II	RD PI N	Vertex name	Vertex ID
se ج [%]	tt-linkag	iation ou	out-affili	ge	in-linka	liation i	in-affi	n	Affiliatic		
NONTITOTIC											

					_																															mund on nevt name
vetem		-										•	•		-	•	-					-							•	•			•			Conti
		>	>	>	>	>	>	>			>					\mathbf{i}			>						>	>	>			\mathbf{i}	>	\mathbf{i}			>	
GD PI									>	>		>	>	>	>		>	>		>	>	>	>	>				>	\mathbf{i}				\mathbf{i}	\mathbf{i}		
Vartav nama		Proton	H_2O	ATP	Phosphate (P)	Proton (periplasmic)	ADP	cyclic-AMP (cAMP)		Crp-cAMP, transcriptional dual regulator			PhoB-P, transcriptional dual regulator		PhoR, sensory histidine kinase (inner membrane)	Leucine		Lrp-Leucine, transcriptional dual regulator	Ferrous (Fe^{2+})		$Fur-Fe^{2+}$	$[Fur-Fe^{2+}]_2$, transcriptional dual regulator	ompF, outer membrane porin F gene	OmpF, outer membrane porin F	$[OmpF]_3$, outer membran porin F complex	CyaA, adenylate cyclase	Guanosine 5'-diphosphate 3'-diphosphate (ppGpp)	ompC, outer membrane porin C gene	OmpC, outer membrane porin C	$[OmpC]_3$, outer membrane porin C complex	Pyrophosphate	H_2O (periplasmic)	ompE, outer membrane porin E gene	OmpE, outer membrane porin E	$[OmpE]_3$, outer membran porin E complex	
Uartey ID		PROTON	WATER	ATP	Pi	$PROTON_p$	ADP	CAMP	RXNO-269_f	CPLX0-226	ADENYLATECYC-RXN	PHOR-RXN	PHOSPHO-PHOB	PHOBR-RXN	PHOSPHO-PHOR_i	LEU	$RXNO-261_f$	MONOMERO-155	FE+2	RXN0-5252_f	CPLX0-7620	CPLX0-7639	EG10671	EG10671-MONOMER	CPLX0-7534_o	ADENYLATECYC-MONOMER	GUANOSINE_TETRAPHOSPHATE	EG10670	EG10670-MONOMER	CPLX0-7533_0	Idd	$WATER_P$	EG10729	MONOMERO-282	CPLX0-7530_0	
Betweenness	CONTRACT CONTRACT	0.29906	0.19894	0.09670	0.08320	0.06466	0.06186	0.05369	0.05362	0.05362	0.05361	0.04082	0.04066	0.04056	0.04052	0.03880	0.03840	0.03839	0.03722	0.03567	0.03567	0.03540	0.03396	0.03395	0.03395	0.03242	0.02743	0.02663	0.02663	0.02663	0.02560	0.02489	0.02432	0.02432	0.02432	

Table S9: Top betweenness central vertices of the integrative E, coli network and involved systems. The central reactions of each system are shaded in the corresponding

			Con	tinued
Betweenness	Vertex ID	Vertex name	GD PI MD System	
0.02319	NAD	NAD ⁺	>	
0.02214	Pi_p	Phosphate (periplasmic)	>	
0.02119	PPPGPPHYDR0-RXN			
0.01890	ATPSYN-RXN_f			
0.01803	PYRUVATE	Pyruvate	► >	
0.01786	GLT	Glutamate	>	
0.01607	CARBON-DIOXIDE	CO-2	•	
0.01603	NADP	NADP ⁺	~	
0.01353	PEPDEPHOS-RXN_2			
0.01252	PEPDEPHOS-RXN_1			
0.01173	ADENYLYLSULFKIN-RXN_r		~	
0.01171	AMMONIUM	NH_4^+	►	
0.01161	CARBAMATE-KINASE-RXN			
0.01144	CD-A	Coenzyme A		
0.01117	ABC-35-RXN_2			
0.01107	ABC-35-RXN_1			
0.01093	EG30063	micF, mRNA-interfering complementary RNA gene		
ATPSYN-R.	$XN: ADP + Pi + 4 PROTON_{p} \leftrightarrow I$	ATP + WATER + 3 PROTON		
ADENYLAT	ECYC-RXN: ADENYLATECYC-MONC	$DMER + ATP \rightarrow CAMP + PPI; RXNO-269; PC00004 + CAM$	$P \rightarrow CPLX0-226$	
PHOR-RXN	: PHOR-MONOMER _i + ATP + Pi -	→ PHOSPHO-PHOR ₁ + ADP; PHOBR-RXN: PHOB-MONOMER +	$PHOSPHO-PHOR_i \rightarrow PHOSPHO-PHOB + PHOR-MONOMER_i$	
RXN0-261	: $PD00353 + LEU \leftrightarrow MONOMERO$	-155		
RXN0-525.	2: PD00260 + FE+2 \leftrightarrow CPLX0-7	7620 -> CPLX0-7639		
EG30063 -	\neg EG10671 \rightarrow EG10671-MONOME1	R -> CPLX0-7534°; EG10670 → EG10670-MONOMER -> CP	$XO-753_{\circ}$; EG10729 \rightarrow MONOMERO-282 -> CPLXO-7530_{\circ}	
РРРСРРНУ	DRO-RXN: GDP-TP + WATER \rightarrow G	UANOSINE_TETRAPHOSPHATE + Pi + PROTON		
PEPDEPHO:	S-RXN: PHOSPHO-ENOL-PYRUVA1	$\Gamma E + ADP + PROTON \leftrightarrow PYRUVATE + ATP$		
CARBAMATI	E-KINASE-RXN: CARBAMOYL-P +	+ ADP + PROTON \rightarrow CARBON-DIOXIDE + AMMONIUM + AT	- Charles	
ADENYLYL.	SULFKIN-RXN: PAPS + ADP + P	$ROTON \leftrightarrow APS + ATP$		

(piBC), respectively. The traversing path systems depict the embedding of system components in the protein interface (Figures 5 and 6, Table S5). Table S10: Top ten key elements of the integrative E. coli network with respect to protein interface-specific degree (piDC) and betweenness centrality

Rank	Vertex ID	Vertex name	piDC	piBC	Traverse path system
1	PTSH-MONOMER	HPr (histidine protein)	1	1	
2	PTSH-PHOSPHORYLATED	HPr-P (phosphorylated HPr)	1	2	-
33	RED-THIOREDOXIN-MONOMER	$_{\rm red}$ Trx1 (reduced thioredoxin 1)	က	က	
က	RED-THIOREDOXIN2-MONOMER	$_{\rm red}$ Trx2 (reduced thioredoxin 2)	က	က	
5	OX-THIOREDOXIN-MONOMER	$_{\rm ox}$ Trx1 (oxidized thioredoxin 1)	3 S	ഹ	
S	OX-THIOREDOXIN2-MONOMER	$_{\rm ox}$ Trx2 (oxidized thioredoxin 2)	33	ഹ	
2	EG50003-MDNDMER	acyl carrier protein (ACP)	x	14	
∞	FLAVODOXIN1-MONOMER	flavodoxin 1	10	17	
6	OX-FLAVODOXIN1	oxidized flavodoxin 1	10	18	
6	PROTEIN-CHEA	chemotaxis protein CheA	6	19	
•••					
12	RIBONUCLEOSIDE-DIP-REDUCTI-CPLX	RDPR1 (ribonucleoside-diphosphate reductase)	7	24	-
•••					
20	RXN0-6718	$EI-P + HPr \rightarrow HPr-P + EI$	63	2	-
•••					
69	ADPREDUCT-RXN_f_1	$NDP + _{red}Trx1 \rightarrow dNDP + _{ox}Trx1 + H_2O$	169	×	
69	ADPREDUCT-RXN_f_2	$NDP + _{red}Trx2 \rightarrow dNDP + _{ox}Trx2 + H_2O$	169	8	
•••					
72	THI OREDOXIN-REDUCT-NADPH-RXN_1	$_{\rm ox} {\rm Trx1} + {\rm NADPH/H^+} \rightarrow _{\rm red} {\rm Trx1} + {\rm NADP^+}$	169	6	
72	THIOREDOXIN-REDUCT-NADPH-RXN_2	$_{\rm ox}{\rm Trx2}$ + NADPH/H ⁺ \rightarrow $_{\rm red}{\rm Trx2}$ + NADP ⁺	169	6	
RXNO-(6718: PTSH-MONOMER + PTSI-PHOSPHORVLATED → PTS UCLEOSIDE-DIP-REDUCTI-CPLX -> RIBONUCLEOSIDE-DI	H-PHOSPHORYLATED + PTSI-MONOMER			



Figure S1: Distribution of intra-domain degree fraction of the integrative *E. coli* network. The yellow shaded area represents the significant low intra-domain degree fractions tested via z-score.

Table S11: Hubs and non-hubs of the integrative *E. coli* network with a significant low intra-domain degree fraction (ξ , tested via z-score) and a total degree (DC) larger than 12, and their domain affiliation (RD, PI and MD).

Ve	ertex name	RD	ΡI	MD	ξ	p-value	DC
	CRP-cAMP DNA-binding transcriptional dual regulator		\checkmark		0.39	0.0042	515
	DksA-ppGpp		\checkmark		1.65	0.0049	121
SC	Cra, transcriptional dual regulator		\checkmark		5.95	0.0071	84
Jul	Guanosine 5'-diphosphate 3'-diphosphate (ppGpp)			\checkmark	18.52	0.0208	81
μ	NsrR-nitric oxide		\checkmark		2.56	0.0052	78
	Lrp-Leucine, transcriptional dual regulator		\checkmark		3.17	0.0055	63
	Lrp, transcriptional dual regulator		\checkmark		3.45	0.0056	58
	${\rm ModE}\text{-}{\rm MoO_4}^{2\text{-}}$ DNA-binding transcriptional dual regulator		\checkmark		4.17	0.006	48
	NtrC-P, transcriptional dual regulator		\checkmark		8.51	0.0089	47
	NagC, transcriptional dual regulator		\checkmark		4.35	0.0061	46
	PdhR, transcriptional dual regulator		\checkmark		4.55	0.0062	44
	ArgR-arginine, transcriptional dual regulator		\checkmark		5	0.0065	40
	DksA RNA polymerase-binding transcription factor		\checkmark		5.13	0.0068	39
	PurR-Hypoxanthine, transcriptional repressor		\checkmark		6.06	0.0071	33
bs	CysB-acetylserine, transcriptional dual regulator		\checkmark		6.25	0.0072	32
-hu	FhlA-Formate, transcriptional activator		\checkmark		6.25	0.0072	32
OD	FadR, transcriptional dual regulator		\checkmark		20.69	0.0245	29
Z	RutR, transcriptional dual regulator		\checkmark		18.18	0.0202	22
	NsrR, transcriptional repressor		\checkmark		14.29	0.0148	21
	CytR, transcriptional repressor		\checkmark		18.75	0.0207	16
	AraC-arabinose, transcriptional activator		\checkmark		12.5	0.0125	16
	GntR, transcriptional repressor		\checkmark		20	0.0228	15
	DnaA-ATP, transcriptional dual regulator		\checkmark		14.29	0.0148	14
	TrpR-Tryptophan, transcriptional repressor		\checkmark		14.29	0.0148	14
	Mlc, transcriptional repressor		\checkmark		15.38	0.0162	13