A note on σ -algebras on sets of affine and measurable maps to the unit interval

Tomáš Crhák

7 Mar 2018

Abstract

In The factorization of the Giry monad Kirk Sturtz considers two σ -algebras on convex spaces of functions to the unit interval. One of them is generated by the Boolean subobjects and the other is the σ -algebra induced by the evaluation maps. The author asserts that, under the assumptions given in the paper, the two σ -algebras coincide. We give examples contradicting this statement.

1 Introduction

The notation and terminology of this paper is mostly based on that of [1]; Cvx is the category of convex spaces with affine maps, Meas is the category of measurable spaces with measurable maps and Set denotes the category of sets.

For a convex space A, the binary operations defining its convex structure will be denoted by $+_r$, $r \in [0,1]$. A subset $E \subseteq A$ is said to be *convex* if it is closed under all $+_r$ and it is said to be a *Boolean subobject* of A if both E and its complement E^{c} are convex. It is immediate that the empty set, A and also the inverse image of a convex set under an affine map are all convex, and similarly for Boolean subobjects.

Depending on the context, the unit interval \mathbb{I} will be viewed either as an object of **Meas** or \mathbf{Cvx} . When considered as a measurable space, the Borel σ -algebra will be assumed. As a convex space, it is endowed with the natural convex structure given by

$$x +_r y = (1 - r)x + ry \qquad \forall r \in [0, 1].$$

The Boolean subobjects of \mathbb{I} are exactly the empty set and the convex sets containing 0 or 1, i.e., the intervals [0, x), [0, x], (x, 1] and [x, 1] for all $x \in \mathbb{I}$.

2 The measurable structure of a convex space

For a convex space A, the Boolean σ -algebra on A is by definition the smallest σ -algebra containing all Boolean subobjects of A, and is denoted by Σ_{bool}^A .

In particular, the Boolean σ -algebra on \mathbb{I} coincides with its Borel σ -algebra. Note that for arbitrary convex spaces A and B, every affine map $A \to B$ is measurable w.r.t. the Boolean σ -algebras on A and B.

For an arbitrary set A, $\mathbf{Set}(A, \mathbb{I})$ may be turned into a convex space by defining $+_r$ pointwise, thus for $f, g \in \mathbf{Set}(A, \mathbb{I})$ and $a \in A$ we have

$$(f +_r g)(a) = f(a) +_r g(a) \quad \forall r \in [0, 1].$$

We are concerned with convex subsets $\mathcal{A} \subseteq \mathbf{Set}(A, \mathbb{I})$. In addition to the Boolean σ -algebra, \mathcal{A} may be endowed with the evaluation σ -algebra, which is induced by the evaluation maps $ev_a : \mathcal{A} \to \mathbb{I}$, $a \in A$, where ev_a is defined by $ev_a(f) = f(a)$. The evaluation σ -algebra will be denoted by $\Sigma_{ev}^{\mathcal{A}}$. Since the evaluations are affine, it follows that

$$\Sigma_{ev}^{\mathcal{A}} \subseteq \Sigma_{bool}^{\mathcal{A}}$$
.

3 Overview of refuted statements

Let A be a convex space, and let $\Sigma A = (A, \Sigma_{bool}^A)$. In [1] the author deals with the Boolean and evaluation σ -algebras on $\mathbf{Cvx}(A, \mathbb{I})$ and $\mathbf{Meas}(\Sigma A, \mathbb{I})$, and asserts:

• Lemma 5.1. For any convex space A

$$(\text{KS-5.1a}) \qquad \qquad \Sigma_{ev}^{\mathbf{Cvx}(A,\mathbb{I})} = \Sigma_{bool}^{\mathbf{Cvx}(A,\mathbb{I})}$$

$$(\text{KS-5.1b}) \qquad \qquad \Sigma_{ev}^{\mathbf{Meas}(\Sigma A, \mathbb{I})} = \Sigma_{bool}^{\mathbf{Meas}(\Sigma A, \mathbb{I})}$$

• Corollary 5.2. For any affine map $P: \mathbf{Meas}(\Sigma A, \mathbb{I}) \to \mathbb{I}$

(KS-5.2)
$$P \text{ is measurable } w.r.t. \ \Sigma_{ev}^{\textit{Meas}(\Sigma A, \mathbb{I})}$$

• Lemma 5.3. For any measurable space X

$$\Sigma_{ev}^{\mathbf{Meas}(X,\mathbb{I})} = \Sigma_{bool}^{\mathbf{Meas}(X,\mathbb{I})}$$

We will show that taking the unit interval for A contradicts (KS-5.2), and hence (KS-5.1b), and thus, for $X = \Sigma A$, disproves (KS-5.3) as well; it is, however, in accordance with (KS-5.1a). Next we show that the free convex space over an uncountable set contradicts (KS-5.1a) and (KS-5.1b).

4 Main results

Throughout this section, A is a convex space, and A is a subset of $\mathbf{Set}(A, \mathbb{I})$. For a function f and a set u, $f|_{u}$, $f^{\rightarrow}(u)$ and $f^{\leftarrow}(u)$ denote the restriction, image and inverse image, respectively.

Let us define $\mathcal{E} \subseteq \mathcal{P}(\mathcal{A})$ by

$$\mathcal{E} = \left\{ E \subseteq \mathcal{A} : \left(\exists u \in [A]^{\leq \omega} \right) \left(\forall f \in E \right) \left(\forall g \in \mathcal{A} \right) \left(f \Big|_{u} = g \Big|_{u} \implies g \in E \right) \right\},\,$$

where $[A]^{\leq \omega}$ denotes the set of countable subsets of A. Then \mathcal{E} has the following properties:

- (1) $\emptyset \in \mathcal{E}, A \in \mathcal{E}$;
- (2) \mathcal{E} is closed under countable unions and intersections;
- (3) for $K \subseteq \mathbb{I}$ and $a \in A$, $ev_a^{\leftarrow}(K) \in \mathcal{E}$.

Proof. (1) is obvious. Regarding (2), let $\mathcal{C} \subseteq \mathcal{E}$ be countable. For each $E \in \mathcal{C}$ fix u_E such that

$$(\forall f \in E) (\forall g \in \mathcal{A}) (f|_{u_E} = g|_{u_E} \implies g \in E)$$

and employ $u = \bigcup \{u_E : E \in \mathcal{C}\}$, which is countable, to show that both $\bigcup \mathcal{C}$ and $\bigcap \mathcal{C}$ belong to \mathcal{E} . The proof of (3) is obvious on setting $u = \{a\}$.

Now it is immediate that $S = \{E \subseteq A : E \in \mathcal{E} \& E^c \in \mathcal{E}\}$ is a σ -algebra and from $(ev_a^{\leftarrow}(K))^c = ev_a^{\leftarrow}(K^c)$ it follows that $ev_a^{\leftarrow}(K) \in \mathcal{S}$. We conclude that $\Sigma_{ev}^{\mathcal{A}} \subseteq \mathcal{S}$, which brings us to

Lemma 1. For every convex space A, and every $A \subseteq Set(A, \mathbb{I})$:

$$\Sigma_{ev}^{\mathcal{A}} \subseteq \mathcal{E}$$
.

We have already mentioned that $\Sigma \mathbb{I} = \mathbb{I}$. From the following two lemmas it follows that $A = \mathbb{I}$ provides a counterexample to (KS-5.1b), though it is in accordance with (KS-5.1a).

Lemma 2. Let $P: \mathbf{Meas}(\mathbb{I}, \mathbb{I}) \to \mathbb{I}$ map every function of $\mathbf{Meas}(\mathbb{I}, \mathbb{I})$ to its Lebesgue integral. Then P is affine but not measurable with respect to the evaluation σ -algebra on $\mathbf{Meas}(\mathbb{I}, \mathbb{I})$.

Proof. We apply Lemma 1 for $A = \mathbb{I}$ and $\mathcal{A} = \mathbf{Meas}(\mathbb{I}, \mathbb{I})$ to show that $P^{-1}(0) \notin \Sigma_{ev}^{\mathbf{Meas}(\mathbb{I},\mathbb{I})}$, from which the conclusion of the lemma follows. For any countable subset $u \subseteq \mathbb{I}$, consider the zero constant function $\overline{0} \in P^{-1}(0)$, and the characteristic function $\chi_{u^c} \in \mathbf{Meas}(\mathbb{I},\mathbb{I})$. Then $\overline{0}|_u = \chi_{u^c}|_u$, but $P(\chi_{u^c}) = 1$, thus $\chi_{u^c} \notin P^{-1}(0)$. Therefore, $P^{-1}(0) \notin \mathcal{E}$.

Lemma 3. The Boolean and evaluation σ -algebras coincide for $Cvx(\mathbb{I}, \mathbb{I})$.

Proof. The affine map $\Phi : \mathbf{Cvx}(\mathbb{I}, \mathbb{I}) \to \mathbb{I} \times \mathbb{I}$ given by

$$\Phi(f) = (f(0), f(1))$$

is an isomorphism of convex spaces. Under this isomorphism, the evaluations ev_0 and ev_1 correspond to the projections of the product, and it is straightforward that $\Sigma_{ev}^{\mathbb{I} \times \mathbb{I}}$ coincides with the usual Borel σ -algebra on $\mathbb{I} \times \mathbb{I}$. Thus, given a Boolean subobject E of $\mathbb{I} \times \mathbb{I}$, our business is to show that E is a Borel set.

Recall that for every pair of elements $p, q \in \mathbb{I} \times \mathbb{I}$ there is a unique affine map $\pi_{p,q} : \mathbb{I} \to \mathbb{I} \times \mathbb{I}$ such that $\pi_{p,q}(0) = p$ and $\pi_{p,q}(1) = q$, which is given explicitly by

$$\pi_{p,q}(r) = p +_r q.$$

From the Closed map lemma it follows that $\pi_{p,q}$ is a closed map, and from this it is clear that the image of every convex subset of \mathbb{I} under $\pi_{p,q}$ is a Borel subset of $\mathbb{I} \times \mathbb{I}$.

Let us now consider the set $\pi_{p,q}^{\leftarrow}(E)$. As it is a Boolean subobject of \mathbb{I} , its image under $\pi_{p,q}$ is a Borel set; we will denote this image by $E_{p,q}$. In the sequel, we will employ the following fact:

$$E_{p,q} \subseteq E$$
 & $\pi_{p,q}^{\rightarrow}(\mathbb{I}) \setminus E_{p,q} \subseteq E^{\mathsf{c}}$.

If $p \in E$, then $0 \in \pi_{p,q}^{\leftarrow}(E)$; the supremum of $\pi_{p,q}^{\leftarrow}(E)$ will be called the dividing point determined by $\pi_{p,q}$. The dividing point determined by $\pi_{p,q}$, denoted for the moment by u, is characterized by the following property:

$$\pi_{p,q}^{\rightarrow}([0,u))\subseteq E\quad \& \quad \pi_{p,q}^{\rightarrow}((u,1])\subseteq E^{\mathsf{c}}.$$

Now that we are through the preliminaries, we will distinguish four cases, depending on how many corners of $\mathbb{I} \times \mathbb{I}$ belong to E and E^{c} .

- (1) If E does not contain any corner, then all of them are contained in E^{c} , and we have $E^{c} = \mathbb{I} \times \mathbb{I}$, so that $E = \emptyset$, hence E is a Borel set.
- (2) Suppose E contains one corner and E^{c} contains the remaining three corners. We proceed for $(0,0) \in E$, the other cases are analogous. Let u and v be the dividing points determined by $\pi_{(0,0),(1,0)}$ and $\pi_{(0,0),(1,0)}$, respectively. Then it is readily checked that

$$E = \{(x, y) \in \mathbb{I} \times \mathbb{I} : uy + vx < uv\} \cup E_{(0, v), (u, 0)},$$

thus E is a Borel set.

(3) Suppose E contains two *neighboring* corners and E^{c} contains the other two corners. We proceed for $(0,0),(1,0) \in E$, the other cases are analogous. Let u and v be the dividing points determined by $\pi_{(0,0),(0,1)}$ and $\pi_{(1,0),(1,1)}$, respectively. Then we have

$$E = \{(x, y) \in \mathbb{I} \times \mathbb{I} : y < (1 - x)u + xv\} \cup E_{(0, u), (1, v)},$$

thus E is a Borel set.

(4) Suppose E contains two *opposite* corners. Then it must also contain at least one of the remaining corners, because otherwise the point $(\frac{1}{2}, \frac{1}{2})$ would belong to both E and E^{c} , a contradiction. But then we deal with a complementary case to (1) or (2).

The next result builds on a free convex space A over an (arbitrary) uncountable set M. Recall that A is the set of maps $\alpha \in \mathbf{Set}(M, \mathbb{I})$ with $\alpha^{-1}(0)$ cofinite in M and $\sum_{m \in M} \alpha(m) = 1$. A is endowed with the pointwise convex structure. As a consequence of the forthcoming lemma we have

- (1) $\Sigma_{ev}^{\mathbf{Cvx}(A,\mathbb{I})} \subsetneq \Sigma_{bool}^{\mathbf{Cvx}(A,\mathbb{I})}$, which contradicts (KS-5.1a);
- (2) $\Sigma_{ev}^{\mathbf{Meas}(\Sigma A, \mathbb{I})} \subsetneq \Sigma_{bool}^{\mathbf{Meas}(\Sigma A, \mathbb{I})}$, which contradicts (KS-5.1b).

Lemma 4. Let $\overline{0}$ be the zero constant function $A \to \mathbb{I}$. For every convex subset $A \subseteq \mathbf{Set}(A, \mathbb{I})$ that includes $\mathbf{Cvx}(A, \mathbb{I})$ we have

$$\{\overline{0}\}\in\Sigma_{bool}^{\mathcal{A}}\setminus\Sigma_{ev}^{\mathcal{A}}.$$

Proof. Convexity of $\{\overline{0}\}$ and its complement is facile, we focus on $\{\overline{0}\} \notin \Sigma_{ev}^{A}$. For a countable subset $u \subseteq A$ let $M_0 = \bigcap \{\alpha^{-1}(0) : \alpha \in u\}$, and define $g: A \to \mathbb{I}$ by

$$g(\alpha) = \sum_{m \in M_0} \alpha(m).$$

Then g is affine, thus $g \in \mathcal{A}$. For $\alpha \in u$ we have $g(\alpha) = 0$, so that $g\big|_u = \overline{0}\big|_u$. However, $g \neq \overline{0}$. To see this, observe that M_0 is non-empty as it is a countable intersection of cofinite subsets of M, which has been chosen to be uncountable. For an arbitrary $m_0 \in M_0$ we take the unique $\alpha_0 \in A$ with $\alpha_0(m_0) = 1$, hence $g(\alpha_0) = 1$. This proves $g \notin \{\overline{0}\}$ and, as u was arbitrary, we conclude that $\{\overline{0}\} \notin \mathcal{E}$. Now apply Lemma 1 to show $\{\overline{0}\} \notin \mathcal{E}_{ev}^A$.

References

[1] Kirk Sturtz, The factorization of the Giry monad (2018), available at arXiv:1707.00488v2[math.CT].