

Boltzmann approach to spin-orbit-induced transport in effective quantum theories

Cong Xiao, Bangguo Xiong, Fei Xue

Department of Physics, The University of Texas, Austin, Texas 78712-0264, USA

In model studies of the spin/anomalous Hall effect, effective Hamiltonians often serve as the starting point. However, a complete effective quantum theory contains not only the effective Hamiltonian but also the relation linking the physical observables to the canonical ones. We construct the semiclassical Boltzmann (SB) transport framework in the weak disorder-potential regime directly in the level of the effective quantum theory, and confirm this construction by formulating a generalized Kohn-Luttinger density matrix transport theory also in this level. The link and difference between the present SB theory and previous phenomenological Boltzmann, quantum kinetic and usual Kubo-Streda theories are clarified. We also present the slightly generalized Kubo-Streda formula in the level of the effective quantum theory. In this level, it is this generalized Kubo-Streda formula rather than the usual one that leads to the same physical interpretations as the present SB theory. In the application to a Rashba 2D effective model, a nonzero spin Hall effect important in the case of strong Rashba coupling but neglected in previous theories is found.

I. INTRODUCTION

In model studies of the spin/anomalous Hall effect – a spin-orbit-induced transverse spin/charge transport effect [1, 2], effective Hamiltonians, such as the two-band Rashba model [3], often serve as the starting point. The spin-orbit interaction (SOI) appearing in the external-perturbation-free effective Hamiltonian is often termed as the band-structure SOI. The effective Hamiltonian is only part of the effective quantum theory, thus may not be enough for predictions relevant to experiments [4, 5]. A complete effective quantum theory contains not only the effective Hamiltonian in the presence of external perturbations but also the relation linking the physical observables to the canonical ones. This is in fact clear in the seminal work of Nozieres and Lewiner [6], who obtained the effective quantum theory for conduction electrons from the parent eight-band Kane model in direct-gap III-V semiconductors. The physical position operator in this effective quantum theory differs from the canonical one by an additional term arising from the projection from the eight-band to the two-band models [4, 5, 7]. This additional term gives rise to sizable effective SOIs with the external electric field and with impurities in the level of the effective quantum theory [6].

Unfortunately, the complicated and phenomenological Boltzmann treatment of Nozieres and Lewiner on the Hall transport did not yield a simple picture for the many terms they obtained. Perhaps partly due to this reason, many subsequent theories using the phenomenological-Boltzmann [8, 9] or usual Kubo-Streda (zero-frequency linear response [10]) diagrammatic [11–13] approaches still considered only the effective Hamiltonian without SOI with the driving external electric field and neglected the change of the physical position. Although some recent phenomenological-Boltzmann [14–16] and quantum kinetic theories [17] considered the complete Nozieres-Lewiner effective model, the link and difference between various theories applied to this simple model have not yet been completely clarified. Moreover, as we will re-

veal, these different considerations may lead to different spin Hall conductivities when the external-perturbation-free effective Hamiltonian has its own internal structure, i.e., the band-structure SOI in the level of the effective theory. Therefore, a simple transport-theory framework with physical insights in the level of the effective quantum theory is highly desirable.

The semiclassical Boltzmann (SB) transport theory is appealing because it is conceptually simple [1, 18, 19] and has the microscopic density matrix approach as its solid foundation [20–24]. When the SB and equivalent theories [24, 25] apply, the spin/anomalous Hall effect can be parsed clearly in the presence of static disorder. Three mechanisms – intrinsic, anomalous quantum (called side-jump in recent reviews [1, 2]) and skew scattering – are defined unambiguously [1, 2]. For the spin/anomalous Hall conductivity, the intrinsic contribution is independent of disorder, the anomalous quantum contribution relies on the disorder but turns out to be independent of the impurity density, whereas the skew scattering contribution from disorder is inversely proportional to the impurity density. However, existing SB theory and the underlying Kohn-Luttinger density matrix theory are only formulated in the level of the full Hamiltonian where the physical observables are just the canonical ones [4, 5].

In this paper we construct the SB framework in the weak disorder-potential regime directly in the level of the effective quantum theory, and confirm this construction by formulating a generalized Kohn-Luttinger theory also in this level [26]. It is shown that the spin/anomalous Hall effect studied in this level can still be parsed into the same categories as in the level of the full Hamiltonian, in the regime where the SB theory works [24]. We discuss the link and difference between the present SB theory and previous phenomenological Boltzmann [8, 9, 14–16], quantum kinetic [27–29] and usual Kubo-Streda diagrammatic [11–13, 30] theories. To help clarify this issue, we also derive the slightly generalized Kubo-Streda formula in the level of the effective quantum theory. In this level, it is this generalized Kubo-Streda formula rather than the usual one [10, 11] that leads to the same physical

picture as the present SB theory.

The SB picture is valid in the Boltzmann regime where the disorder-broadening of bands is quite smaller than the minimal intrinsic energy-scale around the Fermi level [24]. It is easy to reach this regime even in moderately dirty systems if the minimal intrinsic energy-scale around the Fermi level is quite large. However, in the opposite case, the system may be located within the so-called diffusive regime (limit) [31] where a drift-diffusion equation for coupled spin-charge dynamics holds [27, 29]. Both the SB and drift-diffusion theories can only work in limited and different regimes [32] with different physical pictures.

The paper is organized as follows. The SB theory is formulated in Sec. II and III, whereas further model-analysis and comparison with other theories are presented in Sec. IV and V. Before summarizing the paper in Sec. VI we discuss the validity of a widely-accepted idea proposed in a previous phenomenological Boltzmann theory. The generalized Kohn-Luttinger and Kubo-Streda approaches in the level of the effective quantum theory are presented in Appendix A and C, respectively. Appendix B and D contains some other details supporting the discussion in the maintext.

II. EFFECTIVE QUANTUM THEORY

The total single-carrier effective Hamiltonian reads [6]

$$\hat{H}_T = \hat{H}_0 + \hat{V}(\hat{\mathbf{r}}^{phy}) - e\mathbf{E} \cdot \hat{\mathbf{r}}^{phy}, \quad (1)$$

where the physical position operator $\hat{\mathbf{r}}^{phy}$ may differ from the canonical one $\hat{\mathbf{r}}$. Hereafter $\hat{V}(\hat{\mathbf{r}}^{phy})$ and $\hat{V}(\hat{\mathbf{r}})$ are sometimes designated as \hat{W} and \hat{V} , respectively, for brevity. The velocity operator in the presence of disorder and external electric field is $\hat{\mathbf{v}} = \frac{1}{i\hbar} [\hat{\mathbf{r}}^{phy}, \hat{H}_T] = \hat{\mathbf{v}}^{phy} + \delta^V \hat{\mathbf{v}} + \delta^E \hat{\mathbf{v}}$, where $\hat{\mathbf{v}}^{phy} \equiv \frac{1}{i\hbar} [\hat{\mathbf{r}}^{phy}, \hat{H}_0]$, $\delta^V \hat{\mathbf{v}} \equiv \frac{1}{i\hbar} [\hat{\mathbf{r}}^{phy}, \hat{V}(\hat{\mathbf{r}}^{phy})]$ and $\delta^E \hat{\mathbf{v}} \equiv \frac{1}{i\hbar} [\hat{\mathbf{r}}^{phy}, -e\mathbf{E} \cdot \hat{\mathbf{r}}^{phy}]$. For 2D electrons or holes [28],

$$\hat{\mathbf{r}}^{phy} = \hat{\mathbf{r}} + \hat{\mathbf{r}}^a = \hat{\mathbf{r}} + \frac{\lambda_0^2}{4} \hat{\sigma} \times \hat{\mathbf{K}}, \quad (2)$$

where $\hat{\mathbf{K}} = (\hat{k}_x^3 - 3\hat{k}_y^2 \hat{k}_x, 3\hat{k}_x^2 \hat{k}_y - \hat{k}_y^3, 0)$ for 2D holes and $\hat{\mathbf{K}} = \hat{\mathbf{k}}$ for 2D electrons, $\hat{\mathbf{k}} = (\hat{k}_x, \hat{k}_y)$ is the momentum operator. In the band-eigenstate representation of the disorder-free effective Hamiltonian \hat{H}_0 ,

$$(\hat{\mathbf{r}}^{phy})_{ll'} = i\partial_{\mathbf{k}} \delta_{ll'} + i\tilde{\mathbf{J}}_{ll'}, \quad (3)$$

with $i\tilde{\mathbf{J}}_{ll'} \equiv i\mathbf{J}_{ll'} + i\mathbf{J}_{ll'}^a$, $\mathbf{J}_{ll'} = \delta_{\mathbf{k}\mathbf{k}'} \langle u_l | \partial_{\mathbf{k}} | u_{l'} \rangle$ and $i\mathbf{J}_{ll'}^a = \delta_{\mathbf{k}\mathbf{k}'} \frac{\lambda_0^2}{4} \hat{\sigma} \times \mathbf{K}$. $|l\rangle \equiv |\eta\mathbf{k}\rangle = |\mathbf{k}\rangle |u_{\mathbf{k}}^\eta\rangle$ is the eigenstate of \hat{H}_0 with energy $\epsilon_l \equiv \epsilon_{\mathbf{k}}^\eta$, η the band index.

III. SB FRAMEWORK IN THE WEAK DISORDER-POTENTIAL REGIME

Regarding the linear response to a constant electric field \mathbf{E} in non-degenerate multiband electron systems with weak static disorder, in the SB framework the average value of physical quantity A is expressed as

$$A = \sum_l A_l f_l. \quad (4)$$

The semiclassical distribution function $f_l = f_l^0 + g_l + g_l^a$ is expanded up to the linear order of \mathbf{E} around the equilibrium Fermi distribution function f^0 , g_l and g_l^a equilibrate the effects of the driving electric field between and during successive scattering events, respectively [19, 22, 33]. These processes are accounted for by the linearized SB equation in nonequilibrium steady-states [19, 25]

$$e\mathbf{E} \cdot \mathbf{v}_l^0 \frac{\partial f^0}{\partial \epsilon_l} = - \sum_{l'} \omega_{ll'} (g_l - g_{l'}), \quad (5)$$

$$0 = \sum_{l'} \omega_{ll'}^{(2)} \left(g_l^a - g_{l'}^a - \frac{\partial f^0}{\partial \epsilon_l} e\mathbf{E} \cdot \delta \tilde{\mathbf{r}}_{l'l} \right). \quad (6)$$

Here $\mathbf{v}_l^0 = \partial \epsilon_l / \hbar \partial \mathbf{k}$ is the group velocity, $\omega_{ll'} = \frac{2\pi}{\hbar} \langle |T_{ll'}|^2 \rangle \delta(d_{ll'})$ is the scattering rate ($l' \rightarrow l$), $\langle \dots \rangle$ denotes the disorder average. Hereafter we employ the simplified notations $d_{ll'} \equiv \epsilon_l - \epsilon_{l'}$ and $d_{ll'}^\pm \equiv d_{ll'} \pm i\hbar s$. $s \rightarrow 0^+$ appears to be the regularizing factor in the T-matrix theory. The T-matrix is determined by the Lippmann-Schwinger equation $\hat{T} = \hat{W} + \hat{W} (\epsilon_l - \hat{H}_0 + i\hbar s)^{-1} \hat{T}$. In the weak disorder-potential regime $\omega_{ll'}$ is obtained by expanding $\langle |T_{ll'}|^2 \rangle$ in terms of the disorder-potential: $\omega_{ll'} = \omega_{ll'}^{(2)} + \omega_{ll'}^{3a} + \omega_{ll'}^{4a}$. The number and ‘‘a’’ in the superscripts mean the order of disorder potential and the anti-symmetric part ($\omega_{ll'}^a = \frac{1}{2} (\omega_{ll'} - \omega_{l'l})$) of the scattering rate, respectively. In the weak disorder-potential regime the expansion up to the third Born order is sufficient. Taking into account the symmetric part of the higher-order scattering rate only renormalizes the longitudinal distribution function [1, 19] and is not necessary in the Boltzmann regime in the case of weak disorder potential. Therefore, $g_l = g_l^{(-2)} + g_l^{(-1)} + g_l^{(0)}$ and concurrently we can set $g_l^{(-1)} = g_l^{sk}$ and $g_l^{(0)} = g_l^{pair}$ to emphasize the physic related to these distribution functions: g_l^{sk} arises from the skew scattering and g_l^{pair} from scattering off pairs of impurities (details in Refs. [19, 24]). For the coordinate-shift [19] $\delta \tilde{\mathbf{r}}_{l'l}$, the wavepacket derivation of Sinitsyn et al. [22] who assumed $\hat{\mathbf{r}} = \hat{\mathbf{r}}^{phy}$ can be directly generalized to the present case ($\hat{\mathbf{r}} \rightarrow \hat{\mathbf{r}}^{phy}$, $\hat{V}(\hat{\mathbf{r}}) \rightarrow \hat{V}(\hat{\mathbf{r}}^{phy}) = \hat{W}$), yielding

$$\delta \tilde{\mathbf{r}}_{l'l} = i\tilde{\mathbf{J}}_{l'l} - i\tilde{\mathbf{J}}_l - \hat{\mathbf{D}} \arg W_{l'l}, \quad (7)$$

with $i\tilde{\mathbf{J}}_l \equiv i\tilde{\mathbf{J}}_{ll}$ and $\hat{\mathbf{D}} = \partial_{\mathbf{k}'} + \partial_{\mathbf{k}}$.

In Eq. (4), an appropriate expression for A_l is needed. In the conventional SB consideration one takes $A_l = \langle l | \left(\hat{A}^{phy} + \delta^V \hat{A} + \delta^E \hat{A} \right) | l \rangle$. For spin-orbit-induced transport, the electric-field-induced and impurity-induced corrections to $|l\rangle$ contribute qualitatively [23], thus $|l\rangle \rightarrow |l\rangle + |\delta^E l\rangle + |\delta^V l\rangle$. Here $|\delta^E l\rangle = -e\mathbf{E} \cdot \sum_{l' \neq l} |l'\rangle \langle l' | \hat{\mathbf{r}}^{phy} | l \rangle / d_{ll'}^+$ is the electric-field induced correction to the Bloch state, whereas $|\delta^V l\rangle = \left(\epsilon_l - \hat{H}_0 + i\hbar s \right)^{-1} \hat{T} | l \rangle$ is the scattering correction. Therefore we have

$$A_l \equiv \tilde{A}_l^0 + \delta^{in} A_l + \delta^{in,1} A_l + \delta^{sj} A_l + \delta^{sj,1} A_l, \quad (8)$$

where $\tilde{A}_l^0 = \langle l | \hat{A}^{phy} | l \rangle$ and

$$\delta^{in} A_l = 2 \text{Re} \langle l | \hat{A}^{phy} | \delta^E l \rangle, \quad \delta^{in,1} A_l = \langle l | \delta^E \hat{A} | l \rangle, \quad (9)$$

and

$$\begin{aligned} \delta^{sj} A_l &= 2 \text{Re} \left\langle \langle l | \hat{A}^{phy} | \delta^V l \rangle \right\rangle^{off} + \left\langle \langle \delta^V l | \hat{A}^{phy} | \delta^V l \rangle \right\rangle^{off}, \\ \delta^{sj,1} A_l &= 2 \text{Re} \left\langle \langle l | \delta^V \hat{A} | \delta^V l \rangle \right\rangle^{off}. \end{aligned} \quad (10)$$

The superscript “*off*” means that only the terms with off-diagonal elements of \hat{A}^{phy} or $\delta^V \hat{A}$ in the Bloch representation $\{\eta \mathbf{k}\}$ are retained. $\delta^{sj} A_l$ contains η -off-diagonal interband matrix elements of \hat{A}^{phy} [23], whereas $\delta^{sj,1} A_l$ contains \mathbf{k} -off-diagonal matrix elements of $\delta^V \hat{A}$ [8]. In the weak disorder-potential regime, we only preserve $|\delta^V l\rangle$ to the lowest nonzero Born order when calculating $\delta^{sj} A_l$ and $\delta^{sj,1} A_l$. More precisely, we obtain the following expressions

$$\delta^{in} A_l = -\hbar e \mathbf{E} \cdot \sum_{l' \neq l} \delta_{\mathbf{k}\mathbf{k}'} \frac{2 \text{Im} \langle u_l | \hat{\mathbf{v}}^{phy} | u_{l'} \rangle A_{l'l'}^{phy}}{d_{ll'}^2}, \quad (11)$$

$$\begin{aligned} \delta^{sj} A_l &= \sum_{l', l'' \neq l} \frac{\langle W_{ll'} W_{l'l} \rangle A_{l'l''}^{phy}}{d_{ll'}^+ d_{ll''}^+} \\ &+ 2 \text{Re} \sum_{l' \neq l, l''} \frac{\langle W_{l'l''} W_{l'l} \rangle A_{ll''}^{phy}}{d_{ll'}^+ d_{ll''}^+}, \end{aligned} \quad (12)$$

and

$$\delta^{sj,1} A_l = 2 \text{Re} \sum_{l' \neq l} \frac{\langle \langle l | \delta^V \hat{A} | l' \rangle W_{l'l} \rangle}{d_{ll'}^+}. \quad (13)$$

The linear response of A in the weak disorder-potential regime thus reads

$$\begin{aligned} \delta A &= \sum_l \left(g_l^{(-2)} + g_l^{sk} \right) \tilde{A}_l^0 + \sum_l \left(g_l^a + g_l^{pair} \right) \tilde{A}_l^0 \\ &+ \sum_l g_l^{(-2)} \left(\delta^{sj} A_l + \delta^{sj,1} A_l \right) + \sum_l f_l^0 \left(\delta^{in} A_l + \delta^{in,1} A_l \right). \end{aligned} \quad (14)$$

The last term on the right-hand-side (rhs) is the intrinsic contribution. The second and third terms constitute the anomalous quantum contribution mentioned in the introduction, which is also independent of the disorder potential in the weak disorder-potential regime [34].

The SB formalism presented in this section is confirmed by formulating the Kohn-Luttinger density matrix transport approach designed in the weak disorder potential regime [20, 21] directly in the level of the effective quantum theory. This microscopic approach is presented in Appendix A.

A. Anomalous Hall effect

To be more specific, we analyze the anomalous Hall effect. In the level of the full Hamiltonian $\hat{\mathbf{r}} = \hat{\mathbf{r}}^{phy}$, $\delta^{sj} \mathbf{v}_l$ equals the semiclassical side-jump velocity $\mathbf{v}_l^{sj} = \sum_{l'} \omega_{ll'}^{(2)} \delta \mathbf{r}_{l'l}$ [19, 22, 23], and $\delta^{in} \mathbf{v}_l$ is just the Berry-curvature anomalous velocity $\mathbf{v}_l^B = \Omega_l^0 \hat{\mathbf{z}} \times \frac{e}{\hbar} \mathbf{E}$. Here $\delta \mathbf{r}_{l'l} = i\mathbf{J}_{l'} - i\mathbf{J}_l - \hat{\mathbf{D}} \arg V_{l'l}$ and $\Omega_l^0 = \partial_{k_x} (i\mathbf{J}_l)_y - \partial_{k_y} (i\mathbf{J}_l)_x$. In the level of the effective theory, we prove (Appendix B)

$$\delta^{sj} \mathbf{v}_l + \delta^{sj,1} \mathbf{v}_l = \sum_{l'} \omega_{ll'}^{(2)} \delta \tilde{\mathbf{r}}_{l'l} \equiv \tilde{\mathbf{v}}_l^{sj} \quad (15)$$

and (Appendix B)

$$\delta^{in} \mathbf{v}_l + \delta^{in,1} \mathbf{v}_l = \left(\Omega_l^0 + \Omega_l^a \right) \hat{\mathbf{z}} \times \frac{e}{\hbar} \mathbf{E} \equiv \tilde{\mathbf{v}}_l^B, \quad (16)$$

where $\Omega_l^a \equiv \partial_{k_x} (i\mathbf{J}_l^a)_y - \partial_{k_y} (i\mathbf{J}_l^a)_x$. Therefore, the velocity of a semiclassical carrier constructed from an effective quantum theory in the presence of external electric field and static disorder reads

$$\mathbf{v}_l = \tilde{\mathbf{v}}_l^0 + \tilde{\mathbf{v}}_l^{sj} + \tilde{\mathbf{v}}_l^B, \quad (17)$$

which contributes to the electrical current as $\mathbf{j} = e \sum_l \mathbf{v}_l f_l$. The intrinsic anomalous Hall current reads

$$\mathbf{j}^{in} = e \sum_l f_l^0 \tilde{\mathbf{v}}_l^B, \quad (18)$$

and the anomalous quantum contribution is

$$\mathbf{j}^{AQ} = \mathbf{j}^{sj} + \mathbf{j}^{ad} + \mathbf{j}^{pair}, \quad (19)$$

where $\mathbf{j}^{sj} = e \sum_l g_l^{(-2)} \tilde{\mathbf{v}}_l^{sj}$, $\mathbf{j}^{ad} = e \sum_l g_l^a \tilde{\mathbf{v}}_l^0$ and $\mathbf{j}^{pair} = e \sum_l g_l^{pair} \tilde{\mathbf{v}}_l^0$. The anomalous quantum mechanism comprises three contributions [1, 25]: (1) a component of the side-jump velocity transverse to the driving electric field; (2) an anomalous distribution function [22, 33]; (3) scattering off pairs of impurities. Both $\tilde{\mathbf{v}}_l^{sj}$ and g_l^a are related to the coordinate-shift [33], i.e., the so-called side-jump transverse to the incident wave-vector [35]. Besides, the skew scattering contribution reads

$$\mathbf{j}^{sk} = e \sum_l g_l^{sk} \tilde{\mathbf{v}}_l^0. \quad (20)$$

IV. APPLICATION 1: 2D NOZIERES-LEWINER EFFECTIVE MODEL

As the first application, we look into the spin Hall effect in the widely-studied Nozieres-Lewiner effective model where $\hat{H}_0 = \frac{\hbar^2 \mathbf{k}^2}{2m}$ in Eq. (1) [2, 6, 7, 13, 14, 16, 17]. For simplicity we only consider 2D electrons, and the random scalar disorder is modeled by $\langle V(\mathbf{r})V(\mathbf{r}') \rangle = n_{im} V_0^2 \delta(\mathbf{r} - \mathbf{r}')$ with n_{im} the impurity density and V_0 the scattering amplitude. The skew scattering is thus neglected in this section, because it has been thoroughly understood [36]. The z-component of spin is conserved and the two spin channels are independent. In this model $\mathbf{J}_l = 0$, $i\mathbf{J}_{l'}^a - i\mathbf{J}_l^a = -\hat{\mathbf{D}} \arg W_{l'l} = \frac{\eta \lambda_0^2}{4} \hat{\mathbf{z}} \times (\mathbf{k}' - \mathbf{k})$ where $(\hat{\sigma}_z)_{\eta\eta} = \eta$, and

$$\delta \tilde{\mathbf{r}}_{l'l} = 2(i\mathbf{J}_{l'}^a - i\mathbf{J}_l^a) = -2\hat{\mathbf{D}} \arg W_{l'l}. \quad (21)$$

At the same time $\delta^{sj} \mathbf{v}_l = 0$, $\delta^{in} \mathbf{v}_l = 0$ and

$$\tilde{\mathbf{v}}_l^{sj} = \delta^{sj,1} \mathbf{v}_l = \sum_{l'} \omega_{ll'}^{(2)} \delta \tilde{\mathbf{r}}_{l'l}, \quad \tilde{\mathbf{v}}_l^B = \delta^{in,1} \mathbf{v}_l = \Omega_l^a \hat{\mathbf{z}} \times \frac{e}{\hbar} \mathbf{E}.$$

Simple derivations yield $-\mathbf{j}^{z,in} = \mathbf{j}^{z,sj} = \mathbf{j}^{z,ad}$, and $\mathbf{j}^{z,pair} = 0$ in the noncrossing approximation, thus the total $o(n_{im}^0)$ contribution is $\mathbf{j}^z = \mathbf{j}^{z,ad}$.

A phenomenological Boltzmann approach has been employed to analyze this model [6, 14–16]. In that alternative Boltzmann approach, there are three basic pictorial arguments: (1) The contributions from $\delta^V \hat{\mathbf{v}}$ and $\delta^{\mathbf{E}} \hat{\mathbf{v}}$ to $\langle \hat{\mathbf{v}} \rangle_{av}$ cancel out (here $\langle \hat{\mathbf{v}} \rangle_{av} = \sum_{\mathbf{k}} f_{\eta\mathbf{k}} \mathbf{v}_{\eta\mathbf{k}}$), for $\delta^V \hat{\mathbf{v}} + \delta^{\mathbf{E}} \hat{\mathbf{v}} = 2 \frac{\lambda_0^2}{4\hbar} \hat{\sigma}_z (\nabla \hat{V} - e\mathbf{E}) \times \hat{\mathbf{z}}$ and the “net force” $e\mathbf{E} - \nabla \hat{V} = \hbar d\hat{\mathbf{k}}/dt$ acting on electrons equals zero in *steady-states* $\langle d\hat{\mathbf{k}}/dt \rangle_{av} = 0$. (2) The δ -function $\delta(\epsilon_{\mathbf{k}'}^\eta - \epsilon_{\mathbf{k}}^\eta - e\mathbf{E} \cdot \Delta \mathbf{r}_{phy})$ in the presence of the electric field and SOI. (3) The shift of the physical position $\Delta \mathbf{r}_{phy}$ during a specific scattering process from $l = (\eta, \mathbf{k})$ state to $l' = (\eta, \mathbf{k}')$ state is obtained by a time-dependent analysis: $\Delta \mathbf{r}_{phy} = \int dt \mathbf{v}_{phy}$, and the nontrivial part of $\Delta \mathbf{r}_{phy}$ that survives after average over many scatterings is $\Delta \mathbf{r}_{phy} = 2 \frac{\lambda_0^2}{4} (\hat{\sigma}_z)_{\eta\eta} \hat{\mathbf{z}} \times \int dt \frac{d\mathbf{k}}{dt} = 2 \frac{\lambda_0^2}{4} \eta \hat{\mathbf{z}} \times (\mathbf{k}' - \mathbf{k})$, due to $\hat{\mathbf{v}} = \frac{\hbar \mathbf{k}}{m} + 2 \frac{\lambda_0^2}{4} \hat{\sigma}_z \frac{d\mathbf{k}}{dt} \times \hat{\mathbf{z}}$.

Now we point out the correspondence of these three points in the present SB theory. First, point (1) is equivalent to our $\mathbf{j}^{z,in} = -\mathbf{j}^{z,sj}$. As will be detailed in Sec. VI and Appendix D, the argument in above point (1) is not always valid when $\hat{\sigma}_z$ has l -off-diagonal matrix elements. Second, the energy-conservation in the scattering process in the presence of the electric field and SOI is accounted for in the present formalism in another way different from above point (2). The effect of electric-field working $e\mathbf{E} \cdot \Delta \mathbf{r}_{phy}$ during the scattering is compensated by introducing the anomalous distribution function, and the δ -function $\delta(\epsilon_{\mathbf{k}'}^\eta - \epsilon_{\mathbf{k}}^\eta)$ for the energy-conservation

in the semiclassical scattering rate remains unchanged [1, 19, 22]. Finally, $\Delta \mathbf{r}_{phy}$ obtained in above point (3) is equal to our $\delta \tilde{\mathbf{r}}_{l'l} = 2(i\mathbf{J}_{l'}^a - i\mathbf{J}_l^a)$, with the important factor 2.

Regarding this model, there are some more theoretical debates worthwhile to be clarified.

First, if one starts from the effective Hamiltonian (2) with $\hat{H}_0 = \frac{\hbar^2 \mathbf{k}^2}{2m}$ but neglect the change of the physical position operator, then only one half of $\delta \tilde{\mathbf{r}}_{l'l}$ and thus of $\mathbf{j}^{z,ad}$ can be produced in the SB theory. This indicates the fact that the effective Hamiltonian itself is not always enough for consistent predictions with the complete effective quantum theory [4, 5].

Second, in an early influential work, Lyo and Holstein [8] claimed that Berger’s wavepacket analysis on the side-jump [35] is consistent with the original Luttinger theory [21], if in the latter the scalar disorder potential $V(\mathbf{r})$ is replaced by W . Taking into account the scattering correction to the plane-wave state ($\hat{H}_0 = \frac{\hbar^2 \mathbf{k}^2}{2m}$) in the lowest Born order, Lyo-Holstein in fact proved

$$\delta_{LH}^{sj} \mathbf{v}_l \equiv 2 \text{Re} \sum_{l' \neq l} \frac{\left\langle \left(\frac{1}{i\hbar} [\hat{\mathbf{r}}, \hat{W}] \right)_{ll'} W_{l'l} \right\rangle}{d_{ll'}^+} = \sum_{l'} \omega_{ll'}^{(2)} \delta \mathbf{r}_{l'l},$$

where $\delta \mathbf{r}_{l'l} = -\hat{\mathbf{D}} \arg W_{l'l}$. And $\delta_{LH}^{sj} \mathbf{v}_l \tau_l^{tr} = \frac{\eta \lambda_0^2}{4} \mathbf{k} \times \hat{\mathbf{z}}$ gives the sideways shift of Berger [35], with τ_l^{tr} the transport time. According to Eq. (21), the Lyo-Holstein result




FIG. 1. Side-jump diagrams for the spin Hall conductivity in the Nozieres-Lewiner effective model. Squares and circles represent spin-current ($j^{z,eq}$) and charge-current (j^{eq}) vertexes respectively. $j^{z,eq,0}$ and $j^{eq,0}$ are vertexes in the usual Kubo-Streda formalism (Eq. (C7)) whereas $j^{z,eq}$ and j^{eq} are vertexes in the generalized Kubo-Streda formalism (Eq. (C6)). (a) and (b) are spin Hall conductivities arising from the side-jump spin-current and anomalous distribution function respectively discussed in details in the main text.

is nothing else but

$$\delta_{LH}^{sj} \mathbf{v}_l = \frac{1}{2} \delta^{sj,1} \mathbf{v}_l.$$

The other half of $\delta^{sj,1} \mathbf{v}_l$ did not appear in the Lyo-Holstein theory, because the change of the physical position operator was not considered there. The argument by Lyo-Holstein under the weak disorder-potential Born approximation has been generalized directly to the level of full T-matrix [37]. But that generalization has not been confirmed by microscopic quantum transport theories.

Third, the usual Kubo-Streda theory applied to the present model [11–13] in fact starts only from the Hamiltonian $\hat{H} = \frac{\hbar^2 \mathbf{k}^2}{2m} + \hat{W} - e\mathbf{E} \cdot \hat{\mathbf{r}}$. Figure 1 with the current vertexes $j_y^{z,eq,0}$ and $j_x^{eq,0}$ shows the four well-known side-jump diagrams in the diagrammatic approach for the present model. Hereafter the external electric field is applied in x direction. In the language of the SB theory, figure 1 (a) and 1 (b) (with the current vertexes $j_y^{z,eq,0}$ and $j_x^{eq,0}$) represent the spin Hall conductivities $\sigma_{yx}^{z,0,sj} = e \sum_l \frac{\hbar}{2} \eta \left(\delta_{LH}^{sj} \mathbf{v}_l \right)_y \left(-\frac{\partial f^0}{\partial \epsilon_l} \right) \tau_l^{tr} (\mathbf{v}_l^0)_x$ and $\sigma_{yx}^{z,0,ad} = e \sum_l \frac{\hbar}{2} \eta (\mathbf{v}_l^0)_y \frac{\partial f^0}{\partial \epsilon_l} \tau_l^{tr} \left(\delta_{LH}^{sj} \mathbf{v}_l \right)_x$, respectively. Thus $\sigma_{yx}^{z,0,sj} = \sigma_{yx}^{z,0,ad} = \frac{1}{2} \sigma_{yx}^{z,ad} = \frac{1}{2} \sigma_{yx}^{z,sj}$, and the total $o(n_{im}^0)$ contribution $\sigma_{yx}^{z,0,sj} + \sigma_{yx}^{z,0,ad} = \sigma_{yx}^{z,ad}$ coincides with our SB result. However, the physical interpretation given by this diagrammatic approach is very different from that obtained in the quantum kinetic [17] or the SB theories. Being important in the quantum kinetic theory, the SOI with the driving electric field is not incorporated into the derivation of the usual Kubo-Streda formula [10, 11]. Thus for spin-orbit-induced transport in effective quantum theories, the SB and quantum kinetic theories do not always produce the same result as the usual Kubo-Streda formula. In the next section we will provide such an example.

Moreover, in Appendix C we give the slightly generalized Kubo-Streda formula in effective quantum theories, taking into account the change of the physical position operator. That formula yields the same physical picture as the SB theory. The expressions for the current vertex in the corresponding diagrams (Eq. (C6)) are different from those in the usual diagrams (Eq. (C7)), see Appendix C. Besides, the generalized Kubo-Streda formula contains an additional ‘‘Fermi sea’’ term (Eq. (C5)) which results from $\delta^{in,1} \mathbf{j}_l^z$.

V. APPLICATION 2: RASHBA 2D EFFECTIVE MODEL

In the present case $\hat{H}_0 = \frac{\hbar^2 \mathbf{k}^2}{2m} + \alpha_R \hat{\sigma} \cdot (\hat{\mathbf{k}} \times \hat{\mathbf{z}})$ in the effective Hamiltonian (1), α_R is the Rashba SOI parameter. The Rashba spinor in the internal space reads $|u_{\mathbf{k}}^\eta\rangle = \frac{1}{\sqrt{2}} (1, -i\eta \exp(i\phi))^T$. Although the Rashba SOI can be understood as arising also from $\hat{\mathbf{r}}^a$, we here take

a more general viewpoint beyond the context of direct-gap semiconductors, regarding this model as an example where the effective Hamiltonian has its own internal structure. In this model there is no unique definition of the spin current operator [38]. For the convenience of making comparison with most previous researches on this model [27–30, 39], the conventional definition of spin current operator is adopted in the present paper. The results on the conserved spin current proposed in Ref. [38] will be presented elsewhere.

The spin current polarized in z direction is $\hat{j}^z = \frac{1}{2} \left\{ \frac{\hbar}{2} \hat{\sigma}_z, \hat{\mathbf{v}} \right\} = \hat{j}^{z,phy} + \delta^V \hat{j}^z + \delta^E \hat{j}^z$, where $\delta^V \hat{j}^z + \delta^E \hat{j}^z = \frac{\lambda_0^2}{4} \hat{\mathbf{z}} \times (e\mathbf{E} - \nabla \hat{V})$. In this model $\hat{j}^{z,phy} = \frac{\hbar}{2} \frac{\hbar \mathbf{k}}{m} \hat{\sigma}_z$, then $\tilde{\mathbf{j}}_l^{z,0} = 0$, thus $\mathbf{j}^{z,sk} = \mathbf{j}^{z,pair} = \mathbf{j}^{z,ad} = 0$ and

$$\mathbf{j}^z = \sum_l f_l^0 \left(\delta^{in} \mathbf{j}_l^z + \delta^{in,1} \mathbf{j}_l^z \right) + \sum_l g_l^{(2)} \left(\delta^{sj} \mathbf{j}_l^z + \delta^{sj,1} \mathbf{j}_l^z \right).$$

As will be explained in Sec. VI, we have $\sum_l f_l^0 \delta^{in,1} \mathbf{j}_l^z = -\sum_l g_l^{(2)} \delta^{sj,1} \mathbf{j}_l^z$, then

$$\mathbf{j}^z = \sum_l f_l^0 \delta^{in} \mathbf{j}_l^z + \sum_l g_l^{(2)} \delta^{sj} \mathbf{j}_l^z, \quad (22)$$

where $\delta^{in} \mathbf{j}_l^z = -2e\mathbf{E} \cdot \text{Re} \frac{i\tilde{\mathbf{j}}_{\eta\mathbf{k},-\eta\mathbf{k}} \langle u_{\mathbf{k}}^{-\eta} | j^{z,0} | u_{\mathbf{k}}^\eta \rangle}{\epsilon_{\mathbf{k}}^\eta - \epsilon_{\mathbf{k}}^{-\eta}}$ and $\delta^{sj} \mathbf{j}_l^z$ is given by Eq. (12).

We only consider the case of both subbands partially occupied. The corresponding wave number in η band is given as $k_\eta(\epsilon) = -\eta k_R + \alpha_R^{-1} \sqrt{\epsilon_R^2 + 2\epsilon_R \epsilon}$. Here $k_R = m \frac{\alpha_R}{\hbar^2} = k_-(\epsilon) - k_+(\epsilon)$ measures the momentum splitting of two Rashba bands. We assume $(k_R/k_F)^2 \gg (\lambda_0 k_F)^2$, i.e., $(\lambda_0 k_R)^2 \gg (\lambda_0 k_F)^4$, which is the case of strong Rashba SOI. Here $k_F \equiv k_\eta(\epsilon_F) + \eta k_R$. The density of state of η band takes the form $D_\eta(\epsilon) = D_0 \frac{k_\eta(\epsilon)}{k_\eta(\epsilon) + \eta k_R}$, with $D_0 = \frac{m}{2\pi\hbar^2}$.

Equation (12) yields $\delta^{sj} \mathbf{j}_l^z = \frac{\eta}{4k_R \tau_0} \frac{\hbar}{2} \hat{\mathbf{e}}_{\mathbf{k}} \times \hat{\mathbf{z}}$ with $\tau_0^{-1} = \frac{2\pi}{\hbar} n_{im} V_0^2 D_0$, $\hat{\mathbf{e}}_{\mathbf{k}} = (\cos \phi, \sin \phi)$. The longitudinal distribution function is $g_l^{(-2)} = eE_x (-\partial_{\epsilon_l} f^0) (\mathbf{v}_l^0)_x \tau_l^{tr}$, where

$$\frac{\tau_l^{tr}(\epsilon)}{\tau_0} = \frac{D_\eta}{D_0} - \eta \frac{\lambda_0^2}{4} 4k_R k_{-\eta} + 2 \frac{\lambda_0^2}{4} (2k_R)^2 \frac{D_\eta}{D_0}. \quad (23)$$

Thus

$$\frac{\sum_l f_l^0 (\delta^{in} \mathbf{j}_l^z)_y}{E_x} = \frac{-e}{8\pi} - \frac{en_e \lambda_0^2}{4} \left(\frac{1}{2} - \frac{\frac{1}{6} \epsilon_R}{\epsilon_F + \epsilon_R} \right), \quad (24)$$

$$\frac{\sum_l g_l^{(-2)} (\delta^{sj} \mathbf{j}_l^z)_y}{E_x} = \frac{e}{8\pi} + \frac{en_e \lambda_0^2}{4} \left(\frac{1}{2} + \frac{\frac{1}{2} \epsilon_R}{\epsilon_F + \epsilon_R} \right), \quad (25)$$

and then

$$j_y^z = \frac{en_e \lambda_0^2}{4} \frac{\frac{2}{3} \epsilon_R}{\epsilon_F + \epsilon_R} E_x = (\lambda_0 k_R)^2 \frac{e}{6\pi} E_x. \quad (26)$$

This nonzero spin Hall conductivity of order $(\lambda_0 k_R)^2$ did not appear in previous theoretical researches [27, 28, 39] assuming weak Rashba SOI. Strong Rashba SOI energy compared to the Fermi energy is possible, e.g., in heterostructures of non-centrosymmetric semiconductor BiTeX (X=Cl, Br and I) family [40]. Because we do not assume $k_R \ll k_F$, in the case of strong Rashba SOI the magnitude of the above spin Hall conductivity can be comparable to the side-jump spin Hall conductivity in the 2D Nozieres-Lewiner model.

The SB formalism provides alternative explanations to the results in a recent quantum kinetic theory [28]. The second term on the rhs of Eq. (24) arises from the SOI with the external driving electric field, and corresponds to the contribution called ‘‘anomalous spin precession from electric field’’ in Ref. [28]. Whereas the second term on the rhs of Eq. (25) arises from the spin-orbit-scattering-induced correction to the transport time in Eq. (23), and corresponds to the contribution called ‘‘anomalous spin precession from impurities’’ in Ref. [28]. The $o((\lambda_0 k_R)^2)$ contribution is neglected in Ref. [28], where the weak Rashba SOI was assumed.

Now we consider the application of the usual Kubo-Streda formula, which in fact treats the Hamiltonian $\hat{H} = \hat{H}_0 + \hat{W} - e\mathbf{E} \cdot \hat{\mathbf{r}}$, as in Ref. [30]. In the weak disorder-potential regime the diagram calculation can be done in the band-eigenstate representation, where the correspondence to the SB formalism is apparent [1, 23, 25]. In the




FIG. 2. Diagrams for the non-zero contributions to the spin Hall conductivity in the usual Kubo-Streda formalism. Squares represent spin-current ($J^{z,eq,0}$) vertexes. Circles and filled circles represent bare charge-current ($J^{eq,0}$) and renormalized charge-current ($J^{eq,0}$) vertexes respectively. (a) describes intrinsic contributions; (b) describes side-jump contributions from interband coherence response due to band structure SOI; (c) describes side-jump contributions from disorder-induced corrections to the charge-current operator.

language of the SB theory, the diagrams in Fig. 2 yield

$$\sigma_{yx}^{z,0} = \sum_l f_l^0 \frac{(\delta_0^{in} \mathbf{J}_l^z)_y}{E_x} + \sum_l \left(\delta_0^{sj} \mathbf{J}_l^z + \delta_0^{sj,1} \mathbf{J}_l^z \right)_y \frac{g_l^{(2)}}{E_x}, \quad (27)$$

where $\delta_0^{in} \mathbf{J}_l^z = -2e\mathbf{E} \cdot \text{Re} \frac{i\mathbf{J}_{\eta\mathbf{k}, -\eta\mathbf{k}} \langle u_{\mathbf{k}}^{-\eta} | \hat{J}^{z,0} | u_{\mathbf{k}}^{\eta} \rangle}{\epsilon_{\mathbf{k}}^{\eta} - \epsilon_{\mathbf{k}}^{-\eta}}$ and

$$\delta_0^{sj} \mathbf{J}_l^z = \sum_{l', l'' \neq l} \frac{\langle W_{ll'} W_{l'l''} \rangle \mathbf{J}_{l'l''}^{z,0}}{d_{ll'}^- d_{l'l''}^+} + 2\text{Re} \sum_{l' \neq l, l''} \frac{\langle W_{ll'} W_{l'l''} \rangle \mathbf{J}_{l'l''}^{z,0}}{d_{ll'}^+ d_{l'l''}^+},$$

$$\delta_0^{sj,1} \mathbf{J}_l^z = 2\text{Re} \sum_{l' \neq l} \frac{\left\langle |l| \frac{1}{2} \left\{ \frac{\hbar}{2} \hat{\sigma}_z, \frac{1}{i\hbar} [\hat{\mathbf{r}}, \hat{W}] \right\} |l'\rangle W_{ll'} \right\rangle}{d_{ll'}^+}.$$

The first term on the rhs of Eq. (27) corresponds to Fig. 2 (a) ($\sigma_{yx}^{z,II,0} = 0$ in the considered model, see Appendix C), whereas the contributions from $\delta_0^{sj} \mathbf{J}_l^z$ and $\delta_0^{sj,1} \mathbf{J}_l^z$ in the second term on the rhs of Eq. (27) correspond to Fig. 2 (b) and 2 (c), respectively. One can find that $\delta_0^{in} \mathbf{J}_l^z \neq \delta^{in} \mathbf{J}_l^z$, $\delta_0^{sj} \mathbf{J}_l^z = \delta^{sj} \mathbf{J}_l^z$, $\delta_0^{sj,1} \mathbf{J}_l^z = \frac{1}{2} \delta^{sj,1} \mathbf{J}_l^z$ up to the first order of λ_0^2 for the present model. Regarding the spin Hall conductivity j_y^z/E_x we have $\sum_l f_l^0 \frac{(\delta_0^{in} \mathbf{J}_l^z)_y}{E_x} = \frac{-e}{8\pi}$, $\sum_l \left(\delta_0^{sj,1} \mathbf{J}_l^z \right)_y \frac{g_l^{(2)}}{E_x} = -\frac{1}{2} \frac{en_e \lambda_0^2}{4}$, and $\sum_l \left(\delta_0^{sj} \mathbf{J}_l^z \right)_y \frac{g_l^{(2)}}{E_x}$ is the same as Eq. (25). In the calculation we used $\left(\delta_0^{sj,1} \mathbf{J}_l^z \right)_y = -\frac{\hbar}{2} \frac{\lambda_0^2}{4\tau_0} (k_\eta + \eta k_R) \cos \phi$. Then $\sigma_{yx}^{z,0} = (\lambda_0 k_R)^2 \frac{e}{8\pi}$. This result differs from Eq. (26) obtained for the complete effective model.

VI. DISCUSSION AND SUMMARY

We discuss the validity of the widely accepted idea that the effects from corrections to the current operator due to the driving electric field and the gradient of disorder potential always cancel out [14, 16, 17, 28]. This idea indicates $\sum_l g_l^{(-2)} \delta^{sj,1} \mathbf{v}_l = -\sum_l f_l^0 \delta^{in,1} \mathbf{v}_l$ and $\sum_l g_l^{(-2)} \delta^{sj,1} \mathbf{j}_l^z = -\sum_l f_l^0 \delta^{in,1} \mathbf{j}_l^z$. A pictorial argument of this idea is based on the correspondence principle (Ehrenfest’s theorem) [14, 16, 17], as mentioned in Sec. IV. For the Nozieres-Lewiner effective model ($\hat{H}_0 = \frac{\hbar^2 \mathbf{k}^2}{2m}$), this idea has been confirmed by a quantum kinetic theory [17]. However, notice that $\hat{\sigma}_z$ is also involved in $\delta^V \hat{\mathbf{v}} + \delta^{\mathbf{E}} \hat{\mathbf{v}} = 2 \frac{\lambda_0^2}{4\hbar} \hat{\sigma}_z (\nabla \hat{V} - e\mathbf{E}) \times \hat{\mathbf{z}}$ (for 2D electrons). When the off-diagonal matrix elements of $\hat{\sigma}_z$ in the band-eigenstate basis is not zero, the Ehrenfest’s theorem cannot be employed directly. Thus $\sum_l g_l^{(-2)} \delta^{sj,1} \mathbf{v}_l = -\sum_l f_l^0 \delta^{in,1} \mathbf{v}_l$ is not generally valid even for 2D electrons. In Appendix D we confirm this observation by the SB formalism.

On the other hand, considering the conventional spin current polarized in z direction for 2D electrons, $\delta^V \hat{j}^z +$

$\delta^{\mathbf{E}}\hat{J}^z = \frac{\lambda_0^2}{4}\hat{\mathbf{z}} \times (e\mathbf{E} - \nabla\hat{V})$ is free from the above problem. Thus the Ehrenfest's theorem can be applied, validating the pictorial argument [14] for 2D electrons even with band-structure SOI. Specifically, one has $\sum_l f_l^0 (\delta^{in,1}\mathbf{j}_l^z)_y = E_x \frac{e\lambda_0^2}{4} \sum_l f_l^0 = E_x \frac{en_e\lambda_0^2}{4}$ and $\sum_l g_l^{(-2)} (\delta^{sj,1}\mathbf{j}_l^z)_y = 2\frac{\hbar}{2} \sum_{l'V} g_l^{(-2)} \omega_{l'l}^{(2)} \frac{\lambda_0^2}{4} (k'_x - k_x) = \frac{\lambda_0^2}{4} eE_x \sum_l \partial_{k_x} f_l^0 k_x = -\frac{en_e\lambda_0^2}{4} E_x$, where we use the same manipulations as in Eq. (D2).

We note that the special form $\hat{\mathbf{r}}^a = \frac{\lambda_0^2}{4}\hat{\sigma}_z\hat{\mathbf{z}} \times \hat{\mathbf{k}}$ which is linear in the momentum for 2D electrons plays the vital role in the above derivation. But for the conventional spin current in 2D hole systems where $\hat{\mathbf{r}}^a$ is cubic in the momentum, we cannot get the relation $\sum_l g_l^{(-2)} \delta^{sj,1}\mathbf{j}_l^z = -\sum_l f_l^0 \delta^{in,1}\mathbf{j}_l^z$ if no further assumption about the model Hamiltonian is made. Detailed discussions are presented in Appendix D.

In summary, we constructed the SB transport framework directly in the level of the effective quantum theory, confirmed by a generalized Kohn-Luttinger density matrix transport theory also in this level. It was shown that the spin/anomalous Hall effect studied in this level can still be parsed into the same categories as in the level of the full Hamiltonian, in the regime where the SB theory works. We discussed the link and difference between the present SB theory and various previous theories. To help clarify this issue, we also derived the slightly generalized Kubo-Streda formula in the level of the effective quantum theory. This formula leads to the same physical picture as the present SB theory. In a Rashba 2D effective model, a nonzero spin Hall effect important in the case of strong Rashba SOI but neglected in previous theories has been found.

ACKNOWLEDGMENTS

We acknowledge insightful discussions with Q. Niu. C. X. thanks P. Streda and D. Culcer for useful discussions. C. X. and B. X. are supported by DOE (DE-FG03-02ER45958, Division of Materials Science and Engineering), NSF (EFMA-1641101) and Welch Foundation (F-1255). F. X. is supported by the Department of Energy, Office of Basic Energy Sciences under Contract No. DE-FG02-ER45958 and by the Welch foundation under Grant No. TBF1473. The formulation in Sec. III is supported by the DOE grant.

Appendix A: Confirmation of the SB formalism: Kohn-Luttinger density-matrix approach

The original Kohn-Luttinger approach is designed in the level of the full Hamiltonian. Here we formulate a generalized version of it in the level of the effective quantum theory.

The single-carrier Hamiltonian reads $\hat{H}_T = \hat{H}_0 + \hat{W} + \hat{H}_F$, where the field term $\hat{H}_F = \hat{H}_1 e^{st}$ arises from the electric field turned on adiabatically from the remote past $t = -\infty$, with $\hat{H}_1 = -e\mathbf{E} \cdot \hat{\mathbf{r}}^{phy}$. The expectation value of a single-carrier operator \hat{A} is $\langle A \rangle = tr [\hat{A}\hat{\rho}_T]$, where tr denotes the trace operation in the single-carrier Hilbert space, and the single-carrier density matrix $\hat{\rho}_T$ is determined by the quantum Liouville equation $i\hbar \frac{\partial}{\partial t} \hat{\rho}_T = [\hat{H}_T, \hat{\rho}_T]$. In the linear response regime $\hat{\rho}_T = \hat{\rho} + \hat{f}e^{st}$, where $\hat{\rho}$ is the equilibrium density matrix, \hat{f} is linear in the electric field and independent of time. In the band-eigenstate representation of \hat{H}_0 :

$$d_{l'V}^- f_{l'V} = [\hat{f}, \hat{H}_1]_{l'V} + [\hat{\rho}, \hat{H}_1]_{l'V}. \quad (\text{A1})$$

Here $[\hat{A}, \hat{B}]_{l'V} \equiv \sum_{V''} (A_{l'V''} B_{V''V} - B_{l'V''} A_{V''V})$. Defining $C_{l'V} \equiv [\hat{\rho}, \hat{H}_1]_{l'V}$, we have $C_{l'V} = ie\mathbf{E} \cdot [\hat{\mathbf{D}}\rho_{l'V} + [\tilde{\mathbf{J}}, \rho]_{l'V}]$ for $l' \neq l$ and $C_l = ie\mathbf{E} \cdot [\partial_{\mathbf{k}}\rho_l + [\tilde{\mathbf{J}}, \rho]_l]$. Hereafter $\rho_{l'V} \equiv \rho_{l'V}$, $f_{l'V} \equiv f_{l'V}$ and $C_{l'V} \equiv C_{l'V}$. In the effective quantum theory, in the presence of the external electric field \hat{A} may be different from its equilibrium form \hat{A}^{eq} , so $\hat{A} = \hat{A}^{eq} + \delta^{\mathbf{E}}\hat{A}$. The linear response in the weak disorder-potential regime is thus

$$\begin{aligned} \delta A &= tr \langle \hat{f}\hat{A}^{eq} \rangle + tr \langle \hat{\rho}\delta^{\mathbf{E}}\hat{A} \rangle \\ &= \sum_{l'V} \langle f_{l'V} A_{l'V}^{eq} \rangle + \sum_{l'V} \langle \rho_{l'V} \rangle (\delta^{\mathbf{E}}A)_{l'V} \\ &= \sum_l \langle f_l \rangle A_l^{phy} + \sum_{l'V} \langle f_{l'V} A_{l'V}^{eq} \rangle + \sum_l \rho_l^0 (\delta^{\mathbf{E}}\hat{A})_{ll} \end{aligned} \quad (\text{A2})$$

Hereafter the notation \sum' means that all the index equalities should be avoided in the summation.

In the weak disorder-potential regime an iterative solution to Eq. (A1) in ascending powers of the disorder potential is possible:

$$\begin{aligned} f_l &= f_l^{(-2)} + f_l^{(-1)} + f_l^{(0)} + \dots, \\ f_{l'V} &= f_{l'V}^{(-1)} + f_{l'V}^{(0)} + f_{l'V}^{(1)} \dots \quad (l \neq l'), \\ C_{l'V} &= C_{l'V}^{(0)} + C_{l'V}^{(1)} + C_{l'V}^{(2)} + \dots \end{aligned}$$

Then one gets an equation for f_l , as well as expressions for $f_{l'V}$ ($l' \neq l$) in terms of f_l [21]:

$$f_{l'V}^{(-1)} = \frac{f_l^{(-2)} - f_{l'V}^{(-2)}}{d_{l'V}^-} W_{l'V},$$

$$\begin{aligned} f_{l'V}^{(0)} &= \sum_{l'' \neq l, V''} \frac{W_{l'V''} W_{V''V}}{d_{l'V}^-} \left[\frac{f_l^{(-2)} - f_{l''V''}^{(-2)}}{d_{l''V''}^-} - \frac{f_{l''V''}^{(-2)} - f_l^{(-2)}}{d_{l''V''}^-} \right] \\ &+ \frac{f_l^{(-1)} - f_{l'V}^{(-1)}}{d_{l'V}^-} W_{l'V} + \frac{C_{l'V}^{(0)}}{d_{l'V}^-}. \end{aligned}$$

After disorder average one obtains a transport equation for $\langle f_l \rangle$. Comparing Eq. (A3) in Ref. [24] to the present Eq. (A1), one can see that they take the same form, except that $\hat{V} \rightarrow \hat{W}$ and $\mathbf{J}_{ll'} \rightarrow \tilde{\mathbf{J}}_{ll'}$. Thus the derivation of the transport equation for $\langle f_l \rangle$ remains unchanged. Assuming isotropic systems $\sum_{l'} \omega_{ll'}^a = 0$, one has

$$\begin{aligned} 0 &= \frac{1}{i\hbar} C_l^{(0)} + \sum_{l'} \omega_{ll'}^{(2)} \langle f_l^{(-2)} - f_{l'}^{(-2)} \rangle \\ &+ \sum_{l'} \omega_{ll'}^{(2)} \langle f_l^{(-1)} - f_{l'}^{(-1)} \rangle + \sum_{l'} \omega_{ll'}^{(3)} \langle f_l^{(-2)} - f_{l'}^{(-2)} \rangle \\ &+ \sum_{l'} \omega_{ll'}^{(2)} \langle f_l^{(0)} - f_{l'}^{(0)} \rangle + \sum_{l'} \omega_{ll'}^{(3)} \langle f_l^{(-1)} - f_{l'}^{(-1)} \rangle \\ &+ \sum_{l'} \left[\omega_{ll'}^{(4)} + S_{ll'}^{(4)} \right] \langle f_l^{(-2)} - f_{l'}^{(-2)} \rangle + \frac{1}{i\hbar} C_l'', \end{aligned}$$

where the expression [20, 21] for $S_{ll'}^{(4)} = S_{l'l}^{(4)}$ is not necessary here. The symmetric parts of the higher-order scattering rates only contribute to trivial renormalizations to the longitudinal transport, and thus are negligible in the case of weak disorder potential. $\frac{1}{i\hbar} C_l^{(0)} = \frac{1}{\hbar} e\mathbf{E} \cdot \partial_{\mathbf{k}} \rho_l^{(0)}$ is the driving term of the conventional Boltzmann equation, where $\rho_l^{(0)}$ is just the Fermi distribution function [20]. C_l'' contains the combination effects of the electric field and disorder, whose original expression (is of the second order of disorder potential) is given in Ref. [21]. Here the point is that, except the anomalous part $C_{ll'}^{a,a} = -i\hbar e\mathbf{E} \cdot \sum_{l''} \omega_{ll''}^{(2)} \delta \tilde{\mathbf{r}}_{ll''} \partial_{\epsilon_l} \rho_{l''}^{(0)}$, all other terms of $C_{ll'}''$ are present even in the absence of SOI and are trivial renormalizations to the conventional driving term. Therefore, regarding the spin/anomalous Hall effect, the qualitatively and quantitatively important contributions to $\langle f_l \rangle$ in the weak disorder-potential regime are given by

$$\begin{aligned} 0 &= \left[e\mathbf{E} \cdot \mathbf{v}_l^0 \partial_{\epsilon_l} \rho_l^{(0)} + \sum_{l'} \omega_{ll'}^{(2)} \langle f_l^{(-2)} - f_{l'}^{(-2)} \rangle \right] \\ &+ \left[\sum_{l'} \omega_{ll'}^{(2)} \langle f_l^{(-1)} - f_{l'}^{(-1)} \rangle + \sum_{l'} \omega_{ll'}^{3a} \langle f_l^{(-2)} - f_{l'}^{(-2)} \rangle \right] \\ &+ \left[\sum_{l'} \omega_{ll'}^{(2)} \langle f_l^{(0),n} - f_{l'}^{(0),n} \rangle + \sum_{l'} \omega_{ll'}^{3a} \langle f_l^{(-1)} - f_{l'}^{(-1)} \rangle \right. \\ &\quad \left. + \sum_{l'} \omega_{ll'}^{Aa} \langle f_l^{(-2)} - f_{l'}^{(-2)} \rangle \right] \\ &+ \left[\sum_{l'} \omega_{ll'}^{(2)} \langle f_l^{(0),a} - f_{l'}^{(0),a} \rangle - e\mathbf{E} \cdot \sum_{l'} \omega_{ll'}^{(2)} \delta \tilde{\mathbf{r}}_{ll'} \partial_{\epsilon_l} \rho_l^{(0)} \right]. \end{aligned}$$

Here we split $f_l^{(0)}$ into $f_l^{(0)} = f_l^{(0),n} + f_l^{(0),a}$. Setting $\langle f_l^{(-2)} \rangle \equiv g_l^{(-2)}$, $\langle f_l^{(-1)} \rangle \equiv g_l^{(-1)}$, $\langle f_l^{(0),n} \rangle \equiv g_l^{(0)}$ and $\langle f_l^{(0),a} \rangle \equiv g_l^a$, the above equation is just the form of the SB equations (5,6) used in practice [1, 19].

Because the Kohn-Luttinger expansion is basically a bare perturbation theory, some trivial renormalization terms are unavoidable in high orders of this expansion. These terms should be eliminated systematically by a renormalization procedure, if one aims at placing the Kohn-Luttinger theory as a generic foundation for the extended-state transport phenomena. This kind of renormalization treatment has been shown for free electrons without SOI [41]. Although a more complicated procedure is expected to be applicable also in the presence of SOI, it has never been done according to our literature knowledge. In fact, the Kubo linear response approach may be more suitable to serve as the foundation of the extended-state transport, into which the systematic renormalization procedure can be incorporated. On the other hand, we only regard the Kohn-Luttinger approach as a foundation of the SB theory in the case of weak disorder potential. In this case the aforementioned renormalization effects are just much smaller high-order corrections to the longitudinal transport, and can thus be neglected in the Boltzmann regime.

In the weak disorder-potential regime, after disorder average, one has

$$\begin{aligned} \sum_{l'} \langle f_{ll'} A_{l'l}^{eq} \rangle &= \sum_{l'} \langle f_{ll'}^{(0)} \rangle \langle A_{l'l}^{eq} \rangle + \sum_{l'} \langle f_{ll'}^{(-1)} A_{l'l}^{eq} \rangle \\ &= \sum_{l'} C_{ll'}^{(0)} \frac{\langle A_{l'l}^{eq} \rangle}{d_{ll'}^-} \\ &+ \sum_{ll''} \langle W_{ll''} W_{l''l'} \rangle \left\langle \frac{f_l^{(-2)} - f_{l''}^{(-2)}}{d_{ll''}^-} - \frac{f_{l''}^{(-2)} - f_{l'}^{(-2)}}{d_{l''l'}^-} \right\rangle \frac{\langle A_{l'l}^{eq} \rangle}{d_{ll'}^-} \\ &\quad + 2 \text{Re} \sum_{l'} \langle f_l^{(-2)} \rangle \left\langle \frac{W_{l'l} A_{ll'}^{eq}}{d_{ll'}^+} \right\rangle \quad (\text{A3}) \end{aligned}$$

Due to $C_{ll'}^{(0)} = ie\mathbf{E} \cdot \tilde{\mathbf{J}}_{ll'} (\rho_{l'}^{(0)} - \rho_l^{(0)})$ and $\mathbf{v}_{ll'}^{phy} \delta_{\mathbf{k}\mathbf{k}'} = -\frac{1}{i\hbar} (\epsilon_l - \epsilon_{l'}) i\tilde{\mathbf{J}}_{ll'}$ ($l \neq l'$), we have

$$\sum_{l'} C_{ll'}^{(0)} \frac{\langle A_{l'l}^{eq} \rangle}{d_{ll'}^-} = \sum_l f_l^0 \delta^{in} A_l, \quad (\text{A4})$$

$$\begin{aligned} \sum_{ll''} \langle W_{ll''} W_{l''l'} \rangle \left\langle \frac{f_l^{(-2)} - f_{l''}^{(-2)}}{d_{ll''}^-} - \frac{f_{l''}^{(-2)} - f_{l'}^{(-2)}}{d_{l''l'}^-} \right\rangle \frac{\langle A_{l'l}^{eq} \rangle}{d_{ll'}^-} \\ \equiv \sum_l \langle f_l^{(-2)} \rangle \delta^{sj} A_l, \quad (\text{A5}) \end{aligned}$$

and $(\hat{A}^{eq} = \hat{A}^{phy} + \delta^V \hat{A})$

$$2 \text{Re} \sum_{l'} \langle f_l^{(-2)} \rangle \left\langle \frac{W_{l'l} A_{ll'}^{eq}}{d_{ll'}^+} \right\rangle = \sum_l \langle f_l^{(-2)} \rangle \delta^{sj,1} A_l, \quad (\text{A6})$$

with $\delta^{in} A_l$, $\delta^{sj} A_l$ and $\delta^{sj,1} A_l$ given by Sec. III.

Besides, $\sum_l \rho_l^{(0)} \left(\delta^{\mathbf{E}} \hat{A} \right)_u = \sum_l f_l^0 \delta^{in,1} A_l$, thus

$$\begin{aligned} & \sum_{l'} \langle f_{l'} A_{l'}^{eq} \rangle + \sum_l \rho_l^0 \left(\delta^{\mathbf{E}} \hat{A} \right)_u \\ &= \sum_l f_l^0 \left(\delta^{in} A_l + \delta^{in,1} A_l \right) + \sum_l g_l^{(-2)} \left(\delta^{sj} A_l + \delta^{sj,1} A_l \right). \end{aligned} \quad (\text{A7})$$

Appendix B: Proof of Eqs. (15) and (16)

1. Proof of Eq. (15)

Following the same route shown in Appendix A of Ref. [23], we get

$$\begin{aligned} \delta^{sj} \mathbf{v}_l &= \text{Re} \sum_{l'} \frac{2}{\hbar} \frac{\left\langle W_{l'} \left[W, \tilde{\mathbf{J}} \right]_{l'l} \right\rangle_c}{d_{l'}^-} \\ &+ \sum_{l'} \frac{2\pi}{\hbar} \left\langle |W_{l'}|^2 \right\rangle_c \delta(d_{l'}) \left[i\tilde{\mathbf{J}}_{l'} - i\tilde{\mathbf{J}}_l \right] \end{aligned} \quad (\text{B1})$$

with $\left[W, \tilde{\mathbf{J}} \right]_{l'l} \equiv \sum_{l''} \left[W_{l'l''} \tilde{\mathbf{J}}_{l''l} - \tilde{\mathbf{J}}_{l'l''} W_{l''l} \right]$. In the present case we have

$$\left[W, \tilde{\mathbf{J}} \right]_{l'l} = -i \left[\hat{W}, \hat{\mathbf{r}}^{phy} \right]_{l'l} + \hat{\mathbf{D}} W_{l'l}, \quad (\text{B2})$$

then

$$\begin{aligned} \delta^{sj} \mathbf{v}_l &= \text{Re} \sum_{l'} \frac{2}{\hbar} \frac{\left\langle W_{l'} \hat{\mathbf{D}} W_{l'l} \right\rangle_c}{d_{l'}^-} \\ &+ \sum_{l'} \frac{2\pi}{\hbar} \left\langle |W_{l'}|^2 \right\rangle_c \delta(d_{l'}) \left[i\tilde{\mathbf{J}}_{l'} - i\tilde{\mathbf{J}}_l \right] \\ &- \text{Re} \sum_{l'} 2 \frac{\left\langle W_{l'} \left(\frac{1}{i\hbar} \left[\hat{\mathbf{r}}^{phy}, \hat{W} \right] \right)_{l'l} \right\rangle_c}{d_{l'}^-}. \end{aligned}$$

The first term on the rhs can be split into two terms with one related to $\text{Im} \left\langle W_{l'} \hat{\mathbf{D}} W_{l'l} \right\rangle_c = \left\langle |W_{l'}|^2 \right\rangle_c \hat{\mathbf{D}} \arg W_{l'l}$ and the other related to $\hat{\mathbf{D}} \left\langle |W_{l'}|^2 \right\rangle_c$. The first one is related to the phase of the disorder potential and is thus nontrivial. While the latter one, which does not break any symmetry, is just a trivial renormalization to \mathbf{v}_l^0 . It does not contribute to the Hall current in the weak disorder-potential limit and can be ignored. Thus one has

$$\begin{aligned} \delta^{sj} \mathbf{v}_l &= \sum_{l'} \frac{2\pi}{\hbar} \left\langle |W_{l'}|^2 \right\rangle_c \delta(d_{l'}) \left[i\tilde{\mathbf{J}}_{l'} - i\tilde{\mathbf{J}}_l - \hat{\mathbf{D}} \arg W_{l'l} \right] \\ &- \text{Re} \sum_{l'} 2 \frac{\left\langle W_{l'} \left(\frac{1}{i\hbar} \left[\hat{\mathbf{r}}^{phy}, \hat{W} \right] \right)_{l'l} \right\rangle_c}{d_{l'}^-} \end{aligned} \quad (\text{B3})$$

On the other hand

$$\delta^{sj,1} \mathbf{v}_l = 2 \text{Re} \sum_{l' \neq l} \frac{\left\langle \left(\frac{1}{i\hbar} \left[\hat{\mathbf{r}}^{phy}, \hat{W} \right] \right)_{l'} W_{l'l} \right\rangle_c}{d_{l'}^+}, \quad (\text{B4})$$

thus $\delta^{sj} \mathbf{v}_l + \delta^{sj,1} \mathbf{v}_l = \sum_{l'} \omega_{l'}^{(2)} \delta \tilde{\mathbf{r}}_{l'l}$.

2. Proof of Eq. (16)

We choose the external in-plane electric field in x direction, then

$$\begin{aligned} \left(\delta^{in} v_l \right)_y &= -\hbar e E_x \sum_{l' \neq l} \delta_{\mathbf{k}\mathbf{k}'} \frac{2 \text{Im} \langle u_l | \hat{v}_x^{phy} | u_{l'} \rangle \langle u_{l'} | \hat{v}_y^{phy} | u_l \rangle}{d_{l'}^2} \\ &= -\frac{e}{\hbar} E_x \sum_{\eta' \neq \eta} 2 \text{Im} \left[\langle u_{\mathbf{k}}^\eta | (i\partial_{k_x} + \hat{x}^a) | u_{\mathbf{k}}^{\eta'} \rangle \right. \\ &\quad \left. \langle u_{\mathbf{k}}^{\eta'} | (i\partial_{k_y} + \hat{y}^a) | u_{\mathbf{k}}^\eta \rangle \right] \\ &= \frac{e}{\hbar} E_x 2 \text{Im} \langle \partial_{k_y} u_{\mathbf{k}}^\eta | \partial_{k_x} u_{\mathbf{k}}^\eta \rangle - \frac{e}{\hbar} E_x 2 \text{Im} \langle u_{\mathbf{k}}^\eta | \hat{x}^a \hat{y}^a | u_{\mathbf{k}}^\eta \rangle \\ &\quad - \frac{e}{\hbar} E_x 2 \text{Re} \left[\langle u_{\mathbf{k}}^\eta | \hat{x}^a | \partial_{k_y} u_{\mathbf{k}}^\eta \rangle - \langle \partial_{k_x} u_{\mathbf{k}}^\eta | \hat{y}^a | u_{\mathbf{k}}^\eta \rangle \right] \\ &= \frac{e}{\hbar} E_x 2 \text{Im} \langle \partial_{k_y} u_{\mathbf{k}}^\eta | \partial_{k_x} u_{\mathbf{k}}^\eta \rangle \\ &\quad + \frac{e}{\hbar} E_x \left[\partial_{k_x} \langle u_{\mathbf{k}}^\eta | \hat{y}^a | u_{\mathbf{k}}^\eta \rangle - \partial_{k_y} \langle u_{\mathbf{k}}^\eta | \hat{x}^a | u_{\mathbf{k}}^\eta \rangle \right] \\ &\quad - \frac{e}{\hbar} E_x \langle u_{\mathbf{k}}^\eta | (\partial_{k_x} \hat{y}^a - \partial_{k_y} \hat{x}^a) | u_{\mathbf{k}}^\eta \rangle \\ &\quad + i \frac{e}{\hbar} E_x \langle u_{\mathbf{k}}^\eta | [\hat{x}^a, \hat{y}^a] | u_{\mathbf{k}}^\eta \rangle \end{aligned} \quad (\text{B5})$$

On the other hand, $(\delta^{in,1} v_l)_y = (\delta^{\mathbf{E}} \hat{v}_y)_l$ where

$$\begin{aligned} \delta^{\mathbf{E}} \hat{v}_y &= \frac{1}{i\hbar} \left[\hat{y}^{phy}, -e E_x \hat{x}^{phy} \right] \\ &= \frac{e E_x}{\hbar} \left(\frac{\partial \hat{y}^a}{\partial \hat{k}_x} - \frac{\partial \hat{x}^a}{\partial \hat{k}_y} \right) - i \frac{e E_x}{\hbar} [\hat{x}^a, \hat{y}^a], \end{aligned} \quad (\text{B6})$$

thus

$$\begin{aligned} \left(\delta^{in} v_l \right)_y + \left(\delta^{in,1} v_l \right)_y &= \frac{e E_x}{\hbar} \left\{ 2 \text{Im} \langle \partial_{k_y} u_l | \partial_{k_x} u_l \rangle + \right. \\ &\quad \left. \partial_{k_x} \langle u_{\mathbf{k}}^\eta | \hat{y}^a | u_{\mathbf{k}}^\eta \rangle - \partial_{k_y} \langle u_{\mathbf{k}}^\eta | \hat{x}^a | u_{\mathbf{k}}^\eta \rangle \right\} \\ &= \frac{e E_x}{\hbar} \left(\Omega_l^0 + \Omega_l^a \right). \end{aligned} \quad (\text{B7})$$

Appendix C: Kubo-Streda formula in effective quantum theories

In the linear response Eq. (A2) $\delta A = \text{tr} \left\langle \hat{A}^{eq} \hat{f} \right\rangle + \text{tr} \left\langle \hat{\rho} \delta^{\mathbf{E}} \hat{A} \right\rangle$ to the dc uniform external electric field, the first term on the rhs can be found by the Kubo-Streda

formula [10, 11] in terms of the $\hat{A} - \hat{j}$ correlation function, in the presence of static impurities. Starting from the single-particle quantum Liouville equation, one has [10]

$$\hat{f} = \frac{i}{\hbar} \lim_{s \rightarrow 0^+} \int_{-\infty}^0 dt e^{st} e^{i\hat{H}^{eq}t/\hbar} [\hat{\rho}, \hat{H}_1] e^{-i\hat{H}^{eq}t/\hbar}, \quad (\text{C1})$$

where \hat{H}^{eq} is the equilibrium single-carrier Hamiltonian in the presence of disorder. When the physical position operator is just the canonical one, $\hat{H}^{eq} = \hat{H}_0 + \hat{V}(\hat{\mathbf{r}})$ and $\hat{H}_1 = -e\mathbf{E} \cdot \hat{\mathbf{r}}$, then Eq. (C1) leads to the well-known Kubo-Streda formula for the $\hat{A} - \hat{j}$ correlation function [1, 2]. In this case $\delta^{\mathbf{E}}\hat{A} = 0$ when \hat{A} is not the thermal current operator, thus $\delta A = \text{tr} \langle \hat{A}^{eq} \hat{f} \rangle$ is completely determined by the $\hat{A} - \hat{j}$ correlation function.

In effective quantum theories where the physical position operator is not the canonical one, $\hat{H}^{eq} = \hat{H}_0 + \hat{V}(\hat{\mathbf{r}}^{phy})$, $\hat{H}_1 = -e\mathbf{E} \cdot \hat{\mathbf{r}}^{phy}$ and $\delta^{\mathbf{E}}\hat{A} \neq 0$ even when \hat{A} is the electric/spin current operator. In this case we derive the slightly generalized Kubo-Streda formula for δA , by obtaining first the Bastin formula by the method in Ref. [10] and then following the manipulations presented in Ref. [11]. At the low-temperature limit the generalized Kubo-Streda formula for the spin Hall conductivity reads $\sigma_{yx}^z = \sigma_{yx}^{z,I(a)} + \sigma_{yx}^{z,I(b)} + \sigma_{yx}^{z,II} + \sigma_{yx}^{z,in-1}$, where

$$\sigma_{yx}^{z,I(a)} = \frac{\hbar}{2\pi} \text{tr} \left\langle \hat{j}_y^{z,eq} \hat{G}^R(\epsilon_F) \hat{j}_x^{eq} \hat{G}^A(\epsilon_F) \right\rangle, \quad (\text{C2})$$

$$\sigma_{yx}^{z,I(b)} = -\frac{\hbar}{2\pi} \text{Re tr} \left\langle \hat{j}_y^{z,eq} \hat{G}^R(\epsilon_F) \hat{j}_x^{eq} \hat{G}^R(\epsilon_F) \right\rangle, \quad (\text{C3})$$

$$\begin{aligned} \sigma_{yx}^{z,II} &= \frac{\hbar}{2\pi} \text{Re} \int d\epsilon f^0(\epsilon) \\ &\times \text{tr} \left\langle \hat{j}_y^{z,eq} \hat{G}^R(\epsilon) \hat{j}_x^{eq} \frac{d\hat{G}^R(\epsilon)}{d\epsilon} - \hat{j}_y^{z,eq} \frac{d\hat{G}^R(\epsilon)}{d\epsilon} \hat{j}_x^{eq} \hat{G}^R(\epsilon) \right\rangle, \end{aligned} \quad (\text{C4})$$

and $\sigma_{yx}^{z,in-1} = \sum_l f_l^0 \langle l | \frac{1}{2} \left\{ \frac{\hbar}{2} \hat{\sigma}_z, \delta^{\mathbf{E}} \hat{v}_y \right\} | l \rangle / E_x$, i.e.,

$$\sigma_{yx}^{z,in-1} = \sum_l f_l^0 \langle l | \frac{1}{2} \left\{ \frac{\hbar}{2} \hat{\sigma}_z, \frac{-e}{i\hbar} [\hat{y}^{phy}, \hat{x}^{phy}] \right\} | l \rangle. \quad (\text{C5})$$

Here

$$\hat{j}^{eq} = \frac{e}{i\hbar} [\hat{\mathbf{r}}^{phy}, \hat{H}_0 + \hat{W}], \hat{j}^{z,eq} = \frac{1}{2} \left\{ \frac{\hbar}{2e} \hat{\sigma}_z, \hat{j}^{eq} \right\}, \quad (\text{C6})$$

and $\hat{G}^{R/A}(\epsilon) = (\epsilon - \hat{H}^{eq} \pm i0^+)^{-1}$ is the retarded/advanced Green's function operator. In Eq. (C5) only the disorder-free part of $\sigma_{yx}^{z,in-1}$ is retained, because we focus on the weak disorder-potential regime. In this regime $\sigma_{yx}^{z,I(b)}$ can be neglected and in $\sigma_{yx}^{z,II}$ only the disorder-free part is important [1, 2]. Applying Eqs. (C2) - (C5) to the 2D Nozieres-Lewiner effective model, we get $\sigma_{yx}^{z,II} = 0$, $\sigma_{yx}^{z,I(a)} = \sigma_{yx}^{z,sj} + \sigma_{yx}^{z,ad}$,

$\sigma_{yx}^{z,sj} = \sigma_{yx}^{z,ad} = -\sigma_{yx}^{z,in-1}$ and $\sigma_{yx}^z = \sigma_{yx}^{z,ad}$. For the Rashba 2D effective model, one can also obtain the same result as the SB approach.

On the other hand, in usual applications of the Kubo-Streda formula to spin Hall effects in the 2D Nozieres-Lewiner effective model [11-13, 30], $\hat{H}^{eq} = \hat{H}_0 + \hat{W}$ but $\hat{H}_1 = -e\mathbf{E} \cdot \hat{\mathbf{r}}$, and the physical position is not distinguished from the canonical position. Then those authors used $\sigma_{yx}^{z,0} = \sigma_{yx}^{z,I(a),0} + \sigma_{yx}^{z,I(b),0} + \sigma_{yx}^{z,II,0}$ where

$$\sigma_{yx}^{z,I(a),0} = \frac{\hbar}{2\pi} \text{tr} \left\langle \hat{j}_y^{z,eq,0} \hat{G}^R(\epsilon_F) \hat{j}_x^{eq,0} \hat{G}^A(\epsilon_F) \right\rangle,$$

$$\sigma_{yx}^{z,I(b),0} = -\frac{\hbar}{2\pi} \text{Re tr} \left\langle \hat{j}_y^{z,eq,0} \hat{G}^R(\epsilon_F) \hat{j}_x^{eq,0} \hat{G}^R(\epsilon_F) \right\rangle,$$

$$\begin{aligned} \sigma_{yx}^{z,II,0} &= \frac{\hbar}{2\pi} \text{Re} \int d\epsilon f^0(\epsilon) \text{tr} \left\langle \hat{j}_y^{z,eq,0} \hat{G}^R(\epsilon) \hat{j}_x^{eq,0} \frac{d\hat{G}^R(\epsilon)}{d\epsilon} \right. \\ &\quad \left. - \hat{j}_y^{z,eq,0} \frac{d\hat{G}^R(\epsilon)}{d\epsilon} \hat{j}_x^{eq,0} \hat{G}^R(\epsilon) \right\rangle, \end{aligned}$$

with

$$\hat{j}^{eq,0} \equiv \frac{e}{i\hbar} [\hat{\mathbf{r}}, \hat{H}_0 + \hat{W}], \hat{j}^{z,eq,0} \equiv \frac{1}{2} \left\{ \frac{\hbar}{2e} \hat{\sigma}_z, \hat{j}^{eq,0} \right\}. \quad (\text{C7})$$

Applying this usual Kubo-Streda formula to the 2D case of $\hat{H}_0 = \frac{\hbar^2 \hat{\mathbf{k}}^2}{2m}$, one has $\sigma_{yx}^{z,II,0} = 0$, $\sigma_{yx}^{z,I(a),0} = \sigma_{yx}^{z,0,sj} + \sigma_{yx}^{z,0,ad}$ and $\sigma_{yx}^{z,0,sj} = \sigma_{yx}^{z,0,ad} = \frac{1}{2} \sigma_{yx}^{z,0,ad}$, thus $\sigma_{yx}^{z,0} = \sigma_{yx}^{z,ad}$.

Appendix D: Technical details in Sec. VI

For 2D electrons, $\hat{\mathbf{r}}^a = \frac{\lambda_0^2}{4} \hat{\sigma}_z \hat{\mathbf{z}} \times \hat{\mathbf{k}}$ and $i\mathbf{J}_l^a = \frac{\lambda_0^2}{4} (\hat{\sigma}_z)_l \hat{\mathbf{z}} \times \mathbf{k}$. Thus $(\delta^{in,1} \mathbf{v}_l)_y = \frac{eE_x}{\hbar} 2 \frac{\lambda_0^2}{4} (\hat{\sigma}_z)_l$ and $\sum_l f_l^0 (\delta^{in,1} \mathbf{v}_l)_y = \frac{eE_x}{\hbar} 2 \frac{\lambda_0^2}{4} \sum_l f_l^0 (\hat{\sigma}_z)_l$, meanwhile

$$\sum_l g_l^{(-2)} \delta^{sj,1} \mathbf{v}_l \quad (\text{D1})$$

$$= 2 \sum_{l'} g_l^{(-2)} \omega_{l'l}^{(2)} \frac{\lambda_0^2}{4} \text{Re} (\hat{\sigma}_z)_{l'l'} \hat{\mathbf{z}} \times (\mathbf{k}' - \mathbf{k})$$

$$- 4 \frac{\lambda_0^2}{4\hbar} \sum_{l'} g_l^{(-2)} \frac{\langle |V_{l'l}|^2 \rangle}{d_{l'l'}} \text{Im} (\hat{\sigma}_z)_{l'l'} \hat{\mathbf{z}} \times (\mathbf{k}' - \mathbf{k}).$$

In the 2D Nozieres-Lewiner model $(\hat{\sigma}_z)_{l'l'} = \eta \delta_{\eta\eta'}$, thus

$$\sum_l g_l^{(-2)} (\delta^{sj,1} \mathbf{v}_l)_y \quad (\text{D2})$$

$$= 2 \sum_{\mathbf{k}', \mathbf{k}\eta} g_{\eta\mathbf{k}}^{(-2)} \omega_{\eta\mathbf{k}', \eta\mathbf{k}}^{(2)} i \left[\mathbf{J}_{\eta\mathbf{k}'}^a - \mathbf{J}_{\eta\mathbf{k}}^a \right]_y$$

$$= 2 \frac{e}{\hbar} E_x \sum_{\eta\mathbf{k}} \partial_{k_x} f^0 \left(i \mathbf{J}_{\eta\mathbf{k}}^a \right)_y = -\frac{e}{\hbar} E_x \sum_{\eta\mathbf{k}} f_{\eta\mathbf{k}}^0 \Omega_{\eta\mathbf{k}}^a$$

$$= -\sum_l f_l^0 (\delta^{in,1} \mathbf{v}_l)_y.$$

Here $(\delta^{in,1}\mathbf{v}_l)_y = \frac{eE_x}{\hbar}\Omega_{\eta\mathbf{k}}^a$ is only valid for the considered model (Appendix B). But if $\langle u_l | \hat{\sigma}_z | u_{l'} \rangle$ has l -off-diagonal matrix elements or is momentum dependent, we cannot get the relation $\sum_l g_l^{(2)} (\delta^{sj,1}\mathbf{v}_l)_y = -\sum_l f_l^0 (\delta^{\mathbf{E}\mathbf{v}})_l$.

Next we analyze the case of the conventional spin current in 2D hole systems. By $\hat{\mathbf{r}}^a = \frac{\lambda_0^2}{4}\hat{\sigma}_z\hat{\mathbf{z}} \times \hat{\mathbf{K}}$,

$$\sum_l f_l^0 (\delta^{in,1}\mathbf{j}_l^z)_y = E_x \frac{e\lambda_0^2}{4} \sum_l f_l^0 \frac{\frac{\partial K_x}{\partial k_x} + \frac{\partial K_y}{\partial k_y}}{2}, \quad (\text{D3})$$

and $\hat{V}(\hat{\mathbf{r}}^{phy}) = \hat{V}(\hat{\mathbf{r}}) + \frac{1}{2}(\nabla\hat{V} \cdot \hat{\mathbf{r}}^a + \hat{\mathbf{r}}^a \cdot \nabla\hat{V})$ leads to

$$\begin{aligned} (\delta^V \hat{j}_y^z)_{ll'} &= \frac{1}{2i} \frac{\lambda_0^2}{4} V_{ll'} (K_x - K'_x) \\ &+ \frac{i}{4} \frac{\lambda_0^2}{4} V_{ll'} \left((\mathbf{k} - \mathbf{k}') \times \hat{\mathbf{z}} \right) \cdot \left(\frac{\partial \mathbf{K}'}{\partial k'_y} + \frac{\partial \mathbf{K}}{\partial k_y} \right), \quad (\text{D4}) \end{aligned}$$

then

$$\begin{aligned} \sum_l g_l^{(-2)} (\delta^{sj,1}\mathbf{j}_l^z)_y &= \frac{\hbar}{2} \frac{\lambda_0^2}{4} \sum_{ll'} g_l^{(-2)} \omega_{ll'}^{(2)} \left[(K'_x - K_x) \right. \\ &+ (k'_x - k_x) \frac{1}{2} \left(\frac{\partial K'_x}{\partial k'_x} + \frac{\partial K_x}{\partial k_x} \right) \\ &\left. - (k'_y - k_y) \frac{1}{2} \left(\frac{\partial K'_x}{\partial k'_y} + \frac{\partial K_x}{\partial k_y} \right) \right], \quad (\text{D5}) \end{aligned}$$

where we have used $\frac{\partial K_x}{\partial k_x} = \frac{\partial K_y}{\partial k_y}$ for 2D electrons and 2D holes. For 2D electrons, $\hat{\mathbf{K}} = \hat{\mathbf{k}}$ thus the last term in the square brackets on the rhs of Eq. (D5) vanishes and we obtain the previous result. While for 2D holes, $\hat{\mathbf{K}} = (\hat{k}_x^3 - 3\hat{k}_y^2\hat{k}_x, 3\hat{k}_x^2\hat{k}_y - \hat{k}_y^3, 0)$ and $\sum_l g_l^{(-2)} (\delta^{sj,1}\mathbf{j}_l^z)_y$ is unlikely to be equal to $\sum_l f_l^0 (\delta^{in,1}\mathbf{j}_l^z)_y$ in general cases. Even if for slowly varying disorder potentials, the minus sign of the last term in the square brackets on the rhs of Eq. (D5) makes further simplification impossible, unless some special assumptions are made for the model Hamiltonian.

-
- [1] N. Nagaosa, J. Sinova, S. Onoda, A. H. MacDonald, and N. P. Ong, *Rev. Mod. Phys.* **82**, 1539 (2010).
- [2] J. Sinova, S. O. Valenzuela, J. Wunderlich, C. H. Back, and T. Jungwirth, *Rev. Mod. Phys.* **87**, 1213 (2015).
- [3] J. Sinova, D. Culcer, Q. Niu, N. A. Sinitsyn, T. Jungwirth, and A. H. MacDonald, *Phys. Rev. Lett.* **92**, 126603 (2004); E. G. Mishchenko, A. V. Shytov, and B. I. Halperin, *Phys. Rev. Lett.* **93**, 226602 (2004); J. I. Inoue, G. E. W. Bauer, and L. W. Molenkamp, *Phys. Rev. B* **70**, 041303(R) (2004).
- [4] M.C. Chang, Q. Niu, *J. Phys.: Condens. Matter* **20** 193202 (2008).
- [5] D. Xiao, M. C. Chang, and Q. Niu, *Rev. Mod. Phys.* **82**, 1959 (2010).
- [6] P. Nozieres and C. Lewiner, *J. Phys. (Paris)* **34**, 901 (1973).
- [7] H.-A. Engel, B. I. Halperin, and E. I. Rashba, *Phys. Rev. Lett.* **95**, 166605 (2005); X. Liu and J. Sinova, *Phys. Rev. B* **86**, 174301 (2012).
- [8] S. K. Lyo and T. Holstein, *Phys. Rev. Lett.* **29**, 423 (1972).
- [9] S. Zhang, *Phys. Rev. Lett.* **85**, 393 (2000).
- [10] P. Streda, *Phys. Rev. B* **82**, 045115 (2010).
- [11] A. Crepieux and P. Bruno, *Phys. Rev. B* **64**, 014416 (2001).
- [12] V. K. Dugaev, A. Crepieux, and P. Bruno, *Phys. Rev. B* **64**, 104411 (2001).
- [13] W. K. Tse and S. Das Sarma, *Phys. Rev. Lett.* **96**, 056601 (2006).
- [14] E. M. Hankiewicz and G. Vignale, *Phys. Rev. B* **73**, 115339 (2006).
- [15] G. Vignale, *J. Supercond. Novel Magn.* **23**, 3 (2010).
- [16] E. M. Hankiewicz and G. Vignale, *J. Phys.: Condens. Matter* **21**, 253202 (2009).
- [17] D. Culcer, E. M. Hankiewicz, G. Vignale, and R. Winkler, *Phys. Rev. B* **81**, 125332 (2010).
- [18] J. M. Ziman, *Electrons and Phonons* (Clarendon, Oxford, 1960).
- [19] N. A. Sinitsyn, *J. Phys.: Condens. Matter* **20**, 023201 (2008).
- [20] W. Kohn and J. M. Luttinger, *Phys. Rev.* **108**, 590 (1957). The Kohn-Luttinger theory is formulated in the weak disorder potential regime where we work in. In the dilute-impurity regime Luttinger and Kohn designed another approach based on a density expansion, which also lead to a SB theory, see J. M. Luttinger and W. Kohn, *Phys. Rev.* **109**, 1892 (1958). But that SB theory is not same as the present one. More discussions can be found in Ref. [24].
- [21] J. M. Luttinger, *Phys. Rev.* **112**, 739 (1958).
- [22] N. A. Sinitsyn, Q. Niu, and A. H. MacDonald, *Phys. Rev. B* **73**, 075318 (2006).
- [23] C. Xiao and Q. Niu, *Phys. Rev. B* **96**, 045428 (2017).
- [24] C. Xiao, B. Xiong, and F. Xue, arXiv: 1802.09716
- [25] N. A. Sinitsyn, A. H. MacDonald, T. Jungwirth, V. K. Dugaev, and J. Sinova, *Phys. Rev. B* **75**, 045315 (2007).
- [26] It should be stressed that we never bother the matrix ‘‘Boltzmann-like’’ kinetic equations and the ‘‘matrix distribution function’’ (e.g., A. Khaetskii, *Phys. Rev. Lett.* **96**, 056602 (2006); C. Gorini, P. Schwab, R. Raimondi, and A. L. Shelankov, *Phys. Rev. B* **82**, 195316 (2010)).
- [27] R. Raimondi, P. Schwab, C. Gorini, and G. Vignale, *Ann. Phys. (Berlin)* **524**, 153162 (2012).
- [28] X. Bi, P. He, E. M. Hankiewicz, R. Winkler, G. Vignale, and D. Culcer, *Phys. Rev. B* **88**, 035316 (2013).
- [29] K. Shen, R. Raimondi, and G. Vignale, *Phys. Rev. B* **90**, 245302 (2014).
- [30] W. K. Tse and S. Das Sarma, *Phys. Rev. B* **74**, 245309

- (2006).
- [31] C. Xiao and Q. Niu, Phys. Rev. B **96**, 035423 (2017).
 - [32] The Kubo-Streda linear response formula is valid from the Boltzmann regime to the diffusive regime.
 - [33] N. A. Sinitsyn, Q. Niu, J. Sinova, and K. Nomura, Phys. Rev. B **72**, 045346 (2005).
 - [34] S. A. Yang, H. Pan, Y. Yao, and Q. Niu, Phys. Rev. B **83**, 125122 (2011).
 - [35] L. Berger Phys. Rev. B **2**, 4559 (1970).
 - [36] H. Ishizuka and N. Nagaosa, Phys. Rev. B **96**, 165202 (2017). This paper re-quantizes the one-band semiclassical wavepacket motion to obtain an one-band effective quantum theory, thus can be viewed as a specific example of the framework presented in the present paper.
 - [37] A. Fert and P. M. Levy, Phys. Rev. Lett. **106**, 157208 (2011).
 - [38] J. Shi, P. Zhang, D. Xiao, and Q. Niu, Phys. Rev. Lett. **96**, 076604 (2006).
 - [39] E. M. Hankiewicz and G. Vignale, Phys. Rev. Lett. **100**, 026602 (2008).
 - [40] L. Wu, J. Yang, S. Wang, P. Wei, J. Yang, W. Zhang, and L. Chen, Appl. Phys. Lett. **105**, 202115 (2014).
 - [41] E. J. Moore, Phys. Rev. **160**, 607 (1967).