

# Lessons from HAWC PWNe observations: the diffusion constant is not a constant; Pulsars remain the likeliest sources of the anomalous positron fraction; Cosmic rays are trapped for long periods of time in pockets of inefficient diffusion

Stefano Profumo,<sup>1,2,\*</sup> Javier Reynoso-Cordova,<sup>2,3,†</sup> Nicholas Kaaz,<sup>1,‡</sup> and Maya Silverman<sup>1,§</sup>

<sup>1</sup>*Department of Physics, University of California,  
1156 High St. Santa Cruz, CA 95060, United States of America*

<sup>2</sup>*Santa Cruz Institute for Particle Physics, 1156 High St. Santa Cruz, CA 95060, United States of America*

<sup>3</sup>*Departamento de Física, DCI, Campus León, Universidad de Guanajuato, 37150, León, Guanajuato, México*

Recent TeV observations of nearby pulsars with the HAWC telescope have been interpreted as evidence that diffusion of high-energy electrons and positrons within pulsar wind nebulae is highly inefficient compared to the rest of the interstellar medium. If the diffusion coefficient well outside the nebula is close to the value inferred for the region inside the nebula, high-energy electrons and positrons produced by the two observed pulsars could not contribute significantly to the local measured cosmic-ray flux. The HAWC collaboration thus concluded that, under the assumption of isotropic and homogeneous diffusion, the two pulsars are ruled out as sources of the anomalous high-energy positron flux. Here, we argue that since the diffusion coefficient is likely *not* spatially homogeneous, the assumption leading to such conclusion is flawed. We solve the diffusion equation with a radially dependent diffusion coefficient, and show that the pulsars observed by HAWC produce potentially perfect matches to the observed high-energy positron fluxes. We also study the implications of inefficient diffusion within pulsar wind nebulae on Galactic scales, and show that cosmic rays are likely to have very long residence times in regions of inefficient diffusion. We describe how this prediction can be tested with studies of the diffuse Galactic emission.

PACS numbers:

## I. INTRODUCTION

The origin of the cosmic radiation and the mechanisms responsible for cosmic-ray acceleration remain largely, and in detail, unknown. Over the last few years, the discovery by the PAMELA (Payload for Antimatter Matter Exploration and Light-nuclei Astrophysics) detector that the positron fraction, the ratio of the positron to positron-plus-electron flux in the cosmic radiation, *increases* with energy for particle energies greater than about 10 GeV is especially puzzling [1]. This result was subsequently confirmed by the Fermi LAT (Large Area Telescope) [2] and by AMS (the Alpha Magnetic Spectrometer) [3], with increasingly large and convincing statistics.

An increasing high-energy positron fraction is puzzling because in almost any predictive Galactic cosmic-ray model (see e.g. [4]), secondary-to-primary ratios, such as the positron fraction, or B/C, must asymptotically decline with energy – simply as a result of the energy-dependence of the diffusion coefficient, which, in turn, is a macroscopic manifestation of the energy dependence of the Larmor radius. A growing positron fraction is thus likely a signal that the observed “anomalous” positrons, whose energy is observed up to almost 1 TeV, are not

of secondary origin but, rather, are produced as primary particles.

High-energy positrons and electrons cool very efficiently, both via synchrotron and via inverse Compton, both processes responsible for a rate of energy loss quadratic in the energy of the particle. The cooling time of a TeV positron in the Galaxy is, typically, on the order of

$$\tau_e \sim 3 \times 10^5 \text{ yr} \times (1 \text{ TeV}/E_e).$$

Cosmic-ray electrons and positrons diffuse in a random-walk pattern in the magnetic field of the Galaxy. If this process is described with a uniform diffusion coefficient  $D$ , sources producing TeV positrons observed at Earth must lie within a distance

$$d \lesssim \sqrt{D\tau_e} \sim 0.5 \text{ kpc}$$

for a standard diffusion coefficient [4]

$$D_{\text{ISM}} \sim 3 \times 10^{28} \text{ cm}^2/\text{s} (E_e/\text{GeV})^{0.33}.$$

High-energy positrons (and electrons) therefore can only be produced within a few kpc (at most) sphere of our Galactic position.

Of the potential sources of high-energy cosmic-ray positrons and electrons, one stands out as especially intriguing: the annihilation of Galactic dark matter (see e.g. [5–15]): In many microscopic realizations, dark matter particles can pair annihilate into matter-antimatter pairs, such as positrons and electrons. The annihilation rate, additionally, can be in some cases associated with the observed abundance of dark matter. This

\*Electronic address: profumo@ucsc.edu

†Electronic address: reynosoj@fisica.ugto.mx

‡Electronic address: nomahen@ucsc.edu

§Electronic address: massilve@ucsc.edu

possibility has received considerable interest ever since the PAMELA result, but is very tightly constrained by the fact that no associated signal has been observed in gamma rays from dark matter annihilation, for example, in local dwarf spheroidal galaxies [16] or in nearby clusters of galaxies [17]

For many years, an alternate, more mundane explanation has also been put forward [5, 18–23, 23]: that middle-aged ( $\tau \sim 10^5 - 10^6$  yr), nearby pulsars could accelerate, in their magnetosphere and, subsequently, in the surrounding shock with the interstellar medium, (primary) electrons and positrons to very high energy. The energetics of known candidate pulsars, as indicated by the pulsars’ spin-down luminosity, is in the correct range to explain the observed excess positrons, as long as (1) the diffusion coefficient is  $D \sim D_{\text{ISM}}$  between the pulsar and Earth, and (2) a fraction of  $\mathcal{O}(10\%)$  of the spin-down luminosity is injected in positron-electron pairs [20, 23].

Recently, the HAWC (High-Altitude Water Cherenkov) Observatory confirmed earlier results from Milagro [24] and HAWC [25] and observed extended TeV emission surrounding two nearby pulsars, Geminga and Monogem (PSR B0656+14), among the candidate sources for the observed anomalous high-energy positrons [26]. As previously noted in [27], the spectrum and morphology from TeV gamma-ray observations can be used to infer features of the underlying high-energy electron-positron population responsible for the up-scattering of photons to TeV energies. In fact, Ref. [27] argued that HAWC observations available at the time [25] were entirely compatible, and actually supported, the hypothesis that the observed excess high-energy positrons originated from nearby pulsars.

The recent HAWC observations [26] provide detailed information on the spatially extended emission of high-energy, TeV gamma rays from the pulsar wind nebula (PWN) regions surrounding Geminga and Monogem. Since hadronic processes are very unlikely given the environmental gas density in PWNe [28], the emission traces the brightness of inverse Compton emission from high-energy electrons and positrons. As a result of the large Klein-Nishin suppression for the electron-positron scattering off of high-energy photons, the target photon population is predominantly given by the cosmic microwave background, with a spatially uniform density distribution [26]. As a result, the brightness profile of the TeV emission potentially provides direct information on the radial distribution of the high-energy electron-positron population in the observed PWNe.

Following [26], we have also fit the brightness profile of the TeV emission, and indeed confirmed that (1) the emission is well fit by a purely diffusive density distribution for the electron and positron population, (2) the best-fit outward wind value is zero, meaning there is no evidence for non-diffusive transport outwards, and (3) the preferred inferred value of the diffusion constant at the energies of interest is significantly smaller than what inferred from standard cosmic-ray propagation models.

We also concur with the HAWC analysis that the diffusion coefficient needed to fit the data is significantly lower than what inferred from hadronic cosmic rays in the Galaxy [4].

A Bayesian search for best-fit parameters applied to the GALPROP package [4] indicated that different cosmic ray species likely probe very different regions of the interstellar medium, implying that the standard strategy of relying on e.g. the Boron-to-Carbon (B/C) ratio to calibrate propagation parameters might be unreliable [29]. This analysis, however, does not indicate that a difference in the propagation parameters between leptonic and hadronic cosmic rays should exist. Rather, it indicates that homogeneity is likely an incorrect assumption when large-scale Galactic diffusion is considered. Additional support for the lack of homogeneity in the diffusive properties of the interstellar medium comes from magnetohydrodynamic simulations [30] and from models of cosmic ray escape from supernova remnants [31], which generically indicate an exponential suppression of the diffusion coefficient inside the expanding cloud around cosmic-ray accelerators.

The HAWC Collaboration [26] entertained the possibility that the diffusion coefficient inferred from TeV observations is constant (homogeneous and isotropic) all the way to Earth. If that is the case, the contribution to the local flux of cosmic-ray positrons and electrons from Geminga and Monogem, the two pulsars whose nebulae were observed in their TeV emission, is highly suppressed and essentially negligible [26]. Ref. [26] shows that such conclusion holds for a broad range of choices for the pulsar initial spin-down timescale, the energy dependence of the diffusion coefficient as well as the spectral index and flux normalization of the injected cosmic rays.

Hooper and Linden [32] argued that the assumption of isotropic and homogeneous diffusion inside and outside the pulsar nebulae is inconsistent with the detection of high-energy (up to 20 TeV) electrons with H.E.S.S. [33]. The key argument is that a 20 TeV electron cools in around  $t_{\text{cool}} \sim 10,000$  years, and in that time such particles can only travel on the order of  $\sqrt{D \cdot t_{\text{cool}}}$  which, for the diffusion coefficient inferred by the HAWC observations inside the nebulae would indicate the presence of a high-energy cosmic-ray source within 10-20 pc of the Sun’s position. Plausible candidates for such a powerful, nearby source, however, are not known to exist: if such a source existed, it should in fact be readily detectable from its high-energy photon emission [32].

Interestingly, previous theoretical studies have suggested that cosmic-ray gradients can induce suppressed diffusion inside supernova remnants [31]. Alfvén waves generated by cosmic rays induce a net force that suppresses diffusion near the sites of cosmic-ray acceleration and, more generally, where cosmic-ray fluxes are larger [34]. Subsequent studies have elaborated on the role of non-linear diffusion of cosmic rays generated by magnetohydrodynamic turbulence around acceleration sites. Such turbulence induces an effective diffusion coefficient

which depends on the cosmic-ray density and is far from spatially homogeneous [35]. Ref. [36] studies the effect of cosmic-ray induced inhomogeneity in the diffusion coefficient from the gradient in cosmic-ray density induced by the large particle densities in the disc (“near-source regions”) versus the much lower densities in the diffusive halo at large Galactic latitudes; They conclude that the residence time of cosmic rays in regions of inefficient diffusion in the thick Galactic disk is much larger than simple estimates based on global grammage such as the boron-to-carbon ratio. Ref. [37], following the earlier work of [31], studies in a self-consistent picture the back-reaction on transport properties of cosmic-ray out-flux from supernova remnants, showing numerically that streaming instabilities significantly suppress the diffusion coefficient in regions tens of pc from the acceleration site, and for timescales, for high-energy cosmic rays of several TeV of energy, up to millions of years. Finally, ref. [38] studies observational consequences of the “self-confinement” of cosmic rays near their acceleration sites on the diffuse gamma-ray background, using Galactic population models for supernova remnants. Their conclusion are close to what Hooper and Linden find: the fraction of volume of the diffusive halo occupied by pockets of inefficient diffusion is small (less than 1%) if the radius of such pockets is on the order of 30 pc, although it could be substantial if the pockets of inefficient diffusion are on the order of 150 pc [32].

In this study, we build on previous theoretical work, and utilize the HAWC observations to study the effect of non-homogeneous, non-isotropic diffusion near cosmic-ray acceleration sites, and to re-assess whether the HAWC result affects the plausibility of local pulsars as the culprit for the observed positron excess. Our key findings are that physically plausible and theoretically motivated models for the radial dependence of the diffusion coefficient inside and outside the pulsar wind nebulae of Geminga and Monogem likely lead to significant fluxes of high-energy electrons and positrons from those sources to Earth. Models of the Galactic pulsar population and observationally-backed assumptions on the size of pulsar wind nebulae with time indicate that Galactic cosmic rays spend a very significant fraction of their residence time in inefficient diffusion regions; we discuss and propose several observational tests that will allow to test this hypothesis.

The remainder of this work is structured as follows: in Section II we provide details on the algorithm we use to solve the 3D diffusion equation and we describe the parameters and structure of the diffusion models we employ; in Section III we present our results on the differential positron flux; in Section IV we discuss the implications of pockets of inefficient diffusion on large scales and ways to probe their existence, and, finally, we summarize and discuss our results in Section V.

## II. NUMERICAL SOLUTION OF COSMIC-RAY DIFFUSION WITH A NON-HOMOGENEOUS DIFFUSION COEFFICIENT

We describe cosmic-ray diffusion in the Galaxy, as customary, through the partial differential equation

$$\frac{\partial \psi}{\partial t} = \vec{\nabla} \cdot (D(\vec{x}, E) \vec{\nabla} \psi) + \frac{\partial}{\partial E} (P(E) \psi) + Q, \quad (1)$$

where the unknown function  $\psi$  indicates the differential number density of particles in energy, and is a function of time, position and energy,

$$\psi = \psi(t, x, E) = \frac{dN}{dE},$$

$D(\vec{x}, E)$  is the diffusion coefficient, which we take to be time-independent but spatially inhomogeneous,  $P(E)$  describes energy losses (which we assume homogeneous and isotropic), and, finally,  $Q$  is the injection source term. In this work we are interested, for simplicity, in point-like injection sources in time and space, i.e.

$$Q \propto \delta(\vec{x} - \vec{x}_o) \delta(t - t_o).$$

Following Ref. [39], in the spherically symmetric limit we can transform Eq. (1) into a simpler differential equation which only depends on two variables: the radial coordinate and a new variable  $u$  to be defined below. First, we make the substitution  $\psi \rightarrow rPf$ , and define  $T = \int_E^{E_o} 1/P(x) dx$  the time it takes for a particle of initial energy  $E_o$  to cool down to an energy  $E$ , we define  $\tau = T - t$ ,  $z = T$  and, finally, we introduce the variable  $u = \int_0^z D(x) dx$ . The resulting equation in the variables  $r$  and  $u$  is [39]

$$\frac{\partial f}{\partial u} = \frac{\partial^2 f}{\partial r^2}, \quad (2)$$

the standard one dimensional diffusion equation. This equation, for spatially constant diffusion coefficients, can be easily solved by using its well-known Green’s Function and the fact that the source term is a  $\delta$ -like function.  $\psi$  is given by:

$$\psi(t, r, E) = \frac{N_o(E_o) P(E_o)}{\pi^{3/2} P(E) r_{\text{diff}}^3} e^{-r^2/r_{\text{diff}}^2}, \quad (3)$$

where

$$r_{\text{diff}}^2 = \int_E^{E_o} D(x)/P(x) dx, \quad (4)$$

and where  $N_o(E_o)$  is the injection spectrum. Alternately, the equation can also be solved using Monte Carlo methods. These methods have the significant advantage of easily generalizing to the case of non spatially homogeneous diffusion coefficients. We now describe the numerical procedure we adopt, and provide details on the

numerical accuracies compared to the Green's function method in the Appendix.

The algorithm we employ consists of sampling a normal distribution into another normal distribution using a random number generator, a procedure known as the Box-Muller transformation. Consider the simplest, 1D case, where one has a diffusion equation of the type

$$\frac{\partial f}{\partial t} = D \frac{\partial^2 f}{\partial x^2}, \quad (5)$$

where  $D$  is the diffusion coefficient. For a given initial condition  $f_o(x)$  the exact solution is

$$f(t, x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dx' f_o(x') \left\{ \frac{1}{\sqrt{4\pi Dt}} e^{-\frac{(x-x')^2}{4Dt}} \right\}, \quad (6)$$

where the term in parenthesis is the Green's function of the system. The key to solve this equation using a Monte Carlo approach is by noting that the Green's function is itself a Gaussian distribution, which can be thought of as the probability distribution of finding a particle at position  $x$  at time  $t$ . The algorithm then simply consists of "evolving" the system originally at  $t_o, x_o$  via the recipe

$$x = x_o + \eta \sqrt{2D\Delta t}, \quad (7)$$

where  $\eta$  is a set of  $N$  random numbers generated from another normal distribution with mean  $\mu = 0$  and width  $\sigma = 1$ ; the next step is then to store the outcomes  $x$  in histograms and normalize them; the result is finally a mapped distribution now centered at  $t_o + \Delta t$ .

Of course, the method outlined above generalizes to the particular form of the diffusion equation of Eq. (2). In this case, instead of tracking the position of the particle in time, we track it in 'energy steps' in the new variable  $u$ , i.e. the step is now defined through  $\Delta t \rightarrow \Delta u(E)$  where

$$\Delta u(E_i) = \int_{E_i + \Delta E}^{E_i} \frac{D(x)}{P(x)} dx, \quad (8)$$

for a certain energy  $E < E_i < E_o$ . The simulation thus tracks the random motion of particles with initial energy  $E_o$  until they reach a certain final energy  $E$ . Since the equation originally depends on  $x, y$  and  $z$ , to solve it we need to track down the particles independently in the Cartesian coordinate system, that means  $x_i = x_o + \eta \sqrt{2D\Delta t}$ , where  $i$  stands for  $x, y$  and  $z$ .

So far we have not made any assumption about the functional form of  $D$  and  $P$ , but once they are defined, the algorithm uniquely depends on the choice for  $\Delta E$ . We assume that the energy dependence of the diffusion coefficient reads

$$D(E) = D_o \left( \frac{E}{\text{GeV}} \right)^\delta, \quad (9)$$

with typical values of  $D_o \sim 10^{28} \text{cm}^2 \text{s}^{-1}$  and  $\delta = 1/3$  as illustrated in [32, 39, 40].  $P(E) = dE/dt$  accounts

for energy losses, and through this work we consider inverse Compton scattering and synchrotron processes, giving  $P(E) \simeq P_o E^2$ , with  $P_o \sim 1.02 \times 10^{-16}$  (notice that we also implemented Klein-Nishina corrections to the inverse Compton energy losses; we found very small effects, which came at great computational cost; we therefore neglect such corrections for most of the results shown below). Once these quantities are defined, we can solve for the initial energy  $E_o$  a particle must have if we measure it at its final energy  $E$ ,

$$E_o(E) = \max \left( \frac{E}{1 - P_o E t}, 0 \right). \quad (10)$$

The initial injection spectrum of cosmic rays is usually parametrized by a power law  $Q \propto E^{-\alpha}$  possibly with a high-energy exponential cutoff.

As explained in the Introduction, the recent HAWC observations indicate inefficient diffusion in the two observed pulsar wind nebulae, with a preferred value for the diffusion coefficient inside the Geminga PWN at 100 TeV of  $3.2_{-1.0}^{+1.4} \times 10^{27} \text{cm}^2 \text{s}^{-1}$  and inside the Monogem (PSR B0656+14) PWN of  $15_{-9}^{+49} \times 10^{27} \text{cm}^2 \text{s}^{-1}$  [26], to be contrasted with a putative diffusion coefficient for the interstellar medium (ISM) at large from GALPROP<sup>1</sup> analyses [4], at the same energies, of about  $10^{30} \text{cm}^2 \text{s}^{-1}$ ; HAWC therefore indicates a suppression of the diffusion coefficient of a factor between 200-500 (Geminga) and (15-150) inside the PWN compared to the ISM large-scale value.

Such inefficient diffusion yields strongly suppressed diffusion lengths, which following the Green's function solution can be defined as in Eq. (4). We plot the diffusion length, as a function of energy, for three choices of the normalization of the diffusion coefficient  $D_o$  (defined as in Eq.(9) and thus here corresponding to the value at 1 GeV of energy) of  $D_o = 200, 10, 0.7 \times 10^{26} \text{cm}^2 \text{s}^{-1}$ . The vertical axes indicates the distance from the pulsar. The curves on the plot are readily understood from the interplay of the energy dependence of the diffusion coefficient at low energy (below 1 TeV), and from the effects of highly efficient energy losses at higher energies. As a result, the maximal diffusion length, independent of  $D_o$ , corresponds to roughly TeV energies.

Our figure illustrates that for values of the diffusion coefficient close to what indicated by the HAWC observations,  $D_o \sim 0.7 \times 10^{26} \text{cm}^2 \text{s}^{-1}$ , the diffusion length of the injected cosmic-ray electrons and positrons is barely larger than the physical size of the PWN (we shade here only to guide the eye the region at distances smaller than 20 pc as indicative of those distances likely still within the PWN). For intermediate diffusion coefficients the diffusion length is shorter than the distance separation to the Earth (but large enough that a non-trivial flux at Earth of cosmic rays injected from the pulsar is expected) and,

<sup>1</sup> <http://galprop.stanford.edu/>




FIG. 1: The diffusion length  $r_{\text{diff}}$  (see Eq. (4)) for electrons/positrons, for different diffusion constants. A higher diffusion constant allows further distances, whether the low diffusion barely allows some particles to escape the PWN region (shaded green). The shape of the curves depends on the assumed energy dependence of the diffusion coefficient (for energies below one TeV) and on the assumed energy-loss processes at higher energies.

finally, for values of  $D_o$  close to those inferred by global cosmic-ray data the Earth is well within the diffusion length for a large interval of energies.

The outcome of the HAWC collaboration hypothesis that the diffusion coefficient inside and outside the PWN is constant, is therefore entirely unsurprising. However, as explained above, such hypothesis is also unrealistic and theoretically unmotivated. Here, we thus model different diffusive mediums with a simple model where within a few parsecs from the PWN particles experience inefficient diffusion, at the levels observed indirectly by HAWC, followed by a transition from the low diffusion to a high diffusion regime, with a diffusion constant close to the global ISM values.

We assume for simplicity spherical symmetry (unlike for example what assumed in Ref. [36, 38] which also explored spatially dependent diffusion constants but with variations in the axial  $z$  direction) and we explore two possibilities for the functional form of the radial dependence: either the transition is sudden, and modeled by the Heaviside function  $\theta(r)$ , or the transition is smooth; in this latter case we parametrize the radial dependence through an arctan,  $\tan^{-1}(r)$  functional form. In sum-




FIG. 2: The radial dependence of the diffusion coefficient we assume in our study. The plot shows the dependence on position of the coefficient for  $r_T = 30$  pc and different values for  $\delta$ .

mary, we assume the following two functional forms:

$$D_\theta(r) = D_1\theta(r_T - r) + D_2\theta(r - r_T) \quad (11a)$$

$$D_T(r) = D_1 + \frac{(D_2 - D_1)}{\pi} \left( \tan^{-1} \left( \frac{r - r_T}{\delta} \right) + \frac{\pi}{2} \right), \quad (11b)$$

where  $r_T$  stands for the distance from the PWN at which the transition takes place,  $D_1$  and  $D_2$  are the diffusion constants in medium 1 and 2 respectively and  $\delta$  is the width of the transition in the case of it being “smooth”. We display our choices for the radial dependence of the diffusion coefficient in Fig. 2. Naturally, for  $\delta \rightarrow 0$  the two functional forms converge – a fact we use to cross-check our numerical results.

### III. RESULTS: COSMIC RAY POSITRONS AND ELECTRONS FROM THE GEMINGA PULSAR

We solve Eq. (1) using the MC method described above to compute the spectrum at Earth of cosmic rays injected by a single pulsar with a spatially dependent diffusion coefficient. For reference, we focus our study on the case of the Geminga pulsar, and employ a characteristic age of  $t = 3.42 \times 10^5$  yr, a distance to the pulsar  $r = 250$  pc, an initial injection power-law spectrum of  $\alpha = 2.34$ , and a total energy released equal to  $11 \times 10^{48}$  ergs, in agreement with what extrapolated by the HAWC analysis.

Our main results are shown in Fig. 3–5. Fig. 3 shows the positron flux at Earth computed for different choices for the radial dependence of the diffusion coefficient. Here, we use a diffusion coefficient inside the PWN which is a factor of 100 smaller than outside. We also show the positron flux as measured by AMS-02 [3]. The




FIG. 3: The positron spectrum computed using the Monte Carlo algorithm described in section II for multiple choices for the radial dependence of the diffusion constant. We also show the AMS-02 measurement of the positron flux. We use the parameters appropriate for the Geminga pulsar, i.e.  $t = 342000$  yr,  $r = 250$  pc  $\alpha = 2.34$ , and  $D_1 = 3.86 \times 10^{26} \text{cm}^2 \text{s}^{-1}$  and  $D_2 = 3.86 \times 10^{28} \text{cm}^2 \text{s}^{-1}$ .

black dashed line indicates the case of no spatial variation in the diffusion coefficient. The blue and red lines have a step-function transition at 30 and 50 pc from the pulsar, respectively, while the green and dark blue lines show the arctan functional form with a broader and narrower transition region of 5 and 1 pc, respectively; in both cases we assume  $r_T = 30$  pc.

The figure clearly shows how a very significant high-energy flux of positrons is to be expected from e.g. Geminga if indeed diffusion transitions from an inefficient region in the nebula to values typical of the ISM 30 to 50 pc away from the pulsar. The precise spectrum depends, as expected, on the details of the radial dependence of the diffusion coefficient, as well as on the precise location of the transition region. We find for example that a more extended inefficient diffusion region, besides the obvious overall suppression of the flux (in our case study of around a factor 2 going from 30 to 50 pc radius), also enhances the fluxes of higher-energy particles – a consequence of the energy-dependent diffusion length we show in Fig. 1. A “smooth” transition tends to populate lower-energy particle fluxes versus a sharper transition, as visible comparing the 5 and 1 pc widths of the arctan cases, and also the arctan cases versus the sharp transitions.

An additional notable feature of fig. 3 is that for the same total energy injected in electrons and positrons, generally it is not true that a region of inefficient diffusion around the pulsar suppresses the flux at Earth. Rather, we find that for several choices for the radial dependence of the diffusion coefficient, the positron flux from Geminga is actually *enhanced* compared to a uniform, large diffusion coefficient.

We explore and explain this feature by studying, in fig. 4, the integrated flux for different energy ranges as a function of distance. We can appreciate from these plots how the positron number density between 100 GeV and 1000 GeV is higher than the constant diffusion case, as we find in the spectral plot. Our interpretation of these results is that the positrons in a high diffusion constant region reach further distances than the case where a transition is present: Positrons that were injected in a low diffusion constant region are effectively “trapped” until later times and thus, when they reach the high diffusion region, they do not have enough time to effectively diffuse. The greater the distance at which the transition happens, the more the spectrum resembles that of the low diffusion constant.

Fig. 5 provides evidence that a pulsar interpretation of the anomalous positron fraction is perfectly consistent with inefficient diffusion around PWNe. The figure utilizes a transition region with a steep transition profile and of transition radius 35 pc, and an assumed fraction of the pulsar spin-down luminosity converted to electron-positron pairs consistent with the power inferred by the TeV observations of HAWC, i.e. around 40% [26]. We also show a standard secondary positron flux (light blue), and the sum of the positrons from secondary processes plus those from Geminga. The figure clearly shows that with a radially dependent diffusion coefficient positrons from Geminga can dominate the positron flux measure at Earth and give an acceptable fit to the data (notice that for simplicity and for the sake of clarity we do not include additional pulsars, which certainly contribute to the high-energy positron flux as well, see e.g. [27]).

#### IV. MACROSCOPIC EFFECTS OF INEFFICIENT DIFFUSION IN PWNE

In this section we quantify the degree to which the implied inefficient diffusion coefficient inside PWNe,  $D_{\text{PWN}} \sim 10^{-2} D_{\text{ISM}}$  affects cosmic ray diffusion in the Galaxy at macroscopic scales. While we postpone detailed simulations and observational tests for future work, here we employ observationally-motivated relations between the PWN radius,  $R_{\text{PWN}}$ , and the pulsar characteristic age,  $\tau_c$  from the Australia Telescope National Facility (ATNF) catalogue of Galactic pulsars<sup>2</sup>, to infer the total volume in the Galactic diffusive halo where inefficient diffusion is expected,  $V_{\text{PWN}}$ .

PWNe are generated within the supernova remnants accompanying the birth of pulsars, and here we intend to estimate their radial size evolution as a function of time. We consider three distinct phases of the evolutionary stages of the PWNe following the procedure of Ref. [41]. Initially, we assume constant energy output

<sup>2</sup> <http://www.atnf.csiro.au/people/pulsar/psrcat/>




FIG. 4: The integrated flux per square meter per second per steradian for different energy ranges and different diffusion scenarios. The dashed lines represent the cases in which the medium undergoes a transition.  $D_1 = 3.86 \times 10^{26} \text{cm}^2 \text{s}^{-1}$ ,  $D_2 = 3.86 \times 10^{28} \text{cm}^2 \text{s}^{-1}$  and  $t = 342000 \text{ yr}$ .




FIG. 5: The positron flux with the contributions of Geminga and secondary production, compared to AMS-02 data. The transition for this case is set to 35 pc with  $\sim 40\%$   $e^\pm$  energy injection efficiency.

$\dot{E} \sim \dot{E}_0$  while  $t < \tau_0$ , where  $t$  is the age of the pulsar and  $\tau_0$  is the initial spin-down time scale of the pulsar. During this stage, the pulsar is surrounded by the un-shocked ejecta of the surrounding SNR and the PWN expands at supersonic speeds.

As the surrounding SNR accumulates matter from the ISM, the SNR ejecta are decelerated, triggering an inward moving reverse shock in the SNR which will then collide with the expanding PWN, causing the expansion of the PWN to slow down. This stage occurs at some reverse-shock interaction time,  $t_{rs}$ .

After this dynamical period of collision between the PWN and SNR reverse-shock, the PWN again steadily expands, making the assumption  $t_{rs} > \tau_0$ ; at this stage it is no longer warranted to take  $\dot{E} \sim \dot{E}_0$ , and thus the ex-

pansion slows down further. It should be noted that this is a simplified model of PWNe radial evolution; a more detailed picture of the evolution can be found in [42]. Finally, we use here the following equations to estimate  $R_{\text{PWN}}$  as a function of time:

$$R_{\text{PWN}}(t) \propto \begin{cases} t^{6/5}, & \text{for } t \leq \tau_0 \\ t, & \text{for } \tau_0 < t \leq t_{rs} \\ t^{3/10}, & \text{for } t > t_{rs}, \end{cases}$$

where  $t$ ,  $\tau_0$ , and  $t_{rs}$  are related via the following equation [42]:

$$\tau_c \equiv \frac{P}{2\dot{P}} = (\tau_0 + t) \frac{n-1}{2}, \quad (12)$$

$P$  being the spin period,  $n$  is the braking index of the pulsar,  $\tau_0$ ,  $t_{rs}$ , and these values are listed in [41].

The volume of each putative PWN was calculated and then summed to derive the total volume of PWNe in the Milky Way,  $V_{\text{PWN}}$ . We note that it was necessary to account for overlapping PWNe, and so when the volumes were summed, each contiguous region of connected PWNe was checked. If a PWN was isolated, the volume could be calculated in the simple case just using its radius. If two PWNe were connected, their volumes were summed analytically and the overlap region was subtracted to avoid double counting. To account for regions with 3 or more overlapping PWNe, a mesh subdivided into small cubes was created for each contiguous region. We checked these cubes to mark which fall within the radii of one of the PWNe, and the total volume was then estimated by summing these. The net PWNe volume as a function of characteristic age cut off is plotted in Fig. 6.

At the time of our sampling from the ATNF catalogue, 69.4% of pulsars had the necessary published values to






FIG. 6: We plot the net PWNe volume versus the pulsar characteristic age cut off with constants given by the baseline and varied models for PWNe radius evolution as listed in [41]. On the left, the blue region gives the total volume of the PWNe of pulsars sampled from the ATNF catalogue, the red curve is the baseline model adjusted for incompleteness factors in pulsar sampling that account for distance dependent and galactic longitude dependent selection effects [43], and the green region accounts for selection effects due to a pulsar mean beaming fraction of  $\langle f \rangle \sim 0.10$  [44] in addition to the same selection effects as the red curve. On the right, the incomplete volume is plotted with a curve that does not correct for overlapping PWNe to depict the amount of overlap of PWNe at different characteristic age cut offs.

estimate the volumes of their PWNe, leaving us a total sample of 1830 pulsars. To account for the incompleteness, we referred to previous works where a distribution of pulsars as a function of the center of the Galaxy has been suggested [43]; to estimate the total number of pulsars in the Galaxy, we then integrated their radial distribution:

$$\rho(R) = A \left( \frac{R + R_1}{R_\odot + R_1} \right)^a \exp \left[ -b \left( \frac{R - R_\odot}{R_\odot + R_1} \right) \right], \quad (13)$$

where  $\rho(R)$  is a cylindrical surface density with constants  $A = 37.6 \pm 1.9 \text{ kpc}^{-2}$ ,  $a = 1.64 \pm 0.11$ ,  $b = 4.01 \pm 0.14$ ,  $R_1 = 0.55 \pm 0.10 \text{ kpc}$ , and  $R_\odot = 8.5 \text{ kpc}$ , where  $R_\odot$  is the Sun-Galactic Center (GC) distance. Ref. [43] derives the functional form of this distribution by sampling the radial distribution of pulsars in the ATNF catalogue, excluding binary, recycled ( $\dot{P} < 10^{-17}$ ), globular cluster and Large and Small Magellanic Cloud pulsars. At the time, this included 1254 pulsars in total. Correction factors to relate this observed distribution to the total distribution of pulsars in the Milky Way accounted for direction-dependent  $K(l)$  and distance-dependent  $K(r)$  selection effects:

$$\rho(r, R, l(r, R)) = K(l)K(r)\rho_o(r, R, l(r, R)), \quad (14)$$

$\rho_o$  being the observed surface density of pulsars,  $r$  and  $R$  the heliocentric and galactocentric radii (respectively), and  $l$  being the Galactic longitude. Direction-dependent selection effects are accounted for by estimating the effect of background radiation on survey sensitivity as a function of  $l$ . Distance-dependent selection effects account for

minimum detectable flux and pulse broadening, scattering and scintillation. Integrating (12) gives a total pulsar number of  $(24 \pm 3) \times 10^3$ . Accounting for a mean beaming fraction of  $\langle f \rangle \sim 0.10$  [44], the total pulsar number is  $(240 \pm 30) \times 10^3$ . Fig.6 shows the uncorrected baseline model with standard deviation as well as the model of the corrected volume (with and without accounting for beam fraction).

Simpler estimates of the volume fraction of inefficient diffusive regions have been made in the past. For example, Hooper and Linden [27] used one putative single value for the typical size of PWNe of 30 pc, and a typical supernova rate for the Galaxy (0.03 per year) to infer that, for a 20 kpc radius and 200 pc half-height for the diffusion region, the total fraction of inefficient diffusion volume is of about 0.007, or about  $3.5 \text{ kpc}^3$ . This estimate is entirely consistent with our estimates, which indicate similar volumes for the baseline model corrected with beaming effects, for a characteristic age up to roughly 1 Myr.

What is the impact of pockets of inefficient diffusion in the Galaxy, and which diagnostics can be used to detect them? Ref. [38] addressed this question by studying the diffuse gamma-ray emission resulting from cosmic rays trapped for longer-than-usual times inside the “near-source region” where they were accelerated; inelastic collisions of cosmic-ray protons with nuclei in the interstellar medium leads via neutral pion decay to a possibly significant gamma-ray emission from these “halos” with large self-contained cosmic-ray populations. Ref. [38] then proceeds to studying the integrated diffuse gamma-ray emis-



sion in various angular regions. The key issue with this diagnostic is that the emission is basically indistinguishable, spectrally or morphologically, from other sources of diffuse emission; for example, in Ref. [45, 45] it was shown how models of the diffuse emission with an enhanced cosmic-ray source population tracing regions of hypothetically large star formation provide a better global fit to the observed Galactic diffuse gamma-ray emission; such a contribution would be likely highly degenerate with the extended halos considered in Ref. [38]

Before we turn to the observational tools we propose to test whether or not cosmic rays undergo inefficient diffusion near their acceleration sites, we shall discuss first the typical confinement times in these regions (notice that Ref. [38] also discusses escape times, albeit in a completely different diffusion scheme, see their figure 2). We notice that since the mean free path for a particle in the diffusive regime is of the order

$$\langle L \rangle \sim \sqrt{D \cdot t},$$

the ratio of the residence time for cosmic rays in the diffusive regime inside inefficient diffusion regions to that inside the rest of the ISM, is of the order

$$\frac{t_{\text{PWN}}}{t_{\text{ISM}}} \sim \left( \frac{\langle V \rangle_{\text{PWN}}}{\langle V \rangle_{\text{ISM}}} \right)^{2/3} \frac{D_{\text{ISM}}}{D_{\text{PWN}}} \sim 10^{-2} \left( \frac{\langle V \rangle_{\text{PWN}}}{\langle V \rangle_{\text{ISM}}} \right)^{2/3} \quad (15)$$

Thus, for effective diffusive volumes up to 1,000 times larger than our estimated inefficient diffusion volume, cosmic rays in the diffusive regime would spend *more* time inside regions of inefficient diffusion than elsewhere. For a diffusive halo of total volume

$$\langle V \rangle_{\text{ISM}} \simeq 500 \text{ kpc}^3 \left( \frac{R_h}{20 \text{ kpc}} \right)^2 \left( \frac{z_h}{0.2 \text{ kpc}} \right) \quad (16)$$

this means that for  $\langle V \rangle_{\text{PWN}} \gtrsim 0.5 \text{ kpc}^3$  cosmic rays are likely to spend more time trapped within acceleration sites than in the ISM at large. From our fig. 6 this, in turn, corresponds to very likely PWN sizes, corresponding to pulsar ages near approximately  $10^6 \text{ yr}$  of age, thus close to the age of the observed TeV halos of Geminga and Monogem [26].

How can this prediction be tested? The key observables are related to indirect probes of high-energy cosmic rays, such as for example synchrotron and inverse Compton energy losses of high-energy electrons and positrons, or inelastic hadronic processes producing neutral pions. If high-energy cosmic rays indeed spend a significant amount of time in relatively small pockets of inefficient diffusion, they will proportionally lose appreciably more energy in those environments than in the standard case with homogeneous diffusion. As a result, an appropriate diagnostic would consist of a study of the clustering scales of emission at the relevant wavelengths (ranging from radio for synchrotron emission, to X-ray, to soft and hard gamma rays for inverse Compton). The relevant angular

scales are of the order of

$$\theta \sim \frac{R_{\text{PWN}}}{d_{\text{PWN}}}, \quad (17)$$

and thus range from around 5 degrees in the case of the PWN observed by HAWC, down to  $\mathcal{O}(0.5^\circ)$  for pulsars near the center of the Galaxy.

The prediction for this model thus consists of significantly greater emission power on physical scales of the order of PWN sizes, and on angular scales extending from 0.1 to several degrees, as well as a brighter emission in the Galactic regions where more abundant populations of PWNe are present. The emission is expected over a broad range of frequencies, from very high energy gamma rays, down to soft gamma rays, X-rays, and radio frequencies. The diagnostic will consist of, for instance, calculating the angular power spectrum predicted for significant emission from PWN (utilizing both the ATNF pulsar catalogue and corrections thereof) and following the procedure outlines in Ref. [46] to constrain dark matter emission. A similar procedure was used to describe how to constrain decaying dark matter emission at X-ray frequencies with eROSITA in Ref. [47]

In addition to the angular power spectrum, other possible diagnostics include (i) a wavelet analysis, as pioneered again for the case of the diffuse gamma-ray emission in Ref. [48] and there used to distinguish the clustering properties of millisecond pulsars versus dark matter, or (ii) using metrics like the flux probability density function which Ref. [49] utilized to distinguish uniform from non-uniform diffuse emission in the gamma-ray sky. At present, all of the outlined methods are under investigation in their application to the problem of disentangling a contribution from in source, “trapped” cosmic-ray emission from the diffuse emission.

## V. DISCUSSION

In this work we have explored the possibility of a non-homogeneous medium through which cosmic rays diffuse in the vicinity of their acceleration sites as a solution to the low diffusion coefficient inferred by the HAWC Collaboration [26] compared to what inferred from global fits to Galactic cosmic-ray data [4]. We have implemented a numerical solution to the problem of spherically-symmetric diffusion with a radially-dependent diffusion coefficient, which consists of a Monte Carlo integration. We validated our numerical procedure against known analytical solutions in the appropriate limits, and we then proceeded with the exploration of a variety of functional forms for the transition from low to high efficiency in the cosmic-ray diffusion away from their injection sources.

We found that (i) the sharper the transition from inefficient to efficient diffusion, the more abundant the asymptotic relic population of high-energy cosmic rays; (ii) for

the specific case of Geminga, the flux of cosmic rays predicted by scenarios of inefficient-to-efficient diffusion versus uniformly efficient diffusion, with the same injected flux, can actually be larger, especially at higher energies; (iii) the location of the transition region from inefficient to efficient diffusion significantly affects the asymptotic cosmic-ray spectrum, with a larger suppression of low-energy cosmic rays for larger transition radii; (iv) there exist several realization of radially-dependent diffusion from Geminga or Monogem that suitably explain the excess high-energy positron flux at Earth as originating from those nearby, middle-aged pulsars.

We then proceeded to estimate and discuss the *global* implications for Galactic cosmic-ray transport of the existence of pockets of inefficient diffusion around cosmic-ray accelerator. We showed a data-driven approach, based on the ATNF pulsar catalogue and on completeness corrections thereof, and on observationally and theoretically motivated models for the physical size of pulsar wind nebulae, to estimate the total Galactic volume with inefficient diffusion. We showed that such volume is likely at least 1% of the Galactic diffusive halos, and possibly larger. If this is the case, we argued that cosmic rays spend most of their residence time inside inefficient diffusion regions.

We proposed several observational diagnostics to validate our findings, which essentially plan to utilize a study of diffuse emission power on the angular scales relevant for the expected size of regions of inefficient diffusion, ranging from a fraction of a degree to several degrees in the sky.

In conclusion, the key lesson from the HAWC results [26] is evidence that diffusion is not homogeneous in the Galaxy, and that cosmic-ray diffusion is inefficient inside pulsar wind nebulae; this is not an unexpected result, and it had been anticipated by several theoretical studies on self-generated turbulence inside cosmic-ray acceleration sites [31]. Unlike what claimed in Ref. [26], we showed that for several choices for the transition from inefficient to efficient diffusion, nearby pulsars such as Geminga are likely very significant contributors to the local high-energy positron flux. An important implication of the HAWC results is that likely cosmic rays spend most of their Galactic residence time inside hypothetical inefficient diffusion regions. We outlined observational methods to test this possibility which include studying the angular power spectrum of diffuse emission at a variety of wavelengths, wavelet analyses, and non-Poissonian photon statistics.

*Note Added.* During the final stages of this work, Ref. [50] appeared, which deals with a toy model of a two-zone diffusion coefficient and a semi-analytic solution consisting of simple matching of boundary conditions; while the results of Ref. [50] agree with ours when directly comparable, their diffusion coefficient model is unphysical and the accuracy of the solution to the cosmic-ray density questionable.

## Appendix A: Results of the Monte Carlo Simulation

As described in Section II, we used a Monte Carlo (MC) algorithm to solve the 3D diffusion equation by using a Box-Muller transformation. In order to test our method we solved the equation and compared it with the analytical very well known solution Eq. 3. Since the MC method is purely stochastic, we computed the  $1\sigma$  error for the points generated. The results are presented in Fig. 7 for the values of the diffusion regime used through this work. In the figure, the dashed lines indicate the analytical solution for a delta-function injection source, while the blue and red data points show the results of our MC method. The numerical accuracy of our algorithm is clearly very high.

## Acknowledgements

We would like to thank Nicolas Fernandez for helpful discussions. This work was funded by a UCMEXUS-CONACYT collaborative project. SP is partly supported by the U.S. Department of Energy grant number de-sc0010107. JR acknowledges financial support from CONACYT and CONACYT project 182445.




FIG. 7: The accuracy of the Monte Carlo Method. The dashed lines represent the analytical solution of the diffusion equation in the case of a  $\delta$ -like initial injection, the blue and red points were generated through the MC simulation as described in section II. The error bars displayed the probability of finding the point  $1\sigma$  away from the true value. The results indicate that our algorithm is in good agreement with the analytical solutions.

- 
- [1] O. Adriani et al. (PAMELA), *Phys. Rev. Lett.* **105**, 121101 (2010), 1007.0821.
- [2] M. Ackermann et al. (Fermi-LAT), *Phys. Rev.* **D82**, 092004 (2010), 1008.3999.
- [3] M. Aguilar, G. Alberti, B. Alpat, A. Alvino, G. Ambrosi, K. Andeen, H. Anderhub, L. Arruda, P. Azzarello, A. Bachlechner, et al. (AMS Collaboration), *Phys. Rev. Lett.* **110**, 141102 (2013), URL <https://link.aps.org/doi/10.1103/PhysRevLett.110.141102>.
- [4] A. W. Strong and I. V. Moskalenko, *Astrophys. J.* **509**, 212 (1998), astro-ph/9807150.
- [5] I. Cholis and D. Hooper, *Phys. Rev.* **D88**, 023013 (2013), 1304.1840.
- [6] L. Bergstrom, T. Bringmann, and J. Edsjo, *Phys. Rev.* **D78**, 103520 (2008), 0808.3725.
- [7] M. Cirelli and A. Strumia, *PoS IDM2008*, 089 (2008), 0808.3867.
- [8] A. E. Nelson and C. Spitzer, *JHEP* **10**, 066 (2010), 0810.5167.
- [9] N. Arkani-Hamed, D. P. Finkbeiner, T. R. Slatyer, and N. Weiner, *Phys. Rev.* **D79**, 015014 (2009), 0810.0713.
- [10] I. Cholis, D. P. Finkbeiner, L. Goodenough, and N. Weiner, *JCAP* **0912**, 007 (2009), 0810.5344.
- [11] R. Harnik and G. D. Kribs, *Phys. Rev.* **D79**, 095007 (2009), 0810.5557.
- [12] P. J. Fox and E. Poppitz, *Phys. Rev.* **D79**, 083528 (2009), 0811.0399.
- [13] M. Pospelov and A. Ritz, *Phys. Lett.* **B671**, 391 (2009), 0810.1502.
- [14] J. D. March-Russell and S. M. West, *Phys. Lett.* **B676**, 133 (2009), 0812.0559.
- [15] S. Chang and L. Goodenough, *Phys. Rev.* **D84**, 023524 (2011), 1105.3976.
- [16] M. Ackermann et al. (Fermi-LAT), *Phys. Rev. Lett.* **107**, 241302 (2011), 1108.3546.
- [17] M. Ackermann et al. (Fermi-LAT), *Astrophys. J.* **717**, L71 (2010), 1006.0748.
- [18] D. Hooper, P. Blasi, and P. D. Serpico, *JCAP* **0901**, 025 (2009), 0810.1527.
- [19] H. Yuksel, M. D. Kistler, and T. Stanev, *Phys. Rev. Lett.* **103**, 051101 (2009), 0810.2784.
- [20] S. Profumo, *Central Eur. J. Phys.* **10**, 1 (2011), 0812.4457.
- [21] D. Malyshev, I. Cholis, and J. Gelfand, *Phys. Rev.* **D80**, 063005 (2009), 0903.1310.
- [22] D. Grasso et al. (Fermi-LAT), *Astropart. Phys.* **32**, 140 (2009), 0905.0636.
- [23] T. Linden and S. Profumo, *Astrophys. J.* **772**, 18 (2013), 1304.1791.
- [24] A. A. Abdo et al., *Astrophys. J.* **700**, L127 (2009), [Erratum: *Astrophys. J.* 703, L185 (2009)], 0904.1018.
- [25] A. U. Abeysekara et al., *Astrophys. J.* **843**, 40 (2017), 1702.02992.
- [26] A. U. Abeysekara et al. (HAWC), *Science* **358**, 911 (2017), 1711.06223.
- [27] D. Hooper, I. Cholis, T. Linden, and K. Fang, *Phys. Rev.* **D96**, 103013 (2017), 1702.08436.
- [28] F. Paresce, *Astronomical Journal* **89**, 1022 (1984).
- [29] G. Jóhannesson et al., *Astrophys. J.* **824**, 16 (2016), 1602.02243.
- [30] H. Yan and A. Lazarian, *Astrophys. J.* **614**, 757 (2004), astro-ph/0408172.
- [31] M. A. Malkov, P. H. Diamond, R. Z. Sagdeev, F. A. Aharonian, and I. V. Moskalenko, *Astrophys. J.* **768**, 73 (2013), 1207.4728.
- [32] D. Hooper and T. Linden, Submitted to: *JCAP* (2017), 1711.07482.
- [33] H. Abdalla et al. (H.E.S.S.) (2017), 1709.06442.
- [34] A. Achterberg, *Astronomy and Astrophysics* **98**, 195 (1981).
- [35] V. S. Ptuskin, V. N. Zirakashvili, and A. A. Plesser, *Adv. Space Res.* **42**, 486 (2008).
- [36] M. D'Angelo, P. Blasi, and E. Amato, *Phys. Rev.* **D94**, 083003 (2016), 1512.05000.
- [37] L. Nava, S. Gabici, A. Marcowith, G. Morlino, and V. S. Ptuskin, *Mon. Not. Roy. Astron. Soc.* **461**, 3552 (2016), 1606.06902.
- [38] M. D'Angelo, G. Morlino, E. Amato, and P. Blasi, *Mon. Not. Roy. Astron. Soc.* **474**, 1944 (2018), 1710.10937.
- [39] A. M. Atoyan, F. A. Aharonian, and H. J. Völk, *Phys. Rev. D* **52**, 3265 (1995), URL <https://link.aps.org/doi/10.1103/PhysRevD.52.3265>.
- [40] J. Feng and H.-H. Zhang, *Eur. Phys. J.* **C76**, 229 (2016), 1504.03312.
- [41] H. Abdalla et al. (H.E.S.S.) (2017), 1702.08280.
- [42] B. M. Gaensler and P. O. Slane, *Ann. Rev. Astron. Astrophys.* **44**, 17 (2006), astro-ph/0601081.
- [43] I. Yusifov and I. Kucuk, *Astron. Astrophys.* **422**, 545 (2004), astro-ph/0405559.
- [44] T. M. Tauris and R. N. Manchester, *Monthly Notices of the Royal Astronomical Society* **298**, 625 (1998).
- [45] E. Carlson, T. Linden, and S. Profumo, *Phys. Rev.* **D94**, 063504 (2016), 1603.06584.
- [46] M. Fornasa et al., *Phys. Rev.* **D94**, 123005 (2016), 1608.07289.
- [47] F. Zandanel, C. Weniger, and S. Ando, *JCAP* **1509**, 060 (2015), 1505.07829.
- [48] R. Bartels, S. Krishnamurthy, and C. Weniger, *Phys. Rev. Lett.* **116**, 051102 (2016), 1506.05104.
- [49] S. K. Lee, M. Lisanti, and B. R. Safdi, *JCAP* **1505**, 056 (2015), 1412.6099.
- [50] K. Fang, X.-J. Bi, P.-F. Yin, and Q. Yuan (2018), 1803.02640.