

**A SHERMAN MORRISON WOODBURY IDENTITY FOR RANK AUGMENTING MATRICES WITH APPLICATION TO CENTERING\***

KURT S. RIEDEL†

**Abstract.** Matrices of the form  $\mathbf{A} + (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}(\mathbf{V}_2 + \mathbf{W}_2)^*$  are considered where  $\mathbf{A}$  is a singular  $\ell \times \ell$  matrix and  $\mathbf{G}$  is a nonsingular  $k \times k$  matrix,  $k \leq \ell$ . Let the columns of  $\mathbf{V}_1$  be in the column space of  $\mathbf{A}$  and the columns of  $\mathbf{W}_1$  be orthogonal to  $\mathbf{A}$ . Similarly, let the columns of  $\mathbf{V}_2$  be in the column space of  $\mathbf{A}^*$  and the columns of  $\mathbf{W}_2$  be orthogonal to  $\mathbf{A}^*$ . An explicit expression for the inverse is given, provided that  $\mathbf{W}_i^* \mathbf{W}_i$  has rank  $k$ . An application to centering covariance matrices about the mean is given.

**Key words.** Linear Algebra, Schur Matrices, Generalized Inverses

**AMS(MOS) subject classifications.** 65R10, 33A65, 35K05, 62G20, 65P05

The wellknown Sherman-Morrison-Woodbury matrix identity [1]:

$$(\mathbf{A} + \mathbf{X}_1 \mathbf{G} \mathbf{X}_2^T)^{-1} = \mathbf{A}^{-1} - \mathbf{A}^{-1} \mathbf{X}_1 (\mathbf{G}^{-1} + \mathbf{X}_2^T \mathbf{A}^{-1} \mathbf{X}_1)^{-1} \mathbf{X}_2^T \mathbf{A}^{-1} \quad (1)$$

is widely used. Several excellent review articles have appeared recently [2-4]. However (1) is only valid when  $\mathbf{A}$  is nonsingular<sup>1</sup>. In this article, we consider matrix inverses of the form  $\mathbf{A} + \mathbf{X}_1 \mathbf{G} \mathbf{X}_2^T$  where the rank of  $\mathbf{A} + \mathbf{X}_1 \mathbf{G} \mathbf{X}_2^T$  is larger than the rank of  $\mathbf{A}$ .

We decompose the matrix  $\mathbf{X}_1$  into  $\mathbf{V}_1 + \mathbf{W}_1$ , where the columns of  $\mathbf{V}_1$  are contained in the column space of  $\mathbf{A}$  and the columns of  $\mathbf{W}_1$  are orthogonal to it. Similarly, we decompose  $\mathbf{X}_2$  into  $\mathbf{V}_2 + \mathbf{W}_2$ , where the columns of  $\mathbf{V}_2$  are contained in the column space of  $\mathbf{A}^*$  and the columns of  $\mathbf{W}_2$  are orthogonal to  $M(\mathbf{A}^*)$ . We denote the column space of  $\mathbf{A}$  by  $M(\mathbf{A})$ . The Moore-Penrose generalized inverse will be denoted by the superscript  $+$ . We denote the  $k \times k$  matrix  $\mathbf{W}_i^* \mathbf{W}_i$  by  $\mathbf{B}_i$  and define  $\mathbf{C}_i \equiv \mathbf{W}_i (\mathbf{W}_i^* \mathbf{W}_i)^{-1}$ . We will require  $\mathbf{B}_i$  to be nonsingular. However the rank of the perturbation,  $k$ , can be significantly less than the size of the original matrix. We note that  $\mathbf{V}_i^* \mathbf{W}_i = 0$  and  $\mathbf{W}_i^* \mathbf{C}_i = \mathbf{I}_k$ . Finally the projection operator onto the column space of  $\mathbf{W}$  satisfies  $\mathbf{W}_i \mathbf{B}_i^{-1} \mathbf{W}_i^* = \mathbf{W}_1 \mathbf{C}_1^* = \mathbf{C}_2 \mathbf{W}_2^*$ .

**Theorem 1.** Let  $\mathbf{A}$  be a  $\ell \times \ell$  matrix of rank  $\ell_1$ ,  $\ell_1 < \ell$ ,  $\mathbf{V}_i$  and  $\mathbf{W}_i$  be  $\ell \times k$  matrices and  $\mathbf{G}$  be a  $k \times k$  nonsingular matrix. Let the columns of  $\mathbf{V}_1 \in M(\mathbf{A})$  and the columns of  $\mathbf{W}_1$  be orthogonal to  $M(\mathbf{A})$ . Similarly, let the columns of  $\mathbf{V}_2 \in M(\mathbf{A}^*)$  and the columns of  $\mathbf{W}_2$  be orthogonal to  $M(\mathbf{A}^*)$ . Let  $\mathbf{B}_i \equiv \mathbf{W}_i^* \mathbf{W}_i$  have rank  $k$ . The matrix,

$$\Omega \equiv \mathbf{A} + (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}(\mathbf{V}_2 + \mathbf{W}_2)^*,$$

has the following Moore-Penrose generalized inverse:

$$\Omega^+ = \mathbf{A}^+ - \mathbf{C}_2 \mathbf{V}_2^* \mathbf{A}^+ - \mathbf{A}^+ \mathbf{V}_1 \mathbf{C}_1^* + \mathbf{C}_2 (\mathbf{G}^+ + \mathbf{V}_2^* \mathbf{A}^+ \mathbf{V}_1) \mathbf{C}_1^*. \quad (2)$$

\* Received by the editors July 12, 1990; accepted for publication February 19, 1991.

† Courant Institute of Mathematical Sciences, New York University, New York, New York 10012. The work of this author was supported by the U.S. Department of Energy Grant No. DE-FG02-86ER53223.

<sup>1</sup> We denote the transpose of a matrix,  $\mathbf{A}$  by  $\mathbf{A}^T$  and the hermitian or conjugate transpose by  $\mathbf{A}^*$ .

Proof: We recall that the Moore Penrose inverse is the unique generalized inverse which satisfies the following four conditions, (Ref. [5], p.26):

$$(a) \Omega\Omega^+\Omega = \Omega, \quad (b) \Omega^+\Omega\Omega^+ = \Omega^+, \\ (c) (\Omega\Omega^+)^* = \Omega\Omega^+, \quad (d) (\Omega^+\Omega)^* = \Omega^+\Omega.$$

The identity is verified by direct computation,

$$\begin{aligned} \Omega\Omega^+ &\equiv \mathbf{A}\mathbf{A}^+ - \mathbf{A}\mathbf{C}_2\mathbf{V}_2^*\mathbf{A}^+ - \mathbf{A}\mathbf{A}^+\mathbf{V}_1\mathbf{C}_1^* + \mathbf{A}\mathbf{C}_2(\mathbf{G}^+ + \mathbf{V}_2^*\mathbf{A}^+ + \mathbf{V}_1)\mathbf{C}_1^* \\ &+ (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}(\mathbf{V}_2 + \mathbf{W}_2)^*\mathbf{A}^+ - (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}(\mathbf{V}_2 + \mathbf{W}_2)^*\mathbf{C}_2\mathbf{V}_2^*\mathbf{A}^+ \\ &- (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}(\mathbf{V}_2 + \mathbf{W}_2)^*\mathbf{A}^+\mathbf{V}_1\mathbf{C}_1^* \\ &+ (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}(\mathbf{V}_2 + \mathbf{W}_2)^*\mathbf{C}_2(\mathbf{V}_2^*\mathbf{A}^+ + \mathbf{V}_1)\mathbf{C}_1^* \\ &+ (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}(\mathbf{V}_2 + \mathbf{W}_2)^*\mathbf{C}_2\mathbf{G}^+ + \mathbf{C}_1^*. \end{aligned}$$

Since  $\mathbf{W}_2$  is orthogonal to  $\mathbf{A}^*$ , we have  $\mathbf{A}\mathbf{W}_2 = 0$ ,  $\mathbf{W}_2^*\mathbf{A}^+ = 0$ , and  $\mathbf{V}_2^*\mathbf{W}_2 = 0$ , which simplifies the previous expression to

$$\begin{aligned} \Omega\Omega^+ &\equiv \mathbf{A}\mathbf{A}^+ - \mathbf{A}\mathbf{A}^+\mathbf{V}_1\mathbf{C}_1^* + (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}\mathbf{V}_2^*\mathbf{A}^+ \\ &- (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}\mathbf{W}_2^*\mathbf{C}_2\mathbf{V}_2^*\mathbf{A}^+ - (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}\mathbf{V}_2^*\mathbf{A}^+\mathbf{V}_1\mathbf{C}_1^* \\ &+ (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}\mathbf{W}_2^*\mathbf{C}_2\mathbf{V}_2^*\mathbf{A}^+\mathbf{V}_1\mathbf{C}_1^* + (\mathbf{V}_1 + \mathbf{W}_1)\mathbf{G}\mathbf{W}_2^*\mathbf{C}_2\mathbf{G}^+ + \mathbf{C}_1^*. \end{aligned}$$

This expression may be simplified using  $\mathbf{G}\mathbf{W}_2^*\mathbf{C}_2\mathbf{G}^+ + \mathbf{C}_1^* = \mathbf{C}_1^*$ , and  $\mathbf{G}\mathbf{W}_2^*\mathbf{C}_2\mathbf{V}_2^* = \mathbf{G}\mathbf{V}_2^*$ , and  $\mathbf{A}\mathbf{A}^+\mathbf{V}_1 = \mathbf{V}_1$  to

$$\Omega\Omega^+ \equiv \mathbf{A}\mathbf{A}^+ + \mathbf{W}_1\mathbf{C}_1^*,$$

and clearly condition (c) is satisfied.

The corresponding identity for  $\Omega^+\Omega \equiv \mathbf{A}^+\mathbf{A} + \mathbf{C}_2\mathbf{W}_2^*$  requires the decomposition to satisfy  $\mathbf{A}^+\mathbf{W}_1 = 0$ ,  $\mathbf{W}_1^*\mathbf{A} = 0$ ,  $\mathbf{V}_1^*\mathbf{W}_1 = 0$ , and  $\mathbf{V}_2\mathbf{A}^+\mathbf{A} = \mathbf{V}_2$ . In addition, the matrix  $\mathbf{G}$  must satisfy  $\mathbf{C}_2\mathbf{G}^+ + \mathbf{C}_1^*\mathbf{W}_1\mathbf{G} = \mathbf{C}_2$  and  $\mathbf{V}_1\mathbf{C}_1^*\mathbf{W}_1\mathbf{G} = \mathbf{V}_1\mathbf{G}$ . These requirements guarantee that conditions (a), (b) and (d) are also satisfied.  $\square$

Remark: The conditions that  $\mathbf{G}$  and  $\mathbf{W}_i^*\mathbf{W}_i$  have rank  $k$  may be replaced by the somewhat weaker but more complicated conditions that  $\mathbf{G}\mathbf{W}_2^*\mathbf{C}_2\mathbf{G}^+ + \mathbf{C}_1^* = \mathbf{C}_1^*$ ,  $\mathbf{G}\mathbf{W}_2^*\mathbf{C}_2\mathbf{V}_2^* = \mathbf{G}\mathbf{V}_2^*$ ,  $\mathbf{C}_2\mathbf{G}^+ + \mathbf{C}_1^*\mathbf{W}_1\mathbf{G} = \mathbf{C}_2$  and  $\mathbf{V}_1\mathbf{C}_1^*\mathbf{W}_1\mathbf{G} = \mathbf{V}_1\mathbf{G}$ .

Note that the generalized inverse in (2) is singular and tends to infinity as  $\mathbf{W}_i$  approaches to zero. Thus (2) does not reduce to the (1) as the perturbation tends to zero. When the perturbation of the column space of  $\mathbf{A}$  is zero, i.e.  $\mathbf{V} \equiv 0$ , theorem 1 simplifies to

$$\Omega^+ = \mathbf{A}^+ + \mathbf{C}_2\mathbf{G}^+ + \mathbf{C}_1. \quad (3)$$

When  $\mathbf{A}$  is a symmetric matrix, the column spaces of  $\mathbf{A}$  and  $\mathbf{A}^*$  are identical. Thus, for the case of symmetric  $\mathbf{A}$  and  $\Omega$ , Thm. 1 reduces to

**Theorem 2.** Let  $\mathbf{A}$  be a symmetric  $\ell \times \ell$  matrix of rank  $\ell_1$ ,  $\ell_1 < \ell$ ,  $\mathbf{V}$  and  $\mathbf{W}$  be  $\ell \times k$  matrices and  $\mathbf{G}$  be a  $k \times k$  nonsingular matrix. Let  $\mathbf{V} \in M(\mathbf{A})$  and the

columns of  $\mathbf{W}$  be orthogonal to  $M(\mathbf{A})$ . Let  $\mathbf{B} \equiv \mathbf{W}^* \mathbf{W}$  have rank  $k$ . The matrix,

$$\Omega \equiv \mathbf{A} + (\mathbf{V} + \mathbf{W}) \mathbf{G} (\mathbf{V} + \mathbf{W})^* ,$$

has the following Moore-Penrose generalized inverse:

$$\Omega^+ = \mathbf{A}^+ - \mathbf{C} \mathbf{V}^* \mathbf{A}^+ - \mathbf{A}^+ \mathbf{V} \mathbf{C}^* + \mathbf{C} (\mathbf{G}^+ + \mathbf{V}^* \mathbf{A}^+ \mathbf{V}) \mathbf{C}^* . \quad (4)$$

For concreteness, we specialise the preceding identities to the case of rank one perturbations. In this special case,  $k \equiv 1$ , and  $\mathbf{V}_i$  and  $\mathbf{W}_i$  reduce to  $\ell$  vectors,  $v_i$  and  $w_i$ . In the nonsingular case, (1) reduces to Bartlett's identity [6]. It states for an arbitrary nonsingular  $\ell \times \ell$  matrix  $\mathbf{A}$  and  $\ell$  vectors  $v_i$ ,

$$(\mathbf{A} + v_1 v_2^*)^{-1} = \mathbf{A}^{-1} - \frac{(\mathbf{A}^{-1} v_1)(v_2^* \mathbf{A}^{-1})}{(1 + v_2^* \mathbf{A}^{-1} v_1)} . \quad (5)$$

In this case, theorem 1 reduces to the analogous result for an arbitrary singular matrix  $\mathbf{A}$  with a rank one perturbation which contains a component perpendicular to the column space of  $\mathbf{A}$ . Noting that  $\mathbf{G} \equiv 1$  and  $\mathbf{C}_i \equiv w_i/|w_i|^2$ , theorem 1 simplifies to the following result.

**Theorem 3.** Let  $\mathbf{A}$  be a  $\ell \times \ell$  matrix of rank  $\ell_1$ ,  $\ell_1 < \ell$ , and  $v_i, w_i$ ,  $i = 1, 2$  be  $\ell$  vectors. Let  $v_1 \in M(\mathbf{A})$  and  $w_1$  be orthogonal to  $M(\mathbf{A})$ , and  $v_2 \in M(\mathbf{A}^*)$  and  $w_2$  be orthogonal to  $M(\mathbf{A}^*)$ . Assume  $w_2$  is parallel to  $w_1$  and  $w_i \neq 0$ . Let

$$\Omega \equiv \mathbf{A} + (v_1 + w_1)(v_2 + w_2)^* ,$$

The Moore-Penrose generalized inverse is

$$\Omega^+ = \mathbf{A}^+ - \frac{w_2 v_2^* \mathbf{A}^+}{|w_2|^2} - \frac{\mathbf{A}^+ v_1 w_1^*}{|w_1|^2} + (1 + v_2^* \mathbf{A}^+ v_1) \frac{w_2 w_1^*}{|w_1|^2 |w_2|^2} . \quad (6)$$

This generalized inverse is singular and tends to infinity as  $1/|w_1||w_2|$ , as  $w_i$  approaches to zero. Thus (6) does not reduce to Bartlett's identity.

The projection operator onto the row space of  $\Omega$  is

$$P_{X_T} = \mathbf{A}^+ \mathbf{A} + \frac{w_i w_i^*}{|w_i|^2} .$$

The symmetric version of Thm. 3 was originally developed and applied by the author in his statistical analysis of magnetic fusion data [7]. To estimate the regression parameters in ordinary least squares regression, the sum of squares and products (SSP) matrix needs to be inverted. We apply Thm. 3 to determine the inverse of the SSP matrix in terms of the inverse of the covariance matrix of the covariates.

We decompose the independent variable vector,  $x$  into a mean value vector,  $\bar{x}$  and a fluctuating part,  $\tilde{x}$ . Thus the  $i$ -th individual observation has the form

$$x_i = \bar{x} + \tilde{x}_i .$$

Let  $\mathbf{X}$  denote the  $n \times \ell$  data matrix whose rows consist of  $x_i^*$  and  $\tilde{\mathbf{X}}$  be the centered data matrix whose rows consist of  $\tilde{x}_i^*$ .

We assume that some of the independent variables,  $x_k$ , have not been varied. Thus  $\tilde{\mathbf{X}}^* \tilde{\mathbf{X}}$  is singular. The inverse of the uncentered sum of squares and crossproducts

matrix,  $\mathbf{X}^*\mathbf{X}$  can now be expressed in terms of the Moore Penrose generalized inverse of the centered covariance matrix,  $\tilde{\mathbf{X}}^*\tilde{\mathbf{X}}$ .

We decompose a multiple of the mean value vector,  $\sqrt{n}\bar{x}$ , into  $v + w$ , where  $v \in M(\tilde{\mathbf{X}}^*\tilde{\mathbf{X}})$  and  $w \perp M(\tilde{\mathbf{X}}^*\tilde{\mathbf{X}})$ .

The data matrix has the form

$$\mathbf{X}^*\mathbf{X} = \tilde{\mathbf{X}}^*\tilde{\mathbf{X}} + n\bar{x}\bar{x}^T = \tilde{\mathbf{X}}^*\tilde{\mathbf{X}} + (v + w)(v + w)^*.$$

Thus we have rewritten  $\mathbf{X}^*\mathbf{X}$  in a form appropriate to the application of theorem 3.

In conclusion, the application of these matrix identities requires the decomposition of  $\mathbf{X}_i$  into the orthogonal components,  $\mathbf{V}_i$  and  $\mathbf{W}_i$ . Thus our theorems are most useful in situations where the decomposition is trivial.

#### *Acknowledgments*

The helpful comments of the referees are gratefully acknowledged.

### REFERENCES

1. W.J. Duncan, "Some devices for the solution of large sets of simultaneous equations (with an appendix on the reciprocation of partitioned matrices)", *The London, Edinburgh and Dublin Philosophical Magazine and Journal of Science*, Seventh Series, **35**, p. 660, (1944).
2. H.V. Henderson and S.R. Searle, "On deriving the inverse of a sum of matrices", *SIAM Review*, **23**, p.53, (1981).
3. D.V. Ouellete, "Schur complements and statistics", *Journal of Linear Algebra*, **36**, p. 187, (1981).
4. W.W. Hager, "Updating the inverse of a matrix", *SIAM Review*, **31**, p.221, (1989).
5. C.R. Rao, *Linear Statistical Inference and Its Applications*, p. 26,33, J. Wiley and Sons, New York, 1973.
6. M.S. Bartlett, "An inverse matrix adjustment arising in discriminant analysis", *Annals of Mathematical Statistics*, vol. 22, p107, (1951).
7. K.S. Riedel, New York University Report MF-118, National Technical Information Service document no. DOE/ER/53223-98 "Advanced Statistics for Tokamak Transport: Collinearity and Tokamak to Tokamak Variation" (1989).