

Pierre Hodara* and Ioannis Papageorgiou **
 IME, Universidade de Sao Paulo

POINCARÉ TYPE INEQUALITIES FOR COMPACT DEGENERATE PURE JUMP MARKOV PROCESSES.

PIERRE HODARA AND IOANNIS PAPAGEORGIU

ABSTRACT. We aim in proving Poincaré inequalities for a class of pure jump Markov processes inspired by the model introduced in [16] by Galves and Löcherbach to describe the behaviour of interacting brain neurons. In particular, we consider neurons with degenerate jumps, i.e. that lose their memory when they spike, while the probability of a spike depends on the actual position and thus the past of the whole neural system.

1. INTRODUCTION

The aim of this paper is to prove Poincaré inequalities for the semigroup P_t , as well as for the invariant measure, of the model introduced in [16] by Galves and Löcherbach, to describe the activity of a biological neural network. What is in particular interesting about the jump process in question is that it is characterized by degenerate jumps, in the sense that after a particle (neuron) spikes, it loses its memory by jumping to zero. Furthermore, the probability of a spike of a particular neuron at any time depends on its actual position and thus the past of the whole neural system.

For P_t the associated semigroup at first we prove some Poincaré type inequalities of the form

$$\text{Var}_{P_t}(f(x)) \leq \alpha(t) \int_0^t P_s \Gamma(f, f)(x) ds + \beta(t) \sum_{i=1}^n \int_0^t P_s \Gamma(f, f)(\Delta_i(x)) ds.$$

for any possible starting configuration x . We give here the general form of the type of inequalities investigated in this paper, however in order to avoid to overload this

2010 *Mathematics Subject Classification.* 60K35, 26D10, 60G99,

Key words and phrases. Talagrand inequality, Poincaré inequality, brain neuron networks.

Address: Neuromat, Instituto de Matematica e Estatistica, Universidade de Sao Paulo, rua do Matao 1010, Cidade Universitaria, Sao Paulo - SP- Brasil - CEP 05508-090.

Email: *hodarapierre@gmail.com ** ipapageo@ime.usp.br, papyannis@yahoo.com

This article was produced as part of the activities of FAPESP Research, Innovation and Dissemination Center for Neuromathematics (grant 2013/ 07699-0 , S.Paulo Research Foundation);

This article is supported by FAPESP grant (2016/17655-8) * and (2017/15587-8)** .

introduction with technical details we postpone the definitions of classical quantities such as the "carré du champ" and other notations used here to subsection 1.3.

Then, we restrict ourselves to the special case where the initial configuration x in the domain of the invariant measure, and we derive the stronger Poincaré inequality

$$\text{Var}_{P_t}(f(x)) \leq \alpha(t)P_t\Gamma(f, f)(x) + \beta \int_0^t P_s\Gamma(f, f)(x)ds.$$

Then we show a Poincaré inequality for the invariant measure π

$$\text{Var}_{\pi}(f) \leq c\pi(\Gamma(f, f)).$$

Before we describe the model we present the neuroscience framework of the problem.

1.1. the neuroscience framework. The activity of one neuron is described by the evolution of its membrane potential. This evolution presents from time to time a brief and high-amplitude depolarisation called action potential or spike. The spiking probability or rate of a given neuron depends on the value of its membrane potential. These spikes are the only perturbations of the membrane potential that can be transmitted from one neuron to another through chemical synapses. When a neuron spikes, its membrane potential is reset to 0 and the post-synaptic neurons connected to it receive an additional amount of membrane potential.

From a probabilistic point of view, this activity can be described by a simple point process since the whole dynamic is characterised by the jump times. In the literature, Hawkes processes are often used in order to describe systems of interacting neurons, see [9], [14], [15], [16], [18] and [20] for example. The reset to 0 of the spiking neuron provides a variable length memory for the dynamic and therefore point-processes describing these systems are non-Markovian.

On the other hand, it is possible to describe the activity of the network with a process modelling not only the jump times but the whole evolution of the membrane potential of each neuron. This evolution needs then to be specified between the jumps. In [19] the process describing this evolution follows a deterministic drift between the jumps, more precisely the membrane potential of each neuron is attracted with exponential speed towards an equilibrium potential. This process is then Markovian and belongs to the family of Piecewise Deterministic Markov Processes introduced by Davis ([11] and [12]). Such processes are widely used in probability modeling of *e.g.* biological or chemical phenomena (see *e.g.* [10] or [22], see [2] for an overview). The point of view we adopt here is close to this framework, but we work without drift between the jumps. We therefore consider a pure jump Markov process and will make use of the abbreviation PJMP in the rest of the present work.

We consider a process $X_t = (X_t^1, \dots, X_t^N)$, where N is the number of neurons in the network and where for each neuron i , $1 \leq i \leq N$ and each time $t \in \mathbb{R}_+$, each variable X_t^i represents the membrane potential of neuron i at time t . Each membrane potential X_t^i takes value in \mathbb{R}_+ . A neuron with membrane potential x “spikes” with intensity $\phi(x)$, where $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a given intensity function. When a neuron i fires, its membrane potential is reset to 0, interpreted as resting potential, while the membrane potential of any post-synaptic neuron j is increased by $W_{i \rightarrow j} \geq 0$. Between two jumps of the system, the membrane potential of each neuron is constant.

Working with Hawkes processes allows to consider systems with infinitely many neurons, as in [16] or [20]. For our purpose, we need to work in a Markovian framework and therefore our process represents the membrane potentials of the neurons, considering a finite number N of neurons.

1.2. the model. Let $N > 1$ be fixed and $(N^i(ds, dz))_{i=1, \dots, N}$ be a family of *i.i.d.* Poisson random measures on $\mathbb{R}_+ \times \mathbb{R}_+$ having intensity measure $dsdz$. We study the Markov process $X_t = (X_t^1, \dots, X_t^N)$ taking values in \mathbb{R}_+^N and solving, for $i = 1, \dots, N$, for $t \geq 0$,

$$(1.1) \quad \begin{aligned} X_t^i &= X_0^i - \int_0^t \int_0^\infty X_{s-}^i 1_{\{z \leq \phi(X_{s-}^i)\}} N^i(ds, dz) \\ &\quad + \sum_{j \neq i} W_{j \rightarrow i} \int_0^t \int_0^\infty 1_{\{z \leq \phi(X_{s-}^j)\}} N^j(ds, dz). \end{aligned}$$

In the above equation for each $j \neq i$, $W_{j \rightarrow i} \in \mathbb{R}_+$ is the synaptic weight describing the influence of neuron j on neuron i . Finally, the function $\phi : \mathbb{R}_+ \mapsto \mathbb{R}_+$ is the intensity function.

The generator of the process X is given for any test function $f : \mathbb{R}_+^N \rightarrow \mathbb{R}$ and $x \in \mathbb{R}_+^N$ is described by

$$(1.2) \quad \mathcal{L}f(x) = \sum_{i=1}^N \phi(x^i) [f(\Delta_i(x)) - f(x)]$$

where

$$(1.3) \quad (\Delta_i(x))_j = \begin{cases} x^j + W_{i \rightarrow j} & j \neq i \text{ and } x^i + W_{i \rightarrow j} \leq m \\ x^j & j \neq i \text{ and } x^i + W_{i \rightarrow j} > m \\ 0 & j = i \end{cases}$$

for some $m > 0$. With this definition the process remains inside the compact set

$$(1.4) \quad D := \{x \in \mathbb{R}_+^N : x_i \leq m, 1 \leq i \leq N\}.$$

Furthermore, we also assume the following conditions about the intensity function:

$$(1.5) \quad \phi(x) > cx \text{ for } x \in \mathbb{R}_+$$

and

$$(1.6) \quad \phi(x) \geq \delta$$

for some strictly positive constants c and δ .

The probability for a neuron to spike grows with its membrane potential so it is natural to think of the function ϕ as an increasing function. Condition (1.5) implies that this growth is at least linear. Condition (1.6) models the spontaneous activity of the system: whatever the configuration x is, the system will always have a positive spiking rate.

1.3. Poincaré type inequalities. Our purpose is to show Poincaré type inequalities for our PJMP, whose dynamic is similar to the model introduced in [16]. We will investigate Poincaré type inequalities at first for the semigroup P_t and then for the invariant measure π . Concerning the semigroup inequality we will study two different cases. The first, the general one, where the system starts from any possible initial configuration. Then, we restrict to initial configurations that belong to the domain of the invariant measure.

Let us first describe the general framework and define the Poincaré inequalities on a discrete setting (see also [23], [13], [24], [1] and [8]). At first we should note a convention we will widely use. For a function f and measure ν we will write $\nu(f)$ for the expectation of the function f with respect to the measure ν , that is

$$\nu(f) = \int f d\nu.$$

We consider a Markov process $(X_t)_{t \geq 0}$ which is described by the infinitesimal generator \mathcal{L} and the associated Markov semigroup $P_t f(x) = \mathbb{E}^x(f(X_t))$. For a semigroup and its associated infinitesimal generator we will need the following well known relationships: $\frac{d}{ds} P_s = \mathcal{L} P_s = P_s \mathcal{L}$ (see for example [17]).

We define π to be the invariant measure for the semigroup $(P_t)_{t \geq 0}$ if and only if

$$\pi P_t = \pi.$$

Furthermore, we define the "carré du champ" operator by:

$$\Gamma(f, g) := \frac{1}{2}(\mathcal{L}(fg) - f\mathcal{L}g - g\mathcal{L}f).$$

For the PJMP process defined above with the specific generator \mathcal{L} given by (1.2) a simple calculation shows that the carré du champ takes the following form.

$$\Gamma(f, f) = \frac{1}{2}(\mathcal{L}f^2 - 2f\mathcal{L}f) = \frac{1}{2}\left(\sum_{i=1}^N \phi(x^i) [f(\Delta_i(x)) - f(x)]^2\right).$$

We say that a measure ν satisfies a Poincaré inequality if there exists constant $C > 0$ independent of f , such that

$$(SG) \quad \text{Var}_\nu(f) \leq C\nu(\Gamma(f, f))$$

where the variance of a function f with respect to a measure ν is defined with the usual way as: $\text{Var}_\nu(f) = \nu(f - \nu(f))^2$. It should be noted that in the case where the measure ν is the semigroup P_t , then the constant C may depend on t , $C = C(t)$.

In [24], [1] and [8], the Poincaré inequality (SG) for P_t has been shown for some point processes, for a constant that depends on time t , while the stronger log-Sobolev inequality, has been disproved. The general method used in these papers, that will be followed also in the current work, is based on the so called semigroup method which shows the inequality for the semigroup P_t .

The main difficulty here is that, for the pure jump Markov process that we examine in the current paper, the translation property

$$\mathbb{E}^{x+y}f(z) = \mathbb{E}^x f(z + y)$$

used in [24] and [1] does not hold here. This appears to be important because the translation property is a key element in these papers, since it allows to bound the carré du champ by comparing the mean $\mathbb{E}^x f(z)$ where the process starts from position x with the mean $\mathbb{E}^{\Delta_i(x)} f(z)$ where it starts from $\Delta_i(x)$, the jump-neighbour of x . However, we can still obtain Poincaré type inequalities, but with a constant $C(t)$ which is a polynomial of order higher than one. This order is higher than the constant $C(t) = t$, the optimal obtained in [24] for a path space of Poisson point processes.

It should be noted, that the the aforementioned translation property relates with the Γ_2 criterion (see [3] and [4]) for the Poincaré inequality (see discussion in subsection 2.2). Since this is not satisfied in our case we obtain a Poincaré type inequality instead.

Before we present the results of the paper it is important to highlight an important distinction on the nature of the initial configuration from which the process can start. We can classify the initial configurations according to the return probability to them. Recall that the membrane potential x_i of every neuron i takes positive values within a compact set and that whenever a neuron j different than i spikes, the neuron i jumps $W_{i \rightarrow j}$ positions up, while the only other movement it does is jumping to zero when it spikes. That means that every variable x_i can jump down only to zero while after the first jump, can only pass from a finite number of possible positions. Since the neurons stay still between spikes, that implies that there is a finite number of possible configurations to which $X = (X^1, \dots, X^N)$ can return after every neuron has spiked for the first time. This is the domain of the invariant measure π of the semigroup P_t , and we will denote it as \hat{D} . Thus, if the

initial configuration $x = (x^1, \dots, x^N)$ does not belong to \hat{D} , after the process enters \hat{D} , it will never return back to this initial configuration.

It should be noted that it is easy to find initial configurations $x = (x^1, \dots, x^N) \notin \hat{D}$. For example one can consider any x such that at least one of the x^i 's is not a sum of synaptic weights $W_{j \rightarrow i}$, or any x with $x^i = x^j$ for every i and j .

Below we present the Poincaré inequality for the semigroup P_t for general starting configurations.

Theorem 1.1. *Assume the PJMP as described in (1.2)-(1.6). Then, for every $x \in D$, the following Poincaré type inequality holds.*

$$\text{Var}_{E^x}(f(x_t)) \leq \alpha(t) \int_0^t P_s \Gamma(f, f)(x) ds + \beta \sum_{i=1}^n \int_0^t P_s \Gamma(f, f)(\Delta_i(x)) ds$$

with $\alpha(t)$ a second order polynomial of the time t that does not depend on the function f and β a constant.

One notices that since the coefficient $\alpha(t)$ is a polynomial of first order and β is a constant, the first term dominates over the second for big time t , as shown in the next corollary.

Corollary 1.2. *Assume the PJMP as described in (1.2)-(1.6). Then, for every $x \in D$, and t sufficiently large, i.e. $t > \zeta(f)$*

$$\text{Var}_{E^x}(f(x_t)) \leq 2\alpha(t) \int_0^t P_w \Gamma((f, f))(x) dw$$

where $\zeta(f)$ is a constant depending only on f ,

$$\zeta(f) = \left(\frac{6t_0 \sum_{i=1}^n P_t \Gamma(f, f)(\Delta_i(x))}{(1 + C_1) \int_0^t P_s \Gamma(f, f)(x) ds} \right)^{\frac{1}{2}}$$

for some positive constants t_0 , C_1 and M .

One should notice that although the lower value $\zeta(f)$ depends on the function f , the coefficient $2\alpha(t)$ of the inequality does not depend on the function f .

The proof of the Poincaré inequality for the general initial configuration is presented in section 2.

In the special case where x is on the domain of the invariant measure we obtain the stronger inequality

Theorem 1.3. *Assume the PJMP as described in (1.2)-(1.6). Then, there exists a $t_1 > 0$ such that, for every $t > t_1$ and for every $x \in \hat{D}$, the following Poincaré*

type inequality holds.

$$\text{Var}_{E^x}(f(x_t)) \leq \gamma(t)P_t\Gamma(f, f)(x) + 2 \int_0^t P_s\Gamma(f, f)(x)ds.$$

with $\gamma(t)$ a third order polynomial of the time t that do not depend on the function f .

As in the general case, for t large enough we have the following corollary.

Corollary 1.4. *Assume the PJMP as described in (1.2)-(1.6). Then, there exists a $t_1 > 0$ such that, for every for every $x \in \hat{D}$, and t sufficiently large, i.e. $t > \max\{\xi(f), t_1\}$*

$$\text{Var}_{E^x}(f(x_t)) \leq 2\gamma(t)P_t\Gamma(f, f)(x)$$

where $\xi(f)$ is a constant depending only on f ,

$$\xi(f) = \left(\frac{\int_0^t P_s\Gamma(f, f)(x)ds}{4\theta^2 M^2 N P_t\Gamma(f, f)(x)} \right)^{\frac{1}{3}}$$

for some positive constants θ and M .

We conclude this section with the Poincaré inequality for the invariant measure π presented on the next theorem.

Theorem 1.5. *Assume the PJMP as described in (1.2)-(1.6). Then π satisfies a Poincaré inequality*

$$\pi(f - \pi f)^2 \leq C_0\pi(\Gamma(f, f))$$

for some constant $C_0 > 0$.

2. PROOF OF THE POINCARÉ FOR GENERAL INITIAL CONFIGURATIONS.

In this section we focus on neurons that start with values on any possible initial configuration $x \in D$ as described by (1.2)-(1.6), and we prove the local Poincaré inequalities presented in Theorem 1.1 and Corollary 1.2. Let us first state some technical results.

2.1. Technical results. We start by showing properties of the jump probabilities of the degenerate PJMP processes. Since the process is constant between jumps, the set of reachable positions y after a given time t for a trajectory starting from x is discrete. We therefore define

$$\pi_t(x, y) := P_x(X_t = y) \text{ and } D_x := \{y \in D, \pi_t(x, y) > 0\}.$$

This set is finite for the following reasons. On one hand, for each neuron $i \in I$, the set $S_i = \{0\} \cup \{\sum_{k=1}^n W_{j_k \rightarrow i}, n \in \mathbb{N}^*, j_k \in I\}$ is discrete and such that

the intersection with any compact is finite. On the other hand, we have $D_x \subset \left[\prod_{i \in I} (S_i \cup (x_i + S_i)) \right] \cap D$.

The idea is that since the process is constant between jumps, elements of D_x are such that there exists a sequence of jumps leading from x to y . Since we are only interested on the arrival position y , among all jump sequences leading to y , we can consider only sequences with minimal number of jumps and the number of such jump sequences leading to positions inside a compact is finite, due to the fact that each $W_{j \rightarrow i}$ is non-negative.

Since x is also in the compact D , we can have an upper bound for the cardinal of D_x independent from x .

For a given time $s \in \mathbb{R}_+$ and a given position $x \in D$, we denote by $p_s(x)$ the probability that starting at time 0 from position x , the process has no jump in the interval $[0, s]$, and for a given neuron $i \in I$ by $p_s^i(x)$ the probability that the process has exactly one jump of neuron i and no jumps for other neurons.

Introducing the notation $\bar{\phi}(x) = \sum_{j \in I} \phi(x_j)$ and given the dynamics of the model, we have that

$$p_s(x) = e^{-s\bar{\phi}(x)}$$

and

$$(2.1) \quad p_s^i(x) = \int_0^s \phi(x_i) e^{-u\bar{\phi}(x)} e^{-(s-u)\bar{\phi}(\Delta^i(x))} du \\ = \left\{ \begin{array}{ll} \frac{\phi(x_i)}{\bar{\phi}(x) - \bar{\phi}(\Delta^i(x))} \left(e^{-s\bar{\phi}(\Delta^i(x))} - e^{-s\bar{\phi}(x)} \right) & \text{if } \bar{\phi}(\Delta^i(x)) \neq \bar{\phi}(x) \\ s\phi(x_i) e^{-s\bar{\phi}(x)} & \text{if } \bar{\phi}(\Delta^i(x)) = \bar{\phi}(x) \end{array} \right\}.$$

Define

$$(2.2) \quad t_0 = \left\{ \begin{array}{ll} \frac{\ln(\bar{\phi}(x)) - \ln(\bar{\phi}(\Delta^i(x)))}{\bar{\phi}(x) - \bar{\phi}(\Delta^i(x))} & \text{if } \bar{\phi}(\Delta^i(x)) \neq \bar{\phi}(x) \\ \frac{1}{\bar{\phi}(x)} & \text{if } \bar{\phi}(\Delta^i(x)) = \bar{\phi}(x) \end{array} \right\}.$$

As a function of s , $p_s^i(x)$ is continuous, strictly increasing on $(0, t_0)$ and strictly decreasing on $(t_0, +\infty)$ and we have $p_0^i(x) = 0$.

Lemma 2.1. *Assume the PJMP as described in (1.2)-(1.6). There exists positive constants C_1 and C_2 independent of t , x and y such that*

- For all $t > t_0$, we have

$$\sum_{y \in D_x} \frac{\pi_t^2(\Delta^i(x), y)}{\pi_t(x, y)} \leq C_1.$$

- For all $t \leq t_0$, we have

$$\sum_{y \in D_x \setminus \{\Delta^i(x)\}} \frac{\pi_t^2(\Delta^i(x), y)}{\pi_t(x, y)} \leq C_1$$

and

$$\frac{\pi_t^2(\Delta^i(x), \Delta^i(x))}{\pi_t(x, \Delta^i(x))} \leq \frac{\pi_t(\Delta^i(x), \Delta^i(x))}{\pi_t(x, \Delta^i(x))} \leq \frac{C_2}{t}.$$

Proof. As said before, the set D_x is finite so it is sufficient to obtain an upper bound for the ratio $\frac{\pi_t^2(\Delta^i(x), y)}{\pi_t(x, y)}$.

We have for all $s \in (0, t)$

$$(2.3) \quad \frac{\pi_t^2(\Delta^i(x), y)}{\pi_t(x, y)} \leq \frac{(\pi_{t-s}(\Delta^i(x), y)p_s(y) + \sup_{z \in D}(1 - p_s(z)))^2}{p_s^i(x)\pi_{t-s}(\Delta^i(x), y)}.$$

Here we decomposed the numerator according to two events. Either $X_{t-s} = y$ and there no jump in the interval of time $[t-s, t]$ or there is at least one jump in the interval of time $[t-s, t]$, whatever the position $z \in D$ of the process at time $t-s$.

From the previous inequality, we then obtain

$$(2.4) \quad \frac{\pi_t^2(\Delta^i(x), y)}{\pi_t(x, y)} \leq \frac{(\pi_{t-s}(\Delta^i(x), y)p_s(y) + (1 - e^{-sN\phi(m)}))^2}{p_s^i(x)\pi_{t-s}(\Delta^i(x), y)}.$$

where we recall that the constant m appears in the definition of the compact set D introduced in (1.4).

Let us first assume that $t > t_0$. Recall that t_0 is defined in (2.2).

If $\pi_{t-t_0}(\Delta^i(x), y) \geq p_{t_0}^i(x)$, we have

$$(2.5) \quad \frac{\pi_t^2(\Delta^i(x), y)}{\pi_t(x, y)} \leq \frac{1}{(p_{t_0}^i(x))^2}.$$

Assume now that $\pi_{t-t_0}(\Delta^i(x), y) < p_{t_0}^i(x)$ and let us recall that as a function of s , $p_s^i(x)$ is continuous, strictly increasing on $(0, t_0)$ and $p_0^i(x) = 0$.

On the other hand, as a function of s , $\pi_{t-s}(\Delta^i(x), y)$ is continuous and takes value $\pi_t(\Delta^i(x), y) > 0$ for $s = 0$.

We deduce from this that there exists $s_* \in (0, t_0)$ such that $p_{s_*}^i(x) = \pi_{t-s_*}(\Delta^i(x), y)$.

Now (2.4) with $s = s_*$ gives us

$$(2.6) \quad \frac{\pi_t^2(\Delta^i(x), y)}{\pi_t(x, y)} \leq (p_{s_*}(y))^2 + 2p_{s_*}(y) \frac{1 - e^{-s_* N \phi(m)}}{p_{s_*}^i(x)} + \left(\frac{1 - e^{-s_* N \phi(m)}}{p_{s_*}^i(x)} \right)^2.$$

For all $s \in (0, t_0)$, $p_s(y) \leq 1$, and we then study $\frac{1 - e^{-s N \phi(m)}}{p_s^i(x)}$ as a function of $s \in (0, t_0)$.

Using the explicit value of $p_s^i(x)$ given in (2.1) and assumption (1.5), we obtain for all $s \in (0, t_0)$,

$$\frac{1 - e^{-s N \phi(m)}}{p_s^i(x)} \leq \left\{ \begin{array}{ll} \frac{e^{t_0 N \phi(m)}}{\delta} \frac{(\bar{\phi}(x) - \bar{\phi}(\Delta^i(x)))(1 - e^{-s N \phi(m)})}{1 - e^{-s(\bar{\phi}(x) - \bar{\phi}(\Delta^i(x)))}} & \text{if } \bar{\phi}(\Delta^i(x)) \neq \bar{\phi}(x) \\ \frac{e^{t_0 N \phi(m)}}{\delta} \frac{1 - e^{-s}}{s} & \text{if } \bar{\phi}(\Delta^i(x)) = \bar{\phi}(x) \end{array} \right\}.$$

Recall that $\delta > 0$ is defined in assumption (1.6) and satisfies $\phi(x) \geq \delta$ for all $x \in \mathbb{R}_+$.

In both cases, when s is far from zero, we can obtain an upper bound independent of x , and when s goes to zero, the limit of the right hand term is $\frac{N \phi(m) e^{t_0 N \phi(m)}}{\delta}$.

From this, we deduce that there exists a constant M_D such that for all $s \in (0, t_0)$,

$$\frac{1 - e^{-s N \phi(m)}}{p_s^i(x)} \leq M_D.$$

Putting all together, we obtain the announced result for the case where $t > t_0$.

We now consider the case where $t \leq t_0$.

We start by considering the case where $y \neq \Delta^i(x)$ and go back to (2.4).

As a function of s , $\pi_{t-s}(\Delta^i(x), y)$ is continuous and takes values $\pi_t(\Delta^i(x), y) > 0$ and $\pi_0(\Delta^i(x), y) = 0$ respectively for $s = 0$ and $s = t$.

We deduce from this that there exists $s_* \in (0, t) \subset (0, t_0)$ such that $p_{s_*}^i(x) = \pi_{t-s_*}(\Delta^i(x), y)$ and we are back in the previous case so that the same result holds.

Let us assume now that $y = \Delta^i(x)$, we have

$$(2.7) \quad \frac{\pi_t^2(\Delta^i(x), y)}{\pi_t(x, y)} \leq \frac{\pi_t(\Delta^i(x), \Delta^i(x))}{\pi_t(x, \Delta^i(x))} \leq \frac{1}{p_t^i(x)}.$$

Recall the explicit expression of $p_t^i(x)$ given in (2.1) and use (1.6) to bound the intensity function

$$p_t^i(x) = t \phi(x_i) e^{-t \bar{\phi}(x)} \geq t \delta e^{-t_0 \sup_{x \in D} \phi(x)} = Ct$$

for some constant C independent of t and x , which gives us the announced result. \square

Taking under account the last result, we can obtain the first technical bound needed in the proof of the local Poincaré inequality.

Lemma 2.2. *Assume the PJMP as described in (1.2)-(1.6). Then*

$$\left(\int_{t_0}^{t-s} (\mathbb{E}^{\Delta_i(x)} - \mathbb{E}^x) \sum_{j=1}^N \phi(x_u^j) (f(\Delta_j(x_u)) - f(x_u)) du \right)^2 \leq (t-s)(1+C_1)M \int_{t_0}^{t-s} \mathbb{E}^x \Gamma((f, f))(x_u).$$

Proof. Consider $\pi_t(x, y)$ to probability kernel of \mathbb{E}^x , i.e. $\mathbb{E}^x(f(x_t)) = \sum_y \pi_t(x, y) f(y)$. Then we can write

$$\begin{aligned} \mathbf{II}_1 &:= \left(\int_{t_0}^{t-s} (\mathbb{E}^{\Delta_i(x)} - \mathbb{E}^x) \sum_{j=1}^N \phi(x_u^j) (f(\Delta_j(x_u)) - f(x_u)) du \right)^2 = \\ &= \left(\int_{t_0}^{t-s} \sum_y \pi_u(x, y) \left(\frac{\pi_u(\Delta_i(x), y)}{\pi_u(x, y)} - 1 \right) \sum_{j=1}^N \phi_u(y^j) (f(\Delta_j(y)) - f(y)) du \right)^2. \end{aligned}$$

To continue we will use Holder's inequality to pass the second power inside the first integral, which will give

$$\mathbf{II}_1 \leq (t-s) \int_{t_0}^{t-s} \left(\sum_y \pi_u(x, y) \left(\frac{\pi_u(\Delta_i(x), y)}{\pi_u(x, y)} - 1 \right) \sum_{j=1}^N \phi_u(y^j) (f(\Delta_j(y)) - f(y)) \right)^2 du$$

and then we apply the Cauchy-Schwarz inequality for the measure \mathbb{E}^x to get

$$\mathbf{II}_1 \leq (t-s) \int_{t_0}^{t-s} \mathbb{E}^x \left(\frac{\pi_u(\Delta_i(x), y)}{\pi_u(x, y)} - 1 \right)^2 * \mathbb{E}^x \left(\underbrace{\sum_{j=1}^N \phi(y_u^j) (f(\Delta_j(y_u)) - f(y_u))}_{:=\mathbb{S}} \right)^2 du.$$

The first quantity involved in the above integral is bounded from Lemma 2.1 by a constant

$$\mathbb{E}^x \left(\frac{\pi_u(\Delta_i(x), y)}{\pi_u(x, y)} - 1 \right)^2 \leq 1 + E^x \left(\frac{\pi_u(\Delta_i(x), y)}{\pi_u(x, y)} \right)^2 = 1 + \sum_{y \in D_x} \frac{\pi_u^2(\Delta_i(x), y)}{\pi_u(x, y)} \leq 1 + C_1$$

while for the sum involved in the second quantity, since $\phi(x) \geq \delta > 0$ we can use Holder's inequality

$$\begin{aligned} \mathbb{S} &= \left(\sum_{j=1}^N \phi(y_u^j) \right)^2 \left(\sum_{j=1}^N \frac{\phi(y_u^j)}{\sum_{j=1}^N \phi(y_u^j)} (f(\Delta_j(y_u)) - f(y_u)) \right)^2 \leq \\ &\leq M \sum_{j=1}^N \phi(y_u^j) (f(\Delta_j(y_u)) - f(y_u))^2 = M\Gamma(f, f)(x_u) \end{aligned}$$

where $M = \sup_{x \in D} \sum_{i=1}^N \phi(x^i)$, so that

$$\mathbf{II}_1 \leq (t-s)(1+C_1)M \int_{t_0}^{t-s} \mathbb{E}^x \Gamma((f, f))(x_u) du.$$

□

We will now extend the last bound to an integral on a time domain starting at 0.

Lemma 2.3. *For the PJMP as described in (1.2)-(1.6), we have*

$$\begin{aligned} \left(\int_0^{t-s} (\mathbb{E}^{\Delta_i(x)}(\mathcal{L}f(x_u)) - \mathbb{E}^x(\mathcal{L}f(x_u))) du \right)^2 &\leq 16t_0^2 M\Gamma(f, f)(\Delta_i(x)) + \\ &+ c(t-s) \int_0^{t-s} \mathbb{E}^x \Gamma((f, f))(x_u) du \end{aligned}$$

where $c(t) = t_0 8M(C_1 + 1) + 2t(1 + C_1)M$.

Proof. In order to calculate a bound for

$$\mathbb{E}^{\Delta_i(x)}(\mathcal{L}f(x_u)) - \mathbb{E}^x(\mathcal{L}f(x_u))$$

we will need to control the ration $\frac{\pi_u^2(\Delta_i(x), y)}{\pi_u(x, y)}$. As shown in Lemma 2.1, this ratio depends on time when $u \leq t_0$, otherwise it is bounded by a constant. For this reason we will start by breaking the integration variable of the time t into two domains, $(0, t_0)$ and $(t_0, t-s)$.

$$\begin{aligned} \mathbf{I}_1 &:= \left(\int_0^{t-s} (\mathbb{E}^{\Delta_i(x)}(\mathcal{L}f(x_u)) - \mathbb{E}^x(\mathcal{L}f(x_u))) du \right)^2 \leq \\ &2 \underbrace{\left(\int_{t_0}^{t-s} (\mathbb{E}^{\Delta_i(x)} - \mathbb{E}^x) \sum_{i=1}^N \phi(x_u^i) (f(\Delta_i(x_u)) - f(x_u)) du \right)^2}_{\mathbf{II}_1} + \\ (2.8) \quad &+ 2 \underbrace{\left(\int_0^{t_0} (\mathbb{E}^{\Delta_i(x)} - \mathbb{E}^x) \sum_{i=1}^N \phi(x_u^i) (f(\Delta_i(x_u)) - f(x_u)) du \right)^2}_{:=\mathbf{II}_2}. \end{aligned}$$

The first summand \mathbf{II}_1 is upper bounded by the previous lemma. To bound the second term \mathbf{II}_2 on the right hand side of (2.8) we write

$$\begin{aligned}
\mathbf{II}_2 &\leq \\
&2 \underbrace{\left(\int_0^{t_0} (\pi_u(\Delta_i(x), \Delta_i(x)) - \pi_u(x, \Delta_i(x))) \sum_{i=1}^N \phi(\Delta_i(x)^i) (f(\Delta_i(\Delta_i(x))) - f(\Delta_i(x))) du \right)^2}_{:=\mathbf{III}_1} + \\
(2.9) \quad &+ 2 \underbrace{\left(\int_0^{t_0} \left(\sum_{y \in D, y \neq \Delta_i(x)} (\pi_u(\Delta_i(x), y) - \pi_u(x, y)) \sum_{i=1}^N \phi(y^i) (f(\Delta_i(y)) - f(y)) \right) du \right)^2}_{:=\mathbf{III}_2}.
\end{aligned}$$

The distinction on the two cases, whether after time u the neurons configuration is $\Delta_i(x)$ or not, relates to the two different bounds Lemma 2.1 provides for the fraction $\frac{\pi_u^2(\Delta_i(x), y)}{\pi_u(x, y)}$ whether y is $\Delta_i(x)$ or not. We will first calculate the second term on the right hand side. For this term we will work similar to Lemma 2.2. At first we will apply the Holder inequality on the time integral after we first divide with the normalisation constant t_0 . This will give

$$\mathbf{III}_2 \leq t_0 \int_0^{t_0} \left(\sum_{y \in D, y \neq \Delta_i(x)} \left(\frac{\pi_u(\Delta_i(x), y)}{\pi_u(x, y)} - 1 \right) \pi_u(x, y) \sum_{i=1}^N \phi(y^i) (f(\Delta_i(y)) - f(y)) \right)^2 du.$$

Now we will use the Cauchy-Schwarz inequality in the first sum. We will then obtain the following

$$(2.10) \quad \mathbf{III}_2 \leq t_0 \int_0^{t_0} \left[\sum_{y \in D, y \neq \Delta_i(x)} \pi_u(x, y) \left(\frac{\pi_u(\Delta_i(x), y)}{\pi_u(x, y)} - 1 \right)^2 \right] * \left[\sum_{y \in D, y \neq \Delta_i(x)} \pi_u(x, y) \left(\sum_{i=1}^N \phi(y^i) (f(\Delta_i(y)) - f(y)) \right)^2 \right] du.$$

The first term on the last product can be upper bounded from Lemma 2.1

$$(2.11) \quad \sum_{y \in D, y \neq \Delta_i(x)} \pi_u(x, y) \left(\frac{\pi_u(\Delta_i(x), y)}{\pi_u(x, y)} - 1 \right)^2 \leq \sum_{y \in D, y \neq \Delta_i(x)} \left(\frac{\pi_u^2(\Delta_i(x), y)}{\pi_u(x, y)} + \pi_u(x, y) \right) \leq (C_1 + 1).$$

While for the second term involved in the product of (2.10) we can write

$$\begin{aligned}
& \sum_{y \in D, y \neq \Delta_i(x)} \pi_u(x, y) \left(\sum_{i=1}^N \phi(y^i) (f(\Delta_i(y)) - f(y)) \right)^2 \leq \\
& \leq \sum_{y \in D, y \neq \Delta_i(x)} \pi_u(x, y) \left(\sum_{i=1}^N \phi(y^i) \right) \sum_{i=1}^N \phi(y^i) (f(\Delta_i(y)) - f(y))^2 \leq \\
& \leq M \sum_y \pi_u(x, y) \Gamma(f, f)(y) = \\
& = M \mathbb{E}^x \Gamma(f, f)(x_u)
\end{aligned}$$

where for the first bound we made use once more of the Holder inequality, after we divided with the appropriate normalisation constant $\sum_{i=1}^N \phi(x_u^i)$. If we put the last bound together with (2.11) into (2.10), we obtain

$$(2.12) \quad \mathbf{III}_2 \leq t_0 M (C_1 + 1) \int_0^{t_0} \mathbb{E}^x \Gamma(f, f)(x_u) du.$$

We now calculate the first summand of (2.9). Notice that in this case we cannot use the analogue bound from Lemma 2.1, that is $\frac{\pi_i^2(\Delta_i(x), \Delta_i(x))}{\pi_i(x, \Delta_i(x))} \leq \frac{C_2}{t}$, as we did for \mathbf{III}_2 , since that will lead to a final upper bound $\mathbf{III}_1 \leq t_0 M (C_1 + 1) \int_0^{t_0} \frac{1}{u} \mathbb{E}^x \Gamma(f, f)(x_u) du$ which may diverge. Instead, we will bound the \mathbf{III}_1 by the carré du champ of the function after the first jump. We can write

$$\begin{aligned}
\mathbf{III}_1 & \leq 4 \left(\int_0^{t_0} \left(\sum_{i=1}^N \phi(\Delta_i(x)^i) |f(\Delta_i(\Delta_i(x))) - f(\Delta_i(x))| \right) du \right)^2 \\
& \leq 4t_0^2 \left(\sum_{i=1}^N \phi(\Delta_i(x)^i) \right)^2 \left(\sum_{i=1}^N \frac{\phi(\Delta_i(x)^i)}{\sum_{i=1}^N \phi(\Delta_i(x)^i)} |f(\Delta_i(\Delta_i(x))) - f(\Delta_i(x))| \right)^2
\end{aligned}$$

where above we divided with the normalisation constant $\sum_{i=1}^N \phi(\Delta_i(x)^i)$, since $\phi(x) \geq \delta$. We can now apply the Holder inequality on the sum, so that

$$\mathbf{III}_1 \leq 4t_0^2 M \left(\sum_{i=1}^N \phi(\Delta_i(x)^i) (f(\Delta_i(\Delta_i(x))) - f(\Delta_i(x)))^2 \right) = 4t_0^2 M \Gamma(f, f)(\Delta_i(x)).$$

If we combine this together with (2.12) and (2.9) we get the following bound for the second term of (2.8)

$$\mathbf{II}_2 \leq 8t_0^2 M \Gamma(f, f)(\Delta_i(x)) + 4(C_1 + 1)t_0 M \int_0^{t_0} \mathbb{E}^x \Gamma(f, f)(x_u) du.$$

The last one together with the bound shown in Lemma 2.2 for the first term \mathbf{II}_1 of (2.8) gives

$$\begin{aligned} \mathbf{I}_1 &\leq t_0^2 16M\Gamma(f, f)(\Delta_i(x)) + t_0 8M(C_1 + 1) \int_0^{t_0} \mathbb{E}^x \Gamma(f, f)(x_u) du + \\ &\quad + 2(t-s)(1+C_1)M \int_{t_0}^{t-s} \mathbb{E}^x \Gamma((f, f))(x_u) du \leq \\ &\leq t_0^2 16M\Gamma(f, f)(\Delta_i(x)) + 2M(C_1 + 1)(4t_0 + (t-s)) \int_0^{t-s} \mathbb{E}^x \Gamma((f, f))(x_u) du. \end{aligned}$$

since the carré du champ is non negative, as shown below

$$\Gamma(f, f) = \frac{1}{2}(\mathcal{L}(f^2) - 2f\mathcal{L}f) = \lim_{t \downarrow 0} \frac{1}{2t} P_t f^2 - (P_t f)^2 \geq 0$$

by Cauchy-Swartz inequality. \square

We have obtained all the technical results that we need in order to show the Poincaré inequality for the semigroup P_t for general initial configurations.

2.2. proof of Theorem 1.1.

Denote $P_t f(x) = \mathbb{E}^x f(x_t)$. Then

$$(2.13) \quad P_t f^2(x) - (P_t f(x))^2 = \int_0^t \frac{d}{ds} P_s (P_{t-s} f)^2(x) ds = \int_0^t P_s \Gamma(P_{t-s} f, P_{t-s} f)(x) ds$$

since $\frac{d}{ds} P_s = \mathcal{L}P_s = P_s \mathcal{L}$.

We can write

$$(2.14) \quad \Gamma(P_{t-s} f, P_{t-s} f)(x) = \sum_{i=1}^N \phi(x^i) (\mathbb{E}^{\Delta_i(x)} f(x_{t-s}) - \mathbb{E}^x f(x_{t-s}))^2.$$

If we could use the translation property $\mathbb{E}^{x+y} f(z) = \mathbb{E}^x f(z+y)$ used for instance in proving Poincaré and modified log-Sobolev inequalities in [24] and [1], then we could bound relatively easy the carré du champ of the expectation of the functions by the carré du champ of the functions themselves, as demonstrated below

$$\begin{aligned} \sum_{i=1}^N \phi(x^i) (\mathbb{E}^{\Delta_i(x)} f(x_{t-s}) - \mathbb{E}^x f(x_{t-s}))^2 &= \sum_{i=1}^N \phi(x^i) (\mathbb{E}^x f(\Delta_i(x_{t-s})) - \mathbb{E}^x f(x_{t-s}))^2 \\ &\leq P_{t-s} \Gamma(f, f)(x) \end{aligned}$$

The inequality $\Gamma(P_t f, P_t f) \leq P_t \Gamma(f, f)$ for $t > 0$ relates directly with the Γ_2 criterion (see [3] and [4]) which states that if $\Gamma_2(f) := \frac{1}{2} \mathcal{L}(\Gamma(f, f)) - 2\Gamma(f, \mathcal{L}f) \geq 0$

then the Poincaré inequality is true, since

$$\begin{aligned} \frac{d}{ds}(P_s\Gamma(P_{t-s}f, P_{t-s}f)) &= \frac{1}{2}P_s(\mathcal{L}\Gamma(P_{t-s}f, P_{t-s}f)) - 2\Gamma(P_{t-s}f, \mathcal{L}P_{t-s}f) = \\ &= P_s(\Gamma_2(P_{t-s}f)) \geq 0 \end{aligned}$$

implies $\Gamma(P_t f, P_t f) \leq P_t \Gamma(f, f)$ (see also [1]).

Unfortunately this is not the case with our PJMP where the degeneracy of jumps and the memoryless nature of them allows any neuron x_i to jump to zero from any position, with a probability that depends on the current configuration of the neurons. Moreover, contrary on the case of Poisson processes, our intensity also depends on the position.

In order to obtain the carré du champ of the functions we will make use of the Dynkin's formula which will allow us to bound the expectation of a function with the expectation of the infinitesimal generator of the function which is comparable to the desired carré du champ of the function.

So, from Dynkin's formula

$$\mathbb{E}^x f(x_t) = f(x) + \int_0^t \mathbb{E}^x(\mathcal{L}f(x_u))du$$

we get

$$\begin{aligned} (\mathbb{E}^{\Delta_i(x)} f(x_{t-s}) - \mathbb{E}^x f(x_{t-s}))^2 &\leq 2(f(\Delta_i(x)) - f(x))^2 + \\ &+ 2 \left(\int_0^{t-s} (\mathbb{E}^{\Delta_i(x)}(\mathcal{L}f(x_u)) - \mathbb{E}^x(\mathcal{L}f(x_u))) du \right)^2. \end{aligned}$$

In order to bound the second term above we will use the bound shown in Lemma 2.3

$$\begin{aligned} (\mathbb{E}^{\Delta_i(x)} f(x_{t-s}) - \mathbb{E}^x f(x_{t-s}))^2 &\leq 2(f(\Delta_i(x)) - f(x))^2 + 32t_0^2 M \Gamma(f, f)(\Delta_i(x)) + \\ &+ 2c(t-s) \int_0^{t-s} \mathbb{E}^x \Gamma((f, f))(x_u) du \end{aligned}$$

This together with (2.14) gives

$$\begin{aligned} \Gamma(P_{t-s}f, P_{t-s}f)(x) &\leq 2\Gamma(f, f)(x) + 32t_0^2 M \sum_{i=1}^n \phi(x^i) \Gamma(f, f)(\Delta_i(x)) + \\ &+ 2Mc(t-s) \int_0^{t-s} \mathbb{E}^x \Gamma((f, f))(x_u) du. \end{aligned}$$

Finally, plugging this in (2.13) we obtain

$$\begin{aligned} P_t f^2(x) - (P_t f(x))^2 &\leq 2 \int_0^t P_s \Gamma(f, f)(x) ds + 32t_0^2 M \int_0^t P_s \sum_{i=1}^n \phi(x^i) \Gamma(f, f)(\Delta_i(x)) ds + \\ &\quad + 2M \int_0^t c(t-s) P_s \int_0^{t-s} \mathbb{E}^x \Gamma((f, f))(x_u) du ds. \end{aligned}$$

For the second term we can bound ϕ by M . For the last term on the right hand side, since the carré du champ is non negative, we can get

$$\begin{aligned} P_s \int_0^{t-s} \mathbb{E}^x \Gamma((f, f))(x_u) du &= P_s \int_s^t P_{w-s} \Gamma((f, f))(x) dw \leq \\ &\leq P_s \int_0^t P_{w-s} \Gamma((f, f))(x) dw = \int_0^t P_w \Gamma((f, f))(x) dw. \end{aligned}$$

where above we used the property of the Markov semigroup $P_s P_{w-s} = P_w$. Since this last quantity does not depend on s , and $c(t-s) \leq c(t)$ we further get

$$\int_0^t c(t-s) P_s \int_0^{t-s} \mathbb{E}^x \Gamma((f, f))(x_u) du ds \leq c(t) t \int_0^t P_w \Gamma((f, f))(x) dw.$$

Putting everything together we finally obtain

$$\begin{aligned} P_t f^2(x) - (P_t f(x))^2 &\leq (2 + 2Mc(t)t) \int_0^t P_s \Gamma(f, f)(x) ds \\ &\quad + 32t_0^2 M^2 \sum_{i=1}^n \int_0^t P_s \Gamma(f, f)(\Delta_i(x)) ds. \end{aligned}$$

And so, the theorem follows for constants

$$\alpha(t) = 2 + 2Mtc(t) = 2 + 2Mtt_0 8M(C_1 + 1) + 4t^2(1 + C_1)M^2 \text{ and } \beta = 32t_0^2 M^2.$$

3. PROOF OF THE POINCARÉ INEQUALITIES FOR STARTING CONFIGURATION ON THE DOMAIN OF THE INVARIANT MEASURE

We start by showing that in the case where the initial configuration belongs on the domain $x \in \hat{D}$ of the invariant measure, we obtain a strong lower bound for the probabilities $\pi_t(x, y) = P_x(X_t = y)$, for time t big enough, as presented on the following lemma.

Lemma 3.1. *Assume the PJMP as described in (1.2)-(1.6). Then, for every $x \in \hat{D}$ and $y \in D_x$*

$$\pi_t(x, y) \geq \frac{1}{\theta}$$

for $t \geq t_1 = \frac{1}{\delta} + t_0$.

Proof. Since $\hat{D} \subset D$ is finite, we know that there exists a strictly positive constant η , such that $\pi(x) > \eta > 0$ for every $x \in \hat{D}$. Since, $\lim_{t \rightarrow \infty} \pi_t(x, y) = \pi(y)$ for every $x \in \hat{D}$, such that $y \in D_x$, we obtain that there exists a $\theta > 0$ such that $\pi_t(x, y) > \frac{1}{\theta}$ for every $x \in \hat{D}$, such that $y \in D_x$, since $\hat{D} \subset D$ is finite. \square

Taking under account the last result, we can obtain the first technical bound needed in the proof of the local Poincaré inequality, taking advantage of the bounds shown for times bigger than t_1 .

Lemma 3.2. *Assume the PJMP as described in (1.2)-(1.6). Then, for every $z \in \hat{D}$*

$$P_s \left(\int_0^{t-s} (\mathbb{E}^{\Delta_i(z)}(\mathcal{L}f(z_u)\mathcal{I}_{z_u \in \hat{D}}) - \mathbb{E}^z(\mathcal{L}f(z_u)\mathcal{I}_{z_u \in \hat{D}})) du \right)^2 \leq 4\theta^2 t^2 M \mathbb{E}^x(\Gamma(f, f)(x_t)\mathcal{I}_{x_t \in \hat{D}})$$

for every $t \geq t_1$.

Proof. We can compute

$$\begin{aligned} \mathbf{I}_2 &:= P_s \left(\int_0^{t-s} (\mathbb{E}^{\Delta_i(z)}(\mathcal{L}f(z_u)\mathcal{I}_{z_u \in \hat{D}}) - \mathbb{E}^z(\mathcal{L}f(z_u)\mathcal{I}_{z_u \in \hat{D}})) du \right)^2 \leq \\ &\leq 2P_s \left(\int_0^t \sum_{y \in \hat{D}} \pi_u(\Delta_i(z), y) |\mathcal{L}f(y)| du \right)^2 + 2P_s \left(\int_0^t \sum_{y \in \hat{D}} \pi_u(z, y) |\mathcal{L}f(y)| du \right)^2 \end{aligned}$$

Now we will use three times the Cauchy-Schwarz inequality, to pass the square inside the integral and the two sums. We will then obtain

$$\begin{aligned} \mathbf{I}_2 &\leq 2\theta^2 t M \sum_{\omega=z, \Delta_i(z)} \sum_{y \in \hat{D}} \int_0^t P_s \pi_u(\omega, y) \sum_{i=1}^N \phi(y^i) (f(\Delta_i(y)) - f(y))^2 du \\ &= 2\theta^2 t M \sum_{\omega=z, \Delta_i(z)} \sum_{y \in \hat{D}} \int_0^t \pi_{s+u}(\omega, y) \sum_{i=1}^N \phi(y^i) (f(\Delta_i(y)) - f(y))^2 du \end{aligned}$$

where above we used the semigroup property $P_s P_u = P_{s+u}$. Since $t \geq t_1$, we can use Lemma 3.1 to bound for every w and $y \in \hat{D}$, $\pi_{u+s}(w, y) \leq \theta \pi_t(x, y)$. We then obtain

$$\begin{aligned} \mathbf{I}_2 &\leq 4\theta^2 t^2 M \sum_{y \in D} \pi_t(x, y) \sum_{i=1}^N \phi(y^i) (f(\Delta_i(y)) - f(y))^2 \\ &= 4\theta^2 t^2 M P_t \Gamma(f, f)(x). \end{aligned}$$

\square

3.1. proof of Theorem 1.3.

We can now show the Poincaré inequality for the semigroup P_t for initial configurations inside the domain \hat{D} of the invariant measure π .

Proof. We will work as in the proof of the Poincaré inequality of Theorem 1.1 for general initial conditions. As before, we denote $P_t f(x) = \mathbb{E}^x f(x_t)$. Then

$$(3.1) \quad \begin{aligned} \mathbb{E}^x (f^2(x_t) \mathcal{I}_{x_t \in \hat{D}}) - (\mathbb{E}^x (f(x_t) \mathcal{I}_{x_t \in \hat{D}}))^2 &= \int_0^t \frac{d}{ds} P_s \left(\mathbb{E}^{x_s} (f(x_{t-s}) \mathcal{I}_{x_{t-s} \in \hat{D}}) \right)^2 (x) ds = \\ &= \int_0^t P_s \Gamma(\mathbb{E}^{x_s} (f(x_{t-s}) \mathcal{I}_{x_{t-s} \in \hat{D}}), \mathbb{E}^{x_s} (f(x_{t-s}) \mathcal{I}_{x_{t-s} \in \hat{D}}))(x) ds \end{aligned}$$

since $\frac{d}{ds} P_s = \mathcal{L} P_s = P_s \mathcal{L}$. To bound the carré du champ, as in the general case of Theorem 1.1, from (2.14) and Dynkin's formula we obtain the following

$$\begin{aligned} \Gamma(\mathbb{E}^{x_s} (f(x_{t-s}) \mathcal{I}_{x_{t-s} \in \hat{D}}), \mathbb{E}^{x_s} (f(x_{t-s}) \mathcal{I}_{x_{t-s} \in \hat{D}})) &\leq 2\Gamma(f, f)(x_s) + \\ &+ 2 \sum_{i=1}^N \phi(x_s^i) \left(\int_0^{t-s} (\mathbb{E}^{\Delta_i(x_s)} (\mathcal{L} f(x_u) \mathcal{I}_{x_u \in \hat{D}}) - \mathbb{E}^{x_s} (\mathcal{L} f(x_u) \mathcal{I}_{x_u \in \hat{D}})) du \right)^2. \end{aligned}$$

From that and (3.1) and the bound $M = \sup_{x \in D} \sum_{i=1}^N \phi(x^i)$ on ϕ , we then get

$$\begin{aligned} \mathbb{E}^x (f^2(x_t) \mathcal{I}_{x_t \in \hat{D}}) - (\mathbb{E}^x (f(x_t) \mathcal{I}_{x_t \in \hat{D}}))^2 &\leq 2 \int_0^t P_s \Gamma(f, f)(x) ds + \\ &+ 2M \sum_{i=1}^N \int_0^t P_s \left(\int_0^{t-s} (\mathbb{E}^{\Delta_i(x_s)} (\mathcal{L} f(x_u) \mathcal{I}_{x_u \in \hat{D}}) - \mathbb{E}^{x_s} (\mathcal{L} f(x_u) \mathcal{I}_{x_u \in \hat{D}})) du \right)^2 ds. \end{aligned}$$

Since $t \geq t_1$, we can use Lemma 3.2 to bound the second term on the right hand side

$$\begin{aligned} \mathbb{E}^x (f^2(x_t) \mathcal{I}_{x_t \in \hat{D}}) - (\mathbb{E}^x (f(x_t) \mathcal{I}_{x_t \in \hat{D}}))^2 &\leq 2 \int_0^t P_s \Gamma(f, f)(x) ds \\ &+ 8\theta^2 M^2 N t^3 P_t \Gamma(f, f)(x). \end{aligned}$$

□

4. PROOF OF THE POINCARÉ INEQUALITIES FOR THE INVARIANT MEASURE

In this section we prove a Poincaré inequality for the invariant measure π presented in Theorem 1.5, using methods developed in [5], [6] and [7].

Proof. At first assume $\pi(f) = 0$. We can write

$$\text{Var}_\pi(f) = \int f^2 d\pi = \int f^2 \mathcal{I}_{\hat{D}} d\pi.$$

We will follow the method from [23] used to prove Spectral Gap for finite Markov chains. Since $\pi(f) = 0$, we can write

$$\int f^2 \mathcal{I}_{\hat{D}} d\pi = \frac{1}{2} \int \int (f(x) - f(y))^2 \mathcal{I}_{x \in \hat{D}} \mathcal{I}_{y \in \hat{D}} \pi(dx) \pi(dy).$$

Consider $\delta(x, y) = \{i_1, i_2, \dots, i_{|\gamma(x,y)|}\}$ the shortest path from $x \in \hat{D}$ to $y \in \hat{D}$, where the indexes i_k stand for the neuron that spikes. Since \hat{D} is finite $\max\{x, y \in \hat{D} : |\delta(x, y)|\}$ is finite. We denote $\tilde{x}^0 = x$ and $\tilde{x}^k = \Delta_{i_k}(\Delta_{i_{k-1}}(\dots \Delta_{i_1}(x))\dots)$, for $k = 1, \dots, |\delta(x, y)|$, so that $\tilde{x}^{|\delta(x,y)|} = y$. So we can write

$$\begin{aligned} \pi(x)\pi(y)(f(x) - f(y))^2 &\leq \pi(y)\pi(x) \sum_{j=0}^{|\delta(x,y)|} (f(\Delta(\tilde{x}^j)_{i_j}) - f(\tilde{x}^j))^2 \\ &\leq \frac{\pi(y)\pi(x)}{\delta} \sum_{j=0}^{|\delta(x,y)|} \varphi(\tilde{x}_{i_j}^j) (f(\Delta(\tilde{x}^j)_{i_j}) - f(\tilde{x}^j))^2 \end{aligned}$$

where above we used that $\phi \geq \delta$. We can now form the caré du champ

$$\begin{aligned} \pi(x)\pi(y)(f(x) - f(y))^2 &\leq \frac{\pi(y)\pi(x)}{\delta} \sum_{j=0}^{|\delta(x,y)|} \sum_{i \in \hat{D}} \varphi(\tilde{x}_{i_j}^j) (f(\Delta(\tilde{x}^j)_i) - f(\tilde{x}^j))^2 \\ &\leq \frac{\pi(y)\pi(x)}{\min\{x \in \hat{D} : \pi(x)\}\delta} \sum_{j=0}^{|\delta(x,y)|} \pi((\tilde{x}^j)) \Gamma(f, f)(\tilde{x}^j). \end{aligned}$$

We then have

$$\begin{aligned} \int f^2 \mathcal{I}_D d\pi &\leq \frac{N^2}{2 \min\{x \in \hat{D} : \pi(x)\}\delta} \sum_{x \in \hat{D}} \pi(x) \Gamma(f, f)(x) \\ &= \frac{N^2}{2 \min\{x \in \hat{D} : \pi(x)\}\delta} \pi(\Gamma(f, f) \mathcal{I}_{\hat{D}}). \end{aligned}$$

Putting all together leads to

$$\text{Var}_{\pi}(f) \leq \frac{N^2}{2 \min\{x \in \hat{D} : \pi(x)\}\delta} \int \Gamma(f, f) d\pi.$$

□

ACKNOWLEDGEMENTS

The authors thank Eva Löcherbach for careful reading and valuable comments.

REFERENCES

- [1] C. Ane and M. Ledoux, *Rate of convergence for ergodic continuous Markov processes: Lyapunov versus Poincaré*. Probab. Theory Relat. Fields 116, 573-602 (2000).
- [2] R. Azaïs, J.B. Bardet, A. Genadot, N. Krell and P.A. Zitt, *Piecewise deterministic Markov process (pdmps). Recent results*. Proceedings 44, 276-290 (2014).
- [3] D. Bakry, *L'hypercontractivité et son utilisation en théorie des semigroupes*. Ecole d'Eté de Probabilités de St-Flour. Lecture Notes in Math., 1581, 1-114, Springer (1994).
- [4] D. Bakry, *On Sobolev and logarithmic Sobolev inequalities for Markov semigroups*. New trends in Stochastic Analysis, 43-75, World Scientific (1997).
- [5] D. Bakry, P. Cattiaux and A. Guillin, *Lyapunov conditions for super Poincaré inequality*. J Funct Anal 254 (3), 727-759 (2008).
- [6] D. Bakry, I. Gentil, M. Ledoux, *Analysis and geometry of Markov diffusion operators*. Grundlehren der Mathematischen Wissenschaften 348, Springer, Berlin, (2014).
- [7] P. Cattiaux, A. Guillin, F-Y. Wang and L. Wu, *Lyapunov conditions for super Poincaré inequality*. J Funct Anal 256 (6), 1821-1841 (2009).
- [8] D. Chafai, *Entropies, convexity, and functional inequalities*. J. Math. Kyoto Univ. 44 (2), 325 - 363 (2004).
- [9] J. Chevalier, *Mean-field limit of generalized Hawkes processes*. Stochastic Processes and their Applications 127 (12), 3870 - 3912 (2017).
- [10] A. Crudu, A. Debussche, A. Muller and O. Radulescu *Convergence of stochastic gene networks to hybrid piecewise deterministic processes*. The Annals of Applied Probability 22, 1822-1859, (2012).
- [11] M.H.A. Davis *Piecewise-deterministic Markov processes: a general class of nondiffusion stochastic models* J. Roy. Statist. Soc. Ser. B, 46(3) 353 - 388 (1984).
- [12] M.H.A. Davis *Markov models and optimization* Monographs on Statistics and Applied Probability, vol. 49 Chapman & Hall, London. (1993)
- [13] P. Diaconis and L. Saloff-Coste *Logarithmic Sobolev inequalities for finite Markov Chains*. The Annals of Applied Probability 6, 695-750, (1996).
- [14] A. Duarte, E. Löcherbach and G. Ost, *Stability, convergence to equilibrium and simulation of non-linear Hawkes Processes with memory kernels given by the sum of Erlang kernels*. (arxiv).
- [15] A. Duarte and G.Ost, *A model for neural activity in the absence of external stimuli* Markov Processes and Related Fields 22, (2014).
- [16] A. Galves and E. Löcherbach, *Infinite Systems of Interacting Chains with Memory of Variable Length-A Stochastic Model for Biological Neural Nets*. J Stat Phys 151, 896-921 (2013).
- [17] A.Guionnet and B.Zegarlinski, *Lectures on Logarithmic Sobolev Inequalities*, IHP Course 98, 1-134 in Seminaire de Probabilite XXVI, Lecture Notes in Mathematics 1801, Springer (2003).
- [18] N. Hansen, P. Reynaud-Bouret and V. Rivoirard *Lasso and probabilistic inequalities for multivariate point processes*. Bernoulli, 21(1) 83-143 (2015).
- [19] P. Hodara, N. Krell and E. Löcherbach, *Non-parametric estimation of the spiking rate in systems of interacting neurons*. E. Stat Inference Stoch Process, 1-16 (2016).
- [20] P. Hodara and E. Löcherbach, *Hawkes processes with variable length memory and an infinite number of components*. Adv. Appl. Probab 49, 84-107 (2017).
- [21] E. Löcherbach, *Absolute continuity of the invariant measure in piecewise deterministic Markov Processes having degenerate jumps*. Stochastic Processes and their Applications (2017).

- [22] K. Pakdaman, M. Thieulen and G. Wainrib, *Fluid limit theorems for stochastic hybrid systems with application to neuron models*. Adv. Appl. Probab 42, 761-794, (2010).
- [23] L. Saloff-Coste, *Lectures on finite Markov chains*. IHP Course 98, Ecole d' Ete de Probabilites de Saint-Flour XXVI, Lecture Notes in Math. 1665, 301-413, Springer (1996).
- [24] F-Y. Wang and C. Yuan, *Poincaré inequality on the path space of Poisson point processes*. J Theor Probab 23 (3), 824-833 (2010).