

# On graphs with smallest eigenvalue at least $-3$ and their lattices

Jack H. Koolen<sup>†§</sup>, Jae Young Yang<sup>‡</sup>, Qianqian Yang<sup>†</sup>

<sup>†</sup>School of Mathematical Sciences,  
University of Science and Technology of China,  
96 Jinzhai Road, Hefei, 230026, Anhui, PR China

<sup>§</sup>Wen-Tsun Wu Key Laboratory of CAS,  
96 Jinzhai Road, Hefei, 230026, Anhui, PR China

<sup>‡</sup>School of Mathematical Sciences,  
Anhui University,  
111 Jiulong Road, Hefei, 230039, Anhui, PR China

e-mail: koolen@ustc.edu.cn, piez@naver.com, xuanxue@mail.ustc.edu.cn

April 3, 2018

## Abstract

In this paper, we show that a connected graph with smallest eigenvalue at least  $-3$  and large enough minimal degree is 2-integrable. This result generalizes a 1977 result of Hoffman for connected graphs with smallest eigenvalue at least  $-2$ .

**Keywords:**  $s$ -integrability, lattice, root lattice, Hoffman graph, eigenvalue

**AMS classification:** 05C50, 11H99

## 1 Introduction

(For undefined notions, see next section) Graphs with smallest eigenvalue at least  $-2$  are characterized by Cameron et al. [3] in 1976. They showed that

**Theorem 1.1.** *Let  $G$  be a connected graph with smallest eigenvalue at least  $-2$ . Then, either  $G$  is a generalized line graph, or  $G$  has at most 36 vertices.*

Concerning graphs with large minimal degree, Hoffman [8] showed:

**Theorem 1.2.** *For any real number  $\lambda \in (-1 - \sqrt{2}, -2]$ , there exists a constant  $f(\lambda)$  such that if a connected graph  $G$  has smallest eigenvalue at least  $\lambda$  and minimal degree at least  $f(\lambda)$ , then  $G$  is a generalized line graph.*

In this paper, we are going to generalize Theorem 1.2 for graphs with smallest eigenvalue at least  $-3$ . In order to state our main result, we need to introduce the following terminology. Let  $G$  be a graph with smallest eigenvalue  $\lambda_{\min}$ . Define the positive semi-definite matrix  $B$  by  $A - \lfloor \lambda_{\min} \rfloor I$ , where  $A$  is the adjacency matrix of  $G$ . For a positive integer  $s$ , we say that the graph  $G$  is  $s$ -integrable if there exists a matrix  $N$  such that the equality  $B = N^T N$  holds and the matrix  $\sqrt{s}N$  has only integral entries.

Our main result is as follows.

**Theorem 1.3.** *There exists a positive integer  $K$  such that if a connected graph  $G$  has smallest eigenvalue at least  $-3$  and minimal degree at least  $K$ , then  $G$  is 2-integrable.*

The above result partially solves a problem in [1, p111].

**Remark 1.4.** (i) *Theorem 1.3 can be seen as a generalization of Theorem 1.2, as the generalized line graphs are exactly the 1-integrable graphs with smallest eigenvalue at least  $-2$ .*

(ii) *As part of the proof of Theorem 1.3, we can show that there exists a real number  $\varepsilon < -3$ , such that for any real number  $\lambda \in (\varepsilon, -3]$ , there exists a constant  $f(\lambda)$ , such that if a graph  $G$  has smallest eigenvalue at least  $\lambda$  and minimal degree at least  $f(\lambda)$ , then the smallest eigenvalue of  $G$  is at least  $-3$ .*

(iii) *In Example 2.13, we will give a family of non 1-integrable graphs with smallest eigenvalue at least  $-3$  and unbounded minimal degree. This shows that we cannot improve the result in Theorem 1.3 from 2-integrability to 1-integrability.*

(iv) *Denote by  $\overline{McL}$  the complement of the McLaughlin graph. Note that  $\overline{McL}$  is the unique strongly regular graph with parameters  $(275, 162, 105, 81)$ , as the McLaughlin graph is uniquely determined by its parameters (See [6]). In [11], Koolen et al. showed that  $\overline{McL}$  is not 2-integrable. Let  $H$  be the 3-point cone over  $\overline{McL}$ , that is,  $H$  is obtained by adding 3 new vertices to  $\overline{McL}$  and joining these three vertices to all the vertices in  $\overline{McL}$  and to each other. Note that the graph  $H$  has minimal degree 165, smallest eigenvalue  $-3$  and is not 2-integrable, as  $\overline{McL}$  is*

not 2-integrable. This shows that the constant  $K$  in Theorem 1.3 is at least 166. The graph  $H$  was found by Koolen and Munemasa and they also showed that this is a maximal graph with smallest eigenvalue at least  $-3$ , that is, there does not exist a connected graph with smallest eigenvalue at least  $-3$  which contains  $H$  as a proper induced subgraph.

- (v) There exist families of connected non 2-integrable graphs with smallest eigenvalue at least  $-3$  and unbounded number of vertices, as shown by Koolen and Munemasa. For example, let  $H'$  be the cone over  $\overline{McL}$  and join the complete graph  $K_m$  to the cone vertex of  $H'$  to obtain the connected graph  $K(m)$ . Then the connected graph  $K(m)$  with  $m + 276$  vertices has smallest eigenvalue  $-3$  and is not 2-integrable.




Figure 1: The graph  $K(m)$

Now we look at integral lattices with minimal vectors of norm 3.

Let  $\Lambda$  be an (finite-dimensional) integral lattice whose minimal vectors have norm 3, and let  $\Lambda'$  be the sublattice generated by the minimal vectors of  $\Lambda$ . We denote by  $U$  the set of the minimal vectors of  $\Lambda$ , that is,  $U = \{\mathbf{u} \in \Lambda \mid \|\mathbf{u}\|^2 = 3\}$ . Assume that  $U'$  is a maximal subset of  $U$  such that for any two distinct vectors  $\mathbf{u}_1$  and  $\mathbf{u}_2$  in  $U'$ ,  $\mathbf{u}_1 \neq -\mathbf{u}_2$ , and  $L$  is the Gram matrix of  $U'$ . Since any two distinct vectors  $\mathbf{u}_1$  and  $\mathbf{u}_2$  in  $U'$  have inner product 0, 1 or  $-1$ , we find that  $L = 3I + A$ , where  $A$  is the adjacency matrix of some signed graph  $\widehat{G}$ . This concludes that  $A$  (and  $\widehat{G}$ ) has smallest eigenvalue at least  $-3$ . Therefore, in order to understand lattices whose minimal vectors have norm 3, we need to understand signed graphs with smallest eigenvalue at least  $-3$ . In this paper, we will look at connected graphs with smallest eigenvalue at least  $-3$ . We think that graphs and signed graphs behave very similar with respect to the smallest eigenvalue.

Related work in terms of line systems with angles  $\alpha$  satisfying as  $\alpha \in \{0, \pm\frac{1}{3}\}$  has been done by Shult and Yanushka [16]. Bachoc, Nebe and Venkov [13] studied lattices with minimal vectors of norm  $t$ , where  $t$  is an odd integer.

To show our main result, we need the theory of Hoffman graphs as introduced by Woo and Neumaier [19]. In Section 2, we will give a collection

of known results on Hoffman graphs which will be used to prove our main result.

## 2 Definitions and preliminaries

### 2.1 Lattices and graphs

A *lattice*  $\Lambda$  in  $\mathbb{R}^n$  is a set of vectors in  $\mathbb{R}^n$  which is closed under addition and subtraction. We say that a lattice is generated by a set  $X$  of  $\mathbb{R}^n$  if  $\Lambda = \{\sum_{\mathbf{x} \in X} \alpha_{\mathbf{x}} \mathbf{x} \mid \alpha_{\mathbf{x}} \in \mathbb{Z} \text{ for all } \mathbf{x} \in X\}$ , and we write  $\Lambda = \langle X \rangle$ . A lattice  $\Lambda$  is called *integral* if the inner product of any two vectors in  $\Lambda$  is integral. The minimal vectors of an integral lattice are the vectors with minimal positive norm. We say an integral lattice has minimal norm  $t$  if its minimal vectors have norm  $t$ . An integral lattice is called *irreducible* if it is not a direct sum of proper sublattices. An integral lattice  $\Lambda$  is called a *root lattice* if  $\Lambda$  is generated by a set  $X$  of vectors of norm 2, i.e.  $\langle \mathbf{x}, \mathbf{x} \rangle = 2$  for all  $\mathbf{x} \in X$ . The vectors of norm 2 in a root lattice  $\Lambda$  are called the *roots* of  $\Lambda$ .

**Theorem 2.1.** ([18]) *Every irreducible root lattice is isomorphic to one of the following:*

- (i)  $A_n = \{\mathbf{x} \in \mathbb{Z}^{n+1} \mid \sum \mathbf{x}_i = 0\}$  for  $n \geq 1$ ,
- (ii)  $D_n = \{\mathbf{x} \in \mathbb{Z}^n \mid \sum \mathbf{x}_i \text{ is even}\}$  for  $n \geq 4$ ,
- (iii)  $E_8 = D_8 \cup (\mathbf{c} + D_8)$ , where  $\mathbf{c} = \frac{1}{2}(1, 1, 1, 1, 1, 1, 1, 1)$ ,
- (iv)  $E_7 = \{\mathbf{x} \in E_8 \mid \sum \mathbf{x}_i = 0\}$ ,
- (v)  $E_6 = \{\mathbf{x} \in E_7 \mid \mathbf{x}_7 + \mathbf{x}_8 = 0\}$ .

For a positive integer  $s$ , an integral lattice  $\Lambda \subset \mathbb{R}^n$  is called *s-integrable* if  $\sqrt{s}\Lambda$  can be embedded in the standard lattice  $\mathbb{Z}^k$  for some  $k \geq n$ . In other words, an integral lattice  $\Lambda$  is *s-integrable* if and only if every vector in  $\Lambda$  can be described by the form  $\frac{1}{\sqrt{s}}(x_1, \dots, x_k)$  with all  $x_i \in \mathbb{Z}$  in  $\mathbb{R}^k$  for some  $k \geq n$  simultaneously.

**Remark 2.2.** *In Theorem 2.1,  $A_n$  and  $D_n$  are 1-integrable. However,  $E_6$ ,  $E_7$ , and  $E_8$  are not 1-integrable [4, Theorem 2]. To show that they are 2-integrable, we will consider another representation [5] for them. Define 8 vectors  $\mathbf{u}^i$  for  $i = 1, \dots, 8$  as follows:*

$$\mathbf{u}^1 = \frac{1}{\sqrt{2}}(0, 1, 1, 0, 1, 0, 0, 1),$$

$$\begin{aligned}
\mathbf{u}^2 &= \frac{1}{\sqrt{2}}(0, -1, 0, 1, -1, 1, 0, 0), \\
\mathbf{u}^3 &= \frac{1}{\sqrt{2}}(0, 0, -1, 0, 1, -1, 1, 0), \\
\mathbf{u}^4 &= \frac{1}{\sqrt{2}}(1, 0, 0, -1, 0, 1, -1, 0), \\
\mathbf{u}^5 &= \frac{1}{\sqrt{2}}(-1, 1, 0, 0, -1, 0, 1, 0), \\
\mathbf{u}^6 &= \frac{1}{\sqrt{2}}(1, -1, 1, 0, 0, -1, 0, 0), \\
\mathbf{u}^7 &= \frac{1}{\sqrt{2}}(0, 1, -1, 1, 0, 0, -1, 0), \\
\mathbf{u}^8 &= \frac{1}{\sqrt{2}}(-1, -1, 0, 0, 1, 0, -1, 0).
\end{aligned}$$

Then, it is known that

$$\begin{aligned}
E_8 &\simeq \langle \mathbf{u}^i \mid i = 1, \dots, 8 \rangle, \\
E_7 &\simeq \langle \mathbf{u}^i \mid i = 2, \dots, 8 \rangle, \\
E_6 &\simeq \langle \mathbf{u}^i \mid i = 3, \dots, 8 \rangle.
\end{aligned}$$

Hence,  $E_6$ ,  $E_7$ , and  $E_8$  are 2-integrable.

Let  $G$  be a connected graph with adjacency matrix  $A(G)$  and smallest eigenvalue  $\lambda_{\min}(G)$ . For  $t = -\lfloor \lambda_{\min}(G) \rfloor$ , the matrix  $B = A + tI$  is positive semidefinite, and hence there exists a matrix  $N$  such that  $B = N^T N$ . Let  $\Lambda(G)$  be the integral lattice generated by the columns of the matrix  $N$ . Then we find that  $\Lambda(G)$  is an integral lattice generated by vectors of norm  $t$ . Note that the graph  $G$  is  $s$ -integrable if and only if the integral lattice  $\Lambda(G)$  is an  $s$ -integrable lattice.

## 2.2 Hoffman graphs

Now we give definitions and preliminaries of Hoffman graphs. For more details, see [9], [12] and [19].

**Definition 2.3.** A Hoffman graph  $\mathfrak{h}$  is a pair  $(H, \ell)$  where  $H = (V, E)$  is a graph and  $\ell : V \rightarrow \{f, s\}$  is a labeling map satisfying the following conditions:

- (i) Every vertex with label  $f$  is adjacent to at least one vertex with label  $s$ ;

(ii) *Vertices with label  $f$  are pairwise non-adjacent.*

We call a vertex with label  $s$  a *slim vertex*, and a vertex with label  $f$  a *fat vertex*. We denote by  $V_s(\mathfrak{h})$  (resp.  $V_f(\mathfrak{h})$ ) the set of slim (resp. fat) vertices of  $\mathfrak{h}$ .

For a vertex  $x$  of  $\mathfrak{h}$ , we define  $N_{\mathfrak{h}}^s(x)$  (resp.  $N_{\mathfrak{h}}^f(x)$ ) the set of slim (resp. fat) neighbors of  $x$  in  $\mathfrak{h}$ . If every slim vertex of the Hoffman graph  $\mathfrak{h}$  has a fat neighbor, then we call  $\mathfrak{h}$  *fat*.

The *slim graph* of the Hoffman graph  $\mathfrak{h}$  is the subgraph of  $H$  induced on  $V_s(\mathfrak{h})$ . Note that any graph can be considered as a Hoffman graph with only slim vertices, and vice versa. We will not distinguish between Hoffman graphs with only slim vertices and graphs.

A Hoffman graph  $\mathfrak{h}_1 = (H_1, \ell_1)$  is called an (*proper*) *induced Hoffman subgraph* of  $\mathfrak{h} = (H, \ell)$ , if  $H_1$  is an (*proper*) induced subgraph of  $H$  and  $\ell_1(x) = \ell(x)$  holds for all vertices  $x$  of  $H_1$ .

Let  $W$  be a subset of  $V_s(\mathfrak{h})$ . An *induced Hoffman subgraph of  $\mathfrak{h}$  generated by  $W$* , denoted by  $\langle W \rangle_{\mathfrak{h}}$ , is the Hoffman subgraph of  $\mathfrak{h}$  induced by  $W \cup \{f \in V_f(\mathfrak{h}) \mid f \sim w \text{ for some } w \in W\}$ .

**Definition 2.4.** *For a Hoffman graph  $\mathfrak{h} = (H, \ell)$ , there exists a matrix  $C$  such that the adjacency matrix  $A$  of  $H$  satisfies*

$$A = \begin{pmatrix} A_s & C \\ C^T & O \end{pmatrix},$$

where  $A_s$  is the adjacency matrix of the slim graph of  $\mathfrak{h}$ . The special matrix  $Sp(\mathfrak{h})$  of  $\mathfrak{h}$  is the real symmetric matrix  $Sp(\mathfrak{h}) := A_s - CC^T$ . The eigenvalues of  $\mathfrak{h}$  are the eigenvalues of  $Sp(\mathfrak{h})$ , and the smallest eigenvalue of  $\mathfrak{h}$  is denoted by  $\lambda_{\min}(\mathfrak{h})$ .

**Lemma 2.5.** *([19, Lemma 3.4]) Let  $\mathfrak{h}$  be a Hoffman graph and let  $x_i$  and  $x_j$  be two distinct slim vertices of  $\mathfrak{h}$ . The special matrix  $Sp(\mathfrak{h})$  has diagonal entries*

$$Sp(\mathfrak{h})_{x_i, x_i} = -|N_{\mathfrak{h}}^f(x_i)|$$

and off-diagonal entries

$$Sp(\mathfrak{h})_{x_i, x_j} = (A_s)_{x_i, x_j} - |N_{\mathfrak{h}}^f(x_i) \cap N_{\mathfrak{h}}^f(x_j)|.$$

For the smallest eigenvalues of Hoffman graphs and their induced Hoffman subgraphs, Woo and Neumaier showed the following inequality.

**Lemma 2.6.** *([19, Corollary 3.3]) If  $\mathfrak{h}_1$  is an induced Hoffman subgraph of a Hoffman graph  $\mathfrak{h}$ , then  $\lambda_{\min}(\mathfrak{h}_1) \geq \lambda_{\min}(\mathfrak{h})$  holds.*

### 2.2.1 Two canonical fat Hoffman graphs

Given a graph  $H$ , there are two canonical fat Hoffman graphs  $\mathfrak{p}(H)$  and  $\mathfrak{q}(H)$  with  $H$  as their slim graphs. The fat Hoffman graph  $\mathfrak{p}(H)$  has  $H$  as its slim graph and to each slim vertex, a fat vertex is attached such that any two distinct slim vertices do not have a common fat neighbor. The fat Hoffman graph  $\mathfrak{q}(H)$  has  $H$  as its slim graph and one fat vertex that is attached to each slim vertex. Note that  $Sp(\mathfrak{p}(H)) = A(H) - I$  and  $Sp(\mathfrak{q}(H)) = -A(\overline{H}) - I$ , where  $A(H)$  and  $A(\overline{H})$  are the adjacency matrices of  $H$  and its complement  $\overline{H}$ , respectively. Then we have

**Lemma 2.7.** *Let  $H$  be a graph with  $\overline{H}$  as its complement. Denote by  $\lambda_{\min}(H)$  and  $\lambda_{\max}(\overline{H})$  the smallest eigenvalue of  $H$  and the largest eigenvalue of  $\overline{H}$ , respectively. Then  $\lambda_{\min}(\mathfrak{p}(H)) = -1 + \lambda_{\min}(H)$  and  $\lambda_{\min}(\mathfrak{q}(H)) = -1 - \lambda_{\max}(\overline{H})$ .*

*Proof.* It follows immediately from their special matrices.  $\square$

### 2.3 The Hoffman-Ostrowski theorem

In this subsection, we introduce a result of Hoffman and Ostrowski. In order to state the result, we need to introduce the following notations. Suppose  $\mathfrak{h}$  is a Hoffman graph and  $\{f_1, \dots, f_r\}$  is a subset of  $V_f(\mathfrak{h})$ . Let  $\mathfrak{g}^{n_1, \dots, n_r}(\mathfrak{h})$  be the Hoffman graph obtained from  $\mathfrak{h}$  by replacing the fat vertex  $f_i$  by a slim  $n_i$ -clique  $K^{f_i}$ , and joining all the neighbors of  $f_i$  (in  $\mathfrak{h}$ ) with all the vertices of  $K^{f_i}$  for all  $i$ . We will write  $G(\mathfrak{h}, n)$  for the graph  $\mathfrak{g}^{n_1, \dots, n_r}(\mathfrak{h})$ , when  $V_f(\mathfrak{h}) = \{f_1, f_2, \dots, f_r\}$  and  $n_1 = n_2 = \dots = n_r = n$ . With the above notations, we can now state the result of Hoffman and Ostrowski. For a proof of it, see [9, Theorem 2.14]. We will give a closely related result in the next section.

**Theorem 2.8.** *Suppose  $\mathfrak{h}$  is a Hoffman graph with fat vertices  $f_1, f_2, \dots, f_r$ . Then*

$$\lambda_{\min}(\mathfrak{g}^{n_1, \dots, n_r}(\mathfrak{h})) \geq \lambda_{\min}(\mathfrak{h}),$$

and

$$\lim_{n_1, \dots, n_r \rightarrow \infty} \lambda_{\min}(\mathfrak{g}^{n_1, \dots, n_r}(\mathfrak{h})) = \lambda_{\min}(\mathfrak{h}).$$

### 2.4 Lattices and Hoffman graphs

Let  $\mathfrak{h} = (H, \ell)$  be a Hoffman graph with smallest eigenvalue  $\lambda_{\min}(\mathfrak{h})$ , and let  $t$  be a real number satisfying  $\lambda_{\min}(\mathfrak{h}) \geq -t$ . Then the symmetric matrix  $Sp(\mathfrak{h}) + tI$  is positive semidefinite. Therefore, there exists a real matrix

$N$  such that  $Sp(\mathfrak{h}) + tI = N^T N$  holds. We denote by  $\Lambda^{red}(\mathfrak{h}, t)$  the lattice generated by the columns of  $N$ . Note that the isomorphism class of  $\Lambda^{red}(\mathfrak{h}, t)$  depends only on  $t$  and  $Sp(\mathfrak{h})$ , and is independent of  $N$ .

For a real number  $p$ , we define  $L(p)$  to be a diagonal matrix indexed by the vertex set  $V_s(\mathfrak{h}) \cup V_f(\mathfrak{h})$  such that  $L(p)_{x,x} = p$  if  $x \in V_s(\mathfrak{h})$  and  $L(p)_{x,x} = 1$  if  $x \in V_f(\mathfrak{h})$ . Note that the matrix  $A(H) + L(t)$  is also positive semidefinite (see [9, Theorem 2.8]), where  $A(H)$  is the adjacency matrix of  $H$ . Hence, there exists a real matrix  $\tilde{N}$  such that  $A(H) + L(t) = \tilde{N}^T \tilde{N}$  holds. We denote by  $\Lambda(\mathfrak{h}, t)$  the lattice generated by the columns of  $\tilde{N}$ . Note that the isomorphism class of  $\Lambda(\mathfrak{h}, t)$  depends only on  $t$  and  $\mathfrak{h}$ , and is independent of  $\tilde{N}$ .

It is fairly easy to see that the matrices  $N$  and  $\tilde{N}$  of above are closely related. Using this relation, we obtain the following lemma. For the convenience of the readers we give a proof.

**Lemma 2.9.** *Let  $\mathfrak{h} = (H, \ell)$  be a Hoffman graph with smallest eigenvalue  $\lambda_{\min}(\mathfrak{h})$ , and let  $t$  be an integer satisfying  $\lambda_{\min}(\mathfrak{h}) \geq -t$ . The integral lattice  $\Lambda^{red}(\mathfrak{h}, t)$  is  $s$ -integrable if and only if the integral lattice  $\Lambda(\mathfrak{h}, t)$  is  $s$ -integrable.*

*Proof.* Suppose that the lattice  $\Lambda^{red}(\mathfrak{h}, t)$  is  $s$ -integrable. It implies that there exists a matrix  $N$  such that  $Sp(\mathfrak{h}) + tI = N^T N$ , where each entry of  $N$  is in  $\frac{1}{\sqrt{s}}\mathbb{Z}$ . We denote by  $N'$  the fat-slim vertex incidence matrix with respect to

the set of fat vertices of  $\mathfrak{h}$ , that is,  $(N')_{f,x} = \begin{cases} 1 & \text{if } x \text{ is adjacent to } f, \\ 0 & \text{otherwise,} \end{cases}$  for

$x \in V_s(\mathfrak{h})$  and  $f \in V_f(\mathfrak{h})$ . The matrix  $\tilde{N} := \begin{pmatrix} N & O \\ N' & I \end{pmatrix}$  satisfies the equation  $\tilde{N}^T \tilde{N} = A(H) + L(t)$  and this shows that the lattice  $\Lambda(\mathfrak{h}, t)$  is  $s$ -integrable.

Conversely, suppose that the lattice  $\Lambda(\mathfrak{h}, t)$  is  $s$ -integrable. Then there exists a matrix  $\tilde{N}$  such that  $A(H) + L(t) = \tilde{N}^T \tilde{N}$  holds, where each entry of  $\tilde{N}$  is in  $\frac{1}{\sqrt{s}}\mathbb{Z}$ . The matrix  $N$  satisfying  $N^T N = Sp(\mathfrak{h}) + tI$  can be obtained, if for any slim vertex  $x$  of  $\mathfrak{h}$ , we define the column  $N_x$  of  $N$  indexed by  $x$  as follows:

$$N_x := \tilde{N}_x - \sum_{\substack{f \sim x \\ f \in V_f(\mathfrak{h})}} \tilde{N}_f,$$

where  $\tilde{N}_x$  and  $\tilde{N}_f$  are the columns of  $\tilde{N}$  indexed by vertices  $x$  and  $f$ , respectively. Note that all entries of  $N$  are in  $\frac{1}{\sqrt{s}}\mathbb{Z}$  and this shows that the lattice  $\Lambda^{red}(\mathfrak{h}, t)$  is  $s$ -integrable.  $\square$

The following corollary immediately follows from Lemma 2.9.



**Corollary 2.10.** *Let  $\mathfrak{h} = (H, \ell)$  be a Hoffman graph with smallest eigenvalue  $\lambda_{\min}(\mathfrak{h})$ , and let  $H'$  be the slim graph of  $\mathfrak{h}$  with smallest eigenvalue  $\lambda_{\min}(H')$ . Suppose  $\lfloor \lambda_{\min}(\mathfrak{h}) \rfloor = \lfloor \lambda_{\min}(H') \rfloor$  holds. Then the graph  $H'$  is  $s$ -integrable if the integral lattice  $\Lambda(\mathfrak{h}, -\lfloor \lambda_{\min}(\mathfrak{h}) \rfloor)$  is  $s$ -integrable.*

Another result for  $s$ -integrability of Hoffman graphs is given as follows.

**Theorem 2.11.** *Let  $\mathfrak{h}$  be a fat Hoffman graph with smallest eigenvalue  $\lambda_{\min}(\mathfrak{h})$  satisfying the following:*

$$-3 \leq \lambda_{\min}(\mathfrak{h}) < -2.$$

*Then the integral lattice  $\Lambda^{\text{red}}(\mathfrak{h}, 3)$  is 1-integrable if and only if there exists a positive constant  $D$  such that the graph  $G(\mathfrak{h}, n)$  is 1-integrable for all  $n \geq D$ .*

*Proof.* From Theorem 2.8 we have that both  $\lambda_{\min}(G(\mathfrak{h}, n)) \geq \lambda_{\min}(\mathfrak{h})$  and  $\lim_{n \rightarrow \infty} \lambda_{\min}(G(\mathfrak{h}, n)) = \lambda_{\min}(\mathfrak{h})$  hold. It follows that there exists a constant  $D'$  such that  $-3 \leq \lambda_{\min}(G(\mathfrak{h}, n)) < -2$  holds for all  $n \geq D'$ .

Assume  $V_s(\mathfrak{h}) = \{x_1, \dots, x_m\}$  and  $V(K^f) = \{y_1^f, \dots, y_n^f\}$  for any  $f \in V_f(\mathfrak{h})$ , where  $K^f$  is the  $n$ -clique as defined in Subsection 2.3.

If the integral lattice  $\Lambda^{\text{red}}(\mathfrak{h}, 3)$  is 1-integrable, then there exists an integral matrix  $N'$  such that  $Sp(\mathfrak{h}) + 3I = (N')^T N'$  holds. Let  $D := D'$  be the constant. In order to show that  $G(\mathfrak{h}, n)$  is 1-integrable for all  $n \geq D$ , it is sufficient to show that there exists an integral matrix  $N$  such that  $A(G(\mathfrak{h}, n)) + 3I = N^T N$  holds for all  $n \geq D$ . For any vertices  $x_i$  and  $y_j^f$  of  $G(\mathfrak{h}, n)$ , we define the columns  $N_{x_i}$  and  $N_{y_j^f}$  of  $N$  indexed by the vertices  $x_i$  and  $y_j^f$  as follows:

$$N_{x_i} := N'_{x_i} + \sum_{\substack{f \sim x_i \\ f \in V_f(\mathfrak{h})}} \mathbf{e}_f, \quad N_{y_j^f} := \mathbf{e}_f + \mathbf{e}_{y_{j,1}^f} + \mathbf{e}_{y_{j,2}^f},$$

where  $\{\mathbf{e}_f, \mathbf{e}_{y_{j,1}^f}, \mathbf{e}_{y_{j,2}^f} \mid f \in V_f(\mathfrak{h}), j = 1, \dots, n\}$  is a set of orthonormal integral vectors which are orthogonal to the matrix  $N'$ . It is easy to check that the matrix  $N$  satisfies  $A(G(\mathfrak{h}, n)) + 3I = N^T N$ .

Conversely, assume that the graph  $G(\mathfrak{h}, n)$  is 1-integrable for all  $n \geq D$ . Let  $n' := \max\{D, D', 20\}$  be a positive integer. Then there exists an integral matrix  $N$  such that  $A(G(\mathfrak{h}, n')) + 3I = N^T N$  holds. For any vertex  $x$  of  $G(\mathfrak{h}, n')$ , we denote by  $N_x$  the column of  $N$  indexed by  $x$ , and for the column vector  $N_x$ , we denote by  $\underline{N}_x$  the support of  $N_x$ , that is, the set  $\{j \mid (N_x)_j \neq 0\}$ . Note that the size of  $\underline{N}_x$  is 3. Now for any vertex  $x_i$  of  $G(\mathfrak{h}, n')$  and any fat vertex  $f$  adjacent to  $x_i$  in  $\mathfrak{h}$ , we have  $\underline{N}_{x_i} \cap \underline{N}_{y_j^f} \neq \emptyset$  for  $j = 1, \dots, n'$ , as

$x_i$  and  $y_j^f$  are adjacent in  $G(\mathfrak{h}, n')$ . Moreover, as  $n \geq 20$ , we can find four vertices  $y_{j_1}^f, y_{j_2}^f, y_{j_3}^f$  and  $y_{j_4}^f$  satisfying the following conditions:

- there exists an integer  $r(f)$  such that  $r(f) = \cap_{p=1}^4 \underline{N}_{y_{j_p}^f} \cap \underline{N}_{x_i}$ ;
- $(N_{x_i})_{r(f)} = (N_{y_{j_p}^f})_{r(f)} \in \{1, -1\}$  for all  $1 \leq p \leq 4$ ;
- $|\underline{N}_{y_{j_p}^f} \cap \underline{N}_{y_{j_q}^f}| = 1$ , for all  $1 \leq p < q \leq 4$ .

Then, for any  $t \neq i$ , if  $x_t$  is adjacent to  $f$  in  $\mathfrak{h}$ , then we have  $r(f) \in \underline{N}_{x_t}$  and  $(N_{x_t})_{r(f)} = (N_{x_i})_{r(f)}$ , as  $x_t$  is adjacent to all of  $y_{j_1}^f, y_{j_2}^f, y_{j_3}^f$  and  $y_{j_4}^f$  and the size of  $\underline{N}_{x_t}$  is 3. Moreover, if  $x_t$  is not adjacent to  $f$  in  $\mathfrak{h}$ , then we have  $r(f) \notin \underline{N}_{x_t}$ , as  $x_t$  is adjacent to none of  $y_{j_1}^f, y_{j_2}^f, y_{j_3}^f$  and  $y_{j_4}^f$  and the size of  $\underline{N}_{x_t}$  is 3. Now considering the submatrix of  $N$  indexed by the vertices of  $V_s(\mathfrak{h})$ , we change its  $(r(f), x_i)^{\text{th}}$  position from non-zero to zero for  $i = 1, \dots, m$  and obtain a new matrix  $N'$ . It is easy to check that the integral matrix  $N'$  satisfies  $Sp(\mathfrak{h}) + 3I = (N')^T N'$ . This shows that  $\Lambda^{\text{red}}(\mathfrak{h}, 3)$  is 1-integrable.  $\square$

Using the fat Hoffman graph  $\mathfrak{p}(H)$ , as defined in Subsection 2.2.1, we obtain the following corollary.

**Corollary 2.12.** *Assume that  $H$  is a non-complete graph with smallest eigenvalue at least  $-2$ . Then the graph  $H$  is 1-integrable if and only if there exists a constant  $D$  such that the graph  $G(\mathfrak{p}(H), n)$  is 1-integrable for all  $n \geq D$ .*

*Proof.* It follows directly from Theorem 2.11 and Lemma 2.7.  $\square$

**Example 2.13.** *Note that the lattice  $\Lambda(\tilde{E}_6)$  is isomorphic to the lattice  $E_6$ , and hence the graph  $\tilde{E}_6$  is not 1-integrable. Corollary 2.12 tells us that there exists a positive integer  $D$  such that the graph  $G(\mathfrak{p}(\tilde{E}_6), n)$  with smallest eigenvalue at least  $-3$  is not 1-integrable when  $n \geq D$ . As the minimal degree of  $G(\mathfrak{p}(\tilde{E}_6), n)$  is equal to  $n$ , we obtain an infinite family of graphs with smallest eigenvalue at least  $-3$  and unbounded minimal degree.*

## 2.5 Associated Hoffman graphs

In this subsection, we summarize some facts about *associated Hoffman graphs* and *quasi-cliques*, which provide some connection between Hoffman graphs and graphs. For more details, we refer to [10] and [12].

Let  $m$  be a positive integer and let  $G$  be a graph that does not contain  $\tilde{K}_{2m}$  as an induced subgraph, where  $\tilde{K}_{2m}$  is the graph on  $2m + 1$  vertices

consisting of a complete graph  $K_{2m}$  and a vertex  $\infty$  which is adjacent to exactly  $m$  vertices of the  $K_{2m}$ . Let  $n \geq (m+1)^2$  be a positive integer. Let  $\mathcal{C}(n) := \{C \mid C \text{ is a maximal clique of } G \text{ with at least } n \text{ vertices}\}$ . Define the relation  $\equiv_n^m$  on  $\mathcal{C}(n)$  by  $C_1 \equiv_n^m C_2$  if each vertex  $x \in C_1$  has at most  $m-1$  non-neighbors in  $C_2$  and each vertex  $y \in C_2$  has at most  $m-1$  non-neighbors in  $C_1$ . Note that  $\equiv_n^m$  is an equivalence relation.

Let  $[C]_n^m$  denote the equivalence class of  $\mathcal{C}(n)$  of  $G$  under the equivalence relation  $\equiv_n^m$  containing the maximal clique  $C$  of  $\mathcal{C}(n)$ . We define the quasi-clique  $Q([C]_n^m)$  of  $C$  with respect to the pair  $(m, n)$ , as the induced subgraph of  $G$  on the set  $\{x \in V(G) \mid x \text{ has at most } m-1 \text{ non-neighbors in } C\}$ .

Let  $[C_1]_n^m, \dots, [C_r]_n^m$  be the equivalence classes of maximal cliques under  $\equiv_n^m$ . The *associated Hoffman graph*  $\mathfrak{g} = \mathfrak{g}(G, m, n)$  is the Hoffman graph satisfying the following conditions:

- (i)  $V_s(\mathfrak{g}) = V(G)$ ,  $V_f(\mathfrak{g}) = \{f_1, f_2, \dots, f_r\}$ ;
- (ii) The slim graph of  $\mathfrak{g}$  equals  $G$ ;
- (iii) For each  $i$ , the fat vertex  $f_i$  is adjacent to exactly all the vertices of  $Q([C_i]_n^m)$  for  $i = 1, 2, \dots, r$ .

The following result, which is a crucial tool in this paper, is shown in [10, Proposition 4.1].

**Proposition 2.14.** *Let  $G$  be a graph and let  $m \geq 2, \phi, \sigma, p \geq 1$  be integers. There exists a positive integer  $n = n(m, \phi, \sigma, p) \geq (m+1)^2$  such that for any integer  $q \geq n$  and any Hoffman graph  $\mathfrak{h}$  with at most  $\phi$  fat vertices and at most  $\sigma$  slim vertices, the graph  $G(\mathfrak{h}, p)$  is an induced subgraph of  $G$ , provided that the graph  $G$  satisfies the following conditions:*

- (i) *The graph  $G$  does not contain  $\tilde{K}_{2m}$  as an induced subgraph;*
- (ii) *Its associated Hoffman graph  $\mathfrak{g} = \mathfrak{g}(G, m, q)$  contains  $\mathfrak{h}$  as an induced Hoffman subgraph.*

### 3 A limit result

In this section, we will give a limit result on matrices. As a first step, we need the following two inequalities which are direct consequences of the interlacing theorem and the Courant-Weyl inequalities. We denote, for a symmetric real matrix  $M$ , its smallest eigenvalue by  $\lambda_{\min}(M)$ .

**Lemma 3.1.** (i) ([7, Theorem 9.1.1]) *If  $M$  is a real symmetric matrix and  $M'$  a principal submatrix of  $M$ , then  $\lambda_{\min}(M') \geq \lambda_{\min}(M)$ .*

- (ii) ([2, Theorem 2.8.1]) Let  $M_1$  and  $M_2$  be two real symmetric matrices. If  $M := M_1 - M_2$  is positive semidefinite, then  $\lambda_{\min}(M_1) \geq \lambda_{\min}(M_2)$ .

Now we give the limit result. The main idea comes from the proof of [9, Theorem 2.14].

**Theorem 3.2.** Let  $\mathcal{I}$  be a finite set and  $\pi = \{\mathcal{I}_1, \mathcal{I}_2\}$  a partition of  $\mathcal{I}$ .

Assume  $M := \begin{matrix} & \mathcal{I}_1 & \mathcal{I}_2 \\ \mathcal{I}_1 & M^{1,1} & M^{1,2} \\ \mathcal{I}_2 & M^{2,1} & M^{2,2} \end{matrix}$  is a block matrix indexed by  $\mathcal{I}_1 \cup \mathcal{I}_2$ , which has smallest eigenvalue  $\lambda_{\min}(M)$  at most  $-1$ . For a given non-empty set  $\mathcal{I}_3$  disjoint from  $\mathcal{I}$ , say with  $n \geq 1$  elements, define a symmetric block matrix  $\widehat{M}(n)$  indexed by  $\mathcal{I} \cup \mathcal{I}_3$  as follows:

$$\widehat{M}(n) := \begin{matrix} & \mathcal{I}_1 & \mathcal{I}_2 & \mathcal{I}_3 \\ \mathcal{I}_1 & M^{1,1} & M^{1,2} & O \\ \mathcal{I}_2 & M^{2,1} & M^{2,2} + J & J \\ \mathcal{I}_3 & O & J & J - I \end{matrix},$$

where  $O$ ,  $J$  and  $I$  are the zero matrix, the all-ones matrix and the identity matrix, respectively. Denote by  $\mu_n$  the smallest eigenvalue of  $\widehat{M}(n)$ . Then the following holds:

- (i) The sequence  $(\mu_n)_{n \geq 1}$  is non-increasing;
- (ii)  $\mu_n \geq \lambda_{\min}(M)$  and  $\lim_{n \rightarrow \infty} \mu_n = \lambda_{\min}(M)$ ;
- (iii) Let  $\mu \leq -1$  be a real number. If  $\widehat{M}(n)$  has a principal submatrix of order at most  $m$  with smallest eigenvalue less than  $\mu$ , then  $M$  also has a principal submatrix of order at most  $m$  with smallest eigenvalue less than  $\mu$ .

*Proof.* (i) If  $n_1 \geq n_2$ , then  $\widehat{M}(n_2)$  is a principal submatrix of  $\widehat{M}(n_1)$ . By Lemma 3.1 (i), we have  $\mu_{n_2} \geq \mu_{n_1}$  and hence  $(\mu_n)_{n \geq 1}$  is non-increasing.

(ii) As

$$\widehat{M}(n) = \begin{pmatrix} M^{1,1} & M^{1,2} & O \\ M^{2,1} & M^{2,2} & O \\ O & O & -I \end{pmatrix} + \begin{pmatrix} O & O & O \\ O & J & J \\ O & J & J \end{pmatrix},$$

and  $\begin{pmatrix} O & O & O \\ O & J & J \\ O & J & J \end{pmatrix}$  is positive semidefinite, by Lemma 3.1 (ii), we obtain

$$\mu_n \geq \lambda_{\min} \left( \begin{pmatrix} M^{1,1} & M^{1,2} & O \\ M^{2,1} & M^{2,2} & O \\ O & O & -I \end{pmatrix} \right) = \lambda_{\min}(M) \text{ for } n = 1, 2, \dots \quad (1)$$

Since  $(\mu_n)_{n \geq 1}$  is a non-increasing sequence bounded below by  $\lambda_{\min}(M)$ , we see that  $\lim_{n \rightarrow \infty} \mu_n$  exists.

To show  $\lambda_{\min}(M) \geq \lim_{n \rightarrow \infty} \mu_n$ , we will show that

$$\lambda_{\min}(M) \geq \mu_n + \frac{(-\mu_n - 1)|\mathcal{I}_2|}{n - \mu_n - 1} \text{ holds for all } n \geq 2.$$

As  $J - I$  is a principal submatrix of  $\widehat{M}(n)$  and has smallest eigenvalue  $-1$  if  $n \geq 2$ , we see that  $\mu_n \leq -1$  for  $n \geq 2$ . Since the matrix  $\widehat{M}(n) - \mu_n I$  is positive semidefinite, there exist matrices  $\widehat{N}, \widehat{N}^{1,1}, \widehat{N}^{1,2}$  and  $\widehat{N}^{1,3}$  such that the following holds:

- (i)  $\widehat{N} = (\widehat{N}^{1,1}, \widehat{N}^{1,2}, \widehat{N}^{1,3})$ ;
- (ii)  $\widehat{M}(n) - \mu_n I = \widehat{N}^T \widehat{N}$ ;
- (iii)  $(\widehat{N}^{1,1})^T \widehat{N}^{1,1} = M^{1,1} - \mu_n I$ ;
- (iv)  $(\widehat{N}^{1,2})^T \widehat{N}^{1,2} = M^{2,2} + J - \mu_n I$ ; and
- (v)  $(\widehat{N}^{1,3})^T \widehat{N}^{1,3} = J - (\mu_n + 1)I$ .

We will write  $\mathbf{j}$  for a (column) vector with only ones. Set  $\epsilon_1 = 1 - \sqrt{\frac{n}{n - \mu_n - 1}}$ ,  $\epsilon_2 = \sqrt{\frac{-\mu_n - 1}{n - \mu_n - 1}}$ , and  $\mathbf{u} = \frac{1}{\sqrt{n(n - \mu_n - 1)}} \widehat{N}^{1,3} \mathbf{j}$ . Then we have

$$\mathbf{u}^T \mathbf{u} = 1, \quad \mathbf{u}^T \widehat{N}^{1,1} = \mathbf{0}, \quad \mathbf{u}^T \widehat{N}^{1,2} = (1 - \epsilon_1) \mathbf{j}^T, \quad \text{and } \epsilon_2^2 = 2\epsilon_1 - \epsilon_1^2. \quad (2)$$

Let  $B$  be the  $\binom{|\mathcal{I}_2|}{2} \times |\mathcal{I}_2|$  matrix defined by  $B_{\{i,j\},k} = \delta_{i,k} - \delta_{j,k}$ , where  $i, j, k \in \mathcal{I}_2$  and  $i \neq j$ . Then

$$B^T B = |\mathcal{I}_2| I - J. \quad (3)$$

Define the matrix  $N$  as follows:

$$N = \begin{pmatrix} \widehat{N}^{1,1} & \widehat{N}^{1,2} + (\epsilon_1 - 1) \mathbf{u} \mathbf{j}^T \\ \epsilon_2 \sqrt{|\mathcal{I}_2|} I & O \\ O & \epsilon_2 B \end{pmatrix}.$$

By (2) and (3), we find that  $N^T N = M + (-\mu_n + \epsilon_2^2 |\mathcal{I}_2|) I$  holds and this completes the proof.

- (iii) Suppose  $\widehat{M}'(n)$  is a principal submatrix of  $\widehat{M}(n)$  with order  $m$  and smallest eigenvalue less than  $\mu$ , indexed by  $\mathcal{J}_1 \cup \mathcal{J}_2 \cup \mathcal{J}_3$ , where  $\mathcal{J}_i \subseteq \mathcal{I}_i$  for  $i = 1, 2, 3$ . Let  $M'$  be the principal submatrix of  $M$  indexed by  $\mathcal{J}_1 \cup \mathcal{J}_2$ . Then replacing  $M$  and  $\widehat{M}(n)$  in Inequality (1) by  $M'$  and  $\widehat{M}'(n)$ , respectively, we see that  $M'$  has smallest eigenvalue at most  $\mu$ .  $\square$

## 4 Minimal forbidden fat Hoffman graphs

Let  $\mathfrak{G}_3$  be the set of fat Hoffman graphs with smallest eigenvalue at least  $-3$ . Then, by Lemma 2.6, for any  $\mathfrak{h} \in \mathfrak{G}_3$ , every fat induced Hoffman subgraph of  $\mathfrak{h}$  is also contained in  $\mathfrak{G}_3$ . A *minimal forbidden fat Hoffman graph* for  $\mathfrak{G}_3$  is a fat Hoffman graph  $\mathfrak{f} \notin \mathfrak{G}_3$  such that every proper fat induced Hoffman subgraph of  $\mathfrak{f}$  is contained in  $\mathfrak{G}_3$ .

In this section, we will show that the number of isomorphism classes of minimal forbidden fat Hoffman graphs for  $\mathfrak{G}_3$  is finite and all these minimal forbidden fat Hoffman graphs have at most 10 slim vertices. We first need a forbidden matrix result.

### 4.1 A forbidden matrix result

In this subsection, we will show a forbidden matrix result, which will be used later to show that any minimal forbidden fat Hoffman graph for  $\mathfrak{G}_3$  has at most 10 slim vertices. We will use a result of Vijayakumar to show this result.

In [17], Vijayakumar proved that any connected signed graph with smallest eigenvalue less than  $-2$  has an induced signed subgraph with at most 10 vertices and smallest eigenvalue less than  $-2$ . As a consequence of this result, he showed the following:

**Theorem 4.1.** (*[17, Theorem 4.2]*) *Let  $M$  be a real symmetric matrix, whose diagonal entries are 0 and off-diagonal entries are integers. If the smallest eigenvalue of  $M$  is less than  $-2$ , then one of its principal submatrices of order at most 10 also has an eigenvalue less than  $-2$ .*

By Theorem 3.2, we obtain the following generalization of Theorem 4.1.

**Theorem 4.2.** *Let  $M$  be a real symmetric matrix, whose diagonal entries are 0 or  $-1$  and off-diagonal entries are integers. If the smallest eigenvalue of  $M$  is less than  $-2$ , then one of its principal submatrices of order at most 10 also has an eigenvalue less than  $-2$ .*

*Proof.* Let  $\mathcal{I}_1 := \{i \mid M_{ii} = 0\}$  and  $\mathcal{I}_2 := \{i \mid M_{ii} = -1\}$  be two disjoint sets of indices. Without loss of generality, we may assume that  $M$  is the block

matrix 
$$\begin{matrix} & \mathcal{I}_1 & \mathcal{I}_2 \\ \mathcal{I}_1 & \begin{pmatrix} M^{1,1} & M^{1,2} \\ M^{2,1} & M^{2,2} \end{pmatrix} \end{matrix}$$
. Then the matrix  $\widehat{M}(n)$ , as defined in Theorem 3.2, has 0-diagonal and integral off-diagonal entries. Hence the result follows by Theorem 4.1 and Theorem 3.2. □

## 4.2 Forbidden special matrices

In this subsection, we will give some necessary conditions on the matrices that can be the special matrices of minimal forbidden fat Hoffman graphs for  $\mathfrak{G}_3$ .

For two square matrices  $M_1$  and  $M_2$ , we say that  $M_1$  is *equivalent* to  $M_2$ , if there exists a permutation matrix  $P$  such that  $P^T M_1 P = M_2$ . A square matrix  $M$  is called *irreducible* if  $M$  is not equivalent to a block diagonal matrix.

**Proposition 4.3.** *If  $\mathfrak{f}$  is a minimal forbidden fat Hoffman graph for  $\mathfrak{G}_3$ , then the matrix  $M := Sp(\mathfrak{f}) + I$  satisfies the following properties.*

- (i)  $M$  is symmetric and irreducible;
- (ii)  $M_{ij} \in \mathbb{Z}$ ,  $M_{ij} \leq 1$ , and  $M_{ii} \leq 0$  for all  $i, j$ ;
- (iii)  $M_{ij} \geq \max\{M_{ii}, M_{jj}\} - 1$  for all  $i, j$ ;
- (iv) The smallest eigenvalue of  $M$  is less than  $-2$ ;
- (v) Every principal submatrix of  $M$  has smallest eigenvalue at least  $-2$ .

*Proof.* Note that the columns and the rows of  $Sp(\mathfrak{f})$  are indexed by the slim vertices of  $\mathfrak{f}$  and the smallest eigenvalue of  $M$  is less than  $-2$ .

Since  $Sp(\mathfrak{f})$  is symmetric,  $M$  is also symmetric. Let  $U$  be a proper subset of  $V_s(\mathfrak{f})$  and let  $\mathfrak{f}'$  be the Hoffman subgraph generated by  $U$ . Then  $Sp(\mathfrak{f}') + I$  is exactly the principal submatrix indexed by  $U$ . This shows (v) and that  $M$  is irreducible.

By Lemma 2.5, (ii) and (iii) hold. It is clear that (iv) holds. □

Let  $\widehat{\mathcal{M}}$  be the set of square matrices containing all matrices  $M \in \widehat{\mathcal{M}}$  satisfying the five properties of Proposition 4.3.

**Proposition 4.4.** *Let  $M \in \widehat{\mathcal{M}}$  be a matrix.*


- (i) *If  $M$  contains a diagonal entry at most  $-3$ , then  $M = M_1(a)$  for some integer  $a \geq 3$  where  $M_1(a) = \begin{pmatrix} -a \end{pmatrix}$ ;*
- (ii) *If  $M$  contains a diagonal entry  $-2$ , then  $M$  is equivalent to  $M_2(b_1, b_2)$  for some integers  $b_1 \in \{1, -1, -2, -3\}$ , and  $-2 \leq b_2 \leq \min\{0, b_1 + 1\}$ , where  $M_2(b_1, b_2) = \begin{pmatrix} -2 & b_1 \\ b_1 & b_2 \end{pmatrix}$ ;*
- (iii) *If all diagonal entries of  $M$  are  $0$  or  $-1$ , then  $M$  has order at most  $10$ .*

*Proof.* Statements (i) and (ii) are straightforward to check. Statement (iii) follows directly from Theorem 4.2.  $\square$

Now we are in the position to show that there are only finitely many (isomorphism classes of) minimal forbidden fat Hoffman graphs for  $\mathfrak{G}_3$ .



**Proposition 4.5.** *Let  $\mathfrak{f}$  be a minimal forbidden fat Hoffman graph for  $\mathfrak{G}_3$ . Then  $\mathfrak{f}$  satisfies exactly one of the following statements:*

- (i) *There exists a slim vertex of  $\mathfrak{f}$  which contains at least four fat neighbors. In this case,  $\mathfrak{f}$  is isomorphic to*



- (ii) *Every slim vertex of  $\mathfrak{f}$  has at most three fat neighbors. In this case,  $\mathfrak{f}$  has at most 10 slim vertices.*

*In particular, there are finitely many isomorphism classes of minimal forbidden fat Hoffman graphs for  $\mathfrak{G}_3$ .*

*Proof.* Suppose that there exists a slim vertex of  $\mathfrak{f}$  with at least four fat neighbors. Then  $Sp(\mathfrak{f}) + I$  is equivalent to  $M_1(a)$  for some  $a \geq 3$ . If  $a$  is at least 4, then  $\mathfrak{f}$  contains  as a proper induced Hoffman subgraph. By the minimality of  $\mathfrak{f}$ , we have  $a = 3$ . Thus  $Sp(\mathfrak{f}) = \begin{pmatrix} -4 \end{pmatrix}$  and  $\mathfrak{f}$  is isomorphic to .

Now, suppose that every slim vertex of  $\mathfrak{f}$  has at most three fat neighbors. Then  $Sp(\mathfrak{f}) + I$  has order at most 10 by Proposition 4.4 (ii) and (iii). This shows that  $\mathfrak{f}$  has at most 10 slim vertices.  $\square$



## 5 Proof of the main theorem

In this section, we will prove Theorem 1.3.

**Lemma 5.1.** *There exists a positive integer  $p$  such that if  $G$  is a connected graph with smallest eigenvalue  $\lambda_{\min}(G)$  at least  $-3$ , then the associated Hoffman graph  $\mathfrak{g}(G, 12, r)$  has smallest eigenvalue at least  $-3$  for any integer  $r \geq p$ .*

*Proof.* Let  $G$  be a graph with  $\lambda_{\min}(G) \geq -3$ . As  $\lambda_{\min}(\tilde{K}_{24}) < -3$ , it follows that  $G$  does not contain  $\tilde{K}_{24}$  as an induced subgraph. Now it suffices to show that  $\mathfrak{g}(G, 12, r)$  does not contain an isomorphic copy of a minimal forbidden fat Hoffman graph for  $\mathfrak{G}_3$  as an induced Hoffman subgraph. Let  $\{\mathfrak{f}_1, \mathfrak{f}_2, \dots, \mathfrak{f}_s\}$  be the set of the isomorphism classes of the minimal forbidden fat Hoffman graphs for  $\mathfrak{G}_3$ . By Theorem 2.8, we find that for all  $1 \leq i \leq s$ , there exists a positive integer  $p'_i$  such that  $\lambda_{\min}(G(\mathfrak{f}_i, p'_i)) < -3$ , since  $\lambda_{\min}(\mathfrak{f}_i) < -3$ . Let  $p_i = n(12, |V_f(\mathfrak{h}_i)|, |V_s(\mathfrak{h}_i)|, p'_i)$  and  $p = \max_{1 \leq i \leq s} p_i$ . Then for  $r \geq p$ , the Hoffman graph  $\mathfrak{g}(G, 12, r)$  does not contain an isomorphic copy of any  $\mathfrak{f}_i$  as an induced Hoffman subgraph by Proposition 2.14. This completes the proof.  $\square$

Now we recall the Ramsey theorem.

**Theorem 5.2.** ([15]) *Let  $a, b$  be two positive integers. Then there exists a minimum positive integer  $R(a, b)$  such that for any graph  $G$  on  $n \geq R(a, b)$  vertices, the graph  $G$  contains a dependent set of size  $a$  or an independent set of size  $b$ .*

The number  $R(a, b)$  in the previous theorem is called a Ramsey number.

With Lemma 5.1 and Theorem 5.2, we will show the following theorem.

**Theorem 5.3.** *There exists a positive integer  $K$ , such that if a graph  $G$  with smallest eigenvalue at least  $-3$  has minimal degree at least  $K$ , then  $G$  is the slim graph of a fat Hoffman graph with smallest eigenvalue at least  $-3$ .*

*Proof.* Choose  $n = p$ , where  $p$  is such that Lemma 5.1 holds. Let  $K$  be equal to the Ramsey number  $R(n, 10)$ . For any vertex  $x$  of  $G$ , the neighborhood of  $x$  contains a clique with order at least  $n$ , since  $G$  does not contain a 10-claw as an induced subgraph. This implies that the associated Hoffman graph  $\mathfrak{g}(G, 12, n)$  is fat. By Lemma 5.1, the fat Hoffman graph  $\mathfrak{g}(G, 12, n)$  has smallest eigenvalue at least  $-3$  and  $G$  is the slim graph of  $\mathfrak{g}(G, 12, n)$ .  $\square$

Now we are in the position to prove our main result Theorem 1.3.

*Proof of Theorem 1.3.* By using Theorem 1.1, the theorem holds for graphs with smallest eigenvalue at least  $-2$ . Now we may assume that  $-3 \leq \lambda_{\min}(G) < -2$  holds. Let  $K$  be a positive integer such that Theorem 5.3 holds. Then there exists a fat Hoffman graph  $\mathfrak{h}$  with smallest eigenvalue at least  $-3$  which has  $G$  as the slim graph. Note that  $\lfloor \lambda_{\min}(G) \rfloor = \lfloor \lambda_{\min}(\mathfrak{h}) \rfloor = -3$ . By Lemma 2.9 and Corollary 2.10, it is sufficient to show that the integral lattice  $\Lambda^{\text{red}}(\mathfrak{h}, 3)$  is 2-integrable.

Suppose  $N$  is a matrix such that  $Sp(\mathfrak{h}) + 3I = N^T N$  holds. Denote by  $N_{r_1}, N_{r_2}, \dots, N_{r_n}$  all the non-zero columns of  $N$ . Since the Hoffman graph  $\mathfrak{h}$  is fat, then the norm  $\|N_{r_i}\|^2$  of  $N_{r_i}$  is 1 or 2 for  $i = 1, \dots, n$ . This means that the lattice  $\Lambda^{\text{red}}(\mathfrak{h}, 3)$  is a direct sum of a standard lattice and irreducible root lattices. As any irreducible root lattice is 2-integrable, we find that  $\Lambda^{\text{red}}(\mathfrak{h}, 3)$  is also 2-integrable. This completes the proof.  $\square$

**Remark 5.4.** *In Remark 1.4 (ii), we claimed that there exists a real number  $\varepsilon < -3$ , such that for any real number  $\lambda \in (\varepsilon, -3]$ , there exists a constant  $f(\lambda)$ , such that if a graph  $G$  has smallest eigenvalue at least  $\lambda$  and minimal degree at least  $f(\lambda)$ , then the smallest eigenvalue of  $G$  is at least  $-3$ . The real number  $\varepsilon$  can be found as the largest number among the smallest eigenvalues of the minimal forbidden fat Hoffman graphs for  $\mathfrak{G}_3$ . As there are finitely many isomorphism classes of the minimal forbidden fat Hoffman graphs for  $\mathfrak{G}_3$ , the number  $\varepsilon$  exists.*

We finish this paper with two conjectures and one problem.

**Conjecture 5.5.** *There exists a positive integer  $\sigma$  such that any connected graph  $G$  with smallest eigenvalue at least  $-3$  is  $\sigma$ -integrable.*

**Conjecture 5.6.** *There exists a positive integer  $\sigma$  such that for any irreducible integral lattice  $\Lambda$  generated by vectors of norm 3, the lattice  $\Lambda$  is  $\sigma$ -integrable.*

We would also like to formulate the following problem: Find more maximal connected graphs with smallest eigenvalue at least  $-3$ .

## Acknowledgments

J.H. Koolen is partially supported by the National Natural Science Foundation of China (No. 11471009 and No. 11671376).

## References

- [1] A.E. Brouwer, A.M. Cohen, A. Neumaier, *Distance-Regular Graphs*, Springer-Verlag, Berlin, 1989.
- [2] A.E. Brouwer, W.H. Haemers, *Spectra of graphs*, Springer, New York, 2012.
- [3] P.J. Cameron, J.M. Goethals, J.J. Seidel, E.E. Shult, Line graphs, root systems, and elliptic geometry, *J. Algebra*, 43:305–327, 1976.
- [4] J.H. Conway, N.J.A. Sloane, Low-dimensional lattices V. Integral coordinates for integral lattices, *Proc. Roy. Soc. London Ser. A*, 426:211–232, 1989.
- [5] W. Ebeling, Lattices and codes. A course partially based on lectures by F. Hirzebruch. Advanced Lectures in Mathematics. *Friedr. Vieweg & Sohn, Braunschweig*, 1994.
- [6] J.M. Goethals, J.J. Seidel, The regular two-graph on 276 vertices, *Discrete Math.*, 12:143–158, 1975.
- [7] C.D. Godsil, G. Royle, *Algebraic Graph Theory*, Springer-Verlag, New York, 2001.
- [8] A.J. Hoffman, On graphs whose least eigenvalue exceeds  $-1 - \sqrt{2}$ , *Linear Algebra Appl.*, 16:153–165, 1977.
- [9] H.J. Jang, J.H. Koolen, A. Munemasa, T. Taniguchi, On fat Hoffman graphs with smallest eigenvalue at least  $-3$ , *Ars Math. Contemp.*, 7:105–121, 2014.
- [10] H.K. Kim, J.H. Koolen, J.Y. Yang, A structure theory for graphs with fixed smallest eigenvalue, *Linear Algebra Appl.*, 540:1–13, 2016.
- [11] J.H. Koolen, M.U. Rehman, Q. Yang, On the integrability of strongly regular graphs, preprint.
- [12] J.H. Koolen, J.Y. Yang, Q. Yang, A generalization of a theorem of Hoffman, arXiv:1612.07085.
- [13] C. Bachoc, G. Nebe, B. Venkov, Odd unimodular lattices of minimum 4, *Acta Arith.*, 101:151–158, 2002.
- [14] G. Nebe, B. Venkov, On lattices whose minimal vectors form a 6-design, *European J. Combin.*, 30:716–724, 2009.

- [15] F.P. Ramsey, On a problem of formal logic, *Proc. London Math. Soc.*, 30:264–286, 1930.
- [16] E. Shult, A. Yanushka, Near  $n$ -gons and line systems, *Geom. Dedicata*, 9:1–72, 1980.
- [17] G.R. Vijayakumar, Signed graphs represented by  $D_\infty$ , *European J. Combin.*, 8:103–112, 1987.
- [18] E. Witt, Spiegelungsgruppen und Aufzählung halbeinfacher Liescher Ringe, *Abh. Math. Sem. Hansischen Univ.*, 14:289–322, 1941.
- [19] R. Woo, A. Neumaier, On graphs whose smallest eigenvalue is at least  $-1 - \sqrt{2}$ , *Linear Algebra Appl.*, 226–228:577–591, 1995.