arXiv:1804.03314v1 [physics.optics] 10 Apr 2018

Optical solitons in media with focusing and defocusing saturable nonlinearity and a parity-time-symmetric external potential

Pengfei Li¹, Dumitru Mihalache², and Boris A. Malomed^{3,4}

¹Department of Physics, Taiyuan Normal University, Taiyuan, 030031, China

²Horia Hulubei National Institute of Physics and Nuclear Engineering, Magurele, Bucharest, RO-077125, Romania

³Department of Physical Electronics, School of Electrical Engineering,

Faculty of Engineering, Tel Aviv University, Tel Aviv 69978, Israel

⁴ ITMO University, St. Petersburg 197101, Russia

We report results for solitons in models of waveguides with focusing or defocusing saturable nonlinearity and a parity-time (\mathcal{PT})-symmetric complex-valued external potential of the Scarf-II type. The model applies to the nonlinear wave propagation in graded-index optical waveguides with balanced gain and loss. We find both fundamental and multipole solitons for both focusing and defocusing signs of the saturable nonlinearity in such \mathcal{PT} -symmetric waveguides. The dependence of the propagation constant on the soliton's power is presented for different strengths of the nonlinearity saturation, S. The stability of fundamental, dipole, tripole, and quadrupole solitons is investigated by means of the linear-stability analysis and direct numerical simulations of the corresponding (1+1)-dimensional nonlinear Schrödinger-type equation. The results show that the instability of the stationary solutions can be mitigated or completely suppressed, increasing the value of S.

PACS numbers:

I. INTRODUCTION

Dissipative systems are usually governed by nonlinear equations, in which the input of energy supports selforganization, in the form of spontaneously emerging stable non-equilibrium structures [3, 4]. In particular, on the basis of nonlinear dynamics and Prigogine's idea of self-organization, dissipative solitons were studied as a fundamental extension of solitons in conservative media [5, 6]. In this context, it is relevant to stress that \mathcal{PT} -symmetric systems are dissipative ones, as they include symmetrically placed gain and loss; nevertheless, the specific symmetry makes their properties similar to those of conservative systems, such as all-real linear spectra and continuous families of stationary nonlinear modes.

The concept of \mathcal{PT} symmetry has been drawing permanently growing interest in the past two decades since the publication of the pioneering paper by Bender and Boettcher [7]. A simple one-dimensional \mathcal{PT} -symmetric system is described by a Schrödinger-type Hamiltonian with a complex-valued potential, whose real and imaginary parts must be, respectively, even and odd functions of spatial coordinate. Although \mathcal{PT} -symmetric Hamiltonians are non-Hermitian, they may support fully real eigenvalue spectra [7–9]. It is noting that although the introduction of \mathcal{PT} -symmetric models was motivated by the quantum-mechanical setting, the concept of the \mathcal{PT} -symmetry has been extended to a plethora of physical settings in photonics [10–12], Bose-Einstein condensates [13], plasmonic waveguides and metamaterials in which losses appear due to metal absorption [14–16], superconductivity [17], and other areas.

In guided-wave optics, the propagation of light is often modeled, in the paraxial approximation, by equations of the Schrödinger type, with the refractive index of a graded-index optical waveguide represented by a real-valued effective potential. An imaginary part of the potential, if any, stands for effects of loss and gain in the underlying waveguide. When the gain and loss are exactly balanced, the corresponding Schrödinger equation includes a \mathcal{PT} symmetric Hamiltonian. \mathcal{PT} -symmetric optical systems typically arise in coupled optical waveguides with balanced gain and loss, that were firstly investigated theoretically and experimentally in the linear regime [10, 11, 18–20]. A rigorous theoretical formulation of the \mathcal{PT} -symmetry in optics beyond the paraxial approximation, based on the Maxwell's equations, was given in Ref. [21]. Furthermore, the \mathcal{PT} -symmetry concept was extended to nonlinear optical systems, leading to extensive theoretical and experimental studies. Various types of nonlinear stationary modes and their propagation dynamics have been investigated for many types of \mathcal{PT} -symmetric external potentials. In particular, bright, dark, gap, and Bragg solitons were predicted in these models [22–37], see recent reviews in Refs. [38-40]. It is relevant to mention that solitons in one-dimensional \mathcal{PT} -symmetric periodic potentials with saturable nonlinearity were recently considered in Refs. [41, 42], where the existence of gap solitons in optical lattices and superlattices was demonstrated. The existence of solitons in two-dimensional \mathcal{PT} -symmetric periodic potentials was studied too, the results showing that the solitons are stable at moderate powers [43]. Besides that, logarithmically saturable nonlinearity can also support stable solitons in \mathcal{PT} -symmetric periodic potentials [44]. In the present work, we address fundamental and higher-order families of solitons in the one-dimensional model with saturable nonlinearity and a \mathcal{PT} -symmetric potential. We demonstrate that the nonlinearity saturation helps to suppress the instability of fundamental and higher-order solitons.

The rest of the paper is organized as follows. In Sec. 2, the governing model is introduced. In Sec. 3, we show that both fundamental and multipole stationary solutions exist in the (1+1)-dimensional Schrödinger-type equation with both focusing and defocusing saturable nonlinearity. The dependence of the soliton's propagation constant on the power is also investigated. In Sec. 4, the stability of fundamental, dipole, tripole, and quadrupole solitons is systematically investigated in the framework of the linear-stability analysis. The evolution of perturbed stationary solitons is studied by means of direct numerical simulations. The work is concluded by Sec. 5.

II. THE GOVERNING MODEL

In the paraxial approximation, the optical wave propagation in a planar graded-index waveguide with saturable nonlinearity is governed by the (1+1)-dimensional nonlinear Schrödinger equation, written here in the scaled form:

$$i\frac{\partial\psi}{\partial\zeta} + \frac{\partial^2\psi}{\partial\xi^2} + U\left(\xi\right)\psi + \sigma\frac{\left|\psi\right|^2\psi}{1+S\left|\psi\right|^2} = 0,\tag{1}$$

where $\psi(\zeta, \xi)$ is the envelope of the electromagnetic field, $U(\xi) \equiv V(\xi) + iW(\xi)$ is a complex-valued potential, and $\sigma = \pm 1$ represent focusing (+) and defocusing (-) signs of the nonlinearity, respectively. Finally, the coefficient S defines the saturation of the nonlinearity [45, 46].

Stationary solutions are sought for in the usual form, $\psi(\zeta,\xi) = \phi(\xi) e^{i\beta\zeta}$, where $\phi(\xi)$ is a complex function, and β is a real propagation constant. Substitution of this in equation (1) yields

$$\frac{d^2\phi}{d\xi^2} + U(\xi)\phi(\xi) + \sigma \frac{|\phi|^2\phi}{1+S|\phi|^2} - \beta\phi(\xi) = 0.$$
(2)

In this paper, we consider the nonlinear model with the \mathcal{PT} -symmetric Scarf-II potential [47, 48] of the form

$$V\left(\xi\right) = V_0 \operatorname{sech}^2\left(\frac{\xi}{\xi_0}\right),\tag{3}$$

$$W\left(\xi\right) = W_0 \operatorname{sech}\left(\frac{\xi}{\xi_0}\right) \tanh\left(\frac{\xi}{\xi_0}\right),\tag{4}$$

where V_0 and W_0 represent the strengths of its real and imaginary parts, and ξ_0 defines the width of the \mathcal{PT} -symmetric potential.

III. THE ANALYSIS OF THE STATIONARY SOLUTIONS

We seek for solutions to equation (1) with both focusing and defocusing sign of the nonlinearity, i.e., $\sigma = +1$ and $\sigma = -1$, respectively. As an example, which adequately represents the generic case, we choose $V_0 = 8$, $W_0 = 4$, and $\xi_0 = 4$ in equations (3) and (4).

The numerical results demonstrate that there exist both fundamental and higher-order stationary solutions of equation (2). The dependence of the nonlinear propagation constants on the soliton's integral power, $P = \int_{-\infty}^{+\infty} |\phi|^2 d\xi$, is presented in figure 1, where blue and red curves correspond to the case of focusing and defocusing saturable nonlinearities, respectively. Figure 1 demonstrates that the $\beta(P)$ curves originate, at P = 0, from values corresponding to the respective modes generated by the linear Schrödinger equation with the potential (3), (4). Further, figure 1 shows too that the slope of the dependence is positive and negative, i.e., $d\beta/dP > 0$ and $d\beta/dP < 0$, respectively, for the focusing and defocusing sign of the nonlinearity. In either case, the sign complies, severally, with the Vakhitov-Kolokolov (VK) [49, 50] or anti-VK [51] criteria, which are necessary conditions for the stability of solitons created by focusing or defocusing nonlinearity of any type, including the saturable one. The profiles of the corresponding solitons, for P increasing from 0.01 to 10, are depicted in figure 2.

It is also relevant to address the dependence of the propagation constant, β , of fundamental solitons on the integral power, P, for different values of the saturation parameter S. The saturation slows the variation of β with the growth of P, for both the focusing and defocusing signs of the nonlinearity, as shown in figure 3. Naturally, when S is very small (e.g., S = 0.1) in figure 3, the $\beta(P)$ dependence is close to that obtained for the cubic nonlinearity.




FIG. 1: The propagation constants of stationary solutions versus the soliton's integral power. Thick blue solid and dashed curves correspond to the stable and unstable stationary solutions with focusing saturable nonlinearity, and thin red solid and dashed curves correspond to the stable and unstable stationary solutions with defocusing saturable nonlinearities, respectively. Panels (a), (b), (c), and (d) display the results for the fundamental, dipole, tripole, and quadrupole solitons, respectively. Here, parameters in (3) and (4) are $V_0 = 8$, $W_0 = 4$, $\xi_0 = 4$, and S = 1. In fact, results of more extensive numerical calculations demonstrate that relation $W_0 = (1/2)V_0$, implied by these values, adequately represents the generic situation.




FIG. 2: Evolution of profiles of the stationary solutions with the increase of the solitons' power. Panels $(a_{1,2})$, $(b_{1,2})$, $(c_{1,2})$, and $(d_{1,2})$ represent fundamental, dipole, tripole, and quadrupole solitons, respectively, with subscripts 1 and 2 corresponding to the focusing and defocusing sign of the saturable nonlinearity. Parameters are the same as in figure 1.

IV. THE LINEAR-STABILITY ANALYSIS AND PROPAGATION DYNAMICS OF SOLITONS

To examine the stability of the stationary solutions $\phi(\xi)$ with propagation constant β , we added small perturbations $u(\xi)$ and $v(\xi)$ to them, in the usual form,

$$\psi\left(\zeta,\xi\right) = e^{i\beta\zeta} \left[\phi\left(\xi\right) + u\left(\xi\right)e^{\delta\zeta} + v^*\left(\xi\right)e^{\delta^*\zeta}\right],\tag{5}$$




FIG. 3: The propagation constant of fundamental solitons versus their integral power, with different values of saturation parameter S, see Eqs. (3) and (4), for the self-focusing (a) and the defocusing (b) sign of the nonlinearity. Other parameters are the same as in figure 1.

where δ is the (complex) instability growth rate sought for. The substitution of equation (5) into equation (1) and subsequent linearization leads to the eigenvalue problem for δ :

$$i \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \delta \begin{pmatrix} u \\ v \end{pmatrix}, \tag{6}$$

with operators

$$L_{11} = \frac{d^2}{d\xi^2} + U(\xi) - \beta + \frac{2\sigma |\phi|^2}{1 + S |\phi|^2} - \frac{\sigma S |\phi|^4}{\left(1 + S |\phi|^2\right)^2},\tag{7}$$

$$L_{12} = \frac{2\sigma\phi^2}{1+S|\phi|^2} - \frac{\sigma S|\phi|^2 \phi^2}{\left(1+S|\phi|^2\right)^2},$$
(8)

$$L_{21} = \frac{\sigma S |\phi|^2 \phi^{*2}}{\left(1 + S |\phi|^2\right)^2} - \frac{2\sigma \phi^{*2}}{1 + S |\phi|^2},\tag{9}$$

$$L_{22} = -\left[\frac{d^2}{d\xi^2} + U^*\left(\xi\right) - \beta + \frac{2\sigma \left|\phi\right|^2}{1 + S \left|\phi\right|^2} - \frac{\sigma S \left|\phi\right|^4}{\left(1 + S \left|\phi\right|^2\right)^2}\right].$$
(10)

The stationary solution $\phi(\xi)$ is linearly unstable if there is at least one eigenvalue with $\operatorname{Re}(\delta) > 0$.




FIG. 4: Eigenvalues of the linear-stability analysis, produced by the numerical solution of equation (6) for different values of saturation parameter S, see equation (1). Panels (a_1) , (a_2) , (a_3) , and (a_4) depict the largest real part of δ for the fundamental, dipole, tripole, and quadrupole soliton, respectively, in the case of the focusing saturable nonlinearity. Panels (b_1) , (b_2) , (b_3) , and (b_4) show the same for the defocusing nonlinearity. Parameters are the same as in figure 1.

Figure 4 presents the dependence of the largest instability growth rate, $Max(\delta_R)$, on the solitons' integral power, P, as produced by numerical solution of the eigenvalue problem for the fundamental, dipole, tripole, and quadrupole solitons at different values of S and both signs of the nonlinearity, focusing and defocusing (the top and bottom rows in figure 4, respectively). The numerical results displayed in panels $4(a_1)$ and $4(a_2)$ show that, under the action of the self-focusing nonlinearity, $Max(\delta_R)$ is negligibly small for the fundamental and dipole solitons, which indicates that they are linearly stable, irrespective of the value of S. On the other hand, the tripoles and quadrupoles are stable when the solitons' power is sufficiently small, as seen in panels $4(a_3)$ and $4(a_4)$ (in fact, the quadrupoles are almost completely unstable). Further, under the action of self-defocusing nonlinearity, fundamental solitons are linearly stable, irrespective of S, as shown in figure $4(b_1)$, while panels $4(b_2)$, $4(b_3)$, and $4(b_4)$ show that the dipoles, tripoles, and quadrupoles are stable, in this case, only in tiny intervals of very small powers. A general trend, suggested by these results, is that the instability is attenuated with the increase of the saturation strength S.

The results of the linear-stability analysis are confirmed, separately in figures 5 and 6 for the focusing and defocusing nonlinearities, by direct numerical simulations of the perturbed evolution of solitons, performed in the framework of equation (1). In these figures, the results of the simulations are displayed along with (in)stability spectra of the underlying stationary states. The eigenvalue spectra of equation (6) for fundamental, dipole, tripole, and quadrupole solitons are shown on the top row in figure 5. In particular, figures $5(b_1)$ and $5(b_2)$ confirm the stable propagation of the fundamental and dipole solitons in the case of self-focusing nonlinearity, under the action of sufficiently strong random perturbations, with relative amplitudes 10%. Very small nonzero values of $\text{Re}(\delta)$, which are seen, for these solutions, in Figs. $5(a_1)$ and $5(a_2)$, but do not lead to any instability in the direct propagation, are a manifestation of a small inaccuracy of the numerical analysis. On the other hand, the tripoles and quadrupoles are definitely unstable, in accordance with the prediction of the linear-stability analysis, as seen in figures $5(b_3)$ and $5(b_4)$. It is observed that the unstable tripole is spontaneously transformed into a breather, while the quadrupole is, eventually, destroyed by the instability.

Similar results for the defocusing nonlinearity are summarized in figure 6, where, in accordance with the prediction of the linear-stability analysis, the fundamental soliton is stable, while dipole, tripole, and quadrupole are not, transforming into breathers.




FIG. 5: Eigenvalue spectra produced by the linear-stability analysis for different soliton species, and the corresponding evolution plots for the focusing nonlinearity ($\sigma = +1$) and S = 1. Panels (a_1, b_1) , (a_2, b_2) , (a_3, b_3) , and (a_4, b_4) pertain, respectively, to a fundamental soliton with power P = 5, a dipole with P = 10, a tripole with P = 6, and a quadrupole with P = 6. Other parameters are the same as in figure 1.

V. CONCLUSION

We have performed a systematic study of solitons in the models of waveguides with the self-focusing and defocusing saturable nonlinearity and \mathcal{PT} -symmetric complex-valued potential of the Scarf-II type. Both models support families of fundamental, dipole, tripole, and quadrupole solitons. Their stability has been investigated by means of the linear-stability analysis, and confirmed by direct numerical simulations. In particular, fundamental solitons are completely stable for both signs of the nonlinearity and, quite interestingly, dipoles and a part of the tripole family are stable too in the case of self-focusing nonlinearity. Other soliton families are practically completely unstable, including all higher-order solitons under the action of self-defocusing nonlinearity. A general trend is that the increase of the saturation strength partly suppresses the instability.

As an extension of the analysis, it may be interesting to perform it for more general shapes of the \mathcal{PT} -symmetric




FIG. 6: The same as in figure 5, but in the case of the defocusing nonlinearity ($\sigma = -1$). Panels (a_1, b_1) , (a_2, b_2) , (a_3, b_3) , and (a_4, b_4) pertain, respectively, to a fundamental soliton with power P = 10, and dipole, tripole, and quadrupole solitons, all with P = 6.

external potential.

VI. ACKNOWLEDGEMENT

This work was supported by Doctoral Scientific Research Foundation of Taiyuan Normal University No. 1170144, and by the Israel Foundation through grant No. 1287/17.

Prigogine I, Nicolis G. 1967 On Symmetry-Breaking Instabilities in Dissipative Systems. The Journal of Chemical Physics 46, 3542–3550. (https://doi.org/10.1063/1.1841253).

- [2] Prigogine I, Lefever R. 1968 On Symmetry-Breaking Instabilities in Dissipative Systems. II. The Journal of Chemical Physics 48, 1695–1700. (https://doi.org/10.1063/1.1668896).
- [3] Glansdorff P, Prigogine I. 1971 Thermodynamic theory of structure, stability and fluctuations. New York, NY: John Wiley and sons.
- [4] Nicolis G, Prigogine I. 1977 Self-organization in nonequilibrium systems. New York, NY: John Wiley and sons.
- [5] Akhmediev N, Ankiewicz A. 2005 Dissipative Solitons. Verlag Berlin Heidelberg: Springer.
- [6] Akhmediev N, Ankiewicz A. 2008 Dissipative solitons: from optics to biology and medicine. Verlag Berlin Heidelberg: Springer.
- [7] Bender CM, Boettcher S. 1998 Real spectra in non-Hermitian Hamiltonians having *PT* symmetry. *Phys. Rev. Lett.* 80, 5243–5246. (https://doi.org/10.1103/PhysRevLett.80.5243).
- [8] Bender CM, Brody DC, Jones HF. 2002 Complex extension of quantum mechanics. Phys. Rev. Lett. 89, 270401. (https://doi.org/10.1103/PhysRevLett.89.270401).
- [9] Bender CM, Boettcher S, Meisinger PN. 1999 *PT*-symmetric quantum mechanics. J. Math. Phys. 40, 2201–2229. (https://doi.org/10.1063/1.532860)
- [10] Ruschhaupt A, Delgado F, Muga JG. 2005 Physical realization of *PT*-symmetric potential scattering in a planar slab waveguide. J. Phys. A: Math. Gen. 38, L171–L176. (https://doi.org/10.1088/0305-4470/38/9/L03)
- [11] El-Ganainy R, Makris KG, Christodoulides DN, Musslimani ZH. 2007 Optical *PT*-symmetric structures. Opt. Lett. 32, 2632–2634. (https://doi.org/10.1364/OL.32.002632)
- [12] Klaiman, S., Gunther U, Moiseyev N. 2008 Visualization of branch points in *PT*-symmetric waveguides. *Phys. Rev. Lett.* 101, 080402. (https://doi.org/10.1103/PhysRevLett.101.080402)
- [13] Bagnato VS, Frantzeskakis DJ, Kevrekidis PG, Malomed BA, Mihalache D. 2015 Bose-Einstein condensation: twenty years after. Rom. Rep. Phys. 67, 5–50.
- [14] Alaeian H, Dionne JA. 2014 Non-Hermitian nanophotonic and plasmonic waveguides. Phys. Rev. B 89, 075136. (https://doi.org/10.1103/PhysRevB.89.075136)
- [15] Alaeian H, Dionne JA. 2014 Parity-time-symmetric plasmonic metamaterials. Phys. Rev. A 89, 033829. (https://doi.org/10.1103/PhysRevA.89.033829)
- [16] Lazarides N, Tsironis GP. 2013 Gain-driven discrete breathers in *PT*-symmetric nonlinear metamaterials. *Phys. Rev. Lett.* 110, 053901. (https://doi.org/10.1103/PhysRevLett.101.053901)
- [17] Chtchelkatchev NM, Golubov AA, Baturina TI, Vinokur VM. 2012 Stimulation of the fluctuation superconductivity by *PT* symmetry. *Phys. Rev. Lett.* **109**, 150405. (https://doi.org/10.1103/PhysRevLett.109.150405)
- [18] Guo A, Salamo GJ, Duchesne D, Morandotti R, Volatier-Ravat M, Aimez V, Siviloglou GA, Christodoulides DN. 2009 Observation of *PT* -symmetry breaking in complex optical potentials. *Phys. Rev. Lett.* **103**, 093902. (https://doi.org/10.1103/PhysRevLett.103.093902)
- [19] Rüter CE, Makris KG, El-Ganainy R, Christodoulides DN, Segev M, Kip D. 2010 Observation of parity-time symmetry in optics. Nature Physics 6, 192–195. (https://doi.org/10.1038/nphys1515)
- [20] Kottos T. 2010 Optical physics: broken symmetry makes light work. Nature Physics 6, 166–167. (https://doi.org/10.1038/nphys1612)
- [21] Huang CM, Ye FW, Kartashov YV, Malomed BA, 2014 Chen XF. *PT* symmetry in optics beyond the paraxial approximation. Opt. Lett. 39, 5443–5446. (https://doi.org/10.1364/OL.39.005443)
- [22] Musslimani ZH, Makris KG, El-Ganainy R, Christodoulides DN. 2008 Optical solitons in *PT* periodic potentials. *Phys. Rev. Lett.* **100**, 030402. (https://doi.org/10.1103/PhysRevLett.100.030402)
- [23] Abdullaev FK, Kartashov YV, Konotop VV, Zezyulin DA. 2011 Solitons in *PT*-symmetric nonlinear lattices. *Phys. Rev.* A 83, 041805(R). (https://doi.org/10.1103/PhysRevA.83.041805)
- [24] Zhu X, Wang H, Zheng L, Li H, He YJ. 2011 Gap solitons in parity-time complex periodic optical lattices with the real part of superlattices. Opt. Lett. 36, 2680–2682. (https://doi.org/10.1364/OL.36.002680)
- [25] Li HG, Shi ZW, Jiang X, Zhu X. 2011 Gray solitons in parity-time symmetric potentials. Opt. Lett. 36, 3290–3292. (https://doi.org/10.1364/OL.36.003290)
- [26] Driben R, Malomed BA. 2011 Stability of solitons in parity-time-symmetric couplers. Opt. Lett. 36, 4323-4325. (https://doi.org/10.1364/OL.36.004323)
- [27] Hu SM, Ma X, Lu D, Yang Z, Zheng Y, Hu W. 2011 Solitons supported by complex *PT*-symmetric Gaussian potentials. *Phys. Rev. A*, 84, 043818. (https://doi.org/10.1103/PhysRevA.84.043818)
- [28] Midya B, Roychoudhury R. 2013 Nonlinear localized modes in *PT*-symmetric Rosen-Morse potential wells. *Phys. Rev. A* 87, 045803. (https://doi.org/10.1103/PhysRevA.87.045803)
- [29] Nixon S, Ge L, Yang Y. 2012 Stability analysis for solitons in *PT*-symmetric optical lattices. *Phys. Rev. A* 85, 023822. (https://doi.org/10.1103/PhysRevA.85.023822)
- [30] Zezyulin DA, Konotop VV. 2012 Nonlinear modes in the harmonic *PT*-symmetric potential. *Phys. Rev. A* 85, 043840. (https://doi.org/10.1103/PhysRevA.85.043840)
- [31] Achilleos V, Kevrekidis PG, Frantzeskakis DJ, Carretero-González R. 2012 Dark solitons and vortices in *PT*-symmetric nonlinear media: From spontaneous symmetry breaking to nonlinear *PT* phase transitions. *Phys. Rev. A* 86, 013808. (https://doi.org/10.1103/PhysRevA.86.013808)
- [32] Miri MA, Aceves AB, Kottos T, Kovanis V, Christodoulides DN. 2012 Bragg solitons in nonlinear *PT*-symmetric periodic potentials. *Phys. Rev. A* 86, 033801. (https://doi.org/10.1103/PhysRevA.86.033801)
- [33] He YJ, Zhu X, Mihalache D. 2016 Dynamics of spatial solitons in parity-time-symmetric optical lattices: A selection of recent theoretical results. Rom. J. Phys. 61, 595–613.

- [34] Li PF, Liu B, Li L, Mihalache D. 2016 Nonlinear parity-time-symmetry breaking in optical waveguides with complex Gaussian-type potentials. Rom. J. Phys. 61, 577–594.
- [35] Li PF, Li L, Mihalache D. 2018 Optical solitons in \mathcal{PT} -symmetric potentials with competing cubic-quintic nonlinearity: existence, stability, and dynamics. *Rom. Rep. Phys.* **70**, 408.
- [36] Mihalache D. 2017 Multidimensional localized structures in optical and matter-wave media: A topical survey of recent literature. Rom. Rep. Phys. 69, 403.
- [37] Chen Y, Yan Z, Mihalache D, Malomed BA. 2017 Families of stable solitons and excitations in the *PT*-symmetric nonlinear Schrödinger equations with position-dependent effective masses. *Sci. Reports* 7, 1257. (https://doi.org/10.1038/s41598-017-01401-3)
- [38] Konotop VV, Yang J, Zezyulin DA. 2016 Nonlinear waves in *PT*-symmetric systems. *Rev. Mod. Phys.* 88, 035002. (https://doi.org/10.1103/RevModPhys.88.035002)
- [39] Suchkov SV, Sukhorukov AA, Huang J, Dmitriev SV, Lee C, Kivshar YS. 2016 Nonlinear switching and solitons in *PT*-symmetric photonic systems. Laser and Photonics Reviews 10, 177–213. (https://doi.org/lpor.201500227)
- [40] El-Ganainy R, Makris KG, Khajavikhan M, Musslimani ZH, Rotter S, Christodoulides DN. 2018 Non-Hermitian physics and *PT* symmetry. *Nature Physics* 14, 11–19. (https://doi.org/10.1038/nphys4323)
- [41] Cao P, Zhu X, He Y, Li HG. 2014 Gap solitons supported by parity-time-symmetric optical lattices with defocusing saturable nonlinearity. Opt. Commun. 316, 190–197. (https://doi.org/10.1016/j.optcom.2013.12.021)
- [42] Li L, Li H, Lai T. 2015 Defect solitons in parity-time symmetric superlattices with focusing saturable nonlinearity. Opt. Commun. 349, 171–179. (https://doi.org/10.1016/j.optcom.2015.03.067)
- [43] Zhan KY, Tian H, Li X, Xu XF, Jiao ZY, Jia YL. 2016 Solitons in *PT*-symmetric periodic systems with the logarithmically saturable nonlinearity. Sci. Rep. 6, 32990. (https://doi.org/10.1038/srep32990)
- [44] Zhu X, Li H. 2016 Multihump solitons in two-dimensional parity-time-symmetric optical lattices with focusing saturable nonlinearity. Eur. Phys. J. D 70, 14. (https://doi.org/10.1140/epjd/e2015-60434-7)
- [45] Kartashov YV, Vysloukh VA, Torner L. 2004 Soliton trains in photonic lattices. Opt. Express. 12, 2831–2837. (https://doi.org/10.1364/OPEX.12.002831)
- [46] Hu SM, Hu W. 2013 Defect solitons in saturable nonlinearity media with parity-time symmetric optical lattices. *Physica B* 429, 28–32. (https://doi.org/10.1016/j.physb.2013.07.023)
- [47] Scarf FL. 1958 New soluble energy band problem. Phys. Rev. 112, 1137–1140. (https://doi.org/10.1103/PhysRev.112.1137)
- [48] Grosche C. 1993 Path integral discussion of Scarf-like potentials. Nuovo Cim. 108B, 1365–1376. (https://doi.org/10.1007/BF02755190)
- [49] Vakhitov N and Kolokolov A. 1973 Stationary solutions of the wave equation in a medium with nonlinearity saturation. Radiophys. Quantum Electron. 16, 783–789. (https://doi.org/10.1007/BF01031343)
- [50] Bergé L. 1998 Wave collapse in physics: Principles and applications to light and plasma waves. Phys. Rep. 303, 259–370. (https://doi.org/10.1016/S0370-1573(97)00092-6)
- [51] Sakaguchi H, Malomed BA. 2010 Solitons in combined linear and nonlinear lattice potentials. *Phys. Rev.* A 81, 013624. (https://doi.org/10.1103/PhysRevA.81.013624)