AN EXPLICIT FORMULA FOR BERNOULLI POLYNOMIALS WITH A q PARAMETER IN TERMS OF r-WHITNEY NUMBERS

F. A. SHIHA

ABSTRACT. We define the Bernoulli polynomials with a q parameter in terms of r-Whitney numbers of the second kind. Some algebraic properties and combinatorial identities of these polynomials are given. Also, we obtain several relations between the Cauchy and Bernoulli polynomials with a q parameter in terms of r-Whitney numbers of both kinds.

AMS (2010) Subject Classification: 05A15, 05A19, 11B68, 11B73. Key Words. Bernoulli numbers and polynomials, *r*-Whitney numbers, Stirling numbers.

1. INTRODUCTION

The Bernoulli polynomials $B_n(z)$, are defined by the generating function [4]

(1.1)
$$\sum_{n=0}^{\infty} B_n(z) \frac{t^n}{n!} = \frac{t \, e^{zt}}{e^t - 1}.$$

When z = 0, $B_n = B_n(0)$ are called the Bernoulli numbers. Graham et al. [6, P. 560] represented B_n also as

(1.2)
$$B_n = \sum_{k=0}^n (-1)^k \frac{k!}{k+1} S(n,k),$$

where S(n, k) are the Stirling numbers of the second kind [4].

For all integers $n, r \ge 0$, $B_n(r)$ can be written explicitly in terms of r-Stirling numbers of the second kind $S_r(n, k)$ [7]

(1.3)
$$B_n(r) = \sum_{k=0}^n \frac{(-1)^k k!}{k+1} S_r(n+r,k+r).$$

Cenkci et al. [2] introduced the poly-Bernoulli polynomials with a q parameter $B_{n,q}^{(k)}(z)$ as a generalization of the poly-Bernoulli polynomials $B_n^{(k)}(z)$. Let q be a real number with $q \neq 0$, [2] defined the poly-Bernoulli polynomials with a q parameter by

(1.4)
$$\sum_{n=0}^{\infty} B_{n,q}^{(k)}(z) \, \frac{t^n}{n!} = \frac{q \, e^{-zt}}{1 - e^{-qt}} \, \sum_{n=1}^{\infty} \left(\frac{(1 - e^{-qt})}{q} \right)^n \, \frac{1}{n^k}.$$

If z = 0, then $B_{n,q}^{(k)}(0) = B_{n,q}^{(k)}$ are the poly-Bernoulli numbers with a q parameter. If q = 1, then $B_{n,1}^{(k)}(z) = B_n^{(k)}(z)$ are the poly-Bernoulli polynomials [5]. If q = 1, z = 0, then $B_{n,1}^{(k)}(0) = B_n^{(k)}$ are the poly-Bernoulli numbers, defined in [8]. Setting k = 1 in (1.4), and let $B_n^q(z) = B_{n,q}^{(1)}(z)$ denote the Bernoulli polynomials with q parameter, we obtain

(1.5)
$$\sum_{n=0}^{\infty} B_n^q(z) \, \frac{t^n}{n!} = \frac{q \, e^{-zt}}{1 - e^{-qt}} \, \sum_{n=1}^{\infty} \left(\frac{(1 - e^{-qt})}{q} \right)^n \, \frac{1}{n}.$$

If z = 0, then $B_n^q(0) = B_n^q$ are the Bernoulli numbers with q parameter. Note that e^{zt} is replaced by e^{-zt} (see [2, 1, 5]).

In this paper, we are interested in Bernoulli polynomials with a q parameter. New representation for these polynomials in terms of r-Whitney numbers of the second kind and many properties will be given.

For non-negative integers n and k with $0 \le k \le n$, let $w(n,k) = w_{q,r}(n,k)$ and $W(n,k) = W_{q,r}(n,k)$ denote the r-Whitney numbers of the first and second kind, respectively, defined in [10] by

(1.6)
$$q^{n}(x)_{n} = \sum_{k=0}^{n} w(n,k) (qx+r)^{k},$$

(1.7)
$$(qx+r)^n = \sum_{k=0}^n q^k W(n,k) (x)_k.$$

The exponential generating function of W(n,k) is given by [10]

(1.8)
$$\sum_{n\geq k} W(n,k) \frac{t^n}{n!} = \frac{e^{rt}}{k!} \left(\frac{e^{qt}-1}{q}\right)^k.$$

The parameters $r \ge 0$ and q > 0 are usually taken to be integers, but both may also be considered as indeterminates [11].

Komatsu [9] defined the Cauchy polynomials with a q parameter of the first and second kind, denoted by $c_n^q(z)$, $\hat{c}_n^q(z)$, respectively

(1.9)
$$c_n^q(z) = \int_0^1 (x - z|q)_n \, dx \qquad q \neq 0,$$

(1.10)
$$\hat{c}_n^q(z) = \int_0^1 (-x + z|q)_n \, dx \qquad q \neq 0$$

Recently, Shiha [12] gave explicit formulae for $c_n^q(r)$ and $\hat{c}_n^q(r)$ in terms of r-Whitney numbers of the first kind

(1.11)
$$c_n^q(r) = \sum_{k=0}^n w(n,k) \frac{1}{k+1} \qquad q \neq 0,$$

(1.12)
$$\hat{c}_n^q(-r) = \sum_{k=0}^n (-1)^k w(n,k) \frac{1}{k+1} \qquad q \neq 0.$$

The relations between $c_n^q(r),\, \hat{c}_n^q(r)$ and W(n,k) are (see [12])

(1.13)
$$\sum_{k=0}^{n} W(n,k) c_{k}^{q}(r) = \frac{1}{n+1}$$

BERNOULLI POLYNOMIALS WITH A q PARAMETER IN TERMS OF r-WHITNEY NUMBERS

(1.14)
$$\sum_{k=0}^{n} W(n,k) \, \hat{c}_{k}^{q}(-r) = \frac{(-1)^{n}}{n+1}$$

2. The basic results

Theorem 1. We define the Bernoulli polynomials with a q parameter in terms of W(n,k) by

(2.1)
$$B_n^q(r) = \sum_{k=0}^n (-1)^k \, \frac{k!}{k+1} \, W(n,k).$$

Proof. Using (1.8), we get

$$\begin{split} \sum_{n=0}^{\infty} \sum_{k=0}^{n} (-1)^{k} \frac{k!}{k+1} W(n,k) \frac{t^{n}}{n!} &= \sum_{k=0}^{\infty} (-1)^{k} \frac{k!}{k+1} \sum_{n=k}^{\infty} W(n,k) \frac{t^{n}}{n!} \\ &= \sum_{k=0}^{\infty} (-1)^{k} \frac{k!}{k+1} \frac{e^{rt}}{k!} \left(\frac{e^{qt}-1}{q}\right)^{k} \\ &= \frac{q e^{rt}}{1-e^{qt}} \sum_{k=0}^{\infty} \left(\frac{1-e^{qt}}{q}\right)^{k+1} \frac{1}{k+1} \\ &= \frac{q e^{rt}}{1-e^{qt}} \sum_{k=1}^{\infty} \left(\frac{1-e^{qt}}{q}\right)^{k} \frac{1}{k} \\ &= \sum_{n=0}^{\infty} B_{n}^{q}(r) \frac{t^{n}}{n!}. \end{split}$$

Comparing the coefficients of both sides, we get (2.1) (notice that e^{rt} is replaced by e^{-rt} in [2]).

On the other hand,

(2.2)

$$\sum_{n=0}^{\infty} B_n^q(r) \frac{t^n}{n!} = \frac{q \, e^{rt}}{1 - e^{qt}} \sum_{k=1}^{\infty} \left(\frac{1 - e^{qt}}{q}\right)^k \frac{1}{k}$$

$$= \frac{q \, e^{rt}}{1 - e^{qt}} \left(-\ln(1 - (\frac{1 - e^{qt}}{q}))\right)$$

$$= \frac{q \, e^{rt}}{e^{qt} - 1} \left(\ln\frac{q - 1 + e^{qt}}{q}\right).$$

The first few polynomials are

$$\begin{array}{l} B_0^q(r) = 1, \\ B_1^q(r) = r - \frac{1}{2}, \\ B_2^q(r) = r^2 - r - \frac{1}{2}q + \frac{2}{3}, \\ B_3^q(r) = r^3 - \frac{3}{2}r^2 + (2 - \frac{3}{2}q)r - \frac{1}{2}q^2 + 2q - \frac{3}{2}, \\ B_4^q(r) = r^4 - 2r^3 + (4 - 3q)r^2 - 2(q^2 - 4q + 3)r - \frac{1}{2}q^3 + \frac{14}{3}q^2 - 9q + \frac{24}{5}. \end{array}$$

Remark 1. If r = 0, $B_n^q(0) = B_n^q$ are the Bernoulli numbers with q parameter, using (2.2), we get

F. A. SHIHA

(2.3)
$$\sum_{n=0}^{\infty} B_n^q \frac{t^n}{n!} = \frac{q}{e^{qt} - 1} \left(\ln \frac{q - 1 + e^{qt}}{q} \right),$$

 $W_{q,0}(n,k) = q^{n-k} S(n,k)$, then B_n^q can be expressed in terms of S(n,k) as

(2.4)
$$B_n^q = \sum_{k=0}^n (-1)^k \, \frac{k!}{k+1} \, q^{n-k} \, S(n,k),$$

If q = 1, $B_n^1(r) = B_n(r)$ and $W_{1,r}(n,k)$ are reduced to $S_r(n+r,k+r)$ then, we get the explicit formula (1.3), and

(2.5)
$$\sum_{n=0}^{\infty} B_n(r) \frac{t^n}{n!} = \frac{t \, e^{rt}}{e^t - 1}.$$

If q = 1 and r = 0, then $B_n^1(0) = B_n$ and $W_{1,0}(n,k)$ are reduced to S(n,k) then, we get the explicit formula (1.2).

It is known that the r-Whitney numbers satisfy the following orthogonality relation [10]:

(2.6)
$$\sum_{l=k}^{n} w(n,l) W(l,k) = \sum_{l=k}^{n} W(n,l) w(l,k) = \delta_{kn},$$

where δ_{kn} is the Kronecker delta.

Corollary 1. The relation between w(n,k) and $B_n^q(r)$ is given by

(2.7)
$$\sum_{j=0}^{n} (-1)^n w(n,j) B_j^q(r) = \frac{n!}{n+1}.$$

Proof.

$$\sum_{j=0}^{n} (-1)^{n} w(n,j) B_{j}^{q}(r) = \sum_{j=0}^{n} \sum_{k=0}^{j} (-1)^{n+k} \frac{k!}{k+1} w(n,j) W(j,k)$$
$$= \sum_{k=0}^{n} (-1)^{n+k} \frac{k!}{k+1} \sum_{j=k}^{n} w(n,j) W(j,k)$$
$$= \sum_{k=0}^{n} (-1)^{n+k} \frac{k!}{k+1} \delta_{kn} = \frac{n!}{n+1}.$$

The numbers W(n,k) are determined by (see [10]).

(2.8)
$$W(n,k) = \frac{1}{q^k k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} (r+jq)^n,$$

then, we obtain

Theorem 2. For $n \ge 0$, we have

(2.9)
$$B_n^q(r) = \sum_{k=0}^n \frac{1}{q^k (k+1)} \sum_{j=0}^k (-1)^j \binom{k}{j} (r+jq)^n$$

Theorem 3. For $n \ge 0$, we have

(2.10)
$$c_n^q(r) = \sum_{k=0}^n \sum_{j=0}^k \frac{(-1)^k}{k!} w(n,k) w(k,j) B_j^q(r),$$

(2.11)
$$\hat{c}_n^q(-r) = \sum_{k=0}^n \sum_{j=0}^k \frac{1}{k!} w(n,k) w(k,j) B_j^q(r),$$

(2.12)
$$B_n^q(r) = \sum_{k=0}^n \sum_{j=0}^k (-1)^k \, k! \, W(n,k) \, W(k,j) \, c_j^q(r).$$

(2.13)
$$B_n^q(r) = \sum_{k=0}^n \sum_{j=0}^k k! W(n,k) W(k,j) \hat{c}_j^q(-r).$$

Proof. We shall prove the first and the third identities. Other identities are proven similarly. By (2.7), and using (1.11), we have

$$\sum_{k=0}^{n} \sum_{j=0}^{k} \frac{(-1)^{k}}{k!} w(n,k) w(k,j) B_{j}^{q}(r) = \sum_{k=0}^{n} \frac{1}{k!} w(n,k) \sum_{j=0}^{k} (-1)^{k} w(k,j) B_{j}^{q}(r)$$
$$= \sum_{k=0}^{n} \frac{1}{k+1} w(n,k) = c_{n}^{q}(r).$$

By (1.13), and using (2.1), we obtain

$$\sum_{k=0}^{n} \sum_{j=0}^{k} (-1)^{k} k! W(n,k) W(k,j) c_{j}^{q}(r) = \sum_{k=0}^{n} (-1)^{k} k! W(n,k) \sum_{j=0}^{k} W(k,j) c_{j}^{q}(r)$$
$$= \sum_{k=0}^{n} (-1)^{k} k! W(n,k) \frac{1}{k+1} = B_{n}^{q}(r).$$

It is known that (see [3])

(2.14)
$$W_{q,r+s}(n,k) = \sum_{j=k}^{n} \binom{n}{j} r^{n-j} W_{q,s}(j,k).$$

Therefore, we get the following properties, which are similar to those for classical Bernoulli polynomials and numbers.

Theorem 4. For $n \ge 0$, we have

(2.15)
$$B_n^q(r+s) = \sum_{j=0}^n \binom{n}{j} r^{n-j} B_j^q(s).$$

(2.16)
$$B_n^q(r) = \sum_{j=0}^n \binom{n}{j} r^{n-j} B_j^q.$$

Proof.

$$B_n^q(r+s) = \sum_{k=0}^n (-1)^k \frac{k!}{k+1} W_{q,r+s}(n,k)$$

= $\sum_{k=0}^n \sum_{j=k}^n (-1)^k \frac{k!}{k+1} {n \choose j} r^{n-j} W_{q,s}(j,k)$
= $\sum_{j=0}^n {n \choose j} r^{n-j} \sum_{k=0}^j (-1)^k \frac{k!}{k+1} W_{q,s}(j,k) = \sum_{j=0}^n {n \choose j} r^{n-j} B_j^q(s).$

Setting s = 0 in (2.15), we get the second relation (2.16)

Remark 2. Setting q = 1 in (2.15) and (2.16), we obtain the following properties of $B_n(r)$ [13]

(2.17)
$$B_n(r+s) = \sum_{j=0}^n \binom{n}{j} r^{n-j} B_j(s), \text{ and } B_n(r) = \sum_{j=0}^n \binom{n}{j} r^{n-j} B_j.$$

References

- A. Bayad and Y. Hamahata, Polylogarithms and poly-Bernoulli polynomials, Kyushu J. Math. 65, 15–24 (2011).
- [2] M. Cenkci and T. Komatsu, Poly-Bernoulli numbers and polynomials with a q parameter, J. Number Theory 152, 38–54 (2015).
- [3] G. S. Cheon and J. H. Jung, r-Whitney numbers of Dowling lattices, Discrete Math. 312, 2337–2348 (2012).
- [4] L. Comtet, Advanced Combinatorics, Reidel, Dordrecht, 1974.
- [5] M.-A. Coppo and B. Candelpergher, The Arakawa-Kaneko zeta function, Ramanujan J. 22, 153–162 (2010).
- [6] R. L. Graham, D. E. Knuth, and O. Patashnik, Concrete Mathematics-A Foundation for Computer Science, 2nd ed., Addison-Wesley Publishing Company, Reading, MA, 1994.
- [7] B.-N. Guo, I. Mezö, and Feng Qi, An explicit formula for Bernoulli polynomials in terms of r-Stirling numbers of the second kind, Rocky Mountain J. Math. 46 6 (2016), 1919–1923 (2016).
- [8] M. Kaneko, Poly-Bernoulli numbers, J. Théor. Nombres Bordeaux. 9, 199–206. (1997)
- [9] T. Komatsu, Poly-Cauchy numbers with a q parameter, Ramanujan J. 31, 353–371 (2013).
- [10] I. Mezö, A new formula for the Bernoulli polynomials, Results Math. 58, 329–335 (2010).
- [11] M. Shattuck, Identities for Generalized Whitney and Stirling Numbers, J. Integer Seq. 20, Article 17.10.4. (2017)
- [12] F. A. Shiha, Explicit Formulae of Cauchy Polynomials with a q Parameter in Terms of r-Whitney Numbers. Preprint.
- [13] N. M. Temme, Special Functions: An Introduction to the Classical Functions of Mathematical Physics, Wiley–Interscience Publ., New York, 1996.

DEPARTMENT OF MATHEMATICS FACULTY OF SCIENCE, MANSOURA UNIVERSITY MANSOURA, 35516, EGYPT.

E-mail address: fshiha@yahoo.com, fshiha@mans.edu.eg

 $\mathbf{6}$