# Bose–Einstein condensation in the Luttinger–Sy model with contact interaction

Joachim Kerner<sup>1</sup>, Maximilian Pechmann<sup>2</sup>, and Wolfgang Spitzer<sup>3</sup>

Department of Mathematics and Computer Science FernUniversität in Hagen 58084 Hagen Germany

#### Abstract

We study bosons on the real line in a Poisson random potential (Luttinger–Sy model) with contact interaction in the thermodynamic limit at absolute zero temperature. We prove that generalized Bose–Einstein condensation (BEC) occurs almost surely if the intensity  $\nu_N$  of the Poisson potential satisfies  $[\ln(N)]^4/N^{1-2\eta} \ll \nu_N \lesssim 1$ for arbitrary  $0 < \eta \leq 1/3$ . We also show that the contact interaction alters the type of condensation, going from a type-I BEC to a type-III BEC as the strength of this interaction is increased. Furthermore, for sufficiently strong contact interactions and  $0 < \eta < 1/6$  we prove that the mean particle density in the largest interval is almost surely bounded asymptotically by  $\nu_N N^{3/5+\delta}$  for  $\delta > 0$ .

<sup>&</sup>lt;sup>1</sup>E-mail address: joachim.kerner@fernuni-hagen.de

<sup>&</sup>lt;sup>2</sup>E-mail address: maximilian.pechmann@fernuni-hagen.de

<sup>&</sup>lt;sup>3</sup>E-mail address: wolfgang.spitzer@fernuni-hagen.de

### 1 Introduction

Bose–Einstein condensation (BEC) generally refers to a macroscopic occupation of a singleparticle state. In the non-interacting Bose gas and as illustrated by the initiating work of Einstein [Ein24, Ein25], this one-particle state is the ground state of the one-particle Hamiltonian while, in the more general case of an interacting gas, one considers the ground state of the reduced one-particle density matrix instead [PO56]. More general definitions of BEC leading to the notion of generalized BEC require a macroscopic occupation of an arbitrarily narrow energy band of single-particle states [Cas68, vL82, van83, vLP86, vLL86, ZB01] and condensation in this generalized sense is thought to be thermodynamically more stable [Gir60, JPZ10]. More explicitly, BEC is classified into three different types: Type-I or type-II is present whenever finitely or infinitely many single-particle states in this narrow energy band are macroscopically occupied. A generalized BEC without any single-particle state in this band being macroscopically occupied is defined as type-III.

In this paper, we are concerned with (generalized) BEC in the Luttinger–Sy model [LS73b, LS73a] with a contact interaction of strength g of the Lieb–Liniger type [LL63] in the thermodynamic limit at absolute zero temperature. In particular, we are interested in determining its type and in estimating the maximal particle density per interval. Note that the Luttinger–Sy model is a random model with the real line as one-particle configuration space in which singular point impurities (or singular external potentials for that matter) are Poisson distributed and consequently divide the real line into a countable number of disjoint intervals. In the original model, the intensity  $\nu$  of the Poisson distribution was kept fixed but in this paper we will allow it to vary with the number of particles, N. Likewise, the strength g of the contact interaction depends on N. In fact, we always assume that  $g_N$  goes to 0 as N becomes large.

It is interesting to note that, despite its singularity, the Luttinger–Sy model is considered a good approximation to more general Poisson random potentials [Zag07]. Regarding Bose–Einstein condensation, being in accordance with a conjecture for bosonic systems with quite general random potentials [LPZ04], BEC in the non-interacting Luttinger–Sy model is of type-I and leads, in the thermodynamic limit, to an unbounded particle density of order  $\ln(N)$  in the largest interval [Zag07]. Due to the diverging particle density, interactions between bosons cannot be neglected and BEC has to be investigated for the interacting Luttinger–Sy model. So far, however, only a limited amount of rigorous results regarding BEC in (random and non-random) interacting bosonic systems exist, e.g. [KLS88, LS02, LVZ03, LSSY05]. This is even more true for one-dimensional many-particle systems [SYZ12, dS86] or quasi one-dimensional systems such as quantum graphs [BK14, BK16] since Bose-Einstein condensation in one dimension is much more unstable [Hoh67, LW79].

In large parts we follow the paper [SYZ12] by Seiringer, Yngvason, and Zagrebnov, in which an equivalent model is considered, however, with the unit interval as the fixed one-particle configuration space. This implies that the thermodynamic limit is a highdensity limit. There, BEC is exclusively discussed in the sense of a macroscopic occupation of a single-particle state. These authors were able to prove condensation and to make conclusions of its localization among the intervals, assuming a fast decay of the intensity, i.e.,  $1/N \ll \nu_N \ll [\ln(N)]^3/N$ , cf. Appendix B. In the present paper, by considering BEC in the generalized sense, we are able to prove condensation in the almost sure sense and to obtain knowledge of the distribution of the condensate among the intervals for an intensity  $[\ln(N)]^4/N^{1-\eta} \ll \nu_N \lesssim 1$  for any  $0 < \eta \leq 1/3$ , see Theorem 3.1. In addition, we show that the type of the BEC changes with the strength of the contact interaction, see Theorem 3.3.

We also provide an upper bound for the particle density in the largest interval that depends on the strength of the interaction. In particular, we prove that for strong interactions  $g_N \gg \nu_N N^{-1/6} [\ln(N)]^{-2}$ , this density is almost surely non-divergent in case of  $[\ln(N)]^4/N^{1-2\eta} \ll \nu_N \lesssim N^{-3/5-\delta}$  for any  $0 < \eta < 1/6$ ,  $0 < \delta < 2/5 - 2\eta$  and is asymptotically bounded by  $\nu_N N^{3/5+\delta}$  for any  $\delta > 0$  in the case of  $[\ln(N)]^4/N^{1-2\eta} \ll \nu_N \lesssim 1$ ,  $0 < \eta < 1/6$ , see Theorem 3.4.

## 2 The model

We consider bosons on the real line that interact pairwise repulsively via a delta-function potential as in the Lieb-Liniger model [LL63] and with an external random potential as in the Luttinger-Sy model [LS73a]. For given intensity  $\nu > 0$  we think of a Poisson point process X on  $\mathbb{R}$  as a random variable on some probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . We assume that  $X(\omega) = \{x_j(\omega) : j \in \mathbb{Z}\}$  is a strictly increasing sequence of points  $x_j = x_j(\omega) \in \mathbb{R}$ and that  $0 \in (x_0(\omega), x_1(\omega))$ . For any bounded Borel set  $\Lambda \subset \mathbb{R}$  with Lebesgue volume  $|\Lambda|$ , the probability that  $\Lambda$  contains exactly m points  $x_j(\omega)$  is

$$\mathbb{P}(\operatorname{card}(X(\omega) \cap \Lambda) = m) = \frac{(\nu|\Lambda|)^m}{m!} e^{-\nu|\Lambda|}, \quad m \in \mathbb{N}_0.$$
(2.1)

And, if  $\Lambda$  and  $\Lambda'$  are two such subsets which are disjoint, then the events  $\{X(\omega) \cap \Lambda\}$  and  $\{X(\omega) \cap \Lambda'\}$  are (stochastically) independent.

In this paper, we will write  $\mathbb{P} = \mathbb{P}^{\nu}$  and  $\mathbb{E} = \mathbb{E}^{\nu}$  if  $\nu > 0$  is constant. The external random potential we consider is then of the form  $\sigma \sum_{j \in \mathbb{Z}} \delta(z - x_j(\omega))$ . Actually, we restrict the analysis to the case of infinite strength, that is, we set informally  $\sigma = \infty$ . The pair interaction is also described by a delta-function but now of finite strength  $g \ge 0$ ; in fact, g will tend to zero as the number of particles increases. In order to define the full many-boson Hamiltonian we only need (since  $\sigma = \infty$  implies Dirichlet boundary conditions at the endpoints of the intervals) to define the many-boson Hamiltonian on a bounded interval and take their direct sum.

When we perform the thermodynamic limit we define for any particle density  $\rho > 0$  and any particle number  $N \in \mathbb{N}$  the length  $L_N := N/\rho$  and introduce the interval  $\Lambda_N := (-L_N/2, L_N/2)$ . Later, we will define  $W_j = W_j(\omega) := (x_j(\omega), x_{j+1}(\omega)) \cap \Lambda_N$  for any realization  $X(\omega)$  and denote the number of non-empty intervals  $W_j$  by  $k_N$ , which almost surely is finite and has the asymptotic behavior  $\lim_{N\to\infty} k_N/L_N = \nu$  [Zag07], see also Theorem C.2. But for now, we consider an arbitrary interval  $\Lambda \subset \mathbb{R}$ , an arbitrary

set  $\{x_j : j \in \mathbb{Z}\} \subset \mathbb{R}$  such that only finitely many are contained in  $\Lambda$  and define  $W_j := (x_j, x_{j+1}) \cap \Lambda$ . Let  $l_j := |W_j|$  be the (later random) length of the interval  $W_j$ . Note that  $\sum_{j \in \mathbb{Z}} l_j = L_N$  and  $L^2(\Lambda) = \bigoplus_{j \in \mathbb{Z}} L^2(W_j)$ . The inner product is always denoted by  $\langle \cdot, \cdot \rangle$  with norm  $\|\psi\| := \langle \psi, \psi \rangle^{1/2} := (\int |\psi(x)|^2 dx)^{1/2}$ .

We impose Dirichlet boundary conditions for the Laplacian  $-\partial_z^2$  at the end points of an interval W with length l := |W| > 0. For any  $n \in \mathbb{N}, g \ge 0$  we define the *n*-boson Hamiltonian on the interval W,

$$H(n,l,g) := -\sum_{i=1}^{n} \partial_{z_i}^2 + g \sum_{1 \le i < j \le n} \delta(z_i - z_j) , \quad g \ge 0 , \qquad (2.2)$$

acting as a quadratic form on the *n*-fold symmetric tensor product  $\bigotimes_{s}^{n} L^{2}(W)$ . For completeness, we set H(0, l, g) = 0 for any  $l \geq 0$ .

Now, let  $\Lambda, W_j$ , and  $l_j$  be as above. We call  $\mathbf{N} = \{M_j : j \in \mathbb{Z}\}$  an admissible sequence (ad. seq.) of the particle number N if  $M_j \in \mathbb{N}_0$ , if  $M_j = 0$  for any  $j \in \mathbb{Z}$  with  $l_j = 0$ , and if the total number of particles is  $\sum_{j \in \mathbb{Z}} M_j = N$ . Furthermore, we call  $\{M_j\}_{j \in \mathbb{Z}}$  a general admissible sequence (gen. ad. seq.) if it fulfills the first two requirements of but not the last. Then the full N-boson Hamiltonian on the interval  $\Lambda$ ,

$$H(\mathbf{N}, \Lambda, g) := \bigoplus_{j \in \mathbb{Z}} H(M_j, l_j, g), \qquad (2.3)$$

acts on the N-fold symmetric tensor product  $\bigotimes_{s}^{N} L^{2}(\Lambda)$ .

For any (general) admissible sequence  $\{M_j\}_{j\in\mathbb{Z}}$  we denote by  $\mathcal{M}_N^> := \max\{M_j : j \in \mathbb{Z}\}$ the largest particle number  $M_j$ . Sometimes we will also need the (later random) lengths of the intervals  $l_j$  arranged in descending order. We denote them by  $\ell_N^> := l_N^{>,1} \ge l_N^{>,2} \ge \ldots$ ,  $\ell_N^> = \max\{l_j : j \in \mathbb{Z}\}$  being the length of the largest subinterval  $W_j$  of  $\Lambda$ . Note here that we added the particle number N as an index since these numbers will eventually depend on N.

One of our main concerns is the ground-state energy of the full N-boson Hamiltonian in the thermodynamic limit. So, let  $E_0^{\text{QM}}(n, l, g)$  and  $E_0^{\text{QM}}(\mathbf{N}, \Lambda, g)$  be the ground-state energies of the Hamiltonians H(n, l, g) and  $H(\mathbf{N}, \Lambda, g)$ , respectively. That is,

$$E_0^{\text{QM}}(n,l,g) := \inf\left\{ \langle \psi, H(n,l,g)\psi \rangle : \psi \in \bigotimes_s^n \mathrm{H}_0^1(W), \|\psi\| = 1 \right\},$$
(2.4)

$$E_0^{\text{QM}}(\boldsymbol{N},\Lambda,g) := \inf\left\{ \langle \psi, H(\boldsymbol{N},\Lambda,g)\psi \rangle : \psi \in \bigoplus_j \bigotimes_s^{M_j} \mathrm{H}_0^1(W_j), \|\psi\| = 1 \right\}.$$
(2.5)

The latter energy will eventually be random due to the random location of the points  $x_j$  that partition the interval  $\Lambda_N$  into the intervals  $W_j$ .

The relevant quantity is the lowest ground-state energy among all possible distributions of particles in the intervals  $W_i$ , i.e.,

$$E_{\mathrm{LS}}^{\mathrm{QM}}(N,\Lambda,g) := \inf \left\{ E_0^{\mathrm{QM}}(\boldsymbol{N},\Lambda,g) : \boldsymbol{N} \text{ ad. seq. of } N \right\}$$
  
$$= \inf \left\{ \sum_{j \in \mathbb{Z}} E_0^{\mathrm{QM}}(M_j, l_j, g) : \{M_j\}_{j \in \mathbb{Z}} \text{ ad. seq.}, \sum_{j \in \mathbb{Z}} M_j = N \right\}$$
  
$$= \inf \left\{ \sum_{j \in \mathbb{Z}} E_0^{\mathrm{QM}}(M_j, l_j, g) : \{M_j\}_{j \in \mathbb{Z}} \text{ gen. ad. seq.}, \sum_{j \in \mathbb{Z}} M_j \ge N \right\}.$$
  
(2.6)

Note that, for fixed  $\Lambda \subset \mathbb{R}$  and  $g \geq 0$ ,  $E_{\text{LS}}^{\text{QM}}(N, \Lambda, g)$  is non-decreasing in N.

**Remark 2.1.** The energy  $E_{\rm LS}^{\rm QM}(N,\Lambda,g)$  is (for  $\sigma = \infty$ ) the ground-state energy of the self-adjoint operator

$$H(N,\Lambda,g) := -\sum_{i=1}^{N} \partial_{z_i}^2 + g_N \sum_{1 \le i < k \le N} \delta(z_i - z_k) + \sigma \sum_{j \in \mathbb{Z}} \delta(z - x_j)$$
(2.7)  
$$\bigotimes_{i=1}^{N} L^2(\Lambda).$$

defined on  $\bigotimes_{s}^{N}$ 

The ground-state energy  $E_0^{\text{QM}}(N, \Lambda, g)$  will be approximated by the ground-state energy of a mean-field Hamiltonian h: In order to introduce this operator we fix some interval W of length l = |W| > 0 and, as above, let  $q \ge 0$ . For n > 0, the Gross-Pitaevskii (GP) functional  $\mathcal{E}^{\mathrm{GP}}(n,l,g)$  on W with domain  $\{\phi \in \mathrm{H}^1_0(W) : \int_W |\phi(z)|^2 \,\mathrm{d}z = n\}$  then maps a function  $\phi$  from the Sobolev space  $\mathrm{H}^{1}_{0}(W)$  to

$$\mathcal{E}^{\rm GP}(n,l,g)[\phi] := \int_{W} \left( |\phi'(z)|^2 + \frac{g}{2} |\phi(z)|^4 \right) \mathrm{d}z \,.$$
(2.8)

As is well-known [LSY00], there is a unique, non-negative minimizer of  $\mathcal{E}^{\text{GP}}(n, l, g)$ , which we denote by  $\phi_{n,l,g}^{\text{GP}} \in \mathrm{H}_0^1(W)$ . Let

$$E^{\rm GP}(n,l,g) := \mathcal{E}^{\rm GP}(n,l,g)[\phi_{n,l,g}^{\rm GP}] = \inf\left\{\mathcal{E}^{\rm GP}(n,l,g)[\phi] : \phi \in \mathrm{H}_0^1(W), \|\phi\|^2 = n\right\}.$$
 (2.9)

Setting  $e^{\text{GP}}(g) := E^{\text{GP}}(1, 1, g)$ , we obtain by scaling [LSY04]

$$E^{\rm GP}(n,l,g) = nE^{\rm GP}(1,l,ng) = \frac{1}{l^2}E^{\rm GP}(n,1,lg) = \frac{n}{l^2}e^{\rm GP}(nlg) .$$
(2.10)

We also set  $E^{\text{GP}}(0, l, g) := 0$  and  $\phi_{l,g}^{\text{GP}} := \phi_{1,l,g}^{\text{GP}}$  for any  $l, g \ge 0$ . Finally, the self-adjoint (one-particle) mean-field Hamiltonian h = h(l, g) on the oneparticle Hilbert space  $L^2(W)$  shall be

$$h(l,g) := -\partial_z^2 + g|\phi_{l,g}^{\rm GP}|^2 - \frac{g}{2} \int_W |\phi_{l,g}^{\rm GP}(z)|^4 \, \mathrm{d}z \,, \qquad (2.11)$$

and on  $\bigotimes_{s}^{n} L^{2}(W)$  we introduce

$$h^{(i)}(l,g) := \mathbb{1} \otimes \cdots \otimes \mathbb{1} \otimes h(l,g) \otimes \mathbb{1} \otimes \cdots \otimes \mathbb{1}, \qquad (2.12)$$

where h(l, g) acts on the *i*th component in the *n*-fold tensor product.

**Remark 2.2.** The minimizer  $\phi_{l,g}^{\text{GP}}$  is also the ground state of h(l,g) with corresponding ground-state energy  $E^{\text{GP}}(1,l,g)$ .

Now, in a first result we compare the operator H(n, l, g) with the second quantization of h(l, g), i.e.,  $\sum_{i} h^{(i)}(l, g)$ .

**Lemma 2.3.** Let  $n \in \mathbb{N}_0$ , l > 0, and g > 0 be given. Then there exist finite, positive constants  $c, \tilde{c}$  (independent of n and l) such that if  $\tilde{c}(n^{1/3}lg)^{1/2} < 1$  we have, for some  $\tau$  with  $0 < \tau < \frac{\tilde{c}}{2}(n^{1/3}lg)^{1/2}$ ,

$$H(n,l,g) \ge (1-\tau) \sum_{i=1}^{n} h^{(i)}(l,ng) + \tau n l^{-2} e^{\text{GP}}(nlg) - c n^{5/3} g l^{-1} \left( e^{\text{GP}}(nlg) \right)^{1/2} .$$
 (2.13)

This estimate is contained in the proof of [SYZ12, Theorem 2.1] but we take a slightly different route, which is the reason why we recall the main steps.

*Proof.* The statement is trivial for n = 0 and we therefore assume n > 0 in the following. Let  $\epsilon \in (0, 1)$  and b > 0 be given. As demonstrated in [SYZ12], one has the operator inequality

$$-\epsilon \partial_z^2 + g\delta(z) \ge \frac{g}{1 + bg/(2\epsilon)} \delta_b(z) , \qquad (2.14)$$

where  $\delta_b(z) := \frac{1}{2b} \exp(-|z|/b)$  is a function of positive type.

Now, setting  $p_i := -i\partial_{z_i}$  we follow [SYZ12, (7–15)] to obtain

$$H(n, l, g) \geq \sum_{i=1}^{n} \left[ \left( 1 - \frac{\epsilon}{2} \right) p_i^2 \right] + g \left( 1 + \frac{bng}{2\epsilon} \right)^{-1} \sum_{1 \leq i < k \leq N} \delta_b(z_i - z_k)$$
  
$$\geq \sum_{i=1}^{n} \left[ \left( 1 - \frac{\epsilon}{2} \right) p_i^2 + ng |\phi_{l,ng}^{GP}(z_i)|^2 - \frac{ng}{2} \int_{W} |\phi_{l,ng}^{GP}(z)|^4 \, \mathrm{d}z \right] + \sum_{i=1}^{n} \left[ -\frac{b(ng)^2}{2\epsilon} \left( E^{GP}(1, l, ng) \right)^{1/2} - cng \left( E^{GP}(1, l, ng) \right)^{3/4} b^{1/2} - \frac{ng}{4nb} \right]$$
(2.15)

for some constant c > 0. Since

$$\sum_{i=1}^{n} \left[ \left( 1 - \frac{\epsilon}{2} \right) p_i^2 + ng |\phi_{l,ng}^{\text{GP}}(z_i)|^2 - \frac{ng}{2} \int_{W} |\phi_{l,ng}^{\text{GP}}(z)|^4 \, \mathrm{d}z \right]$$

$$\geq \sum_{i=1}^{n} \left[ \left( 1 - \frac{\epsilon}{2} \right) h^{(i)}(l,ng) - \frac{\epsilon}{2} E^{\text{GP}}(1,l,ng) \right] , \qquad (2.16)$$

we obtain

$$H(n,l,g) \ge \sum_{i=1}^{n} \left[ \left( 1 - \frac{\epsilon}{2} \right) h^{(i)}(l,ng) + \frac{\epsilon}{2} E^{\text{GP}}(1,l,ng) \right] - \sum_{i=1}^{n} \left[ \epsilon E^{\text{GP}}(1,l,ng) + \frac{b(ng)^2}{2\epsilon} \left( E^{\text{GP}}(1,l,ng) \right)^{1/2} + cng \left( E^{\text{GP}}(1,l,ng) \right)^{3/4} b^{1/2} + \frac{ng}{4nb} \right].$$
(2.17)

Next, choosing  $\epsilon = 2^{-1/2} b^{1/2} ng \left( E^{\text{GP}}(1,l,ng) \right)^{-1/4}$  and  $b = \tilde{c}^2 n^{-2/3} \left( E^{\text{GP}}(1,l,ng) \right)^{-1/2}$  with some constant  $\tilde{c} > 0$  yields,  $\tau := \epsilon/2$ ,

$$H(n,l,g) \ge (1-\tau) \sum_{i=1}^{n} h^{(i)}(l,ng) + \tau n E^{\text{GP}}(1,l,ng) - \hat{c}n(n)^{2/3} g \left( E^{\text{GP}}(1,l,ng) \right)^{1/2} (2.18)$$

with some constant  $\hat{c} > 0$ .

Finally, note that the bound on  $\tau$  follows from the definitions of  $\epsilon$  and b and from

$$E^{\rm GP}(1,l,ng) = \frac{1}{l^2} e^{\rm GP}(nlg) \ge \frac{nlg}{2l^2} \int_0^1 |\phi_{1,nlg}^{\rm GP}(z)|^4 \, \mathrm{d}z \ge \frac{ng}{2l} \,, \tag{2.19}$$

which holds due to

$$\int_{0}^{1} |\phi_{1,nlg}^{\text{GP}}(z)|^4 \, \mathrm{d}z = \left( \left\| |\phi_{1,nlg}^{\text{GP}}|^2 \right\|_{\mathrm{L}^2(0,1)} \|1\|_{L^2(0,1)} \right)^2 \ge \left( \left\| |\phi_{1,nlg}^{\text{GP}}|^2 \cdot 1 \right\|_{\mathrm{L}^1(0,1)} \right)^2 = 1 \,, \quad (2.20)$$

where we used Hölder's inequality.

In a next step we bound the ground-state energy of each component in (2.3) in terms of the corresponding Gross–Pitaevskii energy.

**Theorem 2.4** (An energy bound). Let  $0 < \epsilon < 1$ ,  $g \ge 0$  and  $N \in \mathbb{N}$  be given. Then, for any general admissible sequence  $\{M_j\}_{j\in\mathbb{Z}}$  with associated lengths  $\{l_j\}_{j\in\mathbb{Z}}$  that fulfills inequality  $[(\mathcal{M}_N^{>})^{1/3}\ell_N^{>}g]^{1/2} < \epsilon / \max\{\sqrt{2}c, \tilde{c}\}$  (where c and  $\tilde{c}$  are the constants from Lemma 2.3), one has

$$M_j E^{\rm GP}(1, l_j, M_j g) \ge E_0^{\rm QM}(M_j, l_j, g) \ge (1 - \epsilon) M_j E^{\rm GP}(1, l_j, M_j g) ,$$
 (2.21)

for all  $j \in \mathbb{Z}$  where  $l_j > 0$ .

*Proof.* Since the inequality (2.21) is trivial for  $M_j = 0$ , we assume  $M_j \ge 1$  in the following.

Upper bound: this follows directly from a standard variational argument using the product state  $\otimes \phi_{l_j,M_jg}^{\text{GP}}$ .

Lower bound: Since  $h^{(i)}(l_j, M_j g) \ge E^{\text{GP}}(1, l_j, M_j g)$  for  $j \in \mathbb{Z}$  with  $M_j \ge 1$  by Remark 2.2 and since

$$\tilde{c} \left[ M_j^{1/3} l_j g \right]^{1/2} \le \tilde{c} \left[ \left( \mathcal{M}_N^{>} \right)^{1/3} \ell_N^{>} g \right]^{1/2} < 1 ,$$
 (2.22)

(2.13) implies

$$E_0^{\text{QM}}(M_j, l_j, g) \ge M_j E^{\text{GP}}(1, l_j, M_j g) \left( 1 - c M_j^{2/3} g \left( E^{\text{GP}}(1, l_j, M_j g) \right)^{-1/2} \right) .$$
(2.23)

Finally, applying inequality (2.19) then yields the statement.

In a next result we estimate the fraction of particles occupying the Gross–Pitaevskii ground state  $\phi_{l_j,M_jg}^{GP}$ . For this note that

$$n_{M_j} := \operatorname{Tr}_{\mathcal{L}^2(W_N^j)}[\rho_{M_j}^{(1)} \phi_{l_j,M_jg}^{\mathrm{GP}}(\phi_{l_j,M_jg}^{\mathrm{GP}}, \cdot)]$$
(2.24)

is the number of particles occupying  $\phi_{l_j,M_jg}^{\text{GP}}$ . Here

$$\rho_{M_j}^{(1)} := \begin{cases} M_j \operatorname{Tr}_{\mathcal{H}_{W_j}^{(M_j - 1)}}[\rho_{M_j}] & \text{if } M_j \ge 2 ,\\ \rho_1 & \text{if } M_j = 1 , \end{cases}$$
(2.25)

with  $\mathcal{H}_{W_j}^{(M_j-1)} := S_{M_j-1} L^2(W_j^{M_j-1})$  and  $S_{M_j-1}$  being the symmetrizer on  $L^2(W_j^{M_j-1})$ , is the reduced one-particle density matrix that is obtained from the many-particle ground state  $\rho_{M_j}$  of  $H(M_j, l_j, g)$  by taking the partial trace. Most importantly,  $\text{Tr}\left[\sum_{i=1}^{M_j} h^{(i)}(l_j, g)\rho_{M_j}\right] = \text{Tr}\left[h(l_j, g)\rho_{M_j}^{(1)}\right]$ , see [Mic07] for more details.

**Theorem 2.5** (Occupation number of single-particle state). Let  $N \in \mathbb{N}$  and a general admissible sequence  $\{M_j\}_{j\in\mathbb{Z}}$  with associated lengths  $\{l_j\}_{j\in\mathbb{Z}}$ . Suppose  $\tilde{c} \left[(\mathcal{M}_N^>)^{1/3}\ell_N^> g\right]^{1/2} < 1/2$  (with the constant  $\tilde{c}$  from Lemma 2.3). Then for any  $j \in \mathbb{Z}$  with  $M_j \geq 1$ ,

$$\left(1 - \frac{n_{M_j}}{M_j}\right) \le \sqrt{7}c \cdot \left\{\ln\left[1 + e^{-2(\pi^2 + 3M_j l_j g)^{1/2}}\right]\right\}^{-1} M_j^{2/3} l_j g$$
(2.26)

with the constant c from Lemma 2.3.

*Proof.* We write  $E_1^{\text{QM}}(1, l_j, M_j g)$  for the second eigenvalue of the mean-field Hamiltonian  $h(1, l_j, M_j g)$ . After tracing (2.13) with the density matrix associated to the ground state of the Hamiltonian  $H(M_j, l_j, g)$  as in [SYZ12, proofs of Theorems 2.1 and 2.2], one obtains

$$E_0^{\text{QM}}(M_j, l_j, g) \ge (1 - \tau) \left[ n_{M_j} E^{\text{GP}}(1, l_j, M_j g) + (M_j - n_{M_j}) E_1^{\text{QM}}(1, l_j, M_j g) \right] + \tau M_j l_j^{-2} e^{\text{GP}}(M_j l_j g) - c M_j^{5/3} g l_j^{-1} \left( e^{\text{GP}}(M_j l_j g) \right)^{1/2} .$$
(2.27)

Note here that inequality (2.26) is slightly improved if compared to [SYZ12, Theorem 2.2] because we omit an estimation of  $(e^{\text{GP}}(M_j l_j g))^{1/2}$  in the last term of this inequality. Then, using the upper bound of Theorem 2.4 we obtain

$$\left(1 - \frac{n_{M_j}}{M_j}\right) \le c(1 - \tau)^{-1} \frac{\left(E^{\rm GP}(1, l_j, M_j g)\right)^{1/2}}{E_1^{\rm QM}(1, l_j, M_j g) - E^{\rm GP}(1, l_j, M_j g)} M_j^{2/3} g \tag{2.28}$$

$$\leq 2c \frac{\left(e^{\text{GP}}(M_j l_j g)\right)^{1/2}}{\left(\pi^2 + 3M_j l_j g\right)^{1/2}} \left\{ \ln\left(1 + \pi e^{-2\sqrt{\pi^2 + 3M_j l_j g}}\right) \right\}^{-1} M_j^{2/3} l_j g \qquad (2.29)$$

for all sufficiently large  $N \in \mathbb{N}$ . For the last estimate we used

$$E_1^{\text{QM}}(1, l, ng) - E^{\text{GP}}(1, l, ng) \ge \frac{\eta}{l^2} \ln\left(1 + \pi e^{-2\eta}\right)$$
(2.30)

with  $\eta = \sqrt{\pi^2 + 3nlg}$  and l, g > 0, see [KS85] and [SYZ12, (A.8)]. The statement then follows with

The statement then follows with

$$\frac{e^{\text{GP}}(M_j l_j g)}{\pi^2 + M_j l_j g} \le \frac{e^{\text{GP}}(M_j l_j g) - e^{\text{GP}}(0)}{M_j l_j g} + \frac{\pi^2}{\pi^2} \le \frac{7}{4} , \qquad (2.31)$$

where we used  $(e^{\text{GP}}(\kappa) - e^{\text{GP}}(0))/\kappa \leq 3/4$  [SYZ12, (31)] for  $\kappa > 0$  as well as  $e^{\text{GP}}(0) = \pi^2$ .

From now on we assume a partition of  $\Lambda_N = (-L_N/2, L_N/2)$  caused by a Poisson point process as described at the beginning. Moreover, we now allow the intensity of impurities  $\nu$  and the pair interaction g to vary with the particle number N. In order to account for variable impurity intensities, we do the following: We introduce a sequence  $(s_N)_{N \in \mathbb{N}} \subset \mathbb{R}$ such that  $s_N \leq 1$  and perform the scaling  $x_j(\omega) \mapsto s_N^{-1} x_j(\omega)$  and we set  $\nu_N := \nu s_N$ . If we consider a constant intensity  $\nu = const$ , we set  $s_N = 1$  for any  $N \in \mathbb{N}$ . We define

$$\tilde{l}_j := |\widetilde{W}_j| := \left| \left( s_N^{-1} x_j(\omega), s_N^{-1} x_{j+1}(\omega) \right) \cap \Lambda_N \right| .$$
(2.32)

We have  $\tilde{l}_j = s_N^{-1} l_j$  for any  $\tilde{l}_j > 0$  except possibly for the first and last subinterval  $\widetilde{W}_j$  within  $\Lambda_N$ . We write  $\tilde{k}_N$  for the number of the scaled subintervals within the window  $\Lambda_N$ . Then  $\lim_{N\to\infty} \tilde{k}_N/(L_N\nu_N) = 1$  almost surely by scaling. We also define  $\tilde{\ell}_N^> := \max{\{\tilde{l}_j : j \in \mathbb{Z}\}}$ .

**Remark 2.6.** Comparing Theorem 2.4 and 2.5 to [SYZ12, Theorems 2.1 and 2.2] one observes that we only require  $(\mathcal{M}_N^{>})^{1/3}\tilde{\ell}_N^{>}g_N$  instead of  $N^{1/3}L_Ng_N$  to converge to zero. This will allow us to consider stronger interactions in the sense that  $g_N$  converges more slowly to zero.

Note that the lengths  $|(x_j(\omega), x_{j+1}(\omega))|$  are exponentially distributed random variables with parameter  $\nu$  for any  $j \in \mathbb{Z} \setminus \{0\}$  [Kin93, Ch. 4]. For any  $N \in \mathbb{N}$  and any  $l, \tilde{\mu} > 0$  we define the function  $l \mapsto N_{g_N,\tilde{\mu}}(l)$  to be the unique minimizer of the Legendre transformation of the map  $\widetilde{N} \mapsto E^{\text{GP}}(\widetilde{N}, l, g_N)$  (see [SYZ12] for details). More explicitly,  $N_{g_N, \widetilde{\mu}}(l)$  is such that

$$E^{\rm GP}(N_{g_N,\widetilde{\mu}}(l),l,g_N) - \widetilde{\mu}N_{g_N,\widetilde{\mu}}(l) = \inf_{\widetilde{N} \ge 0} \left( E^{\rm GP}(\widetilde{N},l,g_N)) - \widetilde{\mu}\widetilde{N} \right) .$$
(2.33)

We remark that  $N_{g_N,\widetilde{\mu}}(l)$  obeys

$$\frac{2}{3}\frac{1}{lg_N}\left[\tilde{\mu}l^2 - \pi^2\right]_+ \le N_{g_N,\tilde{\mu}}(l) \le \frac{1}{lg_N}\left[\tilde{\mu}l^2 - \pi^2\right]_+$$
(2.34)

with  $[x]_+ := \max\{x, 0\}$ . Consequently,

$$\frac{2}{3}\frac{\widetilde{\mu}}{g_N\nu}\xi e^{-\pi\nu/\sqrt{\widetilde{\mu}}} \le \mathbb{E}\left[N_{g_N,\widetilde{\mu}}\right] = \int_0^\infty N_{g_N,\widetilde{\mu}}(l)\,\nu e^{-\nu l}\mathrm{d}l \le \frac{\widetilde{\mu}}{g_N\nu}\xi e^{-\pi\nu/\sqrt{\widetilde{\mu}}} \tag{2.35}$$

with (compare with [SYZ12, (45)])

$$1 \leq \xi := e^{\pi\nu/\sqrt{\tilde{\mu}}} \int_{\pi\nu/\sqrt{\tilde{\mu}}}^{\infty} \left( t - \left(\frac{\pi\nu}{\sqrt{\tilde{\mu}}}\right)^2 t^{-1} \right) e^{-t} dt \leq 2.$$
 (2.36)

Thus we have

$$\mathbb{E}[N_{g_N,\tilde{\mu}}] = \zeta_N \frac{\tilde{\mu}}{g_N \nu} \xi e^{-\pi\nu/\sqrt{\tilde{\mu}}}$$
(2.37)

with  $2/3 \leq \zeta_N \leq 1$  for any  $N \in \mathbb{N}$ . Since  $(\tilde{\mu}/g_N\nu_N)e^{-\pi\nu_N/\sqrt{\tilde{\mu}}}$  is a continuous function of  $\tilde{\mu}$  that converges to zero for  $\tilde{\mu} \searrow 0$  and to  $\infty$  for  $\tilde{\mu} \to \infty$ , for any  $N \in \mathbb{N}$  we are able to choose a  $\mu_N$  in such a way that

$$\mathbb{E}\left[N_{g_{N},\mu_{N}}(s_{N}^{-1}\cdot)\right] = \int_{\pi s_{N}/\sqrt{\mu_{N}}}^{\infty} N_{g_{N},\mu_{N}}(s_{N}^{-1}l) \nu e^{-\nu l} dl = \int_{\pi/\sqrt{\mu_{N}}}^{\infty} N_{g_{N},\mu_{N}}(l) \nu_{N} e^{-\nu_{N}l} dl$$

$$=: \mathbb{E}^{\nu_{N}}\left[N_{g_{N},\mu_{N}}\right] = \zeta_{N} \frac{\mu_{N}}{g_{N}\nu_{N}} \xi_{N} e^{-\pi\nu_{N}/\sqrt{\mu_{N}}} \stackrel{!}{=} \zeta_{N} \xi_{N} \rho \nu_{N}^{-1}$$
(2.38)

with

$$1 \le \xi_N := e^{\pi \nu_N / \sqrt{\mu_N}} \int_{\pi \nu_N / \sqrt{\mu_N}}^{\infty} \left( t - \left( \frac{\pi \nu_N}{\sqrt{\mu_N}} \right)^2 t^{-1} \right) e^{-t} dt \le 2.$$
 (2.39)

Hence

$$\frac{2}{3}\rho\nu_N^{-1} \le \mathbb{E}\left[N_{g_N,\mu_N}(s_N^{-1}\cdot)\right] \le 2\rho\nu_N^{-1} .$$
(2.40)

Similar to [SYZ12, Subsection. 3.3.3], we define the occupation numbers

$$M_j := \left\lceil N_{g_N,\mu_N}(\tilde{l}_j) \right\rceil \tag{2.41}$$

for all intervals  $\widetilde{W}_j$  with  $\tilde{l}_j > 0$  except for  $\widetilde{W}_0$  and the first and last interval within  $\Lambda_N$  and set  $M_j := 0$  else. We remark that we set  $M_0 = 0$  because, unlike  $|(x_j(\omega), x_{j+1}(\omega))|$  for any  $j \in \mathbb{Z} \setminus \{0\}$ , the length  $|(x_0(\omega), x_1(\omega))|$  is not exponentially distributed (cf. waiting time paradox [Kin93]). We also note that  $\{M_j\}_{j \in \mathbb{Z}}$  is a general admissible sequence.

We define

$$\lambda_N := \mathbb{P}\left(l > s_N \pi / \sqrt{\mu_N}\right) = \int_{s_N \pi / \sqrt{\mu_N}}^{\infty} \nu \mathrm{e}^{-\nu l} \,\mathrm{d}l = \mathrm{e}^{-\pi\nu_N / \sqrt{\mu_N}}$$
(2.42)

and note that  $\lambda_N$  is asymptotically equal to the fraction of intervals that are large enough to be, according to our choice (2.41), occupied by at least one particle. Moreover, after taking into account (2.38) and (2.37) we are able to establish the relationship

$$g_N = \rho^{-1} \mu_N \lambda_N = \rho^{-1} \pi^2 \frac{\lambda_N \nu_N^2}{[\ln \lambda_N]^2} .$$
 (2.43)

For the proof of the next theorem we need the following fact about the length  $\ell_N^>$  of the largest scaled subinterval.

**Lemma 2.7.** For  $1/N \ll \nu_N \lesssim 1$ , for any  $\kappa > 4$  and for almost any  $\omega \in \Omega$  there exists an  $\widetilde{N} \in \mathbb{N}$  such that for any  $N \geq \widetilde{N}$  the inequality

$$\tilde{\ell}_N^> \le \kappa \, \nu_N^{-1} \ln(N) \tag{2.44}$$

holds.

This has been proved in [Theorem 6.2, [SYZ12]] for  $\nu_N \equiv const.$  and can be extended to variable  $\nu_N$  by scaling. More precisely, for any  $\kappa > 4$  and almost any  $\omega \in \Omega$  there exists an  $\widetilde{N} \in \mathbb{N}$  such that for any  $N \geq \widetilde{N}$  it is  $\ell_N^> = \max\{|(x_j(\omega), x_{j+1}(\omega)) \cap \Lambda_N : j \in \mathbb{Z}\} \leq \kappa \nu^{-1} \ln(N)$  and therefore

$$\tilde{\ell}_N^{>} = \max\{|W_j| : j \in \mathbb{Z}\} = \max\{|(s_N^{-1}x_j(\omega), s_N^{-1}x_{j+1}(\omega)) \cap \Lambda_N| : j \in \mathbb{Z}\}$$
(2.45)

$$= s_N^{-1} \max\{ |(x_j(\omega), x_{j+1}(\omega)) \cap s_N \Lambda_N| : j \in \mathbb{Z} \} = s_N^{-1} \ell_{s_N N}^{1,>} \le \kappa \nu_N^{-1} \ln(N) .$$
(2.46)

Finally, we establish the main theorem of this section which plays a central role in proving BEC in the following section. For its proof we will use Theorem C.4 which itself is proved in the appendix.

**Theorem 2.8.** Let  $[\ln(N)]^4/N^{1-2\eta} \ll \nu_N \lesssim 1$  and  $g_N \ll \nu_N^2 N^{-\eta} [\ln(N)]^{-2}$  with an  $0 < \eta \leq 1/3$  be given. Then, with  $K(\eta) := 10 \cdot 5^2 \cdot \pi^2 \eta^{-2} + 1$  and  $c_1(\eta) := 90 E^{\text{GP}}(1, 1, K(\eta)) \eta^{-2}$  one obtains

$$\frac{1}{N} E_{\rm LS}^{\rm QM}(N, \Lambda_N, g_N) \le c_1(\eta) \frac{\nu_N^2}{(\ln(N))^2}$$
(2.47)

which holds for all but finitely many  $N \in \mathbb{N}$  almost surely.

Proof. We introduce the new interaction strength  $\hat{g}_N := \rho^{-1} [\pi^2 \nu_N^2 \eta^{-2} \ln(N)^{-2}] N^{-\eta}$  and using relation (2.43) one obtains  $\hat{\mu}_N = \pi^2 \nu_N^2 \eta^{-2} (\ln(N))^{-2}$ . Note that the hat characterizes the corresponding quantities associated with the pair  $(\hat{g}_N, \nu_N)$ . By assumption  $g_N \leq \hat{g}_N$ for all but finitely many  $N \in \mathbb{N}$  and  $\hat{\lambda}_N = N^{-\eta}$ . Furthermore, we choose  $\hat{M}_j$  as in (2.41).

Since  $\hat{\lambda}_N = N^{-\eta} \gg \ln(N) \ln(\nu_N N) (\nu_N N)^{-1/2}$  it is  $1/9 \le N^{-1} \sum_{j \in \mathbb{Z}} \hat{M}_j \le 9$  for all but finitely many  $N \in \mathbb{N}$  almost surely by Theorem C.4.

Defining  $\hat{n}_N := N / \sum_{j \in \mathbb{Z}} \hat{M}_j$  one clearly has  $\sum_{j \in \mathbb{Z}} \left[ \hat{n}_N \hat{M}_j \right] \ge N$  for all but finitely many  $N \in \mathbb{N}$  almost surely. Note that  $\left[ \hat{n}_N \hat{M}_j \right]$  is a general admissible sequence. By Theorem 2.4, because  $\hat{M}_j > 0$  if and only if  $\tilde{l}_j > \pi / \sqrt{\hat{\mu}_N}$  and with (2.6),

$$\frac{1}{N} E_{\rm LS}^{\rm QM}(N,\Lambda_N,g_N) \le \frac{1}{N} \sum_{j \in \mathbb{Z}} E_0^{\rm QM} \left( \left\lceil \hat{n}_N \hat{M}_j \right\rceil, \tilde{l}_j, g_N \right)$$

$$(2.48)$$

$$\leq \frac{1}{N} \sum_{j \in \mathbb{Z} : \tilde{l}_j > \pi/\sqrt{\hat{\mu}_N}} \frac{\left| \widehat{n}_N \hat{M}_j \right|}{\tilde{l}_j^2} E^{\text{GP}} \left( 1, 1, \left| \widehat{n}_N \hat{M}_j \right| \tilde{l}_j g_N \right)$$
(2.49)

$$\leq \frac{\mu_N}{\pi^2} \frac{1}{N} \sum_{j \in \mathbb{Z} : \tilde{l}_j > \pi/\sqrt{\hat{\mu}_N}} \left[ \widehat{n}_N \hat{M}_j \right] E^{\text{GP}} \left( 1, 1, \left[ \widehat{n}_N \hat{M}_j \right] \tilde{l}_j g_N \right)$$
(2.50)

for all but finitely many  $N \in \mathbb{N}$  almost surely.

Finally, we have  $\lceil \hat{n}_N \hat{M}_j \rceil \leq (9\hat{M}_j + 1) \leq 10\hat{M}_j$  and hence  $\lceil \hat{n}_N \hat{M}_j \rceil \tilde{l}_j g_N \leq 10\hat{M}_j \tilde{l}_j \hat{g}_N \leq 10\hat{\mu}_N (\tilde{\ell}_N^{>})^2 + 10\tilde{l}_j \hat{g}_N \leq 10 \cdot 5^2 \pi^2 \eta^{-2} + 1 = K(\eta)$  for any  $j \in \mathbb{Z}$ . This yields

$$\frac{\nu_N^2}{\eta^2 [\ln(N)]^2} \frac{1}{N} \sum_{j \in \mathbb{Z} : \tilde{l}_j > \pi/\sqrt{\hat{\mu}_N}} \left\lceil \hat{n}_N M_j \right\rceil E^{\text{GP}} \left( 1, 1, \left\lceil \hat{n}_N M_j \right\rceil \tilde{l}_j g_N \right)$$
(2.51)

$$\leq \frac{c_1(\eta)}{9} \frac{\nu_N^2}{[\ln(N)]^2} \frac{1}{N} \sum_{j \in \mathbb{Z}} \hat{M}_j \leq c_1(\eta) \frac{\nu_N^2}{[\ln(N)]^2}$$
(2.52)

for all but finitely many  $N \in \mathbb{N}$  almost surely.

#### 3 Main results

Generalized BEC in non-interacting Bose gases is said to occur (almost surely) if, for the sequence  $(\phi_N^j)_{j \in \mathbb{Z}}$  of eigenstates with respective occupation numbers  $n_{\phi_N^j}$  (see Definition 2.24)

and energies  $E_N^j \ge 0$  one has that

$$\lim_{\epsilon \searrow 0} \limsup_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}: E_N^j \le \epsilon} n_{\phi_N^j}$$
(3.1)

is (almost surely) larger than zero [vL82]. By analogy, we say that *generalized BEC* occurs almost surely in our model, the Luttinger–Sy model with interaction, if

$$\rho_0 := \lim_{\epsilon \searrow 0} \limsup_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}: E^{\mathrm{GP}}(1, \tilde{l}_j, N_j g_N) \le \epsilon} n_{N_j} > 0$$
(3.2)

is almost surely larger than zero. Here  $n_{N_j}$  is the number of particles occupying the singleparticle state  $\phi_{\tilde{l}_j,N_jg_N}^{\text{GP}}$ , see Definition (2.24). Consequently, we refer to the case

$$\lim_{\epsilon \searrow 0} \lim_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}: E^{\mathrm{GP}}(1, \tilde{l}_j, N_j g_N) \le \epsilon} n_{N_j} = 1$$
(3.3)

almost surely as almost sure *complete* generalized BEC.

In general, a single-particle state  $\phi$  with respective occupation number  $n_{\phi}$  is called almost surely macroscopically occupied if almost surely  $\lim \sup_{N\to\infty} n_{\phi}/N > 0$ . In this paper we refer to almost sure type-I (type-II) BEC if finitely (infinitely) many states  $\phi_{\tilde{l}_j,N_jg_N}^{\rm GP}$  are almost surely macroscopically occupied. If for almost any  $\omega \in \Omega$  one has  $\rho_0 > 0$  without any state  $\phi_{\tilde{l}_j,N_jg_N}^{\rm GP}$  being macroscopically occupied we speak of almost sure type-III BEC.

**Theorem 3.1.** [Generalized BEC] Assume that  $[\ln(N)]^4/N^{1-2\eta} \ll \nu_N \lesssim 1$  and  $g_N \ll \nu_N^2 N^{-\eta} [\ln(N)]^{-2}$  with  $0 < \eta \leq 1/3$ . Then almost sure complete generalized BEC occurs.

*Proof.* For any  $N \in \mathbb{N}$  let  $\{N_j\}_{j \in \mathbb{Z}}$  be a sequence of occupation numbers of the intervals with respect to the ground state of the Luttinger–Sy model, i.e. one has  $E_{\text{LS}}^{\text{QM}}(N, \Lambda_N, g_N) = \sum_{j \in \mathbb{Z}} E_0^{\text{QM}}(N_j, \tilde{l}_j, g_N)$ . We will show that

$$\rho_0 = \liminf_{\epsilon \searrow 0} \liminf_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z} : E^{\mathrm{GP}}(1, \tilde{l}_j, N_j g_N) \le \epsilon} n_{N_j} = 1$$
(3.4)

which then proves the statement.

Recall that according to Lemma 2.7, there exists a set  $\widehat{\Omega} \subset \Omega$  with  $\mathbb{P}(\widehat{\Omega}) = 1$  and the following property: for any  $\omega \in \widehat{\Omega}$  there exists an  $\widetilde{N}_1(\omega) \in \mathbb{N}$  such that for any  $N \geq \widetilde{N}_1(\omega)$  one has  $\widetilde{\ell}_N^> \leq 5\nu_N^{-1}\ln(N)$ . Moreover, for any  $\omega \in \Omega$  we define  $\widehat{N}_1(\omega)$  such that  $\widehat{N}_1(\omega) \geq \widetilde{N}_1(\omega)$  and  $g_N \leq \nu_N^2 N^{-\eta}[\ln(N)]^{-2}$  for any  $N \geq \widehat{N}_1(\omega)$ .

In a first step we show that for any  $\omega \in \widehat{\Omega}$ , any  $j \in \mathbb{Z}$  as well as any  $N > \widehat{N}_2(\omega) := \max\{3, \widehat{N}_1(\omega), (40(\tilde{c} + \sqrt{2}c)^{8/3} \|\nu_N\|_{\infty})^{1/\eta}\}$  with  $\|\nu_N\|_{\infty} = \max\{\nu_N : N \in \mathbb{N}\}$  the inequality

 $\gamma_j := N_j \tilde{l}_j g_N < [\ln(N)]^{1/2}$  implies  $N_j^{1/3} \tilde{l}_j g_N < 1/(4(\tilde{c} + \sqrt{2}c)^2)$ ,  $\tilde{c}$  and c from Lemma 2.3 : To do this let  $\omega \in \hat{\Omega}$  and  $N > \hat{N}_2(\omega)$  be given. If  $N_j \ge 4^{3/2} \tilde{c}^2 (\ln(N))^{3/4}$  then

$$N_j^{1/3}\tilde{l}_j g_N = N_j^{-2/3} N_j \tilde{l}_j g_N < N_j^{-2/3} [\ln(N)]^{1/2} \le \frac{1}{4(\tilde{c} + \sqrt{2}c)^2} .$$
(3.5)

On the other hand, if  $N_j < 4^{3/2} \tilde{c}^2 (\ln(N))^{3/4}$  then

$$N_j^{1/3} \tilde{l}_j g_N < 2\tilde{c}^{2/3} (\ln(N))^{1/4} \tilde{\ell}_N^> \nu_N^2 N^{-\eta} [\ln(N)]^{-2} \le \frac{1}{4(\tilde{c} + \sqrt{2}c)^2} .$$
(3.6)

We now prove that

$$\lim_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}} N_j = 1$$
(3.7)

almost surely: Suppose there exists a set  $\widetilde{\Omega} \subset \Omega$  with  $\mathbb{P}(\widetilde{\Omega}) > 0$  such that for any  $\omega \in \widetilde{\Omega}$ there is a  $\widetilde{d} > 0$  with  $\limsup_{N \to \infty} N^{-1} \sum_{j \in \mathbb{Z}: \gamma_j \ge (\ln(N))^{1/2}} N_j \ge \widetilde{d}$ . Then for any  $\omega \in \widetilde{\Omega} \cap \widehat{\Omega}$ , using Theorem 2.4 (with  $\epsilon = 1/2$ ), equation (2.10), inequality (2.19) while setting  $c_2 := \widetilde{d}/(8 \cdot 5^2)$ , we obtain

$$\frac{1}{N} E_{\mathrm{LS}}^{\mathrm{QM}}(N, \Lambda_N, g_N) \ge \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j \ge (\ln(N))^{1/2}} E_0^{\mathrm{QM}}(N_j, \tilde{l}_j, g_N)$$
(3.8)

$$\geq \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j \geq (\ln N)^{1/2}} \frac{1}{2} N_j \frac{N_j g_N}{2\tilde{l}_j} \geq c_2 \frac{\nu_N^2}{(\ln(N))^{3/2}}$$
(3.9)

for infinitely many  $N \in \mathbb{N}$ . However, since  $\mathbb{P}(\widetilde{\Omega} \cap \widehat{\Omega}) = \mathbb{P}(\widetilde{\Omega}) > 0$  this is in contradiction with Theorem 2.8.

Next, we prove that for any  $\omega \in \widehat{\Omega}$  and any  $N \ge \max\{\widehat{N}_2(\omega), e^{1/\pi^2}, (5\|\nu_N\|_{\infty})^{2/\eta}, \widehat{N}_3(\omega)\}$  one has

$$\frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}} n_{N_j} \ge \left(1 - \frac{c_3}{\ln(N)}\right) \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}} N_j$$
(3.10)

with  $c_3 := \sqrt{7}c (1 + e^{-2\pi}) e^{4\pi}$  and c > 0 as in Theorem 2.5: By Theorem 2.5 and using that  $\ln(1+x) \ge x/(1+x)$  for x > -1 one infers that if  $\gamma_j < (\ln(N))^{1/2}$  and  $N_j \ge 1$  for some  $j \in \mathbb{Z}$  then

$$\left(1 - \frac{n_{N_j}}{N_j}\right) \le \sqrt{7}c \cdot \left\{\ln\left[1 + e^{-2(\pi^2 + 3N_j\tilde{l}_jg_N)^{1/2}}\right]\right\}^{-1} N_j^{2/3}\tilde{l}_jg_N \tag{3.11}$$

$$\leq \sqrt{7}c \frac{1 + e^{-2\pi}}{e^{-2\sqrt{\pi^2 + 3\gamma_j}}} N_j^{-1/3} \gamma_j .$$
(3.12)

On the one hand, for any  $j \in \mathbb{Z}$  with  $\gamma_j \leq (\ln(N))^{-1}$  it is

$$\sqrt{7}c\left(1 + e^{-2\pi}\right)e^{2\sqrt{\pi^2 + 3\gamma_j}}N_j^{-1/3}\gamma_j \le \sqrt{7}c\left(1 + e^{-2\pi}\right)e^{4\pi}\frac{1}{\ln(N)} = \frac{c_3}{\ln(N)} .$$
(3.13)

On the other hand, for any  $j \in \mathbb{Z}$  with  $(\ln(N))^{-1} \leq \gamma_j < (\ln(N))^{1/2}$  we have

$$N_j N^{-\eta/2} \ge N_j \tilde{\ell}_N^> g_N \ge N_j \tilde{\ell}_j g_N = \gamma_j \ge \frac{1}{\ln(N)} , \qquad (3.14)$$

since  $\tilde{\ell}_N^> \leq 5\nu_N^{-1}\ln(N)$  and  $N \geq (5\|\nu_N\|_{\infty})^{2/\eta}$  imply  $g_N \leq (N^{\eta/2}\tilde{\ell}_N^>)^{-1}$ . Thus  $N_j \geq N^{\eta/2}/\ln(N)$  which leads to

$$\sqrt{7}c\left(1+e^{-2\pi}\right)e^{2\sqrt{\pi^2+3\gamma_j}}N_j^{-1/3}\gamma_j \le \frac{c_3}{\ln(N)}$$
(3.15)

for all  $N \geq \widehat{N}_3(\omega)$ ,  $\widehat{N}_3(\omega)$  some constant. Hence, we have  $(1 - n_{N_j}/N_j) \leq c_3/\ln(N)$  or, equivalently,  $(1 - [c_3/\ln(N)])N_j \leq n_{N_j}$  which implies (3.10).

Our last step is to show that for any  $\epsilon > 0$ 

$$\lim_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}, E_j \ge \epsilon} n_{N_j} = 0$$
(3.16)

almost surely with  $E_j := E^{\text{GP}}(1, \tilde{l}_j, N_j g_N)$ : We assume to the contrary that there exist an  $\epsilon > 0$  and a set  $\tilde{\Omega} \subset \Omega$  with  $\mathbb{P}(\tilde{\Omega}) > 0$  such that for any  $\omega \in \tilde{\Omega}$  there is a constant  $\tilde{r} > 0$  with  $\limsup_{N \to \infty} N^{-1} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}, E_j > \epsilon} n_{N_j} \geq \tilde{r}$ . Then for any  $\omega \in \tilde{\Omega}$  one also has

$$\limsup_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}, E_j > \epsilon} N_j \ge \limsup_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}, E_j > \epsilon} n_{N_j} \ge \tilde{r}$$
(3.17)

since  $N_j \ge n_{N_j}$  for any  $j \in \mathbb{Z}$ ,  $N \in \mathbb{N}$ . Hence, with Theorem 2.4 ( $\epsilon = 1/2$ ), equation (2.10) and inequality (2.19),

$$\frac{1}{N} E_{\mathrm{LS}}^{\mathrm{QM}}(N, \Lambda_N, g_N) \ge \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}, E_j > \epsilon} E_0^{\mathrm{QM}}(N_j, \tilde{l}_j, g_N)$$
(3.18)

$$\geq \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}, E_j > \epsilon} \frac{1}{2} E^{\mathrm{GP}}(N_j, \tilde{l}_j, g_N)$$
(3.19)

$$\geq \frac{\epsilon}{2} \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}, E_j > \epsilon} N_j \geq \frac{\epsilon}{2} \tilde{r}$$
(3.20)

for infinitely many  $N \in \mathbb{N}$  which is again a contradiction to Theorem 2.8. Note that the assumptions of Theorem 2.4 ( $\epsilon = 1/2$ ) are fulfilled according to (3.5) and (3.6).

Altogether we have shown that, using (3.16), (3.10) and (3.7) respectively,

=

$$\lim_{\epsilon \searrow 0} \liminf_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}: E_j \le \epsilon} n_{N_j} \ge \lim_{\epsilon \searrow 0} \liminf_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}, E_j \le \epsilon} n_{N_j}$$
(3.21)

$$= \liminf_{\epsilon \searrow 0} \liminf_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}} n_{N_j}$$
(3.22)

$$\geq \liminf_{N \to \infty} \left( 1 - \frac{c_3}{\ln(N)} \right) \frac{1}{N} \sum_{j \in \mathbb{Z}: \gamma_j < (\ln(N))^{1/2}} N_j = 1 \qquad (3.23)$$

almost surely.

**Remark 3.2.** Whereas in [SYZ12] type-I BEC in probability is shown in the regime where  $1/N \ll \nu_N \ll [\ln(N)]^3/N$ , see Appendix B, we are able to allow for  $\nu_N \equiv (\text{const.})$  which is mainly due to two reasons: Firstly, we consider BEC in the generalized sense. Having proved now complete BEC in a generalized sense we may replace the lim sup in Definition (3.2) by lim. Secondly, instead of the whole system length  $L_N$  and particle number N, we established Theorem 2.4 and Theorem 2.5 containing the occupation numbers  $N_j$  and lengths  $\tilde{l}_j$  of the individual intervals. This eventually enables us to use stronger interactions than in [SYZ12] in the sense that  $g_N$  converges to zero more slowly, see again Appendix B.

**Theorem 3.3** (Transition of condensation). Let  $0 < \eta \leq 1/3$  and  $[\ln(N)]^4/N^{1-2\eta} \ll \nu_N \lesssim 1$  be given. Then, 1) if  $g_N \equiv 0$  then almost surely exactly one single-particle state is macroscopically occupied and hence complete type-I BEC occurs, 2) if  $g_N \ll \nu_N N^{-1}[\ln(N)]^{-2}$  then BEC is almost surely of type I or II, 3) BEC is of type-III almost surely if  $\nu_N N^{-1}[\ln(N)]^{-1} \ll g_N \ll \nu_N^2 N^{-\eta}[\ln(N)]^{-2}$ .

*Proof.* Assume that  $\{N_j\}_{j\in\mathbb{Z}}$  are occupation numbers of the intervals with respect to the ground state of the Luttinger–Sy model, i.e.  $E_{\text{LS}}^{\text{QM}}(N, \Lambda_N, g_N) = \sum_{j\in\mathbb{Z}} E_0^{\text{QM}}(N_j, \tilde{l}_j, g_N)$  for any  $N \in \mathbb{N}$ . As before,  $n_{N_j}$  denotes the number of particles occupying the single-particle state  $\phi_{\tilde{l}_j,N_jg_N}^{\text{GP}}$ .

The first part of the theorem regarding the case  $g_N \equiv 0$  follows readily since almost surely there is only one largest interval and, since the temperature is zero, all particles occupy the ground state corresponding to this length.

Next, we treat the case  $g_N \ll \nu_N N^{-1}[\ln(N)]^{-2}$ : According to Corollary C.8 there exists, for any  $\eta' > 0$  and any  $C_3 > 2e^{\nu/\rho}$ , an  $\widetilde{N}(\eta') \in \mathbb{N}$  such that for  $N \ge \widetilde{N}(\eta')$  it is

$$\mathbb{P}(\Omega_1) > 1 - \eta' \tag{3.24}$$

with

$$\Omega_1 := \left\{ \omega \in \Omega : \tilde{\ell}_N^{>} \ge (1/2)\nu_N^{-1}\ln(\nu_N N), \ \tilde{\ell}_N^{>} - \tilde{\ell}_N^{>,\lceil 2\nu C_3/(C_1\eta')\rceil+1} > \nu_N^{-1}\ln(C_3/(2e^{\nu/\rho})) \right\}$$

and  $C_1 := -\nu/[4\ln(\eta'/2)]$ . Also,  $\tilde{l}_N^{>,\lceil 2\nu C_3/(C_1\eta')\rceil+1}$  is the  $(\lceil 2\nu C_3/(C_1\eta')\rceil+1)$ th largest length of  $\{\tilde{l}_j\}_{j\in\mathbb{Z}}$ . Note that if  $\omega \in \Omega_1$  then there are at most  $(\lceil 2\nu C_3/(C_1\eta')\rceil+1)$  many intervals

that have a length larger than  $\hat{l}_N := \tilde{\ell}_N^2 - \nu_N^{-1} \ln(C_3/(2e^{\nu/\rho}))$ . For convenience we define  $E_0^{\text{QM}}(0,0,g) := 0$  for  $g \ge 0$ . Furthermore, for any  $N \in \mathbb{N}$  and  $j \in \mathbb{Z}$  we write  $\hat{N}^{(1)} := \sum_{j \in \mathbb{Z}: \tilde{l}_j \ge \hat{l}_N} N_j$  and  $\hat{N}^{(2)} := \sum_{j \in \mathbb{Z}: \tilde{l}_j < \hat{l}_N} N_j$ . Let  $\tilde{N}_N^>$  denote the particle number in the largest interval.

Now, let  $\omega \in \Omega_1$  be given. Then

$$E_{\rm LS}^{\rm QM}(N,\Lambda_N,g_N) = \sum_{j\in\mathbb{Z}} E_0^{\rm QM}(N_j,\tilde{l}_j,g_N)$$

$$\geq \sum_{j\in\mathbb{Z}:\tilde{l}_j\geq \hat{l}_N,\tilde{l}_j\neq \tilde{\ell}_N^>} E_0^{\rm QM}(N_j,\tilde{l}_j,g_N) + E_0^{\rm QM}(\tilde{N}_N^>,\tilde{\ell}_N^>,0) + E_0^{\rm QM}(\hat{N}^{(2)},\hat{l}_N,0)$$

$$= \sum_{j\in\mathbb{Z}:\tilde{l}_j\geq \hat{l}_N,\tilde{l}_j\neq \tilde{\ell}_N^>} E_0^{\rm QM}(N_j,\tilde{l}_j,g_N) + \tilde{N}_N^> \frac{\pi^2}{(\tilde{\ell}_N^>)^2} + \hat{N}^{(2)} \frac{\pi^2}{\hat{l}_N^2} .$$
(3.25)

On the other hand, with  $\phi_0$  the ground state of  $H(1,1,0), C := \int_0^1 |\phi_0|^4$ , we can employ a simple variational argument to obtain (see also proof of Theorem 2.4)

$$E_{0}^{\text{QM}}(\widetilde{N}_{N}^{>} + \widehat{N}^{(2)}, \widetilde{\ell}_{N}^{>}, g_{N}) \leq E^{\text{GP}}(\widetilde{N}_{N}^{>} + \widehat{N}^{(2)}, \widetilde{\ell}_{N}^{>}, g_{N})$$

$$\leq \frac{(\widetilde{N}_{N}^{>} + \widehat{N}^{(2)})}{(\widetilde{\ell}_{N}^{>})^{2}} \mathcal{E}^{\text{GP}}(1, 1, (\widetilde{N}_{N}^{>} + \widehat{N}^{(2)})\widetilde{\ell}_{N}^{>}g_{N})[\phi_{0}]$$

$$= \frac{(\widetilde{N}_{N}^{>} + \widehat{N}^{(2)})}{(\widetilde{\ell}_{N}^{>})^{2}} \pi^{2} + \frac{(\widetilde{N}_{N}^{>} + \widehat{N}^{(2)})^{2}g_{N}}{2\widetilde{\ell}_{N}^{>}} C.$$
(3.26)

According to (2.44),  $[\tilde{\ell}_N^> - \kappa \nu_N^{-1} \ln(N)]_+$  converges almost surely to zero and hence in probability to zero for any  $\kappa > 4$ . Therefore, for any  $\eta' > 0$  there exists an  $\widehat{N}(\eta') \in \mathbb{N}$  such that for any  $N \ge \widehat{N}(\eta')$  it is  $\mathbb{P}(\Omega_2) := \mathbb{P}(\widetilde{\ell}_N^> < 5\nu_N^{-1}\ln(N)) > 1 - \eta'$ . Let  $\epsilon, \eta', \eta > 0$  and  $\omega \in \Omega_1 \cap \Omega_2$  be given. Note that  $\mathbb{P}(\Omega_1 \cap \Omega_2) > 1 - 2\eta'$  for any

 $N \geq \max\{\widetilde{N}(\eta'), \widehat{N}(\eta')\}$ . Moreover, let  $N \geq \max\{\widetilde{N}(\eta'), \widehat{N}(\eta')\}$  be such that

$$g_N \le \frac{2\pi^2 \eta}{C} \frac{\nu_N^{-1} \ln(C_3/(2e^{\nu/\rho}))}{(5\nu_N^{-1}\ln(N))^2} \frac{1}{N}$$
(3.27)

and

$$E^{\rm GP}(1,\hat{l}_N,Ng_N) \le \frac{1}{\hat{l}_N^2} \mathcal{E}^{\rm GP}(1,1,N\hat{l}_Ng_N)[\phi_0] = \frac{1}{\hat{l}_N^2} \left(\pi^2 + \frac{N\hat{l}_Ng_N}{2}C\right) \le \epsilon$$
(3.28)

hold.

Now we assume that  $\widehat{N}^{(2)}/N > \eta$  which will lead to a contradiction with the above. Since  $\widetilde{N}_N^> + \widehat{N}^{(2)} \leq N$  we conclude that, for  $N \in \mathbb{N}$  large enough,

$$\frac{N}{\widehat{N}^{(2)}} \frac{(\widetilde{N}_N^{>} + \widehat{N}^{(2)})g_N}{2\pi^2 \widetilde{\ell}_N^{>}} C < \frac{1}{2\pi^2 \eta} \frac{Ng_N}{\widetilde{\ell}_N^{>}} C \le \frac{\nu_N^{-1} \ln(C_3/(2\mathrm{e}^{\nu/\rho}))}{(\widetilde{\ell}_N^{>})^3} \le \frac{(\widetilde{\ell}_N^{>})^2 - (\widehat{\ell}_N)^2}{\widehat{\ell}_N^2 (\widetilde{\ell}_N^{>})^2} \tag{3.29}$$

$$=\frac{1}{\hat{l}_N^2} - \frac{1}{(\tilde{\ell}_N^>)^2}$$
(3.30)

and therefore

$$\widehat{N}^{(2)} \frac{\pi^2}{(\widetilde{\ell}_N^{>})^2} + \frac{(\widetilde{N}_N^{>} + \widehat{N}^{(2)})^2 g_N}{2\widetilde{\ell}_N^{>}} C < \widehat{N}^{(2)} \frac{\pi^2}{\widehat{l}_N^2} .$$
(3.31)

Comparing this with (3.25) and (3.26) we arrive at a contradiction since  $E_{\text{LS}}^{\text{QM}}(N, \Lambda_N, g_N)$  is minimal.

From the assumptions  $\lim_{N\to\infty} N\tilde{\ell}_N^> g_N = 0$  almost surely and with Theorem (2.5),  $\max_{j\in\mathbb{Z}:N_j>0}\{1 - n_{N_j}/N_j\}$  converges to zero almost surely. Therefore, for any  $\epsilon, \eta, \eta' > 0$ and  $C_3 > 2e^{\nu/\rho}$  there exists an  $\tilde{N} \in \mathbb{N}$  such that for any  $N \geq \tilde{N}$  it is, with  $E_j = E^{\mathrm{GP}}(1, \tilde{l}_j, N_j g_N)$ ,

$$\mathbb{P}\left(\left\{\left|\frac{1}{N}\sum_{j\in\mathbb{Z}:\tilde{l}_{j}\geq\hat{l}_{N},E_{j}\leq\epsilon}N_{j}-1\right|\leq\eta\right\}\cap\Omega_{1}\cap\Omega_{2}\right)=\mathbb{P}\left(\Omega_{1}\cap\Omega_{2}\right)>1-2\eta'\qquad(3.32)$$

and

$$\mathbb{P}\left(\left\{\left|\frac{1}{N}\sum_{j\in\mathbb{Z}:\tilde{l}_{j}\geq\hat{l}_{N},E_{j}\leq\epsilon}\left(N_{j}-n_{N_{j}}\right)\right|\leq\eta\right\}\cap\Omega_{1}\cap\Omega_{2}\right)$$
(3.33)

$$\geq \mathbb{P}\left(\left\{\max_{j\in\mathbb{Z}:N_j>0}\left\{1-\frac{n_{N_j}}{N_j}\right\}\frac{1}{N}\sum_{j\in\mathbb{Z}:\tilde{l}_j\geq \hat{l}_N, E_j\leq\epsilon}N_j\leq\eta\right\}\cap\Omega_1\cap\Omega_2\right)>1-3\eta'.$$
 (3.34)

Hence, by the previous two inequalities,

$$\mathbb{P}\left(\frac{n_{\mathcal{N}_{N}^{>}}}{N} \geq \frac{1}{2} \frac{1-2\eta}{\lceil 2\nu C_{3}/(C_{1}\eta')\rceil+1}\right) \geq \mathbb{P}\left(\max_{j\in\mathbb{Z}:\tilde{l}_{j}\geq\tilde{l}_{N},E_{j}\leq\epsilon}\left\{\frac{n_{N_{j}}}{N}\right\} \geq \frac{1-2\eta}{\lceil 2\nu C_{3}/(C_{1}\eta')\rceil+1}\right) \\ \geq \mathbb{P}\left(\left\{\left|\frac{1}{N}\sum_{j\in\mathbb{Z}:\tilde{l}_{j}\geq\tilde{l}_{N},E_{j}\leq\epsilon}n_{N_{j}}-1\right|\leq 2\eta\right\}\cap\Omega_{1}\cap\Omega_{2}\right) \tag{3.35}$$

$$\geq \mathbb{P}\left(\left\{\left|\frac{1}{N}\sum_{j\in\mathbb{Z}:\tilde{l}_{j}\geq\hat{l}_{N},E_{j}\leq\epsilon}\left(n_{N_{j}}-N_{j}\right)\right|\leq\eta\right\}\cap\left\{\left|\frac{1}{N}\sum_{j\in\mathbb{Z}:\tilde{l}_{j}\geq\hat{l}_{N},E_{j}\leq\epsilon}N_{j}-1\right|\leq\eta\right\}\right.\\\left.\cap\Omega_{1}\cap\Omega_{2}\right)>1-5\eta'.$$

Now, suppose there exists a  $0 < C \leq 1$  such that  $\mathbb{P}(\lim_{N \to \infty} n_{\mathcal{N}_N^>}/N = 0) \geq C > 0$ . Then

$$\limsup_{N \to \infty} \mathbb{P}\left(\frac{n_{\mathcal{N}_{N}^{>}}}{N} \geq \frac{1}{4} \frac{1}{\lceil 12\nu C_{3}/(C_{1}C) \rceil + 1}\right) \leq \mathbb{P}\left(\limsup_{N \to \infty} \frac{n_{\mathcal{N}_{N}^{>}}}{N} \geq \frac{1}{4} \frac{1}{\lceil 12\nu C_{3}/(C_{1}C) \rceil + 1}\right)$$
$$\leq \mathbb{P}\left(\neg\left(\lim_{N \to \infty} \frac{n_{\mathcal{N}_{N}^{>}}}{N} = 0\right)\right) \leq 1 - C .$$

$$(3.36)$$

However, we have shown above that, with  $\eta = 1/4$  and  $\eta' = C/6$ , there exists an  $\widetilde{N} \in \mathbb{N}$  such that for any  $N \geq \widetilde{N}$  it is

$$\mathbb{P}\left(\frac{n_{\mathcal{N}_{N}^{>}}}{N} \ge \frac{1}{4} \frac{1}{\lceil 12\nu C_{3}/(C_{1}C) \rceil + 1}\right) \ge 1 - \frac{5C}{6} , \qquad (3.37)$$

see (3.35). Comparing (3.37) with (3.36) one arrives at a contradiction. Hence  $\mathbb{P}(\lim_{N\to\infty} n_{\mathcal{N}_N^>}/N=0) = 0$  and consequently  $\mathbb{P}(\limsup_{N\to\infty} n_{\mathcal{N}_N^>}/N>0) = 1$ . Now we prove the last part of the theorem: We assume to the contrary that there

Now we prove the last part of the theorem: We assume to the contrary that there exists a set  $\widetilde{\Omega} \subset \Omega$  with  $\mathbb{P}(\widetilde{\Omega}) > 0$  such that for any  $\omega \in \widetilde{\Omega}$  there is a  $\widetilde{c} > 0$  with  $\limsup_{N\to\infty} \max_{j\in\mathbb{Z}} \{n_{N_j}/N\} \geq \widetilde{c}$ .

Since  $N_j \ge n_{N_j}$  for any  $N \in \mathbb{N}$  and  $j \in \mathbb{Z}$  one also has  $\limsup_{N \to \infty} \{\mathcal{N}_N^>/N\} \ge \tilde{c}$ . Thus with Theorem 2.4 ( $\epsilon = 1/2$ ), inequality (2.19) and  $\hat{g}_N := \min\{g_N, \nu_N N^{-1/3}[\ln(N)]^{-2}\}$ , for any  $\omega \in \tilde{\Omega} \cap \hat{\Omega}$  (see beginning of proof of Theorem 3.1 for definition of the set  $\hat{\Omega}$ ) it holds

$$\frac{1}{N} E_{\mathrm{LS}}^{\mathrm{QM}}(N, \Lambda_N, g_N) \geq \frac{1}{N} E_0^{\mathrm{QM}}(\mathcal{N}_N^>, \tilde{\ell}_N^>, g_N) \\
\geq \frac{1}{N} E_0^{\mathrm{QM}}(\mathcal{N}_N^>, \tilde{\ell}_N^>, \hat{g}_N) \\
\geq \frac{1}{2} \frac{1}{N} E^{\mathrm{GP}}(\mathcal{N}_N^>, \tilde{\ell}_N^>, \hat{g}_N) \\
\geq \frac{1}{4} \frac{1}{N} \frac{(\mathcal{N}_N^>)^2 \widehat{g}_N}{\tilde{\ell}_N^>} \geq \frac{1}{20} \frac{1}{N} \frac{((\tilde{c}/2)N)^2 \widehat{g}_N}{\nu_N^{-1} \ln(N)}$$
(3.38)

for infinitely many  $N \in \mathbb{N}$ . Note that we inserted the length of the largest interval in the first step which is possible since the energy goes down when increasing the length.

However, since  $\widehat{g}_N \gg \nu_N(\ln(N))^{-1}N^{-1}$  and  $\mathbb{P}(\widehat{\Omega} \cap \widehat{\Omega}) = \mathbb{P}(\widehat{\Omega}) > 0$ , this contradicts Theorem 2.8. Therefore almost surely  $\lim_{N\to\infty} \max_{j\in\mathbb{Z}} \{n_{N_j}/N\} = 0$  and hence the statement follows with Theorem 3.1.

**Theorem 3.4** (Particle density in largest interval). Let  $[\ln(N)]^4/N^{1-2\eta} \ll \nu_N \lesssim 1$  and  $g_N \ll \nu_N^2 N^{-\eta} [\ln(N)]^{-2}$  with an  $0 < \eta \leq 1/3$  be given. Moreover, for any  $N \in \mathbb{N}$  let  $\{N_j\}_{j\in\mathbb{Z}}$  be occupation numbers of the intervals with respect to the ground state of the Luttinger-Sy model, i.e.,  $E_{\text{LS}}^{\text{QM}}(N, \Lambda_N, g_N) = \sum_{j\in\mathbb{Z}} E_0^{\text{QM}}(N_j, \tilde{l}_j, g_N)$ .

Then, with  $c_1(\eta)$  as in Theorem 2.8 one has

$$\frac{\mathcal{N}_N^>}{\tilde{\ell}_N^>} \le \sqrt{8c_1(\eta)} \frac{\nu_N^{3/2}}{\ln(N)} \frac{1}{[\ln(\nu_N N)]^{1/2}} \left( \min\left\{g_N, \frac{\nu_N}{N^{1/3}[\ln(N)]^2}\right\} \right)^{-1/2} N^{1/2}$$
(3.39)

for all but finitely many  $N \in \mathbb{N}$  almost surely. Furthermore, almost surely and for all but finitely many  $N \in \mathbb{N}$  one has

$$\frac{\mathcal{N}_{N}^{>}}{\tilde{\ell}_{N}^{>}} \leq \sqrt{8c_{1}(\eta)} \frac{\nu_{N}}{[\ln(\nu_{N}N)]^{1/2}} N^{3/5+\delta}$$
(3.40)

for any  $\delta > 0$  if  $\eta < 1/6$  and  $g_N \gg \nu_N N^{-1/6} [\ln(N)]^{-2}$ .

Proof. We define  $\widehat{g}_N(\beta) := \nu_N N^{-\beta} [\ln(N)]^{-2}$  for any  $\beta > 0$ . Let  $\omega \in \Omega$  and  $\widetilde{N} = \widetilde{N}(\omega) \in \mathbb{N}$  such that the upper bound (2.44) for the length of the largest interval with  $\kappa = 5$ , inequality  $N^{1/3} \widetilde{\ell}_N^2 \widehat{g}_N(1/3) < (2 \max\{\sqrt{2}c, \widetilde{c}\})^{-2}$  with constants  $c, \widetilde{c} > 0$  from Lemma 2.3, and inequality (2.47) of Theorem 2.8 hold for any  $N \geq \widetilde{N}$ . Then, with Theorem 2.8, Theorem 2.4 ( $\epsilon = 1/2$ ), equation (2.10), and inequality (2.19) we have

$$c_1(\eta) \frac{\nu_N^2}{[\ln(N)]^2} \ge \frac{1}{N} E_{\rm LS}^{\rm QM}(N, \Lambda_N, g_N)$$
 (3.41)

$$\geq \frac{1}{N} E_0^{\text{QM}}(\mathcal{N}_N^>, \tilde{\ell}_N^>, g_N) \tag{3.42}$$

$$\geq \frac{1}{N} E_0^{\text{QM}} \left( \mathcal{N}_N^>, \tilde{\ell}_N^>, \min\left\{ g_N, \hat{g}_N(1/3) \right\} \right)$$
(3.43)

$$\geq \frac{1}{2} \frac{1}{N} E^{\mathrm{GP}} \left( \mathcal{N}_{N}^{>}, \tilde{\ell}_{N}^{>}, \min \left\{ g_{N}, \widehat{g}_{N}(1/3) \right\} \right)$$
(3.44)

$$\geq \frac{1}{4} \frac{1}{N} \frac{(\mathcal{N}_N^{>})^2}{\tilde{\ell}_N^{>}} \min \left\{ g_N, \hat{g}_N(1/3) \right\}$$
(3.45)

and therefore

$$\mathcal{N}_{N}^{>} \leq \sqrt{4c_{1}(\eta)} \frac{\nu_{N}}{\ln(N)} \left( \frac{\tilde{\ell}_{N}^{>}}{\min\{g_{N}, \hat{g}_{N}(1/3)\}} \right)^{1/2} N^{1/2}$$
(3.46)

for any  $N \geq \tilde{N}$ . The first part of the statement then follows taking into account that almost surely  $\tilde{\ell}_N^> \geq (1/2)\nu_N^{-1}\ln(\nu_N N)$  for all but finitely many  $N \in \mathbb{N}$ , see Theorem C.5.

Recall that  $g_N \gg \nu_N N^{-1/6} [\ln(N)]^{-2}$  by assumption. We then define  $\beta_0 := 1/3$  and  $\beta_n := (1 + \beta_{n-1})/6$  for any  $n \in \mathbb{N}$ . We now show by induction that for any  $n \in \mathbb{N}$ ,  $(\mathcal{N}_N^>)^{1/3} \tilde{\ell}_N^> \hat{g}_N(\beta_n)$  converges to zero almost surely: Firstly, one has  $\mathcal{N}_N^> \leq N$  for any  $N \in \mathbb{N}$ .

and  $N^{1/3}\tilde{\ell}_N^> \widehat{g}_N(\beta_0)$  converges to zero almost surely. Next, we assume that for arbitrary  $n \in \mathbb{N}$ ,  $(\mathcal{N}_N^>)^{1/3}\tilde{\ell}_N^> \widehat{g}_N(\beta_n)$  converges to zero almost surely. Then for almost any  $\omega \in \Omega$  there exists an  $\tilde{N}_n = \tilde{N}_n(\omega) \in \mathbb{N}$  such that  $\widehat{g}_N(\beta_n) \leq g_N$ ,  $(\mathcal{N}_N^>)^{1/3}\tilde{\ell}_N^> \widehat{g}_N(\beta_n) < (2\max\{\sqrt{2}c,\tilde{c}\})^{-2}$ , and inequality (2.47) of Theorem 2.8 holds for any  $N \geq \tilde{N}_n$ . We therefore can conclude that

$$c_{1}(\eta) \frac{\nu_{N}^{2}}{[\ln(N)]^{2}} \geq \frac{1}{N} E_{\text{LS}}^{\text{QM}}(N, \Lambda_{N}, g_{N})$$

$$\geq \frac{1}{N} E_{0}^{\text{QM}}(\mathcal{N}_{N}^{>}, \tilde{\ell}_{N}^{>}, g_{N})$$

$$\geq \frac{1}{N} E_{0}^{\text{QM}}(\mathcal{N}_{N}^{>}, \tilde{\ell}_{N}^{>}, \hat{g}_{N}(\beta_{n}))$$

$$\geq \frac{1}{2} \frac{1}{N} E^{\text{GP}}(\mathcal{N}_{N}^{>}, \tilde{\ell}_{N}^{>}, \hat{g}_{N}(\beta_{n})) \geq \frac{1}{4} \frac{1}{N} \frac{(\mathcal{N}_{N}^{>})^{2} \hat{g}_{N}(\beta_{n})}{\tilde{\ell}_{N}^{>}}$$

$$(3.47)$$

or, equivalently,

$$\mathcal{N}_{N}^{>} \leq \sqrt{4c_{1}(\eta)} (\nu_{N} \tilde{\ell}_{N}^{>})^{1/2} N^{(1+\beta_{n})/2}$$
(3.48)

for any  $N \geq \widetilde{N}_n$ . Hence,  $(\mathcal{N}_N^{>})^{1/3} \widetilde{\ell}_N^{>} \widehat{g}_N(\beta_{n+1})$  converges to zero almost surely.

Lastly, note that  $(\beta_n)_{n \in \mathbb{N}_0}$  converges to 1/5. For arbitrary  $\delta > 0$  we choose an  $n \in \mathbb{N}$  such that  $\beta_n \leq 1/5 + 2\delta$ . Hence,

$$\mathcal{N}_{N}^{>} \leq \sqrt{4c_{1}(\eta)} (\nu_{N}\tilde{\ell}_{N}^{>})^{1/2} N^{(1+\beta_{n})/2} \leq c_{4}(\eta) \nu_{N} N^{(1+1/5+2\delta)/2} \tilde{\ell}_{N}^{>} [\ln(\nu_{N}N)]^{-1/2}$$
(3.49)

for all but finitely many  $N \in \mathbb{N}$  almost surely.

Note that Theorem 3.4 implies the following: For interactions  $g_N \gg \nu_N N^{-1/6} [\ln(N)]^{-2}$ , the particle density in the largest interval is almost surely bounded (actually converging to zero) in case of  $[\ln(N)]^4/N^{1-2\eta} \ll \nu_N \lesssim N^{-3/5-\delta}$  for any  $0 < \eta < 1/6$ ,  $0 < \delta < 2/5 - 2\eta$  and it is asymptotically bounded by  $\nu_N N^{3/5+\delta}$  for any  $\delta > 0$  in the case of  $[\ln(N)]^4/N^{1-2\eta} \ll$  $\nu_N \lesssim 1$ . In particular, note that for  $[\ln(N)]^4/N^{1-2\eta} \ll \nu_N \lesssim N^{-3/5-\delta}$  the particle density in the largest interval diverges in the non-interacting model, i.e., if  $g_N \equiv 0$ . Hence, we conclude that the repulsive interaction between the particles is pivotal.

#### A Notation

For two real-valued sequences  $(a_N)_{N \in \mathbb{N}}$ ,  $(b_N)_{N \in \mathbb{N}}$  with all elements positive and unequal to zero we write  $a_N \sim b_N$  if there exist constants c, C > 0 such that  $c \leq a_N/b_N \leq C$  for all but finitely many  $N \in \mathbb{N}$ . We also write  $a_N \ll b_N$  if  $a_N/b_N$  tends to zero. We combine these two possibilities through writing  $a_N \leq b_N$ , meaning either  $a_N \sim b_N$  or  $a_N \ll b_N$ . Moreover, we also write  $a_N \sim b_N$  in the case that  $a_N = b_N = 0$  for all but finitely many  $N \in \mathbb{N}$  to simplify the notation.

## B On the connection to the results of [SYZ12]

We first note that, in contrast to the model discussed in this paper, in [SYZ12] the unit interval is the fixed one-particle configuration space. However, by an appropriate scaling as discussed in [SYZ13, Sec. 4.4] and at the end of this section, the results can be translated into each other.

The Hamiltonian in [SYZ12] is formally given by

$$H = \sum_{i=1}^{N} \left( -\partial_{z_i}^2 + V_{\omega}(z_i) \right) + \frac{\gamma}{N} \sum_{i < j} \delta(z_i - z_j)$$
(B.1)

where

$$V_{\omega}(z) := \sigma \sum_{j} \delta(z - z_{j}^{\omega}) \tag{B.2}$$

with  $\gamma \geq 0$  the coupling parameter for the interaction among the particles,  $\nu$  the density of scatterers  $\{z_j^{\omega}\}, \sigma$  the strength of the scattering potential (note that  $\sigma = \infty$  in our model), and m the number of scatterers in the unit interval. In [SYZ12, Theorem 2.2] which is subsequently used to prove BEC they established the estimate

$$\left(1 - \frac{N_0}{N}\right) \le (const.) \frac{e_0}{e_1 - e_0} N^{-1/3} \min\{\gamma, \gamma^{1/2}\}$$
(B.3)

with N the total number of particles and  $N_0$  the number of particles occupying the minimizer of the Gross-Pitaevskii functional. In addition, [SYZ12, Lemma 5.1] provides the lower bound  $e_1 - e_0 \ge \eta \ln(1 + \pi e^{-2\eta})$  with  $\eta = \sqrt{\pi^2 + 3m\sigma + 3\gamma}$ .

Now, with  $\gamma \ge 1$ ,  $e_0 = E^{\text{GP}}(1, 1, \gamma) \ge \gamma/2$ , and  $\ln(1+x) \le x$  for x > 0 it follows

$$\frac{e_0}{\eta \ln(1 + \pi e^{-2\eta})} N^{-1/3} \min\{\gamma, \gamma^{1/2}\} \gtrsim \frac{\gamma^{3/2}}{\eta} N^{-1/3} e^{2\eta} .$$
(B.4)

In the case of  $\gamma \geq m\sigma$  it is

$$\frac{\gamma^{3/2}}{\eta} N^{-1/3} \mathrm{e}^{2\eta} \gtrsim \frac{\gamma^{3/2}}{\sqrt{\gamma}} N^{-1/3} \mathrm{e}^{2\sqrt{\gamma}} . \tag{B.5}$$

If  $\gamma \geq [\ln(N)]^2$ , this converges to infinity and therefore (B.3) does not prove BEC. On the other hand, if  $\gamma \leq m\sigma$  then

$$\frac{\gamma^{3/2}}{\eta} N^{-1/3} \mathrm{e}^{2\eta} \gtrsim \frac{1}{\sqrt{m\sigma}} N^{-1/3} \mathrm{e}^{2\sqrt{m\sigma}} , \qquad (B.6)$$

which, for  $m\sigma \geq [\ln(N)]^2$ , again converges to infinity. Hence, in order to establish BEC with the estimate (B.3) it must hold that  $m\sigma \leq [\ln(N)]^2$  and therefore  $\gamma \leq [\ln(N)]^2$ .

Furthermore, the assumptions in [SYZ12, Theorem 3.1, Lemma 3.2, Lemma 3.3] are such that  $\nu \to \infty$ ,  $\gamma \to \infty$ , and  $\gamma \gg \nu/(\ln \nu)^2$ . Therefore

$$[\ln(N)]^2 \ge \gamma \gg \nu/(\ln\nu)^2 \gtrsim \nu^{1-\epsilon}$$
(B.7)

for any  $\epsilon > 0$ . This however implies that  $\nu$  must grow slower than  $[\ln(N)]^3$ .

We are now in position to compare with our results: as described in [SYZ13, Sec. 4.4] the above translates to a density  $\nu_N \ll [\ln(N)]^3/N$  when working on the interval  $(-L_N/2, L_N/2)$ . Furthermore, since the condition  $\nu \gg 1$  implies  $\nu_N \gg 1/N$  one concludes that  $1/N \ll \nu_N \ll \ln(N)^3/N$  is a necessary requirement in [SYZ12] to prove BEC. Also,  $\gamma = L_N N g_N \leq [\ln(N)]^2$  implies the requirement  $g_N \lesssim [\ln(N)]^2/N^2$ .

Hence, comparing with Theorem 3.1, Theorem 3.3 and Theorem 3.4 we see that we are able to allow for densities  $\nu_N$  which converge to zero more slowly or even are constant.

#### C Miscellaneous results

Let  $(\Lambda_N)_{N \in \mathbb{N}}$  be an arbitrary sequence of intervals in  $\mathbb{R}$ . For any  $N \in \mathbb{N}$ , we define  $\hat{L}_N := |\hat{\Lambda}_N|$  and  $\kappa_N$  as the number of atoms (impurities) of the Poisson random measure with intensity  $\nu > 0$  within the interval  $\hat{\Lambda}_N$ .

The following (large deviation type) lemma is needed for the proof of the subsequent Theorem C.2. Note here that  $1 - \theta + \theta \ln \theta > 0$  for  $\theta \in (0, \infty)/\{1\}$ .

**Lemma C.1.** Let  $\nu > 0$  and  $N \in \mathbb{N}$  be given. Then for any  $\theta \geq 1$ 

$$\mathbb{P}\left(\kappa_N \ge \theta \nu \hat{L}_N\right) \le e^{-\nu \hat{L}_N (1-\theta+\theta \ln \theta)} , \qquad (C.1)$$

and for any  $0 < \theta \leq 1$ 

$$\mathbb{P}\left(\kappa_N \le \theta \nu \hat{L}_N\right) \le e^{-\nu \hat{L}_N (1-\theta+\theta \ln \theta)} .$$
(C.2)

*Proof.* For  $\theta \geq 1$  we have

$$\mathbb{P}\left(\kappa_{N} \geq \theta \nu \hat{L}_{N}\right) = \sum_{m \geq \theta \nu \hat{L}_{N}} \mathbb{P}(\kappa_{N} = m) = \sum_{m \geq \theta \nu \hat{L}_{N}} e^{-\nu \hat{L}_{N}} \frac{(\nu L_{N})^{m}}{m!}$$
$$\leq \sum_{m \geq \theta \nu \hat{L}_{N}} e^{-\nu \hat{L}_{N}} \frac{(\nu \hat{L}_{N})^{m}}{m!} \theta^{m-\theta \nu \hat{L}_{N}} \leq e^{-\nu \hat{L}_{N}(1-\theta+\theta \ln \theta)}$$

On the other hand, for  $0 < \theta \leq 1$ ,

$$\mathbb{P}\left(\kappa_{N} \leq \theta \nu \hat{L}_{N}\right) = \sum_{m \leq \theta \nu \hat{L}_{N}} \mathbb{P}\left(\kappa_{N} = m\right) = \sum_{m \leq \theta \nu \hat{L}_{N}} e^{-\nu \hat{L}_{N}} \frac{(\nu L_{N})^{m}}{m!}$$
$$\leq \sum_{m \leq \theta \nu \hat{L}_{N}} e^{-\nu \hat{L}_{N}} \frac{(\nu \hat{L}_{N})^{m}}{m!} \theta^{m-\theta \nu \hat{L}_{N}} \leq e^{-\nu \hat{L}_{N}(1-\theta+\theta \ln \theta)}$$

**Theorem C.2.** Let  $(\hat{\Lambda}_N)_{N \in \mathbb{N}}$  with  $\hat{L}_N \gg \ln(N)$  be given. Then, for any  $\epsilon > 0$  and for almost any  $\omega \in \Omega$  there exists an  $\widetilde{N} = \widetilde{N}(\epsilon, \omega) \in \mathbb{N}$  such that for any  $N \ge \widetilde{N}$  we have

$$(1-\epsilon)\nu \hat{L}_N \le \kappa_N \le (1+\epsilon)\nu \hat{L}_N$$
 (C.3)

In particular, almost surely  $\lim_{N\to\infty} k_N/L_N = \nu$  and  $\lim_{N\to\infty} k_N/(\nu_N L_N) = 1$  in case of  $\ln(N)/N \ll \nu_N \lesssim 1$ .

*Proof.* Let  $\epsilon > 0$  be given. Then, with Lemma C.1 we obtain

$$\sum_{N=1}^{\infty} \mathbb{P}\Big(\kappa_N \le (1-\epsilon)\nu\hat{L}_N\Big) < \infty \quad \text{and} \quad \sum_{N=1}^{\infty} \mathbb{P}\Big(\kappa_N \ge (1+\epsilon)\nu\hat{L}_N\Big) < \infty .$$

Hence, the first part of the statement follows with the Borel–Cantelli lemma.

Consequently, for any  $\epsilon > 0$  and almost any  $\omega \in \Omega$ ,

$$\liminf_{N \to \infty} \frac{\kappa_N}{\hat{L}_N} \ge (1 - \epsilon)\nu \quad \text{and} \quad \limsup_{N \to \infty} \frac{\kappa_N}{\hat{L}_N} \le (1 + \epsilon)\nu \;.$$

Setting  $\hat{\Lambda}_N = \Lambda_N$ ,  $N \in \mathbb{N}$ , we conclude that almost surely  $\lim_{N\to\infty} k_N/L_N = \nu$ . Furthermore, recalling that  $\nu_N = s_N \nu$  we obtain  $\lim_{N\to\infty} \tilde{k}_N/(\nu_N L_N) = 1$  almost surely by setting  $\hat{\Lambda}_N = s_N \Lambda_N$ ,  $N \in \mathbb{N}$ . Note here that the assumption  $\ln(N)/N \ll \nu_N \lesssim 1$  implies  $\hat{L}_N \gg \ln(N)$ .

For a Poisson random measure with intensity  $\nu > 0$  we define  $\hat{l}_j := |(x_j(\omega), x_{j+1}(\omega))|$ for any  $j \in \mathbb{Z}$  with  $\{x_j(\omega) : j \in \mathbb{Z}\}$  the strictly increasing sequence of the atoms of the Poisson random measure, see Section 2. Note that  $\{\hat{l}_j : j \in \mathbb{Z} \setminus \{0\}\}$  are independent and identically distributed random variables with common density  $\nu e^{-\nu l}$  [Kin93, Ch. 4]. We also define the set  $J_k := \{-k, -k+1, \ldots, k-1, k\} \setminus \{0\}$  for any  $k \in \mathbb{N}$ .

**Lemma C.3.** Assume that  $\ln(N)/N \ll \nu_N \lesssim 1$  holds. Then, for any  $0 < \epsilon < 1$  and almost any  $\omega \in \Omega$  there exists an  $\widetilde{N} = \widetilde{N}(\epsilon, \omega) \in \mathbb{N}$  such that for any  $N \geq \widetilde{N}$  one has  $\widetilde{l}_j = s_N^{-1} \widehat{l}_j$  for any  $j \in J_{\lceil (1-\epsilon)\nu_N L_N/2 \rceil}$ ,  $\widetilde{l}_j \leq s_N^{-1} \widehat{l}_j$  for any  $j \in J_{\lfloor \nu_N L_N \rfloor}$  and  $\widetilde{l}_j = 0$  for any  $j \in \mathbb{Z} \setminus (J_{\lfloor \nu_N L_N \rfloor} \cup \{0\})$ .

Proof. Let  $0 < \epsilon < 1$  and  $0 < \epsilon' < \epsilon$  be given. For any  $N \in \mathbb{N}$  we divide the window  $(-s_N L_N/2, s_N L_N/2)$  into  $(-s_N L_N/2, 0]$  and  $[0, -s_N L_N/2)$ . Due to Theorem C.2 there exists a set  $\tilde{\Omega} \subset \Omega$  with  $\mathbb{P}(\tilde{\Omega}) = 1$  and the following property: For any  $\omega \in \tilde{\Omega}$  there exists an  $\tilde{N} \in \mathbb{N}$  such that for any  $N \geq \tilde{N}$  one has  $(1/2)(1-\epsilon')\nu_N L_N \leq \kappa_N^{(1)}, \kappa_N^{(2)} \leq (1/2)(1+\epsilon')\nu_N L_N$  with  $\kappa_N^{(1)}$  and  $\kappa_N^{(2)}$  denoting the number of atoms within  $(-s_N L_N/2, 0]$  and  $[0, s_N L_N/2)$ , respectively.

The statement of the lemma now follows since  $s_N^{-1}\hat{l}_j = s_N^{-1}|(x_j(\omega), x_{j+1}(\omega))|$  and  $\tilde{l}_j = s_N^{-1}|(x_j(\omega), x_{j+1}(\omega)) \cap s_N\Lambda_N|$  for any  $j \in \mathbb{Z}$ . Note that we divide the window  $s_N\Lambda_N = (-s_NL_N/2, s_NL_N/2)$  into the two intervals in order to ensure that, for  $N \geq \tilde{N}$ , both intervals with associated lengths  $\tilde{l}_{-\lceil (1-\epsilon)\nu_NL_N/2\rceil}$  and  $\tilde{l}_{\lceil (1-\epsilon)\nu_NL_N/2\rceil}$  are entirely within the window  $s_N\Lambda_N$  and that  $\tilde{l}_j = 0$  for any  $j \leq -\lfloor \nu_N L_N \rfloor - 1$  and for any  $j \geq \lfloor \nu_N L_N \rfloor + 1$ .  $\Box$ 

**Theorem C.4.** Assume that  $\ln(N)/N \ll \nu_N \lesssim 1$  holds. Furthermore, let  $(g_N)_{N \in \mathbb{N}}$  be such that  $\lim_{N \to \infty} \nu_N \lambda_N = 0$  and  $\lambda_N \gg \ln(N) \ln(\nu_N N) (\nu_N N)^{-1/2}$  with  $(\lambda_N)_{N \in \mathbb{N}}$  as in (2.43). Then

$$\liminf_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}} M_j \ge \frac{2}{9} \quad and \quad \limsup_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}} M_j \le 6 \tag{C.4}$$

almost surely with  $M_j$  as in (2.41).

*Proof.* Let  $F_k^{\nu}(l) := (2k)^{-1} \sum_{j \in J_k} \mathbb{1}_{\hat{l}_j \leq l}$  be the empirical distribution function with respect to the random variables  $\{\hat{l}_j : j \in \mathbb{Z} \setminus \{0\}\}$ . Then, for any  $l \in \mathbb{R}$ , there is a unique measure  $F_k^{\nu}(d\ell)$  defined by  $\int_{-\infty}^l F_k^{\nu}(d\ell) := F_k^{\nu}(l)$  and we also set  $F^{\nu}(l) := \int_0^l \mathbb{1}_{(0,\infty)}(l)\nu e^{-\nu x} dx$ .

Then  $\mathbb{P}(\sup_{l\in\mathbb{R}} |F_k^{\nu}(l) - F^{\nu}(l)| > \epsilon) \leq 2e^{-2k\epsilon^2}$  for any  $\epsilon > 0$  and  $k \in \mathbb{N}$  due to Dvoretzky–Kiefer–Wolfowitz inequality [DKW56, Mas90]. Therefore, for any  $\epsilon > 0$  and with  $K \in \mathbb{N}$  such that  $\ln(K) \geq \epsilon^{-2}$  we obtain

$$\sum_{k=K}^{\infty} \mathbb{P}\left(\frac{k^{1/2}}{\ln(k)} \sup_{l \in \mathbb{R}} |F_k^{\nu}(l) - F^{\nu}(l)| > \epsilon\right) \le 2\sum_{k=K}^{\infty} e^{-2\ln(k)} < \infty .$$
(C.5)

Hence,

$$\lim_{k \to \infty} \frac{k^{1/2}}{\ln(k)} \sup_{l \in \mathbb{R}} |F_k^{\nu}(l) - F^{\nu}(l)| = 0$$
(C.6)

almost surely by the Borel–Cantelli lemma.

Note that one has  $\mu_N \geq \pi^2/[5\nu_N^{-1}\ln(N)]^2$  for all but finitely many  $N \in \mathbb{N}$  since otherwise  $\lambda_N \leq N^{-5}$  which then contradicted  $\lambda_N \gg \ln(N)\ln(\nu_N N)(\nu_N N)^{-1/2}$ . Moreover, by Lemma 2.7,  $\ell_N^{>} \leq \kappa \nu^{-1}\ln(N)$  for any  $\kappa > 4$  and therefore  $F_{[\nu_N L_N/4]}^{\nu}(l) = F_{[\nu_N L_N]}^{\nu}(l) = 1$  for  $l \geq 5\nu^{-1}\ln(N)$  for all but finitely many  $N \in \mathbb{N}$  almost surely.

Therefore, almost surely with  $F^{\nu}(d\ell) = \nu e^{-\nu \ell} d\ell$ ,

$$C_{N} := \left| \int_{\pi s_{N}/\sqrt{\mu_{N}}}^{\infty} \frac{s_{N}}{\ell g_{N}} \left( \mu_{N} \frac{\ell^{2}}{s_{N}^{2}} - \pi^{2} \right) \left( F_{\lceil \nu_{N} L_{N}/4 \rceil}^{\nu} (\mathrm{d}\ell) - F^{\nu} (\mathrm{d}\ell) \right) \right|$$

$$\leq \frac{\mu_{N}}{g_{N} s_{N}} \left| \int_{\pi s_{N}/\sqrt{\mu_{N}}}^{5\nu^{-1}\ln(N)} \ell \left( F_{\lceil \nu_{N} L_{N}/4 \rceil}^{\nu} (\mathrm{d}\ell) - F^{\nu} (\mathrm{d}\ell) \right) \right| + \frac{\mu_{N}}{g_{N} s_{N}} \int_{5\nu^{-1}\ln(N)}^{\infty} \ell F^{\nu} (\mathrm{d}\ell)$$

$$+ \pi^{2} \frac{s_{N}}{g_{N}} \left| \int_{\pi s_{N}/\sqrt{\mu_{N}}}^{5\nu^{-1}\ln(N)} \ell^{-1} \left( F_{\lceil \nu_{N} L_{N}/4 \rceil}^{\nu} (\mathrm{d}\ell) - F^{\nu} (\mathrm{d}\ell) \right) \right|$$

$$+ \pi^{2} \frac{s_{N}}{g_{N}} \int_{5\nu^{-1}\ln(N)}^{\infty} \ell^{-1} F^{\nu} (\mathrm{d}\ell) .$$
(C.7)

Since  $F^{\nu}_{\lceil \nu_N L_N/4 \rceil}(l) = 1$  for any  $l \ge 5\nu^{-1} \ln(N)$  we obtain, with an integration by parts in the first and third term,

$$\begin{split} C_{N} &\leq \frac{\mu_{N}}{g_{N}s_{N}} \left( 5\nu^{-1}\ln(N) \right) \left| F_{[\nu_{N}L_{N}/4]}^{\nu} (5\nu^{-1}\ln(N)) - F^{\nu} (5\nu^{-1}\ln(N)) \right| \\ &+ \frac{\mu_{N}}{g_{N}s_{N}} \frac{\pi s_{N}}{\sqrt{\mu_{N}}} \left| F_{[\nu_{N}L_{N}/4]}^{\nu} (\pi s_{N}/\sqrt{\mu_{N}}) - F^{\nu} (\pi s_{N}/\sqrt{\mu_{N}}) \right| \\ &+ \frac{\mu_{N}}{g_{N}s_{N}} \left| \int_{\pi s_{N}/\sqrt{\mu_{N}}}^{5\nu^{-1}\ln(N)} \left( F_{[\nu_{N}L_{N}/4]}^{\nu} (\ell) - F^{\nu}(\ell) \right) d\ell \right| + \frac{\mu_{N}}{g_{N}s_{N}} \int_{5\nu^{-1}\ln(N)}^{\infty} \ell \nu e^{-\nu\ell} d\ell \\ &+ \pi^{2} \frac{s_{N}}{g_{N}} \left( 5\nu^{-1}\ln(N) \right)^{-1} \left| F_{[\nu_{N}L_{N}/4]}^{\nu} (5\nu^{-1}\ln(N)) - F^{\nu} (5\nu^{-1}\ln(N)) \right| \\ &+ \pi^{2} \frac{s_{N}}{g_{N}} \frac{\sqrt{\mu_{N}}}{\pi s_{N}} \left| F_{[\nu_{N}L_{N}/4]}^{\nu} (\pi s_{N}/\sqrt{\mu_{N}}) - F^{\nu} (\pi s_{N}/\sqrt{\mu_{N}}) \right| \\ &+ \pi^{2} \frac{s_{N}}{g_{N}} \left| \int_{\pi s_{N}/\sqrt{\mu_{N}}}^{5\nu^{-1}\ln(N)} \ell^{-2} \left( F_{[\nu_{N}L_{N}/4]}^{\nu} (\ell) d\ell - F^{\nu}(\ell) \right) d\ell \right| \\ &+ \pi^{2} \frac{s_{N}}{g_{N}} \int_{5\nu^{-1}\ln(N)}^{\infty} \ell^{-1} \nu e^{-\nu\ell} d\ell \,. \end{split}$$

Calculating further we obtain

$$C_{N} \leq 3 \frac{5\ln(N)}{\nu_{N}} \frac{\mu_{N}}{g_{N}} \sup_{l \in \mathbb{R}} |F_{\lceil \nu_{N}L_{N}/4 \rceil}^{\nu}(l) - F^{\nu}(l)| + \frac{\mu_{N}}{g_{N}} \frac{1}{\nu_{N}} \frac{5\ln(N) + 1}{N^{5}} + 3 \frac{5\ln(N)}{\nu_{N}} \frac{\mu_{N}}{g_{N}} \sup_{l \in \mathbb{R}} |F_{\lceil \nu_{N}L_{N}/4 \rceil}^{\nu}(l) - F^{\nu}(l)| + \frac{\mu_{N}}{g_{N}} \frac{1}{\nu_{N}} \frac{5\ln(N)}{N^{5}}$$
(C.9)

and since  $\mu_N \ge \pi^2/[5\nu_N^{-1}\ln(N)]^2$  we get

$$C_N \le 6 \cdot 5 \cdot \rho \frac{1}{\nu_N} \frac{\ln(N)}{\lambda_N} \sup_{l \in \mathbb{R}} |F_{\lceil \nu_N L_N/4 \rceil}^{\nu}(l) - F^{\nu}(l)| + \rho \frac{10\ln(N) + 1}{\nu_N \lambda_N} \frac{1}{N^5}$$
(C.10)

with (2.43) for all sufficiently large  $N \in \mathbb{N}$ . Hence, with (C.6) and due to  $\lambda_N \gg \ln(N) \ln(\nu_N N) (\nu_N N)^{-1/2}$  we conclude that  $\lim_{N\to\infty} C_N = 0$ . Since

$$\nu_{N} \int_{\pi s_{N}/\sqrt{\mu_{N}}}^{\infty} \frac{s_{N}}{\ell g_{N}} \left( \mu_{N} \frac{\ell^{2}}{s_{N}^{2}} - \pi^{2} \right) F^{\nu}(\mathrm{d}\ell) = \nu_{N} \mathbb{E}^{\nu} \left[ \frac{s_{N}}{(\cdot)g_{N}} \left[ \mu_{N} \frac{(\cdot)^{2}}{s_{N}^{2}} - \pi^{2} \right]_{+} \right]$$
(C.11)

$$\geq \nu_N \mathbb{E}^{\nu} \left[ N_{g_N,\mu_N}(s_N^{-1}(\cdot)) \right] \geq \frac{2}{3}\rho \tag{C.12}$$

for any  $N \in \mathbb{N}$ , see (2.34) and (2.40). Hence we conclude that almost surely

$$\liminf_{N \to \infty} \left( \nu_N \int_{\pi s_N/\sqrt{\mu_N}}^{\infty} \frac{s_N}{\ell g_N} \left( \mu_N \frac{\ell^2}{s_N^2} - \pi^2 \right) F_{\lceil \nu_N L_N/4 \rceil}^{\nu}(\mathrm{d}\ell) \right)$$

$$= \liminf_{N \to \infty} \left( \nu_N \int_{\pi s_N/\sqrt{\mu_N}}^{\infty} \frac{s_N}{\ell g_N} \left( \mu_N \frac{\ell^2}{s_N^2} - \pi^2 \right) F^{\nu}(\mathrm{d}\ell) \right) \ge \frac{2}{3}\rho .$$
(C.13)

Also, repeating the arguments from above we one can show that almost surely

$$\lim_{N \to \infty} \left| \nu_N \int_{\pi s_N/\sqrt{\mu_N}}^{\infty} \frac{s_N}{\ell g_N} \left( \mu_N \frac{\ell^2}{s_N^2} - \pi^2 \right) \left( F_{\lfloor \nu_N L_N \rfloor}^{\nu} (\mathrm{d}\ell) - F^{\nu} (\mathrm{d}\ell) \right) \right| = 0 , \qquad (C.14)$$

$$\nu_{N} \int_{\pi s_{N}/\sqrt{\mu_{N}}}^{\infty} \frac{s_{N}}{\ell g_{N}} \left( \mu_{N} \frac{\ell^{2}}{s_{N}^{2}} - \pi^{2} \right) F^{\nu}(\mathrm{d}\ell) \leq \frac{3}{2} \nu_{N} \mathbb{E}^{\nu} \left[ N_{g_{N},\mu_{N}}(s_{N}^{-1}(\cdot)) \right] \leq 3\rho \qquad (C.15)$$

for any  $N \in \mathbb{N}$ , see (2.34) and (2.40). Consequently

$$\lim_{N \to \infty} \sup_{N \to \infty} \left( \nu_N \int_{\pi s_N/\sqrt{\mu_N}}^{\infty} \frac{s_N}{\ell g_N} \left( \mu_N \frac{\ell^2}{s_N^2} - \pi^2 \right) F_{\lfloor \nu_N L_N \rfloor}^{\nu}(\mathrm{d}\ell) \right)$$

$$= \limsup_{N \to \infty} \left( \nu_N \int_{\pi s_N/\sqrt{\mu_N}}^{\infty} \frac{s_N}{\ell g_N} \left( \mu_N \frac{\ell^2}{s_N^2} - \pi^2 \right) F^{\nu}(\mathrm{d}\ell) \right) \le 3\rho .$$
(C.16)

Due to Lemma C.3 (with  $\epsilon = 1/2$ ), there exists a set  $\widetilde{\Omega} \subset \Omega$  with  $\mathbb{P}(\widetilde{\Omega}) = 1$  and the following property: For any  $\omega \in \widetilde{\Omega}$  there exists an  $\widehat{N}(\omega) \in \mathbb{N}$  such that for any  $N \geq \widehat{N}(\omega)$  we have  $\widetilde{l}_j = s_N^{-1} \widehat{l}_j$  for any  $j \in J_{\lceil \nu_N L_N/4 \rceil}$ ,  $\widetilde{l}_j \leq s_N^{-1} \widehat{l}_j$  for any  $j \in J_{\lfloor \nu_N L_N \rfloor}$ , and  $\widetilde{l}_j = 0$  for any  $j \in \mathbb{Z} \setminus (J_{\lfloor \nu_N L_N \rfloor} \cup \{0\})$ . Consequently we obtain  $(M_0 = 0)$ , for any  $N \geq \widehat{N}(\omega)$ ,

$$\sum_{j \in J_{\lceil \nu_N L_N/4 \rceil}} \frac{s_N}{\hat{l}_j g_N} \left[ \mu_N \frac{\hat{l}_j^2}{s_N^2} - \pi^2 \right]_+ \leq \sum_{j \in \mathbb{Z}: M_j \ge 1} \frac{1}{\tilde{l}_j g_N} \left[ \mu_N \tilde{l}_j^2 - \pi^2 \right]_+$$

$$\leq \sum_{j \in J_{\lfloor \nu_N L_N \rfloor}} \frac{s_N}{\hat{l}_j g_N} \left[ \mu_N \frac{\hat{l}_j^2}{s_N^2} - \pi^2 \right]_+$$
(C.17)

Lastly, due to (2.41), (2.34), (C.17), (C.13), (C.16), and  $M_0 = 0$ , we have almost surely

$$\begin{split} \liminf_{N \to \infty} \frac{\nu_N}{\nu_N L_N / 2} \sum_{j \in \mathbb{Z}} M_j &\geq \liminf_{N \to \infty} \frac{\nu_N}{\nu_N L_N / 2} \sum_{j \in \mathbb{Z}: M_j \geq 1} N_{g_N, \mu_N}(\tilde{l}_j) \\ &\geq \frac{2}{3} \liminf_{N \to \infty} \frac{\nu_N}{\nu_N L_N / 2} \sum_{j \in \mathbb{Z}: M_j \geq 1} \frac{1}{\tilde{l}_j g_N} \left[ \mu_N \tilde{l}_j^2 - \pi^2 \right]_+ \\ &\geq \frac{2}{3} \liminf_{N \to \infty} \frac{\nu_N}{\nu_N L_N / 2} \sum_{j \in J_{\lceil \nu_N L_N / 4 \rceil}} \frac{s_N}{\hat{l}_j g_N} \left[ \mu_N \frac{\hat{l}_j^2}{s_N^2} - \pi^2 \right]_+ \\ &\geq \frac{2}{3} \liminf_{N \to \infty} \nu_N \int_{\pi s_N / \sqrt{\mu_N}}^{\infty} \frac{s_N}{lg_N} \left( \mu_N \frac{l^2}{s_N^2} - \pi^2 \right) F_{\lceil \nu_N L_N / 4 \rceil}^{\nu} (\mathrm{d}\ell) \\ &\geq \left(\frac{2}{3}\right)^2 \rho \; . \end{split}$$

Similarly,

$$\begin{split} \limsup_{N \to \infty} \frac{\nu_N}{2\nu_N L_N} \sum_{j \in \mathbb{Z}} M_j &\leq \limsup_{N \to \infty} \frac{\nu_N}{2\nu_N L_N} \sum_{j \in \mathbb{Z}: M_j \geq 1} \frac{1}{\tilde{l}_j g_N} \left[ \mu_N \tilde{l}_j^2 - \pi^2 \right]_+ \\ &+ \lim_{N \to \infty} \frac{\nu_N}{2\nu_N L_N} \sum_{j \in \mathbb{Z}: M_j \geq 1} 1 \\ &\leq \limsup_{N \to \infty} \frac{\nu_N}{2\nu_N L_N} \sum_{j \in \mathbb{Z}: M_j \geq 1} \frac{s_N}{\hat{l}_j g_N} \left[ \mu_N \frac{\hat{l}_j^2}{s_N^2} - \pi^2 \right]_+ \\ &+ \lim_{N \to \infty} \frac{\nu_N}{2\nu_N L_N} \sum_{j \in \mathbb{Z}: M_j \geq 1} 1 \\ &\leq \limsup_{N \to \infty} \nu_N \int_{\pi s_N / \sqrt{\mu_N}}^{\infty} \frac{s_N}{lg_N} \left( \mu_N \frac{l^2}{s_N^2} - \pi^2 \right) F_{\lfloor \nu_N L_N \rfloor}^{\nu} (\mathrm{d}\ell) \\ &+ \lim_{N \to \infty} \nu_N \int_{\pi s_N / \sqrt{\mu_N}}^{\infty} F_{\lfloor \nu_N L_N \rfloor}^{\nu} (\mathrm{d}\ell) \leq 3\rho \;, \end{split}$$

since, due to (C.6), (2.42), and our assumptions,

$$\lim_{N \to \infty} \nu_N \int_{\pi s_N/\sqrt{\mu_N}}^{\infty} F_{\lfloor \nu_N L_N \rfloor}^{\nu} (\mathrm{d}\ell) = \lim_{N \to \infty} \nu_N \left[ 1 - F_{\lfloor \nu_N L_N \rfloor}^{\nu} (\pi s_N/\sqrt{\mu_N}) \right]$$

$$= \lim_{N \to \infty} \nu_N \left[ 1 - F^{\nu} (\pi s_N/\sqrt{\mu_N}) \right] = \lim_{N \to \infty} \nu_N \lambda_N = 0 .$$
(C.18)

We therefore obtain almost surely

$$\liminf_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}} M_j = \frac{1}{2\rho} \left( \liminf_{N \to \infty} \frac{\nu_N}{\nu_N L_N/2} \sum_{j \in \mathbb{Z}} M_j \right) \ge \frac{2}{9} , \qquad (C.19)$$

and

$$\limsup_{N \to \infty} \frac{1}{N} \sum_{j \in \mathbb{Z}} M_j \le 6 .$$
 (C.20)

**Theorem C.5.** Let  $1/N \ll \nu_N \lesssim 1$  be given. Then for any  $0 < \epsilon < 1$  and for almost any  $\omega \in \Omega$  there exists an  $\widetilde{N} = \widetilde{N}(\epsilon, \omega) \in \mathbb{N}$  such that for any  $N \geq \widetilde{N}$  we have

$$\tilde{\ell}_N^> > \nu_N^{-1} \Big\{ \ln(L_{\lfloor s_N N \rfloor}) - (1+\epsilon) \ln[\ln(L_{\lfloor s_N N \rfloor})] \Big\} .$$
(C.21)

*Proof.* Let  $0 < \epsilon < 1$  be given. Then, for almost any  $\omega \in \Omega$  there exists an  $\widehat{N}_1 = \widehat{N}_1(\epsilon, \omega) \in \mathbb{N}$  such that

$$\left\{\hat{l}_j: j \in J_{\left\lceil (1-\epsilon)\nu L_N/2 \right\rceil}\right\} \subsetneq \left\{l_j: j \in \mathbb{Z} \setminus \{0\}\right\} \setminus \{0\}$$
(C.22)

for any  $N \geq \hat{N}_1$  by Lemma C.3. Moreover, since  $\{\hat{l}_j : j \in \mathbb{Z}\}$  are mutually independent exponentially distributed random variables one obtains

$$\mathbb{P}\left(\max\left\{\hat{l}_{j}: j \in J_{\lceil (1-\epsilon)\nu L_{N}/2\rceil}\right\} \le \nu^{-1}\left\{\ln(L_{N}) - (1+\epsilon)\ln[\ln(L_{N})]\right\}\right)$$
(C.23)

$$\leq \left(1 - \frac{\left[\ln(L_N)\right]^{1+\epsilon}}{L_N}\right)^{2\left[(1-\epsilon)\nu L_N/2\right]} . \tag{C.24}$$

Moreover, since  $\ln(1-x) \leq -x$  for any 0 < x < 1 we have

$$2\left\lceil \frac{(1-\epsilon)\nu L_N}{2} \right\rceil \cdot \ln\left[ \left( 1 - \frac{[\ln(L_N)]^{1+\epsilon}}{L_N} \right) \right] \le -(1-\epsilon)\nu [\ln(L_N)]^{1+\epsilon} \le -2\ln(N) \quad (C.25)$$

for all but finitely many  $N \in \mathbb{N}$  and therefore

$$\sum_{N=1}^{\infty} \left( 1 - \frac{\left[\ln(L_N)\right]^{1+\epsilon}}{L_N} \right)^{2\left[(1-\epsilon)\nu L_N\right]/2} < \infty .$$
 (C.26)

Hence, with Borel–Cantelli's lemma there exists a set  $\widetilde{\Omega} \subset \Omega$  with  $\mathbb{P}(\widetilde{\Omega}) = 1$  such that for any  $\omega \in \widetilde{\Omega}$  there is an  $\widehat{N}_2 = \widehat{N}_2(\epsilon, \omega) \in \mathbb{N}$  with

$$\ell_N^> \ge \max\left\{\hat{l}_j : j \in J_{\lceil (1-\epsilon)\nu L_N/2\rceil}\right\} > \nu^{-1}\left\{\ln(L_N) - (1+\epsilon)\ln[\ln(L_N)]\right\}$$
(C.27)

for any  $N \geq \widehat{N}_2$ .

Finally, for any  $\omega \in \widetilde{\Omega}$  we define  $\widetilde{N}(\epsilon, \omega)$  such that  $\lfloor s_N N \rfloor \geq \widehat{N}_2(\epsilon, \omega)$  for any  $N \geq \widetilde{N}(\epsilon, \omega)$ . Hence,  $\ell_{\lfloor s_N N \rfloor}^{>}$  denoting the length of the largest interval within the window  $\Lambda_{\lfloor s_N N \rfloor}$ ,

$$s_N^{-1}\ell_{\lfloor s_N N \rfloor}^{>} > \nu_N^{-1} \Big\{ \ln(L_{\lfloor s_N N \rfloor}) - (1+\epsilon) \ln\left[ \ln(L_{\lfloor s_N N \rfloor}) \right] \Big\}$$
(C.28)

for any  $N \ge \widetilde{N}(\epsilon, \omega)$ . The statement then follows since  $\widetilde{\ell}_N^> \ge s_N^{-1} \ell_{\lfloor s_N N \rfloor}^>$ .

**Theorem C.6.** For any  $0 < \eta' < 2$  there exists an  $\widetilde{N}(\eta') \in \mathbb{N}$  such that for any  $N \ge \widetilde{N}(\eta')$  one has

$$\mathbb{P}\left(\ell_N^> > \nu^{-1} \left[\ln(L_N) + \ln(C_1)\right]\right) > 1 - \frac{1}{2}\eta'$$
(C.29)

with  $C_1 := -\nu/[4\ln(\eta'/2)] > 0.$ 

*Proof.* According to Lemma C.3 (with  $\epsilon = 1/2$ ), for almost any  $\omega \in \Omega$  there exists an  $\widehat{N} = \widehat{N}(\omega) \in \mathbb{N}$  such that for any  $N \geq \widehat{N}$  one has

$$\left\{\hat{l}_j: j \in J_{\lceil \nu L_N/4 \rceil}\right\} \subsetneq \left\{l_j: j \in \mathbb{Z} \setminus \{0\}\right\} \setminus \{0\}$$

Hence,  $[\max\{\hat{l}_j : j \in J_{\lceil \nu L_N/4 \rceil}\} - \ell_N^>]_+$  converges to zero almost surely and consequently, for any  $\eta > 0$ ,

$$\lim_{N \to \infty} \mathbb{P}\left(\ell_N^{>} \le \max\left\{\hat{l}_j : j \in J_{\lceil \nu L_N/4 \rceil}\right\} - \eta\right)$$
$$\le \lim_{N \to \infty} \mathbb{P}\left(\left[\max\left\{\hat{l}_j : j \in J_{\lceil \nu L_N/4 \rceil}\right\} - \ell_N^{>}\right]_+ \ge \eta\right) = 0$$

Furthermore, since  $\ln(1-x) = -x$  for 0 < x < 1,

$$\mathbb{P}\left(\max\left\{\hat{l}_{j}: j \in J_{\lceil \nu L_{N}/4 \rceil}\right\} \le \nu^{-1} \left[\ln(L_{N}) + \ln(4C_{1}/3)\right]\right)$$
$$\le \left(1 - \frac{1}{(4/3)L_{N}C_{1}}\right)^{2\lceil \nu L_{N}/4 \rceil} \le e^{-3\nu/(8C_{1})}$$

for all but finitely many  $N \in \mathbb{N}$ . Hence, altogether one obtains, with  $\eta = -\nu^{-1} \ln(3/4)$ ,

$$\mathbb{P}\left(\ell_{N}^{>} > \nu^{-1}\left[\ln(L_{N}) + \ln\left(C_{1}\right)\right]\right) \\
\geq \mathbb{P}\left(\ell_{N}^{>} > \max\left\{\hat{l}_{j} : j \in J_{\left\lceil\nu L_{N}/4\right\rceil}\right\} - \eta\right) \\
+ \mathbb{P}\left(\max\left\{\hat{l}_{j} : j \in J_{\left\lceil\nu L_{N}/4\right\rceil}\right\} > \nu^{-1}\left[\ln(L_{N}) + \ln\left(4C_{1}/3\right)\right]\right) - 1 \\
\geq 1 - \frac{1}{2}\eta'$$

for all but finitely many  $N \in \mathbb{N}$ .

30

Recall that  $\ell_N^> = l_N^{>,1}$  is the largest and  $l_N^{>,k}$ ,  $k \in \mathbb{N}$ , is the *k*th largest length of the set  $\{l_j = |(x_j(\omega), x_{j+1}(\omega)) \cap \Lambda_N| : j \in \mathbb{Z}\}$ . In the same way we define  $\tilde{l}_N^{>,k}$  for the scaled lengths.

**Theorem C.7.** For any  $0 < \eta' < 2$  and any  $C_3 > 2$  there exists an  $\widetilde{N} = \widetilde{N}(\eta', C_3) \in \mathbb{N}$ such that for any  $N \ge \widetilde{N}$ 

$$\mathbb{P}\left(\ell_N^> > \nu^{-1}\ln(L_N C_1), \, l_N^{>,\lceil 2\nu C_3/(C_1\eta')\rceil + 1} \le \nu^{-1}\left[\ln(L_N C_1) - \ln(C_3/2)\right]\right) > 1 - \eta' \quad (C.30)$$

with  $C_1 = -\nu/[4\ln(\eta'/2)].$ 

*Proof.* According to Theorem C.6 there exists a number  $\widetilde{N}(\eta') \in \mathbb{N}$  such that for any  $N \geq \widetilde{N}(\eta')$  one has,  $\Omega_1 := \{\omega \in \Omega : \ell_N^> > \nu^{-1} \ln(C_1 L_N)\}$ ,

$$\mathbb{P}(\Omega_1) > 1 - \frac{1}{2}\eta' . \tag{C.31}$$

Moreover,

$$\mathbb{E}\left[\mathcal{N}_{N}^{\mathrm{I},\omega}(E)\right] \leq \mathcal{N}_{\infty}^{\mathrm{I}}(E) \tag{C.32}$$

for any  $E \ge 0$  and  $N \in \mathbb{N}$  [PF92, Theorem 5.25]. Here,  $\mathcal{N}_N^{\mathrm{I},\omega}(E) = L_N^{-1} |\{i : E_N^{i,\omega} \le E\}|$  is the finite-volume integrated density of states, i.e., the number of eigenvalues of the non-interacting Luttinger–Sy model that are smaller than or equal to E divided by the volume of the system, and

$$\mathcal{N}_{\infty}^{\mathrm{I}}(E) = \nu \frac{\mathrm{e}^{-\nu \pi E^{-1/2}}}{1 - \mathrm{e}^{-\nu \pi E^{-1/2}}} \tag{C.33}$$

is the limiting integrated density of states of the non-interacting Luttinger–Sy model, see e.g. [Zag07, Proposition III.2].

Hence,  $\widetilde{E} := \pi^2 \nu^2 [\ln(C_1 L_N) - \ln(C_3/2)]^{-2}$ ,

$$k \mathbb{P}\left(\left|\left\{i: E_N^{i,\omega} \le \widetilde{E}\right\}\right| \ge k\right) \le \sum_{j\ge 1} j \mathbb{P}\left(\left|\left\{i: E_N^{i,\omega} \le \widetilde{E}\right\}\right| = j\right)$$
$$= \mathbb{E}\left[\left|\left\{i: E_N^{i,\omega} \le \widetilde{E}\right\}\right|\right] \le \nu \frac{C_3}{C_1}$$

for any  $k \in \mathbb{N}$  and all but finitely many  $N \in \mathbb{N}$ . Setting

$$\Omega_2 := \left\{ \omega \in \Omega : \left| \left\{ i : E_N^{i,\omega} \le \widetilde{E} \right\} \right| < \left\lceil \frac{2\nu C_3}{C_1 \eta'} \right\rceil \right\}$$

one obtains

$$\mathbb{P}(\Omega_2) \ge 1 - \frac{1}{2}\eta' . \tag{C.34}$$

for all but finitely many  $N \in \mathbb{N}$ .

Finally,  $\mathbb{P}(\Omega_1 \cap \Omega_2) \ge 1 - \eta'$  for all but finitely many  $N \in \mathbb{N}$ , and for any  $\omega \in \Omega_1 \cap \Omega_2$  one has

$$\ell_N^> > \nu^{-1} \ln(C_1 L_N)$$

and

$$l_N^{>,\lceil 2\nu C_3/(C_1\eta')\rceil+1} \le \nu^{-1} \left[ \ln(C_1 L_N) - \ln\left(\frac{C_3}{2}\right) \right] .$$

Consequently,  $\ell_N^> - l_N^{>,\lceil 2\nu C_3/(C_1\eta')\rceil+1} \ge \nu^{-1} \ln(C_3/2)$  for any  $\omega \in \Omega_1 \cap \Omega_2$ .

**Corollary C.8.** Let  $1/N \ll \nu_N \lesssim 1$  be given. Then for any  $0 < \eta' < 2$  and any  $C_3 > 2e^{\nu/\rho}$  there is an  $\widetilde{N} = \widetilde{N}(\eta', C_3) \in \mathbb{N}$  such that for any  $N \geq \widetilde{N}$  we have

$$\mathbb{P}\left(\tilde{\ell}_{N}^{>} > \nu_{N}^{-1}\ln(C_{1}L_{\lfloor s_{N}N \rfloor}), \,\tilde{\ell}_{N}^{>} - \tilde{\ell}_{N}^{>,\lceil 2\nu C_{3}/(C_{1}\eta')\rceil + 1} > \nu_{N}^{-1}\ln(C_{3}/(2e^{\nu/\rho}))\right) > 1 - \eta' \quad (C.35)$$

with  $C_1 := -\nu/[4\ln(\eta'/2)].$ 

*Proof.* With Theorem C.7 there exists an  $\widehat{N} = \widehat{N}(\eta') \in \mathbb{N}$  such that for any  $N \ge \widehat{N}$  one has

$$\mathbb{P}\left(s_{N}^{-1}\ell_{\lfloor s_{N}N \rfloor}^{>} > \nu_{N}^{-1}\ln(C_{1}L_{\lfloor s_{N}N \rfloor}), s_{N}^{-1}l_{\lfloor s_{N}N \rfloor}^{>,\lceil 2\nu C_{3}/(C_{1}\eta')\rceil+1} < \nu_{N}^{-1}\left[\ln(C_{1}L_{\lfloor s_{N}N \rfloor}) - \ln(C_{3}/2)\right]\right) > 1 - \eta' .$$
(C.36)

As in the proof of Theorem C.5 we use  $\tilde{\ell}_N^> \geq s_N^{-1} \ell_{\lfloor s_N N \rfloor}^>$  as well as  $\tilde{l}_N^{>,\lceil 2\nu C_3/(C_1\eta')\rceil+1} \leq s_N^{-1} \ell_{\lfloor s_N N \rfloor+1}^{>,\lceil 2\nu C_3/(C_1\eta')\rceil+1}$  to obtain

$$\begin{split} \tilde{\ell}_N^> &- \tilde{l}_N^{>,\lceil 2\nu C_3/(C_1\eta')\rceil+1} \ge s_N^{-1} \left( l_{\lfloor s_N N \rfloor}^> - l_{\lfloor s_N N \rfloor+1}^{>,\lceil 2\nu C_3/(C_1\eta')\rceil+1} \right) \\ &\ge s_N^{-1} \left( l_{\lfloor s_N N \rfloor}^> - \left( l_{\lfloor s_N N \rfloor}^{>,\lceil 2\nu C_3/(C_1\eta')\rceil+1} + \rho^{-1} \right) \right) \end{split}$$

Now, using the two inequalities appearing in (C.36) we conclude

$$\tilde{\ell}_N^> - \tilde{l}_N^{>,\lceil 2\nu C_3/(C_1\eta')\rceil + 1} \ge \nu_N^{-1} \ln\left(\frac{C_3}{2\mathrm{e}^{\nu/\rho}}\right)$$

from which the statement readily follows.

By slightly changing the proof we could also allow for  $C_3 > 6$  instead of  $C_3 > 2e^{\nu/\rho}$ . This would replace  $\ln(C_3/2e^{\nu/\rho})$  in (C.35) by  $\ln(C_3/6)$ .

## References

- [BK14] J. Bolte and J. Kerner, Many-particle quantum graphs and Bose-Einstein condensation, J. Math. Phys. 55 (2014), no. 6, 061901.
- [BK16] J. Bolte and J. Kerner, Instability of Bose-Einstein condensation into the one-particle ground state on quantum graphs under repulsive perturbations, J. Math. Phys. 57 (2016), no. 4, 043301.
- [Cas68] H. Casimir, On Bose-Einstein condensation, Fundamental Problems in Statistical Mechanics 3 (1968), 188–196.
- [DKW56] A. Dvoretzky, J. Kiefer, and J. Wolfowitz, Asymptotic minimax character of the sample distribution function and of the classical multinomial estimator, The Annals of Mathematical Statistics (1956), 642–669.
- [dS86] P. de Smedt, The Effect of Repulsive Interactions on Bose-Einstein Condensation, J. Stat. Phys. 45 (1986), 201–213.
- [Ein24] A. Einstein, Quantentheorie des einatomigen idealen Gases, Sitzber. Kgl. Preuss. Akad. Wiss. (1924), 261–267.
- [Ein25] \_\_\_\_, Quantentheorie des einatomigen idealen Gases, II. Abhandlung, Sitzber. Kgl. Preuss. Akad. Wiss. (1925), 3–14.
- [Gir60] M. Girardeau, Relationship between systems of impenetrable bosons and fermions in one dimension, J. Math. Phys. 1 (1960), no. 6, 516–523.
- [Hoh67] P. C. Hohenberg, Existence of Long-Range Order in One and Two Dimensions, Phys. Rev. 158 (1967), 383–386.
- [JPZ10] T. Jaeck, J. V. Pulé, and V. A. Zagrebnov, On the nature of Bose-Einstein condensation enhanced by localization, J. Math. Phys. 51 (2010), no. 10, 103302.
- [Kin93] J. Kingman, Poisson Processes, Clarendon Press, 1993.
- [KLS88] T. Kennedy, E. H. Lieb, and B. S. Shastry, The XY model has long-range order for all spins and all dimensions greater than one, Phys. Rev. Lett. 61 (1988), 2582–2584.
- [KS85] W. Kirsch and B. Simon, Universal lower bounds on eigenvalue splittings for one dimensional Schrödinger operators, Comm. Math. Phys. 97 (1985), no. 3, 453–460.
- [LL63] E. H. Lieb and W. Liniger, Exact Analysis of an Interacting Bose Gas. I. The General Solution and the Ground State, Phys. Rev. 130 (1963), 1605–1616.
- [LPZ04] O. Lenoble, L. A. Pastur, and V. A. Zagrebnov, Bose-Einstein condensation in random potentials, Comptes Rendus Physique 5 (2004), no. 1, 129–142.
- [LS73a] J. M. Luttinger and H. K. Sy, Bose-Einstein Condensation in a One-Dimensional Model with Random Impurities, Phys. Rev. A 7 (1973), no. 2, 712.

- [LS73b] \_\_\_\_, Low-Lying Energy Spectrum of a One-Dimensional Disordered System, Phys. Rev. A 7 (1973), 701–712.
- [LS02] E. H. Lieb and R. Seiringer, Proof of Bose-Einstein Condensation for Dilute Trapped Gases, Phys. Rev. Lett. 88 (2002), 170409.
- [LSSY05] E. H. Lieb, R. Seiringer, J. P. Solovej, and J. Yngvason, The mathematics of the Bose gas and its condensation, Oberwolfach Seminars, vol. 34, Birkhäuser Verlag, Basel, 2005.
- [LSY00] E. Lieb, R. Seiringer, and J. Yngvason, Bosons in a trap: A rigorous derivation of the Gross-Pitaevskii energy functional, Phys. Rev. A 61 (2000), no. 4, 043602.
- [LSY04] E. H. Lieb, R. Seiringer, and J. Yngvason, One-dimensional behavior of dilute, trapped Bose gases, Comm. Math. Phys. 244 (2004), no. 2, 347–393.
- [LVZ03] J. Lauwers, A. Verbeure, and V. A. Zagrebnov, Proof of Bose-Einstein condensation for interacting gases with a one-particle gap, J. Phys. A 36 (2003), 169–174.
- [LW79] L. J. Landau and I. F. Wilde, On the Bose-Einstein condensation of an ideal gas, Comm. Math. Phys. 70 (1979), no. 1, 43–51.
- [Mas90] P. Massart, The tight constant in the Dvoretzky-Kiefer-Wolfowitz inequality, The Annals of Probability (1990), 1269–1283.
- [Mic07] A. Michelangeli, Reduced density matrices and Bose–Einstein condensation, SISSA **39** (2007).
- [PF92] A. L. Pastur and A. Figotin, Spectra of Random and Almost-Periodic Operators, Springer-Verlag, 1992.
- [PO56] O. Penrose and L. Onsager, Bose-Einstein condensation and Liquid Helium, Phys. Rev. 104 (1956), 576–584.
- [SYZ12] R. Seiringer, J. Yngvason, and V. A. Zagrebnov, Disordered Bose-Einstein condensates with interaction in one dimension, J. Stat. Mech.: Theory and Experiment 2012 (2012), no. 11, P11007.
- [SYZ13] R. Seiringer, J. Yngvason, and V. A. Zagrebnov, Disordered Bose-Einstein condensates with interaction, Proceedings of XVIIth International Congress on Mathematical Physics, World Scientific (2013), 610–619.
- [van83] M. van den Berg, On condensation in the free-boson gas and the spectrum of the Laplacian, J. Stat. Phys. 31 (1983), 623–637.
- [vL82] M. van den Berg and J. T. Lewis, On generalized condensation in the free boson gas, Physica A: Statistical Mechanics and its Applications 110 (1982), no. 3, 550–564.
- [vLL86] M. van den Berg, J. T. Lewis, and M. Lunn, On the general theory of Bose-Einstein condensation and the state of the free boson gas, Helvetica Physica Acta 59 (1986), no. 8, 1289–1310.

- [vLP86] M. van den Berg, J. T. Lewis, and J. V. Pulé, A general theory of Bose-Einstein condensation, Helvetica Physica Acta 59 (1986), no. 8, 1271–1288.
- [Zag07] V. A. Zagrebnov, Bose-Einstein condensation in a random media, Journal of Physical Studies 11 (2007), no. 1, 108–121.
- [ZB01] V. A. Zagrebnov and J. B. Bru, The Bogoliubov model of weakly imperfect Bose gas, Phys. Rep. 350 (2001), no. 56, 291–434.