Exoplanet Populations and their Dependence on Host Star Properties

Gijs D. Mulders

Abstract Exoplanets around different types of stars provide a window into the diverse environments in which planets form. This chapter describes the observed relations between exoplanet populations and stellar properties and how they connect to planet formation in protoplanetary disks. Giant planets occur more frequently around more metal-rich and more massive stars. These findings support the core accretion theory of planet formation, in which the cores of giant planets form more rapidly in more solid-rich and more gas-rich protoplanetary disks. Smaller planets, those with sizes roughly between Earth and Neptune, exhibit different scaling relations with stellar properties. These planets orbit stars with a range of metallicities and occur more frequently around lower mass stars, indicating that planet formation takes place in a wide range of environments. Within M dwarfs, both radial velocity and transit surveys show that planets are smaller and located closer to the star when the stellar mass is lower. Additions to the core accretion model, in particular pebble accretion, have shown success in explaining the enhanced planet formation efficiency around low mass stars.

Introduction

Exoplanets are observed around a diverse set of host stars. The first exoplanet discovered around a main-sequence star, 51 Pegasi b, orbits a star enriched in heavy elements (metals) compared to the sun (Mayor and Queloz 1995). In contrast, one of the earliest discovered planets that could conceivably be rocky, Gliese 581e, orbits a metal-poor M dwarf less than a third the mass of the sun (Mayor et al. 2009). While these discoveries represent just two examples of the more than five thou-

Gijs D. Mulders

Facultad de Ingeniería y Ciencias, Universidad Adolfo Ibáñez, Av. Diagonal las Torres 2640, Peñalolén, Santiago, Chile, e-mail: gijs.mulders@uai.cl Millennium Institute for Astrophysics, Chile

sand exoplanets known to date¹, they are indicative of the broader trends between exoplanets and their host stars that have since emerged from exoplanet surveys, illustrated in Figure 1. Giant planets occur more frequently around more massive and more metal-rich stars (e.g. Santos et al. 2004; Johnson et al. 2010; Ghezzi et al. 2018). Sub-Neptunes occur around stars with a wide range of metallicities (Sousa et al. 2008; Buchhave et al. 2012), but occur more frequently around lower mass stars (Howard et al. 2012; Mulders et al. 2015a; Sabotta et al. 2021; Zink et al. 2023).

It is no coincidence that the smallest planets were first discovered around M dwarfs. The lower stellar mass compared to more sun-like stars with spectral types F, G and K facilitates the detection of less massive planets with radial velocity techniques (e.g. Endl et al. 2003). Similarly, the small size of M dwarfs lead to deeper transits for a planet of the same size when compared to sun-like stars (Dressing and Charbonneau 2013). Most exoplanets to date, however, have been discovered around F, G, and K dwarfs because more bright targets are observable. Hence, an understanding of survey detection efficiency and selection biases are crucial to understand trends in the occurrence of the exoplanet population with host star properties.

While the giant planet-metallicity correlation was initially interpreted as pollution of the stellar atmosphere by planetary systems (e.g. Gonzalez 1997), it is now widely accepted that the stellar metallicity is a proxy of the solid inventory of the protoplanetary disks in which planets form. It has been established that the increased occurrence of giant planets around high-metallicity stars arises because giant planet cores are more likely to form in disks with a larger amount of solids (e.g. Ida and Lin 2004; Emsenhuber et al. 2021). Similarly, the lower frequency of giant planets around M dwarfs can be explained by those stars having less massive disks (Laughlin et al. 2004; Burn et al. 2021). The relation between exoplanets and their hosts stars provide stringent constraints on planet formation theory, as properties of exoplanet host stars trace the conditions in protoplanetary disks at the time of planet formation.

These trends, however, breaks down for planets smaller than Neptune, hereafter sub-Neptunes, which poses some urgent questions about the planet formation process. Why is the frequency of sub-Neptunes almost independent of stellar metallicity, even when the initial inventory of condensible solids must have varied by an order of magnitude? Do the elevated planet occurrence rates around M dwarfs, where protoplanetary disk masses were lower, imply that there is something fundamentally different about the planet formation process around low-mass stars?

Not all stars are equally amenable for exoplanets discovery and certain types of stars have been more thoroughly searched than others. To account for these selection and detection biases, planet occurrence rates can be calculated to infer trends in the intrinsic planet population. Variations in the planet occurrence rate with stellar parameters can be estimated from exoplanets surveys under the following conditions:

¹ https://exoplanetarchive.ipac.caltech.edu/



Fig. 1 Trends in the exoplanet population as function of stellar mass and metallicity, illustrating the different behavior of the giant planet population (large pink circles) and planets smaller than Neptune (small cyan circles). The location of the sun is indicated with a yellow star. The location of individual symbols is randomly generated, with the density of point corresponding to the exoplanet occurrence rate. Any resemblance between symbol locations and observed exoplanets is entirely coincidental.

- 1. The survey covers a range of stellar properties, with a sufficient number of planet detections across this range to identify trends.
- 2. Stellar properties are known for the surveyed stars, including those of stars without detected planets, to estimate the *fraction* of stars with a given set of properties hosting planets.
- 3. The survey completeness can be estimated, to separate observation bias from intrinsic trends in the exoplanet population.

The focus of this chapter are trends identified in radial velocity and transit surveys with stellar mass and metallicity, which (mostly) satisfy these three requirements. A notable omission are direct imaging surveys, though it should be noted that they are consistent with the positive stellar mass dependence of giant planets (e.g. Nielsen et al. 2019; Vigan et al. 2021).

Trends for giant planets out to a few au and sub-Neptunes at orbital periods shorter than a few hundred days are discussed seperately. Emphasis is placed on studies that take into account the different observation bias and survey detection efficiency that exist when surveying planets around various type of stars. These trends are then placed into the context of planet formation theory and models. An outlook for current and future surveys that can fill in some of the gaps in the current knowledge of the exoplanets populations around different types of stars is presented towards the end of this chapter, followed by a brief conclusion.

Trends with Stellar Metallicity

There is a general consensus that giant planet occurrence rates increase with host star metallicity, see also the review by Udry and Santos (2007). The giant planetmetallicity relation is seen in radial velocity surveys of sun-like stars, M dwarfs, and evolved stars, and has also been identified for transiting planets. However, Sub-Neptunes are found around stars with a wider range of metallicities, with no clear preference for metal-rich stars. Throughout this chapter, the logarithm of the iron abundance with respect the solar abundance, [Fe/H], is used to represent stellar metallicity.



Fig. 2 Giant planet occurrence rate as function of stellar metallicity, from Fischer and Valenti (2005) figure 5. The red solid line shows a quadratic relation between planet occurrence and stellar metallicity ($\beta = 2$, eq. 1). Figure reproduced from Fischer and Valenti (2005) with permission from the authors.

Positive Giant Planet-Metallicity Correlation

Giant planets occur more frequently around stars with higher metallicities (See Fig. 2). Early indications of a planet-metallicity correlation were found by Gonzalez (1997) based on metallicities of a handful of exoplanet hosts including 51 Peg b. The trend that giant planets are preferentially found around metal-rich host stars was subsequently recovered in larger samples (Gonzalez 1998; Fuhrmann et al. 1998; Santos et al. 2000; Gonzalez et al. 2001; Sadakane et al. 2002; Laws et al. 2003).

As outlined in the preceding section, characterizing star properties of non-planet hosts and detection efficiency of the survey are critical to separate observation bias from intrinsic planet population. Santos et al. (2001, 2003) measured abundances for non-planet hosting stars and found that the giant planet hosts are systematically more metal-rich in a volume-limited sample. The detection frequency of giant planets was shown to increase with metallicity in a volume-limited sample of stars from the Hipparcos catalog (Reid 2002). Santos et al. (2004) estimated planet occurrence rate as function of metallicity and identified a positive correlation at super-solar metallicities. The high occurrence rate of giant planets around metal-rich stars was confirmed by (Fischer and Valenti 2005, see also Fig. 2), who derived stellar abundances of stars in the Keck, Lick, and Anglo-Australian Telescope planet search surveys.

The occurrence rate of giant planets is a strong function of metallicity and scales roughly with the square of the number of iron atoms. At super-solar metallicities, [Fe/H] > 0, where planet detections are plenty, metallicity increases by a factor of 5 from ~ 5% at [Fe/H] = 0 to ~ 25% at [Fe/H] = 0.5. At lower metallicities, the shape of the metallicity distribution is less well quantified due to few planet detections, with a giant planet occurrence rate of approximately ~ 2 – 3% (e.g. Santos et al. 2004). The functional form of the planet occurrence-metallicity correlation is often assumed to be a power-law²

. . .

$$f_{\text{giant}} \propto 10^{\beta [\text{Fe/H}]},$$
 (1)

with index $\beta \approx 2$ (e.g. Fischer and Valenti 2005; Udry and Santos 2007; Sousa et al. 2011). Johnson et al. (2010) showed that such a functional form provides a better fit than a flat distribution at sub-solar metallicities. The planet occurrence rate likely continues to decrease at metallicities below [Fe/H] < -0.5 (Santos et al. 2011; Mortier et al. 2012), consistent with the non-detection of giant planets in metal-poor clusters and halo stars (e.g. Gilliland et al. 2000; Boley et al. 2021). However, Mortier et al. (2013a) also argue that planet statistics at low metallicity are too small to discriminate between a linear function and a power-law.

Transiting Giant Planets The planet-metallicity correlation has also been identified for transiting planets. The biggest challenge in identifying this correlation lies in characterizing stellar properties, in particular for non-planet host stars. The volume of stars searched for transiting planets is much larger than in radial velocity surveys

² Note that [Fe/H] is the logarithm of the iron abundance.

- both in terms of absolute numbers and galactic distance – and characterization of stellar properties with high spectral resolution observations requires a significant investment in observing time. For this reason, most surveys have focused on characterizing planet-hosting stars (Everett et al. 2013; Buchhave et al. 2014; Johnson et al. 2017).

A giant planet-metallicity relation in the *Kepler* survey was first identified based on photometry by Schlaufman and Laughlin (2011), who find that giant planet hosts have systematically redder colors than non-planet hosts, consistent with a metallicity increase in 0.2 dex. Spectroscopic characterization of exoplanet host stars show that planets larger than 4 R_{\oplus} are preferentially found around stars with a super-solar metallicity of 0.15–0.18 dex (Buchhave et al. 2012, 2014; Winn et al. 2017). This result was confirmed by Everett et al. (2013) who found that giant planets only occur around high-metallicity stars ([Fe/H] > -0.05 dex). Wang and Fischer (2015) found 10 times more planets around metal-rich stars based on photometric metallicities, consistent with a power-law index $\beta = 2$ as found in radial-velocity surveys.

Occurrence rate calculations subsequently confirmed the giant planet metallicity relation, using both medium-resolution spectroscopic metallicities from LAMOST (Mulders et al. 2016) and photometric metallicities for the non-planet hosts (e.g. Petigura et al. 2018). The increased occurrence for giant planets has also been detected in ground-based transit data (Osborn and Bayliss 2020), while the hot Jupiter sample from the Transiting Exoplanet Survey Satellite, *TESS*, is still under construction (e.g. Yee et al. 2023).

Dwarfs and Giants The giant planet-metallicity correlation is also present in stars with lower and higher masses than the sun. Low-mass M dwarfs ($\leq 0.5M_{\odot}$) are found to be enhanced in metallicity when they host giant planets (Bonfils et al. 2007; Johnson and Apps 2009; Rojas-Ayala et al. 2012; Terrien et al. 2012). The exponent of the occurrence rate-metallicity correlation, in the range $\beta = [1.26, 2.94]$, is consistent with that of sun-like stars (Neves et al. 2013). The planet-metallicity correlation is less statistically robust than for FGK dwarfs due to a lower number of planet detections (Schlaufman and Laughlin 2010; Gaidos and Mann 2014).

Giant and sub-giant stars that have evolved off the main sequence provide an opportunity to measure planet occurrence rates around higher mass stars ($\gtrsim 1.5M_{\odot}$). The giant planet-metallicity correlation is less well established for these evolved stars than for main-sequence stars. Hekker and Melendez (2007) found the first indications that evolved planet hosts are more metal-rich than non-planet hosts. Subsequent studies did often not find a planet-metallicity correlation (Pasquini et al. 2007; Takeda et al. 2008; Mortier et al. 2013b), showed mixed results (Maldonado et al. 2013; Jofré et al. 2015), or did recover a correlation (Wittenmyer et al. 2017). Limiting this chapter to planet occurrence rate studies, i.e. those that take into account detection efficiency and sample selection, the planet occurrence rate is found to increase with stellar metallicity (Johnson et al. 2010; Reffert et al. 2015; Jones et al. 2016).



Fig. 3 Metallicity of planet host stars as function of planet radius. Points represent spectroscopic metallicities of *Kepler* exoplanet hosts from Buchhave et al. (2014). The average host star metallicity correlates with planet radius, as indicated for a set of discrete radius bins shown in orange (Buchhave et al. 2014) and for a continuous planet radius-metallicity relation (Schlaufman 2015) shown with the dashed purple line. The expected range of planet radii from *In Situ* planet formation models by Dawson et al. (2015) are shown in cyan.

A Wide Range of Stellar Metallicity for sub-Neptunes

Planets smaller than Neptune form around stars with a wide range of metallicities (Sousa et al. 2008; Buchhave et al. 2012). The planet-metallicity correlation identified for giant planets disappears when considering smaller planets (Fig 3, Buchhave et al. 2014).

Neptunes The first indications that Neptune-mass planets are not preferentially found around metal rich stars, as opposed to giant planet hosts, were found by Udry et al. (2006) in a sample including M dwarfs planet hosts, and later confirmed by Sousa et al. (2008). The possibility that a higher planet occurrence rate of Neptune-sized planets around M dwarfs contributed to this correlation was investigated by Ghezzi et al. (2010), who recovered the wide range of stellar metallicities for Neptune-mass planet hosts in a sample of FGK dwarfs. This trend was confirmed by Mayor et al. (2011), who show that planets less massive than 30-40 M_{\oplus}

are equally common around metal-poor and metal-rich stars. The same metallicityindependence was found for M dwarfs hosting Neptune mass and smaller planets (Rojas-Ayala et al. 2012; Neves et al. 2013).

Transiting sub-Neptunes The large number of planets smaller than Neptune discovered by the *Kepler* mission provide a unique opportunity to constrain the metallicity-dependence of planets down to Earth-sizes. Follow-up high resolution spectroscopy of *Kepler* exoplanet hosts confirm that sub-Neptunes form around a wide range of stellar metallicities ($[Fe/H] \approx [-0.6 - 0.5]$) and extend this trend to Earth-sized planets (Buchhave et al. 2012; Everett et al. 2013; Mann et al. 2013).

Buchhave et al. (2014) divided the sample into rocky planets ($R < 1.7R_{\oplus}$) and gas dwarfs ($1.7R_{\oplus} < R < 3.9R_{\oplus}$) and find that the mean metallicity of rocky planets is consistent with solar. On the other hand, the larger gas dwarfs have a mean metallicity of [Fe/H] = 0.05 that is significantly higher than non-planet hosting stars (Buchhave and Latham 2015). Such a trend is consistent with a planet-metallicity correlation for the maximum size/mass of Neptunes (Courcol et al. 2016; Petigura et al. 2017). However, Schlaufman (2015) and Lu et al. (2020) argue that the *Kepler* data is better described by a continuous increase in metallicity with planet radius (Figure 3), though Kutra et al. (2021) argue against such a relation.

Planet occurrence rates as a function of spectroscopic metallicity were calculated by Mulders et al. (2016) for a sample of 20,000 *Kepler* target stars with medium resolution spectroscopy from Frasca et al. (2016). They find no difference in the occurrence rate of sub-Neptunes as a function of metallicity, except at orbital periods smaller than 10 days (see also Wilson et al. 2018; Petigura et al. 2018). This elevated occurrence rate at short orbital periods is consistent with the higher detection frequency of sub-Neptunes around metal-rich stars (Wang and Fischer 2015; Zhu et al. 2016).

Several other papers have pointed out trends in host star metallicity with the planet orbital period distribution (Beaugé and Nesvorný 2013; Adibekyan et al. 2013; Dawson et al. 2015; Dong et al. 2018), though there is some disagreement on the planet radius and orbital period where these transitions occur. The trend identified by Adibekyan et al. (2016) that small ($< 2 R_{\oplus}$) planets interior to the habitable zone may predominantly found in low-metallicity stars is tantalizing, but was found to not be significant when taking into account detection completeness by Mulders et al. (2016).

Trends With Stellar Mass

The correlation between planet occurrence and stellar mass is dependent on planet size. Giant planets occur more frequently around higher-mass stars (Fig. 4, Johnson et al. 2010; Fulton et al. 2021), with a linear dependence that is weaker than the quadratic dependence on metallicity. Sub-Neptunes, those found in abundance

with the *Kepler* survey, occur more frequently around low-mass M dwarfs (Fig 5, Mulders et al. 2015b; Sabotta et al. 2021).



Fig. 4 Giant planet occurrence as function of stellar mass, from Johnson et al. (2010) figure 4. The histogram shows the observed planet occurrence rate. The red red line show the predicted planet occurrence rate based on the metallicity distribution of stars in each stellar mass bin. The blue line shows the stellar-mass dependence at solar metallicity, compare to the predicted relation from the planet formation model by Kennedy and Kenyon (2008). Figure reproduced from Johnson et al. (2010) with permission from the authors.

Giant planets

Giant planets are found less frequently are low-mass M dwarfs than around sun-like stars and more frequently around evolved stars with higher masses (Fig. 4). The giant planet occurrence scales roughly linear with stellar mass, $f_{\text{giant}} \propto M_{\star}$, and is therefore weaker than the planet-metallicity correlation that scales quadratically as $f_{\text{giant}} \propto [\text{Fe}/\text{H}]^2$.

Tentative evidence for a decreased giant planet occurrence around M dwarfs compared to sun-like stars was found by Laws et al. (2003) and Endl et al. (2006).

The giant planet occurrence rate within 2.5 au increases by a factor of \sim 3 from M stars to sun-like stars (Butler et al. 2006; Cumming et al. 2008). Planet occurrence rates for a sample of late K dwarfs support the positive correlation with stellar mass (Gaidos et al. 2013).

Taking metallicity into account, the giant planet occurrence rate increases roughly linear with stellar mass between M dwarfs, GK stars, and retired A stars (Johnson et al. 2007, 2010; Ghezzi et al. 2018, Fig. 4). The trend is also identified in radial velocity samples of MKGF stars (Fulton et al. 2021; Sabotta et al. 2021). The stellar-mass dependence has also been identified for giant planets at longer orbital periods by including radial-velocity trends and micro-lensing data (Montet et al. 2014; Clanton and Gaudi 2014). The planet occurrence rate around giant stars increases with stellar mass up to $\approx 2M_{\odot}$ but decreases at larger stellar mass (Reffert et al. 2015; Jones et al. 2016; Wolthoff et al. 2022).

The giant planet occurrence rate in the *Kepler* transit surveys is low, consistent with the predictions from radial velocity surveys (Dressing and Charbonneau 2013). The occurrence rate of giant planets with orbital periods less than 50 days is more than two times higher for FGK stars than M stars (Mulders et al. 2015b). Using TESS data, Gan et al. (2023) and Bryant et al. (2023) show that he occurrence of transiting giant planets decreases with stellar mass from G dwarfs to early M dwarfs to late M dwarfs. Somewhat surprisingly, the hot Jupiter occurrence rate from TESS also falls towards F and A stars (Beleznay and Kunimoto 2022; Gan et al. 2023). This suggests a peak Hot Jupiter occurrence around one solar mass, in contrast to the colder giant planets from radial velocity surveys peaking around two solar masses (e.g. Wolthoff et al. 2022).

The Sub-Neptune Exoplanet Population

Planet smaller than Neptune far outnumber their larger counterparts (e.g. Howard et al. 2010, 2012). These sub-Neptunes are show a different dependence on host star mass than giant planets: they become more frequent towards lower mass stars (Fig. 5). Neptune-mass and smaller planets are commonly found around M dwarfs in radial velocity surveys, where the smaller mass ratio between star and planet favors planet detection compared to FGK stars. The sub-Neptune exoplanet population is most constrained by *Kepler*, whose detection efficiency reaches down to earth radii and smaller at short orbital periods, supported by *TESS* and radial velocity surveys.

High planet occurrence around low-mass stars The increase in exoplanet occurrence with decreasing effective temperature (a proxy of stellar mass) was discovered by Howard et al. (2012), see also Figure 5. Taking into account differences in detectability between stars of different sizes in the *Kepler* survey, they find that the occurrence rate of planets between 2-4 R_{\oplus} is anti-correlated with effective temperature and increases by a factor 7 between the hottest stars in the sample (late F stars) and the coolest stars (early M dwarfs). This trend was extended down to Earth-sized



Fig. 5 Overview of planet occurrence rates as a function of effective temperature in the literature for planets between $1-4R_{\oplus}$ and P < 50 days. Occurrence rates were re-scaled assuming uniform occurrence in log period and log radius for purpose of this comparison. An increase in planet occurrence from spectral types F to early M is present across all studies. A break in this trends becomes visible towards late M dwarfs, though it should be noted those rates sample only short-period planets (< 10 days) and are potentially lower limits to the occurrence out to 50 days. References – Howard et al. (2012); Mulders et al. (2015b); Hardegree-Ullman et al. (2019); Yang et al. (2020); He et al. (2021); Sabotta et al. (2021); Ment and Charbonneau (2023); Bergsten et al. (2023).

planets by Mulders et al. (2015a), who found an increase in planet occurrence rate between F,G,K, and M type stars at all orbital periods. The occurrence rate of (early) M stars compared to FGK stars is a factor ~ 2-4 higher at planet radii between 1 and ~ $3R_{\oplus}$ (Mulders et al. 2015b; Gaidos et al. 2016; Zink et al. 2023). Radial velocity surveys show a similar increase in the planet occurrence rate around M dwarfs Sabotta et al. (2021). Taking into account system architectures, Yang et al. (2020) and He et al. (2021) show it is the fraction of stars with planetary systems that increases towards lower mass stars, while the number of planets per planetary system remains roughly constant.

Figure 5 shows the occurrence of rate of sub-Neptunes $(1 - 4 R_{\oplus})$ at orbital periods less than 50 days as a function of stellar effective temperature as estimated by different studies. For purposes of this comparison, occurrence rates were rescaled when only estimates for a different range of planet properties were available, assuming a uniform occurrence in log planet radius and log orbital period. While there is significant scatter in occurrence rates at similar effective temperatures, the elevated

planet occurrence rates around M dwarfs compared to FGK stars is clearly present, as well as the more gradual decrease in planet occurrence rate within the FGK stars.



Fig. 6 Radial velocity detected exoplanets around higher-mass (left) and lower mass (right) M dwarfs from the CARMENES survey (Sabotta et al. 2021; Ribas et al. 2023). Aside from a decrease in the number of giant planets, sub-Neptunes tend to be smaller and located closer to the star for lower-mass stars. Figure based on (Sabotta et al. 2021, Fig. 3) with updated data as in (Ribas et al. 2023, Fig. 10).

Trends within M dwarfs M dwarfs are not a homogenous group of stars, but span a large range of stellar masses, radii, and luminosities. Exoplanet properties also vary significantly across this range (see Fig. 6), though it can be hard to characterize the population around the lowest mass M dwarfs due to a lack of stars sufficiently bright for exoplanet detection. At the low-mass end of its stellar population, the Kepler survey was mainly sensitive to early M and late K dwarfs (Dressing and Charbonneau 2015; Bergsten et al. 2023). A small number of even lower-mass mid M dwarfs where also observed, with a planet occurrence rate at orbital periods less than 10 days consistent with that of the early M dwarfs (Hardegree-Ullman et al. 2019). TESS, being an all sky survey, has observed a larger number of M dwarfs, including many lower mass mid and late M dwarfs. Focusing on the nearest and most-well characterized mid-to-late M dwarfs, Ment and Charbonneau (2023) find a high occurrence ($\approx 60\%$) of super-earth sized planets ($< 1.5 R_{\oplus}$) within an orbital period of 7 days. However, the authors find a lack of mini-Neptunes (> 1.5 R_{\oplus}), presenting a break with the ever-increasing sub-Neptune occurrence observed in F, G, K, and early M dwarfs (see Fig. 5).

A similar trend is seen in the CARMENES radial velocity survey of M dwarfs (Fig. 6, Sabotta et al. 2021). Around the more massive M dwarfs, sub-Neptunes have masses between 2 and 20 M_{\oplus} and are mostly located at orbital periods between 10–100 days (left panel of Fig. 6), coinciding with regions of high planet occurrence of transiting sub-Neptunes around early M and FGK stars (e.g. Mulders et al. 2015a). Around less massive M dwarfs (< 0.35 M_{\odot}) planets are on average less massive (1 – 10 M_{\oplus}) and located mostly within an orbital period of 10 days (right panel of Fig. 6).

As such, the planet population around late M dwarfs starts to resemble the archetype of the Trappist-1 system (Gillon et al. 2017), with many (super)Earthsized planets at short orbital periods, but lacking the larger mini-Neptunes at somewhat longer periods so prominent around early M dwarfs and FGK stars.

Constraints On Planet Formation Mechanisms

The dependence of the exoplanet population on host star properties provides constraints on planet formation mechanisms. The positive correlations of giant planet occurrence rate with stellar mass and metallicity support the core accretion scenario of giant planet formation. The constraints provided by the lack of a clear correlation for sub-Neptunes with stellar metallicity and the anti-correlation with stellar mass have yet to be determined, but point to the role of *Pebble Accretion* in the planet formation process. These trends indicate that planet formation is a robust and efficient process that takes place in a variety of environments.

Formation of Giants Planets

The core accretion scenario postulates that giant planets form "bottom up" with the formation of a $\sim 10M_{\oplus}$ solid core followed by a subsequent phase where most of the gas is accreted (Pollack et al. 1996). As the envelope has to be accreted before the protoplanetary disk gas is dispersed, typically ~ 3 million years (e.g Mamajek 2009), the growth of the core has to be sufficiently rapid to allow giant planets to form. Pebble accretion (Ormel and Klahr 2010; Lambrechts and Johansen 2012) is often invoked to facilitate this rapid growth of the core, see Drążkowska et al. (2023) for an extended review. The time scale for core growth depends on the amount of material locally available in the disk, i.e. the solid surface density. Giant planets thus form only in protoplanetary disks with a sufficiently high surface density of solids (e.g. Ikoma et al. 2000; Kokubo and Ida 2002).

The stellar metallicity is a tracer of the solid inventory in protoplanetary disks at the onset of planet formation. Stars and protoplanetary disks inherit the same metallicity from the parental molecular cloud. Stars with a high metallicity formed with disks with a high solid surface density, and are therefore more likely to form giant planets. Numerical simulations of core formation and envelope accretion in disk with different metallicities consistently reproduce the observed giant planetmetallicity correlation (e.g. Ida and Lin 2004; Kornet et al. 2005; Ida and Lin 2008a; Mordasini et al. 2009b).

A similar argument can be made for the dependence of the giant exoplanet population on stellar mass. Protoplanetary disks mass, both gas and solids, scales with stellar mass (see Fig. 7), while giant planets are more likely to form in more massive disks (e.g. Thommes et al. 2008). By extension, the core accretion model predicts a positive correlation between giant planet occurrence and stellar mass. Based on analytical estimates, Laughlin et al. (2004) predict fewer giant planets around M dwarfs. Detailed numerical simulations show a nearly linear dependence of giant planet occurrence on stellar mass (Ida and Lin 2005; Kennedy and Kenyon 2008; Alibert et al. 2011), consistent with the observed trends (Figure 4, however, see Schlecker et al. 2022).

Gravitational instability In the gravitational instability scenario, giant planets form "top down" from the contracting gas in massive protoplanetary disks (Boss 1997). This formation mechanism predicts different dependence on stellar mass and metallicity. A high disk metallicity inhibits cooling and contraction of the gaseous envelope, and therefore giant planets should form more efficiently around low-metallicity stars (Meru and Bate 2010). Gravitational instabilities should also form planets efficiently around M dwarfs (Boss 2006). The observed positive correlations between giant planet occurrence with stellar mass and metallicity indicate that planets at short orbital periods likely did not form through gravitational instability in a protoplanetary disks.

Increasing Stellar Metallicity by Accretion of Planets Accretion of planets can increase the stellar metallicity if planets are more metal-rich than their host star. It was initially suggested that the enhanced metallicity of planet-hosting stars is caused by the accretion of planets or solids (Gonzalez 1997), instead of planet formation being more efficient around more metal-rich stars. The observational signature of planetary accretion is only large enough if the accreted metals are not mixed throughout the entire star, but remain near the surface in the convective zone. In F and A stars, the convective zone is thin enough that the accretion of solids can lead to a metallicity increase that is consistent with observations (Laughlin and Adams 1997). For lower-mass stars the convective zones are deeper and the metallicity signature of accreted planets should drop below detectable levels for G type and earlier stars (Laughlin and Adams 1997). This prediction is inconsistent with the observed giant-planet metallicity relation for these stars (e.g. Fischer and Valenti 2005) as well as for M dwarfs (e.g. Neves et al. 2013). Once stars evolve off the main-sequence, mixing should increase, thereby diluting the metallicity enhancement from planetary accretion. However, the planet metallicity correlation is also observed in evolved stars (e.g. Johnson et al. 2010; Reffert et al. 2015; Jones et al. 2016). Hence, the hypothesis that planetary accretion causes the planet-metallicity correlation is no longer supported by observational evidence.

Formation of sub-Neptunes

The different scaling laws with stellar mass and metallicity indicate a different formation history for giant planets and sub-Neptunes. Indeed, the comparison between the predictions of the core accretion model (Ida and Lin 2008b; Mordasini et al. 2009a) with the population of sub-Neptunes detected in radial velocity surveys (Howard et al. 2010) and the *Kepler* transit survey (Howard et al. 2012) show that the predicted "planet desert" at orbital period less than 50 days is indeed well-populated, highlighting the need to amend planet formation theory for sub-Neptunes (e.g. Emsenhuber et al. 2021; Mulders et al. 2019).

The moniker of 'core accretion' is not particularly useful when discussing sub-Neptunes as they are, almost by definition, the planets that did not accrete massive gaseous envelopes. The planet formation mechanisms discussed here are almost exclusively focused on sub-Neptunes and it should be kept in mind that these new mechanisms are to amend, not replace, core accretion theory.

Several planet formation mechanisms have been proposed to explain the presence of small planets at short orbital periods (e.g. Raymond et al. 2008). The two mechanisms that are of most relevance here are *In Situ* formation and *Planet Migration*. In addition, *Pebble Accretion*, combined with inward radial drift, is increasingly invoked as a mechanism to form close-in sub-Neptunes (see Drążkowska et al. 2023 for a review).

In Situ Formation The *In Situ* Formation scenario for exoplanets is based on terrestrial planet formation in the Solar System. Planetary embryos in the protoplanetary disk can grow through oligarchic growth to a fraction of the final planet mass, typically Mars-size at 1 au (e.g. Lissauer 1987; Kokubo and Ida 2000, 1998) After the gas disk disperses, gravitational interactions increase the protoplanet eccentricities and makes them collide and merge, leading to a phase of giant impacts in which planets grow to their final masses (e.g. Chambers and Wetherill 1998; Wetherill 1985). As the majority of the accreted material is sourced from a region close to the planets final orbit, the planet mass is directly dependent on the local surface density of planetary building blocks (Kokubo and Ida 2002). Chiang and Laughlin (2013) proposed that planetary systems observed with *Kepler* could have formed *In Situ* in disks that are on average more massive than the protoplanetary disk around the sun. N-body simulations of the giant impact phase show that disks with high surface density of solids in the inner regions can indeed form Kepler-like planetary systems (Hansen and Murray 2012, 2013; MacDonald et al. 2020; Mulders et al. 2020).

Pebble Accretion The *Pebble Accretion* hypothesis is based on the rapid growth of planetary embryos through aerodynamically assisted accretion (Ormel and Klahr 2010; Lambrechts and Johansen 2012). Combined with inward radial drift of solids which brings a large amount of pebbles into the inner disk, super-earths can directly form (e.g. Lambrechts et al. 2019). When planetary cores grow massive enough to perturb the surrounding gas, a pressure bump can form that limits the subsequent growth by pebbles. The mass where this happens is called the pebble isolation mass (Lambrechts and Johansen 2014), which in typically in the super-earth or sub-Neptune regime (Liu et al. 2019).

Planet Migration The Planet Migration hypothesis is built on he theoretical expectation that low-mass planets embedded in a gaseous disk undergo rapid inward migration (Type-I migration, Ward 1997). Because planetary embryos can grow to larger sizes in the outer disk where more material is available, *Planet Migration*

does not require disks to be particularly massive (e.g. Swift et al. 2013). The type-I migration time scales are short ($< 10^5$ years) compared to the disk life time of a few million years (e.g. Mamajek 2009), and migration needs to be halted in the inner disk. Possible mechanisms to stall migration include an inner disk cavity (e.g. Terquem and Papaloizou 2007), resonant capture by other planets, and regions of outward migration due to disk density and temperature structure (Dittkrist et al. 2014; Cossou et al. 2014). The largest challenge for planet migration hypothesis is that the observed multi-planet systems are often not in orbital resonances as predicted from convergent migration (Fabrycky et al. 2014), though different mechanisms have been proposed to break resonances after formation (Rein 2012; Goldreich and Schlichting 2014; Izidoro et al. 2017).

These three scenarios are not mutually exclusive. *Planet Migration* models often include a growth phase including pebble accretion, and a giant impact phase during or after migration (e.g. Izidoro et al. 2021). *In Situ* formation models often invoke, explicitly or implicitly, an inward migration phase of solids to increase the amount of planetary building blocks in the inner disk (e.g. Hansen and Murray 2012), similar to the radial drift of pebbles (Moriarty and Fischer 2015; Voelkel et al. 2020). Despite these nuances, *Planet Migration, In Situ* and *Pebble Accretion* remain useful concepts in discussing the origin the observed trends with stellar mass and metallicity.

Metallicity Dependence The stellar metallicity is a direct measure of the amount of condensible solids that was available for planet formation in the disk. The base expectation is that the mass in planetary systems correlates positively with disk metallicity. The In Situ formation simulations in Dawson et al. (2015) show that, for a range of metallicity of a factor 10, the predicted planet radii vary between $1-4R_{\oplus}$, with significant scatter (see also Figure 3). A clear planet size-metallicity relation is not seen in the observed population of small exoplanets. There is tentative evidence for a lack of rocky $(\langle 2R_{\oplus})$ planets at high metallicities at a limited orbital period range (Dawson et al. 2015; Adibekyan et al. 2016), though this trend may not be statistically significant when taking into account survey completeness (Mulders et al. 2016). The predicted lack of sub-Neptunes $(2 - 4R_{\oplus})$ at low metallicity is not observed. However, a planet size-metallicity relation appears to be present for planets more massive than Neptune (Courcol et al. 2016; Petigura et al. 2017). The elevated host star metallicity of transiting sub-Neptunes (Buchhave et al. 2014; Buchhave and Latham 2015) seems to support *In Situ* formation scenario, perhaps with a much wider range in planet radii than predicted by (Dawson et al. 2015), originating from a wide range in disk masses (Kutra et al. 2021). On the other hand, the planet size-metallicity relation inferred for Kepler planets by Schlaufman (2015) is significantly shallower than linear, see Figure 3. Hence, it is clear that the planetmetallicity correlation predicted by *In Situ* formation models is not observed.

In the *Planet Migration* scenario, the mass of planets that form in outer disk is also dependent on disk metallicity. However, the subsequent inward migration may shape the observed distribution of exoplanets in a different way. Cossou et al. (2014) find that super-earths consistently form in a set of simulations varying the dust-to-

gas ratio, a proxy of metallicity, by a factor 4. The total mass of planetary systems show an almost linear dependence on metallicity, and hence does not deviate significantly from the predictions of *In Situ* formation models. Coleman and Nelson (2016) model the growth and migration of super-earths, and find that planets in low-metallicity disks do not reach the mass required for efficient inward migration, and hence close-in super-earths do not form. Instead small mobile bodies (pebbles) must play an important role in the formation of super-earths, though the predictions of such a model have not been explored in detail.

In the *Pebble Accretion* scenario, the pebble isolation mass could potentially play a role at erasing any metallicity dependence. Because this mass scales with gas disk properties, and not with the amount of solids available, an increased metallicity does not directly affect planet size, limiting the role of metallicity in determining final planet properties.



Fig. 7 Average solid mass of planetary systems around stars of different masses. This quantity is calculated by multiplying the planet occurrence rate by an estimate of the planet mass, see text for details. Sub-Neptunes at short orbital periods (blue, purple) are compared to giant planets out to a few au (red) and the median dust masses of Class II protoplanetary disks in the Chamaeleon I star forming region, where the two different slopes of the two lines reflect some of the uncertainties in the derived stellar-mass dependence. An excess of solids in sub-Neptunes around M dwarfs is evident, posing a riddle for planet formation models.

Low Mass Stars The anti-correlation between the occurrence of planets at short orbital period and the stellar mass poses an urgent problem for planet formation theories: How to explain the elevated planet occurrence rates of low mass stars if less material is available in their disks to form planets? Figure 7 illustrates this issue, showing the estimated amounts of solids around stars of different masses in:

- Planets at orbital periods less than 150 days around F, G, K, and early M dwarfs in the *Kepler* survey, from Mulders et al. (2015b), estimated assuming the mass-radius relation from Wolfgang et al. (2016) and a core mass of $20 M_{\oplus}$ per giant planet.
- Transiting planets around mid-to-late M dwarfs observed with TESS from Ment and Charbonneau (2023), assuming an average planet mass of 3 M_{\oplus} .
- Radial velocity detected sub-Neptunes around M dwarfs from Sabotta et al. (2021); Ribas et al. (2023), using an average mass of 3 M_{\oplus} for late M dwarfs and 6 M_{\oplus} for early M dwarfs.
- Giant planets out to 2.5 au around GK stars and retired AF stars from Johnson et al. (2010), assuming $60M_{\oplus}$ of solids per Jupiter-mass giant planet (Thorngren et al. 2016).
- Giant planets around M dwarfs from Ribas et al. (2023) assuming 30 M_{\oplus} of solids per giant planet.
- Protoplanetary disks in the Chamaeleon I star forming region, from Pascucci et al. (2016), probing solids at scales of $\sim 10 100$ au.

While the solids in giant planets and protoplanetary disks show a positive scaling with stellar mass, this relation breaks down for sub-Neptunes at short orbital periods. The estimated amount of solids in M dwarf planetary systems is higher than that in sun-like stars (Mulders et al. 2015b; Gaidos 2017), reflecting the trend in planet occurrence.

A strict *In Situ* formation model for sub-Neptunes, where planet mass is directly related to mass available in disk (e.g. Raymond et al. 2007; Ciesla et al. 2015), is not favored by stellar-mass dependencies as evident from Figure 7. Models that include *Planet Migration* or *Pebble Accretion* move material around radially in the disk, leading to different efficiencies of planet formation at different stellar masses. Figure 8 shows the predictions planet occurrence rates of inner sub-Neptunes and outer giant planets as function of stellar mass.

The planet population synthesis model using planetesimal accretion and planet migration from Burn et al. (2021) show positive correlation between planet occurrence and stellar mass for both sub-Neptunes and Giant planets, the former inconsistent with observed trends. Two *Pebble Accretion* models better explain the occurrence of exoplamets as function of stellar mass. (Mulders et al. 2021) use a growing giant planets outside the snow line to block the flow of pebbles into the inner disk, suppressing the growth of super-earths. Because giant planet form more efficiently around the more massive stars, the occurrence of sub-Neptunes is suppressed more for more massive stars, explaining the observed downturn in planet occurrence towards higher stellar mass. In the model of (Chachan and Lee 2023), planet formation in the inner disk through pebble accretion is most efficient around stars between $0.3 - 0.5 M_{\odot}$, predicting a peak in the planet occurrence for early M dwarfs and a lower occurrence for more massive stars (See fig. 8).



Fig. 8 Relative planet occurrence as function of stellar mass predicted by different planet formation models. The green line shows the Bern planet population synthesis model (Burn et al. 2021) based on planetesimal accretion and planet migration. The orange line (Mulders et al. 2021) and purple line (Chachan and Lee 2023) show pebble accretion models without migration. While all models reproduce the increase in giant planet occurrence rate with stellar mass (left panel), only the pebble accretion models show the observed decrease in the occurrence of sub-Neptunes from late M to F stars.

Both pebble accretion models from predict the observed downturn in planet occurrence toward late M dwarfs (Sabotta et al. 2021; Ment and Charbonneau 2023), though it should be noted that that the model from Burn et al. (2021) predicts a similar scaling relation in this stellar mass range. These pebble accretion models also predict the observed decrease in planet mass for late M dwarfs (see also Liu et al. 2019).

Conclusions

Giant planets occur more frequent around more massive and more metal-rich stars. These trends support the core-accretion scenario for giant planet formation in which accretion of a gaseous envelope starts after a sufficiently rapid assembly of a massive rocky core. The threshold for reaching the critical core mass is reached more easily in protoplanetary disks with a larger amount of condensible solids around metal-rich stars and in more massive disks around more massive stars.

These results stand in contrast to the population of exoplanets smaller than Neptune, those that are found in abundance with *Kepler* and *TESS* and represent the bulk of the exoplanet population. These sub-Neptunes are found around stars with a wide range of metallicities, indicating that planet formation is a robust process that occurs efficiently in a variety of environments. Curiously, sub-Neptunes occur much more frequently around low-mass M dwarfs than around solar-mass stars, though this trend reverses toward the lowest-mass stars, late M dwarfs. Pebble drift and accretion likely plays a key role in boosting the planet formation efficiency for M dwarfs, explaining the abundance of exoplanets around low-mass stars.

Acknowledgements I would like to thank Ignasi Ribas and Silvia Sabotta for providing Figure 6. G.D.M. acknowledges support from FONDECYT project 11221206, from ANID — Millennium Science Initiative — ICN12_009, and the ANID BASAL project FB210003. The results reported herein benefitted from collaborations and/or information exchange within NASA's Nexus for Exoplanet System Science (NEXSS) research coordination network sponsored by NASA's Science Mission Directorate and project "Alien Earths" funded under Agreement No. 80NSSC21K0593

References

- Adibekyan V, Figueira P Santos NC (2016) Which Type of Planets do We Expect to Observe in the Habitable Zone? Origins of Life and Evolution of Biospheres 46(4):351–359
- Adibekyan VZ, Figueira P, Santos NC et al. (2013) Orbital and physical properties of planets and their hosts: new insights on planet formation and evolution. Astronomy & Astrophysics 560:A51
- Alibert Y, Mordasini C Benz W (2011) Extrasolar planet population synthesis. III. Formation of planets around stars of different masses. Astronomy & Astrophysics 526:A63
- Beaugé C Nesvorný D (2013) Emerging Trends in a Period-Radius Distribution of Close-in Planets. The Astrophysical Journal 763(1):12
- Beleznay M Kunimoto M (2022) Exploring the dependence of hot Jupiter occurrence rates on stellar mass with TESS. MNRAS516(1):75–83

Bergsten GJ, Pascucci I, Hardegree-Ullman KK et al. (2023) No Evidence for More Earth-sized Planets in the Habitable Zone of Kepler's M versus FGK Stars. AJ166(6):234

- Boley KM, Wang J, Zinn JC et al. (2021) Searching For Transiting Planets Around Halo Stars. II. Constraining the Occurrence Rate of Hot Jupiters. AJ162(3):85
- Bonfils X, Mayor M, Delfosse X et al. (2007) The HARPS search for southern extra-solar planets. X. A m sin i = 11 Mearth planet around the nearby spotted M dwarf GJ 674. Astronomy & Astrophysics 474(1):293–299

Boss AP (1997) Giant planet formation by gravitational instability. Science 276(5320):1836–1839

- Boss AP (2006) Rapid Formation of Gas Giant Planets around M Dwarf Stars. The Astrophysical Journal 643(1):501–508
- Bryant EM, Bayliss D Van Eylen V (2023) The occurrence rate of giant planets orbiting low-mass stars with TESS. MNRAS521(3):3663–3681
- Buchhave LA Latham DW (2015) The Metallicities of Stars with and without Transiting Planets. The Astrophysical Journal 808(2):187
- Buchhave LA, Batalha N, Latham DW et al. (2012) An abundance of small exoplanets around stars with a wide range of metallicities. Nature 486(7):375–377
- Buchhave LA, Bizzarro M, Latham DW et al. (2014) Three regimes of extrasolar planet radius inferred from host star metallicities. Nature 509(7):593–595
- Burn R, Schlecker M, Mordasini C et al. (2021) The New Generation Planetary Population Synthesis (NGPPS). IV. Planetary systems around low-mass stars. A&A656:A72
- Butler RP, Johnson JA, Marcy GW et al. (2006) A Long-Period Jupiter-Mass Planet Orbiting the Nearby M Dwarf GJ 849 118(8):1685–1689
- Chachan Y Lee EJ (2023) Small Planets around Cool Dwarfs: Enhanced Formation Efficiency of Super-Earths around M Dwarfs. ApJ952(1):L20

- Chambers JE Wetherill GW (1998) Making the Terrestrial Planets: N-Body Integrations of Planetary Embryos in Three Dimensions. Icarus 136(2):304–327
- Chiang E Laughlin G (2013) The minimum-mass extrasolar nebula: in situ formation of close-in super-Earths. Monthly Notices of the Royal Astronomical Society 431(4):3444–3455
- Ciesla FJ, Mulders GD, Pascucci I Apai D (2015) Volatile Delivery to Planets from Water-rich Planetesimals around Low Mass Stars. The Astrophysical Journal 804(1):9
- Clanton C Gaudi BS (2014) Synthesizing Exoplanet Demographics from Radial Velocity and Microlensing Surveys. II. The Frequency of Planets Orbiting M Dwarfs. The Astrophysical Journal 791(2):91
- Coleman GAL Nelson RP (2016) On the formation of compact planetary systems via concurrent core accretion and migration. Monthly Notices of the Royal Astronomical Society 457(3):2480–2500
- Cossou C, Raymond SN, Hersant F Pierens A (2014) Hot super-Earths and giant planet cores from different migration histories. Astronomy & Astrophysics 569:A56
- Courcol B, Bouchy F Deleuil M (2016) An upper boundary in the mass-metallicity plane of exo-Neptunes. Monthly Notices of the Royal Astronomical Society 461(2):1841–1849
- Cumming A, Butler RP, Marcy GW et al. (2008) The Keck Planet Search: Detectability and the Minimum Mass and Orbital Period Distribution of Extrasolar Planets. Publications of the Astronomical Society of Pacific 120(8):531–
- Dawson RI, Chiang E Lee EJ (2015) A metallicity recipe for rocky planets. Monthly Notices of the Royal Astronomical Society 453(2):1471–1483
- Dittkrist KM, Mordasini C, Klahr H, Alibert Y Henning T (2014) Impacts of planet migration models on planetary populations. Effects of saturation, cooling and stellar irradiation. Astronomy & Astrophysics 567:A121
- Dong S, Xie JW, Zhou JL, Zheng Z Luo A (2018) LAMOST telescope reveals that Neptunian cousins of hot Jupiters are mostly single offspring of stars that are rich in heavy elements. Proceedings of the National Academy of Science 115(2):266–271
- Dressing C Charbonneau D (2013) The Occurrence Rate of Small Planets around Small Stars. The Astrophysical Journal 767(1):95
- Dressing C Charbonneau D (2015) The Occurrence of Potentially Habitable Planets Orbiting M Dwarfs Estimated from the Full Kepler Dataset and an Empirical Measurement of the Detection Sensitivity. The Astrophysical Journal 807(1):45
- Drą zkowska J, Bitsch B, Lambrechts M et al. (2023) Planet Formation Theory in the Era of ALMA and Kepler: from Pebbles to Exoplanets. In: Inutsuka S, Aikawa Y, Muto T, Tomida K Tamura M (eds) Protostars and Planets VII, Astronomical Society of the Pacific Conference Series, vol 534, p 717, DOI 10.48550/arXiv.2203.09759
- Emsenhuber A, Mordasini C, Burn R et al. (2021) The New Generation Planetary Population Synthesis (NGPPS). II. Planetary population of solar-like stars and overview of statistical results. A&A656:A70
- Endl M, Cochran WD, Tull RG MacQueen PJ (2003) A Dedicated M Dwarf Planet Search Using The Hobby-Eberly Telescope. The Astronomical Journal 126(6):3099–3107
- Endl M, Cochran WD, Kürster M et al. (2006) Exploring the Frequency of Close-in Jovian Planets around M Dwarfs. The Astrophysical Journal 649(1):436–443
- Everett ME, Howell SB, Silva DR Szkody P (2013) Spectroscopy of Faint Kepler Mission Exoplanet Candidate Host Stars. The Astrophysical Journal 771(2):107
- Fabrycky DC, Lissauer JJ, Ragozzine D et al. (2014) Architecture of Kepler's Multi-transiting Systems. II. New Investigations with Twice as Many Candidates. The Astrophysical Journal 790(2):146
- Fischer DA Valenti J (2005) The Planet-Metallicity Correlation. The Astrophysical Journal 622(2):1102–1117
- Frasca A, Molenda-Zakowicz J, De Cat P et al. (2016) Activity indicators and stellar parameters of the Kepler targets. An application of the ROTFIT pipeline to LAMOST-Kepler stellar spectra. Astronomy & Astrophysics 594:A39

- Fuhrmann K, Pfeiffer MJ Bernkopf J (1998) F- and G-type stars with planetary companions: upsilon Andromedae, rho (1) Cancri, tau Bootis, 16 Cygni and rho Coronae Borealis. Astronomy & Astrophysics 336:942–952
- Fulton BJ, Rosenthal LJ, Hirsch LA et al. (2021) California Legacy Survey. II. Occurrence of Giant Planets beyond the Ice Line. ApJS255(1):14
- Gaidos E (2017) A minimum mass nebula for M dwarfs. Monthly Notices of the Royal Astronomical Society: Letters 470(1):L1–L5
- Gaidos E Mann AW (2014) M Dwarf Metallicities and Giant Planet Occurrence: Ironing Out Uncertainties and Systematics. The Astrophysical Journal 791(1):54
- Gaidos E, Fischer DA, Mann AW Howard AW (2013) An Understanding of the Shoulder of Giants: Jovian Planets around Late K Dwarf Stars and the Trend with Stellar Mass. The Astrophysical Journal 771(1):18
- Gaidos E, Mann AW, Kraus AL Ireland M (2016) They are small worlds after all: revised properties of Kepler M dwarf stars and their planets. Monthly Notices of the Royal Astronomical Society 457(3):2877–2899
- Gan T, Wang SX, Wang S et al. (2023) Occurrence Rate of Hot Jupiters Around Early-type M Dwarfs Based on Transiting Exoplanet Survey Satellite Data. AJ165(1):17
- Ghezzi L, Cunha K, Smith VV et al. (2010) Stellar Parameters and Metallicities of Stars Hosting Jovian and Neptunian Mass Planets: A Possible Dependence of Planetary Mass on Metallicity. The Astrophysical Journal 720(2):1290–1302
- Ghezzi L, Montet BT Johnson JA (2018) Retired A Stars Revisited: An Updated Giant Planet Occurrence Rate as a Function of Stellar Metallicity and Mass. The Astrophysical Journal 860(2):109
- Gilliland RL, Brown TM, Guhathakurta P et al. (2000) A Lack of Planets in 47 Tucanae from a Hubble Space Telescope Search. The Astrophysical Journal 545(1):L47–L51
- Gillon M, Triaud AHMJ, Demory BO et al. (2017) Seven temperate terrestrial planets around the nearby ultracool dwarf star TRAPPIST-1. Nature 542(7):456–460
- Goldreich P Schlichting HE (2014) Overstable Librations can Account for the Paucity of Mean Motion Resonances among Exoplanet Pairs. The Astronomical Journal 147(2):32
- Gonzalez G (1997) The stellar metallicity-giant planet connection. Monthly Notices of the Royal Astronomical Society 285(2):403–412
- Gonzalez G (1998) Spectroscopic analyses of the parent stars of extrasolar planetary system candidates. Astronomy & Astrophysics 334:221–238
- Gonzalez G, Laws C, Tyagi S Reddy BE (2001) Parent Stars of Extrasolar Planets. VI. Abundance Analyses of 20 New Systems. The Astronomical Journal 121(1):432–452
- Hansen BMS Murray N (2012) Migration Then Assembly: Formation of Neptune-mass Planets inside 1 AU. The Astrophysical Journal 751(2):158
- Hansen BMS Murray N (2013) Testing in Situ Assembly with the Kepler Planet Candidate Sample. The Astrophysical Journal 775(1):53
- Hardegree-Ullman KK, Cushing MC, Muirhead PS Christiansen JL (2019) Kepler Planet Occurrence Rates for Mid-type M Dwarfs as a Function of Spectral Type. The Astronomical Journal 158(2):75
- He MY, Ford EB Ragozzine D (2021) Architectures of Exoplanetary Systems. II. An Increase in Inner Planetary System Occurrence toward Later Spectral Types for Kepler's FGK Dwarfs. The Astronomical Journal 161(1):16
- Hekker S Melendez J (2007) Precise radial velocities of giant stars. III. Spectroscopic stellar parameters. Astronomy & Astrophysics 475(3):1003–1009
- Howard AW, Marcy GW, Johnson JA et al. (2010) The Occurrence and Mass Distribution of Closein Super-Earths, Neptunes, and Jupiters. Science 330(6):653–655
- Howard AW, Marcy GW, Bryson ST et al. (2012) Planet Occurrence within 0.25 AU of Solar-type Stars from Kepler. The Astrophysical Journal Supplement 201(2):15
- Ida S Lin DNC (2004) Toward a Deterministic Model of Planetary Formation. II. The Formation and Retention of Gas Giant Planets around Stars with a Range of Metallicities. The Astrophysical Journal 616(1):567–572

Exoplanet Populations and their Dependence on Host Star Properties

- Ida S Lin DNC (2005) Toward a Deterministic Model of Planetary Formation. III. Mass Distribution of Short-Period Planets around Stars of Various Masses. The Astrophysical Journal 626(2):1045–1060
- Ida S Lin DNC (2008a) Toward a Deterministic Model of Planetary Formation. IV. Effects of Type I Migration. The Astrophysical Journal 673(1):487–501
- Ida S Lin DNC (2008b) Toward a Deterministic Model of Planetary Formation. V. Accumulation Near the Ice Line and Super-Earths. The Astrophysical Journal 685(1):584–595
- Ikoma M, Nakazawa K Emori H (2000) Formation of Giant Planets: Dependences on Core Accretion Rate and Grain Opacity. The Astrophysical Journal 537(2):1013–1025
- Izidoro A, Ogihara M, Raymond SN et al. (2017) Breaking the chains: hot super-Earth systems from migration and disruption of compact resonant chains. Monthly Notices of the Royal Astronomical Society 470(2):1750–1770
- Izidoro A, Bitsch B, Raymond SN et al. (2021) Formation of planetary systems by pebble accretion and migration. Hot super-Earth systems from breaking compact resonant chains. A&A650:A152
- Jofré E, Petrucci R, Saffe C et al. (2015) Stellar parameters and chemical abundances of 223 evolved stars with and without planets. Astronomy & Astrophysics 574:A50
- Johnson JA Apps K (2009) On the Metal Richness of M Dwarfs with Planets. The Astrophysical Journal 699(2):933–937
- Johnson JA, Butler RP, Marcy GW et al. (2007) A New Planet around an M Dwarf: Revealing a Correlation between Exoplanets and Stellar Mass. The Astrophysical Journal 670(1):833–840
- Johnson JA, Aller KM, Howard AW Crepp JR (2010) Giant Planet Occurrence in the Stellar Mass-Metallicity Plane. Publications of the Astronomical Society of Pacific 122(8):905–915
- Johnson JA, Petigura EA, Fulton BJ et al. (2017) The California-Kepler Survey. II. Precise Physical Properties of 2025 Kepler Planets and Their Host Stars. AJ154(3):108
- Jones MI, Jenkins JS, Brahm R et al. (2016) Four new planets around giant stars and the massmetallicity correlation of planet-hosting stars. Astronomy & Astrophysics 590:A38
- Kennedy G Kenyon SJ (2008) Planet Formation around Stars of Various Masses: The Snow Line and the Frequency of Giant Planets. The Astrophysical Journal 673(1):502–512
- Kokubo E Ida S (1998) Oligarchic Growth of Protoplanets. Icarus 131(1):171-178
- Kokubo E Ida S (2000) Formation of Protoplanets from Planetesimals in the Solar Nebula. Icarus 143(1):15–27
- Kokubo E Ida S (2002) Formation of Protoplanet Systems and Diversity of Planetary Systems. The Astrophysical Journal 581(1):666–680
- Kornet K, Bodenheimer P, Rozyczka M Stepinski TF (2005) Formation of giant planets in disks with different metallicities. Astronomy & Astrophysics 430(3):1133–1138
- Kutra T, Wu Y Qian Y (2021) Super-Earths and Sub-Neptunes Are Insensitive to Stellar Metallicity. AJ162(2):69
- Lambrechts M Johansen A (2012) Rapid growth of gas-giant cores by pebble accretion. Astronomy & Astrophysics 544:A32
- Lambrechts M Johansen A (2014) Forming the cores of giant planets from the radial pebble flux in protoplanetary discs. Astronomy & Astrophysics 572:A107
- Lambrechts M, Morbidelli A, Jacobson SA et al. (2019) Formation of planetary systems by pebble accretion and migration. How the radial pebble flux determines a terrestrial-planet or super-Earth growth mode. Astronomy & Astrophysics 627:A83
- Laughlin G Adams FC (1997) Possible Stellar Metallicity Enhancements from the Accretion of Planets. The Astrophysical Journal 491(1):L51–L54
- Laughlin G, Bodenheimer P Adams FC (2004) The Core Accretion Model Predicts Few Jovian-Mass Planets Orbiting Red Dwarfs. The Astrophysical Journal 612(1):L73–L76
- Laws C, Gonzalez G, Walker KM et al. (2003) Parent Stars of Extrasolar Planets. VII. New Abundance Analyses of 30 Systems. The Astronomical Journal 125(5):2664–2677
- Lissauer JJ (1987) Timescales for planetary accretion and the structure of the protoplanetary disk. Icarus (ISSN 0019-1035) 69:249–265

- Liu B, Lambrechts M, Johansen A Liu F (2019) Super-Earth masses sculpted by pebble isolation around stars of different masses. Astronomy & Astrophysics 632:A7
- Lu CX, Schlaufman KC Cheng S (2020) An Increase in Small-planet Occurrence with Metallicity for Late-type Dwarf Stars in the Kepler Field and Its Implications for Planet Formation. AJ160(6):253
- MacDonald MG, Dawson RI, Morrison SJ, Lee EJ Khandelwal A (2020) Forming Diverse Super-Earth Systems In Situ. The Astrophysical Journal 891(1):20
- Maldonado J, Villaver E Eiroa C (2013) The metallicity signature of evolved stars with planets. Astronomy & Astrophysics 554:A84
- Mamajek EE (2009) Initial Conditions of Planet Formation: Lifetimes of Primordial Disks. In: EXOPLANETS AND DISKS: THEIR FORMATION AND DIVERSITY: Proceedings of the International Conference. AIP Conference Proceedings, Department of Physics and Astronomy, University of Rochester, Rochester, NY 14627-0171, AIP, pp 3–10
- Mann AW, Gaidos E, Kraus A Hilton EJ (2013) Testing the Metal of Late-type Kepler Planet Hosts with Iron-clad Methods. The Astrophysical Journal 770(1):43
- Mayor M Queloz D (1995) A Jupiter-mass companion to a solar-type star. Nature 378(6):355-359
- Mayor M, Bonfils X, Forveille T et al. (2009) The HARPS search for southern extra-solar planets. XVIII. An Earth-mass planet in the GJ 581 planetary system. Astronomy & Astrophysics 507(1):487–494
- Mayor M, Marmier M, Lovis C et al. (2011) The HARPS search for southern extra-solar planets XXXIV. Occurrence, mass distribution and orbital properties of super-Earths and Neptunemass planets. arXivorg p arXiv:1109.2497
- Ment K Charbonneau D (2023) The Occurrence Rate of Terrestrial Planets Orbiting Nearby Midto-late M Dwarfs from TESS Sectors 1-42. AJ165(6):265
- Meru F Bate MR (2010) Exploring the conditions required to form giant planets via gravitational instability in massive protoplanetary discs. Monthly Notices of the Royal Astronomical Society 406(4):2279–2288
- Montet BT, Crepp JR, Johnson JA, Howard AW Marcy GW (2014) The TRENDS High-contrast Imaging Survey. IV. The Occurrence Rate of Giant Planets around M Dwarfs. The Astrophysical Journal 781(1):28
- Mordasini C, Alibert Y Benz W (2009a) Extrasolar planet population synthesis. I. Method, formation tracks, and mass-distance distribution. Astronomy & Astrophysics 501(3):1139–1160
- Mordasini C, Alibert Y, Benz W Naef D (2009b) Extrasolar planet population synthesis. II. Statistical comparison with observations. Astronomy & Astrophysics 501(3):1161–1184
- Moriarty J Fischer D (2015) Building Massive Compact Planetesimal Disks from the Accretion of Pebbles. The Astrophysical Journal 809(1):94
- Mortier A, Santos NC, Sozzetti A et al. (2012) The frequency of giant planets around metal-poor stars. Astronomy & Astrophysics 543:A45
- Mortier A, Santos NC, Sousa S et al. (2013a) On the functional form of the metallicity-giant planet correlation. Astronomy & Astrophysics 551:A112
- Mortier A, Santos NC, Sousa SG et al. (2013b) New and updated stellar parameters for 71 evolved planet hosts. On the metallicity-giant planet connection. Astronomy & Astrophysics 557:A70
- Mulders GD, Pascucci I Apai D (2015a) A Stellar-mass-dependent Drop in Planet Occurrence Rates. The Astrophysical Journal 798(2):112
- Mulders GD, Pascucci I Apai D (2015b) An Increase in the Mass of Planetary Systems around Lower-mass Stars. The Astrophysical Journal 814(2):130
- Mulders GD, Pascucci I, Apai D, Frasca A Molenda-Zakowicz J (2016) A Super-solar Metallicity for Stars with Hot Rocky Exoplanets. The Astronomical Journal 152(6):187
- Mulders GD, Mordasini C, Pascucci I et al. (2019) The Exoplanet Population Observation Simulator. II. Population Synthesis in the Era of Kepler. The Astrophysical Journal 887(2):157
- Mulders GD, O'Brien DP, Ciesla FJ, Apai D Pascucci I (2020) Earths in Other Solar Systems' Nbody Simulations: The Role of Orbital Damping in Reproducing the Kepler Planetary Systems. The Astrophysical Journal 897(1):72

- Mulders GD, Drą zkowska J, van der Marel N, Ciesla FJ Pascucci I (2021) Why Do M Dwarfs Have More Transiting Planets? ApJ920(1):L1
- Neves V, Bonfils X, Santos NC et al. (2013) Metallicity of M dwarfs. III. Planet-metallicity and planet-stellar mass correlations of the HARPS GTO M dwarf sample. Astronomy & Astrophysics 551:A36
- Nielsen EL, De Rosa RJ, Macintosh B et al. (2019) The Gemini Planet Imager Exoplanet Survey: Giant Planet and Brown Dwarf Demographics from 10 to 100 au. The Astronomical Journal 158(1):13
- Ormel CW Klahr HH (2010) The effect of gas drag on the growth of protoplanets. Analytical expressions for the accretion of small bodies in laminar disks. Astronomy & Astrophysics 520:A43
- Osborn A Bayliss D (2020) Investigating the planet-metallicity correlation for hot Jupiters. MNRAS491(3):4481–4487
- Pascucci I, Testi L, Herczeg GJ et al. (2016) A Steeper than Linear Disk Mass-Stellar Mass Scaling Relation. The Astrophysical Journal 831(2):125
- Pasquini L, Döllinger MP, Weiß A et al. (2007) Evolved stars suggest an external origin of the enhanced metallicity in planet-hosting stars. Astronomy & Astrophysics 473(3):979–982
- Petigura EA, Sinukoff E, Lopez ED et al. (2017) Four Sub-Saturns with Dissimilar Densities: Windows into Planetary Cores and Envelopes. The Astronomical Journal 153(4):142
- Petigura EA, Marcy GW, Winn JN et al. (2018) The California-Kepler Survey. IV. Metal-rich Stars Host a Greater Diversity of Planets. The Astronomical Journal 155(2):89
- Pollack JB, Hubickyj O, Bodenheimer P et al. (1996) Formation of the Giant Planets by Concurrent Accretion of Solids and Gas. Icarus 124(1):62–85
- Raymond SN, Scalo J Meadows VS (2007) A Decreased Probability of Habitable Planet Formation around Low-Mass Stars. The Astrophysical Journal 669(1):606–614
- Raymond SN, Barnes R Mandell AM (2008) Observable consequences of planet formation models in systems with close-in terrestrial planets. Monthly Notices of the Royal Astronomical Society 384(2):663–674
- Reffert S, Bergmann C, Quirrenbach A, Trifonov T Künstler A (2015) Precise radial velocities of giant stars. VII. Occurrence rate of giant extrasolar planets as a function of mass and metallicity. Astronomy & Astrophysics 574:A116
- Reid IN (2002) On the Nature of Stars with Planets 114(7):306-329
- Rein H (2012) Period ratios in multiplanetary systems discovered by Kepler are consistent with planet migration. Monthly Notices of the Royal Astronomical Society: Letters 427(1):L21–L24
- Ribas I, Reiners A, Zechmeister M et al. (2023) The CARMENES search for exoplanets around M dwarfs. Guaranteed time observations Data Release 1 (2016-2020). A&A670:A139
- Rojas-Ayala B, Covey KR, Muirhead PS Lloyd JP (2012) Metallicity and Temperature Indicators in M Dwarf K-band Spectra: Testing New and Updated Calibrations with Observations of 133 Solar Neighborhood M Dwarfs. The Astrophysical Journal 748(2):93
- Sabotta S, Schlecker M, Chaturvedi P et al. (2021) The CARMENES search for exoplanets around M dwarfs. Planet occurrence rates from a subsample of 71 stars. A&A653:A114
- Sadakane K, Ohkubo M, Takeda Y et al. (2002) Abundance Analyses of 12 Parent Stars of Extrasolar Planets Observed with the SUBARU/HDS. Publications of the Astronomical Society of Japan 54(6):911–931
- Santos NC, Israelian G Mayor M (2000) Chemical analysis of 8 recently discovered extra-solar planet host stars. Astronomy & Astrophysics 363:228–238
- Santos NC, Israelian G Mayor M (2001) The metal-rich nature of stars with planets. Astronomy & Astrophysics 373(3):1019–1031
- Santos NC, Israelian G, Mayor M, Rebolo R Udry S (2003) Statistical properties of exoplanets. II. Metallicity, orbital parameters, and space velocities. Astronomy & Astrophysics 398(1):363– 376
- Santos NC, Israelian G Mayor M (2004) Spectroscopic [Fe/H] for 98 extra-solar planet-host stars. Exploring the probability of planet formation. Astronomy & Astrophysics 415(3):1153–1166

- Santos NC, Mayor M, Bonfils X et al. (2011) The HARPS search for southern extrasolar planets. XXV. Results from the metal-poor sample. Astronomy & Astrophysics 526:A112
- Schlaufman KC (2015) A Continuum of Planet Formation between 1 and 4 Earth Radii. The Astrophysical Journal Letters 799(2):L26
- Schlaufman KC Laughlin G (2010) A physically-motivated photometric calibration of M dwarf metallicity. Astronomy & Astrophysics 519:A105
- Schlaufman KC Laughlin G (2011) Kepler Exoplanet Candidate Host Stars Are Preferentially Metal Rich. The Astrophysical Journal 738(2):177
- Schlecker M, Burn R, Sabotta S et al. (2022) RV-detected planets around M dwarfs: Challenges for core accretion models. A&A664:A180
- Sousa SG, Santos NC, Mayor M et al. (2008) Spectroscopic parameters for 451 stars in the HARPS GTO planet search program. Stellar [Fe/H] and the frequency of exo-Neptunes. Astronomy & Astrophysics 487(1):373–381
- Sousa SG, Santos NC, Israelian G, Mayor M Udry S (2011) Spectroscopic stellar parameters for 582 FGK stars in the HARPS volume-limited sample. Revising the metallicity-planet correlation. Astronomy & Astrophysics 533:A141
- Swift JJ, Johnson JA, Morton TD et al. (2013) Characterizing the Cool KOIs. IV. Kepler-32 as a Prototype for the Formation of Compact Planetary Systems throughout the Galaxy. The Astrophysical Journal 764(1):105
- Takeda Y, Sato B Murata D (2008) Stellar Parameters and Elemental Abundances of Late-G Giants. Publications of the Astronomical Society of Japan 60(4):781–802
- Terquem C Papaloizou J (2007) Migration and the Formation of Systems of Hot Super-Earths and Neptunes. The Astrophysical Journal 654(2):1110–1120
- Terrien RC, Mahadevan S, Bender CF et al. (2012) An H-band Spectroscopic Metallicity Calibration for M Dwarfs. The Astrophysical Journal Letters 747(2):L38
- Thommes EW, Matsumura S Rasio FA (2008) Gas Disks to Gas Giants: Simulating the Birth of Planetary Systems. Science 321(5):814–
- Thorngren DP, Fortney JJ, Murray-Clay RA Lopez ED (2016) The Mass-Metallicity Relation for Giant Planets. The Astrophysical Journal 831(1):64
- Udry S Santos NC (2007) Statistical Properties of Exoplanets. Annual Review of Astronomy and Astrophysics 45(1):397–439
- Udry S, Mayor M, Benz W et al. (2006) The HARPS search for southern extra-solar planets. V. A 14 Earth-masses planet orbiting HD 4308. Astronomy & Astrophysics 447(1):361–367
- Vigan A, Fontanive C, Meyer M et al. (2021) The SPHERE infrared survey for exoplanets (SHINE). III. The demographics of young giant exoplanets below 300 au with SPHERE. A&A651:A72
- Voelkel O, Klahr H, Mordasini C, Emsenhuber A Lenz C (2020) Effect of pebble flux-regulated planetesimal formation on giant planet formation. A&A642:A75
- Wang J Fischer DA (2015) Revealing a Universal Planet-Metallicity Correlation for Planets of Different Sizes Around Solar-type Stars. The Astronomical Journal 149(1):14
- Ward WR (1997) Protoplanet Migration by Nebula Tides. Icarus 126(2):261-281
- Wetherill GW (1985) Occurrence of giant impacts during the growth of the terrestrial planets. Science (ISSN 0036-8075) 228(4701):877–879
- Wilson RF, Teske J, Majewski SR et al. (2018) Elemental Abundances of Kepler Objects of Interest in APOGEE. I. Two Distinct Orbital Period Regimes Inferred from Host Star Iron Abundances. AJ155(2):68
- Winn JN, Sanchis-Ojeda R, Rogers L et al. (2017) Absence of a Metallicity Effect for Ultra-shortperiod Planets. The Astronomical Journal 154(2):60
- Wittenmyer RA, Jones MI, Zhao J et al. (2017) The Pan-Pacific Planet Search. VI. Giant Planets Orbiting HD 86950 and HD 222076. The Astronomical Journal 153(2):51
- Wolfgang A, Rogers LA Ford EB (2016) Probabilistic Mass-Radius Relationship for Sub-Neptune-Sized Planets. The Astrophysical Journal 825(1):19

Exoplanet Populations and their Dependence on Host Star Properties

- Wolthoff V, Reffert S, Quirrenbach A et al. (2022) Precise radial velocities of giant stars. XVI. Planet occurrence rates from the combined analysis of the Lick, EXPRESS, and PPPS giant star surveys. A&A661:A63
- Yang JY, Xie JW Zhou JL (2020) Occurrence and Architecture of Kepler Planetary Systems as Functions of Stellar Mass and Effective Temperature. The Astronomical Journal 159(4):164
- Yee SW, Winn JN, Hartman JD et al. (2023) The TESS Grand Unified Hot Jupiter Survey. II. Twenty New Giant Planets. ApJS265(1):1
- Zhu W, Wang J Huang C (2016) Dependence of Small Planet Frequency on Stellar Metallicity Hidden by Their Prevalence. The Astrophysical Journal 832(2):196
- Zink JK, Hardegree-Ullman KK, Christiansen JL et al. (2023) Scaling K2. VI. Reduced Smallplanet Occurrence in High-galactic-amplitude Stars. AJ165(6):262