

ON THE PROBABILITY DISTRIBUTION ASSOCIATED TO COMMUTATOR WORD MAP IN FINITE GROUPS II

TUSHAR KANTA NAIK

ABSTRACT. Let $P(G)$ denotes the set of sizes of fibers of non-trivial commutators of the commutator word map. Here, we prove that $|P(G)| = 1$, for any finite group G of nilpotency class 3 with exactly two conjugacy class sizes. We also show that for given $n \geq 1$, there exists a finite group G of nilpotency class 2 with exactly two conjugacy class sizes such that $|P(G)| = n$.

1. INTRODUCTION

Let \mathbb{G} denote the family of all finite groups. Define $Pr : \mathbb{G} \rightarrow \mathbb{Q} \cap (0, 1]$ as follows:

$$Pr(G) = \frac{|\{(x, y) \in G \times G \mid xy = yx\}|}{|G|^2}, \quad \text{for } G \in \mathbb{G}.$$

$Pr(G)$ is called the commuting probability of G . Various probability distribution associated to commutator word map has been subject of active research in recent years (see [1], [6], [7], [8]). In 2008, Pournaki and Sobhani [7] introduced the notion of $Pr_g(G)$, which is defined as follows:

$$Pr_g(G) = \frac{|\{(x, y) \mid [x, y] = g\}|}{|G|^2}, \quad \text{for } g \in G.$$

For a group G and elements $x, y \in G$, the commutator $[x, y]$ of x and y is defined by $x^{-1}y^{-1}xy$. Note that $Pr_1(G) = Pr(G)$, where 1 denotes the identity element in the group. In [7], Pournaki and Sobhani computed $Pr_g(G)$ for finite groups G , which have only two different irreducible complex character degrees. They also obtained explicit formulas for $Pr_g(G)$, when G is a finite group with $|G'| = p$, where p is a prime integer. We denote by G' the commutator subgroup of G .

A finite group G is said to be of *conjugate type* $(1, m)$, for some integer $m > 1$, if for all non-central element g of G , $|g^G| = m$. Here g^G denotes the conjugacy class of g in G and is defined as $g^G := \{h^{-1}gh \mid h \in G\}$. N. Ito [4] proved that, if G is of conjugate type $(1, m)$, then $m = p^k$, for some prime p and integer $k \geq 1$. Nath and Yadav [6] studied $Pr_g(G)$, when G is either of conjugate type $(1, p^n)$ or a Camina p -group. A finite group G is said to be a *Camina group*, if $[g, G] = G'$, for all $g \in G \setminus G'$. Here $[g, G]$ denotes the set $\{[g, h] \mid h \in G\}$. Note that $g[g, G] = g^G$.

We denote by $K(G)$ the set $\{[x, y] \mid (x, y) \in G \times G\}$. For $g \in K(G)$, we define *fiber* of g as follows:

$$fiber(g) := \{(x, y) \in G \times G \mid [x, y] = g\}.$$

2010 *Mathematics Subject Classification.* 20D60, 20P05.

Key words and phrases. commuting probability, conjugate type, isoclinism, fiber.

Note that, formula for $Pr_g(G)$ can be re-written as follows:

$$Pr_g(G) = \begin{cases} \frac{|fiber(g)|}{|G|^2}, & \text{if } g \in K(G) \\ 0, & \text{otherwise.} \end{cases}$$

Following the notation used by Nath and Yadav [6], we recall the notion of $P(G)$, which is defined as follows:

$$P(G) = \{Pr_g(G) \mid 1 \neq g \in K(G)\}.$$

Note that

$$(1.1) \quad |P(G)| = |\{|fiber(g)| \mid 1 \neq g \in K(G)\}|.$$

In this article, we restrict our attention to finite p -groups of conjugate type $(1, p^n)$. In 2002, K.Ishikawa [3] proved that nilpotency class of such a group can be either 2 or 3. So we consider following two family; \mathcal{G}_2 , the family of finite p -groups of nilpotency class 2 and conjugate type $(1, p^n)$, $n \geq 1$ and \mathcal{G}_3 , the family of finite p -groups of nilpotency class 3 and conjugate type $(1, p^n)$, $n \geq 1$.

Consider the following group G_r , constructed by Ito [4, Page 23], for any positive integer $r \geq 1$ and prime $p > 2$.

$$(1.2) \quad G_r = \langle a_1, \dots, a_{r+1} \mid [a_i, a_j] = b_{ij}, [a_k, b_{ij}] = 1, \\ a_i^p = a_r^p = b_{ij}^p = 1, 1 \leq i < j \leq r+1, 1 \leq k \leq r+1 \rangle.$$

Note that $G_r \in \mathcal{G}_2$, for all $r \geq 1$. In particular, G_r is of conjugate type $(1, p^r)$ and $C_{G_r}(g) = \langle g, Z(G_r) \rangle$, for all non central elements $g \in G_r$. Here $Z(G)$ denotes the center of G , for any group G . Nath and Yadav [6] computed $Pr_g(G_r)$ and proved that $|P(G_r)| = 1$, for all $r \geq 1$. They also gave explicit formula for $Pr_g(G)$, when G is a camina p -group of nilpotency class 2 and proved that $|P(G)| = 1$, for such a group G . Then they asked the following question:

Question. Is it true that $|P(G)| = 1$ for all finite p -groups G of conjugate type $(1, p^n)$ and nilpotency class 2?

In an attmpt to answer this, we prove the following theorem.

Theorem 1.3. *Let $n \geq 1$ be a given positive integer. Then there always exist a group G (depending on n) in \mathcal{G}_2 such that $|P(G)| = n$.*

Recently Naik, Kitture and Yadav [5] proved that finite groups of conjugate type $(1, p^n)$ and nilpotency class 3 exist if and only if n is even and for each positive even integer $2m$, finite p -group of nilpotency class 3 and of conjugate type $(1, p^{2m})$ is unique up to isoclinism (for definition see Section 2). For such groups, we prove the following result.

Theorem 1.4. *Let $G \in \mathcal{G}_3$ be a finite p -group of conjugate type $(1, p^{2n})$. Then for $g \in G'$,*

$$Pr_g(G) = \begin{cases} \frac{p^{3n} + p^{2n} - 1}{p^{5n}}, & \text{if } g = 1 \\ \frac{p^{2n} - 1}{p^{5n}}, & \text{if } 1 \neq g \in G'. \end{cases}$$

Hence $|P(G)| = 1$.

We remark that this theorem rectifies a faulty statement [6, Theorem 5.13], where it was claimed that $|P(G)| > 1$, for a finite p -group G of conjugate type $(1, p^2)$ and nilpotency class 3.

2. KEY RESULTS

We start with definition of *isoclinism*, which was introduced by P. Hall [2] in 1940. Let X be a finite group and $\bar{X} = X/Z(X)$. Then the map $a_X : \bar{X} \times \bar{X} \mapsto X'$ such that $a_X(xZ(X), yZ(X)) = [x, y]$ for $(x, y) \in X \times X$ is well-defined. Two finite groups G and H are said to be *isoclinic* if there exists an isomorphism ϕ of the factor group $\bar{G} = G/Z(G)$ onto $\bar{H} = H/Z(H)$, and an isomorphism θ of the subgroup G' onto H' such that the following diagram is commutative.

$$\begin{array}{ccc} \bar{G} \times \bar{G} & \xrightarrow{a_G} & G' \\ \phi \times \phi \downarrow & & \downarrow \theta \\ \bar{H} \times \bar{H} & \xrightarrow{a_H} & H'. \end{array}$$

The resulting pair (ϕ, θ) is called an *isoclinism* of G onto H . Notice that isoclinism is an equivalence relation among finite groups.

The following result follows from [2].

Lemma 2.1. *Let G be a finite p -group. Then there exists a group H in the isoclinism family of G such that $Z(H) \leq H'$.*

Such a group H is called a *stem group* in its isoclinism class.

Proof of following interesting result can be found in [7, Lemma 3.5] and [6, Theorem 2.3]

Lemma 2.2. *Let G and H be two isoclinic groups with isoclinism (ϕ, θ) . Then $Pr_g(G) = Pr_{\theta(g)}(H)$.*

In the light of the preceding two results, to compute $Pr_g(G)$ or $P(G)$, for any finite group G , we only need to consider a stem group from the isoclinic family of G .

Now we move towards some technical Lemmas. For a finite group G and element $g \in K(G)$, we define

$$\begin{aligned} T_g &= \{x \in G : g \in [x, G]\} \\ &\text{and} \\ TZ_g &= \{xZ(G) \in G/Z(G) : g \in [x, G]\}. \end{aligned}$$

Note that;

$$|T_g| = |Z(G)||TZ_g|.$$

Following useful expression of $Pr_g(G)$ is due to Das and Nath [1].

Lemma 2.3. *Let G be a finite group and $g \in G$. Then*

$$Pr_g(G) = \frac{1}{|G|} \sum_{x \in T_g} \frac{1}{|x^G|}.$$

We remark that for any finite group G and element $1 \neq g \in G$,

$$Pr_1(G) - Pr_g(G) \geq \frac{1}{[G : Z(G)]}.$$

Now, if G is finite group with exactly two conjugacy class sizes, then we can further simplify the formula of $Pr_g(G)$.

Lemma 2.4. *Let G be a finite group of conjugate type $(1, p^n)$ and $g \in K(G)$. Then*

$$Pr_g(G) = \begin{cases} \frac{1}{[G : Z(G)]} \frac{|TZ_g|}{p^n}, & \text{if } g \neq 1 \\ \frac{1}{[G : Z(G)]} \left(1 + \frac{[G : Z(G)] - 1}{p^n}\right), & \text{if } g = 1 \end{cases}$$

Proof. Suppose $1 \neq g \in K(G)$. Then for each $x \in T_g$, $|x^G| = p^n$. By Lemma 2.3, we get

$$Pr_g(G) = \frac{1}{|G|} \sum_{x \in T_g} \frac{1}{p^n} = \frac{|T_g|}{|G|p^n} = \frac{|Z(G)||TZ_g|}{|G|p^n} = \frac{1}{[G : Z(G)]} \frac{|TZ_g|}{p^n}$$

Now let $g = 1$. Then note that $T_g = G$. Again by Lemma 2.3, we get that

$$\begin{aligned} Pr_1(G) &= \frac{1}{|G|} \sum_{x \in G} \frac{1}{|x^G|} \\ &= \frac{1}{|G|} \sum_{x \in Z(G)} \frac{1}{|x^G|} + \frac{1}{|G|} \sum_{x \in G \setminus Z(G)} \frac{1}{|x^G|} \\ &= \frac{|Z(G)|}{|G|} + \frac{1}{|G|} \frac{|G| - |Z(G)|}{p^n} \\ &= \frac{1}{[G : Z(G)]} \left(1 + \frac{[G : Z(G)] - 1}{p^n}\right). \end{aligned}$$

□

Nath and Yadav [6] gave explicit formula for $Pr_g(G_r)$, for $r \geq 1$.

Lemma 2.5. *Let G_r be as defined in 1.2. Then*

$$Pr_g(G_r) = \begin{cases} \frac{p^2 - 1}{p^{2r+1}}, & \text{if } g \neq 1 \\ \frac{p^{r+1} + p^r - 1}{p^{2r+1}}, & \text{if } g = 1 \end{cases}$$

Using preceding Lemma and the formula $Pr_g(G) = \frac{|fiber(g)|}{|G|^2}$, we get an expression for size of fibers in G_r .

Lemma 2.6. *Let G_r be as defined in 1.2. Then*

$$|\text{fiber}(g)| = \begin{cases} (p^2 - 1)p^{r^2+r+1}, & \text{if } g \neq 1 \\ (p^{r+1} + p^r - 1)p^{r^2+r+1}, & \text{if } g = 1 \end{cases}$$

From preceding Lemma, we get an expression for $|K(G)|$.

Lemma 2.7. *Let G_r be as defined in 1.2. Then $|K(G_r)| - 1 = \frac{(p^{r+1} - 1)(p^r - 1)}{p^2 - 1}$.*

Proof. For any finite group G , we know that $\sum_{g \in K(G)} |\text{fiber}(g)| = |G|^2$. Thus we get,

$\sum_{g \in K(G) \setminus 1} |\text{fiber}(g)| = |G|^2 - |\text{fiber}(1)|$. Therefore by Lemma 2.6, we have

$$\begin{aligned} (|K(G_r)| - 1)(p^2 - 1)p^{r^2+r+1} &= p^{r^2+3r+2} - (p^{r+1} + p^r - 1)p^{r^2+r+1} \\ &= (p^{r^2} + r + 1)(p^{2r+1} - p^{r+1} - p^r + 1) \\ &= (p^{r^2} + r + 1)(p^{r+1} - 1)(p^r - 1). \end{aligned}$$

Hence $|K(G_r)| - 1 = \frac{(p^{r+1} - 1)(p^r - 1)}{p^2 - 1}$. □

Lemma 2.8. *Suppose $G = G_{n-1}$ (as defined in 1.2) is generated by a_1, a_2, \dots, a_n , with $n \geq 4$. Then there do not exist $x, y \in G$ such that*

$$(2.9) \quad [x, y] = [a_1, a_2]^{i_1} [a_3, a_4]^{i_2} \dots [a_{2m-1}, a_{2m}]^{i_m},$$

where $2 \leq m \leq \lfloor n/2 \rfloor$ and $i_k \not\equiv 0 \pmod{p}$, for $k = 1, 2, \dots, m$.

Proof. We prove this lemma by the method of contradiction. Suppose that there exist x and $y \in G$ satisfying equation (2.9). Let $x = a_1^{j_1} a_2^{j_2} \dots a_n^{j_n}$ and $y = a_1^{k_1} a_2^{k_2} \dots a_n^{k_n}$ (reading modulo $Z(G)$). As $i_1 \not\equiv 0 \pmod{p}$, at least one of j_1 and k_1 has to be non-zero modulo p . Without loss of generality, we take j_1 to be non-zero. Then we can write y as

$$y = a_1^{k_1} a_2^{k_2} \dots a_n^{k_n} = (a_1^{j_1} a_2^{j_2} \dots a_n^{j_n})^{k_1 j_1^{-1}} (a_2^{l_2} a_3^{l_3} \dots a_n^{l_n} z_1) = x^{k_1 j_1^{-1}} (a_2^{l_2} a_3^{l_3} \dots a_n^{l_n} z_1),$$

where $z_1 \in Z(G)$ and l_2, l_3, l_4 are some suitable integers. Now

$$[x, y] = [x, y_1], \quad \text{where } y_1 = a_2^{l_2} a_3^{l_3} \dots a_n^{l_n}.$$

Computing $[x, y_1]$, we get that $[a_1, a_2^{l_2} a_3^{l_3} \dots a_n^{l_n}]$ is exactly the commutator involving a_1 in left-hand side of equation (2.9). Comparing it with right-hand side of equation (2.9), we see that l_2 has to be non-zero modulo p , in particular $l_2 = i_1 j_1^{-1}$ and l_3, l_4, \dots, l_n have to be zero modulo p . Thus we have

$$y_1 = b^{i_1 j_1^{-1}} \quad \text{with } j_1 \text{ non-zero modulo } p.$$

Now, again computing $[x, y]$, considering commutator involving a_2 and comparing with equation (2.9), we see that j_2, j_3, \dots, j_n have to be zero modulo p . So we have $x = a_1^{j_1}$ and $y_1 = b^{i_1 j_1^{-1}}$ with j_1 non-zero. Therefore $[x, y] = [x, y_1] = [a, b]^{i_1}$, a contradiction. □

Lemma 2.10. *Suppose $G = G_n$ (as defined in 1.2) is generated by a_1, a_2, \dots, a_n , with $n \geq 6$. Then there do not exist x, y, z and $w \in G$ such that*

$$(2.11) \quad [x, y][z, w] = [a_1, a_2]^{i_1} [a_3, a_4]^{i_2} \dots [a_{2m-1}, a_{2m}]^{i_m},$$

where $3 \leq m \leq \lfloor n/2 \rfloor$ and $i_k \not\equiv 0 \pmod{p}$, for $k = 1, 2, \dots, m$.

Proof. We also prove this lemma by the method of contradiction. Suppose that there exist x, y, z and $w \in G$ satisfying equation (2.11). Let $x = a_1^{j_1} a_2^{j_2} \dots a_n^{j_n}$, $y = a_1^{k_1} a_2^{k_2} \dots a_n^{k_n}$, $z = a_1^{l_1} a_2^{l_2} \dots a_n^{l_n}$ and $w = a_1^{t_1} a_2^{t_2} \dots a_n^{t_n}$ (reading modulo $Z(G)$). As $i_1 \not\equiv 0 \pmod{p}$, at least one of j_1, k_1, l_1 and t_1 has to be non-zero modulo p . Without loss of generality, we take j_1 to be non-zero. So $\{x, a_2, a_3, \dots, a_n\}$ also forms a generating set for G .

Let N be the subgroup of G generated by x, a_2 and their commutators with G . Easy to see that N is a normal subgroup. Set $\overline{G} = G/N$. Now $\{\overline{a_3}, \overline{a_4}, \dots, \overline{a_n}\}$ is a generating set for \overline{G} and easy to see that $\overline{G} \cong G_{n-2}$ (as defined in 1.2). In \overline{G} , equation 2.11 reduced to

$$[\overline{z}, \overline{w}] = [\overline{a_3}, \overline{a_4}]^{i_2} \dots [\overline{a_{2m-1}}, \overline{a_{2m}}]^{i_m},$$

where $3 \leq m \leq \lfloor n/2 \rfloor$ and $i_k \not\equiv 0 \pmod{p}$, for $k = 2, 3, \dots, m$. This is not possible by Lemma 2.8. Thus there do not exist x, y, z and $w \in G$ satisfying equation (2.11). This completes the proof. \square

3. PROOF OF THEOREM 1.3

Proof. We know that $|P(G_r)| = 1$ for all $r \geq 1$, from [6, Theorem B]. Therefore, we only need to prove the statement for $n \geq 2$. For given $n \geq 2$, consider $G = G_m$, with $m = n^2 + n - 3$. Note that G is of nilpotency class 2 and conjugate type $(1, p^m)$. Suppose that G is generated by $a_1, a_2, \dots, a_m, a_{m+1}$. Then consider the central subgroup H as follows:

$$\begin{aligned} H = \langle & [a_1, a_2][a_3, a_4], \\ & [a_5, a_6][a_7, a_8], [a_5, a_6][a_9, a_{10}], \\ & [a_{11}, a_{12}][a_{13}, a_{14}], [a_{11}, a_{12}][a_{15}, a_{16}], [a_{11}, a_{12}][a_{17}, a_{18}], \\ & \vdots \\ & [a_{\alpha+1}, a_{\alpha+2}][a_{\alpha+3}, a_{\alpha+4}], [a_{\alpha+1}, a_{\alpha+2}][a_{\alpha+5}, a_{\alpha+6}] \dots [a_{\alpha+1}, a_{\alpha+2}][a_{\alpha+2n-1}, a_{\alpha+2n}] \rangle; \end{aligned}$$

where $\alpha = (n-2)(n+1)$.

Note that $|H| = p^{n(n-1)/2}$ and $\alpha + 2n = n^2 + n - 2 = m + 1$.

Set $\overline{G} := G/H$. Note that $[\overline{a_1}, \overline{a_3}] \neq 1$ in \overline{G} . Thus \overline{G} is non-abelian, in particular nilpotency class of \overline{G} is 2. Our aim is to show that \overline{G} has exactly two conjugacy class sizes and $|P(\overline{G})| = n$. We complete the proof in two steps.

Claim 1: \overline{G} has exactly two conjugacy class sizes.

It is sufficient to prove that each non-central element commutes only with its power (reading modulo center). Let $\overline{x}, \overline{y} \in \overline{G}$ be such that neither is a power of the other. It is not difficult to see that $[x, y] \neq 1$ in G . Hence, if $[\overline{x}, \overline{y}] = 1$ in \overline{G} , then $[x, y] \in H^*$. For

a subgroup H of G , by H^* we denote the set of non-trivial elements of H . Suppose that

$$\begin{aligned}
 1 \neq [x, y] = & ([a_1, a_2][a_3, a_4])^{i_{1,1}} \\
 & ([a_5, a_6][a_7, a_8])^{i_{2,1}} ([a_5, a_6][a_9, a_{10}])^{i_{2,2}} \\
 & ([a_{11}, a_{12}][a_{13}, a_{14}])^{i_{3,1}} ([a_{11}, a_{12}][a_{15}, a_{16}])^{i_{3,2}} ([a_{11}, a_{12}][a_{17}, a_{18}])^{i_{3,3}} \\
 & \vdots \\
 & ([a_{\alpha+1}, a_{\alpha+2}][a_{\alpha+3}, a_{\alpha+4}])^{i_{n-1,1}} ([a_{\alpha+1}, a_{\alpha+2}][a_{\alpha+5}, a_{\alpha+6}])^{i_{n-1,2}} \dots \\
 & ([a_{\alpha+1}, a_{\alpha+2}][a_{\alpha+2n-1}, a_{\alpha+2n}])^{i_{n-1,n-1}}.
 \end{aligned}$$

So at least one $i_{j,k}$ is non-zero (mod p). After simplification, we get that

$$\begin{aligned}
 [x, y] = & [a_1, a_2]^{i_{1,1}} [a_3, a_4]^{i_{1,1}} \\
 & [a_5, a_6]^{i_{2,1}+i_{2,2}} [a_7, a_8]^{i_{2,1}} [a_9, a_{10}]^{i_{2,2}} \\
 & [a_{11}, a_{12}]^{i_{3,1}+i_{3,2}+i_{3,3}} [a_{13}, a_{14}]^{i_{3,1}} [a_{15}, a_{16}]^{i_{3,2}} [a_{17}, a_{18}]^{i_{3,3}} \\
 & \vdots \\
 & [a_{\alpha+1}, a_{\alpha+2}]^{i_{n-1,1}+i_{n-1,2}+\dots+i_{n-1,n-1}} [a_{\alpha+3}, a_{\alpha+4}]^{i_{n-1,1}} [a_{\alpha+5}, a_{\alpha+6}]^{i_{n-1,2}} \dots \\
 & [a_{\alpha+2n-1}, a_{\alpha+2n}]^{i_{n-1,n-1}}.
 \end{aligned}$$

Note that in right-hand side at least two commutator have non-zero power. Thus, by Lemma 2.8, $[x, y] \notin H^\#$ and consequently $[\bar{x}, \bar{y}] \neq 1$ in \overline{G} . Hence \overline{G} has exactly two conjugacy class sizes.

Claim 2: $|P(\overline{G})| = n$.

Here our plan is to show that

$$|\{|\text{fiber}(g)| \mid 1 \neq g \in K(\overline{G})\}| = n.$$

First, note the following elementary facts:

(i) $K(\overline{G}) = \overline{K(G)}$.

(ii) For any $h \in K(G)$, $|\overline{\text{fiber}(h)}| = \frac{|\text{fiber}(h)|}{|H|^2} = \frac{(p^2 - 1)p^{m^2+m+1}}{p^{n(n-1)}}$.

The last equality in (ii) is due to Lemma 2.6.

For each $h \in K(G)$, set $A_h := K(G) \cap hH$. We claim that for each $x \in A_h$,

$$(3.1) \quad \text{fiber}(\bar{x}) = \text{fiber}(\bar{h}) = \cup_{y \in A_h} \overline{\text{fiber}(y)}.$$

The first equality of (3.1) holds because of the fact: $x \in A_h \Rightarrow \bar{x} = \bar{h}$ in \overline{G} . Now, we proceed to show the second equality of (3.1).

Suppose $(\bar{a}, \bar{b}) \in \text{fiber}(\bar{h})$, i.e., $[\bar{a}, \bar{b}] = \bar{h}$. Then there exist some $y \in hH$ such that $[a, b] = y$, i.e., $(a, b) \in \text{fiber}(y)$. This implies $(\bar{a}, \bar{b}) = \overline{(a, b)} \in \overline{\text{fiber}(y)}$. By definition of A_h , $y \in A_h$. Hence $\text{fiber}(\bar{h}) \subseteq \cup_{y \in A_h} \overline{\text{fiber}(y)}$. Similarly reverse inclusion can be shown by backtracking the above steps. This completes the proof of second equality of (3.1) and hence (3.1) holds true.

Now, for $h \in K(G)$ and $x \in A_h$, we get

$$|\text{fiber}(\bar{x})| = |A_h| \frac{(p^2 - 1)p^{m(m-1)/2}}{p^{n(n-1)}}.$$

Hence, to show $|\{|\text{fiber}(\bar{x})| \mid 1 \neq g \in K(\overline{G})\}| = n$, it is sufficient to show that

$$|\{ |A_h| \mid 1 \neq h \in K(G) \}| = n.$$

Note that $h \in A_h$, for all $h \in K(G)$. Now fix some $h \in K(G) \setminus 1$. If $h \neq g \in A_h$, then

$$gh^{-1} = h', \text{ for some } 1 \neq h' \in H.$$

As both $g, h^{-1} \in K(G)$, consider $g = [a, b]$ and $h^{-1} = [c, d]$, for some $a, b, c, d \in G$. Thus, we get

$$[a, b][c, d] = h'.$$

By Lemma 2.10 and presentation of H , this is only possible when h' is of the form

$$h' = [a_{2i-1}, a_{2i}]^\alpha [a_{2j-1}, a_{2j}]^\alpha \text{ or } h' = [a_{2i-1}, a_{2i}]^\alpha [a_{2j-1}, a_{2j}]^{-\alpha},$$

for some suitable index i, j and integer $1 \leq \alpha \leq p-1$, with either $g = [a_{2i-1}, a_{2i}]^\alpha$ and $h^{-1} = [a_{2j-1}, a_{2j}]^{\pm\alpha}$ or vice versa.

Using this observation, presentation of H , Lemma 2.8 and Lemma 2.10, we proceed to compute A_h , for all $h \in K(G) \setminus 1$. First we partition $K(G) \setminus 1$ as $K(G) \setminus 1 = K(G)_1 \cup K(G)_2$; where

$$K(G)_1 = \{[a_{2i-1}, a_{2i}]^\beta \mid 1 \leq \beta \leq p-1, 1 \leq i \leq (m+1)/2\}$$

and $K(G)_2 = (K(G) \setminus 1) \setminus K(G)_1$.

Note that if $h \in K(G)_2$, then $A_h = \{h\}$. For $h \in K(G)_1$, we get the following expressions for A_h . For simplification of notation, now onwards, we denote $[a_i, a_j]$ by $h_{i,j}$ in this proof.

$$\begin{aligned} A_{h_{1,2}^{k_1}} &= \{h_{1,2}^{k_1}, h_{3,4}^{-k_1}\}, \\ A_{h_{5,6}^{k_2}} &= \{h_{5,6}^{k_2}, h_{7,8}^{-k_2}, h_{9,10}^{-k_2}\}, \\ A_{h_{11,12}^{k_3}} &= \{h_{11,12}^{k_3}, h_{13,14}^{-k_3}, h_{15,16}^{-k_3}, h_{17,18}^{-k_3}\}, \\ &\vdots \\ A_{h_{\alpha+1, \alpha+2}^{k_{n-1}}} &= \{h_{\alpha+1, \alpha+2}^{k_{n-1}}, h_{\alpha+5, \alpha+6}^{-k_{n-1}}, h_{\alpha+3, \alpha+4}^{-k_{n-1}} \cdots h_{\alpha+2n-1, \alpha+2n}^{-k_{n-1}}\}, \end{aligned}$$

for all $1 \leq k_i \leq p-1, i = 1, 2, \dots, n-1$.

So, we get that $\{|A_h| \mid 1 \neq h \in K(G)\} = \{1, 2, \dots, n\}$. Hence

$$|P(\overline{G})| = |\{|\text{fiber}(g)| \mid 1 \neq g \in K(\overline{G})\}| = |\{|A_h| \mid 1 \neq h \in K(G)\}| = n.$$

□

4. PROOF OF THEOREM 1.4

We start with a elementary result which follows from [5, Lemmas 3.4, 3.14 and 3.19] and definition of conjugate type.

Lemma 4.1. *Let G be a stem p -group of class 3 with conjugate type $(1, p^{2n})$. Then the following holds true.*

- (i) $|Z(G)| = p^{2n}$, $|G' : Z(G)| = p^n$ and $|G : G'| = p^{2n}$.
- (ii) For each $g \in G \setminus Z(G)$, $|g^G| = [G : C_G(x)] = p^{2n}$ and $|C_G(x) : Z(G)| = p^n$.

Following technical result on finite p -groups of conjugate type $(1, p^{2n})$ and nilpotency class 3 follows from [5, Lemmas 5.8, 5.12, 5.13 and 5.14].

Lemma 4.2. *Let G be a stem p -group of class 3 with conjugate type $(1, p^{2n})$. Then there exists a generating set of G , say $\{a_1, a_2, \dots, a_n, b_1, \dots, b_n\}$ such that the following hold true.*

- (i) $G' = \langle h_1, \dots, h_n, Z(G) \rangle$, where $h_i = [a_i, b_i]$, for $i = 1 \dots n$;
- (ii) $Z(G) = \langle z_1, z_2, \dots, z_{2n} \rangle$, where $[h_1, a_i] = z_i$ and $[h_1, b_i] = z_{n+i}$, for $i = 1 \dots n$.
- (iii) Say $A = \{\prod_{i=1}^n a_i^{k_i} : 0 \leq k_i \leq p-1\}$ and $B = \{\prod_{i=1}^n b_i^{l_i} : 0 \leq l_i \leq p-1\}$.

- (1) $x, y \in A$ or $x, y \in B \Rightarrow [x, y] \in Z(G)$.
- (2) $1 \neq x \in A$ and $1 \neq y \in B \Rightarrow [x, y] \notin Z(G)$.
- (3) For any $1 \neq a \in A$, $\{[a, b_i] : 1 \leq i \leq n\}$ generates G' over $Z(G)$.
- (4) For any $1 \neq b \in B$, $\{[a_i, b] : 1 \leq i \leq n\}$ generates G' over $Z(G)$.

(iv) Say $Z_1 = \langle z_1, \dots, z_n \rangle$ and $Z_2 = \langle z_{n+1}, \dots, z_{2n} \rangle$. Then for any $1 \neq a \in A$, $1 \neq b \in B$ and $h \in G' \setminus Z(G)$,

- (1) $[h, A] = Z_1 = [a, G']$.
- (2) $[h, B] = Z_2 = [b, G']$.

(v) For each $h \in G' - Z(G)$, $[h, G] = Z(G)$.

Set $\overline{G} = G/Z(G)$ and consider Φ to be the canonical homomorphism from G to \overline{G} . For $g \in G \setminus G'$, define

$$(4.3) \quad H_g = \Phi^{-1}(C_{\overline{G}}(\overline{g})).$$

Note that

$$H_g = C_g(G)G'.$$

As $C_g(G) \cap G' = Z(G)$ and $[G_g(G) : Z(G)] = p^n = [G' : Z(G)]$; therefore $[H_g : Z(G)] = p^{2n}$.

Lemma 4.4. *Let $g \in G \setminus G'$. Then for any $x \in H_g \setminus G'$, $[G', x] = [G', g]$.*

Proof. Take an arbitrary element of $[G', x]$, say $[[g, y], x]$, for some $y \in G$. As $x \in H_g$, so $[x, g] \in Z(G)$. Then using Hall-Witt identity, we get $[[g, y], x] = [[y, x], g]$ and so $[G', x] \subseteq [G', g]$. Similarly we can show the reverse inclusion and hence $[G', x] = [G', g]$. \square

Lemma 4.5. *Given any $h \in G' \setminus Z(G)$ and any $ab \neq 1$, for some $a \in A$ and $b \in B$, there always exists some $h^* \in G'$ such that $h \in [abh^*, G]$.*

Proof. Without loss of generality we can assume that $a \neq 1$. By Lemma 4.2(iii)(3), there exists some $b^* \in B$ such that

$$[a, b^*] = h \pmod{Z(G)}.$$

Now $[ab, b^*] = hw_1w_2$, for some $w_1 \in Z_1$ and $w_2 \in Z_2$. By Lemma 4.2(iv)(1), there exists some $h_1 \in G'$ such that $[a, h_1] = w_1^{-1}$. So

$$[ab, b^*h_1] = [ab, b^*][a, h_1][b, h_1] = hw_1w_2w_1^{-1}[b, h_1] = hw_2w_3$$

where $w_3 = [b, h_1] \in Z_2$ (by Lemma 4.2(iv)(2)). Again using Lemma 4.2(iv)(2), we get some $h^* \in G'$ such that $[h^*, b^*] = (w_2w_3)^{-1}$. Therefore

$$[abh^*, b^*h_1] = [ab, b^*h_1][h^*, b^*h_1] = hw_2w_3(w_2w_3)^{-1} = h.$$

□

Proof of Theorem 1.4:

Proof. Let $G \in \mathcal{G}_3$ be a finite p -group of conjugate type $(1, p^{2n})$. Then $|G| = p^{5n}$ and $[G : Z(G)] = p^{3n} = |G'|$. By Lemma 2.4, we get

$$Pr_1(G) = \frac{1}{p^{3n}} + \frac{1}{p^{2n}} \left\{ 1 - \frac{1}{p^{3n}} \right\} = \frac{p^{3n} + p^{2n} - 1}{p^{5n}}.$$

Note that

$$\sum_{g \in K(G)} Pr_g(G) = 1.$$

Thus

$$\sum_{g \in K(G) \setminus 1} Pr_g(G) = 1 - Pr_1(G) = \frac{p^{5n} - p^{3n} - p^{2n} + 1}{p^{5n}} = (p^{3n} - 1) \frac{p^{2n} - 1}{p^{5n}}.$$

Claim: $Pr_g(G) \geq \frac{p^{2n} - 1}{p^{5n}}$, for each $g \in G' \setminus 1$.

By Lemma 2.4, it is sufficient to show that $|TZ_g| \geq p^{2n} - 1$, for all $g \in G' \setminus 1$.

Case 1: $g \in Z(G) \setminus 1$.

Consider any $\alpha \in G' - Z(G)$. By Lemma 4.2(v), there exist some $\beta \in G \setminus G'$ such that $[\alpha, \beta] = g$. Now consider H_β (as defined in 4.3). Take $x \in H_\beta \setminus Z(G)$, then either $x \in H_\beta \setminus G'$ or $x \in G' \setminus Z(G)$. If $x \in H_\beta \setminus G'$, then by Lemma 4.4, $g = [\alpha, \beta] \in [G', \beta] = [G', x]$. If $x \in G' \setminus Z(G)$, then by Lemma 4.1(vii), $g \in [G', x]$. So

$$|TZ_g| \geq \frac{(H_\beta \setminus Z(G))Z(G)}{Z(G)} = p^{2n} - 1.$$

Case 2: $g \in G' \setminus Z(G)$.

By Lemma 4.5, we get

$$|TZ_g| \geq |\{ab : a \in A, b \in B \text{ and } ab \neq 1\}| = p^{2n} - 1.$$

Hence our claim holds true. Now, by counting argument, we get $Pr_g(G) = \frac{p^{2n} - 1}{p^{5n}}$, for each $h \in G' \setminus 1$. This completes the proof. □

Acknowledgement. I sincerely thank my supervisor Dr. Manoj Kumar Yadav for fruitful discussions and carefully reading of the preprint.

REFERENCES

- [1] A. K. Das and R. K. Nath, *On generalized relative commutativity degree of finite groups*, Int. Electronic. J. Algebra, **7**, 140-151, (2010).
- [2] P. Hall, *The classification of prime-power groups*, J. Reine Angew. Math., **182**, 130-141 (1940).
- [3] K. Ishikawa, *On finite p -groups which have only two conjugacy lengths*, Israel J. Math., **129**(1), 119-123 (2002).
- [4] N. Ito, *On finite groups with given conjugate types I*, Nayoga Math. J., **6**, 17-28 (1953).
- [5] T. K. Naik, R. D. Kitture, M. K. Yadav, *Finite p -Groups of Nilpotency Class 3 with Two Conjugacy Class Sizes*, arxiv link: <https://arxiv.org/abs/1708.03245>.
- [6] Rajat K. Nath and M.K. Yadav, *On the probability distribution associated to commutator word map in finite groups*, Internat. J. Algebra and Comput., **25**, 1107-1124 (2015)
- [7] M.R.Pouranki and R.Sobhani, *Probability that the commutator of two group element is equal to a given element*, J. Pure and Applied Algebra, **212**, 727-734 (2008).
- [8] A. Shalev, *Word maps, conjugacy classes and a noncommutative Waring-type theorem* Ann. of Math. **170**(3), 1383-1416 (2009)

HARISH-CHANDRA RESEARCH INSTITUTE, HBNI, CHHATNAG ROAD, JHUNSI, ALLAHABAD-211 019, INDIA
E-mail address: mathematics67@gmail.com and tusharkanta@hri.res.in