

Singularity of the n -th eigenvalue of high dimensional Sturm-Liouville problems

Xijun Hu[†], Lei Liu[†], Li Wu[†] and Hao Zhu^{‡*}

[†]Department of Mathematics, Shandong University
Jinan, Shandong 250100, P. R. China

[‡]Chern Institute of Mathematics, Nankai University
Tianjin, 300071, P. R. China

Abstract

It is natural to consider continuous dependence of the n -th eigenvalue on d -dimensional ($d \geq 2$) Sturm-Liouville problems after the results on 1-dimensional case by Kong, Wu and Zettl [14]. In this paper, we find all the boundary conditions such that the n -th eigenvalue is not continuous, and give complete characterization of asymptotic behavior of the n -th eigenvalue. This renders a precise description of the jump phenomena of the n -th eigenvalue near such a boundary condition. Furthermore, we divide the space of boundary conditions into $2d + 1$ layers and show that the n -th eigenvalue is continuously dependent on Sturm-Liouville equations and on boundary conditions when restricted into each layer. In addition, we prove that the analytic and geometric multiplicities of an eigenvalue are equal. Finally, we obtain derivative formula and positive direction of eigenvalues with respect to boundary conditions.

Key words. High dimensional Sturm-Liouville problems, multiplicity, the n -th eigenvalue, continuity, singularity.

AMS subject classifications. 34B09, 34B24, 34L15

*Corresponding author.

Contents

1	Introduction	2
2	Space of Sturm-Liouville problems	7
3	Basic properties of eigenvalues	11
4	Analysis on 1-dimensional results	13
5	Equality of multiplicities of an eigenvalue	17
6	Continuity of the n -th eigenvalue	19
7	Singularity of the n -th eigenvalue	22
8	Derivative formulas and comparison of eigenvalues	33
	Acknowledgements	35
	References	36

1 Introduction

In this paper, we consider the d -dimensional Sturm-Liouville equation

$$-(Py')' + Qy = \lambda Wy, \quad \text{on } [a, b], \quad (1.1)$$

where $d \geq 2$, λ is the spectral parameter, and P, Q and W satisfy the following assumptions:

Hypothesis 1:

- (i) P, Q and W are $d \times d$ -matrix symmetric real-valued functions on $[a, b]$,
- (ii) $P(t) \geq \mu_1$ and $W(t) \geq \mu_2$ a.e. on $t \in [a, b]$ for some $\mu_1, \mu_2 > 0$,
- (iii) $P, Q, W \in L^\infty([a, b], \mathbb{R}^{d \times d})$.

The self-adjoint boundary condition is given by

$$(A \mid B)Y(a, b) = 0, \quad (1.2)$$

where $Y(a, b) = (-y(a)^T, y(b)^T, (Py')(a)^T, (Py')(b)^T)^T$, A and B are $2d \times 2d$ complex matrices such that

$$\text{rank}(A \mid B) = 2d, \quad AB^* = BA^*, \quad (1.3)$$

A^* is the complex conjugate transpose of A . The spectrum of the Sturm-Liouville problem is bounded from below and consists of discrete eigenvalues, which are ordered in the following non-decreasing sequence

$$\lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \cdots \leq \lambda_n \leq \cdots ,$$

with $\lambda_n \rightarrow \infty$ as $n \rightarrow \infty$, counting repeatedly according to their analytic multiplicities.

When studying the perturbation of eigenvalues of Sturm-Liouville problem (1.1)–(1.2), the n -th eigenvalue changes as coefficients in the equation (1.1) or the boundary condition (1.2) are subjected to perturbations. The indices (i.e. n) of eigenvalues may change drastically in a continuous eigenvalue branch due to the high dimension, and the eigenvalues with the same index may jump from one to another branch in a complex way. The jump phenomena of the n -th eigenvalue in high dimension are more interesting from geometric aspects and more complicated than 1-dimensional Sturm-Liouville problems. Moreover, in computing eigenvalues, their indices are in general unknown and still need to be determined due to the importance of the first few eigenvalues in physical models. The high dimension, however, leads oscillation theory of 1-dimensional Sturm-Liouville problems to becoming invalid and thus makes a difficult task in numerical calculation. So the question “*what the singular (discontinuity) set of the n -th eigenvalue is in high dimensional case*” not only has strong motivation from physics and numerical analysis, but also is theoretical challenging. Indeed, after the previous work in 1-dimensional case [14], it has been an open problem for several years. We shall solve it in this paper and give complete characterization of asymptotic behavior of the n -th eigenvalue.

This question is completely answered for 1-dimensional Sturm-Liouville problems. The first breakthrough is due to Rellich [23]. In 1950, he gave an example for the following 1-dimensional Sturm-Liouville problem at the ICM [24]:

$$-u'' = \lambda u, \quad \text{on } [0, 1],$$

with the boundary condition $u(0) = 0, \kappa u'(1) = u(1)$, where $\kappa \in \mathbb{R}$. Though the n -th eigenvalue is continuous near $\kappa = 0$ from the left direction for each $n \geq 1$, it is discontinuous from the right direction, and has the following asymptotic behavior:

$$\lim_{\kappa \rightarrow 0^+} \lambda_1(\kappa) = -\infty, \quad \lim_{\kappa \rightarrow 0^+} \lambda_n(\kappa) = \lambda_{n-1}(0), \quad n \geq 2.$$

See also Figure 1 in P. 292 of [13]. In 1997, Everitt et al. in [7] investigated 1-dimensional Sturm-Liouville equation (1.1) with separated boundary condition $\cos \alpha y(a) - \sin \alpha (py')(a) = 0$, $\cos \beta y(b) - \sin \beta (py')(b) = 0$. By using Prüfer transformation to (1.1), they obtained that the n -th eigenvalue λ_n is continuous on $\alpha \times \beta \in [0, \pi) \times (0, \pi]$ for each $n \geq 1$, and moreover, for any fixed $\alpha \in [0, \pi)$,

$$\lim_{\beta \rightarrow 0^+} \lambda_1(\alpha, \beta) = -\infty, \quad \lim_{\beta \rightarrow 0^+} \lambda_n(\alpha, \beta) = \lambda_{n-1}(\alpha, \pi), \quad n \geq 2,$$

and for any fixed $\beta \in (0, \pi]$,

$$\lim_{\alpha \rightarrow \pi^-} \lambda_1(\alpha, \beta) = -\infty, \quad \lim_{\alpha \rightarrow \pi^-} \lambda_n(\alpha, \beta) = \lambda_{n-1}(0, \beta), \quad n \geq 2.$$

Kong et al. in [14] regarded the n -th eigenvalue as a function on the space of Sturm-Liouville equations and that of boundary conditions, respectively. They showed that the n -th eigenvalue is continuously dependent on the coefficients in (1.1) for each $n \geq 1$. By using the above results in [7] and some inequalities among eigenvalues of 1-dimensional Sturm-Liouville problems obtained in [6], they found the singular set of the n -th eigenvalue in the space of complex (resp. real) boundary conditions. They also gave all the asymptotic behavior of the n -th eigenvalue near each singular boundary condition. See Theorems 3.39 and 3.76 in [14]. Other discussions of the n -th eigenvalue can be found in [3, 5, 18, 28, 30].

It is worthy to mention that the analytic, algebraic and geometric multiplicities of an eigenvalue are shown to be equal for 1-dimensional Sturm-Liouville problems [6, 15, 16, 27] and some extensions [8, 9, 11, 12, 22, 25]. In particular, Kong et al. showed the equivalence of analytic and geometric multiplicities even if P changes sign in 1-dimensional case [16]. Naimark proved analytic and algebraic multiplicities of an eigenvalue coincide for a class of high-order differential operators [22]. When studying the discontinuity of the n -th eigenvalue, it is always listed according to the analytic multiplicity. From the perspective of application, however, people pay more attention to how many eigenvalue branches jump (tend to infinity) in the sense of geometric multiplicity. So it is necessary to clarify the relationships of these multiplicities of an eigenvalue for high dimensional Sturm-Liouville problems. Motivated by Naimark [22], we shall rigorously prove that the three multiplicities of an eigenvalue are the same in high dimensional case even if P is non-positive.

In this paper, we shall give the set of all the complex (or real) boundary conditions such that the n -th eigenvalue is not continuous in high dimensional case. We call it to be the singular set $\Sigma^{\mathbb{C}}$ (or $\Sigma^{\mathbb{R}}$) in the space of complex (or real) boundary conditions and call any element in $\Sigma^{\mathbb{C}}$ (or $\Sigma^{\mathbb{R}}$) to be a singular boundary condition. We mainly discuss the complex boundary conditions in this paper, and the real ones can be treated in a similar way. When there is truly difference in the discussion, we shall give remarks on providing a feasible way for the real ones. Our inspiration is from the symplectic geometry, especially the structure of Lagrangian-Grassmann manifold [1, 2]. Indeed, we shall prove that in high dimensional case, the singular set, denoted by $\Sigma^{\mathbb{C}}$, consists of all the boundary conditions $\mathbf{A} = [A | B]$ such that

$$n^0(B) > 0, \tag{1.4}$$

where $n^0(B)$ denotes the geometric multiplicity of zero eigenvalue of B .

An accompanying difficulty is how to give and prove asymptotic behavior of the n -th eigenvalue near a singular boundary condition. The strategy based on the Prüfer transformation does not work for separated boundary conditions due to the coupling of the Sturm-Liouville equations. The inequalities argument used in [14] also becomes invalid owing to the complexity of the boundary conditions, for example, the appearance of mixing boundary conditions [26]. Moreover, the directions, from which the boundary conditions tend to a more singular one, are diversified in high dimensional case. All these make the problem nontrivial. Our first task is also to study the topology of the space of boundary conditions. However, instead of using separated, coupled and mixed boundary conditions, we choose the coordinate charts of the Lagrangian-Grassmann manifold introduced by Arnold [1, 2] to describe the space of boundary conditions. We divide $\Sigma^{\mathbb{C}}$ into $2d$ layers such that for any $[A | B]$ in the k -th layer $\Sigma_k^{\mathbb{C}}$,

$$n^0(B) = k, \tag{1.5}$$

where $1 \leq k \leq 2d$. Define $\Sigma_0^{\mathbb{C}}$ to be the complementary set of $\Sigma^{\mathbb{C}}$ in the space of boundary conditions. We then prove that the n -th eigenvalue is continuously dependent on the Sturm-Liouville equations and on the boundary conditions when restricted into $\Sigma_k^{\mathbb{C}}$ for each $n \geq 1$, where $0 \leq k \leq 2d$. In the proof of asymptotic behavior, besides using the above results and the locally uniform property of eigenvalues (Lemma 3.2), our technique is to construct various paths in different parts of the k -th layer in the space of boundary

conditions. The asymptotic behavior of the n -th eigenvalue is proved for the following targets via a step-by-step procedure:

1. A non-coupled Sturm-Liouville equation and a separated boundary condition in a path connected component of $\Sigma_k^{\mathbb{C}}$, see Lemmas 7.3 and 7.4.
2. The non-coupled Sturm-Liouville equation and any boundary condition in the path connected component of $\Sigma_k^{\mathbb{C}}$, see Lemma 7.5.
3. Any fixed Sturm-Liouville equation and any boundary condition in the path connected component of $\Sigma_k^{\mathbb{C}}$, see Lemma 7.6.

Our complete characterization of asymptotic behavior of the n -th eigenvalue near $\mathbf{A} \in \Sigma_k^{\mathbb{C}}$ (or $(\boldsymbol{\omega}, \mathbf{A})$) is given in Theorems 7.1 and 7.2, where $1 \leq k \leq 2d$. The essential characterization is that there exists a neighborhood U of \mathbf{A} in the whole space of boundary conditions such that $U = \cup_{0 \leq i \leq k} U^i$, and for each $0 \leq i \leq k$, $U^i \neq \emptyset$ and

$$\lim_{U^i \ni \mathbf{B} \rightarrow \mathbf{A}} \lambda_n(\mathbf{B}) = -\infty, \quad 1 \leq n \leq i, \quad (1.6)$$

$$\lim_{U^i \ni \mathbf{B} \rightarrow \mathbf{A}} \lambda_n(\mathbf{B}) = \lambda_{n-i}(\mathbf{A}), \quad n > i. \quad (1.7)$$

In order to make the results explicitly, we clarify what U^i is in Corollary 7.1.

In a forthcoming paper, we shall show that $[-\Psi_\lambda \mid \Phi_\lambda]$ defined in (3.3)–(3.4) tends to $[I_{2d} \mid 0_{2d}]$ as $\lambda \rightarrow -\infty$ for a more general Sturm-Liouville system. Based on this result and the theory of Maslov index, we shall give a new proof of the discontinuity of the n -th eigenvalue. Furthermore, we determine the range of the n -th eigenvalue not only on the whole space of boundary conditions but also on the k -th layer $\Sigma_k^{\mathbb{C}}$ (or $\Sigma_k^{\mathbb{R}}$), where $0 \leq k \leq 2d$.

The rest of this paper is organized as follows. In Section 2, topology on space of Sturm-Liouville equations, and that on space of complex boundary conditions are presented. Basic properties of eigenvalues are given in Section 3 and further analysis on 1-dimensional results is provided in Section 4. In Section 5, it is proved that the analytic, algebraic and geometric multiplicities of an eigenvalue are equal. Section 6 is devoted to proving that the n -th eigenvalue is continuous on the space of Sturm-Liouville equations, and on each layer in the space of boundary conditions. Singularity of the n -th eigenvalue is completely characterized in Section 7. In the last section, derivative formula of eigenvalues with respect to the boundary conditions and comparison of eigenvalues are obtained.

2 Space of Sturm-Liouville problems

In this section, we introduce the topology on space of Sturm-Liouville equations, and that on space of complex boundary conditions, respectively.

The space of Sturm-Liouville equations is

$$\Omega := \{(P, Q, W) : P, Q \text{ and } W \text{ satisfy Hypothesis 1}\}$$

with product topology induced by $L^\infty([a, b], \mathbb{R}^{d \times d})$. Following [14], we use bold faced (lower case) Greek letters, such as $\boldsymbol{\omega}$, to stand for elements in Ω .

Two linear algebraic systems

$$(A \mid B)Y(a, b) = 0, \quad (C \mid D)Y(a, b) = 0,$$

represent the same complex boundary condition if and only if there exists a matrix $T \in GL(2d, \mathbb{C})$ such that

$$(C \mid D) = (TA \mid TB),$$

where $GL(2d, \mathbb{C}) := \{2d \times 2d \text{ complex matrix } T : \det T \neq 0\}$. Since each boundary condition (1.2) considered in this paper is self-adjoint, it must satisfy (1.3). So it is natural to take the quotient space

$$\mathcal{B}^{\mathbb{C}} := GL(2d, \mathbb{C}) \backslash \mathcal{L}_{2d, 4d}(\mathbb{C}),$$

equipped with the quotient topology, as the space of complex boundary conditions, where $\mathcal{L}_{2d, 4d}(\mathbb{C}) := \{2d \times 4d \text{ complex matrix } (A \mid B) : \text{rank}(A \mid B) = 2d, AB^* = BA^*\}$. The topology on $\mathcal{L}_{2d, 4d}(\mathbb{C})$ here is induced by that of \mathbb{C}^{8d^2} . The boundary condition represented by (1.2) is denoted by $[A \mid B] := \{(TA \mid TB) : T \in GL(2d, \mathbb{C})\}$. Bold faced capital Latin letters, such as \mathbf{A} , are also used for boundary conditions. See also [15] in 1-dimensional case. $\mathcal{B}^{\mathbb{C}}$ coincides with the complex Lagrangian-Grassmann manifold $\Lambda(2d)$ [2].

Motivated by Arnold [1, 2], we shall give the canonical atlas of local coordinate systems on $\mathcal{B}^{\mathbb{C}}$ in our framework. Let $e_i = (0, \dots, 0, \underset{i}{1}, 0, \dots, 0)_{1 \times 2d}^T$

for any $1 \leq i \leq 2d$, S be a $2d \times 2d$ matrix whose entries and columns are denoted by s_{lj} , $1 \leq l, j \leq 2d$, and $s_j = (s_{1j}, \dots, s_{2dj})^T$, respectively. Set K be a subset of $\{1, 2, \dots, 2d\}$ with $\sharp(K)$ to be the number of elements in K . Define

$$\mathcal{O}_K^{\mathbb{C}} = \{ \mathbf{A} = [A \mid B] = [(a_1, a_2, \dots, a_{2d}) \mid (b_1, b_2, \dots, b_{2d})] : \quad (2.1)$$

$$a_i = \begin{cases} -e_i & \text{if } i \in K, \\ s_i & \text{if } i \in \{1, 2, \dots, 2d\} \setminus K, \end{cases}$$

$$b_i = \begin{cases} s_i & \text{if } i \in K, \\ e_i & \text{if } i \in \{1, 2, \dots, 2d\} \setminus K, \end{cases}$$

for any $2d \times 2d$ Hermitian matrix $S = (s_1, s_2, \dots, s_{2d})$ }

for any $K \subset \{1, 2, \dots, 2d\}$. Below S is written as $S(\mathbf{A})$ when it is necessary to indicate its dependence on \mathbf{A} . The following result gives the topology and geometric structure on $\mathcal{B}^{\mathbb{C}}$.

Theorem 2.1.

$$\mathcal{B}^{\mathbb{C}} = \bigcup_{K \subset \{1, 2, \dots, 2d\}} \mathcal{O}_K^{\mathbb{C}}.$$

Moreover, $\mathcal{B}^{\mathbb{C}}$ is a connected and compact real-analytic manifold of dimension $4d^2$.

Remark 2.1. Similar result is true for $\mathcal{B}^{\mathbb{R}}$ with \mathbb{C} replaced by \mathbb{R} except that the dimension of $\mathcal{B}^{\mathbb{R}}$ is $d(2d + 1)$ as a real-analytic manifold.

Proof. Firstly, we show that

$$\bigcup_{K \subset \{1, 2, \dots, 2d\}} \mathcal{O}_K^{\mathbb{C}} \subset \mathcal{B}^{\mathbb{C}}. \quad (2.2)$$

For any $K \subset \{1, \dots, 2d\}$, choose $[A \mid B] = [(a_1, \dots, a_{2d}) \mid (b_1, \dots, b_{2d})] \in \mathcal{O}_K^{\mathbb{C}}$, where

$$a_j = (a_{1j}, \dots, a_{2dj})^T, b_j = (b_{1j}, \dots, b_{2dj})^T, j = 1, \dots, 2d.$$

Define

$$\delta_{i,K} := \begin{cases} 1, & i \in K, \\ 0, & i \notin K. \end{cases}$$

For any $1 \leq i, k \leq 2d$, by (2.1) we have

$$\begin{aligned}
(AB^*)_{ik} &= \sum_{j=1}^{2d} a_{ij} \bar{b}_{kj} \\
&= \sum_{j \in K} a_{ij} \bar{b}_{kj} + \sum_{j \notin K} a_{ij} \bar{b}_{kj} \\
&= \sum_{j \in K} -\delta_{ij} \bar{s}_{kj} + \sum_{j \notin K} s_{ij} \delta_{kj} \\
&= -\delta_{i,K} \bar{s}_{ki} + (1 - \delta_{k,K}) s_{ik},
\end{aligned} \tag{2.3}$$

$$\begin{aligned}
(BA^*)_{ik} &= \sum_{j=1}^{2d} b_{ij} \bar{a}_{kj} \\
&= \sum_{j \in K} b_{ij} \bar{a}_{kj} + \sum_{j \notin K} b_{ij} \bar{a}_{kj} \\
&= \sum_{j \in K} -s_{ij} \delta_{kj} + \sum_{j \notin K} \delta_{ij} \bar{s}_{kj} \\
&= -\delta_{k,K} s_{ik} + (1 - \delta_{i,K}) \bar{s}_{ki}.
\end{aligned} \tag{2.4}$$

Since $S = S^*$, we have $AB^* = BA^*$. Then (2.2) follows.

Conversely, let

$$[A \mid B] = [(a_1, \dots, a_{2d}) \mid (b_1, \dots, b_{2d})] \in \mathcal{B}^{\mathbb{C}}$$

and $m_0 := \text{rank} A$. Suppose $\text{rank}(a_{k_1}, \dots, a_{k_{m_0}}) = m_0$. Denote $K := \{k_1, \dots, k_{m_0}\}$ and $\{1, 2, \dots, 2d\} \setminus K := \{k_{m_0+1}, \dots, k_{2d}\}$.

Claim: $(a_{k_1}, \dots, a_{k_{m_0}}, b_{k_{m_0+1}}, \dots, b_{k_{2d}})$ is non-degenerate.

Firstly, we can choose $T \in GL(2d, \mathbb{C})$ such that

$$T(A \mid B) = \begin{pmatrix} A_1 & B_1 \\ 0 & B_2 \end{pmatrix} \in \mathcal{L}_{2d, 4d}(\mathbb{C}), \quad A_1, B_1 \in \mathcal{M}_{m_0 \times 2d}, \quad B_2 \in \mathcal{M}_{(2d-m_0) \times 2d},$$

where $\mathcal{M}_{m_0 \times 2d}$ denotes the set of all $m_0 \times 2d$ complex matrices. Let

$$A_1 := (a_{1,1}, \dots, a_{1,2d}), \quad B_2 := (b_{2,1}, \dots, b_{2,2d}),$$

where $a_{1,l} = (a_{1,1l}, \dots, a_{1,m_0l})^T$ and $b_{2,l} = (b_{2,1l}, \dots, b_{2,(2d-m_0)l})^T$, $1 \leq l \leq 2d$. Since $[A \mid B] \in \mathcal{B}^{\mathbb{C}}$, direct computation shows that

$$\begin{aligned} \begin{pmatrix} A_1 \\ 0 \end{pmatrix} \begin{pmatrix} B_1^* & B_2^* \end{pmatrix} &= \begin{pmatrix} B_1 \\ B_2 \end{pmatrix} \begin{pmatrix} A_1^* & 0 \end{pmatrix} \\ \Leftrightarrow A_1 B_1^* &\text{ symmetric and } A_1 B_2^* = 0. \end{aligned}$$

Let

$$\begin{aligned} E &:= (E_1 \mid E_2) = (a_{1,k_1}, \dots, a_{1,k_{m_0}} \mid a_{1,k_{m_0+1}}, \dots, a_{1,k_{2d}}), \\ F &:= (F_1 \mid F_2) = (b_{2,k_1}, \dots, b_{2,k_{m_0}} \mid b_{2,k_{m_0+1}}, \dots, b_{2,k_{2d}}). \end{aligned}$$

Note that $\text{rank} E_1 = \text{rank} E = \text{rank} A_1 = m_0$ and $\text{rank} F = \text{rank} B_2 = 2d - m_0$. Since $A_1 B_2^* = 0$, we have $EF^* = E_1 F_1^* + E_2 F_2^* = 0$. Direct calculation gives

$$(F_2 - F_1 E_1^{-1} E_2) F_2^* = F_2 F_2^* - F_1 E_1^{-1} E_2 F_2^* = F_2 F_2^* + F_1 E_1^{-1} E_1 F_1^* = FF^*,$$

which yields

$$2d - m_0 \geq \text{rank} F_2^* \geq \text{rank} FF^* = \text{rank} F = 2d - m_0.$$

Then $\text{rank} F_2 = 2d - m_0$. Thus

$$(a_{k_1} \cdots, a_{k_{m_0}}, b_{k_{m_0+1}}, \dots, b_{2d}) = T^{-1} \begin{pmatrix} E_1 & * \\ 0 & F_2 \end{pmatrix}$$

is non-degenerate and this claim holds.

Let

$$T_1 := (-a_{k_1}, \dots, -a_{k_{m_0}}, b_{k_{m_0+1}}, \dots, b_{k_{2d}}).$$

Then

$$(T_1^{-1} A \mid T_1^{-1} B) = (a'_1, \dots, a'_{2d}, b'_1, \dots, b'_{2d}),$$

where $a'_{k_i} = -e_i$, $i = 1, \dots, m_0$, $b'_{k_j} = e_j$, $j = m_0 + 1, \dots, 2d$. Let $S = (s_1, \dots, s_{2d}) = (s_{ik}) \in \mathcal{M}_{2d \times 2d}$ and

$$s_l = \begin{cases} a'_l, & l \notin K, \\ b'_l, & l \in K. \end{cases}$$

Since $AB^* = BA^*$, by the calculation in (2.3)–(2.4), we have $s_{ik} = \bar{s}_{ki}$. Thus $[A \mid B] \subset \mathcal{O}_K^{\mathbb{C}}$. Other assertions are direct consequences of properties of Lagrangian-Grassmann manifold [1, 2]. This completes the proof. \square

The product space $\Omega \times \mathcal{B}^{\mathbb{C}}$ is the space of Sturm-Liouville problems, and (ω, \mathbf{A}) is used to stand for an element in $\Omega \times \mathcal{B}^{\mathbb{C}}$.

3 Basic properties of eigenvalues

Firstly, we introduce the following weighted space:

$$L_W^2([a, b], \mathbb{C}^d) = \{y : \int_a^b y(t)^* W(t) y(t) dt < +\infty\},$$

with the inner product $\langle y, z \rangle_W = \int_a^b z(t)^* W(t) y(t) dt$.

Let $(\omega, \mathbf{A}) \in \Omega \times \mathcal{B}^{\mathbb{C}}$ with $\omega = (P, Q, W)$ and $\mathbf{A} = [A \mid B]$. Then the corresponding Sturm-Liouville operator

$$T_{(\omega, \mathbf{A})} y = W^{-1}(-(Py')' + Qy) \quad (3.1)$$

is self-adjoint with the domain

$$D_{\mathbf{A}} = \{y \in L_W^2([a, b], \mathbb{C}^d) : y, Py' \in AC([a, b], \mathbb{C}^d), \\ T_{(\omega, \mathbf{A})} y \in L_W^2([a, b], \mathbb{C}^d), y \text{ satisfies (1.2)}\}. \quad (3.2)$$

For any $\lambda \in \mathbb{C}$, let $\phi_{1,\lambda}, \dots, \phi_{2d,\lambda}$ be the fundamental solutions to (1.1) determined by the initial conditions

$$\begin{pmatrix} \phi_{1,\lambda}(a) & \cdots & \phi_{2d,\lambda}(a) \\ P\phi'_{1,\lambda}(a) & \cdots & P\phi'_{2d,\lambda}(a) \end{pmatrix} = I_{2d}.$$

Denote

$$\Phi_\lambda := \begin{pmatrix} -\phi_{1,\lambda}(a) & \cdots & -\phi_{2d,\lambda}(a) \\ \phi_{1,\lambda}(b) & \cdots & \phi_{2d,\lambda}(b) \end{pmatrix}, \quad (3.3)$$

$$\Psi_\lambda := \begin{pmatrix} P\phi'_{1,\lambda}(a) & \cdots & P\phi'_{2d,\lambda}(a) \\ P\phi'_{1,\lambda}(b) & \cdots & P\phi'_{2d,\lambda}(b) \end{pmatrix}. \quad (3.4)$$

Then Φ_λ and Ψ_λ are entire $2d \times 2d$ -matrix valued functions of λ .

Lemma 3.1. *The spectrum of the d -dimensional Sturm-Liouville problem (ω, \mathbf{A}) consists of isolated eigenvalues, which are all real and bounded from below. Moreover, $\lambda \in \mathbb{R}$ is an eigenvalue of (ω, \mathbf{A}) if and only if λ is a zero of*

$$\Gamma_{(\omega, \mathbf{A})}(\lambda) := \det(A\Phi_\lambda + B\Psi_\lambda). \quad (3.5)$$

Proof. The proof is similar to that of Lemma 4.5 in [29]. \square

Definition 3.1. Let λ be an eigenvalue of $(\boldsymbol{\omega}, \mathbf{A})$. The order of λ as a zero of $\Gamma_{(\boldsymbol{\omega}, \mathbf{A})}$ is called its analytic multiplicity. The number of linearly independent eigenfunctions for λ is called its geometric multiplicity. The dimension of the space $E_\lambda = \{y \in L_W^2([a, b], \mathbb{C}^d) : (T_{(\boldsymbol{\omega}, \mathbf{A})} - \lambda)^k y = 0 \text{ for some integer } k \geq 1\}$ is called its algebraic multiplicity.

We show in Theorem 5.1 that the three multiplicities of an eigenvalue are equal. Thus we shall not distinguish them. The following result is locally continuous dependence of eigenvalues on Sturm-Liouville problems, which can be proved by Rouché's Theorem [4]. See [14, 31] in 1-dimensional case.

Lemma 3.2. Let $r_1 < r_2$ be two real numbers such that neither of them is an eigenvalue of a given Sturm-Liouville problem $(\boldsymbol{\omega}, \mathbf{A})$, and $n \geq 0$ be the number of eigenvalues of $(\boldsymbol{\omega}, \mathbf{A})$ in the interval (r_1, r_2) . Then there exists a neighborhood U of $(\boldsymbol{\omega}, \mathbf{A})$ in $\Omega \times \mathcal{B}^{\mathbb{C}}$ such that each $(\boldsymbol{\sigma}, \mathbf{B}) \in U$ has exactly n eigenvalues in (r_1, r_2) , and neither r_1 nor r_2 is an eigenvalue of $(\boldsymbol{\sigma}, \mathbf{B})$.

Proof. Let $R := \{z \in \mathbb{C} : |z - (r_1 + r_2)/2| < (r_2 - r_1)/2\}$ and $\eta := \min_{\lambda \in \partial R} |\Gamma_{(\boldsymbol{\omega}, \mathbf{A})}(\lambda)|$, where ∂R denotes the boundary of R . Then $\eta > 0$ by Lemma 3.1. By the compactness of ∂R and the uniform continuity of $\Gamma_{(\boldsymbol{\omega}, \mathbf{A})}$ on $(\boldsymbol{\omega}, \mathbf{A})$ and λ , there exists a neighborhood U of $(\boldsymbol{\omega}, \mathbf{A})$ in $\Omega \times \mathcal{B}^{\mathbb{C}}$ such that $|\Gamma_{(\boldsymbol{\sigma}, \mathbf{B})}(\lambda) - \Gamma_{(\boldsymbol{\omega}, \mathbf{A})}(\lambda)| < \eta$ for all $\lambda \in \partial R$ and for all $(\boldsymbol{\sigma}, \mathbf{B}) \in U$, which also implies that $|\Gamma_{(\boldsymbol{\sigma}, \mathbf{B})}(\lambda)| \geq |\Gamma_{(\boldsymbol{\omega}, \mathbf{A})}(\lambda)| - |\Gamma_{(\boldsymbol{\sigma}, \mathbf{B})}(\lambda) - \Gamma_{(\boldsymbol{\omega}, \mathbf{A})}(\lambda)| > 0$. Thus neither r_1 nor r_2 is an eigenvalue of $(\boldsymbol{\sigma}, \mathbf{B}) \in U$. Since $\Gamma_{(\boldsymbol{\sigma}, \mathbf{B})} - \Gamma_{(\boldsymbol{\omega}, \mathbf{A})}$ and $\Gamma_{(\boldsymbol{\omega}, \mathbf{A})}$ are both entire functions of λ , $\Gamma_{(\boldsymbol{\sigma}, \mathbf{B})}$ and $\Gamma_{(\boldsymbol{\omega}, \mathbf{A})}$ have the same number of zeros in R , counting order, by Rouché's Theorem. The proof is complete by the fact of the reality of eigenvalues. \square

The next result is a direct consequence of Lemma 3.2.

Lemma 3.3. Let λ_* be an eigenvalue with multiplicity m of $(\boldsymbol{\omega}, \mathbf{A})$, and $r_1 < r_2$ be two real numbers such that $\lambda_* \in (r_1, r_2)$ is the only eigenvalue of $(\boldsymbol{\omega}, \mathbf{A})$ in the interval $[r_1, r_2]$. Then there exist a connected neighborhood U of $(\boldsymbol{\omega}, \mathbf{A})$ in $\Omega \times \mathcal{B}^{\mathbb{C}}$ and continuous functions $\Lambda_1, \dots, \Lambda_m$ defined on U such that $r_1 < \Lambda_1(\boldsymbol{\sigma}, \mathbf{B}) \leq \dots \leq \Lambda_m(\boldsymbol{\sigma}, \mathbf{B}) < r_2$ for each $(\boldsymbol{\sigma}, \mathbf{B}) \in U$, where $\Lambda_1(\boldsymbol{\sigma}, \mathbf{B}), \dots, \Lambda_m(\boldsymbol{\sigma}, \mathbf{B})$ are eigenvalues of $(\boldsymbol{\sigma}, \mathbf{B})$.

These functions in Lemma 3.3 are locally called continuous eigenvalue branches. When $m = 1$, Λ_1 is called the continuous simple eigenvalue branch. Then we shall make a continuous choice of eigenfunctions for the eigenvalues along a continuous simple eigenvalue branch.

Lemma 3.4. *Let λ_* be a simple eigenvalue of $(\boldsymbol{\omega}, \mathbf{A})$, u_0 be a given eigenfunction for λ_* , and Λ be a continuous simple eigenvalue branch defined on a neighborhood U of $(\boldsymbol{\omega}, \mathbf{A})$ in $\Omega \times \mathcal{B}^{\mathbb{C}}$ through λ_* . Then there exists a neighborhood $U_1 \subset U$ of $(\boldsymbol{\omega}, \mathbf{A})$ such that for any $(\boldsymbol{\sigma}, \mathbf{B}) \in U_1$, there is an eigenfunction $u_{\Lambda(\boldsymbol{\sigma}, \mathbf{B})}$ for $\Lambda(\boldsymbol{\sigma}, \mathbf{B})$ satisfying that $u_{\Lambda(\boldsymbol{\omega}, \mathbf{A})} = u_0$, and $u_{\Lambda(\cdot)}$ and $pu'_{\Lambda(\cdot)}$ are continuous on U_1 in the sense that for any $(\boldsymbol{\sigma}, \mathbf{B}) \in U_1$, $u_{\Lambda(\boldsymbol{\tau}, \mathbf{C})} \rightarrow u_{\Lambda(\boldsymbol{\sigma}, \mathbf{B})}$ and $pu'_{\Lambda(\boldsymbol{\tau}, \mathbf{C})} \rightarrow pu'_{\Lambda(\boldsymbol{\sigma}, \mathbf{B})}$ as $U_1 \ni (\boldsymbol{\tau}, \mathbf{C}) \rightarrow (\boldsymbol{\sigma}, \mathbf{B})$ both uniformly on $[a, b]$.*

Proof. The proof is similar as that of Theorem 3.1 in [17]. □

When $\boldsymbol{\omega}$ (or \mathbf{A}) is fixed, we can get corresponding results on a neighborhood of \mathbf{A} (or $\boldsymbol{\omega}$) as those in Lemmas 3.2–3.4. Then we turn to present the continuity principle for the n -th eigenvalue.

Lemma 3.5. *Let \mathcal{O} be a subset of $\Omega \times \mathcal{B}^{\mathbb{C}}$. If λ_1 is bounded from below on \mathcal{O} , then the restriction of the n -th eigenvalue to \mathcal{O} is continuous for each $n \geq 1$.*

Lemma 3.6. *If \mathcal{O} is a subset of $\Omega \times \mathcal{B}^{\mathbb{C}}$, $(\boldsymbol{\omega}, \mathbf{A}) \notin \mathcal{O}$ is an accumulation point of \mathcal{O} ,*

$$\lim_{\mathcal{O} \ni (\boldsymbol{\sigma}, \mathbf{B}) \rightarrow (\boldsymbol{\omega}, \mathbf{A})} \lambda_n(\boldsymbol{\sigma}, \mathbf{B}) = -\infty$$

for any $n = 1, \dots, m$, where $m \geq 0$, and λ_{m+1} is bounded from below on \mathcal{O} , then

$$\lim_{\mathcal{O} \ni (\boldsymbol{\sigma}, \mathbf{B}) \rightarrow (\boldsymbol{\omega}, \mathbf{A})} \lambda_n(\boldsymbol{\sigma}, \mathbf{B}) = \lambda_{n-m}(\boldsymbol{\omega}, \mathbf{A})$$

for any $n \geq m + 1$.

By using Lemma 3.2, the proofs of Lemmas 3.5–3.6 are similar to those of Theorems 1.40–1.41 in [14], respectively.

4 Analysis on 1-dimensional results

In this section, we reform the singular boundary conditions in the frame (2.1) and refine the results of Theorems 3.39 and 3.76 in [14].

The explicit coordinate systems (2.1) in 1-dimensional case are as follows:

$$\begin{aligned}
\mathcal{O}_\emptyset^{\mathbb{C}} &= \left\{ \begin{bmatrix} s_{11} & s_{12} & 1 & 0 \\ \bar{s}_{12} & s_{22} & 0 & 1 \end{bmatrix} : s_{11}, s_{22} \in \mathbb{R}, s_{12} \in \mathbb{C} \right\}, \\
\mathcal{O}_{\{1\}}^{\mathbb{C}} &= \left\{ \begin{bmatrix} -1 & s_{12} & s_{11} & 0 \\ 0 & s_{22} & \bar{s}_{12} & 1 \end{bmatrix} : s_{11}, s_{22} \in \mathbb{R}, s_{12} \in \mathbb{C} \right\}, \\
\mathcal{O}_{\{2\}}^{\mathbb{C}} &= \left\{ \begin{bmatrix} s_{11} & 0 & 1 & s_{12} \\ \bar{s}_{12} & -1 & 0 & s_{22} \end{bmatrix} : s_{11}, s_{22} \in \mathbb{R}, s_{12} \in \mathbb{C} \right\}, \\
\mathcal{O}_{\{1,2\}}^{\mathbb{C}} &= \left\{ \begin{bmatrix} -1 & 0 & s_{11} & s_{12} \\ 0 & -1 & \bar{s}_{12} & s_{22} \end{bmatrix} : s_{11}, s_{22} \in \mathbb{R}, s_{12} \in \mathbb{C} \right\}.
\end{aligned} \tag{4.1}$$

In order to refine the above results, we need the following notation, which will be used for any $d \geq 1$ in the sequel. For a nonempty subset $K = \{n_1, \dots, n_{m_0}\} \subset \{1, \dots, 2d\}$ and for any $\mathbf{A} \in \mathcal{O}_K^{\mathbb{C}}$, let

$$S_K(\mathbf{A}) = \begin{pmatrix} s_{n_1 n_1} & s_{n_1 n_2} & \cdots & s_{n_1 n_{m_0}} \\ \bar{s}_{n_1 n_2} & s_{n_2 n_2} & \cdots & s_{n_2 n_{m_0}} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{s}_{n_1 n_{m_0}} & \bar{s}_{n_2 n_{m_0}} & \cdots & s_{n_{m_0} n_{m_0}} \end{pmatrix}. \tag{4.2}$$

In 1-dimensional case, $S_{\{i\}}(\mathbf{A}) = (s_{ii})$, $i = 1, 2$, and $S_{\{1,2\}}(\mathbf{A}) = \begin{pmatrix} s_{11} & s_{12} \\ \bar{s}_{12} & s_{22} \end{pmatrix}$.

Let $n^-(S_K(\mathbf{A}))$, $n^0(S_K(\mathbf{A}))$ and $n^+(S_K(\mathbf{A}))$ denote the total multiplicity of negative, zero and positive eigenvalues of $S_K(\mathbf{A})$, respectively. For a nonempty subset $K \subset \{1, \dots, 2d\}$, define

$$\begin{aligned}
&J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)} \\
&:= \{ \mathbf{A} \in \mathcal{O}_K^{\mathbb{C}} \mid n^0(S_K(\mathbf{A})) = n^0, n^+(S_K(\mathbf{A})) = n^+, n^-(S_K(\mathbf{A})) = n^- \}
\end{aligned} \tag{4.3}$$

for three nonnegative integers n^0 , n^+ and n^- satisfying $n^0 + n^+ + n^- = \sharp(K)$. Then we get the following refinement from Theorem 3.76 in [14]. One key fact in the following proposition is that $n^+(S_K(\mathbf{B})) - n^+(S_K(\mathbf{A}))$ is the number of eigenvalues which tend to $-\infty$ as $\mathbf{B} \rightarrow \mathbf{A}$.

Proposition 4.1. (i) *The restriction of λ_n to $\Sigma_k^{\mathbb{C}}$ is continuous for each $n \geq 1$, where $k = 0, 1, 2$.*

(ii) *Consider the restriction of λ_n to $\mathcal{O}_K^{\mathbb{C}}$ for each $n \geq 1$, where $K \subset \{1, 2\}$.*

- (iia) The restriction of λ_n to $\mathcal{O}_0^{\mathbb{C}}$ is continuous.
- (iib) Let K be nonempty, $0 \leq n^0 < n_1^0 \leq \sharp(K)$, $n^+ \geq n_1^+$ and $n^- \geq n_1^-$. Then for any $\mathbf{A} \in J_{\mathcal{O}_K^{\mathbb{C}}}^{(n_1^0, n_1^+, n_1^-)}$, we have

$$\lim_{J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)} \ni \mathbf{B} \rightarrow \mathbf{A}} \lambda_n(\mathbf{B}) = -\infty, \quad n \leq n^+ - n_1^+,$$

$$\lim_{J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)} \ni \mathbf{B} \rightarrow \mathbf{A}} \lambda_n(\mathbf{B}) = \lambda_{n-(n^+-n_1^+)}(\mathbf{A}), \quad n > n^+ - n_1^+.$$

Remark 4.1. Inspired of [19, 20, 21], we provide an intuitional representation of sets in the space of real boundary conditions, which is also helpful to understand the global concept of singular boundary conditions. Let

$$\begin{aligned} \mathcal{O}_{12,++}^{\mathbb{R}} &= \{ \mathbf{A} \in \mathcal{O}_{\{1,2\}}^{\mathbb{R}} : s_{11} > 0, s_{12} \in \mathbb{R}, s_{11}s_{22} > |s_{12}|^2 \}, \\ \mathcal{O}_{12,--}^{\mathbb{R}} &= \{ \mathbf{A} \in \mathcal{O}_{\{1,2\}}^{\mathbb{R}} : s_{11} < 0, s_{12} \in \mathbb{R}, s_{11}s_{22} > |s_{12}|^2 \}, \\ \mathcal{O}_{12,+-}^{\mathbb{R}} &= \{ \mathbf{A} \in \mathcal{O}_{\{1,2\}}^{\mathbb{R}} : s_{12} \in \mathbb{R}, s_{11}s_{22} < |s_{12}|^2 \}, \\ \mathcal{O}_{12,+0}^{\mathbb{R}} &= \{ \mathbf{A} \in \mathcal{O}_{\{1,2\}}^{\mathbb{R}} : s_{11} + s_{22} > 0, s_{12} \in \mathbb{R}, s_{11}s_{22} = |s_{12}|^2 \}, \\ \mathcal{O}_{12,-0}^{\mathbb{R}} &= \{ \mathbf{A} \in \mathcal{O}_{\{1,2\}}^{\mathbb{R}} : s_{11} + s_{22} < 0, s_{12} \in \mathbb{R}, s_{11}s_{22} = |s_{12}|^2 \}. \end{aligned}$$

Consider an element $\mathbf{A} = [-I \mid S]$ in the coordinate chart $(\mathcal{O}_{\{1,2\}}^{\mathbb{R}}, \phi_{12})$, where $\phi_{12} : (\mathbf{A} \rightarrow S := \begin{pmatrix} s_{11} & s_{12} \\ s_{12} & s_{22} \end{pmatrix})$. Firstly we define a map

$$\begin{aligned} \text{rep}_{12} &= \tilde{\Delta}^{-1} \circ \phi_{12}^{-1} : \{S \mid s_{11}, s_{12}, s_{22} \in \mathbb{R}\} \rightarrow (D^1 \times S^1)/f, \\ &S \mapsto [(r, z, \theta)], \end{aligned}$$

where

$$\begin{aligned} [(r, z, \theta)] \in (D^1 \times S^1)/f &= \{(r, z, \theta) \mid (r-2)^2 + z^2 < 1\} \cup \\ &\{(r, z, \theta), (r, z, \theta + \pi)\} \mid (r-2)^2 + z^2 = 1\}, \end{aligned}$$

the map $\tilde{\Delta} : (D^1 \times S^1)/f \rightarrow \text{Lag}(2, \mathbb{R}) \simeq \mathcal{B}^{\mathbb{R}}$ is a homeomorphism defined in the proof of Theorem 1 in [19].

Under the map rep_{12} , we obtain $\text{rep}_{12}(0) = (2, 0, 0)$ and $\text{rep}_{12}(-\tan \frac{\theta}{2} I_2) =$




Figure 1: The intuitional representation of $\mathcal{O}_{12,++}^{\mathbb{R}}, \mathcal{O}_{12,+-}^{\mathbb{R}}, \mathcal{O}_{12,--}^{\mathbb{R}}, \mathcal{O}_{12,+0}^{\mathbb{R}}, \mathcal{O}_{12,-0}^{\mathbb{R}}$.

$(2, 0, \theta)$ with $\theta \in \mathbb{R}/2\pi\mathbb{Z}$. Furthermore,

$$\phi_{12}^{-1}\left(-\tan\frac{\theta}{2}I_2\right) \text{ is } \begin{cases} \text{in } \mathcal{O}_{12,--}^{\mathbb{R}}, \text{ if } \theta \in (0, \pi), \\ \text{Dirichlet boundary condition, if } \theta = 0, \\ \text{in } \mathcal{O}_{12,++}^{\mathbb{R}}, \text{ if } \theta \in (-\pi, 0), \\ \text{converging to Neumann boundary condition,} \\ \text{if } \theta \rightarrow \pm\pi. \end{cases}$$

In Figure 1, the inner torus with two shrinking points denotes two singular cycles, i.e. the part with $x \leq 0$ (or $x \geq 0$) is the set of elements which have common subspace with Neumann (or Dirichlet) boundary condition ($\text{rep}_{12}(-2, 0, 0)$) ($\text{rep}_{12}(2, 0, 0)$). Note that inner torus lies on the outside torus if $x = 0$.

Since the left part ($x \leq 0$) of the inner torus is not in the range of rep_{12} , by the path connectedness of $\mathcal{O}_{12,++}^{\mathbb{R}}, \mathcal{O}_{12,+-}^{\mathbb{R}}, \mathcal{O}_{12,--}^{\mathbb{R}}, \mathcal{O}_{12,+0}^{\mathbb{R}}$ and $\mathcal{O}_{12,-0}^{\mathbb{R}}$ we have Region I denotes $\mathcal{O}_{12,++}^{\mathbb{R}}$, i.e. the inside of the inner torus under Y -axis. The curve $\text{rep}_{12}\left(-\frac{\tan\theta}{2}I_2\right)$, $\theta \in (-\pi, 0)$ is in Region I.

Region II denotes $\mathcal{O}_{12,--}^{\mathbb{R}}$, i.e. the inside of the inner torus upper Y -axis. The curve $\text{rep}_{12}\left(-\frac{\tan\theta}{2}I_2\right)$, $\theta \in (0, \pi)$ is in Region II.

Region III denotes $\mathcal{O}_{12,+-}^{\mathbb{R}}$, i.e. the outside of the inner torus in $(D^1 \times S^1)/f$.

Region IV denotes $\mathcal{O}_{12,+0}^{\mathbb{R}}$ and $\mathcal{O}_{12,-0}^{\mathbb{R}}$. More precisely, the parts of the inner torus with $X > 0, Y < 0$ and with $X, Y > 0$ denote $\mathcal{O}_{12,+0}^{\mathbb{R}}$ and $\mathcal{O}_{12,-0}^{\mathbb{R}}$, respectively.

5 Equality of multiplicities of an eigenvalue

In this section, we show the equivalence of the analytic, algebraic and geometric multiplicities of an eigenvalue.

Theorem 5.1. *The analytic, algebraic and geometric multiplicities of an eigenvalue of $(\boldsymbol{\omega}, \mathbf{A}) \in \Omega \times \mathcal{B}^{\mathbb{C}}$ are equal.*

Proof. Fix λ_* be an eigenvalue of $(\boldsymbol{\omega}, \mathbf{A})$ with $\boldsymbol{\omega} = (P, Q, W)$ and $\mathbf{A} = [A \mid B]$. Since $T_{(\boldsymbol{\omega}, \mathbf{A})}$ is self-adjoint, the algebraic and geometric multiplicities of λ_* coincide. We shall show that the analytic and geometric ones are the same.

By p and κ denote the geometric and analytic multiplicities of λ_* , respectively. Let φ_i , $1 \leq i \leq p$, be the linearly independent eigenfunctions of λ_* . Choose φ_i , $p+1 \leq i \leq 2d$ be the solutions of (1.1) with $\lambda = \lambda_*$ such that φ_i , $1 \leq i \leq 2d$, are linearly independent. Let $y_i(\cdot, \lambda)$ be the solutions of (1.1) with $\lambda \in \mathbb{C}$ such that $y_i(a, \lambda) = \varphi_i(a)$, $P y_i'(a, \lambda) = P \varphi_i'(a)$, $1 \leq i \leq 2d$. Then $y_i(\cdot, \lambda_*) = \varphi_i$ and y_i has the Taylor expansion

$$y_i(x, \lambda) = \sum_{j=0}^{\infty} \varphi_i^{(j)}(x) (\lambda - \lambda_*)^j$$

with $\varphi_i^{(0)} = \varphi_i$, $1 \leq i \leq 2d$. Let $\tilde{\Phi}_\lambda = \begin{pmatrix} y_1 & \cdots & y_{2d} \\ P y_1' & \cdots & P y_{2d}' \end{pmatrix}$. By (3.5), we have

$$\begin{aligned} & \Gamma_{(\boldsymbol{\omega}, \mathbf{A})}(\lambda) \\ &= \det(\tilde{A} + \tilde{B} \tilde{\Phi}_\lambda(b) \tilde{\Phi}_\lambda^{-1}(a)) = \det(\tilde{A} \tilde{\Phi}_\lambda(a) + \tilde{B} \tilde{\Phi}_\lambda(b)) \det(\tilde{\Phi}_\lambda^{-1}(a)), \end{aligned} \quad (5.1)$$

where

$$(\tilde{A}, \tilde{B}) = (A, B) \begin{pmatrix} -I_d & 0 & 0 & 0 \\ 0 & 0 & I_d & 0 \\ 0 & I_d & 0 & 0 \\ 0 & 0 & 0 & I_d \end{pmatrix}.$$

Note that

$$\tilde{A} \begin{pmatrix} \varphi_i(a) \\ P \varphi_i'(a) \end{pmatrix} + \tilde{B} \begin{pmatrix} \varphi_i(b) \\ P \varphi_i'(b) \end{pmatrix} = 0, \quad 1 \leq i \leq p,$$

which yields that the i -th column of $\Gamma_{(\boldsymbol{\omega}, \mathbf{A})}(\lambda)$ must contain the factor $(\lambda - \lambda_*)$. So

$$\Gamma_{(\boldsymbol{\omega}, \mathbf{A})}(\lambda) = (\lambda - \lambda_*)^p \tilde{\Gamma}_{(\boldsymbol{\omega}, \mathbf{A})}(\lambda).$$

It suffices to show that $\tilde{\Gamma}_{(\omega, \mathbf{A})}(\lambda_*) \neq 0$. Let

$$\hat{\Phi} = \begin{pmatrix} \varphi_1^{(1)} & \cdots & \varphi_p^{(1)} & \varphi_{p+1} & \cdots & \varphi_{2d} \\ P\varphi_1^{(1)'} & \cdots & P\varphi_p^{(1)'} & P\varphi_{p+1}' & \cdots & P\varphi_{2d}' \end{pmatrix}.$$

Then

$$\tilde{\Gamma}_{(\omega, \mathbf{A})}(\lambda_*) = \det(\tilde{A}\hat{\Phi}(a) + \tilde{B}\hat{\Phi}(b)) \det(\tilde{\Phi}_{\lambda_*}^{-1}(a)). \quad (5.2)$$

Suppose $\tilde{\Gamma}_{(\omega, \mathbf{A})}(\lambda_*) = 0$. By χ_1, \dots, χ_{2d} denote the columns of $\tilde{A}\hat{\Phi}(a) + \tilde{B}\hat{\Phi}(b)$. Then

$$\sum_{i=1}^{2d} c_i \chi_i = 0 \quad (5.3)$$

for $c_1, \dots, c_{2d} \in \mathbb{C}$ to be not all vanished. We divide the discussion into two cases below.

Case 1. $c_1 = \dots = c_p = 0$.

Let $\psi = \sum_{i=p+1}^{2d} c_i \varphi_i$. Then ψ is a nontrivial solution of (1.1) since $\varphi_{p+1}, \dots, \varphi_{2d}$ are linearly independent solutions of (1.1) with $\lambda = \lambda_*$. By (5.3), we get

$$\tilde{A} \begin{pmatrix} \psi(a) \\ P\psi'(a) \end{pmatrix} + \tilde{B} \begin{pmatrix} \psi(b) \\ P\psi'(b) \end{pmatrix} = 0,$$

which implies that ψ is an eigenfunction for λ_* . Thus $\psi = \sum_{i=1}^p d_i \varphi_i = \sum_{i=p+1}^{2d} c_i \varphi_i$ with d_1, \dots, d_p to be not all zero, which is a contradiction since $\varphi_1, \dots, \varphi_{2d}$ are linearly independent.

Case 2. c_1, \dots, c_p are not all vanished.

Let

$$\tilde{y} = \sum_{i=1}^p c_i y_i + \sum_{i=p+1}^{2d} c_i (\lambda - \lambda_*) y_i.$$

Then \tilde{y} is a nontrivial solution of

$$-(P\tilde{y}')' + Q\tilde{y} = \lambda W\tilde{y}.$$

It is obvious that $\tilde{y}(\cdot, \lambda_*)$ satisfies the boundary condition **A**. Differentiating the above equation by λ , we get

$$-(P(\frac{\partial \tilde{y}}{\partial \lambda})')' + Q \frac{\partial \tilde{y}}{\partial \lambda} = W \tilde{y} + \lambda W \frac{\partial \tilde{y}}{\partial \lambda}. \quad (5.4)$$

Let $\lambda = \lambda_*$. Then $\frac{\partial \tilde{y}}{\partial \lambda}|_{\lambda=\lambda_*}$ is nontrivial since otherwise, by (5.4) $\tilde{y}(\cdot, \lambda_*) = 0$, which is a contradiction. Clearly, $\frac{\partial \tilde{y}}{\partial \lambda}|_{\lambda=\lambda_*} \in L^2_W([a, b], \mathbb{C}^d)$. If in addition $\frac{\partial \tilde{y}}{\partial \lambda}|_{\lambda=\lambda_*}$ satisfies the boundary condition **A**, we have

$$(T_{(\omega, \mathbf{A})} - \lambda_*) \frac{\partial \tilde{y}}{\partial \lambda}|_{\lambda=\lambda_*} = \tilde{y}(\cdot, \lambda_*). \quad (5.5)$$

Now we show that $\frac{\partial \tilde{y}}{\partial \lambda}|_{\lambda=\lambda_*}$ satisfies **A**. Let $\tilde{\psi} = \sum_{i=1}^p c_i \varphi_i^{(1)} + \sum_{i=p+1}^{2d} c_i \varphi_i$. It follows from (5.3) that

$$\tilde{A} \begin{pmatrix} \tilde{\psi}(a) \\ P \tilde{\psi}'(a) \end{pmatrix} + \tilde{B} \begin{pmatrix} \tilde{\psi}(b) \\ P \tilde{\psi}'(b) \end{pmatrix} = 0. \quad (5.6)$$

Note that $\frac{\partial y_i}{\partial \lambda}|_{\lambda=\lambda_*} = \varphi_i^{(1)}$, $1 \leq i \leq p$, and $y_i(\cdot, \lambda_*) = \varphi_i$, $p < i \leq 2d$. Thus

$$\frac{\partial \tilde{y}}{\partial \lambda}|_{\lambda=\lambda_*} = \sum_{i=1}^p c_i \varphi_i^{(1)} + \sum_{i=p+1}^{2d} c_i \varphi_i = \tilde{\psi}.$$

(5.6) implies that $\frac{\partial \tilde{y}}{\partial \lambda}|_{\lambda=\lambda_*}$ satisfies **A**. By (5.5), $\frac{\partial \tilde{y}}{\partial \lambda}|_{\lambda=\lambda_*}$ is a generalized eigenfunction of λ_* , which contradicts the fact that the geometric and algebraic multiplicities of λ_* are equal.

Therefore, $\tilde{\Gamma}_{(\omega, \mathbf{A})}(\lambda_*) \neq 0$ in any case and $\kappa = p$. This completes the proof. \square

Remark 5.1. Here our proof is independent of the condition that P is positive. Thus Theorem 5.1 also holds true when P is invertible and non-positive a.e. on $[a, b]$.

6 Continuity of the n -th eigenvalue

In this section, we prove that the n -th eigenvalue is continuously dependent on the Sturm-Liouville equations and boundary conditions when restricted into the k -th layer, where $0 \leq k \leq 2d$.

Theorem 6.1. *The n -th eigenvalue is continuous on $\Omega \times \Sigma_k^{\mathbb{C}}$ for each $n \geq 1$, where $0 \leq k \leq 2d$.*

Proof. Let $(\boldsymbol{\omega}_1, \mathbf{A}_1) \in \Omega \times \Sigma_k^{\mathbb{C}}$, where $\boldsymbol{\omega}_1 = (P_1, Q_1, W_1)$. By Lemma 3.5, it suffices to show that there exists a neighborhood U_1 of $(\boldsymbol{\omega}_1, \mathbf{A}_1)$ in $\Omega \times \Sigma_k^{\mathbb{C}}$ such that the first eigenvalue λ_1 is bounded from below on U_1 . It is equivalent to show that there exists $\mu \in \mathbb{R}$ such that

$$\langle T_{(\boldsymbol{\omega}, \mathbf{A})} y, y \rangle_W \geq \mu \|y\|_W^2, \quad \forall y \in D_{\mathbf{A}},$$

uniformly for $(\boldsymbol{\omega}, \mathbf{A}) \in U_1$, where $T_{(\boldsymbol{\omega}, \mathbf{A})}$ and $D_{\mathbf{A}}$ are defined in (3.1) and (3.2).

Firstly, let U_1 be chosen sufficiently small that for any $(\boldsymbol{\omega}, \mathbf{A}) \in U_1$ with $\boldsymbol{\omega} = (P, Q, W)$, there exist $\delta_1, \mu_1 > 0$ satisfying $\|\boldsymbol{\omega} - \boldsymbol{\omega}_1\|_{L^\infty \times L^\infty \times L^\infty} < \delta_1$ and

$$P(t) \geq \mu_1, \quad W(t) \geq \mu_1, \quad a.e. \text{ on } t \in [a, b]. \quad (6.1)$$

Direct computations show that for any $(\boldsymbol{\omega}, \mathbf{A}) \in U_1$ with $\boldsymbol{\omega} = (P, Q, W)$ and any $y \in D_{\mathbf{A}}$,

$$\begin{aligned} \langle T_{(\boldsymbol{\omega}, \mathbf{A})} y, y \rangle_W &= \int_a^b (-(Py')' + Qy, y)_d dt \\ &= \int_a^b (Py', y')_d + (Qy, y)_d dt - \left(\begin{pmatrix} Py'(a) \\ Py'(b) \end{pmatrix}, \begin{pmatrix} -y(a) \\ y(b) \end{pmatrix} \right)_{2d}, \end{aligned} \quad (6.2)$$

where $(\cdot, \cdot)_d$ is the usual inner product in \mathbb{C}^d .

Let $\mathbf{A} = [A \mid B] \in \Sigma_k^{\mathbb{C}}$. Choose $T_0 \in GL(2d, \mathbb{C})$ such that

$$T_0(A \mid B) = \begin{pmatrix} A_1 & B_1 \\ A_2 & B_2 \end{pmatrix},$$

where $A_1, B_1 \in \mathcal{M}_{(2d-k) \times 2d}$, $A_2, B_2 \in \mathcal{M}_{k \times 2d}$ and $\text{rank} B_1 = \text{rank} B = 2d - k$. Then exists the unique $E \in \mathcal{M}_{k \times (2d-k)}$ such that $B_2 = EB_1$, and E is locally continuously dependent on B . Direct computation shows that

$$\begin{pmatrix} I_{2d-k} & 0 \\ -E & I_k \end{pmatrix} \begin{pmatrix} A_1 & B_1 \\ A_2 & B_2 \end{pmatrix} = \begin{pmatrix} A_1 & B_1 \\ A_2 - EA_1 & 0 \end{pmatrix}.$$

Applying the QR decomposition (see Theorem 2.1.14 in [10]) on B_1 and $A_2 - EA_1$, we get

$$B_1 = L_1 D_1, \quad A_2 - EA_1 = L_2 C_2,$$

where $L_1 \in GL(2d - k, \mathbb{C})$ and $L_2 \in GL(k, \mathbb{C})$ are lower triangular positive matrices, and $D_1 \in \mathcal{M}_{(2d-k) \times 2d}$ and $C_2 \in \mathcal{M}_{k \times 2d}$ satisfy $D_1 D_1^* = I_{2d-k}$, $C_2 C_2^* = I_k$. Let $C_1 = L_1^{-1} A_1$. Note that L_i, C_i and $D_1, i = 1, 2$, are uniquely determined and locally continuously dependent on \mathbf{A} . Thus we denote them by $L_i(\mathbf{A}), C_i(\mathbf{A})$ and $D_1(\mathbf{A})$. Direct calculation gives

$$\begin{pmatrix} L(\mathbf{A})_1^{-1} & 0 \\ 0 & L(\mathbf{A})_2^{-1} \end{pmatrix} \begin{pmatrix} A_1 & B_1 \\ A_2 - EA_1 & 0 \end{pmatrix} = \begin{pmatrix} C_1(\mathbf{A}) & D_1(\mathbf{A}) \\ C_2(\mathbf{A}) & 0 \end{pmatrix}.$$

Since $\begin{pmatrix} C_1(\mathbf{A}) \\ C_2(\mathbf{A}) \end{pmatrix} (D_1(\mathbf{A})^* \ 0) = \begin{pmatrix} D_1(\mathbf{A}) \\ 0 \end{pmatrix} (C_1(\mathbf{A})^* \ C_2(\mathbf{A})^*)$, we have

$$C_2(\mathbf{A}) D_1(\mathbf{A})^* = 0 \text{ and } C_1(\mathbf{A}) D_1(\mathbf{A})^* = D_1(\mathbf{A}) C_1(\mathbf{A})^*.$$

Thus $E_1(\mathbf{A}) E_1(\mathbf{A})^* = I_{2d}$, which implies that $E_1(\mathbf{A})$ is unitary, where $E_1(\mathbf{A}) := \begin{pmatrix} C_2(\mathbf{A}) \\ D_1(\mathbf{A}) \end{pmatrix}$. It follows from the boundary condition that

$$C_1(\mathbf{A}) \begin{pmatrix} -y(a) \\ y(b) \end{pmatrix} + D_1(\mathbf{A}) \begin{pmatrix} Py'(a) \\ Py'(b) \end{pmatrix} = 0, \quad C_2(\mathbf{A}) \begin{pmatrix} -y(a) \\ y(b) \end{pmatrix} = 0.$$

Then

$$\begin{aligned} & \left(\begin{pmatrix} Py'(a) \\ Py'(b) \end{pmatrix}, \begin{pmatrix} -y(a) \\ y(b) \end{pmatrix} \right)_{2d} \\ &= \left(E_1(\mathbf{A}) \begin{pmatrix} Py'(a) \\ Py'(b) \end{pmatrix}, E_1(\mathbf{A}) \begin{pmatrix} -y(a) \\ y(b) \end{pmatrix} \right)_{2d} \\ &= \left(D_1(\mathbf{A}) \begin{pmatrix} Py'(a) \\ Py'(b) \end{pmatrix}, D_1(\mathbf{A}) \begin{pmatrix} -y(a) \\ y(b) \end{pmatrix} \right)_{2d} \\ &= \left(-C_1(\mathbf{A}) \begin{pmatrix} -y(a) \\ y(b) \end{pmatrix}, D_1(\mathbf{A}) \begin{pmatrix} -y(a) \\ y(b) \end{pmatrix} \right)_{2d} \\ &\leq c \|C_1(\mathbf{A})\|_{\mathcal{M}_{(2d-k) \times 2d}} \|D_1(\mathbf{A})\|_{\mathcal{M}_{(2d-k) \times 2d}} (|y(a)|_d^2 + |y(b)|_d^2) \\ &\leq c \|C_1(\mathbf{A})\|_{\mathcal{M}_{(2d-k) \times 2d}} \|D_1(\mathbf{A})\|_{\mathcal{M}_{(2d-k) \times 2d}} \|y\|_{C^0}^2, \end{aligned} \tag{6.3}$$

where $\|C\|_{\mathcal{M}_{(2d-k) \times 2d}} = \max_{ij} \{ |c_{ij}| \}$ and $\|y\|_{C^0} = \max_{t \in [a, b]} \{ |y(t)|_d \}$. Here and in the sequel, c denotes a generic positive constant and $c(\alpha)$ denotes such a constant depending only on α . We now interpolate $\|y\|_{C^0}$ between

the norms $\|y\|_{L^2}$ and $\|y'\|_{L^2}$. Choose $\varepsilon_1 > 0$ sufficiently small and $c(\varepsilon_1) > 0$ sufficiently large such that

$$c\|C_1(\mathbf{A}_1)\|_{\mathcal{M}_{(2d-k)\times 2d}}\|D_1(\mathbf{A}_1)\|_{\mathcal{M}_{(2d-k)\times 2d}}\varepsilon_1 < \mu_1 \quad (6.4)$$

and

$$\|y\|_{C^0}^2 \leq \varepsilon_1\|y'\|_{L^2}^2 + c(\varepsilon_1)\|y\|_{L^2}^2. \quad (6.5)$$

It follows from (6.4) and the locally continuity of $C_1(\mathbf{A})$ and $D_1(\mathbf{A})$ on \mathbf{A} that U_1 can be shrunk such that for any $(\boldsymbol{\omega}, \mathbf{A}) \in U_1$,

$$c\|C_1(\mathbf{A})\|_{\mathcal{M}_{(2d-k)\times 2d}}\|D_1(\mathbf{A})\|_{\mathcal{M}_{(2d-k)\times 2d}}\varepsilon_1 < \mu_1. \quad (6.6)$$

By (6.1)–(6.3), (6.5)–(6.6) and noting that $\|\cdot\|_{L^2}$ is equivalent to $\|\cdot\|_{L_W^2}$, we have

$$\begin{aligned} & \langle T_{(\boldsymbol{\omega}, \mathbf{A})}y, y \rangle_W \\ & \geq \mu_1\|y'\|_{L^2}^2 - (\|Q_1\|_{L^\infty} + \delta_1)\|y\|_{L^2}^2 - c\|C_1(\mathbf{A})\|_{\mathcal{M}_{(2d-k)\times 2d}}\|D_1(\mathbf{A})\|_{\mathcal{M}_{(2d-k)\times 2d}} \\ & \quad \varepsilon_1\|y'\|_{L^2}^2 - c\|C_1(\mathbf{A})\|_{\mathcal{M}_{(2d-k)\times 2d}}\|D_1(\mathbf{A})\|_{\mathcal{M}_{(2d-k)\times 2d}}c(\varepsilon_1)\|y\|_{L^2}^2 \\ & \geq \mu\|y\|_{L_W^2}^2 \end{aligned}$$

for some $\mu \in \mathbb{R}$ and for all $(\boldsymbol{\omega}, \mathbf{A}) \in U_1$. The proof is complete. \square

7 Singularity of the n -th eigenvalue

In this section, we determine the singular set in the space of boundary conditions for d -dimensional Sturm-Liouville problems, and give complete characterization of asymptotic behavior of the n -th eigenvalue near any fixed singular boundary condition.

Theorem 7.1. *Consider the restriction of λ_n to $\mathcal{O}_K^{\mathbb{C}}$ for each $n \geq 1$, where $K \subset \{1, \dots, 2d\}$.*

(i) *The restriction of λ_n to $\mathcal{O}_\emptyset^{\mathbb{C}}$ is continuous.*

(ii) Let K be nonempty, $0 \leq n^0 < n_0^0 \leq \sharp(K)$, $n^+ \geq n_0^+$ and $n^- \geq n_0^-$. Then for any $\mathbf{A} \in J_{\mathcal{O}_K^{\mathbb{C}}}^{(n_0^0, n_0^+, n_0^-)}$, we have

$$\begin{aligned} \lim_{J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)} \ni \mathbf{B} \rightarrow \mathbf{A}} \lambda_n(\mathbf{B}) &= -\infty, \quad n \leq n^+ - n_0^+, \\ \lim_{J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)} \ni \mathbf{B} \rightarrow \mathbf{A}} \lambda_n(\mathbf{B}) &= \lambda_{n-(n^+-n_0^+)}(\mathbf{A}), \quad n > n^+ - n_0^+, \end{aligned}$$

where $J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ is defined in (4.3).

Consequently, $\Sigma^{\mathbb{C}} = \{\mathbf{A} = [A | B] : n^0(B) > 0\}$.

Proof. (i) holds due to Theorem 6.1 and the fact $\mathcal{O}_\emptyset^{\mathbb{C}} \subset \Sigma_0^{\mathbb{C}}$. We shall give the proof of Theorem 7.1 (ii) step by step via the following results in Lemmas 7.3, 7.4, 7.5 and 7.6. \square

By Theorems 6.1 and 7.1, we give the explicit description of U^i in (1.6) and (1.7):

Corollary 7.1. For each $\mathbf{A} \in J_{\mathcal{O}_K^{\mathbb{C}}}^{(n_0^0, n_0^+, n_0^-)}$ and $0 \leq i \leq n_0^0$, there exists a neighborhood U of \mathbf{A} in $\mathcal{O}_K^{\mathbb{C}}$ such that

$$U^i = \bigcup_{n^+ - n_0^+ = i, n^0 \leq n_0^0, n^- \geq n_0^-} J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)} \cap U,$$

where U^i satisfies (1.6) and (1.7).

Let $K \subset \{1, \dots, 2d\}$ be a nonempty subset below. For each $0 \leq n^0 \leq \sharp(K)$,

$$\mathcal{O}_K^{\mathbb{C}} \cap \Sigma_{n^0}^{\mathbb{C}} \begin{cases} \neq \emptyset & \text{if } 0 \leq n^0 \leq \sharp(K), \\ = \emptyset & \text{if } \sharp(K) < n^0 \leq 2d, \end{cases}$$

and furthermore, $n^0(S_K(\mathbf{A})) = n^0$ for any $\mathbf{A} \in \mathcal{O}_K^{\mathbb{C}} \cap \Sigma_{n^0}^{\mathbb{C}}$. $\mathcal{O}_K^{\mathbb{C}} \cap \Sigma_{n^0}^{\mathbb{C}}$ possesses precisely $\sharp(K) - n^0 + 1$ components as follows:

$$J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}, \quad 0 \leq n^+ \leq \sharp(K) - n^0, \quad n^- = \sharp(K) - n^0 - n^+. \quad (7.1)$$

Next, we show that every $J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$ is a path connected component. In the following discussion, by a path γ to connect x_0 and x_1 in a topological space X , we mean a continuous function $\gamma : [0, 1] \rightarrow X$ such that $\gamma(0) = x_0$ and $\gamma(1) = x_1$.

Lemma 7.1. $J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$ is path connected for each $0 \leq n^0 \leq \sharp(K)$ and $0 \leq n^+ \leq \sharp(K) - n^0$.

Proof. Fix any given $\mathbf{A}_j \in J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$, $j = 1, 2$, and denote

$$\mathbf{A}_j = [A_j \mid B_j] = [(a_1^{(j)}, a_2^{(j)}, \dots, a_{2d}^{(j)}) \mid (b_1^{(j)}, b_2^{(j)}, \dots, b_{2d}^{(j)})],$$

where

$$a_i^{(j)} = \begin{cases} -e_i & \text{if } i \in K, \\ s_i^{(j)} & \text{if } i \in \{1, 2, \dots, 2d\} \setminus K, \end{cases}$$

$$b_i^{(j)} = \begin{cases} s_i^{(j)} & \text{if } i \in K, \\ e_i & \text{if } i \in \{1, 2, \dots, 2d\} \setminus K, \end{cases}$$

for two different $2d \times 2d$ Hermitian matrices $S^{(j)} = (s_1^{(j)}, \dots, s_{2d}^{(j)})$, $s_i^{(j)} = (s_{1i}^{(j)}, \dots, s_{2di}^{(j)})^T$. Let $m_0 := \sharp(K)$ and $S_K(\mathbf{A}_j)$ be defined as that in (4.2). Then there exist matrices $R^{(j)} \in GL(m_0, \mathbb{C})$, $j = 1, 2$, such that

$$S_K(\mathbf{A}_j) = R^{(j)*} \hat{J} R^{(j)}, \hat{J} := \begin{pmatrix} 0_{n^0} & & \\ & I_{n^+} & \\ & & -I_{n^-} \end{pmatrix}. \quad (7.2)$$

Choose a path of $m_0 \times m_0$ matrices: γ to connect $R^{(1)}$ and $R^{(2)}$ such that $\gamma(\tau) \in GL(m_0, \mathbb{C})$, $\tau \in [0, 1]$. Then $\gamma_0(\tau) := \gamma(\tau)^* \hat{J} \gamma(\tau)$ with entries $(\gamma_0(\tau))_{ij}$, $1 \leq i, j \leq m_0$, is a path to connect $S_K(\mathbf{A}_1)$ and $S_K(\mathbf{A}_2)$ such that $n^0(\gamma_0(\tau)) = n^0$ and $n^\pm(\gamma_0(\tau)) = n^\pm$, $\tau \in [0, 1]$. Define a Hermitian $2d \times 2d$ matrix $S^{(\tau)} = (s_1^{(\tau)}, \dots, s_{2d}^{(\tau)})$, $\tau \in [0, 1]$, where $s_i^{(\tau)} = (s_{1i}^{(\tau)}, \dots, s_{2di}^{(\tau)})^T$ with entries

$$s_{li}^{(\tau)} := \begin{cases} (1 - \tau)s_{li}^{(1)} + \tau s_{li}^{(2)} & \text{if } i \notin K \text{ or } l \notin K, \\ (\gamma_0(\tau))_{li} & \text{if } l, i \in K. \end{cases}$$

Thus we can construct a path ξ in $J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ to connect \mathbf{A}_1 and \mathbf{A}_2 :

$$\xi(\tau) = [A^{(\tau)} \mid B^{(\tau)}] = [(a_1^{(\tau)}, a_2^{(\tau)}, \dots, a_{2d}^{(\tau)}) \mid (b_1^{(\tau)}, b_2^{(\tau)}, \dots, b_{2d}^{(\tau)})],$$

where

$$a_i^{(\tau)} = \begin{cases} -e_i & \text{if } i \in K, \\ s_i^{(\tau)} & \text{if } i \in \{1, 2, \dots, 2d\} \setminus K, \end{cases}$$

$$b_i^{(\tau)} = \begin{cases} s_i^{(\tau)} & \text{if } i \in K, \\ e_i & \text{if } i \in \{1, 2, \dots, 2d\} \setminus K. \end{cases}$$

This finishes the proof. \square

Remark 7.1. *The only difference in the proof of Lemma 7.1 for real boundary conditions is that $R^{(j)}$ should be chosen such that $\det R^{(j)} > 0$, $j = 1, 2$. This can be easily done, since otherwise, we can replace $R^{(j)}$ by*

$$\begin{pmatrix} -1 & \\ & I_{m_0-1} \end{pmatrix} R^{(j)}.$$

Lemma 7.2. *Let $0 \leq n_0^0 \leq \sharp(K)$ and $0 \leq n_0^+ \leq \sharp(K) - n_0^0$. Then for any $\mathbf{A} \in J_{\mathcal{O}_K^{\mathbb{C}}}^{(n_0^0, n_0^+, n_0^-)}$, there exists $\varepsilon_1 > 0$ such that for any $0 < \varepsilon < \varepsilon_1$,*

$$\begin{aligned} U_\varepsilon &:= \{\mathbf{B} \in \mathcal{O}_K^{\mathbb{C}} : \|S(\mathbf{B}) - S(\mathbf{A})\|_{\mathcal{M}_{2d \times 2d}} < \varepsilon\} \\ &= \bigcup_{n^0 \leq n_0^0, n^+ \geq n_0^+, n^- \geq n_0^-} U_\varepsilon \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}, \end{aligned} \quad (7.3)$$

and $U_\varepsilon \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ is path connected for any $n^0 \leq n_0^0$, $n^+ \geq n_0^+$ and $n^- \geq n_0^-$.

Proof. Let $\mathbf{A} = [A \mid B]$ be given in (2.1). Then there exists a $m_0 \times m_0$ unitary matrix N such that

$$S_K(\mathbf{A}) = N^* \begin{pmatrix} 0_{n_0^0} & \\ & M \end{pmatrix} N, \quad M = \begin{pmatrix} \mu_1 & & & & \\ & \ddots & & & \\ & & \mu_{n_0^+} & & \\ & & & \nu_1 & \\ & & & & \ddots \\ & & & & & \nu_{n_0^-} \end{pmatrix}, \quad (7.4)$$

where $m_0 = \sharp(K)$ and $\nu_1 \leq \dots \leq \nu_{n_0^-} < 0 < \mu_1 \leq \dots \leq \mu_{n_0^+}$. (7.3) is straightforward from the small perturbation of $S_K(\mathbf{A})$ in (7.4). Fix any $\mathbf{B}_i \in U_\varepsilon \cap J_{\mathcal{O}_K^c}^{(n_0^0, n^+, n^-)}$, $i = 1, 2$. Their entries are given by a similar way as (2.1). Then the connection from $s_{lj}(\mathbf{B}_1)$ to $s_{lj}(\mathbf{B}_2)$ is trivial if $l \notin K$ or $j \notin K$. So it suffices to construct a path connecting $S_K(\mathbf{B}_1)$ and $S_K(\mathbf{B}_2)$. To do so, we only need to show that

$$\mathcal{E} := \{E : \|E - E_0\|_{\mathcal{M}_{m_0 \times m_0}} < \varepsilon_0, E = E^*, n^\pm(E) = n^\pm\}$$

is path connected for $\varepsilon_0 > 0$ sufficiently small, where $E_0 := \begin{pmatrix} 0_{n_0^0} & \\ & M \end{pmatrix}$. Let $E = \begin{pmatrix} E_{11} & E_{12}^* \\ E_{12} & M + E_{22} \end{pmatrix} \in \mathcal{E}$ and define $F_\tau = \begin{pmatrix} & 1 & & 0 \\ -\tau(M + E_{22})^{-1}E_{12} & & & 1 \end{pmatrix}$, where $E_{ii} = E_{ii}^*$, $i = 1, 2$, and $\tau \in [0, 1]$. Then $\gamma(\tau) = F_\tau^* E F_\tau$ is a path in \mathcal{E} from E to $\begin{pmatrix} F & 0 \\ 0 & M + E_{22} \end{pmatrix}$, where $F = E_{11} - E_{12}^*(M + E_{22})^{-1}E_{12}$. Similarly, for another $\tilde{E} = \begin{pmatrix} \tilde{E}_{11} & \tilde{E}_{12}^* \\ \tilde{E}_{12} & M + \tilde{E}_{22} \end{pmatrix} \in \mathcal{E}$, one can construct a path $\tilde{\gamma}$ in \mathcal{E} to connect \tilde{E} to $\begin{pmatrix} \tilde{F} & 0 \\ 0 & M + \tilde{E}_{22} \end{pmatrix}$. Now, we connect $\begin{pmatrix} F & 0 \\ 0 & M + E_{22} \end{pmatrix}$ and $\begin{pmatrix} \tilde{F} & 0 \\ 0 & M + \tilde{E}_{22} \end{pmatrix}$ in \mathcal{E} . $M + (1 - \tau)E_{22} + \tau\tilde{E}_{22}$ is a path from $M + E_{22}$ to $M + \tilde{E}_{22}$. The rest is to connect F and \tilde{F} . Note that $n^\pm(F) = n^\pm(\tilde{F}) = n^\pm - n_0^\pm$. Thus there exists a path γ_1 to connect F and \tilde{F} by a similar strategy used in (7.2) such that $n^\pm(\gamma_1(\tau)) = n^\pm - n_0^\pm$ for each $\tau \in [0, 1]$. However, $\|\gamma_1(\tau)\|$ maybe larger than ε_0 , where $\|\cdot\| := \|\cdot\|_{\mathcal{M}_{n_0^0 \times n_0^0}}$ for convenience. Thus we need to shrink the path γ_1 as follows. Assume that $\|\tilde{F}\| > \|F\|$, otherwise the construction is similar. Connect F and $\frac{\|F\|}{\|\tilde{F}\|}\tilde{F}$ by $\frac{\|F\|}{\|\gamma_1(\tau)\|}\gamma_1(\tau)$, and then connect $\frac{\|F\|}{\|\tilde{F}\|}\tilde{F}$ and \tilde{F} by $\frac{(1-\tau)\|F\| + \tau\|\tilde{F}\|}{\|\tilde{F}\|}\tilde{F}$, $\tau \in [0, 1]$. This completes the proof. \square

In the following discussion, we always assume that $0 < n_0^0 \leq \sharp(K)$, $0 \leq n_0^+ \leq \sharp(K) - n_0^0$, $n^0 < n_0^0$, $n^+ \geq n_0^+$ and $n^- \geq n_0^-$.

Set $K = \{k_1, \dots, k_{m_0}\}$ and define

$$\tilde{\mathbf{A}}_0 := [A_{(n_0^0, n_0^+)} \mid B_{(n_0^0, n_0^+)}] = [-I_{2d} \mid (b_1, b_2, \dots, b_{2d})] \in J_{\mathcal{O}_K^c}^{(n_0^0, n_0^+, n_0^-)}, \quad (7.5)$$

where

$$b_l = \begin{cases} 0 & \text{if } l \in \{k_1, \dots, k_{n_0^0}\}, \\ e_l & \text{if } l \in \{k_{n_0^0+1}, \dots, k_{n_0^0+n_0^+}\} \cup (\{1, \dots, 2d\} \setminus K), \\ -e_l & \text{if } l \in \{k_{n_0^0+n_0^++1}, \dots, k_{m_0}\}. \end{cases}$$

Consider

$$P_0 := \begin{pmatrix} p_{11} & & \\ & \ddots & \\ & & p_{dd} \end{pmatrix}, \quad (7.6)$$

Q_0 and W_0 are defined similarly as (7.6), where $p_{ii}, q_{ii}, w_{ii} \in L^\infty([a, b], \mathbb{R})$, and $p_{ii}, w_{ii} > 0$ a.e. on $[a, b]$ for each $1 \leq i \leq d$. We get the following result from 1-dimensional Sturm-Liouville problems:

Lemma 7.3. *Consider the Sturm-Liouville equation $\omega_0 = (P_0, Q_0, W_0)$ defined in (7.6). Then for $\tilde{\mathbf{A}}_0 \in J_{\mathcal{O}_K^c}^{(n_0^0, n_0^+, n_0^-)}$ defined in (7.5), there exists a path $\tilde{\mathbf{A}}_s \in J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$, $s \in (0, 1]$ such that $\tilde{\mathbf{A}}_s \rightarrow \tilde{\mathbf{A}}_0$ as $s \rightarrow 0^+$, and*

$$\lim_{s \rightarrow 0^+} \lambda_n(\tilde{\mathbf{A}}_s) = -\infty, \quad n \leq n^+ - n_0^+, \quad (7.7)$$

$$\lim_{s \rightarrow 0^+} \lambda_n(\tilde{\mathbf{A}}_s) = \lambda_{n-(n^+-n_0^+)}(\tilde{\mathbf{A}}_0), \quad n > n^+ - n_0^+. \quad (7.8)$$

Proof. $\tilde{\mathbf{A}}_s \in J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$, $s \in (0, 1]$, can be directly constructed by setting

$$\tilde{\mathbf{A}}_s = [-I_{2d} \mid (b_1(s), b_2(s), \dots, b_{2d}(s))] \in J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)},$$

where

$$b_l(s) = \begin{cases} 0 & \text{if } l \in \{k_1, \dots, k_{n^0}\}, \\ se_l & \text{if } l \in \{k_{n^0+1}, \dots, k_{n^0+n^+-n_0^+}\}, \\ -se_l & \text{if } l \in \{k_{n^0+n^+-n_0^++1}, \dots, k_{n^0}\}, \\ e_l & \text{if } l \in \{k_{n^0+1}, \dots, k_{n^0+n^+}\} \cup (\{1, \dots, 2d\} \setminus K), \\ -e_l & \text{if } l \in \{k_{n^0+n^++1}, \dots, k_{m_0}\}, \end{cases}$$

and $b_l(s) := (b_{1l}(s), \dots, b_{2dl}(s))^T$. Since the Sturm-Liouville equation $\boldsymbol{\omega}_0$ and $\tilde{\mathbf{A}}_s$ are equivalent to d one-dimensional Sturm-Liouville equations

$$-(p_{jj}y_j')' + q_{jj}y_j = \lambda w_{jj}y_j, \quad \text{on } [a, b],$$

with boundary conditions

$$y_j(a) + b_{jj}(s)(p_{jj}y_j')(a) = 0, \quad -y_j(b) + b_{d+j, d+j}(s)(p_{jj}y_j')(b) = 0,$$

where $1 \leq j \leq d$. Then (7.7) and (7.8) hold by Proposition 4.1. \square

Next, we consider other paths in $J_{\mathcal{O}_K^c}^{(n_0^0, n_0^+, n_0^-)}$, which tend to $\tilde{\mathbf{A}}_0$ given in (7.5).

Lemma 7.4. *Let $\boldsymbol{\omega}_0 = (P_0, Q_0, W_0)$ be given in (7.6). Then for $\tilde{\mathbf{A}}_0 \in J_{\mathcal{O}_K^c}^{(n_0^0, n_0^+, n_0^-)}$ defined in (7.5) and for any path $\mathbf{A}_s \in J_{\mathcal{O}_K^c}^{(n_0^0, n_0^+, n_0^-)}$, $s \in (0, 1]$ such that $\mathbf{A}_s \rightarrow \tilde{\mathbf{A}}_0$ as $s \rightarrow 0^+$, we have*

$$\lim_{s \rightarrow 0^+} \lambda_n(\mathbf{A}_s) = -\infty, \quad n \leq n^+ - n_0^+, \quad (7.9)$$

$$\lim_{s \rightarrow 0^+} \lambda_n(\mathbf{A}_s) = \lambda_{n-(n^+-n_0^+)}(\tilde{\mathbf{A}}_0), \quad n > n^+ - n_0^+. \quad (7.10)$$

Proof. If $n^+ - n_0^+ = 0$, then it suffices to show that λ_1 is bounded from below on $\{\mathbf{A}_s : s \in (0, 1]\}$ by Lemma 3.5. Suppose otherwise, there exists a sequence $\{s_n\}_{n=1}^\infty$ such that $s_n \rightarrow 0^+$ as $n \rightarrow \infty$ and

$$\lim_{n \rightarrow \infty} \lambda_1(\mathbf{A}_{s_n}) = -\infty. \quad (7.11)$$

Let $r_1 < r_2$ such that $\lambda_1(\tilde{\mathbf{A}}_0)$ with multiplicity m_0 is the only eigenvalue of $(\boldsymbol{\omega}_0, \tilde{\mathbf{A}}_0)$ in (r_1, r_2) and neither r_1 nor r_2 is an eigenvalue of $(\boldsymbol{\omega}_0, \tilde{\mathbf{A}}_0)$. By Lemma 3.2, there exists a neighborhood U_0 of $\tilde{\mathbf{A}}_0$ in \mathcal{O}_K^c such that for each $\mathbf{A} \in U_0$, $(\boldsymbol{\omega}_0, \mathbf{A})$ has exactly m_0 eigenvalues in (r_1, r_2) and neither r_1 nor r_2 is an eigenvalue of $(\boldsymbol{\omega}_0, \mathbf{A})$. Choose $n_1 \in \mathbb{N}$ such that

$$\mathbf{A}_{s_{n_1}} \in U_0 \text{ and } \lambda_1(\mathbf{A}_{s_{n_1}}) < r_1 \quad (7.12)$$

by (7.11). Lemma 7.3 tells us there exists $s' \in (0, 1]$ such that

$$\tilde{\mathbf{A}}_{s'} \in U_0, \text{ and } \lambda_1(\tilde{\mathbf{A}}_{s'}) > r_1. \quad (7.13)$$

Note that $\mathbf{A}_{s_{n_1}}, \tilde{\mathbf{A}}_{s'} \in U_0 \cap J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$, which can be chosen such that it is path connected by Lemma 7.2. Then we choose a path γ_0 in $U_0 \cap J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$ to connect $\mathbf{A}_{s_{n_1}}$ and $\tilde{\mathbf{A}}_{s'}$. By Theorem 6.1, λ_1 is continuous on γ_0 . Then there exists $\tau_0 \in (0, 1)$ such that $\lambda_1(\gamma_0(\tau_0)) = r_1$ by (7.12)–(7.13). However, r_1 is not an eigenvalue of $(\boldsymbol{\omega}_0, \mathbf{A})$ for any $\mathbf{A} \in U_0$. This is a contradiction.

Let $n^+ - n_0^+ > 0$. For any $1 \leq i \leq n^+ - n_0^+$, assume that

$$\lim_{s \rightarrow 0^+} \lambda_j(\mathbf{A}_s) = -\infty, \quad 1 \leq j \leq i - 1, \quad (7.14)$$

we show that

$$\lim_{s \rightarrow 0^+} \lambda_i(\mathbf{A}_s) = -\infty. \quad (7.15)$$

Suppose otherwise, there exists a sequence $\{s_n^{(i)}\}_{n=1}^\infty \subset (0, 1]$ such that $s_n^{(i)} \rightarrow 0^+$ as $n \rightarrow \infty$, and λ_i is bounded from below on $\{\mathbf{A}_{s_n^{(i)}}\}_{n=1}^\infty$. Then $\lim_{n \rightarrow \infty} \lambda_i(\mathbf{A}_{s_n^{(i)}}) = \lambda_1(\tilde{\mathbf{A}}_0)$ by Lemma 3.5 for $i = 1$ and by Lemma 3.6 for $i > 1$. Again from Lemma 7.3, $\lim_{s \rightarrow 0^+} \lambda_i(\tilde{\mathbf{A}}_s) = -\infty$. Thus we can choose $\tilde{\mathbf{A}}_{s'_1}, \mathbf{A}_{s_{n_1}^{(i)}} \in U_0 \cap J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$ such that $\lambda_i(\tilde{\mathbf{A}}_{s'_1}) < r_1 < \lambda_i(\mathbf{A}_{s_{n_1}^{(i)}})$ for some $s'_1 \in (0, 1]$ and some $n_1 \in \mathbb{N}$. Then the choice of r_1 contradicts that $\lambda_i(U_0 \cap J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}) := \{\lambda_i(\mathbf{A}) : \mathbf{A} \in U_0 \cap J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}\}$ is connected by Theorem 6.1 and Lemma 7.2.

To prove (7.10), it is sufficient to show that $\lambda_{n^+ - n_0^+ + 1}$ is bounded from below on $U_0 \cap J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$ by Lemma 3.6. Suppose otherwise, there exists $s'_2 \in (0, 1]$ such that $\mathbf{A}_{s'_2} \in U_0 \cap J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$ and $\lambda_{n^+ - n_0^+ + 1}(\mathbf{A}_{s'_2}) < r_1$. By Lemma 7.3, there exists $s'_3 \in (0, 1]$ such that $\tilde{\mathbf{A}}_{s'_3} \in U_0 \cap J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$ and $\lambda_{n^+ - n_0^+ + 1}(\tilde{\mathbf{A}}_{s'_3}) > r_1$. Connect $\mathbf{A}_{s'_2}$ and $\tilde{\mathbf{A}}_{s'_3}$ by a path γ_1 in $U_0 \cap J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$. Thus there exists $\tau_1 \in (0, 1)$ such that $\lambda_{n^+ - n_0^+ + 1}(\gamma_1(\tau_1)) = r_1$. However, r_1 is not an eigenvalue for any boundary condition in $U_0 \cap J_{\mathcal{O}_K^c}^{(n^0, n^+, n^-)}$, which is a contradiction. \square

From Lemma 7.4, we have shown asymptotic behavior of the n -th eigenvalue near $\tilde{\mathbf{A}}_0$ from all the directions in \mathcal{O}_K^c . Next, we consider other boundary conditions in $J_{\mathcal{O}_K^c}^{(n^0, n_0^+, n_0^-)}$.

Lemma 7.5. *Let $\omega_0 = (P_0, Q_0, W_0)$ be given in (7.6). Then for any $\mathbf{B}_0 \in J_{\mathcal{O}_K^{\mathbb{C}}}^{(n_0^0, n_0^+, n_0^-)}$ and for any path $\mathbf{B}_s \in J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$, $s \in (0, 1]$ such that $\mathbf{B}_s \rightarrow \mathbf{B}_0$ as $s \rightarrow 0^+$, we have*

$$\lim_{s \rightarrow 0^+} \lambda_n(\mathbf{B}_s) = -\infty, \quad n \leq n^+ - n_0^+, \quad (7.16)$$

$$\lim_{s \rightarrow 0^+} \lambda_n(\mathbf{B}_s) = \lambda_{n-(n^+-n_0^+)}(\mathbf{B}_0), \quad n > n^+ - n_0^+. \quad (7.17)$$

Proof. By Lemma 7.1, there exists a path $\tilde{\gamma}$ in $J_{\mathcal{O}_K^{\mathbb{C}}}^{(n_0^0, n_0^+, n_0^-)}$ to connect \mathbf{B}_0 and $\tilde{\mathbf{A}}_0$, which is given in (7.5). λ_1 is continuous on $\tilde{\gamma}$ by Theorem 6.1. Fix any $r_1 < \min \lambda_1(\tilde{\gamma})$. If $n^+ - n_0^+ = 0$, it suffices to show that λ_1 is bounded from below on $\{\mathbf{B}_s : s \in (0, 1]\}$ by Lemma 3.5. Suppose otherwise, there exists a sequence $\{s_n\}_{n=1}^{\infty}$ such that $s_n \rightarrow 0^+$ as $n \rightarrow \infty$ and

$$\lim_{n \rightarrow \infty} \lambda_1(\mathbf{B}_{s_n}) = -\infty. \quad (7.18)$$

For any given \mathbf{A}_s in Lemma 7.4,

$$\lim_{s \rightarrow 0^+} \lambda_1(\mathbf{A}_s) = \lambda_1(\tilde{\mathbf{A}}_0). \quad (7.19)$$

Choose $r_{2,\tau} > r_1$ such that $\lambda_1(\tilde{\gamma}(\tau))$ with multiplicity $m_{1,\tau}$ is the only eigenvalue in $(r_1, r_{2,\tau})$ and $r_{2,\tau}$ is not an eigenvalue of $(\omega_0, \tilde{\gamma}(\tau))$ for each $\tau \in [0, 1]$. Then by Lemma 3.2, for any given $\tau \in [0, 1]$, there exists a neighborhood U_τ of $\tilde{\gamma}(\tau)$ in $\mathcal{O}_K^{\mathbb{C}}$ such that for each $\mathbf{A} \in U_\tau$, (ω_0, \mathbf{A}) has exactly $m_{1,\tau}$ eigenvalues in $(r_1, r_{2,\tau})$ and neither r_1 nor $r_{2,\tau}$ is its eigenvalue. Set $U := \cup_{\tau \in [0,1]} U_\tau$. Then r_1 is not an eigenvalue of (ω_0, \mathbf{A}) for any $\mathbf{A} \in U$. Since $\tilde{\gamma}$ is compact, it is easy to see that $U_{\varepsilon_0} := \{\mathbf{B} \in \mathcal{O}_K^{\mathbb{C}} : \|S(\mathbf{B}) - S(\tilde{\mathbf{A}})\|_{\mathcal{M}_{2d \times 2d}} < \varepsilon_0, \tilde{\mathbf{A}} \in \tilde{\gamma}\} \subset U$ for some $\varepsilon_0 > 0$. It follows from (7.18)–(7.19) that there exist $n_1 \in \mathbb{N}$ and $\tilde{s} \in (0, 1]$ such that $\mathbf{B}_{s_{n_1}}, \mathbf{A}_{\tilde{s}} \in U_{\varepsilon_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ and

$$\lambda_1(\mathbf{B}_{s_{n_1}}) < r_1 < \lambda_1(\mathbf{A}_{\tilde{s}}). \quad (7.20)$$

We can construct a path $\tilde{\gamma}_0$ in $U_{\varepsilon_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ to connect $\mathbf{B}_{s_{n_1}}$ and $\mathbf{A}_{\tilde{s}}$ as follows. Define $s_{ij}(\tilde{\gamma}_1(\tau)) := s_{ij}(\tilde{\gamma}(\tau))$ when $i \notin K$ or $j \notin K$, $\tau \in [0, 1]$. Let $m_0 = \sharp(K)$. Denote $S_K(\tilde{\gamma}(\tau)) = M_\tau^* S_K(\tilde{\mathbf{A}}_0) M_\tau$ to be the path to connect $S_K(\mathbf{B}_0)$ and $S_K(\tilde{\mathbf{A}}_0)$ such that $M_\tau \in GL(m_0, \mathbb{C})$, $\tau \in [0, 1]$. Then define $S_K(\tilde{\gamma}_1(\tau)) := M_\tau^* S_K(\mathbf{A}_{\tilde{s}}) M_\tau$. Note that $\mathbf{A}_{\tilde{s}}$ can be chosen sufficiently close

to $\tilde{\mathbf{A}}_0$ such that $\tilde{\gamma}_1 \subset U_{\varepsilon_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$. Following Lemma 7.2, connect $\tilde{\gamma}_1(0)$ and $\mathbf{B}_{s_{n_1}}$ by $\tilde{\gamma}_2$ in $U_{\varepsilon_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$. Combining $\tilde{\gamma}_i$, $i = 1, 2$, we get the desired path $\tilde{\gamma}_0$. $\lambda_1(\tilde{\gamma}_0)$ is connected by Theorem 6.1. Thus (7.20) contradicts that r_1 is not an eigenvalue for any boundary condition in $\tilde{\gamma}_0$.

If $n^+ - n_0^+ > 0$, for any $1 \leq i \leq n^+ - n_0^+$, assume that $\lim_{s \rightarrow 0^+} \lambda_j(\mathbf{B}_s) = -\infty$, $1 \leq j \leq i - 1$, we show that

$$\lim_{s \rightarrow 0^+} \lambda_i(\mathbf{B}_s) = -\infty. \quad (7.21)$$

Suppose otherwise, there exists a sequence $\{\hat{s}_n^{(i)}\}_{n=1}^{\infty}$ such that $\hat{s}_n^{(i)} \rightarrow 0^+$ as $n \rightarrow \infty$ and λ_i is bounded from below on $\{\mathbf{B}_{\hat{s}_n^{(i)}}\}_{n=1}^{\infty}$. Then $\lim_{n \rightarrow \infty} \lambda_i(\mathbf{B}_{\hat{s}_n^{(i)}}) = \lambda_1(\mathbf{B}_0)$ by Lemmas 3.5 and 3.6. Note that $\lim_{s \rightarrow 0^+} \lambda_i(\mathbf{A}_s) = -\infty$, where \mathbf{A}_s is given in Lemma 7.4. Thus we can choose $\mathbf{B}_{\hat{s}_{n_1}^{(i)}}, \mathbf{A}_{\hat{s}'_1} \in U_{\varepsilon_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ such that

$$\lambda_i(\mathbf{A}_{\hat{s}'_1}) < r_1 < \lambda_i(\mathbf{B}_{\hat{s}_{n_1}^{(i)}}) \quad (7.22)$$

for some $n_1 \in \mathbb{N}$ and some $\hat{s}'_1 \in (0, 1]$. Similar as above, there exists a path $\tilde{\gamma}_3$ in $U_{\varepsilon_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ to connect $\mathbf{B}_{\hat{s}_{n_1}^{(i)}}$ and $\mathbf{A}_{\hat{s}'_1}$. $\lambda_i(\tilde{\gamma}_3)$ is connected by Theorem 6.1. Then the choice of r_1 contradicts (7.22).

Suppose (7.17) is not true. Then we can find $\mathbf{B}_{\hat{s}_2}, \mathbf{A}_{\hat{s}_2} \in U_{\varepsilon_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ such that $\lambda_{n^+ - n_0^+ + 1}(\mathbf{A}_{\hat{s}_2}) > r_1 > \lambda_{n^+ - n_0^+ + 1}(\mathbf{B}_{\hat{s}_2})$. By constructing $\tilde{\gamma}_4$ in $U_{\varepsilon_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ to connect $\mathbf{B}_{\hat{s}_2}$ and $\mathbf{A}_{\hat{s}_2}$, one can get a contradiction as above. This completes the proof. \square

Now, for any $\boldsymbol{\omega} = (P, Q, W) \in \Omega$, we give asymptotic behavior of the n -th eigenvalue near any boundary condition in $J_{\mathcal{O}_K^{\mathbb{C}}}^{(n_0^0, n_0^+, n_0^-)}$, $n_0^0 \geq 1$, from all the directions in $\mathcal{O}_K^{\mathbb{C}}$.

Lemma 7.6. *Let $\boldsymbol{\omega} = (P, Q, W) \in \Omega$. Then for any $\mathbf{C}_0 \in J_{\mathcal{O}_K^{\mathbb{C}}}^{(n_0^0, n_0^+, n_0^-)}$ and for any path $\mathbf{C}_s \in J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$, $s \in (0, 1]$ such that $\mathbf{C}_s \rightarrow \mathbf{C}_0$ as $s \rightarrow 0^+$, we have*

$$\lim_{s \rightarrow 0^+} \lambda_n(\mathbf{C}_s) = -\infty, \quad n \leq n^+ - n_0^+, \quad (7.23)$$

$$\lim_{s \rightarrow 0^+} \lambda_n(\mathbf{C}_s) = \lambda_{n - (n^+ - n_0^+)}(\mathbf{C}_0), \quad n > n^+ - n_0^+. \quad (7.24)$$

Proof. Let

$$\zeta(\tau) := (\tau P_0 + (1 - \tau)P, \tau Q_0 + (1 - \tau)Q, \tau W_0 + (1 - \tau)W), \quad \tau \in [0, 1],$$

where (P_0, Q_0, W_0) is given in (7.6). Then it is easy to verify that $\zeta(\tau) \in \Omega, \tau \in [0, 1]$. $\zeta \times \mathbf{C}_0$ means $\{(\zeta(\tau), \mathbf{C}_0) : \tau \in [0, 1]\}$ and choose $\hat{r}_1 < \min\{\lambda_1(\zeta \times \mathbf{C}_0)\}$.

Let $\hat{r}_{2,\tau} > \hat{r}_1$ such that $\lambda_1(\zeta(\tau), \mathbf{C}_0)$ with multiplicity $\hat{m}_{1,\tau}$ is the only eigenvalue in $(\hat{r}_1, \hat{r}_{2,\tau})$ and $\hat{r}_{2,\tau}$ is not an eigenvalue of $(\zeta(\tau), \mathbf{C}_0)$ for each $\tau \in [0, 1]$. Then by Lemma 3.2, there exists a neighborhood \hat{U}_τ of $(\zeta(\tau), \mathbf{C}_0)$, $\tau \in [0, 1]$, in $\Omega \times \mathcal{O}_K^{\mathbb{C}}$ such that each $(\boldsymbol{\sigma}, \mathbf{C}) \in \hat{U}_\tau$ has exactly $\hat{m}_{1,\tau}$ eigenvalues in $(\hat{r}_1, \hat{r}_{2,\tau})$ and neither \hat{r}_1 nor $\hat{r}_{2,\tau}$ is its eigenvalue. Set $\hat{U} := \cup_{\tau \in [0,1]} \hat{U}_\tau$. Then \hat{r}_1 is not an eigenvalue of $(\boldsymbol{\sigma}, \mathbf{C}) \in \hat{U}$. By the compactness of ζ , there exists a neighborhood $U_{\mathbf{C}_0}$ of \mathbf{C}_0 in $\mathcal{O}_K^{\mathbb{C}}$ such that $(\zeta(\tau), U_{\mathbf{C}_0}) \subset \hat{U}$ for all $\tau \in [0, 1]$.

If $n^+ - n_0^+ = 0$, then by Lemma 7.5, for any $\mathbf{B}_s, s \in (0, 1]$, such that $\mathbf{B}_s \rightarrow \mathbf{C}_0$ as $s \rightarrow 0^+$, λ_1 is continuous on $\{(\boldsymbol{\omega}_0, \mathbf{B}_s) : s \in (0, 1]\} \cup \{(\boldsymbol{\omega}_0, \mathbf{C}_0)\}$. Thus there exists $s_0 \in (0, 1]$ such that $\mathbf{B}_{s_0} \in U_{\mathbf{C}_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ and $\lambda_1(\boldsymbol{\omega}_0, \mathbf{B}_{s_0}) > \hat{r}_1$. Suppose there exists $s_1 \in (0, 1]$ such that $\mathbf{C}_{s_1} \in U_{\mathbf{C}_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ and $\lambda_1(\boldsymbol{\omega}, \mathbf{C}_{s_1}) < \hat{r}_1$. In $(\Omega \times J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}) \cap \hat{U}$, via constructing a path to connect $(\boldsymbol{\omega}_0, \mathbf{B}_{s_0})$ and $(\boldsymbol{\omega}_0, \mathbf{C}_{s_1})$ by Lemma 7.2 then a path to connect $(\boldsymbol{\omega}_0, \mathbf{C}_{s_1})$ and $(\boldsymbol{\omega}, \mathbf{C}_{s_1})$ through ζ , we get a path $\hat{\gamma}$ to connect $(\boldsymbol{\omega}_0, \mathbf{B}_{s_0})$ and $(\boldsymbol{\omega}, \mathbf{C}_{s_1})$. $\lambda_1(\hat{\gamma})$ is connected by Theorem 6.1. This contradicts $\lambda_1(\boldsymbol{\omega}, \mathbf{C}_{s_1}) < \hat{r}_1 < \lambda_1(\boldsymbol{\omega}_0, \mathbf{B}_{s_0})$ and \hat{r}_1 is not an eigenvalue of $(\boldsymbol{\sigma}, \mathbf{C}) \in \hat{\gamma} \subset \hat{U}$.

Let $n^+ - n_0^+ > 0$. Then for any $1 \leq i \leq n^+ - n_0^+$, assuming that we have shown that $\lim_{s \rightarrow 0^+} \lambda_j(\boldsymbol{\omega}, \mathbf{C}_s) = -\infty, 1 \leq j \leq i - 1$, we now show that $\lim_{s \rightarrow 0^+} \lambda_i(\boldsymbol{\omega}, \mathbf{C}_s) = -\infty$. If it is not true, there exists $\{s_n\}_{n=1}^\infty$ such that λ_i is bounded from below on $\{(\boldsymbol{\omega}, \mathbf{C}_{s_n})\}_{n=1}^\infty$ and thus $\lim_{n \rightarrow \infty} \lambda_i(\boldsymbol{\omega}, \mathbf{C}_{s_n}) = \lambda_1(\boldsymbol{\omega}, \mathbf{C}_0)$ by Lemmas 3.5 and 3.6. This, together with Lemma 7.5, implies that there exists $s_2 \in (0, 1]$ and $n_1 \in \mathbb{N}$ such that $\mathbf{B}_{s_2}, \mathbf{C}_{s_{n_1}} \in U_{\mathbf{C}_0} \cap J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$ and

$$\lambda_i(\boldsymbol{\omega}_0, \mathbf{B}_{s_2}) < \hat{r}_1 < \lambda_i(\boldsymbol{\omega}, \mathbf{C}_{s_{n_1}}), \quad (7.25)$$

where $\mathbf{B}_s \in J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}, s \in (0, 1]$ is any path such that $\mathbf{B}_s \rightarrow \mathbf{C}_0$ as $s \rightarrow 0^+$. We can construct a path $\hat{\gamma}_1$ in $(\Omega \times J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}) \cap \hat{U}$ to connect $(\boldsymbol{\omega}_0, \mathbf{B}_{s_2})$ and $(\boldsymbol{\omega}, \mathbf{C}_{s_{n_1}})$ as above. $\lambda_i(\hat{\gamma}_1)$ is connected by Theorem 6.1. Thus (7.25)

contradicts that \hat{r}_1 is not an eigenvalue for any $(\sigma, \mathbf{C}) \in \hat{\gamma}_1$. This completes the proof of (7.23).

(7.24) can be shown similarly as above and the proof is complete. \square

To conclude this section, we combine the Sturm-Liouville equations and boundary conditions to get the following asymptotic behavior of the n -th eigenvalue:

Theorem 7.2. *For any $(\omega, \mathbf{A}) \in \Omega \times J_{\mathcal{O}_K^{\mathbb{C}}}^{(n_0^0, n_0^+, n_0^-)}$ and for any path $(\omega_s, \mathbf{A}_s) \in \Omega \times J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}$, $s \in (0, 1]$ such that $(\omega_s, \mathbf{A}_s) \rightarrow (\omega, \mathbf{A})$ as $s \rightarrow 0^+$, we have*

$$\lim_{s \rightarrow 0^+} \lambda_n(\omega_s, \mathbf{A}_s) = -\infty, \quad n \leq n^+ - n_0^+, \quad (7.26)$$

$$\lim_{s \rightarrow 0^+} \lambda_n(\omega_s, \mathbf{A}_s) = \lambda_{n-(n^+-n_0^+)}(\omega, \mathbf{A}), \quad n > n^+ - n_0^+. \quad (7.27)$$

Proof. The proof is also by a contradiction argument as that used in the proof of Lemmas 7.4, 7.5 and 7.6. So we just sketch the proof here. It follows from Lemma 7.6 that for the path (ω_s, \mathbf{A}_s) , we have the asymptotic behavior (7.26)–(7.27) with ω_s replaced by ω . Thus, to get the contradiction, it suffices to construct a path in $(\Omega \times J_{\mathcal{O}_K^{\mathbb{C}}}^{(n^0, n^+, n^-)}) \cap U$ to connect $(\omega, \mathbf{A}_{s_1})$ and $(\omega_{s_2}, \mathbf{A}_{s_2})$ for some $s_1, s_2 \in (0, 1]$, where U is a sufficiently small neighborhood of (ω, \mathbf{A}) in $\Omega \times \mathcal{O}_K^{\mathbb{C}}$. This can be obtained due to Lemma 7.2. \square

8 Derivative formulas and comparison of eigenvalues

In this section, we give derivative formula of a continuous simple eigenvalue branch with respect to boundary conditions. Then we obtain some inequalities of eigenvalues.

Theorem 8.1. *Fix $\omega \in \Omega$. Let $\lambda_*(\mathbf{A})$ be a simple eigenvalue for \mathbf{A} , y be a normalized eigenfunction for $\lambda_*(\mathbf{A})$, and Λ be a continuous simple eigenvalue branch defined on a neighborhood of \mathbf{A} in $\mathcal{O}_K^{\mathbb{C}}$ through λ_* for some $K \subset \{1, \dots, 2d\}$. Then Λ is differentiable at \mathbf{A} and its Frechet derivative formula is given by*

$$d\Lambda|_{\mathbf{A}}(H) = u^* H u \quad (8.1)$$

for any $2d \times 2d$ Hermitian matrix H , where $u = (u_1, \dots, u_{2d})^T$ is given by

$$u_i = \begin{cases} \rho_i, & i \in \{1, \dots, 2d\} \setminus K, \\ \eta_i, & i \in K, \end{cases}$$

and $(\rho_1, \dots, \rho_{2d}, \eta_1, \dots, \eta_{2d}) := (-y(a)^T, y(b)^T, (Py'(a))^T, (Py'(b))^T)$.

Proof. Let $\mathbf{A} := [A(S) \mid B(S)]$ and $\mathbf{B} := [A(S+H) \mid B(S+H)] \in \mathcal{O}_K^{\mathbb{C}}$, which correspond to Hermitian matrices S and $S+H$, respectively. Then there exists an eigenfunction $\tilde{y} = y_{\Lambda(\mathbf{B})}$ for $\Lambda(\mathbf{B})$ such that $P\tilde{y}' \rightarrow Py'$ as $\mathbf{B} \rightarrow \mathbf{A}$ in the sense of Lemma 3.4.

Note that \tilde{y} and y satisfy

$$-(P\tilde{y}')' + Q\tilde{y} = \lambda(\mathbf{B})W\tilde{y}, \quad -(Py')' + Qy = \lambda(\mathbf{A})Wy.$$

Thus

$$(\lambda(\mathbf{B}) - \lambda(\mathbf{A})) \int_a^b (W\tilde{y}, y)_d dt = \left(\begin{pmatrix} 0 & -I_{2d} \\ I_{2d} & 0 \end{pmatrix} \tilde{Y}(a, b), Y(a, b) \right)_{4d}.$$

Denote

$$Y := Y(a, b), \quad X := \begin{pmatrix} u \\ v \end{pmatrix},$$

where $v := (v_1, \dots, v_{2d})^T$ is given by

$$v_i = \begin{cases} \eta_i, & i \in \{1, \dots, 2d\} \setminus K, \\ -\rho_i, & i \in K. \end{cases}$$

$\tilde{Y}, \tilde{X}, \tilde{u}, \tilde{v}$ can be defined similarly. Then for \mathbf{A} and \mathbf{B} , we have

$$\begin{aligned} [A(S) \mid B(S)]Y &= [S \mid I_{2d}]X \\ &= [S \mid I_{2d}] \begin{pmatrix} u \\ v \end{pmatrix} \\ &= Su + v = 0, \\ [A(S+H) \mid B(S+H)]\tilde{Y} &= [S+H \mid I_{2d}]\tilde{X} \\ &= [S+H \mid I_{2d}] \begin{pmatrix} \tilde{u} \\ \tilde{v} \end{pmatrix} \\ &= (S+H)\tilde{u} + \tilde{v} = 0. \end{aligned}$$

It follows that

$$\begin{aligned}
(\lambda(\mathbf{B}) - \lambda(\mathbf{A})) \int_a^b (W\tilde{y}, y)_d dt &= Y^* \begin{pmatrix} 0 & -I_{2d} \\ I_{2d} & 0 \end{pmatrix} \tilde{Y} \\
&= \sum_{i=1}^{2d} (\bar{\eta}_i \tilde{\rho}_i - \bar{\rho}_i \tilde{\eta}_i) \\
&= \sum_{i \in K} (-\bar{u}_i \tilde{v}_i + \bar{v}_i \tilde{u}_i) + \sum_{i \notin K} (\bar{v}_i \tilde{u}_i - \bar{u}_i \tilde{v}_i) \\
&= \sum_{i=1}^{2d} (\bar{v}_i \tilde{u}_i - \bar{u}_i \tilde{v}_i) \\
&= X^* \begin{pmatrix} 0 & -I_{2d} \\ I_{2d} & 0 \end{pmatrix} \tilde{X} \\
&= (u^*, v^*) \begin{pmatrix} 0 & -I_{2d} \\ I_{2d} & 0 \end{pmatrix} \begin{pmatrix} \tilde{u} \\ \tilde{v} \end{pmatrix} \\
&= (u^*, -u^* S) \begin{pmatrix} 0 & -I_{2d} \\ I_{2d} & 0 \end{pmatrix} \begin{pmatrix} \tilde{u} \\ -(S + H)\tilde{u} \end{pmatrix} \\
&= u^* H \tilde{u}.
\end{aligned}$$

Letting $H \rightarrow 0$, we get (8.1). \square

The positive direction of continuous eigenvalue branches is given in the following result, which is a direct consequence of Theorems 5.1 and 8.1.

Corollary 8.1. *Fix $\omega \in \Omega$. Let $\lambda_*(\mathbf{A})$ be an eigenvalue for \mathbf{A} and Λ be a continuous eigenvalue branch defined on a neighborhood U of \mathbf{A} in $\mathcal{O}_K^{\mathbb{C}}$ through λ_* . Then*

$$\Lambda(\mathbf{A}) \leq \Lambda(\mathbf{B})$$

if $\mathbf{B} \in U$ and $S(\mathbf{B}) - S(\mathbf{A})$ is non-negative.

Remark 8.1. *Theorem 8.1 (it is required that $\lambda_n(\mathbf{A})$ is simple) and Corollary 8.1 also hold true when λ_n is continuous on a neighborhood U of \mathbf{A} in $\mathcal{O}_K^{\mathbb{C}}$.*

Acknowledgements

H. Zhu sincerely thanks Prof. Y. Long for his instructive advice. X. Hu is supported in part by NSFC (No. 11425105, 11790271). H. Zhu is supported in part by CPSF (No. 2018M630266).

References

- [1] V. I. Arnold, Characteristic class entering in quantization conditions, *Funct. Anal. Appl.* 1 (1967) 1–13.
- [2] V. I. Arnold, The complex Lagrangian Grassmannian, (Russian) *Funktsional. Anal. i Prilozhen.* 34 (2000) 63–65; translation in *Funct. Anal. Appl.* 34 (2000) 208–210.
- [3] X. Cao, Q. Kong, H. Wu, A. Zettl, Geometric aspects of Sturm-Liouville problems. III. Level surfaces of the n -th eigenvalue, *J. Comput. Appl. Math.* 208 (2007) 176–193.
- [4] J. Convey, *Functions of One Complex Variable*, 2nd ed., Springer-Verlag, Berlin/New York, 1986.
- [5] R. Courant, D. Hilbert, *Methods of Mathematical Physics*, Interscience Publishers, New York, 1953.
- [6] M. S. P. Eastham, Q. Kong, H. Wu, A. Zettl, Inequalities among eigenvalues of Sturm-Liouville problems, *J. Inequalities Appl.* 3 (1999) 25–43.
- [7] W. N. Everitt, M. Möller, A. Zettl, Discontinuous dependence of the n -th Sturm-Liouville eigenvalues, in “General Inequalities”, Birkhauser, Basel, 1997.
- [8] S. Fu, Z. Wang, The relationships among multiplicities of a J -self-adjoint differential operator’s eigenvalue, *Pac. J. Appl. Math.* 4 (2012) 292–303.
- [9] S. Fu, Z. Wang, Relationships among three multiplicities of a differential operator’s eigenvalue, *Applied Math.* 5 (2014) 2185–2194.
- [10] R. A. Horn, C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, Cambridge, 2013.
- [11] X. Hu, P. Wang, Conditional Fredholm determinant for the S -periodic orbits in Hamiltonian systems, *J. Funct. Anal.* 261 (2011) 3247–3278.
- [12] X. Hu, P. Wang, Eigenvalue problem of Sturm-Liouville systems with separated boundary conditions, *Math. Z.* 283 (2016) 339–348.

- [13] T. Kato, *Perturbation Theory for Linear Operators*, 2nd ed., Springer-Verlag, Berlin/Heidelberg/New York/Tokyo, 1984.
- [14] Q. Kong, H. Wu, A. Zettl, Dependence of the n -th Sturm-Liouville eigenvalue on the problem, *J. Differential Equations* 156 (1999) 328–354.
- [15] Q. Kong, H. Wu, A. Zettl, Geometric aspects of Sturm-Liouville problems, I. Structures on spaces of boundary conditions, *Proc. Roy. Soc. Edinburgh Sect. A* 130 (2000) 561–589.
- [16] Q. Kong, H. Wu, A. Zettl, Multiplicity of Sturm-Liouville eigenvalues, *J. Comput. Appl. Math.* 171 (2004) 291–309.
- [17] Q. Kong, A. Zettl, Eigenvalues of regular Sturm-Liouville problems, *J. Differential Equations* 131 (1996) 1–19.
- [18] Q. Kong, A. Zettl, Dependence of eigenvalues of Sturm-Liouville problems on the boundary, *J. Differential Equations* 126 (1996) 389–407.
- [19] L. Liu, Lagrangian Grassmanian Manifold $\Lambda(2)$, *Front. Math. China* (2018), doi: 10.1007/s11464-018-0683-2.
- [20] Y. Long, The structure of the singular symplectic matrix set, *Sci. China Ser. A* 34 (1991) 897–907.
- [21] Y. Long, *Index Theory for Symplectic Paths with Applications*, *Progr. Math.*, vol. 207, Birkhäuser, Basel, 2002.
- [22] M. A. Naimark, *Linear Differential Operators*, Ungar, New York, 1968.
- [23] F. Rellich, Störungstheorie der Spektralzerlegung V, *Math. Ann.* 118 (1942) 462–484.
- [24] F. Rellich, Störungstheorie der Spektralzerlegung, *Proceedings of the International Congress of Mathematicians* 1 (1950) 606–613.
- [25] D. Shi, Z. Huang, Relationships of multiplicities of eigenvalues of a higher-order ordinary differential operator, *Acta Math. Sinica (Chin. Ser.)* 53 (2010) 763–772.

- [26] A. Wang, J. Sun, A. Zettl, The classification of self-djoint boundary conditions: separated, coupled, and mixed, *J. Funct. Anal.* 255 (2008) 1554–1573.
- [27] Z. Wang, H. Wu, Equalities of multiplicities of a Sturm-Liouville eigenvalue, *J. Math. Anal. Appl.* 306 (2005) 540–547.
- [28] J. Weidmann, *Spectral Theory of Ordinary Differential Operators*, Lecture Notes in Mathematics, vol. 1258, Springer-Verlag, Berlin, 1987.
- [29] A. Zettl, Sturm-Liouville problems, In “Spectral Theory and Computational Methods of Sturm-Liouville Problems” (ed. Hinton and P. Schaefer) (New York: Dekker, 1997).
- [30] A. Zettl, *Sturm-Liouville Theory*, Mathematical Surveys Monographs, vol. 121, Amer. Math. Soc., 2005.
- [31] H. Zhu, Y. Shi, Dependence of eigenvalues on the boundary conditions of Sturm-Liouville problems with one singular endpoint, *J. Differential Equations* 263 (2017) 5582–5609.

Email addresses:

xjhu@sdu.edu.cn (X. Hu),
 201511242@mail.sdu.edu.cn (L. Liu),
 201790000005@sdu.edu.cn (L. Wu),
 haozhu@nankai.edu.cn (H. Zhu).