The generalized numerical range of a set of matrices

Pan-Shun Lau, Chi-Kwong Li, Yiu-Tung Poon, and Nung-Sing Sze

Abstract

For a given set of $n \times n$ matrices \mathcal{F} , we study the union of the C-numerical ranges of the matrices in the set \mathcal{F} , denoted by $W_C(\mathcal{F})$. We obtain basic algebraic and topological properties of $W_C(\mathcal{F})$, and show that there are connections between the geometric properties of $W_C(\mathcal{F})$ and the algebraic properties of C and the matrices in \mathcal{F} . Furthermore, we consider the starshapedness and convexity of the set $W_C(\mathcal{F})$. In particular, we show that if \mathcal{F} is the convex hull of two matrices such that $W_C(A)$ and $W_C(B)$ are convex, then the set $W_C(\mathcal{F})$ is star-shaped. We also investigate the extensions of the results to the joint C-numerical range of an m-tuple of matrices.

AMS classification. 15A60.

Keywords. Numerical range, convex set, star-shaped set.

1 Introduction

Let M_n be the set of all $n \times n$ complex matrices. The numerical range of $A \in M_n$ is defined by

$$W(A) = \{x^*Ax : x \in \mathbb{C}^n, x^*x = 1\},\$$

which is a useful tool for studying matrices and operators; for example, see [8, Chapter 22] and [9, Chapter 1]. In particular, there is an interesting interplay between the geometrical properties of W(A) and the algebraic properties of A.

In this paper, for a nonempty set \mathcal{F} of matrices in M_n , we consider

$$W(\mathcal{F}) = \bigcup \{W(A) : A \in \mathcal{F}\}.$$

We show that there are also interesting connections between the geometrical properties of $W(\mathcal{F})$ and the properties of the matrices in \mathcal{F} .

The study has motivations from different topics. We mention two of them in the following. The first one arises in the study of Crouzeix's conjecture asserting that for any $A \in M_n$,

$$||f(A)|| \le 2 \max\{|f(\mu)| : \mu \in W(A)\}$$

for any complex polynomial f(z), where ||A|| denotes the operator norm of A, see [7]. Instead of focusing on a single matrix $A \in M_n$, one may consider a complex convex set K and show that

$$\|f(A)\| \leq 2\max\{|f(\mu)|: \mu \in K\}$$

whenever $W(A) \subseteq K$. One readily shows that this is equivalent to the Crouzeix's conjecture. In fact, one may focus on the case when K is a convex polygon (including interior) because W(A) can always be approximated by convex polygons from inside or outside.

Another motivation comes from quantum information science. In quantum mechanics, a pure state in M_n is a rank one orthogonal projection of the form xx^* for some unit vector $x \in \mathbb{C}^n$, and a general state is a density matrix, which is a convex combination of pure states. For a measurement operator $A \in M_n$, which is usually Hermitian, the measurement of a state P is computed by $\langle A, P \rangle = \operatorname{tr}(AP)$. As a result,

$$W(A) = \{x^*Ax : x \in \mathbb{C}^n, x^*x = 1\} = \{\text{tr}(Axx^*) : x \in \mathbb{C}^n, x^*x = 1\}$$

can be viewed as the set of all possible measurements on pure states for a given measurement operator A. By the convexity of the numerical range, we have

$$W(A) = \{\operatorname{tr}(Axx^*) : x \in \mathbb{C}^n, x^*x = 1\} = \{\operatorname{tr}(AP) : P \text{ is a general state}\}.$$

So, W(A) actually contains all possible measurements on general states for a given measurement operator A. If we consider all possible measurements under a set \mathcal{F} of measurement operators, then it is natural to study $W(\mathcal{F})$. In fact, if we know the set $W(\mathcal{F})$, we may deduce some properties about the measurement operators in \mathcal{F} . For example, one can show that

- (a) $W(\mathcal{F}) = \{\mu\}$ if and only if $\mathcal{F} = \{\mu I\}$.
- (b) $W(\mathcal{F}) \subseteq \mathbb{R}$ if and only if all matrices in \mathcal{F} are Hermitian.
- (c) $W(\mathcal{F}) \subseteq [0, \infty)$ if and only if all matrices in \mathcal{F} are positive semi-definite.

It is worth pointing out that if $A \in M_n$ is not Hermitian, one may consider the Hermitian decomposition $A = A_1 + iA_2$ such that A_1, A_2 are Hermitian, and identify W(A) as the joint numerical range of (A_1, A_2) defined by

$$W(A_1, A_2) = \{(x^*A_1x, x^*A_2x) : x \in \mathbb{C}^n, x^*x = 1\} \subseteq \mathbb{R}^2.$$

One can study the joint numerical range of k-tuple of Hermitian matrices (A_1, \ldots, A_k) defined by

$$W(A_1, ..., A_k) = \{(x^*A_1x, ..., x^*A_kx) : x \in \mathbb{C}^n, x^*x = 1\} \subseteq \mathbb{R}^k.$$

Accordingly, one may consider the joint numerical range $W(\mathcal{F}) \subseteq \mathbb{R}^k$ of a set \mathcal{F} of k-tuple of Hermitian matrices in M_n .

It turns out that we can study $W(\mathcal{F})$ under a more general setting. For a matrix $C \in M_n$, the C-numerical range of $A \in M_n$ is defined by

$$W_C(A) = \{ \operatorname{tr} (CU^*AU) : U \text{ is unitary } \},$$

which has been studied by many researchers in connection to different topics; see [10] and its references. In particular, the C-numerical range is useful in the study of quantum control

and quantum information; for example, see [16]. Denote by $\{E_{11}, E_{12}, \ldots, E_{nn}\}$ the standard basis for M_n . When $C = E_{11}$, $W_C(A)$ reduces to W(A); if C is a rank m orthogonal projection, then $W_C(A)$ reduces to the mth-numerical range of A; see [8, 9, 12]. For a non-empty set \mathcal{F} of matrices in M_n , we consider

$$W_C(\mathcal{F}) = \bigcup \{W_C(A) : A \in \mathcal{F}\}.$$

In Section 2, we study the connection between the geometric properties of $W_C(\mathcal{F})$ and the properties of the set \mathcal{F} . In Sections 3 and 4, we study conditions for $W_C(\mathcal{F})$ to be star-shaped or convex. In Section 5, we consider the joint C-numerical range of $(A_1, \ldots, A_k) \in M_n^k$ defined by

$$W_C(A_1,\ldots,A_k) = \{(\operatorname{tr}(CU^*A_1U),\ldots,\operatorname{tr}(CU^*A_kU)): U \text{ is unitary}\} \subseteq \mathbb{C}^k.$$

If C, A_1, \ldots, A_k are Hermitian matrices, then $W_C(A_1, \ldots, A_k) \subseteq \mathbb{R}^k$. One may see [10] for the background and references on the C-numerical range.

If $\mu \in \mathbb{C}$ and $C = \mu I$, then

$$W_C(A) = \{ \mu \operatorname{tr} A \}$$
 and $W_C(A_1, \dots, A_k) = \{ \mu (\operatorname{tr} A_1, \dots, \operatorname{tr} A_k) \}.$

We will always assume that C is not a scalar matrix to avoid trivial consideration. For convenience of discussion, we always identify \mathbb{C} with \mathbb{R}^2 .

2 Basic properties of $W_C(A)$ and $W_C(\mathcal{F})$

We first list some basic results in the first two propositions below about the C-numerical range; see [4, 10, 13, 17, 18] and their references.

Proposition 2.1. Let $C, A \in M_n$.

- (a) For any unitary $U, V \in M_n$, $W_{V^*CV}(U^*AU) = W_C(A) = W_A(C)$.
- (b) For any $\alpha, \beta \in \mathbb{C}$,

$$W_C(\alpha A + \beta I) = \alpha W_C(A) + \beta \operatorname{tr} C = \{\alpha \mu + \beta \operatorname{tr} C : \mu \in W_C(A)\}.$$

- (c) The set $W_C(A)$ is compact.
- (d) The set $W_C(A)$ is star-shaped with $(\operatorname{tr} C)(\operatorname{tr} A)/n$ as a star center, i.e.,

$$t\mu + (1-t)(\operatorname{tr} C)(\operatorname{tr} A)/n \in W_C(A)$$
 for any $\mu \in W_C(A)$ and $t \in [0,1]$.

- (e) The set $W_C(A)$ is convex if there is $\gamma \in \mathbb{C}$ such that $\tilde{C} = C \gamma I_n$ satisfies any one of the following conditions.
 - (e.1) \tilde{C} is rank one.

- (e.2) \tilde{C} is a multiple of a Hermitian matrix.
- (e.3) \tilde{C} is unitarily similar to a block matrix of the form $(C_{ij})_{1 \leq i,j \leq m}$ such that C_{11}, \ldots, C_{mm} are square matrices possibly with different sizes, and $C_{ij} = 0$ if $j \neq i + 1$.

Researchers have extended the results on classical numerical range, and showed that there is interesting interplay between the geometry of $W_C(A)$ and the algebraic properties of C and A. A boundary point ν of $W_C(A)$ is a corner point if there is d > 0 such that $W_C(A) \cap \{\mu \in \mathbb{C} : |\mu - \nu| \leq d\}$ is contained in a pointed cone with ν as a vertex.

Proposition 2.2. Let $C = (c_{ij}) \in M_n$ be a non-scalar matrix in lower triangular form with diagonal entries c_1, \ldots, c_n . Suppose $A = (a_{ij}) \in M_n$ is also in lower triangular form with diagonal entries a_1, \ldots, a_n .

- (a) If $\operatorname{tr}(CA) = \sum_{j=1}^{n} c_j a_j$ is a boundary point of $W_C(A)$, then $a_{rs} = 0$ whenever $c_r \neq c_s$ and $c_{pq} = 0$ whenever $a_p \neq a_q$. In particular, if C and A has distinct eigenvalues, then C and A will be in diagonal form, i.e., C and A are normal.
- (b) If ν is a corner point of $W_C(A)$, then there is a unitary V such that V^*AV is lower triangular form with diagonal entries a_{j_1}, \ldots, a_{j_n} such that (j_1, \ldots, j_n) is a permutation of $(1, \ldots, n)$ and

$$\nu = \operatorname{tr}(CV^*AV) = \sum_{\ell=1}^n c_{\ell} a_{j_{\ell}}.$$

- (c) The set $W_C(A)$ is a singleton if and only if A is a scalar matrix.
- (d) The set $W_C(A)$ is a non-degenerate line segment if and only if C and A are non-scalar normal matrices having collinear eigenvalues in \mathbb{C} .
- (e) The set $W_C(A)$ is a convex polygon if and only if

$$W_C(A) = \operatorname{conv}\{(c_1, \dots, c_n)P(a_1, \dots a_n)^t : P \text{ is a permutation matrix}\}.$$

Now, we extend the basic properties of $W_C(A)$ to $W_C(\mathcal{F})$. We always assume that C is not a scalar matrix and \mathcal{F} is a non-empty subset of M_n .

Theorem 2.3. Suppose $C \in M_n$ is non-scalar and $\mathcal{F} \subseteq M_n$ is non-empty.

- (a) If $U \in M_n$ is unitary, then $W_C(\mathcal{F}) = W_C(U^*\mathcal{F}U)$, where $U^*\mathcal{F}U = \{U^*AU : A \in \mathcal{F}\}$.
- (b) For any $\alpha, \beta \in \mathbb{C}$, let $\alpha \mathcal{F} + \beta I = {\alpha A + \beta I : A \in \mathcal{F}}$. Then

$$W_C(\alpha \mathcal{F} + \beta I) = \alpha W_C(\mathcal{F}) + \beta \operatorname{tr} C = \{\alpha \mu + \beta \operatorname{tr} C : \mu \in W_C(\mathcal{F})\}.$$

- (c) If \mathcal{F} is bounded, then so is $W_C(\mathcal{F})$.
- (d) If \mathcal{F} is connected, then so is $W_C(\mathcal{F})$.
- (e) If \mathcal{F} is compact, then so is $W_C(\mathcal{F})$.

Proof. (a) and (b) can be verified readily.

(c) If \mathcal{F} is bounded so that there is R > 0 such that for every $A \in M_n$ we have ||A|| < R, then

$$|\operatorname{tr}(CU^*AU)| \le n||C||||A|| < n||C||R.$$

Thus, $W_C(\mathcal{F})$ is bounded.

- (d) Note that for any $A \in M_n$, $W_C(A)$ is star-shaped with $(\operatorname{tr} C)(\operatorname{tr} A)/n$ as a star center. If $\mu_1 = \operatorname{tr} (CU^*A_1U)$ and $\mu_2 = \operatorname{tr} (CV^*A_2V)$ with $A_1, A_2 \in \mathcal{F}$ and U, V unitary, then there are a line segment with end points μ_1 and $(\operatorname{tr} C)(\operatorname{tr} A_1)/n$, and another line segment with end points μ_2 and $(\operatorname{tr} C)(\operatorname{tr} A_2)/n$. If \mathcal{F} is connected, then so are the sets $\{\operatorname{tr} A : A \in \mathcal{F}\}$ and $\{(\operatorname{tr} A)(\operatorname{tr} C)/n : A \in \mathcal{F}\}$. Hence, there is a path joining μ_1 to $(\operatorname{tr} C)(\operatorname{tr} A_1)/n$, then to $(\operatorname{tr} C)(\operatorname{tr} A_2)/n$ and then to μ_2 .
- (e) Suppose \mathcal{F} is compact. Then \mathcal{F} is bounded and closed. By (c), $W_C(\mathcal{F})$ is also bounded. To show that $W_C(\mathcal{F})$ is closed, let $\{\operatorname{tr}(CU_k^*A_kU_k): k=1,2,\ldots\}$ be a sequence in $W_C(\mathcal{F})$ converging to $\mu_0 \in \mathbb{C}$, where $A_k \in \mathcal{F}$ and U_k is unitary for each k. Since \mathcal{F} is compact, there is a subsequence $\{A_{j_k}: k=1,2,\ldots\}$ of $\{A_k: k=1,2,\ldots\}$ converging to $A_0 \in \mathcal{F}$. We can further consider a subsequence $\{U_{j(\ell)}: \ell=1,2,\ldots\}$ of $\{U_{j_k}: k=1,2,\ldots\}$ converging to U_0 . Thus

$$\{\operatorname{tr}(CU_{i(\ell)}^*A_{j(\ell)}U_{j(\ell)}): k=1,2,\dots\} \to \operatorname{tr}(CU_0^*A_0U_0) = \mu_0 \in W_C(\mathcal{F}).$$

Thus, $W_C(\mathcal{F})$ is closed. As a result, $W_C(\mathcal{F})$ is compact.

The following examples show that none of the converses of the assertions in Theorem 2.3 (c) – (e) is valid, and there are no implications between the conditions that " \mathcal{F} is closed" and " $W_C(\mathcal{F})$ " is closed.

- **Example 2.4.** (a) Suppose $C \in M_n$ is non-scalar and has trace zero, and $\mathcal{F} = \{\mu I : \mu \in \mathbb{C}\}$. Then $W_C(\mathcal{F}) = \{0\}$ is bounded and compact, but \mathcal{F} is not bounded.
 - (b) Suppose $C \in M_n$ is non-scalar, and $\mathcal{F} = \{A_0, A_1\}$ such that $A_0 = 0$ and $A_1 = xy^*$ for a pair of orthonormal vectors x, y. Then $W_C(A_0) = \{0\}$ and $W_C(\mathcal{F}) = W_C(A_1)$ is a circular disk center at the origin with radius

$$R = \max\{|u^*Cv| : \{u, v\} \text{ is an orthonormal set}\}.$$

Thus, $W_C(\mathcal{F})$ is connected, but \mathcal{F} is not.

(c) Let $\mathcal{F} = \text{conv}\mathcal{G}$ with

$$\mathcal{G} = \{2E_{12}\} \cup \{\operatorname{diag}(e^{ir}, e^{-ir}) : r \text{ is a rational number}\}.$$

Then \mathcal{F} is not closed but $W(\mathcal{F}) = W(2E_{12})$ is closed.

(d) Let
$$\mathcal{F} = \left\{ \operatorname{diag}\left(0, x + \frac{i}{x}\right) : x > 0 \right\} \cup \{\operatorname{diag}(0, 0)\}$$
. Then \mathcal{F} is closed, but
$$W(\mathcal{F}) = \{x + iy : x, \ y > 0, \ xy \le 1\} \cup \{0\}$$

is not closed.

Next, we consider the connection between the geometrical properties of $W_C(\mathcal{F})$ and the properties of C and \mathcal{F} . Note that for any subset \mathcal{S} of \mathbb{C} , if $\operatorname{tr} C \neq 0$ and $\mathcal{F} = \{\mu I/\operatorname{tr} C : \mu \in \mathcal{S}\}$, then we have $W_C(\mathcal{F}) = \mathcal{S}$. Thus, the geometrical shape of $W_C(\mathcal{F})$ may be quite arbitrary. Also, if $C = \mu I$ is a scalar matrix, then $W_C(\mathcal{F}) = \{\mu \operatorname{tr} A : A \in \mathcal{F}\}$. Again, $W_C(\mathcal{F})$ does not contain much information about the matrices in \mathcal{F} . Nevertheless, we have the following.

Theorem 2.5. Suppose $C \in M_n$ is non-scalar, and $\mathcal{F} \subseteq M_n$ is non-empty. The following conditions hold.

- (a) The set $W_C(\mathcal{F}) = \{\mu\}$ if and only if $\mathcal{F} = \{\nu I : \nu \operatorname{tr} C = \mu\}$.
- (b) The set $W_C(\mathcal{F})$ is a subset of a straight line L if and only if
 - (i) $\mathcal{F} \subseteq \{ \nu I : \nu \in \mathbb{C}, \nu \text{tr } C \in L \}, \text{ or }$
 - (ii) there are complex units $\alpha, \gamma \in \mathbb{C}$ such that $\gamma(C (\operatorname{tr} C)I/n) \text{ and } \alpha(A (\operatorname{tr} A)I/n) \text{ are Hermitian for all } A \in \mathcal{F},$ and $\{(\operatorname{tr} C)(\operatorname{tr} A)/n : A \in \mathcal{F}\}$ is collinear.
- (c) The set $W_C(\mathcal{F})$ is a convex polygon if and only if $W_C(\mathcal{F}) = \text{conv}\{v_1, \ldots, v_m\}$ where each v_j is of the forms $(c_1, \ldots, c_n)(a_1, \ldots, a_n)^t$. Here c_1, \ldots, c_n are eigenvalues of C, and a_1, \ldots, a_n are eigenvalues of some $A_j \in \mathcal{F}$.

Proof. Condition (a) follows from the fact that $W_C(A) = \{\mu\}$ if and only if $A = \nu I$ with $\nu \operatorname{tr} C = \mu$.

(b) Suppose $W_C(\mathcal{F}) \subseteq L$. If $\mathcal{F} \subseteq \{\mu I : \mu \in \mathbb{C}\}$, then clearly $\mathcal{F} \subseteq \{\nu I : \nu \operatorname{tr} C \in L\}$. Let \mathcal{F} contains a non-scalar matrix A. Then $W_C(A)$ must be a non-degenerate line segment contained in L. Therefore C, A are normal with collinear eigenvalues in \mathbb{C} , see [10, (7.3)]. There exist complex units $\alpha, \gamma \in \mathbb{C}$ such that $\gamma(C - (\operatorname{tr} C)I/n)$ and $\alpha(A - (\operatorname{tr} A)I/n)$ are Hermitian. If $B \in \mathcal{F}$ is a scalar matrix, then $\alpha(B - (\operatorname{tr} B)I/n) = 0$ which is Hermitian. Now assume $B \in \mathcal{F}$ is non-scalar. Let $\tilde{C} = \gamma(C - (\operatorname{tr} C)I/n)$. Then

$$W_{\tilde{C}}(\alpha(B - (\operatorname{tr} B)I/n)) = \gamma \alpha W_{C}(B) + \mu_{B} \subseteq \{\gamma \alpha z + \mu_{B} : z \in L\},\$$

for some constant $\mu_B \in \mathbb{C}$, and

$$W_{\tilde{C}}(\alpha(A - (\operatorname{tr} A)I/n)) = \gamma \alpha W_{C}(A) + \mu_{A} \subseteq \{\gamma \alpha z + \mu_{A} : z \in L\},\$$

for some constant $\mu_A \in \mathbb{C}$. Hence $W_{\tilde{C}}(\alpha(B - (\operatorname{tr} B)I/n))$ is a subset of a line segment parallel to $W_{\tilde{C}}(\alpha(A - (\operatorname{tr} A)I/n)) \subseteq \mathbb{R}$. As $0 \in W_{\tilde{C}}(\alpha(B - (\operatorname{tr} B)I/n))$, we have $W_{\tilde{C}}(\alpha(B - (\operatorname{tr} B)I/n)) \subseteq \mathbb{R}$. Therefore, $\alpha(B - (\operatorname{tr} B)I/n)$ is Hermitian. The last assertion follows from $\{(\operatorname{tr} C)(\operatorname{tr} A)/n : A \in \mathcal{F}\} \subseteq W_C(\mathcal{F}) \subseteq L$. The sufficiency can be verified readily.

(c) Suppose $W_C(\mathcal{F}) = \text{conv}\{v_1, \dots, v_m\}$ is a convex polygon. Then for every v_j , there is $A_j \in \mathcal{F}$ such that $v_j = \text{tr}(CU_j^*A_jU_j) \in W_C(A_j)$ for some unitary $U_j \in M_n$. Since $W_C(A_j) \subseteq W_C(\mathcal{F})$, we see that v_j is a vertex point of $W_C(A_j)$. It follows that v_j has the form $(c_1, \dots, c_n)(a_1, \dots, a_n)^t$, where c_1, \dots, c_n are eigenvalues of C and a_1, \dots, a_n are eigenvalues of A_j arranged in some suitable order. The converse of the assertion is clear.

3 Star-shapedness and Convexity

In this section, we study the star-shapedness and convexity of $W_C(\mathcal{F})$. If \mathcal{F} is not connected, then $W_C(\mathcal{F})$ may not be connected so that $W_C(\mathcal{F})$ is not star-shaped or convex. One might hope that if \mathcal{F} is star-shaped or convex, then $W_C(\mathcal{F})$ will inherit the properties. However, the following examples show that $W_C(\mathcal{F})$ may fail to be convex (star-shaped, resp.) even if \mathcal{F} is convex (star-shaped, resp.).

Example 3.1. Let $C = E_{11}$, A = diag(1 + i, 1 - i) and $\mathcal{F} = \text{conv}\{A, -A\}$. Then

$$W_C(\mathcal{F}) = W(\mathcal{F}) = \bigcup_{t \in [0,1]} W(tA + (1-t)(-A)) = \bigcup_{s \in [-1,1]} sW(A).$$

As $W(A) = \operatorname{conv}\{1+i, 1-i\}$, we have $W_C(\mathcal{F}) = \operatorname{conv}\{0, 1+i, 1-i\} \cup \operatorname{conv}\{0, -1-i, -1+i\}$ which is not convex, see Figure 1.




Figure 1

Example 3.2. Let $C = E_{11}$, A = diag(1 + i, 1 - i), $\mathcal{F}_1 = \text{conv}\{A, -A\}$ and $\mathcal{F}_2 = \text{conv}\{A, -A + 4I\}$. Then $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2$ is star-shaped with star-center A. Since tA + (1 - t)(-A + 4I) = (1 - 2t)(2I - A) + 2I, we have

$$W(\mathcal{F}_2) = \bigcup_{t \in [0,1]} W((1-2t)(-A+2I) + 2I) = \bigcup_{s \in [-1,1]} sW(-A+2I) + 2.$$

Note that $W(-A+2I) = \operatorname{conv}\{1+i, 1-i\} = W(A)$. Then $W(\mathcal{F}_2) = W(\mathcal{F}_1) + 2$ and

$$W_C(\mathcal{F}) = W(\mathcal{F}_1 \cup \mathcal{F}_2) = W(\mathcal{F}_1) \cup W(\mathcal{F}_2)$$

equals $conv\{0, -1 + i, -1 - i\} \cup conv\{0, 2, 1 - i, 1 + i\} \cup conv\{2, 3 + i, 3 - i\}$ which is not star-shaped, see Figure 2.




Figure 2

Notice that in Example 3.1, \mathcal{F} is convex and $W_C(\mathcal{F})$ is star-shaped with star-center at the origin. One may ask if $W_C(\mathcal{F})$ is always star-shaped for a convex set \mathcal{F} . We will study this question in the following, and pay special attention on the case where $\mathcal{F} = \text{conv}\{A_1, \ldots, A_m\}$ for some $A_1, \ldots, A_m \in M_n$.

Denote by $S_C(A)$ the set of all star-centers of $W_C(A)$ and U_n the group of all $n \times n$ unitary matrices. We begin with the following result showing that $W_C(\mathcal{F})$ is star-shaped if C or \mathcal{F} satisfies some special properties.

Proposition 3.3. Suppose $C \in M_n$ and \mathcal{F} is a convex matrix set.

- (a) If \mathcal{F} contains a scalar matrix μI , then $W_C(\mathcal{F})$ is star-shaped with $\mu \text{tr } C$ as a star center.
- (b) Suppose the intersection of all (or any three of) $S_C(A)$ with $A \in \mathcal{F}$ is nonempty. Then $W_C(\mathcal{F})$ is star-shaped with μ as a star center for any $\mu \in \cap \{S_C(A) : A \in \mathcal{F}\}$.
- (c) If $\operatorname{tr} C = 0$, then $W_C(\mathcal{F})$ is star-shaped with 0 as a star-center.
- (d) If all matrices in \mathcal{F} have the same trace ν , then $W_C(\mathcal{F})$ is star-shaped with $\nu \operatorname{tr} C$ as a star-center.

Proof. (a) Suppose \mathcal{F} contains a scalar matrix μI and $B \in \mathcal{F}$. Then

$$\operatorname{conv}\{\operatorname{\mu tr} C, W_C(B)\} \subseteq W_C(\operatorname{conv}\{\operatorname{\mu} I, B\}) \subseteq W_C(\mathcal{F}).$$

The result follows.

(b) Suppose $\mu \in \cap \{S_C(A) : A \in \mathcal{F}\}$. Then for any $\nu \in W_C(\mathcal{F})$, there is $B \in \mathcal{F}$ such that $\nu \in W_C(B)$. As $\mu \in S_C(B)$, the line segment joining μ and ν will lie in $W_C(B) \subseteq W_C(\mathcal{F})$. Thus, $W_C(\mathcal{F})$ is star-shaped with μ as a star center.

If $S_C(A_0) \cap S_C(A_1) \cap S_C(A_2) \neq \emptyset$ for any $A_0, A_1, A_2 \in \mathcal{F}$, then $\cap \{S_C(A) : A \in \mathcal{F}\} \neq \emptyset$ by Helly's Theorem. So, the result follows from the preceding paragraph.

- (c) Note that $\frac{1}{n}(\operatorname{tr} C)(\operatorname{tr} A) \in S_C(A)$ for any $C, A \in M_n$, see [4] or Proposition 2.1 (d). If $\operatorname{tr} C = 0$, then $0 \in \bigcap \{S_C(A) : A \in \mathcal{F}\}$ is a star-center of $W_C(\mathcal{F})$ by (b).
- (d) The assumption implies that $\nu \operatorname{tr} C$ is the common star-center of $W_C(A)$ for all $A \in \mathcal{F}$. Thus, the result follows from (b).

Lemma 3.4. Let $C, A, B \in M_n$ and $\mathcal{F} = \text{conv}\{A, B\}$. If $\mu \in S_C(A) \cap S_C(B)$, then $W_C(\mathcal{F})$ is star-shaped with star-center μ .

Proof. Let $\zeta \in W_C(\mathcal{F})$. There are $V \in \mathcal{U}_n$ and $0 \le t \le 1$ such that $\zeta = \operatorname{tr}(CV^*(tA + (1 - t)B)V)$. It suffices to show $\operatorname{conv}\{\mu, \operatorname{tr}(CV^*AV), \operatorname{tr}(CV^*BV)\} \subseteq W_C(\mathcal{F})$. For any $a, b, c \in \mathbb{C}$, we let $\Delta(a, b, c) = \operatorname{conv}\{a, b\} \cup \operatorname{conv}\{b, c\} \cup \operatorname{conv}\{a, c\}$, i.e., the triangle (without the interior) with the vertices a, b, c. Let $U_A \in \mathcal{U}_n$ such that $\operatorname{tr}(CU_A^*AU_A) = \mu$. As $\mu \in S_C(A) \cap S_C(B)$ we have

$$\operatorname{conv}\{\operatorname{tr}(CV^*AV), \mu\} \cup \operatorname{conv}\{\operatorname{tr}(CV^*BV), \mu\} \subseteq W_C(\mathcal{F}).$$

Moreover we have

$$\operatorname{conv}\{\operatorname{tr}(CV^*AV), \operatorname{tr}(CV^*BV)\} = \{\operatorname{tr}(CV^*(tA + (1-t)B)V) : 0 \le t \le 1\} \subseteq W_C(\mathcal{F}).$$

Hence $\triangle(\operatorname{tr}(CV^*AV), \operatorname{tr}(CV^*BV), \mu) \subseteq W_C(\mathcal{F})$. We shall show that

$$\operatorname{conv}\{\operatorname{tr}(CV^*AV), \operatorname{tr}(CV^*BV), \mu\} \subseteq W_C(\mathcal{F}). \tag{3.1}$$

If $\triangle(\operatorname{tr}(CV^*AV), \operatorname{tr}(CV^*BV), \mu)$ is a line segment or a point, then eq. (3.1) holds clearly. Now assume that $\triangle(\operatorname{tr}(CV^*AV), \operatorname{tr}(CV^*BV), \mu)$ is non-degenerate. As \mathcal{U}_n is path-connected, we define a continuous function $f:[0,1] \to \mathcal{U}_n$ with f(0) = V and $f(1) = U_A$. For $0 \le t \le 1$, let $V_A(t) = \operatorname{tr}(Cf(t)^*Af(t))$ and $V_B(t) = \operatorname{tr}(Cf(t)^*Bf(t))$. Note that for any $t \in [0,1]$, we have

$$\triangle(t) = \triangle(V_A(t), V_B(t), \mu) \subseteq W_C(\mathcal{F}).$$




Figure 3: $\triangle(0)$ and $\triangle(t)$ are triangles in the red and green, respectively. In particular, $\triangle(1)$ degenerates to a line segment in purple.

For any $\zeta \in \text{conv}\Delta(0)$, see Figure 3, let

$$t_0 = \max\{t : \zeta \in \operatorname{conv}\Delta(s) \text{ for all } 0 \le s \le t\}.$$

Since $\triangle(1)$ degenerates, by continuity of f, we have $\zeta \in \triangle(t_0) \subseteq W_C(\mathcal{F})$. Hence the result follows.

One can extend Lemma 3.4 to a more general situation.

Theorem 3.5. Let $C \in M_n$ and \mathcal{G} be a (finite or infinite) family of matrices in M_n . If $\mu \in \bigcap_{A \in \mathcal{G}} S_C(A)$, then $W_C(\mathcal{F})$ is star-shaped with star-center μ for $\mathcal{F} = \text{conv}\mathcal{G}$.

Proof. By Lemma 3.4, the result holds if \mathcal{G} has 2 elements. Suppose $|\mathcal{G}| \geq 3$ and $\mu \in \bigcap_{A \in \mathcal{G}} S_c(A)$. Let $\zeta \in W_C(\mathcal{F})$. Then there exist $A_1, ..., A_m \in \mathcal{G}$, $t_1, ..., t_m > 0$ with $t_1 + \cdots + t_m = 1$ and $U \in \mathcal{U}_n$ such that $\zeta = \operatorname{tr}(CU^*(t_1A_1 + \cdots + t_mA_m)U)$. Let $\zeta_i = \operatorname{tr}(CU^*A_iU)$, i = 1, ..., m. Then $\zeta \in \operatorname{conv}\{\zeta_1, ..., \zeta_m\}$. The half line from μ through ζ intersects a line segment joining some ζ_i, ζ_j with $1 \leq i \leq j \leq m$ such that $\zeta \in \operatorname{conv}\{\mu, \zeta_i, \zeta_j\}$. By Lemma 3.4, we have $\operatorname{conv}\{\mu, \zeta_i, \zeta_j\} \subseteq W_C(\operatorname{conv}\{A_i, A_j\}) \subseteq W_C(\mathcal{F})$.

Note that if $W_C(A)$ is convex, then $W_C(A) = S_C(A)$. So, if $W_C(A)$ is convex for every $A \in \mathcal{G}$, and if $\mu \in \bigcap_{A \in \mathcal{G}} W_C(A)$, then μ is a star-center of $W_C(\operatorname{conv}(\mathcal{G}))$. Checking the condition that $W_C(A)$ is convex for every $A \in \mathcal{G}$ may not be easy. On the other hand, if $C \in M_n$ is such that $W_C(A)$ is always convex, then one can skip the checking process, and we have the following.

Corollary 3.6. Suppose $C \in M_n$ satisfies any one of the conditions (e.1)—(e.3) in Proposition 2.1 (e). If $\mathcal{G} \subseteq M_n$ and $\mu \in \cap_{A \in \mathcal{G}} W_C(A)$, then μ is a star-center of $W_C(\text{conv }\mathcal{G})$.

Next, we show that if $W_C(A)$ and $W_C(B)$ are convex, then $W_C(\mathcal{F})$ is star-shaped for $\mathcal{F} = \text{conv}\{A, B\}$ even when $W_C(A) \cap W_C(B) = \emptyset$.

Theorem 3.7. Let $C \in M_n$. Suppose $A, B \in M_n$ such that $W_C(A)$ and $W_C(B)$ are convex sets with empty intersection. Let $\mathcal{F} = \text{conv}\{A, B\}$.

If $W_C(A) \cup W_C(B)$ lies on a line, then $W_C(\mathcal{F}) = \text{conv}\{W_C(A), W_C(B)\}$ is convex.

Otherwise, there are two non-parallel lines L_1 and L_2 intersecting at μ such that for each $j = 1, 2, L_j$ is a common supporting line of $W_C(A)$ and $W_C(B)$ separating the two convex sets (i.e., $W_C(A)$ and $W_C(B)$ lying on opposite closed half spaces determined by L_j); the set $W_C(\mathcal{F})$ is star-shaped with star-center μ , see Figure 4.




Figure 4

Proof. Assume $W_C(A) \cup W_C(B)$ lies on a line. As \mathcal{F} is connected, by Theorem 2.3, $W_C(\mathcal{F})$ is connected. We have

$$W_C(A) \cup W_C(B) \subseteq W_C(\mathcal{F}) \subseteq \operatorname{conv}\{W_C(A), W_C(B)\}.$$

Therefore, $W_C(\mathcal{F}) = \text{conv}\{W_C(A), W_C(B)\}\$ which is convex.

Otherwise, we may assume that $W_C(A)$ lies on the left open half plane and $W_C(B)$ lies on the right open half plane. For $t \in [-\pi, \pi]$, let L(t) be the support line of $W_C(A)$ with (outward pointing) normal $n(t) = (\cos t, \sin t)$. Let $P(t) = L(t) \cap W_C(A)$. Therefore, for each $t \in [-\pi, \pi]$ and $p \in P(t)$, we have $n(t) \cdot (w - p) \leq 0$ for all $w \in W_C(A)$. Let $q(t) = \min\{n(t) \cdot (w - p) : w \in W_C(B), p \in P(t)\}$. Then we have q(0) > 0 and $q(-\pi) = q(\pi) < 0$. Hence, there exist $-\pi < t_1 < 0 < t_2 < \pi$ such that $q(t_1) = q(t_2) = 0$. Then $L_j = L(t_j)$ for j = 1, 2 will satisfy the requirement.


We claim that $L(t_1)$ and $L(t_2)$ in (a) cannot be parallel. Otherwise, we must have $t_1 = t_2 + \pi$. Therefore, $n(t_1) = -n(t_2)$ and $W_C(A), W_C(B) \subset L_1 = L_2$, is a contradiction. Since L_1 and L_2 are not parallel, they intersect at a point μ .


Now, we show that μ is a star-center of $W_C(\mathcal{F})$. We may apply a suitable affine transform to A and B, and assume that $\mu = 0$, L_1 and L_2 are the x-axis and y-axis respectively. We may further assume that $W_C(A)$ lies in the first quadrant and $W_C(B)$ lies in the third quadrant. For any $\zeta \in W_C(\mathcal{F})$, there are $V \in \mathcal{U}_n$ and $0 \le t \le 1$ such that $\zeta = \operatorname{tr}(CV^*(tA + (1-t)B)V)$. Denote $\zeta_A = \operatorname{tr}(CV^*AV)$ and $\zeta_B = \operatorname{tr}(CV^*BV)$. We claim that $\operatorname{conv}\{\zeta_A, \zeta_B, 0\} \subseteq W_C(\mathcal{F})$. Once the claim holds, we have $\{s\zeta + (1-s) \cdot 0 : 0 \le s \le 1\} \subseteq \operatorname{conv}\{\zeta_A, \zeta_B, 0\} \subseteq W_C(\mathcal{F})$. Then the star-shapedness of $W_C(\mathcal{F})$ follows.


We now show the claim. We denote by $\zeta_1\zeta_2$ the line segment with end points $\zeta_1, \zeta_2 \in \mathbb{C}$. By symmetry, we may assume that the line segment $\overline{\zeta_A\zeta_B}$ intersects the y-axis at (0,b) with $b \geq 0$. The situation is depicted in Figure 5a and Figure 5b.

We shall first show that $\operatorname{conv}\{\zeta_B, bi, 0\} \subseteq W_C(\mathcal{F})$. Let y_B be a point in the intersection of $W_C(B)$ and the y-axis, $U_B \in \mathcal{U}_n$ be such that $y_B = \operatorname{tr}(CU_B^*BU_B)$. Let $y_A = \operatorname{tr}(CU_B^*AU_B)$. Then by the convexity of $W_C(A)$, $W_C(B)$ and \mathcal{F} , we have

$$Q((\zeta_A, y_A); (\zeta_B, y_B)) := \overline{\zeta_A y_A} \cup \overline{\zeta_B y_B} \cup \overline{\zeta_A \zeta_B} \cup \overline{y_A y_B} \subseteq W_C(\mathcal{F}).$$







(b) $Q((\zeta_A, y_A); (\zeta_B, y_B))$ is a union of two triangle.

Figure 5

Note that $Q((\zeta_A, y_A); (\zeta_B, y_B))$ is either a quadrilateral or a union of two triangle, see Figure 5a and Figure 5b. In both case, $\Delta(\zeta_B, bi, 0)$ lies inside the region determined by the closed curve $Q((\zeta_A, y_A); (\zeta_B, y_B))$. Now consider a continuous function $f : [0, 1] \to \mathcal{U}_n$ with f(0) = V and $f(1) = U_B$. Then for $t \in [0, 1]$, let $\zeta'_A(t) = \operatorname{tr}(Cf(t)^*Af(t))$ and $\zeta'_B(t) = \operatorname{tr}(Cf(t)^*Bf(t))$, we have

$$Q((\zeta_A'(t), y_A); (\zeta_B'(t), y_B)) := \overline{\zeta_A'(t)y_A} \cup \overline{\zeta_B'(t)y_B} \cup \overline{\zeta_A'(t)\zeta_B'(t)} \cup \overline{y_Ay_B} \subseteq W_C(\mathcal{F}).$$

Note that $Q((\zeta'_A(1), y_A); (\zeta'_B(1), y_B))$ degenerates, by continuity of f, for any point ζ enclosed by the closed curve $Q((\zeta_A, y_A); (\zeta_B, y_B))$, there exists $0 \le t_0 \le 1$ such that $\zeta \in Q((\zeta'_A(t_0), y_A); (\zeta'_B(t_0), y_B)) \subseteq W_C(\mathcal{F})$. Hence, $\operatorname{conv}\{\zeta_B, bi, 0\} \subseteq W_C(\mathcal{F})$. By symmetry, one can show that $\operatorname{conv}\{\zeta_A, a, 0\} \subseteq W_C(\mathcal{F})$ where a is the intersection point of $\overline{\zeta_A\zeta_B}$ and x-axis. Hence, $\operatorname{conv}\{\zeta_A, \zeta_B, 0\} = \operatorname{conv}\{\zeta_A, a, 0\} \cup \operatorname{conv}\{\zeta_B, bi, 0\} \subseteq W_C(\mathcal{F})$. The claim follows.

4 Additional results on star-shapedness and convexity

It is not easy to extend Theorem 3.7. The following example shows that if $W_C(A)$ and $W_C(B)$ are not convex, then $W_C(\mathcal{F})$ may not be star-shaped for $\mathcal{F} = \text{conv}\{A, B\}$ even when $W_C(A) \cap W_C(B) \neq \emptyset$.

Example 4.1. Let $w = e^{2\pi i/3}$ and $C = \text{diag}(1, w, w^2)$. Suppose $A = C - \frac{1}{6}I$, $B = e^{\pi i/3}C + \frac{1}{6}I$ and $\mathcal{F} = \text{conv}\{A, B\}$. Then $W_{C+I}(\mathcal{F})$ is not star-shaped.

To prove our claim, for $0 \le t \le 1$, let

$$A(t) = tA + (1-t)B = (t + (1-t)e^{\pi i/3})C + \frac{1-2t}{6}I.$$

$$W_{C+I}(A(t)) = W_{C+I}\left((t+(1-t)e^{\pi i/3})C + \frac{1-2t}{6}I\right)$$

$$= (t+(1-t)e^{\pi i/3})W_C(C) + \left(t+(1-t)e^{\pi i/3} + \frac{1-2t}{6}\right)\operatorname{tr} C + \frac{1-2t}{2}$$

$$= (t+(1-t)e^{\pi i/3})W_C(C) + \frac{1-2t}{2}.$$

Therefore,

$$W_{C+I}(\mathcal{F}) = \bigcup \{ (t + (1-t)e^{\pi i/3}) W_C(C) + \frac{1-2t}{2} : 0 \le t \le 1 \}.$$

By [14], $W_C(C)$ is star-shaped with the origin as the unique star-center. Moreover $W_C(C) = e^{2\pi i/3}W_C(C)$ and its boundary is given by

$$\Gamma = \{2e^{i\theta} + e^{-2i\theta} : -\pi \le \theta \le \pi\}.$$

Hence, $W_{C+I}(A(t))$ is star-shaped with $\frac{1-2t}{2}$ as the unique star-center and its boundary is given by

$$\Gamma(t) = \{ (t + (1 - t)e^{\pi i/3}) \left(2e^{i\theta} + e^{-2i\theta} \right) : -\pi \le \theta \le \pi \}. \tag{4.2}$$

For $0 \le t \le 1$ and $-\pi \le \theta \le \pi$, define

$$f(\theta,t) = (t + (1-t)e^{\pi i/3})(2e^{i\theta} + e^{-2i\theta}) + \frac{1-2t}{2}$$

$$= \frac{1+t}{2}(2\cos\theta + \cos 2\theta) - \frac{1-t}{2}\sqrt{3}(2\sin\theta - \sin 2\theta) + \frac{1-2t}{2}$$

$$+ i\left(\frac{1-t}{2}\sqrt{3}(2\cos\theta + \cos 2\theta) + \frac{1+t}{2}(2\sin\theta - \sin 2\theta)\right).$$

Using this description, we can plot $W_{C+I}(\mathcal{F})$ as follows.

First consider the plot of $\Gamma(0)$ and $\Gamma(1)$, which are the boundaries of $W_{C+I}(B)$ and $W_{C+I}(A)$ respectively, see Figure 6.




Figure 6: $W_{C+I}(A) \cup W_{C+I}(B)$

The "vertices" of $\Gamma(0)$ and $\Gamma(1)$ are given by

$$f\left(-\frac{2\pi}{3},0\right) = 2 - \frac{3\sqrt{3}}{2}i \qquad f(0,0) = 2 + \frac{3\sqrt{3}}{2}i \qquad f\left(\frac{2\pi}{3},0\right) = -\frac{5}{2}$$

$$f\left(-\frac{2\pi}{3},1\right) = -2 - \frac{3\sqrt{3}}{2}i \qquad f(0,1) = \frac{5}{2} \qquad \qquad f\left(\frac{2\pi}{3},1\right) = -2 + \frac{3\sqrt{3}}{2}i.$$

As t increases from 0 to 1, $\Gamma(t)$ changes from $W_{C+I}(B)$ to $W_{C+I}(A)$, For $\theta = -\frac{2\pi}{3}, 0, \frac{2\pi}{3}$, the vertex $f(\theta, t)$ moves along the line segment from $f(\theta, 0)$ to $f(\theta, 1)$, see Figure 7.




Figure 7: $W_{C+I}(\mathcal{F})$

Therefore, we see that $W_{C+I}(\mathcal{F})$ is the union of $W_{C+I}(A)$, $W_{C+I}(B)$ and three triangles, $\triangle\left(0.5, 2.5, 2 + \frac{3\sqrt{3}}{2}i\right)$, $\triangle\left(-0.5, -2.5, -2 + \frac{3\sqrt{3}}{2}i\right)$, $\triangle\left(\frac{\sqrt{3}}{2}i, 2 - \frac{3\sqrt{3}}{2}i, -2 - \frac{3\sqrt{3}}{2}i\right)$, see Figure 8.




Figure 8: $W_{C+I}(\mathcal{F})$

Suppose $W_{C+I}(\mathcal{F})$ is star-shaped. By symmetry, $W_{C+I}(\mathcal{F})$ has a star-center c on the imaginary axis. The line segment joining c and f(0,0) must lie below the tangent line to $\Gamma(0)$ at $f(0,0) = 2 + \frac{3\sqrt{3}}{2}i$. By direct calculation, this tangent line is the line through 0.5 and f(0,0). Similarly, The line segment joining c and f(1,0) must lie above the tangent line to $\Gamma(1)$ at f(1,0) = 2.5. By direct calculation, this tangent line is the line through 0.5 and f(1,0). Since these two tangent lines intersect at 0.5, no c on the imaginary axis will satisfy the above conditions.

In the following, we present an example of $C, A_1, A_2, A_3 \in M_n$ such that $W_C(A_1), W_C(A_2), W_C(A_3)$ are convex and $W_C(A_1) \cap W_C(A_2) \cap W_C(A_3) = \emptyset$, but $W_C(\mathcal{F})$ is not star-shaped for $\mathcal{F} = \text{conv}\{A_1, A_2, A_3\}$. In particular, we choose $C = E_{11}$ so that $W_C(\mathcal{F}) = W(\mathcal{F})$.

To describe our example, we first show that the set $W(\mathcal{F})$ is connected to the concept of product numerical range arising in the study of quantum information theory, [15].

Let $A \in M_m \otimes M_n = M_m(M_n)$, the product numerical range of A is

$$W^{\otimes}(A) = \{(u \otimes v)^* A (u \otimes v) : u \in \mathbb{C}^m, \ v \in \mathbb{C}^n \}.$$

We note that $W^{\otimes}(A)$ depends on order of the factors M_m and M_n in the representation of $M_{mn} = M_m \otimes M_n$.

Theorem 4.2. Let $A_1, \ldots, A_m \in M_n$ and $\mathcal{F} = \text{conv}(\{A_1, \ldots, A_m\})$. Then

$$W(\mathcal{F}) = W^{\otimes}(\bigoplus_{i=1}^{m} A_i).$$

Proof. Let $\mu \in W(\mathcal{F})$ Then there exist $t_1, \ldots, t_m \geq 0$ with $\sum_{i=1}^m t_i = 1$ and $v \in \mathbb{C}^n$ such that $\mu = v^* (\sum_{i=1}^m t_i A_i) v$. Let $u = (\sqrt{t_1}, \ldots, \sqrt{t_m})^t$. Then $\mu = (u \otimes v)^* (\bigoplus_{i=1}^m A_i) (u \otimes v) \in W^{\otimes} (\bigoplus_{i=1}^m A_i)$.

Conversely, suppose $\mu \in W^{\otimes}(\bigoplus_{i=1}^{m} A_i)$. Then there exist unit vectors $u \in \mathbb{C}^m$, $v \in \mathbb{C}^n$ such that $\mu = (u \otimes v)^*(\bigoplus_{i=1}^{m} A_i)(u \otimes v)$. Let $u = (u_1, \dots, u_m)^t$. Set $t_i = |u_i|^2$ for $i = 1, \dots, m$. Then $\mu = v^*(\sum_{i=1}^{m} t_i A_i) v \in W(\mathcal{F})$.

Now, we describe the follow example showing that $W(\mathcal{F})$ is not necessarily star-shaped for $\mathcal{F} = \text{conv}\{A_1, A_2, A_3\}$ if $W(A_1) \cap W(A_2) \cap W(A_3) = \emptyset$.

Example 4.3. Suppose $A = \text{diag}\left(e^{i\frac{\pi}{3}}, e^{-i\frac{\pi}{3}}, 0.95e^{i\frac{\pi}{4}}\right)$. Let $A_1 = e^{i\frac{\pi}{3}}A$, $A_2 = e^{-i\frac{\pi}{3}}A$ and $A_3 = 0.95e^{i\frac{\pi}{4}}A$. Then by Theorem 4.2, $W(\text{conv}\{A_1, A_2, A_3\})$ is equal to

$$W^{\otimes}(\bigoplus_{i=1}^{m} A_i) = W^{\otimes}(A \otimes A) = W(A) \cdot W(A) = \{\mu_1 \mu_2 : \mu_1, \mu_2 \in W(A)\}.$$

By the result in [11, Example 3.1], $W(A) \cdot W(A)$ is not star-shaped.

We note that in this example, $W(A_i) \cap W(A_j) \neq \emptyset$ for all i, j, but $W(A_1) \cap W(A_2) \cap W(A_3) = \emptyset$. Therefore, in some sense, the condition in Theorem 3.5 is optimal.

While Theorem 3.5 and Theorem 3.7 provide some sufficient conditions for the star-shapedness of $W_C(\mathcal{F})$, it is a challenging problem to determine whether $W_C(\mathcal{F})$ is star-shaped or not for a given $C \in M_n$ and $\mathcal{F} \subseteq M_n$.

Proposition 4.4. Suppose $\mathcal{F} = \text{conv}\{A_1, \dots, A_m\} \subseteq M_n$. For each unit vector $x \in \mathbb{C}^n$, let $\mathcal{A}_x = \text{diag}(x^*A_1x, \dots, x^*A_mx)$, and let $\hat{\mathcal{F}} = \{\mathcal{A}_x : x \in \mathbb{C}^n, x^*x = 1\}$. Then

$$W(\mathcal{F}) = W(\hat{\mathcal{F}}) = W(\operatorname{conv}\hat{\mathcal{F}}).$$

Proof. Let $\mu \in W(\mathcal{F})$. Then there exist $t_1, \ldots, t_m \geq 0$, $t_1 + \cdots + t_m = 1$ and a unit vector $x \in \mathbb{C}^n$ such that $\mu = x^*(t_1A_1 + \cdots + t_mA_m)x = t_1x^*A_1x + \cdots + t_mx^*A_mx \in W(\hat{\mathcal{F}})$. Therefore, $W(\mathcal{F}) \subseteq W(\hat{\mathcal{F}})$.

Clearly, $W(\hat{\mathcal{F}}) \subseteq W(\operatorname{conv}\hat{\mathcal{F}})$. Finally, we show that $W(\operatorname{conv}\hat{\mathcal{F}}) \subseteq W(\mathcal{F})$. To see this, let $\mu \in W(\operatorname{conv}\hat{\mathcal{F}})$, i.e.,

$$\mu = y^* \left(\sum_{j=1}^r t_j D_j \right) y$$

for a unit vector $y = (y_1, \ldots, y_m)^t$, some $D_j = \operatorname{diag}(x_j^* A_1 x_j, \ldots, x_j^* A_m x_j) \in \hat{\mathcal{F}}$ with $j = 1, \ldots, r$ and $t_1, \ldots, t_r > 0$, $t_1 + \cdots + t_r = 1$. Thus,

$$\mu = \sum_{\ell=1}^{m} |y_{\ell}|^{2} \sum_{j=1}^{r} t_{j} x_{j}^{*} A_{\ell} x_{j} = \sum_{j=1}^{r} t_{j} \left(x_{j}^{*} \left(\sum_{\ell=1}^{m} |y_{\ell}|^{2} A_{\ell} \right) x_{j} \right)$$

$$\in \text{conv}(W(A_{0})) = W(A_{0}) \subseteq W(\mathcal{F})$$

where $A_0 = \sum_{\ell=1}^m |y_\ell|^2 A_\ell \in \mathcal{F}$.

Example 4.5. Suppose that $A_1 = \operatorname{diag}\left(1, e^{i\frac{\pi}{3}}\right)$, $A_2 = \operatorname{diag}\left(e^{i\frac{2\pi}{3}}, e^{i\pi}\right)$, $A_2 = \operatorname{diag}\left(e^{i\frac{4\pi}{3}}, e^{i\frac{5\pi}{3}}\right)$ and $\mathcal{F} = \operatorname{conv}\{A_1, A_2, A_3\}$. Then $W(A_1) \cap W(A_2) \cap W(A_3) = \emptyset$. So the condition in Theorem 3.5 is not satisfied. However, by direct computation, we have

$$\hat{\mathcal{F}} = \left\{ \operatorname{diag}\left(t + (1-t)e^{i\frac{\pi}{3}}, e^{i\frac{2\pi}{3}}(t + (1-t)e^{i\frac{\pi}{3}}), e^{i\frac{4\pi}{3}}(t + (1-t)e^{i\frac{\pi}{3}})\right) : 0 \le t \le 1 \right\},\,$$

where $\hat{\mathcal{F}}$ is defined as in Proposition 4.4. Therefore, $0 \in \bigcap_{A \in \hat{\mathcal{F}}} W(A)$. Hence, by Proposition 4.4 and Theorem 3.5, $W(\mathcal{F}) = W(\hat{\mathcal{F}})$ is star-shaped.

Recall that $W_C(\mathcal{F})$ may fail to be convex even if \mathcal{F} is convex and $W_C(A)$ is convex for all $A \in \mathcal{F}$. In the following, we give a necessary and sufficient condition for $W_C(\mathcal{F})$ to be convex. First, it is easy to see that $W_C(A) \subseteq W_C(\mathcal{F})$ for every $A \in \mathcal{F}$. Thus, $\operatorname{conv}\{\bigcup_{A \in \mathcal{F}} W_C(A)\}$ is the smallest convex set containing $W_C(\mathcal{F})$. As a result, we have the following observation.

Proposition 4.6. Let $C \in M_n$ and $\mathcal{F} \subseteq M_n$. Then $W_C(\mathcal{F})$ is convex if and only if

$$W_C(\mathcal{F}) = \operatorname{conv} \left\{ \bigcup_{A \in \mathcal{F}} W_C(A) \right\}.$$

Proposition 4.7. Suppose $C \in M_n$, $\mathcal{G} \subseteq M_n$ and $\mathcal{F} = \operatorname{conv}\mathcal{G}$ with $|\mathcal{G}| \geq 3$. Then for every $\mu \in W_C(\mathcal{F})$, there are matrices $A_1, A_2, A_3 \in \mathcal{G}$ and a unitary matrix U such that $\mu \in \operatorname{conv}\{\operatorname{tr}(CU^*A_1U), \operatorname{tr}(CU^*A_2U), \operatorname{tr}(CU^*A_3U)\}$. If μ is a boundary point of $W_C(\mathcal{F})$, then there are $B_1, B_2 \in \mathcal{G}$ and a unitary V such that $\mu \in \operatorname{conv}\{\operatorname{tr}(CV^*B_1V), \operatorname{tr}(CV^*B_2V)\}$.

Proof. For any $\mu \in W_C(\mathcal{F})$, we have $\mu = t_1 \operatorname{tr} (CU^*A_1U) + \cdots + t_r \operatorname{tr} (CU^*A_rU)$ for some unitary matrix U and $A_1, \ldots, A_r \in \mathcal{G}$ with $t_1, \ldots, t_r > 0$ summing up to 1. So, $\mu \in \operatorname{conv}\{\operatorname{tr} (CU^*A_1U), \ldots, \operatorname{tr} (CU^*A_rU)\}$. Thus, μ lies in the convex hull of no more than three of the points in $\{\operatorname{tr} (CU^*A_1U), \ldots, \operatorname{tr} (CU^*A_rU)\}$. In case μ is a boundary point of $W_C(\mathcal{F})$,

then μ must be a boundary point of the set conv $\{\operatorname{tr}(CU^*A_1U), \ldots, \operatorname{tr}(CU^*A_rU)\}$. Thus, μ list in the convex hull of no more than two points in the set $\{\operatorname{tr}(CU^*A_1U), \ldots, \operatorname{tr}(CU^*A_rU)\}$. The assertion follows.

Theorem 4.8. Suppose $C \in M_n$, $\mathcal{G} \subseteq M_n$ is compact and $\mathcal{F} = \text{conv}(\mathcal{G})$. The following are equivalent.

- (a) $W_C(\mathcal{F})$ is convex.
- (b) $W_C(\mathcal{F})$ is simply connected and every boundary point $\mu \in \text{conv}W_C(\mathcal{F})$ has the form $\text{tr}(CU^*(tA+(1-t)B)U)$ for some unitary matrix $U, t \in [0,1]$ and $A, B \in \mathcal{G}$.

Proof. Suppose $W_C(\mathcal{F})$ is convex. Then it is clearly simply connected. Now, $\operatorname{conv} W_C(\mathcal{F})$ and $W_C(\mathcal{F})$ have the same boundary. By Proposition 4.7, every boundary point of $W_C(\mathcal{F})$ is a convex combination of $\operatorname{tr}(CU^*AU)$, $\operatorname{tr}(CU^*BU)$ with $A, B \in \mathcal{G}$.

Conversely, suppose (b) holds. We only need to show that each boundary point μ of $conv(W_C(\mathcal{F}))$ lies in $W_C(\mathcal{F})$, which is true by Proposition 4.7 and assumption (b).

5 Extension to the joint C-numerical range

Let $A = A_1 + iA_2 \in M_n$, where $A_1, A_2 \in M_n$ are Hermitian matrices. Then W(A) can be identified as the joint numerical range of (A_1, A_2) defined by

$$W(A_1, A_2) = \{(x^*A_1x, x^*A_2x) : x \in \mathbb{C}^n, x^*x = 1\}.$$

One may consider whether our results can be extended to the joint numerical range of an m-tuple of Hermitian matrices (A_1, \ldots, A_m) defined by

$$W(A_1, \dots, A_m) = \{(x^*A_1x, \dots, x^*A_mx) : x \in \mathbb{C}^n, x^*x = 1\}.$$

Some of the results on classical numerical range are not valid for the joint numerical range. For instance, the joint numerical range of three matrices may not be convex if n = 2.

For
$$A_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$
, $A_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$, $A_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, we have

$$W(A_1, A_2, A_3) = \{(a, b, c) : a, b, c \in \mathbb{R}, a^2 + b^2 + c^2 = 1\}.$$

The following is known, see [1, 12].

- 1. Suppose $A_1, A_2, A_3 \in M_2$ are Hermitian matrices such that $\{I_2, A_1, A_2, A_3\}$ is linearly independent. Then $W(A_1, A_2, A_3)$ is an ellipsoid without interior in \mathbb{R}^3 .
- 2. If $n \geq 3$ and $A_1, A_2, A_3 \in M_n$ are Hermitian matrices, then $W(A_1, A_2, A_3)$ is convex.

3. Suppose $A_1, A_2, A_3 \in M_n$ such that $\{I_2, A_1, A_2, A_3\}$ is linearly independent. Then there is $A_4 \in M_n$ such that $W(A_1, A_2, A_3, A_4)$ is not convex.

There has been study of topological and geometrical properties of $W(A_1, \ldots, A_m)$. Researchers also consider the joint C-numerical range of (A_1, \ldots, A_m) defined by

$$W_C(A_1, \dots, A_m) = \{ (\operatorname{tr}(CU^*A_1U), \dots, \operatorname{tr}(CU^*A_mU)) : U \text{ unitary} \}$$
 (5.3)

for a Hermitian matrix C, for example [5, 6]. In particular, it is known that $W_C(A_1, A_2, A_3)$ is convex for any Hermitian matrices $C, A_1, A_2, A_3 \in M_n$ if $n \geq 3$, see [2]. Of course, one can also consider the C-numerical range of (A_1, \ldots, A_m) for general matrices $C, A_1, \ldots, A_m \in M_n$ defined as in Equation (5.3).

Denote by $\mathbf{A} = (A_1, \dots, A_m)$ an *m*-tuple of matrices in M_n . Let \mathbf{F} be a non-empty subset of M_n^m . We consider

$$W_C(\mathbf{F}) = \bigcup \{W_C(\mathbf{A}) : \mathbf{A} \in \mathbf{F}\}.$$

Evidently, when $\mathbf{F} = \{\mathbf{A}\}$, then $W_C(\mathbf{F}) = W_C(\mathbf{A})$.

5.1 Basic results

We begin with the following results.

Proposition 5.1. Let $C \in M_n$ be non-scalar, and let **F** be a non-empty subset of M_n^m .

1. For any unitary $U, V \in M_n$, we have $W_C(\mathbf{F}) = W_{V^*CV}(U^*\mathbf{F}U)$, where

$$U^*\mathbf{F}U = \{(U^*A_1U, \dots, U^*A_mU) : (A_1, \dots, A_m) \in \mathbf{F}\}.$$

- 2. Let $\gamma_1, \gamma_2 \in \mathbb{C}$ with $\gamma_1 \neq 0$. If $\hat{C} = \gamma_1 C + \gamma_2 I_n$, then for any $\mathbf{A} = (A_1, \dots, A_m) \in M_n^m$, $W_{\hat{C}}(\mathbf{A}) = \{\gamma_1(a_1, \dots, a_m) + \gamma_2(\operatorname{tr} A_1, \dots, \operatorname{tr} A_m) : (a_1, \dots, a_m) \in W_C(\mathbf{A})\}.$
- 3. For any $T = (t_{ij}) \in M_m$ and $f = (f_1, \ldots, f_m)^t$, we can define an affine map R on \mathbb{C}^m by $v \mapsto Tv + f$, and extend the affine map to M_n^m by mapping $\mathbf{A} = (A_1, \ldots, A_m)$ to $\mathbf{B} = (B_1, \ldots, B_m)$ with $B_i = \sum_{j=1}^n t_{ij} A_j + f_i I_n$. Then

$$W_C(\mathbf{B}) = \{ (b_1, \dots, b_m) : (b_1, \dots, b_m)^t = T(a_1, \dots, a_n)^t + (\operatorname{tr} C)(f_1, \dots, f_m)^t, (a_1, \dots, a_m) \in W_C(\mathbf{A}) \}.$$

Consequently,

$$R(W_C(\mathbf{F})) = W_C(R(\mathbf{F})).$$

4. The linear span of $\{A_j - (\operatorname{tr} A_j)/n : j = 1, ..., m\}$ has dimension k if and only if $W_C(\mathbf{F}) \subseteq \mathbf{V} + f$ for a k-dimensional subspace $\mathbf{V} \subseteq \mathbb{C}^m$ and a vector $f \in \mathbb{C}^m$. In particular, $W_C(\mathbf{F})$ is a singleton $\{(\nu_1, ..., \nu_n)\}$ if and only if $\mathcal{F} = \{(\mu_1 I, ..., \mu_m I)\}$ with $(\operatorname{tr} C)(\mu_1, ..., \mu_m) = (\nu_1, ..., \nu_n)$.

Note that if $C \in M_n$ is Hermitian, then $W_C(A_1, \ldots, A_m) \subseteq \mathbb{C}^m$ can be identified as $W_C(X_1, Y_1, X_2, Y_2, \ldots, X_m, Y_m) \subseteq \mathbb{R}^{2m}$, where $X_j = (A_j + A_j^*)/2$ and $Y_j = (A_j - A_j^*)/(2i)$. One can obtain a "real version" of Proposition 5.1 using real scalars γ_1, γ_2 , real matrix T, real vector f, etc.

One can prove the following when $W_C(\mathbf{F})$ is a polyhedral set in \mathbb{C}^m , i.e., a convex combination of a finite set of vertices.

Proposition 5.2. Suppose $C \in M_n$ is non-scalar, and \mathbf{F} is a non-empty set of M_n^m . If $W_C(\mathbf{F})$ is polyhedral, then every vertex has the form (μ_1, \ldots, μ_m) with $\mu_j = \operatorname{tr} V^*CVU^*A_jU$, where $(A_1, \ldots, A_m) \in \mathbf{F}$, $U, V \in M_n$ are unitary so that $V^*CV, U^*A_1U, \ldots, U^*A_mU$ are in lower triangular matrices with diagonal entries

$$c_1, \ldots, c_n, a_1(1), \ldots, a_n(1), \ldots, a_1(m), \ldots, a_n(m)$$

and $\mu_j = \sum_{\ell=1}^n c_\ell a_\ell(j)$. Furthermore, if c_1, \ldots, c_n are distinct, then

$$V^*CV, U^*A_1U, \dots, U^*A_mU$$

are diagonal matrices.

If C has distinct eigenvalues, and if $W_C(\mathbf{A})$ has a conical point (μ_1, \ldots, μ_m) on the boundary, i.e., there is a pointed cone $K \subseteq \mathbb{C}^m \equiv \mathbb{R}^{2m}$ with vertex (μ_1, \ldots, μ_m) such that

$$W_C(\mathbf{A}) \cap \{(\mu_1 + \nu_1, \dots, \mu_m + \nu_m) : |\nu_i| \le \varepsilon\} \subseteq K$$

for some sufficiently small $\varepsilon > 0$, then $\{C, A_1, \ldots, A_m\}$ is a commuting family of normal matrices, and $\operatorname{conv} W_C(\mathbf{A})$ is a polyhedral set, see [3].

Note that if there is $\mathbf{B} \in \mathbf{F}$ such that $W_C(\mathbf{B})$ is a singleton, a subset of a straight line, or a convex polygon, then we can apply the results on $W_C(\mathbf{B})$ to deduce that C and \mathbf{B} has special structure. Then we can deduce results on $W_C(\mathbf{F})$.

We can obtain some topological properties $W_C(\mathbf{F})$.

Proposition 5.3. Let $C \in M_n$ be non-scalar, and $\mathbf{F} \subseteq M_n^m$ be a nonempty set.

- 1. If \mathcal{F} is bounded, then so is $W_C(\mathbf{F})$.
- 2. If \mathcal{F} is connected, then so is $W_C(\mathbf{F})$.
- 3. If \mathcal{F} is compact, then so is $W_C(\mathbf{F})$.

Proof. The proof of the boundedness and compactness are similar to those for $W_C(\mathcal{F})$ in Section 2. The prove connectedness, for any $\mathbf{A}, \mathbf{B} \in \mathbf{F}$ and unitary $U_0, U_1 \in M_n$, there is a path joining U_t with $t \in [0,1]$ joining U_0 and U_t , and hence there is a path joining $(\operatorname{tr} U_0^* C U_0 A_1, \ldots, \operatorname{tr} U_0^* C U_0 A_m)$ to $(\operatorname{tr} U_1^* C U_1 A_1, \ldots, \operatorname{tr} U_1^* C U_1 A_m)$, which is connected to $(\operatorname{tr} U_1^* C U_1 B_1, \ldots, \operatorname{tr} U_1^* C U_1 B_m)$.

5.2 Star-shapedness and convexity

Here we consider whether $W(\mathbf{F})$ is star-shaped or convex. It is known that if $C \in M_n$ is Hermitian, then $W_C(A_1, A_2, A_3)$ is convex for any Hermitian $A_1, A_2, A_3 \in M_n$ with $n \geq 3$, see [2]. One may wonder whether Theorem 3.5 admits an extension to this setting. The following example shows that the answer is negative.

Example 5.4. Let

$$\mathbf{A}_t = \left(\begin{pmatrix} t & & \\ & t & \\ & & 1-t \end{pmatrix}, \begin{pmatrix} 1-t & & \\ & & t & \\ & & & 1-t \end{pmatrix} \begin{pmatrix} 0 & & \\ & t & \\ & & & 1-t \end{pmatrix} \right), \qquad t \in [0,1].$$

Notice that $\operatorname{conv}\{\mathbf{A}_0, \mathbf{A}_1\} = \{\mathbf{A}_t : 0 \le t \le 1\}$. Moreover, $(0,0,0) \in W(\mathbf{A}_0) \cap W(\mathbf{A}_1)$, but (0,0,0) is not a star-center of $W(\operatorname{conv}\{\mathbf{A}_0,\mathbf{A}_1\})$ as $(1/2,1/2,0) \in W(\mathbf{A}_{1/2})$, however, $\operatorname{conv}\{(0,0,0),(1/2,1/2,0)\} \not\subseteq W(\operatorname{conv}\{\mathbf{A}_0,\mathbf{A}_1\})$. Nevertheless, $(1/2,1/2,1/2) \in W(\mathbf{A}_t)$ for all $t \in [0,1]$, and hence (1/2,1/2,1/2) is a star-center of $W(\operatorname{conv}\{\mathbf{A}_0,\mathbf{A}_1\})$. Actually, one can show that $W(\mathbf{A}_t) = \operatorname{conv}\{(t,1-t,0),(t,t,t),(1-t,1-t,1-t)\}$.

Here is an example showing that $W(\mathbf{A})$ may not be star-shaped in general.

Example 5.5. Let $\mathbf{A} = (A_1, A_2, A_3)$ with $A_1 = \text{diag}(0, 1, 0)$, $A_2 = \text{diag}(1, 0, -1)$, $A_3 = I_3$ and let $\mathbf{B} = (B_1, B_2, B_3)$ with $B_1 = \text{diag}(1, 0, 0)$, $B_2 = \text{diag}(0, -1, 1)$, $B_3 = 0_3$. If $\mathbf{F} = \text{conv}\{\mathbf{A}, \mathbf{B}\}$, then $W(\mathbf{F})$ is the union of the triangular disk with vertices

$$(1-t,t,t), (t,t-1,t), (0,1-2t,t).$$

Let $g: \mathbb{R}^3 \to \mathbb{R}^3$ be given by g((a, b, c)) = (a, -b, 1 - c). We have

$$g((1-t,t,t)) = (1-t,-t,1-t) = (1-t,(1-t)-1,1-t)$$

$$g((t, t-1, t)) = (t, 1-t, 1-t) = (1-(1-t), (1-t), (1-t))$$

$$g((0, 1-2t, t)) = (0, 2t-1, 1-t) = (0, 1-2(1-t), (1-t))$$

Therefore, $g(W(\mathbf{F})) = W(\mathbf{F})$. Moreover, we claim that $W(\mathbf{F})$ is not star-shaped.

Suppose the contrary that $W(\mathbf{F})$ is star-shaped with (a,b,t_0) be a star-center. Then $(a,-b,1-t_0)$ is also a star-center. As the set of all star-center of a star-shaped set is convex. We may now assume without loss of generality that b=0 and $t_0=1/2$. Assume now a>0. By assumption, for all $0 \le t \le 1$, $t(a,0,1/2)+(1-t)(1,0,0)=(1-t(1-a),0,t/2) \in W(\mathbf{F})$. By direct computation, given $0 \le t \le 1$, we have

$$\max\{\alpha \in \mathbb{R} : (\alpha, 0, t) \in W(\mathbf{F})\} = t^2 + (1 - t)^2 = 1 - 2t + 2t^2.$$

However for sufficient small t > 0

$$1 - t(1 - a) > 1 - t(1 - t^2/2) > (t/2)^2 + (1 - t/2)^2,$$

which contradicts that $(1 - t(1 - a), 0, t/2) \in W(\mathbf{F})$. Therefore, we have a = 0. However (0, 0, 1/2) is not a star-center as

$$\frac{1}{3}(0,0,1/2) + \frac{2}{3}(0,0,1) = \left(0,0,\frac{5}{6}\right) \notin \operatorname{conv}\left\{\left(\frac{1}{6},\frac{5}{6},\frac{5}{6}\right), \left(\frac{5}{6},-\frac{1}{6},\frac{5}{6}\right), \left(0,-\frac{2}{3},\frac{5}{6}\right)\right\}.$$

Therefore, $W(\mathbf{F})$ is not star-shaped.

It is challenging to determine conditions on \mathbf{F} so that $W(\mathbf{F})$ is star-shaped.

Acknowledgment

Our study was inspired by some discussion at the AIM workshop on Crouzeix's conjecture, July 30 - August 5, 2017, AIM, San Jose. The second and the third authors would like to express their thanks to the organizer and the colleagues at AIM for the well-organized and stimulating workshop.

Li is an honorary professor of the Shanghai University, and an affiliate member of the Institute for Quantum Computing, University of Waterloo; his research was supported by the USA NSF DMS 1331021, the Simons Foundation Grant 351047, and NNSF of China Grant 11571220. Research of Sze and Lau were supported by a PolyU central research grant G-YBKR and a HK RGC grant PolyU 502512. The HK RGC grant also supported the post-doctoral fellowship of Lau at the Hong Kong Polytechnic University.

References

- [1] Y. Au Yeung and Y.T. Poon, A remark on the convexity and positive definiteness concerning Hermitian matrices, Southeast Bull. Math. 3 (1979), 85-92.
- [2] Y. Au Yeung and N.K. Tsing, An extension of the Hausdorff-Toeplitz theorem on the numerical range, Proc. Amer. Math. Soc. 89 (1983), 215-218.
- [3] P. Binding and C. K. Li, Joint ranges of hermitian matrices and simultaneous diagonalization, Linear Algebra Appl. 151 (1991), 157-168.
- [4] W.S. Cheung and N.K. Tsing, The C-numerical range of matrices is star-shaped, Linear and Multilinear Algebra 41 (1996), 245-250.
- [5] M.T. Chien and H. Nakazato, Strict convexity of the joint c-numerical range, Linear Algebra Appl. 438 (2013), 1305-1321.
- [6] M.D. Choi, C.K. Li and Y.T. Poon, Some convexity features associated with unitary orbits, Canad. J. Math. 55 (2003), 91-111.
- [7] M. Crouzeix, Bounds for analytical functions of matrices, Integral Equations Operator Theory 48 (2004), 461477.

- [8] P.R. Halmos, A Hilbert Space Problem Book, Spring-Verlag, New York, 1982.
- [9] R.A. Horn and C.R. Johnson, Topics in Matrix Analysis, Cambridge University Press, Cambridge, 1991.
- [10] C.K. Li, C-numerical ranges and C-numerical radii, Linear and Multilinear Algebra 37 (1994), 51-82.
- [11] C.K. Li, D.C. Pelejo, Y.T. Poon and K.Z. Wang, Minkowski product of convex sets and product numerical range, Operators and Matrices, 10 (2016), 945-965.
- [12] C.K. Li and Y.T. Poon, Convexity of the joint numerical range, SIAM J. Matrix. Anal. Appl. 21 (2000), 668-678.
- [13] C.K. Li and N.K. Tsing, Matrices with circular symmetry on their unitary orbits and C-numerical ranges, Proc. of Amer. Math. Soc. 111 (1991), 19-28.
- [14] H. Nakazato, The C-numerical range of a 3×3 normal matrix, Nihonkai Math. J. 17 (2006), 187-197.
- [15] Z. Puchala, P. Gawron, J.A. Miszczak, L. Skowronek, M.D. Choi, and K. Zy-czkowski, Product numerical range in a space with tensor product structure, Linear Algebra Appl. 434 (2011), 327-342.
- [16] T. Schulte-Herbrüggen, G. Dirr, U.Helmke and S.J. Glaser The significance of the C-numerical range and the local C-numerical range in quantum control and quantum information, Linear and Multilinear Algebra 56 (2008), 3-26.
- [17] N.K. Tsing, The constraint bilinear form and the C-numerical range, Linear Algebra Appl. 56 (1984), 195-162.
- [18] R. Westwick, A theorem on numerical range, Linear and Multilinear Algebra 2 (1975), 311-315.

Addresses

- (P.S. Lau) Department of Applied Mathematics, The Hong Kong Polytechnic University, Hung Hom, Hong Kong. (panshun.lau@polyu.edu.hk)
- (C.K. Li) Department of Mathematics, College of William & Mary, Williamsburg, VA 23185. (ckli@math.wm.edu)
- (Y.T. Poon) Department of Mathematics, Iowa State University, Ames, IA 50011. (ytpoon@iastate.edu)
- (N.S. Sze) Department of Applied Mathematics, The Hong Kong Polytechnic University, Hung Hom, Hong Kong. (raymond.sze@polyu.edu.hk)