

# VOLTERRA–TYPE INTEGRAL OPERATOR ON ANALYTIC FUNCTION SPACES

R. KARGAR

ABSTRACT. In this note we study the radius of convexity (hence univalence) of the Volterra–type integral operator

$$T_g f(z) = \int_0^z f(s)g'(s)ds \quad (|z| < 1),$$

on the spaces of analytic function  $f$  in the open unit disc  $|z| < 1$ .

## 1. VOLTERRA–TYPE INTEGRAL OPERATOR

Let  $\Delta := \{z \in \mathbb{C} : |z| < 1\}$  and  $H(\Delta)$  be the class of all analytic functions in  $\Delta$ . In 1977 Pommerenke [16] introduced an integral operator, called Volterra–type operator as follows:

$$T_g := T_g f(z) = \int_0^z f(s)g'(s)ds \quad (z \in \Delta).$$

He proved that  $T_g$  is a bounded operator on the Hardy space  $H^2$  if, and only if,  $g$  belongs to the class  $BMOA$ . Also, Aleman and Siskakis [1] proved that this characterization (boundedness) is valid on each  $H^p$  for  $1 \leq p < \infty$  and that  $T_g$  is compact on  $H^p$  if, and only if,  $g \in VMOA$ . An another natural integral operator is defined as follows:

$$J_g f(z) := \int_0^z f'(s)g(s)ds \quad (z \in \Delta).$$

It is interesting to know this fact that

$$J_g f(z) + T_g f(z) = M_g f - f(0)g(0),$$

where  $M_g$  is the multiplication operator and is defined by

$$M_g f(z) = g(z)f(z) \quad (f \in H(\Delta), z \in \Delta).$$

Indeed, if  $f$  and  $g$  are two normalized analytic functions, i.e.  $f, g \in H(\Delta)$ ,  $f(0) = g(0) = 0$  and  $f'(0) = g'(0) = 1$ , then

$$J_g f(z) + T_g f(z) = g(z)f(z).$$

We note that the integral operators  $T_g$  and  $J_g$  contain the well-known integral operators in the analytic function theory and geometric function theory, such as the generalized Bernardi–Libera–Livingston linear integral operator (see [2, 7, 8]), Srivastava–Owa fractional derivative operators [11, 12] and the Cesàro integral operator, [18, 19].

Recently many researchers have been studied the integral operators  $T_g$  and  $J_g$ . For example, Li and Stević [6] studied the boundedness and the compactness of  $T_g$  and  $J_g$  on the Zygmund space and the little Zygmund space. We note that there exist some extensions of the integral operators  $T_g$ , see for example [20].

---

2010 *Mathematics Subject Classification.* 47B38; 30C45.

*Key words and phrases.* Starlike; Convex; Radius of convexity; Volterra–type integral operator; Linear–invariant family; Locally univalent; Boundary rotation.

The basis of this article is to study of the integral operator

$$(1.1) \quad T_g f(z) = \int_0^z f(s)g'(s)ds = \int_0^1 f(tz)zg'(tz)dt = \int_0^z fdg.$$

## 2. SOME SUBCLASSES OF ANALYTIC FUNCTIONS

In this section we recall some certain subclasses of analytic functions. Further, we denote by  $\mathcal{A}$  the class of all analytic and normalized functions in the open unit disc  $\Delta$ . The subclass of  $\mathcal{A}$  consisting of univalent (that is, one-to-one) functions is denoted by  $\mathcal{U}$ . We say that the function  $f \in \mathcal{U}$  is starlike of order  $\alpha$ ,  $0 \leq \alpha < 1$  if, and only if

$$\operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} \right\} > \alpha \quad (z \in \Delta).$$

We denote by  $\mathcal{S}^*(\alpha)$  the class of starlike functions of order  $\alpha$ . Also,  $f \in \mathcal{U}$  is convex of order  $\alpha$ ,  $0 \leq \alpha < 1$  iff  $zf'(z) \in \mathcal{S}^*(\alpha)$ . The class of convex functions of order  $\alpha$  is denoted by  $\mathcal{K}(\alpha)$ . Analytically,  $f \in \mathcal{K}(\alpha)$  iff

$$\operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} > \alpha \quad (z \in \Delta).$$

Let  $A$  and  $B$  be two complex numbers such that  $|A| > 1$  and  $|B| \leq 1$ . We say that a function  $f \in \mathcal{A}$  belongs to the class  $\mathcal{S}^*(A, B)$  if it satisfies the following subordination relation:

$$(2.1) \quad z \frac{f'(z)}{f(z)} \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta).$$

We note that for real numbers  $A$  and  $B$ , where  $-1 \leq B < A \leq 1$ , the function  $f$  that satisfies the above subordination relation (2.1) is called "the Janowski starlike function". Also, we say that  $f \in \mathcal{K}(A, B)$  if, and only if,  $zf'(z) \in \mathcal{S}^*(A, B)$ . Indeed, if  $f$  belongs to the class  $\mathcal{K}(A, B)$ , then it satisfies

$$1 + z \frac{f''(z)}{f'(z)} \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta).$$

We remark that  $\mathcal{K}(2, 1)$  and  $\mathcal{K}(2, -1)$  become to the Ozaki conditions. Moreover, by the Lindelöf subordination principle (this principle states that if  $f(z) \prec g(z)$ , then  $|f'(0)| \leq |g'(0)|$  and  $f(\Delta_r) \subset g(\Delta_r)$ ), if  $f \in \mathcal{K}(2, 1)$ , then we have

$$(2.2) \quad \operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} < \frac{3}{2} \quad (z \in \Delta).$$

Also, if  $f \in \mathcal{K}(2, -1)$ , then

$$(2.3) \quad \operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} > -\frac{1}{2} \quad (z \in \Delta).$$

Ozaki proved that if  $f \in \mathcal{A}$  with  $f(z)f'(z)/z \neq 0$ , there, and if either (2.2) or (2.3) holds throughout  $\Delta$ , then  $f$  is univalent and convex in at least one direction in  $\Delta$ , see [13].

Let  $\mathcal{LU}$  denote the family of normalized locally univalent functions in  $\Delta$ . For  $\beta \in \mathbb{R}$ , we consider the class  $\mathcal{G}(\beta)$  consisting of all functions  $f \in \mathcal{LU}$  which satisfy the condition

$$(2.4) \quad \operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} < 1 + \frac{\beta}{2} \quad (z \in \Delta).$$

We note that  $\mathcal{G}(1) \subset \mathcal{U}$  and  $\mathcal{G}(1) \equiv \mathcal{K}(2, 1)$ . Also the functions in the class  $\mathcal{G}(1)$  are starlike of order 0 in  $\Delta$ . The functions class  $\mathcal{G}(\beta)$  was studied extensively by Kargar *et al.* [5] (see also [10]).

Let  $Aut(\Delta)$  be the class of holomorphic automorphisms in  $\Delta$ . Any  $\phi \in Aut(\Delta)$  has the following representation:

$$\phi(z) = e^{i\vartheta} \frac{z+a}{1+\bar{a}z} \quad (\vartheta, a \in \mathbb{R}, z \in \Delta).$$

The family  $\mathcal{F}$  of  $\mathcal{A}$  is called a linear-invariant family (L.I.F.), if  $\mathcal{F} \subset \mathcal{LU}$  and for all  $f \in \mathcal{F}$  and  $\phi \in Aut(\Delta)$

$$F_\phi(f)(z) := \frac{f(\phi(z)) - f(\phi(0))}{f'(\phi(0))\phi'(0)} \in \mathcal{F}.$$

The class  $\mathcal{U}$  and the class  $\mathcal{K}(0)$  on  $\Delta$  are a L.I.F., but the class  $\mathcal{S}^*(0)$  on  $\Delta$  is not a L.I.F.. Also, we define the order of the L.I.F.  $\mathcal{F}$  as

$$\text{ord } f := \sup \left\{ \left| \frac{f''(0)}{2} \right| : f \in \mathcal{F} \right\}$$

and the universal linear invariant family of order  $\gamma \geq 1$  as

$$\mathcal{UL}_\gamma := \{f \in \mathcal{F} : \text{ord } f \leq \gamma\}.$$

We remark that  $\mathcal{UL}_1 \equiv \mathcal{K}(0)$  and  $\mathcal{U} \subset \mathcal{UL}_2$ . For more details about the L.I.F. see [4, Chapter 5].

For fixed  $k \geq 2$ , Let  $V_k$  denote the class of normalized functions  $f \in \mathcal{A}$  such that

$$\int_0^{2\pi} \left| \text{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} \right|_{z=re^{i\theta}} d\theta \leq k\pi.$$

Indeed, a function  $f \in V_k$  maps  $\Delta$  conformally onto an image domain  $f(\Delta)$  of boundary rotation at most  $k\pi$ . The functions class  $V_k$  was introduced by Loewner [9] in 1917 and was developed by Paatero [14] who systematically developed their properties and made an exhaustive study of the class  $V_k$ . In association with the class  $V_k$ , Robertson proved that (see [17, Theorem 1]) the following inequality

$$(2.5) \quad \left| z \frac{f''(z)}{f'(z)} - \frac{2|z|^2}{1-|z|^2} \right| \leq \frac{k|z|}{1-|z|^2} \quad (z \in \Delta, 2 \leq k < \infty)$$

holds true for each  $f \in V_k$ .

### 3. SOME KEY LEMMAS

In this section we recall some lemmas which help us in order to prove of main results.

**Lemma 3.1.** (see [3]) *If  $f \in \mathcal{UL}_\gamma$  and  $\gamma \geq 1$ , then*

$$\left| z \frac{f''(z)}{f'(z)} - \frac{2|z|^2}{1-|z|^2} \right| \leq \frac{2\gamma|z|}{1-|z|^2} \quad (z \in \Delta).$$

The next lemma is due to Pommerenke [15], see also [4, Lemma 5.1.3].

**Lemma 3.2.** *Let  $\mathcal{F}$  be a linear-invariant family and  $\delta = \text{ord } \mathcal{F}$ . Then*

$$(3.1) \quad \delta = \sup_{f \in \mathcal{F}} \sup_{|z| < 1} \left| -\bar{z} + \frac{1}{2}(1-|z|^2) \frac{f''(z)}{f'(z)} \right| \quad (z \in \Delta).$$

The following lemma gives a basic estimate which leads to the distortion theorem for univalent functions.

**Lemma 3.3.** (see [4, p. 15]) *If  $f \in \mathcal{U}$ , then*

$$\left| z \frac{f''(z)}{f'(z)} - \frac{2r^2}{1-r^2} \right| \leq \frac{4r}{1-r^2} \quad (|z| = r < 1).$$

*The estimate is sharp for rotation of Koebe function.*

**Lemma 3.4.** (see [10]) Let  $\beta \in (0, 1]$  be fixed. If  $f \in \mathcal{G}(\beta)$ , then

$$\left| \frac{f''(z)}{f'(z)} \right| \leq \frac{\beta}{1 - |z|} \quad (z \in \Delta).$$

The result is sharp for the function  $f'(z) = (1 - z)^\beta$ .

In the present paper, we obtain the radius of convexity of the Volterra-type integral operator  $T_g$  when the functions  $f$  and  $g$  belonging to the some certain subclasses of analytic functions which are defined in Section 2.

#### 4. MAIN RESULTS

We begin this section with the following result.

**Theorem 4.1.** Let  $A$  and  $B$  be two complex numbers,  $|A| > 1$  and  $|B| \leq 1$  and  $T_g$  be given by (1.1). Also, let  $0 \leq \alpha < 1$  be real number. If  $f \in \mathcal{S}^*(A, B)$  and  $g \in \mathcal{K}(A, B)$ , then the Volterra-type integral operator  $T_g$  is convex of order  $\alpha$  in  $|z| \leq r_c(A, B, \alpha)$ , where

$$r_c(A, B, \alpha) = \begin{cases} \frac{2-\alpha}{2|A|} & B = 0 \\ \frac{|B-A| - |(\alpha-1)B-A|}{\alpha|B|^2 - 2\operatorname{Re}\{A\bar{B}\}} & B \neq 0. \end{cases}$$

*Proof.* Using the definition of convexity of order  $\alpha$ , where  $0 \leq \alpha < 1$  it is enough to find out the largest number  $0 < r < 1$  for which

$$\min_{|z|=r} \operatorname{Re} \left\{ 1 + z \frac{T_g''(z)}{T_g'(z)} \right\} - \alpha \geq 0.$$

Further, from (1.1) we have

$$(4.1) \quad 1 + z \frac{T_g''(z)}{T_g'(z)} - \alpha = z \frac{f'(z)}{f(z)} - \alpha + z \frac{g''(z)}{g'(z)} + 1.$$

On the other hand, since  $f \in \mathcal{S}^*(A, B)$ , by the Lindelöf subordination principle, we get

$$(4.2) \quad \left| z \frac{f'(z)}{f(z)} - \frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right| \leq \frac{|B - A|r}{1 - |B|^2r^2} \quad (|z| = r < 1).$$

From the above inequality (4.2), we obtain

$$(4.3) \quad \begin{aligned} \operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} \right\} - \alpha &\geq \operatorname{Re} \left\{ \frac{1 - A\bar{B}r^2}{1 - |B|^2r^2} \right\} - \frac{|B - A|r}{1 - |B|^2r^2} - \alpha \\ &\geq \frac{1 - \alpha - |B - A|r - (\operatorname{Re}\{A\bar{B}\} - \alpha|B|^2) r^2}{1 - |B|^2r^2}. \end{aligned}$$

Also  $g \in \mathcal{K}(A, B)$  and the above same method imply that

$$(4.4) \quad \operatorname{Re} \left\{ 1 + z \frac{g''(z)}{g'(z)} \right\} \geq \frac{1 - |B - A|r - \operatorname{Re}\{A\bar{B}\}r^2}{1 - |B|^2r^2}.$$

Now from (4.1), (4.3) and (4.4) we get

$$\operatorname{Re} \left\{ 1 + z \frac{T_g''(z)}{T_g'(z)} \right\} - \alpha \geq \frac{2 - \alpha - 2|B - A|r - (2\operatorname{Re}\{A\bar{B}\} - \alpha|B|^2) r^2}{1 - |B|^2r^2} > 0$$

provided  $\phi(r) := 2 - \alpha - 2|B - A|r - (2\operatorname{Re}\{A\bar{B}\} - \alpha|B|^2) r^2 > 0$ . A simple calculation gives that the roots of  $\phi(r)$  are

$$\frac{|B - A| \pm |(\alpha - 1)B - A|}{\alpha|B|^2 - 2\operatorname{Re}\{A\bar{B}\}}.$$

Also, it is clear that if  $B = 0$ , then  $\phi(r)$  yields that  $\phi(r) = 2 - \alpha - 2|A|r$  and it will be positive if  $r < (2 - \alpha)/2|A|$ . This completes the proof.  $\square$

Putting  $A = 2$  and  $B = 1$  in Theorem 4.1, we get.

**Corollary 4.1.** *Let  $\operatorname{Re}\{zf'(z)/f(z)\} < 3/2$  where  $z \in \Delta$  and  $g$  satisfies the condition (2.2). Then the radius of convexity of order  $\alpha$  of the Volterra-type integral operator  $T_g$  is  $\frac{\alpha-2}{\alpha-4}$  where  $(0 \leq \alpha < 1)$ .*

If we take  $A = 2$  and  $B = -1$  in Theorem 4.1, then we have the following.

**Corollary 4.2.** *Let  $\operatorname{Re}\{zf'(z)/f(z)\} > -1/2$  where  $z \in \Delta$  and  $g$  satisfies the condition (2.3). Then the Volterra-type integral operator  $T_g$  is convex of order  $\alpha$   $(0 \leq \alpha < 1)$  in the disc  $|z| \leq (2 - \alpha)/(4 + \alpha)$ .*

**Theorem 4.2.** *Let  $0 \leq \alpha < 1$  and  $\gamma \geq 1$ . If  $f \in \mathcal{S}^*(\alpha)$  and  $g \in \mathcal{LU}_\gamma$ , then the radius of convexity of the Volterra-type integral operator  $T_g$  is*

$$(4.5) \quad r_c^-(\alpha, \gamma) = \frac{\gamma - \sqrt{\alpha^2 + \gamma^2 - 1}}{1 - \alpha}.$$

*Proof.* Let  $f \in \mathcal{S}^*(\alpha)$   $(0 \leq \alpha < 1)$  and  $g$  belongs to the family  $\mathcal{LU}_\gamma$   $(\gamma \geq 1)$ . From (1.1) we get

$$(4.6) \quad 1 + z \frac{T_g''(z)}{T_g'(z)} = z \frac{f'(z)}{f(z)} + z \frac{g''(z)}{g'(z)} + 1.$$

Since  $g \in \mathcal{LU}_\gamma$ , Lemma 3.1 implies that

$$(4.7) \quad \operatorname{Re} \left\{ z \frac{g''(z)}{g'(z)} \right\} \geq \frac{2r^2}{1 - r^2} - \frac{2\gamma r}{1 - r^2}.$$

Now by definition of starlikeness of order  $\alpha$  and using the above inequality (4.7), and also applying the relation (4.6), we obtain

$$\begin{aligned} \operatorname{Re} \left\{ 1 + z \frac{T_g''(z)}{T_g'(z)} \right\} &= \operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} + z \frac{g''(z)}{g'(z)} + 1 \right\} \\ &> \frac{1 + \alpha - 2\gamma r + (1 - \alpha)r^2}{1 - r^2} > 0, \end{aligned}$$

provided  $\varphi(r) = 1 + \alpha - 2\gamma r + (1 - \alpha)r^2 > 0$ . It is easy to see that the roots of  $\varphi(r)$  are

$$r_c^\pm(\alpha, \gamma) = \frac{\gamma \pm \sqrt{\alpha^2 + \gamma^2 - 1}}{1 - \alpha}.$$

Also we see that if  $0 \leq \alpha < 1$  and  $\gamma \geq 1$ , then  $0 < r_c^-(\alpha, \gamma) < 1$  where  $r_c^-(\alpha, \gamma)$  is defined in (4.5). This is the end of proof.  $\square$

Putting  $\alpha = 0$  and  $\gamma = 1$  in Theorem 4.2, we get.

**Corollary 4.3.** *Let  $f$  and  $g$  be starlike and convex univalent functions in the open unit disc  $\Delta$ , respectively. Then the Volterra-type integral operator  $T_g$  is convex univalent function in  $\Delta$ , too.*

**Theorem 4.3.** *Let  $0 \leq \alpha < 1$ . If  $f$  is starlike function of order  $\alpha$  and  $g \in \mathcal{F}$  with  $\operatorname{ord}\mathcal{F} = 1$ , then the Volterra-type integral operator  $T_g$  is convex univalent in the open unit disc  $\Delta$ .*

*Proof.* Since  $f$  is starlike function of order  $\alpha$ , we have

$$(4.8) \quad \operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} \right\} > \alpha \quad (z \in \Delta).$$

Also since  $g \in \mathcal{F}$ , by using the relation (3.1) and with a little calculation we get

$$(4.9) \quad \operatorname{Re} \left\{ 1 + z \frac{g''(z)}{g'(z)} \right\} \geq \frac{1 - 2\delta r + r^2}{1 - r^2} \quad (|z| = r < 1).$$

Now, from (4.6), (4.8) and (4.9) and since  $\operatorname{ord} \mathcal{F} = 1$ , we get

$$\operatorname{Re} \left\{ 1 + z \frac{T_g''(z)}{T_g'(z)} \right\} \geq \frac{1 + \alpha - 2r + (1 - \alpha)r^2}{1 - r^2}$$

and for  $z \in \Delta$  one deduces that

$$\operatorname{Re} \left\{ 1 + z \frac{T_g''(z)}{T_g'(z)} \right\} > 0 \quad (z \in \Delta).$$

Thus by definition we conclude that  $T_g$  is convex univalent in the open unit disc  $\Delta$ . This is the end of proof.  $\square$

The next theorem is the following.

**Theorem 4.4.** *Let  $f$  be starlike function of order  $\alpha$  where  $0 \leq \alpha < 1$  and  $g$  be univalent function. Then the radius of convexity of the Volterra-type integral operator  $T_g$  is*

$$r_c^-(\alpha) = \frac{2 - \sqrt{3 + \alpha^2}}{1 - \alpha} \quad (0 \leq \alpha < 1).$$

*Proof.* Let  $f \in \mathcal{S}^*(\alpha)$  where  $0 \leq \alpha < 1$  and  $g \in \mathcal{U}$ . From Lemma 3.3 and by (4.6), we get

$$\begin{aligned} \operatorname{Re} \left\{ 1 + z \frac{T_g''(z)}{T_g'(z)} \right\} &= \operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} + z \frac{g''(z)}{g'(z)} + 1 \right\} \\ &> \frac{1 + \alpha - 4r + (1 - \alpha)r^2}{1 - r^2} > 0, \end{aligned}$$

when  $|z| \leq (2 - \sqrt{3 + \alpha^2})/(1 - \alpha)$  and concluding the proof.  $\square$

*Remark 4.1.* Taking  $\alpha = 0$  in Theorem 4.4, we see that if  $f$  is starlike univalent function and  $g \in \mathcal{U}$ , then the radius of convexity of the Volterra-type integral operator  $T_g$  is  $2 - \sqrt{3}$ . Indeed, in this case the radius of convexity the Volterra-type integral operator  $T_g$  is equal to familiar radius of convexity for the class  $\mathcal{U}$  (see [4, Theorem 2.2.22]).

**Theorem 4.5.** *Let  $f$  be starlike function of order  $\alpha$  where  $0 \leq \alpha < 1$  and  $g$  be locally univalent function which satisfies the inequality (2.4) where  $0 < \beta \leq 1$ . Then the radius of convexity of the Volterra-type integral operator  $T_g$  is*

$$r_c(\alpha, \beta) = \frac{1 + \alpha}{1 + \alpha + \beta}.$$

*Proof.* Assume that  $f \in \mathcal{S}^*(\alpha)$  and  $g \in \mathcal{G}(\beta)$ , where  $0 \leq \alpha < 1$  and  $0 < \beta \leq 1$ . By Lemma 3.4 and from (4.6) we have

$$\operatorname{Re} \left\{ 1 + z \frac{T_g''(z)}{T_g'(z)} \right\} > \alpha + 1 - \frac{\beta|z|}{1 - |z|} \geq 0,$$

where  $|z| \leq (1 + \alpha)/(1 + \alpha + \beta)$ . This completes the proof.  $\square$

Finally we have.

**Theorem 4.6.** *Let  $f$  be starlike function of order  $\alpha$  where  $0 \leq \alpha < 1$  and  $g \in V_k$  where  $k \geq 2$ . Then the radius of convexity of the Volterra-type integral operator  $T_g$  is*

$$r_c(\alpha, k) = \frac{k - \sqrt{k^2 - 4(1 - \alpha^2)}}{2(1 - \alpha)}.$$

*Proof.* From the definition of starlikeness of order  $\alpha$ , inequality (2.5) and (4.6), one deduces that

$$\begin{aligned} \operatorname{Re} \left\{ 1 + z \frac{T_g''(z)}{T_g'(z)} \right\} &= \operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} + z \frac{g''(z)}{g'(z)} + 1 \right\} \\ &> \frac{(1-\alpha)r^2 - kr + 1 + \alpha}{1-r^2} = \frac{\lambda(r)}{1-r^2} \end{aligned}$$

where

$$\lambda(r) := (1-\alpha)r^2 - kr + 1 + \alpha \quad (k \geq 2, 0 \leq \alpha < 1).$$

It is easy to see that  $\lambda(r) > 0$  if

$$r > \frac{k + \sqrt{k^2 - 4(1-\alpha^2)}}{2(1-\alpha)} \geq 1 \quad (k \geq 2, 0 \leq \alpha < 1)$$

or

$$r < \frac{k - \sqrt{k^2 - 4(1-\alpha^2)}}{2(1-\alpha)} \leq 1 \quad (k \geq 2, 0 \leq \alpha < 1).$$

This completes the proof.  $\square$

#### REFERENCES

- [1] A. Aleman and A. Siskakis, *An integral operator on  $H^p$* , Complex Var. Theory Appl. **28** (1995), 140–158.
- [2] S.D. Bernardi, *Convex and starlike univalent functions*, Trans. Amer. Math. Soc. **135** (1969), 429–446.
- [3] A. Ebadian and R. Kargar, *Univalence of integral operators on neighborhoods of analytic functions*, Iran. J. Sci. Technol. Trans. Sci. **42** (2018), 911–915.
- [4] I. Graham and G. Kohr, *Geometric Function Theory in One and Higher Dimensions*, Marcel Dekker Inc., New York, 2003.
- [5] R. Kargar, N.R. Pascu and A. Ebadian, *Locally univalent approximations of analytic functions*, J. Math. Anal. Appl. **453** (2017), 1005–1021.
- [6] S. Li and S. Stević, *Volterra-type operators on Zygmund spaces*, J. Ineq. Appl., Vol. 2007, Article ID 32124, 10 pages, 2007.
- [7] R.J. Libera, *Some classes of regular univalent functions*, Proc. Amer. Math. Soc. **16** (1965), 755–758.
- [8] A.E. Livingston, *On the radius of univalence of certain analytic functions*, Proc. Amer. Math. Soc. **17** (1966) 352–357.
- [9] C. Loewner, *Untersuchungen über die Verzerrung bei konformen Abbildungen des Einheitskreises  $|z| < 1$ , die durch Funktionen mit nicht verschwindender Ableitung geliefert werden*, Ber. Verh. Sächs. Gess. Wiss. Leipzig, **69** (1917), 89–106.
- [10] M. Obradović, S. Ponnusamy, and K.-J. Wirths, *Coefficient characterizations and sections for some univalent functions*, Sib. Math. J. **54** (2013), 679–696.
- [11] S. Owa, *On the distortion theorems I*, Kyungpook Math. J. **18** (1978), 53–59.
- [12] S. Owa and H.M. Srivastava, *Univalent and starlike generalized hypergeometric functions*, Can. J. Math. **39** (1987), 1057–1077.
- [13] S. Ozaki, *On the theory of multivalent functions*, Sci. Rep. Tokyo Bunrika Daigaku, **4** (1941), 45–86.
- [14] V. Paatero, *Über die konforme Abbildung von Gebieten deren Ränder von beschränkter Drehung sind*, Ann. Acad. Sci. Fenn. Ser. A **33** (1931), 1–77.
- [15] C. Pommerenke, *Linear-invariante familien analytischer funktionen. I*, Math. Ann. **155** (1964), 108–154.
- [16] Ch. Pommerenke, *Schlichte Funktionen und analytische Funktionen von beschränkter mittlerer Oszillation*, Comment. Math. Helv. **52** (1977), 591–602.
- [17] M.S. Robertson, *Coefficients of functions with bounded boundary rotation*, Canad. J. Math. **21** (1969), 1477–1482.
- [18] A.G. Siskakis, *Composition semigroups and the Cesáro operator on  $H^p$* , J. Lond. Math. Soc. **36** (1987), 153–164.
- [19] A.G. Siskakis, *The Cesáro operator is bounded on  $H^1$* , Proc. Amer. Math. Soc. **110** (1990), 461–462.
- [20] S. Stević *On a new integral-type operator from the Bloch space to Bloch-type spaces on the unit ball*, J. Math. Anal. Appl. **354** (2009) 426–434.

YOUNG RESEARCHERS AND ELITE CLUB, ARDABIL BRANCH, ISLAMIC AZAD UNIVERSITY, ARD-  
ABIL, IRAN.

*E-mail address:* [rkargar1983@gmail.com](mailto:rkargar1983@gmail.com)