

VOLTERRA TYPE INTEGRAL OPERATOR AND ANALYTIC FUNCTION SPACES

RAHIM KARGAR

ABSTRACT. We obtain the radius of convexity of the Volterra type integral operator

$$T_g f(z) = \int_0^z f(s)g'(s)ds \quad (|z| < 1),$$

when f and g belong to the some certain subclass of analytic functions.

1. VOLTERRA TYPE OPERATOR

Let $\Delta := \{z \in \mathbb{C} : |z| < 1\}$. In 1977 Pommerenke [17, Lemma 1] introduced an integral operator, called Volterra type operator as follows:

$$(1.1) \quad J_g f(z) := \int_0^z f'(s)g(s)ds \quad (z \in \Delta).$$

Pommerenke proved that $J_g f(z)$ is a bounded operator on the Hardy space H^2 if and only if g belongs to the class $BMOA$. Also Aleman and Siskakis [1] proved that this characterization (boundedness) is valid on each H^p for $1 \leq p < \infty$ and that J_g is compact on H^p if and only if $g \in VMOA$. An another natural integral operator is defined as follows:

$$T_g f(z) := \int_0^z f(s)g'(s)ds \quad (z \in \Delta).$$

It is necessary to refer to the this fact that

$$J_g f(z) + T_g f(z) = M_g f - f(0)g(0),$$

where M_g is the multiplication operator and is defined by

$$(M_g f)(z) = g(z)f(z) \quad (f \in H(\Delta), z \in \Delta)$$

and $H(\Delta)$ denotes the class of all analytic functions on Δ . Indeed, if f and g are two normalized analytic functions, then

$$J_g f(z) + T_g f(z) = g(z)f(z).$$

We note that the integral operators $J_g f(z)$ and $T_g f(z)$ contain the well-known integral operators in the analytic function theory and geometric function theory, such as the generalized Bernardi–Libera–Livingston linear integral operator (see [2, 9, 10]), Srivastava–Owa fractional derivative operators [12, 14] and the Cesàro integral operator, [18, 19].

Recently many researchers have been studied the integral operators $J_g f(z)$ and $T_g f(z)$. For example, Li and Stević [7] studied the boundedness and the compactness of $J_g f(z)$ and $T_g f(z)$ on the Zygmund space and the little Zygmund space. Also, many authors have been studied the essential norm of the integral operators. Laitila *et al.* [6] studied the essential norm of the operator T_g on the Hardy space. Or Liu *et al.* (see [8]) studied the essential norm of the operator T_g on the Bloch

2010 *Mathematics Subject Classification.* 47B38; 30C45.

Key words and phrases. Starlike; Convex; Radius of convexity; Volterra type integral operator; Linear-invariant family; Locally univalent.

space and some other spaces. Zhuo and Ye [20] studied the essential norm of the operator T_g from Morrey spaces to the Bloch space. Finally, Zhou and Zhu studied the essential norm of the operator T_g from Hardy spaces to the BMOA space, Besov spaces, Bergman spaces and Bloch-type spaces, see [21].

The basis of this article is to study of the integral operator

$$(1.2) \quad T_g f(z) = \int_0^z f(s)g'(s)ds = \int_0^1 f(tz)zg'(tz)dt = \int_0^z f dg.$$

2. SOME SUBCLASS OF ANALYTIC FUNCTIONS

In this section we recall some certain subclass of analytic functions. Further, we denote by \mathcal{A} the class of all analytic and normalized functions in the open unit disk Δ . The subclass of \mathcal{A} consisting of univalent (that is, one-to-one) functions is denoted by \mathcal{U} . We say that the function $f \in \mathcal{U}$ is starlike of order $0 \leq \alpha < 1$ if, and only if

$$\operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} \right\} > \alpha \quad (z \in \Delta).$$

We denote by $\mathcal{S}^*(\alpha)$ the class of starlike functions of order α . Also, $f \in \mathcal{U}$ is convex of order $0 \leq \alpha < 1$ iff $zf'(z) \in \mathcal{S}^*(\alpha)$. The class of convex functions of order α is denoted by $\mathcal{K}(\alpha)$. Analytically, $f \in \mathcal{K}(\alpha)$ iff

$$\operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} > \alpha \quad (z \in \Delta).$$

Let A and B be two complex numbers such that $|A| > 1$ and $|B| \leq 1$. We say that the function $f \in \mathcal{A}$ belongs to the class $\mathcal{S}^*(A, B)$ if it satisfies the following subordination relation:

$$z \frac{f'(z)}{f(z)} \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta).$$

We note that if $-1 \leq B < A \leq 1$ and are real, then $\mathcal{S}^*(A, B)$ becomes the family of Janowski starlike functions. Also, we say that $f \in \mathcal{K}(A, B)$ if and only if $zf'(z) \in \mathcal{S}^*(A, B)$. Indeed, if f belongs to the class $\mathcal{K}(A, B)$, then it satisfies

$$1 + z \frac{f''(z)}{f'(z)} \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta).$$

We remark that $\mathcal{K}(2, 1)$ and $\mathcal{K}(2, -1)$ become to the Ozaki conditions. Moreover, by the Lindelöf subordination principle (this principle states that if $f(z) \prec g(z)$, then $|f'(0)| \leq |g'(0)|$ and $f(\Delta_r) \subset g(\Delta_r)$), if $f \in \mathcal{K}(2, 1)$, then we have

$$(2.1) \quad \operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} < \frac{3}{2} \quad (z \in \Delta).$$

Also, if $f \in \mathcal{K}(2, -1)$, then

$$(2.2) \quad \operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} > -\frac{1}{2} \quad (z \in \Delta).$$

Ozaki proved that if $f \in \mathcal{A}$ with $f(z)f'(z)/z \neq 0$, there, and if either (2.1) or (2.2) holds throughout Δ , then f is univalent and convex in at least one direction in Δ , see [15].

Let \mathcal{LU} denote the family of normalized locally univalent functions in Δ . For $\beta \in \mathbb{R}$, we consider the class $\mathcal{G}(\beta)$ consisting of all functions $f \in \mathcal{LU}$ which satisfy the condition

$$(2.3) \quad \operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} < 1 + \frac{\beta}{2} \quad (z \in \Delta).$$

We note that $\mathcal{G}(1) \subset \mathcal{U}$ and $\mathcal{G}(1) \equiv \mathcal{K}(2, 1)$. Also the functions in the class $\mathcal{G}(1)$ are starlike of order zero in Δ . The functions class $\mathcal{G}(\beta)$ was studied extensively by Kargar *et al.* [5] (see also [11]). For more details about the class $\mathcal{G}(\beta)$ see [5] and its references.

Let $Aut(\Delta)$ be the class of holomorphic automorphisms in Δ . Any $\phi \in Aut(\Delta)$ has the following representation:

$$\phi(z) = e^{i\vartheta} \frac{z+a}{1+\bar{a}z} \quad (\vartheta, a \in \mathbb{R}, z \in \Delta).$$

The family \mathcal{F} of \mathcal{A} is called a linear-invariant family (L.I.F.), if $\mathcal{F} \subset \mathcal{LU}$ and for all $f \in \mathcal{F}$ and $\phi \in Aut(\Delta)$

$$(2.4) \quad F_\phi(f)(z) := \frac{f(\phi(z)) - f(\phi(0))}{f'(\phi(0))\phi'(0)} \in \mathcal{F}.$$

The class \mathcal{U} and the class $\mathcal{K}(0)$ on Δ are a L.I.F., but the class $\mathcal{S}^*(0)$ on Δ is not a L.I.F.. Also, we define the order of the L.I.F. \mathcal{F} as

$$\text{ord}f := \sup \left\{ \left| \frac{f''(0)}{2} \right| : f \in \mathcal{F} \right\}$$

and the universal linear invariant family of order $\gamma \geq 1$ as

$$\mathcal{UL}_\gamma := \{f \in \mathcal{F} : \text{ord}f \leq \gamma\}.$$

We remark that $\mathcal{UL}_1 \equiv \mathcal{K}(0)$ and $\mathcal{U} \subset \mathcal{UL}_2$. For more details about the L.I.F. see [4, Chapter 5].

3. SOME KEY LEMMAS

In this section we recall some lemmas which help us in order to prove of main results.

Lemma 3.1. (see [3]) *If $f \in \mathcal{UL}_\gamma$ and $\gamma \geq 1$, then*

$$\left| \frac{zf''(z)}{f'(z)} - \frac{2|z|^2}{1-|z|^2} \right| \leq \frac{2\gamma|z|}{1-|z|^2} \quad (z \in \Delta).$$

The next lemma is due to Pommerenke [16], see also [4, Lemma 5.1.3].

Lemma 3.2. *Let \mathcal{F} be a linear-invariant family and $\delta = \text{ord}\mathcal{F}$. Then*

$$(3.1) \quad \delta = \sup_{f \in \mathcal{F}} \sup_{|z| < 1} \left| -\bar{z} + \frac{1}{2}(1-|z|^2) \frac{f''(z)}{f'(z)} \right| \quad (z \in \Delta).$$

The following lemma gives a basic estimate which leads to the distortion theorem for univalent functions.

Lemma 3.3. (see [4, p. 15]) *If $f \in \mathcal{U}$, then*

$$\left| \frac{zf''(z)}{f'(z)} - \frac{2r^2}{1-r^2} \right| \leq \frac{4r}{1-r^2} \quad (|z| = r < 1).$$

The estimate is sharp for rotation of Koebe function.

Lemma 3.4. (see [11]) *Let $\beta \in (0, 1]$ be fixed. If $f \in \mathcal{G}(\beta)$, then*

$$\left| \frac{f''(z)}{f'(z)} \right| \leq \frac{\beta}{1-|z|} \quad (z \in \Delta).$$

The result is sharp for the function $f'(z) = (1-z)^\beta$.

In the present paper, we obtain the radius of convexity of the Volterra-type integral operator $T_g f(z)$ given by (1.2) when the functions f and g belonging to the some certain subclasses of analytic functions which are defined in the Section 2.

4. MAIN RESULTS

We begin this section with the following result.

Theorem 4.1. *Let A and B be two complex numbers, $|A| > 1$ and $|B| \leq 1$. Also, let $0 \leq \alpha < 1$ be real number. If $f \in \mathcal{S}^*(A, B)$ and $g \in \mathcal{K}(A, B)$, then the Volterra-type integral operator $T_g f(z)$ given by (1.2) is convex of order α in $|z| \leq r_c(A, B, \alpha)$ where*

$$(4.1) \quad r_c(A, B, \alpha) = \begin{cases} \frac{2-\alpha}{2|A|} & B = 0 \\ \frac{|B-A| - |(\alpha-1)B-A|}{\alpha|B|^2 - 2\operatorname{Re}\{AB\}} & B \neq 0. \end{cases}$$

Proof. From now on, for convenience we put $T(z) := T_g f(z)$. By the analytic definition of convexity of order $0 \leq \alpha < 1$ it is enough to find out the largest number $0 < r < 1$ for which

$$(4.2) \quad \min_{|z|=r} \operatorname{Re} \left\{ 1 + z \frac{T''(z)}{T'(z)} \right\} - \alpha \geq 0.$$

Further, from (1.2) we have

$$(4.3) \quad 1 + z \frac{T''(z)}{T'(z)} - \alpha = z \frac{f'(z)}{f(z)} - \alpha + z \frac{g''(z)}{g'(z)} + 1.$$

On the other hand, since $f \in \mathcal{S}^*(A, B)$ we get

$$(4.4) \quad z \frac{f'(z)}{f(z)} \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta).$$

Thus by using the Lindelöf subordination principle, we get

$$(4.5) \quad \left| z \frac{f'(z)}{f(z)} - \frac{1 - A\bar{B}r^2}{1 - |B|^2 r^2} \right| \leq \frac{|B - A|r}{1 - |B|^2 r^2} \quad (|z| = r < 1).$$

From the above inequality (4.5), we obtain

$$(4.6) \quad \begin{aligned} \operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} \right\} - \alpha &\geq \operatorname{Re} \left\{ \frac{1 - A\bar{B}r^2}{1 - |B|^2 r^2} \right\} - \frac{|B - A|r}{1 - |B|^2 r^2} - \alpha \\ &\geq \frac{1 - \alpha - |B - A|r - (\operatorname{Re}\{A\bar{B}\} - \alpha|B|^2) r^2}{1 - |B|^2 r^2}. \end{aligned}$$

Also, because $g \in \mathcal{K}(A, B)$, by the same proof we obtain

$$(4.7) \quad \operatorname{Re} \left\{ 1 + z \frac{g''(z)}{g'(z)} \right\} \geq \frac{1 - |B - A|r - \operatorname{Re}\{A\bar{B}\} r^2}{1 - |B|^2 r^2}.$$

Now from (4.3), (4.6) and (4.7) we get

$$\operatorname{Re} \left\{ 1 + z \frac{T''(z)}{T'(z)} \right\} - \alpha \geq \frac{2 - \alpha - 2|B - A|r - (2\operatorname{Re}\{A\bar{B}\} - \alpha|B|^2) r^2}{1 - |B|^2 r^2} > 0$$

provided $\phi(r) := 2 - \alpha - 2|B - A|r - (2\operatorname{Re}\{A\bar{B}\} - \alpha|B|^2) r^2 > 0$. A simple calculation gives that the roots of $\phi(r)$ are

$$(4.8) \quad \frac{|B - A| \pm |(\alpha - 1)B - A|}{\alpha|B|^2 - 2\operatorname{Re}\{A\bar{B}\}}.$$

Also, it is clear that if $B = 0$, then $\phi(r)$ yields that $\phi(r) = 2 - \alpha - 2|A|r$ and it will be positive if $r < (2 - \alpha)/2|A|$. This completes the proof. \square

If we take $A = 2$ and $B = -1$ in the Theorem 4.1, then we have the following.

Corollary 4.1. *Let $\operatorname{Re}\{z f'(z)/f(z)\} > -1/2$ where $z \in \Delta$ and g satisfies the condition (2.2). Then the Volterra-type integral operator $T_g f(z)$ given by (1.2) is convex of order α ($0 \leq \alpha < 1$) in the disk $|z| \leq (2 - \alpha)/(4 + \alpha)$.*

Putting $A = 2$ and $B = 1$ in the Theorem 4.1, we get.

Corollary 4.2. *Let $\operatorname{Re}\{zf'(z)/f(z)\} < 3/2$ where $z \in \Delta$ and g satisfies the condition (2.1). Then the radius of convexity of order α of the Volterra-type integral operator $T_g f(z)$ given by (1.2) is $\frac{\alpha-2}{\alpha-4}$ where $(0 \leq \alpha < 1)$.*

Theorem 4.2. *Let $0 \leq \alpha < 1$ and $\gamma \geq 1$. If $f \in \mathcal{S}^*(\alpha)$ and $g \in \mathcal{LU}_\gamma$, then the radius of convexity of the Volterra-type integral operator $T_g f(z)$ given by (1.2) is*

$$(4.9) \quad r_c^-(\alpha, \gamma) = \frac{\gamma - \sqrt{\alpha^2 + \gamma^2 - 1}}{2 - (\alpha + 1)}.$$

Proof. Let $f \in \mathcal{S}^*(\alpha)$ ($0 \leq \alpha < 1$) and g belongs to the family \mathcal{LU}_γ ($\gamma \geq 1$). From (1.2) we get

$$(4.10) \quad 1 + z \frac{T''(z)}{T'(z)} = z \frac{f'(z)}{f(z)} + z \frac{g''(z)}{g'(z)} + 1.$$

Since $g \in \mathcal{LU}_\gamma$, the Lemma 3.1 implies that

$$(4.11) \quad \operatorname{Re} \left\{ z \frac{g''(z)}{g'(z)} \right\} \geq \frac{2r^2}{1-r^2} - \frac{2\gamma r}{1-r^2}.$$

Now by definition of starlikeness of order α and using the above inequality (4.11), and applying the relation (4.10) we obtain

$$\begin{aligned} \operatorname{Re} \left\{ 1 + z \frac{T''(z)}{T'(z)} \right\} &= \operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} + z \frac{g''(z)}{g'(z)} + 1 \right\} \\ &> \frac{1 + \alpha - 2\gamma r + (2 - \alpha - 1)r^2}{1 - r^2} > 0, \end{aligned}$$

provided $\varphi(r) = 1 + \alpha - 2\gamma r + (2 - \alpha - 1)r^2 > 0$. It is easy to see that the roots of $\varphi(r)$ are

$$r_c^\pm(\alpha, \gamma) = \frac{\gamma \pm \sqrt{\alpha^2 + \gamma^2 - 1}}{2 - (\alpha + 1)}.$$

Also we see that if $0 \leq \alpha < 1$ and $\gamma \geq 1$, then $0 < r_c^-(\alpha, \gamma) < 1$ where $r_c(\alpha, \gamma)$ defined in (4.9). This is the end of proof. \square

Putting $\alpha = 0$ and $\gamma = 1$ in the Theorem 4.2, we get.

Corollary 4.3. *Let f and g be starlike and convex univalent functions in the open unit disk Δ , respectively. Then the Volterra-type integral operator $T_g f(z)$ is convex univalent function in Δ , too.*

Theorem 4.3. *Let $0 \leq \alpha < 1$. If f is starlike of order α and $g \in \mathcal{F}$ with $\operatorname{ord}\mathcal{F} = 1$, then the Volterra type integral operator $T_g f(z)$ given by (1.2) is convex univalent in the disk Δ .*

Proof. Since f is starlike of order $0 \leq \alpha < 1$ we have

$$(4.12) \quad \operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} \right\} > \alpha \quad (z \in \Delta).$$

Also since $g \in \mathcal{F}$, by using the relation (3.1) and with a little calculation we get

$$(4.13) \quad \operatorname{Re} \left\{ 1 + z \frac{g''(z)}{g'(z)} \right\} \geq \frac{1 - 2\delta r + r^2}{1 - r^2} \quad (|z| = r < 1).$$

Now, from (4.10), (4.12) and (4.13) and since $\operatorname{ord}\mathcal{F} = 1$, we get

$$\operatorname{Re} \left\{ 1 + z \frac{T''(z)}{T'(z)} \right\} \geq \frac{1 + \alpha - 2r + (1 - \alpha)r^2}{1 - r^2}$$

and for $z \in \Delta$ one deduces that

$$\operatorname{Re} \left\{ 1 + z \frac{T''(z)}{T'(z)} \right\} > 0 \quad (z \in \Delta).$$

Thus by definition we conclude that $T(z)$ is convex univalent in the unit disk Δ . This is the end of proof. \square

The next theorem is the following.

Theorem 4.4. *Let f be starlike function of order α and g be univalent function. Then the radius of convexity of the Volterra-type integral operator $T_g f(z)$ is*

$$(4.14) \quad r_c^-(\alpha) = \frac{2 - \sqrt{3 + \alpha^2}}{1 - \alpha} \quad (0 \leq \alpha < 1).$$

Proof. Let $f \in \mathcal{S}^*(\alpha)$ where $0 \leq \alpha < 1$ and $g \in \mathcal{U}$. From Lemma 3.3 and by (4.10), we get

$$\begin{aligned} \operatorname{Re} \left\{ 1 + z \frac{T''(z)}{T'(z)} \right\} &= \operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} + z \frac{g''(z)}{g'(z)} + 1 \right\} \\ &> \frac{1 + \alpha - 4r + (1 - \alpha)r^2}{1 - r^2} > 0, \end{aligned}$$

when $|z| \leq (2 - \sqrt{3 + \alpha^2})/(1 - \alpha)$ and concluding the proof. \square

Remark 4.1. Taking $\alpha = 0$ in the Theorem 4.4, we see that if f is starlike univalent function and $g \in \mathcal{U}$, then the radius of convexity of the Volterra type integral operator is $2 - \sqrt{3}$. Indeed, in this case the radius of convexity the Volterra type integral operator is equal to familiar radius of convexity for the class \mathcal{U} (see [4, Theorem 2.2.22, p. 51]).

Finally we have.

Theorem 4.5. *Let f be starlike function of order $0 \leq \alpha < 1$ and g be locally univalent function which satisfies in (2.3) where $0 < \beta \leq 1$. Then the radius of convexity of the Volterra-type integral operator $T_g f(z)$ is*

$$(4.15) \quad r_c(\alpha, \beta) = \frac{1 + \alpha}{1 + \alpha + \beta}.$$

Proof. Assume that $f \in \mathcal{S}^*(\alpha)$ and $g \in \mathcal{G}(\beta)$, where $0 \leq \alpha < 1$ and $0 < \beta \leq 1$. By Lemma 3.4 and from (4.10) we have

$$\operatorname{Re} \left\{ 1 + z \frac{T''(z)}{T'(z)} \right\} > \alpha + 1 - \frac{\beta|z|}{1 - |z|} \geq 0,$$

where $|z| \leq (1 + \alpha)/(1 + \alpha + \beta)$. This completes the proof. \square

REFERENCES

- [1] A. Aleman and A. Siskakis, *An integral operator on H^p* , Complex Var. Theory Appl. **28** (1995), 140–158.
- [2] S.D. Bernardi, *Convex and starlike univalent functions*, Trans. Amer. Math. Soc. **135** (1969), 429–446.
- [3] A. Ebadian and R. Kargar, *Univalence of integral operators on neighborhoods of analytic functions*, Iran. J. Sci. Technol. Trans. Sci., DOI: 10.1007/s40995-017-0223-z.
- [4] I. Graham and G. Kohr, *Geometric Function Theory in One and Higher Dimensions*, Marcel Dekker Inc., New York, 2003.
- [5] R. Kargar, N.R. Pascu and A. Ebadian, *Locally univalent approximations of analytic functions*, J. Math. Anal. Appl. **453** (2017), 1005–1021.
- [6] J. Laitila, S. Miihkinen and P. Nieminen, *Essential norms and weak compactness of integration operators*, Arch. Mech. **97** (2011), 39–48.

- [7] S. Li and S. Stević, *Volterra-type operators on Zygmund spaces*, J. Ineq. Appl., Vol. 2007, Article ID 32124, 10 pages, 2007.
- [8] J. Liu, Z. Lou and C. Xiong, *Essential norms of integral operators on spaces of analytic functions*, Nonlin. Anal. **75** (2012), 5145–5156.
- [9] R.J. Libera, *Some classes of regular univalent functions*, Proc. Amer. Math. Soc. **16** (1965), 755–758.
- [10] A.E. Livingston, *On the radius of univalence of certain analytic functions*, Proc. Amer. Math. Soc. **17** (1966) 352–357.
- [11] M. Obradović, S. Ponnusamy, and K.-J. Wirths, *Coefficient characterizations and sections for some univalent functions*, Sib. Math. J. **54** (2013), 679–696.
- [12] S. Owa, *On the distortion theorems I*, Kyungpook Math. J. **18** (1978), 53–59.
- [13] S. Owa, *Properties of certain integral operators*, Georgian Math. J. **2** (1995), 535–545.
- [14] S. Owa and H.M. Srivastava, *Univalent and starlike generalized hypergeometric functions*, Can. J. Math. **39** (1987), 1057–1077.
- [15] S. Ozaki, *On the theory of multivalent functions*, Sci. Rep. Tokyo Bunrika Daigaku, **4** (1941), 45–86.
- [16] C. Pommerenke, *Linear-invariante familien analytischer funktionen. I*, Math. Ann. **155** (1964), 108–154.
- [17] Ch. Pommerenke, *Schlichte Funktionen und analytische Funktionen von beschränkter mittlerer Oszillation*, Comment. Math. Helv. **52** (1977), 591–602.
- [18] A.G. Siskakis, *Composition semigroups and the Cesáro operator on H^p* , J. Lond. Math. Soc. **36** (1987), 153–164.
- [19] A.G. Siskakis, *The Cesáro operator is bounded on H^1* , Proc. Amer. Math. Soc. **110** (1990), 461–462.
- [20] Z. Zhuo and S. Ye, *Volterra-type operators from analytic Morrey spaces to Bloch space*, J. Integral Equ. Appl. **27** (2015), 289–309.
- [21] J. Zhou and X. Zhu, *Essential norm of a Volterra-type operator from Hardy spaces to some analytic function spaces*, J. Integral Equ. Appl. **28** (2016), 581–593.

INDEPENDENT RESEARCHER, WITHOUT DEPENDENCE ON THE ORGANIZATION.
E-mail address: rkargar1983@gmail.com