

Existence of multiple periodic solutions to a semilinear wave equation with x -dependent coefficients

Hui Wei^a, Shuguan Ji^{a,b,*}

^a*School of Mathematics and Statistics and Center for Mathematics and Interdisciplinary Sciences, Northeast Normal University, Changchun 130024, P.R. China*

^b*School of Mathematics, Jilin University, Changchun 130012, P.R. China*

Abstract

This paper is concerned with the periodic (in time) solutions to an one-dimensional semilinear wave equation with x -dependent coefficient. Such a model arises from the forced vibrations of a nonhomogeneous string and propagation of seismic waves in nonisotropic media. By combining variational methods with saddle point reduction technique, we obtain the existence of at least three periodic solutions whenever the period is a rational multiple of the length of the spatial interval. Our method is based on a delicate analysis for the asymptotic character of the spectrum of the wave operator with x -dependent coefficients, and the spectral properties play an essential role in the proof.

Keywords: existence, periodic solutions, wave equation

1. Introduction

In this paper, we consider the existence of multiple periodic solutions to the semilinear wave equation with x -dependent coefficients

$$\rho(x)u_{tt} - (\rho(x)u_x)_x = a\rho(x)u + f(t, x, u), \quad t \in \mathbb{R}, \quad 0 < x < \pi, \quad (1.1)$$

with the Dirichlet boundary conditions

$$u(t, 0) = u(t, \pi) = 0, \quad t \in \mathbb{R}, \quad (1.2)$$

*Corresponding author.

Email addresses: weihui01@163.com (Hui Wei), jishuguan@hotmail.com (Shuguan Ji)

and the periodic conditions

$$u(t + T, x) = u(t, x), \quad u_t(t + T, x) = u_t(t, x), \quad t \in \mathbb{R}, \quad 0 < x < \pi, \quad (1.3)$$

where $a > 0$ is a constant and f is a given T -periodic function in time t .

Equation (1.1) originates from the forced vibrations of a bounded nonhomogeneous string and the propagation of seismic waves in nonisotropic media (see e.g. [6, 7, 8, 18, 19, 21, 22, 23, 27, 28] and references therein). More precisely, the vertical displacement $u(t, z)$ at time t and depth z of a plane seismic wave is described by the equation

$$\mu(z)u_{tt} - (\nu(z)u_z)_z = 0 \quad (1.4)$$

with some initial conditions in t and boundary conditions in z , where $\mu(z)$ is the rock density and $\nu(z)$ is the elasticity coefficient. By the change of variable

$$x = \int_0^z \left(\frac{\mu(s)}{\nu(s)} \right)^{1/2} ds,$$

equation (1.4) is transformed into

$$\rho(x)u_{tt} - (\rho(x)u_x)_x = 0,$$

where $\rho = (\mu\nu)^{1/2}$ is called the acoustic impedance function.

It is well known that the case of $\rho \equiv C \neq 0$ (a nonzero constant) corresponds to the classical wave equation, which is called the one with constant coefficients for distinguishing it from the one with x -dependent coefficients discussed here. The problem of finding periodic solutions of nonlinear wave equation with constant coefficients has received a great deal of attention since the original work [24] of Rabinowitz. By using the variational methods, he obtained the existence of periodic solutions for the weakly nonlinear homogeneous string whenever the period T is a rational multiple of the length of the spatial interval. Thereafter, many authors, such as Brézis, Chang, Nirenberg etc., have used and developed the variational methods, topological degree and index theory to obtain a lot of results on the existence and multiplicity of periodic solutions for the problem with various nonlinearities (see e.g. [1, 3, 5, 9, 11, 13, 14, 15, 16, 17, 25, 26, 30] and the references therein).

On the other hand, the problem of finding periodic solutions for the nonlinear wave equation with x -dependent coefficients was studied by Barbu and Pavel in [6, 7, 8] for the first time. In [7], Barbu and Pavel considered the

existence and regularity of periodic solutions for such wave equation with sublinear nonlinearity under the Dirichlet boundary conditions. For the case the nonlinear term having power-law growth, Rudakov [27] proved the existence of periodic solutions under the Dirichlet boundary conditions. Later, Ji and his collaborators obtained some related results for the general Sturm-Liouville boundary value problem [18, 21], and periodic and anti-periodic boundary value problem [19, 22]. In [31], by using topological degree methods, Wang and An obtained an existence result on periodic solution of the problem with resonance and the sublinear nonlinearity. Afterwards, Ji and Li [23] obtained an existence result of periodic solution for $\eta_\rho(x) = 0$ under the Dirichlet boundary conditions, which actually solves an open problem posted in [7]. Recently, Ji et al. [20] obtain the existence and multiplicity of periodic solutions for the Dirichlet-Neumann boundary value problem of a wave equation with x -dependent coefficients by using the Leray-Schauder degree theory. The restriction to such type of boundary value problem is essentially due to the possible loss of the compactness of the inverse operator on its range.

In this paper, we intend to pay close attention to the existence of multiple periodic solutions of wave equation with x -dependent coefficients. By combining variational methods with saddle point reduction technique, we obtain the existence of at least three periodic solutions whenever the period is a rational multiple of the length of the spatial interval. Our method is based on a delicate analysis for the asymptotic character of the spectrum of the wave operator with x -dependent coefficients, and the spectral properties play an essential role in the proof.

Denote $\tilde{f}(t, x, u) = \frac{f(t, x, u)}{\rho(x)}$. Throughout this paper, we assume T is a rational multiple of π which can be rewritten as

$$T = 2\pi \frac{p}{q}$$

for some relatively prime positive integers p and q . Moreover, we make the following assumptions:

(A1) $\rho(x) \in H^2(0, \pi)$ satisfies $\rho(x) > 0$ for $x \in [0, \pi]$, and

$$\rho_0 = \text{ess inf } \eta_\rho(x) > 0,$$

where

$$\eta_\rho(x) = \frac{1}{2} \frac{\rho''}{\rho} - \frac{1}{4} \left(\frac{\rho'}{\rho} \right)^2.$$

(A2) $\tilde{f}(t, x, u) \in C^1(\mathbb{R} \times (0, \pi) \times \mathbb{R})$, $\tilde{f}(t + T, x, u) = \tilde{f}(t, x, u)$, and

$$\tilde{f}(t, x, u) = o(|u|), \quad \text{as } u \rightarrow 0 \quad \text{uniformly in } (t, x), \quad (1.5)$$

and $\tilde{f}(t, x, u)$ is asymptotically linear in u at ∞ in the following sense: there exists a constant $b > 0$ such that

$$\tilde{f}(t, x, u) - bu = o(|u|), \quad \text{as } |u| \rightarrow \infty \quad \text{uniformly in } (t, x). \quad (1.6)$$

The rest of this paper is organized as follows. In Sect. 2, we give some preliminaries and state the main result. In Sect. 3, we first characterize the solutions of problem (1.1)–(1.3) as the critical points of the corresponding variational problem. Then, with the aid of the saddle point reduction technique, we reduce the critical point of the variation problem from an infinite dimensional space to a finite dimensional subspace. In Sect. 4, we prove reduction functional satisfies the $(PS)_c$ condition for any $c \in \mathbb{R}$. In Sect. 5 and Sect. 6, we devote to the proof of the bounds of reduction functional and the main result respectively.

2. Preliminaries and main result

Set $\Omega = (0, T) \times (0, \pi)$, and denote

$$\Psi = \{\psi \in C^\infty(\Omega) : \psi(t, 0) = \psi(t, \pi) = 0, \psi(0, x) = \psi(T, x), \psi_t(0, x) = \psi_t(T, x)\},$$

and

$$L^r(\Omega) = \left\{ u : \|u\|_{L^r(\Omega)}^r = \int_{\Omega} |u(t, x)|^r \rho(x) dt dx < \infty \right\}, \quad r \geq 1.$$

It is well known that Ψ is dense in $L^r(\Omega)$ for any $r \geq 1$, and $L^2(\Omega)$ is a Hilbert space with the inner product

$$\langle u, v \rangle = \int_{\Omega} u(t, x) \overline{v(t, x)} \rho(x) dt dx, \quad \forall u, v \in L^2(\Omega).$$

We rewrite (1.1)–(1.3) on Ω in the following form

$$\rho(x)u_{tt} - (\rho(x)u_x)_x = a\rho(x)u + f(t, x, u), \quad (t, x) \in \Omega, \quad (2.1)$$

$$u(t, 0) = u(t, \pi) = 0, \quad t \in (0, T), \quad (2.2)$$

$$u(0, x) = u(T, x), \quad u_t(0, x) = u_t(T, x), \quad x \in (0, \pi). \quad (2.3)$$

Definition 2.1. A function $u \in L^r(\Omega)$ is called a weak solution of problem (2.1)–(2.3) if it satisfies

$$\int_{\Omega} u(\rho\psi_{tt} - (\rho\psi_x)_x) dt dx - \int_{\Omega} (au + \tilde{f}(t, x, u))\psi\rho dt dx = 0, \quad \forall \psi \in \Psi.$$

Define the linear operator L_0 by

$$L_0\psi = \rho^{-1}(\rho\psi_{tt} - (\rho\psi_x)_x), \quad \forall \psi \in \Psi,$$

and denote its extension in $L^2(\Omega)$ by L . It is known that L is a selfadjoint operator (see [7]), and $u \in L^2(\Omega)$ is a weak solution of problem (2.1)–(2.3) if and only if $Lu = au + \tilde{f}$.

For the study of periodic solutions of problem (2.1)–(2.3), we need to use the following complete orthonormal system of eigenfunctions $\{\phi_m(t)\varphi_n(x) : m \in \mathbb{Z}, n \in \mathbb{N}\}$ in $L^2(\Omega)$ (see [32]), where

$$\phi_m(t) = T^{-\frac{1}{2}}e^{i\mu_m t}, \quad \mu_m = 2m\pi T^{-1}, \quad m \in \mathbb{Z},$$

and $\lambda_n, \varphi_n(x)$ are given by the Sturm-Liouville problem

$$-(\rho(x)\varphi'_n(x))' = \lambda_n^2\rho(x)\varphi_n(x), \quad \varphi_n(0) = \varphi_n(\pi) = 0, \quad n \in \mathbb{N}. \quad (2.4)$$

Lemma 2.1 ([7]). Assume that $\rho(x)$ satisfies (A1), then the eigenvalues of problem (2.4) have the form

$$\lambda_n = n + \theta_n \text{ with } \theta_n \rightarrow 0 \text{ as } n \rightarrow \infty,$$

where

$$0 < \frac{\rho_2}{n} \leq \sqrt{n^2 + \rho_0} - n \leq \theta_n \leq \sqrt{n^2 + \rho_1} - n \leq \frac{\rho_1}{2n}, \quad (2.5)$$

and $\rho_1 = \frac{2}{\pi} \int_0^\pi \eta_\rho(x) dx$, $\rho_2 = \sqrt{\rho_0 + 1} - 1$.

By Lemma 2.1, the eigenvalues of operator L can be rewritten as

$$\lambda_{nm} = \lambda_n^2 - \mu_m^2 = p^{-2}(np + \theta_n p - mq)(np + \theta_n p + mq).$$

Thus, when $np \neq |m|q$, it is easy to verify that $|\lambda_{nm}| \rightarrow \infty$ as $m, n \rightarrow \infty$. On the other hand, when $np = |m|q$, by (2.5) we have

$$2\rho_2 \leftarrow \left(\frac{\rho_2}{n}\right)^2 + 2\rho_2 \leq \lambda_{nm} = \theta_n(2n + \theta_n) \leq \rho_1 + \left(\frac{\rho_1}{2n}\right)^2 \rightarrow \rho_1,$$

as $n \rightarrow \infty$.

Denote the set of eigenvalues of operator L by

$$\Lambda(L) = \{\lambda_{nm} : \lambda_{nm} = \lambda_n^2 - \mu_m^2\}.$$

The above statement shows that $\Lambda(L)$ has at least one accumulation point in $[2\rho_2, \rho_1]$. Therefore we have the following lemma.

Lemma 2.2. *Let assumption (A1) hold. Then*

- (i) *L has at least one essential spectral point, and all of them belong to $[2\rho_2, \rho_1]$;*
- (ii) *If $\lambda \in \Lambda(L)$ and $\lambda \notin [2\rho_2, \rho_1]$, then λ is isolated and its multiplicity is finite.*

If $(a, b) \cap \Lambda(L) \neq \emptyset$ and b satisfies $\rho_1 < a < b$, by Lemma 2.2, we can define

$$b^- = \max\{\lambda \in \Lambda(L) : \lambda < b\}, \quad b^+ = \min\{\lambda \in \Lambda(L) : \lambda > b\}.$$

It is obvious that $b^- < b < b^+$.

Denote $e = b^+ - a$. The main result of this paper is as follows.

Theorem 2.1. *Assume $a, b \notin \Lambda(L)$, $\rho_1 < a < b^+ - b$, $(a, b) \cap \Lambda(L) \neq \emptyset$, and (A1)–(A2) hold. If \tilde{f} is increasing in u and there exists a constant $\kappa > 0$ such that*

$$\frac{\partial \tilde{f}}{\partial u}(t, x, u) \leq e - \kappa, \quad \forall (t, x, u) \in \mathbb{R} \times (0, \pi) \times \mathbb{R}.$$

Then the problem (2.1)–(2.3) has at least three T -periodic solutions.

3. Variational problem and its reduction

In what follows, we always assume a, b satisfy the conditions in Theorem 2.1. Since $a \notin \Lambda(L)$ and $a > \rho_1$, then there exists a constant $\delta > 0$ such that

$$|\lambda_{nm} - a| \geq \delta > 0, \quad m \in \mathbb{Z}, n \in \mathbb{N}. \quad (3.1)$$

We define the working space

$$E = \left\{ u \in L^2(\Omega) : \|u\|_E^2 = \sum_{n,m} |\lambda_{nm} - a| |\alpha_{nm}|^2 < \infty \right\},$$

where α_{nm} denote the Fourier coefficients of $u \in L^2(\Omega)$, i.e.,

$$u = \sum_{n,m} \alpha_{nm} \varphi_n(x) \phi_m(t), \quad \alpha_{nm} = \int_{\Omega} u \varphi_n \bar{\phi}_m \rho dx dt.$$

The estimate (3.1) indicates $\|\cdot\|_E$ is a norm. Furthermore, E is a Hilbert space equipped with the inner product $\langle u, v \rangle_0 = \sum_{n,m} |\lambda_{nm} - a| \alpha_{nm} \bar{\beta}_{nm}$, where α_{nm} and β_{nm} are the Fourier coefficients of u and v respectively.

From (3.1), we have

$$\|u\|_{L^2(\Omega)}^2 = \sum_{n,m} |\alpha_{nm}|^2 \leq \delta^{-1} \sum_{n,m} |\lambda_{nm} - a| |\alpha_{nm}|^2 = \delta^{-1} \|u\|_E^2, \quad (3.2)$$

which implies that the continuous embedding $E \hookrightarrow L^2(\Omega)$. Moreover, for $1 \leq r \leq 2$, the continuous embedding $L^2(\Omega) \hookrightarrow L^r(\Omega)$ implies that there exists a constant $C = C(r)$ such that

$$\|u\|_{L^r(\Omega)} \leq C \|u\|_E, \quad 1 \leq r \leq 2. \quad (3.3)$$

Since $a \in (\rho_1, b^+ - b)$, then

$$e = b^+ - a > b. \quad (3.4)$$

If $\lambda_{nm} > b$, we have

$$\lambda_{nm} - a \geq e > b, \quad m \in \mathbb{Z}, n \in \mathbb{N}. \quad (3.5)$$

Define the energy functional

$$\Phi(u) = \frac{1}{2} \langle (L - a)u, u \rangle - \int_{\Omega} \tilde{F}(t, x, u) \rho dt dx, \quad \forall u \in E, \quad (3.6)$$

where $\tilde{F}(t, x, u) = \int_0^u \tilde{f}(t, x, s) ds$. In addition, by (1.5) and the assumption on \tilde{f} in Theorem 2.1, it is easy to see $\tilde{F}(t, x, u) \geq 0$ for any $u \in E$.

Consequently, Φ is a C^1 functional on E and

$$\langle \Phi'(u), v \rangle = \langle (L - a)u, v \rangle - \int_{\Omega} \tilde{f}(t, x, u) v \rho dt dx, \quad \forall u, v \in E. \quad (3.7)$$

Then u is a weak solution of problem (2.1)–(2.3) if and only if $\Phi'(u) = 0$. Therefore, solutions of problem (2.1)–(2.3) are characterized as critical points of the functional Φ . Since \tilde{f} is C^1 , then Φ is a C^2 functional on E , and

$$\langle \Phi''(u)w, v \rangle = \langle (L - a)w, v \rangle - \int_{\Omega} \frac{\partial \tilde{f}}{\partial u}(t, x, u)vw\rho dt dx, \quad \forall u, v, w \in E.$$

In particular,

$$\langle \Phi''(u)v, v \rangle = \langle (L - a)v, v \rangle - \int_{\Omega} \frac{\partial \tilde{f}}{\partial u}(t, x, u)v^2\rho dt dx, \quad \forall u, v \in E. \quad (3.8)$$

It's not difficult to see that Φ is neither bounded from above nor from below, which shows that we can't obtain the critical points of Φ by a simple minimization or maximization. Here we shall prove our main result by virtue of the following saddle point reduction technique developed by Amann [1], Castro and Lazer [12], and Arcoya and Costa [2] etc.

Lemma 3.1. *Let H be a real Hilbert space with the norm $\|\cdot\|_H$, $\Phi \in C^1(H, \mathbb{R})$, and H_1, H_2 and H_3 be closed subset of H such that $H = H_1 \oplus H_2 \oplus H_3$. If there exists a constant $\gamma > 0$ satisfying*

$$\langle \Phi'(u+w+v_1) - \Phi'(u+w+v_2), v_1 - v_2 \rangle \leq -\gamma \|v_1 - v_2\|_H^2, \quad \forall v_1, v_2 \in H_1, u \in H_2, w \in H_3,$$

and

$$\langle \Phi'(u+w_1+v) - \Phi'(u+w_2+v), w_1 - w_2 \rangle \geq \gamma \|w_1 - w_2\|_H^2, \quad \forall v \in H_1, u \in H_2, w_1, w_2 \in H_3.$$

Then

(i) *There exists a unique continuous mapping $h : H_2 \rightarrow H_1 \oplus H_3$, such that*

$$\Phi(u + h(u)) = \max_{v \in H_1} \min_{w \in H_3} \Phi(u + v + w) = \min_{w \in H_3} \max_{v \in H_1} \Phi(u + v + w);$$

(ii) *Define $\tilde{\Phi}(u) = \Phi(u + h(u))$ for any $u \in H_2$, then $\tilde{\Phi} \in C^1(H_2, \mathbb{R})$, and*

$$\langle \tilde{\Phi}'(u), v \rangle = \langle \Phi'(u + h(u)), v \rangle, \quad \forall u, v \in H_2;$$

(iii) *If $u \in H_2$ is a critical point of $\tilde{\Phi}$, then $u + h(u)$ is a critical point of Φ . On the other hand, if $u + v$ is a critical point of Φ , then $v = h(u)$ and u is a critical point of $\tilde{\Phi}$, where $u \in H_2, v \in H_1 \oplus H_3$;*

(iv) *If Φ satisfies the Palais-Smale condition $(PS)_c$ at the level $c \in \mathbb{R}$, then the functional $\tilde{\Phi}$ also satisfies the $(PS)_c$ condition.*

By an observation of Lemma 3.1, it needs to decompose the working space E into suitable orthogonal subspaces. Noting $a, b \notin \Lambda(L)$ and $a > \rho_1$, the working space E can be decomposed into the following orthogonal subspaces

$$\begin{aligned} E_1 &= \left\{ u \in E : \lambda_{nm} < a, m \in \mathbb{Z}, n \in \mathbb{N} \right\}, \\ E_2 &= \left\{ u \in E : a < \lambda_{nm} < b, m \in \mathbb{Z}, n \in \mathbb{N} \right\}, \\ E_3 &= \left\{ u \in E : \lambda_{nm} > b, m \in \mathbb{Z}, n \in \mathbb{N} \right\}. \end{aligned}$$

Thus we write $E = E_1 \oplus E_2 \oplus E_3$. Moreover, by $(a, b) \cap \Lambda(L) \neq \emptyset$ and Lemma 2.2, we have $E_2 \neq \emptyset$ and $\dim(E_2) < \infty$.

For any $u \in E_1$, we write $u = \sum_{\lambda_{nm} < a} \alpha_{nm} \varphi_n \phi_m$, then

$$\langle (L - a)u, u \rangle = - \sum_{\lambda_{nm} < a} |\lambda_{nm} - a| |\alpha_{nm}|^2 = -\|u\|_E^2. \quad (3.9)$$

For any $u \in E_2 \oplus E_3$, by a similar calculation we have

$$\langle (L - a)u, u \rangle = \|u\|_E^2. \quad (3.10)$$

With the help of (3.9) and (3.10), we have the following lemma.

Lemma 3.2. *Let the assumptions in Theorem 2.1 hold. Then there exists a constant $\gamma > 0$ such that*

$$\langle \Phi'(u + v) - \Phi'(u + w), v - w \rangle \leq -\gamma \|v - w\|_E^2, \quad \forall u \in E_2 \oplus E_3, v, w \in E_1,$$

and

$$\langle \Phi'(u + v) - \Phi'(u + w), v - w \rangle \geq \gamma \|v - w\|_E^2, \quad \forall u \in E_1 \oplus E_2, v, w \in E_3.$$

Proof. For any $u, v, w \in E$ and $s \in \mathbb{R}$, we have

$$\langle \Phi'(u + v) - \Phi'(u + w), v - w \rangle = \int_0^1 \langle \Phi''(u + w + s(v - w))(v - w), v - w \rangle ds. \quad (3.11)$$

From (3.8), we obtain

$$\begin{aligned} & \langle \Phi''(u + w + s(v - w))(v - w), v - w \rangle \\ &= \langle (L - a)(v - w), v - w \rangle - \int_{\Omega} (v - w)^2 \frac{\partial \tilde{f}}{\partial u}(t, x, u + w + s(v - w)) \rho dt dx. \end{aligned} \quad (3.12)$$

The assumption on \tilde{f} in Theorem 2.1 shows $\frac{\partial \tilde{f}}{\partial u} \geq 0$. For the case $v, w \in E_1, u \in E_2 \oplus E_3$, by (3.9) we have

$$\langle \Phi'(u+v) - \Phi'(u+w), v-w \rangle \leq -\|v-w\|_E^2.$$

For the case $v, w \in E_3, u \in E_1 \oplus E_2$, by (3.10) we have

$$\langle (L-a)(v-w), v-w \rangle = \|v-w\|_E^2. \quad (3.13)$$

Moreover, since $0 \leq \frac{\partial \tilde{f}}{\partial u}(t, x, u) \leq e - \kappa$, with the aid of (3.5), a direct calculation yields

$$\int_{\Omega} (v-w)^2 \frac{\partial \tilde{f}}{\partial u} \rho dt dx \leq (e - \kappa) \|v-w\|_{L^2(\Omega)}^2 \leq \frac{(e - \kappa)}{e} \|v-w\|_E^2. \quad (3.14)$$

Inserting (3.13), (3.14) into (3.12), from (3.11) we have

$$\langle \Phi'(u+v) - \Phi'(u+w), v-w \rangle \geq \frac{\kappa}{e} \|v-w\|_E^2.$$

By setting $\gamma = \min\{1, \frac{\kappa}{e}\}$, we arrive at the assertion. \square

Lemma 3.2 verifies all the conditions in Lemma 3.1. Therefore there exists a unique continuous mapping $h : E_2 \rightarrow E_1 \oplus E_3$ such that

$$\tilde{\Phi}(u) = \Phi(u+h(u)) = \max_{v \in E_1} \min_{w \in E_3} \Phi(u+v+w) = \min_{w \in E_3} \max_{v \in E_1} \Phi(u+v+w). \quad (3.15)$$

Moreover, Lemma 3.1 shows the reduction functional $\tilde{\Phi} \in C^1(E_2, \mathbb{R})$, and $u \in E_2$ is a critical point of $\tilde{\Phi}$ if and only if $u+v+w$ is a critical point of Φ , where $v \in E_1, w \in E_3$ and $h(u) = v+w$. Thus, the critical points of Φ on the infinite dimensional space E are transformed into the ones of $\tilde{\Phi}$ on the finite dimensional subspace E_2 . In what follows, we shall apply the variational methods to obtain critical points of the functional $\tilde{\Phi}$ on E_2 .

4. Verification the $(PS)_c$ condition

We will acquire the critical points of $\tilde{\Phi}$ via variational methods, thus it needs to verify $\tilde{\Phi}$ satisfies $(PS)_c$ condition for any $c \in \mathbb{R}$. Furthermore, Lemma 3.1 shows that it suffices to verify that Φ satisfies $(PS)_c$ condition, which means, any sequence $\{u_i\} \subset E$ satisfying $\Phi(u_i) \rightarrow c$ and $\Phi'(u_i) \rightarrow 0$ as $i \rightarrow \infty$ has a convergent subsequence for any $c \in \mathbb{R}$. To this goal, we need the following lemma which provides two estimates for the quadratic forms on different subspaces of E .

Lemma 4.1. *Let a, b satisfy the assumptions in Theorem 2.1. Then there exist $\gamma_1, \gamma_2 > 0$ such that*

$$\langle (L - a - b)u, u \rangle \leq -\gamma_1 \|u\|_E^2, \quad \forall u \in E_1 \oplus E_2, \quad (4.1)$$

$$\langle (L - a - b)u, u \rangle \geq \gamma_2 \|u\|_E^2, \quad \forall u \in E_3. \quad (4.2)$$

Proof. On the one hand, for $u \in E_1 \oplus E_2$, we write $u = \sum_{\lambda_{nm} < b} \alpha_{nm} \phi_m \varphi_n$. A

direct calculation yields

$$\begin{aligned} & \langle (L - a - b)u, u \rangle \\ &= \sum_{\lambda_{nm} < b} (\lambda_{nm} - a) |\alpha_{nm}|^2 - b \sum_{\lambda_{nm} < b} |\alpha_{nm}|^2 \\ &\leq \sum_{\lambda_{nm} < a} (\lambda_{nm} - a) |\alpha_{nm}|^2 + \sum_{a < \lambda_{nm} < b} (\lambda_{nm} - a) |\alpha_{nm}|^2 - b \sum_{a < \lambda_{nm} < b} |\alpha_{nm}|^2. \end{aligned}$$

In virtue of $a < \lambda_{nm} < b$, we have $0 < \lambda_{nm} - a < b^-$. Thus, we have

$$\frac{1}{b^-} \sum_{a < \lambda_{nm} < b} |\lambda_{nm} - a| |\alpha_{nm}|^2 \leq \sum_{a < \lambda_{nm} < b} |\alpha_{nm}|^2.$$

Therefore,

$$\begin{aligned} & \langle (L - a - b)u, u \rangle \\ &\leq - \sum_{\lambda_{nm} < a} |\lambda_{nm} - a| |\alpha_{nm}|^2 - \left(\frac{b}{b^-} - 1 \right) \sum_{a < \lambda_{nm} < b} |\lambda_{nm} - a| |\alpha_{nm}|^2 \\ &\leq -\gamma_1 \|u\|_E^2, \end{aligned}$$

where $\gamma_1 = \min\{1, \frac{b}{b^-} - 1\}$. Noting $0 < b^- < b$, it follows $\gamma_1 > 0$.

On the other hand, for $u \in E_3$, we write $u = \sum_{\lambda_{nm} > b} \alpha_{nm} \phi_m \varphi_n$. Observing $\lambda_{nm} > b$, from (3.5), we obtain

$$\frac{1}{e} \sum_{\lambda_{nm} > b} |\lambda_{nm} - a| |\alpha_{nm}|^2 \geq \sum_{\lambda_{nm} > b} |\alpha_{nm}|^2.$$

Therefore,

$$\begin{aligned} \langle (L - a - b)u, u \rangle &= \sum_{\lambda_{nm} > b} |\lambda_{nm} - a| |\alpha_{nm}|^2 - b \sum_{\lambda_{nm} > b} |\alpha_{nm}|^2 \\ &\geq \left(1 - \frac{b}{e} \right) \sum_{\lambda_{nm} > b} |\lambda_{nm} - a| |\alpha_{nm}|^2. \end{aligned}$$

Let $\gamma_2 = 1 - \frac{b}{\varepsilon}$, by (3.4), it follows $\gamma_2 > 0$. Therefore

$$\langle (L - a - b)u, u \rangle \geq \gamma_2 \|u\|_E^2.$$

The proof is completed. \square

Lemma 4.2. *Let the assumptions in Theorem 2.1 hold. If $\{u_i\} \subset E$ satisfies $\Phi(u_i) \rightarrow c$ and $\Phi'(u_i) \rightarrow 0$ as $i \rightarrow \infty$, then there exists a constant $\tilde{C} > 0$ independent of i such that $\|u_i\|_E \leq \tilde{C}$.*

Proof. Split $u_i = u_i^+ + u_i^-$, where $u_i^+ \in E_3$ and $u_i^- \in E_1 \oplus E_2$, $i = 1, 2, \dots$.

(i) For $u_i^+ \in E_3$, since $\Phi'(u_i) \rightarrow 0$ as $i \rightarrow \infty$, from (3.7), we have

$$\begin{aligned} o(1)\|u_i^+\|_E &\geq \langle \Phi'(u_i), u_i^+ \rangle = \langle (L - a)u_i^+, u_i^+ \rangle - \int_{\Omega} \tilde{f}(t, x, u_i) u_i^+ \rho dt dx \\ &= \langle (L - a - b)u_i^+, u_i^+ \rangle - \int_{\Omega} (\tilde{f}(t, x, u_i) - bu_i) u_i^+ \rho dt dx. \end{aligned} \quad (4.3)$$

From (4.2), we obtain

$$\langle (L - a - b)u_i^+, u_i^+ \rangle \geq \gamma_2 \|u_i^+\|_E^2. \quad (4.4)$$

On the other side, the condition (1.6) shows that for any $\varepsilon > 0$, there exists a constant $C = C(\varepsilon) > 0$ such that

$$|\tilde{f}(t, x, u_i) - bu_i| < \varepsilon |u_i| + C. \quad (4.5)$$

Therefore, by Hölder inequality and (3.2), (3.3), a direct calculation yields

$$\begin{aligned} &\left| \int_{\Omega} (\tilde{f}(t, x, u_i) - bu_i) u_i^+ \rho dt dx \right| \\ &\leq \varepsilon \|u_i^+\|_{L^2(\Omega)} \|u_i\|_{L^2(\Omega)} + C \|u_i^+\|_{L^1(\Omega)} \\ &\leq \frac{\varepsilon}{2\delta} \|u_i^+\|_E^2 + \frac{\varepsilon}{2\delta} \|u_i\|_E^2 + C \|u_i^+\|_E, \end{aligned} \quad (4.6)$$

for some constant C independent of i .

Inserting (4.4), (4.6) into (4.3), we obtain

$$\left(\gamma_2 - \frac{\varepsilon}{2\delta}\right) \|u_i^+\|_E^2 - \frac{\varepsilon}{2\delta} \|u_i\|_E^2 - C \|u_i^+\|_E \leq 0. \quad (4.7)$$

(ii) For $u_i^- \in E_1 \oplus E_2$, by (4.1), we have $\langle (L - a - b)u_i^-, u_i^- \rangle \leq -\gamma_1 \|u_i^-\|_E^2$. Moreover, noting that

$$\begin{aligned} o(1)\|u_i^-\|_E &\geq \langle -\Phi'(u_i), u_i^- \rangle \\ &= -\langle (L - a - b)u_i^-, u_i^- \rangle + \int_{\Omega} (\tilde{f}(t, x, u_i) - bu_i)u_i^- \rho dt dx, \end{aligned}$$

a similar calculation as in (4.6) yields

$$\left| \int_{\Omega} (\tilde{f}(t, x, u_i) - bu_i)u_i^- \rho dt dx \right| \leq \frac{\varepsilon}{2\delta} \|u_i^-\|_E^2 + \frac{\varepsilon}{2\delta} \|u_i\|_E^2 + C\|u_i^-\|_E. \quad (4.8)$$

Consequently,

$$\left(\gamma_1 - \frac{\varepsilon}{2\delta}\right) \|u_i^-\|_E^2 - \frac{\varepsilon}{2\delta} \|u_i\|_E^2 - C\|u_i^-\|_E \leq 0. \quad (4.9)$$

Let $\gamma_0 = \min\{\gamma_1, \gamma_2\}$. Since E_1, E_2 and E_3 are orthogonal subspaces of E , we have $\|u_i\|_E^2 = \|u_i^+\|_E^2 + \|u_i^-\|_E^2$. Therefore, by the fact $\|u_i^+\|_E + \|u_i^-\|_E \leq \sqrt{2}\|u_i\|_E$, the sum of (4.7) and (4.9) yields

$$\left(\gamma_0 - \frac{3\varepsilon}{2\delta}\right) \|u_i\|_E^2 - C\|u_i\|_E \leq 0. \quad (4.10)$$

Taking $\varepsilon \in (0, \frac{2\delta\gamma_0}{3})$ in (4.10), we obtain there exists a constant $\tilde{C} > 0$ independent of i such that $\|u_i\|_E \leq \tilde{C}$. We arrive at the result. \square

Since $E = E_1 \oplus E_2 \oplus E_3$, we can rewrite $E = E_1 \oplus E_1^\perp$ for simplicity, where

$$E_1^\perp = E_2 \oplus E_3 = \left\{ u \in E : \lambda_{nm} > a, m \in \mathbb{Z}, n \in \mathbb{N} \right\}.$$

Denote

$$E_0 = \left\{ u \in L^2(\Omega) : pn = q|m|, m \in \mathbb{Z}, n \in \mathbb{N} \right\}.$$

Remark 4.1. Under the assumptions of Theorem 2.1, the Lemma 2.2 shows E_0 is an infinite dimensional subspace spanned by the eigenfunctions $\phi_m \varphi_n$ for $pn = q|m|$. Moreover, it is easy to see that $\dim(E_1^\perp \cap E_0) < \infty$ and $\dim(E_1 \cap E_0) = \infty$.

Proposition 4.1. For $1 \leq r \leq 2$, the embedding

$$E \ominus E_0 \hookrightarrow L^r(\Omega), \quad (4.11)$$

is compact.

Proof. For $u \in E \ominus E_0$, it can be expanded as $u = \sum_{n,m} \alpha_{nm} \phi_m \varphi_n$ for $pn \neq q|m|$.

The fact $\|u\|_E = \left(\sum_{n,m} |\lambda_{nm} - a| |\alpha_{nm}|^2 \right)^{\frac{1}{2}}$ shows that the mapping

$$I_0 : u = \sum_{n,m} \alpha_{nm} \phi_m \varphi_n \mapsto \{|\lambda_{nm} - a|^{\frac{1}{2}} \alpha_{nm}\}$$

is continuous from $E \ominus E_0$ to l^2 .

Since $|\lambda_{nm} - a| \rightarrow \infty$ as $m, n \rightarrow \infty$, it follows that the mapping

$$I_1 : \{|\lambda_{nm} - a|^{\frac{1}{2}} \alpha_{nm}\} \mapsto \{\alpha_{nm}\}$$

is compact from l^2 to l^2 .

Since $\phi_m \varphi_n$ is a complete orthonormal sequence of $L^2(\Omega)$, then the mapping

$$I_2 : \{\alpha_{nm}\} \mapsto u = \sum_{n,m} \alpha_{nm} \phi_m \varphi_n$$

is continuous from l^2 to $L^2(\Omega)$.

Consequently, the mapping

$$I_2 I_1 I_0 : E \ominus E_0 \rightarrow L^2(\Omega)$$

is compact. Furthermore, for $1 \leq r \leq 2$, the continuous embedding $L^2(\Omega) \hookrightarrow L^r(\Omega)$ implies that the embedding $E \ominus E_0 \hookrightarrow L^r(\Omega)$ is compact. \square

Lemma 4.3. *Let the assumptions of Theorem 2.1 hold. Then Φ satisfies the $(PS)_c$ condition for any $c \in \mathbb{R}$.*

Proof. For any $c \in \mathbb{R}$, assume $\{u_i\} \subset E$ satisfies $\Phi(u_i) \rightarrow c$ and $\Phi'(u_i) \rightarrow 0$ as $i \rightarrow \infty$. Since E is a Hilbert space, by Lemma 4.2, we have $u_i \rightharpoonup u$ as $i \rightarrow \infty$ for some $u \in E$. Decompose $u_i = v_i + y_i + w_i + z_i$ and $u = v + y + w + z$, where v, y, w, z are the weak limits of $\{v_i\}, \{y_i\}, \{w_i\}, \{z_i\}$ respectively, and $v_i, v \in E_1^\perp \ominus E_0, y_i, y \in E_1^\perp \cap E_0, w_i, w \in E_1 \ominus E_0, z_i, z \in E_1 \cap E_0$.

(i) For $v_i, v \in E_1^\perp \ominus E_0$, from (3.10) we have

$$\begin{aligned} \|v_i - v\|_E^2 &= \langle (L - a)(v_i - v), v_i - v \rangle \\ &= \langle (L - a)v_i, v_i - v \rangle - \langle (L - a)v, v_i - v \rangle. \end{aligned} \quad (4.12)$$

In virtue of $v_i \rightharpoonup v$ in E and $E \hookrightarrow L^2(\Omega)$, we have $v_i \rightharpoonup v$ in $L^2(\Omega)$ along with a subsequence of $\{v_i\}$. In fact, by Proposition 4.1, we also have $v_i \rightarrow v$

in $L^2(\Omega)$. We still use $\{v_i\}$ to denote the subsequence for convenience. Thus, it follows

$$\langle (L - a)v, v_i - v \rangle \rightarrow 0, \quad \text{as } i \rightarrow \infty. \quad (4.13)$$

On the other hand, noting $v_i, v \in E_1^\perp \ominus E_0$ and $u_i = v_i + y_i + w_i + z_i$, we have $u_i - v_i \in (E_1^\perp \ominus E_0)^\perp$. Thus

$$\langle (L - a)(u_i - v_i), v_i - v \rangle = 0.$$

By (3.7), we have

$$\begin{aligned} \langle (L - a)v_i, v_i - v \rangle &= \langle (L - a)u_i, v_i - v \rangle \\ &= \langle \Phi'(u_i), v_i - v \rangle + \int_{\Omega} \tilde{f}(t, x, u_i)(v_i - v)\rho dt dx. \end{aligned} \quad (4.14)$$

Since $\Phi'(u_i) \rightarrow 0$ as $i \rightarrow \infty$, we have

$$\langle \Phi'(u_i), v_i - v \rangle \rightarrow 0, \quad \text{as } i \rightarrow \infty. \quad (4.15)$$

By (4.5) and Hölder inequality, a direct calculation yields

$$\left| \int_{\Omega} \tilde{f}(t, x, u_i)(v_i - v)\rho dt dx \right| \leq (b + \varepsilon)\|u_i\|_{L^2(\Omega)}\|v_i - v\|_{L^2(\Omega)} + C\|v_i - v\|_{L^1(\Omega)}.$$

In virtue of $v_i \rightarrow v$ in $L^2(\Omega)$ and the continuous embedding $L^2(\Omega, \rho) \hookrightarrow L^1(\Omega)$, we obtain $v_i \rightarrow v$ in $L^1(\Omega)$. Therefore,

$$\left| \int_{\Omega} \tilde{f}(t, x, u_i)(v_i - v)\rho dt dx \right| \rightarrow 0, \quad \text{as } i \rightarrow \infty. \quad (4.16)$$

Inserting (4.15), (4.16) into (4.14), we have

$$\langle (L - a)v_i, v_i - v \rangle \rightarrow 0, \quad \text{as } i \rightarrow \infty. \quad (4.17)$$

Finally, substituting (4.17), (4.13) into (4.12), we have

$$\|v_i - v\|_E \rightarrow 0, \quad \text{as } i \rightarrow \infty. \quad (4.18)$$

(ii) For $y_i, y \in E_1^\perp \cap E_0$, since $\dim(E_1^\perp \cap E_0) < \infty$ and $y_i \rightharpoonup y$ in E , then there exists a subsequence of $\{y_i\}$ which strongly converges to y in E . The subsequence is still denoted by $\{y_i\}$.

(iii) For $w_i, w \in E_1 \ominus E_0$, by (3.7) and (3.9) we obtain

$$\begin{aligned}\|w_i - w\|_E^2 &= -\langle (L - a)(w_i - w), w_i - w \rangle \\ &= -\langle \Phi'(u_i), w_i - w \rangle - \int_{\Omega} \tilde{f}(t, x, u_i)(w_i - w) \rho dt dx + \langle (L - a)w, w_i - w \rangle.\end{aligned}$$

A similar calculation as in (i) yields

$$\|w_i - w\|_E \rightarrow 0, \text{ as } i \rightarrow \infty. \quad (4.19)$$

(iv) Since $z_i \in E_1 \cap E_0$, then the compact embedding (4.11) is invalid. In addition, since $\dim(E_1 \cap E_0) = \infty$, we can not extract a strong convergence subsequence of $\{z_i\}$ similar to $\{y_i\}$. In what follows, with the aid of the monotone method, we prove $z_i \rightarrow z$ as $i \rightarrow \infty$ in E .

Since $\Phi'(u_i) \rightarrow 0$ and $z_i \rightarrow z$ in E , we have

$$\begin{aligned}\|z_i - z\|_E^2 &= -\langle (L - a)(z_i - z), z_i - z \rangle \\ &= -\langle \Phi'(u_i), z_i - z \rangle - \int_{\Omega} \tilde{f}(t, x, u_i)(z_i - z) \rho dt dx + \langle (L - a)z, z_i - z \rangle \\ &\leq -\int_{\Omega} \tilde{f}(t, x, u_i)(z_i - z) \rho dt dx + o(1).\end{aligned} \quad (4.20)$$

Denote $\tilde{f}(u_i) = \tilde{f}(t, x, u_i)$ for simplicity. We rewrite $\int_{\Omega} \tilde{f}(t, x, u_i)(z_i - z) \rho dt dx$ in the form of inner product and decompose it as follows

$$\begin{aligned}&\int_{\Omega} \tilde{f}(t, x, u_i)(z_i - z) \rho dt dx \quad (4.21) \\ &= \langle \tilde{f}(u_i), z_i - z \rangle \\ &= \langle \tilde{f}(u_i) - \tilde{f}(\tilde{u}_i + z), z_i - z \rangle + \langle \tilde{f}(\tilde{u}_i + z) - \tilde{f}(u), z_i - z \rangle \\ &\quad + \langle \tilde{f}(u), z_i - z \rangle,\end{aligned} \quad (4.22)$$

where $\tilde{u}_i = v_i + y_i + w_i$.

Since \tilde{f} is increasing in u , then

$$\langle \tilde{f}(u_i) - \tilde{f}(\tilde{u}_i + z), z_i - z \rangle \geq 0. \quad (4.23)$$

To continue the discussion, by (4.5), we have $\tilde{f} : u \mapsto \tilde{f}(t, x, u)$ is continuous from $L^2(\Omega)$ to $L^2(\Omega)$. Moreover, from the proof of (i)–(iii), we have

$\tilde{u}_i \rightarrow \tilde{u}$ in E , where $\tilde{u} = v + y + w$. The inequality (3.2) shows $\tilde{u}_i \rightarrow \tilde{u}$ in $L^2(\Omega)$. Thus

$$\langle \tilde{f}(\tilde{u}_i + z) - \tilde{f}(u), z_i - z \rangle \rightarrow 0, \quad \text{as } i \rightarrow \infty. \quad (4.24)$$

Consequently, since $z_i \rightarrow z$ in $L^2(\Omega)$, by (4.20), (4.23) and (4.24), we have

$$\|z_i - z\|_E \rightarrow 0, \quad \text{as } i \rightarrow \infty.$$

We arrive at result. □

5. Bounds of the reduction functional

The following three lemmas are concerned with the bounds of the reduction functional $\tilde{\Phi}$ and play an important role in the proof of Theorem 2.1.

Lemma 5.1. *Let the assumptions of Theorem 2.1 hold. Then*

- (i) *there exists a constant $M > 0$, such that $\tilde{\Phi}(u) < M$, $\forall u \in E_2$;*
- (ii) *there exists a constant $\tilde{R} > 0$ such that $\tilde{\Phi}(u) \leq 0$ for $u \in E_2$ with $\|u\|_E \geq \tilde{R}$.*

Proof. For $u \in E_2$, from (3.15), we have

$$\tilde{\Phi}(u) = \min_{w \in E_3} \max_{v \in E_1} \Phi(u + v + w) \leq \max_{v \in E_1} \Phi(u + v),$$

where

$$\Phi(u+v) = \frac{1}{2} \langle (L-a-b)(u+v), u+v \rangle - \int_{\Omega} \left(\tilde{F}(t, x, u+v) - \frac{b}{2}(u+v)^2 \right) \rho dt dx. \quad (5.1)$$

By (4.5), it follows

$$|\tilde{F}(t, x, u+v) - \frac{b}{2}(u+v)^2| \leq \varepsilon |u+v|^2 + C|u+v|. \quad (5.2)$$

Inserting (5.2) into (5.1), by (4.1), it follows

$$\begin{aligned} \Phi(u+v) &\leq -\frac{\gamma_1}{2} \|u+v\|_E^2 + \int_{\Omega} \left(\varepsilon |u+v|^2 + C|u+v| \right) \rho dt dx \\ &\leq -\frac{\gamma_1}{2} \|u+v\|_E^2 + \varepsilon \|u+v\|_{L^2(\Omega)}^2 + C \|u+v\|_{L^1(\Omega)}, \end{aligned}$$

for some constant C depending on ε .

Taking $\varepsilon = \frac{\delta\gamma_1}{4}$ in above inequality, by (3.2) and (3.3), we obtain

$$\Phi(u+v) \leq -\frac{\gamma_1}{4}\|u+v\|_E^2 + C\|u+v\|_E. \quad (5.3)$$

Therefore, the estimate (5.3) shows there exists $M > 0$ such that $\Phi(u+v) \leq M$ and the assertion (i) is proved.

By the fact $\|u+v\|_E^2 = \|u\|_E^2 + \|v\|_E^2$ and $\|u+v\|_E \leq \|u\|_E + \|v\|_E$, from (5.3), we have

$$\begin{aligned} \Phi(u+v) &\leq -\frac{\gamma_1}{4}\|u\|_E^2 + C\|u\|_E + \left(-\frac{\gamma_1}{4}\|v\|_E^2 + C\|v\|_E\right) \\ &\leq -\frac{\gamma_1}{4}\|u\|_E^2 + C\|u\|_E + C_0, \end{aligned}$$

where $C_0 = \max_{s \geq 0} \{-\frac{\gamma_1}{4}s^2 + Cs\}$. Thus, there exists a constant $\tilde{R} > 0$ such that $\Phi(u+v) \leq 0$ for $\|u\|_E \geq \tilde{R}$. We prove the assertion (ii). \square

Lemma 5.2. *Let the assumptions of Theorem 2.1 hold. Then for any $R_1 > 0$, there exists a constant τ_1 depending on R_1 such that*

$$\tilde{\Phi}(u) \geq \tau_1, \quad \forall u \in E_2 \cap B_{R_1},$$

where $B_{R_1} = \{u \in E : \|u\|_E < R_1\}$.

Proof. For $u \in E_2$, by (3.15), we have

$$\tilde{\Phi}(u) = \max_{v \in E_1} \min_{w \in E_3} \Phi(u+v+w) \geq \min_{w \in E_3} \Phi(u+w),$$

where

$$\Phi(u+w) = \frac{1}{2} \langle (L-a-b)(u+w), u+w \rangle - \int_{\Omega} \left(\tilde{F}(t, x, u+w) - \frac{b}{2}(u+w)^2 \right) \rho dt dx. \quad (5.4)$$

Since E_2 and E_3 are orthogonal subspaces of E , we have

$$\langle (L-a-b)(u+w), u+w \rangle = \langle (L-a-b)u, u \rangle + \langle (L-a-b)w, w \rangle.$$

By (4.2), we have $\langle (L-a-b)w, w \rangle \geq \gamma_2 \|w\|_E^2$. Moreover, noting $\langle (L-a)u, u \rangle = \|u\|_E^2$, by (3.2), we obtain

$$|\langle (L-a-b)u, u \rangle| \leq \|u\|_E^2 + b\|u\|_{L^2(\Omega)}^2 \leq \left(1 + \frac{b}{\delta}\right) \|u\|_E^2 = C_1 \|u\|_E^2,$$

where $C_1 = 1 + \frac{b}{\delta}$. Thus, we have

$$\langle (L - a - b)(u + w), u + w \rangle \geq \gamma_2 \|w\|_E^2 - C_1 \|u\|_E^2. \quad (5.5)$$

On the other hand, a similar calculations as in (5.2) yields

$$|\tilde{F}(t, x, u + w) - \frac{b}{2}(u + w)^2| \leq \varepsilon |u + w|^2 + C|u + w|. \quad (5.6)$$

Inserting (5.5), (5.6) into (5.4) and taking $\varepsilon = \frac{\delta\gamma_2}{4}$, from (3.2) and (3.3), we have

$$\begin{aligned} \Phi(u + w) &\geq \frac{\gamma_2}{2} \|w\|_E^2 - \frac{C_1}{2} \|u\|_E^2 - \frac{\delta\gamma_2}{4} \|u + w\|_{L^2(\Omega)}^2 - C \|u + w\|_{L^1(\Omega)} \\ &\geq \frac{\gamma_2}{2} \|w\|_E^2 - \frac{C_1}{2} \|u\|_E^2 - \frac{\gamma_2}{4} (\|u\|_E^2 + \|w\|_E^2) - C(\|u\|_E + \|w\|_E) \\ &\geq -\left(\frac{C_1}{2} + \frac{\gamma_2}{4}\right) \|u\|_E^2 - C \|u\|_E + C_2, \end{aligned}$$

where $C_2 = \min_{s \geq 0} \left\{ \frac{\gamma_2}{4} s^2 - Cs \right\}$.

For any $R_1 > 0$, the above estimate shows $\tilde{\Phi}(u) \geq \tau_1 = -\left(\frac{C_1}{2} + \frac{\gamma_2}{4}\right) R_1^2 - CR_1 + C_2$ for any $\|u\|_E < R_1$. The proof is completed. \square

Lemma 5.3. *Let the assumptions of Theorem 2.1 hold. Then there exist two constants $\tau_2 > 0$ and $R_2 > 0$ such that $\tilde{\Phi}(u) \geq \tau_2$, for any $u \in E_2$ with $\|u\|_E = R_2$.*

Proof. Firstly, for any $u \in E_2$, $w \in E_3$, by (3.6), we have

$$\Phi(u + w) = \frac{1}{2} \langle (L - a)(u + w), u + w \rangle - \int_{\Omega} \tilde{F}(t, x, u + w) \rho dt dx. \quad (5.7)$$

Since E_2 and E_3 are orthogonal subspaces of E , from (3.10), we obtain

$$\langle (L - a)(u + w), u + w \rangle = \|u\|_E^2 + \|w\|_E^2. \quad (5.8)$$

In addition, it is easy to see

$$\begin{aligned} \int_0^1 \int_0^1 s \frac{\partial \tilde{f}}{\partial \xi}(u + s\theta w) w^2 d\theta ds &= \int_0^1 w \tilde{f}(u + sw) ds - \tilde{f}(u)w \\ &= \tilde{F}(u + w) - \tilde{F}(u) - \tilde{f}(u)w, \end{aligned}$$

where $\tilde{f}(\xi) = \tilde{f}(t, x, \xi)$ and $\tilde{F}(\xi) = \tilde{F}(t, x, \xi)$. Thus,

$$\tilde{F}(u+w) = \int_0^1 \int_0^1 s \frac{\partial \tilde{f}}{\partial \xi}(u+s\theta w) w^2 d\theta ds + \tilde{f}(u)w + \tilde{F}(u). \quad (5.9)$$

In what follows, we estimate the upper bound of $\int_{\Omega} \tilde{F}(u+w) \rho dt dx$. The equation (5.9) shows that it needs to estimate the upper bounds of the following three terms:

$$\int_{\Omega} \left(\int_0^1 \int_0^1 s \frac{\partial \tilde{f}}{\partial \xi}(u+s\theta w) w^2 d\theta ds \right) \rho dt dx, \quad \int_{\Omega} \tilde{f}(u) w \rho dt dx, \quad \int_{\Omega} \tilde{F}(u) \rho dt dx.$$

(i) For $w \in E_3$, the fact $|\lambda_{nm} - a| \geq e$ shows $\|w\|_{L^2(\Omega)}^2 \leq \frac{1}{e} \|w\|_E^2$. Observing $0 \leq \frac{\partial \tilde{f}}{\partial u}(t, x, u) \leq e - \kappa$, we obtain

$$0 \leq \int_{\Omega} \left(\int_0^1 \int_0^1 s \frac{\partial \tilde{f}}{\partial \xi}(u+s\theta w) w^2 d\theta ds \right) \rho dt dx \leq \frac{e - \kappa}{2e} \|w\|_E^2. \quad (5.10)$$

(ii) Fix $r > 1$, by the assumptions (1.5) and (1.6), we obtain that for any $\varepsilon > 0$, there exists a constant $C = C(\varepsilon, r) > 0$ such that

$$|\tilde{f}(u)| \leq \varepsilon |u| + C |u|^r, \quad \forall (t, x, u) \in \Omega \times \mathbb{R}. \quad (5.11)$$

Therefore, a direct calculation yields

$$\begin{aligned} \left| \int_{\Omega} \tilde{f}(u) w \rho dt dx \right| &\leq \varepsilon \|u\|_{L^2(\Omega)} \|w\|_{L^2(\Omega)} + C \|u\|_{L^{2r}(\Omega)}^r \|w\|_{L^2(\Omega)} \\ &\leq \frac{\varepsilon}{2} \|u\|_{L^2(\Omega)}^2 + C \|u\|_{L^{2r}(\Omega)}^{2r} + \varepsilon \|w\|_{L^2(\Omega)}^2. \end{aligned}$$

Since $\dim(E_2) < \infty$, then all norms of E_2 are equivalent. Thus there exists a constant $C > 0$ such that $\|u\|_{L^{2r}(\Omega)}^{2r} \leq C \|u\|_E^{2r}$. Since $\|w\|_{L^2(\Omega)}^2 \leq \frac{1}{e} \|w\|_E^2$, by (3.2), we have

$$\left| \int_{\Omega} \tilde{f}(u) w \rho dt dx \right| \leq \frac{\varepsilon}{2\delta} \|u\|_E^2 + C \|u\|_E^{2r} + \frac{\varepsilon}{e} \|w\|_E^2. \quad (5.12)$$

(iii) Since $\dim(E_2) < \infty$, there exists a constant $C > 0$ such that $\|u\|_{L^{r+1}(\Omega)}^{r+1} \leq C \|u\|_E^{r+1}$. Thus, by (3.2) and (5.11), a direct calculation yields

$$\left| \int_{\Omega} \tilde{F}(u) \rho dt dx \right| \leq \frac{\varepsilon}{2\delta} \|u\|_E^2 + C \|u\|_E^{r+1}, \quad (5.13)$$

for some constant C depending on ε and r .

Therefore, the sum of (5.10), (5.12) and (5.13) yields

$$\begin{aligned} \left| \int_{\Omega} \tilde{F}(u+w) \rho dt dx \right| &\leq \frac{\varepsilon}{\delta} \|u\|_E^2 + C \|u\|_E^{r+1} + C \|u\|_E^{2r} \\ &\quad + \frac{e - \kappa + 2\varepsilon}{2e} \|w\|_E^2. \end{aligned} \quad (5.14)$$

Consequently, inserting (5.8) and (5.14) into (5.7) and taking $\varepsilon = \min\{\frac{\delta}{4}, \frac{\kappa}{4}\}$, we have

$$\begin{aligned} \Phi(u+w) &\geq \left(\frac{1}{2} - \frac{\varepsilon}{\delta}\right) \|u\|_E^2 - C \|u\|_E^{r+1} - C \|u\|_E^{2r} + \left(\frac{1}{2} - \frac{e - \kappa + 2\varepsilon}{2e}\right) \|w\|_E^2 \\ &= \left(\frac{1}{2} - \frac{\varepsilon}{\delta}\right) \|u\|_E^2 - C \|u\|_E^{r+1} - C \|u\|_E^{2r} + \left(\frac{\kappa - 2\varepsilon}{2e}\right) \|w\|_E^2. \\ &\geq \frac{1}{4} \|u\|_E^2 - C \|u\|_E^{r+1} - C \|u\|_E^{2r}. \end{aligned} \quad (5.15)$$

Finally, since $r > 1$, then $\phi(s) = \frac{1}{4}s^2 - Cs^{r+1} - Cs^{2r}$ attains the local minimum at $s = 0$. Therefore there exist two constants $R_2 > 0$ and $\tau_2 > 0$ such that $\phi(R_2) \geq \tau_2$. Since $\tilde{\Phi}(u) \geq \min_{w \in E_3} \Phi(u+w)$, from (5.15), we obtain $\tilde{\Phi}(u) \geq \tau_2$ with $\|u\|_E = R_2$. \square

6. Proof of Theorem 2.1

With the above lemmas in hand, we give the proof of Theorem 2.1.

Proof. Let R_2 and τ_2 be the constants in Lemma 5.3 and $B_{R_2} = \{u \in E_2 : \|u\|_E < R_2\}$.

Firstly, by Lemma 5.1, we know $\tilde{\Phi}$ is bounded from above. Let $\beta_1 = \sup_{u \in E_2} \tilde{\Phi}(u)$. By Lemma 3.1 and Lemma 4.3, $\tilde{\Phi}$ satisfies $(PS)_{\beta_1}$ condition.

Thus, there exists a critical point $u_1 \in E_2$ such that $\tilde{\Phi}(u_1) = \beta_1$ and $\tilde{\Phi}'(u_1) = 0$.

Secondly, recall $\tilde{F}(t, r, u) \geq 0$ for any $u \in E$, then for any $v \in E_1$, from (3.9), we obtain

$$\Phi(v) = \frac{1}{2} \langle (L - a)v, v \rangle - \int_{\Omega} \tilde{F}(t, x, v) \rho dt dx \leq 0.$$

Therefore, we have

$$\tilde{\Phi}(0) = \min_{w \in E_3} \max_{v \in E_1} \Phi(v + w) \leq \max_{v \in E_1} \Phi(v) \leq 0. \quad (6.1)$$

Noting $0 \in B_{R_2}$ and taking $R_1 = R_2$ in Lemma 5.2, in virtue of Lemma 5.2, Lemma 5.3 and $\tilde{\Phi}(0) \leq 0$, we obtain the reduction functional $\tilde{\Phi}$ attains its infimum in B_{R_2} . Let $\beta_2 = \inf_{u \in B_{R_2}} \tilde{\Phi}(u)$, then Lemma 3.1 and Lemma 4.3 show

that $\tilde{\Phi}$ satisfies the $(PS)_{\beta_2}$ condition. Thus, there exists $u_2 \in E_2$ such that $\tilde{\Phi}'(u_2) = 0$ and $\tilde{\Phi}(u_2) = \beta_2$.

Now we claim u_1 and u_2 are two different critical points in E_2 . Observing $0 \in B_{R_2}$, the following inequality

$$\beta_2 = \inf_{u \in B_{R_2}} \tilde{\Phi}(u) \leq \tilde{\Phi}(0) \leq 0 < \tau_2 \leq \inf_{\|u\|_E=R_2} \tilde{\Phi}(u) \leq \sup_{u \in E_2} \tilde{\Phi}(u) = \beta_1 \quad (6.2)$$

shows $u_1 \neq u_2$.

If $\tilde{\Phi}$ possesses another local maximum point which is different from u_1 , then there exist at least three critical points of $\tilde{\Phi}$.

Otherwise, if u_1 is the unique local maximum point of $\tilde{\Phi}$. We shall prove that there exists a critical point of mountain-pass type which is different from u_1 and u_2 .

Taking $u_0 \in E_2$ with $\|u_0\|_E = 1$, by Lemma 5.1, there exists $\tilde{R} > R_2$ such that $\tilde{\Phi}(\tilde{R}u_0) \leq 0$. Moreover, one of the fact holds: for any $s \in [0, 1]$, either $s\tilde{R}u_0 \neq u_1$ or $-s\tilde{R}u_0 \neq u_1$.

Case 1: if $s\tilde{R}u_0 \neq u_1$ for any $s \in [0, 1]$, by Lemma 5.3 and (6.1), we obtain

$$\max\{\tilde{\Phi}(0), \tilde{\Phi}(\tilde{R}u_0)\} \leq 0 < \tau_2 \leq \inf_{\|u\|_E=R_2} \tilde{\Phi}(u).$$

Let

$$c^+ = \inf_{g \in \Delta^+} \max_{s \in [0, 1]} \tilde{\Phi}(g(s)),$$

where $\Delta^+ = \{g \in C([0, 1], E_2) : g(0) = 0, g(1) = \tilde{R}u_0\}$. In addition, we have $\tilde{\Phi}$ is C^1 function and satisfies $(PS)_{c^+}$ condition. By mountain pass lemma [10], we obtain a critical point u_3^+ satisfying $\tilde{\Phi}(u_3^+) = c^+$ and $\tilde{\Phi}'(u_3^+) = 0$. Obviously, $c^+ \geq \tau_2 > 0$.

It is easy to see that $s\tilde{R}u_0 \in \Delta^+$ for any $s \in [0, 1]$. Therefore, the assumption $s\tilde{R}u_0 \neq u_1$ implies

$$c^+ \leq \max_{s \in [0, 1]} \tilde{\Phi}(s\tilde{R}u_0) < \sup_{u \in E_2} \tilde{\Phi}(u) = \beta_1.$$

Thus, the following inequalities

$$\beta_2 = \inf_{u \in B_{R_2}} \tilde{\Phi}(u) \leq 0 < \tau_2 \leq c^+ < \sup_{u \in E_2} \tilde{\Phi}(u) = \beta_1$$

shows $u_1 \neq u_2 \neq u_3^+$.

Case 2: similarly, if $-s\tilde{R}u_0 \neq u_1$ for any $s \in [0, 1]$, we have

$$c^- = \inf_{g \in \Delta^-} \max_{s \in [0, 1]} \tilde{\Phi}(g(s))$$

is the critical value of $\tilde{\Phi}$, where $\Delta^- = \{g \in C([0, 1], E_2) : g(0) = 0, g(1) = -\tilde{R}u_0\}$. Therefore, there exists $u_3^- \in E_2$ satisfying $\tilde{\Phi}(u_3^-) = c^-$ and $\tilde{\Phi}'(u_3^-) = 0$. Furthermore, we have $u_1 \neq u_2 \neq u_3^-$.

Consequently, by Lemma 3.1, we obtain the functional Φ has at least three critical points. The proof is completed. \square

Ethics statement. This work did not involve any active collection of human data.

Data accessibility statement. This work does not have any experimental data.

Competing interests statement. We have no competing interests.

Authors contributions. HW and SJ contributed to the mathematical proof and writing the paper. All authors gave final approval for publication.

Funding. This work was supported by NSFC Grants (nos. 11322105 and 11671071).

References

References

- [1] H. Amann, Saddle points and multiple solutions of differential equations, Math. Z. 169 (1979) 127–166.

- [2] D. Arcoya, D.G. Costa, Nontrivial solutions for a strongly resonant problem, *Differential Integral Equations* 8 (1995) 151–159.
- [3] H. Brézis, Periodic solutions of nonlinear vibrating strings and duality principles, *Bull. Amer. Math. Soc. (N.S.)* 8 (1983) 409–426.
- [4] A.K. Ben-Naoum, J. Berkovits, On the existence of periodic solutions for semilinear wave equation on a ball in \mathbb{R}^n with the space dimension n odd, *Nonlinear Anal.* 24 (1995) 241–250.
- [5] H. Brézis, L. Nirenberg, Forced vibrations for a nonlinear wave equation, *Comm. Pure Appl. Math.* 31 (1978) 1–30.
- [6] V. Barbu, N.H. Pavel, Periodic solutions to one-dimensional wave equation with piece-wise constant coefficients, *J. Differential Equations* 132 (1996) 319–337.
- [7] V. Barbu, N.H. Pavel, Periodic solutions to nonlinear one dimensional wave equation with x -dependent coefficients, *Trans. Amer. Math. Soc.* 349 (1997) 2035–2048.
- [8] V. Barbu, N.H. Pavel, Determining the acoustic impedance in the 1-D wave equation via an optimal control problem, *SIAM J. Control Optim.* 35 (1997) 1544–1556.
- [9] K. Chang, Solutions of asymptotically linear operator equations via Morse theory, *Comm. Pure Appl. Math.* 34 (1981) 693–712.
- [10] K. Chang, *Critical Point Theory and Its Applications*, Shanghai Scientific and Technical Publishers, Shanghai, 1986 (in Chinese).
- [11] K. Chang, S. Wu, S. Li, Multiple periodic solutions for an asymptotically linear wave equation, *Indiana Univ. Math. J.* 31 (1982) 721–731.
- [12] A. Castro, A.C. Lazer, Critical point theory and the number of solutions of a nonlinear Dirichlet problem, *Ann. Mat. Pura Appl.* 120 (1979) 113–137.
- [13] J. Chen, Z. Zhang, Infinitely many periodic solutions for a semilinear wave equation in a ball in \mathbb{R}^n , *J. Differential Equations* 256 (2014) 1718–1734.

- [14] J. Chen, Z. Zhang, Existence of infinitely many periodic solutions for the radially symmetric wave equation with resonance, *J. Differential Equations* 260 (2016) 6017–6037.
- [15] J. Chen, Z. Zhang, Existence of multiple periodic solutions to asymptotically linear wave equations in a ball, *Calc. Var. Partial Differential Equations* 56 (2017) 58.
- [16] W. Craig, C.E. Wayne, Newton’s method and periodic solutions of nonlinear wave equations, *Comm. Pure Appl. Math.* 46 (1993) 1409–1498.
- [17] M. Izydorek, Multiple solutions for an asymptotically linear wave equation, *Differential Integral Equations* 13 (2000) 289–310.
- [18] S. Ji, Time periodic solutions to a nonlinear wave equation with x -dependent coefficients, *Calc. Var. Partial Differential Equations* 32 (2008) 137–153.
- [19] S. Ji, Time-periodic solutions to a nonlinear wave equation with periodic or anti-periodic boundary conditions, *Proc. R. Soc. Lond. Ser. A* 465 (2009) 895–913.
- [20] S. Ji, Y. Gao, W. Zhu, Existence and multiplicity of periodic solutions for Dirichlet-Neumann boundary value problem of a variable coefficient wave equation, *Adv. Nonlinear Stud.* 16 (2016) 765–773.
- [21] S. Ji, Y. Li, Periodic solutions to one-dimensional wave equation with x -dependent coefficients, *J. Differential Equations* 229 (2006) 466–493.
- [22] S. Ji, Y. Li, Time-periodic solutions to the one-dimensional wave equation with periodic or anti-periodic boundary conditions, *Proc. Roy. Soc. Edinburgh Sect. A* 137 (2007) 349–371.
- [23] S. Ji, Y. Li, Time periodic solutions to the one-dimensional nonlinear wave equation, *Arch. Ration. Mech. Anal.* 199 (2011) 435–451.
- [24] P.H. Rabinowitz, Periodic solutions of nonlinear hyperbolic partial differential equations, *Comm. Pure Appl. Math.* 20 (1967) 145–205.
- [25] P.H. Rabinowitz, Free vibrations for a semilinear wave equation, *Comm. Pure Appl. Math.* 31 (1978) 31–68.

- [26] P.H. Rabinowitz, Large amplitude time periodic solutions of a semilinear wave equation, *Comm. Pure Appl. Math.* 37 (1984) 189–206.
- [27] I.A. Rudakov, Periodic solutions of a nonlinear wave equation with non-constant coefficients, *Math. Notes* 76 (2004) 395–406.
- [28] I.A. Rudakov, Periodic solutions of the quasilinear equation of forced vibrations of an inhomogeneous string, *Math. Notes* 101 (2017) 137–148.
- [29] M. Schechter, Rotationally invariant periodic solutions of semilinear wave equations, *Abstr. Appl. Anal.* 3 (1998) 171–180.
- [30] M. Tanaka, Existence of multiple weak solutions for asymptotically linear wave equations, *Nonlinear Anal.* 65 (2006) 475–499.
- [31] P. Wang, Y. An, Resonance in nonlinear wave equations with x -dependent coefficients, *Nonlinear Anal.* 71 (2009) 1985–1994.
- [32] K. Yosida, *Functional Analysis*, sixth ed., Springer-Verlag, Berlin, 1980.