A REFORMULATION OF THE SIEGEL SERIES AND INTERSECTION NUMBERS

SUNGMUN CHO AND TAKUYA YAMAUCHI

Abstract. In this paper, we will explain a conceptual reformulation and inductive formula of the Siegel series. Using this, we will explain that both sides of the local intersection multiplicities of [\[GK93\]](#page-50-0) and the Siegel series have the same inherent structures, beyond matching values.

As an application, we will prove a new identity between the intersection number of two modular correspondences over \mathbb{F}_p and the sum of the Fourier coefficients of the Siegel-Eisenstein series for Sp_4/\mathbb{Q} of weight 2, which is independent of $p (> 2)$. In addition, we will explain a description of the local intersection multiplicities of the special cycles over \mathbb{F}_p on the supersingular locus of the 'special fiber' of the Shimura varieties for $\text{GSpin}(n,2)$, $n \leq 3$ in terms of the Siegel series directly.

CONTENTS

²⁰¹⁰ Mathematics Subject Classification. MSC 11F30, 11F46, 11G18, 14C17, 14G35, 14J15.

Key words and phrases. Siegel series, Gross-Keating invariant, special cycles on an orthogonal Shimura variety.

The first author was supported by JSPS KAKENHI Grant No. 16F16316, Samsung Science and Technology Foundation under Project Number SSTF-BA1802-03, and NRF 2018R1A4A 1023590.

The second author is partially supported by JSPS KAKENHI Grant Number (B) No.19H01778.

1. Introduction

1.1. On the Gross-Keating's formula. In [\[GK93\]](#page-50-0) Gross and Keating studied the arithmetic intersection number of three modular correspondences which are regarded as cycles in the self-product $Y_0(1) \times Y_0(1)/\mathbb{Z}$ of the moduli stack $Y_0(1)$ of elliptic curves over any scheme. They described it purely in terms of certain invariants of a ternary quadratic form created by themselves to formulate it. This invariant has been generalized to quadratic forms of any degree over a local field, and is nowadays called the Gross-Keating invariant. They already expected in the introduction of their paper that the arithmetic intersection number coincides with the sum of the Fourier coefficients of the derivative of the Siegel-Eisenstein series of weight 2 and of degree 3, which has been studied thoroughly in [\[ARGOS07\]](#page-50-2). Kudla in [\[Kud97\]](#page-50-3) later proposed a general program (local version) to make a connection between the local intersection multiplicity of special cycles on an integral model of the Shimura varieties for $\text{GSpin}(n, 2)$ and the derivative of the local factor of a Fourier coefficient of the Siegel-Eisenstein series of weight $(n+2)/2$ and of degree $n+1$. In the latter object, such local factor is called the Siegel series.

The program has been vastly studied by a series of papers [\[Kud97\]](#page-50-3), [\[KRY99\]](#page-50-4),[\[KRY06\]](#page-50-5),[\[KR99\]](#page-50-6),[\[KR00\]](#page-50-7) for $0 \le n \le 3$, and [\[BY\]](#page-50-8) for general n, when the dimension of the arithmetic intersection is zero. A strategy to compute the local intersection multiplicities which had been taken in these papers is to reduce them to the case of Gross-Keating. On the other hand, a computation of the Siegel series side is based on Katsurada's paper [\[Kat99\]](#page-50-9). The relation between both sides then follows by a direct comparison. Note that beyond direct comparison, there had not been known an evidence or structure to conceptually yield the equality between them.

Therefore, in order to understand Kudla's program in conceptual way toward the higher dimensional case of the arithmetic intersection, it would be important to have a better understanding on the relation between the local intersection multiplicity of Gross and Keating in [\[GK93\]](#page-50-0) and the Siegel series. In [\[GK93\]](#page-50-0), a key step is to derive an inductive formula (Lemma 5.6 of [\[GK93\]](#page-50-0)) for the local intersection multiplicities over $W(\overline{\mathbb{F}}_p)$ at a prime p, which involves the local intersection multiplicity on the special fiber at p (Lemma 5.7 of [\[GK93\]](#page-50-0)). As we will compare this inductive formula with our result in the next subsection, we describe the precise form of their inductive formula.

Let (L, q_L) be an anisotropic quadratic lattice over \mathbb{Z}_p of rank 3. Then the Gross-Keating invariant of (L, q_L) consists of three integers $GK(L) = (a_1, a_2, a_3)$ with $a_1 \le a_2 \le a_3$. If we denote by $\alpha_p(a_1, a_2, a_3)$ the local intersection multiplicity associated to (L, q_L) (see (3.18) of [\[GK93\]](#page-50-0)), then it satisfies the following inductive formula, with respect to the Gross-Keating invariant: (cf. the proof of Theorem [6.8\)](#page-33-1):

(1.1)
$$
\alpha_p(a_1, a_2, a_3) = \alpha_p(a_1, a_2, a_3 - 2) + \mathcal{T}_{a_1, a_2}.
$$

Here, \mathcal{T}_{a_1,a_2} is the local intersection multiplicity of two cycles on the special fiber of the setting of Gross-Keating.

1.2. On the Siegel series. On the other hand, the Siegel series is of great importance in automorphic forms, such as the study of conjectures related to automorphic L-functions and the construction of automorphic forms of level 1, so-called Ikeda lift. We refer to the first several paragraphs of [\[IK2\]](#page-50-10) for more introductory discussion about the important usages of the Siegel series in this context. Theories of the Siegel series have been developed by many people such as Kitaoka and Shimura. It was Katsurada in [\[Kat99\]](#page-50-9) who firstly found the explicit formula of the Siegel series for \mathbb{Z}_p . However, as mentioned in the introduction of [\[IK2\]](#page-50-10), his formula is complicated and it is not clear which invariant of a quadratic form determines the Siegel series.

Recently, Ikeda and Katsurada in [\[IK2\]](#page-50-10) obtained the formula of the Siegel series over any finite extension of \mathbb{Z}_p . Furthermore, they show that the Siegel series is completely determined by the Gross-Keating invariant with extra data, called the Extended Gross-Keating datum, for any quadratic form over any finite extension of \mathbb{Z}_p .

The Siegel series is usually described as a polynomial. An explicit formula of the Siegel series given in [\[Kat99\]](#page-50-9) and [\[IK2\]](#page-50-10) is to determine the coefficients of this polynomial. On the other hand, theoretical interpretation of these coefficients had not been known yet.

1.3. Reformulation of the Siegel series. Our first main result is to reformulate the Siegel series over any finite extension of \mathbb{Z}_p in Theorem [3.14](#page-16-0) and Corollary [3.16.](#page-17-1) The Siegel series can be defined as an integral of certain volume form on a p -adic manifold (cf. Definitions [3.3](#page-11-0) and [3.15\)](#page-16-1) associated to a quadratic lattice (L, q_L) over \mathfrak{o} , where \mathfrak{o} is the ring of integers of a finite field extension of \mathbb{Q}_p (for any p). It is usually denoted by $\mathcal{F}_L(X)$, as a polynomial of X. Let n be the rank of L so that the Gross-Keating invariant consists of n-integers $GK(L) = (a_1, \dots, a_n)$ satisfying $a_i \leq a_j$ with $i \leq j$.

Theorem 1.1. (Corollary [3.16\)](#page-17-1) we have the following description of the Siegel series:

$$
\mathcal{F}_L(X) = \n\left(1 - X\right) \cdot \sum_{\substack{0 \le b \le \frac{|{\rm GK}(L)|}{n_0 \le a \le n}}} \left(\# \mathcal{S}_{(L, a^{\pm}, b)} \cdot f^{b \cdot (n+1)} X^{2b} \cdot \left(1 + \chi(a^{\pm}) f^{n-a/2} X\right) \prod_{1 \le i < n-a/2} \left(1 - f^{2i} X^2\right) \right).
$$

Here, $|GK(L)| = a_1 + \cdots + a_n$, n_0 is the number of 0's in $GK(L) = (a_1, \dots, a_n)$ (cf. Proposition [3.13\)](#page-15-1), f is the cardinality of the residue field of \mathfrak{o}, χ is defined just below Equation [\(3.8\)](#page-16-2), and $\mathcal{S}_{(L,a^{\pm},b)}$ is the set of certain (depending on two integers a^{\pm} and b) quadratic lattices containing L, which can be found at Equation [\(3.6\)](#page-16-3).

This gives a conceptual and theoretical interpretation of the coefficients of $\mathcal{F}_L(X)$ as a weighted sum of certain number of quadratic lattices. The method used in Theorem [1.1](#page-2-1) (Corollary [3.16\)](#page-17-1) is based on another geometric nature of the Siegel series involving the stratification of a p-adic scheme, geometric description of each stratum, Grassmannian, and lattice counting argument. This is largely different from the known techniques in this context^{[1](#page-2-2)}.

Using the result of [\[IK2\]](#page-50-10), we then obtain an inductive formula of the Siegel series, with respect to the Gross-Keating invariant, under Conjecture [4.4](#page-19-0) concerning about quadratic forms (which is verified to be true when p is odd or when (L, q_L) is anisotropic over \mathbb{Z}_2 in Lemmas [4.5-](#page-19-1)[4.6\)](#page-20-0).

We describe our inductive formula more precisely. If we choose the integer d characterized by the condition $a_{n-d} < a_{n-d+1} = \cdots = a_n$ \overbrace{d} , then we can associate certain lattice $L^{(d,n)}$ containing L

whose rank is also *n*. To be more precise, for a reduced basis (e_1, \dots, e_n) of L given in Definition [2.5,](#page-8-0) the lattice $L^{(d,n)}$ is spanned by $(e_1, \dots, e_{n-d}, \frac{1}{\pi})$ $\frac{1}{\pi} \cdot e_{n-d+1}, \cdots, \frac{1}{\pi}$ $\frac{1}{\pi} \cdot e_n$ \overbrace{d}). Here, π is a uniformizer

in o. A second main theorem of the current paper is the following:

¹Remark in page 444 of [\[Kat99\]](#page-50-9) says 'it seems very interesting problem to prove Theorems 4.1 and 4.2 directly from the local theory of quadratic forms'. Here, Theorems 4.1 and 4.2 are main results of [\[Kat99\]](#page-50-9), which give an explicit formula of the Siegel series over \mathbb{Z}_p . Our method can be understood in the spirit of the problem proposed by Katsurada.

Theorem 1.2. (Theorem [4.9\)](#page-23-0) Assume that Conjecture [4.4](#page-19-0) is true. If $L^{(d,n)}$ is a quadratic lattice, then we have the following inductive formula, with respect to the Gross-Keating invariant, of the Siegel series $\mathcal{F}_L(X)$:

$$
\mathcal{F}_L(X) = \sum_{m=1}^d \left(c_m \cdot f^{(n+1)m} \cdot X^{2m} \cdot \sum_{L' \in \mathcal{G}_{L,d,m}} \mathcal{F}_{L'}(X) \right) +
$$

$$
(1-X)(1-f^dX)^{-1} \cdot \left(\prod_{i=1}^d (1-f^{2i}X^2) \right) \cdot \mathcal{F}_{L_0^{(d,n)}}(f^dX),
$$

where $c_m = (-1)^{m-1} f^{m(m-1)/2}$. Here, f is the cardinality of the residue field of **o**. For $L' \in \mathcal{G}_{L,d,m}$,

$$
\left\{\begin{array}{l}\operatorname{GK}(L)\succ\operatorname{GK}(L');\\ \left|\operatorname{GK}(L')\right|=\left|\operatorname{GK}(L)\right|-2m;\\ \operatorname{GK}(L_0^{(d,n)})=\operatorname{GK}(L)^{(n-d)}.\end{array}\right.
$$

Note that notion of $L^{(d,n)}$, $\mathcal{G}_{L,d,m}$, and $\binom{m}{k}_f$ can be found at the beginning of Section [4.](#page-17-0) Notion of $L_0^{(d,n)}$ $a_0^{(a,n)}$ can be found at Remark 4.3.(1).

Here, $\mathcal{G}_{L,d,m}$ is identified with Grassmannian to classify the set of m-dimensional subspaces of the vector space of dimension d (given by $L^{(d,n)}/L$) over a finite field $\mathfrak{o}/(\pi)$, whose order is $\binom{d}{n}$ $\binom{d}{m}$ f.

1.4. The comparison between Gross-Keating's formula and the Siegel series. Since both sides, Gross-Keating's formula and the Siegel series, have inductive formulas, it is natural to ask whether or not there is a relation between them.

If we restrict ourselves to an anisotropic quadratic lattice over \mathbb{Z}_p of rank n, which covers the case of Gross-Keating, then we have more refined and simpler inductive formula (Theorem [5.11\)](#page-29-1) as follows:

(1.2)
$$
\mathcal{F}_L(X) = \begin{cases} p^{n+1} \cdot X^2 \cdot \mathcal{F}_{L^{(n)}}(X) + (1-X)(1+pX) \cdot \mathcal{F}_{L_0^{(n)}}(pX) & \text{if } 2 \le n \le 4; \\ p^2 \cdot X^2 \cdot \mathcal{F}_{L^{(1)}}(X) + (1-X)(1+pX) & \text{if } n = 1. \end{cases}
$$

After comparing both inductive formulas, we obtain the following result:

Theorem 1.3. (cf. Theorems [6.7](#page-32-0) and [6.8\)](#page-33-1) The inductive formula of Gross-Keating in Equation [\(1.1\)](#page-1-3) and the derivative of Equation [\(1.2\)](#page-3-2) at $1/p^2$ with $n = 3$ do match each other. As a direct consequence, we have

$$
\begin{cases}\n\alpha_p(a_1, a_2, a_3) = c_1 \cdot \mathcal{F}_L'(1/p^2); \\
\mathcal{T}_{a_1, a_2} = c_2 \cdot \mathcal{F}_{L_0^{(3)}}'(1/p),\n\end{cases}
$$

for explicitly computed constants c_1 and c_2 . Here $GK(L_0^{(3)})$ $\binom{0}{0} = (a_1, a_2).$

This theorem shows that both sides of the local intersection multiplicity and the Siegel series have the same inherent structures, that is, the same inductive formula, beyond matching their values. In addition, it gives us a new observation that the local intersection multiplicity on the special fiber can also be described in terms of the derivative of the Siegel series.

1.5. Applications to intersection numbers over finite fields. Since the above theorem matches both sides on special fibers, we can naturally consider their applications over finite fields. We explain two consequences in this line: intersection numbers over finite fields and the local intersection multiplicities on special fibers of $\text{GSpin}(n,2)$ Shimura varieties with $n \leq 3$ (in the case of zero dimension of the arithmetic intersection).

1.5.1. Intersection numbers over finite fields. Since we obtained a new description of the local intersection multiplicity on the special fiber in the setting of Gross-Keating in terms of the derivative of the Siegel series, it is natural to compute the intersection numbers of two modular correspondences on finite fields.

More precisely, let φ_m be the modular polynomial in $\mathbb{Z}[x, y]$ of degree m whose irreducible factor corresponds to an affine model of the modular curves $Y_0(m/n^2)$ for some $n^2|m$ in $Y_0(1) \times Y_0(1)$ (cf. [\[Vog07\]](#page-51-0)). Then for positive integers m_1, m_2 and a prime p, we define the intersection number over the finite field \mathbb{F}_p or over the complex field as follows:

$$
(1.3) (T_{m_1,p}, T_{m_2,p}) := \text{length}_{\mathbb{F}_p} \mathbb{F}_p[x, y]/(\varphi_{m_1}, \varphi_{m_2}), (T_{m_1,\mathbb{C}}, T_{m_2,\mathbb{C}}) := \text{length}_{\mathbb{C}} \mathbb{C}[x, y]/(\varphi_{m_1}, \varphi_{m_2}).
$$

We compare the above two intersection numbers by using Theorem [1.3](#page-3-3) on the supersingular locus and the theory of quasi-canonical lifts on the ordinary locus. The following theorem is our result:

Theorem 1.4. (Proposition [7.1](#page-34-1) and Theorem [7.3\)](#page-34-2) The intersection number $(T_{m_1,p}, T_{m_2,p})$ is finite if and only if m_1m_2 is not a square. In addition, if m_1m_2 is not a square and p is odd, then

$$
(T_{m_1,p}, T_{m_2,p}) = (T_{m_1,\mathbb{C}}, T_{m_2,\mathbb{C}}).
$$

We refer to Remark 7.17 for a discussion with $p = 2$.

Since $(T_{m_1,\mathbb{C}}, T_{m_2,\mathbb{C}})$ is the sum of the Fourier coefficients of the Siegel-Eisenstein series for Sp_4/\mathbb{Q} by Proposition 2.4 of [\[GK93\]](#page-50-0), the intersection number on the special fiber is also the sum of the Fourier coefficients of the Siegel-Eisenstein series for $Sp_4\mathbb{Q}$. Furthermore the intersection number is independent of the characteristic of a finite field with $p > 2$, whereas the local intersection multiplicities highly depend on p.

The above theorem yields a new interpretation on a classical object $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}$ [x, y]/($\varphi_{m_1}, \varphi_{m_2}$). Note that the two main objects to be analyzed in [\[GK93\]](#page-50-0) are geometric interpretations of

 $\mathbb{C}[x,y]/(\varphi_{m_1}, \varphi_{m_2})$ and $\mathbb{Z}[x,y]/(\varphi_{m_1}, \varphi_{m_2}, \varphi_{m_3}).$

Namely, the dimension of the first object is the intersection number of two modular correspondences over C and (log of) the cardinality of the second object is the arithmetic intersection number of three modular correspondences over \mathbb{Z} . In this context, the \mathbb{Z} -module $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}$ [x, y]/($\varphi_{m_1}, \varphi_{m_2}$) has the following interesting interpretations:

Theorem 1.5. *(Theorem [7.4\)](#page-35-1)*

 (1) $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}][x,y]/(\varphi_{m_1}, \varphi_{m_2})$ is a free $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}$]-module. (2) The rank of $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}][x,y]/(\varphi_{m_1}, \varphi_{m_2}),$ as a $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}$]-module, is equal to 1 288 $\overline{}$ $c(T).$

$$
\begin{array}{cc}\n\text{SO} & T \in \text{Sym}_2(\mathbb{Z}) > 0 \\
\text{diag}(T) = (m_1, m_2)\n\end{array}
$$

Here, $c(T)$ is the Fourier coefficient of the Siegel-Eisenstein series for $Sp_4(\mathbb{Z})$ of weight 2 with respect to the (2×2) - half-integral symmetric matrix T.

1.5.2. The local intersection multiplicities on the special fiber in orthogonal Shimura varieties. Let p be an odd prime. Let T' be a positive definite half-integral symmetric matrix over $\mathbb{Z}_{(p)}$ of size $n+1$ for a non-negative integer n . As explained in subsection [1.1,](#page-1-1) the local intersection multiplicity on the arithmetic intersection $\mathcal{Z} = \mathcal{Z}(T')$ of some special cycles associated to T' on an integral model M over $\mathbb{Z}_{(p)}$ of Shimura varieties for $G = \text{GSpin}(n, 2)$, when the dimension of $\mathcal Z$ is zero, is reduced to that of Gross-Keating. We refer Section [8](#page-45-0) for the notation and the detailed explanation. For any geometric point ξ on $\mathcal Z$ let us define the local intersection multiplicity at $\xi: e(\mathcal Z, \xi) = \text{length}_{\mathbb Z_{(p)}} \mathcal O_{\mathcal Z, \xi}.$ Similarly for any positive definite half-integral symmetric matrix T over $\mathbb{Z}_{(p)}$ of size n one can also define the intersection $\mathcal{Z}(T)_{\mathbb{F}_p}$ of some special cycles associated to T on $\mathcal{M}_{\mathbb{F}_p}$. For any geometric ξ on

 $\mathcal{Z}(T)_{\mathbb{F}_p}$ let us define the local intersection multiplicity at $\xi: e(\mathcal{Z}(T)_{\mathbb{F}_p}, \xi) = \text{length}_{\mathbb{F}_p} \mathcal{O}_{\mathcal{Z}(T)_{\mathbb{F}_p}, \xi}$. Let (a_0, \ldots, a_n) be the GK-invariant of $T' \otimes \mathbb{Z}_p$. It turns out that $e(\mathcal{Z}, \xi)$ depends only on (a_0, \ldots, a_n) and therefore we may put $e(\mathcal{Z}, \xi) = e(a_0, \ldots, a_n)$. Further when $\mathcal{Z}(T')$ is of dimension zero, the initial integer a_0 has to be zero and we will see that $e(a_0, \ldots, a_n) = \alpha_p(0, \ldots, 0, a_1, \ldots, a_n)$ where the zeros in front of a_1 are added until the number of all entries are 3.

In this context, we obtain the following result:

Theorem 1.6. (Proposition [8.3](#page-48-0) and Theorem [8.4\)](#page-49-1) Assume that p is odd and $0 \le n \le 3$.

- (1) Assume that $\mathcal{Z}(T)$ is of dimension zero. Then the local intersection multiplicity $e(\mathcal{Z}, \xi)$ has the same inductive formula which are parallel to the inductive formula induced from Equation (1.[2\)](#page-3-2).
- (2) Assume that T represents 1. Then $\mathcal{Z}(T)_{\mathbb{F}_p}$ is of dimension zero and for any geometric point ξ on $\mathcal{Z}(T)_{\mathbb{F}_p}$, the local intersection multiplicity $e(\mathcal{Z}(T)_{\mathbb{F}_p}, \xi)$ is described in terms of the derivative of the Siegel series for a suitable anisotropic quadratic lattice of rank 2.

Since we have a better understanding on the latter objects that are Siegel series, it is natural to ask if our comparison argument between two inductive formulas in Gross-Keating's case can be extended to the general case.

1.6. Speculation.

1.6.1. As Kudla expected, the local intersection multiplicity in higher dimensional case of the arithmetic intersection on $GSpin(n, 2)$ Shimura varieties is believed to match with the derivative of the Siegel series for a suitable quadratic lattice. The comparison results of Theorem [1.3](#page-3-3) and Theorem [1.6.](#page-5-2)(1) seem to imply that there should be an inductive formula in geometric side which is parallel to that on the Siegel series side. Thus our inductive formula of the Siegel series given in Theorem [1.2](#page-3-4) would be the inductive formula that the local intersection multiplicity is expected to satisfy with.

Recently, Chao Li and Wei Zhang proved the Kudla-Rapoport conjecture in [\[LZ\]](#page-50-11), which is a unitary version of the above Kudla's conjecture. Their proof relies on the principle to compare inductive structures of both sides, rather than to compute them directly.

1.6.2. Since we have an interpretation of the local intersection multiplicity on (the supersingular locus of) the special fiber of $GSpin(n, 2)$ Shimura varieties in terms of the Siegel series in Theorem [1.6,](#page-5-2) we will be able to relate the intersection numbers on the special fiber of $GSpin(n, 2)$ Shimura varieties with the sum of the Fourier coefficients of the Siegel-Eisenstein series with suitable weight and degree. We expect that the intersection number of the special cycles at the special fiber of $\text{GSpin}(n, 2)$ -Shimura variety is independent of p (possibly away from bad primes), which turns to be the sum of the Fourier coefficients of the Siegel-Eisenstein series. This observation is parallel to Theorems [1.4](#page-4-0)[-1.5.](#page-4-1) This would imply that an associated arithmetic intersection is flat over $\mathbb Z$ (possibly away from bad primes).

1.7. Organizations. We will organize this paper as follows. After fixing notations in Section [2,](#page-6-0) we will derive a conceptual study of the Siegel series in Sections [3](#page-10-0)[-4.](#page-17-0) In Section [5,](#page-24-0) we will explain a refined formulation of the Siegel series for anisotropic quadratic lattices over \mathbb{Z}_p . Section [6](#page-29-0) is devoted to compare both sides of the local intersection multiplicity of [\[GK93\]](#page-50-0) and the Siegel series. In Sections [7-](#page-33-0)[8,](#page-45-0) we will explain two applications in the context of intersection numbers (or multiplicities) over a finite field. In Appendix, we list up explicit examples for the intersection numbers related to Section [7.](#page-33-0)

Acknowledgments. We would like to express our deep appreciation to Professors T. Ikeda and H. Katsurada for many fruitful discussions and suggestions. We also thank Professors B. Conrad, B. Gross, B. Howard, R. Schulze-Pillot, and W. Zhang for helpful discussions and corrections in

Theorems [1.4](#page-4-0) and [4.9.](#page-23-0) Special thanks are own to Professor S. Yokoyama for computing intersection numbers for modular correspondences over both a finite field and the complex field in Appendix and also to Professor Chul-hee Lee for pointing out our mistakes on the table in Appendix. Finally we thank the referee for helpful suggestions and comments which substantially helped with the presentation of our paper.

2. Notations

- • Let F be a finite field extension of \mathbb{Q}_p with \mathfrak{o} its ring of integers and κ its residue field. Let π be a uniformizer in **o**. Let f be the cardinality of the finite field $κ$.
- For an element $x \in F$, the exponential order of x with respect to the maximal ideal in $\mathfrak o$ is written by $\mathrm{ord}(x)$.
- Let $e = \text{ord}(2)$. Thus if p is odd, then $e = 0$.
- We consider an $\mathfrak{o}\text{-lattice } L$ with a quadratic form $q_L : L \to \mathfrak{o}$. Here, an $\mathfrak{o}\text{-lattice means a}$ finitely generated free $\mathfrak o$ -module. Such a quadratic form q_L is called an integral quadratic form and such a pair (L, q_L) is called a quadratic lattice. We sometimes say that L is a quadratic lattice by omitting q_L , if this does not cause confusion or ambiguity. Similarly, we define a quadratic space $(V, q_L \otimes_0 1)$ for $V = L \otimes_0 F$ and sometimes say that V is a quadratic space. If there is no ambiguity, then we simply use q_L , rather than $q_L \otimes 1$, to stand for the quadratic form defined on V . We assume that V is nondegenerate with respect to the quadratic form $q_L \otimes 1$. More generally, for a commutative \mathfrak{o} -algebra R, we define a quadratic R-module $L \otimes_{\mathfrak{g}} R$ in the same manner. We sometimes call it a quadratic R-lattice.
- For a quadratic lattice (L, q_L) over **o**, the quadratic form \bar{q}_L on $L \otimes_{\mathfrak{o}} \kappa$ is defined to be $q_L \mod \pi$.
- For two $\mathfrak{o}\text{-}$ -lattices L and M, assume that L and M have the same rank and that $L \supseteq M$. Then we denote by $[L : M]$ the length of the torsion module L/M so that the cardinality of L/M is $f^{[L:M]}$.
- The fractional ideal generated by $q_L(X)$ as X runs through L will be called the norm of L and written $N(L)$.
- For matrices X and Y with entries in F, the matrix $\begin{pmatrix} X & 0 \\ 0 & Y \end{pmatrix}$ $0 Y$ \setminus is denoted by $X \perp Y$ and is called the orthogonal sum of X and Y .
- Let (a_1, \dots, a_m) and (b_1, \dots, b_n) be non-decreasing sequences consisting of non-negative integers in Z. Then $(a_1, \dots, a_m) \cup (b_1, \dots, b_n)$ is defined as the non-decreasing sequence (c_1, \dots, c_{n+m}) such that $\{c_1, \dots, c_{n+m}\} = \{a_1, \dots, a_m\} \cup \{b_1, \dots, b_n\}$ as multisets.
- For $\underline{a} = (a_1, \dots, a_n)$ with each a_i an element of \mathbb{Z} , the sum $a_1 + \dots + a_n$ is denoted by $|\underline{a}|$.
- For $\underline{a} = (a_1, \dots, a_n)$ with each a_i an element of \mathbb{Z} , the first m-tuple (a_1, \dots, a_m) with $m \leq n$ is denoted by $\underline{a}^{(m)}$.
- Let $H =$ $\begin{pmatrix} 0 & 1/2 \\ 1/2 & 0 \end{pmatrix}$ and let H_k be the orthogonal sum of the k-copies of H. Then H_k defines a quadratic lattice of rank 2k. We denote such a lattice by (H_k, q_k) , if this does not cause confusion or ambiguity. Similarly we define the quadratic space $(W_k, q_k \otimes 1)$, where
- $W_k = H_k \otimes_{\mathfrak{o}} F$, and sometimes say that W_k is a quadratic space. If there is no ambiguity, then we simply use q_k , rather than $q_k \otimes 1$, to stand for the quadratic form defined on W_k .
- For a symmetric matrix B of size $n \times n$, we say that B is half-integral over o if each nondiagonal entry multiplied by 2 and each diagonal entry of B are in \mathfrak{o} . We sometimes say that B is half-integral, by omitting 'over \mathfrak{o}' , if it does not cause ambiguity.
- We say that B is non-degenerate if the determinant of B is nonzero.
- For $U \in GL_n(\mathfrak{o})$ and B, where B is a half-integral symmetric matrix over \mathfrak{o} of size $n \times n$, we set $B[U] = {}^{t}UBU$. Here, ${}^{t}U$ is the matrix transpose of U.
- We say that two half-integral matrices B and B' are equivalent if $B' = B[U]$ for a certain $U \in GL_n(\mathfrak{o}).$
- For two quadratic R-lattices L and L' , where R is a commutative \mathfrak{o} -algebra, we say that an R-linear map $f: L \to L'$ is *isometry* if it is injective and preserves the associated quadratic forms, i.e. $q_L(x) = q_{L'}(f(x))$ for any $x \in L$.
- Assume that $k \geq n$, where n is the rank of L. We define $O_0(L, H_k)$ as the affine scheme defined over $\mathfrak o$ such that $O_{\mathfrak o}(L, H_k)(R)$, the set of R-points of $O_{\mathfrak o}(L, H_k)$ for any commutative o-algebra R, is the set of R-linear maps (not necessarily injective) from $L \otimes R$ to $H_k \otimes R$ preserving the associated quadratic forms. If R is a flat \mathfrak{o} -domain, then $O_{\mathfrak{o}}(L, H_k)(R)$ is the set of isometries (i.e. injective) from the quadratic R-lattice $L \otimes R$ to the quadratic R-lattice $H_k \otimes R$, which will be proved in the next lemma.

Lemma 2.1. Assume that $k \geq n$, where n is the rank of L. If R is a flat o-domain, then an R-linear map from $L \otimes R$ to $H_k \otimes R$ preserving the associated quadratic forms is injective. Thus the generic fiber of $O_0(L, H_k)$, denoted by $O_F(V, W_k)$ with $V = L \otimes_{\mathfrak{d}} F$ and $W_k = H_k \otimes_{\mathfrak{d}} F$, represents the set of isometries from the quadratic space V to the quadratic space W_k .

Proof. Let $\varphi: L \otimes R \to H_k \otimes R$ be an R-linear map preserving the associated quadratic forms for a flat \mathfrak{o} -domain R. We choose $v \in L \otimes R$ such that $\varphi(v) = 0$. Our goal is to show that $v = 0$.

Assume that $v \neq 0$ in $L \otimes R$. Let R_0 be the quotient field of R. Note that the characteristic of R_0 is 0 since R is flat over \mathfrak{o} .

If we let $\tilde{v} = v \otimes 1 \in L \otimes R_0$, then \tilde{v} is nonzero. Thus we can choose a basis, say \mathcal{B} , of an R_0 -vector space $L \otimes R_0$ of dimension n involving \tilde{v} . We may assume that the last vector in \mathcal{B} is \tilde{v} .

We write $\tilde{\varphi} = \varphi \otimes 1 : L \otimes R_0 \to H_k \otimes R_0$ such that $\tilde{\varphi}(\tilde{v}) = 0$. If we express an R_0 -linear map $\tilde{\varphi}$ as a matrix T of size $(2k \times n)$ with respect to a basis B, then the last column vector of T is zero. We now consider the following matrix equation:

$$
q_L = {}^tT \cdot q_k \cdot T.
$$

Here, q_L (respectively q_k) is the symmetric matrix associated to $L \otimes R_0$ (respectively $H_k \otimes R_0$) with suitable sets of basis. Thus both q_L and q_k are nondegenerate. This contracts to the given setting since the determinant of q_L is nonzero, whereas that of the right hand side is 0. Therefore $\tilde{v} = 0$ which implies that $v = 0$.

Let B be a non-degenerate half-integral symmetric matrix over $\mathfrak o$ of size $n \times n$. We will define the Gross-Keating invariant for B below. The definition is taken from [\[IK1\]](#page-50-12).

Definition 2.2 (Definitions 0.1 and 0.2 in $[IK1]$). (b_{ij}) . Let $S(B)$ be the set of all non-decreasing sequences $(a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n$ such that

$$
\begin{array}{ll}\n\text{ord}(b_{ii}) \ge a_i & (1 \le i \le n); \\
\text{ord}(2b_{ij}) \ge (a_i + a_j)/2 & (1 \le i \le j \le n).\n\end{array}
$$

Put

$$
\mathbf{S}(\{B\}) = \bigcup_{U \in \mathrm{GL}_n(\mathfrak{o})} S(B[U]).
$$

The Gross-Keating invariant $GK(B)$ of B is the greatest element of $S({B})$ with respect to the lexicographic order \succeq on $\mathbb{Z}_{\geq 0}^n$. Here, the lexicographic order \succeq on $\mathbb{Z}_{\geq 0}^n$ is the following (cf. the paragraph following Definition 0.1 of [\[IK1\]](#page-50-12)). Choose two elements (a_1, \dots, a_n) and (b_1, \dots, b_n) in $\mathbb{Z}_{\geq 0}^n$. Let i be the first integer over which a_i differs from b_i (so that $a_j = b_j$ for any $j < i$). If $a_i > b_i$, then we say that $(a_1, \dots, a_n) \succ (b_1, \dots, b_n)$. Otherwise, we say that $(a_1, \dots, a_n) \prec (b_1, \dots, b_n)$.

(2) The symmetric matrix B is called *optimal* if $GK(B) \in S(B)$.

(3) If B is a symmetric matrix associated to a quadratic lattice (L, q_L) , then $GK(L)$, called the Gross-Keating invariant of (L, q_L) , is defined by $GK(B)$. $GK(L)$ is independent of the choice of a matrix B.

It is known that the set $S({B})$ is finite (cf. [\[IK1\]](#page-50-12)), which explains well-definedness of $GK(B)$. We can also see that $GK(B)$ depends on the equivalence class of B. In general, it is a difficult question to check whether or not a given matrix B is optimal. Ikeda and Katsurada introduced so-called 'reduced form' associated to B and showed that it is optimal. We use a reduced form several times in this paper and thus provide its detailed definition through the following series of definitions [2.3](#page-8-1)[-2.5.](#page-8-0) They are taken from [\[IK1\]](#page-50-12) and [\[IK2\]](#page-50-10) for synchronization.

In [\[IK1\]](#page-50-12), they defined a reduced form when $p = 2$. However, their definition and main theorems hold for any p , which were explained in the initial version of their paper posted on arXiv. Thus, in the following, we will not make restriction on p , unless otherwise stated.

Definition 2.3 (Definition 3.1 in [\[IK2\]](#page-50-10)). Let $\underline{a} = (a_1, \dots, a_n)$ be a non-decreasing sequence of non-negative integers. Write \underline{a} as

$$
\underline{a} = (\underbrace{m_1, \cdots, m_1}_{n_1}, \cdots, \underbrace{m_r, \cdots, m_r}_{n_r})
$$

with $m_1 < \cdots < m_r$ and $n = n_1 + \cdots + n_r$. For $s = 1, 2, \cdots, r$, put

$$
n_s^* = \sum_{u=1}^s n_u,
$$

and

$$
I_s = \{n_{s-1}^* + 1, n_{s-1}^* + 2, \cdots, n_s^*\}.
$$

Here, we let $n_0^* = 0$.

Let \mathfrak{S}_n be the symmetric group of degree *n*. Let $\sigma \in \mathfrak{S}_n$ be an involution i.e. $\sigma^2 = id$.

Definition 2.4 (Definition 3.1 in [\[IK1\]](#page-50-12)). For a non-decreasing sequence of non-negative integers $\underline{a} = (a_1, \cdots, a_n)$, we set

$$
\mathcal{P}^{0} = \mathcal{P}^{0}(\sigma) = \{i | 1 \le i \le n, i = \sigma(i)\},
$$

$$
\mathcal{P}^{+} = \mathcal{P}^{+}(\sigma) = \{i | 1 \le i \le n, a_{i} > a_{\sigma(i)}\},
$$

$$
\mathcal{P}^{-} = \mathcal{P}^{-}(\sigma) = \{i | 1 \le i \le n, a_{i} < a_{\sigma(i)}\}.
$$

We say that an involution $\sigma \in \mathfrak{S}_n$ is an \underline{a} -admissible involution if the following three conditions are satisfied:

(1) \mathcal{P}^0 has at most two elements. If \mathcal{P}^0 has two distinct elements i and j, then $a_i \not\equiv a_j \mod 2$, and

$$
a_i = \max\{a_j | j \in \mathcal{P}^0 \cup \mathcal{P}^+, a_j \equiv a_i \text{ mod } 2\}.
$$

(2) For $s = 1, \dots, r$, we have

$$
#(I_s \cap \mathcal{P}^+) \le 1, \#(I_s \cap \mathcal{P}^-) + #(I_s \cap \mathcal{P}^0) \le 1.
$$

(3) If $i \in \mathcal{P}^-$, then

$$
a_{\sigma(i)} = \min\{a_j | j \in \mathcal{P}^+, a_j > a_i, a_j \equiv a_i \mod 2\}.
$$

Similarly, if $i \in \mathcal{P}^+$, then

$$
a_{\sigma(i)} = \max\{a_j | j \in \mathcal{P}^-, \ a_j < a_i, a_j \equiv a_i \mod 2\}.
$$

Definition 2.5 (Definition 3.2 in [\[IK1\]](#page-50-12)). Write $B = (b_{ij})$. Let $\underline{a} \in S(B)$. Let $\sigma \in \mathfrak{S}_n$ be an \underline{a} admissible involution. We say that B is a reduced form of GK-type (a, σ) if the following conditions are satisfied:

(1) If $i \notin \mathcal{P}^0$, $j = \sigma(i)$, and $a_i \leq a_j$, then

$$
GK\left(\begin{pmatrix}b_{ii}&b_{ij}\\b_{ij}&b_{jj}\end{pmatrix}\right)=(a_i,a_j).
$$

Note that if $p = 2$ then this condition is equivalent to the following condition (by Proposition 2.3 of [\[IK1\]](#page-50-12)).

$$
\begin{cases} \text{ord}(2b_{ij}) = \frac{a_i + a_j}{2} & \text{if } i \notin \mathcal{P}^0, j = \sigma(i); \\ \text{ord}(b_{ii}) = a_i & \text{if } i \in \mathcal{P}^-.\end{cases}
$$

(2) if $i \in \mathcal{P}^0$, then

 $\mathrm{ord}(b_{ii})=a_i.$

(3) If $i \neq i, \sigma(i)$, then

$$
\operatorname{ord}(2b_{ij}) > \frac{a_i + a_j}{2}.
$$

Theorem 2.6 (Corollary 5.1 in [\[IK1\]](#page-50-12)). A reduced form is optimal. More precisely, if B is a reduced form of $G K$ -type (a, σ) , then

$$
GK(B) = \underline{a}.
$$

Remark 2.7. Assume that $p = 2$. In this remark, we will explain the existence of a reduced form and the uniqueness of an involution up to equivalence.

- (1) For any given non-decreasing sequence of non-negative integers $\underline{a} = (a_1, \dots, a_n)$, there always exists an a -admissible involution (cf. the paragraph following Definition 3.1 of $[IK1]$).
- (2) For any given non-degenerate half-integral symmetric matrix B over \mathfrak{o} , there always exist a GK(B)-admissible involution σ and a reduced form of GK type $(GK(B), \sigma)$ which is equivalent to B (cf. Theorem 4.1 of [\[IK1\]](#page-50-12)).
- (3) we say that two *a*-admissible involutions are equivalent if they are conjugate by an element of $\mathfrak{S}_{n_1} \times \cdots \times \mathfrak{S}_{n_r}$. Here, we follow the notation introduced in Definition [2.3](#page-8-1) to specify the integers n_1, \dots, n_r . If σ is an a -admissible involution, then the equivalence class of σ is determined by

(2.1)
$$
\#(\mathcal{P}^+\cap I_s), \quad \#(\mathcal{P}^-\cap I_s), \quad \#(\mathcal{P}^0\cap I_s)
$$

for $1 \leq s \leq r$ (cf. the paragraph following Remark 4.1 in [\[IK1\]](#page-50-12)).

(4) Let σ and τ be GK(B)-admissible involutions associated to reduced forms of GK types $(GK(B), \sigma)$ and $(GK(B), \tau)$, respectively, which are equivalent to a given symmetric matrix B. Then σ and τ are equivalent (cf. Theorem 4.2 of [\[IK1\]](#page-50-12)). Therefore, the above sets in (2.1) for B are independent of the choice of a GK(B)-admissible involution with a reduced form.

We list a few facts about the Gross-Keating invariant below.

- **Remark 2.8.** (1) If p is odd, then a diagonal matrix, whose diagonal entries are $u_i \pi^{a_i}$ with $u_i \in \mathfrak{o}^{\times}$ and $a_i \leq a_j$ for $i < j$, is a reduced form (cf. Remark 1.1 of [\[IK1\]](#page-50-12)) and the Gross-Keating invariant is (a_1, \dots, a_n) . Note that any half-integral symmetric matrix is diagonalizable for p odd.
	- (2) For the half-integral symmetric matrix H_k of rank $2k$, we have

$$
GK(H_k)=(0,\cdots,0).
$$

(3) If there is an isometry from (L, q_L) of rank n to (H_k, q_k) , then $n \leq 2k$ and

$$
GK(L) \succeq GK(H_k)^{(n)} = (0, \cdots, 0)
$$

by Lemma 1.2 of [\[IK1\]](#page-50-12).

- (4) The first integer of $GK(L)$ is the exponential order of a generator of $N(L)$ (cf. Lemma B.1) of [\[Yan04\]](#page-51-1)).
- (5) Consider $L \subseteq L' \subseteq V$ such that $[L':L] = b$. Then

$$
|\mathrm{GK}(L')| = |\mathrm{GK}(L)| - 2b
$$

by Theorem 0.1 of [\[IK1\]](#page-50-12). Here, $|GK(L)| = a_1 + \cdots + a_n$ for $GK(L) = (a_1, \dots, a_n)$.

3. Local densities

The Siegel series of a quadratic lattice (L, q_L) can be defined in terms of the local density associated to two quadratic lattices (L, q_L) and (H_k, q_k) (cf. Definition [3.15\)](#page-16-1). From this section to the end, we assume that $k \geq n$, where n is the rank of L. The purpose of this section is to reformulate the local density (and the Siegel series) in terms of certain lattice counting problem conceptually, whose explicit form is given in Theorem [3.14](#page-16-0) and Corollary [3.16.](#page-17-1)

3.1. Local density and primitive local density. We define the following notions:

 $\sqrt{ }$ \int \overline{a} α F-vector space of quadratic forms defined on V ; \mathfrak{M} : the set of F-linear maps from V to W_k ; \mathfrak{M}_L : the set of $\mathfrak{o}\text{-}linear$ maps from L to H_k ; \mathfrak{Q}_L : the free \mathfrak{o} -module of integral quadratic forms defined on L.

Here, we remind that $V = L \otimes_{\mathfrak{o}} F$ and $W_k = H_k \otimes_{\mathfrak{o}} F$.

Regarding M and $\mathfrak Q$ as varieties over F, let $\omega_{\mathfrak{M},L}$ and $\omega_{\mathfrak{Q},L}$ be nonzero, translation-invariant forms on \mathfrak{M} and \mathfrak{Q} , respectively, with normalizations

$$
\int_{\mathfrak{M}_L} |\omega_{\mathfrak{M},L}| = 1 \text{ and } \int_{\mathfrak{Q}_L} |\omega_{\mathfrak{Q},L}| = 1.
$$

Let \mathfrak{M}^* be the set of injective linear maps from V to W. We can also regard \mathfrak{M}^* as an open subvariety of \mathfrak{M} . Thus \mathfrak{M}^* is a (not necessarily affine) nonsingular variety over F. Define a map $\rho : \mathfrak{M}^* \to \mathfrak{Q}$ by $\rho(m) = q_k \circ m$. Here q_k is the quadratic form defined on W_k . Then the inverse image of $q_L \otimes_{\mathfrak{o}} 1$ along the map ρ is $O_F(V, W_k)$, which represents the set of isometries from the quadratic space V to the quadratic space W_k (cf. Lemma [2.1\)](#page-7-0). Here q_L is the quadratic form defined on L. It is easy to see that the morphism ρ is representable as a morphism of schemes over F and smooth by showing the surjectivity of the differential of ρ over the Zariski tangent space on any closed point.

Definition 3.1. We will define a differential ω_L^{ld} on $O_F(V, W_k)$ associated to $\omega_{\mathfrak{M},L}$ and $\omega_{\mathfrak{Q},L}$. This is taken from Section 3 of [\[GY00\]](#page-50-13). Smoothness of the morphism $\rho : \mathfrak{M}^* \to \mathfrak{Q}$ induces the following short exact sequence of locally free sheaves on \mathfrak{M}^* (cf. Proposition 5 of [\[BLR90\]](#page-50-14)):

$$
0\to \rho^*\Omega_{\mathfrak{Q}/F}\to \Omega_{\mathfrak{M}^*/F}\to \Omega_{\mathfrak{M}^*/\mathfrak{Q}}\to 0.
$$

This gives rise to an isomorphism

$$
\rho^* \left(\bigwedge^{\text{top}} \Omega_{\mathfrak{Q}/F} \right) \otimes \bigwedge^{\text{top}} \Omega_{\mathfrak{M}^*/\mathfrak{Q}} \simeq \bigwedge^{\text{top}} \Omega_{\mathfrak{M}^*/F}.
$$

Let $\omega_L \in$ top
 $\bigwedge^{\text{top}} \Omega_{\mathfrak{M}^*/\mathfrak{Q}}(\mathfrak{M}^*)$ be such that $\rho^* \omega_{\mathfrak{Q},L} \otimes \omega_L = \omega_{\mathfrak{M},L}|_{\mathfrak{M}^*}$. We then denote by ω_L^{ld} the restriction of ω_L to $O_F(V, W_k)$. We sometimes write $\omega_L^{\text{ld}} = \omega_{\mathfrak{M},L}/\rho^* \omega_{\mathfrak{Q},L}$.

Lemma 3.2. We have the following equation:

$$
\int_{\mathcal{O}_{\mathfrak{o}}(L, H_k)(\mathfrak{o})} |\omega_L^{\rm Id}| = \lim_{N \to \infty} f^{-N \cdot dim\mathcal{O}_F(V, W)} \# \mathcal{O}_{\mathfrak{o}}(L, H_k)(\mathfrak{o}/\pi^N \mathfrak{o}),
$$

where the limit stabilizes for N sufficiently large. Here, $dim\mathrm{O}_F(V,W_k)=dim\mathrm{O}(W_k)-dim\mathrm{O}(V^{\perp})=$ $dim\mathfrak{M} - dim\mathfrak{Q} = 2kn - (n^2 + n)/2.$

The lemma is well-known and a generalization of Lemma 3.4 of [\[GY00\]](#page-50-13). If \mathfrak{M} has a group structure, then the proof is explained in page 5 of [\[Han99\]](#page-50-15) or in pages 119-120 of [\[Tam66\]](#page-51-2). The general case is explained in [\[Yu08\]](#page-51-3), which is not available on the internet. Since the proof of the general case is not different from the case handled in [\[Han99\]](#page-50-15) and [\[Tam66\]](#page-51-2), we sketch a main idea of the proof.

Proof. The right hand side can be identified with the following:

$$
\lim_{U_q \to q_L} \frac{\int_{\rho^{-1}(U_q) \cap \mathfrak{M}_L} |\omega_{\mathfrak{M},L}|}{\int_{U_q} |\omega_{\mathfrak{Q},L}|},
$$

where U_q is an open neighborhood of q_L of the form $q_L + \pi^N \mathfrak{Q}_L$ for $N > 0$ so that $\int_{q_L+\pi^N \mathfrak{Q}_L} |\omega_{\mathfrak{Q},L}| =$ $f^{-N \cdot dim\mathfrak{Q}}$ and $\int_{\rho^{-1}(q_L+\pi^N\mathfrak{Q}_L)\cap \mathfrak{M}_L} |\omega_{\mathfrak{M},L}| = f^{-N \cdot dim\mathfrak{M}} \cdot \#O_{\mathfrak{o}}(L, H_k)(\mathfrak{o}/\pi^N \mathfrak{o})$. It is well known that isomorphism classes of quadratic lattices are locally constant. In other words, any quadratic form contained in $q_L + \pi^N \mathfrak{Q}_L$, for sufficiently large integer N, is isometric to q_L by Theorem 2 of [\[Dur44\]](#page-50-16). Thus Fubini's theorem yields that the fraction of these two integrals with sufficiently large integer N is the same as the left hand side. \square

Definition 3.3. The local density associated to the pair of two quadratic lattices L and H_k , denoted by $\alpha(L, H_k)$, is defined as

$$
\alpha(L, H_k) = \int_{\mathcal{O}_\mathfrak{o}(L, H_k)(\mathfrak{o})} |\omega_L^{\mathrm{ld}}|.
$$

We define the subfunctor $O_0^{prim}(L, H_k)$ of $O_0(L, H_k)$ such that $O_0^{prim}(L, H_k)(R)$, the set of Rpoints for a commutative o -algebra R, is the set of elements in $O_0(L, H_k)(R)$ whose at least one $n \times n$ -minor, as a linear map from $L \otimes_{\mathfrak{o}} R$ to $H_k \otimes_{\mathfrak{o}} R$, is a unit in R. In particular, if $R = \mathfrak{o}$, then $O_{\mathfrak{o}}^{prim}(L, H_k)(\mathfrak{o})$ is the set of elements in $O_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$ whose reduction modulo π is injective from $L \otimes_{\mathfrak{o}} \kappa$ to $H_k \otimes_{\mathfrak{o}} \kappa$. Each element in $O_{\mathfrak{o}}^{prim}(L, H_k)(R)$ is called a primitive isometry. We will show that $O_{\mathfrak{o}}^{prim}(L, H_k)$ is an open (not necessarily affine) subscheme of $O_{\mathfrak{o}}(L, H_k)$ (cf. Corollary [3.11\)](#page-14-0) and that $O_{\mathfrak{o}}^{prim}(L, H_k)(\mathfrak{o})$ is open in $O_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$ in terms of inherent p-adic topology (cf. Lemma [3.6\)](#page-12-0).

Consider a lattice L' in V containing L. Let $q_{L'}$ be the quadratic form defined on L' , whose restriction to L is the same as q_L . Here, $q_{L'}$ is abuse of notation since it may not be an integral quadratic form. We can regard $O_0^{prim}(L', H_k)(\mathfrak{o})$ as a subset of $O_0(L, H_k)(\mathfrak{o})$ induced by the restriction to L. Since any linear map in $O_0(L, H_k)(\mathfrak{o})$ is injective (i.e. isometry) by Lemma [2.1,](#page-7-0) we have the following stratification on $O_0(L, H_k)(\mathfrak{o})$:

(3.1)
$$
\mathrm{O}_{\mathfrak{o}}(L, H_k)(\mathfrak{o}) = \bigsqcup_{L \subseteq L' \subseteq V} \mathrm{O}_{\mathfrak{o}}^{prim}(L', H_k)(\mathfrak{o}).
$$

In the following lemma, we will show that the norm $N(L')$ of L' should be contained in the ring \mathfrak{o} , in order that $O_{\mathfrak{o}}^{prim}(L', H_k)(\mathfrak{o})$ is nonempty. This directly implies that the above is a finite disjoint union.

Lemma 3.4. The condition $N(L') \subseteq o$ is equivalent to the existence of a primitive isometry from $(L', q_{L'})$ to (H_k, q_k) .

Proof. If we have a primitive isometry from $(L', q_{L'})$ to (H_k, q_k) , then the fact $N(H_k) = \mathfrak{o}$ yields $N(L') \subseteq \mathfrak{o}.$

Conversely, assume that $N(L') \subseteq \mathfrak{o}$. The symmetric matrix B' of the quadratic lattice L' with respect to any basis is then half-integral over ρ . We consider the half-integral symmetric matrix $\begin{pmatrix} 0 & \frac{1}{2} \end{pmatrix}$ $\begin{array}{cc} 0 & \frac{1}{2} \cdot id_n \\ \frac{1}{2} \cdot id & R' \end{array}$ $rac{1}{2} \cdot id_n \quad B'$ \setminus . Here, id_n is the $(n \times n)$ -identity matrix, where n is the rank of L'. Note that $\int_{0}^{1} 0 \frac{1}{2}$ $\begin{array}{cc} 0 & \frac{1}{2} \cdot id_n \\ \frac{1}{2} \cdot id & 0 \end{array}$ $rac{1}{2} \cdot id_n \qquad 0$ Í. is an half-integral symmetric matrix of the quadratic lattice H_n . We claim that $\int_0^1 0 \frac{1}{2}$ $\begin{array}{cc} 0 & \frac{1}{2} \cdot id_n \\ \frac{1}{2} \cdot id & R' \end{array}$ $rac{1}{2} \cdot id_n$ B' is equivalent to the matrix $\begin{pmatrix} 0 & \frac{1}{2} \\ 1 & 0 \end{pmatrix}$ $\begin{array}{cc} 0 & \frac{1}{2} \cdot id_n \\ \frac{1}{2} \cdot id & 0 \end{array}$ $rac{1}{2} \cdot id_n = 0$ $\big)$ over \mathfrak{o} . This induces the existence of a primitive isometry from $(L', q_{L'})$ to (H_k, q_k) . ′

Our claim follows from the matrix equation:

$$
\begin{pmatrix} 0 & \frac{1}{2} \cdot id_n \\ \frac{1}{2} \cdot id_n & 0 \end{pmatrix} = \begin{pmatrix} id_n & 0 \\ {}^t X & id_n \end{pmatrix} \begin{pmatrix} 0 & \frac{1}{2} \cdot id_n \\ \frac{1}{2} \cdot id_n & B' \end{pmatrix} \begin{pmatrix} id_n & X \\ 0 & id_n \end{pmatrix}.
$$

Here, X is any matrix with entries in \mathfrak{o} such that $X + {}^t X + 2B' = 0$.

This lemma, combined with Remark 2.8 induces the following description of the existence of a primitive isometry in terms of the Gross-Keating invariant:

Corollary 3.5. There exists a primitive isometry from $(L', q_{L'})$ to (H_k, q_k) if and only if

$$
GK(L') \succeq GK(H_k)^{(n)} = (0, \cdots, 0).
$$

Lemma 3.6. Assume that $N(L') \subseteq \mathfrak{o}$. The set $O_{\mathfrak{o}}^{prim}(L', H_k)(\mathfrak{o})$ is open in $O_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$ in terms of the p-adic topology.

Proof. It suffices to show that $O_0(L', H_k)(\mathfrak{o})$ is open in $O_0(L, H_k)(\mathfrak{o})$ and that $O_0^{prim}(L, H_k)(\mathfrak{o})$ is open in $O_0(L, H_k)(\mathfrak{o}).$

Let us identify $\text{End}_{\mathfrak{g}}(L, H_k)(\mathfrak{o})$ with the set of $2k \times n$ -matrices with entries in \mathfrak{o} . Let $d = [L']$: L. Since $L \subset L' \subset \frac{1}{\pi^d}L$, $\text{End}_{\mathfrak{o}}(L', H_k)(\mathfrak{o})$ contains $\text{End}_{\mathfrak{o}}(\frac{1}{\pi^d}L, H_k)(\mathfrak{o})$. The latter is identified with $\pi^d \cdot \text{End}_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$ as a subset of $\text{End}_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$. Here, $\pi^d \cdot \text{End}_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$ is defined as $\{\pi^d X, X \in \text{End}_{\mathfrak{o}}(L, H_k)(\mathfrak{o})\}.$ Therefore, as an $\mathfrak{o}\text{-module}$, $\text{End}_{\mathfrak{o}}(L', H_k)(\mathfrak{o})$ has a finite index in $\text{End}_{\mathfrak{g}}(L, H_k)(\mathfrak{o})$ and so is open.

Since $O_{\mathfrak{o}}(L', H_k)(\mathfrak{o})$ is the intersection of $\text{End}_{\mathfrak{o}}(L', H_k)(\mathfrak{o})$ and $O_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$ inside $\text{End}_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$, it is open in $O_{\mathfrak{o}}(L, H_k)(\mathfrak{o}).$

For the second claim, since $O_{\mathfrak{o}}^{prim}(L, H_k)(\mathfrak{o})$ is the intersection of $\text{End}_{\mathfrak{o}}^{prim}(L, H_k)(\mathfrak{o})$ and $O_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$ inside $\text{End}_{\mathfrak{g}}(L, H_k)(\mathfrak{o})$, as in the above case, it suffices to show that $\text{End}_{\mathfrak{g}}^{prim}(L, H_k)(\mathfrak{o})$ is open in $\text{End}_{\mathfrak{g}}(L, H_k)(\mathfrak{o})$. Here $\text{End}_{\mathfrak{g}}^{prim}(L, H_k)(\mathfrak{o})$ consists of endomorphisms from L to H_k whose at least one $n \times n$ -minor is a unit in \mathfrak{o} . We consider the reduction map modulo π from $\text{End}_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$ to $\text{End}_{\kappa}(L \otimes \kappa, H_k \otimes \kappa)$. The induced quotient topology on the latter is the discrete topology on a finite set. Let $\text{End}_{\kappa}^{prim}(L \otimes \kappa, H_k \otimes \kappa)$ be the (open) subset of $\text{End}_{\kappa}(L \otimes \kappa, H_k \otimes \kappa)$ consisting of endomorphisms from $L \otimes \kappa$ to $H_k \otimes \kappa$ whose at least one $n \times n$ -minor is nonzero in κ . Then $\text{End}_{\mathfrak{o}}^{prim}(L, H_k)(\mathfrak{o})$ is the inverse image of $\text{End}_{\kappa}^{prim}(L \otimes \kappa, H_k \otimes \kappa)$ under the reduction map. Thus it is open.

Proposition 3.7. The local density $\alpha(L, H_k)$ is expressed as the following sum:

(3.2)
$$
\alpha(L, H_k) = \sum_{L \subseteq L' \subseteq V} f^{[L':L] \cdot (n+1-2k)} \int_{O_{\mathfrak{o}}^{prim}(L', H_k)(\mathfrak{o})} |\omega_{L'}^{\text{ld}}|.
$$

Proof. Since the set $O_{\mathfrak{o}}^{prim}(L', H_k)(\mathfrak{o})$ is open in $O_{\mathfrak{o}}(L, H_k)(\mathfrak{o})$ by Lemma [3.6,](#page-12-0) Equation [\(3.1\)](#page-11-1) yields the following identity:

$$
\alpha(L, H_k) = \sum_{L \subseteq L' \subseteq V} \int_{\mathcal{O}_\mathfrak{o}^{prim}(L', H_k)(\mathfrak{o})} |\omega_L^{\mathrm{ld}}|.
$$

Let $d = [L':L]$. Recall that $\omega_L^{\text{ld}} = \omega_{\mathfrak{M},L}/\rho^* \omega_{\mathfrak{Q},L}$ and $\omega_{L'}^{\text{ld}} = \omega_{\mathfrak{M},L'}/\rho^* \omega_{\mathfrak{Q},L'}$ in Definition [3.1.](#page-10-2) By the normalizations of $\omega_{\mathfrak{M},L}$ and $\omega_{\mathfrak{Q},L}$ given at the beginning of this subsection, we have

$$
\begin{cases}\n\omega_{\mathfrak{M},L} = \pi^{2kd} \cdot \omega_{\mathfrak{M},L'}; \\
\omega_{\mathfrak{Q},L} = \pi^{d(n+1)} \cdot \omega_{\mathfrak{Q},L'}.\n\end{cases}
$$

Thus, $\omega_L^{\text{ld}} = \pi^{2kd-(n+1)d} \omega_{L'}^{\text{ld}}$. Equation [\(3.2\)](#page-12-1) follows from this observation.

Definition 3.8. We define the primitive local density associated to the quadratic lattices L and H_k , denoted by $\alpha^{prim}(L, H_k)$, as follows:

$$
\alpha^{prim}(L, H_k) = \int_{\mathcal{O}_\mathfrak{o}^{prim}(L, H_k)(\mathfrak{o})} |\omega_L^{\mathrm{ld}}|.
$$

Thus Equation [\(3.2\)](#page-12-1) is written as follows:

(3.3)
$$
\alpha(L, H_k) = \sum_{L \subseteq L' \subseteq V} f^{[L':L]\cdot (n+1-2k)} \cdot \alpha^{prim}(L', H_k).
$$

Here, the sum runs over all quadratic lattices L' such that $N(L') \subseteq \mathfrak{o}$ and thus is finite. We remark that this formula is well-known in classical literatures using different terminologies (cf. Lemma 3 of [\[Kit83\]](#page-50-17)).

3.2. A formula of the primitive local density. We assume that $N(L) \subseteq \mathfrak{o}$ and thus the set $O_{\mathfrak{o}}^{prim}(L, H_k)(\mathfrak{o})$ is nonempty. In this subsection, we describe a formula of the primitive local density $\alpha^{prim}(L, H_k)$ by proving smoothness of $O_{\mathfrak{o}}^{prim}(L, H_k)$ as a scheme defined over \mathfrak{o} .

We first explain another description of $O_0^{prim}(L, H_k)$ in terms of the fiber of a certain morphism between two smooth schemes over o.

Let $\underline{M}_{\mathfrak{o}}(L, H_k)$ be the functor from the category of flat \mathfrak{o} -algebras to the category of sets such that $M_{\mathcal{O}}(L, H_k)(R)$, the set of R-points for a flat **o**-algebra R, is the set of R-linear maps from $L \otimes_{\mathfrak{o}} R$ to $H_k \otimes_{\mathfrak{o}} R$ by ignoring the associated quadratic forms. Then the functor $M_{\mathfrak{o}}(L, H_k)$ is uniquely represented by a flat \mathfrak{o} -algebra which is a polynomial ring over \mathfrak{o} of $2kn$ variables. Thus we can now talk of $\underline{M}_{\mathfrak{o}}(L, H_k)(R)$ for any (not necessarily flat) \mathfrak{o} -algebra R.

Let $\underline{M}_{\bullet}^{*}(L, H_{k})$ be the subfunctor of $\underline{M}_{\bullet}(L, H_{k})$ such that $\underline{M}_{\bullet}^{*}(L, H_{k})(R)$, the set of R-points for a commutative $\mathfrak o$ -algebra R, is the set of R-linear maps from $L \otimes_{\mathfrak o} R$ to $H_k \otimes_{\mathfrak o} R$ whose at least one $n \times n$ -minor is a unit in R. Then it is easy to see that $\underline{M}_{\mathfrak{o}}^{*}(L, H_{k})$ is an open subscheme of $\underline{M}_{\mathfrak{o}}(L, H_{k})$ which yields smoothness of $\underline{M}_{\mathfrak{o}}^{*}(L, H_{k})$ (cf. Section 3.2 of [\[Cho15\]](#page-50-18)). Note that $\underline{M}_{\mathfrak{o}}^{*}(L, H_{k})$ is not necessarily affine, but has finite affine covers (given by each $n \times n$ -minor). In particular, if $R = \mathfrak{o}$, then $\underline{M}_{\mathfrak{o}}^{*}(L, H_{k})(\mathfrak{o})$ is the set of \mathfrak{o} -linear maps from L to H_{k} whose reduction modulo π is injective from $L \otimes_{\mathfrak{o}} \kappa$ to $H_k \otimes_{\mathfrak{o}} \kappa$.

Let Q_L be the affine space of dimension $n(n+1)/2$ defined over \mathfrak{o} such that $Q_L(R)$, the set of R-points for a commutative $\mathfrak o$ -algebra R, is the set of quadratic forms on $L \otimes_{\mathfrak o} R$ whose coefficients are in R. Similarly, we define Q_{H_k} as the affine space of dimension $2k(2k+1)/2$ defined over \mathfrak{o} .

Let R be a flat **o**-algebra. As a matrix, each element of $Q_L(R)$ is given by a symmetric matrix (a_{ij}) of size $n \times n$ such that each nondiagonal entry a_{ij} with $i \neq j$ is of the form $1/2 \cdot a'_{ij}$ for $a'_{ij} \in R$ and each diagonal entry a_{ii} is contained in R .

Then we consider the following morphism

$$
\underline{Q}_{H_k}\times \underline{M}^*_\mathfrak{o}(L,H_k)\longrightarrow \underline{Q}_L, (q,m)\mapsto q\circ m.
$$

Here, $q \in \underline{Q}_{H_k}(R)$ and $m \in \underline{M}^*_\mathfrak{o}(L, H_k)(R)$ for a flat \mathfrak{o} -algebra R. It is easy to see that the above morphism is well-defined and represented by a morphism of schemes over \mathfrak{o} (cf. the last paragraph of the proof of Theorem 3.4 in [\[Cho16\]](#page-50-19)). The above morphism induces the morphism

$$
\rho: \underline{M}^*_\mathfrak{o}(L, H_k) \longrightarrow \underline{Q}_L, m \mapsto q_k \circ m.
$$

Theorem 3.9. The morphism $\rho : \underline{M}_{\mathfrak{o}}^*(L, H_k) \to \underline{Q}_L$ is smooth.

Proof. The theorem follows from Lemma 5.5.1 of [\[GY00\]](#page-50-13) and the following lemma. \square

Lemma 3.10. The morphism $\rho \otimes \kappa : \underline{M}_{\mathfrak{o}}^*(L, H_k) \otimes \kappa \to \underline{Q}_L \otimes \kappa$ is smooth.

Proof. The proof is based on Lemma 5.5.2 in [\[GY00\]](#page-50-13). It suffices to show that, for any $m \in$ $\underline{M}_{\bullet}^{*}(L, H_{k})(\bar{\kappa})$, the induced map on the Zariski tangent space $\rho_{*,m}: T_{m} \to T_{\rho(m)}$ is surjective. Here, $\bar{\kappa}$ is the algebraic closure of κ .

We introduce another functor on the category of flat \mathfrak{o} -algebras. Define $T(R)$ to be the set of $(n \times n)$ -matrices y such that each entry is of the form $1/2 \cdot y_{ij}$ with $y_{ij} \in R$. Then this functor is represented by an affine space over o.

We now compute the map $\rho_{*,m}$ explicitly. If we identify T_m with $\underline{M}_{\mathfrak{o}}(L, H_k)(\overline{\kappa})$ and $T_{\rho(m)}$ with $Q_L(\bar{\kappa})$, then

$$
\rho_{*,m}: T_m \longrightarrow T_{\rho(m)}, X \mapsto m^t \cdot q_k \cdot X + X^t \cdot q_k \cdot m.
$$

We explain how to compute $X \mapsto m^t \cdot q_k \cdot X + X^t \cdot q_k \cdot m$ explicitly. Firstly, we formally compute $X \mapsto m^t \cdot q_k \cdot X$. It is of the form $1/2 \cdot Y \in T(\bar{\kappa})$, where Y is an $n \times n$ -matrix with entries in $\bar{\kappa}$. Then we formally compute $1/2 \cdot Y + 1/2 \cdot {}^tY$. It is of the form Z, whose diagonal entries, denoted by z_{ii} 's, are in $\bar{\kappa}$ and whose nondiagonal entries are of the form $1/2 \cdot z_{ij}$ with $z_{ij} \in \bar{\kappa}$ such that $z_{ij} = z_{ji}$. More precisely, if we write $Y = (y_{ij})$, then $z_{ii} = y_{ii}$ and $z_{ij} = y_{ij} + y_{ji}$ for $i \neq j$. Thus Z is an element of $Q_L(\bar{\kappa})$.

To prove the surjectivity of $\rho_{*,m}: T_m \longrightarrow T_{\rho(m)}$, it suffices to show the following two statements:

- (1) $X \mapsto m^t \cdot q_k \cdot X$ defines a surjection $\underline{M}_{\mathfrak{o}}(L, H_k)(\overline{\kappa}) \to T(\overline{\kappa});$
- (2) $Y \mapsto {}^t Y + Y$ defines a surjection $T(\bar{\kappa}) \to \underline{Q}_L(\bar{\kappa})$.

These two arguments are direct from the construction of $T(\bar{\kappa})$.

Corollary 3.11. The scheme $O_0^{prim}(L, H_k)$ is an open subscheme $O_0(L, H_k)$. In addition, it is smooth over o.

Proof. The scheme $O_0^{prim}(L, H_k)$ is defined as the fiber of q_L along the smooth morphism ρ . Thus $O_0^{prim}(L, H_k)$ is the intersection of $O_0(L, H_k)$ and $\underline{M}_0^*(L, H_k)$ inside $\underline{M}_0(L, H_k)$. The fact that $M_{\mathfrak{o}}^*(L, H_k)$ is an open subscheme of $M_{\mathfrak{o}}(L, H_k)$ yields the first statement. We note that $O_0^{prim}(L, H_k)$ has finite affine covers given by $n \times n$ -minors, each of which is an open subscheme of $O_{\mathfrak{o}}(L, H_k)$ as well.

For the second statement, since smoothness is stable under base change, $O_0^{prim}(L, H_k)$ is a smooth scheme over \mathfrak{o} .

The special fiber of $O_0^{prim}(L, H_k)$ is $O_{\kappa}^{prim}(\bar{q}_L, \bar{q}_k)$, where \bar{q}_L (= $q_L \mod \pi$) is a quadratic form defined on $L \otimes_{\mathfrak{o}} \kappa$ and \bar{q}_k (= $q_k \mod \pi$) is a quadratic form defined on $H_k \otimes_{\mathfrak{o}} \kappa$. In particular, $O_{\mathfrak{o}}^{prim}(L, H_k)(\kappa)$ is the set of isometries from the quadratic space $(L \otimes_{\mathfrak{o}} \kappa, \bar{q}_L)$ to the quadratic space $(H_k \otimes_{\mathfrak{o}} \kappa, \bar{q}_k).$

We finally state the following formula of the primitive local density.

Theorem 3.12. Assume that $N(L) \subseteq \mathfrak{o}$. Then the primitive local density $\alpha^{prim}(L, H_k)$ is described as follows:

$$
\alpha^{prim}(L, H_k) = f^{-dim \mathcal{O}_F(V, W_k)} \cdot \# \mathcal{O}_\kappa^{prim}(\bar{q}_L, \bar{q}_k)(\kappa).
$$

Here, $\dim O_F(V, W_k) = 2kn - (n^2 + n)/2$ and $\#\mathrm{O}_\kappa^{prim}(\bar{q}_L, \bar{q}_k)(\kappa)$ stands for the cardinality of the set $O_{\kappa}^{prim}(\bar{q}_L, \bar{q}_k)(\kappa)$.

Proof. Theorem [3.9](#page-14-1) yields the following short exact sequence of locally free sheaves on $\underline{M}^*_0(L, H_k)$ as in Definition [3.1](#page-10-2) (cf. Proposition 5 of [\[BLR90\]](#page-50-14)):

$$
0\longrightarrow \rho^*\Omega_{\underline{Q}_L/\mathfrak{o}}\longrightarrow \Omega_{\underline{M}^*_\mathfrak{o}(L,H_k)/\mathfrak{o}}\longrightarrow \Omega_{\underline{M}^*_\mathfrak{o}(L,H_k)/\underline{Q}_L}\longrightarrow 0.
$$

Let $\omega_{\underline{M},L}$ (respectively $\omega_{\underline{Q},L}$) be a differential of top degree on $\underline{M}_{\mathfrak{g}}(L,H_k)$ (respectively \underline{Q}_L) over **o**, whose reduction on the special fiber is nowhere zero. Since $M_0^*(L, H_k)$ is an open subscheme of $\underline{M}_{\mathfrak{o}}(L, H_k)$, the restriction of $\omega_{\underline{M},L}$ to $\underline{M}_{\mathfrak{o}}^*(L, H_k)$ is a differential of top degree on $\underline{M}_{\mathfrak{o}}^*(L, H_k)$ whose reduction on the special fiber is nowhere zero.

We define ω_L^{can} to be $\omega_{M,L}/\rho^* \omega_{Q,L}$. Here, we refer to Definition [3.1](#page-10-2) for the notion of $\omega_{M,L}/\rho^* \omega_{Q,L}$. Then ω_L^{can} is a differential of top degree on $O_{\mathfrak{o}}^{prim}(L, H_k)$ over \mathfrak{o} , whose reduction on the special fiber is nowhere zero.

Since $\underline{M}_{\mathfrak{o}}(L, H_k)(\mathfrak{o}) = \mathfrak{M}_L$ and $\underline{Q}_L(\mathfrak{o}) = \mathfrak{Q}_L$, the differentials $\omega_{\mathfrak{M},L}$ (respectively $\omega_{\mathfrak{Q},L}$) and $\omega_{\underline{M},L}$ (respectively $\omega_{Q,L}$) determine the same measure on M (respectively \mathfrak{Q}). Here, see the beginning of subsection [3.1](#page-10-1) for the notions of $\omega_{\mathfrak{M},L}$, $\omega_{\mathfrak{Q},L}$, $\mathfrak{M}, \mathfrak{Q}_L, \mathfrak{Q}$. Thus the description of ω_L^{ld} given in Definition [3.1](#page-10-2) shows that $\omega_L^{\rm Id}$ and $\omega_L^{\rm can}$ determine the same measure on $O_F(V,W_k)(F)$. This is parallel to Section 7.1 of [\[GY00\]](#page-50-13). Note that a differential of top degree on $O_0^{prim}(L, H_k)$, whose reduction on the special fiber is nowhere zero, is not unique. But such two forms determine the same measure on $O_0^{prim}(L, H_k)(F)$ (= $O_F(V, W_k)(F)$ by Lemma [2.1\)](#page-7-0) due to smoothness of the scheme $O_0^{prim}(L, H_k)$. For more explanation about this, see Section 5 of Chapter 3 in [\[Pop11\]](#page-51-4).

The primitive local density is the volume of $O_0^{prim}(L, H_k)(\mathfrak{o})$ with respect to the measure $|\omega_L^{Id}|$ (cf. Definition [3.8\)](#page-13-1). Since $O_0^{prim}(L, H_k)(\mathfrak{o})$ is the set of \mathfrak{o} -points of the smooth scheme $O_0^{prim}(L, H_k)$ and $\alpha^{prim}(L, H_k)(\kappa) = \mathcal{O}_{\kappa}^{prim}(\bar{q}_L, \bar{q}_k)(\kappa)$, the formula directly follows from Theorem 2.2.5 of [\[Weil82\]](#page-51-5) or Theorem 5.3 of Chapter 3 of [\[Pop11\]](#page-51-4).

3.3. Reformulation of the local density. Using Theorem [3.12](#page-14-2) and Corollary [3.5,](#page-12-2) we can reformulate Equation [\(3.2\)](#page-12-1) of the local density as follows:

(3.4)
$$
\alpha(L, H_k) = f^{-2kn + (n^2 + n)/2} \cdot \sum_{\substack{L \subseteq L' \subseteq V, \\ \text{GK}(L') \succeq (0, \cdots, 0)}} f^{[L':L] \cdot (n+1-2k)} \cdot \#O_{\kappa}^{prim}(\bar{q}_{L'}, \bar{q}_k)(\kappa).
$$

Recall that $O_{\kappa}^{prim}(\bar{q}_{L'}, \bar{q}_k)(\kappa)$ is the set of isometries from the quadratic space $(L' \otimes \kappa, \bar{q}_{L'})$ to the quadratic space $(H_k \otimes \kappa, \bar{q}_k)$. Let $L' \otimes \kappa = \bar{L}'_0 \perp \bar{L}'_1$, where \bar{L}'_1 (= Rad $(L' \otimes \kappa)$) is the radical of $L' \otimes \kappa$ so that the restriction of the quadratic form $\bar{q}_{L'}$ on \bar{L}'_0 is nonsingular. We assign the following notion a^{\pm} with an integer a to \overline{L}' with respect to \overline{L}'_0 :

(3.5)
$$
\begin{cases} a^{+} & \text{if } a = \dim \bar{L}'_{0} \text{ is even and } \bar{L}'_{0} \text{ is split;} \\ a^{-} & \text{if } a = \dim \bar{L}'_{0} \text{ is even and } \bar{L}'_{0} \text{ is nonsplit;} \\ a(= a^{+} = a^{-}) & \text{if } a = \dim \bar{L}'_{0} \text{ is odd.} \end{cases}
$$

Here, $dim \bar{L}'_0$ is the dimension of \bar{L}'_0 as a κ -vector space. If $a = 0$, then we understand $0^+ = 0^-$. By Exercise 4 in Section 5.6 of [\[Kit93\]](#page-50-20), $\#O_{\kappa}^{prim}(\bar{q}_{L'}, \bar{q}_k)(\kappa)$ is completely determined by three ingredients, a^{\pm} , n, and 2k. Thus we can denote it by $\#O_{\kappa}^{prim}(a^{\pm}, n, 2k)$. Here,

 $\sqrt{ }$ \mathbf{J} 2k is the dimension of the nondegenerate split quadratic space $(H_k \otimes \kappa, \bar{q}_{\kappa});$

n is the dimension of the (possibly degenerate) quadratic space $(L' \otimes \kappa, \bar{q}_{L'})$;

 \mathbf{I} a^\pm is as explained above such that a is the dimension of a maximal nonsingular subspace of $L' \otimes \kappa$.

Note that n is the rank of L. In the following proposition, we will describe the integer a in terms of $GK(L)$.

Proposition 3.13. The integer a, which is defined to be the dimension of \bar{L}'_0 as a κ -vector space, is the same as the number of 0 's in $GK(L')$.

Proof. This directly follows from observation of a reduced form associated to the quadratic lattice $L.$

Since each summand in Equation (3.4) is determined by the number of 0's in $GK(L')$ (with the signature \pm) and $[L':L]$, we explain bounds of these two objects in this paragraph. Let $[L':L] = b$ and let n_0 be the number of 0's in $GK(L) = (a_1, \dots, a_n)$. Remark 2.8 yields that $|GK(L')| =$ $|GK(L)| - 2b$. The integer b is then nonnegative and at most $|GK(L)|/2$ since $GK(L') \succeq (0, \dots, 0)$ (cf. Remark 2.8). Let $GK(L') = (a'_1, \dots, a'_n)$ and let a be the number of 0's in $GK(L')$ (cf. Proposition [3.13\)](#page-15-1). If b is positive, then the integer a is at least $max\{n_0, n - |GK(L')|\}$.

We introduce a new notion $\mathcal{S}_{(L,a^{\pm},b)}$ to denote the set of all quadratic lattices L' including L for which the summands in Equation [\(3.4\)](#page-15-2) are equal. More precisely,

(3.6)
$$
\begin{cases} \mathcal{S}_{(L,a^+,b)} = \{L'(\supseteq L) | \text{GK}(L') \succeq (0,\cdots,0), [L':L] = b, a^+ \text{ is assigned to } L' \}; \\ \mathcal{S}_{(L,a^-,b)} = \{L'(\supseteq L) | \text{GK}(L') \succeq (0,\cdots,0), [L':L] = b, a^- \text{ is assigned to } L' \}. \end{cases}
$$

If a is odd or 0, then $\mathcal{S}_{(L,a^+,b)} = \mathcal{S}_{(L,a^-,b)}$. Note that $\mathcal{S}_{(L,a^{\pm},b)}$ is empty if $b > \frac{|\text{GK}(L)|}{2}$ by Remark 2.8. Therefore Equation [\(3.4\)](#page-15-2) is now reformulated as follows:

$$
(3.7) \qquad \alpha(L, H_k) = f^{-2kn + (n^2 + n)/2} \cdot \sum_{\substack{0 \le b \le |\text{GK}(L)|\\n_0 \le a \le n}}, f^{b \cdot (n+1-2k)} \cdot \# \mathcal{S}_{(L, a^{\pm}, b)} \cdot \# \mathcal{O}_{\kappa}^{prim}(a^{\pm}, n, 2k).
$$

Here, if a is odd or 0, then we ignore one of $\mathcal{S}_{(L,a^+,b)}$ or $\mathcal{S}_{(L,a^-,b)}$. If a is even and positive, then we count the summands involving $S_{(L,a^+,b)}$ and $S_{(L,a^-,b)}$ separately.

In the above equation, the number $\#\mathrm{O}_{\kappa}^{prim}(a^{\pm},n,2k)$ is already well-known as follows (cf. Exercise 4 in Section 5.6 of [\[Kit93\]](#page-50-20)):

$$
(3.8) \qquad \mathcal{O}_{\kappa}^{prim}(a^{\pm}, n, 2k) = f^{2kn - (n^2 + n)/2} (1 - f^{-k}) (1 + \chi(a^{\pm}) f^{n - a/2 - k}) \Pi_{1 \le i < n - a/2} (1 - f^{2i - 2k}).
$$

Here,

$$
\chi(a^{\pm}) = \begin{cases}\n0 & \text{if } a \text{ is odd;} \\
1 & \text{if } a \text{ is even and } a^+ \text{ is assigned or if } a = 0; \\
-1 & \text{if } a(> 0) \text{ is even and } a^- \text{ is assigned.} \n\end{cases}
$$

Using Equation [\(3.7\)](#page-16-4) combined with the above description of $\#O_{\kappa}^{prim}(a^{\pm}, n, 2k)$, we finally obtain the local density formula as follows:

Theorem 3.14. For a quadratic lattice L, we have

$$
\begin{split} \alpha(L,H_k) = \\ (1-f^{-k}) \cdot \sum_{\substack{0 \leq b \leq \frac{|\textup{GK}(L)|}{2n}} \ n \geq 2}} \left(\# \mathcal{S}_{(L,a^{\pm},b)} \cdot f^{b \cdot (n+1-2k)} \cdot (1+\chi(a^{\pm}) f^{n-a/2-k}) \prod_{1 \leq i < n-a/2} (1-f^{2i-2k}) \right). \end{split}
$$

Here, if a is odd or 0, then we ignore one of $\mathcal{S}_{(L,a^+,b)}$ or $\mathcal{S}_{(L,a^-,b)}$. If a is even and positive, then we count the summands involving $\mathcal{S}_{(L,a^+,b)}$ and $\mathcal{S}_{(L,a^-,b)}$ separately.

Definition 3.15. For a quadratic lattice L, the Siegel series is defined to be the polynomial $\mathcal{F}_L(X)$ of X such that

$$
\mathcal{F}_L(f^{-k}) = \alpha(L, H_k)
$$

for all integer k such that $k \geq n$, where n is the rank of L.

Corollary 3.16. From the formula of Theorem [3.14,](#page-16-0) we have the following description of the Siegel series:

$$
\mathcal{F}_L(X) =
$$
\n
$$
(1 - X) \cdot \sum_{\substack{0 \le b \le \frac{|{\rm GK}(L)|}{2} \\ n_0 \le a \le n}} \left(\# \mathcal{S}_{(L, a^{\pm}, b)} \cdot f^{b \cdot (n+1)} X^{2b} \cdot (1 + \chi(a^{\pm}) f^{n-a/2} X) \prod_{1 \le i < n-a/2} (1 - f^{2i} X^2) \right).
$$

Thus, each coefficient of the polynomial $\mathcal{F}_L(X)$ is determined by the order of the set $\mathcal{S}_{(L,a^{\pm},b)}$.

4. Inductive formulas of the Siegel series

A main goal of this section is to find an inductive formula of the Siegel series, by detailed investigation of the set of lattices $\mathcal{S}_{(L,a^{\pm},b)}$.

For a quadratic lattice L with $GK(L) = (a_1, \dots, a_n) (\neq (0, \dots, 0)),$ we choose a basis (e_1, \dots, e_n) of L such that with respect to this basis the symmetric matrix of the quadratic lattice L is optimal. Let d be the integer such that $a_{n-d} < a_{n-d+1} = \cdots = a_n$ \overbrace{d} . If $a_1 = \cdots = a_n$, then we let $d = n$. We

denote the lattice $(□ L)$ having a basis

$$
(e_1, \cdots, e_{n-d}, \underbrace{\frac{1}{\pi} \cdot e_{n-d+1}, \cdots, \frac{1}{\pi} \cdot e_n}_{d})
$$

by

$$
\begin{cases}\nL^{(d,n)} & \text{if } d > 1; \\
L^{(n)} \left(\text{or } L^{(1,n)} \right) & \text{if } d = 1.\n\end{cases}
$$

We assume that the quadratic form on $L^{(d,n)}$, naturally induced by the quadratic form on L, is an integral quadratic form, that is, $L^{(d,n)}$ is a quadratic lattice. For example, if $a_n \geq 2$, then $L^{(d,n)}$ is a quadratic lattice. Although it is not required that $a_n \geq 2$, the integers a_i 's should satisfy the condition that $a_n = \cdots = a_{n-d+1} \geq 1$, in order that $L^{(d,n)}$ is a quadratic lattice.

Let $\bar{V}_{L,d} = L^{(d,n)}/L$ be a κ -vector space of dimension d. Then each lattice between L and $L^{(d,n)}$ bijectively corresponds to each subspace of $\bar{V}_{L,d}$. More precisely, the set of all lattices L' between L and $L^{(d,n)}$ with degree $[L': L] = m$, where $0 \leq m \leq d$, equals the set of subspaces of $\bar{V}_{L,d}$ of dimension m. We denote the former set of lattices by $\mathcal{G}_{L,d,m}$. For example, $\mathcal{G}_{L,d,0} = \{L\}$ and $\mathcal{G}_{L,d,d} = \{L^{(d,n)}\}$. The latter set is the Grassmannian, denoted by $G(m,d)$, whose cardinality is well known to be $\binom{d}{m}$ $\binom{d}{m}$. Here,

$$
\binom{d}{m}_f = \frac{[(d)!]_f}{[m!]_f[(d-m)!]_f},
$$

where

$$
[m!]_f = \prod_{t=1}^m \frac{f^t - 1}{f - 1}, \text{ for any positive integer } m.
$$

We write that $[0!]_f = 1$ so that $\binom{d}{d}$ $\left(\begin{smallmatrix} d \\ d \end{smallmatrix}\right)_f = \left(\begin{smallmatrix} d \\ 0 \end{smallmatrix}\right)$ $\binom{d}{0}_f = 1$. For example, if $m = 1$, then $\binom{d}{1}$ $\binom{d}{1}_f = \frac{f^d-1}{f-1}$ $\frac{f^{\alpha}-1}{f-1}$. Thus we have the following formula:

(4.1)
$$
\#\mathcal{G}_{L,d,m} = \binom{d}{m}_{f}.
$$

In the following lemma, we explain a property of a lattice including L , but not an element of $\mathcal{G}_{L,d,m}.$

Lemma 4.1. Consider a lattice L' in V containing L. If L' does not contain any lattice in $\mathcal{G}_{L,d,m}$ for $1 \leq m \leq d$ (equivalently L' does not contain any lattice in $\mathcal{G}_{L,d,1}$), then there exists a direct summand M' of L' such that $L' = M' \oplus \underbrace{\mathfrak{0}e_{n-d+1} \oplus \cdots \oplus \mathfrak{0}e_n}$ \overbrace{d} and $L = M \oplus \underbrace{\mathfrak{oe}_{n-d+1} \oplus \cdots \oplus \mathfrak{oe}_n}$ \overbrace{d} ,

where $M = L \cap M'$.

Proof. Let $l = n - d$. For such L', we denote the image of e_i , with $l + 1 \leq i \leq n$, in $L'/\pi L'$ by \bar{e}_i . Let \bar{V}'_d be the subspace of $L'/\pi L'$, as a κ -vector space, spanned by $\bar{e}_{l+1}, \cdots, \bar{e}_n$. Since L' does not contain any lattice in $\mathcal{G}_{L,d,m}$ for $1 \leq m \leq d$, the vectors $(\bar{e}_{l+1},\cdots,\bar{e}_n)$ are linearly independent and thus the dimension of the vector space \bar{V}'_d is d.

Thus there are *l*-vectors $(\vec{e}'_1, \dots, \vec{e}'_l)$ in $L'/\pi L'$ having $(\vec{e}'_1, \dots, \vec{e}'_l, \vec{e}_{l+1}, \dots, \vec{e}_n)$ as a basis. Choose (e'_1, \dots, e'_l) in L' as preimages of $(\bar{e}'_1, \dots, \bar{e}'_l)$. By Nakayama's lemma, $(e'_1, \dots, e'_l, e_{l+1}, \dots, e_n)$ is a basis of $\overline{L'}$ as an $\mathfrak{o}\text{-module}$.

Let M' be the submodule of L' spanned by (e'_1, \dots, e'_l) so that $L'/M' = \mathfrak{o}e_{l+1} \oplus \dots \oplus \mathfrak{o}e_n$. We consider the following short exact sequence:

$$
1 \to L \cap M' \to L \to L'/M' \to 1.
$$

This short exact sequence splits since there exists a section from L'/M' to L such that e_i maps to e_i with $l+1 \leq i \leq n$. Thus $L \cong (L \cap M') \oplus (\mathfrak{oe}_{l+1} \oplus \cdots \oplus \mathfrak{oe}_n)$ as \mathfrak{o} -modules. Since this isomorphism is induced from the inclusions, we can identify L with $(L \cap M') \oplus (\mathfrak{oe}_{l+1} \oplus \cdots \oplus \mathfrak{oe}_n)$ as submodules of L'. Let $M = L \cap M'$. This completes the proof.

Lemma 4.2. For a basis (e_1, \dots, e_n) of L, consider a direct summand M[†] of L such that $L =$ $M^{\dagger} \oplus (\mathfrak{0} e_{n-d+1} \oplus \cdots \oplus \mathfrak{0} e_n)$. Then there is a basis of M^{\dagger} consisting of the column vectors of the $\overline{matrix \left(\begin{array}{c} id_{n-d} \\ z \end{array} \right)},$ where id_n \boldsymbol{x} \setminus , where id_{n-d} is the identity matrix of size $n-d$ and $x \in M_{d \times (n-d)}(\mathfrak{o})$.

Proof. A basis of $M^{\dagger} \oplus (\mathfrak{oe}_{n-d+1} \oplus \cdots \oplus \mathfrak{oe}_n)$ is given by the column vectors of the matrix $\begin{pmatrix} x_1 & 0 \\ x_2 & id \end{pmatrix}$ x_2 id_d \setminus with entries in \mathfrak{o} , where x_1 is a square matrix of size $n - d$. Since this matrix is invertible, x_1 is invertible over $\mathfrak o$ as well. Thus we can choose another basis for $L = M^{\dagger} \oplus (\mathfrak o e_{n-d+1} \oplus \cdots \oplus \mathfrak o e_n)$, given by the column vectors of the matrix $\begin{pmatrix} x_1 & 0 \\ x_2 & i d \end{pmatrix}$ x_2 id_d $\bigg\{ x_1^{-1} \quad 0$ 0 id_d \setminus = $\int id_{n-d}$ 0 $x_2x_1^{-1}$ id_d $\overline{}$. Let $x = x_2 x_1^{-1}$. This completes the proof. \Box

Remark 4.3. (1) In the situation of Lemma [4.1,](#page-18-0) a direct summand M of L has a basis given by the column vectors of a matrix $\int_0^{\infty} i d_{n-d}$ \overline{x} $\overline{ }$ with $x \in M_{d \times (n-d)}(\mathfrak{o})$ by Lemma [4.2.](#page-18-1) Such a lattice M is denoted by $L_x^{(d,n)}$, in order to emphasize both roles of x and (d, n) , so that

$$
L=L_x^{(d,n)}\oplus \underbrace{\mathfrak{0}e_{n-d+1}\oplus \cdots \oplus \mathfrak{0}e_n}_{d}.
$$

If $d = 1$, then we sometimes write $L_x^{(n)}$ to denote $L_x^{(1,n)}$.

The simplest case of $L_x^{(d,n)}$ is when x is the zero vector. In this case,

$$
L_0^{(d,n)} = \mathfrak{o}e_1 \oplus \cdots \oplus \mathfrak{o}e_{n-d}.
$$

The lattice $L_0^{(d,n)}$ will be crucially used in our inductive formula of the Siegel series (cf. Theorems [4.9](#page-23-0) and [5.11\)](#page-29-1).

(2) A basis of L such that with respect to this basis the symmetric matrix of L is optimal (reduced, respectively) is called an optimal basis (a reduced basis, respectively).

(3) If (e_1, \dots, e_n) is an optimal basis (resp. reduced basis) of L, then Theorem 0.2 (resp. Lemma 4.6) of [\[IK1\]](#page-50-12) yields that

$$
L_x^{(d,n)} \oplus \underbrace{\mathfrak{d}e_{n-d+1} \oplus \cdots \oplus \mathfrak{d}e_n}_{d}
$$

forms an optimal basis (resp. reduced basis).

- (4) Ikeda and Katsurada imposed extra invariant to a quadratic lattice L in addition to the Gross-Keating invariant and called it 'Extended Gross-Keating datum', denoted by $EGK(L)$ in [\[IK1\]](#page-50-12). They defined $EGK(L)$ based on an optimal form of L, which turns to be independent of the choice of an optimal form (cf. Theorem 0.4 and Definition 6.3 of [\[IK1\]](#page-50-12)). For an explicit description of $EGK(L)$, we refer to Definition 6.3 of [\[IK1\]](#page-50-12). A main contribution of their another paper [\[IK2\]](#page-50-10) is to prove that the Siegel series $\mathcal{F}_L(X)$ is completely determined by $EGK(L)$ (cf. Theorem 1.1 of [\[IK2\]](#page-50-10)).
- (5) Since column vectors of the matrix $\begin{pmatrix} id_{n-d} & 0 \\ \vdots & \vdots & \vdots \end{pmatrix}$ $x \quad id_d$ $\overline{ }$ form an optimal basis of L, Theorem 0.3 of [\[IK1\]](#page-50-12) yields that $GK(L_0^{(d,n)})$ $\mathcal{L}_{0}^{(d,n)}$ = GK($L_x^{(d,n)}$). In addition, by Definition 6.3 of [\[IK1\]](#page-50-12), one can easily see that $EGK(L_0^{(d,n)})$ $\mathcal{L}_{0}^{(d,n)}$) = EGK $(L_{x}^{(d,n)})$ since EGK (L) is independent from the choice of an optimal basis. Therefore, the argument of the above (4) implies that

$$
\mathcal{F}_{L_0^{(d,n)}}(X) = \mathcal{F}_{L_x^{(d,n)}}(X).
$$

From now on until the end of this section, we work with the following choice of a basis of a quadratic lattice L:

$$
(e_1, \dots, e_n)
$$
 is $\begin{cases} an \text{ optimal basis} & \text{if } p \text{ is odd;} \\ a \text{ reduced basis} & \text{if } p \text{ is even.} \end{cases}$

Before analyzing $\#\mathcal{S}_{(L,a^{\pm},b)}$ in Proposition [4.8,](#page-21-0) we will introduce one conjecture regarding quadratic forms modulo π in Conjecture [4.4,](#page-19-0) and prove it when p is odd or when L is anisotropic over \mathbb{Z}_2 in Lemmas [4.5](#page-19-1)[-4.6.](#page-20-0)

We write $L = M \oplus N$, where M (respectively N) is spanned by (e_1, \dots, e_{n-d}) (respectively (e_{n-d+1}, \cdots, e_n) \overbrace{d})). We consider a lattice L' containing L in $L \otimes_{\mathfrak{o}} F$ of the form $L' = M' \oplus N$, where

M' is a lattice containing M in $M \otimes_{\mathfrak{o}} F$. Let $q_{L'}$ be the quadratic form on L' which is naturally induced from q_L on L such that $q_{L'}|_L = q_L$. Similarly, we define the quadratic form $q_{M'}$ defined on M' .

In the case that $q_{M'}$ is integral (equivalently, $GK(M') \succeq (0, \dots, 0)$), we define $\overline{M}' = M'/\pi M'$ to be the quadratic space over κ having the quadratic form $q_{M'}$ modulo π . Similarly, in the case that $q_{L'}$ on \tilde{L}' is integral, we define the quadratic space $\bar{L}' = \tilde{L}'/\pi L'$.

If $q_{L'}$ is integral, then $q_{M'}$ is also integral. Conversely, we propose the following conjecture.

Conjecture 4.4. If $q_{M'}$ on M' is integral, then $q_{L'}$ on L' is integral (thus they are equivalent). In this situation, the dimension of \bar{L}' modulo the radical, is the same as the dimension of \bar{M}' modulo the radical. In other words, the number of 0 's in $GK(L')$ is the same as the number of 0 's in $GK(M')$ (cf. Proposition [3.13\)](#page-15-1).

We think that the conjecture is true in the general case. In the following, we will prove it in two cases, when p is odd or when (L, q_L) is an anisotropic \mathbb{Z}_2 -lattice.

Lemma 4.5. The conjecture is true for an odd prime p.

Proof. Let $B =$ $\int a \quad b$ $\begin{pmatrix} a & b \\ t & c \end{pmatrix}$ be an optimal form with respect to (e_1, \dots, e_n) , where the size of a is $(n - d) \times (n - d)$ and the size of c is $d \times d$. A symmetric matrix of L' with respect to the decomposition $L' = M' \oplus N$ such that a basis of N is fixed by (e_{n-d+1}, \dots, e_n)) is of the form

$$
\begin{pmatrix} t_x & 0 \\ 0 & id \end{pmatrix} \cdot \begin{pmatrix} a & b \\ t_b & c \end{pmatrix} \cdot \begin{pmatrix} x & 0 \\ 0 & id \end{pmatrix} = \begin{pmatrix} t_x \cdot a \cdot x & t_x \cdot b \\ t_b \cdot x & c \end{pmatrix}
$$
 for certain $x \in GL_{n-d}(F)$. Here, $t_x \cdot a \cdot x$ is a symmetric matrix of M' . Thus it suffices to show that the exponential order of each entry of $2 \cdot t_x \cdot b$ is at least 1.

Since p is odd, we can choose another basis of M given by the column vectors of $y \in GL_{n-d}(\mathfrak{o})$ such that $t_y \cdot a \cdot y$ is an optimal and diagonal matrix by Remark 2.8.(1). Then by Theorems 0.2-0.3 of [\[IK1\]](#page-50-12), the symmetric matrix $\begin{pmatrix} t_y & a & y & t_y \cdot b \\ t_{b} & y & c \end{pmatrix}$ $\begin{pmatrix} a \cdot y & ^t y \cdot b \\ b \cdot y & c \end{pmatrix}$ is also optimal with respect to a basis forming $L = M \oplus N$. Thus we may and do assume that a is a diagonal matrix with the *i*-th diagonal entry $u_i \pi^{a_i}$, where u_i is a unit in \mathfrak{o} .

Since we assumed that M' is a quadratic lattice containing M in $M \otimes_{\mathfrak{o}} F$ with p odd, the lattice M' is contained in the dual lattice of M, which is defined to be the set $\{v \in M \otimes_{\mathfrak{g}} F | b_{q_M} (v, L) \in \mathfrak{g} \}$. Here, b_{q_M} is the symmetric bilinear form associated to the quadratic form q_M such that $b_{q_M}(v, v) = q_M(v)$. The dual lattice of M is spanned by $(\pi^{-a_1}e_1, \dots, \pi^{-a_{n-d}}e_{n-d})$. Thus, if x_{n-d} is the diagonal matrix of size $n-d$ whose *i*-th diagonal entry is π^{-a_i} , then a matrix x determining M' is of the form $x_{n-d} \cdot x'$, where $x' \in GL_{n-d}(F) \cap M_{n-d}(\mathfrak{o})$.

On the other hand, the exponential order of each entry of 2 · ${}^t x_{n-d} \cdot b$ is at least 1 since B is optimal and $a_i < a_{n-d+1} = a_n$ for any $i \leq n-d$). Therefore, the exponential order of each entry of $2 \cdot {}^t(x') \cdot {}^t x_{n-d} \cdot b$ is at least 1 since $x' \in M_{n-d}(0)$. This completes the proof. of $2 \cdot {}^t(x') \cdot {}^t x_{n-d} \cdot b$ is at least 1 since $x' \in M_{n-d}(0)$. This completes the proof. □

Lemma 4.6. The conjecture is true for an anisotropic quadratic \mathbb{Z}_p -lattice with any p.

Proof. Let L be an anisotropic quadratic lattice over \mathbb{Z}_p and let $GK(L) = (a_1, \dots, a_n)$ so that n is at most 4. It is well known that there is a unique maximal quadratic lattice in $L \otimes_{\mathfrak{g}} F$ by Theorem 91:1 of $[O70]$. Thus it suffices to prove the conjecture when M' is the maximal quadratic lattice inside $M \otimes_{\mathfrak{o}} F$.

We can easily prove that any anisotropic quadratic lattice whose Gross-Keating invariant consists of 0 and 1 is maximal since only two among a_i 's have the same parity (cf. Proposition [5.3\)](#page-25-1).

As in the proof of Lemma [4.5,](#page-19-1) let $B =$ $\int a b$ $\begin{pmatrix} a & b \\ t_b & c \end{pmatrix}$ be a reduced form with respect to (e_1, \dots, e_n) , where the size of a is $(n - d) \times (n - d)$ and the size of c is $d \times d$. Let x_{n-d} be the diagonal matrix of size $n - d$ whose *i*-th diagonal entry is $\pi^{-[a_i/2]}$. The proof of Proposition 3.2 in [\[IK1\]](#page-50-12) implies that the Gross-Keating invariant of ${}^t x_m \cdot a \cdot x_m$ consists of 0 and 1 so that the associated quadratic lattice is maximal. A simple calculation shows that the exponential order of each entry of $2 \cdot {}^t x_m \cdot b$ is at least 1. This completes the proof. \Box

Remark 4.7. In the proof of the above lemma, the only place to use the assumption of $o = \mathbb{Z}_p$ is that any anisotropic quadratic lattice whose Gross-Keating invariant consists of 0 and 1 is maximal which follows from Proposition [5.3.](#page-25-1) If this is true for a general ρ , then the proof of the lemma works so that the conjecture is true for an anisotropic quadratic lattice over \mathfrak{o} .

Using Lemmas [4.1](#page-18-0)[-4.2](#page-18-1) and assuming Conjecture [4.4,](#page-19-0) we will explain an inductive formula of $\#\mathcal{S}_{(L,a^{\pm},b)}$ in the following proposition.

Proposition 4.8. Assume that Conjecture [4.4](#page-19-0) is true. Let b be an integer such that $0 \le b \le \frac{|{\rm GK}(L)|}{2}$ $\frac{\Delta(L)}{2}$. Then for any integer b' with $b' \geq b$, we have the following formula:

$$
\#\mathcal{S}_{(L,a^{\pm},b)} = \sum_{m=1}^{d} \left(c_m \cdot \sum_{L' \in \mathcal{G}_{L,d,m}} \# \mathcal{S}_{(L',a^{\pm},b-m)} \right) + f^{d(b-(n-d)b')} \sum_{x \in M_{d \times (n-d)}(\mathfrak{o}/\pi^{b'} \mathfrak{o})} \# \mathcal{S}_{(L_x^{(d,n)},a^{\pm},b)},
$$

where $c_m = (-1)^{m-1} f^{m(m-1)/2}$. Here, if $b - m < 0$, then we understand $\#\mathcal{S}_{(L',a^{\pm},b-m)} = 0$. And $L_x^{(d,n)}$ for $x \in M_{d \times (n-d)}(\mathfrak{o}/\pi^{b'}\mathfrak{o})$ stands for a lattice associated to any representative of x in $M_{d\times (n-d)}$. It depends on such representative but $\#\mathcal{S}_{(L_{x}^{(d,n)},a^{\pm},b)}$ does not.

If $\#\mathcal{S}_{(L,a^{\pm},b)}=0$, then each $\#\mathcal{S}_{(L',a^{\pm},b-m)}$ and $\#\mathcal{S}_{(L_x^{(d,n)},a^{\pm},b)}$ are zero as well.

Proof. Note that $\mathcal{S}_{(L',a^{\pm},b-m)} \subseteq \mathcal{S}_{(L,a^{\pm},b)}$ for $L' \in \mathcal{G}_{L,d,m}$. We choose a lattice

$$
L^{\dagger} \in \mathcal{S}_{(L,a^{\pm},b)} \setminus \bigcup_{L' \in \mathcal{G}_{L,d,1}} \mathcal{S}_{(L',a^{\pm},b-1)},
$$

if the right hand side is nonempty. By Lemmas [4.1-](#page-18-0)[4.2](#page-18-1) and Remark 4.3.(1), there exists a direct summand $L_x^{(d,n)}$ of L such that

$$
\begin{cases}\nL = L_x^{(d,n)} \oplus \underbrace{\mathfrak{0}e_{n-d+1} \oplus \cdots \oplus \mathfrak{0}e_n}_{d}; \\
L^{\dagger} = (L_x^{(d,n)})^{\dagger} \oplus \underbrace{\mathfrak{0}e_{n-d+1} \oplus \cdots \oplus \mathfrak{0}e_n}_{d},\n\end{cases}
$$

as an o-lattice (not as a quadratic o-lattice). Here, $(L_x^{(d,n)})^{\dagger}$ is a direct summand of L^{\dagger} satisfying the condition that $L \cap (L_x^{(d,n)})^{\dagger} = L_x^{(d,n)}$. Note that if (e_1, \dots, e_n) is an optimal basis (resp. reduced basis) of L , then the above basis for L is also optimal (resp. reduced) by Remark 4.3.(3). Thus we can use Conjecture [4.4](#page-19-0) with $L = L_x^{(d,n)} \oplus \underbrace{\mathfrak{oe}_{n-d+1} \oplus \cdots \oplus \mathfrak{oe}_n}$ \overbrace{d} . More precisely, since

 $[L^{\dagger}:L] = [(L_x^{(d,n)})^{\dagger}:L_x^{(d,n)}] = b$, the lattice $(L_x^{(d,n)})^{\dagger}$ is contained in $\mathcal{S}_{(L_x^{(d,n)},a^{\pm},b)}$ by Conjecture [4.4.](#page-19-0) Thus by Conjecture [4.4](#page-19-0) again, we have that

d

$$
(4.2) \qquad \mathcal{S}_{(L,a^{\pm},b)} \setminus \bigcup_{L' \in \mathcal{G}_{L,d,1}} \mathcal{S}_{(L',a^{\pm},b-1)} = \bigcup_{x \in M_{d \times n-d}(\mathfrak{o})} \mathcal{S}_{(L_x^{(d,n)},a^{\pm},b)} \oplus \underbrace{\mathfrak{o}e_{n-d+1} \oplus \cdots \oplus \mathfrak{o}e_n}_{d}.
$$

Here,
$$
S_{(L_x^{(d,n)},a^{\pm},b)} \oplus \underbrace{\mathfrak{0}e_{n-d+1} \oplus \cdots \oplus \mathfrak{0}e_n}_{d}
$$
 is the set of lattices $\{M' \oplus \underbrace{\mathfrak{0}e_{n-d+1} \oplus \cdots \oplus \mathfrak{0}e_n}_{d}|M' \in$

 $\mathcal{S}_{(L_x^{(d,n)},a^{\pm},b)}$. The above equation holds even when the left hand side is empty, by assuming Conjecture [4.4.](#page-19-0) This confirms the last sentence of the proposition.

Since $[M': L_x^{(d,n)}] = b$ for $M' \in \mathcal{S}_{(L_x^{(d,n)}, a^{\pm}, b)}$, we can see that

$$
\mathcal{S}_{(L_x^{(d,n)},a^{\pm},b)} \oplus \underbrace{\mathfrak{o}e_{n-d+1} \oplus \cdots \oplus \mathfrak{o}e_n}_{d} = \mathcal{S}_{(L_y^{(d,n)},a^{\pm},b)} \oplus \underbrace{\mathfrak{o}e_{n-d+1} \oplus \cdots \oplus \mathfrak{o}e_n}_{d} \text{ if } x \equiv y \mod \pi^b.
$$

Thus Equation [\(4.2\)](#page-21-1) can be expressed as follows:

$$
(4.3) \qquad \mathcal{S}_{(L,a^{\pm},b)} \setminus \bigcup_{L' \in \mathcal{G}_{L,d,1}} \mathcal{S}_{(L',a^{\pm},b-1)} = \bigcup_{x \in M_{d \times (n-d)}(\mathfrak{o}/\pi^{b'}\mathfrak{o})} \mathcal{S}_{(L_x^{(d,n)},a^{\pm},b)} \oplus \underbrace{\mathfrak{o}e_{n-d+1} \oplus \cdots \oplus \mathfrak{o}e_n}_{d}
$$

for any integer b' such that $b' \geq b$. Here, $L_x^{(d,n)}$ for $x \in M_{d \times (n-d)}(\mathfrak{o}/\pi^{b'}\mathfrak{o})$ stands for a lattice associated to any representative of x in $M_{d \times (n-d)}(\mathfrak{o})$. It depends on such representative but $\mathcal{S}_{(L_{x}^{(d,n)},a^{\pm},b)}\oplus \underbrace{\mathfrak{o} e_{n-d+1}\oplus\cdots\oplus \mathfrak{o} e_n}$ \overbrace{a} does not.

d In order to compute the cardinality of the right hand side of Equation [\(4.3\)](#page-21-2), we compare it with

$$
\sum_{x \in M_{d \times (n-d)}(\mathfrak{o}/\pi^{b'} \mathfrak{o})} \# \left(\mathcal{S}_{(L_{x}^{(d,n)},a^{\pm},b)} \oplus \underbrace{\mathfrak{o} e_{n-d+1} \oplus \cdots \oplus \mathfrak{o} e_{n}}_{d} \right).
$$

In the following, we will see how many times a given lattice is counted in this sum.

We choose a lattice L_x^{\dagger} in $\mathcal{S}_{(L_x^{(d,n)},a^{\pm},b)} \oplus \underbrace{\mathfrak{o}e_{n-d+1} \oplus \cdots \oplus \mathfrak{o}e_n}$ \overbrace{d} . The cardinality of the set $\{y \in$

$$
M_{d \times (n-d)}(\mathfrak{o}/\pi^{b'}\mathfrak{o})|L_x^{\dagger} \in \mathcal{S}_{(L_y^{(d,n)},a^{\pm},b)} \oplus \underbrace{\mathfrak{o}e_{n-d+1} \oplus \cdots \oplus \mathfrak{o}e_n}_{d} \} \text{ is then}
$$

$$
\frac{\#(\mathfrak{o}/\pi^{b'-b}\mathfrak{o}) \cdot \cdots \cdot \#(\mathfrak{o}/\pi^{b'-b}\mathfrak{o})}{d} \cdot \frac{\#(\mathfrak{o}/\pi^{b'}\mathfrak{o}) \cdot \cdots \cdot \#(\mathfrak{o}/\pi^{b'}\mathfrak{o})}{d(n-d-1)}} = f^{d(n-d)b'-db}.
$$

Note that the above number is independent of the choice of x. Since

$$
\#\left(\mathcal{S}_{(L_x^{(d,n)},a^\pm,b)}\oplus \underbrace{\mathfrak{o} e_{n-d+1}\oplus \cdots \oplus \mathfrak{o} e_n}_{d}\right)=\#\left(\mathcal{S}_{(L_x^{(d,n)},a^\pm,b)}\right),
$$

we have the following equation:

$$
\#\left(\mathcal{S}_{(L,a^\pm,b)}\setminus \bigcup_{L'\in \mathcal{G}_{L,d,1}} \mathcal{S}_{(L',a^\pm,b-1)}\right) = f^{db-d(n-d)b'}\cdot \sum_{x\in M_{d\times (n-d)}(\mathfrak{o}/\pi^{b'}\mathfrak{o})} \#\mathcal{S}_{(L_x^{(d,n)},a^\pm,b)}.
$$

Thus to complete the proof, it suffices to show that

(4.4)
$$
\#\left(\bigcup_{L' \in \mathcal{G}_{L,d,1}} \mathcal{S}_{(L',a^{\pm},b-1)}\right) = \sum_{m=1}^d \left(c_m \cdot \sum_{L' \in \mathcal{G}_{L,d,m}} \#\mathcal{S}_{(L',a^{\pm},b-m)}\right),
$$

where $c_m = -\left({m \choose 1}_f \cdot c_1 + {m \choose 2}_f \cdot c_2 + \cdots + {m \choose m-1}_f \cdot c_{m-1} \right) + 1$ if $m > 1$ and $c_1 = 1$. This follows from inclusion-exclusion principle using the counting argument of the Grassmannian

given at the beginning of this section.

For the proof^{[2](#page-22-0)} of $c_m = (-1)^{m-1} f^{m(m-1)/2}$, we consider the q-binomial formula

$$
\prod_{k=0}^{m-1} (1 + f^k t) = \sum_{k=0}^{m} f^{k(k-1)/2} \binom{m}{k}_f t^k.
$$

Putting $t = -1$, we have

$$
\sum_{k=0}^{m} (-1)^k f^{k(k-1)/2} \binom{m}{k}_f = 0.
$$

Then by induction, we can easily see that $c_m = (-1)^{m-1} f^{m(m-1)/2}$

We now state our main theorem of this section, an inductive formula of the Siegel series $\mathcal{F}_L(X)$.

. — Первый процесс в постановки программа в серверном становки производительно становки производительно станов
В серверном становки производительно становки производительно становки производительно становки производительн

²This proof was informed by the referee.

Theorem 4.9. Assume that Conjecture [4.4](#page-19-0) is true. Then we have the following inductive formula, with respect to the Gross-Keating invariant, of the Siegel series $\mathcal{F}_L(X)$:

$$
\mathcal{F}_L(X) = \sum_{m=1}^d \left(c_m \cdot f^{(n+1)m} \cdot X^{2m} \cdot \sum_{L' \in \mathcal{G}_{L,d,m}} \mathcal{F}_{L'}(X) \right) +
$$

$$
(1-X)(1-f^dX)^{-1} \cdot \left(\prod_{i=1}^d (1-f^{2i}X^2) \right) \cdot \mathcal{F}_{L_0^{(d,n)}}(f^dX),
$$

where $c_m = (-1)^{m-1} f^{m(m-1)/2}$. Here, for $L' \in \mathcal{G}_{L,d,m}$,

$$
\left\{\begin{array}{l}\operatorname{GK}(L)\succ\operatorname{GK}(L');\\ \left|\operatorname{GK}(L')\right|=\left|\operatorname{GK}(L)\right|-2m;\\ \operatorname{GK}(L_0^{(d,n)})=\operatorname{GK}(L)^{(n-d)}.\end{array}\right.
$$

Note that notion of $L^{(d,n)}$, $\mathcal{G}_{L,d,m}$, and $\binom{m}{k}_f$ can be found at the beginning of this section. Notion of $L_0^{(d,n)}$ $a_0^{(a,n)}$ can be found at Remark 4.3.(1).

Proof. If we plug the formula of Proposition [4.8](#page-21-0) into the formula of Theorem [3.14,](#page-16-0) then we obtain

$$
\alpha(L, H_k) = \sum_{m=1}^d \left(c_m \cdot f^{(n-2k+1)m} \sum_{L' \in \mathcal{G}_{L,d,m}} \alpha(L', H_k) \right) +
$$

$$
f^{-d(n-d)b'} (1 - f^{-k})(1 - f^{-(k-d)})^{-1} \cdot \left(\prod_{i=1}^d (1 - f^{2i-2k}) \right) \cdot \sum_{x \in M_{d \times (n-d)}(\mathfrak{o}/\pi^{b'} \mathfrak{o})} \alpha(L_x^{(d,n)}, H_{k-d}),
$$

where $c_m = -\left(\binom{m}{1}_f \cdot c_1 + \binom{m}{2}_f \cdot c_2 + \cdots + \binom{m}{m-1}_f \cdot c_{m-1}\right) + 1$ if $m > 1$ and $c_1 = 1$.

On the other hand, as mentioned at Remark 4.3.(5), we have that

$$
\alpha(L_x^{(d,n)}, H_{k-d}) = \alpha(L_0^{(d,n)}, H_{k-d})
$$

for any $x \in M_{d \times (n-d)}(\mathfrak{o}/\pi^{b'}\mathfrak{o})$. This completes the proof of the inductive formula. The rest follows from Theorems 0.1 and 0.3 of [\[IK1\]](#page-50-12). \Box

Corollary 4.10. Assume that Conjecture [4.4](#page-19-0) is true. If $a_{n-1} < a_n$ for $GK(L) = (a_1, \dots, a_n)$ so that $d = 1$, then the above inductive formula turns to be

$$
\mathcal{F}_L(X) = f^{n+1} \cdot X^2 \cdot \mathcal{F}_{L^{(n)}}(X) + (1-X)(1+fX) \cdot \mathcal{F}_{L_0^{(n)}}(fX).
$$

Note that notion of $L^{(n)}$ can be found at the beginning of this section and that notion of $L_0^{(n)}$ $\int_0^{(n)}$ can be found at Remark 4.3.(1).

Remark 4.11. We note that Katsurada described several interesting inductive formulas in The-orem 2.6 of [\[Kat99\]](#page-50-9). Theorem 2.6.(1) in loc. cit. is the same as Corollary [4.10](#page-23-1) when p is odd. More generally, our inductive formula in Theorem [4.9](#page-23-0) is directly followed from Proposition 2.2 combined with Proposition 2.5.(1) in loc. cit., when p is odd. On the other hand, when $p = 2$, the inductive formula of Theorem [4.9](#page-23-0) is sharper than that of Theorem 2.6 in [\[Kat99\]](#page-50-9), due to restriction in Proposition 2.5.(1) of loc. cit. Thus our result is a more refined version of Katsurada's formula stated in Theorem 2.6 of [\[Kat99\]](#page-50-9).

The following lemma is used in the main theorems [5.10](#page-28-0)[-5.11](#page-29-1) of the next section.

Lemma 4.12. Let $B =$ $\begin{pmatrix} B' & C \\ ^tC & d \end{pmatrix}$ be a reduced form with $GK(B) = (a_1, \dots, a_{n-1}, a_n)$. Here, B' is of size $(n-1) \times (n-1)$ and $d \in \mathfrak{o}$. We assume that B' is also a reduced form with $GK(B') =$ (a_1, \dots, a_{n-1}) and that B satisfies one of the followings:

$$
\begin{cases} a_n > a_{n-1}; \\ a_n = a_{n-1} > a_{n-2}, \sigma(n-1) = n, \text{ ord}(d) = a_n, \text{ and } \mathfrak{o} = \mathbb{Z}_2. \end{cases}
$$

Let $B_x = {}^t \left(\begin{matrix} id_{n-1} \\ x \end{matrix} \right)$ \boldsymbol{x} \setminus $\cdot B \cdot$ $\int i d_{n-1}$ \boldsymbol{x}), where $x \in M_{1 \times (n-1)}(0)$. Let $\mathcal{F}_{B_x}(X)$ be the Siegel series of the quadratic lattice associated to the symmetric matrix B_x . Then we have

$$
\mathcal{F}_{B_0}(X)=\mathcal{F}_{B_x}(X)
$$

for any x.

Proof. We write

$$
B_x = (id_{n-1} \t t_x) \cdot \begin{pmatrix} B' & C \ t_C & d \end{pmatrix} \cdot \begin{pmatrix} id_{n-1} \ x \end{pmatrix} = B' + (Cx + {}^t(Cx) + d \cdot {}^txx).
$$

The (i, j) -th entry of $(Cx + {}^t(Cx) + d \cdot {}^txx)$ is $c_ix_j+c_jx_i+dx_ix_j$. Here, we write ${}^tC = (c_1, \dots, c_{n-1})$. Then we have the following:

$$
\begin{cases} \operatorname{ord}(c_i x_i + c_i x_i + d x_i x_i) > a_i; \\ \operatorname{ord}(2(c_i x_j + c_j x_i + d x_i x_j)) > \frac{a_i + a_j}{2} \text{ if } i < j. \end{cases}
$$

This, combined with ^{[3](#page-24-1)}Theorems 3.3 and 1.1 of [\[IK2\]](#page-50-10), completes the proof. \square

5. On the Siegel series of anisotropic quadratic lattices

In this section, we will explain a more refined inductive formula of the Siegel series of anisotropic quadratic lattices defined over \mathbb{Z}_p , so as to compare it with an inductive formula of local intersection multiplicities of [\[GK93\]](#page-50-0) in Sections [6](#page-29-0)[-7.](#page-33-0) Thus throughout this section, let (L, q_L) be an anisotropic quadratic lattice over \mathbb{Z}_p . Here, we say that (L, q_L) is anisotropic if $q_L(x) \neq 0$ for any nonzero element $x \in L$. Then the rank of L is at most 4. We first list a few necessary facts about (L, q_L) .

Remark 5.1. (1) Let D be the unique quaternion division algebra over \mathbb{Q}_p . Let q_D be the associated quadratic form, which is defined to be the reduced norm on D. There is a unique maximal order of D , denoted by O_D . The maximal order O_D is characterized as follows:

$$
O_D = \{ v \in D | q_D(v) \in \mathbb{Z}_p \}.
$$

(2) The pair (O_D, q_D) is an anisotropic quadratic lattice of rank 4 over \mathbb{Z}_p . By Exercise 3 of Section 5.2 in [\[Kit93\]](#page-50-20), there is a suitable basis of O_D such that with respect to this basis the symmetric matrix of O_D is described as follows:

$$
\left\{\begin{array}{ll} (1)\perp(-\delta)\perp(p)\perp(-\delta p) & \text{ if } p\neq 2; \\ \begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix}\perp\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} & \text{ if } p=2. \end{array}\right.
$$

Here, δ is a unit in \mathbb{Z}_p (with $p \neq 2$) such that δ modulo p is not a square. These symmetric matrices are reduced forms. Thus we can see that

$$
GK(O_D) = (0, 0, 1, 1).
$$

³Indeed Ikeda and Katsurada assume that $p = 2$ in Theorem 3.3, loc. cit. But this theorem also holds for $p > 2$ and it was explained in the initial version of their paper posted on arXiv.

(3) Any anisotropic quadratic lattice over \mathbb{Z}_p of rank $n \leq 4$) is always embedded into (O_D, q_D) , which can be shown by using Theorem 3.5.1 and Corollary 3.5.4 of [\[Kit93\]](#page-50-20). Thus we may regard (L, q_L) as a sublattice of (O_D, q_D) .

The Gross-Keating invariants of anisotropic quadratic lattices over \mathbb{Z}_p with rank ≤ 3 are well explained in [\[Bou07\]](#page-50-21). In the next subsection, we will explain the Gross-Keating invariants and some properties of anisotropic quadratic lattices of rank 4.

5.1. On an anisotropic quadratic lattice of rank 4.

Lemma 5.2. Consider an anisotropic quadratic lattice of rank 2 over \mathbb{Z}_2 . Choose a basis (e_1, e_2) such that with respect to this basis the half-integral symmetric matrix is $X =$ $\begin{pmatrix} a & b \\ b & c \end{pmatrix}$. Assume that $GK(X) = (a_1, a_1 + 2t)$ for $t \geq 0$. If X is an optimal form, then we have the following:

 $\text{ord}(a) = a_1, \text{ord}(2b) = a_1 + t, \text{ and } \text{ord}(c) = a_1 + 2t.$

Thus any optimal form of an anisotropic quadratic lattice of rank 2 over \mathbb{Z}_2 is a reduced form.

Proof. Assume that X with $GK(X) = (a_1, a_2)$ is diagonalizable. We claim that $a_1 < a_2$. Let $X'=\begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix}$ $0 \t y$ \setminus , where $\text{ord}(x) \leq \text{ord}(y)$, be a diagonal matrix isometric to X. The norm of X is $(2^{\text{ord}(x)})$ so that $a_1 = \text{ord}(x)$ by Remark 2.8.(4). Assume that $\text{ord}(y) > \text{ord}(x)$. Since $GK(X) =$ $(a_1, a_2) \succeq (ord(x), ord(y)),$ we have that $a_2 \geq ord(y) > ord(x) = a_1$ and thus $a_2 > a_1$. Assume that $\text{ord}(x) = \text{ord}(y)$. Consider the following matrix equation:

$$
\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} x & x \\ x & x+y \end{pmatrix}.
$$

Since ord $(x + y)$ is at least $a_1 + 1$, $GK(X) \succeq (a_1, a_1 + 1)$. Thus we have that $a_1 < a_2$.

We first assume that $t = 0$. Since $GK(X) = (a_1, a_1)$, X is not diagonalizable by the above claim. The lemma then follows from Lemma 3.2.(b) of [\[Bou07\]](#page-50-21).

For $t > 0$, it is easy to see that the symmetric matrix with respect to the basis $(e_1, \frac{1}{2^t})$ $\frac{1}{2^t} \cdot e_2$) is an optimal form with $GK = (a_1, a_1)$ by using Remark 2.8.(4)-(5). Using the result of the case $t = 0$ completes the proof. \Box

If we write (a_1, a_2, a_3) to be the Gross-Keating invariant of an anisotropic quadratic lattice of rank 3 over \mathbb{Z}_p for any p, then only two of a_i 's have the same parity by Lemma 5.3 of [\[Bou07\]](#page-50-21). In the following proposition, we prove the same statement in the case of $n = 4$.

Proposition 5.3. Let $GK(L) = (a_1, a_2, a_3, a_4)$. Then only two of a_i 's have the same parity. This holds for any prime p.

Proof. Since L is a sublattice of O_D with the same rank 4, the parity of $|GK(L)|$ is the same as that of $|GK(O_D)|$ by Remark 2.8.(5). Since $GK(O_D) = (0, 0, 1, 1)$ and $|GK(O_D)| = 2$, either only two of a_i 's have the same parity or all of a_i 's have the same parity. Assume that all of a_i 's have the same parity.

If $p \neq 2$, then we choose a basis (e_1, e_2, e_3, e_4) for L, whose associated symmetric matrix is a diagonal matrix having $u_i p^{a_i}$ as the *i*-th diagonal entry, where u_i is a unit in \mathbb{Z}_p , by Remark 2.8.(1). Thus the Gross-Keating invariant of the sublattice L' spanned by (e_1, e_2, e_3) is (a_1, a_2, a_3) . Note that L' is anisotropic with $GK(L') = (a_1, a_2, a_3)$. This is a contradiction since a_1, a_2, a_3 have the same parity.

If $p = 2$, then we choose a reduced basis (e_1, e_2, e_3, e_4) of (L, q_L) . Since a_i 's have the same parity, we can choose an involution σ attached to (a_1, a_2, a_3, a_4) such that $\sigma(1) = 2, \sigma(3) = 4$ (cf. Definition [2.4\)](#page-8-2). Note that σ is unique if $a_2 < a_3$. Let $B = (b_{ij})$ be the associated symmetric matrix. The definition of a reduced form (cf. Defintion [2.5\)](#page-8-0) then tells us that two matrices of size 2

associated to (e_1, e_2) and (e_3, e_4) are also reduced forms. Lemma [5.2](#page-25-2) yields that $\text{ord}(b_{ii}) = a_i$. Thus the symmetric matrix associated to a basis (e_1, e_2, e_3) is also a reduced form whose Gross-Keating invariant is (a_1, a_2, a_3) , where the associated involution permutes 1 and 2. This is a contradiction to the fact that the lattice spanned by (e_1, e_2, e_3) is anisotropic, since a_1, a_2, a_3 have the same \Box

Proposition 5.4. Let B be a reduced form of an anisotropic quadratic lattice L of rank 4 with $GK(L) = (a_1, a_2, a_3, a_4)$. Let (e_1, e_2, e_3, e_4) be a basis of a reduced form B. If p is odd, then we consider B as a diagonal matrix.

Then any (3×3) -submatrix of the matrix B, with respect to a basis (e_i, e_j, e_k) among (e_1, e_2, e_3, e_4) , is a reduced form whose Gross-Keating invariant is (a_i, a_j, a_k) , where $i < j < k$.

Similarly, any (2×2) -submatrix of the matrix B, with respect to a basis (e_i, e_j) among (e_1, e_2, e_3, e_4) , is a reduced form whose Gross-Keating invariant is (a_i, a_j) , where $i < j$.

Proof. If p is odd, then it is clear. When $p = 2$, the proof is similar to the argument used in the proof of the above proposition. Thus we skip it. \Box

Remark 5.5. Let $GK(L) = (a_1, a_2, a_3, a_4)$ with $a_4 \geq 2$ and let $p = 2$. We choose a reduced basis (e_1, e_2, e_3, e_4) with an involution σ . Let $\sigma(4) = i$ and $\sigma(j) = k$ so that j, k are odd and $\{1,2,3,4\} = \{4, i, j, k\}.$ To ease notation, we say $\sigma(a_4) = a_i$ and $\sigma(a_i) = a_k$ if there is no confusion. Then by using a similar argument used in the proof of Proposition [5.3,](#page-25-1) we can easily see that $(e_1, e_2, e_3, 1/p \cdot e_4)$ is a reduced basis of $L^{(4)}$, up to a permutation, such that $GK(L^{(4)}) =$ $(a_1, a_2, a_3) \cup (a_4 - 2)$. A key point of the proof is to use the fact that $\text{ord}(q_L(e_i)) = a_i$. The associated involution to $L^{(4)}$ is compatible with σ so as to exchange $a_4 - 2$ and a_i , and exchange a_i and a_k .

If p is odd, then we can choose a diagonal matrix as a reduced form of L . Then the Gross-Keating invariant consists of orders of each diagonal entries (cf. Remark 2.8.(1)).

We assume that $p = 2$. Let B be an half-integral symmetric matrix associated to (L, q_L) of rank 4. By using Theorem 2.4 of [\[Cho15\]](#page-50-18) and Lemma 3.2.(c) of [\[Bou07\]](#page-50-21), there are three types of B, up to equivalence, as follows:

$$
\begin{cases}\n\text{Case (I)}: B = 2^i \begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix} \perp 2^j \begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix}; \\
\text{Case (II)}: B = 2^i \begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix} \perp (u_1 2^{\mu_1}) \perp (u_2 2^{\mu_2}); \\
\text{Case (III)}: B = (u_1 2^{\mu_1}) \perp (u_2 2^{\mu_2}) \perp (u_3 2^{\mu_3}) \perp (u_4 2^{\mu_4}).\n\end{cases}
$$

Here,

$$
\begin{cases} \n\text{in Case (I), } i \leq j; \\
\text{in Case (II), } u_i \equiv 1 \text{ mod } 2 \text{ and } \mu_1 \leq \mu_2; \\
\text{in Case (III), } u_i \equiv 1 \text{ mod } 2, \mu_i \leq \mu_j \text{ if } i < j, \mu_1 < \mu_3, \text{ and } \mu_2 < \mu_4. \n\end{cases}
$$

In the following theorem, we explain the Gross-Keating invariant of each case.

Theorem 5.6. The determinant of B is a square. The Gross-Keating invariant of B is described as follows:

 (1) In Case (I) , we have

$$
GK(B) = (i, i, j, j),
$$

where *i* and *j* have different parities.

(2) In case (II), μ_1 and μ_2 have the same parity, $u_1 + u_2 \equiv 0 \mod 4$, and

$$
GK(B) = (i, i) \cup (\mu_1, \mu_2 + 2).
$$

Here, i and μ_1 have different parities.

 (3) In case (III),

(a) Assume $\mu_1 \not\equiv \mu_2 \mod 2$. Then μ_3 and μ_4 have different parities and

$$
GK(B) = (\mu_1, \mu_2, \mu_3 + 2, \mu_4 + 2).
$$

(b) Assume that $\mu_1 \equiv \mu_2 \mod 2$ and that $u_1 + u_2 \equiv 2 \mod 4$ or $\mu_2 = \mu_3$. Then μ_3 and μ_4 have the same parity and

$$
GK(B) = (\mu_1, \mu_2 + 1, \mu_3 + 1, \mu_4 + 2).
$$

(c) Assume that $\mu_1 \equiv \mu_2 \mod 2$, $u_1 + u_2 \equiv 0 \mod 4$, and $\mu_3 \ge \mu_2 + 1$. Then

 $GK(B) = (\mu_1, \mu_2 + 2) \cup (\mu_3, \mu_4 + 2).$

Here, μ_3 and μ_4 have the same parity, μ_1 and μ_3 have different parities, and $u_3+u_4 \equiv 0$ mod 4.

Proof. The determinant of B is a square since the determinant of the matrix associated to O_D given at the beginning of this section is a square. Using this, the proof easily follows from Lemma 2.8 and Theorem 3.1 of [\[CIKY2\]](#page-50-22) (or Theorems 3.5-3.8 of [\[CIKY1\]](#page-50-23)) and Proposition [5.3.](#page-25-1)

In the next proposition, we compute the local density $\alpha(L, O_D)$ for an anisotropic quadratic lattice (L, q_L) of rank 4. The definition (and normalization) of the local density follows from Section 5 of [\[IK2\]](#page-50-10).

Proposition 5.7. The local density $\alpha(L, O_D)$ is

$$
\alpha(L, O_D) = [O_D : L] \cdot p^3 \cdot p^{-2} (2(p+1))^2.
$$

Here, $[O_D : L]$ equals to $(GK(L) - 2)/2$.

Proof. By Hilfssatz 17 of [\[Sie35\]](#page-51-7), we have

$$
\alpha(L, O_D) = [O_D : L] \cdot \alpha(O_D, O_D),
$$

where $[O_D: L]$ is the index of L in O_D . The local density $\alpha(O_D, O_D)$ for a single quadratic lattice O_D is fully studied in [\[Cho15\]](#page-50-18) $(p = 2)$ and [\[GY00\]](#page-50-13) $(p \neq 2)$.

By using the matrix description of the quadratic lattice (O_D, q_D) given at Remark 5.1.(2), the local density $\alpha(O_D, O_D)$ (cf. Proposition 6.2.3 and Theorem 7.3 of [\[GY00\]](#page-50-13) when $p \neq 2$, Theorems 4.12 and 5.2 of [\[Cho15\]](#page-50-18) when $p = 2$) is

$$
\alpha(O_D, O_D) = p^3 \cdot (p^{-1}2(p+1))^2.
$$

This completes the proof. \Box

Remark 5.8. In the above proof, indeed Theorems 4.12 and 5.2 of [\[Cho15\]](#page-50-18) when $p = 2$ yield that

$$
\alpha(O_D, O_D) = 1/2 \cdot p^{-3} \cdot p^{-6} \cdot p^4 (2(p+1))^2.
$$

But, using the normalization of the local density explained in Section 5 of [\[IK2\]](#page-50-10), we need to multiply $p⁶$ and to ignore $1/2$ in the above formula.

5.2. The Siegel series of anisotropic quadratic lattice. In this subsection, we will explain a more refined inductive formula of the Siegel series of anisotropic quadratic lattices over \mathbb{Z}_p . Let (L, q_L) be an anisotropic quadratic lattice over \mathbb{Z}_p of rank 4. By Remark 5.1.(3), we may consider it as a sublattice of (O_D, q_D) .

We will work with an exclusively chosen basis (e_1, e_2, e_3, e_4) of L as follows until the end of this subsection:

$$
(e_1, \dots, e_n)
$$
 is $\begin{cases} diagonal and optimal & if p is odd; \\ reduced & if p = 2. \end{cases}$

Let $L^{(4)}$ be the lattice spanned by $(e_1, e_2, e_3, 1/p \cdot e_4)$ and let $L_0^{(4)}$ $_0^{(4)}$ be the lattice spanned by (e_1, e_2, e_3) . If $a_3 < a_4$, then these notions are the same as those introduced at the beginning of Section [4](#page-17-0) and in Remark 4.3.(1). We keep using them even in the case of $a_3 = a_4$.

Lemma 5.9. Assume that (L, q_L) is not maximal. Then $L^{(4)}$ is contained in (O_D, q_D) .

Proof. It suffices to show that $L^{(4)}$ is a quadratic lattice. Let $GK(L) = (a_1, a_2, a_3, a_4)$. Since (L, q_L) is not maximal, $GK(O_D) = (0, 0, 1, 1)$, and only two of a_i 's have the same parity, we have that $a_4 \geq 2$. Using Remark [5.5](#page-26-0) when $p = 2$, we can see that $G(K(L^{(4)}) = (a_1, a_2, a_3) \cup (a_4 - 2) \succeq (0, 0, 0, 0)$. This holds when $p \neq 2$, which completes the proof by Lemma [3.4](#page-11-2) and Corollary [3.5.](#page-12-2)

Recall that $\mathcal{F}_L(X)$ is the Siegel series associated to the quadratic lattice (L, q_L) such that $\mathcal{F}_L(f^{-k}) = \alpha(L, H_k)$ (cf. Definition [3.15\)](#page-16-1). In the following theorem, we will explain an inductive formula of the Siegel series $\mathcal{F}_L(X)$ for an anisotropic quadratic lattice of rank 4 over \mathbb{Z}_2 . This formula is much simpler than that of Theorem [4.9.](#page-23-0)

Theorem 5.10. Assume that (L, q_L) is not maximal. Let $GK(L) = (a_1, a_2, a_3, a_4)$. Then we have the following inductive formula:

$$
\mathcal{F}_L(X) = p^5 \cdot X^2 \cdot \mathcal{F}_{L^{(4)}}(X) + (1-X)(1+pX) \cdot \mathcal{F}_{L_0^{(4)}}(pX).
$$

Here,

$$
\begin{cases} \text{GK}(L^{(4)}) = (a_1, a_2, a_3) \cup (a_4 - 2); \\ \text{GK}(L_0^{(4)}) = (a_1, a_2, a_3). \end{cases}
$$

Proof. If $a_3 < a_4$, then the formula follows from Theorem [4.9.](#page-23-0) Assume that $a_3 = a_4$. Since a_2 and a_3 should have different parities, we have that $a_2 < a_3$.

From our choice of (e_1, e_2, e_3, e_4) , the symmetric matrix with a basis $(e_1, e_2, e_3, \frac{1}{n})$ $\frac{1}{p} \cdot e_4$) (resp. (e_1, e_2, e_3)) is a reduced form whose associated Gross-Keating invariant is $(a_1, a_2, a_3) \cup (a_4 - 2)$ (resp. (a_1, a_2, a_3)) (cf. Proposition [5.4](#page-26-1) and Remark [5.5\)](#page-26-0). Using the argument explained in the proofs of Lemma [4.6](#page-20-0) and Proposition [4.8,](#page-21-0) we have the following formula:

$$
\#\mathcal{S}_{(L,a^\pm,b)}=\#\mathcal{S}_{(L^{(4)},a^\pm,b-1)}+f^{b-3b'}\cdot\sum_{x\in M_{1\times 3}(\mathfrak{o}/\pi^{b'}\mathfrak{o})}\#\mathcal{S}_{(L^{(4)}_x,a^\pm,b)}.
$$

Here b' is any integer with $b' \geq b$ and $L_x^{(4)}$ is a lattice of rank 3 constructed by using the same argument (with $n = 4$ and $d = 1$) given in Remark 4.3.(1) and Proposition [4.8.](#page-21-0)

Assume that $p = 2$. We consider a reduced form B with respect to a basis (e_1, e_2, e_3, e_4) of (L, q_L) . Then B satisfies the assumption of Lemma [4.12,](#page-24-2) by Proposition [5.4](#page-26-1) and Lemma [5.2.](#page-25-2) Thus, if we plug the above formula into the formula of Theorem [3.14](#page-16-0) using the result of Lemma [4.12,](#page-24-2) then we obtain the desired inductive formula.

We now assume that $p \neq 2$. We consider a diagonal matrix $B = (u_1 p^{a_1}) \perp (u_2 p^{a_2}) \perp (u_3 p^{a_3}) \perp$ $(u_4p^{a_4})$ with respect to a basis (e_1, e_2, e_3, e_4) , where $u_i \in \mathbb{Z}_p$ is a unit. Then for $x = (x_1, x_2, x_3) \in \mathbb{Z}_p^3$, we have

$$
B_x = (id_3 \t t_x) \cdot B \cdot \begin{pmatrix} id_3 \\ x \end{pmatrix} = \begin{pmatrix} u_1 p^{a_1} & 0 & 0 \\ 0 & u_2 p^{a_2} & 0 \\ 0 & 0 & u_3 p^{a_3} \end{pmatrix} + u_4 p^{a_4} \begin{pmatrix} x_1 x_1 & x_1 x_2 & x_1 x_3 \\ x_2 x_1 & x_2 x_2 & x_2 x_3 \\ x_3 x_1 & x_3 x_2 & x_3 x_3 \end{pmatrix}.
$$

If $\text{ord}((u_3 + u_4 x_3^2)p^{a_3}) = a_3$, where $(u_3 + u_4 x_3^2)p^{a_3}$ is the $(3, 3)$ -th entry of B_x , then the matrix B_x is a reduced form with $GK(B_x) = (a_1, a_2, a_3)$ by using Theorem 3.3 of [\[IK2\]](#page-50-10) for the 2×2 -minor involving $u_1p^{a_1}$ and $u_2p^{a_2}$ and the definition of a reduced form given in Definition [2.5.](#page-8-0)

As explained in Remark 4.3.(4), the Siegel series is completely determined by the Extended Gross-Keating datum. In our case of anisotropic quadratic lattices, the extended Gross-Keating datum is the same as the Gross-Keating invariant. Thus if $GK(B_x) = (a_1, a_2, a_3)$ for any x, then the associated Siegel series's are all equal. Using a similar argument used in the case $p = 2$, we have the desired formula.

Thus, it suffices to prove that $\text{ord}((u_3 + u_4x_3^2)p^{a_3}) = a_3$, equivalently that $u_3 + u_4x_3^2$ is a unit. For $a \in \mathbb{Z}_p$, let $\left(\frac{a}{p}\right)$ be the Legendre symbol. If $u_3 + u_4x_3^2$ is not a unit, then $\left(\frac{-u_3}{p}\right)$ $=\left(\frac{u_4x_3^2}{p}\right)$ $\Big)$ so that $\left(\frac{-u_3u_4}{p}\right)$ $=\left(\frac{u_4x_3}{p}\right)^2=1$. Since the lattice spanned by e_3 and e_4 is anisotropic, we have that $\left(\frac{-u_3u_4}{p}\right)$ $= -1$ by Lemma 2.8 of [\[Bou07\]](#page-50-21). This is a contradiction. Thus we conclude that $u_3 + u_4x_3^2$ is a unit. \square

The proof of the above theorem also holds for any anisotropic quadratic lattice of rank $n \leq 3$. We state it as the following theorem:

Theorem 5.11. In this theorem, let (L, q_L) be an anisotropic quadratic lattice over \mathbb{Z}_p of rank n. Let $GK(L) = (a_1, \dots, a_n)$. Assume that (L, q_L) is not maximal. Then we have the following inductive formula for $\mathcal{F}_L(X)$:

$$
\mathcal{F}_L(X) = \begin{cases} p^{n+1} \cdot X^2 \cdot \mathcal{F}_{L^{(n)}}(X) + (1-X)(1+pX) \cdot \mathcal{F}_{L_0^{(n)}}(pX) & \text{if } 2 \le n \le 4; \\ p^2 \cdot X^2 \cdot \mathcal{F}_{L^{(1)}}(X) + (1-X)(1+pX) & \text{if } n = 1. \end{cases}
$$

Here, $L^{(n)}$ is spanned by $(e_1, \dots, e_{n-1}, 1/p \cdot e_n)$ and $L_0^{(n)}$ $e_0^{(n)}$ is spanned by (e_1, \dots, e_{n-1}) so that

$$
\begin{cases} \text{GK}(L^{(n)}) = (a_1, \cdots, a_{n-1}) \cup (a_n - 2); \\ \text{GK}(L_0^{(n)}) = (a_1, \cdots, a_{n-1}). \end{cases}
$$

Remark 5.12. As mentioned in the proof of Theorem [5.10,](#page-28-0) for an anisotropic quadratic \mathbb{Z}_p -lattice, it is easy to see from Definition 6.3 of [\[IK1\]](#page-50-12) that the extended Gross-Keating datum is the same as the Gross-Keating invariant. Thus, the Siegel series for an anisotropic quadratic \mathbb{Z}_p -lattice is completely determined by the Gross-Keating invariant by Remark 4.3.(4).

Remark 5.13. Let $GK(L) = (a_1, \dots, a_n)$. Lemma [5.9](#page-28-1) and its proof imply that L of rank 4 is maximal if and only if $a_4 \leq 1$. We claim that this statement holds for any anisotropic quadratic lattice of any rank over \mathbb{Z}_p . This implies that if (L, q_L) is not maximal, then $L^{(n)}$ is integral in Theorem [5.11.](#page-29-1)

If the rank is 1, then it is obvious by Remark $2.8(4)-(5)$. If the rank is 3, then it follows from the fact that only two integers among a_i 's have the same parity. If the rank is 2, then it suffices to show that the quadratic lattice M with $GK(M) = (1, 1)$ is maximal.

Let (e_1, e_2, e_3, e_4) be a basis of (O_D, q_D) given in Remark 5.1.(2). Let M' be a sublattice of O_D spanned by (e_3, e_4) so that $GK(M') = (1, 1)$. If M' is not maximal, then we should be able to find a quadratic lattice \tilde{M} of rank 2 containing M' so that $GK(\tilde{M}) = (0,0)$. Thus, the lattice spanned by (e_1, e_2) and M is also integral whose Gross-Keating invariant is $(0, 0, 0, 0)$, which is a contraction to the fact that (O_D, q_D) is maximal. Thus M' is maximal.

Let's go back to the case M with $GK(M) = (1, 1)$. Thus the Siegel series $\mathcal{F}_M(X)$ should be the same as $\mathcal{F}_{M'}(X)$ by Remark [5.12.](#page-29-2) Assume that M is not maximal. Since M' is maximal, the local density associated to $\mathcal{F}_{M'}(X)$ is the same as the primitive local density for M', whereas the local density associated to $\mathcal{F}_M(X)$ should be the sum of the primitive local density for M and those for quadratic lattices including M . But, the prmitive local densities for M and M' are equal and the primitive local density for a bigger lattice including M is not trivial (cf. Theorem [3.12](#page-14-2) and Equation [\(3.8\)](#page-16-2)). Thus, $\mathcal{F}_M(X)$ cannot be the same as $\mathcal{F}_{M'}(X)$, which is a contradiction.

6. Comparison: the Siegel series and the local intersection multiplicities

Gross and Keating computed the local intersection multiplicities in [\[GK93\]](#page-50-0) and Kudla confirmed that it is the same as the derivative of the Siegel series of an anisotropic quadratic lattice of rank 3 at p^{-2} (cf.[\[ARGOS07\]](#page-50-2)). The method used to show the equality between these two objects is to

compute both sides independently, and then to compare them directly. The calculation of the local intersection multiplicities in [\[GK93\]](#page-50-0) is based on an inductive formula given in Lemma 5.6 in loc. cit.

In this section, we will compare the inductive formula of [\[GK93\]](#page-50-0) with our inductive formula of Theorem [5.11.](#page-29-1) Then we will show that these two are essentially equal, beyond matching values. In addition to that, we will explain a newly discovered equality between the local intersection multiplicity on the special fiber and the derivative of another Siegel series in Theorem [6.7.](#page-32-0) This observation had been missed in both of Siegel series and intersection numbers.

Let us restrict the following situation exclusively in this section:

 $\sqrt{ }$ $\frac{1}{2}$ \mathbf{I} L : anisotropic quadratic lattice over \mathbb{Z}_p of rank 3; M : anisotropic quadratic lattice over \mathbb{Z}_p of rank 2; N : anisotropic quadratic lattice over \mathbb{Z}_p of rank 1.

Since $\mathcal{F}_L(X)$ is determined by $GK(L) = (a_1, a_2, a_3)$ (cf. Remark [5.12\)](#page-29-2), we can write $\mathcal{F}_L(X) =$ $\mathcal{F}_{(a_1,a_2,a_3)}(X)$. Similarly, we can write $\mathcal{F}_M(X) = \mathcal{F}_{(a_1,a_2)}(X)$ with $GK(M) = (a_1,a_2)$ and $\mathcal{F}_N(X) =$ $\mathcal{F}_{(a_1)}(X)$ with $GK(N) = (a_1)$.

As in Section [5.2,](#page-27-0) a basis of each lattice, consisting of e_i 's, is chosen to be

In the following two lemmas, we list the initial values of the Siegel series and its derivative, in order to compare both inductive formulas.

Lemma 6.1. We have

$$
\mathcal{F}_L(1/p^2) = \mathcal{F}_M(1/p) = \mathcal{F}_N(1) = 0.
$$

Proof. These directly follow from Corollary [3.16.](#page-17-1) □

Lemma 6.2. Special values of the derivative of the Siegel series are given as follows:

$$
\mathcal{F}'_M(1/p) = \begin{cases}\n-(p-1) & \text{if } (a_1, a_2) = (0, 0); \\
-2(p-1)(p+1) & \text{if } (a_1, a_2) = (1, 1); \\
-2(p-1) & \text{if } (a_1, a_2) = (0, 1),\n\end{cases}
$$
\n
$$
\mathcal{F}'_N(1) = \begin{cases}\n-1 & \text{if } (a_1) = (0); \\
-(p+1) & \text{if } (a_1) = (1).\n\end{cases}
$$

Proof. We write $\mathcal{F}_{(a_1,a_2)}(X)$ for $\mathcal{F}_M(X)$ with $GK(M) = (a_1,a_2)$. In our situation, a_i is either 0 or 1. Thus M is a maximal quadratic lattice (cf. Remark [5.13\)](#page-29-3). Using Corollary [3.16,](#page-17-1) the Siegel series for a maximal quadratic lattice is described explicitly as follows:

$$
\left\{\begin{array}{l}\mathcal{F}_{(0,0)}(X)=(1-X)(1-pX);\\ \mathcal{F}_{(1,1)}(X)=(1-X)(1+p^2X)(1-p^2X^2);\\ \mathcal{F}_{(0,1)}(X)=(1-X)(1-p^2X^2).\end{array}\right.
$$

Thus

$$
\begin{cases}\n\mathcal{F}'_{(0,0)}(1/p) = -(p-1); \\
\mathcal{F}'_{(1,1)}(1/p) = -2(p-1)(p+1); \\
\mathcal{F}'_{(0,1)}(1/p) = -2(p-1).\n\end{cases}
$$

Similarly, we write $\mathcal{F}_{(a_1)}(X)$ for $\mathcal{F}_N(X)$ with $GK(N) = (a_1)$. Then

$$
\begin{cases}\n\mathcal{F}_{(0)}(X) = 1 - X; \\
\mathcal{F}_{(1)}(X) = (1 - X)(1 + pX).\n\end{cases}
$$

Thus

$$
\begin{cases}\n\mathcal{F}'_{(0)}(1) = -1; \\
\mathcal{F}'_{(1)}(1) = -(p+1).\n\end{cases}
$$

Let $GK(L) = (a_1, a_2, a_3)$ and let (e_1, e_2, e_3) be a chosen basis of L described at the beginning of this section. Let M be $L_0^{(3)}$ $_0^{(5)}$ spanned by (e_1, e_2) so that $GK(M) = (a_1, a_2)$. We have the following inductive formula:

Proposition 6.3. Let $\alpha(L, O_D)$ be the local density of the pair of quadratic lattices (L, q_L) and (O_D, q_D) . Assume that L is not maximal. Then we have the following inductive formula:

$$
\frac{\mathcal{F}'_L(1/p^2)}{\alpha(L, O_D)} = \frac{\mathcal{F}'_{L^{(3)}}(1/p^2)}{\alpha(L^{(3)}, O_D)} + \frac{p-1}{2p} \cdot \mathcal{F}'_M(1/p).
$$

Here, $L^{(3)}$ is a lattice spanned by $(e_1, e_2, \frac{1}{n})$ $rac{1}{p} \cdot e_3$.

Proof. By differentiating the formula of Theorem [5.11](#page-29-1) at p^{-2} using Lemma [6.1,](#page-30-0) we have

$$
\mathcal{F}'_L(1/p^2) = \mathcal{F}'_{L^{(3)}}(1/p^2) + (1 - \frac{1}{p^2})(p+1) \cdot \mathcal{F}'_M(1/p).
$$

On the other hand, for any anisotropic quadratic lattice L of rank 3, we have

$$
\alpha(L, O_D) = 2(p+1)^2 p^{-1}
$$

by Theorem 1.1 of [\[Wed07-1\]](#page-51-8). The desired inductive formula follows from these two. \Box

Let N be $M_0^{(2)}$ $_{0}^{(2)}$ spanned by (e_1) so that $GK(N) = (a_1)$.

Proposition 6.4. Assume that M is not maximal. Then we have the following inductive formulas:

$$
\left\{\begin{array}{l} \mathcal{F}'_M(1/p) = p \cdot \mathcal{F}'_{M^{(2)}}(1/p) + 2(p-1) \cdot \mathcal{F}'_N(1); \\ \mathcal{F}'_N(1) = p^2 \cdot \mathcal{F}'_{N^{(1)}}(1) - (p+1). \end{array}\right.
$$

Here, $M^{(2)}$ is a lattice spanned by $(e_1, \frac{1}{n})$ $\frac{1}{p} \cdot e_2$) and $N^{(1)}$ is a lattice spanned by $(\frac{1}{p})$ $(\frac{1}{p} \cdot e_1)$. In the second equation, we assume that N is not maximal.

Proof. The formulas directly follow by differentiating the formulas of Theorem [5.11](#page-29-1) using Lemma $6.1.$

We write $\mathcal{F}'_{(a_1)}(1) = \mathcal{F}'_N(1)$ with $GK(N) = (a_1)$. The above inductive formula yields the explicit value of $\mathcal{F}'_{(a_1)}(1)$ as follows.

Lemma 6.5. We have that

$$
\mathcal{F}'_{(a_1)}(1) = -(1+p+p^2+\cdots+p^{a_1}).
$$

Proof. Since $GK(N^{(1)}) = (a_1 - 2)$, by Proposition [6.4](#page-31-0) and Lemma [6.2,](#page-30-1) we have that

$$
\mathcal{F}'_{(a_1)}(1) = \begin{cases}\n-p^{a_1} - (p+1)(1+p^2 + \dots + p^{a_1-2}) & \text{if } a_1 \text{ is even;} \\
-(p+1)p^{a_1-1} - (p+1)(1+p^2 + \dots + p^{a_1-3}) & \text{if } a_1 \text{ is odd.}\n\end{cases}
$$

This completes the proof. \square

For an anisotropic lattice M with $GK(M) = (a_1, a_2)$, we define

(6.1)
$$
\mathcal{T}_{a_1,a_2} = \sum_{x=0}^{a_1} \sum_{y=0}^{a_2} p^{\min\{a_1 - x + y, a_2 - y + x\}}.
$$

 \Box

The number \mathcal{T}_{a_1,a_2} is indeed the local intersection multiplicity on the special fiber, defined by Equations (5.3) and (5.16) and Lemma 5.6 of [\[GK93\]](#page-50-0). In the following proposition, we will explain an inductive formula of \mathcal{T}_{a_1,a_2} , motivated by an inductive formula of $\mathcal{F}'_M(1/p)$ in Proposition [6.4](#page-31-0) as they are supposed to match each other. Later in Theorem [6.7,](#page-32-0) \mathcal{T}_{a_1,a_2} will be compared with the derivative of the Siegel series associated to the quadratic lattice M.

Proposition 6.6. If $a_2 \geq 2$, then

$$
\mathcal{T}_{a_1,a_2} = p \cdot \mathcal{T}_{a_1,a_2-2} - 2\mathcal{F}'_{(a_1)}(1).
$$

Proof. We write

$$
\begin{cases}\n\mathcal{T}_{a_1,a_2} = \sum_{x=0}^{a_1} \sum_{y=0}^{a_2} p^{\min\{a_1 - x + y, a_2 - y + x\}}; \\
p \cdot \mathcal{T}_{a_1,a_2-2} = \sum_{x=0}^{a_1} \sum_{y=0}^{a_2 - 2} p^{\min\{a_1 - x + (y+1), a_2 - (y+1) + x\}}.\n\end{cases}
$$

We rewrite the above sums as follows:

$$
\begin{cases}\n\mathcal{T}_{a_1,a_2} = \sum_{x=0}^{a_1} \sum_{y=0}^{x+\frac{a_2-a_1}{2}} p^{a_1-x+y} + \sum_{x=0}^{a_1} \sum_{x+\frac{a_2-a_1}{2} < y}^{a_2} p^{a_2-y+x}; \\
p \cdot \mathcal{T}_{a_1,a_2-2} = \sum_{x=0}^{a_1} \sum_{y=0}^{x+\frac{a_2-a_1}{2}-1} p^{a_1-x+(y+1)} + \sum_{x=0}^{a_1} \sum_{x+\frac{a_2-a_1}{2}-1 < y}^{a_2-2} p^{a_2-(y+1)+x}.\n\end{cases}
$$

Then we have

$$
\begin{cases} \sum_{x=0}^{a_1} \sum_{y=0}^{x+\frac{a_2-a_1}{2}} p^{a_1-x+y} - \sum_{x=0}^{a_1} \sum_{y=0}^{x+\frac{a_2-a_1}{2}-1} p^{a_1-x+(y+1)} = \sum_{x=0}^{a_1} p^{a_1-x};\\ \sum_{x=0}^{a_1} \sum_{x+\frac{a_2-a_1}{2} < y}^{a_2} p^{a_2-y+x} - \sum_{x=0}^{a_1} \sum_{x+\frac{a_2-a_1}{2}-1 < y}^{a_2-(y+1)+x} p^{a_2-(y+1)+x} = \sum_{x=0}^{a_1} p^{a_2-a_2+x}. \end{cases}
$$

Thus we have

$$
\mathcal{T}_{a_1,a_2} - p \cdot \mathcal{T}_{a_1,a_2-2} = 2 \sum_{x=0}^{a_1} p^x = -2 \mathcal{F}'_{(a_1)}(1).
$$

 \Box

We now compare the local intersection multiplicity \mathcal{T}_{a_1,a_2} on the special fiber, with the derivative of the Siegel series associated to the lattice M in the following theorem.

Theorem 6.7. We have the following equality:

$$
\mathcal{T}_{a_1,a_2} = \frac{-1}{p-1} \cdot \mathcal{F}'_M(1/p).
$$

In addition, both sides satisfy the same inductive formula.

Proof. By Propositions [6.4](#page-31-0) and [6.6,](#page-32-1) it suffices to prove that both sides have the same initial values. The initial values of \mathcal{T}_{a_1,a_2} can be computed directly from its definition as follows:

$$
\mathcal{T}_{a_1,a_2} = \begin{cases}\n1 & \text{if } (a_1, a_2) = (0,0); \\
2(p+1) & \text{if } (a_1, a_2) = (1,1); \\
2 & \text{if } (a_1, a_2) = (0,1),\n\end{cases}
$$

On the other hand, by Lemma [6.2,](#page-30-1) we have

$$
\mathcal{F}'_M(1/p) = \begin{cases}\n-(p-1) & \text{if } (a_1, a_2) = (0, 0); \\
-2(p-1)(p+1) & \text{if } (a_1, a_2) = (1, 1); \\
-2(p-1) & \text{if } (a_1, a_2) = (0, 1),\n\end{cases}
$$

Therefore, both \mathcal{T}_{a_1,a_2} and $\frac{-1}{p-1} \cdot \mathcal{F}'_M(1/p)$ have the same initial values.

For an anisotropic quadratic \mathbb{Z}_p -lattice L of rank 3 with $GK(L) = (a_1, a_2, a_3)$, put $\alpha_p(a_1, a_2, a_3) :=$ $\alpha_p(L)$ which is the local intersection multiplicity defined in Equation (5.3) of [\[GK93\]](#page-50-0). We finally compare it with the derivative of the Siegel series associated to the quadratic lattice L in the following theorem.

Theorem 6.8. Let $GK(L) = (a_1, a_2, a_3)$. Then we have the following equality:

$$
\alpha_p(a_1, a_2, a_3) = \frac{-2p}{(p-1)^2} \cdot \frac{\mathcal{F}'_L(1/p^2)}{\alpha(L, O_D)}.
$$

Moreover, both sides satisfy the same inductive formula.

Proof. Let $\widetilde{\mathcal{T}}_{a_1,a_2,a_3} = \frac{-2p}{(p-1)^2}$. $\mathcal{F}_{L}'(1/p^2)$ $\frac{\partial L(1/P)}{\partial L(0,D)}$. Then by Theorem [6.7,](#page-32-0) the formula of Proposition [6.3](#page-31-1) turns to be

$$
\widetilde{\mathcal{T}}_{a_1,a_2,a_3}=\widetilde{\mathcal{T}}_{a_1,a_2,a_3-2}+\mathcal{T}_{a_1,a_2}.
$$

On the other hand, by Lemma 5.6 and Equations (5.16) and (5.18) of [\[GK93\]](#page-50-0), the local intersection multiplicity $\alpha_p(a_1, a_2, a_3)$ satisfies the following inductive formula:

$$
\alpha_p(a_1, a_2, a_3) = \alpha_p(a_1, a_2, a_3 - 2) + \mathcal{T}_{a_1, a_2}.
$$

Therefore, it suffices to show that both $\mathcal{T}_{a_1,a_2,a_3}$ and $\alpha_p(a_1,a_2,a_3)$ have the same initial values.

We write $\mathcal{F}_{(a_1,a_2,a_3)}(X)$ for $\mathcal{F}_L(X)$. If a_i consists of either 0 or 1, then the associated quadratic lattice L is maximal by Remark [5.13.](#page-29-3) Using Corollary [3.16,](#page-17-1) the Siegel series for a maximal quadratic lattice is described explicitly as follows:

$$
\begin{cases}\n\mathcal{F}_{(0,0,1)}(X) = (1-X)(1-p^2X^2)(1-p^2X); \\
\mathcal{F}_{(0,1,1)}(X) = (1-X)(1-p^2X^2)(1-p^4X^2).\n\end{cases}
$$

Thus

$$
\left\{\begin{array}{l} \mathcal{F}'_{(0,0,1)}(\frac{1}{p^2})=-p^2(1-\frac{1}{p^2})^2,\ \frac{-2p}{(p-1)^2}\cdot\frac{\mathcal{F}'_{(0,0,1)}(\frac{1}{p^2})}{\alpha(L,O_D)}=1; \\ \mathcal{F}'_{(0,1,1)}(\frac{1}{p^2})=-2p^2(1-\frac{1}{p^2})^2,\ \frac{-2p}{(p-1)^2}\cdot\frac{\mathcal{F}'_{(0,1,1)}(\frac{1}{p^2})}{\alpha(L,O_D)}=2. \end{array}\right.
$$

Therefore, both sides have the same initial values by Proposition 1.6 of $[Rap07]$.

7. Application 1: The intersection number over a finite field

In this section we revisit the results of Gross-Keating [\[GK93\]](#page-50-0) and give a new identity between certain intersection numbers of cycles over a finite field and the sum of the Fourier coefficients of the Siegel-Eisenstein series for Sp_4/\mathbb{Q} of weight 2. We follow the notation and results in Chapters 3-5 of [\[ARGOS07\]](#page-50-2) (cf. [Gör07-1], [Gör07-2], [\[Wed07-2\]](#page-51-10)).

7.1. Main results. For a positive integer m , we denote by T_m the modular correspondence of degree m defined in [Gör07-2]. It can be regarded as a flat scheme T_m over Z which is explicitly given by $T_m = \text{Spec } \mathbb{Z}[x, y]/(\varphi_m) \subset \text{Spec } \mathbb{Z}[x, y] =: S$, where we think the latter scheme S as the product of two copies of the coarse moduli space $Y_0(1)$ of elliptic curves. Here φ_m is the modular polynomial of degree m (see [\[Vog07\]](#page-51-0) and [Gör07-2]). We consider $T_{m,p} := T_m \otimes \text{Spec } \overline{\mathbb{F}}_p$ and $T_{m,\mathbb{C}} := T_m \otimes \text{Spec } \mathbb{C}$. Two cycles $T_{m_1,\mathbb{C}}$ and $T_{m_2,\mathbb{C}}$ intersect properly if and only if m_1m_2 is not a square (cf. Proposition 2.4 of $[GR93]$). The associated intersection number over $\mathbb C$ is defined by

(7.1)
$$
(T_{m_1,\mathbb{C}}, T_{m_2,\mathbb{C}}) := \text{length}_{\mathbb{C}} \mathbb{C}[x, y]/(\varphi_{m_1}, \varphi_{m_2}).
$$

We first state the following proposition to explain exactly when two cycles over a finite field intersect properly. It turns out to be the same as the situation over C.

Proposition 7.1. For given two positive integers m_1 and m_2 , the cycles $T_{m_1,p}$ and $T_{m_2,p}$ intersect properly in $S \otimes \text{Spec } \overline{\mathbb{F}}_p$ if and only if m_1m_2 is not a square (equivalently, $T_{m_1,\mathbb{C}}$ and $T_{m_2,\mathbb{C}}$ intersect properly inside $S \otimes \text{Spec } \mathbb{C}$ by Proposition 2.4 of $[\text{GK93}].$

Proof. Assume that $T_{m_1,p}$ and $T_{m_2,p}$ intersect properly. If m_1m_2 is square, then we can write $m_i = d_i^2 n$ for some positive integers d_1, d_2, n . By the description of the modular polynomial in p.2 of [\[Vog07\]](#page-51-0) we see that the polynomial ψ_n (see loc.cit. for the notation of ψ_n) is a common divisor of φ_{m_1} and φ_{m_2} as an element of $\mathbb{Z}[X, Y]$. Since $p \nmid \psi(X, Y)$, it defines a curve defined over \mathbb{F}_p . This gives a contradiction with the assumption.

Next assume that m_1m_2 is not a square. If $T_{m_1,p}$ and $T_{m_2,p}$ do not intersect properly, then they share an irreducible common divisor D . Let K be the function field of D which is of transcendental degree one over $\overline{\mathbb{F}}_p$. It follows that there exists a geometric point $x = \text{Spec } \overline{K} = ((E, E'), f_1, f_2)$ with two endomorphisms f_1, f_2 of degree m_1, m_2 between elliptic curves E, E' over K. By the proof of Proposition [7.5](#page-35-2) below $j(E), j(E')$ belong to $K \setminus \overline{\mathbb{F}}_p$. Further it holds that $\text{Hom}(E, E') \otimes_{\mathbb{Z}} \mathbb{Q} \simeq$ End(E) $\otimes_{\mathbb{Z}} \mathbb{Q} = \mathbb{Q}$ by Theorem 12 of [\[CL\]](#page-50-26). Combining this with the Z-freeness of Hom(E, E') (cf. Chapter III, Proposition 4.2-(b) in p.68 of [\[Sil09\]](#page-51-11)) we see that $Hom(E, E') = \mathbb{Z}$. Since f_1, f_2 belong to Hom $(E, E') = \mathbb{Z}$ there exist two (non-zero) integers n_1, n_2 such that $n_1 f_1 = n_2 f_2$. It implies that $n_1^2m_1 = n_2^2m_2$ and also m_1m_2 has to be square. However this yields a contradiction.

Corollary 7.2. If m_1m_2 is not a square, then φ_{m_1} and φ_{m_2} make up a regular sequence in $\mathbb{F}_p[x, y]$ and also in $\bar{\mathbb{F}}_p[[x,y]].$

Proof. It suffices to prove that φ_{m_1} is not a zero divisor in $A := \mathbb{F}_p[x, y]/(\varphi_{m_2})$. Suppose the contrary. Then there exists $\alpha \in \mathbb{F}_p[x, y]$ which is not divided by φ_{m_2} such that φ_{m_2} divides $\alpha \varphi_{m_1}$. Since $\mathbb{F}_p[x,y]$ is UFD, there exists an irreducible common factor h of φ_{m_2} and φ_{m_1} . Proposition [7.1](#page-34-1) implies that $\mathbb{F}_p[x,y]/(\varphi_{m_1}, \varphi_{m_2})$ is Artinian but this gives a contradiction with the existence of h. The same argument works for $\mathbb{F}_p[[x,y]]$. $p[[x, y]].$

Let us take two positive integers m_1 and m_2 such that m_1m_2 is not a square. We write $(T_{m_1,p}, T_{m_2,p})$, the intersection number over a finite field, which is explicitly defined as follows:

(7.2)
$$
(T_{m_1,p}, T_{m_2,p}) := \text{length}_{\mathbb{F}_p} \mathbb{F}_p[x, y]/(\varphi_{m_1}, \varphi_{m_2}).
$$

Our goal is to compute $(T_{m_1,p}, T_{m_2,p})$ explicitly.

Theorem 7.3. Assume that p is odd. Then for any two positive integers m_1 and m_2 such that m_1m_2 is not a square, the intersection number $(T_{m_1,p}, T_{m_2,p})$ is independent of the choice of a prime number p and its explicit value is given as follows:

$$
(T_{m_1,p}, T_{m_2,p}) = \frac{1}{288} \sum_{\substack{T \in \text{Sym}_2(\mathbb{Z}) > 0 \\ \text{diag}(T) = (m_1, m_2)}} c(T) = (T_{m_1,\mathbb{C}}, T_{m_2,\mathbb{C}}).
$$

Here, $c(T)$ is the Fourier coefficient of the Siegel-Eisenstein series for $Sp_4(\mathbb{Z})$ of weight 2 with respect to the (2×2) - half-integral symmetric matrix T (cf. [\[Nag92\]](#page-51-12)).

If we reinterpret Theorem [7.3](#page-34-2) in terms of modular polynomials in Equation [\(7.2\)](#page-34-3), then the $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}$]-module $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}$ [x, y]/($\varphi_{m_1}, \varphi_{m_2}$) satisfies the following interesting properties.

Theorem 7.4. We have the following interpretation about $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}][x,y]/(\varphi_{m_1},\varphi_{m_2})$:

- (1) $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}][x,y]/(\varphi_{m_1},\varphi_{m_2})$ is a free $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}$]-module.
- (2) The rank of $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}][x,y]/(\varphi_{m_1}, \varphi_{m_2})$, as a $\mathbb{Z}[\frac{1}{2}]$ $\frac{1}{2}$]-module, is equal to

$$
\frac{1}{288} \sum_{\substack{T \in \text{Sym}_2(\mathbb{Z})_{>0} \\ \text{diag}(T)=(m_1, m_2)}} c(T).
$$

Here, $c(T)$ is as described in the above theorem.

7.2. Decomposition of the intersection number over a finite field. In what follows let us go into the proof of Theorem [7.3.](#page-34-2) Let m_1, m_2 be positive integers such that m_1m_2 is not a square. We denote by $\text{CLN}_{\overline{\mathbb{F}}_p}$ (resp. $\text{CLN}_{W(\overline{\mathbb{F}}_p)}$) the category of complete local Noetherian \mathbb{F}_p (resp. $W(\overline{\mathbb{F}}_p)$)-algebras with the residue field $\overline{\mathbb{F}}_p$. The local deformation functor for a pair of elliptic curves $x := (E, E')$ over $\overline{\mathbb{F}}_p$ on $\text{CLN}_{\overline{\mathbb{F}}_p}$ is pro-represented by $R_p := \overline{\mathbb{F}}_p[[t, t']]$, which is called the universal deformation ring of x on $\text{CLN}_{\overline{\mathbb{F}}_p}$. Similarly we have the universal deformation ring $R := W(\overline{\mathbb{F}}_p)[[t, t']]$ of x on $\text{CLN}_{W(\overline{\mathbb{F}}_p)}$. Let us clarify the relation between the completion $\widehat{\mathcal{O}}_{S_{\overline{\mathbb{F}}_p}, x}$ of the structure sheaf $\mathcal{O}_{S_{\overline{\mathbb{F}}_p}}$ at x and R_p . Note that $\widehat{\mathcal{O}}_{S_{\overline{\mathbb{F}}_p},x} \simeq \overline{\mathbb{F}}_p[[j-j(E),j'-j(E')]]$. Here, $j(E)$ and $j(E')$ are the *j*-invariants of E and E', respectively. Since $Aut(x) = Aut(E) \times Aut(E')$ acts naturally on the deformation datum (cf. (8.2) of [\[KM85\]](#page-50-27)) we have that $\widehat{\mathcal{O}}_{S_{\overline{\mathbb{F}}_p},x} \simeq R_p^{\text{Aut}(x)} \subset R_p$ and that R_p is a free $\widehat{\mathcal{O}}_{S_{\overline{\mathbb{F}}_p},x}$ -module of rank $\frac{\sharp \text{Aut}(x)}{4}$ (cf. page 21 of [Gör07-2]). It is better to work on R_p instead of $\mathcal{O}_{S_{\overline{\mathbb{F}}_p},x}$ because the moduli space of elliptic curves is not a fine moduli space. The difference between these two objects in the computation of local intersection multiplicity is understood as below. We first consider the decomposition of the modular polynomial φ_m over R_p in terms of the local deformation theory.

Proposition 7.5. For a positive integer m, let (φ_m) be the ideal of R_p generated by the modular polynomial φ_m . Then

$$
(\varphi_m) = \prod_{\substack{f:E \longrightarrow E' \text{ isogeny of} \\ \text{degree } m, \text{ mod} \pm 1}} I_{m,f,p},
$$

where $I_{m,f,p} = (\varphi_{m,f,p})$ with $\varphi_{m,f,p} \in R_p$ is the minimal ideal of R_p such that f lifts to an isogeny over $R_p/I_{m,f,p}$.

Proof. We imitate the proof of Lemma 4.1 of [Gör07-2] for R_p . Let Def_{f,*} for $*\in {\mathbb{F}_p, W(\mathbb{F}_p)}$ be the deformation functor on CLN_* for f. As is similar to the proof of Lemma 4.1 of [Gör07-2], the deformation functor $\mathrm{Def}_{f,\overline{\mathbb{F}}_p}$ is represented by the divisor $\mathrm{div}(\varphi_{f,p})$ in $\mathrm{Spf} R_p$ for some $\varphi_{f,p} \in R_p$. For two isogenies $f, g : E \longrightarrow E'$ of degree m, $\varphi_{f,p}$ and $\varphi_{g,p}$ are coprime unless $f = \pm g$. This is proved as follows. Assume that $\varphi_{f,p}$ and $\varphi_{g,p}$ has a common irreducible factor h. Let \mathbb{E}, \mathbb{E}' be the universal deformations of E, E' over R_p . Since $R_p/(h)$ is of dimension one, the *j*-invariants $j(\mathbb{E} \otimes \mathrm{Spf} R_p/(h)), j(\mathbb{E}' \otimes \mathrm{Spf} R_p/(h))$ are not constant, hence they do not belong to $\overline{\mathbb{F}}_p$. It follows from Theorem 12 of [\[CL\]](#page-50-26) that $\text{End}(\mathbb{E} \otimes \text{Spf}R_p/(h)) = \text{End}(\mathbb{E}' \otimes \text{Spf}R_p/(h)) = \mathbb{Z}$. This implies that

$$
\mathrm{Hom}(\mathbb{E}\otimes \mathrm{Spf}R_p/(h),\mathbb{E}'\otimes \mathrm{Spf}R_p/(h))=\mathbb{Z}.
$$

Since f, g are liftable to isogenies $\tilde{f}, \tilde{g} \in \text{Hom}(\mathbb{E} \otimes \text{Spf}R_p/(h), \mathbb{E}' \otimes \text{Spf}R_p/(h))$ of the same degree, \tilde{f} has to be $\pm \tilde{g}$. Reduction to $\overline{\mathbb{F}}_p$ we have $f = \pm g$.

By Lemma 4.1 of [Gör07-2] which is in the situation with R, we also decompose the ideal φ_mR of R generated by φ_m as follows:

$$
\varphi_m R = \prod_{\substack{f:E \longrightarrow E' \text{ iso. of} \\ \text{degree } m, \text{ mod}\pm 1}} I_{m,f},
$$

where $I_{m,f}$ is the minimal ideal of R such that f lifts to an isogeny over $R/I_{m,f}$. The ideal $I_{m,f}$ is generated by a single element $\varphi_{m,f}$ in R which cannot be divisible by p.

Let $\varphi_{m,f,p}$ be the image of $\varphi_{m,f}$ under the natural projection $R \longrightarrow R_p$. Then we have

$$
\operatorname{div}(\varphi_{f,p})(S) = \operatorname{Def}_{f,\overline{\mathbb{F}}_p}(S) = \operatorname{Def}_{f,W(\overline{\mathbb{F}}_p)}(S) = \operatorname{div}(\varphi_{m,f})(S) = \operatorname{div}(\varphi_{m,f,p})(S)
$$

for any formal scheme S over Spf R_p . Hence we have $I_{m,f,p} = (\varphi_{f,p}) = (\varphi_{m,f,p})$.

Lemma 7.6. For positive integers m_1, m_2 with $m_1 m_2$ non-square and a pair (E, E') of elliptic curves over a finite field, it holds that

$$
\operatorname{length}_{\overline{\mathbb{F}}_p} \overline{\mathbb{F}}_p[[j,j']]_{(j-j(E),j'-j(E'))}/(\varphi_{m_1}, \varphi_{m_2}) =
$$

$$
\sum_{f_1} \sum_{f_2} \frac{4}{\sharp \operatorname{Aut}(E) \sharp \operatorname{Aut}(E')} \operatorname{length}_{\overline{\mathbb{F}}_p} \overline{\mathbb{F}}_p[[t,t']] / (\varphi_{m_1, f_1, p}, \varphi_{m_2, f_2, p})
$$

where $\varphi_{m_i,f_i,p}$ is a factor of φ_{m_i} given in the previous proposition. Here, the sums are over isogenies $f_i: E \to E'$ of degree m_i up to ± 1 .

Proof. By Corollary [7.2,](#page-34-4) φ_{m_1} and φ_{m_2} make up a regular sequence. Using a similar argument of Equation (4.1) in page 23 of [Gör07-2], the claim follows from Lemma 4.2 of [Gör07-2], Proposition [7.5,](#page-35-2) and the fact that R_p is a free $\mathcal{O}_{S_{\overline{\mathbb{F}}_p}}$ -module of rank $\frac{\sharp \text{Aut}(E)\sharp \text{Aut}(E')}{4}$ 4 . — Первый профессиональный профессиональный профессиональный профессиональный профессиональный профессиональн
В собстановки профессиональный профессиональный профессиональный профессиональный профессиональный профессиона

We denote by $(\mathbb{E}, \mathbb{E}')$ the universal pair of elliptic curves over R_p . Let $y = ((E, E'), f_1, f_2)$ be a pair of (E, E') and two isogenies $f_1, f_2 : E \longrightarrow E'$ with $deg(f_i) = m_i, i = 1, 2$. We define I_y as the minimal ideal of R_p such that both f_i 's lift to isogenies $\mathbb{E} \longrightarrow \mathbb{E}'$ of degree m_i 's modulo I_y for $i = 1, 2$, respectively. Put

$$
\mathrm{IM}_{p,y} := \mathrm{length}_{\overline{\mathbb{F}}_p} R_p/I_y,
$$

which is exactly the same as the local contribution in the summation of Lemma [7.6.](#page-36-0)

From now on, for a pair of two elliptic curves (E, E') defined over $\overline{\mathbb{F}}_p$, we use the following notation:

$$
\begin{cases}\n(E, E') : (ord) & \text{if both } E \text{ and } E' \text{ are ordinary;} \\
(E, E') : (ss) & \text{if both } E \text{ and } E' \text{ are supersingular.}\n\end{cases}
$$

The intersection number $(T_{m_1,p}, T_{m_2,p})$ over a finite field is described as follows:

Proposition 7.7. Assume that m_1m_2 is non-square. Then we have

(7.3)
$$
(T_{m_1,p}, T_{m_2,p}) = \sum_{\substack{y = ((E,E'),f_1,f_2) \ (E,F') : (\text{ord})}} \frac{\text{IM}_{p,y}}{\sharp \text{Aut}(E) \sharp \text{Aut}(E')} + \sum_{\substack{y = ((E,E'),f_1,f_2) \ (E,F') : (\text{ss})}} \frac{\text{IM}_{p,y}}{\sharp \text{Aut}(E) \sharp \text{Aut}(E')}.
$$

Proof. LHS is decomposed as follows:

$$
(T_{m_1,p}, T_{m_2,p}) = \sum_{(E,E')} (T_{m_1,p}, T_{m_2,p})_{(j(E,j)(E'))}
$$

$$
= \sum_{(E,E'): (ord)} (T_{m_1,p}, T_{m_2,p})_{(j(E,j)(E'))} + \sum_{(E,E'): (ss)} (T_{m_1,p}, T_{m_2,p})_{(j(E,j)(E'))}.
$$

Using Lemma [7.6,](#page-36-0) we have that

(7.5)
$$
(T_{m_1,p}, T_{m_2,p})_{(j(E,j)(E'))} = \frac{4}{\sharp \text{Aut}(E) \sharp \text{Aut}(E')} \sum_{\substack{f_i: E \longrightarrow E' \text{ iso. of degree } m_i, \text{ mod } \pm 1 \\ \text{degree } m_i, \text{ mod } \pm 1}} \text{IM}_{p,((E,E'),f_1,f_2)}
$$

$$
= \sum_{\substack{f_i: E \longrightarrow E' \text{ iso.} \\ \text{of degree } m_i}} \frac{\text{IM}_{p,((E,E'),f_1,f_2)}}{\sharp \text{Aut}(E) \sharp \text{Aut}(E')}.
$$

This completes the claim.

7.3. The local intersection multiplicity over a finite field. Based on Proposition [7.7,](#page-36-1) we compute $\mathrm{IM}_{p,y}$ by using Serre-Tate theory for the ordinary case and [\[ARGOS07\]](#page-50-2) for the supersingular case. Let us start with a series of the following lemmas.

Lemma 7.8. Let m_1, m_2 be two integers with $m_1 m_2$ non-square and let E, E' be two elliptic curves over a finite field. Then two isogenies $f_1, f_2 : E \longrightarrow E'$ with $\deg(f_i) = m_i$ are linearly independent in the Z-module Hom (E, E') and also in $\text{Hom}(E, E') \otimes_{\mathbb{Z}} \mathbb{Z}_p$.

Proof. Assume that f_1 and f_2 are linearly dependent. Then $n_1f_1 = n_2f_2$ for some integers n_1, n_2 . We may assume that n_1 and n_2 are coprime since $Hom(E, E')$ is a free Z-module (III, Proposition 4.2 of [\[Sil09\]](#page-51-11)). By comparing the degree, we have $n_1^2m_1 = n_2^2m_2$. Observe $(n_1m_1)^2 = n_2^2m_1m_2$. This implies that m_1m_2 is a square, which is a contradiction. For the latter since \mathbb{Z}_p is a Z-flat module, Z-freeness of $Hom(E, E')$ implies that $Hom(E, E') \otimes_{\mathbb{Z}} \mathbb{Z}_p$ is a free \mathbb{Z}_p -module. The degree map on $\text{Hom}(E, E')$ is extended \mathbb{Z}_p -linearly on $\text{Hom}(E, E') \otimes_{\mathbb{Z}} \mathbb{Z}_p$. Then the same argument above works and this completes the proof.

For a *p*-adic integer $z = \sum_{n=0}^{\infty}$ $i=0$ $a_i p^i$ with $a_i \in \{0, ..., p-1\}$ and the indeterminant t, we abusively

define

$$
(1+t)^{z} := \sum_{n=0}^{\infty} \binom{z}{n} t^{n} = \prod_{i=0}^{\infty} (1+t^{p^{i}})^{a_{i}} \text{ modulo } p
$$

as an element in $\mathbb{F}_p[[t]]$, rather than $\mathbb{Z}_p[[t]]$. Here $\begin{pmatrix} z \\ z \end{pmatrix}$ n $\big) := \frac{z(z-1)\cdots(z-n+1)}{1}$ $\frac{(z-n+1)}{n!}$ and set $\begin{pmatrix} z \\ 0 \end{pmatrix}$ $\overline{0}$ $):=$ 1. It follows from the argument in p. 288 of $[Ro]$ (see "THE PROOF OF LEMMA") that $(1 +$ $(t)^{z}(1+t)^{z'} = (1+t)^{z+z'}$ and $((1+t)^{z})^{z'} = (1+t)^{zz'}$ for any $z, z' \in \mathbb{Z}_p$ and these facts yields the above second equality for $(1 + t)^z$.

Lemma 7.9. For $z, w \in \mathbb{Z}_p^{\times}$ and $r \in \mathbb{Z}_{\geq 0}$ define an element $H(t,t') = (1+t)^{zp^r} - (1+t')^{w^r}$ in $\mathbb{F}_p[[t,t']]$. Then there exists an element $f(t)$ in $\mathbb{F}_p[[t]]$ such that $H(t, f(t)) = 0$ and $f(t) \equiv$ $zw^{-1}t^{p^r} \mod(t^{p^r+1}).$

Proof. Apply IV, Lemma 1.2 (Hensel's lemma) of [\[Sil09\]](#page-51-11) with $R = \mathbb{F}_p[[t]]$ (here R is the notation there), $I = (t)$, $F(t') = H(t, t')$, $a = zw^{-1}t^{p^r}$, and $\alpha = -w$. Notice that $F(a) \in I^{p^r+1}$ and $F'(a) = -w(1 + zw^{-1}t^{p^r})^{w-1} \in R^\times = \mathbb{F}_p[[t]]^\times.$

Lemma 7.10. Let e_1, e_2 be two non-negative integers. For any two elements $f, g \in R_p = \overline{\mathbb{F}}_p[[t, t']]$ which are coprime, it holds that

$$
\text{length}_{\overline{\mathbb{F}}_p} R_p/(f^{p^{e_1}}, g^{p^{e_2}}) = p^{e_1+e_2} \cdot \text{length}_{\overline{\mathbb{F}}_p} R_p/(f, g).
$$

Proof. It follows from Lemma 4.2 of [Gör07-2]. \Box

For an isogeny $f: E \longrightarrow E'$ between ordinary elliptic curves over a finite field k, by functoriality we can associate a unique element of

Hom
$$
(\widehat{E}, \widehat{E}') \times \text{Hom}_{\mathbb{Z}_p}(T_p(E) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p/\mathbb{Z}_p, T_p(E') \otimes_{\mathbb{Z}_p} \mathbb{Q}_p/\mathbb{Z}_p) = \text{End}(\widehat{\mathbb{G}}_m/k) \times \text{End}_{\mathbb{Z}_p}(\mathbb{Q}_p/\mathbb{Z}_p) \simeq \mathbb{Z}_p^2
$$

where $\widehat{E}, \widehat{E}'$ are formal groups associated to E, E' respectively and $\widehat{\mathbb{G}}_m$ is the multiplicative formal

group over k. We write $(z(f), w(f)) \in \mathbb{Z}_p^2$ for the element corresponding to f via the above identification.

Proposition 7.11. Let $y = ((E, E'), f_1, f_2)$ as explained in Lemma [7.8.](#page-37-1) Let T be the half-integral symmetric matrix associated to the quadratic lattice spanned by (f_1, f_2) so that $diag(T) = (m_1, m_2)$. Then

- (1) (ord) if (E, E') is a pair of ordinary elliptic curves, then $\text{ord}_p(\text{det}(2T))$ is an even integer and $\mathrm{IM}_{p,y} = p^{r_T}$, where $r_T := \frac{1}{2} \mathrm{ord}_p(\det(2T))$;
- (2) (ss) if (E, E') is a pair of supersingular elliptic curves, then $IM_{p,y} = \mathcal{T}_{a_1,a_2}$. Here, (a_1, a_2) = $GK(T \otimes \mathbb{Z}_p)$. For the definition of \mathcal{T}_{a_1,a_2} , see Equation [\(6](#page-31-2).1).

Proof. The second claim follows from Equation (4.1) , p. 151 of $[Rap07]$. Assume the first case. As in Equation (2.1) of [Gör07-1] there exists $d \in \mathbb{Z}$ such that $d^2d(E) = \det(2T)$, where $d(E)$ is the discriminant of End(E). Since E is ordinary, we see that $p \nmid d(E)$ by Theorem 12, p.182 of [\[La87\]](#page-50-28). We express the matrix T in terms of $(z_i, w_i) := (z(f_i), w(f_i)), i = 1, 2$ as follows:

$$
(7.6) T = \begin{pmatrix} \deg(f_1) & \frac{1}{2} (\deg(f_1 + f_2) - \deg(f_1) - \deg(f_2)) \\ \frac{1}{2} (\deg(f_1 + f_2) - \deg(f_1) - \deg(f_2)) & \deg(f_2) \end{pmatrix}
$$

=
$$
\begin{pmatrix} z_1w_1 & \frac{1}{2} (z_1w_2 + z_2w_1) \\ \frac{1}{2} (z_1w_2 + z_2w_1) & z_2w_2 \end{pmatrix}.
$$

Then we have

$$
\mathrm{ord}_p(\det(2T)) = 2\mathrm{ord}_p(z_1w_2 - z_2w_1).
$$

It follows from Serre-Tate theory, Theorem 2.1-3), p. 148 of [\[Katz81\]](#page-50-29) (see also the observation in p. 140 of [\[Rap07\]](#page-51-9)) that the minimal ideal I_y is given by $I_y = (H_1, H_2)$, where

$$
H_i = H_i(t, t') := (1 + t)^{z_i} - (1 + t')^{w_i}, \ i = 1, 2.
$$

By Lemma [7.10](#page-37-2) and changing z_i and w_i if necessary, we may assume that $w_1 \in \mathbb{Z}_p^{\times}$. Remark that $H_{1,a} := (1+t)^{az_1} - (1+t')^{aw_1} \equiv 0 \mod H_1$ in R_p for any $a \in \mathbb{Z}_p$. By replacing (f_1, f_2) with $(f_1, w_1 f_2)$ (here we view them as a pair of two homomorphisms between \widehat{E} and \widehat{E}') $R_p/(H_1, H_2) \simeq$ $R_p/((1+t)^{z_2w_1} - (1+t')^{w_2w_1}, H_1)$. Then we have that

$$
R_p/(H_1, H_2) \simeq R_p/((1+t)^{z_2w_1} - (1+t')^{w_2w_1}, H_1, H_{1,w_2}) = R_p/((1+t)^{z_1w_2} - (1+t)^{z_2w_1}, H_1).
$$

Lemma [7.9](#page-37-3) yields an existence of $f(t) \in \mathbb{F}_p[[t]]$ such that $H_1(t, f(t)) = 0$. This implies that

$$
R_p/(H_1, H_2) = \overline{\mathbb{F}}_p[[t]]/((1+t)^{z_1w_2} - (1+t)^{z_2w_1}).
$$

Write $z_1w_2 - z_2w_1 = p^{r_T}\alpha$ for $\alpha \in \mathbb{Z}_p^{\times}$. Then $R_p/(H_1, H_2) = \overline{\mathbb{F}}_p[[t]]/(t^{p^{r_T}})$. Hence we have the claim. \Box

7.4. Comparison in the ordinary locus using quasi-canonical lifts: over \mathbb{C} and over $\overline{\mathbb{F}}_p$. Let us recall the canonical lift and quasi-canonical lifts of an ordinary elliptic curve E over $\overline{\mathbb{F}}_p$. BySerre-Tate's theorem ([\[Katz81\]](#page-50-29)), there exists a unique canonical lift $\widetilde{E}^{\text{can}}$ to $W = W(\overline{\mathbb{F}}_p)$ such that the reduction map induces an isomorphism $\text{End}_W(\widetilde{E}^{\text{can}}) \stackrel{\sim}{\longrightarrow} \text{End}(E)$. For the canonical lift, it is known that $\text{End}_{\mathbb{C}}(\tilde{E}^{\text{can}}) = \text{End}_{W(\overline{\mathbb{F}}_p)}(\tilde{E}^{\text{can}}) = \text{End}_{\overline{\mathbb{F}}_p}(E)$ is an order $\mathcal{O}_{K,n}$ of the conductor n which is prime to p and that $K := \text{End}_{\overline{\mathbb{F}}_p}(E) \otimes_{\mathbb{Z}} \mathbb{Q}$ is an imaginary quadratic field for which p is split. Therefore the discriminant $d(\tilde{E}^{\text{can}})$ of $\text{End}_{\mathbb{C}}(\tilde{E}^{\text{can}})$ (which equals that for $\text{End}_{\overline{\mathbb{F}}_p}(E)$) is given

by nD_K , where D_K is the discriminant, and importantly we see that p nD_K (see Theorem 12, p.182 of $[La87]$). Furthermore for two ordinary elliptic curves E, E' , the reduction map induces an isomorphism

$$
\mathrm{Hom}_{\mathbb{C}}(\widetilde{E}^{\mathrm{can}},\widetilde{E}'^{\mathrm{can}}) \simeq \mathrm{Hom}_{W(\overline{\mathbb{F}}_p)}(\widetilde{E}^{\mathrm{can}},\widetilde{E}'^{\mathrm{can}}) \stackrel{\sim}{\longrightarrow} \mathrm{Hom}(E,E')
$$

since the Serre-Tate local coordinates $q(E), q(E')$ for E, E' satisfy $q(E) = q(E') = 1$ respectively (cf. Theorem 2.1 and also last a few lines in p.180 of [\[Katz81\]](#page-50-29)).

On the other hand if an elliptic curve E over $\mathbb C$ has CM by an order in an imaginary quadratic field K, then $\text{End}_{\mathbb{C}}(E) = \mathcal{O}_{K, np^s}$ for some positive integer n which is coprime to p and $s \in \mathbb{Z}_{\geq 0}$. Assume that p is split in K or equivalently that E has a good p-ordinary reduction $E/\overline{\mathbb{F}}_p$ (notice that one can take a smooth integral model over the ring \mathcal{O}_L of integers of some number field L among the isomorphism class of E since its j-invariant is an algebraic integer). Then we see that

(7.7)
$$
\mathcal{O}_{K, np^s} = \mathrm{End}_{\mathbb{C}}(\tilde{E}) = \mathrm{End}_{\mathcal{O}_L}(\tilde{E}) \hookrightarrow \mathrm{End}_{\overline{\mathbb{F}}_p}(E) = \mathcal{O}_{K, n}.
$$

by Theorem 12, p.182 of [\[La87\]](#page-50-28) again. The elliptic curve \tilde{E} is also a lift of an ordinary elliptic curve E and it is called a quasi-canonical lift of level s for E (cf. [\[Gro86\]](#page-50-30), [\[Yu95\]](#page-51-14)). It is known by Section 6 of [\[Gro86\]](#page-50-30) or Proposition 3.5 of [\[Meu07\]](#page-50-31) (see also p.97 of [\[Meu07\]](#page-50-31)) that the number of isomorphism classes of quasi-canonical lifts of level s for an ordinary elliptic curve E is given as follows:

(7.8)
$$
\frac{\sharp \mathcal{O}_{K, np^s}^{\times}}{\sharp \mathcal{O}_{K, n}^{\times}} \sharp (\mathbb{Z}/p^s \mathbb{Z})^{\times} = \frac{\sharp \mathrm{Aut}(\tilde{E})}{\sharp \mathrm{Aut}(E)} (p^s - p^{s-1}).
$$

Let t be the local parameter for the local deformation of quasi-canonical lift of p -divisible group of E and let j be the parameter of the coarse moduli space \mathbb{A}_j defined by j-invariant. Then we have the relation $j - j(E) = t^{\frac{\sharp \text{Aut}(E)}{2}}$. This explains the appearance of $\frac{\sharp \text{Aut}(\tilde{E})}{\sharp \text{Aut}(E)}$ in [\(7.8\)](#page-39-0).

From now on, we will count the number of lifts to quasi-canonical lifts for given two isogenies $f_1, f_2: E \to E'$ where E and E' are ordinary elliptic curves. Let $(z_i, w_i) = (z(f_i), w(f_i))$ be an element of \mathbb{Z}_p^2 for f_i 's with $i = 1, 2$, as explained in the paragraph following Lemma [7.10.](#page-37-2) We write $z_i = u_i p^{a_i}$ and $w_i = v_i p^{b_i}$, where u_i, v_i are units in \mathbb{Z}_p . Let T be the symmetric matrix associated to (f_1, f_2) as Equation [\(7.6\)](#page-38-1). Note that Lemma [7.8](#page-37-1) confirms that T is nonsingular in our situation. Put $r = r_T$. For two isogenies f_1, f_2 given as above we denote by $N(s, s'; f_1, f_2)$ the number of isomorphism classes of quasi-canonical lifts $(\tilde{E}_s, \tilde{E}_{s'})$ of level s, s' respectively $(0 \leq s, s' \leq r)$ such that f_1, f_2 lift to isogenies from E_s to $E_{s'}$.

Proposition 7.12. Keep the notation being as above. Assume that $a_1 = \min\{a_1, a_2, b_1, b_2\}$.

(1) If $b_1 < b_2$ and $a_2 + b_1 \le a_1 + b_2$, then we have that

$$
N(s,s';f_1,f_2) = \begin{cases} \n\frac{\sharp \mathcal{O}_{K,np^s}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \frac{\sharp \mathcal{O}_{K,np^{s'}}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \sharp (\mathbb{Z}/p^s \mathbb{Z})^{\times} & \text{if } 0 \le s \le a_1 \text{ and } 0 \le s' \le b_1; \\
\frac{\sharp \mathcal{O}_{K,np^s}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \frac{\sharp \mathcal{O}_{K,np^{s'}}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \sharp (\mathbb{Z}/p^s \mathbb{Z})^{\times} \cdot p^{b_1} & \text{if } a_1 < s \le a_2 \text{ and } s' = s + b_1 - a_1; \\
\frac{\sharp \mathcal{O}_{K,np^s}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \frac{\sharp \mathcal{O}_{K,np^{s'}}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \sharp (\mathbb{Z}/p^s \mathbb{Z})^{\times} \cdot p^{b_1} & \text{if } a_2 < s \le r - b_1 \\
0 & \text{and } s' = s + b_1 - a_1 = s + b_2 - a_2; \\
0 & \text{otherwise.} \n\end{cases}
$$

(2) If $b_1 < b_2$ and $a_2 + b_1 > a_1 + b_2$, then $r = a_1 + b_2$ and we have that

$$
N(s, s'; f_1, f_2) = \begin{cases} \n\frac{\sharp \mathcal{O}_{K, np^s}^{\times}}{\sharp \mathcal{O}_{K, n}^{\times}} \frac{\sharp \mathcal{O}_{K, np^{s'}}^{\times}}{\sharp \mathcal{O}_{K, n}^{\times}} \sharp (\mathbb{Z}/p^s \mathbb{Z})^{\times} \cdot \sharp (\mathbb{Z}/p^{s'} \mathbb{Z})^{\times} & \text{if } 0 \le s \le a_1 \text{ and } 0 \le s' \le b_1; \\
\frac{\sharp \mathcal{O}_{K, np^s}^{\times}}{\sharp \mathcal{O}_{K, n}^{\times}} \frac{\sharp \mathcal{O}_{K, np^{s'}}^{\times}}{\sharp \mathcal{O}_{K, n}^{\times}} \sharp (\mathbb{Z}/p^s \mathbb{Z})^{\times} \cdot p^{b_1} & \text{if } a_1 < s \le r - b_1 \text{ and } s' = s + b_1 - a_1; \\
0 & \text{otherwise.} \n\end{cases}
$$

(3) If $b_1 \geq b_2$ and $a_2 + b_1 \leq a_1 + b_2$, then we have that

$$
N(s,s';f_1,f_2) = \begin{cases} \frac{\sharp \mathcal{O}_{K,np^s}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \frac{\sharp \mathcal{O}_{K,np^s}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \sharp (\mathbb{Z}/p^s\mathbb{Z})^{\times} \cdot \sharp (\mathbb{Z}/p^{s'}\mathbb{Z})^{\times} & \text{if } 0 \le s' \le b_2 \text{ and } 0 \le s \le a_1; \\ \frac{\sharp \mathcal{O}_{K,np^s}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \frac{\sharp \mathcal{O}_{K,n}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \sharp (\mathbb{Z}/p^s\mathbb{Z})^{\times} \cdot p^{a_1} & \text{if } b_2 < s' \le b_1 \text{ and } s = s' - b_2 + a_2; \\ \frac{\sharp \mathcal{O}_{K,np^s}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \frac{\sharp \mathcal{O}_{K,np^s}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \sharp (\mathbb{Z}/p^s\mathbb{Z})^{\times} \cdot p^{a_1} & \text{if } b_1 < s' \le r - a_1 \\ 0 & \text{otherwise.} \end{cases}
$$

(4) If $b_1 \geq b_2$ and $a_2 + b_1 > a_1 + b_2$, then $r = a_1 + b_2$ and we have that

$$
N(s,s';f_1,f_2) = \begin{cases} \frac{\sharp \mathcal{O}_{K,nps}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \frac{\sharp \mathcal{O}_{K,nps'}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \sharp (\mathbb{Z}/p^s\mathbb{Z})^{\times} \cdot \sharp (\mathbb{Z}/p^{s'}\mathbb{Z})^{\times} & \text{if } 0 \le s' \le b_2 \text{ and } 0 \le s \le a_1; \\ \frac{\sharp \mathcal{O}_{K,nps}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \frac{\sharp \mathcal{O}_{K,nps'}^{\times}}{\sharp \mathcal{O}_{K,n}^{\times}} \sharp (\mathbb{Z}/p^s\mathbb{Z})^{\times} \cdot p^{a_1} & \text{if } b_2 < s' \le r - a_1 \text{ and } s = s' - b_2 + a_2; \\ 0 & \text{otherwise.} \end{cases}
$$

Proof. We choose quasi-canonical lifts \tilde{E}_s and $\tilde{E}'_{s'}$ of level s and s' for E and E' respectively. Let $q_s := q(\tilde{E}_s)$ and $q_{s'} := (\tilde{E}'_{s'})$ be the Serre-Tate coordinates for \tilde{E}_s and $\tilde{E}'_{s'}$ respectively. Then q_s (resp. $q_{s'}$) is a primitive $p^{s'}$ (resp. $p^{s'}$)-th root of unity in $\overline{\mathbb{Q}}_p$ by Proposition 3.5-(3) of [\[Meu07\]](#page-50-31). The condition to lift f_1 and f_2 as an element of $Hom(\tilde{E}_s, \tilde{E}'_{s'})$ is given by the following two equations:

(7.9)
$$
q_s^{z_1} = q_{s'}^{w_1}, \ q_s^{z_2} = q_{s'}^{w_2}.
$$

Here, we follow notations of page 140 of [\[Rap07\]](#page-51-9).

Firstly we consider the case when $b_1 \leq b_2$ and $a_2 + b_1 \leq a_1 + b_2$. If $s \leq a_1$, then $q_s^{z_1} = q_s^{z_2} = 1$ since $a_1 \le a_2$ by the assumption. Thus any $q_{s'}$ with $0 \le s' \le \min\{b_1, b_2\} = b_1$ also satisfies the second equation in [\(7.9\)](#page-40-0). In addition, such a $q_{s'}$ runs over all primitive $p^{s'}$ -roots of unity. Thus we have

$$
N(s,s';f_1,f_2)=\frac{\sharp \mathcal{O}_{K, np^s}^{\times}}{\sharp \mathcal{O}_{K, n}^{\times}}\frac{\sharp \mathcal{O}_{K, np^{s'}}^{\times}}{\sharp \mathcal{O}_{K, n}^{\times}}\sharp (\mathbb{Z}/p^s\mathbb{Z})^{\times}\cdot \sharp (\mathbb{Z}/p^{s'}\mathbb{Z})^{\times}
$$

for s, s' satisfying $0 \le s \le a_1$ and $0 \le s' \le b_1$.

If $a_1 < s \le a_2$, then the first equation in [\(7.9\)](#page-40-0) gives the equality $s' = s + b_1 - a_1$ but there are p^{b_1} numbers of $q_{s'}$ since $q_{s'}^{z_1} = q_{s'}^{w_1} = (q_{s'-b_1}^{v_1})^{p^{b_1}}$. Notice that $q_s^{z_2} = 1$ since $s \leq a_2$. On the other hand $s' = s + b_1 - a_1 \le a_2 + b_1 - a_1 \le b_2$ since we have assumed that $a_2 + b_1 \le a_1 + b_2$. Hence the second equation in [\(7.9\)](#page-40-0) is automatically fulfilled. Thus

$$
N(s,s';f_1,f_2)=\frac{\sharp \mathcal{O}_{K, np^s}^{\times}}{\sharp \mathcal{O}_{K, n}^{\times}}\frac{\sharp \mathcal{O}_{K, np^{s'}}^{\times}}{\sharp \mathcal{O}_{K, n}^{\times}}\sharp (\mathbb{Z}/p^s\mathbb{Z})^{\times}\cdot p^{b_1}
$$

where $s' = s + b_1 - a_1$.

Assume that $s > a_2$. Equation [\(7.9\)](#page-40-0) implies $s = s' - b_1 + a_1 = s' - b_2 + a_2$. In particular $a_2 + b_1 = a_1 + b_2$. In this case $a_2 + b_1 \leq r$ since $r = \text{ord}_p(z_1w_2 - z_2w_1)$. As discussed in the previous case there are p^{b_1} numbers of $q_{s'}$. With the notation fixed we may write $u_1v_2 - u_2v_1 = p^{r-(a_2+b_1)}\alpha$ for some $\alpha \in \mathbb{Z}_p^{\times}$. For such a $q_{s'}$ we compute

$$
q_{s'}^{w_2} = q_{s'-b_2}^{v_2} = q_{s'-b_2}^{v_2v_1v_1^{-1}} = q_{s'-b_1}^{p^{b_2-b_1}v_2v_1v_1^{-1}} = q_{s-a_1}^{p^{b_2-b_1}v_2u_1v_1^{-1}} = q_{s-a_1-b_2+b_1}^{v_2u_1v_1^{-1}} = q_{s-a_2}^{(v_1u_2+p^{r-(a_2+b_1)}\alpha)v_1^{-1}}
$$

$$
= q_{s-a_2}^{u_2+p^{r-(a_2+b_1)}\alpha v_1^{-1}} = q_{s}^{z_2+p^{r-b_1}\alpha v_1^{-1}}.
$$

It follows from this that if Equation [\(7.9\)](#page-40-0) has a solution, then it forces s to be $s \le r - b_1$ (note that $q_s^{z_2} = q_{s'}^{w_2}$ and the above equation imply $q_s^{p^{r-b_1} \alpha v_1^{-1}} = 1$). Otherwise (hence when $s > r - b_1$) there is no solution of [\(7.9\)](#page-40-0).

Next we assume that $b_1 < b_2$ and $a_2 + b_1 > a_1 + b_2$. Then $a_1 + b_2 = r$ since $r = \text{ord}_p(z_1w_2 - z_2w_1)$. A similar computation done before gives the desired results and hence details are omitted.

In the case when $b_1 \geq b_2$ we may replace the role of s with one of s' and carry out a similar \Box computation.

Corollary 7.13. We have the following formula:

$$
\sum_{0 \le s, s' \le r} \frac{N(s, s'; f_1, f_2)}{\sharp \mathrm{Aut}(\tilde{E}_s) \sharp \mathrm{Aut}(\tilde{E}'_s)} = \frac{p^r}{\sharp \mathrm{Aut}(E) \sharp \mathrm{Aut}(E')}
$$

Proof. By exchanging f_1 and f_2 , if necessary, we may assume that either a_1 or b_1 is the minimum among a_1, a_2, b_1, b_2 . We first assume that a_1 is the minimum. Let us treat the case when $b_1 < b_2$ and $a_2 + b_1 \leq a_1 + b_2$. The other cases can be handled similarly so that we may skip them. Using Proposition [7.12,](#page-39-1) the left hand side turns to be

$$
\sum_{0\leq s\leq a_1 \atop 0\leq s'\leq b_1} \frac{1}{\sharp \mathcal{O}_{K,n}^\times \sharp \mathcal{O}_{K,n}^\times} \sharp (\mathbb{Z}/p^s\mathbb{Z})^\times \cdot \sharp (\mathbb{Z}/p^{s'}\mathbb{Z})^\times + \sum_{a_1 < s\leq r-b_1 \atop 0\leq s'\leq b_1} \frac{1}{\sharp \mathcal{O}_{K,n}^\times \sharp \mathcal{O}_{K,n}^\times} \sharp (\mathbb{Z}/p^s\mathbb{Z})^\times \cdot p^{b_1} \newline = \frac{1}{\sharp \mathcal{O}_{K,n}^\times \sharp \mathcal{O}_{K,n}^\times} (p^{a_1+b_1} + p^{b_1}(p^{r-b_1} - p^{a_1})) = \frac{p^r}{\sharp \mathcal{O}_{K,n}^\times \sharp \mathcal{O}_{K,n}^\times} = \frac{p^r}{\sharp \mathrm{Aut}(E)\sharp \mathrm{Aut}(E')}.
$$

We next assume that b_1 is the minimum. Let f_i^{\vee} be the dual isogeny of f_i . Then we can see that $N(s, s'; f_1, f_2) = N(s', s; f_1^{\vee}, f_2^{\vee}).$

Thus the argument used in the above case gives the desired formula.

 \Box

.

7.5. The intersection number in the ordinary locus. The intersection number $(T_{m_1,\mathbb{C}}, T_{m_2,\mathbb{C}})$ is described in Proposition 2.4 of [\[GK93\]](#page-50-0) and it turns out to be the sum of the Fourier coefficients of the Siegel-Eisenstein series for Sp_4/\mathbb{Q} . We first consider the contribution coming from the ordinary part. For a symmetric positive definite (2×2) -half-integral matrix T (namely diagonal entries are integer and anti-diagonal entries are elements in $\frac{1}{2}\mathbb{Z}$), we define $\chi_T(p)$ by

$$
\chi_T(p) = \begin{cases} 1 & \text{(if } p \text{ is split in } \mathbb{Q}(\sqrt{-\det(2T))})\\ 0 & \text{(if } p \text{ is ramified in } \mathbb{Q}(\sqrt{-\det(2T))})\\ -1 & \text{(if } p \text{ is inert in } \mathbb{Q}(\sqrt{-\det(2T))}) \end{cases}
$$

Theorem 7.14. Assume that m_1m_2 is not a square. Then for any prime number p, we have that

$$
\sum_{\substack{y = ((E, E'), f_1, f_2) \\ (E, E') : (\text{ord}) \\ \deg(f_i) = m_i}} \frac{\text{IM}_{p,y}}{\sharp \text{Aut}(E) \sharp \text{Aut}(E')} = \frac{1}{288} \sum_{\substack{T \in \text{Sym}_2(\mathbb{Z}) > 0 \\ \text{diag}(T) = (m_1, m_2), \chi_T(p) = 1}} c(T).
$$

Here $c(T)$ is the Fourier coefficient of the Siegel-Eisenstein series of weight 2 with respect to $\text{Sp}_4(\mathbb{Z})$ (*cf.* [\[Nag92\]](#page-51-12)).

Proof. With the observation in CM elliptic curves having good ordinary reduction at p (which we call it *p*-ordinary and denote it by *p*-ord throughout this proof), we first obtain

(7.10)
$$
\sum_{\substack{(\tilde{E}, \tilde{E}'), f_1, f_2) \\ (\tilde{E}, \tilde{E}') \text{C:CM } p-\text{ord} \\ \deg(f_i)=m_i}} \frac{1}{\sharp \text{Aut}(\tilde{E}) \sharp \text{Aut}(\tilde{E}')} = \frac{1}{288} \sum_{\substack{T \in \text{Sym}_2(\mathbb{Z})_{>0} \\ \text{diag}(T)=(m_1, m_2), \chi_T(p)=1}} c(T)
$$

by (2.19), p. 231 of [\[GK93\]](#page-50-0).

For a pair of p-ordinary elliptic curves (\tilde{E}, \tilde{E}') having two isogenies f_1 and f_2 as in LHS of the above equation, we already observed in Section [7.4](#page-38-0) that $\tilde{E} = \tilde{E}_s$ and $\tilde{E}' = \tilde{E}'_{s'}$ for suitable s and s', where \tilde{E}_s and $\tilde{E}_{s'}$ are quasi-canonical lifts of level s and s' for ordinary elliptic curves E and E' respectively. Since the reduction of endomorphism groups is injective, the reductions of f_1 and f_2 (which are also denoted by f_1 and f_2 respectively) are isogenies from E to E' with the same degrees. On the other hand, any two isogenies $f_1, f_2 : E \to E'$ over \mathbb{F}_p can be lifted to those having the same degree defined over $\mathbb C$ by choosing canonical lifts (cf. Section [7.4\)](#page-38-0). Therefore, using Corollary [7.13,](#page-41-1) LHS of Equation [\(7.10\)](#page-42-1) turns to be

$$
\sum_{\substack{((\tilde{E}, \tilde{E}'), f_1, f_2) \\ (\tilde{E}, \tilde{E}') \text{ (CCM } p \text{ - ord } \\ \deg(f_i) = m_i}}} \frac{1}{\sharp \text{Aut}(\tilde{E}) \sharp \text{Aut}(\tilde{E}')} = \sum_{s, s' \geq 0} \sum_{\substack{((\tilde{E}_s, \tilde{E}'_s), f_1, f_2) \\ \deg(f_i) = m_i \\ (\tilde{E}, E'), f_1, f_2) \\ (\tilde{E}, E'), f_1, f_2) \\ (\tilde{E}, E') \text{ (prod } j \text{ - eq. } f_1, f_2)}} \frac{1}{\sharp \text{Aut}(\tilde{E}) \sharp \text{Aut}(\tilde{E}') } = \sum_{\substack{((E, E'), f_1, f_2) \\ (\tilde{E}, E'), f_1, f_2) \\ (\tilde{E}, E') \text{ (prod } j \text{ - eq. } f_1, f_2) \\ \deg(f_i) = m_i \\ \deg(f_i) = m_i}} \frac{1}{\sharp \text{Aut}(\tilde{E}) \sharp \text{Aut}(\tilde{E}') } = \sum_{\substack{((E, E'), f_1, f_2) \\ (\tilde{E}, E') \text{ (prod } j \text{ - eq. } f_1, f_2) \\ (\tilde{E}, E') \text{ (prod } j \text{ - eq. } f_1, f_2) \\ (\tilde{E}, E') \text{ (prod } j \text{ - eq. } f_1, f_2) \\ \deg(f_i) = m_i}} \frac{1}{\sharp \text{Aut}(\tilde{E}) \sharp \text{Aut}(E')}.
$$

Here, a_i and b_i are as explained in the paragraph just before Proposition [7.12](#page-39-1) associated to (f_1, f_2) . This, combined with Equation (7.10) , completes the proof.

7.6. The intersection number in the supersingular locus. We next consider the contribution coming from the supersingular part of Equation [\(7.3\)](#page-36-2). Let us remind readers that our definition of the Siegel series is the same as Katsurada's one given in [\[Kat99\]](#page-50-9). This convention has been used in Wedhorn's article [\[Wed07-2\]](#page-51-10) and Nagaoka's article [\[Nag92\]](#page-51-12). The difference to Kitaoka's Siegel series given in [\[Kit93\]](#page-50-20) is explained in the last line of Section 4.3, p. 43 of [\[Wed07-2\]](#page-51-10).

Theorem 7.15. Assume that p is odd and that m_1m_2 is not a square. Then we have that

(7.11)
$$
\sum_{\substack{y=(E,E'),f_1,f_2) \ \text{deg}(f_i)=m_i}} \frac{\text{IM}_{p,y}}{\# \text{Aut}(E) \# \text{Aut}(E')} = \frac{1}{288} \cdot \sum_{\substack{T \in \text{Sym}_2(\mathbb{Z})>0 \ \text{diag}(T)=(m_1,m_2),\chi_T(p)=-1,0}} c(T),
$$

where T is a half-integral symmetric matrix.

Proof. We will proceed our proof without assuming $p > 2$. The assumption will be made later when it is needed.

Let (E, E') be a pair of two supersingular elliptic curves and $f_1, f_2 : E \longrightarrow E'$ be isogenies with $deg(f_i) = m_i$. Let T be the (2×2) half-integral symmetric matrix associated to (f_1, f_2) . Since (E, E') are supersingular elliptic curves and $D := \text{Hom}(E) \otimes \mathbb{Q} = \text{End}(E') \otimes \mathbb{Q}$ is ramified at p, the prime p has to be inert or ramified in $\mathbb{Q}(\sqrt{-\det T})$ (cf. Theorem 12, page 182 of [\[La87\]](#page-50-28)). Hence $\chi_T(p) = -1$ or 0.

By the argument explained in page 23 of $[G\ddot{o}r07-2]$ and Proposition [7.11.](#page-38-2)(2), we see that (7.12)

$$
\sum_{\substack{y=(E,E'),f_1,f_2)\\(E,E')/\overline{\mathbb{F}}_p:(\text{ss})\\ \deg(f_i)=m_i}}\frac{\text{IM}_{p,y}}{\sharp \text{Aut}(E)\sharp \text{Aut}(E')}=\sum_{\substack{T\in \text{Sym}_2(\mathbb{Z})>0\\ \text{diag}(T)=(m_1,m_2),\chi_T(p)=-1,0}}\Big(\sum_{(E,E'):(\text{ss})}\frac{R_{\text{Hom}(E,E')}(T)}{\sharp \text{Aut}(E)\sharp \text{Aut}(E')} \mathcal{T}_{a_1,a_2}\Big),
$$

where $(a_1, a_2) = \text{GK}(T \otimes \mathbb{Z}_p)$.

Let $\mathcal{F}'_{T,l}(X)$ be the Siegel series associated to the local completion $T \otimes_{\mathbb{Z}} \mathbb{Z}_l$ for any finite place l. Firstly, by Theorem [6.7](#page-32-0) we have

(7.13)
$$
\mathcal{T}_{a_1,a_2} = \frac{-1}{p-1} \cdot \mathcal{F}'_{T,p}(1/p).
$$

Secondly, by using the argument used in the proof of Theorem 4.3 in [\[Wed07-2\]](#page-51-10) and the Minkowski-Siegel formula (cf. Theorem 4.2 of loc.cit.), we have

$$
(7.14) \quad \sum_{(E,E'):(ss)} \frac{R_{\text{Hom}(E,E')}(T)}{\sharp \text{Aut}(E)\sharp \text{Aut}(E')} = \frac{1}{3^2 \cdot 2^4} \cdot \left(\frac{p-1}{p}\right)^2 \cdot \frac{\pi^{\frac{7}{2}}}{\Gamma(2)\Gamma(3/2)} \cdot \det(T)^{1/2} \cdot \prod_{l < \infty} \alpha_l(T, O_D).
$$

Here O_D is a maximal order in D. Then $O_D \otimes_{\mathbb{Z}} \mathbb{Z}_p$ is the maximal order in the quaternion division algebra over \mathbb{Q}_p and $O_D \otimes_{\mathbb{Z}} \mathbb{Z}_l$ with $l \neq p$ is isomorphic to H_2 , the hyperbolic space of rank 4.

Thus we have

(7.15)
$$
\prod_{l < \infty} \alpha_l(T, O_D) = \alpha_{T, p}(T, O_D) \cdot \prod_{l < \infty, l \neq p} \mathcal{F}_{T, l}(1/l^2).
$$

We now plug in Equations $(7.13)-(7.15)$ $(7.13)-(7.15)$ into Equation (7.12) then we have the following:

$$
(7.16) \sum_{\substack{y=(E,E'),f_1,f_2) \ \deg(f_i)=m_i}} \frac{\text{IM}_{p,y}}{\sharp \text{Aut}(E)\sharp \text{Aut}(E')} =
$$

\n
$$
\sum_{\substack{T\in \text{Sym}_2(\mathbb{Z})>0 \ \deg(T)=(m_1,m_2),\chi_T(p)=-1,0}} \left(-\frac{1}{3^2 \cdot 2^4} \cdot \frac{p-1}{p^2} \cdot \frac{\pi^{\frac{7}{2}}}{\Gamma(2)\Gamma(3/2)} \cdot \det(T)^{1/2} \cdot \alpha_{T,p}(T,O_D) \cdot \mathcal{F}'_{T,p}(1/p) \cdot \prod_{l < \infty, l \neq p} \mathcal{F}_{T,l}(1/l^2)\right).
$$

On the other hand, by the functional equation of the Siegel series (cf. Theorem 4.1 in [\[Ike17\]](#page-50-32)) combined with Theorem 0.1 of [\[IK1\]](#page-50-12) for anisotropic binary quadratic lattice, we have

(7.17)
$$
\mathcal{F}'_{T,p}(1/p) = \begin{cases} -\frac{p^{3+|\mathrm{GK}(T_p)|/2}}{(p-1)(p+1)} \mathcal{F}_{T,p}(\frac{1}{p^2}) & \text{if } |\mathrm{GK}(T_p)| \text{ is even;}\\ -\frac{2p^{4+|\mathrm{GK}(T_p)|-1)/2}}{(p-1)(p+1)^2} \mathcal{F}_{T,p}(\frac{1}{p^2}) & \text{if } |\mathrm{GK}(T_p)| \text{ is odd.} \end{cases}
$$

Here, $T_p = T \otimes \mathbb{Z}_p$ and $|\text{GK}(T_p)| = a_1 + a_2$ if $\text{GK}(T_p) = (a_1, a_2)$.

From now on, we make our assumption of $p > 2$, in order to use Lemma [7.16](#page-44-0) to compute $\alpha_p(L, O_p)$. Then Equation [\(7.16\)](#page-43-3) turns to be (for $p > 2$)

$$
\sum_{\substack{y=(E,E'),f_1,f_2) \ (E,E')\vert\bar{x}_p:(ss)}} \frac{\text{IM}_{p,y}}{\sharp \text{Aut}(E)\sharp \text{Aut}(E')} = \frac{1}{3^2 2^3} \cdot \frac{\pi^{\frac{7}{2}}}{\Gamma(2)\Gamma(3/2)} \cdot \sum_{\substack{T \in \text{Sym}_2(\mathbb{Z})>0 \ \text{diag}(T)=(m_1,m_2),\chi_T(p)=-1,0}} \det(T)^{1/2} \prod_{l<\infty} \mathcal{F}_{T,l}(1/l^2) \newline \text{deg}(f_i)=m_i
$$
\n(7.18)\n
$$
= \frac{\pi^3}{3^2 2^2} \cdot \sum_{\substack{T \in \text{Sym}_2(\mathbb{Z})>0 \ \text{diag}(T)=(m_1,m_2),\chi_T(p)=-1,0}} \det(T)^{1/2} \prod_{l<\infty} \mathcal{F}_{T,l}(1/l^2),
$$

since $\Gamma(3/2) =$ $\sqrt{\pi}$ $\frac{\sqrt{\pi}}{2}$. Notice that the product $\prod_{k \leq x}$ $l<\infty$ $\mathcal{F}_{T,l}(1/l^2)$ coincides with $\lim_{s\to 0} \alpha^{(2)}(2s+2,T)$ defined in Section 1.3 of [\[Nag92\]](#page-51-12) (note that the limit $\lim_{s\to 0}$ will be taken after the analytic continuation given by Kauffold's theorem, cf. Theorem 1.3.1 of [\[Nag92\]](#page-51-12)). The formula (2.5.6), p.84 of [\[Nag92\]](#page-51-12) says

$$
\lim_{s \to 0} \alpha^{(2)}(2s + 2, T) = 36\pi^{-3} \det(T)^{-1/2} \frac{1}{288} c(T)
$$

where $\frac{1}{288}c(T) = \sum_{n=-\infty} dH(\frac{\det(T)}{d^2})$ d |cont (T) $\frac{\partial (2)}{\partial l^2}$) in the notation of [\[Nag92\]](#page-51-12).

In conclusion, we have

(7.19)
$$
\sum_{\substack{y=(E,E'),f_1,f_2)\\(E,E')_{/\overline{\mathbb{F}}_p}:\text{(ss)}} \frac{\text{IM}_{p,y}}{\sharp \text{Aut}(E)\sharp \text{Aut}(E')} = \frac{1}{3^2 2^5} \cdot \sum_{\substack{T \in \text{Sym}_2(\mathbb{Z})_{>0}\\ \text{diag}(T)=(m_1,m_2),\chi_T(p)=-1,0}} c(T).
$$

Here, $c(T)$ is the Fourier coefficient of the Siegel-Eisenstein series for Sp_4 of weight 2, with respect to the half-integral symmetric matrix T .

The proof of Theorem [7.3.](#page-34-2) The identity of Theorem [7.3](#page-34-2) now follows by combining Theorems [7.14](#page-41-2) and [7.15.](#page-42-2)

Lemma 7.16. Let $p > 2$. Let (L, q_L) be an anisotropic quadratic \mathbb{Z}_p -lattice of rank 2. Assume that $(L \otimes_{\mathbb{Z}_p} \mathbb{Q}_p, q_L \otimes_{\mathbb{Z}_p} \mathbb{Q}_p)$ is nondegenerate. Let (O_D, q_D) be the quadratic lattice, where O_D is the maximal order of the quaternion division algebra D over \mathbb{Q}_p with q_D the reduced norm on D. Then the local density $\alpha_p(L, O_D)$ is given as follows:

$$
\alpha_p(L, O_D) = \begin{cases} p^{\frac{-d}{2}} \cdot 2(1 + \frac{1}{p}) & \text{if } d \text{ is even;}\\ p^{\frac{-(d-1)}{2}} \cdot (1 + \frac{1}{p})^2 & \text{if } d = 2d' + 1 \text{ is odd.} \end{cases}
$$

Here, $d = |GK(L)|$.

Proof. Let L' be a sublattice of L of rank 2. Then By Theorem 5.6.4.(d) of [\[Kit93\]](#page-50-20), we have, for any prime p including $p = 2$,

$$
\alpha_p(L', O_D) = p^{-[L:L']}\cdot \alpha_p(L, O_D).
$$

Indeed Theorem 5.6.4.(d) of [\[Kit93\]](#page-50-20) says inequality but in our case of anisotropic lattice, the inequality turns to be the equality.

Thus we may assume that L is a maximal lattice so that $GK(L) = (a_1, a_2)$ has only three possibilities: $(a_1, a_2) = (0, 0), (0, 1), (1, 1)$. We will use the local density formula of Yang in [\[Yan98\]](#page-51-15) $(p > 2)$.

From now on we assume that p is odd. Then L is diagonalizable so that the exponential order of each diagonal entry is a_i . Based on Theorem 7.1 of [\[Yan98\]](#page-51-15), we have

$$
\alpha_p(L, O_D) = \begin{cases}\n2(1 + \frac{1}{p}) & \text{if } (a_1, a_2) = (0, 0); \\
\frac{2}{p} \left(1 + \frac{1}{p}\right) & \text{if } (a_1, a_2) = (1, 1); \\
\left(1 + \frac{1}{p}\right)^2 & \text{if } (a_1, a_2) = (0, 1).\n\end{cases}
$$

This completes the proof. \Box

Remark 7.17. (1) B. Conrad informed us that Theorem [7.3](#page-34-2) is true when $(m_1, m_2) = 1$ with any prime p (cf. [\[Con17\]](#page-50-33)), by proving that the scheme $Spec \mathbb{Z}[x,y]/(\varphi_{m_1}, \varphi_{m_2})$ is flat. On the other hand, Theorem [7.3](#page-34-2) is true when $m_1, m_2 \leq 9$ with any prime p by numerical

calculation given in Appendix [A.](#page-49-0)

(2) We note that Theorem [7.3](#page-34-2) is a combination of Theorems [7.14](#page-41-2) and [7.15,](#page-42-2) and the identity of Theorem [7.14](#page-41-2) holds for $p = 2$. In the proof of Theorem [7.15,](#page-42-2) the only place we make the assumption $p > 2$ is the usage of Lemma [7.16.](#page-44-0)

On the other hand, one can also compute $\alpha_p(L, O_D)$ of Lemma [7.16](#page-44-0) with $p = 2$ by using the local density formula given in [\[Yan04\]](#page-51-1), which is more complicated than that of [\[Yan98\]](#page-51-15) $(p > 2)$. Consequently the explicit computation of $\alpha_p(L, O_D)$ when $p = 2$ would directly yield the identity of Theorem [7.3.](#page-34-2)

8. Application 2: Local intersection multiplicities on the special fiber

In this section we recall the special cycles on the Shimura variety for $\text{GSpin}(n, 2)$, $0 \leq n \leq 3$ defined by Kudla and Rapoport with a collaborator Yang. We refer the articles [\[KRY99\]](#page-50-4),[\[KRY06\]](#page-50-5),[\[KR99\]](#page-50-6),[\[KR00\]](#page-50-7) for $n = 0, 1, 2, 3$ respectively and the readers are supposed to be familiar with these references. Let (n_1, \ldots, n_r) be a partition of $n+1$, hence $n_i \geq 1$, $n_1 + \cdots + n_r = n+1$. We always consider $r = 1$ when $n \leq 1$.

Let V be a quadratic form over $\mathbb Z$ with the signature $(n, 2)$ over $\mathbb R$ considered in each paper. Let $G = \text{GSpin}(V)$ be the generalized spinor group associated to V. Let p be an odd prime so that G is smooth over \mathbb{Z}_p . Then for any neat open compact subgroup $K^p \subset \hat{G}(\mathbb{A}_f^p)$ $_{f}^{p}$) and a hyperspecial open compact subgroup K_p of $G(\mathbb{Q}_p)$ defined by a suitable structure on $G(\mathbb{Q}_p)$ in each paper (cf. p. 704, line 9 of [\[KR00\]](#page-50-7) for $n = 3$, $d_i \in \text{Sym}_{n_i}(\mathbb{Q})$ and open compact subgroups $\omega_i \subset V(\mathbb{A}_f^p)$ $_{f}^{p}$ $)$ ^{n_i} which are invariant under K^p , one can associate the special cycles $\mathcal{Z}(d_i, \omega_i)$ and consider the intersection of them:

$$
\mathcal{Z} = \mathcal{Z}(d_1, \omega_1) \times_{\mathcal{M}} \cdots \times_{\mathcal{M}} \mathcal{Z}(d_r, \omega_r) = \coprod_{\substack{T \in \text{Sym}_{n+1}(\mathbb{Z}_{(p)}) \ge 0 \\ \text{diag}(T) = (d_1, \dots, d_r)}} \mathcal{Z}(T, \omega)
$$

where M is the integral model over $\mathbb{Z}_{(p)}$ for the Shimura variety associated to $(G, K^p K_p)$ and $\omega = {\{\omega_i\}}_{i=1}^r$ (cf. Section 2,3 of [\[KR00\]](#page-50-7) for $n = 3$). Since $n \leq 3$, the Shimura variety M is of PEL type, namely a moduli space of abelian varieties with endomorphism structure by $\mathcal O$ which is a maximal order of $M_2(B_0)$, B_F , B_0 , or K for $n = 3, 2, 1, 0$ respectively. Here B_0 (resp. B_F) is a quaternion algebra over $\mathbb Q$ (resp. over a real quadratic field F) and K is an imaginary quadratic field. Any geometric point $\xi = \text{Spec } k$ on Z in characteristic p consists of quintuple $(A, \iota, \lambda, \overline{\eta}^p, \mathbf{j}')$ satisfying the following conditions;

- (1) A is an abelian variety of dimension 2^n over k considered up to prime to p isogeny;
- (2) $\iota: \mathcal{O}\otimes \mathbb{Z}_{(p)} \hookrightarrow \text{End}_k(A)\otimes \mathbb{Z}_{(p)}$ is a ring homomorphism such that

$$
\det(\iota(c) : \text{Lie}(A)) = N^0(c)^2
$$

for any $c \in \mathcal{O}$ where N^0 is the reduced norm of \mathcal{O} .

- (3) λ is a \mathbb{Z}_{r}^{\times} $\chi_{(p)}^{\times}$ -class of a prime to p isogeny $A \longrightarrow A^*$ such that $n' \lambda$ comes from an ample line bundle on A for some $n' \in \mathbb{Z}$. Here A^* is the dual abelian variety of A;
- (4) $\mathbf{j}' = (\mathbf{j}_1, \ldots, \mathbf{j}_r)$ and $\mathbf{j}_i \in \text{End}_k(A)^{n_i}$ is a vector of special endomorphisms for $i = 1, \ldots, r$ such that $q(\mathbf{j}_i) = d_i$, where the quadratic form q is defined by the Rosati-involution \star with $q(x)$ id_A = $x \circ x^*$ for any $x \in$ End_k(A) (cf. Lemma 2.2 of [\[KR00\]](#page-50-7) for $n = 3$). Here a special endomorphism is defined to be an endomorphism f on A which satisfies $f^* = f$ and $tr^0(f) = 0$ where \star stands for the Rosati-involution with respect to λ and tr^0 means the reduced trace of $\text{End}_k((A, \iota))^{\text{op}}$ (note that special endomorphisms can be regarded as elements in $\text{End}_k((A, \iota))^{\text{op}}$ (cf. (2.13) of [\[KR00\]](#page-50-7) for $n = 3$)).
- (5) $\bar{\eta}^p = {\eta^p k \mid k \in K^p}$ is a K^p -class of a *O*-linear isomorphism $\eta^p : V^p(A) := \prod_{\ell \neq p} T_{\ell}(A) \otimes$ $\mathbb{Q}_{\ell} \stackrel{\sim}{\longrightarrow} \mathcal{O} {\otimes} \mathbb{A}_f^p$ \int_{f}^{p} . Here the action of K^{p} on \mathcal{O} is defined by the Clifford structure of $\mathrm{GSpin}(n,2)$ in each case. It is known that $\text{End}_{\mathcal{O}}(\mathcal{O}\otimes_{\mathbb{Z}} \mathbb{A}_f)$ contains $V(\mathbb{A}_f)$. We require that $(\eta^p)^*(\mathbf{j}_i) :=$ $\eta^p \circ \mathbf{j}_i|_{V^p(A)} \circ (\eta^p)^{-1}$ belongs to ω_i .

By Theorem 0.1 in [\[KR00\]](#page-50-7) for $n = 3$, Theorem 6.1 in [\[KR99\]](#page-50-6) for $n = 2$, Theorem 3.6.1 in [\[KRY06\]](#page-50-5) for $n = 1$, and Proposition 5.9 in [\[KRY99\]](#page-50-4) for $n = 0$ we know a criterion for T which yields that any geometric point of $\mathcal{Z}(T) := \mathcal{Z}(T,\omega)$ is isolated. From now on we assume this condition. For any geometric point ξ on $\mathcal{Z}(T)$ we define the local intersection multiplicity of $\mathcal{Z}(T)$ at ξ by

$$
e(\mathcal{Z}(T),\xi) := \mathrm{length}_{\mathbb{Z}_{(p)}} \mathcal{O}_{\mathcal{Z}(T),\xi}
$$

where $\mathcal{O}_{\mathcal{Z}(T),\xi}$ is the localization of $\mathcal{O}_{\mathcal{Z}(T)}$ at ξ . By the assumption $e(\mathcal{Z}(T),\xi)$ is finite.

To compute $e(\mathcal{Z}(T), \xi)$, we need to consider the formal completion of it along ξ to apply the deformation theory. Put $\mathbf{j}' = (f_1, \ldots, f_{n+1})$ for simplicity. By the assumption of each reference as above, we see that A is isomorphic to a product of supersingular elliptic curves E over $\overline{\mathbb{F}}_p$. Let $G = A$ be the formal group associated to A and we denote by f_i the corresponding special endomorphism on G for each special endomorphism f_i via a natural algebra homomorphism

(8.1)
$$
\mathrm{End}_k((A,\iota))^{\mathrm{op}} \otimes_{\mathbb{Z}} \mathbb{Z}_p \hookrightarrow \mathrm{End}((G,F)) \subset \mathcal{O} \otimes_{\mathbb{Z}} \mathbb{Q}_p
$$

where F is the Frobenius endomorphism and $\text{End}((G, F))$ stands for the set of endomorphisms on G commuting F (cf. (5.13), p. 726 of [\[KR00\]](#page-50-7) for $n = 3$). The \mathbb{Z}_p submodule L' spanned by $\{\widehat{f}_i\}$ in $\text{End}(G)$ endows with the structure as a quadratic space $L' = (L, q_{L'})$ which comes from the Clifford structure. For instance $xy + yx = (x, y)_L'$ for any $x, y \in L'$.

Since p is odd, there exists a basis of L' such that $q_{L'}$ is isometric to $T' = diag(u_1 p^{a_1}, \ldots, u_{n+1} p^{a_{n+1}})$ over \mathbb{Z}_p with integers $a_1 \leq \cdots \leq a_{n+1}$ and with units $u_i, 1 \leq i \leq n+1$ in \mathbb{Z}_p . Then the Gross-Keating invariant of T' is given simply by $GK(L') = GK(T') = (a_1, \ldots, a_{n+1}) \in \mathbb{Z}_{\geq 0}^{n+1}$. Accordingly we can take an optimal basis $\varphi_1, \ldots, \varphi_{n+1}$ in End(G) such that $\varphi_i^2 = q_{L'}(\varphi_i) = u_i p^{a_i}$ $(1 \leq i \leq n+1)$.

Let $R = W(\overline{\mathbb{F}}_p)[[t_1,\ldots,t_n]]$ be the universal deformation ring of G on $\text{CLN}_{W(\overline{\mathbb{F}}_p)}$ which is isomorphic to the strict completion of $\mathcal{O}_\mathcal{M}$ at ξ . This follows from Serre-Tate theorem (Theorem 1.2.1) of [\[Katz81\]](#page-50-29)), Theorem 2.3.3, p. 242 of [\[O70\]](#page-51-6), and the fact that the Shimura variety $\mathcal M$ is a fine moduli space by the assumption on the compact open subgroup K^p . Let $\mathcal G$ be its universal family. Here we make the convention that $R = W(\overline{\mathbb{F}}_p)$ when $n = 0$.

We denote by $I = I(\varphi_1, \ldots, \varphi_{n+1})$ the minimal ideal of R such that all φ_i 's are liftable to special endomorphisms on $\mathcal G$ modulo I . By the theorem of Serre and Tate it is easy to see that

$$
e(\mathcal{Z}(T'),\xi)=\mathrm{length}_{W(\overline{\mathbb{F}}_p)}R/I
$$

(cf. (6.1) of [\[KR00\]](#page-50-7)). By Section 6 of [KR00] for $n = 3$, the proof of Theorem 6.1 of [\[KR99\]](#page-50-6) for $n = 2$, Theorem 3.6.1 of [\[KRY06\]](#page-50-5) for $n = 1$, and Theorem 5.11 of [\[KRY99\]](#page-50-4) for $n = 0$, it turns out that $e(\mathcal{Z}(T'),\xi)$ depends only on $GK(T') = (a_1,\ldots,a_{n+1})$. Hence we may write it for

(8.2)
$$
e(a_1,\ldots,a_{n+1}) := e(\mathcal{Z}(T'),\xi) = \text{length}_{W(\overline{\mathbb{F}}_p)} R/I(\varphi_1,\ldots,\varphi_{n+1}).
$$

This will be checked in Theorem [8.3](#page-48-0) below. When $a_{n+1} \geq 2$, we see that there exist $\varphi'_{n+1} \in \text{End}(G)$ such that $\varphi_{n+1} = p\varphi'_{n+1}$ and that $L' = \langle \varphi_1, \ldots, \varphi_n, \varphi'_{n+1} \rangle$ makes a sublattice of L with $GK(L') =$ $(a_1, ..., a_n) \cup (a_{n+1}-2)$ where $(a_1, ..., a_n) \cup (a_{n+1}-2)$ means the re-ordering of $\{a_1, ..., a_n, a_{n+1}-2\}$ to be the non-decreasing sequence. Therefore we have

length
$$
\text{length}_{W(\overline{\mathbb{F}}_p)} R/I(\varphi_1,\ldots,\varphi'_{n+1})=e((a_1,\ldots,a_n)\cup(a_{n+1}-2)).
$$

Our interest is to understand the difference between

length
$$
W(\overline{\mathbb{F}}_p)R/I(\varphi_1,\ldots,\varphi_{n+1})
$$
 and length $W(\overline{\mathbb{F}}_p)R/I(\varphi_1,\ldots,\varphi'_{n+1})$

in terms of the local intersection multiplicity of special cycles over a finite field. This motivates us to consider the following situation in special cycles in the special fiber $\mathcal{M}_{\mathbb{F}_p}$. Assume that $1 \leq n \leq 3$. Let (n_1, \ldots, n_r) be a partition of n, hence $n_i \geq 1$, $n_1 + \cdots + n_r = n$. We always consider $r = 1$ when $n = 1$. For $d_i \in \text{Sym}_{n_i}(\mathbb{Z}_{(p)})$ $(1 \leq i \leq r)$ let us consider the closed subscheme in $\mathcal{M}_{\mathbb{F}_p}$ given by

$$
\mathcal{Z}(d_1,\omega_1)_{\mathbb{F}_p}\times_{\mathcal{M}_{\mathbb{F}_p}}\cdots\times_{\mathcal{M}_{\mathbb{F}_p}}\mathcal{Z}(d_r,\omega_r)_{\mathbb{F}_p}=\coprod_{T\in\text{Sym}_n(\mathbb{Z}_{(p)})_{\geq 0}\atop \text{diag}(T)=(d_1,\ldots,d_r)}\mathcal{Z}(T,\omega)_{\mathbb{F}_p}
$$

for $1 \leq n \leq 3$. Any geometric point on $\mathcal{Z}(T,\omega)_{\mathbb{F}_p}$ is similarly a quintuple $(A,\iota,\lambda,\overline{\eta}^p,\mathbf{j})$ as before but in this case we replace $n + 1$ endomorphisms with n endomorphisms j which is related to the fourth condition (4). In this section, when $n \leq 3$ and the underlying space $A = A_x$ of a geometric point x is superspecial, we will study that the multiplicity

$$
e(\mathcal{Z}(T,\omega)_{\mathbb{F}_p},x):=\mathrm{length}_{\overline{\mathbb{F}}_p}\mathcal{O}_{\mathcal{Z}(T,\omega)_{\mathbb{F}_p},x}
$$

is finite and that it depends only on $GK(T \otimes \mathbb{Z}_p)$ under some conditions. Let us confirm this as follows.

Recall that G is the formal group of A. Let $R_p = \mathbb{F}_p[[t_1, \ldots, t_n]]$ be the universal deformation of G on $\text{CLN}_{\overline{\mathbb{F}}_p}$ which is isomorphic to the completion of $\mathcal{O}_{\mathcal{M}_{\overline{\mathbb{F}}_p}}$ at x. We put $R_p = \mathbb{F}_p$ when $n = 0$. Clearly $\mathcal{G}_p := \mathcal{G} \otimes \mathbb{F}_p$ is the local deformation of G over R_p . Let $\varphi_1, \ldots, \varphi_n$ be an optimal basis of the lattice consisting of n special endomorphisms j in the data of x via (8.1) .

We define the minimal ideal $I_p = I_p(\varphi_1, \ldots, \varphi_n)$ of R_p such that all φ_i 's $(1 \leq i \leq n)$ are liftable to special endomorphisms on \mathcal{G}_p modulo I_p . As before we put $I_p = \{0\}$ when $n = 0$. By definition we see that

$$
e(\mathcal{Z}(T,\omega)_{\mathbb{F}_p},x)=\text{length}_{\overline{\mathbb{F}}_p}R_p/I_p.
$$

Proposition 8.1. Let T be an element in $\text{Sym}_n(\mathbb{Z}_{(p)})$. For any geometric point $x = (A_x, \iota, \lambda, \overline{\eta}^p, \mathbf{j})$ on $\mathcal{Z}(T,\omega)_{\mathbb{F}_p}$ let $L=(L,q_L)$ be the quadratic space over \mathbb{Z}_p corresponding to the special endomorphisms j with $GK(L) = GK(T \otimes \mathbb{Z}_p) = (a_0, \ldots, a_{n-1})$. Assume that A_x is superspecial and q_L represents 1. Then it holds that

$$
e(\mathcal{Z}(T,\omega)_{\mathbb{F}_p},x)=\mathcal{T}_{b_1,b_2}
$$

where

$$
(b_1, b_2) = \begin{cases} (a_1, a_2) & \text{if } n = 3 \\ (0, a_{n-1}) & \text{if } n = 1, 2 \\ (0, 0) & \text{if } n = 0 \end{cases}.
$$

Proof. Let us first consider the case of $n = 3$. We follow the argument in page 733 of [\[KR00\]](#page-50-7). By our assumption q_L represents 1 over \mathbb{Z}_p . This implies $a_0 = 0$. In this case the formal group G of A_x decomposes into \widehat{A}_0^4 where \widehat{A}_0 is a formal group of dimension 2 and height 4 equipped with a principal quasi polarization $\lambda_{\widehat{A}_0}$ (cf. Section 4 of [\[KR00\]](#page-50-7)). As explained right after (6.3) of [KR00]

there exists $x_0 \in L$ such that $q_L(x_0) = 1$. Then the idempotents $e_0 = \frac{1}{2}$ $\frac{1}{2}(1+x_0), e_1=\frac{1}{2}$ $rac{1}{2}(1-x_0)$ induce the further decomposition of \hat{A}_0 as

$$
\widehat{A}_0 \simeq e_0 \widehat{A}_0 \times e_1 \widehat{A}_0 \simeq G_0^2
$$

where G_0 is a formal group of dimension 1 and height 2. Put $M_0 = \langle x_0 \rangle^{\perp}$ in L which is of rank 2 and $GK(M_0) = (a_1, a_2)$. By the Clifford structure for the special endomorphisms we see that $xx_0 + x_0x = \langle x, x_0 \rangle = 0$ for any $x \in M_0$. It follows from this that $xe_0 = e_1x$. Therefore M_0 can be regarded as a sublattice in $Hom(e_0A_0, e_1A_0) \simeq End(G_0)$ which is the maximal order of a unique quaternion division algebra over \mathbb{Q}_p . Since any principal quasi-polarization deforms automatically the deformation problem on $\text{CLN}_{\overline{\mathbb{F}}_p}$ of x is same as one of (G_0, M_0) . This shows that

(8.3)
$$
e(\mathcal{Z}(T,\omega)_{\mathbb{F}_p},x)=\mathcal{T}_{a_1,a_2}.
$$

Hence we have the claim.

The remaining cases will be done similarly in which the arguments of Theorem 6.1 of [\[KR99\]](#page-50-6), Theorem 3.6.1 of [\[KRY06\]](#page-50-5), and Proposition 5.9 of [KRY06] for $n = 2, 1$ and $n = 0$ should be consulted respectively.

Corollary 8.2. Keep the notation and the assumptions in Proposition [8.1.](#page-47-0) Then the geometric point x is isolated in $\mathcal{Z}(T)_{\overline{\mathbb{F}}_p}$.

Proposition 8.3. Let T' be an element in $Sym_{n+1}(\mathbb{Z}_{(p)})$. For any isolated geometric point $\xi =$ $(A_{\xi},\iota,\lambda,\overline{\eta}^p,\vec{\jmath})$ of $\mathcal{Z}(T')$ let L' be the \mathbb{Z}_p -lattice corresponding to the special endomorphisms $\vec{\jmath}$ with $GK(L') = GK(T' \otimes \mathbb{Z}_p) = (a_0, \ldots, a_n).$ Then for any $T \in Sym_n(\mathbb{Z}_{(p)})$ with $GK(T \otimes \mathbb{Z}_p) =$ (a_0, \ldots, a_{n-1}) such that $T \otimes \mathbb{Z}_p$ comes from a sublattice of L' and a geometric point x of $\mathcal{Z}(T,\omega)_{\mathbb{F}_p}$ whose underlying abelian variety A_x is A_ξ , it holds that

$$
e(\mathcal{Z}(T,\omega)_{\mathbb{F}_p},x)=e(a_0,\ldots,a_n)-e(a_0,\ldots,a_n-2).
$$

Proof. When $n = 3$, by Theorem 0.1 or Corollary 5.15 of [\[KR00\]](#page-50-7), we see that $T' \otimes \mathbb{Z}_p$ represents 1 over \mathbb{Z}_p . This implies $a_0 = 0$. Let L' be the lattice in \mathbb{Q}_p corresponding to T' over \mathbb{Z}_p . Then it turns out that its Gross-Keating invariant becomes $GK(\tilde{L}') = (0, a_1, a_2, a_3)$ with nondecreasing integers $0 \le a_1 \le a_2 \le a_3$ which satisfy the condition that the parities of the three integers never be same. Then the argument in p. 733 of [\[KR00\]](#page-50-7) tells us that

(8.4)
$$
e(0, a_1, a_2, a_3) = \alpha_p(a_1, a_2, a_3)
$$

where α_p is the intersection number in Proposition 5.4 of [\[GK93\]](#page-50-0) as mentioned before. By Lemma 5.6 of [\[GK93\]](#page-50-0) and Proposition [8.1](#page-47-0)

$$
e(0, a_1, a_2, a_3) - e(0, a_1, a_2, a_3 - 2) = \mathcal{T}_{a_1, a_2} = e(\mathcal{Z}(T, \omega)_{\mathbb{F}_p}, x).
$$

When $n = 2$, by Theorem 6.1 of [\[KR99\]](#page-50-6), we see that $T' \otimes \mathbb{Z}_p$ is isometric to diag(1, $u_1 p^{a_1}, u_2 p^{a_2}$) over \mathbb{Z}_p with units u_1, u_2 in \mathbb{Z}_p . Let L' be the lattice in \mathbb{Q}_p corresponding to T' over \mathbb{Z}_p . Then it turns out that its Gross-Keating invariant becomes $GK(L') = (0, a_1, a_2)$ with nondecreasing integers $0 \le a_1 \le a_2$ which satisfy the condition that the parities of $0, a_2, a_3$ never be same. Then the argument in p. 195 loc.cit. shows that

(8.5)
$$
e(0, a_1, a_2) = \alpha_p(0, a_1, a_2)
$$

and similarly we have $e(0, a_1, a_2) - e(0, a_1, a_2 - 2) = \mathcal{T}_{0, a_1} = e(\mathcal{Z}(T, \omega)_{\mathbb{F}_p}, x).$

When $n = 1$, by Theorem 3.6.1 of [\[KRY06\]](#page-50-5), we see that $T' \otimes \mathbb{Z}_p$ is isometric to diag $(u_0 p^{a_0}, u_1 p^{a_1})$ over \mathbb{Z}_p with units u_0, u_1 in \mathbb{Z}_p . Let L be the lattice of rank three in \mathbb{Q}_p corresponding to $\text{diag}(1, T')$ over \mathbb{Z}_p . Then it turns out that its Gross-Keating invariant becomes $GK(L) = (0, a_0, a_1)$ with nondecreasing integers $0 \le a_0 \le a_1$ which satisfy the condition that the parities of $0, a_0, a_1$ never be same. Then Theorem 3.6.1 of loc.cit. shows that

(8.6)
$$
e(a_0, a_1) = \alpha_p(0, a_0, a_1)
$$

and similarly we have $e(a_0, a_1) - e(a_0, a_1 - 2) = \mathcal{T}_{0,a_0} = e(\mathcal{Z}(T,\omega)_{\mathbb{F}_p}, x)$.

Finally we consider the case when $n = 0$. By Proposition 5.9 of [\[KRY06\]](#page-50-5), we see that T' (it is denoted by t in the reference) satisfies that $a_0 := \text{ord}_p(t) \equiv 1 \text{ mod } 2$.

Let L be the lattice of rank three in \mathbb{Q}_p corresponding to diag(1,1,T') over \mathbb{Z}_p . Then it turns out that its Gross-Keating invariant becomes $GK(L) = (0, 0, a_3)$ and it satisfies that a_0 is not even. Then Theorem 5.11 of loc.cit. shows that

(8.7)
$$
e(a_0) = \alpha_p(0, 0, a_0)
$$

Then $e(a_0) - e(a_0 - 2) = 1 = e(\mathcal{Z}(T, \omega)_{\mathbb{F}_p}, x)$ by Lemma 5.6 of [\[GK93\]](#page-50-0) and the claim follows with the convention made when $n = 0$.

Assume that $T \in \text{Sym}_n(\mathbb{Z}_{(p)})$ satisfies the condition in Proposition [8.3.](#page-48-0) For n and Gross-Keating invariant for T in Equations $(8.4)-(8.7)$ $(8.4)-(8.7)$ we take an anisotropic lattice M of rank 2 with

$$
GK(M) = \begin{cases} (a_1, a_2) & \text{if } n = 3\\ (0, a_{n-1}) & \text{if } n = 1, 2\\ (0, 0) & \text{if } n = 0 \end{cases}
$$

Note that in our situation $T \otimes \mathbb{Z}_p$ is always anisotropic and hence we can apply the results in Section [6](#page-29-0) to $T \otimes \mathbb{Z}_p$. Then plugging Proposition [8.3](#page-48-0) with Section [6](#page-29-0) (cf. Theorem [6.7\)](#page-32-0) we have

Theorem 8.4. Keep the assumption in Proposition [8.3.](#page-48-0) Then

$$
e(\mathcal{Z}(T)_{\overline{\mathbb{F}}_p}, x) = \frac{-1}{p-1} \cdot \mathcal{F}'_M(\frac{1}{p}).
$$

Appendix A. The table of intersection numbers

For each positive integer m we denote by ψ_{m/n^2} the polynomial in $\mathbb{Z}[x, y]$ which appears as a factor of $\varphi_m = \prod$ $n^2|m$ ψ_{m/n^2} (cf. p.2 of [\[Vog07\]](#page-51-0)). In this appendix we give a table for

$$
\dim_{\mathbb{C}}\mathbb{C}[x,y]/(\psi_{m_1}, \psi_{m_2}), \ \dim_{\mathbb{F}_p} \mathbb{F}_p[x,y]/(\psi_{m_1}, \psi_{m_2})
$$

for $2 \le m_1 \le m_2 \le 9$ such that $m_1 m_2$ is not a square and $2 \le p < 50$. S. Yokoyama (cf. [\[Yok17\]](#page-51-16)) kindly computed both quantities and checked they coincide directly. From this computation with Corollary [7.2](#page-34-4) and Theorem 2.1 of [\[Vog07\]](#page-51-0) it is easy to see that

$$
(T_{m_1,p}, T_{m_2,p}) = \sum_{n_1^2 \mid m_1, n_2^2 \mid m_2} \dim_{\mathbb{F}_p} \mathbb{F}_p[x, y] / (\psi_{\frac{m_1}{n_1^2}}, \psi_{\frac{m_2}{n_2^2}})
$$

=
$$
\sum_{n_1^2 \mid m_1, n_2^2 \mid m_2} \dim_{\mathbb{C}} \mathbb{C}[x, y] / (\psi_{\frac{m_1}{n_1^2}}, \psi_{\frac{m_2}{n_2^2}}) = (T_{m_1, \mathbb{C}}, T_{m_2, \mathbb{C}})
$$

for m_1, m_2, p above including the case where $p = 2$. Put $d(m_1, m_2) := \dim_{\mathbb{C}} \mathbb{C}[x, y]/(\psi_{m_1}, \psi_{m_2}) =$ $\dim_{\mathbb{F}_p} \mathbb{F}_p[x, y]/(\psi_{m_1}, \psi_{m_2})$. We list up all of them as below:

Let us remark that $d(1, m) = d(m, 1) = \sum$ $d|m$ $\max\{d, \frac{m}{d}\}.$ By using this we see, for example, that $(T_{2,p}, T_{4,p}) = (T_{2,\mathbb{C}}, T_{4,\mathbb{C}}) = d(2,1) + d(2,4) = 4 + 28 = 32$ for any $p < 50$.

REFERENCES

- [ARGOS07] Argos Seminar on Intersections of Modular Correspondences, Held at the University of Bonn, Bonn, 2003-2004. Astérisque No. 312 (2007).
- [BLR90] S. Bosch, W. Lütkebohmert, and M. Raynaud, Néron Models, Ergeb. Math. Grenzgeb.(3) 21, Springer, Berlin, 1990
- [BY] J. Bruinier and T. Yang, Arithmetic degrees of special cycles and derivatives of Siegel Eisenstein series, [arXiv:1802.09489.](http://arxiv.org/abs/1802.09489)
- [Bou07] I. I. Bouw, *Invariants of ternary quadratic forms*, Astérisque 312 (2007) 113-137.
- [Cho15] S. Cho, Group schemes and local densities of quadratic lattices in residue characteristic 2, Compositio Math. Vol. 151 (2015) 793-827.
- [Cho16] S. Cho, Group schemes and local densities of ramified hermitian lattices in residue characteristic 2 Part I, Algebra & Number Theory, 10-3 (2016) 451-532.
- [CIKY1] S. Cho, T. Ikeda, H. Katsurada, and T. Yamauchi, An inductive formula of the Gross-Keating invariant of a quadratic form, preprint.
- [CIKY2] S. Cho, T. Ikeda, H. Katsurada, and T. Yamauchi, Remarks on the Extended Gross-Keating data and the Siegel series of a quadratic form, [arXiv:1709.02772.](http://arxiv.org/abs/1709.02772)
- [CL] P. Clark, Lectures on Shimura curves 1: Endomorphisms of elliptic curves, available at http://math.uga.edu/∼[pete/SC1-endomorphisms.pdf.](http://math.uga.edu/~pete/SC1-endomorphisms.pdf)
- [Con17] B. Conrad, Private communication in April 2017.
- [Dur44] W. H. Durfee, Congruence of quadratic forms over valuation rings, Duke Math. J. 11 (1944) 687-697.
- [GY00] W. T. Gan and J.-K. Yu, Group schemes and local densities, Duke Math. J. 105 (2000) 497-524.
- [Gro86] B-H. Gross, On canonical and quasi-canonical liftings, Invent math 84, 321-326 (1986).
- [GK93] B-H. Gross and K. Keating, On the intersection of modular correspondences, Invent math.112 (1993) 225-245.
- [Gör07-1] U. Görtz, A sum of representation numbers, Astérisque 312 (2007) 9-14.
- [Gör07-2] U. Görtz, Arithmetic intersection numbers, Astérisque 312 (2007) 15-24.
- [Han99] J. P. Hanke, An exact mass formula for orthogonal groups over number fields, Ph.D. dissertation, Princeton Univ. (1999) Princeton.
- [Ike17] T. Ikeda, On the functional equation of the Siegel series, J. Number Theory. 172 (2017) 44-62.
- [IK1] T. Ikeda and H. Katsurada, On the Gross-Keating invariant of a quadratic form over a non-archimedean local field, [arXiv:1504.07330,](http://arxiv.org/abs/1504.07330) to appear in Amer. J. Math.
- [IK2] T. Ikeda and H. Katsurada, Explicit formula for the Siegel series of a half-integral matrix over the ring of integers in a non-archimedian local field, [arXiv:1602.06617.](http://arxiv.org/abs/1602.06617)
- [Kat99] H. Katsurada, An explicit formula for Siegel series, American J. Math. 121 (1999) 415-452.
- [Katz81] N. Katz, Serre-Tate local moduli, Surfaces alg´ebriques, in: Lect. Notes in Math., vol. 868, Springer-Verlag, Berlin, 1981, pp. 138-202.
- [KM85] N. Katz and B. Mazur, Arithmetic of elliptic curves, Annals of Mathematics Studies, 1985.
- [Kit83] Y. Kitaoka, A note on local densities of quadratic forms, Nagoya Math. J. 92 (1983) 145-152.
- [Kit93] Y. Kitaoka, Arithmetic of Quadratic Forms, Cambridge Tracts in Math. 106, Cambridge Univ. Press, Cambridge (1993).
- [Kud97] S. Kudla, Central derivatives of Eisenstein series and height pairings, Ann of math. (2) 146, no.3 (1997) 545-646.
- [KR99] S. Kudla and M. Rapoport, Arithmetic of Hirzebruch-Zagier cycles, J. Reine Angew. Math. 515 (1999) 155-244.
- [KR00] S. Kudla and M. Rapoport, Cycles on Siegel 3-folds and derivatives of Eisenstein series, Ann. Sci. École Norm. Sup. (4) 33, no. 5 (2000) 695-756.
- [KRY99] S. Kudla and M. Rapoport, and T. Yang, On the Derivative of an Eisenstein Series of Weight One, Int. Math. Res. Not., 7 (1999) 347-385.
- [KRY06] S. Kudla and M. Rapoport, and T. Yang, Modular Forms and Special Cycles on Shimura Curves, Annals of Mathematics Studies, vol. 161. Princeton University Press, Princeton (2006).
- [La87] S. Lang, Elliptic functions. With an appendix by J. Tate. Second edition. Graduate Texts in Mathematics, 112. Springer-Verlag, New York, 1987.
- [LZ] C. Li and W. Zhang, Kudla-Rapoport cycles and derivatives of local densities, [arXiv:1908.01701.](http://arxiv.org/abs/1908.01701)
- [Meu07] V. Meusers, Canonical and quasi-canonical liftings in the split case, Astérisque 312 (2007) 87-98.
- [Nag92] S. Nagaoka, A note on the Siegel-Eisenstein series of weight 2 on $Sp_2(\mathbb{Z})$. Manuscripta Math. 77 (1992), no. 1, 71-88.
- [O'Me00] O. T. O'Meara, Introduction to Quadratic Forms, reprint of 1973 ed., Classics Math. (2000), Springer, Berlin
- [O70] F. Oort, Finite group schemes, local moduli for abelian varieties, and lifting problems. Compositio Math. 23 (1971), 265-296.
- [Pop11] M. Popa, Course on p-adic and motivic integration, UIC lecture note (2011), available at https://sites.math. northwestern.edu/empopa/571
- [Rap07] M. Rapoport, *Deformations of isogenies of formal groups*, Astérisque 312 (2007) 139-169.
- [Ro] A-M. Robert, A Course in p-adic Analysis, Graduate Texts in Mathematics, 198. Springer-Verlag, New York, 2000.
- [Sie35] C. Siegel, Über die analytische Theorie der quadratischen Formen, Ann. Math. 36 (1935) 527-606.
- [Sil09] J. Silverman, The Arithmetic of Elliptic Curves, Second edition. Graduate Texts in Mathematics, 106. Springer, Dordrecht, 2009. xx+513 pp.
- [Tam66] T. Tamagawa, Adéles, Algebraic Groups and Discontinuous Subgroups (Proc. Sympos. Pure Math., Boulder, Colo., 1965), vol. 9, Amer. Math. Soc., Providence, R.I. (1966) pp. 113-121.
- [Vog07] G. Vogel, *Modular polynomials*, Astérisque 312 (2007) 1-7.
- [Wed07-1] T. Wedhorn, *Calculation of representation densities*, Astérisque 312 (2007) 179-190.
- [Wed07-2] T. Wedhorn, The genus of the endomorphisms of a supersingular elliptic curve, Astérisque 312 (2007) 25-47.
- [Weil82] A. Weil, Adeles and Algebraic Groups, Progress in Mathematics, vol. 23, Birkhäuser, Boston Mass. (1982) pp. iii+126.
- [Yan98] T. Yang, An explicit formula for local densities of quadratic forms, J. Number Theory 72 (1998) 309-356.
- [Yan04] T. Yang, Local densities of 2-adic quadratic forms, J. Number Theory 108 (2004) 287-345.
- [Yok17] S. Yokoyama, Private communication in March 2017.
- [Yu95] J-.K. Yu, On the moduli of quasi-canonical liftings, Compositio Math. 96 (1995), no. 3, 293-321.
- [Yu08] J-.K. Yu, Tamagawa number Purdue lecture note (2008)

Sungmun Cho, Department of Mathematics, POSTECH, 77, Cheongam-ro, Nam-gu, Pohang-si, Gyeongsangbukdo, 37673, KOREA

E-mail address: sungmuncho12@gmail.com

Takuya Yamauchi, Mathematical Inst. Tohoku Univ., 6-3,Aoba, Aramaki, Aoba-Ku, Sendai 980-8578, JAPAN

E-mail address: tyamauchi@m.tohoku.ac.jp